

Exact finite volume expectation values of local operators in excited states

B. Pozsgay,^a I.M. Szécsényi^{b,c} and G. Takács^{a,d}

^a*MTA-BME “Momentum” Statistical Field Theory Research Group,
Budafoki út 8, 1111 Budapest, Hungary*

^b*Department of Mathematical Sciences, Durham University,
South Road, Durham, DH1 3LE, United Kingdom*

^c*Institute of Theoretical Physics, Eötvös Loránd University,
Pázmány Péter sétány 1/A, 1117 Budapest, Hungary*

^d*Department of Theoretical Physics, Budapest University of Technology and Economics,
Budafoki út 8, 1111 Budapest, Hungary*

E-mail: pozsgay.balazs@gmail.com, szistvan87@gmail.com,
takacsg@eik.bme.hu

ABSTRACT: We present a conjecture for the exact expression of finite volume expectation values in excited states in integrable quantum field theories, which is an extension of an earlier conjecture to the case of general diagonal factorized scattering with bound states and a nontrivial bootstrap structure. The conjectured expression is a spectral expansion which uses the exact form factors and the excited state thermodynamic Bethe Ansatz as building blocks. The conjecture is proven for the case of the trace of the energy-moment tensor. Concerning its validity for more general operators, we provide numerical evidence using the truncated conformal space approach. It is found that the expansion fails to be well-defined for small values of the volume in cases when the singularity structure of the TBA equations undergoes a non-trivial rearrangement under some critical value of the volume. Despite these shortcomings, the conjectured expression is expected to be valid for all volumes for most of the excited states, and as an expansion above the critical volume for the rest.

KEYWORDS: Exact S-Matrix, Bethe Ansatz, Integrable Field Theories

ARXIV EPRINT: [1412.8436](https://arxiv.org/abs/1412.8436)

Contents

1	Introduction	1
2	Finite volume expectation values in excited states: the conjecture	3
3	Equivalence of the form factor series and the TBA for the trace of the stress-energy tensor	7
3.1	$\langle \Theta \rangle$ from TBA	7
3.2	Isolating the singularity density terms	10
3.3	Dressed form factors of Θ	14
4	Finite volume expectation values in the T_2 model	18
4.1	Excited state TBA equations for a single type-1 state	18
4.1.1	General form and solution of the excited TBA equations	18
4.1.2	Zero-momentum state: desingularization in small volume	21
4.2	Densities and the conjecture for states with a single type-1 particle	22
5	Numerical results	23
5.1	Moving one-particle state, $s = 1$	25
5.2	Zero-momentum one-particle state, $s = 0$	25
6	Conclusions and outlook	27
A	Evaluation of connected diagonal form factors	32
A.1	Form factors of the T_2 model	32
A.2	Symmetric form factors	34
A.2.1	Denominator and minimal form factors	34
A.2.2	Symmetric polynomial part	35
A.2.3	Result for symmetric form factor	36
A.3	Evaluation of the connected diagonal form factors	37

1 Introduction

Finite temperature expectation values play an important role in various applications of quantum field theory, and have been intensively studied in recent years in the context of $1 + 1$ dimensional integrable quantum field theories. For the one-point function a series expansion was conjectured by Leclair and Mussardo in [1], based on the thermodynamic Bethe Ansatz [2] as applied to integrable quantum field theories [3], and the exact form factors from the bootstrap program [4–6]. Recently, the Leclair-Mussardo series found applications in quantum quenches [7, 8] and the investigations of one-dimensional quantum gases [9–11].

Another method to obtain finite temperature correlators is based on finite temperature form factors [12, 13]; so far, however, this approach seems limited to free theories such as the Ising model. Other approaches that can be used to construct finite temperature expectation values in integrable models use separation of variables [14, 15], or exploit the hidden Grassmannian/fermionic structure of the XXZ spin chain [16–18].

In [19, 20] a description of form factors in finite volume was introduced, which was subsequently used to prove the Leclair-Mussardo series [8]. This formalism was also used to compute the finite-temperature two-point function [21, 22], for which a systematic expansion was developed in [23, 24]. Besides further applications to two-point functions in condensed matter systems [25, 26], the finite volume form factor formalism have found numerous other applications, in computing one-point functions in the presence of boundaries [27], in the study of quantum quenches in field theories [28–30], and in the context of holographic duality [31, 32]. They also provide a useful tool for testing exact form factor solutions obtained from the form factor bootstrap, recently in the boundary [33] and defect [34] settings.

The finite volume form factor formalism introduced in [19, 20] only included the corrections that decay with a power of the volume; exponential corrections were neglected. However, soon after [19, 20] a method was proposed in [35] to construct certain exponential corrections, the so-called μ -terms which are related to the bootstrap fusion between the particles. It was found that these can be very important in determining matrix elements and resonance parameters [36].

However, for integrable quantum field theories one expects that an exact determination of finite volume form factors is also possible. For the Ising model on a lattice this was known before [37, 38]; however, until recently there have been no such results for generic models.

In [39] an extension of the Leclair-Mussardo series was conjectured to describe exact excited state expectation values (a.k.a. diagonal form factors) in finite volume. The methods of [39] apply to theories like the sinh-Gordon model which have no bound states in their bootstrap, and the thermodynamic Bethe Ansatz equations describing excited states in finite volume have a particularly simple structure [40].

In the present work we extend this conjecture to theories with a diagonal factorized scattering that have a nontrivial bootstrap structure. In such models, the excited state levels in finite volumes are described by excited TBA systems of the type introduced in [41–43]. This conjecture is verified in two ways. First, we show for the trace of stress-energy tensor the conjectured series is equivalent to the result obtained directly from the thermodynamic Bethe Ansatz. Second, we make use of the truncated conformal space approach (TCSA) [44] to get a nontrivial further check of the series. For the latter, we use the methods developed in [45], to which the interested reader is referred to for details.

The outline of the paper is as follows. In section 2 we introduce our notations and state the conjecture. In section 3 we present the proof that the conjectured series gives the same result as the thermodynamic Bethe Ansatz when evaluated for the trace of the stress-energy tensor. In section 4 we turn to the so-called T_2 model used as testing ground, and specify the expansion for the case of the excited state thermodynamic Bethe Ansatz of this particular field theory. The resulting expectation value are then compared to numerical results from

the TCSA in section 5, while section 6 contains our conclusions and outlook. As the method used for numerically evaluating the connected diagonal form factors contains some non-trivial tricks, and could be useful for other applications, it is presented in appendix A.

2 Finite volume expectation values in excited states: the conjecture

The Leclair-Mussardo series for the finite volume vacuum expectation value of a local operator in an integrable model with diagonal scattering and k species of massive particles takes the following form [1]:

$$\langle \mathcal{O} \rangle_L = \sum_{n_1, \dots, n_k=0}^{\infty} \left(\prod_{j=1}^k n_j! \right)^{-1} \int_{-\infty}^{\infty} \prod_{j=1}^{\tilde{N}} \frac{d\theta_j}{2\pi [1 + e^{\varepsilon_{\beta_j}(\theta_j)}]} F_{2n_1, \dots, 2n_k, c}^{\mathcal{O}}(\theta_1, \dots, \theta_{\tilde{N}}) \quad (2.1)$$

where $F_{2n_1, \dots, 2n_k, c}^{\mathcal{O}}$ are the connected diagonal form factors of the operator \mathcal{O} , n_i the number of particles of species i , $\tilde{N} = \sum n_i$ the total number of particles, and the j th particle has rapidity θ_j and species β_j . The $\varepsilon_{\alpha}(\theta)$ are the pseudo-energy functions satisfying the TBA integral equation [3]:

$$\varepsilon_{\alpha}(\theta) = m_{\alpha} L \cosh(\theta) - \sum_{\beta} \int \frac{d\theta'}{2\pi} \varphi_{\alpha\beta}(\theta - \theta') \log(1 + e^{-\varepsilon_{\beta}(\theta')}) \quad (2.2)$$

where the kernels are given by the logarithmic derivatives of the two-particle scattering phases

$$\varphi_{\alpha\beta}(\theta) = -i \frac{\partial}{\partial \theta} \log S_{\alpha\beta}(\theta). \quad (2.3)$$

The finite volume ground state energy is given by

$$E_{\text{TBA}}(L) = E(L) - \mathcal{B}L = - \sum_{\beta} \int \frac{d\theta'}{2\pi} m_{\beta} \cosh(\theta) \log(1 + e^{-\varepsilon_{\beta}(\theta)}) \quad (2.4)$$

where \mathcal{B} is the bulk energy density. The connected diagonal form factors are defined by regularizing the diagonal matrix element

$$F^{\mathcal{O}}(\theta_n + i\pi + \epsilon_n, \dots, \theta_1 + i\pi + \epsilon_1, \theta_1, \dots, \theta_n) \quad (2.5)$$

and retaining the terms which are independent of the ratios ϵ_i/ϵ_j . We remark that the form factor has a finite, but direction dependent limit when all the ϵ_i are taken to zero simultaneously, so the regularized matrix elements can only depend on their ratios.

From the work by Dorey and Tateo [42, 43] it is known that starting from the TBA equation of the ground state one can reach the Riemann surface of excited states by analytic continuation in the volume parameter; the same equations were also obtained in [41] using a different approach. When performing the analytic continuation singularities of the $\log(1 + e^{-\varepsilon_{\beta}})$ terms, corresponding to locations where $Y_{\beta} = e^{\varepsilon_{\beta}} = -1$, cross the integration

contour modifying the TBA equations as

$$\begin{aligned} \varepsilon_\alpha(\theta) &= m_\alpha L \cosh(\theta) - \sum_{j=1}^N \eta_j \log S_{\alpha\alpha_j}(\theta - \bar{\theta}_j) \\ &\quad - \sum_\beta \int \frac{d\theta'}{2\pi} \varphi_{\alpha\beta}(\theta - \theta') \log(1 + e^{-\varepsilon_\beta(\theta')}) \\ E_{\text{TBA}}(L) &= \sum_{j=1}^N i m_{\alpha_j} \eta_j \sinh(\bar{\theta}_j) - \sum_\beta \int \frac{d\theta}{2\pi} m_\beta \cosh(\theta) \log(1 + e^{-\varepsilon_\beta(\theta)}) \end{aligned} \quad (2.6)$$

where the $\bar{\theta}_i$ are the positions of the singularities of the pseudo-energy of species α_i , which satisfy the quantization conditions

$$\varepsilon_{\alpha_j}(\bar{\theta}_j) = i\pi(2I_j + 1) \quad I_j \in \mathbb{Z} \quad (2.7)$$

and the I_j can be viewed as quantum numbers specifying the excited state. Such singularities are called *active*; their contribution further depends on the orientation η_i of the integration contour around the singularity, which take the values ± 1 if the singularity crossed the real axis from above/below, respectively. The number and type of the active singularities depends on the excited state and the position of these singularities at a fixed volume L fully specifies the excited state. Therefore the corresponding finite volume state can be denoted as

$$|\bar{\theta}_1, \dots, \bar{\theta}_N\rangle_L. \quad (2.8)$$

The term $\log(1 + e^{-\varepsilon_\beta})$ also has singularities where the $Y_\beta = e^{\varepsilon_\beta} = 0$. In [42, 43] it was shown that whenever a singularity that corresponds to a zero of a Y function crosses the integration contour, it does not generate new source terms to the TBA equations, but only rearranges the active singularities already present. The only exception is when such a singularity pinches the integration contour; for more details about dealing with this situation see subsection 4.1.2.

The TBA system (2.6) can be recast in a universal functional form called the Y -system [46, 47]

$$\begin{aligned} Y_\alpha\left(\theta - \frac{i\pi}{h}\right) Y_\alpha\left(\theta + \frac{i\pi}{h}\right) &= \prod_{\beta=1}^k (1 + Y_\beta(\theta))^{I_{\alpha\beta}} \\ Y_\alpha(\theta) &= e^{\varepsilon_\alpha(\theta)} \end{aligned} \quad (2.9)$$

where h is the Coxeter number and $I_{\alpha\beta}$ is the incidence matrix of some diagram. In subsection 4.1.2 we shall use the fact that Y -system relates the positions of the two types of logarithmic singularities of the TBA equations.

The analytical continuation is expected to connect not only the energy, but also other quantities such as e.g. expectation values corresponding to the different finite volume levels. It was shown in [39] how to perform the residue integrals over the modified contours and re-sum the terms into a compact form for an analytically continued Leclair-Mussardo

conjecture. This calculation was carried out for the sinh-Gordon theory, where the excited TBA system is still a conjecture [40], but the result passes several consistency checks. Namely, the first e^{-mR} corrections in the infrared limit agree with theoretical expectations and the result also agrees with the TBA results for the trace of the stress-energy tensor.

Let us now state the conjecture for the general form of the finite volume expectation values in excited states. It contains two kind of quantities, the “dressed version” of the diagonal form factors and the densities of the active singularities.

Definition 1. The *dressed diagonal form factors of the local operator \mathcal{O}* are

$$\begin{aligned} \mathcal{D}_\varepsilon^\mathcal{O}(\bar{\theta}_1, \dots, \bar{\theta}_l) &:= \sum_{n_1, \dots, n_k=0}^\infty \frac{1}{\prod_j n_j!} \int_{-\infty}^\infty \prod_{j=1}^{\tilde{N}} \frac{d\theta_j}{2\pi [1 + e^{\varepsilon_{\beta_j}(\theta_j)}]} \\ &\times F_{2l, 2n_1, \dots, 2n_k, c}^\mathcal{O}(\bar{\theta}_1, \dots, \bar{\theta}_l, \theta_1, \dots, \theta_{\tilde{N}}) \\ \tilde{N} &= \sum_{j=1}^k n_j \end{aligned} \tag{2.10}$$

where $\bar{\theta}_i$ are a subset of the active singularities, with the i th one corresponding to species α_i .

To obtain the densities of the active singularities, consider the derivative matrix with respect to the singularity positions

$$\mathcal{K}_{jl} = \frac{\partial Q_j}{\partial \bar{\theta}_l} \tag{2.11}$$

of the quantization conditions (2.7)

$$Q_j = i\eta_j \varepsilon_{\alpha_j}(\bar{\theta}_j | \bar{\theta}_1, \dots, \bar{\theta}_N) \tag{2.12}$$

satisfied by the position of the active singularities. Here the notation $\varepsilon_{\alpha_j}(\bar{\theta}_j | \bar{\theta}_1, \dots, \bar{\theta}_N)$ means that we consider the pseudo-energy functions $\varepsilon_{\alpha_j}(\theta)$ obtained as a solution of the excited TBA system (3.2) for a fixed arrangement of singularity positions $\bar{\theta}_1, \dots, \bar{\theta}_N$, evaluated at $\theta = \bar{\theta}_j$. This is nothing else than the function specifying the exact quantization conditions for the singularity positions via (2.7), written in a more detailed form specifying its dependence on all the singularity positions explicitly.

Definition 2. The *density of active singularities* (in rapidity space) is the determinant of the derivative matrix

$$\rho(\bar{\theta}_1, \dots, \bar{\theta}_N) = \det \mathcal{K}_{jl}. \tag{2.13}$$

Definition 3. For any bipartite partition $\{\bar{\theta}_1, \dots, \bar{\theta}_N\} = \{\bar{\theta}_+\} \cup \{\bar{\theta}_-\}$ of the active singularities, the *restricted density of active singularities in the subset $\{\bar{\theta}_+\}$ relative to $\{\bar{\theta}_-\}$* is defined by

$$\rho(\{\bar{\theta}_+\} | \{\bar{\theta}_-\}) = \det \mathcal{K}_+ \tag{2.14}$$

where \mathcal{K}_+ is the submatrix corresponding to the subset of active singularities $\{\bar{\theta}_+\}$.

Using the above definitions, the main result can be stated as follows:

Conjecture 4. *The exact finite volume expectation values of an operator \mathcal{O} in any finite volume state can be written as*

$$\begin{aligned}
 {}_L \langle \bar{\theta}_1, \dots, \bar{\theta}_N | \mathcal{O} | \bar{\theta}_1, \dots, \bar{\theta}_N \rangle_L &= \frac{1}{\rho(\bar{\theta}_1, \dots, \bar{\theta}_N)} \\
 &\times \sum_{\{\bar{\theta}_+\} \cup \{\bar{\theta}_-\}} \mathcal{D}_\varepsilon^\mathcal{O}(\{\bar{\theta}_+\}) \rho(\{\bar{\theta}_-\} | \{\bar{\theta}_+\}). \quad (2.15)
 \end{aligned}$$

This is a generalization of an earlier conjecture in the sinh-Gordon theory [39]. In that previous work, the expression of the case of one and two particles was obtained by direct calculation of the analytic continuation of the Leclair-Mussardo series (2.1), and conjecturing the general form for any number of particles. For the sinh-Gordon system, the imaginary parts of the singularity positions are fixed as $\text{Im } \bar{\theta}_j = \pi/2$.

However, excited TBA systems of the form (2.6), (2.7) are much more general with the imaginary parts of the singularity positions themselves being dynamically determined by the quantization conditions (2.7). This corresponds to finite volume corrections known as μ -terms, which were considered in the form factor context in [35]. While it is intuitively natural to expect that a direct analytic continuation should lead to the formulae (2.15) in the general case, the details look much more complicated. In particular, we later observe that the naive extension of (2.15) for certain cases where the singularity positions undergo nontrivial rearrangements, is not valid. This shows that implementing the simple intuition above is not straightforward. Therefore we choose other ways to substantiate that the conjecture above is indeed correct:

1. The infrared limit of the formula reproduces previously known results for finite volume diagonal form factors which were obtained in [19, 20]. In large volume the imaginary parts of the active singularities tend to fixed values, which are determined by the poles of the scattering matrix [42, 43], and the real parts $\{\vartheta_j\} = \{\text{Re } \bar{\theta}_i\}$ of the singularity positions can be interpreted as rapidities of on-shell particles, where usually one particle is described by more than one singularity positions, which all have the same real parts. The quantization conditions reduce to the Bethe-Yang equations

$$m_{\alpha_j} \sinh \vartheta_j - i \sum_{k \neq j} \log S_{\alpha_j \alpha_k}(\vartheta_j - \vartheta_k) = 2\pi \bar{I}_j \quad (2.16)$$

where the momentum quantum numbers \bar{I}_j are related to the TBA quantum numbers I_j in (2.7). The density of active singularities specified in Definition 2 reduces to the usual density of states in rapidity space, while the restricted density in Definition 3 turns into the restricted density used in the diagonal form factor formula in [20].

In the same limit, the “dressed” diagonal form factors reduce to connected diagonal form factors, and for theories where particles are represented by a single active singularity, formula (2.15) reduces to the results obtained in [19, 20] for the finite-volume diagonal matrix elements, which are valid up to exponential corrections in the volume.

For theories with a nontrivial bootstrap structure the infrared limit gives even more: it uniquely specifies the correct construction of μ -term corrections, formerly considered in [35, 36]; we return to this point in the conclusions.

2. For the trace of the stress-energy tensor the conjecture is equivalent to the excited state TBA equations for any state, similarly to the Leclair-Mussardo series (2.1) which for the trace of the stress-energy tensor is equivalent to the ground state TBA (2.2), (2.4) [1]. The proof of this equivalence is given in section 3.
3. We stress that in contrast to [39], the generalized formula (2.15) can also be applied to integrable quantum field theories for which the finite volume matrix elements can be evaluated numerically using the truncated conformal space approach. Using the T_2 model considered in section 4, we perform the numerical comparison in section 5, providing evidence that our conjecture is indeed valid for operators different from the trace of the stress-energy tensor.

3 Equivalence of the form factor series and the TBA for the trace of the stress-energy tensor

In this section we present the equivalence of the conjectured form factor series for excited states (2.15) and the TBA equations for Θ the trace of the stress-energy tensor. We proceed in three steps. First we explicitly evaluate the TBA prediction for $\langle \Theta \rangle$, then recast it in a form which can be matched with the dependence of (2.15) on the densities, and then prove that the rest of the formula matches the dressed form factors of Θ .

3.1 $\langle \Theta \rangle$ from TBA

As described in [3] the expectation value of the trace of the stress-energy tensor can be expressed in the following way

$$\langle \Theta \rangle_L = \langle \Theta \rangle_\infty + 2\pi \left[\frac{E_{\text{TBA}}(L)}{L} + \frac{dE_{\text{TBA}}(L)}{dL} \right]. \tag{3.1}$$

For an excited state with N active singularities we obtain

$$\begin{aligned} E_{\text{TBA}}(L) &= \sum_{j=1}^N im_{\alpha_j} \eta_j \sinh(\bar{\theta}_j) - \sum_{\beta} \int \frac{d\theta}{2\pi} m_{\beta} \sinh(\theta) \frac{\partial_{\theta} \varepsilon_{\beta}(\theta)}{1 + e^{\varepsilon_{\beta}(\theta)}} \\ \frac{dE_{\text{TBA}}(L)}{dL} &= \sum_{j=1}^N im_{\alpha_j} \eta_j \cosh(\bar{\theta}_j) \frac{d\bar{\theta}_j}{dL} + \sum_{\beta} \int \frac{d\theta}{2\pi} m_{\beta} \cosh(\theta) \frac{\partial_L \varepsilon_{\beta}(\theta)}{1 + e^{\varepsilon_{\beta}(\theta)}} \end{aligned} \tag{3.2}$$

where we performed a partial integration in the energy expression. The derivatives of the pseudo-energy satisfy the following linear equations

$$\begin{aligned} \partial_\theta \varepsilon_\alpha(\theta) &= m_\alpha L \sinh(\theta) - \sum_{j=1}^N i\eta_j \varphi_{\alpha\alpha_j}(\theta - \bar{\theta}_j) + \sum_\beta \int \frac{d\theta'}{2\pi} \varphi_{\alpha\beta}(\theta - \theta') \frac{\partial_\theta \varepsilon_\beta(\theta')}{1 + e^{\varepsilon_\beta(\theta')}} \\ \partial_L \varepsilon_\alpha(\theta) &= m_\alpha \cosh(\theta) + \sum_{k=1}^N i\eta_k \varphi_{\alpha\alpha_k}(\theta - \bar{\theta}_k) \frac{d\bar{\theta}_k}{dL} \\ &\quad + \sum_\beta \int \frac{d\theta'}{2\pi} \varphi_{\alpha\beta}(\theta - \theta') \frac{\partial_L \varepsilon_\beta(\theta')}{1 + e^{\varepsilon_\beta(\theta')}}. \end{aligned} \tag{3.3}$$

The linearity of the above equations can be exploited by introducing new functions f satisfying the following equations

$$\begin{aligned} f_{s,\alpha}(\theta) &= m_\alpha \sinh(\theta) + \sum_\beta \int \frac{d\theta'}{2\pi} \varphi_{\alpha\beta}(\theta - \theta') \frac{f_{s,\beta}(\theta')}{1 + e^{\varepsilon_\beta(\theta')}} \\ f_{c,\alpha}(\theta) &= m_\alpha \cosh(\theta) + \sum_\beta \int \frac{d\theta'}{2\pi} \varphi_{\alpha\beta}(\theta - \theta') \frac{f_{c,\beta}(\theta')}{1 + e^{\varepsilon_\beta(\theta')}} \\ f_{j,\alpha}(\theta) &= \varphi_{\alpha\alpha_j}(\theta - \bar{\theta}_j) + \sum_\beta \int \frac{d\theta'}{2\pi} \varphi_{\alpha\beta}(\theta - \theta') \frac{f_{j,\beta}(\theta')}{1 + e^{\varepsilon_\beta(\theta')}} \quad j = 1, \dots, N \end{aligned} \tag{3.4}$$

which can be used to express the derivatives as

$$\begin{aligned} \partial_\theta \varepsilon_\alpha(\theta) &= L f_{s,\alpha}(\theta) + \sum_{j=1}^N (-i\eta_j) f_{j,\alpha}(\theta) \\ \partial_L \varepsilon_\alpha(\theta) &= f_{c,\alpha}(\theta) + \sum_{j=1}^N \left(i\eta_j \frac{d\bar{\theta}_j}{dL} \right) f_{j,\alpha}(\theta). \end{aligned} \tag{3.5}$$

Inserting these relation into (3.3)

$$\begin{aligned} E_{\text{TBA}}(L) &= \sum_{j=1}^N i m_{\alpha_j} \eta_j \sinh(\bar{\theta}_j) - \sum_\beta \int \frac{d\theta}{2\pi} m_\beta \sinh(\theta) \frac{L f_{s,\beta}(\theta) + \sum_{j=1}^N (-i\eta_j) f_{j,\beta}(\theta)}{1 + e^{\varepsilon_\beta(\theta)}} \\ \frac{dE_{\text{TBA}}(L)}{dL} &= \sum_{j=1}^N i m_{\alpha_j} \eta_j \cosh(\bar{\theta}_j) \frac{d\bar{\theta}_j}{dL} \\ &\quad + \sum_\beta \int \frac{d\theta}{2\pi} m_\beta \cosh(\theta) \frac{f_{c,\beta}(\theta) + \sum_{j=1}^N \left(i\eta_j \frac{d\bar{\theta}_j}{dL} \right) f_{j,\beta}(\theta)}{1 + e^{\varepsilon_\beta(\theta)}} \end{aligned} \tag{3.6}$$

the expectation value $\langle \Theta \rangle$ takes the form

$$\begin{aligned}
 \frac{\langle \Theta \rangle}{2\pi} &= \frac{\langle \Theta \rangle_\infty}{2\pi} + \sum_{j=1}^N \frac{i}{L} m_{\alpha_j} \eta_j \sinh(\bar{\theta}_j) - \sum_{\beta} \int \frac{d\theta}{2\pi} m_{\beta} \sinh(\theta) \frac{f_{s,\beta}(\theta) + \sum_{k=1}^N \left(-\frac{i}{L} \eta_j\right) f_{j,\beta}(\theta)}{1 + e^{\varepsilon_{\beta}(\theta)}} \\
 &+ \sum_{j=1}^N i m_{\alpha_j} \eta_j \cosh(\bar{\theta}_j) \frac{d\bar{\theta}_j}{dL} + \sum_{\beta} \int \frac{d\theta}{2\pi} m_{\beta} \cosh(\theta) \frac{f_{c,\beta}(\theta) + \sum_{j=1}^N \left(i \eta_j \frac{d\bar{\theta}_j}{dL}\right) f_{k,\beta}(\theta)}{1 + e^{\varepsilon_{\beta}(\theta)}} \\
 &= \frac{\langle \Theta \rangle_\infty}{2\pi} + \sum_{\beta} \int \frac{d\theta}{2\pi} \frac{m_{\beta} \cosh(\theta) f_{c,\beta}(\theta) - m_{\beta} \sinh(\theta) f_{s,\beta}(\theta)}{1 + e^{\varepsilon_{\beta}(\theta)}} \\
 &+ \sum_{j=1}^N \frac{i}{L} \eta_j \left[m_{\alpha_j} \sinh(\bar{\theta}_j) + \sum_{\beta} \int \frac{d\theta}{2\pi} m_{\beta} \sinh(\theta) \frac{f_{j,\beta}(\theta)}{1 + e^{\varepsilon_{\beta}(\theta)}} \right] \\
 &+ \sum_{k=1}^N i \eta_j \frac{d\bar{\theta}_j}{dL} \left[m_{\alpha_j} \cosh(\bar{\theta}_j) + \sum_{\beta} \int \frac{d\theta}{2\pi} m_{\beta} \cosh(\theta) \frac{f_{j,\beta}(\theta)}{1 + e^{\varepsilon_{\beta}(\theta)}} \right]. \tag{3.7}
 \end{aligned}$$

Using that the derivatives of phase shift are even functions $\varphi_{\alpha\beta}(\theta) = \varphi_{\beta\alpha}(-\theta)$, and the definition of f_i and $f_{s,c}$ one can easily see that

$$\begin{aligned}
 \sum_{\beta} \int \frac{d\theta}{2\pi} m_{\beta} \sinh(\theta) \frac{f_{j,\beta}(\theta)}{1 + e^{\varepsilon_{\beta}(\theta)}} &= \sum_{\beta} \int \frac{d\theta}{2\pi} \varphi_{\alpha_j\beta}(\bar{\theta}_j - \theta) \frac{f_{s,\beta}(\theta)}{1 + e^{\varepsilon_{\beta}(\theta)}} \\
 \sum_{\beta} \int \frac{d\theta}{2\pi} m_{\beta} \cosh(\theta) \frac{f_{j,\beta}(\theta)}{1 + e^{\varepsilon_{\beta}(\theta)}} &= \sum_{\beta} \int \frac{d\theta}{2\pi} \varphi_{\alpha_j\beta}(\bar{\theta}_j - \theta) \frac{f_{c,\beta}(\theta)}{1 + e^{\varepsilon_{\beta}(\theta)}} \tag{3.8}
 \end{aligned}$$

and so $\langle \Theta \rangle$ simplifies to

$$\begin{aligned}
 \frac{\langle \Theta \rangle}{2\pi} &= \frac{\langle \Theta \rangle_\infty}{2\pi} + \sum_{\beta} \int \frac{d\theta}{2\pi} \frac{m_{\beta} \cosh(\theta) f_{c,\beta}(\theta) - m_{\beta} \sinh(\theta) f_{s,\beta}(\theta)}{1 + e^{\varepsilon_{\beta}(\theta)}} \\
 &+ \sum_{j=1}^N \frac{i}{L} \eta_j f_{s,\alpha_j}(\bar{\theta}_j) + \sum_{k=1}^N i \eta_j f_{c,\alpha_j}(\bar{\theta}_j) \frac{d\bar{\theta}_j}{dL}. \tag{3.9}
 \end{aligned}$$

The derivatives of the active singularity positions can be expressed using the quantization conditions (2.12)

$$\begin{aligned}
 \frac{dQ_i}{dL} &= \sum_j \frac{\partial Q_i}{\partial \bar{\theta}_j} \frac{d\bar{\theta}_j}{dL} + \frac{\partial Q_i}{\partial L} = 0 \\
 \frac{d\bar{\theta}_i}{dL} &= - \sum_j \mathcal{K}_{ij}^{-1} \frac{\partial Q_j}{\partial L} \tag{3.10}
 \end{aligned}$$

where

$$\begin{aligned} \mathcal{K}_{jl} &= \frac{\partial Q_j}{\partial \bar{\theta}_l} = i\eta_j \frac{\partial \varepsilon_{\alpha_j}(\bar{\theta}_j | \bar{\theta}_1, \dots, \bar{\theta}_N)}{\partial \bar{\theta}_l} \\ &= i\eta_j \begin{cases} Lf_{s,\alpha_j}(\bar{\theta}_j) + \sum_{m \neq j} (-i\eta_m) f_{m,\alpha_j}(\bar{\theta}_j) & j = l \\ (i\eta_l) f_{l,\alpha_j}(\bar{\theta}_j) & j \neq l. \end{cases} \end{aligned} \quad (3.11)$$

Introducing the following combinations

$$\begin{aligned} \mathcal{N}_j &= i\eta_j f_{s,\alpha_j}(\bar{\theta}_j) \\ \mathcal{N}_{\varphi,jl} &= \eta_j \eta_l f_{l,\alpha_j}(\bar{\theta}_j) \end{aligned} \quad (3.12)$$

where $\mathcal{N}_{\varphi,jl} = \mathcal{N}_{\varphi,lj}$, \mathcal{K} can be rewritten as

$$\mathcal{K}_{jl} = \begin{cases} L\mathcal{N}_j + \sum_{m \neq j} \mathcal{N}_{\varphi,jm} & j = l \\ -\mathcal{N}_{\varphi,jl} & j \neq l. \end{cases} \quad (3.13)$$

The explicit volume derivative of the quantization condition is

$$\frac{\partial Q_j}{\partial L} = i\eta_j \frac{\partial \varepsilon_{\alpha_j}(\bar{\theta}_j | \bar{\theta}_1, \dots, \bar{\theta}_N)}{\partial L} = i\eta_j f_{c,\alpha_j}(\bar{\theta}_j). \quad (3.14)$$

Introducing

$$\mathcal{M}_j = i\eta_j f_{c,\alpha_j}(\bar{\theta}_j) \quad (3.15)$$

the derivative of the singularity position takes the following form

$$\frac{d\bar{\theta}_j}{dL} = - \sum_l \mathcal{K}_{jl}^{-1} \mathcal{M}_l \quad (3.16)$$

such as $\langle \Theta \rangle$

$$\begin{aligned} \frac{\langle \Theta \rangle}{2\pi} &= \frac{\langle \Theta \rangle_\infty}{2\pi} + \sum_\beta \int \frac{d\theta}{2\pi} \frac{m_\beta \cosh(\theta) f_{c,\beta}(\theta) - m_\beta \sinh(\theta) f_{s,\beta}(\theta)}{1 + e^{\varepsilon_\beta(\theta)}} \\ &+ \sum_{i=1}^N \frac{\mathcal{N}_i}{L} - \sum_{i,j=1}^N \mathcal{M}_i \mathcal{K}_{ij}^{-1} \mathcal{M}_j. \end{aligned} \quad (3.17)$$

This is our final form for the TBA result for $\langle \Theta \rangle$.

3.2 Isolating the singularity density terms

To see the equivalence of $\langle \Theta \rangle$ to the form factor series (2.15) the terms containing \mathcal{N}_i and \mathcal{M}_i need to be rearranged in order to match the structure of the singularity density terms in (2.15).

Let's start with the term

$$\sum_{i,j=1}^N \mathcal{M}_i \mathcal{K}_{ij}^{-1} \mathcal{M}_j. \quad (3.18)$$

The inverse of \mathcal{K} can be expressed by its co-factor matrix \mathcal{C}

$$\mathcal{K}_{ij}^{-1} = \frac{\mathcal{C}_{ji}}{\det \mathcal{K}}. \quad (3.19)$$

The diagonal elements of the co-factor matrix are just the principal minors of \mathcal{K} :

$$\mathcal{C}_{ii} = \det \mathcal{K}(\{i\}) \quad (3.20)$$

where $\mathcal{K}(I)$ denotes the matrix obtained by omitting from \mathcal{K} the rows and columns that are indexed by the set I . The non-diagonal elements of the co-factor matrix can be expressed with principal minors and sequences of the elements of \mathcal{K} [48]

$$\begin{aligned} \mathcal{C}_{ji} &= \sum_{n=0}^{N-2} \sum_{\{\alpha\}} (-1)^{n+1} \mathcal{K}_{i\alpha_1} \mathcal{K}_{\alpha_1\alpha_2} \dots \mathcal{K}_{\alpha_n j} \det \mathcal{K}(\{j, i, \alpha_1, \dots, \alpha_n\}) \\ &= \sum_{n=0}^{N-2} \sum_{\{\alpha\}} \mathcal{N}_{\varphi, i\alpha_1} \mathcal{N}_{\varphi, \alpha_1\alpha_2} \dots \mathcal{N}_{\varphi, \alpha_n j} \det \mathcal{K}(\{j, i, \alpha_1, \dots, \alpha_n\}) \end{aligned} \quad (3.21)$$

where $\{\alpha\} \subset \{1, \dots, N\} \setminus \{i, j\}$. With the help of (3.20) and (3.21) one can write

$$\begin{aligned} \sum_{i,j=1}^N \mathcal{M}_i \mathcal{K}_{ij}^{-1} \mathcal{M}_j &= \sum_i \frac{\det \mathcal{K}(i)}{\det \mathcal{K}} \mathcal{M}_i \mathcal{M}_i + \sum_{i \neq j} \sum_{n=0}^{N-2} \sum_{\{\alpha\}} \frac{\det \mathcal{K}(j, i, \{\alpha\})}{\det \mathcal{K}} \\ &\quad \times \mathcal{M}_i \mathcal{M}_j \mathcal{N}_{\varphi, i\alpha_1} \mathcal{N}_{\varphi, \alpha_1\alpha_2} \dots \mathcal{N}_{\varphi, \alpha_n j}. \end{aligned} \quad (3.22)$$

Now we turn to rearranging the term

$$\sum_{i=1}^N \frac{\mathcal{N}_i}{L} \quad (3.23)$$

in a similar manner. For this we need the following theorem:

Theorem 5. *If the $N \times N$ matrix $\mathcal{K}^{(N)}$ has the form*

$$\mathcal{K}_{ij}^{(N)} = \begin{cases} L\mathcal{N}_i + \sum_{k \neq i} \mathcal{N}_{\varphi, ik} & i = j \\ -\mathcal{N}_{\varphi, ij} & i \neq j \end{cases} \quad (3.24)$$

its determinant can be expanded as

$$\begin{aligned} \det \mathcal{K}^{(N)} &= L\mathcal{N}_i \det \mathcal{K}^{(N)}(\{i\}) \\ &\quad + \sum_{n=1}^{N-1} \sum_{\{\alpha\}} \mathcal{N}_{\varphi, i\alpha_1} \mathcal{N}_{\varphi, \alpha_1\alpha_2} \dots \mathcal{N}_{\varphi, \alpha_{n-1}\alpha_n} L\mathcal{N}_{\alpha_n} \det \mathcal{K}^{(N)}(\{i, \alpha_1, \dots, \alpha_n\}) \end{aligned} \quad (3.25)$$

where i is any chosen row, $\{\alpha\} \subset \{1, \dots, N\} \setminus \{i\}$ and $\mathcal{K}^{(N)}(I)$ is the submatrix of $\mathcal{K}^{(N)}$ as defined before.

Proof. Up to $N = 3$ it is easy to check the statement by direct evaluation. For $N > 3$ we proceed by induction. Let us suppose the theorem is valid for $N - 1$

$$\det \mathcal{K}^{(N-1)} = LN_i \det \mathcal{K}^{(N-1)}(\{i\}) + \sum_{n=1}^{N-2} \sum_{\{\alpha\}} \left\{ \mathcal{N}_{\varphi, i\alpha_1} \mathcal{N}_{\varphi, \alpha_1\alpha_2} \dots \mathcal{N}_{\varphi, \alpha_{n-1}\alpha_n} LN_{\alpha_n} \right. \\ \left. \times \det \mathcal{K}^{(N-1)}(\{i, \alpha_1, \dots, \alpha_n\}) \right\}. \quad (3.26)$$

The determinant for the matrix $\mathcal{K}^{(N)}$ of size N can be expanded by its row j

$$\det \mathcal{K}^{(N)} = \mathcal{K}_{jj}^{(N)} \mathcal{C}_{jj}^{(N)} + \sum_{i \neq j} \mathcal{K}_{ji}^{(N)} \mathcal{C}_{ji}^{(N)} \quad (3.27)$$

where $\mathcal{C}^{(N)}$ is the co-factor matrix of $\mathcal{K}^{(N)}$. Using (3.20) and (3.21) leads to

$$\det \mathcal{K}^{(N)} = \mathcal{K}_{jj}^{(N)} \det \mathcal{K}^{(N)}(\{j\}) + \sum_{i \neq j} \mathcal{K}_{ji}^{(N)} \sum_{n=0}^{N-2} \sum_{\{\alpha\}} \left\{ (-1)^{n+1} \mathcal{K}_{i\alpha_1} \mathcal{K}_{\alpha_1\alpha_2} \dots \mathcal{K}_{\alpha_n j} \right. \\ \left. \times \det \mathcal{K}^{(N)}(\{j, i, \alpha_1, \dots, \alpha_n\}) \right\} \\ = \left(LN_j + \sum_{i \neq j} \mathcal{N}_{\varphi, ji} \right) \det \mathcal{K}^{(N)}(\{j\}) - \sum_{i \neq j} \left\{ \mathcal{N}_{\varphi, ji} \right. \\ \left. \times \sum_{n=0}^{N-2} \sum_{\{\alpha\}} \mathcal{N}_{\varphi, i\alpha_1} \mathcal{N}_{\varphi, \alpha_1\alpha_2} \dots \mathcal{N}_{\varphi, \alpha_n j} \det \mathcal{K}^{(N)}(\{j, i, \alpha_1, \dots, \alpha_n\}) \right\}. \quad (3.28)$$

Now $\mathcal{K}^{(N)}(\{j\})$ can be related to $\mathcal{K}^{(N-1)}$ by observing that their off-diagonal elements are the same, while $\mathcal{K}_{ii}^{(N)} = \mathcal{K}_{ii}^{(N-1)} + \mathcal{N}_{\varphi, ij}$. Implementing this by shifting $LN_i \rightarrow LN_i + \mathcal{N}_{\varphi, ij}$ in (3.26) one obtains

$$\det \mathcal{K}^N(\{j\}) = \det \mathcal{K}^{(N-1)} \Big|_{LN_i \rightarrow LN_i + \mathcal{N}_{\varphi, ij}} \\ = (LN_i + \mathcal{N}_{\varphi, ij}) \det \mathcal{K}^{(N)}(\{j, i\}) + \sum_{n=1}^{N-2} \sum_{\{\alpha\}} \left\{ \mathcal{N}_{\varphi, i\alpha_1} \mathcal{N}_{\varphi, \alpha_1\alpha_2} \dots \mathcal{N}_{\varphi, \alpha_{n-1}\alpha_n} \right. \\ \left. \times (LN_{\alpha_n} + \mathcal{N}_{\varphi, \alpha_n j}) \det \mathcal{K}^{(N)}(\{j, i, \alpha_1, \dots, \alpha_n\}) \right\} \quad (3.29)$$

and inserting this back to (3.28) gives

$$\begin{aligned}
 \det \mathcal{K}^{(N)} &= L\mathcal{N}_j \det \mathcal{K}^{(N)}(\{j\}) + \sum_{i \neq j} \mathcal{N}_{\varphi,ji} (L\mathcal{N}_i + \mathcal{N}_{\varphi,ij}) \det \mathcal{K}^{(N)}(\{j,i\}) \\
 &+ \sum_{i \neq j} \mathcal{N}_{\varphi,ji} \sum_{n=1}^{N-2} \sum_{\{\alpha\}} \left\{ \mathcal{N}_{\varphi,i\alpha_1} \mathcal{N}_{\varphi,\alpha_1\alpha_2} \cdots \mathcal{N}_{\varphi,\alpha_{n-1}\alpha_n} \right. \\
 &\times (L\mathcal{N}_{\alpha_n} + \mathcal{N}_{\varphi,\alpha_n j}) \det \mathcal{K}^{(N)}(\{j,i,\alpha_1,\dots,\alpha_n\}) \left. \right\} \\
 &- \sum_{i \neq j} \mathcal{N}_{\varphi,ji} \sum_{n=0}^{N-2} \sum_{\{\alpha\}} \mathcal{N}_{\varphi,i\alpha_1} \mathcal{N}_{\varphi,\alpha_1\alpha_2} \cdots \mathcal{N}_{\varphi,\alpha_n j} \det \mathcal{K}^{(N)}(\{i,j,\alpha_1,\dots,\alpha_n\}) \\
 &= L\mathcal{N}_j \det \mathcal{K}^{(N)}(\{j\}) + \\
 &+ \sum_{n=1}^{N-1} \sum_{\{\alpha\}} \mathcal{N}_{\varphi,j\alpha_1} \mathcal{N}_{\varphi,\alpha_1\alpha_2} \cdots \mathcal{N}_{\varphi,\alpha_{n-1}\alpha_n} L\mathcal{N}_{\alpha_n} \det \mathcal{K}^{(N)}(\{j,\alpha_1,\dots,\alpha_n\}) \quad (3.30)
 \end{aligned}$$

which is just the statement we wanted to prove. *Q.e.d.* □

Using the above theorem we can rewrite

$$\begin{aligned}
 \sum_i \frac{\mathcal{N}_i}{L} &= \sum_i \frac{\mathcal{N}_i \det \mathcal{K}}{L \det \mathcal{K}} = \sum_i \mathcal{N}_i \mathcal{N}_i \frac{\det \mathcal{K}(\{i\})}{\det \mathcal{K}} + \sum_{i \neq j} \sum_{n=0}^{N-2} \sum_{\{\alpha\}} \frac{\det \mathcal{K}^N(\{i,j,\alpha_1,\dots,\alpha_n\})}{\det \mathcal{K}} \\
 &\times \mathcal{N}_i \mathcal{N}_j \mathcal{N}_{\varphi,i\alpha_1} \mathcal{N}_{\varphi,\alpha_1\alpha_2} \cdots \mathcal{N}_{\varphi,\alpha_n j} \quad (3.31)
 \end{aligned}$$

which has the same structure as (3.22). Substituting the definitions of \mathcal{N} (3.12) and \mathcal{M} (3.15) into (3.22) and (3.31) we can see that every η_i factor appears twice and so drops out. Therefore the expression for $\langle \Theta \rangle$ (3.17) simplifies to

$$\begin{aligned}
 \frac{\langle \Theta \rangle}{2\pi} &= \frac{\langle \Theta \rangle_\infty}{2\pi} + \sum_\beta \int \frac{d\theta}{2\pi} \frac{m_\beta \cosh(\theta) f_{c,\beta}(\theta) - m_\beta \sinh(\theta) f_{s,\beta}(\theta)}{1 + e^{\varepsilon_\beta(\theta)}} \\
 &+ \sum_i \frac{\det \mathcal{K}(\{i\})}{\det \mathcal{K}} [f_{c,\alpha_i}(\bar{\theta}_i) f_{c,\alpha_i}(\bar{\theta}_i) - f_{s,\alpha_i}(\bar{\theta}_i) f_{s,\alpha_i}(\bar{\theta}_i)] \\
 &+ \sum_{i \neq j} \sum_{n=0}^{N-2} \sum_{\{\alpha\}} \frac{\det \mathcal{K}(\{i,j,\alpha_1,\dots,\alpha_n\})}{\det \mathcal{K}} [f_{c,\alpha_i}(\bar{\theta}_i) f_{c,\alpha_j}(\bar{\theta}_j) - f_{s,\alpha_i}(\bar{\theta}_i) f_{s,\alpha_j}(\bar{\theta}_j)] \\
 &\times f_{\alpha_1,\alpha_i}(\bar{\theta}_i) f_{\alpha_2,\alpha_1}(\bar{\theta}_1) \cdots f_{j,\alpha_n}(\bar{\theta}_n). \quad (3.32)
 \end{aligned}$$

The determinant ratios are exactly the density factors in (2.15); what remains to be shown is that the other terms reproduce the dressed form factors of Θ .

3.3 Dressed form factors of Θ

Theorem 6. *In the absence of active singularities of the TBA equations, the dressed form factors of Θ are given by*

$$\mathcal{D}_\varepsilon^\Theta = \sum_{n_1, \dots, n_k=0}^{\infty} \frac{1}{\prod_i n_i!} \int_{-\infty}^{\infty} \prod_{j=1}^{\tilde{N}} \frac{d\theta_j}{2\pi [1 + e^{\varepsilon\beta_j(\theta_j)}]} F_{2n_1, \dots, 2n_k, c}^\Theta(\theta_1, \dots, \theta_{\tilde{N}}) \quad (3.33)$$

$$\tilde{N} = \sum_{i=1}^k n_i$$

which is equal to

$$\mathcal{D}_\varepsilon^\Theta = \langle \Theta \rangle_\infty + 2\pi \sum_{\beta} \int \frac{d\theta}{2\pi} \frac{m_\beta \cosh(\theta) f_{c,\beta}(\theta) - m_\beta \sinh(\theta) f_{s,\beta}(\theta)}{1 + e^{\varepsilon\beta(\theta)}}. \quad (3.34)$$

Proof. The connected diagonal form factors of Θ are given by [1]

$$F_{2n,c}^\Theta(\theta_1, \dots, \theta_n) = 2\pi \varphi_{12} \varphi_{23} \dots \varphi_{n-1,n} m_{\beta_1} m_{\beta_n} \cosh(\theta_{1n}) + \text{permutations}$$

where $\theta_{ij} = \theta_i - \theta_j$, β_i denotes the species of the i th particle and φ_{ij} is a short-hand for $\varphi_{\beta_i \beta_j}(\theta_{ij})$. (3.33) is symmetric under re-ordering particles of the same species which results in a combinatorial factor $\prod_i n_i!$ canceling the denominators in front of the integrals. To take into account the rest of the permutations we can rewrite $\mathcal{D}_\varepsilon^\Theta$ like

$$\begin{aligned} \mathcal{D}_\varepsilon^\Theta &= 2\pi \sum_{n=0}^{\infty} \sum_{\beta_1, \dots, \beta_n} \int_{-\infty}^{\infty} \prod_{j=1}^n \frac{d\theta_j}{2\pi [1 + e^{\varepsilon\beta_j(\theta_j)}]} \varphi_{12} \varphi_{23} \dots \varphi_{n-1,n} m_{\beta_1} m_{\beta_n} \cosh(\theta_{1n}) \\ &= \sum_{n=0}^{\infty} \mathcal{D}_{\varepsilon,n}^\Theta. \end{aligned} \quad (3.35)$$

Following [1] every $\mathcal{D}_{\varepsilon,n}^\Theta$ can be graphically represented as seen in figure 1a where every node represents a particle with a given rapidity and species, including the integration

$$\int \frac{d\theta_i}{2\pi [1 + e^{\varepsilon\beta_i(\theta_i)}]}$$

and the first and last node is multiplied by its mass. Every horizontal line represents a factor φ_{ij} and the dashed line represents the factor $\cosh \theta_{1n}$. The whole graph is multiplied by 2π to account for the normalization of the operator Θ , and summed over every possible type configuration for the nodes. The empty graph (with zero node) represents $\mathcal{D}_{\varepsilon,0}^\Theta = \langle \Theta \rangle_\infty$. Using hyperbolic addition formulas for the $\cosh \theta_{1n}$ terms every graph can be represented as difference of two chains where the two end nodes instead of being connected by dashed line, are multiplied by \cosh or \sinh of the rapidity at the given node as shown in figure 1b.




Figure 1. Graphical representation of $\mathcal{D}_{\epsilon,n}^\Theta$, $K_{n,\beta}(\theta)$ and $J_{n,\beta}(\theta)$.

Since the functions f_c and f_s in (3.34) satisfy the self-consistent equations (3.4), it is convenient to expand them in the following way

$$\begin{aligned}
 f_{c,\beta}(\theta) &= \sum_{n=0}^{\infty} K_{n,\beta}(\theta) \\
 f_{s,\beta}(\theta) &= \sum_{n=0}^{\infty} J_{n,\beta}(\theta)
 \end{aligned}
 \tag{3.36}$$

where

$$\begin{aligned}
 K_{0,\beta}(\theta) &= m_\beta \cosh(\theta) \\
 J_{0,\beta}(\theta) &= m_\beta \sinh(\theta) \\
 K_{n,\beta}(\theta) &= \sum_{\beta_1, \dots, \beta_n} \int \prod_i \frac{d\theta_i}{2\pi [1 + e^{\epsilon \beta_i(\theta_i)}]} \varphi_{\beta \beta_1} \dots \varphi_{\beta_{n-1} \beta_n} m_{\beta_n} \cosh(\theta_n) \\
 J_{n,\beta}(\theta) &= \sum_{\beta_1, \dots, \beta_n} \int \prod_i \frac{d\theta_i}{2\pi [1 + e^{\epsilon \beta_i(\theta_i)}]} \varphi_{\beta \beta_1} \dots \varphi_{\beta_{n-1} \beta_n} m_{\beta_n} \sinh(\theta_n).
 \end{aligned}
 \tag{3.37}$$

The graphical representation of $K_{n,\beta}(\theta)$ and $J_{n,\beta}(\theta)$ can be seen in figure 1c; the dashed node indicates that the corresponding rapidity integral and the filling fraction belonging to that node is not included in the contribution. Comparing to figure 1b it is clear that $K_{n,\beta}(\theta)$ and $J_{n,\beta}(\theta)$ describe the contribution of the chain between one of the end nodes and a dashed node with type β which is n steps away from the end node. Multiplying $K_{n,\beta}(\theta)$ and $J_{n,\beta}(\theta)$ with $2\pi \sum_\beta \int \frac{d\theta}{2\pi} \frac{m_\beta \cosh(\theta)}{1 + e^{\epsilon \beta(\theta)}}$ and $-2\pi \sum_\beta \int \frac{d\theta}{2\pi} \frac{m_\beta \sinh(\theta) f_{s,\beta}(\theta)}{1 + e^{\epsilon \beta(\theta)}}$ as in (3.34) closes the chains and they become identical to the ones in figure 1b with length $n + 1$, i.e. equal to $\mathcal{D}_{\epsilon,n+1}^\Theta$. The sum for n in f_c and f_s then generates all the contributions in $\mathcal{D}_\epsilon^\Theta$. *Q.e.d.* \square




Figure 2. Graphical representation of $\mathcal{D}_{\epsilon, n m}^{\Theta}(\bar{\theta}_i)$, $K_{n, \alpha_i}(\bar{\theta}_i)$ and $J_{n, \alpha_i}(\bar{\theta}_i)$.

Theorem 7. The dressed form factor of Θ with one active singularity $\bar{\theta}_i$ with type α_i is

$$\mathcal{D}_{\epsilon}^{\Theta}(\bar{\theta}_i) = \sum_{n_1, \dots, n_k=0}^{\infty} \frac{1}{\prod_i n_i!} \int_{-\infty}^{\infty} \prod_{j=1}^{\tilde{N}} \frac{d\theta_j}{2\pi [1 + e^{\epsilon \beta_j(\theta_j)}]} F_{2n_1, \dots, 2n_k, c}^{\Theta}(\bar{\theta}_i, \theta_1, \dots, \theta_{\tilde{N}}) \quad (3.38)$$

$$\tilde{N} = \sum_{i=1}^k n_i$$

which is equal to

$$\mathcal{D}_{\epsilon}^{\Theta}(\bar{\theta}_i) = f_{c, \alpha_i}(\bar{\theta}_i) f_{c, \alpha_i}(\bar{\theta}_i) - f_{s, \alpha_i}(\bar{\theta}_i) f_{s, \alpha_i}(\bar{\theta}_i). \quad (3.39)$$

Proof. The proof follows the ideas used in demonstrating theorem 6. The factor $\prod_i n_i!$ for the non-active singularities cancels as before, but now one must sum over all possible positions of the active singularity:

$$\begin{aligned} \mathcal{D}_{\epsilon}^{\Theta}(\bar{\theta}_i) &= 2\pi \sum_{n, m=0}^{\infty} \sum_{\beta_1, \dots, \beta_{n+m}} \int_{-\infty}^{\infty} \prod_{j=1}^{n+m} \frac{d\theta_j}{2\pi [1 + e^{\epsilon \beta_j(\theta_j)}]} \varphi_{12} \varphi_{23} \dots \varphi_{ni} \\ &\times \varphi_{i, n+1} \dots \varphi_{n+m-1, n+m} m_{\beta_1} m_{\beta_{n+m}} \cosh(\theta_{1, n+m}) = \sum_{n, m=0}^{\infty} \mathcal{D}_{\epsilon, n m}^{\Theta}(\bar{\theta}_i) \end{aligned} \quad (3.40)$$

where the sum runs for the number and the species nodes between the active singularity and the end nodes (for $n = 0$, the active singularity is the end node on the left, while for $m = 0$ it is the end node on the right). The active singularity is marked by a black node in figures 2a and 2b. $K_{n, \alpha_i}(\bar{\theta}_i)$ and $J_{n, \alpha_i}(\bar{\theta}_i)$ are represented in figure 2c; they are equal to the contribution of the chain between one of the end nodes and the active singularity. Multiplying them as in 3.39 it follows that

$$\mathcal{D}_{\epsilon, n m}^{\Theta}(\bar{\theta}_i) = K_{n, \alpha_i}(\bar{\theta}_i) K_{m, \alpha_i}(\bar{\theta}_i) - J_{n, \alpha_i}(\bar{\theta}_i) J_{m, \alpha_i}(\bar{\theta}_i) \quad (3.41)$$

and the summation in both of $f_{c/s}$ the result exactly reproduces $\mathcal{D}_{\epsilon}^{\Theta}(\bar{\theta}_i)$. *Q.e.d.* \square




Figure 3. Graphical representation of the terms of $\mathcal{D}_\varepsilon^\mathcal{O}(\bar{\theta}_1, \dots, \bar{\theta}_N)$ and $L_{n,ji}$.

Theorem 8. The dressed form factor of Θ with N active singularities $\{\bar{\theta}_1, \dots, \bar{\theta}_N\}$ is

$$\begin{aligned} \mathcal{D}_\varepsilon^\mathcal{O}(\bar{\theta}_1, \dots, \bar{\theta}_N) &= \sum_{n_1, \dots, n_k=0}^{\infty} \frac{1}{\prod_i n_i!} \int_{-\infty}^{\infty} \prod_{j=1}^{\tilde{N}} \frac{d\theta_j}{2\pi [1 + e^{\varepsilon\beta_j(\theta_j)}]} \\ &\times F_{2n_1, \dots, 2n_k, c}^\mathcal{O}(\bar{\theta}_1, \dots, \bar{\theta}_N, \theta_1, \dots, \theta_{\tilde{N}}) \\ \tilde{N} &= \sum_{i=1}^k n_i \end{aligned} \quad (3.42)$$

which is equal to

$$\begin{aligned} \mathcal{D}_\varepsilon^\mathcal{O}(\bar{\theta}_1, \dots, \bar{\theta}_N) &= \sum_{i \neq j} \sum_{\{\alpha\}} [f_{c, \alpha_i}(\bar{\theta}_i) f_{c, \alpha_j}(\bar{\theta}_j) - f_{s, \alpha_i}(\bar{\theta}_i) f_{s, \alpha_j}(\bar{\theta}_j)] \\ &\times f_{\alpha_1, \alpha_i}(\bar{\theta}_i) f_{\alpha_2, \alpha_1}(\bar{\theta}_1) \dots f_{j, \alpha_n}(\bar{\theta}_n) \end{aligned} \quad (3.43)$$

where $\{\alpha\} = \{1, \dots, N\} \setminus \{i, j\}$.

Proof. As in the proofs of the previous theorems, (3.42) can be organized into a sum over terms corresponding to individual permutations of the active singularities. For a given permutation, the contribution is the sum of graphs represented in figures 3a and 3b, where the number and type of nodes separating end nodes and the active singularities is varying.

The functions $f_{\alpha_q, \alpha_p}(\bar{\theta}_p)$ can be expanded as $f_{c/s}$ previously, using their definition in (3.4)

$$f_{\alpha_q, \alpha_p}(\bar{\theta}_p) = \sum_{n=0}^{\infty} L_{n, pq} \quad (3.44)$$

where

$$\begin{aligned} L_{0, pq} &= \varphi_{pq} \\ L_{n, pq} &= \sum_{\beta_1, \dots, \beta_n} \int \prod_i \frac{d\theta_i}{2\pi [1 + e^{\varepsilon\beta_i(\theta_i)}]} \varphi_{\alpha_p \beta_1} \dots \varphi_{\beta_n \alpha_q}. \end{aligned} \quad (3.45)$$

$L_{n,pq}$ is represented in figure 3c; it generates all the possible contribution to figure 3b between two active singularities. In a given permutation of the active singularities let us take $\bar{\theta}_i$ and $\bar{\theta}_j$ as the two active singularities closest to the left/right end nodes; then the terms

$$f_{c,\alpha_i}(\bar{\theta}_i) f_{c,\alpha_j}(\bar{\theta}_j) - f_{s,\alpha_i}(\bar{\theta}_i) f_{s,\alpha_j}(\bar{\theta}_j)$$

generate all the contributions between the active singularities and the ends, and

$$f_{\alpha_1,\alpha_i}(\bar{\theta}_i) f_{\alpha_2,\alpha_1}(\bar{\theta}_1) \dots f_{j,\alpha_n}(\bar{\theta}_n)$$

generate all the contributions between the other active singularities. Summing up for all the permutations of the active singularities proves the theorem. *Q.e.d.* \square

Theorems 6, 7 and 8 prove the equivalence of the form factor series (2.15) and the TBA equations for $\langle \Theta \rangle$ in any excited state described by the TBA system (2.6).

4 Finite volume expectation values in the T_2 model

For the numerical validation of the conjecture (2.15) we follow a similar strategy as we did for the Leclair-Mussardo conjecture in [45], where we chose the T_2 model for the numerics. The T_2 model is the perturbation of the $\mathcal{M}_{2,7}$ conformal minimal model [49] by the primary operator $\Phi_{1,3}$.

There are several advantages in this choice. First, all the data necessary to calculate the series (2.15): the scattering theory, the form factors (of primary fields) and the excited state TBA equations, are known for this model. Second, the T_2 model contains operators with known form factors that are not related to the trace of the stress-energy tensor; the trace of the stress energy tensor is not interesting since the conjecture is equivalent to the TBA equation as shown in section 3. Finally for the T_2 model the Truncated Conformal Space Approach (TCSA) improved by renormalization group methods [45] gives an efficient way to evaluate the expectation values directly solving the dynamics of the model. The TCSA was introduced in [44] while renormalization techniques were proposed in [50, 51] and further developed in [52–54]; the development of related methods is now a very active field of investigation [55–58]. For details on TCSA in the T_2 model and the renormalization method we refer the interested reader to [45].

4.1 Excited state TBA equations for a single type-1 state

4.1.1 General form and solution of the excited TBA equations

The simplest excited states in the excited TBA formalism for the T_2 model [43] are those with a single type-1 particle. For these excited states the TBA equations contain only two

active singularities of type-2 with $\eta_1 = -1$, $\eta_2 = 1$:

$$\varepsilon_a(\theta) = m_a L \cosh(\theta) + \log \left(\frac{S_{a2}(\theta - \bar{\theta}_1)}{S_{a2}(\theta - \bar{\theta}_2)} \right) \quad (4.1)$$

$$- \sum_b \int \frac{d\theta'}{2\pi} \varphi_{ab}(\theta - \theta') \log \left(1 + e^{-\varepsilon_b(\theta')} \right)$$

$$e^{-\varepsilon_2(\bar{\theta}_{1/2})} = -1 \quad (4.2)$$

$$E_{\text{TBA}}(L) = -im_2 (\sinh(\bar{\theta}_1) - \sinh(\bar{\theta}_2)) - \sum_a \int \frac{d\theta}{2\pi} m_a \cosh(\theta) \log \left(1 + e^{-\varepsilon_a(\theta)} \right) \quad (4.3)$$

which are related by $\bar{\theta}_2 = \bar{\theta}_1^*$ for states with nonzero momentum, where $*$ is the complex conjugation; or $\bar{\theta}_2 = -\bar{\theta}_1$ for zero momentum.

In large volume (IR limit) the integral term is negligible in (4.1), and the cosh term goes to infinity with the volume, while the value of $\varepsilon_2(\bar{\theta}_1)$ is finite, which forces the imaginary part of the active singularity's position to $\frac{i\pi}{10}$, since S_{22} is singular around $\bar{\theta}_1 - \bar{\theta}_2 \sim \frac{i\pi}{5}$. The position of the active singularity can be written as

$$\bar{\theta}_1 = \tilde{\theta} + i \left(\frac{\pi}{10} + \delta \right) \quad (4.4)$$

where $\tilde{\theta}$ and δ are real; δ is a correction to the imaginary part that decays exponentially in the dimensionless variable $m_1 L$. Substituting this form into condition (4.2) and keeping only the first order corrections in δ , the solution for the position of the active singularity is

$$m_1 L \sinh \tilde{\theta} = 2\pi s$$

$$\delta = \cos(\pi s) \tan \left(\frac{3\pi}{10} \right) \tan^2 \left(\frac{2\pi}{5} \right) e^{-m_2 \cos(\frac{\pi}{10}) \sqrt{m_1^2 L^2 + (2\pi s)^2}} \quad (4.5)$$

where s is an integer number giving the momentum quantum number of the state. Using this solution for TBA energy (4.3) and expanding the $\log(1 + e^{-\varepsilon})$ term in the integral, the energy takes the following form

$$E_{\text{TBA}}(L) = E(L) - \mathcal{B}L$$

$$= 2 \sin \left(\frac{\pi}{10} \right) m_2 \cosh(\tilde{\theta}) + 2m_2 \cosh(\tilde{\theta}) \sin \left(\frac{\pi}{10} \right) \delta$$

$$- \sum_a \int \frac{d\theta}{2\pi} m_a \cosh(\theta) e^{-m_a L \cosh(\theta)} \frac{S_{a2}(\theta - \tilde{\theta} + i\frac{\pi}{10})}{S_{a2}(\theta - \tilde{\theta} - i\frac{\pi}{10})}$$

$$= \sqrt{m_1^2 + \left(\frac{2\pi s}{L} \right)^2} + 2\sqrt{m_2^2 + \left(\frac{2\pi s m_2}{L m_1} \right)^2} \cos \left(\frac{\pi}{10} \right)$$

$$\times \cos(\pi s) \tan \left(\frac{3\pi}{10} \right) \tan^2 \left(\frac{2\pi}{5} \right) e^{-m_2 \cos(\frac{\pi}{10}) \sqrt{m_1^2 L^2 + (2\pi s)^2}}$$

$$- \sum_a \int \frac{d\theta}{2\pi} m_a \cosh(\theta) e^{-m_a L \cosh(\theta)} S_{a1} \left(\theta - \tilde{\theta} + i\frac{\pi}{2} \right) \quad (4.6)$$




Figure 4. Effective central charge for the excited states with momentum quantum number $s = 0, 1$ from TBA and TCSA, with the UV (CFT value) and IR asymptotics (μ and F term).

where the bootstrap identity

$$\frac{S_{a2}(\theta + i\frac{\pi}{10})}{S_{a2}(\theta - i\frac{\pi}{10})} = S_{a1}(\theta + i\frac{\pi}{2}) \tag{4.7}$$

was used. The first term gives L^{-1} corrections related to the kinetic energy of the particle in finite volume, while the second and third terms are the leading exponential corrections, the so-called μ and F terms [59], which for a zero-momentum particle were first derived by Lüscher in [60].

In the small volume (UV) limit the energy is proportional to the effective central charge of the state

$$E_{\text{TBA}}(L) = -\frac{6\pi}{L} c_{\text{eff}}(L) \tag{4.8}$$

which has the ultraviolet limit

$$c_{\text{eff}}(0) = c - 12(\Delta + \bar{\Delta}) \tag{4.9}$$

where $c = -68/7$ is the central charge of the minimal model $\mathcal{M}_{2,7}$ and $\Delta, \bar{\Delta}$ are the left/right conformal weights of the state in the ultraviolet limit. Using the dilogarithm trick introduced in [3] one can confirm the expected effective central charge for these states [43]. In figure 4 we plot c_{eff} for the states $s = 0, 1$ showing that the TBA results match perfectly with the TCSA calculation and also reproduces the expected asymptotics.

The excited state TBA equations are solved numerically by simultaneously iterating eqs. (4.1) and (4.2) in large volume, where the asymptotic of the pole position (4.5) can be used as a starting point. Using this solution the volume is decreased and the equations iterated, and continuing this process the solution can be tracked to small volume. For $s \neq 0$ the numerics is straightforward to perform up to precision of order 10^{-12} , and all the ingredients to calculate the conjecture (2.15) can be readily constructed. For $s = 0$ there exists a critical volume r_c under which it is necessary to be more careful with the numerical calculation.

4.1.2 Zero-momentum state: desingularization in small volume

As described in details in [41–43], for states containing a zero-momentum particle it can happen that under a given critical volume r_c some singularity ends up being on the integration contour. To describe this situation, we recall that the Y -system (2.9) gives relations between positions where the functions $Y_\alpha = e^{\varepsilon_\alpha}$ take the values 0 and -1 , which are the logarithmic singular points of the TBA equations. For the T_2 model the incidence matrix is given by

$$I^{[T_2]} = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} \quad \text{and} \quad h = 5. \quad (4.10)$$

For the excited state containing a single type-1 particle with zero momentum in large volume there are active singularities on the imaginary axis at $\pm\bar{\theta}_1$ with

$$\bar{\theta}_1 = i\frac{\pi}{10} + i\delta \quad (4.11)$$

with $\delta > 0$ where $Y_2 = -1$. From (2.9) it follows that $Y_2 = -1$ at $\bar{\theta}_1 - i\frac{2\pi}{5}$ and $-\bar{\theta}_1 + i\frac{2\pi}{5}$, and $Y_1 = Y_2 = 0$ at positions $\bar{\theta}_1 - i\frac{\pi}{5}$ and $-\bar{\theta}_1 + i\frac{\pi}{5}$. As the volume decreases, the value of δ increases till at some critical value of the volume given by $m_1L = r_c$ it reaches $\delta_c = \frac{\pi}{10}$. At this point $\bar{\theta}_1 = -\bar{\theta}_1 + i\frac{2\pi}{5}$ resulting in a coincidence of singularities, and also of zeros. Decreasing the volume to $m_1L < r_c$ results in the “scattering” of the singularities on each other at right angles pushing them away from the imaginary axis with fixed imaginary part in the form

$$\bar{\theta}_1 = i\frac{\pi}{5} + \alpha. \quad (4.12)$$

As a result, the zeros of Y_1 and Y_2 sit exactly on the integration contour, making the equation for the pseudo-energy (4.1) singular and leading to instabilities in the numerical solution of the TBA equations.

One way to handle the numerical treatment of the problem is to shift the integration contour, while an alternative is to rearrange the self-consistent equations appropriately, which is called desingularization. The latter approach relies on the relation [43]

$$\begin{aligned} \varphi_{ad}(\theta) &= -\varphi_h(\theta) I_{ad}^{[T_2]} + \sum_b \int \frac{d\theta'}{2\pi} \varphi_{ab}(\theta - \theta') \varphi_h(\theta') I_{bd}^{[T_2]} \\ \varphi_h(\theta) &= \frac{h}{2 \cosh\left(\frac{h}{2}\theta\right)} = \pm i \partial_\theta \log\left(\sigma_h\left(\theta \pm i\frac{\pi}{h}\right)\right) \\ \sigma_h(\theta) &= \tanh\left(\frac{h}{4}\theta\right) \end{aligned} \quad (4.13)$$

which allows the TBA equations to be recast as

$$\begin{aligned}
 \hat{\varepsilon}_a(\theta) &= \varepsilon_a(\theta) - \log\left(\sigma(\theta' - \tilde{\theta}_1)\sigma(\theta' - \tilde{\theta}_2)\right) \\
 \hat{\varepsilon}_a(\theta) &= m_a L \cosh(\theta) - \sum_b \int \frac{d\theta'}{2\pi} \varphi_{ab}(\theta - \theta') \log\left(\sigma(\theta' - \tilde{\theta}_1)\sigma(\theta' - \tilde{\theta}_2) + e^{-\hat{\varepsilon}_b(\theta')}\right) \\
 e^{\hat{\varepsilon}_a(\tilde{\theta}_1)} &= i \coth\left(\frac{5}{2}\tilde{\theta}_1 + i\frac{\pi}{4}\right) \\
 E_{\text{TBA}}(L) &= - \sum_a \int \frac{d\theta}{2\pi} m_a \cosh(\theta) \log\left(\sigma(\theta - \tilde{\theta}_1)\sigma(\theta - \tilde{\theta}_2) + e^{-\hat{\varepsilon}_b(\theta)}\right) \tag{4.14}
 \end{aligned}$$

where $\tilde{\theta}_1 = \bar{\theta}_1 - i\frac{\pi}{h}$, $\tilde{\theta}_2 = \bar{\theta}_2 + i\frac{\pi}{h}$. These equations are regular and can be iterated in a stable way, however, the available precision using double precision numbers drops to the order of 10^{-10} . Fortunately, that is still more than sufficient for our purposes.

To calculate the densities and $\langle \Theta \rangle$ under r_c we need to desingularize f_s, f_c, f_i in (3.4) as well. It's easy to see, that the equation for f_s and f_c is regular under r_c , since $\frac{1}{1+e^{\varepsilon_\alpha}}$ at the singularity position is regular, because $Y_\alpha = e^{\varepsilon_\alpha} = 0$. For f_i the source term φ is singular at $\tilde{\theta}_{1,2}$, hence it is necessary to desingularize it:

$$\begin{aligned}
 \hat{f}_{i,\alpha}(\theta) &= f_{i,\alpha}(\theta) + \varphi_h(\theta - \tilde{\theta}_i) \\
 \hat{f}_{i,\alpha}(\theta) &= \sum_\beta \int \frac{d\theta'}{2\pi} \varphi_{\alpha\beta}(\theta - \theta') \frac{e^{\varepsilon_\beta(\theta')} \varphi_h(\theta - \tilde{\theta}_i) + \hat{f}_{i,\beta}(\theta')}{1 + e^{\varepsilon_\beta(\theta')}}. \tag{4.15}
 \end{aligned}$$

For numerical calculations the form

$$\begin{aligned}
 \hat{f}_{1,\alpha}(\theta) &= \sum_\beta \int \frac{d\theta'}{2\pi} \varphi_{\alpha\beta}(\theta - \theta') \frac{\partial\sigma(\theta' - \tilde{\theta}_1)\sigma(\theta' - \tilde{\theta}_2) + e^{-\hat{\varepsilon}_b(\theta')} \hat{f}_{1,\beta}(\theta')}{\sigma(\theta' - \tilde{\theta}_1)\sigma(\theta' - \tilde{\theta}_2) + e^{-\hat{\varepsilon}_b(\theta')}} \\
 \hat{f}_{2,\alpha}(\theta) &= \sum_\beta \int \frac{d\theta'}{2\pi} \varphi_{\alpha\beta}(\theta - \theta') \frac{\sigma(\theta' - \tilde{\theta}_1)\partial\sigma(\theta' - \tilde{\theta}_2) + e^{-\hat{\varepsilon}_b(\theta')} \hat{f}_{1,\beta}(\theta')}{\sigma(\theta' - \tilde{\theta}_1)\sigma(\theta' - \tilde{\theta}_2) + e^{-\hat{\varepsilon}_b(\theta')}} \tag{4.16}
 \end{aligned}$$

is more convenient.

4.2 Densities and the conjecture for states with a single type-1 particle

The derivatives of the quantization condition can be written like (3.11) with the help of the definitions in (3.4), (3.12) for the single type-1 state

$$\frac{\partial(Q_1, Q_2)}{\partial(\bar{\theta}_1, \bar{\theta}_2)} = \begin{pmatrix} L\mathcal{N}_1 + \mathcal{N}_\varphi & -\mathcal{N}_\varphi \\ -\mathcal{N}_\varphi & L\mathcal{N}_2 + \mathcal{N}_\varphi \end{pmatrix} \tag{4.17}$$

where

$$\begin{aligned}
 \mathcal{N}_1 &= -if_{s,2}(\bar{\theta}_1) = -im_2 \sinh(\bar{\theta}_1) - i \sum_{\beta} \int \frac{d\theta'}{2\pi} \varphi_{2\beta}(\bar{\theta}_1 - \theta') \frac{f_{s,\beta}(\theta')}{1 + e^{\varepsilon_{\beta}(\theta')}} \\
 \mathcal{N}_2 &= if_{s,2}(\bar{\theta}_2) = im_2 \sinh(\bar{\theta}_2) + i \sum_{\beta} \int \frac{d\theta'}{2\pi} \varphi_{2\beta}(\bar{\theta}_2 - \theta') \frac{f_{s,\beta}(\theta')}{1 + e^{\varepsilon_{\beta}(\theta')}} \\
 \mathcal{N}_{\varphi} &= -f_{2,2}(\bar{\theta}_1) = -\varphi_{22}(\bar{\theta}_1 - \bar{\theta}_2) - \sum_{\beta} \int \frac{d\theta'}{2\pi} \varphi_{2\beta}(\bar{\theta}_1 - \theta') \frac{f_{2,\beta}(\theta')}{1 + e^{\varepsilon_{\beta}(\theta')}} \\
 &= -f_{1,2}(\bar{\theta}_2) = -\varphi_{22}(\bar{\theta}_2 - \bar{\theta}_1) - \sum_{\beta} \int \frac{d\theta'}{2\pi} \varphi_{2\beta}(\bar{\theta}_2 - \theta') \frac{f_{1,\beta}(\theta')}{1 + e^{\varepsilon_{\beta}(\theta')}}. \tag{4.18}
 \end{aligned}$$

For the case $s = 0$ and $m_1 L < r_c$

$$\begin{aligned}
 \mathcal{N}_{\varphi} &= -\hat{f}_{2,2}(\bar{\theta}_1) + \varphi_h(\bar{\theta}_1 - \bar{\theta}_2) = +\varphi_h(\bar{\theta}_1 - \bar{\theta}_2) \\
 &\quad - \sum_{\beta} \int \frac{d\theta'}{2\pi} \varphi_{\alpha\beta}(\bar{\theta}_1 - \theta') \frac{\sigma(\theta' - \tilde{\theta}_1) \partial\sigma(\theta' - \tilde{\theta}_2) + e^{-\hat{\varepsilon}_b(\theta')} \hat{f}_{1,\beta}(\theta')}{\sigma(\theta' - \tilde{\theta}_1) \sigma(\theta' - \tilde{\theta}_2) + e^{-\hat{\varepsilon}_b(\theta')}} \\
 &= -\hat{f}_{1,2}(\bar{\theta}_2) + \varphi_h(\bar{\theta}_2 - \bar{\theta}_1) = +\varphi_h(\bar{\theta}_2 - \bar{\theta}_1) \\
 &\quad - \sum_{\beta} \int \frac{d\theta'}{2\pi} \varphi_{\alpha\beta}(\bar{\theta}_2 - \theta') \frac{\partial\sigma(\theta' - \tilde{\theta}_1) \sigma(\theta' - \tilde{\theta}_2) + e^{-\hat{\varepsilon}_b(\theta')} \hat{f}_{1,\beta}(\theta')}{\sigma(\theta' - \tilde{\theta}_1) \sigma(\theta' - \tilde{\theta}_2) + e^{-\hat{\varepsilon}_b(\theta')}}. \tag{4.19}
 \end{aligned}$$

With the above densities the conjecture for the form factor series (2.15) for this state takes the form

$$\begin{aligned}
 \langle \bar{\theta}_1, \bar{\theta}_2 | \mathcal{O} | \bar{\theta}_1, \bar{\theta}_2 \rangle_L &= \sum_{n_1, n_2=0}^{\infty} \frac{1}{n_1! n_2!} \int_{-\infty}^{\infty} \prod_{j=1}^{\tilde{N}} \frac{d\theta_j}{2\pi [1 + e^{\varepsilon_{\alpha_j}(\theta_j)}]} \left[\right. \\
 &\quad \left. F_{2n_1, 2n_2, c}^{\mathcal{O}}(\theta_1, \dots, \theta_{\tilde{N}}) + \frac{1}{L^2 \mathcal{N}_1 \mathcal{N}_2 + \mathcal{N}_{\varphi} L (\mathcal{N}_1 + \mathcal{N}_2)} \left\{ \right. \right. \\
 &\quad \left. \left. + (L\mathcal{N}_1 + \mathcal{N}_{\varphi}) F_{2n_1, 2n_2+2, c}^{\mathcal{O}}(\bar{\theta}_1, \theta_1, \dots, \theta_{\tilde{N}}) \right. \right. \\
 &\quad \left. \left. + (L\mathcal{N}_2 + \mathcal{N}_{\varphi}) F_{2n_1, 2n_2+2, c}^{\mathcal{O}}(\bar{\theta}_2, \theta_1, \dots, \theta_{\tilde{N}}) \right. \right. \\
 &\quad \left. \left. + F_{2n_1, 2n_2+4, c}^{\mathcal{O}}(\bar{\theta}_1, \bar{\theta}_2, \theta_1, \dots, \theta_{\tilde{N}}) \right\} \right]. \tag{4.20}
 \end{aligned}$$

5 Numerical results

The last ingredient needed to compute the form factor series (2.15) is the numerical evaluation of the connected diagonal form factors of the T_2 model. This is rather nontrivial to perform in a sufficiently quick and numerically stable way for large number of particles. Since this is a task that is also important for evaluating the spectral series for correlation functions at finite temperature [20, 23, 24, 61], we describe the required tricks in appendix A. The procedure can be straightforwardly generalized to connected diagonal form factors in other integrable models.

label	0	1	2	11	12	111	22	112
# of type-1 integrals	0	1	0	2	1	3	0	2
# of type-2 integrals	0	0	1	0	1	0	2	1
magnitude/L	0	m_1	m_2	$2m_1$	$m_1 + m_2$	$3m_1$	$2m_2$	$2m_1 + m_2$

Table 1. The terms incorporated from the form factor series into the numerical calculation.

In large volume a rough estimate for the magnitude of the terms in the series comes from the behaviour of the filling fractions

$$\prod_i \frac{1}{1 + e^{\varepsilon_{\alpha_j}(\theta_j)}} \lesssim e^{-(\sum_i m_i)L}$$

where m_i are the masses of the particles contained in the given state. Using this estimate we can identify the terms of the series that give the dominant contribution to the expectation value. However, with decreasing volume the ordering of the terms can change depending on the behaviour of the pseudo-energy functions; in addition, to maintain accuracy it is necessary to add progressively more terms. As a result, the form factor series (2.15) is effectively an infrared (low energy/large volume) expansion for the expectation value.

For our numerical calculations we implemented the terms with less than 4 integrals, since for higher terms the number of integrals and the size of the form factors makes the numerical integration too time-consuming; in addition, the terms incorporated already show an excellent agreement with the conjecture. Table 1 shows the terms calculated for numerics.

In the T_2 model there are two primary operators, namely $\Phi_{1,3}$ and $\Phi_{1,2}$. $\Phi_{1,3}$ is the operator perturbing the UV limit of the model, hence it's proportional to Θ

$$\Theta = 2\pi\lambda(2h_{1,3} - 2)\Phi_{1,3} \tag{5.1}$$

where $h_{1,3}$ is the conformal weight of the $\Phi_{1,3}$ and λ is the coupling constant of the perturbation that is proportional to the mass gap [62]

$$\lambda = \kappa m_1^{2-2h_{1,3}} \tag{5.2}$$

with

$$\kappa = -0.04053795542\dots \tag{5.3}$$

Because of this, the form factor series for $\Phi_{1,3}$ is equivalent to the TBA equation as proved in section 3, and the numerical calculation for $\Phi_{1,3}$ is therefore not a real further test for the general validity of the form factor series (2.15). However it is still useful since with its expectation value known from TBA equations one can get an independent check of the numerical precision of TCSA, and the convergence of the form factor series. For $\Phi_{1,2}$ there is only TCSA and the form factor series, with the numerical deviation for $\Phi_{1,3}$ setting the expected precision for the agreement between them.

For the numerical integration we used the Cuba library routines [63], called from inside Wolfram Mathematica [64].




Figure 5. ${}_1\langle\{1\}|\Phi_{1,3}|\{1\}\rangle_1$ evaluated by the form factor series including different contributions, RG extrapolated TCSA and TBA. Here and in all subsequent plots matrix elements are given in units of m_1 .

5.1 Moving one-particle state, $s = 1$

For the moving type-1 excited state with momentum quantum number $s = 1$, figure 5 shows the expectation value ${}_1\langle\{1\}|\Phi_{1,3}|\{1\}\rangle_1$ calculated with RG-extrapolated TCSA, from TBA together with the results from the form factor series (4.20) obtained by adding progressively more terms. The precision of the TBA is of the order 10^{-12} and comparing it with the TCSA data, we find that the precision of the RG-extrapolated TCSA is of order $10^{-6} - 10^{-7}$. Table 2 shows the difference between the form factor series with different terms involved and the TCSA data. For volume $m_1L > 5$ the difference between the form factor series up to and including the 112 term, and the TCSA is in the order of the TCSA error, and including more terms make the agreement better for smaller volume as well.

For $\Phi_{1,2}$ the results for the quantity $i{}_1\langle\{1\}|\Phi_{1,2}|\{1\}\rangle_1$ are shown in figure 6, and the difference between the form factor series and the TCSA is given in table 3. We note that since from (A.14) it follows that the matrix elements of $\Phi_{1,2}$ are imaginary, here and in all subsequent figures and tables concerning $\Phi_{1,2}$ we multiply all data by i . The form factor series shows excellent agreement with the TCSA for volume $m_1L > 5$, and again including more terms the agreement is better for smaller volumes. As noted before, the correctness of the form factor series does not follow from TBA, hence this is a nontrivial verification of the form factor series.

5.2 Zero-momentum one-particle state, $s = 0$

As seen for the $s = 1$ case the form factor series reproduce the expectation value of local operators with very good precision in large volume, even by including only few terms from




Figure 6. $i_1\langle\{1}\|\Phi_{1,2}|\{1}\rangle_1$ evaluated by the form factor series including different contributions and RG-extrapolated TCSA.

the series. The expectation is that for any state in small volume it is necessary to include higher contributions of the series, but for any desired accuracy a finite number of them is sufficient.

As shown below, this expectation is challenged by the nontrivial transition in the TBA equation for standing state at r_c . Figure 7 and 8 shows the result for the expectation values of $\Phi_{1,3}$ and $\Phi_{1,2}$, while table 4 and 5 list the numerical deviations from RG-extrapolated TCSA.

For large volume ($m_1 L \gtrsim 6$) the agreement between the form factor series and the TCSA is again excellent. However, towards the critical volume ($r_c \sim 2.66$) the terms of the form factor series tend to diverge. This can be understood from the fact that the total density of the states

$$\rho_{\text{tot}} = \det \mathcal{K} = L^2 \mathcal{N}_1 \mathcal{N}_2 + \mathcal{N}_\varphi L (\mathcal{N}_1 + \mathcal{N}_2) \quad (5.4)$$

which the denominator of the form factor series (4.20), is zero at r_c . Figure 9 shows the behaviour of ρ_{tot} around r_c ; fitting the location where $\rho_{\text{tot}}(\tilde{r}_c) = 0$, the result is

$$\tilde{r}_c = 2.6646510318 \quad (5.5)$$

which is perfect agreement with the value of r_c obtained in [43].

The reason for ρ_{tot} vanishing at r_c can be understood from the excited TBA. Since the active singularities coincide at this point, the density which is the Jacobi determinant of the quantization condition for the active singularities, is zero due to the degeneracy. Such singularities of the density were observed previously for the finite volume form factors formula in [45]; however in that case including the exponential corrections resolved (or at

$m_1 L$	0	+1	+2	+11	+12	+111	+22	+112
0.5	-0.8115	$-1 \cdot 10^{-2}$	1.039	0.7047	-0.3024	-0.1995	-0.8307	-0.3166
1	-0.6554	-0.1143	0.5094	0.3528	$-7 \cdot 10^{-2}$	$-4 \cdot 10^{-2}$	-0.2652	-0.1089
1.5	-0.5173	-0.1256	0.2649	0.1813	$-2 \cdot 10^{-2}$	$-2 \cdot 10^{-3}$	$-9 \cdot 10^{-2}$	$-4 \cdot 10^{-2}$
2	-0.3966	-0.1079	0.1362	$9 \cdot 10^{-2}$	$-2 \cdot 10^{-3}$	$4 \cdot 10^{-3}$	$-3 \cdot 10^{-2}$	$-1 \cdot 10^{-2}$
2.5	-0.2947	$-8 \cdot 10^{-2}$	$7 \cdot 10^{-2}$	$4 \cdot 10^{-2}$	$5 \cdot 10^{-4}$	$3 \cdot 10^{-3}$	$-1 \cdot 10^{-2}$	$-4 \cdot 10^{-3}$
3	-0.2124	$-6 \cdot 10^{-2}$	$3 \cdot 10^{-2}$	$2 \cdot 10^{-2}$	$6 \cdot 10^{-4}$	$2 \cdot 10^{-3}$	$-3 \cdot 10^{-3}$	$-1 \cdot 10^{-3}$
3.5	-0.1489	$-4 \cdot 10^{-2}$	$1 \cdot 10^{-2}$	$8 \cdot 10^{-3}$	$3 \cdot 10^{-4}$	$6 \cdot 10^{-4}$	$-9 \cdot 10^{-4}$	$-3 \cdot 10^{-4}$
4	-0.102	$-2 \cdot 10^{-2}$	$6 \cdot 10^{-3}$	$3 \cdot 10^{-3}$	$1 \cdot 10^{-4}$	$2 \cdot 10^{-4}$	$-2 \cdot 10^{-4}$	$-5 \cdot 10^{-5}$
4.5	$-7 \cdot 10^{-2}$	$-1 \cdot 10^{-2}$	$3 \cdot 10^{-3}$	$1 \cdot 10^{-3}$	$4 \cdot 10^{-5}$	$8 \cdot 10^{-5}$	$-5 \cdot 10^{-5}$	$-1 \cdot 10^{-5}$
5	$-5 \cdot 10^{-2}$	$-7 \cdot 10^{-3}$	$1 \cdot 10^{-3}$	$4 \cdot 10^{-4}$	$1 \cdot 10^{-5}$	$2 \cdot 10^{-5}$	$-1 \cdot 10^{-5}$	$-2 \cdot 10^{-6}$
5.5	$-3 \cdot 10^{-2}$	$-4 \cdot 10^{-3}$	$5 \cdot 10^{-4}$	$1 \cdot 10^{-4}$	$4 \cdot 10^{-6}$	$7 \cdot 10^{-6}$	$-3 \cdot 10^{-6}$	$-4 \cdot 10^{-7}$
6	$-2 \cdot 10^{-2}$	$-2 \cdot 10^{-3}$	$2 \cdot 10^{-4}$	$4 \cdot 10^{-5}$	$8 \cdot 10^{-7}$	$2 \cdot 10^{-6}$	$-7 \cdot 10^{-7}$	$-2 \cdot 10^{-7}$
6.5	$-1 \cdot 10^{-2}$	$-1 \cdot 10^{-3}$	$7 \cdot 10^{-5}$	$1 \cdot 10^{-5}$	$4 \cdot 10^{-7}$	$6 \cdot 10^{-7}$	$3 \cdot 10^{-8}$	$1 \cdot 10^{-7}$
7	$-8 \cdot 10^{-3}$	$-5 \cdot 10^{-4}$	$3 \cdot 10^{-5}$	$5 \cdot 10^{-6}$	$4 \cdot 10^{-7}$	$4 \cdot 10^{-7}$	$3 \cdot 10^{-7}$	$3 \cdot 10^{-7}$

Table 2. The difference between the evaluations of ${}_1\langle\{1}\mid\Phi_{1,3}\mid\{1}\rangle_1$ from the RG-extrapolated TCSA and the form factor series, depending on the multi-particle contributions included in the latter.

least shifted) the singularity. The present situation is different as all exponential corrections to the density are already included. To resolve the singularity it would be necessary to include every term of the form factor series, to compensate the zero of the denominator.

This conclusion is also supported by the behaviour of the pseudo-energies and the filling fractions around r_c . Approaching r_c the filling fractions no more suppress the higher order terms in the series and the ordering of terms by their magnitude is not valid anymore, i.e. every terms is important in the series. This is consistent with the procedure of desingularization, whereby to describe the excited state level with the TBA equation under r_c it was necessary to redefine the pseudo-energy to have a form which is finite and convergent under iterations.

For the form factor series a similar rearrangement is necessary close to and under r_c . Unfortunately such a rearrangement is not yet known, and this sets the practical validity of the form factor series to IR regions where no nontrivial transitions occur in the TBA equations.

6 Conclusions and outlook

In this paper we presented a conjecture for exact expectation values of local operators in finite volume excited states, in integrable quantum field theories. This conjecture is an extension of an earlier one [39] to models with a non-trivial bootstrap structure. The conjecture was supported by a combination of analytic and numerical evidence. In the latter

$m_1 L$	0	+1	+2	+11	+12	+111	+22	+112
0.5	-1.203	-0.5712	1.113	0.9523	0.1086	0.1457	-0.7935	-0.5066
1	-0.9067	-0.4563	0.5275	0.4431	$7 \cdot 10^{-2}$	$8 \cdot 10^{-2}$	-0.252	-0.1539
1.5	-0.6803	-0.3379	0.2661	0.2171	$4 \cdot 10^{-2}$	$4 \cdot 10^{-2}$	$-9 \cdot 10^{-2}$	$-5 \cdot 10^{-2}$
2	-0.5	-0.2361	0.1325	0.1038	$2 \cdot 10^{-2}$	$2 \cdot 10^{-2}$	$-3 \cdot 10^{-2}$	$-2 \cdot 10^{-2}$
2.5	-0.3581	-0.1561	$6 \cdot 10^{-2}$	$5 \cdot 10^{-2}$	$7 \cdot 10^{-3}$	$9 \cdot 10^{-3}$	$-9 \cdot 10^{-3}$	$-5 \cdot 10^{-3}$
3	-0.2498	$-1 \cdot 10^{-1}$	$3 \cdot 10^{-2}$	$2 \cdot 10^{-2}$	$3 \cdot 10^{-3}$	$3 \cdot 10^{-3}$	$-3 \cdot 10^{-3}$	$-1 \cdot 10^{-3}$
3.5	-0.1701	$-6 \cdot 10^{-2}$	$1 \cdot 10^{-2}$	$8 \cdot 10^{-3}$	$1 \cdot 10^{-3}$	$1 \cdot 10^{-3}$	$-8 \cdot 10^{-4}$	$-3 \cdot 10^{-4}$
4	-0.1136	$-3 \cdot 10^{-2}$	$5 \cdot 10^{-3}$	$3 \cdot 10^{-3}$	$3 \cdot 10^{-4}$	$4 \cdot 10^{-4}$	$-2 \cdot 10^{-4}$	$-6 \cdot 10^{-5}$
4.5	$-7 \cdot 10^{-2}$	$-2 \cdot 10^{-2}$	$2 \cdot 10^{-3}$	$1 \cdot 10^{-3}$	$9 \cdot 10^{-5}$	$1 \cdot 10^{-4}$	$-5 \cdot 10^{-5}$	$-1 \cdot 10^{-5}$
5	$-5 \cdot 10^{-2}$	$-1 \cdot 10^{-2}$	$9 \cdot 10^{-4}$	$4 \cdot 10^{-4}$	$3 \cdot 10^{-5}$	$3 \cdot 10^{-5}$	$-1 \cdot 10^{-5}$	$-3 \cdot 10^{-6}$
5.5	$-3 \cdot 10^{-2}$	$-5 \cdot 10^{-3}$	$4 \cdot 10^{-4}$	$1 \cdot 10^{-4}$	$6 \cdot 10^{-6}$	$8 \cdot 10^{-6}$	$-3 \cdot 10^{-6}$	$-2 \cdot 10^{-6}$
6	$-2 \cdot 10^{-2}$	$-3 \cdot 10^{-3}$	$1 \cdot 10^{-4}$	$4 \cdot 10^{-5}$	$2 \cdot 10^{-7}$	$8 \cdot 10^{-7}$	$-2 \cdot 10^{-6}$	$-2 \cdot 10^{-6}$
6.5	$-1 \cdot 10^{-2}$	$-1 \cdot 10^{-3}$	$5 \cdot 10^{-5}$	$1 \cdot 10^{-5}$	$-1 \cdot 10^{-6}$	$-1 \cdot 10^{-6}$	$-2 \cdot 10^{-6}$	$-2 \cdot 10^{-6}$
7	$-8 \cdot 10^{-3}$	$-6 \cdot 10^{-4}$	$2 \cdot 10^{-5}$	$2 \cdot 10^{-6}$	$-2 \cdot 10^{-6}$	$-2 \cdot 10^{-6}$	$-2 \cdot 10^{-6}$	$-2 \cdot 10^{-6}$

Table 3. The difference between the evaluations of ${}_1\langle\{1\}|\Phi_{1,2}|\{1\}\rangle_1$ from the RG-extrapolated TCSA and the form factor series, depending on the multi-particle contributions included in the latter.

$m_1 L$	0	+1	+2	+11	+12	+111	+22	+112
4	0.3114	-0.4977	-0.9218	-0.4061	0.3695	0.2265	0.494	$9 \cdot 10^{-2}$
4.5	-0.1529	-0.1416	$-8 \cdot 10^{-2}$	$-2 \cdot 10^{-2}$	$3 \cdot 10^{-2}$	$2 \cdot 10^{-2}$	$3 \cdot 10^{-2}$	$3 \cdot 10^{-3}$
5	-0.1637	$-6 \cdot 10^{-2}$	$9 \cdot 10^{-4}$	$5 \cdot 10^{-3}$	$3 \cdot 10^{-3}$	$2 \cdot 10^{-3}$	$1 \cdot 10^{-3}$	$-3 \cdot 10^{-5}$
5.5	-0.114	$-3 \cdot 10^{-2}$	$4 \cdot 10^{-3}$	$2 \cdot 10^{-3}$	$3 \cdot 10^{-4}$	$3 \cdot 10^{-4}$	$-4 \cdot 10^{-5}$	$-3 \cdot 10^{-5}$
6	$-7 \cdot 10^{-2}$	$-1 \cdot 10^{-2}$	$2 \cdot 10^{-3}$	$8 \cdot 10^{-4}$	$5 \cdot 10^{-5}$	$5 \cdot 10^{-5}$	$-2 \cdot 10^{-5}$	$-5 \cdot 10^{-6}$
6.5	$-4 \cdot 10^{-2}$	$-6 \cdot 10^{-3}$	$7 \cdot 10^{-4}$	$2 \cdot 10^{-4}$	$7 \cdot 10^{-6}$	$1 \cdot 10^{-5}$	$-5 \cdot 10^{-6}$	$-9 \cdot 10^{-7}$
7	$-3 \cdot 10^{-2}$	$-3 \cdot 10^{-3}$	$2 \cdot 10^{-4}$	$6 \cdot 10^{-5}$	$1 \cdot 10^{-6}$	$2 \cdot 10^{-6}$	$-1 \cdot 10^{-6}$	$-4 \cdot 10^{-7}$
7.5	$-2 \cdot 10^{-2}$	$-1 \cdot 10^{-3}$	$8 \cdot 10^{-5}$	$2 \cdot 10^{-5}$	$1 \cdot 10^{-7}$	$4 \cdot 10^{-7}$	$-2 \cdot 10^{-7}$	$-1 \cdot 10^{-7}$

Table 4. The difference between the evaluations of ${}_1\langle\{0\}|\Phi_{1,3}|\{0\}\rangle_1$ from the RG-extrapolated TCSA and the form factor series, depending on the multi-particle contributions included in the latter.

regard we note that the generalization of the conjecture was necessary so that it could be evaluated for models in which we have access to non-perturbative numerical techniques (in our particular case, the truncated conformal space approach) to evaluate finite volume expectation values directly from Hamiltonian quantum field theory.




Figure 7. $i_1\langle\{0\}|\Phi_{1,3}|\{0\}\rangle_1$ calculated by the form factor series including different contributions, RG extrapolated TCSA and TBA.




Figure 8. $i_1\langle\{0\}|\Phi_{1,2}|\{0\}\rangle_1$ calculated by the form factor series including different contributions and RG extrapolated TCSA.

m_1L	0	+1	+2	+11	+12	+111	+22	+112
4	-0.5274	-0.3774	-0.6113	-0.3511	0.2018	0.1321	0.3664	0.1129
4.5	-0.5513	-0.139	$-2 \cdot 10^{-2}$	$-9 \cdot 10^{-3}$	$2 \cdot 10^{-2}$	$1 \cdot 10^{-2}$	$2 \cdot 10^{-2}$	$4 \cdot 10^{-3}$
5	-0.3645	$-6 \cdot 10^{-2}$	$1 \cdot 10^{-2}$	$7 \cdot 10^{-3}$	$1 \cdot 10^{-3}$	$1 \cdot 10^{-3}$	$3 \cdot 10^{-4}$	$4 \cdot 10^{-6}$
5.5	-0.2176	$-3 \cdot 10^{-2}$	$7 \cdot 10^{-3}$	$3 \cdot 10^{-3}$	$2 \cdot 10^{-4}$	$2 \cdot 10^{-4}$	$-1 \cdot 10^{-4}$	$-3 \cdot 10^{-5}$
6	-0.1255	$-1 \cdot 10^{-2}$	$3 \cdot 10^{-3}$	$9 \cdot 10^{-4}$	$3 \cdot 10^{-5}$	$5 \cdot 10^{-5}$	$-3 \cdot 10^{-5}$	$-5 \cdot 10^{-6}$
6.5	$-7 \cdot 10^{-2}$	$-7 \cdot 10^{-3}$	$9 \cdot 10^{-4}$	$2 \cdot 10^{-4}$	$5 \cdot 10^{-6}$	$1 \cdot 10^{-5}$	$-6 \cdot 10^{-6}$	$-2 \cdot 10^{-6}$
7	$-4 \cdot 10^{-2}$	$-3 \cdot 10^{-3}$	$3 \cdot 10^{-4}$	$6 \cdot 10^{-5}$	$-2 \cdot 10^{-7}$	$1 \cdot 10^{-6}$	$-2 \cdot 10^{-6}$	$-1 \cdot 10^{-6}$
7.5	$-2 \cdot 10^{-2}$	$-1 \cdot 10^{-3}$	$9 \cdot 10^{-5}$	$1 \cdot 10^{-5}$	$-1 \cdot 10^{-6}$	$-1 \cdot 10^{-6}$	$-2 \cdot 10^{-6}$	$-2 \cdot 10^{-6}$

Table 5. The difference between the evaluations of $i_1 \langle \{0\} | \Phi_{1,2} | \{0\} \rangle_1$ from the RG-extrapolated TCSA and the form factor series, depending on the multi-particle contributions included in the latter.




Figure 9. ρ_{tot} for $s=0$.

The conjecture (2.1) also leads to interesting new results concerning finite volume form factors:

1. In theories where a particle is represented by several active singularities, the particle can be considered as a bound state of the active singularities. In infinite volume this does not make any difference due to bootstrap equations satisfied by the scattering matrix, but in finite volume the composite nature of the particles gives exponential corrections, which are exactly the μ -term corrections to the form factor described in [35]. However, a given particle can generally be represented as a bound states of

others in multiple different ways, which means that the prescription for the μ -terms used in [35] is eventually ambiguous.

However, according to (2.1) the excited state TBA equations give a clear prescription. Namely, for a correct representation of the μ -term, one must take the particular bound state representation of the particle states as specified by the singularity arrangement of the excited TBA equations. Since the singularity arrangement corresponds to the given state (finite volume energy level), one expects that these prescriptions remain valid for non-diagonal matrix elements as well.

2. In line with the usual terminology of finite volume corrections [33, 59, 60], the terms in (2.15) containing rapidity integration, originating from either the quantization conditions (2.7) or the dressed form factors (2.10), give so-called F -term corrections, which describe virtual particle loops winding around the finite volume cylinder. Contrary to μ -terms, however, for the moment it is not clear how to generalize this to non-diagonal matrix elements.

We stress that the series (2.15) is exact, at least for values of the volume where it is convergent; however, there are still some open problems and interesting directions to pursue, as we discuss below.

A lesson learned from studying the generalized conjecture (2.15) is that for certain excited states it is not valid for the whole volume range in its present form. As discussed in subsection 5.2, this happens for states where the corresponding excited TBA singularity configuration undergoes some rearrangement below some critical value of the volume. Note that for the sinh-Gordon model studied in [39] this cannot happen due to the imaginary part of singularity positions being fixed to $\pi/2$. In the series (2.15) this is manifested in a singular behaviour of the terms, which prevent the extension of the series below the critical volume.

For the trace of the stress-energy tensor, the desingularized excited TBA gives such an extension. However, it is presently unclear how to implement the desingularization procedure directly for the series (2.15), so the description cannot be extended to other operators. In order to achieve that, one needs to separate the singularly behaved contributions and re-sum them to all orders, which is not a straightforward task.

In this context we remark that based on the available studies of excited state TBA systems, the problematic rearrangement is only expected to happen for excited states for which some singularities are “stuck in the middle”, as opposed to going to the left/right asymptotic regions logarithmically with decreasing volume. Therefore, for most states the conjecture is expected to be valid for any value of the volume; obviously the smaller is the volume, the larger is the number of terms needed for a given precision. For states with a non-trivial transition in their singularity structure, the conjecture in its present form is expected to be valid for values of the volume above the critical one.

Another interesting issue is to extend the series to theories with non-diagonal scattering, starting with the series introduced in [61]. The formalism of finite volume form factors has been partially extended to these theories [65, 66]. Unfortunately, it is exactly the general form of diagonal matrix elements that is at present not known in full generality,

preventing an application of the methods in [61] to all orders. In this case an extension of the series (2.15) could provide interesting new results.

Furthermore, the form of the terms in the series (2.15) is very suggestive for an extension to finite volume matrix elements between two different states; however, finding the precise form of such an extension is still an open question. This would also give a prescription for the F -terms mentioned above.

Finally, establishing the relation between the present framework, and the approaches based on separation of variables [14, 15] or fermionic structures [16–18] would be of interest, as it could lead to more efficient construction of finite size matrix elements and a deeper understanding of the underlying principles.

Acknowledgments

IMSZ is grateful to Roberto Tateo for the help with excited state TBA numerics and his hospitality in Turin, and also to the INFN and Campus Hungary Scholarship for financial support of the visit. IMSZ was also supported by funding from the People Programme (Marie Curie Actions) of the European Union’s Seventh Framework Programme FP7/2007-2013/ under REA Grant Agreement No 317089 (GATIS). BP and GT were supported by the Momentum grant LP2012-50/2014 of the Hungarian Academy of Sciences.

A Evaluation of connected diagonal form factors

Here we present a fast and efficient way to evaluate the connected diagonal form factors (defined in section 2), which proceeds via the so-called symmetric form factors.

A.1 Form factors of the T_2 model

The form factors of primary fields in the T_2 model were constructed in [67]. For states containing only type-1 particles they are the same as form factors of certain exponential operators in the sinh-Gordon model with a specific value for the coupling constant. The form factors of the exponential operator

$$e^{kg\Phi} \tag{A.1}$$

in the sinh-Gordon model

$$\mathcal{L} = \frac{1}{2} \partial_\mu \Phi \partial^\mu \Phi - \lambda \cosh g\Phi \tag{A.2}$$

have the following form [68]

$$F_n^{(k)}(\theta_1, \dots, \theta_n) = H_n[k] Q_n(x_1, \dots, x_n) \prod_{i < j} \frac{F_{\min}(\theta_{ij})}{x_i + x_j} \tag{A.3}$$

where

$$[n] = \frac{\sin n\pi \frac{B}{2}}{\sin \pi \frac{B}{2}} \quad B = \frac{2g^2}{8\pi + g^2} \tag{A.4}$$

and the minimal two-particle form factor is

$$\begin{aligned}
 F_{\min}(\theta, B) &= \mathcal{N} \exp \left[8 \int_0^\infty \frac{dx \sinh \frac{xB}{4} \sinh \frac{x}{2} \left(1 - \frac{B}{2}\right) \sinh \frac{x}{2}}{x \sinh^2 x} \sin^2 \left[\frac{x(i\pi - \theta)}{2\pi} \right] \right] \\
 &= \mathcal{N} e^{I(\theta, B)} \\
 \mathcal{N} &= \exp \left[-4 \int \frac{dx \sinh \frac{xB}{4} \sinh \frac{x}{2} \left(1 - \frac{B}{2}\right) \sinh \frac{x}{2}}{x \sinh^2 x} \right]
 \end{aligned} \tag{A.5}$$

with

$$\begin{aligned}
 I(\theta, B) &= 8 \int_0^\infty \frac{dx \sinh \frac{xB}{4} \sinh \frac{x}{2} \left(1 - \frac{B}{2}\right) \sinh \frac{x}{2}}{x \sinh^2 x} \sin^2 \left[\frac{x(i\pi - \theta)}{2\pi} \right] = \\
 &= 8 \int_0^\infty \frac{dx \sinh \frac{xB}{4} \sinh \frac{x}{2} \left(1 - \frac{B}{2}\right) \sinh \frac{x}{2}}{x \sinh^2 x} [N+1 - Ne^{-2x}] e^{-2Nx} \sin^2 \left[\frac{x(i\pi - \theta)}{2\pi} \right] + \\
 &+ \sum_{k=0}^{N-1} (k+1) \left\{ \log \nu \left[\theta, k + \frac{1}{2} \right] + \log \nu \left[\theta, k + \frac{3}{2} - \frac{B}{4} \right] + \log \nu \left[\theta, k + 1 + \frac{B}{4} \right] - \right. \\
 &\left. - \log \nu \left[\theta, k + \frac{3}{2} \right] - \log \nu \left[\theta, k + \frac{1}{2} + \frac{B}{4} \right] - \log \nu \left[\theta, k + 1 - \frac{B}{4} \right] \right\}
 \end{aligned} \tag{A.6}$$

and

$$\nu(\theta, a) = 1 + \frac{(i\pi - \theta)^2}{(2\pi)^2 a^2}. \tag{A.7}$$

The normalization factors read

$$H_n = \left(\frac{4 \sin \frac{\pi B}{2}}{F_{\min}(i\pi, B)} \right)^{n/2}. \tag{A.8}$$

Introducing the notations

$$x_i = e^{\theta_i} \quad \text{and} \quad \theta_{ij} = \theta_i - \theta_j \tag{A.9}$$

the Q_n are given in a determinant form

$$\begin{aligned}
 Q_n(k) &= \det M_{ij}(k) \\
 M_{ij}(k) &= [i - j + k] \sigma_{2i-j}^{(n)}, \quad j = 1, \dots, n-1
 \end{aligned} \tag{A.10}$$

where the $\sigma_k^{(n)}$ are the elementary symmetric polynomials of order k in the n variables x_1, \dots, x_n defined by

$$\prod_{i=1}^n (x + x_i) = \sum_k x^{n-k} \sigma_k^{(n)}(x_1, \dots, x_n) \tag{A.11}$$

(this means in particular that $\sigma_n^{(k)} = 0$ for $k > n$ or $k < 0$). To obtain the form factors of local operators in the T_2 model it is necessary to set the coupling as

$$B = -\frac{4}{5}. \tag{A.12}$$

Following the procedure in appendix A of [69], the form factors for type-2 particles can be efficiently calculated with the help of writing the bootstrap fusion in the form

$$F_{2\dots}(\theta, \dots) = \Gamma_{11}^2 F_{11\dots}(\theta - i\bar{u}_{12}^1, \theta + i\bar{u}_{12}^1, \dots) \quad (\text{A.13})$$

where $\bar{u}_{12}^1 = \frac{\pi}{5}$ and $\Gamma_{11}^2 = \sqrt{2 \tan(2\pi/5)}$.

The value of $k = 1$ corresponds to the operator $\Phi_{1,3}$ and $k = 2$ to the operator $\Phi_{1,2}$. The above form factors are normalized so that the vacuum expectation value of the field is 1. To obtain the conformal normalization used in TCSCA, it is necessary to multiply the form factors with the exact vacuum expectation values known from [70]

$$\begin{aligned} \langle \Phi_{1,2} \rangle &= -2.3251365527 \dots \times im_1^{-4/7} \\ \langle \Phi_{1,3} \rangle &= 2.2695506880 \dots \times m_1^{-6/7}. \end{aligned} \quad (\text{A.14})$$

A.2 Symmetric form factors

The symmetric form factors are defined as

$$\begin{aligned} &F_{\underbrace{1, \dots, 1}_n, \underbrace{2, \dots, 2}_m}^s(\theta_1, \dots, \theta_n, \theta_{n+1}, \dots, \theta_{n+m}) \\ &= \lim_{\epsilon \rightarrow 0} F_{\underbrace{2, \dots, 2}_m, \underbrace{1, \dots, 1}_n, \underbrace{1, \dots, 1}_n, \underbrace{2, \dots, 2}_m}(\theta_{n+m} + i\pi + \epsilon, \dots, \theta_{n+1} + i\pi + \epsilon, \\ &\quad, \theta_n + i\pi + \epsilon, \dots, \theta_1 + i\pi + \epsilon, \theta_1, \dots, \theta_n, \theta_{n+1}, \dots, \theta_{n+m}) \end{aligned} \quad (\text{A.15})$$

where there are n numbers of type-1 particles and m numbers of type-2 particles. This definition corresponds to a particular specification for the direction of the limit to the diagonal matrix element. To compute the above expression, we use fusion (A.13) for type-2 particles, and calculate the limit in terms of a form factor with $2(n + 2m)$ type-1 particles.

A.2.1 Denominator and minimal form factors

The denominator has the following form

$$\prod_{i < j}^{2(n+2m)} (\tilde{x}_i - \tilde{x}_j) \quad (\text{A.16})$$

where

$$\tilde{x}_i = \begin{cases} -e^\epsilon x_{n+2m+1-i} & i \leq n + 2m \\ x_{i-n-2m} & i > n + 2m. \end{cases} \quad (\text{A.17})$$

To leading order in ϵ , the denominator of the symmetric form factor takes the form

$$(-\epsilon)^{n+2m} \left[\prod_i^{n+2m} x_i \right] \left[\prod_{i < j}^{n+2m} (x_i^2 - x_j^2)^2 \right]. \quad (\text{A.18})$$

From the minimal form factor part we get an $F_{\min}(i\pi)$ factor for every particle when the rapidities meet with their crossed version, i.e. a factor of $[F_{\min}(i\pi)]^{n+2m}$ altogether.

To simplify the other contribution we use the following relation for the sinh-Gordon form factors [71]

$$F_{\min}(i\pi + \vartheta) F_{\min}(\vartheta) = \frac{\sinh(\vartheta)}{\sinh(\vartheta) + i \sin\left(\frac{\pi B}{2}\right)}. \quad (\text{A.19})$$

There result for two type-1 particle including the denominator term is

$$\frac{F_{\min}(\vartheta_{ij}) F_{\min}(\vartheta_{ji}) F_{\min}(i\pi + \vartheta_{ij}) F_{\min}(i\pi + \vartheta_{ji})}{(x_i^2 - x_j^2)^2} = \frac{1}{(x_i^2 - x_j^2)^2 + 4x_i^2 x_j^2 \sin^2\left(\frac{\pi B}{2}\right)}. \quad (\text{A.20})$$

The result between one type-1 and a type-2 rapidity is

$$\frac{1}{(x_i^2 - x_{j,+}^2)^2 + 4x_i^2 x_{j,+}^2 \sin^2\left(\frac{\pi B}{2}\right)} \times \frac{1}{(x_i^2 - x_{j,-}^2)^2 + 4x_i^2 x_{j,-}^2 \sin^2\left(\frac{\pi B}{2}\right)} \quad (\text{A.21})$$

where $x_{j,\pm} = x_j e^{\pm i\bar{u}_{12}}$. The result for rapidities from the same type-2 particle is ($x_i = x e^{-i\bar{u}_{12}}, x_j = x e^{+i\bar{u}_{12}}$)

$$\frac{[F_{\min}(\vartheta_{ij})]^2 F_{\min}(i\pi + \vartheta_{ij}) F_{\min}(i\pi + \vartheta_{ji})}{(x_i^2 - x_j^2)^2} = \frac{-1}{16x^4 \sin^2\left(\frac{2\pi}{5}\right)}. \quad (\text{A.22})$$

The result for rapidities from different type-2 particles is

$$\begin{aligned} & \frac{[F_{\min}(\theta_{ij}) F_{\min}(i\pi + \theta_{ij})]^2 [F_{\min}(\theta_{ji}) F_{\min}(i\pi + \theta_{ji})]^2}{(x_i^2 - x_j^2)^4 (x_i^2 e^{i2\bar{u}} - x_j^2 e^{-i2\bar{u}})^2 (x_i^2 e^{-i2\bar{u}} - x_j^2 e^{i2\bar{u}})^2} \\ & \times [F_{\min}(\theta_{ij} - 2i\bar{u}) F_{\min}(i\pi + \theta_{ij} - 2i\bar{u})] [F_{\min}(\theta_{ij} + 2i\bar{u}) F_{\min}(i\pi + \theta_{ij} + 2i\bar{u})] \\ & \times [F_{\min}(\theta_{ji} - 2i\bar{u}) F_{\min}(i\pi + \theta_{ji} - 2i\bar{u})] [F_{\min}(\theta_{ji} + 2i\bar{u}) F_{\min}(i\pi + \theta_{ji} + 2i\bar{u})] \\ & = \left[(x_i^2 - x_j^2)^2 + 4x_i^2 x_j^2 \sin^2(2\bar{u}) \right]^{-2} \left[(x_i^2 e^{i2\bar{u}} - x_j^2 e^{-i2\bar{u}})^2 + 4x_i^2 x_j^2 \sin^2(2\bar{u}) \right]^{-1} \\ & \times \left[(x_i^2 e^{-i2\bar{u}} - x_j^2 e^{i2\bar{u}})^2 + 4x_i^2 x_j^2 \sin^2(2\bar{u}) \right]^{-1}. \end{aligned} \quad (\text{A.23})$$

A.2.2 Symmetric polynomial part

There are $2(n+2m)$ type-1 rapidities due to the fusion, so the polynomial part of the form factor is a determinant of a $(2(n+2m)-1) \times (2(n+2m)-1)$ matrix

$$M_{ij}(k) = [i - j + k] \sigma_{2i-j}^{(2(n+2m))}. \quad (\text{A.24})$$

In the $\varepsilon \rightarrow 0$ limit the symmetric polynomials are

$$\begin{aligned} & \sigma_l^{(2p)}(x_1, x_2, \dots, x_p, -e^\varepsilon x_1, -e^\varepsilon x_2, \dots, -e^\varepsilon x_p) \rightarrow \begin{cases} \sigma_l^{(2p)}(x_1, x_2, \dots, x_p, -x_1, -x_2, \dots, -x_p) + \mathcal{O}(\varepsilon) & l \text{ even} \\ \sum_{i=1}^p (-\varepsilon x_i) \sigma_{l-1}^{(2p-1)}(x_1, x_2, \dots, x_i, \dots, x_p, -x_1, -x_2, \dots, -x_{i-1}, -x_{i+1}, \dots, -x_p) + \mathcal{O}(\varepsilon^2). & l \text{ odd} \end{cases} \end{aligned} \quad (\text{A.25})$$

Since every term in the determinant contains $(n + 2m)$ factors of odd symmetrical polynomials, the determinant is proportional to ϵ^{n+2m} , which exactly cancels the ϵ powers in the denominator (A.18). From the definition (A.11) of the elementary symmetric polynomials it is easy to show that

$$\sigma_l^{(2p)}(x_1, x_2, \dots, x_p, -e^\epsilon x_1, -e^\epsilon x_2, \dots, -e^\epsilon x_p) \rightarrow \begin{cases} (-1)^{l/2} \sigma_{l/2}^{(p)}(x_1^2, x_2^2, \dots, x_p^2) & l \text{ even} \\ \sum_{i=1}^p (-\epsilon x_i) (-1)^{(l-1)/2} \sigma_{(l-1)/2}^{(p-1)}(x_1^2, x_2^2, \dots, x_{i-1}^2, x_{i+1}^2, \dots, x_p^2) & l \text{ odd} \end{cases} \quad (\text{A.26})$$

A.2.3 Result for symmetric form factor

Introducing the following definitions

$$\hat{\sigma}_l^p(x_1, x_2, \dots, x_p) = \begin{cases} (-1)^{l/2} \sigma_{l/2}^{(p)}(x_1^2, x_2^2, \dots, x_p^2) & l \text{ even} \\ (-1)^{(l-1)/2} \sum_{i=1}^p x_i \sigma_{(l-1)/2}^{(p-1)}(x_1^2, x_2^2, \dots, x_{i-1}^2, x_{i+1}^2, \dots, x_p^2) & l \text{ odd} \\ x_{i+1}^2, \dots, x_p^2 \end{cases}$$

$$\hat{Q}_{n+2m}(k) = \det \hat{M}_{ij}(k) \quad \hat{M}_{ij}(k) = [i - j + k] \hat{\sigma}_{2i-j}^{(n+m)}$$

$$i, j = 1, \dots, 2(n + 2m) - 1 \quad (\text{A.27})$$

and

$$F_{\min, \text{denom}}(x_1, x_2) = \begin{cases} F_{\min, \text{denom}}^{11}(x_1, x_2) & \text{type } -1 \leftrightarrow \text{type } -1 \\ F_{\min, \text{denom}}^{12}(x_1, x_2) & \text{type } -1 \leftrightarrow \text{type } -2 \\ F_{\min, \text{denom}}^{22, \text{dif}}(x_1, x_2) & \text{type } -2 \leftrightarrow \text{type } -2 \end{cases}$$

$$F_{\min, \text{denom}}^{11}(x_1, x_2) = \frac{1}{(x_i^2 - x_j^2)^2 + 4x_i^2 x_j^2 \sin^2(2\bar{u})}$$

$$F_{\min, \text{denom}}^{12}(x_1, x_2) = F_{\min, \text{denom}}^{11}(x_1, x_2 e^{i\bar{u}}) \times F_{\min, \text{denom}}^{11}(x_1, x_2 e^{-i\bar{u}})$$

$$F_{\min, \text{denom}}^{22, \text{dif}}(x_1, x_2) = [F_{\min, \text{denom}}^{11}(x_1, x_2)]^2 \times F_{\min, \text{denom}}^{11}(x_1 e^{i\bar{u}}, x_2 e^{-i\bar{u}}) \times F_{\min, \text{denom}}^{11}(x_1 e^{-i\bar{u}}, x_2 e^{i\bar{u}})$$

$$F_{\min, \text{denom}}^{22, \text{self}}(x) = \frac{-1}{16x^4 \sin^2(2\bar{u})} \quad (\text{A.28})$$

the symmetric form factor can be rewritten as:

$$F_{\underbrace{1, \dots, 1}_n, \underbrace{2, \dots, 2}_m}^s(\theta_1, \dots, \theta_n, \theta_{n+1}, \dots, \theta_{n+m}) \quad (\text{A.29})$$

$$= [k] \left(-4 \sin \frac{2\pi}{5} \right)^{n+2m} (\Gamma_{11}^2)^{2m} \hat{Q}_{n+2m}(x_1, \dots, x_{n+2m})$$

$$\times \frac{\prod_{i < j}^{n+m} F_{\min, \text{denom}}(x_i, x_j)}{\prod_{i=1}^n x_i} \prod_{j=n+1}^m \frac{F_{\min, \text{denom}}^{22, \text{self}}(x_j)}{x_j^2}.$$

For a large number of particles and/or large rapidities this formula is difficult to evaluate with the required numerical precision, because the determinant \hat{Q} is badly conditioned (the magnitude of its matrix elements differ by many orders). For a better precision it is necessary balance the matrix the following way:

$$\begin{aligned} \hat{Q}_{n+2m}(k) &= \left[\hat{\sigma}_1^{(n+2m)} / (n+2m) \right]^{2(n+2m)^2 - (n+2m)} \det \widetilde{M}_{ij}(k) \\ \widetilde{M}_{ij}(k) &= [i-j+k] \frac{\hat{\sigma}_{2^{i-j}}^{(n+m)}}{\left[\hat{\sigma}_1^{(n+2m)} / (n+2m) \right]^{2i-j}} \\ i, j &= 1, \dots, 2(n+2m) - 1. \end{aligned} \tag{A.30}$$

A.3 Evaluation of the connected diagonal form factors

There are two ways to calculate the connected diagonal form factors using the symmetric form factors. One way is to use the symmetric-connected relations derived in [20] in a recursive manner; this is a lengthy procedure for form factors with several variables and not very convenient for numerical calculations.

However, from the same relations it also follows that the connected diagonal form factor is the only part of the symmetric form factor that is fully periodic in all of its variables with period $i\pi$. This is related to unitarity and crossing invariance, which in theories with self-conjugate particles take the form

$$\begin{aligned} S_{\alpha\beta}(-\theta) &= S_{\alpha\beta}(\theta)^{-1} \\ S_{\alpha\beta}(\theta) &= S_{\alpha\beta}(i\pi - \theta). \end{aligned} \tag{A.31}$$

As a result the kernel functions (2.3) have the anti-periodicity property

$$\varphi_{\alpha\beta}(\theta + i\pi) = -\varphi_{\alpha\beta}(\theta). \tag{A.32}$$

Applying this property to the connected-symmetric relations of [20] leads to

$$\begin{aligned} F_n^c(\theta_1, \dots, \theta_n) &= \frac{1}{2^n} \sum_{\alpha_i=0,1} F_n^s(\theta_1 + \alpha_1 i\pi, \theta_2 + \alpha_2 i\pi, \dots, \theta_n + \alpha_n i\pi) \\ &= \frac{1}{2^{n-1}} \sum_{\alpha_i=0,1} F_n^s(\theta_1, \theta_2 + \alpha_2 i\pi, \dots, \theta_n + \alpha_n i\pi) \end{aligned} \tag{A.33}$$

which gives a faster and numerically much more stable evaluation.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] A. Leclair and G. Mussardo, *Finite temperature correlation functions in integrable QFT*, *Nucl. Phys. B* **552** (1999) 624 [[hep-th/9902075](#)] [[INSPIRE](#)].
- [2] C.-N. Yang and C.P. Yang, *Thermodynamics of one-dimensional system of bosons with repulsive delta function interaction*, *J. Math. Phys.* **10** (1969) 1115 [[INSPIRE](#)].
- [3] A.B. Zamolodchikov, *Thermodynamic Bethe Ansatz in Relativistic Models. Scaling Three State Potts and Lee-yang Models*, *Nucl. Phys. B* **342** (1990) 695 [[INSPIRE](#)].
- [4] M. Karowski and P. Weisz, *Exact Form-Factors in (1+1)-Dimensional Field Theoretic Models with Soliton Behavior*, *Nucl. Phys. B* **139** (1978) 455 [[INSPIRE](#)].
- [5] A.N. Kirillov and F.A. Smirnov, *A Representation of the Current Algebra Connected With the SU(2) Invariant Thirring Model*, *Phys. Lett. B* **198** (1987) 506 [[INSPIRE](#)].
- [6] F. Smirnov, *Form-factors in completely integrable models of quantum field theory*, *Adv. Ser. Math. Phys.* **14** (1992) 1.
- [7] D. Fioretto and G. Mussardo, *Quantum Quenches in Integrable Field Theories*, *New J. Phys.* **12** (2010) 055015 [[arXiv:0911.3345](#)] [[INSPIRE](#)].
- [8] B. Pozsgay, *Mean values of local operators in highly excited Bethe states*, *J. Stat. Mech.* (2011) P01011 [[arXiv:1009.4662](#)] [[INSPIRE](#)].
- [9] M. Kormos, G. Mussardo and A. Trombettoni, *Expectation Values in the Lieb-Liniger Bose Gas*, *Phys. Rev. Lett.* **103** (2009) 210404 [[arXiv:0909.1336](#)] [[INSPIRE](#)].
- [10] M. Kormos, G. Mussardo and A. Trombettoni, *Local Correlations in the Super Tonks-Girardeau Gas*, *Phys. Rev. A* **83** (2011) 013617 [[arXiv:1008.4383](#)] [[INSPIRE](#)].
- [11] B. Pozsgay, *Local correlations in the 1D Bose gas from a scaling limit of the XXZ chain*, *J. Stat. Mech.* (2011) P11017 [[arXiv:1108.6224](#)] [[INSPIRE](#)].
- [12] B. Doyon, *Finite-temperature form-factors in the free Majorana theory*, *J. Stat. Mech.* (2005) P11006 [[hep-th/0506105](#)] [[INSPIRE](#)].
- [13] Y. Chen and B. Doyon, *Form factors in equilibrium and non-equilibrium mixed states of the Ising model*, *J. Stat. Mech.* (2014) P09021 [[arXiv:1305.0518](#)] [[INSPIRE](#)].
- [14] S.L. Lukyanov, *Finite temperature expectation values of local fields in the sinh-Gordon model*, *Nucl. Phys. B* **612** (2001) 391 [[hep-th/0005027](#)] [[INSPIRE](#)].
- [15] N. Grosjean, J.M. Maillet and G. Niccoli, *On the form factors of local operators in the lattice sine-Gordon model*, *J. Stat. Mech.* (2012) P10006 [[arXiv:1204.6307](#)] [[INSPIRE](#)].
- [16] M. Jimbo, T. Miwa and F. Smirnov, *Hidden Grassmann structure in the XXZ model V: sine-Gordon model*, *Lett. Math. Phys.* **96** (2011) 325 [[arXiv:1007.0556](#)] [[INSPIRE](#)].
- [17] S. Negro and F. Smirnov, *On one-point functions for sinh-Gordon model at finite temperature*, *Nucl. Phys. B* **875** (2013) 166 [[arXiv:1306.1476](#)] [[INSPIRE](#)].
- [18] S. Negro, *On sinh-Gordon Thermodynamic Bethe Ansatz and fermionic basis*, *Int. J. Mod. Phys. A* **29** (2014) 1450111 [[arXiv:1404.0619](#)] [[INSPIRE](#)].

- [19] B. Pozsgay and G. Takács, *Form-factors in finite volume I: Form-factor bootstrap and truncated conformal space*, *Nucl. Phys. B* **788** (2008) 167 [[arXiv:0706.1445](#)] [[INSPIRE](#)].
- [20] B. Pozsgay and G. Takács, *Form factors in finite volume. II. Disconnected terms and finite temperature correlators*, *Nucl. Phys. B* **788** (2008) 209 [[arXiv:0706.3605](#)] [[INSPIRE](#)].
- [21] F.H.L. Essler and R.M. Konik, *Finite-temperature lineshapes in gapped quantum spin chains*, *Phys. Rev. B* **78** (2008) 100403 [[arXiv:0711.2524](#)] [[INSPIRE](#)].
- [22] F.H.L. Essler and R.M. Konik, *Finite-temperature dynamical correlations in massive integrable quantum field theories*, *J. Stat. Mech.* (2009) P09018 [[INSPIRE](#)].
- [23] B. Pozsgay and G. Takács, *Form factor expansion for thermal correlators*, *J. Stat. Mech.* (2010) P11012 [[arXiv:1008.3810](#)] [[INSPIRE](#)].
- [24] I.M. Szecsenyi and G. Takács, *Spectral expansion for finite temperature two-point functions and clustering*, *J. Stat. Mech.* (2012) P12002 [[arXiv:1210.0331](#)] [[INSPIRE](#)].
- [25] D.A. Tennant et al., *Anomalous dynamical line shapes in a quantum magnet at finite temperature*, *Phys. Rev. B* **85** (2012) 014402.
- [26] J. Wu, M. Kormos and Q. Si, *Finite temperature spin dynamics in a perturbed quantum critical Ising chain with an E_8 symmetry* *Phys. Rev. Lett.* **113** (2014) 247201 [[arXiv:1403.7222](#)].
- [27] M. Kormos and B. Pozsgay, *One-Point Functions in Massive Integrable QFT with Boundaries*, *JHEP* **04** (2010) 112 [[arXiv:1002.2783](#)] [[INSPIRE](#)].
- [28] G. Mussardo, *Infinite-time Average of Local Fields in an Integrable Quantum Field Theory after a Quantum Quench*, *Phys. Rev. Lett.* **111** (2013) 100401 [[arXiv:1308.4551](#)] [[INSPIRE](#)].
- [29] S. Sotiriadis, G. Takács and G. Mussardo, *Boundary State in an Integrable Quantum Field Theory Out of Equilibrium*, *Phys. Lett. B* **734** (2014) 52 [[arXiv:1311.4418](#)] [[INSPIRE](#)].
- [30] B. Bertini, D. Schuricht and F.H.L. Essler, *Quantum quench in the sine-Gordon model*, *J. Stat. Mech.* (2014) P10035 [[arXiv:1405.4813](#)] [[INSPIRE](#)].
- [31] T. Klose and T. McLoughlin, *Comments on World-Sheet Form Factors in AdS/CFT*, *J. Phys. A* **47** (2014) 055401 [[arXiv:1307.3506](#)] [[INSPIRE](#)].
- [32] Z. Bajnok, R.A. Janik and A. Wereszczynski, *HHL correlators, orbit averaging and form factors*, *JHEP* **09** (2014) 050 [[arXiv:1404.4556](#)] [[INSPIRE](#)].
- [33] Z. Bajnok and R.A. Janik, *Four-loop perturbative Konishi from strings and finite size effects for multiparticle states*, *Nucl. Phys. B* **807** (2009) 625 [[arXiv:0807.0399](#)] [[INSPIRE](#)].
- [34] Z. Bajnok, F. Buccheri, L. Hollo, J. Konczer and G. Takács, *Finite volume form factors in the presence of integrable defects*, *Nucl. Phys. B* **882** (2014) 501 [[arXiv:1312.5576](#)] [[INSPIRE](#)].
- [35] B. Pozsgay, *Lüscher's μ -term and finite volume bootstrap principle for scattering states and form factors*, *Nucl. Phys. B* **802** (2008) 435 [[arXiv:0803.4445](#)] [[INSPIRE](#)].
- [36] G. Takács, *Determining matrix elements and resonance widths from finite volume: The dangerous μ -terms*, *JHEP* **11** (2011) 113 [[arXiv:1110.2181](#)] [[INSPIRE](#)].
- [37] A.I. Bugrij, *The correlation function in two-dimensional Ising model on the finite size lattice. 1.*, *Theor. Math. Phys.* **127** (2001) 528 [[hep-th/0011104](#)] [[INSPIRE](#)].

- [38] A.I. Bugrij, *Form-factor representation of the correlation function of the two-dimensional Ising model on a cylinder*, [hep-th/0107117](#) [INSPIRE].
- [39] B. Pozsgay, *Form factor approach to diagonal finite volume matrix elements in Integrable QFT*, *JHEP* **07** (2013) 157 [[arXiv:1305.3373](#)] [INSPIRE].
- [40] J. Teschner, *On the spectrum of the Sinh-Gordon model in finite volume*, *Nucl. Phys. B* **799** (2008) 403 [[hep-th/0702214](#)] [INSPIRE].
- [41] V.V. Bazhanov, S.L. Lukyanov and A.B. Zamolodchikov, *Integrable quantum field theories in finite volume: Excited state energies*, *Nucl. Phys. B* **489** (1997) 487 [[hep-th/9607099](#)] [INSPIRE].
- [42] P. Dorey and R. Tateo, *Excited states by analytic continuation of TBA equations*, *Nucl. Phys. B* **482** (1996) 639 [[hep-th/9607167](#)] [INSPIRE].
- [43] P. Dorey and R. Tateo, *Excited states in some simple perturbed conformal field theories*, *Nucl. Phys. B* **515** (1998) 575 [[hep-th/9706140](#)] [INSPIRE].
- [44] V.P. Yurov and A.B. Zamolodchikov, *Truncated conformal space approach to scaling Lee-Yang model*, *Int. J. Mod. Phys. A* **5** (1990) 3221 [INSPIRE].
- [45] I.M. Szécsényi, G. Takács and G.M.T. Watts, *One-point functions in finite volume/temperature: a case study*, *JHEP* **08** (2013) 094 [[arXiv:1304.3275](#)] [INSPIRE].
- [46] A.B. Zamolodchikov, *On the thermodynamic Bethe ansatz equations for reflectionless ADE scattering theories*, *Phys. Lett. B* **253** (1991) 391 [INSPIRE].
- [47] F. Ravanini, R. Tateo and A. Valleriani, *Dynkin TBAs*, *Int. J. Mod. Phys. A* **8** (1993) 1707 [[hep-th/9207040](#)] [INSPIRE].
- [48] J.S. Maybee, D.D. Olesky, P. van den Driessche and G. Wiener, *Matrices, digraphs, and determinants*, *SIAM J. Matrix Anal. Appl.* **10** (1989) 500.
- [49] A.A. Belavin, A.M. Polyakov and A.B. Zamolodchikov, *Infinite Conformal Symmetry in Two-Dimensional Quantum Field Theory*, *Nucl. Phys. B* **241** (1984) 333 [INSPIRE].
- [50] R.M. Konik and Y. Adamov, *Numerical Renormalization Group for Continuum One-Dimensional Systems*, *Phys. Rev. Lett.* **98** (2007) 147205.
- [51] R.M. Konik and Y. Adamov, *A Numerical Renormalization Group for Continuum One-Dimensional Systems*, *J. Stat. Mech.* (2008) P03011 [[cond-mat/0701605](#)] [INSPIRE].
- [52] P. Giokas and G. Watts, *The renormalisation group for the truncated conformal space approach on the cylinder*, [arXiv:1106.2448](#) [INSPIRE].
- [53] G.M.T. Watts, *On the renormalisation group for the boundary Truncated Conformal Space Approach*, *Nucl. Phys. B* **859** (2012) 177 [[arXiv:1104.0225](#)] [INSPIRE].
- [54] M. Lencsés and G. Takács, *Excited state TBA and renormalized TCSA in the scaling Potts model*, *JHEP* **09** (2014) 052 [[arXiv:1405.3157](#)] [INSPIRE].
- [55] M. Beria, G.P. Brandino, L. Lepori, R.M. Konik and G. Sierra, *Truncated Conformal Space Approach for Perturbed Wess-Zumino-Witten $SU_{(2)_k}$ Models*, *Nucl. Phys. B* **877** (2013) 457 [[arXiv:1301.0084](#)] [INSPIRE].
- [56] A. Coser, M. Beria, G.P. Brandino, R.M. Konik and G. Mussardo, *Truncated Conformal Space Approach for 2D Landau-Ginzburg Theories*, *J. Stat. Mech.* (2014) P12010 [[arXiv:1409.1494](#)] [INSPIRE].

- [57] M. Hogervorst, S. Rychkov and B.C. van Rees, *Truncated conformal space approach in d dimensions: A cheap alternative to lattice field theory?*, *Phys. Rev. D* **91** (2015) 025005 [[arXiv:1409.1581](#)] [[INSPIRE](#)].
- [58] S. Rychkov and L.G. Vitale, *Hamiltonian Truncation Study of the Φ^4 Theory in Two Dimensions*, [arXiv:1412.3460](#) [[INSPIRE](#)].
- [59] T.R. Klassen and E. Melzer, *On the relation between scattering amplitudes and finite size mass corrections in QFT*, *Nucl. Phys. B* **362** (1991) 329 [[INSPIRE](#)].
- [60] M. Lüscher, *Volume Dependence of the Energy Spectrum in Massive Quantum Field Theories. 1. Stable Particle States*, *Commun. Math. Phys.* **104** (1986) 177 [[INSPIRE](#)].
- [61] F. Buccheri and G. Takács, *Finite temperature one-point functions in non-diagonal integrable field theories: the sine-Gordon model*, *JHEP* **03** (2014) 026 [[arXiv:1312.2623](#)] [[INSPIRE](#)].
- [62] V.A. Fateev, *The exact relations between the coupling constants and the masses of particles for the integrable perturbed conformal field theories*, *Phys. Lett. B* **324** (1994) 45 [[INSPIRE](#)].
- [63] T. Hahn, *CUBA: A library for multidimensional numerical integration*, *Comput. Phys. Commun.* **168** (2005) 78 [[hep-ph/0404043](#)] [[INSPIRE](#)].
- [64] *Mathematica*, Wolfram Research, Inc., Champaign, Illinois, version 10.0 ed., (2014).
- [65] G.Z. Feher, T. Palmai and G. Takács, *sine-Gordon multi-soliton form factors in finite volume*, *Phys. Rev. D* **85** (2012) 085005 [[arXiv:1112.6322](#)] [[INSPIRE](#)].
- [66] T. Pálmai and G. Takács, *Diagonal multisoliton matrix elements in finite volume*, *Phys. Rev. D* **87** (2013) 045010 [[arXiv:1209.6034](#)] [[INSPIRE](#)].
- [67] A. Koubek, *The space of local operators in perturbed conformal field theories*, *Nucl. Phys. B* **435** (1995) 703 [[hep-th/9501029](#)] [[INSPIRE](#)].
- [68] A. Koubek and G. Mussardo, *On the operator content of the sinh-Gordon model*, *Phys. Lett. B* **311** (1993) 193 [[hep-th/9306044](#)] [[INSPIRE](#)].
- [69] B. Pozsgay and G. Takács, *Characterization of resonances using finite size effects*, *Nucl. Phys. B* **748** (2006) 485 [[hep-th/0604022](#)] [[INSPIRE](#)].
- [70] V. Fateev, S.L. Lukyanov, A.B. Zamolodchikov and A.B. Zamolodchikov, *Expectation values of local fields in Bullough-Dodd model and integrable perturbed conformal field theories*, *Nucl. Phys. B* **516** (1998) 652 [[hep-th/9709034](#)] [[INSPIRE](#)].
- [71] A. Fring, G. Mussardo and P. Simonetti, *Form-factors for integrable Lagrangian field theories, the sinh-Gordon theory*, *Nucl. Phys. B* **393** (1993) 413 [[hep-th/9211053](#)] [[INSPIRE](#)].