

Neutrino mass and mixing models with eclectic flavor symmetry $\Delta(27) \times T'$

Gui-Jun Ding,^a Stephen F. King,^b Cai-Chang Li,^{c,d,e} Xiang-Gan Liu^f and Jun-Nan Lu^a

^a*Department of Modern Physics, University of Science and Technology of China, Hefei, Anhui 230026, China*

^b*Physics and Astronomy, University of Southampton, Southampton, SO17 1BJ, U.K.*

^c*School of Physics, Northwest University, Xi'an 710127, China*

^d*Shaanxi Key Laboratory for Theoretical Physics Frontiers, Xi'an 710127, China*

^e*NSFC-SPTP Peng Huanwu Center for Fundamental Theory, Xi'an 710127, China*

^f*Department of Physics and Astronomy, University of California, Irvine, CA 92697-4575, U.S.A.*

E-mail: dinggj@ustc.edu.cn, king@soton.ac.uk, ccli@nwu.edu.cn, xianggal@uci.edu, hitman@mail.ustc.edu.cn

ABSTRACT: The Kähler potentials of modular symmetry models receive unsuppressed contributions which may be controlled by a flavor symmetry, where the combination of the two symmetry types is referred to as eclectic flavor symmetry. After briefly reviewing the consistency conditions of eclectic flavor symmetry models, including those with generalised (g)CP, we perform a comprehensive bottom-up study of eclectic flavor symmetry models based on $\Omega(1) \cong \Delta(27) \times T'$, consisting of the flavor symmetry $\Delta(27)$ in a semi-direct product with the modular symmetry T' . The modular transformations of different $\Delta(27)$ multiplets are given by solving the consistency condition. The eight nontrivial singlets of $\Delta(27)$ are related by T' modular symmetry, and they have to be present or absent simultaneously in any $\Omega(1)$ model. The most general forms of the superpotential and Kähler potential invariant under $\Omega(1)$ are discussed, and the corresponding fermion mass matrices are presented. Based on the eclectic flavor group $\Omega(1)$, two concrete lepton models which can successfully describe the experimental data of lepton masses and mixing parameters are constructed. For the two models without gCP, all six mixing parameters vary in small regions. A nearly maximal atmospheric mixing angle θ_{23} and Dirac CP phase δ_{CP} are obtained in the first model. After considering the compatible gCP symmetry and the

assumption of $\Re\tau = 0$ in the first model, the $\mu - \tau$ reflection symmetry is preserved in the charged lepton diagonal basis. As a consequence, the atmospheric mixing angle and Dirac CP phase are predicted to be maximal, and two Majorana CP phases are predicted to be π .

KEYWORDS: CP Violation, Discrete Symmetries, Flavour Symmetries, Neutrino Mixing

ARXIV EPRINT: [2303.02071](https://arxiv.org/abs/2303.02071)

Contents

1	Introduction	1
2	Formalism of eclectic flavor group	3
2.1	Eclectic flavor group and gCP	5
3	The eclectic flavor group $\Omega(1) \cong \Delta(27) \rtimes T'$	7
3.1	Finite modular group $\Gamma'_3 \cong T'$ and gCP	8
3.2	T' modular transformations of $\Delta(27)$ multiplets	8
4	Effective action invariant under $\Omega(1)$ eclectic flavor symmetry	14
4.1	Superpotential	14
4.1.1	The superpotential of Majorana mass terms	17
4.2	Kähler potential	19
5	$\Omega(1) \cong \Delta(27) \rtimes T'$ eclectic models of lepton mass and mixing	24
5.1	Example model 1	26
5.2	Example model 2	32
6	Conclusion and outlook	34
A	Traditional flavor group $\Delta(27)$	36
B	The finite modular group $\Gamma'_3 \cong T'$	39
C	Modular forms of level $N = 3$	41
D	Alternative method to derive the Majorana superpotential	43

1 Introduction

The modular flavor symmetry is a promising approach to address the flavor puzzle of the standard model (SM) [1]. The superpotential is strongly constrained by the modular symmetry and all higher-dimensional operators are unambiguously fixed in the limit of unbroken supersymmetry. In particular, the modular symmetry enforces the Yukawa couplings and fermion mass matrices to be modular forms which are holomorphic functions of the complex modulus τ . The modular flavor symmetry has been extensively studied from the bottom-up approach, see the recent review [2] and references therein for more details. The modular flavor symmetry allows one to construct quite predictive fermion mass models, all the lepton masses and mixing parameters can be described only in terms of six real free

parameters in the known minimal model [3]. It is remarkable that the modular invariant models exhibit a universal behavior around the fixed points, and the scaling of fermion masses and mixing parameters is independent of the details of the theory [4, 5]. However, the Kähler potential is less constrained by modular symmetry, the most general Kähler potential compatible with modular symmetry is of the following form [6, 7],

$$\mathcal{K} = (-i\tau + i\bar{\tau})^{-k_\psi} \left(\psi^\dagger \psi \right)_1 + \sum_{n, \mathbf{r}_1, \mathbf{r}_2} c^{(n, \mathbf{r}_1, \mathbf{r}_2)} (-i\tau + i\bar{\tau})^{-k_\psi + n} \left(\psi^\dagger Y_{\mathbf{r}_1}^{(n)\dagger} Y_{\mathbf{r}_2}^{(n)} \psi \right)_1, \quad (1.1)$$

where ψ stands for a generic matter field multiple with modular weight k_ψ , and $Y_{\mathbf{r}}^{(n)}$ is a weight $n \neq 0$ modular form in the representation \mathbf{r} of the finite modular group. Usually only the first term of eq. (1.1) is taken, this is the so-called minimal Kähler potential. However, we see that the Kähler potential generally has a lot of terms and the couplings $c^{(n, \mathbf{r}_1, \mathbf{r}_2)}$ are not suppressed. Consequently these additional terms can be as important as the first term, and thus the predictive power of modular flavor symmetry would be reduced [6].

On the other hand, the modular symmetry has been studied in the top-down approach motivated by string theory, and models with modular symmetry have been constructed in heterotic orbifolds [8, 9] and magnetized toroidal compactifications [10–16]. It is found that the modular symmetry and traditional flavor symmetry appear together in top-down constructions. This leads to the concept of eclectic flavor group [17, 18], and it has been developed in a series of papers [8, 9, 17–20]. The eclectic flavor group is a maximal extension of the traditional flavor group by finite modular group, and certain consistency conditions have to be fulfilled in order to consistently combine modular symmetry with traditional flavor symmetry. The scheme of eclectic flavor group is more predictive than the finite modular group and the traditional flavor group alone, and it combines the advantages of both approaches, in which the superpotential and the Kähler potential would be severely restricted. In particular, the Kähler potential would be constrained to be the minimal form plus higher order corrections suppressed by powers of $\langle \Phi \rangle / \Lambda$, where Λ denotes the cutoff scale and $\langle \Phi \rangle$ represents the vacuum expectation value (VEV) of flavons breaking the traditional flavor symmetry. It is known that Kähler potential is suppressed by powers of $\langle \Phi \rangle / \Lambda$ in the traditional flavor symmetry models [21–25]. Flavon fields are necessary in eclectic flavor models to break the traditional flavor symmetry, otherwise the unbroken flavor group would constrain the fermion mixing matrix to be the unit matrix up to possible row and column permutations. The eclectic flavor group considered here is a nontrivial product of the modular symmetry and traditional flavor symmetry. The choices of such eclectic symmetries are quite restrictive, with only three choices of flavor group $Z_3 \times Z_3$, $\Delta(27)$ and $\Delta(54)$ catalogued so far allowing for a non-trivial eclectic extension with a CP-like transformation [17, 18]. This is unlike the quasi-eclectic flavor symmetry which is the direct product of a finite modular symmetry and a traditional flavor symmetry [26], where the modular and traditional flavor groups can be freely chosen, but the Kähler potential is also under control due to the presence of traditional flavor symmetry. Moreover, the first string-derived model based on the eclectic flavor group $\Omega(2) \cong [1944, 3448]$ consisting of the traditional flavor group $\Delta(54)$, the finite modular group T' and a \mathbb{Z}_9^R R -symmetry, has been constructed, and it can give a successful fit of flavor in both the quark and lepton sectors [27].

The eclectic flavor group opens a new approach to understand the flavor puzzle of SM. In the present work, we shall study the eclectic flavor group $\Omega(1) \cong [648, 533]$ in a bottom-up way, here we adopt the naming scheme of GAP [28] and the group order of $\Omega(1)$ is 648. The eclectic group $\Omega(1)$ is the semi-direct product of the traditional flavor group $\Delta(27)$ with the finite modular group $\Gamma'_3 \cong T'$. If the generalized CP (gCP) symmetry is included, the eclectic flavor group $\Omega(1)$ would be enlarged to $[1296, 2891]$ [17]. The finite modular group should be a subgroup of the automorphism group of the traditional flavor group in the scheme of eclectic flavor symmetry. The outer automorphism group of $\Delta(27)$ is $GL(2, 3)$ which really contains the finite modular group T' as a subgroup. We find that the eight nontrivial singlet representations of $\Delta(27)$ transform into each other under T' modular symmetry. Hence there must be seven other fields in the remaining nontrivial singlet representations of $\Delta(27)$, if a $\Omega(1)$ eclectic flavor model contains a field which is a nontrivial singlet of $\Delta(27)$. Consequently the $\Omega(1)$ eclectic flavor symmetry is more restrictive than $\Delta(27)$ flavor symmetry alone and terms allowed by the modular group may be forbidden by the traditional flavor group and vice versa. We shall give the most general form of the superpotential and Kähler potential invariant under $\Omega(1)$. The deviations from the canonical kinetic term are found to be suppressed by the VEVs of flavons. Furthermore, we shall construct two concrete lepton models with $\Omega(1)$ eclectic flavor symmetry. The field content, the modular weights and representations of the fields are assigned at will, while they are derived from the underlying geometry in top-down models. Since there are three generations of quarks and leptons, the matter fields can be assigned to triplets/anti-triplets or trivial singlets of $\Delta(27)$, but nontrivial singlets would have to form a reducible octet, as mentioned above, leading to too many families.

The rest of this paper is organized as follows: we review the formalism of eclectic flavor group in section 2. The eclectic flavor group $\Omega(1)$ is discussed in section 3. The modular transformation and flavor symmetry transformation are closely related through the consistency condition in the eclectic flavor group, and the T' transformations of different $\Delta(27)$ multiplets are determined by solving the consistency condition. Moreover, the gCP transformation matrices are fixed. In section 4 we present the most general form of the $\Omega(1)$ invariant superpotential and Kähler potential for the fermion masses. Two concrete example models are presented in section 5, in which the light neutrino masses are generated by the Weinberg operator and type I seesaw mechanism, respectively. We summarize our results and give the conclusion in section 6. The appendix A and appendix B are for the group theory of $\Delta(27)$ and T' respectively, and we present the modular forms of level 3 in appendix C. The superpotential for Majorana fermion mass is derived in an alternative method by firstly considering T' modular symmetry, then $\Delta(27)$ flavor symmetry in appendix D.

2 Formalism of eclectic flavor group

We shall recapitulate the scheme of eclectic flavor group in this section. The modular group $SL(2, \mathbb{Z})$ denoted as Γ is the group of two dimensional matrices with integer entries and

unit determinant,

$$\mathrm{SL}(2, \mathbb{Z}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \middle| ad - bc = 1, a, b, c, d \in \mathbb{Z} \right\}. \quad (2.1)$$

The modular group is an infinite discrete group generated by the two elements S and T with

$$S = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad (2.2)$$

which satisfy the following relations

$$S^4 = (ST)^3 = \mathbb{1}_2, \quad S^2 T = T S^2, \quad (2.3)$$

where $\mathbb{1}_2$ denotes the 2-by-2 unit matrix. Under the action of a modular transformation, the complex modulus τ and a generic matter field multiplet ψ transform as follow [1]

$$\tau \xrightarrow{\gamma} \gamma\tau \equiv \frac{a\tau + b}{c\tau + d}, \quad \psi \xrightarrow{\gamma} (c\tau + d)^{-k_\psi} \rho(\gamma)\psi, \quad \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}(2, \mathbb{Z}), \quad (2.4)$$

where k_ψ is the modular weight of ψ and ρ is a unitary representation of the finite modular group $\Gamma_N \equiv \Gamma / \pm \Gamma(N)$ or its double covering $\Gamma'_N \equiv \Gamma / \Gamma(N)$, and $\Gamma(N)$ is the so-called principal congruence subgroup of level N . For $N \leq 5$, the defining relations of Γ_N and Γ'_N are given by [1, 29]

$$\begin{aligned} \Gamma_N &= \left\{ S, T \middle| S^2 = (ST)^3 = T^N = 1 \right\}, \\ \Gamma'_N &= \left\{ S, T \middle| S^4 = (ST)^3 = T^N = 1, S^2 T = T S^2 \right\}. \end{aligned} \quad (2.5)$$

Additional relations are needed to render the group finite for $N \geq 6$ [30]. Notice that the representation matrices $\rho(S)$ and $\rho(T)$ should also satisfy the defining relations of Γ_N and Γ'_N in eq. (2.5). Modular invariance constrains the Yukawa couplings to be modular forms which are holomorphic functions of τ . The level N and integer (even) weight k_Y modular forms $Y^{(k_Y)}(\tau)$ can be arranged into multiplets of $\Gamma'_N(\Gamma_N)$ [1, 29]:

$$Y^{(k_Y)}(\tau) \xrightarrow{\gamma} Y^{(k_Y)}(\gamma\tau) = (c\tau + d)^{k_Y} \rho_Y(\gamma) Y^{(k_Y)}(\tau), \quad (2.6)$$

where $\rho_Y(\gamma)$ is a unitary representation of $\Gamma'_N(\Gamma_N)$.

The eclectic flavor group is a nontrivial product of modular group and the traditional flavor group denoted as G_f . The modulus τ as well as modular form $Y^{(k_Y)}(\tau)$ is invariant under the flavor group G_f , and the flavor transformation of the matter multiplet ψ is

$$\psi \xrightarrow{g} \rho(g)\psi, \quad g \in G_f, \quad (2.7)$$

where $\rho(g)$ is a unitary representation of G_f . In order to consistently combine a finite modular group with a traditional flavor group, certainly consistency condition has to be fulfilled [17]. Let us first perform a modular transformation $\gamma \in \Gamma$, subsequently a traditional flavor transformation g and last perform the inverse modular transformation γ^{-1} . Since

the modulus τ is invariant under this chain of transformations, the resulting transformation should be another traditional flavor symmetry transformation g' [17], i.e.

$$\rho(\gamma)\rho(g)\rho^{-1}(\gamma) = \rho(g'), \quad g, g' \in G_f, \quad \gamma \in \Gamma. \quad (2.8)$$

This is the consistency condition between traditional flavor group and finite modular group. Eq. (2.8) implies that the modular transformation γ maps the traditional group element g into another element g' . Therefore eq. (2.8) defines a homomorphism of the traditional flavor group G_f . Furthermore, the modular transformation is bijective. Hence modular transformation is an automorphism of the traditional group G_f . Indeed the possible representation matrices of the finite modular group Γ'_N form a representation of the automorphism group $\text{Aut}(G_f)$ of the traditional flavor group G_f , i.e.

$$\rho(\gamma)\rho(g)\rho^{-1}(\gamma) = \rho(u_\gamma(g)), \quad \forall g \in G_f, \quad (2.9)$$

where $\rho(\gamma)$ represents the automorphism $u_\gamma : G_f \rightarrow G_f$. If u_γ is the trivial identity automorphism with $u_\gamma = 1$ for any modular transformation γ so that $u_\gamma(g) = g$, the modular symmetry transformation would commute with the flavor symmetry transformation and consequently the eclectic flavor group is the direct product $G_f \times \Gamma_N$ or $G_f \times \Gamma'_N$ [17]. Hence one can freely choose both Γ_N , Γ'_N and G_f , the resulting flavor model would be more complex than the traditional flavor symmetry models and modular models. If u_γ is an inner automorphism of G_f , the modular symmetry transformation $\rho(\gamma)$ would coincide with certain flavor symmetry transformation besides the automorphy factor $(c\tau + d)^{-k_\psi}$. Hence u_γ is required to be outer automorphism of G_f in the paradigm of eclectic flavor group [17]. As a consequence, the mathematical structure of the group comprising the traditional flavor symmetry G_f and the finite modular group Γ'_N (Γ_N) is in general a semi-direct product $G_f \rtimes \Gamma'_N$ ($G_f \rtimes \Gamma_N$). Since the finite modular groups Γ_N and Γ'_N can be generated by the two generators S and T , it is sufficient to only discuss the two outer automorphisms u_S and u_T and the consistency condition in eq. (2.9) reduces to

$$\rho(S)\rho(g)\rho^{-1}(S) = \rho(u_S(g)), \quad \rho(T)\rho(g)\rho^{-1}(T) = \rho(u_T(g)). \quad (2.10)$$

Since $\rho(S)$ and $\rho(T)$ are representation matrices of the finite modular group, the outer automorphisms u_S and u_T should also satisfy the multiplication rules eq. (2.5) of the finite modular group Γ'_N (Γ_N):

$$(u_S)^{N_s} = (u_T)^N = (u_S \circ u_T)^3 = 1, \quad (u_S)^2 \circ u_T = u_T \circ (u_S)^2, \quad (2.11)$$

with $N_s = 4$ for Γ'_N and $N_s = 2$ for Γ_N , additional relations are necessary for level $N \geq 6$. Therefore the finite modular group Γ'_N (Γ_N) must be a subgroup of the full automorphism group of the traditional flavor group G_f .

2.1 Eclectic flavor group and gCP

The modular symmetry group Γ can be consistently combined with the gCP symmetry. Then a new generator K_* corresponding to gCP transformation could be introduced and the modular group $\Gamma \cong \text{SL}(2, \mathbb{Z})$ is enhanced to $\Gamma^* \cong \text{GL}(2, \mathbb{Z})$ [31]:

$$\Gamma^* = \left\{ \tau \xrightarrow{S} -1/\tau, \quad \tau \xrightarrow{T} \tau + 1, \quad \tau \xrightarrow{K_*} -\bar{\tau} \right\}, \quad (2.12)$$

where the action of the CP transformation K_* on the modulus τ is represented by the matrix

$$K_* = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (2.13)$$

As a consequence, the action of Γ^* on the complex modulus τ is given by

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma^* : \begin{cases} \tau \rightarrow \frac{a\tau + b}{c\tau + d} & \text{for } ad - bc = 1, \\ \tau \rightarrow \frac{a\bar{\tau} + b}{c\bar{\tau} + d} & \text{for } ad - bc = -1. \end{cases} \quad (2.14)$$

On the other hand, the CP transformation acts on the matter field and the modular form multiplets as

$$\psi(x) \xrightarrow{K_*} \rho(K_*)[\psi^\dagger(t, -\mathbf{x})]^T, \quad Y(\tau) \xrightarrow{K_*} Y(-\bar{\tau}) = \rho(K_*)Y^*(\tau), \quad (2.15)$$

where the unitary matrix $\rho(K_*)$ is the gCP transformation matrix of a generic matter field ψ . Note that the obvious action of CP on the possible spinor indices has been suppressed in eq. (2.15). As the CP transformation K_* is order 2, then one can find

$$\rho(K_*)\rho^*(K_*) = \mathbb{1} \quad \text{or equivalently} \quad \rho^*(K_*) = \rho^{-1}(K_*). \quad (2.16)$$

The consistency between the modular symmetry and gCP symmetry requires the following consistency condition has to be satisfied [31, 32]

$$\rho(K_*)\rho^*(\gamma)\rho^{-1}(K_*) = \rho(\mathbf{u}(\gamma)), \quad \gamma \in \Gamma, \quad (2.17)$$

where $\mathbf{u}(\gamma)$ denotes an outer automorphism of the modular group with

$$\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \rightarrow \mathbf{u}(\gamma) = K_*\gamma K_*^{-1} = \begin{pmatrix} a & -b \\ -c & d \end{pmatrix}. \quad (2.18)$$

It is easy to check that $\mathbf{u}(S) = S^{-1}$ and $\mathbf{u}(T) = T^{-1}$. It is sufficient to impose the consistency condition in eq. (2.17) on the generators S and T of the modular group Γ , then one can obtain [31–33]

$$\rho(K_*)\rho^*(S)\rho^{-1}(K_*) = \rho^{-1}(S), \quad \rho(K_*)\rho^*(T)\rho^{-1}(K_*) = \rho^{-1}(T). \quad (2.19)$$

If one works in the basis where both representation matrices $\rho(S)$ and $\rho(T)$ are unitary and symmetric, $\rho(K_*)$ would reduce to the canonical CP transformation,

$$\rho(K_*) = \mathbb{1}, \quad (2.20)$$

where an arbitrary overall phase is omitted.

Furthermore, it is known that the following consistency condition has to be satisfied when combining the traditional flavor G_f with the gCP symmetry [34–38]:

$$\rho(K_*)\rho^*(g)\rho^{-1}(K_*) = \rho(u_{K_*}(g)), \quad \forall g \in G_f, \quad (2.21)$$

where u_{K_*} is an automorphism of the traditional flavor group G_f . Furthermore, the CP transformation $\rho(K_*)$ with order 2 must be subject to the consistency condition in eq. (2.19). Consequently, the automorphisms u_{K_*} , u_S and u_T of the traditional flavor group G_f should satisfy the following multiplication rules

$$\begin{aligned} (u_S)^{N_s} = (u_T)^N = (u_S \circ u_T)^3 = 1, & \quad (u_S)^2 \circ u_T = u_T \circ (u_S)^2, \\ (u_{K_*})^2 = 1, & \quad u_{K_*} \circ u_S \circ u_{K_*} = u_S^{-1}, \quad u_{K_*} \circ u_T \circ u_{K_*} = u_T^{-1}, \end{aligned} \quad (2.22)$$

for $N \leq 5$ and additional multiplication rules should be imposed for $N > 6$.

From the general formalism sketched above, one could find all possible corresponding eclectic flavor groups and gCP for a given traditional flavor group G_f . It is necessary to find out all outer automorphisms of the traditional flavor group G_f and determine whether there exist particular outer automorphisms u_S , u_T and u_{K_*} satisfying the multiplication rules in eq. (2.22). For a field multiplet transforming in the representation $\rho(g)$ of the traditional flavor group G_f , one cannot freely assign the modular transformation of the fields as in modular symmetry alone. Nevertheless, the modular transformation matrices $\rho(S)$, $\rho(T)$ and the gCP transformation $\rho(K_*)$ should be determined by solving the consistency conditions in eqs. (2.10), (2.19) and (2.21). In general, the representation matrices $\rho(S)$, $\rho(T)$ span a reducible representation of the finite modular group Γ'_N (Γ_N). In the following, we shall study the eclectic flavor group $\Omega(1)$ which is semi-direct product of the traditional flavor group $\Delta(27)$ with the finite modular group T' .

3 The eclectic flavor group $\Omega(1) \cong \Delta(27) \rtimes T'$

In the present work, we shall consider the eclectic flavor group $\Omega(1)$ which is the extension of the traditional flavor symmetry $\Delta(27)$ by finite modular group T' . The mathematical structure of the group $\Omega(1)$ arising from the combination of the flavor symmetry $\Delta(27)$ and the T' modular symmetry is a semi-direct product of the form $\Omega(1) \cong \Delta(27) \rtimes T'$. The group theories of the traditional flavor group $\Delta(27)$ and the finite modular group T' are given in appendix A and appendix B, respectively. The finite modular group compatible with $\Delta(27)$ must be generated by the outer automorphisms of $\Delta(27)$. We find that the outer automorphism group of $\Delta(27)$ is $\text{GL}(2, 3)$:¹

$$\text{Out}(\Delta(27)) \cong \text{Aut}(\Delta(27)) / \text{Inn}(\Delta(27)) \cong \text{GL}(2, 3), \quad (3.1)$$

where $\text{Aut}(\Delta(27))$ and $\text{Inn}(\Delta(27))$ represent the automorphism group and inner automorphism group of $\Delta(27)$, respectively. Their group ID in GAP system are $\text{Aut}(\Delta(27)) \cong [432, 734]$ and $\text{Inn}(\Delta(27)) \cong Z_3 \times Z_3 \cong [9, 2]$. The group $\text{GL}(2, 3)$ contains two finite modular groups $\Gamma_2 \cong S_3$ and $\Gamma'_3 \cong T' \cong \text{SL}(2, 3)$ which can be generated by the outer automorphisms. This implies that the traditional flavor group $\Delta(27)$ can be extended to eclectic flavor group in two ways: $\Delta(27) \rtimes S_3$ or $\Delta(27) \rtimes T'$ [17]. In the present work, we are concerned with the traditional flavor group $\Delta(27)$ and its eclectic extension by $\Gamma'_3 \cong T'$, and the scenarios with/without gCP will be studied.

¹ $\text{GL}(2, 3)$ is the group of all 2×2 invertible matrices whose elements are from the three-member ring Z_3 .

3.1 Finite modular group $\Gamma'_3 \cong T'$ and gCP

The two outer automorphisms u_S and u_T can be taken to be [17]

$$u_S(A) = B^2 A, \quad u_S(B) = B^2 A^2, \quad u_T(A) = BA, \quad u_T(B) = B, \quad (3.2)$$

where A and B are the generators of $\Delta(27)$, see the appendix A for the group theory of $\Delta(27)$. One can check that the outer automorphisms u_S and u_T in eq. (3.2) satisfy the multiplication rules in eq. (2.11) with $N_s = 4$ and $N = 3$. As a consequence, the $\Delta(27)$ flavor symmetry could be extended by the finite modular group is $\Gamma'_3 \cong T'$ in the paradigm of eclectic flavor group. If gCP symmetry is considered further, the corresponding automorphism u_{K_*} should be included. The three automorphisms u_S , u_T and u_{K_*} should satisfy the following multiplication rules

$$\begin{aligned} (u_S)^4 &= (u_T)^3 = (u_S \circ u_T)^3 = 1, & (u_S)^2 \circ u_T &= u_T \circ (u_S)^2, \\ (u_{K_*})^2 &= 1, & u_{K_*} \circ u_S \circ u_{K_*} &= u_S^{-1}, & u_{K_*} \circ u_T \circ u_{K_*} &= u_T^{-1}, \end{aligned} \quad (3.3)$$

where the outer automorphism u_{K_*} acts on the generators A and B as follows

$$u_{K_*}(A) = A^2 B, \quad u_{K_*}(B) = A^2 B A. \quad (3.4)$$

Then the resulting eclectic flavor group is [1296, 2891], it has 1296 group elements. Moreover, the gCP transformation matrix $\rho_r(K_*)$ must satisfy eq. (2.16) and the consistency condition in eq. (2.19). In our basis given in eq. (B.4), the group generators S and T are represented by symmetric matrices for all irreducible representations of T' . As a consequence, the expressions of $\rho_r(K_*)$ are identity matrices for all irreducible representations up to an overall phase, i.e.

$$\rho_r(K_*) = \mathbb{1}_r, \quad r = \mathbf{1}, \mathbf{1}', \mathbf{1}'', \mathbf{2}, \mathbf{2}', \mathbf{2}'', \mathbf{3}. \quad (3.5)$$

The modular forms of level $N = 3$ up to weight 10 are given in appendix C. From the discussion of appendix C, we find that the CP transformation acts on the modular form multiplets $Y_r^{(k)}(\tau)$ as follow [31–33]

$$Y_r^{(k_Y)}(\tau) \xrightarrow{K_*} Y_r^{(k_Y)}(-\bar{\tau}) = \left(Y_r^{(k_Y)}(\tau) \right)^*, \quad (3.6)$$

where $Y_r^{(k_Y)}(\tau)$ represents the weight k_Y and level 3 modular multiplets in the irreducible representation r of the finite modular group T' .

3.2 T' modular transformations of $\Delta(27)$ multiplets

Automorphisms are mappings from a group into itself, and they can be divided into inner automorphisms and outer automorphisms. It is known that all inner automorphism maps each conjugacy class into itself, while certain conjugacy classes are mapped into different ones by the outer automorphisms. In order to determine the explicit form of the modular transformations $\rho(S)$, $\rho(T)$ and the gCP transformation $\rho(K_*)$ corresponding to the outer automorphisms u_S , u_T and u_{K_*} , one needs to know the actions of the three automorphisms u_S , u_T and u_{K_*} on the conjugacy classes and representations of $\Delta(27)$. It is necessary to

consider how these three automorphisms act on the character table of $\Delta(27)$, the results are summarized in table 6.

Firstly, we discuss how the three automorphisms u_S , u_T and u_{K^*} act on the eleven conjugacy classes of $\Delta(27)$. From eqs. (3.2) and (A.4), we see that the outer automorphism $u_S : (A, B) \rightarrow (B^2A, B^2A^2)$ acts on the all conjugacy classes elements of $\Delta(27)$ as

$$u_S : 3C_3^{(3)} \rightarrow 3C_3^{(4)} \rightarrow 3C_3^{(6)} \rightarrow 3C_3^{(2)} \rightarrow 3C_3^{(3)}, \quad 3C_3^{(8)} \rightarrow 3C_3^{(1)} \rightarrow 3C_3^{(7)} \rightarrow 3C_3^{(5)} \rightarrow 3C_3^{(8)}, \quad (3.7)$$

and the elements of the remaining three conjugacy classes are mapped to the elements of the same conjugacy classes under the action of the automorphism u_S . Now let us discuss the action of the outer automorphism u_T on the eleven conjugacy classes of $\Delta(27)$. From eqs. (3.2) and (A.4) we find that the results of the outer automorphism u_T acting on the conjugacy classes of $\Delta(27)$ are given by

$$u_T : 3C_3^{(3)} \rightarrow 3C_3^{(5)} \rightarrow 3C_3^{(4)} \rightarrow 3C_3^{(3)}, \quad 3C_3^{(1)} \rightarrow 3C_3^{(2)} \rightarrow 3C_3^{(6)} \rightarrow 3C_3^{(1)}. \quad (3.8)$$

Furthermore, the outer automorphism u_T acting on the other five conjugacy classes of $\Delta(27)$ will make them unchanged. If gCP symmetry is imposed, we need to consider the actions of the outer automorphism u_{K^*} on the eleven conjugacy classes. We find that the outer automorphism u_{K^*} acts on the conjugacy classes as

$$u_{K^*} : 1C_3^{(1)} \leftrightarrow 1C_3^{(2)}, \quad 3C_3^{(1)} \leftrightarrow 3C_3^{(5)}, \quad 3C_3^{(2)} \leftrightarrow 3C_3^{(3)}, \quad 3C_3^{(4)} \leftrightarrow 3C_3^{(6)}, \quad (3.9)$$

where the symbol “ \leftrightarrow ” denotes that the outer automorphism u_{K^*} interchanges the two conjugacy classes on the two sides of it. The remaining three conjugacy classes of $\Delta(27)$ are invariant under u_{K^*} .

Secondly, let us show how the two automorphisms u_S and u_T act on the eleven irreducible representations of $\Delta(27)$. The consistency condition eq. (2.9) may be understood as the action of an outer automorphism u_γ on an irreducible representation ρ of traditional flavor symmetry as follow

$$u_\gamma : \rho \rightarrow \rho', \quad \text{with} \quad \rho' = \rho \circ u_\gamma. \quad (3.10)$$

Then we have

$$\rho'(u_\gamma^{-1}(g)) = \rho(g). \quad (3.11)$$

This implies that automorphisms are symmetries of the character table. The action of the automorphism on the group elements and irreducible representations is to exchange the rows and columns of the character table. Eq. (3.10) indicates that a modular transformation connects the irreducible representation ρ of traditional flavor symmetry to another irreducible representation $\rho' = \rho \circ u_\gamma$. If $\rho' \neq \rho$, then ρ does not contain all representations that are connected via the outer automorphism u_γ , there is no modular transformation that fulfils the consistency condition (2.9). To implement this modular transformation, the representation ρ therefore has to be enlarged by the missing representations, i.e. we should extend the irreducible representation ρ to a reducible representation which contains the irreducible representations ρ and ρ' at the same time. This is the reason why we have to define

modular transformation on the reducible representation of $\Delta(27)$ sometimes, as shown in the following. Now let us give the actions of outer automorphisms u_S and u_T on irreducible representations of $\Delta(27)$ and show how to choose the representations in which modular transformation fulfils the consistency conditions eq. (2.10). The outer automorphism u_S acts on the eight nontrivial singlet irreducible representations as

$$u_S : \mathbf{1}_{0,1} \rightarrow \mathbf{1}_{2,2} \rightarrow \mathbf{1}_{0,2} \rightarrow \mathbf{1}_{1,1} \rightarrow \mathbf{1}_{0,1}, \quad \mathbf{1}_{1,0} \rightarrow \mathbf{1}_{1,2} \rightarrow \mathbf{1}_{2,0} \rightarrow \mathbf{1}_{2,1} \rightarrow \mathbf{1}_{1,0}. \quad (3.12)$$

In a similar way, we find that the outer automorphism u_T acts on the eight nontrivial singlet representations as

$$u_T : \mathbf{1}_{0,1} \rightarrow \mathbf{1}_{1,1} \rightarrow \mathbf{1}_{2,1} \rightarrow \mathbf{1}_{0,1}, \quad \mathbf{1}_{0,2} \rightarrow \mathbf{1}_{2,2} \rightarrow \mathbf{1}_{1,2} \rightarrow \mathbf{1}_{0,2}, \quad \mathbf{1}_{1,0} \rightarrow \mathbf{1}_{1,0}, \quad \mathbf{1}_{2,0} \rightarrow \mathbf{1}_{2,0}. \quad (3.13)$$

The action of the automorphisms u_S and u_T on the character table of $\Delta(27)$ is shown in table 6. The automorphisms u_S and u_T act on the character table by interchanging its the columns and rows, such that the table keeps invariant, as an outer automorphism should do.

In short, the eight nontrivial one-dimensional representations $\mathbf{1}_{(r,s)}$ of $\Delta(27)$ with $r, s \neq 0$ are related by the T' modular symmetry, in order to fulfil the consistency condition eq. (2.10) with u_S and u_T given by eq. (3.2), the eight nontrivial one-dimensional representations must be extended to a eight-dimensional reducible representation

$$\mathbf{8} = \mathbf{1}_{0,1} \oplus \mathbf{1}_{0,2} \oplus \mathbf{1}_{1,0} \oplus \mathbf{1}_{1,1} \oplus \mathbf{1}_{1,2} \oplus \mathbf{1}_{2,0} \oplus \mathbf{1}_{2,1} \oplus \mathbf{1}_{2,2}. \quad (3.14)$$

Without loss of generality, the eight-dimensional reducible representation matrices $\rho_{\mathbf{8}}(A)$ and $\rho_{\mathbf{8}}(B)$ can take the following form

$$\begin{aligned} \rho_{\mathbf{8}}(A) &= \text{diag}(\rho_{\mathbf{1}_{0,1}}(A), \rho_{\mathbf{1}_{0,2}}(A), \rho_{\mathbf{1}_{1,0}}(A), \rho_{\mathbf{1}_{1,1}}(A), \rho_{\mathbf{1}_{1,2}}(A), \rho_{\mathbf{1}_{2,0}}(A), \rho_{\mathbf{1}_{2,1}}(A), \rho_{\mathbf{1}_{2,2}}(A)) \\ &= \text{diag}(1, 1, \omega, \omega, \omega, \omega^2, \omega^2, \omega^2), \\ \rho_{\mathbf{8}}(B) &= \text{diag}(\rho_{\mathbf{1}_{0,1}}(B), \rho_{\mathbf{1}_{0,2}}(B), \rho_{\mathbf{1}_{1,0}}(B), \rho_{\mathbf{1}_{1,1}}(B), \rho_{\mathbf{1}_{1,2}}(B), \rho_{\mathbf{1}_{2,0}}(B), \rho_{\mathbf{1}_{2,1}}(B), \rho_{\mathbf{1}_{2,2}}(B)) \\ &= \text{diag}(\omega, \omega^2, 1, \omega, \omega^2, 1, \omega, \omega^2). \end{aligned} \quad (3.15)$$

Analogously the outer automorphisms u_S and u_T map the trivial singlet representation $\mathbf{1}_{0,0}$ to itself. Furthermore, the two three-dimensional representations $\mathbf{3}$ and $\bar{\mathbf{3}}$ of $\Delta(27)$ are invariant under the actions of u_S and u_T . Hence the three irreducible representations $\mathbf{1}_{0,0}$, $\mathbf{3}$ and $\bar{\mathbf{3}}$ need not be enlarged to other reducible representations to implement the outer automorphisms u_S and u_T . In order to obtain the solutions of the consistency conditions in eq. (2.10) for the outer automorphisms in eq. (3.2), we should find the solutions of the following equalities

$$\begin{aligned} \rho_{\mathbf{r}}(S) \rho_{\mathbf{r}}(A) \rho_{\mathbf{r}}^{-1}(S) &= \rho_{\mathbf{r}}(B^2 A), & \rho_{\mathbf{r}}(S) \rho_{\mathbf{r}}(B) \rho_{\mathbf{r}}^{-1}(S) &= \rho_{\mathbf{r}}(B^2 A^2), \\ \rho_{\mathbf{r}}(T) \rho_{\mathbf{r}}(A) \rho_{\mathbf{r}}^{-1}(T) &= \rho_{\mathbf{r}}(BA), & \rho_{\mathbf{r}}(T) \rho_{\mathbf{r}}(B) \rho_{\mathbf{r}}^{-1}(T) &= \rho_{\mathbf{r}}(B), \end{aligned} \quad (3.16)$$

where \mathbf{r} can be the three irreducible representations $\mathbf{1}_{0,0}$, $\mathbf{3}$ and $\bar{\mathbf{3}}$, and the reducible eight-dimensional representation $\mathbf{8}$ in eq. (3.14) of $\Delta(27)$. As elements S and T are the

generators of the finite modular group Γ'_3 , then the solutions of $\rho_{\mathbf{r}}(S)$ and $\rho_{\mathbf{r}}(T)$ must satisfy the multiplication rules in eq. (B.1), i.e.

$$\rho_{\mathbf{r}}^4(S) = \rho_{\mathbf{r}}^3(T) = [\rho_{\mathbf{r}}(S)\rho_{\mathbf{r}}(T)]^3 = \mathbb{1}_{\mathbf{r}}, \quad \rho_{\mathbf{r}}^2(S)\rho_{\mathbf{r}}(T) = \rho_{\mathbf{r}}(T)\rho_{\mathbf{r}}^2(S). \quad (3.17)$$

For the trivial singlet $\mathbf{r} = \mathbf{1}_{0,0}$, it is easy to see $\rho_{\mathbf{1}_{0,0}}(S) = 1$ and $\rho_{\mathbf{1}_{0,0}}(T) = 1, \omega$ or ω^2 . As a consequence, a field invariant under $\Delta(27)$ should be T' singlet $\mathbf{1}, \mathbf{1}'$ or $\mathbf{1}''$. For the reducible octet representation $\mathbf{r} = \mathbf{8}$ of $\Delta(27)$, we find that three solutions for $\rho_{\mathbf{r}}(S)$ and $\rho_{\mathbf{r}}(T)$ which fulfill both the consistency conditions in eq. (3.16) and the multiplication rules in eq. (3.17). In the basis of eq. (3.15), the three solutions of $\rho_{\mathbf{r}}(S)$ and $\rho_{\mathbf{r}}(T)$ take the following form:

$$\rho_{\mathbf{8}_k}(S) = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad \rho_{\mathbf{8}_k}(T) = \omega^k \begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \end{pmatrix}, \quad (3.18)$$

with $k = 0, 1, 2$. For the three solutions, the eight-dimensional representation matrices $\rho_{\mathbf{8}_k}(S)$ and $\rho_{\mathbf{8}_k}(T)$ generate the finite modular group $\Gamma'_3 \cong T'$ and the corresponding eight-dimensional representations decompose into different irreducible representations of T' as follows:

$$\mathbf{8}_k = \mathbf{1}^k \oplus \mathbf{2}^{[k+1]} \oplus \mathbf{2}^{[k+2]} \oplus \mathbf{3}, \quad (3.19)$$

in notation of appendix B. The representation matrices of the reducible octet $\mathbf{8}_k$ can be made block-diagonal by performing the similarity transformations $\Omega_{\mathbf{8}_k}\rho_{\mathbf{8}_k}(\gamma)\Omega_{\mathbf{8}_k}^\dagger$ with the unitary matrices $\Omega_{\mathbf{8}_k}$ given by

$$\Omega_{\mathbf{8}_k} = \frac{U_8^k}{2\sqrt{6}} \begin{pmatrix} \sqrt{3} & \sqrt{3} & \sqrt{3} & \sqrt{3} & \sqrt{3} & \sqrt{3} & \sqrt{3} & \sqrt{3} \\ 2\omega & -2\omega & 0 & 2\omega^2 & -2 & 0 & 2 & -2\omega^2 \\ -\sqrt{2} & \sqrt{2} & -i\sqrt{6} & -\sqrt{2} & \sqrt{2} & i\sqrt{6} & -\sqrt{2} & \sqrt{2} \\ -\sqrt{2}\omega & \sqrt{2}\omega & i\sqrt{6}\omega & -\sqrt{2}\omega & \sqrt{2}\omega & -i\sqrt{6}\omega & -\sqrt{2}\omega & \sqrt{2}\omega \\ -2 & 2 & 0 & -2\omega^2 & 2\omega & 0 & -2\omega & 2\omega^2 \\ -\omega^2 & -\omega^2 & 3\omega^2 & -\omega^2 & -\omega^2 & 3\omega^2 & -\omega^2 & -\omega^2 \\ 2\omega & 2\omega & 0 & 2 & 2\omega^2 & 0 & 2\omega^2 & 2 \\ 2 & 2 & 0 & 2\omega & 2\omega^2 & 0 & 2\omega^2 & 2\omega \end{pmatrix}, \quad (3.20)$$

with

$$U_8 = \begin{pmatrix} \mathbb{1}_5 & 0 \\ 0 & P \end{pmatrix}, \quad (3.21)$$

where the matrix P is given in eq. (B.8). Using these similarity transformation, we can obtain the decomposition of the octet $\Phi_{\mathbf{8}_k} = (\phi_1, \phi_2, \phi_3, \phi_4, \phi_5, \phi_6, \phi_7, \phi_8)^T$ into one singlet, two doublets and one triplet of the finite modular symmetry T'

$$\left\{ \begin{array}{l} \mathbf{1}^k : \quad \frac{1}{2\sqrt{2}} (\phi_1 + \phi_2 + \phi_3 + \phi_4 + \phi_5 + \phi_6 + \phi_7 + \phi_8) , \\ \mathbf{2}^{[k+1]} : \quad \frac{1}{2\sqrt{3}} \begin{pmatrix} \sqrt{2} [\omega(\phi_1 - \phi_2) + \omega^2(\phi_4 - \phi_8) - \phi_5 + \phi_7] \\ -\phi_1 + \phi_2 - i\sqrt{3}\phi_3 - \phi_4 + \phi_5 + i\sqrt{3}\phi_6 - \phi_7 + \phi_8 \end{pmatrix} , \\ \mathbf{2}^{[k+2]} : \quad \frac{1}{2\sqrt{3}} \begin{pmatrix} \omega (-\phi_1 + \phi_2 + i\sqrt{3}\phi_3 - \phi_4 + \phi_5 - i\sqrt{3}\phi_6 - \phi_7 + \phi_8) \\ \sqrt{2} [-\phi_1 + \phi_2 - \omega^2(\phi_4 - \phi_8) + \omega(\phi_5 - \phi_7)] \end{pmatrix} , \\ \mathbf{3} : \quad \frac{P^k}{2\sqrt{6}} \begin{pmatrix} -\omega^2(\phi_1 + \phi_2 - 3\phi_3 + \phi_4 + \phi_5 - 3\phi_6 + \phi_7 + \phi_8) \\ 2 [\omega(\phi_1 + \phi_2) + \phi_4 + \phi_8 + \omega^2(\phi_5 + \phi_7)] \\ 2 [\phi_1 + \phi_2 + \omega(\phi_4 + \phi_8) + \omega^2(\phi_5 + \phi_7)] \end{pmatrix} . \end{array} \right. \quad (3.22)$$

Furthermore, for the three-dimensional representation $\mathbf{3}$ of $\Delta(27)$ in eq. (A.5), the modular transformation $\rho_r(S)$ and $\rho_r(T)$ which must fulfil the consistency conditions in eq. (3.16) and the multiplication rules in eq. (3.17) are determined to be

$$\rho_{\mathbf{3}_k}(S) = \frac{i}{\sqrt{3}} \begin{pmatrix} \omega^2 & \omega & \omega \\ \omega & \omega^2 & \omega \\ \omega^2 & \omega^2 & 1 \end{pmatrix}, \quad \rho_{\mathbf{3}_k}(T) = \omega^k \begin{pmatrix} \omega & 0 & 0 \\ 0 & \omega & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (3.23)$$

for $k = 0, 1, 2$. The three dimensional representation $\rho_{\mathbf{3}_k}(S)$ and $\rho_{\mathbf{3}_k}(T)$ generate the finite modular group $\Gamma_3' \cong T'$, and it can be reduced to the direct sum of a one-dimensional and a two-dimensional irreducible representations of T' ,

$$\mathbf{3}_k = \mathbf{1}^{[k+1]} \oplus \mathbf{2}^{[k+2]}. \quad (3.24)$$

The reducible triplets $\Phi_{\mathbf{3}_k} = (\phi_1, \phi_2, \phi_3)$ transforming as $\rho_{\mathbf{3}_k}$ under T' will decompose to one singlet and one doublet of T' , which can be obtained by performing the following similarity transformation

$$\Omega_{\mathbf{3}_k} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -1 & 0 \\ 0 & 0 & \sqrt{2}\omega \\ 1 & 1 & 0 \end{pmatrix}. \quad (3.25)$$

Then the corresponding irreducible singlet and doublet are given by

$$\mathbf{1}^{[k+1]} : \frac{1}{\sqrt{2}}(\phi_1 - \phi_2), \quad \text{and} \quad \mathbf{2}^{[k+2]} : \frac{1}{\sqrt{2}} \begin{pmatrix} \sqrt{2}\omega\phi_3 \\ \phi_1 + \phi_2 \end{pmatrix}. \quad (3.26)$$

For the $\bar{\mathbf{3}}$ representation, the compatible modular transformations $\rho_r(S)$ and $\rho_r(T)$ are denoted as $\rho_{\bar{\mathbf{3}}_k}(S)$ and $\rho_{\bar{\mathbf{3}}_k}(T)$ which are the complex conjugate of $\rho_{\mathbf{3}_k}(S)$ and $\rho_{\mathbf{3}_k}(T)$ in eq. (3.23), respectively. The three reducible three-dimensional representations of T' are labelled as $\bar{\mathbf{3}}_k$ and we find the following decompositions

$$\bar{\mathbf{3}}_k = \mathbf{1}^{[2-k]} \oplus \mathbf{2}^{[1-k]}. \quad (3.27)$$

The corresponding irreducible singlet and doublet for the three reducible triplets $\bar{\mathbf{3}}_k$ are all given by the linear combinations

$$\mathbf{1}^{[2-k]} : \frac{1}{\sqrt{2}}(\phi_1 - \phi_2), \quad \text{and} \quad \mathbf{2}^{[1-k]} : \frac{1}{\sqrt{2}} \begin{pmatrix} \phi_1 + \phi_2 \\ -\sqrt{2}\omega^2\phi_3 \end{pmatrix}. \quad (3.28)$$

Finally, we shall consider the action of the automorphism u_{K_*} on the eleven representations of $\Delta(27)$. Note that the consistency condition eq. (2.21) may be read as a similarity transformation between the representations ρ^* and $\rho \circ u_{K_*}$. Similar to the two automorphisms u_S and u_T , we define the action of the outer automorphism u_{K_*} on representation as $u_{K_*} : \rho \rightarrow \rho \circ u_{K_*}$. Following the discussion above, we find that the automorphism u_{K_*} in eq. (3.4) acts on the irreducible representations of $\Delta(27)$ as

$$u_{K_*} : \mathbf{1}_{0,1} \leftrightarrow \mathbf{1}_{1,1}, \quad \mathbf{1}_{0,2} \leftrightarrow \mathbf{1}_{2,2}, \quad \mathbf{1}_{1,0} \leftrightarrow \mathbf{1}_{2,0}, \quad \mathbf{3} \leftrightarrow \bar{\mathbf{3}}, \quad (3.29)$$

and the other irreducible representations are invariant under the action of the outer automorphism u_{K_*} . The action of the automorphism u_{K_*} on the conjugacy classes and the representations of $\Delta(27)$ is shown in the table 6, and the character table of $\Delta(27)$ is invariant under u_{K_*} . In order to obtain the solution of the gCP transformation $\rho(K_*)$, one should solve the following consistency conditions

$$\begin{aligned} \rho_r(K_*)\rho_r^*(S)\rho_r^{-1}(K_*) &= \rho_r^{-1}(S), & \rho_r(K_*)\rho_r^*(T)\rho_r^{-1}(K_*) &= \rho_r^{-1}(T), & \rho_r^*(K_*) &= \rho_r^{-1}(K_*), \\ \rho_r(K_*)\rho_r^*(A)\rho_r^{-1}(K_*) &= \rho_r(A^2B), & \rho_r(K_*)\rho_r^*(B)\rho_r^{-1}(K_*) &= \rho_r(A^2BA). \end{aligned} \quad (3.30)$$

The automorphism u_{K_*} in combination with the automorphism u_S and u_T implies that the solutions of $\rho_r(S)$, $\rho_r(T)$ and $\rho_r(K_*)$ in the consistency conditions eq. (3.16) and eq. (3.30) can be obtained in the representations $\mathbf{1}_{0,0}$, $\mathbf{3}$, $\bar{\mathbf{3}}$ and $\mathbf{8}$ of $\Delta(27)$. The possible gCP transformation acting on the fields $\Phi_{\mathbf{8}_k}$ which is the reducible octet representation $\mathbf{8}$ of $\Delta(27)$ is found to be of the following form

$$\rho_{\mathbf{8}_k}(K_*) = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & P' & 0 & 0 \\ 0 & 0 & P' & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}, \quad \text{with} \quad P' = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \quad (3.31)$$

up to an overall phase. For the representations $\mathbf{3}_k$, the expressions of $\rho_{\mathbf{3}_k}(K_*)$ are given by

$$\rho_{\mathbf{3}_k}(K_*) = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & \omega \end{pmatrix}. \quad (3.32)$$

For the $\bar{\mathbf{3}}_k$ representations, the CP transformation $\rho_{\bar{\mathbf{3}}_k}(K_*)$ takes the complex conjugate of $\rho_{\mathbf{3}_k}(K_*)$ in eq. (3.32). We see that the gCP transformation matrix $\rho_{\mathbf{8}_k}(K_*)$ and $\rho_{\mathbf{3}_k}(K_*)$ are not diagonal, and the different nontrivial singlets of $\Delta(27)$ are related by gCP symmetry.

4 Effective action invariant under $\Omega(1)$ eclectic flavor symmetry

As shown in eqs. (3.12), (3.13), the eight nontrivial singlets of $\Delta(27)$ are related by the T' modular symmetry. As a consequence, all the eight nontrivial singlets carrying the same SM quantum numbers must be present or absent in a $\Omega(1)$ eclectic model. In other words, if a field is assigned to a nontrivial singlet representation of $\Delta(27)$, then the $\Omega(1)$ eclectic model has to contain other seven fields in the remaining nontrivial singlets of $\Delta(27)$ while transforming in the same way under SM gauge group. Consequently the three generations of quarks and leptons fields can only be assigned to triplets $\mathbf{3}$, $\bar{\mathbf{3}}$ or trivial singlet $\mathbf{1}_{0,0}$ of $\Delta(27)$, and the Higgs fields can only be in $\mathbf{1}_{0,0}$ if no more Higgs fields are introduced. Nevertheless, the flavons can be trivial singlet, triplet and reducible octet $\mathbf{8}$ of the traditional flavor symmetry $\Delta(27)$. The left-handed (LH) quark/lepton doublet, the right-handed (RH) quark/lepton fields and the flavon field are denoted as ψ , ψ^c and Φ , respectively. For illustration, we assume that both ψ and ψ^c transform as $\mathbf{3}$ under $\Delta(27)$ while the Higgs H_u and H_d are invariant under $\Delta(27)$. As shown in eqs. (3.23), (3.24), the T' transformation of both ψ and ψ^c would be the reducible triplet $\mathbf{3}_0$, $\mathbf{3}_1$ or $\mathbf{3}_2$, in the following $\mathbf{3}_0$ is chosen as an example. The results for other assignments of matter fields can be fixed analogously.

In the following, we shall preform a general analysis for the superpotential \mathcal{W} and the Kähler potential \mathcal{K} invariant under the eclectic flavor group $\Omega(1) \cong \Delta(27) \times T'$, and the framework of $\mathcal{N} = 1$ global supersymmetry is adopted. We assume that the level 3 modular form multiplets at weight k_Y comprise all possible irreducible multiplets of T' , i.e.

$$\begin{aligned}
 Y_{\mathbf{1}}^{(k_Y)}(\tau) &= Y_1, & Y_{\mathbf{1}'}^{(k_Y)}(\tau) &= Y_2, & Y_{\mathbf{1}''}^{(k_Y)}(\tau) &= Y_3, & & (4.1) \\
 Y_{\mathbf{2}}^{(k_Y)}(\tau) &= \begin{pmatrix} Y_4 \\ Y_5 \end{pmatrix}, & Y_{\mathbf{2}'}^{(k_Y)}(\tau) &= \begin{pmatrix} Y_6 \\ Y_7 \end{pmatrix}, & Y_{\mathbf{2}''}^{(k_Y)}(\tau) &= \begin{pmatrix} Y_8 \\ Y_9 \end{pmatrix}, & Y_{\mathbf{3}}^{(k_Y)}(\tau) &= \begin{pmatrix} Y_{10} \\ Y_{11} \\ Y_{12} \end{pmatrix}.
 \end{aligned}$$

If certain modular multiplets are absent, one can set the corresponding modular forms Y_i to be zero. If there are several linearly independent modular form multiplets at a given weight, their contributions to the effective action are of similar form and they can be straightforwardly read out from the general results in the following subsections. We shall firstly give the general form of the superpotential which is a holomorphic function of the matter fields and flavons and the couplings are generally expressed in terms of modular forms. Then we shall perform a detailed analysis of the Kähler potential.

4.1 Superpotential

Given the assignment $\psi, \psi^c \sim (\mathbf{3}, \mathbf{3}_0)$,² from the Kronecker products of $\Delta(27)$ in eq. (A.6), we find that the flavon $\Phi \equiv (\phi_1, \phi_2, \phi_3)^T$ transforming as $\mathbf{3}$ under $\Delta(27)$ must be introduced to obtain $\Omega(1)$ invariant Yukawa superpotential, and it should form a T' reducible triplet $\mathbf{3}_k$. The modular transformation of Φ is taken to be $\mathbf{3}_0$ for example. Here the flavon Φ with

²In the notation $(\mathbf{r}, \mathbf{r}')$, the two representations \mathbf{r} and \mathbf{r}' refer to the transformations under $\Delta(27)$ and T' , respectively. For any $\Delta(27)$ representation \mathbf{r} , the consistency condition of eq. (3.16) allows to fix the modular transformation in the representation \mathbf{r}' , as discussed in section 3.

modular weight k_Φ can be understood as a single flavon or combination of several flavons. Then the superpotential for the charged lepton/quark field ψ can be written as

$$\mathcal{W}_D = \frac{1}{\Lambda} \sum_{r,s} c_{r,s} \left(Y_r^{(k_Y)} \Phi \psi^c \psi \right)_{(\mathbf{1}_{\mathbf{0},\mathbf{0}},\mathbf{1}),s} H_{u/d}, \quad (4.2)$$

where $k_Y = k_\psi + k_{\psi^c} + k_\Phi$, the subscripts $\mathbf{1}_{\mathbf{0},\mathbf{0}}$ and $\mathbf{1}$ refer to trivial singlet of $\Delta(27)$ and T' respectively, and one has to sum over all modular multiplets at weight k_Y and all independent singlet contractions s . Invariance under the action of $\Delta(27)$ requires \mathcal{W}_D should be of the following form

$$\begin{aligned} \mathcal{W}_D = & \frac{1}{\Lambda} \sum_r \left(Y_r^{(k_Y)} [c_{r,1}(\psi_1^c \psi_1 \phi_1 + \psi_2^c \psi_2 \phi_2 + \psi_3^c \psi_3 \phi_3) \right. \\ & + c_{r,2}(\psi_1^c \psi_2 \phi_3 + \psi_1^c \psi_3 \phi_2 + \psi_2^c \psi_3 \phi_1 + \psi_2^c \psi_1 \phi_3 + \psi_3^c \psi_1 \phi_2 + \psi_3^c \psi_2 \phi_1) \\ & \left. + c_{r,3}(\psi_1^c \psi_2 \phi_3 - \psi_1^c \psi_3 \phi_2 + \psi_2^c \psi_3 \phi_1 - \psi_2^c \psi_1 \phi_3 + \psi_3^c \psi_1 \phi_2 - \psi_3^c \psi_2 \phi_1) \right] \Big)_{\mathbf{1}} H_{u,d}. \end{aligned} \quad (4.3)$$

We denote the above three $\Delta(27)$ invariant contractions as \mathcal{I}_1 , \mathcal{I}_2 and \mathcal{I}_3 with

$$\begin{aligned} \mathcal{I}_1 &= \psi_1^c \psi_1 \phi_1 + \psi_2^c \psi_2 \phi_2 + \psi_3^c \psi_3 \phi_3 = \left((\psi^c \psi)_{\bar{\mathbf{3}}_{S,1}} \Phi \right)_{\mathbf{1}_{\mathbf{0},\mathbf{0}}}, \\ \mathcal{I}_2 &= \psi_1^c \psi_2 \phi_3 + \psi_1^c \psi_3 \phi_2 + \psi_2^c \psi_3 \phi_1 + \psi_2^c \psi_1 \phi_3 + \psi_3^c \psi_1 \phi_2 + \psi_3^c \psi_2 \phi_1 = \left((\psi^c \psi)_{\bar{\mathbf{3}}_{S,2}} \Phi \right)_{\mathbf{1}_{\mathbf{0},\mathbf{0}}}, \\ \mathcal{I}_3 &= \psi_1^c \psi_2 \phi_3 - \psi_1^c \psi_3 \phi_2 + \psi_2^c \psi_3 \phi_1 - \psi_2^c \psi_1 \phi_3 + \psi_3^c \psi_1 \phi_2 - \psi_3^c \psi_2 \phi_1 = \left((\psi^c \psi)_{\bar{\mathbf{3}}_A} \Phi \right)_{\mathbf{1}_{\mathbf{0},\mathbf{0}}}. \end{aligned} \quad (4.4)$$

Since all the three fields $\psi \equiv (\psi_1, \psi_2, \psi_3)^T$, $\psi^c = (\psi_1^c, \psi_2^c, \psi_3^c)^T$, $\Phi \equiv (\phi_1, \phi_2, \phi_3)^T$ are T' reducible triplet $\mathbf{3}_0$, we find that the modular transformations of $\mathcal{I}_{1,2,3}$ under S and T are given by

$$\begin{aligned} \mathcal{I}_1 &\xrightarrow{S} -\frac{i}{\sqrt{3}} \mathcal{I}_1 - \frac{i}{\sqrt{3}} \omega \mathcal{I}_2, & \mathcal{I}_1 &\xrightarrow{T} \mathcal{I}_1, \\ \mathcal{I}_2 &\xrightarrow{S} -\frac{2i}{\sqrt{3}} \omega^2 \mathcal{I}_1 + \frac{i}{\sqrt{3}} \mathcal{I}_2, & \mathcal{I}_2 &\xrightarrow{T} \omega^2 \mathcal{I}_2, \\ \mathcal{I}_3 &\xrightarrow{S} \mathcal{I}_3, & \mathcal{I}_3 &\xrightarrow{T} \omega^2 \mathcal{I}_3. \end{aligned} \quad (4.5)$$

Hence the combination \mathcal{I}_3 transforms as the singlet $\mathbf{1}''$ under T' , \mathcal{I}_1 and \mathcal{I}_2 span a two-dimensional representation space and they can be arranged into T' doublet $\mathbf{2}'$, i.e.

$$\begin{pmatrix} \mathcal{I}_2 \\ -\sqrt{2} \omega^2 \mathcal{I}_1 \end{pmatrix} \sim \mathbf{2}', \quad \mathcal{I}_3 \sim \mathbf{1}''. \quad (4.6)$$

Therefore only the modular form multiplets $Y_{\mathbf{1}'}^{(k_Y)} = Y_2$ and $Y_{\mathbf{2}''}^{(k_Y)} = (Y_8, Y_9)^T$ can contract with $\mathcal{I}_{1,2,3}$ to form modular invariant combinations under the finite modular group T' . Thus the eclectic symmetry group $\Omega(1)$ severely constrains the superpotential to be

$$\begin{aligned} \mathcal{W}_D = & \frac{1}{\Lambda} \left[i\omega \alpha_1 Y_2 (\psi_1^c \psi_2 \phi_3 - \psi_1^c \psi_3 \phi_2 + \psi_2^c \psi_3 \phi_1 - \psi_2^c \psi_1 \phi_3 + \psi_3^c \psi_1 \phi_2 - \psi_3^c \psi_2 \phi_1) \right. \\ & + \sqrt{2} \alpha_2 Y_8 (\psi_1^c \psi_1 \phi_1 + \psi_2^c \psi_2 \phi_2 + \psi_3^c \psi_3 \phi_3) \\ & \left. + \omega \alpha_2 Y_9 (\psi_1^c \psi_2 \phi_3 + \psi_1^c \psi_3 \phi_2 + \psi_2^c \psi_3 \phi_1 + \psi_2^c \psi_1 \phi_3 + \psi_3^c \psi_1 \phi_2 + \psi_3^c \psi_2 \phi_1) \right] H_{u,d}, \end{aligned} \quad (4.7)$$

where $c_{2'',1} = -\sqrt{2}\omega^2 c_{2'',2} = -\sqrt{2}\alpha_2$ and $c_{1',3} = i\omega\alpha_1$. With the above convention for coupling constants, both α_1 and α_2 would be enforced to be real by the gCP transformation K_* , see eq. (4.12). After the electroweak symmetry breaking, one can read off the mass matrix of the fermion ψ as

$$M_\psi = \frac{v_{u,d}}{\Lambda} \left[i\omega\alpha_1 Y_2 \begin{pmatrix} 0 & \phi_3 & -\phi_2 \\ -\phi_3 & 0 & \phi_1 \\ \phi_2 & -\phi_1 & 0 \end{pmatrix} + \alpha_2 \begin{pmatrix} \sqrt{2}\phi_1 Y_8 & \omega\phi_3 Y_9 & \omega\phi_2 Y_9 \\ \omega\phi_3 Y_9 & \sqrt{2}\phi_2 Y_8 & \omega\phi_1 Y_9 \\ \omega\phi_2 Y_9 & \omega\phi_1 Y_9 & \sqrt{2}\phi_3 Y_8 \end{pmatrix} \right], \quad (4.8)$$

where the mass matrix M_ψ is given in the convention $\psi^c M_\psi \psi$. Note that the first term proportional to α_1 arises from the coupling with $Y_{1'}^{(k_Y)}$ and k_Y should be even, and the second term proportional to α_2 arises from the coupling with $Y_{2''}^{(k_Y)}$ for odd k_Y .

When the gCP transformation K_* is included in the theory, the transformation properties of modular multiplets, matter fields and flavons can be obtained from eqs. (3.6) and (3.32), i.e.

$$K_* : Y_{1'}^{(k_Y)} \rightarrow \left(Y_{1'}^{(k_Y)} \right)^\dagger, \quad Y_{2''}^{(k_Y)} \rightarrow \left(Y_{2''}^{(k_Y)} \right)^{\dagger T},$$

$$\begin{pmatrix} \psi_1 \\ \psi_2 \\ \psi_3 \end{pmatrix} \rightarrow \begin{pmatrix} \psi_2^\dagger \\ \psi_1^\dagger \\ \omega\psi_3^\dagger \end{pmatrix}, \quad \begin{pmatrix} \psi_1^c \\ \psi_2^c \\ \psi_3^c \end{pmatrix} \rightarrow \begin{pmatrix} \psi_2^{c\dagger} \\ \psi_1^{c\dagger} \\ \omega\psi_3^{c\dagger} \end{pmatrix}, \quad \begin{pmatrix} \phi_1 \\ \phi_2 \\ \phi_3 \end{pmatrix} \rightarrow \begin{pmatrix} \phi_2^\dagger \\ \phi_1^\dagger \\ \omega\phi_3^\dagger \end{pmatrix}. \quad (4.9)$$

Consequently under the action of K_* the superpotential \mathcal{W}_D transforms into

$$\begin{aligned} \mathcal{W}_D \xrightarrow{K_*} \frac{1}{\Lambda} \{ & i\omega^2 \alpha_1 Y_2^\dagger \left[\psi_2^{c\dagger} \psi_1^* \phi_3^\dagger - \psi_2^{c\dagger} \psi_3^* \phi_1^\dagger + \psi_1^{c\dagger} \psi_3^\dagger \phi_2^\dagger - \psi_1^{c\dagger} \psi_2^\dagger \phi_3^\dagger + \psi_3^{c\dagger} \psi_2^\dagger \phi_1^\dagger - \psi_3^{c\dagger} \psi_1^\dagger \phi_2^\dagger \right] \\ & + \sqrt{2} \alpha_2 Y_8^\dagger \left[\psi_2^{c\dagger} \psi_2^\dagger \phi_2^\dagger + \psi_1^{c\dagger} \psi_1^\dagger \phi_1^\dagger + \psi_3^{c\dagger} \psi_3^\dagger \phi_3^\dagger \right] + \omega^2 \alpha_2 Y_9^\dagger \left[\psi_2^{c\dagger} \psi_1^\dagger \phi_3^\dagger + \psi_2^{c\dagger} \psi_3^\dagger \phi_1^\dagger \right. \\ & \left. + \psi_1^{c\dagger} \psi_3^\dagger \phi_2^\dagger + \psi_1^{c\dagger} \psi_2^\dagger \phi_3^\dagger + \psi_3^{c\dagger} \psi_2^\dagger \phi_1^\dagger + \psi_3^{c\dagger} \psi_1^\dagger \phi_2^\dagger \right] \} H_{u,d}^\dagger, \end{aligned} \quad (4.10)$$

which must be equal to \mathcal{W}_D^\dagger due to gCP invariance. As a result, the couplings α_1 and α_2 should fulfill

$$\alpha_1 = \alpha_1^*, \quad \alpha_2 = \alpha_2^*. \quad (4.11)$$

Consequently both α_1 and α_2 are real, i.e.

$$\arg(\alpha_1) = 0 \pmod{\pi}, \quad \arg(\alpha_2) = 0 \pmod{\pi}. \quad (4.12)$$

In the same fashion, we can write out the $\Omega(1)$ invariant superpotential and subsequently read off the mass matrix for other assignments of matter fields and flavons. For instance, when the flavon fields are nontrivial singlets of $\Delta(27)$ with $\psi \sim (\mathbf{3}, \mathbf{3}_0)$, $\psi^c \sim (\bar{\mathbf{3}}, \bar{\mathbf{3}}_0)$ and

$\Phi \sim (\mathbf{8}, \mathbf{8}_0)$ under $\Omega(1)$, the fermion mass matrix is determined to be

$$M'_\psi = \frac{v_{u,d}}{\Lambda} \left[i\alpha_1 Y_1 \begin{pmatrix} \omega(\phi_3 + \omega\phi_6) & \phi_2 + \phi_5 + \phi_8 & \phi_1 + \omega^2\phi_4 + \omega\phi_7 \\ \phi_1 + \phi_4 + \phi_7 & \omega^2\phi_3 + \omega\phi_6 & \phi_2 + \omega\phi_5 + \omega^2\phi_8 \\ \phi_2 + \omega^2\phi_5 + \omega\phi_8 & \phi_1 + \omega\phi_4 + \omega^2\phi_7 & \phi_3 + \phi_6 \end{pmatrix} \right. \\ \left. + \omega^2\alpha_2 \begin{pmatrix} -3Y_{10}(\phi_6 + \omega^2\phi_3) & \omega Y_{10}\phi_{258} - 2Y_{11}\phi'_{852} - 2Y_{12}\phi'_{258} & Y_{10}\phi'_{417} - 2\omega^2 Y_{11}\phi_{147} - 2Y_{12}\phi'_{147} \\ \omega Y_{10}\phi_{147} - 2Y_{11}\phi'_{471} - 2Y_{12}\phi'_{174} & -3Y_{10}(\phi_3 + \omega^2\phi_6) & Y_{10}\phi'_{825} - 2\omega^2 Y_{11}\phi_{258} - 2Y_{12}\phi'_{285} \\ Y_{10}\phi'_{528} - 2Y_{11}\phi'_{582} - 2Y_{12}\phi_{258} & Y_{10}\phi'_{714} - 2Y_{11}\phi'_{741} - 2Y_{12}\phi_{147} & -3\omega Y_{10}(\phi_3 + \phi_6) \end{pmatrix} \right], \quad (4.13)$$

for even $k_Y = k_\psi + k_{\psi^c} + k_\Phi$ and

$$M''_\psi = \frac{v_{u,d}}{\Lambda} \left[i\alpha_1 \begin{pmatrix} \sqrt{3}i\omega Y_7(-\phi_3 + \omega\phi_6) & -Y_7\phi_{258} + \sqrt{2}Y_6\phi'_{582} & -\sqrt{2}Y_6\phi'_{471} + Y_7\phi'_{174} \\ Y_7\phi_{147} - \sqrt{2}Y_6\phi'_{741} & \sqrt{3}i\omega Y_7(-\omega\phi_3 + \phi_6) & \sqrt{2}Y_6\phi'_{852} - Y_7\phi'_{258} \\ \sqrt{2}\omega^2 Y_6\phi_{258} - Y_7\phi'_{285} & -\sqrt{2}\omega^2 Y_6\phi_{147} + Y_7\phi'_{147} & \sqrt{3}iY_7(-\phi_3 + \phi_6) \end{pmatrix} \right. \\ \left. + i\omega\alpha_2 \begin{pmatrix} \sqrt{3}iY_8(\phi_3 - \omega\phi_6) & -\omega^2 Y_8\phi_{258} - \sqrt{2}Y_9\phi'_{285} & \sqrt{2}Y_9\phi_{147} + Y_8\phi'_{741} \\ \omega^2 Y_8\phi_{147} + \sqrt{2}Y_9\phi'_{147} & \sqrt{3}iY_8(\omega\phi_3 - \phi_6) & -Y_8\phi'_{582} - \sqrt{2}Y_9\phi_{258} \\ -Y_8\phi'_{852} - \sqrt{2}Y_9\phi'_{258} & Y_8\phi'_{471} + \sqrt{2}Y_9\phi'_{174} & \sqrt{3}i\omega^2 Y_8(\phi_3 - \phi_6) \end{pmatrix} \right], \quad (4.14)$$

for odd $k_Y = k_\psi + k_{\psi^c} + k_\Phi$. Here the functions ϕ_{ijk} and ϕ'_{ijk} are defined as

$$\phi_{ijk} = \phi_i + \phi_j + \phi_k, \quad \phi'_{ijk} = \phi_i + \omega\phi_j + \omega^2\phi_k. \quad (4.15)$$

Furthermore, we summarize the fermion mass matrices for all independent assignments of quarks/leptons and flavons in table 1. The mass matrix would become its transpose, if the transformation properties of ψ and ψ^c are exchanged.

4.1.1 The superpotential of Majorana mass terms

The Majorana mass term for the fermion field ψ^c (e.g., the right-handed neutrinos) can be generally written as

$$\mathcal{W}_M = \sum_{r,s} d_{r,s} \left(Y_r^{(k_Y)} \psi^c \psi^c \Phi \right)_{(\mathbf{1}_0, \mathbf{0}, \mathbf{1}), s}, \quad (4.16)$$

where $k_Y = 2k_{\psi^c} + k_\Phi$, and Φ can be interpreted as one/several flavons or as combination of Higgs fields with flavons.³ For illustration, we consider the case that both ψ^c and Φ transform as $(\mathbf{3}, \mathbf{3}_0)$ under the eclectic symmetry group $\Omega(1)$.

From the tensor product $\mathbf{3} \otimes \mathbf{3} = \bar{\mathbf{3}}_{S,1} \oplus \bar{\mathbf{3}}_{S,2} \oplus \bar{\mathbf{3}}_A$ of $\Delta(27)$ in appendix A, we know the contraction of $\psi^c \psi^c$ gives rise to two symmetric triplets:

$$(\psi^c \psi^c)_{\bar{\mathbf{3}}_{S,1}} = \begin{pmatrix} (\psi_1^c)^2 \\ (\psi_2^c)^2 \\ (\psi_3^c)^2 \end{pmatrix}, \quad (\psi^c \psi^c)_{\bar{\mathbf{3}}_{S,2}} = \begin{pmatrix} \psi_2^c \psi_3^c \\ \psi_1^c \psi_3^c \\ \psi_1^c \psi_2^c \end{pmatrix}. \quad (4.17)$$

³For example, Φ is the combination of two Higgs fields and flavon in the Weinberg operator, ψ^c would be the LH lepton doublet.

\mathcal{W}	$(\Delta(27), T')$	$Y_r^{(k_Y)}$	M_ψ
\mathcal{W}_{D1}	$\psi \sim (\mathbf{3}, \mathbf{3}_0), \Phi \sim (\bar{\mathbf{3}}, \bar{\mathbf{3}}_0)$ $(\psi_1^c, \psi_2^c, \psi_3^c) \sim (\mathbf{1}_{(0,0)}, \mathbf{1}^{[3-k]})$	$Y_{\mathbf{1}^k}^{(k_Y)}$	$\frac{v_{u,d} Y_k}{\Lambda} \begin{pmatrix} \alpha_1 \phi_1 & \alpha_1 \phi_2 & \alpha_1 \phi_3 \\ \alpha_2 \phi_1 & \alpha_2 \phi_2 & \alpha_2 \phi_3 \\ \alpha_3 \phi_1 & \alpha_3 \phi_2 & \alpha_3 \phi_3 \end{pmatrix}$
\mathcal{W}_{D2}	$\psi \sim (\mathbf{3}, \mathbf{3}_0),$ $\psi_m^c \sim (\mathbf{1}_{(0,0)}, \mathbf{1}^{m-1}),$ $\Phi \sim (\bar{\mathbf{3}}, \bar{\mathbf{3}}_k)$	$Y_{\mathbf{1}}^{(k_Y)}$ $Y_{\mathbf{1}'}^{(k_Y)}$ $Y_{\mathbf{1}''}^{(k_Y)}$	$\frac{v_{u,d}}{\Lambda} P^k \begin{pmatrix} \alpha_1 Y_1 \phi_1 & \alpha_1 Y_1 \phi_2 & \alpha_1 Y_1 \phi_3 \\ \alpha_2 Y_3 \phi_1 & \alpha_2 Y_3 \phi_2 & \alpha_2 Y_3 \phi_3 \\ \alpha_3 Y_2 \phi_1 & \alpha_3 Y_2 \phi_2 & \alpha_3 Y_2 \phi_3 \end{pmatrix}$
\mathcal{W}_{D3}	$\psi, \psi^c \sim (\mathbf{3}, \mathbf{3}_0), \Phi \sim (\mathbf{3}, \mathbf{3}_0)$	$Y_{\mathbf{1}'}^{(k_Y)}, Y_{\mathbf{2}''}^{(k_Y)}$	M_ψ in eq. (4.8)
\mathcal{W}_{D4}	$\psi, \psi^c \sim (\mathbf{3}, \mathbf{3}_0), \Phi \sim (\mathbf{3}, \mathbf{3}_1)$	$Y_{\mathbf{1}}^{(k_Y)}, Y_{\mathbf{2}'}^{(k_Y)}$	$M_\psi (Y_2 \rightarrow Y_1, Y_8 \rightarrow Y_6, Y_9 \rightarrow Y_7)$
\mathcal{W}_{D5}	$\psi, \psi^c \sim (\mathbf{3}, \mathbf{3}_0), \Phi \sim (\mathbf{3}, \mathbf{3}_2)$	$Y_{\mathbf{1}''}^{(k_Y)}, Y_{\mathbf{2}'}^{(k_Y)}$	$M_\psi (Y_2 \rightarrow Y_3, Y_8 \rightarrow Y_4, Y_9 \rightarrow Y_5)$
\mathcal{W}_{D6}	$\psi, \psi^c \sim (\bar{\mathbf{3}}, \bar{\mathbf{3}}_0), \Phi \sim (\bar{\mathbf{3}}, \bar{\mathbf{3}}_0)$	$Y_{\mathbf{1}''}^{(k_Y)}, Y_{\mathbf{2}'}^{(k_Y)}$	$M_\psi (Y_2 \rightarrow \omega Y_3, Y_8 \rightarrow Y_7, Y_9 \rightarrow -\omega Y_6)$
\mathcal{W}_{D7}	$\psi, \psi^c \sim (\bar{\mathbf{3}}, \bar{\mathbf{3}}_0), \Phi \sim (\bar{\mathbf{3}}, \bar{\mathbf{3}}_1)$	$Y_{\mathbf{1}}^{(k_Y)}, Y_{\mathbf{2}''}^{(k_Y)}$	$M_\psi (Y_2 \rightarrow \omega Y_1, Y_8 \rightarrow Y_9, Y_9 \rightarrow -\omega Y_8)$
\mathcal{W}_{D8}	$\psi, \psi^c \sim (\bar{\mathbf{3}}, \bar{\mathbf{3}}_0), \Phi \sim (\bar{\mathbf{3}}, \bar{\mathbf{3}}_2)$	$Y_{\mathbf{1}'}^{(k_Y)}, Y_{\mathbf{2}}^{(k_Y)}$	$M_\psi (Y_2 \rightarrow \omega Y_2, Y_8 \rightarrow Y_5, Y_9 \rightarrow -\omega Y_4)$
\mathcal{W}_{D9}	$\psi \sim (\mathbf{3}, \mathbf{3}_0), \psi^c \sim (\bar{\mathbf{3}}, \bar{\mathbf{3}}_0),$ $\Phi \sim (\mathbf{8}, \mathbf{8}_0)$	$Y_{\mathbf{1}}^{(k_Y)}, Y_{\mathbf{3}}^{(k_Y)}$	M'_ψ in eq. (4.13) for even k_Y
		$Y_{\mathbf{2}'}^{(k_Y)}, Y_{\mathbf{2}''}^{(k_Y)}$	M''_ψ in eq. (4.14) for odd k_Y
\mathcal{W}_{D10}	$\psi \sim (\mathbf{3}, \mathbf{3}_0), \psi^c \sim (\bar{\mathbf{3}}, \bar{\mathbf{3}}_0),$ $\Phi \sim (\mathbf{8}, \mathbf{8}_1)$	$Y_{\mathbf{1}''}^{(k_Y)}, Y_{\mathbf{3}}^{(k_Y)}$	$M'_\psi (Y_1 \rightarrow Y_3, Y_3^{(k_Y)} \rightarrow P Y_3^{(k_Y)})$
		$Y_{\mathbf{2}'}^{(k_Y)}, Y_{\mathbf{2}''}^{(k_Y)}$	$M''_\psi (Y_{\mathbf{2}'}^{(k_Y)} \rightarrow Y_{\mathbf{2}}^{(k_Y)}, Y_{\mathbf{2}''}^{(k_Y)} \rightarrow Y_{\mathbf{2}'}^{(k_Y)})$
\mathcal{W}_{D11}	$\psi \sim (\mathbf{3}, \mathbf{3}_0), \psi^c \sim (\bar{\mathbf{3}}, \bar{\mathbf{3}}_0),$ $\Phi \sim (\mathbf{8}, \mathbf{8}_2)$	$Y_{\mathbf{1}'}^{(k_Y)}, Y_{\mathbf{3}}^{(k_Y)}$	$M'_\psi (Y_1 \rightarrow Y_2, Y_3^{(k_Y)} \rightarrow P^2 Y_3^{(k_Y)})$
		$Y_{\mathbf{2}}^{(k_Y)}, Y_{\mathbf{2}''}^{(k_Y)}$	$M''_\psi (Y_{\mathbf{2}'}^{(k_Y)} \rightarrow \omega Y_{\mathbf{2}''}^{(k_Y)}, Y_{\mathbf{2}''}^{(k_Y)} \rightarrow \omega^2 Y_{\mathbf{2}}^{(k_Y)})$

Table 1. The mass matrices of the charged leptons/quarks ψ for all independent assignments of ψ, ψ^c and Φ under the eclectic symmetry group $\Omega(1) \equiv \Delta(27) \times T'$, where P is permutation matrix given in eq. (B.8). In the third column, we list the modular form multiplets which give nonzero contributions. In our convention, all couplings in M_ψ are fixed to be real by the gCP transformation K_* .

Consequently there are only two $\Delta(27)$ invariant contractions for the Majorana mass term $\psi^c \psi^c \Phi$,

$$\begin{aligned}
 \left((\psi^c \psi^c)_{\bar{\mathbf{3}}_{S,1}} \Phi \right)_{\mathbf{1}_{0,0}} &= (\psi_1^c)^2 \phi_1 + (\psi_2^c)^2 \phi_2 + (\psi_3^c)^2 \phi_3, \\
 \left((\psi^c \psi^c)_{\bar{\mathbf{3}}_{S,2}} \Phi \right)_{\mathbf{1}_{0,0}} &= \psi_1^c \psi_2^c \phi_3 + \psi_1^c \psi_3^c \phi_2 + \psi_2^c \psi_3^c \phi_1.
 \end{aligned} \tag{4.18}$$

The modulus and modular forms are invariant under $\Delta(27)$ flavor symmetry, thus $\Delta(27)$ invariance constrains the superpotential \mathcal{W}_M to be

$$\mathcal{W}_M = \sum_r \left(Y_r^{(k_Y)} \left[d_{r,1} \left((\psi_1^c)^2 \phi_1 + (\psi_2^c)^2 \phi_2 + (\psi_3^c)^2 \phi_3 \right) + d_{r,2} \left(\psi_1^c \psi_2^c \phi_3 + \psi_1^c \psi_3^c \phi_2 + \psi_2^c \psi_3^c \phi_1 \right) \right] \right)_{\mathbf{1}}. \tag{4.19}$$

The two $\Delta(27)$ singlet contractions in eq. (4.18) can be arranged into a T' doublet $\mathbf{2}'$:

$$\begin{pmatrix} \psi_1^c \psi_2^c \phi_3 + \psi_1^c \psi_3^c \phi_2 + \psi_2^c \psi_3^c \phi_1 \\ -\omega^2 \left((\psi_1^c)^2 \phi_1 + (\psi_2^c)^2 \phi_2 + (\psi_3^c)^2 \phi_3 \right) / \sqrt{2} \end{pmatrix} \sim \mathbf{2}'. \tag{4.20}$$

\mathcal{W}	$(\Delta(27), T')$	$Y_r^{(k_Y)}$	M_{ψ^c}
\mathcal{W}_{M1}	$\psi^c \sim (\mathbf{3}, \mathbf{3}_0), \Phi \sim (\mathbf{3}, \mathbf{3}_0)$	$Y_{2''}^{(k_Y)}$	M_{ψ^c} in eq. (4.22)
\mathcal{W}_{M2}	$\psi^c \sim (\mathbf{3}, \mathbf{3}_0), \Phi \sim (\mathbf{3}, \mathbf{3}_1)$	$Y_{2'}^{(k_Y)}$	$M_{\psi^c} (Y_8 \rightarrow Y_6, Y_9 \rightarrow Y_7)$
\mathcal{W}_{M3}	$\psi^c \sim (\mathbf{3}, \mathbf{3}_0), \Phi \sim (\mathbf{3}, \mathbf{3}_2)$	$Y_2^{(k_Y)}$	$M_{\psi^c} (Y_8 \rightarrow Y_4, Y_9 \rightarrow Y_5)$
\mathcal{W}_{M4}	$\psi^c \sim (\bar{\mathbf{3}}, \bar{\mathbf{3}}_0), \Phi \sim (\bar{\mathbf{3}}, \bar{\mathbf{3}}_0)$	$Y_{2'}^{(k_Y)}$	$M_{\psi^c} (Y_8 \rightarrow Y_7, Y_9 \rightarrow -\omega Y_6)$
\mathcal{W}_{M5}	$\psi^c \sim (\bar{\mathbf{3}}, \bar{\mathbf{3}}_0), \Phi \sim (\bar{\mathbf{3}}, \bar{\mathbf{3}}_1)$	$Y_{2''}^{(k_Y)}$	$M_{\psi^c} (Y_8 \rightarrow Y_9, Y_9 \rightarrow -\omega Y_8)$
\mathcal{W}_{M6}	$\psi^c \sim (\bar{\mathbf{3}}, \bar{\mathbf{3}}_0), \Phi \sim (\bar{\mathbf{3}}, \bar{\mathbf{3}}_2)$	$Y_2^{(k_Y)}$	$M_{\psi^c} (Y_8 \rightarrow Y_5, Y_9 \rightarrow -\omega Y_4)$

Table 2. The mass matrices of the Majorana fermion ψ^c for possible independent assignments of ψ^c and Φ under the eclectic symmetry group $\Omega(1) \equiv \Delta(27) \rtimes T'$. The singlet assignments of ψ^c and Φ are omitted, as $\Omega(1)$ would play less rule. In the third column, we list the modular form multiplets giving nonzero contributions. In our convention, all couplings in M_{ψ^c} are fixed to be real by the gCP transformation K_* .

Consequently modular invariance of \mathcal{W}_M requires $d_{2'',2} = -\sqrt{2}\omega d_{2'',1}$ and only the modular multiplet $Y_{2''}^{(k_Y)}(\tau) = (Y_8, Y_9)^T$ is relevant. Thus the eclectic flavor symmetry $\Omega(1)$ fixes the superpotential \mathcal{W}_M to be

$$\mathcal{W}_M = \alpha \left\{ \sqrt{2}Y_8 \left[(\psi_1^c)^2 \phi_1 + (\psi_2^c)^2 \phi_2 + (\psi_3^c)^2 \phi_3 \right] + 2\omega Y_9 (\psi_1^c \psi_2^c \phi_3 + \psi_1^c \psi_3^c \phi_2 + \psi_2^c \psi_3^c \phi_1) \right\}, \quad (4.21)$$

where $\alpha = \omega^2 d_{2'',2}/2$ is a general complex number. Thus we can read out the Majorana mass matrix of ψ^c as follow

$$M_{\psi^c} = \alpha \begin{pmatrix} \sqrt{2}Y_8\phi_1 & \omega Y_9\phi_3 & \omega Y_9\phi_2 \\ \omega Y_9\phi_3 & \sqrt{2}Y_8\phi_2 & \omega Y_9\phi_1 \\ \omega Y_9\phi_2 & \omega Y_9\phi_1 & \sqrt{2}Y_8\phi_3 \end{pmatrix}. \quad (4.22)$$

If the gCP transformation K_* is considered, the superpotential \mathcal{W}_M transforms as

$$\mathcal{W}_M \xrightarrow{K_*} \alpha \left\{ \sqrt{2}Y_8^\dagger \left[(\psi_2^{c\dagger})^2 \phi_2^\dagger + (\psi_1^{c\dagger})^2 \phi_1^\dagger + (\psi_3^{c\dagger})^2 \phi_3^\dagger \right] + 2\omega^2 Y_9^\dagger (\psi_2^{c\dagger} \psi_1^{c\dagger} \phi_3^\dagger + \psi_2^{c\dagger} \psi_3^{c\dagger} \phi_1^\dagger + \psi_1^{c\dagger} \psi_3^{c\dagger} \phi_2^\dagger) \right\}, \quad (4.23)$$

which must equal to \mathcal{W}_M^\dagger . Then we have $\alpha^* = \alpha$ and consequently

$$\arg(\alpha) = 0 \pmod{\pi}. \quad (4.24)$$

For other assignments of ψ^c and Φ , the corresponding predictions for the mass matrix of ψ^c are listed in table 2.

4.2 Kähler potential

The Kähler potential is a hermitian function of the modulus τ , the chiral superfields of quarks/leptons ψ , ψ^c and the flavon superfield Φ as well as their complex conjugates τ^\dagger , ψ^\dagger , $\psi^{c\dagger}$ and Φ^\dagger . As shown in eq. (1.1), many terms on the same footing as the minimal Kähler potential are compatible with modular symmetry [6]. They can affect the fermion

masses and mixing parameters significantly, see refs. [6, 7] for explicit examples. In the paradigm of eclectic flavor symmetry, the Kähler potential has to be invariant under both modular symmetry and the traditional flavor symmetry transformations. In particular, the traditional flavor symmetry can impose severe constraints on the Kähler potential so that it can be expanded in power of $\langle \Phi \rangle / \Lambda$, where Φ denotes the VEVs of the flavon fields. As explained in previous section, the matter fields ψ and ψ^c can only be singlet $\mathbf{1}_{0,0}$ or triplets $\mathbf{3}$ and $\bar{\mathbf{3}}$ of $\Delta(27)$. As an example, we assume that both ψ and ψ^c transform as $(\mathbf{3}, \mathbf{3}_0)$ under $\Omega(1)$ eclectic flavor group in this section. Since the Kähler potential of ψ has a very similar form to that of ψ^c , consequently we shall focus on ψ in the following.

At leading order, the Kähler potential for the matter fields ψ can be written as⁴

$$\mathcal{K}_{\text{LO}} = \sum_{n, \mathbf{r}_1, \mathbf{r}_2, s} (-i\tau + i\bar{\tau})^{-k_\psi + n} \left(Y_{\mathbf{r}_1}^{(n)\dagger} Y_{\mathbf{r}_2}^{(n)} \psi^\dagger \psi \right)_{(\mathbf{1}_{0,0,1}), s}, \quad (4.25)$$

where the couplings of each independent term are dropped, and we should sum over the weights $n \in \mathbb{N}$ and representation \mathbf{r} of all linearly independent modular multiplets $Y_{\mathbf{r}}^{(n)}(\tau)$ and all $\Omega(1)$ singlet contractions labelled by the index s . The modular form with weight zero is a constant and it is taken to be $Y_{\mathbf{r}}^{(0)} = 1$ without loss of generality. The minimal Kähler potential corresponds to the term of $n = 0$. Because the modular forms are invariant under $\Delta(27)$ flavor symmetry, $\psi^\dagger \psi$ has to be a $\Delta(27)$ trivial singlet, i.e.

$$\left(\psi^\dagger \psi \right)_{\mathbf{1}_{0,0}} = \psi_1^\dagger \psi_1 + \psi_2^\dagger \psi_2 + \psi_3^\dagger \psi_3 = \left(\psi^\dagger \psi \right)_{(\mathbf{1}_{0,0,1})}, \quad (4.26)$$

which is also an invariant singlet $\mathbf{1}$ of the T' modular symmetry. Hence the invariance of \mathcal{K} under $\Delta(27)$ requires the leading order Kähler potential in eq. (4.25) to be of the following form

$$\mathcal{K}_{\text{LO}} = \sum_{n, \mathbf{r}_1, \mathbf{r}_2} (-i\tau + i\bar{\tau})^{-k_\psi + n} \left(Y_{\mathbf{r}_1}^{(n)\dagger} Y_{\mathbf{r}_2}^{(n)} \right)_{(\mathbf{1}_{0,0,1})} \left(\psi^\dagger \psi \right)_{(\mathbf{1}_{0,0,1})}. \quad (4.27)$$

Note that the non-vanishing contraction $\left(Y_{\mathbf{r}_1}^{(n)\dagger} Y_{\mathbf{r}_2}^{(n)} \right)_{(\mathbf{1}_{0,0,1})}$ requires that \mathbf{r}_1 and \mathbf{r}_2 should be the same representation $\mathbf{r}_1 = \mathbf{r}_2 = \mathbf{r}$ of T' . Moreover, from the Kronecker products and Clebsch-Gordon (CG) coefficients of T' in appendix B, we see that there is only a unique singlet contraction $\left(Y_{\mathbf{r}}^{(n)\dagger} Y_{\mathbf{r}}^{(n)} \right)_{(\mathbf{1}_{0,0,1})}$ for any irreducible representation of T' , i.e.

$$\left(Y_{\mathbf{r}}^{(n)\dagger} Y_{\mathbf{r}}^{(n)} \right)_{(\mathbf{1}_{0,0,1})} = Y_{\mathbf{r},1}^{(n)\dagger} Y_{\mathbf{r},1}^{(n)} + Y_{\mathbf{r},2}^{(n)\dagger} Y_{\mathbf{r},2}^{(n)} + \dots \quad (4.28)$$

Then the most general leading order Kähler potential invariant under $\Omega(1)$ for ψ takes the form

$$\begin{aligned} \mathcal{K}_{\text{LO}} &= \sum_{n, \mathbf{r}} (-i\tau + i\bar{\tau})^{-k_\psi + n} \left(Y_{\mathbf{r}}^{(n)\dagger} Y_{\mathbf{r}}^{(n)} \right)_{(\mathbf{1}_{0,0,1})} \left(\psi^\dagger \psi \right)_{(\mathbf{1}_{0,0,1})} \\ &= \sum_{n, \mathbf{r}} (-i\tau + i\bar{\tau})^{-k_\psi + n} \left(Y_{\mathbf{r}}^{(n)\dagger} Y_{\mathbf{r}}^{(n)} \right)_{(\mathbf{1}_{0,0,1})} \left(\psi_1^\dagger \psi_1 + \psi_2^\dagger \psi_2 + \psi_3^\dagger \psi_3 \right). \end{aligned} \quad (4.29)$$

⁴The modular transformation of $-i\tau + i\bar{\tau}$ is

$$-i\tau + i\bar{\tau} \xrightarrow{\gamma} |c\tau + d|^{-2} (-i\tau + i\bar{\tau}).$$

Hence the Kähler metric is diagonal and it is proportional to a unit matrix at leading order. In order to get canonical kinetic terms, we have to rescale the matter fields ψ . The effect of such rescaling on fermion masses can be absorbed into the parameters of the superpotential. Thus leading order Kähler potential invariant under $\Omega(1)$ does not introduce any corrections to the flavor observables due to the traditional flavor symmetry $\Delta(27)$. If the three generations of matter fields ψ transform as $(\mathbf{1}_{\mathbf{0},\mathbf{0}}, \mathbf{1}^k)$ under the eclectic flavor group $\Delta(27) \rtimes T'$, they are generally distinguished by the different charges under auxiliary abelian symmetry, so that the Kähler potential would be still diagonal.

Then we proceed to consider the next-to-leading-order (NLO) corrections to the Kähler potential of ψ . It involve only one flavon Φ and its most general form is given by

$$\mathcal{K}_{\text{NLO}} = \sum_{m,n,r_1,r_2,s} (-i\tau + i\bar{\tau})^{-k_\psi+m} \left(Y_{r_1}^{(m)\dagger} Y_{r_2}^{(n)} \psi^\dagger \psi \Phi \right)_{(\mathbf{1}_{\mathbf{0},\mathbf{0}}, \mathbf{1}),s} + \text{h.c.}, \quad (4.30)$$

where the modular weight k_Φ of the flavon Φ should fulfill

$$k_\Phi = n - m. \quad (4.31)$$

If some auxiliary symmetry is imposed, obviously the flavon Φ should be chargeless under the auxiliary group. From the $\Delta(27)$ tensor product $\bar{\mathbf{3}} \otimes \mathbf{3} = \sum_{r,s=0}^2 \mathbf{1}_{r,s}$ in eq. (A.6), we know the contractions of $\psi^\dagger \psi$ are

$$\begin{aligned} (\psi^\dagger \psi)_{(\mathbf{1}_{\mathbf{0},\mathbf{0}}, \mathbf{1})} &= \psi_1^\dagger \psi_1 + \psi_2^\dagger \psi_2 + \psi_3^\dagger \psi_3, \\ (\psi^\dagger \psi)_{(\mathbf{8}, \mathbf{8}_0)} &= \begin{pmatrix} \psi_1^\dagger \psi_2 + \psi_2^\dagger \psi_3 + \psi_3^\dagger \psi_1 \\ \psi_1^\dagger \psi_3 + \psi_2^\dagger \psi_1 + \psi_3^\dagger \psi_2 \\ \psi_1^\dagger \psi_1 + \omega^2 \psi_2^\dagger \psi_2 + \omega \psi_3^\dagger \psi_3 \\ \omega^2 \psi_1^\dagger \psi_2 + \omega \psi_2^\dagger \psi_3 + \psi_3^\dagger \psi_1 \\ \omega \psi_1^\dagger \psi_3 + \psi_2^\dagger \psi_1 + \omega^2 \psi_3^\dagger \psi_2 \\ \psi_1^\dagger \psi_1 + \omega \psi_2^\dagger \psi_2 + \omega^2 \psi_3^\dagger \psi_3 \\ \omega \psi_1^\dagger \psi_2 + \omega^2 \psi_2^\dagger \psi_3 + \psi_3^\dagger \psi_1 \\ \omega^2 \psi_1^\dagger \psi_3 + \psi_2^\dagger \psi_1 + \omega \psi_3^\dagger \psi_2 \end{pmatrix}. \end{aligned} \quad (4.32)$$

Invariance under $\Delta(27)$ require $\psi^\dagger \psi \Phi$ should be a trivial singlet of $\Delta(27)$, so that the flavon Φ_i must transform as $\mathbf{1}_{\mathbf{0},\mathbf{0}}$ or $\mathbf{8}$ under $\Delta(27)$. In the case of $\Phi \sim (\mathbf{1}_{\mathbf{0},\mathbf{0}}, \mathbf{1}^k)$, we find

$$\begin{aligned} (\psi^\dagger \psi \Phi)_{(\mathbf{1}_{\mathbf{0},\mathbf{0}}, \mathbf{1}^k)} &= (\psi^\dagger \psi)_{(\mathbf{1}_{\mathbf{0},\mathbf{0}}, \mathbf{1})} \Phi = (\psi_1^\dagger \psi_1 + \psi_2^\dagger \psi_2 + \psi_3^\dagger \psi_3) \Phi, \\ (Y_{r_1}^{(m)\dagger} Y_{r_2}^{(n)} \psi^\dagger \psi \Phi)_{(\mathbf{1}_{\mathbf{0},\mathbf{0}}, \mathbf{1})} &= (Y_{r_1}^{(m)\dagger} Y_{r_2}^{(n)})_{(\mathbf{1}_{\mathbf{0},\mathbf{0}}, \mathbf{1}^{[3-k]})} (\psi^\dagger \psi)_{(\mathbf{1}_{\mathbf{0},\mathbf{0}}, \mathbf{1})} \Phi, \end{aligned} \quad (4.33)$$

which is proportional to $\psi_1^\dagger \psi_1 + \psi_2^\dagger \psi_2 + \psi_3^\dagger \psi_3$. As a result, the corresponding contribution can be absorbed into the LO Kähler potential in eq. (4.29), and it yields no corrections to the observable fermion masses and mixing. If the flavon transform as $\Phi \sim (\mathbf{8}, \mathbf{8}_k)$, the NLO Kähler potential \mathcal{K}_{NLO} invariant under the eclectic flavor group $\Omega(1)$ would be of the

following form

$$\begin{aligned}
 \mathcal{K}_{\text{NLO}} = & \sum_{m,n,r_1,r_2} (-i\tau + i\bar{\tau})^{-k_\psi+m} \left(Y_{r_1}^{(m)\dagger} Y_{r_2}^{(n)} \right)_{(\mathbf{10},\mathbf{0},\mathbf{1}^{[3-k]})} \left((\psi^\dagger \psi)_{(\mathbf{8},\mathbf{8}_0)} \Phi \right)_{(\mathbf{10},\mathbf{0},\mathbf{1}^{[k]})} \\
 & + \sum_{m,n,r_1,r_2} (-i\tau + i\bar{\tau})^{-k_\psi+m} \left((Y_{r_1}^{(m)\dagger} Y_{r_2}^{(n)})_{(\mathbf{10},\mathbf{0},\mathbf{2}^{[2-k]})} \left((\psi^\dagger \psi)_{(\mathbf{8},\mathbf{8}_0)} \Phi \right)_{(\mathbf{10},\mathbf{0},\mathbf{2}^{[1+k]})} \right)_{(\mathbf{10},\mathbf{0},\mathbf{1})} \\
 & + \sum_{m,n,r_1,r_2} (-i\tau + i\bar{\tau})^{-k_\psi+m} \left((Y_{r_1}^{(m)\dagger} Y_{r_2}^{(n)})_{(\mathbf{10},\mathbf{0},\mathbf{2}^{[1-k]})} \left((\psi^\dagger \psi)_{(\mathbf{8},\mathbf{8}_0)} \Phi \right)_{(\mathbf{10},\mathbf{0},\mathbf{2}^{[2+k]})} \right)_{(\mathbf{10},\mathbf{0},\mathbf{1})} \\
 & + \sum_{m,n,r_1,r_2} (-i\tau + i\bar{\tau})^{-k_\psi+m} \left((Y_{r_1}^{(m)\dagger} Y_{r_2}^{(n)})_{(\mathbf{10},\mathbf{0},\mathbf{3})} \left((\psi^\dagger \psi)_{(\mathbf{8},\mathbf{8}_0)} \Phi \right)_{(\mathbf{10},\mathbf{0},\mathbf{3})} \right)_{(\mathbf{10},\mathbf{0},\mathbf{1})} + \text{h.c.} .
 \end{aligned} \tag{4.34}$$

The general formulas for the $\Delta(27)$ trivial singlet contractions of two octets $\chi \sim (\mathbf{8}, \mathbf{8}_k)$ and $\varphi \sim (\mathbf{8}, \mathbf{8}_{k'})$ are given by:

$$\begin{aligned}
 (\chi\varphi)_{(\mathbf{10},\mathbf{0},\mathbf{1}^{[k+k']})} &= \chi_1\varphi_2 + \chi_2\varphi_1 + \chi_3\varphi_6 + \chi_4\varphi_8 + \chi_5\varphi_7 + \chi_6\varphi_3 + \chi_7\varphi_5 + \chi_8\varphi_4, \\
 (\chi\varphi)_{(\mathbf{10},\mathbf{0},\mathbf{2}^{[k+k'+1]})} &= \begin{pmatrix} \sqrt{2}(\omega\chi_1\varphi_2 - \omega\chi_2\varphi_1 + \omega^2\chi_4\varphi_8 - \chi_5\varphi_7 + \chi_7\varphi_5 - \omega^2\chi_8\varphi_4) \\ -\chi_1\varphi_2 + \chi_2\varphi_1 - i\sqrt{3}\chi_3\varphi_6 - \chi_4\varphi_8 + \chi_5\varphi_7 + i\sqrt{3}\chi_6\varphi_3 - \chi_7\varphi_5 + \chi_8\varphi_4 \end{pmatrix}, \\
 (\chi\varphi)_{(\mathbf{10},\mathbf{0},\mathbf{2}^{[k+k'+2]})} &= \begin{pmatrix} \chi_1\varphi_2 - \chi_2\varphi_1 - i\sqrt{3}\chi_3\varphi_6 + \chi_4\varphi_8 - \chi_5\varphi_7 + i\sqrt{3}\chi_6\varphi_3 + \chi_7\varphi_5 - \chi_8\varphi_4 \\ \sqrt{2}(\omega^2\chi_1\varphi_2 - \omega^2\chi_2\varphi_1 + \omega\chi_4\varphi_8 - \chi_5\varphi_7 + \chi_7\varphi_5 - \omega\chi_8\varphi_4) \end{pmatrix}, \\
 (\chi\varphi)_{(\mathbf{10},\mathbf{0},\mathbf{3})} &= P^{k+k'} \begin{pmatrix} \chi_1\varphi_2 + \chi_2\varphi_1 - 3\chi_3\varphi_6 + \chi_4\varphi_8 + \chi_5\varphi_7 - 3\chi_6\varphi_3 + \chi_7\varphi_5 + \chi_8\varphi_4 \\ -2(\omega^2\chi_1\varphi_2 + \omega^2\chi_2\varphi_1 + \omega\chi_4\varphi_8 + \chi_5\varphi_7 + \chi_7\varphi_5 + \omega\chi_8\varphi_4) \\ -2(\omega\chi_1\varphi_2 + \omega\chi_2\varphi_1 + \omega^2\chi_4\varphi_8 + \chi_5\varphi_7 + \chi_7\varphi_5 + \omega^2\chi_8\varphi_4) \end{pmatrix},
 \end{aligned} \tag{4.35}$$

with the permutation matrix P given in eq. (B.8). Using the contraction rules in eq. (4.35), we can write out the explicit expression of \mathcal{K}_{NLO} , yet it is too lengthy to show here. We find that \mathcal{K}_{NLO} will give rise to off-diagonal elements of the Kähler metric and they are suppressed by $\langle \Phi \rangle / \Lambda$ in comparison with \mathcal{K}_{LO} .

If the $\Omega(1)$ eclectic model does not contain a flavon Φ transforming as $(\mathbf{8}, \mathbf{8}_k)$ under $\Delta(27) \times T'$, the off-diagonal contributions to the Kähler metric of the matter field ψ could arise from the next-to-next-to-leading order (NNLO) terms which can be written as:⁵

$$\mathcal{K}_{\text{NNLO}} = \sum_{m,n,r_1,r_2,s} (-i\tau + i\bar{\tau})^{-k_\psi - k_\Theta + m} \left(Y_{r_1}^{(m)\dagger} Y_{r_2}^{(n)} \psi^\dagger \psi \Theta^\dagger \Phi \right)_{(\mathbf{10},\mathbf{0},\mathbf{1}),s} + \text{h.c.} . \tag{4.36}$$

We see that two generic flavons Φ and Θ are involved and they could be identical with $\Phi = \Theta$. The total modular weight of each term should be vanishing and consequently the

⁵Depending on the transformation properties of the matter fields and flavons under $\Omega(1)$, $\mathcal{K}_{\text{NNLO}}$ can possibly admit the terms of form $\sum_{m,n,r_1,r_2,s} (-i\tau + i\bar{\tau})^{-k_\psi - k_\Theta + m} \left(Y_{r_1}^{(m)\dagger} Y_{r_2}^{(n)} \psi^\dagger \psi \Theta^\dagger \Phi \right)_{(\mathbf{10},\mathbf{0},\mathbf{1}),s} + \text{h.c.}$, which can be analyzed in a similar way.

modular weight should fulfill the following relation

$$m - n = k_\Theta - k_\Phi. \quad (4.37)$$

The traditional flavor symmetry $\Delta(27)$ requires the combination $\psi^\dagger\psi\Theta^\dagger\Phi$ be a $\Delta(27)$ singlet $\mathbf{1}_{0,0}$, thus we have the contractions

$$\left(\psi^\dagger\psi\Theta^\dagger\Phi\right)_{(\mathbf{1}_{0,0},r)} = \left(\psi^\dagger\psi\right)_{(\mathbf{1}_{0,0},1)} \left(\Theta^\dagger\Phi\right)_{(\mathbf{1}_{0,0},r)} \quad \text{or} \quad \left[\left(\psi^\dagger\psi\right)_{(\mathbf{8},\mathbf{8}_0)} \left(\Theta^\dagger\Phi\right)_{(\mathbf{8},\mathbf{8}_k)}\right]_{(\mathbf{1}_{0,0},r)}, \quad (4.38)$$

which contracts with the modular form $Y_{r_1}^{(m)\dagger}Y_{r_2}^{(n)}$ to form T' invariants, here r are certain representations of T' modular group. As can be seen from the expressions of $\left(\psi^\dagger\psi\right)_{(\mathbf{1}_{0,0},1)}$ and $\left(\psi^\dagger\psi\right)_{(\mathbf{8},\mathbf{8}_0)}$ in eq. (4.32), the first contraction $\left(\psi^\dagger\psi\right)_{(\mathbf{1}_{0,0},1)} \left(\Theta^\dagger\Phi\right)_{(\mathbf{1}_{0,0},r)}$ leads to a diagonal Kähler metric and its contribution can be absorbed in the overall normalization while the second contraction $\left[\left(\psi^\dagger\psi\right)_{(\mathbf{8},\mathbf{8}_0)} \left(\Theta^\dagger\Phi\right)_{(\mathbf{8},\mathbf{8}_k)}\right]_{(\mathbf{1}_{0,0},r)}$ gives rise to an off-diagonal Kähler metric. Moreover, we see that $\Theta^\dagger\Phi$ should be the invariant singlet $\mathbf{1}_{0,0}$ or the reducible octet $\mathbf{8}$ of the $\Delta(27)$ flavor symmetry. The flavons Θ and Φ can transform as $\mathbf{1}_{0,0}$, $\mathbf{3}$, $\bar{\mathbf{3}}$ or $\mathbf{8}$ under $\Delta(27)$, and they can contribute to the Kähler potential $\mathcal{K}_{\text{NNLO}}$ only for the following assignments.

- i) $\Theta \sim \mathbf{1}_{0,0}$, $\Phi \sim \mathbf{1}_{0,0}$

In this case, only the first contraction in eq. (4.38) is allowed, and the resulting Kähler potential is of the same form as the minimal Kähler potential.

- ii) $\Theta, \Phi \sim \mathbf{3}$ or $\Theta, \Phi \sim \bar{\mathbf{3}}$

In terms of the components of the two $\Delta(27)$ triplets $\Theta = (\vartheta_1, \vartheta_2, \vartheta_3)^T$, $\Phi = (\phi_1, \phi_2, \phi_3)^T$, we have

$$\begin{aligned} \left(\Theta^\dagger\Phi\right)_{(\mathbf{1}_{0,0},1^k)} &= \vartheta_1^\dagger\phi_1 + \vartheta_2^\dagger\phi_2 + \vartheta_3^\dagger\phi_3, \\ \left(\Theta^\dagger\Phi\right)_{(\mathbf{8},\mathbf{8}_k)} &= \begin{pmatrix} \vartheta_1^\dagger\phi_2 + \vartheta_2^\dagger\phi_3 + \vartheta_3^\dagger\phi_1 \\ \vartheta_1^\dagger\phi_3 + \vartheta_2^\dagger\phi_1 + \vartheta_3^\dagger\phi_2 \\ \vartheta_1^\dagger\phi_1 + \omega^2\vartheta_2^\dagger\phi_2 + \omega\vartheta_3^\dagger\phi_3 \\ \omega^2\vartheta_1^\dagger\phi_2 + \omega\vartheta_2^\dagger\phi_3 + \vartheta_3^\dagger\phi_1 \\ \omega\vartheta_1^\dagger\phi_3 + \vartheta_2^\dagger\phi_1 + \omega^2\vartheta_3^\dagger\phi_2 \\ \vartheta_1^\dagger\phi_1 + \omega\vartheta_2^\dagger\phi_2 + \omega^2\vartheta_3^\dagger\phi_3 \\ \omega\vartheta_1^\dagger\phi_2 + \omega^2\vartheta_2^\dagger\phi_3 + \vartheta_3^\dagger\phi_1 \\ \omega^2\vartheta_1^\dagger\phi_3 + \vartheta_2^\dagger\phi_1 + \omega\vartheta_3^\dagger\phi_2 \end{pmatrix} \quad \text{for } \Theta, \Phi \sim \mathbf{3}, \end{aligned}$$

$$\left(\Theta^\dagger \Phi \right)_{(\mathbf{8}, \mathbf{8}_k)} = \begin{pmatrix} \vartheta_1^\dagger \phi_3 + \vartheta_2^\dagger \phi_1 + \vartheta_3^\dagger \phi_2 \\ \vartheta_1^\dagger \phi_2 + \vartheta_2^\dagger \phi_3 + \vartheta_3^\dagger \phi_1 \\ \vartheta_1^\dagger \phi_1 + \omega^2 \vartheta_2^\dagger \phi_2 + \omega \vartheta_3^\dagger \phi_3 \\ \vartheta_1^\dagger \phi_3 + \omega^2 \vartheta_2^\dagger \phi_1 + \omega \vartheta_3^\dagger \phi_2 \\ \vartheta_1^\dagger \phi_2 + \omega^2 \vartheta_2^\dagger \phi_3 + \omega \vartheta_3^\dagger \phi_1 \\ \vartheta_1^\dagger \phi_1 + \omega \vartheta_2^\dagger \phi_2 + \omega^2 \vartheta_3^\dagger \phi_3 \\ \vartheta_1^\dagger \phi_3 + \omega \vartheta_2^\dagger \phi_1 + \omega^2 \vartheta_3^\dagger \phi_2 \\ \vartheta_1^\dagger \phi_2 + \omega \vartheta_2^\dagger \phi_3 + \omega^2 \vartheta_3^\dagger \phi_1 \end{pmatrix} \quad \text{for } \Theta, \Phi \sim \bar{\mathbf{3}}. \quad (4.39)$$

Then both singlet and octet contractions in eq. (4.38) are allowed, and the explicit expressions can be obtained from eqs. (4.32), (4.35), (4.39). Obviously these contributions depend on the flavon VEVs and are suppressed by $\langle \Phi \rangle^2 / \Lambda^2$ with respect to the leading order term \mathcal{K}_{LO} .

- iii) $\Theta \sim \mathbf{1}_{0,0}, \Phi \sim \mathbf{8}$ or $\Theta \sim \mathbf{8}, \Phi \sim \mathbf{1}_{0,0}$

Only the octet contractions in eq. (4.38) is allowed in this case, and it leads to off-diagonal Kähler metric of the matter fields.

- iv) $\Theta \sim \mathbf{8}$ and $\Phi \sim \mathbf{8}$

For this assignment, eight independent singlet contractions $\left(\Theta^\dagger \Phi \right)_{(\mathbf{1}_{0,0}, \mathbf{1}^k)}$ and seven independent octet contractions $\left(\Theta^\dagger \Phi \right)_{(\mathbf{8}, \mathbf{8}_k)}$ can be built. They generally induce corrections to the canonical kinetic terms of matter fields.

In short, generally the Kähler potential $\mathcal{K}_{\text{NNLO}}$ can yield deviations from canonical kinetic terms of quark/lepton fields after the flavons develop VEVs, unless all the flavons Θ and Φ are invariant singlet of $\Delta(27)$. However, the induced corrections to the quark/lepton mixing parameters are suppressed by $\langle \Phi \rangle^2 / \Lambda^2$ which are usually small enough to be negligible. In the following example models, we do not introduce a flavon transforming as $(\mathbf{8}, \mathbf{8}_k)$ under $\Delta(27) \rtimes T'$. Hence the corrections to the minimal Kähler potential arise from $\mathcal{K}_{\text{NNLO}}$.

5 $\Omega(1) \cong \Delta(27) \rtimes T'$ eclectic models of lepton mass and mixing

In this section, we shall construct two concrete lepton models based on the eclectic flavor group $\Omega(1) \cong \Delta(27) \rtimes T'$ in the framework of rigid $\mathcal{N} = 1$ global supersymmetry. Then modular invariance requires that the modular weight of each term in the Lagrangian should be vanishing. In the two example models, the neutrinos are assumed to be Majorana particles and their masses are generated through the Weinberg operator and the type I seesaw mechanism, respectively. The model construction is guided by the principle of minimality and simplicity, and consequently we shall introduce as few flavon fields as possible besides the complex modulus τ . The VEVs of flavons spontaneously break the $\Delta(27)$ flavor symmetry, and the breaking of T' modular symmetry can arise from the VEVs of both flavons and modulus τ . Without loss of generality, the VEV of τ is limited in the

Z_3^g	Res. CP	$\langle\phi_{\mathbf{3}}\rangle/v_\phi$	$\langle\phi_{\bar{\mathbf{3}}}\rangle/v_\phi$	$\langle\phi_{\mathbf{8}}\rangle/v_\phi$
Z_3^A	—	$(1, 1, 1)^T$	$(1, 1, 1)^T$	$v_{8_1} = (1, x, 0, 0, 0, 0, 0, 0)^T$
Z_3^B	A	$(1, 0, 0)^T$	$(1, 0, 0)^T$	$v_{8_2} = (0, 0, \omega^2, 0, 0, -\omega x, 0, 0)^T$
	AB	$(\omega^2, 0, 0)^T$	$(\omega, 0, 0)^T$	v_{8_2}
	BAB^2	$(\omega, 0, 0)^T$	$(\omega^2, 0, 0)^T$	v_{8_2}
Z_3^{AB}	$BAB^2 A^2$	$(1, \omega^2, 1)^T$	$(1, \omega, 1)^T$	$v_{8_3} = (0, 0, 0, 0, 1 - ix, 0, 1 + ix, 0)^T$
	$ABA^2 B^2$	$\omega(1, \omega^2, 1)^T$	$\omega^2(1, \omega, 1)^T$	v_{8_3}
	AB	$\omega^2(1, \omega^2, 1)^T$	$\omega(1, \omega, 1)^T$	v_{8_3}
Z_3^{BA}	$BAB^2 A^2$	$\omega(1, 1, \omega^2)^T$	$\omega^2(1, 1, \omega)^T$	v_{8_3}
	$ABA^2 B^2$	$\omega^2(1, 1, \omega^2)^T$	$\omega(1, 1, \omega)^T$	v_{8_3}
	BA	$(1, 1, \omega^2)^T$	$(1, 1, \omega)^T$	v_{8_3}
Z_3^{ABA}	—	$(1, \omega^2, \omega^2)^T$	$(1, \omega, \omega)^T$	$v_{8_4} = (0, 0, 0, 1, 0, 0, 0, x)^T$
$Z_3^{A^2 B}$	—	$(1, 1, \omega)^T$	$(1, 1, \omega^2)^T$	v_{8_4}
$Z_3^{BA^2}$	—	$(1, \omega, 1)^T$	$(1, \omega^2, 1)^T$	v_{8_4}
$Z_3^{BAB^2}$	—	$(1, \omega, \omega^2)^T$	$(1, \omega^2, \omega)^T$	v_{8_1}
$Z_3^{B^2 AB}$	—	$(1, \omega^2, \omega)^T$	$(1, \omega, \omega^2)^T$	v_{8_1}
$Z_3^{A^2 BA}$	A^2	$(0, 1, 0)^T$	$(0, 1, 0)^T$	$v_{8_2}^*$
	$B^2 A^2 B$	$\omega(0, 1, 0)^T$	$\omega^2(0, 1, 0)^T$	$v_{8_2}^*$
	$BA^2 B^2$	$\omega^2(0, 1, 0)^T$	$\omega(0, 1, 0)^T$	$v_{8_2}^*$
$Z_3^{ABA^2}$	$BAB^2 A^2$	$(0, 0, 1)^T$	$(0, 0, 1)^T$	$v_{8_5} = (0, 0, 1, 0, 0, x, 0, 0)^T$
	$ABA^2 B^2$	$\omega(0, 0, 1)^T$	$\omega^2(0, 0, 1)^T$	v_{8_5}
	$AB^2 A^2$	$\omega^2(0, 0, 1)^T$	$\omega(0, 0, 1)^T$	v_{8_5}
$Z_3^{A^2 BA^2}$	$BAB^2 A^2$	$(\omega^2, 1, 1)^T$	$(\omega, 1, 1)^T$	v_{8_3}
	$ABA^2 B^2$	$\omega(\omega^2, 1, 1)^T$	$\omega^2(\omega, 1, 1)^T$	v_{8_3}
	$A^2 BA^2$	$\omega^2(\omega^2, 1, 1)^T$	$\omega(\omega, 1, 1)^T$	v_{8_3}
$Z_3^{BAB^2 A^2}$	—	—	—	general 8-D vector

Table 3. The most general vacuum alignment invariants under the Z_3 subgroups of $\Delta(27)$ and residual CP symmetry, where $\phi_{\mathbf{3}}$, $\phi_{\bar{\mathbf{3}}}$ and $\phi_{\mathbf{8}}$ denote the flavons which transform as $\mathbf{3}$, $\bar{\mathbf{3}}$ and reducible $\mathbf{8}$ respectively under $\Delta(27)$, and x is a real free parameter. The residual CP symmetry in the second column refers to $\rho(g)\rho(K_*)$ with $g \in \Delta(27)$, and we only present the element g for simplicity. The overall VEV v_ϕ is constrained to be real by the residual CP symmetry.

fundamental domain \mathcal{D} of $\text{SL}(2, \mathbb{Z})$:

$$\mathcal{D} = \{ \tau \mid |\Re\tau| \leq 1/2, \Im\tau > 0, |\tau| \geq 1 \} . \quad (5.1)$$

Other values of τ are related by modular transformations to those in \mathcal{D} . In both example models, the Higgs doublets $H_{u,d}$ are invariant under both $\Delta(27)$ and T' with vanishing modular weights. The flavon fields transform as trivial singlet $\mathbf{1}_{0,0}$ or triplet $\mathbf{3}, \bar{\mathbf{3}}$ under $\Delta(27)$, and we don't introduce any flavon in nontrivial singlet representation of $\Delta(27)$ otherwise there would be a flavon octant, as explained in section 3. Extensive studies of the discrete flavor symmetry approach to the flavor problem have revealed that the flavon fields have to develop a set of particularly VEVs [1]. Hence we will consider the vacuum configuration of flavon which preserve certain subgroup of $\Delta(27)$ and the most general invariant VEVs of all Z_3 subgroups of $\Delta(27)$ are given in table 3. It is known that rather elaborate scalar potentials of flavons should be designed to get the correct vacuum alignment in traditional discrete flavor symmetry approach. On the other hand, it is still an open question to determine the VEV of τ from first principles in modular flavor symmetry, this is the so-called modulus stabilisation problem [39, 40]. The presence of both flavons and complex modulus in eclectic flavor symmetry exacerbates the vacuum selection problem mentioned above. Hence we will not attempt to build the most general supersymmetric and eclectic symmetry invariant scalar potential for the modulus τ and flavons. Both τ and flavons will be treated as spurions, and the VEV of τ is freely varied in the fundamental domain to adjust the agreement with the data.

In previous section, we have performed a general analysis for the $\Omega(1)$ invariant superpotential and Kähler potential. Given the representation assignments of the lepton fields and flavons in the two example models, we find that the nontrivial corrections to the minimal Kähler potential arise at the NNLO. The corresponding contributions to the lepton masses and mixing parameters are suppressed by $\langle \Phi \rangle^2 / \Lambda^2$ such that they are negligible. We shall present the two example models based on the eclectic flavor symmetry $\Omega(1) \cong \Delta(27) \times T'$ in the following.

5.1 Example model 1

The light neutrino masses are generated by the Weinberg operator in this model. All the fields and their classifications under the traditional flavor group $\Delta(27)$ and the finite modular group T' are summarized in table 4. The LH leptons L , RH charged leptons E^c , the flavon ϕ in the charged lepton sector and the flavons φ and χ in the neutrino sector are all assumed to be triplet $\mathbf{3}$ of $\Delta(27)$. Then these fields must transform as reducible three-dimensional representations $\mathbf{3}_0, \mathbf{3}_1$ or $\mathbf{3}_2$, and the corresponding representation matrices of generators S and T are given in eq. (3.23). Furthermore, one $\Delta(27)$ trivial singlet flavon ξ is introduced in the model. In order to eliminate the unwanted operators, we introduce the auxiliary symmetry $Z_2 \times Z_3$ in the model. Then we can read out the superpotential for the charged lepton and neutrino masses, which is invariant under the symmetry of the model,

$$\begin{aligned} \mathcal{W} = & \frac{\alpha}{\Lambda} \left(E^c L \phi Y_{\mathbf{2}'}^{(5)} \right)_{(\mathbf{1}_{0,0,1})} H_d + \frac{\beta}{\Lambda^2} \left(E^c L \xi \phi Y_{\mathbf{1}}^{(4)} \right)_{(\mathbf{1}_{0,0,1})} H_d \\ & + \frac{g_1}{2\Lambda^2} \left(LL \varphi Y_{\mathbf{2}''}^{(5)} \right)_{(\mathbf{1}_{0,0,1})} H_u H_u + \frac{g_2}{2\Lambda^2} \left(LL \chi Y_{\mathbf{2}'}^{(7)} \right)_{(\mathbf{1}_{0,0,1})} H_u H_u . \end{aligned} \quad (5.2)$$

Fields	L	E^c	H_u	H_d	ϕ	φ	χ	ξ	$Y_r^{(k_Y)}$
$SU(2)_L \times U(1)_Y$	$(\mathbf{2}, -\frac{1}{2})$	$(\mathbf{1}, 1)$	$(\mathbf{2}, \frac{1}{2})$	$(\mathbf{2}, -\frac{1}{2})$	$(\mathbf{1}, 0)$	$(\mathbf{1}, 0)$	$(\mathbf{1}, 0)$	$(\mathbf{1}, 0)$	$(\mathbf{1}, 0)$
$\Delta(27)$	$\mathbf{3}$	$\mathbf{3}$	$\mathbf{1}_{0,0}$	$\mathbf{1}_{0,0}$	$\mathbf{3}$	$\mathbf{3}$	$\mathbf{3}$	$\mathbf{1}_{0,0}$	$\mathbf{1}_{0,0}$
$\Gamma'_3 \cong T'$	$\mathbf{3}_0$	$\mathbf{3}_0$	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{3}_1$	$\mathbf{3}_0$	$\mathbf{3}_1$	$\mathbf{1}$	\mathbf{r}
modular weight	0	0	0	0	5	5	7	-1	k_Y
Z_2	1	-1	1	1	-1	1	1	1	1
Z_3	ω	ω^2	1	1	1	ω	ω	1	1

Table 4. The matter fields, flavon fields of the example model 1 and their transformation properties under $\Delta(27)$ and T' , where $Y_r^{(k_Y)}$ refers to the level 3 and weight k_Y modular multiplet in the irreducible representation \mathbf{r} of T' .

The first and second rows of eq. (5.2) are the mass terms of charged leptons and neutrinos, respectively. The charged lepton superpotential is the case of \mathcal{W}_{D4} in table 1, and the two neutrino mass terms correspond to \mathcal{W}_{M1} and \mathcal{W}_{M2} in table 2, respectively. Notice that the second term contains two flavons and it is suppressed by $\langle \xi \rangle / \Lambda$ with respect to the first term. We assume that the traditional flavor symmetry $\Delta(27)$ is broken down to $Z_3^{A^2 B A^2}$ in the charged lepton sector by the VEV of ϕ , and the VEVs of the flavons φ and χ entering in neutrino sector are invariant under $Z_3^{A B A^2}$ and Z_3^B , respectively. As a consequence, the residual traditional flavor symmetry $Z_3^{A^2 B A^2}$ is preserved in the charged lepton sector after symmetry breaking, while the $\Delta(27)$ flavor symmetry is completely broken in the neutrino sector.⁶ As shown in table 3, the assumed residual symmetry allows us to fix VEVs of flavons ξ , ϕ , φ and χ as follows,

$$\langle \xi \rangle = v_\xi, \quad \langle \phi \rangle = (\omega^2, 1, 1)^T v_\phi, \quad \langle \varphi \rangle = (0, 0, 1)^T v_\varphi, \quad \langle \chi \rangle = (1, 0, 0)^T v_\chi, \quad (5.3)$$

where v_ξ , v_ϕ , v_φ and v_χ are undetermined and generally complex. After the electroweak and flavor symmetry breaking in the way of eq. (5.3), we obtain the charged lepton mass matrix and the light neutrino mass matrix as follows

$$m_l = \frac{\alpha v_\phi v_d}{\Lambda} \begin{pmatrix} \sqrt{2}\omega^2 Y_{2',1}^{(5)} & \omega Y_{2',2}^{(5)} & \omega Y_{2',2}^{(5)} \\ \omega Y_{2',2}^{(5)} & \sqrt{2}Y_{2',1}^{(5)} & Y_{2',2}^{(5)} \\ \omega Y_{2',2}^{(5)} & Y_{2',2}^{(5)} & \sqrt{2}Y_{2',1}^{(5)} \end{pmatrix} + \frac{i\beta Y_{\mathbf{1}}^{(4)} v_\xi v_\phi v_d}{\Lambda^2} \begin{pmatrix} 0 & \omega & -\omega \\ -\omega & 0 & 1 \\ \omega & -1 & 0 \end{pmatrix},$$

$$m_\nu = \frac{g_1 v_\varphi v_u^2}{\Lambda^2} \begin{pmatrix} 0 & \omega Y_{2'',2}^{(5)} & 0 \\ \omega Y_{2'',2}^{(5)} & 0 & 0 \\ 0 & 0 & \sqrt{2}Y_{2'',1}^{(5)} \end{pmatrix} + \frac{g_2 v_\chi v_u^2}{\Lambda^2} \begin{pmatrix} \sqrt{2}Y_{2',1}^{(7)} & 0 & 0 \\ 0 & 0 & \omega Y_{2',2}^{(7)} \\ 0 & \omega Y_{2',2}^{(7)} & 0 \end{pmatrix}, \quad (5.4)$$

where we use $Y_{r,i}^{(k_Y)}$ to represent the i th component of modular multiple $Y_r^{(k_Y)}$. Obviously we have $\rho_3^\dagger(A^2 B A^2) m_l^\dagger m_l \rho_3(A^2 B A^2) = m_l^\dagger m_l$ which implies that the charged lepton mass ma-

⁶The residual symmetry of the Majorana neutrino mass matrix can only be Z_2 or K_4 for three nonvanishing light neutrino masses [41, 42]. Since the $\Delta(27)$ flavor group does not have Z_2 and K_4 subgroups, no residual flavor symmetry should be preserved in the neutrino sector.

trix is invariant under the residual flavor symmetry $Z_3^{A^2BA^2}$. Therefore the unitary transformation U_l diagonalizing the hermitian combination $m_l^\dagger m_l$ via $U_l^\dagger m_l^\dagger m_l U_l = \text{diag}(m_e^2, m_\mu^2, m_\tau^2)$ is of the form:

$$U_l = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & \omega^2 & \omega \\ \omega^2 & \omega^2 & \omega^2 \\ 1 & \omega & \omega^2 \end{pmatrix}. \quad (5.5)$$

The charged lepton masses are given by

$$\begin{aligned} m_e &= \left| \sqrt{2}Y_{2',1}^{(5)} - Y_{2',2}^{(5)} - \frac{\sqrt{6}\beta v_\xi Y_1^{(4)}}{\alpha\Lambda} \right| \frac{\alpha v_\phi v_d}{\Lambda}, \\ m_\mu &= \left| \sqrt{2}Y_{2',1}^{(5)} - Y_{2',2}^{(5)} + \frac{\sqrt{6}\beta v_\xi Y_1^{(4)}}{\alpha\Lambda} \right| \frac{\alpha v_\phi v_d}{\Lambda}, \\ m_\tau &= \left| \sqrt{2}Y_{2',1}^{(5)} + 2Y_{2',2}^{(5)} \right| \frac{\alpha v_\phi v_d}{\Lambda}. \end{aligned} \quad (5.6)$$

Since L and E^c are assigned to be triplet of $\Delta(27)$, both couplings α and β are relevant to the three charged lepton masses. Consequently some fine-tuning in the values of τ and $\beta v_\xi/(\alpha\Lambda)$ is necessary to accommodate charged lepton mass hierarchies. We transfer to the charged lepton diagonal basis by redefining the LH lepton fields $L \rightarrow U_l L$, then the light neutrino mass matrix is given by

$$\begin{aligned} m'_\nu &= U_l^T m_\nu U_l \\ &= \frac{g_1 v_\phi v_u^2}{\Lambda^2} \begin{pmatrix} \sqrt{2}Y_{2'',1}^{(5)} + 2Y_{2'',2}^{(5)} & \omega \left(\sqrt{2}Y_{2'',1}^{(5)} - Y_{2'',2}^{(5)} \right) & \omega^2 \left(\sqrt{2}Y_{2'',1}^{(5)} - Y_{2'',2}^{(5)} \right) \\ \omega \left(\sqrt{2}Y_{2'',1}^{(5)} - Y_{2'',2}^{(5)} \right) & \omega^2 \left(\sqrt{2}Y_{2'',1}^{(5)} + 2Y_{2'',2}^{(5)} \right) & \sqrt{2}Y_{2'',1}^{(5)} - Y_{2'',2}^{(5)} \\ \omega^2 \left(\sqrt{2}Y_{2'',1}^{(5)} - Y_{2'',2}^{(5)} \right) & \sqrt{2}Y_{2'',1}^{(5)} - Y_{2'',2}^{(5)} & \omega \left(\sqrt{2}Y_{2'',1}^{(5)} + 2Y_{2'',2}^{(5)} \right) \end{pmatrix} \\ &+ \frac{g_2 v_\chi v_u^2}{\Lambda^2} \begin{pmatrix} \sqrt{2}Y_{2',1}^{(7)} + 2Y_{2',2}^{(7)} & \omega^2 \left(\sqrt{2}Y_{2',1}^{(7)} - Y_{2',2}^{(7)} \right) & \omega \left(\sqrt{2}Y_{2',1}^{(7)} - Y_{2',2}^{(7)} \right) \\ \omega^2 \left(\sqrt{2}Y_{2',1}^{(7)} - Y_{2',2}^{(7)} \right) & \omega \left(\sqrt{2}Y_{2',1}^{(7)} + 2Y_{2',2}^{(7)} \right) & \sqrt{2}Y_{2',1}^{(7)} - Y_{2',2}^{(7)} \\ \omega \left(\sqrt{2}Y_{2',1}^{(7)} - Y_{2',2}^{(7)} \right) & \sqrt{2}Y_{2',1}^{(7)} - Y_{2',2}^{(7)} & \omega^2 \left(\sqrt{2}Y_{2',1}^{(7)} + 2Y_{2',2}^{(7)} \right) \end{pmatrix}. \end{aligned} \quad (5.7)$$

Let us proceed to perform a numerical analysis for the model. We first consider the scenario without gCP. This model involves 6 real free parameters besides the complex modulus τ . All the six lepton masses, three lepton mixing angles and three CP violation phases rely on the following six dimensionless real parameters

$$\Re\tau, \quad \Im\tau, \quad |\beta v_\xi/(\alpha\Lambda)|, \quad \arg(\beta v_\xi/(\alpha\Lambda)), \quad |g_2 v_\chi/(g_1 v_\phi)|, \quad \arg(g_2 v_\chi/(g_1 v_\phi)), \quad (5.8)$$

and two overall scales $\alpha v_d v_\phi/\Lambda$ and $g_1 v_u^2 v_\phi/\Lambda^2$ which can be taken to real. In order to quantitatively estimate how well the model can describe the experimental data on the following seven dimensionless observable quantities

$$m_e/m_\mu, \quad m_\mu/m_\tau, \quad \sin^2 \theta_{12}, \quad \sin^2 \theta_{13}, \quad \sin^2 \theta_{23}, \quad \delta_{CP}, \quad \Delta m_{21}^2/\Delta m_{31}^2, \quad (5.9)$$

which only depend on the six dimensionless real parameters in eq. (5.8), we perform a conventional χ^2 analysis to determine the best fit values of the free parameters as well

as the corresponding predictions for the flavor observables. We adopt the following the experimental data and errors to build the χ^2 function:

$$\begin{aligned}
 \frac{m_e}{m_\mu}\Big|_{\text{bf}+1\sigma} &= 0.004737_{-0.00004}^{+0.00004}, & \frac{m_\mu}{m_\tau}\Big|_{\text{bf}+1\sigma} &= 0.05876_{-0.000465}^{+0.000465}, & \sin^2\theta_{12}\Big|_{\text{bf}+1\sigma} &= 0.303_{-0.012}^{+0.012}, \\
 \sin^2\theta_{13}\Big|_{\text{bf}+1\sigma} &= 0.02225_{-0.00059}^{+0.00056}, & \sin^2\theta_{23}\Big|_{\text{bf}+1\sigma} &= 0.451_{-0.016}^{+0.019}, \\
 \frac{\delta_{CP}}{\pi}\Big|_{\text{bf}+1\sigma} &= 1.289_{-0.14}^{+0.20}, & \frac{\Delta m_{21}^2}{\Delta m_{31}^2}\Big|_{\text{bf}+1\sigma} &= 0.0294_{-0.00088}^{+0.00088},
 \end{aligned} \tag{5.10}$$

where the data of the lepton mixing angles and CP phase δ_{CP} is taken from NuFIT5.2 [43], the charged lepton mass ratios are taken from [44] with $M_{\text{SUSY}} = 10$ TeV and $\tan\beta = 5$. We notice that this model yields a successful fit to current experimental data for certain values of the input parameters. We find the minimum value of the χ^2 function is $\chi^2_{\text{min}} = 10.68$, and the best fit values of the free parameters are

$$\begin{aligned}
 \Re\langle\tau\rangle &= 0.00157, & \Im\langle\tau\rangle &= 1.120, & |\beta v_\xi/(\alpha\Lambda)| &= 0.0478, & \arg(\beta v_\xi/(\alpha\Lambda)) &= 1.036\pi, \\
 |g_2 v_\chi/(g_1 v_\varphi)| &= 0.9786, & \arg(g_2 v_\chi/(g_1 v_\varphi)) &= 1.005\pi, \\
 \alpha v_\xi v_\phi v_d/\Lambda^2 &= 267.2 \text{ MeV}, & g_1 v_\varphi v_u^2/\Lambda^2 &= 5.40 \text{ meV}.
 \end{aligned} \tag{5.11}$$

The overall scales $\alpha v_d v_\phi/\Lambda$ and $g_1 v_u^2 v_\varphi/\Lambda^2$ are fixed by the measured electron mass and the solar neutrino mass square difference Δm_{21}^2 . We see that the real part of modulus $\Re\langle\tau\rangle$ is quite small, and both phases $\arg(\beta v_\xi/(\alpha\Lambda))$ and $\arg(g_2 v_\chi/(g_1 v_\varphi))$ are very close to π . At the best fit point, the lepton masses and mixing parameters are determined to be

$$\begin{aligned}
 \sin^2\theta_{13} &= 0.02252, & \sin^2\theta_{12} &= 0.3283, & \sin^2\theta_{23} &= 0.4954, & \delta_{CP} &= 1.435\pi, \\
 \alpha_{21} &= 0.961\pi, & \alpha_{31} &= 0.927\pi, & m_1 &= 15.12 \text{ meV}, & m_2 &= 17.40 \text{ meV}, \\
 m_3 &= 52.30 \text{ meV}, & \sum_{i=1}^3 m_i &= 84.82 \text{ meV}, & m_{\beta\beta} &= 5.617 \text{ meV}, \\
 m_e &= 0.511 \text{ MeV}, & m_\mu &= 107.9 \text{ MeV}, & m_\tau &= 1.836 \text{ GeV},
 \end{aligned} \tag{5.12}$$

where m_i ($i = 1, 2, 3$), m_j ($j = e, \mu, \tau$) and $m_{\beta\beta}$ refers to the absolute masses of the three light neutrinos, the masses of the three charged leptons and the effective mass in neutrinoless double beta decay, respectively. The effective Majorana mass is determined to be $m_{\beta\beta} = 5.617$ meV which is much below the current most stringent limit $m_{\beta\beta} < 36 - 156$ meV from KamLAND-Zen [45]. The next generation neutrinoless double decay experiments of ton-scale can improve the sensitivity to $m_{\beta\beta}$ considerably. For example, the SNO+ Phase II could probe the effective mass $m_{\beta\beta}$ in the range of 19 – 46 meV [46]. The LEGEND experiment is expected to reach the sensitivity of (15–50) meV by operating 1000 kg of detectors for 10 years [47]. The experiment nEXO will be able to probe $m_{\beta\beta}$ down to 5.7 – 17.7 meV after 10 years of data taking [48]. Thus our prediction for $m_{\beta\beta}$ is a bit below but close to the future sensitivity. Moreover, the three neutrino mass sum is predicted to be 84.82 meV which is compatible with the upper bound $\sum_{i=1}^3 m_i < 120$ meV on neutrino mass sum from Planck [49].

If the gCP symmetry is imposed on the model, all the couplings would be enforced to be real, as discussed in section 4. Moreover, certain residual CP symmetry require the flavons vacuum parameters v_ξ , v_ϕ , v_φ and v_χ in eq. (5.3) be real, as shown in table 3. Thus the two phase parameters $\arg(\beta v_\xi/(\alpha\Lambda))$ and $\arg(g_2 v_\chi/(g_1 v_\varphi))$ in eq. (5.8) are 0 or π because of the gCP symmetry. Furthermore, if the modulus τ lies on the imaginary axis with $\Re\tau = 0$, the VEV of τ preserves the gCP symmetry K_* , and $q = e^{2\pi i\tau}$ as well as the modular forms $Y_1(\tau)$ and $Y_2(\tau)$ in eq. (C.9) would be real. Hence all the integer weight modular forms of level 3 are real for pure imaginary τ , as can be seen from appendix C. As a consequence, the neutrino mass matrix m'_ν in the charged lepton diagonal basis in eq. (5.7) satisfies

$$(m'_\nu)_{ee} = (m'_\nu)_{ee}^*, \quad (m'_\nu)_{\mu\tau} = (m'_\nu)_{\mu\tau}^*, \quad (m'_\nu)_{e\tau} = (m'_\nu)_{e\mu}^*, \quad (m'_\nu)_{\mu\mu} = (m'_\nu)_{\tau\tau}^*, \quad (5.13)$$

which is exactly the requirement of a theory invariant under $\mu - \tau$ reflection symmetry [50–52]. In other words, the neutrino mass matrix m'_ν in eq. (5.7) really fulfills the $\mu - \tau$ reflection symmetry

$$P_{\nu\tau}^T m'_\nu P_{\nu\tau} = (m'_\nu)^*, \quad (5.14)$$

with

$$P_{\nu\tau} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}. \quad (5.15)$$

It is known that the $\mu - \tau$ reflection symmetry leads to maximal atmospheric angle, maximal Dirac CP phase and trivial Majorana CP phases: $\theta_{23} = \pi/4$, $\delta_{CP} = \pm\pi/2$, $\alpha_{21}, \alpha_{31} = 0, \pi$ [50–52]. After imposing the gCP symmetry and the assumption of $\Re\tau = 0$, we find that all the seven dimensionless flavor observables in eq. (5.9) would only depend on three dimensionless real parameters in the model,

$$\Im\tau, \quad \beta v_\xi/(\alpha\Lambda), \quad g_2 v_\chi/(g_1 v_\varphi). \quad (5.16)$$

Analogously performing a numerical analysis, we find that the minimum value of χ^2 function is $\chi_{\min}^2 = 11.75$ when the free parameters take the following values

$$\begin{aligned} \langle\tau\rangle &= 1.120i, & \beta v_\xi/(\alpha\Lambda) &= -0.0482, & g_2 v_\chi/(g_1 v_\varphi) &= -0.9804, \\ \alpha v_d v_\xi v_\phi/\Lambda^2 &= 267.3 \text{ MeV}, & g_1 v_u^2 v_\varphi/\Lambda^2 &= 5.409 \text{ meV}. \end{aligned} \quad (5.17)$$

Accordingly the lepton masses and mixing parameters are determined to be

$$\begin{aligned} \sin^2 \theta_{13} &= 0.02239, & \sin^2 \theta_{12} &= 0.3266, & \sin^2 \theta_{23} &= 0.5, & \delta_{CP} &= 1.5\pi, \\ \alpha_{21} &= \pi, & \alpha_{31} &= \pi, & m_1 &= 15.17 \text{ meV}, & m_2 &= 17.44 \text{ meV}, \\ m_3 &= 52.42 \text{ meV}, & \sum_i m_i &= 85.03 \text{ meV}, & m_{\beta\beta} &= 5.595 \text{ meV}, \\ m_e &= 0.511 \text{ MeV}, & m_\mu &= 107.9 \text{ MeV}, & m_\tau &= 1.836 \text{ GeV}. \end{aligned} \quad (5.18)$$

In order to show the viability and predictions of this model with/without gCP, we scan over the parameter space of the model, the six (three) dimensionless real parameters in




Figure 1. The predicted correlations among the input free parameters, neutrino mixing angles and CP violation phases in the example model 1. The green and red regions denote points compatible with experimental data for models without gCP and with gCP plus $\Re\tau = 0$, respectively. The light blue bounds represent the 1σ and 3σ ranges respectively with SK atmospheric data [43]. The orange (blue) dashed lines in the last panel represent the most general allowed regions for NO (IO) neutrino masses respectively, where the neutrino oscillation parameter are varied within their 3σ ranges. The present upper limit $m_{\beta\beta} < (36 - 156)$ meV from KamLAND-Zen [45] is shown by horizontal grey band. The vertical grey exclusion band denotes the current bound from the cosmological data of $\sum_i m_i < 0.120$ eV at 95% confidence level given by the Planck collaboration [49]. Notice that we have sharp prediction for the effective mass $m_{\beta\beta}$, the green region (without gCP) almost coincides with the red region (with gCP) in the last panel.

eq. (5.8) (in eq. (5.16)) are taken as random numbers. Then we can evaluate the predictions for lepton masses and mixing parameters and only keep the points for which the flavor observables are within their experimentally allowed 3σ regions. Some interesting correlations among the observables and free parameters are obtained. The results are shown in figure 1. Here the results without gCP and with gCP plus $\Re\tau = 0$ are displayed in green and red, respectively. In the case that gCP is considered with $\Re\tau = 0$, the $\mu - \tau$ reflection symmetry is reproduced, the predictions for θ_{23} , δ_{CP} , α_{21} and α_{31} are labelled with a red pentagram. From figure 1, we see that all the mixing parameters and $m_{\beta\beta}$ are predicted to lie in narrow ranges:

$$\begin{aligned}
 \sin^2 \theta_{13} &\in [0.02172, 0.02398], & \sin^2 \theta_{12} &\in [0.319, 0.341], & \sin^2 \theta_{23} &\in [0.486, 0.514], \\
 \delta_{CP} &\in [1.31\pi, 1.69\pi], & \alpha_{21} &\in [0.881\pi, 1.119\pi], & \alpha_{31} &\in [0.778, 1.218], \\
 m_{\beta\beta} &\in [5.328 \text{ meV}, 6.109 \text{ meV}], & & & &
 \end{aligned}
 \tag{5.19}$$

Fields	L	E^c	N^c	H_u	H_d	ϕ	φ	ξ	$Y_r^{(k_Y)}$
$SU(2)_L \times U(1)_Y$	$(\mathbf{2}, -\frac{1}{2})$	$(\mathbf{1}, 1)$	$(\mathbf{1}, 0)$	$(\mathbf{2}, \frac{1}{2})$	$(\mathbf{2}, -\frac{1}{2})$	$(\mathbf{1}, 0)$	$(\mathbf{1}, 0)$	$(\mathbf{1}, 0)$	$(\mathbf{1}, 0)$
$\Delta(27)$	$\bar{\mathbf{3}}$	$\bar{\mathbf{3}}$	$\bar{\mathbf{3}}$	$\mathbf{1}_{0,0}$	$\mathbf{1}_{0,0}$	$\bar{\mathbf{3}}$	$\bar{\mathbf{3}}$	$\mathbf{1}_{0,0}$	$\mathbf{1}_{0,0}$
$\Gamma'_3 \cong T'$	$\bar{\mathbf{3}}_0$	$\bar{\mathbf{3}}_0$	$\bar{\mathbf{3}}_0$	$\mathbf{1}$	$\mathbf{1}$	$\bar{\mathbf{3}}_0$	$\bar{\mathbf{3}}_1$	$\mathbf{1}''$	\mathbf{r}
modular weight	0	0	1	0	0	3	-1	7	k_Y
Z_2	1	1	-1	1	1	1	-1	1	1
Z_3	ω	ω	ω	1	1	ω	ω	1	1

Table 5. The matter fields, flavon fields of example model 2 and their transformation properties under the eclectic flavor symmetry $\Delta(27) \times T'$ and the auxiliary symmetry $Z_2 \times Z_3$.

for the case without gCP, and

$$\begin{aligned} \sin^2 \theta_{23} &= 0.5, & \delta_{CP} &= 1.5\pi, & \alpha_{21} &= \alpha_{31} = \pi, \\ \sin^2 \theta_{13} &\in [0.02172, 0.02282], & \sin^2 \theta_{12} &\in [0.324, 0.331], & m_{\beta\beta} &\in [5.328 \text{ meV}, 5.776 \text{ meV}], \end{aligned} \tag{5.20}$$

for the case with gCP plus $\Re\tau = 0$. Furthermore, this model predicts a nearly maximal atmospheric mixing angle θ_{23} and a nearly maximal Dirac CP phase δ_{CP} when gCP is not imposed. Notice that the model predicts a nearly maximal atmospheric mixing angle θ_{23} even if gCP isn't imposed. It is remarkable that the variation region of the effective Majorana neutrino mass $m_{\beta\beta}$ is very small because of the linear correlation between α_{21} and α_{31} shown in figure 1.

The NLO correction to the minimal Kähler potential is of the form

$$\begin{aligned} \mathcal{K}_{\text{NLO}} &= \sum_{n,r} \frac{1}{\Lambda} (-i\tau + i\bar{\tau})^n \left[Y_r^{(n)\dagger}(\tau) Y_r^{(n-1)}(\tau) L^\dagger L \xi \right]_{(\mathbf{1}_{0,0,1})} \\ &\quad + \frac{1}{\Lambda} (-i\tau + i\bar{\tau})^n \left[Y_r^{(n)\dagger}(\tau) Y_r^{(n-1)}(\tau) E^{c\dagger} E^c \xi \right]_{(\mathbf{1}_{0,0,1})}. \end{aligned} \tag{5.21}$$

The corresponding Kähler metric for both LH lepton doublet L and RH charged leptons E^c is a unit matrix so that its contribution can be absorbed into the minimal Kähler potential. Hence the non-diagonal contributions of Kähler metric come from NNLO corrections and they are suppressed by $\langle \Phi \rangle^2 / \Lambda^2$. Therefore the contributions of Kähler potential to the lepton mixing parameters and masses are suppressed by $\langle \Phi \rangle^2 / \Lambda^2$ and can be neglected.

5.2 Example model 2

In the second model, the neutrino masses are described by the type I seesaw mechanism with three RH neutrinos. This model is also based on the eclectic flavor group $\Omega(1)$ with the extra symmetry $Z_2 \times Z_3$ to eliminate undesired operators. The three generations of LH doublet leptons L , the RH charged leptons E^c and the RH neutrinos N^c are all embedded into the triplet $\bar{\mathbf{3}}$ of $\Delta(27)$, and we introduce three flavons ϕ , φ and ξ which transform as $\bar{\mathbf{3}}$, $\bar{\mathbf{3}}$ and $\mathbf{1}_{0,0}$ under $\Delta(27)$ respectively.⁷ The fields content and their classification under the

⁷Similar results can be obtained if the representation $\mathbf{3}$ instead of $\bar{\mathbf{3}}$ is used.

eclectic flavor group $\Omega(1)$ and the auxiliary symmetry $Z_2 \times Z_3$ are listed in table 5. The superpotential for the leptons can be written as

$$\begin{aligned} \mathcal{W} = & \frac{\alpha}{\Lambda} \left(E^c L \phi Y_{\mathbf{2}' }^{(3)} \right)_{(10,0,1)} H_d + \frac{\beta}{\Lambda^2} \left(E^c L \phi \xi Y_{\mathbf{1} }^{(10)} \right)_{(10,0,1)} H_d + \frac{g_1}{\Lambda} (N^c L \varphi)_{(10,0,1)} H_u \\ & + \frac{g_2}{\Lambda^2} \left(N^c L \varphi \xi Y_{\mathbf{2}' }^{(7)} \right)_{(10,0,1)} H_u + \frac{h}{2} \left(N^c N^c \phi Y_{\mathbf{2}' }^{(5)} \right)_{(10,0,1)}. \end{aligned} \quad (5.22)$$

In this model we assume that the VEV of flavon ϕ breaks the traditional flavor symmetry $\Delta(27)$ down to $Z_3^{A^2B}$ while the subgroup Z_3^A is preserved by vacuum of flavon φ . Hence the VEVs of flavons ξ , ϕ and φ take the following form:

$$\langle \xi \rangle = v_\xi, \quad \langle \phi \rangle = (1, 1, \omega^2)^T v_\phi, \quad \langle \varphi \rangle = (1, 1, 1)^T v_\varphi. \quad (5.23)$$

From table 3, we see that no nontrivial residual CP transformation is consistent with the residual flavor symmetries $Z_3^{A^2B}$ and Z_3^A . Therefore we do not consider gCP symmetry in this model, thus all coupling constants and the VEVs v_ξ , v_ϕ , v_φ are generally complex. After flavor symmetry and electroweak symmetry breaking, we can straightforwardly read out the charged lepton mass matrix, the Dirac neutrino mass matrix and the Majorana neutrino mass matrix,

$$\begin{aligned} m_l &= \frac{\alpha v_d v_\phi}{\Lambda} \begin{pmatrix} \sqrt{2} Y_{\mathbf{2}',2}^{(3)} & -\omega Y_{\mathbf{2}',1}^{(3)} & -\omega^2 Y_{\mathbf{2}',1}^{(3)} \\ -\omega Y_{\mathbf{2}',1}^{(3)} & \sqrt{2} Y_{\mathbf{2}',2}^{(3)} & -\omega^2 Y_{\mathbf{2}',1}^{(3)} \\ -\omega^2 Y_{\mathbf{2}',1}^{(3)} & -\omega^2 Y_{\mathbf{2}',1}^{(3)} & \sqrt{2} \omega^2 Y_{\mathbf{2}',2}^{(3)} \end{pmatrix} + \frac{i\beta Y_{\mathbf{1}}^{(10)} v_\xi v_d v_\phi}{\Lambda} \begin{pmatrix} 0 & \omega & -\omega^2 \\ -\omega & 0 & \omega^2 \\ \omega^2 & -\omega^2 & 0 \end{pmatrix}, \\ m_D &= \frac{i\omega^2 g_1 v_u v_\varphi}{\Lambda} \begin{pmatrix} 0 & 1 & -1 \\ -1 & 0 & 1 \\ 1 & -1 & 0 \end{pmatrix} + \frac{g_2 v_u v_\xi v_\varphi}{\Lambda^2} \begin{pmatrix} \sqrt{2} Y_{\mathbf{2},2}^{(7)} & -\omega^2 Y_{\mathbf{2},1}^{(7)} & -\omega^2 Y_{\mathbf{2},1}^{(7)} \\ -\omega^2 Y_{\mathbf{2},1}^{(7)} & \sqrt{2} Y_{\mathbf{2},2}^{(7)} & -\omega^2 Y_{\mathbf{2},1}^{(7)} \\ -\omega^2 Y_{\mathbf{2},1}^{(7)} & -\omega^2 Y_{\mathbf{2},1}^{(7)} & \sqrt{2} Y_{\mathbf{2},2}^{(7)} \end{pmatrix}, \\ m_N &= h v_\phi \begin{pmatrix} \sqrt{2} Y_{\mathbf{2}',2}^{(5)} & -\omega Y_{\mathbf{2}',1}^{(5)} & -\omega^2 Y_{\mathbf{2}',1}^{(5)} \\ -\omega Y_{\mathbf{2}',1}^{(5)} & \sqrt{2} Y_{\mathbf{2}',2}^{(5)} & -\omega^2 Y_{\mathbf{2}',1}^{(5)} \\ -\omega^2 Y_{\mathbf{2}',1}^{(5)} & -\omega^2 Y_{\mathbf{2}',1}^{(5)} & \sqrt{2} \omega^2 Y_{\mathbf{2}',2}^{(5)} \end{pmatrix}. \end{aligned} \quad (5.24)$$

The light neutrino mass matrix is then given by the seesaw relation:

$$m_\nu = -m_D^T m_N^{-1} m_D. \quad (5.25)$$

Since both flavons ϕ and φ enter into the neutrino mass terms and their VEVs preserve different Z_3 subgroups of $\Delta(27)$, the traditional flavor group $\Delta(27)$ is completely broken in the neutrino sector. One can easily check that the hermitian combination $m_l^\dagger m_l$ is invariant under the action of $\rho_{\mathbf{3}}(A^2B)$. It implies that the residual flavor symmetry is $Z_3^{A^2B}$ in the charged lepton sector. We find that the experimental data can be accommodated rather well for certain values of the input parameters. The best fit point is found to be given by

$$\begin{aligned} \Re\langle \tau \rangle &= 0.256, & \Im\langle \tau \rangle &= 0.987, & |\beta v_\xi / (\alpha \Lambda)| &= 0.127, & \arg(\beta v_\xi / (\alpha \Lambda)) &= 0.459\pi, \\ |g_2 v_\xi / (g_1 \Lambda)| &= 0.0416, & \arg(g_2 v_\xi / (g_1 \Lambda)) &= 0.552\pi, \\ \alpha v_d v_\xi v_\phi / \Lambda^2 &= 226.1 \text{ MeV}, & g_1 v_u^2 v_\varphi / \Lambda^2 &= 29.25 \text{ meV}, \end{aligned} \quad (5.26)$$




Figure 2. The predicted correlations among the input free parameters, neutrino mixing angles, CP violation phases and the effective mass in neutrinoless double beta decay in the seesaw model.

with $\chi^2_{\min} = 1.940$. Accordingly the lepton masses and mixing parameters are

$$\begin{aligned}
 \sin^2 \theta_{13} &= 0.02222, & \sin^2 \theta_{12} &= 0.291, & \sin^2 \theta_{23} &= 0.465, & \delta_{CP} &= 1.432\pi, \\
 \alpha_{21} &= 1.159\pi, & \alpha_{31} &= 0.186\pi, & m_1 &= 0.950 \text{ meV}, & m_2 &= 8.649 \text{ meV}, \\
 m_3 &= 50.02 \text{ meV}, & \sum_i m_i &= 59.61 \text{ meV}, & m_{\beta\beta} &= 2.981 \text{ meV}, \\
 m_e &= 0.511 \text{ MeV}, & m_\mu &= 107.9 \text{ MeV}, & m_\tau &= 1.836 \text{ GeV}.
 \end{aligned}
 \tag{5.27}$$

We perform a numerical analysis for this model, and the results are plotted in figure 2. It is remarkable that the phenomenologically viable parameter space is actually small. We find that the mixing angles θ_{13} and θ_{12} can take any value in their 3σ ranges, the atmospheric mixing angle $\sin^2 \theta_{23}$ is predicted to be in the range $[0.420, 0.529]$, the Dirac CP violation phase δ_{CP} is limited in the range $[1.247\pi, 1.627\pi]$, and the two Majorana CP violation phases lie in narrow regions $\alpha_{21} \in [1.113\pi, 1.187\pi]$ and $\alpha_{31} \in [0.097\pi, 0.222\pi]$. The forthcoming long-baseline neutrino oscillation experiments DUNE [53] and T2HK [54] can significantly improve the sensitivity to θ_{23} and δ_{CP} . It is expected that a 5σ discovery of CP violation can be reached after ten years of data taking. We expect that the predictions for θ_{23} and δ_{CP} can be tested at DUNE and T2HK.

6 Conclusion and outlook

In the present work, we elaborate how to construct models based on the so-called eclectic flavor group [17] which is in general a semi-direct product of the traditional flavor group

with the finite modular group in the bottom-up approach. The concept of eclectic flavor group originated in top-down model building motivated by string theory. In the scheme of eclectic flavor groups, which allows us to combine the advantages of modular symmetries and traditional flavor symmetries while avoiding their limitations, the superpotential and the Kähler potential are both severely restricted at the same time. However, there are no remaining exact flavor symmetries in nature, flavon fields should be introduced in this kind of models, and their VEVs break the flavor symmetries. It turns out that the choices for such eclectic symmetries are quite restrictive, with only three choices of flavor group $Z_3 \times Z_3$, $\Delta(27)$ and $\Delta(54)$ catalogued so far allowing for a non-trivial eclectic extension with a CP-like transformation [17, 18].

We perform a comprehensive analysis of the superpotential and Kähler potential of models based on the eclectic flavor group $\Omega(1)$ which is semi-direct product of the traditional flavor group $\Delta(27)$ with the finite modular group T' , and find three generations of quarks and leptons can only be assigned to triplet, antitriplet or trivial singlet of $\Delta(27)$. From the discussion in section 4, the general form of fermion mass matrices can be obtained for any modular weight and any number of flavons. If a $\Delta(27)$ octet flavon which is invariant under auxiliary cyclic symmetries is imposed in a model, we find that the contributions to the fermion masses and fermion mixings from higher order corrections of Kähler potential are suppressed by powers of $\langle \Phi \rangle / \Lambda$, or they will be suppressed by $\langle \Phi \rangle^2 / \Lambda^2$.

In order to extract general lessons from the eclectic flavor symmetry scheme, we construct two concrete lepton models invariant under the eclectic flavor group $\Omega(1) \cong \Delta(27) \rtimes T'$ in the framework of $\mathcal{N} = 1$ global supersymmetry. In the two models, all lepton fields are assigned to $\Delta(27)$ triplet and the three light neutrino masses are generated by the Weinberg operator and the type I seesaw mechanism, respectively. To break the traditional flavor group $\Delta(27)$, three (two) triplet flavons and a trivial singlet flavon are introduced in the first (second) model. Then the higher order corrections from Kähler potential to the lepton mixing parameters and masses are suppressed by $\langle \Phi \rangle^2 / \Lambda^2$ and can be neglected. For the first model, we assume that the traditional flavor group $\Delta(27)$ is broken down to the subgroup $Z_3^{A^2 B A^2}$ in the charged lepton sector and is broken completely in the neutrino sector. For the second model, the subgroup $Z_3^{A^2 B}$ is preserved by the VEVs of the flavon in the charged lepton mass terms and neutrino Majorana mass term, while the subgroup Z_3^A is preserved by vacuum of flavon in the neutrino Dirac mass terms. Furthermore, we perform a comprehensive analysis of the two models for lepton masses and mixing parameters. In both of the two models, the six lepton masses and six lepton mixing parameters depend on two complex coupling constants, two real coupling constants and the complex modulus τ in the case of without gCP. From the numerical scan of the parameter space, we find that all six mixing parameters vary in very small regions, and a nearly maximal atmospheric mixing angle θ_{23} and Dirac CP phase δ_{CP} are obtained in the first model. When the conditions gCP and $\Re\tau = 0$ are imposed in the first model, the experimental data of lepton masses and mixing parameters can be successfully described in terms of five real parameters, and the $\mu - \tau$ reflection symmetry is preserved in the charged lepton diagonal basis. Then atmospheric mixing angle and Dirac CP phase are predicted to be maximal, and two Majorana CP phases are predicted to be π . Furthermore, the predictions for the effective mass in neutrinoless double beta decay are obtained in both of the two models.

All the predictions of the two models are compatible with the experimental data and could be tested at forthcoming long baseline neutrino experiments and neutrinoless double beta decay experiments.

In general, the main advantage of eclectic flavor symmetry is that it leads to predictive lepton mass and mixing models with the Kähler potential under control. Also, the representation of the matter fields under the flavor symmetry is constrained by the eclectic flavor symmetry. For example, in the considered models, this means that non-trivial singlet representations are not permitted since they would have to form a reducible octet, which is not phenomenologically possible. Also, we have seen that the modular symmetry transformation is determined by the flavor symmetry transformation through the consistency condition.

Looking ahead, it would be interesting to explore how to implement top-down ingredients, such as the constrained choice of modular weights, in the future bottom-up eclectic models. Also, the problem of modulus stabilisation in theory based on modular symmetry still persists, and the presence of flavons only exacerbates the problem. Such problems are left for future work. In the present paper, we have developed the formalism of eclectic flavor symmetry from the bottom-up point of view, focusing on an attractive class of models based on $\Omega(1)$, where the imposition of gCP symmetry leads to a predictive model with $\mu - \tau$ reflection symmetry.

Acknowledgments

GJD and XGL are grateful to professor Ferruccio Feruglio for sharing the his insightful notes on eclectic flavor group. CCL is supported by the National Natural Science Foundation of China under Grant Nos. 12005167, 12247103. XGL is supported in part by U.S. National Science Foundation under Grant No. PHY-2210283. JNL is supported by the Grants No. NSFC-12147110 and the China Post-doctoral Science Foundation under Grant No. 2021M70. GJD is supported by the National Natural Science Foundation of China under Grant Nos. 11975224, 11835013. SFK acknowledges the STFC Consolidated Grant ST/L000296/1 and the European Union’s Horizon 2020 Research and Innovation programme under Marie Skłodowska-Curie grant agreement HIDDEN European ITN project (H2020-MSCA-ITN-2019//860881-HIDDEN).

A Traditional flavor group $\Delta(27)$

The $\Delta(27)$ group as flavor symmetry has been studied in the literature, see [55–58] and references therein. It is a non-Abelian group of order 27 with GAP ID [27, 3] [59], and the group structure of $\Delta(27)$ is $\Delta(27) \cong (Z_3 \times Z_3) \rtimes Z_3$. In detail, $\Delta(27)$ can be generated by two generators A and B which satisfy the multiplication rules

$$A^3 = B^3 = (AB)^3 = (AB^2)^3 = 1. \tag{A.1}$$

The multiplication rules of $\Delta(27)$ in refs. [58, 59] are given by:

$$a^3 = a'^3 = b^3 = 1, \quad aa' = a'a, \quad bab^{-1} = a^{-1}a'^{-1}, \quad ba'b^{-1} = a. \tag{A.2}$$

Note that the chosen generators A and B in the present work are related to the generators a , a' and b of refs. [58, 59] via

$$a = BAB^2, \quad a' = A, \quad b = B. \quad (\text{A.3})$$

The traditional flavor group $\Delta(27)$ has eleven conjugacy classes and they are

$$\begin{aligned} 1C_1 &= \{1\}, & 3C_3^{(1)} &= \{A^2B^2, B^2A^2, AB^2A\}, \\ 3C_3^{(2)} &= \{A^2B, ABA, BA^2\}, & 3C_3^{(3)} &= \{A, BAB^2, B^2AB\}, \\ 3C_3^{(4)} &= \{AB^2, BAB, B^2A\}, & 3C_3^{(5)} &= \{AB, BA, A^2BA^2\}, \\ 3C_3^{(6)} &= \{A^2, B^2A^2B, BA^2B^2\}, & 3C_3^{(7)} &= \{B^2, AB^2A^2, A^2B^2A\}, \\ 3C_3^{(8)} &= \{B, A^2BA, ABA^2\}, & 1C_3^{(1)} &= \{BAB^2A^2\}, \quad 1C_3^{(2)} = \{ABA^2B^2\}, \end{aligned} \quad (\text{A.4})$$

where kC_n denotes a conjugacy class which contains k elements with order n . Group $\Delta(27)$ contains seventeen nontrivial abelian subgroups in total: thirteen Z_3 subgroups and four $Z_3 \times Z_3$ subgroups. In terms of the generators A and B , the concrete forms of these abelian subgroups are as follows

- Z_3 subgroups

$$\begin{aligned} Z_3^A &= \{1, A, A^2\}, & Z_3^{BAB^2} &= \{1, BAB^2, BA^2B^2\}, & Z_3^{B^2AB} &= \{1, B^2AB, B^2A^2B\}, \\ Z_3^B &= \{1, B, B^2\}, & Z_3^{A^2BA} &= \{1, A^2BA, A^2B^2A\}, & Z_3^{ABA^2} &= \{1, ABA^2, AB^2A^2\}, \\ Z_3^{AB} &= \{1, AB, B^2A^2\}, & Z_3^{BA} &= \{1, BA, A^2B^2\}, & Z_3^{AB^2A} &= \{1, AB^2A, A^2BA^2\}, \\ Z_3^{A^2B} &= \{1, A^2B, B^2A\}, & Z_3^{AB^2} &= \{1, AB^2, BA^2\}, & Z_3^{ABA} &= \{1, ABA, BAB\}, \\ Z_3^{BAB^2A^2} &= \{1, BAB^2A^2, ABA^2B^2\}. \end{aligned}$$

The thirteen Z_3 subgroups in each line above are related with each other by group conjugation.

- $Z_3 \times Z_3$ subgroups

$$\begin{aligned} Z_3^A \times Z_3^{BAB^2} &= \{1, A, BAB^2A^2, A^2, BAB^2, ABA^2B^2, B^2A^2B, B^2AB, BA^2B^2\}, \\ Z_3^B \times Z_3^{ABA^2} &= \{1, B, BAB^2A^2, B^2, A^2BA, ABA^2B^2, AB^2A^2, ABA^2, A^2B^2A\}, \\ Z_3^{BA} \times Z_3^{B^2A^2} &= \{1, BAB^2A^2, AB, ABA^2B^2, BA, A^2B^2, A^2BA^2, B^2A^2, AB^2A\}, \\ Z_3^{B^2A} \times Z_3^{AB^2} &= \{1, BAB^2A^2, ABA^2B^2, A^2B, AB^2, ABA, BAB, BA^2, B^2A\}. \end{aligned}$$

All the four $Z_3 \times Z_3$ subgroups are related with each other by group conjugation.

As the number of irreducible representation is equal to the number of conjugacy class, the traditional flavor group $\Delta(27)$ has eleven inequivalent irreducible representations with nine singlets labelled as $\mathbf{1}_{(r,s)}$ ($r, s = 0, 1, 2$) and two triplets labelled as $\mathbf{3}$ and $\bar{\mathbf{3}}$. The character

	E	A^2B^2	A^2B	A	AB^2	AB	A^2	B^2	B	BAB^2A^2	ABA^2B^2
	$1C_1$	$3C_3^{(1)}$	$3C_3^{(2)}$	$3C_3^{(3)}$	$3C_3^{(4)}$	$3C_3^{(5)}$	$3C_3^{(6)}$	$3C_3^{(7)}$	$3C_3^{(8)}$	$1C_3^{(1)}$	$1C_3^{(2)}$
$\mathbf{1}_{0,0}$	1	1	1	1	1	1	1	1	1	1	1
$\mathbf{1}_{0,1}$	1	ω^2	ω	1	ω^2	ω	1	ω^2	ω	1	1
$\mathbf{1}_{0,2}$	1	ω	ω^2	1	ω	ω^2	1	ω	ω^2	1	1
$\mathbf{1}_{1,0}$	1	ω^2	ω^2	ω	ω	ω	ω^2	1	1	1	1
$\mathbf{1}_{1,1}$	1	ω	1	ω	1	ω^2	ω^2	ω^2	ω	1	1
$\mathbf{1}_{1,2}$	1	1	ω	ω	ω^2	1	ω^2	ω	ω^2	1	1
$\mathbf{1}_{2,0}$	1	ω	ω	ω^2	ω^2	ω^2	ω	1	1	1	1
$\mathbf{1}_{2,1}$	1	1	ω^2	ω^2	ω	1	ω	ω^2	ω	1	1
$\mathbf{1}_{2,2}$	1	ω^2	1	ω^2	1	ω	ω	ω	ω^2	1	1
$\mathbf{3}$	3	0	0	0	0	0	0	0	0	3ω	$3\omega^2$
$\bar{\mathbf{3}}$	3	0	0	0	0	0	0	0	0	$3\omega^2$	3ω

Table 6. The character table of $\Delta(27)$, where ω is the cube root of unity $\omega = e^{2\pi i/3}$. The arrowed lines show the transformation of the irreducible representations and conjugacy classes of $\Delta(27)$ under the actions of the outer automorphisms u_S (blue), u_T (red) and u_{K^*} (green).

table of $\Delta(27)$ is shown in table 6. The first line of table 6 indicates representatives of the eleven conjugacy classes in the second line. In our working basis, the eleven irreducible representations of $\Delta(27)$ are taken to be

$$\begin{aligned}
 \mathbf{1}_{r,s} : \quad & \rho_{\mathbf{1}_{r,s}}(A) = \omega^r, & \rho_{\mathbf{1}_{r,s}}(B) = \omega^s, & \text{with } r, s = 0, 1, 2, \\
 \mathbf{3} : \quad & \rho_{\mathbf{3}}(A) = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, & \rho_{\mathbf{3}}(B) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega & 0 \\ 0 & 0 & \omega^2 \end{pmatrix}, \\
 \bar{\mathbf{3}} : \quad & \rho_{\bar{\mathbf{3}}}(A) = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, & \rho_{\bar{\mathbf{3}}}(B) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega^2 & 0 \\ 0 & 0 & \omega \end{pmatrix}.
 \end{aligned} \tag{A.5}$$

From the character table, it is easy to calculate the Kronecker products of two different irreducible representations of $\Delta(27)$ as

$$\begin{aligned}
 \mathbf{1}_{r,s} \otimes \mathbf{1}_{r',s'} &= \mathbf{1}_{[r+r'], [s+s']}, & \mathbf{1}_{r,s} \otimes \mathbf{3} &= \mathbf{3}, & \mathbf{1}_{r,s} \otimes \bar{\mathbf{3}} &= \bar{\mathbf{3}}, \\
 \mathbf{3} \otimes \mathbf{3} &= \bar{\mathbf{3}}_{S,1} \otimes \bar{\mathbf{3}}_{S,2} \otimes \bar{\mathbf{3}}_A, & \bar{\mathbf{3}} \otimes \bar{\mathbf{3}} &= \mathbf{3}_{S,1} \otimes \mathbf{3}_{S,2} \otimes \mathbf{3}_A, & \mathbf{3} \otimes \bar{\mathbf{3}} &= \sum_{r,s=0}^2 \mathbf{1}_{r,s},
 \end{aligned} \tag{A.6}$$

where $r, s, r', s' = 0, 1, 2$, the integer $[n] \equiv n \pmod{3}$, and the subscript S (A) denotes symmetric (antisymmetric) combination. In the following, all CG coefficients are presented in the form $\alpha \otimes \beta$, where α_i denotes the elements of the first representation and β_j stands

for the elements of the second representation of the tensor product. In the following, we shall adopt the convention $\beta_{[3]} = \beta_0 \equiv \beta_3$.

$\mathbf{1}_{r,s} \otimes \mathbf{1}_{r',s'} = \mathbf{1}_{[r+r'],[s+s']}$	$\mathbf{1}_{r,s} \otimes \mathbf{3} = \mathbf{3}$	$\mathbf{1}_{r,s} \otimes \bar{\mathbf{3}} = \bar{\mathbf{3}}$
$\mathbf{1}_{[r+r'],[s+s]} : \alpha_1 \beta_1$	$\mathbf{3} : \alpha_1 \begin{pmatrix} \beta_{[1-s]} \\ \omega^r \beta_{[2-s]} \\ \omega^{2r} \beta_{[3-s]} \end{pmatrix}$	$\bar{\mathbf{3}} : \alpha_1 \begin{pmatrix} \beta_{[1+s]} \\ \omega^r \beta_{[2+s]} \\ \omega^{2r} \alpha_1 \beta_{[3+s]} \end{pmatrix}$
$\mathbf{3} \otimes \bar{\mathbf{3}} = \sum_{k,r=0}^2 \mathbf{1}_{k,r}$	$\mathbf{3} \otimes \mathbf{3} = \bar{\mathbf{3}}_{S_1} \oplus \bar{\mathbf{3}}_{S_2} \oplus \bar{\mathbf{3}}_A$	$\bar{\mathbf{3}} \otimes \bar{\mathbf{3}} = \mathbf{3}_{S_1} \oplus \mathbf{3}_{S_2} \oplus \mathbf{3}_A$
$\mathbf{1}_{0,0} : \alpha_1 \beta_1 + \alpha_2 \beta_2 + \alpha_3 \beta_3$ $\mathbf{1}_{0,1} : \alpha_1 \beta_3 + \alpha_2 \beta_1 + \alpha_3 \beta_2$ $\mathbf{1}_{0,2} : \alpha_1 \beta_2 + \alpha_2 \beta_3 + \alpha_3 \beta_1$ $\mathbf{1}_{1,0} : \alpha_1 \beta_1 + \omega^2 \alpha_2 \beta_2 + \omega \alpha_3 \beta_3$ $\mathbf{1}_{1,1} : \alpha_1 \beta_3 + \omega^2 \alpha_2 \beta_1 + \omega \alpha_3 \beta_2$ $\mathbf{1}_{1,2} : \alpha_1 \beta_2 + \omega^2 \alpha_2 \beta_3 + \omega \alpha_3 \beta_1$ $\mathbf{1}_{2,0} : \alpha_1 \beta_1 + \omega \alpha_2 \beta_2 + \omega^2 \alpha_3 \beta_3$ $\mathbf{1}_{2,1} : \alpha_1 \beta_3 + \omega \alpha_2 \beta_1 + \omega^2 \alpha_3 \beta_2$ $\mathbf{1}_{2,2} : \alpha_1 \beta_2 + \omega \alpha_2 \beta_3 + \omega^2 \alpha_3 \beta_1$	$\bar{\mathbf{3}}_{S_1} : \begin{pmatrix} \alpha_1 \beta_1 \\ \alpha_2 \beta_2 \\ \alpha_3 \beta_3 \end{pmatrix}$ $\bar{\mathbf{3}}_{S_2} : \begin{pmatrix} \alpha_2 \beta_3 + \alpha_3 \beta_2 \\ \alpha_3 \beta_1 + \alpha_1 \beta_3 \\ \alpha_1 \beta_2 + \alpha_2 \beta_1 \end{pmatrix}$ $\bar{\mathbf{3}}_A : \begin{pmatrix} \alpha_2 \beta_3 - \alpha_3 \beta_2 \\ \alpha_3 \beta_1 - \alpha_1 \beta_3 \\ \alpha_1 \beta_2 - \alpha_2 \beta_1 \end{pmatrix}$	$\mathbf{3}_{S_1} : \begin{pmatrix} \alpha_1 \beta_1 \\ \alpha_2 \beta_2 \\ \alpha_3 \beta_3 \end{pmatrix}$ $\mathbf{3}_{S_2} : \begin{pmatrix} \alpha_2 \beta_3 + \alpha_3 \beta_2 \\ \alpha_3 \beta_1 + \alpha_1 \beta_3 \\ \alpha_1 \beta_2 + \alpha_2 \beta_1 \end{pmatrix}$ $\mathbf{3}_A : \begin{pmatrix} \alpha_2 \beta_3 - \alpha_3 \beta_2 \\ \alpha_3 \beta_1 - \alpha_1 \beta_3 \\ \alpha_1 \beta_2 - \alpha_2 \beta_1 \end{pmatrix}$

B The finite modular group $\Gamma'_3 \cong T'$

The finite modular group $\Gamma'_3 \cong T'$ contains 24 elements and it can be generated by two generators S and T with [7, 29, 60, 61]

$$S^4 = T^3 = (ST)^3 = 1, \quad S^2 T = T S^2. \quad (\text{B.1})$$

The 24 elements of T' can be divided into the seven conjugacy classes as follows,

$$\begin{aligned}
 1C_1 &= \{1\}, & 1C_2 &= \{S^2\}, & 6C_4 &= \{S, T^2 ST, TST^2, T^2 S^3 T, TS^3 T^2, S^3\}, \\
 4C_3^{(1)} &= \{T^2, S^3 T^2, T^2 S^3, TS^3 T\}, & 4C_3^{(2)} &= \{T, TS, ST, T^2 ST^2\}, \\
 4C_6^{(1)} &= \{TS^2, TS^3, S^3 T, T^2 S^3 T^2\}, & 4C_6^{(2)} &= \{ST^2, T^2 S, TST, T^2 S^2\},
 \end{aligned} \quad (\text{B.2})$$

where the conjugacy class is denoted by nC_k with n elements and the subscript k is the order of the elements. The finite modular group T' has three singlet representations $\mathbf{1}$, $\mathbf{1}'$ and $\mathbf{1}''$, three doublet representations $\mathbf{2}$, $\mathbf{2}'$ and $\mathbf{2}''$, and one triplet representation $\mathbf{3}$. The character table of T' group is given in table 7. The Kronecker products between different irreducible representations can be obtained from the character table

$$\begin{aligned}
 \mathbf{1}^a \otimes \mathbf{1}^b &= \mathbf{1}^{[a+b]}, & \mathbf{1}^a \otimes \mathbf{2}^b &= \mathbf{2}^{[a+b]}, & \mathbf{1}^a \otimes \mathbf{3} &= \mathbf{3}, & \mathbf{2}^a \otimes \mathbf{2}^b &= \mathbf{1}^{[a+b]} \oplus \mathbf{3}, \\
 \mathbf{2}^a \otimes \mathbf{3} &= \mathbf{2} \oplus \mathbf{2}' \oplus \mathbf{2}'', & \mathbf{3} \otimes \mathbf{3} &= \mathbf{1} \oplus \mathbf{1}' \oplus \mathbf{1}'' \oplus \mathbf{3}_S \oplus \mathbf{3}_A,
 \end{aligned} \quad (\text{B.3})$$

Classes	$1C_1$	$1C_2$	$4C_3^{(1)}$	$4C_3^{(2)}$	$6C_4$	$4C_6^{(1)}$	$4C_6^{(2)}$
1	1	1	1	1	1	1	1
1'	1	1	ω^2	ω	1	ω	ω^2
1''	1	1	ω	ω^2	1	ω^2	ω
2	2	-2	-1	-1	0	1	1
2'	2	-2	$-\omega^2$	$-\omega$	0	ω	ω^2
2''	2	-2	$-\omega$	$-\omega^2$	0	ω^2	ω
3	3	3	0	0	-1	0	0

Table 7. Character table of the finite modular group T' .

where $a, b = 0, 1, 2$ and we denote $\mathbf{1}^0 \equiv \mathbf{1}$, $\mathbf{1}^1 \equiv \mathbf{1}'$, $\mathbf{1}^2 \equiv \mathbf{1}''$ and similarly for the doublet representations. On the right-hand-side $[a + b]$ is defined as $a + b$ modulo 3. In the present work, the seven irreducible representations are taken to be

$$\begin{aligned}
 \mathbf{1}^a : \quad & \rho_{\mathbf{1}^a}(S) = 1, & \rho_{\mathbf{1}^a}(T) &= \omega^a, \\
 \mathbf{2}^a : \quad & \rho_{\mathbf{2}^a}(S) = \frac{i}{\sqrt{3}} \begin{pmatrix} 1 & \sqrt{2} \\ \sqrt{2} & -1 \end{pmatrix}, & \rho_{\mathbf{2}^a}(T) &= \omega^{a+1} \begin{pmatrix} 1 & 0 \\ 0 & \omega \end{pmatrix}, \\
 \mathbf{3} : \quad & \rho_{\mathbf{3}}(S) = \frac{1}{3} \begin{pmatrix} -1 & 2 & 2 \\ 2 & -1 & 2 \\ 2 & 2 & -1 \end{pmatrix}, & \rho_{\mathbf{3}}(T) &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega & 0 \\ 0 & 0 & \omega^2 \end{pmatrix}, \quad (\text{B.4})
 \end{aligned}$$

with $a = 0, 1, 2$. From the irreducible representation matrices of T' in eq. (B.4), one can easily check that

$$\begin{aligned}
 \rho_{\mathbf{2}}^*(T) &= U_2^\dagger \rho_{\mathbf{2}}(T) U_2, & \rho_{\mathbf{2}}^*(S) &= U_2^\dagger \rho_{\mathbf{2}}(S) U_2, \\
 \rho_{\mathbf{2}'}^*(T) &= U_2^\dagger \rho_{\mathbf{2}''}(T) U_2, & \rho_{\mathbf{2}'}^*(S) &= U_2^\dagger \rho_{\mathbf{2}''}(S) U_2, \\
 \rho_{\mathbf{3}}^*(T) &= U_3^\dagger \rho_{\mathbf{3}}(T) U_3, & \rho_{\mathbf{3}}^*(S) &= U_3^\dagger \rho_{\mathbf{3}}(S) U_3,
 \end{aligned} \quad (\text{B.5})$$

with

$$U_2 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad U_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}. \quad (\text{B.6})$$

It implies that the irreducible representations $\mathbf{2}'$ and $\mathbf{2}''$ are complex conjugated to each other by a unitary transformation U_2 , and the irreducible representations $\mathbf{2}$ and $\mathbf{3}$ are real representation. Thus we have

$$\begin{aligned}
 \alpha &= (\alpha_1, \alpha_2)^T \sim \mathbf{2}, & \rightarrow & \quad (-\alpha_2^*, \alpha_1^*)^T \sim \mathbf{2}, \\
 \alpha &= (\alpha_1, \alpha_2)^T \sim \mathbf{2}', & \rightarrow & \quad (-\alpha_2^*, \alpha_1^*)^T \sim \mathbf{2}'', \\
 \alpha &= (\alpha_1, \alpha_2)^T \sim \mathbf{2}'', & \rightarrow & \quad (-\alpha_2^*, \alpha_1^*)^T \sim \mathbf{2}', \\
 \alpha &= (\alpha_1, \alpha_2, \alpha_3)^T \sim \mathbf{3}, & \rightarrow & \quad (\alpha_1^*, \alpha_3^*, \alpha_2^*)^T \sim \mathbf{3}.
 \end{aligned} \quad (\text{B.7})$$

In the end, we present the CG coefficients of T' in the chosen basis. As the same as the CG coefficients of $\Delta(27)$, we use the notation α_i (β_i) to denote the elements of the first (second) representation. Furthermore, we shall adopt the following notation to facilitate the expressions of CG coefficients

$$P = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}. \quad (\text{B.8})$$

Then all the CG coefficients of T' for the decomposition of product representations can be written as

$\mathbf{1}^a \otimes \mathbf{2}^b = \mathbf{2}^{[a+b]}$	$\mathbf{1}^a \otimes \mathbf{3} = \mathbf{3}$	$\mathbf{2}^a \otimes \mathbf{2}^b = \mathbf{1}^{[a+b]} \oplus \mathbf{3}$
$\mathbf{2}^{[a+b]} : \alpha_1 \begin{pmatrix} \beta_1 \\ \beta_2 \end{pmatrix}$	$\mathbf{3} : \alpha_1 P^a \begin{pmatrix} \beta_1 \\ \beta_2 \\ \beta_3 \end{pmatrix}$	$\mathbf{1}^{[a+b]} : \alpha_1 \beta_2 - \alpha_2 \beta_1$ $\mathbf{3} : P^{[a+b]} \begin{pmatrix} \alpha_1 \beta_2 + \alpha_2 \beta_1 \\ \sqrt{2} \alpha_2 \beta_2 \\ -\sqrt{2} \alpha_1 \beta_1 \end{pmatrix}$
$\mathbf{3} \otimes \mathbf{3} = \mathbf{1} \oplus \mathbf{1}' \oplus \mathbf{1}'' \oplus \mathbf{3}_S \oplus \mathbf{3}_A$		$\mathbf{2}^a \otimes \mathbf{3} = \mathbf{2} \oplus \mathbf{2}' \oplus \mathbf{2}''$
$\mathbf{1} : \alpha_1 \beta_1 + \alpha_2 \beta_3 + \alpha_3 \beta_2$ $\mathbf{1}' : \alpha_1 \beta_2 + \alpha_2 \beta_1 + \alpha_3 \beta_3$ $\mathbf{1}'' : \alpha_1 \beta_3 + \alpha_2 \beta_2 + \alpha_3 \beta_1$ $\mathbf{3}_S : \begin{pmatrix} 2\alpha_1 \beta_1 - \alpha_2 \beta_3 - \alpha_3 \beta_2 \\ 2\alpha_3 \beta_3 - \alpha_1 \beta_2 - \alpha_2 \beta_1 \\ 2\alpha_2 \beta_2 - \alpha_1 \beta_3 - \alpha_3 \beta_1 \end{pmatrix}$ $\mathbf{3}_A : \begin{pmatrix} \alpha_2 \beta_3 - \alpha_3 \beta_2 \\ \alpha_1 \beta_2 - \alpha_2 \beta_1 \\ \alpha_3 \beta_1 - \alpha_1 \beta_3 \end{pmatrix}$		$\mathbf{2}^a : \begin{pmatrix} \sqrt{2} \alpha_2 \beta_3 + \alpha_1 \beta_1 \\ \sqrt{2} \alpha_1 \beta_2 - \alpha_2 \beta_1 \end{pmatrix}$ $\mathbf{2}^{[1+a]} : \begin{pmatrix} \sqrt{2} \alpha_2 \beta_1 + \alpha_1 \beta_2 \\ \sqrt{2} \alpha_1 \beta_3 - \alpha_2 \beta_2 \end{pmatrix}$ $\mathbf{2}^{[2+a]} : \begin{pmatrix} \sqrt{2} \alpha_2 \beta_2 + \alpha_1 \beta_3 \\ \sqrt{2} \alpha_1 \beta_1 - \alpha_2 \beta_3 \end{pmatrix}$

C Modular forms of level $N = 3$

The modular forms of weight k and level 3 are holomorphic functions of the complex modulus τ and they span a linear space $\mathcal{M}_k(\Gamma(3))$ with dimension

$$\dim \mathcal{M}_k(\Gamma(3)) = k + 1, \quad k \geq 1, \quad (\text{C.1})$$

where the modular weight k is a generic non-negative integer. Each modular form $f_i^{(k)}(\tau)$ of weight k and level 3 is invariant under the traditional flavor symmetry and transforms under the action of $\Gamma(3)$ in the following way

$$f_i^{(k)}(\tau) \rightarrow f_i^{(k)}(\gamma\tau) = (c\tau + d)^k f_i^{(k)}(\tau), \quad \forall \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma(3). \quad (\text{C.2})$$

Furthermore, the modular forms $f_i^{(k)}(\tau)$ can be arranged into some modular multiplets $Y_{\mathbf{r}}^{(k)}(\tau) \equiv (f_1(\tau), f_2(\tau), \dots)^T$ which transform as certain irreducible representation \mathbf{r} of the finite modular group T' [1, 29]

$$Y_{\mathbf{r}}^{(k)}(\gamma\tau) = (c\tau + d)^k \rho_{\mathbf{r}}(\gamma) Y_{\mathbf{r}}^{(k)}(\tau), \quad \forall \gamma \in \Gamma, \quad (\text{C.3})$$

where γ is the representative element of the coset $\gamma\Gamma(N)$ in T' , and $\rho_{\mathbf{r}}(\gamma)$ is the representation matrix of γ in the irreducible representation \mathbf{r} of T' .

The modular forms of level 3 can be generated by the products of Dedekind function $\eta(\tau)$, and there are only two linearly independent weight 1 and level 3 modular forms which can be taken to be [29]:

$$\hat{e}_1(\tau) = \frac{\eta^3(3\tau)}{\eta(\tau)}, \quad \hat{e}_2(\tau) = \frac{\eta^3(\tau/3)}{\eta(\tau)}, \quad (\text{C.4})$$

where the Dedekind function $\eta(\tau)$ is defined as

$$\eta(\tau) = q^{1/24} \prod_{n=1}^{\infty} (1 - q^n), \quad q \equiv e^{2\pi i\tau}. \quad (\text{C.5})$$

Under the actions of the two generators S and T of Γ , $\eta(\tau)$ transforms as

$$\eta(\tau) \xrightarrow{S} \eta(-1/\tau) = \sqrt{-i\tau} \eta(\tau), \quad \eta(\tau) \xrightarrow{T} \eta(\tau + 1) = e^{i\pi/12} \eta(\tau), \quad (\text{C.6})$$

which lead to the following transformation properties [29]

$$\begin{aligned} \hat{e}_1(\tau) &\xrightarrow{S} 3^{-3/2}(-i\tau)\hat{e}_2(\tau), & \hat{e}_2(\tau) &\xrightarrow{S} 3^{3/2}(-i\tau)\hat{e}_1(\tau). \\ \hat{e}_1(\tau) &\xrightarrow{T} \omega\hat{e}_1(\tau), & \hat{e}_2(\tau) &\xrightarrow{T} 3(1 - \omega)\hat{e}_1 + \hat{e}_2. \end{aligned} \quad (\text{C.7})$$

The two independent weight 1 modular forms of level 3 can be arranged into a doublet $\mathbf{2}''$ of T' in our basis given in appendix B and the corresponding modular multiplet $Y_{\mathbf{2}''}^{(1)}(\tau)$ is defined as

$$Y_{\mathbf{2}''}^{(1)}(\tau) \equiv \begin{pmatrix} Y_1(\tau) \\ Y_2(\tau) \end{pmatrix} = \begin{pmatrix} 3\hat{e}_1(\tau) + \hat{e}_2(\tau) \\ 3\sqrt{2}\hat{e}_1(\tau) \end{pmatrix}. \quad (\text{C.8})$$

The expressions of the q-expansions of $Y_i(\tau)$ are given by

$$\begin{aligned} Y_1(\tau) &= 1 + 6q + 6q^3 + 6q^4 + 12q^7 + 6q^9, \\ Y_2(\tau) &= 3\sqrt{2}q^{1/3}(1 + q + 2q^2 + 2q^4 + q^5 + 2q^6 + q^8 + 2q^9). \end{aligned} \quad (\text{C.9})$$

The higher weight modular multiplets can be constructed by the tensor products of lower weight modular multiplets. One can find the modular multiplets of any integral weight by using the weight 1 modular multiplet $Y_{\mathbf{2}''}^{(1)}(\tau)$ and the CG coefficients of T' in appendix B. We summarize the modular multiplets of level $N = 3$ up to weight 10 in table 8. Their

weight	Modular form $Y_r^{(k)}$	weight	Modular form $Y_r^{(k)}$
$k = 1$	$Y_{2''}^{(1)}$	$k = 2$	$Y_3^{(2)}$
$k = 3$	$Y_{2'}^{(3)}, Y_{2''}^{(3)}$	$k = 4$	$Y_1^{(4)}, Y_{1'}^{(4)}, Y_3^{(4)}$
$k = 5$	$Y_2^{(5)}, Y_{2'}^{(5)}, Y_{2''}^{(5)}$	$k = 6$	$Y_1^{(6)}, Y_{3i}^{(6)}, Y_{3ii}^{(6)}$
$k = 7$	$Y_2^{(7)}, Y_{2'}^{(7)}, Y_{2''i}^{(7)}, Y_{2''ii}^{(7)}$	$k = 8$	$Y_1^{(8)}, Y_{1'}^{(8)}, Y_{1''}^{(8)}, Y_{3i}^{(8)}, Y_{3ii}^{(8)}$
$k = 9$	$Y_2^{(9)}, Y_{2'i}^{(9)}, Y_{2''ii}^{(9)}, Y_{2''i}^{(9)}, Y_{2''ii}^{(9)}$	$k = 10$	$Y_1^{(10)}, Y_{1'}^{(10)}, Y_{3i}^{(10)}, Y_{3ii}^{(10)}, Y_{3iii}^{(10)}$

Table 8. The higher integral weight modular multiplets of level 3 up to weight 10, the subscript r denotes the transformation property under homogeneous finite group $\Gamma_3' \cong T'$.

explicit forms of these modular multiplets are

$$\begin{aligned}
Y_3^{(2)} &= \left(Y_{2''}^{(1)} Y_{2''}^{(1)} \right)_3, & Y_{2'}^{(3)} &= \left(Y_{2''}^{(1)} Y_3^{(2)} \right)_{2'}, & Y_{2''}^{(3)} &= \left(Y_{2''}^{(1)} Y_3^{(2)} \right)_{2''}, \\
Y_1^{(4)} &= \left(Y_3^{(2)} Y_3^{(2)} \right)_1, & Y_{1'}^{(4)} &= \left(Y_3^{(2)} Y_3^{(2)} \right)_{1'}, & Y_3^{(4)} &= \left(Y_3^{(2)} Y_3^{(2)} \right)_3, \\
Y_2^{(5)} &= \left(Y_3^{(2)} Y_{2'}^{(3)} \right)_2, & Y_{2'}^{(5)} &= \left(Y_3^{(2)} Y_{2'}^{(3)} \right)_{2'}, & Y_{2''}^{(5)} &= \left(Y_3^{(2)} Y_{2'}^{(3)} \right)_{2''}, \\
Y_{1'}^{(6)} &= \left(Y_{2'}^{(3)} Y_{2''}^{(3)} \right)_1, & Y_{3i}^{(6)} &= \left(Y_{2'}^{(3)} Y_{2''}^{(3)} \right)_3, & Y_{3ii}^{(6)} &= \left(Y_{2'}^{(3)} Y_{2''}^{(3)} \right)_3, \\
Y_2^{(7)} &= \left(Y_{2'}^{(3)} Y_3^{(4)} \right)_2, & Y_{2'}^{(7)} &= \left(Y_{2'}^{(3)} Y_1^{(4)} \right)_{2'}, & Y_{2''i}^{(7)} &= \left(Y_{2'}^{(3)} Y_{1'}^{(4)} \right)_{2''}, \\
Y_{2''ii}^{(7)} &= \left(Y_{2'}^{(3)} Y_3^{(4)} \right)_{2''}, & Y_1^{(8)} &= \left(Y_1^{(4)} Y_1^{(4)} \right)_1, & Y_{1'}^{(8)} &= \left(Y_1^{(4)} Y_1^{(4)} \right)_{1'}, \\
Y_{1''}^{(8)} &= \left(Y_{1'}^{(4)} Y_{1'}^{(4)} \right)_{1''}, & Y_{3i}^{(8)} &= \left(Y_1^{(4)} Y_3^{(4)} \right)_3, & Y_{3ii}^{(8)} &= \left(Y_{1'}^{(4)} Y_3^{(4)} \right)_3, \\
Y_2^{(9)} &= \left(Y_{2''}^{(1)} Y_{1'}^{(8)} \right)_2, & Y_{2'i}^{(9)} &= \left(Y_{2''}^{(1)} Y_{1''}^{(8)} \right)_{2'}, & Y_{2''ii}^{(9)} &= \left(Y_{2''}^{(1)} Y_{3i}^{(8)} \right)_{2''}, \\
Y_{2''i}^{(9)} &= \left(Y_{2''}^{(1)} Y_1^{(8)} \right)_{2''}, & Y_{2''ii}^{(9)} &= \left(Y_{2''}^{(1)} Y_{3ii}^{(8)} \right)_{2''}, & Y_1^{(10)} &= \left(Y_3^{(2)} Y_{3i}^{(8)} \right)_1, \\
Y_{1'}^{(10)} &= \left(Y_3^{(2)} Y_{3ii}^{(8)} \right)_{1'}, & Y_{3i}^{(10)} &= \left(Y_3^{(2)} Y_1^{(8)} \right)_3, & Y_{3ii}^{(10)} &= \left(Y_3^{(2)} Y_{1'}^{(8)} \right)_3, \\
Y_{3iii}^{(10)} &= \left(Y_3^{(2)} Y_{1''}^{(8)} \right)_3.
\end{aligned} \tag{C.10}$$

D Alternative method to derive the Majorana superpotential

As shown in section 4, it is convenient to first consider the constraints of traditional flavor symmetry and then the modular symmetry in the paradigm of eclectic flavor symmetry. Likewise we can first analyze the constraints of modular symmetry and subsequently the traditional flavor symmetry. In the following, we shall revisit the superpotential \mathcal{W}_M for the Majorana fermion mass in this alternative method. The general expression of \mathcal{W}_M is given in eq. (4.16), where ψ^c and Φ are assumed to transform as $(\mathbf{3}, \mathbf{3}_0)$ under the eclectic symmetry group $\Delta(27) \times T'$.

Let us first study the T' invariance of the superpotential \mathcal{W}_M . Notice that $\mathbf{3}_0$ can be decomposed into a singlet $\mathbf{1}'$ and a doublet $\mathbf{2}''$ of T' with $\mathbf{3}_0 = \mathbf{1}' \oplus \mathbf{2}''$, as shown in

eq. (3.24). With the modular form multiplets defined in eq. (4.1), the T' invariant terms of \mathcal{W}_M result from the following T' tensor products

$$\mathbf{1} \subset (\mathbf{1} \oplus \mathbf{1}' \oplus \mathbf{1}'' \oplus \mathbf{2} \oplus \mathbf{2}' \oplus \mathbf{2}'' \oplus \mathbf{3}) \otimes (\mathbf{1}' \oplus \mathbf{2}'') \otimes (\mathbf{1}' \oplus \mathbf{2}'') \otimes (\mathbf{1}' \oplus \mathbf{2}'') . \quad (\text{D.1})$$

From the Kronecker product of T' in eq. (B.3) we know

$$(\mathbf{1}' \oplus \mathbf{2}'') \otimes (\mathbf{1}' \oplus \mathbf{2}'') = \mathbf{1}'' \oplus \mathbf{2} \oplus \mathbf{2} \oplus \mathbf{1}' \oplus \mathbf{3} . \quad (\text{D.2})$$

Since the contraction $(\psi^c \psi^c)_{\mathbf{1}'} = 0$ and the two doublet contractions $(\psi^c \psi^c)_{\mathbf{2}}$ should be symmetrized, the non-zero T' contraction of $(\psi^c \psi^c)_r$ are

$$(\mathbf{1}' \oplus \mathbf{2}'') \otimes (\mathbf{1}' \oplus \mathbf{2}'') = \mathbf{1}'' \oplus \mathbf{2} \oplus \mathbf{3} . \quad (\text{D.3})$$

Consequently the trivial singlet $\mathbf{1}$ resulting from the tensor products of T' representations are

$$\begin{aligned} \mathbf{1} &\subset (\mathbf{1} \oplus \mathbf{1}' \oplus \mathbf{1}'' \oplus \mathbf{2} \oplus \mathbf{2}' \oplus \mathbf{2}'' \oplus \mathbf{3}) \otimes (\mathbf{1}'' \oplus \mathbf{2} \oplus \mathbf{3}) \otimes (\mathbf{1}' \oplus \mathbf{2}'') \\ &\subset (\mathbf{1} \oplus \mathbf{1}' \oplus \mathbf{1}'' \oplus \mathbf{2} \oplus \mathbf{2}' \oplus \mathbf{2}'' \oplus \mathbf{3}) \otimes (\mathbf{1} \oplus \mathbf{2}' \oplus \mathbf{3} \oplus \mathbf{2}' \oplus \mathbf{1}'' \oplus \mathbf{3} \oplus \mathbf{2} \oplus \mathbf{2}' \oplus \mathbf{2}'') . \end{aligned} \quad (\text{D.4})$$

Hence, we can obtain nine T' invariant contractions when the modular form multiplets run over all possible irreducible multiplets of T' . The results are given by

$$\begin{aligned} \mathcal{W}_M &= \alpha_1 Y_1 (\psi_1^c - \psi_2^c)^2 (\phi_1 - \phi_2) - 2\sqrt{2}\omega\alpha_2 Y_2 (\psi_1^c - \psi_2^c) [\phi_3 (\psi_1^c + \psi_2^c) - \psi_3^c (\phi_1 + \phi_2)] \\ &\quad + \alpha_3 \left(\sqrt{2}\psi_3^c Y_5 - \omega^2 Y_4 (\psi_1^c + \psi_2^c) \right) (\psi_3^c (\phi_1 + \phi_2) - \phi_3 (\psi_1^c + \psi_2^c)) \\ &\quad + \alpha_4 \left[2\psi_3^c Y_6 \phi_3 (\psi_1^c + \psi_2^c) - \frac{\omega Y_7 (\psi_1^c + \psi_2^c)^2 (\phi_1 + \phi_2)}{\sqrt{2}} + \psi_3^{c2} \left(Y_6 (\phi_1 + \phi_2) + \sqrt{2}\omega Y_7 \phi_3 \right) \right] \\ &\quad - \frac{\alpha_5}{\sqrt{2}} (\psi_1^c - \psi_2^c)^2 \left[\sqrt{2} Y_9 \phi_3 - \omega^2 Y_8 (\phi_1 + \phi_2) \right] \\ &\quad + 2\alpha_6 (\psi_1^c - \psi_2^c) (\phi_1 - \phi_2) \left[Y_8 (\psi_1^c + \psi_2^c) - \sqrt{2}\omega\psi_3^c Y_9 \right] \\ &\quad + \frac{\alpha_7}{4} \left[\sqrt{2} Y_8 \left((\psi_1^c + \psi_2^c)^2 (\phi_1 + \phi_2) + 4\psi_3^{c2} \phi_3 \right) + 2\omega Y_9 (\psi_1^c + \psi_2^c) (\phi_3 (\psi_1^c + \psi_2^c) + 2\psi_3^c (\phi_1 + \phi_2)) \right] \\ &\quad + 2\sqrt{2}\alpha_8 (\psi_1^c - \psi_2^c) \left[\omega Y_{11} (\phi_3 (\psi_1^c + \psi_2^c) + \psi_3^c (\phi_1 + \phi_2)) + Y_{10} (\psi_1^c + \psi_2^c) (\phi_1 + \phi_2) - 2\omega^2 \psi_3^c Y_{12} \phi_3 \right] \\ &\quad + \frac{\alpha_9}{2} (\phi_1 - \phi_2) \left[-2\omega^2 \psi_3^c Y_{11} (\psi_1^c + \psi_2^c) - \omega Y_{10} (\psi_1^c + \psi_2^c)^2 + 42\psi_3^{c2} Y_{12} \right] . \end{aligned} \quad (\text{D.5})$$

Furthermore, invariance under $\Delta(27)$ requires that \mathcal{W}_M must be a trivial singlet $\mathbf{1}_{\mathbf{0},\mathbf{0}}$ of $\Delta(27)$. As both matter field ψ^c and flavon field Φ are assigned to $\Delta(27)$ triplet $\mathbf{3}$, while the modular forms are invariant under the actions of $\Delta(27)$. Consequently the trivial singlet $\mathbf{1}_{\mathbf{0},\mathbf{0}}$ resulting from the tensor products of $\Delta(27)$ representations are

$$\mathbf{1}_{\mathbf{0},\mathbf{0}} \subset \mathbf{3} \otimes \mathbf{3} \otimes \mathbf{3} = \left(\bar{\mathbf{3}}_{S,1} \oplus \bar{\mathbf{3}}_{S,2} \oplus \bar{\mathbf{3}}_A \right) \otimes \mathbf{3} = \left(\bar{\mathbf{3}}_{S,1} \oplus \bar{\mathbf{3}}_{S,2} \right) \otimes \mathbf{3} .$$

Hence there are only two $\Delta(27)$ invariant terms in \mathcal{W}_M ,

$$\mathcal{W}_M = \sum_{i=1}^{12} Y_i \left[a_i \left((\psi_1^c)^2 \phi_1 + (\psi_2^c)^2 \phi_2 + (\psi_3^c)^2 \phi_3 \right) + a'_i (\psi_1^c \psi_2^c \phi_3 + \psi_1^c \psi_3^c \phi_2 + \psi_2^c \psi_3^c \phi_1) \right] . \quad (\text{D.6})$$

We find that the $\Delta(27)$ flavor symmetry imposes the following restrictions on the coefficients α_i in eq. (D.5):

$$\alpha_1 = \alpha_2 = \alpha_3 = \alpha_4 = \alpha_8 = \alpha_9 = 0, \quad \alpha_5 = \sqrt{2}\omega\alpha_6 = \frac{\omega}{2}\alpha_7 \equiv \omega\alpha. \quad (\text{D.7})$$

Thus the superpotential \mathcal{W}_M reduces to

$$\mathcal{W}_M = \alpha \left[\sqrt{2}Y_8 \left((\psi_1^c)^2 \phi_1 + (\psi_2^c)^2 \phi_2 + (\psi_3^c)^2 \phi_3 \right) + 2\omega Y_9 (\psi_1^c \psi_2^c \phi_3 + \psi_1^c \psi_3^c \phi_2 + \psi_2^c \psi_3^c \phi_1) \right], \quad (\text{D.8})$$

where the complex coupling α is enforced to be real by the gCP transformation K_* . We obtain the same superpotential \mathcal{W}_M as that of eq. (4.21), and only the modular form doublet $Y_{\mathbf{2}''}^{(k_Y)}(\tau) = (Y_8, Y_9)^T$ gives non-vanishing contribution.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited. SCOAP³ supports the goals of the International Year of Basic Sciences for Sustainable Development.

References

- [1] F. Feruglio, *Are neutrino masses modular forms?*, in *From My Vast Repertoire...: Guido Altarelli's Legacy*, A. Levy et al. eds., World Scientific (2019), p. 227–266 [[DOI:10.1142/9789813238053_0012](https://doi.org/10.1142/9789813238053_0012)] [[arXiv:1706.08749](https://arxiv.org/abs/1706.08749)] [[INSPIRE](#)].
- [2] F. Feruglio and A. Romanino, *Lepton flavor symmetries*, *Rev. Mod. Phys.* **93** (2021) 015007 [[arXiv:1912.06028](https://arxiv.org/abs/1912.06028)] [[INSPIRE](#)].
- [3] G.-J. Ding, X.-G. Liu and C.-Y. Yao, *A minimal modular invariant neutrino model*, *JHEP* **01** (2023) 125 [[arXiv:2211.04546](https://arxiv.org/abs/2211.04546)] [[INSPIRE](#)].
- [4] F. Feruglio, *The irresistible call of $\tau = i$* , [arXiv:2211.00659](https://arxiv.org/abs/2211.00659) [[INSPIRE](#)].
- [5] F. Feruglio, *Fermion masses, critical behavior and universality*, *JHEP* **03** (2023) 236 [[arXiv:2302.11580](https://arxiv.org/abs/2302.11580)] [[INSPIRE](#)].
- [6] M.-C. Chen, S. Ramos-Sánchez and M. Ratz, *A note on the predictions of models with modular flavor symmetries*, *Phys. Lett. B* **801** (2020) 135153 [[arXiv:1909.06910](https://arxiv.org/abs/1909.06910)] [[INSPIRE](#)].
- [7] J.-N. Lu, X.-G. Liu and G.-J. Ding, *Modular symmetry origin of texture zeros and quark lepton unification*, *Phys. Rev. D* **101** (2020) 115020 [[arXiv:1912.07573](https://arxiv.org/abs/1912.07573)] [[INSPIRE](#)].
- [8] A. Baur, H.P. Nilles, A. Trautner and P.K.S. Vaudrevange, *Unification of Flavor, CP, and Modular Symmetries*, *Phys. Lett. B* **795** (2019) 7 [[arXiv:1901.03251](https://arxiv.org/abs/1901.03251)] [[INSPIRE](#)].
- [9] A. Baur, H.P. Nilles, A. Trautner and P.K.S. Vaudrevange, *A String Theory of Flavor and CP*, *Nucl. Phys. B* **947** (2019) 114737 [[arXiv:1908.00805](https://arxiv.org/abs/1908.00805)] [[INSPIRE](#)].
- [10] T. Kobayashi, S. Nagamoto and S. Uemura, *Modular symmetry in magnetized/intersecting D-brane models*, *PTEP* **2017** (2017) 023B02 [[arXiv:1608.06129](https://arxiv.org/abs/1608.06129)] [[INSPIRE](#)].
- [11] T. Kobayashi et al., *Modular symmetry and non-Abelian discrete flavor symmetries in string compactification*, *Phys. Rev. D* **97** (2018) 116002 [[arXiv:1804.06644](https://arxiv.org/abs/1804.06644)] [[INSPIRE](#)].
- [12] T. Kobayashi and S. Tamba, *Modular forms of finite modular subgroups from magnetized D-brane models*, *Phys. Rev. D* **99** (2019) 046001 [[arXiv:1811.11384](https://arxiv.org/abs/1811.11384)] [[INSPIRE](#)].

- [13] Y. Kariyazono et al., *Modular symmetry anomaly in magnetic flux compactification*, *Phys. Rev. D* **100** (2019) 045014 [[arXiv:1904.07546](#)] [[INSPIRE](#)].
- [14] H. Ohki, S. Uemura and R. Watanabe, *Modular flavor symmetry on a magnetized torus*, *Phys. Rev. D* **102** (2020) 085008 [[arXiv:2003.04174](#)] [[INSPIRE](#)].
- [15] S. Kikuchi et al., *Revisiting modular symmetry in magnetized torus and orbifold compactifications*, *Phys. Rev. D* **102** (2020) 105010 [[arXiv:2005.12642](#)] [[INSPIRE](#)].
- [16] Y. Almumin et al., *Metaplectic Flavor Symmetries from Magnetized Tori*, *JHEP* **05** (2021) 078 [[arXiv:2102.11286](#)] [[INSPIRE](#)].
- [17] H.P. Nilles, S. Ramos-Sánchez and P.K.S. Vaudrevange, *Eclectic Flavor Groups*, *JHEP* **02** (2020) 045 [[arXiv:2001.01736](#)] [[INSPIRE](#)].
- [18] H.P. Nilles, S. Ramos-Sánchez and P.K.S. Vaudrevange, *Lessons from eclectic flavor symmetries*, *Nucl. Phys. B* **957** (2020) 115098 [[arXiv:2004.05200](#)] [[INSPIRE](#)].
- [19] H.P. Nilles, S. Ramos-Sánchez and P.K.S. Vaudrevange, *Eclectic flavor scheme from ten-dimensional string theory — I. Basic results*, *Phys. Lett. B* **808** (2020) 135615 [[arXiv:2006.03059](#)] [[INSPIRE](#)].
- [20] H.P. Nilles, S. Ramos-Sánchez and P.K.S. Vaudrevange, *Eclectic flavor scheme from ten-dimensional string theory — II detailed technical analysis*, *Nucl. Phys. B* **966** (2021) 115367 [[arXiv:2010.13798](#)] [[INSPIRE](#)].
- [21] S.F. King and I.N.R. Peddie, *Canonical normalization and Yukawa matrices*, *Phys. Lett. B* **586** (2004) 83 [[hep-ph/0312237](#)] [[INSPIRE](#)].
- [22] S.F. King et al., *Kähler corrections and softly broken family symmetries*, *JHEP* **07** (2005) 049 [[hep-ph/0407012](#)] [[INSPIRE](#)].
- [23] S. Antusch, S.F. King and M. Malinsky, *Third Family Corrections to Quark and Lepton Mixing in SUSY Models with non-Abelian Family Symmetry*, *JHEP* **05** (2008) 066 [[arXiv:0712.3759](#)] [[INSPIRE](#)].
- [24] M.-C. Chen, M. Fallbacher, M. Ratz and C. Staudt, *On predictions from spontaneously broken flavor symmetries*, *Phys. Lett. B* **718** (2012) 516 [[arXiv:1208.2947](#)] [[INSPIRE](#)].
- [25] M.-C. Chen et al., *Predictivity of models with spontaneously broken non-Abelian discrete flavor symmetries*, *Nucl. Phys. B* **873** (2013) 343 [[arXiv:1302.5576](#)] [[INSPIRE](#)].
- [26] M.-C. Chen et al., *Quasi-eclectic modular flavor symmetries*, *Phys. Lett. B* **824** (2022) 136843 [[arXiv:2108.02240](#)] [[INSPIRE](#)].
- [27] A. Baur et al., *The first string-derived eclectic flavor model with realistic phenomenology*, *JHEP* **09** (2022) 224 [[arXiv:2207.10677](#)] [[INSPIRE](#)].
- [28] The GAP Group, *GAP — Groups, Algorithms, and Programming, Version 4.12.2*, (2022), <https://www.gap-system.org>.
- [29] X.-G. Liu and G.-J. Ding, *Neutrino Masses and Mixing from Double Covering of Finite Modular Groups*, *JHEP* **08** (2019) 134 [[arXiv:1907.01488](#)] [[INSPIRE](#)].
- [30] R. de Adelhart Toorop, F. Feruglio and C. Hagedorn, *Finite Modular Groups and Lepton Mixing*, *Nucl. Phys. B* **858** (2012) 437 [[arXiv:1112.1340](#)] [[INSPIRE](#)].
- [31] P.P. Novichkov, J.T. Penedo, S.T. Petcov and A.V. Titov, *Generalised CP Symmetry in Modular-Invariant Models of Flavour*, *JHEP* **07** (2019) 165 [[arXiv:1905.11970](#)] [[INSPIRE](#)].

- [32] G.-J. Ding, F. Feruglio and X.-G. Liu, *CP symmetry and symplectic modular invariance*, *SciPost Phys.* **10** (2021) 133 [[arXiv:2102.06716](#)] [[INSPIRE](#)].
- [33] B.-Y. Qu, X.-G. Liu, P.-T. Chen and G.-J. Ding, *Flavor mixing and CP violation from the interplay of an S_4 modular group and a generalized CP symmetry*, *Phys. Rev. D* **104** (2021) 076001 [[arXiv:2106.11659](#)] [[INSPIRE](#)].
- [34] W. Grimus and M.N. Rebelo, *Automorphisms in gauge theories and the definition of CP and P*, *Phys. Rept.* **281** (1997) 239 [[hep-ph/9506272](#)] [[INSPIRE](#)].
- [35] F. Feruglio, C. Hagedorn and R. Ziegler, *Lepton Mixing Parameters from Discrete and CP Symmetries*, *JHEP* **07** (2013) 027 [[arXiv:1211.5560](#)] [[INSPIRE](#)].
- [36] G.-J. Ding, S.F. King, C. Luhn and A.J. Stuart, *Spontaneous CP violation from vacuum alignment in S_4 models of leptons*, *JHEP* **05** (2013) 084 [[arXiv:1303.6180](#)] [[INSPIRE](#)].
- [37] M. Holthausen, M. Lindner and M.A. Schmidt, *CP and Discrete Flavour Symmetries*, *JHEP* **04** (2013) 122 [[arXiv:1211.6953](#)] [[INSPIRE](#)].
- [38] M.-C. Chen et al., *CP Violation from Finite Groups*, *Nucl. Phys. B* **883** (2014) 267 [[arXiv:1402.0507](#)] [[INSPIRE](#)].
- [39] K. Ishiguro, T. Kobayashi and H. Otsuka, *Landscape of Modular Symmetric Flavor Models*, *JHEP* **03** (2021) 161 [[arXiv:2011.09154](#)] [[INSPIRE](#)].
- [40] P.P. Novichkov, J.T. Penedo and S.T. Petcov, *Modular flavour symmetries and modulus stabilisation*, *JHEP* **03** (2022) 149 [[arXiv:2201.02020](#)] [[INSPIRE](#)].
- [41] P. Chen, C.-C. Li and G.-J. Ding, *Lepton Flavor Mixing and CP Symmetry*, *Phys. Rev. D* **91** (2015) 033003 [[arXiv:1412.8352](#)] [[INSPIRE](#)].
- [42] P. Chen, C.-Y. Yao and G.-J. Ding, *Neutrino Mixing from CP Symmetry*, *Phys. Rev. D* **92** (2015) 073002 [[arXiv:1507.03419](#)] [[INSPIRE](#)].
- [43] I. Esteban et al., *The fate of hints: updated global analysis of three-flavor neutrino oscillations*, *JHEP* **09** (2020) 178 [[arXiv:2007.14792](#)] [[INSPIRE](#)].
- [44] S. Antusch and V. Maurer, *Running quark and lepton parameters at various scales*, *JHEP* **11** (2013) 115 [[arXiv:1306.6879](#)] [[INSPIRE](#)].
- [45] KAMLAND-ZEN collaboration, *Search for the Majorana Nature of Neutrinos in the Inverted Mass Ordering Region with KamLAND-Zen*, *Phys. Rev. Lett.* **130** (2023) 051801 [[arXiv:2203.02139](#)] [[INSPIRE](#)].
- [46] SNO+ collaboration, *Current Status and Future Prospects of the SNO+ Experiment*, *Adv. High Energy Phys.* **2016** (2016) 6194250 [[arXiv:1508.05759](#)] [[INSPIRE](#)].
- [47] LEGEND collaboration, *The Large Enriched Germanium Experiment for Neutrinoless Double Beta Decay (LEGEND)*, *AIP Conf. Proc.* **1894** (2017) 020027 [[arXiv:1709.01980](#)] [[INSPIRE](#)].
- [48] NEXO collaboration, *Sensitivity and Discovery Potential of nEXO to Neutrinoless Double Beta Decay*, *Phys. Rev. C* **97** (2018) 065503 [[arXiv:1710.05075](#)] [[INSPIRE](#)].
- [49] PLANCK collaboration, *Planck 2018 results. VI. Cosmological parameters*, *Astron. Astrophys.* **641** (2020) A6 [*Erratum ibid.* **652** (2021) C4] [[arXiv:1807.06209](#)] [[INSPIRE](#)].
- [50] P.F. Harrison and W.G. Scott, *Symmetries and generalizations of tri-bimaximal neutrino mixing*, *Phys. Lett. B* **535** (2002) 163 [[hep-ph/0203209](#)] [[INSPIRE](#)].

- [51] W. Grimus and L. Lavoura, *A Nonstandard CP transformation leading to maximal atmospheric neutrino mixing*, *Phys. Lett. B* **579** (2004) 113 [[hep-ph/0305309](#)] [[INSPIRE](#)].
- [52] Y. Farzan and A.Y. Smirnov, *Leptonic CP violation: Zero, maximal or between the two extremes*, *JHEP* **01** (2007) 059 [[hep-ph/0610337](#)] [[INSPIRE](#)].
- [53] DUNE collaboration, *Long-Baseline Neutrino Facility (LBNF) and Deep Underground Neutrino Experiment (DUNE): Conceptual Design Report, Volume 2: The Physics Program for DUNE at LBNF*, [arXiv:1512.06148](#) [[INSPIRE](#)].
- [54] HYPER-KAMIOKANDE PROTO- collaboration, *Physics potential of a long-baseline neutrino oscillation experiment using a J-PARC neutrino beam and Hyper-Kamiokande*, *PTEP* **2015** (2015) 053C02 [[arXiv:1502.05199](#)] [[INSPIRE](#)].
- [55] I. de Medeiros Varzielas, S.F. King and G.G. Ross, *Neutrino tri-bi-maximal mixing from a non-Abelian discrete family symmetry*, *Phys. Lett. B* **648** (2007) 201 [[hep-ph/0607045](#)] [[INSPIRE](#)].
- [56] G.C. Branco, I. de Medeiros Varzielas and S.F. King, *Invariant approach to CP in unbroken $\Delta(27)$* , *Nucl. Phys. B* **899** (2015) 14 [[arXiv:1505.06165](#)] [[INSPIRE](#)].
- [57] I. de Medeiros Varzielas, *$\Delta(27)$ family symmetry and neutrino mixing*, *JHEP* **08** (2015) 157 [[arXiv:1507.00338](#)] [[INSPIRE](#)].
- [58] P. Chen et al., *Warped flavor symmetry predictions for neutrino physics*, *JHEP* **01** (2016) 007 [[arXiv:1509.06683](#)] [[INSPIRE](#)].
- [59] H. Ishimori et al., *Non-Abelian Discrete Symmetries in Particle Physics*, *Prog. Theor. Phys. Suppl.* **183** (2010) 1 [[arXiv:1003.3552](#)] [[INSPIRE](#)].
- [60] C.-C. Li, X.-G. Liu and G.-J. Ding, *Modular symmetry at level 6 and a new route towards finite modular groups*, *JHEP* **10** (2021) 238 [[arXiv:2108.02181](#)] [[INSPIRE](#)].
- [61] G.-J. Ding, F.R. Joaquim and J.-N. Lu, *Texture-zero patterns of lepton mass matrices from modular symmetry*, *JHEP* **03** (2023) 141 [[arXiv:2211.08136](#)] [[INSPIRE](#)].