

Localization and attraction

Pietro Benetti Genolini ^a, Jerome P. Gauntlett ^b, Yusheng Jiao ^b, Alice Lüscher^c
and James Sparks^c

^a*Département de Physique Théorique, Université de Genève,
24 quai Ernest-Ansermet, 1211 Genève, Suisse*

^b*Blackett Laboratory, Imperial College,
Prince Consort Rd., London, SW7 2AZ, U.K.*

^c*Mathematical Institute, University of Oxford,
Andrew Wiles Building, Radcliffe Observatory Quarter,
Woodstock Road, Oxford, OX2 6GG, U.K.*

E-mail: Pietro.benettigenolini@unige.ch, j.gauntlett@imperial.ac.uk,
yusheng.jiao17@imperial.ac.uk, Alice.Luscher@maths.ox.ac.uk,
sparks@maths.ox.ac.uk

ABSTRACT: We use equivariant localization to construct off-shell entropy functions for supersymmetric black holes in $\mathcal{N} = 2$, $D = 4$ gauged supergravity coupled to matter. This allows one to compute the black hole entropy without solving the supergravity equations of motion and provides a novel generalization of the attractor mechanism. We consider magnetically charged black holes in AdS_4 which have an $AdS_2 \times M_2$ near horizon geometry, where M_2 is a sphere or a spindle, and we also obtain entropy functions for ungauged supergravity as a simple corollary. We derive analogous results for black strings and rings in $D = 5$ supergravity which have an $AdS_3 \times M_2$ near horizon geometry, and in this setting we derive an off-shell expression for the central charge of the dual $\mathcal{N} = (0, 2)$, $d = 2$ SCFT.

KEYWORDS: AdS-CFT Correspondence, Supergravity Models, Black Holes in String Theory

ARXIV EPRINT: [2401.10977](https://arxiv.org/abs/2401.10977)

Contents

1	Introduction	2
2	$D = 5, \mathcal{N} = 2$ gauged supergravity	6
2.1	The AdS_3 ansatz	7
2.2	Equivariantly closed forms	8
2.3	Localization	9
2.4	The $b_0 \rightarrow 0$ limit	11
2.5	STU model	12
2.6	Uplifting to $AdS_3 \times Y_7$ solutions	13
3	$D = 5$ STU model with hypermultiplets	15
3.1	The AdS_3 ansatz	17
3.2	Equivariantly closed forms and localization	17
4	$D = 5, \mathcal{N} = 2$ ungauged supergravity	20
5	$D = 4, \mathcal{N} = 2$ gauged supergravity	22
5.1	The AdS_2 ansatz	24
5.2	Equivariantly closed forms and localization	25
5.3	STU model	28
5.4	Uplifting to $AdS_2 \times Y_9$ solutions	30
6	$D = 4$ STU model with a hypermultiplet	31
6.1	The AdS_2 ansatz	32
6.2	Equivariantly closed forms and localization	33
7	$D = 4, \mathcal{N} = 2$ ungauged supergravity	36
8	Discussion	38
A	Reduction of $D = 5$ Killing spinors	39
A.1	Killing spinor equations	39
A.2	Bilinear equations	40
A.3	STU with hypermultiplet	40
B	Reduction of $D = 4$ Killing spinors	42
B.1	Killing spinor equations	43
B.2	Bilinear equations	44
B.3	STU with hypermultiplet	45

1 Introduction

It has recently been shown that supersymmetric solutions of supergravity with an R-symmetry Killing vector are equipped with a set of equivariantly closed forms which can be constructed from bilinears in the Killing spinor [1–3]. The localization fixed point formula [4, 5] then allows one to compute various physical observables without needing to solve the supergravity equations of motion. Furthermore, the formalism naturally gives rise to results that are off-shell with, generically, a simple extremization required in order to get the final result. Within the context of holography this latter feature is precisely dual to an extremization problem in the corresponding supersymmetric conformal field theory.

Here we further investigate the new equivariant calculus in the context of magnetically charged black holes of $\mathcal{N} = 2$, $D = 4$ gauged supergravity and black strings (or rings) of $D = 5$ gauged supergravity. More precisely, by analysing the near horizon geometries we are able to derive novel off-shell entropy functions, and furthermore provide a new perspective on the attractor mechanism. In both cases we allow for arbitrary numbers of vector multiplets and we also consider hypermultiplets. Interestingly, off-shell entropy functions for the associated *ungauged* supergravity theories will also follow as simple corollaries.

We now elaborate a little, first discussing the case of black holes in $\mathcal{N} = 2$, $D = 4$ gauged supergravity. By analysing spinor bilinears, we construct an off-shell entropy function for classes of supersymmetric AdS_4 black holes which have an $AdS_2 \times M_2$ near horizon geometry and carry magnetic charge on M_2 . We take M_2 to be topologically a two-sphere or a spindle with an azimuthal symmetry. The off-shell entropy function, explicitly given in (1.1) below, depends on the magnetic charges, as well as the values of certain scalar fields and a metric warp function at the two poles of the horizon M_2 . Extremizing over this data then gives the entropy of the black hole solution, provided the solution actually exists.¹ This extremization problem is actually implied by extremizing the action itself, and hence our results can be viewed as a kind of supersymmetric refinement of the approach of Sen [6].

Our results also provide a new way of thinking about some aspects of the attractor mechanism [7–9], which can be viewed² as a way of characterizing the near horizon geometry as the end-point of a flow, as well as providing a generalization thereof. Consider, for example, static, magnetically charged black holes with spherical horizons and with supersymmetry preserved via a topological twist. A standard procedure is to assume that the sphere³ has a constant curvature metric which allows one to construct an ansatz for the metric and matter fields that only depends on a radial variable. Preservation of supersymmetry then leads

¹To be more precise: our extremization procedure gives a result for the $AdS_2 \times M_2$ solution, assuming it exists. If, furthermore, this solution arises as the near horizon limit of an AdS_4 black hole solution, it will compute the entropy of the black hole.

²In ungauged supergravity the attractor mechanism also reveals that the near horizon geometry is independent of scalar moduli at the asymptotically flat boundary. By contrast in gauged supergravity the values of the scalars at the asymptotic AdS boundary are generically fixed by the potential. The attractor mechanism has been extensively investigated in gauged supergravity; see [10] for a recent discussion. For the black holes we focus on here, some early references include [11–13]. A review of the attractor mechanism in ungauged $D = 5$ supergravity is given in [14].

³One can also consider constant curvature metrics on other Riemann surfaces, but as they do not have suitable Killing vectors our new equivariant techniques do not provide further insight.

to a set of BPS equations, which consist of a set of first order flow equations in the radial variable. Assuming one approaches an $AdS_2 \times S^2$ near horizon solution, the scalars and warp factor are necessarily constant at the horizon and the flow equations lead to conditions which express the warp factor and some of the constant scalars on the horizon in terms of the magnetic charges. Moreover, one can, in effect, obtain the entropy of the putative black holes by carrying out an extremization of an entropy function depending on the black hole horizon data, which has been shown, in the case of the STU model, to be dual to \mathcal{I} -extremization in the dual field theory [15, 16].

Here, instead, we consider a more general class of magnetically charged AdS_4 black holes. For spherical horizons, we just assume that the near horizon geometry has an $AdS_2 \times S^2$ factor with the S^2 having a Killing vector. That is, we do not assume the S^2 horizon has a constant curvature metric nor do we assume that the S^2 at the AdS_4 boundary has such a metric. Furthermore, while the near horizon geometry is static we do not need to assume the black hole is static.⁴ The presence of the Killing vector at the horizon is sufficient data to use our equivariant techniques to derive an off-shell entropy function, which is precisely the same as that arising in the standard attractor mechanism. Our approach therefore increases the scope of the attractor mechanism in that it covers all black hole solutions that approach $AdS_2 \times S^2$ in the near horizon, with the S^2 having a Killing vector. In particular, this allows for the possibility that the metric on the S^2 at the AdS_4 boundary is arbitrary, a possibility that was also investigated from a different point of view in [17].

We also consider black holes with spindle horizons [18]. Recall that a spindle is topologically a two-sphere but with quantized conical deficit angles at the two poles. Such black holes are known to arise when acceleration is present, with the conical deficits at the poles associated with the acceleration [19]. Since a spindle does not admit a constant curvature metric, the standard attractor mechanism cannot be deployed. However, assuming that the spindle has a Killing vector, we can again utilize equivariant localization to obtain an off-shell entropy function. Explicitly, for a spindle horizon, which locally looks like $\mathbb{R}^2/\mathbb{Z}_{n_{\pm}}$ at the two poles, the off-shell entropy function is given by

$$S_{\text{BH}} = \frac{\pi}{2G_4} \frac{1}{2b_0} \left[i\mathcal{F}(x_+^I) - \sigma i\mathcal{F}(x_-^I) \right]. \tag{1.1}$$

Here \mathcal{F} is the prepotential of the $\mathcal{N} = 2$, $D = 4$ gauged supergravity, with $i\mathcal{F}$ real, and $\sigma = \pm 1$ labels whether supersymmetry is preserved via a twist or an anti-twist [20]. The localization is performed with respect to an R-symmetry Killing vector bilinear ξ , which exists for any supersymmetric solution. Here we write

$$\xi = b_0 \partial_{\varphi}, \tag{1.2}$$

with ∂_{φ} generating rotations of the horizon, φ having canonical period 2π , and b_0 being a real parameter. The x_{\pm}^I are real scalar quantities evaluated at the two poles of the spindle,⁵

⁴This is particularly relevant in $D = 5$ supergravity, discussed below, which can admit non-static black ring solutions, but with locally $AdS_3 \times S^2$ horizons.

⁵The x_{\pm}^I are also values of equivariant first Chern classes at the two poles.

where the index I labels the number of vector multiplets present, and satisfy the constraints

$$\xi_I x_+^I = 2 - 2\frac{b_0}{n_+}, \quad \xi_I x_-^I = 2 + 2\sigma\frac{b_0}{n_-}, \quad \mathbf{p}^I = \frac{1}{2b_0}(x_+^I - x_-^I). \quad (1.3)$$

Here the constants $\xi_I \in \mathbb{R}$ determine the Fayet-Iliopoulos (FI) gauging of the theory, while \mathbf{p}^I are the magnetic charges. The black hole entropy is then obtained by extremizing (1.1) over the variables x_+^I, x_-^I, b_0 , subject to the constraints (1.3), where we hold the spindle data n_\pm, σ and magnetic charges \mathbf{p}^I fixed. Our results then provide a derivation of “gravitational block” formulae of gauged supergravity that have been conjectured in the literature [21] (see also [22–24]).

With regard to the results for S^2 horizons discussed above, we will find it most convenient to obtain them as a limiting case $b_0 \rightarrow 0$ of the results for spindle horizons, by setting $n_\pm = 1$ and $\sigma = 1$, and we obtain

$$S_{\text{BH}} = \frac{\pi}{2G_4} \mathbf{p}^I \frac{\partial}{\partial x^I} \mathcal{F}(x^I), \quad (1.4)$$

with the x^I and magnetic charges constrained via $\xi_I x^I = 2$ and $\xi_I \mathbf{p}^I = -2$, in agreement with [12] for purely electric gauging.

Of particular interest is the $D = 4$ STU model, an example of $\mathcal{N} = 2$ gauged supergravity coupled to three vector multiplets, since it can be obtained as a consistent truncation of $D = 11$ supergravity on S^7 . For the STU model our results for the off-shell entropy functions associated with $AdS_2 \times M_2$ horizons, where M_2 is a two-sphere or spindle, and three real scalar fields, precisely recover the known results in [15] and [23, 25, 26], respectively. We can also take a $D = 11$ perspective by uplifting on S^7 to obtain $AdS_2 \times M_9$ solutions of $D = 11$ supergravity, with M_9 being an S^7 bundle over M_2 and having a GK geometry [27, 28]. We then show that our new entropy functions (1.1) and (1.4) are equivalent to the entropy functions and gravitational block formulae that were derived within the context of GK geometry, using a higher-dimensional supergravity perspective⁶ and very different techniques for spherical horizons in [31, 32] and spindle horizons [33, 34], respectively.

We also extend our $D = 4$ analysis to study a particular model involving hypermultiplets. Specifically, we consider a model that extends the STU model with a complex scalar field, which also arises as a consistent truncation of $D = 11$ supergravity on S^7 [35, 36]. The model has two AdS_4 vacua, one dual to the ABJM theory and the other dual to the mABJM theory [37–41]. In this setting the new off-shell entropy function is actually the same as above, but with additional constraints on the x_\pm^I and fluxes \mathbf{p}^I associated with the fact that the charged fields in the hypermultiplet break some of the gauge symmetries. In particular, we recover the known off-shell entropy function for $AdS_2 \times S^2$ black hole horizons considered in [35]. For spindle horizons the new off-shell entropy function gives on-shell results consistent with those found for [36], which were obtained by a direct analysis of the BPS equations. Here we will also provide examples with properly quantized magnetic flux.

⁶Gravitational block formulae have also been obtained in a higher-dimensional context in [29, 30] using the equivariant volume of symplectic toric orbifolds. However, spinor bilinears and the associated canonical procedure for going off-shell that are utilised here and in [1–3] were not considered in [29, 30].

Having analysed gauged supergravity, we can obtain results for $\mathcal{N} = 2$, $D = 4$ ungauged supergravity by simply turning off the gauge parameters. We construct off-shell entropy functions that immediately eliminate the possibility of having spindle horizons, leaving just the possibility of $AdS_2 \times S^2$ near horizon geometry. We show how the entropy function for the standard attractor mechanism can then be recovered.

The above discussion focused on $\mathcal{N} = 2$, $D = 4$ gauged supergravity, but an analogous story unfolds for $D = 5$ gauged supergravities coupled to vector and hypermultiplets. In this case, associated with magnetically charged black strings (or rings) in AdS_5 with a horizon⁷ that is (locally) $AdS_3 \times M_2$, we derive an off-shell central charge function using the equivariant calculus, and this computes the central charge of the $d = 2$, $\mathcal{N} = (0, 2)$ SCFT dual to the $AdS_3 \times M_2$ solution. In a similar manner to the $D = 4$ case, our work generalizes the standard attractor mechanism equations for the case when M_2 is an S^2 and, in addition, it is also applicable to spindle horizons. Explicitly, for the spindle horizon, we find the off-shell central charge

$$c = -\frac{3\pi}{2G_5} \frac{1}{b_0} \left[\mathcal{F}(x_+^I) - \mathcal{F}(x_-^I) \right], \tag{1.5}$$

where \mathcal{F} is the real prepotential of the $D = 5$ gauged supergravity. The x_{\pm}^I now satisfy the constraints $\xi_I x_+^I = 2 - \frac{b_0}{n_+} P_+$ and $\xi_I x_-^I = 2 + \frac{b_0}{n_-} P_-$, where the constants $\xi_I \in \mathbb{R}$ determine the FI gauging of the theory, and $P_{\pm} = 1$ or $P_{\pm} = -1$ determine the chirality of the Killing spinor at the poles of the spindle with $\sigma = P_+/P_-$ then labelling twist and anti-twist. The magnetic charges are given by $\mathbf{p}^I = -\frac{1}{b_0}(x_+^I - x_-^I)$. For S^2 horizons we find

$$c = \frac{3\pi}{2G_5} \mathbf{p}^I \frac{\partial}{\partial x^I} \mathcal{F}(x^I), \tag{1.6}$$

with the x^I and magnetic charges constrained via $\xi_I x^I = 2$ and $\xi_I \mathbf{p}^I = 2P_+$, in agreement with [44].

For the special case of the $D = 5$ STU model we make contact with known results for S^2 [44] and the spindle [18, 20]. Furthermore, after uplifting on S^5 to type IIB we make contact with the off-shell central charge computed in GK geometry for $AdS_3 \times M_7$ solutions of type IIB with M_7 an S^5 bundle over M_2 , for spheres [45, 46] and spindles [34].

We also extend our $D = 5$ analysis to study a model involving hypermultiplets, finding results consistent with [47] for S^2 horizons. For spindle horizons we obtain an off-shell central charge that after extremization gives rise to the on-shell central charge found in [48] obtained just by manipulating the BPS equations.

Finally, we also obtain results for $D = 5$ ungauged supergravity by simply turning off the gauge parameters. We construct off-shell entropy functions that immediately eliminate the possibility of having spindle horizons, leaving just the possibility of $AdS_3 \times S^2$ near horizon geometry. We show how the entropy function for the standard attractor mechanism can then be recovered.

The plan of the rest of paper is as follows. Since the analysis is slightly simpler, we first discuss the five-dimensional examples before turning to those in four dimensions. In section 2

⁷Black rings of ungauged supergravity have a near horizon limit that is locally isometric to $AdS_3 \times S^2$ [42, 43]. Black rings in AdS_5 are not known, but if they exist our results can be utilized.

we discuss $D = 5$ gauged supergravity, coupled to an arbitrary number of Abelian vector multiplets, including the $D = 5$ STU model as a special case. Section 3 analyses a model that extends the $D = 5$ STU model with a hypermultiplet, and section 4 summarizes the results obtained for ungauged supergravity. We then move to the parallel analysis in four dimensions, discussing $\mathcal{N} = 2$, $D = 4$ gauged supergravity coupled to Abelian vector multiplets in section 5, including the $D = 4$ STU model as an example, and then the STU model with a hypermultiplet in section 6. Section 7 discusses $\mathcal{N} = 2$, $D = 4$ ungauged supergravity and we conclude with some discussion in section 8. The details of the construction of the equivariantly closed forms in terms of spinor bilinears are presented in the appendices A and B, in $D = 5$ and $D = 4$, respectively.

2 $D = 5$, $\mathcal{N} = 2$ gauged supergravity

We begin by considering $\mathcal{N} = 2$ gauged supergravity in $D = 5$ dimensions, coupled to an arbitrary number n of Abelian vector multiplets. This theory was originally constructed in [49], but we will largely follow the conventions of [50–52].

The bosonic part of the action is given by

$$S_5 = \frac{1}{16\pi G_5} \int_{M_5} \left(R_5 - \mathcal{V} - \mathcal{G}_{ij} \partial_\mu \varphi^i \partial^\mu \varphi^j - \frac{1}{2} G_{IJ} F_{\mu\nu}^I F^{J\mu\nu} + \frac{1}{24} C_{IJK} \varepsilon^{\mu\nu\rho\sigma\lambda} F_{\mu\nu}^I F_{\rho\sigma}^J A_\lambda^K \right) \text{vol}_5. \quad (2.1)$$

Here we use signature $(-1, 1, \dots, 1)$, with spacetime indices $\mu, \nu = 0, \dots, 4$, R_5 denotes the Ricci scalar, vol_5 is the Lorentzian volume form, and G_5 is the Newton constant. Recall that minimal $D = 5$, $\mathcal{N} = 2$ gauged supergravity contains a graviphoton A , which in (2.1) has been combined with the n additional vector multiplet gauge fields into A^I with field strengths $F^I = dA^I$, where $I = 0, 1, \dots, n$. The n real scalars φ^i , $i = 1, \dots, n$, have likewise been repackaged into $X^I = X^I(\varphi^i)$, where the latter satisfy the constraint

$$\mathcal{F}(X^I) \equiv \frac{1}{3!} C_{IJK} X^I X^J X^K = 1, \quad (2.2)$$

where \mathcal{F} is the prepotential with C_{IJK} a totally symmetric constant tensor, which also specifies the Chern-Simons coupling in (2.1).

The metrics for the kinetic terms for the gauge fields and scalars are respectively

$$G_{IJ} = -\frac{1}{2} \partial_I \partial_J \log \mathcal{F} \Big|_{\mathcal{F}=1}, \quad \mathcal{G}_{ij} = \partial_i X^I \partial_j X^J G_{IJ} \Big|_{\mathcal{F}=1}, \quad (2.3)$$

where we use the notation $\partial_I \equiv \partial_{X^I}$, $\partial_i \equiv \partial_{\varphi^i}$. The scalar potential is

$$\mathcal{V} \equiv \xi_I \xi_J \left(\mathcal{G}^{ij} \partial_i X^I \partial_j X^J - \frac{4}{3} X^I X^J \right), \quad (2.4)$$

where ξ_I are constant Fayet-Iliopoulos (FI) gauging parameters. This vector multiplet geometry is known as *very special geometry* [53], and the following definition and relations will prove useful for what follows:

$$X_I \equiv \frac{2}{3} G_{IJ} X^J, \quad \partial_i X_I = -\frac{2}{3} G_{IJ} \partial_i X^J, \quad (2.5)$$

together with the expression

$$G^{IJ} = \mathcal{G}^{ij} \partial_i X^I \partial_j X^J + \frac{2}{3} X^I X^J, \quad (2.6)$$

for the inverse of G_{IJ} . In particular note $X^I X_I = 1$ follows from the constraint (2.2).

A solution is supersymmetric if there is a non-trivial solution ϵ to the Killing spinor and gaugino equations:

$$\begin{aligned} 0 &= \left[\nabla_\mu - \frac{i}{2} \xi_I A_\mu^I + \frac{1}{6} W \Gamma_\mu + \frac{i}{8} X_I F_{\nu\rho}^I (\Gamma_\mu^{\nu\rho} - 4\delta_\mu^\nu \Gamma^\rho) \right] \epsilon, \\ 0 &= \left[-\frac{i}{2} \mathcal{G}_{ij} \partial_\mu \varphi^j \Gamma^\mu + \frac{i}{2} \partial_i W + \frac{3}{8} \partial_i X_I F_{\mu\nu}^I \Gamma^{\mu\nu} \right] \epsilon. \end{aligned} \quad (2.7)$$

Here Γ_μ generate Cliff(1,4) in an orthonormal frame, ∇_μ is the spin connection, and we have defined the superpotential

$$W \equiv \xi_I X^I. \quad (2.8)$$

Notice that we may then write the potential in (2.4) as

$$\mathcal{V} = \mathcal{G}^{ij} \partial_i W \partial_j W - \frac{4}{3} W^2. \quad (2.9)$$

If the $D = 5$ theory admits a supersymmetric AdS_5 solution with radius R , then it is dual to a $d = 4$, $\mathcal{N} = 1$ SCFT with central charge given (in the large N limit) by the usual formula

$$a = \frac{\pi R^3}{8G_5}. \quad (2.10)$$

2.1 The AdS_3 ansatz

We consider supersymmetric solutions of the warped product form

$$ds_5^2 = e^{2\lambda} \left[ds^2(AdS_3) + ds^2(M_2) \right]. \quad (2.11)$$

Here the metric on AdS_3 is taken to have unit radius, λ and the scalar fields are taken to be functions on M_2 , with the gauge fields A^I taken to be (pull-backs of) gauge fields on M_2 .

Inserting this ansatz into the $D = 5$ equations of motion gives rise to $D = 2$ equations of motion. The latter can be obtained by varying a two-dimensional action S_2 for the fields on M_2 , which can in turn be obtained by substituting the ansatz into the $D = 5$ action (2.1) and writing $S_5 = \frac{\text{Vol}(AdS_3)}{16\pi G_5} S_2$. Specifically, we have

$$\begin{aligned} S_2 &= \int_{M_2} \left[e^{3\lambda} \left(R - 6 + 12(\nabla\lambda)^2 - \mathcal{G}_{ij} \partial_\mu \varphi^i \partial^\mu \varphi^j \right) - e^{5\lambda} \mathcal{V} \right. \\ &\quad \left. - \frac{1}{2} e^\lambda G_{IJ} F_{\mu\nu}^I F^{J\mu\nu} \right] \text{vol}, \end{aligned} \quad (2.12)$$

where R , ∇ and vol are respectively the Ricci scalar, Levi-Civita connection and volume form on M_2 . The $D = 2$ Maxwell equation for A^I reads

$$d \left[e^\lambda G_{IJ} (*F^J) \right] = 0 \quad \implies \quad d \left[e^\lambda G_{IJ} F_{12}^J \right] = 0, \quad \forall I = 0, \dots, n. \quad (2.13)$$

Here we write $F^I = F_{12}^I \text{vol}$, where F_{12}^I is a function on M_2 . The equation of motion for the warp factor λ is

$$e^{3\lambda} \left[R - 6 + 12(\nabla\lambda)^2 - \mathcal{G}_{ij} \partial_\mu \varphi^i \partial^\mu \varphi^j \right] = \frac{5}{3} e^{5\lambda} \mathcal{V} + \frac{1}{6} e^\lambda G_{IJ} F_{\mu\nu}^I F^{J\mu\nu} + \text{total derivative}. \quad (2.14)$$

The trace of the $D = 2$ Einstein equation reads

$$\frac{2}{3} e^{5\lambda} \mathcal{V} - \frac{1}{3} e^\lambda G_{IJ} F_{\mu\nu}^I F^{J\mu\nu} = -4e^{3\lambda} + \text{total derivative}. \quad (2.15)$$

Imposing the equation of motion for the warp factor⁸ (2.14) then leads to the following form of the partially off-shell (POS)⁹ two-dimensional action:

$$S_2|_{\text{POS}} = \frac{2}{3} \int_{M_2} \left(e^{5\lambda} \mathcal{V} \text{vol} - e^\lambda G_{IJ} F_{12}^I F^J \right). \quad (2.16)$$

We next note that if we were to also impose the trace of the $D = 2$ Einstein equation (2.15) we would get the *on-shell* action

$$S_2|_{\text{OS}} = -4 \int_{M_2} e^{3\lambda} \text{vol}. \quad (2.17)$$

The central charge of the dual $d = 2$, $\mathcal{N} = (0, 2)$ SCFT is given by $c = 3/2G_3$, where G_3 is the $D = 3$ Newton constant. By analogy with c -extremization in field theory [54] we then introduce the “trial” central charge function

$$c = -\frac{3}{8G_5} S_2|_{\text{POS}} = -\frac{3a}{\pi R^3} S_2|_{\text{POS}}. \quad (2.18)$$

This has the property that for an on-shell AdS_3 solution c is the central charge. In the last equality we have used (2.10) to express the result for c in terms of the central charge a of the $D = 4$ SCFT (when the $D = 5$ model admits an AdS_5 vacuum).

Our goal is to compute the central charge of the $d = 2$ SCFT for any AdS_3 solution without having an explicit solution, but instead using localization. To do this we will implement localization using the partially off-shell action $S_2|_{\text{POS}}$ and then carry out a final residual extremization in order to get the on-shell action and hence central charge c .

2.2 Equivariantly closed forms

With the ansatz (2.11), the Killing spinor ϵ that solves (2.7) may be decomposed as

$$\epsilon = \vartheta \otimes e^{\lambda/2} \zeta. \quad (2.19)$$

Here ϑ is a Killing spinor on AdS_3 , ζ is a spinor on M_2 and the warp factor is included for convenience. The Killing spinor equation (A.3) can then be reduced, leading to a set of differential and algebraic equations that ζ must satisfy, as given in appendix A.

⁸Equivalently, this is the same as imposing the trace of the $D = 5$ Einstein equations.

⁹The acronym also stands for partially on-shell, for the more optimistic reader.

It is next convenient to introduce the following real bilinears in ζ :

$$S = \zeta^\dagger \zeta, \quad P = \zeta^\dagger \gamma_3 \zeta, \quad K = \zeta^\dagger \gamma_{(1)} \zeta, \quad \xi^b = -i \zeta^\dagger \gamma_{(1)} \gamma_3 \zeta. \quad (2.20)$$

Here we denote $\gamma_{(n)} = \frac{1}{n!} \gamma_{\mu_1 \dots \mu_n} dx^{\mu_1} \wedge \dots \wedge dx^{\mu_n}$, and $\gamma_3 = -i \gamma_1 \gamma_2$ is the chirality operator on M_2 . These bilinears satisfy a set of algebraic relations and differential conditions that follow from the reduced Killing spinor equations (A.3). Importantly, one deduces that the vector field ξ dual to the one-form ξ^b is a Killing vector on M_2 and we also have

$$d\xi^b = -2(2 + PS^{-1}e^\lambda W)P \text{ vol}. \quad (2.21)$$

In appendix A it is also shown that the multi-forms

$$\Phi^{F^I} \equiv F^I - X^I e^\lambda P, \quad (2.22)$$

as well as

$$\Phi^{\text{vol}} \equiv e^{5\lambda} \mathcal{V} \text{ vol} - e^{4\lambda} WS, \quad (2.23)$$

are both equivariantly closed under $d_\xi = d - \xi \lrcorner$. Using the two-dimensional Maxwell equation (2.13), we can immediately construct another equivariantly closed form,

$$\begin{aligned} \Phi^S &\equiv \frac{2}{3} \left(\Phi^{\text{vol}} - e^\lambda G_{IJ} F_{12}^I \Phi^{F^J} \right) \\ &= \frac{2}{3} \left(e^{5\lambda} \mathcal{V} \text{ vol} - e^\lambda G_{IJ} F_{12}^I F^J - e^{4\lambda} WS + e^{2\lambda} G_{IJ} F_{12}^I X^J P \right), \end{aligned} \quad (2.24)$$

and we notice the top-form is precisely that appearing in the partially off-shell two-dimensional action (2.16).

It is also shown in appendix A that the scalar bilinear S is constant, and without loss of generality we normalize $S \equiv 1$.

2.3 Localization

From the definitions (2.20) one can show that $\|\xi\|^2 = \|K\|^2 = S^2 - P^2$. At a point p where $\xi|_p = 0$ it immediately follows that $P|_p = \pm S|_p = \pm 1$, where this sign is the chirality of the spinor ζ at the point p . Provided ξ is not identically zero,¹⁰ near the point p we may model M_2 as $\mathbb{R}^2/\mathbb{Z}_{d_p}$, where we allow for potential orbifold singularities, and in these local coordinates write $\xi = \epsilon_p \partial_{\phi_p}$, where ϕ_p is a polar coordinate on $\mathbb{R}^2/\mathbb{Z}_{d_p}$ with period $2\pi/d_p$. Notice that the orientation here is determined by the global orientation on M_2 induced by the volume form. We refer to ϵ_p as the *weight* of ξ at the fixed point p .

¹⁰If ξ is identically zero then so is K , and $S \equiv P \equiv 1$. One quickly deduces from the equations in appendix A that λ and X^I are constant on M_2 , and localization plays no immediate role in this case. In particular this is the case for the topological twist solutions in which M_2 is a constant curvature Riemann surface. However, when $M_2 = S^2$ we are still able to obtain a non-trivial result for the off-shell action using localization by taking a limit of our fixed point formula, and also can recover the result that ξ is identically zero on S^2 — see section 2.4.

With this notation to hand, we may apply the Berline-Vergne-Atiyah-Bott fixed point formula to evaluate integrals of the equivariantly closed forms in (2.22) and (2.24) over M_2 . We have the magnetic charges¹¹

$$\mathfrak{p}^I \equiv \int_{M_2} \frac{F^I}{2\pi} = \int_{M_2} \frac{\Phi^{F^I}}{2\pi} = -\frac{1}{2\pi} \sum_{\text{fixed } p} \frac{1}{d_p} \frac{2\pi}{\epsilon_p} (X^I e^\lambda P) \Big|_p. \quad (2.25)$$

We may then similarly localize the action (2.16) using (2.24)

$$\begin{aligned} S_2|_{\text{POS}} &= \int_{M_2} \Phi^S = -\frac{2}{3} \sum_{\text{fixed } p} \frac{1}{d_p} \frac{2\pi}{\epsilon_p} \left(e^{4\lambda} W S - e^{2\lambda} G_{IJ} F_{12}^I X^J P \right) \Big|_p \\ &= 2 \sum_{\text{fixed } p} \frac{1}{d_p} \frac{2\pi}{\epsilon_p} e^{3\lambda} P \Big|_p, \end{aligned} \quad (2.26)$$

where in the last equality we have used the algebraic equation (A.7), and then used that $P^2|_p = S^2|_p = 1$. The final constraint comes from the equation for $d\xi^b$ in (2.21). Since at a fixed point $d\xi^b|_p = 2\epsilon_p \text{vol}$, in terms of the weight ϵ_p of ξ at p , we deduce that

$$e^\lambda W P|_p = -(2 + \epsilon_p P_p), \quad (2.27)$$

where recall that $P_p = \pm 1$, with the sign according to the spinor chirality at p .

In order to evaluate these expressions further, it is helpful to first introduce the new scalar fields

$$x^I \equiv -X^I e^\lambda P \quad \implies \quad \Phi^{F^I} = F^I + x^I. \quad (2.28)$$

Using the constraint $\mathcal{F}(X^I) = 1$ in (2.2), at a fixed point p we may then compute

$$\mathcal{F}(x^I)|_p = \frac{1}{3!} C_{IJK} x^I x^J x^K|_p = -e^{3\lambda} P|_p, \quad (2.29)$$

where we have again used $P^2|_p = 1$. We may then rewrite (2.26), (2.25), (2.27) as

$$S_2|_{\text{POS}} = -2 \sum_{\text{fixed } p} \frac{2\pi}{d_p \epsilon_p} \mathcal{F}(x_p^I), \quad \mathfrak{p}^I = \sum_{\text{fixed } p} \frac{1}{d_p \epsilon_p} x_p^I, \quad \xi_I x_p^I = 2 + \epsilon_p P_p, \quad (2.30)$$

respectively, where $x_p^I \equiv x^I|_p$. This makes manifest that the action may be computed knowing only the values of the scalar fields x_p^I at the fixed points, where these are constrained to satisfy the equations in (2.30), involving the magnetic charges \mathfrak{p}^I and the data d_p, ϵ_p, P_p at the fixed points.

If M_2 is compact without boundary, then the only such space admitting a $U(1)$ isometry with fixed points is topologically a spindle $M_2 \cong \mathbb{WCP}_{[n_+, n_-]}^1$.¹² Here there are two fixed

¹¹In general there is no reason to quantize the magnetic charges. For models that can be uplifted to string/M-theory, such as the STU model discussed in section 2.5, the gauge fields are associated with $U(1)$ bundles and there is a quantization condition.

¹²One could in principle have $M_2 = T^2$, but in this case any non-trivial Killing vector is fixed point free, and our formulae immediately imply that the magnetic charges P^I are zero, and the action is zero, ruling out such solutions. Note that this argument is not in contradiction with the existence of the T^2 solutions constructed in [55] — it only says that such solutions do not have a non-zero R-symmetry Killing vector along the T^2 . Unlike the case of S^2 discussed in footnote 10 and section 2.4, we cannot deduce more about the T^2 case using our approach.

points, which we label by $p = \pm$, with $d_{\pm} = n_{\pm}$ and we take $n_{\pm} \geq 1$. We may write the Killing vector field globally as

$$\xi = b_0 \partial_{\varphi}, \tag{2.31}$$

where φ is an azimuthal coordinate on the spindle with period 2π . Then without loss of generality we write $\epsilon_+ = -b_0/n_+$, $\epsilon_- = b_0/n_-$, but leave the signs of P_{\pm} arbitrary, where recall $|P_{\pm}| = 1$. The constraint equations in (2.30) now read

$$\mathbf{p}^I = -\frac{1}{b_0}(x_+^I - x_-^I), \quad \xi_I x_+^I = 2 - \frac{b_0}{n_+} P_+, \quad \xi_I x_-^I = 2 + \frac{b_0}{n_-} P_-, \tag{2.32}$$

and combining these equations one deduces

$$\xi_I \mathbf{p}^I = \frac{P_+}{n_+} + \frac{P_-}{n_-} = \frac{n_- P_+ + n_+ P_-}{n_- n_+}. \tag{2.33}$$

Since from (2.7) the Killing spinor ϵ is charged under precisely the combination $\xi_I A^I/2$, equation (2.33) gives the magnetic flux of the R-symmetry gauge field through the spindle M_2 . The form of the right hand side of (2.33) was proven very generally in [20]. Introducing $\sigma \equiv P_+/P_-$, the cases $\sigma = 1$, $\sigma = -1$ correspond to the spinor chirality at the two poles being either the same or opposite, respectively; in turn, these were called the twist and anti-twist in [20], respectively.¹³ The off-shell action itself takes the compact form

$$S_2|_{\text{POS}} = \frac{4\pi}{b_0} \left[\mathcal{F}(x_+^I) - \mathcal{F}(x_-^I) \right], \tag{2.34}$$

and using (2.18) gives our final expression for the off-shell central charge.

To obtain the on-shell central charge, we need to extremize over the undetermined data and we can proceed as follows. First, the x_-^I are determined by the x_+^I and magnetic charges \mathbf{p}^I via the first equation in (2.32):

$$x_-^I = x_+^I + b_0 \mathbf{p}^I. \tag{2.35}$$

The magnetic charges \mathbf{p}^I are constrained to satisfy the twist/anti-twist condition (2.33), while the x_+^I satisfy a single constraint in (2.32) (which is a constraint on the superpotential). This leaves n degrees of freedom in the x_+^I , and b_0 and one then varies the action in (2.34) over these degrees of freedom to obtain the final on-shell result. We will carry out this extremization explicitly for the STU model further below.

2.4 The $b_0 \rightarrow 0$ limit

Notice that substituting (2.35) into (2.34) allows us to compute the limit

$$\lim_{b_0 \rightarrow 0} S_2|_{\text{POS}} = -4\pi \mathbf{p}^I \frac{\partial}{\partial x^I} \mathcal{F}(x^I) = -2\pi C_{IJK} \mathbf{p}^I x^J x^K. \tag{2.36}$$

¹³More precisely, from equation (3.21) of [20] we see that P_+ , P_- are equivalent to the variables η_1, η_2 in [20], which are also ± 1 .

Here in the limit $x_-^I = x_+^I = x^I$ is constant on M_2 . This is of course consistent with our comments in footnote 10. In addition, from (2.32) we have that the x^I are subject to the constraint $\xi_I x^I = 2$. Our general procedure implies that for an on-shell solution one should extremize the quadratic function of the x^I in (2.36), subject to the constraint $\xi_I x^I = 2$. In particular setting $n_+ = n_- = 1$ this recovers the result in [44] for S^2 solutions with a topological twist, where the magnetic charges \mathbf{p}^I are constrained via (2.33) with $\sigma = +1$ to satisfy $\xi_I \mathbf{p}^I = 2P_+$.

It is remarkable that we have obtained these non-trivial results for $AdS_3 \times S^2$ solutions with a topological twist which have $b_0 = 0$, not by taking $b_0 = 0$ at the beginning but instead assuming $b_0 \neq 0$ and then taking the limit $b_0 \rightarrow 0$.

2.5 STU model

The STU model is a consistent truncation of type IIB supergravity on S^5 , with the $U(1)^3 = SO(2)^3 \subset SO(6)$ isometry gauged in $D = 5$. In the conventions used in [20], this may be obtained from the general $D = 5$, $\mathcal{N} = 2$ gauged supergravity theory by setting $n = 2$, $\xi_I = (1, 1, 1)$ and where the only non-zero component of C_{IJK} (up to permutations) is $C_{123} = 1$. This model then has an AdS_5 vacuum which has unit radius, i.e. $R = 1$ in (2.10), which is dual to $\mathcal{N} = 4$ SYM theory with a central charge given by $a = N^2/4$. The model is explicitly written down in section 3 (where one should set $\rho = 0$ there). As discussed in section 2.6, the quantization condition on the magnetic fluxes that enables one to uplift the solutions to $D = 10$ on S^5 is such that if we write $\mathbf{p}^I = p_I/(n_+ n_-)$ we have $p_I \in \mathbb{Z}$.

Our result for the off-shell central charge function (2.18) in this set-up reads

$$c = -\frac{3N^2}{b_0} \left(x_+^1 x_+^2 x_+^3 - x_-^1 x_-^2 x_-^3 \right), \quad (2.37)$$

with

$$\mathbf{p}^I = -\frac{1}{b_0} (x_+^I - x_-^I), \quad \sum_{I=1}^3 x_+^I = 2 - \frac{b_0}{n_+} P_+, \quad \sum_{I=1}^3 x_-^I = 2 + \frac{b_0}{n_-} P_-, \quad (2.38)$$

and the sum of the magnetic charges constrained to obey $\sum_{I=1}^3 \mathbf{p}^I = \frac{n_- P_+ + n_+ P_-}{n_- n_+}$, or equivalently, $p_1 + p_2 + p_3 = n_- P_+ + n_+ P_-$.

It is straightforward to extremize over the independent variables b_0, x_+^1, x_+^2 , say, and we find the following on-shell results. For c and b_0 we now get

$$c = \frac{6p_1 p_2 p_3}{n_+ n_- s} N^2, \quad b_0 = \frac{2n_+ n_- (n_- P_+ - n_+ P_-)}{s}, \quad (2.39)$$

where

$$s \equiv n_+^2 + n_-^2 - (p_1^2 + p_2^2 + p_3^2). \quad (2.40)$$

For the variables x_\pm^I evaluated at the poles we find

$$x_\pm^I = \frac{2p_I (n_\pm P_\mp - p_I)}{s}. \quad (2.41)$$

Recall that we took $n_{\pm} \geq 1$ and $|P_{\pm}| = 1$, so from (2.28) we have $P_+ x_+^I < 0$ and $P_- x_-^I < 0$. From (2.17) we also have $c > 0$. We find that for the twist case where $\sigma = P_+/P_- = +1$ there are no solutions with $P_+ = P_- = 1$, but for $P_+ = P_- = -1$ the above inequalities reduce to precisely two of the p_I being positive, which since $p_1 + p_2 + p_3 < 0$ in this case then implies that the third p_I is negative. For the anti-twist case with $\sigma = P_+/P_- = -1$ we instead find that the inequalities reduce simply to $p_I > 0$.

These results are in precise agreement with those found for the explicit supergravity solutions given in [20]. Notice that we are then able to obtain expressions for various other quantities, without having to solve any supergravity equations directly. For example, one can compute the warp factors at the poles of the spindle:

$$e^{3\lambda} P|_{\pm} = \frac{8 \prod_{I=1}^3 p_I (p_I - n_{\mp} P_{\pm})}{s^3}. \tag{2.42}$$

2.6 Uplifting to $AdS_3 \times Y_7$ solutions

The STU solutions of the previous subsection may be uplifted on S^5 to solutions of type IIB supergravity, where the anti-twist class were first constructed and uplifted in [22, 56]. The metric on the S^5 internal space takes the fibred form

$$ds_{S^5}^2 = \sum_{I=1}^3 (X^I)^{-1} \left[d\mu_I^2 + \mu_I^2 (d\psi_I + A^I)^2 \right]. \tag{2.43}$$

Here (μ_I, ψ_I) form a system of polar coordinates on $S^5 \subset \mathbb{R}^2 \oplus \mathbb{R}^2 \oplus \mathbb{R}^2$, where correspondingly $\sum_{I=1}^3 \mu_I^2 = 1$ and the angular coordinates ψ_I each have period 2π . The solutions of type IIB then take the form $AdS_3 \times Y_7$, where Y_7 is the total space of a fibration

$$S^5 \hookrightarrow Y_7 \rightarrow M_2. \tag{2.44}$$

Here $M_2 \cong \mathbb{WCP}_{[n_+, n_-]}^1$ is a spindle, and from (2.43) one can see that S^5 is fibred as an associated bundle for the action of $U(1)^3$ on $S^5 \subset \mathbb{R}^2 \oplus \mathbb{R}^2 \oplus \mathbb{R}^2$, where each $U(1)$ is fibred with connection one-form given by the $D = 5$ supergravity gauge field A^I , $I = 1, 2, 3$. The magnetic charges $\mathbf{p}^I = p_I / (n_+ n_-)$, with $p_I \in \mathbb{Z}$, are then precisely Chern numbers for this fibration, with their quantization condition being equivalent to (2.44) being a well-defined orbifold fibration [20]. The solutions have only the RR five-form flux turned on in type IIB, and may be interpreted as the near-horizon limits of N D3-branes wrapped over the spindle M_2 .

On the other hand, supersymmetric $AdS_3 \times Y_7$ solutions of type IIB supergravity with only RR five-form flux were first studied in [57]. The results of [28] show that the existence of a solution to the corresponding Killing spinor equation on Y_7 is equivalent to imposing certain geometric conditions on Y_7 . The resulting geometric structure was dubbed ‘‘GK geometry’’ in [58]. In particular Y_7 is equipped with an R-symmetry Killing vector field ξ that is constructed as a bilinear in the Killing spinor. In [58] an *a priori* different approach to defining an off-shell central charge function was introduced in this setting. There one first introduces an action for the supergravity fields on Y_7 , where solutions to the equations of motion extremize this action; one then imposes that the fields are such that a solution to the Killing spinor equation exists, which as mentioned is equivalent to imposing a certain

geometric structure (an “off-shell GK geometry”, in the terminology of [58]). The resulting supersymmetric action (i.e. the action with supersymmetry imposed) was shown to be a function of a trial R-symmetry vector ξ , where solutions to the equations of motion extremize this function over the choice of ξ . Moreover, at the critical point this action is proportional to the central charge of the AdS_3 solution.

We refer the reader to [34, 58] for further details, here just quoting the result when the internal space Y_7 takes the fibred form (2.44). From section 5.1 of [34] we can write down the supersymmetric action/central charge function in our conventions¹⁴

$$\mathcal{Z} = -\frac{3N^2}{b_0} \left(R_+^1 R_+^2 R_+^3 - R_-^1 R_-^2 R_-^3 \right). \quad (2.45)$$

Here geometrically the R_\pm^I parametrize the R-symmetry Killing vector on the copies of S^5/\mathbb{Z}_{n_\pm} that lie over the poles of the spindle:

$$\xi|_\pm = \sum_{I=1}^3 R_\pm^I \partial_{\psi_I}. \quad (2.46)$$

These obey the constraints (see section 7.4 of [34])

$$R_+^I - R_-^I = -b_0 \frac{M^I}{N} = -b_0 \mathbf{p}^I, \quad \sum_{I=1}^3 R_+^I = 2 + \frac{b_0}{n_+}, \quad \sum_{I=1}^3 R_-^I = 2 - \frac{b_0}{\sigma n_-}. \quad (2.47)$$

Here in the type IIB solutions the M^I arise as quantized fluxes of the RR five-form through certain five-cycles in Y_7 , but in this particular case these can also be straightforwardly identified with the Chern numbers $M^I/N = \mathbf{p}^I = p_I/(n_+ n_-)$ of the fibration in (2.44). In [34] it is also shown that N is divisible by $n_+ n_-$, so that the $M^I \in \mathbb{Z}$ are correctly Dirac quantized fluxes. The constraints (2.47) precisely agree with (2.38) provided we set $P_+ = -1$ and recall $\sigma = P_+/P_-$, with $\sigma = \pm 1$ being the twist and anti-twist respectively, and identify

$$R_\pm^I = x_\pm^I. \quad (2.48)$$

Moreover, the trial central charge function (2.45) in GK geometry precisely matches our trial central charge function (2.37), defined in $D = 5$ STU gauged supergravity.

Another set of variables was also introduced in [34], so as to match to both a field theory expression and corresponding conjectures for gravitational block formulae in [22, 23]. We write

$$x_\pm^I = \phi^I \mp \frac{b_0}{2} \mathbf{p}^I. \quad (2.49)$$

By virtue of (2.32), (2.33), the ϕ^I variables then satisfy the constraint

$$\sum_{I=1}^3 \phi^I = 2 - \frac{b_0}{2} \frac{n_- P_+ - n_+ P_-}{n_+ n_-}. \quad (2.50)$$

¹⁴Specifically, $b_0^{\text{there}} = -b_0$, $m_\pm^{\text{there}} = n_\pm$. Although the variables R_\pm^I were already introduced in [34], they are related to the variables b_I^\pm used more extensively in that paper via $R_\pm^1 = b_1^\pm - b_2^\pm - b_3^\pm$, $R_\pm^2 = b_2^\pm$, $R_\pm^3 = b_3^\pm$. This simply corresponds to a different choice of basis for the $U(1)^3$ action on S^5 in which the Killing spinor has charge $b_1/2 = 1$ under the first $U(1)$ direction and is uncharged under the other two. We refer to [34] for further details of these variables.

The central charge expression (2.37) then reads

$$c = 3N^2 \left[\mathbf{p}^1 \phi^2 \phi^3 + \mathbf{p}^2 \phi^3 \phi^1 + \mathbf{p}^3 \phi^1 \phi^2 + \frac{b_0^2}{4} \mathbf{p}^1 \mathbf{p}^2 \mathbf{p}^3 \right], \quad (2.51)$$

in precise agreement with (5.11) of [34].

The agreement of the off-shell functions (2.37), (2.45), with the identification of variables (2.48), seems to be something of a miracle, since the former is defined purely in $D = 5$ gauged supergravity, while the latter is defined in terms of the GK geometry of the internal space Y_7 . However, the identification (2.48) itself is straightforward to explain geometrically. The equivariantly closed forms $\Phi^{F^I}/2\pi$ introduced in (2.22) are by construction representatives of the equivariant extensions of the first Chern class of the complex line bundles L^I on which the A^I are connection one-forms. Denoting these equivariant first Chern classes $c_1^\xi(L^I)$, at a fixed point p we then have the standard property of equivariant Chern classes

$$c_1^\xi(L^I) \Big|_p = \frac{\epsilon_p(L^I)}{2\pi}, \quad (2.52)$$

where $\epsilon_p(L^I)$ is the *weight* of the action of the vector field ξ on the fixed complex line over the point p . On the other hand, from (2.28) we then have

$$\epsilon_\pm(L^I) = x_\pm^I. \quad (2.53)$$

The variables x_\pm^I are thus precisely weights of the lifted R-symmetry vector field on the three fixed complex lines $\mathbb{C} \oplus \mathbb{C} \oplus \mathbb{C} = \mathbb{R}^6$ over each pole of the spindle, which is also precisely what the variables R_\pm^I are in GK geometry. That is, upon uplifting the STU model solution to a IIB solution, the R-symmetry vector ξ acting on M_2 is lifted to the R-symmetry vector acting on Y_7 . This at least partially explains why the formula (2.37) for the off-shell central charge matches the expression (2.45) taken from [34], in terms of these weights being identified.

Clearly this argument is not restricted to the STU model: if there is an uplift of a gauged supergravity, then the values of the scalars x^I at the fixed points in M_2 should always be identified with the weights of the R-symmetry vector in the solution of the uplifted theory.

3 $D = 5$ STU model with hypermultiplets

We now consider extending the results of the previous section to include hypermultiplets. We will restrict our considerations to a specific model that extends the STU model, for which $AdS_3 \times M_2$ solutions, with M_2 a spindle, were analysed in [48].

We start by considering a truncation of maximal $SO(6)$ gauged supergravity discussed in [59], extending [60]. One first considers a $\mathbb{Z}_2 \times \mathbb{Z}_2 \subset SO(6)$ invariant sector which gives rise to an $\mathcal{N} = 2$ gauged supergravity theory with two vector multiplets and 4 hypermultiplets. Then one utilizes an additional $\mathbb{Z}_4 \subset SO(6) \times SL(2)$ symmetry, as in [60], to further truncate the hypermultiplets. This gives a theory whose bosonic content consists of a metric, three gauge fields A^I , two real and neutral scalars φ^i that live in the $\mathcal{N} = 2$ vector multiplets, and four complex and charged scalar fields Z_j that are maintained from the hypermultiplets and parametrise the coset $SU(1, 1)/U(1)$. For our purposes, as in [48], we further truncate

to a single complex scalar Z , and also set the $D = 5$ Chern-Simons term to zero, which is sufficient for the solutions that we study.

In effect we have the STU model coupled to one complex scalar field, $Z = \rho e^{i\theta}$, that lives in a hypermultiplet and is charged under a certain linear combination of the gauge fields. The bosonic Lagrangian in a mostly plus signature is given¹⁵ by

$$\mathcal{L} = \frac{1}{16\pi G} \sqrt{-g} \left[R - \mathcal{V} - \frac{1}{2} \sum_{i=1}^2 (\partial\varphi^i)^2 - \frac{1}{4} \sum_{I=1}^3 (X^I)^{-2} (F^I)^2 - \frac{1}{2} (\partial\rho)^2 - \frac{1}{2} \sinh^2 \rho (D\theta)^2 \right]. \quad (3.1)$$

Here A^I are three U(1) gauge fields, $I = 1, 2, 3$, with field strengths $F^I = dA^I$. The X^I satisfy the constraint $X^1 X^2 X^3 = 1$ and are given by

$$X^1 = e^{-\frac{\varphi^1}{\sqrt{6}} - \frac{\varphi^2}{\sqrt{2}}}, \quad X^2 = e^{-\frac{\varphi^1}{\sqrt{6}} + \frac{\varphi^2}{\sqrt{2}}}, \quad X^3 = e^{\frac{2\varphi^1}{\sqrt{6}}}. \quad (3.2)$$

The potential is given by

$$\mathcal{V} = 2 \left[\sum_{i=1}^2 (\partial_{\varphi^i} W)^2 + (\partial_{\rho} W)^2 \right] - \frac{4}{3} W^2, \quad (3.3)$$

where the superpotential W is given by

$$W = \sum_{I=1}^3 X^I + \sinh^2 \frac{\rho}{2} (\zeta_I X^I), \quad (3.4)$$

with $\zeta_I = (1, 1, -1)$. In addition the complex scalar is charged with respect to the linear combination of gauge fields given by $\zeta_I A^I$, and we have

$$D\theta \equiv d\theta - \zeta_I A^I. \quad (3.5)$$

For a bosonic solution to preserve supersymmetry, we require

$$\begin{aligned} \left[\nabla_{\mu} - \frac{i}{2} Q_{\mu} + \frac{1}{6} W \Gamma_{\mu} + \frac{i}{24} \sum_I (X^I)^{-1} F_{\nu\rho}^I (\Gamma_{\mu}^{\nu\rho} - 4\delta_{\mu}^{\nu} \Gamma^{\rho}) \right] \epsilon &= 0, \\ \left[\Gamma^{\mu} \partial_{\mu} \varphi^i - 2\partial_{\varphi^i} W + \frac{i}{2} \sum_{I=1}^3 \partial_{\varphi^i} (X^I)^{-1} F_{\mu\nu}^{(I)} \Gamma^{\mu\nu} \right] \epsilon &= 0, \\ \left[\Gamma^{\mu} \partial_{\mu} \rho - 2\partial_{\rho} W + 2i\partial_{\rho} Q_{\mu} \Gamma^{\mu} \right] \epsilon &= 0, \end{aligned} \quad (3.6)$$

where

$$Q_{\mu} = \sum_{I=1}^3 A_{\mu}^I - \sinh^2 \frac{\rho}{2} D_{\mu} \theta. \quad (3.7)$$

¹⁵We have obtained this from [48] by taking $\alpha \rightarrow \frac{1}{2\sqrt{6}}\varphi^1$, $\beta \rightarrow -\frac{1}{2\sqrt{2}}\varphi^2$, $\varphi \rightarrow \frac{1}{2}\rho$, $g_{\mu\nu} \rightarrow -g_{\mu\nu}$, $A \rightarrow \frac{c_1}{2}A$, $g \rightarrow 2c_2$, $\gamma^{\mu} \rightarrow ic_3\gamma^{\mu}$ with $c_i = \pm 1$ and $c_1 c_2 = -1$, $c_2 c_3 = +1$. We have also redefined $W \rightarrow -\frac{1}{2}W$, $\mathcal{P} \rightarrow \mathcal{V}/4$.

The model admits an AdS_5 vacuum solution with unit radius and vanishing scalar fields, which is dual to $\mathcal{N} = 4$ SYM theory. The $D = 5$ Newton constant is given by $\frac{1}{G_5} = \frac{2N^2}{\pi}$ so that $a_{\mathcal{N}=4} = \frac{N^2}{4}$. There is another AdS_5 vacuum with

$$e^{\frac{\sqrt{3}}{2}\varphi^1} = 2, \quad \varphi^2 = 0, \quad e^\rho = 3, \tag{3.8}$$

and radius $L_{LS} = 3/2^{5/3}$. After uplifting to type IIB supergravity [61] this solution is dual to the $d = 4$, $\mathcal{N} = 1$ Leigh-Strassler (LS) SCFT [62]; the latter arises as the IR limit of an RG flow from $\mathcal{N} = 4$ SYM theory deformed by a mass deformation and the corresponding holographic solution was found in [63]. The central charge of the LS SCFT, in the large N limit, is given by $a_{LS} = \frac{27}{32}a_{\mathcal{N}=4} = \frac{27}{128}N^2$.

3.1 The AdS_3 ansatz

We again consider supersymmetric solutions of the form (2.11) with the warp factor λ , the scalars and the gauge fields all defined on M_2 and $ds^2(AdS_3)$ having unit radius. We can obtain a $D = 2$ action by substituting this ansatz into the $D = 5$ action, and this then gives rise to the correct $D = 2$ equations of motion. Furthermore, after imposing the $D = 2$ equations of motion for λ , or equivalently the trace of the $D = 5$ Einstein equations, we obtain the partially off-shell action (cf. (2.16))

$$S_2|_{\text{POS}} = \frac{2}{3} \int_{\mathcal{M}^2} \left[e^{5\lambda} \mathcal{V} \text{vol} - \frac{1}{2} \sum_{I=1}^3 e^\lambda (X^I)^{-2} F_{12}^I F^I \right], \tag{3.9}$$

where $F^I = F_{12}^I \text{vol}$. We may then similarly define a trial central charge function

$$c = -\frac{3}{\pi} a_{\mathcal{N}=4} S_2|_{\text{POS}} = -\frac{32}{9\pi} a_{LS} S_2|_{\text{POS}}, \tag{3.10}$$

which on-shell is the central charge of the AdS_3 solution.

3.2 Equivariantly closed forms and localization

The ansatz for the Killing spinor is exactly as discussed in section 2.2 and, correspondingly, so are the spinor bilinears. We analyse these further in the appendix where we show that the one-form dual to the Killing vector again satisfies

$$d\xi^b = -2(2 + PS^{-1}e^\lambda W)P \text{vol}. \tag{3.11}$$

We also use the analysis in appendix A to construct the following equivariantly closed forms. For the gauge fields we again find that

$$\Phi^{F^I} = F^I - X^I e^\lambda P, \tag{3.12}$$

are equivariantly closed. A new feature in the presence of the hypermultiplet scalars is that a specific linear combination of the Φ^{F^I} is equivariantly exact:

$$d_\xi D\theta = -\zeta_I \Phi^{F^I}. \tag{3.13}$$

The polyform that we considered before without hypermultiplets, $\Phi^{\text{vol}} \equiv e^{5\lambda} \mathcal{V} \text{vol} - e^{4\lambda} WS$, is no longer equivariantly closed when $\rho \neq 0$. However, the following polyform associated with the action (3.9),

$$\begin{aligned} \Phi^S &= \frac{2}{3} \left[\Phi^{\text{vol}} - \frac{1}{2} \sum_{I=1}^3 e^\lambda (X^I)^{-2} F_{12}^I \Phi^{F^I} \right] \\ &= \frac{2}{3} \left[e^{5\lambda} \mathcal{V} \text{vol} - \frac{1}{2} \sum_{I=1}^3 e^\lambda (X^I)^{-2} F_{12}^I F^I - e^{4\lambda} WS + \frac{1}{2} e^{2\lambda} P \sum_{I=1}^3 (X^I)^{-1} F_{12}^I \right], \end{aligned} \tag{3.14}$$

is equivariantly closed.

The localization then proceeds as in section 2.3 with a few small, but important, changes. We consider M_2 to be a spindle with Killing vector globally defined as $\xi = b_0 \partial_\varphi$, with $\Delta\varphi = 2\pi$, as before. It is again convenient to define

$$x^I = -e^\lambda X^I P. \tag{3.15}$$

For the fluxes, localization gives

$$\mathfrak{p}^I \equiv \int_{M_2} \frac{F^I}{2\pi} = -\frac{1}{b_0} (x_+^I - x_-^I), \tag{3.16}$$

and for the spindle we have the quantization condition

$$\mathfrak{p}^I = \frac{p_I}{n_- n_+}, \quad p_I \in \mathbb{Z}. \tag{3.17}$$

This quantization condition, as in the STU model, is imposed so that when the solutions are uplifted on S^5 we get a well-defined $AdS_3 \times M_7$ solution of type IIB supergravity with M_7 an S^5 fibred over the spindle M_2 . We emphasise that for solutions in which the charged scalar in the hypermultiplet is non-zero, we break a $U(1) \subset U(1)^3$ symmetry and, correspondingly, we will lose a Killing vector on the S^5 in the uplifted solution, but this does not alter the fact that we still need to impose a quantization condition on the fluxes to ensure we have a good S^5 fibration over M_2 , which is a topological constraint.

We can also integrate the new expression (3.13) over M_2 . Since the left hand side is equivariantly exact, the integral vanishes and we immediately deduce that there is a new constraint on the fluxes

$$\sum_{I=1}^3 \zeta_I \mathfrak{p}^I = 0, \tag{3.18}$$

where ζ_I were defined in (3.4). In addition, we can consider the zero-form component of (3.13). By regularity, if $\rho|_{\pm} \neq 0$, we must have $D\theta|_{\pm} = 0$ in order that the complex scalar is regular at the poles (in the orbifold sense), and hence we deduce the constraints

$$\zeta_I x_+^I = \zeta_I x_-^I = 0. \tag{3.19}$$

Notice, in particular this implies at the poles $\zeta_I X^I|_{\pm} = 0$ and hence from (3.4), we find that the value of the superpotential W at the poles only depends on the scalars in the vector

multiplets as in the STU model:

$$W|_{\pm} = \sum_{I=1}^3 X^I|_{\pm}. \tag{3.20}$$

We can now use this to obtain expressions for the weights ϵ_p at the fixed points after recalling that, by definition, $d\xi^b|_p = 2\epsilon_p \text{vol}$ as well as using (3.11) to find

$$\sum_{I=1}^3 x_+^I = 2 - \frac{b_0}{n_+} P_+, \quad \sum_{I=1}^3 x_-^I = 2 + \frac{b_0}{n_-} P_-, \tag{3.21}$$

which, combined with (3.16), gives

$$\sum_{I=1}^3 \mathbf{p}^I = \frac{n_- P_+ + n_+ P_-}{n_- n_+}, \tag{3.22}$$

all as in the STU model without the hypermultiplet.

The localization of the $D = 2$ action proceeds exactly as in the STU model and we again find

$$S_2|_{\text{POS}} = \frac{4\pi}{b_0} [\mathcal{F}(x_+^I) - \mathcal{F}(x_-^I)]. \tag{3.23}$$

The action depends on seven variables, b_0, x_{\pm}^I . We can eliminate x_-^I using (3.16). Also, for given spindle data n_{\pm} and P_{\pm} , there is only one independent flux due to (3.18) and (3.22), which we can take to be $\mathbf{p}_F \equiv \mathbf{p}^1 - \mathbf{p}^2$. We also write $\mathbf{p}_F \equiv p_F/(n_+ n_-)$. Similarly, there is only one independent x_+^I due to (3.19) and (3.21), which we can take to be x_+^1 .

Thus, for given spindle data n_{\pm} , choice of spinor chiralities at the poles P_{\pm} , and independent flux parameter \mathbf{p}_F , we need to vary the action over two variables, which we can take to be x_+^1 and b_0 . Extremizing then leads to the central charge, as well as the value of the warp factor and scalars in the vector multiplet at the poles. Notice, however, that this procedure does not fix the value of the complex scalar ρ in the hypermultiplet at the poles of the spindle. This exactly mirrors the results found in [48] which were obtained by a direct analysis of the BPS equations.

In order to compare to the results of [48] we now take the anti-twist case $P_+ = 1, P_- = \sigma = -1$, and $n_{\pm} \geq 1$, and extremize over x_+^1, b_0 . The extremal value of b_0 is given by

$$b_0 = \frac{n_+ n_- (n_+ + n_-) [3(n_+ - n_-)^2 + 4p_F^2]}{(n_+ - n_-)^2 (n_+^2 + n_+ n_- + n_-^2) + 4n_+ n_- p_F^2}, \tag{3.24}$$

and we can also explicitly write down x_+^1 and hence all of the x_{\pm}^I . The extremal value for the action allows us to write down the on-shell central charge from (3.10)

$$c = \frac{3(n_- - n_+) [(n_+ - n_-)^2 - 4p_F^2] [3(n_+ - n_-)^2 + 4p_F^2]}{32n_+ n_- [(n_+ - n_-)^2 (n_+^2 + n_+ n_- + n_-^2) + 4n_+ n_- p_F^2]} N^2. \tag{3.25}$$

Notice from (3.15) that with $P_+ = 1, P_- = \sigma = -1$ we have $x_-^I < 0, x_+^I > 0$ and we also have $c > 0$. From the explicit expressions we find that these conditions are satisfied if and only if

$$n_- - n_+ > 2|p_F|, \tag{3.26}$$

and $b_0 > 0$. The flux conditions require that $n_- - n_+$ is even and that p_3, p_F are both even or both odd.

For the case of the twist, with $P_+/P_- = \sigma = +1$ but with potentially $P_+ = P_- = 1$ or $P_+ = P_- = -1$, we can similarly extremize over b_0, x_{\pm}^1 and obtain the extremal values for b_0, x_{\pm}^I and c . Imposing $\{x_+^I < 0, x_-^I < 0, c > 0\}$ or $\{x_+^I > 0, x_-^I > 0, c > 0\}$ leads to no solutions.

All of these results are in precise alignment with those¹⁶ found in [48], that were also obtained without explicitly solving the BPS equations, but instead just analysing their structure. The results here thus provide an elegant rederivation using the calculus of equivariant localization.

4 $D = 5, \mathcal{N} = 2$ ungauged supergravity

Starting from the general analysis in section 2, it is straightforward to obtain results for ungauged $D = 5, \mathcal{N} = 2$ supergravity coupled to arbitrary vector multiplets, by simply turning off the FI gauging parameters $\xi_I = 0$. Although we shall only recover known results, we do so in a novel way that is likely to generalize further.¹⁷

We begin by noting that setting $\xi_I = 0$ means that the superpotential $W \equiv 0$ in (2.8), and correspondingly also the potential $\mathcal{V} \equiv 0$ in (2.9). Making the same $AdS_3 \times M_2$ ansatz (2.11), everything goes through as written and we may localize on $M_2 \cong \mathbb{WCP}_{[n_+, n_-]}^1$. The R-symmetry flux in (2.33) is however now zero, which gives the equation

$$n_- P_+ + n_+ P_- = 0. \tag{4.1}$$

Since $n_{\pm} \geq 1$ and $|P_{\pm}| = 1$ we necessarily have (without loss of generality) $P_+ = 1, P_- = -1$ and $n_+ = n_- = n$. But then $M_2 = S^2/\mathbb{Z}_n$ and we may always lift to the simply-connected covering space S^2 with $n = 1$, which we henceforth do. In particular we immediately deduce that supersymmetric spindle solutions do not exist in this ungauged supergravity theory, which is a new result.

Next the constraints (2.32) immediately force $b_0 = 2$, and thus the R-symmetry Killing vector $\xi = 2\partial_{\varphi}$ is fixed to rotate the $M_2 = S^2$ horizon with weight 2. The partially off-shell action now reads

$$S_2|_{\text{POS}} = 2\pi \left[\mathcal{F}(x_+^I) - \mathcal{F}(x_-^I) \right], \tag{4.2}$$

where we have defined

$$x_+^I = x^I - \mathbf{p}^I, \quad x_-^I = x^I + \mathbf{p}^I, \tag{4.3}$$

and $S_2|_{\text{POS}}$ is to be extremized over the constant variables x^I . Using the form of the prepotential (2.2), the extremal equations read

$$C_{IJK} x^J \mathbf{p}^K = 0, \tag{4.4}$$

¹⁶Note the typo in (3.56) of [48], viz. there should be no factor of $n_N n_S$ on the right hand side.

¹⁷For the specific model with hypermultiplets considered in section 3, we find that switching off the gauging in this sector by taking $\zeta_I = 0$ implies $d_{\xi} D\theta = 0$ from (3.13) and hence one no longer deduces the constraints (3.18) and (3.19).

for each I , where recall that C_{IJK} is totally symmetric. Substituting (4.4) back into (4.2) we may immediately compute the on-shell central charge

$$c = -\frac{3\pi}{2G_5} \frac{1}{b_0} [\mathcal{F}(x_+) - \mathcal{F}(x_-)] = \frac{\pi}{4G_5} C_{IJK} \mathbf{p}^I \mathbf{p}^J \mathbf{p}^K. \quad (4.5)$$

We may read the equation (4.4) as saying that x^J is in the null-space of the symmetric matrix $C_{IJK} \mathbf{p}^K$. If the latter is invertible then certainly the only solution is $x^I = 0$. On the other hand, notice that (2.28) gives the values of the physical scalars X^I at the fixed poles of the S^2 as

$$X^I|_{\pm} = \mp(x^I \mp \mathbf{p}^I) e^{-\lambda}|_{\pm}. \quad (4.6)$$

For solutions where the warp factor and scalars are independent of the S^2 coordinates, then we immediately see that this assumption forces the $x^I = 0$ solution above. Proceeding with this solution, the constraint (2.2) then fixes

$$e^{3\lambda}|_{\pm} = \frac{1}{3!} C_{IJK} \mathbf{p}^I \mathbf{p}^J \mathbf{p}^K, \quad (4.7)$$

and we deduce

$$X^I|_{\pm} = \frac{\mathbf{p}^I}{(\frac{1}{3!} C_{IJK} \mathbf{p}^I \mathbf{p}^J \mathbf{p}^K)^{1/3}}, \quad (4.8)$$

which is a well-known formula that may be obtained from the standard attractor mechanism in $D = 5$ for supersymmetric black strings when we have a round metric on S^2 ; see, for example, [14].

This theory is the low-energy effective theory one obtains by compactifying M-theory on a Calabi-Yau three-fold X_3 . Including the graviphoton, the number of Abelian vector multiplets is $h^{1,1}$, while the number of hypermultiplets (here turned off) is $h^{2,1} + 1$, where $h^{i,j} = h^{i,j}(X_3)$ denote Hodge numbers. Introducing a basis of two-cycles Ω^I , $I = 1, \dots, h^{1,1}$, generating the free part of $H_2(X_3, \mathbb{Z})$, and a dual basis of four-cycles Ω_I with intersection numbers $\Omega^I \cap \Omega_J = \delta^I_J$, then the scalar fields are

$$X^I = \int_{\Omega^I} \mathcal{J}, \quad (4.9)$$

where \mathcal{J} is the Kähler form on the Calabi-Yau three-fold X_3 . The C_{IJK} are the intersection numbers

$$C_{IJK} = \Omega_I \cap \Omega_J \cap \Omega_K. \quad (4.10)$$

The supersymmetric extremal black string solutions we have discussed above arise by wrapping an M5-brane over the four-cycle $\mathbf{p}^I \Omega_I$, and we see that the central charge (4.5) simply counts the triple intersection number of this wrapped M5-brane.

More generally in our approach nothing forces the warp factor and scalars to be constant over the horizon, and depending on C_{IJK} one may wonder if there are more general $AdS_3 \times S^2$ solutions than the homogeneous choice $x^I = 0$ above; in particular, is it possible¹⁸ to have

¹⁸We thank James Lucietti for discussion on this.

solutions with an inhomogeneous metric on the S^2 ? We have assumed that there is at least an axial symmetry on the S^2 , so in the case of minimal $\mathcal{N} = 2$ ungauged supergravity it is known [64] that the only possibility is $AdS_3 \times S^2$ with a round S^2 ; it would be interesting to extend [64] to include vector multiplets. We can also consider the uniqueness results of [65, 66]; since any $AdS_3 \times M_2$ can be interpreted as a near horizon geometry with horizon $S^1 \times M_2$, by writing AdS_3 in Gaussian null coordinates, the results of [65, 66] imply that the metric on S^2 should be the round metric, provided that the R-symmetry Killing vector is null on a hypersurface on AdS_3 . This leaves the possibility of solutions with non-round metrics on S^2 if the R-symmetry Killing vector is strictly timelike, which is not something we have considered in our analysis. It would be interesting if this argument could be extended to conclude that any supersymmetric $AdS_3 \times S^2$ solution must have a round metric on the S^2 .

5 $D = 4, \mathcal{N} = 2$ gauged supergravity

In this section we discuss $\mathcal{N} = 2$ gauged supergravity in four dimensions, coupled to Abelian vector multiplets [67]. Note that we slightly modify the normalization of the gauge fields and gauge coupling compared to [68, 69] (see [70] as well for a review).

The bosonic content of the graviton supermultiplet is the metric and a graviphoton A^0 , while that of the n Abelian vector multiplets consists of gauge fields A^i and complex scalar fields z^i , where $i = 1, \dots, n$. It is convenient to introduce the index $I = 0, 1, \dots, n$ to unify the description of the gauge fields. The scalars parametrize a special Kähler manifold of complex dimension n , with Kähler potential \mathcal{K} and metric $\mathcal{G}_{i\bar{j}} = \partial_i \partial_{\bar{j}} \mathcal{K}$, that is the base of a symplectic bundle with covariantly holomorphic sections

$$e^{\mathcal{K}/2} \begin{pmatrix} X^I \\ \mathcal{F}_I \end{pmatrix}. \tag{5.1}$$

Here the scalars $X^I = X^I(z^i)$. These obey a symplectic constraint that allows us to express the Kähler potential in terms of the sections

$$e^{\mathcal{K}} \left(X^I \bar{\mathcal{F}}_I - \mathcal{F}_I \bar{X}^I \right) = i \quad \Rightarrow \quad \mathcal{K} = -\log \left[i \left(\mathcal{F}_I \bar{X}^I - X^I \bar{\mathcal{F}}_I \right) \right]. \tag{5.2}$$

We shall assume there exists a holomorphic prepotential $\mathcal{F} = \mathcal{F}(X^I)$, which is homogeneous of degree two, such that $\mathcal{F}_I = \frac{\partial \mathcal{F}}{\partial X^I}$.¹⁹ We define

$$\mathcal{N}_{IJ} = \bar{\mathcal{F}}_{IJ} + i \frac{N_{IK} X^K N_{JL} X^L}{N_{NM} X^N X^M}, \tag{5.3}$$

where $\mathcal{F}_{IJ} = \partial_I \partial_J \mathcal{F}$ and $N_{IJ} = 2 \text{Im} \mathcal{F}_{IJ}$.

The bosonic part of the action, which is determined by the prepotential, is given by

$$S_4 = \frac{1}{16\pi G_4} \int_{M_4} \left(R_4 + \frac{1}{4} \text{Im} \mathcal{N}_{IJ} F_{\mu\nu}^I F^{J\mu\nu} + \frac{1}{4i} \text{Re} \mathcal{N}_{IJ} F_{\mu\nu}^I \tilde{F}^{J\mu\nu} - 2\mathcal{G}_{i\bar{j}} \partial^\mu z^i \partial_\mu \bar{z}^{\bar{j}} - \mathcal{V} \right) \text{vol}_4, \tag{5.4}$$

¹⁹In general the existence of the prepotential depends on the choice of symplectic sections, see [71].

where $\tilde{F}^{I\mu\nu} \equiv -\frac{i}{2}\epsilon^{\mu\nu\rho\sigma}F_{\rho\sigma}^I$. Here we have assumed that there is no gauging of the special Kähler isometries, and the potential is given by

$$\mathcal{V} \equiv \xi_I \xi_J e^{\mathcal{K}} \left(\mathcal{G}^{i\bar{j}} \nabla_i X^I \nabla_{\bar{j}} \bar{X}^J - 3X^I \bar{X}^J \right), \quad (5.5)$$

where the Kähler covariant derivatives of X^I are

$$\nabla_i X^I \equiv (\partial_i + \partial_i \mathcal{K}) X^I, \quad (5.6)$$

and we have chosen to work with U(1) Fayet-Ilioupoulos gauging with $\xi_I \in \mathbb{R}$. It is useful to introduce the holomorphic superpotential

$$W \equiv \xi_I X^I, \quad (5.7)$$

in terms of which the scalar potential is written as

$$\mathcal{V} = e^{\mathcal{K}} \left(\mathcal{G}^{i\bar{j}} \nabla_i W \nabla_{\bar{j}} \bar{W} - 3W \bar{W} \right). \quad (5.8)$$

For use in later sections, it is also helpful to define the real superpotential

$$\mathcal{W} = -\sqrt{2} e^{\mathcal{K}/2} |W|, \quad (5.9)$$

and the potential then reads

$$\mathcal{V} = 2\mathcal{G}^{i\bar{j}} \partial_i \mathcal{W} \partial_{\bar{j}} \mathcal{W} - \frac{3}{2} \mathcal{W}^2. \quad (5.10)$$

A solution is supersymmetric if there exists a non-vanishing Dirac spinor ϵ satisfying the following equations²⁰

$$\begin{aligned} 0 = & \nabla_\mu \epsilon + \frac{i}{2} \mathcal{A}_\mu \Gamma_5 \epsilon - \frac{i}{4} \xi_I A_\mu^I \epsilon + \frac{1}{2\sqrt{2}} \Gamma_\mu e^{\mathcal{K}/2} (\text{Im } W + i \text{Re } W \Gamma_5) \epsilon \\ & - \frac{i}{4\sqrt{2}} \text{Im } \mathcal{N}_{IJ} F_{\nu\rho}^J \Gamma^{\nu\rho} \Gamma_\mu e^{\mathcal{K}/2} (\text{Im } X^I + i \text{Re } X^I \Gamma_5) \epsilon, \end{aligned} \quad (5.11)$$

$$\begin{aligned} 0 = & \frac{1}{2\sqrt{2}} \text{Im } \mathcal{N}_{IJ} F_{\mu\nu}^J \Gamma^{\mu\nu} e^{\mathcal{K}/2} \left[i \text{Im}(\mathcal{G}^{i\bar{j}} \nabla_j X^I) - \text{Re}(\mathcal{G}^{i\bar{j}} \nabla_j X^I) \Gamma_5 \right] \epsilon \\ & + \Gamma^\mu \left(\text{Re } \partial_\mu z^i - i \text{Im } \partial_\mu z^i \Gamma_5 \right) \epsilon \\ & + \frac{i}{\sqrt{2}} e^{\mathcal{K}/2} \left[i \text{Im}(\mathcal{G}^{i\bar{j}} \nabla_j W) - \text{Re}(\mathcal{G}^{i\bar{j}} \nabla_j W) \Gamma_5 \right] \epsilon, \end{aligned} \quad (5.12)$$

where Γ_μ generate Cliff(1,3) and $\Gamma_5 \equiv i\Gamma_{0123}$, and the U(1) connection \mathcal{A} is defined by

$$\mathcal{A}_\mu = -\frac{i}{2} \left(\partial_i \mathcal{K} \partial_\mu z^i - \partial_{\bar{i}} \mathcal{K} \partial_\mu \bar{z}^{\bar{i}} \right). \quad (5.13)$$

²⁰The supersymmetry variations of the gravitino and the gauginos are commonly written in terms of two Weyl spinors of definite chirality. For our purposes it is more convenient to combine them in a single Dirac spinor, as done in [70] (and recall we have modified the normalization of the gauge fields and gauge coupling).

5.1 The AdS_2 ansatz

We consider supersymmetric $AdS_2 \times M_2$ solutions with a warped product metric

$$ds_4^2 = e^{2\lambda} \left[ds^2(AdS_2) + ds^2(M_2) \right]. \quad (5.14)$$

The metric on AdS_2 has unit radius, λ and the scalar fields are functions on M_2 , and the gauge fields A^I are (pull-backs of) gauge fields on M_2 . These solutions can arise as the near horizon limit of magnetically charged static extremal supersymmetric black holes.

We can now proceed as in section 2.1. We first insert the ansatz in the $D = 4$ equations of motion, obtaining $D = 2$ equations of motion, which can be obtained from the effective action (which can also be obtained by substituting the ansatz directly in (5.4))

$$S_2 = \int_{M_2} \left[e^{2\lambda} \left(R - 2 + 6(\nabla\lambda)^2 - 2\mathcal{G}_{i\bar{j}} \partial^\mu z^i \partial_\mu \bar{z}^{\bar{j}} \right) - e^{4\lambda} \mathcal{V} + \frac{1}{4} \text{Im} \mathcal{N}_{IJ} F_{\mu\nu}^I F^{J\mu\nu} \right] \text{vol}, \quad (5.15)$$

where R is the Ricci scalar of M_2 , and ∇ and vol are the Levi-Civita connection and the volume form. We are interested in the Maxwell equation for A^I and the equation for λ , which read

$$d \left(\text{Im} \mathcal{N}_{IJ} * F^J \right) = 0 \quad \Rightarrow \quad d \left(\text{Im} \mathcal{N}_{IJ} F_{12}^J \right) = 0, \quad \forall I, \quad (5.16)$$

$$e^{2\lambda} \left(R - 2 + 6(\nabla\lambda)^2 - 2\mathcal{G}_{i\bar{j}} \partial^\mu z^i \partial_\mu \bar{z}^{\bar{j}} \right) = 2e^{4\lambda} \mathcal{V} + \text{total derivative}. \quad (5.17)$$

Imposing the equation for the warp factor leads to the following partially off-shell two-dimensional action

$$S_2|_{\text{POS}} = \int_{M_2} \left(e^{4\lambda} \mathcal{V} \text{vol} + \frac{1}{2} \text{Im} \mathcal{N}_{IJ} F_{12}^I F^J \right). \quad (5.18)$$

The trace of the Einstein equation derived from the action S_2 is

$$e^{4\lambda} \mathcal{V} + \frac{1}{4} \text{Im} \mathcal{N}_{IJ} F_{\mu\nu}^I F^{J\mu\nu} = -2e^{2\lambda}, \quad (5.19)$$

and imposing this gives us an expression for the on-shell action

$$S_2|_{\text{OS}} = -2 \int_{M_2} e^{2\lambda} \text{vol}. \quad (5.20)$$

If (5.14) is the near horizon limit of an extremal black hole, then $S_2|_{\text{OS}}$ is related to its Bekenstein-Hawking entropy

$$S_{\text{BH}} = -\frac{1}{8G_4} S_2|_{\text{OS}}. \quad (5.21)$$

We will compute this quantity for arbitrary M_2 without explicitly solving the equations of motion, but instead defining an off-shell entropy function

$$S_{\text{BH}} = -\frac{1}{8G_4} S_2|_{\text{POS}}, \quad (5.22)$$

which we compute using the BVAB localization theorem. We can then extremize over the remaining parameters in order to get $S_2|_{\text{OS}}$, and thus the entropy of the black hole.

What we just showed suggests a relation with Euclidean quantum gravity. Recall that $S_2|_{\text{POS}}$ can be obtained by imposing the trace of the Einstein equations of motion in S_4 (5.4), so one has the following

$$S_{\text{BH}} = -\frac{2\pi}{\text{Vol}(AdS_2)} S_4|_{\text{trace } E_{g_4}} = \frac{2\pi}{\text{Vol}(\mathbb{H}_2)} S_4^E|_{\text{trace } E_{g_4}} = -S_4^E|_{\text{trace } E_{g_4}}. \quad (5.23)$$

Here, the first equality follows from the definition, the second one is just a Wick rotation, and the third one uses the renormalized volume of Euclidean AdS_2 . Therefore, at least at the formal level, we find a relation between the entropy function of the black hole and the four-dimensional Euclidean on-shell action. Of course, this relation is suggestive of the canonical relation between the entropy and the Euclidean on-shell action for magnetically charged black holes: $S_{\text{BH}} = -I$. However, there are various caveats. First, in order to establish the latter we would have to include the Gibbons-Hawking-York term and the holographic renormalization counterterms, which are not known for an arbitrary gauged supergravity model. Moreover, notice that (5.23) is a relation between the entropy and the action of the *near horizon* geometry $AdS_2 \times M_2$, whereas the thermodynamics would require the on-shell action of the entire black hole spacetime. This concern is in fact addressed by the equivariant localization itself. As we shall see, S_{BH} is computed using the BVAB theorem, and therefore receives contributions only from fixed point sets, with the result depending only on the topology of spacetime. It is then conceivable that the extremal black hole geometry arises from a limit of a family of geometries with topology $\text{disc} \times M_2$, in which case the action of the entire family of solutions would be the same, independent of the limit parameter. Whilst this is only a conjecture for the model currently studied, it can be proved for the case of minimal $\mathcal{N} = 2$ gauged supergravity, both with spindle [72] and with Riemann surface horizon [73].²¹ At the same time, this also resolves the concern over the divergence of the on-shell action due to the AdS_2 infinite throat. We leave solving these issues and establishing the suggestive relation (5.23) to future work.

5.2 Equivariantly closed forms and localization

Following the same procedure as in section 2.2, we impose a spinor ansatz consistent with (5.14), namely

$$\epsilon = \vartheta \otimes e^{\lambda/2} \zeta, \quad (5.24)$$

where ζ is a spinor on M_2 , and now ϑ is a Killing spinor on AdS_2 . This allows us to reduce the spinor equations (5.11) and (5.12) to spinor equations on M_2 , as we show in appendix B. We only have magnetic fields on M_2 , so in particular $F^I \wedge F^J = 0$; it is then consistent to

²¹In the latter case, namely universal black holes with Riemann surface horizon, the relation $S_{\text{BH}} = -I$ has also been established by finding families of solutions both with supersymmetry [74] and which break supersymmetry [75–77]. The existence of supersymmetric deformations of the black hole with closed interior has been discussed within the STU model in [69].

now set the symplectic sections X^I to be purely imaginary²²

$$z^i \in \mathbb{R}, \quad X^I \in i\mathbb{R}. \quad (5.25)$$

As shown in appendix B, we then define the same real bilinears as in (2.20), and again a canonical analysis of the bilinear equations leads to the conclusion that ξ is a Killing vector, and that the following polyforms are equivariantly closed

$$\begin{aligned} \Phi^{F^I} &\equiv F^I + 2\sqrt{2}e^\lambda e^{\mathcal{K}/2} iX^I P, \\ \Phi^{\text{vol}} &\equiv e^{4\lambda} \mathcal{V} \text{vol} + \sqrt{2}e^{3\lambda} e^{\mathcal{K}/2} iWS, \end{aligned} \quad (5.26)$$

where the bilinear S is a constant that we henceforth set to 1. We can then immediately define another equivariantly closed form Φ^S given by

$$\Phi^S = \Phi^{\text{vol}} + \frac{1}{2} \text{Im} \mathcal{N}_{IJ} F_{12}^I \Phi^{F^J}. \quad (5.27)$$

Having constructed the equivariantly closed polyforms, and assuming that ξ acts on M_2 with only isolated fixed points, then we may appeal to the Berline-Vergne-Atiyah-Bott theorem to evaluate their integrals, as done in section 2.3. Indeed, we find the magnetic charges²³

$$\mathfrak{p}^I = \int_{M_2} \frac{\Phi^{F^I}}{4\pi} = \frac{1}{4\pi} \sum_{\text{fixed } p} \frac{1}{d_p} \frac{2\pi}{\epsilon_p} \left(2\sqrt{2}e^\lambda e^{\mathcal{K}/2} iX^I P \right) \Big|_p, \quad (5.28)$$

and action

$$\begin{aligned} S_2|_{\text{POS}} &= \int_{M_2} \Phi^S \\ &= \sum_{\text{fixed } p} \frac{1}{d_p} \frac{4\pi}{\epsilon_p} \frac{1}{\sqrt{2}} \left(e^{3\lambda} e^{\mathcal{K}/2} iWS + e^\lambda \text{Im} \mathcal{N}_{IJ} F_{12}^I e^{\mathcal{K}/2} iX^J P \right) \Big|_p \\ &= - \sum_{\text{fixed } p} \frac{1}{d_p} \frac{4\pi}{\epsilon_p} P e^{2\lambda} \Big|_p. \end{aligned} \quad (5.29)$$

Analogously to the evaluation of (2.26), we used the Maxwell equation (5.16) and the algebraic relation (B.15) obtained from the spinor equations.

Finally, we introduce the scalars x^I that are the degree-zero components of the equivariant Chern classes (5.26), namely

$$x^I \equiv -2\sqrt{2}e^\lambda e^{\mathcal{K}/2} iX^I P. \quad (5.30)$$

In terms of these, the equation (B.12) for $d\xi^b$ reads

$$d\xi^b = 2 \left(1 - \frac{1}{2} \xi_I x^I \right) P \text{vol} \quad \Rightarrow \quad \xi_I x_p^I = 2 - 2P_p \epsilon_p, \quad (5.31)$$

²²Using a U(1) rotation, associated with the gauge field (5.13), it is possible to change the phase of the symplectic section if we correspondingly rotate the Killing spinors, as discussed around (4.16) in [13] (see also [77]). We also highlight that in section 5.3 we will in fact utilize such a rotation to discuss the STU model. More details are in appendix B.

²³In general there is no reason to quantize the magnetic charges. For models that can be uplifted to string/M-theory, such as the STU model discussed in section 5.3, the suitably normalized gauge fields are associated with U(1) bundles and there is a quantization condition.

and the magnetic charges are

$$\mathbf{p}^I = -\frac{1}{2} \sum_{\text{fixed } p} \frac{1}{d_p \epsilon_p} x_p^I. \quad (5.32)$$

To conclude, we look again at the symplectic constraint (5.2) under the reality assumptions (5.25). This is only consistent if $\mathcal{F}(X^I)$ is pure imaginary, and using homogeneity of \mathcal{F} leads to

$$e^{\mathcal{K}} i\mathcal{F}(X^I) = -\frac{1}{4}. \quad (5.33)$$

Using again homogeneity, we find that

$$i\mathcal{F}(x^I)|_p = 2e^{2\lambda}|_p, \quad (5.34)$$

and thus the reduced action (5.29) takes the form

$$S_2|_{\text{POS}} = - \sum_{\text{fixed } p} \frac{2\pi}{d_p \epsilon_p} P_p i\mathcal{F}(x_p^I). \quad (5.35)$$

Notice that (5.32) and (5.35) have the same form as their analogues (2.30). Therefore, for the four-dimensional solutions of type (5.14), again we can compute the action and the magnetic charges knowing only the values of the scalars x_p^I at the fixed points, together with the data of the circle action, namely d_p , ϵ_p and P_p .

Having established these results, we now consider $M_2 \cong \mathbb{W}\mathbb{P}_{[n_+, n_-]}$ to be a spindle. There are two fixed points labelled by \pm with the order of the orbifold group $d_{\pm} = n_{\pm}$, where we take $n_{\pm} \geq 1$, and weights $\epsilon_{\pm} = \mp b_0/n_{\pm}$, where b_0 is the weight of the azimuthal circle action. We also set $P_- = \sigma P_+$. After application of the BV theorem the form of the integrals for \mathbf{p}^I and $S_2|_{\lambda}$ EOM are the same as in section 2.3, so we can now follow the same steps as we did there.

We find that the magnetic charges (5.32) are given by

$$\mathbf{p}^I = \frac{1}{2b_0}(x_+^I - x_-^I), \quad (5.36)$$

and the constraint (5.31), which now reads

$$\xi_I x_+^I = 2 + P_+ \frac{2b_0}{n_+}, \quad \xi_I x_-^I = 2 - \sigma P_+ \frac{2b_0}{n_-}, \quad (5.37)$$

leads us to an expression for the total magnetic flux of the R-symmetry through the spindle of the form

$$\xi_I \mathbf{p}^I = P_+ \frac{n_- + \sigma n_+}{n_- n_+}. \quad (5.38)$$

Similarly, the off-shell entropy function obtained from the reduced action takes the form

$$S_{\text{BH}} = -\frac{1}{8G_4} S_2|_{\text{POS}} = -\frac{\pi}{2G_4} \frac{1}{2b_0} P_+ \left[i\mathcal{F}(x_+^I) - \sigma i\mathcal{F}(x_-^I) \right]. \quad (5.39)$$

As in the discussion of (2.34), these constraints imply that we have a variational problem with n degrees of freedom.

We can also consider the limit that $b_0 \rightarrow 0$ in order to obtain non-trivial results for the case $M_2 = S^2$. Specifically, we can substitute x_-^I using (5.36) and take the $b_0 \rightarrow 0$ limit in the $\sigma = 1$ case to get

$$\lim_{b_0 \rightarrow 0} S_{\text{BH}} = -P_+ \frac{\pi}{2G_4} \mathbf{p}^I \frac{\partial}{\partial x^I} \mathfrak{i}\mathcal{F}(x^I), \tag{5.40}$$

where note that $x_+^I = x_-^I \equiv x^I$ and they are subject to the constraint $\xi_I x^I = 2$. In the case $n_+ = n_- = 1$, the constraint (5.38) on the fluxes reads $\xi_I \mathbf{p}^I = 2P_+$ and is associated with the standard topological twist on S^2 . This expression for the off-shell entropy has also been obtained by expressing the BPS equations of gauged supergravity as attractor equations [11, 12].

5.3 STU model

The four-dimensional gauged STU model is an $\mathcal{N} = 2$ supergravity theory containing $n = 3$ vector multiplets with $\xi_I = 1$ for all values of I , and prepotential

$$\mathfrak{i}\mathcal{F} = 2\sqrt{X^0 X^1 X^2 X^3}. \tag{5.41}$$

It is a consistent truncation of eleven-dimensional supergravity on S^7 , where the Cartan subgroup $U(1)^4 \subset SO(8)$ is gauged [78].

In contrast to the choice made previously in (5.25), for the STU model we now choose

$$z^i \in \mathbb{R}, \quad X^I \in \mathbb{R}, \tag{5.42}$$

in order to facilitate some comparisons with the literature (e.g. [20]). Specifically, as in [79] we choose

$$\begin{aligned} X^0 &= e^{\frac{1}{2}(\varphi^1 + \varphi^2 + \varphi^3)}, & X^1 &= e^{\frac{1}{2}(\varphi^1 - \varphi^2 - \varphi^3)}, \\ X^2 &= e^{\frac{1}{2}(-\varphi^1 + \varphi^2 - \varphi^3)}, & X^3 &= e^{\frac{1}{2}(-\varphi^1 - \varphi^2 + \varphi^3)}. \end{aligned} \tag{5.43}$$

Here we have used a standard parametrization of the X^I in terms of three real scalars φ^i , as opposed to the scalars z^i used in the general case. As already mentioned in footnote 22, and discussed further in appendix B, we can make this alternative choice provided that we carry out a $U(1)$ rotation. This rotation introduces some changes to the Killing spinor equation on M_2 that one needs to analyse, but otherwise the analysis of the previous subsection goes through unchanged. With this new gauge choice, we have $X^0 X^1 X^2 X^3 = 1$, so $\mathfrak{i}\mathcal{F} = 2$ and $\mathcal{K} = -\log 8$, so that $e^{\mathcal{K}} \mathfrak{i}\mathcal{F}(X^I) = 1/4$ (in contrast with (5.33)). The conventions we use for the STU model are precisely the same as those used in the next section (upon setting $\rho = \theta = 0$). We also recall that the AdS_4 vacuum solution of the STU model is dual to the ABJM theory (with $k = 1$) and the four-dimensional Newton constant is normalised as

$$\frac{1}{G_4} = \frac{2\sqrt{2}}{3} N^{3/2}, \tag{5.44}$$

where N is the rank of the gauge group of ABJM.

Following the procedure in the previous sections, specifically (5.21) and (5.39), we find that the off-shell entropy function for the STU model is given by

$$S_{\text{BH}} = -P_+ \frac{\pi\sqrt{2}}{3} N^{3/2} \frac{1}{b_0} \left(\sqrt{x_+^0 x_+^1 x_+^2 x_+^3} - \sigma \sqrt{x_-^0 x_-^1 x_-^2 x_-^3} \right), \quad (5.45)$$

together with the constraints (5.37) and (5.38), which read

$$\sum_{I=0}^3 x_+^I = 2 + P_+ \frac{2b_0}{n_+}, \quad \sum_{I=0}^3 x_-^I = 2 - \sigma P_+ \frac{2b_0}{n_-}, \quad \sum_{I=0}^3 \mathbf{p}^I = P_+ \frac{n_- + \sigma n_+}{n_- n_+}. \quad (5.46)$$

Extremizing this function leads to the entropy of the accelerating black holes, with various different subfamilies (and/or near horizon limits) studied in [19, 25, 26, 33, 72] (see also [80] for earlier work on accelerating black holes in STU gauged supergravity). The discussion of the most general case is quite involved [25], although the near horizon solutions have been analysed in detail in [26]. However, it is straightforward to obtain simple analytic formulae in subfamilies of examples, and here for illustration we look at the simplest case of minimal $\mathcal{N} = 2$ gauged supergravity. This is obtained setting $\varphi^i = 0$ for all i and $A^I \equiv A$ for all I . Upon choosing $P_+ = -1$ (we could also choose $P_+ = +1$), the off-shell entropy function becomes

$$S_{\text{BH}} = \frac{\pi\sqrt{2}}{3} N^{3/2} \frac{1}{4b_0} \left(1 - \sigma - 2b_0 \frac{n_- + n_+}{n_+ n_-} + b_0^2 \frac{n_-^2 - \sigma n_+^2}{n_+^2 n_-^2} \right), \quad (5.47)$$

where we have solved the constraints (5.46), and the magnetic charge satisfies

$$\mathbf{p} = -\frac{n_- + \sigma n_+}{4n_- n_+}. \quad (5.48)$$

Extremizing S_{BH} leads to

$$b_{0\pm} = \pm \frac{n_+ n_- \sqrt{1 - \sigma}}{\sqrt{n_-^2 - \sigma n_+^2}}. \quad (5.49)$$

Therefore, in the twist case we obtain the extremum at $b_{0*} = 0$, a case associated with S^2 , which we examine further below. In the anti-twist case we find a positive entropy for b_{0+} :

$$S_{\text{BH}} = \frac{\pi\sqrt{2}}{3} N^{3/2} \frac{\sqrt{2(n_+^2 + n_-^2)} - (n_+ + n_-)}{2n_+ n_-}. \quad (5.50)$$

This reproduces the entropy of the explicit family of supersymmetric, accelerating, magnetically charged black holes in [19]. Notice that these anti-twist solutions have $n_+ < b_0 < n_-$, so $\mathbf{p} < 0$, with

$$x_{\pm} = \frac{1}{2} \left(1 - n_{\mp} \sqrt{\frac{2}{n_-^2 + n_+^2}} \right), \quad e^{2\lambda}|_{\pm} = x_{\pm}^2, \quad (5.51)$$

and furthermore $x_+ < 0$, $x_- > 0$, as required (see (6.15)).

We can also consider taking the $b_0 \rightarrow 0$ limit, and set $n_{\pm} = 1 = \sigma$ to recover the S^2 case, as done in (5.40). Taking this limit of (5.45), with $P_+ = -1$, we obtain the entropy function

$$\lim_{b_0 \rightarrow 0} S_{\text{BH}} = \frac{\pi\sqrt{2}}{3} N^{3/2} \sqrt{x^0 x^1 x^2 x^3} \sum_{I=0}^3 \frac{\mathbf{p}^I}{x^I}. \tag{5.52}$$

The variables are now constrained to obey $\sum_{I=0}^3 x^I = 2 = -\sum_{I=0}^3 \mathbf{p}^I$. This precisely matches the result of [15]. In particular, when comparing with the topologically twisted index in the dual field theory one identifies²⁴

$$\mathbf{p}^I \leftrightarrow \mathbf{n}^I, \quad x^I \leftrightarrow \Delta^I. \tag{5.53}$$

Here \mathbf{n}^I are the magnetic fluxes of the background R-symmetry gauge fields, and Δ^I are chemical potentials for the ABJM vector multiplets. The first identification is clear, being the same flux. The second identification, instead, relies on the existence of the uplift on S^7 , in which case the values of the x^I at the fixed points should be identified with the weights of the R-symmetry vector, and thus with the R-charges, as explained in section 2.6. Thus, in writing (5.53) we are identifying the chemical potentials Δ^I for the topologically twisted index with R-charges in the gravity dual; this has of course been noticed before, but the way we have arrived at it here seems to provide a new perspective, that might be worth investigating further.

5.4 Uplifting to $AdS_2 \times Y_9$ solutions

The spindle STU solutions just discussed can be uplifted on S^7 to solutions of eleven-dimensional supergravity [19, 25]. The discussion of the uplift and off-shell geometry are quite similar to that in section 2.6, so we shall be brief.

The eleven-dimensional solution has the form $AdS_2 \times Y_9$, where Y_9 is the total space of a S^7 fibration over M_2 . More specifically, in analogy to (2.43), S^7 is the sphere bundle in \mathbb{R}^8 acted upon by $U(1)^4$, where each $U(1)$ acts on one of the transverse two-planes in \mathbb{R}^8 and the fibration introduces a connection one-form A^I that uplifts the $D = 4$ gauged supergravity field. The magnetic charges (5.28) are then quantized as $\mathbf{p}^I = p^I/n_+n_-$, with $p^I \in \mathbb{Z}$, and they correspond to the Chern numbers of the Y_9 fibration [34]. By construction, the flux of the $D = 11$ four-form vanishes on Y_9 , so these solutions describe the near horizon limit of N M2-branes wrapping M_2 .

The resulting geometries are again the class of GK geometries introduced in [27, 28] and studied in [34, 58]. More specifically, the existence of a Killing spinor guarantees the existence of an R-symmetry vector ξ , and it is possible to introduce a functional \mathcal{S} of the supergravity fields on Y_9 with the following properties. Upon imposing the supersymmetry conditions, which do *not* imply all the equations of motions for the fields, it is a function of ξ , and further extremizing the functional over the choice of ξ is implied by imposing the remaining equations of motion, and \mathcal{S} agrees with the area of M_2 , and thus is related to the Bekenstein-Hawking entropy of the black hole with near horizon geometry $AdS_2 \times M_2$.

²⁴This mapping uses the notation of [35]. To match [15], one needs to rescale Δ^I by π and change the sign of \mathbf{n}^I .

It is also possible to introduce a further change of variables, in order to match other expressions in the literature. We can introduce ϕ^I defined by

$$x_{\pm}^I = \phi^I \pm b_0 \mathbf{p}^I, \tag{5.54}$$

which by construction are constrained by

$$\sum_{I=0}^3 \phi^I = 2 - b_0 \frac{n_- - \sigma n_+}{n_- n_+}. \tag{5.55}$$

Upon substitution of these variables in (5.45), the resulting expression matches (5.41) in [34], and thus the conjectures in [22, 23].

As discussed in section 2.6 for uplifts of five-dimensional solutions, there is *a priori* no reason for the correspondence of the gauged supergravity functional S_{BH} and the internal space one \mathcal{S} . Nonetheless, the geometric discussion surrounding (2.52) works in the same way, *mutatis mutandis*, so again we can identify x_p^I , the values of the zero-degree components of the equivariant first Chern classes at a fixed point, with the weights of the circle action. Since in eleven dimensions the circle action on M_2 uplifts to the R-symmetry action on Y_9 , we can identify x_p^I with the R-symmetry charges.

6 $D = 4$ STU model with a hypermultiplet

The model we will consider was given in [36]. It is a subtruncation of the $U(1)^2 \subset SU(3) \subset SO(6) \subset SO(8)$ invariant consistent truncation of $D = 4$ maximal $SO(8)$ gauged supergravity of [35]. The latter is an $\mathcal{N} = 2$ gauged supergravity theory with three vector multiplets and one hypermultiplet and we consistently truncate one of the two complex scalar fields in the hypermultiplet to zero, and further only considering solutions with $F^I \wedge F^J = 0$ we can take the three complex scalars in the vector multiplets to be real [36]. The model extends the $U(1)^4$ STU model of section 5.3 with the addition of an extra complex scalar field $\rho e^{i\theta}$.

We use the results of [36], correcting the typos noticed in [81]. The bosonic part of the Lagrangian, in a mostly plus signature, is given by²⁵

$$\mathcal{L} = \frac{1}{16\pi G_4} \sqrt{-g} \left[R - \frac{1}{2} \sum_{i=1}^3 (\partial\varphi^i)^2 - \frac{1}{4} \sum_{I=0}^3 (X^I)^{-2} F_{\mu\nu}^I F^{I\mu\nu} - \mathcal{V} - \frac{1}{2} (\partial\rho)^2 - \frac{1}{2} \sinh^2 \rho (D\theta)^2 \right], \tag{6.1}$$

where $X^I(\varphi^i)$, $I = 0, 1, 2, 3$ are given in (5.43). In addition, we have

$$D\theta \equiv d\theta + \sum_{I=0}^3 \zeta_I A^I, \tag{6.2}$$

where $\zeta_I = \frac{1}{2}(1, -1, -1, -1)$. The scalar potential \mathcal{V} is given by

$$\mathcal{V} = 2 \left(\frac{\partial\mathcal{W}}{\partial\rho} \right)^2 + 2 \sum_{i=1}^3 \left(\frac{\partial\mathcal{W}}{\partial\varphi^i} \right)^2 - \frac{3}{2} \mathcal{W}^2, \tag{6.3}$$

²⁵Starting from [81], we have taken $\lambda_i \rightarrow \frac{1}{2}\varphi^i$, $\varphi \rightarrow \frac{1}{2}\rho$, $A^I \rightarrow -\frac{1}{\sqrt{2}}A^I$, and $g \rightarrow \frac{1}{\sqrt{2}}$.

where \mathcal{W} is the real superpotential defined by

$$\mathcal{W} = -\frac{1}{2} \sum_{I=0}^3 X^I + \zeta_I X^I \sinh^2 \frac{\rho}{2}. \quad (6.4)$$

A solution preserves some of the supersymmetry of the maximal gauged supergravity theory if we can solve

$$\begin{aligned} \left[\nabla_\mu - \frac{i}{2} Q_\mu - \frac{1}{4} \mathcal{W} \Gamma_\mu + \frac{i}{16} \sum_{I=0}^3 (X^I)^{-1} F_{\nu\rho}^I \Gamma^{\nu\rho} \Gamma_\mu \right] \epsilon &= 0, \\ \left[\Gamma^\mu \partial_\mu \varphi^i + 2\partial_{\varphi^i} \mathcal{W} + \frac{i}{2} \sum_{I=0}^3 \partial_{\varphi^i} (X^I)^{-1} F_{\mu\nu}^I \Gamma^{\mu\nu} \right] \epsilon &= 0, \\ \left[\Gamma^\mu \partial_\mu \rho + 2\partial_\rho \mathcal{W} + 2i\partial_\rho Q_\mu \Gamma^\mu \right] \epsilon &= 0, \end{aligned} \quad (6.5)$$

where ϵ is a complex $D = 4$ Dirac spinor and

$$Q_\mu \equiv \frac{1}{2} \sum_{I=0}^3 A_\mu^I - \frac{1}{2} (\cosh \rho - 1) D_\mu \theta. \quad (6.6)$$

Since this model extends the STU model, it admits the maximally supersymmetric AdS_4 vacuum solution discussed around (5.44), with vanishing matter fields and the AdS_4 having radius squared equal to $R_{\text{ABJM}}^2 = 1$. This solution uplifts to the $AdS_4 \times S^7$ solution dual to ABJM theory. The model also admits a supersymmetric AdS_4 solution [37] with $R_{\text{mABJM}}^2 \equiv \frac{4}{3\sqrt{3}}$, $e^{\frac{1}{2}\varphi^i} = 3^{1/4}$ and $\tanh \frac{\rho}{2} = \frac{1}{\sqrt{3}}$ and vanishing gauge fields. This solution preserves $SU(3) \times U(1)_R$ global symmetry and after uplifting to $D = 11$ on S^7 is dual to the $d = 3$, $\mathcal{N} = 2$ mABJM SCFT that arises as the IR fixed point in the RG flow of mass-deformed ABJM theory [38–41].

The action is normalized with the $D = 4$ Newton constant as given in (5.44). The free energy of the ABJM theory on S^3 is given by $F_{S^3} = \frac{\pi R_{\text{ABJM}}^2}{2G_4} = \frac{\pi\sqrt{2}}{3} N^{3/2}$. Similarly, the free energy of the mABJM theory on S^3 is given by $F_{S^3} = \frac{\pi R_{\text{mABJM}}^2}{2G_4} = \frac{4\sqrt{2}\pi}{9\sqrt{3}} N^{3/2}$.

6.1 The AdS_2 ansatz

We look for $AdS_2 \times M_2$ solutions with metric again of the form (5.14), with the gauge fields A^I and the scalars $\lambda, \varphi^i, \rho, \theta$ all defined just on M_2 . We can obtain the equations of motion for all of the fields by extremizing a $D = 2$ action obtained by reducing the $D = 4$ action, analogous to the discussion in section 5.1. If we further impose the trace of the $D = 4$ Einstein equation or, equivalently, the $D = 2$ equation of motion for λ , we obtain the partially off-shell action

$$S_2|_{\text{POS}} = \int_{M_2} \left[e^{4\lambda} \mathcal{V} \text{vol} - \frac{1}{2} \sum_{I=0}^3 (X^I)^{-2} F_{12}^I F^I \right]. \quad (6.7)$$

As in section 5.1, we further notice that on-shell we have

$$S_2|_{\text{OS}} = -2 \int_{M_2} e^{2\lambda} \text{vol}, \quad (6.8)$$

which is proportional to the Bekenstein-Hawking entropy, which again takes the form (5.21).

The $D = 4$ Dirac spinor can be written as (5.24) exactly as in the STU model. The Killing spinors therefore take the same form and satisfy the same algebraic equations. Some analysis of the differential conditions is given in appendix B.3. In particular, as in the STU model we have

$$d\xi^b = 2 \left(1 + e^\lambda \mathcal{W} P S^{-1} \right) P \text{ vol}. \quad (6.9)$$

We also need the relation

$$\frac{1}{2} (X^I)^{-1} F_{12}^I = -e^{2\lambda} \mathcal{W} P S^{-1} - 2e^\lambda. \quad (6.10)$$

6.2 Equivariantly closed forms and localization

There are several equivariant polyforms, Φ , that can be constructed on M_2 , satisfying $d_\xi \Phi \equiv (d - \xi \lrcorner) \Phi = 0$. Associated with the four gauge fields, as in the STU model we have

$$\Phi^{F^I} = F^I - X^I e^\lambda P. \quad (6.11)$$

A new feature compared with the STU case is that a specific linear combination of the Φ^{F^I} , corresponding to the specific $U(1)$ that the complex scalar transforms under, is equivariantly exact:

$$d_\xi D\theta = \zeta_I \Phi^{F^I}. \quad (6.12)$$

Associated with the action we have that

$$\Phi^S = e^{4\lambda} \mathcal{V} \text{ vol} - \frac{1}{2} \sum_{I=0}^3 (X^{(I)})^{-2} F_{12}^I F^I + e^{3\lambda} \mathcal{W} S + \frac{1}{2} e^\lambda P \sum_{I=0}^3 (X^{(I)})^{-1} F_{12}^I, \quad (6.13)$$

is equivariantly closed. Unlike the STU case, with $\rho \neq 0$ this is no longer the sum of two equivariantly closed forms, due to the presence of charged matter in the Maxwell equation. From (6.8) and (5.44), we can define the off-shell entropy function analogously to (5.22) as

$$S_{\text{BH}} = -\frac{N^{3/2}}{6\sqrt{2}} \int_{M_2} \Phi^S. \quad (6.14)$$

We now assume that the M_2 has the topology of a spindle, with Killing vector ξ . The localization procedure proceeds in a similar way to the $D = 5$ case with hypermultiplets discussed in section 3. The orbifold singularities of M_2 are given by two integers $n_\pm \geq 1$ and we write the Killing vector as $\xi = b_0 \partial_\varphi$, with $\Delta\varphi = 2\pi$. We also write $\epsilon_\pm = \mp b_0 / n_\pm$ and leave P_+ and $P_+/P_- \equiv \sigma$ arbitrary for now.

We set $S = 1$ and define

$$x^I \equiv X^I e^\lambda P. \quad (6.15)$$

Using localization to obtain the fluxes, we get

$$\mathbf{p}^I \equiv \frac{p^I}{n_+ n_-} \equiv \frac{1}{4\pi} \int_{M_2} F^I = \frac{1}{2b_0} (x_+^I - x_-^I), \quad p^I \in \mathbb{Z}. \quad (6.16)$$

Here we demand that the magnetic charges are suitably quantized, so that when the solutions are uplifted on S^7 we get a well-defined $AdS_2 \times M_9$ solution of $D = 11$ supergravity with M_9 an S^7 fibred over the spindle M_2 . For the STU model, the factor of $1/(4\pi)$ in the quantization condition can be obtained²⁶ using the uplifting formulae in [79]. It is also important to emphasize that for solutions in which the charged scalar in the hypermultiplet is non-zero, we break a $U(1) \subset U(1)^4$ symmetry and correspondingly, we will lose a Killing vector on the S^7 in the uplifted solution, but this does not alter the fact that we still need to impose a quantization condition on the fluxes to ensure we have a good S^7 fibration over M_2 , which is a topological constraint.

Next, if we integrate (6.12) over M_2 we deduce that the charges satisfy the constraint

$$\sum_{I=0}^3 \zeta_I p^I = 0. \tag{6.17}$$

From (6.2) we see that the magnetic flux associated with the $U(1) \subset U(1)^4$ for which the complex scalar is charged, must vanish. Now if the complex scalar is non-vanishing at the poles, $\rho_{\pm} \neq 0$, regularity implies that $D\theta|_{\pm} = 0$ and then, considering the zero-form part of (6.12), we deduce the following constraints on x_{\pm}^I :

$$\sum_{I=0}^3 \zeta_I x_+^I = \sum_{I=0}^3 \zeta_I x_-^I = 0. \tag{6.18}$$

At the poles we therefore have $\zeta_I X^I|_{\pm} = 0$ and hence, from (6.4), we have

$$\mathcal{W}|_{\pm} = -\frac{1}{2} \sum_{I=0}^3 X^I|_{\pm}. \tag{6.19}$$

Next, recalling that the weights ϵ_{\pm} of the action of the Killing vector at the poles are given by $d\xi^{\flat}|_{\pm} = 2\epsilon_{\pm} \text{vol}_2$, using (6.9) we deduce

$$\sum_{I=0}^3 x_+^I = 2 + P_+ \frac{2b_0}{n_+}, \quad \sum_{I=0}^3 x_-^I = 2 - \sigma P_+ \frac{2b_0}{n_-}. \tag{6.20}$$

Using the definition of the fluxes in (6.16), we then deduce the following constraint on the R-symmetry flux

$$\sum_{I=0}^3 \mathfrak{p}^I = P_+ \frac{n_- + \sigma n_+}{n_- n_+}, \quad \Leftrightarrow \quad \sum_{I=0}^3 p^I = P_+ (n_- + \sigma n_+). \tag{6.21}$$

The localization of the action proceeds as in the $D = 4$ STU model of section 5.3 and we find

$$\int_{M_2} \Phi^S = \frac{4\pi}{b_0} P_+ \left(\sqrt{x_+^0 x_+^1 x_+^2 x_+^3} - \sigma \sqrt{x_-^0 x_-^1 x_-^2 x_-^3} \right), \tag{6.22}$$

²⁶We also observe that ϵ is a section of a Spin^c bundle, as can be gleaned from its charge under the A^I in (6.5), and our quantization condition is in agreement with general analysis for supersymmetric spindles in [20].

and hence

$$S_{\text{BH}} = -P_+ \frac{\pi\sqrt{2}}{3} N^{3/2} \frac{1}{b_0} \left(\sqrt{x_+^0 x_+^1 x_+^2 x_+^3} - \sigma \sqrt{x_-^0 x_-^1 x_-^2 x_-^3} \right), \quad (6.23)$$

exactly the same as (5.45). Thus, the expression of the off-shell entropy function does not change upon including a hypermultiplet. Nonetheless, the extremization leads to a different result, since the constraints are different. The x_{\pm}^I satisfy (6.16) and (6.20), which implies the constraint on the R-symmetry flux (6.21), all as in the STU model. In addition we now need to impose (6.18) which imply the vanishing of the broken U(1) flux in (6.17). The action depends on 9 variables b_0, x_{\pm}^I , for given n_{\pm}, σ . We can eliminate x_-^I using (6.16). With the constraints on the R-symmetry flux in (6.21) and the vanishing of the flux associated with the broken U(1) in (6.17), we are left with two independent fluxes which we can take to be, for example, p_1, p_2 .

One can now carry out an explicit extremization looking for solutions satisfying all of the relevant positivity conditions. An example was presented in [36] by studying the BPS equations directly, but it does not satisfy the correct quantization conditions on the fluxes.²⁷

For the anti-twist class, we have found various explicit solutions that satisfy the appropriate positivity constraints, including $S_{\text{BH}} > 0$. We choose $P_+ = -1$, as we did for the STU model, and summarize some examples in table 1, where we give the values of (n_+, n_-) and the magnetic fluxes (p^1, p^2) with (p^0, p^3) determined via

$$p^0 = -\frac{1}{2}(n_- - n_+), \quad p^3 = p^0 - p^1 - p^2. \quad (6.24)$$

With $P_{\pm} = \mp 1$ from (6.15) we have $x_+ < 0$ and $x_- > 0$. From (6.20), with $n_{\pm} \geq 1$ we deduce $n_- > b_0 > n_+$ and in turn, from (6.16) that $p^I < 0$. We highlight that there is a symmetry that permutes p_1, p_2, p_3 and this can be utilized to obtain other cases. For example, for $(n_-, n_+) = (9, 1)$ we can have $p^I = -(4, 1, 1, 2)$ and $p^I = -(4, 1, 2, 1)$ and $p^I = -(4, 2, 1, 1)$. We expect that for all these cases supergravity solutions exist: for the special cases with $p^1 = p^2 = -\frac{1}{6}(n_- - n_+)$, we have explicit solutions of minimal $\mathcal{N} = 2$ gauged supergravity and for remaining cases one could check by numerically solving the BPS equations as in [36]. As an illustration, we can be more explicit for a particular example by considering $(n_-, n_+) = (15, 3)$ and $p^I = -(6, 3, 2, 1)$, finding

$$\begin{aligned} b_0 &= 4.23685, \\ x_-^0 &= 0.717543, & x_-^1 &= 0.331002, & x_-^2 &= 0.249196, & x_-^3 &= 0.137345, \\ x_+^0 &= -0.412284, & x_+^1 &= -0.233912, & x_+^2 &= -0.127413, & x_+^3 &= -0.0509594, \\ e^{2\lambda}|_- &= 0.0901606, & e^{2\lambda}|_+ &= 0.0250233, & S_{\text{BH}} &= 0.0402617N^{3/2}. \end{aligned} \quad (6.25)$$

Interestingly, the integers (n_-, n_+) in this example are not co-prime and moreover, we find that there are no solutions, with properly quantized fluxes, if we consider the \mathbb{Z}_3 quotient of the spindle and take $(n_-, n_+) = (5, 1)$. The cases with $(n_-, n_+) = (8, 2), (10, 2)$ and $(9, 3)$

²⁷In our language, if we take $(n_-, n_+) = (1, 8)$ and fluxes $p^I = (-\frac{7}{2}, -\frac{1}{6} - \frac{1}{\sqrt{3}}, -\frac{13}{6} - \frac{1}{\sqrt{3}}, -\frac{7}{6} + \frac{2}{\sqrt{3}})$, we precisely recover the numerical values for the entropy just below (3.51) in [36], as well as the values for the scalar fields at the poles and k given in (3.47) of [36], with $b_0 = 1/k$.

(n_-, n_+)	Value of $-(p_1, p_2)$
(7,1)	(1,1)*
(9,1)	(1,1)
(11,1)	(1,1); (2,2)
(13,1)	(1,1); (1,2); (2,2)*
(15,1)	(1,1); (1,3); (1,4); (1,5); (2,2)
(8,2)	(1,1)*
(10,2)	(1,1)
(9,3)	(1,1)*

Table 1. Some values for spindle data in the anti-twist class with quantized magnetic fluxes p^I . Cases with $p^1 = p^2 = -\frac{1}{6}(n_- - n_+)$, marked with *, are associated with explicit solutions in minimal $\mathcal{N} = 2$ gauged supergravity. The values of p^0, p^3 are determined from (6.24), and there are additional examples that can be obtained by permuting (p^1, p^2, p^3) .

in the table have the same feature. In previous works on spindles this possibility seems to have been overlooked.

We have made a numerical search for any possible solutions in the twist class but have not found any. It is therefore natural to conjecture that the twist class is obstructed.

We can also consider taking the $b_0 \rightarrow 0$ limit, and set $n_{\pm} = 1 = \sigma$ to recover the S^2 case. This leads to the same entropy function (5.52) as in the STU case, but with the additional hypermultiplet constraints $\zeta_I x^I = \zeta_I p^I = 0$, and precisely matches the result in [35].

7 $D = 4, \mathcal{N} = 2$ ungauged supergravity

Similar to the discussion in section 4 regarding $D = 5$ supergravity, it is straightforward to set the FI gaugings to zero and derive results for $D = 4$ ungauged supergravity coupled to an arbitrary number of vector multiplets.

The key difference with the discussion of the $D = 4$ gauged supergravity theory in section 5 is that the scalar potential \mathcal{V} in (5.5) vanishes. However, we can still study the $AdS_2 \times M_2$ spacetime (5.14) with $M_2 \cong \mathbb{WCP}_{[n_+, n_-]}$, and most of the derivations in section 5.2 go through, with the equivariantly closed form Φ^{vol} (5.26) now being identically zero and correspondingly with a simplified form Φ^S in (5.27).

We now find the R-symmetry flux (5.38) vanishes, thus imposing the equation

$$n_- P_+ + n_+ P_- = 0. \tag{7.1}$$

With $n_{\pm} \geq 1$, we have $\sigma = P_+ P_- = -1$ and without loss of generality, this implies we can take $P_+ = -1, P_- = 1$ and $n_+ = n_- = n$. Thus, we deduce the new result that ungauged supergravity does not admit any $AdS_2 \times M_2$ solutions with M_2 a spindle. Furthermore, as in section 4, by going to the covering space we find that M_2 can always be taken to be S^2 with $n = 1$. The constraints on x^I (5.31) lead to $b_0 = 2$, so that ξ rotates S^2 with weight 2. The off-shell entropy is

$$S_{\text{BH}} = \frac{\pi}{8G_4} \left[i\mathcal{F}(x_+^I) + i\mathcal{F}(x_-^I) \right], \tag{7.2}$$

and we can introduce constants κ^I via

$$x_+^I = 2(\kappa^I + \mathbf{p}^I), \quad x_-^I = 2(\kappa^I - \mathbf{p}^I), \quad (7.3)$$

so that the extremization of S_{BH} has to be done over these κ^I , obtaining the equation

$$\partial_J \mathcal{F}(x_+^I) = -\partial_J \mathcal{F}(x_-^I). \quad (7.4)$$

This equation may be solved by taking $\kappa^I = 0$. Similarly to the discussion in section 4, $\kappa^I = 0$ necessarily holds for solutions which are invariant under the $\text{SO}(3)$ isometry group acting on S^2 .²⁸ With this solution, we then find

$$S_{\text{BH}} = \frac{\pi}{G_4} \mathbf{i}\mathcal{F}(\mathbf{p}^I). \quad (7.5)$$

At the poles, for these $AdS_2 \times S^2$ solutions we find from the definition of x^I that

$$-\sqrt{2}e^{\mathcal{K}/2} \mathbf{i}X^I|_{\pm} = \mp (\kappa^I \pm \mathbf{p}^I) e^{-\lambda}|_{\pm}, \quad (7.6)$$

and for the case of $\kappa^I = 0$, the constraint (5.34) gives

$$e^{\mathcal{K}/2} \mathbf{i}X^I|_{\pm} = -\frac{\mathbf{p}^I}{2\sqrt{\mathbf{i}\mathcal{F}(\pm\mathbf{p}^I)}}. \quad (7.7)$$

We immediately see one of the conclusions of the attractor mechanism (see e.g. [7, 82]): the value of the scalar fields at the horizon only depends on the charges, and not on the values at infinity. Our simple formula for the black hole entropy (7.5) in the purely magnetically charged case we have studied does not appear in [7, 82], but it is straightforward to see that it agrees with those results. From equation (3.2) of [82] we have (restoring the Newton constant in their formula)

$$S_{\text{BH}} = \frac{\pi}{G_4} |Z_{\text{hor}}|^2, \quad (7.8)$$

where Z is the central charge function, which is evaluated at the horizon in (7.8). On the other hand equation (2.13) of [82] gives the general formula (setting the electric charges to zero)

$$|Z|^2 = e^{\mathcal{K}} |\mathbf{p}^I \mathcal{F}_I(X^J)|^2, \quad (7.9)$$

where recall $\mathcal{F}_I \equiv \partial_{X^I} \mathcal{F}$. From (7.7) we may write $X^I = \gamma \mathbf{p}^I$, where the proportionality constant γ in fact drops out of both sides of (7.7), using $e^{\mathcal{K}} \mathbf{i}\mathcal{F}(X^I) = -\frac{1}{4}$ together with the fact that \mathcal{F} is homogeneous degree 2. We then compute $\gamma \mathbf{p}^I \mathcal{F}_I(X^J) = 2\mathcal{F}(X^J)$ using homogeneity and Euler's formula, where we are evaluating at the extremal point $X^J = \gamma \mathbf{p}^J$. Combing these formulae then gives

$$S_{\text{BH}} = \frac{\pi}{G_4} e^{\mathcal{K}} \cdot \left| \frac{2}{\gamma} \mathcal{F}(X^I) \right|^2 = \frac{\pi}{G_4} \frac{\mathbf{i}\mathcal{F}(X^I)}{\gamma^2} = \frac{\pi}{G_4} \mathbf{i}\mathcal{F}(\mathbf{p}^I), \quad (7.10)$$

which agrees with (7.5).

²⁸Also as in section 4 we leave investigation of possible additional solutions for future work.

8 Discussion

Building on [1–3] we have shown that equivariant localization provides a powerful tool to study black holes within gauged and ungauged supergravity. Using the BVAB theorem we have constructed off-shell entropy functions for magnetically charged black holes of $\mathcal{N} = 2$, $D = 4$ supergravity theories that have $AdS_2 \times M_2$ horizons. We have also obtained analogous results for magnetically charged black strings and rings (when they exist) of $\mathcal{N} = 2$, $D = 5$ supergravity theories that have $AdS_3 \times M_2$ horizons. This provides a new perspective and also extends the attractor mechanism in both gauged and ungauged supergravity.

In order to use the localization technology we require M_2 to have a Killing vector and so we considered M_2 to be a spindle, S^2 or T^2 . Naively, one might expect our approach to be useful only when the Killing vector on M_2 is an R-symmetry Killing vector constructed as a spinor bilinear. This is certainly the case when M_2 is a spindle and for this class we obtained many new results including showing that there are no solutions within ungauged supergravity. For the case of T^2 the two Killing vectors do not arise as Killing spinor bilinears and our approach does not immediately lead to any new results. For the case of $AdS_2 \times S^2$ or $AdS_3 \times S^2$ solutions of gauged supergravity, it is also the case that the Killing vectors on S^2 are not arising from Killing spinor bilinears; nevertheless by considering off-shell solutions where we consider this to be the case, we are able to construct off-shell entropy and central charges by taking a suitable limit.

We considered supergravity models with arbitrary number of vector multiplets and illustrated the formalism with some specific examples. We studied the STU model for both $D = 4$ and $D = 5$ gauged supergravity and recovered results consistent with known explicit supergravity solutions. The solutions of the STU models are of particular interest as they can be uplifted on S^7 and S^5 to obtain exact solutions of $D = 11$ and type IIB, respectively. The uplifted solutions are of the form $AdS_2 \times M_9$ and $AdS_3 \times M_7$, respectively, with M_7, M_9 admitting a GK geometry. We showed that our off-shell entropy functions and off-shell central charge are precisely equivalent to the gravitational block formulae derived in [33, 34] using a very different approach in GK geometry.

We also considered the STU model extended with a complex scalar arising from a hypermultiplet in both $D = 4$ and $D = 5$. We showed that the off-shell entropy and central charge functions that one needs to extremize are the same as in the STU model, but the presence of the hypermultiplets imposes additional constraints on the variables, as well as restricting the allowed magnetic fluxes. For the case when M_2 is a spindle we illuminated and clarified the results of [36, 48] which have been obtained by studying the BPS equations directly. It would be interesting to study other models that can also be uplifted to string/M-theory. For example, one should be able to make contact with the solutions discussed in [83] and [84] with S^2 and spindle horizons, respectively. More generally, it would certainly be of interest to extend our analysis to supergravity theories with arbitrary hypermultiplets.

In this paper we have discussed black objects that are magnetically charged, but we fully expect that the technology will generalize to include electric charge and non-trivial rotation. We have also just discussed two-derivative supergravity, and it would be interesting to extend our analysis to include higher derivative corrections and further investigate the ideas presented in [85].

Acknowledgments

We thank Seyed Morteza Hosseini, David Katona, Gabriel Lopes Cardoso, James Lucietti and Minwoo Suh for helpful discussions. This work was supported in part by STFC grants ST/T000791/1 and ST/T000864/1, and EPSRC grant EP/R014604/1. JPG is supported as a Visiting Fellow at the Perimeter Institute. PBG is supported by the SNSF Ambizione grant PZ00P2_208666. AL is supported by the Palmer scholarship in Mathematical Physics of Merton college. JPG and JFS would like to thank the Isaac Newton Institute for Mathematical Sciences, Cambridge, for support and hospitality during the programme “Black holes: bridges between number theory and holographic quantum information” where work on this paper was undertaken.

A Reduction of $D = 5$ Killing spinors

In this appendix we reduce the $D = 5$ Killing spinor equations (2.7) and (3.6) via the $AdS_3 \times M_2$ ansatz (2.11), and use these to deduce various algebraic and differential equations for bilinears in the Killing spinor ζ on M_2 .

A.1 Killing spinor equations

The decomposition of the spinor ϵ and $Cliff(1, 4)$ matrices is

$$\epsilon = \vartheta \otimes e^{\lambda/2} \zeta, \quad \Gamma_{\mathbf{i}} = \beta_{\mathbf{i}} \otimes \gamma_3, \quad \Gamma_{a+2} = \mathbb{1} \otimes \gamma_a, \quad (\text{A.1})$$

where we denote frame indices on AdS_3 by $\mathbf{i} = 0, 1, 2$ and those on M_2 by $a = 1, 2$. Here $\beta_{\mathbf{i}}$ generate $Cliff(1, 2)$ and γ_a generate $Cliff(2)$, with $\gamma_3 = -i\gamma_1\gamma_2$. The Killing spinor ϑ on AdS_3 satisfies

$$\nabla_{\mathbf{i}} \vartheta = \frac{1}{2} \beta_{\mathbf{i}} \vartheta. \quad (\text{A.2})$$

Inserting this ansatz into (2.7) leads to the reduced spinor equations

$$\begin{aligned} \nabla_a \zeta &= \left[\frac{1}{2} \left(1 - e^{-\lambda} G_{IJ} X^I F_{12}^J \right) \gamma_a \gamma_3 + \frac{i}{2} Q_a \right] \zeta, \\ 0 &= \left[\not{\partial} \lambda + \frac{1}{3} e^{\lambda} W + \left(1 - \frac{1}{3} e^{-\lambda} G_{IJ} X^I F_{12}^J \right) \gamma_3 \right] \zeta, \\ 0 &= \left[\mathcal{G}_{ij} \not{\partial} \varphi^j - e^{\lambda} \partial_i W + e^{-\lambda} G_{IJ} \partial_i X^I F_{12}^J \gamma_3 \right] \zeta. \end{aligned} \quad (\text{A.3})$$

Using these one can derive the following useful relations, where \mathbb{A} is any element of $Cliff(2)$:

$$\begin{aligned} \nabla_a (\zeta^\dagger \mathbb{A} \zeta) &= \frac{1}{2} \left(1 - e^{-\lambda} G_{IJ} X^I F_{12}^J \right) \zeta^\dagger [\mathbb{A}, \gamma_a \gamma_3]_- \zeta, \\ 0 &= (\partial_a \lambda) \zeta^\dagger [\mathbb{A}, \gamma^a]_{\pm} \zeta + \frac{1}{3} (1 \pm 1) e^{\lambda} W \zeta^\dagger \mathbb{A} \zeta \\ &\quad + \left(1 - \frac{1}{3} e^{-\lambda} G_{IJ} X^I F_{12}^J \right) \zeta^\dagger [\mathbb{A}, \gamma_3]_{\pm} \zeta, \\ 0 &= \mathcal{G}_{ij} (\partial_a \varphi^j) \zeta^\dagger [\mathbb{A}, \gamma^a]_{\pm} \zeta - (1 \pm 1) e^{\lambda} \partial_i W \zeta^\dagger \mathbb{A} \zeta \\ &\quad + e^{-\lambda} G_{IJ} \partial_i X^I F_{12}^J \zeta^\dagger [\mathbb{A}, \gamma_3]_{\pm} \zeta, \end{aligned} \quad (\text{A.4})$$

where $[\cdot, \cdot]_{\pm}$ denote anti-commutator and commutator, respectively.

A.2 Bilinear equations

Given the bilinear definitions (2.20) we begin by noting the following identities:

$$\xi \lrcorner \text{vol} = -K, \quad F_{12}^I K = -\xi \lrcorner F^I. \quad (\text{A.5})$$

The relations in (A.4) then lead to the following differential equations

$$\begin{aligned} dS &= 0, \quad dK = 0, \quad d\xi^b = -2 \left(2 + e^\lambda W P S^{-1} \right) P \text{vol}, \\ d(e^\lambda P) &= -\frac{2}{3} G_{JK} X^J (\xi \lrcorner F^K), \quad d(e^{4\lambda} S) = \frac{4}{3} e^{5\lambda} W (\xi \lrcorner \text{vol}), \\ dX^I &= -e^{-\lambda} \mathcal{G}^{ij} G_{JK} \partial_i X^I \partial_j X^J (\xi \lrcorner F^K) P^{-1} \\ &= -e^\lambda \mathcal{G}^{ij} \partial_i X^I \partial_j W (\xi \lrcorner \text{vol}) S^{-1}, \end{aligned} \quad (\text{A.6})$$

and algebraic equation

$$e^{-\lambda} G_{IJ} X^I F_{12}^J = 3 + e^\lambda W P S^{-1}. \quad (\text{A.7})$$

Using equation (A.6) we compute

$$\begin{aligned} d(X^I e^\lambda P) &= (dX^I) e^\lambda P + X^I d(e^\lambda P) \\ &= -G_{JK} \left(\mathcal{G}^{ij} \partial_i X^I \partial_j X^J + \frac{2}{3} X^I X^J \right) (\xi \lrcorner F^K) \\ &= -G_{JK} G^{IJ} (\xi \lrcorner F^K) = -\xi \lrcorner F^I, \end{aligned} \quad (\text{A.8})$$

where we have used (2.6). This proves that the multi-form $\Phi^{F^I} = F^I - X^I e^\lambda P$ is equivariantly closed.

Next we similarly compute

$$\begin{aligned} d(e^{4\lambda} W S) &= \xi_I (dX^I) e^{4\lambda} S + W d(e^{4\lambda} S) \\ &= -e^{5\lambda} \left(\mathcal{G}^{ij} \partial_i W \partial_j W - \frac{4}{3} W^2 \right) (\xi \lrcorner \text{vol}) \\ &= -e^{5\lambda} \mathcal{V} (\xi \lrcorner \text{vol}), \end{aligned} \quad (\text{A.9})$$

where we have used (2.9). This proves that the multi-form $\Phi^{\text{vol}} = e^{5\lambda} \mathcal{V} \text{vol} - e^{4\lambda} W S$ is equivariantly closed.

A.3 STU with hypermultiplet

We now turn to the STU model with hypermultiplets considered in section 3. Compared to the general vector model presented in the previous subsection, this amounts to setting $\xi_I = (1, 1, 1)$ and considering an additional complex scalar field $\rho e^{i\theta}$. Every equation from the previous section follows through directly, except for the last relation (A.9). There are also some additional constraints for ρ . Note that with the STU prepotential $\mathcal{F} = X^1 X^2 X^3$, the metrics defined in (2.3) are simply

$$G_{IJ} = \frac{1}{2} \begin{pmatrix} (X^1)^{-2} & 0 & 0 \\ 0 & (X^2)^{-2} & 0 \\ 0 & 0 & (X^3)^{-2} \end{pmatrix}, \quad \mathcal{G}_{ij} = \frac{1}{2} \delta_{ij}. \quad (\text{A.10})$$

Under the ansätze (2.11) and (A.1) the Killing spinor equations (3.6) reduce to

$$\begin{aligned}
 \nabla_a \zeta &= \left[\frac{1}{2} \left(1 - \frac{1}{2} e^{-\lambda} \sum_I (X^I)^{-1} F_{12}^I \right) \gamma_a \gamma_3 + \frac{i}{2} Q_a \right] \zeta, \\
 0 &= \left[\not{\partial} \lambda + \frac{1}{3} e^\lambda W + \left(1 - \frac{1}{6} e^{-\lambda} \sum_I (X^I)^{-1} F_{12}^I \right) \gamma_3 \right] \zeta, \\
 0 &= \left[\not{\partial} \varphi^i - 2e^\lambda \partial_i W - e^{-\lambda} \sum_I \partial_i (X^I)^{-1} F_{12}^I \gamma_3 \right] \zeta \\
 0 &= [\not{\partial} \rho - 2e^\lambda \partial_\rho W + 4i(\partial_\rho Q_a) \gamma^a] \zeta,
 \end{aligned} \tag{A.11}$$

which are the same equations as (A.3), but where the last line is new. Again these give the constraints (A.4) plus the following relation for ρ where $\mathbb{A} \in \text{Cliff}(2)$

$$0 = (\partial_a \rho) \zeta^\dagger [\mathbb{A}, \gamma^a]_\pm \zeta - 2(1 \pm 1) e^\lambda \partial_\rho W \zeta^\dagger \mathbb{A} \zeta + 4i(\partial_\rho Q_a) \zeta^\dagger [\mathbb{A}, \gamma^a]_\mp \zeta. \tag{A.12}$$

From these we deduce the same differential constraints as before, plus a new one on the last line:

$$\begin{aligned}
 dS &= 0, \quad dK = 0, \quad d\xi^b = -2 \left(2 + e^\lambda W P S^{-1} \right) P \text{vol}, \\
 d(e^\lambda P) &= -\frac{1}{3} \sum_I (X^I)^{-1} (\xi \lrcorner F^I), \quad d(e^{4\lambda} S) = \frac{4}{3} e^{5\lambda} W (\xi \lrcorner \text{vol}), \\
 dX^I &= -\sum_{i,J} e^{-\lambda} \partial_i X^I \partial_i (X^J)^{-1} (\xi \lrcorner F^J) P^{-1}, \\
 d\varphi^i &= -2e^\lambda \partial_i W (\xi \lrcorner \text{vol}) S^{-1}, \\
 d\rho &= -2e^\lambda \partial_\rho W (\xi \lrcorner \text{vol}) S^{-1} - 4(*\partial_\rho Q) P S^{-1},
 \end{aligned} \tag{A.13}$$

such that

$$dW = -2e^\lambda \left[\sum_i (\partial_i W)^2 + (\partial_\rho W)^2 \right] (\xi \lrcorner \text{vol}) S^{-1} - 4(\partial_\rho W) (*\partial_\rho Q) P S^{-1}, \tag{A.14}$$

where

$$\partial_\rho W = \frac{1}{2} \zeta_I X^I \sinh \rho, \quad \partial_\rho Q = -\frac{1}{4} D\theta \sinh \rho. \tag{A.15}$$

We also record the Maxwell equation

$$d \left[e^\lambda (X^I)^{-2} F_{12}^I \right] = e^{3\lambda} \zeta_I \sinh^2 \rho (*D\theta). \tag{A.16}$$

As before Φ^{F^I} is equivariantly closed:

$$\begin{aligned}
 d(X^I e^\lambda P) &= (dX^I) e^\lambda P + X^I d(e^\lambda P) \\
 &= -\sum_{i,J} \left[\partial_i X^I \partial_i (X^J)^{-1} + \frac{1}{3} X^I (X^J)^{-1} \right] (\xi \lrcorner F^J) \\
 &= -\sum_J \left(\sum_i \varsigma_{Ii} \varsigma_{Ji} + \frac{1}{3} \right) X^I (X^J)^{-1} (\xi \lrcorner F^J) = -\xi \lrcorner F^I,
 \end{aligned} \tag{A.17}$$

where we have summarized (3.2) as

$$X^I = \exp\left(\sum_i \varsigma_{Ii} \varphi^i\right), \quad \left(\sum_i \varsigma_{Ii} \varsigma_{Ji} + \frac{1}{3}\right) = \delta_{IJ}. \quad (\text{A.18})$$

On the other hand using (A.14) and the definition (3.3) of the scalar potential

$$d(e^{4\lambda} WS) = -e^{5\lambda} \mathcal{V} (\xi \lrcorner \text{vol}) - 4e^{4\lambda} (\partial_\rho W) (*\partial_\rho Q) P, \quad (\text{A.19})$$

we find that $\Phi^{\text{vol}} = e^{5\lambda} \mathcal{V} \text{vol} - e^{4\lambda} WS$ is not equivariantly closed. Instead using (A.15) and (A.16) we obtain that

$$\begin{aligned} d_\xi \Phi^{\text{vol}} &= 4e^{4\lambda} (\partial_\rho W) (*\partial_\rho Q) P \\ &= -\frac{1}{2} \sum_I d \left[e^\lambda (X^I)^{-2} F_{12}^I \right] X^I e^\lambda P. \end{aligned} \quad (\text{A.20})$$

This proves that the multi-form having the action as the top form,

$$\Phi^S = \frac{2}{3} \left[\Phi^{\text{vol}} - \frac{1}{2} \sum_I \left(e^\lambda (X^I)^{-2} F_{12}^I \right) \Phi^{F^I} \right], \quad (\text{A.21})$$

is equivariantly closed.

Finally we show that $\zeta_I \Phi^{F^I}$ is equivariantly exact. Indeed (A.12) gives

$$\xi \lrcorner (\partial_\rho Q) = \frac{1}{2} e^\lambda (\partial_\rho W) P \implies \xi \lrcorner D\theta = -\zeta_I X^I e^\lambda P, \quad (\text{A.22})$$

where we used the explicit expressions (A.15), so that

$$d_\xi D\theta = -\zeta_I F^I + \zeta_I X^I e^\lambda P = -\zeta_I \Phi^{F^I}. \quad (\text{A.23})$$

B Reduction of $D = 4$ Killing spinors

In this appendix we reduce the $D = 4$ Killing spinor equations (5.11) and (5.12) via the ansatz (5.14), (5.24). This is analogous to the procedure described in appendix A, and will lead to differential and algebraic equations for the bilinears in ζ . Because of the similarities, we shall be brief.

One relevant difference with the five-dimensional case is the presence of the Abelian R-symmetry factor mentioned in footnote 22. Concretely, this transformation acts on the fields by

$$\epsilon \rightarrow e^{-i\frac{\alpha}{2}\Gamma_5} \epsilon, \quad e^{\mathcal{K}/2} X^I \rightarrow e^{-i\alpha} e^{\mathcal{K}/2} X^I, \quad \mathcal{A}_\mu \rightarrow \mathcal{A} + \partial_\mu \alpha, \quad (\text{B.1})$$

and under this transformation the equations (5.11) and (5.12) transform covariantly, that is, schematically

$$(\delta\psi_\mu, \delta\lambda) \left(e^{-i\frac{\alpha}{2}\Gamma_5} \epsilon, e^{-i\alpha} e^{\mathcal{K}/2} X^I, \mathcal{A} + \partial_\mu \alpha \right) = e^{-i\frac{\alpha}{2}\Gamma_5} (\delta\psi_\mu, \delta\lambda) \left(\epsilon, e^{\mathcal{K}/2} X^I, \mathcal{A} \right), \quad (\text{B.2})$$

where ψ_μ and λ are the gravitino and gaugino, respectively. We shall see in B.3 that this transformation allows us to change the reality of the sections.

B.1 Killing spinor equations

We decompose the spinor and Clifford algebra $\text{Cliff}(1, 3)$ as

$$\epsilon = \vartheta \otimes e^{\lambda/2} \zeta, \quad \Gamma_i = \beta_i \otimes \gamma_3, \quad \Gamma_{a+1} = \mathbb{1} \otimes \gamma_a, \quad (\text{B.3})$$

where β_i , with $i = 0, 1$, generate $\text{Cliff}(1, 1)$ with $\beta_3 \equiv -\beta_{01}$, and γ_a generate $\text{Cliff}(2)$, with $\gamma_3 \equiv -i\gamma_{12}$. We also assume that

$$\nabla_i \vartheta = -\frac{1}{2} \beta_i \vartheta. \quad (\text{B.4})$$

Inserting this ansatz in (5.11) gives

$$\begin{aligned} 0 = \vartheta \otimes & \left[\nabla_a \zeta + \frac{1}{2} \gamma_a \not{\partial} \lambda \zeta - \frac{i}{4} \xi_I A_a^I \zeta + \frac{1}{2\sqrt{2}} e^\lambda e^{\mathcal{K}/2} \text{Im } W \gamma_a \zeta \right. \\ & \left. + \frac{1}{2\sqrt{2}} e^{-\lambda} \text{Im } \mathcal{N}_{IJ} F_{12}^J e^{\mathcal{K}/2} \text{Im } X^I \gamma_3 \gamma_a \zeta \right] \\ & + \frac{i}{2} \beta_3 \vartheta \otimes \left(\mathcal{A}_a \gamma_3 \zeta + \frac{1}{\sqrt{2}} e^\lambda e^{\mathcal{K}/2} \text{Re } W \gamma_a \gamma_3 \zeta \right. \\ & \left. - \frac{1}{\sqrt{2}} e^{-\lambda} \text{Im } \mathcal{N}_{IJ} F_{12}^J e^{\mathcal{K}/2} \text{Re } X^I \gamma_a \zeta \right), \end{aligned} \quad (\text{B.5})$$

$$\begin{aligned} 0 = \frac{1}{2} \beta_\lambda \vartheta \otimes & \left[-\zeta + \gamma_3 \not{\partial} \zeta + \frac{1}{\sqrt{2}} e^\lambda e^{\mathcal{K}/2} \text{Im } W \gamma_3 \zeta + \frac{1}{\sqrt{2}} e^{-\lambda} \text{Im } \mathcal{N}_{IJ} F_{12}^J e^{\mathcal{K}/2} \text{Im } X^I \zeta \right] \\ & + \frac{i}{2} e^\lambda \beta_\lambda \beta_3 \vartheta \otimes e^{\mathcal{K}/2} \left[\frac{1}{\sqrt{2}} \text{Re } W \zeta + \frac{1}{\sqrt{2}} e^{-2\lambda} \text{Im } \mathcal{N}_{IJ} F_{12}^J \text{Re } X^I \gamma_3 \zeta \right]. \end{aligned} \quad (\text{B.6})$$

The gaugino variation (5.12), instead, gives

$$\begin{aligned} 0 = \vartheta \otimes & \left[-\frac{1}{\sqrt{2}} e^{-2\lambda} \text{Im } \mathcal{N}_{IJ} F_{12}^J e^{\mathcal{K}/2} \text{Im}(\mathcal{G}^{\bar{i}j} \nabla_j X^I) \gamma_3 \zeta + e^{-\lambda} \not{\partial} \text{Re } z^i \zeta - \frac{1}{\sqrt{2}} \text{Im}(\mathcal{G}^{\bar{i}j} \nabla_j W) \zeta \right] \\ & + i \beta_3 \vartheta \otimes \left[-\frac{1}{\sqrt{2}} e^{-2\lambda} \text{Im } \mathcal{N}_{IJ} F_{12}^J e^{\mathcal{K}/2} \text{Re}(\mathcal{G}^{\bar{i}j} \nabla_j X^I) \zeta - e^{-\lambda} \not{\partial} \text{Im } z^i \gamma_3 \zeta \right. \\ & \left. - \frac{1}{\sqrt{2}} \text{Re}(\mathcal{G}^{\bar{i}j} \nabla_j W) \gamma_3 \zeta \right]. \end{aligned} \quad (\text{B.7})$$

Motivated by known examples (e.g. [13]), in order to solve the equations above we impose the following conditions

$$z^i \in \mathbb{R}, \quad X^I \in i\mathbb{R}, \quad (\text{B.8})$$

and so for instance we may write $\text{Im } W = -iW$. These conditions imply that the equations simplify to

$$\begin{aligned} 0 = \nabla_a \zeta - \frac{i}{4} \xi_I A_a^I \zeta + \left(-\frac{1}{2} - \frac{1}{\sqrt{2}} i e^{-\lambda} e^{\mathcal{K}/2} \text{Im } \mathcal{N}_{IJ} F_{12}^J X^I \right) \gamma_3 \gamma_a \zeta, \\ 0 = \not{\partial} \zeta - \frac{i}{\sqrt{2}} e^\lambda e^{\mathcal{K}/2} W \zeta + \left(-1 - \frac{1}{\sqrt{2}} i e^{-\lambda} \text{Im } \mathcal{N}_{IJ} F_{12}^J e^{\mathcal{K}/2} X^I \right) \gamma_3 \zeta, \\ 0 = \frac{1}{\sqrt{2}} i e^{-2\lambda} \text{Im } \mathcal{N}_{IJ} F_{12}^J e^{\mathcal{K}/2} \mathcal{G}^{\bar{i}j} \nabla_j X^I \gamma_3 \zeta + e^{-\lambda} \not{\partial} z^i \zeta + \frac{i}{\sqrt{2}} \mathcal{G}^{\bar{i}j} \nabla_j W \zeta. \end{aligned} \quad (\text{B.9})$$

If \mathbb{A} is any element of Cliff(2), then the following relations hold

$$\begin{aligned}
 \nabla_a(\zeta^\dagger \mathbb{A} \zeta) &= \left(-\frac{1}{2} - \frac{1}{\sqrt{2}} i e^{-\lambda} e^{\mathcal{K}/2} \text{Im} \mathcal{N}_{IJ} F_{12}^J X^I \right) \zeta^\dagger [\gamma_3 \gamma_a, \mathbb{A}]_- \zeta, \\
 0 &= (\partial_a \lambda) \zeta^\dagger [\mathbb{A}, \gamma^a]_{\pm} \zeta - \frac{1 \pm 1}{\sqrt{2}} i e^{\lambda} e^{\mathcal{K}/2} W \zeta^\dagger \mathbb{A} \zeta \\
 &\quad + \left(-1 - \frac{1}{\sqrt{2}} i e^{-\lambda} \text{Im} \mathcal{N}_{IJ} F_{12}^J e^{\mathcal{K}/2} X^I \right) \zeta^\dagger [\mathbb{A}, \gamma_3]_{\pm} \zeta, \\
 0 &= e^{-\lambda} \partial_a z^i \zeta^\dagger [\mathbb{A}, \gamma^a]_{\pm} \zeta + \frac{1 \pm 1}{\sqrt{2}} i e^{\mathcal{K}/2} \mathcal{G}^{ij} \nabla_j W \zeta^\dagger \mathbb{A} \zeta \\
 &\quad + \frac{1}{\sqrt{2}} i e^{-2\lambda} \text{Im} \mathcal{N}_{IJ} F_{12}^J e^{\mathcal{K}/2} \mathcal{G}^{ij} \nabla_j X^I \zeta^\dagger [\mathbb{A}, \gamma_3]_{\pm} \zeta. \tag{B.10}
 \end{aligned}$$

B.2 Bilinear equations

Using the definitions (2.20) for the bilinears constructed using ζ , we find the following differential relations

$$dS = 0, \quad d(e^\lambda P) = \frac{1}{\sqrt{2}} \text{Im} \mathcal{N}_{IJ} e^{\mathcal{K}/2} \text{Im} X^I \xi \lrcorner F^J, \quad dK = 0, \tag{B.11}$$

$$d\xi^b = 2P \left(1 + \sqrt{2} i P S^{-1} e^\lambda e^{\mathcal{K}/2} W \right) \text{vol}, \tag{B.12}$$

$$d(e^\lambda S) = -\frac{i}{\sqrt{2}} e^{2\lambda} e^{\mathcal{K}/2} W \xi \lrcorner \text{vol}, \tag{B.13}$$

$$\begin{aligned}
 d(e^{\mathcal{K}/2} X^I) &= e^{\mathcal{K}/2} \nabla_i X^I dz^i = -\frac{1}{\sqrt{2}} i P^{-1} e^{-\lambda} \text{Im} \mathcal{N}_{JL} e^{\mathcal{K}} \mathcal{G}^{ij} \nabla_i X^I \nabla_j X^L \xi \lrcorner F^J \\
 &= \frac{1}{\sqrt{2}} S^{-1} i e^\lambda e^{\mathcal{K}} \mathcal{G}^{ij} \nabla_i X^I \nabla_j W \xi \lrcorner \text{vol}, \tag{B.14}
 \end{aligned}$$

and the algebraic condition

$$-\frac{1}{\sqrt{2}} e^{-\lambda} \text{Im} \mathcal{N}_{IJ} F_{12}^I e^{\mathcal{K}/2} i X^J = 1 + \frac{i}{\sqrt{2}} P S^{-1} e^\lambda e^{\mathcal{K}/2} W. \tag{B.15}$$

This allows us to prove the following

$$\begin{aligned}
 d(e^{\mathcal{K}/2} i X^I e^\lambda P) &= \frac{1}{\sqrt{2}} \text{Im} \mathcal{N}_{JL} e^{\mathcal{K}} \left(\mathcal{G}^{ij} \nabla_i X^I \nabla_j X^L + X^I X^L \right) \xi \lrcorner F^J \\
 &= \frac{1}{2\sqrt{2}} \xi \lrcorner F^I, \tag{B.16}
 \end{aligned}$$

where we used the identity

$$e^{\mathcal{K}} \left(\mathcal{G}^{i\bar{j}} \nabla_i X^I \nabla_{\bar{j}} \bar{X}^J + \bar{X}^I X^J \right) = -\frac{1}{2} [(\text{Im} \mathcal{N})^{-1}]^{IJ}. \tag{B.17}$$

This relation implies that Φ^{F^I} in (5.26) is equivariantly closed.

Similarly

$$\begin{aligned}
 d\left(\sqrt{2} e^{3\lambda} e^{\mathcal{K}/2} i W S\right) &= -e^{4\lambda} e^{\mathcal{K}} \left(\mathcal{G}^{ij} \nabla_i W \nabla_j W - 3W^2 \right) \xi \lrcorner \text{vol} \\
 &= e^{4\lambda} \mathcal{V} \xi \lrcorner \text{vol}, \tag{B.18}
 \end{aligned}$$

using the definition (5.8). This relation, together with Maxwell's equations, implies that Φ^{vol} in (5.26) is equivariantly closed.

B.3 STU with hypermultiplet

In this appendix we present the bilinear equations and show that the polyform Φ^S in (6.13) is equivariantly closed. This model differs from the previous discussion of the theory with an arbitrary number of vector multiplets because there is an additional supersymmetry equation coming from the variation of the hyperino, and because of a difference choice of conventions (as mentioned below (5.43), there is a U(1) phase rotation that is required to go between the two), so we repeat some steps for the reader's convenience.

First, we clarify the latter point for the STU model (that is, in absence of the hypermultiplet). As remarked earlier, the phase of the sections X^I is immaterial, since the U(1) transformation (B.1) can be used to trade it for a change of phase for the Killing spinor according to (B.2). Concretely, if instead of choosing the ansatz (B.3), we choose the spinor $\epsilon' = e^{-i\frac{\pi}{4}\Gamma_5}\epsilon$, then we find the following equations instead of (B.5), (B.6) and (B.7)

$$\begin{aligned}
 0 = \vartheta \otimes & \left[\nabla_a \zeta + \frac{1}{2} \gamma_a \not{\partial} \lambda \zeta - \frac{i}{4} \xi_I A_a^I \zeta + \frac{1}{2\sqrt{2}} e^\lambda e^{\mathcal{K}/2} \text{Re } W \gamma_a \zeta \right. \\
 & \left. + \frac{1}{2\sqrt{2}} e^{-\lambda} \text{Im } \mathcal{N}_{IJ} F_{12}^J e^{\mathcal{K}/2} \text{Re } X^I \gamma_3 \gamma_a \zeta \right] \\
 & + \frac{i}{2} \beta_3 \vartheta \otimes \left(\mathcal{A}_a \gamma_3 \zeta - \frac{1}{\sqrt{2}} e^\lambda e^{\mathcal{K}/2} \text{Im } W \gamma_a \gamma_3 \zeta \right. \\
 & \left. + \frac{1}{\sqrt{2}} e^{-\lambda} \text{Im } \mathcal{N}_{IJ} F_{12}^J e^{\mathcal{K}/2} \text{Im } X^I \gamma_a \zeta \right), \tag{B.19}
 \end{aligned}$$

$$\begin{aligned}
 0 = \frac{1}{2} \beta_\lambda \vartheta \otimes & \left[-\zeta + \gamma_3 \not{\partial} \zeta + \frac{1}{\sqrt{2}} e^\lambda e^{\mathcal{K}/2} \text{Re } W \gamma_3 \zeta + \frac{1}{\sqrt{2}} e^{-\lambda} \text{Im } \mathcal{N}_{IJ} F_{12}^J e^{\mathcal{K}/2} \text{Re } X^I \zeta \right] \\
 & + \frac{i}{2} e^\lambda \beta_\lambda \beta_3 \vartheta \otimes e^{\mathcal{K}/2} \left[-\frac{1}{\sqrt{2}} \text{Im } W \zeta - \frac{1}{\sqrt{2}} e^{-2\lambda} \text{Im } \mathcal{N}_{IJ} F_{12}^J \text{Im } X^I \gamma_3 \zeta \right], \tag{B.20}
 \end{aligned}$$

$$\begin{aligned}
 0 = \vartheta \otimes & \left[-\frac{1}{\sqrt{2}} e^{-2\lambda} \text{Im } \mathcal{N}_{IJ} F_{12}^J e^{\mathcal{K}/2} \text{Re}(\mathcal{G}^{\bar{i}j} \nabla_j X^I) \gamma_3 \zeta + e^{-\lambda} \not{\partial} \text{Re } z^i \zeta - \frac{1}{\sqrt{2}} \text{Re}(\mathcal{G}^{\bar{i}j} \nabla_j W) \zeta \right] \\
 & + i \beta_3 \vartheta \otimes \left[\frac{1}{\sqrt{2}} e^{-2\lambda} \text{Im } \mathcal{N}_{IJ} F_{12}^J e^{\mathcal{K}/2} \text{Im}(\mathcal{G}^{\bar{i}j} \nabla_j X^I) \zeta - e^{-\lambda} \not{\partial} \text{Im } z^i \gamma_3 \zeta \right. \\
 & \left. + \frac{1}{\sqrt{2}} \text{Im}(\mathcal{G}^{\bar{i}j} \nabla_j W) \gamma_3 \zeta \right]. \tag{B.21}
 \end{aligned}$$

These are simplified by the choice $z^i \in \mathbb{R}$ and $X^I \in \mathbb{R}$ (compare with (B.8)), assuming which we find

$$\begin{aligned}
 0 = \nabla_a \zeta - \frac{i}{4} \xi_I A_a^I \zeta + \left(-\frac{1}{2} + \frac{1}{\sqrt{2}} e^{-\lambda} e^{\mathcal{K}/2} \text{Im } \mathcal{N}_{IJ} F_{12}^J X^I \right) \gamma_3 \gamma_a \zeta, \\
 0 = \not{\partial} \zeta + \frac{1}{\sqrt{2}} e^\lambda e^{\mathcal{K}/2} W \zeta - \left(1 - \frac{1}{\sqrt{2}} e^{-\lambda} \text{Im } \mathcal{N}_{IJ} F_{12}^J e^{\mathcal{K}/2} X^I \right) \gamma_3 \zeta, \\
 0 = -\frac{1}{\sqrt{2}} e^{-2\lambda} \text{Im } \mathcal{N}_{IJ} F_{12}^J e^{\mathcal{K}/2} \mathcal{G}^{\bar{i}j} \nabla_j X^I \gamma_3 \zeta + e^{-\lambda} \not{\partial} \text{Re } z^i \zeta - \frac{1}{\sqrt{2}} \mathcal{G}^{\bar{i}j} \nabla_j W \zeta. \tag{B.22}
 \end{aligned}$$

Moving now to the concrete case of the STU model with a hypermultiplet, the supersym-

metry equations are (6.5) and inserting the ansätze (B.3), (B.4) leads to the equations

$$\begin{aligned}
0 &= \nabla_a \zeta - \frac{i}{2} Q_a \zeta + \frac{1}{2} \left(-1 - \frac{1}{2} e^{-\lambda} \sum_I (X^I)^{-1} F_{12}^I \right) \gamma_3 \gamma_\alpha \zeta, \\
0 &= \not{\partial} \lambda \zeta - \frac{1}{2} \mathcal{W} e^\lambda \zeta + \left(-1 - \frac{1}{4} e^{-\lambda} \sum_I (X^I)^{-1} F_{12}^I \right) \gamma_3 \zeta, \\
0 &= -e^{-2\lambda} \sum_I \partial_{\varphi^i} (X^I)^{-1} F_{12}^I \gamma_3 \zeta + e^{-\lambda} \not{\partial} \varphi^i \zeta + 2 \partial_{\varphi^i} \mathcal{W} \zeta, \\
0 &= \partial_\rho \mathcal{W} \zeta + \frac{1}{2} e^{-\lambda} (\partial_a \rho + 2i \partial_\rho Q_a) \gamma^a \zeta.
\end{aligned} \tag{B.23}$$

The first three are the same as (B.22), and the last one arises from the variation of the hyperino.

From these equations, using the same techniques that lead to (B.11)–(B.14) we find the following relations

$$\begin{aligned}
dS &= 0, \quad d(e^\lambda P) = -\frac{1}{4} \sum_I (X^I)^{-1} \xi \lrcorner F^I, \quad dK = 0, \\
d\xi^b &= 2P \left(1 + PS^{-1} e^\lambda \mathcal{W} \right) \text{vol}, \\
d(e^\lambda S) &= -e^{2\lambda} \frac{\mathcal{W}}{2} \xi \lrcorner \text{vol}, \\
dX^I &= \sum_i \partial_{\varphi^i} X^I d\varphi^i = \frac{1}{2} e^{-\lambda} P^{-1} \sum_i \varsigma_{Ii} X^I \sum_J \partial_{\varphi^i} (X^J)^{-1} \xi \lrcorner F^J \\
&= S^{-1} e^\lambda \sum_i \varsigma_{Ii} X^I \partial_{\varphi^i} \mathcal{W} \xi \lrcorner \text{vol},
\end{aligned} \tag{B.24}$$

and

$$\begin{aligned}
S d\rho &= 2e^\lambda \partial_\rho \mathcal{W} \xi \lrcorner \text{vol} - P \sinh \rho * D\theta, \\
\xi \lrcorner D\theta &= P e^\lambda \zeta_I X^I,
\end{aligned} \tag{B.25}$$

as well as the algebraic condition

$$e^{-\lambda} \sum_I (X^I)^{-1} F_{12}^I = -4 - PS^{-1} e^\lambda \mathcal{W}. \tag{B.26}$$

In these expression we have summarized (5.43) as

$$X^I = \exp \left(\sum_i \frac{1}{2} \varsigma_{Ii} \varphi^i \right), \tag{B.27}$$

and used the definitions of \mathcal{W} and Q_μ in (6.6) and (6.4), respectively. Compared to the discussion in the previous appendix, we highlight that (B.25) are new equations arising from (B.23).

Once again, we find that F^I is the top form of an equivariantly closed polyform. Using (B.24), we have

$$\begin{aligned}
d(X^I e^\lambda P) &= -\frac{1}{4} X^I \sum_J \left(1 + \sum_i \varsigma_{Ii} \varsigma_{Ji} \right) (X^J)^{-1} \xi \lrcorner F^J \\
&= -\xi \lrcorner F^I,
\end{aligned} \tag{B.28}$$

which implies that Φ^{F^I} in (6.11) is equivariantly closed.

We find that

$$\begin{aligned} d(e^{3\lambda}\mathcal{W}S) &= 3e^{2\lambda}\mathcal{W}d(e^\lambda S) + e^{3\lambda}\left(\partial_\rho\mathcal{W}Sd\rho + \sum_i\partial_{\varphi^i}\mathcal{W}Sd\varphi^i\right) \\ &= e^{4\lambda}\mathcal{V}\xi\lrcorner\text{vol} + \frac{1}{2}Pe^{3\lambda}\sinh^2\rho\sum_I\zeta_I X^I * D\theta, \end{aligned} \tag{B.29}$$

having used the definition (6.3) of the scalar potential in terms of the real superpotential. In order to relate this to the integrand of the action in (6.7), we also need Maxwell's equations, which now include a matter term

$$d\left[(X^I)^{-2}F_{12}^I\right] = -\zeta_I e^{2\lambda}\sinh^2\rho * D\theta. \tag{B.30}$$

Combining this with (B.28) and (B.29), we find

$$d\left(e^{3\lambda}\mathcal{W}S + \frac{1}{2}e^\lambda P\sum_I(X^I)^{-1}F_{12}^I\right) = \xi\lrcorner\left(e^{4\lambda}\mathcal{V}\text{vol} - \frac{1}{2}\sum_I(X^I)^{-2}F_{12}^I F^I\right), \tag{B.31}$$

thus proving that Φ^S in (6.13) is equivariantly closed.

Finally, we can show that there is a combination of the Φ^{F^I} that is equivariantly exact. This follows directly from (B.25) and the definition (6.2):

$$(d - \xi\lrcorner)D\theta = \zeta_I\Phi^{F^I}. \tag{B.32}$$

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] P. Benetti Genolini, J.P. Gauntlett and J. Sparks, *Equivariant Localization in Supergravity*, *Phys. Rev. Lett.* **131** (2023) 121602 [[arXiv:2306.03868](https://arxiv.org/abs/2306.03868)] [[INSPIRE](#)].
- [2] P. Benetti Genolini, J.P. Gauntlett and J. Sparks, *Localizing wrapped M5-branes and gravitational blocks*, *Phys. Rev. D* **108** (2023) L101903 [[arXiv:2308.10933](https://arxiv.org/abs/2308.10933)] [[INSPIRE](#)].
- [3] P. Benetti Genolini, J.P. Gauntlett and J. Sparks, *Equivariant localization for AdS/CFT*, *JHEP* **02** (2024) 015 [[arXiv:2308.11701](https://arxiv.org/abs/2308.11701)] [[INSPIRE](#)].
- [4] N. Berline and M. Vergne, *Classes caractéristiques équivariantes. Formules de localisation en cohomologie équivariante*, *C. R. Acad. Sc. Paris* **295** (1982) 539.
- [5] M.F. Atiyah and R. Bott, *The moment map and equivariant cohomology*, *Topology* **23** (1984) 1 [[INSPIRE](#)].
- [6] A. Sen, *Black hole entropy function and the attractor mechanism in higher derivative gravity*, *JHEP* **09** (2005) 038 [[hep-th/0506177](https://arxiv.org/abs/hep-th/0506177)] [[INSPIRE](#)].
- [7] S. Ferrara, R. Kallosh and A. Strominger, *N = 2 extremal black holes*, *Phys. Rev. D* **52** (1995) R5412 [[hep-th/9508072](https://arxiv.org/abs/hep-th/9508072)] [[INSPIRE](#)].
- [8] A. Strominger, *Macroscopic entropy of N = 2 extremal black holes*, *Phys. Lett. B* **383** (1996) 39 [[hep-th/9602111](https://arxiv.org/abs/hep-th/9602111)] [[INSPIRE](#)].

- [9] S. Ferrara and R. Kallosh, *Supersymmetry and attractors*, *Phys. Rev. D* **54** (1996) 1514 [[hep-th/9602136](#)] [[INSPIRE](#)].
- [10] M. David, N. Ezroua and F. Larsen, *The attractor flow for AdS_5 black holes in $\mathcal{N} = 2$ gauged supergravity*, *JHEP* **08** (2023) 090 [[arXiv:2306.05206](#)] [[INSPIRE](#)].
- [11] S.L. Cacciatori and D. Klemm, *Supersymmetric $AdS(4)$ black holes and attractors*, *JHEP* **01** (2010) 085 [[arXiv:0911.4926](#)] [[INSPIRE](#)].
- [12] G. Dall’Agata and A. Gnechchi, *Flow equations and attractors for black holes in $N = 2$ $U(1)$ gauged supergravity*, *JHEP* **03** (2011) 037 [[arXiv:1012.3756](#)] [[INSPIRE](#)].
- [13] K. Hristov and S. Vandoren, *Static supersymmetric black holes in AdS_4 with spherical symmetry*, *JHEP* **04** (2011) 047 [[arXiv:1012.4314](#)] [[INSPIRE](#)].
- [14] F. Larsen, *The Attractor Mechanism in Five Dimensions*, *Lect. Notes Phys.* **755** (2008) 249 [[hep-th/0608191](#)] [[INSPIRE](#)].
- [15] F. Benini, K. Hristov and A. Zaffaroni, *Black hole microstates in AdS_4 from supersymmetric localization*, *JHEP* **05** (2016) 054 [[arXiv:1511.04085](#)] [[INSPIRE](#)].
- [16] F. Benini, K. Hristov and A. Zaffaroni, *Exact microstate counting for dyonic black holes in AdS_4* , *Phys. Lett. B* **771** (2017) 462 [[arXiv:1608.07294](#)] [[INSPIRE](#)].
- [17] N. Bobev, F.F. Gautason and K. Parmentier, *Holographic Uniformization and Black Hole Attractors*, *JHEP* **06** (2020) 095 [[arXiv:2004.05110](#)] [[INSPIRE](#)].
- [18] P. Ferrero et al., *$D3$ -Branes Wrapped on a Spindle*, *Phys. Rev. Lett.* **126** (2021) 111601 [[arXiv:2011.10579](#)] [[INSPIRE](#)].
- [19] P. Ferrero et al., *Accelerating black holes and spinning spindles*, *Phys. Rev. D* **104** (2021) 046007 [[arXiv:2012.08530](#)] [[INSPIRE](#)].
- [20] P. Ferrero, J.P. Gauntlett and J. Sparks, *Supersymmetric spindles*, *JHEP* **01** (2022) 102 [[arXiv:2112.01543](#)] [[INSPIRE](#)].
- [21] S.M. Hosseini, K. Hristov and A. Zaffaroni, *Gluing gravitational blocks for AdS black holes*, *JHEP* **12** (2019) 168 [[arXiv:1909.10550](#)] [[INSPIRE](#)].
- [22] S.M. Hosseini, K. Hristov and A. Zaffaroni, *Rotating multi-charge spindles and their microstates*, *JHEP* **07** (2021) 182 [[arXiv:2104.11249](#)] [[INSPIRE](#)].
- [23] F. Faedo and D. Martelli, *$D4$ -branes wrapped on a spindle*, *JHEP* **02** (2022) 101 [[arXiv:2111.13660](#)] [[INSPIRE](#)].
- [24] S.M. Hosseini, *Gravitational blocks: Symplectic covariance unveiled*, *Phys. Lett. B* **843** (2023) 138044 [[arXiv:2303.07365](#)] [[INSPIRE](#)].
- [25] P. Ferrero, M. Inglese, D. Martelli and J. Sparks, *Multicharge accelerating black holes and spinning spindles*, *Phys. Rev. D* **105** (2022) 126001 [[arXiv:2109.14625](#)] [[INSPIRE](#)].
- [26] C. Couzens, *A tale of $(M)2$ twists*, *JHEP* **03** (2022) 078 [[arXiv:2112.04462](#)] [[INSPIRE](#)].
- [27] N. Kim and J.-D. Park, *Comments on AdS_2 solutions of $D = 11$ supergravity*, *JHEP* **09** (2006) 041 [[hep-th/0607093](#)] [[INSPIRE](#)].
- [28] J.P. Gauntlett and N. Kim, *Geometries with Killing Spinors and Supersymmetric AdS Solutions*, *Commun. Math. Phys.* **284** (2008) 897 [[arXiv:0710.2590](#)] [[INSPIRE](#)].
- [29] D. Martelli and A. Zaffaroni, *Equivariant localization and holography*, *Lett. Math. Phys.* **114** (2024) 15 [[arXiv:2306.03891](#)] [[INSPIRE](#)].

- [30] E. Colombo, F. Faedo, D. Martelli and A. Zaffaroni, *Equivariant volume extremization and holography*, *JHEP* **01** (2024) 095 [[arXiv:2309.04425](#)] [[INSPIRE](#)].
- [31] S.M. Hosseini and A. Zaffaroni, *Proving the equivalence of c-extremization and its gravitational dual for all toric quivers*, *JHEP* **03** (2019) 108 [[arXiv:1901.05977](#)] [[INSPIRE](#)].
- [32] J.P. Gauntlett, D. Martelli and J. Sparks, *Toric geometry and the dual of \mathcal{I} -extremization*, *JHEP* **06** (2019) 140 [[arXiv:1904.04282](#)] [[INSPIRE](#)].
- [33] A. Boido, J.P. Gauntlett, D. Martelli and J. Sparks, *Entropy Functions For Accelerating Black Holes*, *Phys. Rev. Lett.* **130** (2023) 091603 [[arXiv:2210.16069](#)] [[INSPIRE](#)].
- [34] A. Boido, J.P. Gauntlett, D. Martelli and J. Sparks, *Gravitational Blocks, Spindles and GK Geometry*, *Commun. Math. Phys.* **403** (2023) 917 [[arXiv:2211.02662](#)] [[INSPIRE](#)].
- [35] N. Bobev, V.S. Min and K. Pilch, *Mass-deformed ABJM and black holes in AdS_4* , *JHEP* **03** (2018) 050 [[arXiv:1801.03135](#)] [[INSPIRE](#)].
- [36] M. Suh, *Spindle black holes from mass-deformed ABJM*, [arXiv:2211.11782](#) [[INSPIRE](#)].
- [37] N.P. Warner, *Some New Extrema of the Scalar Potential of Gauged $N = 8$ Supergravity*, *Phys. Lett. B* **128** (1983) 169 [[INSPIRE](#)].
- [38] C.-H. Ahn and J. Paeng, *Three-dimensional SCFTs, supersymmetric domain wall and renormalization group flow*, *Nucl. Phys. B* **595** (2001) 119 [[hep-th/0008065](#)] [[INSPIRE](#)].
- [39] R. Corrado, K. Pilch and N.P. Warner, *An $N = 2$ supersymmetric membrane flow*, *Nucl. Phys. B* **629** (2002) 74 [[hep-th/0107220](#)] [[INSPIRE](#)].
- [40] M. Benna, I. Klebanov, T. Klose and M. Smedback, *Superconformal Chern-Simons Theories and AdS_4/CFT_3 Correspondence*, *JHEP* **09** (2008) 072 [[arXiv:0806.1519](#)] [[INSPIRE](#)].
- [41] I. Klebanov, T. Klose and A. Murugan, *AdS_4/CFT_3 Squashed, Stretched and Warped*, *JHEP* **03** (2009) 140 [[arXiv:0809.3773](#)] [[INSPIRE](#)].
- [42] H. Elvang, R. Emparan, D. Mateos and H.S. Reall, *A supersymmetric black ring*, *Phys. Rev. Lett.* **93** (2004) 211302 [[hep-th/0407065](#)] [[INSPIRE](#)].
- [43] J.P. Gauntlett and J.B. Gutowski, *General concentric black rings*, *Phys. Rev. D* **71** (2005) 045002 [[hep-th/0408122](#)] [[INSPIRE](#)].
- [44] S.L. Cacciatori, D. Klemm and W.A. Sabra, *Supersymmetric domain walls and strings in $D = 5$ gauged supergravity coupled to vector multiplets*, *JHEP* **03** (2003) 023 [[hep-th/0302218](#)] [[INSPIRE](#)].
- [45] J.P. Gauntlett, D. Martelli and J. Sparks, *Toric geometry and the dual of c-extremization*, *JHEP* **01** (2019) 204 [[arXiv:1812.05597](#)] [[INSPIRE](#)].
- [46] J.P. Gauntlett, D. Martelli and J. Sparks, *Fibred GK geometry and supersymmetric AdS solutions*, *JHEP* **11** (2019) 176 [[arXiv:1910.08078](#)] [[INSPIRE](#)].
- [47] N. Bobev, K. Pilch and O. Vasilakis, *$(0, 2)$ SCFTs from the Leigh-Strassler fixed point*, *JHEP* **06** (2014) 094 [[arXiv:1403.7131](#)] [[INSPIRE](#)].
- [48] I. Arav, J.P. Gauntlett, M.M. Roberts and C. Rosen, *Leigh-Strassler compactified on a spindle*, *JHEP* **10** (2022) 067 [[arXiv:2207.06427](#)] [[INSPIRE](#)].
- [49] M. Gunaydin, G. Sierra and P.K. Townsend, *Gauging the $d = 5$ Maxwell-Einstein Supergravity Theories: More on Jordan Algebras*, *Nucl. Phys. B* **253** (1985) 573 [[INSPIRE](#)].
- [50] K. Behrndt, M. Cvetič and W.A. Sabra, *Nonextreme black holes of five-dimensional $N = 2$ AdS supergravity*, *Nucl. Phys. B* **553** (1999) 317 [[hep-th/9810227](#)] [[INSPIRE](#)].

- [51] K. Behrndt, A.H. Chamseddine and W.A. Sabra, *BPS black holes in $N = 2$ five-dimensional AdS supergravity*, *Phys. Lett. B* **442** (1998) 97 [[hep-th/9807187](#)] [[INSPIRE](#)].
- [52] A.H. Chamseddine and W.A. Sabra, *Magnetic strings in five-dimensional gauged supergravity theories*, *Phys. Lett. B* **477** (2000) 329 [[hep-th/9911195](#)] [[INSPIRE](#)].
- [53] B. de Wit and A. Van Proeyen, *Broken sigma model isometries in very special geometry*, *Phys. Lett. B* **293** (1992) 94 [[hep-th/9207091](#)] [[INSPIRE](#)].
- [54] F. Benini and N. Bobev, *Exact two-dimensional superconformal R-symmetry and c-extremization*, *Phys. Rev. Lett.* **110** (2013) 061601 [[arXiv:1211.4030](#)] [[INSPIRE](#)].
- [55] A. Almuhairi and J. Polchinski, *Magnetic AdS $\times \mathbb{R}^2$: Supersymmetry and stability*, [arXiv:1108.1213](#) [[INSPIRE](#)].
- [56] A. Boido, J.M.P. Ipiña and J. Sparks, *Twisted D3-brane and M5-brane compactifications from multi-charge spindles*, *JHEP* **07** (2021) 222 [[arXiv:2104.13287](#)] [[INSPIRE](#)].
- [57] N. Kim, *AdS₃ solutions of IIB supergravity from D3-branes*, *JHEP* **01** (2006) 094 [[hep-th/0511029](#)] [[INSPIRE](#)].
- [58] C. Couzens, J.P. Gauntlett, D. Martelli and J. Sparks, *A geometric dual of c-extremization*, *JHEP* **01** (2019) 212 [[arXiv:1810.11026](#)] [[INSPIRE](#)].
- [59] N. Bobev, A. Kundu, K. Pilch and N.P. Warner, *Supersymmetric Charged Clouds in AdS₅*, *JHEP* **03** (2011) 070 [[arXiv:1005.3552](#)] [[INSPIRE](#)].
- [60] A. Khavaev and N.P. Warner, *A Class of $N = 1$ supersymmetric RG flows from five-dimensional $N = 8$ supergravity*, *Phys. Lett. B* **495** (2000) 215 [[hep-th/0009159](#)] [[INSPIRE](#)].
- [61] K. Pilch and N.P. Warner, *A new supersymmetric compactification of chiral IIB supergravity*, *Phys. Lett. B* **487** (2000) 22 [[hep-th/0002192](#)] [[INSPIRE](#)].
- [62] R.G. Leigh and M.J. Strassler, *Exactly marginal operators and duality in four-dimensional $N = 1$ supersymmetric gauge theory*, *Nucl. Phys. B* **447** (1995) 95 [[hep-th/9503121](#)] [[INSPIRE](#)].
- [63] D.Z. Freedman, S.S. Gubser, K. Pilch and N.P. Warner, *Renormalization group flows from holography supersymmetry and a c theorem*, *Adv. Theor. Math. Phys.* **3** (1999) 363 [[hep-th/9904017](#)] [[INSPIRE](#)].
- [64] H.K. Kunduri and J. Lucietti, *Static near-horizon geometries in five dimensions*, *Class. Quant. Grav.* **26** (2009) 245010 [[arXiv:0907.0410](#)] [[INSPIRE](#)].
- [65] H.S. Reall, *Higher dimensional black holes and supersymmetry*, *Phys. Rev. D* **68** (2003) 024024 [*Erratum* *ibid.* **70** (2004) 089902] [[hep-th/0211290](#)] [[INSPIRE](#)].
- [66] J.B. Gutowski, *Uniqueness of five-dimensional supersymmetric black holes*, *JHEP* **08** (2004) 049 [[hep-th/0404079](#)] [[INSPIRE](#)].
- [67] B. de Wit and A. Van Proeyen, *Potentials and Symmetries of General Gauged $N = 2$ Supergravity: Yang-Mills Models*, *Nucl. Phys. B* **245** (1984) 89 [[INSPIRE](#)].
- [68] D.Z. Freedman and A. Van Proeyen, *Supergravity*, Cambridge University Press (2012) [[DOI:10.1017/cbo9781139026833](#)].
- [69] N. Bobev, A.M. Charles and V.S. Min, *Euclidean black saddles and AdS₄ black holes*, *JHEP* **10** (2020) 073 [[arXiv:2006.01148](#)] [[INSPIRE](#)].
- [70] S.L. Cacciatori, D. Klemm, D.S. Mansi and E. Zorzan, *All timelike supersymmetric solutions of $N = 2$, $D = 4$ gauged supergravity coupled to abelian vector multiplets*, *JHEP* **05** (2008) 097 [[arXiv:0804.0009](#)] [[INSPIRE](#)].

- [71] A. Ceresole, R. D'Auria, S. Ferrara and A. Van Proeyen, *Duality transformations in supersymmetric Yang-Mills theories coupled to supergravity*, *Nucl. Phys. B* **444** (1995) 92 [[hep-th/9502072](#)] [[INSPIRE](#)].
- [72] D. Cassani, J.P. Gauntlett, D. Martelli and J. Sparks, *Thermodynamics of accelerating and supersymmetric AdS_4 black holes*, *Phys. Rev. D* **104** (2021) 086005 [[arXiv:2106.05571](#)] [[INSPIRE](#)].
- [73] P. Benetti Genolini, J.M. Pérez Ipiña and J. Sparks, *Localization of the action in AdS/CFT*, *JHEP* **10** (2019) 252 [[arXiv:1906.11249](#)] [[INSPIRE](#)].
- [74] P. Benetti Genolini and C. Toldo, *Magnetic charge and black hole supersymmetric quantum statistical relation*, *Phys. Rev. D* **107** (2023) L121902 [[arXiv:2304.00605](#)] [[INSPIRE](#)].
- [75] F. Azzurli et al., *A universal counting of black hole microstates in AdS_4* , *JHEP* **02** (2018) 054 [[arXiv:1707.04257](#)] [[INSPIRE](#)].
- [76] N. Halmagyi and S. Lal, *On the on-shell: the action of AdS_4 black holes*, *JHEP* **03** (2018) 146 [[arXiv:1710.09580](#)] [[INSPIRE](#)].
- [77] A. Cabo-Bizet et al., *Entropy functional and the holographic attractor mechanism*, *JHEP* **05** (2018) 155 [[arXiv:1712.01849](#)] [[INSPIRE](#)].
- [78] A. Azizi, H. Godazgar, M. Godazgar and C.N. Pope, *Embedding of gauged STU supergravity in eleven dimensions*, *Phys. Rev. D* **94** (2016) 066003 [[arXiv:1606.06954](#)] [[INSPIRE](#)].
- [79] M. Cvetič et al., *Embedding AdS black holes in ten-dimensions and eleven-dimensions*, *Nucl. Phys. B* **558** (1999) 96 [[hep-th/9903214](#)] [[INSPIRE](#)].
- [80] H. Lü and J.F. Vázquez-Poritz, *C-metrics in Gauged STU Supergravity and Beyond*, *JHEP* **12** (2014) 057 [[arXiv:1408.6531](#)] [[INSPIRE](#)].
- [81] I. Arav, J.P. Gauntlett, Y. Jiao, M. Roberts and C. Rosen, *Superconformal Monodromy Defects in ABJM and mABJM Theory*, to appear.
- [82] K. Behrndt et al., *Classical and quantum $N = 2$ supersymmetric black holes*, *Nucl. Phys. B* **488** (1997) 236 [[hep-th/9610105](#)] [[INSPIRE](#)].
- [83] N. Halmagyi, M. Petrini and A. Zaffaroni, *BPS black holes in AdS_4 from M-theory*, *JHEP* **08** (2013) 124 [[arXiv:1305.0730](#)] [[INSPIRE](#)].
- [84] K. Hristov and M. Suh, *Spindle black holes in $AdS_4 \times SE_7$* , *JHEP* **10** (2023) 141 [[arXiv:2307.10378](#)] [[INSPIRE](#)].
- [85] K. Hristov, *$4d \mathcal{N} = 2$ supergravity observables from Nekrasov-like partition functions*, *JHEP* **02** (2022) 079 [[arXiv:2111.06903](#)] [[INSPIRE](#)].