

RECEIVED: March 19, 2011

REVISED: June 13, 2011

ACCEPTED: June 22, 2011

PUBLISHED: July 5, 2011

## Right-handed sneutrino dark matter in supersymmetric $B - L$ model

---

Shaaban Khalil,<sup>a,b</sup> Hiroshi Okada<sup>a</sup> and Takashi Toma<sup>c</sup>

<sup>a</sup>Centre for Theoretical Physics, The British University in Egypt,  
El Sherouk City, Postal No. 11837, P.O. Box 43, Egypt

<sup>b</sup>Department of Mathematics, Faculty of Science, Ain Shams University,  
Cairo 11566, Egypt

<sup>c</sup>Institute for Theoretical Physics, Kanazawa University,  
Kanazawa 920-1192, Japan

E-mail: [skhalil@Bue.edu.eg](mailto:skhalil@Bue.edu.eg), [HOkada@Bue.edu.eg](mailto:HOkada@Bue.edu.eg),  
[t-toma@hep.s.kanazawa-u.ac.jp](mailto:t-toma@hep.s.kanazawa-u.ac.jp)

**ABSTRACT:** We show that the lightest right-handed sneutrino in TeV scale supersymmetric  $B - L$  model with inverse seesaw mechanism is a viable candidate for cold dark matter. We find that it accounts for the observed dark matter relic abundance in a wide range of parameter space. The spin-independent cross section of  $B - L$  right-handed sneutrino is consistent with the recent results CDMS II and XENON experiments and it is detectable in future direct detection experiments. Although the  $B - L$  right-handed sneutrinos annihilate into leptons, the PAMELA results can not be explained in this model unless a huge boost factor is considered. Also the muon flux generated by  $B - L$  right-handed sneutrino in the galactic center is smaller than Super-Kamiokande's upper bound.

**KEYWORDS:** Supersymmetry Phenomenology

**ARXIV EPRINT:** [1102.4249](https://arxiv.org/abs/1102.4249)

---

**Contents**

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Supersymmetric <math>B - L</math> model with inverse seesaw mechanism</b>	<b>3</b>
<b>3</b>	<b><math>B - L</math> right-handed sneutrino</b>	<b>5</b>
<b>4</b>	<b>Relic abundance of <math>B - L</math> right-handed sneutrino</b>	<b>8</b>
<b>5</b>	<b>Direct detection</b>	<b>10</b>
<b>6</b>	<b>Indirect detection</b>	<b>13</b>
	6.1 PAMELA and Fermi-LAT experiments	13
	6.2 Muon flux measurement from Super-Kamiokande	13
<b>7</b>	<b>Conclusions</b>	<b>15</b>

---

**1 Introduction**

The experimental verifications of non-vanishing neutrino masses and the alluring hints of dark matter's (DM's) existence are serious indications for new physics beyond the Standard Model (SM). Supersymmetry (SUSY) is an attractive candidate for new physics at TeV scale that provides an elegant solution for the SM gauge hierarchy problem and stabilize the SM Higgs mass at the electroweak scale. The minimal supersymmetric standard model (MSSM) is the simplest extension of the SM. In order to account for the observed neutrino masses and mixing, SM singlets (right-handed neutrinos) are usually introduced.

In the MSSM with  $R$ -parity conservation and universal soft SUSY breaking terms, the lightest neutralino, which is typically bino dominated, is an attractive candidate for cold DM. However, the current experimental constraints on SUSY particles lead to overproduction of bino relic abundance, in contradiction with the observational limits of the Wilkinson Microwave Anisotropy Probe (WMAP) [1]. In addition, the recent results of Cryogenic DM Search (CDMS II) [2] set an upper limit on the DM-nucleon elastic scattering spin independent cross section of order  $3.8 \times 10^{-44} \text{cm}^2$  for DM mass of 70 GeV, which imposes stringent limits on the lightest neutralino of the MSSM, even if it consists of gaugino-Higgsino mixture. Therefore, one concludes that the DM constraints severely reduce the allowed range in the parameter space of the MSSM. It is also worth mentioning that the observed anomalies in the cosmic rays may favor the type of dark matter that annihilates into leptons not to quarks, unlike the lightest neutralino in MSSM.

TeV scale right-handed neutrinos can be naturally implemented in supersymmetric  $B - L$  extension of the SM (SUSY  $B - L$ ), which is based on the gauge group  $G_{B-L} \equiv$

$SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_{B-L}$  [3]. In this model, three SM singlet fermions arise quite naturally due to the  $U(1)_{B-L}$  anomaly cancellation conditions. These particles are accounted for right-handed neutrinos, and hence a natural explanation for the seesaw mechanism is obtained [3–5]. This work was based on earlier papers in ref. [6–8]. In this class of model, the scale of  $B-L$  symmetry breaking is related to supersymmetry breaking scale [9]. Therefore, the right-handed neutrino masses are naturally of order TeV scale. This has initiated a considerable interest in analyzing the phenomenological implications of these models and their possible signatures at the LHC [10–23].

TeV scale SUSY  $B-L$  is one of the simplest models that provides viable and testable solution to the two puzzles of the DM and the neutrino masses. However, in order to fulfill the experimental measurements for the light-neutrino masses, one of the following scenarios must be adopted: (i) Type I seesaw mechanism with very small Dirac neutrino Yukawa couplings,  $Y_\nu < \mathcal{O}(10^{-7})$  [3]. (ii) Inverse seesaw mechanism with order one Yukawa couplings and small mass scale  $\sim \mathcal{O}(1)$  KeV, that corresponds to scale of breaking a remnant discrete symmetry,  $(-1)^{L+S}$  [26]. This work was based on earlier papers in ref. [27, 28]. In the first case, due to the smallness of Dirac Yukawa couplings, the right-handed neutrino sector has a very suppressed interaction with the SM particle. Therefore, the prediction of SUSY  $B-L$  remains close to the MSSM ones. It turns out that the DM candidate of this model is still the lightest neutralino [29], which is a kind of a mixture of three neutral gauginos and four neutral Higgsinos.

In this paper we consider the scenario where the right-handed sneutrino in SUSY  $B-L$  with inverse seesaw is the lightest SUSY particle (LSP) and stable, so that it can be a cold DM candidate [30–35]. It is worth mentioning that in MSSM the left-handed sneutrino is the only stable weakly coupled neutral boson that can be a DM candidate [36–40]. However, the current limits in direct detection experiments rule out this possibility, since left-handed sneutrino has a tree level interaction with the  $Z$  gauge boson, hence its elastic scattering cross section with nucleons is quite large. In case of MSSM extension with TeV right-handed neutrino ( $N$ ) superfields, the interaction of  $N$  with the SM particles can be obtained from the superpotential:

$$\mathcal{W} = \mathcal{W}_{\text{MSSM}} + Y_\nu N^c L H_2. \quad (1.1)$$

Due to smallness of  $Y_\nu$ , the annihilation cross section of the right-handed sneutrinos is very suppressed, hence its relic density is larger than the WMAP limit. Moreover, since they cannot couple to the quark sectors, the direct detection experiments such as CDMS II cannot be tested in this case.

We will show that in the SUSY  $B-L$  model with  $Y_\nu \sim \mathcal{O}(1)$  the dominant annihilation channel for right-handed sneutrino is given by the four point interactions, leading to  $h^0$  and  $h^0$ . While the effective couplings of right-handed sneutrinos with quarks are obtained through the exchange of TeV scale  $B-L$  gauge boson,  $Z_{B-L}$ . Thus, one finds that the WMAP results of the DM relic abundance and CDMS II/XENON for direct detection can be accommodated in SUSY  $B-L$  simultaneously [41–44].

The paper is organized as follows. In section 2 we analyze the supersymmetric  $B-L$  model with Inverse Seesaw Mechanism. We study the neutrino sector and the neutrino

mass eigenstates. In section 3 we analyze the  $B - L$  right-handed sneutrino mass and interactions. Section 4 is devoted for the analysis of the relic abundance of  $B - L$  right-handed sneutrino. We show that the WMAP limits can be easily satisfied in a wide range of parameter space. In section 5 we discuss the direct detection rate of  $B - L$  right-sneutrino DM. We show that the elastic cross section of our DM candidate with nucleon is consistent with the recent results of CDMS II/XENON experiment and it is detectable in near future experiments. In section 6 we discuss the indirect detection rate of  $B - L$  right-handed sneutrino. We show that the annihilation channels of right-handed sneutrino into leptons are subdominant, therefore it cannot account for the controversial PAMELA results. We also show that the muon flux generated from the right-handed sneutrino in the galactic center is much smaller than the Super-Kamiokande's limits. Finally we give our conclusions in section 7.

## 2 Supersymmetric $B - L$ model with inverse seesaw mechanism

The proposed TeV scale supersymmetric  $B - L$  extension of the SM is based on the gauge group  $SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_{B-L}$ , where the  $U(1)_{B-L}$  is spontaneously broken by a chiral singlet superfield  $\chi_1$  with  $B - L$  charge = +1 and  $\chi_2$  with  $B - L$  charge = -1. As in the conventional  $B - L$  model, a gauge boson  $Z_{B-L}$  and three chiral singlet superfields  $N_i$  with  $B - L$  charge = -1 are introduced for the consistency of the model. Finally, three chiral singlet superfields  $S_1$  with  $B - L$  charge = +2 and three chiral singlet superfields  $S_2$  with  $B - L$  charge = -2 are considered to implement the inverse seesaw mechanism. The superpotential of the leptonic sector of this model is given by

$$\mathcal{W} = Y_e E^c L H_1 + Y_\nu N^c L H_2 + Y_S N^c \chi_1 S_2 + \mu H_1 H_2 + \mu' \chi_1 \chi_2. \quad (2.1)$$

It is worth noting that the chiral singlet superfields  $\chi_2$  and  $N$  have the same  $B - L$  charge. Therefore, one may impose a discrete symmetry in order to distinguish them and to prohibit other terms beyond those given in eq. (2.1). In this case, the relevant soft SUSY breaking terms, assuming the usual universality assumptions, are as follows

$$\begin{aligned}
 -\mathcal{L}_{\text{soft}} = & \sum_{\phi} \tilde{m}_{\phi}^2 |\phi|^2 + Y_\nu^A \tilde{N}^c \tilde{L} H_2 + Y_e^A \tilde{E}^c \tilde{L} H_1 + Y_S^A \tilde{N}^c \tilde{S}_2 \chi_1 + B\mu H_1 H_2 + B\mu' \chi_1 \chi_2 \\
 & + \frac{1}{2} M_1 \tilde{B} \tilde{B} + \frac{1}{2} M_2 \tilde{W}^a \tilde{W}^a + \frac{1}{2} M_3 \tilde{g}^a \tilde{g}^a + \frac{1}{2} M_{B-L} \tilde{Z}_{B-L} \tilde{Z}_{B-L} + h.c., \quad (2.2)
 \end{aligned}$$

where the sum in the first term runs over  $\phi = H_1, H_2, \chi_1, \chi_2, \tilde{L}, \tilde{E}^c, \tilde{N}^c, \tilde{S}_1, \tilde{S}_2$  and  $Y_L^A \equiv Y_L A_L$  ( $L = e, \nu, S$ ) is the trilinear associated with lepton Yukawa coupling. In order to prohibit a possible large mass term  $M S_1 S_2$  in the above, we assume that the particles,  $N_i^c$ ,  $\chi_{1,2}$ , and  $S_2$  are even under matter parity, while  $S_1$  is an odd particle. The  $B - L$  symmetry is radiatively broken by the non-vanishing vacuum expectation values (VEVs)  $\langle \chi_1 \rangle = v'_1$  and  $\langle \chi_2 \rangle = v'_2$  [9]. The tree level potential  $V(\chi_1, \chi_2)$  is given by

$$V(\chi_1, \chi_2) = \frac{1}{2} g_{B-L}^2 (|\chi_2|^2 - |\chi_1|^2)^2 + \mu_1^2 |\chi_1|^2 + \mu_2^2 |\chi_2|^2 - \mu_3^2 (\chi_1 \chi_2 + h.c.). \quad (2.3)$$

At GUT scale,  $\mu_i^2 = m_0^2 + \mu'^2$ ,  $i = 1, 2$  and  $\mu_3^2 = -B\mu'$ . However, they have different evolution from GUT scale to TEV scale and  $\mu_2$  becomes negative, so that  $B - L$  is spontaneously broken [9]. The minimization of  $V(\chi_1, \chi_2)$  leads to the following condition:

$$v'^2 = (v_1'^2 + v_2'^2) = \frac{(\mu_1^2 - \mu_2^2) - (\mu_1^2 + \mu_2^2) \cos 2\theta}{2g_{B-L}^2 \cos 2\theta}, \quad (2.4)$$

The angle  $\theta$  is defined as  $\tan \theta = v_1'/v_2'$ . The minimization conditions also leads to

$$\sin 2\theta = \frac{2\mu_3^2}{\mu_1^2 + \mu_2^2}. \quad (2.5)$$

After  $B - L$  breaking, the  $Z_{B-L}$  gauge boson acquires a mass [3]:  $M_{Z_{B-L}}^2 = 4g_{B-L}^2 v'^2$ . The high energy experimental searches for an extra neutral gauge boson impose lower bounds on this mass. The stringent constraint on  $U(1)_{B-L}$  obtained from LEP II result, which implies [47]

$$\frac{M_{Z_{B-L}}}{g_{B-L}} > 6 \text{ TeV}. \quad (2.6)$$

Now we turn to the neutrino sector and show how the observed light-neutrino masses can be obtained with  $\mathcal{O}(1)$  Dirac neutrino Yukawa coupling. As can be seen from eq. (2.1), after  $B - L$  and EW symmetry breaking, the neutrino Yukawa interaction terms lead to the following mass terms:

$$\mathcal{L}_m^\nu = m_D \bar{\nu}_L N^c + M_N N^c S_2, \quad (2.7)$$

where  $m_D = Y_\nu v \sin \beta$  and  $M_N = Y_S v' \sin \theta$ . From this Lagrangian, one can easily observe that although the lepton number is broken through the spontaneous  $B - L$  symmetry breaking, a remnant symmetry:  $(-1)^{L+S}$  is survived, where  $L$  is the lepton number and  $S$  is the spin. After this global symmetry is broken at much lower scale, a mass term for  $S_2$  (and possibly for  $S_1$  as well) is generated. Therefore, the Lagrangian of neutrino masses, in the flavor basis, is given by:

$$\mathcal{L}_m^\nu = m_D \bar{\nu}_L N^c + M_N N^c S_2 + \mu_{S_2} S_2^2 (+\mu_{S_1} S_1^2). \quad (2.8)$$

In the basis  $\{\nu_L, N^c, S_2\}$ , the  $3 \times 3$  neutrino mass matrix of one generation takes the form:

$$\begin{pmatrix} 0 & m_D & 0 \\ m_D & 0 & M_N \\ 0 & M_N & \mu_{S_2} \end{pmatrix}. \quad (2.9)$$

The mixing matrix( $O$ ) for this mass matrix leads to the following light and heavy neutrino masses respectively in the limit of  $\mu_{S_i} \ll m_D, M_N$  (where  $i = 1, 2$ ):

$$m_{\nu_\ell} = \frac{m_D^2 \mu_{S_2}}{M_N^2 + m_D^2}, \quad m_{\nu_{H,H'}} = \pm \sqrt{M_N^2 + m_D^2} + \frac{1}{2} \frac{M_N^2 \mu_{S_2}}{M_N^2 + m_D^2}, \quad (2.10)$$

where

$$O \simeq \begin{pmatrix} \frac{M_N}{\sqrt{M_N^2+m_D^2}} & \frac{1}{\sqrt{2}} \frac{m_D}{\sqrt{M_N^2+m_D^2}} + \frac{3}{4\sqrt{2}} \frac{M_N^2 m_D \mu_{S_2}}{(M_N^2+m_D^2)^2} & \frac{1}{\sqrt{2}} \frac{m_D}{\sqrt{M_N^2+m_D^2}} - \frac{3}{4\sqrt{2}} \frac{M_N^2 m_D \mu_{S_2}}{(M_N^2+m_D^2)^2} \\ \frac{M_N m_D \mu_{S_2}}{(M_N^2+m_D^2)^{3/2}} & -\frac{1}{\sqrt{2}} - \frac{1}{4\sqrt{2}} \frac{M_N^2 \mu_{S_2}}{(M_N^2+m_D^2)^{3/2}} & \frac{1}{\sqrt{2}} - \frac{1}{4\sqrt{2}} \frac{M_N^2 \mu_{S_2}}{(M_N^2+m_D^2)^{3/2}} \\ -\frac{m_D}{\sqrt{M_N^2+m_D^2}} & \frac{1}{\sqrt{2}} \frac{M_N}{\sqrt{M_N^2+m_D^2}} - \frac{1}{4\sqrt{2}} \frac{M_N(M_N^2+4m_D^2)\mu_{S_2}}{(M_N^2+m_D^2)^2} & \frac{1}{\sqrt{2}} \frac{M_N}{\sqrt{M_N^2+m_D^2}} + \frac{1}{4\sqrt{2}} \frac{M_N(M_N^2+4m_D^2)\mu_{S_2}}{(M_N^2+m_D^2)^2} \end{pmatrix}. \quad (2.11)$$

Thus, the light neutrino mass can be of order eV, as required by the oscillation data, for a TeV scale  $M_N$ , provided  $\mu_{S_2}$  is sufficiently small,  $\mu_{S_2} \ll M_N$ . In this case, there is no any restriction imposed on the value of Dirac mass  $m_D$ . Therefore, the possibility of testing this type of model in LHC is quite feasible. Note that in the limit  $\mu_{S_2} \rightarrow 0$  which corresponds to the unbroken  $(-1)^{L+S}$  symmetry, we have massless light neutrinos. Therefore, a small non-vanishing  $\mu_{S_2}$  can be considered as a slight breaking of a this global symmetry. Hence, according to 't Hooft criteria, the smallness of  $\mu_{S_2}$  is natural. The possibility of generating small  $\mu_{S_2}$  radiatively has been discussed in ref. [45].

Finally, it is worth mentioning that the light neutrinos  $\nu_l$  have suppressed mixing (of order  $m_D \mu_{S_2} / (M_N^2 + m_D^2)$ ) with one type of the heavy neutrinos (say  $\nu_{H'}$ ) and a rather small mixing (of order  $m_D / M_N$ ) with the other type of heavy neutrinos ( $\nu_H$ ) by choosing appropriate parameters. The mixing between the heavy neutrino  $\nu_H$  and  $\nu_{H'}$  is maximal. In general, the physical neutrino states are given in terms of  $\nu_L$ ,  $N^c$ , and  $S_2$  as follows:

$$\nu_l \simeq \frac{M_N}{\sqrt{M_N^2+m_D^2}} \nu_L + \frac{M_N m_D \mu_{S_2}}{(M_N^2+m_D^2)^{3/2}} N^c - \frac{m_D}{\sqrt{M_N^2+m_D^2}} S_2 \simeq \nu_L + a_1 N^c - a_2 S_2 \quad (2.12)$$

$$\nu_H \simeq -\frac{1}{\sqrt{2}} \frac{m_D}{\sqrt{M_N^2+m_D^2}} \nu_L + \frac{1}{\sqrt{2}} N^c - \frac{1}{\sqrt{2}} \frac{M_N}{\sqrt{M_N^2+m_D^2}} S_2 \simeq \alpha (-a_2 \nu_L + N^c - S_2) \quad (2.13)$$

$$\nu_{H'} \simeq \frac{1}{\sqrt{2}} \frac{m_D}{\sqrt{M_N^2+m_D^2}} \nu_L + \frac{1}{\sqrt{2}} N^c + \frac{1}{\sqrt{2}} \frac{M_N}{\sqrt{M_N^2+m_D^2}} S_2 \simeq \alpha (a_2 \nu_L + N^c + S_2). \quad (2.14)$$

For  $m_D \simeq 100$  GeV,  $M_N \simeq 1$  TeV and  $\mu_{S_2} \simeq 1$  KeV, one finds that  $a_1 \sim \mathcal{O}(10^{-10})$ ,  $a_2 \sim \mathcal{O}(0.1)$ ,  $a_3 \sim \mathcal{O}(0.07)$  and  $\alpha \sim \sin \pi/4$ . In this respect, the gauge eigenstates for neutrinos can be expressed in terms of the mass eigenstates as follow:

$$\begin{pmatrix} \nu_L \\ N^c \\ S_2 \end{pmatrix} = O \begin{pmatrix} \nu_l \\ \nu_H \\ \nu_{H'} \end{pmatrix} \simeq \begin{pmatrix} 1 & \alpha a_2 & \alpha a_2 \\ a_1 & -\alpha & \alpha \\ -a_2 & \alpha & \alpha \end{pmatrix} \begin{pmatrix} \nu_l \\ \nu_H \\ \nu_{H'} \end{pmatrix}. \quad (2.15)$$

### 3 B – L right-handed sneutrino

In our model, the sneutrino mass matrix of one generation is given by  $8 \times 8$  matrix, which can be decomposed into the following two mass matrices: (i)  $6 \times 6$  mass matrix in basis of  $(\tilde{\nu}_L, \tilde{\nu}_L^\dagger, \tilde{N}, \tilde{N}^\dagger, \tilde{S}_2, \tilde{S}_2^\dagger)^T$ . (ii)  $2 \times 2$  mass matrix in basis of  $(\tilde{S}_1, \tilde{S}_1^\dagger)^T$ . The  $\tilde{S}_1$ 's are decoupled and have now interactions with the SM particles. Therefore, one can neglect it and focus

on the  $6 \times 6$  sneutrino mass matrix. In the flavor basis;  $\tilde{\nu} \equiv (\tilde{\nu}_L, \tilde{\nu}_L^\dagger, \tilde{N}, \tilde{N}^\dagger, \tilde{S}_2, \tilde{S}_2^\dagger)^T$ , the sneutrino mass matrix  $\tilde{M}_{\tilde{\nu}}^2$  is given as

$$M_{\tilde{\nu}}^2 = \begin{pmatrix} M_{\tilde{\nu}_L^\dagger \tilde{\nu}_L}^2 & 0 & (M_{\tilde{N}^\dagger \tilde{\nu}_L}^2)^\dagger & 0 & (M_{\tilde{S}_2^\dagger \tilde{\nu}_L}^2)^\dagger & 0 \\ 0 & (M_{\tilde{\nu}_L^\dagger \tilde{\nu}_L}^2)^T & 0 & (M_{\tilde{N}^\dagger \tilde{\nu}_L}^2)^T & 0 & (M_{\tilde{S}_2^\dagger \tilde{\nu}_L}^2)^T \\ M_{\tilde{N}^\dagger \tilde{\nu}_L}^2 & 0 & M_{\tilde{N}^\dagger \tilde{N}}^2 & 0 & (M_{\tilde{S}_2^\dagger \tilde{N}}^2)^\dagger & (M_{\tilde{S}_2 \tilde{N}}^2)^\dagger \\ 0 & (M_{\tilde{N}^\dagger \tilde{\nu}_L}^2)^* & 0 & (M_{\tilde{N}^\dagger \tilde{N}}^2)^T & (M_{\tilde{S}_2 \tilde{N}}^2)^T & (M_{\tilde{S}_2^\dagger \tilde{N}}^2)^T \\ M_{\tilde{S}_2^\dagger \tilde{\nu}_L}^2 & 0 & M_{\tilde{S}_2^\dagger \tilde{N}}^2 & (M_{\tilde{S}_2 \tilde{N}}^2)^* & M_{\tilde{S}_2^\dagger \tilde{S}_2}^2 & (M_{\tilde{S}_2 \tilde{S}_2}^2)^\dagger \\ 0 & (M_{\tilde{S}_2^\dagger \tilde{\nu}_L}^2)^* & M_{\tilde{S}_2 \tilde{N}}^2 & (M_{\tilde{S}_2^\dagger \tilde{N}}^2)^* & M_{\tilde{S}_2 \tilde{S}_2}^2 & (M_{\tilde{S}_2^\dagger \tilde{S}_2}^2)^T \end{pmatrix}, \quad (3.1)$$

where

$$M_{\tilde{\nu}_L^\dagger \tilde{\nu}_L}^2 = \tilde{m}_{\nu_L}^2 + v^2 \cos^2 \beta Y^{e\dagger} Y^e + M_D^\dagger M_D + \frac{m_Z^2}{2} \cos 2\beta - \frac{M_{Z_{B-L}}^2}{4} (1 - \cot^2 \theta), \quad (3.2)$$

$$M_{\tilde{N}^\dagger \tilde{N}}^2 = \tilde{m}_N^2 + v'^2 \sin^2 \theta Y^S Y^{S\dagger} + M_D M_D^\dagger - \frac{M_{Z_{B-L}}^2}{4} (1 - \cot^2 \theta), \quad (3.3)$$

$$M_{\tilde{S}_2^\dagger \tilde{S}_2}^2 = \tilde{m}_{S_2}^2 + |\mu_{S_2}|^2 + M_N^\dagger M_N - \frac{M_{Z_{B-L}}^2}{2} (1 - \cot^2 \theta), \quad (3.4)$$

$$M_{\tilde{N}^\dagger \tilde{\nu}_L}^2 = \mu^* v \cos \beta Y^\nu + v \sin \theta Y_A^\nu, \quad M_{\tilde{S}_2^\dagger \tilde{\nu}_L}^2 = v v' \sin \theta \sin \beta Y^{S\dagger} Y^\nu, \quad (3.5)$$

$$M_{\tilde{S}_2^\dagger \tilde{N}}^2 = \mu' v' \cos \theta Y^{S\dagger} + v' \sin \theta Y_A^{\nu\dagger}, \quad M_{\tilde{S}_2 \tilde{S}_2}^2 = B' \mu_{S_2}, \quad M_{\tilde{S}_2 \tilde{N}}^2 = \mu_{S_2} v' \sin \theta Y_S^\dagger. \quad (3.6)$$

In the case of existing the mixing of  $\tilde{S}_2 \tilde{S}_2$  or  $\tilde{S}_2 \tilde{N}$ , the complex scalar DM splits in two real scalar and the lighter one is DM. If the mass split is small as well as the momentum of DM, inelastic scattering can be considered.

The mass matrix is diagonalized by unitary matrix  $\Gamma$  as

$$\Gamma^\dagger M_{\tilde{\nu}}^2 \Gamma = \text{diag}(m_{\tilde{\nu}_1^m}^2, m_{\tilde{\nu}_2^m}^2, m_{\tilde{\nu}_3^m}^2, m_{\tilde{\nu}_4^m}^2, m_{\tilde{\nu}_5^m}^2, m_{\tilde{\nu}_6^m}^2). \quad (3.7)$$

Thus, the mass eigenstates  $\tilde{\nu}^m$  are defined as  $\tilde{\nu}_i = \Gamma_{ij} \tilde{\nu}_j^m$ . In general, the lightest sneutrino can be written as a linear combination of the sneutrino mass eigenstate, as follows:


$$\tilde{\nu}_1^m = \Gamma_{11}^\dagger \tilde{\nu}_L + \Gamma_{12}^\dagger \tilde{\nu}_L^\dagger + \Gamma_{13}^\dagger \tilde{N} + \Gamma_{14}^\dagger \tilde{N}^\dagger + \Gamma_{15}^\dagger \tilde{S}_2 + \Gamma_{16}^\dagger \tilde{S}_2^\dagger. \quad (3.8)$$

However if one considers large  $\tan \beta$  and small  $\tan \theta$  limits, one finds that the lightest sneutrino is mainly obtained from the  $(\tilde{N}, \tilde{S}_2)$  sector. Therefore, it can be expressed as

$$\tilde{\nu}_1 \simeq \Gamma_{13}^\dagger \tilde{N} + \Gamma_{14}^\dagger \tilde{N}^\dagger + \Gamma_{15}^\dagger \tilde{S}_2 + \Gamma_{16}^\dagger \tilde{S}_2^\dagger. \quad (3.9)$$

The mass of this particle, which we call “ $B - L$  right-handed sneutrino” depends on the universal soft scalar mass  $m_0$  and on the parameters  $\mu$  and  $\mu'$ . In figure 1, we display the mass range of  $B - L$  right-handed sneutrino as function of the most relevant parameters  $m_0$  and  $\mu$ . If this particle is the lightest SUSY particle, then it is stable and can be considered as an interesting candidate for DM.

Now we consider the relevant interactions of the  $B - L$  right-handed sneutrino. From the superpotential  $\mathcal{W}$  in eq. (2.1) one gets the following interacting Lagrangian of  $\tilde{N}$  in the



**Figure 1.** A contour plot for the mass of  $B - L$  right-handed sneutrino in terms of the most relevant parameters: soft mass term  $m_0$  and  $\mu$  term and for  $\theta \simeq \beta \simeq \pi/4$  and  $Y_\nu \simeq 1$ . The red region represents  $m_{\tilde{\nu}_1} < 115$  GeV, which is kinematically excluded because DM mass is lighter than the SM Higgs mass. The green region represents  $115 \text{ GeV} \leq m_{\tilde{\nu}_1} < 200$  GeV. The blue region represents  $200 \text{ GeV} \leq m_{\tilde{\nu}_1} < 300$  GeV.

flavor basis:

$$\begin{aligned} \mathcal{L}_{\text{int}}^W = & Y_{\nu ij} \tilde{N}_i^\dagger \left[ (\tilde{H}_2^0)^c P_L \nu_{Lj} - (\tilde{H}_2^+)^c P_L \ell_{Lj}^- \right] + Y_{Sij} \tilde{N}_i^\dagger (\tilde{S}_{2j})^c P_L \tilde{\chi}_1 \\ & + Y_{\nu ij}^A \tilde{N}_i^\dagger \left( \tilde{\nu}_{Lj} H_2^0 - \tilde{\ell}_{Lj}^- H_2^+ \right) + Y_{Sij}^A \tilde{N}_i^\dagger \tilde{S}_{2j} \chi_1 + h.c., \end{aligned} \quad (3.10)$$

Also from the  $F$ -term contributions to the scalar potential one finds the following interaction terms:

$$\begin{aligned} \mathcal{L}_{\text{int}}^F = & -|H_1^-|^2 \tilde{\nu}_{Li}^\dagger (Y_e^\dagger Y_e)_{ij} \tilde{N}_j - (|H_2^+|^2 + |H_2^0|^2) \tilde{N}_i^\dagger (Y_\nu^\dagger Y_\nu)_{ij} \tilde{N}_j \\ & - (H_1^{0\dagger} H_2^+ + H_1^- \dagger H_2^0) \tilde{N}_i^\dagger (Y^\nu Y^{e\dagger})_{ij} \tilde{\ell}_{Rj} - (\tilde{N}^\dagger Y_\nu \tilde{\nu}_L) (\tilde{\nu}_L^\dagger Y_\nu^\dagger \tilde{N}) - \mu^* H_1^{0\dagger} (\tilde{N}^\dagger Y_\nu \tilde{\nu}_L) \\ & - (\tilde{N}^\dagger Y_\nu \tilde{\ell}_L) (\tilde{\ell}_L^\dagger Y_\nu^\dagger \tilde{N}) - \mu^* H_1^- \dagger (\tilde{N}^\dagger Y_\nu \tilde{\ell}_L) - (\tilde{S}_2^\dagger Y_S^\dagger \tilde{N}) (\tilde{N}^\dagger Y_S \tilde{S}_2) - \mu^* \chi_2^\dagger (\tilde{N}^\dagger Y_S \tilde{S}_2) \\ & - |\chi_1|^2 \tilde{N}_i^\dagger (Y_S Y_S^\dagger) \tilde{N}_j - \mu_{S_2}^* \chi_1 \tilde{N}_i^\dagger Y_{Sij} \tilde{S}_{2j}^\dagger + h.c.. \end{aligned} \quad (3.11)$$

Next the interactions of  $\tilde{N}$  with the gauge fields lead to the following Lagrangian:

$$\begin{aligned} \mathcal{L}_{\text{int}}^G = & g_{B-L}^2 Z_{B-L}^2 \tilde{N}_i^\dagger \tilde{N}_i - ig_{B-L} Z_{B-L}^\mu \left( \tilde{N}_i \partial_\mu \tilde{N}_i^\dagger - \tilde{N}_i^\dagger \partial_\mu \tilde{N}_i \right) \\ & - i\sqrt{2} g_{B-L} \tilde{N}_i N_i^c \tilde{Z}_{B-L} + h.c.. \end{aligned} \quad (3.12)$$

Finally, the  $D$ -term implies that

$$\begin{aligned} \mathcal{L}_{\text{int}}^D = & g_{B-L}^2 (\tilde{N}^\dagger \tilde{N}) \left( -|\tilde{\nu}_L|^2 - |\tilde{N}|^2 + \frac{|\tilde{u}_L|^2}{3} + \frac{|\tilde{d}_L|^2}{3} + \frac{|\tilde{u}_R|^2}{3} + \frac{|\tilde{d}_R|^2}{3} \right. \\ & \left. - |\tilde{\ell}_L|^2 - |\tilde{\ell}_R|^2 + 2|\tilde{S}_1|^2 - 2|\tilde{S}_2|^2 + |\tilde{\chi}_1|^2 - |\tilde{\chi}_2|^2 \right). \end{aligned} \quad (3.13)$$



In our analysis for  $B - L$  right-handed sneutrino ( $\tilde{\nu}_1$ ) DM, the following assumptions are considered for simplification: (i) Each sector consists of one generation only. (ii)  $m_{SM} < m_{DM} < m_{SUSY}$ ,  $m_{h^\pm}$ ,  $m_{\chi_{1,2}}$ ,  $M_{Z_{B-L}}$ ,  $m_{S_{1,2}}$ , where  $m_{SM}$  is standard model particles including of the lightest neutral Higgs boson ( $h_0$ ),  $m_{SUSY}$  is the supersymmetric particles, and  $m_{DM}$  is the mass of the lightest SUSY sneutrinos. (iii) In chargino sector,  $h^\pm$  mass is approximately given by  $\mu$ , in the limit of  $\mu$ ,  $M_2 \gg m_W$ , where  $m_W$  is the SM charged weak gauge boson mass. Under these assumptions, the relevant interacting Lagrangian of  $\tilde{\nu}_1$  consists of

$$\mathcal{L}_{\text{int}} = \mathcal{L}_{\text{int}}^W + \mathcal{L}_{\text{int}}^F + \mathcal{L}_{\text{int}}^G. \quad (3.14)$$

Note that the D-term is kinematically irrelevant, while the  $W, F, G$  interactions take the following simplified form:

$$\mathcal{L}_{\text{int}}^W = Y'_\nu \Gamma_{41} \tilde{\nu}_1^\dagger \left[ \overline{\tilde{h}^{0c}} P_L \nu_L - \overline{\tilde{h}^{+c}} P_L \ell_L^- \right] + h.c., \quad (3.15)$$

$$\mathcal{L}_{\text{int}}^F = -(Y_\nu^\dagger Y_\nu) \Gamma_{41} \Gamma_{31} |h^0|^2 \tilde{\nu}_1^\dagger \tilde{\nu}_1, \quad (3.16)$$

$$\mathcal{L}_{\text{int}}^G = -ig_{B-L} Z_{B-L}^\mu \Gamma_{41} \Gamma_{31} \left( \tilde{\nu}_1 \partial_\mu \tilde{\nu}_1^\dagger - \tilde{\nu}_1^\dagger \partial_\mu \tilde{\nu}_1 \right), \quad (3.17)$$

where  $Y'_\nu$  is considered to be included in the mixing of Higgsino and Chargino. Hereafter we use it as  $Y_\nu$ . Eq. (3.15) might be relevant to the indirect detection, which will be discussed in the section 6. Eq. (3.16) is more relevant to the WMAP experiment, which will be discussed in the next section. Eq. (3.17) is applied to analyze the direct detection as CDMS II/XENON, which will be also discussed in the section 5.

#### 4 Relic abundance of $B - L$ right-handed sneutrino


In this section, we compute the relic abundance of  $B - L$  right-handed sneutrino DM. We consider the standard computation of the cosmological abundance, where  $\tilde{\nu}_1$  is assumed to be in thermal equilibrium with the SM particles in the early universe and decoupled when it was non-relativistic. Therefore, the  $\tilde{\nu}_1$  density can be obtained by solving the Boltzmann equation:

$$\frac{dn_{\tilde{\nu}_1}}{dt} + 3Hn_{\tilde{\nu}_1} = - \langle \sigma_{\tilde{\nu}_1}^{\text{ann}} v \rangle [(n_{\tilde{\nu}_1})^2 - (n_{\tilde{\nu}_1}^{\text{eq.}})^2], \quad (4.1)$$

where  $n_{\tilde{\nu}_1}$  is  $\tilde{\nu}_1$  number density with  $\rho_{\tilde{\nu}_1} = m_{\tilde{\nu}_1} n_{\tilde{\nu}_1}$ . One usually defines  $\Omega_{\tilde{\nu}_1} = \rho_{\tilde{\nu}_1} / \rho_c$ , where  $\rho_c$  is the critical mass density. In addition,  $\langle \sigma_{\tilde{\nu}_1}^{\text{ann}} v \rangle$  is the thermal averaged of the total cross section for  $\tilde{\nu}_1$  annihilation into SM lighter particles times the DM relative velocity  $v$ . For non-relativistic  $\tilde{\nu}_1$ , the thermal averaged annihilation cross section,  $\langle \sigma_{\tilde{\nu}_1}^{\text{ann}} v \rangle$ , can be approximated as follows [51]:

$$\langle \sigma_{\tilde{\nu}_1}^{\text{ann}} v \rangle \simeq a_{\tilde{\nu}_1} + b_{\tilde{\nu}_1} v^2, \quad (4.2)$$

where  $a_{\tilde{\nu}_1}$   $b_{\tilde{\nu}_1}$  are the coefficients coming from  $s$ -wave and  $p$ -wave of  $\tilde{\nu}_1 \tilde{\nu}_1$  annihilation, respectively.



**Figure 2.** Possible annihilation channels of  $\tilde{\nu}_1$ . The second diagram gives a sub-dominant contribution, however it may be relevant for indirect detection processes.

From eqs. (3.15)–(3.17), one finds that the dominant annihilation channels of  $\tilde{\nu}_1$  are given in figure 2. It turns out the annihilation of  $B-L$  right-handed sneutrino into SM-like neutral Higgs, through the four point interaction vertex, gives the dominant contribution. The tree level annihilation channel  $\tilde{\nu}_1 \tilde{\nu}_1 \rightarrow \ell^+ \ell^-$  is suppressed by the mass of the chargino exchanged particle. This channel may be relevant for the indirect detection processes which will be discussed later. Our computation for the annihilation cross section leads to the following  $a_{\tilde{\nu}_1}$   $b_{\tilde{\nu}_1}$ :

$$a_{\tilde{\nu}_1} = \frac{\beta'_{h^0}}{32\pi m_{\tilde{\nu}_1}^2} |Y_\nu \Gamma_{31} \Gamma_{41}|^4, \tag{4.3}$$

$$b_{\tilde{\nu}_1} = \frac{\beta'_{h^0} (x_{h^0}^2 - 1)}{128\pi m_{\tilde{\nu}_1}^2} |Y_\nu \Gamma_{31} \Gamma_{41}|^4, \tag{4.4}$$

where

$$z_a = \frac{m_a}{m_{\tilde{\nu}_1}}, \quad \beta_a'^2 = 1 - z_a^2, \quad x_a^2 = \frac{z_a^2}{2(1 - z_a^2)}, \tag{4.5}$$

here we define that  $a$  is a final-state particle. From eq. (4.1) one finds that the relic abundance  $\Omega_{\tilde{\nu}_1} h^2$  is given by [52]


$$\Omega_{\tilde{\nu}_1} h^2 \simeq \frac{8.76 \times 10^{-11} \text{GeV}^{-2}}{g_*^{1/2}(T_F) (a_{\tilde{\nu}_1}/x_F + 3b_{\tilde{\nu}_1}/x_F^2)}, \tag{4.6}$$

where

$$x_F = \ln \frac{0.0955 m_{\text{pl}} m_{\tilde{\nu}_1} (a_{\tilde{\nu}_1} + 6b_{\tilde{\nu}_1}/x_F)}{(g_*^{1/2}(T_F) x_F)^{1/2}}. \tag{4.7}$$

Here  $m_{\text{pl}}$  is the Planck mass ( $1.22 \times 10^{19}$  GeV) and  $g_*(T_F)$  enumerates the degrees of freedom of relativistic particles at the freeze out temperature  $T_F$ , which can be fixed as  $g_*^{1/2}(T_F) = 10$ . From the above expressions, one notes that  $\tilde{\nu}_1$  relic abundance depends only on the right-sneutrino mass and the annihilation cross section coefficients  $a_{\tilde{\nu}_1}$  and  $b_{\tilde{\nu}_1}$ .

Given  $a_{\tilde{\nu}_1}$   $b_{\tilde{\nu}_1}$  we can determine the freeze out temperature  $T_F$ , below which the  $B-L$  right-handed sneutrino annihilation rate is smaller than the expansion rate of the universe



**Figure 3.** Relic density  $\Omega_{\tilde{\nu}_1} h^2$  as function of the  $B - L$  right-handed sneutrino mass  $m_{\tilde{\nu}_1}$  for three values of Dirac neutrino Yukawa coupling. The region between two lines is allowed by the experiment of WMAP [1].

and then computing the relic density  $\Omega_{\tilde{\nu}_1} h^2$ . In our numerical computation we assume a universal soft mass ( $m_0$ ) and fix the other parameter as follows:

$$\begin{aligned} m_{h^0} &= 115 \text{ GeV}, & M_{Z_{B-L}} &= 6 \text{ TeV}, & M_N &= 1 \text{ TeV}, & \mu_{S_{1,2}} &= 1 \text{ KeV}, & v &= 175 \text{ GeV}, \\ m_D &= 100 \text{ GeV}, & B'_{1,2} = v' = \mu^{(\prime)} &= 1 \text{ TeV}, & Y_S &= 0.1, & Y^e &= 0.05, & Y_\nu^A = Y_S^A &= 0.1 \text{ TeV}, \end{aligned} \quad (4.8)$$

In addition we analyze the relic density of  $\tilde{\nu}_1$  in the following regions of the parameter  $\mu$ ,  $\beta$ , and  $\theta$ :

$$0 < \theta, \quad \beta < \pi, \quad 50 \text{ GeV} < \mu, \quad m_0 < 300 \text{ GeV}. \quad (4.9)$$

In figure 3, we present the values of relic abundance of  $B - L$  right-handed sneutrino  $\Omega_{\tilde{\nu}_1} h^2$  as a function of  $m_{\tilde{\nu}_1}$  for the following values of Dirac neutrino Yukawa coupling:  $Y_\nu = 0.5, 0.7, 1.0$ . We require  $B - L$  right-handed sneutrino relic density to be  $0.09 < \Omega_{\tilde{\nu}_1} h^2 < 0.12$  in order to be consistent with WMAP results at  $3\sigma$  [53]. As can be seen from this figure, smaller values of Dirac neutrino Yukawa coupling are favored and lead to more allowed points that satisfy the WMAP observational limits of DM relic density. Also we find that the smaller  $Y_\nu$  is considered, the smaller DM mass one obtains.


## 5 Direct detection

In this section we discuss the possibility to detect our  $B - L$  right-handed sneutrino in direct detection experiments such as CDMS (II) [2] and XENON 100 experiment [54]. The general form of the elastic scattering cross section between DM  $\tilde{\nu}_1$  and nuclei  $N$  is given by [51, 59]

$$\sigma_{\tilde{\nu}_1-N}^{\text{vec}} = \frac{m_r^2}{16\pi} |b_N|^2. \quad (5.1)$$

The reduced mass  $m_r$  is defined as

$$m_r = \left( \frac{1}{m_{\tilde{\nu}_1}} + \frac{1}{M} \right)^{-1}, \quad (5.2)$$



**Figure 4.** Our dominant diagram for the direct detection.

where  $M$  is the nuclei mass. The coefficient  $b_N$  is given by

$$b_N = (A - Z)b_n + Zb_p, \quad b_p = 2b_u + b_d, \quad b_n = b_u + 2b_d, \quad (5.3)$$

Here  $A$  and  $Z$  are the mass number and the atomic number, respectively. The effective Lagrangian parameters  $b_u$  and  $b_d$  are defined as

$$\mathcal{L}_{\text{eff}} = b_q X^\mu \bar{q} \gamma^\mu q, \quad q = (u, d). \quad (5.4)$$

Here  $X^\mu$  is a general form of the vector current. In case of fermionic DM  $X^\mu$  is given by  $X^\mu \simeq \bar{f} \gamma^\mu f$ . While for bosonic DM, it is defined as  $X^\mu \simeq ib^\dagger \partial^\mu b - ib \partial^\mu b^\dagger$ .

In our  $B - L$  case, the elastic scattering cross section of the right-handed sneutrino with a given nuclei has a spin-independent contribution arising from  $Z_{B-L}$  gauge boson exchange diagrams, as can be seen in figure 4. The interactions between the right-handed sneutrino  $\tilde{\nu}_1$  and  $Z_{B-L}$  boson in  $B - L$  model are

$$\mathcal{L} \supset -ig_{B-L} Z_{B-L}^\mu \Gamma_{41} \Gamma_{31} \left( \tilde{\nu}_1 \partial_\mu \tilde{\nu}_1^\dagger - \tilde{\nu}_1^\dagger \partial_\mu \tilde{\nu}_1 \right) - \frac{1}{3} g_{B-L} Z_{B-L}^\mu \bar{u} \gamma_\mu u - \frac{1}{3} g_{B-L} Z_{B-L}^\mu \bar{d} \gamma_\mu d. \quad (5.5)$$

Writing down the effective interaction, one obtains


$$\begin{aligned} i\mathcal{L}_{\text{eff}} \supset & -\frac{|\Gamma_{41} \Gamma_{31}|^2 g_{B-L}^2}{M_{Z_{B-L}}^2} \frac{1}{3} \left( \tilde{\nu}_1 \partial_\mu \tilde{\nu}_1^\dagger - \tilde{\nu}_1^\dagger \partial_\mu \tilde{\nu}_1 \right) \bar{u} \gamma^\mu u \\ & -\frac{|\Gamma_{41} \Gamma_{31}|^2 g_{B-L}^2}{M_{Z_{B-L}}^2} \frac{1}{3} \left( \tilde{\nu}_1 \partial_\mu \tilde{\nu}_1^\dagger - \tilde{\nu}_1^\dagger \partial_\mu \tilde{\nu}_1 \right) \bar{d} \gamma^\mu d. \end{aligned} \quad (5.6)$$

Assuming  $|\Gamma_{41} \Gamma_{31}|^2 \simeq 1$  for simplicity, one finds

$$b_p = b_n = i \frac{g_{B-L}^2}{M_{Z_{B-L}}^2}. \quad (5.7)$$

Therefore,  $b_N$  is given by

$$b_N = iA \frac{g_{B-L}^2}{M_{Z_{B-L}}^2}. \quad (5.8)$$



**Figure 5.** Elastic scattering cross section between a nuclei and the  $B - L$  right-handed sneutrino DM as function of the DM mass for  $g_{B-L} = 1$ .

Thus, the elastic scattering cross section of  $B - L$  right-handed sneutrino is given by

$$\sigma_{\tilde{\nu}_1-N}^{Z_{B-L}} = \frac{m_r^2}{16\pi} |A|^2 \frac{g_{B-L}^4}{M_{Z_{B-L}}^4}. \tag{5.9}$$

In figure 5 we depict the relation between the cross section and  $B - L$  right-sneutrino mass. As can be seen from this figure, the following upper bound on  $\sigma_{\tilde{\nu}_1-N}^{Z_{B-L}}$  can be obtained:

$$\sigma_{\tilde{\nu}_1-N}^{Z_{B-L}} \leq 6.2 \times 10^{-45} \text{ cm}^2. \tag{5.10}$$

It is also remarkable that the elastic cross section is quite insensitive to the  $B - L$  right-handed sneutrino mass  $m_{\tilde{\nu}_1}$ . However, one observes that a light sneutrino  $\sim \mathcal{O}(100)$  GeV is more favored by direct detection experimental results. The current limits from CDMS II and XENON experiments indicate to a lower-bound of order  $3.7 \times 10^{-44} \text{ cm}^2$ . This suggests that our  $B - L$  right-handed sneutrino DM is expected to be detected in the direct detection experiments in near future.

Before concluding this section, it is worth mentioning that the XENON 100 [55] experiment has recently presented new limits on the WIMP-nucleon cross section for inelastic DM. These limits are due to a DM run with 100.9 live-days of data, taken from January to June 2010. It was shown that  $\sigma_{\chi-N} < 10^{-41} \text{ cm}^2$  can be extracted for  $m_\chi > 100 \text{ GeV}$ . Where  $\chi$  is a generic dark matter. The bound rules out the explanation of controversial DAMA/LIBRA modulation results, as being due to inelastic DM. The bound obtained on  $\sigma_{\chi-N}$  from the inelastic DM analysis of XENON 100 should be considered carefully, since a minimum velocity for DM to scatter in a detector is introduced, hence a large amount of fiducial is needed. Nevertheless, as can be seen from eq. (52), the cross section of our  $B - L$  sneutrino is well below the XENON 100 bound.

## 6 Indirect detection

### 6.1 PAMELA and Fermi-LAT experiments


As advocated in the introduction, the indirect searches for DM by the Space Observatory PAMELA [64] and Fermi-LAT [65, 66], indicate that the DM may contribute to the positron flux by direct annihilation in  $\ell^+\ell^-$ . This is one of the main feature of our DM candidate  $B - L$  right-handed sneutrino, therefore, it important to investigate the possibility that it accounts for these results. PAMELA collaboration reported excess flux between 8 and 80 GeV, with no excess in the corresponding anti-proton flux. Also ATIC and Fermi-LAT balloon experiments have shown excess electron and positron flux at energies around 10–1000 GeV. While there are plausible astrophysical explanations for these excesses, such as local pulsars and supernovas remnants, they could also result from DM annihilation. Note that in order to explain the Fermi-LAT experiment by DM annihilation, the DM mass must be of order  $\mathcal{O}(\text{TeV})$ . As shown in the previous section, such heavy DM mass is not favored by of direct detections. Therefore, in our analysis, we assume that the Fermi-LAT experiment may be saturated by considering an astrophysical background and we will focus on PAMELA measurement.

It is known that if the DM annihilation is to explain the observed anomalous flux, a large annihilation cross section,  $\langle\sigma v\rangle \simeq 10^{-24} \text{ cm}^3\text{s}^{-1}$ , is required to fit the excess flux, which is incompatible with straightforward estimates of the relic DM abundance in conventional cosmological models. Otherwise, a huge, unexplained, boost factor must be introduced. In our SUSY  $B - L$  model, the  $B - L$  right-handed sneutrino annihilates into  $\ell^+\ell^-$  channels, as shown in the second diagram of figure 2. However, as discussed in the previous section that these channels give sub-dominate contribution to the annihilation process. Therefore, the corresponding annihilation cross section is  $< 10^{-27} \text{ cm}^3\text{s}^{-1}$ . In this case, one requires a huge boost factor  $\mathcal{O}(10^{5-6})$  at least in order to account for PAMELA results. In general, it is known that there are two mechanisms to enhance the cross section and may justify this large boost factor. The first is Breit-Wigner mechanism [67, 68] and the second is the Sommer-feld [69–71] enhancement. Breit-Wigner mechanism can not be implemented in our model, since there is no any diagrams with  $s$ -channel. On the other hand, the Sommer-feld enhancement requires higher DM mass compared to the mediated particles in order to obtain enough large boost factor  $> \mathcal{O}(10^{5-6})$ . This assumption we have already avoided in order to not spoil the direct detection. As a result, it is difficult for our  $B - L$  right-handed neutrino to explain the controversial results of PAMELA experiment.

### 6.2 Muon flux measurement from Super-Kamiokande

The high energy neutrinos induced by DM annihilations in the earth, the sun, and the galactic center is an important signal for the indirect detection of DM. Such energetic neutrinos induce upward through-going muons from charged current interactions provide the most effective signatures in Super-Kamiokande. The neutrino-induced muon flux is evaluated from the neutrino flux [75, 76] as

$$F_{\mu^+\mu^-}^{(\text{ann})} \simeq 5.9 \times 10^{-15} \text{ cm}^{-2}\text{s}^{-1} \times \sum_F S_F \left( \frac{\langle\sigma v\rangle_F}{10^{-23} \text{ cm}^3\text{s}^{-1}} \right) \left( \frac{\langle J_2 \rangle_{\Omega} \Delta\Omega}{10} \right). \quad (6.1)$$



**Figure 6.** Feynman diagram of  $\tilde{\nu}_1 \tilde{\nu}_1 \rightarrow \ell^- \ell^+$ . Note that, in general, there is another contribution of  $\tilde{\nu}_1, \tilde{\nu}_1 \rightarrow \tilde{h}^0 \rightarrow \nu, \nu$  process. However, due to the massless limit of  $\nu$ , the cross section of this channel vanishes.

where we fix the typical values  $\langle J_2 \rangle_\Omega \Delta \Omega \sim 10$  for  $\psi_{\max} = 5^\circ$  in case of the Navarro-Frenk-White (NFW) halo density profile [77], and  $F$  collectively denotes the primary annihilation mode (e.g.,  $\tau^+ \tau^-$ , etc.). Notice that model dependence comes from  $\langle \sigma v \rangle_F$ , which will be shown later.  $S_F$  is defined as

$$S_F = \sum_{\nu_i} \int_{E_{\min}}^{E_{\text{in}}} \frac{dN_F^{(\nu_i)}}{dE} P_{\nu_i \nu_\mu} \left( \frac{E}{E_{\text{in}}} \right)^2 dE, \quad (6.2)$$

where  $E_{\text{in}} = m_{\tilde{\nu}_1}$ , and  $E_{\min}$  is the threshold energy above which the muons can be detected.  $P_{\nu_i \nu_\mu}$  denotes the probability that the  $\nu_i$  at the production is observed as  $\nu_\mu$  at the Earth due to the effect of neutrino oscillation. Regardless to the complicated expression of  $S_F$ , it is found as a fixed value depending on each of the  $F$  particle, which is, e.g., 0.2 for  $\mu$  pair, 0.14 – 0.18 for  $\tau$  pair, 0.78 for  $\nu_\tau$  pair, etc [72]. The limits from Super-Kamiokande are given in the ref. [78].

In SUSY  $B - L$  model, the relevant Lagrangian is given by

$$\mathcal{L}_{\text{int}}^W = Y_\nu \Gamma_{41} \tilde{\nu}_1^\dagger \left[ \tilde{h}^{0c} P_L \nu_L - \tilde{h}^{+c} P_L \ell_L^- \right] + h.c. \quad (6.3)$$


One can show that in the limit  $v \rightarrow 0$ , the thermally averaged cross section is given by

$$\begin{aligned} \langle \sigma v \rangle_F |_{v \rightarrow 0} &\simeq \sum_{F=\tau, \nu_\tau} \frac{3\beta'_F}{16\pi m_{\tilde{\nu}_1}^2} |Y_\nu \Gamma_{31} \Gamma_{41}|^4 z_F^2 (2 - z_F^2) \times \left( r_{F, \tilde{h}^+}^2 + r_{F, \tilde{h}^-}^2 + r_{F, \tilde{h}^-} r_{F, \tilde{h}^+} + r_{F, \tilde{h}^0}^2 \right) \\ &\simeq \sum_{F=\tau} \frac{9\beta'_F}{16\pi m_{\tilde{\nu}_1}^2} |Y_\nu \Gamma_{31} \Gamma_{41}|^4 z_F^2 (2 - z_F^2) \times r_{F, \tilde{h}^+}^2. \end{aligned} \quad (6.4)$$

Note that the term proportional to  $r_{F, \tilde{h}^0}^2$  vanishes in the massless limit of  $\nu$ . The parameters  $z_F$  and  $\beta'_F$  are as defined in eq. (4.5). The other parameters are defined as follows:

$$w_\alpha = \frac{m_\alpha}{m_{\tilde{\nu}_1}}, \quad r_{F\alpha} = (1 - z_F^2 + w_\alpha^2)^{-1}. \quad (6.5)$$

Here  $F$  refers to the final-state particle and  $\alpha$  denotes the mediated-particle. In our analysis for the muon flux, we consider the same set of inputs that we have used in the previous



**Figure 7.** Muon flux induced from the  $B - L$  annihilation in the galactic center. The horizontal line refers to the Super-Kamiokande upper bound which is given by  $\sim 5 \times 10^{-15} \text{cm}^{-2} \text{sec}^{-1}$ .

sections that leads to relic abundance within the WMAP limits:  $0.09 < \Omega h^2 < 0.12$ . Moreover, since we are considering the effect one generation, we assume  $\tau$  final state only. Using  $S_F$  that corresponds to  $\tau$  final state, one finds  $\langle \sigma v \rangle$

$$\begin{aligned}
 F_{\mu^+\mu^-}^{(\text{ann})} &\simeq 5.9 \times 10^{-15} \text{ cm}^{-2} \text{ s}^{-1} \times \sum_{F=\tau^\pm} S_F \left( \frac{\langle \sigma v \rangle_F}{8.56 \times 10^{-7}} \right) \left( \frac{\langle J_2 \rangle_\Omega \Delta \Omega}{10} \right) \\
 &\simeq 6.9 \times 10^{-9} \text{ cm}^{-2} \text{ s}^{-1} \times \sum_{F=\tau^\pm} S_F \left( \frac{9\beta'_F}{16\pi m_{\tilde{\nu}_1}^2} |Y_\nu \Gamma_{31} \Gamma_{41}|^4 z_F^2 (2 - z_F^2) \times r_{F, \tilde{h}^+}^2 \right). \quad (6.6)
 \end{aligned}$$

As we fixed the cone-half angle from the galactic center:  $\psi_{\text{max}} = 5^\circ$  which is maximum in the case of NFW profile, one finds that the Super-Kamiokande limit should be less than  $5 \times 10^{-15} \text{cm}^{-2} \text{sec}^{-1}$  (see figures of ref. [72]).

In figure 7, we plot the muon flux induced from the  $B - L$  annihilation in the galactic center. As can be seen from this figure, the result of the induced muon flux is  $\lesssim 10^{-20} \text{cm}^{-2} \text{sec}^{-1}$ , which is few order of magnitudes smaller than the upper bound of Super-Kamiokande. This result is expected since the thermally averaged cross cross is much smaller than the typical value required by indirect detections.

## 7 Conclusions

In this paper we considered the supersymmetric  $B - L$  model with inverse seesaw mechanism. We demonstrated that the lightest right-handed sneutrino in this model can be stable and a viable candidate for cold dark matter. We studied the relic abundance of the  $B - L$  right-handed sneutrino and showed that the WMAP result,  $\Omega h^2 \simeq 0.11$ , can be satisfied in a wide range of the parameter space. We emphasized that the dominate annihilation channel of  $B - L$  right-handed sneutrino is given by  $\tilde{\nu}_1 \tilde{\nu}_1 \rightarrow h^0 h^0$ , where  $h^0$  is the SM-like Higgs. We also studied the direct detection rate of  $B - L$  right-handed sneutrino. We found that its elastic cross section is consistent with the upper bounds of current experiments,



such as CDMS II and XENON. Our result of  $B - L$  right-handed sneutrino direct detection is promising and indicates that it can be detectable in near future experiment.

In addition, we have analyzed the indirect detection rate of  $B - L$  right-handed sneutrino. In particular, we focused on the observation of the Space Observatory PAMELA for positron flux and also on the neutrino-induced upward through-going muons in the Super-Kamiokande detector. We showed that although the  $B - L$  right-handed sneutrino annihilates at tree level into leptons, the corresponding cross section is much smaller than the required one for accommodating PAMELA results. Also the neutrino flux induced by the  $B - L$  right-handed sneutrino annihilations in the galactic center is much smaller than the Super-Kamiokande's limits.

Finally it is worth mentioning that our  $B - L$  right-handed sneutrino may be produced at the Large Hadron Collider (LHC) through the channel is  $q\bar{q} \rightarrow Z_{B-L} \rightarrow \tilde{\nu}_1^\dagger \tilde{\nu}_1$ . However, the amplitude of this channel vanishes identically due to the fact that left and right quarks or leptons has the same  $B - L$  quantum numbers. However, slepton/left-handed sneutrino may decay to right-handed sneutrino, which escapes the detector and gives the missing energy signal similar to other examples of cold DM.

## Acknowledgments

We would like to thank Y. Kajiyama for fruitful discussions. S. K. and H. O. acknowledge partial support from the Science and Technology Development Fund (STDF) project ID 437 and the ICTP project ID 30.

## References

- [1] WMAP collaboration, D.N. Spergel et al., *Wilkinson Microwave Anisotropy Probe (WMAP) three year results: Implications for cosmology*, *Astrophys. J. Suppl.* **170** (2007) 377 [[astro-ph/0603449](#)] [[SPIRES](#)].
- [2] THE CDMS-II collaboration, Z. Ahmed et al., *Dark Matter Search Results from the CDMS II Experiment*, *Science* **327** (2010) 1619 [[arXiv:0912.3592](#)] [[SPIRES](#)].
- [3] S. Khalil, *Low scale  $B - L$  extension of the Standard Model at the LHC*, *J. Phys.* **G 35** (2008) 055001 [[hep-ph/0611205](#)] [[SPIRES](#)].
- [4] M. Abbas and S. Khalil, *Neutrino masses, mixing and leptogenesis in TeV scale  $B - L$  extension of the standard model*, *JHEP* **04** (2008) 056 [[arXiv:0707.0841](#)] [[SPIRES](#)].
- [5] W. Emam and S. Khalil, *Higgs and  $Z'$  Phenomenology in  $B-L$  extension of the Standard Model at LHC*, *Eur. Phys. J.* **C 55** (2007) 625 [[arXiv:0704.1395](#)] [[SPIRES](#)].
- [6] R.N. Mohapatra and R.E. Marshak, *Local  $B-L$  Symmetry of Electroweak Interactions, Majorana Neutrinos and Neutron Oscillations*, *Phys. Rev. Lett.* **44** (1980) 1316 [Erratum *ibid* **44** (1980) 1643] [[SPIRES](#)].
- [7] F. Borzumati and A. Masiero, *Large Muon and electron Number Violations in Supergravity Theories*, *Phys. Rev. Lett.* **57** (1986) 961 [[SPIRES](#)].
- [8] C.S. Aulakh, A. Melfo, A. Rasin and G. Senjanović, *See-saw and supersymmetry or exact  $R$ -parity*, *Phys. Lett.* **B 459** (1999) 557 [[hep-ph/9902409](#)] [[SPIRES](#)].

- [9] S. Khalil and A. Masiero, *Radiative  $B - L$  symmetry breaking in supersymmetric models*, *Phys. Lett. B* **665** (2008) 374 [[arXiv:0710.3525](#)] [[SPIRES](#)].
- [10] K. Huitu, S. Khalil, H. Okada and S.K. Rai, *Signatures for right-handed neutrinos at the Large Hadron Collider*, *Phys. Rev. Lett.* **101** (2008) 181802 [[arXiv:0803.2799](#)] [[SPIRES](#)].
- [11] L. Basso, A. Belyaev, S. Moretti and C.H. Shepherd-Themistocleous, *Phenomenology of the minimal  $B-L$  extension of the Standard model:  $Z'$  and neutrinos*, *Phys. Rev. D* **80** (2009) 055030 [[arXiv:0812.4313](#)] [[SPIRES](#)].
- [12] L. Basso, A. Belyaev, S. Moretti and G.M. Pruna, *Probing the  $Z'$  sector of the minimal  $B - L$  model at future Linear Colliders in the  $e^+e^- \rightarrow \mu^+\mu^-$  process*, *JHEP* **10** (2009) 006 [[arXiv:0903.4777](#)] [[SPIRES](#)].
- [13] J.A. Aguilar-Saavedra, *Heavy lepton pair production at LHC: model discrimination with multi-lepton signals*, *Nucl. Phys. B* **828** (2010) 289 [[arXiv:0905.2221](#)] [[SPIRES](#)].
- [14] P. Fileviez Perez, T. Han and T. Li, *Testability of Type I Seesaw at the CERN LHC: Revealing the Existence of the  $B-L$  Symmetry*, *Phys. Rev. D* **80** (2009) 073015 [[arXiv:0907.4186](#)] [[SPIRES](#)].
- [15] S. Iso, N. Okada and Y. Orikasa, *The minimal  $B - L$  model naturally realized at TeV scale*, *Phys. Rev. D* **80** (2009) 115007 [[arXiv:0909.0128](#)] [[SPIRES](#)].
- [16] F.M.L. de Almeida et al., *Double seesaw mechanism in a left-right symmetric model with TeV neutrinos*, *Phys. Rev. D* **81** (2010) 053005 [[arXiv:1001.2162](#)] [[SPIRES](#)].
- [17] P. Nath et al., *The Hunt for New Physics at the Large Hadron Collider*, *Nucl. Phys. Proc. Suppl.* **200-202** (2010) 185 [[arXiv:1001.2693](#)] [[SPIRES](#)].
- [18] P. Fileviez Perez and M.B. Wise, *Baryon and Lepton Number as Local Gauge Symmetries*, *Phys. Rev. D* **82** (2010) 011901 [Erratum *ibid* **D 82** (2010) 079901] [[arXiv:1002.1754](#)] [[SPIRES](#)].
- [19] L. Basso, A. Belyaev, S. Moretti and G.M. Pruna, *Tree Level Unitarity Bounds for the Minimal  $B - L$  Model*, *Phys. Rev. D* **81** (2010) 095018 [[arXiv:1002.1939](#)] [[SPIRES](#)].
- [20] D. Cogollo, H. Diniz and C.A. de S. Pires, *Triple seesaw mechanism*, *Phys. Lett. B* **687** (2010) 400 [[arXiv:1002.1944](#)] [[SPIRES](#)].
- [21] L. Basso, A. Belyaev, S. Moretti, G.M. Pruna and C.H. Shepherd-Themistocleous,  *$Z'$  discovery potential at the LHC in the minimal  $B - L$  extension of the Standard Model*, *Eur. Phys. J. C* **71** (2011) 1613 [[arXiv:1002.3586](#)] [[SPIRES](#)].
- [22] S.K. Majee and N. Sahu, *Dilepton Signal of a Type-II Seesaw at CERN LHC: Reveals a TeV Scale  $B - L$  Symmetry*, *Phys. Rev. D* **82** (2010) 053007 [[arXiv:1004.0841](#)] [[SPIRES](#)].
- [23] A. Ibarra, E. Molinaro and S.T. Petcov, *TeV Scale See-Saw Mechanisms of Neutrino Mass Generation, the Majorana Nature of the Heavy Singlet Neutrinos and  $(\beta\beta)_{0\nu}$ -Decay*, *JHEP* **09** (2010) 108 [[arXiv:1007.2378](#)] [[SPIRES](#)].
- [24] Q.-H. Cao, S. Khalil, E. Ma and H. Okada, *Observable  $T_7$  Lepton Flavor Symmetry at the Large Hadron Collider*, *Phys. Rev. Lett.* **106** (2011) 131801 [[arXiv:1009.5415](#)] [[SPIRES](#)].
- [25] L. Basso, S. Moretti and G.M. Pruna, *Phenomenology of the minimal  $B - L$  extension of the Standard Model: the Higgs sector*, *Phys. Rev. D* **83** (2011) 055014 [[arXiv:1011.2612](#)] [[SPIRES](#)].

- [26] S. Khalil, *TeV Scale Gauged B-L With Inverse Seesaw Mechanism*, [arXiv:1004.0013](#) [SPIRES].
- [27] R.N. Mohapatra, *Mechanism for understanding small neutrino mass in superstring theories*, *Phys. Rev. Lett.* **56** (1986) 561 [SPIRES].
- [28] R.N. Mohapatra and J.W.F. Valle, *Neutrino mass and baryon-number nonconservation in superstring models*, *Phys. Rev.* **D 34** (1986) 1642 [SPIRES].
- [29] S. Khalil and H. Okada, *Dark Matter in B – L Extended MSSM Models*, *Phys. Rev.* **D 79** (2009) 083510 [[arXiv:0810.4573](#)] [SPIRES].
- [30] N. Arkani-Hamed, L.J. Hall, H. Murayama, D. Tucker-Smith and N. Weiner, *Neutrino masses at  $v^{3/2}$* , [hep-ph/0007001](#) [SPIRES].
- [31] N. Arkani-Hamed, L.J. Hall, H. Murayama, D. Tucker-Smith and N. Weiner, *Small neutrino masses from supersymmetry breaking*, *Phys. Rev.* **D 64** (2001) 115011 [[hep-ph/0006312](#)] [SPIRES].
- [32] Y. Grossman and H.E. Haber, *Sneutrino mixing phenomena*, *Phys. Rev. Lett.* **78** (1997) 3438 [[hep-ph/9702421](#)] [SPIRES].
- [33] D. Tucker-Smith and N. Weiner, *Inelastic dark matter*, *Phys. Rev.* **D 64** (2001) 043502 [[hep-ph/0101138](#)] [SPIRES].
- [34] D. Tucker-Smith and N. Weiner, *Inelastic dark matter at DAMA, CDMS and future experiments*, *Nucl. Phys. Proc. Suppl.* **124** (2003) 197 [[astro-ph/0208403](#)] [SPIRES].
- [35] D. Tucker-Smith and N. Weiner, *The status of inelastic dark matter*, *Phys. Rev.* **D 72** (2005) 063509 [[hep-ph/0402065](#)] [SPIRES].
- [36] L.J. Hall, T. Moroi and H. Murayama, *Sneutrino cold dark matter with lepton-number violation*, *Phys. Lett.* **B 424** (1998) 305 [[hep-ph/9712515](#)] [SPIRES].
- [37] L.E. Ibáñez, *The Scalar Neutrinos as the Lightest Supersymmetric Particles and Cosmology*, *Phys. Lett.* **B 137** (1984) 160 [SPIRES].
- [38] J.R. Ellis, J.S. Hagelin, D.V. Nanopoulos, K.A. Olive and M. Srednicki, *Supersymmetric relics from the big bang*, *Nucl. Phys.* **B 238** (1984) 453 [SPIRES].
- [39] J.S. Hagelin, G.L. Kane and S. Raby, *Perhaps Scalar Neutrinos Are the Lightest Supersymmetric Partners*, *Nucl. Phys.* **B 241** (1984) 638 [SPIRES].
- [40] K. Freese, *Can Scalar Neutrinos Or Massive Dirac Neutrinos Be the Missing Mass?*, *Phys. Lett.* **B 167** (1986) 295 [SPIRES].
- [41] S. Khalil and O. Seto, *Sterile neutrino dark matter in B – L extension of the standard model and galactic 511 keV line*, *JCAP* **10** (2008) 024 [[arXiv:0804.0336](#)] [SPIRES].
- [42] N. Okada and O. Seto, *Higgs portal dark matter in the minimal gauged  $U(1)_{B-L}$  model*, *Phys. Rev.* **D 82** (2010) 023507 [[arXiv:1002.2525](#)] [SPIRES].
- [43] T. Li and W. Chao, *Neutrino Masses, Dark Matter and B – L Symmetry at the LHC*, *Nucl. Phys.* **B 843** (2011) 396 [[arXiv:1004.0296](#)] [SPIRES].
- [44] S. Kanemura, O. Seto and T. Shimomura, *Masses of dark matter and neutrino from TeV scale spontaneous  $U(1)_{B-L}$  breaking*, [arXiv:1101.5713](#) [SPIRES].
- [45] E. Ma, *Radiative inverse seesaw mechanism for nonzero neutrino mass*, *Phys. Rev.* **D 80** (2009) 013013 [[arXiv:0904.4450](#)] [SPIRES].

- [46] R. Allahverdi, B. Dutta, K. Richardson-McDaniel and Y. Santoso, *Sneutrino Dark Matter and the Observed Anomalies in Cosmic Rays*, *Phys. Lett. B* **677** (2009) 172 [[arXiv:0902.3463](#)] [[SPIRES](#)].
- [47] M.S. Carena, A. Daleo, B.A. Dobrescu and T.M.P. Tait, *Z-prime gauge bosons at the Tevatron*, *Phys. Rev. D* **70** (2004) 093009 [[hep-ph/0408098](#)] [[SPIRES](#)].
- [48] Y. Kajiyama, S. Khalil and M. Raidal, *Electron EDM and soft leptogenesis in supersymmetric B-L extension of the standard model*, *Nucl. Phys. B* **820** (2009) 75 [[arXiv:0902.4405](#)] [[SPIRES](#)].
- [49] F. Bazzocchi, D.G. Cerdeno, C. Muñoz and J.W.F. Valle, *Calculable inverse-seesaw neutrino masses in supersymmetry*, *Phys. Rev. D* **81** (2010) 051701 [[arXiv:0907.1262](#)] [[SPIRES](#)].
- [50] P.S.B. Dev and R.N. Mohapatra, *TeV Scale Inverse Seesaw in SO(10) and Leptonic Non-Unitarity Effects*, *Phys. Rev. D* **81** (2010) 013001 [[arXiv:0910.3924](#)] [[SPIRES](#)].
- [51] K. Griest, *Cross sections, relic abundance and detection rates for neutralino dark matter*, *Phys. Rev. D* **38** (1988) 2357 [Erratum *ibid* **D 39** (1989) 3802] [[SPIRES](#)].
- [52] K. Griest, M. Kamionkowski and M.S. Turner, *Supersymmetric Dark Matter Above the W Mass*, *Phys. Rev. D* **41** (1990) 3565 [[SPIRES](#)].
- [53] I. Antoniadis, E. Dudas, D.M. Ghilencea and P. Tziveloglou, *Beyond the MSSM Higgs with D = 6 effective operators*, *Nucl. Phys. B* **848** (2011) 1 [[arXiv:1012.5310](#)] [[SPIRES](#)].
- [54] XENON100 collaboration, E. Aprile et al., *First Dark Matter Results from the XENON100 Experiment*, *Phys. Rev. Lett.* **105** (2010) 131302 [[arXiv:1005.0380](#)] [[SPIRES](#)].
- [55] XENON100 collaboration, E. Aprile et al., *Implications on Inelastic Dark Matter from 100 Live Days of XENON100 Data*, [arXiv:1104.3121](#) [[SPIRES](#)].
- [56] B. de Carlos and J.R. Espinosa, *Cold dark matter candidate in a class of supersymmetric models with an extra U(1)*, *Phys. Lett. B* **407** (1997) 12 [[hep-ph/9705315](#)] [[SPIRES](#)].
- [57] V. Barger et al., *Recoil detection of the lightest neutralino in MSSM singlet extensions*, *Phys. Rev. D* **75** (2007) 115002 [[hep-ph/0702036](#)] [[SPIRES](#)].
- [58] P. Gondolo et al., *DarkSUSY: computing supersymmetric dark matter properties numerically*, *JCAP* **07** (2004) 008 [[astro-ph/0406204](#)] [[SPIRES](#)].
- [59] M.W. Goodman and E. Witten, *Detectability of certain dark-matter candidates*, *Phys. Rev. D* **31** (1985) 3059 [[SPIRES](#)].
- [60] C. Arina and N. Fornengo, *Sneutrino cold dark matter, a new analysis: Relic abundance and detection rates*, *JHEP* **11** (2007) 029 [[arXiv:0709.4477](#)] [[SPIRES](#)].
- [61] DAMA collaboration, R. Bernabei et al., *First results from DAMA/LIBRA and the combined results with DAMA/NaI*, *Eur. Phys. J. C* **56** (2008) 333 [[arXiv:0804.2741](#)] [[SPIRES](#)].
- [62] S. Khalil, H.-S. Lee and E. Ma, *Bound on Z' Mass from CDMS II in the Dark Left-Right Gauge Model II*, *Phys. Rev. D* **81** (2010) 051702 [[arXiv:1002.0692](#)] [[SPIRES](#)].
- [63] H.-S. Lee, *Lightest U-parity Particle (LUP): a hidden sector dark matter candidate*, *AIP Conf. Proc.* **1078** (2009) 569 [[arXiv:0808.3600](#)] [[SPIRES](#)].
- [64] PAMELA collaboration, O. Adriani et al., *An anomalous positron abundance in cosmic rays with energies 1.5 – 100 GeV*, *Nature* **458** (2009) 607 [[arXiv:0810.4995](#)] [[SPIRES](#)].

- [65] THE FERMI LAT collaboration, A.A. Abdo et al., *Measurement of the Cosmic Ray  $e^+ + e^-$  spectrum from 20 GeV to 1 TeV with the Fermi Large Area Telescope*, *Phys. Rev. Lett.* **102** (2009) 181101 [[arXiv:0905.0025](#)] [[SPIRES](#)].
- [66] THE FERMI LAT collaboration, M. Ackermann et al., *Fermi LAT observations of cosmic-ray electrons from 7 GeV to 1 TeV*, [arXiv:1008.3999](#) [[SPIRES](#)].
- [67] M. Ibe, H. Murayama and T.T. Yanagida, *Breit-Wigner Enhancement of Dark Matter Annihilation*, *Phys. Rev. D* **79** (2009) 095009 [[arXiv:0812.0072](#)] [[SPIRES](#)].
- [68] D. Feldman, Z. Liu and P. Nath, *PAMELA Positron Excess as a Signal from the Hidden Sector*, *Phys. Rev. D* **79** (2009) 063509 [[arXiv:0810.5762](#)] [[SPIRES](#)].
- [69] J. Hisano, S. Matsumoto, M.M. Nojiri and O. Saito, *Non-perturbative effect on dark matter annihilation and gamma ray signature from galactic center*, *Phys. Rev. D* **71** (2005) 063528 [[hep-ph/0412403](#)] [[SPIRES](#)].
- [70] M. Pospelov and A. Ritz, *Astrophysical Signatures of Secluded Dark Matter*, *Phys. Lett. B* **671** (2009) 391 [[arXiv:0810.1502](#)] [[SPIRES](#)].
- [71] J.L. Feng, M. Kaplinghat and H.-B. Yu, *Sommerfeld Enhancements for Thermal Relic Dark Matter*, *Phys. Rev. D* **82** (2010) 083525 [[arXiv:1005.4678](#)] [[SPIRES](#)].
- [72] J. Hisano, M. Kawasaki, K. Kohri and K. Nakayama, *Neutrino Signals from Annihilating/Decaying Dark Matter in the Light of Recent Measurements of Cosmic Ray Electron/Positron Fluxes*, *Phys. Rev. D* **79** (2009) 043516 [[arXiv:0812.0219](#)] [[SPIRES](#)].
- [73] J.F. Beacom, N.F. Bell and G.D. Mack, *General upper bound on the dark matter total annihilation cross section*, *Phys. Rev. Lett.* **99** (2007) 231301 [[astro-ph/0608090](#)] [[SPIRES](#)].
- [74] H. Yuksel, S. Horiuchi, J.F. Beacom and S. Ando, *Neutrino Constraints on the Dark Matter Total Annihilation Cross Section*, *Phys. Rev. D* **76** (2007) 123506 [[arXiv:0707.0196](#)] [[SPIRES](#)].
- [75] G. Jungman, M. Kamionkowski and K. Griest, *Supersymmetric dark matter*, *Phys. Rept.* **267** (1996) 195 [[hep-ph/9506380](#)] [[SPIRES](#)].
- [76] S. Ritz and D. Seckel, *Detailed Neutrino Spectra From Cold Dark Matter Annihilations In The Sun*, *Nucl. Phys. B* **304** (1988) 877 [[SPIRES](#)].
- [77] J.F. Navarro, C.S. Frenk and S.D.M. White, *The Structure of Cold Dark Matter Halos*, *Astrophys. J.* **462** (1996) 563 [[astro-ph/9508025](#)] [[SPIRES](#)].
- [78] SUPER-KAMIOKANDE collaboration, S. Desai et al., *Search for dark matter WIMPs using upward through-going muons in Super-Kamiokande*, *Phys. Rev. D* **70** (2004) 083523 [*Erratum ibid* **D 70** (2004) 109901] [[hep-ex/0404025](#)] [[SPIRES](#)].