

Symmetry properties of Wilson loops with a Lagrangian insertion

Dmitry Chicherin^a and Johannes M. Henn^b

^a*LAPTh, Université Savoie Mont Blanc, CNRS,
B.P. 110, F-74941 Annecy-le-Vieux, France*

^b*Max-Planck-Institut für Physik, Werner-Heisenberg-Institut,
D-80805 München, Germany*

E-mail: chicherin@lapth.cnrs.fr, henn@mpp.mpg.de

ABSTRACT: Null Wilson loops in $\mathcal{N} = 4$ super Yang-Mills are dual to planar scattering amplitudes. This duality implies hidden symmetries for both objects. We consider closely related infrared finite observables, defined as the Wilson loop with a Lagrangian insertion, normalized by the Wilson loop itself. Unlike ratio and remainder functions studied in the literature, this observable is non-trivial already for four scattered particles and bears close resemblance to (finite parts of) scattering processes in non-supersymmetric Yang-Mills theory. Moreover, by integrating over the insertion point, one can recover information on the amplitude, as was recently done to compute the full four-loop cusp anomalous dimension. We study the general structure of the Wilson loop with a Lagrangian insertion, focusing in particular on its leading singularities and their (hidden) symmetry properties. Thanks to the close connection of the observable to integrands of MHV amplitudes, it is natural to expect that its leading singularities can be written as certain Grassmannian integrals. The latter are manifestly dual conformal. They also have a conformal symmetry, up to total derivatives. We find that, surprisingly, the conformal symmetry becomes an invariance in the frame where the Lagrangian insertion point is sent to infinity. Furthermore, we use integrability methods to study how higher Yangian charges act on the Grassmannian integral. We evaluate the n -particle observable both at tree- and at one-loop level, finding compact analytic formulas. These results are explicitly written in the form of conformal leading singularities, multiplied by transcendental functions. We then compare these formulas to known expressions for all-plus amplitudes in pure Yang-Mills theory. We find a remarkable new connection: the Wilson loop with Lagrangian insertion in $\mathcal{N} = 4$ super Yang-Mills appears to predict the maximal weight terms of the planar pure Yang-Mills all-plus amplitude. We test this relationship for the two-loop n -point Yang-Mills amplitude, as well as for the three-loop four-point amplitude.

KEYWORDS: Scattering Amplitudes, Wilson, 't Hooft and Polyakov loops

ARXIV EPRINT: [2202.05596](https://arxiv.org/abs/2202.05596)

Contents

1	Introduction	1
2	Recap of key features of the Wilson loop/scattering amplitudes duality	4
3	Null Wilson loop with a Lagrangian insertion	6
3.1	Definition, kinematics, and key properties	6
3.2	n -point tree-level formula in different representations	10
3.3	Conformal symmetry of the leading singularities	12
3.4	Two Grassmannian formulas for the leading singularities	14
4	Higher order symmetries and integrability	16
4.1	Proof of conformal and higher-order symmetries of the leading singularities r_n	17
4.1.1	Yangian generators	17
4.1.2	Spin-chain picture of the Yangian symmetry	18
4.1.3	Summary and discussion of symmetry relations	22
4.2	Formulation of higher-order symmetries of r_n in momentum space	23
4.3	Higher-order symmetries of R_n and integrability	24
5	Results for F_n in perturbation theory	26
5.1	Result for F_n at one loop	27
5.2	Structure of the two-loop five-point result	29
6	Connection to all-plus amplitudes in pure Yang-Mills theory	30
6.1	Matching of leading singularities and one-loop all-plus amplitude	30
6.2	Connection to two-loop all-plus amplitudes	31
7	Discussion and outlook	33
A	Leading singularities B in terms of space-time variables	35
B	Details of symmetry proofs for r_n	36
B.1	Dual-conformal algebra	36
B.2	Variable change in the conformal boost generator	36
B.3	Implications of Yangian cyclicity in space-time coordinates	37
B.4	From R-operators to the Grassmannian	38
C	Details of integrability proofs for R_n	39
C.1	R-operators, Lax-operators, and the monodromy matrix	39
C.2	From R-operators to the Grassmannian	41
C.3	Cyclicity of the Yangian	42
C.4	Lemmas	42
C.5	Yangian invariances of R_n	44

1 Introduction

Wilson lines are of fundamental importance in gauge theories. They describe many physical situations, especially in singular limits, where they provide an effective, often universal description. For example, correlation functions of Wilson loops describe the key part of infrared divergences of scattering amplitudes [1]. In planar $\mathcal{N} = 4$ super Yang-Mills (sYM), null polygonal Wilson loops even describe the full scattering amplitudes, including the finite part [2]. This duality implies many useful properties for both the scattering amplitudes and the Wilson loops. The former have an intrinsic (super)conformal symmetry, which is augmented by a (dual) superconformal symmetry of the Wilson loops [3]. Together the symmetries combine to a Yangian symmetry [4].

Since the discovery of the Wilson loop/scattering amplitude duality countless important further developments occurred. Very remarkably, information about the planar loop integrand of the scattering amplitudes is readily available: its leading singularities are described by a Grassmannian [5, 6]; it can be obtained from a loop-level version of the BCFW on-shell recursions [7], and there is a dual geometric description in terms of the Amplituhedron [8]. Moreover, loop integrands at high loop orders have been obtained from soft and collinear consistency relations [9, 10], and via a connection to correlation functions [11–13].

In parallel, integrated planar amplitudes were obtained to high loop orders via bootstrap methods [14, 15], relying in part on observations about the symbol alphabet [16], novel cluster algebraic structures [17] and analytic properties [18, 19] of the integrated answer (for reviews see [20, 21]). However, apart from initial calculations used to jump-start this program, it is not known how to go directly from the integrands to the integrated answers. Filling this gap would signify important progress in our understanding and ability of performing loop calculations, and would very likely be useful beyond $\mathcal{N} = 4$ sYM [21].

So we would like to use $\mathcal{N} = 4$ sYM as a training ground for learning how to go from the loop integrand of scattering amplitudes to the integrated answer. Indeed, the setting seems promising: both the planar integrand and the integrated answer are known to staggering loop orders. However, there are two immediate concerns that should be addressed before getting started. The first has to do with the kind of finite observables studied so far, namely ratio and remainder functions [22]. We use the word observable in the sense of a finite, gauge-invariant and scheme-independent quantity. For example, for scattering amplitudes, a suitably defined finite part can be identified as a particular contribution to a cross section, see e.g. [23]. Unfortunately, by construction, the ratio and remainder functions are non-trivial only starting from six external particles. However, in realistic gauge theories, already the four- and five-particle scattering processes are very complicated and of enormous interest. Therefore, tools for studying such processes directly are desirable. The second issue has to do with the fact that scattering amplitudes have infrared divergences. The form of the latter is known in principle (and often can be given explicitly to very high loop orders), so that only the finite part of the scattering amplitudes is new. Moreover, one expects that simplifications occur for the finite part, while intermediate, regulated results may be considerably more complicated. Unfortunately, despite knowing the form of

the divergences, it is not in general known how to set up the calculation of the finite part without needing regularization of intermediate steps (for recent related work see [24]).

In ref. [25] a new class of observables was studied that has the potential to solve these two issues. The authors considered correlation functions of Wilson loops with local operator insertions. More precisely, one takes

$$\frac{1}{\pi^2} F_n = \frac{\langle W_n \mathcal{L}(x_0) \rangle}{\langle W_n \rangle}, \quad (1.1)$$

where W_n is the n -sided null Wilson loop dual to scattering amplitudes, and \mathcal{L} is the Lagrangian, inserted at an arbitrary position x_0 . We take the Wilson loop to be in the fundamental representation of the gauge group. This quantity has a number of desirable properties. Firstly, it is easy to see that cusp divergences of the Wilson loop (dual to infrared divergences of scattering amplitudes) drop out in the ratio in eq. (1.1). Secondly, upon integrating over x_0 , one recovers (the logarithmic derivative of) the logarithm of the Wilson loop itself, and via the duality, the logarithm of the planar scattering amplitude. Therefore, F_n can be thought of as the integrand of (the logarithm of) the scattering amplitude, where all but one integrations have been carried out, in such a way as to give a finite answer. This is reminiscent of [26], where a finite web integrand function is defined for exponentiated Wilson loops, and where divergences only come from further integrations over the web integrand. Thirdly, in addition to the dual conformal symmetry of the Wilson loop, one might expect F_n to have further symmetries, thanks to the amplitude duality. This however is rather subtle, and is a central point of study in the present paper.

In view of the dual conformal symmetry, the ratio in (1.1) is a function of $3n - 11$ variables [25]. In the literature, it has been mostly studied in the case $n = 4$, where it is given by a non-trivial function of one variable, in analogy with four-particle scattering amplitudes. That function is known at weak coupling to three loops, and at leading order in strong coupling [25, 27, 28].

Quite remarkably, the Amplituhedron description of the planar loop integrand gives a natural way of discussing the logarithm of the amplitude (or the Wilson loop) [29]. It can be seen that this allows one to define the integrand for F_n directly from geometric equations, analogous to the original definition of the Amplituhedron. In this way the planar integrand for F_n is arranged into a novel form, where each term is manifestly finite in four dimensions [30]. This representation is explored further for $n = 4$ in [30].

In the present paper we study the structure of the n -particle function F_n , with a particular focus on its leading singularities and their symmetry properties. Our motivation is to inform future bootstrap approaches for this quantity, and to lay the basis for future studies of anomalous symmetries.

We profit from known results for integrands of scattering amplitudes [5, 7]. More precisely, using convenient expressions for the integrand of the (logarithm of) the one- and two-loop MHV amplitudes, we are able to determine the analytic result for F_n at tree-level and at one loop. We find that the leading singularities appearing in these expressions are governed by contour integrals over a Grassmannian $\text{Gr}(2, 2 + n)$. The Grassmannian formula was first obtained in the context of one-loop MHV amplitudes in [5, 7]. We find

it suggestive that this formula is valid to higher loops as well. Indeed, further evidence towards this conjecture is provided by an explicit two-loop calculation of F_5 [31].

The planar loop integrand of $\mathcal{N} = 4$ sYM is known to have a Yangian symmetry. However, this symmetry is somewhat formal, in the sense that it is only valid when the integrations are considered as well. For example, the leading singularities of planar amplitudes in $\mathcal{N} = 4$ sYM are Yangian invariant. In our case, however, we do not integrate over the insertion point x_0 , so that the Yangian invariance could be obscured by total derivative terms. This is not an issue for the dual conformal part of the symmetry. In our setup, it corresponds to the native symmetry of the polygonal Wilson loop, and the Lagrangian transforms covariantly under it. However, the same is not true for the conformal symmetry of amplitudes. We find that the situation becomes easier if we choose a dual conformal frame where $x_0 \rightarrow \infty$, which can be done without loss of generality. In that case, the formulas for the leading singularities simplify, and we find that they are *invariant* under conformal transformations. In other words, the symmetry is manifest, as opposed to being obscured by potential total derivative terms. We first observe this property at $n = 4, 5$, and then provide a proof that the Grassmannian formula is conformally invariant.

Indeed there is much encouraging work in the context of scattering amplitudes where symmetry properties could be proven starting from Grassmannian formulas, see e.g. [32–36]. It turns out, rather surprisingly, that the formula for the leading singularities of F_n resembles very closely a special case of a Grassmannian formula for tree-level Amplituhedron volume functions considered in [35–37].

Motivated by the additional symmetry, we further explore the structure of the higher Yangian charges, using integrability methods. We do this both in the dual conformal frame $x_0 \rightarrow \infty$, and for generic x_0 . We find that the leading singularities are annihilated by half of the Yangian generators, and identify among them the conformal symmetry. The proof uses methods of integrability to construct Yangian generators and to study their action on the Grassmannian representation. Specifically, we rely on constructions borrowed from the quantum inverse scattering method [38, 39] and integrable spin chains. Similar considerations to formulate the Yangian symmetry in QFT context has been previously applied to the tree-level scattering amplitudes in $\mathcal{N} = 4$ sYM [40–43] and ABJM theory [44], form factors of composite operators [45, 46], form factors of Wilson lines [47], kernels of QCD parton evolutions [48], Amplituhedron volume functions [36], splitting amplitudes [49], and Fishnet diagrams [50, 51].

Next, we analyze further the full expression for the observable F_n , and find that it is closely related to all-plus helicity scattering amplitudes in pure Yang-Mills theory. In the frame $x_0 \rightarrow \infty$, we write the leading singularities in a standard momentum-space notation, and make contact with conformal invariants that appeared in all-plus Yang-Mills amplitudes [52, 53]. But not only the leading singularities are the same — in fact we find that the lowest order expression for the amplitude matches precisely that of our correlator!

Motivated by the leading order results, we compute F_n for arbitrary n at the one-loop order, again finding the expected structure of conformal leading singularities, multiplied by transcendental functions, which we compute explicitly. We then compare F_n to the two-loop all-plus helicity amplitudes [54]. We find evidence that the Wilson loop with

the Lagrangian insertion at L loops agrees with the $(L + 1)$ -loop all-plus amplitude in pure Yang-Mills theory! This is at the level of leading transcendental terms, and up to scheme-dependent terms. If true in general, this relationship can be used as a prediction for yet to be computed scattering amplitudes.

The outline of this paper is as follows. In section 2, we give a lightning review of key features of the Wilson loop/scattering amplitudes duality that are relevant to this paper. Section 3 is dedicated to the Wilson loop with a Lagrangian insertion. We discuss this observable at Born level, and propose Grassmannian formulas for its leading singularities at loop level. We observe that the leading singularities have a hidden, conformal symmetry, which we prove using the Grassmannian representation. In section 4, we study higher order symmetries of the leading singularities. In section 5 we provide evidence that we correctly identified all leading singularities of the loop corrections. In particular, we present the full analytic one-loop result for F_n . In section 6, we compare F_n to the maximal-weight piece of two-loop all-plus amplitudes in pure Yang-Mills theory, and find evidence for a new duality. We discuss future research directions in section 7.

2 Recap of key features of the Wilson loop/scattering amplitudes duality

In this section we quickly review aspects of the duality between Wilson loops and scattering amplitudes that provide useful background information for the topic developed in the paper.

Unless otherwise stated, we will be discussing the planar limit, where the ‘t Hooft coupling

$$\lambda = g_{\text{YM}}^2 N_c, \tag{2.1}$$

stays fixed, while taking the rank of the gauge group N_c to infinity, and letting the Yang-Mills coupling g_{YM} go to zero. We will find it convenient to express results in terms of the combination

$$g^2 = \frac{g_{\text{YM}}^2 N_c}{16\pi^2}, \tag{2.2}$$

which is often used in the literature.

While there are indications that a version of the duality may hold for non-planar cases [55], it is certainly best understood in the planar limit [56–58]. For reviews, see [21, 59]. In its simplest form, the duality relates maximally-helicity-violating (MHV) scattering amplitudes and null polygonal Wilson loops. On the one hand, the scattering amplitudes depend on n on-shell momenta $p_i = \lambda_i \tilde{\lambda}_i$, obeying momentum conservation $\sum_{i=1}^n p_i = 0$. On the other hand, the Wilson loops are defined on a null polygonal contour, with cusps x_i . The duality identifies the kinematics according to

$$x_{i+1} - x_i = p_{i+1}, \tag{2.3}$$

where $x_{n+1} := x_1$. The statement of the duality is that

$$\langle W(x_1, \dots, x_n) \rangle A_{\text{tree}}^{\text{MHV}} \sim A_n^{\text{MHV}}, \tag{2.4}$$

where A_n^{MHV} is the color-ordered MHV amplitude, and $A_{\text{tree}}^{\text{MHV}}$ is its tree-level value. It is the well-known Parke-Taylor factor which ensures that both objects have the same helicity weights. The symbol \sim indicates that the objects are dual to each other, but not strictly equal. The reason is that the objects have different divergences: the Wilson left-hand side of eq. (2.4) has ultraviolet divergences (short-distance divergences arising near the cusp points) [3], while the amplitude has infrared divergences (i.e. soft and collinear divergences in momentum space) [60].

It turns out that there are two ways one can sharpen the duality and replace the equivalence by an equal sign. The first is to go to appropriately defined finite versions of the objects entering the duality. This idea has led to studying ratio and remainder functions [22]. The second is to consider both sides of eq. (2.4) at the level of four-dimensional loop integrands. This version will be very useful for the purpose of this paper. On one side of the duality, we have the planar L -loop integrand of MHV scattering amplitudes (with the tree-level factor removed). Thanks to its planarity, it can be expressed in terms of region variables, removing the ambiguity of loop momentum relabelling. On the other side of the duality, let us consider a derivative in the coupling,

$$g_{\text{YM}}^2 \partial_{g_{\text{YM}}^2} \langle W(x_1, \dots, x_n) \rangle \sim \int d^D y \langle W(x_1, \dots, x_n) \mathcal{L}(y) \rangle. \tag{2.5}$$

Expanding this relation in the coupling, we see that integrating over a Lagrangian insertion generates one-loop corrections to the Wilson loop. Iterating this procedure leads in a natural way to consider

$$\langle W(x_1, \dots, x_n) \mathcal{L}(y_1) \dots \mathcal{L}(y_L) \rangle, \tag{2.6}$$

evaluated at the lowest perturbative order, as the L -loop integrand of the Wilson loop.

In principle, it is not obvious that the two quantities, i.e. eq. (2.6) and the integrand of scattering amplitudes, should coincide, since loop integrands are defined up to total derivatives. Indeed, taking alternative formulations of the Lagrangian can lead to such differences. However it was noticed in refs. [12, 61] that when \mathcal{L} is taken to be the chiral Lagrangian, the duality holds at integrand level. In the following, we will define a finite observable based on the same integrands, which however involves loop integrations and therefore non-trivial transcendental functions. It will be very useful for us to rely on insights discovered in the context of the Wilson loop/scattering amplitudes duality.

The duality implies hidden symmetries for both objects. The scattering amplitudes have a conformal symmetry, which is augmented by a dual conformal symmetry of the Wilson loop. The symmetries are anomalous at loop level but have a high predictive power: they completely fix the form of four- and five-particle scattering amplitudes, and heavily constrain higher-particle amplitudes. In a supersymmetrized version of the duality, both objects have a superconformal and dual superconformal symmetry [22, 62], which combine to a Yangian symmetry [4].

Conjecturally, scattering amplitudes in $\mathcal{N} = 4$ sYM can be written in terms of transcendental functions (of transcendental weight $2L$ at L loops), multiplied by certain algebraic prefactors. The latter are computed using so-called leading singularities. Several dual

formulations for computing leading singularities exist, for example in terms of residues of certain Grassmannian formulas, and in terms of plabic graphs. This helps to make the Yangian symmetry of the leading singularities manifest. Moreover, in some cases it leads to a complete classification of what form the leading singularities can take. For example, for planar MHV amplitudes, there is only a single leading singularity, the Parke-Taylor factor. This implies that the integrated answer is a linear combination of transcendental functions, with coefficients in \mathbb{Q} , i.e. independent of the kinematics.

3 Null Wilson loop with a Lagrangian insertion

3.1 Definition, kinematics, and key properties

We define the ratio of the Wilson loop correlation function with the Lagrangian \mathcal{L} and the vacuum expectation value of the Wilson loop $W_F[\mathcal{C}] = 1/N_F \text{tr}_F P \exp(i \oint_{\mathcal{C}} dx \cdot A(x))$ in the fundamental representation,

$$\frac{1}{\pi^2} F_n(x_1, \dots, x_n; x_0) = \frac{\langle W_F[x_1, \dots, x_n] \mathcal{L}(x_0) \rangle}{\langle W_F[x_1, \dots, x_n] \rangle}, \quad (3.1)$$

where the Wilson loop contour $\mathcal{C} = [x_1, \dots, x_n]$ is an n -cusp polygon with the light-like edges, that is

$$(x_{i+1} - x_i)^2 = 0, \quad i = 1, \dots, n, \quad (3.2)$$

embedded in Minkowski space.

There are several different forms of the $\mathcal{N} = 4$ sYM Lagrangian, which all coincide modulo total derivatives and equations of motion. Here we take \mathcal{L} to be the chiral on-shell Lagrangian. An explicit expression for the latter can be found in eqs. (3.37) and (A.13) of [63]. Another choice of the Lagrangian would result in the shift of F_n by a total derivative term (in x_0). Our choice of the Lagrangian is motivated by the fact that via the correlator/amplitude duality [63–66] it exactly corresponds to the amplitude loop integrands generated by means of the loop BCFW recursion [7]. Moreover, with this choice of the operator, \mathcal{L} is a component of the stress-tensor supermultiplet.

A major motivation for considering the ratio (3.1) is that it is a finite quantity. This can be understood as follows. It is known that cusp divergences of Wilson loops exponentiate [67, 68]. This means that while at L loops, $\langle W[x_1, \dots, x_n] \rangle$ has poles up to ϵ^{-2L} in dimensional regularization with $D = 4 - 2\epsilon$, its logarithm has only double poles at most. Moreover, the exponent, $\log \langle W \rangle$, can be computed directly from certain web diagrams that are free of subdivergences (see [26] and references therein). Let us now consider the following derivative,

$$g_{\text{YM}}^2 \partial_{g_{\text{YM}}^2} \log \langle W[x_1, \dots, x_n] \rangle = \int d^D x_0 F_n(x_1, \dots, x_n; x_0). \quad (3.3)$$

This means that F_n can be considered as the *integrand* of the exponentiated Wilson loop. We already know from the discussion of webs that $\log \langle W \rangle$ is free of subdivergences. In other words the cusp divergences only arise via the final integration in x_0 (more precisely




Figure 1. Representative Feynman diagrams contributing to $F_4^{(0)}$, i.e. at order g_{YM}^2 .

from integration regions where x_0 approaches the light-like contour), but the integrand F_n is free of divergences.

We will therefore consider F_n in $D = 4$ dimensions. Note that this is not sufficient in order to use eq. (3.3) to obtain the full $\log\langle W[x_1, \dots, x_n] \rangle$ in dimensional regularization, but this is not our aim here. (Nevertheless, it was shown that this relation can be used to efficiently compute the cusp anomalous dimension, see [28].) Rather we use this to motivate the finiteness of the observable F_n that we study.

Let us consider the perturbative expansion of F_n in the coupling g_{YM} . At the lowest perturbative order, least two gluons need to be emitted from the Wilson loop contour to connect to the Lagrangian insertion (see figure 1). This means that the expansion starts at order $\mathcal{O}(g_{\text{YM}}^2)$, and we define accordingly,

$$F_n = \sum_{L \geq 0} (g^2)^{1+L} F_n^{(L)}. \tag{3.4}$$

The expected uniform transcendental weight property of $\mathcal{N} = 4$ sYM theory suggests that the terms of the perturbative expansion should be of the following form

$$F_n^{(L)} = \sum_s R_{n,s} g_{n,s}^{(L)}. \tag{3.5}$$

Here $\{R_{n,s}\}$ is a set of n -cusp leading singularities, which are rational functions of x_1, \dots, x_n and of x_0 , and where the accompanying $g_{n,s}^{(L)}$ are pure functions of transcendental weight $2L$. In particular, at the Born-level ($L = 0$) we have $g_{n,s}^{(0)} = 1$, and $F_n^{(0)}$ is a rational function.

For example, evaluating the Feynman diagrams of figure 1, one finds (see [64, 65] for a similar Lagrangian insertion calculation for a correlation function of scalar operators)

$$F_4^{(0)} = -\frac{x_{13}^2 x_{24}^2}{x_{01}^2 x_{02}^2 x_{03}^2 x_{04}^2}. \tag{3.6}$$

The light-like geometry of the contour is preserved by dual-conformal transformations. The quantum corrections to the Wilson loop render the dual-conformal symmetry anomalous. In the ratio (3.1), the anomalies of the dual-conformal symmetry cancel out, and this object transforms covariantly under the dual conformal transformations. It carries zero dual-conformal weight with respect to each of the cusp points x_1, \dots, x_n , and dual-conformal weight 4 with respect to the Lagrangian point x_0 . It is easy to see these properties for eq. (3.6), for example.

The leading singularities $R_{n,s}$ in eq. (3.5) are dual-conformal covariant, while the functions $g_{n,s}^{(L)}$ depend on dual conformal invariant cross-ratios build from x_1, \dots, x_n, x_0 . Given the kinematic setup of a null polygon, plus one additional point x_0 , it turns out that there are $3n - 11$ such dual conformal invariants [69].

Note that unlike the Wilson loop $\langle W_n \rangle$, F_n has both a parity-even and odd part, due to the Lagrangian insertion. This means that in addition to the $3n - 11$ parity-even cross ratios just mentioned, F_n depends on parity-odd Lorentz invariants. Since the square of a parity-odd variable is parity even, each parity-odd invariant can be expressed in terms of the cross-ratios, up to a sign.

For example, at five points, there is one parity-odd invariant, which we write as [13]

$$\epsilon_{123450} = -\frac{i}{6} \sum_{\sigma \in \mathcal{S}_6} (-1)^\sigma x_{\sigma_1}^2 \epsilon(x_{\sigma_2}, x_{\sigma_3}, x_{\sigma_4}, x_{\sigma_5}), \quad (3.7)$$

where $\epsilon(a, b, c, d) = \epsilon_{\mu\nu\rho\sigma} a^\mu b^\nu c^\rho d^\sigma$, $(-1)^\sigma$ stands for the signature of σ , and where the summation runs over all permutations of $\{1, 2, 3, 4, 5, 0\}$. This expression is totally antisymmetric in the points x_0, x_1, \dots, x_5 . It is dual-conformally covariant with weight -1 with respect to each of the points [13]. We remark that its square is parity even, and can be written as follows,

$$(\epsilon_{123450})^2 = -\frac{1}{24} \sum_{\sigma, \tau \in \mathcal{S}_5} (-1)^\sigma (-1)^\tau x_{0\sigma_1}^2 x_{0\tau_1}^2 x_{\sigma_2\tau_2}^2 x_{\sigma_3\tau_3}^2 x_{\sigma_4\tau_4}^2 x_{\sigma_5\tau_5}^2, \quad (3.8)$$

where the summation is over all permutations of $\{1, 2, 3, 4, 5\}$.

With this, we have all ingredients to write the full result for the Born-level five-particle case. According to [64], it is given by

$$F_5^{(0)} = -\frac{1}{2} \frac{x_{24}^2 x_{35}^2 x_{10}^2 + x_{14}^2 x_{35}^2 x_{20}^2 + x_{14}^2 x_{25}^2 x_{30}^2 + x_{13}^2 x_{25}^2 x_{40}^2 + x_{13}^2 x_{24}^2 x_{50}^2 + \epsilon_{123450}}{x_{10}^2 x_{20}^2 x_{30}^2 x_{40}^2 x_{50}^2}. \quad (3.9)$$

This expression allows us to make a number of comments:

- The expression is manifestly dual-conformally covariant, as expected.
- The expression is manifestly local, i.e. it has poles only at expected locations $x_{i0}^2 = 0$. Below we will present different representations where individual terms have spurious poles, but where a hidden conformal symmetry is manifest.
- The expression is a sum of a parity even and parity odd piece. One might think that the parity odd piece is unimportant, for several reasons: firstly, upon carrying out the x_0 integration, it vanishes, since $\log W$ is parity even; secondly, in principle it depends on the precise definition of the Lagrangian insertion. Nonetheless, we will find that it is only due to the specific combination of parity even and odd terms that F has a (hidden) conformal symmetry.
- Some readers might notice some structural similarities to all-plus amplitudes in pure Yang-Mills theory. This is indeed the case, and we will explore this connection and find hints for a duality in section 6 below.

We saw that dual conformal symmetry is manifest when using the x -coordinates. We can use dual-conformal transformations, without loss of generality, to send x_0 to infinity,

$$f_n(x_1, \dots, x_n) := \lim_{x_0 \rightarrow \infty} (x_0^2)^4 F_n(x_1, \dots, x_n; x_0). \quad (3.10)$$

This object contains the same amount of information as F_n , and dependence on x_0 can be easily restored in the given f_n . We also define the leading singularities of F_n in this frame,

$$r_n(x_1, \dots, x_n) := \lim_{x_0 \rightarrow \infty} (x_0^2)^4 R_n(x_1, \dots, x_n; x_0). \quad (3.11)$$

The frame $x_0 \rightarrow \infty$ has the advantage that it is easy to connect the kinematics to familiar momentum space terminology. This is seen as follows. In this frame, the leading singularities are Poincaré-invariant functions of n points separated by light-like distances, i.e. they are invariant under space-time shifts and Lorentz transformations. Equivalently, we can say that they are functions of n light-like momenta, cf. eq. (2.3). Since we have already fixed dual conformal symmetry, this is exactly the same kinematics as for a non-dual-conformal scattering amplitude. Indeed, we will see that the known results bear close resemblance to scattering amplitudes in non-supersymmetric Yang-Mills theory.

We extensively use Lorentz spinor notations for space-time coordinates $x_{\alpha\dot{\alpha}} := x_\mu \sigma_{\alpha\dot{\alpha}}^\mu$ and $x^{\dot{\alpha}\alpha} := x^\mu \tilde{\sigma}_\mu^{\dot{\alpha}\alpha}$, which are defined with the help of 2×2 Pauli matrices, $\sigma^\mu = (1, \vec{\sigma})$ and $\tilde{\sigma}^\mu = (1, -\vec{\sigma})$. The spinor indices are raised and lowered with the help of the antisymmetric tensors $\epsilon_{\alpha\beta}$ and $\epsilon^{\dot{\alpha}\dot{\beta}}$, such that $x^{\dot{\alpha}\alpha} = \epsilon^{\dot{\alpha}\beta} \epsilon^{\alpha\beta} x_{\beta\dot{\beta}}$ and $x_{\alpha\dot{\alpha}} = \epsilon_{\alpha\beta} \epsilon_{\dot{\alpha}\dot{\beta}} x^{\beta\dot{\beta}}$. The product of matrices x is the diagonal matrix $x_{\alpha\dot{\alpha}} x^{\dot{\alpha}\beta} = x^2 \delta_\alpha^\beta$. The difference of the space-time coordinates of two adjacent cusps is a light-like vector (2.3), which factorizes in a pair of helicity spinors $\lambda, \tilde{\lambda}$,

$$p_i^{\dot{\alpha}\alpha} = \lambda_i^\alpha \tilde{\lambda}_i^{\dot{\alpha}}. \quad (3.12)$$

We also use a short-hand notation for the Lorentz invariant $\langle a|x y|b \rangle = \lambda_a^\alpha x_{\alpha\dot{\alpha}} y^{\dot{\alpha}\beta} \lambda_{b\beta}$ and $\langle \lambda_a \lambda_b \rangle = \lambda_a^\alpha \lambda_{b\alpha}$.

We prefer to use momentum twistor [70, 71] variables instead of the space-time coordinates x_1, \dots, x_n, x_0 , since the former automatically resolve the null polygon constraints (2.3). We introduce n momentum twistors Z_1, \dots, Z_n which are points in \mathbb{P}^3 . A pair of twistors Z_i and Z_{i+1} defines a line in the twistor space \mathbb{P}^3 , which corresponds to a space-time point $x_i \sim Z_i \wedge Z_{i+1}$,

$$Z_i^I = (\lambda_i^\alpha, x_i^{\dot{\alpha}\beta} \lambda_{i\beta}), \quad I = \alpha, \dot{\alpha}. \quad (3.13)$$

The collection of n pairwise intersecting lines in the twistor space corresponds to n light-like separated space-time points x_1, \dots, x_n .

In order to describe the Lagrangian space-time coordinate x_0 , we introduce a pair of momentum twistors Z_A and Z_B specifying a line in \mathbb{P}^3 ,

$$Z_A^I = (\lambda_A^\alpha, x_0^{\dot{\alpha}\beta} \lambda_{A\beta}), \quad Z_B^I = (\lambda_B^\alpha, x_0^{\dot{\alpha}\beta} \lambda_{B\beta}). \quad (3.14)$$

The auxiliary spinors λ_A and λ_B are redundant in this formalism, since only the line $x_0 \sim Z_A \wedge Z_B$ has a geometric meaning. In the following we deal with functions of the line, which are invariant under shifts $Z_A^I \rightarrow Z_A^I + \epsilon Z_B^I$ and $Z_B^I \rightarrow Z_B^I + \epsilon Z_A^I$.

The four-bracket of momentum twistors provides a natural dual-conformal invariant. The twistor four-brackets with consecutive indices are space-time distances squared, i.e.

$$x_{ij}^2 = \frac{\langle ii+1jj+1 \rangle}{\langle ii+1 \rangle \langle jj+1 \rangle}, \quad x_{i0}^2 = \frac{\langle ABii+1 \rangle}{\langle ii+1 \rangle \langle AB \rangle}. \quad (3.15)$$

The four-brackets with generic indices, e.g. $\langle ABij \rangle$, contain both parity even and parity odd contributions. A twistor expression of the parity odd invariant (3.7) is provided in (A.3).

Because of the nonzero dual-conformal weight of F_n at the Lagrangian point x_0 , we cannot rewrite F_n and its leading singularities R_n (see eq. (3.5)) in $SU(2,2)$ invariant form solely in terms of twistor variables. In order to compensate nontrivial helicity weights, we have to introduce prefactors

$$\langle AB \rangle = \langle \lambda_A, \lambda_B \rangle = \epsilon_{\alpha\beta} \lambda_A^\alpha \lambda_B^\beta, \quad (3.16)$$

in the leading singularities. In the following, we tacitly assume that the appropriate factor $\langle AB \rangle$ is taken into account when transforming between the twistor and space-time coordinates. Slightly abusing notations, we use the same letters to denote space-time and twistor objects, e.g. for leading singularities

$$R(x_1, \dots, x_n; x_0) = \langle AB \rangle^4 R(Z_1, \dots, Z_n; Z_A, Z_B), \quad (3.17)$$

hoping that the distinction will be obvious from the context, i.e. the variables used. The momentum twistor version of the leading singularities R is invariant under local rescalings $Z_i \rightarrow tZ_i$, $i = 1, \dots, n$, and scales as $R \rightarrow t^{-4}R$ at $Z_A \rightarrow tZ_A$ or $Z_B \rightarrow tZ_B$.

As we have already mentioned, the frame $x_0 \rightarrow \infty$ is very helpful. The twistor counterpart of the dual-conformal transformation $x_0 \rightarrow \infty$ is a $SU(2,2)$ transformation which sets $Z_A \wedge Z_B$ to be the infinity bi-twistor [71]. In other words, to reach that frame we substitute $Z_A^I = (0, \mu_A^{\dot{\alpha}})$ and $Z_B^I = (0, \mu_B^{\dot{\alpha}})$ where $[\mu_A, \mu_B] = 1$, so that the only nonzero components of $Z_A \wedge Z_B$ are $\epsilon^{\dot{\alpha}\dot{\beta}}$.

3.2 n -point tree-level formula in different representations

After having seen first examples for $n = 4, 5$ (eqs. (3.6) and (3.9)), here we present the tree-level results for F_n for arbitrary number of particles. We present the formula in different ways, as we expect one representation to be more suitable for higher-loop generalization, and for making symmetries manifest.

We can benefit from the fact that the planar loop integrand for F_n at L loops can be obtained from $(L+1)$ -loop amplitude integrands. Different forms of the one-loop n -point MHV integrand are available in the literature. We do not use the original box representation [72] since the parity odd integrand terms (that integrate to zero for the amplitude, but are relevant here) are not kept there. Using instead eq. (3.10) of [73], we




Figure 2. Left: ‘magic’ pentagon representation [73]. Right: ‘kermit’ representation of one-loop MHV integrand [7].

immediately find

$$F_n^{(0)} = - \sum_{1 < i < j < n} \frac{\langle ij1n \rangle \langle AB | (i-1ii+1) \cap (j-1jj+1) \rangle}{\langle ABi-1i \rangle \langle ABii+1 \rangle \langle ABj-1j \rangle \langle ABjj+1 \rangle \langle AB1n \rangle}. \quad (3.18)$$

This is a representation in terms of pentagon integrands [73], see figure 2. Here $(abc) \cap (def)$ stands for the line defined by the intersection of the planes (abc) and (def) , given by

$$\begin{aligned} \langle xy(abc) \cap (def) \rangle &= \langle xabc \rangle \langle ydef \rangle - \langle yabc \rangle \langle xdef \rangle \\ &= \langle xyab \rangle \langle cdef \rangle + \langle xybc \rangle \langle adef \rangle + \langle xyca \rangle \langle bdef \rangle. \end{aligned} \quad (3.19)$$

Eq. (3.18) has the expected dual conformal weights, and it has only local poles. Although it is not obvious from the formula, the result does have the expected cyclic symmetry. One may verify that the $n = 4, 5$ cases agree with eqs. (3.6) and (3.9), respectively, when converting them to momentum twistors.

For our purposes, we prefer an alternative formula based on the BCFW representation of the one-loop MHV amplitude [7] (dubbed ‘kermit’ formula, see figure 2), which yields,

$$F_n^{(0)} = - \sum_{1 < i < j < n} \frac{\langle AB | (1ii+1) \cap (1jj+1) \rangle^2}{\langle AB1i \rangle \langle ABii+1 \rangle \langle ABi+1j \rangle \langle AB1j \rangle \langle ABjj+1 \rangle \langle ABj+1n \rangle}. \quad (3.20)$$

In distinction to eq. (3.18), the representation (3.20) contains spurious poles that cancel between the different terms. This is a normal feature of BCFW representations.

Looking at eq. (3.20) it is natural to define the following five-point momentum twistor expression,

$$B_{ijklm} = \frac{\langle AB(mij) \cap (jkl) \rangle^2}{\langle ABij \rangle \langle ABjk \rangle \langle ABkl \rangle \langle ABlj \rangle \langle ABjm \rangle \langle ABmi \rangle}. \quad (3.21)$$

Furthermore, for certain ‘boundary terms’ in eq. (3.20), this formula simplifies, so that it is convenient to introduce the four-point twistor function

$$B_{ijkl} = \frac{\langle ijkl \rangle^2}{\langle ABij \rangle \langle ABjk \rangle \langle ABkl \rangle \langle ABli \rangle}. \quad (3.22)$$

Let us rewrite eq. (3.20) with explicit summation ranges, and writing out the boundary terms explicitly,

$$F_n^{(0)} = \sum_{i=1}^{n-3} B_{i,i+1,n-1,n} - \sum_{i=1}^{n-4} B_{i,i+1,n-2,n-1,n} - \sum_{k=2}^{n-4} \sum_{i=1}^{n-k-3} B_{i,i+1,i+k,i+k+1,n}. \quad (3.23)$$

n	4	5	6	7	8	9	10
$\# B_{ijkl}$	1	5	15	35	70	126	210
$\# B_{ijklm}$	0	1	6	21	56	126	252
$\#$ linearly independent	1	6	20	50	105	196	336

Table 1. Counting of the linearly independent twistor functions (3.21) and (3.22) in the n -cusp case assuming the cyclic ordering of the indices. Conjecturally, these rational functions give a basis of the leading singularities R_n at any loop order.

This formula will be a useful starting point for the following, in several respects. As we will see presently, the B 's are given by a Grassmannian formula, and can be thought of as leading singularities of F_n . While we do not have a general proof that all leading singularities take the form of the B 's, we will provide evidence for this at higher loops. Moreover, the form (3.23) will allow us to discover hidden conformal properties, and a connection to all-plus amplitudes in pure Yang-Mills theory. Finally, the B terms provide a good starting point for a supersymmetric extension [74].

Let us close this subsection by returning to the four- and five-cusp examples that we wrote earlier in x notation. Using the momentum twistor language, they are now expressed as follows,

$$F_4^{(0)} = B_{1234}, \tag{3.24}$$

and

$$F_5^{(0)} = B_{1245} + B_{2345} - B_{12345}. \tag{3.25}$$

Note that individual terms in the linear combination (3.25) have spurious singularities which cancel out in the sum. In other words, $F_5^{(0)}$ contains only local poles, that is singularities when some x_{ij}^2 vanishes.

The benefits of using momentum twistors to describe the B 's becomes clear if we consider that when written in terms of the space-time variables x_{ij}^2 , their expressions look rather complicated. See appendix A for details.

It is interesting to enumerate the linearly independent functions. In the n -point case we take a collection of $\binom{n}{4}$ functions B_{ijkl} with $1 \leq i < j < k < l \leq n$, and a collection of $\binom{n}{5}$ functions B_{ijklm} with $1 \leq i < j < k < l < m \leq n$. Here we have imposed the cyclic ordering of indices which reflects the geometry of the Wilson loop contour. Both collections are linearly independent separately. Combining them together we find a number of linear relations counted in table 1 for $n \leq 10$. We observe that the number of linear-independent n -point dual-conformal invariants for the chosen cyclic ordering equals to $(n-1)(n-2)^2(n-3)/12$. We checked this counting up to $n = 15$.

3.3 Conformal symmetry of the leading singularities

We now wish to reveal the conformal symmetry of the leading singularities B_{ijkl} and B_{ijklm} given in eqs. (3.21) and (3.22). In order to do so, we consider the frame $x_0 \rightarrow \infty$, which

in twistor notation corresponds to $Z_A \wedge Z_B \rightarrow \epsilon^{\dot{\alpha}\dot{\beta}}$. This will yield an expression written explicitly in the momentum space notation, using the usual $\lambda, \tilde{\lambda}$ spinor variables. We denote

$$b = B|_{Z_A \wedge Z_B \rightarrow \epsilon^{\dot{\alpha}\dot{\beta}}}. \quad (3.26)$$

Substituting each twistor Z_i by $(\lambda_i, x_i \lambda_i)$, and evaluating the four-brackets as determinants, we arrive at

$$b_{ijkl} = \frac{(\langle ijkl \rangle)^2}{\langle ij \rangle \langle jk \rangle \langle kl \rangle \langle li \rangle}, \quad (3.27)$$

$$b_{ijklm} = \frac{(\langle ij \rangle \langle jklm \rangle - \langle jm \rangle \langle ijkl \rangle)^2}{\langle ij \rangle \langle jk \rangle \langle kl \rangle \langle kj \rangle \langle jm \rangle \langle mi \rangle}, \quad (3.28)$$

with $1 \leq i < j < k < l \leq n$ and $1 \leq i < j < k < l < m \leq n$, respectively, where

$$\begin{aligned} \langle abcd \rangle &= \langle a|x_a x_b|b \rangle \langle cd \rangle - \langle a|x_a x_c|c \rangle \langle b \rangle + \langle a|x_a x_d|d \rangle \langle bc \rangle \\ &+ \langle b|x_b x_c|c \rangle \langle ad \rangle - \langle b|x_b x_d|d \rangle \langle ac \rangle + \langle c|x_c x_d|d \rangle \langle ab \rangle. \end{aligned} \quad (3.29)$$

Recall the notation $\langle a|x_a x_b|b \rangle = \lambda_a^\alpha (x_a)_{\alpha\dot{\alpha}} (x_b)^{\dot{\alpha}\beta} \lambda_b^\beta$.

Since the functions b_{ijkl} and b_{ijklm} carry zero helicity weight with respect to each λ_i , they can be also expressed solely in terms of the coordinates x_1, \dots, x_n . The counting of linearly independent b 's is the same as that of B 's in table 1. In the following, when discussing conformal symmetry and connection with non-supersymmetric amplitudes, it will be convenient to normalize the b 's with Parke-Taylor factors, which carry nontrivial helicity weights,

$$\text{PT}_n = \frac{1}{\langle 12 \rangle \langle 23 \rangle \dots \langle n1 \rangle}. \quad (3.30)$$

Let us consider as an example $n = 5$. There are six linearly independent b 's, see table 1. After normalizing by the Parke-Taylor prefactor PT_5 (3.30), they take the following simple form in spinor helicity notation,

$$\begin{aligned} \text{PT}_5 b_{1234} &= \frac{[23]^2}{\langle 45 \rangle \langle 51 \rangle \langle 14 \rangle}, & \text{PT}_5 b_{2345} &= \frac{[34]^2}{\langle 51 \rangle \langle 12 \rangle \langle 25 \rangle}, \\ \text{PT}_5 b_{1345} &= \frac{[45]^2}{\langle 12 \rangle \langle 23 \rangle \langle 31 \rangle}, & \text{PT}_5 b_{1245} &= \frac{[51]^2}{\langle 23 \rangle \langle 34 \rangle \langle 42 \rangle}, \\ \text{PT}_5 b_{1235} &= \frac{[12]^2}{\langle 34 \rangle \langle 45 \rangle \langle 53 \rangle}, & \text{PT}_5 b_{12345} &= \frac{[13]^2}{\langle 24 \rangle \langle 45 \rangle \langle 52 \rangle}. \end{aligned} \quad (3.31)$$

We are now in a position to find something remarkable. The above expressions are conformally invariant. This can be seen as follows. The conformal boost operator in momentum space is given the following second order generator [75],

$$\mathbb{K}_{\alpha\dot{\alpha}} = \sum_{i=1}^n \frac{\partial^2}{\partial \lambda_i^\alpha \partial \tilde{\lambda}_i^{\dot{\alpha}}}. \quad (3.32)$$

The quantities in eq. (3.31) are obviously conformal (for generic kinematics), i.e. they are annihilated by the operator in eq. (3.32).

As we discussed earlier, the fact that F_n has an underlying conformal symmetry is not completely surprising, as it is a feature of the amplitudes loop integrand, but this symmetry is obvious only when integrating over x_0 , which is not the case for F_n . What we find here is that the symmetry is realized in a simple way in the dual conformal frame where $x_0 \rightarrow \infty$, which we can reach without loss of generality. In section 4, we show that the conformal symmetry persists at arbitrary number of cusps n .

We wish to emphasize that the individual summands in eq. (3.9), when multiplied by PT_5 , are *not* conformally invariant. The same is true for the individual terms of the local version of F_n given in eq. (3.18). In contrast, the conformal invariance at n points is true term by term for eqs. (3.20) and (3.23), when multiplied by PT_n and $x_0 \rightarrow \infty$.

We now discuss Grassmannian formulas for the leading singularities. The latter will allow us to prove the conformal invariance of the b 's for arbitrary n .

3.4 Two Grassmannian formulas for the leading singularities

In ref. [7], an interesting Grassmannian formula was presented for the leading singularities of n -particle one-loop MHV integrands, as a function of the external variables, and of the integration point. Indeed, the different terms in the MHV formula (3.20) above, and hence the B 's, correspond to certain residues in that Grassmannian integral. In this section, we review this formula, and derive a second form that is useful in the frame $x_0 \rightarrow \infty$.

The relevant auxiliary geometry is the Grassmannian $\text{Gr}(2, n+2)$ of 2-planes in \mathbb{C}^{n+2} . We specify a 2-plane by a pair of \mathbb{C}^{n+2} vectors subjected to $\text{GL}(2)$ transformations and assemble their coordinates in $2 \times (n+2)$ matrix

$$C = \begin{pmatrix} c_A^1 & c_B^1 & c_1^1 & \dots & c_n^1 \\ c_A^2 & c_B^2 & c_1^2 & \dots & c_n^2 \end{pmatrix}. \quad (3.33)$$

Following [7], we introduce contour integrals over the complex Grassmannian manifold $\text{Gr}(2, n+2)$,

$$R_n(Z_1, \dots, Z_n; Z_A, Z_B) = \int D^{2n}C \frac{(AB)^2}{(12)(23) \dots (n1)} \prod_{r=1}^2 \delta^4 \left(\sum_{i \in \{A, B, 1, \dots, n\}} c_i^r Z_i \right), \quad (3.34)$$

where (ab) denotes 2×2 minors of the matrix C (3.33). The notation for the integration measure is a shorthand for

$$D^{2n}C = \frac{d^{2(n+2)}C}{\text{vol}(\text{GL}(2))}. \quad (3.35)$$

Explicitly, it is given by

$$\begin{aligned} \frac{d^{2(n+2)}C}{\text{vol}(\text{GL}(2))} &= \frac{1}{(n+2)!^2} \epsilon^{i_1 \dots i_{n+2}} \epsilon^{j_1 \dots j_{n+2}} (i_{n+1} i_{n+2})(j_{n+1} j_{n+2}) \cdot \\ &\quad \times dc_{i_1}^1 \wedge \dots \wedge dc_{i_n}^1 \wedge dc_{j_1}^2 \wedge \dots \wedge dc_{j_n}^2, \end{aligned} \quad (3.36)$$

where the summation $i_1, \dots, i_{n+2}, j_1, \dots, j_{n+2}$ is over $A, B, 1, \dots, n$. The measure $D^{2n}C$ is covariant under local $GL(2)$ transformations, namely $D^{2n}(gC) = (\det(g))^{n+2}D^{2n}C$ with $g \in GL(2)$. Taking into account $GL(2)$ transformations of the minors in (3.34), one can easily see that the integration is well-defined over the Grassmannian manifold. We use the letter R_n to indicate that we expect (3.34) to give the leading singularities of F_n .

The integral (3.34) is a function of the twistor line $Z_A \wedge Z_B$, since any $GL(2)$ rotations of this pair of twistors can be compensated by a $GL(2)$ rotation of the first two columns of C (see. eq. (3.33)). Indeed, the measure $D^{2n}C$ is covariant upon global $GL(n+2)$ rotations acting on C (3.33) from the right. The compensating global $GL(2)$ transformation is embedded into $GL(n+2)$.

Let us count the dual conformal weights of the contour integral (3.34). We rescale one of the momentum twistors, $Z_i \rightarrow tZ_i$ with $i = 1, \dots, n$. The compensating rescaling of the i -th column of matrix C , i.e. $\vec{c}_i \rightarrow t^{-1}\vec{c}_i$, preserves $\vec{c}_i Z_i$ in the delta function argument in (3.34). The measure rescaling $D^{2n}C \rightarrow t^{-2}D^{2n}C$ is balanced by rescaling of the minors $(i-1i)(ii+1) \rightarrow t^{-2}(i-1i)(ii+1)$. Thus, R_n is invariant under the rescaling of the Z_i . On the other hand, rescaling one of Z_a with $a = A, B$, i.e. $Z_a \rightarrow tZ_a$, the compensating transformation $\vec{c}_a \rightarrow t^{-1}\vec{c}_a$ results in $R_n \rightarrow t^{-4}R_n$. Indeed, the nontrivial weight comes from rescaling of the measure and $(AB)^2 \rightarrow t^{-2}(AB)^2$. This counting agrees with the discussion of the dual conformal weights of the leading singularities mentioned after eq. (3.17).

The dual-conformal symmetry of R_n , namely covariance upon $GL(4)$ transformations of the momentum twistors, is also obvious in the contour integral representation (3.34). In order to evaluate the contour integral, we can choose the coordinate patch

$$C \xrightarrow{GL(2)} \begin{pmatrix} 1 & 0 & c_1^1 & \dots & c_n^1 \\ 0 & 1 & c_1^2 & \dots & c_n^2 \end{pmatrix}, \tag{3.37}$$

and then $D^{2n}C \rightarrow \prod_{r=1,2} \prod_{i=1}^n dc_i^r$. The bosonic delta functions freeze 8 integrations, and the remaining $(2n-8)$ complex integrations are done by taking residues.

We find it easier to study the residues in a different version of this formula that we derive presently. Indeed, without loss of generality, let us study the contour integral representation (3.34) in the dual conformal frame $x_0 \rightarrow \infty$. In twistor variables this amounts to

$$r_n(Z_1, \dots, Z_n) = R_n(Z_1, \dots, Z_n; Z_A, Z_B)|_{Z_A \wedge Z_B \rightarrow \epsilon^{\dot{\alpha}\dot{\beta}}} \tag{3.38}$$

where we use notations $Z_i = (\lambda_i^\alpha, \mu_i^{\dot{\alpha}})$ for the twistor components split according to the choice of the infinity bi-twistor. Making this substitution, we obtain

$$r_n(Z_1, \dots, Z_n) = \int D^{2n}C \frac{(AB)^2}{(12)(23)\dots(n1)} \prod_{r=1}^2 \delta^2 \left(\sum_{i=1}^n c_i^r \lambda_i \right) \times \delta \left(c_A^r + \sum_{i=1}^n c_i^r [\mu_i, \mu_B] \right) \delta \left(c_B^r + \sum_{i=1}^n c_i^r [\mu_A, \mu_i] \right), \tag{3.39}$$

where we can choose a patch on the Grassmannian and integrate out c_A^r, c_B^r with the help of the delta-constraints. The remaining integrations are over $(2n-4)$ -dimensional

Grassmannian manifold $\text{Gr}(2, n)$, with the measure $D^{2(n-2)}C$, see eq. (3.36). In other words, doing the dual-conformal transformation $x_0 \rightarrow \infty$ we contract the auxiliary geometry,

$$\text{Gr}(2, n + 2) \rightarrow \text{Gr}(2, n). \tag{3.40}$$

Finally, we arrive at the contour integral representation for the leading singularities in the frame $x_0 \rightarrow \infty$,

$$r_n(Z_1, \dots, Z_n) = \int D^{2(n-2)}C \frac{\left(\sum_{i < j} (ij)[\mu_i \mu_j]\right)^2}{(12)(23) \dots (n1)} \prod_{r=1,2} \delta^2\left(\sum_{i=1}^n c_i^r \lambda_i\right). \tag{3.41}$$

The substitution $\mu_i^{\dot{\alpha}} = x_i^{\dot{\alpha}\alpha} \lambda_{i\alpha}$ converts the previous expression to space-time coordinates,

$$[\mu_i \mu_j] = \langle i | x_i x_j | j \rangle. \tag{3.42}$$

It may seem that the resulting expression is not invariant under space-time shifts. In fact, the integral in (3.41) is invariant under the Poincaré part of the dual conformal algebra (see eqs. (B.2) and (B.3)). Using the delta-function constraints from (3.41) we achieve the translation invariance

$$\sum_{i < j} (ij) \langle i | x_i x_j | j \rangle \rightarrow \sum_{i < j} (ij) \langle i | x_{ik} x_{kj} | j \rangle, \tag{3.43}$$

with any k . However, since we have chosen the frame $x_0 \rightarrow \infty$, the dual-conformal symmetry of (3.41) is broken.

The delta-function constraints in (3.41) freeze four integrations, and the remaining $2n - 8$ integrations are done by the residue calculus. The residues are linear combinations of b 's (3.26), and counting of the linearly independent residues r_n and the functions b_{ijkl} and b_{ijklm} is the same as in table 1.

Certain choices of contours in eqs. (3.34) and (3.41) give the B 's and b 's, respectively. This follows from the connection to one-loop MHV scattering amplitudes, which correspond to F_n at Born level. We find it plausible that at loop level these formulas govern the leading singularities of F_n . In the following we will present evidence for this hypothesis.

The contour integral representation (3.41) of the b 's will be instrumental in proving their conformal symmetry and higher charge Yangian symmetries. Moreover, the contour integral admits a BCFW-bridge-like decomposition in a sequence of twistor shifts, thereby making a connection with the cell decomposition of the Grassmannian and plabic graphs [6]. Another remarkable feature of this concise expression for the conformal invariants is that for a particular integration contour it gives a non-supersymmetric loop amplitude, as will be discussed in section 6.

4 Higher order symmetries and integrability

In this section, we study higher order symmetries of the Grassmannian formulas we proposed for r_n and R_n , respectively. Recall that due to the close connection to scattering amplitudes,

it is known that integrating over $Z_A Z_B$ (for example, on a cycle giving leading singularities of the amplitudes), one obtains Yangian invariants. This means that R_n (and hence also r_n , which corresponds to a particular dual conformal frame choice of R_n), should ‘know about’ the Yangian symmetry: upon Yangian transformations acting on the external twistors, it should turn into a total derivative w.r.t. $Z_A Z_B$. On the one hand, it is unclear to us how useful this assertion is. On the other hand, in section 3.3 we have uncovered hints for a conformal invariance of r_n . This motivates us to look for a proof of this symmetry, and to look for higher-order constraints on both R_n and r_n . We do so starting from the Grassmannian formulas.

We provide a proof of the conformal symmetry of the Grassmannian formula (3.41) for r_n , and hence for the four-point and five-point functions b given in eqs. (3.27) and (3.28). We also identify additional higher-order symmetries.

Furthermore, we carry out a similar analysis in the general frame, i.e. for arbitrary x_0 . In this way, we can formulate analogues of the conformal symmetry relation, as well as of the higher-order equations found, directly for the rational functions R_n and B . We find that this is closely related to differential equations found in ref. [36] in the context of the tree-level Amplituhedron.

4.1 Proof of conformal and higher-order symmetries of the leading singularities r_n

4.1.1 Yangian generators

The dual-conformal algebra generators have a very simple representation when written in momentum twistors $Z^I = (\lambda^\alpha, \mu^{\dot{\alpha}})$, namely¹

$$\mathfrak{J}^I{}_J = Z^I \frac{\partial}{\partial Z^J} - \frac{\delta^I_J}{4} Z^L \frac{\partial}{\partial Z^L}, \tag{4.1}$$

where $I = \alpha, \dot{\alpha}$ and $J = \beta, \dot{\beta}$. Indeed, in this formula the 4×4 matrix $\mathfrak{J}^I{}_J$ comprises the dual conformal generators D, P, M, \bar{M}, K (see appendix B.1),

$$\mathfrak{J}^I{}_J = \begin{pmatrix} M^\alpha{}_\beta - \frac{\delta^\alpha_\beta}{2} D & P^\alpha{}_{\dot{\beta}} \\ K^{\dot{\alpha}}{}_\beta & \bar{M}^{\dot{\alpha}}{}_{\dot{\beta}} + \frac{\delta^{\dot{\alpha}}_{\dot{\beta}}}{2} D \end{pmatrix}. \tag{4.2}$$

In the following we wish to act on the external momentum twistors Z_i , with $i = 1, \dots, n$. We denote by \mathfrak{J}_i the local generator acting on the i -th coordinate, and we represent the dual-conformal generators as sums of the local ones

$$\begin{aligned} \mathfrak{J}^I{}_J &= \sum_{i=1}^n (\mathfrak{J}_i)^I{}_J, \\ (\mathfrak{J}_i)^I{}_J &= Z_i^I \frac{\partial}{\partial Z_i^J} - \frac{\delta^I_J}{4} Z_i^L \frac{\partial}{\partial Z_i^L}. \end{aligned} \tag{4.3}$$

¹We could have chosen to supplement $su(2,2) \approx sl(4)$ with the central element $H = Z^L \frac{\partial}{\partial Z^L}$ and consider $gl(4)$ generators $\mathfrak{J}^I{}_J = Z^I \frac{\partial}{\partial Z^J}$. Indeed, each local instance of the central element annihilates the leading singularities $H_i r_n = 0$, $i = 1, \dots, n$, and commutes with other operators.

Let us start by discussing the obvious symmetries. The leading singularities r_n given in eq. (3.41) are Poincaré invariant,

$$P_{\alpha\dot{\beta}} r_n = M_{\alpha\beta} r_n = \bar{M}_{\dot{\alpha}\dot{\beta}} r_n = 0, \quad D r_n = 2 r_n, \quad (4.4)$$

but due to our frame choice $x_0 \rightarrow \infty$ the dual conformal invariance is broken,

$$K_{\dot{\alpha}\beta} r_n \neq 0. \quad (4.5)$$

In order to study higher symmetries of the leading singularities r_n , we introduce an infinite-dimensional Yangian algebra which extends the dual-conformal algebra. The Yangian of $su(2, 2)$ is a Hopf algebra with an infinite tower of generators grouped into levels. The zeroth level generators $\mathfrak{J}^{(0)} := \mathfrak{J}$ are those of $su(2, 2)$, and the level-one generators $\widehat{\mathfrak{J}}^{(1)} := \widehat{\mathfrak{J}}$ transform in the adjoint representation of $su(2, 2)$,

$$[\widehat{\mathfrak{J}}^I{}_J, \widehat{\mathfrak{J}}^K{}_L] = \delta_J^K \widehat{\mathfrak{J}}^I{}_L - \delta_L^I \widehat{\mathfrak{J}}^K{}_J. \quad (4.6)$$

Higher-level generators are defined as repeated commutators of the level-one generators, subjected to the Serre relations [76].

The coproduct structure of the Hopf algebra suggests that we represent the level-one generators $\widehat{\mathfrak{J}}$ of the Yangian as sums of bi-local terms formed from the local generators \mathfrak{J}_i ,

$$\widehat{\mathfrak{J}}^I{}_J = \sum_{1 \leq i < j \leq n} (\mathfrak{J}_i)^I{}_K (\mathfrak{J}_j)^K{}_J. \quad (4.7)$$

4.1.2 Spin-chain picture of the Yangian symmetry

As mentioned in the Introduction, here we use constructions borrowed from the quantum inverse scattering method and integrable spin chains to formulate the Yangian symmetry.

Let us define the *Lax operator* $L(u)$, which is a 4×4 matrix with first order differential operator entries (dual-conformal generators $\mathfrak{J}^I{}_J$ (4.1)),

$$[L(u)]^I{}_J = u \delta_J^I + Z^I \frac{\partial}{\partial Z^J} - \frac{\delta_J^I}{4} Z^K \frac{\partial}{\partial Z^K}, \quad (4.8)$$

and $u \in \mathbb{C}$ is called the spectral parameter. Let us denote by \mathcal{H}_i the space of homogeneous functions of Z_i^I with homogeneity degree zero, i.e. functions which are invariant under rescaling $G(Z_i^I) = G(tZ_i^I)$. Then $L_i(u)$ acts on $\mathcal{H}_i \otimes \mathbb{C}^4$.

We associate a quantum spin chain to the leading singularities r_n , such that each i -th site of the spin chain carrying representation space \mathcal{H}_i corresponds to the i -th edge of the Wilson loop. We assign a Lax operator to each site of the spin chain and we form the matrix product of n Lax operators, which is called the *monodromy matrix*,

$$[T(u)]^I{}_J := [L_1(u)]^I{}_{K_1} [L_2(u)]^{K_1}{}_{K_2} \dots [L_n(u)]^{K_{n-1}}{}_J. \quad (4.9)$$

The monodromy matrix $T(u)$ acts in the space

$$\mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \dots \otimes \mathcal{H}_n \otimes \mathbb{C}^4. \quad (4.10)$$

Expanding the monodromy matrix in the spectral parameter u we recover the Yangian generators in the evaluation representation [77]: the dual-conformal generators $\mathfrak{J} := \mathfrak{J}^{(0)}$ (4.1) at the zeroth level, first level generators $\widehat{\mathfrak{J}} := \mathfrak{J}^{(1)}$ (4.7) given by the sum of bi-local terms, as well as higher level generators $\mathfrak{J}^{(2)}, \dots, \mathfrak{J}^{(n-1)}$,

$$[\mathbb{T}(u)]^I{}_J = u^n \delta^I{}_J + u^{n-1} [\mathfrak{J}^{(0)}]^I{}_J + u^{n-2} [\mathfrak{J}^{(1)}]^I{}_J + \sum_{k=2}^{n-1} u^{n-k-1} [\mathfrak{J}^{(k)}]^I{}_J, \quad (4.11)$$

with k -th level generator being the sum of ordered products of $k + 1$ local generators (4.3),

$$[\mathfrak{J}^{(k)}]^I{}_J = \sum_{1 \leq i_1 < i_2 < \dots < i_{k+1} \leq n} (\mathfrak{J}_{i_1})^I{}_{L_1} (\mathfrak{J}_{i_2})^{L_1}{}_{L_2} \dots (\mathfrak{J}_{i_{k+1}})^{L_k}{}_J. \quad (4.12)$$

In the following we denote $\mathfrak{J}^{(0)}$ and $\mathfrak{J}^{(1)}$ as \mathfrak{J} and $\widehat{\mathfrak{J}}$, respectively, to simplify notations. In eq. (4.12) a specific choice of origin of the sum was made. Below we also discuss the cyclically related choices.

Let us note that in the evaluation representation (4.9) of the Yangian induced by (4.8) the infinite tower of Yangian generators is truncated at the level n . Studying the Yangian symmetry, we work with the monodromy matrix (4.9), which encompasses all Yangian generators in the chosen representation. Instead of acting with individual Yangian generators $\mathfrak{J}^{(k)}$ on r_n , we would like to calculate the action of the monodromy matrix $\mathbb{T}(u)$ on r_n . Moreover, we would like to reduce the evaluation of $\mathbb{T}(u) r_n$ to manipulations with local objects, those which act nontrivially on a single or a pair of spin chain sites. By definition (4.9), the monodromy matrix is a product of local blocks. Therefore, we will also need a decomposition of r_n into local blocks.

We define an R-operator which acts on a pair of the spin chain sites $\mathcal{H}_i \otimes \mathcal{H}_j$ and transforms a test function G as follows,

$$[\mathbb{R}_{ij} G](Z_i, Z_j) := \int \frac{dt}{t} G(Z_i + tZ_j, Z_j). \quad (4.13)$$

We imply that the integration contour lies in the complex plane and it is closed, but we do not fix it. Let us note that the R-operator preserves homogeneity, hence it does not map out of $\mathcal{H}_i \otimes \mathcal{H}_j$. The R-operator (4.13) commutes with the matrix product of the pair of Lax operators (4.8),

$$\mathbb{R}_{ij} L_i(u) L_j(u) = L_i(u) L_j(u) \mathbb{R}_{ij}, \quad (4.14)$$

$$\mathbb{R}_{ji} L_i(u) L_j(u) = L_i(u) L_j(u) \mathbb{R}_{ji}, \quad (4.15)$$

as can be easily verified. Let us stress that the R-operator acts trivially in the matrix space of the previous equations. Thus, the R-operator acting in adjacent spin chain sites commutes with the monodromy matrix (4.9),

$$\mathbb{R}_{i+1} \mathbb{T}(u) = \mathbb{T}(u) \mathbb{R}_{i+1}, \quad \mathbb{R}_{i+1} \mathbb{T}(u) = \mathbb{T}(u) \mathbb{R}_{i+1}, \quad i = 1, \dots, n-1. \quad (4.16)$$

Now we use a corollary proven in section B.4. There we show that the contour integral (3.41) over the Grassmannian $\text{Gr}(2, n)$ can be rewritten as a product of $2(n-2)$

R-operators acting on a function $|\Omega_n\rangle$,

$$r_n(Z_1, \dots, Z_n) = R_{32} R_{43} \dots R_{n n-1} R_{n-2 n-1} \dots R_{23} R_{12} |\Omega_n\rangle. \quad (4.17)$$

Each of the R-operators from the product in (4.17) introduces one-dimensional integration. We call $|\Omega_n\rangle$ a pseudo-vacuum state of the n -site spin chain,

$$|\Omega_n\rangle := \delta^2(\lambda_1) \delta^2(\lambda_n) [\mu_1 \mu_n]^2. \quad (4.18)$$

One can easily see that the distribution $|\Omega_n\rangle$ is of homogeneity degree zero, i.e. it is an element of the space $\mathcal{H}_1 \otimes \dots \otimes \mathcal{H}_n$.

The employed notations are very suggestive. The pseudo-vacuum $|\Omega_n\rangle$ of the spin chain represents a zero-dimensional cell of the Grassmannian $\text{Gr}(2, n)$. Each ‘excitation’ over the pseudo-vacuum introduced by the R-operator is a BCFW bridge [6], which maps to a Grassmannian cell of higher dimension. There are $2(n-2)$ ‘excitations’ in (4.17) that correspond to the top cell of the Grassmannian.

Since the monodromy matrix $T(u)$ commutes with individual R-operators (4.16), then it also commutes with their products in (4.17). Thus, in order to calculate $T(u) r_n$ it remains to find how the 4×4 monodromy matrix (4.9) acts on the pseudo-vacuum $|\Omega_n\rangle$ (4.18). We find that it takes a block-triangular form

$$T(u) |\Omega_n\rangle = u^{n-2} \begin{pmatrix} u^2 - 2u + 1 & 0 \\ 0 & u^2 + 2u - 1 \end{pmatrix} |\Omega_n\rangle + u^{n-2} \begin{pmatrix} 0 & 0 \\ f + ug & h \end{pmatrix}, \quad (4.19)$$

where dependence of the r.h.s. on the spectral parameter u is explicit; f, g, h are 2×2 matrices which depend on twistor variables Z_1 and Z_n , and matrix h is traceless, $\text{tr}(h) = 0$. The explicit expressions for f, g, h will be irrelevant in what follows. Eq. (4.19) expression is easy to work out, since by acting on the pseudo-vacuum successively with each of the constituent Lax operators $L(u)$ given in eq. (4.8), we obtain either diagonal or block-triangular matrices.

Finally, eqs. (4.19) and (4.17) enable us to find how the monodromy matrix acts on the leading singularities r_n , namely

$$T(u) r_n = u^{n-2} \begin{pmatrix} u^2 - 2u + 1 & 0 \\ 0 & u^2 + 2u - 1 \end{pmatrix} r_n + u^{n-2} \begin{pmatrix} 0 & 0 \\ \mathcal{O}(u) & \mathcal{O}(u^0) \end{pmatrix}, \quad (4.20)$$

where the last matrix term of the r.h.s. is traceless.

These equations embody our statement on the partial Yangian symmetry of r_n . They contain the complete collection of the Yangian invariances satisfied by r_n with Yangian generators of the form (4.12).

Expanding eq. (4.20) in the spectral parameter u according to (4.11), we find how the Yangian generators of all levels (4.12) act on r_n . At order u^{n-1} in (4.20), we recover Poincaré invariance (4.4) with $\mathfrak{J} = \mathfrak{J}^{(0)}$,

$$\mathfrak{J}^\alpha{}_\beta r_n = -2\delta^\alpha_\beta r_n, \quad \mathfrak{J}^\alpha{}_{\dot{\beta}} r_n = 0, \quad \mathfrak{J}^{\dot{\alpha}}{}_\beta r_n = 2\delta^{\dot{\alpha}}_\beta r_n. \quad (4.21)$$

At order u^{n-2} , we find that a half of the level-one generators $\widehat{\mathfrak{J}}^{(1)} = \widehat{\mathfrak{J}}$ provide invariances,

$$\widehat{\mathfrak{J}}^\alpha{}_\beta r_n = \delta^\alpha_\beta r_n, \quad \widehat{\mathfrak{J}}^\alpha{}_{\dot{\beta}} r_n = 0, \quad \widehat{\mathfrak{J}}^{\dot{\alpha}}{}_\alpha r_n = -2r_n. \quad (4.22)$$

However, the level one-generators $\widehat{\mathfrak{J}}^{\alpha}_{\beta}$ and the traceless part of $\widehat{\mathfrak{J}}^{\alpha}_{\beta}$ are not symmetries of r_n . In other words, this part of the Yangian symmetry is broken.

We observe that there are no terms u^{n-1-k} with $k = 2, \dots, n-1$ on the r.h.s. of (4.20). Consequently, all Yangian generators at the level two and higher provide invariance relations,

$$\left(\widehat{\mathfrak{J}}^{(k)}\right)^I{}_J r_n = 0, \quad k = 2, \dots, n-1. \quad (4.23)$$

In a similar fashion, we can prove additional relations. Let us introduce the second order differential operators

$$D_I^{(i)} := \frac{\partial}{\partial Z_i^J} \left(\sum_{j=1}^n Z_j^J \frac{\partial}{\partial Z_j^I} \right) - 2 \frac{\partial}{\partial Z_i^I}, \quad i = 1, \dots, n. \quad (4.24)$$

They will play the role of the monodromy matrix from the previous discussion. Indeed, we can check by explicit calculation that the operators $D_I^{(i)}$ commute with any R-operator (4.13) and annihilate the pseudo-vacuum $|\Omega_n\rangle$ (4.18),

$$\begin{aligned} D_I^{(i)} |\Omega_n\rangle &= 0, \\ [\mathbf{R}_{kl}, D_I^{(i)}] &= 0, \quad i, k, l = 1, \dots, n. \end{aligned} \quad (4.25)$$

Thus, we can pull $D_I^{(i)}$ through the product of R-operators in representation (4.17) and find that it annihilates the leading singularities r_n ,

$$D_I^{(i)} r_n = 0, \quad i = 1, \dots, n. \quad (4.26)$$

These relations can also be formulated in terms of the dual-conformal algebra generators (4.3) as follows,

$$\left(\widehat{\mathfrak{J}}_i\right)^K{}_J \widehat{\mathfrak{J}}^J{}_I r_n = 2 \left(\widehat{\mathfrak{J}}_i\right)^K{}_I r_n, \quad i = 1, \dots, n. \quad (4.27)$$

In appendix B.3, we rewrite eq. (4.27) in terms of the dual-conformal generators acting on the space-time variables. This enable us to see in more details how the frame choice $x_0 \rightarrow \infty$ for r_n breaks the dual-conformal symmetry.

Another way to interpret eqs. (4.26) and (4.27) is to consider the cyclic shift transformations of the Yangian. The ordering of the Wilson loop edges is irrelevant for the level-zero generators $\widehat{\mathfrak{J}}^{(0)}$. However, the level-one and higher-level generators of the Yangian (4.12) employ the cyclic ordering with the specific choice of the sum origin. At the same time, the contour integral representing the leading singularities r_n is invariant under cyclic shifts $Z_i \rightarrow Z_{i+1}$, see (3.41). Thus, we can formulate the Yangian symmetries of r_n using a set of cyclically shifted generators. Then we sum in (4.12) over cyclically shifted labels. We provide the transformation properties of the Yangian generators upon the cyclic shift in appendix C.3. Let us consider the simplest nontrivial cyclic-shift transformation. Indeed, along with the level-one generator $\widehat{\mathfrak{J}}^{(1)} = \widehat{\mathfrak{J}}$ (4.12) we introduce the level-one generator $\widehat{\mathfrak{J}}'$ which employs the ordering $n, 1, 2, \dots, n-1$. The level-one Yangian invariances from (4.22)

hold with the cyclically shifted generator $\widehat{\mathfrak{J}}^i$ as well. For example, the following difference of the generators annihilates r_n ,

$$\left[\widehat{\mathfrak{J}}^\alpha{}_\beta - \widehat{\mathfrak{J}}^{\prime\alpha}{}_\beta \right] r_n = 0. \quad (4.28)$$

Evaluating the difference of generators on the space of homogeneous functions of degree zero, i.e. $(\mathfrak{J}_n)^C{}_C = 0$,

$$\widehat{\mathfrak{J}}^A{}_B - \widehat{\mathfrak{J}}^{\prime A}{}_B = 4(\mathfrak{J}_n)^A{}_B + \mathfrak{J}^A{}_C (\mathfrak{J}_n)^C{}_B - (\mathfrak{J}_n)^A{}_C \mathfrak{J}^C{}_B, \quad (4.29)$$

and choosing the indices A, B as in (4.28), we conclude that (4.28) is equivalent to (4.26) with $i = n$. Choosing other cyclic shifts we can reproduce all n relations (4.26). Let us also note that the cyclic consistency of the remaining Yangian invariances (4.22) does not provide new relations.

Despite of the fact that a half of the level-one Yangian generators $\mathfrak{J}^{(1)} = \widehat{\mathfrak{J}}$ is broken, we can formulate several invariance relations for r_n involving the broken generators and the dual-conformal generators, e.g.

$$\mathfrak{J}^\alpha{}_\gamma \widehat{\mathfrak{J}}^{\dot{\gamma}}{}_\beta r_n = 0, \quad \mathfrak{J}^{\dot{\alpha}}{}_\gamma \widehat{\mathfrak{J}}^{\dot{\gamma}}{}_\beta r_n = 2\delta_{\dot{\beta}}^{\dot{\alpha}} r_n + 4\widehat{\mathfrak{J}}^{\dot{\alpha}}{}_\beta r_n, \quad (4.30)$$

$$\mathfrak{J}^\alpha{}_\gamma \widehat{\mathfrak{J}}^{\dot{\gamma}}{}_\beta r_n = 4\delta_{\dot{\beta}}^\alpha r_n, \quad \mathfrak{J}^{\dot{\alpha}}{}_\gamma \widehat{\mathfrak{J}}^{\dot{\gamma}}{}_\beta r_n = 4\widehat{\mathfrak{J}}^{\dot{\alpha}}{}_\beta r_n. \quad (4.31)$$

In other words, even though the Yangian variations $\widehat{\mathfrak{J}}^{\dot{\alpha}}{}_\beta$ and $\widehat{\mathfrak{J}}^{\dot{\alpha}}{}_{\dot{\beta}}$ of r_n are not covariant, contracting the variations with the dual-conformal generators we find symmetry relations for r_n . We deduce them from the commutation relations among \mathfrak{J} and $\widehat{\mathfrak{J}}$ (4.6) and the invariance relations (4.22). Relations (4.30) and (4.31) are closely related to differential equations found in [36], as will be discussed in section 4.3.

4.1.3 Summary and discussion of symmetry relations

Using integrability methods, we have proven the Yangian invariance relations (4.22), (4.23) as well as the bonus relations (4.27), for the Grassmannian formula (3.41) for r_n . Let us now discuss these results.

We see that eqs. (4.22) involve half of the level-one Yangian generators. This may be due to, at least in part, to our choice of fixing the dual conformal frame $x_0 \rightarrow \infty$.

Among the level-one Yangian invariances in eq. (4.22), the one with $\widehat{\mathfrak{J}}^{\dot{\alpha}}{}_\beta$ has a transparent meaning. It is a conformal invariance in momentum space. In order to see this, let us make contact with the conformal boost generator $\mathbb{K}_{\alpha\dot{\beta}}$ given in eq. (3.32), which is written in terms of the spinor-helicity variables $(\lambda, \tilde{\lambda})$, see appendix B.2 and [32]. There it is shown that

$$\widehat{\mathfrak{J}}^{\dot{\alpha}}{}_\beta r_n(Z_1, \dots, Z_n) = -\langle 12 \rangle \langle 23 \rangle \dots \langle n1 \rangle \mathbb{K}^\alpha{}_{\dot{\beta}} \frac{r_n(p_1, \dots, p_n)}{\langle 12 \rangle \langle 23 \rangle \dots \langle n1 \rangle}. \quad (4.32)$$

In view of eq. (4.21) this implies

$$\mathbb{K}^\alpha{}_{\dot{\beta}} \frac{r_n(p_1, \dots, p_n)}{\langle 12 \rangle \langle 23 \rangle \dots \langle n1 \rangle} = 0. \quad (4.33)$$

Therefore we have proven the conformal invariance of the leading singularities r_n . This is one of the main results of this section.

4.2 Formulation of higher-order symmetries of r_n in momentum space

Above we constructed higher charge symmetries (4.12) of r_n out of the local dual-conformal generators. Then we identified the conformal boost symmetry of r_n (4.33) among the level-one Yangian generators. We could wonder whether it is possible to construct higher charge symmetries out of the conformal generators, which act on the momentum variables. In this section we briefly comment on this possibility.

Indeed, let us assemble the conformal algebra generators written in spinor helicity variables [75] in the 4×4 matrix

$$\mathbb{J}^I_J = \begin{pmatrix} \lambda^\alpha \frac{\partial}{\partial \lambda^\beta} & -\lambda^\alpha \tilde{\lambda}^{\dot{\beta}} \\ \frac{\partial^2}{\partial \lambda^{\dot{\alpha}} \partial \lambda^\beta} & -\frac{\partial}{\partial \lambda^{\dot{\alpha}}} \tilde{\lambda}^{\dot{\beta}} \end{pmatrix} = \begin{pmatrix} \lambda^\alpha \\ \frac{\partial}{\partial \lambda^{\dot{\alpha}}} \end{pmatrix} \otimes \begin{pmatrix} \frac{\partial}{\partial \lambda^\beta} & -\tilde{\lambda}^{\dot{\beta}} \end{pmatrix} \quad (4.34)$$

where $I = (\alpha, \dot{\alpha})$ and $J = (\beta, \dot{\beta})$, and for the sake of brevity we omitted index $i = 1, 2, \dots, n$ which labels the particles. Summing local generators \mathbb{J}_i over i , we obtain the conformal symmetries of r_n . In particular $I = \dot{\alpha}$ and $J = \beta$ corresponds to the conformal boost (3.32),

$$\mathbb{J}^{(0)}_{\dot{\alpha}\beta} := \sum_{i=1}^n (\mathbb{J}_i)_{\dot{\alpha}\beta} = \mathbb{K}_{\dot{\alpha}\beta}. \quad (4.35)$$

In analogy with (4.12), we define higher charge non-local generators out of the local conformal generators (4.34). The level- k generators are the following sums of the ordered products of (4.34),

$$\left[\mathbb{J}^{(k)} \right]^I_J := \sum_{1 \leq i_1 < i_2 < \dots < i_{k+1} \leq n} (\mathbb{J}_{i_1})^I_{L_1} (\mathbb{J}_{i_2})^{L_1}_{L_2} \dots (\mathbb{J}_{i_{k+1}})^{L_k}_{J}. \quad (4.36)$$

It would be interesting to systematically study the properties of r_n under these generators, using methods similar to the last subsection. For the sake of this paper however, we limit ourselves to ‘experiment’ with the differential operators (4.36), and find, conjecturally, a set of equations that the r_n satisfy.

Let us consider $I = \dot{\alpha}$ and $J = \beta$ in (4.36), which is given explicitly by the following differential operator in terms of helicity spinors,

$$\begin{aligned} \mathbb{J}^{(k)}_{\dot{\alpha}\beta} &= \sum_{1 \leq i_1 < i_2 < \dots < i_{k+1} \leq n} \frac{\partial}{\partial \tilde{\lambda}_{i_1}^{\dot{\alpha}}} Q_{i_1 i_2} \dots Q_{i_k i_{k+1}} \frac{\partial}{\partial \lambda_{i_{k+1}}^\beta}, \\ Q_{ij} &:= \lambda_j^\gamma \frac{\partial}{\partial \lambda_i^\gamma} - \tilde{\lambda}_i^{\dot{\gamma}} \frac{\partial}{\partial \tilde{\lambda}_j^{\dot{\gamma}}}. \end{aligned} \quad (4.37)$$

These are level- k counterparts of the conformal boost generator. We conjecture that the following equations hold,

$$\mathbb{J}^{(k)}_{\dot{\alpha}\beta} \frac{r_n(p_1, \dots, p_n)}{\langle 12 \rangle \langle 23 \rangle \dots \langle n1 \rangle} \Bigg|_{\substack{\tilde{\lambda}_1 \rightarrow \frac{1}{\langle n1 \rangle} (p_2 + \dots + p_{n-1}) | n \\ \tilde{\lambda}_n \rightarrow \frac{1}{\langle 1n \rangle} (p_2 + \dots + p_{n-1}) | 1}} = 0, \quad k = 0, 1, \dots, n. \quad (4.38)$$

In this formula it is understood that $\tilde{\lambda}_1$ and $\tilde{\lambda}_n$ are eliminated with the help of momentum conservation prior to differentiations (this choice is related to the ordering of local generators in (4.36)).

In order to check eq. (4.38), we choose r_n to be the four-point b_{ijkl} (3.27) and five-point b_{ijklm} (3.28) invariants written in spinor helicity variables $(\lambda, \tilde{\lambda})$ with $\tilde{\lambda}_1$ and $\tilde{\lambda}_n$ eliminated as specified above. Thus, the four-point and five-point invariants nontrivially depend on all n momenta. Following this approach, we checked eq. (4.38) up to $n = 15$.

It would be interesting to find a proof of eq. (4.38) for any n which employs the integrability constructions above. As mentioned before, in order to achieve this goal one would need a momentum space analogue of the contour integral representation (3.41), e.g. similar to the formulas in [5] for scattering amplitudes.

4.3 Higher-order symmetries of R_n and integrability

In the previous subsections we have established higher symmetries of the leading singularities r_n in the dual conformal frame $x_0 \rightarrow \infty$. In this and the next subsections we perform a similar analysis directly for R_n .

Before doing so, let us comment on a rather surprising connection. It turns out that the formula for the leading singularities R_n resembles very closely a special case of a Grassmannian formula for tree-level Amplituhedron volume functions considered in [35, 36] (see also [37]). Indeed, consider eq. (2.4) of [36] for the volume functions $\Omega_{n,k}^{(m)}(Y, Z)$. Upon setting $k = m = 2$ there and re-interpreting their Y_α^I as $Y_{\alpha=1}^I = (Z_A)^I$ and $Y_{\alpha=2}^I = (Z_B)^I$, this formula equals our formula for R_n . Very interestingly, the authors of [35, 36] show that certain derivatives w.r.t. Y of $\Omega_{n,k}^{(m)}(Y, Z)$ satisfy Yangian differential equations. Moreover, the latter are related to so-called Capelli differential equations [78]. We comment on how these equations look like in our notations for R_n below.

Momentum twistors are the most appropriate variables to formulate the symmetries of R_n , so we write

$$R_n = R_n(Z_1, \dots, Z_n; Z_A, Z_B). \quad (4.39)$$

We introduce the weight-counting operator

$$H_a := Z_a^J \frac{\partial}{\partial Z_a^J}, \quad (4.40)$$

which commutes with the dual-conformal transformations. Let us recall that R_n carries nonzero dual-conformal weight with respect to Z_A and Z_B ,

$$H_A R_n = -4R_n, \quad H_B R_n = -4R_n, \quad H_i R_n = 0, \quad i = 1, \dots, n. \quad (4.41)$$

Instead of dealing with the $su(2, 2)$ generators, it will be more convenient for us to supplement the dual-conformal algebra with the central element H , thus we effectively deal with the $gl(4)$ symmetry. Namely, the local generators \mathfrak{J}_a acting on the a -th coordinate take the following form in the momentum twistor variables,

$$(\mathfrak{J}_a)^I{}_J = Z_a^I \frac{\partial}{\partial Z_a^J}. \quad (4.42)$$

Here $a = A, B, 1, 2, \dots, n$. Momentum twistors Z_A and Z_B specifying position of the Lagrangian and Z_1, \dots, Z_n specifying the light-contour do not appear on equal footing in the Grassmannian integral representation of R_n (3.34). Then, it is reasonable to distinguish them when studying symmetries of R_n . Thus, we define two sets of the dual-conformal generators as sums of the local ones (4.42) with respect to coordinates $1, \dots, n$ and A, B ,

$$\mathfrak{J}^I_J := \sum_{i=1}^n (\mathfrak{J}_i)^I_J, \quad \bar{\mathfrak{J}}^I_J := (\mathfrak{J}_A)^I_J + (\mathfrak{J}_B)^I_J. \quad (4.43)$$

The statement of the dual-conformal invariance of R_n takes the following form

$$[\mathfrak{J}^I_J + \bar{\mathfrak{J}}^I_J] R_n = -2\delta^I_J R_n. \quad (4.44)$$

In order to study the higher-order symmetries we define the Yangian generators of k -th level in the same way as in eq. (4.12) with the local \mathfrak{J}_a from (4.42),

$$[\mathfrak{J}^{(k)}]^I_J = \sum_{1 \leq i_1 < i_2 < \dots < i_{k+1} \leq n} (\mathfrak{J}_{i_1})^I_{L_1} (\mathfrak{J}_{i_2})^{L_1}_{L_2} \dots (\mathfrak{J}_{i_{k+1}})^{L_k}_J. \quad (4.45)$$

They act nontrivially on the variables of the Wilson loop contour, but the coordinates Z_A, Z_B of the Lagrangian are inert with respect to them. Let us recall that the level-zero generators are the dual-conformal generators from eq. (4.43), i.e. $\mathfrak{J}^{(0)} = \mathfrak{J}$, and we use the shorthand notations $\hat{\mathfrak{J}}$ for the level-one Yangian generators,

$$\hat{\mathfrak{J}}^I_J := [\mathfrak{J}^{(1)}]^I_J = \sum_{1 \leq i < j \leq n} (\mathfrak{J}_i)^I_K (\mathfrak{J}_j)^K_J. \quad (4.46)$$

We derive higher-order symmetries in appendix C. Let us summarize our findings here. We find that the leading singularities R_n are annihilated by the level-two and higher-level Yangian generators

$$[\mathfrak{J}^{(k)}]^I_J R_n = 0, \quad k = 2, \dots, n-1. \quad (4.47)$$

The level-one Yangian generators $\hat{\mathfrak{J}}$ do not annihilate R_n . Nevertheless, there are several symmetry relations involving them. Projecting $\hat{\mathfrak{J}}^I_J$ with the tensor $\langle Z_A, Z_B, *, * \rangle$ we achieve the invariance,

$$\epsilon_{IKLM} Z_A^K Z_B^L \hat{\mathfrak{J}}^I_J R_n = \epsilon_{JKLM} Z_A^K Z_B^L R_n. \quad (4.48)$$

In other words, certain linear combinations of the level-one Yangian generators $\hat{\mathfrak{J}}$ are symmetries of R_n .

Another possibility to arrive at a symmetry relation with the level-one generators is to introduce the dual-conformal variations $\bar{\mathfrak{J}}$ (4.43) which act on the Lagrangian coordinate,

$$[\bar{\mathfrak{J}}^I_K \hat{\mathfrak{J}}^K_J + 2\hat{\mathfrak{J}}^I_J + \mathfrak{J}^I_J] R_n = 0. \quad (4.49)$$

This relation is equivalent to eq. (4.2) of [36] found for the tree-level Amplituhedron volume function.

Let us also notice that projecting (4.49) with the tensor $\langle Z_A, Z_B, *, * \rangle$ we reproduce (4.48). Let us also note that taking the trace of the level-one Yangian generators leads to an invariance,

$$\widehat{\mathfrak{J}}^I_I R_n = 0. \quad (4.50)$$

In fact, the latter relation is a direct consequence of the dual-conformal invariance (4.44).

Besides eq. (4.48), we also find another bi-local combination of the dual-conformal generators which annihilates R_n ,

$$\left[(\mathfrak{J}_i)^K_J \bar{\mathfrak{J}}^J_I + 4 (\mathfrak{J}_i)^K_I \right] R_n = 0, \quad i = 1, \dots, n, \quad (4.51)$$

or more explicitly,

$$\left[\frac{\partial}{\partial Z_i^J} Z_A^J \frac{\partial}{\partial Z_A^I} + \frac{\partial}{\partial Z_i^J} Z_B^J \frac{\partial}{\partial Z_B^I} + 4 \frac{\partial}{\partial Z_i^I} \right] R_n = 0, \quad i = 1, \dots, n. \quad (4.52)$$

Similar to eq. (4.27), we can trace back the origin of (4.51) into the cyclic shift transformations of the Yangian. Indeed, the cyclic shift $Z_i \rightarrow Z_{i+1}$ does not change the Grassmannian representation for R_n , see eq. (3.34). However, it changes the Yangian generators $\widehat{\mathfrak{J}}$ (4.46). Consistency of eqs. (4.48) and (4.49) under the cyclic shift is guaranteed by (4.51).

We notice that the found symmetries (4.47), (4.48), (4.49), (4.50), (4.51) of R_n are consistent with those of r_n from section 4.1.2. Choosing $Z_A \wedge Z_B$ as the infinity bi-twistor brings R_n down to r_n , see eq. (3.38), and the symmetry relations for R_n presented in this section boil down to those for r_n from section 4.1.2. This is almost trivial for those symmetry relations which do not involve differentiation of the Lagrangian coordinate, namely those without $\bar{\mathfrak{J}}$. Indeed, the level- k (with $k \geq 2$) Yangian invariances of R_n in eq. (4.47) turn into those of r_n in eq. (4.23). The trace of level-one generators in eq. (4.50) is consistent with values of traces in eq. (4.22). The frame $Z_A \wedge Z_B \rightarrow \epsilon^{\alpha\dot{\beta}}$ choice in eq. (4.47) restricts indices $I = \alpha$ and $J = (\beta, \dot{\beta})$ that leads to the first two eqs. in (4.22). Eq. (4.51) implies eq. (4.27). In order to see it, we eliminate $\bar{\mathfrak{J}}$ in (4.51) owing to the dual-conformal invariance (4.44) of R_n , and then we can choose $Z_A \wedge Z_B$ as the infinity bi-twistor. Likewise, we can show that eq. (4.49) implies the four eqs. in (4.30) and (4.31).

We delegated the proofs of the relations given above to appendix C. Similar to considerations in subsection 4.1, they rely on the quantum-inverse scattering method constructions [38]. There are slight modifications w.r.t. subsection 4.1 to take into account that R_n carries nonzero dual-conformal weights (4.41).

5 Results for F_n in perturbation theory

In this section we provide evidence that the twistor dual-conformal functions B (3.21) and (3.22), or equivalently their space-time versions b (3.27) and (3.28), provide a basis of leading singularities for F_n at higher loops. This is important both in view of future bootstrap applications and a supersymmetric generalization of the observable [74].

$$[\log \mathcal{A}]_{\text{MHV}}^{2\text{-loop}} = - \sum_{i < k < j < l < i} \text{Diagram}$$

Figure 3. Two-loop logarithm of the MHV amplitude. Figure from ref. [73].




Figure 4. The chiral pentagon one-loop integral I_{ij}^{cp} (5.4), on the left. The one-loop box in the two-mass easy configuration $F_{ij}^{2\text{me}}$ (5.6), on the right.

5.1 Result for F_n at one loop

In ref. [73] a compact expression for the logarithm of MHV amplitudes is given, see figure 3. From this expression it is easy to read off an expression for F_n in a manifestly finite way. The formula reads²

$$F_n^{(1)} = -\frac{1}{2} \sum_{1 \leq i < k < j < l \leq n} \left[I_{ij}^{\text{cp}} LS_{ijkl} + I_{kl}^{\text{cp}} LS_{klij} \right], \quad (5.1)$$

where the rational factors LS are the leading singularities, which depend on a number of twistors specifying the Wilson loop contour and the Lagrangian insertion at (AB) ,

$$LS_{ijkl} := \frac{\langle ijkl \rangle \langle AB | (k-1kk+1) \cap (l-1ll+1) \rangle}{\langle ijAB \rangle \langle ABk-1k \rangle \langle ABkk+1 \rangle \langle ABl-1l \rangle \langle ABll+1 \rangle}, \quad (5.2)$$

and the dual-conformal chiral one-loop pentagon I_{cp} is a finite pure function, figure 4,

$$I_{ij}^{\text{cp}} = \int_{CD} \frac{\langle CD | (i-1ii+1) \cap (j-1jj+1) \rangle \langle ijAB \rangle}{\langle CDi-1i \rangle \langle CDii+1 \rangle \langle CDj-1j \rangle \langle CDjj+1 \rangle \langle CDAB \rangle}. \quad (5.3)$$

Here, starting from the double pentagon integrand in figure 3, a factor $1 = \langle ijAB \rangle / \langle ijAB \rangle$ was split between the leading singularity and the pentagon integral, in order to make the latter a pure function. The chiral pentagon integral evaluates to a pure function

$$I_{ij}^{\text{cp}} = -\frac{1}{2} \log^2(u_3^{ij}) - \text{Li}_2(1 - u_1^{ij}) - \text{Li}_2(1 - u_2^{ij}) \\ - \text{Li}_2\left(1 - \frac{u_1^{ij}}{u_3^{ij}}\right) - \text{Li}_2\left(1 - \frac{u_2^{ij}}{u_3^{ij}}\right) + \text{Li}_2\left(1 - \frac{u_1^{ij} u_2^{ij}}{u_3^{ij}}\right), \quad (5.4)$$

²We explicitly symmetrized the two-loop formula in AB and CD .




Figure 5. The three terms in eq. (5.7) correspond to the three polygons on the figure.

of the three cross-ratios

$$\begin{aligned}
 u_1^{ij} &= \frac{\langle i-1j j+1 \rangle \langle ii+1AB \rangle}{\langle i-1iAB \rangle \langle ii+1j j+1 \rangle}, \\
 u_2^{ij} &= \frac{\langle ii+1j-1j \rangle \langle jj+1AB \rangle}{\langle ii+1j j+1 \rangle \langle j-1jAB \rangle}, \\
 u_3^{ij} &= \frac{\langle i-1ij-1j \rangle \langle ii+1AB \rangle \langle jj+1AB \rangle}{\langle i-1iAB \rangle \langle ii+1j j+1 \rangle \langle j-1jAB \rangle}.
 \end{aligned} \tag{5.5}$$

So we have obtained a closed form analytic expression for $F_n^{(1)}$.

The formula for the pentagon is slightly rewritten from the form given in [73], so that boundary terms from (5.1) with u_1 or u_2 vanishing can be considered without using dilogarithm identities. Let us note that the pure function (5.3) is the two-mass-easy one-loop box function F_{ij}^{2me} . Indeed, in the frame $x_0 \rightarrow \infty$, or equivalently $Z_A \wedge Z_B \rightarrow \epsilon^{\dot{\alpha}\dot{\beta}}$, the dual-conformal chiral pentagon turns into the one-loop box in the easy-mass configuration of the external legs, figure 4,

$$F_{ij}^{2me} := I_{ij}^{cp} \left(u_1^{ij} = \frac{x_{i-1j}^2}{x_{ij}^2}, u_2^{ij} = \frac{x_{ij-1}^2}{x_{ij}^2}, u_3^{ij} = \frac{x_{i-1j-1}^2}{x_{ij}^2} \right). \tag{5.6}$$

The ‘magic’ numerator of (5.4) turns into a pair of chiral fermion propagators, which suppress IR-divergences of the box integral.

We would like to see whether the formula for the one-loop n -point $F_n^{(1)}$ can be expressed in terms of the conformal invariants B (3.21) and (3.22) introduced earlier. In order to achieve this, we rewrite coefficients of the pentagon integrals I_{cp} in terms of the k -point functions $F_k^{(0)}$ (3.23), which are invariant upon the cyclic shift of their arguments. We find,

$$\begin{aligned}
 F_n^{(1)} = & \sum_{\substack{1 \leq i < j \leq n \\ 1 < |i-j| < n-1}} \left[F_n^{(0)}(1, 2, \dots, n) - F_{j-i+1}^{(0)}(i, i+1, \dots, j-1, j) \right. \\
 & \left. - F_{n+1+i-j}^{(0)}(j, j+1, \dots, n-1, n, 1, \dots, i-1, i) \right] I_{ij}^{cp}.
 \end{aligned} \tag{5.7}$$

In order to explain notations, let us graphically interpret eq. (5.7). Consider a polygon with vertices $1, 2, \dots, n$ in figure 5, and attach twistor Z_k to each vertex k . Draw the diagonal (i, j) . The diagonal splits the polygon into two sub-polygons: $i, i+1, \dots, j-1, j$ and $j, j+1, \dots, n-1, n, 1, 2, \dots, i-1, i$. The three terms in (5.7) correspond to the

three polygons. More precisely, we evaluate the twistor function (3.23), which is a linear combination of B 's, on the set of cyclically ordered twistors specified by the polygons. Let us stress that since the diagonal (ij) is not light-like, then the last two terms in (5.7) could not be interpreted as the Born-level observables; instead, they are understood as $(j - i + 1)$ -point and $(n + 1 + i - j)$ -point instances of (3.23). Nevertheless, since (3.23) is a linear combination of the four- and five-point B -functions, all leading singularities in $F_n^{(1)}$ can be expressed in terms of the B functions. Let us also note that (3.23) is invariant under cyclic shifts, and the three terms in the square brackets in (5.7) are invariant under cyclic shifts. We explicitly checked eq. (5.7) up to $n = 15$.

Let us write out the results (5.7) for five and six points explicitly. At $n = 5$, we find

$$F_5^{(1)} = F_5^{(0)} (I_{13}^{\text{cp}} + I_{14}^{\text{cp}} + I_{24}^{\text{cp}} + I_{25}^{\text{cp}} + I_{35}^{\text{cp}}) - B_{1234}I_{14}^{\text{cp}} - B_{2345}I_{25}^{\text{cp}} - B_{1345}I_{13}^{\text{cp}} - B_{1245}I_{24}^{\text{cp}} - B_{1235}I_{35}^{\text{cp}}, \quad (5.8)$$

where the Born-level $F_5^{(0)}$ expression is given in terms of the B functions in eq. (3.25).

At $n = 6$, we have the following explicit expression for the Born-level $F_6^{(0)}$, see eq. (3.23),

$$F_6^{(0)} = B_{1256} + B_{2356} + B_{3456} - B_{12346} - B_{12456} - B_{23456}. \quad (5.9)$$

The one-loop result can be written as follows,

$$F_6^{(1)} = F_6^{(0)} (I_{13}^{\text{cp}} + I_{14}^{\text{cp}} + I_{15}^{\text{cp}} + I_{24}^{\text{cp}} + I_{25}^{\text{cp}} + I_{26}^{\text{cp}} + I_{35}^{\text{cp}} + I_{36}^{\text{cp}} + I_{46}^{\text{cp}}) - (B_{1234} + B_{1456})I_{14}^{\text{cp}} - (B_{1256} + B_{2345})I_{25}^{\text{cp}} - (B_{1236} + B_{3456})I_{36}^{\text{cp}} + (B_{13456} - B_{1356} - B_{3456})I_{13}^{\text{cp}} + (B_{12345} - B_{1245} - B_{2345})I_{15}^{\text{cp}} + (B_{12456} - B_{1256} - B_{2456})I_{24}^{\text{cp}} + (B_{23456} - B_{2356} - B_{3456})I_{26}^{\text{cp}} + (B_{12356} - B_{1256} - B_{2356})I_{35}^{\text{cp}} + (B_{12346} - B_{1246} - B_{2346})I_{46}^{\text{cp}}. \quad (5.10)$$

We hope that these expression may prove useful in future investigations into the structure of the loop corrections for this interesting observable.

5.2 Structure of the two-loop five-point result

Although the Grassmannian formula was derived for $F_n^{(0)}$ only, one may hope that this formula also covers the leading singularities at loop level. We have already seen that this is the case for $F_n^{(1)}$. Here we mention further evidence for this for $n = 5$ at two loops.

In [31] we consider five-cusp planar observable F_5 in the two-loop approximation, namely $F_5^{(L)}$ with $L = 2$ in eq. (3.4). The explicit perturbative calculations reveals that

$$F_5^{(L)} = F_5^{(0)} g_0^{(L)}(\mathbf{u}) + B_{2345}g_1^{(L)}(\mathbf{u}) + B_{1345}g_2^{(L)}(\mathbf{u}) + B_{1245}g_3^{(L)}(\mathbf{u}) + B_{1235}g_4^{(L)}(\mathbf{u}) + B_{1234}g_5^{(L)}(\mathbf{u}), \quad (5.11)$$

where $g_s^{(L)}(\mathbf{u})$ with $s = 0, 1, \dots, 5$ are pure functions of transcendental weight $2L$ of four dual-conformal invariant cross-ratios build from x_1, \dots, x_5, x_0 . There are four independent cross-ratios. They can be chosen as follows

$$\mathbf{u} = \left\{ \frac{x_{13}^2 x_{20}^2 x_{50}^2}{x_{25}^2 x_{10}^2 x_{30}^2}, \frac{x_{14}^2 x_{20}^2 x_{50}^2}{x_{25}^2 x_{10}^2 x_{40}^2}, \frac{x_{24}^2 x_{50}^2}{x_{25}^2 x_{40}^2}, \frac{x_{35}^2 x_{20}^2}{x_{25}^2 x_{30}^2} \right\}. \quad (5.12)$$

The pure functions $\{g_s^{(L)}(\mathbf{u})\}_{s=1\dots 5}$, as well as the accompanying four-point rational prefactors $B_{1234}, B_{2345}, B_{1345}, B_{1245}, B_{1235}$, are related by cyclic shifts: $g_{s+1}^{(L)} = g_s^{(L)}|_{Z_i \rightarrow Z_{i+1}}$. The Born-level $F_5^{(0)}$, which was given in eq. (3.25) in terms of the B 's, is cyclically-symmetric. Let us recall that eq. (5.8) is a one-loop instance $L = 1$ of eq. (5.11).

6 Connection to all-plus amplitudes in pure Yang-Mills theory

The light-like polygon geometry is dual to massless scattering kinematics. One might wonder whether there is a closer analogy to physical scattering processes. We find that this is indeed the case. As we will show presently, the leading singularities r_n appear in n -gluon scattering in the all-plus helicity configuration in pure Yang-Mills theory. What is more, we find a relationship connecting $F_n^{(L)}$ and the $(L + 1)$ -loop all-plus amplitude in pure Yang-Mills theory.

6.1 Matching of leading singularities and one-loop all-plus amplitude

The all-plus amplitude in pure Yang-Mills theory vanishes at tree-level. At one loop, it is a rational function of helicity spinors [79]. At two loops, it contains both polylogarithmic and rational terms, with the maximal transcendental weight being two [52, 54].

In [53, 80, 81] a concise representation of the one-loop all-plus amplitude was given as the sum of C_{kmn} functions defined in eq. (2.9) of that ref. [53]. These functions were found to be conformally invariant, and moreover have a certain directional dual conformal symmetry.

This motivates us to look for a connection between these coefficients, and the conformal invariants discussed in this paper. To match the kinematics, we first go to the frame $x_0 \rightarrow \infty$. We then find that the functions C_{kmn} with $3 \leq k \leq n - 1$ and $k + 1 \leq m \leq n$ are linear combinations of the conformal invariants $\{\text{PT}_n \cdot b_{ijkl}\}$ and $\{\text{PT}_n \cdot b_{ijklm}\}$ defined in the present work in eqs. (3.27) and (3.28) which are normalized with the Parke-Taylor factor (3.30).

In particular, this means that the functions C_{kmn} have a contour integral representation as discussed in section 3.4, and all symmetry statements from section 4.1 hold for C_{kmn} .

Since the one-loop all-plus n -gluon YM amplitude [79] is a linear combination of conformal invariants C_{kmn} , we can rewrite it in terms of the b invariants of the present paper. We find the following explicit formula for the one-loop color-ordered partial amplitude $A_{\text{YM},n}^{(1)}$,³

$$A_{\text{YM},n}^{(1)} = \text{PT}_n a_n(1, 2, \dots, n), \tag{6.1}$$

where the linear combination a_n of the conformal invariants b is completely analogous to the linear combinations of the dual-conformal B 's in (3.23),

$$a_n(1, 2, \dots, n) = \sum_{i=1}^{n-3} b_{i,i+1,n-1,n} - \sum_{i=1}^{n-4} b_{i,i+1,n-2,n-1,n} - \sum_{k=2}^{n-4} \sum_{i=1}^{n-k-3} b_{i,i+1,i+k,i+k+1,n}. \tag{6.2}$$

³For the sake of simplicity, we slightly changed normalization of the amplitude as compared to that of refs. [79] and [54], $A_{\text{YM},n}^{(L)\text{ref.}[79]} = -\frac{i}{3} A_{\text{YM},n}^{(L)\text{our}}$.

Eq. (6.1) allows us to make two interesting observations: firstly, taking into account expression (3.23) for $F_n^{(0)}$ in terms of the B 's, we find that the one-loop all-plus n -gluon YM amplitude is obtained by taking the limit $x_0 \rightarrow \infty$ in the tree-level observable $F_n^{(0)}$,

$$A_{\text{YM},n}^{(1)} = \text{PT}_n \cdot F_n^{(0)} \Big|_{Z_A \wedge Z_B \rightarrow \epsilon^{\dot{\alpha}\dot{\beta}}}. \tag{6.3}$$

Secondly, recalling the contour integral representation (3.41) of the conformal invariants, we could say that the all-plus one-loop Yang-Mills amplitude corresponds to

$$A_{\text{YM},n}^{(1)} = \text{PT}_n \int_{\Gamma \subset \text{Gr}(2,n)} D^{2(n-2)} C \frac{\left(\sum_{i < j} (ij) \langle \lambda_i | x_i \tilde{x}_j | \lambda_j \rangle \right)^2}{(12)(23) \dots (n1)} \prod_{r=1,2} \delta^2 \left(\sum_{i=1}^n c_i^r \lambda_i \right), \tag{6.4}$$

for an appropriate choice of integration cycle Γ on the Grassmannian manifold $\text{Gr}(2, n)$.

6.2 Connection to two-loop all-plus amplitudes

Is the resemblance of the all-plus Yang-Mills amplitude and the observable F_n restricted to the lowest perturbative approximation? To answer this question, let us proceed and compare our analytic one-loop result for $F_n^{(1)}$ (5.7) to the two-loop all-plus Yang-Mills amplitude. Ref. [54] contains the polylogarithmic part of the leading-color two-loop all-plus n -particle amplitude in Yang-Mills theory. More precisely, the leading-color L -loop partial amplitude $A_{\text{YM},n}^{(L)}$ evaluated in dimensional regularization $D = 4 - 2\epsilon$ is defined in eqs. (1.1) and (1.2) of that paper, and the two-loop amplitude has the following expansion in the dimensional regularization parameter,

$$A_{\text{YM},n}^{(2)} = A_{\text{YM},n}^{(1)} I_n^{(1)} + P_{\text{YM},n}^{(2)} + R_{\text{YM},n}^{(2)} + \mathcal{O}(\epsilon) \tag{6.5}$$

where $I_n^{(1)}$ contains dipole infrared divergences (see also [82]),

$$I_n^{(1)} = -\frac{1}{\epsilon^2} \sum_{i=1}^n \left(\frac{\mu^2}{-2p_i \cdot p_{i+1}} \right)^\epsilon + \frac{n\pi^2}{12}. \tag{6.6}$$

The finite part consists of the polylogarithmic $P_{\text{YM},n}^{(2)}$ and rational $R_{\text{YM},n}^{(2)}$ terms. The transcendental terms $P_{\text{YM},n}^{(2)}$ are given in eq. (1.9) of ref. [54]. We find that the finite maximally transcendental piece of the two-loop amplitude can be rewritten in the following suggestive form (we changed normalization of the amplitude as given in footnote 3)

$$P_{\text{YM},n}^{(2)} = -\text{PT}_n \sum_{\substack{1 \leq i < j \leq n \\ 1 < |i-j| < n-1}} \left[a_{j-i+1}(i, i+1, \dots, j-1, j) + a_{n+1+i-j}(j, j+1, \dots, n-1, n, 1, \dots, j-1, j) \right] F_{ij}^{2\text{me}}, \tag{6.7}$$

with the two-mass-easy function $F_{ij}^{2\text{me}}$ (5.6). Let us note that in the four-point case $n = 4$ the coefficient in the square bracket is trivially zero and $P_{\text{YM},4}^{(2)} = 0$ as expected [83].

For the sake of clarity, let us mention that the terms a_{j-i+1} and $a_{n+1+i-j}$ in eq. (6.7) are $(j-i+1)$ -point and $(n+1+i-j)$ -point instances of the function (6.2), respectively, evaluated along the two subpolygons in figure 5. The twistor function (6.2) is a linear combination of the b 's. Let us stress that a_{j-i+1} and $a_{n+1+i-j}$ from (6.7) cannot be interpreted as the one-loop amplitudes because of the momentum conservation.

We would like to match the polylogarithmic finite part $P_{\text{YM},n}^{(2)}$ of the YM two-loop amplitude (6.7) with the one-loop observable $F_n^{(1)}$ (5.7).

Comparing their accompanying leading singularities in the frame $x_0 \rightarrow \infty$, we observe they are almost identical, with the difference being proportional to the one-loop YM amplitude $A_{\text{YM},n}^{(1)}$ (6.3),

$$P_{\text{YM},n}^{(2)} + A_{\text{YM},n}^{(1)} \left(\sum_{\substack{1 \leq i < j \leq n \\ 1 < |i-j| < n-1}} F_{ij}^{2\text{me}} \right) = \text{PT}_n \cdot F_n^{(1)} \Big|_{Z_A \wedge Z_B \rightarrow \epsilon^{\alpha\beta}}. \quad (6.8)$$

This discrepancy can be compensated by a finite IR subtraction, namely by redefinition of $I_n^{(1)}$ (6.6) in eq. (6.5).

Looking more closely at eq. (6.8), one realizes that the additional term in parenthesis on the left-hand-side is exactly the (finite part of the) one-loop MHV amplitude. Recalling the Wilson loop/scattering amplitude duality, one realizes that this factor appears in the denominator of the definition of F_n , cf. eq. (1.1). In this way we arrive at the following duality relation,

$$\text{PT}_n \langle W_n \mathcal{L}(x_0) \rangle_{x_0 \rightarrow \infty} \sim A_n^{\text{YM}, \text{all-plus}}. \quad (6.9)$$

This relation is to be understood for the leading transcendental parts of the two objects in the planar limit, up to scheme differences (as usual in Wilson loop/scattering amplitude dualities, these scheme differences arises due to the fact that formally, the Wilson loop has ultraviolet divergences, while the amplitudes have infrared divergences). We have just shown that eq. (6.9) holds up to order g_{YM}^4 in perturbation theory, for arbitrary n .

Let us present another version of this formula, directly in terms of finite quantities, similar to eq. (6.8). The perturbative expansion of the all-plus Yang-Mills amplitude starts from the one-loop approximation $A_{\text{YM},n}^{(1)}$, which is the rational function given in eq. (6.1), up to $\mathcal{O}(\epsilon)$ corrections. The perturbative expansion of the renormalized planar all-plus amplitude can be written in the following factorized form,

$$A_n^{\text{YM}} = \mathcal{Z}_{\text{IR}}^{\text{YM}} A_{\text{YM},n}^{(1)} \mathcal{H}_n^{\text{YM}}. \quad (6.10)$$

Here the finite remainder $\mathcal{H}_n^{\text{YM}}$ is known for $n = 4$ up to the three-loop order [84, 85]. The infrared divergences are taken into account by the factor $\mathcal{Z}_{\text{IR}}^{\text{YM}}$, which implements the minimal subtraction. Next we invoke the Wilson loop/amplitude duality (2.4)

$$\langle W_n \rangle \sim \frac{A_n^{\text{MHV}}}{A_{n,\text{tree}}^{\text{MHV}}} = \mathcal{Z}_{\text{IR}}^{\text{MHV}} \mathcal{H}_n^{\text{MHV}}, \quad (6.11)$$

where the factor $\mathcal{Z}_{\text{IR}}^{\text{MHV}}$ minimally subtracts infrared divergences, and $\mathcal{H}_n^{\text{MHV}}$ is the finite remainder. We can identify the pure Yang-Mills and $\mathcal{N} = 4$ super-Yang-Mills infrared factors, $\mathcal{Z}_{\text{IR}}^{\text{MHV}} \sim \mathcal{Z}_{\text{IR}}^{\text{YM}}$. Indeed, in (6.9), we compare only the leading transcendental terms of the two objects. The infrared factors coincide in this approximation, in agreement with the maximal transcendentality principle [86]. Taking the logarithm on both sides of eq. (6.9), we then obtain the following relation between three finite objects: the finite remainder of the all-plus amplitude, the finite remainder of the MHV amplitude, and the ratio F_n ,

$$\log \mathcal{H}_n^{\text{MHV}} + \log \left(F_n / F_n^{(0)} \right)_{x_0 \rightarrow \infty} \sim \log \mathcal{H}_n^{\text{YM}} + \mathcal{O}(\epsilon). \quad (6.12)$$

As before, this is to be understood at the level of the leading transcendental terms. Let us now check eq. (6.12) for $n = 4$. Note that since A_n^{YM} vanishes at tree-level, the quantities on the r.h.s. of eq. (6.12) enter at one loop order higher than the quantities on its l.h.s. Substituting the available perturbative data into this equation, namely the ABDK/BDS ansatz [60, 87] expression for the four-point MHV amplitude, the known expression for F_4 up to two loops, and the three-loop all-plus four-point amplitude [27, 84, 85] into this equation, we find perfect agreement.

Let us discuss these results. Previously, it had been noted that *integrand*s of MHV amplitudes in $\mathcal{N} = 4$ sYM and the integrands of the all-plus pure YM amplitudes are related. At one-loop, they are related by a dimensional shift [88], or, equivalently, the pure YM integrands are obtained from the sYM ones via a particular insertion of $(D - 4)$ -dimensional components of the loop momenta into the integrand. At two loops, part of the pure YM integrand is again obtained from the sYM one, albeit through a slightly more complicated insertion [89, 90].

Here, we have provided evidence for a connection between three quantities: the L -loop observable $F_n^{(L)}$ and the (finite-part of the) L -loop MHV amplitude in $\mathcal{N} = 4$ sYM, and the (finite part of the) $(L + 1)$ -loop all-plus amplitude in pure YM theory. This is a relation between integrated quantities.

7 Discussion and outlook

In this paper, we studied a class of finite observables in maximally supersymmetric Yang-Mills theory. The bosonic observable, which is essentially a Wilson loop with a Lagrangian insertion, had mainly been studied for four-sided Wilson loops. We performed, for the first time, a systematic analysis of the n -particle case.

Already at Born level, we uncovered interesting new structures. While the analytic n -particle expression has the expected dual conformal symmetry (following from the definition of the correlator in position space), we found that there is also an unexpected conformal symmetry, i.e. a conformal symmetry in momentum space.

This conformal symmetry is present at Born level, and also at loop level, where it manifests itself in the leading singularities that multiply the transcendental functions. Conjecturally, the leading singularities are given by a Grassmannian formula, for which we gave two versions, one for general kinematics, and one in the dual conformal frame $x_0 \rightarrow \infty$.

We studied the symmetry properties in the Grassmannian description, which allowed us to prove the conformal symmetry, and to establish several higher-order symmetries related to a Yangian. Curiously, a very similar Grassmannian formula also appeared independently in another context, namely the canonical form for the tree-level Amplituhedron [35, 36].

We also computed the full one-loop n -particle observable and provided a compact analytic formula for it. This confirms our leading singularity conjecture for this case. The two-loop five-particle case will be discussed in a separate publication [31].

There are striking similarities between F_n in the dual conformal frame $x_0 \rightarrow \infty$ and all-plus amplitudes in pure Yang-Mills theory. We found that for all cases investigated, the leading weight terms of the two objects are dual to each other. The origin of this duality is unknown. If true in general, it could be used to predict the leading transcendental terms of planar all-plus amplitudes in pure Yang-Mills theory: the four-point four-loop case could be deduced from [91], and the five-point three-loop case from the results of [31].

There are a number of open questions and natural extensions of this work:

1. The hidden conformal symmetry we found in the frame $x_0 \rightarrow \infty$ ought to have a fundamental explanation, which is yet to be found. Moreover, in the case of scattering amplitudes, writing both conformal and dual conformal symmetry generators in the same variables allowed one to see that their closure gives a Yangian algebra [4]. What is the full symmetry of the Wilson loop with Lagrangian insertion?
2. We find it very intriguing that similar Grassmannian formulas appeared in the context of tree-level Amplituhedron volume functions [35–37]. It would be interesting to find a momentum-space version that would make it possible to study higher Yangian charges directly in momentum space, as discussed in section 4.2.
3. In ref. [30], a novel decomposition of F_4 in terms of certain ‘negative geometries’ was given, generalizing ideas from the Amplituhedron. In this way, the contributions at different loop orders could be expressed in terms of manifestly finite Feynman integrals. In this way, certain all-order contributions could be summed and compared to strong coupling results. It would be very interesting to generalize these results to $n > 4$ [92].
4. In the case of scattering amplitudes, (broken) superconformal symmetry, in particular the \overline{Q} -equation, played an important role in understanding the structure of the perturbative results [93]. It would be very interesting to define supersymmetrised version of our observable, and study its \overline{Q} symmetry [74].

We are looking forward to seeing progress on these exciting questions in the near future.

Acknowledgments

The authors are indebted to Jaroslav Trnka and Nima Arkani-Hamed for multiple discussions and insightful comments, and to Livia Ferro and Simone Zoia for helpful comments on the draft. This research received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 725110), *Novel structures in scattering amplitudes*. DC is supported by the French National Research Agency in the framework of the *Investissements d'avenir* program (ANR-15-IDEX-02).

A Leading singularities B in terms of space-time variables

In this appendix we show how the dual-conformal invariants B from the main text (eqs. (3.21) and (3.22)) look like when expressed in terms of space-time variables.

Let us consider as an example B_{1234} , and set $n = 5$. Firstly, we make the denominator parity-even

$$B_{1234}|_{n=5} = \frac{\langle 1234 \rangle}{\langle AB12 \rangle \langle AB23 \rangle \langle AB34 \rangle} \cdot \frac{\langle 1234 \rangle \langle AB(\overline{14}) \rangle}{\langle AB14 \rangle \langle AB(\overline{14}) \rangle} \quad (\text{A.1})$$

where $(\overline{14}) = (512) \cap (345)$ is the conjugate of (14) , see e.g. [73] and eq. (3.19).

The twistor four-brackets of the type $\langle ii+1jj+1 \rangle$ and $\langle ABii+1 \rangle$ are provided in (3.15). The the parity-even combination in the denominator of (A.1) can be written in x coordinates with the help of

$$\begin{aligned} \langle ABij \rangle \langle AB(\overline{ij}) \rangle &= \langle AB \rangle \langle i-1i \rangle \langle ii+1 \rangle \langle j-1j \rangle \langle jj+1 \rangle \\ &\times \left[-x_{0i-1}^2 x_{0j-1}^2 x_{ij}^2 + x_{0i-1}^2 x_{0j}^2 x_{ij-1}^2 + x_{0i}^2 x_{0j-1}^2 x_{i-1j}^2 - x_{0i}^2 x_{0j}^2 x_{i-1j-1}^2 \right]. \end{aligned} \quad (\text{A.2})$$

The numerator of (A.1) contains both the parity even and the parity odd pieces. They are revealed as follows

$$\langle AB(\overline{14}) \rangle \langle 1234 \rangle - \langle AB14 \rangle \langle 2351 \rangle \langle 2345 \rangle = \langle AB \rangle (\text{PT}_5)^{-1} \epsilon_{123450}, \quad (\text{A.3})$$

$$\begin{aligned} &\langle AB(\overline{14}) \rangle \langle 1234 \rangle + \langle AB14 \rangle \langle 2351 \rangle \langle 2345 \rangle \\ &= \langle AB \rangle (\text{PT}_5)^{-1} \left(x_{13}^2 x_{24}^2 x_{50}^2 + x_{13}^2 x_{25}^2 x_{40}^2 - x_{14}^2 x_{25}^2 x_{30}^2 + x_{14}^2 x_{20}^2 x_{35}^2 - x_{10}^2 x_{24}^2 x_{35}^2 \right), \end{aligned} \quad (\text{A.4})$$

where PT_5 is the Parke-Taylor factor (3.30), which balances the spinor helicity weight on both sides of the previous equation.

Substituting the previous identities into eq. (A.1) we obtain space-time expression for one of the dual-conformal invariants at $n = 5$, where the numerator is split in the parity-even and parity-odd pieces,

$$B_{1234}|_{n=5} = \frac{1}{2} \frac{x_{13}^2 (x_{13}^2 x_{24}^2 x_{50}^2 + x_{13}^2 x_{25}^2 x_{40}^2 - x_{14}^2 x_{25}^2 x_{30}^2 + x_{14}^2 x_{20}^2 x_{35}^2 - x_{10}^2 x_{24}^2 x_{35}^2 + \epsilon_{123450})}{x_{10}^2 x_{20}^2 x_{30}^2 (x_{10}^2 x_{35}^2 x_{40}^2 - x_{13}^2 x_{40}^2 x_{50}^2 + x_{14}^2 x_{30}^2 x_{50}^2)}. \quad (\text{A.5})$$

B Details of symmetry proofs for r_n

B.1 Dual-conformal algebra

The dual-conformal algebra is an instance of $su(2, 2)$. The algebra comprises Lorentz rotations split into a pair of symmetric tensors $M_{\alpha\beta}$ and $\bar{M}_{\dot{\alpha}\dot{\beta}}$, dilatation D which counts dimension of the remaining generators, space-time shifts $P_{\alpha\dot{\alpha}}$ and the conformal boost $K_{\alpha\dot{\alpha}}$, which satisfy the following commutation relation

$$[K_{\alpha\dot{\alpha}}, P^{\dot{\beta}\beta}] = \delta_{\alpha}^{\beta} \delta_{\dot{\alpha}}^{\dot{\beta}} D + M_{\alpha}^{\beta} \delta_{\dot{\alpha}}^{\dot{\beta}} + \bar{M}_{\dot{\alpha}}^{\dot{\beta}} \delta_{\alpha}^{\beta}. \quad (\text{B.1})$$

Representation of the dual-conformal algebra generators in the momentum twistor variables $Z^I = (\lambda^{\alpha}, \mu^{\dot{\alpha}})$ can be read off from the matrix (4.1). It is also instructive to provide generators in the coordinate space

$$P_{\alpha\dot{\alpha}} = \frac{\partial}{\partial x^{\alpha\dot{\alpha}}}, \quad K^{\dot{\alpha}\alpha} = x^{\dot{\alpha}\beta} x^{\beta\alpha} \frac{\partial}{\partial x^{\dot{\beta}\beta}} + x^{\dot{\alpha}\beta} \lambda_{\beta} \frac{\partial}{\partial \lambda_{\alpha}}, \quad (\text{B.2})$$

$$D = x_{\alpha\dot{\alpha}} \frac{\partial}{\partial x_{\alpha\dot{\alpha}}} + \frac{1}{2} \lambda_{\alpha} \frac{\partial}{\partial \lambda_{\alpha}}, \quad M_{\alpha\beta} = x^{\dot{\gamma}}_{(\alpha} \frac{\partial}{\partial x^{\dot{\gamma}\beta)} + \lambda_{(\alpha} \frac{\partial}{\partial \lambda^{\beta)}, \quad \bar{M}_{\dot{\alpha}\dot{\beta}} = x^{\gamma}_{(\dot{\alpha}} \frac{\partial}{\partial x^{\dot{\beta})\gamma)}. \quad (\text{B.3})$$

B.2 Variable change in the conformal boost generator

There is a variable change that maps the conformal boost generator $\mathbb{K}_{\alpha\dot{\alpha}}$ (3.32) to the level-one generator $\widehat{\mathfrak{J}}$ of the Yangian (4.7) built from the dual-conformal algebra. We used this fact in (4.32) in order to establish the conformal symmetry of r_n . Let us give some more details on this variable change $(\lambda, \tilde{\lambda}) \rightarrow (\lambda, \mu)$, which is given by the following equations,

$$\begin{aligned} \mu_j^{\dot{\alpha}} &= x_n^{\dot{\alpha}\alpha} \lambda_{j\alpha} - \sum_{k=j+1}^n \langle \lambda_k \lambda_j \rangle \tilde{\lambda}_k^{\dot{\alpha}}, \quad j = 1, \dots, n-1, \\ \mu_n^{\dot{\alpha}} &= x_n^{\dot{\alpha}\alpha} \lambda_{n\alpha}, \end{aligned} \quad (\text{B.4})$$

where x_n is fixed. Differentiations in the helicity spinors $(\lambda, \tilde{\lambda})$ are reexpressed into derivatives with respect to the twistor variables $Z = (\lambda, \mu)$ as follows

$$\frac{\partial}{\partial \lambda_{j\alpha}} = \frac{\partial}{\partial \lambda_{j\alpha}} - x_j^{\dot{\alpha}\alpha} \frac{\partial}{\partial \mu_j^{\dot{\alpha}}} - \tilde{\lambda}_j^{\dot{\alpha}} \sum_{k=1}^{j-1} \lambda_k^{\alpha} \frac{\partial}{\partial \mu_k^{\dot{\alpha}}}, \quad (\text{B.5})$$

$$\frac{\partial}{\partial \tilde{\lambda}_j^{\dot{\alpha}}} = - \sum_{k=1}^{j-1} \langle \lambda_j \lambda_k \rangle \frac{\partial}{\partial \mu_k^{\dot{\alpha}}}, \quad j = 1, \dots, n. \quad (\text{B.6})$$

Substituting (B.5) and (B.6) in the second order differential operator of the conformal boost \mathbb{K} (3.32) one finds how it acts on a function of momentum twistors $f(\lambda_1, \mu_1, \dots, \lambda_n, \mu_n)$ normalized by the Parke-Taylor factor (3.30). The calculation is done in detail in [32]. One finds,

$$\begin{aligned} & \left(\sum_{j=1}^n \frac{\partial^2}{\partial \lambda_j^{\alpha} \partial \tilde{\lambda}_j^{\dot{\alpha}}} \right) \frac{f(\lambda, \mu)}{\langle 12 \rangle \dots \langle n1 \rangle} \\ &= - \frac{1}{\langle 12 \rangle \dots \langle n1 \rangle} \sum_{1 \leq j < k \leq n} \left(\lambda_{j\alpha} \frac{\partial}{\partial \lambda_j^{\beta}} \lambda_k^{\beta} \frac{\partial}{\partial \mu_k^{\dot{\alpha}}} + \lambda_{j\alpha} \frac{\partial}{\partial \mu_j^{\dot{\beta}}} \mu_k^{\dot{\beta}} \frac{\partial}{\partial \mu_k^{\dot{\alpha}}} \right) f(\lambda, \mu), \end{aligned} \quad (\text{B.7})$$

provided that f is annihilated by the dual-conformal shift $P_{\alpha\dot{\alpha}}$,

$$\left(\sum_{j=1}^n \lambda_j^\alpha \frac{\partial}{\partial \mu_j^{\dot{\alpha}}} \right) f(\lambda, \mu) = 0, \quad (\text{B.8})$$

and that f is a homogeneous function of the momentum twistor variables with zero weight with respect to each twistor,

$$\left(\lambda_j^\alpha \frac{\partial}{\partial \lambda_j^\alpha} + \mu_j^{\dot{\alpha}} \frac{\partial}{\partial \mu_j^{\dot{\alpha}}} \right) f(\lambda, \mu) = 0, \quad j = 1, \dots, n. \quad (\text{B.9})$$

The leading singularities r_n satisfy both these requirements, thus (B.7) justifies eq. (4.32).

B.3 Implications of Yangian cyclicity in space-time coordinates

In this section we provide space-time form of eq. (4.27). As we showed in section 4.1.2, this equation originates from the cyclic shift transformations of the Yangian generators.

Eq. (4.27) at $I = \dot{\alpha}$ is an immediate consequence of the Poincaré invariance (4.4). At $I = \alpha$, eq. (4.27) takes the following form,

$$\frac{\partial}{\partial \mu_i^{\dot{\beta}}} K^{\dot{\beta}\alpha} r_n(Z_1, \dots, Z_n) = 4 \frac{\partial}{\partial \lambda_i^\alpha} r_n(Z_1, \dots, Z_n), \quad i = 1, \dots, n, \quad (\text{B.10})$$

with the dual-conformal boost generator in momentum twistors variables

$$K^{\dot{\alpha}\alpha} = \sum_{i=1}^n \mu_i^{\dot{\alpha}} \frac{\partial}{\partial \lambda_i^\alpha}. \quad (\text{B.11})$$

It is instructive to rewrite relation (B.10) in the space-time coordinates,

$$\left[\lambda_{i+1}^\gamma \frac{\partial}{\partial x_i^{\dot{\gamma}\gamma}} K^{\dot{\gamma}\alpha} - 4 \lambda_{i+1}^\gamma x_i^{\dot{\gamma}\alpha} \frac{\partial}{\partial \tilde{x}_i^{\dot{\gamma}\gamma}} + 4(i+1) \frac{\partial}{\partial \lambda_{i\alpha}} \right] r_n(\lambda_1, x_1, \dots, \lambda_n, x_n) = 0, \quad (\text{B.12})$$

where $K^{\dot{\alpha}\alpha}$ is the dual-conformal boost generator in the space-time coordinate representation, see (B.2),

$$K^{\dot{\alpha}\alpha} = \sum_{i=1}^n \left[x_i^{\dot{\alpha}\beta} x_i^{\dot{\beta}\alpha} \frac{\partial}{\partial x_i^{\dot{\beta}\beta}} + x_i^{\dot{\alpha}\beta} \lambda_{i\beta} \frac{\partial}{\partial \lambda_{i\alpha}} \right]. \quad (\text{B.13})$$

Contracting (B.12) with $\lambda_{i\alpha}$ and taking into account that r_n carries zero helicity weight with respect to each point and the twistor incidence relation $\lambda_i^\alpha \frac{\partial r_n}{\partial x_i^{\dot{\alpha}\alpha}} = 0$, we obtain an interesting implication of (B.10),

$$\frac{\partial}{\partial x_i^{\dot{\alpha}\alpha}} \left(K^{\dot{\alpha}\alpha} r_n \right) = 4 x_i^{\dot{\alpha}\alpha} \frac{\partial}{\partial x_i^{\dot{\alpha}\alpha}} r_n, \quad i = 1, \dots, n. \quad (\text{B.14})$$

We can think of (B.14) as an anomaly equation for the dual-conformal symmetry, which is broken by the frame fixing $x_0 \rightarrow \infty$, see (4.5).

B.4 From R-operators to the Grassmannian

In the Yangian symmetry considerations of the leading singularities r_n , we rely upon their R-operator decomposition (4.17). In order to prove this relation, we would like to show that it is equivalent to the Grassmannian representation (3.41). In other words,

$$\int \omega(C) \Delta(C, Z) = \prod_{i=2}^{\overrightarrow{n-1}} R_{i+1i} \cdot \prod_{i=1}^{\overleftarrow{n-2}} R_{ii+1} |\Omega_n\rangle \tag{B.15}$$

where the arrows specify ordering of the R-operator product; C is a $2 \times n$ matrix representing a 2-plane in \mathbb{C}^n ,

$$C = \begin{pmatrix} c_1^1 & \dots & c_n^1 \\ c_1^2 & \dots & c_n^2 \end{pmatrix} := (\vec{c}_1 \dots \vec{c}_n) \tag{B.16}$$

with 2×2 minors denoted as (ab) ; $\omega(C)$ is a measure on the Grassmannian $\text{Gr}(2, n)$ (see (3.35) and (3.36)) for the fixed ordering,

$$\omega(C) := \frac{D^{2(n-2)} C}{(12)(23) \dots (n1)}, \tag{B.17}$$

and distribution $\Delta(C, Z)$ is defined as

$$\Delta(C, Z) := \left(\sum_{i < j} (ij) [\mu_i \mu_j] \right)^2 \prod_{r=1,2} \delta^2 \left(\sum_{i=1}^n c_i^r \lambda_i \right). \tag{B.18}$$

We note that R_{ij} (see eq. (4.13)) acts on Z_i^I as the shift $Z_i \rightarrow Z_i + tZ_j$. In view of the identity,

$$\vec{c}_i(Z_i + tZ_j) + \sum_{k \neq i} \vec{c}_k Z_k = (\vec{c}_j + t\vec{c}_i)Z_j + \sum_{k \neq j} \vec{c}_k Z_k, \tag{B.19}$$

the action of R_{ij} onto $\Delta(C, Z)$ (B.18) by the shift of twistor variables is equivalent to the column transformation $\vec{c}_j \rightarrow \vec{c}_j + t\vec{c}_i$ of matrix C (B.16).

Each R-operator in the product (B.15) contains one-fold integration (see (4.13)). Let us label the corresponding integration variables as $t_1, t_2, \dots, t_{2n-4}$ from right to the left. Let us start with matrix C_0

$$C_0 = \begin{pmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & 0 & \dots & 0 & 1 \end{pmatrix} \tag{B.20}$$

which represents a dimension zero cell of the Grassmannian $\text{Gr}(2, n)$ and corresponds to the pseudo-vacuum state of the spin-chain (4.18),

$$\Delta(C_0, Z) = |\Omega_n\rangle. \tag{B.21}$$

Each R-operator from (B.15) implements a column transformation of C_0 introducing one extra t -parameter and increments dimension of the Grassmannian cell. Acting with all

$(2n - 4)$ R-operators from (B.15) we arrive at the top cell of the Grassmannian. The measure $\omega(C)$ (B.17) takes the canonical form in the t -parametrization, and each R-operator (4.13) contributes one $d \log$ factor,

$$\omega(C) = \frac{dt_1}{t_1} \wedge \frac{dt_2}{t_2} \wedge \dots \wedge \frac{dt_{2n-4}}{t_{2n-4}}. \tag{B.22}$$

The described procedure is equivalent to the BCFW-decomposition and positroid stratification of the Grassmannians in the context of $\mathcal{N} = 4$ sYM amplitudes [6].

Let us also note that the decomposition (B.15) is not unique. We could represent the Grassmanian top cell by another sequence of R-operators. However, not every decomposition is compatible with the monodromy (4.9). In other words, the Yangian symmetry could be not immediately obvious if we use a decomposition other than (B.15).

It is also instructive to consider the decomposition (B.15) at $n = 4$,

$$r_4 = \underbrace{R_{32}}_{t_4} \underbrace{R_{43}}_{t_3} \underbrace{R_{23}}_{t_2} \underbrace{R_{12}}_{t_1} |\Omega_4\rangle = \int \omega(C_4) \Delta(C_4) \tag{B.23}$$

where we explicitly indicate the labeling of the one-fold integrations (4.13), and $|\Omega_4\rangle = [\mu_1 \mu_4]^2 \delta^2(\lambda_1) \delta^2(\lambda_4)$ (4.18). The sequence of R-operators from (B.23) corresponds to the following sequence of C_0 column transformations (B.20), such that R_{ij} acts as $\vec{c}_j \rightarrow \vec{c}_j + t\vec{c}_i$,

$$\begin{aligned} C_0 &:= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \xrightarrow{R_{12}} C_1 := \begin{pmatrix} 1 & t_1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \xrightarrow{R_{23}} C_2 := \begin{pmatrix} 1 & t_1 & t_1 t_2 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \rightarrow \\ &\xrightarrow{R_{43}} C_3 := \begin{pmatrix} 1 & t_1 & t_1 t_2 & 0 \\ 0 & 0 & t_3 & 1 \end{pmatrix} \xrightarrow{R_{32}} C_4 := \begin{pmatrix} 1 & t_1(1 + t_2 t_4) & t_1 t_2 & 0 \\ 0 & t_3 t_4 & t_3 & 1 \end{pmatrix}, \end{aligned} \tag{B.24}$$

and C_i represents i -th dimensional cell of $\text{Gr}(2, 4)$; and $\omega(C_4)$ is the canonical form (B.22).

C Details of integrability proofs for R_n

In this appendix we provide proofs of the symmetry relations for the leading singularities R_n presented in section 4.3. The line of argumentation is similar to that of section 4.1.2 where the Yangian symmetries of r_n are considered, but requires several extra technical details. We split the proof in several steps. In C.1 we provide the spin-chain formulation of the Yangian algebra generators and define the R-operators. The latter are employed in C.2 to represent R_n as a product of R-operators acting on the pseudo-vacuum state of the spin-chain. In C.3 we discuss the cyclic shift transformations of the Yangian. In C.4 we provide several technical statements which are used in the proof of the Yangian invariance relations in C.5.

C.1 R-operators, Lax-operators, and the monodromy matrix

Let us denote $\mathcal{H}_i^{(\delta_i)}$ the space of homogeneous functions of Z_i of the homogeneity degree δ_i , namely $H_i = \delta_i$ is the dual-conformal weight (4.40). An elementary block in the

decomposition of the leading singularities R_n will be the R-operator. We will need to introduce the spectral parameter u in the R-operator,

$$\left[\mathbf{R}_{ij}^{(u)} G \right] (Z_i, Z_j) := \int \frac{dt}{t^{1+u}} G(Z_i + tZ_j, Z_j). \quad (\text{C.1})$$

The R-operator (C.1) acts in the tensor product $\mathcal{H}_i^{(\delta_i)} \otimes \mathcal{H}_j^{(\delta_j)}$ and maps it into $\mathcal{H}_i^{(\delta_i-u)} \otimes \mathcal{H}_j^{(\delta_j+u)}$. In other words, the R-operator does not preserve the dual-conformal weight,

$$\left[H_1, \mathbf{R}_{12}^{(u)} \right] = -u \mathbf{R}_{12}^{(u)}, \quad \left[H_2, \mathbf{R}_{12}^{(u)} \right] = u \mathbf{R}_{12}^{(u)}. \quad (\text{C.2})$$

In order to construct the Yangian generators (4.45), we combine the dual-conformal generators (4.42) in the 4×4 matrix depending on the spectral parameter u , and call it the Lax operator,

$$[\mathbf{L}(u)]^I{}_J = u \delta_J^I + Z^I \frac{\partial}{\partial Z^J}. \quad (\text{C.3})$$

The Lax operator is a local object acting in one site of the spin chain. We also decorate the Lax operator with the dual-conformal weight, $\mathbf{L}_i^{(\delta)}(u)$, to indicate that it acts onto $\mathcal{H}_i^{(\delta)} \otimes \mathbb{C}^4$. The matrix product of a pair of Lax operators acting in the i -th and j -th sites has nice exchange relations with the R-operators (C.1),

$$\mathbf{R}_{ij}^{(a)} \mathbf{L}_i^{(\delta_i)}(u) \mathbf{L}_j^{(\delta_j)}(v) = \mathbf{L}_i^{(\delta_i-a)}(v) \mathbf{L}_j^{(\delta_j+a)}(u) \mathbf{R}_{ij}^{(a)}, \quad a = v - u \quad (\text{C.4})$$

and

$$\mathbf{R}_{ji}^{(a)} \mathbf{L}_i^{(\delta_i)}(u) \mathbf{L}_j^{(\delta_j)}(v) = \mathbf{L}_i^{(\delta_i+a)}(u) \mathbf{L}_j^{(\delta_j-a)}(v) \mathbf{R}_{ji}^{(a)}, \quad a = v - u - \delta_i + \delta_j. \quad (\text{C.5})$$

These local exchange relations are exactly what we need to establish the symmetries of R_n .

A simple corollary of the exchange relation (C.4) is the dual-conformal invariance (4.42) of the R-operator

$$\left[\mathbf{R}_{ij}^{(a)}, \mathfrak{J}_i + \mathfrak{J}_j \right] = 0. \quad (\text{C.6})$$

We form the monodromy matrix $\mathbf{T}(u)$ as the matrix product of the Lax operators (C.3) acting on spin-chain sites $1, 2, \dots, n$,

$$[\mathbf{T}(u)]^I{}_J := [\mathbf{L}_1(u)]^I{}_{K_1} [\mathbf{L}_2(u)]^{K_1}{}_{K_2} \dots [\mathbf{L}_n(u)]^{K_{n-1}}{}_J. \quad (\text{C.7})$$

The monodromy matrix is a polynomial in the spectral parameter u . The coefficients of the polynomial are the Yangian generators (4.45) acting in the spin-chain sites $1, 2, \dots, n$,

$$[\mathbf{T}(u)]^I{}_J = u^n \delta_J^I + u^{n-1} [\mathfrak{J}^{(0)}]^I{}_J + u^{n-2} [\mathfrak{J}^{(1)}]^I{}_J + \sum_{k=2}^{n-1} u^{n-k-1} [\mathfrak{J}^{(k)}]^I{}_J, \quad (\text{C.8})$$

where we recall $\mathfrak{J}^{(0)} = \mathfrak{J}$ (4.43) and $\mathfrak{J}^{(1)} = \widehat{\mathfrak{J}}$ (4.46).

C.2 From R-operators to the Grassmannian

We will need an analogue of the R-operator decomposition (4.17) for R_n . The presentation in this section parallels the one in section B.4. We define the pseudo-vacuum of the spin chain which belongs to $\mathcal{H}_1^{(-4)} \otimes \mathcal{H}_n^{(-4)}$,

$$|\Omega_n\rangle := \delta^4(Z_1) \delta^4(Z_n). \quad (\text{C.9})$$

In order to represent the leading singularities R_n , we act onto the pseudo-vacuum by the product of $2n$ R-operators (C.1) as follows,

$$R_n = R_{AB}^{(-1)} R_{BA}^{(-1)} |\Phi_n\rangle, \quad (\text{C.10})$$

$$|\Phi_n\rangle := R_{1A}^{(-4)} R_{nB}^{(-4)} \prod_{i=2}^{n-1} R_{i+1i}^{(0)} \cdot \prod_{i=1}^{n-2} R_{ii+1}^{(0)} |\Omega_n\rangle, \quad (\text{C.11})$$

where for the sake of further convenience we split the product into two factors containing two and $2n - 2$ R-operators, and we introduced a shorthand notation $|\Phi_n\rangle$ for the latter product acting on the pseudo-vacuum. The dual-conformal weight counting of the R-operators (C.2) enables us to find the weights of R_n and $|\Phi_n\rangle$,

$$R_n, |\Phi_n\rangle \in \mathcal{H}_A^{(-4)} \otimes \mathcal{H}_B^{(-4)} \otimes \mathcal{H}_1^{(0)} \otimes \dots \otimes \mathcal{H}_n^{(0)}. \quad (\text{C.12})$$

The resulting expression in (C.10) coincides with the contour integral (3.34) representation of R_n ,

$$R_n = \int \omega(C) \Delta(C, Z). \quad (\text{C.13})$$

Here we use notations $\omega(C)$ for the measure (see (3.35) and (3.36))

$$\omega(C) := \frac{(AB)^2 D^{2n} C}{(12)(23) \dots (n1)}, \quad (\text{C.14})$$

and for the delta-function constraints $\Delta(C, Z)$,

$$\Delta(C, Z) := \prod_{r=1,2} \delta^4 \left(\sum_{i \in \{A, B, 1, \dots, n\}} c_i^r Z_i \right). \quad (\text{C.15})$$

The pseudo-vacuum state (C.9) $|\Omega_n\rangle = \Delta(C_0, Z)$ corresponds to the following $2 \times (2n + 2)$ -matrix,

$$C_0 = \begin{pmatrix} 0 & 0 & 1 & 0 & \dots & 0 & 0 \\ 0 & 0 & 0 & 0 & \dots & 0 & 1 \end{pmatrix}. \quad (\text{C.16})$$

Each R-operator in the products (C.10) and (C.11) contains one-fold integration. Let us label the corresponding integration variables as t_1, t_2, \dots, t_{2n} from right to the left. Each shift $Z_i \rightarrow Z_i + tZ_j$ produced by R_{ij} induces the shift $\vec{c}_j \rightarrow \vec{c}_j + t\vec{c}_i$ in the matrix C_0 . Thus the sequence of R-operators in (C.10) and (C.11) brings us from C_0 to the top-cell of

$\text{Gr}(2, n + 2)$ parametrized with t_1, \dots, t_{2n} . The measure $\omega(C)$ (C.14) takes the following form

$$\omega(C) = \frac{(t_{2n-2}t_{2n-3})^3}{\prod_{i=1}^{2n-4} t_i} \bigwedge_{i=1}^{2n} dt_i. \quad (\text{C.17})$$

which agrees with values of spectral parameters carried by R-operators (C.1) in (C.10) and (C.11).

C.3 Cyclicity of the Yangian

The Yangian generators are entries of the monodromy matrix, which is a matrix product of the Lax operators. We will need the cyclic shift transformations of the Yangian generators, which we formulate in terms of the monodromy matrix.

If a homogeneous function $|\Psi\rangle$ of momentum twistors Z_1, \dots, Z_m is annihilated by the off-diagonal entries of the 4×4 matrix formed as the matrix product of m Lax operators (C.3) and the action of the diagonal entries results in an eigenvalue $\varphi(u, \delta)$,

$$\left[L_1^{(\delta_1)}(u_1) \dots L_{m-1}^{(\delta_{m-1})}(u_{m-1}) L_m^{(\delta_m)}(u_m) \right]^I{}_J |\Psi\rangle = \varphi(u, \delta) \delta_J^I |\Psi\rangle \quad (\text{C.18})$$

then the matrix product of the cyclically shifted Lax operators has a similar action on $|\Psi\rangle$

$$\left[L_m^{(\delta_m)}(u_m - 4) L_1^{(\delta_1)}(u_1) \dots L_{m-1}^{(\delta_{m-1})}(u_{m-1}) \right]^I{}_J |\Psi\rangle = \varphi'(u, \delta) \delta_J^I |\Psi\rangle \quad (\text{C.19})$$

with another eigenvalue $\varphi'(u, \delta)$,

$$\varphi'(u, \delta) = \varphi(u, \delta) \cdot \frac{(u_m - 4)(u_m - 1 + \delta_m)}{u_m(u_m - 1 + \delta_m) - \delta_m}. \quad (\text{C.20})$$

One can check this statement following [40].

C.4 Lemmas

Here we collect several technical statements which will be instrumental in section C.5. In the following we deal with multiple products of the dual-conformal generators and we adopt the matrix notations,

$$(\tilde{\mathfrak{J}}_a \tilde{\mathfrak{J}}_b)^I{}_J := (\tilde{\mathfrak{J}}_a)^I{}_K (\tilde{\mathfrak{J}}_b)^K{}_J, \quad (\tilde{\mathfrak{J}}_a \tilde{\mathfrak{J}}_b \tilde{\mathfrak{J}}_c)^I{}_J := (\tilde{\mathfrak{J}}_a)^I{}_K (\tilde{\mathfrak{J}}_b)^K{}_L (\tilde{\mathfrak{J}}_c)^L{}_J, \dots \quad (\text{C.21})$$

We observe that the products of several local dual-conformal generators $\tilde{\mathfrak{J}}_a$ (4.42) acting at the same spin-chain site labeled by a are proportional to $\tilde{\mathfrak{J}}_a$,

$$(\tilde{\mathfrak{J}}_a \tilde{\mathfrak{J}}_a)^I{}_J = (3 + H_a) (\tilde{\mathfrak{J}}_a)^I{}_J, \quad (\text{C.22})$$

and the proportionality coefficient can be considered as a number. Indeed, the dual-conformal weight counting operator H_a (4.40) commutes with the dual-conformal generators. The latter equation enables us to invert the Lax operator (C.3),

$$\left[L_a^{(\delta)}(-u - 3 - \delta) L_a^{(\delta)}(u) \right]^I{}_J = u(-u - 3 - \delta) \delta_J^I. \quad (\text{C.23})$$

Simplifications of the products of the dual-conformal generators $\tilde{\mathfrak{J}}_A$ and $\tilde{\mathfrak{J}}_B$ also appear when they act on functions of the twistor line $Z_A \wedge Z_B$,

$$Z_A \wedge Z_B : \quad (\tilde{\mathfrak{J}}_A \tilde{\mathfrak{J}}_B)^I{}_J = -(\tilde{\mathfrak{J}}_A)^I{}_J, \quad (\tilde{\mathfrak{J}}_B \tilde{\mathfrak{J}}_A)^I{}_J = -(\tilde{\mathfrak{J}}_B)^I{}_J. \quad (\text{C.24})$$

In particular, we will need the product of the dual-conformal transformations $\tilde{\mathfrak{J}}$ (4.43), which acts on the spin-chain sites A and B , restricted to this space,

$$Z_A \wedge Z_B : \quad (\tilde{\mathfrak{J}} \tilde{\mathfrak{J}})^I{}_J = -2\tilde{\mathfrak{J}}^I{}_J. \quad (\text{C.25})$$

A special role is played by the product of two R-operators which appear on the r.h.s. of (C.10), namely $\mathbb{R}_{AB}^{(-1)} \mathbb{R}_{BA}^{(-1)}$. In view of (C.6), the product is invariant under the dual-conformal transformations with $\tilde{\mathfrak{J}}$ (4.43),

$$\left[\mathbb{R}_{AB}^{(-1)} \mathbb{R}_{BA}^{(-1)}, \tilde{\mathfrak{J}}^I{}_J \right] = 0, \quad (\text{C.26})$$

and it does not alter the dual-conformal weight

$$\left[H_a, \mathbb{R}_{AB}^{(-1)} \mathbb{R}_{BA}^{(-1)} \right] = 0, \quad a = A, B. \quad (\text{C.27})$$

Given an arbitrary function $G(Z_A, Z_B)$ with equal dual-conformal weights w.r.t. both twistor points A and B ,

$$H_A G = H_B G, \quad (\text{C.28})$$

one can show that $\mathbb{R}_{AB}^{(-1)} \mathbb{R}_{BA}^{(-1)} G(Z_A, Z_B)$ is a function of the twistor line $Z_A \wedge Z_B$, namely it is invariant under shifts $Z_A \rightarrow Z_A + \epsilon Z_B$ and $Z_B \rightarrow Z_B + \epsilon Z_A$. Let us apply this observation to eq. (C.10). Indeed, $|\Phi_n\rangle$ depends on twistors Z_A and Z_B , and it carries equal dual-conformal weights at points A and B , see eq. (C.12). Then R_n given in (C.10) is a function of the line $Z_A \wedge Z_B$ in the twistor space.

The local exchange relations (C.4) and (C.5) result in the exchange relation satisfied by $\mathbb{R}_{AB}^{(-1)} \mathbb{R}_{BA}^{(-1)}$ with the product of Lax operators $L_A L_B$,

$$\mathbb{R}_{AB}^{(-1)} \mathbb{R}_{BA}^{(-1)} \left[L_A^{(-4)}(v+1) L_B^{(-4)}(v) \right]^I{}_J = \left[L_A^{(-4)}(v) L_B^{(-4)}(v+1) \right]^I{}_J \mathbb{R}_{AB}^{(-1)} \mathbb{R}_{BA}^{(-1)}. \quad (\text{C.29})$$

Expanding the latter equation in powers of the spectral parameter v , we reproduce the statement of the dual-conformal invariance (C.26) as well as the following exchange relation, which is quadratic in the dual-conformal generators (4.42),

$$\mathbb{R}_{AB}^{(-1)} \mathbb{R}_{BA}^{(-1)} [\tilde{\mathfrak{J}}_B + \tilde{\mathfrak{J}}_A \tilde{\mathfrak{J}}_B]^I{}_J = [\tilde{\mathfrak{J}}_A + \tilde{\mathfrak{J}}_A \tilde{\mathfrak{J}}_B]^I{}_J \mathbb{R}_{AB}^{(-1)} \mathbb{R}_{BA}^{(-1)}. \quad (\text{C.30})$$

In order to obtain more exchange relations which involve products of the dual-conformal generators, we contract the latter relation with the dual-conformal generators $\tilde{\mathfrak{J}}$ (4.43), which commute with the product of the R-operators, see (C.26),

$$\mathbb{R}_{AB}^{(-1)} \mathbb{R}_{BA}^{(-1)} [\tilde{\mathfrak{J}} \tilde{\mathfrak{J}}_B + \tilde{\mathfrak{J}} \tilde{\mathfrak{J}}_A \tilde{\mathfrak{J}}_B]^I{}_J = [\tilde{\mathfrak{J}} \tilde{\mathfrak{J}}_A + \tilde{\mathfrak{J}} \tilde{\mathfrak{J}}_A \tilde{\mathfrak{J}}_B]^I{}_J \mathbb{R}_{AB}^{(-1)} \mathbb{R}_{BA}^{(-1)}. \quad (\text{C.31})$$

In the following we work on the space of homogeneous functions carrying the dual-conformal weight equal to -4 . The products of the dual-conformal generators from (C.31) simplify as follows on this space

$$\mathcal{H}_A^{(-4)} \otimes \mathcal{H}_B^{(-4)} : \quad \begin{aligned} \bar{\mathfrak{J}} \mathfrak{J}_B + \bar{\mathfrak{J}} \mathfrak{J}_A \mathfrak{J}_B &= -\mathfrak{J}_B + \mathfrak{J}_B \mathfrak{J}_A \mathfrak{J}_B, \\ \bar{\mathfrak{J}} \mathfrak{J}_A + \bar{\mathfrak{J}} \mathfrak{J}_A \mathfrak{J}_B &= -\mathfrak{J}_A + \mathfrak{J}_B \mathfrak{J}_A - \mathfrak{J}_A \mathfrak{J}_B + \mathfrak{J}_B \mathfrak{J}_A \mathfrak{J}_B, \end{aligned} \quad (\text{C.32})$$

in view of (C.22). We take the sum of (C.30) and (C.31) and restrict the relation to the space of homogeneous functions, since the dual-conformal weights are not altered by $R_{AB}^{(-1)} R_{BA}^{(-1)}$, see (C.27),

$$\mathcal{H}_A^{(-4)} \otimes \mathcal{H}_B^{(-4)} : \quad R_{AB}^{(-1)} R_{BA}^{(-1)} [\mathfrak{J}_A \mathfrak{J}_B + \mathfrak{J}_B \mathfrak{J}_A \mathfrak{J}_B]^I{}_J = [\mathfrak{J}_B \mathfrak{J}_A + \mathfrak{J}_B \mathfrak{J}_A \mathfrak{J}_B]^I{}_J R_{AB}^{(-1)} R_{BA}^{(-1)} \quad (\text{C.33})$$

According to the discussion around eq. (C.28), the r.h.s. of (C.33) is a function of the twistor line $Z_A \wedge Z_B$. The product of the dual-conformal generators $\mathfrak{J}_B \mathfrak{J}_A + \mathfrak{J}_B \mathfrak{J}_A \mathfrak{J}_B$ from the r.h.s. of (C.33) vanishes on this space, see (C.24). Finally, after rewriting the product of the dual-conformal generators in the l.h.s. of (C.28),

$$\mathcal{H}_A^{(-4)} \otimes \mathcal{H}_B^{(-4)} : \quad \mathfrak{J}_A \mathfrak{J}_B + \mathfrak{J}_B \mathfrak{J}_A \mathfrak{J}_B = 2\mathfrak{J}_A \mathfrak{J}_B + \bar{\mathfrak{J}} \mathfrak{J}_A \mathfrak{J}_B, \quad (\text{C.34})$$

eq. (C.28) takes the following form

$$\mathcal{H}_A^{(-4)} \otimes \mathcal{H}_B^{(-4)} : \quad R_{AB}^{(-1)} R_{BA}^{(-1)} [2\mathfrak{J}_A \mathfrak{J}_B + \bar{\mathfrak{J}} \mathfrak{J}_A \mathfrak{J}_B]^I{}_J = 0. \quad (\text{C.35})$$

The relations established in this section are required in the next section.

C.5 Yangian invariances of R_n

We are going to establish how the Yangian generators from the monodromy matrix expansion (C.8) act on the leading singularities R_n . To this end, we form an auxiliary monodromy matrix out of the Lax operators (C.3) taken in the order $A, 1, 2, \dots, n, B$. The auxiliary monodromy matrix acts diagonally on the pseudo-vacuum state $|\Omega_n\rangle$ (C.9),

$$\left[L_A^{(0)}(u) L_1^{(-4)}(u) L_2^{(0)}(u) \dots L_{n-1}^{(0)}(u) L_n^{(-4)}(u+4) L_B^{(0)}(u) \right]^I{}_J |\Omega_n\rangle = u^n (u-1)(u+3) \delta_J^I |\Omega_n\rangle, \quad (\text{C.36})$$

since each of the constituent Lax operators acts diagonally on the delta-functions, see (C.3),

$$[L_i(u)]^I{}_J \delta^4(Z_j) = u \delta_J^I \delta^4(Z_j), \quad i \neq j \quad (\text{C.37})$$

$$[L_i(u)]^I{}_J \delta^4(Z_i) = (u-1) \delta_J^I \delta^4(Z_i). \quad (\text{C.38})$$

Then we act with $(2n-2)$ R-operators from (C.11) onto (C.36) and pull them through the products of Lax operators by means of the local exchange relations (C.4) and (C.5),

$$\left[L_A^{(-4)}(u) L_1^{(0)}(u) L_2^{(0)}(u) \dots L_{n-1}^{(0)}(u) L_n^{(0)}(u) L_B^{(-4)}(u+4) \right]^I{}_J |\Phi_n\rangle = u^n (u-1)(u+3) \delta_J^I |\Phi_n\rangle. \quad (\text{C.39})$$

We apply the cyclicity of appendix C.3 to the previous monodromy relation cyclically shifting the ordering of the Lax operators from $A, 1, 2, \dots, n, B$ to $B, A, 1, 2, \dots, n$ and replacing the monodromy matrix $L_1 \dots L_n$ with the shorthand notation (C.7),

$$\left[L_B^{(-4)}(u) L_A^{(-4)}(u) T(u) \right]^I | \Phi_n \rangle = u^n (u-1)^2 \delta_J^I | \Phi_n \rangle, \quad (\text{C.40})$$

and we invert L_A and L_B on the l.h.s. of the previous equation by means of (C.23),

$$[T(u)]^I | \Phi_n \rangle = u^{n-2} \left[L_A^{(-4)}(1-u) L_B^{(-4)}(1-u) \right]^I | \Phi_n \rangle. \quad (\text{C.41})$$

We need to act with $R_{AB}^{(-1)} R_{BA}^{(-1)}$ on both sides of (C.41) in order to promote $|\Phi_n\rangle$ to the leading singularities R_n according to (C.10). This is straightforward in the l.h.s. of (C.41). If we could pull this R-operator product through $L_A L_B$ in the r.h.s. of (C.41) then we would find that R_n is invariant under all Yangian generators. However, the latter is not true. The obstacle is that values of spectral parameters in $L_A L_B$ in the r.h.s. of (C.41) are not adjusted like in (C.29).

In order to obtain symmetry relations for R_n , we expand (C.41) in the spectral parameter taking into account definitions in (C.8), (4.43) and (4.46),

$$u^{n-1} : \quad \mathfrak{J}^I | \Phi_n \rangle = -\tilde{\mathfrak{J}}^I | \Phi_n \rangle - 2\delta_J^I | \Phi_n \rangle, \quad (\text{C.42})$$

$$u^{n-2} : \quad \hat{\mathfrak{J}}^I | \Phi_n \rangle = \delta_J^I | \Phi_n \rangle + \tilde{\mathfrak{J}}^I | \Phi_n \rangle + (\tilde{\mathfrak{J}}_A \tilde{\mathfrak{J}}_B)^I | \Phi_n \rangle, \quad (\text{C.43})$$

$$u^{n-1-k} : \quad \left(\hat{\mathfrak{J}}^{(k)} \right)^I | \Phi_n \rangle = 0, \quad k = 2, \dots, n-1. \quad (\text{C.44})$$

We act with $R_{AB}^{(-1)} R_{BA}^{(-1)}$ onto (C.42) and (C.44) and replace there $|\Phi_n\rangle$ with R_n according to (C.10). Indeed, the R-operators are dual-conformal invariant, see eq. (C.26). Thus, (C.42) is equivalent to the dual-conformal invariance of the leading singularities R_n (C.6), while (C.44) is equivalent to the Yangian invariance of R_n with the level-two and higher-level Yangian generators (4.47). Only (C.43) cannot be immediately interpreted as the Yangian invariance relation, since $R_{AB}^{(-1)} R_{BA}^{(-1)}$ does not commute nicely with $\tilde{\mathfrak{J}}_A \tilde{\mathfrak{J}}_B$ in the last term on the r.h.s. of (C.43).

Upon contraction of (C.43) with the tensor $\langle Z_A, Z_B, *, \star \rangle$ the unwanted terms drop out,

$$\epsilon_{IKLM} Z_A^K Z_B^L \hat{\mathfrak{J}}^I | \Phi_n \rangle = \epsilon_{JKLM} Z_A^K Z_B^L | \Phi_n \rangle, \quad (\text{C.45})$$

and the latter relation is equivalent to (4.48) since both R_{AB} and R_{BA} , which act as shifts on twistors Z_A and Z_B , commute with $\langle Z_A, Z_B, *, \star \rangle$.

Another way to acquire a symmetry relation with the level-one Yangian generators $\hat{\mathfrak{J}}$ from (C.43) is to cook up a polynomial in the local dual-conformal generators $\tilde{\mathfrak{J}}_A$ and $\tilde{\mathfrak{J}}_B$ which nicely commutes with $R_{AB}^{(-1)} R_{BA}^{(-1)}$. We have established in (C.35) that this is the case for $\tilde{\mathfrak{J}} \tilde{\mathfrak{J}}_A \tilde{\mathfrak{J}}_B + 2\tilde{\mathfrak{J}}_A \tilde{\mathfrak{J}}_B$. Indeed, contracting (C.43) with $\tilde{\mathfrak{J}}^K | \Phi_n \rangle + 2\delta_J^K | \Phi_n \rangle$ we organize the unwanted terms into this polynomial,

$$\left(\tilde{\mathfrak{J}} \hat{\mathfrak{J}} + 2\hat{\mathfrak{J}} \right)^I | \Phi_n \rangle = 2\delta_J^I | \Phi_n \rangle + 3\tilde{\mathfrak{J}}^I | \Phi_n \rangle + \left(\tilde{\mathfrak{J}} \tilde{\mathfrak{J}} \right)^I | \Phi_n \rangle + \left(\tilde{\mathfrak{J}} \tilde{\mathfrak{J}}_A \tilde{\mathfrak{J}}_B + 2\tilde{\mathfrak{J}}_A \tilde{\mathfrak{J}}_B \right)^I | \Phi_n \rangle. \quad (\text{C.46})$$

We act on the previous relation with $R_{AB}^{(-1)}R_{BA}^{(-1)}$ pulling it through $\tilde{\mathfrak{J}}$ (C.26), replacing $\tilde{\mathfrak{J}}\tilde{\mathfrak{J}} \rightarrow -2\tilde{\mathfrak{J}}$ according to (C.25), and throwing out the last term on the r.h.s. in view of (C.35),

$$\left(\tilde{\mathfrak{J}}\hat{\mathfrak{J}} + 2\hat{\mathfrak{J}}\right)^I{}_J R_n = 2\delta^I{}_J R_n + \tilde{\mathfrak{J}}^I{}_J R_n. \quad (\text{C.47})$$

The latter is the symmetry relation (4.49).

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited. SCOAP³ supports the goals of the International Year of Basic Sciences for Sustainable Development.

References

- [1] O. Almeliid, C. Duhr and E. Gardi, *Three-loop corrections to the soft anomalous dimension in multileg scattering*, *Phys. Rev. Lett.* **117** (2016) 172002 [[arXiv:1507.00047](https://arxiv.org/abs/1507.00047)] [[INSPIRE](#)].
- [2] L.F. Alday and J.M. Maldacena, *Gluon scattering amplitudes at strong coupling*, *JHEP* **06** (2007) 064 [[arXiv:0705.0303](https://arxiv.org/abs/0705.0303)] [[INSPIRE](#)].
- [3] J.M. Drummond, J. Henn, G.P. Korchemsky and E. Sokatchev, *Conformal Ward identities for Wilson loops and a test of the duality with gluon amplitudes*, *Nucl. Phys. B* **826** (2010) 337 [[arXiv:0712.1223](https://arxiv.org/abs/0712.1223)] [[INSPIRE](#)].
- [4] J.M. Drummond, J.M. Henn and J. Plefka, *Yangian symmetry of scattering amplitudes in $N = 4$ super Yang-Mills theory*, *JHEP* **05** (2009) 046 [[arXiv:0902.2987](https://arxiv.org/abs/0902.2987)] [[INSPIRE](#)].
- [5] N. Arkani-Hamed, F. Cachazo, C. Cheung and J. Kaplan, *A Duality For The S Matrix*, *JHEP* **03** (2010) 020 [[arXiv:0907.5418](https://arxiv.org/abs/0907.5418)] [[INSPIRE](#)].
- [6] N. Arkani-Hamed, J.L. Bourjaily, F. Cachazo, A.B. Goncharov, A. Postnikov and J. Trnka, *Grassmannian Geometry of Scattering Amplitudes*, Cambridge University Press, Cambridge, U.K. (2016) [[DOI](#)] [[arXiv:1212.5605](https://arxiv.org/abs/1212.5605)] [[INSPIRE](#)].
- [7] N. Arkani-Hamed, J.L. Bourjaily, F. Cachazo, S. Caron-Huot and J. Trnka, *The All-Loop Integrand For Scattering Amplitudes in Planar $N = 4$ SYM*, *JHEP* **01** (2011) 041 [[arXiv:1008.2958](https://arxiv.org/abs/1008.2958)] [[INSPIRE](#)].
- [8] N. Arkani-Hamed and J. Trnka, *The Amplituhedron*, *JHEP* **10** (2014) 030 [[arXiv:1312.2007](https://arxiv.org/abs/1312.2007)] [[INSPIRE](#)].
- [9] J.L. Bourjaily, A. DiRe, A. Shaikh, M. Spradlin and A. Volovich, *The Soft-Collinear Bootstrap: $N = 4$ Yang-Mills Amplitudes at Six and Seven Loops*, *JHEP* **03** (2012) 032 [[arXiv:1112.6432](https://arxiv.org/abs/1112.6432)] [[INSPIRE](#)].
- [10] J.L. Bourjaily, P. Heslop and V.-V. Tran, *Amplitudes and Correlators to Ten Loops Using Simple, Graphical Bootstraps*, *JHEP* **11** (2016) 125 [[arXiv:1609.00007](https://arxiv.org/abs/1609.00007)] [[INSPIRE](#)].
- [11] B. Eden, P. Heslop, G.P. Korchemsky and E. Sokatchev, *Hidden symmetry of four-point correlation functions and amplitudes in $N = 4$ SYM*, *Nucl. Phys. B* **862** (2012) 193 [[arXiv:1108.3557](https://arxiv.org/abs/1108.3557)] [[INSPIRE](#)].

- [12] B. Eden, P. Heslop, G.P. Korchemsky and E. Sokatchev, *Constructing the correlation function of four stress-tensor multiplets and the four-particle amplitude in $N = 4$ SYM*, *Nucl. Phys. B* **862** (2012) 450 [[arXiv:1201.5329](#)] [[INSPIRE](#)].
- [13] R.G. Ambrosio, B. Eden, T. Goddard, P. Heslop and C. Taylor, *Local integrands for the five-point amplitude in planar $N = 4$ SYM up to five loops*, *JHEP* **01** (2015) 116 [[arXiv:1312.1163](#)] [[INSPIRE](#)].
- [14] L.J. Dixon, J.M. Drummond and J.M. Henn, *Bootstrapping the three-loop hexagon*, *JHEP* **11** (2011) 023 [[arXiv:1108.4461](#)] [[INSPIRE](#)].
- [15] J.M. Drummond, G. Papathanasiou and M. Spradlin, *A Symbol of Uniqueness: The Cluster Bootstrap for the 3-Loop MHV Heptagon*, *JHEP* **03** (2015) 072 [[arXiv:1412.3763](#)] [[INSPIRE](#)].
- [16] A.B. Goncharov, M. Spradlin, C. Vergu and A. Volovich, *Classical Polylogarithms for Amplitudes and Wilson Loops*, *Phys. Rev. Lett.* **105** (2010) 151605 [[arXiv:1006.5703](#)] [[INSPIRE](#)].
- [17] J. Golden, A.B. Goncharov, M. Spradlin, C. Vergu and A. Volovich, *Motivic Amplitudes and Cluster Coordinates*, *JHEP* **01** (2014) 091 [[arXiv:1305.1617](#)] [[INSPIRE](#)].
- [18] S. Caron-Huot, L.J. Dixon, A. McLeod and M. von Hippel, *Bootstrapping a Five-Loop Amplitude Using Steinmann Relations*, *Phys. Rev. Lett.* **117** (2016) 241601 [[arXiv:1609.00669](#)] [[INSPIRE](#)].
- [19] S. Caron-Huot, L.J. Dixon, F. Dulat, M. Von Hippel, A.J. McLeod and G. Papathanasiou, *The Cosmic Galois Group and Extended Steinmann Relations for Planar $\mathcal{N} = 4$ SYM Amplitudes*, *JHEP* **09** (2019) 061 [[arXiv:1906.07116](#)] [[INSPIRE](#)].
- [20] S. Caron-Huot et al., *The Steinmann Cluster Bootstrap for $N = 4$ Super Yang-Mills Amplitudes*, *PoS CORFU2019* (2020) 003 [[arXiv:2005.06735](#)] [[INSPIRE](#)].
- [21] J.M. Henn, *What Can We Learn About QCD and Collider Physics from $N = 4$ Super Yang-Mills?*, *Ann. Rev. Nucl. Part. Sci.* **71** (2021) 87 [[arXiv:2006.00361](#)] [[INSPIRE](#)].
- [22] J.M. Drummond, J. Henn, G.P. Korchemsky and E. Sokatchev, *Dual superconformal symmetry of scattering amplitudes in $N = 4$ super-Yang-Mills theory*, *Nucl. Phys. B* **828** (2010) 317 [[arXiv:0807.1095](#)] [[INSPIRE](#)].
- [23] S. Weinzierl, *Does one need the $O(\epsilon)$ - and $O(\epsilon^2)$ -terms of one-loop amplitudes in an NNLO calculation?*, *Phys. Rev. D* **84** (2011) 074007 [[arXiv:1107.5131](#)] [[INSPIRE](#)].
- [24] C. Anastasiou and G. Sterman, *Removing infrared divergences from two-loop integrals*, *JHEP* **07** (2019) 056 [[arXiv:1812.03753](#)] [[INSPIRE](#)].
- [25] L.F. Alday, P. Heslop and J. Sikorowski, *Perturbative correlation functions of null Wilson loops and local operators*, *JHEP* **03** (2013) 074 [[arXiv:1207.4316](#)] [[INSPIRE](#)].
- [26] O. Erdoğan and G. Sterman, *Gauge Theory Webs and Surfaces*, *Phys. Rev. D* **91** (2015) 016003 [[arXiv:1112.4564](#)] [[INSPIRE](#)].
- [27] L.F. Alday, J.M. Henn and J. Sikorowski, *Higher loop mixed correlators in $N = 4$ SYM*, *JHEP* **03** (2013) 058 [[arXiv:1301.0149](#)] [[INSPIRE](#)].
- [28] J.M. Henn, G.P. Korchemsky and B. Mistlberger, *The full four-loop cusp anomalous dimension in $\mathcal{N} = 4$ super Yang-Mills and QCD*, *JHEP* **04** (2020) 018 [[arXiv:1911.10174](#)] [[INSPIRE](#)].
- [29] N. Arkani-Hamed and J. Trnka, *Into the Amplituhedron*, *JHEP* **12** (2014) 182 [[arXiv:1312.7878](#)] [[INSPIRE](#)].

- [30] N. Arkani-Hamed, J. Henn and J. Trnka, *Nonperturbative negative geometries: amplitudes at strong coupling and the amplituhedron*, *JHEP* **03** (2022) 108 [[arXiv:2112.06956](#)] [[INSPIRE](#)].
- [31] D. Chicherin and J. Henn, *Pentagon Wilson loop with Lagrangian insertion at two loops in $\mathcal{N} = 4$ super Yang-Mills theory*, [arXiv:2204.00329](#) [[INSPIRE](#)].
- [32] J.M. Drummond and L. Ferro, *Yangians, Grassmannians and T-duality*, *JHEP* **07** (2010) 027 [[arXiv:1001.3348](#)] [[INSPIRE](#)].
- [33] J.M. Drummond and L. Ferro, *The Yangian origin of the Grassmannian integral*, *JHEP* **12** (2010) 010 [[arXiv:1002.4622](#)] [[INSPIRE](#)].
- [34] G.P. Korchemsky and E. Sokatchev, *Superconformal invariants for scattering amplitudes in $N = 4$ SYM theory*, *Nucl. Phys. B* **839** (2010) 377 [[arXiv:1002.4625](#)] [[INSPIRE](#)].
- [35] L. Ferro, T. Lukowski, A. Orta and M. Parisi, *Towards the Amplituhedron Volume*, *JHEP* **03** (2016) 014 [[arXiv:1512.04954](#)] [[INSPIRE](#)].
- [36] L. Ferro, T. Łukowski, A. Orta and M. Parisi, *Yangian symmetry for the tree amplituhedron*, *J. Phys. A* **50** (2017) 294005 [[arXiv:1612.04378](#)] [[INSPIRE](#)].
- [37] Y. Bai, S. He and T. Lam, *The Amplituhedron and the One-loop Grassmannian Measure*, *JHEP* **01** (2016) 112 [[arXiv:1510.03553](#)] [[INSPIRE](#)].
- [38] L.D. Faddeev, *How algebraic Bethe ansatz works for integrable model*, in *Les Houches School of Physics: Astrophysical Sources of Gravitational Radiation*, Les Houches, France, September 26–October 6, 1995, pp. 149–219 (1996) [[hep-th/9605187](#)] [[INSPIRE](#)].
- [39] E.K. Sklyanin, *Quantum inverse scattering method. Selected topics*, [hep-th/9211111](#) [[INSPIRE](#)].
- [40] D. Chicherin, S. Derkachov and R. Kirschner, *Yang-Baxter operators and scattering amplitudes in $N = 4$ super-Yang-Mills theory*, *Nucl. Phys. B* **881** (2014) 467 [[arXiv:1309.5748](#)] [[INSPIRE](#)].
- [41] R. Frassek, N. Kanning, Y. Ko and M. Staudacher, *Bethe Ansatz for Yangian Invariants: Towards Super Yang-Mills Scattering Amplitudes*, *Nucl. Phys. B* **883** (2014) 373 [[arXiv:1312.1693](#)] [[INSPIRE](#)].
- [42] N. Kanning, T. Lukowski and M. Staudacher, *A shortcut to general tree-level scattering amplitudes in $\mathcal{N} = 4$ SYM via integrability*, *Fortsch. Phys.* **62** (2014) 556 [[arXiv:1403.3382](#)] [[INSPIRE](#)].
- [43] J. Broedel, M. de Leeuw and M. Rosso, *A dictionary between R-operators, on-shell graphs and Yangian algebras*, *JHEP* **06** (2014) 170 [[arXiv:1403.3670](#)] [[INSPIRE](#)].
- [44] T. Bargheer, Y.-t. Huang, F. Loebbert and M. Yamazaki, *Integrable Amplitude Deformations for $N = 4$ Super Yang-Mills and ABJM Theory*, *Phys. Rev. D* **91** (2015) 026004 [[arXiv:1407.4449](#)] [[INSPIRE](#)].
- [45] L.V. Bork and A.I. Onishchenko, *On soft theorems and form factors in $\mathcal{N} = 4$ SYM theory*, *JHEP* **12** (2015) 030 [[arXiv:1506.07551](#)] [[INSPIRE](#)].
- [46] R. Frassek, D. Meidinger, D. Nandan and M. Wilhelm, *On-shell diagrams, Grassmannians and integrability for form factors*, *JHEP* **01** (2016) 182 [[arXiv:1506.08192](#)] [[INSPIRE](#)].
- [47] L.V. Bork and A.I. Onishchenko, *Wilson lines, Grassmannians and gauge invariant off-shell amplitudes in $\mathcal{N} = 4$ SYM*, *JHEP* **04** (2017) 019 [[arXiv:1607.02320](#)] [[INSPIRE](#)].

- [48] J. Fuksa and R. Kirschner, *Correlators with sl_2 Yangian symmetry*, *Nucl. Phys. B* **914** (2017) 1 [[arXiv:1608.04912](#)] [[INSPIRE](#)].
- [49] R. Kirschner and G. Savvidy, *Yangian and SUSY symmetry of high spin parton splitting amplitudes in generalised Yang-Mills theory*, *Mod. Phys. Lett. A* **32** (2017) 1750121 [[arXiv:1701.06660](#)] [[INSPIRE](#)].
- [50] D. Chicherin, V. Kazakov, F. Loebbert, D. Müller and D.-l. Zhong, *Yangian Symmetry for Bi-Scalar Loop Amplitudes*, *JHEP* **05** (2018) 003 [[arXiv:1704.01967](#)] [[INSPIRE](#)].
- [51] D. Chicherin, V. Kazakov, F. Loebbert, D. Müller and D.-l. Zhong, *Yangian Symmetry for Fishnet Feynman Graphs*, *Phys. Rev. D* **96** (2017) 121901 [[arXiv:1708.00007](#)] [[INSPIRE](#)].
- [52] S. Badger et al., *Analytic form of the full two-loop five-gluon all-plus helicity amplitude*, *Phys. Rev. Lett.* **123** (2019) 071601 [[arXiv:1905.03733](#)] [[INSPIRE](#)].
- [53] J. Henn, B. Power and S. Zoia, *Conformal Invariance of the One-Loop All-Plus Helicity Scattering Amplitudes*, *JHEP* **02** (2020) 019 [[arXiv:1911.12142](#)] [[INSPIRE](#)].
- [54] D.C. Dunbar, G.R. Jehu and W.B. Perkins, *The two-loop n -point all-plus helicity amplitude*, *Phys. Rev. D* **93** (2016) 125006 [[arXiv:1604.06631](#)] [[INSPIRE](#)].
- [55] R. Ben-Israel, A.G. Tumanov and A. Sever, *Scattering amplitudes — Wilson loops duality for the first non-planar correction*, *JHEP* **08** (2018) 122 [[arXiv:1802.09395](#)] [[INSPIRE](#)].
- [56] J.M. Drummond, G.P. Korchemsky and E. Sokatchev, *Conformal properties of four-gluon planar amplitudes and Wilson loops*, *Nucl. Phys. B* **795** (2008) 385 [[arXiv:0707.0243](#)] [[INSPIRE](#)].
- [57] A. Brandhuber, P. Heslop and G. Travaglini, *MHV amplitudes in $N = 4$ super Yang-Mills and Wilson loops*, *Nucl. Phys. B* **794** (2008) 231 [[arXiv:0707.1153](#)] [[INSPIRE](#)].
- [58] Z. Bern et al., *The Two-Loop Six-Gluon MHV Amplitude in Maximally Supersymmetric Yang-Mills Theory*, *Phys. Rev. D* **78** (2008) 045007 [[arXiv:0803.1465](#)] [[INSPIRE](#)].
- [59] L.F. Alday and R. Roiban, *Scattering Amplitudes, Wilson Loops and the String/Gauge Theory Correspondence*, *Phys. Rept.* **468** (2008) 153 [[arXiv:0807.1889](#)] [[INSPIRE](#)].
- [60] Z. Bern, L.J. Dixon and V.A. Smirnov, *Iteration of planar amplitudes in maximally supersymmetric Yang-Mills theory at three loops and beyond*, *Phys. Rev. D* **72** (2005) 085001 [[hep-th/0505205](#)] [[INSPIRE](#)].
- [61] D. Chicherin et al., *Correlation functions of the chiral stress-tensor multiplet in $\mathcal{N} = 4$ SYM*, *JHEP* **06** (2015) 198 [[arXiv:1412.8718](#)] [[INSPIRE](#)].
- [62] N. Berkovits and J. Maldacena, *Fermionic T -duality, Dual Superconformal Symmetry, and the Amplitude/Wilson Loop Connection*, *JHEP* **09** (2008) 062 [[arXiv:0807.3196](#)] [[INSPIRE](#)].
- [63] B. Eden, P. Heslop, G.P. Korchemsky and E. Sokatchev, *The super-correlator/super-amplitude duality: Part I*, *Nucl. Phys. B* **869** (2013) 329 [[arXiv:1103.3714](#)] [[INSPIRE](#)].
- [64] B. Eden, G.P. Korchemsky and E. Sokatchev, *From correlation functions to scattering amplitudes*, *JHEP* **12** (2011) 002 [[arXiv:1007.3246](#)] [[INSPIRE](#)].
- [65] B. Eden, G.P. Korchemsky and E. Sokatchev, *More on the duality correlators/amplitudes*, *Phys. Lett. B* **709** (2012) 247 [[arXiv:1009.2488](#)] [[INSPIRE](#)].
- [66] B. Eden, P. Heslop, G.P. Korchemsky and E. Sokatchev, *The super-correlator/super-amplitude duality: Part II*, *Nucl. Phys. B* **869** (2013) 378 [[arXiv:1103.4353](#)] [[INSPIRE](#)].

- [67] R.A. Brandt, F. Neri and M.-a. Sato, *Renormalization of Loop Functions for All Loops*, *Phys. Rev. D* **24** (1981) 879 [INSPIRE].
- [68] I.A. Korchemskaya and G.P. Korchemsky, *On lightlike Wilson loops*, *Phys. Lett. B* **287** (1992) 169 [INSPIRE].
- [69] O.T. Engelund and R. Roiban, *On correlation functions of Wilson loops, local and non-local operators*, *JHEP* **05** (2012) 158 [arXiv:1110.0758] [INSPIRE].
- [70] A. Hodges, *Eliminating spurious poles from gauge-theoretic amplitudes*, *JHEP* **05** (2013) 135 [arXiv:0905.1473] [INSPIRE].
- [71] L.J. Mason and D. Skinner, *Dual Superconformal Invariance, Momentum Twistors and Grassmannians*, *JHEP* **11** (2009) 045 [arXiv:0909.0250] [INSPIRE].
- [72] Z. Bern, L.J. Dixon, D.C. Dunbar and D.A. Kosower, *One loop n point gauge theory amplitudes, unitarity and collinear limits*, *Nucl. Phys. B* **425** (1994) 217 [hep-ph/9403226] [INSPIRE].
- [73] N. Arkani-Hamed, J.L. Bourjaily, F. Cachazo and J. Trnka, *Local Integrals for Planar Scattering Amplitudes*, *JHEP* **06** (2012) 125 [arXiv:1012.6032] [INSPIRE].
- [74] D. Chicherin, J.M. Henn and A. Tumanov, in preparation.
- [75] E. Witten, *Perturbative gauge theory as a string theory in twistor space*, *Commun. Math. Phys.* **252** (2004) 189 [hep-th/0312171] [INSPIRE].
- [76] V.G. Drinfeld, *Hopf algebras and the quantum Yang-Baxter equation*, *Sov. Math. Dokl.* **32** (1985) 254 [INSPIRE].
- [77] A. Molev, M. Nazarov and G. Olshansky, *Yangians and classical Lie algebras*, *Russ. Math. Surveys* **51** (1996) 205 [hep-th/9409025] [INSPIRE].
- [78] L. Ferro, T. Łukowski and M. Staudacher, *$\mathcal{N} = 4$ scattering amplitudes and the deformed Grassmannian*, *Nucl. Phys. B* **889** (2014) 192 [arXiv:1407.6736] [INSPIRE].
- [79] Z. Bern, G. Chalmers, L.J. Dixon and D.A. Kosower, *One loop N gluon amplitudes with maximal helicity violation via collinear limits*, *Phys. Rev. Lett.* **72** (1994) 2134 [hep-ph/9312333] [INSPIRE].
- [80] C.R. Mafra and O. Schlotterer, *The Structure of n -Point One-Loop Open Superstring Amplitudes*, *JHEP* **08** (2014) 099 [arXiv:1203.6215] [INSPIRE].
- [81] S. He, R. Monteiro and O. Schlotterer, *String-inspired BCJ numerators for one-loop MHV amplitudes*, *JHEP* **01** (2016) 171 [arXiv:1507.06288] [INSPIRE].
- [82] S. Catani, *The Singular behavior of QCD amplitudes at two loop order*, *Phys. Lett. B* **427** (1998) 161 [hep-ph/9802439] [INSPIRE].
- [83] Z. Bern, A. De Freitas and L.J. Dixon, *Two loop helicity amplitudes for gluon-gluon scattering in QCD and supersymmetric Yang-Mills theory*, *JHEP* **03** (2002) 018 [hep-ph/0201161] [INSPIRE].
- [84] Q. Jin and H. Lüo, *Analytic Form of the Three-loop Four-gluon Scattering Amplitudes in Yang-Mills Theory*, arXiv:1910.05889 [INSPIRE].
- [85] F. Caola, A. Chakraborty, G. Gambuti, A. von Manteuffel and L. Tancredi, *Three-Loop Gluon Scattering in QCD and the Gluon Regge Trajectory*, *Phys. Rev. Lett.* **128** (2022) 212001 [arXiv:2112.11097] [INSPIRE].

- [86] A.V. Kotikov and L.N. Lipatov, *DGLAP and BFKL equations in the $N = 4$ supersymmetric gauge theory*, *Nucl. Phys. B* **661** (2003) 19 [Erratum *ibid.* **685** (2004) 405] [[hep-ph/0208220](#)] [[INSPIRE](#)].
- [87] C. Anastasiou, Z. Bern, L.J. Dixon and D.A. Kosower, *Planar amplitudes in maximally supersymmetric Yang-Mills theory*, *Phys. Rev. Lett.* **91** (2003) 251602 [[hep-th/0309040](#)] [[INSPIRE](#)].
- [88] Z. Bern, L.J. Dixon, D.C. Dunbar and D.A. Kosower, *One loop selfdual and $N = 4$ superYang-Mills*, *Phys. Lett. B* **394** (1997) 105 [[hep-th/9611127](#)] [[INSPIRE](#)].
- [89] S. Badger, H. Frellesvig and Y. Zhang, *A Two-Loop Five-Gluon Helicity Amplitude in QCD*, *JHEP* **12** (2013) 045 [[arXiv:1310.1051](#)] [[INSPIRE](#)].
- [90] S. Badger, G. Mogull and T. Peraro, *Local integrands for two-loop all-plus Yang-Mills amplitudes*, *JHEP* **08** (2016) 063 [[arXiv:1606.02244](#)] [[INSPIRE](#)].
- [91] B. Mistlberger, *Higgs boson production at hadron colliders at N^3LO in QCD*, *JHEP* **05** (2018) 028 [[arXiv:1802.00833](#)] [[INSPIRE](#)].
- [92] N. Arkani-Hamed, D. Chicherin, J.M. Henn and J. Trnka, in preparation.
- [93] S. Caron-Huot and S. He, *Jumpstarting the All-Loop S-matrix of Planar $N = 4$ Super Yang-Mills*, *JHEP* **07** (2012) 174 [[arXiv:1112.1060](#)] [[INSPIRE](#)].