

3d dualities from 4d dualities for orthogonal groups

Ofer Aharony,^{a,b} Shlomo S. Razamat,^a Nathan Seiberg^a and Brian Willett^a

^a*School of Natural Sciences, Institute for Advanced Study,
Princeton, NJ 08540, U.S.A.*

^b*Department of Particle Physics and Astrophysics, Weizmann Institute of Science,
Rehovot 76100, Israel*

E-mail: Ofer.Aharony@weizmann.ac.il, razamat@ias.edu, seiberg@ias.edu,
bwillett@ias.edu

ABSTRACT: We extend recent work on the relation of 4d and 3d IR dualities of supersymmetric gauge theories with four supercharges to the case of orthogonal gauge groups. The distinction between different $SO(N)$ gauge theories in 4d plays an important role in this relation. We show that the 4d duality leads to a 3d duality between an $SO(N_c)$ gauge theory with N_f flavors and an $SO(N_f - N_c + 2)$ theory with N_f flavors and extra singlets, and we derive its generalization in the presence of Chern-Simons terms. There are two different $O(N)$ theories in 3d, which we denote by $O(N)_{\pm}$, and we also show that the $O(N_c)_{-}$ gauge theory is dual to a $Spin(N_f - N_c + 2)$ theory, and derive from 4d the known duality between $O(N_c)_{+}$ and $O(N_f - N_c + 2)_{+}$. We verify the consistency of these 3d dualities by various methods, including index computations.

KEYWORDS: Supersymmetry and Duality, Duality in Gauge Field Theories, Supersymmetric gauge theory, Field Theories in Lower Dimensions

ARXIV EPRINT: [1307.0511](https://arxiv.org/abs/1307.0511)

Contents

1	Introduction	1
2	Background	4
2.1	Monopole operators and Coulomb branch coordinates for $g = so(N_c)$	4
2.2	The quantum moduli space of $3d \mathcal{N} = 2$ theories with $g = so(N_c)$	8
2.3	Baryon-monopole operators	9
2.4	On $3d O(N_c)$ theories	11
3	The Coulomb branch of $4d so(N_c)$ theories on \mathbb{S}^1	11
3.1	$4d \text{Spin}(N_c)$ theories on \mathbb{S}^1	11
3.2	$4d \text{SO}(N_c)$ theories on \mathbb{S}^1	14
4	The $4d$ duality on a circle	16
4.1	Dual theories on a circle	16
4.2	A test of the duality and of the Coulomb branch mapping	17
5	$3d$ dualities	20
5.1	Reduction of the $\text{SO}(N)_+$ duality to $3d$	20
5.2	Reduction of the $\text{Spin}(N_c) \leftrightarrow \text{SO}(\tilde{N}_c)_-$ duality to $3d$	23
5.3	Dualities with Chern-Simons terms	24
5.4	The special case of $\text{SO}(2) = \text{U}(1)$ with $N_f = 2$: a triality of dualities	26
6	Partition functions and indices for $so(N)$ dualities	29
6.1	The partition function on $\mathbb{S}^2 \times \mathbb{S}^1$	29
6.2	The partition functions on \mathbb{S}^3	34

1 Introduction

A crucial role in gauge theory dynamics is played by various dualities. They give a weakly coupled description of some strong coupling phenomena (like confinement and chiral symmetry breaking), and may point to a deep structure underlying the theory. In different situations these dualities manifest themselves differently. Some $4d$ superconformal theories like $\mathcal{N} = 4$ and certain $\mathcal{N} = 2$ supersymmetric theories exhibit exact electric/magnetic duality, leading to several distinct descriptions of the same theory, with different values of the coupling constant and sometimes even different gauge groups. Many four-dimensional $\mathcal{N} = 1, 2$, three-dimensional $\mathcal{N} = 2, 3, \dots$, and certain two-dimensional theories have IR dualities, relating different theories with the same IR limit [1]. In some situations, including $4d \mathcal{N} = 1 \text{SO}(N)$ dualities, it is clear that these are also related to electric/magnetic

duality [1–3]; when the gauge group is broken to $SO(2)$ they reduce to an Abelian electric/magnetic duality, and they exchange Wilson lines with 't Hooft-Wilson lines [4, 5].

In $3d$ there are several known examples of such IR dualities, both with $\mathcal{N} = 2$ and with higher supersymmetries. In a previous paper we argued that most and perhaps all such dualities in $3d$ originate from ancestor dualities in $4d$ [6] (see also [7]). The purpose of this note is to extend this discussion to theories with orthogonal gauge groups.

The discussion in [6] starts with any $4d$ $\mathcal{N} = 1$ duality, and by carefully compactifying it on a circle, it leads to a clear prescription for how to generate from it a corresponding $3d$ duality. For example, we can start with the characteristic example of a $4d$ $\mathcal{N} = 1$ duality. This is the duality between an $SU(N_c)$ gauge theory with N_f flavors Q^i and $\tilde{Q}_{\tilde{i}}$, and its dual $SU(N_f - N_c)$ gauge theory with N_f dual quarks q_i and $\tilde{q}^{\tilde{i}}$ and elementary gauge neutral “mesons” $M_{\tilde{i}}^i$ and a superpotential $W = M_{\tilde{i}}^i q_i \tilde{q}^{\tilde{i}}$ [1]. It is common to refer to these theories as the electric and the magnetic theories, but we will refer to them as theory A and theory B. A naive dimensional reduction of any of these two dual theories to $3d$ leads to a theory with an additional “axial” $U(1)$ global symmetry. This is the symmetry that is anomalous in $4d$, but is preserved in $3d$. The prescription of [6] is to modify the naive dimensionally reduced theory by adding to its Lagrangian a suitable operator, generated by non-perturbative effects in the theory on a circle, which explicitly breaks this anomalous $U(1)$ symmetry. In theory A we add a superpotential

$$W_A = \eta Y, \tag{1.1}$$

where $\eta = \Lambda^{b_0}$ is the instanton factor [3] of theory A, and Y is its monopole operator. In theory B, which already had a superpotential in $4d$, we have

$$W_B = M_{\tilde{i}}^i q_i \tilde{q}^{\tilde{i}} + \tilde{\eta} \tilde{Y}, \tag{1.2}$$

where $\tilde{\eta} = \tilde{\Lambda}^{\tilde{b}_0} = (-1)^{N_f - N_c} \eta^{-1}$ is the instanton factor of theory B, and \tilde{Y} is its monopole operator.¹ The arguments of [6] imply that the two $3d$ theories (1.1), (1.2) are equivalent at low energies.

Once such a $3d$ duality is established one can find many additional $3d$ dualities, which follow from it. First, we can turn on relevant operators in the two sides of the duality and flow to the IR. Second, we can gauge any of the global symmetries of the theories and generate new dual pairs. These two tools were used in [6] to reproduce all the known dualities between $3d$ $\mathcal{N} = 2$ theories with $SU(N_c)$, $U(N_c)$ and $USp(2N_c)$ gauge groups, and to generate many new dualities.

However, the application of this procedure to theories with orthogonal gauge groups turns out to need more care. In fact, already in $4d$ $\mathcal{N} = 1$ theories the IR dualities for orthogonal groups are significantly more subtle than for unitary or symplectic gauge

¹The first term in (1.2) contains already in $4d$ a scale μ [3], which is related to the normalization of the kinetic terms of M , q and \tilde{q} . It is natural to normalize M such that it is identified with the composite operator $M = Q\tilde{Q}$ in theory A. We will choose this normalization. When we reduce to $3d$ the parameter μ is still present, and there are also various factors of the radius. Here we ignore this normalization, which is irrelevant in the IR. We fix arbitrarily (but self-consistently) the coefficient of the first term in (1.2) ($\mu = 1$) and in the analogous expressions for $SO(N_c)$.

groups [1–3, 8]. One underlying reason for this complexity was recently identified in [5]. It is known that if the Lie algebra of the gauge symmetry is $so(N_c)$, the gauge group can be $\text{Spin}(N_c)$ or $\text{SO}(N_c)$ (and it could even have disconnected components, making it $\text{Pin}(N_c)$ or $O(N_c)$). The main point of [5] is that even when the gauge group is $\text{SO}(N_c)$, there are two distinct $4d$ gauge theories with that gauge group, denoted by $\text{SO}(N_c)_\pm$. In the Euclidean path integral they are distinguished by a new term in the Lagrangian — a certain \mathbb{Z}_2 -valued theta-like-angle, associated with the Pontryagin square $\mathcal{P}(w_2)$ of the Stiefel-Whitney class w_2 of the gauge bundle.

A simple physical way to distinguish between the three gauge theories $\text{SO}(N_c)_\pm$ and $\text{Spin}(N_c)$ already in \mathbb{R}^4 is to study their line operators. The $\text{Spin}(N_c)$ theory has a Wilson loop W in a spinor representation.² Its square W^2 can be screened by dynamical fields and we will view it as trivial. The two $\text{SO}(N_c)$ theories do not have a Wilson loop in a spinor representation. Instead, they have 't Hooft loops carrying smaller magnetic charge than is allowed in $\text{Spin}(N_c)$. The $\text{SO}(N_c)_+$ theory has a purely magnetic 't Hooft loop operator H , and the $\text{SO}(N_c)_-$ has the non-trivial loop operator HW . For a closely related earlier discussion, see [9].

For $N_c = 3$ the distinction between $\text{SO}(3)_\pm$ can be understood by extending the range of the ordinary theta-angle to be in $[0, 4\pi)$, and then [9]

$$\text{SO}(3)_+^\theta = \text{SO}(3)_-^{\theta+2\pi}. \tag{1.3}$$

Similarly, for $N_c = 4$ we have $\text{Spin}(4) = \text{SU}(2) \times \text{SU}(2)$ and $\text{SO}(4) = (\text{SU}(2) \times \text{SU}(2))/\mathbb{Z}_2$. Hence, the $so(4)$ theory has two theta-angles, one for each $\text{SU}(2)$, and

$$\text{SO}(4)_+^{\theta_1, \theta_2} = \text{SO}(4)_-^{\theta_1, \theta_2+2\pi} = \text{SO}(4)_-^{\theta_1+2\pi, \theta_2} = \text{SO}(4)_+^{\theta_1+2\pi, \theta_2+2\pi}. \tag{1.4}$$

For higher values of N_c the distinction between the two $\text{SO}(N_c)$ theories cannot be absorbed in extending the range of θ , and the only way to describe it at the level of the Lagrangian is by using the Pontryagin square $\mathcal{P}(w_2)$.

The $4d$ $\mathcal{N} = 1$ IR duality relates [1–3, 8] an $so(N_c)$ gauge theory with N_f vectors Q^i to an $so(\tilde{N}_c = N_f - N_c + 4)$ gauge theory, with N_f vectors q_i and elementary gauge singlet mesons M^{ij} , and with a superpotential $W = \frac{1}{2}M^{ij}q_iq_j$. When either N_c or \tilde{N}_c are 2, 3, 4 a more careful discussion is needed. The analysis of [5], which took into account the global structure,³ identified this duality as

$$\begin{aligned} \text{Spin}(N_c) &\rightarrow \text{SO}(\tilde{N}_c)_-, \\ \text{SO}(N_c)_- &\rightarrow \text{Spin}(\tilde{N}_c), \\ \text{SO}(N_c)_+ &\rightarrow \text{SO}(\tilde{N}_c)_+. \end{aligned} \tag{1.5}$$

Subtleties associated with the line operators in the $4d$ theory translate into subtleties with the local operators when the theory is compactified on a circle to $3d$. In particular,

²In this note we will study $so(N_c)$ gauge theories with matter fields in the vector representation. Therefore, for many purposes we can identify Wilson loops in different spinor representations. For a more detailed discussion see [5].

³For an earlier related discussion see [4].

a $4d$ 't Hooft line operator H wrapping this circle turns into a local monopole operator Y in $3d$. Hence, the choice of line operators in $4d$ becomes a choice of local operators in $3d$, which has more dramatic consequences, as we will see in our discussion below.

An additional subtlety in the analysis of orthogonal groups is that the corresponding $4d$ supersymmetric QCD (SQCD) theories on \mathbb{S}^1 have a Coulomb branch that is not lifted by quantum corrections. This did not occur in any of the cases analyzed in [6], and the mapping of the Coulomb branch across the duality turns out to be non-trivial.

In section 2 we discuss some classical and quantum properties of $3d$ $\mathcal{N} = 2$ theories with orthogonal gauge groups. We identify the coordinates on their moduli space of vacua, paying particular attention to the global structure (the distinction between $\text{Spin}(N)$ and $\text{SO}(N)$). In section 3 we discuss the $4d$ gauge theories on $\mathbb{R}^3 \times \mathbb{S}^1$ and their moduli space of vacua. Here the distinction between the three different theories with the same Lie algebra $\mathfrak{so}(N_c)$ is crucial. In section 4 we follow [6] and consider two dual $4d$ theories compactified on a circle, and carefully identify their moduli spaces.

In section 5 we derive the main result of this paper. By taking an appropriate limit of the $4d$ theory on a circle we derive $3d$ dualities. In particular, the $\text{SO}(N_c)$ SQCD theory with N_f vectors Q is dual to an $\text{SO}(N_f - N_c + 2)$ gauge theory with N_f vectors q , with gauge singlet fields M and Y , and with an appropriate superpotential. Here the elementary fields M and Y are identified with the composite meson QQ and monopole operator of the original $\text{SO}(N_c)$ theory. This is closely related to dualities of $O(N_c)$ SQCD theories, previously found in [10–12]. We also find that the $\text{Spin}(N_c)$ theory is dual to an $O(N_f - N_c + 2)_-$ theory, where $O(N)_-$ is a novel $3d$ $O(N)$ theory that we introduce in section 2.4. We perform various tests of these dualities, and deform them to obtain dualities for theories with Chern-Simons terms. Additional detailed tests are performed in section 6, where we discuss the $\mathbb{S}^2 \times \mathbb{S}^1$ and the \mathbb{S}^3 partition functions of all these theories.

2 Background

Much of the necessary background for this paper is found in the preceding paper [6], and in references therein. We will assume here familiarity with that paper, and discuss only the new issues which arise for orthogonal gauge groups. Some aspects of the theory depend on the precise choice of gauge group, while others depend only on the gauge algebra $\mathfrak{g} = \mathfrak{so}(N_c)$, and we will try to distinguish the two in the following.

2.1 Monopole operators and Coulomb branch coordinates for $\mathfrak{g} = \mathfrak{so}(N_c)$

Three dimensional $\mathcal{N} = 2$ gauge theories have classical Coulomb branches, where the adjoint scalar σ in the vector multiplet gets an expectation value, generically breaking the gauge group G to $U(1)^{r_G}$ (where r_G is the rank of G). On this branch we can dualize the r_G photons to scalars a_i , or supersymmetrically dualize the r_G $U(1)$ vector multiplets to chiral multiplets Y_i . The expectation values of these chiral multiplets label the classical Coulomb branch of the theory. The chiral multiplets Y_i are “monopole operators” in the effective low-energy theory, creating a $U(1)^{r_G}$ magnetic flux around them. In some cases they arise as low-energy limits of microscopic “monopole operators”. The allowed spectrum of monopole

operators, and thus the appropriate coordinates on the Coulomb branch, depends on the choice of the gauge group; this choice determines the allowed Wilson line operators, and the monopole operators need to be mutually local with respect to these Wilson lines.

For theories based on the Lie algebra $g = so(N_c)$, when N_c is even, $N_c = 2r_G$ and when N_c is odd, $N_c = 2r_G + 1$. We write the adjoint matrix σ as a matrix in the vector representation of $so(N_c)$, and we can always diagonalize it. For every non-zero eigenvalue, there is another eigenvalue of equal magnitude and opposite sign. For even values of N_c we write the eigenvalues as $\{\sigma_1, \dots, \sigma_{r_G}, -\sigma_{r_G}, \dots, -\sigma_1\}$. By a Weyl transformation we can always choose

$$N_c \text{ even : } \quad \sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_{r_G-1} \geq |\sigma_{r_G}|. \tag{2.1}$$

If our gauge group includes reflections (namely, it is $G = O(N_c)$ or $G = Pin(N_c)$ rather than $G = SO(N_c)$ or $G = Spin(N_c)$) then we can also set $\sigma_{r_G} \geq 0$, while otherwise we cannot do this in general. For odd values of N_c we can write the eigenvalues of σ as $\{\sigma_1, \dots, \sigma_{r_G}, 0, -\sigma_{r_G}, \dots, -\sigma_1\}$, and by a Weyl transformation we can always choose

$$N_c \text{ odd : } \quad \sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_{r_G-1} \geq \sigma_{r_G} \geq 0. \tag{2.2}$$

The magnetic charges carried by the Coulomb branch coordinates should be thought of as charges in the magnetic-dual algebra to $so(N_c)$. For even N_c , this algebra is $so(N_c)$, and for odd N_c , it is $usp(N_c - 1)$. There are always operators carrying the charges of the roots of this algebra, and when the gauge group is $G = Spin(N_c)$, these are the only allowed charges. We can then write the Coulomb branch coordinates semi-classically as

$$\begin{aligned}
 Y_i &\approx \exp\left(\frac{\sigma_i - \sigma_{i+1}}{\hat{g}_3^2} + i(a_i - a_{i+1})\right), \quad (i = 1, \dots, r_G - 1), \\
 Y_{r_G} &\approx \begin{cases} \exp\left(\frac{\sigma_{r_G-1} + \sigma_{r_G}}{\hat{g}_3^2} + i(a_{r_G-1} + a_{r_G})\right) & \text{for even } N_c, \\ \exp\left(\frac{2\sigma_{r_G}}{\hat{g}_3^2} + i(2a_{r_G})\right) & \text{for odd } N_c. \end{cases} \tag{2.3}
 \end{aligned}$$

Here $\hat{g}_3^2 = g_3^2/4\pi$, where g_3 is the gauge coupling constant of the $3d$ gauge theory, normalized as in [6]. The dependence of these operators on the σ 's that we wrote is valid far out on the Coulomb branch, and gets quantum corrections, while their dependence on the dual photons a_i is exact. As usual, the global symmetry charges of these operators can be determined by summing over the charges of the fermions in chiral and vector multiplets, which are coupled to the corresponding σ 's [13]. The $SO(N_c)$ and $Spin(N_c)$ theories have a global charge conjugation symmetry \mathbb{Z}_2^C , which is gauged in the $O(N_c)$ and $Pin(N_c)$ theories. In the theories with even N_c , this symmetry exchanges the Coulomb branch coordinates Y_{r_G-1} and Y_{r_G} .

As discussed in [6, 14], some of the Coulomb branch coordinates are low-energy limits of microscopic monopole operators. These are defined so that their insertion at a point x generates some magnetic flux on the \mathbb{S}^2 surrounding x , and takes the $\sigma(y)$'s pointing in the direction of the flux to $+\infty$ as $y \rightarrow x$. In the $Spin(N_c)$ theory, the monopole operators all carry charges corresponding to roots of the dual magnetic group. The minimal monopole

operator Y_{Spin} turns on one unit of flux, breaking $so(N_c) \rightarrow so(N_c-2) \times u(1)$, and takes one of the eigenvalues of σ to ∞ . On the moduli space at low energies, using (2.1) and (2.2), this monopole Y_{Spin} looks semi-classically like

$$Y_{\text{Spin}} \approx \exp\left(\frac{2\sigma_1}{\hat{g}_3^2} + 2ia_1\right). \tag{2.4}$$

It is a combination of the Coulomb branch coordinates described in (2.3),

$$Y_{\text{Spin}} = \begin{cases} Y_1^2 Y_2^2 \cdots Y_{r_G-2}^2 Y_{r_G-1} Y_{r_G} & \text{for even } N_c, \\ Y_1^2 Y_2^2 \cdots Y_{r_G-1}^2 Y_{r_G} & \text{for odd } N_c. \end{cases} \tag{2.5}$$

Another monopole operator that will play a role in our discussion is the one that takes two eigenvalues of σ to infinity together, breaking $so(N_c) \rightarrow so(N_c-4) \times u(2)$. This monopole semi-classically looks like

$$Z \approx \exp\left(\frac{\sigma_1 + \sigma_2}{\hat{g}_3^2} + i(a_1 + a_2)\right), \tag{2.6}$$

and we will see that it will play an important role in the discussion of $4d$ $so(N_c)$ theories on a circle. It obeys $Y_{\text{Spin}} = Y_1 Z$, and

$$Z = \begin{cases} Y_1 Y_2^2 Y_3^2 \cdots Y_{r_G-2}^2 Y_{r_G-1} Y_{r_G} & \text{for even } N_c, \\ Y_1 Y_2^2 Y_3^2 \cdots Y_{r_G-1}^2 Y_{r_G} & \text{for odd } N_c. \end{cases} \tag{2.7}$$

For $G = \text{SO}(N_c)$, Wilson lines carrying spinor charge are not allowed (we will always assume that Wilson lines in the vector representation are allowed, since we will be interested here in theories with matter fields in the vector representation). This means that extra Coulomb branch coordinates and monopole operators are allowed, carrying weights which are not roots of the dual magnetic group. For even N_c they are allowed to carry weights in the vector representation of the dual $so(N_c)$, and for odd N_c in the fundamental representation of the dual $usp(N_c-1)$. The basic monopole operator in this case behaves semi-classically as [10, 12]

$$Y \approx \exp\left(\frac{\sigma_1}{\hat{g}_3^2} + ia_1\right), \tag{2.8}$$

and it obeys

$$Y^2 = Y_{\text{Spin}} = Y_1 Z. \tag{2.9}$$

All other “new” monopole operators that exist in this case may be written as products of Y with the operators corresponding to roots of the dual gauge group. Note that while for the monopole operators corresponding to roots, there is a classical ’t Hooft-Polyakov monopole solution (which is an instanton of the $3d$ theory) that is associated with them, there is no such solution for Y of (2.8). But this is not related to the definition of this operator, both microscopically and in the low-energy effective action.

The fact that the quantum numbers of the Coulomb branch coordinates are determined by those of the matter fields implies that they change when some matter fields go to

infinite mass and decouple. In such cases we have a relation between the Coulomb branch coordinates in the high-energy and in the low-energy theories, which can usually be uniquely determined by matching their quantum numbers. For example, if we start from a $\text{Spin}(N_c)$ theory with N_f chiral superfields in the vector representation, and give a mass m to one of them in the superpotential, we have a relation of the form

$$(Y_{\text{Spin}})_{N_c, N_f} = m(Y_{\text{Spin}})_{N_c, N_f - 1}. \quad (2.10)$$

Similarly, if we start from such a theory and break the gauge group to $\text{Spin}(N_c - 1)$ with $(N_f - 1)$ flavors, by giving an expectation value to one of the chiral superfields Q in the vector representation, we have a relation

$$(Y_{\text{Spin}})_{N_c, N_f} \langle Q^2 \rangle = (Y_{\text{Spin}})_{N_c - 1, N_f - 1}. \quad (2.11)$$

For low values of N_c , $N_c < 5$, some modifications are needed in our discussion.

For $N_c = 2$ the gauge group $G = \text{Spin}(2) = \text{U}(1)$, and instead of the operator Y_{Spin} we have the two standard $\text{U}(1)$ Coulomb branch coordinates [13]

$$\hat{V}_+ \approx \exp\left(\frac{2\sigma}{\hat{g}_3^2} + 2ia\right), \quad \hat{V}_- \approx \exp\left(-\frac{2\sigma}{\hat{g}_3^2} - 2ia\right), \quad (2.12)$$

parameterizing the parts of the Coulomb branch with σ positive and negative, respectively. Note that in $\text{Spin}(2)$ we have particles of charge $\pm 1/2$ under the $\text{U}(1)$ group, and hence the normalization of the monopole operators is twice the usual normalization. For $G = \text{SO}(2)$ all particles have integer $\text{U}(1)$ charge, and we have monopole operators carrying half the charge of (2.12), given by

$$V_+ \approx \exp\left(\frac{\sigma}{\hat{g}_3^2} + ia\right), \quad V_- \approx \exp\left(-\frac{\sigma}{\hat{g}_3^2} - ia\right). \quad (2.13)$$

The charge conjugation symmetry $\mathbb{Z}_2^{\mathcal{C}}$ exchanges \hat{V}_+ and \hat{V}_- (or V_+ and V_-), and we will find it convenient to define the linear combinations

$$W_{\pm} \equiv V_+ \pm V_-, \quad \hat{W}_{\pm} \equiv \hat{V}_+ \pm \hat{V}_-, \quad (2.14)$$

that are even and odd under $\mathbb{Z}_2^{\mathcal{C}}$.

For $N_c = 3$, we have $\text{Spin}(3) = \text{SU}(2)$, and the monopole Y_{Spin} that we defined above is the standard Coulomb branch coordinate / monopole operator of the $\text{SU}(2)$ theory (usually denoted by Y [13]). When the gauge group is $\text{SO}(3) = \text{SU}(2)/\mathbb{Z}_2$, there exists a monopole operator of lower charge, which we denoted by Y above.

For $N_c = 4$, the group $\text{Spin}(4)$ is equivalent to $\text{SU}(2) \times \text{SU}(2)$, and we can then have separate Coulomb branch coordinates and monopole operators in the two $\text{SU}(2)$ factors. The Coulomb branch coordinates of the two $\text{SU}(2)$'s correspond to $\sigma_1 \pm \sigma_2$ in our notations above. Thus, the Coulomb branch coordinates of the two $\text{SU}(2)$'s, which we will denote by $Y^{(1)}$ and $Y^{(2)}$, look semi-classically like the operators Y_1 and Z discussed above. The operator Y_{Spin} in this case is the product of these two $\text{SU}(2)$ operators. When the gauge

group is $G = \text{SO}(4) = (\text{SU}(2) \times \text{SU}(2))/\mathbb{Z}_2$, one does not allow Wilson loops carrying a charge under the center of each $\text{SU}(2)$ separately, but only under both $\text{SU}(2)$'s. In this case the operator Y that we defined in (2.8) exists, and squares to the product of the two $\text{SU}(2)$ monopole operators, $Y^2 = Y^{(1)}Y^{(2)}$. The charge conjugation symmetry \mathbb{Z}_2^C exchanges the two $\text{SU}(2)$ factors, exchanging $Y^{(1)}$ with $Y^{(2)}$. As in (2.14), it is convenient to define the combinations

$$Y_{\pm} \equiv Y^{(1)} \pm Y^{(2)}, \tag{2.15}$$

which are even and odd under \mathbb{Z}_2^C .

2.2 The quantum moduli space of $3d \mathcal{N} = 2$ theories with $g = so(N_c)$

In the quantum theory, most of the Coulomb branch described above is lifted. Whenever two of the eigenvalues of σ come together at a non-zero value, the corresponding $\text{U}(1)^2$ symmetry is enhanced to $\text{U}(2)$. As shown in [15], the corresponding 't Hooft Polyakov monopole-instanton solutions generate a superpotential in this case, which drives the eigenvalues apart. For even values of N_c we have this effect whenever σ_i approaches σ_{i+1} for $i = 1, 2, \dots, r_G - 1$, and also when $\sigma_{r_G - 1}$ approaches $-\sigma_{r_G}$. Thus, in the $3d$ pure $g = so(N_c)$ theory with N_c even we have an effective quantum superpotential

$$N_c \text{ even : } \quad W = \sum_{i=1}^{r_G-1} \frac{1}{Y_i} + \frac{1}{Y_{r_G}}, \tag{2.16}$$

which completely lifts the Coulomb branch. For odd values of N_c there is always one eigenvalue at $\sigma = 0$, and when σ_{r_G} approaches 0, the corresponding $u(1)$ is enhanced to $so(3)$ from 3 eigenvalues coming together at $\sigma = 0$. There is a similar superpotential arising here, with a different normalization [16], so that

$$N_c \text{ odd : } \quad W = \sum_{i=1}^{r_G-1} \frac{1}{Y_i} + \frac{2}{Y_{r_G}}. \tag{2.17}$$

Again this completely lifts the Coulomb branch, so that the pure $3d$ supersymmetric Yang-Mills (SYM) theory has a runaway with no supersymmetric vacua.

We can follow the reasoning used in [6] for $\text{SU}(N)$ gauge theories to show that (2.16) and (2.17) are in fact exact. More precisely, they are exact as functions of the chiral superfields Y_i , but the Y_i are complicated functions of σ_i and a_i .

Similar superpotentials lift the Coulomb branch also for $g = so(3) = su(2)$, and for $g = so(4) = su(2) \oplus su(2)$, where we have a separate superpotential of this type in each $su(2)$ factor. For $g = so(2) = u(1)$ there is no such effect, and the Coulomb branch of the (free) pure gauge theory is simply a cylinder, labeled by $\hat{V}_+ = 1/\hat{V}_-$ for $G = \text{Spin}(2)$, and by $V_+ = 1/V_-$ for $G = \text{SO}(2)$.

In theories with flavors in the vector representation that have no real mass, these flavors become massless whenever some eigenvalues of σ vanish. This gives extra zero modes to monopole-instantons corresponding to eigenvalues coming together at $\sigma = 0$, such that they no longer generate superpotentials; for odd N_c this happens for the monopole Y_{r_G} , and for

even N_c it happens either for $Y_{r_{G-1}}$ or for Y_{r_G} , depending on the sign of σ_{r_G} . In these theories the moduli space is not completely lifted, but a one (complex) dimensional branch remains, where only a single eigenvalue σ_1 is turned on (and $\sigma_2 = \sigma_3 = \dots = \sigma_{r_G} = 0$). In the $\text{Spin}(N_c)$ theories, this quantum Coulomb branch may be parameterized by the operator Y_{Spin} of (2.4), and for $G = \text{SO}(N_c)$ theories it can be parameterized by Y of (2.8). Note that because of the quantum superpotentials (2.16), (2.17), the operator Z is not a chiral operator in the low-energy 3d theory,⁴ but Y and Y_{Spin} still are. As discussed in [10, 12], for low numbers of flavors, $N_f < N_c - 2$, there are additional quantum effects that lift the Coulomb branch, while for $N_f \geq N_c - 2$ the Coulomb branch is not lifted. For $N_f \geq N_c - 1$, the quantum moduli space is the same as the classical moduli space (the discussion in [10, 12] is just for $G = O(N_c)$, but most of it can be generalized also to the other gauge groups with $g = so(N_c)$).

The 3d SQCD theory with N_f chiral multiplets Q_i in the vector representation has a global $\text{SU}(N_f) \times \text{U}(1)_A \times \text{U}(1)_R$ symmetry. We can choose the flavors Q_i to transform in the fundamental of $\text{SU}(N_f)$, with one unit of $\text{U}(1)_A$ charge and no $\text{U}(1)_R$ charge. In this case, for $G = \text{SO}(N_c)$ theories the Coulomb branch coordinate Y is a singlet of $\text{SU}(N_f)$, with $(-N_f)$ units of $\text{U}(1)_A$ charge and $(N_f - N_c + 2)$ units of $\text{U}(1)_R$ charge. For $G = \text{Spin}(N_c)$ the operator Y_{Spin} has the same charges as Y^2 in the $G = \text{SO}(N_c)$ theories.

Note that for $N_c = 2$ our theories with flavors are the same as the $\text{U}(1)$ theories with flavors discussed in [13]. However, for $N_c = 3$ our $so(3)$ theories have matter in the triplet (adjoint) representation, so they are not the same as $su(2)$ theories with fundamental matter. In particular, for $N_c = 3$ and $N_f = 1$, the SQCD theory has enhanced supersymmetry, and it is the same as the 3d $\mathcal{N} = 4$ SYM theory discussed in [17]. For $N_c = 4$ the matter fields are charged under both $\text{SU}(2)$'s, and couple them together. Depending on where we are in the moduli space, the superpotential involving the Coulomb branch coordinate of one of the $\text{SU}(2)$'s is lifted by the matter zero modes, while the other one remains. For $N_c = 3, 4$ we can still parameterize the remaining part of the Coulomb branch by Y_{Spin} or Y , as for higher values of N_c .

2.3 Baryon-monopole operators

In the $g = so(N_c)$ SQCD theory with N_f flavors Q_i , the list of chiral multiplets includes the monopole operators discussed above, the mesons $M_{ij} = Q_i Q_j$ (symmetric in i, j) and (for $N_f \geq N_c$, and for $G = \text{SO}(N_c)$ or $G = \text{Spin}(N_c)$) the baryons $B = Q^{N_c}$, contracted with an epsilon symbol in $so(N_c)$. The operator B^2 may be written as a combination of products of N_c mesons M , but B is an independent operator, charged under \mathbb{Z}_2^C . For $G = O(N_c)$ and $G = \text{Pin}(N_c)$ the operator B does not exist, since the charge conjugation symmetry is gauged.

Consider now a monopole operator like Y in $\text{SO}(N_c)$, which breaks

$$\text{SO}(N_c) \rightarrow S\left(O(N_c - 2) \times O(2)\right). \tag{2.18}$$

⁴ Z is proportional to two supercharges acting on $(Y_{\text{Spin}} Y_1 K_{Y_1})$, where K_{Y_1} is the derivative of the Kähler potential with respect to Y_1 .

Note that this includes transformations with determinant (-1) both in $\text{SO}(N_c - 2)$ and in $\text{SO}(2)$. This means that the gauge-invariant operator Y must be charge conjugation even in $\text{SO}(2)$, and thus it reduces to the operator W_+ (2.14) in this group. However, we can also build a “baryon-monopole” operator involving W_- in $\text{SO}(2)$ (2.14), by defining

$$\beta = Q^{N_c-2}W_-, \tag{2.19}$$

with the indices of the flavors contracted by an epsilon symbol in the $\text{SO}(N_c - 2)$ that is unbroken by the monopole. The product Q^{N_c-2} is invariant under the $\text{SO}(N_c - 2) \times \text{SO}(2)$ subgroup of $\text{SO}(N_c)$ that is left unbroken by the monopole, and its product with W_- is invariant also under the extra \mathbb{Z}_2 , so that (2.19) defines a gauge-invariant operator in $\text{SO}(N_c)$. Note that the operators (2.19) exist for any $N_f \geq N_c - 2$. The standard matching of quantum numbers for monopole operators, generalizing (2.11), implies that when we give a VEV to Q breaking $\text{SO}(N_c)$ to $\text{SO}(2)$, β reduces directly to W_{low} in the low-energy $\text{SO}(2)$ theory (with no extra factors of $\langle Q \rangle$).

As we discussed above, when the gauge group is $\text{Spin}(N_c)$, the monopole operators Y and W_{\pm} do not exist. But we can still repeat the above discussion using the operator Y_{Spin} (which reduces to \hat{W}_+), and define a baryon-monopole

$$\beta_{\text{Spin}} = Q^{N_c-2}\hat{W}_- \tag{2.20}$$

as above. Note that in an $\text{SO}(2)$ theory, $\hat{W}_- = V_+^2 - V_-^2 = (V_+ - V_-)(V_+ + V_-)$, so in $\text{SO}(N_c)$ theories $\beta_{\text{Spin}} = \beta Y$.

The operator β satisfies an interesting chiral ring relation. Consider the $3d$ $\text{SO}(N_c)$ gauge theory with $N_f = N_c - 2$. At a generic point on the moduli space of this theory, we break $\text{SO}(N_c) \rightarrow \text{SO}(2)$. The monopole operator Y reduces in the low-energy $\text{SO}(2)$ theory to W_{low} ; the standard mapping of monopoles (2.11) implies that $W_{\text{low}} = Y\sqrt{\det(M)}$. The low-energy $\text{SO}(2)$ theory has no massless flavors, and hence $V_{\text{low}}V_{\text{low}} = 1$. Therefore, in this vacuum

$$\beta^2 = W_{\text{low}}^2 = V_{\text{low}}^2 - 2V_{\text{low}}V_{\text{low}} + V_{\text{low}}^2 = W_{\text{low}}^2 - 4 = Y^2\det(M) - 4. \tag{2.21}$$

This reflects an exact chiral ring relation

$$\beta^2 = Y^2\det(M) - 4, \tag{2.22}$$

which is valid in every vacuum of this theory. Classically this theory has a point at the origin of its moduli space where $Y = M = \beta = 0$, but we see that quantum mechanically the moduli space is deformed and obeys (2.22). This is similar to the deformation of the classical moduli space in some $4d$ theories [18] and in some $3d$ theories [13].

Similarly, we can use the extra monopole operators of $\text{SO}(4)$, by having monopoles breaking $\text{SO}(N_c)$ to $\text{SO}(N_c - 4) \times \text{U}(2)$. More precisely, to define such monopoles we first choose some $S(O(N_c - 4) \times O(4)) \subset \text{SO}(N_c)$, and then turn on a monopole like (2.6) in the $\text{SO}(4)$ factor, breaking it to $\text{U}(2)$. As in our discussion above, the monopole Z on its own reduces to the even operator Y_+ in $\text{SO}(4)$ (2.15). But we can now consider instead the operator

$$b = Q^{N_c-4}Y_-, \tag{2.23}$$

where the quarks are contracted with the epsilon symbol of $\text{SO}(N_c-4)$. As in our discussion above, this operator b is gauge-invariant in the $G = \text{SO}(N_c)$ theory, including also the gauge transformations of determinant (-1) in the two factors. The same discussion applies also to $\text{Spin}(N_c)$ theories. The baryon-monopole b exists for any $N_f \geq N_c - 4$ (for $N_c = 4$ it is simply Y_-). As in our discussion of Z above, due to quantum effects b is not really a chiral operator in $3d$ SQCD theories, but we will see that it still plays a role in our analysis.

We cannot generalize this construction to breaking $\text{SO}(N_c) \rightarrow S(O(N_c-2n) \times O(2n))$ with $n > 2$, because there is no obvious monopole operator in $\text{SO}(2n)$ that is odd under charge conjugation.

2.4 On $3d$ $O(N_c)$ theories

We mentioned above that one can obtain $O(N_c)$ theories by gauging the charge conjugation symmetry of $\text{SO}(N_c)$, but in fact there are two different $O(N_c)$ theories in $3d$ that will play a role in this paper. In one $O(N_c)$ theory the minimal monopole operator Y of $\text{SO}(N_c)$, which is charge-conjugation-even in the $\text{SO}(N_c)$ theory, is gauge-invariant, and is the minimal monopole operator also for $O(N_c)$. We will denote this theory by $O(N_c)_+$; this is the theory that was discussed in previous papers about $3d$ $O(N_c)$ theories and their dualities. However, as in [5], one can also define a second $O(N_c)_-$ theory, in which the monopole operator Y is charge-conjugation-odd (it changes sign under gauge transformations whose determinant is (-1)). In this $O(N_c)_-$ theory, Y and B are both not gauge-invariant, but their product, as well as the operators Y_{Spin} and β , are gauge-invariant (note that β is not gauge-invariant in the standard $O(N_c)_+$ theory).

In the Lagrangian language, the two theories differ by a discrete theta angle, analogous to the one that distinguishes the $4d$ $\text{SO}(N_c)_\pm$ theories [5]. The relevant term in the Lagrangian is proportional to $w_1 \wedge w_2$, where $w_i \in H^i(X, \mathbb{Z}_2)$ are \mathbb{Z}_2 -valued characteristic classes of the $O(N_c)$ bundle on a manifold X ; w_1 is non-zero when the $O(N_c)$ bundle cannot be written as an $\text{SO}(N_c)$ bundle, while w_2 is non-zero when the $O(N_c)$ bundle cannot be written as a $\text{Pin}(N_c)$ bundle. In particular, w_2 is non-zero on a two-sphere around an insertion of the operator Y . Note that the two options exist only for $O(N_c)$ gauge groups, not for $\text{SO}(N_c)$, $\text{Spin}(N_c)$ or $\text{Pin}(N_c)$ (in which either w_1 , or w_2 , or both, are trivial).⁵

3 The Coulomb branch of $4d$ $so(N_c)$ theories on \mathbb{S}^1

3.1 $4d$ $\text{Spin}(N_c)$ theories on \mathbb{S}^1

As discussed in [6], when one compactifies a $4d$ gauge theory on \mathbb{S}^1 and goes to low energies, naively one gets the same gauge theory in $3d$, but there are two important differences. The first is that the Coulomb branch coordinates now come from holonomies of the gauge field on a circle, so the coordinates σ described above are periodic and the Coulomb branch is compact. The second is that there is an extra monopole-instanton in the theory on a circle, that gives an extra term in the effective superpotential. We will start by discussing these

⁵More generally, it may be possible to add other terms such as $w_1 \wedge w_1 \wedge w_1$, but we will not discuss this here.

aspects for the reduction of a $4d$ theory with $G = \text{Spin}(N_c)$, and consider $G = \text{SO}(N_c)$ theories in the next subsection. We begin with the case of $N_c \geq 5$.

In the $4d$ theory on a circle of radius r , the scalars σ described in the previous section originate from A_3 , but only the eigenvalues of $U = P\exp(i \oint A_3)$ are gauge-invariant, so there are relations between different values of σ associated with large gauge transformations. In particular, the eigenvalues of U in the vector representation are $\exp(\pm 2\pi i r \sigma_i)$, so each σ_i gives the same holonomy in this representation as $\sigma_i + 1/r$. Note that the eigenvalues of U in the spinor representation are $\exp(2\pi i r (\pm \sigma_1/2 + \dots))$, so the periodicity of each σ_i for $G = \text{Spin}(N_c)$ is actually $2/r$ (or one can shift two σ_i 's together by $1/r$). But the masses of W-bosons and matter fields in the vector representation are periodic in σ_i with periodicity $1/r$. In particular, whenever all the σ_i are integer multiples of $1/r$, the gauge group is unbroken and any matter fields in the vector representation are massless.

In the $4d$ theory on a circle, we can get an enhanced non-Abelian symmetry not just by having $\sigma_i \rightarrow \sigma_{i+1}$, but also by having eigenvalues meet the images of other eigenvalues. When σ_1 meets the image $-\sigma_1$ at $\langle \sigma_1 \rangle = 1/2r$ there is no enhanced non-Abelian symmetry, since we just have an enhancement of $U(1)$ to $SO(2)$ or $\text{Spin}(2)$. However, when σ_1 meets the image of $-\sigma_2$, when $\sigma_1 = -\sigma_2 + 1/r$, there is an enhancement of $U(1)^2$ to $U(2)$ (if this happens at $\sigma_1 = 1/2r$ then there is even an enhancement to $SO(4)$ or $\text{Spin}(4)$). The same computation yielding the monopole-instanton contributions described above [15], thus gives in the theory on a circle an extra superpotential. The analogy with the $1/Y$ superpotential of [15] implies that semi-classically the extra superpotential looks like

$$\frac{1}{\exp\left(\frac{1-\sigma_1-\sigma_2}{r g_3^2} - i(a_1 + a_2)\right)} \approx \eta Z, \tag{3.1}$$

where Z was defined in (2.6), and $\eta \equiv \Lambda^{b_0} = \exp(-8\pi^2/g_4^2) = \exp(-4\pi/g_3^2 r)$ is the strong coupling scale of the $4d$ gauge theory ($b_0 = 3(N_c - 2) - N_f$ is the $4d$ one-loop beta function coefficient, and we set the $4d$ theta angle to zero and the renormalization scale to one for simplicity). The precise form (3.1) follows by carefully analyzing all the instantons, as in [16, 17, 19–21]. From the point of view of the effective $3d$ theory, (3.1) breaks precisely the global $U(1)$ symmetry that is anomalous in the $4d$ theory. Note that in the $3d$ theory Z is not a chiral operator, but in the $4d$ theory on a circle, it can no longer be written as a descendant, due to the extra superpotential (3.1).

In the pure SYM theory, the extra term (3.1) stabilizes the runaway caused by (2.16) and (2.17), and leads to a finite number of supersymmetric vacua, obtained by solving the F-term equations for the Y_i . One can check that, both for even and odd values of N_c , this leads to $(N_c - 2)$ supersymmetric vacua, with

$$(2\eta \langle Z \rangle)^{N_c - 2} = 16\eta. \tag{3.2}$$

This is the same number of vacua as in the $4d$ theory, as expected in this case [5], and the value of the superpotential also agrees with its $4d$ value. As discussed in [6], the $4d$ chiral operator $S \propto \text{tr}(W_\alpha^2)$ reduces in the theory on a circle to Z , with a chiral ring relation

$$S = \eta Z, \tag{3.3}$$

which is consistent with (3.2). Note that in the $4d$ theory on a circle, the monopole and baryon-monopole operators discussed in the previous section do not exist microscopically (due to the compactness of the Coulomb branch), but we can give a microscopic definition to Z using (3.3).

Moving next to the theories with flavors, note that unlike in the cases of $G = \text{SU}(N_c)$ and $G = \text{USp}(2N_c)$ discussed in [6], here the extra superpotential (3.1) is not proportional to the coordinate Y_{Spin} along the unlifted Coulomb branch of the $3d$ $\text{Spin}(N_c)$ SQCD theory. Thus, this superpotential does not lift the Coulomb branch, but just adds extra interactions. The coordinate Y_{Spin} on this Coulomb branch is uncharged under the continuous $\text{SU}(N_f) \times \text{U}(1)_R$ global symmetry that is preserved by (3.1).

The global structure of the moduli space is interesting. First, σ_1 should be identified with $(-\sigma_1 + 2/r)$ because they lead to the same holonomy, so we can take $0 \leq \sigma_1 \leq 1/r$. In the quantum theory the moduli space is parameterized by Y_{Spin} (2.4), and we identify the point $Y_{\text{Spin}} = 0$ with $\sigma_1 = 0$, and the point $Y_{\text{Spin}} = \infty$ with $\sigma_1 = 1/r$. Classically, at generic values of Y_{Spin} the $\text{Spin}(N_c)$ symmetry is broken, and at the two special points $Y_{\text{Spin}} = 0, \infty$ the full $\text{Spin}(N_c)$ symmetry is preserved.

Second, if our $\text{Spin}(N_c)$ gauge theory does not couple to matter fields in a spinor representation, the compactified theory has a *global \mathbb{Z}_2 symmetry*, acting on the moduli space by $\sigma_1 \rightarrow -\sigma_1 + 1/r$. The point is that these two values of σ_1 represent two different holonomies in $\text{Spin}(N_c)$, but this difference is not felt by any dynamical field.⁶ The two special points on the moduli space $Y_{\text{Spin}} = 0, \infty$ are not identified. Instead, they are exchanged by the global \mathbb{Z}_2 symmetry, which acts on the moduli space as

$$Y_{\text{Spin}} \longleftrightarrow \frac{1}{\eta^2 Y_{\text{Spin}}}. \tag{3.4}$$

In the $4d$ $G = \text{Spin}(N_c)$ theory there are several baryonic operators defined in [1, 2],

$$B = Q^{N_c}, \quad \mathcal{W}_\alpha = W_\alpha Q^{N_c-2}, \quad b_{4d} = W_\alpha^2 Q^{N_c-4}, \tag{3.5}$$

which all involve contractions with the epsilon symbol of $\text{Spin}(N_c)$. The first operator obviously reduces to the same baryon operator in $3d$. The second operator is useful when

⁶More generally, whenever we have a \mathbb{Z}_p gauge theory in d dimensions that does not couple to charged fields, the theory compactified to $d-1$ dimensions has both a gauge \mathbb{Z}_p symmetry and a global \mathbb{Z}_p symmetry. One way to see that is to represent the \mathbb{Z}_p gauge theory by a one-form $\text{U}(1)$ gauge field $A^{(1)}$ and a $(d-2)$ -form gauge field $A^{(d-2)}$ with a Lagrangian given by $\frac{p}{2\pi} A^{(1)} \wedge dA^{(d-2)}$ [22, 23]. Reducing this system to $d-1$ dimensions leads to four fields, with the Lagrangian $\frac{p}{2\pi} (A^{(1)} \wedge dA^{(d-3)} + A^{(0)} \wedge dA^{(d-2)})$. The first term represents a \mathbb{Z}_p gauge theory, and the second term describes a \mathbb{Z}_p global symmetry (and a gauge symmetry for $A^{(d-2)}$). The global symmetry acts as $A^{(0)} \rightarrow A^{(0)} + 2\pi/p$, with the identification $A^{(0)} \sim A^{(0)} + 2\pi$. If the original d dimensional theory includes matter fields charged under the \mathbb{Z}_p gauge symmetry, the global symmetry $A^{(0)} \rightarrow A^{(0)} + 2\pi/p$ in the $d-1$ dimensional theory is explicitly broken, but the identification $A^{(0)} \sim A^{(0)} + 2\pi$ is preserved. A famous example of this phenomenon, due to Polyakov, is the compactification of a $4d$ $\text{SU}(N)$ gauge theory without matter on \mathbb{S}^1 . The resulting $3d$ theory has a global \mathbb{Z}_N symmetry, which originates from the $4d$ $\mathbb{Z}_N \subset \text{SU}(N)$ gauge symmetry. The order parameter for its spontaneous breaking is the Polyakov loop $e^{iA^{(0)}} \propto \frac{1}{N} \text{tr}(e^{i \oint A})$. In our case, the relevant \mathbb{Z}_2 gauge symmetry is the subgroup of the center of $\text{Spin}(N_c)$ that acts on spinors.

$(N_c - 2)$ quarks get expectation values breaking the gauge group to $g = so(2)$, to label the remaining unbroken $so(2) \subset so(N_c)$; in the effective $3d$ theory the same role is played by β or β_{Spin} . Similarly, when the gauge group is broken to $g = so(4)$, we have a relation between $S_1 - S_2$ (of $SU(2) \times SU(2)$ in $4d$) and Y_- of $so(4)$ (2.15) that is analogous to (3.3), and that implies that the $4d$ baryon b_{4d} goes down in the low-energy effective theory to η times the baryon-monopole b defined in (2.23).

As in the $3d$ discussion, there are some modifications to this story for low values of N_c . For $N_c = 2$ there is no extra superpotential on the circle, and the moduli space for $G = U(1)$ was discussed in section 4.2 of [6]. For $N_c = 3$, there is only a single non-trivial eigenvalue of σ , and instead of (3.1) we get a superpotential from the fact that when $\sigma_1 \rightarrow 1/r$ the gauge group is enhanced again to $so(3)$. The full superpotential of the $4d$ Spin(3) pure SYM theory on a circle takes the form $W = 1/Y_{\text{Spin}} + \eta^2 Y_{\text{Spin}}$, consistent with the global symmetry (3.4) that acts also in this case. Note that in this special case the instanton factor of Spin(3) is actually $\eta_{\text{Spin}(3)} = \eta^2 = \Lambda^{2b_0}$ (keeping our general definition above for η), so this discussion is consistent with the standard discussion of $4d$ SU(2) theories on a circle [17]. The $4d$ Spin(3) SYM theory on a circle has two vacua at $Y_{\text{Spin}} = \pm 1/\eta = \pm 1/\sqrt{\eta_{\text{Spin}(3)}}$, which are fixed points of the \mathbb{Z}_2 symmetry (3.4) (though they are related by a \mathbb{Z}_4 global R-symmetry transformation).

For $N_c = 4$ with Spin(4) = SU(2) \times SU(2), we have in the pure $4d$ gauge theory on a circle two copies of the discussion of the previous paragraph,

$$W = \frac{1}{Y^{(1)}} + \frac{1}{Y^{(2)}} + \eta Y^{(1)} + \eta Y^{(2)} \tag{3.6}$$

(consistent with the identification of Z discussed in section 2.3). Note that the two SU(2)'s have $\eta_1 = \eta_2 = \eta$. There are 4 supersymmetric vacua at $Y^{(1)} = \pm 1/\sqrt{\eta}$, $Y^{(2)} = \pm 1/\sqrt{\eta}$. In two of these vacua $Y_{\text{Spin}} = Y^{(1)}Y^{(2)} = 1/\eta$, and in the other two $Y_{\text{Spin}} = -1/\eta$; they are all fixed points of the global symmetry transformation (3.4).

3.2 $4d$ SO(N_c) theories on \mathbb{S}^1

We saw in the previous section that an important difference between $G = \text{Spin}(N_c)$ and $G = \text{SO}(N_c)$ is that in the latter case there is an extra monopole operator Y that can be used to label the Coulomb branch. The Coulomb branch of the SO(N_c) theory is a double cover of that of the Spin(N_c) theory, which is labeled by $Y_{\text{Spin}} = Y^2$. When we discuss the $4d$ theory on a circle, another difference is that the global symmetry transformation (3.4) becomes a large gauge transformation in SO(N_c), with the gauge transformation parameter periodic around the circle in SO(N_c) but not in Spin(N_c). Thus, the points on the Coulomb branch related by (3.4) are identified. This is related to the fact that we no longer have a spinor Wilson line, so the σ_i have periodicity $1/r$, and we can restrict the range of σ_1 to $0 \leq \sigma_1 \leq 1/2r$.

Let us see in more detail how the \mathbb{Z}_2 large gauge transformation acts on Y . Given the relation $Y^2 = Y_{\text{Spin}}$, and the action on Y_{Spin} (3.4), there are two possible actions on Y : it could act either as $Y \longleftrightarrow 1/\eta Y$, or as $Y \longleftrightarrow -1/\eta Y$. These two options are related to the fact that there are two distinct SO(N_c) $4d$ gauge theories [5], SO(N_c) $_{\pm}$. As explained

in [5] and reviewed in the introduction, the difference between the three $4d$ gauge theories $\text{Spin}(N_c)$ and $\text{SO}(N_c)_\pm$ is in the choice of line operators. Y_{Spin} has the same magnetic quantum numbers as the minimal $4d$ 't Hooft loop of the $\text{Spin}(N_c)$ theory H^2 , wrapped on the circle. Y in the $\text{SO}(N_c)_+$ theory is related to the wrapped $4d$ 't Hooft loop H , and Y in the $\text{SO}(N_c)_-$ theory is related to the wrapped $4d$ 't Hooft-Wilson loop HW . (The wrapped 't Hooft loops are not BPS operators, but they reduce to chiral superfields in the low-energy effective action.) Correspondingly, the large gauge transformation in $\text{SO}(N_c)_\pm$ acts on Y and leads to the identification (recalling that the wrapped spinor Wilson line W is odd under this transformation)

$$\text{SO}(N_c)_+ : \quad Y \sim \frac{1}{\eta Y}, \quad \text{SO}(N_c)_- : \quad Y \sim -\frac{1}{\eta Y}. \quad (3.7)$$

In the $3d$ dimensionally reduced theory which arises as $\eta \rightarrow 0$, we are left just with the region near $Y = 0$ so there is no longer any distinction between $\text{SO}(N_c)_+$ and $\text{SO}(N_c)_-$, and there is just a single $3d$ $\text{SO}(N_c)$ gauge theory.

In both cases, in the theories with flavors, the Coulomb branch of the $4d$ gauge theory on a circle is labeled by Y with the identification (3.7). These theories also have a \mathbb{Z}_2 global symmetry taking $Y \rightarrow -Y$, which acts on the Coulomb branch (this symmetry, acting on the non-trivial wrapped 't Hooft lines, is not present in the $\text{Spin}(N_c)$ theory). Note that this symmetry remains also in the $3d$ limit, and that the baryon-monopole β of (2.19) is odd under it (while β_{Spin} of (2.20) is even).

To summarize, in the $\text{SO}(N_c)_\pm$ theories we have a gauge identification on the Coulomb branch given by (3.7), and in all 3 theories we have a \mathbb{Z}_2 global symmetry changing the sign of the non-trivial line operator wrapped on the circle. We denote this symmetry by \mathbb{Z}_2^M . It acts on the Coulomb branch as:

$$\text{Spin}(N_c) : \quad Y_{\text{Spin}} \longleftrightarrow \frac{1}{\eta^2 Y_{\text{Spin}}}, \quad \text{SO}(N_c)_\pm : \quad Y \longleftrightarrow -Y. \quad (3.8)$$

We can now look at the pure $so(N_c)$ SYM theory on a circle, and see how many vacua we have in the different $so(N_c)$ theories [5]. Solving the F-term equations of the $\text{Spin}(N_c)$ theory as above gives $(N_c - 2)$ different solutions for Z (3.2), which all have $\langle Y_{\text{Spin}} \rangle = \langle Y_1 Z \rangle = 1/\eta$. In the $\text{Spin}(N_c)$ theory, these are all fixed points of \mathbb{Z}_2^M . In the $\text{SO}(N_c)$ theories, we can additionally characterize each vacuum by the expectation value of Y , which can be $\pm 1/\sqrt{\eta}$. In the $\text{SO}(N_c)_+$ theory, each of the two possibilities maps to itself under the gauge identification (3.7), and thus each solution of (3.2) splits into two separate vacua, giving a total of $2(N_c - 2)$ supersymmetric vacua. On the other hand, in the $\text{SO}(N_c)_-$ theory, the two options for the sign of Y are mapped to each other by the gauge identification (3.7), and thus there are only $(N_c - 2)$ supersymmetric vacua. This shows the consistency of our identifications (3.7) with the counting of vacua presented in [5].

While much of our discussion remains the same also for low values of N_c , it is interesting to verify its consistency with the relations (1.3) and (1.4). For $N_c = 3$ we mentioned that $\eta_{\text{SO}(3)} = \eta^2$, so shifting the $\text{SO}(3)$ theta angle by 2π as in (1.3) takes $\eta \rightarrow -\eta$, consistent with (3.7). In this case there is no extra monopole operator Z , and Y carries the full

information about the vacuum. Each of the Spin(3) vacua at $Y_{\text{Spin}} = \pm 1/\sqrt{\eta_{\text{Spin}(3)}}$ splits into two possible SO(3) vacua at $Y = \pm\sqrt{Y_{\text{Spin}}}$. In the $4d$ SO(3)₊ SYM theory on a circle two of these are identified by the large gauge transformation (3.7), and the other two are not, and the situation is reversed in the SO(3)₋ theory. Both theories thus have 3 vacua [5]. Similarly, for $N_c = 4$ the two vacua at $Y_{\text{Spin}} = 1/\eta$ each split into two vacua with $Y = \pm 1/\sqrt{\eta}$, and the two vacua at $Y_{\text{Spin}} = -1/\eta$ each split into two vacua with $Y = \pm i/\sqrt{\eta}$. In the SO(4)₊ theory the latter two vacua are identified by (3.7), so that we have 6 overall vacua, and similarly in the SO(4)₋ theory the vacua from the first splitting are identified, and again we have 6 overall vacua [5]. The transformation shifting (say) the first theta angle in (1.4) changes the sign of $Y^{(1)}$ and thus of Y_{Spin} , and exchanges the two types of vacua, consistent with its exchanging SO(4)₊ and SO(4)₋.

4 The 4d duality on a circle

4.1 Dual theories on a circle

As discussed in [6], whenever we take two theories that are IR-dual in $4d$, compactify them on a circle of radius r , and go to low energies (compared to the scales $1/r$, Λ and $\tilde{\Lambda}$), the resulting low-energy $3d$ theories are IR-dual as well. We can start from the $4d$ duality reviewed in the introduction (with $N_c \geq 4$ and $N_f \geq N_c$), between theory A with gauge group G (which can be Spin(N_c), SO(N_c)₊ or SO(N_c)₋), with N_f chiral multiplets Q^i in the vector representation and $W_A = 0$, and theory B with gauge group \tilde{G} (which is SO(\tilde{N}_c)₋, SO(\tilde{N}_c)₊ or Spin(\tilde{N}_c), respectively, with $\tilde{N}_c = N_f - N_c + 4$), with N_f chiral multiplets q_i in the vector representation, $N_f(N_f + 1)/2$ singlets M^{ij} and a superpotential $W_B = \frac{1}{2}M^{ij}q_iq_j$.⁷ The continuous global symmetries in both theories are $SU(N_f) \times U(1)_R$, with Q in the $(\mathbf{N}_f)_{((N_f - N_c + 2)/N_f)}$ representation in theory A, and q in the $(\tilde{\mathbf{N}}_f)_{((N_c - 2)/N_f)}$ representation in theory B. Both theories have a $\mathbb{Z}_{2N_f} \times \mathbb{Z}_2^{\mathcal{C}}$ discrete symmetry. The charge conjugation symmetry $\mathbb{Z}_2^{\mathcal{C}}$, generated by \mathcal{C} , under which the baryonic operators (3.5) are odd, maps to itself under the duality. The \mathbb{Z}_{2N_f} symmetry of theory A (which is a subgroup of the anomalous axial U(1) symmetry) is generated by $g : Q \rightarrow \exp(2\pi i/2N_f)Q$, and that of theory B by $\tilde{g} : q \rightarrow \exp(-2\pi i/2N_f)q$. They are mapped by the duality as $g \leftrightarrow \tilde{g}\mathcal{C}$ [1, 2].

When we compactify the two theories on a circle, we generate extra superpotentials

$$W_A = \eta Z, \quad W_B = \frac{1}{2}M^{ij}q_iq_j + \tilde{\eta}\tilde{Z}, \quad (4.1)$$

where the $4d$ duality implies $\eta\tilde{\eta} = (-1)^{N_f - N_c}/256$ [2, 3]. We also get the compact Coulomb branches on both sides, described in the previous section. The general arguments of [6] imply that the low-energy theories with these extra superpotentials and compact Coulomb branches should be dual at low energies.

The mapping between the chiral operators involving the flavors, and the associated flat directions, is the same in the theory on a circle as in $4d$; M^{ij} in theory B is identified with Q^iQ^j , and the baryonic operators B , \mathcal{W}_α and b_{4d} are identified with \tilde{b}_{4d} , $\tilde{\mathcal{W}}_\alpha$ and \tilde{B} ,

⁷We begin by discussing connected gauge groups, we will discuss the disconnected $O(N)$ and $Pin(N)$ cases below.

respectively [1, 2]. But on a circle we have the extra Coulomb branch that we need to identify, which is not lifted by the extra superpotential (4.1), and we have the associated chiral operators that parameterize it. Note that the Coulomb branch coordinates are not charged under any continuous global symmetries, so these do not constrain their mapping.

However, there are discrete symmetries acting on the Coulomb branch (3.8), and these should map to each other under the duality (this follows from the mapping of the corresponding $4d$ non-trivial line operators [4, 5]). In fact, the full mapping of the Coulomb branches is uniquely determined by requiring that we have a single-valued meromorphic transformation between them (after identifying by the large gauge transformations (3.7) for $SO(N)$), which correctly maps the global \mathbb{Z}_2 symmetries (3.8), together with the extra requirement (for the $SO(N)_+$ case) that the mapping is non-trivial (this follows from the analysis of the next subsection). The mapping between the Coulomb branch coordinate Y_{Spin} of $\text{Spin}(N_c)$, and the coordinate \tilde{Y} of the dual $SO(\tilde{N}_c)_-$ theory, takes the form

$$\eta Y_{\text{Spin}} = \left[\frac{i + \sqrt{\tilde{\eta}} \tilde{Y}}{1 + i\sqrt{\tilde{\eta}} \tilde{Y}} \right]^2. \tag{4.2}$$

The two choices of sign for $\sqrt{\tilde{\eta}}$ are related by the global symmetry $\tilde{Y} \rightarrow -\tilde{Y}$, which maps in the dual theory to the global symmetry $Y_{\text{Spin}} \rightarrow 1/\eta^2 Y_{\text{Spin}}$. The two identified points \tilde{Y} and $(-1/\tilde{\eta} \tilde{Y})$ map to the same value of Y_{Spin} , as they should.

The map between the coordinate Y of the $SO(N_c)_-$ theory, and the coordinate \tilde{Y}_{Spin} of the dual $\text{Spin}(\tilde{N}_c)$ theory, is simply the inverse of this map (consistent with the fact that performing the duality twice should bring us back to the same point). The map between the two $SO(N)_+$ theories is just the square root of (4.2),

$$\sqrt{\eta} Y = \frac{i + \sqrt{\tilde{\eta}} \tilde{Y}}{1 + i\sqrt{\tilde{\eta}} \tilde{Y}}. \tag{4.3}$$

Again, one can check that it is consistent with the discrete gauge and global symmetries on both sides.

For low numbers of flavors, $N_c = 3$ or $\tilde{N}_c = 3$, there are extra terms appearing in the dual superpotential [2], and the mapping of η to $\tilde{\eta}$ is somewhat different because of the different instanton factor in $so(3)$ theories. Both for $N_c = 3$ and $N_c = 4$ there are also extra “triviality” relations between the $4d$ theories [2, 5, 8], because of the relations (1.3) and (1.4). In any case, the extra superpotential factors and dualities do not introduce any new issues when reduced on a circle, so we will not discuss them further here.

Note that we can get the dualities for the $O(N)$ and $Pin(N)$ theories just by gauging the charge conjugation symmetries in the dualities for $SO(N)$ and $\text{Spin}(N)$. This is true both in $4d$, and for the $4d$ theories on a circle. The main difference in the non-connected cases is that we do not have the baryon operators on both sides, so we have fewer distinguishable vacua and fewer chiral operators.

4.2 A test of the duality and of the Coulomb branch mapping

As a consistency check for our mappings (4.2), (4.3), let us analyze what happens far on the Higgs branch of theory A. In this theory we can turn on a vacuum expectation value

(VEV) for $M^{ij} = Q^i Q^j$ of rank N_c , such that the gauge group is completely broken. For each such VEV there are two supersymmetric vacua, differing by the sign of the non-zero component of the baryon $B = Q^{N_c}$ (which squares to $\det_{N_c \times N_c}(M)$). This is true both in $4d$ and in the theory on a circle. In the latter case, since the gauge group is completely broken by the quarks, we must be at a point on the Coulomb branch where classically the gauge group is unbroken, namely $Y_{\text{Spin}} = 0$ or $Y_{\text{Spin}} = \infty$ for $\text{Spin}(N_c)$, and $Y = 0$ for $\text{SO}(N_c)$.

We now discuss these vacua in theory B. The meson VEV gives a mass to N_c quarks, leaving $(N_f - N_c) = \tilde{N}_c - 4$ massless quarks. The low-energy theory is $so(\tilde{N}_c)$ with $\tilde{N}_c - 4$ massless flavors q , with a scale $\tilde{\eta}_{\text{low}} = \tilde{\eta} \det_{N_c \times N_c}(M)$. Let us first ignore the singlets and the superpotential. We then have classically a moduli space for the q 's, where generically the gauge group is broken to $so(4)$, with no light charged fields. We can think of the $so(4)$ theory as an $su(2) \oplus su(2)$ gauge theory, where each $su(2)$ factor has an instanton factor $\tilde{\eta}_{su(2)}$ related to that of the original $so(\tilde{N}_c)$ theory by $\tilde{\eta}_{su(2)} = \tilde{\eta}_{\text{low}} / \det(qq)$. In the $4d$ theory, gaugino condensation in each $su(2)$ factor leads to an effective superpotential $W = 2(\pm 1 \pm 1) \tilde{\eta}_{su(2)}^{1/2}$ for the q 's, which vanishes (and leads to a supersymmetric vacuum for the original $so(\tilde{N}_c)$ theory), if and only if we choose opposite signs for the two gaugino condensates. The operator \tilde{b}_{4d} of theory B is then equal to

$$\tilde{b}_{4d} = q^{\tilde{N}_c - 4} \tilde{W}_\alpha^2 = \pm 2 \sqrt{\det(qq)} \sqrt{\tilde{\eta}_{su(2)}}, \tag{4.4}$$

where at generic points on the moduli space this \tilde{W}_α^2 is the difference between the two gaugino bilinears of the two $su(2)$ factors. It does not vanish in the SUSY vacua. This operator obeys

$$\tilde{b}_{4d}^2 = 4 \det(qq) \tilde{\eta}_{su(2)} = 4 \tilde{\eta}_{\text{low}} = 4 \tilde{\eta} \det_{N_c \times N_c}(M). \tag{4.5}$$

Clearly, this relation is true for any value of the q 's and is an exact ring relation. Hence, in the full $4d$ theory B we find that \tilde{b}_{4d} is non-zero when M has rank N_c , and obeys a similar relation to B of theory A, so that we can identify

$$\tilde{b}_{4d} = 2 \sqrt{\tilde{\eta}} B \tag{4.6}$$

(when we normalize the superpotential of theory B to be $\frac{1}{2} Mqq$ with no extra factors). We thus identify the two supersymmetric vacua discussed above for this VEV of M in theory B. Similarly, one can show that if $\text{rank}(M) > N_c$ there is no supersymmetric vacuum in theory B.

Let us now repeat our discussion of theory B, when it is compactified on a circle. Most of the discussion is the same for $\text{Spin}(N_c)$ and $\text{SO}(N_c)$, so first we will not distinguish between them. Again we turn on a VEV of rank N_c for M , leaving in theory B $(\tilde{N}_c - 4)$ massless flavors. The matching between the high-energy and low-energy Coulomb branch coordinates implies that $\tilde{Z}_{\text{low}} = \tilde{Z} / \det_{N_c \times N_c}(M)$ (as in (2.10)), so that the low-energy superpotential (4.1) includes $\tilde{\eta} \tilde{Z} = \tilde{\eta}_{\text{low}} \tilde{Z}_{\text{low}}$. Again, let us ignore for a moment the extra $\frac{1}{2} Mqq$ superpotential in theory B, and imagine that we turn on generic VEVs for the remaining massless q 's, breaking the gauge symmetry to $so(4)$. Each of the $su(2)$ factors

in $so(4)$ now has a Coulomb branch coordinate $\tilde{Y}^{(j)}$, and, as in (2.11), the relation of the low-energy and high-energy coordinates is

$$\tilde{Y}_+ = \tilde{Y}^{(1)} + \tilde{Y}^{(2)} = \tilde{Z}_{\text{low}} \det(qq). \tag{4.7}$$

The full low-energy superpotential, including the Affleck-Harvey-Witten superpotentials [15] of the two $su(2)$'s, is thus

$$W_B = \frac{1}{\tilde{Y}^{(1)}} + \frac{1}{\tilde{Y}^{(2)}} + \frac{\tilde{\eta}_{\text{low}}}{\det(qq)} (\tilde{Y}^{(1)} + \tilde{Y}^{(2)}), \tag{4.8}$$

leading to four states with

$$\tilde{Y}^{(j)} = \pm \sqrt{\frac{\det(qq)}{\tilde{\eta}_{\text{low}}}} = \pm \frac{1}{\sqrt{\tilde{\eta}_{su(2)}}}. \tag{4.9}$$

(More precisely, in counting the physical states we should take into account the global aspects of whether our gauge group is $\text{Spin}(\tilde{N}_c)$ or $\text{SO}(\tilde{N}_c)$. We will do this momentarily.) Note that this is consistent with our discussion in the previous paragraph, and with the relation $\eta Y = S$ for $\text{SU}(2)$ theories on a circle [6]; in this case we have (in the chiral ring) $\tilde{\eta}_{su(2)} \tilde{Y}^{(j)} = S_j$. As in $4d$, in order not to turn on a superpotential for the q 's we need the expectation values to obey $\tilde{Y}^{(1)} = -\tilde{Y}^{(2)}$. So, as in $4d$, we find two supersymmetric vacua in theory B, in which

$$\tilde{Y}^{(1)} \tilde{Y}^{(2)} = -\frac{1}{\tilde{\eta}_{su(2)}} = -\frac{\det(qq)}{\tilde{\eta}_{\text{low}}}. \tag{4.10}$$

Note that the two choices for the sign of $\tilde{Y}^{(j)}$ are distinguished by the sign of the baryons $\tilde{b} = q^{\tilde{N}_c-4} \tilde{Y}_-$.⁸ The discussion above implies that \tilde{b}^2 has an expectation value equal to $4 \det_{N_c \times N_c}(M)/\tilde{\eta}$, so we can identify

$$\tilde{b} = \frac{1}{\tilde{\eta}} \tilde{b}_{4d} = \frac{2}{\sqrt{\tilde{\eta}}} B. \tag{4.11}$$

For every value of M of rank N_c in theory A there are two choices of B , and they are mapped under the duality to the two choices of \tilde{b} in theory B.

But we mentioned above that for fixed M of rank N_c and fixed sign of B there could be either one or two vacua in theory A. Are they mapped correctly? For that we should be more careful about our gauge group.

When theory A is $\text{Spin}(N_c)$, for each VEV of M of rank N_c and each sign of B there are also two choices for the monopole operator: $Y_{\text{Spin}} = 0$ or $Y_{\text{Spin}} = \infty$ (related by \mathbb{Z}_2^M). In this case theory B is $\text{SO}(\tilde{N}_c)_-$, and we have

$$\tilde{Y}^2 = \tilde{Y}_{\text{Spin}} = \frac{\det_{N_c \times N_c}(M)}{\det(qq)} \tilde{Y}_{\text{Spin,low}} = \frac{\det_{N_c \times N_c}(M)}{\det(qq)} \tilde{Y}^{(1)} \tilde{Y}^{(2)} = -\frac{1}{\tilde{\eta}}. \tag{4.12}$$

The two possible values of $\tilde{Y} = \pm \frac{i}{\sqrt{\tilde{\eta}}}$ are fixed points of the identification $\tilde{Y} \sim -\frac{1}{\tilde{\eta}\tilde{Y}}$ (3.7), and hence lead to two different vacua. They are related by the global \mathbb{Z}_2^M symmetry whose

⁸Like the operator Z , the b baryon-monopole operators in the low-energy effective action cannot be written as descendants in the $4d$ theory on a circle, even though they are descendants in $3d$ SQCD.

generator takes $\tilde{Y} \rightarrow -\tilde{Y}$, and are the dual of the choice of $Y_{\text{Spin}} = 0, \infty$ in theory A. This interpretation is consistent with our mappings (4.2).

When theory A is $\text{SO}(N_c)_-$, for each value of M there are still two values of B , but there is a single choice, $Y = 0$ (which is a fixed point of \mathbb{Z}_2^M). Correspondingly, theory B is $\text{Spin}(\tilde{N}_c)$ in which there are two values of \tilde{b} , as in (4.11), but no additional freedom in the VEV of the monopole operator \tilde{Y}_{Spin} in (4.12).

When theory A is $\text{SO}(N_c)_+$ there is freedom only in the sign of B but not in $Y = 0$. In this case theory B is $\text{SO}(\tilde{N}_c)_+$, and there are again two possible values for \tilde{Y} , which using (4.12) are $\tilde{Y} = \pm i/\sqrt{\tilde{\eta}}$. But these two vacua are actually identified by (3.7), so we have a single vacuum, agreeing with the situation in theory A (and with (4.3)).

Thus, the mapping (1.5) leads to a consistent mapping of all these vacua far on the Higgs branch in the $4d$ theory on a circle, using (4.2) and (4.3).

5 3d dualities

5.1 Reduction of the $\text{SO}(N)_+$ duality to 3d

The duality we found up to now is not purely a $3d$ duality, since it involves the compact moduli spaces that we get in the $4d$ theory on a circle. In this subsection we will see how we can turn it into a bona fide duality of $3d$ gauge theories.

Consider the duality between $\text{SO}(N_c)_+$ and $\text{SO}(\tilde{N}_c)_+$. The $\text{SO}(N_c)_+$ theory discussed above differs from the $3d$ $\text{SO}(N_c)$ theory by the superpotential (4.1) and by the compactness of the moduli space. Let us focus on theory A near the origin of the moduli space $Y = 0$, and keep $|\eta Y^2| \ll 1$. In the dual theory B, this means that we are (using (4.3)) near $\tilde{Y} = i/\sqrt{\tilde{\eta}}$ (or equivalently $\tilde{Y} = -i/\sqrt{\tilde{\eta}}$), with $|\sqrt{\tilde{\eta}}\tilde{Y} - i| \ll 1$.

If we look at the low-energy superpotential in theory A we still have $W_A = \eta Z$, though the effects of this superpotential are very small in the region we are now discussing. In theory B we break the $\text{SO}(\tilde{N}_c)$ theory at this value of \tilde{Y} to $\text{SO}(N_f - N_c + 2) \times \text{SO}(2)$.⁹ The operator \tilde{Y} maps at low energies to the Coulomb branch coordinate V_+ of the $\text{SO}(2)$, and we can consider a new \tilde{Y}_{low} Coulomb branch coordinate for the low-energy $\text{SO}(N_f - N_c + 2)$ (defined as in (2.8)). In the low-energy superpotential of theory B we have contributions from the original W_B of (4.1). The semi-classical forms of the monopole operators (2.6), (2.8), imply that $\tilde{Z} = \tilde{Y}\tilde{Y}_{\text{low}}$. In addition we have an Affleck-Harvey-Witten superpotential related to the breaking of the $\text{SO}(\tilde{N}_c)$ gauge group, which is proportional to $\tilde{Y}_{\text{low}}/\tilde{Y}$. Thus, the full low-energy superpotential near this point is

$$W_B = \frac{1}{2}Mqq + \frac{\tilde{Y}_{\text{low}}}{\tilde{Y}} + \tilde{\eta}\tilde{Y}_{\text{low}}\tilde{Y}. \tag{5.1}$$

⁹Naively, one may think that the symmetry is broken to $S(O(N_f - N_c + 2) \times O(2))$, with an extra \mathbb{Z}_2 factor. However, the identification on the moduli space (3.7) uses the Weyl transformation $\sigma_1 \rightarrow -\sigma_1$, which is in $O(2)$, but not in $\text{SO}(2)$. This means that the unbroken gauge symmetry in the $\text{SO}(\tilde{N}_c)_+$ theory around $\tilde{Y} = i/\sqrt{\tilde{\eta}}$ is only $\text{SO}(N_f - N_c + 2) \times \text{SO}(2)$.

We can now use the mapping (4.3) between \tilde{Y} and Y to rewrite this in terms of Y , which is now an elementary field in theory B:

$$W_B = \frac{1}{2}Mqq + 4\sqrt{\eta\tilde{\eta}}\tilde{Y}_{\text{low}}\frac{Y}{1+\eta Y^2}. \quad (5.2)$$

The choice of sign for the square root is arbitrary (the two choices are related by the global symmetry $Y \rightarrow -Y$). We can now simply take $\eta \rightarrow 0$ on both sides (keeping $\eta\tilde{\eta} = (-1)^{N_f-N_c}/256$ fixed); in theory A this is allowed since the effect of the superpotential smoothly goes to zero in the region we are keeping, and in theory B the same is also true (since $|\eta Y^2| \ll 1$). In this limit we find in theory A an $\text{SO}(N_c)$ 3d theory with a non-compact Coulomb branch and with $W_A = 0$, and in theory B an $\text{SO}(N_f - N_c + 2)$ 3d theory, again with a non-compact Coulomb branch, and with

$$W_B = \frac{1}{2}Mqq + \frac{i^{N_f-N_c}}{4}\tilde{Y}_{\text{low}}Y, \quad (5.3)$$

where Y is now an elementary singlet in this low-energy theory, and \tilde{Y}_{low} is its standard Coulomb branch coordinate (2.8).

We can now lift this to a high-energy 3d duality between these two gauge theories, by replacing \tilde{Y}_{low} by the appropriate microscopic monopole operator \tilde{y} of $\text{SO}(N_f - N_c + 2)$, and the superpotential of theory B with

$$W_B = \frac{1}{2}Mqq + \frac{i^{N_f-N_c}}{4}\tilde{y}Y. \quad (5.4)$$

Note that unlike in other cases discussed in [6], here we did not need to perform any real mass deformation in order to obtain the duality for the standard 3d SQCD theory from 4d, but just to take the 3d limit carefully. In the 3d limit we have an extra global $\text{U}(1)_A$ symmetry, that was anomalous in 4d. The quantum numbers of the various operators are consistent with the superpotential (5.4); using a specific choice for the 3d R-symmetry, they are

	$\text{SO}(N_c)$	$\text{SU}(N_f)$	$\text{U}(1)_A$	$\text{U}(1)_R$	$\mathbb{Z}_2^{\mathcal{C}}$	$\mathbb{Z}_2^{\mathcal{M}}$
Q	\mathbf{N}_c	\mathbf{N}_f	1	0		
M	$\mathbf{1}$	$\mathbf{N}_f(\mathbf{N}_f + \mathbf{1})/2$	2	0	+1	+1
Y	$\mathbf{1}$	$\mathbf{1}$	$-N_f$	$N_f - N_c + 2$	+1	-1
B	$\mathbf{1}$	$(\mathbf{N}_f)_A^{N_c}$	N_c	0	-1	+1
β	$\mathbf{1}$	$(\mathbf{N}_f)_A^{N_c-2}$	$N_c - N_f - 2$	$N_f - N_c + 2$	-1	-1

(5.5)

in theory A, and

	$\text{SO}(N_f - N_c + 2)$	$\text{SU}(N_f)$	$\text{U}(1)_A$	$\text{U}(1)_R$	$\mathbb{Z}_2^{\mathcal{C}}$	$\mathbb{Z}_2^{\tilde{\mathcal{M}}}$
q	$\mathbf{N}_f - \mathbf{N}_c + \mathbf{2}$	$\bar{\mathbf{N}}_f$	-1	1		
M	$\mathbf{1}$	$\mathbf{N}_f(\mathbf{N}_f + \mathbf{1})/\mathbf{2}$	2	0	$+1$	$+1$
Y	$\mathbf{1}$	$\mathbf{1}$	$-N_f$	$N_f - N_c + 2$	$+1$	-1
\tilde{y}	$\mathbf{1}$	$\mathbf{1}$	N_f	$N_c - N_f$	$+1$	-1
\tilde{B}	$\mathbf{1}$	$(\mathbf{N}_f)_A^{N_c-2}$	$N_c - N_f - 2$	$N_f - N_c + 2$	-1	$+1$
$\tilde{\beta}$	$\mathbf{1}$	$(\mathbf{N}_f)_A^{N_c}$	N_c	0	-1	-1

(5.6)

in theory B. Note that $\text{SU}(N_f) \times \text{U}(1)_A$ is really $\text{U}(N_f)$. $(\mathbf{N}_f)_A^{N_c-2}$ and $(\mathbf{N}_f)_A^{N_c}$ denote totally antisymmetric products. $\mathbb{Z}_2^{\mathcal{C}}$ is the charge conjugation symmetry, generated by \mathcal{C} , and $\mathbb{Z}_2^{\mathcal{M}}$ and $\mathbb{Z}_2^{\tilde{\mathcal{M}}}$ are the global symmetries (3.8), generated by \mathcal{M} and $\tilde{\mathcal{M}}$, respectively. We included their action only on the gauge singlets. The composites \tilde{B} and $\tilde{\beta}$ in theory B are defined as in theory A (see (2.19)), and their identification in theory A will be discussed below.

The three symmetries $\text{U}(1)_A$, $\mathbb{Z}_2^{\mathcal{C}}$ and $\mathbb{Z}_2^{\tilde{\mathcal{M}}}$ are actually not independent. In theory A with gauge group $\text{SO}(N_c)$, the action of $e^{\pi i A}$ (which is in $\text{SU}(N_f)$ for even values of N_f) on Q is part of the gauge group for even values of N_c , and is the same as \mathcal{C} for odd values of N_c . The action of $e^{\pi i A}$ on Y is the same as \mathcal{M}^{N_f} . Thus, on gauge-invariant operators we have

$$e^{\pi i A} \mathcal{C}^{N_c} \mathcal{M}^{N_f} = 1, \tag{5.7}$$

and $\text{SU}(N_f) \times \text{U}(1)_A \times \mathbb{Z}_2^{\mathcal{C}} \times \mathbb{Z}_2^{\tilde{\mathcal{M}}}$ is really $(\text{U}(N_f) \times \mathbb{Z}_2^{\mathcal{C}} \times \mathbb{Z}_2^{\tilde{\mathcal{M}}})/\mathbb{Z}_2$. In the dual theory we have $e^{\pi i A} \mathcal{C}^{N_f - N_c + 2} \tilde{\mathcal{M}}^{N_f} = 1$, implying that for odd values of N_f ,

$$\mathcal{M} \longleftrightarrow \tilde{\mathcal{M}} \mathcal{C}. \tag{5.8}$$

We will see below that this must be true for even values of N_f as well.

The duality we find is very similar to the one discussed for $O(N_c)$ theories (more precisely, $O(N_c)_+$ theories) in [10–12]. Indeed, if we now gauge the charge conjugation symmetry $\mathbb{Z}_2^{\mathcal{C}}$ on both sides, we obtain precisely that duality, so our discussion is a derivation of this duality from $4d$. But we obtain a duality also for $\text{SO}(N_c)$ groups, meaning that there should be a consistent mapping of the charge-conjugation-odd baryons between the two sides. We can follow what happens to the $4d$ baryon mapping by our reduction procedure. In $4d$ the baryon $B = Q^{N_c}$ mapped to $\tilde{b}_{4d}/2\sqrt{\tilde{\eta}}$ (4.11). In the reduction on the circle we say that this first becomes equal to $\sqrt{\tilde{\eta}}\tilde{b}/2$ (4.11), where the latter operator (2.23) involves a monopole operator in $so(4)$. When we go onto the Coulomb branch as above, this monopole operator \tilde{Y}_- becomes $i/\sqrt{\tilde{\eta}}$ (from the VEV of \tilde{Y}) times the odd monopole operator \tilde{W}_- of $\text{SO}(2)$, so we find that B maps to $i\tilde{\beta}/2$, with $\tilde{\beta}$ defined as in (2.19). This is consistent with their global symmetry quantum numbers as in (5.5), (5.6).

The $4d$ operator b_{4d} goes to zero in the $\eta \rightarrow 0$ limit that we described, as does its $4d$ dual $\tilde{B}_{4d} = q^{N_f - N_c + 4}$ (since only $(N_f - N_c + 2)$ components of the quarks remain massless in

the limit we took in theory B). However, we now get a new relation (required by consistency of the duality), mapping β to the $3d$ baryon $\tilde{B} = q^{N_f - N_c + 2}$. We cannot derive this duality directly from $4d$, but on the part of the moduli space where we break both gauge groups to $SO(2)$, it follows by dualizing the vector multiplets (3.5) in the $4d$ relation $\mathcal{W}_\alpha \leftrightarrow \tilde{\mathcal{W}}_\alpha$ into chiral multiplets (taking into account again the VEV of \tilde{Y} in theory B). It is also consistent with the global symmetries as in (5.5), (5.6). We conclude that the baryons map in the $3d$ duality between $SO(N_c)$ and $SO(N_f - N_c + 2)$ by

$$\{B, \beta\} \quad \longleftrightarrow \quad \left\{ \frac{i}{2} \tilde{\beta}, -2i \tilde{B} \right\}. \quad (5.9)$$

Note that this mapping requires that the \mathbb{Z}_2^M symmetry (3.8), which takes $B \rightarrow B$ and $\beta \rightarrow -\beta$, maps under the duality by (5.8) for all values of N_f . As we mentioned above, in the $3d$ theory b and Z are not chiral, so they do not have a simple mapping under the duality.

We can perform many tests of this duality, comparing moduli spaces, chiral operators, deformations, and so on, but most of these tests are identical to tests of the $O(N_c)_+$ duality that were already performed in [12]. We can find new tests by involving also the baryon operators. For instance, suppose that we turn on a VEV for M of rank N_c , as in our discussion of the previous section. In theory A we still have two vacua for every such M , with $B^2 = \det_{N_c \times N_c}(M)$. In theory B we now give a mass to N_c quarks, so that we are left at low energies with $N_f - N_c = \tilde{N}_c - 2$ massless quarks, and with a low-energy Coulomb branch coordinate $\tilde{Y}_{N_f - N_c} = \tilde{y} / \sqrt{\det_{N_c \times N_c}(M)}$. Our discussion around (2.22) implies that in this low-energy theory, in which the superpotential sets its meson $qq = 0$, there is a relation $\tilde{\beta}_{N_f - N_c}^2 = -4$. Translating this into the high-energy theory (using (2.10)) we find $\tilde{\beta}^2 = -4 \det_{N_c \times N_c}(M)$, so that we can indeed identify $\tilde{\beta}$ with $(-2iB)$.

For low values of N_c and of $N_f - N_c$ there are slight modifications of this discussion, as in the $4d$ duality [2] and in the analysis of the pure $3d$ theory [12], but these do not raise any new issues so we will not discuss them in detail here.

5.2 Reduction of the $Spin(N_c) \leftrightarrow SO(\tilde{N}_c)_-$ duality to $3d$

We can similarly obtain the dual of the $3d$ $Spin(N_c)$ theory, by starting from the $4d$ duality between $Spin(N_c)$ and $SO(\tilde{N}_c)_-$. We can again focus on the same points $Y_{Spin} = 0$ and $\tilde{Y} = -i/\sqrt{\tilde{\eta}}$ in the moduli space, and obtain the low-energy superpotential (5.1). However, now we are at a fixed point of (3.7), so the discussion in footnote 9 implies that the unbroken gauge symmetry in theory B is $S(O(N_f - N_c + 2) \times O(2))$. The moduli space coordinate \tilde{Y} is the monopole operator of the $SO(2)$ factor; when expressed in terms of \tilde{Y} , the $SO(2) \subset O(2)$ is not visible. Focusing on the region around $\tilde{Y} = -i/\sqrt{\tilde{\eta}}$, it is convenient to change variables from the approximate free field \tilde{Y} to y defined by

$$4\sqrt{\eta\tilde{\eta}} \frac{y}{1 + \eta y^2} = \frac{1}{\tilde{Y}} + \tilde{\eta} \tilde{Y}. \quad (5.10)$$

Now we expand around $y = 0$, the mapping from theory A to theory B (4.2) identifies $Y_{Spin} = y^2$, and the extra \mathbb{Z}_2 gauge identification acts as $y \rightarrow -y$. This \mathbb{Z}_2 also acts on the

$SO(N_f - N_c + 2)$ theory as charge conjugation, and it also changes the sign of the monopole operator $\tilde{y} = \tilde{Y}_{\text{low}}$ of $SO(N_f - N_c + 2)$. Thus, we recognize the gauge theory we get as an $O(N_f - N_c + 2)_-$ theory, with the elementary field y odd under $\mathbb{Z}_2^{\tilde{C}} \subset O(N_f - N_c + 2)$. The low energy superpotential after taking $\eta \rightarrow 0$ is

$$W_B = \frac{1}{2} M q q + \frac{i^{N_f - N_c}}{4} \tilde{y} y . \tag{5.11}$$

The bottom line is that the dual of the $\text{Spin}(N_c)$ SQCD theory is similar to (5.4), but the dual gauge group is $O(N_f - N_c + 2)_-$, and y and \tilde{y} are odd under its $\mathbb{Z}_2^{\tilde{C}}$ subgroup.

The $\mathbb{Z}_2^{\tilde{C}}$ global symmetry of theory A is mapped under this duality to $\mathbb{Z}_2^{\tilde{M}}$. In the $\text{Spin}(N_c)$ theory B is a $\mathbb{Z}_2^{\tilde{C}}$ -odd gauge-invariant operator, while Y and β do not exist. In the dual $O(N_f - N_c + 2)_-$ theory $\tilde{\beta}$ is a $\mathbb{Z}_2^{\tilde{M}}$ -odd gauge-invariant operator (mapped to B), while y , \tilde{y} and \tilde{B} are not gauge-invariant. The operator β_{Spin} of (2.20) is present in theory A, and maps to the gauge-invariant operator $\tilde{B}y$ in theory B.

In the discussion above we expanded around the point $Y_{\text{Spin}} = 0$, which is mapped to $\tilde{Y} = -i/\sqrt{\tilde{\eta}}$. Instead we could expand around $\tilde{Y} = 0$, which corresponds to $Y_{\text{Spin}} = -1/\eta$. Here the $SO(N_f - N_c + 4)$ gauge group of theory B is unbroken, while the gauge group of theory A is broken as $\text{Spin}(N_c) \rightarrow (\text{Spin}(N_c - 2) \times \text{Spin}(2))/\mathbb{Z}_2$, where the \mathbb{Z}_2 acts on the spinors in both groups. Next we analyze the low energy dynamics of theory A, as we did in theory B above, focusing on the $\text{Spin}(2)$ dynamics. It is important that there are no massless fields charged under this group. Hence, its Wilson loops become trivial at long distances. Further, we can map each $(\text{Spin}(N_c - 2) \times \text{Spin}(2))/\mathbb{Z}_2$ bundle to an $SO(N_c - 2)$ bundle by simply ignoring the $\text{Spin}(2)$ transition functions. Hence, when we integrate out the $\text{Spin}(2)$ dynamics we are left with an $SO(N_c - 2)$ gauge theory. The dual of the $\text{Spin}(2)$ gauge field, Y_{Spin} , is an almost free field, which we can replace using (4.2) by an elementary field \tilde{Y} , identified with the monopole operator of theory B. Finally, we can identify the gauge-invariant operator $(iY_{\text{low,Spin}}/\sqrt{\tilde{\eta}}Z)$ in theory A (where $Y_{\text{low,Spin}}$ is the Coulomb branch coordinate of $\text{Spin}(N_c - 2)$) with Y_{low} of $SO(N_c - 2)$, and, as in our analysis above, the various monopole-instantons couple \tilde{Y} to Y_{low} . For $\tilde{\eta} \rightarrow 0$ we find a superpotential term proportional to $Y_{\text{low}}\tilde{Y}$, as in (5.3). The duality we derive this way is precisely the inverse of the $SO(N)$ duality that we derived in section 5.1. This is a non-trivial consistency check on our web of dualities, because in section 5.1 we derived this duality from the compactification of a different 4d duality.

We can also obtain a dual for $\text{Pin}(N_c)$, by gauging the global symmetry $\mathbb{Z}_2^{\tilde{M}}$ in the duality for $O(N_c)_+$ groups. The fact that in the $O(N_c)_+$ duality the symmetry $\mathbb{Z}_2^{\tilde{M}}$ maps to itself implies that the $\text{Pin}(N_c)$ theory is dual to a $\text{Pin}(N_f - N_c + 2)$ theory.

5.3 Dualities with Chern-Simons terms

As in [6], we can flow from the duality above to a duality with Chern-Simons terms. We can obtain an $SO(N_c)$ theory with N_f flavors and a Chern-Simons term at level $k > 0$ by starting from the theory with $N_f + k$ flavors and giving k flavors a positive real mass, by turning on a background field for the $U(N_f + k)$ global symmetry. In the dual $SO(N_f + k - N_c + 2)$ theory, this maps to giving k flavors a negative real mass, giving the

mesons they couple to a positive real mass, and also giving a negative real mass to the singlet Y . Integrating out the massive fields we find an $SO(N_f + k - N_c + 2)$ theory with N_f flavors, level $(-k)$, and a $W_B = \frac{1}{2}Mqq$ superpotential. This is precisely the duality conjectured in [24] for the $O(N)$ theories (more precisely, $O(N)_+$ theories), and here we see that it is true also for $SO(N)$.

The difference between $O(N)_+$ and $SO(N)$ is that now we need to understand also how to map the baryon operators in the two sides, and this is more complicated (as in the discussion of $SU(N_c)$ Chern-Simons-matter theories in [6]) since they involve monopoles. In theory A we still have the baryon operator $B = Q^{N_c}$, while the baryon-monopole $\beta = Q^{N_c-2}W_-$ is no longer gauge-invariant in the presence of the Chern-Simons term, and similarly in theory B. We claim, similar to what we found for $SU(N_c)$ Chern-Simons-matter theories, that the dual of B is now given by a monopole operator $\tilde{\beta}' = q^{N_f - N_c} \tilde{W}_- (\tilde{W}_\alpha)^k$, which is gauge-invariant. The quarks are contracted with an epsilon symbol, and break $SO(N_f + k - N_c + 2)$ to $SO(k + 2)$. The monopole operator \tilde{W}_- breaks $SO(k + 2)$ to $SO(k) \times U(1)$, and because of the Chern-Simons term it carries a charge $(-k)$ under the $U(1)$. The gluinos \tilde{W}_α are off-diagonal gluinos in $SO(k) \times U(1)$ which cancel this charge, and carry k different vector indices of $SO(k)$ that are contracted by an epsilon symbol, such that the total operator is gauge-invariant. One can verify that the global symmetry charges of this $\tilde{\beta}'$ exactly match with those of B . Similarly, we can construct an operator β' in theory A that matches with $\tilde{B} = q^{N_f + k - N_c + 2}$.

In theory A we have the relation $B^2 = \det_{N_c \times N_c}(M)$. To see this in theory B we turn on a VEV of rank N_c for M , leaving $N_f - N_c$ massless flavors q , and we then ignore for a moment the superpotential and imagine giving an expectation value to the remaining massless flavors. This breaks the gauge group to $SO(k + 2)$ with level $(-k)$ and no massless flavors. At low energies this is a purely topological theory, in which we can construct a singlet operator $\tilde{\beta}' = \tilde{W}_- \tilde{W}_\alpha^k$ as above, which is charged under the charge conjugation symmetry of this theory, and argue that it squares to one (similar to our discussion of the $SU(k)$ theory at level $(-k)$ in [6]). Lifting this to the high-energy theory using (2.10) we get precisely the expected relation (which turns out to be independent of the VEVs of the q 's, so it is valid even for $q = 0$).

For $N_f = 0$ our duality reduces to a duality of pure supersymmetric Chern-Simons theories, $SO(N_c)_k$ being identified with $SO(|k| - N_c + 2)_{-k}$. This is just the standard level-rank duality of $SO(N)$ Chern-Simons theories. At low energies we can integrate out the gauginos, shifting the $SO(N_c)$ level to $k \rightarrow k - (N_c - 2)\text{sign}(k)$. We then obtain the standard level-rank relation [25–27]

$$SO(n)_m \longleftrightarrow SO(m)_{-n}, \tag{5.12}$$

for $n, m > 0$, that can be proven by studying nm real fermions in two dimensions.

Similarly, we can flow from our $\text{Spin}(N_c)_k$ duality to find a duality between $\text{Spin}(N_c)_k$ and $(O(N_f + |k| - N_c + 2)_{-k})_{-k}$ (and a corresponding non-supersymmetric level-rank duality taking $\text{Spin}(n)_m$ to $(O(m)_{-})_{-n}$), and a duality between the supersymmetric $\text{Pin}(N_c)_k$ and $\text{Pin}(N_f + |k| - N_c + 2)_{-k}$ theories (with a level-rank duality taking $\text{Pin}(n)_m$ to $\text{Pin}(m)_{-n}$).

5.4 The special case of $SO(2) = U(1)$ with $N_f = 2$: a triality of dualities

We now have two different dualities for $SO(2) = U(1)$ gauge theories with N_f flavors, which we refer to as theory A. First, we can view the gauge group as $U(1)$ and find a dual theory based on $U(N_f - 1)$ [28]. We will refer to this dual theory as B1. Alternatively, as in this paper, we can view it as $SO(2)$ and find a dual theory based on $SO(N_f)$. We will refer to it as B2. The B1 dual exhibits the full $SU(N_f) \times SU(N_f) \times U(1)_A \times U(1)_J \times U(1)_R$ global symmetry, while the B2 dual exhibits explicitly only $SU(N_f) \times U(1)_A \times U(1)_R$, and the other symmetries arise as accidental symmetries at low energies.

The $SO(2)$ theory with $N_f = 2$ deserves special attention. In this case the gauge groups of theories A, B1 and B2 are all $U(1)$, and they all have two flavors. Furthermore, in this case there is also a mirror theory, that also has gauge group $U(1)$ and two flavors [29, 30]–[13, 31]. We will refer to this theory as B3.

Let us understand the relation between these dual descriptions (see also [12]). We begin with theory A. We can think of it either as a $U(1)$ theory with two flavors $Q^a, \tilde{Q}^{\tilde{a}}$ ($a, \tilde{a} = 1, 2$), or as an $SO(2)$ theory with two doublets P_i ($i = 1, 2$). Let us work out the translation between these two languages. In the $U(1)$ description of this theory, the chiral operators are the magnetic monopoles V_{\pm} (2.13) and four mesons $\hat{M}^{a\tilde{b}} = Q^a \tilde{Q}^{\tilde{b}}$. The translation to $SO(2)$ variables, if we keep the standard normalization for the kinetic terms, is by

$$Q^i = \frac{1}{\sqrt{2}}(P_i^1 + iP_i^2), \quad \tilde{Q}^i = \frac{1}{\sqrt{2}}(P_i^1 - iP_i^2). \quad (5.13)$$

Defining the standard $SO(2)$ mesons $M_{ij} = P_i \cdot P_j$, the symmetric part of \hat{M} is related to M by $\hat{M}_{ij} + \hat{M}_{ji} = M_{ij}$. The anti-symmetric part of \hat{M} is related to the $SO(2)$ baryon $B \equiv P_1^1 P_2^2 - P_1^2 P_2^1$ by $\hat{M}_{12} - \hat{M}_{21} = -iB$. The natural monopole-related operators in the $SO(2)$ language are

$$Y \equiv V_+ + V_-, \quad \beta \equiv V_+ - V_-. \quad (5.14)$$

The former is the basic monopole in the $SO(2)$ language, and the latter is the baryon operator β (2.19) in this special case. The charges of the different objects under the global $U(1)_A \times U(1)_R$ symmetry that is visible in all descriptions are (using our standard conventions):

	$U(1)_A$	$U(1)_R$
Q, \tilde{Q}, P	1	0
Y, β	-2	2
B, M	2	0

(5.15)

The dual description which has all the symmetries of theory A manifest is B1. This is a $U(1)$ theory with two flavors $q_a^{(1)}, \tilde{q}_{\tilde{a}}^{(1)}$ ($a, \tilde{a} = 1, 2$) and with additional singlet fields $\hat{M}_{(1)}^{a\tilde{b}}$ and $V_{\pm}^{(1)}$. The superpotential is [28]

$$W_{B1} = q^{(1)} \hat{M}^{(1)} \tilde{q}^{(1)} + V_+^{(1)} \tilde{V}_-^{(1)} + V_-^{(1)} \tilde{V}_+^{(1)}. \quad (5.16)$$

For comparison with other duals it will be useful to translate this to the SO(2) language, as we did above. We define as above

$$\tilde{Y}^{(1)} \equiv \tilde{V}_+^{(1)} + \tilde{V}_-^{(1)}, \quad \tilde{\beta}^{(1)} \equiv \tilde{V}_+^{(1)} - \tilde{V}_-^{(1)}, \quad \tilde{B}^{(1)} \equiv i(q_1^{(1)}\tilde{q}_2^{(1)} - q_2^{(1)}\tilde{q}_1^{(1)}). \quad (5.17)$$

Translating the quarks q, \tilde{q} to SO(2) quarks p as above, the superpotential (5.16) becomes

$$W_{B1} = \frac{1}{4}M_{ab}^{(1)} p_a^{(1)} p_b^{(1)} - \frac{1}{2}B^{(1)} \tilde{B}^{(1)} + \frac{1}{2}Y^{(1)} \tilde{Y}^{(1)} - \frac{1}{2}\beta^{(1)}\tilde{\beta}^{(1)}, \quad (5.18)$$

with the singlets $B^{(1)}$ and $M^{(1)}$ related to $\hat{M}^{(1)}$, and the singlets $\beta^{(1)}$ and $Y^{(1)}$ related to $V_{\pm}^{(1)}$, as in theory A. These singlets are identified with the corresponding operators in theory A. The $U(1)_A \times U(1)_R$ charges of the different objects are:

	$U(1)_A$	$U(1)_R$
$q^{(1)}, \tilde{q}^{(1)}, p$	-1	1
$\tilde{Y}^{(1)}, \tilde{\beta}^{(1)}$	2	0
$\tilde{B}^{(1)}$	-2	2

(5.19)

The normalization of the first term in (5.18) is half of the normalization in the standard SO(N_c) duality (5.4), and it will be easier to compare them if we have the same normalization in both cases. Since the p 's do not appear in any gauge-invariant chiral operator, we can simply rescale them to new variables $\hat{p}_i^{(1)} = p_i^{(1)}/\sqrt{2}$. This also rescales the baryon $\tilde{B}^{(1)}$ to $\tilde{\tilde{B}}^{(1)} = \tilde{B}^{(1)}/2$, and because of the relation of the quantum numbers of the monopoles to those of the quarks, the latter are also rescaled to $\tilde{\tilde{Y}}^{(1)} = 2\tilde{Y}^{(1)}$, $\tilde{\tilde{\beta}}^{(1)} = 2\tilde{\beta}^{(1)}$. We can now write (5.18) as

$$W_{B1} = \frac{1}{2}M_{ab}^{(1)} \hat{p}_a^{(1)} \hat{p}_b^{(1)} - B^{(1)} \tilde{\tilde{B}}^{(1)} + \frac{1}{4}Y^{(1)} \tilde{\tilde{Y}}^{(1)} - \frac{1}{4}\beta^{(1)}\tilde{\tilde{\beta}}^{(1)}. \quad (5.20)$$

The dual description B2, with gauge group SO(2), is quite similar to B1. The difference is that we do not have the singlet fields B and β , and the superpotential is (5.4)

$$W_{B2} = \frac{1}{2}M_{ab}^{(2)} p_a^{(2)} p_b^{(2)} + \frac{1}{4}Y^{(2)} \tilde{Y}^{(2)}. \quad (5.21)$$

The map between the chiral operators here is

$$M \rightarrow M^{(2)}, \quad B \rightarrow \frac{i}{2}\tilde{\beta}^{(2)}, \quad \beta \rightarrow -2i\tilde{B}^{(2)}, \quad Y \rightarrow Y^{(2)}. \quad (5.22)$$

Note that in description B2 the $U(1)_J$ symmetry is not present in the UV gauge theory, as the singlet Y mixes with β under this symmetry. Moreover only an $SU(2) \subset SU(2) \times SU(2)$ flavor symmetry is visible in the UV. The symmetries broken in this UV description should appear as accidental symmetries of the IR physics.

Finally, in description B3 we do not have M_{12} and Y , and the superpotential is

$$W_{B3} = \hat{M}_{(3)}^{11} q_1^{(3)} \tilde{q}_1^{(3)} + \hat{M}_{(3)}^{22} q_2^{(3)} \tilde{q}_2^{(3)} = \frac{1}{4}M_{11}^{(3)} p_1^{(3)} p_1^{(3)} + \frac{1}{4}M_{22}^{(3)} p_2^{(3)} p_2^{(3)}. \quad (5.23)$$

Using the fact that mirror symmetry exchanges the monopoles V_{\pm} with the off-diagonal mesons $\hat{M}^{12}, \hat{M}^{21}$ with coefficient one, the map between the chiral operators here is

$$M_{\ell\ell} \rightarrow M_{\ell\ell}^{(3)}, \quad B \rightarrow i\tilde{\beta}^{(3)}, \quad \beta \rightarrow -i\tilde{B}^{(3)}, \quad Y \rightarrow p_1^{(3)} p_2^{(3)}, \quad M_{12} \rightarrow \tilde{Y}^{(3)}. \quad (5.24)$$

Here $U(1)_J$ is present in the UV description and is identified with part of the dual flavor group, but only $U(1) \times U(1) \subset SU(2) \times SU(2)$ is a symmetry of the UV theory. As in the B1 theory, it is convenient to rescale the p 's by $\sqrt{2}$, and in the rescaled variables defined as above we have

$$W_{B3} = \frac{1}{2} M_{11}^{(3)} \hat{p}_1^{(3)} \hat{p}_1^{(3)} + \frac{1}{2} M_{22}^{(3)} \hat{p}_2^{(3)} \hat{p}_2^{(3)}, \quad (5.25)$$

with the new mapping

$$M_{\ell\ell} \rightarrow M_{\ell\ell}^{(3)}, \quad B \rightarrow \frac{i}{2} \tilde{\beta}^{(3)}, \quad \beta \rightarrow -2i\tilde{B}^{(3)}, \quad Y \rightarrow 2\hat{p}_1^{(3)} \hat{p}_2^{(3)}, \quad M_{12} \rightarrow \frac{1}{2} \tilde{Y}^{(3)}. \quad (5.26)$$

Note that the mapping between the B 's and β 's is now the same as in (5.22).

Let us now relate these theories, by understanding their deformations. We claim that the IR superconformal field theory that all these theories flow to has eight marginal deformations, and that two of them are exactly marginal. In the B1 description the eight marginal operators are $\hat{M}_{(1)}^{a\bar{a}} V_{\pm}^{(1)}$. Denote their coupling constants by $\lambda_{a\bar{a}}^{\pm}$. The space of exactly marginal deformations is generally given by the (complexified) quotient of the space of marginal deformations by the global symmetries, which act on it non-trivially [32]. This can be found by noting the invariant combinations constructed out of $\lambda_{a\bar{a}}^{\pm}$. All these deformations preserve $U(1)_A$, but they are charged under $SU(2) \times SU(2) \times U(1)_J$. There are two non-trivial invariants of this group: $\epsilon^{ab} \epsilon^{\bar{a}\bar{b}} \lambda_{a\bar{a}}^+ \lambda_{b\bar{b}}^-$ and $\left(\epsilon^{ab} \epsilon^{\bar{a}\bar{b}} \lambda_{a\bar{a}}^+ \lambda_{b\bar{b}}^+ \right) \left(\epsilon^{cd} \epsilon^{\bar{c}\bar{d}} \lambda_{c\bar{c}}^- \lambda_{d\bar{d}}^- \right)$. This shows that there are two exactly marginal deformations. Equivalently, the global symmetry $SU(2) \times SU(2) \times U(1)_J$ is not completely broken on the space of couplings, but a $U(1)$ always remains (which is a subgroup of the diagonal $SU(2)$).

By a global symmetry transformation, we can choose the two exactly marginal deformations to be given by (in the $SO(2)$ language)

$$\delta W = \gamma B \beta + \rho M_{12} Y. \quad (5.27)$$

Note that these deformations are invariant under \mathbb{Z}_2^C , but they break \mathbb{Z}_2^M (3.8).

Suppose we first add the term with γ to theory A. In the B1 dual description, the superpotential becomes after integrating out the massive singlet fields

$$W_{B1} + \gamma B^{(1)} \beta^{(1)} \rightarrow \frac{1}{2} M_{ab}^{(1)} \hat{p}_a^{(1)} \hat{p}_b^{(1)} + \frac{1}{4} Y \tilde{Y}^{(1)} - \frac{1}{4\gamma} \tilde{\beta}^{(1)} \tilde{B}^{(1)}. \quad (5.28)$$

This is exactly the same as the superpotential W_{B2} , deformed by $\frac{-1}{4\gamma} \tilde{B}^{(2)} \tilde{\beta}^{(2)}$. On the other hand, adding $\gamma B \beta$ to theory A translates in theory B2 to adding $\gamma \tilde{B}^{(2)} \tilde{\beta}^{(2)}$. These two descriptions should be equivalent, so we conclude that our deformed IR superconformal field theory possesses an exact duality under $\gamma \longleftrightarrow (-1/4\gamma)$.

Next we add the term with ρ to theory A. The superpotential in the dual B2 description becomes, after integrating out massive singlet fields,

$$W_{B2} + \rho M_{12}^{(2)} Y^{(2)} \rightarrow \frac{1}{2} M_{11}^{(2)} p_1^{(2)} p_1^{(2)} + \frac{1}{2} M_{22}^{(2)} p_2^{(2)} p_2^{(2)} - \frac{1}{4\rho} \tilde{Y}^{(2)} p_1^{(2)} p_2^{(2)}. \quad (5.29)$$

This is exactly the same as the superpotential W_{B3} , deformed by $-\frac{1}{4\rho} \tilde{Y}^{(3)} \hat{p}_1^{(3)} \hat{p}_2^{(3)}$; as we mentioned, the baryon mappings are also consistent. On the other hand, adding $\rho M_{12} Y$ to theory A translates in theory B3 to adding $\rho \tilde{Y}^{(3)} \hat{p}_1^{(3)} \hat{p}_2^{(3)}$. Thus we conclude that our IR superconformal field theory also possesses an exact duality $\rho \longleftrightarrow (-1/4\rho)$.

Assuming the three dualities, we deduced that the exactly marginal couplings enjoy dualities taking $\gamma \longleftrightarrow (-1/4\gamma)$ and $\rho \longleftrightarrow (-1/4\rho)$. Alternatively, if one could prove the duality of the marginal deformations, one could deduce all three duals of U(1) with two flavors from knowing any one of them.

6 Partition functions and indices for $so(N)$ dualities

A set of useful checks of dualities is given by comparisons of supersymmetric partition functions of the putative dual pair: if the two $3d$ UV theories describe the same IR physics, their S^3 and $S^2 \times S^1$ supersymmetric partition functions should agree. In this section we will discuss these checks for the dualities of the previous section.

6.1 The partition function on $S^2 \times S^1$

Let us start by discussing the matching of the partition function on $S^2 \times S^1$, also known as the supersymmetric index. The indices for the $O(N_c)_+$ versions of the dualities discussed in this paper were checked to match in [11, 33]. The index is sensitive to the global structure of the gauge group, and thus the matching of the indices for $SO(N_c)$ does not directly follow from these computations.¹⁰ We will check here that the supersymmetric indices match also for the $SO(N_c)$ dualities. In the process of doing this, we will see that, since the index contains information about local operators, it can test the proposed mapping of the baryon operators to the baryon-monopole operators discussed in the preceding sections.

First, let us briefly review the definition of the $3d$ supersymmetric index. It is defined by the following trace over states on $S^2 \times \mathbb{R}$ (see [34–37]–[6, 38] for details):

$$\mathcal{I}(x; \{u_a\}) = \text{Tr} \left[(-1)^{2J_3} x^{\Delta+J_3} \prod_a u_a^{e_a} \right]. \quad (6.1)$$

Here Δ is the energy in units of the S^2 radius (related to the conformal dimension for superconformal field theories), J_3 is the Cartan generator of the Lorentz $SO(3)$ isometry of S^2 , and e_a are charges under U(1) global symmetries (which could be subgroups of non-Abelian global symmetries). The states that contribute to this index satisfy $\Delta - R - J_3 = 0$, where R is the R-charge (that is used in the compactification on S^2).

¹⁰We will discuss below a generalization of the $O(N_c)_+$ index, which contains the same information as the $SO(N_c)$ index.

This index can be computed by a partition function on $\mathbb{S}^2 \times \mathbb{S}^1$, and localization dictates that the index gets contributions only from BPS configurations. For example, for a $U(1)$ gauge multiplet, we can take the gauge field to have a holonomy $z \in U(1)$ around the \mathbb{S}^1 and magnetic flux $m \in \mathbb{Z}$ on the \mathbb{S}^2 , which then determines the configurations of the other fields in the gauge multiplet. The 1-loop determinant of a chiral multiplet of R-charge R coupled with unit charge to this gauge multiplet is:

$$\mathcal{I}_\chi^R(x; z; m) \equiv (x^{1-R} z^{-1})^{|m|/2} \prod_{j=0}^{\infty} \frac{1 - (-1)^m z^{-1} x^{|m|+2-R+2j}}{1 - (-1)^m z x^{|m|+R+2j}}. \quad (6.2)$$

For a general gauge theory with gauge group G of rank r_G , one introduces fugacities z_i ($i = 1, \dots, r_G$) parameterizing the maximal torus of G , with corresponding GNO magnetic fluxes m_i on \mathbb{S}^2 . One can similarly introduce fugacities u_a and fluxes n_a for background gauge multiplets coupled to global symmetries. The 1-loop determinant in such a configuration is given by taking the product of the contributions (6.2) of the chiral multiplets, along with a contribution from the vector multiplet:

$$\mathcal{I}_V(x; z_i; m_i) \equiv \prod_{\alpha \in R} x^{-|\alpha(m)|/2} (1 - (-1)^{\alpha(m)} z^\alpha x^{|\alpha(m)|}), \quad (6.3)$$

where the product is over the roots of the gauge group. One can also include Chern-Simons terms for background or dynamical gauge multiplets, whose contribution, for instance, for a level k term for a $U(1)$ gauge multiplet, is z^{km} . Finally, the partition function is given by integrating over the gauge parameters z_i and summing over the gauge fluxes m_i .

We will be interested in the $SO(N_c)$ gauge theory with N_f chiral multiplets Q_a of R-charge R in the vector representation of $SO(N_c)$, and with a Chern-Simons term at level k . We include fugacities and fluxes, μ_a and n_a , $a = 1, \dots, N_f$, for the $U(N_f)$ flavor symmetry, as well as fugacities $\zeta = \pm 1$ for the global symmetry \mathbb{Z}_2^M , and $\chi = \pm 1$ for the charge conjugation symmetry \mathbb{Z}_2^C .

Let us write down explicitly the relevant indices. The index with $\chi = +1$ is given by:

$$\begin{aligned} \mathcal{I}_{SO(N_c)}^A(x; \mu_a; n_a; \zeta, \chi = +1) &= \zeta^{k\zeta \sum_a n_a} \prod_a \mu_a^{k_F n_a} \sum_{\{m_1, \dots, m_{r_G}\}} \zeta^{\sum_i m_i} \frac{1}{|W_{\{m_\ell\}}|} \\ &\oint \prod_{\ell=1}^{r_G} \left(\frac{dz_\ell}{2\pi i z_\ell} z_\ell^{k m_\ell} \right) \prod_{a=1}^{N_f} \left((\mathcal{I}_\chi^R(x; \mu_a; n_a))^\epsilon \prod_{i=1}^{r_G} \mathcal{I}_\chi^R(x; z_i \mu_a; m_i + n_a) \mathcal{I}_\chi^R(x; z_i^{-1} \mu_a; -m_i + n_a) \right) \\ &\prod_{i < j}^{r_G} \left(x^{-|m_i - m_j|} (1 - (-1)^{m_i - m_j} z_i z_j^{-1} x^{|m_i - m_j|}) (1 - (-1)^{m_i - m_j} z_i^{-1} z_j x^{|m_i - m_j|}) \right. \\ &\quad \left. x^{-|m_i + m_j|} (1 - (-1)^{m_i + m_j} z_i z_j x^{|m_i + m_j|}) (1 - (-1)^{m_i + m_j} z_i^{-1} z_j^{-1} x^{|m_i + m_j|}) \right) \\ &\left(\prod_{i=1}^{r_G} x^{-|m_i|} (1 - (-1)^{m_i} z_i x^{|m_i|}) (1 - (-1)^{m_i} z_i^{-1} x^{|m_i|}) \right)^\epsilon. \end{aligned} \quad (6.4)$$

Here $N_c = 2r_G + \epsilon$ with $\epsilon = 0, 1$. The integers m_i run over the Weyl-inequivalent GNO charges, and $|W_{\{m\}}|$ is the order of the residual Weyl group [35]. In the first term on the

first line we have introduced background Chern-Simons terms for the global symmetries. k_F is the level of a background Chern-Simons term for the $U(N_f)$ global symmetry [39, 40],¹¹ and k_ζ (obeying $k_\zeta \sim k_\zeta + 2$) is a similar term mixing the discrete $\mathbb{Z}_2^{\mathcal{M}}$ symmetry with $U(1)_A$.¹² We are free to choose the values of these terms, as long as parity anomalies are canceled so that the index is well defined, namely, it has an expansion in fugacities with integer powers.¹³ This requires

$$k_\zeta \in \{0, 1\}, \quad k_F \in \mathbb{Z} + \frac{1}{2} N_c, \quad (6.5)$$

where the second requirement is the standard parity anomaly [41, 42].

Next we want to compute the index with $\chi = -1$, where we should sum over holonomies of $O(N_c)$ that have determinant (-1) . The computation is different for the cases of odd and even N_c (see [11, 33] and also [43]). First, let us discuss the odd N_c case. A general $O(2r_G + 1)$ holonomy of determinant χ can be brought to the form

$$\text{diag}(z_1, z_1^{-1}, \dots, z_{r_G}, z_{r_G}^{-1}, \chi). \quad (6.6)$$

Thus, the indices with $\chi = -1$ are given by¹⁴

$$\begin{aligned} \mathcal{I}_{\text{SO}(N_c)}^A(x; \mu_a; n_a; \zeta, \chi = -1) &= \left(\zeta^{k_\zeta} \chi^{k_\chi} \right)^{\sum_a n_a} \prod_a \mu_a^{k_F n_a} \sum_{\{m_1, \dots, m_{r_G}\}} \zeta^{\sum_i m_i} \frac{1}{|W_{\{m_\ell\}}|} \\ &\oint \prod_{\ell=1}^{r_G} \left(\frac{dz_\ell}{2\pi i z_\ell} z_\ell^{k m_\ell} \right) \prod_{a=1}^{N_f} \left(\mathcal{I}_\chi^R(x; \chi \mu_a; n_a) \prod_{i=1}^{r_G} \mathcal{I}_\chi^R(x; z_i \mu_a; m_i + n_a) \mathcal{I}_\chi^R(x; z_i^{-1} \mu_a; -m_i + n_a) \right) \\ &\prod_{i < j}^{r_G} \left(x^{-|m_i - m_j|} (1 - (-1)^{m_i - m_j} z_i z_j^{-1} x^{|m_i - m_j|}) (1 - (-1)^{m_i - m_j} z_i^{-1} z_j x^{|m_i - m_j|}) \right. \\ &\quad \left. x^{-|m_i + m_j|} (1 - (-1)^{m_i + m_j} z_i z_j x^{|m_i + m_j|}) (1 - (-1)^{m_i + m_j} z_i^{-1} z_j^{-1} x^{|m_i + m_j|}) \right) \\ &\left(\prod_{i=1}^{r_G} x^{-|m_i|} (1 + (-1)^{m_i} z_i x^{|m_i|}) (1 + (-1)^{m_i} z_i^{-1} x^{|m_i|}) \right). \end{aligned} \quad (6.7)$$

We introduced here a background Chern-Simons term with coefficient k_χ ($k_\chi \sim k_\chi + 2$) mixing the charge conjugation symmetry $\mathbb{Z}_2^{\mathcal{C}}$ with $U(1)_A$, and for the partition function to be well-defined we must have

$$k_\chi \in \left\{ -\frac{1}{2}, \frac{1}{2} \right\}. \quad (6.8)$$

¹¹We could also introduce different levels for $U(1)_A$ and $SU(N_f)$.

¹²If we describe the background $\mathbb{Z}_2^{\mathcal{M}}$ gauge symmetry by two $U(1)$ gauge fields $A_{1,2}$ with an action given by an off-diagonal Chern-Simons term at level two [22, 23], we can write k_ζ as the coefficient of an ordinary Chern-Simons term that mixes A_1 with the background $U(1)_A$ gauge field.

¹³Note that, in the building blocks defining the index, there appear half-integer powers of the fugacities (see, e.g. (6.2)), and so these factors are not well-defined individually. However, when we expand the index as a series and include the appropriate background Chern-Simons terms, this expansion has only integer powers of the fugacities, and thus is well-defined.

¹⁴In the expression of the index here and in the even N_c case below, we write the last eigenvalue of the holonomy as χ in the contribution of the chiral multiplets, in order to keep track of the fractional powers of χ appearing in the intermediate expressions in a consistent way.

For even $N_c = 2r_G$, any holonomy of determinant $\chi = -1$ can be brought to the form

$$\text{diag} (z_1, z_1^{-1}, \dots, z_{r_G-1}, z_{r_G-1}^{-1}, 1, -1) . \quad (6.9)$$

Thus, the index is given by

$$\begin{aligned} \mathcal{I}_{\text{SO}(N_c)}^A(x; \mu_a; n_a; \zeta, \chi = -1) &= \left(\zeta^{k\zeta} \chi^{k\chi} \right)^{\sum_a n_a} \prod_a \mu_a^{k_F n_a} \sum_{\{m_1, \dots, m_{r_G-1}\}} \zeta^{\sum_i m_i} \frac{1}{|W_{\{m_\ell\}}|} \\ &\oint \prod_{\ell=1}^{r_G-1} \left(\frac{dz_\ell}{2\pi i} z_\ell^{km_\ell} \right) \prod_{a=1}^{N_f} \left(\prod_{i=1}^{r_G-1} \left(\mathcal{I}_\chi^R(x; z_i \mu_a; m_i + n_a) \mathcal{I}_\chi^R(x; z_i^{-1} \mu_a; -m_i + n_a) \right) \right. \\ &\left. \mathcal{I}_\chi^R(x; \mu_a; n_a) \mathcal{I}_\chi^R(x; \chi \mu_a; n_a) \right) \prod_{j=1}^{r_G-1} x^{-2|m_j|} (1 - z_j^{-2} x^{2|m_j|}) (1 - z_j^2 x^{2|m_j|}) \\ &\prod_{i < j < r_G}^{r_G-1} \left(x^{-|m_i - m_j|} (1 - (-1)^{m_i - m_j} z_i z_j^{-1} x^{|m_i - m_j|}) (1 - (-1)^{m_i - m_j} z_i^{-1} z_j x^{|m_i - m_j|}) \right. \\ &\left. x^{-|m_i + m_j|} (1 - (-1)^{m_i + m_j} z_i z_j x^{|m_i + m_j|}) (1 - (-1)^{m_i + m_j} z_i^{-1} z_j^{-1} x^{|m_i + m_j|}) \right), \end{aligned} \quad (6.10)$$

with the same quantization conditions on the background Chern-Simons terms as above.

The dual theory has an $\text{SO}(\tilde{N}_c = N_f + |k| - N_c + 2)$ gauge group and Chern-Simons level $(-k)$, with N_f chiral multiplets q_a in the vector representation, $N_f(N_f + 1)/2$ singlet mesons M_{ab} , and, for $k = 0$, also a singlet Y . The index of the dual theory with $\chi = 1$, with the charges and the background terms mapped appropriately across the duality, is:

$$\begin{aligned} \mathcal{I}_{\text{SO}(\tilde{N}_c)}^B(x; \mu_a; n_a; \zeta, \chi = +1) &= \zeta^{(k\zeta - k\chi) \sum_a n_a} \prod_a \mu_a^{k_F n_a} \\ &\zeta^{-\frac{\text{sign}(k)}{2} \sum_a n_a} \left(\prod_a (\mu_a x^R)^{\frac{1}{2}(kn_a - \text{sign}(k) \sum_a n_a)} \right) x^{\frac{\text{sign}(k)}{2} (N_f + 1 - N_c) \sum_a n_a} \\ &\left(\prod_{a \leq b}^{N_f} \mathcal{I}_\chi^{2R}(z; \mu_a \mu_b, n_a + n_b) \right) \left[\mathcal{I}_\chi^{N_f + 2 - N_c - N_f R}(z; \zeta^{-1} \prod_a \mu_a^{-1}, -\sum_a n_a) \right]^{\delta_{k,0}} \\ &\sum_{\{m_1, \dots, m_{\tilde{r}_G}\}} \zeta^{\sum_i m_i} \frac{1}{|W_{\{m_\ell\}}|} \oint \prod_{\ell=1}^{\tilde{r}_G} \left(\frac{dz_\ell}{2\pi i} z_\ell^{-km_\ell} \right) \\ &\prod_{a=1}^{N_f} \left((\mathcal{I}_\chi^{1-R}(x; \mu_a^{-1}, -n_a))^{\tilde{\epsilon}} \prod_{i=1}^{\tilde{r}_G} \mathcal{I}_\chi^{1-R}(x; z_i \mu_a^{-1}; m_i - n_a) \mathcal{I}_\chi^{1-R}(x; z_i^{-1} \mu_a^{-1}; -m_i - n_a) \right) \\ &\prod_{i < j}^{\tilde{r}_G} \left(x^{-|m_i - m_j|} (1 - (-1)^{m_i - m_j} z_i z_j^{-1} x^{|m_i - m_j|}) (1 - (-1)^{m_i - m_j} z_i^{-1} z_j x^{|m_i - m_j|}) \right. \\ &\left. x^{-|m_i + m_j|} (1 - (-1)^{m_i + m_j} z_i z_j x^{|m_i + m_j|}) (1 - (-1)^{m_i + m_j} z_i^{-1} z_j^{-1} x^{|m_i + m_j|}) \right) \\ &\left(\prod_{i=1}^{\tilde{r}_G} x^{-|m_i|} (1 - (-1)^{m_i} z_i x^{|m_i|}) (1 - (-1)^{m_i} z_i^{-1} x^{|m_i|}) \right)^{\tilde{\epsilon}}, \end{aligned} \quad (6.11)$$

where again $\tilde{N}_c = N_f + |k| - N_c + 2 = 2\tilde{r}_G + \tilde{\epsilon}$. Note that the parameter ζ now also appears in the contribution of the elementary field Y , since the $\mathbb{Z}_2^{\tilde{M}}$ symmetry in theory B also acts on this singlet. The factors on the first and the second line represent the contribution of background Chern-Simons terms. The background Chern-Simons terms on the second line are the relative ones, which must be included when $k \neq 0$ [10] (here we defined $\text{sign}(k) = 0$ for $k = 0$) for the duality to work. The expressions for $\mathcal{I}_{\text{SO}(\tilde{N}_c)}^B$ with $\chi = -1$ are obtained in an analogous way to our discussion of theory A above.

The dualities discussed in this paper imply the following equality for the indices,

$$\mathcal{I}_{\text{SO}(N_c)}^A(x; \mu_a; n_a; \zeta, \chi) = \mathcal{I}_{\text{SO}(N_f + |k| - N_c + 2)}^B(x; \mu_a; n_a; \zeta, \zeta \chi). \quad (6.12)$$

We have checked this equality for various values of the discrete parameters k , n_a , ζ and χ , by expanding both sides in a power series in x and comparing the leading coefficients.

We also can write the indices for other orthogonal gauge groups. In the $\text{SO}(N_c)$ index computation we introduced a fugacity $\chi = \pm 1$ for the global charge conjugation symmetry \mathbb{Z}_2^C . Similarly, we can introduce in the computation for an $O(N_c)$ gauge group a discrete theta-like parameter $\chi' = \pm 1$, determining whether we project on even or odd states under \mathbb{Z}_2^C . The $O(N_c)_+$ result for $\chi' = 1$ is half of the sum of the $\text{SO}(N_c)$ results with $\chi = 1, -1$, and the $O(N_c)_+$ result for $\chi' = -1$ is half of their difference. Thus, allowing for arbitrary χ and χ' one can relate the $\text{SO}(N_c)$ and the $O(N_c)_+$ expressions.¹⁵ In the $O(N_c)_-$ case we need to change the sign of the projection for states charged under \mathbb{Z}_2^M . Similarly, the $\text{Spin}(N_c)$ and $\text{Pin}(N_c)$ indices are given by summing over the sectors with different ζ , and we can define for them an index with $\zeta' = 1$ that projects on the \mathbb{Z}_2^M -even states (which make up the standard $\text{Spin}(N_c)$ and $\text{Pin}(N_c)$ theories), and an index with $\zeta' = -1$ that projects on the odd states. We then have:

$$\begin{aligned} \mathcal{I}_{\text{Spin}(N_c)}(x; \mu_a; n_a; \zeta', \chi) &= \\ & \frac{1}{2} (\mathcal{I}_{\text{SO}(N_c)}(x; \mu_a; n_a; \zeta = +1, \chi) + \zeta' \mathcal{I}_{\text{SO}(N_c)}(x; \mu_a; n_a; \zeta = -1, \chi)) , \\ \mathcal{I}_{O(N_c)_+}(x; \mu_a; n_a; \zeta, \chi') &= \\ & \frac{1}{2} (\mathcal{I}_{\text{SO}(N_c)}(x; \mu_a; n_a; \zeta, \chi = +1) + \chi' \mathcal{I}_{\text{SO}(N_c)}(x; \mu_a; n_a; \zeta, \chi = -1)) , \\ \mathcal{I}_{O(N_c)_-}(x; \mu_a; n_a; \zeta, \chi') &= \\ & \frac{1}{2} (\mathcal{I}_{\text{SO}(N_c)}(x; \mu_a; n_a; \zeta, \chi = +1) + \chi' \mathcal{I}_{\text{SO}(N_c)}(x; \mu_a; n_a; -\zeta, \chi = -1)) , \\ \mathcal{I}_{\text{Pin}(N_c)}(x; \mu_a; n_a; \zeta', \chi') &= \\ & \frac{1}{2} (\mathcal{I}_{\text{Spin}(N_c)}(x; \mu_a; n_a; \zeta', \chi = +1) + \chi' \mathcal{I}_{\text{Spin}(N_c)}(x; \mu_a; n_a; \zeta', \chi = -1)) . \end{aligned} \quad (6.13)$$

Our tests of the $\text{SO}(N_c)$ duality (6.12) with general ζ and χ provide tests also for the dualities between two $O(N_c)_+$ theories, between $\text{Spin}(N_c)$ and $O(\tilde{N}_c)_-$ theories, and between two $\text{Pin}(N_c)$ theories.¹⁶

¹⁵This is a discrete version of the situation for $U(N_c)$ gauge groups discussed in [6, 44, 45], where one can go between $SU(N_c)$ and $U(N_c)$ by gauging a $U(1)$ or by “ungauging” a topological $U(1)_J$ symmetry.

¹⁶Note that the standard indices of these theories map to each other, but the twisted indices which project onto odd states map only up to a sign, because of the non-trivial mapping of the \mathbb{Z}_2 global symmetries that is embodied in (6.12).

When the indices are expanded as a series in the various fugacities, the terms in the series represent the contributions of BPS operators with the corresponding charges, as follows from the definition of the index (6.1). Thus we can use the expressions above to attempt to trace how the baryons map between the two dual SO descriptions (6.12).¹⁷ On the electric side the usual baryons are Q^{N_c} , with an anti-symmetric product of some choice of N_c flavors. For example, let us define B to be the operator constructed out of the flavors $a = 1, \dots, N_c$. In the dual side, this maps to a monopole-baryon operator, $\tilde{\beta}'$, as described in section 5.3. Specifically, in this case:

$$B = \prod_{a=1}^{N_c} Q_a \quad \rightarrow \quad \tilde{\beta}' = e^{\tilde{\sigma}_1/\hat{g}_3^2 + i\tilde{a}_1} \left(\prod_{\alpha=3}^{k+2} \tilde{\lambda}_{1\alpha} \right) \left(\prod_{a=N_c+1}^{N_f} q_a \right). \quad (6.14)$$

Here $e^{\tilde{\sigma}_1/\hat{g}_3^2 + i\tilde{a}_1}$ is the basic $SO(\tilde{N}_c)$ monopole with GNO charges $(1, 0, 0, \dots, 0)$, $\tilde{\lambda}_{\alpha\beta}$ are the gluinos, and q_a are the dual quarks, with their color indices contracted anti-symmetrically in the subgroup $SO(N_f - N_c) \subset SO(\tilde{N}_c)$ that is left unbroken by the monopole and the gluinos. To test this mapping in the index, note that each chiral multiplet $Q_{a,\alpha}$ ($\alpha = 1, \dots, N_c$, $a = 1, \dots, N_f$) contributes a factor¹⁸ $x^R w_\alpha \mu_a$, so the operator B contributes a term $x^{RN_c} \prod_{a=1}^{N_c} \mu_a$ to the index. On the dual side, $q_{a,\beta}$ contributes $x^{1-R} w_\beta \mu_a^{-1}$, each gluino contributes $-x w_\alpha/w_1$, and the monopole background contributes $(-1)^k w_1^k x^{N_c - N_f - k + N_f R} \prod_{a=1}^{N_f} \mu_a$. Putting this together we find that the contribution of $\tilde{\beta}'$ matches that of B (there is also a factor of χ in theory A, and a factor of $\zeta\chi$ in theory B). The baryons of theory B q^{N_c} are mapped in a similar way to $\beta' = e^{\sigma_1/\hat{g}_3^2 + ia_1} \left(\prod_{\alpha=3}^{k+2} \lambda_{1\alpha} \right) Q^{N_c - k - 2}$ in theory A.

6.2 The partition functions on \mathbb{S}^3

Let us now comment on the \mathbb{S}^3 partition functions. The partition functions on \mathbb{S}^3 for $\mathcal{N} = 1$ SQCD with $O(N_c)_+$ gauge group were computed and found to agree for the $O(N_c)_+$ dualities discussed in [10, 11, 24, 33]. In fact, the equality of the partition functions of the theories with $SO(N_c)$ (and $Spin(N_c)$) gauge groups discussed in the preceding sections follows directly from the equality of partition functions for theories with $O(N_c)_+$ gauge groups. The \mathbb{S}^3 partition functions are computed by a matrix integral over the Lie algebra, which is the same in all these cases, and thus the partition functions differ only by overall factors of 2 due to the different volumes of the gauge groups. Hence, the results of [10, 11, 24, 33] straightforwardly imply that the $SO(N_c)$ dual pairs discussed in this paper have the same \mathbb{S}^3 partition functions.

In certain cases, e.g. the dualities discussed in [6], the equality of the partition functions on \mathbb{S}^3 of the $3d$ theories follows in a simple way from the equality of the $4d$ partition

¹⁷This statement comes with an obvious caveat. The index counts operators with signs in such a way that long multiplets cancel out, so we cannot exclude the possibility that there may be multiple states contributing to a given term.

¹⁸Here we use the notation $w_{2j-1} = z_j$ and $w_{2j} = z_j^{-1}$, $j = 1, \dots, r_G$, and $w_{N_c} = 1$ for N_c odd, where z_j , $j = 1, \dots, r_G$ span the maximal torus of $SO(N_c)$ as in (6.4). Thus the w_α , $\alpha = 1, \dots, N_c$ run over the weights of the vector representation, with $\prod_{\alpha=1}^{N_c} w_\alpha = 1$.

functions on $\mathbb{S}^3 \times \mathbb{S}^1$ (the supersymmetric index) of the $4d$ theories from which these $3d$ theories descend. However, this is more subtle in the case of dualities with orthogonal groups, as we will now explain.

First, let us briefly outline how the $3d$ partition functions are obtained from the $4d$ indices: for more details see [6, 7, 46–48]. The partition function of a $4d$ theory on $\mathbb{S}^3 \times \mathbb{S}^1$ can be thought of explicitly as an \mathbb{S}^3 partition function of the dimensionally reduced theory with all the KK modes included.¹⁹ The (inverse) radius of the \mathbb{S}^1 appears in the \mathbb{S}^3 partition function as a real mass for the $U(1)$ symmetry associated with the rotation around the circle. Taking the small radius limit corresponds to taking this real mass to be large, and thus decoupling the massive KK modes. The fugacities for the $4d$ global symmetries become real mass parameters in $3d$. Some of the classical symmetries of the $4d$ gauge theories are anomalous, but the $3d$ theories obtained by dimensional reduction of the matter content of the $4d$ ones do have these symmetries at the full quantum level. The $4d$ index cannot be refined with fugacities for the anomalous symmetries, and thus the $3d$ partition functions obtained by this reduction procedure are not refined with the corresponding real mass parameters. This is an indication that the $3d$ theory obtained by the reduction has a superpotential breaking the symmetry that is anomalous in $4d$ [6].

The above discussion presumes that the dimensional reduction produces a well-defined and finite \mathbb{S}^3 partition function. This presumption is true for the cases discussed in [6], but it is not true for the $SO(N_c)$ theories discussed in this paper: the reduction of the $4d$ index²⁰ for $SO(N_c)$ SQCD produces a divergent $3d$ partition function. The divergence can be explained physically by the fact that not all of the Coulomb branch is lifted when putting the theory on the circle, as discussed in the previous sections. In particular, in the $4d$ theory on \mathbb{S}^1 , the operators Y or Y_{Spin} , parameterizing the Coulomb branch, have no continuous global symmetry charges and no R-charge, and the presence of such a field leads the $3d$ partition function to diverge.²¹ We have seen in the previous sections that due to the intricate moduli space on the circle, the $3d$ $SO(N_c)$ dualities are obtained by focusing on certain regions of the Coulomb branch. It is possible that this more intricate procedure can also be mimicked at the level of the index,²² and we leave this question to future investigations.

¹⁹For the free chiral field, the representation of the $4d$ index as a product of $3d$ partition functions of KK modes is the physical content of [49], as explained in appendix B of [6].

²⁰The equality of $4d$ indices of dualities with $so(N_c)$ Lie algebras were checked in [50]. See also [51] for a related discussion.

²¹One can see this problem directly in $3d$. The partition functions on \mathbb{S}^3 and $\mathbb{S}^2 \times \mathbb{S}^1$, of $SO(N_c)$ SQCD with general real mass parameters and R-charges, are finite and well defined. However, on the subspace corresponding to the $4d$ R-symmetry, and where we turn on only real masses for non-anomalous symmetries in $4d$, both partition functions diverge.

²²A possible way to obtain a finite $3d$ partition function is to take a “double scaling” limit, of a small radius of the circle together with taking some of the real masses to be large.

Acknowledgments

We would like to thank R. Dijkgraaf, D. Freed, K. Intriligator, G. Moore, J. Park, I. Shamir, Y. Tachikawa and E. Witten for useful discussions. OA is the Samuel Sebbas Professorial Chair of Pure and Applied Physics, and he is supported in part by a grant from the Rosa and Emilio Segre Research Award, by an Israel Science Foundation center for excellence grant, by the German-Israeli Foundation (GIF) for Scientific Research and Development, by the Minerva foundation with funding from the Federal German Ministry for Education and Research, and by the I-CORE program of the Planning and Budgeting Committee and the Israel Science Foundation (grant number 1937/12). OA gratefully acknowledges support from an IBM Einstein Fellowship at the Institute for Advanced Study. SSR gratefully acknowledges support from the Martin A. Chooljian and Helen Chooljian membership at the Institute for Advanced Study. The research of SSR was also partially supported by NSF grant number PHY-0969448. The work of NS was supported in part by DOE grant DE-SC0009988 and by the United States-Israel Binational Science Foundation (BSF) under grant number 2010/629. The research of BW was supported in part by DOE Grant DE-SC0009988.

References

- [1] N. Seiberg, *Electric-magnetic duality in supersymmetric nonAbelian gauge theories*, *Nucl. Phys. B* **435** (1995) 129 [[hep-th/9411149](#)] [[INSPIRE](#)].
- [2] K.A. Intriligator and N. Seiberg, *Duality, monopoles, dyons, confinement and oblique confinement in supersymmetric $SO(N_c)$ gauge theories*, *Nucl. Phys. B* **444** (1995) 125 [[hep-th/9503179](#)] [[INSPIRE](#)].
- [3] K.A. Intriligator and N. Seiberg, *Lectures on supersymmetric gauge theories and electric-magnetic duality*, *Nucl. Phys. Proc. Suppl.* **45BC** (1996) 1 [[hep-th/9509066](#)] [[INSPIRE](#)].
- [4] M.J. Strassler, *Duality, phases, spinors and monopoles in $SO(N)$ and $spin(N)$ gauge theories*, *JHEP* **09** (1998) 017 [[hep-th/9709081](#)] [[INSPIRE](#)].
- [5] O. Aharony, N. Seiberg and Y. Tachikawa, *Reading between the lines of four-dimensional gauge theories*, [arXiv:1305.0318](#) [[INSPIRE](#)].
- [6] O. Aharony, S.S. Razamat, N. Seiberg and B. Willett, *3d dualities from 4d dualities*, *JHEP* **07** (2013) 149 [[arXiv:1305.3924](#)] [[INSPIRE](#)].
- [7] V. Niarchos, *Seiberg dualities and the 3d/4d connection*, *JHEP* **07** (2012) 075 [[arXiv:1205.2086](#)] [[INSPIRE](#)].
- [8] K.A. Intriligator and N. Seiberg, *Phases of $N = 1$ supersymmetric gauge theories and electric-magnetic triality*, [hep-th/9506084](#) [[INSPIRE](#)].
- [9] D. Gaiotto, G.W. Moore and A. Neitzke, *Framed BPS States*, [arXiv:1006.0146](#) [[INSPIRE](#)].
- [10] F. Benini, C. Closset and S. Cremonesi, *Comments on 3d Seiberg-like dualities*, *JHEP* **10** (2011) 075 [[arXiv:1108.5373](#)] [[INSPIRE](#)].
- [11] C. Hwang, K.-J. Park and J. Park, *Evidence for Aharony duality for orthogonal gauge groups*, *JHEP* **11** (2011) 011 [[arXiv:1109.2828](#)] [[INSPIRE](#)].

- [12] O. Aharony and I. Shamir, *On $O(N_c)$ $D = 3$ $N = 2$ supersymmetric QCD Theories*, *JHEP* **12** (2011) 043 [[arXiv:1109.5081](#)] [[INSPIRE](#)].
- [13] O. Aharony, A. Hanany, K.A. Intriligator, N. Seiberg and M. Strassler, *Aspects of $N = 2$ supersymmetric gauge theories in three-dimensions*, *Nucl. Phys. B* **499** (1997) 67 [[hep-th/9703110](#)] [[INSPIRE](#)].
- [14] K. Intriligator and N. Seiberg, *Aspects of 3d $N = 2$ Chern-Simons-Matter Theories*, [arXiv:1305.1633](#) [[INSPIRE](#)].
- [15] I. Affleck, J.A. Harvey and E. Witten, *Instantons and (Super)Symmetry Breaking in $(2+1)$ -Dimensions*, *Nucl. Phys. B* **206** (1982) 413 [[INSPIRE](#)].
- [16] N.M. Davies, T.J. Hollowood and V.V. Khoze, *Monopoles, affine algebras and the gluino condensate*, *J. Math. Phys.* **44** (2003) 3640 [[hep-th/0006011](#)] [[INSPIRE](#)].
- [17] N. Seiberg and E. Witten, *Gauge dynamics and compactification to three-dimensions*, [hep-th/9607163](#) [[INSPIRE](#)].
- [18] N. Seiberg, *Exact results on the space of vacua of four-dimensional SUSY gauge theories*, *Phys. Rev. D* **49** (1994) 6857 [[hep-th/9402044](#)] [[INSPIRE](#)].
- [19] K.-M. Lee and P. Yi, *Monopoles and instantons on partially compactified D-branes*, *Phys. Rev. D* **56** (1997) 3711 [[hep-th/9702107](#)] [[INSPIRE](#)].
- [20] K.-M. Lee, *Instantons and magnetic monopoles on $R^3 \times S^1$ with arbitrary simple gauge groups*, *Phys. Lett. B* **426** (1998) 323 [[hep-th/9802012](#)] [[INSPIRE](#)].
- [21] N.M. Davies, T.J. Hollowood, V.V. Khoze and M.P. Mattis, *Gluino condensate and magnetic monopoles in supersymmetric gluodynamics*, *Nucl. Phys. B* **559** (1999) 123 [[hep-th/9905015](#)] [[INSPIRE](#)].
- [22] J.M. Maldacena, G.W. Moore and N. Seiberg, *D-brane charges in five-brane backgrounds*, *JHEP* **10** (2001) 005 [[hep-th/0108152](#)] [[INSPIRE](#)].
- [23] T. Banks and N. Seiberg, *Symmetries and Strings in Field Theory and Gravity*, *Phys. Rev. D* **83** (2011) 084019 [[arXiv:1011.5120](#)] [[INSPIRE](#)].
- [24] A. Kapustin, *Seiberg-like duality in three dimensions for orthogonal gauge groups*, [arXiv:1104.0466](#) [[INSPIRE](#)].
- [25] S.G. Naculich, H. Riggs and H. Schnitzer, *Group level duality in wzw models and Chern-Simons theory*, *Phys. Lett. B* **246** (1990) 417 [[INSPIRE](#)].
- [26] E. Mlawer, S.G. Naculich, H. Riggs and H. Schnitzer, *Group level duality of WZW fusion coefficients and Chern-Simons link observables*, *Nucl. Phys. B* **352** (1991) 863 [[INSPIRE](#)].
- [27] T. Nakanishi and A. Tsuchiya, *Level rank duality of WZW models in conformal field theory*, *Commun. Math. Phys.* **144** (1992) 351 [[INSPIRE](#)].
- [28] O. Aharony, *IR duality in $D = 3$ $N = 2$ supersymmetric $USp(2N_c)$ and $U(N_c)$ gauge theories*, *Phys. Lett. B* **404** (1997) 71 [[hep-th/9703215](#)] [[INSPIRE](#)].
- [29] K.A. Intriligator and N. Seiberg, *Mirror symmetry in three-dimensional gauge theories*, *Phys. Lett. B* **387** (1996) 513 [[hep-th/9607207](#)] [[INSPIRE](#)].
- [30] J. de Boer, K. Hori, H. Ooguri and Y. Oz, *Mirror symmetry in three-dimensional gauge theories, quivers and D-branes*, *Nucl. Phys. B* **493** (1997) 101 [[hep-th/9611063](#)] [[INSPIRE](#)].

- [31] J. de Boer, K. Hori, Y. Oz and Z. Yin, *Branes and mirror symmetry in $N = 2$ supersymmetric gauge theories in three-dimensions*, *Nucl. Phys. B* **502** (1997) 107 [[hep-th/9702154](#)] [[INSPIRE](#)].
- [32] D. Green, Z. Komargodski, N. Seiberg, Y. Tachikawa and B. Wecht, *Exactly Marginal Deformations and Global Symmetries*, *JHEP* **06** (2010) 106 [[arXiv:1005.3546](#)] [[INSPIRE](#)].
- [33] C. Hwang, H. Kim, K.-J. Park and J. Park, *Index computation for 3d Chern-Simons matter theory: test of Seiberg-like duality*, *JHEP* **09** (2011) 037 [[arXiv:1107.4942](#)] [[INSPIRE](#)].
- [34] J. Bhattacharya, S. Bhattacharyya, S. Minwalla and S. Raju, *Indices for Superconformal Field Theories in 3,5 and 6 Dimensions*, *JHEP* **02** (2008) 064 [[arXiv:0801.1435](#)] [[INSPIRE](#)].
- [35] S. Kim, *The Complete superconformal index for $N = 6$ Chern-Simons theory*, *Nucl. Phys. B* **821** (2009) 241 [Erratum *ibid.* **B 864** (2012) 884] [[arXiv:0903.4172](#)] [[INSPIRE](#)].
- [36] Y. Imamura and S. Yokoyama, *Index for three dimensional superconformal field theories with general R-charge assignments*, *JHEP* **04** (2011) 007 [[arXiv:1101.0557](#)] [[INSPIRE](#)].
- [37] A. Kapustin and B. Willett, *Generalized Superconformal Index for Three Dimensional Field Theories*, [arXiv:1106.2484](#) [[INSPIRE](#)].
- [38] T. Dimofte, D. Gaiotto and S. Gukov, *3-Manifolds and 3d Indices*, [arXiv:1112.5179](#) [[INSPIRE](#)].
- [39] C. Closset, T.T. Dumitrescu, G. Festuccia, Z. Komargodski and N. Seiberg, *Contact Terms, Unitarity and F-Maximization in Three-Dimensional Superconformal Theories*, *JHEP* **10** (2012) 053 [[arXiv:1205.4142](#)] [[INSPIRE](#)].
- [40] C. Closset, T.T. Dumitrescu, G. Festuccia, Z. Komargodski and N. Seiberg, *Comments on Chern-Simons Contact Terms in Three Dimensions*, *JHEP* **09** (2012) 091 [[arXiv:1206.5218](#)] [[INSPIRE](#)].
- [41] A. Niemi and G. Semenoff, *Axial Anomaly Induced Fermion Fractionization and Effective Gauge Theory Actions in Odd Dimensional Space-Times*, *Phys. Rev. Lett.* **51** (1983) 2077 [[INSPIRE](#)].
- [42] A. Redlich, *Parity Violation and Gauge Noninvariance of the Effective Gauge Field Action in Three-Dimensions*, *Phys. Rev. D* **29** (1984) 2366 [[INSPIRE](#)].
- [43] B.I. Zwiebel, *Charging the Superconformal Index*, *JHEP* **01** (2012) 116 [[arXiv:1111.1773](#)] [[INSPIRE](#)].
- [44] A. Kapustin, *Three-dimensional Avatars of Seiberg Duality*, talk given at *Simons Summer Workshop in Mathematics and Physics 2011*, http://media.scgp.stonybrook.edu/video/video.php?f=20110810_1_qtp.mp4.
- [45] J. Park and K.-J. Park, *Seiberg-like Dualities for 3d $N = 2$ Theories with $SU(N)$ gauge group*, [arXiv:1305.6280](#) [[INSPIRE](#)].
- [46] F. Dolan, V. Spiridonov and G. Vartanov, *From 4d superconformal indices to 3d partition functions*, *Phys. Lett. B* **704** (2011) 234 [[arXiv:1104.1787](#)] [[INSPIRE](#)].
- [47] A. Gadde and W. Yan, *Reducing the 4d Index to the S^3 Partition Function*, *JHEP* **12** (2012) 003 [[arXiv:1104.2592](#)] [[INSPIRE](#)].
- [48] Y. Imamura, *Relation between the 4d superconformal index and the S^3 partition function*, *JHEP* **09** (2011) 133 [[arXiv:1104.4482](#)] [[INSPIRE](#)].

- [49] G. Felder and A. Varchenko, *The elliptic gamma function and $SL(3, \mathbb{Z}) \times Z^3$* , *Adv. Math.* **156** (2000) 44 [[math/9907061](#)].
- [50] F. Dolan and H. Osborn, *Applications of the Superconformal Index for Protected Operators and q -Hypergeometric Identities to $N = 1$ Dual Theories*, *Nucl. Phys. B* **818** (2009) 137 [[arXiv:0801.4947](#)] [[INSPIRE](#)].
- [51] V. Spiridonov and G. Vartanov, *Elliptic hypergeometry of supersymmetric dualities II. Orthogonal groups, knots and vortices*, [arXiv:1107.5788](#) [[INSPIRE](#)].