

Quantum finite-size effects for dyonic magnons in the $AdS_4 \times CP^3$

Changrim Ahn,^a Minkyoo Kim^b and Bum-Hoon Lee^b

^a*Department of Physics and Institute for the Early Universe, Ewha Womans University, Seoul 120-750, S. Korea*

^b*Department of Physics and Center for Quantum Spacetime, Sogang University, Seoul 121-742, S. Korea*

E-mail: ahn@ewha.ac.kr, mkim80@sogang.ac.kr, bhl@sogang.ac.kr

ABSTRACT: We compute quantum corrections to finite-size effects for various dyonic giant magnons in the $AdS_4 \times CP^3$ in two different approaches. The off-shell algebraic curve method is used to quantize the classical string configurations in semi-classical way and to compute the corrections to the string energies. These results are compared with the F -term Lüscher formula based on the S -matrix of the AdS_4/CFT_3 . The fact that the two results match exactly provides another stringent test for the all-loop integrability conjecture and the exact S -matrix based on it.

KEYWORDS: AdS-CFT Correspondence, Integrable Field Theories

ARXIV EPRINT: [1007.1598](https://arxiv.org/abs/1007.1598)

Contents

1	Introduction	1
2	Algebraic curve method	2
2.1	GMs in the algebraic curve	2
2.2	Off-shell frequency of Magnons	4
2.3	One-loop shifts of dyonic GMs	7
3	<i>S</i>-matrix and Lüscher formula	8
3.1	<i>S</i> -matrix	8
3.2	Small dyonic magnon (\mathbb{CP}^2)	11
3.3	Pair of small dyonic magnon (\mathbb{RP}^3)	12
3.4	Big magnon	12
4	Multi dyonic magnons	13
5	Conclusions	14

1 Introduction

The AdS/CFT correspondence [1–3] between string theories on AdS backgrounds and supersymmetric conformal gauge theories has been studied very actively and produced many exciting developments in understanding non-perturbative structures of the string and gauge theories. In the AdS_4/CFT_3 duality the three-dimensional conformal field theory is $\mathcal{N} = 6$ super Chern-Simons (CS) theory with $SU(N) \times SU(N)$ gauge symmetry and level k . This model, which was first proposed by Aharony, Bergman, Jafferis, and Maldacena (ABJM) [4], is believed to be dual to M-theory on $AdS_4 \times S^7/Z_k$. Furthermore, in the planar limit of $N, k \rightarrow \infty$ with a fixed value of 't Hooft coupling $\lambda = N/k$, the $\mathcal{N} = 6$ CS is believed to be dual to type IIA superstring theory on $AdS_4 \times \mathbb{CP}^3$.

In the strong coupling limit $\lambda \gg 1$, the string theory on $AdS_4 \times \mathbb{CP}^3$ is classically integrable [5–7] and probably maintains the integrability for any value of λ which would lead to the all-loop Bethe ansatz equations (BAEs) as conjectured in [8]. A discrepancy in one-loop correction to the energy of a folded spinning string from the all-loop Bethe ansatz results [9–11] can be resolved by a non-zero one-loop correction in the central interpolating function $h(\lambda)$ [12]. Based on the assumption that the ABJM model in the planar limit is quantum integrable along with centrally extended $su(2|2)$ symmetry, the exact *S*-matrix has been conjectured and used to confirm the above exact BAEs in [13]. However, this working hypothesis of all-loop integrability needs to be checked in all available methods. An efficient way is to compute and compare finite-size effects from both *S*-matrix and

string theory sides. A most successful way for S -matrix side is the Lüscher formula which computes small energy shift of on-shell particle states due to the finite-size of the system.

In the classical string theory side, various classical string configurations which correspond to the on-shell particle states have been identified. In addition to such classical string configurations as small giant magnon (GM) (\mathbb{CP}^1) [14] and small dyonic GM (\mathbb{CP}^2) [15, 16] which have also found in AdS_5/CFT_4 , new type of classical string solutions like pair of small GM (\mathbb{RP}^2) [14, 17, 18], pair of small dyonic GM (\mathbb{RP}^3) [19], and big GM [20–23] have been also discovered in $AdS_4 \times \mathbb{CP}^3$. The finite-size effects for these string configurations have been computed by either solving the superstring sigma model directly such as Neumann-Rosochatius reduction or the algebraic curve method [15, 19, 20, 24]. Leading finite-size corrections for the AdS_4/CFT_3 string configurations have been confirmed by the μ -term Lüscher formula for non-dyonic (\mathbb{CP}^1 and \mathbb{RP}^2) GM [25] and dyonic (\mathbb{RP}^3) GM [26].

The purpose of this paper is to compute quantum finite-size effects for dyonic magnons on $AdS_4 \times \mathbb{CP}^3$ and provide some more stringent tests of the exact S -matrix in the strong coupling limit. Quantum effects for the string theory on nontrivial background can be computed from quadratic fluctuations of fields in an effective Lagrangian (see, for example, [27]). However, this method has not been applied to the finite-size (dyonic) GM solutions on $AdS_4 \times \mathbb{CP}^3$ so far due to some technical difficulties. A more efficient way which utilizes the classical integrability of the string sigma-model is the algebraic curve method [28–30]. More specifically, we implement the off-shell quantization of the algebraic curve method developed in [28] for the various string configurations. These are compared with the F -term Lüscher formula in the S -matrix side. These two independent computations are shown to be identical, which provides a stringent test for both S -matrix and all-loop integrability conjecture. This paper is organized as follows. We will present in section 2 the quantum corrections for various classical string configurations using the off-shell algebraic curve formulation. The S -matrix and the F -term Lüscher formula based on it are used in section 3 to derive the finite-size effects. We will also generalize the one loop energy shifts to general multi-magnons in section 4. We conclude the paper with a few remarks in section 5.

2 Algebraic curve method

2.1 GMs in the algebraic curve

In the algebraic curve framework, different string solutions are mapped to different sets of eigenvalues of the classical transfer matrix. These are realized through a set of quasi-momenta $q_i(x)$ $i = 1, \dots, 10$ depending on a spectral parameter x , which are defined on a multi-sheet Riemann surface with particular analytic properties. In particular, solutions living mostly in \mathbb{CP}^3 have the following quasi-momenta as proposed in [7, 8, 24]:

$$\begin{aligned}
 q_1 &= -q_{10} = \frac{\alpha x}{x^2 - 1} \\
 q_2 &= -q_9 = \frac{\alpha x}{x^2 - 1} \\
 q_3 &= -q_8 = \frac{\alpha x}{x^2 - 1} + G_u(0) - G_u\left(\frac{1}{x}\right) + G_v(0) - G_v\left(\frac{1}{x}\right) + G_r(x) - G_r(0) + G_r\left(\frac{1}{x}\right)
 \end{aligned}$$

$$q_4 = -q_7 = \frac{\alpha x}{x^2 - 1} + G_u(x) + G_v(x) - G_r(x) + G_r(0) - G_r\left(\frac{1}{x}\right)$$

$$q_5 = -q_6 = G_u(x) - G_u(0) + G_u\left(\frac{1}{x}\right) - G_v(x) + G_v(0) - G_v\left(\frac{1}{x}\right).$$

Here α is related to the energy Δ by $\alpha = \frac{\Delta}{2g}$. This ansatz satisfies several analytic properties [7].

At large x , quasi-momenta are related with physical conserved quantities:

$$\lim_{x \rightarrow \infty} \begin{pmatrix} q_1(x) \\ q_2(x) \\ q_3(x) \\ q_4(x) \\ q_5(x) \end{pmatrix} \simeq \frac{1}{2gx} \begin{pmatrix} \Delta \\ \Delta \\ J_1 \\ J_2 \\ J_3 \end{pmatrix}. \tag{2.1}$$

J 's are angular momenta of strings moving in \mathbb{CP}^3 . Because we are interested in string solutions moving mostly in \mathbb{CP}^3 , we have set spins in AdS to zero. These quasi-momenta are related with each other by inversion symmetry:

$$\begin{pmatrix} q_1(1/x) \\ q_2(1/x) \\ q_3(1/x) \\ q_4(1/x) \\ q_5(1/x) \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ \pi m \\ \pi m \\ 0 \end{pmatrix} + \begin{pmatrix} -q_2(x) \\ -q_1(x) \\ -q_4(x) \\ -q_3(x) \\ +q_5(x) \end{pmatrix}. \tag{2.2}$$

Virasoro constraints in string σ -model manifest in the algebraic curve through the synchronization of poles at $x = \pm 1$ [7].

$$\lim_{x \rightarrow \pm 1} \begin{pmatrix} q_1(x) \\ q_2(x) \\ q_3(x) \\ q_4(x) \\ q_5(x) \end{pmatrix} \simeq \frac{1}{2(x \mp 1)} \begin{pmatrix} \alpha_{\pm} \\ \alpha_{\pm} \\ \alpha_{\pm} \\ \alpha_{\pm} \\ 0 \end{pmatrix}. \tag{2.3}$$

In contrast with the $AdS_5 \times S^5$ case, we need three resolvents G_u, G_v and G_r . Subscripts u, v, r correspond to different kinds of excitations, which are also related with Dynkin labels of $su(4)$. One can reproduce the charges of different string solutions by choosing different resolvents. The resolvents for the GM solutions, in particular, are log-cut distributions of Bethe roots and can be thought of as degeneration of two cuts [31–33]. At infinite J the GM resolvents are defined as [31]

$$G_{\text{magnon}} = -i \log \left(\frac{x - X^+}{x - X^-} \right),$$

while at finite size we have to think of the GM as a degenerate case of a two-cut solution with a resolvent [24, 32, 33]

$$G_{\text{finite}} = -2i \log \left(\frac{\sqrt{x - X^+} + \sqrt{x - Y^+}}{\sqrt{x - X^-} + \sqrt{x - Y^-}} \right).$$

There are three kinds of GM solutions in \mathbb{CP}^3 [14, 17, 20–23]. If $G_u = G_{\text{magnon}}$ and $G_v = G_r = 0$, then we obtain the small GM solution on \mathbb{CP}^2 . If we consider the v -type resolvent, we obtain small magnon with a reversed sign in the charge Q . If we set $G_u = G_v = G_{\text{magnon}}$ and $G_r = 0$, then we obtain a pair of small GMs, also called a \mathbb{RP}^3 solution. This solution corresponds to a GM excitation in each $su(2)$ sector of the gauge theory, and in particular we will consider only configurations with the same momentum in each sector. Finally, the big GM solution corresponds to $G_u = G_v = G_r = G_{\text{magnon}}$, which is two-parameter but one-charge solution because Q is zero [15].

Finite-size corrections of GMs in the algebraic curve can be computed from fluctuations of quasi-momenta. When we compute classical finite-size corrections, we use G_{finite} and consider log-cut distribution of Bethe roots with small square-root cut at the tips. As deviation from usual log-cut is very small, we could take the leading term of the fluctuation of quasi-momenta [32, 33]. On the other hand, if we are interested in quantum corrections, we have to add extra poles to quasi-momenta with G_{magnon} [29]. Such extra poles give fluctuations of energy by summing over on-shell frequencies. As we will explain in detail in section 2.2, on-shell frequencies are efficiently obtained from the off-shell frequencies. We first compute the fluctuation frequencies by using the off-shell method [28] in section 2.2 then evaluate energy shift in section 2.3.

2.2 Off-shell frequency of Magnons

One-loop computation using the algebraic curve method has been developed in [29, 30]. As explained above, fluctuations of superstring fields are mapped onto quasi-momenta fluctuations. These correspond to adding extra poles connecting sheets of the Riemann surface on which quasi-momenta are defined [29]. From the quasi-momenta perturbations, we can compute fluctuation frequencies.

There are two ways to compute fluctuation frequencies. In the on-shell method, we compute fluctuation frequencies by adding extra small poles to all physical polarization pairs. The position of these extra poles is determined by $q_i(x_n^{ij}) - q_j(x_n^{ij}) = 2\pi n_{ij}$. But this method involves repeated lengthy calculations for general solution, such as generic two-cut ones. More efficient way is the off-shell method where we need to consider only additional poles of quasi-momenta for particular polarization pair [28]. In this way, we don't fix the position of extra pole which we consider. The general ansatz for fluctuations of quasi-momenta can be written using analytic properties of algebraic curve. Then, we evaluate fluctuations of quasi-momenta at large x to determine the physical conserved charges. We can easily obtain off-shell fluctuation frequencies from these equations.

We will use this efficient off-shell method to compute one-loop effects for dyonic GMs. Fluctuation frequencies Ω_{ij} of all three types of GMs of $\text{AdS}_4/\text{CFT}_3$ are closely related with those of $\text{AdS}_5/\text{CFT}_4$ with one distinctive difference that there are two kinds of physical modes in $\text{AdS}_4/\text{CFT}_3$ — heavy and light modes. Heavy modes, however, do not contribute to one-loop leading term since it is suppressed exponentially.

Because functional form of off-shell frequencies of all three GMs are identical, we only show off-shell frequencies for the small magnon. A small magnon is obtained by $G_u = G_{\text{magnon}}$ and $G_v = G_r = 0$ [15, 24]. The quasi-momenta of small magnon are given

by

$$\begin{aligned}
 q_1 &= -q_{10} = \frac{\alpha x}{x^2 - 1} \\
 q_2 &= -q_9 = \frac{\alpha x}{x^2 - 1} \\
 q_3 &= -q_8 = \frac{\alpha x}{x^2 - 1} - i \log \left(\frac{X^+}{X^-} \right) + i \log \left(\frac{\frac{1}{x} - X^+}{\frac{1}{x} - X^-} \right) + \tau \\
 q_4 &= -q_7 = \frac{\alpha x}{x^2 - 1} - i \log \left(\frac{x - X^+}{x - X^-} \right) + \tau \\
 q_5 &= -q_6 = -i \log \left(\frac{x - X^+}{x - X^-} \right) + i \log \left(\frac{X^+}{X^-} \right) - i \log \left(\frac{\frac{1}{x} - X^+}{\frac{1}{x} - X^-} \right). \tag{2.4}
 \end{aligned}$$

Here, τ is a twist which is introduced to make single magnon satisfy the usual periodic boundary condition. In the case of small magnon, $\tau = -\frac{p}{2}$. (The twist for both a pair of small magnons and a big magnon is $-p$.)

On-shell fluctuation energy is defined as follows:

$$\Omega_n^{ij} = -\kappa_{ij} \delta_{i,1} + 2g \lim_{x \rightarrow \infty} x \delta_n^{ij} q_1(x).$$

Here, $\kappa_{ij} = 2$ for $(i, j) = (1, 10), (2, 9)$ and $\kappa_{ij} = 1$ for other pairs. Also, n_{ij} are mode numbers and moduli of the algebraic curve. They satisfy that $q_i^+ - q_j^- = 2\pi n_{ij}$. To obtain off-shell energy from on-shell energy, we just change n -dependence in the above expression to quasi-momentum dependence as follows [28]:

$$\Omega^{ij}(y) = \Omega_n^{ij} \Big|_{n \rightarrow \frac{q_i(y) - q_j(y)}{2\pi}}.$$

There are 16 polarization modes for type IIA superstring in $AdS_4 \times CP^3$. They consist of 8 light modes which are $(i, 5)$ or $(i, 6)$ pairs and 8 heavy modes for other pairs. We have to compute all types of fluctuation frequencies for these polarization modes. But, as explained first in [28] and studied in [34] for AdS_4/CFT_3 , off-shell frequencies are related to each other by the inversion symmetry. The light modes are related by

$$\begin{aligned}
 \Omega_{i5}(x) &= \Omega_{i6}(x) \\
 \Omega_{25}(x) &= \Omega_{15}(0) - \Omega_{15}\left(\frac{1}{x}\right) \\
 \Omega_{35}(x) &= \Omega_{45}(0) - \Omega_{45}\left(\frac{1}{x}\right),
 \end{aligned}$$

and heavy modes by

$$\begin{aligned}
 \Omega_{17}(x) &= \Omega_{15}(x) + \Omega_{57}(x) = \Omega_{15}(x) + \Omega_{45}(x) \\
 \Omega_{18}(x) &= \Omega_{15}(x) + \Omega_{58}(x) = \Omega_{15}(x) + \Omega_{35}(x) \\
 \Omega_{19}(x) &= \Omega_{15}(x) + \Omega_{59}(x) = \Omega_{15}(x) + \Omega_{25}(x) \\
 \Omega_{110}(x) &= \Omega_{15}(x) + \Omega_{15}(x) = 2\Omega_{15}(x)
 \end{aligned}$$

$$\begin{aligned}
\Omega_{27}(x) &= \Omega_{25}(x) + \Omega_{57}(x) = \Omega_{25}(x) + \Omega_{45}(x) \\
\Omega_{28}(x) &= \Omega_{25}(x) + \Omega_{58}(x) = \Omega_{25}(x) + \Omega_{35}(x) \\
\Omega_{29}(x) &= \Omega_{25}(x) + \Omega_{59}(x) = 2\Omega_{25}(x) \\
\Omega_{37}(x) &= \Omega_{35}(x) + \Omega_{57}(x) = \Omega_{35}(x) + \Omega_{45}(x).
\end{aligned}$$

All Ω_{ij} 's can be written in terms of only Ω_{15} and Ω_{45} .

Now we calculate the off-shell frequencies by considering fluctuations of quasi-momenta in the form of extra poles [28–30]. The quasi-momenta fluctuations can be determined by properties (2.2) and (2.3). For (1,5) polarization, we must add extra poles to q_1 and q_5 . If we write $\delta q_1, \delta q_4, \delta q_5$ to satisfy the analytic properties, then δq_2 and δq_3 can be automatically written by inversion symmetry. Note that δq_5 has self-inversion symmetry. The ansatz is then written as follows:

$$\begin{aligned}
\delta q_1 &= \frac{A_+}{x+1} + \frac{A_-}{x-1} + \frac{\alpha(y)}{x-y} \\
\delta q_2 &= -\frac{A_+}{\frac{1}{x}+1} - \frac{A_-}{\frac{1}{x}-1} - \frac{\alpha(y)}{\frac{1}{x}-y} \\
\delta q_3 &= -\frac{A_+}{\frac{1}{x}+1} - \frac{A_-}{\frac{1}{x}-1} - \frac{B^+}{\frac{1}{x}-X^+} - \frac{B^-}{\frac{1}{x}-X^-} \\
\delta q_4 &= \frac{A_+}{x+1} + \frac{A_-}{x-1} + \frac{B^+}{x-X^+} + \frac{B^-}{x-X^-} \\
\delta q_5 &= -\frac{\alpha(y)}{x-y} - \frac{\alpha(y)}{\frac{1}{x}-y} - \frac{\alpha(y)}{y} + \frac{B^+}{x-X^+} + \frac{B^-}{x-X^-} \\
&\quad + \frac{B^+}{\frac{1}{x}-X^+} + \frac{B^-}{\frac{1}{x}-X^-} + \frac{B^+}{X^+} + \frac{B^-}{X^-},
\end{aligned}$$

where $\alpha(x) = \frac{1}{2g} \frac{x^2}{x^2-1}$. The unknown constants A_{\pm} and B^{\pm} in the above ansatz are determined by the following equations coming from the asymptotic behaviors of the fluctuations in (2.1):

$$\begin{aligned}
A_+ - A_- &= \frac{\alpha(y)}{y} \\
\frac{B^+}{X^+} + \frac{B^-}{X^-} &= \frac{\alpha(y)}{y} \\
A_+ + A_- + \frac{\alpha(y)}{y^2} &= \frac{\delta\Delta}{2g} \\
A_+ + A_- + \frac{B^+}{X^{+2}} + \frac{B^-}{X^{-2}} &= 0 \\
A_+ + A_- + B^+ + B^- &= 0.
\end{aligned} \tag{2.5}$$

In the above equations, we used the N_{ij} in [20]. The fluctuation energy $\delta\Delta$ is given by the frequencies Ω_{ij}

$$\delta\Delta = \sum_{ij,n} N_{ij}^n \Omega_{ij}^n. \tag{2.6}$$

For (1, 5) polarization, all N_{ij} are zero except N_{15} . Also, we add just one pole ($N_{15} = 1$). Then, from (2.6), we have $\delta\Delta = \Omega_{15}(y)$. Now, we can solve $\delta\Delta$ from (2.5),

$$\delta\Delta = \frac{1}{y^2 - 1} \left(1 - y \frac{X^+ + X^-}{X^+ X^- + 1} \right).$$

With the similar argument, we can also compute $\Omega_{45}(y)$ which turns out to be the same as $\Omega_{15}(y)$. Using $\Omega_{15}(y)$ and $\Omega_{45}(y)$, we can obtain frequencies

$$\Omega_{ij}^{\text{light}}(y) = \frac{1}{y^2 - 1} \left(1 - y \frac{X^+ + X^-}{X^+ X^- + 1} \right), \tag{2.7}$$

for the light modes, and

$$\Omega_{ij}^{\text{heavy}}(y) = \frac{2}{y^2 - 1} \left(1 - y \frac{X^+ + X^-}{X^+ X^- + 1} \right), \tag{2.8}$$

for the heavy modes, which are twice of those of light modes.

These are exactly the same results as those of [20]. Similarly, off-shell frequencies for other cases (pair of small and big magnon) can be evaluated. In other cases, we also consider the most general ansatz for fluctuations of quasi-momenta and compare their asymptotic with the conserved charges. Then, we get a set of similar equations as (2.5). The result for other magnons is exactly same than that of the small magnon. We can use the off-shell frequencies of the small magnon (2.7) and (2.8) for all other magnons.

2.3 One-loop shifts of dyonic GMs

The leading part of one-loop energy shift is given by the sum of fluctuation frequencies.

$$\delta\Delta_{\text{one-loop}} = \frac{1}{2} \sum_{ij} \sum_n (-1)^{F_{ij}} \Omega_{ij}^n = \int \frac{dx}{2\pi i} \partial_x \Omega(x) \sum_{ij} \gamma_{ij} (-1)^{F_{ij}} e^{-i(q_i - q_j)}.$$

Here, $\gamma_{ij} = 1$ for light modes and $\gamma_{ij} = 2$ for heavy modes and $\Omega(x)$ in the last expression is off-shell energy. When we change from infinite summation over n to integration over x , there are some non-trivial steps which we have to be careful [20, 30, 49].

When we evaluate the above integral by using saddle-point approximation, heavy modes can be suppressed because of the factor 2 in exponent. This is related to the fact there are no $\frac{\alpha x}{x^2 - 1}$ term in q_5 or q_6 . Hence, only light modes are important in this computation. This fact is consistent with the observation in [35] that only 8 light degrees of freedom are physical. The other 8 heavy degrees of freedom of superstring theory are unstable.

So we need to compute $\sum_{ij} (-1)^{F_{ij}} e^{-i(q_i - q_j)}$ where the sum over (i, j) pairs includes only the light modes. Using quasi-momenta of the small magnon in (2.4), we obtain

$$\begin{aligned} \sum_{ij} (-1)^{F_{ij}} e^{-i(q_i - q_j)} &= e^{\frac{-i\alpha x}{x^2 - 1}} \left[2 \frac{\frac{1}{x} - X^+ x - X^+ X^+}{\frac{1}{x} - X^- x - X^- X^-} + 2 \frac{\frac{1}{x} - X^- x - X^- X^-}{\frac{1}{x} - X^+ x - X^+ X^+} \right. \\ &\quad - \frac{\frac{1}{x} - X^+}{\frac{1}{x} - X^-} \sqrt{\frac{X^-}{X^+}} - \frac{x - X^-}{x - X^+} \sqrt{\frac{X^+}{X^-}} - \left(\frac{\frac{1}{x} - X^+}{\frac{1}{x} - X^-} \right)^2 \frac{x - X^+}{x - X^-} \left(\frac{X^-}{X^+} \right)^{\frac{3}{2}} \\ &\quad \left. - \frac{\frac{1}{x} - X^-}{\frac{1}{x} - X^+} \left(\frac{x - X^+}{x - X^-} \right)^2 \left(\frac{X^+}{X^-} \right)^{\frac{3}{2}} \right]. \tag{2.9} \end{aligned}$$

The pair of small magnon is obtained by the following ansatz:

$$\begin{aligned}
 q_1 &= -q_{10} = \frac{\alpha x}{x^2 - 1} \\
 q_2 &= -q_9 = \frac{\alpha x}{x^2 - 1} \\
 q_3 &= -q_8 = \frac{\alpha x}{x^2 - 1} - 2i \log \left(\frac{X^+}{X^-} \right) + 2i \log \left(\frac{\frac{1}{x} - X^+}{\frac{1}{x} - X^-} \right) - p \\
 q_4 &= -q_7 = \frac{\alpha x}{x^2 - 1} - 2i \log \left(\frac{x - X^+}{x - X^-} \right) - p \\
 q_5 &= -q_6 = 0,
 \end{aligned}$$

which leads to

$$\sum_{ij} (-1)^{F_{ij}} e^{-i(q_i - q_j)} = 2e^{\frac{-i\alpha x}{x^2 - 1}} \left[2 - \left(\frac{\frac{1}{x} - X^+}{\frac{1}{x} - X^-} \right)^2 \frac{X^-}{X^+} - \left(\frac{x - X^-}{x - X^+} \right)^2 \frac{X^+}{X^-} \right]. \quad (2.10)$$

For the big magnon, using the quasi-momenta

$$\begin{aligned}
 q_1 &= -q_{10} = \frac{\alpha x}{x^2 - 1} \\
 q_2 &= -q_9 = \frac{\alpha x}{x^2 - 1} \\
 q_3 &= -q_8 = \frac{\alpha x}{x^2 - 1} - i \log \left(\frac{X^+}{X^-} \right) + i \log \left(\frac{\frac{1}{x} - X^+}{\frac{1}{x} - X^-} \right) - i \log \left(\frac{x - X^+}{x - X^-} \right) - p \\
 q_4 &= -q_7 = \frac{\alpha x}{x^2 - 1} - i \log \left(\frac{X^+}{X^-} \right) + i \log \left(\frac{\frac{1}{x} - X^+}{\frac{1}{x} - X^-} \right) - i \log \left(\frac{x - X^+}{x - X^-} \right) - p \\
 q_5 &= -q_6 = 0,
 \end{aligned} \quad (2.11)$$

we get

$$\sum_{ij} (-1)^{F_{ij}} e^{-i(q_i - q_j)} = 4e^{\frac{-i\alpha x}{x^2 - 1}} \left[1 - \frac{\frac{1}{x} - X^+}{\frac{1}{x} - X^-} \frac{x - X^-}{x - X^+} \right]. \quad (2.12)$$

The term $-p$ in (2.10) and (2.11) are the twists of \mathbb{RP}^3 magnon and big magnon.

3 S-matrix and Lüscher formula

We use the Ahn-Nepomechie S -matrix of the $\text{AdS}_4/\text{CFT}_3$ [13] in the F -term Lüscher formula to compute the quantum corrections for the various dyonic GM states.

3.1 S-matrix

The $\mathcal{N} = 6$ Chern-Simons theory contains two types of fundamental excitations, denoted by A and B , which belong to a fundamental representation of the centrally extended $su(2|2)$. The S -matrices among these states are given by

$$\begin{aligned}
 S^{AA}(p_1, p_2) &= S^{BB}(p_1, p_2) = S_0(p_1, p_2) \hat{S}(p_1, p_2) \\
 S^{AB}(p_1, p_2) &= S^{BA}(p_1, p_2) = \tilde{S}_0(p_1, p_2) \hat{S}(p_1, p_2),
 \end{aligned}$$

where \hat{S} is the $su(2|2)$ -invariant S -matrix [36, 37]. Main feature of the S -matrix is encoded into the two scalar factors,

$$S_0(p_1, p_2) = \frac{1 - \frac{1}{x_1^+ x_2^-}}{1 - \frac{1}{x_1^- x_2^+}} \sigma(p_1, p_2)$$

$$\tilde{S}_0(p_1, p_2) = \frac{x_1^- - x_2^+}{x_1^+ - x_2^-} \sigma(p_1, p_2),$$

where $\sigma(p_1, p_2)$ is the BES dressing factor [38]. These scalar factors have been checked classically in [39].

Here we compute the leading one-loop correction using the F -term Lüscher formula which is given by [40–42]

$$\delta E_F = - \int \frac{dq}{2\pi} \left[1 - \frac{\varepsilon'_Q(p)}{\varepsilon'_1(q^*)} \right] e^{-iq^*L} \sum_b (-1)^{F_b} \left(S_{ba}^{ba}(q^*, p) - 1 \right). \quad (3.1)$$

The integrand of the formula consists of the kinematic and the S -matrix factors.

For the kinematical factor we consider first the \mathbb{CP}^2 small magnon whose energy dispersion relation is given by

$$\Delta - J/2 = \varepsilon_Q(p) = \sqrt{\frac{Q^2}{4} + 16g^2 \sin^2 \frac{p}{2}}.$$

It is convenient to introduce X^\pm and y^\pm variables defined by

$$X^+ - X^- + \frac{1}{X^+} - \frac{1}{X^-} = \frac{iQ}{2g}$$

$$y^+ - y^- + \frac{1}{y^+} - \frac{1}{y^-} = \frac{i}{2g}$$

$$\frac{X^+}{X^-} = e^{ip}, \quad \frac{y^+}{y^-} = e^{iq^*}. \quad (3.2)$$

To compare the kinematic factor with algebraic curve results, we introduce a variable x defined by [25, 30].

$$x + \frac{1}{x} \pm \frac{i}{4g} = y^\pm + \frac{1}{y^\pm},$$

which leads to

$$y^\pm = x \pm \frac{ix^2}{4g(x^2 - 1)}$$

at strong coupling limit. The dispersion relation of virtual fundamental GM in this limit becomes

$$\varepsilon_1(q^*) = \frac{g}{i} \left(y^+ - y^- - \frac{1}{y^+} + \frac{1}{y^-} \right) = \frac{1}{2} \frac{x^2 + 1}{x^2 - 1}.$$

From (3.2), one gets

$$e^{iq^*} \simeq 1 + iq^* = 1 + i \frac{x}{g(x^2 - 1)}.$$

Also from $q^2 + \varepsilon'^2 (q^*) = 0$, q can be written as

$$q = \frac{i x^2 + 1}{2 x^2 - 1}.$$

From these, we can get

$$\begin{aligned} \varepsilon'_Q(p) &= g \left(\frac{X^+ + X^-}{X^+ X^- + 1} \right) \\ \varepsilon'_1(q^*) &= g \left(\frac{2x}{x^2 + 1} \right). \end{aligned}$$

Inserting these into the kinematic factor and changing the integration variable to x , one can get the same kinematic factor as that of the algebraic curve computation,

$$\delta E_{\text{one-loop}}^F = \int \frac{dx}{2\pi i} \partial_x \Omega(x) e^{-\frac{i x J}{2g(x^2-1)}} \sum_b (-1)^{F_b} S_{b1Q}^{b1Q}(q^*, p).$$

For other types of dyonic GMs, the results stand in the same way.

The S -matrix factor in the Lüscher formula contains scatterings between virtual and physical particles. In the leading order, only fundamental particles contribute to the sum over virtual particles $b = 1, 2, 3, 4$ of both A and B types. The physical particles are dyonic GMs in $su(2)$ sector which are bound states of Q number of bosonic particles with $su(2|2)$ index $a = 1_Q = (11 \dots 1)$. Since the physical particles carry $su(2|2)$ index 1, the relevant S -matrix elements are

$$\begin{aligned} \hat{S}_{11}^{11} &= a_1(p_1, p_2), \quad \hat{S}_{21}^{21} = a_1(p_1, p_2) + a_2(p_1, p_2), \quad \hat{S}_{31}^{31} = \hat{S}_{41}^{41} = a_6(p_1, p_2), \\ a_1(p_1, p_2) &= \frac{x_2^- - x_1^+ \eta_1 \eta_2}{x_2^+ - x_1^- \tilde{\eta}_1 \tilde{\eta}_2} \\ a_2(p_1, p_2) &= \frac{(x_1^- - x_1^+) (x_2^- - x_2^+) (x_2^- - x_1^+) \eta_1 \eta_2}{(x_1^- - x_2^+) (x_2^- x_1^- - x_2^+ x_1^+) \tilde{\eta}_1 \tilde{\eta}_2} \\ a_6(p_1, p_2) &= \frac{x_1^+ - x_2^+ \eta_2}{x_1^- - x_2^- \tilde{\eta}_2}. \end{aligned}$$

We choose the phase factors η_i in the string frame [36]

$$\frac{\eta_1}{\tilde{\eta}_1} = \sqrt{\frac{x_2^+}{x_2^-}}, \quad \frac{\eta_2}{\tilde{\eta}_2} = \sqrt{\frac{x_1^-}{x_1^+}}.$$

The relevant S -matrix elements for the dyonic magnons are $S_{b1Q}^{AA b1Q}$ and $S_{b1Q}^{BA b1Q}$ which are given by [26]

$$S_{b1Q}^{AA b1Q} = \prod_{k=1}^Q \left(\frac{1 - \frac{1}{y^+ x_k^-}}{1 - \frac{1}{y^- x_k^+}} \sigma_{BES}(y, x_k) \hat{S}_{b1}^{b1}(y, x_k) \right) \quad (3.3)$$

$$S_{b1Q}^{BA b1Q} = \prod_{k=1}^Q \left(\frac{y^- - x_k^+}{y^+ - x_k^-} \sigma_{BES}(y, x_k) \hat{S}_{b1}^{b1}(y, x_k) \right). \quad (3.4)$$

From these, we obtain

$$\begin{aligned}\sum_b (-1)^{F_b} S^{AA b_1 Q}_{b_1 Q} &= \sigma_{BES}(y, X) \frac{\eta(X)}{\tilde{\eta}(X)} \left(\frac{\eta(y)}{\tilde{\eta}(y)} \right)^Q (s_1 + s_2 - s_3 - s_4) S_{BDS}^Q \\ \sum_b (-1)^{F_b} S^{BA b_1 Q}_{b_1 Q} &= \sigma_{BES}(y, X) \frac{\eta(X)}{\tilde{\eta}(X)} \left(\frac{\eta(y)}{\tilde{\eta}(y)} \right)^Q (s_1 + s_2 - s_3 - s_4).\end{aligned}$$

where we defined $x_1^- = X^-$, $x_Q^+ = X^+$, and $x_k^- = x_{k-1}^+$. The s_b s defined by

$$s_b = \prod_{k=1}^Q \frac{\hat{S}_{b_1}^{b_1}(y, x_k)}{\hat{S}_{11}^{11}(y, x_k)}$$

can be written as

$$\begin{aligned}s_1 &= 1, & s_2 &= \left(\frac{y^+ - X^+}{y^+ - X^-} \right) \left(\frac{1 - \frac{1}{y^- X^+}}{1 - \frac{1}{y^- X^-}} \right) \\ s_3 &= s_4 = \left(\frac{y^+ - X^+}{y^+ - X^-} \right) \frac{\tilde{\eta}(X)}{\eta(X)}.\end{aligned}\tag{3.5}$$

The dressing factor and S_{BDS} of bound-state in this limit are given by [30, 44]

$$\begin{aligned}\sigma_{BES}(y, X) &= \left(\frac{y - \frac{1}{X^+}}{y - \frac{1}{X^-}} \right) e^{\frac{-ix}{2g(x^2-1)}(\Delta - J/2 - Q)} \\ S_{BDS}^Q &= \frac{(y^+ - X^-) \left(1 - \frac{1}{y^+ X^-}\right) (y^- - X^-) \left(1 - \frac{1}{y^- X^-}\right)}{(y^- - X^+) \left(1 - \frac{1}{y^- X^+}\right) (y^+ - X^+) \left(1 - \frac{1}{y^+ X^+}\right)}.\end{aligned}\tag{3.6}$$

3.2 Small dyonic magnon (\mathbb{CP}^2)

Dyonic small magnon is a bound-state of A -particles or B -particles. Because the S -matrices are invariant under $A \leftrightarrow B$, we can only consider a A -particle without loss of generality. Then we can write the S -matrix elements for small dyonic magnon as below:

$$\sum_b (-1)^{F_b} S_{b_1 Q}^{b_1 Q}(q^*, p) = \sum_b (-1)^{F_b} \left(S^{AA b_1 Q}_{b_1 Q} + S^{BA b_1 Q}_{b_1 Q} \right).\tag{3.7}$$

We can approximate $y^\pm = x$ in the strong coupling limit for the leading F -term integration and get

$$\frac{\eta(X)}{\tilde{\eta}(X)} = e^{\frac{ip}{2}}, \quad \left(\frac{\eta(y)}{\tilde{\eta}(y)} \right)^Q = e^{-i \frac{xQ}{2g(x^2-1)}}.\tag{3.8}$$

These lead to the one-loop correction as follows:

$$\begin{aligned}\delta\Delta_F &= \int \frac{dx}{2\pi i} \partial_x \Omega(x) e^{-i\Delta \frac{x}{2g(x^2-1)}} \left[\left(\frac{x - \frac{1}{X^+}}{x - \frac{1}{X^-}} \right) e^{\frac{ip}{2}} + \left(\frac{x - X^-}{x - X^+} \right) e^{\frac{ip}{2}} \right. \\ &\quad + \left(\frac{x - \frac{1}{X^-}}{x - \frac{1}{X^+}} \right) \left(\frac{x - X^-}{x - X^+} \right)^2 e^{\frac{ip}{2}} + \left(\frac{x - \frac{1}{X^+}}{x - \frac{1}{X^-}} \right)^2 \left(\frac{x - X^+}{x - X^-} \right) e^{\frac{ip}{2}} \\ &\quad \left. - 2 \left(\frac{x - \frac{1}{X^+}}{x - \frac{1}{X^-}} \right) \left(\frac{x - X^+}{x - X^-} \right) - 2 \left(\frac{x - \frac{1}{X^-}}{x - \frac{1}{X^+}} \right) \left(\frac{x - X^-}{x - X^+} \right) \right].\end{aligned}\tag{3.9}$$

It can be easily seen that the expression in bracket with the exponential factor in front is exactly the same as the result in (2.9) in algebraic curve for small dyonic magnon.

3.3 Pair of small dyonic magnon (\mathbb{RP}^3)

In case of \mathbb{RP}^3 dyonic magnon, we need to consider a pair of A and B type dyonic GMs in the physical state with the same momentum $p_1 = p_2 = p$ [26]. The F -term Lüscher formula becomes [25]

$$\begin{aligned} \sum_b (-1)^{F_b} S_{b1Q}^{b1Q}(q^*, p) &= \sum_b (-1)^{F_b} \left(S^{AA b1Q}_{b1Q} S^{AB b1Q}_{b1Q} + S^{BA b1Q}_{b1Q} S^{BB b1Q}_{b1Q} \right) \\ &= 2 \sum_b (-1)^{F_b} S^{AA b1Q}_{b1Q} S^{BA b1Q}_{b1Q}. \end{aligned} \quad (3.10)$$

Using the S -matrix elements in (3.3) and (3.4), we obtain

$$\delta\Delta_F = \int \frac{dx}{2\pi i} \partial_x \Omega(x) e^{-i\Delta \frac{x}{2g(x^2-1)}} \left[4 - 2 \left(\frac{x - \frac{1}{X^+}}{x - \frac{1}{X^-}} \right)^2 e^{ip} - 2 \left(\frac{x - X^-}{x - X^+} \right)^2 e^{ip} \right].$$

By comparing this with (2.10) of algebraic curve, one can confirm that the energy shift from both methods agree each other.

3.4 Big magnon

Particle interpretation of big magnon is not clear until now. We propose that big magnon may be superposition of a small magnon and an “anti-small magnon”. The anti-small magnon has the same momentum as the usual small magnon but the quantum number $-Q$. Then, we can compute S -matrix elements as below:

$$\sum_b (-1)^{F_b} S_{ba}^{ba}(q^*, p) = \sum_b (-1)^{F_b} \left(S^{AA b1Q}_{b1Q} S^{AA' b1-Q}_{b1-Q} + S^{BA b1Q}_{b1Q} S^{BA' b1-Q}_{b1-Q} \right).$$

From these S -matrix elements, we have

$$\begin{aligned} \sum_b (-1)^{F_b} \left(S_{ba}^{ba}(q^*, p) \right) &= 2 \prod_{k=1}^Q \left[\left(\frac{1 - \frac{1}{y^+ X_k^-}}{1 - \frac{1}{y^- X_k^+}} \right) \left(\frac{y^+ - X_k^-}{y^- - X_k^+} \right) \left(\frac{1 - \frac{X_k^+}{y^+}}{1 - \frac{X_k^-}{y^-}} \right) \left(\frac{y^+ - \frac{1}{X_k^+}}{y^- - \frac{1}{X_k^-}} \right) \right] \\ &\times \left(\frac{y - \frac{1}{X_Q^+}}{y - \frac{1}{X_Q^-}} \right) \left(\frac{y - X_Q^-}{y - X_Q^+} \right) \left(\frac{\eta(X^Q)}{\tilde{\eta}(X^Q)} \right)^2 e^{-i(\Delta-J) \frac{x}{2g(x^2-1)}} (1 + s_2 s'_2 - 2s_3 s'_3). \end{aligned}$$

The primed quantities, A', s'_2 and s'_3 , are related to those unprimed by $X^\pm \rightarrow \frac{1}{X^\mp}$ which is equivalent to changing $Q \rightarrow -Q$. From the S -matrix element, the one-loop correction of big magnon is given by

$$\delta\Delta_F = \int \frac{dx}{2\pi i} \partial_x \Omega(x) e^{-i\Delta \frac{x}{2g(x^2-1)}} 4 \left[1 - \left(\frac{x - \frac{1}{X^+}}{x - \frac{1}{X^-}} \right) \left(\frac{x - X^-}{x - X^+} \right) e^{ip} \right].$$

This result matches exactly with that of the algebraic curve (2.12). The one-loop corrections for pair of small magnon and big magnon are exactly the same in the non-dyonic limit. While the physical meaning of the anti-magnon state is not clear, the one-loop correction analysis suggests how to express the big magnon state in terms of the on-shell particles.

4 Multi dyonic magnons

As authors of [24, 32, 33] first explain, in algebraic curve technology, the case of (N, M) -type magnon is realized as a sum of N resolvents of u and M resolvents of v type. The meaning of the r -type resolvent is not clear in the S -matrix interpretation. So, we set the r -type resolvent to zero and consider only N -number of u -type excitations and M -number of v -type excitations. These correspond to general excitations of N -number of A -type dyonic magnons and M -number of B -type dyonic magnons in spin chain. \mathbb{CP}^2 and \mathbb{RP}^3 magnons are special cases of this general multi-magnon state. To compute one-loop finite size effects for such configurations of magnons, we use the following quasi-momenta ansatz:

$$\begin{aligned}
 q_1 &= -q_{10} = \frac{\alpha x}{x^2 - 1} \\
 q_2 &= -q_9 = \frac{\alpha x}{x^2 - 1} \\
 q_3 &= -q_8 = \frac{\alpha x}{x^2 - 1} + \sum_{k=1}^N \left(G_u^k(0) - G_u^k\left(\frac{1}{x}\right) \right) + \sum_{k=N+1}^{N+M} \left(G_v^k(0) - G_v^k\left(\frac{1}{x}\right) \right) + \sum_{i=1}^{N+M} \tau_i \\
 q_4 &= -q_7 = \frac{\alpha x}{x^2 - 1} + \sum_{k=1}^N G_u^k(x) + \sum_{k=N+1}^{N+M} G_v^k(x) + \sum_{i=1}^{N+M} \tau_i \\
 q_5 &= -q_6 = \sum_{k=1}^N \left(G_u^k(x) - G_u^k(0) + G_u^k\left(\frac{1}{x}\right) \right) + \sum_{k=N+1}^{N+M} \left(-G_v^k(x) + G_v^k(0) - G_v^k\left(\frac{1}{x}\right) \right).
 \end{aligned}$$

Here, $G_u^k(x)$ and $G_v^k(x)$ are all the same with $G_{\text{magnon}} = -i \log \left(\frac{x - X_k^+}{x - X_k^-} \right)$ and $\tau_i = -\frac{p_i}{2}$.

The problem here is that unknown functions in the fluctuations of quasi-momenta can not be completely determined by the constraint equations coming from asymptotic limit. So we get some undetermined functions α_l which satisfy $\sum_l \alpha_l = 1$ where the index l runs from 1 to $N + M$. This feature of the fluctuation frequencies for multi-particle states is noticed first in [45] for $\mathcal{N} = 4$ SYM. In $\text{AdS}_4/\text{CFT}_3$, off-shell frequencies are as below.

$$\begin{aligned}
 \Omega_{ij}^{\text{light}}(x) &= \frac{1}{x^2 - 1} \left(1 - \sum_l \alpha_l \left(x \frac{X_l^+ + X_l^-}{X_l^+ X_l^- + 1} \right) \right) = \Omega_{\text{multi}}(x) \\
 \Omega_{ij}^{\text{heavy}}(x) &= \frac{2}{x^2 - 1} \left(1 - \sum_l \alpha_l \left(x \frac{X_l^+ + X_l^-}{X_l^+ X_l^- + 1} \right) \right).
 \end{aligned}$$

We obtain the integral representation of the one-loop effect as before by using $\Omega_{\text{multi}}(x)$ and quasi-momenta for multi-magnons configurations.

$$\begin{aligned}
 \sum_{ij} (-1)^{F_{ij}} e^{-i(q_i - q_j)} &= \left(\prod_{i=1}^N \left(\frac{x - \frac{1}{X_i^-}}{x - \frac{1}{X_i^+}} \right)^2 \left(\frac{x - X_i^-}{x - X_i^+} \right)^2 + \prod_{j=N+1}^{N+M} \left(\frac{x - \frac{1}{X_j^-}}{x - \frac{1}{X_j^+}} \right)^2 \left(\frac{x - X_j^-}{x - X_j^+} \right)^2 \right) \\
 &\quad \times \left(1 + \prod_{i=1}^{M+N} \left(\frac{x - \frac{1}{X_i^+}}{x - \frac{1}{X_i^-}} \right) \left(\frac{x - X_i^+}{x - X_i^-} \right) - 2 \prod_{j=1}^{N+M} \left(\frac{x - X_j^+}{x - X_j^-} \right) \sqrt{\frac{X_i^-}{X_i^+}} \right)
 \end{aligned}$$

$$\times \prod_{i=1}^{N+M} \left(\frac{x - \frac{1}{X_i^+}}{x - \frac{1}{X_i^-}} \right) \sqrt{\frac{X_i^+}{X_i^-}}. \tag{4.1}$$

Here (i, j) pairs include only light modes. To compare this with multi-particle Lüscher formula, we need to compute the S -matrix factor which includes scatterings between a virtual fundamental magnon and multi dyonic magnons. The S -matrix factor is given by

$$\begin{aligned} \sum (-1)^{F_b} S_{\text{multi}} &= \sigma_{BES} \left(y, X_1^{Q_1} \right) \cdots \sigma_{BES} \left(y, X_{N+M}^{Q_{N+M}} \right) \frac{\eta \left(X_1^{Q_1} \right)}{\tilde{\eta} \left(X_1^{Q_1} \right)} \cdots \frac{\eta \left(X_{N+M}^{Q_{N+M}} \right)}{\tilde{\eta} \left(X_{N+M}^{Q_{N+M}} \right)} \\ &\times \left(\prod_{i=1}^N S_{BDS} \left(y, X_i^{Q_i} \right) + \prod_{i=N+1}^{N+M} S_{BDS} \left(y, X_i^{Q_i} \right) \right) \left(\frac{\eta(y)}{\tilde{\eta}(y)} \right)^{\sum Q_i} \sum_b (-1)^{F_b} \prod_{i=1}^{N+M} s_b(p_i). \end{aligned}$$

Using $s_b(p)$ defined in (3.5) and S_{BDS} in (3.6), we can obtain the final expression

$$\begin{aligned} \delta \Delta_F^{\text{multi}} &= \int \frac{dx}{2\pi i} \partial_x \Omega_{\text{multi}}(x) e^{-i\Delta_{\text{total}} \frac{x}{2g(x^2-1)}} \times \prod_{i=1}^{N+M} \left(\frac{x - \frac{1}{X_i^+}}{x - \frac{1}{X_i^-}} \sqrt{\frac{X_i^+}{X_i^-}} \right) \\ &\times \left(\prod_{i=1}^N \left(\frac{x - X_i^-}{x - X_i^+} \right)^2 \left(\frac{x - \frac{1}{X_i^-}}{x - \frac{1}{X_i^+}} \right)^2 + \prod_{j=N+1}^{N+M} \left(\frac{x - X_j^-}{x - X_j^+} \right)^2 \left(\frac{x - \frac{1}{X_j^-}}{x - \frac{1}{X_j^+}} \right)^2 \right) \\ &\times \left(1 + \prod_{k=1}^{N+M} \left(\frac{x - X_k^+}{x - X_k^-} \right) \left(\frac{x - \frac{1}{X_k^+}}{x - \frac{1}{X_k^-}} \right) - 2 \prod_{l=1}^{N+M} \left(\frac{x - X_l^+}{x - X_l^-} \right) \sqrt{\frac{X_l^-}{X_l^+}} \right). \end{aligned}$$

Here we have used multi-particle Lüscher formula [45]. It is straightforward to check that this result matches with the algebraic curve result (4.1).

5 Conclusions

In this paper we have computed the quantum finite-size effects for \mathbb{CP}^2 , \mathbb{RP}^3 , and big magnons and general combination of these magnons. We have computed the off-shell frequencies of magnons from the fluctuations of quasi-momenta and summed over all the physical polarizations. We have shown that these results agree with the F -term Lüscher formula based on the AdS_4/CFT_3 S -matrix proposed in [13]. This provides a stringent check for the S -matrix up to the one-loop order as well as the off-shell algebraic curve method.

In the computations, we have noticed that the unstable heavy modes should be suppressed in the $AdS_4 \times \mathbb{CP}^3$ algebraic curve. We also provide an on-shell particle interpretation of the big magnon, which was found in the context of the algebraic curve [20] and the dressing methods [21–23], as a bound-state of two small dyonic magnons with $u(1)$ charges Q and $-Q$, respectively. Our result shows that the Lüscher F -term based on this particle interpretation correctly reproduces the one-loop energy shift from the algebraic curve method. It is not clear at this moment if this is just mathematical coincidence or has a deeper physical meaning. In the AdS_5/CFT_4 context, it has been argued that a

dyonic magnon state with $-Q$ is an anti-magnon state which is obtained by reversing time direction [46]. It will be interesting if this argument also applies to AdS_4/CFT_3 .

The undetermined functions α_l arising in the algebraic curve method for multi-magnon states should be determined in the S -matrix approach. For example, the multi-particle Lüscher formula in [47] doesn't include any such functions. In fact, these functions are related with the variation of Bethe-Yang equations. Therefore, we can compute α_l from Bethe-Yang equations. In the case of the strong coupling limit, terms that contribute to α_l vanish in the F -term integration for multi-magnons by using saddle point method. However, it becomes important to determine these α_l at weak coupling regime such as in the five-loop Konishi computation [48].

Note added. While typing this article, we have found a paper [49] whose results on the off-shell algebraic curves overlap partially with ours.

Acknowledgments

We thank D. Bombardelli and I. Aniceto for informing us their result [49] and valuable comments. This work was supported in part by KRF-2007-313- C00150, WCU Grant No. R32-2008-000-101300 (C. A. and M. K.), and by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MEST) through the Center for Quantum Spacetime (CQUeST) of Sogang University with grant number 2005-0049409 (M. K. and B.-H. L.).

References

- [1] J.M. Maldacena, *The large- N limit of superconformal field theories and supergravity*, *Adv. Theor. Math. Phys.* **2** (1998) 231 [*Int. J. Theor. Phys.* **38** (1999) 1113] [[hep-th/9711200](#)] [[SPIRES](#)].
- [2] S.S. Gubser, I.R. Klebanov and A.M. Polyakov, *Gauge theory correlators from non-critical string theory*, *Phys. Lett. B* **428** (1998) 105 [[hep-th/9802109](#)] [[SPIRES](#)].
- [3] E. Witten, *Anti-de Sitter space and holography*, *Adv. Theor. Math. Phys.* **2** (1998) 253 [[hep-th/9802150](#)] [[SPIRES](#)].
- [4] O. Aharony, O. Bergman, D.L. Jafferis and J. Maldacena, *$N = 6$ superconformal Chern-Simons-matter theories, M2-branes and their gravity duals*, *JHEP* **10** (2008) 091 [[arXiv:0806.1218](#)] [[SPIRES](#)].
- [5] G. Arutyunov and S. Frolov, *Superstrings on $AdS_4 \times CP^3$ as a Coset σ -model*, *JHEP* **09** (2008) 129 [[arXiv:0806.4940](#)] [[SPIRES](#)].
- [6] B. Stefanski, jr, *Green-Schwarz action for Type IIA strings on $AdS_4 \times CP^3$* , *Nucl. Phys. B* **808** (2009) 80 [[arXiv:0806.4948](#)] [[SPIRES](#)].
- [7] N. Gromov and P. Vieira, *The AdS_4/CFT_3 algebraic curve*, *JHEP* **02** (2009) 040 [[arXiv:0807.0437](#)] [[SPIRES](#)].
- [8] N. Gromov and P. Vieira, *The all loop AdS_4/CFT_3 Bethe ansatz*, *JHEP* **01** (2009) 016 [[arXiv:0807.0777](#)] [[SPIRES](#)].

- [9] T. McLoughlin and R. Roiban, *Spinning strings at one-loop in $AdS_4 \times CP^3$* , *JHEP* **12** (2008) 101 [[arXiv:0807.3965](#)] [[SPIRES](#)].
- [10] L.F. Alday, G. Arutyunov and D. Bykov, *Semiclassical Quantization of Spinning Strings in $AdS_4 \times CP^3$* , *JHEP* **11** (2008) 089 [[arXiv:0807.4400](#)] [[SPIRES](#)].
- [11] C. Krishnan, *AdS_4/CFT_3 at One Loop*, *JHEP* **09** (2008) 092 [[arXiv:0807.4561](#)] [[SPIRES](#)].
- [12] T. McLoughlin, R. Roiban and A.A. Tseytlin, *Quantum spinning strings in $AdS_4 \times P^3$: testing the Bethe Ansatz proposal*, *JHEP* **11** (2008) 069 [[arXiv:0809.4038](#)] [[SPIRES](#)].
- [13] C. Ahn and R.I. Nepomechie, *$N=6$ super Chern-Simons theory S-matrix and all-loop Bethe ansatz equations*, *JHEP* **09** (2008) 010 [[arXiv:0807.1924](#)] [[SPIRES](#)].
- [14] D. Gaiotto, S. Giombi and X. Yin, *Spin Chains in $N = 6$ Superconformal Chern-Simons-Matter Theory*, *JHEP* **04** (2009) 066 [[arXiv:0806.4589](#)] [[SPIRES](#)].
- [15] M.C. Abbott, I. Aniceto and O. Ohlsson Sax, *Dyonic Giant Magnons in CP^3 : Strings and Curves at Finite J* , *Phys. Rev. D* **80** (2009) 026005 [[arXiv:0903.3365](#)] [[SPIRES](#)].
- [16] T.J. Hollowood and J.L. Miramontes, *A New and Elementary CP^n Dyonic Magnon*, *JHEP* **08** (2009) 109 [[arXiv:0905.2534](#)] [[SPIRES](#)].
- [17] G. Grignani, T. Harmark and M. Orselli, *The $SU(2) \times SU(2)$ sector in the string dual of $N = 6$ superconformal Chern-Simons theory*, *Nucl. Phys. B* **810** (2009) 115 [[arXiv:0806.4959](#)] [[SPIRES](#)].
- [18] G. Grignani, T. Harmark, M. Orselli and G.W. Semenoff, *Finite size Giant Magnons in the string dual of $N = 6$ superconformal Chern-Simons theory*, *JHEP* **12** (2008) 008 [[arXiv:0807.0205](#)] [[SPIRES](#)].
- [19] C. Ahn, P. Bozhilov and R.C. Rashkov, *Neumann-Rosochatius integrable system for strings on $AdS_4 \times CP^3$* , *JHEP* **09** (2008) 017 [[arXiv:0807.3134](#)] [[SPIRES](#)].
- [20] I. Shenderovich, *Giant magnons in AdS_4/CFT_3 : dispersion, quantization and finite-size corrections*, [arXiv:0807.2861](#) [[SPIRES](#)].
- [21] T.J. Hollowood and J.L. Miramontes, *Magnons, their Solitonic Avatars and the Pohlmeyer Reduction*, *JHEP* **04** (2009) 060 [[arXiv:0902.2405](#)] [[SPIRES](#)].
- [22] C. Kalousios, M. Spradlin and A. Volovich, *Dyonic Giant Magnons on CP^3* , *JHEP* **07** (2009) 006 [[arXiv:0902.3179](#)] [[SPIRES](#)].
- [23] R. Suzuki, *Giant Magnons on CP^3 by Dressing Method*, *JHEP* **05** (2009) 079 [[arXiv:0902.3368](#)] [[SPIRES](#)].
- [24] T. Lukowski and O. Ohlsson Sax, *Finite size giant magnons in the $SU(2) \times SU(2)$ sector of $AdS_4 \times CP^3$* , *JHEP* **12** (2008) 073 [[arXiv:0810.1246](#)] [[SPIRES](#)].
- [25] D. Bombardelli and D. Fioravanti, *Finite-Size Corrections of the CP^3 Giant Magnons: the Lüscher terms*, *JHEP* **07** (2009) 034 [[arXiv:0810.0704](#)] [[SPIRES](#)].
- [26] C. Ahn and P. Bozhilov, *Finite-size Effect of the Dyonic Giant Magnons in $N = 6$ super Chern-Simons Theory*, *Phys. Rev. D* **79** (2009) 046008 [[arXiv:0810.2079](#)] [[SPIRES](#)].
- [27] M. Beccaria, G.V. Dunne, V. Forini, M. Pawellek and A.A. Tseytlin, *Exact computation of one-loop correction to energy of spinning folded string in $AdS_5 \times S^5$* , *J. Phys. A* **43** (2010) 165402 [[arXiv:1001.4018](#)] [[SPIRES](#)].
- [28] N. Gromov, S. Schäfer-Nameki and P. Vieira, *Efficient precision quantization in AdS/CFT* , *JHEP* **12** (2008) 013 [[arXiv:0807.4752](#)] [[SPIRES](#)].

- [29] N. Gromov and P. Vieira, *The $AdS_5 \times S^5$ superstring quantum spectrum from the algebraic curve*, *Nucl. Phys. B* **789** (2008) 175 [[hep-th/0703191](#)] [[SPIRES](#)].
- [30] N. Gromov, S. Schäfer-Nameki and P. Vieira, *Quantum Wrapped Giant Magnon*, *Phys. Rev. D* **78** (2008) 026006 [[arXiv:0801.3671](#)] [[SPIRES](#)].
- [31] J.A. Minahan, A. Tirziu and A.A. Tseytlin, *Infinite spin limit of semiclassical string states*, *JHEP* **08** (2006) 049 [[hep-th/0606145](#)] [[SPIRES](#)].
- [32] J.A. Minahan and O. Ohlsson Sax, *Finite size effects for giant magnons on physical strings*, *Nucl. Phys. B* **801** (2008) 97 [[arXiv:0801.2064](#)] [[SPIRES](#)].
- [33] O. Ohlsson Sax, *Finite size giant magnons and interactions*, *Acta Phys. Polon. B* **39** (2008) 3143 [[arXiv:0810.5236](#)] [[SPIRES](#)].
- [34] M.A. Bandres and A.E. Lipstein, *One-Loop Corrections to Type IIA String Theory in $AdS_4 \times CP^3$* , *JHEP* **04** (2010) 059 [[arXiv:0911.4061](#)] [[SPIRES](#)].
- [35] K. Zarembo, *Worldsheet spectrum in AdS_4/CFT_3 correspondence*, *JHEP* **04** (2009) 135 [[arXiv:0903.1747](#)] [[SPIRES](#)].
- [36] G. Arutyunov, S. Frolov and M. Zamaklar, *The Zamolodchikov-Faddeev algebra for $AdS_5 \times S^5$ superstring*, *JHEP* **04** (2007) 002 [[hep-th/0612229](#)] [[SPIRES](#)].
- [37] N. Beisert, *The $SU(2|2)$ dynamic S-matrix*, *Adv. Theor. Math. Phys.* **12** (2008) 945 [[hep-th/0511082](#)] [[SPIRES](#)].
- [38] N. Beisert, B. Eden and M. Staudacher, *Transcendentality and crossing*, *J. Stat. Mech.* (2007) P01021 [[hep-th/0610251](#)] [[SPIRES](#)].
- [39] Y. Hatsuda and H. Tanaka, *Scattering of Giant Magnons in CP^3* , *JHEP* **02** (2010) 085 [[arXiv:0910.5315](#)] [[SPIRES](#)].
- [40] M. Lüscher, *Volume Dependence of the Energy Spectrum in Massive Quantum Field Theories. 1. Stable Particle States*, *Commun. Math. Phys.* **104** (1986) 177 [[SPIRES](#)].
- [41] T.R. Klassen and E. Melzer, *On the relation between scattering amplitudes and finite size mass corrections in QFT*, *Nucl. Phys. B* **362** (1991) 329 [[SPIRES](#)].
- [42] R.A. Janik and T. Lukowski, *Wrapping interactions at strong coupling – the giant magnon*, *Phys. Rev. D* **76** (2007) 126008 [[arXiv:0708.2208](#)] [[SPIRES](#)].
- [43] N. Beisert, *The Analytic Bethe Ansatz for a Chain with Centrally Extended $su(2|2)$ Symmetry*, *J. Stat. Mech.* (2007) P01017 [[nlin/0610017](#)].
- [44] Y. Hatsuda and R. Suzuki, *Finite-Size Effects for Dyonically Giant Magnons*, *Nucl. Phys. B* **800** (2008) 349 [[arXiv:0801.0747](#)] [[SPIRES](#)].
- [45] Y. Hatsuda and R. Suzuki, *Finite-Size Effects for Multi-Magnon States*, *JHEP* **09** (2008) 025 [[arXiv:0807.0643](#)] [[SPIRES](#)].
- [46] N. Dorey, D.M. Hofman and J.M. Maldacena, *On the singularities of the magnon S-matrix*, *Phys. Rev. D* **76** (2007) 025011 [[hep-th/0703104](#)] [[SPIRES](#)].
- [47] Z. Bajnok and R.A. Janik, *Four-loop perturbative Konishi from strings and finite size effects for multiparticle states*, *Nucl. Phys. B* **807** (2009) 625 [[arXiv:0807.0399](#)] [[SPIRES](#)].
- [48] Z. Bajnok, A. Hegedus, R.A. Janik and T. Lukowski, *Five loop Konishi from AdS/CFT* , *Nucl. Phys. B* **827** (2010) 426 [[arXiv:0906.4062](#)] [[SPIRES](#)].
- [49] M.C. Abbott, I. Aniceto and D. Bombardelli, *Quantum Strings and the AdS_4/CFT_3 Interpolating Function*, [arXiv:1006.2174](#) [[SPIRES](#)].