

# New discovery modes for a light charged Higgs boson at the LHC

---

**A. Arhrib,<sup>a</sup> R. Benbrik,<sup>b</sup> M. Krab,<sup>c</sup> B. Manaut,<sup>c</sup> S. Moretti,<sup>d</sup> Yan Wang<sup>e</sup> and Qi-Shu Yan<sup>f,9</sup>**

<sup>a</sup>*Faculty of Sciences and techniques, Abdelmalek Essaadi University, Tanger, Morocco*

<sup>b</sup>*Faculté Polydisciplinaire de Safi, Laboratoire de Physique Fondamentale et Appliquée Safi, Sidi Bouzid, B.P. 4162, Safi, Morocco*

<sup>c</sup>*Research Team in Theoretical Physics and Materials (RTTPM), Polydisciplinary Faculty, Sultan Moulay Slimane University, Beni Mellal, 23000, Morocco*

<sup>d</sup>*School of Physics and Astronomy, University of Southampton, Southampton, SO17 1BJ, U.K.*

<sup>e</sup>*College of Physics and Electronic Information, Inner Mongolia Normal University, Hohhot 010022, P.R. China*

<sup>f</sup>*Center for Future High Energy Physics, Chinese Academy of Sciences, Beijing 100049, P.R. China*

<sup>9</sup>*School of Physics Sciences, University of Chinese Academy of Sciences, Beijing 100039, P.R. China*

*E-mail: [aarhrib@gmail.com](mailto:aarhrib@gmail.com), [r.benbrik@uca.ma](mailto:r.benbrik@uca.ma), [mohamed.krab@usms.ac.ma](mailto:mohamed.krab@usms.ac.ma), [b.manaut@usms.ma](mailto:b.manaut@usms.ma), [s.moretti@soton.ac.uk](mailto:s.moretti@soton.ac.uk), [wangyan@imnu.edu.cn](mailto:wangyan@imnu.edu.cn), [yanqishu@ucas.ac.cn](mailto:yanqishu@ucas.ac.cn)*

ABSTRACT: At the Large Hadron Collider (LHC), both the ATLAS and CMS Collaborations have been searching for light charged Higgs bosons via top (anti)quark production and decays channels, like  $pp \rightarrow t\bar{t}$  with one top (anti)quark decaying into a charged Higgs boson and a  $b$  (anti)quark, when the decay is kinematically open (i.e., when  $m_{H^\pm} \lesssim m_t$ ). In this paper, we propose new searches at the LHC involving light charged Higgs bosons via their pair production channels like  $pp \rightarrow H^\pm h/A$  and  $pp \rightarrow H^+ H^-$  in the 2-Higgs Doublet Model (2HDM) Type-I and -X scenarios. By focusing on the case where the heavy  $H$  state plays the role of the Standard Model (SM)-like Higgs boson with a mass near 125 GeV, we study the aforementioned Higgs boson pair production channels and investigate their bosonic decays, such as  $H^\pm \rightarrow W^\pm h$  and/or  $H^\pm \rightarrow W^\pm A$ . We demonstrate that for a light charged Higgs boson state, with  $m_{H^\pm} \lesssim m_t$ , at the LHC, such di-Higgs production and decay channels can give rise to signatures with event rates much larger than those emerging from  $pp \rightarrow t\bar{t} \rightarrow t\bar{b}H^- + \text{c.c.}$  We specifically study  $h/A \rightarrow b\bar{b}$  and  $\tau^+\tau^-$  decays. We, therefore, claim that the discussed combination of new production and decay modes can result in an alternative discovery channel for charged Higgs bosons lighter than the top (anti)quark at the LHC within the above two 2HDM Types. Finally, in order to motivate experimentalists in ATLAS and CMS to search for such signatures, we propose 16 Benchmark Points (BPs) which are compatible with both theoretical and experimental constraints.

KEYWORDS: Beyond Standard Model, Higgs Physics

ARXIV EPRINT: [2106.13656](https://arxiv.org/abs/2106.13656)

---

**Contents**

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>A review of the 2HDM</b>	<b>3</b>
<b>3</b>	<b>Theoretical and experimental constraints</b>	<b>5</b>
<b>4</b>	<b>Numerical results</b>	<b>5</b>
<b>5</b>	<b>Benchmark points</b>	<b>11</b>
5.1	Benchmark points for Type-I	12
5.2	Benchmark points for Type-X	14
<b>6</b>	<b>Conclusions</b>	<b>15</b>

---

**1 Introduction**

Following the discovery of a 125 GeV Higgs boson in the first run of the LHC [1, 2], several studies of its properties were undertaken. The current situation is that the measured Higgs signal rates in all production and decay channels agree with the SM predictions at the  $\sim 2\sigma$  level [3, 4]. However, although the current LHC Higgs data are consistent with the SM, there is still the possibility that the observed Higgs state could be part of a model with an extended Higgs sector including, e.g., an extra doublet, singlet and/or triplet. Such a possibility is understandable since most of these extended Higgs sector scenarios possess a decoupling limit and give back the SM after integrating out the heavier states. Typically, most of the higher Higgs representations with an extra doublet predict one or more charged Higgs bosons. Therefore, it will be a smoking gun for new physics if a charged Higgs bosons were to be found. Indeed, one of the main goals of future LHC runs, in addition to raising the precision of the existing measurements of the discovered Higgs boson properties, is to pursue direct searches for new Higgs states in the quest to discover the evidence of new physics.

The 2-Higgs Doublet Model (2HDM) is one of the simplest extensions of the SM. It contains two Higgs doublets, labelled as  $\Phi_1$  and  $\Phi_2$ , which can generate masses to all fermions and gauge bosons. The particle spectrum of the 2HDM includes two CP even Higgs bosons ( $h$  and  $H$ , with  $m_h < m_H$ ), one CP odd Higgs boson ( $A$ ) and a pair of charged Higgs bosons ( $H^\pm$ ).<sup>1</sup> Within such a framework, at hadron colliders, a charged Higgs boson can be produced through several channels. In particular, a light charged Higgs boson (with  $m_{H^\pm} \leq m_t - m_b$ ) can be copiously produced from the  $pp \rightarrow t\bar{t}$  process

---

<sup>1</sup>Herein, we assume that the  $H$  state of the 2HDM is the discovered SM-like Higgs boson state.

via the top decay  $t \rightarrow bH^+$  (or the equivalent antitop mode). When kinematically allowed,  $pp \rightarrow t\bar{t} \rightarrow b\bar{b}H^-W^+ + \text{c.c.}$  provides the largest production rate for a light charged Higgs bosons. However, other production mechanisms have been studied in the literature, listed below.

- Single production:  $gb \rightarrow tH^- + \text{c.c.}$  and  $gg \rightarrow t\bar{b}H^-$ , the latter containing the  $t\bar{t}$  production and decay mode, appropriately combined [5–13].
- Associated production with a  $W^\pm$  gauge boson:  $gg \rightarrow W^\pm H^\mp$  and  $b\bar{b} \rightarrow W^\pm H^\mp$  [14–16].
- Production through  $W^\pm b$  scattering:  $qb \rightarrow q'H^+b$  [17, 18].
- Resonant production via  $c\bar{s}, c\bar{b} \rightarrow H^+$  [19–21].
- Associate production with a neutral Higgs:  $q\bar{q}' \rightarrow H^\pm\phi$  where  $\phi$  denotes one of the three neutral Higgs bosons,  $\phi = h, H$  or  $A$  [22] (see also refs. [23–27]).
- Pair production:  $gg \rightarrow H^+H^-$  and  $q\bar{q} \rightarrow H^+H^-$  [28, 29].

(See also refs. [30–33] for a review of all available  $H^\pm$  hadro-production modes in 2HDMs.)

Before the opening of the  $t\bar{b}$  channel, a charged Higgs boson decays predominantly in  $\tau\nu$  followed by  $c\bar{s}$ , with  $c\bar{b}$  being Cabibbo-Kobayashi-Maskawa (CKM) suppressed. However, it has been shown that, in the presence of a light CP-even,  $h$ , or CP-odd,  $A$ , Higgs boson, the bosonic decays of the charged Higgs bosons  $H^\pm \rightarrow W^\pm h/A$  would compete with the  $\tau\nu$  and  $c\bar{s}$  modes and could even be dominant in some cases [34, 35]. Once the  $t\bar{b}$  channel is open,  $\tau\nu$  and  $c\bar{s}$  would be suppressed, so that  $t\bar{b}$  would compete with the bosonic decay channels such as  $H^\pm \rightarrow W^\pm h/A$ .

A charged Higgs boson has been searched for at the Tevatron through top decays,  $pp \rightarrow t\bar{t} \rightarrow b\bar{b}H^-W^+$ , followed by either  $\tau\nu$  or  $c\bar{s}$  decays. Negative searches have been used to set a limit on  $\text{BR}(t \rightarrow bH^+) \times \text{BR}(H^+ \rightarrow \tau\nu)$  [36, 37]. Alongside Tevatron searches, charged Higgs bosons have been searched for also at LEP using  $e^+e^- \rightarrow \gamma^*, Z^* \rightarrow H^+H^-$  followed by either  $H^\pm \rightarrow \tau\nu$ ,  $H^\pm \rightarrow cs$  or  $H^\pm \rightarrow W^\pm A$  [38]. If the charged Higgs boson decays dominantly to  $\tau\nu$  or  $c\bar{s}$ , LEP2 sets a lower bound on the mass of the order of 80 GeV while in the case where charged Higgs decays are dominated by  $W^\pm A$ , via a light CP-odd Higgs state ( $m_A \approx 12$  GeV), the lower bound on the charged Higgs mass is about 72 GeV [38]. At the LHC, both ATLAS and CMS have been searching for charged Higgs bosons either from top decay, in case  $t \rightarrow bH^+$  is open, or from the first production mechanisms listed above,  $pp \rightarrow tH^-$  or  $pp \rightarrow t\bar{b}H^-$ , otherwise. For a light charged Higgs, ATLAS and CMS experiments have already set an exclusion on the product of Branching Ratios (BRs),  $\text{BR}(t \rightarrow bH^+) \times \text{BR}(H^+ \rightarrow \tau^+\nu)$  [39–42], of order percent in the mass range of 80 to 160 GeV. Other channels, such as  $H^+ \rightarrow c\bar{s}$ , have also been searched for by ATLAS and CMS. Assuming that  $\text{BR}(H^+ \rightarrow c\bar{s}) = 100\%$ , one can set a limit on  $\text{BR}(t \rightarrow bH^+)$  in the range 1.68% to 0.25% for a charged Higgs boson mass between 90 and 160 GeV [43–45]. In the case of a heavy charged Higgs boson, ATLAS and CMS provide a limit on cross section times  $\text{BR}(H^\pm \rightarrow \tau\nu_\tau)$  or  $\text{BR}(H^+ \rightarrow tb)$  [39, 40, 46] (but these are not relevant here).

The aim of this letter is to revisit the last two  $H^\pm$  production modes listed above in the context of the upcoming LHC Run 3, to assess to what extent they could supplement  $H^\pm$  searches via  $pp \rightarrow t\bar{t} \rightarrow b\bar{b}W^+H^- + c.c.$ . We will show that the production rates of such alternative di-Higgs production channels, via  $pp \rightarrow H^\pm h/A$  and  $pp \rightarrow H^+H^-$ , can be overwhelmingly larger than those of the top quark associated production channels. Specifically, we will show that the bosonic decays of a light charged Higgs boson, via  $H^\pm \rightarrow W^\pm h/A$ , could be dominant and lead to accessible signatures alternative to those emerging from the top-antitop production and decay. Hence, such modes may serve as new discovery channels for light  $H^\pm$  states at the LHC. We further stress that, just like  $pp \rightarrow t\bar{t}$ , also  $pp \rightarrow H^\pm h/A$  and  $pp \rightarrow H^+H^-$  are model independent as they are predominantly mediated by  $s$ -channel gauge boson exchange with no 2HDM parameters involved.<sup>2</sup>

The paper is organised as follows. In section 2 we give a brief review of the 2HDM and the Yukawa couplings used. In section 3 we list the theoretical and experimental constraints that will be used during our study. Our numerical results and a set of Benchmark Points (BPs) are given in section 4. Our conclusion is given in section 5.

## 2 A review of the 2HDM

The general 2HDM is obtained by extending the SM Higgs sector, which has one doublet scalar field  $\Phi_1$  with  $Y = +1$ , with an additional doublet scalar field  $\Phi_2$  with  $Y = +1$ . The most general renormalisable potential which is invariant under  $SU(2)_L \times U(1)_Y$  is given by [47]:

$$\begin{aligned}
 V_{\text{Higgs}}(\Phi_1, \Phi_2) = & \lambda_1(\Phi_1^\dagger\Phi_1)^2 + \lambda_2(\Phi_2^\dagger\Phi_2)^2 + \lambda_3(\Phi_1^\dagger\Phi_1)(\Phi_2^\dagger\Phi_2) + \lambda_4(\Phi_1^\dagger\Phi_2)(\Phi_2^\dagger\Phi_1) + \\
 & + \frac{1}{2} \left[ \lambda_5(\Phi_1^\dagger\Phi_2)^2 + \text{h.c.} \right] + \left\{ \left[ \lambda_6(\Phi_1^\dagger\Phi_1) + \lambda_7(\Phi_2^\dagger\Phi_2) \right] (\Phi_1^\dagger\Phi_2) + \text{h.c.} \right\} \\
 & - \left\{ m_{11}^2\Phi_1^\dagger\Phi_1 + m_{22}^2\Phi_2^\dagger\Phi_2 + \left[ m_{12}^2\Phi_1^\dagger\Phi_2 + \text{h.c.} \right] \right\}. \tag{2.1}
 \end{aligned}$$

By the hermiticity of eq. (2.1),  $\lambda_{1,2,3,4}$  as well as  $m_{11}^2$  and  $m_{22}^2$  are real-valued while the parameters  $\lambda_5, \lambda_6, \lambda_7$  and  $m_{12}^2$  are in general complex and can generate CP violation in the Higgs sector. After Electro-Weak Symmetry Breaking (EWSB) takes place, of the 8 degrees of freedom initially present in  $\Phi_1$  and  $\Phi_2$ , 3 correspond to the longitudinal components of  $W^\pm$  and  $Z^0$  while the remaining 5 appear as the physical Higgs fields mentioned in the previous section:  $h, H, A$  and  $H^\pm$ .

We finally end up with 7 free parameters in the 2HDM, which here we choose to be

$$m_h, m_H, m_A, m_{H^\pm}, \alpha, \beta \text{ and } m_{12}^2. \tag{2.2}$$

One of the CP-even Higgs bosons must be the one discovered at the LHC in 2012 (as intimated, we assume this to be the  $H$  state). Furthermore, in the list above,  $\alpha$  is the mixing angle between the two CP-even Higgs bosons while  $\tan \beta$  is the ratio of the Vacuum Expectation Values (VEVs) of the two Higgs doublet fields.

---

<sup>2</sup>In fact, even the  $pp \rightarrow H^\pm h$  channel is effectively model independent because of the properties of the SM-like Higgs boson discovered at the LHC, as we shall discuss later.

	$\kappa_u^h$	$\kappa_d^h$	$\kappa_l^h$	$\kappa_u^H$	$\kappa_d^H$	$\kappa_l^H$	$\kappa_u^A$	$\kappa_d^A$	$\kappa_l^A$
Type-I	$c_\alpha/s_\beta$	$c_\alpha/s_\beta$	$c_\alpha/s_\beta$	$s_\alpha/s_\beta$	$s_\alpha/s_\beta$	$s_\alpha/s_\beta$	$c_\beta/s_\beta$	$-c_\beta/s_\beta$	$-c_\beta/s_\beta$
Type-X	$c_\alpha/s_\beta$	$c_\alpha/s_\beta$	$-s_\alpha/c_\beta$	$s_\alpha/s_\beta$	$s_\alpha/s_\beta$	$c_\alpha/c_\beta$	$c_\beta/s_\beta$	$-c_\beta/s_\beta$	$s_\beta/c_\beta$

**Table 1.** Higgs couplings to fermions in the 2HDM Type-I and -X.

It is well known that the presence of a single doublet in the SM plays a dual role, of generating the mass of both the gauge bosons and fermions (the latter via the introduction of Yukawa couplings). This dynamic produces two strong constraints. The first constraint is linked to the bosonic sector:  $\rho = m_{W^\pm}^2/(m_Z^2 \cos^2 \theta_W^2) \approx 1$  which is in good agreement with experiments. The second constraint is linked to the fermionic sector and guarantees the absence of Flavour Changing Neutral Currents (FCNCs). So, any extension of the SM must be done by preserving these two constraints. In the framework of the 2HDM, the first constraint is satisfied because of the doublet representation of the Higgs fields, so long certain Higgs mass relations are respected. In the Yukawa sector, though, with the presence of two Higgs doublets, though, if we allow for EWSB like in the SM, we end up with FCNCs in the Yukawa sector at the tree level already. To avoid the latter, the most elegant solution is dictated by the Glashow-Weinberg theorem [48]: FCNCs can be avoided at tree level if we impose a discrete  $Z_2$  symmetry such that each fermion type couples only to one of the Higgs doublets. Thus, there are four Types of 2HDM [47]. In this study, though, we are interested only in two of these, Type-I and -X. In the 2HDM Type-I, only the doublet  $\Phi_2$  couples to all the fermions exactly like in the SM while in the 2HDM Type-X (or lepton-specific) all the quarks couple to  $\Phi_2$  and the charged leptons couple to  $\Phi_1$ . If we extend such a  $Z_2$  symmetry to the scalar potential, wherein  $\Phi_1 \rightarrow +\Phi_1$  and  $\Phi_2 \rightarrow -\Phi_2$ , this requires  $\lambda_6 = \lambda_7 = 0$ . Note that, however, the  $m_{12}^2$  term therein, being of dimension-2, can remain, as it would only break the  $Z_2$  symmetry softly, thereby allowing for FCNCs compliant with experimental constraints.

The neutral and charged Higgs couplings to fermions can be obtained from the Yukawa Lagrangian and are given by [47]

$$\begin{aligned}
 -\mathcal{L}_{\text{Yukawa}} = \sum_{f=u,d,l} & \left( \frac{m_f}{v} \kappa_f^h \bar{f} f h + \frac{m_f}{v} \kappa_f^H \bar{f} f H - i \frac{m_f}{v} \kappa_f^A \bar{f} \gamma_5 f A \right) + \\
 & \left( \frac{V_{ud}}{\sqrt{2}v} \bar{u} (m_u \kappa_u^A P_L + m_d \kappa_d^A P_R) d H^+ + \frac{m_l \kappa_l^A}{\sqrt{2}v} \bar{\nu}_L l_R H^+ + \text{H.c.} \right), \quad (2.3)
 \end{aligned}$$

where  $\kappa_f^S$  are the Yukawa couplings in the 2HDM, which are listed in table 1 for the two Types of interest here. Here,  $V_{ud}$  represents a CKM matrix element.

As one can read from such a table, the Higgs couplings to quarks are identical in 2HDM Type-I and -X, the only difference being the coupling to leptons. From the phenomenological point of view, the most relevant features are that, in the 2HDM Type-X, the Higgs couplings to leptons enjoy an enhancement for large  $\tan \beta$  while, in the 2HDM Type-I, the couplings to all fermions are suppressed if  $\tan \beta \gg 1$ .

### 3 Theoretical and experimental constraints

The 2HDM parameter space is limited by both theoretical and experimental constraints. The theoretical constraints that have been imposed on the scalar potential are as follows.

- Vacuum stability conditions, which require the Higgs potential to be bounded from below [49], are given by the following inequalities:

$$\lambda_1 > 0, \quad \lambda_2 > 0, \quad \lambda_3 > -\sqrt{\lambda_1 \lambda_2}, \quad \lambda_3 + \lambda_4 - |\lambda_5| > -\sqrt{\lambda_1 \lambda_2}. \quad (3.1)$$

- Perturbativity constraints imply that all quartic coefficients of the scalar potential satisfy the condition  $|\lambda_i| \leq 8\pi$  ( $i = 1, \dots, 5$ ).
- Perturbative unitarity constraints [50, 51] require that scattering processes involving gauge bosons and Higgs states remain unitary at high energy.

We also take into account experimental constraints from direct Higgs boson searches at LEP, Tevatron, LHC as well as Electroweak Precision Observables (EWPOs) plus flavour physics data are also used.

- Measurements of the oblique parameters  $S$ ,  $T$  and  $U$  [52] can constrain several new physics models and, in the case of the 2HDM, they can be used to set a limit on the mass splitting between the physical Higgs bosons (specifically, a charged and a neutral Higgs state) and on the mixing angles as well. The 2HDM contributions to  $S$ ,  $T$  and  $U$  parameters are taken from refs. [53–55].
- Exclusion limits at 95% Confidence Level (CL) from Higgs searches at colliders (LEP, Tevatron and LHC) via `HiggsBounds-5.9.0` [56] and compliance with SM-like Higgs boson measurements via `HiggsSignals-2.6.0` [57] are enforced.
- B-physics observables are calculated using the public code `SuperIso v4.1` [58]. Specifically, we have used the following observables:

1.  $\text{BR}(B \rightarrow X_s \gamma) = (3.32 \pm 0.15) \times 10^{-4}$  [59],
2.  $\text{BR}(B_s \rightarrow \mu^+ \mu^-) = (3.0 \pm 0.6 \pm 0.25) \times 10^{-9}$  [60],
3.  $\text{BR}(B_u \rightarrow \tau \nu) = (1.06 \pm 0.19) \times 10^{-4}$  [59].

Note that we have used the public code `2HDMC` [61] to check the theoretical constraints such as perturbative unitarity, perturbativity of the  $\lambda_i$ , vacuum stability of the scalar potential as well as the oblique parameters  $S$ ,  $T$  and  $U$ .

### 4 Numerical results

As previously mentioned, in what follows, we assume that the SM-like Higgs particle discovered at CERN in 2012 is the  $H$  state, for which  $m_H = 125 \text{ GeV}$ , so that  $m_h$  would be smaller in comparison. We then perform a systematic scan over the 2HDM parameter space as indicated in table 2, for both Type-I and -X.

Parameters	2HDM-I, -X
$m_h$	[10, 120]
$m_H$	125
$m_A$	[10, 120]
$m_{H^\pm}$	[80, 170]
$s_{\beta-\alpha}$	[-0.3, -0.05]
$\tan \beta$	[2, 60]
$m_{12}^2$	$[0, m_H^2 \sin \beta \cos \beta]$
$\lambda_6 = \lambda_7$	0

**Table 2.** 2HDM Type-I and -X parameter scans adopted (all masses are in GeV).

Since we assume that  $m_H = 125$  GeV, then the  $S$ ,  $T$  and  $U$  constraints will force the whole Higgs boson spectrum of the 2HDM to be rather light. The charged Higgs is taken in the range 80–170 GeV and the CP-odd is presumed to be in the range 10–120 GeV. Having assumed that  $H$  is the known Higgs particle, taking into account all LHC data will force the  $H$  couplings to SM particles to be rather SM-like. As a consequence, the coupling of the  $H$  state to gauge bosons  $W^+W^-$  (and  $ZZ$ ), which is proportional to  $\cos(\beta - \alpha)$ , would be SM-like if  $\cos(\beta - \alpha) \approx 1$ . This trigonometric function is the same entering the coupling  $W^+H^-h$ , which is then essentially independent of the  $\alpha$  and  $\beta$  parameters of the 2HDM, as already remarked (just like  $W^+H^-A$ ).


We are interested in light charged Higgs bosons, which could come either from (anti)top decay or Higgs pair production: i.e.,  $gg, q\bar{q} \rightarrow t\bar{b}H^- + \text{c.c.}$ ,  $q\bar{q} \rightarrow H^+H^-$ , and  $q\bar{q}' \rightarrow H^+h/A + \text{c.c.}$ <sup>3</sup> We compute cross sections at Leading Order (LO) only (i.e., at tree level), though Next-to-LO (NLO) corrections from Quantum Chromo-Dynamics (QCD) exist for all processes: see refs. [62–64] for the top (anti)quark process while those for Higgs pair production can be accounted for through to the use of NLO Parton Distribution Functions (PDFs). The former (30%) is somewhat larger than the latter (20%) for a light  $H^\pm$  state, so such different relative effects from QCD should be borne in mind while comparing the two types of processes, but their inclusion will not change our conclusions.

We now examine the production rates of light charged Higgs boson at the LHC with  $\sqrt{s} = 14$  TeV from the above processes. This is done for the 2HDM Type-I in figure 1, over the range  $160 \text{ GeV} \lesssim m_{H^\pm} \lesssim 350 \text{ GeV}$ ,<sup>4</sup> for the following values of the other BSM parameters:  $m_h = 94$  GeV,  $m_A = 105$  GeV,  $\sin(\beta - \alpha) = -0.09$  and  $m_{12}^2 = 294 \text{ GeV}^2$ . Here, the left(right) frame refers to  $\tan \beta = 3(30)$ . Let us start with the rates for  $\sigma(q\bar{q}, gg \rightarrow t\bar{t})$

<sup>3</sup>Therefore, in our approach, we privilege the use of the  $2 \rightarrow 3$  description of the first channel, i.e., in the 4-Flavour Scheme (4FS), as it is the most suitable to model the so-called ‘transition region’  $m_{H^\pm} \approx m_t$  [11, 65]. However, as it is customary in many experimental analyses, we will also present results for the process  $pp \rightarrow t\bar{t}$  followed by  $t \rightarrow bH^+$  (and c.c.), wherein the (anti)top quark is treated in Narrow Width Approximation (NWA), again, in LO approximation.

<sup>4</sup>This is done for illustrative purposes, though, as the maximum allowed value for  $m_{H^\pm}$  in the 2HDM Type-I and -X setups is actually 197 GeV since, for higher values, one would no longer satisfy  $S$ ,  $T$  and  $U$  constraints.





**Figure 1.** Cross section for  $gg, q\bar{q} \rightarrow t\bar{b}H^- + c.c.$  (black),  $gg, q\bar{q} \rightarrow t\bar{t} \times \text{BR}(t \rightarrow bH^+) + c.c.$  (orange),  $q\bar{q} \rightarrow H^+H^-$  (red) and  $q\bar{q}' \rightarrow H^\pm h/A$  (green/blue) at  $\sqrt{s} = 14$  TeV, as a function of  $m_{H^\pm}$  for  $\tan\beta = 3$  (left) and 30 (right). All cross sections are calculated using the code MadGraph [66] at LO. Note that we have used the MSTW2008 PDF set from [67] with factorisation/renormalisation scale set to  $\sqrt{\hat{s}}$ , i.e., the centre-of-mass energy at parton level.

$\times \text{BR}(t \rightarrow bH^+) + c.c.$ , which are clearly only appearing for  $m_{H^\pm} \leq m_t - m_b$  and coincide with those for  $\sigma(q\bar{q}, gg \rightarrow t\bar{b}H^- + c.c.)$  in the limit  $m_{H^\pm} \ll m_t$ . The latter, of course, extends beyond such a transition region, though the absence of any resonant contribution in this (heavy) mass region reflects a significant drop in cross section. Clearly, as shown in literature and as the plots in figure 1 demonstrate, the  $pp \rightarrow t\bar{b}H^- + c.c.$  process is important for charged Higgs boson phenomenology at the LHC. However, it is worth noticing that the  $\text{BR}(t \rightarrow bH^+)$  can only be really significant in a parameter space region with small  $\tan\beta$  due to the enhancement of the coupling  $H^+\bar{t}b$  in this model configuration. As can be seen from figure 1 (left panel), 2HDM Type-I predicts a scenario where, for  $m_{H^\pm} \ll m_t$ ,  $\sigma(q\bar{q}, gg \rightarrow t\bar{t}) \times \text{BR}(t \rightarrow bH^+) + c.c.$  (and  $\sigma(q\bar{q}, gg \rightarrow t\bar{b}H^- + c.c.)$ ) could reach values around 30 pb and, in fact, even for very large charged Higgs masses, the rates for the  $pp \rightarrow t\bar{b}H^- + c.c.$  channel remain larger than those for  $pp \rightarrow H^\pm h$ ,  $pp \rightarrow H^\pm A$  as well as  $pp \rightarrow H^+H^-$ . However, as demonstrated in the right panel of figure 1, when the parameter  $\tan\beta$  is large, the top (anti)quark channels are suppressed by the Yukawa couplings, so that the Higgs pair production channels can largely dominate the  $H^\pm$  phenomenology, no matter the value of its mass. Quite interestingly, once can then exploit the decay channels  $H^\pm \rightarrow W^\pm\phi$ , with  $\phi = h$  or  $A$ . Finally, being the neutral Higgs bosons emerging therein rather light, it is also noted that they finally dominantly decay into two fermions, either  $b$ 's or  $\tau$ 's.


With this in mind, in table 3, we tabulate the typical final states (left column) and corresponding production and decay patterns (right column) for top pair production and decay, charged Higgs boson associated production with top plus bottom and di-Higgs processes. Notice that the first two channels always have  $2W$  bosons in the final states, similarly to the charged Higgs boson pair production channel. In contrast, the di-Higgs production

	Top pair production and decay chain
$\sigma_{2t}^{h_i}(2W+2b+2f)$	$2 \sigma_{t\bar{t}} \times \text{BR}(t \rightarrow bH^+) \times \text{BR}(\bar{t} \rightarrow \bar{b}W^-) \times \text{BR}(H^\pm \rightarrow W^\pm h_i) \times \text{BR}(h_i \rightarrow f\bar{f})$
	Associated production with top plus bottom and decay chain
$\sigma_t^{h_i}(2W+2b+2f)$	$\sigma(pp \rightarrow t\bar{b}H^-) \times \text{BR}(t \rightarrow bW^+) \times \text{BR}(H^\pm \rightarrow W^\pm h_i) \times \text{BR}(h_i \rightarrow f\bar{f})$
	Di-Higgs production and decay chain
$\sigma_{h_j}^{h_i}(2W+2f+2f')$	$\frac{1}{1+\delta_{ff'}} \sigma(H^+H^-) \times \text{BR}(H^\pm \rightarrow W^\pm h_i) \times \text{BR}(H^\pm \rightarrow W^\pm h_j) \times (\text{BR}(h_i \rightarrow f\bar{f}) \times \text{BR}(h_j \rightarrow f'\bar{f}') + \text{BR}(h_j \rightarrow f\bar{f}) \times \text{BR}(h_i \rightarrow f'\bar{f}'))$
$\sigma_{h_j}^{h_i}(W+2f+2f')$	$\frac{1}{1+\delta_{ff'}} \sigma(H^\pm h_i) \times \text{BR}(H^\pm \rightarrow W^\pm h_j) \times (\text{BR}(h_i \rightarrow f\bar{f}) \times \text{BR}(h_j \rightarrow f'\bar{f}') + \text{BR}(h_j \rightarrow f\bar{f}) \times \text{BR}(h_i \rightarrow f'\bar{f}'))$


**Table 3.** The discussed production processes of charged Higgs bosons are shown here, where the main decay chains which can lead to the given final states are also provided. Here, we define  $i, j = 1, 2$  and have  $h_1 = h$  and  $h_2 = A$ . The symbol  $f(f')$  denotes fermions, like  $b$  or  $\tau$ .

channels  $pp \rightarrow H^\pm A/h$  lead to only a single  $W$  in the final states. Below, we will focus on three types of final states, the  $4b$  final states (including  $2W+4b$  and  $W+4b$ ), the  $2b2\tau$  final states (including  $2W+2b2\tau$  and  $W+2b2\tau$ ) and the  $4\tau$  final state (including only  $W+4\tau$ ).

In figure 2, again, for the 2HDM Type-I scenario and the parameter ranges given in table 2, we show the relative magnitudes of the production and decay rates of  $4b$  final states from  $pp \rightarrow H^\pm h \rightarrow W^\pm h h$  and from the two top (anti)quark processes. In the left(right) panel, the cross sections of the di-Higgs process  $\sigma_h^h(W+4b)$  are compared with those of the process  $\sigma_{2t}(2W+4b)(\sigma_t(2W+4b))$ . The red points shown in these two plots demonstrate cases where also the  $pp \rightarrow H^+H^-$  process yielding  $\sigma_h^h(2W+4b)$  can be larger than the top (anti)quark processes. In figure 3, similar to figure 2, we show the corresponding results of the production of  $4b$  final states from  $pp \rightarrow H^\pm A \rightarrow W^\pm AA$ . It is therefore clear that the combination of charged and neutral current di-Higgs production can provide a significant alternative discovery channel for  $H^\pm$  states at the LHC in the 2HDM Type-I. This conclusion can in fact be extended to the case of the  $\sigma_h^h(2W+2b2\tau)$  and the  $\sigma_h^h(W+2b2\tau)$  signatures versus the  $\sigma_{2t}(2W+2b2\tau)$  or  $\sigma_t(2W+2b2\tau)$  ones, too, as demonstrated in figures 4–5. In fact, these last two figures also report 2HDM Type-X rates for  $\sigma(pp \rightarrow H^\pm h) \times \text{BR}(H^\pm \rightarrow W^\pm h) \times \text{BR}(h \rightarrow \tau^+\tau^-)^2$  (figure 4) and  $\sigma(pp \rightarrow H^\pm A) \times \text{BR}(H^\pm \rightarrow W^\pm A) \times \text{BR}(A \rightarrow \tau^+\tau^-)^2$  (figure 5), showing that, for this Yukawa structure, the  $W+4\tau$  signature is the one of interest, as here  $\text{BR}(h/A \rightarrow b\bar{b})$  is very small compared to  $\text{BR}(h/A \rightarrow \tau^+\tau^-)$ . Conversely, the  $W+4\tau$  signature is of little relevance for the 2HDM Type-I. As for the  $2W+4\tau$  case, this is not accessible in either scenario. Needless to say, for all the parameter space points considered, in both 2HDM Type-I and -X, the decay channels of the charged Higgs boson are dominated by  $H^\pm \rightarrow Wh/A$  with a BR that could reach 100% in the alignment limit [31, 32, 34].




**Figure 2.** Values of  $\sigma(pp \rightarrow H^\pm h) \times \text{BR}(H^\pm \rightarrow W^\pm h) \times \text{BR}(h \rightarrow b\bar{b})^2$  are compared with those of  $2 \sigma_{t\bar{t}} \times \text{BR}(t \rightarrow bH^+) \times \text{BR}(\bar{t} \rightarrow \bar{b}W^-) \times \text{BR}(H^\pm \rightarrow W^\pm h) \times \text{BR}(h \rightarrow b\bar{b})$  (left) and  $\sigma(pp \rightarrow t\bar{b}H^-) \times \text{BR}(t \rightarrow bW^+) \times \text{BR}(H^\pm \rightarrow W^\pm h) \times \text{BR}(h \rightarrow b\bar{b})$  (right). The red points identify the values of  $\sigma(H^+H^-) \times \text{BR}(H^\pm \rightarrow W^\pm h)^2 \times \text{BR}(h \rightarrow b\bar{b})^2$  which also exceeds those of the latter two top mediated processes, respectively. The colour bar denotes the mass of the charged Higgs boson. Results are for the 2HDM Type-I.




**Figure 3.** Values of  $\sigma(pp \rightarrow H^\pm A) \times \text{BR}(H^\pm \rightarrow W^\pm A) \times \text{BR}(A \rightarrow b\bar{b})^2$  are compared with those of  $2 \sigma_{t\bar{t}} \times \text{BR}(t \rightarrow bH^+) \times \text{BR}(\bar{t} \rightarrow \bar{b}W^-) \times \text{BR}(H^\pm \rightarrow W^\pm A) \times \text{BR}(A \rightarrow b\bar{b})$  (left) and  $\sigma(pp \rightarrow t\bar{b}H^-) \times \text{BR}(t \rightarrow bW^+) \times \text{BR}(H^\pm \rightarrow W^\pm A) \times \text{BR}(A \rightarrow b\bar{b})$  (right). The red points identify the values of  $\sigma(H^+H^-) \times \text{BR}(H^\pm \rightarrow W^\pm A)^2 \times \text{BR}(A \rightarrow b\bar{b})^2$  which also exceeds those of the latter two top mediated processes, respectively. The colour bar denotes the mass of the charged Higgs boson. Results are for the 2HDM Type-I.

Finally, it should be pointed out that, when a final state is specified, the di-Higgs signature should sum over all processes which can produce such a final state. For example, when  $W + 4b$  is considered, the cross section from di-Higgs processes should be summed over those of the intermediate states  $W^\pm + hh$ ,  $W^\pm + hA$  (or  $W^\pm + Ah$ ) and  $W^\pm + AA$ , in



**Figure 4.** Values of  $2 \sigma(pp \rightarrow H^\pm h) \times \text{BR}(H^\pm \rightarrow W^\pm h) \times \text{BR}(h \rightarrow b\bar{b}) \times \text{BR}(h \rightarrow \tau^+\tau^-)$  are compared with those of  $2 \sigma_{t\bar{t}} \times \text{BR}(t \rightarrow bH^+) \times \text{BR}(\bar{t} \rightarrow \bar{b}W^-) \times \text{BR}(H^\pm \rightarrow W^\pm h) \times \text{BR}(h \rightarrow \tau^+\tau^-)$  (left) and  $\sigma(pp \rightarrow t\bar{b}H^-) \times \text{BR}(t \rightarrow bW^+) \times \text{BR}(H^\pm \rightarrow W^\pm h) \times \text{BR}(h \rightarrow \tau^+\tau^-)$  (right). The red points identify the values of  $\sigma(H^+H^-) \times \text{BR}(H^\pm \rightarrow W^\pm h)^2 \times \text{BR}(h \rightarrow b\bar{b}) \times \text{BR}(h \rightarrow \tau^+\tau^-)$  which also exceeds those of the latter two top mediated processes, respectively. The colour bar denotes the mass of the charged Higgs boson. Results are for the 2HDM Type-I with the exception of the black points, which refer to the 2HDM Type-X rates for  $\sigma(pp \rightarrow H^\pm h) \times \text{BR}(H^\pm \rightarrow W^\pm h) \times \text{BR}(h \rightarrow \tau^+\tau^-)^2$ .



**Figure 5.** Values of  $2 \sigma(pp \rightarrow H^\pm A) \times \text{BR}(H^\pm \rightarrow W^\pm A) \times \text{BR}(A \rightarrow b\bar{b}) \times \text{BR}(A \rightarrow \tau^+\tau^-)$  are compared with those of  $2 \sigma_{t\bar{t}} \times \text{BR}(t \rightarrow bH^+) \times \text{BR}(\bar{t} \rightarrow \bar{b}W^-) \times \text{BR}(H^\pm \rightarrow W^\pm A) \times \text{BR}(A \rightarrow \tau^+\tau^-)$  (left) and  $\sigma(pp \rightarrow t\bar{b}H^-) \times \text{BR}(t \rightarrow bW^+) \times \text{BR}(H^\pm \rightarrow W^\pm A) \times \text{BR}(A \rightarrow \tau^+\tau^-)$  (right). The red points identify the values of  $\sigma(H^+H^-) \times \text{BR}(H^\pm \rightarrow W^\pm A)^2 \times \text{BR}(A \rightarrow b\bar{b}) \times \text{BR}(A \rightarrow \tau^+\tau^-)$  which also exceeds those of the latter two top mediated processes, respectively. The colour bar denotes the mass of the charged Higgs boson. Results are for the 2HDM Type-I with the exception of the black points, which refer to the 2HDM Type-X rates for  $\sigma(pp \rightarrow H^\pm A) \times \text{BR}(H^\pm \rightarrow W^\pm A) \times \text{BR}(A \rightarrow \tau^+\tau^-)^2$ .

	Allowed by theoretical constraints	Allowed by HiggsBounds	Allowed by HiggsSignals	Allowed by all constraints
Type-I	5.36%	17.96%	55.36%	0.53%
Type-X	5.34%	1.37%	7.50%	0.0055%

**Table 4.** Details of the allowed points in parameter space after imposing theoretical and experimental constraints. Note that EWPO and flavour constraints are essentially always satisfied in both 2HDM Types. Here, we have used a dataset with  $5 \times 10^6$  sample points.

turn emerging from the processes  $pp \rightarrow H^\pm h/A$ . Obviously, the ability to rely on different subchannels all leading to the same final state is a welcome feature when we tension the new discovery modes to the established ones. However, notice that in the ‘mixed decay chain’, i.e., when an intermediate state of the type  $W^\pm + hA$  (or  $W^\pm + Ah$ ) is produced, from a selection analysis perspective, it would not be possible to enforce an equal mass constraint on the fermionic systems comprised by two  $b\bar{b}$  pairs or  $b\bar{b}$  and  $\tau^+\tau^-$  pairs, so we do expect the experimental efficiency of this case to be less than when two identical neutral Higgs bosons emerge from the production and decay stages of the di-Higgs processes. This is why we have refrained from presenting the equivalent of figures 2–5 for the mixed decay chain.

## 5 Benchmark points

In order to examine the salient features of the advocated parameter space and to encourage experimentalists to search for such new discovery channels at the LHC (and/or future colliders), we propose eight Benchmark Points (BPs) for both the 2HDM Type-I and -X scenarios. These BPs are presented in tables 5 and 6, respectively. BPs satisfying both conditions  $m_{H^\pm} + m_b > m_t$  and  $m_{H^\pm} + m_b < m_t$  are chosen.

Before we list our BPs in detail, we first show the fractions of parameter space that survive theoretical and experimental constraints. This information is given in table 4 for both Type-I and -X. In the Type-I scenario, some 0.53% of the points generated in our scan can survive after all constraints have been imposed. In contrast, in the Type-X scenario, only 0.0055% survive. It is found that the fractions of points that survive the theoretical constraints for these two scenarios are almost equal, which is understandable since they share the same Higgs potential. Compared with the Type-I scenario, in Type-X, fewer points pass HiggsBounds constraints enforcing limits from nil searches for additional Higgs bosons (beyond the SM-like one) performed at LEP, Tevatron and the LHC. Similarly, for the Type-X scenario, again, fewer points than in Type-I pass the constraints from the SM-like Higgs boson data gathered by the LHC experiments and included in HiggsSignals. The reason can be attributed to the fact that, when the couplings to quarks are suppressed and those to leptons are enhanced, it is crucial to predict the correct production rate for  $gg \rightarrow H$  and the correct decay fractions for  $H \rightarrow b\bar{b}$  and  $H \rightarrow \gamma\gamma$  and, in the light of the couplings in table 1, this is more easily achieved in Type-I than -X.

For the BPs given in tables 5 and 6, it is observed that the mass splittings between the two lightest neutral Higgs bosons ( $h$  and  $A$ ) are not allowed to be too large, which is

also due to the constraints from EWPOs. It should also be emphasised that these BPs have relatively light neutral Higgs bosons, which can also be searched for at future lepton colliders, like the 240 GeV ‘Higgs factories’ of CEPC, FCC-ee and ILC, where the search modes would involve four fermion final states, like  $4b$ ,  $2b2\tau$  and  $4\tau$ , while at the LHC such signal events might be overwhelmingly concealed by the huge QCD background.<sup>5</sup> Finally, notice that, for all our BPs, to compute the production and decay rates, we have taken  $m_H = 125$  GeV,  $m_t = 172.5$  GeV,  $m_b = 4.75$  GeV and  $m_\tau = 1.77684$  GeV.

### 5.1 Benchmark points for Type-I

In table 5, we present eight BPs for the 2HDM Type-I scenario. Here, BP1,2,3 and 5 satisfy the condition  $m_{H^\pm} + m_b < m_t$  while the rest of the points satisfy the condition  $m_{H^\pm} + m_b > m_t$ . For every BP in the table, we provide the cross sections of the  $W + 4b$  and  $W + 2b2\tau$  signatures from both di-Higgs and top (anti)quark processes, to gauge the relative yields against alternative numerical estimates. Let us now discuss the features of each BP. There are two observations to make regarding BP1.

1. The total production rate of  $2W + 4b$  from top (anti)quark processes is around 5 fb,<sup>6</sup> while the total production rate of  $2W + 4b$  from Higgs pair production of charged Higgs bosons is  $\approx 23.1$  fb. Furthermore, the production rate of  $W + 4b$  from di-Higgs (charged current) production channels can reach to 174 fb, which is 7 times larger than the final state of  $2W + 4b$  from the di-Higgs (neutral current) production channels.
2. While for the total production rate of  $2W + 2b2\tau$  the top (anti)quark processes can yield only 0.46 fb, pair production of charged Higgs bosons yields 2.13 fb. Furthermore, the production rate of  $W + 2b2\tau$  from di-Higgs processes can even reach 16 fb.

Such two observations also hold for BP2, 3 and 5. For BP4,6 and 8, the charged Higgs bosons are so heavy that a top (anti)quark cannot decay into a charged Higgs boson plus a  $b$  (anti)quark. Therefore, the corresponding  $\sigma_{2t}(2W + 4b)$  rates vanish due to this kinematic reason while  $\sigma_t(2W + 4b)$  is generally rather small, below the fb level. For BP4, the production rate of  $2W + 4b$  from top (anti)quark processes yields a cross section of 3.88 fb while that from Higgs-pair production involving charged Higgs bosons yields 20.24 fb and that of  $W + 4b$  yields 169.35 fb. For the final state  $2W + 2b2\tau$ , the production rate from top (anti)quark processes is 0.36 fb while that from pair production of charged Higgs bosons is 1.87 fb. Finally, the signature  $W + 2b2\tau$  from di-Higgs channels is 15.63 fb. (Such features are similar to those of BP6, 7 and 8.)

---

<sup>5</sup>In this sense, the triggering possibilities (e.g., by exploiting  $W^\pm \rightarrow l^\pm \nu_l$  decays, with  $l = e, \mu$ ) in our advocated LHC signatures offered by the presence of one or two  $W^\pm$ 's are crucial to establishing the corresponding signatures, involving such four fermion combinations.

<sup>6</sup>Here, we refer to the  $q\bar{q}, gg \rightarrow t\bar{b}H^- + \text{c.c.}$  rates, as, to remind the reader, the  $q\bar{q}, gg \rightarrow t\bar{t} \rightarrow t\bar{b}H^-$  case is provided only for reference purposes, as this channel is what is typically being used in experimental analyses for the case  $m_{H^\pm} < m_t$ .

Parameters	BP1	BP2	BP3	BP4	BP5	BP6	BP7	BP8
$m_h$ (GeV)	91.00	96.84	103.34	99.61	95.57	94.00	94.00	94.00
$m_H$ (GeV)	125.00	125.00	125.00	125.00	125.00	125.00	125.00	125.00
$m_A$ (GeV)	102.04	112.35	93.80	88.98	94.41	105.00	105.00	105.00
$m_{H^\pm}$ (GeV)	167.02	166.34	161.02	169.46	167.02	176.00	186.00	196.00
$s_{\beta-\alpha}$	-0.18	-0.11	-0.19	-0.06	-0.09	-0.09	-0.09	-0.09
$\tan\beta$	40.87	58.17	54.79	39.10	32.44	30.00	30.00	30.00
$m_{12}^2$ (GeV <sup>2</sup> )	204.22	161.85	196.73	252.94	277.81	294.00	294.00	294.00
$\sigma_{2t}^h(2W+4b)$	2.30	1.65	2.06	–	2.42	–	–	–
$\sigma_t^h(2W+4b)$	3.85	2.35	2.26	0.85	3.84	5.03	4.68	3.52
$\sigma_{2t}^A(2W+4b)$	0.70	0.25	4.63	–	2.47	–	–	–
$\sigma_t^A(2W+4b)$	1.17	0.36	5.07	3.03	3.92	0.83	0.44	1.08
$\sigma_h^h(2W+4b)$	13.58	15.99	2.29	0.97	5.38	14.08	13.27	7.35
$\sigma_A^h(2W+4b)$	4.13	2.44	5.14	3.46	5.50	2.32	1.25	2.24
$\sigma_A^A(2W+4b)$	1.26	0.37	11.55	12.35	5.62	0.38	0.12	0.68
$\sigma_h^A(2W+4b)$	4.13	2.44	5.14	3.46	5.50	2.32	1.25	2.24
$\sigma_h^h(W+4b)$	75.88	77.61	26.47	17.68	46.00	73.25	68.00	48.81
$\sigma_A^h(W+4b)$	23.07	11.86	59.44	63.04	47.00	12.07	6.42	14.90
$\sigma_A^A(W+4b)$	17.48	6.12	64.39	69.22	43.51	9.16	4.91	11.45
$\sigma_h^A(W+4b)$	57.51	40.06	28.68	19.41	42.59	55.59	52.02	37.51
$\sigma_{2t}^h(2W+2b2\tau)$	0.21	0.15	0.19	–	0.22	–	–	–
$\sigma_t^h(2W+2b2\tau)$	0.35	0.22	0.21	0.08	0.35	0.46	0.43	0.32
$\sigma_{2t}^A(2W+2b2\tau)$	0.07	0.02	0.43	–	0.23	–	–	–
$\sigma_t^A(2W+2b2\tau)$	0.11	0.03	0.47	0.28	0.37	0.08	0.04	0.10
$\sigma_h^h(2W+2b2\tau)$	1.24	1.48	0.21	0.09	0.50	1.29	1.22	0.67
$\sigma_A^h(2W+2b2\tau)$	0.38	0.23	0.48	0.32	0.51	0.21	0.11	0.21
$\sigma_A^A(2W+2b2\tau)$	0.12	0.04	1.08	1.14	0.52	0.04	0.01	0.07
$\sigma_h^A(2W+2b2\tau)$	0.39	0.24	0.48	0.32	0.51	0.22	0.12	0.21
$\sigma_h^h(W+2b2\tau)$	6.93	7.17	2.47	1.64	4.24	6.73	6.24	4.48
$\sigma_A^h(W+2b2\tau)$	2.18	1.14	5.54	5.81	4.38	1.15	0.61	1.42
$\sigma_A^A(W+2b2\tau)$	1.66	0.59	6.00	6.38	4.06	0.87	0.47	1.09
$\sigma_h^A(W+2b2\tau)$	5.25	3.70	2.68	1.80	3.92	5.10	4.78	3.44

**Table 5.** Mass spectra and mixing angles in the Type-I scenario, alongside cross sections (in fb) from different production channels, are shown.

Parameters	BP1	BP2	BP3	BP4	BP5	BP6	BP7	BP8
$m_h$ (GeV)	83.66	83.23	100.04	115.35	95.12	84.84	103.41	86.87
$m_H$ (GeV)	125.00	125.00	125.00	125.00	125.00	125.00	125.00	125.00
$m_A$ (GeV)	113.60	109.52	93.55	79.30	101.38	108.83	90.46	112.97
$m_{H^\pm}$ (GeV)	166.22	169.14	166.18	158.67	169.99	176.64	186.78	195.68
$s_{\beta-\alpha}$	-0.10	-0.13	-0.17	-0.10	-0.13	-0.12	-0.13	-0.12
$\tan\beta$	18.57	14.41	10.51	17.42	13.90	15.37	15.36	14.53
$m_{12}^2$ (GeV <sup>2</sup> )	367.17	408.42	801.13	728.57	645.95	437.53	631.00	456.00
$\sigma_{2t}^h(2W + 2b2\tau)$	2.42	–	2.54	0.18	–	–	–	–
$\sigma_t^h(2W + 2b2\tau)$	3.61	11.33	3.68	0.19	1.77	13.23	1.84	19.07
$\sigma_{2t}^A(2W + 2b2\tau)$	0.08	–	4.74	6.43	–	–	–	–
$\sigma_t^A(2W + 2b2\tau)$	0.11	0.29	6.88	6.72	0.99	0.30	13.35	0.97
$\sigma_h^h(W + 4\tau)$	17.29	48.02	4.48	0.19	6.29	66.34	7.55	80.44
$\sigma_A^h(W + 4\tau)$	0.54	1.25	8.36	6.72	3.51	1.48	54.84	4.07
$\sigma_A^A(W + 4\tau)$	0.38	0.93	9.33	10.68	3.31	1.14	64.36	3.14
$\sigma_h^A(W + 4\tau)$	12.12	35.62	5.00	0.29	5.94	50.98	8.86	62.04

**Table 6.** Mass spectra and mixing angles in the Type-X scenario, alongside cross sections (in fb) from different production channels, are shown.

## 5.2 Benchmark points for Type-X

In table 6, eight BPs for the Type-X scenario are presented. In this scenario, the light neutral Higgs bosons dominantly decay into  $\tau$ 's due to the structure and size of the Yukawa couplings, as seen in table 1. Here, for the advocated new discovery channels, we only tabulate the results for  $W + 4\tau$  final states, since (as previously shown) this is the only signature available. Specifically, notice that the only viable channel, in this Type, is the associated production of a charged Higgs boson with a neutral Higgs,  $h$  or  $A$ , i.e.,  $pp \rightarrow H^\pm h/A$ , which would lead to a  $W^\pm + 4\tau$  final state. For BP1,3 and 4, the condition  $m_{H^\pm} + m_b < m_t$  is satisfied. For BP2,5,6,7 and 8, the condition  $m_{H^\pm} + m_b > m_t$  is satisfied (so that charged Higgs boson production from top (anti)quark decay is forbidden). For BP1, the total production rate of  $2W + 2b2\tau$  from top (anti)quark channels is 3.72 fb and that of  $W + 4\tau$  from di-Higgs production is 30.33 fb, around 6 times larger. Similarly for BP3 and BP4. For BP2, the total production rate of  $2W + 2b2\tau$  from top (anti)quark channels is 11.62 fb and that of  $W + 4\tau$  from the di-Higgs production channels is 85.82 fb. Hence, the  $W + 4\tau$  signature from the aforementioned di-Higgs model is always more sizeable than the  $2W + 2b2\tau$  one from top (anti)quark process dynamics.



## 6 Conclusions

By adding a Higgs doublet to the SM, one can address some fundamental issues of the latter, like the structure of EWSB, the flavour puzzle, the need for additional CP violation, the dynamics of EW baryogenesis, the provision of suitable Dark Matter (DM) candidates and so on. This addition generates a 2HDM, which has been the subject of extensive studies in the literature. Here, we wanted to examine the phenomenology of the 2HDM in regions of its parameter space that can offer us new ways of detecting the charged Higgs boson state emerging in this BSM construct, above and beyond what is already established in such a literature.

Specifically, the experimental teams at the LHC are presently searching for  $H^\pm$  states in 2HDM realisations by assuming these to be produced in association with top (anti)quarks. The advantage of exploiting such production processes is twofold. On the one hand, (anti)top quarks are produced (in pairs) via QCD interactions, which thus have no BSM parameter dependence. On the other hand, being QCD induced, the corresponding cross sections at the LHC are rather large. However, when it comes to the top (anti)quark interacting with the charged Higgs bosons, via a  $H^+\bar{t}b$  vertex, a BSM parameter dependence becomes manifest. This can be ascribed to the different ways the two Higgs doublets of this BSM scenario can give mass to elementary fermions, quarks and leptons, via Yukawa couplings. There are typically four possible choices for doing so, referred to as Types. Here, we have been concerned with two of these, known as Type-I and -X. In these two realisations of the 2HDM, the  $H^\pm$  production and decay rates associated with processes involving top (anti)quarks can be surpassed by those in which charged Higgs bosons are produced via EW interactions which only gauge dependent (i.e., they do not depend on 2HDM parameters), when the  $H^\pm$  states are searched for through their  $H^\pm \rightarrow W^\pm h/A$  decays and the SM-like Higgs boson discovered at the LHC is identified with the heavy CP-even state of the 2HDM,  $H$ , so that the light CP-even one,  $h$ , and CP-odd one,  $A$ , are lighter in comparison.

Specifically, in this work, we have investigated charged Higgs boson production via  $pp \rightarrow H^\pm \phi$  ( $\phi = h, A$ ) and  $pp \rightarrow H^+ H^-$  at the LHC with  $\sqrt{s} = 14$  TeV in the 2HDM Type-I and -X, after taking into account all up-to-date theoretical and experimental constraints. Using these production mechanisms and allowing for  $H^\pm \rightarrow W^\pm h/A$  decays in all possible combinations, we have examined the final states  $2W + 4b$ ,  $2W + 2\tau 2b$ ,  $W + 4b$ ,  $W + 2\tau 2b$  and  $W + 4\tau$  as potential discovery channels. We have thus compared their yield against that of the channels involving top (anti)quarks and shown that the former can be larger than the latter for a large interval of  $m_{H^\pm}$  values, up to 197 GeV or so when  $\tan\beta$  is large. However, the most interesting region of 2HDM parameter space in both Types studied here is the one where  $m_{H^\pm} \lesssim m_t$ , as it gives potentially detectable event rates for most of the aforementioned final states. Thus, in order to enable the ATLAS and CMS Collaborations to confirm or disprove this possibility, we have finally presented a variety of BPs in both 2HDM Type-I and -X amenable to experimental investigation.

## Acknowledgments

The work of AA, RB, MK and BM is supported by the Moroccan Ministry of Higher Education and Scientific Research MESRSFC and CNRST Project PPR/2015/6. The work of SM is supported in part through the NExT Institute and the STFC Consolidated Grant No. ST/L000296/1. Y. W. is supported by the ‘Scientific Research Funding Project for Introduced High-level Talents’ of the Inner Mongolia Normal University Grant No. 2019YJRC001. Q.-S. Yan’s work is supported by the Natural Science Foundation of China Grant No. 11875260.

**Open Access.** This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

## References

- [1] ATLAS collaboration, *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC*, *Phys. Lett. B* **716** (2012) 1 [[arXiv:1207.7214](https://arxiv.org/abs/1207.7214)] [[INSPIRE](#)].
- [2] CMS collaboration, *Observation of a New Boson at a Mass of 125 GeV with the CMS Experiment at the LHC*, *Phys. Lett. B* **716** (2012) 30 [[arXiv:1207.7235](https://arxiv.org/abs/1207.7235)] [[INSPIRE](#)].
- [3] ATLAS collaboration, *Combined measurements of Higgs boson production and decay using up to 80 fb<sup>-1</sup> of proton-proton collision data at  $\sqrt{s} = 13$  TeV collected with the ATLAS experiment*, *Phys. Rev. D* **101** (2020) 012002 [[arXiv:1909.02845](https://arxiv.org/abs/1909.02845)] [[INSPIRE](#)].
- [4] CMS collaboration, *Combined measurements of Higgs boson couplings in proton-proton collisions at  $\sqrt{s} = 13$  TeV*, *Eur. Phys. J. C* **79** (2019) 421 [[arXiv:1809.10733](https://arxiv.org/abs/1809.10733)] [[INSPIRE](#)].
- [5] V.D. Barger, R.J.N. Phillips and D.P. Roy, *Heavy charged Higgs signals at the LHC*, *Phys. Lett. B* **324** (1994) 236 [[hep-ph/9311372](https://arxiv.org/abs/hep-ph/9311372)] [[INSPIRE](#)].
- [6] J.F. Gunion, H.E. Haber, F.E. Paige, W.-K. Tung and S.S.D. Willenbrock, *Neutral and Charged Higgs Detection: Heavy Quark Fusion, Top Quark Mass Dependence and Rare Decays*, *Nucl. Phys. B* **294** (1987) 621 [[INSPIRE](#)].
- [7] R.M. Barnett, H.E. Haber and D.E. Soper, *Ultraheavy Particle Production from Heavy Partons at Hadron Colliders*, *Nucl. Phys. B* **306** (1988) 697 [[INSPIRE](#)].
- [8] J.L. Diaz-Cruz and O.A. Sampayo, *Contribution of gluon fusion to the production of charged Higgs at hadron colliders*, *Phys. Rev. D* **50** (1994) 6820 [[INSPIRE](#)].
- [9] S. Moretti and D.P. Roy, *Detecting heavy charged Higgs bosons at the LHC with triple b tagging*, *Phys. Lett. B* **470** (1999) 209 [[hep-ph/9909435](https://arxiv.org/abs/hep-ph/9909435)] [[INSPIRE](#)].
- [10] D.J. Miller, S. Moretti, D.P. Roy and W.J. Stirling, *Detecting heavy charged Higgs bosons at the CERN LHC with four b quark tags*, *Phys. Rev. D* **61** (2000) 055011 [[hep-ph/9906230](https://arxiv.org/abs/hep-ph/9906230)] [[INSPIRE](#)].
- [11] M. Guchait and S. Moretti, *Improving the discovery potential of charged Higgs bosons at Tevatron run II*, *JHEP* **01** (2002) 001 [[hep-ph/0110020](https://arxiv.org/abs/hep-ph/0110020)] [[INSPIRE](#)].
- [12] J. Alwall, C. Biscarat, S. Moretti, J. Rathsmann and A. Sopczak, *The  $p\bar{p} \rightarrow tbH^\pm$  process at the Tevatron in HERWIG and PYTHIA simulations*, *Eur. Phys. J. C* **39S1** (2005) 37 [[hep-ph/0312301](https://arxiv.org/abs/hep-ph/0312301)] [[INSPIRE](#)].

- [13] J. Alwall and J. Rathsmann, *Improved description of charged Higgs boson production at hadron colliders*, *JHEP* **12** (2004) 050 [[hep-ph/0409094](#)] [[INSPIRE](#)].
- [14] A.A. Barrientos Bendezu and B.A. Kniehl,  *$W^\pm H^\mp$  associated production at the large hadron collider*, *Phys. Rev. D* **59** (1999) 015009 [[hep-ph/9807480](#)] [[INSPIRE](#)].
- [15] S. Moretti and K. Odagiri, *The Phenomenology of  $W^\pm H^\mp$  production at the large hadron collider*, *Phys. Rev. D* **59** (1999) 055008 [[hep-ph/9809244](#)] [[INSPIRE](#)].
- [16] A.A. Barrientos Bendezu and B.A. Kniehl, *Quark loop amplitudes for  $W^\pm H^\mp$  associated hadroproduction*, *Phys. Rev. D* **61** (2000) 097701 [[hep-ph/9909502](#)] [[INSPIRE](#)].
- [17] S. Moretti and K. Odagiri, *Production of charged Higgs bosons of the minimal supersymmetric standard model in  $b$  quark initiated processes at the large hadron collider*, *Phys. Rev. D* **55** (1997) 5627 [[hep-ph/9611374](#)] [[INSPIRE](#)].
- [18] A. Arhrib, K. Cheung, J.S. Lee and C.-T. Lu, *Enhanced charged Higgs production through  $W$ -Higgs fusion in  $W$ - $b$  scattering*, *JHEP* **05** (2016) 093 [[arXiv:1509.00978](#)] [[INSPIRE](#)].
- [19] J. Hernandez-Sanchez, S. Moretti, R. Noriega-Papaqui and A. Rosado, *Off-diagonal terms in Yukawa textures of the Type-III 2-Higgs doublet model and light charged Higgs boson phenomenology*, *JHEP* **07** (2013) 044 [[arXiv:1212.6818](#)] [[INSPIRE](#)].
- [20] J. Hernández-Sánchez, C.G. Honorato, S. Moretti and S. Rosado-Navarro, *Charged Higgs boson production via  $cb$ -fusion at the Large Hadron Collider*, *Phys. Rev. D* **102** (2020) 055008 [[arXiv:2003.06263](#)] [[INSPIRE](#)].
- [21] S. Dittmaier, G. Hiller, T. Plehn and M. Spannowsky, *Charged-Higgs Collider Signals with or without Flavor*, *Phys. Rev. D* **77** (2008) 115001 [[arXiv:0708.0940](#)] [[INSPIRE](#)].
- [22] S. Kanemura and C.P. Yuan, *Testing supersymmetry in the associated production of  $CP$  odd and charged Higgs bosons*, *Phys. Lett. B* **530** (2002) 188 [[hep-ph/0112165](#)] [[INSPIRE](#)].
- [23] R. Enberg, W. Klemm, S. Moretti, S. Munir and G. Wouda, *Charged Higgs boson in the  $W^\pm$  Higgs channel at the Large Hadron Collider*, *Nucl. Phys. B* **893** (2015) 420 [[arXiv:1412.5814](#)] [[INSPIRE](#)].
- [24] R. Enberg, W. Klemm, S. Moretti, S. Munir and G. Wouda,  *$H^\pm$  in the  $W^\pm h$  channel at the LHC Run 2*, in *2nd Toyama International Workshop on Higgs as a Probe of New Physics*, 6, 2015 [[arXiv:1506.04409](#)] [[INSPIRE](#)].
- [25] R. Enberg, W. Klemm, S. Moretti and S. Munir, *Electroweak production of Higgs boson pairs in 2HDMs*, *PoS ICHEP2016* (2017) 1174 [[arXiv:1704.06405](#)] [[INSPIRE](#)].
- [26] R. Enberg, W. Klemm, S. Moretti and S. Munir, *Signatures of the Type-I 2HDM at the LHC*, *PoS CORFU2018* (2018) 013 [[arXiv:1812.08623](#)] [[INSPIRE](#)].
- [27] R. Enberg, W. Klemm, S. Moretti and S. Munir, *Electroweak production of multiple (pseudo)scalars in the 2HDM*, *Eur. Phys. J. C* **79** (2019) 512 [[arXiv:1812.01147](#)] [[INSPIRE](#)].
- [28] S. Moretti, *Pair production of charged Higgs scalars from electroweak gauge boson fusion*, *J. Phys. G* **28** (2002) 2567 [[hep-ph/0102116](#)] [[INSPIRE](#)].
- [29] S. Moretti and J. Rathsmann, *Pair production of charged Higgs bosons in association with bottom quark pairs at the Large Hadron Collider*, *Eur. Phys. J. C* **33** (2004) 41 [[hep-ph/0308215](#)] [[INSPIRE](#)].
- [30] M. Aoki, R. Guedes, S. Kanemura, S. Moretti, R. Santos and K. Yagyu, *Light Charged Higgs bosons at the LHC in 2HDMs*, *Phys. Rev. D* **84** (2011) 055028 [[arXiv:1104.3178](#)] [[INSPIRE](#)].
- [31] A.G. Akeroyd et al., *Prospects for charged Higgs searches at the LHC*, *Eur. Phys. J. C* **77** (2017) 276 [[arXiv:1607.01320](#)] [[INSPIRE](#)].

- [32] A. Arhrib, R. Benbrik, H. Harouiz, S. Moretti and A. Rouchad, *A Guidebook to Hunting Charged Higgs Bosons at the LHC*, [arXiv:1810.09106](#) [INSPIRE].
- [33] B. Coleppa, A. Sarkar and S.K. Rai, *Charged Higgs boson discovery prospects*, *Phys. Rev. D* **101** (2020) 055030 [[arXiv:1909.11992](#)] [INSPIRE].
- [34] A. Arhrib, R. Benbrik and S. Moretti, *Bosonic Decays of Charged Higgs Bosons in a 2HDM Type-I*, *Eur. Phys. J. C* **77** (2017) 621 [[arXiv:1607.02402](#)] [INSPIRE].
- [35] A.G. Akeroyd, *Three body decays of Higgs bosons at LEP-2 and application to a hidden fermiophobic Higgs*, *Nucl. Phys. B* **544** (1999) 557 [[hep-ph/9806337](#)] [INSPIRE].
- [36] D0 collaboration, *Combination of  $t\bar{t}$  cross section measurements and constraints on the mass of the top quark and its decays into charged Higgs bosons*, *Phys. Rev. D* **80** (2009) 071102 [[arXiv:0903.5525](#)] [INSPIRE].
- [37] CDF collaboration, *Search for charged Higgs bosons from top quark decays in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$ -TeV*, *Phys. Rev. Lett.* **96** (2006) 042003 [[hep-ex/0510065](#)] [INSPIRE].
- [38] ALEPH, DELPHI, L3, OPAL and LEP collaborations, *Search for Charged Higgs bosons: Combined Results Using LEP Data*, *Eur. Phys. J. C* **73** (2013) 2463 [[arXiv:1301.6065](#)] [INSPIRE].
- [39] ATLAS collaboration, *Search for charged Higgs bosons decaying via  $H^\pm \rightarrow \tau^\pm \nu$  in fully hadronic final states using  $pp$  collision data at  $\sqrt{s} = 8$  TeV with the ATLAS detector*, *JHEP* **03** (2015) 088 [[arXiv:1412.6663](#)] [INSPIRE].
- [40] CMS collaboration, *Search for a charged Higgs boson in  $pp$  collisions at  $\sqrt{s} = 8$  TeV*, *JHEP* **11** (2015) 018 [[arXiv:1508.07774](#)] [INSPIRE].
- [41] CMS collaboration, *Search for charged Higgs bosons with the  $H^\pm \rightarrow \tau^\pm \nu_\tau$  decay channel in the fully hadronic final state at  $\sqrt{s} = 13$  TeV*, *CMS-PAS-HIG-16-031* (2016).
- [42] ATLAS collaboration, *Search for charged Higgs bosons decaying via  $H^\pm \rightarrow \tau^\pm \nu_\tau$  in the  $\tau$ +jets and  $\tau$ +lepton final states with  $36\text{fb}^{-1}$  of  $pp$  collision data recorded at  $\sqrt{s} = 13$  TeV with the ATLAS experiment*, *JHEP* **09** (2018) 139 [[arXiv:1807.07915](#)] [INSPIRE].
- [43] ATLAS collaboration, *A Search for a light charged Higgs boson decaying to  $cs^-$  in  $pp$  collisions at  $\sqrt{s} = 7$  TeV with the ATLAS detector*, *ATLAS-CONF-2011-094* (2011).
- [44] CMS collaboration, *Search for a light charged Higgs boson in the  $H^\pm \rightarrow cs$  channel in proton-proton collisions at  $\sqrt{s} = 13$  TeV*, *Phys. Rev. D* **102** (2020) 072001 [[arXiv:2005.08900](#)] [INSPIRE].
- [45] PARTICLE DATA GROUP collaboration, *Review of Particle Physics*, *Phys. Rev. D* **98** (2018) 030001.
- [46] CMS collaboration, *Search for charged Higgs bosons in the  $H^\pm \rightarrow \tau^\pm \nu_\tau$  decay channel in proton-proton collisions at  $\sqrt{s} = 13$  TeV*, *JHEP* **07** (2019) 142 [[arXiv:1903.04560](#)] [INSPIRE].
- [47] G.C. Branco, P.M. Ferreira, L. Lavoura, M.N. Rebelo, M. Sher and J.P. Silva, *Theory and phenomenology of two-Higgs-doublet models*, *Phys. Rept.* **516** (2012) 1 [[arXiv:1106.0034](#)] [INSPIRE].
- [48] S.L. Glashow and S. Weinberg, *Natural Conservation Laws for Neutral Currents*, *Phys. Rev. D* **15** (1977) 1958 [INSPIRE].
- [49] N.G. Deshpande and E. Ma, *Pattern of Symmetry Breaking with Two Higgs Doublets*, *Phys. Rev. D* **18** (1978) 2574 [INSPIRE].

- [50] S. Kanemura, T. Kubota and E. Takasugi, *Lee-Quigg-Thacker bounds for Higgs boson masses in a two doublet model*, *Phys. Lett. B* **313** (1993) 155 [[hep-ph/9303263](#)] [[INSPIRE](#)].
- [51] A.G. Akeroyd, A. Arhrib and E.-M. Naimi, *Note on tree level unitarity in the general two Higgs doublet model*, *Phys. Lett. B* **490** (2000) 119 [[hep-ph/0006035](#)] [[INSPIRE](#)].
- [52] M.E. Peskin and T. Takeuchi, *Estimation of oblique electroweak corrections*, *Phys. Rev. D* **46** (1992) 381 [[INSPIRE](#)].
- [53] H.-J. He, N. Polonsky and S.-f. Su, *Extra families, Higgs spectrum and oblique corrections*, *Phys. Rev. D* **64** (2001) 053004 [[hep-ph/0102144](#)] [[INSPIRE](#)].
- [54] W. Grimus, L. Lavoura, O.M. Ogreid and P. Osland, *The Oblique parameters in multi-Higgs-doublet models*, *Nucl. Phys. B* **801** (2008) 81 [[arXiv:0802.4353](#)] [[INSPIRE](#)].
- [55] H.E. Haber and D. O’Neil, *Basis-independent methods for the two-Higgs-doublet model III: The CP-conserving limit, custodial symmetry, and the oblique parameters S, T, U*, *Phys. Rev. D* **83** (2011) 055017 [[arXiv:1011.6188](#)] [[INSPIRE](#)].
- [56] P. Bechtle et al., *HiggsBounds-5: Testing Higgs Sectors in the LHC 13 TeV Era*, *Eur. Phys. J. C* **80** (2020) 1211 [[arXiv:2006.06007](#)] [[INSPIRE](#)].
- [57] P. Bechtle, S. Heinemeyer, T. Klingl, T. Stefaniak, G. Weiglein and J. Wittbrodt, *HiggsSignals-2: Probing new physics with precision Higgs measurements in the LHC 13 TeV era*, *Eur. Phys. J. C* **81** (2021) 145 [[arXiv:2012.09197](#)] [[INSPIRE](#)].
- [58] F. Mahmoudi, *SuperIso v2.3: A Program for calculating flavor physics observables in Supersymmetry*, *Comput. Phys. Commun.* **180** (2009) 1579 [[arXiv:0808.3144](#)] [[INSPIRE](#)].
- [59] HFLAV collaboration, *Averages of b-hadron, c-hadron, and  $\tau$ -lepton properties as of summer 2016*, *Eur. Phys. J. C* **77** (2017) 895 [[arXiv:1612.07233](#)] [[INSPIRE](#)].
- [60] LHCb collaboration, *Measurement of the  $B_s^0 \rightarrow \mu^+ \mu^-$  branching fraction and effective lifetime and search for  $B^0 \rightarrow \mu^+ \mu^-$  decays*, *Phys. Rev. Lett.* **118** (2017) 191801 [[arXiv:1703.05747](#)] [[INSPIRE](#)].
- [61] D. Eriksson, J. Rathsman and O. Stal, *2HDMC: Two-Higgs-Doublet Model Calculator Physics and Manual*, *Comput. Phys. Commun.* **181** (2010) 189 [[arXiv:0902.0851](#)] [[INSPIRE](#)].
- [62] S.-h. Zhu, *Complete next-to-leading order QCD corrections to charged Higgs boson associated production with top quark at the CERN large hadron collider*, *Phys. Rev. D* **67** (2003) 075006 [[hep-ph/0112109](#)] [[INSPIRE](#)].
- [63] T. Plehn, *Charged Higgs boson production in bottom gluon fusion*, *Phys. Rev. D* **67** (2003) 014018 [[hep-ph/0206121](#)] [[INSPIRE](#)].
- [64] E.L. Berger, T. Han, J. Jiang and T. Plehn, *Associated production of a top quark and a charged Higgs boson*, *Phys. Rev. D* **71** (2005) 115012 [[hep-ph/0312286](#)] [[INSPIRE](#)].
- [65] K.A. Assamagan, M. Guchait and S. Moretti, *Charged Higgs bosons in the transition region  $M_{H^\pm} \sim m_t$  at the LHC*, in *3rd Les Houches Workshop on Physics at TeV Colliders*, Les Houches France (2004) [[hep-ph/0402057](#)] [[INSPIRE](#)].
- [66] T. Stelzer and W.F. Long, *Automatic generation of tree level helicity amplitudes*, *Comput. Phys. Commun.* **81** (1994) 357 [[hep-ph/9401258](#)] [[INSPIRE](#)].
- [67] H. Plathow-Besch, *PDFLIB: A Library of all available parton density functions of the nucleon, the pion and the photon and the corresponding  $\alpha_s$  calculations*, *Comput. Phys. Commun.* **75** (1993) 396 [[INSPIRE](#)].