

Isospin-breaking effects in the two-pion contribution to hadronic vacuum polarization

Gilberto Colangelo,^a Martin Hoferichter,^a Bastian Kubis^b and Peter Stoffer^{c,d}

^aAlbert Einstein Center for Fundamental Physics, Institute for Theoretical Physics, University of Bern, Sidlerstrasse 5, 3012 Bern, Switzerland

^bHelmholtz-Institut für Strahlen- und Kernphysik (Theorie) and Bethe Center for Theoretical Physics, Universität Bonn, 53115 Bonn, Germany

^cPhysik-Institut, Universität Zürich, Winterthurerstrasse 190, 8057 Zürich, Switzerland

^dPaul Scherrer Institut, 5232 Villigen PSI, Switzerland

E-mail: gilberto@itp.unibe.ch, hoferichter@itp.unibe.ch, kubis@hiskp.uni-bonn.de, stoffer@physik.uzh.ch

ABSTRACT: Isospin-breaking (IB) effects in the two-pion contribution to hadronic vacuum polarization (HVP) can be resonantly enhanced, if related to the interference of the $\rho(770)$ and $\omega(782)$ resonances. This particular IB contribution to the pion vector form factor and thus the line shape in $e^+e^- \rightarrow \pi^+\pi^-$ can be described by the residue at the ω pole — the ρ - ω mixing parameter ϵ_ω . Here, we argue that while in general analyticity requires this parameter to be real, the radiative channels $\pi^0\gamma$, $\pi\pi\gamma$, $\eta\gamma$ can induce a small phase, whose size we estimate as $\delta_\epsilon = 3.5(1.0)^\circ$ by using a narrow-width approximation for the intermediate-state vector mesons. We then perform fits to the $e^+e^- \rightarrow \pi^+\pi^-$ data base and study the consequences for the two-pion HVP contribution to the anomalous magnetic moment of the muon, its IB part due to ρ - ω mixing, and the mass of the ω resonance. We find that the global fit does prefer a non-vanishing value of $\delta_\epsilon = 4.5(1.2)^\circ$, close to the narrow-resonance expectation, but with a large spread among the data sets, indicating systematic differences in the ρ - ω region.

KEYWORDS: Chiral Lagrangian, Precision QED

ARXIV EPRINT: [2208.08993](https://arxiv.org/abs/2208.08993)

Contents

1	Introduction	1
2	Radiative channels and phase in the ρ-ω mixing parameter	2
2.1	Dispersive representation	2
2.2	Radiative channels	4
3	Fits to $e^+e^- \rightarrow 2\pi$ data	6
4	Isospin-breaking contribution to a_μ from ρ-ω mixing	8
5	Consequences for the ω mass	10
6	Conclusions	11

1 Introduction

The two-pion channel gives the dominant contribution to hadronic vacuum polarization (HVP) in the low-energy region most relevant for the anomalous magnetic moment of the muon [1], adding about 70% of the total leading-order (LO) effect [2–7]

$$a_\mu^{\text{HVP, LO}}|_{e^+e^-} = 693.1(4.0) \times 10^{-10}. \quad (1.1)$$

Its contribution needs to be understood at a level of at least 0.3% to match the final precision expected from the Fermilab E989 experiment [8]. The nominal combined sensitivity of the 2π data sets entering eq. (1.1) — from SND [9, 10], CMD-2 [11–14], BESIII [15], CLEO [16], and dominated by the precision data sets from BaBar [17, 18] and KLOE [19–22] — does reach 0.4%, but becomes diluted due to a tension between BaBar and KLOE, inflating the 2π uncertainty included in eq. (1.1) to 0.7%. In dispersive approaches [4–6, 23, 24] also space-like data [25, 26] can be used, and while stabilizing the extrapolation to the space-like region, their impact on the time-like HVP integral is minor. More recently, new data from SND [27] have become available, lying in between BaBar and KLOE, but not at a comparable level of precision that would allow one to resolve the tension. Such new precision measurements are expected from CMD-3 [28], BaBar [29], BESIII [30], and Belle II [31] in the future.

Improved understanding of the 2π channel has further become critical to address the emerging tension between lattice QCD [32–35] and e^+e^- data at least for the intermediate window quantity [36], with immediate consequences for the current 4.2σ discrepancy for the anomalous magnetic moment of the muon between experiment [37–41] and the prediction in the Standard Model [1–7, 42–59] when the HVP contribution is derived from

$e^+e^- \rightarrow$ hadrons cross-section data. While the detailed comparison to lattice QCD as well as related observables defines an important path forward [60–67], so does renewed scrutiny of the data-driven approach.

For the 2π channel, new precision data sets constitute the clear first priority, but another aspect concerns the role of radiative corrections [29, 68], in particular, the question in which cases the use of a point-like approximation [69–73] for the pion might miss relevant effects, as recently observed in the forward-backward asymmetry [74, 75], and currently under study for the C -even contributions [76]. In this work, we study a complementary point, i.e., not isospin-breaking (IB) effects that manifest themselves as final- or initial-state radiation, but corrections that are typically absorbed into the pion vector form factor (VFF) itself.¹ The most prominent such correction arises from ρ - ω mixing. From a dispersive point of view the fact that the $\omega(782)$ resonance is so narrow allows one to describe this interference in terms of a single real parameter: the ρ - ω mixing parameter ϵ_ω . But given the extraordinary precision requirements for the 2π channel together with the resonance enhancement in the ρ - ω region, even higher-order effects may affect the value of this parameter and generate non-negligible effects in $a_\mu^{\text{HVP, LO}}$. Most notably, the radiative channels $\pi^0\gamma$, $\pi\pi\gamma$, $\eta\gamma$, all of which couple to both ρ and ω , can induce imaginary parts in the mixing and thereby an effective small phase δ_ϵ in the parameter ϵ_ω to which e^+e^- data might be sensitive.

To derive the phenomenological consequences of this phase we first generalize the dispersive representation of the pion VFF from ref. [4] and estimate its size based on a narrow-resonance approach, see section 2. We then perform fits, to individual data sets and globally, allowing for a free phase δ_ϵ , to assess consistency both among the data sets and with the narrow-resonance expectation, see section 3. Consequences for the IB contribution to a_μ due to ρ - ω mixing and the ω mass are discussed in sections 4 and 5, respectively, before concluding in section 6.

2 Radiative channels and phase in the ρ - ω mixing parameter

2.1 Dispersive representation

Dispersive representations for the pion VFF, $F_\pi^V(s)$, that link the matrix element of the electromagnetic current $j_{\text{em}}^\mu = (2\bar{u}\gamma^\mu u - \bar{d}\gamma^\mu d - \bar{s}\gamma^\mu s)/3$,

$$\langle \pi^\pm(p') | j_{\text{em}}^\mu(0) | \pi^\pm(p) \rangle = \pm(p' + p)^\mu F_\pi^V((p' - p)^2), \quad (2.1)$$

to $\pi\pi$ scattering have been used for a long time in the literature [4, 6, 23, 82–91], not only for the HVP application, but also for hadronic light-by-light scattering (HLbL), where the extrapolation into the space-like region enters [50, 51, 92–95].

Here, we build upon the representation from ref. [4] (in turn based on refs. [83, 84]), whose main features can be summarized as follows: the VFF is decomposed into three

¹Such corrections were studied before in the context of relating VFF measurements in $e^+e^- \rightarrow \pi^+\pi^-$ to $\tau^\pm \rightarrow \pi^\pm\pi^0\nu_\tau$ data [77–81]. Here, we aim instead at a rigorous implementation of ρ - ω mixing in a dispersive framework, both to quantify its impact on $a_\mu^{\text{HVP, LO}}$ and as another powerful consistency check on the e^+e^- data base.

factors

$$F_\pi^V(s) = \Omega_1^1(s) G_\omega(s) G_{\text{in}}^N(s), \quad (2.2)$$

corresponding to 2π , 3π , and higher intermediate states, respectively. The Omnès function [96]

$$\Omega_1^1(s) = \exp \left\{ \frac{s}{\pi} \int_{4M_\pi^2}^{\infty} ds' \frac{\delta_1^1(s')}{s'(s'-s)} \right\} \quad (2.3)$$

implements 2π singularities in terms of the isospin $I = 1$ elastic $\pi\pi$ phase shift $\delta_1^1(s)$ in the isospin limit. The phase shift is further constrained by $\pi\pi$ Roy equations [97], which are solved with the phase shifts at $s_0 = (0.8 \text{ GeV})^2$ and $s_1 = (1.15 \text{ GeV})^2$ as free parameters [98, 99]. Systematic errors from the asymptotic continuation of δ_1^1 beyond s_1 are treated as described in ref. [4].

The focus of this work is the second factor, G_ω , which takes into account the effect of 3π intermediate states. Here, the parameterization from ref. [4] reads

$$G_\omega(s) = 1 + \frac{s}{\pi} \int_{9M_\pi^2}^{\infty} ds' \frac{\text{Im } g_\omega(s')}{s'(s'-s)} \left(\frac{1 - \frac{9M_\pi^2}{s'}}{1 - \frac{9M_\pi^2}{s}} \right)^4, \quad (2.4)$$

with

$$g_\omega(s) = 1 + \epsilon_\omega \frac{s}{(M_\omega - \frac{i}{2}\Gamma_\omega)^2 - s}. \quad (2.5)$$

The dispersive reconstruction in eq. (2.4) ensures both the absence of unphysical imaginary parts below $s = 9M_\pi^2$ (for real ϵ_ω) and the correct threshold behavior above [83]. In this formulation, ϵ_ω is an effective parameter tightly related to the residue at the ω pole. The latter, however, is complex in general, but its phase is expected to be tiny: $\delta_\epsilon \simeq \arctan \Gamma_\omega/M_\omega \simeq 0.6^\circ$ arising from the analytic continuation from the real axis to the pole position in the complex plane. Such a small difference is of no concern and if we take it as a measure of the systematic uncertainty in the phase of ϵ_ω , allows us to view the latter as the residue at the ω pole. With the threshold behavior and the pole parameters determined, the resulting $G_\omega(s)$ is then largely insensitive to the parameterization of $g_\omega(s)$, e.g., the numerator could be taken to a constant without any relevant changes to the fit outcome. The main observation in this paper is that the assumption of a real ϵ_ω no longer holds if further IB effects due to radiative channels, $X = \pi^0\gamma, \pi\pi\gamma, \eta\gamma, \dots$, are considered, and these imaginary parts, despite being small, can alter the fit parameters in a significant way, as only the modulus $|F_\pi^V(s)|^2$ is probed by the fit to the cross-section data.

Finally, for the inelastic channels we continue to use a conformal polynomial, whose phase is constrained by the Eidelman–Łukaszuk bound [100, 101]. Its threshold is chosen as the $\omega\pi^0$ threshold, below which inelasticities are negligibly small. After removing the S -wave cusp, $G_{\text{in}}^N(s)$ involves $N - 1$ free parameters, which together with ϵ_ω and $\delta_1^1(s_0)$, $\delta_1^1(s_1)$ are to be constrained in the fit. The fit range is restricted to $s \leq 1 \text{ GeV}^2$, going beyond would require including the effects of ρ' , ρ'' resonances along the lines of refs. [86, 102], an extension left for future work.

2.2 Radiative channels

Both ρ and ω possess non-negligible branching fractions into radiative channels, of which $\pi^0\gamma$ yields the largest contribution. The corresponding imaginary part in the pion VFF can be expressed as

$$\text{Im } F_\pi^V(s)|_{\pi^0\gamma} = \frac{\alpha(s - M_{\pi^0}^2)^3}{48s} F_{\pi^0\gamma^*\gamma^*}(s, 0) (f_1(s))^*, \quad (2.6)$$

where $F_{\pi^0\gamma^*\gamma^*}$ is the pion transition form factor normalized according to

$$F_{\pi^0\gamma^*\gamma^*}(0, 0) = F_{\pi\gamma\gamma} = \sqrt{\frac{4\Gamma[\pi^0 \rightarrow \gamma\gamma]}{\pi\alpha^2 M_{\pi^0}^3}} \quad (2.7)$$

and f_1 denotes the P -wave projection of the $\gamma\pi \rightarrow \pi\pi$ amplitude, see refs. [103–107] for detailed discussions of these amplitudes. To map this imaginary part onto $\text{Im } \epsilon_\omega$, we first write the full pion VFF in the approximation

$$F_\pi^V(s) = \left(1 + \epsilon_\omega \frac{s}{M_\omega^2 - s - i\epsilon}\right) \Omega_1^1(s), \quad (2.8)$$

where we have neglected inelastic corrections for the time being to focus on the interplay of ρ and ω resonances, with the ω approximated in the narrow-width limit for simplicity. As a first step, we show that an imaginary part in ϵ_ω in this form is actually compatible with unitarity. Applying Cutkosky rules to eq. (2.8), we have

$$\begin{aligned} \frac{1}{2i} \text{disc } F_\pi^V(s)|_{2\pi} &= \left(1 + \epsilon_\omega \frac{s}{M_\omega^2 - s - i\epsilon}\right) \text{Im } \Omega_1^1(s), \\ \frac{1}{2i} \text{disc } F_\pi^V(s)|_{3\pi} &= \epsilon_\omega^* s \pi \delta(s - M_\omega^2) (\Omega_1^1(s))^*, \\ \frac{1}{2i} \text{disc } F_\pi^V(s)|_{\pi^0\gamma} &= \text{Im } \epsilon_\omega \frac{s}{M_\omega^2 - s - i\epsilon} (\Omega_1^1(s))^*. \end{aligned} \quad (2.9)$$

These equations are only consistent as long as the sum of these discontinuities is purely imaginary. Collecting all terms, this consistency check is indeed satisfied,

$$\begin{aligned} &\text{Im} \left[\frac{1}{2i} \text{disc } F_\pi^V(s)|_{2\pi} + \frac{1}{2i} \text{disc } F_\pi^V(s)|_{3\pi} + \frac{1}{2i} \text{disc } F_\pi^V(s)|_{\pi^0\gamma} \right] \\ &= \text{Im } \Omega_1^1(s) \left(\text{Im } \epsilon_\omega \frac{s}{M_\omega^2 - s} + \text{Re } \epsilon_\omega s \pi \delta(s - M_\omega^2) \right) \\ &\quad + s \pi \delta(s - M_\omega^2) \left(-\text{Re } \epsilon_\omega \text{Im } \Omega_1^1(s) - \text{Re } \Omega_1^1(s) \text{Im } \epsilon_\omega \right) \\ &\quad + \text{Im } \epsilon_\omega \left(-\frac{s}{M_\omega^2 - s} \text{Im } \Omega_1^1(s) + \text{Re } \Omega_1^1(s) s \pi \delta(s - M_\omega^2) \right) \\ &= 0, \end{aligned} \quad (2.10)$$

so that as long as imaginary parts are avoided below the respective thresholds a phase in ϵ_ω is indeed possible.

Next, the comparison of eqs. (2.6) and (2.9) points towards a strategy for a practical implementation, with $\Omega_1^1(s)$ corresponding to $f_1(s)$, and the ω propagator to the ω contribution in $F_{\pi^0\gamma^*\gamma^*}(s, 0)$. The latter is given by

$$F_{\pi^0\gamma^*\gamma^*}(s, 0) \simeq \frac{g_{\omega\pi\gamma}}{g_{\omega\gamma}} \frac{M_\omega^2}{M_\omega^2 - s - i\epsilon} \quad (2.11)$$

(see, e.g., ref. [108]), while the former can be approximated by [106]

$$f_1(s) \simeq \frac{2g_{\rho\pi\gamma}g_{\rho\pi\pi}}{M_\rho^2 - s - iM_\rho\Gamma_\rho}, \quad \Omega_1^1(s) \simeq \frac{M_\rho^2}{M_\rho^2 - s - iM_\rho\Gamma_\rho}, \quad (2.12)$$

leading to

$$\text{Im } \epsilon_\omega \simeq \frac{\alpha(s - M_{\pi^0}^2)^3}{24s} \frac{g_{\omega\pi\gamma}g_{\rho\pi\gamma}g_{\rho\pi\pi}}{g_{\omega\gamma}M_\rho^2}. \quad (2.13)$$

Inserting the expressions for the radiative decay widths,

$$\Gamma[V \rightarrow \pi^0\gamma] = \frac{\alpha(M_V^2 - M_{\pi^0}^2)^3}{24M_V^3} |g_{V\pi\gamma}|^2, \quad V = \omega, \rho, \quad (2.14)$$

as well as the VMD predictions $g_{\rho\pi\pi} = g_{\rho\gamma} = g_{\omega\gamma}/3$, and evaluating eq. (2.13) at $s = M_V^2 \simeq M_\rho^2 \simeq M_\omega^2$, we find

$$\text{Im } \epsilon_\omega \simeq \frac{\sqrt{\Gamma[\omega \rightarrow \pi^0\gamma]\Gamma[\rho \rightarrow \pi^0\gamma]}}{3M_V}. \quad (2.15)$$

In fact, in the narrow-width limit the same relation can be established for an arbitrary intermediate state X , leading to estimates for the phases around 2.8° ($\pi^0\gamma$), 0.2° ($\eta\gamma$), and 0.02° ($\pi^0\pi^0\gamma$) when using the averages from ref. [109] for branching fractions and masses. For the charged channel $\pi^+\pi^-\gamma$ one needs to take into account the fact that the presence of the Born-term contribution leads to an infrared divergence, in such a way that branching fractions are typically quoted with a cut $E_\gamma = 50$ MeV in the photon energy [110, 111]. However, combined with virtual corrections calculated in a scalar-QED approximation one can define an infrared-safe decay width as

$$\Gamma[V \rightarrow \pi^+\pi^-\gamma] = \Gamma[V \rightarrow \pi^+\pi^-] \frac{\alpha}{\pi} \eta(M_V^2), \quad (2.16)$$

where explicit expressions for the function η can be found in refs. [69–72]. This procedure gives an estimate of 0.4° for the $\pi^+\pi^-\gamma$ channel, subject to minor corrections from non-Born contributions [111]. As for the relative signs, VMD arguments show that the $\pi^0\gamma$ and $\eta\gamma$ channels enter with the same sign (in the standard phase conventions, both intermediate states couple with the same sign to ρ and ω), and Born-term dominance for $\rho, \omega \rightarrow \pi^+\pi^-\gamma$, as well as the positive sign of $\text{Re } \epsilon_\omega$, suggest that its contribution should also add to the other two. Taking further potential corrections due to the analytic continuation to the ω pole as the uncertainty, we conclude that the range $\delta_\epsilon = 3.5(1.0)^\circ$ should give a realistic estimate of the phase in ϵ_ω that can be expected.

Finally, the discussion of the $\pi^0\gamma$ channel suggests that the dominant threshold can be reproduced by implementing the imaginary part in ϵ_ω via

$$G_\omega(s) = 1 + \frac{s}{\pi} \int_{9M_\pi^2}^\infty ds' \frac{\text{Re } \epsilon_\omega}{s'(s' - s)} \text{Im} \left[\frac{s'}{(M_\omega - \frac{i}{2}\Gamma_\omega)^2 - s'} \right] \left(\frac{1 - \frac{9M_\pi^2}{s'}}{1 - \frac{9M_\pi^2}{M_\omega^2}} \right)^4 + \frac{s}{\pi} \int_{M_{\pi^0}^2}^\infty ds' \frac{\text{Im } \epsilon_\omega}{s'(s' - s)} \text{Re} \left[\frac{s'}{(M_\omega - \frac{i}{2}\Gamma_\omega)^2 - s'} \right] \left(\frac{1 - \frac{M_{\pi^0}^2}{s'}}{1 - \frac{M_{\pi^0}^2}{M_\omega^2}} \right)^3, \quad (2.17)$$

but we also checked that the fit results are largely insensitive to the details of the implementation, such as the inclusion of the explicit $\pi^0\gamma$ threshold in the unphysical region of the pion VFF. In fact, even replacing $G_\omega(s)$ with $g_\omega(s)$ only leads to small changes as long as the imaginary part in ϵ_ω is kept, in line with the expectation that it is solely the resonance enhancement that makes these higher-order effects relevant.

3 Fits to $e^+e^- \rightarrow 2\pi$ data

To gauge the impact of a possible phase in ϵ_ω on the HVP contribution to a_μ , we generalize the global fits from ref. [4], including a free imaginary part via the prescription (2.17), and express our results in terms of $\text{Re } \epsilon_\omega$ and δ_ϵ . In particular, we now include the BESIII data [15] and the SND measurement [27], which became available after ref. [4].² The results for the fits are shown in table 1 (single experiments) and table 2 (combinations), in terms of the most relevant parameters: goodness of fit, the ω mass, real part and phase of ϵ_ω , and the contribution to a_μ . In table 3, we also provide the decomposition into the Euclidean windows from ref. [36].

In most cases, we observe a moderate improvement when a non-vanishing phase is admitted, the main exception being the SND20 data, which we cannot describe with our dispersive representation otherwise. Accordingly, in this case the resulting phase comes out around 10° and thus much larger than can be justified via radiative intermediate states. A similarly large phase is also found for the previous energy-scan experiments SND06 and CMD-2, but in these cases good fits can still be found when imposing a realistic size of δ_ϵ . Even if a large phase is admitted in the fit to the SND20 data, the fit quality remains rather poor.³ As long as the reason for this behavior, which might point towards underestimated systematic effects, is not understood, we will therefore take the global fit to all experiments apart from SND20 as our new central result, i.e.

$$\begin{aligned} \text{Re } \epsilon_\omega &= 1.97(3) \times 10^{-3}, & \delta_\epsilon &= 4.5(1.2)^\circ, \\ a_\mu^{\pi\pi}|_{\leq 1 \text{ GeV}} &= 494.6(2.3) \times 10^{-10}, & M_\omega &= 782.09(12) \text{ MeV}. \end{aligned} \quad (3.1)$$

²In the case of ref. [15] the corrected covariance matrix was critical for the inclusion of this data set in a statistically meaningful way. For the uncertainty of the energy calibration at the ρ peak we use $\Delta E = 0.6 \text{ MeV}$ [112] and $\Delta E = 0.26 \text{ MeV}$ [113], respectively.

³The fit presented in ref. [27] in terms of a sum of Breit-Wigner functions for $V = \rho, \omega, \rho'$ displays a slightly better fit quality, $\chi^2/\text{dof} = 47/30 = 1.57$, with p -value of 2.5%, but such a representation cannot be reconciled with the analytic properties of the pion VFF.

	χ^2/dof	p -value	M_ω [MeV]	$10^3 \times \text{Re } \epsilon_\omega$	δ_ϵ [°]	$10^{10} \times a_\mu^{\pi\pi} _{\leq 1 \text{ GeV}}$
SND06	1.40	5.3%	781.49(32)(2)	2.03(5)(2)		499.7(6.9)(4.1)
	1.08	35%	782.11(32)(2)	1.98(4)(2)	8.5(2.3)(0.3)	497.8(6.1)(4.9)
CMD-2	1.18	14%	781.98(29)(1)	1.88(6)(2)		496.9(4.0)(2.3)
	1.01	45%	782.64(33)(4)	1.85(6)(4)	11.4(3.1)(1.0)	495.8(3.7)(4.2)
BaBar	1.14	5.7%	781.86(14)(1)	2.04(3)(2)		501.9(3.3)(2.0)
	1.14	5.5%	781.93(18)(4)	2.03(4)(1)	1.3(1.9)(0.7)	501.9(3.3)(1.8)
KLOE	1.36	7.4×10^{-4}	781.82(17)(4)	1.97(4)(2)		492.0(2.2)(1.8)
	1.27	6.7×10^{-3}	782.50(25)(6)	1.94(5)(2)	6.8(1.8)(0.5)	491.0(2.2)(2.0)
KLOE''	1.20	3.1%	781.81(16)(3)	1.98(4)(1)		491.8(2.1)(1.8)
	1.13	10%	782.42(23)(5)	1.95(4)(2)	6.1(1.7)(0.6)	490.8(2.0)(1.7)
BESIII	1.12	25%	782.18(51)(7)	2.01(19)(9)		490.8(4.8)(3.9)
	1.02	44%	783.05(60)(2)	1.99(19)(7)	17.6(6.9)(1.2)	490.3(4.5)(3.1)
SND20	2.93	3.3×10^{-7}	781.79(30)(6)	2.04(6)(3)		494.2(6.7)(9.0)
	1.87	4.1×10^{-3}	782.37(28)(6)	2.02(5)(2)	10.1(2.4)(1.4)	494.9(5.3)(3.1)

Table 1. Comparison of fits to single experiments with and without a phase δ_ϵ in ϵ_ω . Note that BESIII has only a few data points in the interference region and hence is not able to put a strong constraint on δ_ϵ (the corresponding line is indicated in gray). The first error is the fit uncertainty, inflated by $\sqrt{\chi^2/\text{dof}}$, the second error is the combination of all systematic uncertainties.

	χ^2/dof	p -value	M_ω [MeV]	$10^3 \times \text{Re } \epsilon_\omega$	δ_ϵ [°]	$10^{10} \times a_\mu^{\pi\pi} _{\leq 1 \text{ GeV}}$
Energy scan w/o SND20	1.28	2.1%	781.75(22)(1)	1.97(4)(2)		498.5(3.4)(2.6)
	1.05	33%	782.39(23)(2)	1.93(4)(3)	9.9(1.8)(0.4)	497.3(3.1)(3.9)
Energy scan	1.65	6.3×10^{-7}	781.74(17)(2)	2.01(3)(3)		497.4(3.0)(4.4)
	1.19	5.2%	782.37(16)(3)	1.97(3)(3)	10.1(1.3)(0.7)	496.0(2.6)(5.5)
All e^+e^- w/o SND20	1.25	1.8×10^{-5}	781.70(9)(4)	2.02(2)(3)		494.5(1.5)(2.3)
	1.20	3.3×10^{-4}	782.10(12)(4)	1.96(2)(2)	4.5(9)(8)	494.2(1.4)(2.1)
NA7 + all e^+e^- w/o SND20	1.23	3.0×10^{-5}	781.69(9)(3)	2.02(2)(3)		494.8(1.4)(2.1)
	1.19	4.8×10^{-4}	782.09(12)(4)	1.97(2)(2)	4.5(9)(8)	494.6(1.5)(1.7)
All e^+e^-	1.36	1.0×10^{-9}	781.71(8)(3)	2.02(2)(3)		495.0(1.4)(2.4)
	1.30	2.3×10^{-7}	782.09(10)(4)	1.97(2)(2)	4.5(8)(8)	494.6(1.4)(2.1)
NA7 + all e^+e^-	1.34	2.5×10^{-9}	781.71(8)(3)	2.02(2)(3)		495.2(1.4)(2.2)
	1.28	4.5×10^{-7}	782.09(10)(4)	1.97(2)(2)	4.5(8)(8)	494.9(1.4)(1.8)

Table 2. Comparison of fits to combinations of experiments with and without a phase δ_ϵ in ϵ_ω . The first error is the fit uncertainty, inflated by $\sqrt{\chi^2/\text{dof}}$, the second error is the combination of all systematic uncertainties.

For BESIII, the preferred central value for δ_ϵ comes out even larger, yet with a very large uncertainty that reflects the limited sensitivity to δ_ϵ , resulting from a relatively small number of data points in the ρ - ω region (accordingly, this line is indicated in light gray in tables 1 and 3). Finally, the KLOE fits produce a phase slightly larger than expected, while the BaBar data are even consistent with $\delta_\epsilon = 0$. We thus observe a large spread in the results for the phase of δ_ϵ , pointing towards systematic differences among the data sets in the ρ - ω region.

	δ_ϵ [°]	SD window	int window	LD window	$10^{10} \times a_\mu^{\pi\pi} _{\leq 1 \text{ GeV}}$
SND06		13.9(2)(1)	140.0(2.0)(1.0)	345.8(4.7)(3.0)	499.7(6.9)(4.1)
	8.5(2.3)(0.3)	13.9(2)(1)	139.6(1.8)(1.2)	344.3(4.1)(3.6)	497.8(6.1)(4.9)
CMD-2		13.9(1)(0)	139.5(1.1)(0.4)	343.6(2.7)(1.8)	496.9(4.0)(2.3)
	11.4(3.1)(1.0)	13.9(1)(1)	139.4(1.0)(0.9)	342.6(2.5)(3.2)	495.8(3.7)(4.2)
BaBar		14.0(1)(0)	140.6(1.0)(0.5)	347.3(2.2)(1.5)	501.9(3.3)(2.0)
	1.3(1.9)(0.7)	14.0(1)(0)	140.6(1.0)(0.5)	347.3(2.3)(1.3)	501.9(3.3)(1.8)
KLOE''		13.6(1)(1)	137.3(6)(6)	340.9(1.4)(1.2)	491.8(2.1)(1.8)
	6.1(1.7)(0.6)	13.6(1)(0)	137.1(6)(4)	340.2(1.4)(1.3)	490.8(2.0)(1.7)
BESIII		13.7(1)(0)	138.0(1.4)(0.5)	339.0(3.3)(3.4)	490.8(4.8)(3.9)
	17.6(6.9)(1.2)	13.7(1)(0)	137.8(1.3)(0.4)	338.8(3.1)(2.6)	490.3(4.5)(3.1)
SND20		13.9(2)(1)	139.4(1.9)(1.5)	340.9(4.6)(7.4)	494.2(6.7)(9.0)
	10.1(2.4)(1.4)	13.8(2)(0)	139.2(1.5)(0.5)	341.9(3.7)(2.6)	494.9(5.3)(3.1)
NA7 + all e^+e^- w/o SND20		13.7(0)(0)	138.3(4)(5)	342.7(1.0)(1.6)	494.8(1.4)(2.1)
	4.5(9)(8)	13.7(0)(0)	138.3(4)(4)	342.5(1.0)(1.3)	494.6(1.5)(1.7)

Table 3. Decomposition of $10^{10} \times a_\mu^{\pi\pi}|_{\leq 1 \text{ GeV}}$ into the Euclidean windows from ref. [36]. The first error is the fit uncertainty, inflated by $\sqrt{\chi^2/\text{dof}}$, the second error is the combination of all systematic uncertainties.

In addition, we confirm a correlation between δ_ϵ and M_ω , as already observed in ref. [18]: the larger the phase, the larger the extracted value of M_ω . However, as discussed in more detail in section 5, the size of the phase permitted by radiative intermediate states, roughly in line with the result of the global fit shown in table 2, does not suffice to remove the tension with ω -mass determinations from $e^+e^- \rightarrow 3\pi$ and $e^+e^- \rightarrow \pi^0\gamma$.

In this regard, we also observe that the BESIII data suggest larger values of M_ω than all other data sets, with the result for $\delta_\epsilon = 0$ close to the global fit with non-vanishing phase. Within uncertainties there is still consistency, but it is noteworthy that the size and direction of the effect echo a similar tension in $e^+e^- \rightarrow 3\pi$ [12, 114–116] and $\eta' \rightarrow \pi^+\pi^-\gamma$ [117, 118].

4 Isospin-breaking contribution to a_μ from ρ - ω mixing

Based on the dispersive representation (2.2) we can quantify $a_\mu^{\rho-\omega}$ — the IB contribution to a_μ due to ϵ_ω — by contrasting the full result to the HVP integral evaluated with $\epsilon_\omega = 0$. In principle, there is some ambiguity due to final-state radiation (FSR), but in practice this effect comes out well below 0.1×10^{-10} . For definiteness, in table 4 we show the variant without FSR, to isolate the pure $\mathcal{O}(\epsilon_\omega)$ terms.

In general, $a_\mu^{\rho-\omega}$ is sensitive to the assumed line shape [119]. However, we find that the dispersive representations (2.4) or (2.17) are quite robust in that regard, i.e., with the threshold behavior and the properties close to the ω pole determined, the remaining interpolation only has a marginal effect, e.g., changing $s \rightarrow M_\omega^2$ in the numerator of eq. (2.5) changes the outcome for $a_\mu^{\rho-\omega}$ again by less than 0.1×10^{-10} . In contrast, whether or not a phase in ϵ_ω is permitted does change the resulting value for $a_\mu^{\rho-\omega}$ in a significant way, and we show results for both scenarios (the difference in the FSR contribution is of $\mathcal{O}(e^2\epsilon_\omega)$ and negligible). Since ϵ_ω in the global fit comes out close to the narrow-resonance

	SD window	int window	LD window	total
$10^{10} \times a_\mu^{\rho-\omega}, \delta_\epsilon = 0$	0.08(0)(0)	1.06(1)(2)	3.23(3)(5)	4.37(4)(7)
$10^{10} \times a_\mu^{\pi\pi, \text{FSR}}, \delta_\epsilon = 0$	0.11(0)(0)	1.12(0)(0)	3.00(1)(1)	4.23(1)(2)
$10^{10} \times a_\mu^{\rho-\omega}, \delta_\epsilon = 4.5(1.2)^\circ$	0.05(0)(0)	0.83(5)(4)	2.79(9)(6)	3.68(14)(10)
$10^{10} \times a_\mu^{\pi\pi, \text{FSR}}, \delta_\epsilon = 4.5(1.2)^\circ$	0.11(0)(0)	1.12(0)(0)	3.00(1)(1)	4.24(1)(2)

Table 4. IB contribution to $a_\mu^{\pi\pi}|_{\leq 1 \text{ GeV}}$ due to ρ - ω mixing, compared to the effect of FSR and split into the different Euclidean windows from ref. [36]. We only include the linear effects, i.e., $\mathcal{O}(\epsilon_\omega)$ for the ρ - ω -mixing contribution and the $\mathcal{O}(e^2)$ effect for FSR — $\mathcal{O}(e^2\epsilon_\omega)$ effects give very small corrections. The results correspond to the combined fit to all experiments apart from SND20. The first error is the fit uncertainty, inflated by $\sqrt{\chi^2/\text{dof}}$, the second error is the combination of all systematic uncertainties.

expectation, we quote the variant with non-vanishing δ_ϵ as our preferred result, which has already been used as input in estimating the three-flavor quark-disconnected contribution to a_μ in ref. [120].⁴ Finally, we also provide the breakdown of $a_\mu^{\rho-\omega}$ onto the Euclidean windows from ref. [36].

Further, for the comparison to lattice QCD it is also of interest to study the decomposition of ϵ_ω into its $\mathcal{O}(e^2)$ and $\mathcal{O}(m_u - m_d)$ pieces, as was discussed in the context of resonance chiral perturbation theory in ref. [121]. Translated to our normalization one has the prediction [121]

$$\tilde{\epsilon}_\omega = \frac{2}{3R} \frac{M_{K^*} - M_V}{M_V} - \frac{e^2}{|g_{\omega\gamma}|^2}, \quad R = \frac{m_s - \hat{m}}{m_d - m_u}, \quad \hat{m} = \frac{m_u + m_d}{2}, \quad (4.1)$$

where we have written the electromagnetic component in terms of the ω - γ coupling, as this is the quantity that enters directly in the corresponding diagram. However, this latter diagram produces a one-particle-reducible correction, and would thus be subtracted when vacuum polarization is removed from the $e^+e^- \rightarrow \pi^+\pi^-$ cross sections. Accordingly, we have that our conventions are related to the ones of ref. [121] by

$$\tilde{\epsilon}_\omega = \epsilon_\omega + \tilde{\epsilon}_\omega|_{e^2}, \quad \tilde{\epsilon}_\omega|_{e^2} = -\frac{e^2}{|g_{\omega\gamma}|^2} = -0.34(1) \times 10^{-3}, \quad (4.2)$$

using $\Gamma[\omega \rightarrow e^+e^-] = 4\pi\alpha^2 M_\omega / (3|g_{\omega\gamma}|^2)$ and the average of ref. [109] for the $\omega \rightarrow e^+e^-$ branching fraction.

The prediction for the $\mathcal{O}(m_u - m_d)$ part of $\tilde{\epsilon}_\omega$, which coincides with our ϵ_ω , is far less robust, as already from higher-order quark-mass and SU(3)-breaking corrections one would expect an accuracy around 30%. Using the $N_f = 2 + 1$ and $N_f = 2 + 1 + 1$ averages from ref. [122], $R = 38.1(1.5)$ [123–128] and $R = 35.9(1.7)$ [129–133], respectively, and identifying the vector mesons with the neutral ρ and K^* resonances, the predictions for the strong IB contribution become $\tilde{\epsilon}_\omega|_{m_u - m_d}^{2+1} = 2.71(11) \times 10^{-3}$, $\tilde{\epsilon}_\omega|_{m_u - m_d}^{2+1+1} = 2.88(14) \times 10^{-3}$,

⁴The tiny difference to the number for $a_\mu^{\rho-\omega}$ quoted therein as private communication originates from the improved implementation of the $\pi^0\gamma$ threshold (2.17).

Reference	$e^+e^- \rightarrow 3\pi$	$e^+e^- \rightarrow \pi^0\gamma$	$e^+e^- \rightarrow 2\pi$	PDG average
Ref. [5]	782.631(28)			
Ref. [46]		782.584(28)		
Ref. [4]			781.68(10)	
This work, $\delta_\epsilon = 0$			781.69(9)	
This work, $\delta_\epsilon = 4.5(1.2)^\circ$			782.09(12)	
Ref. [109]				782.53(13)

Table 5. Dispersive determinations of M_ω in MeV from e^+e^- reactions, compared to the global average from ref. [109]. In all cases, vacuum-polarization corrections are not included, and the average from ref. [109] has been adjusted accordingly using $\Delta M_\omega = 0.13$ MeV [117].

about 40% larger than results from our fit to the $e^+e^- \rightarrow \pi^+\pi^-$ data. This is in line with subsequent work on vector mesons in chiral perturbation theory [134, 135], which concluded that higher-order corrections can be substantial. This includes photon loops, short-distance corrections, and meson loops, parts of which scale with e^2 and thus lead to electromagnetic effects not subtracted when removing vacuum polarization in the definition of the bare cross section.

However, from the LO expression (4.1) it still follows that the ρ - ω -mixing contribution to a_μ should be considered primarily a quark-mass effect,

$$a_\mu^{\rho-\omega}[e^2, \text{LO}] = 0, \quad a_\mu^{\rho-\omega}[m_u - m_d, \text{LO}] = 3.68(17) \times 10^{-10}, \quad (4.3)$$

which is expected to yield the dominant strong IB contribution to a_μ .⁵ This number agrees well with a recent estimate from SU(3) chiral perturbation theory, $a_\mu[m_u - m_d]_{[136]} = 3.32(89) \times 10^{-10}$, where the required low-energy constant is determined from hadronic τ decays. Both indicate a somewhat larger central value than the lattice-QCD result $a_\mu[m_u - m_d]_{[32]} = 1.9(1.2) \times 10^{-10}$.

5 Consequences for the ω mass

The correlation between δ_ϵ and M_ω discussed in section 3 affects the resulting determination of M_ω from $e^+e^- \rightarrow 2\pi$. In table 5 we compare our updated extraction from the 2π data to analogous ones from $e^+e^- \rightarrow 3\pi$, $e^+e^- \rightarrow \pi^0\gamma$, as well as the average from ref. [109]. As discussed in more detail in refs. [4, 5, 46], the PDG average involves a cancellation between determinations from $e^+e^- \rightarrow \pi^0\gamma$ [137] and $\bar{p}p \rightarrow \omega\pi^0\pi^0$ [138], while dominated by Breit-Wigner-based extractions from $e^+e^- \rightarrow 3\pi$ [12, 116] that are in agreement with the dispersive result given in table 5 (further confirmed by the recent BaBar measurement [115], while BESIII suggests a larger value [114]). Our updated value for $\delta_\epsilon = 0$ changes only marginally compared to ref. [4], leading to the same 5σ tension with the PDG value observed therein. Allowing a finite value for δ_ϵ instead removes about half the discrepancy, but we

⁵Resonance-enhanced threshold effects in the $\bar{K}K$ channels largely cancel between K^+K^- and $K_S K_L$.

emphasize that this effect cannot explain the entire tension as it would require a size of δ_ϵ that cannot be reconciled with the strength of the radiative channels giving rise to a phase in ϵ_ω in the first place.

6 Conclusions

In this work we performed a detailed study of ρ - ω mixing in $e^+e^- \rightarrow \pi^+\pi^-$, based on a dispersive representation of the pion vector form factor. In particular, we investigated the role of imaginary parts that can be generated by radiative intermediate states coupling ω and ρ resonances, estimated their size by narrow-width arguments, and devised a strategy to include their effect in fits to the $e^+e^- \rightarrow \pi^+\pi^-$ data base. We found that while the size of the phase of the ρ - ω mixing parameter in a global fit does come out in agreement with narrow-resonance expectations, see eq. (3.1) for the central results, there is a substantial spread among the different data sets, ranging from a vanishing phase to values as large as 10° . As applications, we derived the isospin-breaking part of the HVP contribution to a_μ originating from ρ - ω mixing and quantified the changes in the extracted value of the ω mass when a non-vanishing phase is permitted.

Our work reveals systematic differences in the low-energy hadronic cross sections that go beyond the well-known BaBar-KLOE tension in the $e^+e^- \rightarrow \pi^+\pi^-$ total cross section, including the spread in the phase of the ρ - ω mixing parameter and discrepancies in the ω mass extracted from different decay channels, both of which can be unambiguously defined in terms of pole parameters and residues. While of course resolving the tension in the HVP integral itself carries the highest priority, forthcoming high-precision data on $e^+e^- \rightarrow \pi^+\pi^-$ should also allow one to address the tensions pointed out here, and thus increase confidence that the hadronic cross sections are understood at the level required for robust data-driven evaluations of the HVP contribution to the anomalous magnetic moment of the muon.

Acknowledgments

We thank Christoph Hanhart for valuable discussions that led to the derivation in section 2.2, and Kim Maltman for encouragement to write up these results. We further thank Pablo Sánchez-Puertas for pointing out the issue of one-photon-reducible contributions to ϵ_ω , and Hans Bijnens for discussions on ϵ_ω in resonance chiral perturbation theory. Financial support by the DFG through the funds provided to the Sino-German Collaborative Research Center TRR110 “Symmetries and the Emergence of Structure in QCD” (DFG Project-ID 196253076 — TRR 110) and the SNSF (Project Nos. 200020_175791, PCEFP2_181117, and PCEFP2_194272) is gratefully acknowledged.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited. SCOAP³ supports the goals of the International Year of Basic Sciences for Sustainable Development.

References

- [1] T. Aoyama et al., *The anomalous magnetic moment of the muon in the Standard Model*, *Phys. Rept.* **887** (2020) 1 [[arXiv:2006.04822](#)] [[INSPIRE](#)].
- [2] M. Davier, A. Hoecker, B. Malaescu and Z. Zhang, *Reevaluation of the hadronic vacuum polarisation contributions to the Standard Model predictions of the muon $g - 2$ and $\alpha(m_Z^2)$ using newest hadronic cross-section data*, *Eur. Phys. J. C* **77** (2017) 827 [[arXiv:1706.09436](#)] [[INSPIRE](#)].
- [3] A. Keshavarzi, D. Nomura and T. Teubner, *Muon $g - 2$ and $\alpha(M_Z^2)$: a new data-based analysis*, *Phys. Rev. D* **97** (2018) 114025 [[arXiv:1802.02995](#)] [[INSPIRE](#)].
- [4] G. Colangelo, M. Hoferichter and P. Stoffer, *Two-pion contribution to hadronic vacuum polarization*, *JHEP* **02** (2019) 006 [[arXiv:1810.00007](#)] [[INSPIRE](#)].
- [5] M. Hoferichter, B.-L. Hoid and B. Kubis, *Three-pion contribution to hadronic vacuum polarization*, *JHEP* **08** (2019) 137 [[arXiv:1907.01556](#)] [[INSPIRE](#)].
- [6] M. Davier, A. Hoecker, B. Malaescu and Z. Zhang, *A new evaluation of the hadronic vacuum polarisation contributions to the muon anomalous magnetic moment and to $\alpha(m_Z^2)$* , *Eur. Phys. J. C* **80** (2020) 241 [Erratum *ibid.* **80** (2020) 410] [[arXiv:1908.00921](#)] [[INSPIRE](#)].
- [7] A. Keshavarzi, D. Nomura and T. Teubner, *$g - 2$ of charged leptons, $\alpha(M_Z^2)$, and the hyperfine splitting of muonium*, *Phys. Rev. D* **101** (2020) 014029 [[arXiv:1911.00367](#)] [[INSPIRE](#)].
- [8] MUON $g - 2$ collaboration, *Muon $g - 2$ Technical Design Report*, [arXiv:1501.06858](#) [[INSPIRE](#)].
- [9] M.N. Achasov et al., *Study of the process $e^+e^- \rightarrow \pi^+\pi^-$ in the energy region $400 < \sqrt{s} < 1000$ MeV*, *J. Exp. Theor. Phys.* **101** (2005) 1053 [[hep-ex/0506076](#)] [[INSPIRE](#)].
- [10] M.N. Achasov et al., *Update of the $e^+e^- \rightarrow \pi^+\pi^-$ cross-section measured by SND detector in the energy region $400 < \sqrt{s} < 1000$ MeV*, *J. Exp. Theor. Phys.* **103** (2006) 380 [[hep-ex/0605013](#)] [[INSPIRE](#)].
- [11] CMD-2 collaboration, *Measurement of $e^+e^- \rightarrow \pi^+\pi^-$ cross-section with CMD-2 around ρ -meson*, *Phys. Lett. B* **527** (2002) 161 [[hep-ex/0112031](#)] [[INSPIRE](#)].
- [12] CMD-2 collaboration, *Reanalysis of hadronic cross-section measurements at CMD-2*, *Phys. Lett. B* **578** (2004) 285 [[hep-ex/0308008](#)] [[INSPIRE](#)].
- [13] V.M. Aul'chenko et al., *Measurement of the $e^+e^- \rightarrow \pi^+\pi^-$ cross section with the CMD-2 detector in the 370–520 MeV c.m. energy range*, *JETP Lett.* **84** (2006) 413 [[hep-ex/0610016](#)] [[INSPIRE](#)].
- [14] CMD-2 collaboration, *High-statistics measurement of the pion form factor in the ρ -meson energy range with the CMD-2 detector*, *Phys. Lett. B* **648** (2007) 28 [[hep-ex/0610021](#)] [[INSPIRE](#)].
- [15] BESIII collaboration, *Measurement of the $e^+e^- \rightarrow \pi^+\pi^-$ cross section between 600 and 900 MeV using initial state radiation*, *Phys. Lett. B* **753** (2016) 629 [Erratum *ibid.* **812** (2021) 135982] [[arXiv:1507.08188](#)] [[INSPIRE](#)].
- [16] T. Xiao, S. Dobbs, A. Tomaradze, K.K. Seth and G. Bonvicini, *Precision Measurement of the Hadronic Contribution to the Muon Anomalous Magnetic Moment*, *Phys. Rev. D* **97** (2018) 032012 [[arXiv:1712.04530](#)] [[INSPIRE](#)].

- [17] BABAR collaboration, *Precise measurement of the $e^+e^- \rightarrow \pi^+\pi^-(\gamma)$ cross section with the Initial State Radiation method at BABAR*, *Phys. Rev. Lett.* **103** (2009) 231801 [[arXiv:0908.3589](#)] [[INSPIRE](#)].
- [18] BABAR collaboration, *Precise Measurement of the $e^+e^- \rightarrow \pi^+\pi^-(\gamma)$ Cross Section with the Initial-State Radiation Method at BABAR*, *Phys. Rev. D* **86** (2012) 032013 [[arXiv:1205.2228](#)] [[INSPIRE](#)].
- [19] KLOE collaboration, *Measurement of $\sigma(e^+e^- \rightarrow \pi^+\pi^-\gamma(\gamma))$ and the dipion contribution to the muon anomaly with the KLOE detector*, *Phys. Lett. B* **670** (2009) 285 [[arXiv:0809.3950](#)] [[INSPIRE](#)].
- [20] KLOE collaboration, *Measurement of $\sigma(e^+e^- \rightarrow \pi^+\pi^-)$ from threshold to 0.85 GeV^2 using Initial State Radiation with the KLOE detector*, *Phys. Lett. B* **700** (2011) 102 [[arXiv:1006.5313](#)] [[INSPIRE](#)].
- [21] KLOE collaboration, *Precision measurement of $\sigma(e^+e^- \rightarrow \pi^+\pi^-\gamma)/\sigma(e^+e^- \rightarrow \mu^+\mu^-\gamma)$ and determination of the $\pi^+\pi^-$ contribution to the muon anomaly with the KLOE detector*, *Phys. Lett. B* **720** (2013) 336 [[arXiv:1212.4524](#)] [[INSPIRE](#)].
- [22] KLOE-2 collaboration, *Combination of KLOE $\sigma(e^+e^- \rightarrow \pi^+\pi^-\gamma(\gamma))$ measurements and determination of $a_\mu^{\pi^+\pi^-}$ in the energy range $0.10 < s < 0.95 \text{ GeV}^2$* , *JHEP* **03** (2018) 173 [[arXiv:1711.03085](#)] [[INSPIRE](#)].
- [23] B. Ananthanarayan, I. Caprini and D. Das, *Pion electromagnetic form factor at high precision with implications to $a_\mu^{\pi\pi}$ and the onset of perturbative QCD*, *Phys. Rev. D* **98** (2018) 114015 [[arXiv:1810.09265](#)] [[INSPIRE](#)].
- [24] D. Stamen, D. Hariharan, M. Hoferichter, B. Kubis and P. Stoffer, *Kaon electromagnetic form factors in dispersion theory*, *Eur. Phys. J. C* **82** (2022) 432 [[arXiv:2202.11106](#)] [[INSPIRE](#)].
- [25] E.B. Dally et al., *Elastic Scattering Measurement of the Negative Pion Radius*, *Phys. Rev. Lett.* **48** (1982) 375 [[INSPIRE](#)].
- [26] NA7 collaboration, *A Measurement of the Space-Like Pion Electromagnetic Form-Factor*, *Nucl. Phys. B* **277** (1986) 168 [[INSPIRE](#)].
- [27] SND collaboration, *Measurement of the $e^+e^- \rightarrow \pi^+\pi^-$ process cross section with the SND detector at the VEPP-2000 collider in the energy region $0.525 < \sqrt{s} < 0.883 \text{ GeV}$* , *JHEP* **01** (2021) 113 [[arXiv:2004.00263](#)] [[INSPIRE](#)].
- [28] A.E. Ryzhnenkov et al., *Overview of the CMD-3 recent results*, *J. Phys. Conf. Ser.* **1526** (2020) 012009 [[INSPIRE](#)].
- [29] G. Abbiendi et al., *Mini-Proceedings of the STRONG2020 Virtual Workshop on “Space-like and Time-like determination of the Hadronic Leading Order contribution to the Muon $g - 2$ ”*, [[arXiv:2201.12102](#)] [[INSPIRE](#)].
- [30] BESIII collaboration, *Future Physics Programme of BESIII*, *Chin. Phys. C* **44** (2020) 040001 [[arXiv:1912.05983](#)] [[INSPIRE](#)].
- [31] BELLE-II collaboration, *The Belle II Physics Book*, *PTEP* **2019** (2019) 123C01 [Erratum *ibid.* **2020** (2020) 029201] [[arXiv:1808.10567](#)] [[INSPIRE](#)].
- [32] S. Borsányi et al., *Leading hadronic contribution to the muon magnetic moment from lattice QCD*, *Nature* **593** (2021) 51 [[arXiv:2002.12347](#)] [[INSPIRE](#)].
- [33] M. Cè et al., *Window observable for the hadronic vacuum polarization contribution to the muon $g - 2$ from lattice QCD*, [[arXiv:2206.06582](#)] [[INSPIRE](#)].

- [34] C. Alexandrou et al., *Lattice calculation of the short and intermediate time-distance hadronic vacuum polarization contributions to the muon magnetic moment using twisted-mass fermions*, [arXiv:2206.15084](#) [INSPIRE].
- [35] FERMILAB LATTICE, HPQCD and MILC collaborations, *Windows on the hadronic vacuum polarisation contribution to the muon anomalous magnetic moment*, [arXiv:2207.04765](#) [INSPIRE].
- [36] RBC and UKQCD collaborations, *Calculation of the hadronic vacuum polarization contribution to the muon anomalous magnetic moment*, *Phys. Rev. Lett.* **121** (2018) 022003 [[arXiv:1801.07224](#)] [INSPIRE].
- [37] MUON G-2 collaboration, *Final Report of the Muon E821 Anomalous Magnetic Moment Measurement at BNL*, *Phys. Rev. D* **73** (2006) 072003 [[hep-ex/0602035](#)] [INSPIRE].
- [38] MUON G-2 collaboration, *Measurement of the Positive Muon Anomalous Magnetic Moment to 0.46 ppm*, *Phys. Rev. Lett.* **126** (2021) 141801 [[arXiv:2104.03281](#)] [INSPIRE].
- [39] MUON G-2 collaboration, *Magnetic-field measurement and analysis for the Muon $g - 2$ Experiment at Fermilab*, *Phys. Rev. A* **103** (2021) 042208 [[arXiv:2104.03201](#)] [INSPIRE].
- [40] MUON G-2 collaboration, *Beam dynamics corrections to the Run-1 measurement of the muon anomalous magnetic moment at Fermilab*, *Phys. Rev. Accel. Beams* **24** (2021) 044002 [[arXiv:2104.03240](#)] [INSPIRE].
- [41] MUON G-2 collaboration, *Measurement of the anomalous precession frequency of the muon in the Fermilab Muon $g - 2$ Experiment*, *Phys. Rev. D* **103** (2021) 072002 [[arXiv:2104.03247](#)] [INSPIRE].
- [42] T. Aoyama, M. Hayakawa, T. Kinoshita and M. Nio, *Complete Tenth-Order QED Contribution to the Muon $g - 2$* , *Phys. Rev. Lett.* **109** (2012) 111808 [[arXiv:1205.5370](#)] [INSPIRE].
- [43] T. Aoyama, T. Kinoshita and M. Nio, *Theory of the Anomalous Magnetic Moment of the Electron*, *Atoms* **7** (2019) 28 [INSPIRE].
- [44] A. Czarnecki, W.J. Marciano and A. Vainshtein, *Refinements in electroweak contributions to the muon anomalous magnetic moment*, *Phys. Rev. D* **67** (2003) 073006 [Erratum *ibid.* **73** (2006) 119901] [[hep-ph/0212229](#)] [INSPIRE].
- [45] C. Gnendiger, D. Stöckinger and H. Stöckinger-Kim, *The electroweak contributions to $(g - 2)_\mu$ after the Higgs boson mass measurement*, *Phys. Rev. D* **88** (2013) 053005 [[arXiv:1306.5546](#)] [INSPIRE].
- [46] B.-L. Hoid, M. Hoferichter and B. Kubis, *Hadronic vacuum polarization and vector-meson resonance parameters from $e^+e^- \rightarrow \pi^0\gamma$* , *Eur. Phys. J. C* **80** (2020) 988 [[arXiv:2007.12696](#)] [INSPIRE].
- [47] A. Kurz, T. Liu, P. Marquard and M. Steinhauser, *Hadronic contribution to the muon anomalous magnetic moment to next-to-next-to-leading order*, *Phys. Lett. B* **734** (2014) 144 [[arXiv:1403.6400](#)] [INSPIRE].
- [48] K. Melnikov and A. Vainshtein, *Hadronic light-by-light scattering contribution to the muon anomalous magnetic moment revisited*, *Phys. Rev. D* **70** (2004) 113006 [[hep-ph/0312226](#)] [INSPIRE].
- [49] P. Masjuan and P. Sánchez-Puertas, *Pseudoscalar-pole contribution to the $(g_\mu - 2)$: a rational approach*, *Phys. Rev. D* **95** (2017) 054026 [[arXiv:1701.05829](#)] [INSPIRE].

- [50] G. Colangelo, M. Hoferichter, M. Procura and P. Stoffer, *Rescattering effects in the hadronic-light-by-light contribution to the anomalous magnetic moment of the muon*, *Phys. Rev. Lett.* **118** (2017) 232001 [[arXiv:1701.06554](#)] [[INSPIRE](#)].
- [51] G. Colangelo, M. Hoferichter, M. Procura and P. Stoffer, *Dispersion relation for hadronic light-by-light scattering: two-pion contributions*, *JHEP* **04** (2017) 161 [[arXiv:1702.07347](#)] [[INSPIRE](#)].
- [52] M. Hoferichter, B.-L. Hoid, B. Kubis, S. Leupold and S.P. Schneider, *Pion-pole contribution to hadronic light-by-light scattering in the anomalous magnetic moment of the muon*, *Phys. Rev. Lett.* **121** (2018) 112002 [[arXiv:1805.01471](#)] [[INSPIRE](#)].
- [53] M. Hoferichter, B.-L. Hoid, B. Kubis, S. Leupold and S.P. Schneider, *Dispersion relation for hadronic light-by-light scattering: pion pole*, *JHEP* **10** (2018) 141 [[arXiv:1808.04823](#)] [[INSPIRE](#)].
- [54] A. Gérardin, H.B. Meyer and A. Nyffeler, *Lattice calculation of the pion transition form factor with $N_f = 2 + 1$ Wilson quarks*, *Phys. Rev. D* **100** (2019) 034520 [[arXiv:1903.09471](#)] [[INSPIRE](#)].
- [55] J. Bijnens, N. Hermansson-Truedsson and A. Rodríguez-Sánchez, *Short-distance constraints for the HLbL contribution to the muon anomalous magnetic moment*, *Phys. Lett. B* **798** (2019) 134994 [[arXiv:1908.03331](#)] [[INSPIRE](#)].
- [56] G. Colangelo, F. Hagelstein, M. Hoferichter, L. Laub and P. Stoffer, *Short-distance constraints on hadronic light-by-light scattering in the anomalous magnetic moment of the muon*, *Phys. Rev. D* **101** (2020) 051501 [[arXiv:1910.11881](#)] [[INSPIRE](#)].
- [57] G. Colangelo, F. Hagelstein, M. Hoferichter, L. Laub and P. Stoffer, *Longitudinal short-distance constraints for the hadronic light-by-light contribution to $(g - 2)_\mu$ with large- N_c Regge models*, *JHEP* **03** (2020) 101 [[arXiv:1910.13432](#)] [[INSPIRE](#)].
- [58] T. Blum et al., *Hadronic Light-by-Light Scattering Contribution to the Muon Anomalous Magnetic Moment from Lattice QCD*, *Phys. Rev. Lett.* **124** (2020) 132002 [[arXiv:1911.08123](#)] [[INSPIRE](#)].
- [59] G. Colangelo, M. Hoferichter, A. Nyffeler, M. Passera and P. Stoffer, *Remarks on higher-order hadronic corrections to the muon $g - 2$* , *Phys. Lett. B* **735** (2014) 90 [[arXiv:1403.7512](#)] [[INSPIRE](#)].
- [60] M. Passera, W.J. Marciano and A. Sirlin, *The Muon $g - 2$ and the bounds on the Higgs boson mass*, *Phys. Rev. D* **78** (2008) 013009 [[arXiv:0804.1142](#)] [[INSPIRE](#)].
- [61] A. Crivellin, M. Hoferichter, C.A. Manzari and M. Montull, *Hadronic Vacuum Polarization: $(g - 2)_\mu$ versus Global Electroweak Fits*, *Phys. Rev. Lett.* **125** (2020) 091801 [[arXiv:2003.04886](#)] [[INSPIRE](#)].
- [62] A. Keshavarzi, W.J. Marciano, M. Passera and A. Sirlin, *Muon $g - 2$ and $\Delta\alpha$ connection*, *Phys. Rev. D* **102** (2020) 033002 [[arXiv:2006.12666](#)] [[INSPIRE](#)].
- [63] B. Malaescu and M. Schott, *Impact of correlations between a_μ and α_{QED} on the EW fit*, *Eur. Phys. J. C* **81** (2021) 46 [[arXiv:2008.08107](#)] [[INSPIRE](#)].
- [64] G. Colangelo, M. Hoferichter and P. Stoffer, *Constraints on the two-pion contribution to hadronic vacuum polarization*, *Phys. Lett. B* **814** (2021) 136073 [[arXiv:2010.07943](#)] [[INSPIRE](#)].
- [65] M. Cè et al., *The hadronic running of the electromagnetic coupling and the electroweak mixing angle from lattice QCD*, *JHEP* **08** (2022) 220 [[arXiv:2203.08676](#)] [[INSPIRE](#)].

- [66] G. Colangelo et al., *Prospects for precise predictions of a_μ in the Standard Model*, [arXiv:2203.15810](#) [INSPIRE].
- [67] G. Colangelo et al., *Data-driven evaluations of Euclidean windows to scrutinize hadronic vacuum polarization*, *Phys. Lett. B* **833** (2022) 137313 [[arXiv:2205.12963](#)] [INSPIRE].
- [68] WORKING GROUP ON RADIATIVE CORRECTIONS, MONTE CARLO GENERATORS FOR LOW ENERGIES collaboration, *Quest for precision in hadronic cross sections at low energy: Monte Carlo tools vs. experimental data*, *Eur. Phys. J. C* **66** (2010) 585 [[arXiv:0912.0749](#)] [INSPIRE].
- [69] A. Hofer, J. Gluza and F. Jegerlehner, *Pion pair production with higher order radiative corrections in low energy e^+e^- collisions*, *Eur. Phys. J. C* **24** (2002) 51 [[hep-ph/0107154](#)] [INSPIRE].
- [70] H. Czyż, A. Grzebińska, J.H. Kühn and G. Rodrigo, *The Radiative return at ϕ and B factories: FSR for muon pair production at next-to-leading order*, *Eur. Phys. J. C* **39** (2005) 411 [[hep-ph/0404078](#)] [INSPIRE].
- [71] J. Gluza, A. Hofer, S. Jadach and F. Jegerlehner, *Measuring the FSR inclusive $\pi^+\pi^-$ cross-section*, *Eur. Phys. J. C* **28** (2003) 261 [[hep-ph/0212386](#)] [INSPIRE].
- [72] Y.M. Bystritskiy, E.A. Kuraev, G.V. Fedotovitch and F.V. Ignatov, *The Cross sections of the muons and charged pions pairs production at electron-positron annihilation near the threshold*, *Phys. Rev. D* **72** (2005) 114019 [[hep-ph/0505236](#)] [INSPIRE].
- [73] F. Campanario et al., *Standard model radiative corrections in the pion form factor measurements do not explain the a_μ anomaly*, *Phys. Rev. D* **100** (2019) 076004 [[arXiv:1903.10197](#)] [INSPIRE].
- [74] F. Ignatov and R.N. Lee, *Charge asymmetry in $e^+e^- \rightarrow \pi^+\pi^-$ process*, *Phys. Lett. B* **833** (2022) 137283 [[arXiv:2204.12235](#)] [INSPIRE].
- [75] G. Colangelo, M. Hoferichter, J. Monnard and J. Ruiz de Elvira, *Radiative corrections to the forward-backward asymmetry in $e^+e^- \rightarrow \pi^+\pi^-$* , *JHEP* **08** (2022) 295 [[arXiv:2207.03495](#)] [INSPIRE].
- [76] J. Monnard, *Radiative corrections for the two-pion contribution to the hadronic vacuum polarization contribution to the muon $g - 2$* , Ph.D. Thesis, Bern University (2020) [<https://boristheses.unibe.ch/2825/>].
- [77] R. Alemany, M. Davier and A. Hoecker, *Improved determination of the hadronic contribution to the muon ($g - 2$) and to $\alpha(M_Z^2)$ using new data from hadronic τ decays*, *Eur. Phys. J. C* **2** (1998) 123 [[hep-ph/9703220](#)] [INSPIRE].
- [78] V. Cirigliano, G. Ecker and H. Neufeld, *Isospin violation and the magnetic moment of the muon*, *Phys. Lett. B* **513** (2001) 361 [[hep-ph/0104267](#)] [INSPIRE].
- [79] V. Cirigliano, G. Ecker and H. Neufeld, *Radiative τ decay and the magnetic moment of the muon*, *JHEP* **08** (2002) 002 [[hep-ph/0207310](#)] [INSPIRE].
- [80] M. Davier et al., *The Discrepancy Between τ and e^+e^- Spectral Functions Revisited and the Consequences for the Muon Magnetic Anomaly*, *Eur. Phys. J. C* **66** (2010) 127 [[arXiv:0906.5443](#)] [INSPIRE].
- [81] F. Jegerlehner and R. Szafron, *$\rho^0 - \gamma$ mixing in the neutral channel pion form factor F_π^e and its role in comparing e^+e^- with τ spectral functions*, *Eur. Phys. J. C* **71** (2011) 1632 [[arXiv:1101.2872](#)] [INSPIRE].

- [82] J.F. de Trocóniz and F.J. Ynduráin, *Precision determination of the pion form-factor and calculation of the muon $g - 2$* , *Phys. Rev. D* **65** (2002) 093001 [[hep-ph/0106025](#)] [[INSPIRE](#)].
- [83] H. Leutwyler, *Electromagnetic form-factor of the pion*, in *Continuous Advances in QCD 2002/ARKADYFEST (honoring the 60th birthday of Prof. Arkady Vainshtein)*, pp. 23–40 (2002) [[DOI](#)] [[hep-ph/0212324](#)] [[INSPIRE](#)].
- [84] G. Colangelo, *Hadronic contributions to a_μ below one GeV*, *Nucl. Phys. B Proc. Suppl.* **131** (2004) 185 [[hep-ph/0312017](#)] [[INSPIRE](#)].
- [85] J.F. de Trocóniz and F.J. Ynduráin, *The Hadronic contributions to the anomalous magnetic moment of the muon*, *Phys. Rev. D* **71** (2005) 073008 [[hep-ph/0402285](#)] [[INSPIRE](#)].
- [86] C. Hanhart, *A New Parameterization for the Pion Vector Form Factor*, *Phys. Lett. B* **715** (2012) 170 [[arXiv:1203.6839](#)] [[INSPIRE](#)].
- [87] B. Ananthanarayan, I. Caprini, D. Das and I. Sentitemsu Imsong, *Two-pion low-energy contribution to the muon $g - 2$ with improved precision from analyticity and unitarity*, *Phys. Rev. D* **89** (2014) 036007 [[arXiv:1312.5849](#)] [[INSPIRE](#)].
- [88] B. Ananthanarayan, I. Caprini, D. Das and I. Sentitemsu Imsong, *Precise determination of the low-energy hadronic contribution to the muon $g - 2$ from analyticity and unitarity: An improved analysis*, *Phys. Rev. D* **93** (2016) 116007 [[arXiv:1605.00202](#)] [[INSPIRE](#)].
- [89] M. Hoferichter, B. Kubis, J. Ruiz de Elvira, H.-W. Hammer and U.-G. Meißner, *On the $\pi\pi$ continuum in the nucleon form factors and the proton radius puzzle*, *Eur. Phys. J. A* **52** (2016) 331 [[arXiv:1609.06722](#)] [[INSPIRE](#)].
- [90] C. Hanhart, S. Holz, B. Kubis, A. Kupść, A. Wirzba and C.W. Xiao, *The branching ratio $\omega \rightarrow \pi^+\pi^-$ revisited*, *Eur. Phys. J. C* **77** (2017) 98 [*Erratum ibid.* **78** (2018) 450] [[arXiv:1611.09359](#)] [[INSPIRE](#)].
- [91] G. Colangelo, M. Hoferichter, B. Kubis, M. Niehus and J. Ruiz de Elvira, *Chiral extrapolation of hadronic vacuum polarization*, *Phys. Lett. B* **825** (2022) 136852 [[arXiv:2110.05493](#)] [[INSPIRE](#)].
- [92] M. Hoferichter, G. Colangelo, M. Procura and P. Stoffer, *Virtual photon-photon scattering*, *Int. J. Mod. Phys. Conf. Ser.* **35** (2014) 1460400 [[arXiv:1309.6877](#)] [[INSPIRE](#)].
- [93] G. Colangelo, M. Hoferichter, M. Procura and P. Stoffer, *Dispersive approach to hadronic light-by-light scattering*, *JHEP* **09** (2014) 091 [[arXiv:1402.7081](#)] [[INSPIRE](#)].
- [94] G. Colangelo, M. Hoferichter, B. Kubis, M. Procura and P. Stoffer, *Towards a data-driven analysis of hadronic light-by-light scattering*, *Phys. Lett. B* **738** (2014) 6 [[arXiv:1408.2517](#)] [[INSPIRE](#)].
- [95] G. Colangelo, M. Hoferichter, M. Procura and P. Stoffer, *Dispersion relation for hadronic light-by-light scattering: theoretical foundations*, *JHEP* **09** (2015) 074 [[arXiv:1506.01386](#)] [[INSPIRE](#)].
- [96] R. Omnès, *On the Solution of certain singular integral equations of quantum field theory*, *Nuovo Cim.* **8** (1958) 316 [[INSPIRE](#)].
- [97] S.M. Roy, *Exact integral equation for pion-pion scattering involving only physical region partial waves*, *Phys. Lett. B* **36** (1971) 353 [[INSPIRE](#)].
- [98] B. Ananthanarayan, G. Colangelo, J. Gasser and H. Leutwyler, *Roy equation analysis of $\pi\pi$ scattering*, *Phys. Rept.* **353** (2001) 207 [[hep-ph/0005297](#)] [[INSPIRE](#)].

- [99] I. Caprini, G. Colangelo and H. Leutwyler, *Regge analysis of the $\pi\pi$ scattering amplitude*, *Eur. Phys. J. C* **72** (2012) 1860 [[arXiv:1111.7160](#)] [[INSPIRE](#)].
- [100] L. Łukaszuk, *A generalization of the Watson theorem*, *Phys. Lett. B* **47** (1973) 51 [[INSPIRE](#)].
- [101] S. Eidelman and L. Łukaszuk, *Pion form-factor phase, $\pi\pi$ elasticity and new e^+e^- data*, *Phys. Lett. B* **582** (2004) 27 [[hep-ph/0311366](#)] [[INSPIRE](#)].
- [102] G. Chanturia, *A two-potential formalism for the pion vector form factor*, *PoS Regio2021* (2022) 030 [[INSPIRE](#)].
- [103] M. Hoferichter, B. Kubis and D. Sakkas, *Extracting the chiral anomaly from $\gamma\pi \rightarrow \pi\pi$* , *Phys. Rev. D* **86** (2012) 116009 [[arXiv:1210.6793](#)] [[INSPIRE](#)].
- [104] S.P. Schneider, B. Kubis and F. Niecknig, *The $\omega \rightarrow \pi^0\gamma^*$ and $\phi \rightarrow \pi^0\gamma^*$ transition form factors in dispersion theory*, *Phys. Rev. D* **86** (2012) 054013 [[arXiv:1206.3098](#)] [[INSPIRE](#)].
- [105] M. Hoferichter, B. Kubis, S. Leupold, F. Niecknig and S.P. Schneider, *Dispersive analysis of the pion transition form factor*, *Eur. Phys. J. C* **74** (2014) 3180 [[arXiv:1410.4691](#)] [[INSPIRE](#)].
- [106] M. Hoferichter, B. Kubis and M. Zanke, *Radiative resonance couplings in $\gamma\pi \rightarrow \pi\pi$* , *Phys. Rev. D* **96** (2017) 114016 [[arXiv:1710.00824](#)] [[INSPIRE](#)].
- [107] M. Niehus, M. Hoferichter and B. Kubis, *The $\gamma\pi \rightarrow \pi\pi$ anomaly from lattice QCD and dispersion relations*, *JHEP* **12** (2021) 038 [[arXiv:2110.11372](#)] [[INSPIRE](#)].
- [108] M. Zanke, M. Hoferichter and B. Kubis, *On the transition form factors of the axial-vector resonance $f_1(1285)$ and its decay into e^+e^-* , *JHEP* **07** (2021) 106 [[arXiv:2103.09829](#)] [[INSPIRE](#)].
- [109] PARTICLE DATA GROUP collaboration, *Review of Particle Physics*, *PTEP* **2020** (2020) 083C01 [[INSPIRE](#)].
- [110] S.I. Dolinsky et al., *Summary of experiments with the neutral detector at the e^+e^- storage ring VEPP-2M*, *Phys. Rept.* **202** (1991) 99 [[INSPIRE](#)].
- [111] B. Moussallam, *Unified dispersive approach to real and virtual photon-photon scattering at low energy*, *Eur. Phys. J. C* **73** (2013) 2539 [[arXiv:1305.3143](#)] [[INSPIRE](#)].
- [112] C.F. Redmer, private communication (2022).
- [113] A.S. Kupich, private communication (2020).
- [114] BESIII collaboration, *Measurement of the $e^+e^- \rightarrow \pi^+\pi^-\pi^0$ Cross Section from 0.7 GeV to 3.0 GeV via Initial-State Radiation*, [arXiv:1912.11208](#) [[INSPIRE](#)].
- [115] BABAR and BABAR collaborations, *Study of the process $e^+e^- \rightarrow \pi^+\pi^-\pi^0$ using initial state radiation with BABAR*, *Phys. Rev. D* **104** (2021) 112003 [[arXiv:2110.00520](#)] [[INSPIRE](#)].
- [116] M.N. Achasov et al., *Study of the process $e^+e^- \rightarrow \pi^+\pi^-\pi^0$ in the energy region \sqrt{s} below 0.98 GeV*, *Phys. Rev. D* **68** (2003) 052006 [[hep-ex/0305049](#)] [[INSPIRE](#)].
- [117] S. Holz, C. Hanhart, M. Hoferichter and B. Kubis, *A dispersive analysis of $\eta' \rightarrow \pi^+\pi^-\gamma$ and $\eta' \rightarrow \ell^+\ell^-\gamma$* , *Eur. Phys. J. C* **82** (2022) 434 [[arXiv:2202.05846](#)] [[INSPIRE](#)].
- [118] BESIII collaboration, *Precision Study of $\eta' \rightarrow \gamma\pi^+\pi^-$ Decay Dynamics*, *Phys. Rev. Lett.* **120** (2018) 242003 [[arXiv:1712.01525](#)] [[INSPIRE](#)].
- [119] C.E. Wolfe and K. Maltman, *Models of Isospin Breaking in the Pion Form Factor: Consequences for the Determination of $\Pi_{\rho\omega}(m_\rho^2)$ and $(g-2)_\mu/2$* , *Phys. Rev. D* **80** (2009) 114024 [[arXiv:0908.2391](#)] [[INSPIRE](#)].

- [120] D. Boito, M. Golterman, K. Maltman and S. Peris, *Evaluation of the three-flavor quark-disconnected contribution to the muon anomalous magnetic moment from experimental data*, *Phys. Rev. D* **105** (2022) 093003 [[arXiv:2203.05070](#)] [[INSPIRE](#)].
- [121] R. Urech, ρ^0 - ω mixing in chiral perturbation theory, *Phys. Lett. B* **355** (1995) 308 [[hep-ph/9504238](#)] [[INSPIRE](#)].
- [122] Y. Aoki et al., *FLAG Review 2021*, [arXiv:2111.09849](#) [[INSPIRE](#)].
- [123] RBC and UKQCD collaborations, *Domain wall QCD with physical quark masses*, *Phys. Rev. D* **93** (2016) 074505 [[arXiv:1411.7017](#)] [[INSPIRE](#)].
- [124] S. Dürr et al., *Lattice QCD at the physical point: light quark masses*, *Phys. Lett. B* **701** (2011) 265 [[arXiv:1011.2403](#)] [[INSPIRE](#)].
- [125] S. Dürr et al., *Lattice QCD at the physical point: Simulation and analysis details*, *JHEP* **08** (2011) 148 [[arXiv:1011.2711](#)] [[INSPIRE](#)].
- [126] MILC collaboration, *MILC results for light pseudoscalars*, *PoS CD09* (2009) 007 [[arXiv:0910.2966](#)] [[INSPIRE](#)].
- [127] ALPHA collaboration, *Light quark masses in $N_f = 2 + 1$ lattice QCD with Wilson fermions*, *Eur. Phys. J. C* **80** (2020) 169 [[arXiv:1911.08025](#)] [[INSPIRE](#)].
- [128] Z. Fodor et al., *Up and down quark masses and corrections to Dashen's theorem from lattice QCD and quenched QED*, *Phys. Rev. Lett.* **117** (2016) 082001 [[arXiv:1604.07112](#)] [[INSPIRE](#)].
- [129] EUROPEAN TWISTED MASS collaboration, *Up, down, strange and charm quark masses with $N_f = 2 + 1 + 1$ twisted mass lattice QCD*, *Nucl. Phys. B* **887** (2014) 19 [[arXiv:1403.4504](#)] [[INSPIRE](#)].
- [130] A. Bazavov et al., *B- and D-meson leptonic decay constants from four-flavor lattice QCD*, *Phys. Rev. D* **98** (2018) 074512 [[arXiv:1712.09262](#)] [[INSPIRE](#)].
- [131] FERMILAB LATTICE and MILC collaborations, *Charmed and Light Pseudoscalar Meson Decay Constants from Four-Flavor Lattice QCD with Physical Light Quarks*, *Phys. Rev. D* **90** (2014) 074509 [[arXiv:1407.3772](#)] [[INSPIRE](#)].
- [132] D. Giusti et al., *Leading isospin-breaking corrections to pion, kaon and charmed-meson masses with Twisted-Mass fermions*, *Phys. Rev. D* **95** (2017) 114504 [[arXiv:1704.06561](#)] [[INSPIRE](#)].
- [133] MILC collaboration, *Lattice computation of the electromagnetic contributions to kaon and pion masses*, *Phys. Rev. D* **99** (2019) 034503 [[arXiv:1807.05556](#)] [[INSPIRE](#)].
- [134] J. Bijnens and P. Gosdzinsky, *Electromagnetic contributions to vector meson masses and mixings*, *Phys. Lett. B* **388** (1996) 203 [[hep-ph/9607462](#)] [[INSPIRE](#)].
- [135] J. Bijnens, P. Gosdzinsky and P. Talavera, *Vector meson masses in chiral perturbation theory*, *Nucl. Phys. B* **501** (1997) 495 [[hep-ph/9704212](#)] [[INSPIRE](#)].
- [136] C.L. James, R. Lewis and K. Maltman, *ChPT estimate of the strong-isospin-breaking contribution to the anomalous magnetic moment of the muon*, *Phys. Rev. D* **105** (2022) 053010 [[arXiv:2109.13729](#)] [[INSPIRE](#)].
- [137] CMD-2 collaboration, *Study of the processes $e^+e^- \rightarrow \eta\gamma$, $\pi^0\gamma \rightarrow 3\gamma$ in the c.m. energy range 600–1380 MeV at CMD-2*, *Phys. Lett. B* **605** (2005) 26 [[hep-ex/0409030](#)] [[INSPIRE](#)].
- [138] CRYSTAL BARREL collaboration, *Antiproton-proton annihilation at rest into $\omega\pi^0\pi^0$* , *Phys. Lett. B* **311** (1993) 362 [[INSPIRE](#)].