

New physics in $b \rightarrow s\mu^+\mu^-$: CP-violating observables

Ashutosh Kumar Alok,^a Alakabha Datta,^b Amol Dighe,^c Murugeswaran Duraisamy,^b Diptimoy Ghosh^c and David London^a

^a*Physique des Particules, Université de Montréal,
C.P. 6128, succ. centre-ville, Montréal, QC - H3C 3J7, Canada*

^b*Department of Physics and Astronomy, University of Mississippi,
108 Lewis Hall, Oxford, MS 38677-1848, U.S.A.*

^c*Tata Institute of Fundamental Research,
Homi Bhabha Road, Mumbai 400005, India*

E-mail: alok@lps.umontreal.ca, datta@phy.olemiss.edu,
amol@theory.tifr.res.in, duraism@phy.olemiss.edu,
diptimoyghosh@theory.tifr.res.in, london@lps.umontreal.ca

ABSTRACT: We perform a comprehensive study of the impact of new-physics operators with different Lorentz structures on CP-violating observables involving the $b \rightarrow s\mu^+\mu^-$ transition. We examine the effects of new vector-axial vector (VA), scalar-pseudoscalar (SP) and tensor (T) interactions on the CP asymmetries in the branching ratios and forward-backward asymmetries of $\bar{B}_s^0 \rightarrow \mu^+\mu^-$, $\bar{B}_d^0 \rightarrow X_s\mu^+\mu^-$, $\bar{B}_s^0 \rightarrow \mu^+\mu^-\gamma$, $\bar{B}_d^0 \rightarrow \bar{K}\mu^+\mu^-$, and $\bar{B}_d^0 \rightarrow \bar{K}^*\mu^+\mu^-$. In $\bar{B}_d^0 \rightarrow \bar{K}^*\mu^+\mu^-$, we also explore the direct CP asymmetries in the longitudinal polarization fraction f_L and the angular asymmetries $A_T^{(2)}$ and A_{LT} , as well as the triple-product CP asymmetries $A_T^{(im)}$ and $A_{LT}^{(im)}$. We find that, in almost all cases, the CP-violating observables are sensitive only to new physics which involves VA operators. The VA new physics may therefore be unambiguously identified by a combined analysis of future measurements of these CP-violating observables.

KEYWORDS: Beyond Standard Model, B-Physics

ARXIV EPRINT: [1103.5344](https://arxiv.org/abs/1103.5344)

Contents

1	Introduction	1
2	$b \rightarrow s\mu^+\mu^-$ operators	3
2.1	Effective Hamiltonian in the SM and with NP	3
2.2	Constraints on NP couplings	3
2.3	CP-violating effects	5
3	$\bar{B}_s^0 \rightarrow \mu^+\mu^-$	6
4	$\bar{B}_d^0 \rightarrow X_s\mu^+\mu^-$	7
5	$\bar{B}_s^0 \rightarrow \mu^+\mu^-\gamma$	8
6	$\bar{B}_d^0 \rightarrow \bar{K}\mu^+\mu^-$	9
7	$\bar{B}_d^0 \rightarrow \bar{K}^*\mu^+\mu^-$	11
7.1	Direct CP asymmetries in the DBR and A_{FB}	12
7.2	Direct CP asymmetry in the polarization fraction f_L	13
7.3	Direct CP asymmetries in the angular asymmetries $A_T^{(2)}$ and A_{LT}	14
7.4	CP-violating triple-product asymmetries	16
8	Discussion and summary	18

1 Introduction

The B factories have taken us to the luminosity frontier with more than a billion B^+/B_d mesons, and the Tevatron experiments have provided us with invaluable data on B_s mesons. We have now entered the precision era of B physics. The Standard Model (SM) has been successful in explaining most of the data to date. However, this is now the time to look forward to precision tests, with the ATLAS and CMS experiments already running, the LHCb expected to start recording data soon, and the Super- B factories on their way. One can now be ambitious and not only look for new-physics (NP) effects, but also try to identify the kind of NP involved.

Though there is no unambiguous signal of NP so far in all of the B decays we have observed, some possible hints of NP have recently surfaced in modes involving $b \rightarrow s$ transitions. These include measurements of CP-averaged quantities such as the large transverse polarization in $B \rightarrow \phi K^*$ [1, 2], and the anomalous forward-backward asymmetry in $B \rightarrow K^*\mu^+\mu^-$ [3-5]. There are also measurements of CP-violating quantities such as the difference between the mixing-induced CP asymmetries seen in $b \rightarrow s$ penguin decays and

in $B_d \rightarrow J/\psi K_S$ [6–8], the large CP asymmetry in $B_s \rightarrow J/\psi \phi$ [9], and the anomalous CP asymmetry in like-sign dimuon signals [10].

In the companion paper [11], we performed a general analysis with all possible Lorentz structures of NP in the transition $b \rightarrow s\mu^+\mu^-$. We included NP vector-axial vector (VA), scalar-pseudoscalar (SP), and tensor (T) $b \rightarrow s\mu^+\mu^-$ operators, and explored their possible effects on the decays $\bar{B}_s^0 \rightarrow \mu^+\mu^-$, $\bar{B}_d^0 \rightarrow X_s\mu^+\mu^-$, $\bar{B}_s^0 \rightarrow \mu^+\mu^-\gamma$, $\bar{B}_d^0 \rightarrow \bar{K}\mu^+\mu^-$, and $\bar{B}_d^0 \rightarrow \bar{K}^*\mu^+\mu^-$. We focused on CP-conserving observables such as differential branching ratios, forward-backward asymmetries, polarization fractions, and the asymmetries $A_T^{(2)}$, A_{LT} in $\bar{B}_d^0 \rightarrow \bar{K}^*\mu^+\mu^-$. Because we only considered CP-conserving observables, all the NP couplings were taken to be real. We computed the effects of all NP operators, individually and in all combinations, on these observables.

The CP-violating observables in various $b \rightarrow s\mu^+\mu^-$ decays in the SM as well as in some NP models have been studied in refs. [12–27]. In this paper, we explore the CP-violating quantities that may be measured in the same decay modes by allowing the new couplings to be complex. The introduction of complex couplings has two effects. First, some quantities which were taken to be CP-conserving above now display CP-violation, i.e. the quantities take different values in the CP-conjugate decays. The difference between the value of a measurement in a decay and in its CP-conjugate counterpart is then a CP-violating observable. Second, new observables appear which vanish in the CP-conserving limit. (These were not considered in ref. [11] for this reason.) These essentially correspond to the CP-violating triple-product asymmetries $A_T^{(im)}$ and $A_{LT}^{(im)}$ in $\bar{B}_d^0 \rightarrow \bar{K}^*\mu^+\mu^-$, which may be obtained from the angular distribution in this decay. Our goal is to identify those quantities for which there may be large effects due to the presence of NP. In such cases, we try to find salient features of the effects of NP, which may help us identify the Lorentz structure of the NP involved.

Here we have taken the NP to be present only in the effective $b \rightarrow s\mu^+\mu^-$ operator. While this can, in principle, contribute to CP violation in B_d - \bar{B}_d and B_s - \bar{B}_s mixing, it is a higher-order effect, and hence negligible compared to the SM contribution. We therefore neglect mixing-induced (indirect) CP violation in this work, and focus only on CP violation in the decay. In the SM, such CP violation is expected to be close to zero in $b \rightarrow s$ transitions. A naive estimate indicates that this asymmetry will be $\sim 10^{-3}$ [20, 23], but even if next-to-leading order (NLO) QCD corrections and hadronic uncertainties are included, it is observed that the CP asymmetry will not exceed 1% [24, 25, 28]. Thus, if a large CP-violating effect, more than a few percent, is observed in any of the $b \rightarrow s\mu^+\mu^-$ channels, this will therefore be a clear signature of NP. In this paper, we go further and explore the extent to which the Lorentz structure of NP can be ascertained from the CP-violating measurements.

The paper is organized as follows. We begin in section 2 by describing the effective Hamiltonian with NP operators and new couplings. Although the formalism is the same as that used in ref. [11], the constraints on the NP couplings are now more relaxed since the couplings are allowed to be complex. We also present an overview of the types of CP-violating observables which are examined. In section 3 we note that there are essentially no measurable CP-violating quantities in the mode $\bar{B}_s^0 \rightarrow \mu^+\mu^-$. We then consider the decays $\bar{B}_d^0 \rightarrow X_s\mu^+\mu^-$ (section 4), $\bar{B}_s^0 \rightarrow \mu^+\mu^-\gamma$ (section 5), and $\bar{B}_d^0 \rightarrow \bar{K}\mu^+\mu^-$ (section 6). In

these sections we examine the same observables as in ref. [11], this time looking at the asymmetries between these processes and their CP-conjugates. In section 7, we study the CP asymmetries in $\bar{B}_d^0 \rightarrow \bar{K}^* \mu^+ \mu^-$ for the observables considered in ref. [11], and in addition we explore new observables that vanish in the CP-conserving limit (triple products). We summarize our findings in section 8 and discuss their implications.

2 $b \rightarrow s \mu^+ \mu^-$ operators

2.1 Effective Hamiltonian in the SM and with NP

Our formalism is identical to that used in ref. [11]. We repeat it here briefly for the sake of completeness. Within the SM, the effective Hamiltonian for the quark-level transition $b \rightarrow s \mu^+ \mu^-$ is

$$\begin{aligned} \mathcal{H}_{\text{eff}}^{SM} = & -\frac{4G_F}{\sqrt{2}} V_{ts}^* V_{tb} \left\{ \sum_{i=1}^6 C_i(\mu) \mathcal{O}_i(\mu) + C_7 \frac{e}{16\pi^2} [\bar{s} \sigma_{\mu\nu} (m_s P_L + m_b P_R) b] F^{\mu\nu} \right. \\ & \left. + C_9 \frac{\alpha_{em}}{4\pi} (\bar{s} \gamma^\mu P_L b) \bar{\mu} \gamma_\mu \mu + C_{10} \frac{\alpha_{em}}{4\pi} (\bar{s} \gamma^\mu P_L b) \bar{\mu} \gamma_\mu \gamma_5 \mu \right\} + h.c., \end{aligned} \quad (2.1)$$

where $P_{L,R} = (1 \mp \gamma_5)/2$. The operators \mathcal{O}_i ($i = 1, \dots, 6$) correspond to the P_i in ref. [29], and $m_b = m_b(\mu)$ is the running b -quark mass in the $\overline{\text{MS}}$ scheme. We use the SM Wilson coefficients (C_i) as given in ref. [25].

The effective Hamiltonian in the presence of NP is

$$\mathcal{H}_{\text{eff}}(b \rightarrow s \mu^+ \mu^-) = \mathcal{H}_{\text{eff}}^{SM} + \mathcal{H}_{\text{eff}}^{VA} + \mathcal{H}_{\text{eff}}^{SP} + \mathcal{H}_{\text{eff}}^T + h.c., \quad (2.2)$$

where

$$\begin{aligned} \mathcal{H}_{\text{eff}}^{VA} = & -\frac{4G_F}{\sqrt{2}} \frac{\alpha_{em}}{4\pi} V_{ts}^* V_{tb} \left\{ R_V (\bar{s} \gamma^\mu P_L b) \bar{\mu} \gamma_\mu \mu + R_A (\bar{s} \gamma^\mu P_L b) \bar{\mu} \gamma_\mu \gamma_5 \mu \right. \\ & \left. + R'_V (\bar{s} \gamma^\mu P_R b) \bar{\mu} \gamma_\mu \mu + R'_A (\bar{s} \gamma^\mu P_R b) \bar{\mu} \gamma_\mu \gamma_5 \mu \right\}, \end{aligned} \quad (2.3)$$

$$\begin{aligned} \mathcal{H}_{\text{eff}}^{SP} = & -\frac{4G_F}{\sqrt{2}} \frac{\alpha_{em}}{4\pi} V_{ts}^* V_{tb} \left\{ R_S (\bar{s} P_R b) \bar{\mu} \mu + R_P (\bar{s} P_R b) \bar{\mu} \gamma_5 \mu \right. \\ & \left. + R'_S (\bar{s} P_L b) \bar{\mu} \mu + R'_P (\bar{s} P_L b) \bar{\mu} \gamma_5 \mu \right\}, \end{aligned} \quad (2.4)$$

$$\mathcal{H}_{\text{eff}}^T = -\frac{4G_F}{\sqrt{2}} \frac{\alpha_{em}}{4\pi} V_{ts}^* V_{tb} \left\{ C_T (\bar{s} \sigma_{\mu\nu} b) \bar{\mu} \sigma^{\mu\nu} \mu + i C_{TE} (\bar{s} \sigma_{\mu\nu} b) \bar{\mu} \sigma_{\alpha\beta} \mu \epsilon^{\mu\nu\alpha\beta} \right\} \quad (2.5)$$

are the new contributions. Here, $R_V, R_A, R'_V, R'_A, R_S, R_P, R'_S, R'_P, C_T$ and C_{TE} are the NP effective couplings. In our numerical analysis in this paper, we take all NP couplings to be complex. As in ref. [11], we do not include NP through the $O_7 = \bar{s} \sigma^{\alpha\beta} P_R b F_{\alpha\beta}$ operator or its chirally-flipped counterpart $O'_7 = \bar{s} \sigma^{\alpha\beta} P_L b F_{\alpha\beta}$.

2.2 Constraints on NP couplings

The constraints on the NP couplings in $b \rightarrow s \mu^+ \mu^-$ come mainly from the upper bound on the branching ratio $B(\bar{B}_s^0 \rightarrow \mu^+ \mu^-)$ and the measurements of the total branching ratios

$B(\bar{B}_d^0 \rightarrow X_s \mu^+ \mu^-)$ and $B(\bar{B}_d^0 \rightarrow \bar{K} \mu^+ \mu^-)$ [30–32]:

$$B(\bar{B}_s^0 \rightarrow \mu^+ \mu^-) < 4.70 \times 10^{-8} \quad (90\% \text{ C.L.}) , \quad (2.6)$$

$$B(\bar{B}_d^0 \rightarrow X_s \mu^+ \mu^-) = \begin{cases} (1.60 \pm 0.50) \times 10^{-6} & (\text{low } q^2) \\ (0.44 \pm 0.12) \times 10^{-6} & (\text{high } q^2) \end{cases} , \quad (2.7)$$

$$B(\bar{B}_d^0 \rightarrow \bar{K} \mu^+ \mu^-) = (4.5_{-1.0}^{+1.2}) \times 10^{-7} , \quad (2.8)$$

where the low- q^2 and high- q^2 regions correspond to $1 \text{ GeV}^2 \leq q^2 \leq 6 \text{ GeV}^2$ and $q^2 \geq 14.4 \text{ GeV}^2$, respectively. Here q^2 is the invariant mass squared of the two muons.

We consider all the NP couplings R_i to be complex and parametrize them as

$$R_i = |R_i| e^{i\phi_{R_i}} , \quad (2.9)$$

where $i = V, A, S, P, T, TE$ and $-\pi \leq \phi_{R_i} \leq \pi$. The bounds on these couplings will in general depend on which operators are present. While we take the correlations in these constraints into account in our numerical calculations, for the sake of simplicity we only give the bounds when the NP operators (VA, SP, T) are present individually.

If the only NP couplings present are $R_{V,A}$, we obtain

$$\frac{|\text{Re}(R_V) + 2.8|^2}{(6.3)^2} + \frac{|\text{Im}(R_V)|^2}{(6.0)^2} \lesssim 1.0 , \quad \frac{|\text{Re}(R_A) - 4.1|^2}{(6.1)^2} + \frac{|\text{Im}(R_A)|^2}{(6.0)^2} \lesssim 1.0 . \quad (2.10)$$

If the only NP couplings present are $R'_{V,A}$, the constraints are

$$\frac{|\text{Re}(R'_V)|^2}{(3.5)^2} + \frac{|\text{Im}(R'_V)|^2}{(4.0)^2} \lesssim 1.0 , \quad \frac{|\text{Re}(R'_A)|^2}{(3.5)^2} + \frac{|\text{Im}(R'_A)|^2}{(4.0)^2} \lesssim 1.0 . \quad (2.11)$$

For the SP operators, the present upper bound on $B(\bar{B}_s^0 \rightarrow \mu^+ \mu^-)$ provides the limit

$$|R_S - R'_S|^2 + |R_P - R'_P|^2 \lesssim 0.44 . \quad (2.12)$$

This constitutes a severe constraint on the NP couplings if only $R_{S,P}$ or $R'_{S,P}$ are present. However, if both types of operators are present, these bounds can be evaded due to cancellations between the $R_{S,P}$ and $R'_{S,P}$. In that case, $B(\bar{B}_d^0 \rightarrow X_s \mu^+ \mu^-)$ and $B(\bar{B}_d^0 \rightarrow \bar{K} \mu^+ \mu^-)$ can still bound these couplings. The stronger bound is obtained from the measurement of the latter quantity, which yields

$$|R_S|^2 + |R_P|^2 \lesssim 9 , \quad R_S \approx R'_S , \quad R_P \approx R'_P . \quad (2.13)$$

Finally, the constraints on the NP tensor operators come entirely from $B(\bar{B}_d^0 \rightarrow X_s \mu^+ \mu^-)$. When only the T operators are present,

$$|C_T|^2 + 4|C_{TE}|^2 \lesssim 1.0 . \quad (2.14)$$

The constraints are not affected significantly if more than one type (VA, SP or T) of NP operators is present simultaneously.

2.3 CP-violating effects

All CP-violating effects are due to the interference of (at least) two amplitudes with a relative weak phase. In principle, there can be three types of interference: SM-SM, SM-NP, NP-NP. In the SM, all contributions to the $b \rightarrow s\mu^+\mu^-$ modes are proportional to the Cabibbo-Kobayashi-Maskawa (CKM) factors $V_{tb}^*V_{ts}$, $V_{cb}^*V_{cs}$, or $V_{ub}^*V_{us}$. The term $V_{cb}^*V_{cs}$ can be eliminated in terms of the other two using the unitarity of the CKM matrix. Furthermore, although $V_{ub}^*V_{us}$ has a large weak phase, its magnitude is greatly suppressed relative to that of $V_{tb}^*V_{ts}$. Thus, to a good approximation, all nonzero SM contributions have the same weak phase, and so all CP-violating effects are predicted to be tiny in the SM.

There are two types of CP violation. The first is direct CP-violating asymmetries. Suppose that a particular \bar{B} decay has two contributing amplitudes: $i\mathcal{M}(\bar{B} \text{ decay}) = \mathcal{A}_1 + \mathcal{A}_2$. Each amplitude has both a weak and a strong phase. The matrix element $i\overline{\mathcal{M}}$ for the CP-conjugate decay is the same as $i\mathcal{M}$, except that the weak phases change signs. CP violation is indicated by a nonzero value of $|\mathcal{M}|^2 - |\overline{\mathcal{M}}|^2$. It is straightforward to show that this is proportional to $\sin\phi_w \sin\delta$, where ϕ_w and δ are, respectively, the relative weak and strong phases between \mathcal{A}_1 and \mathcal{A}_2 . Direct CP-violating asymmetries therefore require that the interfering amplitudes have both a nonzero relative weak and strong phase.

The second type of CP violation is triple-product (TP) asymmetries. Suppose that the matrix element for the \bar{B} decay takes the form $i\mathcal{M}(\bar{B} \text{ decay}) = \mathcal{A}_1 + i\mathcal{A}_2\epsilon_{\mu\nu\rho\sigma}p_B^\mu v_1^\nu v_2^\rho v_3^\sigma$, where the v_i are spins or momenta of the final-state particles. The difference $|\mathcal{M}|^2 - |\overline{\mathcal{M}}|^2$ is proportional to $m_{\bar{B}}\vec{v}_1 \cdot (\vec{v}_2 \times \vec{v}_3) \sin\phi_w \cos\delta$. By measuring the TP $\vec{v}_1 \cdot (\vec{v}_2 \times \vec{v}_3)$ in both \bar{B} and B decays, the TP asymmetry can be obtained. Note that the measurement of a nonzero TP in the \bar{B} decay alone is not sufficient to establish CP violation, i.e. it does not necessarily imply a nonzero weak phase. A fake, CP-conserving TP can be produced if \mathcal{A}_1 and \mathcal{A}_2 have a relative strong phase. It is only by measuring the difference of TPs in \bar{B} and B decays that the fake TP can be eliminated and a true, CP-violating signal produced [33].

Let us first turn to direct CP violation, which requires both a relative weak and strong phase between two interfering amplitudes. Now, strong phases are generated through the rescattering of the operators in the effective Hamiltonian. The NP strong phases involve only the (constrained) NP operators, and are therefore small [34]. Thus, direct CP asymmetries can never arise from NP-NP interference.

On the other hand, the SM strong phase is not so small. It is generated because the Wilson coefficient C_9^{eff} , which gets a contribution from a $c\bar{c}$ quark loop, has an imaginary piece. (C_9^{eff} also gets a contribution from a $u\bar{u}$ quark loop. But this is proportional to $V_{ub}^*V_{us}$, and hence negligible.) The quantity C_9^{eff} can be written as [25]

$$\begin{aligned}
 C_9^{\text{eff}} &= C_9(m_b) + h(z, \hat{m}_c) \left(\frac{4}{3}C_1 + C_2 + 6C_3 + 60C_5 \right) \\
 &\quad - \frac{1}{2}h(z, \hat{m}_b) \left(7C_3 + \frac{4}{3}C_4 + 76C_5 + \frac{64}{3}C_6 \right) \\
 &\quad - \frac{1}{2}h(z, 0) \left(C_3 + \frac{4}{3}C_4 + 16C_5 + \frac{64}{3}C_6 \right) + \frac{4}{3}C_3 + \frac{64}{9}C_5 + \frac{64}{27}C_6.
 \end{aligned}
 \tag{2.15}$$

Here $z \equiv q^2/m_b^2$, and $\hat{m}_q \equiv m_q/m_b$ for all quarks q . The function $h(z, \hat{m})$ represents the one-loop correction to the four-quark operators O_1 - O_6 and is given by [23, 25, 35]

$$h(z, \hat{m}) = -\frac{8}{9} \ln \frac{m_b}{\mu_b} - \frac{8}{9} \ln \hat{m} + \frac{8}{27} + \frac{4}{9}x \tag{2.16}$$

$$-\frac{2}{9}(2+x)|1-x|^{1/2} \begin{cases} \left(\ln \left| \frac{\sqrt{1-x+1}}{\sqrt{1-x-1}} \right| - i\pi \right), & \text{for } x \leq 1, \\ 2 \arctan \frac{1}{\sqrt{x-1}}, & \text{for } x > 1, \end{cases}$$

where $x \equiv 4\hat{m}^2/z$. Thus, a nontrivial strong phase is generated when $z \geq 4\hat{m}^2$. This leads to the complex nature of C_9^{eff} in the SM. For example, typical values of C_9^{eff} in the low- and high- q^2 regions are $C_9^{\text{eff}}(m_b) = 4.75+0.09i$ ($z = 0.1$), $C_9^{\text{eff}}(m_b) = 4.76+0.88i$ ($z = 0.7$). C_9^{eff} therefore has a nontrivial imaginary component, which implies that direct CP asymmetries can arise due to SM-NP interference. Since the SM operator (C_9^{eff}) is of VA type, the NP operator must also be VA in order to generate a significant direct CP asymmetry. Other NP operators can also interfere with the SM, but the effect is suppressed by m_μ/m_b , and hence very small. Note that, although this argument has used the total decay rate for illustration, we could have used (almost) any observable which is related to the square of the matrix element. This includes the differential branching ratio, forward-backward asymmetry, polarization asymmetries, etc.

The TP asymmetries, on the other hand, do not need a difference in strong phases between two amplitudes. Indeed, they are proportional to $\cos \delta$, though they do require a weak-phase difference. This means that a TP asymmetry can be produced by either SM-NP or NP-NP interference. Given that all SM operators are of VA type, the NP must also be VA if SM-NP interference is the reason for the TP. On the other hand, if NP-NP interference is involved, this can arise due to new SP and T operators (other NP-NP interference are possible, but the effects are suppressed by m_μ/m_b).

In this paper, we explore both sources of CP asymmetries, direct CP violation and TPs. While we have checked the effects of SP and T NP operators on all the observables, we find them to be insignificant in most places (as expected from the arguments above), and we will mention them only during the discussion of TP asymmetries, where, in principle, they may play a significant role.

3 $\bar{B}_s^0 \rightarrow \mu^+ \mu^-$

We begin by considering the direct CP asymmetry in $\bar{B}_s^0 \rightarrow \mu^+ \mu^-$. Helicity conservation in the decay of B_s or \bar{B}_s implies that the only final states can be $\mu_L^+ \mu_L^-$ or $\mu_R^+ \mu_R^-$, which are CP conjugates. The only CP-violating observables that can be constructed are then

$$A_{CP}^{RL}(t) \equiv \frac{B(\bar{B}_s^0(t) \rightarrow \mu_R^+ \mu_R^-) - B(B_s^0(t) \rightarrow \mu_L^+ \mu_L^-)}{B(\bar{B}_s^0(t) \rightarrow \mu_R^+ \mu_R^-) + B(B_s^0(t) \rightarrow \mu_L^+ \mu_L^-)},$$

$$A_{CP}^{LR}(t) \equiv \frac{B(\bar{B}_s^0(t) \rightarrow \mu_L^+ \mu_L^-) - B(B_s^0(t) \rightarrow \mu_R^+ \mu_R^-)}{B(\bar{B}_s^0(t) \rightarrow \mu_L^+ \mu_L^-) + B(B_s^0(t) \rightarrow \mu_R^+ \mu_R^-)}. \tag{3.1}$$

The CP asymmetry in the longitudinal polarization fraction A_{LP} may be written in terms of these two CP asymmetries. The measurement of either of these CP asymmetries requires

the measurement of muon polarization, which will be an impossible task for the upcoming experiments [11]. And even if this were doable, the lack of any sources for different strong phases in the two CP-conjugate final states implies that the direct CP asymmetry would vanish even with NP. We therefore do not study CP violation in $\bar{B}_s^0 \rightarrow \mu^+\mu^-$.

4 $\bar{B}_d^0 \rightarrow X_s\mu^+\mu^-$

A model-independent analysis of the CP asymmetry in the differential branching ratio (DBR) of $\bar{B}_d^0 \rightarrow X_s\mu^+\mu^-$ was previously carried out in ref. [14]. There, the CP asymmetry in the DBR was predicted for some specific values of the NP couplings. However, no experimental constraints on the parameters were used. In this paper we study the CP asymmetry in the DBR, taking into account the constraints from the present measurements of other related observables. Moreover, in addition to the CP asymmetry in the DBR, we also study the CP asymmetry in the forward-backward asymmetry.

The CP asymmetry in DBR of $\bar{B}_d^0 \rightarrow X_s\mu^+\mu^-$ is defined as

$$A_{\text{CP}}(q^2) = \frac{(dB/dz) - (d\bar{B}/dz)}{(dB/dz) + (d\bar{B}/dz)}, \quad (4.1)$$

where $z \equiv q^2/m_b^2$, and dB/dz and $d\bar{B}/dz$ are the DBRs of $\bar{B}_d^0 \rightarrow X_s\mu^+\mu^-$ and its CP-conjugate $B_d^0 \rightarrow X_s\mu^+\mu^-$, respectively. The expression for (dB/dz) has been given in ref. [11].

The CP asymmetry in the forward-backward asymmetry A_{FB} is defined as

$$\Delta A_{FB}(q^2) \equiv A_{FB}(q^2) - \bar{A}_{FB}(q^2), \quad (4.2)$$

where the definition of A_{FB} is given in ref. [11], and \bar{A}_{FB} is the analogous quantity for the CP-conjugate decay. Note that while the relevant angle θ in $\bar{B}_d^0 \rightarrow X_s\mu^+\mu^-$ is defined relative to the direction of μ^+ , for the CP-conjugate decay one should define θ in relation to the direction of μ^- , and similarly for A_{FB} in other $b \rightarrow s\mu^+\mu^-$ decay modes below.

Figure 1 shows $A_{\text{CP}}(q^2)$ and $\Delta A_{FB}(q^2)$ for $\bar{B}_d^0 \rightarrow X_s\mu^+\mu^-$ in the presence of new VA couplings. We make the following observations:

- When only $R_{V,A}$ couplings are present, $A_{\text{CP}}(q^2)$ can be enhanced up to 6% at low q^2 . On the other hand, its value at high q^2 can be as high as 12%. $A_{\text{CP}}(q^2)$ can have either sign at both low and high q^2 . At high q^2 , the magnitude of $A_{\text{CP}}(q^2)$ is almost independent of q^2 .
- When only $R'_{V,A}$ couplings are present, $A_{\text{CP}}(q^2)$ cannot be enhanced above the SM value. This is because $R'_{V,A}$ couplings do not contribute to the numerator of $A_{\text{CP}}(q^2)$ in eq. (4.1). They can only affect the DBR, which may be enhanced by up to 50%, thus decreasing $A_{\text{CP}}(q^2)$.
- In the presence of $R_{V,A}$ couplings, ΔA_{FB} can be enhanced up to 3% at low q^2 . At high q^2 , the enhancement can be up to 12%. The impact of $R'_{V,A}$ couplings is negligible ($< 1\%$).




Figure 1. The left (right) panels of the figure show $A_{CP}(q^2)$ and ΔA_{FB} for $\bar{B}_d^0 \rightarrow X_s \mu^+ \mu^-$ in the low- q^2 (high- q^2) region, in the scenario where only (R_V, R_A) couplings are present. The green line corresponds to the SM prediction. The other lines show predictions for some representative values of the NP parameters (R_V, R_A) . For example, the blue curve in the low- q^2 and high- q^2 regions for the A_{CP} plot corresponds to $(5.68e^{i2.13}, 2.64e^{-i0.04})$ and $(4.29e^{i1.68}, 4.15e^{-i0.26})$, respectively, whereas the blue curve in the low- q^2 and high- q^2 regions for the ΔA_{FB} plot corresponds to $(1.80e^{i2.91}, 5.45e^{i0.90})$ and $(1.69e^{-i3.08}, 6.83e^{-i0.91})$, respectively.

5 $\bar{B}_s^0 \rightarrow \mu^+ \mu^- \gamma$

Although $\bar{B}_s^0 \rightarrow \mu^+ \mu^- \gamma$ requires the emission of an additional photon as compared to $\bar{B}_s^0 \rightarrow \mu^+ \mu^-$, which suppresses the branching ratio (BR) by a factor of α_{em} , the photon emission also frees it from helicity suppression, making its BR much larger than $\bar{B}_s^0 \rightarrow \mu^+ \mu^-$. The SM prediction for the BR in the range $q^2 \leq 9.5 \text{ GeV}^2$ and $q^2 \geq 15.9 \text{ GeV}^2$ is $\approx 18.9 \times 10^{-9}$ [36]. As argued in ref. [11], if we choose $2 \text{ GeV}^2 \leq q^2 \leq 6 \text{ GeV}^2$ and $14.4 \text{ GeV}^2 \leq q^2 \leq 25 \text{ GeV}^2$ as the low- q^2 and high- q^2 regions, respectively, then the dominating contribution comes from the diagrams in which the final-state photon is emitted either from the b or the s quark, and the $\bar{B}_s^0 \rightarrow \mu^+ \mu^- \gamma$ decay is governed by the same $b \rightarrow s \mu^+ \mu^-$ effective Hamiltonian as the other decays considered in this paper.

The CP asymmetry in $\bar{B}_s^0 \rightarrow \mu^+ \mu^- \gamma$ is given in eq. (4.1), where dB/dx_γ and $d\bar{B}/dx_\gamma$ are the DBRs of $\bar{B}_s^0 \rightarrow \mu^+ \mu^- \gamma$ and its CP-conjugate $B_s^0 \rightarrow \mu^+ \mu^- \gamma$, respectively. The expression for (dB/dx_γ) has been given in ref. [11]. The CP asymmetry in A_{FB} is given in eq. (4.2), where the definition of A_{FB} is given in ref. [11], and \bar{A}_{FB} is the analogous quantity for the CP-conjugate decay.

The CP asymmetry in the DBR of $B_s \rightarrow \mu \mu \gamma$ was studied in refs. [17, 18], albeit only for the new-physics cases where $C_7 = -C_7^{\text{SM}}, C_9 = -C_9^{\text{SM}}$ and $C_{10} = -C_{10}^{\text{SM}}$, and naturally

only for VA operators. Here, we include a complete discussion of the possible enhancement of the asymmetry for all allowed values of C_9 and C_{10} , and in the presence of SP and T operators. In addition, we study the CP-violating asymmetry in A_{FB} , which also turns out to give possibly significant NP signals.

Figure 2 shows $A_{CP}(q^2)$ and $\Delta A_{FB}(q^2)$ for $\bar{B}_s^0 \rightarrow \mu^+ \mu^- \gamma$ in the presence of new VA couplings. We make the following observations:

- When only $R_{V,A}$ couplings are present, at low q^2 the magnitude of $A_{CP}(q^2)$ can be enhanced up to 30% at certain q^2 values. At high q^2 , the magnitude of $A_{CP}(q^2)$ is almost independent of q^2 , and can be enhanced to about 13%. The asymmetry can have either sign at both low and high q^2 .
- When only $R'_{V,A}$ couplings are present, $A_{CP}(q^2)$ cannot be enhanced in magnitude to more than 1.5% at low q^2 , or more than 3% at high q^2 . The detection of NP of this kind is therefore expected to be very difficult in this channel. When both primed and unprimed VA couplings are present, the results are the same as those obtained with only $R_{V,A}$ couplings.
- The behaviour of $\Delta A_{FB}(q^2)$ is similar to that of $A_{CP}(q^2)$. This quantity can be enhanced up to 40% for some values in the low- q^2 region. It can be as high as 18% throughout the high- q^2 region. The impact of $R'_{V,A}$ couplings is negligible ($< 1\%$).

The new VA operators can therefore enhance the asymmetries $A_{CP}(q^2)$ and $\Delta A_{FB}(q^2)$ in $\bar{B}_s^0 \rightarrow \mu^+ \mu^- \gamma$ to $\sim 10\%$ throughout the q^2 region. For a branching ratio of $O(2 \times 10^{-8})$, a measurement of a CP asymmetry of 10% at the 3σ level would require $\sim 10^{10}$ B mesons. It should therefore be possible to measure a CP asymmetry at the level of a few per cent at future colliders such as the Super- B factories [37–39].

6 $\bar{B}_d^0 \rightarrow \bar{K} \mu^+ \mu^-$

The CP asymmetry in $\bar{B}_d^0 \rightarrow \bar{K} \mu^+ \mu^-$ is defined in a manner similar to that in eq. (4.1), where dB/dz and $d\bar{B}/dz$ are the DBRs of $\bar{B}_d^0 \rightarrow \bar{K} \mu^+ \mu^-$ and its CP-conjugate $B_d^0 \rightarrow K \mu^+ \mu^-$, respectively. The expression for (dB/dz) has been given in ref. [11]. A model-independent analysis of the CP asymmetry in the DBR, with specific chosen values of VA operators, was carried out in ref. [19]. However, the constraints on the NP operators, coming from the measured branching ratio of $\bar{B}_d^0 \rightarrow X_s \mu^+ \mu^-$, were not taken into account. Here, in addition to taking these constraints into account, we also consider new SP and T operators, and extend the analysis to study the CP asymmetry in A_{FB} .

The CP asymmetry in A_{FB} is given in eq. (4.2), where the definition of A_{FB} is as given in ref. [11], while \bar{A}_{FB} is the analogous quantity for the CP-conjugate decay. Now, the decay mode $\bar{B}_d^0 \rightarrow \bar{K} \mu^+ \mu^-$ is unique because the forward-backward asymmetry of muons is predicted to vanish exactly in the SM. This is due to the fact that the $\bar{B}_d^0 \rightarrow \bar{K}$ hadronic matrix element does not have any axial-vector contribution. A_{FB} can therefore have a nonzero value only if it receives a contribution from new physics. However, even in the presence of NP, the expressions in ref. [11] indicate that the only term contributing to




Figure 2. The left (right) panels of the figure show $A_{CP}(q^2)$ and ΔA_{FB} for $\bar{B}_s^0 \rightarrow \mu^+ \mu^- \gamma$ in the low- q^2 (high- q^2) region, in the scenario where only (R_V, R_A) couplings are present. For example, the blue curve in the low- q^2 and high- q^2 regions for the A_{CP} plot corresponds to $(2.95e^{-i0.38}, 4.56e^{-i0.04})$, whereas the blue curve in the low- q^2 and high- q^2 regions for the ΔA_{FB} plot corresponds to $(1.60e^{-i0.08}, 4.14e^{-i0.12})$.

$\Delta A_{FB}(q^2)$ is that with VA+SP NP operators, and this is suppressed by the factor m_μ/m_b . As a result, one does not expect a significant enhancement in ΔA_{FB} from any Lorentz structure of NP.

Figure 3 shows $A_{CP}(q^2)$ for $\bar{B}_d^0 \rightarrow \bar{K} \mu^+ \mu^-$ in the presence of new VA couplings. We make the following observations:

- When only $R_{V,A}$ couplings are present, $A_{CP}(q^2)$ can be enhanced up to 7% at low q^2 . On the other hand, its value at high q^2 can be as high as 12%. $A_{CP}(q^2)$ can have either sign at both low and high q^2 , and its magnitude is almost independent of q^2 in these regions.
- When only $R'_{V,A}$ couplings are present, $A_{CP}(q^2)$ can be enhanced up to 4% at low q^2 . On the other hand, its value at high q^2 can be as high as 12%. $A_{CP}(q^2)$ can have either sign at both low and high q^2 , and its magnitude is almost independent of q^2 in these regions.
- When both primed and unprimed VA couplings are present, the results are the same as those obtained with only $R_{V,A}$ couplings.

For a $\bar{B}_d^0 \rightarrow \bar{K} \mu^+ \mu^-$ branching ratio of $O(0.5 \times 10^{-6})$, a measurement of a CP asymmetry of 1% at the 3σ level would require $\sim 10^{11}$ B mesons. It should therefore be possible




Figure 3. The left (right) panel of the figure shows $A_{CP}(q^2)$ for $\bar{B}_d^0 \rightarrow \bar{K}^* \mu^+ \mu^-$ in the low- q^2 (high- q^2) region, in the scenario where only (R_V, R_A) terms are present. The green line corresponds to the SM prediction. The other lines show predictions for some representative values of the NP parameters (R_V, R_A) . For example, the blue curve in the low- q^2 and high- q^2 regions corresponds to $(5.97e^{i2.23}, 3.08e^{-i0.05})$ and $(6.47e^{i2.30}, 3.11e^{i0.48})$, respectively.

to measure a CP asymmetry at the level of a few per cent at future colliders such as the Super- B factories [37–39].

7 $\bar{B}_d^0 \rightarrow \bar{K}^* \mu^+ \mu^-$

The complete three-angle distribution for the decay $\bar{B}^0 \rightarrow \bar{K}^{*0}(\rightarrow K^- \pi^+) \mu^+ \mu^-$ in the presence of NP can be expressed in terms of q^2 , two polar angles θ_μ and θ_K , and the azimuthal angle ϕ between the planes of the dimuon and $K\pi$ decays:

$$\begin{aligned} \frac{d^4 \Gamma^{\bar{B}}}{dq^2 d \cos \theta_\mu d \cos \theta_K d \phi} = N_F \left\{ \right. & \cos^2 \theta_K \left(I_1^0 + I_2^0 \cos 2\theta_\mu + I_3^0 \cos \theta_\mu \right) \\ & + \sin^2 \theta_K \left(I_1^T + I_2^T \cos 2\theta_\mu + I_3^T \cos \theta_\mu \right. \\ & \left. + I_4^T \sin^2 \theta_\mu \cos 2\phi + I_5^T \sin^2 \theta_\mu \sin 2\phi \right) + \sin 2\theta_K \left(I_1^{LT} \sin 2\theta_\mu \cos \phi \right. \\ & \left. + I_2^{LT} \sin 2\theta_\mu \sin \phi + I_3^{LT} \sin \theta_\mu \cos \phi + I_4^{LT} \sin \theta_\mu \sin \phi \right) \left. \right\}. \quad (7.1) \end{aligned}$$

The expressions for the normalization N_F and the I 's are given in ref. [11]. The I 's are functions of the couplings, kinematic variables and form factors. The definitions of the angles in $\bar{B}_d^0 \rightarrow \bar{K}^* \mu^+ \mu^-$ involve the directions of the μ^+ and \bar{K}^* . For the CP-conjugate decay $B^0 \rightarrow K^{*0}(\rightarrow K^+ \pi^-) \mu^+ \mu^-$, one defines these angles relative to the directions of the μ^- and K^* . The \bar{I} 's can be obtained from the I 's by replacing $\theta_\mu \rightarrow \theta_\mu - \pi$ and $\phi \rightarrow -\phi$, and changing the signs of the weak phases.

The CP asymmetries in the branching ratio and forward-backward asymmetry were analyzed in ref. [23] with the measurement of $B \rightarrow X_s \gamma$ and the limit on the $\bar{B}_d^0 \rightarrow \bar{K}^* \mu^+ \mu^-$ branching ratio available then. An analysis of CP asymmetries in $\bar{B}_d^0 \rightarrow \bar{K}^* \mu^+ \mu^-$ in the low- q^2 region was also performed earlier in ref. [25]. We extend this analysis by including T operators, and present our results for all asymmetries, in both the low- q^2 and high- q^2 regions.

A detailed discussion of the CP-conserving observables in this decay distribution can be found in ref. [11]. In this section we consider the direct CP asymmetries in the differential branching ratio (DBR), the forward-backward asymmetry A_{FB} , the longitudinal polarization fraction f_L , and the angular asymmetries $A_T^{(2)}$ and A_{LT} . We also examine the triple-product CP asymmetries $A_T^{(im)}$ and $A_{LT}^{(im)}$, which were not considered in ref. [11] since they identically vanish in the CP-conserving limit (no strong or weak phases), regardless of the presence of NP.

7.1 Direct CP asymmetries in the DBR and A_{FB}

The direct CP asymmetry in the differential branching ratio is defined as

$$A_{CP}(q^2) = \frac{(d\Gamma^{\bar{B}}/dq^2) - (d\Gamma^B/dq^2)}{(d\Gamma^{\bar{B}}/dq^2) + (d\Gamma^B/dq^2)}, \quad (7.2)$$

where

$$\frac{d\Gamma^{\bar{B}}}{dq^2} = \frac{8\pi N_F}{3}(A_L^{\bar{B}} + A_T^{\bar{B}}). \quad (7.3)$$

Here the longitudinal and transverse polarization amplitudes $A_L^{\bar{B}}$ and $A_T^{\bar{B}}$ are obtained from eq. (7.1):

$$A_L^{\bar{B}} = \left(I_1^0 - \frac{1}{3}I_2^0 \right), \quad A_T^{\bar{B}} = 2 \left(I_1^T - \frac{1}{3}I_2^T \right). \quad (7.4)$$

The expressions for A_L^B and A_T^B of the CP-conjugate mode can be obtained by replacing the I 's with \bar{I} 's.

The forward-backward asymmetry in $\bar{B}_d^0 \rightarrow \bar{K}^* \mu^+ \mu^-$ has recently been measured, and shows features that may indicate a deviation from the SM. This measured quantity is actually the CP-averaged forward-backward asymmetry A_{FB} . However, the difference between the measurement of this quantity in $\bar{B}_d^0 \rightarrow \bar{K}^* \mu^+ \mu^-$ and its CP-conjugate mode may also reveal the presence of NP. This CP asymmetry is quantified as

$$\Delta A_{FB}(q^2) = A_{FB}^{\bar{B}}(q^2) - A_{FB}^B(q^2), \quad (7.5)$$

where

$$A_{FB}^{\bar{B}(B)}(q^2) = \frac{\int_0^1 d \cos \theta_\mu \frac{d^2 \Gamma^{\bar{B}(B)}}{dq^2 d \cos \theta_\mu} - \int_{-1}^0 d \cos \theta_\mu \frac{d^2 \Gamma^{\bar{B}(B)}}{dq^2 d \cos \theta_\mu}}{\int_0^1 d \cos \theta_\mu \frac{d^2 \Gamma^{\bar{B}(B)}}{dq^2 d \cos \theta_\mu} + \int_{-1}^0 d \cos \theta_\mu \frac{d^2 \Gamma^{\bar{B}(B)}}{dq^2 d \cos \theta_\mu}}. \quad (7.6)$$

It can be obtained by integrating over the two angles θ_K and ϕ in eq. (7.1).

Figure 4 shows $A_{CP}(q^2)$ and $\Delta A_{FB}(q^2)$ for $\bar{B}_d^0 \rightarrow \bar{K}^* \mu^+ \mu^-$ in the presence of new VA couplings. We make the following observations:

- If only $R_{V,A}$ couplings are present, $A_{CP}(q^2)$ can be enhanced up to 5% at low q^2 , and up to 14 % at high q^2 . $\Delta A_{FB}(q^2)$ can be enhanced up to 3% at low q^2 , and up to 11 % at high q^2 . Both $A_{CP}(q^2)$, and $\Delta A_{FB}(q^2)$ can have either sign at both low and high q^2 .




Figure 4. The left (right) panels of the figure show $A_{CP}(q^2)$ and $\Delta A_{FB}(q^2)$ for $\bar{B}_d^0 \rightarrow \bar{K}^* \mu^+ \mu^-$ in the low- q^2 (high- q^2) region, in the scenario where (R_V, R_A, R'_V, R'_A) terms are all present. The green line corresponds to the SM prediction. The other lines show predictions for some representative values of the NP parameters. For example, the blue curve for $A_{CP}(q^2)$ in the low- q^2 and high- q^2 regions corresponds to $(2.77e^{i1.83}, 2.08e^{i0.5}, 3.8e^{i0.08}, 1.23e^{-i2.74})$ and $(5.88e^{i2.29}, 1.66e^{i0.82}, 3.49e^{i0.36}, 1.02e^{i0.98})$, respectively. The blue curve for $\Delta A_{FB}(q^2)$ in the low- q^2 and high- q^2 regions corresponds to $(1.56e^{-i2.59}, 1.80e^{-i0.35}, 4.23e^{i0.67}, 1.29e^{i1.43})$ and $(3.21e^{i2.61}, 1.38e^{i2.26}, 5.55e^{i0.69}, 3.03e^{i1.92})$, respectively.

- If only $R'_{V,A}$ couplings are present, $A_{CP}(q^2)$ can be enhanced up to 3% at low q^2 , and up to 7% at high q^2 . $\Delta A_{FB}(q^2)$ can be enhanced up to 1% at low q^2 , and up to 4% at high q^2 . Both $A_{CP}(q^2)$, and $\Delta A_{FB}(q^2)$ can have either sign at both low and high q^2 .
- When both primed and unprimed VA couplings are present, $A_{CP}(q^2)$ can be enhanced up to 9% at low q^2 , and up to 14% at high q^2 . $\Delta A_{FB}(q^2)$ can be enhanced up to 6% at low q^2 , and up to 19% at high q^2 . Both $A_{CP}(q^2)$, and $\Delta A_{FB}(q^2)$ can have either sign at both low and high q^2 (see figure 4).

These observations are consistent with the rough expectations in ref. [23] about the effect of VA operators.

7.2 Direct CP asymmetry in the polarization fraction f_L

The CP asymmetry in the longitudinal polarization fraction f_L is defined as

$$\Delta f_L = f_L^{\bar{B}} - f_L^B, \quad (7.7)$$




Figure 5. The left (right) panel of the figure shows $\Delta f_L(q^2)$ for $\bar{B}_d^0 \rightarrow \bar{K}^* \mu^+ \mu^-$ in the low- q^2 (high- q^2) region, in the scenario where (R_V, R_A, R'_V, R'_A) terms are all present. For example, the blue curve in the low- q^2 and high- q^2 regions corresponds to $(2.78e^{i2.98}, 2.19e^{-i0.77}, 6.91e^{-i0.29}, 3.34e^{-i0.56})$.

where

$$f_L^{\bar{B}(B)} = \frac{A_L^{\bar{B}(B)}}{A_L^{\bar{B}(B)} + A_T^{\bar{B}(B)}}. \tag{7.8}$$

figure 5 shows $\Delta f_L(q^2)$ for $\bar{B}_d^0 \rightarrow \bar{K}^* \mu^+ \mu^-$ in the presence of new VA couplings. We make the following observations:

- If only $R_{V,A}$ couplings are present, $\Delta f_L(q^2)$ can be enhanced up to 2% at very low q^2 . On the other hand, $\Delta f_L(q^2)$ is almost the same as the SM at high q^2 . It can have either sign at both low and high q^2 .
- If only $R'_{V,A}$ couplings are present, $\Delta f_L(q^2)$ can be enhanced up to 2% at both low and high q^2 . It can have either sign at both low and high q^2 .
- When both primed and unprimed VA couplings are present, $\Delta f_L(q^2)$ can be enhanced up to 9% at low q^2 , and up to 6% at high q^2 . It can have either sign at both low and high q^2 (see figure 5).

7.3 Direct CP asymmetries in the angular asymmetries $A_T^{(2)}$ and A_{LT}

The transverse asymmetry $A_T^{(2)\bar{B}(B)}$ is defined [40] through the double differential decay rate as

$$\frac{d^2\Gamma^{\bar{B}(B)}}{dq^2 d\phi} = \frac{1}{2\pi} \frac{d\Gamma^{\bar{B}(B)}}{dq^2} \left[1 + f_T^{\bar{B}(B)} \left(A_T^{(2)\bar{B}(B)} \cos 2\phi + A_T^{(im)\bar{B}(B)} \sin 2\phi \right) \right]. \tag{7.9}$$

It can be obtained by integrating eq. (7.1) over the two polar angles θ_μ and θ_K . Here $A_T^{(im)\bar{B}(B)}$ is a triple product, and is discussed separately below. In terms of the coupling constants and matrix elements defined in ref. [11], $A_T^{(2)\bar{B}(B)}$ can be expressed as

$$A_T^{(2)\bar{B}} = \frac{4I_4^T}{3A_T^{\bar{B}}}, \quad A_T^{(2)B} = \frac{4\bar{I}_4^T}{3A_T^B}. \tag{7.10}$$




Figure 6. The left (right) panel of the figure shows $\Delta A_T^{(2)}(q^2)$ for $\bar{B}_d^0 \rightarrow \bar{K}^* \mu^+ \mu^-$ in the low- q^2 (high- q^2) region, in the scenario where (R_V, R_A, R'_V, R'_A) terms are all present. The green line corresponds to the SM prediction. The other lines show predictions for some representative values of the NP parameters. For example, the blue curve in the low- q^2 and high- q^2 regions corresponds to $(0.11e^{i2.18}, 2.66e^{-i1.31}, 4.3e^{i0.03}, 0.23e^{-i2.27})$ and $(2.32e^{i2.51}, 4.89e^{i1.27}, 3.12e^{i0.42}, 0.14e^{-i1.55})$, respectively.

While $A_T^{(2)\bar{B}} (A_T^{(2)B})$ is finite even in the CP-conserving limit (and was discussed in ref. [11]), a CP asymmetry may be defined through the difference

$$\Delta A_T^{(2)} \equiv A_T^{(2)\bar{B}} - A_T^{(2)B} . \quad (7.11)$$

figure 6 shows $\Delta A_T^{(2)}$ for $\bar{B}_d^0 \rightarrow \bar{K}^* \mu^+ \mu^-$ in the presence of new VA couplings. We make the following observations:

- If only $R_{V,A}$ couplings are present, $\Delta A_T^{(2)}$ cannot be enhanced more than 1% at both low and high q^2 . It can have either sign at both low and high q^2 .
- If only $R'_{V,A}$ couplings are present, $\Delta A_T^{(2)}$ can be enhanced up to 4% at low q^2 , and up to 6% high q^2 . It can have either sign at both low and high q^2 .
- When both primed and unprimed VA couplings are present, $\Delta A_T^{(2)}$ can be enhanced up to 11% at low q^2 , and up to 12% at high q^2 . It can have either sign at both low and high q^2 (see figure 6).

The longitudinal-transverse asymmetry $A_{LT}^{\bar{B}(B)}$ is defined through

$$\frac{d^2\Gamma_{LT}^{\bar{B}(B)}}{dq^2 d\phi} = \frac{d\Gamma^{\bar{B}(B)}}{dq^2} \left(A_{LT}^{(re)\bar{B}(B)} \cos \phi + A_{LT}^{(im)\bar{B}(B)} \sin \phi \right) , \quad (7.12)$$

where

$$\frac{d^2\Gamma_{LT}^{\bar{B}(B)}}{dq^2 d\phi} = \int_0^1 d \cos \theta_K \frac{d^3\Gamma^{\bar{B}(B)}}{dq^2 d \cos \theta_K d\phi} - \int_{-1}^0 d \cos \theta_K \frac{d^3\Gamma^{\bar{B}(B)}}{dq^2 d \cos \theta_K d\phi} . \quad (7.13)$$

Here $A_{LT}^{(im)\bar{B}(B)}$ is a triple product, and is discussed separately below. In terms of the coupling constants and matrix elements defined in ref. [11], $A_{LT}^{(re)\bar{B}(B)}$ can be expressed as

$$A_{LT}^{(re)\bar{B}} = \frac{I_3^{LT}}{4(A_L^{\bar{B}} + A_T^{\bar{B}})} , \quad A_{LT}^{(re)B} = -\frac{\bar{I}_3^{LT}}{4(A_L^B + A_T^B)} . \quad (7.14)$$

Note that $A_{LT}^{(re)B} = -A_{LT}^{(re)\bar{B}}$ in the CP-conserving limit. Thus, a CP asymmetry may be defined through the sum

$$\Delta A_{LT}(q^2) \equiv A_{LT}^{(re)\bar{B}}(q^2) + A_{LT}^{(re)B}(q^2). \quad (7.15)$$

We now assume the presence of new VA couplings. However, we find that these couplings cannot enhance $\Delta A_{LT}(q^2)$ to more than 3% at both low and high q^2 .

Note that $\Delta A_{LT}(q^2)$ is related to the observable A_5^D in ref. [24]: $\Delta A_{LT}(q^2) \approx A_5^D/4$. Our limit of 3% on the maximum value of $\Delta A_{LT}(q^2)$ is then consistent with the limit of 0.07 on the average value $\langle A_5^D \rangle$ over the low- q^2 region, as calculated in ref. [24].

7.4 CP-violating triple-product asymmetries

In this subsection, we consider the triple products (TPs) in the decays $\bar{B}^0 \rightarrow \bar{K}^{*0}(\rightarrow K^-\pi^+)\mu^+\mu^-$ and $B^0 \rightarrow K^{*0}(\rightarrow K^+\pi^-)\mu^+\mu^-$. For the decaying \bar{B} meson, the TP is proportional to $(\hat{n}_K \times \hat{n}_\mu) \cdot \hat{n}_z$ in its rest frame, where the unit vectors are given in terms of the momenta of the final-state particles as

$$\hat{n}_K = \frac{\hat{p}_{K^-} \times \hat{p}_{\pi^+}}{|\hat{p}_{K^-} \times \hat{p}_{\pi^+}|}, \quad \hat{n}_z = \frac{\hat{p}_{K^-} + \hat{p}_{\pi^+}}{|\hat{p}_{K^-} + \hat{p}_{\pi^+}|}, \quad \hat{n}_\mu = \frac{\hat{p}_{\mu^-} \times \hat{p}_{\mu^+}}{|\hat{p}_{\mu^-} \times \hat{p}_{\mu^+}|}. \quad (7.16)$$

In terms of the azimuthal angle ϕ , one gets

$$\cos \phi = \hat{n}_K \cdot \hat{n}_\mu, \quad \sin \phi = (\hat{n}_K \times \hat{n}_\mu) \cdot \hat{n}_z, \quad (7.17)$$

and hence the quantities that are coefficients of $\sin \phi$ (or of $\sin 2\phi = 2 \sin \phi \cos \phi$) are the TPs.

As noted above, while the angular distribution for the \bar{B} decay involves ϕ , for B it involves $-\phi$. Thus, the CP-violating triple-product asymmetry is proportional to the *sum* of \bar{B} and B TPs.

The first TP is $A_T^{(im)\bar{B}(B)}$, introduced above in eq. (7.9). In terms of the coupling constants and matrix elements defined in ref. [11], $A_T^{(im)\bar{B}(B)}$ can be written as

$$A_T^{(im)\bar{B}} = \frac{4I_5^T}{3A_T^{\bar{B}}}, \quad A_T^{(im)B} = -\frac{4\bar{I}_5^T}{3A_T^{\bar{B}}}. \quad (7.18)$$

We observe that $A_T^{(im)}$ depends only on the VA couplings. The CP-violating triple-product asymmetry is

$$A_T^{(im)} = \frac{1}{2}(A_T^{(im)\bar{B}} + A_T^{(im)B}). \quad (7.19)$$

figure 7 shows $A_T^{(im)}(q^2)$ for $\bar{B}_d^0 \rightarrow \bar{K}^{*0}\mu^+\mu^-$ in the presence of new VA couplings. We make the following observations:

- If only $R_{V,A}$ couplings are present, $A_T^{(im)}(q^2)$ can be enhanced up to 5% at low q^2 and can have either sign. On the other hand, $A_T^{(im)}(q^2)$ is almost same as the SM prediction ($\simeq 0$) at high q^2 .




Figure 7. The left (right) panel of the figure shows $A_T^{(im)}(q^2)$ for $\bar{B}_d^0 \rightarrow \bar{K}^* \mu^+ \mu^-$ in the low- q^2 (high- q^2) region, in the scenario where (R_V, R_A, R'_V, R'_A) terms are all present. The green line corresponds to the SM prediction. The other lines show predictions for some representative values of the NP parameters. For example, the blue curve in the low- q^2 and high- q^2 regions corresponds to $(1.33e^{-i2.96}, 0.78e^{i2.47}, 0.83e^{-i0.27}, 3.15e^{i1.75})$ and $(2.15e^{-i2.77}, 0.7e^{-i2.43}, 8.20e^{-i0.16}, 4.8e^{-i1.62})$, respectively.

- If only $R'_{V,A}$ couplings are present, $A_T^{(im)}(q^2)$ can be enhanced up to 49% at low q^2 , and up to 46% at high q^2 . It can have either sign at both low and high q^2 .
- When both primed and unprimed VA couplings are present, the results for $A_T^{(im)}(q^2)$ are almost the same as those obtained with only $R'_{V,A}$ couplings (see figure 7).

The second TP is $A_{LT}^{(im)\bar{B}(B)}$, introduced above in eq. (7.12). In terms of the coupling constants and matrix elements defined in ref. [11], $A_{LT}^{(im)\bar{B}(B)}$ can be written as

$$A_{LT}^{(im)\bar{B}} = \frac{I_4^{LT}}{4(A_L^{\bar{B}} + A_T^{\bar{B}})}, \quad A_{LT}^{(im)B} = \frac{\bar{I}_4^{LT}}{4(A_L^B + A_T^B)}. \quad (7.20)$$

We observe that A_{LT} depends on the VA couplings, as well as on V-S and SP-T interference terms. The CP-violating triple-product asymmetry is

$$A_{LT}^{(im)} = \frac{1}{2}(A_{LT}^{(im)\bar{B}} - A_{LT}^{(im)B}). \quad (7.21)$$

figure 8 shows $A_{LT}^{(im)}(q^2)$ for $\bar{B}_d^0 \rightarrow \bar{K}^* \mu^+ \mu^-$ in the presence of new VA couplings. We make the following observations:

- If only $R_{V,A}$ couplings are present, $A_{LT}^{(im)}(q^2)$ can be enhanced up to 6% at very low q^2 , and is almost same as the SM prediction (≈ 0) at high q^2 . It can have either sign at both low and high q^2 .
- If only $R'_{V,A}$ couplings are present, $A_{LT}^{(im)}(q^2)$ can be enhanced up to 8% at low q^2 and is almost same as the SM prediction (≈ 0) at high q^2 . It can have either sign at both low and high q^2 .
- When both primed and unprimed VA couplings are present, $A_{LT}^{(im)}(q^2)$ can be enhanced up to 10% at low q^2 and up to 0.5% at high q^2 . It can have either sign at both low and high q^2 (see figure 8).




Figure 8. The figure shows $A_{LT}^{(im)}(q^2)$ for $\bar{B}_d^0 \rightarrow \bar{K}^* \mu^+ \mu^-$ in the low- q^2 region, in the scenario where (R_V, R_A, R'_V, R'_A) terms are all present. The green line corresponds to the SM prediction. The other lines show predictions for some representative values of the NP parameters. For example, the blue curve corresponds to $(1.68e^{i1.92}, 2.27e^{i0.53}, 4.22e^{i0.28}, 0.14e^{-i1.91})$.

Note that $A_{LT}^{(im)}(q^2)$ is related to the observable A_7^D in ref. [24]: $A_{LT}^{(im)}(q^2) \approx A_7^D/8$. Our limit of 10% on the maximum value of $A_{LT}^{(im)}(q^2)$ is then consistent with the limit of 0.76 on the average value $\langle A_7^D \rangle$ over the low- q^2 region, as calculated in ref. [24]. However, in addition we present the full q^2 -dependence of this quantity.

In principle, $A_{LT}^{(im)\bar{B}(B)}$ can be generated due to NP SP-T interference. However, we find that the effect is tiny: $A_{LT}^{(im)}(q^2)$ can be enhanced up to 0.4% at low q^2 and can have either sign; $A_{LT}^{(im)}(q^2)$ is same as the SM ($\simeq 0$) at high q^2 .

8 Discussion and summary

Even after the successful start of the LHC that will search for new physics (NP) at the TeV scale and beyond, B decays still remain one of the best avenues of detecting indirect NP signals. The copious amount of data on B decays, expected from future experiments like the LHC and super- B factories, will allow us to explore in detail many decay modes that are currently considered to be rare. The combined analysis of many such decay modes will allow us to look for NP in a model-independent manner.

We consider all possible Lorentz structures of new physics (NP) in the $b \rightarrow s \mu^+ \mu^-$ transition, and analyze their effects on the CP-violating observables in (i) $\bar{B}_s^0 \rightarrow \mu^+ \mu^-$, (ii) $\bar{B}_d^0 \rightarrow X_s \mu^+ \mu^-$, (iii) $\bar{B}_s^0 \rightarrow \mu^+ \mu^- \gamma$, (iv) $\bar{B}_d^0 \rightarrow \bar{K} \mu^+ \mu^-$, (v) $\bar{B}_d^0 \rightarrow \bar{K}^* \mu^+ \mu^-$, and their CP-conjugate modes. These are the same modes we explored in the companion paper [11], where we considered only CP-conserving quantities. We find that for $\bar{B}_s^0 \rightarrow \mu^+ \mu^-$, the only CP-violating quantities that can be constructed even in principle require the measurement of muon polarization, a task not possible in foreseeable detectors. Therefore, we do not dwell on this mode further. For the rest of the modes, we focus on

- CP violation in the differential branching ratio (A_{CP}), and
- CP violation in the forward-backward asymmetry (ΔA_{FB}).

In addition, for $\bar{B}_d^0 \rightarrow \bar{K}^* \mu^+ \mu^-$, we analyze

- the CP asymmetry in the longitudinal polarization fraction (Δf_L),
- the CP asymmetries $\Delta A_T^{(2)}$ and ΔA_{LT} arising in the angular distributions, and
- the triple-product (TP) CP asymmetries $\Delta A_T^{(im)}$ and $\Delta A_{LT}^{(im)}$.

We determine the constraints on the coupling constants in the effective NP operators by using the currently available data. On the basis of these limits and general arguments, we expect that the CP-violating quantities in most of the modes can only be sensitive to the vector-axial vector (VA) couplings, while the scalar-pseudoscalar (SP) and the tensor (T) NP operators can only contribute, if at all, to certain TP asymmetries. Our later detailed exploration of the allowed parameter space for all the NP couplings vindicates this argument. The effects of SP and T NP operators are therefore discussed only briefly in this paper.

On the other hand, the VA operators can have a significant impact on the CP-violating observables. (See table 1). The SM predicts $A_{CP}(q^2) \lesssim 10^{-3}$ for all the modes, while VA NP operators allow this quantity to be as large as $\sim 10\%$ (for $\bar{B}_d^0 \rightarrow X_s \mu^+ \mu^-$, $\bar{B}_d^0 \rightarrow \bar{K} \mu^+ \mu^-$ and $\bar{B}_d^0 \rightarrow \bar{K}^* \mu^+ \mu^-$) and even up to $\sim 30\%$ for $\bar{B}_s^0 \rightarrow \mu^+ \mu^- \gamma$. Even ΔA_{FB} , expected to be $\lesssim 10^{-4}$ in the SM, can be enhanced up to $\sim 10\%$ (for $\bar{B}_d^0 \rightarrow X_s \mu^+ \mu^-$) and up to $\sim 40\%$ (for $\bar{B}_s^0 \rightarrow \mu^+ \mu^- \gamma$). While ΔA_{FB} in $\bar{B}_d^0 \rightarrow \bar{K} \mu^+ \mu^-$ stays zero even with VA NP, its value in $\bar{B}_d^0 \rightarrow \bar{K}^* \mu^+ \mu^-$ may be enhanced to $\sim 10\%$ from its SM expectation of $\lesssim 10^{-4}$.

In $\bar{B}_d^0 \rightarrow \bar{K}^* \mu^+ \mu^-$ the SM predicts $\Delta f_L \lesssim 10^{-4}$, while VA NP operators allow this quantity to be enhanced up to $\sim 10\%$. $\Delta A_T^{(2)}$, ΔA_{LT} , $A_T^{(im)}$ and $A_{LT}^{(im)}$ are all zero in the SM. VA NP operators can enhance $\Delta A_T^{(2)}$ up to $\sim 12\%$, $A_T^{(im)}$ even up to $\sim 50\%$, and $A_{LT}^{(im)}$ up to $\sim 10\%$. ΔA_{LT} can not be enhanced more than $\sim 3\%$ even in the presence of VA NP operators. Note that while in almost all the cases the impact of the left-handed VA NP couplings $R_{V,A}$ is dominant, for the TP asymmetry $\Delta A_T^{(im)}$, the $R'_{V,A}$ couplings play a dominating role.

TP's can also be generated by NP-NP interference. However, we do not find large effects. The interference of SP-T operators can increase $A_{LT}^{(im)}(q^2)$ up to only 0.4% at low q^2 .

It is quite possible that if the NP is of the VA type, its presence would first be indicated through the CP-conserving/CP-averaged quantities considered in ref. [11]. However, the CP-violating signals considered in this paper are so robust (orders of magnitude more than the SM predictions) that these may be the ones that will unambiguously establish the presence of NP of the VA kind. Moreover, hadronic uncertainties play a very minor role in the CP-violating asymmetries considered in this paper. A combined analysis of CP-violating and CP-conserving signals may allow even the determination of the magnitudes and phases of the NP coupling constants, in addition to confirming the NP Lorentz structure.

Acknowledgments

We thank S. Uma Sankar for helpful collaboration on several parts of this analysis. The work of AKA and DL was financially supported by NSERC of Canada. The work of A. Datta and M. Duraisamy was supported by the US-Egypt Joint Board on Scientific and

Observable	SM	Only new VA	Only new SP	Only new T
$\bar{B}_d^0 \rightarrow X_s \mu^+ \mu^-$ A_{CP} ΔA_{FB}	$\bullet 10^{-3} \rightarrow 10^{-4}$ (low→high q^2) $10^{-4} \rightarrow 10^{-5}$ (low→high q^2)	$\bullet (6 \rightarrow 12)\%$ (low→high q^2) $\bullet (3 \rightarrow 12)\%$ (low→high q^2)	\bullet Marginal S $\bullet < 1\%$	\bullet Marginal S/E No effect
$\bar{B}_s^0 \rightarrow \mu^+ \mu^- \gamma$ A_{CP} ΔA_{FB}	$\bullet 10^{-3} \rightarrow 10^{-4}$ (low→high q^2) $10^{-4} \rightarrow 10^{-5}$ (low→high q^2)	$\bullet (30 \rightarrow 13)\%$ (low→high q^2) $\bullet (40 \rightarrow 18)\%$ (low→high q^2)	No effect No effect	$\bullet < 1\%$ $\bullet < 1\%$
$\bar{B}_d^0 \rightarrow \bar{K} \mu^+ \mu^-$ A_{CP} ΔA_{FB}	$\bullet 10^{-3} \rightarrow 10^{-4}$ (low→high q^2) Zero	$\bullet (7 \rightarrow 12)\%$ (low→high q^2) No effect	\bullet Marginal S $\bullet < 1\%$	\bullet Marginal S/E No effect
$\bar{B}_d^0 \rightarrow \bar{K}^* \mu^+ \mu^-$ A_{CP} ΔA_{FB} Δf_L $\Delta A_T^{(2)}$ ΔA_{LT} $A_T^{(im)}$ $A_{LT}^{(im)}$	$\bullet 10^{-3} \rightarrow 10^{-4}$ (low→high q^2) $\bullet 10^{-4} \rightarrow 10^{-6}$ (low→high q^2) $\bullet 10^{-4} \rightarrow 10^{-7}$ (low→high q^2) Zero Zero Zero Zero	$\bullet (9 \rightarrow 14)\%$ (low→high q^2) $\bullet (6 \rightarrow 19)\%$ (low→high q^2) $\bullet (9 \rightarrow 16)\%$ (low→high q^2) $\bullet \sim 12\%$ $\bullet < 3\%$ $\bullet \sim 50\%$ $\bullet \sim 10\%$	No effect No effect No effect No effect No effect No effect	$\bullet < 1\%$ $\bullet < 1\%$ $\bullet < 1\%$ No effect No effect No effect No effect

Table 1. The effect of NP couplings on observables. E: enhancement, S: suppression. The numbers given are optimistic estimates.

Technological Co-operation award (Project ID: 1855) administered by the US Department of Agriculture.

References

- [1] BABAR collaboration, B. Aubert et al., *Rates, polarizations and asymmetries in charmless vector-vector B meson decays*, *Phys. Rev. Lett.* **91** (2003) 171802 [[hep-ex/0307026](#)] [[INSPIRE](#)].
- [2] BELLE collaboration, K. Chen et al., *Measurement of branching fractions and polarization in $B \rightarrow \phi K^* (*)$ decays*, *Phys. Rev. Lett.* **91** (2003) 201801 [[hep-ex/0307014](#)] [[INSPIRE](#)].
- [3] BELLE collaboration, J.-T. Wei et al., *Measurement of the Differential Branching Fraction and Forward-Backward Asymmetry for $B \rightarrow K^{(*)} \ell^+ \ell^-$* , *Phys. Rev. Lett.* **103** (2009) 171801 [[arXiv:0904.0770](#)] [[INSPIRE](#)].

- [4] BABAR collaboration, B. Aubert et al., *Angular Distributions in the Decays $B \rightarrow K^* \ell^+ \ell^-$* , *Phys. Rev. D* **79** (2009) 031102 [[arXiv:0804.4412](#)] [[INSPIRE](#)].
- [5] A.K. Alok et al., *New-physics contributions to the forward-backward asymmetry in $B \rightarrow K^* \mu^+ \mu^-$* , *JHEP* **02** (2010) 053 [[arXiv:0912.1382](#)] [[INSPIRE](#)].
- [6] H.-Y. Cheng, C.-K. Chua and A. Soni, *CP-violating asymmetries in B_0 decays to $K^+ K^- K_{S(L)}^0$ and $K_S^0 K_S^0 K_{S(L)}^0$* , *Phys. Rev. D* **72** (2005) 094003 [[hep-ph/0506268](#)] [[INSPIRE](#)].
- [7] G. Buchalla, G. Hiller, Y. Nir and G. Raz, *The Pattern of CP asymmetries in $B \rightarrow s$ transitions*, *JHEP* **09** (2005) 074 [[hep-ph/0503151](#)] [[INSPIRE](#)].
- [8] E. Lunghi and A. Soni, *Hints for the scale of new CP-violating physics from B-CP anomalies*, *JHEP* **08** (2009) 051 [[arXiv:0903.5059](#)] [[INSPIRE](#)].
- [9] CDF collaboration, T. Aaltonen et al., D0 collaboration, V.M. Abazov et al., *Combination of D0 and CDF Results on $\Delta\Gamma_s$ and the CP-Violating Phase $\beta_s^{J/\psi\phi}$* , CDF Note CDF/PHYS/BOTTOM/CDFR/9787 (2009), D0 Note 5928-CONF (2009).
- [10] D0 collaboration, V.M. Abazov et al., *Evidence for an anomalous like-sign dimuon charge asymmetry*, *Phys. Rev. D* **82** (2010) 032001 [[arXiv:1005.2757](#)] [[INSPIRE](#)].
- [11] A.K. Alok et al., *New Physics in $b \rightarrow s \mu^+ \mu^-$: CP-Conserving Observables*, [arXiv:1008.2367](#) [[INSPIRE](#)].
- [12] D.-S. Du and M.-Z. Yang, *CP violation for $b \rightarrow X_s \ell^+ \ell^-$ including long distance effects*, *Phys. Rev. D* **54** (1996) 882 [[hep-ph/9510267](#)] [[INSPIRE](#)].
- [13] T. Aliev, D.A. Demir, E. Iltan and N. Pak, *The CP asymmetry in $b \rightarrow s \ell^+ \ell^-$ decay*, *Phys. Rev. D* **54** (1996) 851 [[hep-ph/9511352](#)] [[INSPIRE](#)].
- [14] S. Fukae, *CP asymmetry of $B \rightarrow X_s \ell^+ \ell^-$ in low invariant mass region*, *Phys. Rev. D* **64** (2001) 054010 [[hep-ph/0102041](#)] [[INSPIRE](#)].
- [15] A.K. Alok, A. Dighe and S. Ray, *CP asymmetry in the decays $B \rightarrow (X_s, X_d) \mu^+ \mu^-$ with four generations*, *Phys. Rev. D* **79** (2009) 034017 [[arXiv:0811.1186](#)] [[INSPIRE](#)].
- [16] A. Soni, A.K. Alok, A. Giri, R. Mohanta and S. Nandi, *SM with four generations: Selected implications for rare B and K decays*, *Phys. Rev. D* **82** (2010) 033009 [[arXiv:1002.0595](#)] [[INSPIRE](#)].
- [17] I. Balakireva, D. Melikhov, N. Nikitin and D. Tlisov, *Forward-backward and CP-violating asymmetries in rare $B_{d,s} \rightarrow (\phi, \gamma) l^+ l^-$ decays*, *Phys. Rev. D* **81** (2010) 054024 [[arXiv:0911.0605](#)] [[INSPIRE](#)].
- [18] I. Balakireva, N. Nikitin and D. Tlisov, *Asymmetries in Rare Radiative Leptonic and Semileptonic Decays of B-Mesons*, *Yad. Fiz.* **73** (2010) 1762 [[INSPIRE](#)].
- [19] T. Aliev, S. Choudhury, A.S. Cornell and N. Gaur, *CP violation in the $B \rightarrow K \ell^+ \ell^-$ decay*, *Eur. Phys. J. C* **49** (2007) 657 [[hep-ph/0506188](#)] [[INSPIRE](#)].
- [20] F. Krüger, L.M. Sehgal, N. Sinha and R. Sinha, *Angular distribution and CP asymmetries in the decays $\bar{B} \rightarrow K^- \pi^+ e^- e^+$ and $\bar{B} \rightarrow \pi^- \pi^+ e^- e^+$* , *Phys. Rev. D* **61** (2000) 114028 [*Erratum ibid* **D 63** (2001) 019901] [[hep-ph/9907386](#)] [[INSPIRE](#)].
- [21] T. Aliev, D.A. Demir and M. Savci, *Probing the sources of CP-violation via $B \rightarrow K^* \ell^+ \ell^-$ decay*, *Phys. Rev. D* **62** (2000) 074016 [[hep-ph/9912525](#)] [[INSPIRE](#)].
- [22] G. Buchalla, G. Hiller and G. Isidori, *Phenomenology of nonstandard Z couplings in exclusive semileptonic $b \rightarrow s$ transitions*, *Phys. Rev. D* **63** (2000) 014015 [[hep-ph/0006136](#)] [[INSPIRE](#)].

- [23] F. Krüger and E. Lunghi, *Looking for novel CP-violating effects in $\bar{B} \rightarrow K^* \ell^+ \ell^-$* , *Phys. Rev. D* **63** (2001) 014013 [[hep-ph/0008210](#)] [[INSPIRE](#)].
- [24] C. Bobeth, G. Hiller and G. Piranishvili, *CP Asymmetries in $\bar{B} \rightarrow \bar{K}^*(\rightarrow \bar{K}\pi)\bar{\ell}\ell$ and Untagged $\bar{B}_s, B_s \rightarrow \phi(\rightarrow K^+K^-)\bar{\ell}\ell$ Decays at NLO*, *JHEP* **07** (2008) 106 [[arXiv:0805.2525](#)] [[INSPIRE](#)].
- [25] W. Altmannshofer, P. Ball, A. Bharucha, A.J. Buras, D.M. Straub and M. Wick, *Symmetries and Asymmetries of $B \rightarrow K^* \mu^+ \mu^-$ Decays in the Standard Model and Beyond*, *JHEP* **01** (2009) 019 [[arXiv:0811.1214](#)] [[INSPIRE](#)].
- [26] U. Egede, T. Hurth, J. Matias, M. Ramon and W. Reece, *New physics reach of CP-violating observables in the decay $\bar{B}_d \rightarrow \bar{K}^* \ell^+ \ell^-$* , *PoS(EPS-HEP 2009)184* [[arXiv:0912.1349](#)] [[INSPIRE](#)].
- [27] U. Egede, T. Hurth, J. Matias, M. Ramon and W. Reece, *New physics reach of the decay mode $\bar{B} \rightarrow \bar{K}^{*0} \ell^+ \ell^-$* , *JHEP* **10** (2010) 056 [[arXiv:1005.0571](#)] [[INSPIRE](#)].
- [28] C. Bobeth, G. Hiller and D. van Dyk, *More Benefits of Semileptonic Rare B Decays at Low Recoil: CP-violation*, *JHEP* **07** (2011) 067 [[arXiv:1105.0376](#)] [[INSPIRE](#)].
- [29] C. Bobeth, M. Misiak and J. Urban, *Photonic penguins at two loops and m_t dependence of $BR(B \rightarrow X_s \ell^+ \ell^-)$* , *Nucl. Phys. B* **574** (2000) 291 [[hep-ph/9910220](#)] [[INSPIRE](#)].
- [30] PARTICLE DATA GROUP collaboration, C. Amsler et al. *Review of particle physics*, *Phys. Lett. B* **667** (2008) 1 [[INSPIRE](#)].
- [31] HEAVY FLAVOR AVERAGING GROUP collaboration, E. Barberio et al., *Averages of b-hadron and c-hadron Properties at the End of 2007*, [arXiv:0808.1297](#) [[INSPIRE](#)].
- [32] T. Huber, T. Hurth and E. Lunghi, *Logarithmically Enhanced Corrections to the Decay Rate and Forward Backward Asymmetry in $\bar{B} \rightarrow X_s \ell^+ \ell^-$* , *Nucl. Phys. B* **802** (2008) 40 [[arXiv:0712.3009](#)] [[INSPIRE](#)].
- [33] A. Datta and D. London, *Triple-product correlations in $B \rightarrow V_1 V_2$ decays and new physics*, *Int. J. Mod. Phys. A* **19** (2004) 2505 [[hep-ph/0303159](#)] [[INSPIRE](#)].
- [34] A. Datta and D. London, *Measuring new physics parameters in B penguin decays*, *Phys. Lett. B* **595** (2004) 453 [[hep-ph/0404130](#)] [[INSPIRE](#)].
- [35] A.J. Buras and M. Münz, *Effective Hamiltonian for $B \rightarrow X_s \ell^+ \ell^-$ beyond leading logarithms in the NDR and HV schemes*, *Phys. Rev. D* **52** (1995) 186 [[hep-ph/9501281](#)] [[INSPIRE](#)].
- [36] D. Melikhov and N. Nikitin, *Rare radiative leptonic decays $B_{d,s} \rightarrow \ell^+ \ell^- \gamma$* , *Phys. Rev. D* **70** (2004) 114028 [[hep-ph/0410146](#)] [[INSPIRE](#)].
- [37] T.E. Browder, T. Gershon, D. Pirjol, A. Soni and J. Zupan, *New Physics at a Super Flavor Factory*, *Rev. Mod. Phys.* **81** (2009) 1887 [[arXiv:0802.3201](#)] [[INSPIRE](#)].
- [38] SUPERB collaboration, M. Bona et al., *SuperB: A High-Luminosity Asymmetric e^+e^- Super Flavor Factory. Conceptual Design Report*, [arXiv:0709.0451](#) [[INSPIRE](#)].
- [39] SUPERB collaboration, B. O’Leary et al., *SuperB Progress Reports – Physics*, [arXiv:1008.1541](#) [[INSPIRE](#)].
- [40] F. Krüger and J. Matias, *Probing new physics via the transverse amplitudes of $B^0 \rightarrow K^{*0}(\rightarrow K^-\pi^+)\ell^+\ell^-$ at large recoil*, *Phys. Rev. D* **71** (2005) 094009 [[hep-ph/0502060](#)] [[INSPIRE](#)].