

$\mathcal{N} = 1$ sigma models in AdS_4

Allan Adams,^a Hans Jockers,^{b,c} Vijay Kumar^c and Joshua M. Lapan^c

^aCenter for Theoretical Physics, Massachusetts Institute of Technology,
Cambridge, MA 02139, U.S.A.

^bDepartment of Physics, Stanford University,
Stanford, CA 94305-4060, U.S.A.

^cKavli Institute for Theoretical Physics, University of California,
Santa Barbara, CA 93106, U.S.A.

E-mail: awa@mit.edu, jockers@kitp.ucsb.edu, vijayk@kitp.ucsb.edu,
lapan@kitp.ucsb.edu

ABSTRACT: We study sigma models in AdS_4 with global $\mathcal{N} = 1$ supersymmetry and find that they differ significantly from their flat-space cousins — the target space is constrained to be a Kähler manifold with an exact Kähler form, the superpotential transforms under Kähler transformations, the space of supersymmetric vacua is generically a set of isolated points even when the superpotential vanishes, and the R -symmetry is classically broken by the cosmological constant. Remarkably, the exactness of the Kähler class is also required for the sigma model to arise as a decoupling limit of $\mathcal{N} = 1$ supergravity, and ensures the vanishing of gravitational anomalies. As applications of these results, we argue that fields with AdS_4 scale masses are ubiquitous in, for example, type IIB $\mathcal{N} = 1$ AdS_4 vacua stabilized near large volume; we also present a schematic argument that the Affleck-Dine-Seiberg runaway of $N_f < N_c$ SQCD can be regulated by considering the theory in AdS_4 .

KEYWORDS: Supersymmetric Effective Theories, Sigma Models, Supergravity Models

ARXIV EPRINT: [1104.3155](https://arxiv.org/abs/1104.3155)

Contents

1	Introduction and conclusions	1
2	Supersymmetric Lagrangians for chiral multiplets in AdS₄	4
2.1	The $\mathcal{N} = 1$ supersymmetric sigma model in Minkowski space	4
2.2	Chiral multiplets in AdS ₄	6
2.3	AdS ₄ supersymmetric sigma model	8
2.4	Supersymmetric vacua in AdS ₄	10
2.5	Target space geometry for $\mathcal{N} = 1$ sigma models in AdS ₄	11
2.6	Derivation through supergravity	13
3	Lessons for constraints in flat space and beyond	14
3.1	Anomalies and constraints on (de-)coupling gravity	15
3.2	The background principle	16
4	Comments on moduli stabilization	17
5	Gauge theories: regulating the Affleck-Dine-Seiberg runaway in AdS	22
A	Geometry of the enlarged target space \hat{X}	24

1 Introduction and conclusions

The considerations in this paper are driven by two general observations. First is the curious fact that large radius compactifications to four-dimensional anti-de Sitter space with $\mathcal{N} = 1$ supersymmetry very often come with anomalously light moduli whose masses scale with the AdS₄ curvature (see [1–3] for reviews of moduli stabilization scenarios and AdS₄ vacua in string theory). This is not a deep obstruction to phenomenological model building, as one can certainly give all moduli parametrically large masses by for example breaking supersymmetry or stepping away from a perturbative large-radius limit, which successful models such as KKLT [4] of course do. Nevertheless, it is instructive to take this issue seriously as it will reveal features of rigid sigma models in AdS₄, and their coupling to supergravity, that depart from naive flat-space intuition.

Second, it has recently been argued that consistent flat-space decoupling limits of supergravity lead to very special rigid supersymmetric theories [5]. More precisely, it has been argued that the target space of an $\mathcal{N} = 1$ sigma model in flat space must have an exact Kähler form in order to couple it to linearized supergravity; when applied to typical classes of string compactifications, this implies the inevitable existence of massless moduli in any smooth decoupling limit to flat space that preserves supersymmetry. It would be interesting to understand how and when such arguments apply in supersymmetric AdS₄ compactifications.

In this paper, we will argue that both of these properties — the ubiquity of light moduli in $\mathcal{N} = 1$ AdS₄ compactifications and the constraints on the topology and geometry of sigma models arising in the rigid limit of supergravity theories — follow from basic properties of supersymmetry in AdS₄. More precisely, we will study $\mathcal{N} = 1$ AdS₄ sigma models with Kähler target spaces X both as theories with global AdS₄ supersymmetry and as decoupling limits of consistent supergravity theories. As we shall explain, both of the above observations follow from simple, but surprisingly constraining, consistency conditions and kinematic properties of rigid AdS₄ sigma models.

Let us briefly summarize these constraints and properties. A supersymmetric sigma model in AdS₄ (with cosmological constant $-3\lambda^2$) is specified by a Kähler target space X with Kähler potential $K(\varphi, \bar{\varphi})$, and by a holomorphic superpotential $W(\varphi)$ governing relevant interactions. A key property of these rigid theories in AdS₄ is that, as in supergravity, the superpotential is not an invariant object but rather mixes with the Kähler potential under Kähler transformations,

$$K(\varphi, \bar{\varphi}) \rightarrow K(\varphi, \bar{\varphi}) + f(\varphi) + \bar{f}(\bar{\varphi}), \quad W(\varphi) \rightarrow W(\varphi) - \lambda f(\varphi).$$

The usual flat-space formula for the scalar potential, $g^{i\bar{j}}W_i\bar{W}_{\bar{j}}$, is not invariant under Kähler transformations, and must therefore be modified in AdS₄. Indeed, we find the form

$$V(\varphi, \bar{\varphi}) = g^{i\bar{j}}(W_i + \lambda K_i)(\bar{W}_{\bar{j}} + \lambda K_{\bar{j}}) - 3\lambda\bar{W} - 3\lambda W - 3\lambda^2 K.$$

The Kähler invariance of the action leads to a host of constraints on the possible form of the theory. In particular, supersymmetry further requires that the target X have a *trivial Kähler class*, $[\omega] = 0$. If we wish to build a sigma model starting with a manifold containing compact holomorphic cycles, we must fiber additional scalars over the geometry to trivialize all these cycles; otherwise, the theory itself will spontaneously decompactify them. If we started with a Hodge manifold then one such scalar would suffice, but generally we are forced to introduce $h^{1,1}(X)$ independent scalars.

Interestingly, the consistency conditions we find for the target space of a rigid $\mathcal{N} = 1$ sigma model in AdS₄ are (a) identical to those required for a rigid *flat-space* sigma model to arise as a decoupling limit of supergravity [5], and (b) imply the vanishing of all (mixed) gravitational anomalies upon coupling the rigid sigma model to $\mathcal{N} = 1$ supergravity. Thus, working in a non-trivial classical background makes this quantum constraint classically manifest, as in [6]. It is pleasing to see these constraints go over smoothly as $\lambda \rightarrow 0$.

A second surprising fact follows from the mixing of the Kähler potential and superpotential in rigid AdS₄: the right hand side of the supersymmetry variations are proportional not simply to W_i , as in flat space, but to $W_i + \lambda K_i$. Thus, even when the superpotential is zero, the vanishing of the fermion variations impose n equations on the n sigma model coordinates, implying that the supersymmetric vacua of this sigma model are generically a set of isolated points on X ¹ This differs sharply from familiar intuition from flat-space sigma models, where the moduli space for a vanishing superpotential is the full manifold, X . However, these two results pair naturally: away from these isolated points, the scalar

¹For non-generic models, there can be flat directions.

potential is non-zero but scales as λ — the supersymmetric points being gentle attractors — implying that we recover the expected moduli space in the flat-space limit.² It is useful to think of AdS_4 as a homogenous box inside of which massless modes are gapped: even if the target space is non-compact, the zero mode on the target feels a harmonic potential due to the constant negative curvature and is therefore massive.

The results above can be used to build a cartoon argument for the ubiquity of light moduli in large-radius compactifications respecting an unbroken $\mathcal{N} = 1$ supersymmetry. (Such $\mathcal{N} = 1$ compactifications play important roles in a variety of moduli stabilization scenarios, including for example the KKLT scenario [4], where the ensuing light moduli can subsequently be lifted by the supersymmetry-breaking uplifting stage.) Consider a large-radius flux compactification on some Calabi-Yau (or a decorated version thereof) to AdS_4 with suitably small cosmological constant, λ . By “large radius” we mean a manifold that is large compared to the four-dimensional Planck scale, allowing for a consistent perturbative expansion in $1/M_{\text{pl}}$ around the rigid limit; by “suitably small” we mean $\lambda \ll M_{\text{pl}}$.

At leading order in this perturbative expansion (i.e., in the decoupling limit), the results above tell us that there must be light fields in the theory with masses of order λ as long as supersymmetry is unbroken. Now, if perturbation theory in λ/M_{pl} is valid, leading corrections from supergravity will shift the masses of these light moduli by $\mathcal{O}\left(\frac{\lambda}{M_{\text{pl}}}\right)$ — i.e., not by very much. For these pesky moduli to be lifted above the AdS_4 scale, perturbation theory around large radius must *not* be reliable — we would need corrections that are non-perturbative in $1/M_{\text{pl}}$ — in which case we should not overcommit ourselves to perturbation theory around large radius with unbroken $\mathcal{N} = 1$ supersymmetry. This is, of course, what successful models of stabilization already do, for example by breaking supersymmetry and uplifting. In large volume compactifications of type IIB with $\mathcal{N} = 1$ AdS_4 vacua, we will show that the moduli are light with masses necessarily proportional to λ .

The fact that sigma models in AdS_4 have moduli spaces composed of isolated points suggests that the study of $\mathcal{N} = 1$ gauge theories may also considerably simplify in AdS_4 . We shall present a schematic argument to this effect by studying the example of $\text{SU}(N_c)$ SQCD with $N_f < N_c$ fundamental quarks in AdS_4 . In flat space, this theory famously suffers from an Affleck-Dine-Seiberg (ADS) runaway [7] in which various mesons run off to infinity in field space. In AdS_4 , however, the zero mode of a sigma model is generically lifted, suggesting that the ADS runaway may be lifted in AdS. We will marshal evidence for this picture, arguing that the meson field should be stabilized at a finite vev controlled by the ratio of the confinement scale Λ_c to the cosmological constant, $\langle m \rangle \sim \left(\frac{\Lambda_c}{\lambda}\right)^{\frac{1}{2}}$. A detailed study of $\mathcal{N} = 1$ gauge theories in AdS_4 is beyond the scope of this paper but is, clearly, of considerable interest.³

²Note that we can, of course, turn on additional superpotential terms to shift the masses of the light fields and shift around the supersymmetric points; we cannot, however, make the vanishing of the fermion variations a holomorphic condition, unlike in flat space, unless the Kähler potential is special Kähler (or simply trivial). This raises an interesting question about $\mathcal{N} = 2$ sigma models in AdS_4 , but that is beyond the scope of this discussion.

³See [8] for a discussion of Seiberg-Witten theory in AdS. We thank D. Tong for interesting and valuable discussions on these and related issues.

We thus see that simple consistency conditions for $\mathcal{N} = 1$ sigma models in AdS_4 lead to the constraints for weakly coupling to supergravity in the flat space limit [5], to the ubiquity of light moduli in large-radius $\mathcal{N} = 1$ compactifications to AdS_4 , and to a surprising set of features of rigid $\mathcal{N} = 1$ sigma models and gauge theories in AdS_4 . The remainder of the paper is devoted to deriving and explicating the above results, approaching them from two directions. First, in section 2, we study the structure of rigid $\mathcal{N} = 1$ sigma models in AdS_4 , deriving many of their properties directly and exploring a set of illustrative examples. We also show how the AdS_4 supersymmetric Lagrangian can be derived from supergravity through a decoupling limit. In section 3, we discuss the constraints on sigma models, particularly the triviality of the Kähler class, in the context of work by Bagger & Witten [9] and Komargodski & Seiberg [5] on the decoupling limits of supergravity theories. We also prove that the triviality of the Kähler class implies that there are no mixed gravitational anomalies when the sigma model is coupled to supergravity. The $[\omega] = 0$ constraint can, therefore, be derived by looking at mixed gravitational anomalies around a flat background, or through purely classical considerations of sigma models in an AdS_4 background. We suggest that this coincidence is not an accident but a consequence of a more general “Background Principle”, which we briefly discuss. In section 4, we present a simple argument for the existence of AdS_4 scale moduli in string compactifications at large volume that preserve supersymmetry; this suggests that one must break supersymmetry, or move away from large volume, in order to give large masses to the moduli, as is of course done in all phenomenologically successful models of moduli stabilization such as KKLT. We end in section 5 with a discussion of how the low-energy behavior of supersymmetric gauge theories, specifically the Affleck-Dine-Seiberg runaway, may be altered by AdS.

2 Supersymmetric Lagrangians for chiral multiplets in AdS_4

In this section, we construct the most general supersymmetric Lagrangian describing the interactions of chiral multiplets in AdS_4 . We show that the superpotential shifts under Kähler transformations and then derive various consequences from this fact: generally, there are no moduli spaces of supersymmetric vacua in AdS_4 ; the Kähler class of the target space must be trivial. We also present an alternate derivation of the sigma model Lagrangian as a decoupling limit of supergravity. For earlier work on rigid supersymmetric quantum field theories in AdS_4 , see [10–20].

2.1 The $\mathcal{N} = 1$ supersymmetric sigma model in Minkowski space

The well-known four-dimensional sigma model describes the general effective action (usually up to two derivatives) governing the interactions of massless scalar fields. It is given by a map $\varphi : M \rightarrow X$ from the spacetime M into a target space X , which is also equipped with a (positive-definite) metric g . Then the sigma model action reads

$$S_{\text{kin}} = f_\pi^2 \int d^4x \sqrt{-\gamma} \gamma^{mn} (\varphi^* g)(\partial_m, \partial_n), \quad (2.1)$$

which in local target-space coordinates, φ^I , corresponding to the (massless) scalar fields of the sigma model, has the familiar form

$$S_{\text{kin}} = f_\pi^2 \int d^4x \sqrt{-\gamma} g_{IJ}(\varphi) \partial_m \varphi^I \partial^m \varphi^J . \quad (2.2)$$

Here γ_{mn} is the spacetime metric with respect to the basis of tangent vectors ∂_m , and φ^*g denotes the pullback of the metric $g = g_{IJ} d\varphi^I d\varphi^J$ of the target space X to the spacetime M . There is a characteristic energy scale, f_π , that controls the strength of the scalar self-interactions. Due to unitarity constraints on low-energy scattering amplitudes (discussed, for example, in [21]), the effective action has a UV cutoff $\Lambda_\sigma \lesssim 4\pi f_\pi$.

The mass dimensions of the sigma model fields are $[\varphi^I] = 0$, $[\partial_m] = 1$, $[f_\pi] = 1$. The kinetic term (2.2) (without the normalization factor f_π^2) has mass dimension two. We can include interaction terms with mass dimension less than or equal to two, which are then relevant or marginal with respect to the scalar kinetic term (e.g., mass terms $(m^2)_{IJ} \varphi^I \varphi^J$).

To set the stage, and for later reference, we collect some well-known properties of the $\mathcal{N} = 1$ sigma model in Minkowski space $\mathbb{R}^{1,3}$, which describes the interactions of $\mathcal{N} = 1$ chiral multiplets. As before, the kinetic term for the bosonic scalars in the chiral multiplets is given by the action (2.1), but $\mathcal{N} = 1$ supersymmetry requires the target space X to be a complex Kähler manifold with Kähler metric g [22]. In local complex target space coordinates φ^i , which are identified with the complex scalars φ^i of the $\mathcal{N} = 1$ chiral multiplets, the bosonic kinetic term reads

$$S_{\text{kin}}^{\text{bos}} = f_\pi^2 \int d^4x g_{i\bar{j}}(\varphi, \bar{\varphi}) \partial_m \varphi^i \partial^m \bar{\varphi}^{\bar{j}} . \quad (2.3)$$

The Kähler potential $K(\varphi, \bar{\varphi})$ is a real function of the complex scalars φ , and it specifies locally the Kähler metric g and the Kähler (1,1)-form ω as

$$g_{i\bar{j}}(\varphi, \bar{\varphi}) = \frac{\partial^2}{\partial \varphi^i \partial \bar{\varphi}^{\bar{j}}} K(\varphi, \bar{\varphi}) , \quad \omega_{i\bar{j}}(\varphi, \bar{\varphi}) = i \frac{\partial^2}{\partial \varphi^i \partial \bar{\varphi}^{\bar{j}}} K(\varphi, \bar{\varphi}) . \quad (2.4)$$

Note that the Kähler metric g , the Kähler form ω , and consequently the supersymmetric sigma model itself, are invariant under Kähler transformations

$$K(\varphi, \bar{\varphi}) \rightarrow K(\varphi, \bar{\varphi}) + f(\varphi) + \bar{f}(\bar{\varphi}) , \quad (2.5)$$

for arbitrary holomorphic functions $f(\varphi)$.

The Kähler potential allows us to express the whole supersymmetric sigma model action (also including the fermionic terms) in global $\mathcal{N} = 1$ superspace [22]

$$S_{\text{kin}} = f_\pi^2 \int d^4x d^4\theta K(\Phi, \bar{\Phi}) . \quad (2.6)$$

Here, the arguments of the Kähler potential are the chiral superfields Φ associated to the complex scalars φ and the integral is taken over the whole $\mathcal{N} = 1$ superspace.

Relevant interactions in the $\mathcal{N} = 1$ supersymmetric sigma model (such as mass terms) are encoded in the superpotential P , which is a holomorphic function on the Kähler target space X . In the action, the superpotential P yields (locally) the bosonic interaction terms

$$S_{\text{int}}^{\text{bos}} = f_\pi^2 \int d^4x g^{i\bar{j}}(\varphi, \bar{\varphi}) P_i(\varphi) \bar{P}_{\bar{j}}(\bar{\varphi}) , \quad (2.7)$$

with $P_i \equiv \frac{\partial P}{\partial \varphi^i}$. These bosonic interactions pair with the fermionic terms and are conveniently expressed in terms of the $\mathcal{N} = 1$ superspace superpotential interaction

$$S_{\text{int}} = f_\pi^2 \int d^4x d^2\theta P(\Phi) + \text{c.c.} . \tag{2.8}$$

The Kähler potential $K(\varphi, \bar{\varphi})$, defined on a local patch in the target space X , need not be extendable to a function that is well-defined over the entire space X . Instead, in order to yield a globally well-defined Kähler metric g , the various local Kähler potentials K may differ by Kähler transformations (2.5) on overlapping regions of the local patches. In fact, the obstruction to extending a local Kähler potential to one defined over all of X (without the use of Kähler transformations on overlaps) is measured by the Dolbeault cohomology class in $H^{1,1}(X)$ of the Kähler (1,1)-form ω . Thus, unless the form ω is exact — i.e., $\omega = d\theta$ globally — the Kähler manifold X does not admit a globally defined Kähler potential K .

Note that an exact Kähler form has strong implications on the Kähler target space geometry and topology. Recall that compact holomorphic submanifolds of Kähler manifolds are calibrated by the Kähler class ω — i.e., the volume of a compact holomorphic submanifold S of complex dimension n is given by (e.g., [23])

$$\text{vol}(S) = \frac{1}{n!} \int_S \omega^n . \tag{2.9}$$

This innocent looking property has important consequences for Kähler manifolds with an exact Kähler form $\omega = d\theta$: exactness of the integrand implies that the integral of ω^n over any compact submanifold vanishes. For exact Kähler forms, the calibration condition (2.9) thus implies that the only compact holomorphic submanifolds are points (since any compact submanifold of dimension greater than zero must have a finite volume), a consequence of which is that *the Kähler manifold itself must be non-compact*.

2.2 Chiral multiplets in AdS₄

Consider AdS₄ with radius λ^{-1} , e.g., as a hyperboloid $-x_-^2 - x_0^2 + x_1^2 + x_2^2 + x_3^2 = -\frac{1}{\lambda^2}$ embedded in $\mathbb{R}^{2,3}$. The associated $\mathcal{N} = 1$ AdS₄ superalgebra, denoted by $osp(1,4)$,⁴ reads [10, 11, 13]

$$\begin{aligned} \{Q_\alpha, \bar{Q}_{\dot{\alpha}}\} &= -2\sigma_{\alpha\dot{\alpha}}^a R_a, & \{Q_\alpha, Q^\beta\} &= 2i\lambda(\sigma^{ab})_\alpha{}^\beta M_{ab}, \\ [R_a, Q_\alpha] &= -\frac{1}{2}\lambda(\sigma_a \bar{Q})_\alpha, & [M_{ab}, Q_\alpha] &= -i(\sigma_{ab})_\alpha{}^\beta Q_\beta, \\ [R_a, R_b] &= -i\lambda^2 M_{ab}, & [M_{ab}, R_c] &= i(\eta_{ac} R_b - \eta_{bc} R_a). \end{aligned} \tag{2.10}$$

The vector indices a, b, \dots , and the spinor indices $\alpha, \dot{\alpha}, \dots$, refer to the local Lorentz frame, and we use Wess and Bagger [22] two-component spinor notation throughout. Q_α and $\bar{Q}_{\dot{\alpha}}$ are the $\mathcal{N} = 1$ fermionic supersymmetry generators, whereas the bosonic generators R_a and M_{ab} generate AdS₄ translations and local Lorentz transformations, respectively. These

⁴We can, of course, remove λ from the supersymmetry algebra by a simple rescaling, $Q_\alpha \rightarrow \sqrt{\lambda} Q_\alpha$ and $R_a \rightarrow \lambda R_a$. We work with the chosen normalization to make the flat space limit manifest.

operators combine into the generators $(\lambda R_a, M_{ab})$, which are the generators of the group $SO(2,3)$ that acts on the AdS_4 hyperboloid. Note that in the limit $\lambda \rightarrow 0$, the generators R_a and M_{ab} become the usual translation and rotation operators of four-dimensional Minkowski space and the AdS_4 superalgebra (2.10) reduces to the familiar four-dimensional $\mathcal{N} = 1$ superalgebra of flat space.

Starting from the superalgebra (2.10), we can derive the representations on the fields following, for instance, [24]. The supersymmetry transformation of a field Φ is defined as

$$\delta_\xi \Phi(x) = -i[\xi Q + \bar{\xi} \bar{Q}, \Phi(x)] . \tag{2.11}$$

The chiral multiplet of AdS_4 supersymmetry is defined as a multiplet whose lowest component is a complex scalar that is annihilated by $\bar{Q}_{\dot{\alpha}}$, i.e., $[\bar{Q}_{\dot{\alpha}}, \varphi] = 0$. By acting on the lowest component with the “raising operator” Q_α , we can derive the other components of the multiplet and their supersymmetry transformations. As in flat space, the chiral multiplet consists of a complex scalar φ , a Weyl fermion χ , and a complex auxiliary field F . The transformation laws are listed below:

$$\begin{aligned} \delta_\xi \varphi^i &= \sqrt{2} \xi \chi^i , \\ \delta_\xi \chi^i &= \sqrt{2} F^i \xi + i\sqrt{2} \sigma^m \bar{\xi} \partial_m \varphi^i , \\ \delta_\xi F^i &= -\sqrt{2} \lambda \xi \chi^i + i\sqrt{2} \bar{\xi} \bar{\sigma}^m \nabla_m \chi^i . \end{aligned} \tag{2.12}$$

The algebra closes on these fields only if the supersymmetry parameter ξ satisfies the Killing spinor equation,

$$(\nabla_m \xi)^\alpha = \frac{i\lambda}{2} (\bar{\xi} \bar{\sigma}_m)^\alpha , \quad (\nabla_m \bar{\xi})_{\dot{\alpha}} = \frac{i\lambda}{2} (\xi \sigma_m)_{\dot{\alpha}} . \tag{2.13}$$

In the $\lambda \rightarrow 0$ limit, when the AdS_4 superalgebra reduces to the Poincaré superalgebra, the transformations (2.12) reduce to the usual supersymmetry transformations of a chiral multiplet in flat space, with the supersymmetry parameter ξ a constant spinor.

Note that the AdS_4 superalgebra (2.10) does not enjoy an R -symmetry. The trouble is particularly clear in the Killing spinor equation, (2.13), which relates ξ to $\bar{\xi}$ and thus does not allow a chiral rotation of ξ . However, as long as we have a consistent flat space limit, the broken R -symmetry can still be useful. To see this, imagine our rigid AdS_4 theory came from a decoupling limit of some $\mathcal{N} = 1$ supergravity (we discuss this in more detail in section 2.6). In this case, λ is the vev of the superpotential, $\lambda = M_{\text{pl}}^{-2} \langle P \rangle$. Since the superpotential carries R -charge 2, this vev breaks the R -symmetry. We may thus treat λ as an R -charge 2 spurion controlling the breaking of R -symmetry — indeed, this charge assignment restores the R -covariance of the killing spinor equation (2.13) while also forbidding the rescaling discussed above that removes λ from the supersymmetry algebra.

Just how far such a spurion analysis can take us is an extremely interesting question. For example, can we use λ to define an invariant holomorphic superpotential even away from a flat space limit? Can we use the would-be R -symmetry to constrain patterns of supersymmetry breaking, extending the arguments of Nelson and Seiberg [25] on spontaneous supersymmetry breaking in Wess-Zumino models to the AdS_4 context? Do standard

arguments, which rely heavily on (possibly anomalous) R -symmetries, extend to $\mathcal{N} = 1$ theories in AdS_4 ? For now, we simply note these questions and postpone a more detailed discussion to future work.⁵

2.3 AdS_4 supersymmetric sigma model

Quantum field theories in AdS_4 are naturally regulated in the IR by the AdS_4 scale λ (see [26] for an excellent discussion). In the remainder of this section, we focus on $\mathcal{N} = 1$ sigma models in AdS_4 , which also enjoy a UV cutoff, $4\pi f_\pi$, with a hierarchy thus given by

$$\lambda \ll \Lambda_\sigma \lesssim 4\pi f_\pi. \quad (2.14)$$

To derive the most general couplings of chiral multiplets in AdS_4 , we first write down all the possible terms with mass dimension no greater than two. The relevant assignments of the mass dimensions are given by

$$[\varphi^i] = 0, \quad [\chi^i] = \frac{1}{2}, \quad [F^i] = 1, \quad [\lambda] = 1, \quad [\partial_m] = 1. \quad (2.15)$$

Since in the $\lambda \rightarrow 0$ limit the AdS_4 superalgebra becomes the usual Poincaré superalgebra, we expect that the Lagrangian for the supersymmetric sigma model in AdS_4 smoothly goes over to the supersymmetric sigma model Lagrangian in flat space. The flat-space Lagrangian (including interaction terms induced from the superpotential $P(\varphi)$) is given by the superspace action (2.6) and (2.8), which yields the component expression [22]

$$\begin{aligned} \mathcal{L}(\lambda \rightarrow 0) = & g_{i\bar{j}} F^i \bar{F}^{\bar{j}} - g_{i\bar{j}} \partial_m \varphi^i \partial^m \bar{\varphi}^{\bar{j}} - i g_{i\bar{j}} \bar{\chi}^{\bar{j}} \bar{\sigma}^m \mathfrak{D}_m \chi^i - \frac{1}{2} P_{ij} \chi^i \chi^j - \frac{1}{2} \bar{P}_{\bar{i}\bar{j}} \bar{\chi}^{\bar{i}} \bar{\chi}^{\bar{j}} \\ & - F^i \left(\frac{1}{2} g_{i\bar{j},\bar{k}} \bar{\chi}^{\bar{j}} \bar{\chi}^{\bar{k}} - P_i \right) - \bar{F}^{\bar{i}} \left(\frac{1}{2} g_{j\bar{i},k} \chi^j \chi^k - \bar{P}_{\bar{i}} \right) + \frac{1}{4} g_{i\bar{j},k\bar{l}} \chi^i \chi^k \bar{\chi}^{\bar{j}} \bar{\chi}^{\bar{l}}. \end{aligned}$$

Here $\mathfrak{D}_m \chi^i \equiv \partial_m \chi^i + \partial_m \varphi^k \Gamma_{jk}^i \chi^j$ is the Kähler covariant derivative with respect to the Kähler target space connection and, as in (2.4), the Kähler metric can locally be derived from a Kähler potential K .

According to (2.15), all terms in the above Lagrangian have scaling dimension less than or equal to two. In the AdS_4 background, the dimensionful parameter λ allows for other relevant terms. Requiring that the Lagrangian has a smooth $\lambda \rightarrow 0$ limit, we arrive at the following ansatz

$$\begin{aligned} \mathcal{L}(\lambda) = & g_{i\bar{j}} F^i \bar{F}^{\bar{j}} - g_{i\bar{j}} \partial_m \varphi^i \partial^m \bar{\varphi}^{\bar{j}} - i g_{i\bar{j}} \bar{\chi}^{\bar{j}} \bar{\sigma}^m \mathfrak{D}_m \chi^i - \frac{1}{2} P_{ij} \chi^i \chi^j - \frac{1}{2} \bar{P}_{\bar{i}\bar{j}} \bar{\chi}^{\bar{i}} \bar{\chi}^{\bar{j}} \\ & - F^i \left(\frac{1}{2} g_{i\bar{j},\bar{k}} \bar{\chi}^{\bar{j}} \bar{\chi}^{\bar{k}} - P_i \right) - \bar{F}^{\bar{i}} \left(\frac{1}{2} g_{j\bar{i},k} \chi^j \chi^k - \bar{P}_{\bar{i}} \right) + \frac{1}{4} g_{i\bar{j},k\bar{l}} \chi^i \chi^k \bar{\chi}^{\bar{j}} \bar{\chi}^{\bar{l}} \\ & + \lambda (r + F^i t_i + \bar{F}^{\bar{i}} \bar{t}_{\bar{i}} + u_{ij} \chi^i \chi^j + \bar{u}_{\bar{i}\bar{j}} \bar{\chi}^{\bar{i}} \bar{\chi}^{\bar{j}}). \end{aligned} \quad (2.16)$$

The additional terms on the last line are all proportional to λ . The new parameters we have introduced, r, t_i, u_{ij} , are all functions of φ^i and $\bar{\varphi}^{\bar{j}}$. The parameter r is real with mass

⁵A.A. thanks J. Thaler for discussions on this and related topics.

dimension one, while t_i and u_{ij} are complex with mass dimension zero. Also note that in the χ kinetic term, the covariant derivative \mathfrak{D}_m now includes AdS₄ spin-connection in addition to the target-space connection.

We now demand that the above Lagrangian be invariant under the AdS₄ supersymmetry variations (2.12). Acting with these supersymmetry variations on the Lagrangian (2.16), we find,

$$\begin{aligned} \frac{1}{\sqrt{2}}\delta_\xi \mathcal{L} = & \lambda \left[\xi \chi^i (r_i - 3P_i - 3\lambda t_i) + (\xi \chi^i)(\bar{\chi}^{\bar{j}} \bar{\chi}^{\bar{k}}) \left(\frac{1}{2} g_{i\bar{j},\bar{k}} + \bar{u}_{\bar{j}\bar{k},i} \right) + (\xi \chi^i)(\chi^j \chi^k) u_{jk,i} \right. \\ & + \xi \chi^i F^j (t_{j,i} + 2u_{ij}) + \xi \chi^i \bar{F}^{\bar{j}} (-g_{i\bar{j}} + \bar{t}_{\bar{j},i}) \\ & \left. + i \xi \sigma^m \bar{\chi}^{\bar{i}} (\nabla_m \varphi^j (g_{j\bar{i}} - t_{j,\bar{i}}) + \nabla_m \bar{\varphi}^{\bar{j}} (\bar{t}_{\bar{j}} + 2\bar{u}_{i\bar{j}})) \right] + \text{c.c.} . \end{aligned} \quad (2.17)$$

In deriving this variation we have integrated by parts and used the Killing spinor equation (2.13).⁶ Note that all non-vanishing terms in the variation are proportional to λ since our Lagrangian was constructed to be supersymmetric in the limit $\lambda \rightarrow 0$. Requiring that the variation (2.17) vanishes for general λ then yields the conditions⁷

$$r = 3(W(\varphi) + \bar{W}(\varphi) + \lambda K(\varphi, \bar{\varphi})), \quad u_{ij} = -\frac{1}{2}(K_{ij}(\varphi, \bar{\varphi}) + \Delta_{ij}(\varphi)), \quad t_i = (K_i(\varphi, \bar{\varphi}) + \Delta_i(\varphi)), \quad (2.18)$$

where $\Delta(\varphi)$ is an undetermined holomorphic function and where we have defined the ‘‘AdS₄ superpotential’’

$$W(\varphi) = P(\varphi) + \lambda \Delta(\varphi), \quad (2.19)$$

which is convenient since the action can be written only in terms of W , with no reference to P or Δ separately.

Inserting the conditions (2.18) into the ansatz (2.16), we arrive at the supersymmetric AdS₄ Lagrangian

$$\begin{aligned} \mathcal{L}(\lambda) = & -g_{i\bar{j}} \partial_m \varphi^i \partial^m \bar{\varphi}^{\bar{j}} - i g_{i\bar{j}} \bar{\chi}^{\bar{m}} \mathfrak{D}_m \chi^i + g_{i\bar{j}} F^i \bar{F}^{\bar{j}} - F^i \left(\frac{1}{2} g_{i\bar{j},\bar{k}} \bar{\chi}^{\bar{j}} \bar{\chi}^{\bar{k}} - (W_i + \lambda K_i) \right) \\ & - \bar{F}^{\bar{i}} \left(\frac{1}{2} g_{j\bar{i},k} \chi^j \chi^k - (\bar{W}_{\bar{i}} + \lambda K_{\bar{i}}) \right) + \frac{1}{4} g_{i\bar{j},k\bar{l}} \chi^i \chi^k \bar{\chi}^{\bar{j}} \bar{\chi}^{\bar{l}} - \frac{1}{2} (W_{ij} + \lambda K_{ij}) \chi^i \chi^j \\ & - \frac{1}{2} (\bar{W}_{\bar{i}\bar{j}} + \lambda K_{\bar{i}\bar{j}}) \bar{\chi}^{\bar{i}} \bar{\chi}^{\bar{j}} + 3\lambda \bar{W} + 3\lambda W + 3\lambda^2 K . \end{aligned} \quad (2.20)$$

The equation of motion for the auxiliary field F^i is

$$g_{i\bar{i}} F^i = \left(\frac{1}{2} g_{i\bar{m}} \Gamma_{jk}^m \chi^j \chi^k - (\bar{W}_{\bar{i}} + \lambda K_{\bar{i}}) \right), \quad (2.21)$$

⁶Throughout this paper, we will be cavalier about the boundary conditions satisfied by our fields. At the present step, we do not expect any obstructions to finding boundary conditions that are compatible with the supersymmetry variations described. In other steps, however, we must be more careful since the necessity of choosing certain boundary conditions can be surprisingly restrictive; for example, they can be shown to forbid the existence of charged chiral fermions in AdS₄ [18, 19, 26]. In general, a careful treatment of boundary conditions, such as in [27], is an important additional step which we defer to future work.

⁷Another consequence is that the functions $u_{jk,i}$ are symmetric with respect to all their indices. Therefore, the term $(\xi \chi^i)(\chi^j \chi^k) u_{jk,i}$ in (2.17) automatically vanishes due to a Fierz identity.

so we can integrate it out to obtain

$$\begin{aligned} \mathcal{L}(\lambda) = & -g_{i\bar{j}}\partial_m\varphi^i\partial^m\bar{\varphi}^{\bar{j}} - ig_{i\bar{j}}\bar{\chi}^{\bar{j}}\bar{\sigma}^m\mathfrak{D}_m\chi^i - g^{i\bar{i}}(W_i + \lambda K_i)(\bar{W}_{\bar{i}} + \lambda K_{\bar{i}}) + 3\lambda\bar{W} + 3\lambda W + 3\lambda^2 K \\ & - \frac{1}{2}\mathfrak{D}_i(W_j + \lambda K_j)\chi^i\chi^j - \frac{1}{2}\bar{\mathfrak{D}}_{\bar{i}}(\bar{W}_{\bar{j}} + \lambda K_{\bar{j}})\bar{\chi}^{\bar{i}}\bar{\chi}^{\bar{j}} + \frac{1}{4}\mathcal{R}_{i\bar{j}k\bar{l}}\chi^i\chi^k\bar{\chi}^{\bar{j}}\bar{\chi}^{\bar{l}}. \end{aligned} \quad (2.22)$$

The scalar potential, which is simply $g^{i\bar{j}}W_i\bar{W}_{\bar{j}}$ in the flat space case, is

$$V(\varphi, \bar{\varphi}) = g^{i\bar{i}}(W_i + \lambda K_i)(\bar{W}_{\bar{i}} + \lambda K_{\bar{i}}) - 3\lambda\bar{W} - 3\lambda W - 3\lambda^2 K. \quad (2.23)$$

The resulting Lagrangians (2.20) and (2.22) are not invariant under a Kähler transformation of the Kähler potential alone, but they are invariant if supplemented with a shift of the AdS₄ superpotential:

$$K(\varphi, \bar{\varphi}) \rightarrow K(\varphi, \bar{\varphi}) + f(\varphi) + \bar{f}(\bar{\varphi}), \quad W(\varphi) \rightarrow W(\varphi) - \lambda f(\varphi). \quad (2.24)$$

As a consequence, the undetermined function $\Delta(\varphi)$ in (2.18) could be absorbed, through a Kähler transformation, into the Kähler potential. However, for a particular Kähler potential K (chosen in the flat space limit $\lambda \rightarrow 0$), there are distinct choices to extend the supersymmetric flat space Lagrangian to inequivalent supersymmetric AdS₄ Lagrangians. For a fixed Kähler potential K , these choices are distinguished by the holomorphic shift Δ in the AdS₄ superpotential W .

2.4 Supersymmetric vacua in AdS₄

The supersymmetric vacua of this theory, preserving the full AdS₄ invariance, satisfy

$$Q_\alpha|0\rangle = 0 \iff \langle \delta\chi^i \rangle = 0 \iff F^i = 0 \iff W_i + \lambda K_i = 0. \quad (2.25)$$

These constitute $n = \dim(X)$ equations for n complex variables and generically result in a *discrete* set of vacua. Even when $W \equiv 0$, the moduli space of an $\mathcal{N} = 1$ sigma model in AdS₄ is *not* a copy of the target space X , but rather a set of isolated points *on* X , with the masses of generic scalar fields being proportional to λ . Since we have assumed that $\lambda \ll 4\pi f_\pi$ is an IR scale, the fields are still “light”. The conditions for a supersymmetric vacuum in AdS₄ were first written down and analyzed in [12] for the Wess-Zumino model, and for general supersymmetric sigma models in [20].

The fact that the would-be moduli are lifted in AdS₄ should not come as a surprise. In gravity, AdS₄ is often described as a homogeneous box for gravity, with light traveling to the boundary and back in finite observer time. Importantly, this is a good description whether the metric is dynamical or not. The fact that the spectrum of our sigma model is gapped in AdS₄ with the moduli space generically reduced to a set of isolated points is simply a reflection of the effectively compact nature of AdS₄.

Note that the vacuum energy at a supersymmetric vacuum in AdS₄ need not vanish. The scalar potential (2.23) at a supersymmetric vacuum is

$$V|_{\text{susy}} = -3\lambda(W + \bar{W} + \lambda K), \quad (2.26)$$

where the quantities on the right hand side are evaluated at the supersymmetric vacuum. Different supersymmetric vacua of the same theory can, in general, have different vacuum energies in AdS₄. This is another difference from Poincaré supersymmetry where all supersymmetric vacua have identical vacuum energies, conventionally taken to be zero. This is not a problem since the absolute energy of a state carries no invariant meaning in the absence of gravity. In weakly coupling an AdS₄ supersymmetric theory to gravity, however, the vacuum energy of the globally supersymmetric sector results in a tadpole for the metric fluctuation. The AdS₄ scale receives an M_{pl} suppressed correction,

$$\delta\lambda^2 = \frac{V|_{\text{susy}}}{M_{\text{pl}}^2}, \tag{2.27}$$

which can be disregarded in the limit of weak gravity. Furthermore, in the rigid limit the vacuum energy can be absorbed into a constant shift of the superpotential,

$$W \rightarrow W + \frac{1}{6\lambda} V|_{\text{susy}}. \tag{2.28}$$

With this freedom, we can set the vacuum energy of any given supersymmetric vacuum to zero and then weakly couple to supergravity to avoid the gravitational tadpole.

Example. $X = \mathbb{C}$, $K(\varphi, \bar{\varphi}) = \bar{\varphi}\varphi$.

The Lagrangian in the simple case of a free chiral multiplet with no superpotential is

$$\mathcal{L} = -\partial_m\varphi\partial^m\bar{\varphi} - i\bar{\chi}\bar{\sigma}^m\nabla_m\chi + 2\lambda^2\varphi\bar{\varphi}. \tag{2.29}$$

We see that even in the absence of a superpotential, the complex scalar has a tachyonic mass $m^2 = -2\lambda^2$, which is above the Breitenlohner-Freedman bound $-(9/4)\lambda^2$ for AdS₄ [14]. In the flat space case, the free complex scalar has a moduli space of vacua parameterized by $\langle\varphi\rangle$. In the AdS₄ case, we see that there is a single vacuum at $\langle\varphi\rangle = 0$.

2.5 Target space geometry for $\mathcal{N} = 1$ sigma models in AdS₄

Thus far we have examined the local structure of supersymmetric AdS₄ sigma models, so let us now consider their global properties. In this regard, the Kähler transformation (2.24) of the superpotential has crucial implications on the global structure of the Kähler target space geometry X : the superpotential W can only be extended to a non-singular holomorphic superpotential on the whole Kähler target space X if the target space Kähler form ω is cohomologically trivial, $[\omega] = 0$.⁸ To argue this formally, we cover the Kähler target space by holomorphic patches U_α decorated with local Kähler potentials K_α and local (holomorphic) superpotentials W_α in each patch. On double overlaps of two patches, α and β , the holomorphic Kähler transition functions obey $\lambda f_{\alpha\beta}(\varphi) = W_\beta(\varphi) - W_\alpha(\varphi)$, in

⁸The easiest way to see this is to note that the combination $K + \lambda^{-1}(W + \bar{W})$ is a function on all of X that is also a potential for the Kähler form ω . There are therefore two options: first, the Kähler form is exact; second, the combination $K + \lambda^{-1}(W + \bar{W})$ has branch cuts or singularities, necessarily creating divergences in the scalar potential (2.23) that effectively decompactify the target space by confining the scalars to the regions away from infinite potential energy.

accordance with (2.24), implying that the Čech cocycle $(\delta f)_{\alpha\beta\gamma}$ vanishes identically on all triple overlaps,

$$(\delta f)_{\alpha\beta\gamma} = f_{\alpha\beta} - f_{\alpha\gamma} + f_{\beta\gamma} = 0 .$$

If the Kähler form is non-trivial, the associated Čech cocycle $(\delta f)_{\alpha\beta\gamma}$ cannot vanish identically on all triple overlaps [9]. We thus conclude that the Kähler form ω must be cohomologically trivial for rigid supersymmetric AdS₄ sigma models.

Example. $X = \mathbb{P}^1$, Fubini-Study metric.

To illustrate this point, we attempt to construct a supersymmetric sigma model in AdS₄ with target space $X = \mathbb{P}^1$, endowed with the Fubini-Study Kähler metric. Covering the target space \mathbb{P}^1 with the two patches $U_0, U_\infty \cong \mathbb{C}$ of the local coordinates z and w that arise from the stereographic projection of \mathbb{P}^1 , we have the Kähler metric and potentials

$$\begin{aligned} z \in U_0 : \quad g_{z\bar{z}} &= \frac{n}{(1+z\bar{z})^2} = \partial_z \partial_{\bar{z}} K_0(z, \bar{z}), & K_0(z, \bar{z}) &= n \log(1+z\bar{z}), \\ w \in U_\infty : \quad g_{w\bar{w}} &= \frac{n}{(1+w\bar{w})^2} = \partial_w \partial_{\bar{w}} K_\infty(w, \bar{w}), & K_\infty(w, \bar{w}) &= n \log(1+w\bar{w}). \end{aligned} \quad (2.30)$$

The (non-trivial) Kähler form $\omega = i\partial\bar{\partial}K$ generates $H^{1,1}(\mathbb{P}^1)$ with $\int_{\mathbb{P}^1} \omega = 2\pi n$. On the overlap $U_0 \cap U_\infty$ (where $z = w^{-1}$), there is a non-trivial Kähler transformation $f_{\infty 0}$

$$K_\infty(z^{-1}, \bar{z}^{-1}) - K_0(z, \bar{z}) = f_{\infty 0}(z) + \bar{f}_{\infty 0}(\bar{z}), \quad f_{\infty 0}(z) = -n \log z = n \log w . \quad (2.31)$$

Due to (2.24) we know that the superpotential must obey

$$W_\infty(z^{-1}) - W_0(z) = -\lambda f_{\infty 0}(z) = \lambda n \log z . \quad (2.32)$$

For the local superpotential $W_0(z) \equiv 0$, we arrive at the (bosonic) Lagrangian in terms of the chiral field z in the patch U_0

$$\mathcal{L}_{U_0 \subset \mathbb{P}^1} = -n \frac{\partial_m z \partial^m \bar{z}}{(1+z\bar{z})^2} - \lambda^2 (n z \bar{z} - 3n \log(1+z\bar{z})) , \quad (2.33)$$

with supersymmetric vacuum for $\langle z \rangle = 0$. In the U_∞ patch, with the (singular) superpotential $W_\infty(w) = -n\lambda \log w$, the Lagrangian becomes

$$\mathcal{L}_{U_\infty \subset \mathbb{P}^1} = -n \frac{\partial_m w \partial^m \bar{w}}{(1+w\bar{w})^2} - \lambda^2 \left(\frac{n}{w\bar{w}} + 3n \log \left(1 + \frac{1}{w\bar{w}} \right) \right) . \quad (2.34)$$

Note that due to the infinity of the superpotential W_∞ and the resulting infinity in the scalar potential at $w = 0$, a fluctuation in the neighborhood around $w = 0$ has arbitrarily high energy. Therefore removing all fluctuations above a certain UV cutoff effectively removes the point $w = 0$ (or $z = \infty$) from the target space $X = \mathbb{P}^1$. Thus, the resulting effective AdS₄ sigma model has the trivialized target space $\mathbb{C} \simeq \mathbb{P}^1 \setminus \{w = 0\}$ with a cohomologically trivial Kähler form $\omega|_{U_0}$.

2.6 Derivation through supergravity

We have shown that the target space X of a rigid $\mathcal{N} = 1$ sigma model in AdS_4 has an exact Kähler form. Since this is also true for sigma models in supergravity that admit weak-coupling limits [5, 9], this gives us an easy route to the AdS_4 sigma model Lagrangian (2.22) via a decoupling limit of supergravity. As we shall now verify, this approach leads to precisely the same theories and the same constraints as found above through a direct analysis of the rigid supersymmetry algebra.

We start with the Lagrangian for supergravity coupled to a set of chiral multiplets that parameterize a Kähler target space X (e.g., [22]). For the present argument it suffices to restrict attention to the bosonic terms of the $\mathcal{N} = 1$ supergravity action, but the derivation goes through for the fermionic terms as well. The general form of the (bosonic) $\mathcal{N} = 1$ supergravity action of interacting chiral multiplets is completely determined in terms of the (local) Kähler potential $K(\varphi, \bar{\varphi})$ and the superpotential P

$$\begin{aligned}
 S_{\text{sugra}}^{\text{bos}} &= - \int d^4x \sqrt{-\gamma} \left[\frac{1}{2} M_{\text{pl}}^2 R + g_{i\bar{j}} \partial_m \varphi^i \partial^m \bar{\varphi}^{\bar{j}} + V(\varphi, \bar{\varphi}) \right], \\
 V_{\text{sugra}}(\varphi, \bar{\varphi}) &= e^{K/M_{\text{pl}}^2} \left[g^{i\bar{j}} \left(P_i + \frac{K_i}{M_{\text{pl}}^2} P \right) \left(\bar{P}_{\bar{j}} + \frac{K_{\bar{j}}}{M_{\text{pl}}^2} \bar{P} \right) - \frac{3}{M_{\text{pl}}^2} |P|^2 \right],
 \end{aligned} \tag{2.35}$$

where M_{pl} is the four-dimensional Planck mass. The mass dimensions of the fields are chosen as in (2.15), and the Kähler potential K and the superpotential P have dimensions two and three, respectively.

The dynamics of the sigma model are controlled by the cutoff scale Λ_σ and various mass scales $\mu \lesssim \Lambda_\sigma$ that appear in the superpotential. In order to decouple gravity, we assume that all these scales are much smaller than the Planck scale M_{pl} . This prevents, for example, low-energy scattering of sigma model fields from producing final state gravitons, as the graviton coupling (via the stress tensor) is suppressed by inverse powers of M_{pl} . Graviton self-interactions are also suppressed in the low-energy regime. The dynamics at energies $E \ll \Lambda_\sigma \lesssim 4\pi f_\pi$ therefore consists of a sector of noninteracting soft gravitons and a decoupled sector of interacting sigma model fields.

The metric satisfies the Einstein equations and is sourced by the stress-energy tensor of the matter sector. If we assume that all the energy scales associated to the sigma model sector are small, as we argued in the previous paragraph, the metric equation of motion would yield Minkowski space as a solution. In order to obtain AdS_4 , we introduce a constant term to the supergravity superpotential⁹

$$P_{\text{sugra}}(\varphi) = \lambda M_{\text{pl}}^2 + W(\varphi), \tag{2.36}$$

which gives rise to a negative cosmological constant in the supergravity scalar potential. The quantity W will turn out to be the superpotential that appears in the global supersymmetric AdS_4 sigma model Lagrangian. With the addition of the constant term, the scalar potential in an M_{pl}^{-2} expansion takes the form

$$V_{\text{sugra}} = -3M_{\text{pl}}^2 \lambda^2 + \left(g^{i\bar{j}} (W_i + \lambda K_i) (\bar{W}_{\bar{j}} + \lambda K_{\bar{j}}) - 3\lambda (W + \bar{W} + \lambda K) \right) + \mathcal{O}\left(\frac{1}{M_{\text{pl}}^2}\right) \tag{2.37}$$

⁹Even in the absence of a matter sector we may introduce a constant supergravity superpotential $P = \lambda M_{\text{pl}}^2$, describing supergravity in a background with a negative cosmological constant.

The metric equation of motion, to leading order in M_{pl} , is

$$R_{mn} = 3\lambda^2 \gamma_{mn}, \tag{2.38}$$

which gives AdS₄ with the radius λ^{-1} . The terms at order M_{pl}^0 above agree precisely with the AdS₄ scalar potential in (2.23). Moreover, the complete Lagrangian (2.22), including all the fermion terms, can be reproduced through the decoupling procedure we have described.

The supergravity action is invariant under Kähler-Weyl transformations

$$K \rightarrow K + f(\varphi) + \bar{f}(\bar{\varphi}), \quad P \rightarrow \exp(-f(\varphi)/M_{\text{pl}}^2) P, \tag{2.39}$$

where the fermions are also rotated by a phase dependent upon $\text{Im}(f)$. Note that the supergravity superpotential P transforms as a holomorphic section of a line bundle over the target space [9], which is necessary in order for the scalar potential $V(\varphi, \bar{\varphi})$ to remain invariant. The modified Kähler invariance (2.24) of the AdS₄ Lagrangian can be derived in the gravity decoupling limit from the supergravity Kähler invariance above. Heuristically, we assume that the mass scales of the Kähler transformation functions $f(\varphi)$, which are of mass dimension two in our conventions, are all associated with the sigma model scales and are much smaller than M_{pl} . This is a somewhat vague restriction on the class of allowed Kähler transformations and it can be made mathematically more precise. Here we avoid presenting all the necessary technical details, as it is intuitive — due to the mentioned separation of scales — that the Kähler transformation of $P = \lambda M_{\text{pl}}^2 + W$, (2.39), expanded in powers of M_{pl} , yields the AdS₄ Kähler transformations (2.24).

3 Lessons for constraints in flat space and beyond

In the previous section, we studied the consistency conditions required for $\mathcal{N} = 1$ supersymmetry of sigma models in AdS₄. It is instructive to consider these results in relation to the consistency conditions for $\mathcal{N} = 1$ sigma models in flat space, as well as to general results on supergravity in four dimensions.

In flat space, rigid $\mathcal{N} = 1$ supersymmetry does not impose any topological conditions on the cohomology class of the Kähler form, nor does the superpotential transform under Kähler transformations. When supersymmetry is gauged, on the other hand, the work of Bagger and Witten [9] shows that the target space must have an even integral Kähler class ω in $H^2(X, \mathbb{Z})$ (normalized by M_{pl}). Therefore when the Kähler form ω is not exact, the dimensionful irrelevant couplings of the sigma model that are associated to the non-trivial holomorphic cycles of the Kähler target space are necessarily quantized in units of M_{pl} . For example, when $X = \mathbb{P}^1$ the sigma model is characterized by a single dimensionful scale f_π , so the Bagger-Witten analysis shows that the ratio $\frac{f_\pi^2}{M_{\text{pl}}^2}$ is an even integer. We cannot, in such a situation, dial the scales f_π and M_{pl} independently. As a result of this quantization, only Planck-scale experiments would be able to probe the curvature of the \mathbb{P}^1 target space, or the irrelevant interactions specific to the \mathbb{P}^1 sigma model. At energies much lower than the Planck scale, the sigma model is essentially trivialized to a local patch $\mathbb{C} \subset \mathbb{P}^1$. Such field theories do not have the interpretation of being weakly coupled to gravity and are intrinsically gravitational.

This raises the interesting question of which rigid supersymmetric field theories can be weakly coupled to gravity. Komargodski and Seiberg recently approached this question by studying the conditions under which the stress tensor and supercurrent could fit into a single flat-space supersymmetry multiplet [5] (see [28] for a very recent extension of these arguments to AdS₄). They showed that the standard Ferrara-Zumino multiplet [29] is only globally well-defined when the Kähler class of X , $[\omega]$, is trivial. When $[\omega] \neq 0$, one can find a different set of supersymmetric current multiplets that can be coupled to supergravity with an additional linear multiplet. When the target space X is a Hodge manifold, this additional linear multiplet can be dualized to a chiral superfield such that the enlarged geometry \hat{X} has a trivial Kähler class. The enlarged geometry \hat{X} is then a \mathbb{C}^* -fiber bundle over X of the form discussed in appendix A.

As is by now clear, these conditions are all equivalent to the conditions for unbroken $\mathcal{N} = 1$ supersymmetry in an AdS₄ background. Remarkably, these conditions also imply the vanishing of (mixed) gravitational anomalies, even around flat space, as we will shortly explain. The remainder of this section is devoted to explaining this connection with anomalies and expanding this lesson into a general conjecture about the conditions for a general rigid $\mathcal{N} = 1$ theory to arise as the decoupling limit of some $\mathcal{N} = 1$ supergravity.

3.1 Anomalies and constraints on (de-)coupling gravity

As we found in section 2.6, the classical conditions for rigid $\mathcal{N} = 1$ supersymmetry in AdS₄ spacetimes are equivalent to the conditions for the rigid theory to arise as a decoupling limit of $\mathcal{N} = 1$ supergravity. However, since the supergravity multiplet contains a gravitino, any such decoupling limit alters the chiral spectrum. It is thus possible that decoupling (or re-coupling!) gravity, while classically straightforward, is quantum-mechanically obstructed by anomalies in either the global or local supersymmetric theory. In the remainder of this section, we shall check for such potential obstructions to (de-)coupling gravity and our sigma model by studying the possible mixed-gravitational anomalies in both local and global theories.¹⁰

In a globally supersymmetric $\mathcal{N} = 1$ sigma model $\varphi : \Sigma \rightarrow X$ from a four-dimensional spacetime background Σ into a Kähler target space X , the fermions χ in the chiral multiplets transform as spinor-valued sections of the pullback tangent bundle φ^*TX . As a consequence, the global $\mathcal{N} = 1$ sigma model is anomaly-free when the six-form anomaly polynomial vanishes [30],

$$P_{\text{global}}^{(\Sigma, X)} = \hat{A}(\Sigma) \wedge \tilde{\varphi}^* \text{ch}(X) \Big|_{(6\text{-form})} = \tilde{\varphi}^* \text{ch}_3(X) - \frac{1}{24} \tilde{\varphi}^* c_1(X) \wedge p_1(\Sigma) . \quad (3.1)$$

Here $\text{ch}(X)$ denotes the total Chern character of the target space manifold X , and $\hat{A}(\Sigma)$ and $p_1(\Sigma)$ are the A -roof genus and the first Pontryagin class of the spacetime, Σ , respectively.¹¹

¹⁰Note that, in general, this analysis should include a careful discussion of potential boundary terms which can have important effects on the chiral spectrum. For example, it is impossible to find boundary conditions which allow chiral matter to couple to a massless gauge field in AdS₄.

¹¹ $\hat{A}(\Sigma) = 1 - \frac{p_1(\Sigma)}{24}$ for four-dimensional manifolds Σ .

The map $\tilde{\varphi}$ is directly related to the sigma model map φ , for a detailed definition of which we refer the reader to [30].

For the $\mathcal{N} = 1$ supersymmetric sigma model $\varphi : \Sigma \rightarrow X$ in four spacetime dimensions coupled to gravity, chiral fermions are spinor-valued sections of the bundle $\varphi^*(TX \otimes \mathcal{K})$, where \mathcal{K} is the Kähler line bundle obeying $c_1(\mathcal{K}) \simeq \frac{1}{2}\omega$ and ω is the Kähler form of the target space X .¹² In addition, the gravitino ψ_μ transforms as a spinor-valued section of $(T\Sigma \ominus \mathbf{1}) \otimes \varphi^*\mathcal{K}^{-1}$ [31]. Therefore the resulting six-form anomaly polynomial for the $\mathcal{N} = 1$ supersymmetric sigma model coupled to gravity reads [31]

$$\begin{aligned}
 P_{\text{local}}^{(\Sigma, X)} &= \hat{A}(\Sigma) \wedge \left[\text{ch } \tilde{\varphi}^*(TX \otimes \mathcal{K}) - \text{ch} \left((T\Sigma \ominus \mathbf{1}) \otimes \varphi^*\mathcal{K}^{-1} \right) \right] \Big|_{(6\text{-form})} \\
 &= \tilde{\varphi}^* \text{ch}_3(X) - \frac{1}{24} \tilde{\varphi}^* c_1(X) \wedge p_1(\Sigma) + \tilde{\varphi}^* c_1(\mathcal{K}) \left(\tilde{\varphi}^* \text{ch}_2(X) + \frac{21-n}{24} p_1(\Sigma) \right) \\
 &\quad + \frac{1}{2} \tilde{\varphi}^* c_1(\mathcal{K})^2 \wedge \tilde{\varphi}^* c_1(X) + \frac{n+3}{6} \tilde{\varphi}^* c_1(\mathcal{K})^3 .
 \end{aligned} \tag{3.2}$$

Here n is the complex dimension of the target space manifold X .

The important observation is now that the local anomaly $P_{\text{local}}^{(\Sigma, X)}$ decomposes as

$$\begin{aligned}
 P_{\text{local}}^{(\Sigma, X)} &= P_{\text{global}}^{(\Sigma, X)} + \Delta P^{(\Sigma, X)} , \\
 \Delta P^{(\Sigma, X)} &= \tilde{\varphi}^* c_1(\mathcal{K}) \left[\left(\tilde{\varphi}^* \text{ch}_2(X) + \frac{21-n}{24} p_1(\Sigma) \right) \right. \\
 &\quad \left. + \frac{1}{2} \tilde{\varphi}^* c_1(\mathcal{K}) \wedge \tilde{\varphi}^* c_1(X) + \frac{n+3}{6} \tilde{\varphi}^* c_1(\mathcal{K})^2 \right] ,
 \end{aligned} \tag{3.3}$$

where the contribution $\Delta P^{(\Sigma, X)}$ is proportional to $c_1(\mathcal{K})$ and therefore to the Kähler class $[\omega]$ of the target space X . Thus, if the target space X has a cohomologically trivial Kähler form, then the process of weakly coupling to gravity does not change the sigma model anomaly. In particular, if the global $\mathcal{N} = 1$ sigma model is anomaly free then the addition of gravity does not introduce an additional anomaly.

For global $\mathcal{N} = 1$ supersymmetric AdS₄ sigma models, we observed in section 2.5 that the target space Kähler form must be cohomologically trivial. Thus if the global $\mathcal{N} = 1$ supersymmetric AdS₄ sigma model (of chiral multiplets) is free of anomalies, then there are no further anomaly constraints in coupling to gravity in an AdS₄ background. We conclude then that the vanishing of gravitational anomalies is already guaranteed by the *classical* consistency of these models!

3.2 The background principle

It is quite a remarkable fact that the analysis of section 2, which is completely classical, implies the vanishing of quantum anomalies above, i.e. that the Kähler form must be exact. There is a simple reason for this: the (mixed) gravitational anomalies tell us the conditions for consistently coupling a microscopic theory to gravity. Any theory that

¹²In $\mathcal{N} = 1$ supergravity, the target X must be a Hodge manifold with an even integral Kähler form ω [9].

can be consistently coupled to gravity should also be able to be expanded around a non-trivial metric which solves the equations of motion and preserves the same symmetries. For this purpose, AdS_4 is peculiarly well-suited as it is maximally symmetric, preserves supersymmetry, and arises as a one-parameter deformation of the theory in flat space — including the supersymmetry algebra and its representation theory. We thus expect the classical conditions for $\mathcal{N} = 1$ supersymmetry in AdS_4 to correspond to necessary conditions for consistently coupling the flat-space theory to supergravity, which is exactly what we found.

Similar effects obtain in other contexts. For example, consider the bounds on the signs of leading irrelevant operators as discussed in [6]. These bounds can be identified in two ways: when expanding around the trivial vacuum, these constraints can only be seen from a dispersion relation for the quantum S -matrix elements; on the other hand, when expanding about a suitable classical background, these constraints are visible classically and at low energies. Roughly speaking, working in a non-trivial background takes a microscopic (quantum) effect and exponentiates it via multiple scattering off the classical background.

All of this entices us to make a more general conjecture, which we will call the “Background Principle”: any rigid $\mathcal{N} = 1$ theory in Minkowski space which can be consistently, quantum-mechanically coupled to $\mathcal{N} = 1$ supergravity — or, conversely, which can arise as the decoupling limit of a well-defined $\mathcal{N} = 1$ supergravity theory — must also behave smoothly, *as a classical theory*, under a deformation of the rigid Minkowski spacetime to AdS. In this paper, we have shown this to be the case for conventional sigma models containing only chiral superfields; we conjecture this to be true for all rigid $\mathcal{N} = 1$ theories.

If true, this principle affords both a straightforward route to identifying four-dimensional QFTs which cannot be coupled to supergravity, and leads to a strong statement about the moduli spaces of theories which can be coupled to supergravity and admit a UV completion: their $\lambda \rightarrow 0$ moduli spaces, governed by long-distance sigma models, must necessarily be non-compact lest supersymmetry be broken when expanding about a rigid AdS_4 background.

4 Comments on moduli stabilization

We use the $\mathcal{N} = 1$ AdS_4 sigma model to study the moduli sector of a large class of string compactifications, concluding, on general grounds, that these compactifications necessarily have moduli whose masses are proportional to the AdS_4 scale. Such $\mathcal{N} = 1$ compactifications arise, for example, in the first stages of KKLT/KKLMNT-type scenarios [4, 32], where the resulting light moduli can be dealt with by a supersymmetry-breaking uplift. The lesson of our analysis, which depends only on simple properties of $\mathcal{N} = 1$ sigma models in AdS_4 , is that such light moduli arise very generally in a model independent fashion, and that lifting them requires either moving away from large volume or breaking supersymmetry, as in the specific scenarios of [4, 32].

We begin with a lightning review of type IIB supersymmetric AdS_4 flux vacua (see [1–3] for a general review of various moduli stabilization scenarios). In the large volume regime of type IIB Calabi-Yau compactifications, the moduli fields can heuristically be divided into

the Kähler and complex structure moduli of the Calabi-Yau threefold, the complexified axio-dilaton, and the brane moduli. In the presence of spacetime filling D3 branes, the latter moduli include the D3 position moduli in the compactification space. In these large volume scenarios, the complex structure and the axio-dilaton are typically stabilized at weak coupling by turning on R-R and NS-NS fluxes that thread 3-cycles in the Calabi-Yau manifold. These fluxes give relatively large masses to the moduli through the flux-induced superpotential, and they introduce a warp factor [33].¹³ At this stage, in the sketched approximation, the Kähler moduli and the D3 brane moduli are still massless. However nonperturbative effects, such as Euclidean D3 instantons or gaugino condensation on 7-branes, induce a nonperturbative superpotential that can stabilize the remaining moduli and yield an $\mathcal{N} = 1$ AdS₄ vacuum. In this scheme, the masses of the D3 brane and Kähler moduli (in string units) are exponentially small in the volume of the internal space, while the masses of the complex structure moduli and the axio-dilaton are relatively larger as they depend on the volume through an inverse power law.

Scenarios of moduli stabilization that make use of the supergravity approximation are consistent in the limit that the volume of the internal space and all of its cycles are large in string units. This is precisely the gravity decoupling limit discussed in earlier sections since the four-dimensional Planck mass is related to the volume of the internal space in string units as

$$M_{\text{pl}}^2 = \frac{\text{Vol}}{g_s^2} M_s^2, \tag{4.1}$$

$\text{Vol} \rightarrow \infty$ thus implies that $M_{\text{pl}}/M_s \rightarrow \infty$.¹⁴ In this limit, the light moduli can be modeled as a supersymmetric sigma model in AdS₄. Another consequence of the large volume limit is that only the leading nonperturbative effects appear in the superpotential. We will argue that for a generic supersymmetric AdS flux vacuum at large volume (in type IIB scenarios), the masses of the light moduli are all proportional to λ . Our conclusions clearly do not apply to the scenarios discussed in [36], since the AdS minima in those constructions are non-supersymmetric. For a detailed analysis of the moduli spectrum in these models, see [37].

The structure of this section is as follows: first, we consider an example with a single Kähler modulus, then we consider an example with additional brane moduli. We point out that the mass matrix of the light modes in these scenarios is proportional to λ^2 , and we end with a general argument that shows the existence of light moduli even when one allows for multiple Kähler moduli.

Example 1. (Kähler modulus): $K = -\log(y + \bar{y})$.

This is the sigma model encountered in the case of a single Kähler modulus, where $\rho \equiv y + \bar{y}$ measures the volume of the internal space. (We set $M_{\text{pl}} = 1$ throughout this section.) The imaginary part $a \equiv \text{Im}(y)$ is an axion that enjoys a continuous shift

¹³Strictly speaking, the process of turning on background fluxes does not just introduce a warp factor, but also requires compactification spaces beyond Calabi-Yau manifolds. In the context of type II compactifications to supersymmetric AdS₄ vacua, such generalizations are discussed in [34, 35].

¹⁴Note that we cannot strictly take this limit since then the internal space decompactifies.

symmetry. The supersymmetric vacua of the theory are determined by solving (2.25), which yields a supersymmetric vacuum at $\rho \rightarrow \infty$. This runaway is typical of the no-scale structure where the scalar potential vanishes as the internal space decompactifies. Of course, y can be naturally stabilized at large volume by including a “small” superpotential term. There is, however, a more general argument for a superpotential term — since this sigma model is part of a consistent quantum gravity theory, the continuous shift symmetry $a \rightarrow a+c$ must be broken by non-perturbative effects. This motivates a superpotential term $W(y) = \mu^3 \exp[-y]$. The negative exponential ensures that the superpotential vanishes, as it should, in the limit of large volume. Solving the supersymmetry conditions (2.25), we find a single supersymmetric vacuum at $\rho = \rho_0$ and $a = \pi + 3 \arg(\mu)$, where

$$\rho_0 \exp(-\rho_0/2) = \frac{\lambda}{|\mu|^3} . \tag{4.2}$$

Since the superpotential scale $\mu \lesssim \Lambda_\sigma \ll M_{\text{pl}} = 1$, this suggests that a supersymmetric vacuum exists at large ρ_0 if and only if the AdS scale λ is exponentially small (in units of M_{pl}). The masses of fluctuations of ρ and a can be easily computed and are both $\approx \rho_0^2 \lambda^2$. Note that the masses are proportional to λ and hence these moduli are light.¹⁵ The superpotential could, in general, include higher order exponentials $\exp(-ny)$ for $n \in \mathbb{Z}^+$. By working in the large volume limit, we can drop these terms and keep only the leading exponential.

As an example, for $\rho_0 = 50$ and $\mu = 1$, we find

$$\frac{\lambda}{M_{\text{pl}}} \approx 7 \times 10^{-10} , \quad m_a^2 = 2350 \lambda^2 , \quad m_\rho^2 = 2448 \lambda^2 . \tag{4.3}$$

Example 2. Mobile D3 brane + Kähler modulus.

This sigma model describes the coupling of the D3 brane position to the overall volume modulus in a IIB compactification. The Kähler potential for such a scenario was first written down in [38] and derived in [39, 40]:

$$\hat{K}(y, \bar{y}, z^i, \bar{z}^{\bar{i}}) = -\log [y + \bar{y} - f^2 k(z^i, \bar{z}^{\bar{i}})] , \tag{4.4}$$

where $k(z^i, \bar{z}^{\bar{i}})$ is the Kähler potential on the brane moduli space, X . In the above formula, we set $M_{\text{pl}} = 1$ and take $f \ll 1$, which should be thought of as setting the KK scale [41]. We denote the space spanned by the coordinates y and z^i by \hat{X} . This space is a \mathbb{C}^* fibration over X [32] with an exact Kähler form derived from (4.4).

We discuss in detail various aspects of the geometry of \hat{X} in appendix A. There we show that such a \mathbb{C}^* fibration over a compact space X is allowed only when it is a Hodge manifold, i.e. when the (normalized) Kähler class is an integral class in $H^2(X, \mathbb{Z}) \cap H^{1,1}(X, \mathbb{C})$. When this is not the case, there are at least two ways in which one can create a total space \hat{X} with an exact Kähler form: first, we could couple the sigma model on X to a single linear multiplet and avoid the quantization condition mentioned above; second, we could add multiple \mathbb{C}^* fibers, up to $h^{1,1}(X)$ of them, to trivialize the Kähler form of the total space of the fibration.

¹⁵Since $\rho_0 \gg 1$, however, the masses of the light moduli are parametrically larger than the AdS scale.

When X has a non-trivial Kähler form, Komargodski and Seiberg used an analysis of supercurrent multiplets in Minkowski space to show that it is impossible to stabilize the Kähler modulus while leaving the brane moduli massless [5]. This followed from the fact that the theory with target space \hat{X} has an exact Kähler form and hence a well-defined Ferrara-Zumino multiplet. Then if y became massive while the z^i remained massless, we could integrate out y and be left with a sigma model on X . Since the Kähler form on X is non-trivial, the resulting supercurrent multiplet would then not be well-defined over the moduli space, but it is not possible for an RG flow to take well-defined operators from the UV to ill-defined operators in the IR. Thus, the inclusion of a nonperturbative superpotential would necessarily have to lift both the Kähler modulus and the brane moduli. This result matches with the original stringy arguments of [32]. We will be led to similar conclusions in AdS₄ sigma models by simply appealing to the conditions necessary for the existence of supersymmetric vacua at large volume.

In analogy with the previous example, the analysis will be carried out in terms of the variables

$$\rho \equiv y + \bar{y} - f^2 k(z^i, \bar{z}^i), \quad a \equiv \text{Im}(y) = \frac{y - \bar{y}}{2i}. \quad (4.5)$$

As explained in [39, 40], the variable ρ measures the overall volume of the space X . In the large volume limit $\rho \rightarrow \infty$, it is consistent to include only the leading exponential in the nonperturbative effective superpotential,

$$W(y, z^i) = p(z^i) \exp(-ny). \quad (4.6)$$

As in the previous example, the superpotential breaks the shift symmetry of a , but since y is not a good coordinate on the total space \hat{X} , there is a z^i dependent prefactor [42]. $p(z^i)$ is a holomorphic section of a line bundle on X with transition functions chosen precisely to cancel those of the \mathbb{C}^* section $\exp(-ny)$. The resulting $W(y, z^i)$ is simply a function on \hat{X} .

The conditions for a supersymmetric vacuum in AdS₄ (2.25) are

$$nW = -\frac{\lambda}{\rho}, \quad \frac{p_i}{p} W = \frac{\lambda f^2 k_i}{\rho}, \quad (4.7)$$

where the subscript i denotes a derivative with respect to z^i . Substituting for W in the second equation using the first, we have

$$np_i(z) + f^2 k_i(z, \bar{z}) p(z) = 0. \quad (4.8)$$

The vacuum expectation values for the z^i are completely determined by this equation, independent of ρ and λ . The first equation in (4.7) then has a solution with large ρ only when λ is exponentially small.

The moduli masses can now be computed from the two-derivative matrix of the scalar potential evaluated at the supersymmetric vacuum,

$$\begin{aligned} V_{ab} &= -\lambda(W_{ab} + \lambda K_{ab}), \\ V_{a\bar{b}} &= g^{c\bar{d}}(W_{ac} + \lambda K_{ac})(\bar{W}_{\bar{b}\bar{d}} + \lambda K_{\bar{b}\bar{d}}) - 2\lambda^2 g_{a\bar{b}}, \\ V_{\bar{a}\bar{b}} &= -\lambda(\bar{W}_{\bar{a}\bar{b}} + \lambda K_{\bar{a}\bar{b}}). \end{aligned} \quad (4.9)$$

The indices a, b, \dots run through the coordinates (y, z^i) . This mass matrix is proportional to λ^2 if W_{ab} is proportional to λ for all a, b . Here we find

$$W_{yy} = n^2 W = -\frac{n\lambda}{\rho}, \quad W_{yi} = -n \frac{p_i}{p} W = \frac{n\lambda f^2 k_i}{\rho}, \quad W_{ij} = \frac{p_{ij}}{p} W = -\frac{\lambda}{n\rho} \frac{p_{ij}}{p}, \quad (4.10)$$

which indeed shows that W_{ab} is proportional to λ . The moduli masses are obtained by canonically normalizing the kinetic terms for the fluctuations and then computing the eigenvalues. The eigenvalues are clearly proportional to λ , but the question is whether the prefactor, which depends on the vevs of y, z^i , can modify this scaling. We noted earlier that the vevs of the z^i are completely independent of λ . The vev of ρ does depend on λ , but only in a logarithmic manner, and so we expect the moduli masses to be exponentially small (in the volume).

General argument. This argument for the appearance of light fields easily generalizes. To this end, let us assume that at the classical/perturbative level the analyzed scenarios have a set of shift symmetries $\text{Im}(y^A) \rightarrow \text{Im}(y^A) + \text{const}$ with respect to the moduli fields y^A , and we denote the remaining moduli in the theory by z^i . In type II compactifications, for instance, the moduli fields y^A could arise from complexified Kähler moduli while the fields z^i may represent complex structure moduli of the internal Calabi-Yau spaces. Nonrenormalization theorems for the superpotential severely constrain the form of the effective superpotential such that no perturbative contributions to the effective superpotential can break these shift symmetries. However, there still may be nonperturbative corrections that appear as exponentials $\exp(-ny^A)$ in the effective superpotential. As a consequence, the leading order terms of the (nonperturbatively generated) effective superpotential take the form

$$W(y, z) = \sum_A C^A(y^A, z) = \sum_A p^A(z) \exp(-y^A). \quad (4.11)$$

Here we have absorbed any numerical factors in the exponential into y^A . To avoid runaways, we require that the p^A , which are general functions of the z^i , do not vanish identically for any A . The Kähler potential $K(y, \bar{y}, z, \bar{z})$ is such that y^A appear in the combination $y^A + \bar{y}^{\bar{A}}$ so that the shift symmetry is only broken by the superpotential W . In addition, we require that the boundaries $y^A \rightarrow \pm\infty$ of the target space are at infinite distance with respect to the Kähler metric, a condition that we discuss further in appendix A.

Before we move on, let us point out a few caveats to the arguments that led us to (4.11): first, for general AdS theories we cannot unambiguously identify a unique superpotential since a Kähler transformation can shift (part of) the superpotential into the Kähler potential and vice versa; however, when our theory has a well-defined flat space limit, $\lambda \rightarrow 0$, we *can* unambiguously define the superpotential as the surviving part in the limit $\lambda \rightarrow 0$. Second, in arguing for the structure of the superpotential we relied upon nonrenormalization theorems for the superpotential, but it is not clear to what extent such theorems are applicable in the context of the global AdS sigma models; again, we are simply guided by the limit $\lambda \rightarrow 0$ and the intuition gained from phenomena in supersymmetric gauge theories [43]. In further defense of the form (4.11), note that it also agrees with the expected

structure of nonperturbative effects that arise in string compactifications [42]. How far such a naive analysis can be pushed in the general AdS₄ setting is an interesting question to which we hope to return elsewhere.

The conditions to have a supersymmetric vacuum (2.25) read

$$C^A = \lambda K_A, \quad \sum_A C_i^A + \lambda K_i = 0. \quad (4.12)$$

The subscripts denote derivatives with respect to the corresponding fields. The second derivatives of the superpotential W evaluated in a supersymmetric vacuum are given by

$$\begin{aligned} W_{AA}|_{\text{susy}} &= C^A|_{\text{susy}} = \lambda K_A|_{\text{susy}}, & W_{AB}|_{\text{susy}} &= 0 \quad \text{for } A \neq B, \\ W_{Ai}|_{\text{susy}} &= -C_i^A|_{\text{susy}} = -\lambda \frac{p_i^A}{p^A} K_A|_{\text{susy}}, & W_{ij}|_{\text{susy}} &= \sum_A C_{ij}^A|_{\text{susy}} = \lambda \sum_A \frac{p_{ij}^A}{p^A} K_A|_{\text{susy}}. \end{aligned} \quad (4.13)$$

The argument is virtually identical as in the previous example. As a result, the moduli masses are proportional to λ with a prefactor that depends only logarithmically on λ .

In summary, we find that for KKL-like scenarios of large-volume moduli stabilization with $\mathcal{N} = 1$ AdS₄ supersymmetry, there are always light moduli with masses proportional to the AdS scale.

5 Gauge theories: regulating the Affleck-Dine-Seiberg runaway in AdS

So far, we have focused on $\mathcal{N} = 1$ sigma models in AdS₄, their consistency conditions, and their implications for certain moduli stabilization scenarios. It would be interesting to extend this analysis to $\mathcal{N} = 1$ gauge theories and their moduli spaces, as well. While a systematic treatment is beyond the scope of this paper,¹⁶ our analysis above suggests some interesting predictions on the moduli spaces of $\mathcal{N} = 1$ gauge theories in AdS₄. For example, placing $N_f < N_c$ SQCD in AdS₄ can regulate the Affleck-Dine-Seiberg runaway [7]. In this section, we present a schematic analysis of this system. We hope to return to a more detailed discussion of gauge theories in AdS₄ in the future.

It was suggested by Callan and Wilczek [26] that AdS₄ serves as an infrared regulator for theories that would otherwise have incurable divergences in flat space. They analyzed the case of the XY model in AdS₂ where the properties of the high temperature phase, in which the vortices are deconfined, can be calculated reliably in a dilute gas approximation. They also proposed that the confining phase of QCD could be studied at weak coupling in AdS₄ since the usual IR divergences associated with nonperturbative computations would be regulated.

In this paper, we have studied the Lagrangian for interacting chiral multiplets in AdS₄. In some cases, an asymptotically free gauge theory at energies below the confinement scale, Λ_c , can be described by an effective Lagrangian consisting of the bound state mesons and baryons (as in the case of QCD). For example, consider the case of four-dimensional $\mathcal{N} = 1$

¹⁶A complete analysis must take into account boundary conditions, which have a significant effect in AdS₄ [20], and a more thorough treatment of possible corrections to the Kähler potential.

$SU(N_c)$ SQCD with N_f quarks in the fundamental representation. The dynamics of the gauge theory depends on the ratio N_f/N_c (see [43]); in the case $N_f < N_c$, an effective superpotential is generated [7]

$$W_{\text{eff}} = (N_c - N_f) \left(\frac{\Lambda_c^{3N_c - N_f}}{\det M} \right)^{\frac{1}{N_c - N_f}} . \tag{5.1}$$

Here M is an $N_f \times N_f$ matrix of meson superfields. As explained in [43], this superpotential does not violate nonrenormalization theorems since it is generated by non-perturbative effects (instantons when $N_f = N_c - 1$ and gaugino condensation in the unbroken $SU(N_c - N_f)$ gauge group when $N_f < N_c - 1$). The classical theory has a moduli space of vacua along which various mesons acquire expectation values, thus Higgsing the gauge group. Quantum mechanically, the entire moduli space is lifted and the theory has no vacuum; instead, it has a runaway $M \rightarrow \infty$.

We can study the behavior of this theory in globally supersymmetric AdS_4 by arranging a hierarchy of scales $\Lambda_c \gg \lambda$, where λ is the inverse AdS_4 radius. The space is effectively Minkowski at the confinement scale, so we expect that the effects of the AdS_4 curvature are negligible at those energies (we will discuss corrections below). For convenience, we specialize to the case $N_f = 1$, $N_c = 2$. Then in terms of the dimensionless meson field $m \equiv M/\Lambda_c^2$, the condition for a supersymmetric vacuum reads

$$\frac{\partial W}{\partial m} + \lambda \frac{\partial K(m, \bar{m})}{\partial m} = 0 . \tag{5.2}$$

In contrast to the supersymmetry conditions in Minkowski space, the existence of a supersymmetric vacuum depends crucially on the Kähler potential in AdS_4 . For example, if we assume that the Kähler potential remains canonical (more on this below), $K = 2\Lambda_c^2|m|$, we find a single supersymmetric vacuum located at

$$\langle m \rangle = \left(\frac{\Lambda_c}{\lambda} \right)^{1/2} \gg 1 . \tag{5.3}$$

The masses of the meson fluctuations around the supersymmetric vacuum, which can be computed from the scalar potential, are proportional to λ . Since $\lambda \ll \Lambda_c$, we can treat the mesonic fields as light propagating degrees of freedom below the confinement scale, so the Affleck-Dine-Seiberg runaway can apparently be regulated by the AdS_4 scale.

Of course, this discussion ignores boundary conditions and quantum corrections to both the Kähler potential and the superpotential. The effects of boundary conditions are quite subtle and beg for further study, which we leave for future work; for now we will simply assume their effects are negligible in the limit $\lambda/\Lambda_c \rightarrow 0$. We have argued that in AdS there is no invariant distinction between the superpotential and the Kähler potential. As a consequence, it is difficult to say whether the vacuum we have found is stable against quantum corrections without a detailed calculation. A correction to the Kähler potential, for example, of the form $\Lambda_c^j \lambda^k |M|^{1-j/2-k/2} = \Lambda_c^2 |m| \times \left[\left(\frac{\lambda}{\Lambda_c} \right)^k |m|^{-j/2-k/2} \right]$ for $j, k \in \mathbb{Z}$ and $j, k \geq 0$ does not destabilize the vacuum. Logarithmic corrections, on the other hand,

are potentially dangerous if the coefficients are sufficiently large. In this note, we do not further examine the structure of such quantum corrections, but we hope that the presented arguments serve as a motivation to study these theories in greater detail.

Our understanding of supersymmetric gauge theories in flat space was greatly advanced by the holomorphy arguments pioneered by Seiberg [44]. In the AdS₄ case, as we have seen, these arguments can become quite subtle because of the mixing between the superpotential and the Kähler potential under Kähler transformations. An extremely important task, then, is to understand how AdS₄ modifies the rich phase structure of nonabelian gauge theories in Minkowski space. We hope to return to this important problem in future work.

Acknowledgments

We thank J. Distler, T. Faulkner, S. Franco, D. Freedman, S. Giddings, Z. Komargodski, J. Louis, D. Marolf, J. Polchinski, A. Van Proeyen, D. Robbins, E. Silverstein, C. Tamarit, J. Thaler, D. Tong, G. Torroba, and S. Trivedi, for useful discussions and correspondences. We would especially like to thank S. Kachru, L. McAllister and N. Seiberg for enlightening discussions, and S. Kachru for helpful comments on the draft. A.A. thanks the 2010 Amsterdam Summer Workshop for an enlightening setting in which this work was initiated, and the Stanford Institute for Theoretical Physics and the Tata Institute of Fundamental Research for hospitality while this work was being completed. V.K. thanks the organizers of the Advanced String School 2010 in Puri, India, where a part of this research was carried out. A.A. is supported by the DOE under contract #DE-FC02-94ER40818. H.J., V.K., and J.M.L., are supported by the Kavli Institute for Theoretical Physics and in part by the NSF Grant PHY05-51164. H.J. is also supported in part by the Stanford Institute for Theoretical Physics and the NSF Grant 0244728. J.M.L. is also supported in part by the NSF Grant No. PHY07-57035.

A Geometry of the enlarged target space \hat{X}

Since the global target space structure of supersymmetric AdS₄ sigma models require us (even without coupling to gravity) to consider target space geometries with trivial Kähler forms, we discuss next in some detail how to obtain an enlarged target space \hat{X} that lacks compact holomorphic cycles (suitable for an AdS₄ sigma model) from a target space geometry X that contains compact holomorphic cycles. In this process, we are naturally led to the same target space enlargement \hat{X} as discussed in [5]. Hence our results for the Kähler target space apply equally well for both scenarios.

Consider an arbitrary Kähler manifold X with a non-trivial Kähler form ω .¹⁷ In a patch $U_\alpha \subset X$, we can define the Kähler potential $k_\alpha(\varphi, \bar{\varphi})$, where the φ are holomorphic coordinates in the patch. On the intersection of two patches, the Kähler potential undergoes a Kähler transformation specified by the holomorphic function $f_{\alpha\beta}(\varphi)$

$$k_\alpha(\varphi, \bar{\varphi}) - k_\beta(\varphi, \bar{\varphi}) = f_{\alpha\beta}(\varphi) + \bar{f}_{\alpha\beta}(\bar{\varphi}) . \tag{A.1}$$

¹⁷In this work we consider smooth target space manifolds. Various aspects of singular target spaces are discussed in [45].

This determines the $f_{\alpha\beta}(\varphi)$ up to imaginary constants

$$f_{\alpha\beta}(\varphi) \sim f_{\alpha\beta}(\varphi) + 2\pi i c_{\alpha\beta} . \tag{A.2}$$

Moreover, due to the relation (A.1) the transformation functions $f_{\alpha\beta}(\varphi)$ must obey

$$f_{\alpha\beta}(\varphi) - f_{\alpha\gamma}(\varphi) + f_{\beta\gamma}(\varphi) + \bar{f}_{\alpha\beta}(\bar{\varphi}) - \bar{f}_{\alpha\gamma}(\bar{\varphi}) + \bar{f}_{\beta\gamma}(\bar{\varphi}) = 0 .$$

Then on triple overlaps, we can define the real constants

$$\omega_{\alpha\beta\gamma} = \frac{1}{2\pi i} (f_{\alpha\beta}(\varphi) - f_{\alpha\gamma}(\varphi) + f_{\beta\gamma}(\varphi)) . \tag{A.3}$$

These real constants $\omega_{\alpha\beta\gamma}$ are defined modulo the real constants $(\delta c)_{\alpha\beta\gamma} = c_{\alpha\beta} - c_{\alpha\gamma} + c_{\beta\gamma}$, according to the ambiguity (A.2), and they obey $\omega_{\alpha\beta\gamma} - \omega_{\alpha\beta\delta} + \omega_{\alpha\gamma\delta} - \omega_{\beta\gamma\delta} = 0$. Thus the constants $\omega_{\alpha\beta\gamma}$ furnish an element in Čech cohomology group $\check{H}^2(X, \mathbb{R})$, which may be identified (by the Čech-de Rham-isomorphism) with the non-exact Kähler $(1, 1)$ -form ω .

We construct a new space \hat{X} as a fibration over X in the following way: for each U_α we add a fiber coordinate $y_\alpha \in \mathbb{C}$. On the overlap of two patches, the local coordinates y_α are related through the transition functions $f_{\alpha\beta}(\varphi)$ as

$$y_\alpha - y_\beta = f_{\alpha\beta}(\varphi) . \tag{A.4}$$

Since X has a non-trivial Kähler form, we have to make sure that these transformations are consistent on triple overlaps:

$$y_\alpha - y_\beta = (y_\alpha - y_\gamma) - (y_\beta - y_\gamma) = f_{\alpha\gamma}(\varphi) - f_{\beta\gamma}(\varphi) = f_{\alpha\beta}(\varphi) - 2\pi i \omega_{\alpha\beta\gamma} . \tag{A.5}$$

Then we see that y_α is a good fiber coordinate on the triple overlap only if we identify $y_\alpha \sim y_\alpha + 2\pi i \omega_{\alpha\beta\gamma}$. For the chiral field y to be a well-defined periodic field, and hence to ensure a geometric interpretation in terms of a fibration, we require that the constants $\omega_{\alpha\beta\gamma}$ are all mutually commensurate — i.e., there exists a constant $\kappa \in \mathbb{R}$ for which all the $\kappa \omega_{\alpha\beta\gamma}$ are integers.¹⁸ Then we get a periodic chiral field y with periodicity

$$y \sim y + 2\pi i \kappa . \tag{A.6}$$

The requirement that we obtain a well-defined periodic chiral field y transforming as in (A.4) imposes a geometric condition on the Kähler target space geometry X . Namely, the Kähler form ω needs to be quantized with respect to some positive real constant κ ,

$$\kappa \omega \in H^{1,1}(X) \cap H^2(X, \mathbb{Z}) . \tag{A.7}$$

Such Kähler manifolds with integral Kähler forms are called Kähler manifolds of restricted type or Hodge manifolds (see, e.g., [47]). (Again, the Hodge condition follows from an

¹⁸When this condition is not satisfied, there is no geometric interpretation of this space since the fiber is periodically identified on a dense set. From the physics point of view, at least in Minkowski space, there is a sensible dual interpretation in terms of a linear multiplet coupled to a sigma model with target space X . Only when a Dirac quantization condition — i.e., when the identifications are integrally related — is obeyed are we allowed to dualize linear and chiral multiplets [46].

attempt to trivialize the Kähler class of X by fibering a single line over it. The more generic situation is discussed in a paragraph below.)

From (A.1) and (A.4), we can define a *global* real coordinate $\rho_\alpha \equiv y_\alpha + \bar{y}_\alpha - k_\alpha(\varphi, \bar{\varphi})$ since the transformations on the overlaps precisely cancel (we henceforth drop the α label on ρ). Using this coordinate, we can construct a globally well-defined Kähler potential on the enlarged space \hat{X} consisting of the coordinates φ^i, y ,

$$\hat{K}(\varphi, \bar{\varphi}, y, \bar{y}) = -H(\rho) = -H(y + \bar{y} - k(\varphi, \bar{\varphi})) . \tag{A.8}$$

Note that this ansatz for the Kähler potential respects the continuous shift symmetry $y \rightarrow y + 2\pi i c$. This indicates that there is a dual formulation in terms of a linear multiplet by dualizing the chiral multiplet y [46]. We obtain a positive definite Kähler metric \hat{G} , represented by the line element

$$ds_{\hat{X}}^2 = -H''(\rho)(dy - k_i(\varphi, \bar{\varphi})d\varphi^i)(d\bar{y} - k_{\bar{j}}(\varphi, \bar{\varphi})d\bar{\varphi}^{\bar{j}}) + H'(\rho)g_{i\bar{j}}(\varphi, \bar{\varphi})d\varphi^i d\bar{\varphi}^{\bar{j}} , \tag{A.9}$$

as long as H (in a domain of ρ) is a smooth real function obeying

$$H'(\rho) > 0, \quad H''(\rho) < 0 . \tag{A.10}$$

There are a few remarks in order before we further analyze the geometry. For a compact Kähler target space X with an integral basis ω_A of non-trivial $(1,1)$ -forms, the Hodge condition (A.7) imposes that the (rescaled) Kähler form $\kappa\omega$ is an integral linear combination of the integral basis ω_A . When $H^{1,1}(X)$ is generated by a single non-trivial $(1,1)$ -form, we can always rescale the Kähler form ω with an appropriate constant κ to achieve integrality. When $\dim(H^{1,1}(X)) > 1$, the Hodge condition is not met for Kähler forms arising as linear combinations with irrational coefficients relative to each other. In such a situation, we can trivialize the Kähler form by introducing additional periodic chiral fields. For instance, we could, following [48], introduce a periodic chiral field for each $(1,1)$ -form ω_A to compensate its contribution to the Kähler form ω . Then the failure of the Hodge condition would be reflected in the fact that at least two of the introduced periodic chiral fields would have periodicities that are irrational relative to each other. Of course, this is exactly what one must do at a generic point in the Kähler moduli space of X .

In order to now exhibit the geometric structure of the constructed Kähler target space \hat{X} with the Kähler potential (A.8), we exponentiate the chiral field ϕ and introduce the reparametrized chiral field U

$$U = \exp\left(\frac{y}{\kappa}\right) . \tag{A.11}$$

Due to (A.6), the chiral field U yields a local single valued \mathbb{C}^* coordinate in each local patch. Therefore, the constructed target space \hat{X} is a \mathbb{C}^* -fiber bundle over the original target space X ,

$$\mathbb{C}^* \longrightarrow \hat{X} \xrightarrow{\pi} X . \tag{A.12}$$

Note that under Kähler transformations $\phi \rightarrow \phi + f(A)$ of the base, the \mathbb{C}^* -fiber coordinate U transforms as

$$U \rightarrow g(\varphi)U, \quad g(\varphi) = \exp\left(\frac{f(\varphi)}{\kappa}\right) . \tag{A.13}$$

Here the holomorphic transition functions $g(\varphi)$ are locally non-vanishing. We can think of the \mathbb{C}^* -bundle as arising from a complex line bundle with its zero section removed. Therefore, analogously to line bundles, we characterize the \mathbb{C}^* -bundle by its first Chern class

$$c_1(\mathbb{C}^*) = [\kappa\omega] \in H^{1,1}(X) \cap H^2(X, \mathbb{Z}), \tag{A.14}$$

which by construction is equal to the integral cohomology representative of the rescaled Kähler $(1, 1)$ -form $\kappa\omega$.¹⁹

Let us pause to analyze how the \mathbb{C}^* -fiber bundle structure (A.12) of the Kähler target space \hat{X} manages to trivialize its Kähler form $\hat{\omega}$. First of all, a straightforward calculation shows that a globally well-defined trivializing one form $\hat{\theta}$ exists, with $\hat{\omega} = d\hat{\theta}$:

$$\begin{aligned} \hat{\theta} &= \frac{i}{2} H'(\rho) \left(\kappa \frac{dU}{U} - \kappa \frac{d\bar{U}}{\bar{U}} - (K_i(\varphi, \bar{\varphi}) d\varphi^i - K_{\bar{j}}(\varphi, \bar{\varphi}) d\bar{\varphi}^{\bar{j}}) \right) \\ &= \frac{i}{2} H'(\rho) (d\phi - d\bar{\phi} - (K_i(\varphi, \bar{\varphi}) d\varphi^i - K_{\bar{j}}(\varphi, \bar{\varphi}) d\bar{\varphi}^{\bar{j}})) . \end{aligned} \tag{A.15}$$

The relationship between the cohomology classes of the base space X and the whole space \hat{X} sheds more light on the exactness of the Kähler form $\hat{\omega}$. We can compute the cohomology classes of the \mathbb{C}^* -fibration \hat{X} with the Leray spectral sequence [49]. In particular, we find that the two-form class associated to the first Chern class of the \mathbb{C}^* -bundle is removed from the two-form cohomology of \hat{X} since it becomes exact,²⁰ i.e.,

$$H^2(\hat{X}) \simeq H^2(X)/c_1(\mathbb{C}^*) . \tag{A.16}$$

This is in agreement with the observation that in the construction of the global one-form $\hat{\theta}$ in (A.15), the \mathbb{C}^* -fiber directions play an essential role.

We can also check that the Kähler manifold \hat{X} satisfies the general geometric criteria (collected at the end of section 2.1) for a Kähler manifold to admit an exact Kähler form. Namely, the Kähler manifold is non-compact since the \mathbb{C}^* -fibers are non-compact. Furthermore, since the \mathbb{C}^* -fibers are non-compact we can only try to construct compact holomorphic subspaces (of dimension greater than zero) in the base. However, any compact holomorphic subspace of the base X ceases to be compact in the total space \hat{X} . Owing to the fact that we require a positive-definite Kähler metric, the \mathbb{C}^* -fibration restricts to a non-trivial \mathbb{C}^* -fibration over any compact holomorphic subspace. Moreover, a non-trivial \mathbb{C}^* -fibration does not admit any global holomorphic sections. Therefore, the considered compact subspace of the base X cannot be holomorphically embedded in the \mathbb{C}^* -fibered target space \hat{X} .

¹⁹The transition functions $g_{\alpha\beta}(\varphi)$ give rise to the first Chern class via the map $H^1(X, \mathcal{O}^*) \rightarrow H^2(X, \mathbb{Z})$ arising from the long exact sequence of the exponential exact sequence of sheaves $0 \rightarrow 2\pi i \mathbb{Z} \hookrightarrow \mathcal{O} \xrightarrow{\exp} \mathcal{O}^* \rightarrow 0$ [47]. Under this map, one finds the κ -rescaled Čech representative $\kappa\omega_{\alpha\beta\gamma}$ of (A.3) for the first Chern class.

²⁰The E_2 term in the Leray spectral sequence reads $E_2^{p,q} = H^p(X) \otimes H^q(\mathbb{C}^*) \simeq \begin{cases} \mathbb{R}^{b_p} & q = 0, 1 \\ 0 & \text{else} \end{cases}$ in terms of the Betti numbers $b_p = \dim H^p(X)$. The spectral sequence applied to \mathbb{C}^* -bundles degenerates at $E_3^{p,q} = H^{p+q}(\hat{X})$, and the only non-trivial differential is $d_2 : E_2^{p-2,1} \rightarrow E_2^{p,0}, \vartheta \mapsto (-1)^{p+1} \pi^* c_1(\mathbb{C}^*) \wedge \pi^* \vartheta$; together, these imply that the two-form $\pi^* c_1(\mathbb{C}^*) = -d_2(0\text{-form on } \mathbb{C}^*)$ is exact in the space \hat{X} .

We end this section with a discussion on the metric (A.9) of the target space \hat{X} and, in particular, on the structure of the real and sufficiently smooth function $H(\rho)$. As observed in (A.10), $H(\rho)$ must be a monotonically increasing and negatively curved function in order to yield a non-degenerate Kähler metric. The latter conditions on $H(\rho)$ are necessary to arrive at a sigma model with a well-defined, non-tachyonic kinetic term. Furthermore, because of the non-compactness of the target space \hat{X} (due to the non-compact \mathbb{C}^* -fibers), we may choose the function $H(\rho)$ such that the target space boundary is at infinite distance.

To arrive at the conditions for infinitely far boundaries, we first note that the domain of function $H(\rho)$ is a real interval (ρ_-, ρ_+) . The boundary of this domain corresponds to the target space boundary in the fiber direction.²¹ To determine the distance to the boundary, it is convenient to rewrite the metric (A.9) in terms of the variables ρ and $a = \text{Im}(y)$,

$$ds_{\hat{X}}^2 = -\frac{H''(\rho)}{4}d\rho^2 - H''(\rho)(da - \text{Im}(k_i d\varphi^i))^2 + H'(\rho)g_{i\bar{j}}d\varphi^i d\bar{\varphi}^{\bar{j}}. \quad (\text{A.17})$$

Then the conditions for the boundary to be at infinity read

$$\int_{\rho_-}^{\rho_0} ds_{\hat{X}} = -\frac{1}{4} \int_{\rho_-}^{\rho_0} d\rho \sqrt{H''(\rho)} = \infty, \quad \int_{\rho_0}^{\rho_+} ds_{\hat{X}} = -\frac{1}{4} \int_{\rho_0}^{\rho_+} d\rho \sqrt{H''(\rho)} = \infty, \quad (\text{A.18})$$

where ρ_0 is an arbitrary point in the interval (ρ_-, ρ_+) . These constraints, together with (A.10), are met by the function $H(\rho) = \log \rho$, for example, with the domain $(0, +\infty)$. This choice actually appears in the context of large volume compactifications in various string scenarios, as we discussed in section 4.

References

- [1] M.R. Douglas and S. Kachru, *Flux compactification*, *Rev. Mod. Phys.* **79** (2007) 733 [[hep-th/0610102](#)] [[INSPIRE](#)].
- [2] R. Blumenhagen, B. Körs, D. Lüst and S. Stieberger, *Four-dimensional string compactifications with D-branes, orientifolds and fluxes*, *Phys. Rept.* **445** (2007) 1 [[hep-th/0610327](#)] [[INSPIRE](#)].
- [3] F. Denef, *Les Houches lectures on constructing string vacua*, [arXiv:0803.1194](#) [[INSPIRE](#)].
- [4] S. Kachru, R. Kallosh, A.D. Linde and S.P. Trivedi, *de Sitter vacua in string theory*, *Phys. Rev. D* **68** (2003) 046005 [[hep-th/0301240](#)] [[INSPIRE](#)].
- [5] Z. Komargodski and N. Seiberg, *Comments on supercurrent multiplets, supersymmetric field theories and supergravity*, *JHEP* **07** (2010) 017 [[arXiv:1002.2228](#)] [[INSPIRE](#)].
- [6] A. Adams, N. Arkani-Hamed, S. Dubovsky, A. Nicolis and R. Rattazzi, *Causality, analyticity and an IR obstruction to UV completion*, *JHEP* **10** (2006) 014 [[hep-th/0602178](#)] [[INSPIRE](#)].
- [7] I. Affleck, M. Dine and N. Seiberg, *Dynamical supersymmetry breaking in supersymmetric QCD*, *Nucl. Phys. B* **241** (1984) 493.
- [8] O. Aharony, M. Berkooz, D. Tong, and S. Yankielowicz, work in progress.

²¹Note that ρ_- and ρ_+ could also be $-\infty$ or $+\infty$, respectively. Actually, in order to obtain a smooth function with $H'(\rho) > 0$ and $H''(\rho) < 0$ with a boundary at infinite distance, it turns out that one boundary of the domain of $H(\rho)$ must be infinite.

- [9] E. Witten and J. Bagger, *Quantization of Newton's constant in certain supergravity theories*, *Phys. Lett. B* **115** (1982) 202.
- [10] B. Keck, *An alternative class of supersymmetries*, *J. Phys. A* **8** (1975) 1819.
- [11] B. Zumino, *Nonlinear realization of supersymmetry in Anti de Sitter space*, *Nucl. Phys. B* **127** (1977) 189.
- [12] E. Ivanov and A.S. Sorin, *Wess-Zumino model as linear σ -model of spontaneously broken conformal and $OSp(1,4)$ supersymmetries*, *Sov. J. Nucl. Phys.* **30** (1979) 440 [INSPIRE].
- [13] E. Ivanov and A.S. Sorin, *Superfield formulation of $Osp(1,4)$ supersymmetry*, *J. Phys. A* **13** (1980) 1159 [INSPIRE].
- [14] P. Breitenlohner and D.Z. Freedman, *Positive energy in anti-de Sitter backgrounds and gauged extended supergravity*, *Phys. Lett. B* **115** (1982) 197 [INSPIRE].
- [15] C.J. Burges, D.Z. Freedman, S. Davis and G. Gibbons, *Supersymmetry in anti-de Sitter space*, *Annals Phys.* **167** (1986) 285 [INSPIRE].
- [16] B. de Wit and I. Herger, *Anti-de Sitter supersymmetry*, *Lect. Notes Phys.* **541** (2000) 79 [hep-th/9908005] [INSPIRE].
- [17] D. McKeon and C. Schubert, *Supersymmetry on AdS_3 and AdS_4* , *Class. Quant. Grav.* **21** (2004) 3337 [hep-th/0301225] [INSPIRE].
- [18] B. Gripaios, H.D. Kim, R. Rattazzi, M. Redi and C. Scrucca, *Gaugino mass in AdS space*, *JHEP* **02** (2009) 043 [arXiv:0811.4504] [INSPIRE].
- [19] R. Rattazzi and M. Redi, *Gauge boson mass generation in AdS_4* , *JHEP* **12** (2009) 025 [arXiv:0908.4150] [INSPIRE].
- [20] O. Aharony, D. Marolf and M. Rangamani, *Conformal field theories in Anti-de Sitter space*, *JHEP* **02** (2011) 041 [arXiv:1011.6144] [INSPIRE].
- [21] J. Donoghue, E. Golowich, and B. R. Holstein, *Dynamics of the standard model*, *Camb. Monogr. Part. Phys. Nucl. Phys. Cosmol.* **2** (1992) 1.
- [22] J. Wess and J. Bagger, *Supersymmetry and supergravity*, Princeton University Press, Princeton U.S.A. (1992).
- [23] R. Harvey and H.B. Lawson, *Calibrated geometries*, *Acta Mathematica* **148** (1982) 47.
- [24] S. Weinberg, *The quantum theory of fields. Vol. 3: supersymmetry*, Cambridge University Press, Cambridge U.K. (2000).
- [25] A.E. Nelson and N. Seiberg, *R symmetry breaking versus supersymmetry breaking*, *Nucl. Phys. B* **416** (1994) 46 [hep-ph/9309299] [INSPIRE].
- [26] C.G. Callan Jr. and F. Wilczek, *Infrared behavior at negative curvature*, *Nucl. Phys. B* **340** (1990) 366 [INSPIRE].
- [27] A.J. Amsel and D. Marolf, *Supersymmetric multi-trace boundary conditions in AdS*, *Class. Quant. Grav.* **26** (2009) 025010 [arXiv:0808.2184] [INSPIRE].
- [28] D. Butter and S.M. Kuzenko, *$\mathcal{N} = 2$ AdS supergravity and supercurrents*, *JHEP* **07** (2011) 081 [arXiv:1104.2153] [INSPIRE].
- [29] S. Ferrara and B. Zumino, *Transformation properties of the supercurrent*, *Nucl. Phys. B* **87** (1975) 207 [INSPIRE].

- [30] G.W. Moore and P.C. Nelson, *The etiology of sigma model anomalies*, *Commun. Math. Phys.* **100** (1985) 83.
- [31] J. Distler and E. Sharpe, *Quantization of Fayet-Iliopoulos parameters in supergravity*, *Phys. Rev. D* **83** (2011) 085010 [[arXiv:1008.0419](#)] [[INSPIRE](#)].
- [32] S. Kachru, R. Kallosh, A.D. Linde, J.M. Maldacena, L.P. McAllister, et al., *Towards inflation in string theory*, *JCAP* **10** (2003) 013 [[hep-th/0308055](#)] [[INSPIRE](#)].
- [33] S.B. Giddings, S. Kachru and J. Polchinski, *Hierarchies from fluxes in string compactifications*, *Phys. Rev. D* **66** (2002) 106006 [[hep-th/0105097](#)] [[INSPIRE](#)].
- [34] D. Lüüst and D. Tsimplis, *Classes of AdS₄ type IIA / IIB compactifications with SU(3) × SU(3) structure*, *JHEP* **04** (2009) 111 [[arXiv:0901.4474](#)] [[INSPIRE](#)].
- [35] D. Lüüst and D. Tsimplis, *New supersymmetric AdS₄ type-II vacua*, *JHEP* **09** (2009) 098 [[arXiv:0906.2561](#)] [[INSPIRE](#)].
- [36] V. Balasubramanian, P. Berglund, J.P. Conlon and F. Quevedo, *Systematics of moduli stabilisation in Calabi-Yau flux compactifications*, *JHEP* **03** (2005) 007 [[hep-th/0502058](#)] [[INSPIRE](#)].
- [37] J.P. Conlon, F. Quevedo and K. Suruliz, *Large-volume flux compactifications: moduli spectrum and D3/D7 soft supersymmetry breaking*, *JHEP* **08** (2005) 007 [[hep-th/0505076](#)] [[INSPIRE](#)].
- [38] O. DeWolfe and S.B. Giddings, *Scales and hierarchies in warped compactifications and brane worlds*, *Phys. Rev. D* **67** (2003) 066008 [[hep-th/0208123](#)] [[INSPIRE](#)].
- [39] P.G. Camara, L. Ibáñez and A. Uranga, *Flux induced SUSY breaking soft terms*, *Nucl. Phys. B* **689** (2004) 195 [[hep-th/0311241](#)] [[INSPIRE](#)].
- [40] M. Graña, T.W. Grimm, H. Jockers and J. Louis, *Soft supersymmetry breaking in Calabi-Yau orientifolds with D-branes and fluxes*, *Nucl. Phys. B* **690** (2004) 21 [[hep-th/0312232](#)] [[INSPIRE](#)].
- [41] D. Baumann, A. Dymarsky, I.R. Klebanov, J.M. Maldacena, L.P. McAllister, et al., *On D3-brane potentials in compactifications with fluxes and wrapped D-branes*, *JHEP* **11** (2006) 031 [[hep-th/0607050](#)] [[INSPIRE](#)].
- [42] E. Witten, *Nonperturbative superpotentials in string theory*, *Nucl. Phys. B* **474** (1996) 343 [[hep-th/9604030](#)] [[INSPIRE](#)].
- [43] K.A. Intriligator and N. Seiberg, *Lectures on supersymmetric gauge theories and electric-magnetic duality*, *Nucl. Phys. Proc. Suppl.* **45BC** (1996) 1 [[hep-th/9509066](#)] [[INSPIRE](#)].
- [44] N. Seiberg, *Naturalness versus supersymmetric nonrenormalization theorems*, *Phys. Lett. B* **318** (1993) 469 [[hep-ph/9309335](#)] [[INSPIRE](#)].
- [45] S. Hellerman and E. Sharpe, *Sums over topological sectors and quantization of Fayet-Iliopoulos parameters*, [arXiv:1012.5999](#) [[INSPIRE](#)].
- [46] P. Adamietz, P. Binetruy, G. Girardi and R. Grimm, *Supergravity and matter: linear multiplet couplings and Kähler anomaly cancellation*, *Nucl. Phys. B* **401** (1993) 257 [[INSPIRE](#)].
- [47] P. Griffiths and J. Harris, *Principles of algebraic geometry*, John Wiley & Sons, New York U.S.A. (1994).

- [48] T. Banks and N. Seiberg, *Symmetries and strings in field theory and gravity*, *Phys. Rev. D* **83** (2011) 084019 [[arXiv:1011.5120](#)] [[INSPIRE](#)].
- [49] R. Bott and L.W. Tu, *Differential forms in algebraic topology*, *Graduate Texts in Mathematics*, Springer, Berlin Germany (1982).