

# The Inferior Colliculus



Jeffery A. Winer • Christoph E. Schreiner  
Editors

 Springer

# The Inferior Colliculus

Jeffery A. Winer Christoph E. Schreiner  
Editors

# The Inferior Colliculus

With 168 Illustrations

 Springer

Jeffery A. Winer  
Room 289, Life Sciences Addition  
Division of Neurobiology  
Department of Molecular and  
Cell Biology  
University of California at Berkeley  
Berkeley, CA 94720-3200, USA

Christoph E. Schreiner  
Department of Otolaryngology  
University of California at San Francisco  
Coleman Memorial Laboratory and  
W.M. Keck Center for  
Integrative Neuroscience  
513 Parnassus Avenue HSE-824  
San Francisco, CA 94143-0732, USA

*Cover illustration:* A transverse section through the center of the cat inferior colliculus showing its principal subdivisions. See page 20 of the text.

Library of Congress Cataloging-in-Publication Data

The inferior colliculus / Jeffery A. Winer, Christoph E. Schreiner, editors.  
p. cm.

Includes biographical references and index.

ISBN 0-387-22038-0 (alk. paper)

1. Inferior colliculus. I. Winer, Jeffery A., 1945– II. Schreiner, Christoph.

QM455.I53 2004

612.8'264—dc22

2004049192

ISBN 0-387-22038-0

Printed on acid-free paper.

© 2005 Springer Science+Business Media, Inc.

All rights reserved. This work may not be translated or copied in whole or in part without the written permission of the publisher (Springer Science+Business Media, Inc., 233 Spring Street, New York, NY 10013, USA), except for brief excerpts in connection with reviews or scholarly analysis. Use in connection with any form of information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed is forbidden.

The use in this publication of trade names, trademarks, service marks, and similar terms, even if they are not identified as such, is not to be taken as an expression of opinion as to whether or not they are subject to proprietary rights.


Printed in the United States of America. (WBG/MVY)

9 8 7 6 5 4 3 2 1

SPIN 10941461

springeronline.com





Photograph by courtesy of T. Stewart

Dedicated to the memory of  
Jerzy E. Rose (1909–1992)  
Scholar, scientist, colleague, mentor, friend

# Preface

The purpose of this volume is twofold. First, it offers an in-depth analysis of current approaches and issues in the study of the auditory system. By concentrating on one structure, the inferior colliculus, a focused and coherent treatment of many aspects of auditory neural processing is possible. The position of the inferior colliculus is unique, as its study offers insights into the influence of the peripheral auditory system and at the same time reveals the initial stages of central processing principles. By providing, in the first chapter, an overview of auditory system function and structure, a framework is given that guides the interpretation of operational mechanisms and rules. Second, the book provides a state-of-the-art reference tool for researchers working on the inferior colliculus. The last such treatment appeared in the mid-1980s (Aitkin 1986) and was 246 pages long with 507 citations; since then, more than 1900 articles on the inferior colliculus have been published, and there has been no inclusive summary of facts and ideas about this critical junction in the auditory pathway. In this period, there has been substantial progress on the many facets of inferior colliculus function that constitute the subject matter for this volume. The mere accretion of publications alone would not in itself justify a new volume devoted to the auditory midbrain. The rationale, then, is to summarize recent advances in this discipline from the perspective of some of the many researchers who have engendered this progress. As a case in point, consider the growing body of data on the role of the inferior colliculus in seizure genesis and as a model system for the study of epilepsy (Chapter 21), an area that has grown considerably since 1986 and that has had a significant impact on diverse areas including sensory-to-motor transformations and the possible role of GABAergic neurons in kindling and seizure control. Any consideration of GABAergic neurons, of course, must include their role in local processing as putative interneurons (Oliver et al. 1991) as well as their ascending projection to the medial geniculate body (Winer et al. 1996), the differential, subdivision-specific concentration of GABAergic neurons and axon terminals (Oliver et al. 1994), and the maturation of GABAergic transmission (Yigit et al. 2003); each of these topics is of moment, each crosses interdisciplinary boundaries that can range from development to pathology, and none could have received the appropriate attention in prior synthe-

ses. Because we could not in conscience exclude a particular subject, we attempted to include all that seemed to us to capture best the sense of flux and excitement of current approaches. There remain, of course, many gaps: for example, the subject of synaptic organization has received less attention than might have been expected, and it remains an area that will require further scrutiny if we are to understand how signals arising from the many medullary auditory centers and converging onto the inferior colliculus are transformed locally before they ascend to the auditory thalamus or descend within the brain stem. Likewise, developmental studies are at their earliest stage other than the purely descriptive, and we have little knowledge of how closely the cellular ontogenetic molecules and migratory processes that shape the midbrain follow principles established in the cerebral cortex (Molnár and Blakemore 1995).

Other conceptual approaches that have not been included explicitly are those that can be subsumed under the umbrella of computational neuroscience. There are a few modeling approaches to aspects of temporal coding or binaural processing that were designed to reflect properties specific to the inferior colliculus (e.g., Hewitt and Meddis 1994; Cai et al. 1998; Shackleton et al. 2000; Borisyuk et al. 2002). But it is difficult, and perhaps premature, to assemble a coherent theoretically oriented treatment of inferior colliculus properties, mechanisms, and function.

Where it was possible, we asked that authors propose an agenda for the future in which the salient questions for their discipline are enumerated as an organic part of their exposition. To keep the reference list within manageable limits, we requested that authors cite only the most recent work when this was possible; this strategy acknowledges the historical and intellectual value of Aitkin's (1986) volume.

The conceptual framework for this volume is integrative and reflects a systems perspective. In this context, integrative implies that we sought authors who would collaborate with peers who often held a different perspective, thus producing what we hope are balanced accounts of a given area that are free of parochialism. Where there is only one author, it was our view that the consensus of opinion (or the limited knowledge) in that particular area could be captured by the author we chose. Likewise, chapter length was guided by the literature available and whether the issues at hand were perceived as volatile or matters of settled opinion. The systems viewpoint construes the brain in terms of interacting neural networks whose separate elements contribute to the abstraction of larger entities related to hearing, perception, and binding the disparate streams from independent neural channels into a coherent experience. Perhaps the next volume devoted to the auditory midbrain can realize that goal.

Berkeley, California  
San Francisco, California

JEFFERY A. WINER  
CHRISTOPH E. SCHREINER

## REFERENCES

- Aitkin L (1986) *The Auditory Midbrain. Structure and Function in the Central Auditory Pathway*. Humana Press, Clifton.
- Borisjuk A, Semple MN, and Rinzel J (2002) Adaptation and inhibition underlie responses to time-varying interaural phase cues in a model of inferior colliculus neurons. *Journal of Neurophysiology* **88**:2134–2146.
- Cai H, Carney LH, and Colburn HS (1998) A model for binaural response properties of inferior colliculus neurons. I. A model with interaural time difference-sensitive excitatory and inhibitory inputs. *Journal of the Acoustical Society of America* **103**:475–493.
- Hewitt MJ and Meddis R (1994) A computer model of amplitude-modulation sensitivity of single units in the inferior colliculus. *Journal of the Acoustical Society of America* **95**:2145–2159.
- Molnár Z and Blakemore C (1995) How do thalamic axons find their way to the cortex? *Cerebral Cortex* **18**:389–397.
- Oliver DL, Kuwada S, Yin TCT, Haberly LB, and Henkel CK (1991) Dendritic and axonal morphology of HRP-injected neurons in the inferior colliculus of the cat. *Journal of Comparative Neurology* **303**:75–100.
- Oliver DL, Winer JA, Beckius GE, and Saint Marie RL (1994) Morphology of GABAergic cells and axon terminals in the cat inferior colliculus. *Journal of Comparative Neurology* **340**:27–42.
- Shackleton TM, McAlpine D, and Palmer AR (2000) Modelling convergent input onto interaural-delay-sensitive inferior colliculus neurones. *Hearing Research* **149**:199–215.
- Winer JA, Saint Marie RL, Larue DT, and Oliver DL (1996) GABAergic feedforward projections from the inferior colliculus to the medial geniculate body. *Proceedings of the National Academy of Sciences of the United States of America* **93**:8005–8010.
- Yigit M, Keipert C, and Backus KH (2003) Muscarinic acetylcholine receptors potentiate the GABAergic transmission in the developing rat inferior colliculus. *Neuropharmacology* **45**:504–513.

# Acknowledgments

Many friends, collaborators, and students generously contributed to this book. In lieu of a list that might include some and omit others just as worthy of mention, we simply thank all whose labors made this volume possible. The chapter authors, in particular, deserve high praise for their responsiveness and for their willingness to work together in the best collegial fashion.

On a more individual basis, we are grateful to our editor Janet Slobodien of Springer for her many helpful efforts to see a complex venture to fruition. Dr. Arthur N. Popper was instrumental in encouraging and shaping our early efforts to frame our objectives. We thank David T. Larue for his help with the many figures that constitute the heart of this volume. Dr. Hanna M. Sobkowicz kindly gave guidance in selecting the photograph of Dr. Jerzy E. Rose, and Dr. Tom C.T. Yin arranged the permission for publication and obtained the photographs. Ms. Xuan Yen Thi Phan helped to prepare the Author Index. Ms. Deborah Masi provided meticulous and conscientious care throughout the production process.

On a personal note, we acknowledge the aid, support, and encouragement of our families. Eileen G. Winer was unfailingly generous and gave much moral guidance and love. Marcia Raggio and Christina Schreiner inspired, as always, through their spirit, wisdom, and love.

Berkeley, California  
San Francisco, California

JEFFERY A. WINER  
CHRISTOPH E. SCHREINER

# Contents

Preface . . . . .	vii
Acknowledgments . . . . .	xi
Contributors . . . . .	xvii
Chapter 1 The Central Auditory System: A Functional Analysis. . . . .	1
JEFFERY A. WINER AND CHRISTOPH E. SCHREINER	
Chapter 2 Neuronal Organization in the Inferior Colliculus. . . . .	69
DOUGLAS L. OLIVER	
Chapter 3 Projections from the Cochlear Nuclear Complex to the Inferior Colliculus. . . . .	115
NELL BEATTY CANT	
Chapter 4 Superior Olivary Complex and Lateral Lemniscal Connections of the Auditory Midbrain. . . . .	132
BRETT R. SCHOFIELD	
Chapter 5 Intrinsic and Commissural Connections of the Inferior Colliculus. . . . .	155
ENRIQUE SALDAÑA AND MIGUEL A. MERCHÁN	
Chapter 6 Descending Connections of the Auditory Midbrain. . . . .	182
ANN M. THOMPSON	
Chapter 7 The Tectothalamic System. . . . .	200
JEFFREY J. WENSTRUP	
Chapter 8 Three Systems of Descending Projections to the Inferior Colliculus. . . . .	231
JEFFERY A. WINER	

Chapter 9	Pharmacology of the Inferior Colliculus . . . . .	248
	JACK B. KELLY AND DONALD M. CASPARY	
Chapter 10	Biophysical Properties of Inferior Colliculus Neurons. . . . .	282
	SHU HUI WU	
Chapter 11	Spectral and Intensity Coding in the Auditory Midbrain . . . . .	312
	GÜNTER EHRET AND CHRISTOPH E. SCHREINER	
Chapter 12	Temporal Coding in the Auditory Midbrain . . . . .	346
	ADRIAN REES AND GERALD LANGNER	
Chapter 13	Binaural and Spatial Coding in the Inferior Colliculus . . . . .	377
	ALAN R. PALMER AND SHIGEYUKI KUWADA	
Chapter 14	Coding of Communication Sounds in the Inferior Colliculus . . . . .	411
	CHRISTINE V. PORTFORS AND DONAL G. SINEX	
Chapter 15	Acoustic Behavior and Midbrain Function . . . . .	426
	TOM C.T. YIN AND BRADFORD J. MAY	
Chapter 16	Auditory Midbrain of Fish, Amphibians, and Reptiles: Model Systems for Understanding Auditory Function. . . . .	459
	ANDREW H. BASS, GARY J. ROSE, AND MICHAEL B. PRITZ	
Chapter 17	The Auditory Midbrain in Bats and Birds . . . . .	493
	ELLEN COVEY AND CATHERINE E. CARR	
Chapter 18	Development of Auditory Afferents to the Central Nucleus of the Inferior Colliculus . . . . .	537
	JUDY K. BRUNSO-BECHTOLD AND CRAIG K. HENKEL	
Chapter 19	Inferior Colliculus: Aging and Plasticity. . . . .	559
	ROBERT D. FRISINA AND RAMESH RAJAN	
Chapter 20	Hearing Loss and the Inferior Colliculus . . . . .	585
	JAMES F. WILLOTT	
Chapter 21	The Midbrain and Audiogenic Seizures . . . . .	603
	CARL L. FAINGOLD	

Chapter 22 The Inferior Colliculus: Past, Present, and Future . . . . . 626  
JOHN H. CASSEDAY, CHRISTOPH E. SCHREINER, AND  
JEFFERY A. WINER

Subject Index . . . . . 641

Author Index . . . . . 675



# Contributors

ANDREW H. BASS

Section on Neurobiology and Behavior, Cornell University, Ithaca, NY 14853,  
USA

JUDY K. BRUNSO-BECHTOLD

Department of Neurobiology and Anatomy, Wake Forest University School of  
Medicine, Bowman Gray School of Medicine, Winston-Salem, NC 27157-1010,  
USA

NELL BEATTY CANT

Department of Neurobiology, Duke University School of Medicine, Durham,  
NC 27710, USA

CATHERINE E. CARR

Department of Zoology, University of Maryland, College Park, MD 20742-4415,  
USA

DONALD M. CASPARY

Department of Pharmacology, Southern Illinois University School of Medicine,  
Springfield, IL 62794-9230, USA

JOHN H. CASSEDAY

Department of Psychology, University of Washington, Seattle, WA 98195-1525,  
USA

ELLEN COVEY

Department of Psychology, University of Washington, Seattle, WA 98195-1525,  
USA

GÜNTER EHRET

Abteilung Neurobiologie, Universität Ulm, D-89069 Ulm, Germany

CARL L. FAINGOLD

Department of Pharmacology, Southern Illinois University School of Medicine,  
Springfield, IL 62794-9629, USA

ROBERT D. FRISINA

Division of Otolaryngology, University of Rochester School of Medicine Roch-  
ester, NY 14642-8629, USA

CRAIG K. HENKEL

Department of Neurobiology and Anatomy, Wake Forest University School of  
Medicine, Winston-Salem, NC 27157-1010, USA

JACK B. KELLY

Laboratory of Sensory Neuroscience, Carleton University, Ottawa, ON K1S  
5B6, Canada

SHIGEYUKI KUWADA

Department of Neuroscience, University of Connecticut Health Center, Far-  
mington, CT 06030-3401, USA

GERALD LANGNER

Institut für Zoologie/Neuroakustik, Technische Universität Darmstadt, D 64287  
Darmstadt, Germany

BRADFORD J. MAY

Center for Hearing Science, Johns Hopkins University School of Medicine, Bal-  
timore, MD 21205-2195, USA

MIGUEL A. MERCHÁN

Cellular Biology and Pathology, Avenida Campo Charro, University of Sala-  
manca, Salamanca 37007, Spain

DOUGLAS L. OLIVER

Department of Neuroscience, University of Connecticut Health Center, Far-  
mington, CT 06030-3401, USA

ALAN R. PALMER

Medical Research Council Institute of Hearing Research University Park, Uni-  
versity of Nottingham, Nottingham NG7 2RD, England

CHRISTINE V. PORTFORS

School of Biological Sciences, Washington State University, Vancouver, WA  
98686, USA

MICHAEL B. PRITZ

Section of Neurological Surgery, Indiana University School of Medicine, Indianapolis, IN 46202-5124, USA

RAMESH RAJAN

Department of Physiology, Monash University, Clayton Victoria 3168, Australia

ADRIAN REES

School of Neurology, Neurobiology and Psychiatry, University of Newcastle upon Tyne, Framlington Place, Newcastle upon Tyne, NE2 4HH, England

GARY J. ROSE

Department of Biology, University of Utah, Salt Lake City, UT 84112-0840, USA

ENRIQUE SALDAÑA

Departamento de Biología Celular, Universidad de Salamanca, Facultad de Medicina, 37007 Salamanca, Spain

BRETT R. SCHOFIELD

Department of Anatomical Sciences and Neurobiology, University of Louisville School of Medicine, Louisville, KY 40292, USA

CHRISTOPH E. SCHREINER

Department of Otolaryngology, University of California at San Francisco, School of Medicine, Coleman Memorial Laboratory and W.M. Keck Center for Integrative Neuroscience, San Francisco, CA 94143-0732, USA

DONAL G. SINEX

Department of Psychology, Utah State University, Logan, UT 84322-2810, USA

ANN M. THOMPSON

Department of Otorhinolaryngology, University of Oklahoma College of Medicine, Oklahoma City, OK 73190-3048, USA

JEFFREY J. WENSTRUP

Department of Neurobiology, Northeastern Ohio Universities College of Medicine, Rootstown, OH 44272, USA

JAMES F. WILLOTT

Department of Psychology, University of South Florida, Tampa, FL 33620, and The Jackson Laboratory, Bar Harbor, ME 04609, USA

JEFFERY A. WINER

Division of Neurobiology, Department of Molecular and Cell Biology, University of California at Berkeley, Berkeley, CA 94720-3200, USA

SHU HUI WU

Department of Psychology, Carleton University, Ottawa, ON K1S 5B6, Canada

TOM C.T. YIN

Department of Physiology, University of Wisconsin, Madison, WI 53706, USA

# Chapter 1

## The Central Auditory System: A Functional Analysis

JEFFERY A. WINER AND CHRISTOPH E. SCHREINER

### 1. INTRODUCTION

Auditory neurons must encode and decipher the spectral, spatial, and temporal properties of sound. These processes are a prerequisite to the subsequent extraction of biologically meaningful signals from a noisy environment and to establish perceptual attributes. The goal of this survey is to provide a framework for thinking about how the auditory system performs these tasks. Such a framework seems essential when considering a network of neurons that extends from the external ear to the cochlea and through many synaptic relays before reaching the cerebral cortex (Figs. 1.1A and 1.2). The internal complexity of this system is apparent in each of the nuclei that comprise the auditory pathway: a given nucleus contains thousands of neurons, each connected to and sharing information with nearby neurons as well as many other, remote nuclei. These neurons thus interact with themselves and their neighbors through many neurotransmitters, a host of synaptic mechanisms, and a wide range of membrane channels. These neurochemical and physiologic features endow the various neuronal classes with an enormous range of response repertoires individually as well as the functional capacity to decipher the temporal, amplitude, and spatial aspects of the auditory signal. In each nucleus, the several kinds of cells each have a particular constellation of response properties that defines them. Some discharge most strongly to sound onset, others respond to noise or frequency-modulated sound, and still others prefer changes in loudness, to name just a few of the many stimulus dimensions that the auditory system routinely extracts, encodes, and represents. Considering the diversity of response types, the possible multiplicity of their connections, and the many levels of interaction between nuclei from the medulla to the cortex, the auditory system presents a formidable challenge to, and an opportunity for, reductionistic thinking.

To confront such a challenge, this system is considered from the perspective of its operating elements, the neurons, and the circuits they form. This viewpoint is useful as the many cells and circuits and the operations they perform have now been characterized in sufficient detail that they can be described in a biological context. This level of discourse is among the first steps toward a mature




Figure 1.1. Primary components of the central acoustic system (**A**) and level-specific distribution of components of the auditory evoked potential (**B**). (**A**) The major nuclei of the auditory pathway; for clarity, the cochlea has been omitted (see Fig. 1.2), as has the posteroventral cochlear nucleus (see Fig. 1.3A). *I*: The cochlear nucleus receives its ascending input via the cochlear nerve. Cochlear nucleus axons then decussate (present example) or terminate ipsilaterally (Fig. 1.2: cochlear nuclei). Intrinsic cochlear nucleus connections are not shown, and the totality of cochlear nucleus targets is extensive,

science of hearing, as these elements constrain the performance of the system by defining its permissible operations. This brief introduction concentrates on the central auditory system from the cochlear nucleus to the cerebral cortex; the cochlea is excluded because it merits a separate and extended treatment in its own right and because the issues it entails are beyond the scope of the themes addressed here.

The emphasis is explicitly functional and it treats the auditory pathway as a network of six interrelated parts whose organization will be explored in a neuro-anatomical and a neurophysiological context. The parts are:

- Cochlear nucleus: genesis of basic response patterns and emergence of parallel pathways.
- Olivary complex: construction of binaural pathways and establishment of time lines.
- Lateral lemniscal nuclei: emergence of chemically specific nuclei.
- Inferior colliculus: site of brain stem convergence and multisensory integration.
- Medial geniculate body: modulation of auditory information by cortical and limbic systems.
- Auditory cortex: interface of hearing and higher order communication and cognitive networks.

Unless noted otherwise, the cat is the reference species because the literature available is so extensive. Surveys of basic auditory system organization should be consulted for other topics (Brodal 1981; Edelman et al. 1988; Popper and Fay 1992; Webster et al. 1992).

---

Figure 1.1. *Continued*

involving up to eight nuclei (Fig. 1.7). 2: Input to the lateral superior olive (*LSO*) is topographic and ends in regular arrays (Feng and Vater 1985) that preserve the tonotopic arrangement in the cochlear nuclei (Fig. 1.3C, D). 3: The *LSO* projects to the contralateral central nucleus of the *IC*; it conserves this topography (Servièrè et al. 1984) and may selectively target specific aural subregions in the *IC* (Ross et al. 1988; Brückner and RübSamen 1995). 4: The tectothalamic input is bilateral (not shown), with the ipsilateral component dominant (González-Hernández et al. 1991). 5: The projection to auditory cortex is entirely ipsilateral, and highly divergent, involving several primary fields (Niimi and Matsuoka 1979) and has a specific laminar arrangement (Sousa-Pinto 1973); the latter feature could redirect thalamocortical influences to parts of the ipsilateral cortico-cortical system (Rouiller et al. 1991) or to the corticofugal network (Prieto and Winer 1999). Although the auditory system is depicted as hierarchical, its operations involve direct descending projections (Fig. 1.18) at almost every level (Aitkin 1986) as well as extensive connections with auditory-related areas of the neocortex (Seltzer and Pandya 1994). **(B)** The brain stem evoked response at seven levels (I to VII) in the auditory system. In humans, the superior olivary complex is in the pons and the cochlear nucleus at the junction of the pons and medulla. The arrival time of each waveform appears on the ordinate, while the abscissa shows their amplitude. For all figures, please refer to the list of abbreviations in the text. [Redrawn and modified from original sources (Chusid 1982; Webster and Garey 1990).]




Figure 1.2. Schematic view of the principal ascending connections of the central auditory pathways. Local circuit arrangements and descending connections have been omitted for clarity, as have smaller nuclei such as the interstitial nucleus of the cochlear nerve root (Hutson and Morest 1996). Several important principles are embodied even in this highly schematic picture of connectivity. First, the output of almost every nucleus targets many centers and is highly divergent; this is accomplished either via branching axons from a single projection cell (Friauf and Ostwald 1988) or by different cochlear nucleus cell types targeting separate nuclei (Cant 1982; Thompson 1998). A second principle is connectional convergence: most nuclei receive multiple sources of input, of which the inferior colliculus is among the best examples, with more than 10 different projections to the central nucleus (Roth et al. 1978; Brunso-Bechtold et al. 1981). The acoustic striae are the main output tracts of the cochlear nuclei; their fibers converge ventral and medial to the superior olivary complex as the trapezoid body (Fig. 1.8A), in which many small nuclei are embedded (Brownell 1975). Third, commissural projections are present at most synaptic stations in the auditory system (Aitkin and Phillips 1984a). Not shown are commissural interconnections between the cochlear nuclei (Cant and Gaston 1982); an exception is the medial geniculate body, which has no commissural pathway. Fourth, there is an array of internal circuitry in every nucleus, and a nucleus- and sometimes a



## 2. THE COCHLEAR NUCLEI

In the cochlear nuclei the principles emerge that will dominate the auditory system at subsequent levels of processing (Osen 1988; Rhode and Greenberg 1992). The major themes are that there is (1) an orderly input that preserves the topography of frequency established in the cochlea (Osen 1970); (2) massive divergence of auditory nerve projections creates multiple new and nonequivalent maps from a single cochlear origin (Osen 1972); (3) great variety in structure and function among the specialized postsynaptic neurons that process the output of the auditory nerve (Osen 1969; Cant 1992); (4) an abundant number of inhibitory neurons to constrain, modulate, and refine the primary excitatory input (Adams and Mugnaini 1987); and (5) an array of descending projections from higher order auditory structures that provide feedback (Kane and Conlee 1979). The cochlear nucleus will be considered in slightly more detail because a large literature is available and because the functional principles that govern it are repeated and elaborated at many further levels.

The cochlear nuclei are among the first parts of the central auditory system to have evolved, and they are present in every vertebrate (Baird 1974). In amphibians and reptiles they form a semicircular mass of a few hundred neurons on the lateral edge of the medulla (Gregory 1974), and in carnivores and primates they represent a distinct lobe of highly differentiated cells beside the cochlea and the cerebellum. The auditory nerve is thus among the shortest cranial nerves in most species, and the thick coat of myelin on primary afferent axons and even covering ganglion cell perikarya ensures rapid transmission of acoustic impulses to the medulla. The auditory nerve axons spiral as they traverse the cochlea to reach the brain. They enter the cochlear nucleus rostroventrally, and then divide into descending and ascending branches (Ryugo 1992). This bifurcation redistributes the single sheet of the cochlear sensory epithelium across a volume; because each point (or, more properly, strip) represents a limited range of frequency (Fig. 1.3) and because each frequency is in topographic registration with its neighbors, the cochlear nucleus contains at least two independent maps of the frequency spectrum (Rose et al. 1959) (Fig. 1.4). The volume that corresponds to a frequency is termed isofrequency, and an isofrequency contour may contain several thousand neurons arrayed in flattened sheets and across which the cochlear axons are distributed more or less evenly (Fekete et al. 1984) (Fig. 1.3). This or analogous arrangements in which a distal map

---

Figure 1.2. *Continued*

species-specific distribution of interneurons and their synapses (Winer and Larue 1996). Fifth, many projections are reciprocal (not shown), especially those between the brain stem nuclei (Conlee and Kane 1982) and those of the thalamus and cortex (Colwell 1975). [Redrawn and modified from original sources: main diagram (Gulick et al. 1989); cochlea (Brödel 1946).]




Figure 1.3. Some physiologic and neurochemical aspects of cochlear nucleus organization. (**A**, **B**) Cochlear nucleus subdivisions (Osen 1969; Brawer et al. 1974). In this and subsequent figures relating to other auditory nuclei, only the principal subdivisions are shown. (**C**) Spatial and nuclear arrangement of characteristic frequency as expressed in isofrequency lines (*dashed*) in the anteroventral and posteroventral cochlear nuclei (*AVCN* and *PVCN*). The systematic representation of frequency reflects the topography of cochlear nerve input (Leake-Jones and Snyder 1982) and it is conserved in both the *AVCN* and *PVCN*. Noteworthy features include the large spatial representation of frequencies <10 kHz, and the complexity of laminar overlap near nuclear borders and in regions where cochlear nerve axons bifurcate and their complex spatial geometry (Osen 1970) distorts a simple laminar organization; elsewhere in the central auditory system a spatial

of frequency is expanded, contracted, or distorted will be repeated many times between the cochlear nuclei and the cortex.

A constraint on a perfectly uniform distribution of the frequency spectrum is the biological significance of particular subregions: frequencies essential in communication are likely to have an enlarged representation (Suga and Jen 1976), and the degree of this expansion can match that in the cochlea itself (Feng and Vater 1985). This principle recalls the disproportionate sensory and motor neural representation of the primate hand relative to that of the foot (Kaas et al. 1984), or the central (foveal) part of the retina compared to its more peripheral regions (Tusa et al. 1981).

The divergence of the auditory nerve is not simply a bifurcation: it entails a structural reorganization of the postsynaptic neurons as well. The axons of the descending branch terminate in delicate boutons that will make synapses on the dendrites and cell bodies of neurons in the dorsal cochlear nucleus (DCN). These synapses—their number, location, and shape—will profoundly influence the physiologic behavior of the postsynaptic neurons. For example, synapses closest to the axon hillock, which is the electrogenic membrane controlling the discharge behavior of the cell, will have the maximum efficacy (Peters et al. 1991). The divergence of connectivity multiplies the channels of information exiting the cochlear nuclei and which, after reaching a variety of medullary auditory centers, converge on the inferior colliculus (IC). (See Fig. 1.5.)

The potential functional ramifications of the diverging projections from the auditory nerve are magnified by the great variety of morphologic cell classes that is targeted. As a consequence of morphology, convergence, and local circuitry, these cell classes differ functionally in distinct ways including their temporal response pattern, spectral selectivity, and discharge regularity. To highlight differences in their response properties, we next compare two types of cochlear nucleus neurons. Among the most conspicuous of the many types of DCN neu-

---

Figure 1.3. *Continued*

compression of higher frequencies occurs (Morest 1965). This distortion is partly an artifact of collapsing a three-dimensional geometry onto two dimensions. [(C, D) Redrawn and modified from original sources (Bourk et al. 1981 with permission from Elsevier Science).] (D) In the AVCN the expanded volume devoted to low frequencies is even more pronounced than in the dorsal cochlear nucleus (DCN). (E) The distribution of GABAergic and glycinergic neurons follows specific patterns in the cochlear nucleus. In the DCN, both are concentrated in layers I to III; both neurons and axon terminals are plentiful here and much sparser elsewhere. GABAergic neurons may also colocalize glycine (Osen et al. 1990) and there is evidence for simultaneous corelease of both molecules (Jonas et al. 1998). In this and other nuclei, the vast majority of the other neurons are glutamatergic (Saint Marie 1996) or aspartatergic (Altschuler et al. 1981), as is the cochlear nerve (Wentholt 1985). [(E, F) Redrawn and modified from original GABA observations (Adams and Mugnaini 1987); redrawn and modified from original glycine observations (Wentholt et al. 1987).] (F) In the AVCN, there are few GABAergic or glycinergic neurons.




Figure 1.4. Representation of two basic principles of auditory brain stem connectivity—divergence and convergence—and their complementary functional roles. 1, 2: Spiral ganglion cell axons divide and project to subdivisions of the cochlear nucleus (Lorente de Nó 1933, 1981). This creates independent maps of characteristic frequency (Rose et al. 1959). 3, 4: Topographically matched projections from each anteroventral cochlear nucleus (AVCN) have different targets in the medulla: the ipsilateral projection ends in the lateral superior olive (LSO) (Cant and Casseday 1986), while the contralateral AVCN projects to the medial nucleus of the trapezoid body (MNTB) (Warr 1966), whose axons converge on the other primary dendritic trunk of the same postsynaptic neuron (Spangler et al. 1985; Vater et al. 1995). This creates binaural receptive fields among superior olivary complex neurons (Tsuchitani and Boudreau 1966; Tsuchitani and Johnson 1991).

rons is the fusiform cell; it has a pyramidal cell body and an extensive dendritic tree (Brawer et al. 1974). These processes act as postsynaptic sites: their thick trunks and slender spines are studded with presynaptic axon terminals, most from the auditory part of the eighth cranial nerve (Cant 1992). A consequence of this pattern of axodendritic organization is that it requires spatiotemporal summation to elicit a postsynaptic response that differs significantly from the rate of spontaneous discharge. A constraint is that the postsynaptic dendritic propagation of all-or-none signals is decremental because the dendritic membrane typically lacks the regenerative channels found in myelinated axons and that permit them to send spike codes over long distances while conserving amplitude or fidelity. Thus, inputs to the most distal dendrites usually have correspondingly less influence than terminals near the axon hillock (Jack et al. 1975), unless they are amplified at dendritic branch points (Stuart et al. 1997). Dendritic input may require a coordinated volley of presynaptic impulses arriving optimally to achieve the spatial and temporal summation necessary to cross the threshold of the postsynaptic neuron. The threshold and the electrotonic properties of the dendrites act as adaptive filters to reduce spontaneous discharges in an auditory environment filled with ambient noise. Many neurons respond best to changes in sound pressure level (dynamic response), while others prefer steady-state (tonic) conditions (Young 1984). These extremes effectively reduce noise yet preserve the ability to respond to static stimuli or to signals with a wide dynamic range.

The physiologic types of cochlear nucleus neurons have been determined using tonal or noise stimuli to generate peristimulus time histograms that provide insight into basic coding properties (Fig. 1.6). This strategy was applied originally in the auditory nerve to classify the responses of cochlear ganglion cell fibers (Kiang et al. 1965), and it has revealed the temporal profile of neuronal responses in many regions (Kiang et al. 1973). The discharge pattern of intracellularly identified fusiform cells in the DCN shows a pauser/buildup pattern (Fig. 1.6D, E). This consists of a delayed discharge after stimulus onset, a burst of spikes concentrated in the first few milliseconds, an abrupt decline to near-zero spikes, and, finally, a subsequent slow growth in spike rate punctuated by briefer reductions; the unit never reestablishes the spike density at onset, and the discharge can persist or even increase long after the poststimulus period (Rhode et al. 1983b).

As viewed in the electron microscope, many of these physiologic attributes have ultrastructural correlates that could support them. Thus, fusiform cells receive synapses indicative of independent excitatory and inhibitory input on their distal dendrites. The excitatory endings (Oliver et al. 1983) are axodendritic synapses of eighth nerve origin and terminals from granule cell parallel fibers, which project in long rows across these dendrites (Fig. 1.7B: pf). Prospective inhibitory input arises from four types of neuron—each with a specific synaptic arrangement on the postsynaptic cell. Stellate, vertical, Golgi, and cartwheel cells each project to fusiform cell dendrites (Berrebi and Mugnaini 1991). Some of these neurons may be glycinergic (Wenthold and Hunter 1990) and others




Figure 1.5. Components of the striae of Held and von Monakow and the trapezoid body (Fig. 1.2) shown schematically. Their medullary target nuclei each project to the inferior colliculus. **(A)** Trapezoid body, ventral and medial components. **(B)** Intermediate acoustic stria and trapezoid body. [Redrawn and modified from original sources (Warr 1982; Irvine 1986).] Key for this and subsequent schematics: *open boxes with heavy outlines*, origins of projections; *solid boxes with white lettering*, targets of projection; *lines*, axons; *dotted line*, midline; *brackets*, fiber tracts. Ascending pathways are shown in these diagrams unless indicated otherwise (Fig. 1.18).

$\gamma$ -aminobutyric acid (GABA)-ergic (Adams and Mugnaini 1987); this suggests cell-specific interactions that mediate the discharge properties and influence the receptive field profile of postsynaptic neurons. Although the exact nature of the dynamic interactions among these convergent inputs on fusiform cells is not known, they could contribute to the periodic fluctuations and gradual decline in rate seen in pauser/buildup units. Overall, then, fusiform cells perform a profound and complex spectral and temporal transformation of the auditory nerve input whose ultimate functional purpose may contribute to several, still largely elusive, central auditory tasks.

In the anteroventral cochlear nucleus (AVCN), the response of a different type of neuron, the bushy cell, can be contrasted with, and complements that of, the DCN fusiform cell. Here, ascending auditory nerve fibers end on bushy cell somata. These neurons differ fundamentally from fusiform cells (Lorente de N3 1981). Bushy cells have only one or two short dendrites, and their predominant synaptic ending is an auditory nerve axon, the endbulb of Held, which forms a massive cuplike terminal that encloses the perikaryon, with several such endings and with virtually identical tuning curves converging onto a cell (Cant and Morest 1979). Thus, the dendritic arbors of these cells are modest, while the most potent synaptic input from the endbulbs terminates near the axon hillock. The net probability of presynaptic signals evoking a postsynaptic discharge therefore is effectively unity (Rhode et al. 1983a). Moreover, these neurons discharge preferentially to the earliest, onset-related features of the stimulus. The tonotopic representation in the AVCN is unusual: frequencies  $<5$  kHz are mapped selectively and expansively. The electrophysiologic profile of the response is also unique because of the size of the endbulb and the large-amplitude prepotential in the afferent fiber, neither of which is present in the DCN (Bourk et al. 1981). Frequency tuning, rate-intensity function, and discharge regularity of bushy cells closely resemble their auditory nerve inputs. Hence, they perform only a minimal transformation of the input and, rather, ensure a faithful information transmission beneficial for processes in the following stations, such as binaural comparators. The fusiform and bushy cells are just two examples of the wide range of structural/functional patterns represented in the cochlear nucleus, and they embody only some of the nuclear- and cell-specific response patterns (Rhode 1991). More than 20 kinds of neuron have been identified in the cochlear nucleus (Brawer et al. 1974).

The presence of local circuitry within the cochlear nucleus itself further refines signal processing. These circuits may function to reject unwanted signals, amplify weak inputs, provide inhibition that enhances the signal-to-noise ratio (Berrebi and Mugnaini 1991), or serve a disinhibitory role. Because the responses of auditory nerve axons are sharply tuned to specific frequencies with no inhibition, and because stimulation at unfavorable frequencies elicits a negative or no response (Kiang et al. 1965), such circuitry intuitively might not seem necessary. In the cochlear nucleus, however, the convergence of input from many different fibers onto a single neuron could blur the fine spatial and temporal patterns of activity, or help to retune or focus the cell's response area to a particular aspect of the stimulus, such as in monaural echo suppression




Figure 1.6. A categorical description of cochlear nucleus physiologic response profiles and their correlation with neuronal types. The frequency-response areas (1, 3 . . . 11) depict the tuning curve types, while the peristimulus time-histogram (2, 4 . . . 12) shows the interplay of intrinsic membrane properties and the excitatory–inhibitory interactions that shape the cell’s dynamic response to tones. The inhibitory contribution to the receptive field is also visible in the frequency-response area (D: 7, E: 9, F: 11). Although



(Wickesberg and Oertel 1990). Inhibition can impose unique filter properties on the tuning behavior and modify the neural response profiles. Thus, type II DCN interneurons are sharply tuned, with large excitatory response areas flanked by lateral zones of inhibition, and they have little spontaneous activity (Young and Voigt 1982). Type II cells are thought to have local feedforward projections onto type IV neurons, the fusiform or giant cells. Thus, type IV cells cease firing when the type II units increase their spike rate. The type II discharge creates powerful inhibitory subregions in the tuning curve of the type IV cell; the tight temporal cross correlation in spike discharge indicates that these neurons may be linked monosynaptically. Such circuits could underlie the type IV unit's robust response to broad-band noise and the inhibitory effects of pure tones (Young 1984). This circuit is analogous to one made in the cerebral neocortex by inhibitory interneurons (basket cells) on pyramidal cells (White and Keller 1989; Prieto et al. 1994a) and by similar kinds of IC cells (Oliver et al. 1994). Such circuits may be a common feature in many nuclei for regulating the output of projection neurons and for reorganizing their receptive fields in response to environmental demands (Bjardahl et al. 1998; Kilgard and Merzenich 1998a). It remains to elucidate in equal detail the functional connectivities among the many other cochlear nucleus neuronal populations.

Already in this first nucleus of the auditory pathway, several themes are established that have major consequences in subsequent processing and reflect equivalent mechanisms throughout the auditory system. Many of the different cell types distribute the transformed information to specific proximal targets. Although there is not necessarily a one-to-one relationship between type of projecting neuron and target nucleus or cell type, at least six parallel streams of diversely transformed or modulated auditory information emerge. Each stream embodies a different preprocessing principle that is likely best suited for further, specific central processing tasks. Accordingly, the auditory signal is decomposed not just in terms of its frequency content but also in terms of a variety of other stimulus features that are composed of complex aspects of spectral, temporal, and spatial signal attributes. The particular contributions of most of these concurrent streams originating in the cochlear nucleus to the process of feature extraction, object recognition, and, ultimately, perception remain to be established.

---

Figure 1.6. *Continued*

a detailed treatment of the individual categories is beyond the scope of this chapter, the range of responses in the primary auditory nuclei suggests that peripheral receptor diversity contributes little to coding, while in the somatic sensory system peripheral receptor diversity plays a major role in the establishment of separate processing streams (Sinclair 1981). A major basis for auditory coding is thus local circuitry, although diversity in hair cell responses (Ashmore 1991) and spiral ganglion cell behavior (Kim and Molnar 1979) and differential projections (Brown and Ledwith 1990) must contribute to this pattern. [Redrawn and modified from the original sources (Evans 1982 with the permission of Cambridge University Press; Webster and Garey 1990).]




Figure 1.7. Some projections of cochlear nucleus neurons that may contribute to parallel processing in the superior olivary complex, lateral lemniscal nuclei, and inferior colliculus. (A) In a parasagittal schematic, cartoons of the principal cochlear nucleus neurons demonstrate that they have diverse ipsi- and contralateral targets. Differences in the temporal coding among these cell types (Fig. 1.6: 2, 4 . . . 12) suggest that the streams leaving the cochlear nucleus are integrated in higher centers (when they converge) or contribute to parallel pathways (when they diverge) (Warr 1982). [Redrawn and modified from

### 3. THE OLIVARY COMPLEX

The olivary complex consists of four (humans) to nine (cats) nuclei (Moore and Moore 1971) that have several complementary physiologic roles. Neurons in the LSO integrate the monaural input of the cochlear nuclei to derive intensity-difference sensitive binaural signals, mainly for high frequencies, and to send this information to the lateral lemniscal nuclei (LLN) and the IC (Schwartz 1992). Neurons in the medial superior olive (MSO) encode phase relationships and delay sensitivity from the two ears (Fig. 1.8D), mainly from lower frequencies; these signals are essential for accurate spatial localization (Yin and Chan 1990). Neurons in the medial nucleus of the trapezoid body (MNTB) contribute to the creation of binaural subtypes via their inhibitory input to the LSO (Guinan and Li 1990). The construction and variety of binaural interactions represent a situation in which emergent function can be directly related to a particular neural circuit (Tsuchitani and Johnson 1991). The species-specific variability in the size, shape, and disposition of the olivary nuclei (Schofield and Cant 1991) suggests that ecologic factors (Masterton et al. 1975) might induce structural changes related to evolutionary adaptations in hearing (Papez 1929). All subdivisions have a tonotopic organization (Fig. 1.8B).

The striking precision of connectivity in the central auditory system is exemplified in the ascending projection onto principal cell lateral dendrites in the LSO. In this example, one primary dendrite is postsynaptic to input from the ipsilateral AVCN, whose output cells may be aspartatergic and are likely excitatory (Oliver et al. 1983). The other dendrite is the target of axons from cells in the MNTB (Cant and Casseday 1986), which are glycinergic and probably inhibitory (Bledsoe et al. 1990). The contralateral AVCN provides the excitatory drive to the MNTB. This convergence has important functional consequences, as olivary principal cells integrate ipsilateral (excitatory) and contralateral (inhibitory) input whose chemical signals can facilitate or suppress postsynaptic neurons. This convergence creates the first binaural neurons in the auditory system and their discharge pattern integrates excitation and inhibition.

---

Figure 1.7. *Continued*

Aitkin (1989).] **(B)** Schematic view of local circuits in layers 1 to 3 of the dorsal cochlear nucleus, showing the main types of neurons and their synaptic relationships. Some connections, for example, between stellate cells (3), have not yet been demonstrated. The major relationships shown here in the guinea pig depict cartwheel cells (4) projecting to the superficial and deep dendrites of pyramidal cells (5); the cartwheel cell input to the vertical cell (6) is likewise unconfirmed. Parallel fibers (*pf*) arising from granule cells (1) provide excitatory input to the layer 1 inhibitory neurons and to the apical dendrites of pyramidal cells. A '?' indicates a possible projection. Arrowhead, gap junction between stellate cell dendrites. [Redrawn and modified from original sources (Berrebi and Mugnaini 1991).] Neurons in *black* are considered inhibitory (Osen et al. 1990).




Figure 1.8. Basic organizational features of the superior olivary complex. (A) The individual nuclei of the superior olivary complex are well developed in the cat. As these nuclei have species-specific sizes and shapes (Harrison and Feldman 1970; Schofield and Cant 1991), proposals for homologies require careful analysis (Carr and Friedman 1999). (B) The principal nuclei each have a tonotopic organization in which frequencies <10 kHz are overrepresented at the expense of neurons with higher characteristic frequencies, a trend most evident in the medial superior olive. [Material in B, C redrawn from original source (Guinan et al. 1972b, Taylor and Francis Ltd. <http://www.tandf.co.uk/journals>).] (C) Summary profile of the main response types found in the superior olivary complex. The classification scheme (“Primary-like” . . . “Long latency”) integrates the frequency response areas (Fig. 1.6, left) and the peristimulus response histograms (Fig. 1.6, right), which resemble those derived for neurons in cochlear nucleus sub-

An interesting issue is the existence of the MNTB in humans (Richter et al. 1983). Its absence would be unexpected because human performance in a test for discriminating the minimum audible azimuthal angle in sound localization is superior to that of all rodents and carnivores, and better than that of the porpoise for both noise and clicks (Fay 1988). Perhaps nuclei other than the MNTB have assumed the functions it once may have supported, or humans use other pathways for accurate localization, or the MNTB is undistinguished anatomically. There are certainly massive species differences among medullary nuclei and that undoubtedly have behavioral correlates (Harrison and Irving 1966). A second theme, which is preserved at subsequent levels of synaptic processing, is that two (or more) representations of a sensory attribute are interleaved across the several spatial axes of the LSO. One is related to frequency and has approximately a medial-to-lateral orientation; the other is concerned with aurality and runs roughly dorsoventral or orthogonal to the axis of frequency. It is highly probable that single nuclei do not represent one function only. A third idea that

---

Figure 1.8. *Continued*

divisions. [Redrawn and modified from original sources (Evans 1982); some material redrawn from other sources (Webster and Garey 1990).] The responses in the superior olivary complex not shown in this key are “Off” (in which stimulus onset suppresses all spontaneous activity, much like “Inhibitory” in Fig. 1.6F) and “Long latency” (in which the stimulus evokes a delayed response). **(D)** Circuit summary of axons in the medial superior olive from the anteroventral cochlear nucleus [Redrawn from original source (Beckius et al. 1999. Copyright by the Society for Neuroscience.).] The original, Jeffress, model is indicated by the *solid lines* (Jeffress 1948). Contralaterally, the shortest axons reach the central and caudal medial superior olive; ipsilaterally, the most caudal part was the target of the short fibers. Some axons (*dashed lines*) had a more restricted pattern of termination. **(E)** A neurochemical profile of the superior olivary complex, showing transmitter specific differences among nuclei. Some nuclei have glycinergic neurons only (medial nucleus of the trapezoid body), and others have both GABAergic and glycinergic cells (lateral superior olive). Projections of specific cell types to the inferior colliculus and elsewhere may be segregated in some cases and convergent in others (Oliver et al. 1997). Even within a nucleus such as the lateral superior olive, the distribution of GABAergic neurons and axon terminals (puncta) can be nonuniform, with gradients of immunoreactivity that are parallel to or cross the tonotopic axis (Winer et al. 1995). Observations on GABA taken from an original source (Adams and Mugnaini 1990). The periolivary cholinergic neurons have a special role in hearing as they are the primary source of the olivocochlear system. These neurons form a heterogeneous group (Vetter and Mugnaini 1992), most of which are dispersed on the margins of the ipsilateral lateral superior olive (about 50%) and medial to the medial superior olive (11%), while the crossed component is situated less in periolivary regions (15%) and medial to the medial superior olive (Lieberman and Brown 1986). This system thus has multiple nuclear origins and at least some neurons receive descending input from the inferior colliculus (Vetter et al. 1993). Although the olivocochlear system is numerically small—ranging from 475 (mouse) to 2346 neurons (guinea pig)—it has widespread effects on cochlear processing (Warr 1992).

is reinforced by this convergence is that cell shape can have implications for functional arrangements in some cases. Thus, some principal cells in the LSO have polarized dendritic fields ideal for segregating ipsi- and contralateral afferents. A fourth observation is that neurons with binaural properties will be found at most subsequent levels of the auditory pathway. This implies that the processing at each synaptic station extracts different information from a stimulus; it may serve to explain the otherwise bewildering multiplicity of nuclei, maps, and representations in audition. A fifth issue is developmental, as the accuracy of innervation from the converging inputs must be high, and errors in connectivity low, for optimum performance (Sanes et al. 1990). Finally, the emergence early in the ascending auditory pathway of binaural neurons underscores a fundamental difference between it and the visual system, where the appearance of binocular neurons is deferred until the cerebral cortex (Ferster and LeVay 1978; LeVay et al. 1978).

The LSO has an important redistributive role: it sends axons bilaterally to the IC, with the most lateral, low-frequency (Tsuchitani 1977) projections ending ipsilaterally and the more medial, high-frequency portions terminating contralaterally (Elverland 1978; Glendenning and Masterton 1983). This projection has been characterized as an acoustic chiasm, a decussation like that in the visual and somatic sensory systems (Glendenning et al. 1985) in which a sensory hemifield is represented contralaterally within the brain. The idea of parallel pathways carrying different types of synaptic information is reinforced by the finding that the ipsilateral input has two neurochemical components, one of which is glycinergic and constitutes 41% to 60% of the projection neurons, while the contralateral pathway has only 4% glycinergic cells (Saint Marie et al. 1989). Such neurochemically specialized projections appear to be absent in the cochlear nucleus, excepting a glycinergic projection with a proposed role in monaural echo suppression (Wickesberg and Oertel 1990), and they are absent in the somatic sensory and visual systems.

A further novel feature of the olivary nuclei is the emergence of the olivocochlear efferent pathway. This system arises from fewer than 2000 neurons in the periolivary cell groups, a collection of small nuclei that are the origin of the centrifugal projection to peripheral receptors, principally to the outer hair cells. These primarily cholinergic (Vetter and Mugnaini 1990) centrifugal cells (yet another example of auditory neurochemical specificity) may influence the micromechanical properties of outer hair cells, thereby affecting basilar membrane dynamics and influencing the gain of the inner hair cells (Liberman and Brown 1986; Warr 1992). The olivocochlear system exemplifies the neurochemical and connectional diversity of the olivary complex in particular and the auditory system in general. Again, there are no parallels to the centrifugal pathways in the visual or somatic sensory systems in mammals.

In addition, another descending projection arises in the olivary complex, from the lateral and medial nuclei of the trapezoid body. These neurons project heavily, bilaterally, and topographically onto most parts of the cochlear nucleus (Spangler et al. 1987). This theme of prominent descending projections is pre-

served in the IC and the auditory cortex (and less so in the thalamus), and it is a robust feature in the subcortical auditory system. Such descending connections are absent in visual and somatic sensory systems.

The olivary nuclei, then, are the clearest examples of task-specific structures in the auditory system yet identified. The LSO and the MSO contain prime examples of computational maps, that is, systematic construction and representation of aspects of sound sources that cannot be derived from each ear in isolation but require convergence of different information streams. The mechanisms that lead to these maps, namely the use of coincidence detection and delay lines, also reflect general elements of neuronal processing evident in many other stations and a variety of tasks throughout the auditory system.

#### 4. THE LATERAL LEMNISCAL NUCLEI

The lateral lemniscus (Gr., *ribbon*) consists of three nuclei (dorsal, intermediate, and ventral) and the axons in which they lie along the lateral surface of the brain stem, near the transition from the pons to the midbrain. The nuclei extend from the rostral pole of the LSO to the ventral tip of the inferior colliculus. Many of their connections and physiologic properties resemble those of collicular neurons (Aitkin et al. 1970); however, certain structural and neurochemical arrangements are unique, such as the virtually pure GABAergic population of the dorsal nucleus (Adams and Mugnaini 1984) and the prominence of glycinergic cells in the ventral nucleus (Winer et al. 1995). These nuclei appear to be present in all mammals studied so far (Schwartz 1992). They have as wide a range in size and shape in different species as any auditory nucleus, from modest in rats (Bajo et al. 1993; Merchán et al. 1994) to well developed in cats (Berman 1968) (Fig. 1.9B: 2) to hypertrophied in some echolocating bats (Pollak and Casseday 1989). They do not represent a mandatory synaptic relay as some projections from the cochlear nuclei (Oliver 1984) or superior olivary complex (Glendenning and Masterton 1983) bypass the LLN en route to the IC (Fig. 1.10B: 1). Perhaps the LL nuclei are essential only in some facets of auditory processing. They differ from the IC and more rostral auditory centers in having functional and connectional affiliations that are exclusively auditory. Current thinking suggests that they are part of a brain stem binaural and monaural system parallel to those in the LSO (Irvine 1986) and intimately related to the olivary nuclei and the IC.

A striking feature that sets the LLN apart from many other brain stem nuclei is their neurochemical specificity and heterogeneity. About 85% of dorsal nucleus neurons are GABAergic, only 18% of intermediate nucleus cells are either GABAergic or glycinergic, and >80% of ventral nucleus cells are glycinergic (Saint Marie et al. 1997) (Fig. 1.11D, E). These values should be interpreted with caution as GABAergic neurons may be immunoreactive for glycine, and vice versa; the significance of this colocalization is unknown (Wenthold and Hunter 1990; Winer et al. 1995; Saint Marie et al. 1997). Much the same pattern




Figure 1.9. Connections between the cochlear nuclei and the lateral lemniscal nuclei, and the nuclei of the lateral lemniscus and the inferior colliculus; descending projections from the inferior colliculus to the cochlear nucleus are also present. **(A)** Each cochlear nucleus subdivision has different targets, and a different strength of projection, to the lateral lemniscal nuclei. The dorsal nucleus of the lateral lemniscus receives stronger input from the superior olivary complex (Fig. 1.10B: 2) than from the cochlear nuclei. **[A, B** redrawn from the original source (Irvine 1986).] **(B)** The nuclei of the lateral lemniscus share many projections with the inferior colliculus, for example, from the cochlear nucleus (Fig. 1.13B: 3) and the superior olivary complex (Fig. 1.14B: 2). It is unknown whether the postsynaptic targets of these projections are also segregated or if they converge onto single neurons (Oliver et al. 1997). **1:** Each posteroventral cochlear nucleus projects to the ventral nucleus of the lateral lemniscus, which then sends fibers



is seen in the mustache bat, with three differences: when both types of immunoreactivity are present, they may be segregated spatially. Second, the intermediate nucleus has clusters of immunoreactive cells associated with physiologic subregions: glycinergic neurons predominate in the presumptive areas of the 30- and 90-kHz components of the bat's echolocation call, while the constant-frequency, 60-kHz territory has many GABAergic neurons (Winer et al. 1995). This suggests a functional dissociation between the dorsal nucleus, which provides binaural inhibition to both ICs (Markovitz and Pollak 1994), and the ventral nucleus, which is part of a monaural system (Fig. 1.11C). Third, neurons in the dorsal and ventral nuclei of the lateral lemniscus show interesting temporal properties with regard to spiking precision (Covey and Casseday 1991), onset, and duration, suggesting modulatory or even instructive roles in the shaping of timing information in their respective targets. Fourth, there is strong evidence for further physiologic subdivisions within the bat LLN as indicated by the spectral tuning, spontaneous activity, and timing of neural discharge in the columnar part of the bat ventral nucleus of the lateral lemniscus. Despite these nuclear and physiologic differences among the LLN, their specific role in hearing remains largely uncertain. Overall, the LLN add further branches of segregated processing streams and additional diversity and complexity to the expression and role of functional stream segregation and information integration.

## 5. THE INFERIOR COLLICULUS

The IC (La., *small posterior hill*) is a dome-shaped structure that is among the largest auditory nuclei in the vertebrate brain and it is virtually an obligatory synaptic terminus for ascending input to the medial geniculate body (MGB) (Aitkin and Phillips 1984b). Its size and its many connections suggest that it has critical roles in both the ascending and descending limbs of the auditory system. In reptiles and amphibia, its homologue, the torus semicircularis, is a massive prominence that forms the roof of the caudal midbrain (Foster and Hall 1978). The importance of the IC is accentuated by the relatively reduced size of the homologue of the MGB in frogs (Hall and Feng 1987) and alligators (Pritz 1974a,b).

---

Figure 1.9. *Continued*

to the central nucleus of the inferior colliculus. 2: The dorsal nucleus of the lateral lemniscus, which receives input mainly from cochlear nucleus subdivisions outside the posteroventral cochlear nucleus (A), also projects onto the central nucleus. The central nucleus thus receives direct and indirect input from all cochlear nucleus subdivisions. 3: Besides its integrative role in the ascending auditory system, the inferior colliculus sends descending projections to the cochlear nucleus (CN). Thus, the dorsal cortex (DC) of the inferior colliculus, which receives only sparse ascending input from the brain stem, projects to the dorsal cochlear nucleus (Weedman and Ryugo 1996a). (C) Schematic of tectocochlear projection. [Redrawn from the original source (Conlee and Kane 1982).]




Figure 1.10. Ascending projections from the olivary complex to the lateral lemniscal nuclei and the inferior colliculus. **(A)** Schematic view of connections. Each lateral lemniscal nucleus has both common and unique input from subdivisions of the superior olivary complex. [Redrawn from original source (Irvine 1986).] **(B)** The lateral superior olive (*LSO*), the medial superior olive (*MSO*), and the medial nucleus of the trapezoid body (*MNTB*) each have convergent projections to the dorsal nucleus of the lateral lemniscus. *1*: The input to the dorsal nucleus of the lateral lemniscus is heterogeneous chemically as well: some contralateral projections from the lateral superior olive (*LSO*) may be glutamatergic (Glendenning et al. 1992; Saint Marie 1996); parts of the ipsilateral input are GABAergic (*black cells*) while the neurons in the *MNTB* are glycinergic (Bledsoe et al. 1990; Spirou and Berrebi 1997). *2*: The *MNTB* also projects to the intermediate nucleus of the lateral lemniscus. *3*: GABAergic superior olivary complex neurons also end in the central nucleus (*CN*) of the inferior colliculus, reaching it presumably before the parallel GABAergic projection from the dorsal nucleus of the lateral lemniscus.

A second feature is that the IC may have the most diverse connections of any auditory structure. These include input from almost all parts of the cochlear nucleus (Oliver 1984, 1987), from much of the olivary complex (Glendenning et al. 1992), from each of the LLN (Saint Marie et al. 1997), and from every auditory cortical area (Winer et al. 1998). Besides these extrinsic projections, there is a rich array of commissural (Aitkin and Phillips 1984a) and intrinsic projections (Oliver et al. 1991). The IC itself projects to almost all brain stem nuclei that project to it (Huffman and Henson 1990), and it sends axons bilaterally to the auditory thalamus (Andersen et al. 1980). Most of these connections are topographic; perhaps they are related to processes embedded in the tonotopic arrangement of characteristic frequency in the central nucleus.

The IC contains three principal divisions: central nucleus, lateral nucleus, and dorsal cortex (Fig. 1.12A). Each has several nuclei that differ in neuronal structure (Morest and Oliver 1984) and connections (Rockel and Jones 1973a,b) and probably in their ontogenetic sequence (Aitkin and Moore 1975; Aitkin and Reynolds 1975; Binns et al. 1995) and in their functional (Aitkin 1979) roles. The central nucleus is exclusively auditory (Aitkin et al. 1994) and it is essential for normal hearing (Jenkins and Masterton 1982); the lateral nucleus is multisensory (Aitkin et al. 1978) and the target of considerable nonauditory input (Morest and Oliver 1984), the dorsal cortex receives most of its projections from the cerebral cortex (Winer et al. 1998) and its role in hearing is unknown (Oliver and Huerta 1992).

The arrangement of afferent input from brain stem neurons to the central nucleus preserves the topographic relations in the various afferent populations (Figs. 1.13 and 1.14). Thus, low-frequency (lateral) parts of the LSO (Guinan et al. 1972a,b) project to the dorsal part of the central nucleus (Glendenning and Masterton 1983), where low frequencies are represented preferentially (Merzenich and Reid 1974). Here, projections from the most dorsal, low-frequency parts of the dorsal nucleus of the lateral lemniscus (Aitkin et al. 1970) also terminate (Shneiderman et al. 1988) among those from other brain stem nuclei (Oliver et al. 1995). Because there is only one map of the basilar membrane in the central nucleus of the IC (Aitkin 1986), and because input from intrinsic (Oliver et al. 1991), commissural (Hutson et al. 1991), and many other sources (González-Hernández et al. 1996) converges along with brain stem input, perhaps each projection has a considerable topographic precision, or elaborate local inhibitory connections that could suppress the activity of tonotopically inappropriate input. The evidence available favors the first interpretation but does not necessarily exclude the second.

A feature of IC organization that has attracted considerable attention is the structural basis for frequency-specific arrangements, the isofrequency or frequency-band lamina, which contains thousand of neurons with a similar characteristic frequency (Fig. 1.12B). One mechanism for enhancing the precision of ascending axons ending within the inferior colliculus is to restrict spatially the afferent input available to the postsynaptic neuron by adjusting the size and shape of the postsynaptic cell's dendritic domain and that of the associated




Figure 1.11. The lateral lemniscal nuclei. (A) Nuclear subdivisions; the functional status of the cuneiform (*Cu*) and sagulum (*Sa*) nuclei is uncertain, although the latter has widespread connections with the IC, the MGB, and the auditory cortex (Beneyto et al. 1998). [A, B: Adapted from published data (Aitkin et al. 1970).] (B) Distribution of characteristic frequency in a transverse section of the dorsal nucleus of the lateral lemniscus (Aitkin et al. 1970). (C) Mosaic pattern of characteristic frequency within the ventral lateral lemniscal subdivisions is consistent with a complex topographic projection pattern. [Redrawn and modified from the original source (Malmierca et al. 1998)]; color values in original converted to grayscale. The intermediate and ventral nuclei (B) are here combined (C: 1). The periodic distribution of characteristic frequency is noteworthy

afferent axonal plexus. Thus, central nucleus neurons are variable in size, packed densely, and form long rows between which are interposed fascicles of ascending fibers. One of the two main classes of neuron has a disc-shaped dendritic field 200 to 800  $\mu\text{m}$  long and only 50 to 70  $\mu\text{m}$  wide; the long processes are oriented parallel to the axis of the afferent axons. Together, the preterminal afferent fibers and the polarized dendritic arbors of these neurons constitute fibrodendritic laminae that extend through the central nucleus. These resemble the laminated parts of the DCN and the sheetlike dendritic configurations in the MSO and the laminae in the ventral division of the MGB. This arrangement may be a common device for frequency-specific representation and segregation elsewhere in the central auditory system. Laminae are not necessarily uniform physiologically despite their narrow frequency range, as recordings along the primary tonotopic axis reveal systematic, small discontinuities that could correspond to critical bands (Schreiner and Langner 1997), and recordings within laminae reveal systematic shifts of several response properties, including onset latency (Langner et al. 1987) and response threshold (Stiebler and Ehret 1985). A task for future work is to understand more fully the range of computational possibilities within a frequency-band lamina, and the interactions between them. These neural ensembles have much in common with the laminae in the DCN, cell columns in the ventral nucleus of the lateral lemniscus, and frequency-specific columnar regions in auditory cortex. These are analogous to laminae in the lateral geniculate body and the ventrobasal complex (Jones 1985), and to the columnar organization of the sensory neocortex (Mountcastle 1997).

A second variety of central nucleus neuron, the stellate cell, has a spherical dendritic field and radiating processes that cross central nucleus laminae. Other neuronal subtypes vary in size but share features of dendritic architecture. The smallest such neurons have dendritic arbors  $<200$   $\mu\text{m}$  wide, and presumably intercept influences from narrower spatial domains than the large disc-shaped neurons, while some stellate cells span territories of approximately 500  $\mu\text{m}$ . Other stellate neurons have dendritic orientations at right angles to the fibrodendritic axis and could integrate afferent input across a wider range of the frequency spectrum (or other possible physiologic axes) than neurons with narrower dendritic domains. Central nucleus subdivisions exist that vary mainly on the basis of local differences in laminar organization and neuronal density.

---

Figure 1.11. *Continued*

(1, 2). **(D)** Aural classes and their distribution within the lateral lemniscal nuclei. The segregation of response properties on a nuclear basis in the mustache bat suggests that they serve different functional roles (Markovitz and Pollak 1993), that there may be species differences in physiologic properties (Markovitz and Pollak 1994), and at least one nucleus (e.g., dorsal nucleus) has connectional subdivisions (Yang et al. 1996). **(E)** GABAergic neurons have a differential distribution within the lateral lemniscal nuclei and sagulum. [**E**, **F**: The lateral lemniscal nuclei redrawn from the original source (Saint Marie et al. 1997); for the sagulum (Beneyto et al. 1998).] **(F)** Proportion of glycinergic neurons.




Figure 1.12. Physiologic and neurochemical aspects of IC organization. **(A)** Subdivisions of the IC. **(B)** Presumptive laminae (*dashed lines*) have a characteristic dorsomedial-to-ventrolateral orientation that, in the central nucleus, recapitulates the distribution of characteristic frequency (Rose et al. 1963) and the arrangement of the fibrodendritic laminae (Rockel and Jones 1973a). There is a substantial low-frequency representation; the precise disposition of frequency in the dorsal cortex is uncertain as it has a laminated structure (Morest and Oliver 1984), and the orientation of the cytoarchitecturally defined layers may not be in register with that of the isofrequency laminae. More than one map of characteristic frequency may occur as the discharge properties of dorsal cortex neurons

Intracellular filling studies reveal that the axons of most central nucleus cell types have extensive local branches, thus resembling interneurons elsewhere in the brain, while their collaterals often contribute to tectofugal projections. The local axonal network may run parallel to, or cross, the fibrodendritic laminae. A smaller subset of cells has only local connections within the central nucleus and could function as short axon Golgi type II cells (Oliver et al. 1991).

Although inhibitory neurons can enhance tonotopic organization, the laminated structure of the central nucleus imposes a physical arrangement that does not necessarily require the participation of interneurons in that process. If interneuronal effects are important in tonotopic organization, then their unique contribution cannot be discerned without more intracellular recordings (Nelson and Erulkar 1963; Oliver et al. 1991). This underscores an important difference between the central nucleus of the IC and the anterior part of the AVCN: in the latter, there are few or no GABAergic inhibitory interneurons (Adams and Mugnaini 1987) and the tuning curves of bushy cells are as, or more sharply, tuned than those in the central nucleus (Bourk et al. 1981), a process that would seem to depend on operations of the auditory nerve alone (Kiang et al. 1965). A related point is that there appears to be more than one way to organize topographic input (Kaas 1997).

There is evidence for other forms of organization in the central nucleus besides tonotopy. The representations of monaural inputs seem to vary spatially with regard to onset latency, receptive field bandwidth, and response threshold (Stiebler and Ehret 1985; Langner et al. 1987; Schreiner and Langner 1988). The binaural neurons represent several physiologic subclasses (Irvine 1986). In

---

Figure 1.12. *Continued*

differ from those in the central and lateral nuclei (Aitkin et al. 1994). [Redrawn and modified from the original source (Merzenich and Reid 1974 with permission from Elsevier Science).] (C) The percentage of aural subtypes in the central nucleus of the IC, showing the dominance of contralateral excitatory-occluded/monaural (*EO/mon*) and excitatory/inhibitory (*EI*) neurons. This is congruent with the notion that there is a spatial segregation of response types within the mustache bat central nucleus (Wenstrup et al. 1986). Whether these patterns are preserved in the MGB is unknown (see Fig. 1.15C). [Adapted and modified from the original source (Irvine 1986).] (D) Regional distribution and concentration of prospective neurotransmitters in IC subdivisions. In all subdivisions, glutamatergic neurons are numerous; whether tectothalamic and other projection neurons are glutamatergic requires further experimental study because of the structural role of glutamate/aspartate. Glycinergic axon terminals, but not neurons, are present in each part. [Interpreted from original observations (Oliver et al. 1994) for GABA; for glycine and glutamate/aspartate (Winer, unpublished observations); for somatostatin (Spangler and Morley 1987).] (E) Summary of comparisons of physiologic responses in IC subdivisions. The central nucleus is the principal component of the lemniscal pathway; the lateral nucleus integrates auditory and somatic sensory information, and too little is known of the details of dorsal cortex functional organization to reach firm conclusions. [Adapted and modified from the original sources (Aitkin 1986; Aitkin et al. 1994).]




Figure 1.13. Some pathways between the cochlear nucleus, the IC, and the MGB. **(A)** Schematic view of cochlear nucleus projections to the central nucleus. [Adapted and modified from the original source (Irvine 1986).] **(B)** The contralateral input is strongest. The ascending convergence to the IC (Fig. 1.14A) arises from all major parts of the cochlear nucleus. 1: Dorsal cochlear nucleus (DCN) and posteroventral cochlear nucleus (PVCN) projections reach the central nucleus. [Interpreted from original sources (Warr 1969; Oliver 1984).] 2: The anteroventral cochlear nucleus (AVCN) projects topographically to the central nucleus laminae. [Adapted from the original source (Oliver 1987).]



the mustache bat, populations of binaural and monaural neurons appear to be segregated regionally within the central nucleus, forming vertical columns 250  $\mu\text{m}$  long with common binaural properties and confined to an isofrequency region (Wenstrup et al. 1985). This suggests that, although central nucleus neurons are ordered across the entire frequency spectrum, other internal representations may also be present.

A new kind of physiologic organization arises in the auditory midbrain that enlarges its functional role and distinguishes it from medullary auditory nuclei. This pertains to the lateral nucleus and the dorsal cortex, structures whose role in inferior collicular processing is not well understood. The lateral nucleus receives input primarily from the medullary dorsal column nuclei and the trigeminal system in the pons (Aitkin et al. 1981), which represent the somatic sensory system for the body and head, respectively. Other projections come from the auditory cortex (Winer et al. 1998); there are no or very few axons from the lateral lemniscus (Rockel and Jones 1973b). What auditory influence it receives may arise principally from intracollicular connections (Aitkin and Phillips 1984a). In any case, the lateral nucleus has no tonotopic map, its neurons respond with long latencies compared to those of central nucleus cells, and they have very broad tuning curves. In the lateral nucleus of guinea pigs, contralateral azimuth position appears to be represented systematically, suggesting a role for this station in sound localization (Binns et al. 1992). Moreover, there is a coarse map of the body surface that overlaps with the auditory representation (Aitkin et al. 1978). A plausible function for the lateral nucleus might be to represent the position of the body with respect to sound sources in extrapersonal space. In the barn owl auditory midbrain (the lateral mesencephalic nucleus), for example, there are independent representations of the basilar membrane and a map of auditory space (Knudsen and Konishi 1978). These maps may be independent in the owl because prey acquisition involves rapid spatial processing and nearly ballistic motor estimates of the precise position of acoustic signals (Konishi 1973a,b). Such computations may be best performed independently by specific nuclei. In the cat, in contrast, the maps may not require the precision and independence that are manifest in the owl, perhaps behaving as combinatorial gates that require independent signals to converge to elicit a response.

---

Figure 1.13. *Continued*

[3: Adapted and modified from the original sources for intrinsic ipsilateral and commissural projections (Aitkin and Phillips 1984a); for tectothalamic connections (Calford and Aitkin 1983).] These circuits link parts of the IC that receive differential degrees of brain stem and cortical input. Some projections are reciprocal (e.g., commissural), others asymmetric (e.g., between the lateral and central nuclei). This could permit cortical access to nuclei that receive indirect cortical input via polysynaptic pathways in the mustache bat (Zhang et al. 1997; Jen et al. 1998). 4: The dorsal cortex of the IC receives little ascending auditory input (Irvine 1986) and a substantial projection from many areal origins (Fig. 1.18G) of cerebral cortex (Winer et al. 1998). The mustache bat tectothalamic pathway likely includes GABAergic and nonGABAergic projection neurons (Winer et al. 1996).




Figure 1.14. Some connections between the olivary complex, the IC, and the MGB. (A) Schematic of some ascending input to the central nucleus of the IC; only a few of the many projections are shown. [Adapted and modified from the original source (Irvine 1986).] 1: It is unknown whether the projections from the lateral superior olive (*LSO*) and medial superior olive (*MSO*) are segregated onto single postsynaptic cells in the central nucleus; connectational studies suggest considerable spatial overlap in their terminal targets (Glendenning and Masterton 1983; Oliver 1987; Shneiderman and Henkel 1987). 2: The IC projection to the MGB is bilateral from the central nucleus (Andersen et al. 1980; Calford and Aitkin 1983. Copyright by the Society for Neuroscience.). Descending pathways from the central nucleus to the DCN have been omitted (Conlee and Kane 1982).

Broad tuning is not confined to the lateral nucleus and it even occurs among some neurons in the posteroventral cochlear nucleus (PVCN) (Rhode and Greenberg 1992). Moreover, the DCN has a somatic sensory representation (Young et al. 1995). This suggests that even at the earliest stages of central neural processing, broad tuning and noncochlear input are both present, although the dominant functional influence in both cases remains auditory. In the lateral nucleus, however, a new principle emerges: auditory function is only one of many and any laminar organization differs fundamentally from that in the central nucleus (Morest and Oliver 1984; Oliver and Morest 1984). The emerging picture of lateral nucleus function suggests a multisensory integrative role in acoustico-motor behavior.

The dorsal cortex resembles the lateral nucleus as its neurons are tuned broadly, and they strongly prefer vocal stimuli to noise (Aitkin et al. 1994). Although isofrequency laminae seem to be absent, the neurons do form layers (Aitkin 1986), suggesting that further levels of functional organization may be present. The layers appear to have a closer relationship to corticocollicular, commissural, and intrinsic projections than to specific maps (Winer et al. 1998). Thus, auditory projections to the dorsal cortex are limited exclusively to indirect, polysynaptic auditory input from intrinsic and commissural midbrain neurons (Aitkin and Phillips 1984a) and the auditory cortex (Winer et al. 1998). Behavioral data, although limited, suggest that lesions that include the central nucleus lead to deficits in sound localization in the contralateral hemifield (Jenkins and Masterton 1982), while damage to the dorsal cortex affects attention and vigilance more severely than auditory discrimination behavior (Jane et al. 1965). Other dorsal cortex projections terminate in the paralemniscal and perireticular regions medial to the lateral lemniscus in the tegmental midbrain. Stimulation here can elicit a variety of vocalizations in which the cortex may participate (Schuller and Radtke-Schuller 1990). Neurons here have also been implicated in the startle reflex (Aitkin 1986). Substantial cortical input also ends in the nearby central gray, a center concerned with ascending and descending modulation of nociceptive influences and in vocalization related to emotive and visceral aspects of communication (Aitkin 1986; Winer et al. 1998).

A final example illustrates the breadth of functional arrangements that the IC mediates. In genetically epilepsy-prone rats, somatic motor seizures can be induced by acoustic stimulation (Faingold et al. 1989). GABAergic neurons, which are plentiful in the IC (Oliver et al. 1994), may participate in seizure genesis as blocking GABA uptake reduces seizure development (Faingold et al. 1994). This demonstrates the manifold roles of inhibitory neurons and the complex behaviors mediated by the IC and adjoining tegmental and reticular parts of the midbrain, which range from exclusively auditory processing (Aitkin 1986) to possible participation in vocalization (Suga and Yajima 1988) or the startle reflex (Yeomans and Frankland 1996). These properties are specific to the auditory midbrain and emphasize its pivotal role in auditory processing and the breadth of its functional impact.

## 6. THE MEDIAL GENICULATE BODY

The mammalian thalamus (Gr., *inner chamber*) consists of two oval masses that extend from the midbrain to the forebrain. The thalamus contains different nuclei that have been classified as sensory, motor, association, or intralaminar on the basis of their functions, connections, cell structure, and evolutionary history (Jones 1985). Sensory nuclei represent one modality, such as the MGB for hearing and the lateral geniculate body (LGB) for vision. Motor nuclei receive input from the cerebellum (and other structures) and participate in preparation for movement and ongoing motor behavior. Association nuclei are tied less securely to one modality, their internal maps may not be as fine-grained or as topographic as those of sensory nuclei, and their role is probably integrative. The intralaminar nuclei are conserved phylogenetically, they receive input from several modalities, and project broadly on the cerebral cortex and to subcortical nuclei; they may have a role in mediating shifts in behavioral state, for example, between sleeping and waking. Corresponding thalamic nuclei in different modalities are not necessarily equivalent functionally and there are significant species differences in size and internal circuitry. Thus, destroying one LGB in rodents causes a severe visual defect (Thompson 1969), whereas bilateral destruction of the medial geniculate body does not materially impair sound localization (Kelly and Judge 1985). In contrast, even unilateral damage in cats has serious consequences for auditory localization (Jenkins and Masterton 1982). Such nonequivalence between systems and across species reflects the concentration of thalamic visual function and a broader distribution of auditory function across brain stem nuclei.

The MGB differs from the IC in several ways (Winer 1991). First, its projections are almost entirely ascending and ipsilateral. Second, the inputs target not only auditory cortex but also subcortical limbic forebrain structures such as the amygdala and caudate nucleus; thus, the role of the MGB may be more than only auditory. Third, the MGB has a wide range of species-specific variability: it is relatively small in reptiles and amphibia, even when the midbrain homologue of the IC is well developed, and massive in carnivores. Fourth, MGB internal circuitry, especially the number of Golgi type II/GABAergic neurons, differs across species, suggesting that some forms of processing might be species-specific. Fifth, the strength of corticofugal input to the MGB is far greater than that of the corresponding corticocollicular projection. This implies a tighter coupling between thalamic and cortical oscillatory behavior than between the cortex and the auditory midbrain. Sixth, there are no internal connections linking MGB divisions, which accentuates their independence, while these pathways are plentiful in the IC and cochlear nucleus.

The main sensory nucleus of the MGB is the ventral division (Fig. 1.15), whose bushy tufted principal cells are arranged in fibrodendritic laminae (Morest 1965; Cetas et al. 2001) across which there is a systematic representation of characteristic frequency (Imig and Morel 1984) (Fig. 1.15B). Neurons with diverse response profiles are present, including specific binaural properties (Clarey

et al. 1992). The primary input to the ventral division arises from the central nucleus of the IC (Calford and Aitkin 1983) (Fig. 1.16B: 1, 2) and from the auditory cortex (Diamond et al. 1969; Winer et al. 2001). There are many parallels between the central nucleus of the IC (Aitkin 1986) and the ventral division (Winer 1992), including its laminated cytologic and orderly physiologic organizations. An important difference is the degree of cortical input, which is massive in the thalamus and sparser in the central nucleus (Winer et al. 1998). A second difference relates to intrinsic organization. In the ventral division there are Golgi type II cells whose axons are confined entirely to the thalamus and that are GABAergic (Rouiller et al. 1990), while in the IC many of the GABAergic neurons—especially the largest—project to the MGB (Winer et al. 1996). Perhaps there are no or few interneurons in the IC (Oliver et al. 1994), while all ventral division GABAergic neurons are type II cells as none have projections which leave the thalamus (Huang et al. 1999). The ventral division has a specialized synaptic arrangement, the glomerulus, in which dendrites and axons from Golgi type II cells are presynaptic to bushy tufted cell dendrites and ascending afferents, and this ensemble is surrounded by a glial sheath that isolates it from neighboring influences (Morest 1975). This synaptic pattern does not seem to occur in the IC (Rockel and Jones 1973c). The function of the glomeruli is not known (Morest 1971) but they may be a module for local intrathalamic processing, as they are present in all the corresponding sensory thalamic nuclei in all modalities (Rakic 1975).

Finally, there are species-specific differences in the proportion of GABAergic cells, ranging from <1% in some bats (Vater et al. 1992) to >25% in cat (Huang et al. 1999) and monkey (Smith et al. 1987). Such differences are far more marked in the thalamus than in the auditory midbrain (Oliver et al. 1994) or cerebral cortex (Prieto et al. 1994a), where these neurons represent 20% to 25% of the populations. The evolution of thalamic GABAergic circuitry may be relatively recent and system specific, as in other thalamic nuclei there is much more homogeneity with regard to Golgi type II cells (Winer and Larue 1996).

Ventral division response properties have much in common with those in the central nucleus of the IC, the main input source. Consequently, no clear hypothesis has emerged regarding the nature and purpose of functional transformation at this processing stage. One noticeable difference between the central nucleus of the IC and the MGBv is an increased local diversity of response properties, including best frequency (Morel et al. 1987). Similar to the central nucleus, local gradients can be seen for a variety of response properties. Gradients in the rostrocaudal axis include temporal response pattern, the ability to synchronize to repetitive stimuli, responsiveness to broad-band stimuli, and the monotonicity of rate-level functions (Rodrigues-Dagaeff et al. 1989). Binaural response classes also show some differences along the rostrocaudal domain and are characterized by clustering in the dorsoventral domain (Middlebrooks and Zook 1983; Rodrigues-Dagaeff et al. 1989).

The dorsal division receives its primary brain stem input from the dorsal cortex of the IC and from the lateral tegmental area, near the IC (Calford and




Figure 1.15. Basic features of medial geniculate body organization. **(A)** Subdivisions of the medial geniculate complex include the ventral division, which is present in all mammals and essential for normal hearing, and the dorsal and medial divisions, which are far more variable in size and have roles in auditory integration and polysensory processing, respectively. Not shown are the caudal dorsal and rostral dorsal nuclei or the rostral pole nucleus; the latter may be an extension of the ventral division (Imig and Morel 1985a,b). Some prior accounts considered the ventral and dorsal divisions collectively as the principal division (Rose and Woolsey 1958; Rinvik et al. 1987). **(B)** The schematic organization of unit best frequency in the ventral division (Imig and Morel 1985b) is in good accord with the configuration of anatomical fibrodendritic laminae (Morest 1965). **(C)** Orthogonal to the fibrodendritic laminae and interdigitated with them may be various binaural subregions. [Adapted and modified from the original source

Aitkin 1983). Like the cells in the dorsal cortex of the IC, dorsal division neurons have wide tuning curves, long latencies, and an uncertain tonotopic arrangement. They also have extended, oscillating slow-wave activity whose origins and functions are unknown but that is specific to this division (Aitkin and Dunlop 1968, 1969). Dorsal division projections to auditory cortex differ in two ways from those of the ventral division neurons. First, their targets include many more areas—up to nine—while the ventral division projects to only five areas (Huang and Winer 2000). Second, the cortical targets of the dorsal division are heterogeneous functionally and widely dispersed spatially (Shinonaga et al. 1994), while ventral division projections end in areas with similar auditory properties (Reale and Imig 1980) and which adjoin. Dorsal division function(s) appear independent of those in the tonotopic system; perhaps these nuclei affect attentional processes or global thalamic control of cortical excitability (Winer and Morest 1983). Responsiveness to structured signals, such as vocalizations (Shipley et al. 1988), may be more prominent in the dorsal division than responses to simpler stimuli (Buchwald et al. 1988). Damage to the cortical targets of the dorsal division impairs the perception of temporal patterns (Kelly 1973; Colavita et al. 1974).

The medial division has few of the features described here as characteristic of an auditory nucleus. It has a coarse tonotopic organization (more properly, a gradient) (Rouiller et al. 1989), and its afferent input includes nonauditory sources such as the vestibular system (Blum et al. 1979) as well as fibers from the lateral nucleus of the IC (Calford and Aitkin 1983). The medial division has the widest spectrum of targets in the MGB: it projects to every part of auditory cortex (Niimi and Matsuoka 1979), to nonauditory cortex (Jones and Powell 1973), and to the amygdala (Shinonaga et al. 1994). The thalamoamygdaloid pathway is unique to the auditory system (LeDoux et al. 1985), and it is essential for autonomic learning based on auditory cues (LeDoux et al. 1986). Because medial division neurons can show long-term potentiation (Gerren and Weinberger 1983), they may play a role in higher functions. A final difference with regard to the medial division is that its projections terminate mainly in layers I

---

Figure 1.15. *Continued*

(Middlebrooks and Zook 1983).] For locus of this arrangement, see (B), box. (D) Neurochemical profile of the auditory thalamus. GABAergic neurons represent about 26% of the cells on average, ranging from 33% in the ventral division to 18% in the medial division (Huang et al. 1999). Choline acetyltransferase-positive axon terminals are concentrated in the rat ventral division (Levey et al. 1987). Enkephalinergic neurons are found mainly in the dorsal division (Coveñas et al. 1986) and glutamatergic and/or aspartatergic cells are abundant in all divisions in the rat (Winer 1991). The chemical heterogeneity of the auditory thalamus does not include glycinergic neurons and axon terminals, which are concentrated in the caudal brain stem. (E) Summary of the attributes of MGB divisions. [Adapted and modified from the original source (Calford 1983).]




Figure 1.16. Some ascending projections of the auditory midbrain and thalamus. **(A)** A schematic picture of midbrain input to the MGB; the ipsilateral projection is dominant, and there are smaller inputs from the sagulum and central gray whose significance is obscure. Some auditory thalamic subdivisions such as the caudal dorsal nucleus receive little ascending auditory input and receive projections chiefly from the cerebral cortex (not shown) or from midbrain nuclei (such as the sagulum) whose function is uncertain. **(B)** The tectothalamocortical pathway. 1: Neurons in the central nucleus (CN) of the IC project topographically to the ventral division of the MGB (Kudo and Niimi 1980). 2: Ventral division neurons project ipsilaterally to auditory cortex subdivisions. 3: All four primary auditory cortical regions (areas AI, AAF, P, and Ve) receive topographic input (Morel and Imig 1987); each area has a unique redistributive pattern of corticocortical connectivity (Fig. 1.17E). [Modified from data in Calford and Aitkin 1983; Parts 2 and 3 from Niimi and Matsuoka 1979.]



and VI, whereas those from the ventral division end largely in layers III and IV (Huang and Winer 2000). Information reaching the cortex from the ventral and medial divisions is segregated and may target different postsynaptic populations, as the neuronal composition and output of the target layers are so varied.

Overall, the MGB is characterized by several parallel projection systems that, at this stage, appear less distinguishable by their auditory task specificity than by nonsensory influences. Strong modulatory influences from cortical feedback and many extracortical sources have been described in all sensory nuclei of the thalamus (Sherman and Guillery 2000). Accordingly, thalamic function may be less related to local receptive field transformation or implementation of feature extraction than to state-dependent modulation and gating of activity forwarded to the cortex (Miller and Schreiner 2000; Winer et al. 2001).

## 7. THE AUDITORY CORTEX

The cerebral cortex (La., *bark*) is a structure whose internal order, stereotyped neuronal organization, and high degree of local microcircuitry are largely conserved from insectivore to primate, despite vast differences in scale (Winer 1992). Its dynamic ability to modify itself through learning and memory suggests that static schemes to subdivide it may be inherently too limited to capture its functional complexity (Weinberg 1997). In the human cortex, for example, from about 50 to several hundred areas have been recognized (von Economo and Koskinas 1925; Brazier 1968). If it were smaller or less critical in behavior, these differences would reflect largely semantic distinctions with little conceptual import. But the human cerebral cortex is the largest organ in the brain: dissected and smoothed, it would be almost 1 m<sup>2</sup> in area and 3600  $\mu\text{m}$  thick with perhaps  $10^5$  neurons/mm<sup>2</sup>. If the human MGB has some  $5 \times 10^5$  neurons, the auditory cortex has millions more, irrespective of how its boundaries are defined, and the functional attributes of most parts remain unknown because of technical limitations. Moreover, because the cortex is interconnected extensively with itself in recurrent and feedforward networks (White and Keller 1989), the number of prospective neuronal and synaptic interactions assumes impressive proportions relative to the smaller number of permissible connections in the thalamus.

A second feature—laminar organization—distinguishes the cortex from most subcortical centers and adds another dimension to areal classification. Although the types of neuron are presumptively similar from area to area, they are largely segregated on a laminar basis. Horizontal cells, for example, are found only in neocortical layers I and VI and mediate lateral intracortical connectivity, whereas the largest pyramidal cells that may be critical for initiating rapid changes in subcortical targets are exclusively in layer V (Kelly and Wong 1981). This suggests that, within an area, layer-specific processes occur that resemble those in subdivisions of the cochlear nucleus, IC, or MGB. In a sense, the layers then correspond to nuclei in the brain stem, while areas subserve global functions.

The primary auditory cortex (AI) is the best studied of the tonotopic areas

(Brugge and Reale 1985; Aitkin 1990). Like all of the neocortex, it has six layers, with layers III and IV receiving axons arising from cells in the ventral division of the MGB, while layers I and VI are targets of the medial division (Huang and Winer 2000). The layers are linked vertically by interneurons whose axons have intralaminar projections, for example, from layer VI to layer IV (Prieto and Winer 1999) or by specific populations of pyramidal cells with intrinsic connections (Ojima et al. 1992). These pathways are analogous to the layer-specific connections in the DCN between inhibitory interneurons and fusiform cells. Although there are extensive ipsilateral corticocortical connections between areas with similar properties (Imig and Reale 1980) and robust interhemispheric commissural networks (Code and Winer 1985), perhaps >75% of the synapses (LeVay and Gilbert 1976) arise from an immense web of local connections (Matsubara and Phillips 1988; Read et al. 2001). The rules governing these interconnection are known best in the visual cortex, where vertical and horizontal modules are linked in intricate spatiotemporal patterns (Fitzpatrick et al. 1985; Hübener et al. 1997). Thalamic information in AI reaches layer I both directly (via thalamocortical projections) (Mitani et al. 1984) and along short axon polysynaptic intracortical intralaminar connections (Mitani and Shimokouchi 1985). These intrinsic vertical inputs and the monosynaptic thalamocortical projection of ventral division neurons to layer IV (Huang and Winer 2000) contact the long apical dendrites of many cells with perikarya in layers V and VI. Layer V and VI cells give rise to the corticofugal projections to the IC (Kelly and Wong 1981; Winer et al. 1998), thalamus (Winer et al. 2001), olivary complex (Doucet et al. 2002), striatum (Beneyto and Prieto 2001), amygdala (Romanski and LeDoux 1993), and cochlear nucleus (Weedman and Ryugo 1996a), to name just some of their targets (Wong 1991). Thus, thalamic projections to the cortex reach corticofugal neurons with different targets and influence prospectively almost every subcortical auditory nucleus. The complexity of the neural networks involved in this brief summary emphasizes the difficulty of explaining cortical function, as some connections seem to act in a highly redistributive way, whereas others are far more specific. Tracing these patterns of connectivity to the next level is a formidable task (Briggs and Callaway 2001).

A useful way to examine auditory cortex function is to enumerate and assess its physiologic representations. A primary auditory field with a tonotopic representation was found in every species studied (Luethke et al. 1988). In cat primary auditory cortex the gradient of tonotopic organization seen in barbiturate-anesthetized animals is for the most part quite strict (Merzenich et al. 1975) (Fig. 1.17C) but it is perhaps less ordered in awake preparations (Evans and Whitfield 1964; Evans et al. 1965) and at its dorsal and ventral margins (Schreiner and Sutter 1992). Across the tonotopic sequence is arrayed a series of orthogonal bands representing binaural subclasses (Imig and Adrián 1977; Kitzes et al. 1980; Middlebrooks et al. 1980) (Fig. 1.17C: EE, EI). A complete picture of the functional design would include other axes of organization interleaved among or embedded within the tonotopic and binaural axes (Ehret 1997). Evidence for amplitopic organization (Schreiner et al. 1992; Phillips et al. 1995;

Schreiner 1995) as well as representations of sharpness of tuning (Schreiner and Mendelson 1990; Heil et al. 1994; Schreiner 1995) or a systematic distribution of responses to frequency modulated tones (Mendelson et al. 1993; Eggermont 1998) have been seen in primary auditory cortex; these are reminiscent of the multiple, overlapping maps in primary visual cortex (Tusa et al. 1981). Cortical cells generally do not represent amplitude modulation as precisely as thalamic neurons (Creutzfeldt et al. 1980; Miller et al. 2001), and their response areas have a range of sizes and shapes (Phillips et al. 1985). Some subcortical aspects of spectral and temporal processing are further refined in the cortex to extract new information (Miller et al. 2001). The proportion of neurons sensitive to interaural level or phase or with azimuthal sensitivity is equal to or exceeds that in the MGB (Clarey et al. 1992). Many cortical cells respond selectively to the spatial location of tone bursts, with hemifield units among the most numerous (Middlebrooks and Pettigrew 1981). Inhibitory processes profoundly shape their receptive fields (Hendry and Jones 1991; Phillips et al. 1991; Prieto et al. 1994a,b) (Fig. 1.17F). More than 10 auditory cortex subdivisions are known in the cat (Winer, 1992); in primates, whose auditory cortical representation has been explored less fully, almost as many areas have been described (Merzenich and Brugge 1973; Morel and Kaas 1992; Morel et al. 1993; Kaas and Hackett 2000; Cheung et al. 2001). It remains for future work to understand how areas differ. These data will have important consequences for competing paradigms of cortical function. For example, the serial processing model envisages the cortex as a multistage decoding device whereas parallel processing schemes emphasize largely independent processing streams that are conveyed concurrently to different postsynaptic targets (Cowey 1981; Diamond 1983; Rouiller et al. 1991). Although elements of both models are present in auditory cortex (Romanski et al. 1999; Rauschecker and Tian 2000), more explicit tests of the differential predictions of each model will require a more complete picture of cortical operations than is now available.

A central issue is how the cortex affects subcortical centers. Stimulating or inactivating bat auditory cortex strongly affects the delay tuning of IC neurons, reducing the auditory-evoked response to subsequent afferent volleys by 90% (Yan and Suga 1996) and significantly altering the tuning properties of postsynaptic thalamic and collicular neurons (Zhang et al. 1997). Perhaps cortex exerts executive control on subcortical neurons for critical processing in demanding acoustic environments. Such corticofugal control is not limited to primary areas. Indeed, studies of these projections from nonprimary, nontopographic areas show that they are as strong, focal, and topographic, as that of their primary, tonotopic counterparts (Winer et al. 1998, 1999, 2001). Contemporary models of sensory processing emphasize biological and perceptual roles for the corticofugal systems commensurate with their size (Deschênes et al. 1998; Przybylski 1998) (Fig. 1.18).

The notion that topical representations in the cortex are rigid and invariant has recently been challenged in two complementary ways. Exposure to specific sounds in a training paradigm alters the distribution of cortical characteristic




Figure 1.17. Some aspects of auditory cortical organization. **(A)** Areal borders in a lateral view as derived from physiologic (Imig and Reale 1980) and architectonic (Rose 1949; Wallace et al. 1991) investigations. The primary fields are AI, AAF, P, Ve, and VP. Each has a tonotopic organization; the neurons respond mainly to auditory stimuli and they receive topographic projections from the auditory thalamus. Secondary areas are periauditory, include AII (Schreiner and Cynader 1984) and SF (He et al. 1997), and probably are not essential for sound localization (Diamond et al. 1962; Neff et al. 1975). The tertiary areas include limbic-related territories, Ins and Te, near the pseudosylvian sulcus, respond to auditory and nonauditory stimuli (Sindberg and Thompson 1962), and have complex ascending connections with thalamic association nuclei (Niimi and Matsuoka 1979; Winer and Morest 1983). The other fields in the posterior ectosylvian gyrus (*EP*)

frequency (Recanzone et al. 1993), and even relatively brief experience in a ventriloquism task shifts perceptual estimates of sound spatial localization (Recanzone 1998). This implies a rapid retuning of functional domains. A second approach used a stimulation paradigm involving subcortical nonauditory interactions between the cortex and the nucleus basalis, a forebrain limbic center that contains populations of cholinergic and GABAergic neurons; the former project widely to neocortex and elsewhere (Jones et al. 1976). Nucleus basalis stimulation, when paired with appropriate training, has robust and temporally enduring effects on gross tonotopy, single-neuron tuning, receptive field structure, and

---

Figure 1.17. *Continued*

have relationships with periauditory-perivisual thalamic nuclei (Bowman and Olson 1988). **(B)** Major sulci in cat auditory cortical areas (Kawamura 1971). **(C)** Selected features of AI functional organization from physiologic studies. Locus of observations is shown in **(B)**. Binaural representation (Middlebrooks et al. 1980) includes alternating bands (*ovals*) superimposed on and crossing the isofrequency contours (*dashed lines*) (Merzenich et al. 1975) throughout AI. Neurons with a preferred direction for frequency-modulated (*FM*) sweeps (Mendelson et al. 1993) also are localized across the isofrequency gradient (*arrows*). The dispersion of characteristic frequency values (CF) is larger in the ventral part of AI (*small black circles*) (Schreiner and Sutter 1992). Other representations are omitted for purposes of clarity. [Redrawn and modified from the original (EHret 1997).] **(D)** Intrinsic organization of a module in AI (see **B**: *D*). Units with narrowband (*NB*) and broadband (*BB*) frequency selectivities segregate spatially into multiple clusters across the surface of AI. Spatially segregated narrowband clusters are connected via long-range (>1 mm) intrinsic horizontal circuitry as shown with the retrograde labeling pattern (*white circles* denote somata; *lines* denote axons) which aligns with a prominent dorsal cluster of narrowband units (d1–d4). Quality factors (*Q*) at 40 dB above spike threshold were derived from unit responses to pure tone stimuli over a 5-octave range. *Light gray*: low  $Q_{40\text{dB}}$  (wide bandwidth); *white*: high  $Q_{40\text{dB}}$  (narrow bandwidth). The injection and retrograde labeling are largely confined to the white narrowband regions. *a–c*: Sublayers of layer V with differential corticofugal projections. [Adapted and modified from the original source (Read et al. 2002).] The distribution of thalamic laminar projections (right side, white numbers) from type 1 (ventral division) and type 2 (medial division) input is shown numerically as the percentage of boutons/layer after thalamic deposits of tracers. [Adapted and modified from the original source (Huang and Winer 2000).] **(E)** Some ipsilateral corticocortical connections between core, belt, and parabelt auditory areas. Each area receives thalamocortical, ipsilateral corticocortical, and commissural input (Winer 1992). *Arrows* are proportional to the strength of interareal connections. AI acts as a major redistributive hub since it receives projections from a few, mainly lemniscal thalamic nuclei and redistributes their output to other primary fields (*AAF*), to belt areas (*AII*), and to remote parabelt cortex (*EP*). In most instances strong feedback connections are present as well. [Adapted and modified from original sources (Kawamura 1973; Rouiller et al. 1991; Clarke et al. 1993).] **(F)** Concentration of putative neurotransmitters in AI, of which GABAergic neurons comprise about 20% of the population (Prieto et al. 1994a,b), and glutamatergic or aspartatergic cells represent the majority. The serotonergic innervation consists of axon terminals only (Mulligan and Törk 1988), many of which form perisomatic baskets onto pyramidal cells.




Figure 1.18. Some descending connections of auditory cortex. (A–E) Main targets of corticofugal auditory system. (A) Cortical auditory areas in the cat (Rose 1949; Imig and Reale 1980, 1981). (B) All subdivisions of the MGB receive auditory corticofugal projections (see Fig. 1.18G: MGB); most are topographic, even when they involve thalamic nuclei devoid of fine tonotopic organization (Diamond et al. 1969; Pontes et al. 1975). Few neurons project to both the MGB and the IC (Wong and Kelly 1981); this suggests that the different corticofugal projections have specific laminar origins in rat (Games and Winer 1988) and cat (Prieto and Winer 1999). (C) The complementary corticofugal projections to the IC (G: inferior colliculus) differ fundamentally from those to the MGB. Thus, input to the lemniscal part (the central nucleus) is sparse, whereas

topical representation in primary auditory cortex (Weinberger 1997; Kilgard and Merzenich 1998b; Kilgard et al. 2001; Kilgard and Merzenich 2002). Thus, even topographically secure areas can show physiologic lability under circumstances that might approximate normal subcortical–cortical interactions (Ohl and Scheich 1997).

In humans several auditory areas have been defined with physiologic (Celesia 1976; Howard et al. 2000), imaging (Belin et al. 2000), neurochemical (Hutsler and Gazzaniga, 1996), connectional (Tardif and Clarke 2001), and histochemical (Hackett et al. 2001) methods. These regions correspond, at least superficially, to their presumed homologues in primates (Brugge 1982). For example, chimpanzees exhibit planum temporale asymmetry (Gannon et al. 1998) much like that in humans (Geschwind and Levitsky 1968), and macaque species-specific vocalizations are impaired by a left, but not a right, hemisphere lesion in a way that recalls expressive aphasia seen after human cortical damage (Heffner and Heffner 1984).

The question of how auditory cortex output influences other parts of the brain can be examined from a neurologic perspective. Cortical disconnection syndromes from infarcts or other trauma leave some functions isolated but otherwise largely intact (Geschwind 1965). In primates the output of the auditory cortex and associated polysensory areas along the superior temporal gyrus reaches wide areas of the parietal lobe and prefrontal cortex (Jones and Powell 1973). Although questions remain about the identity of homologous areas related to language in humans and nonhuman primates (Aboitiz and García 1997), the linguistic deficits in humans consequent to destruction of Wernicke's area—impairments in repetition, comprehension, and naming—are consistent clinical features (Benson 1988) and leave little doubt that auditory cortex functions in-

---

Figure 1.18. *Continued*

the ventral division of the medial geniculate body receives a much larger projection (Winer et al. 1998). The other main midbrain targets—the intercollicular tegmentum and lateral nucleus—have roles related to spatial orientation to sound (Oliver and Huerta 1992). A third difference is that giant axon terminals of cortical origin are present in the MGB (Winer et al. 1999) and not in the IC (Diehl and Winer 1996). Cortical influence on nuclei in the ascending auditory system is nucleus-, and perhaps areally, specific. **(D)** The dorsal cochlear nucleus (*DCN*) receives a corticofugal projection. [Adapted and modified from the original sources (Weedman and Ryugo 1996a,b).] **(E)** The corpus striatum and associated nuclei (e.g., claustrum) each receive input from almost all subdivisions of auditory cortex **(G, lower part)**. [Adapted and modified from the original source (Beneyto and Prieto 2001).] **(F)** Other corticofugal projections reach the sagulum and pontine nuclei. [Adapted and modified from the original source (Brodal 1972).] **(G)** Summary of auditory corticofugal input to the MGB (Winer et al. 2001), IC (Winer et al. 1998), olivary region (Doucet et al. 2002), and cochlear nucleus (Weedman and Ryugo 1996a). The range in the strength, specificity, and origin of the input suggests that the cortex has many roles. 1, Medial dorsal nucleus; 2, lateral mesencephalic nucleus; 3, lateral geniculate body.

clude an essential prelinguistic role analogous to that of the somatic sensory cortex in planning motor behavior. Corticofugal projections thus permit cortical access to some of the earliest parts of the central synaptic sequence in hearing.

By the same token, the auditory cortex has massive feedforward connections that convey its output signals to other cortical and subcortical areas with a role in comprehending and initiating speech. The case can be made for a disjunction between the brain stem auditory system, which encodes the spatial and temporal and spectral qualities of sound, and the thalamus and forebrain, which use this information to derive higher-order representations, namely, biological significance, pattern recognition, and planning for action. The temporal aspects of biological signals contain abundant information that the cortex might extract (Phillips et al. 1991) and that cortical lesions surely impair (Albert and Bear 1974). The primal sketch derived by the brain stem imposes temporal structure on these premotor and cognitive aspects of hearing. The auditory cortex, in conjunction with the frontal lobes and speech sensory-motor cortex, is responsible for the fluidity, precision, and salience of speech in the context of language.

The role of the auditory system up to and including primary cortex can be viewed as to establish a stable representation of a multidimensional feature space that allows extraction of object form, attribute, and location from the sound background for a reconstruction of the environmental scene. Higher sensory cortical areas utilize these representations for the recognition of objects and the formation of reference-independent categories of objects, their relationship to environment and self, multisensory integration, assignment and evaluation of behavioral significance according to context, and the transition to action (Dinse and Schreiner 2002).

## 8. THEMES FOR DISCUSSION

1. Most auditory centers receive projections from other areas (convergence) and project to many others (divergence). This can entail multiple levels of function within a nucleus so that more than one processing task is superimposed on single neurons, or functions are allocated to ensembles of cells specialized for a specific computational role.
2. The diverse structure of auditory neurons and the rich array of auditory nuclei contrast with the relative homogeneity of peripheral input and the remarkably small number of receptors.
3. Parallel and serial forms of processing emerge as organizing principles at the level of the cochlear nucleus, and they prevail throughout the auditory system. Corticofugal feedback and feedforward connections violate any simple or unidirectional principles of serial organization.
4. Single auditory neurons participate in several representations. Their complete receptive field profile should include, but not be limited to, considerations of aurality, onset/offset behavior, participation in tonotopic or amplitude representations, and sharpness of tuning. This range is a chal-



- lenge to single neuron feature detection models of sensory processing. How many and which of these dimensions are required to classify neurons?
5. The auditory forebrain has two forms of inhibition: intrinsic processes are local to a nucleus, and extrinsic inhibition links remote nuclei/areas. The latter projections are robust in the auditory system, and virtually nonexistent in other modalities. The role that remote inhibition plays in auditory processing is not yet understood.
  6. The forebrain auditory system has powerful and direct connections with premotor neurons in the striatum and the pons, each of which is involved in ongoing movement. What is the impact of auditory information on these structures?
  7. One role of the auditory system might be as the anticipatory stage for the higher-order communication system, analogous to the function of the premotor cortex in primate movement.
  8. It may not be possible or even appropriate to delegate functions to nuclei, as functions are global constructs, while neurons and nuclei and circuits are restricted to local operations.
  9. The auditory “system” is composed of several subsystems (for example, olivocochlear, thalamocortical, etc.), each with a distinct connectivity, function, structure, and physiology. The apparent unity of these operations is an inference, not necessarily an implicit or emergent property of these neurons. The presence of so many nuclei is consistent with the view that function is widely dispersed yet highly localized.
  10. Some auditory nuclei are strongly conserved in phylogeny, while others are far more variable (Butler and Hodos 1996; Glendenning and Masterton 1998). This principle applies both to size and to internal circuitry. The cochlea is, arguably, the most variable of all the structures. The ancestral forms common to mammals must have diverged rapidly to occupy specific behavioral niches (Baird 1974).
  11. The descending auditory connections are among the largest and the most elaborate in any modality. They reach the earliest stage (cochlear nucleus) monosynaptically and increase progressively in size (corticofugal) until they are one of the largest projections in the brain. How do the actions of the several descending systems differ?
  12. Much of the central auditory system has no, or only a rudimentary, tonotopic organization. This principle seems to reach its apex in the forebrain, where only a part of the representation for hearing has maps like those so prevalent in the brain stem. This feature appears as early as the PVCN, where broadly tuned cells occur. Are these classes analogous to the two visual systems in mammals (Schneider 1969)?
  13. Comparatively few types of neuron are present in the auditory system. These are conserved across nuclei and analogous types are found in different species. Some neurons closely resemble their visual and somatic sensory counterparts, especially in the thalamus and the cortex. Other features of auditory system organization, such as the endbulbs and calyces of Held, are unique

to it and the vestibular system. What is the relationship between these types of neuron and the information they process?

14. The auditory system is the only sensory modality with two chemically distinct central neurotransmitters (GABA and glycine) that may serve an inhibitory/disinhibitory role. Although these transmitters occur in other pathways, they are nowhere as prominent (especially glycine) as they are in the auditory system. Is more than one form of inhibition operating in the auditory system?
15. GABA is present, in varying amounts, through the entire auditory pathway from cochlea to cortex. Glycine as a transmitter is limited to the locus between the cochlear nucleus and the inferior colliculus, and all glycinergic neurons are subcollicular. What is the significance of the nucleus-specific distribution of these and other putative transmitters?
16. The concept of a relay nucleus requires critical scrutiny as all central nuclei transform and modify the information that passes through them (Stanley et al. 1999). It remains to be seen how processing in the IC differs from that in the cerebral cortex (Langner 1992; Eggermont 1998, 2001).
17. The auditory system is the only modality with direct projections from the hippocampus. These could integrate working memory with prior auditory experience. Audition is the only sensory modality with significant direct limbic affiliations.
18. The MGB has monosynaptic projections to the amygdala in the limbic system. This permits polysensory parts of the auditory system to influence smooth muscle as amygdalohypothalamic pathways have significant impact on the autonomic nervous system. It suggests a special saliency for auditory input.
19. There are species differences in the size and internal organization of auditory nuclei, especially in the cochlear nucleus and superior olivary complex. Some of these differences extend to neural circuitry and may entail species-specific patterns of processing. This has implications for the evolution of this circuitry and for understanding species differences.
20. Considerations of tonotopy have dominated thinking about the functional organization of hearing at the expense of considering how other aspects of the stimulus or response patterns such as spontaneous rate, amplitude, latency, and others are organized globally.
21. The auditory system seems to contain three types of neurons as defined on connectional and neurochemical grounds. Two types are common throughout the nervous system: a large projection cell that uses glutamate or aspartate as a transmitter (type I), and a smaller local circuit neuron that uses GABA or glycine as their main transmitter (type II). The third type resembles the projection neuron in size, but uses GABA or glycine as a transmitter, and projects remotely (type III). Neurons with a type III profile are rare or nonexistent in other modalities. What role do they have in audition?
22. Only the auditory system has a robust centrifugal pathway that is conserved phylogenetically. In other modalities it is either nonexistent in all taxa (so-

matic sensory system) or present in a highly selective way in a few species only, such as avian isthmo optic projections to retinal ganglion cells (Cowan et al. 1961). A possible parallel is the projection from spinal  $\gamma$  motoneurons onto intrafusal fibers in the somatic motor system (Matthews 1972). Why are these centrifugal systems so prominent in hearing?

23. Localization theory suggests that functions reside in particular sites that are linked by connections. Computational theory emphasizes interactions between areas. Localization theory depends on maps but the nature of their interactions or the properties that underlie synaptic plasticity are largely unknown. Computation theory depends on operations that cannot readily be related to either neuronal connections or realistic behavior. There is evidence that neural coding properties are labile and experience dependent, but the nature of the code is a matter of debate (Eggermont 2001).
24. Biologically naturalistic stimuli can reveal facets of neural response profiles that more artificial stimuli cannot (Phillips 1989). Studies using pure tone stimuli in anesthetized preparations must be extended to include awake, behaving animals.
25. A general theory of communication should encompass the range of auditory behavior from cricket stridulation (Robert et al. 1992) to song learning in birds (Margoliash 1987) to language acquisition in humans (Nobre and Plunkett 1997). If a grand unified theory is impossible, then the common and unique features of the several communication systems should be identified.

## Acknowledgments

This research was supported by United States Public Health Service grants R01 DC02319-24 (J.A.W.) and DC02260-8 (C.E.S.). We thank Mr. David T. Larue and Ms. Tania J. Bettis for essential assistance in executing the figures. Dr. Steven Greenberg gave helpful advice and counsel.

## Abbreviations

AAF	anterior auditory field
AC	anterior commissure
aes	anterior ectosylvian sulcus
AI	primary auditory cortex
AII	second auditory cortical area
Am	amygdaloid nuclei
AMa	anterior amygdaloid nucleus
AnLc	ansa lenticularis
AS	dorsal and intermediate subdivisions of the acoustic striae
as	anterior suprasylvian sulcus

AVCN	anteroventral cochlear nucleus
BB	broadband
Ca	caudate nucleus
CC	caudal cortex of the inferior colliculus <i>or</i> corpus callosum
CF	characteristic frequency
CG	central gray
ChAt	choline acetyltransferase
CiC	commissure of the inferior colliculus
CiG	cingulate gyrus
Cl	claustrum
CN	central nucleus of the inferior colliculus
cNB	central narrowband region
Cu	cuneiform nucleus
$C_z^+$	positive interelectrode difference
$C_z^-$	negative interelectrode difference
D	dorsal division of the medial geniculate body <i>or</i> dorsal
DC	dorsal cortex of the inferior colliculus
DC I-IV	layers of the dorsal cortex of the inferior colliculus
DCa	caudal pole of the medial geniculate body
DCN	dorsal cochlear nucleus
DD	deep dorsal nucleus of the medial geniculate body
DLPO	dorsolateral periolivary nucleus
DMPO	dorsomedial periolivary nucleus
DNLL	dorsal nucleus of the lateral lemniscus
DS	dorsal superficial nucleus of the medial geniculate body
EE	excitatory/excitatory
EI	excitatory/inhibitory
EO/F	excitatory/occluded/facilitated
EO/mon	excitatory/occluded/monaural
EPD	posterior ectosylvian gyrus, dorsal part
EPI	posterior ectosylvian gyrus, intermediate part
EPV	posterior ectosylvian gyrus, ventral part
FM	frequency modulated
GABA	$\gamma$ -aminobutyric acid
IC	inferior colliculus
ICa	internal capsule
IcT	intercollicular tegmentum
INLL	intermediate nucleus of the lateral lemniscus
Ins	insular cortex
LD	lateral dorsal thalamic nucleus
LGB	lateral geniculate body
LLN	lateral lemniscal nuclei
LN	lateral nucleus of the inferior colliculus
LNTB	lateral nucleus of the trapezoid body
LP	lateral posterior nucleus

LSO	lateral superior olive
M	medial division of the medial geniculate body <i>or</i> medial
MCP	middle cerebellar peduncle
MGB	medial geniculate body
ML	medial lemniscus
MNTB	medial nucleus of the trapezoid body
MRF	mesencephalic reticular formation
MSO	medial superior olive
mss	middle suprasylvian sulcus
NB	narrowband
NRTP	reticular tegmental nucleus of the pons
OC	optic chiasm
OCA	octopus cell area
Ov	<i>pars ovoidea</i> of the ventral division of the medial geniculate body
P	posterior auditory field
pf	parallel fiber
pes	posterior ectosylvian sulcus
PL	posterior limitans nucleus
PN	pontine nuclei
Pol	lateral part of the posterior group
Pro	preoptic area
ps	pseudosylvian sulcus
pss	posterior suprasylvian sulcus
Pu	putamen
Pul	pulvinar nucleus
Py	pyramidal tract
PVCN	posteroventral cochlear nucleus
RP	rostral pole of the inferior colliculus
Sa	sagulum
SC	superior colliculus
SeN	septal nuclei
SF	suprasylvian fringe
Sgl	suprageniculate nucleus, lateral part
Sgm	suprageniculate nucleus, medial part
Spf	subparafascicular nucleus
SpN	suprapeduncular nucleus
Te	temporal cortex
V	ventral division of the medial geniculate body <i>or</i> trigeminal nerve <i>or</i> ventral
V <sub>dc</sub>	ventral division, dorsocaudal part
Ve	ventral auditory field
V <sub>l</sub> , V <sub>l</sub>	ventral division, lateral part
V <sub>o</sub>	ventral division, oral part
VMPO	ventromedial periolivary nucleus

VNLL	ventral nucleus of the lateral lemniscus
VP	ventral posterior auditory field
I–VI	components of auditory brain stem evoked potential <i>or</i> layers of auditory cortex

## References

- Aboitiz F and García VR (1997) The evolutionary origins of the language areas of the human brain. A neuroanatomical perspective. *Brain Research Reviews* **25**:381–396.
- Adams JC and Mugnaini E (1984) Dorsal nucleus of the lateral lemniscus: a nucleus of GABAergic projection neurons. *Brain Research Bulletin* **14**:585–590.
- Adams JC and Mugnaini E (1987) Patterns of glutamate decarboxylase immunostaining in the feline cochlear nucleus complex studied with silver enhancement and electron microscopy. *Journal of Comparative Neurology* **262**:375–401.
- Adams JC and Mugnaini E (1990) Immunocytochemical evidence for inhibitory and disinhibitory circuits in the superior olive. *Hearing Research* **49**:281–298.
- Aitkin L (1979) The auditory midbrain. *Trends in Neurosciences* **2**:308–310.
- Aitkin L (1986) *The Auditory Midbrain. Structure and Function in the Central Auditory Pathway*. Humana Press, Clifton.
- Aitkin LM (1989) The auditory system. In: Björklund A, Hökfelt T, and Swanson LW (eds). *Handbook of Chemical Neuroanatomy, Volume 7: Integrated Systems of the CNS, Part II: Central Visual, Auditory, Somatosensory, Gustatory*. Elsevier, Amsterdam, pp. 165–218.
- Aitkin L (1990) *The Auditory Cortex: Structural and Functional Bases of Auditory Perception*. Chapman and Hall, London.
- Aitkin LM and Dunlop CW (1968) Interplay of excitation and inhibition in the cat medial geniculate body. *Journal of Neurophysiology* **31**:44–61.
- Aitkin LM and Dunlop CW (1969) Inhibition in the medial geniculate body of the cat. *Experimental Brain Research* **7**:68–83.
- Aitkin LM and Moore DR (1975) Inferior colliculus, II. Development of tuning characteristics and tonotopic organization in central nucleus of the neonatal cat. *Journal of Neurophysiology* **38**:1208–1216.
- Aitkin LM and Phillips SC (1984a) The interconnections of the inferior colliculi through their commissure. *Journal of Comparative Neurology* **228**:210–216.
- Aitkin LM and Phillips SC (1984b) Is the inferior colliculus an obligatory relay in the cat auditory system? *Neuroscience Letters* **44**:259–264.
- Aitkin LM and Reynolds A (1975) Development of binaural responses in the kitten inferior colliculus. *Neuroscience Letters* **1**:315–319.
- Aitkin LM, Anderson DJ, and Brugge JF (1970) Tonotopic organization and discharge characteristics of single neurons in nuclei of the lateral lemniscus of the cat. *Journal of Neurophysiology* **33**:421–440.
- Aitkin LM, Dickhaus H, Schult W, and Zimmerman M (1978) External nucleus of the inferior colliculus: auditory and spinal somatosensory afferents and their interactions. *Journal of Neurophysiology* **41**:837–847.
- Aitkin LM, Kenyon CE, and Philpott P (1981) The representation of the auditory and somatosensory systems in the external nucleus of the cat inferior colliculus. *Journal of Comparative Neurology* **196**:25–40.

- Aitkin L, Tran L, and Syka J (1994) The responses of neurons in subdivisions of the inferior colliculus of cats to tonal, noise and vocal stimuli. *Experimental Brain Research* **98**:53–64.
- Albert ML and Bear D (1974) Time to understand. A case study of word deafness with reference to the role of time in auditory comprehension. *Brain* **97**:373–384.
- Altschuler RA, Neises GR, Harmison GG, Wenthold RJ, and Fex J (1981) Immunocytochemical localization of aspartate aminotransferase immunoreactivity in cochlear nucleus of the guinea pig. *Proceedings of the National Academy of Sciences of the United States of America* **78**:6553–6557.
- Andersen RA, Roth GL, Aitkin LM, and Merzenich MM (1980) The efferent projections of the central nucleus of the inferior colliculus in the cat. *Journal of Comparative Neurology* **194**:649–662.
- Ashmore JF (1991) The electrophysiology of hair cells. *Annual Review of Physiology* **53**:465–476.
- Baird IL (1974) Some aspects of the comparative anatomy and evolution of the inner ear of submammalian vertebrates. *Brain, Behavior and Evolution* **10**:11–36.
- Bajo VM, Merchán MA, López DE, and Rouiller EM (1993) Neuronal morphology and efferent projections of the dorsal nucleus of the lateral lemniscus in the rat. *Journal of Comparative Neurology* **334**:241–262.
- Beckius GE, Batra R, and Oliver DL (1999) Axons from the anteroventral cochlear nucleus that terminate in medial superior olive of cat: observations related to delay lines. *Journal of Neuroscience* **19**:3146–3161.
- Belin P, Zatorre RJ, Lafaille P, Pierre A, and Pike B (2000) Voice-selective areas in human auditory cortex. *Nature* **403**:309–312.
- Beneyto M and Prieto JJ (2001) Connections of the auditory cortex with the claustrum and endopiriform nucleus in the cat. *Brain Research Bulletin* **54**:485–498.
- Beneyto M, Winer JA, Larue DT, and Prieto JJ (1998) Auditory connections and neurochemistry of the sagulum. *Journal of Comparative Neurology* **401**:329–351.
- Benson DF (1988) Classical syndromes of aphasia. In: Goodglass H (ed). *Handbook of Neuropsychology*, Volume 1, Section 3: *Language and Aphasia*, Part 1. Elsevier, Amsterdam, pp. 267–280.
- Berman AL (1968) *The Brain Stem of the Cat: A Cytoarchitectonic Atlas with Stereotaxic Coordinates*. University of Wisconsin Press, Madison.
- Berberi AS and Mugnaini E (1991) Distribution and targets of the cartwheel cell axon in the dorsal cochlear nucleus of the guinea pig. *Anatomy and Embryology* **183**:427–454.
- Binns KE, Grant S, Withington DJ, and Keating MJ (1992) A topographic representation of auditory space in the external nucleus of the inferior colliculus of the guinea-pig. *Brain Research* **589**:231–242.
- Binns KE, Withington DJ, and Keating MJ (1995) The developmental emergence of the representation of auditory azimuth in the external nucleus of the inferior colliculus of the guinea-pig: the effects of visual and auditory deprivation. *Developmental Brain Research* **85**:14–24.
- Bjordahl TS, Dimyan MA, and Weinberger NM (1998) Induction of long-term receptive field plasticity in the auditory cortex of the waking guinea pig by stimulation of the nucleus basalis. *Behavioral Neuroscience* **112**:467–479.
- Bledsoe SC Jr, Snead CR, Helfert RH, Prasad V, Wenthold RJ, and Altschuler RA (1990) Immunocytochemical and lesion studies support the hypothesis that the projection from

- the medial nucleus of the trapezoid body to the lateral superior olive is glycinergic. *Brain Research* **517**:189–194.
- Blum PS, Abraham LD, and Gilman S (1979) Vestibular, auditory and somatic input to the posterior thalamus of the cat. *Experimental Brain Research* **34**:1–9.
- Bourk TR, Mielcarz JP, and Norris BE (1981) Tonotopic organization of the anteroventral cochlear nucleus of the cat. *Hearing Research* **4**:215–241.
- Bowman EM and Olson CR (1988) Visual and auditory association areas of the cat's posterior ectosylvian gyrus: thalamic afferents. *Journal of Comparative Neurology* **272**:15–29.
- Brawer JR, Morest DK, and Kane EC (1974) The neuronal architecture of the cochlear nucleus of the cat. *Journal of Comparative Neurology* **155**:251–300.
- Brazier MAB (1968) Architectonics of the cerebral cortex: research in the 19th century. In: Brazier MAB and Petsche H (eds). *Architectonics of the Cerebral Cortex*. Raven Press, New York, pp. 9–29.
- Briggs F and Callaway EM (2001) Layer-specific input to distinct cell types in layer 6 of monkey primary visual cortex. *Journal of Neuroscience* **21**:3600–3608.
- Brodal A (1981) *Neurological Anatomy in Relation to Clinical Medicine*. Oxford University Press, New York.
- Brodal P (1972) The corticopontine projection in the cat. The projection from the auditory cortex. *Archives Italiennes de Biologie* **110**:119–144.
- Brödel M (1946) *Three Unpublished Drawings of the Anatomy of the Human Ear*. W.B. Saunders, Philadelphia.
- Brown MC and Ledwith III JV (1990) Projections of thin (type-II) and thick (type-I) auditory-nerve fibers into the cochlear nucleus of the mouse. *Hearing Research* **49**:105–118.
- Brownell WE (1975) Organization of the cat trapezoid body and the discharge characteristics of its fibers. *Brain Research* **94**:413–433.
- Brückner S and Rübsamen R (1995) Binaural response characteristics in isofrequency sheets of the gerbil inferior colliculus. *Hearing Research* **86**:1–14.
- Brugge JF (1982) Auditory cortical areas in primates. In: Woolsey CN (ed). *Cortical Sensory Organization, Volume 3: Multiple Auditory Areas*. Humana Press, Clifton, pp. 59–70.
- Brugge JF and Reale RA (1985) Auditory cortex. In: Peters A and Jones EG (eds). *Cerebral Cortex, Volume 4: Association and Auditory Cortices*. Plenum Press, New York, pp. 229–271.
- Brunso-Bechtold JK, Thompson GC, and Masterton RB (1981) HRP study of the organization of auditory afferents ascending to central nucleus of inferior colliculus in cat. *Journal of Comparative Neurology* **197**:705–722.
- Buchwald J, Dickerson L, Harrison J, and Hinman C (1988) Medial geniculate body unit responses to cat cries. In: Syka J and Masterton RB (eds). *Auditory Pathway: Structure and Function*. Plenum Press, New York, pp. 319–322.
- Butler AB and Hodos W (1996) *Comparative Vertebrate Neuroanatomy: Evolution and Adaptation*. John Wiley & Sons, New York.
- Calford MB (1983) The parcellation of the medial geniculate body of the cat defined by the auditory response properties of single units. *Journal of Neuroscience* **3**:2350–2364.
- Calford MB and Aitkin LM (1983) Ascending projections to the medial geniculate body of the cat: evidence for multiple, parallel auditory pathways through the thalamus. *Journal of Neuroscience* **3**:2365–2380.



- Cant NB (1982) Identification of cell types in the anteroventral cochlear nucleus that project to the inferior colliculus. *Neuroscience Letters* **32**:241–246.
- Cant NB (1992) The cochlear nucleus: neuronal types and their synaptic organization. In: Webster DB, Popper AN, and Fay RR (eds). *Springer Handbook of Auditory Research*, Volume 1: *The Mammalian Auditory Pathways: Neuroanatomy*. Springer-Verlag, New York, pp. 66–116.
- Cant NB and Casseday JH (1986) Projections from the anteroventral cochlear nucleus to the lateral and medial superior olivary nuclei. *Journal of Comparative Neurology* **247**:457–476.
- Cant NB and Gaston KC (1982) Pathways connecting the right and left cochlear nuclei. *Journal of Comparative Neurology* **212**:313–326.
- Cant NB and Morest DK (1979) The bushy cells in the anteroventral nucleus of the cat: a study with the electron microscope. *Neuroscience* **4**:1925–1945.
- Carr CE and Friedman MA (1999) Evolution of time coding systems. *Neural Computation* **11**:1–20.
- Celesia GG (1976) Organization of auditory cortical areas in man. *Brain* **99**:403–414.
- Cetas JS, Price RO, Velenovsky DS, Sinex DG, and McMullen NT (2001) Frequency organization and cellular lamination in the medial geniculate body of the rabbit. *Hearing Research* **155**:113–123.
- Cheung SW, Bedenbaugh PH, Nagarajan SS, and Schreiner CE (2001) Functional organization of squirrel monkey primary auditory cortex: responses to pure tones. *Journal of Neurophysiology* **85**:1732–1749.
- Chusid JG (1982) *Correlative Neuroanatomy and Functional Neurology*. Lange Medical Publications, Los Altos.
- Clarey JC, Barone P, and Imig TJ (1992) Physiology of thalamus and cortex. In: Popper AN and Fay RR (eds). *Springer Handbook of Auditory Research*, Volume 2: *The Mammalian Auditory Pathway: Neurophysiology*. Springer-Verlag, New York, pp. 232–334.
- Clarke S, de Ribaupierre F, Rouiller EM, and de Ribaupierre Y (1993) Several neuronal and axonal types form long intrinsic connections in the cat primary auditory cortical field (AD). *Anatomy and Embryology* **188**:117–138.
- Code RA and Winer JA (1985) Commissural neurons in layer III of cat primary auditory cortex (AD): pyramidal and non-pyramidal cell input. *Journal of Comparative Neurology* **242**:485–510.
- Colavita FB, Szeligo FV, and Zimmer SD (1974) Temporal pattern discrimination in cats with insular-temporal lesions. *Brain Research* **79**:153–156.
- Colwell S (1975) Thalamocortical-corticothalamic reciprocity: a combined anterograde-retrograde tracer technique. *Brain Research* **92**:443–449.
- Conlee JW and Kane ES (1982) Descending projections from the inferior colliculus to the dorsal cochlear nucleus in the cat: an autoradiographic study. *Neuroscience* **7**:161–178.
- Coveñas R, Romo R, Cherymy A, Cesselin F, and Conrath M (1986) Immunocytochemical study of enkephalin-like cell bodies in the thalamus of the cat. *Brain Research* **377**:355–361.
- Covey E and Casseday JH (1991) The monaural nuclei of the lateral lemniscus in an echolocating bat: parallel pathways for analyzing temporal features of sound. *Journal of Neuroscience* **11**:3456–3470.
- Cowan WM, Adamson L, and Powell TPS (1961) An experimental study of the avian visual system. *Journal of Anatomy (London)* **95**:545–563.
- Cowey A (1981) Why are there so many visual areas? In: Schmitt FO, Worden FG,

- Adelman G, and Dennis SG (eds). *The Organization of the Cerebral Cortex*. MIT Press, Cambridge, pp. 395–413.
- Creutzfeldt O, Hellweg F-C, and Schreiner C (1980) Thalamocortical transformation of responses to complex auditory stimuli. *Experimental Brain Research* **39**:87–104.
- Deschênes M, Veinante P, and Zhang Z-W (1998) The organization of corticothalamic projections: reciprocity versus parity. *Brain Research Reviews* **28**:286–308.
- Diamond IT (1983) Parallel pathways in the auditory, visual and somatic systems. In: Macchi G, Rustioni A, and Spreafico R (eds). *Somatosensory Integration in the Thalamus*. Elsevier, Amsterdam, pp. 251–272.
- Diamond IT, Goldberg JM, and Neff WD (1962) Tonal discrimination after ablation of auditory cortex. *Journal of Neurophysiology* **25**:223–235.
- Diamond IT, Jones EG, and Powell TPS (1969) The projection of the auditory cortex upon the diencephalon and brain stem of the cat. *Brain Research* **15**:305–340.
- Diehl JJ and Winer JA (1996) Contrasting auditory corticofugal projections to the inferior colliculus and medial geniculate body. *Proceedings of the Society for Neuroscience* **22**:1069.
- Dinse HE and Schreiner CE (2002) Do primary sensory areas play analogous roles in different sensory modalities? In: Schuez A and Miller R (eds). *Cortical Areas: Unity and Diversity*. Taylor & Francis, London, pp. 273–310.
- Doucet JR, Rose L, and Ryugo DK (2002) The cellular origin of corticofugal projections to the superior olivary complex in the rat. *Brain Research* **925**:28–41.
- Edelman GM, Gall WE, and Cowan WM (eds) (1988) *Auditory Function. Neurobiological Bases of Hearing*. John Wiley & Sons, New York.
- Eggermont JJ (1998) Is there a neural code? *Neuroscience and Biobehavioral Reviews* **22**:355–370.
- Eggermont JJ (2001) Between sound and perception: reviewing the search for a neural code. *Hearing Research* **157**:1–42.
- Ehret G (1997) The auditory cortex. *Journal of Comparative Physiology A* **181**:547–557.
- Elverland HH (1978) Ascending and intrinsic projections of the superior olivary complex in the cat. *Experimental Brain Research* **32**:117–134.
- Evans EF (1982) Functional anatomy of the auditory system. In: Barlow HB and Mollon JM (eds). *The Senses*. Cambridge University Press, Cambridge, pp. 251–306.
- Evans EF and Whitfield IC (1964) Classification of unit responses in the auditory cortex of unanaesthetized and unrestrained cat. *Journal of Physiology (London)* **171**:476–493.
- Evans EF, Ross HF, and Whitfield IC (1965) The spatial distribution of unit characteristic frequency in the primary auditory cortex of the cat. *Journal of Physiology (London)* **179**:238–247.
- Faingold CL, Millan MH, Boersma Anderson CA, and Meldrum BS (1989) Induction of audiogenic seizures in normal and genetically epilepsy-prone rats following focal microinjection of an excitant amino acid into reticular formation and auditory nuclei. *Epilepsy Research* **3**:199–205.
- Faingold CL, Randall ME, and Boersma Anderson CA (1994) Blockade of GABA uptake with tiagabine inhibits audiogenic seizures and reduces neuronal firing in the inferior colliculus of the genetically epilepsy-prone rat. *Experimental Neurology* **126**:225–232.
- Fay RR (1988) *Hearing in Vertebrates: A Psychophysics Databook*. Hill-Fay Associates, Winnetka.
- Fekete DM, Rouiller EM, Liberman MC, and Ryugo DK (1984) The central projections

- of intracellularly labeled auditory nerve fibers in cats. *Journal of Comparative Neurology* **229**:432–450.
- Feng AS and Vater M (1985) Functional organization of the cochlear nucleus of rufous horseshoe bats (*Rhinolophus rouxi*): frequencies and internal connections are arranged in slabs. *Journal of Comparative Neurology* **235**:529–553.
- Ferster D and LeVay S (1978) The axonal arborizations of lateral geniculate neurons in the striate cortex of the cat. *Journal of Comparative Neurology* **182**:923–944.
- Fitzpatrick D, Lund JS, and Blasdel GG (1985) Intrinsic connections of macaque striate cortex: afferent and efferent connections of lamina 4C. *Journal of Neuroscience* **5**:3329–3349.
- Foster RE and Hall WC (1978) The organization of central auditory pathways in a reptile, *Iguana iguana*. *Journal of Comparative Neurology* **178**:783–832.
- Friauf E and Ostwald J (1988) Divergent projections of physiologically characterized rat ventral cochlear nucleus neurons as shown by intra-axonal injection of horseradish peroxidase. *Experimental Brain Research* **73**:263–284.
- Games KD and Winer JA (1988) Layer V in rat auditory cortex: projections to the inferior colliculus and contralateral cortex. *Hearing Research* **34**:1–26.
- Gannon PJ, Holloway RL, Broadfield DC, and Braun AR (1998) Asymmetry of chimpanzee planum temporale: humanlike pattern of Wernicke's brain language area homolog. *Science* **279**:220–222.
- Gerren RA and Weinberger NM (1983) Long term potentiation in the magnocellular medial geniculate nucleus of the anesthetized cat. *Brain Research* **265**:138–142.
- Geschwind N (1965) Disconnexion syndromes in animals and man. *Brain* **88**:237–294, 585–644.
- Geschwind N and Levitsky W (1968) Human brain: left-right asymmetries in temporal speech region. *Science* **161**:186–187.
- Glendenning KK and Masterton RB (1983) Acoustic chiasm: efferent projections of the lateral superior olive. *Journal of Neuroscience* **3**:1521–1537.
- Glendenning KK and Masterton RB (1998) Comparative morphometry of mammalian central auditory systems: variation in nuclei and form of the ascending system. *Brain, Behavior and Evolution* **51**:59–89.
- Glendenning KK, Hutson KA, Nudo RJ, and Masterton RB (1985) Acoustic chiasm II: anatomical basis of binaurality in lateral superior olive of cat. *Journal of Comparative Neurology* **232**:261–285.
- Glendenning KK, Baker BN, Hutson KA, and Masterton RB (1992) Acoustic chiasm V: inhibition and excitation in the ipsilateral and contralateral projections of LSO. *Journal of Comparative Neurology* **319**:100–122.
- González-Hernández TH, Galindo-Mireles D, Castañeyra-Perdomo A, and Ferres-Torres R (1991) Divergent projections of projecting neurons of the inferior colliculus to the medial geniculate body and the contralateral inferior colliculus in the rat. *Hearing Research* **52**:17–22.
- González-Hernández TH, Mantolán-Sarmiento B, González-González B, and Pérez-González H (1996) Sources of GABAergic input to the inferior colliculus of the rat. *Journal of Comparative Neurology* **372**:309–326.
- Gregory KM (1974) The stato-acoustic nuclear complex and the nucleus cerebelli of the frog. A Golgi study. *Brain, Behavior and Evolution* **10**:146–156.
- Guinan JJ Jr and Li RY-S (1990) Signal processing in brainstem auditory neurons which

- receive giant endings (calyces of Held) in the medial nucleus of the trapezoid body of the cat. *Hearing Research* **49**:321–334.
- Guinan JJ Jr, Guinan SS, and Norris BE (1972a) Single auditory units in the superior olivary complex. I: Responses to sounds and classifications based on physiological properties. *International Journal of Neuroscience* **4**:101–120.
- Guinan JJ Jr, Norris BE, and Guinan SS (1972b) Single auditory units in the superior olivary complex. II: Locations of unit categories and tonotopic organization. *International Journal of Neuroscience* **4**:147–166.
- Gulick WL, Gescheider GA, and Frisina RD (1989) *Hearing. Physiological Acoustics, Neural Coding, and Psychoacoustics*. Oxford University Press, New York.
- Hackett TA, Preuss TM, and Kaas JH (2001) Architectonic identification of the core region in auditory cortex of macaques, chimpanzees, and humans. *Journal of Comparative Neurology* **441**:197–222.
- Hall JC and Feng AS (1987) Evidence for parallel processing in the frog's auditory thalamus. *Journal of Comparative Neurology* **258**:407–419.
- Harrison JM and Feldman ML (1970) Anatomical aspects of the cochlear nucleus and superior olivary complex. In: Neff WD (ed). *Contributions to Sensory Physiology*. Academic Press, New York, pp. 95–142.
- Harrison JM and Irving R (1966) Visual and nonvisual auditory systems in mammals. *Science* **154**:738–743.
- He J, Hashikawa T, Ojima H, and Kinouchi Y (1997) Temporal integration and duration tuning in the dorsal zone of cat auditory cortex. *Journal of Neuroscience* **17**:2615–2625.
- Heffner HE and Heffner RS (1984) Temporal lobe lesions and perception of species-specific vocalizations by macaques. *Science* **226**:75–76.
- Heil P, Rajan R, and Irvine DRF (1994) Topographic representation of tone intensity along the isofrequency axis of cat primary auditory cortex. *Hearing Research* **76**:188–202.
- Hendry SHC and Jones EG (1991) GABA neuronal subpopulations in cat primary auditory cortex—co-localization with calcium binding proteins. *Brain Research* **543**:45–55.
- Howard MA, Volkov IO, Mirsky R, Garell PC, Noh MD, Granner M, Damasio H, Steinschneider M, Reale RA, Hind JE, and Brugge JF (2000) Auditory cortex on the human posterior superior temporal gyrus. *Journal of Comparative Neurology* **416**:79–92.
- Huang CL and Winer JA (2000) Auditory thalamocortical projections in the cat: laminar and areal patterns of input. *Journal of Comparative Neurology* **427**:302–331.
- Huang CL, Larue DT, and Winer JA (1999) GABAergic organization of the cat medial geniculate body. *Journal of Comparative Neurology* **415**:368–392.
- Hübener M, Shoham D, Grinvald A, and Bonhoeffer T (1997) Spatial relationships among three columnar systems in cat area 17. *Journal of Neuroscience* **17**:9270–9284.
- Huffman RF and Henson OW Jr (1990) The descending auditory pathway and acousticomotor systems: connections with the inferior colliculus. *Brain Research Reviews* **15**:295–323.
- Hutsler JJ and Gazzaniga MS (1996) Acetylcholinesterase staining in human auditory and language cortices: regional variation of structural features. *Cerebral Cortex* **6**:260–270.
- Hutson KA and Morest DK (1996) Fine structure of the cell clusters in the cochlear nerve root: stellate, granule, and mitt cells offer insights into the synaptic organization of local circuit neurons. *Journal of Comparative Neurology* **371**:397–414.
- Hutson KA, Glendenning KK, and Masterton RB (1991) Acoustic chiasm IV: eight mid-

- brain decussations of the auditory system in the cat. *Journal of Comparative Neurology* **312**:105–131.
- Imig TJ and Adrián HO (1977) Binaural columns in the primary auditory field (A1) of cat auditory cortex. *Brain Research* **138**:241–257.
- Imig TJ and Morel A (1984) Topographic and cytoarchitectonic organization of thalamic neurons related to their targets in low-, middle-, and high-frequency representations in cat auditory cortex. *Journal of Comparative Neurology* **227**:511–539.
- Imig TJ and Morel A (1985a) Tonotopic organization in lateral part of posterior group of thalamic nuclei in the cat. *Journal of Neurophysiology* **53**:836–851.
- Imig TJ and Morel A (1985b) Tonotopic organization in ventral nucleus of medial geniculate body in the cat. *Journal of Neurophysiology* **53**:309–340.
- Imig TJ and Reale RA (1980) Patterns of cortico-cortical connections related to tonotopic maps in cat auditory cortex. *Journal of Comparative Neurology* **192**:293–332.
- Imig TJ and Reale RA (1981) Ipsilateral corticocortical projections related to binaural columns in cat primary auditory cortex. *Journal of Comparative Neurology* **203**:1–14.
- Irvine DRF (1986) The auditory brainstem. A review of the structure and function of auditory brainstem processing mechanisms. In: Autrum H, Ottoson D, Perl ER, Schmidt RF, Shimazu H, and Willis WD (eds). *Progress in Sensory Physiology* **7**:1–279.
- Jack JJB, Noble D, and Tsien RW (1975) *Electric Current Flow in Excitable Cells*. Oxford University Press, London.
- Jane JA, Masterton RB, and Diamond IT (1965) The function of the tectum for attention to auditory stimuli in the cat. *Journal of Comparative Neurology* **125**:165–192.
- Jeffress LA (1948) A place theory of sound localization. *Journal of Comparative and Physiological Psychology* **41**:35–39.
- Jen PH-S, Chen QC, and Sun XD (1998) Corticofugal regulation of auditory sensitivity in the bat inferior colliculus. *Journal of Comparative Physiology A* **183**:683–697.
- Jenkins WM and Masterton RB (1982) Sound localization: effects of unilateral lesions in central auditory system. *Journal of Neurophysiology* **47**:987–1016.
- Jonas P, Bischofberger J, and Sandkühler J (1998) Corelease of two fast neurotransmitters at a central synapse. *Science* **281**:419–424.
- Jones EG (1985) *The Thalamus*. Plenum Press, New York.
- Jones EG and Powell TPS (1970) An anatomical study of converging sensory pathways within the cerebral cortex of the monkey. *Brain* **93**:793–820.
- Jones EG and Powell TPS (1973) Anatomical organization of the somatosensory cortex. In: Iggo A (ed). *Handbook of Sensory Physiology*, Volume II: *Somatosensory System*. Springer-Verlag, Berlin, pp. 579–620.
- Jones EG, Burton H, Saper CB, and Swanson LW (1976) Midbrain, diencephalic and cortical relationships of the basal nucleus of Meynert and associated structures in primates. *Journal of Comparative Neurology* **167**:385–420.
- Kaas JH (1997) Topographic maps are fundamental to sensory processing. *Brain Research Bulletin* **44**:107–112.
- Kaas JH and Hackett TA (2000) Subdivisions of auditory cortex and processing streams in primates. *Proceedings of the National Academy of Sciences of the United States of America* **97**:11793–11799.
- Kaas JH, Nelson RJ, Sur M, Dykes RW, and Merzenich MM (1984) The somatotopic organization of the ventroposterior thalamus of the squirrel monkey, *Saimiri sciureus*. *Journal of Comparative Neurology* **226**:111–140.

- Kane ES and Conlee JW (1979) Descending inputs to the caudal cochlear nucleus of the cat: degeneration and autoradiographic studies. *Journal of Comparative Neurology* **187**:759–784.
- Kawamura K (1971) Variations of the cerebral sulci of the cat. *Acta Anatomica* **80**:204–221.
- Kawamura K (1973) Corticocortical fiber connections of the cat cerebrum. I. The temporal region. *Brain Research* **51**:1–21.
- Kelly JB (1973) The effects of insular and temporal lesions in cats on two types of auditory pattern discrimination. *Brain Research* **62**:71–87.
- Kelly JB and Judge PW (1985) Effects of medial geniculate lesions on sound localization by the rat. *Journal of Neurophysiology* **53**:361–372.
- Kelly JB and Wong D (1981) Laminar connections of the cat's auditory cortex. *Brain Research* **212**:1–15.
- Kiang NY-S, Watanabe T, Thomas EC, and Clark LF (1965) *Discharge Patterns of Single Fibers in the Cat's Auditory Nerve*. MIT Press, Cambridge.
- Kiang NYS, Morest DK, Godfrey DA, Guinan JJ Jr, and Kane EC (1973) Stimulus coding at caudal levels of the cat's auditory nervous system. I. Response characteristics of single neurons. In: Møller AR (ed). *Basic Mechanisms in Hearing*. Academic Press, New York, pp. 455–478.
- Kilgard MP and Merzenich MM (1998a) Cortical map reorganization enabled by nucleus basalis activity. *Science* **279**:1714–1718.
- Kilgard MP and Merzenich MM (1998b) Plasticity of temporal information processing in the primary auditory cortex. *Nature Neuroscience* **1**:727–731.
- Kilgard MP and Merzenich MM (2002) Order-sensitive plasticity in adult primary auditory cortex. *Proceedings of the National Academy of Sciences of the United States of America* **99**:3205–3209.
- Kilgard MP, Pandya PK, Vasquez J, Gehi A, Schreiner CE, and Merzenich MM (2001) Sensory input directs spatial and temporal plasticity in primary auditory cortex. *Journal of Neurophysiology* **86**:326–338.
- Kim DO and Molnar CE (1979) A population study of cochlear nerve fibers: comparison of spatial distributions of average-rate and phase-locking measures of responses to single tones. *Journal of Neurophysiology* **42**:16–30.
- Kitzes LM, Wrege KS, and Cassady JM (1980) Patterns of responses of cortical cells to binaural stimulation. *Journal of Comparative Neurology* **192**:455–472.
- Knudsen EI and Konishi M (1978) A neural map of auditory space in the owl. *Science* **200**:795–797.
- Konishi M (1973a) Locatable and non-locatable acoustic signals for barn owls. *American Naturalist* **107**:775–785.
- Konishi M (1973b) How the owl tracks its prey. *American Scientist* **61**:414–427.
- Kudo M and Niimi K (1980) Ascending projections to the inferior colliculus in the cat: an autoradiographic study. *Journal of Comparative Neurology* **191**:545–556.
- Langner G (1992) Periodicity coding in the auditory system. *Hearing Research* **60**:115–142.
- Langner G, Schreiner CE, and Merzenich MM (1987) Covariation of latency and temporal resolution in the inferior colliculus of the cat. *Hearing Research* **31**:197–201.
- Leake-Jones PA and Snyder RL (1982) Uptake and transport of horseradish peroxidase by cochlear spiral ganglion neurons. *Hearing Research* **8**:199–223.
- LeDoux JE, Ruggiero DA, and Reis DJ (1985) Projections to the subcortical forebrain from anatomically defined regions of the medial geniculate body in the rat. *Journal of Comparative Neurology* **242**:182–213.

- LeDoux JE, Iwata J, Pearl D, and Reis DJ (1986) Disruption of auditory but not visual learning by destruction of intrinsic neurons in the rat medial geniculate body. *Brain Research* **371**:395–399.
- LeVay S and Gilbert CD (1976) Laminar patterns of geniculocortical projection in the cat. *Brain Research* **113**:1–19.
- LeVay S, Stryker MP, and Shatz CJ (1978) Ocular dominance columns and their development in layer IV of the cat's visual cortex: a quantitative study. *Journal of Comparative Neurology* **179**:223–244.
- Levey AI, Hallanger AE, and Wainer BH (1987) Choline acetyltransferase immunoreactivity in the rat thalamus. *Journal of Comparative Neurology* **257**:317–332.
- Liberman MC and Brown MC (1986) Physiology and anatomy of single olivocochlear neurons in the cat. *Hearing Research* **24**:17–36.
- Lorente de N6 R (1933) Anatomy of the eighth nerve: the central projection of the nerve endings of the inner ear. *Laryngoscope* **43**:1–38.
- Lorente de N6 R (1981) *The Primary Acoustic Nuclei*. Raven Press, New York.
- Luetheke LE, Krubitzer LA, and Kaas JH (1988) Cortical connections of electrophysiologically and architectonically defined subdivisions of auditory cortex in squirrels. *Journal of Comparative Neurology* **268**:181–203.
- Malmierca MS, Leergaard TB, Bajo VM, Bjaalie JG, and Merchán MA (1998) Anatomic evidence of a three-dimensional mosaic pattern of tonotopic organization in the ventral complex of the lateral lemniscus in cat. *Journal of Neuroscience* **18**:10603–10618.
- Margoliash D (1987) Neural plasticity in birdsong learning. In: Rauschecker JP and Marler P (eds). *Imprinting and Cortical Plasticity*. John Wiley & Sons, New York, pp. 23–54.
- Markovitz NS and Pollak GD (1993) The dorsal nucleus of the lateral lemniscus in the mustache bat: monaural properties. *Hearing Research* **71**:51–63.
- Markovitz NS and Pollak GD (1994) Binaural processing in the dorsal nucleus of the lateral lemniscus. *Hearing Research* **73**:121–140.
- Masterton B, Thompson GC, Bechtold JK, and RoBards MJ (1975) Neuroanatomical basis of binaural phase-difference analysis for sound localization: a comparative study. *Journal of Comparative and Physiological Psychology* **89**:379–386.
- Matsubara JA and Phillips DP (1988) Intracortical connections and their physiological correlates in the primary auditory cortex (AI) of the cat. *Journal of Comparative Neurology* **268**:38–48.
- Matthews PBC (1972) *Mammalian Muscle Receptors and Their Central Actions*. Arnold, London.
- Mendelson JR, Schreiner CE, Sutter ML, and Grasse KL (1993) Functional topography of cat primary auditory cortex: responses to frequency-modulated sweeps. *Experimental Brain Research* **94**:65–87.
- Merchán MA, Saldaña E, and Plaza I (1994) Dorsal nucleus of the lateral lemniscus in the rat: concentric organization and tonotopic projection to the inferior colliculus. *Journal of Comparative Neurology* **342**:259–278.
- Merzenich MM and Brugge JF (1973) Representation of the cochlear partition on the superior temporal plane of the macaque monkey. *Brain Research* **50**:275–296.
- Merzenich MM and Reid MD (1974) Representation of the cochlea within the inferior colliculus of the cat. *Brain Research* **77**:397–415.
- Merzenich MM, Knight PL, and Roth GL (1975) Representation of cochlea within primary auditory cortex in the cat. *Journal of Neurophysiology* **38**:231–249.
- Middlebrooks JC and Pettigrew JD (1981) Functional classes of neurons in primary

- auditory cortex of the cat distinguished by sensitivity to sound location. *Journal of Neuroscience* **1**:107–120.
- Middlebrooks JC and Zook JM (1983) Intrinsic organization of the cat's medial geniculate body identified by projections to binaural response-specific bands in the primary auditory cortex. *Journal of Neuroscience* **3**:203–225.
- Middlebrooks JC, Dykes RW, and Merzenich MM (1980) Binaural response-specific bands in primary auditory cortex (AI) of the cat: topographic organization orthogonal to isofrequency contours. *Brain Research* **181**:31–48.
- Miller LM and Schreiner CE (2000) Stimulus based state control in the thalamocortical system. *Journal of Neuroscience* **20**:7011–7016.
- Miller LM, Escabí MA, Read HL, and Schreiner CE (2001) Functional convergence of response properties in the auditory thalamocortical system. *Neuron* **32**:151–160.
- Mitani A and Shimokouchi M (1985) Neuronal connections in the primary auditory cortex: an electrophysiological study in the cat. *Journal of Comparative Neurology* **235**:417–429.
- Mitani A, Itoh K, Nomura S, Kudo M, Kaneko T, and Mizuno N (1984) Thalamocortical projections to layer I of the primary auditory cortex in the cat: a horseradish peroxidase study. *Brain Research* **310**:347–350.
- Moore JK and Moore RY (1971) A comparative study of the superior olivary complex in the primate brain. *Folia Primatologica (Basel)* **16**:35–51.
- Morel A and Imig TJ (1987) Thalamic projections to fields A, AI, P, and VP in the cat auditory cortex. *Journal of Comparative Neurology* **265**:119–144.
- Morel A and Kaas JH (1992) Subdivisions and connections of auditory cortex in owl monkeys. *Journal of Comparative Neurology* **318**:27–63.
- Morel A, Garraghty PE, and Kaas JH (1993) Tonotopic organization, architectonic fields, and connections of auditory cortex in macaque monkeys. *Journal of Comparative Neurology* **335**:437–459.
- Morel A, Rouiller E, de Ribaupierre Y, and de Ribaupierre F (1987) Tonotopic organization in the medial geniculate body (MGB) of lightly anesthetized cats. *Experimental Brain Research* **69**:24–42.
- Morest DK (1965) The laminar structure of the medial geniculate body of the cat. *Journal of Anatomy (London)* **99**:143–160.
- Morest DK (1971) Dendrodendritic synapses of cells that have axons: the fine structure of the Golgi type II cell in the medial geniculate body of the cat. *Zeitschrift für Anatomie und Entwicklungsgeschichte* **133**:216–246.
- Morest DK (1975) Synaptic relationships of Golgi type II cells in the medial geniculate body of the cat. *Journal of Comparative Neurology* **162**:157–194.
- Morest DK and Oliver DL (1984) The neuronal architecture of the inferior colliculus in the cat: defining the functional anatomy of the auditory midbrain. *Journal of Comparative Neurology* **222**:209–236.
- Mountcastle VB (1997) The columnar organization of the neocortex. *Brain* **120**:701–722.
- Mulligan KA and Törk I (1988) Serotonergic innervation of the cat cerebral cortex. *Journal of Comparative Neurology* **270**:86–110.
- Neff WD, Diamond IT, and Casseday JH (1975) Behavioral studies of auditory discrimination: central nervous system. In: Keidel WD and Neff WD (eds). *Handbook of Sensory Physiology*, Volume V, Part 2: *Auditory System, Anatomy, Physiology (Ear)*. Springer-Verlag, Berlin, pp. 307–400.
- Nelson PG and Erulkar SD (1963) Synaptic mechanisms of excitation and inhibition in the central auditory pathway. *Journal of Neurophysiology* **26**:908–923.



- Niimi K and Matsuoka H (1979) Thalamocortical organization of the auditory system in the cat studied by retrograde axonal transport of horseradish peroxidase. *Advances in Anatomy, Embryology and Cell Biology* **57**:1–56.
- Nobre AC and Plunkett K (1997) The neural system of language: structure and development. *Current Opinion in Neurobiology* **7**:262–268.
- Ohl FW and Scheich H (1997) Learning-induced dynamic receptive field changes in primary auditory cortex (AI) of the unanesthetized Mongolian gerbil. *Journal of Comparative Physiology A* **181**:685–696.
- Ojima H, Honda CN, and Jones EG (1992) Characteristics of intracellularly injected infragranular pyramidal neurons in cat primary auditory cortex. *Cerebral Cortex* **2**:197–216.
- Oliver DL (1984) Dorsal cochlear nucleus projections to the inferior colliculus in the cat. A light and electron microscopic study. *Journal of Comparative Neurology* **224**:155–172.
- Oliver DL (1987) Projections to the inferior colliculus from the anteroventral cochlear nucleus in the cat: possible substrates for binaural interaction. *Journal of Comparative Neurology* **264**:24–46.
- Oliver DL and Huerta MF (1992) Inferior and superior colliculi. In: Webster DB, Popper AN, and Fay RR (eds). *Springer Handbook of Auditory Research*, Volume 1: *The Mammalian Auditory Pathway: Neuroanatomy*. Springer-Verlag, New York, pp. 168–221.
- Oliver DL and Morest DK (1984) The central nucleus of the inferior colliculus in the cat. *Journal of Comparative Neurology* **222**:237–264.
- Oliver DL, Potashner SJ, Jones DR, and Morest DK (1983) Selective labeling of spiral ganglion and granule cells with D-aspartate in the auditory system of cat and guinea pig. *Journal of Neuroscience* **3**:455–472.
- Oliver DL, Kuwada S, Yin TCT, Haberly LB, and Henkel CK (1991) Dendritic and axonal morphology of HRP-injected neurons in the inferior colliculus of the cat. *Journal of Comparative Neurology* **303**:75–100.
- Oliver DL, Winer JA, Beckius GE, and Saint Marie RL (1994) Morphology of GABAergic cells and axon terminals in the cat inferior colliculus. *Journal of Comparative Neurology* **340**:27–42.
- Oliver DL, Beckius GE, and Shneiderman A (1995) Axonal projections from the lateral and medial superior olive to the inferior colliculus of the cat: a study using electron microscopic autoradiography. *Journal of Comparative Neurology* **360**:17–32.
- Oliver DL, Beckius GE, Bishop DC, and Kuwada S (1997) Simultaneous anterograde labeling of axonal layers from lateral superior olive and dorsal cochlear nucleus in the inferior colliculus of cat. *Journal of Comparative Neurology* **382**:215–229.
- Osen KK (1969) Cytoarchitecture of the cochlear nuclei in the cat. *Journal of Comparative Neurology* **136**:453–484.
- Osen KK (1970) Course and termination of the primary afferents in the cochlear nuclei of the cat. An experimental anatomical study. *Archives Italiennes de Biologie* **108**:21–51.
- Osen KK (1972) Projection of the cochlear nuclei on the inferior colliculus in the cat. *Journal of Comparative Neurology* **144**:355–372.
- Osen KK (1988) Anatomy of the mammalian cochlear nuclei; a review. In: Syka J and Masterton RB (eds). *Auditory Pathway: Structure and Function*. Plenum Press, New York, pp. 65–75.
- Osen KK, Ottersen OP, and Storm-Mathisen J (1990) Colocalization of glycine-like and GABA-like immunoreactivities: a semiquantitative study in the dorsal cochlear nucleus

- of the cat. In: Ottersen OP and Storm-Mathisen J (eds). *Glycine Neurotransmission*. John Wiley & Sons, Chichester, pp. 417–451.
- Papez JW (1929) *Comparative Neurology. A Manual and Text for the Study of the Nervous System of Vertebrates*. Hafner, New York.
- Peters A, Palay SL, and Webster HdeF (1991) *The Fine Structure of the Nervous System. The Neurons and Their Supporting Cells*. Oxford University Press, New York.
- Phillips DP (1989) The neural coding of simple and complex sounds in the auditory cortex. In: Lund JS (ed). *Sensory Processing in the Mammalian Brain. Neural Substrates and Experimental Strategies*. Oxford University Press, New York, pp. 172–203.
- Phillips DP, Orman SS, Musicant AD, and Wilson GF (1985) Neurons in the cat's primary auditory cortex distinguished by their responses to tones and wide-spectrum noise. *Hearing Research* **18**:73–86.
- Phillips DP, Reale RA, and Brugge JF (1991) Stimulus processing in auditory cortex. In: Altschuler RA, Bobbin RP, Clopton BM, and Hoffman DW (eds). *Neurobiology of Hearing: The Central Auditory System*. Raven Press, New York, pp. 335–364.
- Phillips DP, Semple MN, and Kitzes LM (1995) Factors shaping the tone level sensitivity of single neurons in posterior field of cat auditory cortex. *Journal of Neurophysiology* **73**:674–686.
- Pollak GD and Casseday JH (1989) *The Neural Basis of Echolocation in Bats*. Springer-Verlag, Berlin.
- Pontes C, Reis FF, and Sousa-Pinto A (1975) The auditory cortical projections onto the medial geniculate body in the cat: an experimental anatomical study with silver and autoradiographic methods. *Brain Research* **91**:43–63.
- Popper AN and Fay RR (eds) (1992) *Springer Handbook of Auditory Research, Volume 2: The Mammalian Auditory Pathway: Neurophysiology*. Springer-Verlag, New York.
- Prieto JJ and Winer JA (1999) Neurons of layer VI in cat primary auditory cortex (AI): Golgi study and sublaminal origins of projection neurons. *Journal of Comparative Neurology* **404**:332–358.
- Prieto JJ, Peterson BA, and Winer JA (1994a) Morphology and spatial distribution of GABAergic neurons in cat primary auditory cortex (AI). *Journal of Comparative Neurology* **344**:349–382.
- Prieto JJ, Peterson BA, and Winer JA (1994b) Laminar distribution and neuronal targets of GABAergic axon terminals in cat primary auditory cortex (AI). *Journal of Comparative Neurology* **344**:383–402.
- Pritz MB (1974a) Ascending connections of a midbrain auditory area in a crocodile, *Caiman crocodilus*. *Journal of Comparative Neurology* **153**:179–198.
- Pritz MB (1974b) Ascending connections of a thalamic auditory area in a crocodile, *Caiman crocodilus*. *Journal of Comparative Neurology* **153**:199–214.
- Przybylski AW (1998) Does top-down processing help us to see? *Current Biology* **8**:R135–R139.
- Rakic P (ed.) (1975) Local circuit neurons. *Neuroscience Research Program Bulletin* **13**: 291–446.
- Rauschecker JP and Tian B (2000) Mechanisms and streams for processing of “what” and “where” in auditory cortex. *Proceedings of the National Academy of Sciences of the United States of America* **97**:11800–11806.
- Read HL, Winer JA, and Schreiner CE (2001) Modular organization of intrinsic connections associated with spectral tuning in cat auditory cortex. *Proceedings of the National Academy of Sciences of the United States of America* **98**:8042–8047.

- Read HL, Winer JA, and Schreiner CE (2002) Functional architecture of auditory cortex. *Current Opinion in Neurobiology* **12**:433–440.
- Reale RA and Imig TJ (1980) Tonotopic organization in auditory cortex of the cat. *Journal of Comparative Neurology* **182**:265–291.
- Recanzone GH (1998) Rapidly induced auditory plasticity: the ventriloquism aftereffect. *Proceedings of the National Academy of Sciences of the United States of America* **95**:869–875.
- Recanzone GH, Schreiner CE, and Merzenich MM (1993) Plasticity in the frequency representation of primary auditory cortex following discrimination training in adult owl monkeys. *Journal of Neuroscience* **13**:87–103.
- Rhode WS (1991) Physiological-morphological properties of the cochlear nucleus. In: Altschuler RA, Bobbin RP, Clopton BM, and Hoffman DW (eds). *Neurobiology of Hearing: The Central Auditory System*. Raven Press, New York, pp. 47–78.
- Rhode WS and Greenberg S (1992) Physiology of the cochlear nuclei. In: Popper AN and Fay RR (eds). *Springer Handbook of Auditory Research, Volume 2: The Mammalian Auditory System: Neurophysiology*. Springer-Verlag, New York, pp. 94–152.
- Rhode WS, Oertel D, and Smith PH (1983a) Physiological response properties of cells labeled intracellularly with horseradish peroxidase in cat dorsal cochlear nucleus. *Journal of Comparative Neurology* **213**:426–447.
- Rhode WS, Oertel D, and Smith PH (1983b) Physiological response properties of cells labeled intracellularly with horseradish peroxidase in cat ventral cochlear nucleus. *Journal of Comparative Neurology* **213**:448–463.
- Richter EA, Norris BE, Fullerton BC, Levine RA, and Kiang NY-S (1983) Is there a medial nucleus of the trapezoid body in humans? *American Journal of Anatomy* **68**:157–166.
- Rinvik E, Ottersen OP, and Storm-Mathisen J (1987) Gamma-aminobutyrate-like immunoreactivity in the thalamus of the cat. *Neuroscience* **21**:787–805.
- Robert D, Amoroso J, and Hoy RR (1992) The evolutionary convergence of hearing in a parasitoid fly and its cricket host. *Science* **258**:1135–1137.
- Rockel AJ and Jones EG (1973a) The neuronal organization of the inferior colliculus of the adult cat. I. The central nucleus. *Journal of Comparative Neurology* **147**:11–60.
- Rockel AJ and Jones EG (1973b) The neuronal organization of the inferior colliculus of the adult cat. II. The pericentral nucleus. *Journal of Comparative Neurology* **149**:301–334.
- Rockel AJ and Jones EG (1973c) Observations on the fine structure of the central nucleus of the inferior colliculus of the cat. *Journal of Comparative Neurology* **147**:61–92.
- Rodrigues-Dagaeff C, Simm G, de Ribaupierre Y, Villa A, de Ribaupierre F, and Rouiller EM (1989) Functional organization of the ventral division of the medial geniculate body of the cat: evidence for a rostro-caudal gradient of response properties and cortical projections. *Hearing Research* **39**:103–125.
- Romanski LM and LeDoux JE (1993) Information cascade from primary auditory cortex to the amygdala: corticocortical and corticoamygdaloid projections of temporal cortex in the rat. *Cerebral Cortex* **3**:515–532.
- Romanski LM, Tian B, Fritz J, Mishkin M, Goldman-Rakic PS, and Rauschecker JP (1999) Dual streams of auditory afferents target multiple domains in the primate prefrontal cortex. *Nature Neuroscience* **2**:1131–1136.
- Rose JE (1949) The cellular structure of the auditory region of the cat. *Journal of Comparative Neurology* **91**:409–440.
- Rose JE and Woolsey CN (1958) Cortical connections and functional organization

- of thalamic auditory system of cat. In: Harlow HF and Woolsey CN (eds). *Biological and Biochemical Bases of Behavior*. University of Wisconsin Press, Madison, pp. 127–150.
- Rose JE, Galambos R, and Hughes JR (1959) Microelectrode studies of the cochlear nuclei of the cat. *Bulletin of the Johns Hopkins Hospital* **104**:211–251.
- Rose JE, Gross NB, Geisler CD, and Hind JE (1963) Some neural mechanisms in the inferior colliculus of the cat which may be relevant to localization of a sound source. *Journal of Neurophysiology* **26**:288–314.
- Ross LS, Pollak GD, and Zook JM (1988) Origin of ascending projections to an iso-frequency region of the mustache bat's inferior colliculus. *Journal of Comparative Neurology* **270**:488–505.
- Roth GL, Aitkin LM, Andersen RA, and Merzenich MM (1978) Some features of the spatial organization of the central nucleus of the inferior colliculus of the cat. *Journal of Comparative Neurology* **182**:661–680.
- Rouiller EM, Rodrigues-Dagaëff C, Simm GM, de Ribaupierre Y, Villa AEP, and de Ribaupierre F (1989) Functional organization of the medial division of the medial geniculate body of the cat: tonotopic organization, spatial distribution of response properties and cortical connections. *Hearing Research* **39**:127–146.
- Rouiller EM, Capt M, Hornung JP, and Streit P (1990) Correlation between regional changes in the distributions of GABA-containing neurons and unit response properties in the medial geniculate body of the cat. *Hearing Research* **49**:249–258.
- Rouiller EM, Simm GM, Villa AEP, de Ribaupierre Y, and de Ribaupierre F (1991) Auditory corticocortical interconnections in the cat: evidence for parallel and hierarchical arrangement of the auditory cortical areas. *Experimental Brain Research* **86**:483–505.
- Ryugo DK (1992) The auditory nerve: peripheral innervation, cell body morphology, and central projections. In: Webster DB, Popper AN, and Fay RR (eds). *Springer Handbook of Auditory Research*, Volume 1: *The Mammalian Auditory Pathway: Neuroanatomy*. Springer-Verlag, New York, pp. 23–65.
- Saint Marie RL (1996) Glutamatergic connections of the auditory midbrain: selective uptake and axonal transport of D-[<sup>3</sup>H]aspartate. *Journal of Comparative Neurology* **373**:255–270.
- Saint Marie RL, Ostapoff E-M, Morest DK, and Wenthold RJ (1989) Glycine-immunoreactive projections of the cat lateral superior olive: possible role in midbrain ear dominance. *Journal of Comparative Neurology* **279**:382–396.
- Saint Marie RL, Shneiderman A, and Stanforth DA (1997) Patterns of  $\gamma$ -aminobutyric acid and glycine immunoreactivities reflect structural and functional differences of the cat lateral lemniscal nuclei. *Journal of Comparative Neurology* **389**:264–276.
- Sanes DH, Goldstein NA, Ostad M, and Hillman DE (1990) Dendritic morphology of central auditory neurons correlates with their tonotopic position. *Journal of Comparative Neurology* **294**:443–454.
- Schneider GE (1969) Two visual systems. *Science* **163**:895–902.
- Schofield BR and Cant NB (1991) Organization of the superior olivary complex in the guinea pig. I. Cytoarchitecture, cytochrome oxidase histochemistry, and dendritic morphology. *Journal of Comparative Neurology* **314**:645–670.
- Schreiner CE (1995) Order and disorder in auditory cortical maps. *Current Opinion in Neurobiology* **5**:489–496.
- Schreiner CE and Cynader MS (1984) Basic functional organization of second auditory cortical field (AII) of the cat. *Journal of Neurophysiology* **51**:1284–1305.

- Schreiner CE and Langner G (1988) Periodicity coding in the inferior colliculus of the cat. II. Topographical organization. *Journal of Neurophysiology* **60**:1823–1840.
- Schreiner CE and Langner G (1997) Laminar fine structure of frequency organization in auditory midbrain. *Nature* **388**:383–386.
- Schreiner CE and Mendelson JR (1990) Functional topography of cat primary auditory cortex: distribution of integrated excitation. *Journal of Neurophysiology* **64**:1442–1459.
- Schreiner CE and Sutter ML (1992) Topography of excitatory bandwidth in cat primary auditory cortex: single-neuron versus multiple-neuron recordings. *Journal of Neurophysiology* **68**:1487–1502.
- Schreiner CE, Mendelson JR, and Sutter ML (1992) Functional topography of cat primary auditory cortex: representation of tone intensity. *Experimental Brain Research* **92**:105–122.
- Schuller G and Radtke-Schuller S (1990) Neural control of vocalization in bats: mapping of brainstem areas with electrical microstimulation eliciting species-specific echolocation calls in the rufous horseshoe bat. *Experimental Brain Research* **79**:192–206.
- Schwartz IR (1992) The superior olivary complex and lateral lemniscal nuclei. In: Webster DB, Popper AN, and Fay RR (eds). *Springer Handbook of Auditory Research, Volume 1: The Mammalian Auditory Pathway: Neuroanatomy*. Springer-Verlag, New York, pp. 117–167.
- Seltzer B and Pandya DN (1994) Parietal, temporal and occipital projections to cortex of the superior temporal sulcus in the rhesus monkey: a retrograde tracer study. *Journal of Comparative Neurology* **343**:445–463.
- Servièrè J, Webster WR, and Calford MB (1984) Isofrequency labelling revealed by a combined [<sup>14</sup>C]-2-deoxyglucose, electrophysiological, and horseradish peroxidase study of the inferior colliculus of the cat. *Journal of Comparative Neurology* **228**:463–477.
- Sherman SM and Guillery RW (2000) *Exploring the Thalamus*. Academic Press, Orlando.
- Shinonaga Y, Takada M, and Mizuno N (1994) Direct projections from the non-laminated divisions of the medial geniculate nucleus to the temporal polar cortex and amygdala in the cat. *Journal of Comparative Neurology* **340**:405–426.
- Shiple C, Buchwald JS, and Carterette EC (1988) The role of auditory feedback in the vocalizations of cats. *Experimental Brain Research* **69**:431–438.
- Shneiderman A and Henkel CK (1987) Banding of lateral superior olivary nucleus afferents to the inferior colliculus: a possible substrate for sensory integration. *Journal of Comparative Neurology* **266**:519–534.
- Shneiderman A, Oliver DL, and Henkel CK (1988) Connections of the dorsal nucleus of the lateral lemniscus: an inhibitory parallel pathway in the ascending auditory system? *Journal of Comparative Neurology* **276**:188–208.
- Sinclair D (1981) *Mechanisms of Cutaneous Sensation*. Oxford University Press, Oxford.
- Sindberg RM and Thompson RF (1962) Auditory response fields in ventral temporal and insular cortex of cat. *Journal of Neurophysiology* **25**:21–28.
- Smith Y, Séguela P, and Parent A (1987) Distribution of GABA-immunoreactive neurons in the thalamus of the squirrel monkey (*Saimiri sciureus*). *Neuroscience* **22**:579–591.
- Sousa-Pinto A (1973) Cortical projections of the medial geniculate body in the cat. *Advances in Anatomy, Embryology and Cell Biology* **48**:1–42.
- Spangler KM and Morley BJ (1987) Somatostatin-like immunoreactivity in the midbrain of the cat. *Journal of Comparative Neurology* **260**:87–97.

- Spangler KM, Warr WB, and Henkel CK (1985) The projections of principal cells of the medial nucleus of the trapezoid body of the cat. *Journal of Comparative Neurology* **238**:249–261.
- Spangler KM, Cant NB, Henkel CK, Farley GR, and Warr WB (1987) Descending projections from the superior olivary complex to the cochlear nucleus of the cat. *Journal of Comparative Neurology* **259**:452–465.
- Spirou GA and Berrebi AS (1997) Glycine immunoreactivity in the lateral nucleus of the trapezoid body of the cat. *Journal of Comparative Neurology* **383**:473–488.
- Stanley GB, Li FF, and Dan Y (1999) Reconstruction of natural scenes from ensemble responses in the lateral geniculate nucleus. *Journal of Neuroscience* **19**:8036–8042.
- Stiebler I and Ehret G (1985) Inferior colliculus of the house mouse. I. A quantitative study of tonotopic organization, frequency representation, and tone-threshold distribution. *Journal of Comparative Neurology* **238**:65–76.
- Stuart G, Spruston N, Sakmann B, and Häusser M (1997) Action potential initiation and backpropagation in neurons of the mammalian CNS. *Trends in Neurosciences* **20**:125–131.
- Suga N and Jen PH-S (1976) Disproportionate tonotopic representation for processing CF-FM sonar signals in the mustache bat auditory cortex. *Science* **194**:542–544.
- Suga N and Yajima Y (1988) Auditory-vocal integration in the midbrain of the mustached bat: periaqueductal gray and reticular formation. In: Newman JD (ed). *The Physiological Control of Mammalian Vocalization*. Plenum Press, New York, pp. 87–107.
- Tardif E and Clarke S (2001) Intrinsic connectivity of human auditory areas: a tracing study with DiI. *European Journal of Neuroscience* **13**:1045–1050.
- Thompson AM (1998) Heterogeneous projections of the cat posteroventral cochlear nucleus. *Journal of Comparative Neurology* **390**:439–453.
- Thompson R (1969) Localization of the “visual memory system” in the white rat. *Journal of Comparative and Physiological Psychology* **69**:1–29.
- Tsuchitani C (1977) Functional organization of lateral cell groups of cat superior olivary complex. *Journal of Neurophysiology* **40**:296–318.
- Tsuchitani C and Boudreau JC (1966) Single unit analysis of cat superior olive S segment with tonal stimuli. *Journal of Neurophysiology* **29**:684–697.
- Tsuchitani C and Johnson DH (1991) Binaural cues and signal processing in the superior olivary complex. In: Altschuler RA, Bobbin RP, Clopton BM, and Hoffman DW (eds). *Neurobiology of Hearing: The Central Auditory System*. Raven Press, New York, pp. 163–194.
- Tusa RJ, Palmer LA, and Rosenquist AC (1981) Multiple cortical visual areas: visual field topography in the cat. In: Woolsey CN (ed). *Cortical Sensory Organization, Volume 2, Multiple Visual Areas*. Humana Press, Clifton, pp. 1–31.
- Vater M, Casseday JH, and Covey E (1995) Convergence and divergence of ascending binaural and monaural pathways from the superior olives of the mustached bat. *Journal of Comparative Neurology* **351**:632–646.
- Vater M, Habbicht H, Kössl M, and Grothe B (1992) The functional role of GABA and glycine in monaural and binaural processing in the inferior colliculus of horseshoe bats. *Journal of Comparative Physiology A* **171**:541–553.
- Vetter DE and Mugnaini E (1990) An evaluation of retrograde tracing methods for the identification of chemically distinct cochlear efferent neurons. *Archives Italiennes de Biologie* **128**:331–353.
- Vetter DE and Mugnaini E (1992) Distribution and dendritic features of three groups of rat olivocochlear neurons. *Anatomy and Embryology* **185**:1–16.

- Vetter DE, Saldaña E, and Mugnaini E (1993) Input from the inferior colliculus to medial olivocochlear neurons in the rat: a double label study with PHA-L and cholera toxin. *Hearing Research* **70**:173–186.
- von Economo C and Koskinas GN (1925) *Die Cytoarchitectonik der Hirnrinde des erwachsenen Menschen*. Springer-Verlag, Berlin.
- Wallace MN, Kitzes LM, and Jones EG (1991) Chemoarchitectonic organization of the cat primary auditory cortex. *Experimental Brain Research* **86**:518–526.
- Warr WB (1966) Fiber degeneration following lesions in the anterior ventral cochlear nucleus of the cat. *Experimental Neurology* **14**:453–474.
- Warr WB (1969) Fiber degeneration following lesions in the posteroventral cochlear nucleus of the cat. *Experimental Neurology* **23**:140–155.
- Warr WB (1982) Parallel ascending pathways from the cochlear nucleus: neuroanatomical evidence of functional specialization. In: Neff WD (ed). *Contributions to Sensory Physiology*. Academic Press, New York, pp. 1–38.
- Warr WB (1992) Organization of olivocochlear efferent systems in mammals. In: Webster DB, Popper AN, and Fay RR (eds). *Springer Handbook of Auditory Research*, Volume 1: *The Mammalian Auditory Pathway: Neuroanatomy*. Springer-Verlag, New York, pp. 410–448.
- Webster DB, Fay RR, and Popper AN (eds) (1992) *The Evolutionary Biology of Hearing*. Springer-Verlag, New York.
- Webster WR and Garey LJ (1990) Auditory system. In: Paxinos G (ed). *The Human Nervous System*. Academic Press, San Diego, pp. 899–944.
- Weedman DL and Ryugo DK (1996a) Pyramidal cells in primary auditory cortex project to cochlear nucleus in rat. *Brain Research* **706**:97–102.
- Weedman DL and Ryugo DK (1996b) Projections from auditory cortex to the cochlear nucleus in rats: synapses on granule cell dendrites. *Journal of Comparative Neurology* **371**:311–324.
- Weinberg RJ (1997) Are topographic maps fundamental to sensory processing? *Brain Research Bulletin* **44**:113–116.
- Weinberger NM (1997) Learning-induced receptive field plasticity in the auditory cortex. *Seminars in Neuroscience* **9**:59–67.
- Wenstrup JJ, Ross LS, and Pollak GD (1985) A functional organization of binaural responses in the inferior colliculus. *Hearing Research* **17**:191–195.
- Wenstrup JJ, Ross LS, and Pollak GD (1986) Binaural response organization within a frequency-band representation of the inferior colliculus: implications for sound localization. *Journal of Neuroscience* **6**:962–973.
- Wenthold RJ (1985) Glutamate and aspartate as neurotransmitters of the auditory nerve. In: Drescher DG (ed). *Auditory Biochemistry*. Charles C Thomas, Springfield, pp. 125–140.
- Wenthold RJ and Hunter C (1990) Immunocytochemistry of glycine and glycine receptors in the central nervous system. In: Ottersen OP and Storm-Mathisen J (eds). *Glycine Neurotransmission*. John Wiley & Sons, Chichester, pp. 391–416.
- Wenthold RJ, Huie D, Altschuler RA, and Reeks KA (1987) Glycine immunoreactivity localized in the cochlear nucleus and superior olivary complex. *Neuroscience* **22**:897–912.
- White EL and Keller A (1989) *Cortical Circuits. Synaptic Organization of the Cerebral Cortex. Structure, Function, and Theory*. Birkhäuser, Boston.
- Wickesberg RE and Oertel D (1990) Delayed, frequency-specific inhibition in the cochlear nuclei of mice: a mechanism for monaural echo suppression. *Journal of Neuroscience* **10**:1762–1768.

- Winer JA (1991) Anatomy of the medial geniculate body. In: Altschuler RA, Bobbin RP, Clopton BM, and Hoffman DW (eds). *Neurobiology of Hearing: The Central Auditory System*. Raven Press, New York, pp. 293–333.
- Winer JA (1992) The functional architecture of the medial geniculate body and the primary auditory cortex. In: Webster DB, Popper AN, and Fay RR (eds). *Springer Handbook of Auditory Research*, Volume 1: *The Mammalian Auditory Pathway: Neuroanatomy*. Springer-Verlag, New York, pp. 222–409.
- Winer JA and Larue DT (1996) Evolution of GABAergic circuitry in the mammalian medial geniculate body. *Proceedings of the National Academy of Sciences of the United States of America* **93**:3083–3087.
- Winer JA and Morest DK (1983) The neuronal architecture of the dorsal division of the medial geniculate body of the cat. A study with the rapid Golgi method. *Journal of Comparative Neurology* **221**:1–30.
- Winer JA, Diehl JJ, and Larue DT (2001) Projections of auditory cortex to the medial geniculate body of the cat. *Journal of Comparative Neurology* **430**:27–55.
- Winer JA, Larue DT, and Huang CL (1999) Two systems of giant axon terminals in the cat medial geniculate body: convergence of cortical and GABAergic inputs. *Journal of Comparative Neurology* **413**:181–197.
- Winer JA, Larue DT, and Pollak GD (1995) GABA and glycine in the central auditory system of the mustache bat: structural substrates for inhibitory neuronal organization. *Journal of Comparative Neurology* **355**:317–353.
- Winer JA, Larue DT, Diehl JJ, and Hefti BJ (1998) Auditory cortical projections to the cat inferior colliculus. *Journal of Comparative Neurology* **400**:147–174.
- Winer JA, Saint Marie RL, Larue DT, and Oliver DL (1996) GABAergic feedforward projections from the inferior colliculus to the medial geniculate body. *Proceedings of the National Academy of Sciences of the United States of America* **93**:8005–8010.
- Wong D (1991) Cellular organization of the cat's auditory cortex. In: Altschuler RA, Bobbin RP, Clopton BM, and Hoffman DW (eds). *Neurobiology of Hearing: The Central Auditory System*. Raven Press, New York, pp. 367–387.
- Wong D and Kelly JP (1981) Differentially projecting cells in individual layers of the auditory cortex: a double-labeling study. *Brain Research* **230**:362–366.
- Yan J and Suga N (1996) Corticofugal modulation of time-domain processing of biosonar information in bats. *Science* **273**:1100–1103.
- Yang L, Liu Q, and Pollak GD (1996) Afferent connections to the dorsal nucleus of the lateral lemniscus of the mustache bat: evidence for two functional subdivisions. *Journal of Comparative Neurology* **373**:575–592.
- Yeomans JS and Frankland PW (1996) The acoustic startle reflex: neurons and connections. *Brain Research Reviews* **21**:301–314.
- Yin TCT and Chan JCK (1990) Interaural time sensitivity in medial superior olive of cat. *Journal of Neurophysiology* **64**:465–488.
- Young ED (1984) Response characteristics of neurons of the cochlear nuclei. In: Berlin CI (ed). *Hearing Science, Recent Advances*. College-Hill Press, San Diego, pp. 423–460.
- Young ED and Voigt HF (1982) Response properties of type II and type III units in dorsal cochlear nucleus. *Hearing Research* **6**:159–182.
- Young ED, Nelken I, and Conley RA (1995) Somatosensory effects on neurons in dorsal cochlear nucleus. *Journal of Neurophysiology* **73**:743–765.
- Zhang Y, Suga N, and Yan J (1997) Corticofugal modulation of frequency processing in bat auditory system. *Nature* **387**:900–903.



# Chapter 2

## Neuronal Organization in the Inferior Colliculus

DOUGLAS L. OLIVER

### 1. KEY ISSUES REGARDING INFERIOR COLLICULUS NEURONAL ORGANIZATION

An understanding of the neuronal organization of the inferior colliculus (IC) requires an exploration of how the types of neurons, the microcircuitry, and the synaptic organization of the IC interact to define functional zones. The IC was originally divided using anatomical methods to identify the neurons and their inputs with the hope that these subdivisions would correspond to functional zones. The central nucleus, the largest of the subdivisions, has been the focus of most studies of IC neuronal organization and its neuron types and the inputs are best known. Understanding the neuronal organization of the IC in terms of subdivisions has been a problem when it is uncertain how other parts differ from the central nucleus. Because IC subdivisions must be related to the constituent neurons and their inputs, the first part of this chapter discusses how the central nucleus and surrounding structures differ in this regard. One way to address the possible functional differences is through the study of IC microanatomy, our second topic. Within the central nucleus the microanatomical arrangements suggest that the inputs may create modules with specific combinations of inputs. Because a module may contain neurons performing similar types of processing, the types of functional zones present in each subdivision will define the function of that subdivision and clarify the differences between it and others. A third aspect of neuronal organization is the definition of the neuron type. Neurons have a functional as well as a morphological profile. Hence, neurons defined by their axonal targets, neurotransmitter content, and intrinsic membrane properties should reveal fundamental aspects of IC neuronal organization. The final facet of neuronal organization to be discussed is the synaptic organization—the types of synapses and patterns of synaptic input that are manifest on different neuron types or in different functional modules. Synaptic organization leads to the final question regarding neuronal organization. How is auditory processing influenced by the interaction of synaptic inputs with the intrinsic properties of the IC neurons?

## 2. THE PROBLEM OF SUBDIVIDING THE INFERIOR COLLICULUS

The IC is one of the largest structures in the mammalian midbrain. It forms the posterior half of the corpora quadrigemina in the midbrain tectum: four bulges on the dorsal surface of the midbrain (Fig. 2.1). Grossly, the IC is readily distinguished from the superior colliculus that constitutes the anterior half of the tectum. Likewise, it is distinguished from the midbrain tegmentum readily in frontal or transverse sections. Tectum and tegmentum develop from different stem cells (Senut and Alvarado-Mallart 1987). The tectum is an alar (sensory) plate derivative whereas the tegmentum is a basal plate derivative, which suggests a motor function.

Despite the ease with which the IC is distinguished from other midbrain structures, understanding the internal or functional organization of the IC has not been so obvious. Like many other brain structures, sharply defined internal boundaries are not always evident.

### *2.1. THE CENTRAL NUCLEUS AND A SURROUNDING CORTEX ARE THE MAJOR INFERIOR COLLICULUS SUBDIVISIONS*


The simplest parcellation recognizes a central nucleus (ICC; Fig. 2.2) that is surrounded by a cortex. Most historical accounts as well as most modern studies follow this basic design. For example, Ramón y Cajal (1995) referred to a nucleus of the IC covered by a dorsal cortex (DC; Fig. 2.2) that includes the commissural connections. He described a lateral cortex (LC; Fig. 2.2) which is a thin layer of gray matter beside ICC that is covered by the brachium of the IC (BI; Fig. 2.2). The brachium contains axons that project to the medial geniculate body in the diencephalon and descending fibers to the IC from the neocortex. Although the name “central nucleus” has been in use for many years, the precise IC region associated with it has depended on the methodological approach and its interpretation.

The anatomical subdivisions of the cat (Oliver and Shneiderman 1991; Oliver and Huerta 1992) and human (Waitzman and Oliver 2002) IC have been reviewed and the references cited include a discussion of adjoining tegmental structures. Here, we address the IC structure from a comparative perspective and evaluate different views on the organization of the subdivisions.

### *2.2. MIDBRAIN SUBDIVISIONS REFLECT THE METHODS USED TO IDENTIFY THEM*

Neuroanatomical studies have traditionally relied on Nissl stains or myelin stains to identify subdivisions (Figs. 2.2 to 2.4). These stains are complementary, as Nissl stains show cell bodies while myelin stains reveal axons, and increased

Figure 2.1. Gross anatomy of the cat (A) and rat (B) midbrain. *AI*, Primary auditory cortex; *BI*, brachium of the inferior colliculus; *CN*, cochlear nucleus; *IC*, inferior colliculus; *LL*, lateral lemniscus; *MG*, medial geniculate body; *SC*, superior colliculus. The scale bar represents 2 mm.



axon density usually reflects lower neuronal cell body density (Berman 1968; Olszewski and Baxter 1982; Paxinos and Watson 1998). Nissl and myelin stains, although widely used, can only hint at the functional subdivisions and internal complexity in the IC.

Golgi impregnations reveal details of IC neurons that are not discernible by other methods. They reveal more distinguishing features of unique neuron types, local differences in the structure of the neuronal distribution, and the differences in local dendritic neuropil organization. Golgi studies have been essential to identifying more specialized regions in the IC (Rockel and Jones 1973a,b; Fitzpatrick 1975; Oliver 1984b; Morest and Oliver 1984; Faye-Lund and Osen 1985; Meininger et al. 1986; Ramón y Cajal 1995). Unfortunately, Golgi stains are incompatible with many histological approaches. Thus, more refined methods that identify morphologically distinct cell types must be used to recognize subdivisions and to confirm their identity independently.

Molecular methods may ultimately identify subdivisions. For example, parvalbumin, a calcium binding protein, has a high concentration in ICC whereas calbindin and calretinin, other calcium binding proteins, concentrate in the dorsal cortex (ICD; see Section 2.6). Cytochrome oxidase, a metabolic marker, concentrates in the ICC (Dezso et al. 1993; González-Lima and Cada 1994; González-Lima et al. 1997; Poremba et al. 1997) rather than in the ICD.





Figure 2.2. Nissl cytoarchitecture of the cat IC in the Horsley–Clarke stereotaxic plane. The *inset* shows the planes of section. (A) Mid-IC level through the DNLL. (B) A more rostral section at the level of the rostral pole nucleus (RP). BA, Nucleus of the brachium; BI, brachium of the inferior colliculus; CG, central gray; CM, commissural, commissure; CU, cuneiform nucleus; DC, dorsal cortex; DI, dorsal intercollicular tegmentum; DNLL, dorsal nucleus of the lateral lemniscus; ICC, central nucleus; LC, lateral cortex; LI, lateral intercollicular tegmentum; MI, medial intercollicular tegmentum; SC, superior colliculus; 1–4, layers of the dorsal cortex; VL, ventral lateral nucleus. The scale bar represents 1 mm.

Figure 2.3. Cat IC in the anatomical transverse plane of section. The *inset* shows the plane of #2. (A) Nissl stain. (B) Heidenhain fiber stain. The scale bar represents 1 mm.



Subdivisions can also be identified electrophysiologically. Microelectrode studies showed that ICC neurons had low thresholds and vigorous responses to simple acoustic stimuli, and their spectral properties and response latencies often distinguished them from ICD (see Chapter 11). ICC neurons have the sharpest tuning and shortest latencies (Aitkin et al. 1975), suggesting its functional primacy.

However, the presence of a tonotopic map does not distinguish the ICC from the surrounding cortex. In other brain regions there is often a reversal in the sequence of frequencies at a subdivision border. Such a reversal is found at the border of the ICC and lateral cortex but *not* at the border of the ICC and ICD or caudal cortex. This has frustrated any sharp distinction between the ICC and dorsal cortex in electrophysiological studies, as the best frequency of a neuron cannot conveniently determine electrode position relative to the ICC border. Electrophysiological methods must often be complemented by other independent methods such as post hoc histology to identify recording sites in smaller subdivisions.

Two methods combine elements of histological approaches and acoustic stimulation and may distinguish subdivisions. The first uses the accumulation of 2-deoxyglucose (2DG) in presynaptic endings during synaptic activity (Nudo and Masterton 1986). A second method is the detection of the Fos protein, a nuclear transcription factor, or its mRNA (Sheng et al. 1993; Fields et al. 1997). Both approaches can reveal activity evoked by acoustic stimuli in presynaptic endings (2DG) or in single neurons (Fos). The challenge in using these methods to identify IC subdivisions is the choice of an appropriate stimulus. One approach uses different spectral, temporal, and binaural features of acoustic stimuli to evoke differential responses in the ICC and surrounding structures.




Figure 2.4. Rat IC. (A) Caudal central nucleus. 1–3, layers of the lateral cortex. (B) Rostral central nucleus. (C) Myelin stain at the level of rostral pole. (D) Nissl preparation. The scale bar represents 1 mm.

### 2.3. HOW IS THE CENTRAL NUCLEUS DEFINED?

The main subdivision of the mammalian IC is the central nucleus (Figs. 2.2 to 2.4). Its main feature is the fibrodendritic lamina, an entity comprised of disc-shaped neurons and the laminar plexus of afferent axons terminating in it.

#### 2.3.1. Morphology of Central Nucleus Neurons

The main feature of ICC neurons is their highly oriented dendritic field. These disc-shaped neurons are by far the most common cell type and have been observed in almost every species studied. Similar neurons have been described in Golgi preparations of the cat (Oliver 1984b), mouse (Meininger et al. 1986), bat

(Zook et al. 1985), rat (Faye-Lund and Osen 1985; Malmierca et al. 1993. Reprinted with permission of Wiley-Liss, Inc., a subsidiary of John Wiley & Sons, Inc.), and human (Geniec and Morest 1971) and after intracellular injections of horseradish peroxidase (Kuwada et al. 1997b; Oliver et al. 1991). In the cat most ICC neurons are disc-shaped and have dendritic fields parallel to one another (Figs. 2.5A and 2.6). This arrangement imparts a distinct appearance in Golgi stains that distinguishes the ICC. Disc-shaped neurons have dendritic



Figure 2.5. Disc-shaped (**A**) and stellate (**B**) cell types in the ICC labeled by intracellular injections of horseradish peroxidase in the adult cat. The approximate position of the cells is shown in the *inset* (closed circle). Laminar orientation is shown by arrow in (**A**) and by lines in (**B**). *a*, axon. The scale bar represents 100  $\mu\text{m}$ . (From Oliver et al. 1991.)




Figure 2.6. Two disc-shaped cells in the ICC of the cat after intracellular injection of horseradish peroxidase. (A, B) The cells are shown *en face* in the original parasagittal plane of section. (C, D) The same cells after 90° rotation to show the most highly oriented edge-on view of the dendritic field. (E) Location of the neuron. (From Kuwada et al. 1997.)

fields about 50 to 70  $\mu\text{m}$  in diameter at the narrowest dimension although the longest axis may be a millimeter (Fig. 2.6). These dimensions may be species-specific. Disc-shaped cells form rows with cells stacked end-to-end or overlapping with long axes of the dendritic fields in parallel. In the rat the high-frequency part of ICC has layers where most disc-shaped neurons, called flat cells in that species, alternate with the less oriented less-flat cells (an oriented



stellate neuron is 100  $\mu\text{m}$  in diameter) (Malmierca et al. 1993). Here, the laminar unit is proposed to be one flat cell and one less-flat neuron, yielding a laminar thickness of about 150 to 170  $\mu\text{m}$  (Fig. 2.7). In the lower-frequency part in the rat and throughout the cat ICC, alternating cell types have not been observed and the laminar unit is likely a multiple of the diameter of the disc-shaped dendritic field. In Nissl stains the disc-shaped morphology is not apparent, as




Figure 2.7. Alternating flat and less-flat cells create laminae in the rat central nucleus. (A) The *en face* view shows less-flat cells (*top*) and flat cells (*bottom*) with different-sized dendritic fields. (B) The on edge view shows the narrowest dimension of the neurons after 90° rotation. (From Malmierca et al. 1993.)

the dendrites are invisible but the somatic orientation is often still evident. Most ICC neurons have dispersed Nissl bodies (granular endoplasmic reticulum) and a smooth nuclear envelope, and presumably correspond to disc-shaped cells (Oliver 1984b; Ribak and Roberts 1986).

A second major cell type defined by morphology in ICC is the stellate (cat; Fig. 2.5B) or less-flat (rat; Fig. 2.7) neurons, which represent <25% of the population and often have radiating dendritic fields that are spherical. Dendrites typically extend beyond the single fibrodendritic lamina into adjacent laminae. Other cell types have ovoid dendritic fields oriented perpendicular to the long axis of the disc-shaped cells (cat) or parallel to the flat cells (the less-flat cells in rat). Nissl stains reveal that presumptive stellate cells have more Nissl bodies (stacks of granular endoplasmic reticulum) and an irregular or infolded nuclear envelope.

The identification of functionally relevant neuron types remains a challenge to research in this field. There are probably more than two types of ICC cells distinguished only on anatomical grounds. They can also be discriminated by size (disc-shaped cells are large to medium or small) and by the complexity of the dendritic branching (stellate cells have simple or complex dendritic branching patterns) (Malmierca et al. 1993). The incidence of dendritic spines (Paloff et al. 1992) and the number and types of axosomatic synaptic input may vary (Oliver 1984b; Ribak and Roberts 1986; Paloff et al. 1989). Observations on the neurotransmitter content, axonal targeting, and intrinsic membrane properties (see Section 4) support the notion that the definition of functional cell types must include data beyond classical morphological models.


### 2.3.2. Synaptic Inputs to the Central Nucleus

Inputs to the ICC ascend from neurons in the lower auditory brain stem, including the cochlear nuclear complex consisting of the dorsal (DCN; Figs. 2.8 and 2.9) anteroventral (AVCN) and posteroventral (PVCN) cochlear nuclei (see



Figure 2.8. Projections from cat dorsal cochlear nucleus to the inferior colliculus. **(A)** Injection of biotinylated dextran amine in dorsal cochlear nucleus. **(B)** A single layer of dextran-labeled axons in the central nucleus of the inferior colliculus. **(C)** Higher magnification view of labeling. The scale represents 1 mm in **(A, B)**; the scale bar represents 0.5 mm in **(C)**. (From Oliver et al. 1997.)

Figure 2.9. Projection from dorsal cochlear nucleus (*DCN*) to the inferior colliculus in the rat. **(A)** Band of labeled axons in the ICC (*arrow*). The lateral band of axons in the external (lateral) cortex is orthogonal to the medial band. **(B)** Injection of dextran in the 4.5 kHz region of dorsal cochlear nucleus. The scale bar represents 0.5 mm. (From Malmierca et al. 2002.)



Chapter 3), and from the superior olivary complex, which includes the medial superior olive (MSO; Fig. 2.10) and the lateral superior olivary nuclei (LSO; Chapter 4). Smaller inputs originate in the medial nucleus of the trapezoid body and from periolivary nuclei, including the superior paraolivary nucleus, and the lateral and ventromedial nuclei of the trapezoid body. The lateral lemniscal nuclei (dorsal and ventral nuclei; DNLL and VNLL, respectively) also provide many afferent axons to ICC (Brunso-Bechtold et al. 1981; Oliver and Shneiderman 1991) and represent the largest single source of input fibers. These projections have been described in small mammals (Merchán et al. 1994; Merchán and Berbel 1996; Malmierca et al. 1998).

Excitatory inputs to the ICC outnumber inhibitory inputs. Electron microscopic autoradiographic studies have identified the synaptic endings from different brain stem sources. These afferent endings are defined as excitatory inputs by the morphology of their synapses—their endings contain clear, round synaptic vesicles (R-type) that make asymmetrical synaptic contacts. It is possible to determine the relative prevalence of different types of axonal boutons from different sources. The MSO provides the largest single source of excitatory synaptic inputs (37%) in the cat followed by the ipsilateral LSO (26%), the contralateral LSO (18%) (Oliver et al. 1995), and the cochlear nucleus (DCN: 11%; AVCN: 13% to 18%) (Oliver 1984a, 1985, 1987). The sum of R-type endings from all sources is about 60% of the ICC axonal endings.

Inhibitory inputs are substantial and are a remarkable feature of ICC neuronal organization. Synaptic terminals that contain  $\gamma$ -aminobutyric acid (GABA) or




Figure 2.10. Medial superior olivary projections in the cat. (A) Plots of dextran-labeled axons terminating in the most dorsolateral ICC laminae corresponding to the lowest frequency representation. A few fibers cross the border into the dorsal cortex. Others may terminate in the most dorsal part of the ventrolateral nucleus. (B) Injection of dextran amines in the dorsal 150 Hz region of the medial superior olive. (C) Adjacent Nissl-stained section. (From Loftus et al. 2004.)

glycine as a neurotransmitter comprise 40% of the presynaptic endings (Oliver 2000). Electron micrographic studies find many endings with the morphology associated with inhibitory synapses—pleomorphic synaptic vesicles (PL-type) that make symmetrical synaptic contacts (Shneiderman and Oliver 1989). Many of these contain GABA (Shneiderman et al. 1993; Oliver et al. 1994) and originate in the DNLL (Hutson 1988; Shneiderman et al. 1988; Zhang et al. 1998; Chen et al. 1999) and they account for 26% of the PL-type endings (Shneiderman and Oliver 1989). A second source of inhibitory endings is the ipsilateral LSO projection, which contains glycine (Saint Marie et al. 1989; Saint Marie and Baker 1990; Glendenning et al. 1992) and also accounts for 26% of ICC endings (Oliver et al. 1995). A third and probably inhibitory projection is the VNLL, whose neurons contain both glycine and GABA (Saint Marie et al. 1997; Riquelme et al. 2001). These endings have not been identified at the electron microscopic level nor have the local axons from IC GABAergic neurons (see Section 4.2).

Although the inputs to ICC have been identified at the electron microscopic level, their synaptic role in processing auditory information remains an area of intense interest. For example, it is unclear whether these inputs are distributed homogeneously. Further evidence on ICC microanatomy suggests that inputs are segregated and raises the possibility that excitatory and inhibitory inputs have different densities in different functional modules (see Section 3.2.2). Likewise,

as the definitions of cell types expand to include transmitter content and intrinsic properties, it will be important to determine whether different cell types receive the same proportions of excitatory and inhibitory input.

### 2.3.3. Axonal Components of the Fibrodendritic Laminae

The major axonal components of the fibrodendritic laminae are lemniscal axons. Many axons in Golgi preparations run parallel to the dendrites of the disc-shaped neurons but the identity of these axons cannot be determined. In experimental material the contribution of the ascending fibers to the fibrodendritic laminae can be determined. Axons from lower brain stem sources form ventrolateral-to-dorsomedial oriented rows parallel to the dendritic fields of the disc-shaped neurons. Small injections of clusters of neurons in auditory brain stem structures with anterograde tracers (biotinylated dextran amines, [<sup>3</sup>H]leucine) reveal axonal bundles that run from the caudal to rostral direction through much of the ICC (Figs. 2.8 to 2.10). In essence a single band of axons represents the narrowest laminar dimension, about 200  $\mu\text{m}$  wide. In the cat the axonal laminae can extend 4 mm rostrocaudally and reach 2 mm in height (Fig. 2.11). These run from caudolateral-to-rostromedial at approximately 45° to the rostrocaudal plane and from ventrolateral to dorsomedial in the dorsoventral plane. The rostrocaudal laminar length may be species-specific. However, the width of single bands of axons from the DCN (compare Figs. 2.8 and 2.9) is similar in cat and rat and resembles that for AVCN and for cat LSO and MSO axons (Fig. 2.10).

A third component of the fibrodendritic laminae are axons from local IC neurons (see Chapter 5). Axons from both disc-shaped and stellate cells form local collaterals before leaving the IC. Disc-shaped cells injected with intracellular deposits of horseradish peroxidase (HRP) have axons that remain in the same lamina as the parent cell and the local axonal plexus is parallel to the dendritic field (Fig. 2.5A) (Oliver et al. 1991). IC neurons may also project to other ipsilateral laminae. Stellate neuron axons in Golgi impregnations in the cat or after intracellular filling in the rat contribute to fibrodendritic laminae other than the one in which the cell body resides (Fig. 2.12A). Thus, ICC neurons of all types have axons that contribute to ICC laminar organization. An important unresolved issue here is whether the ICC contains true interneurons with axons that are confined solely to the ipsilateral IC (see Chapter 22). Although there is evidence for local axonal collaterals, it is more difficult to prove that the axon remains within the ICC. No neuron with a unique morphology has been identified as a true interneuron and the evidence from HRP intracellular filling suggests that all neurons have axons with local collaterals, even those whose targets are outside of the IC or in the contralateral IC (Oliver et al. 1991).

A fourth component of the fibrodendritic lamina is commissural fibers (see Chapter 5). Small injections of *Phaseolus vulgaris* leucoagglutinin (PHA-L) or biotinylated dextran amines (BDA) on one side of the IC can label single contralateral laminae (Saldaña and Merchán 1992; Malmierca et al. 1995). The contralateral laminar labeling is at the same best frequency as the injected lam-

ina. Thus, the homotypical laminae on each side of the central nucleus appear to be interconnected.

#### 2.4. THE LATERAL CORTEX

The lateral cortex is lateral to the central nucleus (LC; Fig. 2.2A) and has also been called the external nucleus (Berman 1968), external cortex (Faye-Lund and Osen 1985), and lateral zone (Geniec and Morest 1971). Part of it may receive input from the lateral lemniscal nuclei, but it lacks disc-shaped cells. In the cat and human the LC includes the lateral nucleus and the ventrolateral nucleus (Geniec and Morest 1971; Morest and Oliver 1984).

The LC has a fibrous outer layer 1 and a small-celled layer 2. In the cat it is easily defined relative to the low-frequency parts of ICC by its lower neuronal density (Fig. 2.2A: LC). The LC is also apparent in myelin stains because the fibers that become the brachium of the IC aggregate and the outer fibrous layer thickens rostrally (Fig. 2.3). Layer 2 of the LC is apparent in many species because of high acetylcholinesterase concentrations (rat: Paxinos and Watson 1998). The main inputs to the lateral nucleus are from the ipsilateral ICC, the auditory cortex, the spinal cord, and dorsal column nuclei of the somatic sensory system (Morest and Oliver 1984; Oliver and Huerta 1992). These terminate primarily in LC layer 2 and lateral lemniscus fibers do not enter layers 1 or 2 (Oliver et al. 1999) (see below).

In the cat and human the lateral cortex includes the ventrolateral nucleus (Figs. 2.2 and 2.3: VL) wedged between the ventral part of the lateral nucleus and the fibers of the lateral lemniscus as they enter the ventral central nucleus (also called the interstitial zone). VL has large and small cells and a cellular density intermediate to that in the central or lateral nuclei. Thus, it can appear as a third layer beneath the small-celled second layer of the lateral nucleus in the ventral IC. Rostrally, VL is usually segregated from ICC by the lateral lemniscal axons. More caudally, however, lemniscal fibers are fewer and the separation is less obvious. The cat VL receives lemniscal inputs from the cochlear nucleus and the MSO and LSO (Shneiderman and Henkel 1987). These are laminar projections. When labeled experimentally they form a "medial band" of axons in ICC parallel to the disc-shaped neurons, and a shorter "lateral band" in VL oriented orthogonal to those in ICC (Shneiderman and Henkel 1987).

In the rat the LC (Fig. 2.4A, B: LC) is called the external cortex (Faye-Lund and Osen 1985; Paxinos and Watson 1998). The many studies of rats make it useful to compare this region to that in the cat. The external cortex is a three-layered structure. Layers 1 and 2 resemble the cat LC, the fibrous layer and the small cellular layer, respectively. Layer 3 of rat LC corresponds to the ventrolateral nucleus. Studies of both the local connections and the lemniscal inputs to the IC in rat show laminated input to both the ICC and to layer 3 of the external cortex (see Chapter 5). As in the cat ICC and VL, the medial band of lemniscal fibers terminates parallel to the ICC laminae and the lateral band of fibers in the external cortex is nearly orthogonal (Fig. 2.9). Single lateral bands

are as wide as the medial bands. The lemniscal inputs to the external cortex are tonotopically organized with lower frequency bands located dorsally (Loftus et al. 2004b).

One explanation for the different appearances of the IC in the rat and cat is the absence of laminae in rat ICC and lateral cortex tuned to frequencies  $<1$  kHz and the attenuation of laminae tuned to 1 to 5 kHz. In the rat, the ICC is separated from the lateral nucleus by the VL (layer 3 of LC). In contrast, in the cat the frequencies  $<5$  kHz and especially  $<1$  kHz occupy much of the dorso-lateral ICC. The portions of the IC lateral to this low-frequency ICC region have little or no ventrolateral nucleus intercalated between it and the lateral nucleus, and lateral bands of lemniscal axons tuned to frequencies  $<2$  kHz diminish. Because in the rat few IC responses are  $<3$  kHz (Clopton and Winfield 1973), the comparable low-frequency part of ICC is presumably absent or attenuated, perhaps explaining the species difference. A similar effect may be present in mice, whose low-frequency hearing is likewise minimal.

Questions remain concerning the LC particularly. Not least among these is the persistent use of different names for what may be the same structure. We propose that the term “lateral cortex” be adopted as a substitute for external cortex, as it clearly denotes this lateral portion of the IC, it distinguishes it from other rostral, medial, and dorsal regions, and it identifies its cortical affiliations.

The function of the LC is uncertain. This may be because observations from the deeper portion (VL nucleus, layer 3), which receives lemniscal inputs, are often combined with data from the superficial portion, which receives little ascending input but is a major target of ICC. Electrophysiological studies have not characterized these areas systematically and consequently there is little basis on which to compare their neuronal response properties. One possible function, suggested below, is that the lateral nucleus of the LC extends rostrally as the nucleus of the brachium of the IC and this amalgamated structure has a role in multimodal integration (Malmierca et al. 2002).

## 2.5. STRUCTURES ROSTRAL TO THE CENTRAL NUCLEUS

Rostral to the ICC the laminae disappear and the nucleus of the rostral pole (RP; Figs. 2.2B and 2.4C, D) emerges. In many species the RP receives the most anterior lemniscal fibers and these terminate on stellate neurons. The RP is evidently related to auditory function owing to its lemniscal inputs and it projects to the superior colliculus, providing auditory signals to the visual–motor system (Harting and Van Lieshout 2000). However, few studies have considered the responses of these neurons to sound stimuli.

The intercollicular tegmentum surrounds the RP (Figs. 2.2B and 2.4C, D) and it is named according to its location in the dorsal (DI), medial (MI), or lateral (LI) intercollicular tegmentum (Morest and Oliver 1984). These complex tegmental structures are the part of the mesencephalic reticular formation that separates the IC from the superior colliculus. The borders of ICC, RP and tegmental structures are best seen in sagittal or horizontal planes where the fibrous teg-

mental architecture contrasts with the denser cell packing of the ICC. The functions of the intercollicular tegmentum at the level of the IC remain obscure, but it may be involved with multimodal integration. Like the second layer of LC, the intercollicular tegmentum receives inputs from the somatosensory system (RoBards et al. 1976; RoBards 1979; Wiberg et al. 1987), from the auditory nerve root (Lopez et al. 1999), and more rostral portions of the mesencephalic reticular formation that are involved in visual-motor function (Waitzman and Oliver 2002). The intercollicular tegmentum may therefore be involved with multimodal integration.

The nucleus of the brachium of the IC (BI; Figs. 2.2B and 2.4C, D) may be involved in gaze control. Whereas the brachium of the IC (BI; Figs. 2.2B, 2.4C, D) contains the tectothalamic fibers from the IC and corticotectal fibers from the neocortex (Kudo and Niimi 1980; Winer et al. 1998), the BI is a cellular structure that receives inputs primarily from the IC. It is continuous caudally with the LC. The nucleus of the brachium projects in a reciprocal topographical manner to the superior colliculus (King et al. 1998; Doubell et al. 2000). The BI may also serve as an interface for routing auditory information from the ICC to the superior colliculus. Because of this role and its position in the auditory pathway the nucleus of the brachium (perhaps including the lateral nucleus) may be homologous to the “external cortex” as defined in the barn owl, where convergent inputs from ICC create a spatial map of sound location that is conveyed to the optic tectum (Knudsen 1983a,b). The nature of the auditory signals sent by BI to the superior colliculus in the mammal is unknown.

## 2.6. STRUCTURES DORSAL AND CAUDAL TO THE CENTRAL NUCLEUS

Dorsal to the ICC is the dorsal cortex (DC; Figs. 2.2A, 2.3, and 2.4B, C). Earlier studies (Berman 1968; Rockel and Jones 1973b) recognized a “pericentral nucleus” that is largely confined to the outer margin of the IC and resembles the LC in that it has a fibrous outer layer and a parvocellular inner layer. These layers correspond to layers 1 and 2 of the cat’s DC (Morest and Oliver 1984). The deeper layers of DC (layers 3 and 4; Figs. 2.2A and 2.3) were considered the dorsomedial part of ICC in some prior accounts (Rockel and Jones 1973b). Layer 3 is continuous with the fibers of the commissure of the IC and the deeper layers have successively larger cells. The caudal cortex is limited to the superficial 200 to 300  $\mu\text{m}$  IC caudal to ICC (Morest and Oliver 1984; Oliver and Morest 1984). Much like the pericentral nucleus, it has two outer layers but the most superficial fibrous layer is less prominent. Where it is caudal to the DC in the dorsomedial IC, layer 2 of the caudal cortex borders layer 3 of the DC.

Golgi studies have shown that the deeper layers of DC continue beneath the pericentral region until it meets the ICC. Their border is clear, as the orientation of the dendritic fields shows disc-shaped neurons of the fibrodendritic laminae in the ICC ending at the border of the DC, where nonoriented neurons become prominent (see Fig. 11 in Morest and Oliver 1984). Fiber stains and anterograde



transport studies reveal the transition from the ICC, as the axonal populations change here. For example, DC neurons have unoriented axons while disc-shaped ICC neurons have axons confined to a lamina (Oliver et al. 1991). Inputs to ICC are primarily from the brain stem with minor (or no) inputs from the neocortex and a modest input from the contralateral IC (see earlier). This pattern is reversed in the deepest layer of dorsal cortex (layer 4), where the neocortical and commissural inputs are predominant (Chapters 5 and 8). Only modest inputs from the DCN and the DNLL reach DC layer 4 (Oliver 1984a; Shneiderman et al. 1988). These fibers are often smaller collaterals of the lemniscal axons whose principal terminal fields are in the ICC. Input to the DC from the SOC is absent.

Defining the dorsal border of ICC experimentally remains problematic and there is a question as to whether there is a sharp functional border. As noted earlier, there is no reversal of the tonotopic map at this border. However, changes in the populations of inputs and possibly in the intrinsic properties of the neurons (Smith 1992; Li et al. 1998) suggest that the response properties of neurons in these regions differ. For example, DC neurons have broader tuning than ICC neurons (Aitkin et al. 1975, 1994).

Certain histochemical stains and immunocytochemical probes may be useful to identify this border. In many species a high parvalbumin concentration delineates the ICC from the DC and the latter has high concentrations of calbindin and calretinin. Comparisons are available for human, gerbil, bat, guinea pig, macaque, dolphin, rat, and mouse (Seto-Ohshima et al. 1990; Ohshima et al. 1991; Coleman et al. 1992; Vater and Braun 1994; Yasuhara et al. 1994; Caicedo et al. 1996; Lohmann and Friauf 1996; Glezer et al. 1998; Idrizbegovic et al. 1999; Spencer et al. 2002; Tardif et al. 2003). Molecules such as cytochrome oxidase also have high concentrations in the ICC (Dezso et al. 1993; González-Lima and Cada 1994; González-Lima et al. 1997; Poremba et al. 1997) and NADPH-diaphorase is concentrated in the dorsal and lateral cortex (Paxinos 1999). These latter proteins may be expressed in the presynaptic lemniscal axons in the IC and these axons may not all terminate precisely at the ICC border, suggesting a concentration gradient at the DC border.

## *2.7. FUTURE RESEARCH ON DEFINING FUNCTIONAL SUBDIVISIONS IN THE INFERIOR COLLICULUS*

Other endogenous proteins may better serve as architectonic guides to IC parcellation. The premise that IC subdivisions represent homogeneous functional zones based on either unique neuron types or fixed combinations of inputs suggests that cell-specific molecules or inputs may identify the subdivisions. At this point, however, single IC subdivisions remain to be identified by single molecules. Moreover, the premise that the identity of an IC subdivision is represented as a single functional module is tenuous.

A complementary approach that deserves attention is the molecular specificity of different IC cell types or cells of different lineage. Thus, if DC neurons originate from different stem cells or develop at different times than those in

the ICC it could explain why the dendrites of their neurons differentiate later (Morest 1969). Because ICC and DC neurons often differ in dendritic morphology, perhaps there are molecules related to this phenotype that would demarcate IC subdivisions. Different IC neurons have different patterns of ion channel expression, as suggested by firing patterns related to unique combinations of potassium and calcium currents (Sivaramakrishnan and Oliver 2001). Thus, probes for ion channel subunits or combinations of subunits may specify subdivisions.

### 3. MICROANATOMY OF THE CENTRAL NUCLEUS

There are units of IC neuronal organization finer than the subdivisions described earlier. The microanatomy within each subdivision may define functional modules that control important aspects of information processing. We have proposed that synaptic domains are functional modules formed by the segregation of in-




Figure 2.11. A model of ICC synaptic domains in the ICC. Each functional module is denoted by a different shade and represents a different excitatory brain stem input (*light gray*, medial superior olive; *dark gray*, cochlear nucleus; *black*, lateral superior olive). Inhibitory inputs (lateral superior olive, *white spheres*; dorsal nucleus of the lateral lemniscus, *black spheres*) terminate in particular domains and avoid others. The distribution of some modules is highly related to the tonotopic map since some inputs are absent at the ends of the frequency ranges.




Figure 2.12. Microanatomy of ICC. **(A)** Disc-shaped neurons (D) align to form laminae that are reinforced by the local axons of these cells and parallel to the brain stem laminar afferents. Stellate cells (S) have dendrites and axons that interact with several laminae. Both disc-shaped and stellate cells can synthesize GABA (*black cells*) while others may be glutamatergic. GABAergic neurons can project to the medial geniculate body as do the nonGABAergic cells (*not shown*). **(B)** Golgi-impregnated section of the cat IC shows the subdivisions and ICC laminae at lower magnification. Most laminae run from ventrolateral to dorsomedial. For other abbreviations see Figs. 2.1 and 2.2. (From Oliver 1984. Reprinted with permission of Wiley-Liss, Inc., a subsidiary of John Wiley & Sons, Inc.; Oliver et al. 1994.)

puts from different sources (Oliver and Huerta 1992; Oliver 2000). Synaptic domains are a basic feature of organization in the inferior and superior colliculus. Thus, small groups of neurons share a common set of synaptic inputs and are likely to share a similar function. In the ICC a single lamina may actually be composed of two or more synaptic domains (Fig. 2.11). Each DC layer may represent a functional module or a layer may contain multiple domains similar to the deep layers of the superior colliculus. Because the LC, VL, and RP nuclei also have unique combinations of inputs, they should be considered functional modules. Future study may show that they contain more than one domain. ICC functional modules have been the most thoroughly studied. Consequently, this section focuses on ICC modules and their role in processing auditory information.

The ICC has a three-dimensional organization defined by its microanatomy. Its laminae consist of arrays of disc-shaped neurons and parallel axons from brain stem and local sources (Fig. 2.12; cf. Section 2.3.2). In the cat the laminae are long sheets running obliquely to the rostrocaudal plane and several millimeters high. The bands of labeled axons seen after small injections in the brain stem or intracellular injections of single IC neurons suggest that single laminae are discrete units approximately 200  $\mu\text{m}$  wide (Figs. 2.8 to 2.11). This unit organization is probably driven by the axonal components, as single dendritic

fields are usually one half to one third the diameter of the axonal bands (Oliver and Morest 1984; Malmierca et al. 1993).

Given this stereotyped ICC microanatomy, the synaptic domains may be related to the dimensions of a single lamina. The most obvious functional parameter related to the laminae is tonotopic organization. This varies in the dimension parallel to the width of the laminae. We first discuss the evidence that the spectral properties of ICC cells are related to the laminar microanatomy. Whether synaptic domains create other functional modules will be the second topic of this section.

### *3.1. WHAT IS THE STRUCTURAL BASIS OF TONOTOPIC ORGANIZATION?*

The elementary organization of the ICC tonotopic map is clear from auditory physiology and experimental neuroanatomical studies. The lowest frequencies are represented dorsolaterally and the highest frequencies are ventromedial (Merzenich and Reid 1974; Schreiner and Langner 1997). Anatomical experiments show that brain stem neurons tuned to different frequencies project to corresponding positions of the ICC tonotopic map and these axonal laminae form discrete units approximately 200  $\mu\text{m}$  wide.

Studies with 2DG show the activation pattern of axons activated by pure tones. Because 2DG is taken up by presynaptic axons during synaptic activity (Nudo and Masterton 1986), this method reveals the three-dimensional organization of the axons when they are activated by sound (Brown et al. 1997b). Their spatial pattern recapitulates the morphology of the axonal laminae. Most sound-activated axons are in the medial band in ICC and the lateral band in the VL is weaker or unlabeled. These studies also show that the width of the axonal band activated by tone increases with stimulus intensity (Brown et al. 1997a). The minimum width of the band activated by near-threshold stimuli should be related to the spatial dimensions of the axons. However, it is difficult to relate the activation pattern to the microanatomy of the laminae because 2DG labeling is usually performed with carbon-14 emission, producing substantial cross-scatter that reduces resolution of the method to a level too coarse to resolve single axons. Because the 2DG accumulates in the presynaptic axons only, the firing of single neurons is not detected.

Microelectrode recordings from single and multiple neurons provide higher resolution and are consistent with a frequency organization based on discrete laminae. The characteristic frequency changes in steps of  $175 \pm 83 \mu\text{m}$  in penetrations perpendicular to the tonotopic axis (the dorsolateral to ventromedial direction) and each step is approximately 0.28 octaves (Schreiner and Langner 1997; see Chapter 11). Such frequency steps would constrain the maximum number of laminae available for ICC frequency coding. Thus, 35 to 40 laminae would be required to encode the 9 octaves audible in the cat (Schreiner and Langner 1997). This study suggests that an entire lamina may represent a narrow

range of characteristic frequencies, a “frequency-band lamina” rather than representing a single frequency, the “iso-frequency lamina” concept. However, this method does not address whether the laminar organization is effective in organizing the spectral properties of ICC at sound intensity levels above threshold.

It is now possible to relate the activity of the entire population of the ICC laminae to frequency coding using sound-activated Fos protein. Most neurons produce this nuclear transcription factor when stimuli induce regular bursts of firing for prolonged periods. The production of cFos mRNA and Fos protein is proportional to the amount of firing in the cochlear nucleus and dorsal root ganglion (Sheng et al. 1993; Fields et al. 1997; Saint Marie et al. 1999; Yang et al. 2003b). Consequently, the amount of protein reflects stimulus intensity (Yang et al. 2003b).

The sound activation of Fos protein can address whether the population of neurons activated by single tones is confined to a single lamina, independent of the stimulus intensity (Yang et al. 2003a). Interestingly the maximum band of neurons activated by a single sinusoidally amplitude modulated (SAM) tone (8 or 16 kHz; 14 Hz modulation) is limited to approximately 180  $\mu\text{m}$  wide (Fig. 2.13A). The width of the band is virtually the same for 47- to 80-dB stimuli, although at 27 dB the band narrows (Fig. 2.13B, C). Because the laminar dimension is similar to that suggested by the microelectrode recordings in the cat,




Figure 2.13. Fos protein in the rat IC activated by sounds of two different frequencies. (A) Montage showing Fos-immunopositive neural nuclei in the ICC, DC, and LC. The *inset* shows nuclei at higher magnification. Labeled ICC and LC neurons form bands. Isodensity contours show clusters of cells within bands activated by monaural 80-dB amplitude-modulated tones. (B) Histogram showing density of labeled cells in the box in A. Peaks correspond to the bands of labeled cells. (C) Smoothed curves fitted to the histograms generated by stimuli of different intensities. (From Yang et al. 2003a.)

the microanatomy of ICC laminae may constrain the spread of excitation in the frequency domain. Lateral inhibition may limit the number of neurons activated by a tone to those within the same lamina; however, the mechanisms for this inhibition are still unclear.

A second question addressed by sound activation of Fos is the laminar role in processing spectral information. For example, what frequency separation is necessary for two stimuli to activate independent ICC neural populations? Two populations of neurons were activated by two SAM tones (as above) one octave apart (Yang et al. 2003a, 2004). At levels from 27 to 80 dB the centers of the bands of labeled neurons were separated by 240  $\mu\text{m}$ . Similarly two bands of neurons were evident after stimulation with 37 dB tones separated by 0.5 octave and their centers were half the distance of those in the one octave cases. However, when two tones were separated only by 0.25 octave at 27 to 37 dB the sound-activated neurons formed one band approximately 200  $\mu\text{m}$  wide.

These experiments suggest that ICC laminar microanatomy may play an important role in spectral information processing and may be related to critical bands (Ehret and Merzenich 1985; Schreiner and Langner 1997). The laminae could permit independent neural populations to respond to stimuli separated by at least 0.5 octave across a relatively broad range of stimulus intensities. Two stimuli spectrally more separated are less likely to interact, as they activate discrete populations. For stimuli within 0.25 octave the activation is confined to a single population of neurons within a lamina. This population may respond to all stimuli within this range of frequencies while also encoding subtle differences in the activity pattern related to the stimulus spectrum. Stimuli that activate the same population of neurons in a lamina are likely to be identical perceptually because the neurons cannot distinguish them.

### *3.2. HOW DOES MICROANATOMY ORGANIZE INFERIOR COLLICULUS NEURONS WITH DIFFERENT BINAURAL RESPONSE PROPERTIES?*

Because there is only one tonotopic map in the ICC, the processing of other types of auditory information faces a severe problem. All neurons with a similar best frequency (within 0.25 octave) may be within the same frequency-band lamina. Thus, all ICC neurons with similar tuning are at the same point on the tonotopic map. If each receives identical inputs from all of the lower auditory brain stem, they might have the same response properties. However, this is clearly not the case (see Chapters 11 to 14).

Functional classes of neurons have been identified using several sets of criteria. Early studies classified IC neurons based on interaural level differences (ILD) (EE: bilateral excitation; EI: ipsilateral suppression; EO: monaural only) (Semple and Aitkin 1979; Irvine 1986, 1992). Further studies have clarified the binaural response properties and enlarged the earlier classifications (Kelly and Sally 1993). Binaural neurons sensitive to interaural time disparity (ITD) com-

prise at least two types, both related to ITD sensitivity in the SOC. For peak-type neurons the ITD functions across frequencies align at or near maximal discharge (Kuwada and Yin 1983; Yin and Kuwada 1983a,b; Kuwada et al. 1984, 1987; Stanford et al. 1992) similar to those in MSO (Yin and Chan 1990; Batra et al. 1997a,b). In contrast, trough-type IC neurons align at or near the minimal discharge (Batra et al. 1993) like those in the LSO (Joris and Yin 1995; Joris 1996). In contrast, monaural neurons (Irvine 1986, 1992) must constitute a separate class, as a monaural response would be impossible with the convergence of monaural and binaural inputs. Spectral response properties have been used to classify these neurons (Ramachandran et al. 1999; Le Beau et al. 2001). The frequency-tuning curve shape (V-, I-, or O-shaped) may be correlated with certain binaural inputs (Ramachandran and May 2002). For example, low-BF neurons with V-shaped tuning curves often show peak-type responses, consistent with an input from MSO. I- and O-shaped response classes may receive their primary excitatory synaptic inputs from LSO and DCN, respectively (Ramachandran et al. 1999; Davis 2002).

A critical unresolved issue is how the combination of brain stem inputs contributes to the response properties of IC neurons. One clue is that the inputs to the IC differ by their frequency ranges. For example, MSO is a low-frequency structure with most of its neurons in the range from 100 Hz to about 5 kHz (Guinan et al. 1972). By comparison, the DCN is a high-frequency structure with little representation <1 kHz (Spirou et al. 1993). A particularly complex pattern is shown by the LSO, as the contralateral projection to the IC is heavier at high frequencies and the ipsilateral projection is stronger for low frequencies (Glendenning and Masterton 1983). Such differences in frequency range suggest one reason why synaptic domains across the IC will likely receive different combinations of inputs.

The synaptic domain hypothesis predicts that neurons with similar response properties will occupy the same functional module and receive similar brain stem inputs. For example, IC neurons aggregate when they have similar properties for coding interaural level differences (Roth et al. 1978; Semple and Aitkin 1979). The following sections consider the evidence for synaptic domains as defined by the major excitatory inputs to the IC from the MSO, LSO, and cochlear nucleus.

### 3.2.1. The Medial Superior Olivary Domain and the Representation of Azimuth

One type of synaptic domain in the ICC may be dominated by MSO input. MSO neurons compute ITD and their receptive fields indicate the location of azimuthal sound sources (Yin and Chan 1990; Batra et al. 1997a,b). The MSO projections neurons are the major source of ITD information to the midbrain. Despite the importance of this pathway, how ITD information is passed from the MSO to the IC is unclear. For example, if the ITD is mapped in the MSO along the rostrocaudal axis, as is often assumed, is this map transmitted to the




Figure 2.14. Three-dimensional reconstructions of MSO axonal boutons terminating in a lamina in the cat IC. (A, B) Single laminae from three animals. (C) Lateral view of boutons in a lamina. (D) *En face* view of the bouton distribution in (C) shows the absence of a simple density gradient related to injection site location. In (A), (C), and (D) the boutons are from two injections of different dextran anterograde transport markers in the same MSO at the same best frequency, one rostral and the other caudal. Boutons are labeled with either biotinylated dextran amines (BDA) or dextran labeled with tetramethylrhodamine (TMR). (From Oliver et al. 2003.)

rostrocaudal dimension of the IC laminae? The evidence for spatial topography in the MSO–IC projections was examined in the cat by making recordings of ITD sensitivity and small injections of BDA in different MSO locations and tracing the axons to the IC laminae (Figs. 2.14 and 2.15; Oliver et al. 2003). Two deposits in the same animal allow overlap and segregation of terminal fields to be studied. Different rostrocaudal locations on the same MSO frequency plane do not terminate at restricted points on the IC lamina. Instead, the MSO axons terminate along the length of a lamina (Figs. 2.14A to C, 2.15C). Within the lamina the axonal boutons are not distributed at uniform density or in a linear gradient (Fig. 2.14D). Consequently, there are neither point-to-point connections






Figure 2.15. Axons from two points in the MSO with the same best frequency converge on the same IC lamina. (A) Injections of different dextrans. (B) Three-dimensional reconstruction of the colliculus to show the laminar labeling from these injections. (C) Axonal labeling in serial sections through the ICC (see *inset*). (From Oliver et al. 2003.)

nor a gradient of connections to convey information from a map of space in the MSO to a comparable map in the IC laminae.

These results suggest that the dimension of the lamina orthogonal to the frequency axis does not necessarily represent a monotonic spatial map of a sensory dimension. For the MSO projections to the IC, the transmission of information about azimuthal sound location does not rely on a rerepresentation of a spatial map of azimuth in the MSO that is conveyed to the IC. Consequently, the azimuthal response of an IC neuron may be unrelated to its laminar position. Instead, the results suggest that a complex nontopographical neural network codes the position of a sound source in the IC. The MSO domain is a key component of this network for the IC, as it identifies a population of neurons on a lamina that receive inputs and participate in the network to code sound location.

### 3.2.2. Lateral Superior Olivary Domain vs. the Medial Superior Olivary Domain

Do MSO inputs remain segregated from other major excitatory inputs to the ICC laminae? Excitatory LSO inputs have different binaural properties. As mentioned earlier, binaural response properties that mimic these inputs would be difficult to find in IC neurons if excitatory LSO and MSO inputs always converged on the same IC neuron. For example, peak-type ITD responses would be expected in IC neurons that receive excitatory inputs from MSO, but not excitatory trough-type ITD inputs from the LSO. To be consistent with the synaptic domain hypothesis, excitatory inputs from the MSO and the LSO should remain segregated. To test the hypothesis that the inputs from MSO and LSO would not converge in the same IC lamina, ITD-sensitive MSO and LSO cells with a similar characteristic frequency, but different binaural responses, were labeled with different anterograde tracers to trace the axons that project from these sites to the ICC.

The results suggest that the excitatory inputs from MSO and contralateral LSO remain separate (Loftus et al. 2002, 2004a). The MSO projections on one side were compared to the projections of the opposite LSO, as MSO axons project only to the ipsilateral IC, while the LSO excitatory projections are mostly contralateral. The LSO target in IC is more rostral than the MSO target (Fig. 2.11, black; Fig. 2.16A). The MSO termination zone is longer and has more boutons than the rostral LSO target (Fig. 2.11, light gray; Fig. 2.16B, C). This is consistent with injections of retrograde tracers in the caudal, low-frequency ICC, where neurons have peak-type ITD responses. Both the retrograde and anterograde labeling suggests that IC neurons with peak-type responses to ITD are in synaptic domains with excitatory inputs from the MSO whereas neurons with trough-type ITD responses are in another synaptic domain receiving mainly excitatory contralateral LSO input.




Figure 2.16. Summary of projections from the superior olives to the IC. One injection was made in the left MSO, and the second in the right LSO at an overlapping frequency range. Presumed excitatory ipsilateral MSO inputs (*open circles*) dominate (**B, C**) except in the most rostral levels (**A**) where excitatory inputs from the contralateral LSO dominate (*closed triangles*). Presumed inhibitory inputs from the ipsilateral LSO (*gray filled squares*) are superimposed on those from the medial superior olive. LN, Lateral nucleus; other labels as in Figs. 2.1 and 2.2. (From Loftus et al. 2004.)

These experiments also suggested that the MSO and LSO domains may have other forms of organization as well, including different inhibitory influences. The LSO projects to both the contralateral and ipsilateral IC, but only the contralateral projection is excitatory. The ipsilateral LSO projection is glycinergic. A comparison of both LSO projections was made by superimposing the bouton data from the right IC onto the left IC for each section. In contrast to the excitatory LSO and MSO inputs, which remained separate, the ipsilateral inhibitory LSO projection is denser, terminates in a broader area, and largely overlaps the MSO target zone (Fig. 2.16). Thus, there appears to be one synaptic domain with excitatory inputs from the MSO, and postsynaptic IC neurons also receive inhibitory input from the LSO. There appears to be a second synaptic domain where excitatory LSO inputs are prevalent. These are summarized in the model of the synaptic domains (Fig. 2.11) as the MSO domain (lightly shaded) and LSO domain (black). The inhibitory LSO input is shown as white spheres.

### 3.2.3. Comparing Central Nucleus Monaural and Binaural Domains

The third major excitatory input to the IC originates in the cochlear nucleus. It arises from multiple cell types in each major division of the cochlear nucleus. Although it is tempting to consider the cochlear nucleus inputs to the IC as a single entity, that would be a gross oversimplification (see Chapters 1 and 3) and DCN, AVCN, and PVCN neurons may not participate in the same synaptic domains. How the inputs from the divisions of the cochlear nucleus complex are related to each other in the IC remains an unresolved issue.

Because a proportion of IC neurons are monaural they likely receive direct input from the cochlear nucleus. They should lack input from binaural lower auditory brain stem neurons. This characteristic is most likely to occur in the high-frequency parts of ICC devoid of MSO afferents. In the rat IC cochlear nucleus axons were labeled with anterograde transport of BDA while tectothalamic IC neurons in the same animals were identified by retrograde transport from the medial geniculate body, and their dendrites were labeled by subsequent intracellular injection of Lucifer Yellow in formaldehyde-fixed brain slices. Cochlear nucleus inputs terminate directly on tectothalamic neurons (Oliver et al. 1999) and this synaptic input fulfills the first requirement for a monaural pathway extending from the cochlear nucleus to the thalamus.

The second requirement for a monaural pathway and for monaural IC neurons is a synaptic domain without binaural input. To address this issue the distribution of afferent axons from the DCN and LSO to the contralateral IC was compared (Oliver et al. 1997). DCN and LSO neurons were characterized by responses to monaural and binaural acoustic stimulation and a deposit was made in each structure with different markers. Both injection sites had cells with overlapping best frequencies. The results showed that DCN and LSO axons are superimposed in part of the contralateral ICC, in laminae in the ventral part of the central nucleus (Fig. 2.11, high frequency, black). However, in the dorsal part of the

same layer LSO axons are absent (Fig. 2.11, dark gray). These data suggest two types of synaptic domains in the high-frequency ICC laminae. One contains binaural neurons that combine the properties of inputs from the contralateral LSO and DCN. A second functional module may contain monaural neurons that have cochlear nucleus inputs and none from binaural structures.

### *3.3. UNRESOLVED ISSUES ABOUT MICROANATOMY AND FUNCTIONAL ZONES IN THE CENTRAL NUCLEUS*

The synaptic domain hypothesis remains a viable explanation for the anatomical organization of the inputs to the ICC. Unresolved issues remain concerning the microanatomy and the organization of the inputs to functionally defined neuron types. Chief among them is the relative contribution of inhibitory inputs to the microanatomy and to each synaptic domain. It is not clear whether modules defined by excitatory inputs receive different combinations of inhibitory input. Inhibitory inputs to the IC are GABAergic axons from the DNLL and glycinergic axons from the ipsilateral LSO. Lemniscal and olivary nuclei influences on ITD sensitivity may differ (Kuwada et al. 1997). Some ITD-sensitive neurons receive predominantly DNLL inputs while others receive mostly ipsilateral LSO inputs. Future experiments will determine if separate groups of IC neurons are postsynaptic to these two inhibitory brain stem inputs.

The contribution of inhibitory inputs to monaural processing also remains unclear. The VNLL is actually a complex of nuclei whose neurons have predominantly monaural responses (Aitkin et al. 1970; Covey 1993; Batra and Fitzpatrick 1999), although one subregion has binaural responses (Batra and Fitzpatrick 2002). Many VNLL neurons contain glycine, GABA, or both and they represent a major input to the ICC that is probably inhibitory (Brunso-Bechtold et al. 1981; Merchán and Berbel 1996; Malmierca et al. 1998). Anatomical studies suggest that the ventral nucleus inputs are laminar rather than diffuse, as suggested earlier (Whitley and Henkel 1984). However, it is unclear to which synaptic domains they contribute.

The local components of the functional modules also remain uncertain. It is unknown how the synaptic domains relate to ICC local axons. Although the existence of Golgi type II interneurons is unresolved (see Section 2.3.2 and Chapter 22) there could be local intra- or intermodular input. It is also unclear how the local environment within a module influences the postsynaptic neurons that reside there. Perhaps the combination of inputs that define a module can also influence number and types of synaptic receptors and ion channels in the postsynaptic neurons (see below).

While synaptic domains are defined largely by the excitatory inputs from binaural or monaural sources, the present relationship of these zones to other facets of auditory processing remains to be clarified. Because inputs from different brain stem sources may differ in their spectral properties, the specific combination of inputs may be related to the spectral properties of single IC neurons. Likewise, the brain stem inputs to IC may differ in their ability to code

complex temporal patterns. Whether functional zones that differ in binaural properties also differ in their spectral and temporal properties remains an open question.

A final question is how ICC synaptic domains relate to functional modules in other subdivisions of IC. Because a frequency reversal does not mark the border between the ICC and the DC, other response properties may distinguish them from the upper tier of ICC synaptic domains. Only the basic inputs of these regions have been determined. The predominant inputs to the upper domains in the ICC are from the cochlear nucleus and nuclei of the lateral lemniscus. In contrast, the predominant input to the adjacent deep dorsal cortex is from the neocortex. Much remains to be learned about the microanatomy of the dorsal cortex and how it differs from that in the ICC.

## 4. FUNCTIONAL DEFINITIONS OF NEURON TYPES IN THE CENTRAL NUCLEUS

Up to this point the discussion of functional modules has focused almost exclusively on the inputs. However, the postsynaptic neurons in each module will integrate information from each input. Different types of IC neurons may influence how the inputs are integrated. Although IC neurons were originally defined primarily by their dendritic morphology into disc-shaped and stellate categories, this binary classification does not begin to account for all the functional properties observed. IC neurons also differ in their axonal targeting, the types of neurotransmitters produced, and in their electrical properties. These factors must be considered in any definition of IC neuron types to better understand their role in auditory information processing.

### *4.1. NEURON CLASSIFICATIONS BASED ON AXONAL TARGETING*

One criterion for IC neuron types is a unique axonal target. The main IC target is the ipsilateral medial geniculate body (MGB) (see Chapter 7). ICC axons terminate primarily in the ventral division whereas axons from the dorsal cortex project to the deep part of the dorsal division. Some neurons from all subdivisions have axons that terminate in the medial division of MGB. Other IC targets are the ipsilateral nucleus of the brachium, the contralateral IC (see Chapter 5), and the contralateral MGB. Descending IC projections terminate in the periolivary nuclei of the superior olive and the cochlear nucleus (see Chapter 6).

Despite the fact that the targets of the IC are known, we have only a rudimentary knowledge of the cell types that participate in these connections. For example, both disc-shaped and stellate neurons project to the ipsilateral MGB (Oliver 1984b). It is not clear whether these two cell types carry the same information to the MGB and whether their termination patterns are similar. Even less is known about the cell types that participate in tectal or descending projections. Understanding which cell types participate in each IC projection will

allow us to determine whether the same information is transmitted from the IC to each target.

#### 4.2. GABA PROJECTION NEURONS IN THE INFERIOR COLLICULUS

A second functional criterion for defining IC neurons is the neurotransmitter content. Two types of IC neurons project to the MGB. One type is excitatory and probably uses glutamate as the neurotransmitter. The second type is inhibitory and uses GABA as a neurotransmitter. In the cat about 20% of the IC projection neurons to the MGB are GABAergic (Winer et al. 1996). In the rat 40% of the IC neurons projecting to the MGB contain GABA (Peruzzi et al. 1997). GABAergic projection neurons are distributed throughout the IC subdivisions except in the caudal cortex and intercollicular tegmentum (Fig. 2.17). This matches the distribution of GABA immunostaining without reference to axonal target. Some of the IC neurons projecting to the contralateral IC also contain GABA (González-Hernández et al. 1996). However, all of the targets have not been investigated and it is not known whether these two populations of IC neurons project to other targets.

This dual population of GABAergic and presumed glutamatergic projection neurons is a distinctive feature of the IC. GABAergic neurons are often the largest neurons in the IC (Oliver et al. 1994). Most GABA-containing neurons are disc-shaped neurons although some with stellate morphology are also seen.



Figure 2.17. Distribution of GABAergic and nonGABAergic IC neurons projecting to the rat MGB. (A) Caudal section showing ICC and caudal cortex. (B) Section through middle ICC. Tectothalamic neurons are labeled by retrograde transport and are GABA-positive (*black circles*) or GABA-negative neurons (*open squares*); there are also GABA-positive neurons (*open triangles*) that are not retrogradely labeled. (From Peruzzi et al. 1997.)

However, the dendritic morphology of GABAergic neurons has not been correlated with projections. It is suspected that disc-shaped and stellate GABAergic neurons project from the IC to the MGB.

GABAergic IC neurons may have a special role in auditory processing at the thalamic level. Monosynaptic inhibitory postsynaptic potentials (IPSPs) from the IC often reach the MGB before the excitatory postsynaptic potentials (EPSP) (Peruzzi et al. 1997; Bartlett and Smith 2002). Additional IPSPs are seen after the EPSP. The fast IC GABAergic input may actively control the timing of the onset of excitation in the MGB and the slow GABAergic input may contribute to the suppression of firing along with other mechanisms such as GABAergic inhibition from the thalamic reticular nucleus. These data suggest that the IC projection to the thalamus is more complex than the thalamic inputs in other sensory systems (Bartlett et al. 2000). Retinal inputs to the lateral geniculate and the spinal cord or dorsal column inputs to the ventrobasal complex are assumed to be purely excitatory. Only in the thalamic motor nuclei is there a mixture of inhibitory and excitatory inputs and an interplay between them that seems to underlie the temporal responses of the thalamic neurons (Peruzzi et al. 1997).

There are several questions regarding GABAergic neurons. As mentioned earlier, there are both disc-shaped and stellate GABA-containing IC neurons. However, it is unknown if both project to the same MGB targets. Further, the GABAergic IC neuron and its role as an inhibitory neuron has not been related to the auditory responses of IC neurons. For example, in the context of binaural response types (see Section 2.3.2), it is unknown whether GABAergic neurons favor MSO domains, LSO domains, or monaural domains.

### *4.3. NEURON TYPES DEFINED BY INTRINSIC MEMBRANE PROPERTIES*

A third basis for defining IC neurons is their intrinsic membrane properties. ICC neurons have distinct discharge patterns to intracellular current injection (Peruzzi et al. 2000) and their responses are correlated with distinctive current/voltage relationships (Sivaramakrishnan and Oliver 2001). Six physiological classes of IC cell are revealed by intracellular recording with either sharp electrodes or whole-cell patch-clamp techniques in rat brain slices. Neurons formed two groups based on their responses to the offset of hyperpolarizing currents. One group exhibits a rebound calcium-dependent depolarization in addition to sodium action potentials (Sivaramakrishnan and Oliver 2001), while the other group lacks the rebound response (Fig. 2.18). The rebound neurons form three subtypes based on their responses to depolarizing currents—a transient response (few action potentials, rapidly adapting), a sustained slowly adapting response, and a regular nonadapting response (Fig. 2.18, rebound). Nonrebound neurons also have three subtypes—an onset response with one spike (Fig. 2.18, onset), a sustained-regular response, and a pause-buildup response (Fig. 2.18, pause-buildup). The latter appears only when the depolarization is preceded by a hyperpolarization (see Chapter 10).






Figure 2.18. Firing patterns of neurons in the IC brain slice preparation. For each type the discharge pattern to depolarizing and hyperpolarizing intracellular current injections are shown (**left**). Neurons (**middle**) whose responses are shown were filled with neurobiotin (rebound onset) or Lucifer yellow (pause-buildup). IC location of the neuron appears on the inset (**right**). (From Peruzzi et al. 2000. Reprinted from Neuroscience with permission from Elsevier.)

The six IC firing patterns are generated by a unique  $K^+$  current and set of cellular parameters (Sivaramakrishnan and Oliver 2001; cf. Chapter 10). For example, the IC rebound-adapting cell likely has an SK-type  $K^+$  channel as it has a  $Ca^{2+}$ -activated  $K^+$  current that is blocked by apamin. The rebound-transient cell also has a  $Ca^{2+}$ -activated  $K^+$  current, but of the BK-type, as it is

blocked by charybdotoxin. Likewise, the pause-buildup cell has a response that can be directly related to an A-current  $K^+$  current, while the onset response requires a high-threshold delayed rectifying  $K^+$  current such as *Kv3.1* (Perney et al. 1992). The rebound-regular and the sustained-regular cells have primarily delayed rectifier currents and show no evidence for the additional currents mentioned earlier. However, the rebound-regular cell has a rebound response that may reflect a high density of a T-type  $Ca^{2+}$  channel that produces a  $Ca^{2+}$  influx after hyperpolarization offset.

The intrinsic properties of IC neurons entail several questions. There is little understanding of the molecular basis of these membrane properties and the distinct combinations of ionic currents are still poorly understood. Such currents likely depend on the precise subunit combinations that compose the channel and different subunit combinations may produce subtle differences in their current properties. Identification of expression patterns will have an enormous practical impact beyond the immediate issue of membrane properties, as they will identify intrinsic molecular markers for functional IC cell types.

The relationship of morphology to the firing patterns needs to be resolved. The cell types based on intrinsic properties do not correspond to the simple disc-shaped or stellate morphological classifications. Most neurons in the small sample were flat cells (Fig. 2.18) and the data suggests subtypes with different intrinsic properties (Peruzzi et al. 2000; Sivaramakrishnan and Oliver 2001). A small sample of less-flat rat stellate cells does not allow conclusions about the firing patterns. However, some morphological features besides dendritic orientation may be related to discharge pattern. Dendritic branching patterns, size, or the incidence of spines may be correlated with the intrinsic properties and the dendritic electrotonus of different cell types.

It is also not known how the firing patterns relate to the cell's projection. In some brain regions intrinsic properties are associated with cell types with different targets, as in the cochlear nucleus, where the spherical bushy cells project to the MSO while the stellate cells project to the IC, and octopus neurons project to the VNLL but not to the main targets of the other cells. In the midbrain the tectothalamic neurons may have a different firing pattern than cells projecting to the superior olive. One possible purpose of correlated firing properties and targeting is that each neuron type can convey different types of information to its target.

Finally, the relationship of the firing pattern to GABA needs to be clarified. Elsewhere in the nervous system GABAergic and glycinergic neurons often play a special role in the microcircuitry. If GABAergic neurons have firing properties that distinguish them from other cells, the temporal pattern of inhibition may differ from that of excitation. To resolve this issue the organization of the synapses of the types of IC cells must be determined.

## 5. SYNAPTIC ORGANIZATION OF CENTRAL NUCLEUS NEURONS

The previous section suggests that IC neurons are more complex physiologically than their early morphological designations indicated. If different types of neurons also receive different types of synaptic inputs it would further enhance the functional differences between them. Understanding synaptic inputs to the different IC neuron types and how the intrinsic membrane properties of these cells further modify the synaptic input to shape a neural response is vital.

### *5.1. DO INFERIOR COLLICULUS NEURON TYPES RECEIVE SIMILAR SYNAPTIC INPUTS?*

Synaptic input is important for IC signal processing but little is known about the synaptic inputs to the different neuron types. Here synaptic input refers specifically to the excitatory or inhibitory postsynaptic current (EPSC or IPSC) elicited in the IC cell by stimulating the presynaptic ending and its interaction with ligand-gated receptors on the postsynaptic cell. IC neurons have various receptors for glutamate, glycine, and GABA (see Chapter 9) and the function of the synapses will depend on their subunit composition and their effect on the postsynaptic IC cell, as it does elsewhere in the brain (Barnes-Davies and Forsythe 1995; Wang et al. 1998; Gardner et al. 2001; Schmid et al. 2001). Thus, the GABA<sub>A</sub> receptors are responsible for fast Cl<sup>-</sup>-mediated IPSCs at GABAergic IC synapses. The patterns of expression for GABA<sub>A</sub> subunits provide the best evidence for neurons of different size having different combinations of GABA<sub>A</sub> subunits (Shiraishi et al. 2001). Such differences suggest that there may be different types of GABA synapses in the IC just as in the neocortex (Gupta et al. 2000).

A related issue is activity-dependent changes in IC synapses that differ among the different IC cell types. Glutamate inputs to IC neurons can show long-term potentiation (LTP) (Zhang and Wu 2000; Wu et al. 2002) similar to *N*-methyl D-aspartate- (NMDA) dependent LTP elsewhere in the nervous system. It is important to resolve whether such changes are equally common to all cell types.

The important implication for IC neuronal organization is that synaptic properties are vital parameters for the function of neuron types. Each type of neuron must be defined by the interactions of multiple parameters—the location in a particular synaptic domain, the axonal target, the cell's morphology, the intrinsic membrane properties, and the types of synaptic input. Because of unique AMPA, NMDA, or GABA<sub>A</sub> subunit compositions, activity could modify synaptic strength or the nature of the synaptic inputs, and consequently output could vary by the cell type. The relative number and location of glutamate, GABA, and glycine synapses also vary according to cell type. Each factor would contribute to the information processing carried out by the different IC neuron types.

## 5.2. CAN INTRINSIC AND SYNAPTIC PROPERTIES INTERACT TO CREATE THE RESPONSES OF INFERIOR COLLICULUS NEURON TYPES?

The cellular mechanisms that regulate synaptic integration are essential to understanding IC function in sound processing. Membrane properties can facilitate or degrade the temporal pattern of synaptic inputs evoked by sound and thus shape the synaptic inputs to which the cell ultimately responds. Neurons defined by different intrinsic properties may interact with the synaptic inputs in different ways that may especially influence the timing of responses.

The IC represents a transition stage for temporal coding in the auditory system. Timing here is better suited for coding the envelopes of complex signals (SAM tones; Batra et al. 1989), communication calls (Klug et al. 2002), and duration (Casseday et al. 2000) information than for coincidence detection (Fitzpatrick et al. 1997). Although the sources of input to IC neurons may phase lock and fire at high rates (Batra et al. 1997a,b; Spitzer 1998; Spitzer and Semple 1998), synaptic properties or intrinsic membrane properties of IC neurons could transform this information. Synaptic inputs may alter time codes because facilitation or depression occurs in some cell types, related perhaps to the relative contributions and kinetics of NMDA and AMPA components (see Chapter 10). Membrane properties are also important, as some IC cells follow stimuli at higher rates (Peruzzi et al. 2000).

We do not yet know the relative contributions of synaptic inputs and membrane properties to temporal processing in the different IC cells. Ultimately, the IC cell's response may reflect both the intrinsic membrane properties (voltage-gated and  $\text{Ca}^{2+}$ -activated ion channels) and synaptic inputs (presynaptic mechanisms and postsynaptic ligand-gated receptors).  $\text{Ca}^{2+}$ -activated conductances in particular can disrupt the normal cadence of neuronal firing in response to synaptic input, as they often enhance adaptation related to increased  $\text{Ca}^{2+}$  influx. The presence of  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  currents in some cell types suggests that their membrane properties may alter the response to synaptic input, transforming afferent input and shaping the temporal aspects of the response.

A final question relates to both synaptic inputs and intrinsic membrane properties and it is whether the neural responses to sound *in vivo* reflect different synaptic inputs or different membrane properties. The link between brain slice data, usually obtained from young animals, and sound processing in the adult is tenuous. Experiments are necessary in which IC cell types are identified in adult animals with the same criteria used in the brain slice experiments. Such an experiment may explain how the IC neuronal response is related to the cell-specific synaptic input and its sources.

## 6. SUMMARY

How are the types of IC neurons related to the synaptic domains? The synaptic domain hypothesis defines functional IC modules and seeks to establish a relationship between the auditory response properties of a neuron and the sources of its brain stem inputs. Within these domains the cell types are defined by their membrane properties, dendritic morphology,  $K^+$  channel subunits, and transmitter content. Whether particular IC cell types are associated with specific synaptic inputs remains to be resolved in future experiments. If so, it suggests a special relationship between the sources of the brain stem projections, the synaptic inputs, and the postsynaptic IC cells that coexist within a synaptic domain.

## Acknowledgments

The work in our laboratory has been supported by NIH grant DC00189. It could not have been accomplished without the outstanding contributions of Ms. Debbie Bishop, Ms. Gretchen Beckius, Drs. Shobhana Sivaramakrishnan, Ami Shneiderman, Dan Peruzzi, and William Loftus.

## Abbreviations

a	axon
AMPA	$\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazole propionate
AVCN	anteroventral cochlear nucleus
BI	brachium of the inferior colliculus
CG	central gray
DC	dorsal cortex
DCN	dorsal cochlear nucleus
2DG	2-deoxyglucose
DI	dorsal intercollicular tegmentum
DLL, DNLL	dorsal nucleus of the lateral lemniscus
EPSP	excitatory postsynaptic potential
GABA	$\gamma$ -aminobutyric acid
IC	inferior colliculus
ICC	central nucleus of the inferior colliculus
ILD	interaural level differences
IPSP	inhibitory postsynaptic potential
LC	lateral cortex
LI	lateral intercollicular tegmentum
LL	lateral lemniscus
LSO	lateral superior olive
LTB	lateral nucleus of the trapezoid body

MGB	medial geniculate body
MI	medial intercollicular tegmentum
MSO	medial superior olive
NMDA	<i>N</i> -methyl-D-aspartate
PVCN	posteroventral cochlear nucleus
RP	rostral pole
SAM	sinusoidally amplitude modulated
VL	ventrolateral nucleus
VNLL	ventral nucleus of the lateral lemniscus

## References

- Aitkin LM, Anderson DJ, and Brugge JF (1970) Tonotopic organization and discharge characteristics of single neurons in nuclei of the lateral lemniscus of the cat. *Journal of Neurophysiology* **33**:421–440.
- Aitkin LM, Tran L, and Syka J (1994) The responses of neurons in subdivisions of the inferior colliculus of cats to tonal noise and vocal stimuli. *Experimental Brain Research* **98**:53–64.
- Aitkin LM, Webster WR, Veale JL, and Crosby DC (1975) Inferior colliculus. I. Comparison of response properties of neurons in central, pericentral, and external nuclei of adult cat. *Journal of Neurophysiology* **38**:1196–1207.
- Barnes-Davies M and Forsythe ID (1995) Pre- and postsynaptic glutamate receptors at a giant excitatory synapse in rat auditory brainstem slices. *Journal of Physiology (London)* **488**:387–406.
- Bartlett EL and Smith PH (2002) Effects of paired-pulse and repetitive stimulation on neurons in the rat medial geniculate body. *Neuroscience* **113**:957–974.
- Bartlett EL, Stark JM, Guillery RW, and Smith PH (2000) Comparison of the fine structure of cortical and collicular terminals in the rat medial geniculate body. *Neuroscience* **100**:811–828.
- Batra R and Fitzpatrick DC (1999) Discharge patterns of neurons in the ventral nucleus of the lateral lemniscus of the unanesthetized rabbit. *Journal of Neurophysiology* **82**:1097–1113.
- Batra R and Fitzpatrick DC (2002) Processing of interaural temporal disparities in the medial division of the ventral nucleus of the lateral lemniscus. *Journal of Neurophysiology* **88**:666–675.
- Batra R, Kuwada S, and Fitzpatrick DC (1997a) Sensitivity to interaural temporal disparities of low- and high-frequency neurons in the superior olivary complex. I. Heterogeneity of responses. *Journal of Neurophysiology* **78**:1222–1236.
- Batra R, Kuwada S, and Fitzpatrick DC (1997b) Sensitivity to interaural temporal disparities of low- and high-frequency neurons in the superior olivary complex. II. Coincidence detection. *Journal of Neurophysiology* **78**:1237–1247.
- Batra R, Kuwada S, and Stanford TR (1989) Temporal coding of envelopes and their interaural delays in the inferior colliculus of the unanesthetized rabbit. *Journal of Neurophysiology* **61**:257–268.
- Batra R, Kuwada S, and Stanford TR (1993) High-frequency neurons in the inferior colliculus that are sensitive to interaural delays of amplitude-modulated tones: evidence for dual binaural influences. *Journal of Neurophysiology* **70**:64–80.

- Berman AL (1968) *The Brain Stem of the Cat. A Cytoarchitectonic Atlas with Stereotaxic Coordinates*. The University of Wisconsin Press, Madison.
- Brown M, Webster WR, and Martin RL (1997a) Intensity and frequency functions of [<sup>14</sup>C]2-deoxyglucose labelling in the central nucleus of the inferior colliculus in the cat. *Hearing Research* **104**:73–89.
- Brown M, Webster WR, and Martin RL (1997b) The three-dimensional frequency organization of the inferior colliculus of the cat: a 2-deoxyglucose study. *Hearing Research* **104**:57–72.
- Brunso-Bechtold JK, Thompson GC, and Masterton RB (1981) HRP study of the organization of auditory afferents ascending to central nucleus of inferior colliculus in cat. *Journal of Comparative Neurology* **197**:705–722.
- Caicedo A, d'Aldin C, Puel JL, and Eybalin M (1996) Distribution of calcium-binding protein immunoreactivities in the guinea pig auditory brainstem. *Anatomy and Embryology* **194**:465–487.
- Casseday JH, Ehrlich D, and Covey E (2000) Neural measurement of sound duration: control by excitatory-inhibitory interactions in the inferior colliculus. *Journal of Neurophysiology* **84**:1475–1487.
- Chen L, Kelly JB, and Wu SH (1999) The commissure of Probst as a source of GABAergic inhibition. *Hearing Research* **138**:106–114.
- Clopton BM and Winfield JA (1973) Tonotopic organization in the inferior colliculus of the rat. *Brain Research* **56**:355–358.
- Coleman JR, McDonald AJ, Pinek B, and Zrull MC (1992) The inferior colliculus: calbindin and parvalbumin immunoreactivity in neural grafts. *Experimental Neurology* **115**:142–145.
- Covey E (1993) The monaural nuclei of the lateral lemniscus: parallel pathways from cochlear nucleus to the midbrain. In: Merchán MA, Juiz JM, Godfrey DA, and Mugnaini E (eds). *The Mammalian Cochlear Nuclei*. Plenum Press, New York, pp. 321–334.
- Davis KA (2002) Evidence of a functionally segregated pathway from dorsal cochlear nucleus to inferior colliculus. *Journal of Neurophysiology* **87**:1824–1835.
- Dezso A, Schwarz DWF, and Schwarz IE (1993) A survey of the auditory midbrain, thalamus and forebrain in the chicken (*Gallus domesticus*) with cytochrome oxidase histochemistry. *Journal of Otolaryngology* **22**:391–396.
- Doubell TP, Baron J, Skaliora I, and King AJ (2000) Topographical projection from the superior colliculus to the nucleus of the brachium of the inferior colliculus in the ferret: convergence of visual and auditory information. *European Journal of Neuroscience* **12**:4290–4308.
- Ehret G and Merzenich MM (1985) Auditory midbrain responses parallel spectral integration phenomena. *Science* **227**:1245–1247.
- Faye-Lund H and Osen KK (1985) Anatomy of the inferior colliculus in rat. *Anatomy and Embryology* **171**:1–20.
- Fields RD, Eshete F, Stevens B, and Itoh K (1997) Action potential-dependent regulation of gene expression: temporal specificity in Ca<sup>2+</sup> cAMP-responsive element binding proteins and mitogen-activated protein kinase signaling. *Journal of Neuroscience* **17**:7252–7266.
- Fitzpatrick DC, Batra R, Stanford TR, and Kuwada S (1997) A neuronal population code for sound localization. *Nature* **388**:871–874.
- Fitzpatrick KA (1975) Cellular architecture and topographic organization of the inferior colliculus of the squirrel monkey. *Journal of Comparative Neurology* **164**:185–208.
- Gardner SM, Trussell LO, and Oertel D (2001) Correlation of AMPA receptor subunit

- composition with synaptic input in the mammalian cochlear nuclei. *Journal of Neuroscience* **21**:7428–7437.
- Geniec P and Morest DK (1971) The neuronal architecture of the human posterior colliculus. *Acta Otolaryngologica Supplement* **295**:1–33.
- Glendenning KK and Masterton RB (1983) Acoustic chiasm: efferent projections of the lateral superior olive. *Journal of Neuroscience* **3**:1521–1537.
- Glendenning KK, Baker BN, Hutson KA, and Masterton RB (1992) Acoustic chiasm V: inhibition and excitation in the ipsilateral and contralateral projections of LSO. *Journal of Comparative Neurology* **319**:100–122.
- Glazer II, Hof PR, and Morgane PJ (1998) Comparative analysis of calcium-binding protein-immunoreactive neuronal populations in the auditory and visual systems of the bottlenose dolphin (*Tursiops truncatus*) and the macaque monkey (*Macaca fascicularis*). *Journal of Chemical Neuroanatomy* **15**:203–237.
- González-Hernández T, Mantolán-Sarmiento B, González-González B, and Pérez-González H (1996) Sources of GABAergic input to the inferior colliculus of the rat. *Journal of Comparative Neurology* **372**:309–326.
- González-Lima F and Cada A (1994) Cytochrome oxidase activity in the auditory system of the mouse: a qualitative and quantitative histochemical study. *Neuroscience* **63**:559–578.
- González-Lima F, Valla J, and Matos-Collazo S (1997) Quantitative cytochemistry of cytochrome oxidase and cellular morphometry of the human inferior colliculus in control and Alzheimer's patients. *Brain Research* **752**:117–126.
- Guinan JJ Jr, Norris BE, and Guinan SS (1972) Single auditory units in the superior olivary complex. II: Locations of unit categories and tonotopic organization. *International Journal of Neuroscience* **4**:147–166.
- Gupta A, Wang Y, and Markram H (2000) Organizing principles for a diversity of GABAergic interneurons and synapses in the neocortex. *Science* **287**:273–278.
- Harting JK and Van Lieshout DP (2000) Projections from the rostral pole of the inferior colliculus to the cat superior colliculus. *Brain Research* **881**:244–247.
- Hutson KA (1988) *Connections of the Auditory Midbrain: Efferent Projections of the Dorsal Nucleus of the Lateral Lemniscus, the Nucleus Sagulum, and the Origins of the GABAergic Commissure of Probst*. Doctoral thesis. Florida State University, Tallahassee.
- Idrizbegovic E, Bogdanovic N, and Canlon B (1999) Sound stimulation increases calcium-binding protein immunoreactivity in the inferior colliculus in mice. *Neuroscience Letters* **259**:49–52.
- Irvine DRF (1986) The auditory brainstem. A review of the structure and function of auditory brainstem processing mechanisms. In: Autrum H, Ottoson D, Perl ER, Schmidt RF, Shimazu H, and Willis WD (eds). *Progress in Sensory Physiology* **7**:1–279.
- Irvine DRF (1992) Physiology of the auditory brainstem. In: Popper AN and Fay RR (eds). *Springer Handbook of Auditory Research, Volume 2: The Mammalian Auditory Pathway: Neurophysiology*. Springer-Verlag, New York, pp. 153–231.
- Joris PX (1996) Envelope coding in the lateral superior olive. II. Characteristic delays and comparison with responses in the medial superior olive. *Journal of Neurophysiology* **76**:2137–2156.
- Joris PX and Yin TCT (1995) Envelope coding in the lateral superior olive. I. Sensitivity to interaural time differences. *Journal of Neurophysiology* **73**:1043–1062.
- Kelly JB and Sally SL (1993) Effects of superior olivary complex lesions on binaural responses in rat auditory cortex. *Brain Research* **605**:237–250.



- King AJ, Jiang ZD, and Moore DR (1998) Auditory brainstem projections to the ferret superior colliculus: anatomical contribution to the neural coding of sound azimuth. *Journal of Comparative Neurology* **390**:342–365.
- Klug A, Bauer EE, Hanson JT, Hurley L, Meitzen J, and Pollak GD (2002) Response selectivity for species-specific calls in the inferior colliculus of Mexican free-tailed bats is generated by inhibition. *Journal of Neurophysiology* **88**:1941–1954.
- Knudsen EI (1983a) Early auditory experience aligns the auditory map of space in the optic tectum of the barn owl. *Science* **222**:939–942.
- Knudsen EI (1983b) Subdivisions of the inferior colliculus in the barn owl (*Tyto alba*). *Journal of Comparative Neurology* **218**:174–186.
- Kudo M and Niimi K (1980) Ascending projections of the inferior colliculus in the cat: an autoradiographic study. *Journal of Comparative Neurology* **191**:545–556.
- Kuwada S and Yin TCT (1983) Binaural interaction in low-frequency neurons in inferior colliculus of the cat. I. Effects of long interaural delays, intensity, and repetition rate on interaural delay function. *Journal of Neurophysiology* **50**:981–999.
- Kuwada S, Yin TCT, Syka J, Buunen TJF, and Wickesberg RE (1984) Binaural interaction in low-frequency neurons in inferior colliculus of the cat. IV. Comparison of monaural and binaural response properties. *Journal of Neurophysiology* **51**:1306–1325.
- Kuwada S, Stanford TR, and Batra R (1987) Interaural phase-sensitive units in the inferior colliculus of the unanesthetized rabbit: effects of changing phase. *Journal of Neurophysiology* **57**:1338–1360.
- Kuwada S, Batra R, and Fitzpatrick DC (1997) Neural processing of binaural temporal cues. In: Gilkey RH and Anderson TR (eds). *Binaural and Spatial Hearing in Real and Virtual Environments*. Lawrence Erlbaum Associates, Mahwah, pp. 399–425.
- Le Beau FEN, Malmierca MS, and Rees A (2001) Iontophoresis *in vivo* demonstrates a key role for GABA<sub>A</sub> and glycinergic inhibition in shaping frequency response areas in the inferior colliculus of guinea pig. *Journal of Neuroscience* **21**:7303–7312.
- Li Y, Evans MS, and Faingold CL (1998) *In vitro* electrophysiology of neurons in subnuclei of rat inferior colliculus. *Hearing Research* **121**:1–10.
- Loftus WC, Bishop DC, Saint Marie RL, and Oliver DL (2002) Segregation of excitatory inputs from the medial superior olive (MSO) and the lateral superior olive (LSO) in laminae of the low frequency inferior colliculus (IC). *Society for Neuroscience Abstracts* **28**:845.4.
- Loftus WC, Bishop DC, Saint Marie RL, and Oliver DL (2004a) Organization of binaural excitatory and inhibitory inputs to the inferior colliculus from the superior olive. *Journal of Comparative Neurology* **472**:330–344.
- Loftus WC, Malmierca MS, Bishop DC, and Oliver DL (2004b) Ascending inputs to the lateral cortex of the inferior colliculus (IC) in rat and cat. *Association for Research in Otolaryngology Abstracts* **27**:109.
- Lohmann C and Friauf E (1996) Distribution of the calcium-binding proteins parvalbumin and calretinin in the auditory brainstem of adult and developing rats. *Journal of Comparative Neurology* **367**:90–109.
- Lopez DE, Saldaña E, Nodal FR, Merchán MA, and Warr WB (1999) Projections of cochlear root neurons, sentinels of the rat auditory pathway. *Journal of Comparative Neurology* **415**:160–174.
- Malmierca MS, Blackstad TW, Osen KK, Karagülle T, and Molowny RL (1993) The central nucleus of the inferior colliculus in rat: a Golgi and computer reconstruction study of neuronal and laminar structure. *Journal of Comparative Neurology* **333**:1–27.

- Malmierca MS, Rees A, Le Beau FEN, and Bjaalie JG (1995) Laminar organization of frequency-defined local axons within and between the inferior colliculi of the guinea pig. *Journal of Comparative Neurology* **357**:124–144.
- Malmierca MS, Leergaard TB, Bajo VM, Bjaalie JG, and Merchán MA (1998) Anatomic evidence of a three-dimensional mosaic pattern of tonotopic organization in the ventral complex of the lateral lemniscus in cat. *Journal of Neuroscience* **18**:10603–10618.
- Malmierca MS, Merchán MA, Henkel CK, and Oliver DL (2002) Direct projections from cochlear nuclear complex to auditory thalamus in the rat. *Journal of Neuroscience* **22**:10891–10897.
- Meininger V, Pol D, and Derer P (1986) The inferior colliculus of the mouse. A Nissl and Golgi study. *Neuroscience* **17**:1159–1179.
- Merchán MA and Berbel P (1996) Anatomy of the ventral nucleus of the lateral lemniscus in rats: a nucleus with a concentric laminar organization. *Journal of Comparative Neurology* **372**:245–263.
- Merchán MA, Saldaña E, and Plaza I (1994) Dorsal nucleus of the lateral lemniscus in the rat: concentric organization and tonotopic projection to the inferior colliculus. *Journal of Comparative Neurology* **342**:259–278.
- Merzenich MM and Reid MD (1974) Representation of the cochlea within the inferior colliculus of the cat. *Brain Research* **77**:397–415.
- Morest DK (1969) The growth of dendrites in the mammalian brain. *Zeitschrift für Anatomie und Entwicklungsgeschichte* **128**:290–317.
- Morest DK and Oliver DL (1984) The neuronal architecture of the inferior colliculus in the cat: defining the functional anatomy of the auditory midbrain. *Journal of Comparative Neurology* **222**:209–236.
- Nudo RJ and Masterton RB (1986) Stimulation-induced [<sup>14</sup>C]2-deoxyglucose labeling of synaptic activity in the central auditory system. *Journal of Comparative Neurology* **245**:553–565.
- Ohshima T, Endo T, and Onaya T (1991) Distribution of parvalbumin immunoreactivity in the human brain. *Journal of Neurology* **238**:320–322.
- Oliver DL (1984a) Dorsal cochlear nucleus projections to the inferior colliculus in the cat: a light and electron microscopic study. *Journal of Comparative Neurology* **224**:155–172.
- Oliver DL (1984b) Neuron types in the central nucleus of the inferior colliculus that project to the medial geniculate body. *Neuroscience* **11**:409–424.
- Oliver DL (1985) Quantitative analyses of axonal endings in the central nucleus of the inferior colliculus and distribution of 3H-labeling after injections in the dorsal cochlear nucleus. *Journal of Comparative Neurology* **237**:343–359.
- Oliver DL (1987) Projections to the inferior colliculus from the anteroventral cochlear nucleus in the cat: possible substrates for binaural interaction. *Journal of Comparative Neurology* **264**:24–46.
- Oliver DL (2000) Ascending efferent projections of the superior olivary complex. *Microscopy Research and Technique* **51**:355–363.
- Oliver DL and Huerta MF (1992) Inferior and superior colliculi. In: Webster DB, Popper AN, and Fay RR (eds). *Springer Handbook of Auditory Research, Volume 2: The Mammalian Auditory Pathway: Neurophysiology*. Springer-Verlag, New York, pp. 168–221.
- Oliver DL and Morest DK (1984) The central nucleus of the inferior colliculus in the cat. *Journal of Comparative Neurology* **222**:237–264.
- Oliver DL and Shneiderman A (1991) The anatomy of the inferior colliculus: cellular basis for integration of monaural and binaural information. In: Altschuler RA, Bobbin

- RP, Clopton BM, and Hoffman DW (eds). *Neurobiology of Hearing: The Central Auditory System*. Raven Press, New York, pp. 195–222.
- Oliver DL, Kuwada S, Yin TCT, Haberly LB, and Henkel CK (1991) Dendritic and axonal morphology of HRP-injected neurons in the inferior colliculus of the cat. *Journal of Comparative Neurology* **303**:75–100.
- Oliver DL, Winer JA, Beckius GE, and Saint Marie RL (1994) Morphology of GABAergic neurons in the inferior colliculus of the cat. *Journal of Comparative Neurology* **340**:27–42.
- Oliver DL, Beckius GE, and Shneiderman A (1995) Axonal projections from the lateral and medial superior olive to the inferior colliculus of the cat: a study using electron microscopic autoradiography. *Journal of Comparative Neurology* **360**:17–32.
- Oliver DL, Beckius GE, Bishop DC, and Kuwada S (1997) Simultaneous anterograde labeling of axonal layers from lateral superior olive and dorsal cochlear nucleus in the inferior colliculus of cat. *Journal of Comparative Neurology* **382**:215–229.
- Oliver DL, Ostapoff E-M, and Beckius GE (1999) Direct innervation of identified tectothalamic neurons in the inferior colliculus by axons from the cochlear nucleus. *Neuroscience* **93**:643–658.
- Oliver DL, Beckius GE, Loftus WC, and Batra R (2003) Topography of interaural temporal disparity coding in projections of medial superior olive to inferior colliculus. *Journal of Neuroscience* **23**:7438–7449.
- Olszewski J and Baxter D (1982) *Cytoarchitecture of the Human Brain Stem*. Karger, New York.
- Paloff AM, Usunoff KG, Hinova-Palova DV, and Ivanov DP (1989) The fine structure of the inferior colliculus in the cat. I. Neuronal perikarya in the central nucleus. *Journal für Hirnforschung* **30**:69–90.
- Paloff AM, Usunoff KG, and Hinova-Palova DV (1992) Ultrastructure of Golgi-impregnated and gold-toned neurons in the central nucleus of the inferior colliculus in the cat. *Journal für Hirnforschung* **33**:361–407.
- Paxinos G (1999) *Chemoarchitectonic Atlas of the Rat Brainstem*. Academic Press, San Diego.
- Paxinos G and Watson C (1998) *The Rat Brain in Stereotaxic Coordinates*. Academic Press, San Diego.
- Perney TM, Marshall J, Martin KA, Hockfield S, and Kaczmarek LK (1992) Expression of the mRNAs for the KV3.1 potassium channel gene in the adult and developing rat brain. *Journal of Neurophysiology* **68**:756–766.
- Peruzzi D, Bartlett E, Smith PH, and Oliver DL (1997) A monosynaptic GABAergic input from the inferior colliculus to the medial geniculate body in rat. *Journal of Neuroscience* **17**:3766–3777.
- Peruzzi D, Sivaramakrishnan S, and Oliver DL (2000) Identification of cell types in brain slices of the inferior colliculus. *Neuroscience* **101**:403–416.
- Poremba A, Jones D, and González-Lima F (1997) Metabolic effects of blocking tone conditioning on the rat auditory system. *Neurobiology of Learning and Memory* **68**:154–171.
- Ramachandran R and May BJ (2002) Functional segregation of ITD sensitivity in the inferior colliculus of decerebrate cats. *Journal of Neurophysiology* **88**:2251–2261.
- Ramachandran R, Davis KA, and May BJ (1999) Single-unit responses in the inferior colliculus of decerebrate cats I. Classification based on frequency response maps. *Journal of Neurophysiology* **82**:152–163.
- Ramón y Cajal S (1995) *Histology of the Nervous System of Man and Vertebrates*. Oxford University Press, New York.

- Ribak CE and Roberts RC (1986) The ultrastructure of the central nucleus of the inferior colliculus of the Sprague-Dawley rat. *Journal of Neurocytology* **15**:421–438.
- Riquelme R, Saldaña E, Osen KK, Ottersen OP, and Merchán MA (2001) Colocalization of GABA and glycine in the ventral nucleus of the lateral lemniscus in rat: an in situ hybridization and semiquantitative immunocytochemical study. *Journal of Comparative Neurology* **432**:409–424.
- RoBards MJ (1979) Somatic neurons in the brainstem and neocortex projecting to the external nucleus of the inferior colliculus: an anatomical study in the opossum. *Journal of Comparative Neurology* **184**:547–565.
- RoBards MJ, Watkins DW III, and Masterton RB (1976) An anatomical study of some somesthetic afferents to the intercollicular terminal zone of the midbrain of the opossum. *Journal of Comparative Neurology* **170**:499–524.
- Rockel AJ and Jones EG (1973a) The neuronal organization of the inferior colliculus of the adult cat. I. The central nucleus. *Journal of Comparative Neurology* **147**:11–60.
- Rockel AJ and Jones EG (1973b) The neuronal organization of the inferior colliculus of the adult cat. II. The pericentral nucleus. *Journal of Comparative Neurology* **149**:301–334.
- Roth GL, Aitkin LM, Andersen RA, and Merzenich MM (1978) Some features of the spatial organization of the central nucleus of the inferior colliculus of the cat. *Journal of Comparative Neurology* **182**:661–680.
- Saint Marie RL and Baker RA (1990) Neurotransmitter-specific uptake and retrograde transport of [3H]glycine from the inferior colliculus by ipsilateral projections of the superior olivary complex and nuclei of the lateral lemniscus. *Brain Research* **524**:244–253.
- Saint Marie RL, Ostapoff E-M, Morest DK, and Wenthold RJ (1989) Glycine-immunoreactive projection of the cat lateral superior olive: possible role in midbrain ear dominance. *Journal of Comparative Neurology* **279**:382–396.
- Saint Marie RL, Shneiderman A, and Stanforth DA (1997) Patterns of  $\gamma$ -aminobutyric acid and glycine immunoreactivities reflect structural and functional differences of the cat lateral lemniscal nuclei. *Journal of Comparative Neurology* **389**:264–276.
- Saint Marie RL, Luo L, and Ryan AF (1999) Effects of stimulus frequency and intensity on c-fos mRNA expression in the adult rat auditory brainstem. *Journal of Comparative Neurology* **404**:258–270.
- Saldaña E and Merchán MA (1992) Intrinsic and commissural connections of the rat inferior colliculus. *Journal of Comparative Neurology* **319**:417–437.
- Schmid S, Guthmann A, Ruppertsberg JP, and Herbert H (2001) Expression of AMPA receptor subunit flip/flop splice variants in the rat auditory brainstem and inferior colliculus. *Journal of Comparative Neurology* **430**:160–171.
- Schreiner CE and Langner G (1997) Laminar fine structure of frequency organization in auditory midbrain. *Nature* **388**:383–386.
- Semple MN and Aitkin LM (1979) Representation of sound frequency and laterality by units in central nucleus of cat inferior colliculus. *Journal of Neurophysiology* **42**:1626–1639.
- Senut MC and Alvarado-Mallart RM (1987) Cytodifferentiation of quail tectal primordium transplanted homotopically into the chick embryo. *Brain Research* **429**:187–205.
- Seto-Ohshima A, Aoki E, Semba R, Emson PC, and Heizmann CW (1990) Parvalbumin immunoreactivity in the central auditory system of the gerbil: a developmental study. *Neuroscience Letters* **119**:60–63.

- Sheng HZ, Fields RD, and Nelson PG (1993) Specific regulation of immediate early genes by patterned neuronal activity. *Journal of Neuroscience Research* **35**:459–467.
- Shiraishi S, Shiraishi Y, Oliver DL, and Altschuler RA (2001) Expression of GABA<sub>A</sub> receptor subunits in the rat central nucleus of the inferior colliculus. *Molecular Brain Research* **96**:122–132.
- Shneiderman A and Henkel CK (1987) Banding of lateral superior olivary nucleus afferents in the inferior colliculus: a possible substrate for sensory integration. *Journal of Comparative Neurology* **266**:519–534.
- Shneiderman A and Oliver DL (1989) EM autoradiographic study of the projections from the dorsal nucleus of the lateral lemniscus: a possible source of inhibitory inputs to the inferior colliculus. *Journal of Comparative Neurology* **286**:28–47.
- Shneiderman A, Oliver DL, and Henkel CK (1988) Connections of the dorsal nucleus of the lateral lemniscus: an inhibitory parallel pathway in the ascending auditory system? *Journal of Comparative Neurology* **276**:188–208.
- Shneiderman A, Chase MB, Rockwood JM, Benson CG, and Potashner SJ (1993) Evidence for a GABAergic projection from the dorsal nucleus of the lateral lemniscus to the inferior colliculus. *Journal of Neurochemistry* **60**:72–82.
- Sivaramakrishnan S and Oliver DL (2001) Distinct K currents result in physiologically distinct cell types in the inferior colliculus of the rat. *Journal of Neuroscience* **21**:2861–2877.
- Smith PH (1992) Anatomy and physiology of multipolar cells in the rat inferior collicular cortex using the in vitro brain slice technique. *Journal of Neuroscience* **12**:3700–3715.
- Spencer RF, Shaia WT, Gleason AT, Sismanis A, and Shapiro SM (2002) Changes in calcium-binding protein expression in the auditory brainstem nuclei of the jaundiced Gunn rat. *Hearing Research* **171**:129–141.
- Spirou GA, May BJ, Wright DD, and Ryugo DK (1993) Frequency organization of the dorsal cochlear nucleus in cats. *Journal of Comparative Neurology* **329**:36–52.
- Spitzer MW and Semple MN (1998) Transformation of binaural response properties in the ascending auditory pathway: influence of time-varying interaural phase disparity. *Journal of Neurophysiology* **80**:3062–3076.
- Stanford TR, Kuwada S, and Batra R (1992) A comparison of the interaural time sensitivity of neurons in the inferior colliculus and thalamus of the unanesthetized rabbit. *Journal of Neuroscience* **12**:3200–3216.
- Tardif E, Chiry O, Probst A, Magistretti PJ, and Clarke S (2003) Patterns of calcium-binding proteins in human inferior colliculus: identification of subdivisions and evidence for putative parallel systems. *Neuroscience* **116**:1111–1121.
- Vater M and Braun K (1994) Parvalbumin calbindin D-28k and calretinin immunoreactivity in the ascending auditory pathway of horseshoe bats. *Journal of Comparative Neurology* **341**:534–558.
- Waitzman DM and Oliver DL (2002) The midbrain. In: Ramachandran VS (ed). *Encyclopedia of the Human Brain*. Academic Press, San Diego, pp. 43–68.
- Wang YX, Wenthold RJ, Ottersen OP, and Petralia RS (1998) Endbulb synapses in the anteroventral cochlear nucleus express a specific subset of AMPA-type glutamate receptor subunits. *Journal of Neuroscience* **18**:1148–1160.
- Whitley JM and Henkel CK (1984) Topographical organization of the inferior collicular projection and other connections of the ventral nucleus of the lateral lemniscus in the cat. *Journal of Comparative Neurology* **229**:257–270.

- Wiberg M, Westman J, and Blomqvist A (1987) Somatosensory projection to the mesencephalon: an anatomical study in the monkey. *Journal of Comparative Neurology* **264**:92–117.
- Winer JA, Saint Marie RL, Larue DT, and Oliver DL (1996) GABAergic feedforward projections from the inferior colliculus to the medial geniculate body. *Proceedings of the National Academy of Sciences of the United States of America* **93**:8005–8010.
- Winer JA, Larue DT, Diehl JJ, and Hefti BJ (1998) Auditory cortical projections to the cat inferior colliculus. *Journal of Comparative Neurology* **400**:147–174.
- Wu SH, Ma CL, Sivaramkrishnan S, and Oliver DL (2002) Synaptic modification in neurons of the central nucleus of the inferior colliculus. *Hearing Research* **168**:43–54.
- Yang Y, Saint Marie RL, and Oliver DL (2003a) Frequency resolution in the auditory brainstem determined by sound-induced expression of Fos protein. *Society for Neuroscience Abstracts* **29**:592.11.
- Yang Y, Saint Marie RL, and Oliver DL (2003b) Threshold sensitivities of neuronal populations in cochlear nucleus (CN) determined by sound-induced FOS expression: evidence for acoustic stimulation of granule cells. *Association for Research in Otolaryngology Abstracts* **26**:701.
- Yang Y, Saint Marie RL, and Oliver DL (2004) The effect of binaural stimulation on frequency resolution in the rat inferior colliculus (IC). *Association for Research in Otolaryngology Abstracts* **27**:48.
- Yasuhara O, Akiyama H, McGeer EG, and McGeer PL (1994) Immunohistochemical localization of hyaluronic acid in rat and human brain. *Brain Research* **635**:269–282.
- Yin TCT and Chan JCK (1990) Interaural time sensitivity in medial superior olive of cat. *Journal of Neurophysiology* **64**:465–488.
- Yin TCT and Kuwada S (1983a) Binaural interaction in low-frequency neurons in inferior colliculus of the cat. III. Effects of changing frequency. *Journal of Neurophysiology* **50**:1020–1042.
- Yin TCT and Kuwada S (1983b) Binaural interaction in low-frequency neurons in inferior colliculus of the cat. II. Effects of changing rate and direction of interaural phase. *Journal of Neurophysiology* **50**:1000–1019.
- Zhang DX, Li L, Kelly JB, and Wu SH (1998) GABAergic projections from the lateral lemniscus to the inferior colliculus of the rat. *Hearing Research* **117**:1–12.
- Zhang Y and Wu SH (2000) Long-term potentiation in the inferior colliculus studied in rat brain slice. *Hearing Research* **147**:92–103.
- Zook JM, Winer JA, Pollak GD, and Bodenhamer RD (1985) Topology of the central nucleus of the mustache bat's inferior colliculus: correlation of single unit properties and neuronal architecture. *Journal of Comparative Neurology* **231**:530–546.

# Chapter 3

## Projections from the Cochlear Nuclear Complex to the Inferior Colliculus

NELL BEATTY CANT

### 1. INTRODUCTION

The substantial projections from the cochlear nuclei to the inferior colliculus (IC) do not seem to have been generally recognized as a common mammalian characteristic until about the late 1960s. Although they were described in some of the earliest experimental studies of the auditory pathways (Woollard and Harpman 1940, *q.v.*, for a discussion of the earlier literature on the subject; Barnes et al. 1943), the prevailing view appeared to be that most of the axons leaving the cochlear nucleus terminated in the superior olivary complex or nuclei of the lateral lemniscus and that the projections to the IC were sparse. Certainly, the matter appeared to be unresolved in 1953, as Stotler stated flatly in his influential article on brain stem auditory pathways in the cat that “all neurons reaching the level of the inferior colliculus are of the third order.” It was not until the late 1960s and early 1970s that studies with sensitive degeneration techniques (*cf.* Nauta 1993) demonstrated conclusively that both the dorsal and ventral cochlear nuclei project directly to the contralateral inferior colliculus (cat: Warr 1966, 1969, 1972; Fernandez and Karapas 1967; van Noort 1969; Osen 1972; Rhesus monkey: Strominger and Strominger 1971; Strominger 1973; chimpanzee: Strominger et al. 1977; kangaroo rat: Browner and Webster 1975). In some cases, sparse ipsilateral projections were also reported.

Degeneration studies established that the main target of both the dorsal and ventral cochlear nuclei in the IC is the central nucleus, that the projections from these two sources overlap, and that they are topographically organized, with the cochlea systematically represented from apex to base. Since the late 1970s, neuroanatomical methods based on retrograde and anterograde axonal transport of various tracers have allowed detailed studies of the projections from the cochlear nuclei to the IC in a wide variety of species (Table 3.1). These tracing studies, supplemented by electron microscopy, have provided descriptions of both the cell types contributing to the projections and also the distribution and arborization patterns of their axons within the IC. An overview of the projections from the cochlear nuclei to the IC is presented in Section 2. The discussion is based mainly on the results of large injections of retrograde or anterograde tracers in

Table 3.1. Neuroanatomical studies of projections from the cochlear nucleus to the inferior colliculus by species, 1978–2003.

Species	Authors and year of study
CAT	Roth, Aitkin, Andersen, and Merzenich 1978 Adams 1979, 1983 Aitkin, Kenyon, and Philpott 1981 Brunso-Bechtold, Thompson, and Masterton 1981 Oliver 1984, 1985, 1987 Aitkin and Schuck 1985 Maffi and Aitkin 1987 Oliver and Beckius 1993 Oliver, Beckius, Bishop, and Kuwada 1997
RAT	Beyerl 1978 Druga and Syka 1984 Tokunaga, Sugita, and Otani 1984 Coleman and Clerici 1987 Alibardi 1998, 1999 Oliver, Ostapoff, and Beckius 1999
MOUSE	Ryugo, Willard, and Fekete 1981 Ryugo and Willard 1985 Frisina, Walton, Lynch-Armour, and Byrd 1998
MARSUPIAL	Aitkin and Kenyon 1981 Willard and Martin 1983 Aitkin, Byers, and Nelson 1986
GERBIL	Nordeen, Killackey, and Kitzes 1983a,b Moore and Kitzes 1985
FERRET	Moore 1988
MOLE	Kudo, Nakamura, Tokuno, and Kitao 1990
GUINEA PIG	Schofield and Cant 1996 Alibardi 2000
CHINCHILLA	Josephson and Morest 1998
BAT: Mustache bat	Zook and Casseday 1982, 1985, 1987 Ross, Pollak, and Zook 1988 Frisina, O'Neill, and Zettel 1989 Ross and Pollak 1989 Wenstrup, Mittmann, and Grose 1999
Horseshoe bat	Schweizer 1981 Vater and Feng 1990



the IC or cochlear nuclei, respectively. In Section 3, the laminar organization of inputs to the IC as revealed by small tracer injections is discussed.

## 2. PROJECTIONS FROM THE COCHLEAR NUCLEUS TO THE IC


### 2.1. COMPARISON WITH OTHER SOURCES OF ASCENDING INPUT

The major sources of ascending auditory inputs to the IC are the cochlear nuclei, the nuclei of the superior olivary complex, and the nuclei of the lateral lemniscus (see Chapter 1). Some of the relevant neuroanatomical studies (those that contain information about the projections from the cochlear nuclei) are listed in Table 3.1. In some of these studies, counts were made of the numbers of neurons labeled in brain stem nuclei after large injections in the IC of retrograde tracers (substances that are taken up by axon terminals and transported through the parent axons back to the cell body of origin). Results from these studies have been replotted on common axes in Fig. 3.1 so that they can be compared directly. The figure shows that the pattern of labeling in the brain stem after large tracer injections is quite consistent across studies and species. (The results of small tracer injections are considerably more variable [see Section 3.3]). In terms of the numbers of neurons that are labeled, the projections from the contralateral cochlear nuclei are matched only by the projections from the ipsilateral ventral nucleus of the lateral lemniscus in most species. In the Japanese mole and the mustache bat (Fig. 3.1F, G), the number of labeled cells in the cochlear nuclei and superior olivary complex may be more nearly equal than in the other species, but the apparent differences among species could also be a function of differences in the sizes of the injection sites (see Section 4.3). The number of cells in the contralateral ventral cochlear nucleus that project to the IC is consistently larger than the number in the contralateral dorsal cochlear nucleus.

The contralateral projections are topographically organized. Cells in the ventral (low-frequency) parts of the cochlear nuclei project to the dorsolateral IC, and cells in the more dorsal (high-frequency) parts of the cochlear nuclei project to the ventromedial IC. Osen (1972) noted that the parts of the cochlear nucleus that represent middle frequencies project more caudally in the IC than do the parts with the lowest and highest frequency representations. In most species, a few cells in both the dorsal and ventral cochlear nuclei project to the ipsilateral IC. The ipsilateral projections appear to be mainly to the dorsolateral (or low-frequency) part (Nordeen et al. 1983a; Oliver 1984, 1987). In the chimpanzee, the ipsilateral projections were described as “substantial” (Strominger et al. 1977).

### 2.2. CELL TYPES THAT PROJECT TO THE INFERIOR COLLICULUS

The cochlear nuclear complex contains a number of well-characterized cell types that can be distinguished based on a wide variety of criteria (reviewed by Cant



Sources of ascending input (see caption for key)

Figure 3.1. Summary of results of 10 studies in seven species in which large injections of retrograde axonal tracers were placed in the IC. In each study, counts were made of the number of labeled cells in some or all of the sources of ascending inputs to the IC. For those studies in which the counts were reported as the numbers of labeled cells in each structure, the data have been converted to percent of total labeled cells counted in the structures shown. In some cases, counts were also made in other sources of input, such as the contralateral IC. If possible, these were not included in the calculation of the percent of labeled cells. **(A)** Cat (Adams 1979). The result from pressure injection of horseradish peroxidase (HRP) in ventral IC. Counts made in a sample of one twelfth of the total tissue;  $N = 3799$ . **(B)** Cat (Roth et al. 1978). The result from pressure injection of HRP in dorsal IC. The largest injection (in terms of numbers of labeled cells) in a series of six; 2-kHz region; counts made in 60% of the tissue;  $N = 6305$ . **(C)** Cat (Kudo and Nakamura 1988, quoted in Kudo et al. 1990). The result from a pressure injection of HRP conjugated to wheat germ agglutinin (HRP-WGA). Information about the counts not available. **(D)** Rat (Tokunaga et al. 1984). The result from pressure injection of HRP in the IC. The largest injection (in terms of numbers of labeled cells) in a series of four; counts made in every other section;  $N = 4754$ . **(E)** Ferret (Moore 1988). The result from multiple pressure injections of HRP-WGA in the IC. Counts made in every other section; numbers corrected for uncounted sections;  $N = 41,605$ . This does not include 15,435 neurons labeled in the contralateral IC; those cells were not included in the calculation of the percent labeled cells shown here. **(F)** Japanese mole (Kudo et al. 1990). The result from a pressure injection of HRP-WGA in IC. Counts made in every other section; average of counts from four different animals; total  $N = 12,291$ . **(G)** Mustache bat (Ross et al. 1988). The result from an iontophoretic injection filling most of the 60-kHz region of the IC. Counts made in every other section; estimated  $N = 2000-2300$ . A small percentage of the counted labeled cells were in the contralateral IC; data not included here. **(H)** Rat (Druga and Syka 1984). Pressure injection of HRP in the IC. Counts made in three animals and summed; sampling rate not given; total  $N = 3608$ . **(I)** Guinea pig (Schofield and Cant 1992). The result from a pressure injection of fluorescent tracers. Counts made in every sixth section;  $N = 4122$ . **(J)** Gerbil (Nordeen et al. 1983a). The result from a pressure injection of HRP in the IC. Counts in every other section; estimated  $N = 4300$ . For all panels, the *black bars* show the cell counts for the cochlear nucleus: *a*, ipsilateral dorsal cochlear nucleus (DCN); *b*, contralateral DCN; *c*, ipsilateral ventral cochlear nucleus (VCN); *d*, contralateral VCN. *Gray bars (middle)* show the cell counts for nuclei in the superior olivary complex: *e*, ipsilateral lateral superior olivary nucleus (LSO); *f*, contralateral LSO; *g*, ipsilateral medial superior olivary nucleus (MSO). *White bars* show the cell counts for nuclei of the lateral lemniscus: *h*, ipsilateral ventral nucleus of the lateral lemniscus; *i*, ipsilateral dorsal nucleus of the lateral lemniscus (DNLL); *j*, contralateral DNLL. *Gray bars (end)* show cell counts in other nuclei counted in specific studies: *k*, periolivary nuclei (PO); *l*, contralateral MSO (a substantial projection in the mole); *m*, ipsilateral PO; *n*, contralateral PO. Note the change in scale for panels **(H-J)**. Labeled cells in the nuclei of the lateral lemniscus were not counted in these studies, resulting in larger percentages in the other nuclei.

and Benson 2003). Even in studies based on degeneration techniques, it was recognized that only some of these cell types project to the inferior colliculus (Warr 1966, 1969, 1972; van Noort 1969). In all species studied (Table 3.1), it has been demonstrated that both fusiform and giant cells of the dorsal cochlear nucleus send axons to the IC. In fact, it appears that all of the large cells in the dorsal cochlear nucleus participate in the projection (Ryugo et al. 1981; Moore 1988). The projection from the ventral cochlear nucleus arises from neurons throughout its extent, including all but the most anterior part of the anteroventral cochlear nucleus and the octopus cell area in the posteroventral cochlear nucleus. It is generally agreed that multipolar (or stellate) cells are the major source of projections from the ventral cochlear nucleus, that octopus cells do not terminate in the IC, and that few if any globular cells or large spherical cells project that far. The case is less clear for small spherical cells in the anteroventral cochlear nucleus. A number of authors report that small, round cells in this part of the complex project to the IC, and some refer to these as small spherical cells, implying that they have a bushy cell morphology (cf. Cant and Benson 2003). Because the dendrites of the cells usually are not filled with the tracer, it is generally not possible to be sure that these really are spherical bushy cells, although Adams (1979) did illustrate one well-filled spherical bushy cell labeled after a large injection of horseradish peroxidase in the IC of the cat. Electron microscopic studies have provided no evidence for such a projection, however. In three species, all cells labeled retrogradely from the IC were identified in the electron microscope as type I stellate (or multipolar) cells (cat: Cant 1982; chinchilla: Josephson and Morest 1998; rat: Alibardi 1998). The cells that project from the ipsilateral cochlear nucleus have been identified as fusiform, giant, and multipolar cells (Adams 1979; Oliver 1984, 1987).

The ventral cochlear nucleus contains several types of multipolar cells. Those that project to the contralateral IC appear to be the type I multipolar cells (or type I stellate cells; cf. Cant and Benson 2003). It seems likely that most, if not all, of the type I cells participate in this projection (Josephson and Morest 1998). The same neurons in the ventral cochlear nucleus that project to the IC send collateral branches to the dorsal cochlear nucleus (Adams 1983). Multipolar cells in the ventral cochlear nucleus also give rise to projections to the superior olivary complex and the ventral nucleus of the lateral lemniscus (see Chapter 4). Although they have not been identified with certainty, the cells that give rise to these projections are probably also type I multipolar cells (Smith et al. 1993), but it is not known whether individual cells project to all of the targets. Some fusiform or giant cells in the dorsal cochlear nucleus may send axonal branches to both the ipsilateral and contralateral IC, but bilateral projections from cells in the ventral cochlear nucleus have not been described (Schofield and Cant 1996).

Axons arising from the fusiform and giant cells leave the cochlear nucleus in the dorsal acoustic stria. After crossing the midline, they enter the lateral lemniscus and travel in its medial part or in the adjacent tegmentum (Osen 1972; Willard and Martin 1983; Oliver 1984). The projections from the contralateral

ventral cochlear nucleus travel mainly in the trapezoid body, where they make up part of the thin fiber component, cross the midline, and enter the lateral part of the lateral lemniscus (Warr 1966; Osen 1972; Willard and Martin 1983). Ipsilateral projections from the cochlear nucleus travel in a small fiber bundle known as the lateral trapezoid body tract (Warr 1972).

### 2.3. *SYNAPTIC TERMINALS IN THE INFERIOR COLLICULUS*

Electron microscopic studies of the terminals from both the anteroventral and dorsal cochlear nuclei in the IC demonstrate that they contain round synaptic vesicles and make asymmetrical contacts with their targets, the morphology associated with excitatory synapses (Oliver 1984, 1985, 1987). The fine structure of the inputs from the posteroventral cochlear nucleus to the IC has not been reported, but presumably they are also excitatory, as the collaterals of their axons that project into the dorsal cochlear nucleus (Adams 1983) give rise to terminals with round synaptic vesicles (Smith and Rhode 1989). Alibardi (1998) also concluded that the neurons in the posteroventral cochlear nucleus that project to the IC are excitatory, based on the variable presence of immunoreactivity for glutamate and the absence of immunoreactivity for the inhibitory neurotransmitters glycine and  $\gamma$ -aminobutyric acid (GABA). The ventral nucleus of the lateral lemniscus contains almost as many neurons that project to the IC as the cochlear nuclei (Fig. 3.1), but because a large proportion of the cells in the ventral nucleus of the lateral lemniscus are inhibitory (Riquelme et al. 2001), it is apparent that, in terms of numbers of neurons, the contralateral cochlear nucleus represents the major brain stem source of excitatory inputs to the IC.

Although the number of cells projecting to the IC from the cochlear nuclei may be much greater than the number of cells projecting from the main superior olivary nuclei, many of which are also excitatory (e.g., Oliver et al. 1995), the number of terminals that is made by axons from the different sources may be more comparable. For example, in areas where the terminals from the ipsilateral medial superior olivary nucleus terminate most densely, they may account for up to 36% of the terminals with round synaptic vesicles in that area. Terminals with round vesicles from the ipsilateral and contralateral lateral superior olivary nucleus can reach 26% and 18%, respectively. These values can be compared to the maximum density of terminals from the contralateral anteroventral cochlear nucleus (13%) and the contralateral dorsal cochlear nucleus (11%). Interestingly, in the lateral part of the central nucleus, the synapses formed by the ipsilateral anteroventral cochlear nucleus can contribute up to 18% of the terminals with round vesicles, although the number of neurons projecting from the ipsilateral cochlear nucleus is always very small (Fig. 3.1). It is not yet known how the regions with the maximum density of inputs from any one source overlap with those from other sources (see Section 3).

The inputs from the cochlear nuclei form synapses on the dendrites of both disc-shaped cells and stellate cells, the main types in the central nucleus (see Chapter 2). Occasionally, terminals are also found on the cell bodies of the

stellate cells. Little is known about differential innervation of different types of IC neurons (defined in terms of morphology, physiological response properties, neurotransmitter chemistry, connections, other sources of input, or any other criteria). Oliver et al. (1999) showed that at least some of the inputs from the contralateral cochlear nucleus terminate directly on neurons that project to the medial geniculate nucleus.

### 3. DISTRIBUTION OF COCHLEAR NUCLEAR INPUTS WITHIN THE INFERIOR COLLICULUS

The topography related to frequency is the most obvious organizational principle in the IC, and most, if not all, inputs to the IC are organized with respect to frequency. The part of each brain stem nucleus that represents the apex of the cochlea (low frequencies) projects to the dorsolateral IC and the part that represents the base of the cochlea (high frequencies) projects to the ventromedial IC, with a systematic progression of the intermediate regions (e.g., Osen 1972; Adams 1979). This topography has given rise to the concept that the IC is organized into a series of laminae, each of which receives inputs from a limited frequency range and is continuous across the major subdivisions of the IC (cf. Oliver and Morest 1984; Saldaña and Merchán 1992; Brown et al. 1997). Within this framework of tonotopic organization, however, there is evidence for other levels of organization. Some studies have focused on differences in the organization of inputs from one subdivision to the next, while others have provided evidence for partial segregation of inputs within the central nucleus of the IC.

#### *3.1. PROJECTIONS TO THE MAJOR SUBDIVISIONS OF THE INFERIOR COLLICULUS*

The main target of both the ventral and dorsal cochlear nuclei in the IC is the central nucleus, where the projections from these two parts of the cochlear nuclear complex appear to overlap almost completely (Osen 1972; Oliver 1984). In addition, both the dorsal and ventral cochlear nuclei send projections into the deep layers of the dorsal cortex, although the terminations appear to be more sparse than those to the central nucleus (Oliver 1984; Coleman and Clerici 1987; Zook and Casseday 1987). In the dorsal cortex, the projections from the ventral cochlear nucleus may arborize more widely than those from the dorsal cochlear nucleus (Oliver 1987). The dorsal cochlear nucleus also projects to the external cortex of the IC in rats (Coleman and Clerici 1987; Oliver et al. 1999), mice (Ryugo et al. 1981), and opossums (Willard and Martin 1983) but not in the cat (Aitkin et al. 1981). Like those to the central nucleus, the projections to both the dorsal and external cortices appear to be topographically organized (Ryugo et al. 1981; Oliver 1984, 1987).

Because the main sources of inputs to each subdivision of the IC are different (cf. Chapter 1), the inputs from the cochlear nuclei must converge with those from different sources in the various parts of the IC. For example, in the central nucleus, the projections from the cochlear nuclei overlap to a large extent with the projections from the superior olivary complex, the other major source of ascending excitatory inputs to the IC (Oliver et al. 1995; Oliver 2000; see Chapter 4). However, the superior olivary complex does not project into the dorsal or external cortices of the IC (Henkel and Spangler 1983; Shneiderman and Henkel 1987; Oliver et al. 1997) so that in some parts of the IC, inputs from the olivary nuclei and the cochlear nuclei may be convergent and in other parts, cells could be influenced by the cochlear nuclei but not by the olivary nuclei. Likewise, the main targets in the IC of the auditory cortex are the dorsal and external cortices (Chapter 8). Here, the neurons that receive inputs from the cochlear nuclei might be more subject to descending control than those in the central nucleus, which receives relatively little cortical input.

### 3.2. ORGANIZATION WITHIN THE CENTRAL NUCLEUS

The proportion of labeled cells located in each source of ascending input to the IC is quite consistent both across studies and also across species after large injections of tracers (Fig. 3.1). However, when small (usually iontophoretic) injections of retrograde tracers are made into the central nucleus of the IC, the results are considerably more variable in terms of the proportions or even the presence of labeled cells in the different sources (Roth et al. 1978; Brunso-Bechtold et al. 1981; Aitkin and Schuck 1985; Maffi and Aitkin 1987; Ross and Pollak 1989; Wenstrup et al. 1999; cf. Oliver et al. 1995). For example, in some small regions, projections from the superior olivary nuclei may dominate. In the most extreme example reported, after an injection of a tracer in the lateral IC, 98% of the labeled cells were located in the ipsilateral medial superior olivary nucleus (Aitkin and Shuck 1985). In general, however, it is rare that so many labeled cells are located in only one source of input. In other cases, cells in the cochlear nuclei and nuclei of the lateral lemniscus might be labeled with little or no labeling in the superior olive. Across species and studies, almost every possible combination of inputs has been seen, although some are more common than others.

Studies with anterograde tracers also provide evidence for segregation of inputs within the central nucleus. Many inputs to the IC from the cochlear nucleus form bands that are parallel to the isofrequency laminae (Oliver 1984, 1987). In some cases, the bands from two sources overlap, as is the case for inputs from the contralateral dorsal cochlear nucleus and the contralateral lateral superior olivary nucleus (Oliver et al. 1997). Other inputs appear to lie in adjacent and possibly nonoverlapping bands, as is the case for the inputs from the ipsilateral and contralateral lateral superior olivary nuclei (Shneiderman and Henkel 1987). The inputs from the ipsilateral medial superior olivary nucleus and the ipsilateral cochlear nucleus may remain partially separate in the dorsolateral part of the IC

(Oliver and Beckius 1993). In that same part of the IC, some of the inputs from the contralateral and ipsilateral cochlear nuclei also appear to remain separate (C.G. Benson and N.B. Cant, unpublished results in the gerbil).

A dramatic example of segregation of inputs within a frequency lamina is provided by studies of the mustache bat, which exploited the fact that the 60-kHz “layer” is greatly expanded in this species (Wenstrup et al. 1986; Ross et al. 1988; Ross and Pollak 1989). When tracer injections filled most of the 60-kHz region, the pattern of labeling was very similar to that seen in other species (Fig. 3.1G), although more labeled cells were located in the nuclei of the lateral lemniscus than in the cochlear nuclei and the numbers of cells in the cochlear nucleus and the superior olivary complex were approximately equal (Ross et al. 1988). (Whether these differences from the common mammalian pattern are a result of specializations in the bat or occur because the injections are still relatively small is not clear.) However, when small, discrete injections were made systematically throughout the 60-kHz lamina, Ross and Pollak (1989) found different patterns of projections in four different regions that could be related to differences in the physiological response properties of the neurons in each region. Their results show that both connectivity and physiological response properties change systematically across this isofrequency representation.

In one of the first studies of projections to the IC employing tracer injections (and physiological recording), Roth and colleagues (1978) suggested that, “Within the cochleotopic and laminar framework of the central nucleus, it would seem that other rules of order must exist.” Other investigators have reached the same conclusion. Maffi and Aitkin (1987) proposed that the central nucleus is made up of “core zones” in which inputs from specific brain stem nuclei are dominant. Based on their demonstration of a systematic difference in the proportions of labeled cells in the cochlear nuclei vs. the superior olivary complex when horseradish peroxidase injections were made in the caudal vs. rostral IC, Brunso-Bechtold and colleagues (1981) concluded that an organization based on “nucleotopy” was superimposed on the cochleotopic organization of the IC. The apparent differential organization of inputs forms the basis for the hypothesis that the central nucleus is organized into “synaptic domains” as put forth by Oliver and Huerta (1992; Oliver 2000). In this conception, the synaptic domains are defined as small groups of neurons that share the same populations of synaptic inputs. A given synaptic domain may lie next to another synaptic domain with the same frequency characteristics but with different sets of inputs and, presumably, different physiological properties (see Chapter 2).

Consistent with the conclusions based on neuroanatomical studies, the physiological response properties of IC neurons have been shown to vary with location, and neurons with similar properties tend to cluster together (Roth et al. 1978; Semple and Aitkin 1979; Wenstrup et al. 1986; Brückner and RübSamen 1995). As noted earlier for the mustache bat, differences in the sources of inputs covary with differences in the physiological types found in various parts of the central nucleus. A similar progression of response types across isofrequency laminae was demonstrated in the gerbil (Brückner and RübSamen 1995). The



properties of some unit types in the IC of the cat appear to reflect a dominant input from only one of the many brain stem sources of inputs (Davis et al. 1999; Ramachandran et al. 1999; Davis 2002).

## 4. CONCLUSIONS

Neurons in the IC receive inputs from a wide variety of brain stem, forebrain, commissural, and intrinsic sources. Learning exactly how these inputs are organized at the level of the individual neurons that make up the IC represents a major challenge for future studies. Although at the light microscopic level, many of the inputs overlap to a greater or lesser extent, it is not known whether individual cells receive synapses from all of the axons in their vicinity or only from some of them. Although the potential for convergence of multiple inputs at the level of the IC is generally emphasized, there is ample potential for divergence and segregation of inputs from different sources at the level of the individual cells, especially because the IC may contain over five times as many neurons as the cochlear nuclei, superior olivary complex, and nuclei of the lateral lemniscus combined (Kulesza et al. 2002).

Given the many sources of inputs to the IC, the possibilities for combinations of inputs to any particular neuron are staggering. Even if the inputs from the cochlear nuclei alone are considered, there are a number of possible combinations. The cell types in the cochlear nuclei that project to the IC—the fusiform and giant cells in the dorsal cochlear nucleus and multipolar cells in the ventral cochlear nucleus—each have different physiological response properties and appear to encode different types of information (Rhode and Greenberg 1992). It is curious that no major differences in the organization of the terminal fields of these three cell types have been described, but it is certainly possible that there are important differences at the level of the synaptic organization of the cells in the IC. Potential interactions between ipsilateral and contralateral inputs from the cochlear nuclei add to the number of possibilities, as all three major cell types that project contralaterally also project ipsilaterally. Although the number of cells projecting to the IC from the ipsilateral cochlear nucleus is small, they may provide a substantial proportion of the inputs in the dorsolateral region. The inputs from the cochlear nuclei to the IC could also be combined in many different ways with the ascending inputs from the superior olivary complex and nuclei of the lateral lemniscus (see Chapter 4), commissural connections (see Chapter 5), and descending projections from the forebrain (see Chapter 8).

Adding to the complexity of organization made possible by the multitude of inputs is the intrinsic complexity afforded by the many different cell types that make up the IC. As discussed elsewhere, neurons in the IC may differ in their morphology (see Chapter 2), projection targets (see Chapters 5 and 6), transmitter expression (see Chapter 9), biophysical properties (see Chapter 10), and physiological response properties (see Chapters 11 to 13). It is reasonable to expect that cell types defined based on these differences will also differ in their

sources of input and synaptic organization, but at present very little is known about the synaptic organization of any particular cell class.

Neuroanatomical methods based on axonal transport of tracers have constantly improved since their introduction in the 1970s, and can still be used profitably to address many of the outstanding questions about the auditory pathways. Double and triple labeling experiments, combined with electron microscopy or physiological recording and marking techniques, can answer many important questions about the specific sources of inputs to specific cell types and provide substantive hypotheses for further work. These are tried and true methods, but they are technically difficult and progress is slow. It is exciting, therefore, that in the near future, new methods based on work in genetics and molecular biology can be expected to have a major impact on our ability to examine neural circuitry. For example, transgenic mice have been produced that express axonally transported tracers under the control of cell-type specific promoters (Yoshihara et al. 1999). In these animals, the projections of one specific cell type may be studied in isolation from those of other cell types. Similar methods are already being applied to some sensory systems (Braz et al. 2002; Kinoshita et al. 2002), and there is every reason to expect that such technology will vastly improve our ability to trace specific circuits in the complex pathways of the auditory system.

Knowledge of the connectivity and synaptic organization of specific cell types is essential for advancing our understanding of how physiological response properties emerge and, ultimately, of how each pathway contributes to overall auditory function. Although questions remain that can be addressed with currently available neuroanatomical techniques, we can look forward to an exciting new era of investigations of brain stem auditory circuitry as new techniques of molecular biology are used to address these questions.

## Acknowledgments

Dr. Christina Benson made many essential contributions to the writing of this chapter. Research in our laboratory is supported by the National Institutes on Deafness and Other Communication Disorders (DC00135).

## References

- Adams JC (1979) Ascending projections to the inferior colliculus. *Journal of Comparative Neurology* **183**:519–538.
- Adams JC (1983) Multipolar cells in the ventral cochlear nucleus project to the dorsal cochlear nucleus and the inferior colliculus. *Neuroscience Letters* **37**:205–208.
- Aitkin LM and Kenyon CE (1981) The auditory brain stem of a marsupial. *Brain, Behavior and Evolution* **19**:126–143.
- Aitkin LM and Schuck D (1985) Low frequency neurons in the lateral central nucleus of the cat inferior colliculus receive their input predominantly from the medial superior olive. *Hearing Research* **17**:87–93.

- Aitkin LM, Kenyon CE, and Philpott P (1981) The representation of the auditory and somatosensory systems in the external nucleus of the cat inferior colliculus. *Journal of Comparative Neurology* **196**:25–40.
- Aitkin LM, Byers M, and Nelson JE (1986) Brain stem auditory nuclei and their connections in a carnivorous marsupial, the Northern native cat (*Dasyurus hallucatus*). *Brain, Behavior and Evolution* **29**:1–16.
- Alibardi L (1998) Ultrastructural and immunocytochemical characterization of neurons in the rat ventral cochlear nucleus projecting to the inferior colliculus. *Annals of Anatomy* **180**:415–426.
- Alibardi L (1999) Fine structure, synaptology and immunocytochemistry of large neurons in the rat dorsal cochlear nucleus connected to the inferior colliculus. *Journal für Hirnforschung* **39**:429–439.
- Alibardi L (2000) Cytology of large neurons in the guinea pig dorsal cochlear nucleus contacting the inferior colliculus. *European Journal of Histochemistry* **44**:365–375.
- Barnes WT, Magoun HW, and Ranson SW (1943) The ascending auditory pathway in the brain stem of the monkey. *Journal of Comparative Neurology* **79**:129–152.
- Beyerl BD (1978) Afferent projections to the central nucleus of the inferior colliculus in the rat. *Brain Research* **145**:209–223.
- Braz JM, Rico B, and Basbaum AI (2002) Transneuronal tracing of diverse CNS circuits by Cre-mediated induction of wheat germ agglutinin in transgenic mice. *Proceedings of the National Academy of Sciences of the United States of America* **99**:15148–15153.
- Brown M, Webster WR, and Martin RL (1997) The three-dimensional frequency organization of the inferior colliculus of the cat: a 2-deoxyglucose study. *Hearing Research* **104**:57–72.
- Browner RH and Webster DB (1975) Projections of the trapezoid body and the superior olivary complex of the kangaroo rat (*Dipodomys merriami*). *Brain, Behavior and Evolution* **11**:322–354.
- Brückner S and Rübnsamen R (1995) Binaural response characteristics in isofrequency sheets of the gerbil inferior colliculus. *Hearing Research* **86**:1–14.
- Brunso-Bechtold JK, Thompson GC, and Masterton RB (1981) HRP study of the organization of auditory afferents ascending to central nucleus of inferior colliculus in cat. *Journal of Comparative Neurology* **197**:705–722.
- Cant NB (1982) Identification of cell types in the anteroventral cochlear nucleus that project to the inferior colliculus. *Neuroscience Letters* **32**:241–246.
- Cant NB and Benson CG (2003) Parallel auditory pathways: projection patterns of the different neuronal populations in the dorsal and ventral cochlear nuclei. *Brain Research Bulletin* **60**:457–474.
- Coleman JR and Clerici WJ (1987) Sources of projections to subdivisions of the inferior colliculus in the rat. *Journal of Comparative Neurology* **262**:215–226.
- Davis KA (2002) Evidence of a functionally segregated pathway from dorsal cochlear nucleus to inferior colliculus. *Journal of Neurophysiology* **87**:1824–1835.
- Davis KA, Ramachandran R, and May BJ (1999) Single-unit responses in the inferior colliculus of decerebrate cats. II. Sensitivity to interaural level differences. *Journal of Neurophysiology* **82**:164–175.
- Druga R and Syka J (1984) Ascending and descending projections to the inferior colliculus in the rat. *Physiologia Bohemoslovaca* **33**:31–42.
- Fernandez C and Karapas F (1967) The course and termination of the striae of Monakow and Held in the cat. *Journal of Comparative Neurology* **131**:371–386.

- Frisina RD, O'Neill WE, and Zettel ML (1989) Functional organization of mustached bat inferior colliculus: II. Connections of the FM<sub>2</sub> region. *Journal of Comparative Neurology* **284**:85–107.
- Frisina RD, Walton JP, Lynch-Armour MA, and Byrd JD (1998) Inputs to a physiologically characterized region of the inferior colliculus of the young adult CBA mouse. *Hearing Research* **115**:61–81.
- Henkel CK and Spangler KM (1983) Organization of the efferent projections of the medial superior olivary nucleus in the cat as revealed by HRP and autoradiographic tracing methods. *Journal of Comparative Neurology* **221**:416–428.
- Josephson EM and Morest DK (1998) A quantitative profile of the synapses on the stellate cell body and axon in the cochlear nucleus of the chinchilla. *Journal of Neurocytology* **27**:841–864.
- Kinoshita N, Mizuno T, and Yoshihara Y (2002) Adenovirus-mediated WGA gene delivery for transsynaptic labeling of mouse olfactory pathways. *Chemical Senses* **27**:215–223.
- Kudo M, Nakamura Y, Tokuno H, and Kitao Y (1990) Auditory brainstem in the mole (*Mogera*): nuclear configurations and the projections to the inferior colliculus. *Journal of Comparative Neurology* **298**:400–412.
- Kulesza RJ Jr, Viñuela A, Saldaña E, and Berrebi AS (2002) Unbiased stereological estimates of neuron number in subcortical auditory nuclei of the rat. *Hearing Research* **168**:12–24.
- Maffi CL and Aitkin LM (1987) Differential neural projections to regions of the inferior colliculus of the cat responsive to high frequency sounds. *Hearing Research* **26**:211–219.
- Moore DR (1988) Auditory brainstem of the ferret: sources of projections to the inferior colliculus. *Journal of Comparative Neurology* **269**:342–354.
- Moore DR and Kitzes LM (1985) Projections from the cochlear nucleus to the inferior colliculus in normal and neonatally cochlea-ablated gerbils. *Journal of Comparative Neurology* **240**:180–195.
- Nauta WJH (1993) Some early travails of tracing axonal pathways in the brain. *Journal of Neuroscience* **13**:1337–1345.
- Nordeen KW, Killackey HP, and Kitzes LM (1983a) Ascending auditory projections to the inferior colliculus in the adult gerbil, *Meriones unguiculatus*. *Journal of Comparative Neurology* **214**:131–143.
- Nordeen KW, Killackey HP, and Kitzes LM (1983b) Ascending projections to the inferior colliculus following unilateral cochlear ablation in the neonatal gerbil, *Meriones unguiculatus*. *Journal of Comparative Neurology* **214**:144–153.
- Oliver DL (1984) Dorsal cochlear nucleus projections to the inferior colliculus in the cat: a light and electron microscopic study. *Journal of Comparative Neurology* **224**:155–172.
- Oliver DL (1985) Quantitative analyses of axonal endings in the central nucleus of the inferior colliculus and distribution of <sup>3</sup>H-labeling after injections in the dorsal cochlear nucleus. *Journal of Comparative Neurology* **237**:343–359.
- Oliver DL (1987) Projections to the inferior colliculus from the anteroventral cochlear nucleus in the cat: possible substrates for binaural interaction. *Journal of Comparative Neurology* **264**:24–46.
- Oliver DL (2000) Ascending efferent projections of the superior olivary complex. *Microscopy Research and Technique* **51**:355–363.

- Oliver DL and Beckius GE (1993) Ascending projections from the cochlear nucleus to the inferior colliculus and their interactions with projections from the superior olivary complex. In: Merchán MA, Juiz JM, Godfrey DA, and Mugnaini E (eds). *The Mammalian Cochlear Nuclei. Organization and Function*. Plenum Press, New York, pp. 335–347.
- Oliver DL and Huerta MF (1992) Inferior and superior colliculi. In: Webster DB, Popper AN, and Fay RR (eds). *Springer Handbook of Auditory Research, Volume 1: The Mammalian Auditory Pathway: Neuroanatomy*. Springer-Verlag, New York, pp. 168–221.
- Oliver DL and Morest DK (1984) The central nucleus of the inferior colliculus in the cat. *Journal of Comparative Neurology* **222**:237–264.
- Oliver DL, Beckius GE, and Shneiderman A (1995) Axonal projections from the lateral and medial superior olive to the inferior colliculus of the cat: a study using electron microscopic autoradiography. *Journal of Comparative Neurology* **360**:17–32.
- Oliver DL, Beckius GE, Bishop DC, and Kuwada S (1997) Simultaneous anterograde labeling of axonal layers from lateral superior olive and dorsal cochlear nucleus in the inferior colliculus of cat. *Journal of Comparative Neurology* **382**:215–229.
- Oliver DL, Ostapoff E-M, and Beckius GE (1999) Direct innervation of identified tectothalamic neurons in the inferior colliculus by axons from the cochlear nucleus. *Neuroscience* **93**:643–658.
- Osen KK (1972) Projection of the cochlear nuclei on the inferior colliculus in the cat. *Journal of Comparative Neurology* **144**:355–372.
- Ramachandran R, Davis KA, and May BJ (1999) Single-unit responses in the inferior colliculus of decerebrate cats I. Classification based on frequency response maps. *Journal of Neurophysiology* **82**:152–163.
- Rhode WS and Greenberg S (1992) Physiology of the cochlear nuclei. In: Popper AN and Fay RR (eds). *Springer Handbook of Auditory Research, Volume 2: The Mammalian Auditory Pathway: Neurophysiology*. Springer-Verlag, New York, pp. 94–152.
- Riquelme R, Saldaña E, Osen KK, Ottersen OP, and Merchán MA (2001) Colocalization of GABA and glycine in the ventral nucleus of the lateral lemniscus in rat: an in situ hybridization and semiquantitative immunocytochemical study. *Journal of Comparative Neurology* **432**:409–424.
- Ross LS and Pollak GD (1989) Differential ascending projections to aural regions in the 60 kHz contour of the mustache bat's inferior colliculus. *Journal of Neuroscience* **9**: 2819–2834.
- Ross LS, Pollak GD, and Zook JM (1988) Origin of ascending projections to an iso-frequency region of the mustache bat's inferior colliculus. *Journal of Comparative Neurology* **270**:488–505.
- Roth GL, Aitkin LM, Andersen RA, and Merzenich MM (1978) Some features of the spatial organization of the central nucleus of the inferior colliculus of the cat. *Journal of Comparative Neurology* **182**:661–680.
- Ryugo DK and Willard FH (1985) The dorsal cochlear nucleus of the mouse: a light microscopic analysis of neurons that project to the inferior colliculus. *Journal of Comparative Neurology* **242**:381–396.
- Ryugo DK, Willard FH, and Fekete DM (1981) Differential afferent projections to the inferior colliculus from the cochlear nucleus in the albino mouse. *Brain Research* **210**: 342–349.
- Saldaña E and Merchán MA (1992) Intrinsic and commissural connections of the rat inferior colliculus. *Journal of Comparative Neurology* **319**:417–437.

- Schofield BR and Cant NB (1992) Organization of the superior olivary complex in the guinea pig: II. Patterns of projection from the periolivary nuclei to the inferior colliculus. *Journal of Comparative Neurology* **317**:438–455.
- Schofield BR and Cant NB (1996) Projections from the ventral cochlear nucleus to the inferior colliculus and the contralateral cochlear nucleus in guinea pigs. *Hearing Research* **102**:1–14.
- Schweizer H (1981) The connections of the inferior colliculus and the organization of the brainstem auditory system in the greater horseshoe bat (*Rhinolophus ferrum-equinum*). *Journal of Comparative Neurology* **201**:25–49.
- Semple MN and Aitkin LM (1979) Representation of sound frequency and laterality by units in central nucleus of cat inferior colliculus. *Journal of Neurophysiology* **42**:1626–1639.
- Shneiderman A and Henkel CK (1987) Banding of lateral superior olivary nucleus afferents in the inferior colliculus: a possible substrate for sensory integration. *Journal of Comparative Neurology* **266**:519–534.
- Smith PH and Rhode WS (1989) Structural and functional properties distinguish two types of multipolar cells in the ventral cochlear nucleus. *Journal of Comparative Neurology* **282**:595–616.
- Smith PH, Joris PX, Banks MI, and Yin TCT (1993) Responses of cochlear nucleus cells and projections of their axons. In: Merchán MA, Juiz JM, Godfrey DA, and Mugnaini E (eds). *The Mammalian Cochlear Nuclei. Organization and Function*. Plenum Press, New York, pp. 349–360.
- Stotler WA (1953) An experimental study of the cells and connections of the superior olivary complex of the cat. *Journal of Comparative Neurology* **98**:401–431.
- Strominger NL (1973) The origins, course and distribution of the dorsal and intermediate acoustic striae in the rhesus monkey. *Journal of Comparative Neurology* **147**:209–234.
- Strominger NL and Strominger AI (1971) Ascending brain stem projections of the anteroventral cochlear nucleus in the Rhesus monkey. *Journal of Comparative Neurology* **143**:217–242.
- Strominger NL, Nelson LR, and Dougherty WJ (1977) Second order auditory pathways in the chimpanzee. *Journal of Comparative Neurology* **172**:349–366.
- Tokunaga A, Sugita S, and Otani K (1984) Auditory and non-auditory subcortical afferents to the inferior colliculus in the rat. *Journal für Hirnforschung* **25**:461–472.
- van Noort J (1969) *The Structure and Connections of the Inferior Colliculus. An Investigation of the Lower Auditory System*. Van Gorcum, Assen.
- Vater M and Feng AS (1990) Functional organization of ascending and descending connections of the cochlear nucleus of horseshoe bats. *Journal of Comparative Neurology* **292**:373–395.
- Warr WB (1966) Fiber degeneration following lesions in the anterior ventral cochlear nucleus of the cat. *Experimental Neurology* **14**:453–474.
- Warr WB (1969) Fiber degeneration following lesions in the posteroventral cochlear nucleus of the cat. *Experimental Neurology* **23**:140–155.
- Warr WB (1972) Fiber degeneration following lesions in the multipolar and globular cell areas in the ventral cochlear nucleus of the cat. *Brain Research* **40**:247–270.
- Wenstrup JJ, Ross LS, and Pollak GD (1986) Binaural response organization within a frequency-band representation of the inferior colliculus: implications for sound localization. *Journal of Neuroscience* **6**:962–973.
- Wenstrup JJ, Mittmann DH, and Grose CD (1999) Inputs to combination-sensitive neurons of the inferior colliculus. *Journal of Comparative Neurology* **409**:509–528.

- Willard FH and Martin GF (1983) The auditory brainstem nuclei and some of their projections to the inferior colliculus in the North American opossum. *Neuroscience* **10**:1203–1232.
- Woollard HH and Harpman JA (1940) The connexions of the inferior colliculus and of the dorsal nucleus of the lateral lemniscus. *Journal of Anatomy (London)* **74**:441–458.
- Yoshihara Y, Mizuno T, Nakahira M, Kawasaki M, Watanabe Y, Kagamiyama H, Jishage K, Ueda O, Suzuki H, Tabuchi K, Sawamoto K, Okano H, Noda T, and Mori K (1999) A genetic approach to visualization of multisynaptic neural pathways using plant lectin transgene. *Neuron* **22**:33–41.
- Zook JM and Casseday JH (1982) Origin of ascending projections to inferior colliculus in the mustache bat, *Pteronotus parnellii*. *Journal of Comparative Neurology* **207**:14–28.
- Zook JM and Casseday JH (1985) Projections from the cochlear nuclei in the mustache bat, *Pteronotus parnellii*. *Journal of Comparative Neurology* **237**:307–324.
- Zook JM and Casseday JH (1987) Convergence of ascending pathways at the inferior colliculus of the mustache bat, *Pteronotus parnellii*. *Journal of Comparative Neurology* **261**:347–361.

# Chapter 4

## Superior Olivary Complex and Lateral Lemniscal Connections of the Auditory Midbrain

BRETT R. SCHOFIELD

### 1. INTRODUCTION

The inferior colliculus (IC) receives the majority of its ascending auditory inputs from the superior olivary complex (SOC) and the nuclei of the lateral lemniscus (NLL). These nuclei receive their inputs from the cochlear nucleus (CN), which also projects directly to the IC (see Chapter 3). Thus, there are parallel ascending pathways from the CN to the IC. These pathways analyze different aspects of acoustic information and send that information to the IC, where it is combined in ways that remain largely mysterious. Indeed, some of the most pressing questions in auditory research concern the functions of these parallel pathways and the nature of their convergence in the IC.

An important tool for understanding the functional organization of the ascending pathways has been conceptualizing them as monaural and binaural systems (Kudo and Nakamura 1987; Pollak and Casseday 1989; Helfert and Aschoff 1997; Casseday et al. 2002). Although this approach has been useful, it has been limited by a paucity of data on many of the relevant nuclei, especially the periolivary nuclei and the ventral and intermediate nuclei of the lateral lemniscus. The purpose of the present chapter is to examine new data on these nuclei and to present a new view of the monaural and binaural pathways. We begin with a brief review of the nuclei of the SOC and NLL and their projections to the IC, with an emphasis on recently acquired data. We then discuss the monaural and binaural systems and consider several aspects of their convergence in the IC. For additional information, and particularly for discussions of the identification of nuclei in different species, several reviews of the SOC and NLL are available (Schwartz 1992; Helfert and Aschoff 1997; Oliver 2000; Thompson and Schofield 2000; Smith and Spirou 2002; Casseday et al. 2002; Oertel and Wickesberg 2002).



## 2. THE SUPERIOR OLIVARY COMPLEX

Figure 4.1A illustrates the SOC nuclei. We consider the medial superior olivary nucleus (MSO) and the lateral superior olivary nucleus (LSO) to be the principal nuclei, and the remainder to constitute the periolivary nuclei (PON). Following a brief description of MSO and LSO circuitry (reviewed by Oliver 2000), we




Figure 4.1. The superior olivary complex and the circuits of the principal nuclei. (A) Schematic drawing (outlined from a guinea pig brain) showing the location of the major nuclei of the SOC. (B, C) Schematic drawings showing the major inputs to the MSO and the LSO. In this and subsequent drawings, *arrows* indicate major projections, with *line thickness* indicating relative size. The major neurotransmitter (or the postsynaptic effect) is indicated by the legend. GABA and glycine are generally inhibitory; the excitatory neurotransmitter(s) are less well known but for many pathways are believed to be glutamate or a related substance. The midline is represented by a *thin, dashed vertical line*.

focus on the PON and especially on the superior paraolivary nucleus (SPN), which is emerging as an important element in this system (Fig. 4.1A).

### 2.1. *THE MEDIAL SUPERIOR OLIVE*

The MSO receives bilateral excitation and inhibition (Fig. 4.1B). The excitation originates from spherical bushy (SB) cells, while the inhibition arises from globular bushy (GB) cells whose axons activate glycinergic cells in the medial or lateral nuclei of the trapezoid body (MNTB and LNTB, respectively), that project in turn to the MSO. The projections are organized tonotopically, with a relative overrepresentation of low frequencies.

The majority of MSO cells are excited by both ears and are particularly sensitive to interaural time differences. These cells likely play an important role in encoding azimuthal location of a sound.

Most MSO cells project ipsilaterally to the IC, where the axons terminate in bands in the central nucleus (ICc). Although a few MSO cells may use  $\gamma$ -aminobutyric acid (GABA) or glycine (Helfert et al. 1989; Winer et al. 1995), most appear to use an excitatory neurotransmitter (Kumoi et al. 1993; Oliver et al. 1995).

### 2.2. *THE LATERAL SUPERIOR OLIVE*

The major excitatory input to the LSO arises from SB cells in the ipsilateral VCN (Fig. 4.1C). The major inhibitory input arises from the MNTB, which is driven by GB cells in the contralateral VCN. Additional inputs, presumed to be inhibitory, arise from the ipsilateral LNTB and contralateral VNTB. Finally, small inputs arise from GB cells in the ipsilateral VCN and from multipolar cells in the VCN on both sides (not shown; Helfert and Aschoff 1997; Doucet and Ryugo 2003). As for the MSO, the inputs to the LSO are organized tonotopically; however, in the LSO the system is biased toward high frequencies.

Most LSO cells are excited by ipsilateral stimulation and inhibited by contralateral stimulation. These cells are particularly sensitive to interaural level differences (and less so to time differences). This sensitivity and the bias toward high frequencies have led to the idea that the LSO analyzes interaural differences that are critical for azimuthal localization of high-frequency sounds. A few binaurally excited cells are also present.

The projections from the LSO to the IC are bilateral, with ipsilateral and contralateral projections originating from different cells. In cats and several other species, the ipsilateral projection arises preferentially from the lateral limb of the LSO, while the contralateral projection arises more heavily from the medial limb (Helfert and Aschoff 1997). The opposite bias is true in ferrets (Henkel and Brunso-Bechtold 1993); the significance of this species difference remains to be determined. In any case, the crossed and uncrossed projections help to establish what Glendenning and colleagues (1981) called an “acoustic chiasm,”

referring to the fact that most cells in the midbrain and above respond to sounds in the contralateral sound field. The contralateral projection from the LSO is excitatory, whereas the ipsilateral projection contains a large glycinergic component and a smaller, presumably excitatory, component (Saint Marie et al. 1989). A small, bilateral projection may also originate from GABAergic cells (González-Hernández et al. 1996). Both the ipsilateral and contralateral projections terminate in bands in the ICc and in the deeper layers of the dorsal cortex of the IC (reviewed by Oliver 2000).

The LSO and MSO are widely considered as the first site of binaural convergence, even though binaural effects are observable in the CN and the cochlea. However, the LSO and MSO are the “lowest” nuclei to exhibit exquisite sensitivity to interaural disparities. These nuclei, along with their inputs and outputs, form the core of the binaural system and play an essential role in sound localization.

### 2.3. THE PERIOLIVARY NUCLEI

The periolivary nuclei (PON) are a group of nuclei surrounding the LSO and the MSO. They contain a variety of cell types that vary in morphology, inputs, projections, immunocytochemical properties, and physiologic responses (Schwartz 1992; Helfert and Aschoff 1997). Collectively, the PON are a significant source of projections to the IC (Fig. 4.2A). We first describe projections of the PON in general and then consider the SPN in detail.

Ascending inputs to the PON originate primarily from multipolar, octopus, and globular bushy cells of the VCN (reviewed by Thompson and Schofield 2000). Additional ascending inputs to particular PON can originate from other SOC nuclei.

All the periolivary nuclei project to the ipsilateral IC (Fig. 4.2B) (Adams 1983). Many of the nuclei, particularly the LNTB, send a smaller projection to the contralateral IC as well. Little is known regarding termination patterns in the IC. Ipsilateral projections from all nuclei appear to terminate in the central nucleus, and there is apparently a small projection to the dorsal and external cortices (González-Hernández et al. 1996).

There have been few studies of the responses of periolivary cells to acoustic stimuli (Guinan et al. 1972a,b; Irvine, 1986; Finlayson et al. 1997; Kuwada and Batra 1999). Binaural effects are common, and include various combinations of excitation and inhibition driven from either ear. Guinan et al. (1972a,b) distinguished 10 types of responses; in general, each unit type could be found among multiple nuclei, and each nucleus contained units of different response types. The heterogeneity suggests that the periolivary nuclei send a variety of signals to higher levels.

In few cases are the projections of the recorded PON cells known (Irvine 1986; Helfert and Aschoff 1997). The latter comment applies to principal cells of the MNTB and to medial olivocochlear cells; because neither of these cell types project to the IC, these studies provide no insight into the responses of




Figure 4.2. Periolivary projections to the inferior colliculus. **(A)** Graph showing the distribution of labeled cells in the SOC after injection of a retrograde tracer into one IC in a guinea pig (Schofield and Cant 1992). Cells in the PON were totaled to illustrate that olivocollicular cells in the PON can outnumber those in the principal nuclei (MSO and LSO). **(B)** Summary diagram showing general pattern of periolivary projections to the IC. *Line thickness* indicates in relative terms the number of cells giving rise to each projection (transmitters are not indicated in this figure). **(C)** Schematic diagram showing the major inputs to the SPN.

olivocollicular cells. One exception to this rule is the SPN, which provides a substantial projection to the IC and is discussed in detail later.

Several neurotransmitters have been implicated in periolivary projections to the IC. GABA is present in ipsilaterally projecting cells in the SPN (Kulesza and Berrebi, 2000) and in “other periolivary nuclei” (apparently including both VNTB and LNTB; González-Hernández et al. 1996) in rats. Presumptive glycinergic cells projecting to the ipsilateral IC have been identified in the SPN (guinea pig and chinchilla; Saint Marie and Baker, 1990) and, in smaller numbers, in the LNTB and MNTB (chinchilla; Saint Marie and Baker 1990). Other prospectively neuroactive substances, such as somatostatin and enkephalin, are associated with PON cells, but it is not known if they are present in olivocollicular cells. These substances are considered neuromodulators and could play an important role in modifying activity in the IC.

#### 2.4. THE SUPERIOR PARAOLIVARY NUCLEUS

Inputs to the SPN are distinctly different from those to the MSO and LSO. The largest input arises from multipolar (Mp) and octopus (Oct) cells of the contralateral VCN, with additional inputs from Mp cells in the ipsilateral VCN (Fig. 4.2C; Schofield 1995). Additional inputs arrive from the ipsilateral MNTB and MSO (Kuwabara and Zook 1992; Smith et al. 1998).

Several reports describe response properties of SPN units (rat: Finlayson and Adam, 1997; Kulesza et al. 2003; rabbit: Kuwada and Batra 1999; gerbil: Behrend et al. 2002; Dehmel et al. 2002). Common denominators are a clear best frequency, and that the units appeared to be arranged tonotopically. There are considerable differences in the reported patterns of responses, with offset responses prevailing in some studies and rare in others. In most studies, monaural responses were more common than binaural responses. The binaural responses are of particular interest for the present discussion because they were quite different from those typical of the classic "binaural" nuclei (i.e., MSO and LSO). Few SPN cells are reported as sensitive to interaural differences of time or intensity (a hallmark of MSO and LSO units) (Behrend et al. 2002). Furthermore, many SPN units have transient responses (at onset or offset) and respond well to sinusoidally amplitude-modulated tones. These properties have led to a general conclusion that SPN units are specialized for analysis of temporal features of sound (Behrend et al. 2002; Kulesza et al. 2003).

Most SPN cells appear to be inhibitory. In rats, potentially all SPN cells are GABAergic (Kulesza and Berrebi 2000); in other species, there also appear to be glycinergic cells (Helfert et al. 1989). Both GABAergic and glycinergic cells appear to project to the IC (Saint Marie and Baker 1990; González-Hernández et al. 1996).

The SPN sends a large projection to the ipsilateral IC (rat: Coleman and Clerici 1987; mouse: Ollo and Schwartz 1979; Willard and Ryugo 1983; gerbil: Nordeen et al. 1983); in some species, a small contralateral projection is also present (guinea pig: Schofield 1991; chinchilla: Saint Marie and Baker 1990; North American opossum: Willard and Martin 1983). The projections appear to be organized tonotopically (Kelly et al. 1998; Saldaña and Berrebi 2000). Although there is general agreement that the projections terminate in the central nucleus, there are reports of input to the dorsal cortex and external cortex (rat: González-Hernández et al. 1996; Coleman and Clerici 1987; guinea pig and gerbil: Cant et al. 1999).

Several authors have distinguished the SPN from other PON (González-Hernández et al. 1996; Saldaña and Berrebi 2000) and the latter authors have stressed the similarities of the SPN to other principal nuclei. In rats (the only species for which cell counts are available), the number of SPN neurons supports this argument (Table 4.1). SPN contains about 2400 neurons per side, compared to 2500 neurons in the LSO and 1100 in MSO. Because virtually all SPN neurons project to the IC in rats (Saldaña and Berrebi 2000), the SPN certainly qualifies as a substantial source of projections to the IC.

Table 4.1. Summary of the number of neurons in six nuclei in the SOC of the rat.

Nucleus	Number of neurons <sup>a</sup>	Percentage
MNTB	6000	32.2
MSO	1100	5.9
LSO	2500	13.4
SPN	2400	12.9
VNTB	4500	24.2
LNTB	2100	11.3
Total	18,600	100

<sup>a</sup>Unbiased stereologic estimates of the number of neurons, rounded to nearest hundred, in each nucleus. (From Kulesza et al. 2002.)


Is the SPN equivalent to the dorsomedial periolivary nucleus (DMPO)? A prominent SPN is present in many animals, including rat, guinea pig, gerbil, mouse, mole, North American opossum, and chinchilla (Willard and Martin 1983; Willard and Ryugo 1983; Kudo et al. 1990; Saint Marie and Baker 1990; Schwartz 1992). In other animals, a much less prominent nucleus in the same location has been called the DMPO (e.g., cat, ferret, primate; Moore 1988; Schwartz 1992). Kuwabara et al. (1991) suggest that the SPN and the DMPO are distinct: the DMPO receives input from GB cells of the contralateral CN (cf. Morest, 1968; cat), whereas such inputs are absent in the SPN.

In summary, the SPN, and the PON overall, are a significant source of projections to the IC. Most of these projections terminate ipsilaterally and appear to use inhibitory neurotransmitters. Where do the PON fit into the monaural/binaural scheme? The PON have traditionally been included with the MSO and LSO within the binaural system. Without question, several of the PON play essential roles in the binaural pathways; the best known example is the MNTB and its projections to the MSO and LSO. We believe that the MNTB plays an essential role in the monaural pathways as well, and that some PON should be considered components of both monaural and binaural systems. Other nuclei, such as the SPN, are more appropriately considered part of the monaural system. We return to these issues after reviewing the nuclei of the lateral lemniscus.


### 3. NUCLEI OF THE LATERAL LEMNISCUS

The lateral lemniscus contains ventral, intermediate, and dorsal nuclei (VNLL, INLL, and DNLL, respectively) that differ in cytoarchitecture, connections, and immunocytochemical properties. DNLL and VNLL are recognized across species. The INLL is sometimes considered a dorsal division of the VNLL (see discussion in Oertel and Wickesberg 2002), but we will treat it as a separate structure. The VNLL is often further divided into two or more parts (Fig 4.3A).


**A Lemniscal nuclei**



**B VNLL**



**C INLL**



**D DNLL**





Figure 4.3. Lateral lemniscal nuclei and their connections. (A) Schematic diagram of the NLL. (B–D) Schematic diagrams showing the inputs and collicular projections of the two major subdivisions of the VNLL (B), the INLL (C), and the DNLL (D). Conventions as in Fig. 4.2. #, includes GABAergic and excitatory projections; \*, includes GABAergic and glycinergic projections.

We will discuss two large subdivisions, dorsal (VNLLd) and ventral (VNLLv), as described in guinea pigs (Schofield and Cant, 1997).

### 3.1. VENTRAL NUCLEUS OF THE LATERAL LEMNISCUS

The VNLL receives bilateral inputs from the VCN and ipsilateral input from the MNTB (Fig. 4.3B; Oertel and Wickesberg 2002). Note that, although most inputs are shared by the two subdivisions of the VNLL, VCN octopus cells project only to the contralateral VNLLv. The octopus cell axons have large endings (Adams 1997; Schofield and Cant 1997) and presumably play an important role in distinguishing the response properties of cells in the two subdivisions. Additional inputs (not shown) arise in several periolivary nuclei (Glendenning et al. 1981).

In contrast to most other brain stem auditory nuclei, there is no clear tonotopic order in the VNLL (Glendenning and Hutson 1998). Most VNLL cells are excited by contralateral stimulation; many are also affected by ipsilateral stimulation, and are thus binaural (Batra and Fitzpatrick 2002). Some binaural cells are sensitive to interaural time disparities, but the transient responses of these cells distinguish them from MSO cells. A variety of response patterns have been described, including onset, offset, and sustained.

The VNLL of some bats is divided into two parts with distinct cell types and response properties (Covey and Casseday 1991). The multipolar subdivision has primarily sustained responses, whereas the columnar division has onset responses. Furthermore, these cells appear to be exclusively monaural. These properties may be particularly important for bat echolocation, and may explain the specialized appearance of the VNLL in bats. However, the basic analyses, focused on temporal processing, are likely to be useful to nonecholocating animals as well. In fact, proposed functions for the VNLL (and INLL) include “timing” (Glendenning and Hutson 1998), pattern recognition (Oertel and Wickesberg 2002), and duration and delay tuning (Casseday et al. 2002). The emphasis on temporal features reflects, in part, the inputs from octopus cells, which are specialized to respond as onset detectors across a wide range of frequencies (Trussell 2002).

A majority of VNLL cells appear to use glycine or GABA as a neurotransmitter, suggesting that the VNLL provides an inhibitory projection to the IC (Thompson et al. 1985; Saint Marie and Baker 1990; Winer et al. 1995). In addition, a smaller number of cells may use the excitatory neurotransmitter glutamate (Saint Marie 1996). It is interesting that these presumptive glutamatergic cells are located medially in the VNLL (Fig. 3 of Saint Marie 1996), in a region that Batra and Fitzpatrick (1997) have characterized as containing a large percentage of binaural cells that are sensitive to interaural time disparities.

Numerous studies have identified the VNLL as the brain stem nucleus that contains the greatest number of labeled cells after injection of a retrograde tracer into the IC (e.g., Brunso-Bechtold et al. 1981). The projections from the VNLL are diffuse (as opposed to banded) and are widespread, terminating heavily in



the central nucleus and dorsal cortex and less so in the lateral nucleus (Whitley and Henkel 1984; reviewed by Helfert and Aschoff 1997). These projections are almost exclusively ipsilateral. The many GABAergic and glycinergic cells in this nucleus make the VNLL the largest single source of inhibition to the IC.

### 3.2. *THE INTERMEDIATE NUCLEUS OF THE LATERAL LEMNISCUS*

The INLL receives input primarily from the contralateral VCN and the ipsilateral MNTB (Fig. 4.3C) (Oertel and Wickesberg 2002).

The most detailed information on INLL responses comes from studies on bats (Covey and Casseday 1991). INLL units are excited by contralateral stimulation and respond throughout the stimulus duration. These properties resemble those of the multipolar subdivision of the VNLL in the same bats (discussed earlier), leading Covey and Casseday (1999) to conclude that these two nuclei convey to the IC similar information about stimulus duration and intensity.

Immunohistochemical studies suggest that INLL contains some cells that use GABA or glycine, but most probably use an excitatory neurotransmitter.

The projection from the INLL to the IC is largely ipsilateral and terminates diffusely in the central nucleus and dorsal cortex (Whitley and Henkel 1984). The pathway appears to be organized tonotopically (Zook and Casseday 1982).

Both the VNLL and the INLL have been implicated in temporal processing. As discussed later, this broad category of analysis has been applied generally to the monaural system, of which INLL and VNLL are considered major components. In contrast, the DNLL differs in its inputs and its response properties, and is considered part of the binaural system.

### 3.3. *THE DORSAL NUCLEUS OF THE LATERAL LEMNISCUS*

The DNLL receives inputs bilaterally from the VCN and SOC (Fig. 4.3D). The olivary inputs arise from the LSO on both sides and from the MSO on the same side (see Pollak et al. 2003). Additional inputs may arise from the ipsilateral VNLL and INLL (not shown). All ascending axons reach the DNLL via the lateral lemniscus. An additional input arises from the contralateral DNLL, whose axons arrive via the commissure of Probst.

Most DNLL cells are binaural, and excited by the contralateral ear and inhibited by the ipsilateral ear (EI cells). A small number of cells are binaurally excited. DNLL appears to be organized tonotopically, although the precise arrangement may vary between species (Helfert and Aschoff 1997).

The DNLL contains primarily, perhaps exclusively, GABAergic cells (Adams and Mugnaini 1984). These cells comprise separate populations that project to the ipsilateral or contralateral IC and are organized tonotopically. In the contralateral IC, DNLL axons form bands that terminate in the central nucleus and extend into the dorsal cortex. The projection terminates more diffusely ipsilaterally (Shneiderman et al. 1988; reviewed by Helfert and Aschoff 1997). Projections to external cortex (lateral nucleus) may be present.

As mentioned earlier, the DNLL has been associated with binaural processing and sound localization. In the next section, we review the monaural and binaural systems and consider how recent data impact these ideas.

#### 4. ORGANIZATION OF THE ASCENDING PATHWAYS

The ascending pathways to the IC have been conceptualized as monaural and binaural systems (Kudo and Nakamura 1987; Pollak and Casseday 1989; Helfert and Aschoff 1997; Casseday et al. 2002). In most accounts, the monaural system has included the direct projections from the CN as well as indirect projections from multipolar and octopus cells in the VCN through the VNLL and INLL. The binaural system has included spherical bushy and globular bushy cells in the VCN as well as the nuclei of the SOC (particularly the MSO, LSO, and MNTB) and the DNLL. We present here a modified view of these systems, taking into account new data, particularly concerning the SPN and the VNLL. While maintaining the concept of monaural and binaural systems, we propose to dissect the SOC further and to consider its components individually. Some SOC nuclei belong to the monaural system and others to the binaural system. Further, some nuclei, and some CN cell types, should be considered common elements of both systems.




Figure 4.4. Summary of the binaural pathways through the SOC and NLL to the inferior colliculus. Conventions as in Fig. 4.2.

#### 4.1. NUCLEI OF THE BINAURAL SYSTEM

Our view of the binaural system is similar to previous proposals except that we do not include the entire SOC (Fig. 4.4). The key nuclei are the MSO, LSO, and DNLL. The inputs originate from spherical bushy and globular bushy cells in the VCN. Cells in both the MNTB and LNTB contribute to binaural as well as monaural pathways. The binaural pathways include crossed and uncrossed projections from the SOC and provide for both excitation and inhibition (the latter including both GABAergic and glycinergic projections) in the IC.

#### 4.2. NUCLEI OF THE MONAURAL SYSTEM


The INLL and VNLL are customarily included with direct projections from the CN to the IC as part of the monaural system. In an earlier study, we discussed numerous similarities between the SPN and VNLL (Schofield and Cant 1997), parallels that led Helfert and Aschoff (1997) to include the SPN (which they called the dorsomedial periolivary nucleus) with the monaural system. Subsequent work has expanded the list of similarities between SPN and VNLL. Like the VNLL, the SPN has widespread projections to all IC subdivisions (Cant et al. 1999). Inputs to both nuclei arise largely from the contralateral CN, and less so from the ipsilateral CN and SOC. The CN inputs arise from multipolar cells bilaterally and from octopus cells contralaterally. Both nuclei have a preponderance of cells with monaural responses. Furthermore, the binaural responses in each of these nuclei are different from those in the binaural system (i.e., MSO, LSO, and DNLL). Not surprisingly, the data have led to similar conclusions regarding functions of the SPN and the VNLL.

The monaural pathways include the direct projections from the CN (Fig. 4.5A; see Chapter 3) and consist of SPN, VNLL, and INLL. Note the variety of VCN cell types involved; of these, the multipolar and octopus cells are most prominent in the monaural pathways. Most information arriving at the IC via these pathways originates from the contralateral ear. The monaural pathways, like the binaural ones, provide excitatory as well as inhibitory projections to the IC.

#### 4.3. NUCLEI OF THE MONAURAL AND BINAURAL SYSTEMS

The MNTB has often been considered a principal nucleus of the SOC, and is generally considered part of the binaural system. Unlike the MSO and the LSO, the MNTB is not a major source of projections to the IC. In fact, the small projection it does send to the IC originates from nonprincipal cells (Schofield 1994), about which we know very little. The MNTB principal cells, in contrast, are well known as the targets of globular bushy cell calyces, and as the source of projections to the MSO and LSO. The principal cells also project to each of the nuclei associated with the monaural system as well as to the VCN (Fig. 4.6; Spangler et al. 1985; Kuwabara and Zook 1992; Schofield 1994; Smith et al. 1998). It seems appropriate to consider the MNTB (or at least the majority of

**A Direct monaural pathways**



**B Indirect monaural pathways**




Figure 4.5. Summary of monaural pathways to the IC. (A) Direct projections from the cochlear nuclei (see Chapter 3). (B) Indirect monaural projections through the SOC and NLL. Conventions as in Fig. 4.2.

its neuronal population, the principal cells) as contributing to both the monaural and binaural systems, and not to assign it exclusively to one or the other.

**4.4. CONNECTIONS BETWEEN THE MONAURAL AND BINAURAL SYSTEMS**

There are multiple points of interaction between the monaural and binaural systems. VCN multipolar cells are probably the most numerous single input to the monaural pathways, but they also project to the LSO (Doucet and Ryugo 2003) and they are probably the source of VCN projections to the DNLL. Both VNLL and INLL project to the DNLL (Glendenning et al. 1981). Thus, there are several




Figure 4.6. Schematic diagram showing projections from MNTB principal cells and non-principal cells. Nuclei associated with the monaural pathways are shown in *dark gray* and binaural pathways in *light gray*.

unexpected instances of “monaural” components projecting to binaural nuclei. There are also examples of the converse arrangement, that is, binaural inputs to monaural nuclei. For example, spherical bushy cells project not only to the MSO and LSO but also to the VNLL. Finally, the MSO projects to the SPN. Identifying the functions of these interactions will be important for understanding the ascending systems and should temper categorical assignments of specific cell types to particular systems with more caution than they ordinarily receive.

#### 4.5. PROPOSED FUNCTIONS OF MONAURAL AND BINAURAL SYSTEMS

The direct projections from the CN are the largest source of ascending excitatory inputs to the IC. The indirect monaural pathways provide excitatory inputs as well as the largest source of ascending inhibitory inputs to the IC (via the VNLL). The binaural system also gives rise to excitatory and inhibitory projections. Despite their interconnections, the differences in physiologic responses of the monaural vs. the binaural nuclei indicate that the two systems send different information to the IC. For each system, the proposed functions are suggested by the physiologic properties of both the constituent nuclei and those of the CN cells that provide inputs.

The binaural pathways have long been considered integral to analysis of interaural differences that are important for sound localization (particularly with regard to azimuth). A question that arises (for both binaural and monaural systems) is, Why are there so many nuclei and multiple, seemingly redundant,

pathways? In the binaural system, both the LSO and the DNLL contain many EI cells with inhibitory projections to the ipsilateral IC (Fig. 4.4). Pollak et al. (2003) demonstrate that these projections are not redundant. In fact, the DNLL projection confers new binaural properties on the target IC cells, allowing the IC cells to respond to stimuli to which they would not otherwise respond. These new properties could contribute to a variety of functions, such as analysis of stimulus motion.

Functional roles for the monaural pathways have generally received less attention than those of the binaural system. The VNLL and INLL have been implicated in “timing” (Glendenning and Hutson 1998), pattern recognition (Oertel and Wickesberg 2002), and temporal processing and duration and delay tuning (Casseday et al. 2002). The latter authors discuss mechanisms by which the VNLL and INLL could contribute to tuning for sound duration, rate of frequency or amplitude modulation, delay between an echolocation pulse and its echo, and selectivity for upward or downward frequency modulations. The similarities between the SPN and the VNLL lead to the conclusion that the SPN may be involved in some of these same functions. Both the SPN and the VNLL are substantial sources of inhibitory projections to the ipsilateral IC. Are these pathways redundant? We hypothesize that, like the DNLL and LSO projections in the binaural system, the SPN and VNLL projections confer different properties on their targets in the IC. Future experiments will be needed to test this idea.

## 5. RELATIONSHIP BETWEEN AFFERENTS AND THE INFERIOR COLLICULUS

### 5.1. LARGE-SCALE ORDER: NUCLEOTOPIC ORGANIZATION

The majority of data pertinent to this issue are from the ICc, and the following discussion is focused on that division. Results from both anterograde and retrograde tracing studies indicate that some projections to the ICc terminate more broadly than others. Projections from the VNLL, for example, terminate widely in ICc. MSO and LSO projections are more dense rostrally, whereas those from the CN are more dense caudally. This organization has been called “nucleotopic” (Brunso-Bechtold et al. 1981). In addition, the ICc itself and most of its inputs are organized tonotopically. Thus, projections from the low-frequency-biased MSO target the low-frequency (dorsal) part of ICc, whereas those from the LSO are denser in the high-frequency (ventral) regions. The consequence of nucleotopic and tonotopic organization is that different parts of the ICc receive different combinations of inputs, leading to local order and small regions with distinct circuitry.

### 5.2. LOCAL ORDER AND MICROCIRCUITRY

A frequent finding in physiologic studies is the clustering of units with similar response types (Roth et al. 1978). The scale of this clustering is much smaller

and finer than that of nucleotopic order discussed earlier, and occurs within an isofrequency lamina. Furthermore, the responses of many clusters are similar to those of lower centers, suggesting that IC cells simply “mirror” a dominant input. This applies particularly to temporal response patterns (e.g., onset vs. sustained) and to binaural interactions. Interestingly, some IC cells may, in fact, mirror a single, dominant input. Aitkin and Schuck (1985) recorded from a small low-frequency region of the IC in which the cells had response properties resembling those of the MSO. A small injection of retrograde tracer into this region labeled cells primarily in the MSO, consistent with the idea that the IC cells were mirroring MSO inputs. Using different techniques, Davis (2002) suggested that some IC cells are dominated by input from the contralateral dorsal cochlear nucleus (DCN). These projections are part of the monaural system, and thus there have been suggestions of mirroring in both monaural and binaural systems.

However, “mirroring” cannot explain all instances of similar responses. Kuwada et al. (1997) have shown that the same temporal discharge pattern can be generated by the intrinsic membrane properties of a neuron or by convergent excitatory and inhibitory inputs as well as by mirroring. The significance is that different mechanisms could result in different responses to more complex stimuli; for example, different types of onset units should respond differently in a reverberant environment. A similar situation applies to binaural properties. An EI cell (excited by the contralateral ear and inhibited by the ipsilateral ear) may mirror an EI input or integrate inputs that carry the excitation and inhibition separately (Pollak et al. 2003).

Several patterns of convergence have been suggested by experiments combining physiologic with anatomical and/or pharmacologic methods. IC cells could receive convergent input from the contralateral DCN and the ipsilateral MSO, LSO, or DNLL, or the contralateral DNLL (Semple and Aitkin 1981; Aitkin and Schuck 1985). Inputs from the ipsilateral MSO have been suggested to converge with those from ipsilateral LSO and ipsilateral or contralateral DNLL (Aitkin and Schuck 1985; Vater et al. 1995; Kuwada et al. 1997). Finally, inputs from the contralateral DNLL may converge with a contralateral monaural input (likely arising from DCN or VCN; Pollak et al. 2003). Anatomical tracing after very small injections in the IC suggest additional possibilities (Kudo and Nakamura 1987; Ross and Pollak 1989). Understanding and testing these possibilities, which essentially represent functional subdomains, will require a clear picture of circuitry within a lamina.

An important insight into microcircuitry followed from the discovery that several large projections to the IC terminate in bands. As described earlier, projections from the CN, MSO, LSO, and DNLL are banded whereas those from the VNLL and INLL are not (patterns for the PON are not known). The bands are on the order of a few hundred micrometers wide, providing a small-scale order of segregation (or overlap or periodicity) of specific inputs. Studies employing two different anterograde tracers have shown that the bands from the ipsilateral LSO and the contralateral LSO are interdigitated (Shneiderman and Henkel 1987). A disc-shaped cell situated in the middle of a band may receive just one of the inputs whereas a stellate cell, with dendrites that span

multiple bands, could receive convergent inputs. The detailed arrangement of bands from different sources, and their overlap with nonbanded inputs, gives rise to small regions, “synaptic domains,” with unique patterns of microcircuitry (see Chapter 2).

A great deal of work remains to be done to identify the various patterns of convergence, to understand how they give rise to IC response patterns, and to recognize their contributions to auditory function.

## 6. FUTURE DIRECTIONS

We have reviewed the olivary and lemniscal inputs to the IC, and have suggested some revisions to the classic dichotomous scheme of monaural and binaural categorical pathways. These pathways ascend in parallel to the direct projections from the CN, which provide the IC with a massive, excitatory drive. The indirect pathways through the SOC and the NLL provide additional excitatory inputs as well as major inhibitory inputs that use both GABA and glycine. We conclude by enumerating a few of the questions that still remain.

1. To what extent, and to what purpose, do the monaural and binaural pathways, and their individual components, converge in the IC?
2. With respect to the principal SOC nuclei, what function is served by contralateral, or bilateral, projections from individual MSO cells? What purpose might be served by excitatory projection from the LSO to the ipsilateral IC: does it exist, and if so, what does it do? What function do binaurally excited LSO cells serve?
3. Does the SPN serve a unique function? What cells serve the same purpose in animals without an SPN? Is the SPN absent in some species? What about projections from other PON?
4. MNTB principal cells are well studied and are the source of the connections reviewed in the preceding (Fig. 4.6). However, it is the nonprincipal cells in the MNTB that project directly to the IC (Schofield 1994). These cells probably receive input from the contralateral VCN and thus form a small, disynaptic pathway from the VCN to the IC. A similar disynaptic pathway appears to traverse other periolivary nuclei, including the dorsal periolivary nucleus, the VNTB, and the LNTB (Thompson and Schofield 2000). Very little is known about the physiologic properties of the “periolivocollicular” cells, so it is premature to assign them to monaural or binaural systems or to reach definitive conclusions as to their role.

We have focused on the ICc because most of the available data concern that subdivision. Difficulties in establishing criteria for distinguishing the IC subdivisions have complicated efforts to understand it. Such criteria will be essential before we can more fully understand IC function.

5. Can identification of SOC and NLL cell types provide functional insight, as it has for the VCN and for olivocochlear cells?



6. How do the ascending pathways converge, merge, and segregate from intrinsic and commissural projections (see Chapter 5), and how are these many pathways integrated with information from descending projections from the auditory cortex and thalamus (Chapter 8)?
7. How do the various ascending pathways relate to the outputs of the IC? This issue appears to be underappreciated. The ascending IC projections are often viewed from the perspective of transmission of information to the thalamus and thus to cortex. The IC projects to numerous nonthalamic targets, including the SOC and CN, reticular formation, and superior colliculus and pontine nuclei, and has motor as well as sensory functions (see Chapter 6). It seems likely that the diversity of physiologic properties characteristic of IC cells may be a reflection of the different functions served by the IC. One could predict that different pathways comprise units with different response properties (or at least differing proportions of the major response types).

## Acknowledgments

We thank Nell Cant, Diana Coomes, and Ark Slusarczyk for helpful comments. This work was supported by NIH Grant DC04391.

## Abbreviations

CN	cochlear nucleus
DCN	dorsal cochlear nucleus
DLPO	dorsolateral periolivary nucleus
DMPO	dorsomedial periolivary nucleus
DNLL	dorsal nucleus of the lateral lemniscus
DPO	dorsal periolivary nucleus
EE	excited by both ears
EI	excited by contralateral ear, inhibited by ipsilateral ear
GABA	$\gamma$ -aminobutyric acid
GB	globular bushy cell of the ventral cochlear nucleus
IC	inferior colliculus
ICc	central nucleus of the inferior colliculus
INLL	intermediate nucleus of the lateral lemniscus
LNTB	lateral nucleus of the trapezoid body
LSO	lateral superior olivary nucleus
MNTB	medial nucleus of the trapezoid body
Mp	multipolar cell of the ventral cochlear nucleus
MSO	medial superior olivary nucleus
NLL	nuclei of the lateral lemniscus
Oct	octopus cell of the ventral cochlear nucleus

PON	periolivary nuclei
SB	spherical bushy cell of the ventral cochlear nucleus
SOC	superior olivary complex
SPN	superior paraolivary nucleus
VCN	ventral cochlear nucleus
VNLL	ventral nucleus of the lateral lemniscus
VNLLd	dorsal division of ventral nucleus of the lateral lemniscus
VNLLv	ventral division of ventral nucleus of the lateral lemniscus
VNTB	ventral nucleus of the trapezoid body

## References

- Adams JC (1983) Cytology of periolivary cells and the organization of their projections in the cat. *Journal of Comparative Neurology* **215**:275–289.
- Adams JC (1997) Projections from octopus cells of the posteroventral cochlear nucleus to the ventral nucleus of the lateral lemniscus in cat and human. *Auditory Neuroscience* **3**:335–350.
- Adams JC and Mugnaini E (1984) Dorsal nucleus of the lateral lemniscus: a nucleus of GABAergic projection neurons. *Brain Research Bulletin* **13**:585–590.
- Aitkin L and Schuck D (1985) Low frequency neurons in the lateral central nucleus of the cat inferior colliculus receive their input predominantly from the medial superior olive. *Hearing Research* **17**:87–93.
- Batra R and Fitzpatrick DC (1997) Neurons sensitive to interaural temporal disparities in the medial part of the ventral nucleus of the lateral lemniscus. *Journal of Neurophysiology* **78**:511–515.
- Batra R and Fitzpatrick DC (2002) Processing of interaural temporal disparities in the medial division of the ventral nucleus of the lateral lemniscus. *Journal of Neurophysiology* **88**:666–675.
- Behrend O, Brand A, Kapfer C, and Grothe B (2002) Auditory response properties in the superior paraolivary nucleus of the gerbil. *Journal of Neurophysiology* **87**:2915–2928.
- Brunso-Bechtold JK, Thompson GC, and Masterton RB (1981) HRP study of the organization of auditory afferents ascending to central nucleus of inferior colliculus in cat. *Journal of Comparative Neurology* **197**:705–722.
- Cant NB, Benson CG, and Schofield BR (1999) Projections from the superior paraolivary nucleus to the inferior colliculus in guinea pigs and gerbils. *Association for Research in Otolaryngology Abstracts* **19**:221–222.
- Casseday JH, Fremouw T, and Covey E (2002) The inferior colliculus: a hub for the central auditory system. In: Oertel D, Fay RR, and Popper AN (eds). *Springer Handbook of Auditory Research*, Volume 15: *Integrative Functions in the Mammalian Auditory Pathway*. Springer-Verlag, New York, pp. 238–318.
- Coleman JR and Clerici WJ (1987) Sources of projections to subdivisions of the inferior colliculus in the rat. *Journal of Comparative Neurology* **262**:215–226.
- Covey E and Casseday JH (1991) The monaural nuclei of the lateral lemniscus in an echolocating bat: parallel pathways for analyzing temporal features of sound. *Journal of Neuroscience* **11**:3456–3470.

- Covey E and Casseday JH (1999) Timing in the auditory system of the bat. *Annual Review of Physiology* **61**:457–476.
- Davis KA (2002) Evidence of a functionally segregated pathway from dorsal cochlear nucleus to inferior colliculus. *Journal of Neurophysiology* **87**:1824–1835.
- Dehmel S, Kopp-Scheinpflug C, Dorrscheidt GJ, and Rübnsamen R (2002) Electrophysiological characterization of the superior paraolivary nucleus in the Mongolian gerbil. *Hearing Research* **172**:18–36.
- Doucet JR and Ryugo DK (2003) Axonal pathways to the lateral superior olive labeled with biotinylated dextran amine injections in the dorsal cochlear nucleus of rats. *Journal of Comparative Neurology* **461**:452–465.
- Finlayson PG and Adam TJ (1997) Excitatory and inhibitory response adaptation in the superior olive complex affects binaural acoustic processing. *Hearing Research* **103**:1–18.
- Glendenning KK and Hutson KA (1998) Lack of topography in the ventral nucleus of the lateral lemniscus. *Microscopy Research and Technique* **41**:298–312.
- Glendenning KK, Brunso-Bechtold JK, Thompson GC, and Masterton RB (1981) Ascending auditory afferents to the nuclei of the lateral lemniscus. *Journal of Comparative Neurology* **197**:673–703.
- González-Hernández T, Mantolán-Sarmiento B, González-González B, and Pérez-González H (1996) Sources of GABAergic input to the inferior colliculus of the rat. *Journal of Comparative Neurology* **372**:309–326.
- Guinan JJ Jr, Guinan SS, and Norris BE (1972a) Single auditory units in the superior olivary complex I. Responses to sounds and classification based on physiological properties. *International Journal of Neuroscience* **4**:101–120.
- Guinan JJ Jr, Norris BE, and Guinan SS (1972b) Single auditory units in the superior olivary complex II. Locations of unit categories and tonotopic organization. *International Journal of Neuroscience* **4**:147–166.
- Helfert RH and Aschoff A (1997) Superior olivary complex and nuclei of the lateral lemniscus. In: Ehret G and Romand R (eds). *The Central Auditory System*. Oxford University Press, New York, pp. 193–258.
- Helfert RH, Bonneau JM, Wenthold RJ, and Altschuler RA (1989) GABA and glycine immunoreactivity in the guinea pig superior olivary complex. *Brain Research* **501**:269–286.
- Henkel CK and Brunso-Bechtold JK (1993) Laterality of superior olive projections to the inferior colliculus in adult and developing ferret. *Journal of Comparative Neurology* **331**:458–468.
- Irvine DRF (1986) The auditory brainstem. A review of the structure and function of auditory brainstem processing mechanisms. In: Autrum H, Ottoson D, Perl ER, Schmidt RF, Shimazu H, and Willis WD (eds). *Progress in Sensory Physiology* **7**:1–279.
- Kelly JB, Liscum A, van Adel B, and Ito M (1998) Projections from the superior olive and lateral lemniscus to tonotopic regions of the rat's inferior colliculus. *Hearing Research* **116**:43–54.
- Kudo M and Nakamura K (1987) Organization of the lateral lemniscal fibers converging onto the inferior colliculus in the cat: an anatomical review. In: Syka J and Masterton RB (eds). *Auditory Pathway: Structure and Function*. Plenum Press, New York, pp. 171–183.
- Kudo M, Kitao Y, and Nakamura Y (1990) Differential organization of crossed and uncrossed projections from the superior olive to the inferior colliculus in the mole. *Neuroscience Letters* **117**:26–30.

- Kulesza RJ Jr, and Berrebi AS (2000) Superior paraolivary nucleus of the rat is a GABAergic nucleus. *Journal of the Association for Research in Otolaryngology* **1**:255–269.
- Kulesza RJ Jr, Spirou GA, and Berrebi AS (2003) Physiological response properties of neurons in the superior paraolivary nucleus of the rat. *Journal of Neurophysiology* **89**: 2299–2312.
- Kumoi K, Saito N, and Tanaka C (1993) Immunohistochemical localization of gamma-aminobutyric acid- and aspartate-containing neurons in the guinea pig superior olivary complex. *Hearing Research* **68**:173–179.
- Kuwabara N and Zook JM (1992) Projections to the medial superior olive from the medial and lateral nuclei of the trapezoid body in rodents and bats. *Journal of Comparative Neurology* **324**:522–538.
- Kuwabara N, DiCaprio RA, and Zook JM (1991) Afferents to the medial nucleus of the trapezoid body and their collateral projections. *Journal of Comparative Neurology* **314**: 684–706.
- Kuwada S and Batra R (1999) Coding of sound envelopes by inhibitory rebound in neurons of the superior olivary complex in the unanesthetized rabbit. *Journal of Neuroscience* **19**:2273–2287.
- Kuwada S, Batra R, Yin TCT, Oliver DL, Haberly LB, and Stanford TR (1997) Intracellular recordings in response to monaural and binaural stimulation of neurons in the inferior colliculus of the cat. *Journal of Neuroscience* **17**:7565–7581.
- Moore DR (1988) Auditory brainstem of the ferret: sources of projections to the inferior colliculus. *Journal of Comparative Neurology* **269**:342–354.
- Morest DK (1968) The collateral system of the medial nucleus of the trapezoid body of the cat, its neuronal architecture and relation to the olivo-cochlear bundle. *Brain Research* **9**:288–311.
- Nordeen KW, Killackey HP, and Kitzes LM (1983) Ascending auditory projections to the inferior colliculus in the adult gerbil, *Meriones unguiculatus*. *Journal of Comparative Neurology* **214**:131–143.
- Oertel D and Wickesberg RE (2002) Ascending pathways through ventral nuclei of the lateral lemniscus and their possible role in pattern recognition in natural sounds. In: Oertel D, Fay RR, and Popper AN (eds). *Springer Handbook of Auditory Research, Volume 15: Integrative Functions in the Mammalian Auditory Pathway*. Springer-Verlag, New York, pp. 207–237.
- Oliver DL (2000) Ascending efferent projections of the superior olivary complex. *Microscopy Research and Technique* **51**:355–363.
- Oliver DL, Beckius GE, and Shneiderman A (1995) Axonal projections from the lateral and medial superior olive to the inferior colliculus of the cat: a study using electron microscopic autoradiography. *Journal of Comparative Neurology* **360**:17–32.
- Ollo C and Schwartz IR (1979) The superior olivary complex in C57BL/6 mice. *American Journal of Anatomy* **155**:349–374.
- Pollak GD and Casseday JH (1989) *The Neural Basis of Echolocation in Bats*. Springer-Verlag, New York.
- Pollak GD, Burger RM, and Klug A (2003) Dissecting the circuitry of the auditory system. *Trends in Neurosciences* **26**:33–39.
- Ross LS and Pollak GD (1989) Differential ascending projections to aural regions in the 60 kHz contour of the mustache bat's inferior colliculus. *Journal of Neuroscience* **9**: 2819–2834.

- Roth GL, Aitkin LM, Andersen RA, and Merzenich MM (1978) Some features of the spatial organization of the central nucleus of the inferior colliculus of the cat. *Journal of Comparative Neurology* **182**:661–680.
- Saint Marie RL (1996) Glutamatergic connections of the auditory midbrain: selective uptake and axonal transport of D-[<sup>3</sup>H]aspartate. *Journal of Comparative Neurology* **373**:255–270.
- Saint Marie RL and Baker RA (1990) Neurotransmitter-specific uptake and retrograde transport of [<sup>3</sup>H]glycine from the inferior colliculus by ipsilateral projections of the superior olivary complex and nuclei of the lateral lemniscus. *Brain Research* **524**:244–253.
- Saint Marie RL, Ostapoff E-M, Morest DK, and Wenthold RJ (1989) Glycine-immunoreactive projection of the cat lateral superior olive: possible role in midbrain ear dominance. *Journal of Comparative Neurology* **279**:382–396.
- Saldaña E and Berrebi AS (2000) Anisotropic organization of the rat superior paraolivary nucleus. *Anatomy and Embryology* **202**:265–279.
- Schofield BR (1991) Superior paraolivary nucleus in the pigmented guinea pig: separate classes of neurons project to the inferior colliculus and the cochlear nucleus. *Journal of Comparative Neurology* **312**:68–76.
- Schofield BR (1994) Projections to the cochlear nuclei from principal cells in the medial nucleus of the trapezoid body in guinea pigs. *Journal of Comparative Neurology* **344**:83–100.
- Schofield BR (1995) Projections from the cochlear nucleus to the superior paraolivary nucleus in guinea pigs. *Journal of Comparative Neurology* **360**:135–149.
- Schofield BR and Cant NB (1997) Ventral nucleus of the lateral lemniscus in guinea pigs: cytoarchitecture and inputs from the cochlear nucleus. *Journal of Comparative Neurology* **379**:363–385.
- Schwartz IR (1992) The superior olivary complex and lateral lemniscal nuclei. In: Webster DB, Popper AN, and Fay RR (eds). *Springer Handbook of Auditory Research, Volume 1: The Mammalian Auditory Pathway: Neuroanatomy*. Springer-Verlag, New York, pp. 117–167.
- Semple MN and Aitkin LM (1981) Integration and segregation of input to the cat inferior colliculus. In: Syka J and Aitkin LM (eds). *Neuronal Mechanisms of Hearing*. Plenum Press, New York, pp. 155–161.
- Shneiderman A and Henkel CG (1987) Banding of lateral superior olivary nucleus afferents in the inferior colliculus: a possible substrate for sensory integration. *Journal of Comparative Neurology* **266**:519–534.
- Shneiderman A, Oliver DL, and Henkel CK (1988) Connections of the dorsal nucleus of the lateral lemniscus: an inhibitory parallel pathway in the ascending auditory system? *Journal of Comparative Neurology* **276**:188–208.
- Smith PH and Spirou GA (2002) Cochlea to cortex and back. In: Oertel D, Fay RR, and Popper AN (eds). *Springer Handbook of Auditory Research, Volume 15: Integrative Functions in the Mammalian Auditory Pathway*. Springer-Verlag, New York, pp. 6–71.
- Smith PH, Joris PX, and Yin TCT (1998) Anatomy and physiology of principal cells of the medial nucleus of the trapezoid body (MNTB) of the cat. *Journal of Neurophysiology* **79**:3127–3142.
- Spangler KM, Warr WB, and Henkel CK (1985) The projections of principal cells of the medial nucleus of the trapezoid body in the cat. *Journal of Comparative Neurology* **238**:249–262.

- Thompson AM and Schofield BR (2000) Afferents of the superior olivary complex. *Microscopy Research and Technique* **51**:330–354.
- Thompson GC, Cortez AM, and Lam DM-K (1985) Localization of GABA immunoreactivity in the auditory brainstem of guinea pigs. *Brain Research* **339**:119–122.
- Trussell LO (2002) Cellular mechanisms for information coding in auditory brainstem nuclei. In: Oertel D, Fay RR, and Popper AN (eds). *Springer Handbook of Auditory Research*, Volume 15: *Integrative Functions in the Mammalian Auditory Pathway*. Springer-Verlag, New York, pp. 72–98.
- Vater M, Casseday JH, and Covey E (1995) Convergence and divergence of ascending binaural and monaural pathways from the superior olives of the mustached bat. *Journal of Comparative Neurology* **351**:632–646.
- Whitley JM and Henkel CK (1984) Topographical organization of the inferior collicular projection and other connections of the ventral nucleus of the lateral lemniscus in the cat. *Journal of Comparative Neurology* **229**:257–270.
- Willard FH and Martin GF (1983) The auditory brainstem nuclei and some of their projections to the inferior colliculus in the North American opossum. *Neuroscience* **10**:1203–1232.
- Willard FH and Ryugo DK (1983) Anatomy of the central auditory system. In: Willott JF (ed). *The Auditory Psychobiology of the Mouse*. Charles C Thomas, Springfield, pp. 201–304.
- Winer JA, Larue DT, and Pollak GD (1995) GABA and glycine in the central auditory system of the mustache bat: structural substrates for inhibitory neuronal organization. *Journal of Comparative Neurology* **355**:317–353.
- Zook JM and Casseday JH (1982) Origin of ascending projections to inferior colliculus in the mustache bat, *Pteronotus parnellii*. *Journal of Comparative Neurology* **207**:14–28.

# Chapter 5

## Intrinsic and Commissural Connections of the Inferior Colliculus

ENRIQUE SALDAÑA AND MIGUEL A. MERCHÁN

### 1. INTRODUCTION

#### *1.1. INTRACOLLICULAR CONNECTIONS*

The inferior colliculus (IC) is regarded as a nearly obligatory relay center for the information ascending from subcollicular auditory centers. Few axons bypass the IC en route to the auditory thalamus (Ramón y Cajal 1904; Henkel 1983; Malmierca et al. 2002), and most lateral lemniscal fibers end in the IC. Because of these convergent projections, the complexity of the inputs to the IC (see Chapters 3 and 4) contrasts with the apparent simplicity of midbrain projections to the thalamus (see Chapter 7).

The rat IC contains approximately 350,000 neurons, almost five times that in auditory subcollicular nuclei, and five times the number in the medial geniculate body (MGB) (Kulesza et al. 2002). These numbers suggest that, before arriving at the MGB, auditory information is extensively processed in the IC. Such processing is shaped by interactions between IC neurons, which are mediated by intracollicular connections.

Intracollicular connections, which are the focus of this chapter, are monosynaptic interactions between ipsilateral or contralateral IC neurons. Connections between neurons in the same IC are intrinsic, whereas those between the colliculi are commissural and travel in the commissure of the IC (CoIC). In this account intrinsic and commissural connections are considered together, as most if not all IC neurons with commissural connections also have intrinsic connections. Moreover, intrinsic and commissural connections share similar patterns of distribution. The chapter also contains data about the organization of intracollicular connections of the cortical subdivisions of the rat IC, and the distribution of commissural IC neurons. All references are to the rat unless stated otherwise.

We use the now-classic cytoarchitectural parcellation of the rat IC (Faye-Lund and Osen 1985) with slight modifications (Saldaña and Merchán 1992). The IC contains a central nucleus (CN) surrounded dorsally and dorsocaudally by the dorsal cortex (DC), and rostrolaterally by the external cortex. For clarity, we




Figure 5.1. Micrographs of glutaraldehyde-fixed, 60- $\mu$ m-thick coronal sections through the rat midbrain tectum. Sections were postfixed with osmium tetroxide, which stained myelinated fibers black. The numbers at the bottom of the section indicate the approximate rostrocaudal coordinate of the section in the interaural coronal plane. Fiber fascicles of CoC are well defined and the IC neuropil is fiber rich, in contrast to the lightly stained periaqueductal gray matter. The *white arrowheads (left side)* indicate lateral lemniscal fascicles entering the IC.



refer to external cortex rostral to the CN as the rostral cortex (RC), and to the portion lateral to the CN as the external nucleus (EN).

## 1.2. THE COMMISSURE OF THE INFERIOR COLLICULUS

### 1.2.1. Anatomy and Topography

Despite subtle interspecies variations in volume and anatomical relationships, the main structural features of the CoIC are shared by most mammals (Figs. 5.1 and 5.2; see Chapter 22). The CoIC is a prominent transverse fiber tract that, at very caudal levels, consists of thin fascicles that constitute the roof of the recess of the fourth ventricle (Fig. 5.1C). Rostrally, it becomes a thick tract corresponding to the enlarged caudal portion of the tectal commissure (Faye-Lund and Osen 1985) and it is separated from the cerebral aqueduct by the thin rim of caudal periaqueductal gray matter (Fig. 5.1A, B). The CoIC is largest in the rostral third of the IC (Fig. 5.1A, B), where most commissural axons decussate. Whereas most CoIC axons project horizontally and in parallel, more caudal fascicles have variable, ascending or descending, oblique trajectories (Fig. 5.1C).

The rostral position of the CoIC is best seen in horizontal sections (Fig. 5.2) which show commissural axons traveling rostrally across the midline and then turning caudally in the contralateral IC. The horizontal plane shows that commissural axons cross in a nearly transverse plane (Fig. 5.2D), and that there is direct continuity between the rostroventral CoIC and the commissure of the superior colliculus (Fig. 5.2C).

### 1.2.2. Commissural Projections of the Inferior Colliculus

CoIC carries information bidirectionally (Held 1891, 1893; Moore and Goldberg 1966; van Noort 1969) and most CoIC axons belong to IC neurons. However, the CoIC contains some axons from other sources or destined for other targets. Tracer deposits in the MGB or the severed brachium of the IC (BIC) label projections to the contralateral MGB (Kudo and Niimi 1978; Aitkin and Phillips 1984; Oliver 1984; Hutson et al. 1991). Injection of anterograde tracers into the IC confirms crossed tectothalamic projections traveling in CoIC (Kudo and Niimi 1980; Andersen et al. 1980a; Hutson et al. 1991). It is unknown whether such axons also project in the contralateral IC before reaching the brachium.

The CoIC may contain also axons from subcollicular auditory centers (Ramón y Cajal 1904). Tracing studies in mice, rats, and cats suggest that the CoIC contains sagulum and/or lateral lemniscal fibers (González-Hernández et al. 1987; Henkel and Shneiderman 1988; Hutson et al. 1991; Bajo et al. 1993). Rat superior paraolivary nucleus cells project to the ipsilateral IC and proceed dorsomedially over the CoIC before ending in the contralateral DC (Fuentes et al. 1999). We are not aware of reports of axons in the CoIC from neural centers caudal to the superior olivary complex. The CoIC contains also auditory corticocollicular axons (Saldaña et al. 1996).




Figure 5.2. (A, B) Glutaraldehyde-fixed, 80- $\mu$ m-thick horizontal sections of the rat mid-brain tectum. Sections were postfixed with osmium tetroxide. Rostral is at the *top*. The *dotted white line* (C) indicates the border between the CoIC and the commissure of the superior colliculus. (D) Detail of the CoIC in the same section shown in (A).

Whether the projections from different sources are segregated within the CoIC is unknown. There is rostrocaudal or dorsoventral topography for commissural IC axons with more rostral IC neurons crossing the CoIC more rostrally (Saldaña and Merchán, unpublished observations). Likewise, CN axons of neurons with low characteristic frequency (dorsolateral neurons) course more dorsally than higher characteristic frequency neurons (Saldaña and Merchán 1992; Figs. 5.5, 5.6).

## 2. INTRACOLLICULAR PROJECTIONS OF THE CENTRAL NUCLEUS AND THE DORSAL CORTEX

### 2.1. CENTRAL NUCLEUS PROJECTIONS

A study of the organization of the intrinsic and commissural connections of the albino rat (Saldaña and Merchán 1992) injected the anterograde tracer *Phaseolus vulgaris*-leucoagglutinin (PHA-L) into the CN. Axons (Fig. 5.3) from a discrete CN deposit formed two ipsilateral, rostrocaudally oriented laminar plexuses of terminal fibers in IC, a main (medial) plexus (iMP), and an external (lateral) plexus (iLP), and two more plexuses in the contralateral IC, a main (medial) plexus (cMP), and an external (lateral) plexus (cLP). Few terminals were outside the plexuses. The iMP extended dorsomedially and ventrolaterally from the injection site, parallel to the CN fibrodendritic laminae of the CN, a well-known cytoarchitectonic arrangement in which flattened terminal fields of incoming fibers intermingle with similarly oriented, flattened dendritic trees of the resident cells (see Chapter 2). This iMP was continuous in CN and DC and extended rostrally into the RC. The iLP was narrower than iMP and in the EN, iLP was nearly parallel to the surface of the nucleus, with its ventral pole deeper than its dorsal end. Each contralateral plexus was symmetrical with the ipsilateral plexus. The main and the external plexuses met caudally at their ventral poles, forming a dorsally open three-dimensional structure. The fiber density was variable in the plexuses: iLPs were most prominent.

These features were common to all experiments (Saldaña and Merchán 1992), irrespective of the CN injection site. Individual differences in plexus thickness, extent, position, and orientation were related to the features of the injection site. First, plexus size and density was a function of injection site size. Second, plexus size and shape depended on the deposit's location in the CN tonotopic axis (Fig. 5.4). Injections in the ventromedial CN, whose neurons have a high characteristic frequency, labeled the iMP ventromedially, near the IC-periaqueductal gray matter border, while the iLP was ventrolateral in EN, near the pial surface. With more dorsolateral injection sites at lower characteristic frequency, the iMP moved dorsolateral, and the iLP was more dorsal and more medial.

Such intracollicular projections are not unique to the rat, and a similar organization is present in the IC of mouse (González-Hernández et al. 1986; Frisina et al. 1997, 1998), gerbil (Doroshenko and Cant 1994), chinchilla (Caspary and




Figure 5.3. Camera lucida drawings of coronal sections showing the terminal intracollicular labeling pattern from a small PHA-L injection in the right CN. The numbers (*right side*) are the distance from the caudal pole of the IC in microns. In the IC, *lines* represent fibers of passage, and *stippling* denotes terminal fields. The injection site is near C. Intracollicular terminal fields are symmetrical, and the commissural labeling is weaker. The four axonal plexuses of terminal fibers span most of the IC rostrocaudally, and enter the RC. The iMP and cMP are continuous across the CN and DC. (Adapted from Saldaña and Merchán 1992. Reprinted by permission of Wiley-Liss, Inc., a subsidiary of John Wiley & Sons, Inc.)




Figure 5.4. Schematic representation of the topographic distribution of intracollicular connections of the CN. The *black*, *gray*, and *white* areas represent territories targeted by discrete groups of CN neurons (*star* in **B**). *Black areas*: territories with input from ventromedial (high-frequency) CN neurons. *Gray areas*: targets of central (medium-frequency) CN neurons. *White areas*: targets of dorsolateral (low-frequency) CN neurons. The distribution of intracollicular fibers suggest that the laminae cross the divisions and involve most of the IC. These fibrocellular laminae curve around a point located dorsally. (Adapted from Saldaña and Merchán 1992. Reprinted by permission of Wiley-Liss, Inc., a subsidiary of John Wiley & Sons, Inc.)

Helfert 1993; Saint Marie 1996), guinea pig (Malmierca et. 1995), cat (Brunso-Bechtold et al. 1981; Malmierca et al. 1998), and bat (Vater and Feng 1990).

Intracollicular connections studied with biocytin deposits into physiologically identified CN loci showed a similar pattern of intracollicular CN projections in the guinea pig and the rat (Malmierca et al. 1995). Three-dimensional reconstructions of the plexuses provided a clearer picture of the near-concentric organization in IC (Fig. 5.5).

Further analysis of earlier results (Saldaña and Merchán 1992) and injections of PHA-L or biotinylated dextran (BDA) in rat CN show that plexuses labeled by injections in rostral isofrequency regions are denser rostrally than caudally, and the converse is true for the caudal deposits (Saldaña and Merchán, unpublished observations). Likewise, plexuses labeled by a ventral injection are denser ventrally. Thus, despite the widely divergent nature of the intracollicular projections of the CN, they involve preferentially neurons at rostrocaudal and dorsoventral levels near those of the parent cell bodies, a topographic arrangement whose functional significance will require further investigation.

## 2.2. DORSAL CORTEX PROJECTIONS

Because the ipsi- and contralateral targets of intracollicular fibers from the CN extend into the DC, it is of interest how DC intracollicular projections relate to those of CN. A PHA-L injection into rat central DC spared both the fiber-rich superficial layer of the DC and the CN (Fig. 5.6). It was centered at medium or medium-to-low characteristic frequency (CF). The labeled fibers created a medial and a lateral laminar plexus in each IC that spanned the IC rostrocaudally and merged caudally at their ventral ends, a pattern like that of the intracollicular CN projections. These plexuses were denser dorsally (compare Figs. 5.3 and 5.6). With more ventromedial (higher CF) or more dorsolateral (lower CF) DC injections, the plexuses resembled those labeled by injections in comparable CN tonotopic regions (Fig. 5.7A).

The organization of the DC intracollicular projections is remarkably similar to that of CN intracollicular projections, which accentuates the connectional parallels between the CN and the DC. Accordingly, the DC sends topographic (tonotopic) projections to all four ipsi- and contralateral IC subdivisions.

## 2.3. IMPLICATIONS FOR INFERIOR COLLICULUS ORGANIZATION

The pattern of intrinsic and commissural connections described in the preceding for CN and DC leads to several conclusions.

### 2.3.1. Strength of Intracollicular Inputs

Intracollicular projections are robust and may constitute one of the major inputs to IC neurons. They are an essential component of CN fibrodendritic laminae revealed by studies of the morphology of single intrinsic Golgi impregnated




Figure 5.5. Three-dimensional reconstructions of guinea pig fiber plexuses of intrinsic CN axons at CFs of 0.5 kHz (A), 6 kHz (B), and 21 kHz (C), respectively. The medial and lateral plexuses merge and form a complex, dorsally open structure. (Adapted from Malmierca et al. 1995. Reprinted by permission of Wiley-Liss, Inc., a subsidiary of John Wiley & Sons, Inc.)

axons (Herrera et al. 1988, 1989) or neurons and axons filled with horseradish peroxidase (HRP) injected intracellularly (Oliver et al. 1991).

### 2.3.2. Divergence of Intracollicular Inputs

Neurons in specific CN and DC regions project divergently to the four divisions of the ipsi- and contralateral IC (CN, DC, EN, and RC), confirming the idea from retrograde tracing experiments that the commissural projections chiefly connect homotopic IC regions (Adams 1980; Schweizer 1981; Zook and Caseday 1982; Druga and Syka 1984; Tokunaga et al. 1984; Coleman and Clerici 1987; Ross et al. 1988; Ross and Pollak, 1989; Frisina et al. 1989; González-Hernández et al. 1996; Zhang et al. 1998). The commissural projection is more than merely reciprocal or homotopic because the contralateral IC targets includes areas homotopic to the iLPs (González-Hernández et al. 1986; Malmierca et al. 1995; Frisina et al. 1998).

### 2.3.3. Topography of Intracollicular Inputs

The topographic (tonotopic) arrangement of the main plexuses show that CN and DC neurons project to CN and DC bilaterally at similar CF. We thus propose that the intracollicular projections enhance frequency specificity (see later), and that they may have other roles as well.

### 2.3.4. Possible Internal Topographies in Nuclei

The topography in the CN and DC projections to the EN raises the possibility that the EN possesses an unexpected tonotopic arrangement, whose isofrequency axes would parallel the IC surface, and whose main tonotopic axis is approximately mediolateral.

### 2.3.5. Contribution to Laminar Organization

The IC contains fibrocellular laminae. Each lamina is defined as the territory receiving input from CN and DC neurons with similar CF. Such laminae resemble the CN and DC plexuses or intracollicular input which curves around a point situated dorsally. Every fibrocellular lamina contains neurons and a precisely ordered axonal plexus. Fibrodendritic CN laminae may represent the central part of the fibrocellular laminae, but the latter are larger and extend into the other IC subdivisions. IC intrinsic and commissural projections are an essential component of the fibrocellular laminae, although other projection systems may also contribute. The distribution of the corticocollicular projections mimics that of the intracollicular connections (Fig. 5.7C), and this may apply to the cat (Andersen et al. 1980b).

Various lemniscal projections may follow a similar pattern, including those from the dorsal cochlear nucleus (Willard and Martin 1983; Malmierca et al. 2002) and the superior paraolivary nucleus (SPN) (Willard and Martin 1983; Fuentes et al. 1999). Input from the SPN forms two laminar plexuses in the






Figure 5.6. DC intracollicular projections after a small injection of PHA-L. (A-D) Micrographs of coronal sections. The numbers are the distance from interaural zero. The distribution of the terminal fibers plexus resembles that formed by CN axons (compare with Figs. 5.3 and 5.4).

ipsilateral IC, whose topography matches that of the iMP and iLP (Fig. 5.7B), and a weak CoIC projection. The SPN projection to the IC is topographic, and the size and shape of the terminal plexuses matches that of the intracollicular and corticocollicular projections. Thus, specific SPN cells project divergently to topographically matched neurons in the IC fibrocellular lamina.

### 3. INTRACOLLICULAR PROJECTIONS OF THE EXTERNAL NUCLEUS

Do EN neurons contribute to the same fibrocellular laminar distribution of intrinsic, commissural, ascending, and descending projections? The labeling in a representative experiment with a PHA-L deposit in EN (Fig. 5.8) showed terminal ipsilateral IC input to the EN extending rostrocaudally and dorsoventrally; in the DC and the dorsal part of the RC; and along the ventromedial IC border. Commissural terminal fields were sparser, and their distribution mirrored the ipsilateral projection. CN contained few fibers of passage bilaterally, and had few terminals. Thus, the EN projects to EN, DC, and RC bilaterally, and not to the CN. EN intracollicular projections do not contribute to the fibrocellular laminae of the IC, accentuating connective differences between the medial (CN and DC) and the lateral (EN) IC.

Comparing PHA-L injections of EN with DC or CN deposits suggests that the intracollicular projections of the EN are weaker. That the EN innervates preferentially peripheral regions of the IC is of interest because these regions are the main recipients of the corticocollicular projections (Saldaña et al. 1996) and the EN is an early link in the “belt” auditory pathway.

The absence of significant projections from the EN to the CN is in partial agreement with studies using [<sup>3</sup>H]leucine in the cat (Kudo and Niimi 1980) and with work in the rat (Chernock and Winer 2003) that show few cells retrogradely labeled in rat EN after CN injections of gold conjugated to cholera toxin  $\beta$  subunit. In contrast, several accounts find abundant retrogradely labeled cell bodies in the ipsi- and/or contralateral EN following HRP injections into the CN (Willard and Martin 1983; Willard and Ryugo 1983; Druga and Syka 1984; Tokunaga et al. 1984; Coleman and Clerici 1987; Frisina et al. 1998). A parsimonious explanation may be methodological: HRP injection sites tend to be large, so the possibility that the tracer spread or was incorporated by fibers of passage cannot be dismissed.

## 4. INTRACOLLICULAR CELLS OF ORIGIN

### 4.1. NEURONS WITH INTRINSIC CONNECTIONS

Golgi studies of the IC show that most neurons with well-impregnated axons have ipsilateral collaterals (cat: Rockel and Jones 1973a,b; rat: Herrera et al.




Figure 5.7. Schematic IC representation of the distribution of the terminal plexuses formed by axons from the right DC (A), the right superior paraolivary nucleus (B), and the right primary AC (C). *Black, gray, and white areas* represent the IC territories targeted by a discrete group of neurons (*star* in the DC in A, *shadowing* in insets B and C). *Black areas*: territories receiving high-frequency projections. *Gray areas*: zones with medium-frequency input. *White areas*: loci targeted by low-frequency neurons. The distribution of IC ascending and descending projections mimics the topography of the CN and DC intracollicular projections except in the contralateral EN, which does not receive cortical or superior paraolivary nucleus input. (Adapted from Saldaña et al. 1996. Reprinted by permission of Wiley-Liss, Inc., a subsidiary of John Wiley & Sons, Inc.)




Figure. 5.8. Camera lucida drawings of bilateral coronal sections showing intracollicular terminal labeling from an EN PHA-L injection. *Numbers* are the rostrocaudal locus. *Lines* represent fibers of passage, stippling the terminal fields. The injection site center is marked (*asterisk*). (**B**) The main features are input to the ipsilateral EN, DC, and RC; a weak, symmetrical commissural projection; and no ipsi- or contralateral CN projection.

1988, 1989; González-Hernández et al. 1989). Intracellular injection in cats and rodents demonstrates that most IC neurons with filling of distal axon segments, including those whose axons enter the BIC, the CoIC or the lateral lemniscus, possess intrinsic axonal collaterals (Oliver et al. 1991; Smith 1992; Wagner 1994, 1996; Reetz and Ehret 1999; Peruzzi et al. 2000). Therefore, with the reservations imposed by the limited sample inherent to the Golgi method and to intracellular filling in mind, many if not all IC neurons have intrinsic connections. This is supported by the IC injection of retrogradely transported tracers, which label IC neurons belonging to virtually all morphologic types in rodents, bats, cats, and other small mammals (Tokunaga et al. 1984; Coleman and Clerici 1987; Saint-Marie 1996; Frisina et al. 1998).

Two conclusions can be drawn. Numerically, intrinsic connections may constitute the largest source of input to IC neurons, as the number of IC-projecting neurons in other auditory centers falls short of the number of labeled intrinsic IC neurons; and IC neurons with intrinsic connections are diverse in morphology, connections, neurochemistry, and physiology (see Chapter 2). This diverse morphology is evident in two representative cat IC neurons with very different intrinsic axonal arborization (Figs. 5.9 and 5.10) labeled by intracellular HRP injection (Oliver et al. 1991).

#### 4.2. NEURONS WITH COMMISSURAL CONNECTIONS

Although there are no estimates of the total number of IC commissural neurons, nor has the number of CoIC axons been determined, the contralateral IC is surely among the main IC projection sources. For instance, the number of ferret IC neurons labeled after HRP injections in the contralateral IC was more than twice the number of neurons labeled in any other auditory center (Moore 1988).

Much like neurons with intrinsic connections, IC commissural neurons are heterogeneous in form and distribution. This is apparent in mammals as diverse as opossum (Willard and Martin 1983), mouse (González-Hernández et al. 1986; Frisina et al. 1998), rat (Druga and Syka 1984; Tokunaga et al. 1984; Coleman and Clerici 1987; González-Hernández et al. 1996), chinchilla (Saint Marie 1996), cat (Adams 1980), ferret (Moore 1988), and bat (Schweizer 1981). Although principal (disc-shaped) CN neurons have crossed projections, most commissural neurons are described as multipolar or stellate. While this trend may reflect differences in the relative proportion of different neuron types, multipolar neurons might also possess more extensive contralateral axonal arbors, or take up tracer more efficiently.

Four commissural neuron types were distinguished in the albino mouse on the basis of somatic size and their dendritic pattern: small-to-medium sized spiny cells, small-to-medium sized spineless cells, medium-sized sparsely spinous neurons, and large cells (González-Hernández et al. 1986). The profusion of intrinsic axon collaterals of the commissural neurons was also marked.

Commissural neuron projections may have various target nuclei. Rat IC contains some neurons that project to both the contralateral IC and the ipsilateral




Figure 5.9. A cat CN neuron labeled in vivo by intracellular HRP injection. The neuron is shown in the coronal plane and its IC position is indicated (*inset*). *Black arrow*, the orientation of CN fibrodendritic laminae. The highly oriented dendritic tree and the profusely ramified axonal arborization (*thinner lines*) both parallel the isofrequency laminae. The calibration bar represents 100  $\mu\text{m}$ . (From Oliver et al. 1991. Reprinted by permission of Wiley-Liss, Inc., a subsidiary of John Wiley & Sons, Inc.)

MGB (González-Hernández et al. 1991). The axonal branching of some mouse IC commissural neurons suggests that they project in the ipsilateral brachium and/or lateral lemniscus (Reetz and Ehret 1999). Some commissural axons send collaterals to CoIC to end in the tectal commissural column (Saldaña and Viñuela 2002).




Figure 5.10. The axonal arborization of a cat CN neuron labeled by an intracellular injection of HRP in vivo. The neuron is shown in the sagittal plane, the cell body is *cross-hatched*, and the dendrites are omitted to reveal the axonal arbor. The main axon (*arrowhead*) courses laterally toward the brachium of the IC. This neuron's intrinsic axonal branches bear more than 2000 presumed synaptic boutons. The calibration bar represents 500  $\mu\text{m}$ . (From Oliver et al. 1991. Reprinted by permission of Wiley-Liss, Inc., a subsidiary of John Wiley & Sons, Inc.)

### 4.3. DISTRIBUTION OF COMMISSURAL NEURONS

Early degeneration studies speculated that the commissural projections might originate throughout the IC (van Noort 1969), a possibility now confirmed. Large HRP injections that fill most of the IC-labeled neurons retrogradely in all contralateral IC subdivisions (Schweizer 1981; Ross et al. 1988; Fig. 5.11). Experiments with discrete injection of retrograde tracers invariably found retrogradely labeled neurons in contralateral IC areas that included, but were not necessarily restricted to, the area homotopic to the deposit. Moreover, all anterograde transport experiments label contralateral IC fibers, irrespective of the location of the ipsilateral IC injection (Kudo and Niimi 1980; Malmierca et al. 1995).

Although all IC regions are a source of commissural projections, commissural neurons are distributed nonuniformly. To label all cat CoIC neurons retrogradely, HRP was deposited into a cut made in the CoIC (Aitkin and Phillips 1984; Hutson et al. 1991) and commissural neurons were concentrated ventrolaterally at posterior levels, and more dorsally at anterior levels. A similar trend is found in the rat (Fig. 5.11).

## 5. THE FUNCTION OF INTRACOLLICULAR CONNECTIONS

The density of intracollicular connections could enable IC neurons to modulate the activity of other collicular neurons. Many IC neurons may receive lateral lemniscal information through local connections rather than through lemniscal synaptic input (Smith 1992; Wagner 1996; Moore et al. 1998; Reetz and Ehret 1999).

It seems unlikely that the CoIC is associated with some aspect of sound localization, as transecting the CoIC and/or the corpus callosum of trained cats had little or no effect on the animal's ability to localize a sound source (Moore et al. 1974). This result is consistent with results in the ferret after unilateral IC ablation (Kelly and Kavanagh 1994).

### 5.1. THE PHYSIOLOGIC ROLE OF COMMISSURAL CONNECTIONS

The morphologic diversity of IC neurons with commissural projections suggests that they might be equally varied neurochemically and electrophysiologically, and the intrinsic and commissural projections of the IC may contain excitatory and inhibitory fibers (Saldaña and Merchán 1992). Preliminary immunoelectron microscopic results in the chinchilla suggested that the commissural axons are immunoreactive for  $\gamma$ -aminobutyric acid (GABA) and glutamate (Caspary and Helfert 1993). Later morphologic data disagree as to the neurochemical status of these projections. A combined tract tracing and the immunocytochemical approach found many double-labeled cells in the contralateral IC (González-Hernández et al. 1996). This is consistent with the presence of fibers immuno-






Figure 5.11. Distribution of commissural neurons. (*Inset*) A BDA deposit (*black area*) spared only the lateral and dorsolateral CN and involved most of the DC. (**A**) A coronal section of the contralateral IC at the level of the injection site. Retrogradely labeled neurons are embedded in anterogradely labeled commissural fibers. (**B–E**) Commissural neurons in sections 400  $\mu$ m apart. A *dot* represents one neuron; the position of each labeled cell body was noted. Drawing made at  $\times 193$  with a plan  $\times 10$  objective (numerical aperture 0.30).

reactive for glutamic acid decarboxylase, the synthesizing enzyme for GABA (Vetter and Mugnaini 1985). However, similar experiments combining retrograde transport of a fluorescent tracer and immunocytochemistry for GABA revealed no commissural double-labeled neurons, despite their abundance in the lateral lemniscal nuclei (Zhang et al. 1998).

Using a different approach, many chinchilla neurons are retrogradely labeled following deposits of D- $^3\text{H}$ aspartate in the contralateral IC, which suggests that they are glutamatergic and that (at least some parts of) the commissural projections are excitatory (Saint Marie 1996). Some commissural neurons are immunopositive for enkephalin and neuropeptide Y (Nakagawa et al. 1995).

Support for a mixed excitatory–inhibitory commissural projection comes from electrophysiologic studies of living slices *in vitro*, which examined the effects on IC cells of electrical stimulation of the CoIC. DC neurons respond with an early inhibitory postsynaptic potential (IPSP) followed by a longer latency excitatory postsynaptic potential (EPSP); the pharmacologic evidence suggests that the IPSP reflects a direct, commissural GABAergic input, while the EPSP arises from direct, crossed input that activates non-*N*-methyl-D-aspartate-type (non-NMDA) glutamate receptors (Smith 1992). Similarly, most gerbil CN neurons receive both excitatory and inhibitory CoIC input, although GABAergic inhibition dominates (Moore et al. 1998). Furthermore, CoIC shock stimulation evokes short-latency PSPs in mouse IC, and these potentials may be excitatory or inhibitory (Reetz and Ehret 1999).

That the CoIC exerts both excitatory and inhibitory effects is corroborated by recent investigations *in vivo*. CN electrical stimulation and the injection of the glutamate-receptor agonist NMDA into CN both produced excitatory or inhibitory effects on the acoustically evoked activity of contralateral RC neurons, and this effect occurred in both normal and genetically epilepsy-prone rats (Chakravarty and Faingold 1997). These manipulations changed the neural discharge rate but did not affect sustained or onset discharge patterns. By the same token, silencing one IC reversibly by local microinjections of the glutamate receptor blocker kynurenic acid modified the spectral properties of contralateral IC neurons (Malmierca et al. 2003). In most neurons, blocking the contralateral IC increased the frequency response area to monaural (contralateral) or binaural stimulation, although decreased frequency response areas were also recorded (Fig. 5.12). One limitation of these *in vivo* studies is that they do not reveal if the neurons whose activity is modified by the contralateral IC manipulation receive direct commissural input or if other, interposed neurons are required. Moreover, IC stimulation or blockade alters the stream of information ascending to the cerebral cortex, and so may interfere with descending control of collicular activity.

## 5.2. NEOCORTICAL CONTROL OF INTRACOLLICULAR CIRCUITS

The corticofugal system modulates the sensitivity of IC neurons to sound frequency, amplitude, and duration (Suga and Ma 2003). For frequency selectivity,




Figure 5.12. Blocking reversibly the activity of one IC with kynurenic acid changes the frequency response area of commissural neurons. In unit 115-2 (*upper half*), the contralateral blockade reduced the frequency response areas elicited by monaural (A1–A3) and binaural stimulation (B1–B3). In unit 118-2 (*lower half*), blockade increased the frequency response areas for monaural (C1–C3) and binaural stimulation (D1–D3). (Reproduced from Malmierca et al. 2003.)

the auditory neocortical activity evokes both facilitation and inhibition of auditory responses. The amount of facilitation and inhibition varies as a function of the relationship in frequency tuning between neocortical and IC neurons. A neuron's response is augmented at its CF and inhibited at other frequencies (see Chapter 8). Corticocollicular and intracollicular projections are topographically (tonotopically) matched (Andersen et al. 1980b; Saldaña et al. 1996). Corticocollicular projections are excitatory (Feliciano and Potashner 1995), and intracollicular projections have both excitatory and inhibitory effects (see Section 5.1), so that the facilitation and inhibition of IC neurons produced by auditory cortex activation can be best explained in terms of interactions mediated by intracollicular connections governed by corticocollicular modulation.

## 6. THEMES FOR DISCUSSION

1. Different IC neurons receive their main inputs from different sources. How do the IC intrinsic and commissural connections relate to the origin of their inputs?
2. Available data on the commissural connections come from bulk transport experiments. What is the morphology and distribution of individual commissural axons?
3. Within the IC, different afferent projections terminate preferentially on different parts of the target neurons (cell body, proximal or distal dendrites). For example, some lateral lemniscal inputs provide a significant input to more proximal portions of IC neurons, including cell bodies, than do corticocollicular fibers. What part(s) of the neuron is/are targeted preferentially by intracollicular connections?
4. The EN receives topographic ascending as well as intracollicular and descending projections. Are the efferent projections of the EN topographically organized?
5. The EN is usually regarded as a center for multisensory integration. How do the nonauditory projections intersperse within the EN with the topographic projection from the CN and the DC?
6. The CN has been implicated in the genesis of audiogenic seizures and the EN/RC in the propagation of the epileptic crises. Does the CN-to-EN projection play a role in the physiopathology of these seizures?
7. Neurons within an IC isofrequency region differ in other physiologic parameters. The distribution of some of these parameters within the IC is orthogonal to the isofrequency planes of the nucleus (Schreiner and Langner 1988). Is there a relationship between the orderly pattern of intracollicular connections and the representation of parameters other than frequency?
8. What information is conveyed by commissural fibers? What is the biologic significance of the CoIC?

9. The results reviewed here suggest that the IC consists of concentric fibrocellular laminae that involve all collicular subdivisions. Principal (disc-shaped) neurons are likely integral elements of these CN laminae. What neurons form the cellular substrate of the fibrocellular laminae in the cortical subdivisions of the IC?
10. Are there comparative variations in the organization of intracollicular projections? Some big brown bat EN neurons are excited by corticocollicular fibers and inhibit CN neurons, which are facilitated by corticofugal input (Jen et al. 1998). However, the morphologic data suggest that, in the rat, the EN does not project significantly to the CN.

## Acknowledgments

This research was supported by Spanish Ministry of Science and Technology Grants BFI2000-1358 (E.S.) and BFI2000-1369 (M.A.M.), and by Junta de Castilla y León Grants SA079/01 (E.S.) and SA084/01 (M.A.M.). We thank Ms. M.-Auxiliadora Aparicio for skillful assistance with processing the osmicated sections.

## Abbreviations

BDA	biotinylated dextran amine
BIC	brachium of the inferior colliculus
CN	central nucleus of the inferior colliculus
CoIC	commissure of the inferior colliculus
DC	dorsal cortex of the inferior colliculus
EN	external nucleus of the inferior colliculus
EPSP	excitatory postsynaptic potential
GABA	$\gamma$ -aminobutyric acid
GAD	glutamic acid decarboxylase
HRP	horseradish peroxidase
IC	inferior colliculus
IPSP	inhibitory postsynaptic potential
LSO	lateral superior olive
MNTB	medial nucleus of the trapezoid body
NMDA	<i>N</i> -methyl-D-aspartate
PHA-L	<i>Phaseolus vulgaris</i> -leucoagglutinin
RC	rostral cortex of the inferior colliculus
SC	superior colliculus
SPON	superior paraolivary nucleus
Te1	primary auditory area of rat cerebral cortex

## References

- Adams JC (1980) Crossed and descending projections to the inferior colliculus. *Neuroscience Letters* **19**:1–5.
- Aitkin LM and Phillips SC (1984) The interconnections of the inferior colliculi through their commissure. *Journal of Comparative Neurology* **228**:210–216.
- Andersen RA, Roth GL, Aitkin LM, and Merzenich MM (1980a) The efferent projections of the central nucleus and the pericentral nucleus of the inferior colliculus in the cat. *Journal of Comparative Neurology* **194**:649–662.
- Andersen RA, Snyder RL, and Merzenich MM (1980b) The topographic organization of corticocollicular projections from physiologically identified loci in the AI, AII, and anterior auditory cortical fields of the cat. *Journal of Comparative Neurology* **191**:479–494.
- Bajo VM, Merchán MA, López DE, and Rouiller EM (1993) Neuronal morphology and efferent projections of the dorsal nucleus of the lateral lemniscus in the rat. *Journal of Comparative Neurology* **334**:241–262.
- Brunso-Bechtold JK, Thompson GC, and Masterton RB (1981) HRP study of the organization of auditory afferents ascending to central nucleus of inferior colliculus in cat. *Journal of Comparative Neurology* **197**:705–722.
- Caspary DM and Helfert RH (1993) Two populations of commissural projections connecting like regions of the inferior colliculus immunostain for different amino acid transmitters. *Society for Neuroscience Abstracts* **19**:1425.
- Chakravarty DN and Faingold CL (1997) Aberrant neuronal responsiveness in the genetically epilepsy-prone rat: acoustic responses and influences of the central nucleus upon the external nucleus of inferior colliculus. *Brain Research* **761**:263–270.
- Chernock ML and Winer JA (2003) Intrinsic and commissural connections of the rat inferior colliculus. *Association for Research in Otolaryngology Abstracts* **26**:787.
- Coleman JR and Clerici WJ (1987) Sources of projections to subdivisions of the inferior colliculus in the rat. *Journal of Comparative Neurology* **262**:215–226.
- Doroshenko NZ and Cant NB (1994) Morphology of commissural axons in the inferior colliculus of the gerbil. *Society for Neuroscience Abstracts* **20**:926.
- Druga R and Syka J (1984) Ascending and descending projections to the inferior colliculus in the rat. *Physiologica Bohemoslovaca* **33**:31–42.
- Faye-Lund H and Osen KK (1985) Anatomy of the inferior colliculus in rat. *Anatomy and Embryology* **171**:1–20.
- Feliciano M and Potashner SJ (1995) Evidence for a glutamatergic pathway from the guinea pig auditory cortex to the inferior colliculus. *Journal of Neurochemistry* **65**:1348–1357.
- Frisina RD, O'Neill WE, and Zettel ML (1989) Functional organization of mustached bat inferior colliculus: II. Connections of the FM<sub>2</sub> region. *Journal of Comparative Neurology* **284**:85–107.
- Frisina RD, Walton JP, Lynch-Armour MA, and Klotz DA (1997) Efferent projections of a physiologically characterized region of the inferior colliculus of the young adult CBA mouse. *Journal of the Acoustical Society of America* **101**:2741–2753.
- Frisina RD, Walton JP, Lynch-Armour MA, and Byrd JD (1998) Inputs to a physiologically characterized region of the inferior colliculus of the young adult CBA mouse. *Hearing Research* **115**:61–81.

- Fuentes V, Berrebi AS, and Saldaña E (1999) Trajectory, morphology and distribution of axons of the superior paraolivary nucleus (SPON) that innervate the inferior colliculus in the rat. *Association for Research in Otolaryngology Abstracts* **22**:222.
- González-Hernández TH, Meyer G, and Ferres-Torres R (1986) The commissural interconnections of the inferior colliculus in the albino mouse. *Brain Research* **368**:268–276.
- González-Hernández TH, Meyer G, Ferres-Torres R, Castañeyra-Perdomo A, and del Mar Perez Delgado MM (1987) Afferent connections of the inferior colliculus in the albino mouse. *Journal für Hirnforschung* **28**:315–323.
- González-Hernández TH, Meyer G, and Ferres-Torres R (1989) Development of neuronal types and laminar organization in the central nucleus of the inferior colliculus in the cat. *Neuroscience* **30**:127–141.
- González-Hernández TH, Galindo-Mireles D, Castañeyra-Perdomo A, and Ferres-Torres R (1991) Divergent projections of projecting neurons of the inferior colliculus to the medial geniculate body and the contralateral inferior colliculus in the rat. *Hearing Research* **52**:17–21.
- González-Hernández T, Mantolán-Sarmiento B, González-González B, and Pérez-González H (1996) Sources of GABAergic input to the inferior colliculus of the rat. *Journal of Comparative Neurology* **372**:309–326.
- Held H (1891) Die centralen Bahnen des Nervus acusticum bei der Katze. *Archiv für Anatomie und Physiologie. Anatomische Abteilung* **15**:271–290.
- Held H (1893) Die centrale Gehörleitung. *Archiv für Anatomie und Physiologie. Anatomische Abteilung* **17**:201–248.
- Henkel CK (1983) Evidence of sub-collicular auditory projections to the medial geniculate nucleus in the cat: an autoradiographic and horseradish peroxidase study. *Brain Research* **259**:21–30.
- Henkel CK and Shneiderman A (1988) Nucleus sagulum: projections of a lateral tegmental area to the inferior colliculus in the cat. *Journal of Comparative Neurology* **271**:577–588.
- Herrera M, Correa J, Sánchez del Campo F, and Ruiz A (1988) Stellate cells and their axonal patterns in the central nucleus of the inferior colliculus of the cat (*Felis domesticus*). *Journal für Hirnforschung* **29**:393–402.
- Herrera M, Sánchez del Campo F, Puchades A, and Correa J (1989) Axonal patterns of disc-shaped cells in the central nucleus of the cat inferior colliculus. *Zeitschrift für mikroskopisch-anatomische Forschung* **103**:515–525.
- Hutson KA, Glendenning KK, and Masterton RB (1991) Acoustic chiasm IV: eight mid-brain decussations of the auditory system in the cat. *Journal of Comparative Neurology* **312**:105–131.
- Jen PH-S, Chen QC, and Sun XD (1998) Corticofugal regulation of auditory sensitivity in the bat inferior colliculus. *Journal of Comparative Physiology A* **183**:683–697.
- Kelly JB and Kavanagh GL (1994) Sound localization after unilateral lesions of inferior colliculus in the ferret (*Mustela putorius*). *Journal Neurophysiology* **71**:1078–1087.
- Kudo M and Niimi K (1978) Ascending projections of the inferior colliculus onto the medial geniculate body in the cat studied by anterograde and retrograde tracing techniques. *Brain Research* **155**:113–117.
- Kudo M and Niimi K (1980) Ascending projections of the inferior colliculus in the cat: an autoradiographic study. *Journal of Comparative Neurology* **191**:545–556.

- Kulesza RJ Jr, Viñuela A, Saldaña E, and Berrebi AS (2002) Unbiased stereological estimates of neuron number in subcortical auditory nuclei of the rat. *Hearing Research* **168**:12–24.
- Malmierca MS, Rees A, Le Beau FEN, and Bjaalie JG (1995) Laminar organization of frequency-defined local axons within and between the inferior colliculi of the guinea pig. *Journal of Comparative Neurology* **357**:124–144.
- Malmierca MS, Leergaard TB, Bajo VM, Bjaalie JG, and Merchán MA (1998) Anatomic evidence of a three-dimensional mosaic pattern of tonotopic organization in the ventral complex of the lateral lemniscus in cat. *Journal of Neuroscience* **18**:10603–10618.
- Malmierca MS, Merchán MA, Henkel CK, and Oliver DL (2002) Direct projections from cochlear nuclear complex to auditory thalamus in the rat. *Journal of Neuroscience* **22**:10891–10897.
- Malmierca MS, Hernández O, Falconi A, López-Poveda EA, Merchán M, and Rees A (2003) The commissure of the inferior colliculus shapes frequency response areas in rat: an *in vivo* study using reversible blockade with microinjection of kynurenic acid. *Experimental Brain Research* **153**:522–529.
- Moore CN, Casseday JH, and Neff WD (1974) Sound localization: the role of the commissural pathways of the auditory system of the cat. *Brain Research* **82**:13–26.
- Moore DR (1988) Auditory brainstem of the ferret: sources of projections to the inferior colliculus. *Journal of Comparative Neurology* **269**:342–354.
- Moore DR, Kotak VC, and Sanes DH (1998) Commissural and lemniscal synaptic input to the gerbil inferior colliculus. *Journal of Neurophysiology* **80**:2229–2236.
- Moore RY and Goldberg JM (1966) Projections of the inferior colliculus in the monkey. *Experimental Neurology* **14**:429–438.
- Nakagawa H, Ikeda M, Houtani T, Ueyama T, Baba K, Kondoh A, Yamamoto T, Yamashita T, and Sugimoto T (1995) Immunohistochemical evidence for enkephalin and neuropeptide Y in rat inferior colliculus neurons that provide ascending or commissural fibers. *Brain Research* **690**:236–240.
- Oliver DL (1984) Neuron types in the central nucleus of the inferior colliculus that project to the medial geniculate body. *Neuroscience* **11**:409–424.
- Oliver DL, Kuwada S, Yin TCT, Haberly LB, and Henkel CK (1991) Dendritic and axonal morphology of HRP-injected neurons in the inferior colliculus of the cat. *Journal of Comparative Neurology* **303**:75–100.
- Peruzzi D, Sivaramakrishnan S, and Oliver DL (2000) Identification of cell types in brain slices of the inferior colliculus. *Neuroscience* **101**:403–416.
- Ramón y Cajal S (1904) *Textura del Sistema Nervioso del Hombre y de los Vertebrados*. Nicolás Moya, Madrid.
- Reetz G and Ehret G (1999) Inputs from three brainstem sources to identified neurons of the mouse inferior colliculus slice. *Brain Research* **816**:527–543.
- Rockel AJ and Jones EG (1973a) The neuronal organization of the inferior colliculus of the adult cat. I. The central nucleus. *Journal of Comparative Neurology* **147**:11–60.
- Rockel AJ and Jones EG (1973b) The neuronal organization of the inferior colliculus of the adult cat. II. The pericentral nucleus. *Journal of Comparative Neurology* **149**:301–334.
- Ross LS and Pollak GD (1989) Differential ascending projections to aural regions in the 60 kHz contour of the mustache bat's inferior colliculus. *Journal of Neuroscience* **9**:2819–2834.
- Ross LS, Pollak GD, and Zook JM (1988) Origin of ascending projections to an iso-frequency region of the mustache bat's inferior colliculus. *Journal of Comparative Neurology* **270**:488–505.



- Saint Marie RL (1996) Glutamatergic connections of the auditory midbrain: selective uptake and axonal transport of D-[3H]aspartate. *Journal of Comparative Neurology* **373**:255–270.
- Saldaña E and Merchán MA (1992) Intrinsic and commissural connections of the rat inferior colliculus. *Journal of Comparative Neurology* **319**:417–437.
- Saldaña E and Viñuela A (2002) Projections from the inferior colliculus to the tectal commissural column. *Association for Research in Otolaryngology Abstracts* **25**:176.
- Saldaña E, Feliciano M, and Mugnaini E (1996) Distribution of descending projections from primary auditory neocortex to inferior colliculus mimics the topography of intracollicular projections. *Journal of Comparative Neurology* **371**:15–40.
- Schreiner CE and Langner G (1988) Periodicity coding in the inferior colliculus of the cat. II. Topographical organization. *Journal of Neurophysiology* **60**:1823–1840.
- Schweizer H (1981) The connections of the inferior colliculus and the organization of the brainstem auditory system in the greater horseshoe bat (*Rhinolophus ferrum-equinum*). *Journal of Comparative Neurology* **201**:25–49.
- Smith PH (1992) Anatomy and physiology of multipolar cells in the rat inferior collicular cortex using the in vitro brain slice technique. *Journal of Neuroscience* **12**:3700–3715.
- Suga N and Ma X (2003) Multiparametric corticofugal modulation and plasticity in the auditory system. *Nature Reviews Neuroscience* **4**:783–794.
- Tokunaga A, Sugita S, and Otani K (1984) Auditory and non-auditory subcortical afferents to the inferior colliculus in the rat. *Journal für Hirnforschung* **25**:461–472.
- van Noort J (1969) *The Structure and Connections of the Inferior Colliculus. An Investigation of the Lower Auditory System*. Van Gorcum, Assen.
- Vater M and Feng AS (1990) Functional organization of ascending and descending connections of the cochlear nucleus of horseshoe bats. *Journal of Comparative Neurology* **292**:373–395.
- Vetter DE and Mugnaini E (1985) Discrete bilateral GABAergic neuron pools at the commissures of superior and inferior colliculi in the rat. *Society for Neuroscience Abstracts* **11**:246.
- Wagner T (1994) Intrinsic properties of identified neurones in the central nucleus of mouse inferior colliculus. *NeuroReport* **6**:89–93.
- Wagner T (1996) Lemniscal input to identified neurons of the central nucleus of mouse inferior colliculus: an intracellular brain slice study. *European Journal of Neuroscience* **8**:1231–1239.
- Willard FH and Martin GF (1983) The auditory brainstem nuclei and some of their projections to the inferior colliculus in the North American opossum. *Neuroscience* **10**:1203–1232.
- Willard FH and Ryugo DK (1983) Anatomy of the central auditory system. In: Willott JF (ed). *The Auditory Psychobiology of the Mouse*. Charles C Thomas, Springfield, pp. 201–304.
- Zhang DX, Li L, Kelly JB, and Wu SH (1998) GABAergic projections from the lateral lemniscus to the inferior colliculus of the rat. *Hearing Research* **117**:1–12.
- Zook JM and Casseday JH (1982) Origin of ascending projections to inferior colliculus in the mustache bat, *Pteronotus parnellii*. *Journal of Comparative Neurology* **207**:14–28.

# Chapter 6

## Descending Connections of the Auditory Midbrain

ANN M. THOMPSON

### 1. INTRODUCTION

The inferior colliculus (IC) integrates acoustic information from many auditory nuclei and relays the processed information to higher centers. In addition, the IC is the source of robust projections to lower auditory nuclei, as well as to a number of nonauditory brain regions. Identification of these subcollicular targets and the behavioral effects of manipulating the IC have implicated the IC in modulating certain ascending and descending auditory pathways as well as premotor centers. However, the functional impact of the descending IC pathways remain poorly understood. The following overview presents some of the descending IC projections within a functional context.

This chapter begins with descriptions of the IC projections to lower auditory nuclei and the functional auditory circuits that are likely recipients of these projections. Then, the nonauditory brain stem centers that receive direct and indirect projections of the IC are described. The potential IC inputs to brain circuits mediating various behaviors are also discussed. It should be noted that the pathways presented have been described mostly in the rat, but also in a few other mammalian species. For clarity, a generalized mammalian scheme is presented but species differences likely exist. Further, some projections have been found only in species with specialized hearing, for example, bats and owls (see Chapter 17), and these potentially unique projections are noted where appropriate.

### 2. PRIMARY AUDITORY TARGETS

#### *2.1. NUCLEI OF THE LATERAL LEMNISCUS*

Axons of IC neurons exit the midbrain ventrolaterally and enter the ipsilateral lateral lemniscus (LL). As the fibers descend in the LL, many terminate in the dorsal nucleus (DNLL) (Caicedo and Herbert 1993; Thompson and Thompson 1993; Wenstrup et al. 1994; Malmierca et al. 1996). As the fibers continue to

descend, minor terminations are made in the intermediate (INLL) and then in the ventral nucleus of the lateral lemniscus (VNLL) (Thompson and Thompson 1993; Malmierca et al. 1996). The part of the IC giving rise to the projection to the ipsilateral nuclei of the lateral lemniscus is the central nucleus (ICc; Fig. 6.1B; Caicedo and Herbert 1993; Malmierca et al. 1996). On the contralateral side, the only IC projection is a minor one to the VNLL arising from the ICc or the dorsal cortex (ICd) (Thompson and Thompson 1993).

Among the projections to LL nuclei, those to the DNLL predominate. The DNLL projects to the central nucleus of the IC bilaterally (see Chapter 2); many




Figure 6.1. Descending IC auditory projections. The predominant targets are indicated by *thicker arrows* and *darker shading*; see the text for details. (A) The dorsal cortex of the IC (*gray*) projects to the DCN bilaterally. (B) The main target of the central nucleus is the ipsilateral DNLL and secondarily the GCD of the DCN. (C) The central and lateral IC nuclei project to the SOC and the CN. The major SOC target is the ipsilateral VNTB and the RPO. Minor periolivary targets are the ipsilateral CPO, the LNTB, the DMPO, and the contralateral rostral periolivary region (RPO) and the ventral nucleus of the trapezoid body (VNTB). Within the CN, the DCN is the main target of the ICc and ICi. The projection is bilateral.

of its cells use  $\gamma$ -aminobutyric acid (GABA) as a transmitter and help to shape the responses of IC neurons (Ehret 1997). It is also notable that the projection to the DNLL is the only projection to LL nuclei that is tonotopic (Caicedo and Herbert 1993; Malmierca et al. 1996).

Descending IC fibers also target neuronal cell groups situated near the LL nuclei. One such cell group, the sagulum, lies lateral to the DNLL. IC axons enter the sagulum near the beginning of their descent in the LL. The IC areas projecting to the sagulum include the lateral nucleus (ICl), ICd, and ICc (Caicedo and Herbert 1993; Malmierca et al. 1996; Beneyto et al. 1998). Although the ipsilateral projection predominates, the ICd and/or the ICc also project to the contralateral sagulum. Descending IC axons also terminate in the horizontal cell group located ventral to the dorsal nucleus of the lateral lemniscus as described in the rat (Caicedo and Herbert 1993). Other IC axon collaterals terminate in ipsilateral perilemniscal regions, for example, within the LL but outside its primary nuclei. The source of the perilemniscal projections is the ICc and/or the ICd (Caicedo and Herbert 1993; Thompson and Thompson 1993).

## 2.2. SUPERIOR OLIVARY COMPLEX INPUTS AND CONNECTIONS WITH THE OLIVOCOCHLEAR SYSTEM

Some IC fibers descending in the LL enter the trapezoid body and terminate in the superior olivary complex (SOC). Not all of the SOC nuclei receive IC input; the IC projects predominantly to the periolivary regions with little or no input to the principal nuclei (Fig. 6.1C). On entering the rostral part of the SOC, IC fibers terminate in the rostral periolivary region (RPO) (Caicedo and Herbert 1993; Thompson and Thompson 1993; Malmierca et al. 1996; Schofield and Cant 1999). Many other fibers travel ventrally to enter and terminate in the ventral nucleus of the trapezoid body (VNTB) (Thompson and Thompson 1993; Vetter et al. 1993; Malmierca et al. 1996; Schofield and Cant 1999). Some fibers end in the lateral nucleus of the trapezoid body and the caudal periolivary region (Caicedo and Herbert 1993; Thompson and Thompson 1993). Other fibers maintain their dorsal position in the trapezoid body and terminate in the dorsomedial periolivary region (superior paraolivary nucleus of rodent; Thompson and Thompson 1993; Schofield and Cant 1999). Although not often observed and comparatively sparse, IC projections to the margins of the lateral superior olive (LSO), the LSO itself (Thompson and Thompson 1993), and the medial nucleus of the trapezoid body (Carey and Webster 1971) have been noted. The several origins of the descending IC pathway to the SOC include the ICl (especially layer 3) and ICc nuclei (Caicedo and Herbert 1993; Malmierca et al. 1996). These descending IC cells are glutamatergic and likely have an excitatory influence on SOC target neurons (Saint Marie 1996).

Two periolivary regions, the VNTB and the RPO, also receive input from the contralateral IC (Malmierca et al. 1996; Schofield and Cant 1999). The axons cross the midline in the ventral part of the trapezoid body. The contralateral projections are sparser than the ipsilateral projections.

The periolivary regions, where most IC axons terminate, contain the efferent (centrifugal) neurons of the medial olivocochlear system (MOC) (Warr 1992). MOC neurons project to the cochlea where they innervate outer hair cells (OHCs). The terminal endings of descending IC neurons make synaptic contacts with MOC neurons (Vetter et al. 1992; Thompson and Thompson 1993). This indicates that, by its descending projections to the SOC, the IC may influence cochlear function mediated by MOC neurons and OHCs. This is further supported by neurophysiologic studies indicating that the IC, by exciting MOC neurons, influences OHC responses (Scates et al. 1999; Braun 2000; Mulders and Robertson 2000, 2002). Because the IC projection to the SOC is mainly ipsilateral, and because most MOC neurons project to the contralateral cochlea, the IC likely influences the cochlea of the opposite ear (Fig. 6.2). The IC receives most of its direct ascending auditory input from the contralateral side and, therefore, this can be considered a feedback pathway. In addition, MOC neurons send collaterals to the cochlear nucleus (CN), allowing for IC modulation of these neurons.

Periolivary regions also contain olivocochlear nucleus neurons (OCNs) that project to the CN complex. Olivocochlear nucleus neurons are a specific target of the descending IC system (Schofield and Cant 1999). Olivocochlear nucleus neurons that receive IC inputs project bilaterally and hence form a route by which the IC may modulate the ascending auditory pathway simultaneously and bilaterally at the level of the cochlear nuclear complex (Fig. 6.2). The effect may be excitatory if glutamate is the neurotransmitter of this pathway and the synaptic targets of these cells are themselves excitatory.




Figure 6.2. Descending IC projections to the centrifugal auditory pathways innervating the CN and cochlea. The MOCs near the SOC are the main target. MOCs project to the cochlea bilaterally, mainly contralaterally, and terminate on outer hair cells (OHCs). Via its projections to MOCs, the IC may modulate cochlear processing. The IC also projects to olivocochlear nucleus (OCN) neurons in the SOC that, in turn, project to the CN. It is unknown whether IC projections to the CN reach cells that themselves project to the IC.

### 2.3. *INPUTS TO EXTRINSIC AND INTRINSIC CIRCUITS OF THE COCHLEAR NUCLEUS COMPLEX*

Of the descending IC projections, the one to the CN complex has been the most studied. Descending IC axons enter the CN complex via the LL and then the trapezoid body. As the fibers reach the ventral CN within the trapezoid body, they turn dorsally, between the descending trigeminal nerve root and the anteroventral cochlear nucleus (AVCN), to enter the dorsal cochlear nucleus (DCN). Among the main cochlear nuclear complex subdivisions, the DCN is the major target of descending IC fibers (Fig. 6.1A, C; Vater and Feng 1990; Saint Marie 1996; Schofield 2001). The projection to the DCN is bilateral, with a slight ipsilateral preponderance (Kane and Conlee 1979; Caicedo and Herbert 1993; Schofield 2001). At least one component of the IC pathway to the DCN is GABAergic (Alibardi 2002).

Within the ipsilateral DCN, most IC axons terminate in the fusiform cell layer and deeper layers (Kane and Conlee 1979; Conlee and Kane 1982; Shore et al. 1991; Malmierca et al. 1996). In the ipsilateral fusiform cell layer, IC terminal endings are associated with granule cell dendrites and these terminals have small, round vesicles and short, asymmetric densities (type 1a glomerular endings; Kane 1977). Besides those scattered throughout the DCN, granule cells within the granule cell domain (GCD) may also receive IC input (Malmierca et al. 1996; Shore and Moore 1998). The IC projection to the GCD is bilateral. The small cell cap, adjacent to the GCD, may also receive IC input (Malmierca et al. 1996).

Through its terminations onto the dendrites of granule cells in the fusiform cell layer and input to the granule cell domain, the IC influences the intrinsic circuitry of the CN complex (Fig. 6.3). The cochlear nucleus intrinsic circuits are comprised of a number of cell types, including granule cells (see Chapter 1). Granule cells are remarkable as they receive a wide variety of inputs from within the auditory system (including auditory cortex, type II auditory nerve fibers, medial olivocochlear efferents), and also from the vestibular and somatosensory systems (Wright and Ryugo 1996), the pontine gray (Ohlrogge et al. 2001), and the brain stem serotonergic system (Hurley and Thompson 2001). Therefore, the descending IC pathway to granule cells is but one part of an intricate, multimodal circuit that likely has a role in modulating the responses of CN neurons.

The IC also provides inputs to giant cells and the dendrites of fusiform cells situated in the deeper layers of the DCN (Kane 1977). The ipsilateral IC projects to the innermost part of the molecular layer where the axons terminate onto the fusiform cell apical dendrites (Kane and Conlee 1979; Shore et al. 1991; Malmierca et al. 1996). Both fusiform and giant cells are principal neurons of the DCN and project to the contralateral IC (see Chapter 3). Therefore, by its descending inputs to fusiform and giant cells, the IC can influence the ascending auditory pathway that originates in the DCN. Moreover, because the granule cells project directly and indirectly (via cartwheel cells) to fusiform cells (Young




Figure 6.3. Interaction of descending connections with ascending and intrinsic CN circuits. The DCN is a major auditory target of the IC. Both the ipsilateral and contralateral IC provide input to giant and fusiform cells (*dashed arrows*), both of which receive direct input from the cochlear type I auditory afferents and send monosynaptic input to the contralateral IC (*solid arrows*). The IC also projects to an intrinsic cochlear nucleus circuit through the ipsilateral GCD, and to the fusiform cell layer where other granule cells are found. Granule cells receive many acoustic and nonacoustic inputs and project directly and indirectly (via cartwheel cells) to fusiform cells, possibly propagating IC influences to the ascending CN output via the fusiform cell. Although the function of fusiform cells is not fully understood, they may be involved in vertical sound localization.

and Davis 2002), granule cells may be an indirect route for the IC to influence the ascending output of the DCN.

ICc and ICd neurons are the primary source of the descending projections to the CN complex (Caicedo and Herbert 1993; Schofield 2001). Some cell-specific pathways have been identified. Giant neurons in the deep DCN receive input from ipsilateral ICc neurons (Conlee and Kane 1982; Shore et al. 1991). The ICc is also the source of input to the granule cell domain (predominantly ipsilaterally; Shore and Moore 1998). The ICd projects to the CN complex bilaterally (Ostapoff et al. 1990; Schofield 2001). Some of the projections arise solely from the ICd whereas others terminate solely in the DCN (Fig. 6.1A). The CN granule cell domain receives only ICc input (Fig. 6.1B).

The types of IC neurons that are the source of descending projections to the CN complex have also been identified in the guinea pig. The projections originate from multipolar (stellate) cells in the ICc and ICI, from disc-shaped ICc neurons (Schofield 2001), and multipolar neurons residing mostly in ICd layers 2 and 3.

The projections from the ICc to the DCN are topographic and/or tonotopic in rodents and carnivores (Kane and Conlee 1979; Saldaña et al. 1989; Caicedo and Herbert 1993; Malmierca et al. 1996). Unlike those arising from the ICc, the ICI (external cortex) projections terminating in the CN complex are diffuse, and probably not tonotopic (Caicedo and Herbert 1993). These descending inputs to the DCN likely modulate fusiform cells to enhance vertical sound localization (Fujino and Oertel 2003).

The projections to the contralateral DCN are independent of those to the ipsilateral side (Schofield 2001). Electron microscopic evidence supports the independence of the contralateral projection because, unlike those on the ipsilateral side, the majority of IC synapses in the contralateral fusiform cell layer appear to be axodendritic (type 5a) and to terminate on fusiform cells (Kane 1977).

There is little evidence that the IC projects to the ventral cochlear nucleus (VCN). Interpretation of some anatomical findings is problematic because collicular axons traverse the VCN en route to the DCN. The IC has a sparse projection bilaterally to the most anterior part of the octopus cell area in the posteroventral cochlear nucleus (PVCN) (Kane and Conlee 1979). Other results (Caicedo and Herbert 1993; Saint Marie 1996) indicate a sparse projection (at least from the ICc and/or ICI) to the AVCN and PVCN.

### 3. PROJECTIONS TO MOTOR PATHWAYS AND STRUCTURES INVOLVED IN ORIENTATION BEHAVIORS

#### 3.1. *THE SUPERIOR COLLICULUS AND PINNA MOVEMENTS*

Various lines of evidence indicate that the IC controls pinna movements. The IC does not project directly to the facial (VII) motor nucleus (FMN), which innervates the pinna muscles, but there are several indirect routes whereby the IC could influence these muscles. The predominant route for such influence is through the ipsilateral deep superior colliculus (SC) (Fig. 6.4).

The deep SC receives convergent visual, auditory, and somatosensory input and it is involved in the control of orienting movements of the head, eyes, and pinnae in response to stimuli from different modalities (Irvine 1992; Oliver and Huerta 1992). IC fibers enter the brachium of the SC and terminate in the deep layers (Frisina et al. 1989; Appell and Behan 1990; Jiang et al. 1997). The IC source of most projections to the deep SC is the ICI (Covey et al. 1987; Thiele et al. 1996; King et al. 1998). The SC also receives sparse inputs from the ICc, in particular the dorsomedial part (Covey et al. 1987; Zhang et al. 1987; Frisina et al. 1997). A major source of input to the deep SC may be the rostral pole (rostral nucleus) of the IC (Harting and Van Lieshout 2000).

The IC also projects to deep layers of the contralateral caudal SC (Appell and Behan 1990). These crossed projections are sparser than the ipsilateral projections. The IC inputs to the contralateral deep SC originate from the ICc






Figure 6.4. IC inputs to premotor centers involved in pinna movements. The paralemniscal zone (*PLN*) is the premotor center for pinna movements and it projects to the facial motor nuclei (*FMN*) to innervate pinna muscles. The IC influence on pinna movements involves the superior colliculus. All three IC regions project to the deep SC with the lateral nucleus projection the heaviest. The deep SC, in turn, projects to the PLN ipsilaterally. The PLN projects predominantly to the ipsilateral FMN. A smaller projection from the central and lateral nuclei of the IC to the ipsilateral PLN may be an alternate route to the FMN.

(ventromedial part) and ICI subdivisions in several species (Covey et al. 1987; Zhang et al. 1987; Jiang et al. 1997; King et al. 1998). The projections from the IC to the deep SC may be glutamatergic and thus excitatory (Saint Marie 1996).

From the deep SC, the indirect IC pathway to FMN continues to the ipsilateral lateral paralemniscal zone (PLN) (Stein et al. 1984; Metzner 1996). The PLN, in turn, projects to the contralateral FMN (Fig. 6.4; Henkel and Edwards 1978; May et al. 1990; Takeuchi et al. 1993; Chen et al. 1995; Schuller et al. 1997).

The IC itself projects to the PLN. As described in cat, the PLN extends from the retrorubral nucleus (associated with the substantia nigra) to the VNLL (Henkel 1981; Herbert et al. 1997). The cat PLN receives input from the ICI ipsilaterally (Henkel 1981). A PLN has been defined in the horseshoe bat that is rostral and medial to the DNLL and INLL and it may receive input from the ipsilateral ICc (Metzner 1993, 1996).

It is also possible that IC inputs are relayed to pinna muscles via descending projections to other targets. Some lower IC targets project to the PLN and/or the FMN directly. For example, both the FMN (Hinrichsen and Watson

1983) and PLN (Takeuchi et al. 1993) receive projections from the periaqueductal gray. Other IC targets that project to the PLN are the sagulum and the dorsomedial periolivary nucleus of the SOC (Henkel 1981). Deep layers of the SC also project to the contralateral FMN (Hattox et al. 2002).

### 3.2. PREMOTOR/MOTOR CIRCUITS AND THE ACOUSTIC STARTLE RESPONSE

The acoustic startle response is characterized by the coordinated contraction of many craniofacial and skeletomotor muscle groups in response to a loud, brief sound. Sounds initiating the response enter the acoustic startle system via cochlear root neurons (CRNs). CRNs are found in the CN complex and they project to the ventrocaudal pontine reticular formation (VPRF) (Fig. 6.5; Lee et al. 1996; Nodal and López 2003). CRNs are not the only input to the VPRF; the latter




Figure 6.5. Connections within acoustic startle reflex pathways. The IC is one of two auditory centers providing acoustic input to a premotor center, the ventrocaudal pontine reticular formation, involved in the acoustic startle reflex. Cochlear root neurons provide for the rapid reflex component. Unlike the direct input of cochlear root neurons, the path of IC neurons to the ventrocaudal pontine reticular formation is indirect and involves three routes: these are via the ventrolateral tegmental nucleus, the dorsal cochlear nucleus, and the pedunculopontine tegmentum (via the deep superior colliculus). Each receives input primarily from the lateral and central IC nuclei. Each IC target projects to the ventrocaudal pontine reticular formation, the premotor center providing input to the spinal cord and facial motor nucleus, the brain stem hubs for the startle response.

receives many types and sources of input that could modulate this reflex. As an essential premotor center of the startle reflex, the VPRF integrates the information it receives and redistributes it to motoneurons in both the FMN and spinal cord, which control the muscles involved in the reflex.

Several studies have shown that the IC has a role in modulating the startle reflex (Leuman et al. 2001; Swerdlow et al. 2001; Yeomans et al. 2002). Like the FMN, spinal cord motor centers do not receive direct input from the IC. The FMN and the spinal cord probably receive input from the IC indirectly via the VPRF. The IC projections to the VPRF also may be indirect and may be relayed through at least three brain regions, including the ventrolateral tegmental nucleus, the pedunculo-pontine tegmental nucleus, and the DCN (Fig. 6.5). The ventrolateral tegmental nucleus is medial to the LL at the level of the VNLL and anterior to the rostral periolivary region of the SOC (Kandler and Herbert 1991). Neurons in the ICI project to the ventrolateral tegmental nucleus. The DCN is also a candidate as it receives descending IC input (as described earlier), and it projects to the VPRF. The third potential route is a bit more indirect, traveling through the deep SC and then to the pedunculo-pontine tegmentum.

Besides the anatomical evidence, behavioral data confirm the role of the IC and these three relay stations in the acoustic startle (Leuman et al. 2001; Swerdlow et al. 2001; Yeomans et al. 2002). Through one or more of these pathways, the IC may be involved in suppressing the acoustic startle response. In addition, the IC may modulate the startle response via its projections to the pontocerebellar pathway (Kandler and Herbert 1991). Finally, the IC may serve as a gating mechanism to coordinate the startle response with descending input from higher brain centers (see Chapter 8).

### *3.3. INPUTS TO PONTINE NUCLEI AND THE COORDINATION OF RESPONSES TO SOUND*

Some IC axons descend in the ipsilateral LL to enter the pontine gray. These are probably independent of axons that enter the SOC and that target auditory nuclei (Schuller et al. 1991). In the ipsilateral pontine gray, specific IC targets include the lateral, dorsolateral, and ventrolateral nuclei in several species (Mower et al. 1979; Burne et al. 1981; Aas 1989; Frisina et al. 1989; Wenstrup et al. 1994). Although the primary source of the IC projections to the pontine nuclei is the ICc, the ICI and ICd also contribute (Frisina et al. 1989; Schuller et al. 1991; Caicedo and Herbert 1993). The IC projection to the ipsilateral dorsolateral pontine nuclei is glutamatergic (Saint Marie 1996). In the mustached bat, the projections from the ICc to the ipsilateral dorsolateral and lateral pontine nuclei are topographic (Wenstrup et al. 1994).

Through its descending projections to the pontine gray, the IC (principally the ipsilateral central nucleus but also ICI) forms part of the pontocerebellar auditory pathway. Other auditory inputs to the pontine gray originate in belt

areas of the auditory cortex (Schuller et al. 1991) and the AVCN and PVCN (Kandler and Herbert 1991; Thompson 1998). It is thought that, in the cerebellum, sensory information is coordinated with motor feedback from the eyes, head, and body to control their movements (Schuller et al. 1991). Therefore, the pontine nuclei are believed to be the route by which the auditory system projects to the cerebellum to enable coordinated responses to sound.

It has also been hypothesized that, through its projections to the deep SC, the IC plays a role in sound orientation (Olazábal and Moore 1989). Neurons from the anterodorsal IC, in particular, may be involved. Interestingly, both the deep SC and this region of the IC receive input from the substantia nigra, which is presumed to have a modulatory role in orienting movements to visual and auditory stimuli.

#### 4. PROJECTIONS TO THE PERIAQUEDUCTAL GRAY AND VOCALIZATION

There is also evidence that the IC projects to the periaqueductal or central gray (PAG) (Schweizer 1981). The origin of this projection is the ICc bilaterally (Frisina et al. 1989). The dorsomedial ICc is a prominent source of input (Frisina et al. 1997). The ICc projection does not appear to terminate throughout the entire PAG, as the ipsilateral IC projections are restricted to its dorsolateral part. The neurons that project ipsilaterally may also project to the contralateral IC via collaterals that leave the main IC axons before they enter the commissure (Kudo and Niimi 1980).

It has been suggested that the direct IC projections to the PAG have a role in modulating vocalizations (Jürgens 2002). The PAG enables vocalizations through its projections to the nucleus retroambiguus, which projects to abdominal muscle motoneurons that increase abdominal pressure, and to nucleus ambiguus motor neurons that innervate the soft palate, pharynx, and larynx. It is hypothesized that the IC influences vocalizations implicated in communication and echolocation calls in certain bats (Casseday and Covey 1996; Fenzl and Schuller 2002).

There are also indirect routes by which the IC may affect vocalizations. One such potential route is via the deep SC, which projects to the nucleus ambiguus (Metzner 1996). Another indirect route is via the PLN, which also projects to the nucleus ambiguus (Metzner 1996). The reticular formation and the nucleus of the brachium of the IC may also contribute to the vocalization circuits (Pieper and Jürgens 2003).

#### 5. PROJECTIONS TO STRUCTURES INVOLVED IN EMOTIVE BEHAVIORS

The amygdala processes many types of aversive inputs and has been considered as a critical nexus for decision-making related to aversion (Brandão et al. 2003). The IC may affect input to the amygdala via its connections with the deep SC

and the dorsal PAG (Casseday et al. 2002), although the IC does not project directly to the amygdala. The likely IC source of the polysynaptic projections that ultimately reach the amygdala is the ICc (Pandossio and Brandão 1999). Therefore, the ICc, via its projections to the deep SC and the dorsal PAG, could influence the cascade of responses to aversive stimuli. Another potential route is from IC structures other than the ICc and that terminate in parts of the dorsal and medial divisions of the medial geniculate body (MGB) (Calford and Aitkin 1983; Wenstrup et al. 1994), which project in turn to the amygdala (Shinonaga et al. 1994).

## 6. INPUTS TO THE PARAGIGANTOCELLULAR NUCLEUS AND AROUSAL

Another set of descending IC projections end in caudal areas of the ipsilateral pontine reticular formation. These target the rostral part of the paragigantocellular nucleus, which is medial to the FMN (juxtafacial), and the gigantocellular reticular nucleus (Andrezik et al. 1981; Van Bockstaele et al. 1989; Caicedo and Herbert 1993). The IC inputs to the paragigantocellular nucleus arise from the ICc, ICd, and ICI (Caicedo and Herbert 1993; Van Bockstaele et al. 1993), whereas the projections to the gigantocellular reticular nucleus are more restricted, originating only from ICI.

The rostral paragigantocellular nucleus itself receives projections from many brain systems including somatic sensory, autonomic, and auditory systems (Van Bockstaele et al. 1993). Auditory input to it may be from the IC directly or from the IC indirectly via the PLN. The IC, perhaps with the PLN, may thus provide auditory input to the rostral paragigantocellular nucleus, where it is integrated with other sensory input. The integrated sensory information is likely then relayed to the locus coeruleus, a critical center for coordinating arousal (Berridge and Waterhouse 2003).

## 7. CONCLUSIONS

This review summarizes several monosynaptic and multisynaptic descending projections of the IC to the brain stem. Within the ascending auditory system, the IC targets the DCN and the DNLL. Most of the projections to these nuclei originate in the ICc, activate inhibitory neurons, and are arranged tonotopically and/or topographically. Both the DNLL and the DCN project to the IC. Therefore, these two descending IC pathways form a reciprocal circuit that could enhance auditory signal processing necessary for sound localization and speech processing. The descending IC pathway to the SOC feeds into the olivocochlear system and thus could modulate cochlear function.

Among other targets of descending IC pathways, the most notable is the deep SC. It is an integral component of the descending IC input to various targets.

This is likely the route by which the IC influences reflexive and other motor behaviors including pinna movements, acoustic startle reflexes, responses to aversive stimuli, and vocalizations.

The IC is also probably involved in modulating proprioceptive adjustments to sound by its descending projections to the cerebellum, via the pontine nuclei. This permits auditory access to many brain stem and spinal pathways that affect muscle tone and the initiation of movement. The IC also likely contributes to global arousal by forwarding acoustic information, albeit indirect, to a major arousal center, the locus coeruleus.

In conclusion, the descending IC connections may serve many roles, including auditory processing. They provide key information to various brain regions necessary for survival. The functions of the descending IC pathways should also be considered in context with the incoming projections from the forebrain (see Chapter 8). This circuitry connects the IC with higher and lower parts of the central nervous system (Huffman and Henson 1990). The IC serves an important role in gating information from higher to lower centers of the brain (Ehret 1997). More studies are necessary to delineate further the function of the descending IC system. In particular, the components and connections of the brain circuits necessary for survival and that rely on the descending IC pathways for acoustic information compel further investigation.

## Abbreviations

AVCN	anteroventral cochlear nucleus
CN	cochlear nucleus
CPO	caudal periolivary region
CRN	cochlear root neuron
DCN	dorsal cochlear nucleus
DMPO	dorsomedial periolivary nucleus
DNLL	dorsal nucleus of the lateral lemniscus
FMN	facial motor nucleus
GCD	granule cell domain
INLL	intermediate nucleus of the lateral lemniscus
LNTB	lateral nucleus of the trapezoid body
LSO	lateral superior olive
MGB	medial geniculate body
MOC	medial olivocochlear system
MSO	medial superior olive
OCN	olivocochlear nucleus neuron
OHC	outer hair cells
PAG	periaqueductal gray
PLN	paralemniscal zone
PVCN	posteroventral cochlear nucleus
RPO	rostral periolivary region
SC	superior colliculus

SOC	superior olivary complex
VNLL	ventral nucleus of the lateral lemniscus
VNTB	ventral nucleus of the trapezoid body
VPRF	ventrocaudal pontine reticular formation

## References

- Aas JE (1989) Subcortical projections to the pontine nuclei in the cat. *Journal of Comparative Neurology* **282**:331–354.
- Alibardi L (2002) Putative inhibitory collicular boutons contact large neurons and their dendrites in the dorsal cochlear nucleus of the rat. *Journal of Submicroscopic Cytology and Pathology* **34**:433–446.
- Andrezik JA, Chan-Palay V, and Palay SL (1981) The nucleus paragigantocellularis lateralis in the rat. Demonstration of afferents by the retrograde transport of horseradish peroxidase. *Anatomy and Embryology* **161**:373–390.
- Appell PP and Behan M (1990) Sources of subcortical GABAergic projections to the superior colliculus in the cat. *Journal of Comparative Neurology* **302**:143–158.
- Beneyto M, Winer JA, Larue DT, and Prieto JJ (1998) Auditory connections and neurochemistry of the sagulum. *Journal of Comparative Neurology* **401**:329–351.
- Berridge CW and Waterhouse BD (2003) The locus coeruleus-noradrenergic system: modulation of behavioral state and state-dependent cognitive processes. *Brain Research Reviews* **42**:33–84.
- Brandão ML, Troncoso AC, de Souza Silva MA, and Huston JP (2003) The relevance of neuronal substrates of defense in the midbrain tectum to anxiety and stress: empirical and conceptual considerations. *European Journal of Pharmacology* **463**:225–233.
- Braun M (2000) Inferior colliculus as candidate for pitch extraction: multiple support from statistics of bilateral spontaneous otoacoustic emissions. *Hearing Research* **145**:130–140.
- Burne RA, Azizi SA, Mihailoff GA, and Woodward DJ (1981) The tectopontine projection in the rat with comments on visual pathways to the basilar pons. *Journal of Comparative Neurology* **202**:287–307.
- Caicedo A and Herbert H (1993) Topography of descending projections from the inferior colliculus to auditory brainstem nuclei in the rat. *Journal of Comparative Neurology* **328**:377–392.
- Calford MB and Aitkin LM (1983) Ascending projections to the medial geniculate body of the cat: evidence for multiple, parallel auditory pathways through the thalamus. *Journal of Neuroscience* **3**:365–380.
- Carey C and Webster D (1971) Ascending and descending projections of the inferior colliculus in the kangaroo rat (*Dipodomys merriami*). *Brain, Behavior and Evolution* **4**:401–412.
- Casseday JH and Covey E (1996) A neuroethological theory of the operation of the inferior colliculus. *Brain, Behavior and Evolution* **47**:311–336.
- Casseday JH, Fremouw T, and Covey E (2002) The inferior colliculus: a hub for the central auditory system. In: Oertel D, Fay RR, and Popper AN (eds). *Springer Handbook of Auditory Research*, Volume 15: *Integrative Functions in the Mammalian Auditory Pathway*. Springer-Verlag, New York, pp. 238–318.
- Chen XH, Itoh M, Miki T, Ichiyama M, Fujimoto Y, Sun W, and Takeuchi Y (1995) Midbrain paralemniscal projections to the facial nucleus: an anatomical and immunohistochemical study. *Okajimas Folia Anatomica Japonica* **72**:69–79.

- Conlee JW and Kane ES (1982) Descending projections from the inferior colliculus to the dorsal cochlear nucleus in the cat: an autoradiographic study. *Neuroscience* **7**:161–178.
- Covey E, Hall WC, and Kobler JB (1987) Subcortical connections of the superior colliculus in the mustache bat, *Pteronotus parnellii*. *Journal of Comparative Neurology* **263**:179–197.
- Ehret G (1997) The auditory midbrain, a 'shunting-yard' of acoustical information processing. In: Ehret G and Romand R (eds). *The Central Auditory System*. Oxford University Press, New York, pp. 259–316.
- Fenzl T and Schuller G (2002) Periaqueductal gray and the region of the paralemniscal area have different functions in the control of vocalization in the neotropical bat, *Phyllostomus discolor*. *European Journal of Neuroscience* **16**:1974–1986.
- Frisina RD, O'Neill WE, and Zettel ML (1989) Functional organization of mustached bat inferior colliculus: II. Connections of the FM<sub>2</sub> region. *Journal of Comparative Neurology* **284**:85–107.
- Frisina RD, Walton JP, Lynch-Armour MA, and Klotz DA (1997) Efferent projections of a physiologically characterized region of the inferior colliculus of the young adult CBA mouse. *Journal of the Acoustical Society of America* **101**:2741–2753.
- Fujino K and Oertel D (2003) Bidirectional synaptic plasticity in the cerebellum-like mammalian dorsal cochlear nucleus. *Proceedings of the National Academy of Sciences of the United States of America* **100**:265–270.
- Harting JK and Van Lieshout DP (2000) Projections from the rostral pole of the inferior colliculus to the cat superior colliculus. *Brain Research* **881**:244–247.
- Hattox AM, Priets CA, and Keller A (2002) Functional circuitry involved in the regulation of whisker movements. *Journal of Comparative Neurology* **442**:266–276.
- Henkel CK (1981) Afferent sources of a lateral midbrain tegmental zone associated with the pinnae in the cat as mapped by retrograde transport of horseradish peroxidase. *Journal of Comparative Neurology* **203**:213–226.
- Henkel CK and Edwards SB (1978) The superior colliculus control of pinna movements in the cat: possible anatomical connections. *Journal of Comparative Neurology* **182**:763–776.
- Herbert H, Klepper A, and Ostwald J (1997) Afferent and efferent connections of the ventrolateral tegmental area in the rat. *Anatomy and Embryology* **196**:235–259.
- Hinrichsen CF and Watson CD (1983) Brain stem projections to the facial nucleus of the rat. *Brain, Behavior and Evolution* **22**:153–163.
- Huffman RF and Henson OW Jr (1990) The descending auditory pathways and acousticomotor systems: connections with the inferior colliculus. *Brain Research Reviews* **15**:295–323.
- Hurley L and Thompson AM (2001) Serotonergic innervation of the auditory brainstem of the Mexican free-tailed bat, *Tadarida brasiliensis*. *Journal of Comparative Neurology* **435**:78–88.
- Irvine DRF (1992) Auditory brainstem processing. In: Popper AN and Fay RR (eds) *Springer Handbook of Auditory Research, Volume 2: The Mammalian Auditory Pathway: Neurophysiology*. Springer-Verlag, New York, pp. 153–231.
- Jiang ZD, Moore DR, and King AJ (1997) Sources of subcortical projections to the superior colliculus in the ferret. *Brain Research* **755**:279–292.
- Jürgens U (2002) Neural pathways underlying vocal control. *Neuroscience and Biobehavioral Reviews* **26**:235–258.



- Kandler K and Herbert H (1991) Auditory projections from the cochlear nucleus to pontine and mesencephalic reticular nuclei in the rat. *Brain Research* **562**:230–242.
- Kane ES (1977) Descending inputs to the cat dorsal cochlear nucleus: an electron microscopic study. *Journal of Neurocytology* **6**:583–605.
- Kane ES and Conlee JW (1979) Descending inputs to the caudal cochlear nucleus of the cat: degeneration and autoradiographic studies. *Journal of Comparative Neurology* **187**:759–784.
- King AJ, Jiang ZD, and Moore DR (1998) Auditory brainstem projections to the ferret superior colliculus: anatomical contribution to the neural coding of sound azimuth. *Journal of Comparative Neurology* **390**:342–365.
- Kudo M and Niimi K (1980) Ascending projections of the inferior colliculus in the cat: an autoradiographic study. *Journal of Comparative Neurology* **191**:545–556.
- Lee Y, López DE, Meloni EG, and Davis M (1996) A primary acoustic startle pathway: obligatory role of cochlear root neurons and the nucleus reticularis pontis caudalis. *Journal of Neuroscience* **16**:3775–3789.
- Leumann L, Sterchi D, Vollenweider F, Ludewig K, and Früh H (2001) A neural network approach to the acoustic startle reflex and prepulse inhibition. *Brain Research Bulletin* **56**:101–110.
- Malmierca MS, Le Beau FEN, and Rees A (1996) The topographical organization of descending projections from the central nucleus of the inferior colliculus in guinea pig. *Hearing Research* **93**:167–180.
- May PJ, Vidal PP, and Baker R (1990) Synaptic organization of tectal-facial pathways in cat. II. Synaptic potentials following midbrain tegmental stimulation. *Journal of Neurophysiology* **64**:381–402.
- Metzner W (1993) An audio-vocal interface in echolocating horseshoe bats. *Journal of Neuroscience* **13**:1899–1915.
- Metzner W (1996) Anatomical basis for audio-vocal integration in echolocating horseshoe bats. *Journal of Comparative Neurology* **368**:252–269.
- Mower G, Gibson A, and Glickstein M (1979) Tectopontine pathway in the cat: laminar distribution of cells of origin and visual properties of target cells in dorsolateral pontine nucleus. *Journal of Neurophysiology* **42**:1–15.
- Mulders WHAM and Robertson D (2000) Effects on cochlear responses of activation of descending pathways from the inferior colliculus. *Hearing Research* **149**:11–23.
- Mulders WHAM and Robertson D (2002) Inputs from the cochlea and the inferior colliculus converge on olivocochlear neurones. *Hearing Research* **167**:206–213.
- Nodal FR and López DE (2003) Direct input from cochlear root neurons to pontine reticulospinal neurons in albino rat. *Journal of Comparative Neurology* **460**:80–93.
- Ohlroge M, Doucet JR, and Ryugo DK (2001) Projections of the pontine nuclei to the cochlear nucleus in rats. *Journal of Comparative Neurology* **436**:290–303.
- Olazábal UE and Moore JK (1989) Nigrotectal projection to the inferior colliculus: horseradish peroxidase transport and tyrosine hydroxylase immunohistochemical studies in rats, cats, and bats. *Journal of Comparative Neurology* **282**:98–118.
- Oliver DL and Huerta MF (1992) Inferior and superior colliculi. In: Webster DB, Popper AN, and Fay RR (eds). *Springer Handbook of Auditory Research*, Volume 1: *The Mammalian Auditory Pathway: Neuroanatomy*. Springer-Verlag, New York, pp. 168–221.
- Ostapoff E-M, Morest DK, and Potashner SJ (1990) Uptake and retrograde transport of [<sup>3</sup>H]GABA from the cochlear nucleus to the superior olive in the guinea pig. *Journal of Chemical Neuroanatomy* **3**:285–295.

- Pandossio JE and Brandão ML (1999) Defensive reactions are counteracted by midazolam and muscimol and elicited by activation of glutamate receptors in the inferior colliculus of rats. *Psychopharmacology* **142**:360–368.
- Pieper F and Jürgens U (2003) Neuronal activity in the inferior colliculus and bordering structures during vocalization in the squirrel monkey. *Brain Research* **979**:153–164.
- Saint Marie RL (1996) Glutamatergic connections of the auditory midbrain: selective uptake and axonal transport of D-[<sup>3</sup>H]aspartate. *Journal of Comparative Neurology* **373**:255–270.
- Saldaña E, Bajo VM, López DE, and Malmierca MS (1989) Is the descending projection from the central nucleus of the inferior colliculus to the cochlear nuclear complex tonotopically organized? *European Journal of Neuroscience Supplement* **2**:262.
- Scates KW, Woods CI, and Azerado WJ (1999) Inferior colliculus stimulation and changes in 2f1-f2 distortion product otoacoustic emissions in the rat. *Hearing Research* **128**:51–60.
- Schofield BR (2001) Origins of projections from the inferior colliculus to the cochlear nucleus in guinea pigs. *Journal of Comparative Neurology* **429**:206–220.
- Schofield BR and Cant NB (1999) Descending auditory pathways: projections from the inferior colliculus contact superior olivary cells that project bilaterally to the cochlear nuclei. *Journal of Comparative Neurology* **409**:210–223.
- Schuller G, Covey E, and Casseday JH (1991) Auditory pontine grey: connections and response properties in the horseshoe bat. *European Journal of Neuroscience* **3**:648–662.
- Schuller G, Fischer S, and Schweizer H (1997) Significance of the paralemiscal tegmental area for audio-motor control in the moustached bat, *Pteronotus p. parnellii*: the afferent and efferent connections of the paralemiscal area. *European Journal of Neuroscience* **9**:342–355.
- Schweizer H (1981) The connections of the inferior colliculus and the organization of the brainstem auditory system in the greater horseshoe bat (*Rhinolophus ferrum-equinum*). *Journal of Comparative Neurology* **201**:25–49.
- Shinonaga Y, Takada M, and Mizuno N (1994) Direct projections from the non-laminated divisions of the medial geniculate nucleus to the temporal polar cortex and amygdala in the cat. *Journal of Comparative Neurology* **340**:405–426.
- Shore SE and Moore JK (1998) Sources of input to the cochlear granule cell region in the guinea pig. *Hearing Research* **116**:33–42.
- Shore SE, Helfert RH, Bledsoe SC, Altschuler RA, and Godfrey DA (1991) Descending projections to the dorsal and ventral divisions of the cochlear nucleus in guinea pig. *Hearing Research* **52**:255–268.
- Stein BE, Spencer RF, and Edwards SB (1984) Efferent projections of the neonatal cat superior colliculus: facial and cerebellum-related brainstem structures. *Journal of Comparative Neurology* **230**:47–54.
- Swerdlow NR, Geyer MA, and Braff DL (2001) Neural circuit regulation of prepulse inhibition of startle in the rat: current knowledge and future challenges. *Psychopharmacology* **156**:194–215.
- Takeuchi Y, Chen XH, Fukui Y, Itoh M, Miki T, and Miyoshi S (1993) Midbrain tegmentum as an intermediate relay station of the periaqueductal-facial pathway in the cat. *Brain Research Bulletin* **31**:57–65.
- Thiele A, Rübsamen R, and Hoffmann K-P (1996) Anatomical and physiological investigation of auditory input to the superior colliculus of the echolocating megachiropteran bat *Rousettus aegyptiacus*. *Experimental Brain Research* **112**:223–236.

- Thompson AM (1998) Heterogeneous projections of the cat posteroventral cochlear nucleus. *Journal of Comparative Neurology* **390**:439–453.
- Thompson AM and Thompson GC (1993) Relationship of descending inferior colliculus projections to olivocochlear neurons. *Journal of Comparative Neurology* **335**:402–412.
- Van Bockstaele EJ, Pieribone VA, and Aston-Jones G (1989) Diverse afferents converge on the nucleus paragigantocellularis in the rat ventrolateral medulla: retrograde and anterograde tracing studies. *Journal of Comparative Neurology* **290**:561–584.
- Van Bockstaele EJ, Akaoka H, and Aston-Jones G (1993) Brainstem afferents to the rostral (juxtafacial) nucleus paragigantocellularis: integration of exteroceptive and interoceptive sensory inputs in the ventral tegmentum. *Brain Research* **603**:1–18.
- Vater M and Feng AS (1990) Functional organization of ascending and descending connections of the cochlear nucleus of horseshoe bats. *Journal of Comparative Neurology* **292**:373–395.
- Vetter DE, Saldaña E, and Mugnaini E (1992) Input from the inferior colliculus to medial olivocochlear neurons in the rat: a double label study with PHA-L and cholera toxin. *Hearing Research* **70**:173–186.
- Warr WB (1992) Organization of olivocochlear efferent systems in mammals. In: Webster DB, Popper AN, and Fay RR (eds). *Springer Handbook of Auditory Research*, Volume 1: *The Mammalian Auditory Pathway: Neuroanatomy*. Springer-Verlag, New York, pp. 410–448.
- Wenstrup JJ, Larue DT, and Winer JA (1994) Projections of physiologically defined subdivisions of the inferior colliculus in the mustached bat: targets in the medial geniculate body and extrathalamic nuclei. *Journal of Comparative Neurology* **346**:207–236.
- Wright DD and Ryugo DK (1996) Mossy fiber projections from the cuneate nucleus to the cochlear nucleus in the rat. *Journal of Comparative Neurology* **365**:159–172.
- Yeomans JS, Li L, Scott LL, and Frankland PW (2002) Tactile, acoustic and vestibular systems sum to elicit the startle reflex. *Neuroscience and Biobehavioral Reviews* **26**: 1–11.
- Young ED and Davis KA (2002) Circuitry and function of the dorsal cochlear nucleus. In: Oertel D, Fay RR, and Popper AN (eds). *Springer Handbook of Auditory Research*, Volume 15: *Integrative Functions in the Mammalian Auditory Pathway*. Springer-Verlag, New York, pp. 160–206.
- Zhang SQ, Sun XD, and Jen PH-S (1987) Anatomical study of neural projections to the superior colliculus of the big brown bat, *Eptesicus fuscus*. *Brain Research* **416**:375–380.

# Chapter 7

## The Tectothalamic System

JEFFREY J. WENSTRUP

### 1. INTRODUCTION

As the midbrain auditory center, the inferior colliculus (IC) integrates ascending auditory brain stem projections with descending projections from the auditory cortex, aminergic inputs, and, in some species, limbic projections. Although physiologic properties of IC neurons clearly reflect their inputs, many neurons display novel responses arising by integration of these inputs. The IC then projects to the auditory thalamus, to lower auditory nuclei, and to nuclei at the sensorimotor interface. This chapter focuses on the projection of the IC to the medial geniculate body (MGB), its main target.

Like the IC, the MGB integrates input from multiple ascending and descending sources, but these are more restricted. Its major input is from the IC but it also receives auditory brain stem input and nonauditory input. Its principal output is to auditory cortical areas, while other significant targets include the amygdala and the frontal cortex.

Representations of acoustic signals differ among MGB subdivisions. This chapter considers the IC contribution to subdivisional differences. Are these a function of different ascending inputs, or the different operations each subdivision performs? How does the organization of acoustic responses differ between the IC and the MGB? For information on MGB structure and function, please see reviews (Clarey et al. 1992; Winer 1992; Wenstrup 1995; de Ribaupierre 1997).

### 2. PROJECTIONS OF THE INFERIOR COLLICULUS TO THE MGB

#### *2.1. ORGANIZATION OF THE TECTOTHALAMIC PROJECTION*

The MGB contains dorsal (D), ventral (V), and medial (M) divisions (Fig. 7.1). V is the MGB component of the ascending lemniscal pathway. Its connections relate it to the central nucleus of the IC (CN) and primary auditory cortex (AI).




Figure 7.1. Neuronal architecture of MGB in cat, based on Golgi-impregnated neurons. The section at left is more caudal. *D*, Dorsal nucleus; *DD*, deep dorsal nucleus; *DS*, superficial dorsal nucleus; *M*, medial division; *MZ*, marginal zone; *Ov*, ovoid part of the ventral division; *Sg*, supragenulate nucleus; *VL*, ventrolateral nucleus of the dorsal division; *VI*, lateral part of the ventral division. (Adapted from Morest 1964.)

Physiologically, its cells have robust, short-latency auditory responses with sharp frequency tuning. The dorsal division contains, at minimum, dorsal (*D*), deep dorsal (*DD*), superficial dorsal (*DS*), ventrolateral (*VL*), and supragenulate (*Sg*) nuclei. *D* nuclei have diverse inputs, outputs, and functional properties, and show major species variations. They receive most input from regions outside the CN and project to secondary auditory cortical areas. *Sg* and the rostral dorsal nucleus also project to the amygdala, mediating emotional responses to auditory stimuli. Auditory responses in *D* are typically weaker, more broadly tuned, and more plastic than in *V*, but the diversity and species variability also distinguish it. *M*, rarely subdivided, receives input from the lateral (*La*) or external nucleus of the IC and brain stem auditory nuclei. It projects widely to the auditory cortex and the amygdala. Auditory responses in *M*, again relative to *V*, usually have longer latency and broader frequency tuning.

The IC projection to MGB is complex (Fig. 7.2). Restricted tracer deposits in mustached bat CN show projections to four or more MGB subdivisions, including *V*, *M*, and at least two *D* nuclei (Wenstrup and Grose 1995). Across species, the IC projects to each MGB subdivision, but the projections of IC subdivisions differ. The CN projects most strongly to *V*, *M*, and *DD*. The dorsal cortex (*DC*) projects to *D*, including *D* and *VL* nuclei. The *La* projects to *D* and *M*. The IC also projects to perigeniculate regions including the cat posterior thalamus (Andersen et al. 1980), rat posterior paralaminar nucleus (LeDoux et al. 1987; Linke 1999), and the cat and bat pretectum (Kudo et al. 1983; Frisina et al. 1989; Wenstrup et al. 1994; Fig. 7.2B). The tectothalamic projection shows considerable species variability (Table 7.1).




Figure 7.2. Projections to MGB from two frequency-band representations of the central nucleus of IC in the mustached bat. In IC sections (*left*) are electrode penetrations, characteristic frequencies at multiunit recording sites (in kHz), and tracer deposit sites (*black regions*). Anterograde MGB labeling (*dots*) was plotted in a caudal (*left*) to rostral (*right*) series of transverse sections. Numbers (*lower left*) show the IC or MGB section level. (**A**) Widespread projections from the 29- to 30-kHz IC representation terminate in the midbrain and thalamus. Significant MGB labeling occurs in V (lateral part), M, D, Sg, and the rostral pole nucleus (of the dorsal division). (**B**) Different input to MGB and pretectum from the 85- to 86-kHz IC representation. Labeling in MGB includes the medial part of V, M, D, and two zones of RP. *bic*, Brachium of the IC; *Pt*, pretectum; *RP*, rostral pole nucleus of the dorsal division. (Adapted from Wenstrup and Grose 1995.)

The IC–MGB projection is predominantly ipsilateral. In cats and rats, 5% to 10% of tectothalamic cells are contralateral (Aitkin and Phillips 1984; Rouiller and de Ribaupierre 1985; González-Hernández et al. 1991). V receives the strongest contralateral IC input. In mice, guinea pigs, and bats, there is little or no evidence of a contralateral projection (Wenstrup 1995; Frisina et al. 1997; Malmierca et al. 1997).

What sets of IC and other inputs are received by each MGB subdivision? This question is best addressed by retrograde transport studies using restricted tracer deposits, but such studies have been rare. Additional evidence comes from

Table 1: Species Differences in IC Inputs to MGB Subdivisions

MGB Nucleus	Species <sup>1</sup>	CN	DC	La	MGB Nucleus	Species	CN	DC	La
Ventral	All	●	•	•	Dorsal	C	•	●	•
Medial Division	C	●	—	●	MB	●	—	—	—
	MB	●	×	×	R	—	•	●	—
	R	•	•	●	TS <sup>2</sup>	—	—	—	—
	TS	•	•	●	Deep Dorsal	C	●	•	●
Suprageniculate					MB <sup>3</sup>	●	—	—	—
					R	×	×	×	—
					TS	—	●	—	—
					Ventrolateral	C	—	●	•
				MB	×	×	×	—	
				R	—	●	●	●	
				TS	—	●	—	—	

• ● ● Relative strength of projection  
 — No projection to MGB subdivision  
 × Undetermined

<sup>1</sup>C, cat; MB, mustached bat; R, rat; TS, tree shrew  
<sup>2</sup>DC input is to the anterodorsal nucleus  
<sup>3</sup>Correspondence to DD suggested by Olsen and Suga, 1991a and Wenstrup 1999

anterograde transport studies, in which the convergence of input must be assessed across several experiments. Nonetheless, it is clear that MGB subdivisions receive different but overlapping sets of inputs.

Alone among MGB divisions, V receives a nearly unitary ascending projection. Strong input is from the CN, with weak DC and La projections (Calford and Aitkin 1983; LeDoux et al. 1985; Rouiller and de Ribaupierre 1985), a conclusion corroborated by anterograde transport and lesion studies (Oliver and Hall 1978; Andersen et al. 1980; LeDoux et al. 1985). The CN to V projection is topographic/tonotopic, although it has been documented in few species. In cat (Kudo and Niimi 1978; Andersen et al. 1980; Calford and Aitkin 1983) and mustached bat (Wenstrup et al. 1994), low-frequency CN representations end in the lateral part of V, with higher frequency input medially. This differs in the guinea pig: low-frequency dorsal CN regions target dorsal and medial V, while higher frequency, ventral CN regions project to the ventrolateral V (Malmierca et al. 1997). These projection patterns agree with the tonotopy revealed in physiologic studies of V (see Section 5).

While preserving tonotopy, the CN–V projection may alter the representation of bands of different frequencies. In mustached bats, frequency organization is preserved but proportional representation is substantially altered. Thus, CN frequency bands analyzing frequency-modulated components of sonar signals send only a small projection to V; most output terminates in D (Frisina et al. 1989; Wenstrup et al. 1994; Wenstrup 1999). Such an unequal division of CN output to different MGB subdivisions may occur in other species. In cats, for example,

the DD nucleus receives input from high-frequency but not low-frequency CN regions (Andersen et al. 1980; Calford and Aitkin 1983; Rouiller and de Ribaupierre 1985).

The CN projection may contribute to regional differences in responses across V (Rodrigues-Dageaff et al. 1989; Cetas et al. 2001), but reorganization of CN output is not well documented. In cats, single CN tracer deposits form discontinuous labeling patterns in V (Andersen et al. 1980), suggesting that CN loci project divergently. Such patterns may confer different binaural response regions onto V (Andersen et al. 1980; Middlebrooks and Zook 1983). In the guinea pig discontinuities in the CN–V projection are not apparent (Malmierca et al. 1997).

IC inputs to D are more complex because each subdivision receives distinct sets of auditory midbrain inputs and major species variations occur. In cats and tree shrews, the DC projects to rostral parts of the D nucleus (Oliver and Hall 1978; Andersen et al. 1980; Calford and Aitkin 1983; Rouiller and de Ribaupierre 1985). In rats, the La projects more strongly than the DC to the D nucleus (LeDoux et al. 1985). Mustached bat CN frequency bands corresponding to constant-frequency elements of sonar signals have the strongest input to the D nucleus (Wenstrup et al. 1994; Wenstrup and Grose 1995).

The DD nucleus receives input from the CN, DC, and La, with major species differences. In cats input is from the higher frequency CN, with lesser DC and La input (Andersen et al. 1980; Calford 1983; Rouiller and de Ribaupierre 1985). Similarly, the mustached bat has strong, higher frequency CN input to a region corresponding to the DD (Olsen and Suga 1991a; Wenstrup 1999); these regions receive lesser input from La and the pericollicular tegmentum and none from the DC (Wenstrup et al. 1994; Wenstrup and Grose 1995). In contrast, tree shrews (Oliver and Hall 1978) and rodents (Frisina et al. 1997; Malmierca et al. 1997) send little CN input to DD, with the main input from the DC (Oliver and Hall 1978).

The suprageniculate nucleus (Sg) receives variable input across species. In cats, Sg retrograde tracer deposits labeled only the nucleus of the brachium of IC (nBIC) and the superior colliculus (SC) (Calford and Aitkin 1983), but such limited projections may reflect tracer insensitivity. Degeneration or anterograde transport methods show brain stem inputs traversing the lateral tegmentum (Morest 1965), some from the vicinity of the ventral nucleus of the lateral lemniscus (Henkel 1983). In mustached bats, the Sg receives SC input and significant auditory midbrain and brain stem inputs, including those from the nucleus of the central acoustic tract (NCAT) (Casseday et al. 1989), part of the La or the midbrain tegmentum (Gordon and O'Neill, 2000), and the CN (Frisina et al. 1989; Wenstrup et al. 1994). In rats, the major input is from the SC and specific layers of the La (LeDoux et al. 1987; Linke 1999), with a smaller DC input and little from the CN. Both visual and somatic sensory inputs converge on the Sg (LeDoux et al. 1987).

M integrates a broad range of inputs, with major projections from the La and the CN (Calford 1983) and other auditory projections from the lateral tegmental system (Morest 1965; Oliver and Hall 1978), sagulum (Hutson et al. 1991),



ventral nucleus of the lateral lemniscus (Henkel 1983), and the dorsal cochlear nucleus (Malmierca et al. 2002). There is no evidence for tonotopic projections to M (Wenstrup et al. 1994; Malmierca et al. 2002) but there is a coarse frequency map in the cat (Rouiller et al. 1989). This may arise from the La, which receives topographic CN input (Saldaña and Merchán 1992). Like Sg, M receives multisensory input, including somatic sensory (LeDoux et al. 1987) and vestibular (Blum et al. 1979) projections.

## 2.2. CELLULAR FEATURES OF THE TECTOTHALAMIC PROJECTION

Disc-shaped IC neurons form the major tectothalamic cell type. In the cat CN, 80% of projection neurons are varieties of disc-shaped cells; stellate cells are also present (Oliver 1984). In rats, only disc-shaped CN cells project (Oliver et al. 1999), but both stellate and disc-shaped cells project from the La and the DC (Oliver et al. 1999). Sagulum neurons projecting to the MGB include small stellate and larger multipolar neurons (Beneyto et al. 1998).

Although most tectothalamic input is likely glutamatergic, a major finding was the discovery of a  $\gamma$ -aminobutyric acid (GABA)-ergic IC–MGB projection. In cats 20% (CN) to 30% (La) of tectothalamic neurons are GABAergic (Fig. 7.3A) (Winer et al. 1996). In rats about 45% of CN tectothalamic cells are GABAergic (Fig. 7.3A) (Peruzzi et al. 1997). Tectothalamic GABAergic neurons had sizes and morphologies corresponding to those of IC GABAergic cell types (Oliver et al. 1994; Winer et al. 1996). On average, GABAergic BIC axons were larger than nonGABAergic axons (Saint Marie et al. 1997). These results suggest that IC monosynaptic feedforward inhibition to MGB may precede excitatory input. This is a major departure from the patterns of thalamic somatic sensory and visual inputs, where ascending GABAergic inputs are sparse (De Biasi et al. 1990; Pow et al. 1994). The functional properties of this GABAergic input are considered below.

The morphology of tracer-filled tectothalamic axons corroborate earlier Golgi and degeneration findings (Morest 1964; Majorossy and Réthelyi 1968). IC terminals in the MGB are knoblike or en passant endings (Fig. 7.3) with a range of sizes. In V, both types occur, although knoblike clusters dominate (Fig. 7.3B: 1) (Pallas and Sur 1994; Wenstrup et al. 1994; Bartlett et al. 2000). These endings are complex, as observed in ferrets and mustached bats (Fig. 7.3B: 3). As in cats (Morest 1965), preterminal axons in mustached bats are aligned with dendrites of tufted neurons (Fig. 7.3B: 3) (Wenstrup et al. 1994; Winer and Wenstrup 1994a,b). This arrangement is thought to conserve a precise frequency representation in V. In the rabbit MGB, a close relationship between dendritic/cellular laminae and the fine frequency representation of V has been documented (Cetas et al. 2001, 2002).

D neurons receive small, en passant terminals and larger knoblike endings (Fig. 7.3B: 2). These are similar across species, even when the input source differs, with terminals in cats, rats, and bats matching closely, despite the fact that the major input arises in different IC subdivisions (Morest 1964; Wenstrup




Figure 7.3. Cellular features of the tectothalamic projection. (A) GABAergic IC neurons projecting to MGB in cats and rats. *Filled squares (left)* show cells labeled both by MGB injections of a retrograde tracer and by anti-GABA antibody. *Small dots*, cells labeled only by the retrograde tracer. (Adapted from Winer et al. 1996. Copyright 1996 National Academy of Sciences, USA.) *Right, filled circles* show cells labeled both by MGB injections of retrograde tracer and by anti-GABA antibody. *Open squares*, cells labeled only by the retrograde tracer; *open triangles*, GABA-positive cells. (Adapted from Peruzzi et al. 1997.) (B) IC inputs to the MGB. Boutons from rat ventral (1) and dorsal (2) divisions after IC tracer deposits. (Adapted from Bartlett et al. 2000.) 3, Labeled terminals and preterminal axons after tracer deposits in lower frequency representation of the mus-

et al. 1994; Bartlett et al. 2000). Likewise, the cat Sg is dominated by fine, en passant boutons from the SC and the nBIC (Calford and Aitkin 1983), while CN afferents to Sg in mustached bats are strikingly similar (Wenstrup et al. 1994). The relationship between these morphologic features and response properties remains unclear.

Terminals in the rat MGB were studied with electron microscopy, GABA immunohistochemistry, and tract tracing (Fig. 7.3B: 4) (Bartlett et al. 2000). Tectothalamic axodendritic endings are plentiful, and axosomatic terminals sparse, as in earlier studies (Morest 1964). Their distribution onto MGB neurons depends partly on their size and neurotransmitter. Larger glutamatergic terminals end on primary through tertiary dendrites; a few smaller, presumptive glutamatergic terminals are axosomatic; and most small terminals end on distal dendrites. In D, there are more small, presumptive glutamatergic IC terminals on thinner dendrites. GABAergic terminals are typically smaller and target secondary and higher order dendrites. Thus, some GABAergic IC input may be slow or weak, although GABAergic IC axons are larger than nonGABAergic axons (Saint Marie et al. 1997). Such IC termination patterns may underlie temporal patterns of excitation and inhibition (Bartlett and Smith 1999).

Noncollicular terminals are either large GABAergic terminals, probably from thalamic reticular nucleus cells, or glutamatergic corticothalamic terminals (Winer et al. 1999; Bartlett et al. 2000). Compared to tectothalamic inputs, corticothalamic terminals are smaller and target higher order dendrites. There are species differences in the confluence of collicular, thalamic reticular, and cortical input in MGB. In cats and monkeys, triadic synaptic arrangements involving GABAergic interneurons, collicular afferents, and corticothalamic terminals are common (Jones and Rockel 1971; Morest 1975; Majorossy and Kiss 1976). In contrast, rats and some bats have few intrinsic GABAergic MGB neurons (Winer and Larue 1988, 1996; Vater et al. 1992; Winer et al. 1992) and consequently few triads (Bartlett et al. 2000).

In summary, the tectothalamic projection shows common features across species, but also major differences that challenge understanding of the structure–function relationships of this projection. Most puzzling is the lack of consistent alignments among phylogenetic groups with respect to anatomic features of the projection. For example, considering the projection from CN to D, rodents are at one extreme (no CN input), mustached bats at the other (strong CN input),

---

Figure 7.3. *Continued*

tached bat's IC follow the laminar arrangement of MGB principal cell dendrites. (Adapted from Wenstrup et al. 1994). The scale bar applies to 1–3. 4, Schematic distribution of rat IC and other terminals onto MGB neurons show few axosomatic contacts. *Squares*, IC inputs of different sizes; *circles*, auditory cortical terminals of different sizes; *triangles*, large inputs of unknown origin; *filled symbols*, GABAergic terminals; *open symbols*, glutamatergic terminals. Numbers in **(B)**: 4 indicate order of dendrites. *CB*, Cell body; *HO*, higher order dendrites. (Adapted from Bartlett et al. 2000. Copyright 2000 with permission from Elsevier.)

and cats are intermediate. GABAergic MGB neurons are rare in rats and some bats, while cats and monkeys have >20%. The functional implications of these species differences are not understood.

### 3. SYNAPTIC AND INTRINSIC PROPERTIES OF MEDIAL GENICULATE BODY NEURONS

It has long been recognized that auditory input to the MGB triggers interactions between excitation and inhibition (Nelson and Erulkar 1963; Aitkin and Dunlop 1968). Cellular bases for these interactions were investigated using rat MGB explant (Hu et al. 1994; Hu 1995) or slice (Peruzzi et al. 1997; Bartlett and Smith 1999) preparations. When the BIC is shocked, V and D neurons show fast, reliable excitatory postsynaptic potentials (EPSPs) that are insensitive to changes in stimulation rate and have little temporal jitter. In the absence of inhibition, EPSP latencies average <2.5 ms and are completely blocked by simultaneous application of antagonists to *N*-methyl-D-aspartate (NMDA) and  $\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) glutamate receptors. Such EPSPs reflect monosynaptic glutamatergic IC inputs to V and D, and probably M (Webber et al. 1999).

In V, all IC-evoked EPSP or spike responses were blocked by AMPA receptor antagonists, but few were reduced by NMDA antagonists (Fig. 7.4A) (Hu et al. 1994). In contrast, EPSPs in D were reduced by both NMDA and AMPA receptor antagonists (Fig. 7.4A). The NMDA-mediated responses could sometimes evoke action potentials in D without AMPA receptor-mediated depolarization. In M, AMPA and NMDA receptor antagonists each blocked most action potentials evoked by BIC stimulation (Webber et al. 1999).

Postsynaptic effects of IC GABAergic input to the MGB are varied (Fig. 7.4B) (Peruzzi et al. 1997; Bartlett and Smith 1999). In about one third of V and D cells there was no evidence of GABA<sub>A</sub> receptor-mediated IPSPs before or after the EPSP evoked by BIC stimulation (type EX/0). Most such neurons also lacked GABA<sub>B</sub> receptor-mediated inhibition activated by the IC afferents. In some 38% of V and D cells, GABA<sub>A</sub>-mediated IPSPs occurred before the glutamatergic EPSP (type IN/EX), while in 14% the IPSPs followed the EPSP (type EX/IN). In 11%, IPSPs were not accompanied by EPSPs after BIC activation (type 0/IN). Neurons showing GABA<sub>A</sub>-mediated inhibitory postsynaptic potentials (IPSPs) also had longer latency and longer GABA<sub>B</sub>-mediated IPSPs.

MGB action potentials elicited by BIC activation result from interaction among excitatory, inhibitory, and intrinsic cellular properties. Like other thalamic neurons (Jahnsen and Llinás 1984; see Sherman 2001), MGB neurons have distinct firing modes associated with levels of alertness (Fig. 7.4C) (Steriade and Llinás 1988; McCormick and Feeseer 1990). For MGB *in vitro* preparations (Hu et al. 1994; Tennigkeit et al. 1996), one mode is tonic discharge to intracellular current pulses, followed by a return to near-resting membrane potentials.

Discharge latency is short (a few milliseconds) following EPSPs or applied current, and the interspike interval is longer. Tonic mode occurs in neurons at slightly depolarized resting membrane potentials. At slightly hyperpolarized potentials, they may still fire tonically but latency increases significantly and spike probability declines. If the membrane potential is further hyperpolarized, cells discharge in burst mode;  $\text{Na}^+/\text{K}^+$  action potentials at short interspike intervals ride atop a slower depolarization resulting from a low-threshold  $\text{Ca}^{2+}$ -dependent spike (Steriade et al. 1990; Hu 1995; Tennigkeit et al. 1996). Action potential latency is longer and more variable than in the tonic mode.

In both rat explant (Hu et al. 1994; Hu 1995) and anesthetized guinea pig preparations (He and Hu 2002), lemniscal V neurons fired tonically, whereas nonlemniscal D neurons fired in burst mode. Hu and colleagues related several physiologic characteristics of D neurons to the burst mode firing: longer EPSPs, more NMDA receptor-mediated glutamate excitation, more negative resting membrane potentials, and no depolarizing “sag” mediated by the  $I_h$  current. They concluded that such intrinsic and synaptic properties explain major differences between MGB lemniscal and nonlemniscal systems, including firing mode, latency, regularity of firing, and perhaps plasticity.

Other studies have challenged these conclusions. In rat slices, single V neurons show a range of firing modes, latencies, and temporal discharge patterns that are dependent on the cell's membrane potential and the applied current (Fig. 7.4C) (Tennigkeit et al. 1996, 1997). Cells shift from burst to tonic mode with additional stimulation during the burst mode response, and can be influenced by metabotropic receptors to glutamate, GABA, and acetylcholine (Mooney et al. 1995; Tennigkeit et al. 1998, 1999). Few differences in intrinsic membrane properties were seen in V and D neurons, or even between stellate and tufted neurons (Bartlett and Smith 1999). An exception was the concentration in V of the hyperpolarization-activated depolarizing  $I_h$  current, which may impede burst mode firing by reducing membrane hyperpolarization. These studies do not support a sharp distinction between D and V neurons with regard to intrinsic properties or firing mode. Perhaps the firing mode is more indicative of alertness levels (Tennigkeit et al. 1996, 1997) as in other thalamic nuclei. Burst mode firing may have an alerting function that provides lower quality information to the cerebral cortex, whereas tonic mode firing may typify high-quality analyses of auditory features in awake animals (Tennigkeit et al. 1997; Sherman 2001). Such a speculation cannot be tested *in vitro*.

Response properties of guinea pig MGB neurons were tested in awake, sleeping, and anesthetic states with electroencephalograph (EEG) monitoring. Bursts occurred in awake animals, but increased during slow-wave sleep and were most common under barbiturate anesthesia (Massaux and Edeline 2003). In awake animals, there was more V burst firing than in D, M, and the lateral part of the posterior complex (Pol), but no differences during slow-wave sleep. Frequency selectivity during slow-wave or paradoxical sleep was sharper than in awake animals, although sensitivity and spike number declined (Edeline et al. 2000). There was no correlation between burst mode firing and changes in frequency




Figure 7.4. Physiologic properties of MGB neurons. (A–C, E) Synaptic and intrinsic behavior from in vitro preparations. (A) Postsynaptic potentials in D neuron after electrical stimulation of BIC. Bath-application of CNQX and APV separately remove early

receptive fields. These studies strongly suggest that burst firing mode is part of the repertoire of both lemniscal and nonlemniscal MGB neurons, and demonstrate firing mode dependence on levels of alertness. They imply that the information about acoustic stimuli conveyed by burst firing neurons is as precise (for sound frequency) as in neurons firing tonically.

The interplay of synaptic and intrinsic properties in MGB neurons thus transforms their temporal discharge pattern. Such patterns are shaped by a neuron's membrane potential and regulated both by intrinsic properties and short and longer term actions of synaptic inputs. Multiple firing modes occur throughout the MGB and do not distinguish any subdivision, even though neurons in a subdivision may burst more under particular experimental conditions or levels of alertness.

#### 4. TRANSFORMATIONS IN FUNCTIONAL PROPERTIES

MGB auditory responses differ across subdivisions. Typically, V neurons display sharp tuning, short latencies, and reliable responses to sound. D neurons have longer latencies, broader tuning, and less reliable and more labile responses. M neurons vary, with population means often between those of V and D. Some D and M neurons respond to other sensory modalities. These differences hold across species and different levels of anesthesia, although species differences and anesthesia effects can be profound. This contributes to the view that both the tectothalamic projection and MGB functional properties embody parallel (but incompletely segregated) ascending systems whose origin, termination, and functional properties differ. Physiologic support for this hypothesis is plentiful (Calford 1983; Lennartz and Weinberger 1992; Bordi and LeDoux 1994a; Edeline et al. 1999).

This section considers whether MGB neural responses depend on *de novo* processing within MGB or reflect the IC responses or other inputs. Addressing

---

Figure 7.4. *Continued*

and late synaptic response components, respectively. Combined, they eliminate the post-synaptic response, showing that glutamate mediates IC excitatory input. (Adapted from Hu et al. 1994.) **(B)** Postsynaptic responses after BIC stimulation show excitatory and inhibitory IC inputs. The vertical scale differs across four neurons. (Adapted from Bartlett and Smith 1999.) **(C)** Firing modes of the V neuron depend on intracellular potential. (Adapted from Tennigkeit et al. 1997.) **(D)** First-spike latency difference in CN and V neurons from *in vivo* recording of free-tailed bats (Adapted from Klug et al. 2000.) **(E)** IC GABAergic input stabilizes first-spike latency of MGB neurons. EX/0 neuron (*above*) and IN/EX neuron (*below*) spikes elicited by BIC stimulation. Each cell was tested over a range of membrane potentials, which alter neuronal firing mode. The EX/0 neuron displays large changes in first-spike latency over a range of membrane potentials, whereas the IN/EX cell has no change in latency over a range of membrane potentials. *sa*, stimulus artifact. (Adapted from Bartlett and Smith 1999.)

this issue requires two types of experiments, of which few examples exist. The first includes *in vivo* MGB microiontophoretic studies that test auditory responses before and during application of neurotransmitter antagonists, a method that has identified novel transformations in the IC. The second type of study directly compares IC and MGB recordings in the same species under identical anesthetic and recording conditions. Because both species and experimental variables create significant response disparities, comparisons in which these factors differ are problematic.

#### 4.1. TEMPORAL PROPERTIES

##### 4.1.1. Latency and Temporal Response Patterns

MGB neurons differ significantly in response latencies, from <10 ms to >100 ms. Likewise, there are different temporal response patterns, including onset (ON), sustained, offset (OFF), ON–OFF, and variable patterns (Calford 1983; Rodrigues-Dagaëff et al. 1989; Rouiller et al. 1989; Edeline et al. 1999). For both first spike timing and the temporal spiking pattern, differences exist among subdivisions. Thus, V latencies are shortest and least variable, and temporal response patterns the most homogeneous. D latencies are longest and most variable, while M responses and those in Pol are intermediate. Large differences exist even in V, where latency, temporal response pattern, and other features form caudorostral gradients (Rodrigues-Dagaëff et al. 1989).

Regional differences may be amplified by anesthesia. Large differences in latencies were found in barbiturate-anesthetized cats (Calford 1983). In awake guinea pigs whose level of alertness was monitored with EEG recordings, smaller regional differences persist (Edeline et al. 1999); V, M, and Pol latencies are longer than in the cat while D latencies are substantially shorter. Species differences, and possibly experimental procedures, confound comparisons from awake guinea pigs and little brown bats. In this bat, regional latency differences are small, averaging 8.9 ms in V and 12.8 ms in D and M (Llano and Feng 1999).

In awake free-tailed bats, latencies in V range from 15.9 to 12.8 ms for characteristic frequency (CF) sounds 10 to 50 dB above threshold, respectively, exceeding values in CN by 3 to 4 ms (Fig. 7.4D) (Klug et al. 2000). Axonal and synaptic delays may explain such latency differences, as average EPSP latencies in V after BIC electrical stimulation are 1.1 to 3.8 ms, depending on early GABAergic inhibition (Bartlett and Smith 1999). In awake mustached bats, different types of D neurons (Olsen and Suga 1991b; O'Neill and Brimijoin 2002) reveal strong auditory responses at latencies only a little longer than auditory midbrain or brain stem responses (Portfors and Wenstrup 1999; Gordon and O'Neill 2000). Thus, in awake animals there is little evidence of major response latency transformations except those introduced by axonal and synaptic delays. This is supported by an extensive physiologic survey of IC properties



(Syka et al. 2000) in ketamine/xylazine-anesthetized guinea pigs. Although the use of anesthesia precludes a direct comparison to an MGB study (Edeline et al. 1999), each IC subdivision displays a broad range of latencies. Thus, the IC input latencies, brief synaptic delays, and the effects of GABAergic IC inputs probably account for the range of latencies across MGB subdivisions. In addition, GABAergic input can maintain constant response latencies under different firing modes (Fig. 7.4E; Bartlett and Smith 1999).

In all subdivisions in awake guinea pigs, ON responses form the largest group although these are fewer in D (Edeline et al. 1999). Sustained responders, substantially fewer than ON cells, form the next largest group. OFF, ON-OFF, and other categories are least common. A similar preponderance of ON responses occurs in awake little brown bats (Llano and Feng 1999), barbiturate-anesthetized guinea pigs (He 2001), and cats under barbiturate or light nitrous oxide anesthesia (Calford 1983; Rodrigues-Dageaff et al. 1989). Although IC discharge patterns vary across species and anesthetic regimens, sustained response patterns seem more common than in MGB (Bock et al. 1972; Ehret and Moffat 1985; Semple and Kitzes 1985; Aitkin et al. 1994). This suggests that one tectothalamic transformation increases responsiveness to dynamic rather than static stimuli.

Such transformations in temporal discharge properties appear unrelated to state-dependent changes in discharge mode caused by neuronal intrinsic and synaptic properties (see Section 3). It has been suggested that ON patterns elicited by auditory stimuli in anesthetized animals indicate burst mode firing (Tennigkeit et al. 1996). This does not appear to be the case, however. Even in EEG-monitored awake guinea pigs, in which most responses are in “tonic” firing mode, ON patterns predominate (Edeline et al. 1999, 2000; Massaux and Edeline 2003). Discharges during tonic mode firing are probably restricted either by inhibitory inputs or by the ON pattern of excitatory inputs, rather than by mode of firing.

#### 4.1.2. Response to Signal Periodicities

Temporal sensitivity to signal periodicities has been studied using periodic amplitude modulations, repetitive click or tone burst trains, or responsiveness to paired click or tonal stimuli. Although many central auditory neurons discharge synchronously to stimulus modulations or repetitive stimuli, this capacity diminishes at successively higher levels (Langner 1992). MGB neurons show modest ability to synchronize to temporal events in sounds (Aitkin and Dunlop 1968; Rodrigues-Dageaff et al. 1989) and this correlates with sharp frequency tuning (Lennartz and Weinberger 1992). However, there are few direct comparisons between IC and MGB responses in awake animals.

In awake little brown bats, temporal sensitivity to rate variations in tone burst trains was examined in the IC, the MGB, and the auditory cortex. Among MGB cells, modulation rate transfer functions based on response synchrony had a low-

pass cutoff at about 64 pulses/s. V neurons had the highest cutoffs, but these were different only from M neurons (Llano and Feng 1999). Comparable studies of IC and auditory cortex found synchronization cutoffs of 163 and 37 pulses/s, respectively (Condon et al. 1994, 1997), suggesting a progressive reduction in the preservation of fine temporal information in the tectothalamocortical pathway. In a few MGB neurons, the response to a second tone was facilitated but discharge synchrony decreased. Responsiveness to pulse trains improved when a slower amplitude modulation was imposed on the pulse train (Llano and Feng 1999). This response, which was more noticeable in AI, may reflect a change from temporal to rate coding of signal periodicities at higher levels of the auditory system.

MGB neurons recorded *in vitro* show either response depression or facilitation to the second of two shocks to the BIC (Bartlett and Smith 2002). The effect was related to the presence and timing of IC glutamatergic and GABA<sub>A</sub> receptor-mediated events. Type EX/O neurons in MGB, for which a large IC glutamatergic EPSP is unaccompanied by IC GABA<sub>A</sub>-mediated IPSPs, showed only paired-pulse depression. These are thought to originate from larger glutamatergic IC synaptic inputs. The depression occurred despite chemical blockade of GABA<sub>A</sub> and GABA<sub>B</sub> receptors and was apparent for either AMPA or NMDA receptor-mediated EPSPs. Paired-pulse MGB depression is likely mediated presynaptically. In other thalamic nuclei with little ascending GABAergic input, paired-pulse depression is the rule (Eysel 1976; Mishima 1992). In MGB, however, V and D neurons could show paired-pulse facilitation, the dominant effect among IN/EX neurons. The facilitation depended on NMDA receptors, but its relation to GABA<sub>A</sub>-mediated inhibition is unclear. Such neurons had smaller EPSPs and presumably smaller IC glutamatergic synapses on the MGB cell. These results suggest a complex relationship between synaptic ending size, glutamate pharmacology, and the presence and timing of inhibition.

Functional magnetic resonance imaging (fMRI) in awake humans shows decreased responsiveness to higher modulation rates (Giraud et al. 2000; Harms and Melcher 2002), supporting earlier work in various animal models (see Langner 1992). The major change between IC and MGB is that fMRI IC activation is nearly continuous during noise burst trains from 1 to 35 Hz. The MGB response at higher rates is largest at the burst train onset. The change in fMRI activation between IC and MGB is part of a transformation from the IC to the cortex, in which activation at higher repetition rates is phasic.

In summary, compared to IC cells, MGB neurons have longer latencies, greater discharge probability to sound onset, and lower discharge synchrony to repetitive stimuli. There is little experimental evidence that latencies extend beyond predicted axonal and synaptic delays. However, other temporal features such as onset responses and synchronized responses to transients are likely modified by MGB processing and interactions between excitatory and inhibitory IC inputs. Finally, a major transformation in the MGB appears to be the state-dependent modification of discharge mode, which affects latency, temporal discharge pattern, and perhaps other temporal response indices.

## 4.2. SPECTRAL PROPERTIES

Cells in V and Pol typically have the sharpest frequency tuning, while in D and M tuning is broader (Calford 1983; Bordi and LeDoux 1994a; Edeline et al. 2000). “Broad” tuning includes multiple forms of spectrally complex responses, such as preferences for broad- vs. narrow-band stimuli, multi-peaked tuning curves, inhibitory frequency bands, and combination sensitivity. The anesthetic and the behavioral state affect frequency tuning, such that the complexity of spectral responses, even in V, is often greater in lightly anesthetized or awake animals (Allon et al. 1981; Morel et al. 1987; Edeline et al. 2000). Frequency tuning differences persist across divisions in awake guinea pigs but are weaker (Edeline et al. 1999, 2000). Despite divisional differences in frequency tuning, little evidence exists of transformations in tuning sharpness and spectrally complex responses from the IC to the MGB.

### 4.2.1. Sharpness of Frequency Tuning

One study has directly addressed how MGB inhibitory interactions contribute to frequency tuning. Blockade of GABA<sub>A</sub> receptor-mediated inhibition affects frequency tuning of sharply tuned neurons in the MGB of awake mustached bats (Suga et al. 1997); recording sites were in either V or D, both of which receive their major input from CN (Wenstrup et al. 1994; Wenstrup and Grose 1995). Local bicuculline application broadened frequency tuning in 37% of neurons and often increased dynamic range at CF. The authors concluded that inhibitory interactions sharpened frequency tuning in a minority of cells. The small number of GABAergic MGB cells in this bat (Vater et al. 1992; Winer et al. 1992) suggests that the inhibition is of midbrain origin. Because tuning sharpness in MGB neurons was intermediate between auditory nerve and the AC, Suga and colleagues suggested that frequency tuning is sharpened at several levels of the ascending auditory pathway.

### 4.2.2. Spectral Integration: Responses to Two-Tone Stimuli

Two-tone stimuli, combining tones of different frequencies, permit an assessment of neural responses to complex sounds. Two-tone responses were studied in the context of monaural sound localization cues in barbiturate-anesthetized cats (Imig et al. 1997). Monaurally directional neurons, using monaural cues to create spatial selectivity, are about 17% of cells in V and Pol. They derive their spatial selectivity from inhibitory spectral interactions; their complex frequency receptive fields have one or more inhibitory bands and some display multi-peaked excitatory tuning curves. In comparable IC recordings, slightly more monaural IC directional neurons were found than in MGB, but with somewhat weaker sideband inhibition (Poirier et al. 2003). Thus, the spectral interactions characteristic of monaurally directional MGB and auditory cortex (AC) neurons are present in IC. Although the barbiturate anesthetic used in these studies may

enhance inhibitory response elements, the results support the conclusion (Suga et al. 1997) that some frequency sharpening occurs in the MGB.

#### 4.2.3. Combination Sensitivity

Combination sensitivity refers to neural responses showing preference for particular spectrotemporal combinations of acoustic elements (Suga et al. 1978; Fuzessery and Feng 1983; Margoliash and Fortune 1992; Rauschecker et al. 1995; Ohlemiller et al. 1996). These neurons require distinct frequency-tuned inputs and special temporal properties to achieve their response selectivities. Such integrative features are usually viewed as a forebrain attribute for audition, vision, or somatic sensation. In the mustached bat, however, this integrative feature originates in IC or at subcollicular levels (Mittmann and Wenstrup 1995; Portfors and Wenstrup 1999, 2001b; Wenstrup and Leroy 2001).

A comparison of IC and MGB found sharper temporal sensitivity and stronger facilitation among combination-sensitive MGB neurons (Yan and Suga 1996). In contrast, other studies in the IC (Portfors and Wenstrup 1999) and the MGB (Wenstrup 1999) found no significant differences in facilitatory interactions. They suggested that the earlier IC data reflect multiple-unit responses that, summed across different neurons, failed to show the strength and sharp temporal features of facilitation observed in single-unit recordings. Additional studies (Leroy and Wenstrup 2000) found that other forms of spectrally complex responses observed in MGB (Wenstrup, 1999) are also seen in the IC, including interactions between frequency bands not used in biosonar and noise-selective responses. Thus, most evidence suggests that basic features of combination-sensitive responses in the mustached bat originate in IC or below and undergo little MGB modification.

#### 4.2.4. Frequency Modulation

Responses to temporal modulations of stimulus frequency have rarely been investigated in the MGB, limiting comparison to the more extensive IC literature. Studies of selectivity for direction and rate of frequency sweeps in the nuclei of the lateral lemniscus (NLL), IC, and MGB of awake mustached bats (Gordon and O'Neill 2000; O'Neill and Brimijoin 2002) found that a ventral region of the external nucleus of the IC (ICvx) more strongly prefers upward frequency sweeps than either NLL or CN neurons, although responses to CF tones are generally strong. Directional preferences likely result from a low-threshold inhibitory sideband above the unit's CF. ICvx, with short latency responses in the range of 5 to 8 ms, receives its main input from the nucleus of the central acoustic tract (NCAT) and projects to both the SC and the Sg, which also are NCAT targets. More Sg neurons display preferences for upward than downward frequency sweeps, but respond well to CF tones, like ICvx neurons. Overall, the preference for upward sweeps is less in Sg than in ICvx. Perhaps this reflects the range of inputs to Sg from NCAT (sweep selectivity unknown), ICvx (upward preference), and CN (less directional preference) (O'Neill and Brimijoin

2002). The range of Sg response latencies (5 to 25 ms) is consistent with this range of inputs. Upward sweep selectivity may contribute to the bat's analysis of social vocalizations. These studies suggest that diverse response properties within an MGB subdivision result from diverse auditory inputs, rather than from diverse computations performed on the tectothalamic input.

#### 4.3. *BINAURAL SENSITIVITY AND SPATIAL SELECTIVITY*

MGB binaural and spatial processing has been studied mostly in cats, but these studies are far fewer in number than those in IC and AC. All MGB subdivisions show a range of responses to binaural sounds (Calford 1983). Sensitivity to interaural phase disparities (IPDs) is rare in D, common in V and M, and abundant in Pol (Ivarsson et al. 1988). Because the latter study employed nitrous oxide anesthesia, comparison with IC studies using barbiturate anesthesia is problematic (Kuwada and Yin 1983; Irvine and Gago 1990; Delgutte et al. 1999). However, the MGB may have a more continuous distribution of IPD sensitivity, whereas IPD-sensitive IC neurons favor the contralateral field (Kuwada and Yin 1983). Similarly, more neurons are driven by the ipsilateral ear, although the numbers are small. The increased representation of the ipsilateral sound field may result from contralateral auditory tectothalamic projection in cats (Clarey et al. 1992).

Monaural and binaural mechanisms create spatial selectivity in higher frequency neurons of the cat IC (Poirier et al. 2003), MGB (Imig et al. 1997; Samson et al. 2000), and AC (Samson et al. 1993, 1994). Monaural directional neurons retain azimuthal location selectivity with one ear plugged. Monaural cues from frequency-dependent head-related transfer functions may be encoded by IC, MGB, and AC units with distinct excitatory and inhibitory frequency domains revealed by two-tone stimuli (Imig et al. 1997). Binaural directional neurons lose azimuthal selectivity when an ear is plugged. They use binaural level cues encoded through inhibitory and/or facilitatory mechanisms that compare binaural inputs. The relative proportion of directional neurons based on monaural vs. binaural cues is constant in the IC, MGB (V and Pol), and AC (area AI). Further, there is little change in numbers of inhibitory frequency domains of monaurally directional neurons or in the strength of spectral inhibition. The chief alteration between IC and MGB is an increased proportion of neurons with enhanced responses to ipsilateral sound.

## 5. *PHYSIOLOGIC ORGANIZATION*

Do organizational features change between IC subdivisions and their MGB counterparts? In V, as in CN, the dominant feature is tonotopy. CF increases from lateral to medial and caudal to rostral in V for cats (Aitkin and Webster 1972; Imig and Morel 1985), squirrel monkeys (Gross et al. 1974), and mustached bats (Wenstrup 1999) (Fig. 7.5A). In lagomorphs and rodents, CF

increases from dorsal to ventral, with species variants (Redies and Brandner 1991; Bordi and LeDoux 1994a; Cetas et al. 2001). These patterns agree with each species' organization of CN inputs and with the arrangement of dendrites and tectothalamic terminals, insofar as these are known (see Sections 2.1, 2.2).

In the mustached bat, however, the IC–MGB projection reorganizes frequency. In the CN, the frequency organization is disproportionate but in accord with the cochlear map (Zook et al. 1985; O'Neill et al. 1989). However, anatomic (Frisina et al. 1989; Wenstrup et al. 1994; Wenstrup and Grose 1995) and physiologic (Wenstrup 1999) evidence shows that the tectothalamic projection radically alters this organization. V maintains an ordered CF progression, but certain frequency bands that enjoy large CN representations, those devoted to analyzing frequency-modulated (FM) elements of sonar signals, are diminished in V (Fig. 7.5A) and are, instead, represented in the rostral part of D. Whether such alterations occur in other species is unknown. This seems unlikely in rodents as CN projects mainly to V. In cats, divergent CN projections to V, DD, Pol, and M, and the high-frequency bias in DD neurons, raise this possibility.

Within V, several response gradients or zones of response segregation occur in cats (Calford and Webster 1981; Rodrigues-Dagaëff et al. 1989), guinea pigs (Redies and Brandner 1991), and rabbits (Cetas et al. 2001). In cats, there are related caudorostral gradients of responsiveness to clicks, noise and tone bursts, sharpness of tuning, temporal response pattern, and responsiveness to repetitive stimulation. More rostral cells perform more precise spectral and temporal analyses; they usually are more sharply tuned, respond to a broader range of stimuli, have shorter latencies, display ON responses to noise and tone bursts, and respond better to click trains. These gradients are independent of CF, suggesting that they occur within frequency band representations. Binaural response categories have a patchy distribution. The caudorostral functional gradient was matched by differential connections to AC. Whether these gradients reflect IC arrangements or are imposed by reorganizing the tectothalamic projection is unknown (Rodrigues-Dagaëff et al. 1989).

Few reports of response organization in D are available, although He (2001) described a topographic distribution of OFF and ON–OFF responses in barbiturate-anesthetized guinea pigs. About 18% of D cells had invariant OFF responses to auditory stimuli, whereas 22% showed ON–OFF properties. OFF neurons lay in parts of D, M, and the shell subdivision surrounding V. This may be a novel MGB organization because it is absent in the IC of ketamine-anesthetized guinea pigs (Syka et al. 2000). Interpretation of these results is not straightforward, however, as anesthetic regimens affect temporal response patterns. In the MGB of awake guinea pigs, OFF cells comprise only 3% of the recorded population, with ON–OFF cells forming another 3% (Edeline et al. 1999).

In the mustached bat, D contains an organization of frequency bands representing higher harmonics of the FM component of the sonar call (from 48 to 59 kHz, from 72 to 89 kHz, and from 96 to 119 kHz) (Wenstrup 1999). Two




Figure 7.5. MGB functional organization. **(A)** Schematic tonotopy in V of four mammals in transverse sections. Caudal sections are on the *left*, rostral on the *right*. *Numbers*, characteristic frequency. A tonotopic pattern is present in each species, but the orientation varies. (Cat, adapted from Imig and Morel 1985; mustached bat, adapted from Wenstrup 1999; guinea pig, adapted from Redies and Brandner 1991; rabbit, adapted from Cetas et al. 2001.) **(B)** Transformation of frequency tuning and delay sensitivity from IC (*left*) to MGB (*right*) in the mustached bat. Frequency reorganization shown for three bands corresponding to harmonic components of FM sonar signal (FM2, 48 to 59 kHz; FM3, 72 to 89 kHz; FM4, 96 to 118 kHz). Reorganization of delay tuning is shown only for FM3 representation. *Dots*, best delay of neurons to combinations of the fundamental FM component in the sonar pulse (FM1, 24 to 29 kHz) and the delayed FM3 component of echoes. (Adapted from Portfors and Wenstrup 2001a. Copyright 2001 with permission from Elsevier.)

features are noteworthy (Fig. 7.5B). First, for each of the three frequency bands, there are two segregated maps in the rostral D and they are not arranged in a frequency gradient. Nearly all the neurons are combination-sensitive, tuned to delays between a 24 to 29 kHz signal (the fundamental FM component of sonar signals) and a later higher harmonic FM signal crossing the unit's CF. A second feature concerns delay sensitivity. Neurons within each frequency map display a rough organization of delay sensitivity not seen in CN single-unit studies (Portfors and Wenstrup 2001a). Thus, novel organizations of frequency and delay sensitivity may be created by the tectothalamic projection, setting the stage for thalamocortical projections that result in functionally distinct AC areas. The cortical neurons tuned to both sonar harmonics and delay after a 24 to 29 kHz signal are thought to analyze the sonar echo delays (O'Neill and Suga 1982; Suga et al. 1983; Fitzpatrick et al. 1998).

In M of lightly anesthetized cats, there is a tonotopic gradient, especially rostrally (Rouiller et al. 1989). It is weaker than in V, with no organization of response properties orthogonal to the tonotopic axis. M response properties vary widely but no other systematic representation is reported.

In the rostral and medial MGB, responses to other sensory modalities occur (Wepsic 1966; Phillips and Irvine 1979). Somatic sensory responses occur in rat Sg, M, rostral parts of D, the adjacent posterior intralaminar nuclei, and the posterior complex (Bordi and LeDoux 1994b). These and multimodal (auditory and somatic sensory) responses concentrate in the rostral MGB, in agreement with the distribution of somatic sensory input to the auditory thalamus (LeDoux et al. 1987). Some neurons could be activated antidromically from the lateral amygdala, indicating a role in conditioned emotional responses to sounds (Bordi and LeDoux 1994b).

In summary, anatomic and physiologic studies in several species show that the tonotopic organization present in CN is conferred on V, and it may be substantially altered by the tectothalamic projection. Within V, auditory responses and cortical connections show significant caudorostral variation, suggesting different functional roles. In other MGB divisions, there are few indications that responses are organized or segregated, in part because it has been difficult to identify, especially in D, the stimuli that optimally excite the neuronal population.

## 6. SYNTHESIS AND PERSPECTIVE

Multiple auditory pathways through the thalamus form multiple streams of auditory information processing that serve many behavioral and perceptual capabilities. Underlying these auditory streams are distinctive anatomic and physiologic features of different MGB subdivisions. The major question is the origin of these streams. Are they created *de novo* in the MGB or do lower auditory nuclei impose their functional properties and organization onto MGB neurons? There are several conclusions related to this question.



### *6.1. TRANSFORMATIONS IN TEMPORAL PROCESSING FROM MIDBRAIN TO THALAMUS*

The transformations noted earlier are common throughout the ascending auditory pathway and do not distinguish MGB integration. Thus, MGB latencies are longer, the neurons synchronize less well to repetitive acoustic transients, and transient discharge patterns may increase. Feedforward GABAergic inhibition may contribute to these phenomena. Such inhibition characterizes many auditory nuclei, but it is unusual in the thalamus. In contrast, multiple firing modes are a hallmark of thalamic neurons and distinguish them from other auditory nuclei. These state-dependent MGB firing modes alter both latency and discharge pattern.

A mystery related to inhibition and temporal properties among MGB neurons is the role of intrinsic vs. tectothalamic GABAergic inhibition (see Chapters 1, 2, and 22). There are species differences in the source and size of these inputs. How are differences in these forms of GABAergic inhibition manifest in temporal or other encoding features of MGB neurons?

### *6.2. SERIAL TRANSFORMATIONS IN OTHER INTEGRATIVE RESPONSE PROPERTIES*

There is no evidence that novel forms of spectrotemporal integration such as multiple excitatory or inhibitory tuning peaks, combination sensitivity, duration tuning, directional selectivity to FM sweeps, or of binaural integration are created in the MGB, and the evidence that response selectivities are substantially sharpened is modest. Creation or extensive modification of response properties are more characteristic of processing in the IC and the AC.

There are several caveats to this conclusion. First, an exception may be the auditory and somatic sensory responses of the rostral and medial parts of the MGB. Because these regions receive both auditory and somatic sensory input, some neurons may integrate inputs to form bimodal responses. However, at least one input, the IC lateral nucleus, is multimodal, making it difficult to discern whether such integration occurs in the MGB. A second caveat is limited understanding of the functional relevance of auditory responses in M and D, regions that have received little attention and require further scrutiny. Identifying their optimal stimuli, delineating the functional properties of their cortical or subcortical targets, and understanding the genesis of their responses as related to their inputs remain as future challenges.

### *6.3. EMERGENT RESPONSE PROPERTIES*

The reorganization of IC and other projections in MGB creates pathway-specific assortments of response properties that distinguish projections to auditory cortex and other targets. This reorganization takes several forms. One is the rearrangement of IC outputs to create novel organization in MGB subdivisions and, in

turn, cortical areas. This is evident in the mustached bat, where projections from CN reorganize to create functionally distinct MGB subdivisions that then project to specific cortical regions associated with the analyses of particular information-bearing elements in biosonar signals. Another form is the convergence of input from multiple sources that creates regions with a broad range of response properties, as in the medial and rostral MGB. In this view, the MGB does not create novel response properties, but instead creates novel ensembles of response properties. A major challenge is to understand the functional roles played by these ensembles of neurons.

## Acknowledgment

This work was supported by National Institutes of Health Research Grant R01 DC00937 from the National Institute on Deafness and Other Communication Disorders.

## Abbreviations

AC	auditory cortex
AMPA	$\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid
BIC, bic	brachium of the inferior colliculus
CF	characteristic frequency
CG	central gray
CN	central nucleus of the inferior colliculus
CUN	cuneiform nucleus
D	dorsal division <i>or</i> dorsal nucleus of the medial geniculate body
DC	dorsal cortex of the inferior colliculus
DD	deep dorsal nucleus of the medial geniculate body
DS	superficial dorsal nucleus of the medial geniculate body
EPSP	excitatory postsynaptic potential
FM	frequency-modulated
fMRI	functional magnetic resonance imaging
GABA	$\gamma$ -aminobutyric acid
IC	inferior colliculus
ICvx	ventral region of the external nucleus of the inferior colliculus
IPD	interaural phase disparity
IPSP	inhibitory postsynaptic potential
La	lateral nucleus of the inferior colliculus
M	medial division of the medial geniculate body
MGB	medial geniculate body
nBIC	nucleus of the brachium of the inferior colliculus
NBICm	nucleus of the brachium of the inferior colliculus, medial part
NCAT	nucleus of the central acoustic tract

NLL	nuclei of the lateral lemniscus
NMDA	<i>N</i> -methyl-D-aspartate
PAG	periaqueductal gray
Pol	lateral part of the posterior complex of the thalamus
Pt	pretectum
SC	superior colliculus
Sg	supragenulate nucleus of the medial geniculate body
V	ventral division <i>or</i> ventral nucleus of the medial geniculate body
VL	ventrolateral nucleus of the medial geniculate body
VI	ventral division, lateral part
Vm	ventral division, medial part

## References

- Aitkin LM and Dunlop CW (1968) Interplay of excitation and inhibition in the cat medial geniculate body. *Journal of Neurophysiology* **31**:44–61.
- Aitkin LM and Webster WR (1972) Medial geniculate body of the cat: organization and responses to tonal stimuli of neurons in ventral division. *Journal of Neurophysiology* **35**:365–380.
- Aitkin LM and Phillips SC (1984) The interconnections of the inferior colliculi through their commissure. *Journal of Comparative Neurology* **228**:210–216.
- Aitkin L, Tran L, and Syka J (1994) The responses of neurons in subdivisions of the inferior colliculus of cats to tonal, noise and vocal stimuli. *Experimental Brain Research* **98**:53–64.
- Allon N, Yeshurun Y, and Wollberg Z (1981) Responses of single cells in the medial geniculate body of awake squirrel monkeys. *Experimental Brain Research* **41**:222–232.
- Andersen RA, Roth GL, Aitkin LM, and Merzenich MM (1980) The efferent projections of the central nucleus and the pericentral nucleus of the inferior colliculus in the cat. *Journal of Comparative Neurology* **194**:649–662.
- Bartlett EL and Smith PH (1999) Anatomic, intrinsic, and synaptic properties of dorsal and ventral division neurons in rat medial geniculate body. *Journal of Neurophysiology* **81**:1999–2016.
- Bartlett EL and Smith PH (2002) Effects of paired-pulse and repetitive stimulation on neurons in the rat medial geniculate body. *Neuroscience* **113**:957–974.
- Bartlett EL, Stark JM, Guillery RW, and Smith PH (2000) Comparison of the fine structure of cortical and collicular terminals in the rat medial geniculate body. *Neuroscience* **100**:811–828.
- Beneyto M, Winer JA, Larue DT, and Prieto JJ (1998) Auditory connections and neurochemistry of the sagulum. *Journal of Comparative Neurology* **401**:329–351.
- Blum PS, Abraham LD, and Gilman S (1979) Vestibular, auditory, and somatic input to the posterior thalamus of the cat. *Experimental Brain Research* **34**:1–9.
- Bock GR, Webster WR, and Aitkin LM (1972) Discharge patterns of single units in inferior colliculus of the alert cat. *Journal of Neurophysiology* **35**:265–277.
- Bordi F and LeDoux JE (1994a) Response properties of single units in areas of rat auditory thalamus that project to the amygdala. I. Acoustic discharge patterns and frequency receptive fields. *Experimental Brain Research* **98**:261–274.

- Bordi F and LeDoux JE (1994b) Response properties of single units in areas of rat auditory thalamus that project to the amygdala. II. Cells receiving convergent auditory and somatosensory inputs and cells antidromically activated by amygdala stimulation. *Experimental Brain Research* **98**:275–286.
- Calford MB (1983) The parcellation of the medial geniculate body of the cat defined by the auditory response properties of single units. *Journal of Neuroscience* **3**:2350–2364.
- Calford MB and Aitkin LM (1983) Ascending projections to the medial geniculate body of the cat: evidence for multiple, parallel auditory pathways through thalamus. *Journal of Neuroscience* **3**:2365–2380.
- Calford MB and Webster WR (1981) Auditory representation within principal division of cat medial geniculate body: an electrophysiological study. *Journal of Neurophysiology* **45**:1013–1028.
- Casseday JH, Kobler JB, Isbey SF, and Covey E (1989) Central acoustic tract in an echolocating bat: an extralemniscal auditory pathway to the thalamus. *Journal of Comparative Neurology* **287**:247–259.
- Cetas JS, Price RO, Velenovsky DS, Sinex DG, and McMullen NT (2001) Frequency organization and cellular lamination in the medial geniculate body of the rabbit. *Hearing Research* **155**:113–123.
- Cetas JS, Price RO, Velenovsky DS, Crowe JJ, Sinex DG, and McMullen NT (2002) Cell types and response properties of neurons in the ventral division of the medial geniculate body of the rabbit. *Journal of Comparative Neurology* **445**:78–96.
- Clarey JC, Barone P, and Imig TJ (1992) Physiology of thalamus and cortex. In Fay RR and Popper AN (eds). *Springer Handbook of Auditory Research, Volume 2: The Mammalian Auditory Pathway: Neurophysiology*. Springer-Verlag, New York, pp. 232–334.
- Condon CJ, White KR, and Feng AS (1994) Processing of amplitude-modulated signals that mimic echoes from fluttering targets in the inferior colliculus of the little brown bat, *Myotis lucifugus*. *Journal of Neurophysiology* **71**:768–784.
- Condon CJ, Galazyuk A, and Feng AS (1997) Neurons in the auditory cortex of the little brown bat exhibit selectivity for complex-amplitude modulated signals that mimic echoes from fluttering insects. *Auditory Neuroscience* **3**:269–287.
- De Biasi S and Rustioni A (1990) Ultrastructural immunocytochemical localization of excitatory amino acids in the somatosensory system. *Journal of Histochemistry and Cytochemistry* **38**:1745–1754.
- Delgutte B, Joris PX, Litovsky RY, and Yin TCT (1999) Receptive fields and binaural interactions for virtual-space stimuli in the cat inferior colliculus. *Journal of Neurophysiology* **81**:2833–2851.
- de Ribaupierre F (1997) Acoustical information processing in the auditory thalamus and cerebral cortex. In: Ehret G and Romand R (eds). *The Central Auditory System*. Oxford University Press, New York, pp. 317–397.
- Edeline J-M, Manunta Y, Nodal FR, and Bajo VM (1999) Do auditory responses recorded from awake animals reflect the anatomical parcellation of the auditory thalamus? *Hearing Research* **131**:135–152.
- Edeline JM, Manunta Y, and Hennevin E (2000) Auditory thalamus neurons during sleep: changes in frequency selectivity, threshold, and receptive field size. *Journal of Neurophysiology* **84**:934–952.
- Ehret G and Moffat AJM (1985) Inferior colliculus of the house mouse. II. Single unit responses to tones, noise and tone-noise combinations as a function of sound intensity. *Journal of Comparative Physiology A* **156**:619–635.

- Eysel UT (1976) Quantitative studies of intracellular postsynaptic potentials in the lateral geniculate nucleus of the cat with respect to optic tract stimulus response latencies. *Experimental Brain Research* **25**:469–486.
- Fitzpatrick DC, Olsen JF, and Suga N (1998) Connections among functional areas in the mustached bat auditory cortex. *Journal of Comparative Neurology* **391**:366–396.
- Frisina RD, O'Neill WE, and Zettel ML (1989) Functional organization of mustached bat inferior colliculus: II. Connections of the FM<sub>2</sub> region. *Journal of Comparative Neurology* **284**:85–107.
- Frisina RD, Walton JP, Lynch-Armour MA, and Klotz DA (1997) Efferent projections of a physiologically characterized region of the inferior colliculus of the young adult CBA mouse. *Journal of the Acoustical Society of America* **101**:2741–2753.
- Fuzessery ZM and Feng AS (1983) Mating call selectivity in the thalamus and midbrain of the leopard frog (*Rana p. pipiens*): single and multiunit analyses. *Journal of Comparative Physiology A* **150**:333–344.
- Giraud A-L, Lorenzi C, Ashburner J, Wable J, Johnsrude I, Frackowiak R, and Kleinschmidt A (2000) Representation of the temporal envelope of sounds in the human brain. *Journal of Neurophysiology* **84**:1588–1598.
- González-Hernández TH, Galindo-Mireles D, Castaneyra-Perdomo A, and Ferres-Torres R (1991) Divergent projections of projecting neurons of the inferior colliculus to the medial geniculate body and the contralateral inferior colliculus in the rat. *Hearing Research* **52**:17–21.
- Gordon M and O'Neill WE (2000) An extralemniscal component of the mustached bat inferior colliculus selective for direction and rate of linear frequency modulations. *Journal of Comparative Neurology* **426**:165–181.
- Gross NB, Lifschitz WS, and Anderson DJ (1974) The tonotopic organization of the auditory thalamus of the squirrel monkey (*Saimiri sciureus*). *Brain Research* **65**:323–332.
- Harms MP and Melcher JR (2002) Sound repetition rate in the human auditory pathway: representations in the waveshape and amplitude of fMRI activation. *Journal of Neurophysiology* **88**:1433–1450.
- He J (2001) ON and OFF pathways segregated at the auditory thalamus of the guinea pig. *Journal of Neuroscience* **21**:8672–8679.
- He J and Hu B (2002) Differential distribution of burst and single-spike responses in auditory thalamus. *Journal of Neurophysiology* **88**:2152–2156.
- Henkel CK (1983) Evidence of sub-collicular auditory projections to the medial geniculate nucleus in the cat: an autoradiographic and horseradish peroxidase study. *Brain Research* **259**:21–30.
- Hu B (1995) Cellular basis of temporal synaptic signalling: an in vitro electrophysiological study in rat auditory thalamus. *Journal of Physiology (London)* **483**:167–182.
- Hu B, Senatorov V, and Mooney D (1994) Lemniscal and non-lemniscal synaptic transmission in rat auditory thalamus. *Journal of Physiology (London)* **479**:217–231.
- Hutson KA, Glendenning KK, and Masterton RB (1991) Acoustic chiasm IV: eight mid-brain decussations of the auditory system in the cat. *Journal of Comparative Neurology* **312**:105–131.
- Imig TJ and Morel A (1985) Tonotopic organization in ventral nucleus of medial geniculate body in the cat. *Journal of Neurophysiology* **53**:309–340.
- Imig TJ, Poirier P, Irons WA, and Samson FK (1997) Monaural spectral contrast mechanism for neural sensitivity to sound direction in the medial geniculate body of the cat. *Journal of Neurophysiology* **78**:2754–2771.

- Irvine DRF and Gago G (1990) Binaural interaction in high-frequency neurons in inferior colliculus of the cat: effects of variations in sound pressure level on sensitivity to interaural intensity differences. *Journal of Neurophysiology* **63**:570–591.
- Ivarsson C, de Ribaupierre Y, and de Ribaupierre F (1988) Influence of auditory localization cues on neuronal activity in the auditory thalamus of the cat. *Journal of Neurophysiology* **59**:586–606.
- Jahnsen H and Llinás R (1984) Electrophysiological properties of guinea-pig thalamic neurones: an *in vitro* study. *Journal of Physiology (London)* **349**:205–226.
- Jones EG and Rockel AJ (1971) The synaptic organization in the medial geniculate body of afferent fibres ascending from the inferior colliculus. *Zeitschrift für Zellforschung und Mikroskopische Anatomie* **113**:44–66.
- Klug A, Khan A, Burger RM, Bauer EE, Hurley LM, Yang L, Grothe B, Halvorsen MB, and Park TJ (2000) Latency as a function of intensity in auditory neurons: influences of central processing. *Hearing Research* **148**:107–123.
- Kudo M and Niimi K (1978) Ascending projections of the inferior colliculus onto the medial geniculate body in the cat studied by anterograde and retrograde tracing techniques. *Brain Research* **155**:113–117.
- Kudo M, Itoh K, Kawamura S, and Mizuno N (1983) Direct projections to the pretectum and the midbrain reticular formation from auditory relay nuclei in the lower brainstem of the cat. *Brain Research* **288**:13–19.
- Kuwada S and Yin TCT (1983) Binaural interaction in low-frequency neurons in inferior colliculus of the cat. I. Effects of long interaural delays, intensity, and repetition rate on interaural delay function. *Journal of Neurophysiology* **50**:981–999.
- Langner G (1992) Periodicity coding in the auditory system. *Hearing Research* **60**:115–142.
- LeDoux JE, Ruggiero DA, and Reis DJ (1985) Projections to the subcortical forebrain from anatomically defined regions of the medial geniculate body in the rat. *Journal of Comparative Neurology* **242**:182–213.
- LeDoux JE, Ruggiero DA, Forest R, Stornetta R, and Reis DJ (1987) Topographic organization of convergent projections to the thalamus from the inferior colliculus and spinal cord in the rat. *Journal of Comparative Neurology* **264**:123–146.
- Lennartz RC and Weinberger NM (1992) Frequency selectivity is related to temporal processing in parallel thalamocortical auditory pathways. *Brain Research* **583**:81–92.
- Leroy SA and Wenstrup JJ (2000) Spectral integration in the inferior colliculus of the mustached bat. *Journal of Neuroscience* **20**:8533–8541.
- Linke R (1999) Differential projection patterns of superior and inferior collicular neurons onto posterior paralaminar nuclei of the thalamus surrounding the medial geniculate body in the rat. *European Journal of Neuroscience* **11**:187–203.
- Llano DA and Feng AS (1999) Response characteristics of neurons in the medial geniculate body of the little brown bat to simple and temporally-patterned sounds. *Journal of Comparative Physiology A* **184**:371–385.
- Majorossy K and Kiss A (1976) Types of interneurons and their participation in the neuronal network of the medial geniculate body. *Experimental Brain Research* **26**:19–37.
- Majorossy K and Réthelyi M (1968) Synaptic architecture in the medial geniculate body (ventral division). *Experimental Brain Research* **6**:306–323.
- Malmierca MS, Rees A, and Le Beau FEN (1997) Ascending projections to the medial geniculate body from physiologically identified loci in the inferior colliculus. In: Syka

- J (ed). *Acoustical Signal Processing in the Central Auditory System*. Plenum, New York, pp. 295–302.
- Malmierca MS, Merchán MA, Henkel CK, and Oliver DL (2002) Direct projections from cochlear nuclear complex to auditory thalamus in the rat. *Journal of Neuroscience* **22**: 10891–10897.
- Margoliash D and Fortune ES (1992) Temporal and harmonic combination-sensitive neurons in the zebra finch's HVc. *Journal of Neuroscience* **12**:4309–4326.
- Massaux A and Edeline J-M (2003) Bursts in the medial geniculate body: a comparison between anesthetized and unanesthetized states in guinea pig. *Experimental Brain Research* **153**:573–578.
- McCormick DA and Feeser HR (1990) Functional implications of burst firing and single spike activity in lateral geniculate relay neurons. *Neuroscience* **39**:103–113.
- Middlebrooks JC and Zook JM (1983) Intrinsic organization of the cat's medial geniculate body identified by projections to binaural response-specific bands in the primary auditory cortex. *Journal of Neuroscience* **3**:203–224.
- Mishima K (1992) Facilitatory and inhibitory processes in the thalamic ventrobasal nucleus of the rat. *Japanese Journal of Physiology* **42**:193–210.
- Mittmann DH and Wenstrup JJ (1995) Combination-sensitive neurons in the inferior colliculus. *Hearing Research* **90**:185–191.
- Mooney DM, Hu B, and Senatorov VV (1995) Muscarine induces an anomalous inhibition of synaptic transmission in rat auditory thalamic neurons in vitro. *Journal of Pharmacology and Experimental Therapeutics* **275**:838–844.
- Morel A, Rouiller E, de Ribaupierre Y, and de Ribaupierre F (1987) Tonotopic organization in the medial geniculate body (MGB) of lightly anesthetized cats. *Experimental Brain Research* **69**:24–42.
- Morest DK (1964) The neuronal architecture of the medial geniculate body of the cat. *Journal of Anatomy (London)* **98**:611–630.
- Morest DK (1965) The laminar structure of the medial geniculate body of the cat. *Journal of Anatomy (London)* **99**:143–160.
- Morest DK (1975) Synaptic relationships of Golgi type II cells in the medial geniculate body of the cat. *Journal of Comparative Neurology* **162**:157–194.
- Nelson PG and Erulkar SD (1963) Synaptic mechanisms of excitation and inhibition in the central auditory pathway. *Journal of Neurophysiology* **26**:908–923.
- Ohlemiller KK, Kanwal JS, and Suga N (1996) Facilitative responses to species-specific calls in cortical FM-FM neurons of the mustached bat. *NeuroReport* **7**:1749–1755.
- Oliver DL (1984) Neuron types in the central nucleus of the inferior colliculus that project to the medial geniculate body. *Neuroscience* **11**:409–424.
- Oliver DL and Hall WC (1978) The medial geniculate body of the tree shrew (*Tupaia glis*). I. Cytoarchitecture and midbrain connections. *Journal of Comparative Neurology* **182**:423–458.
- Oliver DL, Beckius GE, and Ostapoff E-M (1994) Connectivity of neurons in identified auditory circuits studied with transport of dextran and microspheres plus intracellular injection of lucifer yellow. *Journal of Neuroscience Methods* **53**:23–27.
- Oliver DL, Ostapoff E-M, and Beckius GE (1999) Direct innervation of identified tectothalamic neurons in the inferior colliculus by axons from the cochlear nucleus. *Neuroscience* **93**:643–658.
- Olsen JF and Suga N (1991a) Combination-sensitive neurons in the medial geniculate body of the mustached bat: encoding of relative velocity information. *Journal of Neurophysiology* **65**:1254–1274.

- Olsen JF and Suga N (1991b) Combination-sensitive neurons in the medial geniculate body of the mustached bat: encoding of target range information. *Journal of Neurophysiology* **65**:1275–1296.
- O'Neill WE and Brimijoin WO (2002) Directional selectivity for FM sweeps in the suprageniculate nucleus of the mustached bat medial geniculate body. *Journal of Neurophysiology* **88**:172–187.
- O'Neill WE and Suga N (1982) Encoding of target range and its representation in the auditory cortex of the mustached bat. *Journal of Neuroscience* **2**:17–31.
- O'Neill WE, Frisina RD, and Gooler DM (1989) Functional organization of mustached bat inferior colliculus: I. Representation of FM frequency bands important for target ranging revealed by 14C-2-deoxyglucose autoradiography and single unit mapping. *Journal of Comparative Neurology* **284**:60–84.
- Pallas SL and Sur M (1994) Morphology of retinal axon arbors induced to arborize in a novel target, the medial geniculate nucleus. II. Comparison with axons from the inferior colliculus. *Journal of Comparative Neurology* **349**:363–376.
- Peruzzi D, Bartlett E, Smith PH, and Oliver DL (1997) A monosynaptic GABAergic input from the inferior colliculus to the medial geniculate body in rat. *Journal of Neuroscience* **17**:3766–3777.
- Phillips DP and Irvine DRF (1979) Acoustic input to single neurons in pulvinar-posterior complex of cat thalamus. *Journal of Neurophysiology* **42**:123–136.
- Poirier P, Samson FK, and Imig TJ (2003) Spectral shape sensitivity contributes to the azimuth tuning of neurons in the cat's inferior colliculus. *Journal of Neurophysiology* **89**:2760–2777.
- Portfors CV and Wenstrup JJ (1999) Delay-tuned neurons in the inferior colliculus of the mustached bat: implications for analyses of target distance. *Journal of Neurophysiology* **82**:1326–1338.
- Portfors CV and Wenstrup JJ (2001a) Topographical distribution of delay-tuned responses in the mustached bat inferior colliculus. *Hearing Research* **151**:95–105.
- Portfors CV and Wenstrup JJ (2001b) Responses to combinations of tones in the nuclei of the lateral lemniscus. *Journal of the Association for Research in Otolaryngology* **2**:104–117.
- Pow DV, Crook DK, and Wong ROL (1994) Early appearance and transient expression of putative amino acid neurotransmitters and related molecules in the developing rabbit retina: an immunocytochemical study. *Visual Neuroscience* **11**:1115–1134.
- Rauschecker JP, Tian B, and Hauser M (1995) Processing of complex sounds in the macaque nonprimary auditory cortex. *Science* **268**:111–114.
- Redies H and Brandner S (1991) Functional organization of the auditory thalamus in the guinea pig. *Experimental Brain Research* **86**:384–392.
- Rodrigues-Dageaff C, Simm G, de Ribaupierre Y, Villa A, de Ribaupierre F, and Rouiller EM (1989) Functional organization of the ventral division of the medial geniculate body of the cat: evidence for a rostro-caudal gradient of response properties and cortical projections. *Hearing Research* **39**:103–126.
- Rouiller EM and de Ribaupierre F (1985) Origin of afferents to physiologically defined regions of the medial geniculate body of the cat: ventral and dorsal divisions. *Hearing Research* **19**:97–114.
- Rouiller EM, de Ribaupierre Y, Toros-Morel A, and de Ribaupierre F (1981) Neural coding of repetitive clicks in the medial geniculate body of cat. *Hearing Research* **5**:81–100.
- Rouiller EM, Rodrigues-Dageaff C, Simm G, de Ribaupierre Y, Villa A, and de Ribaupierre F (1989) Functional organization of the medial division of the medial geniculate



- body of the cat: tonotopic organization, spatial distribution of response properties and cortical connections. *Hearing Research* **39**:127–142.
- Saint Marie RL, Stanforth DA, and Jubelier EM (1997) Substrate for rapid feedforward inhibition of the auditory forebrain. *Brain Research* **765**:173–176.
- Saldaña E and Merchán MA (1992) Intrinsic and commissural connections of the rat inferior colliculus. *Journal of Comparative Neurology* **19**:417–437.
- Samson FK, Clarey JC, Barone P, and Imig TJ (1993) Effects of ear plugging on single-unit azimuth sensitivity in cat primary auditory cortex. I. Evidence for monaural directional cues. *Journal of Neurophysiology* **70**:492–511.
- Samson FK, Barone P, Clarey JC, and Imig TJ (1994) Effects of ear plugging on single-unit azimuth sensitivity in cat primary auditory cortex. II. Azimuth tuning dependent upon binaural stimulation. *Journal of Neurophysiology* **71**:2194–2216.
- Samson FK, Barone P, Irons WA, Clarey JC, Poirier P, and Imig TJ (2000) Directionality derived from differential sensitivity to monaural and binaural cues in the cat's medial geniculate body. *Journal of Neurophysiology* **84**:1330–1345.
- Semple MN and Kitzes LM (1985) Single-unit responses in the inferior colliculus: different consequences of contralateral and ipsilateral auditory stimulation. *Journal of Neurophysiology* **53**:1467–1482.
- Sherman SM (2001) Tonic and burst firing: dual modes of thalamocortical relay. *Trends in Neurosciences* **24**:122–126.
- Steriade M and Llinás RR (1988) The functional states of the thalamus and the associated neuronal interplay. *Physiological Reviews* **68**:649–742.
- Steriade M, Datta S, Paré D, Oakson G, and Curró Dossi RC (1990) Neuronal activities in brain-stem cholinergic nuclei related to tonic activation processes in thalamocortical systems. *Journal of Neuroscience* **10**:2541–2559.
- Suga N, O'Neill WE, and Manabe T (1978) Cortical neurons sensitive to combinations of information-bearing elements of biosonar signals in the mustache bat. *Science* **200**:778–781.
- Suga N, O'Neill WE, Kujirai K, and Manabe T (1983) Specificity of combination-sensitive neurons for processing of complex biosonar signals in auditory cortex of the mustached bat. *Journal of Neurophysiology* **49**:1573–1626.
- Suga N, Zhang Y, and Yan J (1997) Sharpening of frequency tuning by inhibition in the thalamic auditory nucleus of the mustached bat. *Journal of Neurophysiology* **77**:2098–2114.
- Syka J, Popelář J, Kvašňák E, and Astl J (2000) Response properties of neurons in the central nucleus and external and dorsal cortices of the inferior colliculus in guinea pig. *Experimental Brain Research* **133**:254–266.
- Tennigkeit F, Schwarz DWF, and Puil E (1996) Mechanisms for signal transformation in lemniscal auditory thalamus. *Journal of Neurophysiology* **76**:3597–3608.
- Tennigkeit F, Puil E, and Schwarz DWF (1997) Firing modes and membrane properties in lemniscal auditory thalamus. *Acta Otolaryngologica* **117**:254–257.
- Tennigkeit F, Schwarz DWF, and Puil E (1998) GABA<sub>B</sub> receptor activation changes membrane and filter properties of auditory thalamic neurons. *Hearing Research* **122**:18–24.
- Tennigkeit F, Schwarz DWF, and Puil E (1999) Effects of metabotropic glutamate receptor activation in auditory thalamus. *Journal of Neurophysiology* **82**:718–729.
- Vater M, Kössl M, and Horn AKE (1992) GAD- and GABA-immunoreactivity in the ascending auditory pathway of horseshoe and mustached bats. *Journal of Comparative Neurology* **325**:183–206.
- Webber TJ, Green EJ, Winters RW, Schneiderman N, and McCabe PM (1999) Contri-

- bution of NMDA and non-NMDA receptors to synaptic transmission from the brachium of the inferior colliculus to the medial subdivision of the medial geniculate nucleus in the rabbit. *Experimental Brain Research* **124**:295–303.
- Wenstrup JJ (1995) The auditory thalamus in bats. In: Popper AN and Fay RR (eds). *Springer Handbook of Auditory Research*, Volume 4: *Hearing by Bats*. Springer-Verlag, New York, pp. 368–415.
- Wenstrup JJ (1999) Frequency organization and responses to complex sounds in the medial geniculate body of the mustached bat. *Journal of Neurophysiology* **82**:2528–2544.
- Wenstrup JJ and Grose CD (1995) Inputs to combination-sensitive neurons in the medial geniculate body of the mustached bat: the missing fundamental. *Journal of Neuroscience* **15**:4693–4711.
- Wenstrup JJ and Leroy SA (2001) Spectral integration in the inferior colliculus: role of glycinergic inhibition in response facilitation. *Journal of Neuroscience* **21** RC124:1–6.
- Wenstrup JJ, Larue DT, and Winer JA (1994) Projections of physiologically defined subdivisions of the inferior colliculus in the mustached bat: targets in the medial geniculate body and extrathalamic nuclei. *Journal of Comparative Neurology* **346**:207–236.
- Wepsic JG (1966) Multimodal sensory activation of cells in the magnocellular medial geniculate nucleus. *Experimental Neurology* **15**:299–318.
- Winer JA (1992) The functional architecture of the medial geniculate body and the primary auditory cortex. In: Webster DB, Popper AN, and Fay RR (eds). *Springer Handbook of Auditory Research*, Volume 1: *The Mammalian Auditory Pathway: Neuroanatomy*, Springer-Verlag, New York, pp. 222–409.
- Winer JA and Larue DT (1988) Anatomy of glutamic acid decarboxylase immunoreactive neurons and axons in the rat medial geniculate body. *Journal of Comparative Neurology* **278**:47–68.
- Winer JA and Larue DT (1996) Evolution of GABAergic circuitry in the mammalian medial geniculate body. *Proceedings of the National Academy of Sciences of the United States of America* **93**:3083–3087.
- Winer JA and Wenstrup JJ (1994a) Cytoarchitecture of the medial geniculate body in the mustached bat (*Pteronotus parnellii*). *Journal of Comparative Neurology* **346**:161–182.
- Winer JA and Wenstrup JJ (1994b) The neurons of the medial geniculate body in the mustached bat (*Pteronotus parnellii*). *Journal of Comparative Neurology* **346**:183–206.
- Winer JA, Larue DT, and Huang CL (1999) Two systems of giant axon terminals in the cat medial geniculate body: convergence of cortical and GABAergic inputs. *Journal of Comparative Neurology* **413**:181–197.
- Winer JA, Wenstrup JJ, and Larue DT (1992) Patterns of GABAergic immunoreactivity define subdivisions of the mustached bat's medial geniculate body. *Journal of Comparative Neurology* **319**:172–190.
- Winer JA, Saint Marie RL, Larue DT, and Oliver DL (1996) GABAergic feedforward projections from the inferior colliculus to the medial geniculate body. *Proceedings of the National Academy of Sciences of the United States of America* **93**:8005–8010.
- Yan J and Suga N (1996) The midbrain creates and the thalamus sharpens echo-delay tuning for the cortical representation of target-distance information in the mustached bat. *Hearing Research* **93**:102–110.
- Zook JM, Winer JA, Pollak GD, and Bodenhamer RD (1985) Topology of the central nucleus of the mustache bat's inferior colliculus: correlation of single unit response properties and neuronal architecture. *Journal of Comparative Neurology* **231**:530–546.

# Chapter 8

## Three Systems of Descending Projections to the Inferior Colliculus

JEFFERY A. WINER

### 1. INTRODUCTION

The inferior colliculus (IC) is the first site in which the several parallel streams established in the cochlear nucleus (Warr 1982) converge (Aitkin and Phillips 1984) before sending this information to the auditory thalamus (see Chapter 7) and cortex (Winer 1992). There is parallel convergence of ascending inputs from  $\gamma$ -aminobutyric acid-positive (GABAergic) and glycinergic subcollicular pathways that reach the IC (Winer et al. 1995) and terminate there (Saint Marie and Baker 1990) or GABAergic inputs that reach the medial geniculate body (MGB) (Winer et al. 1996). The emphasis on the IC as the midbrain hub for ascending input has obscured the fact that it is the target of descending forebrain projections from three systems. The first arises in 11 of the 12 areas that constitute the cat auditory cortex (Winer et al. 1998) and is the corticocollicular system (CC); its physiological role is not well understood despite the fact that the cortex can affect many facets of IC operations. A second, smaller system also has widespread origins but represents far fewer neurons: the thalamotectal system (TT) arises from a widely distributed and diverse set of phylogenetically ancient nuclei in the posterior and intralaminar thalamic regions (Kuwabara and Zook 2000). Nothing is known about the physiological impact of the TT system. A third input arises from the bat lateral amygdaloid nucleus (Marsh et al. 2002), which is a multisensory limbic center and itself the target of auditory and somatic sensory cortical projections in rat (Romanski et al. 1993) and of ascending input from two MGB subdivisions (Shinonaga et al. 1994). No physiologic evidence is available on the role of the amygdalocollicular system (AC). It is not implausible to suggest that the CC, TT, and AC systems collectively represent parallel descending influences that converge on IC subdivisions largely outside the lemniscal pathway represented in the central nucleus. Ultimately, the cortical output targets many of the same subcortical nuclei that provide input to the cortex (Winer et al. 2001), and to other more remote nuclei early in the ascending chain of serial processing in rat (Doucet et al. 2002).

A reconsideration of descending input to the IC is timely, as the CC and TT pathways are rarely compared. The CC input merits attention because it arises

from so many subdivisions of auditory cortex and is thus massive, while the TT system is widespread, diffuse, and far smaller. This CC input to some IC subdivisions is larger than the ascending auditory input, suggesting a role other than modulatory. Physiologic studies find powerful and diverse cortical influences on IC neural responses. Because the posterior thalamus targets many of the same IC subdivisions as does the auditory cortex input, there may be local interactions between corticofugal and thalamofugal systems. The auditory corticofugal projection is comparable in size to the corticospinal pathway; perhaps it has subcortical effects analogous to those in the motor system, except within an auditory frame of reference. The TT pathway remains to be integrated within the family of descending systems. Finally, the AC pathway implies a new kind of functional centrality for the IC as inputs from the cortex, posterior thalamic nuclei, and limbic forebrain, as well as classic ascending projections, all converge in the midbrain subdivisions. Unless noted otherwise, citations refer to the cat.

## 2. ORIGINS OF THE CORTICOLICULAR SYSTEM

At least 11 cortical areas with some auditory affiliation project onto the IC. This input varies on an areal basis (Fig. 8.4A). The strongest CC projection is from the primary (tonotopic) subdivisions of auditory cortex, although nontopographic regions have more targets in the midbrain and a wider range in projection strength. Limbic-affiliated cortical areas, for example, differ in their CC projections, suggesting that otherwise comparable regions have distinctive corticofugal patterns (Winer et al. 1998).

Layer V is the only laminar source of CC input from the primary auditory cortex (AI). Layer V in AI has three subdivisions, and the CC system arises in sublayers Vb and Vc (Winer and Prieto 2001). The most superficial and the deepest parts of layer V give rise to corticogeniculate axons (Prieto and Winer 1999). Layer V thus projects in the corticocollicular and the corticothalamic systems, besides participating in the commissural (Code and Winer 1985), corticopontine (Brodal 1972), and corticostriatal (Reale and Imig 1983; Beneyto and Prieto 2001) pathways. Thus, despite the laminar specificity of layer V projections, the CC system is only one of its several outputs.

### 2.1. TYPES OF NEURONS

All CC neurons are pyramidal in rat (Games and Winer 1988), cat (Winer and Prieto 2001), and monkey (Winer et al. 2002). CC pyramidal cells range from medium-sized to the largest in AI, and CC axons may have corresponding differences in form, conduction velocity and, perhaps their postsynaptic targets (cf. Section 2.3). These neurons are glutamatergic (Feliciano and Potashner 1995). Many large pyramidal neurons have apical dendrites that extend to layer I to receive both specific (layers III to IV) and nonspecific (layer I) thalamocortical input (White and Hersch 1982; Huang and Winer 2000). Such cells can fire in




Figure 8.1. Corticofugal projections to the IC (A–D; F–I) and MGB (E, J) from deposits in AI (black area in A) and temporal (black area in F) cortices. Both projections are topographically specific and produce focal terminal labeling in different IC subdivisions. MGB-labeled neurons (J, open circles) mingle with anterogradely labeled axon terminals. (Winer et al. 1998, 2001.)

an intrinsic bursting mode and have a sparser local projection in AI, whereas those with shorter apical dendrites have more regular spiking behavior and intricate local projections in rat (Hefti and Smith 2000).

## 2.2. TOPOGRAPHY

The distribution of CC axon terminals in the IC is topographic, irrespective of the cortical origin of the projection (Winer et al. 1998). Because most auditory cortical areas have no tonotopic arrangement (Imig and Reale 1980; Schreiner and Cynader 1984), the projections from a limbic-related area (Fig. 8.1F–I) were as orderly and focal as those from tonotopic areas (Fig. 8.1A–D). Perhaps organizing principles other than tonotopy regulate the CC projection topography, and nuclear specificity is a general quality of corticofugal projections.

A second feature is that the CC system targets IC subdivisions that themselves receive modest input from subcollicular auditory structures (Brunso-Bechtold et al. 1981). The wealth of brain stem input to the IC (Warr 1982) is thus partly segregated from cortical projections. This is not the case in the MGB (Calford and Aitkin 1983; Winer et al. 2001).

## 2.3. CORTICOCOLLICULAR AXONS

CC axons vary in size and shape and may have a distribution specific to each target. For example, fibers targeting the lateral nucleus are much thicker and may form elaborate terminal nests (Fig. 8.2A: 7). In contrast, the input to the central nucleus is unique because it is conspicuously lighter (Fig. 8.2C: 4), because the axons are so slender ( $<1 \mu\text{m}$  in diameter; Fig. 8.2B: 1) and have relatively few boutons, and because many fibers parallel the axis of central nucleus laminae (Fig. 8.2B: 7). The density of input and the distribution of the axons corroborates results found with other tracers (Fig. 8.1A–D, F–I). Perhaps the cortical neurons projecting to separate midbrain subdivisions and their terminal architecture is specific to each nuclear target. Further subtypes of layer V neurons may exist that are analogous to the subclasses of layer VI pyramidal neurons in visual cortex (Briggs and Callaway 2001). Combined, these characteristics support the idea of multiple parallel descending pathways.

## 3. CORTICAL EFFECTS ON MIDBRAIN PHYSIOLOGY

About one half of a sample of rat IC cells responded to electrical pulses delivered to the cortex; most had brief excitatory responses, and in some this preceded a longer lasting inhibition. Inhibition suppressed spontaneous or sound-evoked discharge (Syka and Popelář 1984). AI stimulation evoked excitatory postsynaptic potentials, inhibitory postsynaptic potentials, or a sequence of both in IC neurons projecting to the MGB; the inhibitory potentials had a longer latency (Mitani et al. 1983). Perhaps colliculogeniculate neurons are un-




Figure 8.2. Corticofugal axons in IC subdivisions (C: 3,4) after deposits of biotinylated dextran amines in AI (C: 1) or AI and AAF (C: 2). (A) Axons in the lateral nucleus. Four sections were required for this panel (C: 3, small squares). 1, Vertically oriented fibers; 2, thick preterminal axon; 3, clustered boutons; 4, a massive terminal; 5, fibers cross the plexus at right angles; 6, medium-sized (1–2  $\mu\text{m}$  thick) fibers; 7, terminals forming complex nests; 8, fibers with a corkscrew trajectory. (B) Central nucleus axons were sparser, thinner, and had fewer boutons than fibers in other subdivisions. Seven sections were required for this panel (C: 4, small squares). 1, The main type is a fine fiber (<1  $\mu\text{m}$  thick) with sparse beads, and oriented parallel to the fibrodendritic laminae; 2, thicker preterminal branch; 3, elaborate terminal appendages; 4, grape-like boutons; 5, the delicate terminal plexus of a thin fiber (cf. 1); 6, extremely fine fibers about 0.5  $\mu\text{m}$  thick; 7, laminar orientation of fibers; 8, more terminal boutons of fine axons (cf. 5, 6); 9, corkscrews in the neuropil. Protocol for (A, B): planapochromat, numerical aperture 1.32,  $\times 2000$ . (C) Deposit sites (black) in AI (1) and AAF/AI (2). 3, 4, Locus of IC axons in the lateral nucleus (LN) and central nucleus (CN).

der direct cortical control, and the cortex can modulate thalamic neurons through three routes: directly via corticogeniculate projections and indirectly, through the IC and the thalamic reticular nucleus (Crabtree 1998).

Echolocating bats have been the subject of physiologic work on the CC system (Suga et al. 2000). Enhancement of local cortical activity by electrical stimulation in the big brown bat decreased collicular rate-intensity functions and the size of the spatial response area, increased thresholds, and sharpened tuning. Suppression of local cortical activity with lidocaine elevated rate-intensity functions and increased the size of the response area, lowered threshold, and broadened the frequency tuning curve (Sun et al. 1996). Iontophoretic studies using GABA or bicuculline in the IC elicited many of these effects by local pharmacologic administration (Jen et al. 1998). Perhaps cortical axons preferentially target local IC inhibitory circuits responsible for these processes. In the mustached bat, echo-delay processing is enhanced by CC stimulation (Yan and Suga 1996a), while MGB neurons sharpen the delay-tuning curves significantly (Yan and Suga 1996b). Frequency specific shifts in IC frequency reorganization occur in mice after cortical stimulation (Yan and Ehret 2001), and in bats cortically evoked shifts in IC organization emerge before those in auditory cortex (Gao and Suga 2000). Overall, both global and specific changes in receptive field properties and physiologic organization are under CC control.

## 4. MODELS OF CORTICOFUGAL FUNCTIONAL ORGANIZATION

The corticofugal projections to the auditory midbrain and thalamus might exceed those ascending from the brain stem. This raises two questions: are these descending pathways parallel and as specific functionally as their ascending counterparts, and how do they differ in species in which the corticofugal system is reduced?

### 4.1. FEEDBACK

The classic view of corticothalamic projections emphasizes their feedback role (Frigyesi et al. 1972). Although this is a reasonable hypothesis it does not explain why these projections are necessary or reciprocal or how they interact (Deschênes et al. 1998), nor can it explain why some axons have larger zones of influence than the topographically matched retinal input (Murphy et al. 1999). With respect to the CC system, the feedback hypothesis is even less compelling because the cortex cannot regulate temporal processes that are briefer than the CC conduction time (cf. Section 4.2), and because many IC neurons that the cortex might be expected to influence, for example, those in the central nucleus, likely receive more extensive intracollicular than cortical input, at least in rodents (Saldaña and Merchán 1992; Malmierca et al. 1995).



#### 4.2. *STEPS TOWARD A THEORY OF DESCENDING CONTROL*

There is no global theory of corticofugal operations that places corticothalamic, CC, and other corticofugal projections in a unified context. Such a theory is vital because the targets of this system include the striatum and the pons and other centers that do not have the same functions. Moreover, the corticothalamic and the corticocollicular projections must have different roles because the auditory thalamus and midbrain are connected bidirectionally. A global view should relate corticofugal to corticocortical connections because these systems interact. A model is proposed that postulates a predictive role in which CC input provides instructive commands for processes such as dynamic retuning of receptive fields, postural adjustments, preparations for head and neck reflexes concerned with spatial localization, and the integration of descending auditory and ascending somatic sensory information in the IC. Alternative hypotheses emphasize descending control (Przybylski 1998) or signal selection (King 1997). The present account differs in its emphasis on the anticipatory and predictive aspects of corticofugal control, and it is consistent with the notion that feedback responses are temporally rapid in other neural systems and may influence events earlier in the chain of local processing (Hupé et al. 2001).

Several ideas flow from this model. To be predictive, the outflow must be temporally remote from immediate sensory events; this is the case with the time course of evoked potentials. Second, there should be, and there are, significant differences in the corticothalamic and CC projections; each has unique laminar origins and axonal structure, and they differ in their strength of input to the main targets, namely, the ventral division of the MGB (the input to which is massive) and the central nucleus of the IC (whose input is far smaller). Much cortical input to the central nucleus may synapse first in the lateral nucleus (or external cortex), perhaps influencing multisensory operations that are poorly understood. A third requirement is that the cortical areas with descending projections should have extensive feedforward inputs to the higher order areas, which they do. This enables cortex to craft strategic responses long after the initial sensory events, and far in advance of actual movement, much as the premotor cortex enables movement sequence preparation. This theory must account for auditory cortex projections to the basal ganglia, which might coordinate musculoskeletal and audiospinal frames of reference for complex behaviors such as prey acquisition, predator avoidance, mating, freezing, startle, and motor aspects of vocalization dependent on sensory feedback. A fifth prediction is that dynamic midbrain processes such as vigilance or state-dependent changes might require neurons capable of firing in the bursting mode; such neurons are abundant in layer V of AI. These requirements cannot confirm the validity of predictive autonomy as a principle of midbrain function, but they do provide a context for further exploration.

## 5. THE THALAMOTECTAL SYSTEM

The discovery of the thalamotectal (TT) system was fortuitous and unexpected. It was fortuitous because labeled neurons were found in the auditory thalamus after IC tracer deposits designed to label ascending midbrain axons (Adams 1979) and unexpected because there are no analogous descending projections in other modalities.

Elegant work in a slice preparation first revealed TT organization by filling cells in the medial or dorsal divisions of the gerbil MGB and tracing their axons to the IC (Kuwabara and Zook 2000). Some of the strongest projections were from the medial division to the external nucleus (50% of 14 intracellularly filled neurons), and descending axons were traced to regions near the superior olivary complex.

The TT system arises from the MGB, posterior thalamus, and rostral midbrain, and it targets the central nucleus, dorsal cortex, and lateral nucleus (external cortex) of the IC in rat, cat, and monkey (Winer et al. 2002). Ipsilateral origins are the medial division of the MGB, central gray, and substantia nigra. Bilateral sources include the peripeduncular nuclei, the posterior intralaminar thalamic nuclei, the nucleus of the brachium, the lateral tegmentum, and the deep superior colliculus. TT neurons are diverse in form (Fig. 8.3), often having long, smooth, thin dendrites that branch rarely and resemble thalamic intralaminar cells (Winer et al. 1988). Some TT cells also receive input from the IC, suggesting a reciprocal relationship between ascending and descending systems.

### 5.1. SPECULATION ON THE THALAMOTECTAL SYSTEM

The role of the TT system is unknown. Thus, functional hypotheses must largely be based on the properties of the projection neurons and their IC targets. A few examples demonstrate the prospective breadth of these relationships (Casseday and Covey 1996).

Neurons in the medial division of the MGB have broad tuning curves, respond both to auditory and nonauditory input (Aitkin 1973), show long-term potentiation (Gerren and Weinberger 1983), project to many layers in all subdivisions of auditory cortex (Huang and Winer 2000), and receive cortical inputs of variable strength from several areas (Winer et al. 2001). In contrast, the deep layers of the superior colliculus integrate multimodal input (Huerta and Harting 1984), its projections reach the nucleus of the brachium of the IC (Doubell et al. 2000) and the intralaminar thalamic nuclei (Krout et al. 2001), and they may enable certain types of seizures via their connections with the IC (Yang et al. 2001). The propensity of posterior thalamic neurons for oscillatory and temporally protracted activity suggests roles in processes ranging from seizure to cognition (Llinás and Paré 1997) to autonomic function (Endepols and Walkowiak 1999).



Figure 8.3. Corticocollicular (**A**) and thalamotectal (**B**) neurons after IC tracer deposits in the rat. (**A**) Corticocollicular neurons are exclusively pyramidal (1–3) and some have dendrites that reach layer I (3). *Inset*, deposit site (*solid*) and tracer diffusion (*stippled*). Protocol as in Fig. 8.2. (**B**) Neurons projecting to the IC from perithalamic territories include cells in the superior colliculus (1), nucleus of the brachium (2), and posterior limitans nucleus (3, 4). Many (1–4) have long, rather smooth dendrites and a simple branching pattern reminiscent of tegmental cells.

## 5.2. THE AMYGDALOCOLLICULAR SYSTEM

This pathway is documented in two bat species, one of which (*Antrozous pallidus*) is highly visual and uses passive echolocation (Fuzessery et al. 1993) and gleaning (Brown et al. 1978) to acquire prey, while the other (*Pteronotus p. parnellii*) has a remarkable echolocation repertoire (Suga 1964; Fitzpatrick et al. 1998). Neurons in the magnocellular basal amygdaloid nucleus were labeled bilaterally after tracer deposits in IC subdivisions that included the central nucleus (Marsh et al. 2002); the number of labeled cells approximated that in auditory cortex, suggesting a massive, rather than a modulatory, projection.

## 6. A NEW KIND OF CONVERGENCE

The three descending systems considered here terminate preferentially in the external nucleus and dorsal cortex of the IC. This implies that there is convergence of descending influences analogous to that reported for ascending pathways, that the descending pathways act as parallel systems that complement their ascending counterparts, and that there are multiple forebrain systems in hearing without a precise analogue in the visual and somatic sensory systems.

## 7. DIRECTIONS FOR FUTURE RESEARCH

Understanding how the CC and TT systems accomplish their functional tasks will require a number of complementary approaches.

### 7.1. THE CORTICOCOLLICULAR SYSTEM

Direct optical imaging of the neurons during behavioral tasks could provide insight into the sublaminar pattern of activation in the CC system. If there are parallel CC pathways, such an approach might identify them on a temporal basis.

Pharmacologic inactivation of single or small groups of cortical neurons in specific sublayers may have contrasting effects on the midbrain.

Mapping corticocolliculofugal connections with transneuronal tracers could demonstrate the downstream targets of CC neurons. Such experiments address issues of connectional and functional parallelism.

Comparing the synaptic arrangements in different IC subdivisions is an essential step in understanding how midbrain neurons and their associated circuits are under cortical control. For example, cortical influence on the central nucleus of the IC and the ventral division of the MGB may differ. Insights into intrinsic IC circuits could relate them directly to CC input.

The massive CC convergence should be correlated with inactivation studies of single areas, sets of areas, and of hemispheres.




Figure 8.4. (A) Summary of auditory corticofugal anterograde projections (origins: white letters at top; targets: left side) to the MGB (upper half) and inferior colliculus (lower half), and thalamotectal projections (origins: white letters at bottom; terminations: gray stippling). In the thalamus, the ventral division receives a heavy input, while the central nucleus (CN) of the IC receives less. Likewise, the limbic components (Ins, Te) of non-tonotopic auditory cortex have different thalamic and midbrain projection patterns (Winer et al. 1998, 2001). (B) Schematic showing possible nuclear convergence of cortical (upper panel) and thalamic (lower panel) projections to the lateral nucleus of the IC. Such input might reach the CN and affect ultimately pontine neurons, not to mention prospective commissural and tectofugal effects. The links between the amygdala and the inferior colliculus have been demonstrated only in bats (Marsh et al. 2002) and are here projected onto cat amygdaloid subdivisions (Krettek and Price 1978).

## 7.2. *THE THALAMOTECTAL SYSTEM*

Selective, reversible inactivation of TT projections could shed light on how thalamic neurons affect IC cells. This should be contrasted with the effects of cortical blockade alone and of simultaneous midbrain and cortical inactivation.

Creating transgenic species devoid of TT projections for physiologic study might reveal the functional contributions of this system and the effects of its genetic deletion.

Exploring the interactions between descending systems and ascending, commissural, and intrinsic pathways could reveal polysynaptic relationships. As the resolution of imaging methods improve, the locus and order of temporal activation patterns should be explored.

## 7.3. *THE AMYGDALOCOLLICULAR SYSTEM*

Dual simultaneous recordings in the lateral amygdala and the IC during feeding, procreation, and social behavior including vocalizations should reveal their interactions. Reversible inactivation of the amygdala through cooling or pharmacologic means could elucidate its role in complex social behaviors.

Transneuronal tract tracing with rabies virus should permit visualization of tectofugal projection neurons that themselves are the target of AC input, thus connecting subsequent synaptic chains into a more explicit and extended functional linkage that might include tectopontine or tectocochlear centers.

## Abbreviations

AAF	anterior auditory field
AC	amygdalocollicular system
AI	primary auditory cortex
APt	anterior pretectum
AII	second auditory cortex
BL	basolateral amygdaloid nucleus
BM	basomedial amygdaloid nucleus
C	caudal
CC	corticocollicular system <i>or</i> caudal cortex of the inferior colliculus
Cel	lateral division of central amygdaloid nucleus
Cem	medial division of central amygdaloid nucleus
CG	central gray
CIC	commissure of the inferior colliculus
Cl	claustrum
CN	central nucleus of the inferior colliculus
CP	cerebral peduncle
Cu	cuneiform nucleus
D	dorsal nucleus of the medial geniculate body <i>or</i> dorsal

DCa	caudal dorsal nucleus of the medial geniculate body
DC	dorsal cortex of the inferior colliculus
DCI–IV	dorsal cortex layers I–IV
DD	deep dorsal nucleus of the medial geniculate body
DNLL	dorsal nucleus of the lateral lemniscus
DS	dorsal superficial nucleus of the medial geniculate body
EC	external cortex of the inferior colliculus
EPD	posterior ectosylvian gyrus, dorsal part
EPI	posterior ectosylvian gyrus, intermediate part
EPV	posterior ectosylvian gyrus, ventral part
Ep	entopeduncular nucleus
GABA	$\gamma$ -aminobutyric acid
GP	globus pallidus
IC	inferior colliculus
IcT	intercollicular tegmentum
INLL	intermediate nucleus of the lateral lemniscus
Ins	insular cortex
La	lateral nucleus of the amygdala
LD	lateral dorsal nucleus
LGB	lateral geniculate body
LL	lateral lemniscus
LMN	lateral mesencephalic nucleus
LN	lateral nucleus of the inferior colliculus
LP	lateral posterior nucleus
M	medial division
MCP	middle cerebellar peduncle
Me	medial amygdaloid nucleus
MGB	medial geniculate body
ML	medial lemniscus
MLF	medial longitudinal fasciculus
MRF	mesencephalic reticular formation
NBIC	nucleus of the brachium of the inferior colliculus
NRTPT	reticular tegmental nucleus of the pons
Ov	<i>pars ovoidea</i> of the ventral division
P	posterior auditory field <i>or</i> putamen
PL	posterior limitans nucleus
PN	pontine nuclei
Pol	rostral pole of the medial geniculate body
Pul	pulvinar nucleus
Pu	putamen
Py	pyramidal tract
R	rostral <i>or</i> red nucleus
Ra	raphe
RN	red nucleus
RP	rostral pole nucleus of the inferior colliculus

Sa	nucleus sagulum
SC	superior colliculus
SCP	superior cerebellar peduncle
SCPX	decussation of the superior cerebellar peduncle
SF	suprasylvian fringe area
Sg	suprageniculate nucleus
Sgl	suprageniculate nucleus, lateral part
Sgm	suprageniculate nucleus, medial part
SNL	substantia nigra, <i>pars lateralis</i>
SNR	substantia nigra, <i>pars reticulata</i>
Spf	subparafascicular nucleus
SpN	suprapeduncular nucleus
Te	temporal cortex
Tr	trochlear nerve
TT	thalamotectal system
V	<i>pars lateralis</i> of the ventral division <i>or</i> ventral <i>or</i> trigeminal nerve
Ve	ventral auditory area
VI	ventrolateral nucleus of the medial geniculate body
VP	ventral posterior auditory area
I–IV	layers of the dorsal cortex

## Acknowledgments

I thank Christoph E. Schreiner, Michelle L. Chernock, and David T. Larue for helpful comments. This work was supported by United States Public Health Service Grant R01 DC02319-25.

## References

- Adams JC (1979) Ascending projections to the inferior colliculus. *Journal of Comparative Neurology* **183**:519–538.
- Aitkin LM (1973) Medial geniculate body of the cat: responses to tonal stimuli of neurons in medial division. *Journal of Neurophysiology* **36**:275–283.
- Aitkin LM and Phillips SC (1984) Is the inferior colliculus an obligatory relay in the cat auditory system? *Neuroscience Letters* **44**:259–264.
- Beneyto M and Prieto JJ (2001) Connections of the auditory cortex with the claustrum and endopiriform nucleus in the cat. *Brain Research Bulletin* **54**:485–498.
- Briggs F and Callaway EM (2001) Layer-specific input to distinct cell types in layer 6 of monkey primary visual cortex. *Journal of Neuroscience* **21**:3600–3608.
- Brodal P (1972) The corticopontine projection in the cat. The projection from the auditory cortex. *Archives Italiennes de Biologie* **110**:119–144.
- Brown PE, Grinnell AD, and Harrison JB (1978) The development of hearing in the pallid bat, *Antrozous pallidus*. *Journal of Comparative Physiology A* **126**:169–182.
- Brunso-Bechtold JK, Thompson GC, and Masterton RB (1981) HRP study of the organization of auditory afferents ascending to central nucleus of inferior colliculus in cat. *Journal of Comparative Neurology* **197**:705–722.



- Calford MB and Aitkin LM (1983) Ascending projections to the medial geniculate body of the cat: evidence for multiple, parallel auditory pathways through the thalamus. *Journal of Neuroscience* **3**:2365–2380.
- Casseday JH and Covey E (1996) A neuroethological theory of the operation of the inferior colliculus. *Brain, Behavior and Evolution* **47**:311–336.
- Code RA and Winer JA (1985) Commissural neurons in layer III of cat primary auditory cortex (AI): pyramidal and non-pyramidal cell input. *Journal of Comparative Neurology* **242**:485–510.
- Crabtree JW (1998) Organization in the auditory sector of the cat's thalamic reticular nucleus. *Journal of Comparative Neurology* **390**:167–182.
- Deschênes M, Veinante P, and Zhang Z-W (1998) The organization of corticothalamic projections: reciprocity versus parity. *Brain Research Reviews* **28**:286–308.
- Doubell TP, Baron J, Skaliora I, and King AJ (2000) Topographical projection from the superior colliculus to the nucleus of the brachium of the inferior colliculus in the ferret: convergence of visual and auditory information. *European Journal of Neuroscience* **12**:4290–4308.
- Doucet JR, Rose L, and Ryugo DK (2002) The cellular origin of corticofugal projections to the superior olivary complex in the rat. *Brain Research* **925**:28–41.
- Endepols H and Walkowiak W (1999) Influence of descending forebrain projections on processing of acoustic signals and audiomotor integration in the anuran midbrain. *European Journal of Morphology* **37**:182–184.
- Feliciano M and Potashner SJ (1995) Evidence for a glutamatergic pathway from the guinea pig auditory cortex to the inferior colliculus. *Journal of Neurochemistry* **65**:1348–1357.
- Fitzpatrick DC, Suga N, and Olsen JF (1998) Distribution of response types across entire hemispheres of the mustached bat's auditory cortex. *Journal of Comparative Neurology* **391**:353–365.
- Frigyesi T, Rinvik E, and Yahr MD (eds) (1972) *Corticothalamic Projections and Sensorimotor Activities*. Raven Press, New York.
- Fuzessery ZM, Buttenhoff P, Andrews B, and Kennedy JM (1993) Passive sound localization of prey by the pallid bat (*Antrozous p. pallidus*). *Journal of Comparative Physiology A* **171**:767–777.
- Games KD and Winer JA (1988) Layer V in rat auditory cortex: projections to the inferior colliculus and contralateral cortex. *Hearing Research* **34**:1–26.
- Gao E and Suga N (2000) Experience-dependent plasticity in the auditory cortex and the inferior colliculus of bats: role of the corticofugal system. *Proceedings of the National Academy of Sciences of the United States of America* **97**:8081–8086.
- Gerren RA and Weinberger NM (1983) Long term potentiation in the magnocellular medial geniculate nucleus of the anesthetized cat. *Brain Research* **265**:138–142.
- Hefti BJ and Smith PH (2000) Anatomy, physiology, and synaptic responses of rat layer V auditory cortical cells and effects of intracellular GABA<sub>A</sub> blockade. *Journal of Neurophysiology* **83**:2626–2638.
- Huang CL and Winer JA (2000) Auditory thalamocortical projections in the cat: laminar and areal patterns of input. *Journal of Comparative Neurology* **427**:302–331.
- Huerta MF and Harting JK (1984) The mammalian superior colliculus: studies of its morphology and connections. In: Vanegas H (ed). *Comparative Neurology of the Optic Tectum*. Plenum Press, New York, pp. 687–773.
- Hupé J-M, James AC, Girard P, Lomber SG, Payne BR, and Bullier J (2001) Feedback connections act on the early part of the responses in monkey visual cortex. *Journal of Neurophysiology* **85**:134–145.

- Imig TJ and Reale RA (1980) Patterns of cortico-cortical connections related to tonotopic maps in cat auditory cortex. *Journal of Comparative Neurology* **192**:293–332.
- Jen PH-S, Chen QC, and Sun XD (1998) Corticofugal regulation of auditory sensitivity in the bat inferior colliculus. *Journal of Comparative Physiology A* **183**:683–697.
- King AJ (1997) Signal selection by cortical feedback. *Current Biology* **7**:R85–R88.
- Krettek JE and Price JL (1978) Amygdaloid projections to subcortical structures within the basal forebrain and brainstem in the rat and cat. *Journal of Comparative Neurology* **178**:225–253.
- Krout KE, Loewy AD, Westby GWM, and Redgrave P (2001) Superior colliculus projections to midline and intralaminar thalamic nuclei of the rat. *Journal of Comparative Neurology* **431**:198–216.
- Kuwabara N and Zook JM (2000) Geniculo-collicular descending projections in the gerbil. *Brain Research* **878**:79–87.
- Llinás R and Paré D (1997) Coherent oscillations in specific and nonspecific thalamo-cortical networks and their role in cognition. In: Steriade M, Jones EG, and McCormick DA (eds). *Thalamus*, Volume II: *Experimental and Clinical Aspects*. Elsevier, Amsterdam, pp. 501–516.
- Malmierca MS, Rees A, Le Beau FEN, and Bjaalie JG (1995) Laminar organization of frequency-defined local axons within and between the inferior colliculi of the guinea pig. *Journal of Comparative Neurology* **357**:124–144.
- Marsh RA, Fuzessery ZM, Grose CD, and Wenstrup JJ (2002) Projection to the inferior colliculus from the basal nucleus of the amygdala. *Journal of Neuroscience* **22**:10449–10460.
- Mitani A, Shimokouchi M, and Nomura S (1983) Effects of stimulation of the primary auditory cortex upon colliculogeniculate neurons in the inferior colliculus of the cat. *Neuroscience Letters* **42**:185–189.
- Murphy PC, Duckett SG, and Sillito AM (1999) Feedback connections to the lateral geniculate nucleus and cortical response properties. *Science* **286**:1552–1554.
- Prieto JJ and Winer JA (1999) Neurons of layer VI in cat primary auditory cortex (AI): Golgi study and sublaminar origins of projection neurons. *Journal of Comparative Neurology* **404**:332–358.
- Przybylski AW (1998) Does top-down processing help us to see? *Current Biology* **8**:R135–R139.
- Reale RA and Imig TJ (1983) Auditory cortical field projections to the basal ganglia of the cat. *Neuroscience* **8**:67–86.
- Romanski LM, Clugnet M, Bordi F, and LeDoux JE (1993) Somatosensory and auditory convergence in the lateral nucleus of the amygdala. *Behavioral Neuroscience* **107**:444–450.
- Saint Marie RL and Baker RA (1990) Neurotransmitter-specific uptake and retrograde transport of [3H]glycine from the inferior colliculus by ipsilateral projections of the superior olivary complex and nuclei of the lateral lemniscus. *Brain Research* **524**:244–253.
- Saldaña E and Merchán MA (1992) Intrinsic and commissural connections of the rat inferior colliculus. *Journal of Comparative Neurology* **319**:417–437.
- Schreiner CE and Cynader MS (1984) Basic functional organization of second auditory cortical field (AII) of the cat. *Journal of Neurophysiology* **51**:1284–1305.
- Shinonaga Y, Takada M, and Mizuno N (1994) Direct projections from the non-laminated divisions of the medial geniculate nucleus to the temporal polar cortex and amygdala in the cat. *Journal of Comparative Neurology* **340**:405–426.

- Suga N (1964) Single unit activity in cochlear nucleus and inferior colliculus of echolocating bats. *Journal of Physiology (London)* **172**:449–474.
- Suga N, Gao E, Zhang Y, Ma X, and Olsen JF (2000) The corticofugal system for hearing: recent progress. *Proceedings of the National Academy of Sciences of the United States of America* **97**:11807–11814.
- Sun X, Chen QC, and Jen PH-S (1996) Corticofugal control of central auditory sensitivity in the big brown bat, *Eptesicus fuscus*. *Neuroscience Letters* **212**:131–134.
- Syka J and Popelář J (1984) Inferior colliculus in the rat: neuronal responses to stimulation of the auditory cortex. *Neuroscience Letters* **51**:235–240.
- Warr WB (1982) Parallel ascending pathways from the cochlear nucleus: neuroanatomical evidence of functional specialization. In: Neff WD (ed). *Contributions to Sensory Physiology*. Academic Press, New York, pp. 1–38.
- White EL and Hersch SM (1982) A quantitative study of thalamocortical and other synapses involving the apical dendrites of corticothalamic projection cells in mouse SmI cortex. *Journal of Neurocytology* **11**:137–158.
- Winer JA (1992) The functional architecture of the medial geniculate body and the primary auditory cortex. In: Webster DB, Popper AN, and Fay RR (eds). *Springer Handbook of Auditory Research*, Volume 1: *The Mammalian Auditory Pathway: Neuroanatomy*. Springer-Verlag, New York, pp. 222–409.
- Winer JA and Prieto JJ (2001) Layer V in cat primary auditory cortex (AI): cellular architecture and identification of projection neurons. *Journal of Comparative Neurology* **434**:379–412.
- Winer JA, Morest DK, and Diamond IT (1988) A cytoarchitectonic atlas of the medial geniculate body of the opossum, *Didelphys virginiana*, with a comment on the posterior intralaminar nuclei of the thalamus. *Journal of Comparative Neurology* **274**:422–448.
- Winer JA, Larue DT, and Pollak GD (1995) GABA and glycine in the central auditory system of the mustache bat: structural substrates for inhibitory neuronal organization. *Journal of Comparative Neurology* **355**:317–353.
- Winer JA, Saint Marie RL, Larue DT, and Oliver DL (1996) GABAergic feedforward projections from the inferior colliculus to the medial geniculate body. *Proceedings of the National Academy of Sciences of the United States of America* **93**:8005–8010.
- Winer JA, Larue DT, Diehl JJ, and Hefti BJ (1998) Auditory cortical projections to the cat inferior colliculus. *Journal of Comparative Neurology* **400**:147–174.
- Winer JA, Diehl JJ, and Larue DT (2001) Projections of auditory cortex to the medial geniculate body of the cat. *Journal of Comparative Neurology* **430**:27–55.
- Winer JA, Chernock ML, Larue DT, and Cheung SW (2002) Descending projections to the inferior colliculus from the posterior thalamus and the auditory cortex in rat, cat, and monkey. *Hearing Research* **168**:181–195.
- Yan J and Ehret G (2001) Corticofugal reorganization of the midbrain tonotopic map in mice. *NeuroReport* **12**:3313–3316.
- Yan J and Suga N (1996a) Corticofugal modulation of time-domain processing of biosonar information in bats. *Science* **273**:1100–1103.
- Yan J and Suga N (1996b) The midbrain creates and the thalamus sharpens echo-delay tuning for the cortical representation of target-distance information in the mustached bat. *Hearing Research* **93**:102–110.
- Yang L, Long C, and Faingold CL (2001) Neurons in the deep layers of the superior colliculus are a requisite component of the neuronal network for seizures during ethanol withdrawal. *Brain Research* **920**:134–141.

# Chapter 9

## Pharmacology of the Inferior Colliculus

JACK B. KELLY AND DONALD M. CASPARY

### 1. INTRODUCTION

The inferior colliculus (IC) receives convergent projections from a diversity of sources within the auditory pathway and is an important midbrain structure for processing acoustic information. The characteristics of the neurotransmitters and neuromodulators acting at synapses on the somata and dendrites of IC neurons are critically important in shaping responses to both monaural and binaural stimuli. The ways in which IC neurons encode information about complex sounds are determined in large measure by the neurotransmitters and receptors that regulate and control synaptic responses as well as the circuitry and biophysical properties of the neurons and their processes (see Chapter 10). The present chapter examines how the principal amino acid neurotransmitters, acting through ionotropic receptors, influence IC coding. We briefly consider how other identified neurotransmitters and neuromodulators influence the response properties of IC neurons and circuits.

We first provide an overview of the major transmitters released from the principal extrinsic projections and intrinsic pathways and then describe how these neurotransmitters shape the acoustic message as it ascends. The primary excitatory neurotransmitter, released onto IC neurons from ascending, descending, and perhaps intrinsic inputs is the excitatory amino acid, glutamate, acting at both  $\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) and *N*-methyl-D-aspartate (NMDA) receptors (Adams and Wenthold 1979; Bergman et al. 1989; Faingold et al. 1989a). Many extrinsic and intrinsic IC circuits utilize the amino acid,  $\gamma$ -aminobutyric acid (GABA), as the principal inhibitory neurotransmitter. Also, the inhibitory amino acid glycine is released from several ascending IC inputs (Faingold et al. 1991a; Winer et al. 1995). Other neuroactive substances released onto IC neurons and their dendrites include the catecholamines, serotonin and norepinephrine, as well as acetylcholine, and many peptides. Our emphasis, however, is on the pharmacology of the amino acid neurotransmitters: glutamate, GABA, and glycine.

## 2. GLUTAMATERGIC INPUTS INTO THE INFERIOR COLLICULUS

There remains little doubt that glutamate is the primary excitatory transmitter throughout the IC (Adams and Wenthold 1979; Faingold et al. 1989b; Bergman et al. 1989; Saint Marie 1996). Neurochemical and functional studies document its release, uptake, and action on both AMPA and NMDA receptors (Feldman and Knudsen 1994; Suneja et al. 2000; Kelly and Zhang 2002; Ma et al. 2002a; Wu et al. 2002). Other studies have specified the subunit makeup of the AMPA receptor and the NMDA receptor in the different IC subdivisions (Marianowski et al. 2000; Parks 2000) and have suggested that the NMDA receptor plays a role in regulating response plasticity (Feldman et al. 1996; Kelly and Zhang 2002; Wu et al. 2002). Physiologic studies have begun to assess the relative impact of parallel ascending excitatory inputs onto neurons in the central nucleus of the inferior colliculus (ICC) (Davis 2002; Davis et al. 2003).

### 2.1. ASCENDING EXCITATORY INPUTS

The ventral cochlear nucleus (VCN) is generally considered the main excitatory projection to the IC (Helfert et al. 1991; see Chapter 3). Multipolar/stellate cells in both anterior and posterior VCN project directly to the contralateral IC with collateral projections to other auditory nuclei en route. Most of these neurons are likely excitatory and glutamatergic (Wenthold 1991; Wenthold et al. 1993; Romand and Avan 1997).

Fusiform cells of the dorsal cochlear nucleus (DCN) project to the IC directly and contribute specific excitatory response properties to subpopulations of IC neurons (Davis 2002). Fusiform cells display strongly nonmonotonic rate-intensity functions around their characteristic frequency (CF) and have stereotyped temporal response properties that include buildup or pauser-buildup response patterns observed in their post-stimulus time histograms (PSTHs), that is, the temporal distribution of their action potentials over the time course of stimulus presentation. Fusiform cells show temporal locking to the envelope of sinusoidally amplitude modulated sounds with a greater discharge synchrony than expected from the modulation waveform, that is, they have modulation transfer function gains  $>1$ . This temporal locking is not altered by GABA<sub>A</sub> receptor blockade in ICC, suggesting that the high level of response synchrony is established in DCN. Some, but not all, of these temporal properties are reflected in the responses of a subpopulation of contralateral IC neurons, which provides functional evidence for a direct input from DCN fusiform cells (Casparly et al. 2002; Davis 2002). Ascending excitatory inputs from the contralateral lateral superior olive (LSO) and the ipsilateral medial superior olive (MSO) are generally considered to be sources of glutamatergic activation of IC neurons

(Casseday et al. 1988; Glendenning et al. 1992; Feldman and Knudsen 1994; Suneja et al. 2000; Kelly and Zhang 2002; Wu et al. 2002). LSO projections convey binaural information to recipient IC neurons, shaping, at least in part, their responses to interaural intensity differences, especially for high-frequency sounds (Li and Kelly 1992; Sally and Kelly 1992; Park and Pollak 1994). MSO projections impart information about interaural timing differences/interaural phase differences especially pertaining to low-frequency binaural coding. It is important to note that additional shaping of binaural responses occurs in the IC and thus the coding is not complete at the superior olive (Faingold et al. 1989a, 1993; Li and Kelly 1992; Sally and Kelly 1992; Kelly and Li 1997).

## 2.2. *DESCENDING EXCITATORY INPUTS*

The corticocollicular inputs and, to a lesser extent, geniculocollicular inputs likely are glutamatergic. However, the functional nature of this excitation, especially in the ICC, is not well understood. Layer V pyramidal neurons in auditory cortex, which project to the IC dorsal cortex (ICD), external nucleus (ICX), and central nucleus (ICC), likely possess fairly robust excitatory responses within a conventional V-shaped frequency tuning curve (Turner et al. 2003). Whether these descending inputs are in alignment/register with the frequency tuning of their midbrain targets is unknown. Whether inhibitory neurons in the ICD and the ICX convert these descending excitatory inputs into inhibition in the ipsilateral and contralateral ICC also remains an open question (Jen et al. 2001).

## 2.3. *EXTRINSIC INHIBITORY INPUTS TO THE INFERIOR COLLICULUS*

In cats and mustache bats about 20% of the IC neurons are GABAergic (Oliver et al. 1994; Winer et al. 1995). In rats 20% to 40% of the neurons in the three major subdivisions of the IC are GABA- or glutamic acid decarboxylase (GAD)-positive (Mugnaini and Oertel 1985; Caspary et al. 1990) and approximately 44% of the synapses are GABAergic (Helfert et al. 1999). GABAergic extrinsic, intrinsic, and commissural projections terminate as either focused or diffuse inhibitory inputs on IC neurons (Adams and Mugnaini 1984; Shneiderman et al. 1988, 1993; Helfert et al. 1989; Saint Marie et al. 1989; Shneiderman and Oliver 1989; Huffman and Henson 1990). Particularly high levels of GABA exist in the IC (Tachibana and Kuriyama 1974; Banay-Schwartz et al. 1989b) and many neurons that label for GABA/GAD likely affect neuronal responses through intrinsic or commissural projections (Fisher and Davies 1976; Adams 1979; Contreras and Bachelard 1979; Mugnaini and Oertel 1985; Nagai et al. 1985; Thompson et al. 1985; Moore and Moore 1987; Roberts and Ribak 1987a,b; Carr et al. 1989; Wenthold 1991; Smith 1992; Vater et al. 1992; Oliver et al. 1994; González-Hernández et al. 1996; Zhang et al. 1998; Jen et al. 2001; see Chapter 10).

GABAergic inputs arise from several extrinsic sources, including a projection from the contralateral dorsal nucleus of the lateral lemniscus (DNLL) and a smaller ipsilateral DNLL projection (Adams and Mugnaini 1984; Li and Kelly 1992; Vater et al. 1992b; Faingold et al. 1993; Shneiderman et al. 1993). In the rat, the superior paraolivary nucleus of the superior olivary complex sends direct GABAergic input to the ipsilateral ICC (Kulesza et al. 2003). Nearly half of the neurons in LSO are glycinergic and send direct input to the ipsilateral IC (Helfert et al. 1989; Saint Marie et al. 1989; Glendenning et al. 1992; Vater et al. 1992b; Vater 1995; Ostapoff et al. 1997). Inhibitory input also arises from the medial geniculate body (Vater et al. 1992b) and perhaps the auditory cortex (Adams and Wenthold 1979; Jen et al. 2001). Connectional studies using small injections of tracers into tonotopically identified target areas suggest that GABAergic projections from the DNLL and the collicular commissure may be tonotopically aligned with their ICC targets (Saldaña and Merchán 1992; Merchán et al. 1994; Malmierca et al. 1996; Bajo et al. 1999). These observations have important implications for the shaping of frequency-response tuning curves and the coding of modulated stimuli.

#### 2.4. ROLE OF $\gamma$ -AMINO BUTYRIC ACID IN THE CENTRAL NUCLEUS

All criteria needed to establish GABA as an IC neurotransmitter have been satisfied. High levels of the synthetic enzyme GAD and the degradative enzyme GABA transaminase are found in the ICC (Fisher and Davies 1976; Adams and Wenthold 1979; Contreras and Bachelard 1979; Nagai et al. 1985; Vater et al. 1992b; Raza et al. 1994). GABA is released from ICC slices/punches in a  $\text{Ca}^{2+}$ -dependent manner (López-Colomé et al. 1978; Banay-Schwartz et al. 1989b; Caspary et al. 1990; Shneiderman et al. 1993) and a high-affinity GABA uptake system exists in the ICC (Shneiderman et al. 1993; Raza et al. 1994; Suneja et al. 1998). Immunocytochemical probes for conjugated GABA and GAD show dense labeling in somata, fibers, and axon terminal puncta within the ICC, which has among the highest GAD labeling in the auditory brain stem (Tachibana and Kuriyama 1974; Mugnaini and Oertel 1985; Thompson et al. 1985; Moore and Moore 1987; Carr et al. 1989; Caspary et al. 1990).

The GABA<sub>A</sub> receptor was initially demonstrated in binding studies (Glendenning and Baker 1988), and high GABA<sub>A</sub> receptor binding levels are seen in the rat brain, especially for the benzodiazepine, flunitrazepam (Fig. 9.1A), and the GABA<sub>A</sub> receptor channel marker at the picrotoxin site, *t*-butylbicyclo-orthobenzoate (TBOB) (Fig. 9.1C). Muscimol binding in the IC (Fig. 9.1B) is less dramatic than in the cerebellum, where it may selectively bind different GABA<sub>A</sub> subunit constructs.

The GABA<sub>A</sub> receptor is a heteromeric pentamer with >20 subunits. The adult wild-type GABA<sub>A</sub> receptor likely consists of  $2\alpha_1 2\beta_2 \gamma_2$  (Chebib and Johnston 1999) and the GABA<sub>A</sub> receptor subunit construct will partly determine the response to presynaptically released GABA at IC neuronal synapses. The subunit makeup of the GABA<sub>A</sub> receptor in IC subregions is known (Milbrandt et al.




Figure 9.1. Receptor binding of three GABA<sub>A</sub> receptor ligands to sites in rat IC (for abbreviations, see the list). Flunitrazepam binds benzodiazepine sites located between the  $\alpha$  and  $\gamma$  GABA<sub>A</sub> subunits. TBOB (*t*-[<sup>3</sup>H]butylbicycloorthobenzoate) is thought to bind in the GABA<sub>A</sub> receptor channel at the picrotoxin or cage convulsant site, and muscimol is thought to bind at the GABA binding loci on the  $\alpha$  subunit. The distribution of these three ligands in a sagittal view of the rat brain is very different. Both TBOB and flunitrazepam show very high levels of binding in IC relative to other brain structures and their distributions within the IC are different.



1997), as is the distribution of GABA<sub>A</sub> receptor subunits, for different ICC cell types (Shiraishi et al. 2001). GABA<sub>B</sub> receptor binding occurs throughout the rat's IC (Milbrandt et al. 1994) and is prominent in the dorsomedial IC of the bat (Fubara et al. 1996). Whether these receptors are presynaptic sites capable of modulating excitatory or inhibitory inputs or postsynaptic sites opposite GABA terminals is unknown (Bormann 2000). Physiologic studies provide functional and pharmacologic support for the importance of GABA<sub>A</sub> and GABA<sub>B</sub> receptors in ICC.

### 3. BRAIN SLICE STUDIES OF SYNAPTIC PHARMACOLOGY

In vitro brain slice methods are a powerful tool for investigating synaptic pharmacology. Stable intracellular recordings from one cell over an extended period allow the effects of drugs, for example, selective receptor agonists and/or antagonists, to be evaluated by adding them to the solution bathing the tissue slice. Drug concentration can be precisely controlled and the drug effects can usually be reversed by washing the slice in normal saline. Furthermore, the neurons that provide physiologic data can be filled intracellularly with biocytin, neurobiotin, or Lucifer yellow. Recordings made in either current-clamp or voltage-clamp modes using either sharp electrodes or whole-cell patch-clamp electrodes allow manipulation of the intracellular environment by modifying the solution in the recording pipette to assess membrane properties in detail.

In brain slice studies, synaptic responses are usually evoked by current-pulse stimulation of the tracts that terminate on the neurons from which recordings are made. Such stimulation usually elicits complex synaptic effects involving multiple inputs from potentially diverse sources. Moreover, each current pulse activates synaptic inputs synchronously and may not resemble the timing or input combinations associated with natural acoustic stimulation. Nevertheless, the brain slice procedure provides important insight into the pharmacology of neural circuits that contribute to hearing.

Electrical stimulation of the lateral lemniscal pathways projecting to the ICC evokes excitatory and inhibitory postsynaptic responses converging on a common neural population (Wagner 1996; Moore et al. 1998; Reetz and Ehret 1999; Ma et al. 2002a). Most ICC neurons receive both excitatory and inhibitory input from ascending fibers.

Excitatory responses are mediated by AMPA and NMDA receptors, which are usually present on the same postsynaptic neuron, possibly at the same terminals, and can be activated by the same current pulse stimulus (Ma et al. 2002a). With the brain slice in normal saline, two postsynaptic excitatory response components can be distinguished based on their time of occurrence following current-pulse stimulation. An initial early response with short latency is followed by a more slowly developing late response that produces a second peak in the excitatory postsynaptic potential (EPSP) or postsynaptic current (EPSC). Application of AMPA receptor antagonists, for example, 6-cyano-7-

nitroquinoxaline-2,3-dione (CNQX) or 1,2,3,4-tetrahydro-6-nitro-2,3-dioxo-benzo[*f*]quinoxaline-7-sulfonamide (NBQX) causes a decrease in the amplitude of the early synaptic response. The early response can be blocked by these AMPA antagonists without eliminating the later response.

The later postsynaptic response can be selectively blocked by the NMDA receptor antagonists, ( $\pm$ )-2-amino-5-phosphonovaleric acid (APV) or ( $\pm$ )-3-(2-carboxypiperazin-4-yl)-propyl-1-phosphonic acid (CPP). These drugs can eliminate the late response while leaving the early response intact. Thus, the early and late excitatory synaptic currents in the ICC can be separated pharmacologically. A more accurate assessment of the timing of these excitatory currents comes from recordings in which inhibitory responses are blocked by receptor antagonists. AMPA receptor-mediated EPSPs have shorter rise times than NMDA receptor-mediated EPSPs (means: 5.4 and 22.8 ms, respectively) and shorter decay times (35.6 and 92.1 ms), shorter times to peak response (19.5 ms and 62.9 ms), and smaller half-widths (42.6 and 120.8 ms) (Ma et al. 2002a). The NMDA receptor-mediated response is greatly enhanced in a Mg-free saline solution and, with Mg, the response size is dependent on the cell's membrane voltage. In contrast, the AMPA receptor-mediated response is unaffected by Mg concentration and is not voltage dependent. The size of the late NMDA receptor-mediated response is increased by depolarization and decreased by hyperpolarization of the cell membrane relative to the resting potential. The sensitivity of ICC neurons to membrane voltage manipulation varies considerably from cell to cell, suggesting cell populations with different input-output characteristics (physiologic gain) (Ma et al. 2002b).

For most ICC neurons, both AMPA and NMDA receptor-mediated responses can be elicited at the cell's resting potential and it is unnecessary to activate AMPA receptors or depolarize the cell membrane to demonstrate a long-lasting NMDA receptor-mediated response. An NMDA receptor-mediated current can often be found even when the AMPA receptor-mediated response is blocked by AMPA receptor antagonists, suggesting that both AMPA and NMDA receptor-mediated excitatory responses are available at or near the threshold for hearing and play important roles in processing sensory information even at low stimulus intensity. This expectation has been confirmed by *in vivo* recordings from ICC neurons in response to acoustic stimulation (Zhang and Kelly 2001, 2003).

Inhibitory responses in the ICC elicited by lemniscal stimulation are mediated by GABA<sub>A</sub> or glycine receptors (Moore et al. 1998; Reetz and Ehret 1999; Ma et al. 2002a). Bath application of the selective GABA<sub>A</sub> receptor antagonists, bicuculline or picrotoxin, reduces or eliminates inhibitory postsynaptic potentials (IPSPs) in most ICC neurons. However, the glycine receptor antagonist, strychnine, also reduces postsynaptic inhibition in many cases. Probable sources of glycinergic inhibition are the ipsilateral LSO and the ventral nucleus of the lateral lemniscus (VNLL) (Vater et al. 1997; Riquelme et al. 2001). Probable sources of GABAergic input are the lateral lemniscal nuclei, VNLL and DNLL (Adams and Mugnaini 1984; Li and Kelly 1992; Vater et al. 1992b; Faingold et al. 1993; Shneiderman et al. 1988, 1993), the superior paraolivary nucleus (Ku-

lesza et al. 2003), or intrinsic connections within the ICC. Any of these could contribute to postsynaptic responses depending on the placement of the stimulating electrode.

The glycinergic projection from the ipsilateral LSO and the GABAergic projection from the contralateral DNLL play functionally parallel roles in inhibiting ICC neurons. Because of their unique binaural innervation pattern, LSO neurons are strongly excited by ipsilateral acoustic stimulation and inhibited by contralateral stimulation. ICC target neurons are excited by crossed (glutamatergic) efferent projections and inhibited by uncrossed (glycinergic) LSO projections. Thus a sound source on the left would thus excite neurons in the right ICC and inhibit those in the left ICC. In contrast, DNLL neurons are typically excited by contralateral and inhibited by ipsilateral stimulation. The crossed GABAergic projection from the DNLL inhibits responses in the opposite ICC. Thus, a sound on the left would excite the right DNLL, which would in turn inhibit the left ICC. Both the ipsilateral (glycinergic) LSO projection and the contralateral (GABAergic) DNLL projection serve to enhance the contrast in neural activity between left and right ICC through a lateral inhibitory process and amplify (or at least maintain) differences in the response strength on each side of the brain. Both inhibitory projections would reinforce the functional contralaterality of auditory processing in the midbrain and above.

Many ICC neurons receive convergent input from the lateral lemniscus and the opposite IC via commissural projections (Moore et al. 1998). In the gerbil, current-pulse stimulation of either the lateral lemniscus or the commissure of the IC can evoke postsynaptic responses in the same ICC neuron. Commissural evoked responses can be either excitatory or inhibitory; the former are exclusively GABAergic and can be blocked by bicuculline. Thus, the IC commissure might also contribute to the functional laterality of responses.

GABA<sub>B</sub> receptors also contribute to synaptic responses in ICC. In brain slice studies, when AMPA and NMDA receptor-mediated excitatory responses and glycinergic inhibitory responses are eliminated pharmacologically, the remaining inhibitory responses can be completely blocked by bicuculline and are not affected by the GABA<sub>B</sub> antagonist, phaclofen (Ma et al. 2002c). On the other hand, the GABA<sub>B</sub> receptor agonist, baclofen, decreases IPSP amplitude and this effect can be blocked by phaclofen. Baclofen did not affect the membrane conductance of ICC neurons whereas the GABA<sub>A</sub> receptor agonist, muscimol, greatly increased conductance. This result suggests that GABA<sub>B</sub> receptors in the ICC influence synaptic transmission primarily by presynaptic modulation of transmitter release. Presynaptic GABA<sub>B</sub> receptors can affect synaptic inhibition by controlling the amount of transmitter acting at postsynaptic GABA<sub>A</sub> receptor sites.

Also, presynaptic GABA<sub>B</sub> receptors can regulate excitatory neurotransmitter release in the ICC (Wu et al. 2004) as well as in other structures in the brain (Potashner 1979; Misgeld et al. 1995; Brenowitz et al. 1998; Lim et al. 2000).

Results from brain slice recordings show that ICC PSPs can be potentiated by strong repetitive (tetanic) stimulation of the lateral lemniscus (Zhang and Wu 2000). These effects persist over several hours of recording and resemble, in




Figure 9.2. Schematic view of GABA and glutamate receptor sites in the ICC based on physiologic recordings and pharmacologic manipulations in brain slice studies of auditory midbrain neurons. The magnitude and sign of postsynaptic potentials (*PSP*) are determined by the balance and timing of excitatory and inhibitory inputs from glutamatergic (*Glu*) and GABAergic terminals. Excitation is mediated by both AMPA and NMDA receptors eliciting relatively early and late excitatory postsynaptic potentials, respectively (upward deflections in the schematic representation of typical responses to current-pulse stimulation). Inhibition is mediated by GABA acting on GABA<sub>A</sub> receptors, resulting in an inhibitory postsynaptic potential (a downward deflection in the representative traces). The presynaptic release of both GABA and glutamate is regulated by the activation of GABA<sub>B</sub> receptors located on the axon terminals. The normal mechanism for activation of the GABA<sub>B</sub> receptors is known. GABA release is likely regulated by GABA<sub>B</sub> receptors at the synaptic cleft (autoreceptors), but regulation by presynaptic inhibitory terminals cannot be excluded. Glutamate release might be regulated by presynaptic GABAergic terminals (as indicated), but receptor sites without corresponding axon terminals cannot be excluded. Ionotropic (AMPA, NMDA, and GABA<sub>A</sub>) receptors are shown as *nonshaded transmembrane pores*. Metabotropic (GABA<sub>B</sub>) receptors appear as *solid trapezoidal symbols*. Not shown are the receptors for glycine and other neurotransmitters/modulators that affect auditory midbrain physiologic responses.

most respects, the long-term potentiation (LTP) reported elsewhere in the brain (Bliss and Collingridge 1993; Bear and Malenka 1994). The NMDA receptor antagonist, APV, blocks LTP induction in this preparation, indicating that the NMDA receptor plays a key role in ICC synaptic plasticity as it does in other brain centers. GABA<sub>B</sub> receptor antagonists block LTP induction, whereas the GABA<sub>B</sub> receptor agonist, baclofen, facilitates it, probably by a downward regulation of presynaptic GABA release (Zhang and Wu 2000).

Figure 9.2 shows the probable convergence of synaptic input onto ICC neurons and the potential roles of various receptor types in regulating excitation and inhibition as determined from brain slice studies.

Recordings from brain slice preparations of the IC have examined the dorsal cortex (ICD) and external nucleus (ICX) as well as the ICC (Pierson et al. 1989; Smith 1992; Li et al. 1998, 1999), but reveal little evidence of pharmacologic differences at the synaptic level. In the ICX, ICD, and ICC, excitatory convergence is mediated by glutamate and inhibition largely by GABA. Whether these synapses exhibit unique characteristics that reflect each subdivision's functional specialization will require further study.

#### 4. IN VIVO STUDIES OF THE INFERIOR COLLICULUS

The pharmacology of ICC neurons has been studied *in vivo* by recording sound-evoked responses before, during, and after drug release at the recording site, usually by iontophoresis. These studies have the advantage of showing the contribution of specific receptors to activity elicited by natural acoustic rather than direct electrical stimulation. Moreover, adult, peripherally altered, and aged animals can be used. The chief disadvantage is the difficulty of controlling drug concentration at the recording site. Although application of the agent is proportional to the current or pressure magnitude over a range of values, an accurate assessment of drug concentration at its site of action is problematic. Concentration decreases as a cubic function of distance, causing a diminishing concentration gradient as the drug spreads from the pipette tip (Foeller et al. 2001). The site of action of a drug on a given neuron is dependent on the size of the molecule, on whether the ligand is actively incorporated by neurons or glia, and on the location of the pipette relative to receptors being studied. Because many IC neurons have substantial dendritic arbors that extend far from the recording site, distal receptors are likely less affected than more proximal dendritic and somatic receptors. Morphologic knowledge of the receptor location is important for interpreting iontophoretic results.

Early studies of *in vivo* pharmacology show the significant contribution of glutamate and GABA in determining neural acoustic response properties in the IC (see Faingold et al. 1991b). Glutamate plays a major role in evoking excitatory responses to tones or other acoustic stimuli and GABA mediates most inhibitory responses. Application of the GABA<sub>A</sub> antagonist, bicuculline, increases sound-evoked firing rate in most neurons, as does glutamate, whereas

glutamate antagonists (e.g., APV) reduce it. Other transmitters, however, affect the level of neural activity, with glycine contributing to the inhibition of responses and acetylcholine, serotonin (5-hydroxytryptamine), and norepinephrine modifying the responses in the IC.

#### *4.1. AMPA AND NMDA RECEPTORS AND THEIR ANTAGONISTS*

Contralateral acoustic stimulation of neurons in the rat ICC involves both AMPA and NMDA glutamate receptor subtypes (Zhang and Kelly 2001, 2003). Iontophoretic release of the selective AMPA receptor antagonist, NBQX, decreased excitatory responses to contralaterally presented 100-ms tone bursts in most IC neurons and blocked responses entirely in some. The blocking effect was present during both early (20-ms) and late (20- to 120-ms) periods of a sustained response to the tone bursts. The selective NMDA receptor antagonist, CPP, also had a blocking effect on most ICC neurons and this effect was strongest during the late part of the response. Thus, the AMPA and NMDA receptor antagonists had distinct and selective effects on specific components of a sustained spike discharge to a contralateral tone, as expected from the different time constants of these receptor types noted in brain slice preparations. The NMDA receptor contribution to auditory responses in rat ICC has been described previously (Faingold et al. 1989b, 1991b), and the differential effect of NMDA and non-NMDA glutamate receptor types has been reported for the barn owl's auditory midbrain homologue of the mammalian IC (Feldman and Knudsen 1994).

Both AMPA and NMDA receptors contribute to responses evoked by acoustic stimulation at all sound pressure levels above threshold, as indicated by rate-intensity curves derived from responses to contralateral tone bursts (Zhang and Kelly 2001; Kelly and Zhang 2002). AMPA and NMDA receptor antagonists reduced the firing rate of most ICC neurons throughout their dynamic ranges. These data show that NMDA receptors can mediate responses at low to moderate levels of acoustic stimulation, and therefore probably play an important role in processing sensory information.

AMPA and NMDA receptors also contribute to the responses of ICC neurons to sinusoidal amplitude-modulated (AM) tones (Zhang and Kelly 2003). The selective antagonists (NBQX for AMPA, CPP for NMDA receptors) reduced the firing rate of most ICC neurons to AM stimuli across a wide range of modulation frequencies, but did not have selective effects at different modulation frequencies. The shape of the modulation transfer functions for firing rate ( $MTF_{FR}$ ) was unchanged during drug application although overall firing rate was greatly reduced. Furthermore, the synchrony of spike discharges to the modulation cycle was largely unaffected by either receptor antagonist. Thus, the modulation transfer functions for vector strength (a measure of response synchrony,  $MTF_{VS}$ ) were usually not changed. This suggests that temporal locking to the envelope of AM signals is coded below the level of the IC (Fig. 9.3).




Figure 9.3. The contribution of AMPA and NMDA receptors to excitatory responses recorded from the ICC. (A) Both the AMPA receptor antagonist, NBQX, and the NMDA receptor antagonist, CPP, reduced the number of spikes in response to a 120-ms tone burst presented to the contralateral ear. The AMPA receptor antagonist reduced the response during both early (first 20 ms) and late (subsequent 100 ms) parts of the response with a slightly greater early effect, whereas the NMDA receptor antagonist had a greater effect on the late response. (B, C) Both NBQX and CPP reduced firing across a wide range of tone burst intensities as indicated by the rate-level curves (*panels on the left side*). Responses to sinusoidal AM tones were also reduced by both AMPA and NMDA antagonists as shown by the modulation transfer functions for firing rate ( $MTF_{FR}$ ) (*middle panels*) for NBQX and CPP. Neither drug substantially reduced the synchrony of firing to the period of the modulation envelope, as indicated by the modulation transfer functions for vector strength ( $MTF_{VS}$ ) (*panels on the right side*). The arrows (*left panels*) (rate level curves) indicate the level at which modulated sounds were presented. (Adapted from Zhang and Kelly 2001, 2003.)

#### *4.2. GABA SHAPES RECEPTIVE FIELDS AND FREQUENCY-RESPONSE TUNING CURVES*

The importance of GABA in regulating IC physiologic responses is well established. GABA application onto cat or rat IC neurons reduces the tone-evoked firing rate for almost all cells that respond to acoustic stimulation (Watanabe and Simada 1973; Faingold et al. 1989a, 1991a,b). Similar results are found with application of the GABA uptake inhibitor, nipecotic acid. The shapes of the rate-intensity functions for CF tones were altered from nonmonotonic to more monotonic during application of GABA<sub>A</sub> receptor blockers (Faingold et al. 1991a). Near-threshold responses were less affected by GABA<sub>A</sub> receptor blockade than above-threshold responses (Faingold et al. 1991a; Yang et al. 1992; Le Beau et al. 1996, 2001; Palombi and Caspary 1996a,b).

Similar studies in bat, guinea pig, and chinchilla describe receptive field (frequency tuning curve) changes after blockade of GABA<sub>A</sub> receptors in the IC (Vater et al. 1992a; Yang et al. 1992; Palombi and Caspary 1996a; Le Beau et al. 2001). There are two populations of ICC neurons, in roughly equal numbers, that respond differently to GABA<sub>A</sub> antagonists. One response type displays significant firing rate changes only in the excitatory response area of the monaural (contralateral) receptive field. These changes resemble the nonmonotonic CF rate-intensity curves that become more monotonic on GABA<sub>A</sub> receptor blockade (Faingold et al. 1991a). Such neurons had large increases in near-CF discharge rates as stimulus intensity within their receptive fields was increased, with little or no change in firing rate outside or lateral to the boundaries of the predrug receptive field (Vater et al. 1992a; Yang et al. 1992; Palombi and Caspary 1996a; Le Beau et al. 2001). Thus, one large population of IC neurons did not display GABA-mediated lateral inhibition (Palombi and Caspary 1996a). GABA<sub>A</sub> receptor blockade in the second group resulted in similarly large near-CF changes in firing rate at higher intensities within the receptive field, but with a broadening of the frequency range represented in the excitatory tuning curve. Some neurons had significant low-side expansion of their frequency tuning curves (Yang et al. 1992; Le Beau et al. 2001).

Two response types occur in guinea pig IC, one with V-shaped tuning curves and the other with more complex frequency tuning properties (Le Beau et al. 2001). GABA<sub>A</sub> receptor blockade did not alter frequency tuning of cells with V-shaped frequency-response curves, although it did increase firing rate within the excitatory response area. The frequency-response properties of the complex IC neurons were altered by GABA<sub>A</sub> receptor blockade, becoming more V-like and showing increased discharge rate within the excitatory response area.

There is no compelling evidence from these studies for the presence of neurons in IC with "pure" lateral inhibition, that is, cells with a change in firing confined to frequencies above or below the excitatory response area with no change inside this area (Fig. 9.4). There are no neurons for which GABA antagonists increase firing outside the boundaries of the predrug excitatory tuning






Figure 9.4. Three possible models of the relationship in the frequency domain between excitatory and inhibitory inputs converging onto an ICC neuron. Theoretical frequency tuning curves (*top*) are shown for the excitatory (*black*) and inhibitory (*gray*) input neurons. The *horizontal line* indicates a mid-level intensity. Theoretical isointensity contours (*bottom*) are shown before and during bicuculline methiodide application at this intensity. In (**A**), the putative inhibitory inputs are aligned tonotopically with the excitatory inputs; blockade of GABA<sub>A</sub> receptors by bicuculline would increase the rate of firing within the excitatory response area, with the greatest effect near CF, but without increasing the range of effective frequencies. This first response type was observed in 30% to 50% of ICC neurons in the studies reviewed in the text. (**B**) The inhibitory inputs are aligned with, but extend lateral to, the excitatory inputs; blockade by bicuculline would increase both the firing rate within the excitatory response area and the range of effective frequencies, especially on the low-frequency side. This second response type was observed for 30% to 50% of the neurons tested in the reviewed studies. (**C**) Inhibitory inputs flank excitatory inputs; bicuculline blockade increases the response area width with no discharge rate increase near CF. No iontophoretic studies of ICC neurons showed this response pattern following GABA<sub>A</sub> receptor blockade.

curve without also increasing the firing to frequencies within the tuning curve, particularly near the cell's characteristic frequency.

The absence of neurons with "pure" lateral inhibition suggests that inhibitory inputs arising from neurons with CFs exclusively above or below (i.e., tonotopically misaligned with) the excitatory response areas of their IC target neurons do not exist (Palombi and Caspary 1996a). Collectively, these contralateral monaural findings suggest at least two populations of IC neurons classified by response type (Fig. 9.4). Both frequency-response curve types show powerful GABA<sub>A</sub> receptor-mediated inhibition within the excitatory receptive field, near-CF, suggesting that extrinsic and intrinsic inhibitory inputs onto ICC neurons are in tonotopic alignment with excitatory target cells and have tuning curves with similar breadth (i.e., similar  $Q$  values) (Fig. 9.4A). The second response curve type reveals some "lateral inhibition" likely due to inhibitory inputs arising from projection neurons with broader tuning (lower  $Q$  values) than their excitatory IC target neurons (Fig. 9.4B). In some cases high  $Q$ -value inhibitory neurons with slightly offset CFs could increase the complexity of ICC frequency-response maps (Le Beau et al. 2001).

### 4.3. GABA SHAPES BINAURAL RESPONSES

In vivo and in vitro studies indicate that some binaural coding can occur within the IC itself. Binaural responses shaped by excitatory–inhibitory interactions (usually elicited by stimulation of the contra- and ipsilateral ear, respectively) are altered by local IC GABA<sub>A</sub> receptor blockade (Vater et al. 1992a; Park and Pollak 1993; Klug et al. 1995; Zhang et al. 1999). One probable origin of the inhibitory binaural influence on these cells is through the crossed GABAergic DNLL projection (Adams and Mugnaini 1984; Chen et al. 1999). Cutting the crossed DNLL projection, that is, severing the commissure of Probst, or pharmacologic block of excitation of DNLL cells that give rise to this projection, results in a release from binaural inhibition in the rat IC (Li and Kelly 1992; Faingold et al. 1993; Kidd and Kelly 1996; van Adel et al. 1999; Kelly and Kidd 2000). This projection could supplement or sharpen binaural responses originating in subcollicular nuclei or create *de novo* binaural interactions within the IC (Park and Pollak 1993; Kelly and Li 1997). Local GABAergic circuits could also contribute to binaural processing in the IC.

Selective responses to the direction of virtual motion generated by binaural phase cues can be recorded from IC neurons (Spitzer and Semple 1998). Similar results are found with dynamic binaural level cues (Sanes et al. 1998). Blocking GABAergic inhibition by local application of bicuculline increases the IC neurons' sensitivity to dynamic binaural motion cues (McAlpine and Palmer 2002). This effect is attributed to adaptation-of-excitation, a process by which the initial level of excitation influences the magnitude of subsequent binaural responses (McAlpine et al. 2000). The release from inhibition by bicuculline enhances the level of excitation and creates an exaggerated response to dynamically changing binaural phase cues.

The role of GABA in shaping responses to binaural time differences was examined in the barn owl's IC (Fujita and Konishi 1991). Bicuculline increased firing rate and decreased the selectivity of responses to binaural input and GABA had the opposite effect. These results suggest that GABAergic inhibition in IC sharpens neural responses to interaural time differences. The underlying mechanism for such sharpening is probably similar to that proposed for mammalian IC neurons. GABAergic inhibition in the barn owl's auditory midbrain also plays a role in adjusting responses to the spatial location of a sound source through sensory experience (Zheng and Knudsen 1999). Furthermore, in the frog's auditory tectum (IC) bicuculline altered the direction-dependent tuning of neurons to sound frequency (Zhang et al. 1999).

#### *4.4. GABAERGIC EFFECTS ON RESPONSES TO MODULATED SOUNDS*

Neurons in the IC are responsive to both sinusoidal AM and frequency-modulated (FM) stimuli (Rees and Møller 1983). Bicuculline substantially increased discharge rates to modulated sounds, as expected from studies with unmodulated tone bursts. Some neurons increased their firing equally across all modulation frequencies with no change in the selectivity for particular rates of modulation. For other cells the increase in firing rate was proportional to the excitatory response produced prior to drug release (Le Beau et al. 1996; Burger and Pollak 1998; Koch and Grothe 1998; Caspary et al. 2002; Zhang and Kelly 2003). In these cells, the effect of bicuculline at specific modulation rates was greatest at modulation frequencies that produced the strongest excitation before bicuculline. Still other neurons increased their activity preferentially at low modulation frequencies (Caspary et al. 2002; Zhang and Kelly 2003; Fig. 9.5).

The pattern of increased firing produced by bicuculline at different modulation rates might be determined in part by the receptor mechanism underlying the excitatory response. Excitation based on AMPA receptor activation would likely be level independent so that the release from GABAergic inhibition would cause a constant increase in firing rate. On the other hand, excitation based on NMDA receptors would depend on the background level of activity and the extent of membrane depolarization. Thus, the release from inhibition produced by bicuculline would likely result in a larger increase in firing at modulation frequencies that produced more excitation before drug release. The finding that most neurons in rat IC are driven by both AMPA and NMDA receptors lends some support to this idea (Zhang and Kelly 2003). The proportion of AMPA to NMDA receptor-mediated excitation for an ICC cell might determine its response gain to increasing levels of activity due to acoustic stimulation or GABAergic release from inhibition. The response to modulated or unmodulated sounds would depend on the combined influence of AMPA, NMDA, and GABA<sub>A</sub> receptor-mediated interactions.

Some neurons in ICC have band-reject MTF<sub>FR</sub> profiles, that is, they respond well to high- and low-modulation frequencies but have low response rates to




Figure 9.5. The effects of bicuculline on responses to CF tone bursts and sinusoidal AM tones. Iontophoretic application of bicuculline methiodide increased firing rates to both tones and modulated sounds, but had little effect on the synchrony of discharges to the envelope of the AM tone as indicated in the rate level functions (*RLF*) and modulation transfer functions for firing rate (*MTF<sub>FR</sub>*) and vector strength (*MTF<sub>VS</sub>*) (panels from left to right) for three ICC neurons (A–C). For the neuron in (A), bicuculline increased firing by about the same amount for all modulation frequencies. For the two neurons in (B) and (C), the increase in firing was more nearly proportional to the firing level before drug application. For the neuron in (B) the increase was more pronounced for modulation rates below than above best-modulation frequency. For all panels the predrug level is indicated by filled squares (■), the drug condition by crosses (X) and the postdrug recovery by open squares (□). (Adapted from Zhang and Kelly 2003.)

middle frequencies (Krishna and Semple 2000; Zhang and Kelly 2003). For these neurons, bicuculline increased the overall firing rate, but did not eliminate the band-reject feature of the  $MTF_{FR}$ . In some cases, the increase in activity was constant across a wide range of modulation frequencies, but in others it was greater at modulation rates that produced high firing levels before drug application (Fig. 9.6). These results suggest that the consequences of release from inhibition might be related to the type of underlying excitation, that is, whether the excitatory input is based on AMPA or NMDA receptors and in what pro-




Figure 9.6. The effect of bicuculline on two neurons (**A**, **B**) with band-reject modulation transfer functions ( $MTF_{FR}$ ) in response to a sinusoidal AM tone presented to the contralateral ear. Both neurons responded well to high and low modulation frequencies before drug application, but had much weaker responses to intermediate modulation frequencies. For the neuron shown in (**A**), bicuculline methiodide increased firing by nearly equal amounts at all AM frequencies. In contrast, for the neuron shown in (**B**), the increase in firing was nearly proportional to the level of activity prior to drug application. The band-reject profile of the  $MTF_{FR}$  was preserved in both cases. In neither case was the discharge synchrony to the modulation envelope greatly affected by the drug. For other conventions, see Fig. 9.5. (Adapted from Zhang and Kelly 2003.)

portion. The band-reject feature of the  $MTF_{FR}$  for many ICC neurons appears to be shaped primarily by neural circuits at lower levels of the auditory system rather than local GABAergic synapses.

Bicuculline blockade of  $GABA_A$  receptors did not strongly affect the synchrony of discharges locked to the modulation envelope of AM tones, even when firing rate was strongly affected (Zhang and Kelly 2003). Vector strength remained high at most modulation frequencies and the temporal pattern of synchronous discharge was similar before and during bicuculline release; similar results were seen in the mustached bat (Burger and Pollak 1998) and chinchilla (Caspary et al. 2002). Thus, response synchrony is not disrupted by local imbalances in excitation and  $GABA_A$  receptor-mediated inhibition in IC created by receptor antagonists. Response synchrony is likely governed by the synchronized discharge of inputs from extrinsic sources (likely the DCN), local circuits in the ICC, and/or the intrinsic membrane properties of the neurons from which recordings are made.

$GABA_A$  receptors help shape ICC responses to sinusoidal FM tones in the big brown bat (Koch and Grothe 1998, 2000).  $GABA_A$  receptor blockade reduced the sharpness of tuning to the rate of FM in 60% of the cells, affecting the upper and lower cutoff frequencies of the MTF. The sharpness of tuning was increased in 30% of the cells. Bicuculline also reduced response synchrony to the modulation envelope in many cells owing primarily to a prolonged discharge during the modulation cycle. Temporal selectivity to modulation rate was affected by the strength of binaural inhibition in about half the neurons in the ICC, and local  $GABA_A$  receptor blockade altered this binaural dependency in some cells (Koch and Grothe 2000).

Many pallid bat ICC cells have directional FM sensitivity dependent on local  $GABA_A$  receptors, and this can be abolished or reduced with bicuculline application (Fuzessery and Hall 1996). Koch and Grothe (1998) also reported that bicuculline abolished the FM directional sensitivity of some neurons in the big brown bat ICC.

#### 4.5. GABAERGIC EFFECTS ON RESPONSES TO SOUND DURATION

Many IC neurons in bats (Fuzessery and Hall 1999; Casseday et al. 2002) and other mammalian species (chinchilla: Chen 1998; mouse: Brand et al. 2000) are selectively tuned to sounds of a specific duration. Some cells respond maximally to short-duration sounds (short-pass tuning) and others prefer a specific duration rather than longer or shorter durations (band-pass tuning). A “coincidence model” has been proposed for ICC neural duration tuning in the big brown bat based on the convergence of delayed ascending excitatory and inhibitory inputs. Bicuculline application locally in ICC greatly reduces or eliminates the response selectivity of duration-tuned neurons (Casseday et al. 1994, 2000).

Likewise, many cells in the pallid bat IC are duration tuned. Bicuculline often abolishes duration selectivity, emphasizing the importance of local  $GABA$  receptors. However, in some cells duration selectivity is not abolished by bicu-

culline. A revised coincidence model was proposed to explain the response properties of these duration-tuned cells (Fuzessery and Hall 1999).

#### 4.6. GLYCINERGIC INHIBITION

Although both anatomical and brain slice studies show that there are glycinergic inputs to the ICC, only a few studies have examined in detail how glycine receptors shape responses to acoustic stimulation. Both GABA and glycine contributed to binaural responses in the mustached bat's ICC (Klug et al. 1995) and their respective antagonists, bicuculline and strychnine, were each effective in partially blocking binaural inhibition (see also Vater et al. 1992a). Furthermore, glycinergic inhibition influenced responses to FM stimuli much like GABA, but less strongly (Koch and Grothe 1998, 2000). Strychnine altered frequency tuning curves in guinea pig IC (Le Beau et al. 2001; see Section 4.2) and affected duration tuning of IC neurons in the big brown bat much like bicuculline, but more weakly (Casseday et al. 1994, 2000). The greater efficacy of the GABA<sub>A</sub> receptor blockade when compared with glycine receptor blockade is consistent with brain slice and binding studies of the afferent projections to ICC, which show a GABA<sub>A</sub> receptor dominance (Zarbin et al. 1981; Milbrandt et al. 1996; Moore et al. 1998; Ma et al. 2002c). However, the relative expression of GABAergic and glycinergic inhibition might be related to the recording site location within the IC (see Chapters 2 and 5).

#### 4.7. ROLE OF THE GABA<sub>B</sub> RECEPTOR

The presence of GABA<sub>B</sub> receptors in the IC is supported by the actions of the GABA<sub>B</sub> receptor agonist, baclofen, and the receptor antagonist, phaclofen (Faingold et al. 1989a, 1991a), as well as receptor binding (Milbrandt et al. 1994, 1996) and brain slice (Zhang and Wu 2000; Ma et al. 2002c) studies. Baclofen reduced firing moderately and phaclofen slightly increased responses to acoustic stimulation (Faingold et al. 1989a, 1991a; see also Burger and Pollak 1998). The reduced firing produced by baclofen, however, cannot be attributed with certainty to a postsynaptic inhibitory action. Indeed, there is no indication of a postsynaptic effect of the GABA<sub>B</sub> agonist baclofen in a brain slice preparation of the rat's IC. Rather, baclofen had an effect on GABAergic inhibition through presynaptic modulation of inhibitory transmitter release (Ma et al. 2002c; see Fig. 9.2).

Activation of GABA<sub>B</sub> receptors can also reduce the strength of synaptic excitation in the rat's IC as shown by dose-dependent suppressive effects of baclofen on excitatory postsynaptic currents recorded in brain slices with whole-cell patch clamp techniques (Wu et al. 2004). Thus, the decreased response to acoustic stimulation produced by baclofen *in vivo* might be due to a presynaptic GABA<sub>B</sub> receptor-mediated reduction in excitatory transmitter release rather than postsynaptic inhibition.

#### 4.8. SEROTONIN

Serotonin [5-hydroxytryptamine (5-HT)] has a modulatory effect on the responses of many neurons in the ICC (see Hurley et al. 2002). Simple and complex response properties were altered by serotonin application (Hurley and Pollak 1999, 2001), which reduced tone-evoked responses in 60% of the IC neurons in the Mexican free-tailed bat. The firing decrease was often uniform across frequency, although some frequency-specific effects did occur. The role of serotonin in arousal and attention to particular sounds remains to be examined. Further studies with specific serotonin receptor antagonists are needed to determine the receptor specificity of the effects, and behavioral studies are called for to assess the extent to which the action of serotonin in IC depends on the animal's central state.

### 5. MODIFICATION OF SYNAPTIC FUNCTION

#### 5.1. EFFECTS OF DECREASED PERIPHERAL ACOUSTIC INPUT ON NORMAL INHIBITORY FUNCTION

Studies of animals exposed to extremely high levels of noise show increased IC evoked potential amplitude and increased single-unit activity to suprathreshold acoustic stimuli following exposure (Willott and Lu 1982; Salvi et al. 1990; Gerken et al. 1991; Szczepaniak and Møller 1995; Wang et al. 1996; Syka and Rybalko 2000; see Chapters 19 and 20). These findings suggest a global loss of synaptic inhibition, as acoustic nerve and VCN show reduced responses to suprathreshold stimuli after noise exposure (Gerken et al. 1991). Damage to the auditory periphery evokes a selective down-regulation of normal inhibitory GABAergic IC function. A loss of bicuculline sensitivity is seen in IC surface-recorded evoked potentials from noise-exposed animals (Szczepaniak and Møller 1995). Bilateral deafness decreases in vivo GABA release and fewer neurons show electrically evoked suppression of activity (Bledsoe et al. 1995). GAD levels were reduced 2 to 30 days following noise exposure (Abbott et al. 1999; Milbrandt et al. 2000) and at 24 hours and 7 days after cochleotomy (Mossop et al. 2000). Studies of GABA uptake and release following ossicle removal or cochlear ablation reveal complex changes in GABA and glycine neurochemistry (Suneja et al. 1998). Collectively, these studies show significant changes in normal GABA function after peripheral manipulations that decrease acoustic input.

#### 5.2. AGE-RELATED CHANGES IN THE INFERIOR COLLICULUS

It is unknown whether age-related changes in GABA neurotransmission reflect *de novo* adjustment in the central auditory system or embody a gradual loss of peripheral input, or both. A full discussion is beyond the scope of this review (see Chapter 19). However, the significant decreases in endogenous IC GABA



and GABA release are not evident in CN (Banay-Schwartz et al. 1989a,b; Caspary et al. 1990; Raza et al. 1994). GABA levels, GAD levels, and GABA<sub>B</sub> receptor binding each decrease with age, and there is neuropil rearrangement in aging Fischer 344 (F344) rats (Banay-Schwartz et al. 1989a,b; Caspary et al. 1990; Gutiérrez et al. 1994; Milbrandt et al. 1994, 1996; Raza et al. 1994; Helfert et al. 1999). Comparison of single-unit recordings from ICC in anesthetized young (3 to 4 months) and aged (24 months) F344 rats shows significantly more monotonic ICC rate-intensity responses in the aged animals, and an 18% increase in the breadth of isointensity functions at 30 dB above threshold (Palombi and Caspary 1996a–c). Analogous physiologic changes occur in C57 and CBA mice (Willott 1986; Willott et al. 1988a,b, 1991; McFadden and Willott 1994; Walton et al. 1998).

### 5.3. AGE-RELATED GABA<sub>A</sub> RECEPTOR SUBUNIT CHANGES

The GABA<sub>A</sub> receptor is a heterogeneous family of ligand-gated Cl<sup>-</sup> ion channel receptors that generally mediate inhibition in the adult brain. GABA<sub>A</sub> receptors likely exist as pentameric subunit complexes that can be allosterically modulated by numerous pharmacologic agents (Sieghart 1992a,b, 1995; Wafford et al. 1993; Rabow et al. 1995; Mohler et al. 2002). Molecular cloning reveals the presence of 6- $\alpha$ , 4- $\beta$ , 3- $\gamma$ , 1- $\delta$ , and 2- $\epsilon$  GABA<sub>A</sub> receptor subunits (Macdonald and Olsen 1994; Rabow et al. 1995; Sieghart 1995). Specific subunits have regionally selective distributions in several species (Burt and Kamatchi 1991; Luddens and Wisden 1991; Doble and Martin 1992; Persohn et al. 1992; Poulter et al. 1992; Wisden et al. 1992; Wynne et al. 1995; McKernan and Whiting 1996). It is thought that as few as 10 combinations exist in significant numbers in the CNS (McKernan and Whiting 1996). In situ hybridization studies describe age-related changes in GABA<sub>A</sub> receptor subunit composition in rat IC (Milbrandt et al. 1997), with significant age-related increases in  $\alpha_2$  and  $\gamma_1$  subunit mRNA along with small declines in  $\alpha_1$ ,  $\beta_2$ , and  $\gamma_2$  subunit mRNA in two strains (Caspary et al. 1999). Coexpression of the  $\gamma_1$  subunit with  $\alpha_1$  and  $\beta_2$  subunits in oocytes produces a GABA<sub>A</sub> receptor complex with more sensitivity to GABA than receptors with the  $\gamma_2$  subunit (Ducic et al. 1995), and binding studies find significant age-related enhancement of GABA modulation of binding at the picrotoxin GABA<sub>A</sub> receptor site (Milbrandt et al. 1996). These findings suggest altered GABA<sub>A</sub> receptor pharmacology in aged animals. Such changes seem to differ in the three main IC subregions (Milbrandt et al. 1997). Altered GABA<sub>A</sub> receptor expression levels are supported by immunocytochemical measures of protein levels for selected GABA<sub>A</sub> receptor subunits, with significant increases in  $\gamma_1$  subunit protein and a decrease in  $\alpha_1$  subunit protein with age (Caspary et al. 1999).

A direct functional measure of this subunit change was made by examining the ability of GABA to flux Cl<sup>-</sup> ions in microsac/synaptosome preparations from rat IC. The GABA-mediated influx was significantly increased in samples from

aged animals (Caspary et al. 1999). These findings contrast with previous whole-brain synaptosome chloride uptake studies, which found reduced  $\text{Cl}^-$  uptake with aging (Concas et al. 1988). The  $\text{Cl}^-$  flux findings strongly support an age-related change in  $\text{GABA}_A$  receptor subunit composition likely resulting in altered coding in the IC of aged animals.  $\text{GABA}_A$  receptor changes may reflect a compensatory up-regulation of postsynaptic inhibitory function after significant loss of presynaptic GABA release.

#### 5.4. QUANTITATIVE RECEPTOR AUTORADIOGRAPHY

There is significant muscimol binding in the IC (Glendenning and Baker 1988), and several ligands related to normal  $\text{GABA}_{A \text{ and } B}$  receptor pharmacology bind with high affinity in the IC of young and aged animals (Milbrandt et al. 1994, 1996). No significant age-related changes in IC  $\text{GABA}_A$  receptor binding are found with single concentrations of tritiated  $\text{GABA}_A$  receptor ligands (muscimol, TBOB, and flunitrazepam) (Milbrandt et al. 1996). However, as noted earlier, modulation of binding at the picrotoxin site with different concentrations of bath-applied GABA caused a dose-dependent shift to the left in the GABA modulation curve, indicating an age-related increase in the potency of GABA inhibition at this site. In contrast to the complex changes at the  $\text{GABA}_A$  receptor,  $\text{GABA}_B$  binding was significantly reduced in the ICs of aged rats, likely indicative of the loss of normal presynaptic GABA function (Milbrandt et al. 1994).

## 6. CONCLUSIONS AND DIRECTIONS FOR FUTURE RESEARCH

It is evident that the excitatory and inhibitory amino acids and their respective receptors are critically involved in shaping the response properties of neurons in the three subdivisions of the IC. Extrinsic projections and intrinsic circuits releasing glutamate and GABA are decisive in determining the activity of neurons projecting from the IC to the medial geniculate body and auditory forebrain. Glycine also plays a role in shaping responses to acoustic stimulation, and other neurotransmitters or neuromodulators, such as serotonin, can influence the responses of IC neurons. The coding properties of IC cells are determined by the relative balance and timing of their excitatory and inhibitory inputs as well as their inherent characteristics, for example, membrane properties and biophysics. The amino acid neurotransmitter systems and the other modulators show plastic changes as the acoustic input changes over time with age or with selective peripheral damage. Such changes appear to be a compensatory increase in gain, ensuing from a loss of excitatory input, that remodels the relative balance of excitation and inhibition centrally. It remains to be determined whether these plastic changes are reversible or amenable to further adjustment using pharmacologic probes.

An important goal for future research is the identification of specific cell types within the IC that receive projections from various sources and the further characterization of their receptor pharmacology and circuitry. For example, do different cell types in IC show variations in the proportion of AMPA and NMDA receptors that might determine the gain of excitatory responses? Do the GABAergic and glycinergic projections from subcollicular auditory nuclei target different IC cell types, and do these transmitters serve the same physiologic functions? Do the ipsilateral and contralateral DNLL projections make synaptic contact with the same or different ICC neuronal populations? How do the ascending projections from cochlear nucleus, superior olive, and lateral lemniscal neurons interact with commissural projections or descending projections from the medial geniculate or cortex? A major challenge for future research will be to provide definitive answers to these intriguing questions.

## Acknowledgments

This research was supported by grants from the Natural Sciences and Engineering Research Council of Canada (J.B.K.), the Hearing Foundation of Canada (J.B.K.), and the United States Public Health Service (D.M.C., Grant RO1 DC000151). We thank Drs. Shu Hui Wu and Huiming Zhang for their helpful contributions and Ms. Judy Bryan for her dedication in preparing the manuscript for publication.

## Abbreviations

AM	amplitude-modulated
AMPA	$\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid
APV	( $\pm$ )-2-amino-5-phosphonovaleric acid
CF	characteristic frequency
CN	cochlear nucleus
CNQX	6-cyano-7-nitroquinoxaline-2,3-dione
CPP	( $\pm$ )-3-(2-carboxypiperazin-4-yl)-propyl-1-phosphonic acid
DCN	dorsal cochlear nucleus
EPSC	excitatory postsynaptic current
EPSP	excitatory postsynaptic potential
FM	frequency-modulated
GABA	$\gamma$ -aminobutyric acid
GAD	glutamic acid decarboxylase
IC	inferior colliculus
ICC	central nucleus of the inferior colliculus
ICD	dorsal cortex <i>or</i> dorsal nucleus of the inferior colliculus
ICX	external nucleus of the inferior colliculus
IPSP	inhibitory postsynaptic potential

LSO	lateral superior olive
MSO	medial superior olive
MTF <sub>FR</sub>	modulation transfer function for firing rate
MTF <sub>VS</sub>	modulation transfer function for vector strength
NBQX	1,2,3,4-tetrahydro-6-nitro-2,3-dioxo-benzo[f]quinoxaline-7-sulfonamide disodium
NMDA	<i>N</i> -methyl-D-aspartate
PSTH	post-stimulus time histogram
RLF	rate level function
TBOB	<i>t</i> -[ <sup>3</sup> H]butylbicycloorthobenzoate
VCN	ventral cochlear nucleus

## References

- Abbott SD, Hughes LF, Bauer CA, Salvi R, and Caspary DM (1999) Detection of glutamate decarboxylase isoforms in rat inferior colliculus following acoustic exposure. *Neuroscience* **93**:1375–1381.
- Adams JC (1979) Ascending projections to the inferior colliculus. *Journal of Comparative Neurology* **183**:519–538.
- Adams JC and Mugnaini E (1984) Dorsal nucleus of the lateral lemniscus: a nucleus of GABAergic projection neurons. *Brain Research Bulletin* **13**:585–590.
- Adams JC and Wenthold RJ (1979) Distribution of putative amino acid transmitters, choline acetyltransferase and glutamate decarboxylase in the inferior colliculus. *Neuroscience* **4**:947–951.
- Bajo VM, Merchán MA, Malmierca MS, Nodal FR, and Bjaalie JG (1999) Topographic organization of the dorsal nucleus of the lateral lemniscus in the cat. *Journal of Comparative Neurology* **407**:349–366.
- Banay-Schwartz M, Lajtha A, and Palkovits M (1989a) Changes with aging in the levels of amino acids in rat CNS structural elements. I. Glutamate and related amino acids. *Neurochemical Research* **14**:555–562.
- Banay-Schwartz M, Lajtha A, and Palkovits M (1989b) Changes with aging in the levels of amino acids in rat CNS structural elements. II. Taurine and small neutral amino acids. *Neurochemical Research* **14**:563–570.
- Bear MF and Malenka RC (1994) Synaptic plasticity: LTP and LTD. *Current Opinion in Neurobiology* **4**:389–399.
- Bergman M, Staatz-Benson C, and Potashner SJ (1989) Amino acid uptake and release in the guinea pig cochlear nucleus after inferior colliculus ablation. *Journal of Neurophysiology* **42**:283–291.
- Bledsoe SC Jr, Nagase S, Miller JM, and Altschuler RA (1995) Deafness-induced plasticity in the mature central auditory system. *NeuroReport* **7**:225–229.
- Bliss TVP and Collingridge GL (1993) A synaptic model of memory: long-term potentiation in the hippocampus. *Nature* **361**:31–39.
- Bormann J (2000) The ‘ABC’ of GABA receptors. *Trends in Pharmacological Sciences* **21**:16–19.
- Brand A, Urban R, and Grothe B (2000) Duration tuning in the mouse auditory midbrain. *Journal of Neurophysiology* **84**:1790–1799.

- Brenowitz S, David J, and Trussell L (1998) Enhancement of synaptic efficacy by pre-synaptic GABA<sub>B</sub> receptors. *Neuron* **20**:135–141.
- Burger RM and Pollak GD (1998) Analysis of the role of inhibition in shaping responses to sinusoidally amplitude-modulated signals in the inferior colliculus. *Journal of Neurophysiology* **80**:1686–1701.
- Burt DR and Kamatchi GL (1991) GABA<sub>A</sub> receptor subtypes: from pharmacology to molecular biology. *Federation of American Society for Experimental Biology Journal* **5**:2916–2923.
- Carr CE, Fujita I, and Konishi M (1989) Distribution of GABAergic neurons and terminals in the auditory system of the barn owl. *Journal of Comparative Neurology* **286**:190–207.
- Caspary DM, Raza A, Armour BAL, Phippen J, and Arnerić SP (1990) Immunocytochemical and neurochemical evidence for age related loss of GABA in the inferior colliculus: implications for neural presbycusis. *Journal of Neuroscience* **10**:2363–2372.
- Caspary DM, Holder TM, Hughes LF, Milbrandt JC, McKernan RM, and Naritoku DK (1999) Age-related changes in GABA<sub>A</sub> receptor subunit composition and function in rat auditory system. *Neuroscience* **93**:307–312.
- Caspary DM, Palombi PS, and Hughes LF (2002) GABAergic inputs shape responses to amplitude modulated stimuli in the inferior colliculus. *Hearing Research* **168**:163–173.
- Casseday JH, Covey E, and Vater M (1988) Connections of the superior olivary complex in the rufous horseshoe bat *Rhinolophus rouxi*. *Journal of Comparative Neurology* **278**:313–329.
- Casseday JH, Ehrlich D, and Covey E (1994) Neural tuning for sound duration: role of inhibitory mechanisms in the inferior colliculus. *Science* **264**:847–850.
- Casseday JH, Ehrlich D, and Covey E (2000) Neural measurement of sound duration: control by excitatory-inhibitory interactions in the inferior colliculus. *Journal of Neurophysiology* **84**:1475–1487.
- Casseday JH, Fremouw T, and Covey E (2002) The inferior colliculus: a hub for the central auditory system. In: Oertel D, Fay RR, and Popper AN (eds). *Springer Handbook of Auditory Research*, Volume 15: *Integrative Functions in the Mammalian Auditory Pathway*. Springer-Verlag, New York, pp. 238–318.
- Chebib M and Johnston GA (1999) The ‘ABC’ of GABA receptors: a brief review. *Clinical Experimental Pharmacology and Physiology* **26**:937–940.
- Chen GD (1998) Effects of stimulus duration on responses of neurons in the chinchilla inferior colliculus. *Hearing Research* **122**:142–150.
- Chen L, Kelly JB, and Wu SH (1999) The commissure of Probst as a source of GABAergic inhibition. *Hearing Research* **138**:106–114.
- Concas A, Pepitoni S, Atsoggiu T, Toffano G, and Biggio G (1988) Aging reduces the GABA-dependent <sup>36</sup>Cl<sup>-</sup> flux in rat brain membrane vesicles. *Life Sciences* **43**:1761–1771.
- Contreras NEIR and Bachelard HS (1979) Some neurochemical studies on auditory regions of mouse brain. *Experimental Brain Research* **36**:573–584.
- Davis KA (2002) Evidence of a functionally segregated pathway from dorsal cochlear nucleus to inferior colliculus. *Journal of Neurophysiology* **87**:1824–1835.
- Davis KA, Ramachandran R, and May BJ (2003) Auditory processing of spectral cues for sound localization in the inferior colliculus. *Journal of the Association for Research in Otolaryngology* **4**:148–163.

- Doble A and Martin IL (1992) Multiple benzodiazepine receptors: no reason for anxiety. *Trends in Pharmacological Sciences* **13**:76–81.
- Ducic I, Caruncho HJ, Zhu WJ, Vicini S, and Costa E (1995).  $\gamma$ -Aminobutyric acid gating of  $\text{Cl}^-$  channels in recombinant  $\text{GABA}_A$  receptors. *Journal of Pharmacology and Experimental Therapeutics* **272**:438–445.
- Faingold CL, Gehlbach G, and Caspary DM (1989a) On the role of GABA as an inhibitory neurotransmitter in inferior colliculus neurons: iontophoretic studies. *Brain Research* **500**:302–312.
- Faingold CL, Hoffmann WE, and Caspary DM (1989b) Effects of excitant amino acids on acoustic responses of inferior colliculus neurons. *Hearing Research* **40**:127–136.
- Faingold CL, Boersma Anderson CA, and Caspary DM (1991a) Involvement of GABA in acoustically-evoked inhibition in inferior colliculus neurons. *Hearing Research* **52**: 201–216.
- Faingold CL, Gehlbach G, and Caspary DM (1991b) Functional pharmacology of inferior colliculus neurons. In: Altschuler RA, Bobbin RP, Clopton BM, and Hoffman DW (eds). *Neurobiology of Hearing: The Central Auditory System*. Raven Press, New York, pp. 223–251.
- Faingold CL, Anderson CAB, and Randall ME (1993) Stimulation or blockade of the dorsal nucleus of the lateral lemniscus alters binaural and tonic inhibition in contralateral inferior colliculus neurons. *Journal of Neurophysiology* **69**:98–106.
- Feldman DE and Knudsen EI (1994) NMDA and non-NMDA glutamate receptors in auditory transmission in the barn owl inferior colliculus. *Journal of Neuroscience* **14**: 5939–5958.
- Feldman DE, Brainard MS, and Knudsen EI (1996) Newly learned auditory responses mediated by NMDA receptors in the owl inferior colliculus. *Science*. **271**:525–528.
- Fisher SK and Davies WE (1976) GABA and its related enzymes in the lower auditory system of the guinea pig. *Journal of Neurochemistry* **27**:1145–1155.
- Foeller E, Vater M, and Kössl M (2001) Laminar analysis of inhibition in the gerbil primary auditory cortex. *Journal of the Association for Research in Otolaryngology* **2**:279–296.
- Fubara BM, Casseday JH, Covey E, and Schwartz-Bloom RD (1996) Distribution of  $\text{GABA}_A$ ,  $\text{GABA}_B$ , and glycine receptors in the central auditory system of the big brown bat, *Eptesicus fuscus*. *Journal of Comparative Neurology* **369**:83–92.
- Fujita I and Konishi M (1991) The role of GABAergic inhibition in processing of interaural time difference in the owls auditory system. *Journal of Neuroscience* **11**:722–739.
- Fuzessery ZM and Hall JC (1996) Role of GABA in shaping frequency tuning and creating FM sweep selectivity in the inferior colliculus. *Journal of Neurophysiology* **76**:1059–1073.
- Fuzessery ZM and Hall JC (1999) Sound duration selectivity in the pallid bat inferior colliculus. *Hearing Research* **137**:137–154.
- Gerken GM, Solecki JM, and Boettcher FA (1991) Temporal integration of electrical stimulation of auditory nuclei in normal-hearing and hearing-impaired cat. *Journal of Neurophysiology* **53**:101–112.
- Glendenning KK and Baker BN (1988) Neuroanatomical distribution of receptors for three potential inhibitory neurotransmitters in the brainstem auditory nuclei of the cat. *Journal of Comparative Neurology* **275**:288–308.

- Glendenning KK, Baker BN, Hutson KA, and Masterton RB (1992) Acoustic chiasm V: inhibition and excitation in the ipsilateral and contralateral projections of LSO. *Journal of Comparative Neurology* **319**:100–122.
- González-Hernández T, Mantolán-Sarmiento B, González-González B, and Pérez-González H (1996) Sources of GABAergic input to the inferior colliculus of the rat. *Journal of Comparative Neurology* **372**:309–326.
- Gutiérrez A, Khan ZU, Morris SJ, and De Blas AL (1994) Age-related decrease of GABA<sub>A</sub> receptor subunits and glutamic acid decarboxylase in the rat inferior colliculus. *Journal of Neuroscience* **14**:7469–7477.
- Helfert RH, Bonneau JM, Wenthold RJ, and Altschuler RA (1989) GABA and glycine immunoreactivity in the guinea pig superior olivary complex. *Brain Research* **501**:269–286.
- Helfert RH, Snead CR, and Altschuler RA (1991) The ascending auditory pathways. In: Altschuler RA, Bobbin RP, Clopton BM, and Hoffman DW (eds). *Neurobiology of Hearing: The Central Auditory System*. Raven Press, New York, pp. 1–25.
- Helfert RH, Sommer TJ, Meeks J, Hofstetter P, and Hughes LF (1999) Age-related synaptic changes in the central nucleus of the inferior colliculus of Fischer-344 rats. *Journal of Comparative Neurology* **406**:285–298.
- Huffman RF and Henson OW Jr (1990) The descending auditory pathway and acousticomotor systems: connections with the inferior colliculus. *Brain Research Reviews* **15**:295–323.
- Hurley LM and Pollak GD (1999) Serotonin differentially modulates responses to tones and frequency-modulated sweeps in the inferior colliculus. *Journal of Neuroscience* **19**:8071–8082.
- Hurley LM and Pollak GD (2001) Serotonin effects on frequency tuning of inferior colliculus neurons. *Journal of Neurophysiology* **85**:828–842.
- Hurley LM, Thompson AM, and Pollak GD (2002) Serotonin in the inferior colliculus. *Hearing Research* **168**:1–11.
- Jen PH-S, Sun X, and Chen QC (2001) An electrophysiological study of neural pathways for corticofugally inhibited neurons in the central nucleus of the inferior colliculus of the big brown bat, *Eptesicus fuscus*. *Experimental Brain Research* **137**:292–302.
- Kelly JB and Kidd SA (2000) NMDA and AMPA receptors in the dorsal nucleus of the lateral lemniscus shape binaural responses in rat inferior colliculus. *Journal of Neurophysiology* **83**:1403–1414.
- Kelly JB and Li L (1997) Two sources of inhibition affecting binaural evoked responses in the rat's inferior colliculus: the dorsal nucleus of the lateral lemniscus and the superior olivary complex. *Hearing Research* **104**:112–126.
- Kelly JB and Zhang H (2002) Contribution of AMPA and NMDA receptors to excitatory responses in the inferior colliculus. *Hearing Research* **168**:35–42.
- Kidd SA and Kelly JB (1996) Contribution of the dorsal nucleus of the lateral lemniscus to binaural responses in the inferior colliculus of the rat: interaural time delays. *Journal of Neuroscience* **16**:7390–7397.
- Klug A, Park TJ, and Pollak GD (1995) Glycine and GABA influence binaural processing in the inferior colliculus of the mustache bat. *Journal of Neurophysiology* **74**:1701–1713.
- Koch U and Grothe B (1998) GABAergic and glycinergic inhibition sharpens tuning for frequency modulations in the inferior colliculus of the big brown bat. *Journal of Neurophysiology* **80**:71–82.

- Koch U and Grothe B (2000) Interdependence of spatial and temporal coding in the auditory midbrain. *Journal of Neurophysiology* **83**:2300–2314.
- Krishna BS and Semple MN (2000) Auditory temporal processing: responses to sinusoidally amplitude-modulated tones in the inferior colliculus. *Journal of Neurophysiology* **84**:255–273.
- Kulesza RJ Jr, Spirou GA, and Berrebi AS (2003) Physiological response properties of neurons in the superior paraolivary nucleus of the rat. *Journal of Neurophysiology* **89**:2299–2312.
- Le Beau FEN, Rees A, and Malmierca MS (1996) Contribution of GABA- and glycine-mediated inhibition to the monaural temporal response properties of neurons in the inferior colliculus. *Journal of Neurophysiology* **75**:902–919.
- Le Beau FEN, Malmierca MS, and Rees A (2001) Iontophoresis *in vivo* demonstrates a key role for GABA<sub>A</sub> and glycinergic inhibition in shaping frequency response areas in the inferior colliculus of guinea pig. *Journal of Neuroscience* **21**:7303–7312.
- Li L and Kelly JB (1992) Inhibitory influence of the dorsal nucleus of the lateral lemniscus on binaural responses in the rat's inferior colliculus. *Journal of Neuroscience* **12**:4530–4539.
- Li Y, Evans MS, and Faingold CL (1998) *In vitro* electrophysiology of neurons in subnuclei of rat inferior colliculus. *Hearing Research* **121**:1–10.
- Li Y, Evans MS, and Faingold CL (1999) Synaptic response patterns of neurons in the cortex of rat inferior colliculus. *Hearing Research* **137**:15–28.
- Lim D, Oh S, Chang SO, and Kim CS (2000) Auditory responses of field L to electrical stimulation. *Advances in Otorhinolaryngology* **57**:15–17.
- López-Colomé AM, Tapia R, Saleda R, and Pasantes-Morales H (1978) K<sup>+</sup>-stimulated release of labeled  $\gamma$ -aminobutyrate, glycine and taurine in slices of several regions of rat central nervous system. *Neuroscience* **3**:1069–1074.
- Luddens H and Wisden W (1991) Function and pharmacology of multiple GABA<sub>A</sub> receptor subunits. *Trends in Pharmacological Sciences* **12**:49–51.
- Ma CL, Kelly JB, and Wu SH (2002a) AMPA and NMDA receptors mediate synaptic excitation in the rat's inferior colliculus. *Hearing Research* **168**:25–34.
- Ma CL, Kelly JB, and Wu SH (2002b) Physiological properties of NMDA receptors in the rat's inferior colliculus. *Association for Research in Otolaryngology Abstracts* **25**:174.
- Ma CL, Kelly JB, and Wu SH (2002c) Presynaptic modulation of GABAergic inhibition by GABA<sub>B</sub> receptors in the rat's inferior colliculus. *Neuroscience* **114**:207–215.
- Macdonald RL and Olsen RW (1994) GABA<sub>A</sub> receptor channels. *Annual Review of Neuroscience* **17**:569–602.
- Malmierca MS, Le Beau FEN, and Rees A (1996) The topographical organization of descending projections from the central nucleus of the inferior colliculus in guinea pig. *Hearing Research* **93**:167–180.
- Marianowski R, Liao WH, Van Den Abbeele T, Fillit P, Herman P, Frachet B, and Huy PT (2000) Expression of NMDA, AMPA and GABA<sub>A</sub> receptor subunit mRNAs in the rat auditory brainstem. I. Influence of early auditory deprivation. *Hearing Research* **150**:1–11.
- McAlpine D and Palmer AR (2002) Blocking GABAergic inhibition increases sensitivity to sound motion cues in the inferior colliculus. *Journal of Neuroscience* **22**:1443–1453.
- McAlpine D, Jiang D, Shackleton TM, and Palmer AR (2000) Responses of neurons in the inferior colliculus to dynamic interaural phase cues: evidence for a mechanism of binaural adaptation. *Journal of Neurophysiology* **83**:1356–1365.



- McFadden SL and Willott JF (1994) Responses of inferior colliculus neurons in C57BL/6J mice with and without sensorineural hearing loss: effects of changing the azimuthal location of an unmasked pure-tone stimulus. *Journal of Neurophysiology* **78**:115–131.
- McKernan RM and Whiting PJ (1996) Which GABA<sub>A</sub>-receptor subtypes really occur in the brain? *Trends in Neurosciences* **19**:139–143.
- Merchán MA, Saldaña E, and Plaza I (1994) Dorsal nucleus of the lateral lemniscus in the rat: concentric organization and tonotopic projection to the inferior colliculus. *Journal of Comparative Neurology* **342**:259–278.
- Milbrandt JC, Albin RL, and Caspary DM (1994) Age-related decrease in GABA<sub>B</sub> receptor binding in the Fischer-344 rat inferior colliculus. *Neurobiology of Aging* **15**:699–703.
- Milbrandt JC, Albin RL, Turgeon SM, and Caspary DM (1996) GABA<sub>A</sub> receptor binding in the aging rat inferior colliculus. *Neuroscience* **73**:449–458.
- Milbrandt JC, Hunter C, and Caspary DM (1997) Alterations of GABA<sub>A</sub> receptor subunit mRNA levels in the aging Fischer 344 rat inferior colliculus. *Journal of Comparative Neurology* **379**:455–465.
- Milbrandt JC, Holder TM, Wilson MC, Salvi RJ, and Caspary DM (2000) GAD levels and muscimol binding in rat inferior colliculus following acoustic trauma. *Hearing Research* **147**:251–260.
- Misgeld U, Bijak M, and Jarolimek W (1995) A physiological role for GABA<sub>B</sub> receptors and the effects of baclofen in the mammalian central nervous system. *Progress in Neurobiology* **46**:423–462.
- Mohler H, Fritschy JM, and Rudolph U (2002) A new benzodiazepine pharmacology. *Journal of Pharmacology and Experimental Therapeutics* **300**:2–8.
- Moore DR, Kotak VC, and Sanes DH (1998) Commissural and lemniscal synaptic input to the gerbil inferior colliculus. *Journal of Neurophysiology* **80**:2229–2236.
- Moore JK and Moore RY (1987) Glutamic acid decarboxylase-like immunoreactivity in brainstem auditory nuclei of the rat. *Journal of Comparative Neurology* **260**:157–174.
- Mossop JE, Wilson MJ, Caspary DM, and Moore DR (2000). Down-regulation of inhibition following unilateral deafening. *Hearing Research* **147**:183–187.
- Mugnaini E and Oertel WH (1985) An atlas of the distribution of GABAergic neurons and terminals in the rat CNS as revealed by GAD immunohistochemistry. In: Björklund A and Hökfelt T (eds). *Handbook of Chemical Neuroanatomy*, Volume 4: *GABA and Neuropeptides in the CNS*, Part I, Elsevier, Amsterdam, pp. 436–608.
- Nagai T, Maeda T, Imai H, McGeer PL, and McGeer EG (1985) Distribution of GABA-T-intensive neurons in the rat hindbrain. *Journal of Comparative Neurology* **231**:260–269.
- Oliver DL, Winer JA, Beckius GE, and Saint Marie RL (1994) Morphology of GABAergic neurons in the inferior colliculus of the cat. *Journal of Comparative Neurology* **340**:27–42.
- Ostapoff E-M, Benson CG, and Saint Marie RL (1997) GABA- and glycine-immunoreactive projections from the superior olivary complex to the cochlear nucleus in guinea pig. *Journal of Comparative Neurology* **381**:500–512.
- Palombi PS and Caspary DM (1996a) GABA inputs control discharge rate primarily within frequency receptive fields of inferior colliculus neurons. *Journal of Neurophysiology* **75**:2211–2219.
- Palombi PS and Caspary DM (1996b) Responses of young and aged Fischer 344 rat inferior colliculus neurons to binaural tonal stimuli. *Hearing Research* **100**:59–67.

- Palombi PS and Caspary DM (1996c) Physiology of the aged Fischer 344 rat inferior colliculus: responses to contralateral monaural stimuli. *Journal of Neurophysiology* **76**: 3114–3125.
- Park TJ and Pollak GD (1993) GABA shapes a topographic organization of response latency in the mustache bat's inferior colliculus. *Journal of Neuroscience* **13**:5172–5187.
- Park TJ and Pollak GD (1994) Azimuthal receptive fields are shaped by GABAergic inhibition in the inferior colliculus of the mustache bat. *Journal of Neurophysiology* **72**:1080–1102.
- Parks TN (2000) The AMPA receptors of auditory neurons. *Hearing Research* **147**:77–91.
- Persohn E, Malherbe P, and Richards JG (1992) Comparative molecular neuroanatomy of cloned GABA<sub>A</sub> receptor subunits in the rat CNS. *Journal of Comparative Neurology* **326**:193–216.
- Pierson MG, Smith KL, and Swann JW (1989) A slow NMDA-mediated synaptic potential underlies seizures originating from midbrain. *Brain Research* **486**:381–386.
- Potashner SJ (1979) Baclofen: effects on amino acid release and metabolism in slices of guinea pig cerebral cortex. *Journal of Neurochemistry* **32**:103–109.
- Poulter MO, Barker JL, O'Carroll AM, Lolait SJ, and Mahan LC (1992) Differential and transient expression of GABA<sub>A</sub> subunit mRNAs in the developing rat CNS. *Journal of Neuroscience* **12**:2888–2900.
- Rabow LE, Russek SJ, and Farb DH (1995) From ion currents to genomic analysis: recent advances in GABA<sub>A</sub> receptor research. *Synapse* **21**:189–274.
- Raza A, Milbrandt JC, Arnerić SP, and Caspary DM (1994) Age-related changes in brainstem auditory neurotransmitters: measures of GABA and acetylcholine function. *Journal of Neurophysiology* **77**:221–230.
- Rees A and Møller AR (1983) Responses of neurons in the inferior colliculus of the rat to AM and FM tones. *Hearing Research* **10**:301–330.
- Reetz G and Ehret G (1999) Inputs from three brainstem sources to identified neurons of the mouse inferior colliculus slice. *Brain Research* **816**:527–543.
- Riquelme R, Saldaña E, Osen KK, Ottersen OP, and Merchán MA (2001) Colocalization of GABA and glycine in the ventral nucleus of the lateral lemniscus in rat: an in situ hybridization and semiquantitative immunocytochemical study. *Journal of Comparative Neurology* **432**:409–424.
- Roberts RC and Ribak CE (1987a) An electron microscopic study of GABAergic neurons and terminals in the central nucleus of the inferior colliculus of the rat. *Journal of Neurocytology* **16**:333–345.
- Roberts RC and Ribak CE (1987b) GABAergic neurons and axon terminals in the brainstem auditory nuclei of the gerbil. *Journal of Comparative Neurology* **258**:267–280.
- Romand R and Avan P (1997) Anatomical and functional aspects of the cochlear nucleus. In: Ehret G and Romand R (eds). *The Central Auditory System*. Oxford University Press, New York, pp. 97–192.
- Saint Marie RL (1996) Glutamatergic connections of the auditory midbrain: selective uptake and axonal transport of D-[<sup>3</sup>H]aspartate. *Journal of Comparative Neurology* **373**:255–270.
- Saint Marie RL, Ostapoff E-M, Morest DK, and Wenthold RJ (1989) Glycine-immunoreactive projection of the cat lateral superior olive: possible role in midbrain ear dominance. *Journal of Comparative Neurology* **279**:382–396.

- Saldaña E and Merchán MA (1992) Intrinsic and commissural connections of the rat inferior colliculus. *Journal of Comparative Neurology* **319**:417–437.
- Sally SL and Kelly JB (1992) Effects of superior olivary complex lesions on binaural responses in rat inferior colliculus. *Brain Research* **572**:5–18.
- Salvi RJ, Saunders SS, Gratton MA, Arehole S, and Powers N (1990) Enhanced evoked response amplitudes in the inferior colliculus of the chinchilla following acoustic trauma. *Hearing Research* **50**:245–257.
- Sanes DH, Malone BJ, and Semple MN (1998) Role of synaptic inhibition in processing of dynamic binaural level stimuli. *Journal of Neuroscience* **18**:794–803.
- Shiraishi S, Shiraishi Y, Oliver DL, and Altschuler RA (2001) Expression of GABA<sub>A</sub> receptor subunits in the rat central nucleus of the inferior colliculus. *Molecular Brain Research* **96**:122–132.
- Shneiderman A and Oliver DL (1989) EM autoradiographic study of the projections from the dorsal nucleus of the lateral lemniscus: a possible source of inhibitory inputs to the inferior colliculus. *Journal of Comparative Neurology* **286**:28–47.
- Shneiderman A, Oliver DL, and Henkel CK (1988) Connections of the dorsal nucleus of the lateral lemniscus: an inhibitory parallel pathway in the ascending auditory system? *Journal of Comparative Neurology* **276**:188–208.
- Shneiderman A, Chase MB, Rockwood JM, Benson CG, and Potashner SJ (1993) Evidence for a GABAergic projection from the dorsal nucleus of the lateral lemniscus to the inferior colliculus. *Journal of Neurochemistry* **60**:72–82.
- Sieghart W (1992a) GABA<sub>A</sub> receptors: ligand-gated Cl<sup>-</sup> ion channels modulated by multiple drug-binding sites. *Trends in Pharmacological Sciences* **13**:446–450.
- Sieghart W (1992b) Molecular basis of pharmacological heterogeneity of GABA<sub>A</sub> receptors. *Cellular Signalling* **4**:231–237.
- Sieghart W (1995) Structure and pharmacology of gamma-aminobutyric acid<sub>A</sub> receptor subtypes. *Pharmacological Reviews* **47**:181–234.
- Smith PH (1992) Anatomy and physiology of multipolar cells in the rat inferior collicular cortex using the in vitro brain slice technique. *Journal of Neuroscience* **12**:3700–3715.
- Spitzer MW and Semple MN (1998) Transformation of binaural response properties in the ascending auditory pathway: influence of time-varying interaural phase disparity. *Journal of Neurophysiology* **80**:3062–3076.
- Suneja SK, Potashner SJ, and Benson CG (1998) Plastic changes in glycine and GABA release and uptake in adult brainstem auditory nuclei after unilateral middle ear ossicle removal and cochlear ablation. *Experimental Neurology* **151**:273–288.
- Suneja SK, Potashner SJ, and Benson CG (2000) AMPA receptor binding in adult guinea pig brain stem auditory nuclei after unilateral cochlear ablation. *Experimental Neurology* **165**:355–369.
- Syka J and Rybalko N (2000) Threshold shifts and enhancement of cortical evoked responses after noise exposure in rats. *Hearing Research* **139**:59–68.
- Szczepaniak WS and Møller AR (1995) Evidence of decreased GABAergic influence on temporal integration in the inferior colliculus following acute noise exposure: a study of evoked potentials in the rat. *Neuroscience Letters* **196**:77–80.
- Tachibana M and Kuriyama K (1974) Gamma-aminobutyric acid in the lower auditory pathway of the guinea pig. *Brain Research* **69**:370–374.
- Thompson GC, Cortez AM, and Lam DM-K (1985) Localization of GABA immunoreactivity in the auditory brainstem of guinea pigs. *Brain Research* **339**:119–122.

- Turner JG, Hughes LF, and Caspary DM (2003) Response characteristics of layer V neurons in rat primary auditory cortex. *Association for Research in Otolaryngology Abstracts* **26**:52–53.
- van Adel BA, Kidd SA, and Kelly JB (1999) Contribution of the commissure of Probst to binaural evoked responses in the rat's inferior colliculus: interaural time differences. *Hearing Research* **130**:115–130.
- Vater M (1995) Ultrastructural and immunocytochemical observations on the superior olivary complex of the mustached bat. *Journal of Comparative Neurology* **358**:155–180.
- Vater M, Habbicht H, Kössl M, and Grothe B (1992a) The functional role of GABA and glycine in monaural and binaural processing in the inferior colliculus of horseshoe bats. *Journal of Comparative Physiology A* **171**:541–553.
- Vater M, Kössl M, and Horn AKE (1992b) GAD- and GABA-immunoreactivity in the ascending auditory pathway of horseshoe and mustached bats. *Journal of Comparative Neurology* **325**:183–206.
- Vater M, Covey E, and Casseday JH (1997) The columnar region of the ventral nucleus of the lateral lemniscus in the big brown bat (*Eptesicus fuscus*): synaptic arrangements and structural correlates of feedforward inhibitory function. *Cell and Tissue Research* **289**:223–233.
- Wafford KA, Bain CJ, Whiting PJ, and Kemp JA (1993) Functional comparison of the role of  $\gamma$  subunits in recombinant human  $\gamma$ -aminobutyric acid<sub>A</sub>/benzodiazepine receptors. *Molecular Pharmacology* **44**:437–442.
- Wagner T (1996) Lemniscal input to identified neurons of the central nucleus of mouse inferior colliculus: an intracellular brain slice study. *European Journal of Neuroscience* **8**:1231–1239.
- Walton JP, Frisina RD, and O'Neill WE (1998) Age-related alteration in processing of temporal sound features in the auditory midbrain of the CBA mouse. *Journal of Neuroscience* **18**:2764–2776.
- Wang J, Salvi RJ, and Powers N (1996) Plasticity of response properties of inferior colliculus neurons following acute cochlear damage. *Journal of Neurophysiology* **75**:171–183.
- Watanabe T and Simada Z (1973) Pharmacological properties of cat's collicular auditory neurons. *Japanese Journal of Physiology* **23**:291–308.
- Wenthold RJ (1991) Neurotransmitters of brainstem auditory nuclei. In: Altschuler RA, Bobbin RP, Clopton BM, and Hoffman DW (eds). *Neurobiology of Hearing: The Central Auditory System*. Raven Press, New York, pp. 121–139.
- Wenthold RJ, Hunter C, and Petralia RS (1993) Excitatory amino acid in the rat cochlear nucleus. In: Merchán MA, Juiz JS, Godfrey DA, and Mugnaini E (eds). *The Mammalian Cochlear Nuclei. Organization and Function*. Plenum Press, New York, pp. 179–194.
- Willott JF (1986) Effects of aging, hearing loss, and anatomical location on thresholds of inferior colliculus neurons in C57BL/6 and CBA mice. *Journal of Neurophysiology* **56**:391–408.
- Willott JF and Lu SM (1982) Noise-induced hearing loss can alter neural coding and increase excitability in the central nervous system. *Science* **216**:1331–1334.
- Willott JF, Parham K, and Hunter KP (1988a) Response properties of inferior colliculus neurons in middle-aged C57BL/6J mice with presbycusis. *Hearing Research* **37**:15–27.

- Willott JF, Hunter KP, and Coleman JR (1988b) Aging and presbycusis: effects on 2-deoxy-D-glucose uptake in the mouse auditory brain stem in quiet. *Experimental Neurology* **99**:615–621.
- Willott JF, Parham K, and Hunter KP (1991) Comparison of the auditory sensitivity of neurons in the cochlear nucleus and inferior colliculus of young and aging C57BL/6J and CBA/J mice. *Journal of Neurophysiology* **53**:78–94.
- Winer JA, Larue DT, and Pollak GD (1995) GABA and glycine in the central auditory system of the mustache bat: structural substrates for inhibitory neuronal organization. *Journal of Comparative Neurology* **355**:317–353.
- Wisden W, Laurie DJ, Monyer H, and Seeburg PH (1992) The distribution of 13 GABA<sub>A</sub> receptor subunit mRNAs in the rat brain. I: Telencephalon, diencephalon, mesencephalon. *Journal of Neuroscience* **12**:1040–1062.
- Wu SH, Ma CL, Sivaramakrishnan S, and Oliver DL (2002) Synaptic modification in neurons of the central nucleus of the inferior colliculus. *Hearing Research* **168**:43–54.
- Wu SH, Ma CL, and Kelly JB (2004) GABA<sub>B</sub> receptors modulate synaptic excitation in the rat's central and dorsal nuclei of the inferior colliculus. *Association for Research in Otolaryngology Abstracts* **27**:600.
- Wynne B, Harvey AR, Robertson D, and Sirinathsinghji DJS (1995) Neurotransmitter and neuromodulator systems of the rat inferior colliculus and auditory brainstem studied by *in situ* hybridization. *Journal of Chemical Neuroanatomy* **9**:289–300.
- Yang L, Pollak GD, and Resler C (1992) GABAergic circuits sharpen tuning curves and modify response properties in the mustache bat inferior colliculus. *Journal of Neurophysiology* **68**:1760–1774.
- Zarbin MA, Wamsley JK, and Kuhar MJ (1981) Glycine receptor: light microscopic autoradiographic localization with [<sup>3</sup>H]strychnine. *Journal of Neuroscience* **1**:532–547.
- Zhang DX, Li L, Kelly JB, and Wu SH (1998) GABAergic projections from the lateral lemniscus to the inferior colliculus of the rat. *Hearing Research* **117**:1–12.
- Zhang H and Kelly JB (2001) AMPA and NMDA receptors regulate responses of neurons in the rat's inferior colliculus. *Journal of Neurophysiology* **86**:871–880.
- Zhang H and Kelly JB (2003) Glutamatergic and GABAergic regulation of neural responses in inferior colliculus to amplitude modulated sounds. *Journal of Neurophysiology* **90**:477–490.
- Zhang H, Xu J, and Feng AS (1999) Effects of GABA-mediated inhibition on direction-dependent frequency tuning in the frog inferior colliculus. *Journal of Comparative Physiology A* **184**:85–98.
- Zhang Y and Wu SH (2000) Long-term potentiation in the inferior colliculus studied in rat brain slice. *Hearing Research* **147**:92–103.
- Zheng W and Knudsen EI (1999) Functional selection of adaptive auditory space map by GABA<sub>A</sub>-mediated inhibition. *Science* **284**:962–965.

# Chapter 10

## Biophysical Properties of Inferior Colliculus Neurons

SHU HUI WU

### 1. INTRODUCTION

The inferior colliculus (IC) is a pivotal nucleus in the central auditory pathway. It receives and integrates ascending afferent projections from almost all of the major auditory nuclei in the lower brain stem (see Chapters 3 and 4). Auditory information that has been encoded in specific temporal or spatial configurations in the lower brain stem nuclei is further analyzed, relayed, or transformed in the IC. The wealth of afferent inputs includes excitatory and inhibitory projections. The cochlear nucleus (CN), medial superior olive (MSO), and contralateral lateral superior olive (LSO) are sources for major excitatory inputs. The dorsal nucleus of the lateral lemniscus (DNLL), ipsilateral LSO, and ventral nucleus of the lateral lemniscus (VNLL) are sources for major inhibitory inputs. The IC also receives descending projections from the auditory cortex (AC) and medial geniculate body (MGB) (see Chapters 7 and 8), and local inputs from neurons within the IC (see Chapter 5). In addition, the IC receives converging auditory, somatosensory, visual, and motor information. Thus, IC neurons are influenced by ascending, descending, and internal excitatory and inhibitory synaptic inputs while processing auditory and other signals.

Information processed by IC neurons is sent to the MGB and the AC. The synaptic potentials of IC neurons are therefore transformed into action potentials that will further excite or inhibit their target neurons. The input–output transformation in a neuron is a result of interaction between biophysical properties of the neuron and synaptic inputs to it. The biophysical properties, that is, properties of intrinsic electrical membrane and synaptic receptors, shape the output signals from the IC neurons.

Electrical membrane properties of the neuron have passive and active components. The passive electrical properties, such as the resting potential, membrane input resistance and time constant, are relatively invariant, but crucial for electrical signaling. They determine whether a synaptic potential will elicit a suprathreshold depolarization at the spike-generating zone at the axon hillock. Passive properties also influence the conduction speed of an action potential. The voltage-gated ion channels and synaptic receptors on the cell membrane are

active components that allow rapid changes in membrane potential caused by ionic currents flowing across the cell membrane. The input signals activate ion channels and synaptic membrane receptors, which in turn produce excitatory or inhibitory synaptic potentials. This is a dynamic process in which cellular intrinsic properties interact with neural connections to produce a complex response profile that is arguably unique to each class of neuron.

In this chapter the passive and active membrane properties of IC neurons are summarized, and the question of how IC neurons integrate and process incoming signals based on their biophysical properties is considered. Suggested directions for future research in the cellular biophysics of IC neurons will provide a guide for better understanding neuronal mechanisms of auditory processing.

## 2. ELECTRICAL MEMBRANE PROPERTIES

Electrical membrane properties are usually studied by using intracellular recordings from neurons in rodent IC brain slices. Firing patterns and voltage changes across the cell membrane are examined with current-clamp recordings by injection of depolarizing and hyperpolarizing current. Ionic currents underlying the firing pattern are investigated with voltage-clamp recording.

### 2.1. FIRING PATTERNS AND POTASSIUM CURRENTS

Different IC neurons respond selectively and specifically to intracellular injection of depolarizing and hyperpolarizing current (Wagner 1994; Li et al. 1998; Reetz and Ehret 1999; Peruzzi et al. 2000; Wu et al. 2002). Depolarizing current injection activates voltage-sensitive ion channels and generates action potentials, eliciting either onset, sustained-regular, or adapting firing patterns (Wagner 1994; Peruzzi et al. 2000). Some cells show a depolarizing rebound after injection of hyperpolarizing current and others show a pause in response to injection of depolarizing current following hyperpolarizing current. Different firing patterns are correlated with different combinations of specific  $K^+$  channels. Many of the details of the firing patterns discussed in Sections 2.1.1 to 2.1.4 are based on a study by Sivaramakrishnan and Oliver (2001) in which the firing pattern and associated  $K^+$  currents were examined in single neurons of the central nucleus of the IC (ICC) with both current- and voltage-clamp recordings.

#### 2.1.1. Sustained-Regular Pattern

In the study by Sivaramakrishnan and Oliver (2001) neurons with a regular discharge pattern showed sustained firing with constant interspike intervals throughout the period of depolarizing current injection (Fig. 10.1A). Membrane depolarization caused a similar increase in both firing frequency and the magnitude of the 4-aminopyridine (4-AP)- and tetraethylammonium (TEA)-sensitive  $K^+$  currents, suggesting that the sustained-regular firing pattern results from




Figure 10.1. The six firing patterns of ICC neurons (for abbreviations, please see the list). In each panel, the *bottom two traces* are the injected depolarizing and hyperpolarizing current pulses, respectively, and the *top two traces* are the voltage responses to each current pulse, respectively. In (C), the thick and thin current traces represent the current that elicited the top and bottom voltage traces, respectively. *Dotted lines* indicate resting membrane potentials. *b*, buildup response; *p*, pause in firing; *r*,  $\text{Ca}^{2+}$ -dependent rebound. (Adapted from Sivaramakrishnan and Oliver 2001. Copyright 2001 by the Society for Neuroscience.)



4-AP- and TEA-sensitive delayed rectifying  $K^+$  currents (Rudy et al. 1999). In vivo these neurons can readily encode information about sound duration and intensity by a linear increase in firing rate because of their intrinsic linear rate-current function.

### 2.1.2. Rebound Pattern

Rebound neurons had a period of depolarization after cell membrane hyperpolarization. This rebound was eliminated by reducing extracellular  $Ca^{2+}$  concentration. With  $Na^+$  and  $K^+$  channel blockers, depolarizing current injection into the rebound cell produced regenerative  $Ca^{2+}$  spikes far longer (around 100 ms) than typical  $Na^+$  action potentials.

Rebound cells responded to depolarizing stimulation with regular, adapting, or transient firing (Fig. 10.1D–F). Like the sustained-regular cells, rebound-regular cells had a delayed rectifying  $K^+$  current, which helps to repolarize the cell during an action potential and to maintain regular firing during depolarizing stimulation. These cells may encode sound intensity because their linear rate-current function resembles that of the sustained-regular cells.

Rebound-adapting neurons fired action potentials with nonuniform interspike intervals to depolarizing current injection. Spikes were initially close together, but became more separate during the injection period. The outward  $K^+$  currents in these neurons were partially blocked by apamin, a blocker for small-conductance, voltage-independent  $Ca^{2+}$ -activated  $K^+$  current (Hugues et al. 1982), and reduced by low extracellular  $Ca^{2+}$  concentration. The smallest apamin-sensitive currents were evoked at the membrane potential for which the greatest adaptation occurred, suggesting that apamin-sensitive  $Ca^{2+}$ -activated  $K^+$  currents contribute to sustained firing during membrane depolarization.

Rebound-transient cells fired transiently during depolarizing current injection. Transient firing was converted to sustained firing by carybdotoxin (CTX), a blocker of the voltage-dependent BK channel. This type of  $K^+$  channel has a large single-channel conductance and is activated by increase in intracellular  $Ca^{2+}$  (Vergara et al. 1998). In rebound-transient ICC neurons the outward  $K^+$  currents had a  $Ca^{2+}$ -dependent component that was rapidly inactivated after activation and was blocked by CTX. Thus, CTX-sensitive  $Ca^{2+}$ -activated  $K^+$  currents are likely responsible for transient firing in rebound-transient neurons.

The depolarizing rebound increases the probability of a  $Na^+$ -dependent anode-break spike following membrane hyperpolarization, which may underlie the OFF response to acoustic stimulation noted in IC neurons in vivo (Covey et al. 1996; Kuwada et al. 1997). The rebound mechanism may permit precise temporal coding of ongoing complex sounds such as an amplitude-modulated tone (Kuwada and Batra 1999).

The inhibition-induced rebound mechanism may also underlie sound duration tuning (Casseday et al. 1994, 2000; Covey et al. 1996; Ehrlich et al. 1997). Some IC neurons in bat, rat, and mouse respond preferentially to a specific range of sound durations (Chen 1998; Brand et al. 2000). These neurons may have

excitatory and inhibitory input components that occur in a certain temporal sequence (see Chapter 17). For a preferred sound duration, the OFF excitation that follows the onset inhibition coincides and summates with the delayed excitation, generating a suprathreshold response (Casseday et al. 1994). The OFF excitation is likely the depolarizing rebound observed in rebound neurons *in vitro*.

### 2.1.3. Buildup-Pause Pattern

When depolarizing current injection was preceded by a hyperpolarizing current step, buildup-pause neurons delayed the onset of sustained firing or developed a pause between first spike and subsequent firing (Fig. 10.1C). As the prehyperpolarization increased, the pause and magnitude of the A-type  $K^+$  current ( $I_A$ ), which is characterized as a fast transient and rapidly inactivating  $K^+$  current (Connor and Stevens 1971), increased in parallel. Perhaps the delay or pause at the beginning of sustained firing follows activation of  $I_A$  after removal of its inactivation by prehyperpolarization.

As the  $I_A$  in the buildup-pause neurons decayed to zero about 80 to 100 ms after activation, the pause would be elicited only if the interval between inhibitory and excitatory stimuli were  $<100$  ms. Like sustained-regular and rebound regular cells, the buildup-pause neuron can encode sound intensity because of its linear rate-current function. The buildup-pause neurons are suited to register successive excitatory stimuli or pairs of inhibitory and excitatory stimuli (Sivaramakrishnan and Oliver 2001).

### 2.1.4. Onset Pattern

Onset cells fire only once at the onset of a depolarizing current injection. The  $K^+$  current activated at  $-50$  to  $-40$  mV, known as low-threshold  $K^+$  current, probably underlies the onset firing pattern. The high-threshold ( $-10$  to  $+10$  mV)  $K^+$  current also exists in the onset neurons. It may help them to repolarize rapidly even if the cell responds to synaptic inputs with a large excitatory potential. Basta and Vater's (2003) study shows that some onset neurons in the mouse's ICC had onset firing to depolarizing current injection only when the neuron was at a hyperpolarizing membrane potential (near  $-80$  mV). In these neurons the inward current that was sensitive to tetrodotoxin, a specific blocker of voltage-gated  $Na^+$  channels, appeared only at  $-80$  mV. The results suggest that the onset firing in these neurons could be attributed to the prehyperpolarization-activated  $Na^+$  current. Only 14% of IC neurons in their study showed a prehyperpolarization-evoked onset pattern. Nevertheless, this  $Na^+$  conductance may enable gating of excitatory synaptic inputs in a narrow time window after inhibitory inputs (Basta and Vater 2003).

In summary, IC neurons represent at least six physiological types based on firing patterns to combinations of depolarizing and hyperpolarizing current injections. Sustained-regular and rebound-regular neurons have only the delayed rectifier  $K^+$  current, and other neurons have more specific  $K^+$  currents besides

the delayed rectifier  $K^+$  current (Table 10.1). In Sivaramakrishnan and Oliver's study (2001) the  $Ca^{2+}$  rebound was seen in 57% of cells after hyperpolarizing current injection. Nearly half of these showed a rebound-adapting firing pattern. The regular and transient patterns in the rebound cells were 11% and 21%, respectively. Sustained-regular and buildup-pause patterns comprised 19% and 15% of the population, respectively, while onset cells were 9%. Other studies find that the sustained discharge pattern is prevalent in IC neurons (Table 10.2) (Li et al. 1998; Reetz and Ehret 1999; Bal et al. 2002).

Synaptic inputs may be processed differently by each physiological neuron type (Sivaramakrishnan and Oliver 2001). Input signals can be modified substantially in rebound-adapting, rebound-transient, buildup-pause, and onset neurons, and less so in rebound-regular and sustained-regular neurons. The firing pattern can be changed when inhibition precedes excitation in rebound and buildup-pause neurons, but not in onset and sustained-regular cells. Sustained-regular cells can relay information about sound intensity and duration, and onset cells likely encode the precise onset of sound signals.

Onset, sustained-regular, and rebound-adapting neurons are all found in the IC cortex (Li et al. 1998; Peruzzi et al. 2000). The active membrane properties of IC cortex neurons resemble those of ICC neurons (Smith 1992). Most IC cortex neurons generate  $Na^+$  overshooting action potentials followed by a biphasic afterhyperpolarization (AHP) with a fast and a slow component (Li et al. 1998). The AHP is attributed to  $Ca^{2+}$ -activated  $K^+$  currents because it is abolished by CTX and its shape is altered by extracellular  $Ca^{2+}$  removal (Smith 1992; Sivaramakrishnan and Oliver 2001). An obvious feature of intrinsic membrane properties of IC cortex neurons is a slow depolarizing  $Ca^{2+}$ -

Table 10.1. Specific  $K^+$  current present in IC neurons with different firing patterns.<sup>a</sup>

Firing pattern	Apamin-sensitive $I_k(Ca)$	CTX-sensitive $I_k$	High-threshold $I_k$ (0.2 mM TEA-sensitive)	A-type of $I_k$
Sustained-regular	Not present	Not present	Not present	Not present
Rebound-regular	Not present	Not present	Not present	Not present
Rebound-adapting	30% of total $K^+$ current	Not present	Not present	Not present
Rebound-transient	Not present	55% of total $K^+$ current	Not present	Not present
Buildup-pause	9% of total $K^+$ current	Not present	Not present	56.5% of total $K^+$ current
Onset	No $Ca^{2+}$ -dependent $K^+$ current	No $Ca^{2+}$ -dependent $K^+$ current	45% of total $K^+$ current	Not present

<sup>a</sup>From Sivaramakrishnan and Oliver (2001). Neurons with any of these six firing patterns all have delayed rectifier  $K^+$  current which can be blocked by 2 mM 4-AP or TEA.

Table 10.2. Proportion of sustained and onset IC neurons.

Study	Type of recording	Sustained <sup>a,b</sup>	Onset
Wagner (1994)	Intracellular	62	38
Li et al. (1998)	Intracellular	77	23
Reetz and Ehret (1999)	Intracellular	55	45
Peruzzi et al. (2000)	Intracellular	71	29
	Whole-cell patch clamp	66	34
Sivaramakrishnan and Oliver (2001)	Whole-cell patch clamp	91	9
Bal et al. (2002)	Intracellular	84	16

<sup>a</sup>Sustained neurons include regular, adapting, and transient types.

<sup>b</sup>Values are percentages.

dependent hump in response to depolarizing current injection. This hump is often observed in IC dorsal cortex (ICD) neurons, less so in IC external cortex (ICX) cells, and never in ICC neurons. The hump reflects  $\text{Ca}^{2+}$  current flowing through  $\text{Ca}^{2+}$  channels (Smith 1992). The absence of the hump in ICC cells may be caused by a strong  $\text{Na}^+$  current that masks a weak  $\text{Ca}^{2+}$  current (Li et al. 1998), as ICC neurons do exhibit  $\text{Ca}^{2+}$  spikes after  $\text{Na}^+$  and  $\text{K}^+$  channels are blocked (Sivaramakrishnan and Oliver 2001). In any event, the voltage-dependent  $\text{Ca}^{2+}$  current and  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  current contribute to action potential shape and firing behavior in a similar way for ICC and ICX cells. The apparent calcium hump in ICD neurons indicates more  $\text{Ca}^{2+}$  channel activity, which may partly explain their lower firing rate.

## 2.2. PASSIVE MEMBRANE PROPERTIES

The average range of input resistance of IC neurons is 44 to 116  $\text{M}\Omega$  in intracellular recordings (Wagner 1994; Reetz and Ehret 1999; Bal et al. 2002) and 240 to 644  $\text{M}\Omega$  in whole-cell patch clamp recordings (Sivaramakrishnan and Oliver 2001). The higher input resistance with the latter is consistent with the smaller leak current in the whole-cell patch configuration. In whole-cell patch clamp recordings by Sivaramakrishnan and Oliver (2001) the onset neurons had the highest input resistance. The sustained-regular, buildup-pause, and rebound-regular cells had similar input resistances, which were intermediate to onset and rebound-adapting-transient groups. The rebound-adapting and rebound-transient cells had similar input resistances, which were the lowest among IC cell types. The membrane time constant of many IC neurons is 2 to 8 ms in intracellular recordings (Wagner 1994; Reetz and Ehret 1999), resembling that of many auditory brain stem neurons (Hirsch and Oertel 1988; Wu and Kelly 1991, 1995). However, onset neurons have a far shorter time constant than the sustained cells (Wagner 1994), and a shorter time constant for depolarizing than to hyperpolarizing current injection. These membrane properties allow onset cells to respond to input signals faster than the other cell types. Onset neurons can follow

a short train of current pulses at approximately 200 Hz, whereas rebound cells saturate at 50 Hz (Peruzzi et al. 2000).

Differences in passive membrane properties and firing characteristics among ICC, ICX, and ICD neurons have been noted (Li et al. 1998). The input resistance of sustained-firing type of neurons in ICX and ICD is higher than that of ICC neurons. The time constant of sustained type of ICD neurons is longer than that of ICX and ICC neurons. Among the three subdivisions, ICX neurons have the highest spontaneous activity and firing rate in response to depolarizing current injection. The threshold for sustained firing is closer (10 to 20 mV) to the resting potential in ICX and ICC than in ICD neurons. These differences suggest that ICX neurons are more excitable than ICC and ICD neurons, which is consistent with their higher audiogenic seizure susceptibility (Li et al. 1998).

### 3. THE RELATIONSHIP BETWEEN CELL PHYSIOLOGY AND MORPHOLOGY

#### 3.1. THE CENTRAL NUCLEUS

ICC neurons have been classified morphologically as disc-shaped (or flat cells) and stellate cells (or less-flat cells) on the basis of their dendritic shape and branching (Peruzzi et al. 2000; see Chapter 2). Electrophysiologically identified categories, that is, rebound, buildup-pause, and so forth do not correlate with the morphologic classification of cells (Peruzzi et al. 2000; Bal et al. 2002). The passive membrane properties of flat and less-flat cells show no significant differences for input resistance, time constant, membrane capacitance, or resting potential (Reetz and Ehret 1999; Bal et al. 2002). However, other morphologic features correlate with electrophysiological profiles (Peruzzi et al. 2000). For example, rebound cells have larger somatic areas and diameters than buildup-pause neurons, which may explain their lower input resistance. Buildup-pause neurons have fewer and simpler dendritic branches than onset and rebound neurons. Axons of some biocytin-labeled IC neurons in mouse brain slices in Reetz and Ehret's study (1999) showed branches, which were parallel to ICC isofrequency laminae and could be traced into the IC brachium, commissure of Probst, ICX, ICD, lateral lemniscus, and IC commissure. But a correlation between the intrinsic electrophysiology and axon arborization has not been found.

#### 3.2. THE INFERIOR COLLICULUS CORTEX

The relationship between the membrane properties and morphology of IC cortex neurons in rat has been examined only in multipolar cells (Smith 1992). The labeled neurons show similar electrical membrane characteristics, responding to depolarizing current injection with a sustained, adapting firing pattern and depolarizing rebound following hyperpolarizing current injection. The results indicate the presence of voltage-dependent  $\text{Na}^+$ ,  $\text{Ca}^{2+}$  and  $\text{Ca}^{2+}$ -activated  $\text{K}^+$

channels in these neurons (Smith 1992). Future study of physiological properties in different morphologic types of cells is essential for understanding how IC cortical neurons process ascending and descending inputs and influence intra- and intercollicular targets.

## 4. ION CHANNEL CHARACTERISTICS

The intrinsic membrane conductance of an IC neuron, which is attributed to the type, number, and kinetic properties of ion channels, plays a key role in preserving and modulating auditory signals. Electrophysiological, pharmacological, and molecular–biological studies have revealed the features of a wealth of ion channels in IC neurons.

### 4.1. POTASSIUM CHANNELS

Traditionally, K<sup>+</sup> channels have been classified on the basis of their electrophysiological characteristics. However, molecular–biologic advances have allowed the cloning of the  $\alpha$  subunit genes of the rapidly activating, delayed-rectifier (i.e., the Kv channel), which has four subfamilies: *Kv1* to *Kv4* (Coetzee et al. 1999). Each subfamily has different genetic isoforms, for example, *Kv1.1* to *1.7*, *Kv2.1* to *2.2*, *Kv3.1* to *3.4*, and *Kv4.1* to *4.4* subunits, that may encode distinct K<sup>+</sup> channels (Blaine and Ribera 1998; Song 2002).

An in situ hybridization study showed expression of both *Kv1.1* and *1.2* in mouse IC (Grigg et al. 2000). An immunocytochemical study in big brown bat found a gradient of *Kv1.1* labeling from ventrolateral to dorsomedial IC (Rosenberger et al. 2003). The labeled cells were multipolar and confined to the ventral IC. The low-threshold K<sup>+</sup> current in rat IC onset neurons resembles the current through *Kv1.1* channels, which are expressed strongly in CN bushy cells and octopus cells, and in principal cells of the medial nucleus of the trapezoid body (MNTB) (Grigg et al. 2000). This K<sup>+</sup> channel activates rapidly at membrane potentials of  $-50$  to  $-30$  mV, and then slowly inactivates (Brew and Forsythe 1995; Dodson et al. 2002). The low-threshold K<sup>+</sup> current in these auditory neurons accounts for the steep decrease in membrane input resistance near the resting potential and the brevity of synaptic potentials, both specializations for conveying precise timing information (Wu and Oertel 1984; Manis and Marx 1991; Wu and Kelly 1991; Golding et al. 1999; Svirskis et al. 2002). Whether IC neurons with *Kv1.1* also process precise temporal information should be examined in future studies.

IC neurons have weaker *Kv1.1* and *Kv1.2* expression than MNTB neurons, but stronger *Kv1.2* expression than CN bushy and octopus cells (Grigg et al. 2000). The physiological observation that a small number of IC neurons (onset type) have a low-threshold K<sup>+</sup> current probably flowing through the *Kv1.1* channels (Sivaramakrishnan and Oliver 2001) seems consistent with the finding of weak *Kv1.1* expression in the IC. Most IC neurons have a sustained firing pattern

and may express the *Kv1.2* channel. As these channels activate more slowly and at a more positive voltage than *Kv1.1* channels, the *Kv1.2* channel in the sustained firing neurons would allow temporal summation of multiple synaptic inputs, but not precise synaptic transmission of phase-locked information (Grigg et al. 2000).

A high level of *Kv3.1* mRNA expression has been observed in rat IC neurons (Perney et al. 1992; Weiser et al. 1994; Li et al. 2001). Liu and Kaczmarek's study (1998) shows that depolarization caused parallel increases in *Kv3.1* mRNA level and in the amplitude of a voltage-dependent high-threshold, noninactivating  $K^+$  current, strongly suggesting the presence of functional *Kv3.1* channels in IC neurons. *Kv3.1* currents have a high threshold ( $-10$  to  $-20$  mV) for activation, and fast kinetics for both activation and deactivation. These currents are very sensitive to external TEA and 4-AP (Luneau et al. 1991; Critz et al. 1993; Perney and Kaczmarek 1997; Wang et al. 1998). In IC onset neurons the outward  $K^+$  current has a high-threshold component whose activation kinetics and pharmacologic properties resemble those of the *Kv3.1* channel. This current likely flows through *Kv3.1* channels (Sivaramakrishnan and Oliver 2001).

The physiological profile of *Kv3.1* current is ideal to allow neurons to fire action potentials at high frequencies, as these channels open quickly once their threshold is reached, often near the peak of an action potential. The  $K^+$  current influx then rapidly repolarizes the cell membrane and truncates action potential duration. On repolarization, *Kv3.1* channels close rapidly because of deactivation, thus minimizing the action potential refractory period and enabling the generation of subsequent action potentials (Li et al. 2001). Some auditory neurons that fire at high rates express *Kv3.1* channels strongly (Perney and Kaczmarek 1997; Wang et al. 1998). For example, CN bushy cells and MNTB principal cells can fire action potentials in response to electrical stimulation of their synaptic inputs up to about 500 Hz (Wu and Kelly 1993; Oertel 1997), and both express a high level of *Kv3.1* mRNA (Li et al. 2001). Similarly, IC onset neurons, which probably have *Kv3.1* channels, can also follow a train of short depolarizing current pulses at frequencies up to 200 Hz (Peruzzi et al. 2000).

*Kv3.3* mRNA is also strongly expressed in the rat IC (Weiser et al. 1994; Li et al. 2001). The *Kv3.3* channel has activation kinetics, voltage sensitivity, and pharmacology similar to those of the *Kv3.1* channel, but differs in inactivation rate (Rudy et al. 1999). *Kv3.1* transcripts express a delayed-rectifier type current with very little inactivation, while *Kv3.3* transcripts express transient currents with moderate inactivation (Weiser et al. 1994). Although *Kv3.3* mRNA is expressed in IC neurons, the inactivating high-threshold  $K^+$  current flowing through *Kv3.3* channels has not been detected (Liu and Kaczmarek 1998).

*Kv9.1*, one of the genes that encode the  $\alpha$  subunit of the voltage-dependent  $K^+$  channel, is co-expressed in IC but cannot form functional potassium channels alone. However, if *Kv9.1* is coexpressed with the *Kv2.1* channel that generates slowly inactivating delayed rectifier  $K^+$  currents, the *Kv2.1* channel activation rate decreases, and the voltage dependence of inactivation and activation shifts

toward more negative membrane potentials (Richardson and Kaczmarek 2000). Because *Kv2.1* is expressed in IC (Hwang et al. 1993), the *Kv2.1* channel may be modulated by *Kv9.1*, which might affect firing patterns significantly (Richardson and Kaczmarek 2000).

The *Kv4* gene subfamily has been proposed as the key component of the classic A-type  $K^+$  channel for A-current ( $I_A$ ). *Kv4* channels activate and inactivate at subthreshold potentials, and recover from inactivation more quickly than other inactivating *Kv* channels. An in situ hybridization study (Serôdio and Rudy 1998) found high *Kv4* expression in the dorsal cochlear nucleus (DCN) and a low level in the IC. The DCN pyramidal cells express the fast A-type current with kinetic properties appropriate to support their pause-buildup firing pattern (Kanold and Manis 1999). Only about 15% of IC neurons have buildup-pause firing pattern (Sivaramakrishnan and Oliver 2001), which may explain the weak expression of mRNA of the *Kv4* gene.

#### 4.2. CALCIUM-ACTIVATED POTASSIUM CHANNEL

IC neurons express two types of  $Ca^{2+}$ -activated  $K^+$  channel, a large conductance, BK-type (or Maxi-type) and a small conductance, SK-type. The characteristics of BK- and SK-type currents were investigated in adapting- and transient-rebound neurons (Sivaramakrishnan and Oliver 2001). Neurons with a transient firing pattern had a voltage-dependent BK current that was rapidly inactivated after activation and blocked by CTX. Elimination of the BK current abolished AHP and altered the transient discharge to more sustained firing. Thus, the transient firing behavior and the AHP are attributed to the BK current. Neurons with an adapting firing pattern possessed an SK current, which showed little voltage dependence, had a threshold comparable to that of the total  $K^+$  current, and was blocked by apamin. The SK current determined the degree of firing adaptation and maintained sustained firing to a depolarizing current injection.

How BK and SK currents contribute to the spike AHP of IC neurons has not been systematically investigated. IC neurons have two types of spike AHP with different time courses. The fast AHP (1 to 10 ms) was common, whereas the slower AHP (10 to 100 ms) was found in only 20% of IC neurons (Li et al. 1998). The duration of spike AHP is one of the determinants for the maximal neural firing frequency. The AHP time course in different IC neurons and identification of the responsible  $Ca^{2+}$ -activated  $K^+$  currents require further study.

#### 4.3. OTHER POTASSIUM CHANNELS

Immunocytochemical and cloning studies have revealed other  $K^+$  channels besides the *Kv* channels in IC neurons. For example, mRNA for rat *eag2*, a member of the *ether-à-go-go* family of  $K^+$  channel genes, is expressed abundantly in the rat IC and midbrain (Ludwig et al. 2000). This gene encodes the prototype of a family of voltage-gated  $K^+$  channels distinct from those of the *Kv* channel group. These channels have a strong voltage-dependence of activation, which is



sensitive to extracellular  $Mg^{2+}$  concentration (Terlau et al. 1996), and can be blocked by low cytosolic  $Ca^{2+}$  (Meyer and Heinemann 1998). The significance of plentiful IC rat *eag2* gene expression is not clear.

The ATP-sensitive  $K^+$  channel ( $K_{ATP}$ ) and inward rectifier ( $K_{IR}$ ) belong to another superfamily of  $K^+$  channels. Outward  $K^+$  currents activated by the purinoceptor agonists are seen in cultured rat IC neurons (Ikeuchi and Nishizaki 1995b). The purine receptors in IC have been identified as the  $P_{2Y}$  subtype. Unlike  $K_{ATP}$  channels in other neurons, in which the  $K_{ATP}$  channel is activated by a G-protein-coupled second messenger system (Barnard et al. 1994; Ikeuchi and Nishizaki 1995a), the  $P_{2Y}$  purinoceptor gated  $K^+$  channels in IC neurons are activated by direct coupling to the  $\beta\gamma$  subunits of G-protein (Ikeuchi and Nishizaki 1995b). The physiological significance of the  $K_{ATP}$  channel in IC neurons is unknown.

The IC also expresses an inward rectifier  $K^+$  channel ( $K_{IR}$ ). Immunocytochemical and in situ hybridization studies show staining of G-protein gated inward rectifier  $K^+$  channel protein, the GIRK1 protein (Bausch et al. 1995), and its mRNA (Karschin et al. 1994) in IC neuron somata and dendrites.  $K_{IR}$  channels conduct very little current at physiological potentials but pass far more current when the cell is at a membrane potential negative to the  $K^+$  equilibrium potential. Opening these channels also depends on extracellular  $K^+$  concentration. GIRK1 and  $\mu$  opioid receptors are colocalized over the same neurons in many brain regions, including the IC. These results suggest that  $\mu$  opioid receptors may regulate  $K^+$  conductance by activation of GIRK1 (Bausch et al. 1995). The significance of  $K_{IR}$  channels in IC neurons remains to be determined.

#### 4.4. CALCIUM CHANNELS

$Ca^{2+}$  currents are implicated in the generation of rebound after hyperpolarization and in the depolarizing hump elicited by subthreshold positive current injection in IC neurons (Smith 1992; Sivaramakrishnan and Oliver 2001). A biophysical and pharmacologic study of  $Ca^{2+}$  currents found several types of voltage-sensitive  $Ca^{2+}$  channels in the 2- to 8-day-old rat IC (N'Gouemo and Rittenhouse 2000). All neurons expressed a high-threshold voltage-activated  $Ca^{2+}$  current that activated around  $-40$  mV. Low- (activated at  $-70$  mV) and mid-threshold (activated at  $-50$  mV) voltage-activated  $Ca^{2+}$  currents were present in 33% and 40% of the recorded neurons, respectively. In IC there are four types of high threshold  $Ca^{2+}$  channels: L-type, N-type, P/Q-type, and a type that is resistant to a combination of the antagonists for all three types of high-threshold  $Ca^{2+}$  channels. How the various  $Ca^{2+}$  channels contribute to membrane excitability, action potential genesis, discharge pattern, firing threshold, and neurotransmitter release in the IC is not clear. Nevertheless, intracellular  $Ca^{2+}$  can influence many neural processes by regulating membrane and cytosolic proteins, such as ion channels, exchangers, pumps, and enzymes. Therefore, the significance of  $Ca^{2+}$  current in electrical activity, biochemical processes, and plasticity in IC should be investigated further.

## 5. SYNAPTIC RECEPTORS


This section focuses on the physiological properties of synaptic receptors in IC neurons activated by excitatory and inhibitory amino acids, including glutamate,  $\gamma$ -aminobutyric acid (GABA) and glycine; biogenic amines, such as serotonin (5-hydroxytryptamine, 5-HT); and acetylcholine (Ach).

### 5.1. IONOTROPIC GLUTAMATE RECEPTORS

Excitatory synaptic transmission in the IC is probably mediated by glutamate through ionotropic glutamate receptors. Intracellular and patch clamp recordings from ICC neurons in rat, mouse, and gerbil brain slices have found that both  $\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) and *N*-methyl-D-aspartate (NMDA) receptors, the two main ionotropic glutamate receptors, mediate excitatory synaptic responses evoked by electrical stimulation of the lateral lemniscus or the IC commissure (Wagner 1996; Moore et al. 1998; Ma et al. 2002a). Excitatory commissural transmission to the ICD is also mediated by AMPA and NMDA receptors (Li et al. 1999).

Excitatory postsynaptic potentials (EPSPs) examined by Ma et al. (2002a) had two components—the early and the late. They were blocked by the AMPA antagonists, for example, 6-cyano-7-nitroquinoxaline-2,3-dione (CNQX) or 6,7-dinitroquinoxaline-2,3-dione (DNQX), and the NMDA receptor antagonist, D, L-2-amino-5-phosphonovaleric acid (APV), respectively (Fig. 10.2). These results suggest that the early part of the EPSP is mediated by AMPA receptors, and the later part by NMDA receptors. The temporal properties of these two types of glutamate receptors in ICC neurons are distinct (Ma et al. 2002a). Rise time, decay time, and half-width of AMPA component were 20% to 30% of the values for the NMDA component. Removing extracellular  $Mg^{2+}$  greatly enhanced only the NMDA component in the EPSP, suggesting that the NMDA receptors are suppressed by extracellular  $Mg^{2+}$ . AMPA and NMDA receptors in ICC neurons also differ in voltage dependence. The NMDA receptor-mediated response was voltage dependent and it was smaller when the membrane was hyperpolarized. The current-voltage (*I/V*) relationship for AMPA receptor-mediated EPSPs was linear, whereas that for the NMDA receptor-mediated EPSPs was nonlinear (Fig. 10.3) (Ma et al. 2002a). The rectification of the *I/V* curve for NMDA responses is attributable to a voltage-dependent block of the NMDA receptors by extracellular  $Mg^{2+}$ . These features of AMPA and NMDA receptors in ICC neurons resemble those of ionotropic glutamate receptors in other neurons, for example, hippocampal pyramidal cells (Mayer et al. 1984; Hestrin et al. 1990). However, unlike the glutamate receptors in hippocampal pyramidal neurons, in which NMDA receptors usually do not activate at the cell's resting potential (Hestrin et al. 1990), NMDA receptor-mediated responses in ICC are elicited at the cell's resting membrane potential and are not eliminated by blocking AMPA receptor-mediated responses (Ma et al. 2002a). NMDA re-

Figure 10.2. Excitatory synaptic response of ICC neurons is mediated by both AMPA and NMDA receptors. (A) In a brain slice perfused with normal saline an EPSP was elicited by electrical stimulation of the lateral lemniscus with an early and fast, and a later and slower, component from an ICC neuron. (B) APV application ( $100 \mu\text{M}$ ) suppressed the later part of the EPSP. (C) The early part of the EPSP was blocked by CNQX ( $10 \mu\text{M}$ ). The brain slice was perfused with strychnine ( $0.5 \mu\text{M}$ ) and bicuculline ( $10 \mu\text{M}$ ) while APV and CNQX were applied. Arrow, stimulus artifact.



ceptors in the IC may have lower sensitivity to  $\text{Mg}^{2+}$  blockade, therefore requiring less membrane depolarization for activation, as demonstrated in hippocampal stratum radiatum giant neurons (Kirson and Yaari 2000).

The recombinant heterodimeric NMDA receptors that contain NR2A or NR2B subunits are more strongly blocked by  $\text{Mg}^{2+}$  than those with NR2C or NR2D subunits (Kuner and Schoepfer 1996). Thus, the lower  $\text{Mg}^{2+}$  sensitivity of IC NMDA receptors may be attributable to expression of more NR2C and NR2D subunits than NR2A and NR2B subunits. How subunit composition affects  $\text{Mg}^{2+}$  sensitivity and voltage dependence of NMDA receptors in IC neurons requires further investigation. NMDA receptor subunits also undergo maturational changes, that is, down-regulation of NR2B and NR2D subunits, and up-regulation of NR2A (Kirson et al. 1999). In rat ICC neurons, the NMDA components decline relative to AMPA components during 1 to 2 weeks postnatal and then stabilize at 13 to 16 days (Ma et al. 2002a) and probably throughout adulthood. The ontogeny of the subunit composition of NMDA receptors in IC neurons is unknown.




Figure 10.3. Current-voltage relationship of AMPA and NMDA receptor-mediated EPSCs in an ICC neuron. **(A) Left panel:** AMPA receptor-mediated EPSCs at various membrane potentials after blocking the NMDA receptor-mediated EPSCs with APV ( $100 \mu\text{M}$ ). **Right panel:** The *I/V* curve obtained from the peak amplitude of the isolated AMPA EPSCs was linear. **(B)** The NMDA receptor-mediated EPSCs from the same neuron at various membrane potentials after blocking the AMPA receptor-mediated EPSCs with CNQX ( $15 \mu\text{M}$ ). The ensuing current-voltage curve was highly nonlinear at negative voltages. In both conditions the slice was perfused with strychnine ( $0.5 \mu\text{M}$ ) and bicuculline ( $10 \mu\text{M}$ ). (Adapted from Ma et al. 2002a. Reprinted from Hearing Research with permission from Elsevier.)

The function of NMDA receptors in the IC is not fully understood. NMDA receptors are required for induction of long-term potentiation (LTP), a long-lasting increase in synaptic strength evoked by tetanic stimulation, in rat IC brain slices (Zhang and Wu 2000; Wu et al. 2002). Activation of NMDA receptors influences expression of newly learned auditory responses in the barn owl IC homologue (Feldman and Knudsen, 1998) and is involved in auditory responses in rat (Zhang and Kelly 2001; Kelly and Zhang 2002; see Chapter 9). In Zhang and Kelly's study (2001) iontophoretic application of AMPA or NMDA antagonists onto ICC cells of adult rats reduced responses to tone bursts and amplitude-modulated sounds. AMPA receptor antagonists selectively reduced the number of spikes evoked during the early response whereas the NMDA receptor antagonists reduced spikes evoked later. The NMDA receptor may amplify stimulus-evoked responses when the cell is depolarized by AMPA receptors and set the gain of neuronal responses to increasing stimulus intensity. The slow kinetics of NMDA receptors may provide a wide window for processing the inputs converging onto IC neurons at variable delays and with different time courses (Kelly and Zhang 2002). These hypotheses can be examined *in vivo* with direct application of NMDA.

NMDA receptors are implicated in audiogenic seizures (see Chapter 21). ICC and ICD neurons generate sustained or paroxysmal depolarization and epileptiform activity (burst discharges) after reduction of extracellular  $Mg^{2+}$  or application of NMDA (Pierson et al. 1989; Smith 1992; Wagner 1996; Li et al. 1998; Ma et al. 2002a). NMDA receptors in the IC may also be involved in the propagation of generalized tonic-clonic seizures (Yasuda et al. 2000). The increased audiogenic seizure susceptibility in some mice correlates with the overexpression of NR2C subunits (Marianowski et al. 1995) that reduce  $Mg^{2+}$  block of the NMDA receptor, thus promoting epileptic events.

## 5.2. $\gamma$ -AMINO BUTYRIC ACID RECEPTORS

GABAergic inhibition plays a vital role in normal IC auditory processing as well as in audiogenic seizure genesis. It affects firing rate, tuning curves, rate-level functions, duration tuning, and binaural responses (Faingold et al. 1989; Vater et al. 1992; Yang et al. 1992; Park and Pollak 1993, 1994; Pollak and Park 1993; Casseday et al. 1994; Klug et al. 1995; Fuzessery and Hall 1996; Le Beau et al. 1996; Palombi and Caspary 1996; Burger and Pollak 1998; Koch and Grothe 1998; Zhang et al. 1999). Blocking GABA uptake in the ICC or applying GABA<sub>A</sub> or GABA<sub>B</sub> receptor agonists significantly reduces audiogenic seizure severity in genetically epilepsy-prone rats (Faingold et al. 1994a,b). Conversely, bicuculline application to the ICC of normal rats can produce a short-term susceptibility to audiogenic seizures (Millan et al. 1986). GABA-mediated inhibition is thus a critical mechanism in genetic and induced forms of audiogenic seizures (Faingold 2002).

Intracellular studies in rat and gerbil brain slices have found that IPSPs of ICC neurons evoked by lateral lemniscal or commissural stimulation are medi-

ated predominantly by GABA<sub>A</sub> receptors (Smith 1992; Moore et al. 1998; Ma et al. 2002a). The reversal potential of synaptically evoked GABAergic inhibitory postsynaptic currents (IPSCs) and GABA-activated currents corresponds to the equilibrium potential of Cl<sup>-</sup> ions, indicating that ICC GABAergic inhibition is mediated by Cl<sup>-</sup> ions (Hosomi et al. 1997). The kinetics of GABA<sub>A</sub> receptors resemble those of AMPA receptors (Wu et al. 2004a). GABAergic IPSCs rise rapidly and peak within 10 ms, and have a half-width of only 20 to 30 ms. The fast decay time constant (about 30 ms) of GABAergic spontaneous IPSCs in juvenile IC cells (Kraushaar and Backus 2002) may reflect a high level of expression of the  $\alpha_1$  subunit of GABA<sub>A</sub> receptors (Sato et al. 2000; Kraushaar and Backus 2002).

The time course of the GABAergic response resembles that of the AMPA receptor-mediated excitatory response, allowing IC neurons to integrate excitatory and inhibitory synaptic inputs arriving simultaneously. When GABAergic inputs arrive earlier than excitatory inputs, activation of GABA<sub>A</sub> receptors can drive the cell's membrane potential toward the GABA reversal potential by pre-opening Cl<sup>-</sup> channels (Wagner 1996). Therefore, the GABAergic response not only counteracts the AMPA component, but also suppresses the late and slow activation of the NMDA receptors. As shown in whole-cell patch clamp recordings (Ma et al. 2002a; Wu et al. 2004a), eliminating GABAergic inhibition enhanced the AMPA receptor mediated response and prolonged the NMDA receptor-mediated response, resulting in epileptic-like discharges (Fig. 10.4). Similar results have been found in ICX and ICD neurons (Smith 1992; Li et al. 1998). Thus, GABAergic inhibition normally suppresses NMDA receptor-mediated depolarization. Reducing this inhibition depolarizes the neuron over a longer time course and leads to epileptic-like discharges.

Iontophoresis of GABA or the GABA<sub>A</sub> agonist, muscimol, to IC neurons in brain slices of rat and gerbil pups (<14 days old) can elicit biphasic responses, that is, membrane de- and hyperpolarization. This is interesting because GABA usually acts on the GABA<sub>A</sub> receptor, hyperpolarizing the membrane potential. The membrane depolarization by GABA or muscimol in young IC neurons was accompanied by an increase in intracellular Cl<sup>-</sup> and Ca<sup>2+</sup> (Lo et al. 1998; Frech et al. 1999). The Ca<sup>2+</sup> influx activated by GABA was dependent on extracellular HCO<sub>3</sub><sup>-</sup> (Frech et al. 1999), which implies an HCO<sub>3</sub><sup>-</sup> efflux during GABA activation. Active Cl<sup>-</sup> uptake and HCO<sub>3</sub><sup>-</sup> efflux could shift the GABA reversal potential to a more positive level, which may account for the depolarizing effect of GABA during development. Increased intracellular Ca<sup>2+</sup> by GABA may be secondary to activation of voltage-dependent Ca<sup>2+</sup> channels, which can influence the growth and refinement of prospective inhibitory synaptic connections (Segal 1993).

The possibility that GABA<sub>B</sub> receptors are involved in synaptic transmission in the IC has been studied in *in vivo* and *in vitro* experiments. The GABA<sub>B</sub> receptor agonist, baclofen, did not affect membrane input resistance (Ma et al. 2002b), resting potential, or K<sup>+</sup> membrane conductance (Hosomi et al. 1997). These negative effects along with the short time course of the GABAergic IPSPs




Figure 10.4. Strong suppression of synaptic excitation by GABAergic inhibition in an ICC neuron. **(A)** In a brain slice perfused with normal saline, lateral lemniscal stimulation elicited a small early EPSP, an IPSP, and a late EPSP. **(B)** After strychnine ( $0.5 \mu M$ ; Stry) application there was little change in the synaptic response. **(C)** With bicuculline ( $10 \mu M$ ; Bic) the EPSP grew larger and longer, and multiple spikes appeared. **(D)** The late EPSP and associated action potentials were eliminated by CPP ( $20 \mu M$ ), the NMDA receptor antagonist, but an early EPSP and a short latency action potential were still present. **(E)** The early EPSP and its related spike were completely blocked by the AMPA receptor antagonist, NBQX ( $2 \mu M$ ). (Adapted from Ma et al. 2002a.)

and the IPSP reversal potential, which is close to the cell's resting potential, suggest that GABAergic synaptic transmission in the ICC is not mediated by metabotropic GABA<sub>B</sub> receptors. However, baclofen reduced the GABAergic IPSP amplitudes of ICC neurons in brain slices and phaclofen reversed this effect (Ma et al. 2002b). These results suggest that GABAergic inhibition is modulated by presynaptic GABA<sub>B</sub> receptors via regulation of GABA release (Ma et al. 2002b). GABA<sub>B</sub> receptors also affect synaptic plasticity. LTP induction in IC brain slices requires GABA<sub>B</sub> receptor activation (Zhang and Wu 2000). Electrophysiological studies in vivo show that GABA<sub>B</sub> receptors influence excitatory transmission in the IC (Faingold et al. 1991; Vaughn et al. 1996). Baclofen reduced sound-evoked responses, while GABA<sub>B</sub> receptor antagonists increased them. Systemic baclofen application reversed potentiation of click-evoked responses to intense sound (Szczepaniak and Møller 1996). A regulatory effect of GABA<sub>B</sub> receptors on glutamate release was proposed (Kelly and Zhang 2002) and tested in in vitro brain slices. EPSCs of IC neurons were reduced by baclofen, and this effect was reversed by the GABA<sub>B</sub> receptor antagonist, CGP 35348 (Wu et al. 2004b). Thus, GABA<sub>B</sub> receptors can modulate GABA<sub>A</sub> receptor-mediated inhibition as well as glutamatergic excitation in the IC.

### 5.3. OTHER SYNAPTIC RECEPTORS

#### 5.3.1. Glycine Receptors

Electrophysiological studies suggest that glycinergic inhibition plays a role analogous to GABAergic inhibition in shaping frequency-response areas, sharpening tuning for frequency modulation, sound duration coding, and binaural processing (Klug et al. 1995; Koch and Grothe 1998; Casseday et al. 2000; Le Beau et al. 2001; Lu and Jen 2001). However, differential effects for the GABA<sub>A</sub> and glycine receptor antagonists, bicuculline and strychnine, were reported. Excitatory tuning curves in some IC neurons in the horseshoe bat were changed by bicuculline, but not by strychnine (Vater et al. 1992). Bicuculline was more effective in abolishing the inhibitory frequency tuning curves and influencing the excitatory frequency tuning curves than strychnine (Lu and Jen 2001). Most of the bicuculline-sensitive cells were located more dorsally than those influenced by strychnine (Lu and Jen 2001), consistent with regional differences in GABA<sub>A</sub> and glycine receptor density (Sanes et al. 1987; Fubara et al. 1996). Physiological properties of glycine receptors resemble those of GABA<sub>A</sub> receptors in juvenile and mature IC neurons: activation of both GABA<sub>A</sub> and glycine receptors inhibits mature neurons by triggering Cl<sup>-</sup> influx and hyperpolarizing the cell's membrane. During embryonic and neonatal development, glycine and GABA depolarize IC neurons and increase intracellular Ca<sup>2+</sup> concentration (Ehrlich et al. 1999; Kotak and Sanes 2000).

However, IC glycine receptors develop more slowly than GABA<sub>A</sub> receptors. In immature rats (3 to 8 days postnatal) glycine receptors activate and deactivate significantly more slowly (Kraushaar and Backus 2002) than those in ventral



CN and spinal motor neurons (Jonas et al. 1998; Harty and Manis 1998). Slow kinetics may reflect immature glycine receptors with  $\alpha 2$  subunits (Takahashi et al. 1992; Singer et al. 1998). Slowly deactivating glycine receptors may prolong depolarization and enhance intracellular  $\text{Ca}^{2+}$  stores, perhaps affecting inhibitory synaptic ontogeny (Kraushaar and Backus 2002). Slowly developed glycine receptors may retard glycinergic relative to GABAergic neurotransmission, which matches the later arrival of glycinergic afferents from the lower auditory brain stem.

### 5.3.2. Serotonin Receptors

Serotonin (5-HT) modulates neural responses in bat to tone bursts and complex, behaviorally relevant sounds such as FM sweeps and species-specific vocalizations (Hurley et al. 2002). Iontophoretic application of serotonin onto IC neurons in vivo often depressed tone-evoked responses (Faingold et al. 1991; Hurley and Pollak 1999), facilitated or suppressed FM sweep responses, and had frequency-specific effects on response areas (Hurley and Pollak 2001). These effects are thought to be more complex than gain control because serotonin can change frequency tuning (Hurley et al. 2002). Autoradiographic labeling of the 5-HT<sub>1A</sub> receptor agonist shows the highest level of binding in the IC compared to other auditory brain stem nuclei (Thompson et al. 1994). The 5-HT<sub>1A</sub> and 5-HT<sub>2</sub> receptors are implicated in defensive behavioral responses after direct agonist injection in the auditory midbrain (Melo and Brandão 1995).

Serotonin acts on 5-HT receptor subtypes through various ionic mechanisms (Peroutka 1988; Bobker and Williams 1990), for example, activation of the 5-HT<sub>1A</sub> receptor hyperpolarizes the cell's membrane potential, which is mediated by increased conductance to  $\text{K}^+$  ions. The 5-HT<sub>1B</sub> and 5-HT<sub>1D</sub> receptors may have presynaptic inhibitory effects on glutamate, GABA, and serotonin release. Activation of the 5-HT<sub>2</sub> receptor has a depolarizing excitatory effect owing to decreased  $\text{K}^+$  conductance. Activating ionotropic 5-HT<sub>3</sub> receptors causes depolarization by directly opening the receptor for cations. Serotonin enhances the cationic current,  $I_h$  (which activates over hundreds of milliseconds on hyperpolarization), via undefined 5-HT receptor subtypes. Whether the complex effects of exogenous 5-HT on IC neurons reflect activation of different 5-HT receptor subtypes is not clear. Future studies should characterize the physiological properties of 5-HT receptor subtypes and determine regional differences in receptor physiology because 5-HT input and receptors are concentrated in the peripheral IC (Thompson et al. 1994; Hurley and Thompson 2001).

### 5.3.3. Acetylcholine Receptors

Nicotinic and muscarinic Ach receptors in IC are revealed by labeling receptors and mRNA of receptor subunits (Schwartz et al. 1982; Cortés and Palacios 1986; Regenold et al. 1989; Morley and Happe 2000). The role of Ach in IC physiology has been studied by iontophoretic application of Ach receptor agonists and antagonists in in vivo electrophysiological studies (Farley et al. 1983; Fain-

gold et al. 1991). Discharges elicited by the nonspecific agonists, Ach and carbachol, were enhanced in some neurons and decreased in others; temporal patterns, tuning curves, and rate-level functions were unaffected. Application of the nicotinic receptor antagonist, hexamethonium, enhanced activity in most cells and decreased tone-evoked activity in others. The nonspecific muscarinic antagonist atropine and the specific M1 receptor antagonist, pirenzepine, decreased responses in most neurons whereas the M2 receptor antagonist, gallamine, often increased activity. These results suggest that cholinergic inputs constitute a dynamic regulatory system, which controls the overall firing rate and modulates more specific excitatory and inhibitory circuits (Habbicht and Vater 1996). Perhaps Ach modulates acoustic responses via subtypes of Ach receptors that are differentially expressed pre- or postsynaptically. Further investigation should characterize physiological properties of Ach receptor subtypes in the IC.

In summary, excitatory synaptic transmission in the IC is glutamatergic via AMPA and NMDA receptors. The AMPA receptors subserve the early and fast part of the synaptic response, whereas NMDA receptors control the later and slower component. Fast inhibitory transmission is mediated by GABA and glycine via GABA<sub>A</sub> and glycine receptors. Glutamatergic excitation and GABAergic inhibition can be modulated by presynaptic GABA<sub>B</sub> receptors. Serotonin and Ach receptors have a regional modulatory role in synaptic transmission, perhaps through various receptor subtypes.

## 6. FUTURE DIRECTIONS

Understanding the biophysical profile of IC neurons has just begun. Only basic information about intrinsic electrical membrane properties and synaptic receptor characteristics is available. Much more work is needed in order to understand how the biophysical features of IC neurons contribute to auditory processing.

Ion channels that underlie the intrinsic membrane properties play a key role in shaping neural input–output functions. However, the output signal from a neuron embodies the cell's intrinsic electrical membrane properties and the activation of specific synaptic receptors. Therefore, synaptic inputs and synaptic receptors (or subtypes) in IC neurons with specific intrinsic membrane properties should be defined. Molecular–biologic study combined with physiological analysis of ion channels and synaptic receptors in IC neurons will drive such work. As a profile of subtypes and subunits of ion channels and synaptic receptors emerges, biophysical study of the channels and receptors will clarify the molecular mechanisms of auditory signal processing. Systematic studies of membrane and synaptic properties, combined with the anatomy of IC neurons in the three major subdivisions, are essential to understand how cell structure, physiological characteristics, and synaptic connections interact and are integrated.

## Abbreviations

Ach	acetylcholine
AHP	afterhyperpolarization
AMPA	$\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid
4-AP	4-aminopyridine
Bic	bicuculline
CN	cochlear nucleus
CNQX	6-cyano-7-nitroquinoxaline-2,3-dione
CPP	( $\pm$ )-3-(2-carboxypiperazin-4-yl)-propyl-1-phosphonic acid
CTX	charybdotoxin
DNLL	dorsal nucleus of the lateral lemniscus
DNQX	6,7-dinitroquinoxaline-2,3-dione
EPSC	excitatory postsynaptic current
EPSP	excitatory postsynaptic potential
FM	frequency-modulated
GABA	$\gamma$ -aminobutyric acid
GIRK1	G-protein gated inward rectifier K <sup>+</sup> channel
5-HT	5-hydroxytryptamine (serotonin)
IC	inferior colliculus
ICC	central nucleus of the inferior colliculus
ICD	dorsal cortex of the inferior colliculus
ICX	external nucleus of the inferior colliculus
IPSC	inhibitory postsynaptic current
IPSP	inhibitory postsynaptic potential
K <sub>IR</sub>	inward rectifier K <sup>+</sup> channel
LSO	lateral superior olive
LTP	long-term potentiation
MNTB	medial nucleus of the trapezoid body
MSO	medial superior olive
NBQX	1,2,3,4-tetrahydro-6-nitro-2,3-dioxo-benzo[f]quinoxaline-7-sulfonamide
NMDA	<i>N</i> -methyl-D-aspartate
SAM	sinusoidal amplitude modulation
Stry	strychnine
TEA	tetraethylammonium
VNLL	ventral nucleus of the lateral lemniscus

## Acknowledgments

This work was supported by research grants from the Natural Sciences and Engineering Research Council (NSERC) of Canada and the Hearing Foundation of Canada. The author thanks Dr. J.A. Winer and Dr. J.B. Kelly for careful reading of the manuscript and helpful comments and suggestions.

## References

- Bal R, Green GGR, Rees A, and Sanders DJ (2002) Firing patterns of inferior colliculus neurons—histology and mechanism to change firing patterns in rat brain slices. *Neuroscience Letters* **317**:42–46.
- Barnard EA, Burnstock G, and Webb TE (1994) G-protein-coupled receptors for ATP and other nucleotides: a new receptor family. *Trends in Pharmacological Sciences* **15**: 67–70.
- Basta D and Vater M (2003) Membrane-based gating mechanism for auditory information in the mouse inferior colliculus. *Brain Research* **968**:171–178.
- Bausch SB, Patterson TA, Ehrenguber MU, Lester HA, Davidson N, and Chavkin C (1995) Colocalization of  $\mu$  opioid receptors with GIRK1 potassium channels in the rat brain: an immunocytochemical study. *Receptors and Channels* **3**:221–241.
- Blaine JT and Ribera AB (1998) Heteromultimeric potassium channels formed by members of the Kv2 subfamily. *Journal of Neuroscience* **18**:9585–9593.
- Bobker DH and Williams JT (1990) Ion conductances affected by 5-HT receptor subtypes in mammalian neurons. *Trends in Neurosciences* **13**:169–173.
- Brand A, Urban R, and Grothe B (2000) Duration tuning in the mouse auditory midbrain. *Journal of Neurophysiology* **84**:1790–1799.
- Brew HM and Forsythe ID (1995) Two voltage-dependent  $K^+$  conductances with complementary functions in postsynaptic integration at a central auditory synapse. *Journal of Neuroscience* **15**:8011–8022.
- Burger RM and Pollak GD (1998) Analysis of the role of inhibition in shaping responses to sinusoidally amplitude-modulated signals in the inferior colliculus. *Journal of Neurophysiology* **80**:1686–1701.
- Casseday JH, Ehrlich D, and Covey E (1994) Neural tuning for sound duration: role of inhibitory mechanisms in the inferior colliculus. *Science* **264**:847–850.
- Casseday JH, Ehrlich D, and Covey E (2000) Neural measurement of sound duration: control by excitatory-inhibitory interactions in the inferior colliculus. *Journal of Neurophysiology* **84**:1475–1487.
- Chen GD (1998) Effects of stimulus duration on responses of neurons in the chinchilla inferior colliculus. *Hearing Research* **122**:142–150.
- Coetzee WA, Amarillo Y, Chiu J, Chow A, Lau D, McCormack T, Moreno H, Nadal MS, Ozaita A, Pountney D, Saganich M, Vega-Saenz de Miera E, and Rudy B (1999) Molecular diversity of  $K^+$  channel. *Annals of the New York Academy of Sciences* **868**: 233–285.
- Connor JA and Stevens CF (1971) Voltage clamp studies of a transient outward membrane current in gastropod neural somata. *Journal of Physiology (London)* **213**: 21–30.
- Cortés R and Palacios JM (1986) Muscarinic cholinergic receptor subtypes in rat brain. I. Quantitative autoradiographic studies. *Brain Research* **362**:227–238.
- Covey E, Kauer JA, and Casseday JH (1996) Whole-cell patch-clamp recording reveals subthreshold sound-evoked postsynaptic currents in the inferior colliculus of awake bats. *Journal of Neuroscience* **16**:3009–3018.
- Critz SD, Wible BA, Lopez HS, and Brown AM (1993) Stable expression and regulation of a rat brain  $K^+$  channel. *Journal of Neurochemistry* **60**:1175–1178.
- Dodson PD, Barker MC, and Forsythe ID (2002) Two heteromeric Kv1 potassium channels differentially regulate action potential firing. *Journal of Neuroscience* **22**:6953–6961.

- Ehrlich D, Casseday JH, and Covey E (1997) Neural tuning to sound duration in the inferior colliculus of the big brown bat, *Eptesicus fuscus*. *Journal of Neurophysiology* **77**:2360–2372.
- Ehrlich I, Löhre S, and Friauf E (1999) Shift from depolarizing to hyperpolarizing glycine action in rat auditory neurones is due to age-dependent  $\text{Cl}^-$  regulation. *Journal of Physiology (London)* **520**:121–137.
- Faingold CL (2002) Role of GABA abnormalities in the inferior colliculus pathophysiology – audiogenic seizures. *Hearing Research* **168**:223–237.
- Faingold CL, Gehlbach G, and Caspary DM (1989) On the role of GABA as an inhibitory neurotransmitter in inferior colliculus neurons: iontophoretic studies. *Brain Research* **500**:302–312.
- Faingold CL, Gehlbach G, and Caspary DM (1991) Functional pharmacology of inferior colliculus neurons. In: Altschuler RA, Bobbin RP, Clopton BM, and Hoffman DW (eds). *Neurobiology of Hearing. The Central Auditory System*. Raven Press, New York, pp. 223–251.
- Faingold CL, Marcinczyk MJ, Casebeer DJ, Randall ME, Arnerić SP, and Browning RA (1994a) GABA in the inferior colliculus plays a critical role in control of audiogenic seizures. *Brain Research* **640**:40–47.
- Faingold CL, Randall ME, and Anderson CA (1994b) Blockade of GABA uptake with tiagabine inhibits audiogenic seizures and reduces neuronal firing in the inferior colliculus of the genetically epilepsy-prone rat. *Experimental Neurology* **126**:225–232.
- Farley GR, Foley BJ, Javel E, and Gorga MP (1983) Single-unit responses to cholinergic agents in the rat inferior colliculus. *Hearing Research* **11**:73–91.
- Feldman DE and Knudsen EI (1998) Pharmacological specialization of learned auditory responses in the inferior colliculus of the barn owl. *Journal of Neuroscience* **18**:3073–3087.
- Frech MJ, Deitmer JW, and Backus KH (1999) Intracellular chloride and calcium transients evoked by  $\gamma$ -aminobutyric acid and glycine in neurons of the rat inferior colliculus. *Journal of Neurobiology* **40**:386–396.
- Fubara BM, Casseday JH, Covey E, and Schwartz-Bloom RD (1996) Distribution of  $\text{GABA}_A$ ,  $\text{GABA}_B$ , and glycine receptors in the central auditory system of the big brown bat, *Eptesicus fuscus*. *Journal of Comparative Neurology* **369**:83–92.
- Fuzessery ZM and Hall JC (1996) Role of GABA in shaping frequency tuning and creating FM sweep selectivity in the inferior colliculus. *Journal of Neurophysiology* **76**:1059–1073.
- Golding NL, Ferragamo MJ, and Oertel D (1999) Role of intrinsic conductances underlying responses to transients in octopus cells of the cochlear nucleus. *Journal of Neuroscience* **19**:2897–2905.
- Grigg JJ, Brew HM, and Tempel BL (2000) Differential expression of voltage-gated potassium channel genes in auditory nuclei of the mouse brainstem. *Hearing Research* **140**:77–90.
- Habbicht H and Vater M (1996) A microiontophoretic study of acetylcholine effects in the inferior colliculus of horseshoe bats: implications for a modulatory role. *Brain Research* **724**:169–179.
- Harty TP and Manis PB (1998) Kinetic analysis of glycine receptor currents in ventral cochlear nucleus. *Journal of Neurophysiology* **79**:1891–1901.
- Hestrin S, Nicoll RA, Perkel DJ, and Sah P (1990) Analysis of excitatory synaptic action in pyramidal cells using whole-cell recording from rat hippocampal cell. *Journal of Physiology (London)* **422**:203–225.

- Hirsch JA and Oertel D (1988) Intrinsic properties of neurones in the dorsal cochlear nucleus of mice, *in vitro*. *Journal of Physiology (London)* **396**:535–548.
- Hosomi H, Mori M, Amatsu M, and Okada Y (1997) GABA-activated conductance in cultured rat inferior colliculus neurons. *Journal of Neurophysiology* **77**:994–1002.
- Hugues M, Romey G, Duval D, Vincent JP, and Lazdunski M (1982) Apamin as a selective blocker of the calcium-dependent potassium channel in neuroblastoma cells: voltage-clamp and biochemical characterization of the toxin receptor. *Proceedings of the National Academy of Sciences of the United States of America* **79**:1308–1312.
- Hurley LM and Pollak GD (1999) Serotonin differentially modulates responses to tones and frequency-modulated sweeps in the inferior colliculus. *Journal of Neuroscience* **19**:8071–8082.
- Hurley LM and Pollak GD (2001) Serotonin effects on frequency tuning of inferior colliculus neurons. *Journal of Neurophysiology* **85**:828–842.
- Hurley LM and Thompson AM (2001) Serotonergic innervation of the auditory brainstem of the Mexican free-tailed bat, *Tadarida brasiliensis*. *Journal of Comparative Neurology* **435**:78–88.
- Hurley LM, Thompson AM, and Pollak GD (2002) Serotonin in the inferior colliculus. *Hearing Research* **168**:1–11.
- Hwang PM, Fotuhi M, Brecht DS, Cunningham AM, and Snyder SH (1993) Contrasting immunohistochemical localizations in rat brain of two novel K<sup>+</sup> channels of the *Shab* subfamily. *Journal of Neuroscience* **13**:1569–1576.
- Ikeuchi Y and Nishizaki T (1995a) ATP-evoked potassium currents in rat striatal neurons are mediated by a P<sub>2</sub> purinergic receptor. *Neuroscience Letters* **190**:89–92.
- Ikeuchi Y and Nishizaki T (1995b) The P<sub>2Y</sub> purinoceptor-operated potassium channel is possibly regulated by the  $\beta\gamma$  subunits of a pertussis toxin-insensitive G-protein in cultured rat inferior colliculus neurons. *Biochemical and Biophysical Research Communications* **214**:589–596.
- Jonas P, Bischofberger J, and Sandkühler J (1998) Corelease of two fast neurotransmitters at a central synapse. *Science* **281**:419–424.
- Kanold PO and Manis PB (1999) Transient potassium currents regulate the discharge patterns of dorsal cochlear nucleus pyramidal cells. *Journal of Neuroscience* **19**: 2195–2208.
- Karschin C, Schreibmayer W, Dascal N, Lester H, Davidson N, and Karschin A (1994) Distribution and localization of a G-protein-coupled inwardly rectifying K<sup>+</sup> channel in the rat. *Federation of European Biochemical Society Letters* **348**:139–144.
- Kelly JB and Zhang H (2002) Contribution of AMPA and NMDA receptors to excitatory responses in the inferior colliculus. *Hearing Research* **168**:35–42.
- Kirson ED and Yaari Y (2000) Unique properties of NMDA receptors enhances synaptic excitation of radiatum giant cells in rat hippocampus. *Journal of Neuroscience* **20**: 4844–4854.
- Kirson ED, Schirra C, Konnerth A, and Yaari Y (1999) Early postnatal switch in magnesium sensitivity of NMDA receptors in rat CA1 pyramidal cells. *Journal of Physiology (London)* **521**:99–111.
- Klug A, Park TJ, and Pollak GD (1995) Glycine and GABA influence binaural processing in the inferior colliculus of the mustache bat. *Journal of Neurophysiology* **74**:1701–1713.
- Koch U and Grothe B (1998) GABAergic and glycinergic inhibition sharpens tuning for frequency modulations in the inferior colliculus of the big brown bat. *Journal of Neurophysiology* **80**:71–82.

- Kotak VC and Sanes DH (2000) Long-lasting inhibitory synaptic depression is age- and calcium-dependent. *Journal of Neuroscience* **20**:5820–5826.
- Kraushaar U and Backus KH (2002) Characterization of GABA<sub>A</sub> and glycine receptors in neurons of the developing rat inferior colliculus. *Pflügers Archiv European Journal of Physiology* **445**:279–288.
- Kuner T and Schoepfer R (1996) Multiple structural elements determine subunit specificity of Mg<sup>2+</sup> block in NMDA receptor channels. *Journal of Neuroscience* **16**:3549–3558.
- Kuwada S and Batra R (1999) Coding of sound envelopes by inhibitory rebound in neurons of the superior olivary complex in the unanesthetized rabbit. *Journal of Neuroscience* **19**:2273–2287.
- Kuwada S, Batra R, Yin TCT, Oliver DL, Haberly LB, and Stanford TR (1997) Intracellular recordings in response to monaural and binaural stimulation of neurons in the inferior colliculus of the cat. *Journal of Neuroscience* **17**:7565–7581.
- Le Beau FEN, Rees A, and Malmierca MS (1996) Contribution of GABA- and glycine-mediated inhibition to the monaural temporal response properties of neurons in the inferior colliculus. *Journal of Neurophysiology* **75**:902–919.
- Le Beau FEN, Malmierca MS, and Rees A (2001) Iontophoresis *in vivo* demonstrates a key role for GABA<sub>A</sub> and glycinergic inhibition in shaping frequency response areas in the inferior colliculus of guinea pig. *Journal of Neuroscience* **21**:7303–7312.
- Li Y, Evans MS, and Faingold CL (1998) *In vitro* electrophysiology of neurons in sub-nuclei of rat inferior colliculus. *Hearing Research* **121**:1–10.
- Li Y, Evans MS, and Faingold CL (1999) Synaptic response patterns of neurons in the cortex of rat inferior colliculus. *Hearing Research* **137**:15–28.
- Li W, Kaczmarek LK, and Perney TM (2001) Localization of two high-threshold potassium channel subunits in the rat central auditory system. *Journal of Comparative Neurology* **437**:196–218.
- Liu SQ and Kaczmarek LK (1998) Depolarization selectively increases the expression of the Kv3.1 potassium channel in developing inferior colliculus neurons. *Journal of Neuroscience* **18**:8758–8769.
- Lo Y-J, Rao SC, and Sanes DH (1998) Modulation of calcium by inhibitory systems in the developing auditory midbrain. *Neuroscience* **83**:1075–1084.
- Lu Y and Jen PH-S (2001) GABAergic and glycinergic neural inhibition in excitatory frequency tuning of bat inferior collicular neurons. *Experimental Brain Research* **141**:331–339.
- Ludwig J, Weseloh R, Karschin C, Liu Q, Netzer R, Engeland B, Stansfeld C, and Pongs O (2000) Cloning and functional expression of rat eag2, a new member of the *ether-à-go-go* family of potassium channels and comparison of its distribution with that of eag1. *Molecular and Cellular Neuroscience* **16**:59–70.
- Luneau CJ, Williams JB, Marshall J, Levitan ES, Oliva C, Smith JS, Antanavage J, Folander K, Stein RB, Swanson R, Kaczmarek LK, and Buhrow SA (1991) Alternative splicing contributes to K<sup>+</sup> channel diversity in the mammalian central nervous system. *Proceedings of the National Academy of Sciences of the United States of America* **88**:3932–3936.
- Ma CL, Kelly JB, and Wu SH (2002a) AMPA and NMDA receptors mediate synaptic excitation in the rat's inferior colliculus. *Hearing Research* **168**:25–34.
- Ma CL, Kelly JB, and Wu SH (2002b) Presynaptic modulation of GABAergic inhibition by GABA<sub>B</sub> receptors. *Neuroscience* **114**:207–215.
- Manis PB and Marx SO (1991) Outward currents in isolated ventral cochlear nucleus neurons. *Journal of Neuroscience* **11**:2865–2880.

- Marianowski R, Pollard H, Moreau J, Després G, Ben Ari Y, Tran Ba Huy P, and Romand R (1995) *N*-methyl-D-aspartate receptor subunits NR1 and NR2C are overexpressed in the inferior colliculus of audiogenic mice. *Neuroscience Letters* **189**:190–194.
- Mayer ML, Westbrook GL, and Guthrie PB (1984) Voltage-dependent block by  $Mg^{++}$  of NMDA responses in spinal cord neurones. *Nature* **309**:261–263.
- Melo LL and Brandão ML (1995) Involvement of 5-HT<sub>1A</sub> and 5-HT<sub>2</sub> receptors of the inferior colliculus in aversive states induced by exposure of rats to the elevated plus-maze test. *Behavioural Pharmacology* **6**:413–417.
- Meyer R and Heinemann SH (1998) Characterization of an *eag*-like potassium channel in human neuroblastoma cells. *Journal of Physiology (London)* **508**:49–56.
- Millan MH, Meldrum BS, and Faingold CL (1986) Induction of audiogenic seizure susceptibility by focal infusion of excitant amino acid or bicuculline into the inferior colliculus of normal rats. *Experimental Neurology* **91**:634–639.
- Moore DR, Kotak VC, and Sanes DH (1998) Commissural and lemniscal synaptic input to the gerbil inferior colliculus. *Journal of Neurophysiology* **80**:2229–2236.
- Morley BJ and Happe HK (2000) Cholinergic receptors: dual roles in transduction and plasticity. *Hearing Research* **147**:104–112.
- N'Gouemo P and Rittenhouse AR (2000) Biophysical and pharmacological characterization of voltage-sensitive calcium currents in neonatal rat inferior colliculus neurons. *Neuroscience* **96**:753–765.
- Oertel D (1997) Encoding of timing in the brain stem auditory nuclei of vertebrates. *Neuron* **19**:959–962.
- Palombi PS and Caspary DM (1996) GABA inputs control discharge rate primarily within frequency receptive fields of inferior colliculus neurons. *Journal of Neurophysiology* **75**:2211–2219.
- Park TJ and Pollak GD (1993) GABA shapes sensitivity to interaural intensity disparities in the mustache bat's inferior colliculus: implications for encoding sound location. *Journal of Neuroscience* **13**:2050–2067.
- Park TJ and Pollak GD (1994) Receptive fields in the inferior colliculus of the mustache bat are shaped by GABAergic inhibition. *Journal of Neurophysiology* **72**:1080–1102.
- Perney TM and Kaczmarek LK (1997) Localization of a high threshold potassium channel in the rat cochlear nucleus. *Journal of Comparative Neurology* **386**:178–202.
- Perney TM, Marshall J, Martin KA, Hockfield S, and Kaczmarek LK (1992) Expression of the mRNAs for the Kv3.1 potassium channel gene in the adult and developing rat brain. *Journal of Neurophysiology* **68**:756–766.
- Peroutka SJ (1988) 5-Hydroxytryptamine receptor subtypes: molecular, biochemical and physiological characterization. *Trends in Neurosciences* **11**:496–500.
- Peruzzi D, Sivaramakrishnan S, and Oliver DL (2000) Identification of cell types in brain slices of the inferior colliculus. *Neuroscience* **101**:403–416.
- Pierson MG, Smith KL, and Swann JW (1989) A slow NMDA-mediated synaptic potential underlies seizures originating from midbrain. *Brain Research* **486**:381–386.
- Pollak GD and Park TJ (1993) The effects of GABAergic inhibition on monaural response properties of neurons in the mustache bat's inferior colliculus. *Hearing Research* **65**:99–117.
- Reetz G and Ehret G (1999) Inputs from three brainstem sources to identified neurons in the mouse inferior colliculus slice. *Brain Research* **816**:527–543.



- Regenold W, Araujo DM, and Quirion R (1989) Quantitative autoradiographic distribution of [<sup>3</sup>H]AF-DX116 muscarinic M<sub>2</sub> receptor binding sites in rat brain. *Synapse* **4**: 115–125.
- Richardson FC and Kaczmarek LK (2000) Modification of delayed rectifier potassium currents by the Kv9.1 potassium channel subunit. *Hearing Research* **147**:21–30.
- Rosenberger MH, Fremouw T, Casseday JH, and Covey E (2003) Expression of the Kv1.1 ion channel subunit in the auditory brainstem of the big brown bat, *Eptesicus fuscus*. *Journal of Comparative Neurology* **462**:101–120.
- Rudy B, Chow A, Lau D, Amarillo Y, Ozaita A, Saganich M, Moreno H, Nadal MS, Hernandez-Pineda R, Hernandez-Cruz A, Erisir A, Leonard C, and Vega-Saenz de Miera E (1999) Contributions of Kv3 channel to neuronal excitability. *Annals of the New York Academy of Sciences* **868**:304–343.
- Sanes DH, Geary WA, Wooten GF, and Rubel EW (1987) Quantitative distribution of the glycine receptor in the auditory brain stem of the gerbil. *Journal of Neuroscience* **7**:3793–3802.
- Sato K, Shiraishi S, Nakagawa H, Kuriyama H, and Altschuler RA (2000) Diversity and plasticity in amino acid receptor subunits in the rat auditory brain stem. *Hearing Research* **147**:137–144.
- Schwartz RD, McGeer R, and Kellar KJ (1982) Nicotinic cholinergic receptors labelled by [<sup>3</sup>H]acetylcholine in rat brain. *Molecular Pharmacology* **22**:56–62.
- Segal M (1993) Calcium and neuronal plasticity. *Israel Journal of Medical Sciences* **29**: 543–548.
- Serôdio P and Rudy R (1998) Differential expression of Kv4 K<sup>+</sup> channel subunits mediating subthreshold transient K<sup>+</sup> (A-type) currents in rat brain. *Journal of Neurophysiology* **79**:1081–1091.
- Singer JH, Talley EM, Bayliss DA, and Berger AJ (1998) Development of glycinergic synaptic transmission to rat brain stem motoneurons. *Journal of Neurophysiology* **80**: 2608–2620.
- Sivaramakrishnan S and Oliver DL (2001) Distinct K currents result in physiologically distinct cell types in the inferior colliculus of the rat. *Journal of Neuroscience* **21**: 2861–2877.
- Smith PH (1992) Anatomy and physiology of multipolar cells in the rat inferior colliculus cortex using the in vitro brain slice technique. *Journal of Neuroscience* **12**:3700–3715.
- Song WJ (2002) Genes responsible for native depolarization-activated K<sup>+</sup> currents in neurons. *Neuroscience Research* **42**:7–14.
- Svirskis G, Kotak V, Sanes DH, and Rinzel J (2002) Enhancement of signal-to-noise ratio and phase locking for small inputs by a low-threshold outward current in auditory neurons. *Journal of Neuroscience* **22**:11019–11025.
- Szczepaniak WS and Møller AR (1996) Effects of (–)-baclofen, clonazepam, and diazepam on tone exposure-induced hyperexcitability of the inferior colliculus in the rat: possible therapeutic implications for pharmacological management of tinnitus and hyperacusis. *Hearing Research* **97**:46–53.
- Takahashi T, Momiyama A, Hirai K, Hishinuma F, and Akagi H (1992) Functional correlation of fetal and adult forms of glycine receptors with developmental changes in inhibitory synaptic receptor channels. *Neuron* **9**:1155–1161.
- Terlau H, Ludwig J, Steffan R, Pong O, Stühmer W, and Heinemann SH (1996) Extracellular Mg<sup>++</sup> regulates activation of rat *eag* potassium channel. *Pflügers Archiv European Journal of Physiology* **432**:301–312.

- Thompson GC, Thompson AM, Garrett KM, and Britton BH (1994) Serotonin and serotonin receptors in the central auditory system. *Otolaryngology Head and Neck Surgery* **110**: 93–102.
- Vater M, Habbicht H, Kössl M, and Grothe B (1992) The functional role of GABA and glycine in monaural and binaural processing in the inferior colliculus of horseshoe bats. *Journal of Comparative Physiology A* **171**:541–553.
- Vaughn MD, Pozza MF, and Lingenhöhl K (1996) Excitatory acoustic responses in the inferior colliculus of the rat are increased by GABA<sub>B</sub> receptor blockade. *Neuropharmacology* **35**:1761–1767.
- Vergara C, Latorre R, Marrion NV, and Adelman JP (1998) Calcium-activated potassium channels. *Current Opinion in Neurobiology* **8**:321–329.
- Wagner T (1994) Intrinsic properties of identified neurons in the central nucleus of mouse inferior colliculus. *NeuroReport* **30**:89–93.
- Wagner T (1996) Lemniscal input to identified neurons of the central nucleus of mouse inferior colliculus: an intracellular brain slice study. *European Journal of Neuroscience* **8**:1231–1239.
- Wang LY, Gan L, Forsythe ID, and Kaczmarek LK (1998) Contribution of the Kv3.1 potassium channel to high-frequency firing in mouse auditory neurones. *Journal of Physiology (London)* **509**:183–194.
- Weiser M, Vega-Saenz de Miera E, Kentros C, Moreno H, Franzen L, Hillman D, Baker H, and Rudy B (1994) Differential expression of *Shaw*-related K<sup>+</sup> channels in the rat central nervous system. *Journal of Neuroscience* **14**:949–972.
- Wu SH and Kelly JB (1991) Physiological properties of neurons in the mouse superior olive: membrane characteristics and postsynaptic responses studied in vitro. *Journal of Neurophysiology* **65**:230–246.
- Wu SH and Kelly JB (1993) Response of neurons in the lateral superior olive and medial nucleus of the trapezoid body to repetitive stimulation: intracellular and extracellular recordings from mouse brain slice. *Hearing Research* **68**:189–201.
- Wu SH and Kelly JB (1995) In vitro brain slice studies of the rat's dorsal nucleus of the lateral lemniscus. I. Membrane and synaptic response properties. *Journal of Neurophysiology* **73**:780–793.
- Wu SH and Oertel D (1984) Intracellular injection with horseradish peroxidase of physiologically characterized stellate and bushy cells in slices of mouse anteroventral cochlear nucleus. *Journal of Neuroscience* **4**:1577–1588.
- Wu SH, Ma CL, Sivaramakrishnan S, and Oliver DL (2002) Synaptic modification in neurons of the central nucleus of the inferior colliculus. *Hearing Research* **168**:43–54.
- Wu SH, Ma CL, and Kelly JB (2004a) Contribution of AMPA, NMDA and GABA<sub>A</sub> receptors to temporal pattern of postsynaptic responses in rat's inferior colliculus. *Journal of Neuroscience* **24**:4625–4634.
- Wu SH, Ma CL, and Kelly JB (2004b) GABA<sub>B</sub> receptors modulate synaptic excitation in the rat's central and dorsal nuclei of the inferior colliculus. *Association for Research in Otolaryngology Abstracts* **27**:600.
- Yang L, Pollak GD, and Resler C (1992) GABAergic circuits sharpen tuning curves and modify response properties in the mustache bat's inferior colliculus. *Journal of Neurophysiology* **68**:1760–1774.
- Yasuda S, Ishida N, Higashiyama A, Morinobu S, and Kato N (2000) Characterization of audiogenic-like seizures in naive rats evoked by activation of AMPA and NMDA receptors in the inferior colliculus. *Experimental Neurology* **164**:396–406.

- Zhang H and Kelly JB (2001) AMPA and NMDA receptors regulate responses of neurons in the rat's inferior colliculus. *Journal of Neurophysiology* **86**:871–880.
- Zhang Y and Wu SH (2000) Long-term potentiation in the inferior colliculus studied in rat brain slice. *Hearing Research* **147**:92–103.
- Zhang H, Xu J, and Feng AS (1999) Effects of GABA-mediated inhibition and direction-dependent frequency tuning in the frog inferior colliculus. *Journal of Comparative Physiology A* **184**:92–103.

# Chapter 11

## Spectral and Intensity Coding in the Auditory Midbrain

GÜNTER EHRET AND CHRISTOPH E. SCHREINER

### 1. INTRODUCTION

This overview emphasizes general properties of spectral and intensity processing in the inferior colliculus (IC) common to mammals such as the cat, mouse, rat, guinea pig, Mongolian gerbil, chinchilla, rhesus monkey, and various species of bats. Considerable data are available for the domestic cat (*Felis catus*) and mice (*Mus musculus*, laboratory strains) so that these species can serve as representative unless specified otherwise (for special spectral processing regimens in bats see Chapters 14 and 17). Functional distinctions exist among IC subdivisions (Aitkin 1986). Accordingly, we employ a basic anatomical framework of three major divisions: the central nucleus (ICC), the external nucleus (EN), and the dorsal cortex (DC). In this framework, the ICC contains layer IV of the DC as defined in Golgi studies of the cat (Morest and Oliver 1984) and mouse (Meininger et al. 1986).

#### *1.1. PRINCIPLES OF SPECTRAL ANALYSIS AND CODING*

A defining step for the organization of the entire auditory system is the spectral decomposition of the acoustic waveform in the cochlea (Patuzzi 1996). All subsequent operations directly reflect this peripheral process or are a consequence of specific central auditory filtering and transformations acting on the residue of this initial process. The orderly frequency representation along the basilar membrane, the tonotopy, is the basis of the concomitant order for characteristic frequency (CF) found in many nuclei and centers of the auditory pathways, including the ICC and extending to the primary auditory cortex. Thus, the cochlear tonotopy is the first and most general principle of spectral analysis and coding. Although the cochlear transformation of the signal is not identical to a Fourier analysis, the parallels between them are sufficiently compelling to warrant analysis of the resulting transmission channels as a bank of parallel spectral filters. Therefore, many characterizations of auditory neuronal elements are cast in the language of frequency filtering, using metrics such as center frequency,

frequency selectivity or filter bandwidth, and response magnitude. The response timing of the individual frequency channels—reflected in phase-locking of action potentials to individual cycles of a frequency—also affects auditory processing, but is confined to relatively low frequencies, and is not considered further. Here, we deal with properties of frequency filtering of IC neurons. A profile of the spectral receptive field (SRF) is a first analytic step in characterizing a neuron type or a nucleus.

## 1.2. PRINCIPLES OF INTENSITY ANALYSIS AND CODING

Relative and absolute stimulus magnitudes are two parameters in the spectral analysis of sounds with profound influence on auditory processing and perception. Mean sound pressure level (SPL) may influence neural responses in three ways: it can change spectral filter width, alter neural response strength in terms of firing rate, and modify temporal response parameters such as latency and precision of action potential timing relative to the stimulus waveform. Intensity effects on filter bandwidth are closely related to spectral receptive-field shape. Response magnitude effects are commonly discussed in terms of rate-level functions (RLs), that is, the total number of spikes/stimulus plotted against mean stimulus level. From RLs, measures of response threshold, intensity selectivity, and dynamic range can be derived.

Another aspect of stimulus magnitude is the amplitude distribution across the sound spectrum. The relative intensity of spectral minima and maxima, or the “contrast” with regard to their mean amplitudes, and the time course of these amplitude changes across the spectrum, must be encoded and represented. It is evident that amplitude coding in a complex spectral environment, the most natural listening condition, is inseparable from coding in the spectral domain.

## 2. SPECTRAL RECEPTIVE-FIELD PROPERTIES

### 2.1. EXCITATORY/INHIBITORY/FACILITATORY AREAS

Traditionally, spectrally simple sounds that are readily parameterized and manipulated are used for the exploration of receptive-field (RF) characteristics. Pure tones of various frequency and intensity combinations can demonstrate the excitatory SRF (eSRF) by identifying the parameter combinations that elevate firing rate above the spontaneous activity. The border of the eSRF represents the excitatory frequency tuning curve (eFTC) (Fig. 11.1A) and its shape is assessed by several measures:

- Characteristic frequency (CF): frequency producing a response at the lowest sound pressure levels (SPL)
- Best frequency (BF): frequency evoking the most spikes at a given SPL (CF and BF are not always identical)
- Minimum threshold (MT): SPL at CF




Figure 11.1. Spectral receptive fields (*SRFs*) of neurons in the central nucleus of the inferior colliculus (ICC). (A) Map of the *SRF* of a mouse neuron. Spike rates are shown as short vertical bars. The excitatory frequency tuning curve (*eFTC*) borders an area of increased responsiveness (above the spontaneous activity), the excitatory *SRF* (*eSRF*). Shaded gray areas: inhibitory *SRFs* (*iSRF*; suppressed spontaneous activity). (M.A. Ego-

- Bandwidth (BW): frequency width of FTC at various levels above MT
- Quality factor ( $Q$ ): the relative sharpness of tuning CF/BW at a fixed stimulus level (e.g.,  $Q/10\text{dB}$  is based on BW 10 dB above MT)
- The slopes of the low- and the high-frequency sides of the eFTC (expressed in dB/octave); also related to sharpness of tuning

A tone at a frequency and/or level outside the eSRF may have no effect on response rate or decrease it below spontaneous activity. The latter effect is called “one-tone inhibition” and reveals inhibitory areas surrounding the eSRF (Fig. 11.1A, B) if the spontaneous activity is sufficient to reliably detect an activity decrease (Suga 1964; Willott and Urban 1978; Wang et al. 1996; Ramachandran et al. 1999).

A more comprehensive approach uses a “two-tone paradigm” (e.g., Egorova et al. 2001): one tone excites the neuron at its CF (10 to 20 dB above threshold) while a second tone of variable frequency and level is presented simultaneously, with or shortly before the CF tone. This second tone can increase excitation to the constant CF tone (summation or facilitation of excitation; Fig. 11.1D), decrease or suppress the response to the CF tone (inhibition; Fig. 11.1C, D), or it can have no visible effect. Thus, the two-tone paradigm can identify the superposition of inhibitory and facilitatory receptive fields (iSRFs, fSRFs, respectively) with the eSRF. Inhibitory areas outside the eFTC are called “inhibitory side bands.” The shapes of iSRFs and fSRFs are quantified by similar measures as eSRFs and eFTCs.

Facilitatory SRFs have rarely been studied (Fuzessery and Feng 1983; Ehret and Merzenich 1988a; Egorova et al. 2001; Jen et al. 2002) except in the highly specialized mustached bat IC (Mittman and Wenstrup 1995; Wenstrup et al. 1999; Portfors and Wenstrup 2002) (see Chapter 17). Few cells show spectral facilitation (about 11% in the mouse, 18% in the big brown bat). In the mouse, facilitation is often found within about 1 octave of CF (Egorova et al. 2001).

---

Figure 11.1. *Continued*

rova and G. Ehret, unpublished observations). **(B)** Response rates to tone bursts relative to spontaneous activity (horizontal lines at five sound levels) at various frequency-level combinations indicate a narrow excitatory area (*black*) around the characteristic frequency (*CF*) bordered by inhibitory areas (*gray*) of a cat type I neuron. Lines demarcate the borders of the eSRF and the two iSRFs. (Modified from Ramachandran et al. 1999.) **(C, D)** Spike rates (*black bars*) of two mouse neurons in response to *(a)* tone bursts of various frequency-level combinations, *(b)* combinations of two tones, one at the CF and 10 dB above minimum threshold, the other at various frequency-level combinations. (Modified from Egorova et al. 2001.) *(c)* eSRFs are marked in various gray levels (proportional to response rate); iSRFs are marked with different hatching reflecting grades of inhibitory strength. The *black area* in **(C)** indicates a facilitatory SRF (fSRF).  $CF_E$ , Excitatory CF;  $CF_{IL}$ ,  $CF_{IH}$ , inhibitory CFs below and above the  $CF_E$ , respectively. The neuron in **(D)** has two excitatory CFs,  $CF_{EH}$  and  $CF_{EL}$ , both separated by a central inhibitory area with a  $CF_{IC}$ . *dB att*, Attenuation steps in dB; *MT*, minimum threshold.

Facilitation in the IC indicates that spectrally related tones can enhance responses.

## 2.2. SHAPES OF EXCITATORY AND INHIBITORY RECEPTIVE FIELDS

Basic differences have been found between IC subdivisions with regard to SRF and FTC shapes. In the DC and especially the EN, the frequency selectivity is low, with broad eFTCs, low  $Q$ -values, and irregularly shaped eFTCs (Fig. 11.2) (Aitkin et al. 1975, 1978; Willott and Urban 1978; Binns et al. 1992; Syka et al. 2000). Studies specifying iSRFs and iFTCs for DC and EN neurons remain to be done.

In the ICC, eSRFs have been studied extensively (Rose et al. 1963; Suga 1969; Gersuni et al. 1971; Ehret and Moffat 1985a; Casseday and Covey 1992; Wang et al. 1996; Ramachandran et al. 1999; Egorova et al. 2001; Le Beau et al. 2001). iSRFs measured in a two-tone paradigm have been investigated in comparatively few studies (Suga 1969; Ehret and Merzenich 1988a; Vater et al. 1992; Fuzessery 1994; Egorova et al. 2001; Lu and Jen 2001). eSRFs and iSRFs have varied shapes (Figs. 11.1, 11.3, and 11.6A) that sometimes reflect differences in stimulus presentation (monaural vs. binaural, short vs. long durations or rise times, simultaneous vs. asynchronous two-tone presentation) and data evaluation (different criteria for thresholds of excitatory and inhibitory thresholds). However, the rather uniform eFTC shapes of auditory nerve fibers (Kiang et al. 1965; Liberman 1978; Fig. 11.3A) are transformed into ICC FTCs that deviate substantially from the peripheral frequency filtering (Figs. 11.3 and




Figure 11.2. Excitatory frequency tuning curves from the external nucleus of the cat inferior colliculus. (Modified from Aitkin et al. 1975.)






Figure 11.3. (A) Examples of excitatory tuning curves of auditory nerve fibers with a low and a high characteristic frequency (*CF*). Areas of lateral suppression are shaded. (Modified from Ehret 1996.) (B) Classification (classes I to IV) of ICC neurons according to the shapes of their excitatory frequency tuning curves (*eFTCs*). Some variability of the tuning-curve slopes and shapes within each class is indicated. (Modified from Egorova et al. 2001.)

11.6A). With the simultaneous two-tone paradigm, grades of inhibition and heterogeneous iSRF shapes are seen in almost all ICC neurons. Owing to the glycinergic and  $\gamma$ -aminobutyric acid-positive (GABAergic) projections to the ICC (see Chapters 2, 4, and 9) and the intrinsic network with many GABAergic neurons (see Chapters 2, 5, and 9), inhibition is a prominent feature (Oliver et al. 1994; Winer et al. 1995; Le Beau et al. 1996, 2001).

The eSRFs and iSRFs of mouse ICC neurons have properties that define four classes of cells (Egorova et al. 2001) (Figs. 11.3B and 11.6A). All may have fSRFs.

Class I neurons closely resemble the excitatory and lateral suppression areas of auditory nerve fibers (Sachs and Kiang 1968). Their essential properties are: (1) strongly asymmetric eFTCs with steep high-frequency slopes and shallow low-frequency slopes; (2) inhibitory areas below and above CF that are asymmetric with higher inhibitory thresholds below the CF; (3) inhibitory areas that can overlap eSRFs only partially.

Class II neurons are dominated by inhibition. Their main features are: (1) inhibitory areas that are nearly symmetrical, with similar inhibitory thresholds below and above CF and that always invade eSRFs, often completely overlapping them; (2) the eFTCs have either steep slopes on both sides, are skewed toward the low- or high-frequency side at high levels, or are closed (with an upper response threshold); (3) inhibitory and excitatory thresholds and CFs are in close register.

Class III neurons have weak inhibition. Their other attributes are: (1) symmetric and shallow eFTC slopes; (2) inhibitory areas are small, inhibition is weak, and inhibitory areas are often only on one side of the eSRF; (3) excitatory and inhibitory thresholds are unrelated.

Class IV neurons have complex eSRFs and iSRFs, always with several eSRFs or multiple CFs.

iSRFs have not been determined in most ICC studies. Profiles of sound processing in the spectral domain are, however, incomplete without such measures and considerations of the graded influence of inhibition implicit in any RF classification. The gradient of eFTC slopes is an estimate of the extent and strength of inhibition, without direct measurement of iSRFs (Egorova et al. 2001). Most (93%) class I to III neurons can be classified by the slope criterion alone. Class I cells have low-frequency slopes  $<150$  dB/octave and high-frequency slopes  $>250$  dB/octave. Class II neuron low-frequency slopes are  $>150$  dB/octave and high-frequency slopes  $>250$  dB/octave. Class III neurons have low-frequency slopes  $<150$  dB/octave and high-frequency slopes of  $<250$  dB/octave. The critical slope values for classifying neurons may be species-specific. It is encouraging that the assignment of ICC neurons into four classes extends to other ICC studies for comparative purposes, and for functional considerations such as amount and extent of inhibition. Inhibition-dominated class II neurons have similar proportions in mice (27% to 28%; Egorova et al. 2001; Ehret et al. 2003), cats (27%; Ehret and Merzenich 1988a), and guinea pigs (26%; Le Beau et al. 2001). This suggests comparable inhibitory innervation patterns and functions. Class IV neurons are sparse in mice (6% to 10%; Egorova et al. 2001;

Ehret et al. 2003), guinea pigs (2%; Le Beau et al. 2001), and big brown bats (3%; Lu and Jen 2001), and high (40%) in the mustached bat (Portfors and Wenstrup 2002). Combinations of the harmonics of biosonar calls play a decisive role in echolocation, and selective adaptations may create combination-sensitive neurons and irregular FTCs (Mittmann and Wenstrup 1995; Leroy and Wenstrup 2000).

In the decerebrate cat, a related tuning curve classification (Ramachandran et al. 1999) is based on excitatory FTCs and inhibition of spontaneous activity (one-tone inhibition). Type “I” neurons combine a narrowly tuned I-shaped excitatory SRF with flanking inhibitory regions (Fig. 11.1B), like some class II neurons defined previously. Type “V” neurons show more broadly tuned, V-shaped excitatory areas with no inhibition of spontaneous activity (much like some class I/III neurons). Type “O” units have frequency response maps dominated by inhibition except for a circumscribed (O-shaped) island of excitation near CF and MT, also resembling some class II cells. The distribution of these classes seems to be method-specific: type O neurons comprise some 70% of ICC neurons in decerebrate cats (Ramachandran et al. 1999) but only 5% in nondecerebrated and anesthetized cats (Ehret and Merzenich 1988a) and 8% in nondecerebrated and anesthetized mice (Egorova et al. 2001). Differences between the two classification schemes will likely resolve when similar methodologies are applied across species.

Detailed descriptions of RFs in DC and EN are sparse, especially for inhibitory properties. Excitatory SRFs are broadly V-shaped or irregular (cat: Aitkin et al. 1975; guinea pig: Syka et al. 2000), suggesting many class III and IV neurons. Because class IV receptive fields are rare in the dorsal cochlear nucleus (Rhode and Smith 1986; Spirou and Young 1991) and superior olivary complex (Guinan et al. 1972) and ICC, they are likely the result of interactions between ascending (and descending) inputs.

### 3. FREQUENCY ORGANIZATION

#### 3.1. THE CENTRAL NUCLEUS

Virtually all ICC neurons are frequency tuned, although tuning sharpness differs from auditory nerve fibers. The one-dimensional cochlear frequency map is transformed into a three-dimensional map in the ICC (Fig. 11.4A). ICC tonotopy consists of two frequency gradients. The first is steep with stepwise changes from low frequencies (dorsal and dorsolateral ICC) to high frequencies (ventromedial ICC; Fig. 11.4B). This gradient, typical for most mammals, is orthogonal to the cellular laminae (see Chapter 2) and their input projections (see Chapters 3 to 5) and has been demonstrated with electrophysiologic measurements of single- or multiunit CFs (Merzenich and Reid 1974; Fitzpatrick 1975; Roth et al. 1978; Semple and Aitkin 1979; Stiebler and Ehret 1985; Poon et al. 1990; Romand and Ehret 1990; Casseday and Covey 1992; Brückner and RübSamen 1995; Schreiner and Langner 1997), with activity-dependent glucose labeling



by [ $^{14}\text{C}$ ]2-deoxyglucose autoradiography (Servièrè et al. 1984; Webster et al. 1984; Huang and Fex 1986; Martin et al. 1988; Ehret and Romand 1994; Brown et al. 1997), and with immediate-early gene activation using c-Fos immunocytochemistry (Ehret and Fischer 1991; Friauf 1992; Reimer 1993). This tonotopic organization forms a stack of “isofrequency sheets” of variable sizes and shapes covering the frequency spectrum of hearing (Fig. 11.4A). Isofrequency sheets parallel the ascending projections to the ICC (see Chapter 2). Cochlear CF space is often distorted in the ICC with frequency ranges important in communication or echolocation overrepresented (bat: Casseday and Covey 1992; mouse: Stiebler and Ehret 1985).

The second frequency gradient, described in cat ICC, is a shallow continuum within the isofrequency sheets (Schreiner and Langner 1997). Low CFs are dorsomedial and higher CFs lateral and ventrolateral (Fig. 11.4A: arrows). The frequency range covered within an isofrequency sheet is about one critical band (Schreiner and Langner 1997; see below). Thus, the sheets of the classic dorsoventral tonotopic gradient constitute “frequency-band laminae” (Merzenich and Reid 1974; Schreiner and Langner 1997). They are larger than synaptic domains, the microarchitectonic compartments of ICC laminae (see Chapter 2).

### 3.2. THE DORSAL CORTEX AND EXTERNAL NUCLEUS

The frequency representations in DC and EN have not been studied in the same detail as in the ICC. Historical differences in the definition of the DC have confounded a clear picture of its CF organization. Some studies show a frequency gradient from high (dorsal) to low (ventral at the border to the ICC) (Merzenich and Reid 1974; Servièrè et al. 1984), whereas others indicate that the ICC frequency-band laminae encroach dorsomedially into the DC, thus creating a continuous tonotopic organization for both subnuclei, at least in DC layers III and IV (Stiebler and Ehret 1985; Schreiner and Langner 1988; Romand and Ehret 1990). This physiologic ambiguity has a parallel anatomical component, with inputs from the lateral superior olive and other medullary structures (Oliver 1987) encroaching far beyond the architectonically recognized border of ICC (Morest and Oliver 1984; Oliver and Morest 1984) into layer IV of

Figure 11.4. (A) Three-dimensional plot of frequency-band laminae (10, 30, 50, 60 kHz) of the left-side ICC of the mouse. (Modified from Stiebler and Ehret 1985.) The *long thick lines* on the 10- and 30-kHz lamina indicate the direction of the shallow gradient of frequency increase (mainly from medial to lateral) within a lamina, as found in the cat (Schreiner and Langner 1997.) The concentric increase in the darkness of shading (10-kHz lamina) relates to the increase of the minimum threshold (*MT*) of neurons on a lamina (Stiebler 1986). (B) Electrode penetration from dorsal to ventral through the ICC of the cat showing a stepwise increase of the neuronal characteristic frequency (*CF*). The CF plateaus comprise frequency-band laminae. (Modified from Schreiner and Langner 1997 and unpublished data.)



the DC and perhaps beyond. In small mammals such as the mouse, a clear CF-based distinction between the DC and the ICC may not have been seen because of the thin DC (only layers I to III) (Meininger et al. 1986). Consequently, the ICC itself may not contain a representation of the lowest CF range because this area may be regarded as DC on the basis of anatomical criteria. Further study is necessary.

In the EN, a frequency gradient runs from high (lateral) to low (medial) in the cat (Rose et al. 1963; Roth et al. 1978) and in the mouse from low (lateral and dorsal) to high (medial and ventral) (Stiebler and Ehret 1985). Despite species differences there is agreement from studies in the possum (Aitkin et al. 1978), rat (Clopton and Winfield 1973), and gerbil (Kites 1984) that EN tonotopy is separate from that of ICC. Moreover, the CF ranges of DC and EN neurons are incomplete, with the high-frequency part absent or underrepresented. In the mouse EN, for example, CFs  $>35$  kHz have not been found while in the ICC CFs extend to  $>60$  kHz (Stiebler and Ehret 1985; Romand and Ehret 1990).

## 4. SPECTRAL RESOLUTION AND INTEGRATION

A basic principle of auditory processing is the decomposition of complex sounds into perceptually distinct frequency components or narrow frequency bands such as formants (Fletcher 1940; Scharf 1970; Patterson 1974). Each cochlear hair cell and most central auditory neurons respond only to a limited part of the spectrum, the RF bandwidth, and can distinguish or “resolve” these frequencies from those outside its range. Frequency components within an RF cannot be distinguished, and their acoustic energy is summed for further processing, a process defined as spectral integration. Much of the ascending auditory pathway can thus be interpreted as a bank of band-pass filters. At any given CF, the bandwidths of auditory nerve fibers are uniform, although intensity dependent (Rhode and Smith 1985). In view of the various excitatory, facilitatory, and inhibitory shapes of IC receptive fields, the uniformity of the peripheral band-pass filters and thus a common mechanism of spectral resolution, is either lost in the IC or transformed. The choice of experimental stimuli and of analytic methods influences the picture of spectral integration. We discuss three approaches: responses to pure tones, to tones in noise, and to various spectral envelopes of broadband signals.

### 4.1. SHARPNESS OF PURE-TONE TUNING CURVES

A common metric of the neuronal frequency resolution is the quality factor  $Q$ . Small  $Q$ -values reflect broad tuning or low frequency resolution, and large values narrow tuning and high resolution. The range of  $Q_{10\text{dB}}$ -values in IC neurons is enormous, from  $<0.05$  to  $>43$  (cat: Aitkin et al. 1994; rat: Kelly et al. 1991; guinea pig: Syka et al. 2000; mouse: Egorova et al. 2001). A portion of the

large variability (Fig. 11.5A) stems from the frequency dependence of  $Q$ -values established in the cochlear filtering.

Frequency resolution differs between subdivisions. The median  $Q_{10\text{dB}}$  in the ICC for cat is 4 (Aitkin et al. 1994), for guinea pig 2.5 (Syka et al. 2000) and for the big brown bat 4.4 (Jen et al. 2002), which is higher than in the EN of these species (cat: 2; guinea pig: 1.1; big brown bat: 2.2).

The categories of ICC tuning curve shapes (see Section 2) also reflect differences in frequency resolution (Egorova et al. 2001). Class II neurons have high (5 to 10)  $Q$ -values that are largely intensity independent (Fig. 11.5A).  $Q$ -values of class I neurons resemble those of class II neurons at 10 dB above threshold, but decrease at 80 dB. Class III  $Q$ -values decrease from about 6.5 to 0.5 (Fig. 11.5A) with increasing sound intensity. Similarly, sharply tuned cat ICC type I neurons have  $Q_{10\text{dB}}$ -values from 2 to 8 and  $Q_{40\text{dB}}$ -values from 1 to 7 with little




Figure 11.5. Sharpness of frequency tuning. (A) Average  $Q$ -values (see Section 2.1) as a function of the sound level for class I (filled circles), class II (open circles), and class III neurons (triangles) in the mouse ICC. (Modified from Egorova et al. 2001.) (B) View on the horizontal projection of a frequency-band lamina of the cat ICC. The lines are iso- $Q_{10\text{dB}}$  contours. The average  $Q_{10\text{dB}}$  decreases from about 8 near the center to around 2 near the peripheral margin of the lamina. (Modified from Schreiner and Langner 1988.) (C) Distribution of neurons of the tuning-curve classes I to IV as a function of their location in the central, pericentral, or peripheral area of frequency-band laminae of the mouse ICC. (Modified from Ehret et al. 2003.)

intensity dependence, whereas type V neurons show much broader and intensity-dependent tuning ( $Q_{10\text{dB}} < 3$ ;  $Q_{40\text{dB}} < 1.5$ ; Ramachandran et al. 1999). This range in  $Q$  behavior suggests that frequency resolution may serve different roles in different functional tasks.

The processing of sensory information through filters of different widths, a multiresolution analysis, is a general concept in sensory processing (Graham 1989). Such a multitask filtering operation may be expressed in the spatial organization of tuning sharpness and eFTC shape in the ICC. Multiunit recordings show a continuous spatial eFTC bandwidth gradient across frequency-band laminae (Stiebler 1987; Schreiner and Langner 1988; Fig. 11.5B), with higher frequency resolution in the center and declining resolution toward the laminar periphery, suggesting processing subdomains within laminae. This bandwidth gradient likely embodies the distribution of the different eFTC classes (Ehret et al. 2003; Hage and Ehret 2003). An abundance of sharply tuned class II neurons in the laminar center decreases toward the laminar periphery, while the number of broadly tuned class III neurons increases from center to periphery (Fig. 11.5C). Frequency resolution gradients are also present in the thalamocortical system (Read et al. 2001), suggesting that they are a general feature in central auditory organization.

#### 4.2. CRITICAL BANDS: SPECTRAL RESOLUTION

Psychophysical considerations have led to the concept of a bank of auditory filters or “critical bands” that subservise many perceptual aspects such as tone discrimination in noisy environments and the summation of loudness (Fletcher 1940; Zwicker et al. 1957; Scharf 1970; Moore 1997). Critical bands determined from narrow-band masking experiments cover equal distances of about 0.7 to 1 mm of the basilar membrane (BM) (Greenwood 1961, 1990; Ehret 1983, 1988). Thus, mice whose BM is short (about 7 mm; Ehret and Frankenreiter 1977) have fewer critical bands ( $< 10$ ; Ehret 1976) and poorer spectral resolution than cats (some 22 mm BM; Schuknecht 1960; some 24 critical bands; Pickles 1975, 1979; Nienhuys and Clark 1979), and humans (around 32 mm BM; von Békésy 1960; approximately 41 critical bands; Schafer et al. 1950; Margolis and Small 1975).

Auditory filter shape, bandwidth, and level dependence rests on the choice of test stimuli and psychophysical method. Accordingly, filter estimates in neurons must faithfully emulate psychophysical test conditions to be useful. For the ICC, two sets of data fulfill this condition (cats: Ehret and Merzenich 1985, 1988a; mice: Vartanian et al. 1999; Egorova et al. 2002). They show that ICC neural critical bandwidths follow the same rules as psychophysical critical bands (i.e., frequency dependence and level tolerance) when determined with a CF-centered tone and narrowband noise masker (Fig. 11.6A, B). Thus, critical-band related perceptual phenomena such as masking of tonal signals by noise (Fletcher 1940; Scharf 1970), the transition from consonance to dissonance evoked by shifting frequency components in a tone complex (Plomp and Levelt 1965), and changes





in the identification of vowels and animal vocalizations by shifts of frequency components (Flanagan 1972; Ehret and Haack 1982) are embedded in the spectral filter properties of ICC neurons. Perceptual critical-band properties are absent in auditory nerve fibers (Ehret 1995), and human auditory brain stem and middle-latency responses suggest that they first emerge in the IC (Zerlin 1986; Burrows and Barry 1990).

Several anatomical properties of the ICC are congruent with spectral integration within the frequency range of critical bands. For example, ICC frequency-band laminae cover a CF range similar to one critical bandwidth. Also, for any point within a frequency-band lamina, the CF distance to the neighboring laminae is about one critical bandwidth (Schreiner and Langner 1997; Fig. 11.4B). ICC neurons with disc-shaped or flat dendritic configurations constrain their integration to a limited part of the isofrequency domain (cf. Chapter 2) (Oliver and Morest 1984; Malmierca et al. 1993) and may process spectral components within a critical band. By contrast, multipolar or stellate cells with dendritic trees orthogonal to frequency-band laminae (Oliver and Morest 1984; Meininger et al. 1986; Wagner 1994; Reetz and Ehret 1999) may process information across frequency-band laminae (Oliver et al. 1991; Reetz and Ehret 1999), a feature perhaps essential for the analysis of spectral shape and other perceptual phenomena (Hall et al. 1984; Moore 1997; O'Connor et al. 1999). Critical band analyses of neurons in the DC and EN are not yet available.

#### 4.3. SPECTRAL ENVELOPE CODING: SPECTRAL INTEGRATION

The structural complexity of natural sounds and nonlinear mechanisms in auditory processing disbar simple stimuli alone, such as pure tones, from providing a complete characterization of spectral integration (Smolders et al. 1979; Nelken et al. 1999; Theunissen et al. 2000; Escabí and Schreiner 2002). Broadband stimuli derived from a few essential components in natural sounds can be effective for a more comprehensive characterization of spectral RF properties. Broadband sounds with sinusoidal spectral envelopes (“ripple spectra”) represent a class of complex stimuli intermediate between narrow-band artificial sounds and broadband natural vocalizations (Schreiner and Calhoun 1994; Shamma et al. 1995; Kowalski et al. 1996; Escabí and Schreiner 2002). They retain white noise criteria necessary for measuring reverse correlations (Eggermont et al. 1983): these include a flat power spectrum and an impulsive autocorrelation function and can probe the responsiveness of neurons in an unbiased manner. The resulting spectrotemporal receptive field (STRF) is a profile of the spectral and temporal envelope features to which a neuron responds. The SRF or spectral component of the STRF profile can quantify neuronal responses to complex spectra (e.g., formants speech) and can reveal the RF arrangement of excitation and inhibition (Kowalski et al. 1996; Calhoun and Schreiner 1998; Versnel and Shamma 1998; Miller et al. 2002; Qiu et al. 2003). This approach allows the study of spectral integration properties to dynamic broadband sounds with a rich spectral structure and thus under more natural conditions. The ability of IC



neurons to represent specific spectrotemporal feature combinations is a relatively common property in central auditory stations in species with specialized acoustic behavior (see Chapters 14, 16, and 17). With the STRF approach, stimulus feature selectivity can be generalized and applied to acoustically less specialized species.

The spectral integration properties found with ripple spectra parallel and expand those seen with tones. In cat ICC neurons tuned to 8 to 16 kHz, SRF bandwidths varied from 0.14 to 4.8 octaves with most (93%) bandwidths below about 2.0 octaves (Qui et al. 2003). Unlike pure-tone estimates, these bandwidth estimates were not strongly level dependent (Escabí et al. 2003), recalling critical-band measurements demonstrating spectral integration to be level independent (Ehret and Merzenich 1985, 1988a). SRF bandwidth distribution was unimodal (mean 1 octave). The best ripple density, the number of spectral envelope peaks/octave that produces a maximal neural response (Schreiner and Calhoun 1994; Klein et al. 2000; Escabí and Schreiner 2002) was low (mean = 0.61 cycles/octave) for most ICC neurons, indicating a preference for broadly spaced spectral features. The wide range of best ripple-density (0.02 to 2.11 cycles/octave), also reflects a wide range of spectral integration/selectivity. It is unclear how these measures relate to psychophysical or physiologic estimates of critical bands. Comparison of spectral envelope preferences in the ICC, the ventral division of the medial geniculate body, and primary auditory cortex reveals that general spectral integration properties seem well conserved in the lemniscal pathway between the IC and the auditory cortex (Miller et al. 2001, 2002; Qiu et al. 2003).

#### 4.4. ORIGINS OF SPECTRAL TUNING AND RESOLUTION

Cochlear filtering represented in the RFs of auditory nerve fibers is the basis for spectral tuning in higher centers of the auditory pathways. Careful removal of a narrow frequency channel at the level of the spiral ganglion can result in a narrow gap at corresponding frequencies in the eFTCs of ICC neurons (Snyder et al. 2000; Snyder and Sinex 2002). Auditory nerve fibers have relatively uniform eSRFs with areas of lateral suppression (Sachs and Kiang 1968; Schmiedt

Figure 11.6. (A) Excitatory spectral receptive fields (*eSRFs*, *cross-hatched areas*) bordered by the eFTC (*thin lines*), inhibitory SRFs (*iSRFs*, *shaded areas*), critical bandwidths (horizontal hatching bordered by thick lines), and facilitation areas (*arrows*) of class I to IV neurons of the cat ICC. The critical bandwidths are nearly independent of the FTC shapes. (Modified from Ehret and Merzenich 1988a and unpublished observations.) (B) Frequency dependence of critical bandwidth from cat ICC neurons (*open symbols*, measurements at various tone levels) and from behavioral measurements (*closed symbols*; Pickles 1975, 1979). Average behavioral and neural regression lines (*thick lines*) follow virtually the same functions if the neural data are adjusted to compensate for overmasking (*dashed line*). (Modified from Ehret and Merzenich 1985.)

1982; Delgutte 1990) but no facilitation areas (Fig. 11.3A). Brain stem neurons have far more diverse eSRF (eFTC) shapes, and reflect the presence of inhibition (Guinan et al. 1972; Young et al. 1988; Rhode 1991) and facilitation (Palmer et al. 1995; Winter and Palmer 1995). These diverse eSRF, iSRF, and fSRF shapes converge onto the IC and, in some cases, may be directly inherited by IC neurons (Ramachandran et al. 1999; Egorova et al. 2001).

The SRF of a given ICC neuron may be dominated by brain stem input as well as intrinsic biophysical properties (see Chapter 10). For example, many class I (I-type) ICC neurons (Ramachandran et al. 1999; Egorova et al. 2001) may inherit eSRF properties of type I cells in the ventral cochlear nucleus (Young et al. 1988) projecting to the ICC (Romand and Avan 1997). Similarly, many closed eSRFs of ICC neurons (type O; Ramachandran et al. 1999) are derived from O-shaped eSRFs of dorsal cochlear nucleus neurons, as blocking this projection reduces the incidence of O-neurons in the ICC by nearly 80% (Davis 2002). Further, medial superior olive neurons with broader tuning (Goldberg and Brown 1969; Guinan et al. 1972) project to the lateral ICC (Aitkin and Schuck 1985; Kudo and Nakamura 1988) and may engender the many broadly tuned class III neurons there. Pharmacologic blockade of GABAergic input to ICC neurons showed that inhibitory interactions can arise from subcollicular cells projecting to the ICC and from intrinsic ICC neurons. These inputs may affect SRFs in class I and II neurons (Yang et al. 1992; Pollak and Park 1993; Fuzessery and Hall 1996; Palombi and Caspary 1996; Lu and Jen 2001). Further, neurons with complex FTCs (class IV) likely arise from the convergence of neurons with class I and/or class II properties and somewhat different CFs (Egorova et al. 2001). Thus, SRFs of ICC neurons can be inherited from ascending projections and generated locally by excitatory–inhibitory interactions. The weight of both factors on a given neuron may depend on the location of the neuron in the three-dimensional fibrodendritic (see Chapter 2) and critical band (Schreiner and Langner 1997) space in the ICC. The origin of neural critical bandwidths with properties of psychophysical critical bands is associated with ICC neural filtering properties. Perhaps the genesis of critical-band properties reflects the shape of converging neuronal SRFs. This seems at first to be the case only for neurons with level-tolerant, sharp tuning (some class II or I-type neurons) because of the differences in shapes of FTCs and critical-band filters in class I, III, IV (and some class II) neurons (Fig. 11.6A). The steep slopes of the critical-band borders may derive from class I and class III eSRFs; however, they are shaped by inhibitory and facilitatory iSRF and fSRF influences (Ehret 1995). The suggestion that neural critical bands are sculpted from eSRFs mainly by inhibition (Ehret and Merzenich 1988a) has been supported in the mouse ICC (Egorova et al. 2002, 2003), where a strong correlation between critical-band cutoff frequencies and the CFs of inhibition (Figs. 11.1C and 11.6A, class I and II) occurs. In class II neurons, the border frequencies of the critical bands align almost perfectly with the peak frequencies of the iFTCs below and above the CF.

In summary, both spectral filtering (FTCs) and spectral resolution/integration are based on the same kind of processing in the ICC. The shapes of eFTCs can be considered a special case of “critical-band” filters for a one-tone stimulus. Whenever a multitone or complex spectrum is processed, inhibitory and facilitatory influences are activated from off-CF components that refine the neural response into a critical band. Accordingly, the properties of the composite SRF of a neuron (eSRF + iSRF + fSRF) determine its functional capabilities as spectral integrator.

## 5. INTENSITY CODING

The total dynamic range of the human ear comprises about 130 dB SPL in the frequency range of best hearing. Within this range, loudness can be matched across frequencies, and small intensity steps can be discriminated. Because loudness estimates are rather independent of the sound’s spectral composition (Zwicker and Feldtkeller 1967; Scharf 1978; Moore 1997), the neural coding of sound level and level differences seems to be largely independent of the type of sound.

The coding of sound level in the auditory nerve rests on three mechanisms: (1) there is an approximately 60-dB range of response threshold across AN fibers with similar CFs (Lieberman 1978); increasing sound level recruits more auditory nerve fibers with similar CFs. (2) AN fiber RLs vary in shape and dynamic range between 20 and 70 dB for CF tones (Sachs and Abbas 1974; Jackson and Relkin 1998). Dynamic ranges for tones above CF can exceed 70 dB. Thus, some AN fibers vary their response rate to tones over many levels. (3) With increasing amplitude, excitation spreads along the basilar membrane, especially toward the high-frequency end so that louder sounds recruit more AN fibers of increasing CFs (Zwicker and Feldtkeller 1967).

For the first two mechanisms, sound level at a given cochlear position (or CF) is coded over the 130-dB dynamic range by jointly considering the number of activated fibers and their activity levels. The code for AN sound intensity could equal the integrated activity of a small neural ensemble of similar CF. The third mechanism may enhance this population code at higher levels (Chatterjee and Zwislocki 1998) and may underlie the just-noticeable-intensity differences in psychoacoustical tests, which decrease with growing sound level (Maiwald 1967; Ehret 1989). Can the distribution of response thresholds and shapes of RLs illuminate how sound intensity is coded in the IC?

### 5.1. RESPONSE SENSITIVITY (THRESHOLDS)

The stimulus with the lowest SPL to evoke a neural response characterizes neural sensitivity (minimum threshold, MT) to sound pressure level. The behavior of response thresholds in psychophysics and for onset-responses, from auditory

nerve fibers to cortical neurons, is well described by the temporally integrated pressure envelope of a stimulus, in particular its onset envelope (Heil and Neubauer 2001, 2003). The site of this integration process is postulated to be at the inner hair cells and auditory nerve fiber's synapse (Heil and Neubauer 2003). Usually, neuronal responses are expressed as a function of the steady-state SPL although a metric that emphasizes dynamic stimulus aspects may be more appropriate (Heil 1997).

IC minimum thresholds for CF tones range over 60 to 80 dB in the cat (Aitkin 1991) and 35 to 45 dB in mice (Ehret and Moffat 1985a; Stiebler 1986). Because MTs may vary with the physiological state and anesthesia (Ehret and Moffat 1985b; Zurita et al. 1994), a large threshold range at a given CF may embody the animal's status rather than the actual MT range in a behaving animal. MTs are frequency dependent and typically follow the behavioral detection thresholds (Ehret and Moffat 1985a; Kelly et al. 1991; Egorova et al. 2001). The lowest contralateral values are  $-10$  and  $+10$  dB SPL for most species, and ipsilateral thresholds are about 10 to 20 dB higher (Aitkin 1991). Neurons of different spectral response types can also be characterized by MT differences: cat type V neurons have MTs 8 to 10 dB above that of type I and type O cells (Ramachandran et al. 1999). Mouse class III neurons have MTs 10 to 15 dB higher than cells of other classes (Egorova et al. 2001). The median MTs in the DC and the EN are 4 to 7 dB above ICC values (Palombi and Caspary 1996; Syka et al. 2000). Considering the spectral integration within critical bands, MTs for complex sounds such as noises are much lower than pure-tone MTs, if expressed as spectrum level (level/Hz) (Ehret and Moffat 1985a).

Tone MTs of cells in a frequency-band lamina vary systematically in mouse ICC, with lowest MTs in the center and progressively higher values peripherally (Stiebler and Ehret 1985; Stiebler 1986; Fig. 11.4A). Analogous topographic variations occur in the auditory cortex (Sutter and Schreiner 1995). Systematic spatial variations of intensity parameters may contribute a central (cortical) place-code component for loudness representation (Heil et al. 1994).

## 5.2. INTENSITY SELECTIVITY

The shape of RLs for superthreshold tones reveals several characteristics to increasing stimulus amplitude (Fig. 11.7). A segment of monotonic increase of firing rate with intensity shows a near-linear relationship between rate and SPL in all ICC neurons. However, few neurons maintain such growth at higher stimulus levels (Fig. 11.7A; Table 11.1). Most saturate at a certain intensity (plateau RLs, Fig. 11.7B), decline in rate (nonmonotonic RLs, Fig. 11.7C), or show variable behavior (complex RLs, Fig. 11.7D). Nonmonotonic RLs are intensity tuned, as the maximum rates are limited to a narrow range of stimulus levels. In most studies, these represent the preponderant tonal RL type in ICC (Table 11.1). Each type of RL to tones (Fig. 11.7) is found in varying proportions in the DC and EN (Table 11.1).




Figure 11.7. Examples of rate-level functions (RLs) in response to tone bursts from the cat ICC showing monotonic (**A**), plateau (**B**), nonmonotonic (**C**), and complex (**D**) shapes. *MT*, Minimum threshold. (Modified from Ehret and Merzenich 1988b and unpublished data.)

Table 11.1. Ranges of percentages for pure-tone rate-level functions of four basic types in IC subdivisions in cats (Ehret and Merzenich 1988b; Aitkin 1991; Aitkin et al. 1994), mice (Ehret and Moffat 1985a), rats (Palombi and Caspary 1996), and guinea pigs (Rees and Palmer 1988; Syka et al. 2000).

	Monotonic	Nonmonotonic	Plateau	Complex
ICC	6–28	15–66	5–35	7–26
EN	32–43	24–32	24–30	11–16
DC	6–39	21–59	4–58	0–31

The level eliciting maximal discharge is the “best” level. Best levels of nonmonotonic neurons occur at any superthreshold level in the cat (Ehret and Merzenich 1988b; Irvine and Gago 1990; Aitkin 1991) and have a range 10 to 40 dB above MT in rat and guinea pig (Rees and Palmer 1988; Palombi and Caspary 1996). Dynamic range is the extent of superthreshold sound levels over which an increased response rate occurs (Fig. 11.7). Dynamic ranges of monotonic neurons can exceed 80 dB (Ehret and Merzenich 1988b; Irvine and Gago 1990), while those of other RL types fall within about 50 dB (Aitkin 1991).

The large variability of the RL type percentages (Table 11.1) has several causes, including methodologic differences between studies and possible species differences. Studies available are constrained by a limited range of sound levels (<85 dB) tested; failure to consider the tone burst repetition rate (Galazyuk et al. 2000); or the fact that RLs were not calculated separately from the onset and the sustained response of neurons, which results in different RL shapes (Rees 1992). Moreover, differential criteria for assigning neurons to different RL types have been used. RL shapes often depend on stimulus type (Ehret and Merzenich 1988b; Aitkin 1991; Syka et al. 2000; Palombi and Caspary 1996), on the sound angle incidence (Semple and Kitzes 1985; Irvine and Gago 1990), on behavioral state (Ryan and Miller 1977), and on corticofugal influences (Syka et al. 1988; Zhang and Suga 1997; Jen et al. 1998; Suga et al. 2000; Zhou and Jen 2000; Yan and Ehret 2001, 2002; see Chapter 8). Hence, the shapes of RLs in response to tones and other stimuli are diverse and reflect many variables.

RL shape may be predicted in part from the SRF shape. Class II or type O neurons with a closed eSRF or class II neurons with an eSRF skewed toward higher or lower frequencies at high sound levels (Fig. 11.3) must have nonmonotonic (or complex) RLs to CF-tone bursts because, at higher SPLs, the CF may be outside the eSRF border. Thus, the RL shape embodies the same factors molding RFs (convergence of excitatory, inhibitory, and facilitatory input). Intracellular recordings and current injection of IC cells showed that monotonic, nonmonotonic, and plateau RLs may be locally produced by different ionic regimens interacting with membrane properties (Smith 1992; Wagner 1994; Reetz and Ehret 1999; Peruzzi et al. 2000; Sivaramakrishnan and Oliver 2001; Bal et al. 2002).



### 5.3. CONTRAST SENSITIVITY

The ability of the auditory system to encode amplitude differences has been studied almost exclusively in the context of intensity discrimination and loudness coding (Evans and Palmer 1980; Ehret and Merzenich 1988b; Viemeister 1988). The range of amplitude values present during a short segment of natural sounds, its contrast, can be substantial (Attias and Schreiner 1998), and varies considerably over time (e.g., during speech) and between sound environments (Escabí et al. 2003). The decibel amplitude distribution of spectrotemporal contrast in natural sounds is approximately normal (Fig. 11.8A). Perhaps the operating range of IC neurons, reflected in the RLs, is matched to the natural amplitude statistics, and thus can efficiently process sounds and thereby detect and segregate signals. ICC neurons are tuned to naturally occurring contrast distributions (Escabí et al. 2003). Moreover, mean intensity and contrast coding appear to be independent. Like RLs for stimulus level, rate-contrast functions are monotonic or nonmonotonic but are independent of the type of RL function in the cell (Fig. 11.8B). This independent response behavior for two stimulus amplitude measures (mean and variance) suggests that both are relevant features of the acoustic landscape and its neural representation. Understanding the role of contrast in




Figure 11.8. (A) Probability distributions of amplitude values in three different sound classes. The width of the distributions corresponds to the contrast of the sounds. Note the differences in width between white noise and natural signals. (B) Firing rate profiles for three different neurons in the ICC for “ripple-spectra” (broadband stimuli with noise-like spectral envelopes) that were systematically and independently varied in mean intensity and contrast. (Adapted from Escabí et al. 2003.)

auditory processing from the perspective of natural contrast distributions must receive further attention.

#### 5.4. MODELS OF SOUND INTENSITY CODING

The available evidence for sound intensity coding at the IC does not allow conclusive statements about how sound intensity, level or loudness, and intensity differences or contrasts are coded. At present, three fundamental ideas about intensity coding are plausible (Irvine 1992; Ehret 1997): (1) Small groups of neurons with monotonic, sound-type independent RLs over 80 dB or more code sound intensity and level variation by their integrated spike response. (2) Neurons with nonmonotonic RLs aggregate by their best levels at ICC loci in frequency-band laminae and map sound intensity by their location. A similar spatial code for sound level occurs in a specialized field of the mustached bat's auditory cortex (Suga 1977; Suga and Manabe 1982). (3) From maps of absolute response thresholds in frequency-band laminae (Stiebler 1986), a model of the transfer of cochlear sound parameters to the three-dimensional ICC space predicts a spatial code for sound intensity, that is, a shift of the maximum of the neural response from the center of a frequency-band lamina to its periphery with increasing amplitude (Herrnberger et al. 2002). Implications of this model include systematic neural arrangements with monotonic and nonmonotonic RLs in frequency-band laminae. These three propositions about midbrain intensity coding incorporate RL variability and predict that an average-rate code for sound intensity derived from a large proportion of ICC neurons is unlikely (Ehret and Merzenich 1988b). Onset and sustained responses could have unique contributions to intensity and contrast coding. The biological significance or semantic content of environmental and communication sound is often coded independently of intensity. This suggests as yet unidentified mechanisms in the auditory pathways that separate the influence of the total sound level from that of biologically significant level differences or contrasts. However, as total level can mediate arousal, emotions, and motivations, information about it must be coded, and may be transmitted, independently of semantic information, to higher brain centers.

## 6. CONCLUSIONS AND QUESTIONS FOR FUTURE RESEARCH

The coding of sound frequency and intensity are fundamental tasks of auditory processing and have been extensively studied in the IC. Despite the impressive body of studies available for RF properties, many aspects of the midbrain contribution to various sound processing tasks are unknown. The convergence of input from many sources and the several types of spectral/intensity RFs creates an interpretational dilemma. The segregation of subcollicular pathways in the

cochlear nucleus complex, superior olivary complex, and nuclei of the lateral lemniscus can be construed in terms of specialized tasks that require certain dedicated structural and functional substrates of neural organization. After the massive convergence at the midbrain level, this paradigm is simply no longer appropriate without a deeper grasp of the nature, purpose, or task-specificity of the convergence. Parallel output from an abundance of subcollicular nuclei appears compressed in the ICC onto a single tonotopic substrate, giving a new meaning to the term convergence, with no counterpart in either the visual or somatosensory systems (Ehret 1997). The challenge still is to decipher how the neurons of this substrate are specialized in their structure, physiology, and network properties to serve the many functional demands of sound processing.

We note here just a few immediate tasks directly related to matters of spectral and intensity processing:

1. What are the relationships between the morphologic and physiologic types of neurons and functional maps within ICC frequency-band laminae and those in the DC and the EN?
2. How are the composition and pattern of input projections from lower auditory centers related to the type of IC neuron (morphologic, physiologic, biophysical, functional)? Are there unique circuitries between neighboring neurons and potential laminar and subnuclear synaptic domains?
3. IC neurons are modulated by neurotransmitter-specific inputs such as the cholinergic and noradrenergic systems (Faingold et al. 1991; see Chapter 9) and by descending projections from the auditory cortex and thalamus (see Chapter 8). How do these influence sound processing under natural conditions in awake and behaving animals, and what role do they play in auditory scene analysis, hearing impairment, and neural plasticity?

## Abbreviations

BF	best frequency
CF	characteristic frequency
DC	dorsal cortex of the inferior colliculus
eFTC	excitatory frequency tuning curve
EN	external nucleus of the inferior colliculus
eSRF	excitatory spectral receptive field
fSRF	facilitatory spectral receptive field
FTC	frequency tuning curve
GABA	$\gamma$ -aminobutyric acid
IC	inferior colliculus
ICC	central nucleus of the inferior colliculus
iFTC	inhibitory frequency tuning curve
iSRF	inhibitory spectral receptive field
MT	minimum threshold
RF	receptive field

RL	rate-level function
SPL	sound pressure level
SRF	spectral receptive field
STRF	spectrotemporal receptive field

## References

- Aitkin L (1986) *The Auditory Midbrain. Structure and Function in the Central Auditory Pathway*. Humana Press, Clifton.
- Aitkin L (1991) Rate-level functions of neurons in the inferior colliculus of cats measured with the use of free-field sound stimuli. *Journal of Neurophysiology* **65**:383–392.
- Aitkin L and Schuck D (1985) Low frequency neurons in the lateral central nucleus of the cat inferior colliculus receive their input predominantly from the medial superior olive. *Hearing Research* **17**:87–93.
- Aitkin L, Tran L, and Syka J (1994) The response of neurons in subdivisions of the inferior colliculus of cats to tonal, noise and vocal stimuli. *Experimental Brain Research* **98**:53–64.
- Aitkin LM, Webster WR, Veale JL, and Crosby DC (1975) Inferior colliculus. I. Comparison of response properties of neurons in central, pericentral, and external nuclei of adult cat. *Journal of Neurophysiology* **38**:1196–1207.
- Aitkin LM, Dickhaus H, Schult W, and Zimmermann M (1978) External nucleus of inferior colliculus: auditory and spinal somatosensory afferents and their interactions. *Journal of Neurophysiology* **41**:837–847.
- Attias H and Schreiner CE (1998) Coding of naturalistic stimuli by auditory midbrain neurons. In: Jordan MI (ed). *Advances in Neural Information Processing Systems 10*. MIT Press, Cambridge, pp. 103–109.
- Bal R, Green GGR, Rees A, and Sanders DJ (2002) Firing patterns of inferior colliculus neurons—histology and mechanism to change firing patterns in rat brain slices. *Neuroscience Letters* **317**:42–46.
- Binns KE, Grant S, Withington DJ, and Keating MJ (1992) A topographic representation of auditory space in the external nucleus of the inferior colliculus of the guinea-pig. *Brain Research* **589**:231–242.
- Brown M, Webster WR, and Martin RL (1997) The three-dimensional frequency organization of the inferior colliculus of the cat: a 2-deoxyglucose study. *Hearing Research* **104**:57–72.
- Brückner S and Rübsamen R (1995) Binaural response characteristics in iso-frequency sheets of the gerbil inferior colliculus. *Hearing Research* **86**:1–14.
- Burrows DL and Barry SJ (1990) Electrophysiological evidence for the critical band in humans: middle-latency responses. *Journal of the Acoustical Society of America* **88**:180–184.
- Calhoun BM and Schreiner CE (1998) Spectral envelope coding in cat primary auditory cortex: linear and non-linear effects of stimulus characteristics. *European Journal of Neuroscience* **10**:926–940.
- Casseday JH and Covey E (1992) Frequency tuning properties of neurons in the inferior colliculus of an FM bat. *Journal of Comparative Neurology* **319**:34–50.
- Chatterjee M and Zwislocki JJ (1998) Cochlear mechanisms of frequency and intensity coding. II. Dynamic range and the code for loudness. *Hearing Research* **124**:170–181.

- Clopton BM and Winfield JA (1973) Tonotopic organization in the inferior colliculus of the rat. *Brain Research* **56**:355–358.
- Davis KA (2002) Evidence of a functionally segregated pathway from dorsal cochlear nucleus to inferior colliculus. *Journal of Neurophysiology* **87**:1824–1835.
- Delgutte B (1990) Two-tone rate suppression in auditory-nerve fibers: dependence on suppressor frequency and level. *Hearing Research* **49**:225–246.
- Eggermont JJ, Aertsen AMHJ, and Johannesma PIM (1983) Prediction of the responses of auditory neurons in the midbrain of the grass frog based on the spectro-temporal receptive field. *Hearing Research* **10**:191–202.
- Egorova MA, Ehret G, Vartanian IA, and Esser K-H (2001) Frequency response areas of neurons in the mouse inferior colliculus I. Threshold and tuning characteristics. *Experimental Brain Research* **140**:145–161.
- Egorova MA, Vartanyan IA, and Ehret G (2002) Neural critical bands and inhibition in the auditory midbrain of house mouse (*Mus domesticus*). *Dokladi Biological Sciences* **382**:5–7 (in Russian).
- Egorova MA, Ehret G, and Vartanyan I (2003) Critical bandwidths and inhibition in auditory midbrain neurons of house mice. In: Elsner N and Zimmermann H (eds). *The Neurosciences from Basic Research to Therapy*. Thieme, Stuttgart, pp. 1080–1081.
- Ehret G (1976) Critical bands and filter characteristics in the ear of the house mouse (*Mus musculus*). *Biological Cybernetics* **24**:35–42.
- Ehret G (1983) Psychophysics. In: Willott JF (ed). *The Auditory Psychobiology of the Mouse*. Charles C Thomas, Springfield, pp. 13–56.
- Ehret G (1988) Frequency resolution, spectral filtering and integration on the neuronal level. In: Edelman GM, Gall WE, and Cowan WM (eds). *Auditory Function. Neurobiological Bases of Hearing*. John Wiley & Sons, New York, pp. 363–384.
- Ehret G (1989) Hearing in the mouse. In: Dooling RJ and Hulse SH (eds). *The Comparative Psychology of Audition: Perceiving Complex Sounds*. Lawrence Erlbaum, Hillsdale, pp. 3–32.
- Ehret G (1995) Auditory frequency resolution in mammals: from neuronal representation to perception. In: Manley GA, Klump GM, Köppl C, Fastl H, and Oeckinghaus H (eds). *Advances in Hearing Research*. World Scientific, Singapore, pp. 387–397.
- Ehret G (1996) Auditorische Systeme. In: Dudel J, Menzel R, and Schmidt RF (eds). *Neurowissenschaft*. Springer-Verlag, Berlin, pp. 353–381.
- Ehret G (1997) The auditory midbrain, a ‘shunting yard’ of acoustical information processing. In: Ehret G and Romand R (eds). *The Central Auditory System*. Oxford University Press, New York, pp. 259–316.
- Ehret G and Fischer R (1991) Neuronal activity and tonotopy in the auditory system visualized by c-fos gene expression. *Brain Research* **567**:350–354.
- Ehret G and Frankenreiter M (1977) Quantitative analysis of cochlear structures in the house mouse in relation to mechanisms of acoustical information processing. *Journal of Comparative Physiology A* **122**:65–85.
- Ehret G and Haack B (1982) Ultrasound recognition in house mice: key-stimulus configuration and recognition mechanisms. *Journal of Comparative Physiology A* **148**:245–251.
- Ehret G and Merzenich MM (1985) Auditory midbrain responses parallel spectral integration phenomena. *Science* **227**:1245–1247.
- Ehret G and Merzenich MM (1988a) Complex sound analysis (frequency resolution, filtering and spectral integration) by single units of the inferior colliculus of the cat. *Brain Research Reviews* **13**:139–163.

- Ehret G and Merzenich MM (1988b) Neuronal discharge rate is unsuitable for encoding sound intensity at the inferior colliculus level. *Hearing Research* **35**:1–8.
- Ehret G and Moffat AJM (1985a) Inferior colliculus of the house mouse. II. Single unit responses to tones, noise and tone-noise combinations as a function of sound intensity. *Journal of Comparative Physiology A* **156**:619–635.
- Ehret G and Moffat AJM (1985b) Inferior colliculus of the house mouse. III. Response probabilities and thresholds of single units to synthesized mouse calls compared to tone and noise bursts. *Journal of Comparative Physiology A* **156**:637–644.
- Ehret G and Romand R (1994) Development of tonotopy in the inferior colliculus. II. 2-DG measurements in kitten. *European Journal of Neuroscience* **6**:1589–1595.
- Ehret G, Egorova MA, Hage SR, and Müller BA (2003) Spatial map of frequency tuning-curve shapes in the mouse inferior colliculus. *NeuroReport* **14**:1365–1369.
- Escabí MA and Schreiner CE (2002) Nonlinear spectrotemporal sound analysis by neurons in the auditory midbrain. *Journal of Neuroscience* **22**:4114–4131.
- Escabí MA, Miller LM, Read HL, and Schreiner CE (2003) Naturalistic auditory contrast improves spectrotemporal coding in the cat inferior colliculus. *Journal of Neuroscience* **23**:11489–11504.
- Evans EF and Palmer AR (1980) Relationship between the dynamic range of cochlear nerve fibres and their spontaneous activity. *Experimental Brain Research* **40**:115–118.
- Faingold CL, Gehlbach G, and Caspary DM (1991) Functional pharmacology of inferior colliculus neurons. In: Altschuler RA, Bobbin RP, Clopton BM, and Hoffman DW (eds). *Neurobiology of Hearing: The Central Auditory System*. Raven Press, New York, pp. 223–251.
- Fitzpatrick KA (1975) Cellular architecture and topographic organization of the inferior colliculus of the squirrel monkey. *Journal of Comparative Neurology* **164**:185–208.
- Flanagan JL (1972) *Speech Analysis, Synthesis and Perception*. Springer-Verlag, Berlin.
- Fletcher H (1940) Auditory patterns. *Reviews of Modern Physics* **12**:47–65.
- Friauf E (1992) Tonotopic order in the adult and developing auditory system of the rat as shown by c-fos immunocytochemistry. *European Journal of Neuroscience* **4**:798–812.
- Fuzessery ZM (1994) Response selectivity for multiple dimensions of frequency sweeps in the pallid bat inferior colliculus. *Journal of Neurophysiology* **72**:1061–1079.
- Fuzessery ZM and Feng AS (1983) Mating call selectivity in the thalamus and midbrain of the leopard frog (*Rana pipiens*): single and multiunit analyses. *Journal of Comparative Physiology A* **150**:333–344.
- Fuzessery ZM and Hall JC (1996) Role of GABA in shaping frequency tuning and creating FM sweep selectivity in the inferior colliculus. *Journal of Neurophysiology* **76**:1059–1073.
- Galazyuk AV, Llano D, and Feng AS (2000) Temporal dynamics of acoustic stimuli enhance amplitude tuning of inferior colliculus neurons. *Journal of Neurophysiology* **83**:128–138.
- Gersuni GV, Altman JA, Maruseva EA, Radionova EA, Ratnikova GI, and Vartanian IA (1971) Functional classification of neurons in the inferior colliculus of the cat according to their temporal characteristics. In: Gersuni GF (ed). *Sensory Processes at the Neuronal and Behavioral Level*. Academic Press, New York, pp. 157–179.
- Goldberg JM and Brown PB (1969) Response of binaural neurons of dog superior olivary complex to dichotic tonal stimuli: some physiological mechanisms of sound localization. *Journal of Neurophysiology* **32**:613–636.
- Graham N (1989) *Visual Pattern Analyzers*. Oxford University Press, Oxford.

- Greenwood DD (1961) Auditory masking and the critical band. *Journal of the Acoustical Society of America* **33**:484–502.
- Greenwood DD (1990) A cochlear frequency-position function for several species—29 years later. *Journal of the Acoustical Society of America* **87**:2592–2605.
- Guinan JJ Jr, Guinan SS, and Norris BE (1972) Single auditory units in the superior olivary complex. I: responses to sounds and classifications based on physiological properties. *International Journal of Neuroscience* **4**:101–120.
- Hage SR and Ehret G (2003) Mapping responses to frequency sweeps and tones in the inferior colliculus of house mice. *European Journal of Neuroscience* **18**:2301–2312.
- Hall JW, Haggard MP, and Fernandes MA (1984) Detection in noise by spectro-temporal pattern analysis. *Journal of the Acoustical Society of America* **76**:50–56.
- Heil P (1997) Aspects of temporal processing of FM stimuli in primary auditory cortex. *Acta Otolaryngologica (Stockholm) Supplement* **532**:99–102.
- Heil P and Neubauer H (2001) Temporal integration of sound pressure determines thresholds of auditory-nerve fibers. *Journal of Neuroscience* **21**:7404–7415.
- Heil P and Neubauer H (2003) A unifying basis of auditory thresholds based on temporal summation. *Proceedings of the National Academy of Sciences of the United States of America* **100**:6151–6156.
- Heil P, Rajan R, and Irvine DRF (1994) Topographic representation of tone intensity along the isofrequency axis of cat primary auditory cortex. *Hearing Research* **76**:188–202.
- Herrnberger B, Kempf S, and Ehret G (2002) Basic maps of the auditory midbrain. *Biological Cybernetics* **87**:231–240.
- Huang C and Fex J (1986) Tonotopic organization in the inferior colliculus of the rat demonstrated with the 2-deoxyglucose method. *Experimental Brain Research* **61**:506–512.
- Irvine DRF (1992) Physiology of the auditory brainstem. In: Popper AN and Fay RR (eds). *Springer Handbook of Auditory Research, Volume 2: The Mammalian Auditory Pathway: Neurophysiology*. Springer-Verlag, New York, pp. 153–231.
- Irvine DRF and Gago G (1990) Binaural interaction in high-frequency neurons in inferior colliculus of the cat: effects of variations in sound pressure level on sensitivity to interaural intensity differences. *Journal of Neurophysiology* **63**:570–591.
- Jackson BS and Relkin EM (1998) A frequency-dependent saturation evident in rate-intensity functions of the chinchilla auditory nerve. *Hearing Research* **126**:75–83.
- Jen PH-S, Chen QC, and Sun XD (1998) Corticofugal regulation of auditory sensitivity in the bat inferior colliculus. *Journal of Comparative Physiology A* **183**:683–697.
- Jen PH-S, Wu FJ, and Chen QC (2002) The effect of two-tone stimulation on responses of two simultaneous recorded neurons in the inferior colliculus of the big brown bat, *Eptesicus fuscus*. *Hearing Research* **168**:139–149.
- Kelly JB, Glenn L, and Beaver CJ (1991) Sound frequency and binaural response properties of single neurons in rat inferior colliculus. *Hearing Research* **56**:273–280.
- Kiang NY-S, Watanabe T, Thomas EC, and Clark LF (1965) *Discharge Patterns of Single Fibers in the Cat's Auditory Nerve*. MIT Press, Cambridge.
- Kitzes LM (1984) Some physiological consequences of neonatal cochlear destruction in the inferior colliculus of the gerbil, *Meriones unguiculatus*. *Brain Research* **306**:171–178.
- Klein DJ, Depireux DA, Simon JZ, and Shamma SA (2000) Robust spectrotemporal reverse correlation for the auditory system: optimizing stimulus design. *Journal of Computational Neuroscience* **9**:85–111.

- Kowalski N, Depireux DA, and Shamma SA (1996) Analysis of dynamic spectra in ferret primary auditory cortex. I. Characteristics of single-unit responses to moving ripple spectra. *Journal of Neurophysiology* **76**:3503–3523.
- Kudo M and Nakamura Y (1988) Organization of the lateral lemniscal fibers converging onto the inferior colliculus in the cat: an anatomical review. In: Syka J and Masterton RB (eds). *Auditory Pathway: Structure and Function*. Plenum Press, New York, pp. 171–183.
- Le Beau FEN, Rees A, and Malmierca MS (1996) Contribution of GABA- and glycine-mediated inhibition to the monaural temporal response properties of neurons in the inferior colliculus. *Journal of Neurophysiology* **75**:902–919.
- Le Beau FEN, Malmierca MS, and Rees A (2001) Iontophoresis in vivo demonstrates a key role for GABA<sub>A</sub> and glycinergic inhibition in shaping frequency response areas in the inferior colliculus of the guinea pig. *Journal of Neuroscience* **21**:7303–7312.
- Leroy SA and Wenstrup JJ (2000) Spectral integration in the inferior colliculus of the mustached bat. *Journal of Neuroscience* **20**:8533–8541.
- Liberman MC (1978) Auditory-nerve response from cats raised in a low-noise chamber. *Journal of the Acoustical Society of America* **63**:442–455.
- Lu Y and Jen PH-S (2001) GABAergic and glycinergic neural inhibition in excitatory frequency tuning of bat inferior collicular neurons. *Experimental Brain Research* **141**: 331–339.
- Maiwald D (1967) Ein Funktionsschema des Gehörs zur Beschreibung der Erkennbarkeit kleiner Frequenz- und Amplitudenänderungen. *Acustica* **18**:81–92.
- Malmierca MS, Blackstad TW, Osen KK, Karagülle T, and Molowny RL (1993) The central nucleus of the inferior colliculus in rat: a Golgi and computer reconstruction study of neuronal and laminar structure. *Journal of Comparative Neurology* **333**:1–27.
- Margolis RH and Small AM (1975) The measurement of critical masking bands. *Journal of Speech and Hearing Research* **18**:571–587.
- Martin RL, Webster WR, and Servière J (1988) The frequency organization of the inferior colliculus of the guinea pig: a (14C)-2-deoxyglucose study. *Hearing Research* **33**:245–256.
- Meininger V, Pol D, and Derer P (1986) The inferior colliculus of the mouse. A Nissl and Golgi study. *Neuroscience* **17**:1159–1179.
- Merzenich MM and Reid MD (1974) Representation of the cochlea within the inferior colliculus of the cat. *Brain Research* **77**:397–415.
- Miller LM, Escabí MA, Read HL, and Schreiner CE (2001) Functional convergence of response properties in the auditory thalamocortical system. *Neuron* **32**:151–160.
- Miller LM, Escabí MA, Read HL, and Schreiner CE (2002) Spectrotemporal receptive fields in the lemniscal auditory thalamus and cortex. *Journal of Neurophysiology* **87**: 516–527.
- Mittman DH and Wenstrup JJ (1995) Combination-sensitive neurons in the inferior colliculus. *Hearing Research* **90**:185–191.
- Moore BCJ (1997) *An Introduction to the Psychology of Hearing*. Academic Press, San Diego.
- Morest DK and Oliver DL (1984) The neuronal architecture of the inferior colliculus in the cat: defining the functional anatomy of the auditory midbrain. *Journal of Comparative Neurology* **222**:209–236.
- Nelken I, Rotman Y, and Yosef OB (1999). Responses of auditory-cortex neurons to structural features of natural sounds. *Nature* **397**:154–156.



- Nienhuys TGW and Clark GM (1979) Critical bands following the selective destruction of cochlear inner and outer hair cells. *Acta Otolaryngologica (Stockholm)* **88**:350–358.
- Oliver DL (1987) Projections to the inferior colliculus from the anteroventral cochlear nucleus in the cat: possible substrates for binaural interaction. *Journal of Comparative Neurology* **264**:24–46.
- Oliver DL and Morest DK (1984) The central nucleus of the inferior colliculus in the cat. *Journal of Comparative Neurology* **222**:237–264.
- Oliver DL, Kuwada S, Yin TCT, Haberly LB, and Henkel CK (1991) Dendritic and axonal morphology of HRP-injected neurons in the inferior colliculus of the cat. *Journal of Comparative Neurology* **303**:75–100.
- Oliver DL, Winer JA, Beckius GE, and Saint Marie RL (1994) Morphology of GABAergic neurons in the inferior colliculus of the cat. *Journal of Comparative Neurology* **340**:27–42.
- O'Connor KN, Barruel P, Hajalilou R, and Sutter ML (1999) Auditory temporal integration in the rhesus macaque. *Journal of the Acoustical Society of America* **106**:954–965.
- Palmer AR, Winter IM, Jiang D, and James N (1995) Across-frequency integration by neurones in the ventral cochlear nucleus. In: Manley GA, Klump GM, Köppl C, Fastl H, and Oeckinghaus H (eds). *Advances in Hearing Research*. World Scientific, Singapore, pp. 250–263.
- Palombi PS and Caspary DM (1996) GABA inputs control discharge rate primarily within frequency receptive fields of inferior colliculus neurons. *Journal of Neurophysiology* **75**:2211–2219.
- Patterson RD (1974) Auditory filter shape. *Journal of the Acoustical Society of America* **55**:802–809.
- Patuzzi R (1996) Cochlear micromechanics and macromechanics. In: Dallos P, Popper AN, and Fay RR (eds). *Springer Handbook of Auditory Research, Volume 8: The Cochlea*. Springer-Verlag, New York, pp. 186–257.
- Peruzzi D, Sivaramakrishnan S, and Oliver DL (2000) Identification of cell types in brain slices of the inferior colliculus. *Neuroscience* **101**:403–416.
- Pickles JO (1975) Normal critical bands in the cat. *Acta Otolaryngologica* **80**:245–254.
- Pickles JO (1979) Psychophysical frequency resolution in the cat as determined by simultaneous masking and its relation to auditory-nerve resolution. *Journal of the Acoustical Society of America* **66**:1725–1732.
- Plomp R and Levelt WJM (1965) Tonal consonance and critical bandwidth. *Journal of the Acoustical Society of America* **38**:548–560.
- Pollak GD and Park TJ (1993) The effects of GABAergic inhibition on monaural response properties of neurons in the mustache bat's inferior colliculus. *Hearing Research* **65**:99–117.
- Poon PWF, Sun X, Kamada T, and Jen PH-S (1990) Frequency and space representation in the inferior colliculus of the FM bat, *Eptesicus fuscus*. *Experimental Brain Research* **79**:83–91.
- Portfors CV and Wenstrup JJ (2002) Excitatory and facilitatory frequency response areas in the inferior colliculus of the mustached bat. *Hearing Research* **168**:131–138.
- Qiu A, Schreiner CE, and Escabi MA (2003) Gabor analysis of auditory midbrain receptive fields: spectro-temporal and binaural composition. *Journal of Neurophysiology* **90**:456–476.

- Ramachandran R, Davis KA, and May BJ (1999) Single-unit responses in the inferior colliculus of decerebrate cats I. Classification based on frequency response maps. *Journal of Neurophysiology* **82**:152–163.
- Read HL, Winer JA, and Schreiner CE (2001) Modular organization of intrinsic connections associated with spectral tuning in cat auditory cortex. *Proceedings of the National Academy of Sciences of the United States of America* **98**:8042–8047.
- Rees A (1992) The coding of intensity in the inferior colliculus: onset and sustained responses. In: Cazals Y, Demany L, and Horner K (eds). *Auditory Physiology and Perception*. Pergamon Press, Oxford, pp. 263–269.
- Rees A and Palmer AR (1988) Rate-intensity functions and their modification by broadband noise for neurons in the guinea pig inferior colliculus. *Journal of the Acoustical Society of America* **83**:1488–1498.
- Reetz G and Ehret G (1999) Inputs from three brainstem sources to identified neurons of the mouse inferior colliculus slice. *Brain Research* **816**:527–543.
- Reimer K (1993) Simultaneous demonstration of Fos-like immunoreactivity and 2-deoxyglucose uptake in the inferior colliculus of the mouse. *Brain Research* **616**:339–343.
- Rhode WS (1991) Physiological-morphological properties of the cochlear nucleus. In: Altschuler RA, Bobbin RP, Clopton BM, and Hoffman DW (eds). *Neurobiology of Hearing: The Central Auditory System*. Raven Press, New York, pp. 47–77.
- Rhode WS and Smith PH (1985) Characteristics of tone-pip response patterns in relationship to spontaneous rate in cat auditory nerve fibers. *Hearing Research* **18**:159–168.
- Rhode WS and Smith PH (1986) Physiological studies on neurons in the dorsal cochlear nucleus of cat. *Journal of Neurophysiology* **56**:287–307.
- Romand R and Avan P (1997) Anatomical and functional aspects of the cochlear nucleus. In: Ehret G and Romand R (eds). *The Central Auditory System*. Oxford University Press, New York, pp. 97–191.
- Romand R and Ehret G (1990) Development of tonotopy in the inferior colliculus. I. Electrophysiological mapping in house mice. *Developmental Brain Research* **54**:221–234.
- Rose JE, Greenwood DD, Goldberg JM, and Hind JE (1963) Some discharge characteristics of single neurons in the inferior colliculus of the cat. I. Tonotopical organization, relation of spike-counts to tone intensity, and firing patterns of single elements. *Journal of Neurophysiology* **26**:294–320.
- Roth GL, Aitkin LM, Andersen RA, and Merzenich MM (1978) Some features of the spatial organization of the central nucleus of the inferior colliculus of the cat. *Journal of Comparative Neurology* **182**:661–680.
- Ryan A and Miller J (1977) Effects of behavioral performance on single-unit firing patterns in inferior colliculus of the rhesus monkey. *Journal of Neurophysiology* **40**:943–956.
- Sachs MB and Abbas PJ (1974) Rate versus level functions for auditory-nerve fibers in cats: tone-burst stimuli. *Journal of the Acoustical Society of America* **56**:1835–1847.
- Sachs MB and Kiang NY-S (1968) Two-tone inhibition in auditory-nerve fibers. *Journal of the Acoustical Society of America* **43**:1120–1128.
- Schafer TH, Gales RS, Shewmaker CA, and Thompson PO (1950) The frequency selectivity of the ear as determined by masking experiments. *Journal of the Acoustical Society of America* **22**:490–496.

- Scharf B (1970) Critical bands. In: Tobias JV (ed). *Foundations of Modern Auditory Theory*, Volume I. Academic Press, New York, pp. 159–202.
- Scharf B (1978) Loudness. In: Carterette EC and Friedman MP (eds). *Handbook of Perception*, Volume IV: *Hearing*. Academic Press, New York, pp. 187–242.
- Schmiedt RA (1982) Boundaries of two-tone rate suppression of cochlear-nerve activity. *Hearing Research* **7**:335–351.
- Schreiner CE and Calhoun BM (1994) Spectral envelope coding in cat primary auditory cortex: properties of ripple transfer functions. *Auditory Neuroscience* **1**:39–62.
- Schreiner CE and Langner G (1988) Periodicity coding in the inferior colliculus of the cat: II. Topographical organization. *Journal of Neurophysiology* **60**:1823–1840.
- Schreiner CE and Langner G (1997) Laminar fine structure of frequency organization in auditory midbrain. *Nature* **388**:383–386.
- Schuknecht HF (1960) Neuroanatomical correlates of auditory sensitivity and pitch discrimination in the cat. In: Rasmussen GL and Windle WF (eds). *Neural Mechanisms of the Auditory and Vestibular Systems*. Charles C Thomas, Springfield, pp. 76–90.
- Semple MN and Aitkin LM (1979) Representation of sound frequency and laterality by units in central nucleus of cat inferior colliculus. *Journal of Neurophysiology* **42**:1626–1639.
- Semple MN and Kitzes LM (1985) Single-unit responses in the inferior colliculus: different consequences of contralateral and ipsilateral auditory stimulation. *Journal of Neurophysiology* **53**:1467–1482.
- Servièrè J, Webster WR, and Calford MB (1984) Iso-frequency labelling revealed by a combined (<sup>14</sup>C)-2-deoxyglucose, electrophysiological, and horseradish peroxidase study of the inferior colliculus of the cat. *Journal of Comparative Neurology* **228**:463–477.
- Shamma SA, Versnel H, and Kowalski N (1995) Ripple analysis in ferret primary auditory cortex. I. Response characteristics of single units to sinusoidally rippled spectra. *Auditory Neuroscience* **1**:233–254.
- Sivaramakrishnan S and Oliver DL (2001) Distinct K currents result in physiologically distinct cell types in the inferior colliculus of the rat. *Journal of Neuroscience* **21**:2861–2877.
- Smith PH (1992) Anatomy and physiology of multipolar cells in the rat inferior collicular cortex using the in vitro brain slice technique. *Journal of Neuroscience* **12**:3700–3715.
- Smolders JWT, Aertsen AMHJ, and Johannesma PIM (1979) Neural representation of the acoustic biotope. *Biological Cybernetics* **35**:11–20.
- Snyder RL and Sinex DG (2002) Immediate changes in tuning of inferior colliculus neurons following acute lesions of cat spiral ganglion. *Journal of Neurophysiology* **87**:434–452.
- Snyder RL, Sinex DG, McGee JD, and Walsh EW (2000) Acute spiral ganglion lesions change the tuning and tonotopic organization of cat inferior colliculus neurons. *Hearing Research* **147**:200–220.
- Spirou GA and Young ED (1991) Organization of dorsal cochlear nucleus type IV unit response maps and their relationship to activation by bandlimited noise. *Journal of Neurophysiology* **65**:1750–1768.
- Stiebler I (1986) Tone-threshold mapping in the inferior colliculus of the house mouse. *Neuroscience Letters* **65**:336–340.
- Stiebler I (1987) *Frequenzrepräsentation und Schallempfindlichkeit im Colliculus inferior und auditorischen Cortex der Hausmaus (Mus musculus)*. Konstanzer Dissertationen, Volume 173, Hartung-Gorre, Konstanz.

- Stiebler I and Ehret G (1985) Inferior colliculus of the house mouse. I. A quantitative study of tonotopic organization, frequency representation and tone-threshold distribution. *Journal of Comparative Neurology* **238**:65–76.
- Suga N (1964) Single unit activity in cochlear nucleus and inferior colliculus of echolocating bats. *Journal of Physiology (London)* **172**:449–474.
- Suga N (1969) Classification of inferior collicular neurons of bats in terms of responses to pure tones, FM sounds and noise bursts. *Journal of Physiology (London)* **200**:555–574.
- Suga N (1977) Amplitude-spectrum representation in the Doppler-shifted CF processing area of the auditory cortex of the mustache bat. *Science* **196**:64–67.
- Suga N and Manabe T (1982) Neural basis of amplitude-spectrum representation in auditory cortex of the mustached bat. *Journal of Neurophysiology* **47**:225–255.
- Suga N, Gao E, Zhang Y, and Olsen JF (2000) The corticofugal system for hearing: recent progress. *Proceedings of the National Academy of Sciences of the United States of America* **97**:11807–11814.
- Sutter ML and Schreiner CE (1995) Topography of intensity tuning in cat primary auditory cortex: single-neuron versus multiple-neuron recordings. *Journal of Neurophysiology* **73**:190–204.
- Syka J, Popelář J, Druga R, and Vlková A (1988) Descending central auditory pathway—structure and function. In: Syka J and Masterton RB (eds). *Auditory Pathway. Structure and Function*. Plenum Press, New York, pp. 279–292.
- Syka J, Popelář J, and Kvašňák E (2000) Response properties of neurons in the central nucleus and external and dorsal cortices of the inferior colliculus in guinea pig. *Experimental Brain Research* **133**:254–266.
- Theunissen FE, Sen K, and Doupe AJ (2000) Spectral-temporal receptive fields of nonlinear auditory neurons obtained using natural sounds. *Journal of Neuroscience* **20**:2315–2331.
- Vartanian IA, Egorova MA, and Ehret G (1999) Expression of the main properties of critical bands in the neuronal activity of posterior quadrigemini in mice. *Doklady Biological Sciences* **368**:437–439 (in Russian).
- Vater M, Kössl M, and Horn AKE (1992) GAD- and GABA-immunoreactivity in the ascending auditory pathway of horseshoe and mustache bats. *Journal of Comparative Neurology* **325**:183–206.
- Versnel H and Shamma SA (1998). Spectral-ripple representation of steady-state vowels in primary auditory cortex. *Journal of the Acoustical Society of America* **103**:2502–2514.
- Viemeister NF (1988) Psychophysical aspects of auditory intensity coding. In: Edelman GM, Gall WE, and Cowan WM (eds). *Auditory Function. Neurobiological Bases of Hearing*. John Wiley & Sons, New York, pp. 213–241.
- von Békésy G (1960) *Experiments in Hearing*. McGraw-Hill, New York.
- Wagner T (1994) Intrinsic properties of identified neurones in the central nucleus of mouse inferior colliculus. *NeuroReport* **6**:89–93.
- Wang J, Salvi RJ, and Powers N (1996) Plasticity of response properties of inferior colliculus neurons following acute cochlear damage. *Journal of Neurophysiology* **75**:171–183.
- Webster WR, Servièrre J, Crewther D, and Crewther S (1984) Iso-frequency 2-DG contours in the inferior colliculus of the awake monkey. *Experimental Brain Research* **56**:425–437.

- Wenstrup JJ, Mittmann DH, and Grose CD (1999) Inputs to combination-sensitive neurons of the inferior colliculus. *Journal of Comparative Neurology* **409**:509–528.
- Willott JF and Urban GP (1978) Response properties of neurons in nuclei of the mouse inferior colliculus. *Journal of Comparative Physiology A* **127**:175–184.
- Winer JA, Larue DT, and Pollak GD (1995) GABA and glycine in the central auditory system of the mustache bat: structural substrates for inhibitory neuronal organization. *Journal of Comparative Neurology* **355**:317–353.
- Winter IM and Palmer AR (1995) Level dependence of cochlear nucleus onset unit responses and facilitation by second tones or broad-band noise. *Journal of Neurophysiology* **73**:141–159.
- Yan J and Ehret G (2001) Corticofugal reorganization of the midbrain tonotopic map in mice. *NeuroReport* **12**:3313–3316.
- Yan J and Ehret G (2002) Corticofugal modulation of midbrain sound processing in the house mouse. *European Journal of Neuroscience* **16**:119–128.
- Yang L, Pollak GD, and Resler C (1992) GABAergic circuits sharpen tuning curves and modify response properties in the mustache bat inferior colliculus. *Journal of Neurophysiology* **68**:1760–1774.
- Young ED, Shofner WP, White JA, Robert JM, and Voigt HF (1988) Response properties of cochlear nucleus neurons in relationship to physiological mechanisms. In: Edelman GM, Gall WE, and Cowan WM (eds). *Auditory Function. Neurobiological Bases of Hearing*. John Wiley & Sons, New York, pp. 277–312.
- Zerlin S (1986) Electrophysiological evidence for the critical band in humans. *Journal of the Acoustical Society of America* **79**:1612–1616.
- Zhang Y and Suga N (1997) Corticofugal amplification of subcortical responses to single-tone stimuli in the mustached bat. *Journal of Neurophysiology* **78**:3489–3492.
- Zhou X and Jen PH-S (2000) Corticofugal inhibition compresses all types of rate-intensity functions of inferior collicular neurons in the big brown bat. *Brain Research* **881**:62–68.
- Zurita P, Villa AEP, de Ribaupierre Y, de Ribaupierre F, and Rouiller EM (1994) Changes of single unit activity in the cat's auditory cortex associated to different anesthetic conditions. *Neuroscience Research* **19**:303–316.
- Zwicker E and Feldtkeller R (1967) *Das Ohr als Nachrichtenempfänger*. Hirzel, Stuttgart.
- Zwicker E, Flottorp G, and Stevens SS (1957) Critical band width in loudness summation. *Journal of the Acoustical Society of America* **29**:548–557.

# Chapter 12

## Temporal Coding in the Auditory Midbrain

ADRIAN REES AND GERALD LANGNER

### 1. INTRODUCTION

#### *1.1. THE BIOLOGICAL SIGNIFICANCE OF TEMPORAL CODING*

The cochlear frequency analyzer is a remarkable biological machine, and within the approximately 30,000 parallel spectral channels of its output in the cochlear nerve temporal stimulus components are highly conserved. In many communication sounds such as speech, it is often the changes in amplitude and frequency that carry information rather than their absolute values (Fig. 12.1; Rosen 1992; Shannon et al. 1995; Smith et al. 2002). Here, we consider the temporal modulations essential for auditory perception and show that the inferior colliculus (IC) is critical in the representation and transformation of temporal information from peripheral levels.

#### *1.2. TEMPORAL INFORMATION CONTENT OF NATURAL SOUNDS*

Temporal information in sound stimuli occurs over a wide time scale from a few Hertz up to about 1000 Hz (Joris et al. 2004). Temporal information is important in auditory perception in two ways. Dynamic changes bear information and denote differences between sounds, as in speech consonants in which the direction of a frequency change distinguishes consonants while changes in voice fundamental frequency, or prosody, denote interrogatives or indicate mood. Temporal information may also help to separate sound sources by stream segregation.

#### *1.3. TEMPORAL CODING FOR STIMULUS SEGREGATION AND GROUPING*

The auditory system can separate and identify concurrent sound sources that contain many, overlapping frequency components. Frequency decomposition in the auditory periphery operates on the sum of all the sound waves reaching the ear. The recovery of a sound source requires a grouping or streaming of its frequency components, allowing them to be perceived as a whole (Bregman




Figure 12.1. Amplitude modulation in human speech. **(A)** Waveforms of utterances from adult male speaker low-pass filtered to show their amplitude envelopes. (Adapted from Rosen 1992.) **(B)** Average modulation envelope spectra extracted from 1 minute of connected speech from 10 male speakers. Modulation index for one-third octave bands centered on the carrier frequency shown adjacent to each curve. *Arrows* indicate the dominant modulation frequency associated with different speech components. (Replotted from Plomp 1983.)

1990). Cues contributing to this identification process include onset and offset times and ongoing temporal envelopes (Darwin and Carlyon 1995). Frequency glides may fuse components into a stream, and tracking the fundamental frequencies of harmonic complexes is important in stream segregation (Moore 2003).

Other psychophysical phenomena suggest interactions between the envelopes of components widely separated in carrier frequency, including comodulation masking release (Hall et al. 1984) and modulation detection interference (Yost and Sheft 1989, 1993). Both phenomena imply that interactions must occur between different frequency channels to extract common envelope patterns. The IC may play a crucial role in this process.

#### *1.4. TEMPORAL CODING OF PITCH*

When the frequency range of harmonic sounds is sufficiently broad to activate different frequency channels, the neural responses in many channels will be elicited by a waveform that is a superposition of the several frequency components within the channel. Because all harmonic sound components are integer multiples of its fundamental frequency, their superposition periodicity on the basilar membrane will correspond to that of the fundamental frequency or a multiple of it. Analyzing this periodicity and comparing temporal information over different frequency channels enhances the auditory system's capacity to detect communication signals in a noisy environment (Cherry and Sayers 1956).

A by-product of this temporal analysis is periodicity pitch. The timbre of a periodically modulated sound depends on its spectral envelope while the sound's perceived pitch may be identical to a pure tone whose frequency equals that of the sound's temporal envelope. Pitch depends also on the periodic sound's spectral content, as in the first and second effect of pitch shift with inharmonic amplitude modulation (AM) signals (Schouten 1940, 1970; Schouten et al. 1962). Periodicity pitch may be perceived similarly by humans and animals (cats: Chung and Colavita 1976; Heffner and Whitfield 1976; birds: Cynx and Shapiro 1986; monkeys: Tomlinson and Schwarz 1988; Wright et al. 2000; chinchilla: Shofner and Yost 1997; Shofner 2000). Neuronal mechanisms for periodicity pitch likely rely on the temporal encoding up to 5 kHz in the auditory nerve (Greenberg 1988), which is the upper limit of musical pitch (Semal and Demany 1990).

## **2. ASPECTS OF TEMPORAL INFORMATION CODING**

### *2.1. TEMPORAL RESPONSES OF INFERIOR COLLICULUS NEURONS TO TONES AND NOISE*

Before considering IC neural responses to temporally varying stimuli, we note their temporal firing patterns to constant pure tones or noise bursts, which have



been studied in many species (Rose et al. 1963; Ryan and Miller 1978; Syka et al. 2000).

The most obvious distinction is between cells that fire only to the stimulus onset and those with sustained firing patterns. The proportions of units in these classes vary between studies, perhaps because of species variations, anesthesia, stimulus duration, repetition rate, or the cell's locus in the IC. Onset units comprise about 50% of the cells in anesthetized cats (Rose et al. 1963) and tranquillized mice (Walton et al. 1998) and 27% in the guinea pig central nucleus of the IC (CNIC) (Syka et al. 2000).

Pauser, on-sustained, chopper, and sustained/buildup response subgroups exist within the broader classes (Willott and Urban 1978; Ehret and Moffat 1985; Le Beau et al. 1996; Rees et al. 1997; Syka et al. 2000). However, response patterns are not fixed for a neuron under all stimulus conditions and response patterns may differ for frequencies remote from the characteristic frequency (CF) or for other stimulus levels (Ehret and Moffat 1985).

IC response patterns are strongly influenced by both  $\gamma$ -aminobutyric acid (GABA) and glycinergic inhibition (Faingold et al. 1989; Vater et al. 1992; Le Beau et al. 1996). Blockade of inhibition by bicuculline or strychnine elevates firing rate and may change its response pattern. Half of the cells changed their firing pattern with bicuculline, of which 70% of these became choppers (Le Beau et al. 1996). Intracellular recordings in vitro (see Chapters 2, 9, and 10) or in vivo (Nelson and Erulkar 1963; Covey et al. 1996; Kuwada et al. 1997) are revealing the underlying mechanisms for these changes. The inhibitory currents driving the response patterns frequently precede the excitatory input to the cell (Nelson and Erulkar 1963; Covey et al. 1996; Pedemonte et al. 1997) and specific ion channels influence the intrinsic neural membrane properties and their response patterns.

## 2.2. SYNCHRONIZATION CODE

Temporal stimulus waveform information can be represented by the temporal synchronization of neuronal responses to the stimulus envelope. Period histograms reveal the spike-timing relative to the stimulus modulation period. A quantitative measure derived by circular statistics represents the period of a signal as a circle, a spike by a vector  $\vec{a}_i$  with unit length, and the phase angle by vector orientation (Goldberg and Brown 1969). The sum  $\vec{R} = \sum_{i=1}^N \vec{a}_i$  over all  $N$  spike vectors of a response to a periodic signal is the mean synchronization, and the mean phase angle. Normalization gives the so-called vector strength  $r = \left| \sum_{i=1}^N \vec{a}_i \right| / N$ , a measure of average synchronization, which equals 1 for maximal and 0 for no synchronization.

If the IC participates in temporal processing, its input should be synchronized to stimulus envelope fluctuations. Synchronized responses are not "reproductions




Figure 12.2. Synchronized response to amplitude modulation for a single neuron in IC of guinea pig. Period histograms show distribution of spike firing over one cycle of modulation for three modulation frequencies ( $f_{\text{mod}}$ ). Center: Modulation transfer function for synchronized response. (Adapted from Rees and Sarbaz 1997.)

of the waveform” (Møller 1974), because they are restricted to a small portion of the modulation cycle. Envelope fluctuations of acoustic signals are amplified in neuronal response fluctuations in the rat IC, with a gain up to 20 dB (Rees and Møller 1983; Rees and Palmer 1989; Fig. 12.2). Similarly, potentials evoked by modulation frequencies  $<100$  Hz in the frog midbrain show enhanced AM coding relative to auditory nerve evoked potentials (Hillery 1984). Synchronization deteriorates at higher levels in the auditory system, significantly reducing temporal resolution. However, some midbrain neurons can synchronize to modulation frequencies up to 800 or 1000 Hz (Schuller 1979, 1984; Rose and Capranica 1985; Epping and Eggermont 1986; Lesser et al. 1986, 1990; Langner and Schreiner 1988) or even up to 1200 Hz (Langner 1983).

### 2.3. RATE CODE

A second feature for neuronal periodicity coding is average response rate. Many IC neurons fire maximally for a particular modulation frequency (Langner 1983; Müller-Preuss et al. 1994). In cat most IC units were rate-tuned, some up to 1000 Hz, while tuning in synchronization above 300 Hz was rare (Langner and Schreiner 1988; Fig. 12.3), suggesting a midbrain synchronization-to-rate trans-




Figure 12.3. Normalized band-pass modulation transfer functions of neurons in cat CNIC. In contrast to more peripheral auditory nuclei, many units in the CNIC are tuned in terms of average firing rate. Numbers above the curves indicate the CF of the unit that also served as carrier frequency of the AM stimuli and the maximal firing rate corresponding to the tip of the MTF. (Adapted from Langner and Schreiner 1988.)

formation for higher modulation frequencies. In guinea pig and gerbil, fewer units show rate tuning and modulation cutoff frequencies match those for synchronization measures (Rees and Palmer 1989; Krishna and Semple 2000).

Neurons in the dorsal and posteroventral cochlear nucleus show near-constant average rates across modulation frequency (Kim et al. 1990). The auditory mid-brain might transform synchronized envelope information into a rate/space representation (Langner 1981; Epping and Eggermont 1986; Pinheiro et al. 1991; Langner 1992) as seen when guinea pig IC mean rates were largest at modulation frequencies that elicited maximal synchronization (Rees and Palmer 1989). Additional representation of temporal information in the timing of neural activity or the transmission of low ( $< 200$  Hz) modulation frequencies to higher centers may also be present.

#### 2.4. ACROSS-FREQUENCY ANALYSIS

Psychoacoustic experiments show that across-frequency-channel processes occur (see Section 1.3) with a particular role for harmonicity (Treurniet and Boucher 2001). Bat IC (and higher level) neurons may respond to frequencies outside

their pure tone response area when presented in conjunction with other components of their sonar or social calls (Wenstrup and Grose 1995; Mittmann and Wenstrup 1995; Misawa and Suga 2001). This combination sensitivity can be generated within the IC (Wenstrup and Leroy 2001). Guinea pig IC neurons receive input from a wide frequency range that may sum nonlinearly for two-tone stimuli (Thornton and Rees 2001). In chinchilla IC, integration related to harmonically structured social calls is common (Biebel and Langner 2002).

### 3. AMPLITUDE MODULATION CODING

Sinusoidal amplitude modulations (SAMs) are useful test stimuli for investigating periodicity coding. A signal's AM spectrum contains the carrier frequency ( $f_c$ ) and two side-bands,  $f_c + f_m$  and  $f_c - f_m$  which cochlear frequency analysis cannot resolve at low modulation frequencies ( $f_m$ ). The SAM envelope frequency corresponds naturally to  $f_m$  and, when the carrier is an integer multiple of the modulation frequency, it may be "harmonic." Therefore, SAM stimuli contain critical features of temporally varying sounds.

#### 3.1. MODULATION TRANSFER FUNCTIONS

For linear systems the average response to an AM is independent of the modulation frequency as seen in the auditory nerve (Rose and Capranica 1985) and for many dorsal cochlear nucleus (DCN) neurons and posteroventral cochlear nucleus (PVCN) neurons (Schreiner and Snyder 1987; Kim et al. 1990), but not for IC neurons (Langner 1983; Langner and Schreiner 1988; Krishna and Semple 2000).

For nonlinear systems such as the auditory system, the best way to measure the modulation transfer function (MTF) is with AM, presenting one modulation frequency at a time (Schroeder 1981). Synchronization MTFs are derived by calculating vector strength, whereas rate MTFs are measured from the average discharge rate. Responses to AM show changes in synchronization and average discharge rate as a function of modulation frequency described as band-pass or low-pass MTFs (Rees and Møller 1983; Reimer 1987; Langner and Schreiner 1988; Rees and Palmer 1989). The maximum of a band-pass MTF or the cutoff frequency of a low-pass MTF is the "best modulation frequency" (BMF). In cat IC, 33% of the cells had band-pass synchronization MTFs and 48% had low-pass characteristics (Langner and Schreiner 1988). Seventy percent of cat IC single units had rate MTFs with a band-pass characteristics, 7% low-pass characteristics (Langner and Schreiner 1988). Other MTF-types distinguished are high-pass, band-reject, or nonselective/complex (Bibikov and Gorodetskaya 1980; Walkowiak 1984; Rose and Capranica 1985; Langner and Schreiner 1988).

If the temporal code representation of periodicity is partly transformed into a rate code in the midbrain, then activation strength signals stimulus periodicity, just as pure tone frequency is signalled by the response at a particular place in the tonotopic map (Langner et al. 2002). Because the CNIC is tonotopically and

periodotopically organized, neural activation signals simultaneously a certain carrier frequency and a certain envelope periodicity.

### 3.2. *FACTORS AFFECTING AMPLITUDE MODULATION CODING*

Many aspects of midbrain AM responses are insensitive to signal intensity variations and modulation depth. Kitten CNIC cells synchronize to AM waveforms across stimulus levels and modulation depths, and mean firing phase during the modulation cycle is nearly independent of level and modulation depth (Brugge et al. 1993). Other cells show changes in MTF shape and BMF with mean intensity as MTFs shift from band-pass to low-pass at low sound intensities (Rees and Møller 1987; Krishna and Semple 2000). Modulation depth changes do not affect MTF shape and peak value although IC cells in many species are sensitive to AM and may synchronize to modulation depths of only 2% (Schuller 1979; Bibikov and Nizamov 1996; Biebel et al. 1998).

To encode AM information temporally, a frequency filter needs a bandwidth exceeding twice the modulation frequency to avoid sideband attenuation. Because peripheral tuning curve width increases with CF, an increased upper frequency limit of encoded modulations is expected and has been found (Langner and Schreiner 1988). At CFs < 4 kHz, virtually all units have BMF values below CF/4, which corresponds to the upper boundary of the psychophysical existence region of periodicity pitch (Terhardt 1970). In conclusion, spectral tuning width of auditory nerve fibers may restrict the upper frequency limit of modulation and periodicity pitch coding. It is difficult to see the relationship between CF and BMF in a sample of neurons with only few BMFs > 100 Hz (Rees and Møller 1987; Krishna and Semple 2000).

AM signal carrier frequency may also influence MTFs. Small changes of the carrier frequency around a neuron's BF not only affect slopes of MTF shapes but also may alter BMFs systematically (Langner 1983; Langner and Schreiner 1988).

### 3.3. *INFLUENCE OF BACKGROUND NOISE ON AMPLITUDE MODULATION CODING*

At stimulus levels that generate a band-pass MTF, broadband noise reduces the modulation gain at low modulation frequencies, changing the MTF from band-pass to low-pass. It is as if adding noise reduces the effective level of the modulated stimulus.

Noise shifts IC pure-tone rate-level functions toward higher stimulus levels (Rees and Palmer 1988). For a modulated stimulus at an average level that gives a band-pass MTF in quiet, broadband noise shifts the rate-level function and moves the stimulus lower on the cell's input-output function. This change yields a low-pass transfer function, like that for a low-level stimulus in quiet (Fig. 12.4). A concomitant change in modulation gain at low modulation frequency with the shift in the rate-level function supports this idea (Rees and Palmer 1989).




Figure 12.4. Influence of broadband noise on the synchronized responses to sinusoidal amplitude modulation in guinea pig IC. **(A)** Synchronization modulation transfer functions at two different carrier levels (9 dB, *solid line*, and 39 dB re threshold, *long dash*) and for the 39 dB carrier level presented in different levels of broadband noise ( $N$  parameter, noise spectrum level). Carrier frequency and unit best frequency = 4.25 kHz, modulation depth = 50%. **(B, C)** Schematic representations of the rate-level functions **(B)** and MTFs **(C)** to explain the shift from band-pass to low-pass MTF in **(A)**. Point 1 on the rate-level function shows the position of the carrier giving a low-pass MTF. An increase in the level of the carrier (*solid arrow*) shifts the modulated stimulus to a higher level on the rate-level function **(B:2)**, and gives rise to a band-pass MTF **(C:2)**. Addition of broadband noise causes the rate-level function to shift to the right (*dashed function*) and the modulated stimulus now falls on a lower position on the rate-level function **(B:3 and dashed arrow)** resulting in the MTF changing from band-pass to low-pass MTF **(C:3 and dashed arrow)**. (Adapted from Rees and Palmer 1989.)

### 3.4. *RESPONSES TO NONSINUSOIDAL MODULATIONS*

Most studies of AM encoding use sinusoidal modulation waveforms, but advantages accrue in dissociating stimulus periodicity from envelope parameters such as waveform rise-time and asymmetry. That such properties are important can be seen in response histograms to SAM that often have a more peaked than sinusoidal appearance (Nelson et al. 1966; Rees and Møller 1983; Krishna and Semple 2000). At maximal modulation depths, peak position may change relative to the stimulus waveform and even become bimodal, consistent with the neurons tracking the amplitude envelope at low depths and other parameters, such as the amplitude change rate, at high depths. Sensitivity to rate of amplitude change is evident for exponential and triangular modulation envelopes (Poon and Chiu 1997; Fig. 12.5). Some responses to exponential envelopes showed two histogram peaks compared to one peak for triangular modulation. In most cells spikes occurred during the rising and falling phase, suggesting that these effects embody nonmonotonic or skewed frequency response areas. Four AM unit types were identified: type I units followed the modulation envelope; type II units followed only to the rising phase and were dependent on AM intensity, velocity, and range; type III combined features of types I and II while type IV responded to rising and falling intensity (Chiu and Poon 2000). Asymmetries in the rate and temporal responses of IC neurons are also seen for ramped and damped sinusoids (Neuert et al. 2001). Units with onset firing patterns showed high asymmetry to these stimuli, with most unresponsive to damped stimuli except at the shortest half-life. Sustained units showed little response asymmetry. More cells had asymmetrical rate responses to such stimuli than in the ventral cochlear nucleus (Pressnitzer et al. 2000), demonstrating enhanced IC response asymmetry.

The sensitivity of rat IC neurons to AM periodicity is similar for sinusoidal and for nonsinusoidal modulations studied with tones amplitude modulated with pseudorandom noise (Møller and Rees 1986; Rees and Møller 1987). MTFs obtained by cross correlating neural discharge with the noise modulator estimate the linear response component and closely resemble the SAM transfer function (Møller and Rees 1986). Responses to pseudorandom AM reveal that the linear component explains less of the total response than it does for cochlear nucleus cells (Møller and Rees 1986). Most nonlinear IC responses were even-order and might therefore represent asymmetries to rising and falling amplitude. Comparison of responses to a speech utterance in cat auditory nerve, cochlear nucleus, and IC with responses predicted by the linear component of the site's modulation response found that only IC responses departed significantly from the linear prediction (Delgutte et al. 1998).

### 3.5. *RELATIONSHIP OF AMPLITUDE MODULATION CODING TO OTHER NEURONAL PROPERTIES*

IC neurons show several temporal firing patterns when stimulated with pure tones (see Section 2.1). Onset and sustained neurons respond to continuous AM

tones under anesthesia (Rees and Palmer 1989) even at modulation depths well below 100%. In awake chinchilla, a phasic response often predicts narrowly tuned MTFs, whereas an additional tonic component indicates broader MTFs (Langner et al. 2002). However, in bats and rats onset units are the least likely to respond to AM and onset units were least responsive to AM (Condon et al. 1996; Palombi et al. 2001). These differences may represent species differences or anesthesia effects.

IC neurons also differ in the firing regularity of their responses to pure tones (Rees et al. 1997). Some fire regularly with preferred interspike intervals and their interval period is highly correlated with temporal best modulation frequency (Rees and Sarbaz 1997). In contrast, cells with band-pass modulation rate tuning usually fired irregularly to tones.

### 3.6. SPATIAL REPRESENTATION OF ENVELOPE PERIODICITY

#### 3.6.1. Periodotopy


By providing a substrate for relative position and spatial shifts of excitation, neuronal maps optimize information processing in the corresponding signal space (Suga and O'Neill 1979). The periodotopic map based on the temporal analysis of periodicity information in the CNIC is such a map (Fig. 12.6).

Systematic distributions of responses to AM rate are found in several species (Langner and Schreiner 1988; Langner et al. 1992, 2002; Heil et al. 1995). Periodotopy occurs in CNIC frequency-band laminae. In cat, low BMFs are represented medially and high BMFs laterally; even more lateral subareas with low BMFs, probably in the external nucleus of the IC (ENIC), resulted in the interpretation of concentric periodicity representation (Schreiner and Langner 1988). However, it turned out that recordings in CNIC alone show a simple mediolateral gradient orthogonal to the main dorsoventral tonotopic gradient (Langner et al. 2002; Fig. 12.6). The local BMF increased from about 20 Hz medially to CF/4 (for CF <4 kHz) laterally. Moreover, within the frequency-band laminae, the periodicity map is organized orthogonal to the intralaminar tonotopic fine structure (Langner 1992; Fig. 12.7).

Single-unit findings are supported by *c-fos* mapping (Büttner et al. 1997) and metabolic labeling with [<sup>14</sup>C]2-deoxyglucose (Langner 2004). Spatial representation of periodicity information is also transferred to higher centers of the avian forebrain (Hose et al. 1987), cat cortex (Langner 1997), and human auditory cortex (Langner et al. 1997).

Figure 12.5. Responses of IC neuron to triangular and exponential amplitude modulation envelopes. (A) Triangular modulation waveforms (arbitrary units, a.u.) of a pure tone at 4.8 kHz and 30 dB above threshold (*upper*) and neuronal responses to this stimulus shown as a dot raster plot (*middle*) and a histogram of spike counts (*spk cnt, lower*). The response is single-peaked for the triangular, and double-peaked for the exponential, envelope indicative of a response to the rate of change of amplitude. (Adapted from Poon and Chiu 1997.)








Figure 12.6. Mapping of periodicity and latency in the CNIC. **(A)** Distribution of 52 BMFs in a frequency-band of the chinchilla CNIC. The plane corresponds to the frequency-band lamina of 6 kHz, while the vertical axis represents approximately 5 octaves of BMFs. Low-BMF neurons are located medially and high-BMF neurons laterally. **(B)** Latency map for the same 52 neurons plus latencies from an additional 31 units. Neurons with long latencies are located medially and neurons with short latencies laterally. Note the different viewing orientations in **(A)** and **(B)**.




Figure 12.7. Orthogonality of tonotopy and periodotopy in a frequency-band lamina of cat CNIC. In this particular lamina, CFs increased from approximately 2.1 kHz rostromedially to CFs approximately 2.5 kHz located laterally; lowest BMFs were observed in a caudomedial corner and highest BMFs in the rostralateral corner.

Periodicity, like frequency, seems to be mapped logarithmically in the CNIC, providing scaling invariance in mapping and integration of information from broad band harmonic sounds (Fig. 12.8). Together with the orthogonality of tonotopic and periodotopic gradients, this suggests the periodotopic axis as a second functional axis of the auditory system (Langner 2004).

### 3.6.2. Relationship of Periodotopy to Other Neuronal Maps

CNIC maps for other auditory parameters, such as onset latency for pure tone responses, also exist. The spatial gradient for onset latency closely resembles that of periodicity maps and, accordingly, is orthogonal to the tonotopic map in many species (Schreiner and Langner 1988; Hattori and Suga 1997; Walton et al. 1998; Biebel and Langner 2002). On average, neurons with low BMFs have long onset latencies and neurons with high BMFs have short onset latencies (Langner et al. 1987, 2002; Condon et al. 1996). Because of the linear relation between latency and  $1/\text{BMF}$  (Langner et al. 1987) it would, therefore, suffice to measure latency and calculate a periodicity map by latency information. In bats the latencies within the CNIC could correspond to the echo delays (Yan and Suga 1996). However, no map of IC best echo delays was found, although latency was organized topographically (Portfors and Wenstrup 2001).

Other parameters mapped in the CNIC include threshold (Stiebler and Ehret 1985), tuning curve bandwidth, and binaural input characteristics (Schreiner and Langner 1988; see Chapter 11). However, none of these parameters is represented in a monotonic map with a single gradient as are periodicity and latency.

### 3.7. PUTATIVE MECHANISMS FOR AMPLITUDE MODULATION CODING

Models using a combination of short-term adaptation, refractory properties, intrinsic oscillations, coincidence detection, or temporal integration can explain




Figure 12.8. A functional and topographic model of periodicity processing. **(Left)** A black-box version of a correlation model. After the frequency analysis of the cochlea and further processing in DCN and VCN periodicity information converges with different delays on coincidence neurons in the CNIC. Oscillator (chopper) in the VCN and reducer (pauser) responses in the DCN are extremely broadly tuned, but precisely synchronized to the signal envelopes by means of a trigger (onset-) neuron. The relatively long delay—resulting from the integration time of the reducer circuit—has to be compensated by the period of the signal and the period of the oscillation for the coincidence unit to be activated and is crucial to its BMF. Although the coincidence of inputs from DCN and VCN provides a correlation between the fine-structure (or carrier) of the signal and its envelope modulation an additional synchronized inhibition provided by neurons in the VNLL shapes the resulting comb-filter into a band-pass MTF. To cover the relevant spectral and temporal ranges many such circuits are necessary in parallel for the periodicity analysis. **(Right)** A topographic model of the CNIC. In accordance with anatomical findings the model suggests that coincidence units (disc-cells) are arranged in neuronal laminae, oriented approximately orthogonal to the main frequency gradient. In each of about 30 to 40 frequency-band laminae, CF increases over a small range of frequencies orthogonal to the main frequency gradient of the CNIC, while BMF increases along an isofrequency-line from about 10Hz to  $CF/4$  ( $<1000$  Hz) with the highest BMFs located at the lateral border of the CNIC. The CF-ranges given for each of the five selected laminae result from a mapping study of the CNIC of cat (Schreiner and Langner 1997).

many aspects of modulation coding (Langner 1983, 1992; van Stokkum 1987, 1989).

A “cancellation model” for a hypothetical “neural comb filter” employs neural delay lines, as already suggested by Licklider (1951), and inhibitory synapses (de Cheveigné 1993). A similar model network that matches the periodicity of tuned neurons in the fish auditory midbrain (*Pollimyrus*) uses coincidence of excitatory and inhibitory input and post-inhibitory rebound, enhanced by neural oscillations (Large and Crawford 2002).

To explain why the periodicity pitch of AM signals (Schouten 1970) and the BMFs of midbrain neurons depend on the carrier and the modulation frequency (Langner 1983; Langner and Schreiner 1988), a model was proposed which

performs a correlation between signal fine-structure and envelope (Langner 1981, 1983, 1988, 1992; Borst et al. 2004). Neuronal elements include a trigger, an oscillator, a reducer unit (reducing high-frequency information), and a coincidence detector. These elements may correspond to ventral cochlear nucleus (VCN) on-type chopper neurons, DCN pauser neurons, and disc-shaped CNIC neurons (Langner and Schreiner 1996). The trigger unit is activated by each modulation cycle and synchronizes the oscillator and the reducer. The oscillator responds with regular interspike intervals for each modulation cycle. The reducer makes use of synchronized activity of many nerve fibers in accordance with the volley principle (Wever 1949) and provides regular intervals calibrated to signal fine structure with phase delays equal to an integer multiple of the carrier period. The output is from coincidence units that receive input from the oscillator and the reducer circuit. Therefore, the coincidence unit responds when carrier and envelope frequency are correlated and the envelope period matches the reducer delay. Finally, to avoid responses to periods corresponding to integer fractions of the preferred period resulting in comb-filter type MTFs (Langner 2004), the coincidence unit requires additional inhibitory input synchronized to the envelope period. An appropriate inhibitory input to the IC may arise from neurons in the ventral nucleus of the lateral lemniscus (VNLL) (Vater et al. 1997; Covey and Casseday 1999; Riquelme et al. 2001). In support of this idea, VNLL cells receive giant synapses mainly from on-type neurons in cat, human (Adams 1997), and bat (Vater et al. 1997) CN.

In an alternative model, BMF preferences of CNIC neurons reflect convergence of inputs from VCN choppers with the same chopping frequency and, therefore, sensitivity to the same modulation frequency (Hewitt and Meddis 1994). Indeed, in many CNIC neurons a chopper-like intrinsic oscillation occurs after signal onset and often to each modulation cycle (Langner 1983; Langner and Schreiner 1988; Condon et al. 1996). However, the intervals of these onset oscillations are usually only a fraction (about 1/6) of the BMF periods of the neurons and their oscillation alone cannot explain unit BMF preference (Langner et al. 2002). However, after a certain latency in the onset response additional spike clusters occur at regular intervals in ongoing responses (Langner 1983). In contrast to onset oscillations, these intervals correlate with the neuron's BMF (Rees and Sarbaz 1997).

## 4. FREQUENCY MODULATION CODING

Frequency modulations (FMs) are important elements of many natural sounds including vocalizations and speech and of bat sonar signals. The earliest reports of IC function described FM-selective IC neurons, including some that respond to dynamic frequency changes but not to pure tones.

### *4.1. RESPONSES TO SINUSOIDAL FREQUENCY MODULATIONS*

Many units responding to sinusoidal FM (SFM) do so with an approximately sinusoidal modulation of their neural discharge (Nelson et al. 1966; Casseday

et al. 1997). Such neurons show great sensitivity to FM with a substantial modulation of the response at depths of <10% of the carrier. As with AM sounds, rat IC neurons have lower best modulation frequencies (<100 Hz) for SFM than cochlear nucleus cells, and similar BMF ranges for the two types of modulation (Rees and Møller 1983). Indeed, as an FM sweeps through the neuron's tuning curve it effectively generates an AM.

Some bat neurons are SFM-selective and responded best or only to SFM (as opposed to frequency sweeps); the larger group of nonselective units had best-SFM frequencies <180 Hz (Casseday et al. 1997). Some units were also modulation-depth selective. As for many IC responses, it is difficult to establish whether such specializations arise within the IC or below it. However, similar stimuli analyzed in the lateral lemniscal nuclei show significantly higher BMFs (Huffman et al. 1998), suggesting that further processing occurs in the IC. Similar findings are reported for SAM (Yang and Pollak 1997).

IC inhibition helps to limit neural responses to low SFM rates. GABA<sub>A</sub> or glycine receptor antagonists increased the upper, and/or decreased the lower, cutoff values of the SFM rate response, but in only one-third of the units was the change >50% (Koch and Grothe 1998). Whether these mechanisms alone can account for SFM selectivity differences between the IC neurons and those in more peripheral structures is unknown.

#### 4.2. RESPONSES TO FREQUENCY MODULATION SWEEPS

Frequency sweeps are powerful tools for studying FM responses in the IC (Suga 1965; Clopton and Winfield 1974; Fuzessery 1994; Gordon and O'Neill 1998; Hage and Ehret 2003). These stimuli, unlike SFM, enable rate of frequency change to be dissociated from stimulus periodicity and permit identification of directional preference for frequency glides. They are important for studying responses in echolocating bats whose calls contain frequency sweeps. Evidence for directional selectivity as well as specificity to FM was first reported by Suga (1965, 1969). Specific IC regions contain higher percentages of FM-sensitive neurons. In the pallid bat (*Antrozous p. pallidus*), which combines echolocation and passive listening, about 50% of the ventral (high frequency) IC neurons prefer the descending frequency sweeps in its echolocation call and about a third responded only to them and not to pure tones or noise bursts. In contrast, in the dorsal, low-frequency IC, few such neurons were found (Fuzessery 1994). In the mustached bat (*Pteronotus p. parnellii*) many units responding to upward frequency glides were found in a specific part of the external nucleus (Gordon and O'Neill 2000).

Two studies have addressed the relationship between the response to sweeps and IC frequency band laminae. In rat, 34% of units responded to FM sweeps but not to *continuous* pure tones and 43% responded to both (Poon et al. 1992). Cells responding only to FM preferred higher rates and wider ranges of frequency change than other neurons. Most FM-specialized cells also had a strong preference for the FM sweep direction. Anatomically, FM-specialized cells were

multipolar with larger, more extensively branched dendritic fields and more dendritic spines. FM cells were found in all three IC subdivisions, with a higher proportion in EN and fewest in DC. FM response differences may thus depend on cellular morphology and IC location. The best FM ranges were highly correlated with the degree of dendritic extension along the tonotopic IC axis (Poon et al. 1992). However, other rat data found little evidence for FM selective cells and only small changes in response strength for different directions of sweep (Felsheim and Ostwald 1996) and selectivity for sweep speed showed band-pass tuning in many neurons. The use of exponential rather than linear sweeps in the latter study might account for discrepancies between them.

Selectivity for frequency sweep speed is represented topographically in mouse IC (Hage and Ehret 2003). The proportion of cells responding to high sweep speeds increased with distance from the laminar center, as did the firing rate to high, and to a lesser extent, medium sweep speeds. These distributions correlated with those for the units' frequency response area type, sharpness of tuning, and temporal response pattern (see Chapter 11). The authors propose that a common feature underlying all these distributions is a gradient of increasing inhibition from the lamina edge to its center. Some spatial segregation of directional selectivity is seen, although its precise form and correspondence with sweep speed representation is less clear (Hage and Ehret 2003).

#### 4.3. PUTATIVE MECHANISMS FOR FREQUENCY MODULATION CODING

Asymmetric side band inhibition has long been hypothesized to be involved in determining FM directional specificity (Suga 1965; Gordon and O'Neill 1998). The principle is that an FM-selective cell fails to respond when the frequency trajectory of a stimulus crosses an inhibitory region of its frequency receptive field before entering the excitatory region. Variants on this theme have been proposed to explain complexities such as the unresponsiveness of such FM-specialized neurons to pure tones (Fuzessery and Hall 1996). These accounts either postulate two subthreshold excitatory inputs with different latencies that sum only when a frequency sweep travels in one direction, or combinations of excitatory and inhibitory inputs. The appropriate time delays may be created by inputs with different latencies (Fuzessery 1994; Casseday et al. 1997) or by differences in spatial distributions of frequency specific inputs along the cell's dendrites (Rall 1962; Erulkar et al. 1968). A frequency sweep in one direction results in greater spatiotemporal integration of the postsynaptic potential than an opposite sweep.

The postulated role of inhibition opens the model to testing with transmitter antagonists. Blockade of GABAergic inhibition in the pallid bat reduced directional selectivity for FM sweeps in some cells without changing selectivity for FM over tones or noise (Fuzessery and Hall 1996). However, in another bat (*Eptesicus fuscus*), few cells changed their directional selectivity to SFM with

GABA or glycinergic antagonists (Koch and Grothe 1998). These results suggest differences in FM processing between species, or stimulus differences (SFM vs. linear sweeps) but limitations of iontophoresis may also be significant.

Serotonin also modulates responses to FM stimuli in free-tailed bats (*Tadarida brasiliensis*) (Hurley and Pollak 1999). Both depression and facilitatory effects were observed, depending on the stimulus parameters. Thus, IC responses may be influenced by neuromodulatory systems.

## 5. OTHER ASPECTS OF TEMPORAL CODING

### 5.1. RESPONSES INFLUENCED BY TEMPORAL GAPS

The ability to detect a gap in an otherwise continuous signal is a useful psychophysical measure of temporal resolution (Viemeister and Plack 1993; Moore 2003). The advantage of such measures of temporal resolution is that, for a broadband noise, the magnitude spectrum is unaffected by the introduction of a brief gap allowing temporal acuity to be assessed without spectral changes.

The effect of temporal gaps on IC responses has been studied in mouse (Barsz et al. 1998; Wilson and Walton 2002). Mouse behavioral gap detection thresholds resembled thresholds derived from CNIC responses (Fig. 12.9). Onset units responded to each noise burst in the stimulus, while tonic units showed a reduced discharge during the gap and increased firing at the second noise burst onset (Walton et al. 1997). Phasic off-units and inhibitory units responded during the gap. Onset and primary-like tonic units showed the shortest detection thresholds of 1 to 2 ms. Neural gap thresholds for phasic-off and tonic inhibitory units were 3 to 6 times higher than for other unit types (Wilson and Walton 2002).

Gap thresholds for most units increase as the fall- and rise-times of the noise on either side of the gap are increased and the gap blurs (Barsz et al. 1998). Neural gap thresholds are higher in aged mice (Walton et al. 1998) which may provide a model for investigating the mechanisms underlying similar age-related changes in human temporal acuity (Viemeister and Plack 1993). A loss of inhibition may contribute to these changes, as behaviorally measured gap detection thresholds in aged gerbils improved after systemic drug application that enhanced the GABAergic system (Gleich et al. 2003).

### 5.2. RESPONSES AFFECTED BY TONE DURATION AND ECHO DELAYS

In echolocating bat, neurons sensitive to sound duration intervals (Pinheiro 1991; Fuzessery and Hall 1999) and echo delay (Feng et al. 1978; Yan and Suga 1996) exist in the IC. Bats vary the duration of their biosonar calls during different stages of hunting with the sounds usually becoming shorter and more frequent as they acquire their prey. The echo delay-time allows the bat to compute target distance.






Figure 12.9. Similarity of behavioral (*closed symbol*) and neurophysiological (*open symbols*) measurements of responses to gaps in broadband noise. Behavioral data obtained from five mice using inhibition of the acoustic startle reflex as the paradigm. Neurophysiological data taken from two neuronal types (onset  $n = 13$ , primary-like  $n = 3$ ) in CNIC in unanesthetized mouse. For onset units, the response is the reciprocal of the increase in firing immediately following the gap. The response for the primary-like units is the decrease in firing during the gap. In both cases the data are normalized to the equivalent time window in the response to noise with no gap present. (Adapted from Walton et al. 1997.)

Approximately 30% of big brown bat neurons increase firing rate significantly with sound duration. Some duration sensitive neurons responded to pure tones, others only to frequency sweeps. The tuning functions were band-pass and the duration selectivity matched the durations of the sounds emitted with most cells preferring durations  $< 10$  ms (Casseday et al. 1994; Ehrlich et al. 1997). Similar results were seen in pallid bats (Fuzessery and Hall 1999).

Inhibition is essential for duration tuning (Casseday et al. 1994). GABA<sub>A</sub> or glycine antagonists reduce or eliminate duration tuning, but in cells tested with both GABA predominated (Casseday et al. 2000). Intracellular measurement of synaptic currents *in vivo*, using whole-cell patch clamp, shows an excitatory–inhibitory interplay consistent with duration tuning (Covey et al. 1996). Sound onset evokes a short latency sustained inhibition and long latency excitation. The end of the inhibition is followed by an intrinsic rebound excitation or other source of excitation associated with offset. However, the cell fires only when this rebound coincides with the long-latency excitation evoked by stimulus onset.

At other latencies the excitation is either suppressed by inhibition or occurs after the rebound excitation. The cell's preferred duration is thus determined by the sound latency evoked by the long latency excitation (Ehrlich et al. 1997; see Chapter 17).

Evidence for duration tuned neurons also comes from studies of chinchilla and mouse (Chen 1998; Brand et al. 2000). About half the mouse neurons showed long, short or band-pass functions for duration, but most varied with other stimulus properties. It is unknown whether the underlying mechanisms resemble those in bat, but intracellular current measurements in cat IC neurons found one essential component of the model, namely that some cells receive inhibition before excitation (Nelson and Erulkar 1963). It has been suggested that the mechanism for sensitivity to duration might underlie psychophysical masking. In the big brown bat the unit response to the best duration tone was suppressed when it preceded or followed a tone that evoked only inhibition. Thus, inhibition with a shorter latency than excitation may contribute to backward masking (Faure et al. 2003).

Bat IC neurons may be sensitive to the delay between the emission of a biosonar signal and the echo return (Feng et al. 1978; Mittmann and Wenstrup 1995). Their preferred delays (<10 ms) are within the range a bat would experience while echolocating. Mechanisms, like those proposed for periodicity analysis (see Section 3.7), involving time-dependent excitation and inhibition (Saitoh and Suga 1995), that elicit firing when coincidences occur between the emitted sound and an echo component, could explain how these detectors operate.

## 6. CONCLUSIONS AND DIRECTIONS FOR FUTURE RESEARCH

Hearing depends on the accurate perception of temporal information. From pitch perception to sound localization and speech discrimination, time and timing are critical in auditory perception. We have focussed on how temporal *changes* in amplitude, frequency and duration of sounds are encoded in the responses of IC neurons.

Several themes are noteworthy. First is the emergence of response properties that are not apparent below the IC. For AM coding, one theme is the appearance of modulation frequency selectivity signalled by average firing rate, besides synchronized firing to the modulation envelope at lower auditory stations. Although this implies that temporal information is recoded as a rate-based stream at the IC, we emphasize that many IC neurons have strongly synchronized responses to AM envelopes, sometimes well above 100 Hz, and many neurons do not show tuned responses to modulation frequency when transfer functions are derived from average rate measures. Indeed, for processing sounds with temporal information (as in source streaming and grouping) it might be important to retain

a temporal component in the response to identify different frequency components that share common temporal features.

Coding information by average rate may compensate for the reduction in frequency limit for synchronized firing to modulation envelopes for IC neurons compared to auditory nerve fibers and CN neurons. This trend, apparent in the periodic responses to both AM and FM, continues in the thalamus and the cortex. A likely extension of coding information by rate is that the coded parameter is represented topographically across the IC laminae. Evidence has been presented for such representations of AM rate and frequency sweep speed, and these may underlie the generation of specific auditory percepts. A particular pitch may originate when the appropriate frequency channels are stimulated with the appropriate envelope periodicity. Periodicity pitch and frequency information—defining sound timbre—are orthogonal percepts, a relationship in accord with the finding that periodotopy and tonotopy have complementary IC representations.

For both FM and AM there is evidence that the IC is the first site in the auditory pathway for neural selectivity to the direction of frequency or amplitude change, and some units may respond exclusively to frequency dynamics rather than to static frequencies. Similarly, duration tuning appears to first emerge at the IC. An extensive inhibitory component present in both the synaptic input to IC and in a significant proportion of its neurons plays an important role in the genesis, and perhaps the spatial representation, of these selectivities. Concomitant with the specialization and refinement in the temporal response properties of IC neurons, is the robust representation of nonlinear components in their responses. These have to date been described in general terms, such as the nonsinusoidal responses of many cells to sinusoidal modulations. A key aspect of future studies will be a more rigorous and quantitative analyses of these nonlinearities.

With respect to stimuli, most studies have used relatively simple sounds such as sinusoidal or linear modulations of pure tones often centered on the BF of the unit under study. Given the significant nonlinearities in IC responses, some features of temporal processing in the IC will likely become apparent only with the use of more complex sounds that contain multiple frequency components and combine more than one temporal property. This approach will be vital for addressing hypotheses about across-frequency interactions within the IC and the creation of auditory streams. A fundamental requirement for addressing many of the questions about temporal information processing in the IC is an understanding of how its circuits are organized, and how the membrane properties of neurons shape their responses. Auditory information coded in many nuclei and by numerous brain stem neuronal types converges in the IC. For some models, like the periodicity model of Langner, one constraint is that different inputs to the IC contact the same neurons and that their spatial and temporal coincidence is essential for auditory processing. On the other hand, other inputs may target specific domains within the IC to create local networks representing specific stimulus parameters. For example, neurons preferentially coding localization and

others coding periodicity may participate in such local networks and may be important for auditory streaming.

To resolve these conundrums, we need a clearer and more complete description of IC microcircuitry to specify the functional and structural relationships between its neurons, the different subdivisions in which they reside, and the synaptic inputs they receive. Some of this information is emerging from *in vitro* studies; complementary intracellular *in vivo* analyses, that enable the temporal characteristics of both excitatory and inhibitory postsynaptic currents to be measured in response to physiological stimuli, will be vital to this agenda.

## Abbreviations

AM	amplitude modulation
BMF	best modulation frequency
CF	characteristic frequency
CN	cochlear nucleus
CNIC	central nucleus of the inferior colliculus
DCN	dorsal cochlear nucleus
ENIC	external nucleus of the inferior colliculus
FM	frequency modulation
GABA	$\gamma$ -aminobutyric acid
IC	inferior colliculus
MTF	modulation transfer function
PVCN	posteroventral cochlear nucleus
SAM	sinusoidal amplitude modulation
SFM	sinusoidal frequency modulation
VCN	ventral cochlear nucleus

## References

- Adams JC (1997) Projections from octopus cells of the posteroventral cochlear nucleus to the ventral nucleus of the lateral lemniscus in cat and human. *Auditory Neuroscience* **3**:335–350.
- Barsz K, Benson PK, and Walton JP (1998) Gap encoding by inferior collicular neurons is altered by minimal changes in signal envelope. *Hearing Research* **115**:13–26.
- Bibikov NG and Gorodetskaya MM (1980) Single unit responses in the auditory center of the frog mesencephalon to amplitude modulated tones. *Neirofiziologiya*. **12**:185–191.
- Bibikov NG and Nizamov SV (1996) Temporal coding of low-frequency amplitude modulation in the torus semicircularis of the grass frog. *Hearing Research* **101**:23–44.
- Biebel UW and Langner G (2002) Evidence for interactions across frequency channels in the inferior colliculus of awake chinchilla. *Hearing Research* **169**:151–168.
- Biebel UW, Tomlinson RWW, Bibikov N, and Langner G (1998) Responses to low modulation depth tones in single units of inferior colliculus in the alert chinchilla. In:

- Elsner N and Wehner R (eds). *Proceedings of the 26th Göttingen Neurobiology Conference*, Volume II. Thieme, Stuttgart, pp. 344.
- Borst M, Palm G, and Langner G (2004) A biologically motivated neural network for phase extraction from complex sounds. *Biological Cybernetics* **90**: 98–104.
- Brand A, Urban A, and Grothe B (2000) Duration tuning in the mouse auditory midbrain. *Journal of Neurophysiology* **84**:1790–1799.
- Bregman AS (1990) *Auditory Scene Analysis: The Perceptual Organization of Sound*. Bradford Books, Cambridge.
- Brugge JF, Blatchley B, and Kudoh M (1993) Encoding of amplitude-modulated tones by neurons of the inferior colliculus of the kitten. *Brain Research* **615**:199–217.
- Büttner C, Braun S, Jähn-Siebert TK, and Langner G (1997) Periodotopic organization in the inferior colliculus of chinchillas revealed by c-fos labelling due to electrical and acoustical stimulation. In: Elsner E and Wässle H (eds). *Proceedings of the 25th Göttingen Neurobiology Conference*, Volume II. Thieme, Stuttgart, pp. 369.
- Casseday JH, Ehrlich D, and Covey E (1994) Neural tuning for sound duration—role of inhibitory mechanisms in the inferior colliculus. *Science* **264**:847–850.
- Casseday JH, Covey E, and Grothe B (1997) Neural selectivity and tuning for sinusoidal frequency modulations in the inferior colliculus of the big brown bat, *Eptesicus fuscus*. *Journal of Neurophysiology* **77**:1595–1605.
- Casseday JH, Ehrlich D, and Covey E (2000) Neural measurement of sound duration: control by excitatory–inhibitory interactions in the inferior colliculus. *Journal of Neurophysiology* **84**:1475–1487.
- Chen GD (1998) Effects of stimulus duration on responses of neurons in the chinchilla inferior colliculus. *Hearing Research* **122**:142–150.
- Cherry EC and Sayers BMcA (1956) Human ‘cross-correlator’—A technique for measuring certain parameters of speech perception. *Journal of the Acoustical Society of America* **28**:889–895.
- Chung DY and Colavita FB (1976) Periodicity pitch perception and its upper frequency limit in cats. *Perception and Psychophysics* **20**:433–437.
- Clopton BM and Winfield JA (1974) Unit responses in the inferior colliculus of rat to temporal auditory patterns of tone sweeps and noise bursts. *Experimental Neurology* **42**:532–540.
- Condon CJ, White KR, and Feng AS (1996) Neurons with different temporal firing patterns in the inferior colliculus of the little brown bat differentially process sinusoidal amplitude-modulated signals. *Journal of Comparative Physiology A* **178**:147–157.
- Covey E and Casseday JH (1999) Timing in the auditory system of the bat. *Annual Review of Physiology* **61**:457–476.
- Covey E, Kauer JA, and Casseday JH (1996) Whole-cell patch-clamp recording reveals subthreshold sound-evoked postsynaptic currents in the inferior colliculus of awake bats. *Journal of Neuroscience* **16**:3009–3018.
- Cynx J and Shapiro M (1986) Perception of missing fundamental by a species of songbird (*Sturnus vulgaris*). *Journal of Comparative Psychology* **100**:356–360.
- Darwin CJ and Carlyon RP (1995) Auditory grouping. In: Moore BCJ (ed). *The Handbook of Perception and Cognition*, Volume 6: *Hearing*. Academic Press, San Diego, pp. 387–424.
- de Cheveigné A (1993) Separation of concurrent harmonic sounds: fundamental frequency estimation and a time-domain cancellation model of auditory processing. *Journal of the Acoustical Society of America* **93**:3271–3290.

- Delgutte B, Hammond BM, and Cariani PA (1998) Neural coding of the temporal envelope of speech: relation to modulation transfer functions. In: Palmer AR, Rees A, Summerfield AQ, and Meddis R (eds). *Psychophysical and Physiological Advances in Hearing*. Whurr, London, pp. 595–603.
- Ehret G and Moffat AJM (1985) Inferior colliculus of the house mouse. 2. Single unit responses to tones, noise and tone-noise combinations as a function of sound intensity. *Journal of Comparative Physiology A* **156**:619–635.
- Ehrlich D, Casseday JH, and Covey E (1997) Neural tuning to sound duration in the inferior colliculus of the big brown bat, *Eptesicus fuscus*. *Journal of Neurophysiology* **77**:2360–2372.
- Epping WJM and Eggermont JJ (1986) Sensitivity of neurons in the auditory midbrain of the grassfrog to temporal characteristics of sound. II. Stimulation with amplitude modulated sounds. *Hearing Research* **24**:55–72.
- Eruikar SD, Butler RA, and Gerstein GL (1968) Excitation and inhibition in the cochlear nucleus. II. Frequency modulated tones. *Journal of Neurophysiology* **31**:537–548.
- Faingold CL, Gehlbach G, and Caspary DM (1989) On the role of GABA as an inhibitory neurotransmitter in inferior colliculus neurons—iontophoretic studies. *Brain Research* **500**:302–312.
- Faure PA, Fremouw T, Casseday JH, and Covey E (2003) Temporal masking reveals properties of sound-evoked inhibition in duration-tuned neurons of the inferior colliculus. *Journal of Neuroscience* **23**:3052–3065.
- Felsheim C and Ostwald J (1996) Responses to exponential frequency modulations in the rat inferior colliculus. *Hearing Research* **98**:137–151.
- Feng AS, Simmons JA, and Kick SA (1978) Echo detection and target-ranging neurons in the auditory system of the bat *Eptesicus fuscus*. *Science* **202**:645–648.
- Fuzessery ZM (1994) Response selectivity for multiple dimensions of frequency sweeps in the pallid bat inferior colliculus. *Journal of Neurophysiology* **72**:1061–1079.
- Fuzessery ZM and Hall JC (1996) Role of GABA in shaping frequency tuning and creating FM sweep selectivity in the inferior colliculus. *Journal of Neurophysiology* **76**:1059–1073.
- Fuzessery ZM and Hall JC (1999) Sound duration selectivity in the pallid bat inferior colliculus. *Hearing Research* **137**:137–154.
- Gleich O, Hamann I, Klump GM, Kittel M, and Strutz J (2003) Boosting GABA improves impaired auditory temporal resolution in the gerbil. *NeuroReport* **14**:1877–1880.
- Goldberg JM and Brown PB (1969) Response of binaural neurons of dog superior olivary complex to dichotic tonal stimuli: some physiological mechanisms of sound localization. *Journal of Neurophysiology* **32**:613–636.
- Gordon M and O'Neill WE (1998) Temporal processing across frequency channels by FM selective auditory neurons can account for FM rate selectivity. *Hearing Research* **122**:97–108.
- Gordon M and O'Neill WE (2000) An extralemniscal component of the mustached bat inferior colliculus selective for direction and rate of linear frequency modulations. *Journal of Comparative Neurology* **426**:165–181.
- Greenberg S (1988) Acoustic transduction in the auditory periphery. *Journal of Phonetics* **16**:3–17.
- Hage SR and Ehret G (2003) Mapping responses to frequency sweeps and tones in the inferior colliculus of house mice. *European Journal of Neuroscience* **18**:2301–2312.
- Hall JW, Haggard MP, and Fernandes MA (1984) Detection in noise by spectro-temporal pattern analysis. *Journal of the Acoustical Society of America* **76**:50–56.

- Hattori T and Suga N (1997) The inferior colliculus of the mustached bat has the frequency-vs-latency coordinates. *Journal of Comparative Physiology A* **180**:271–284.
- Heffner HE and Whitfield IC (1976) Perception of the missing fundamental by cats. *Journal of the Acoustical Society of America* **59**:915–919.
- Heil P, Schulze H, and Langner G (1995) Ontogenetic development of periodicity coding in the inferior colliculus of the Mongolian gerbil. *Auditory Neuroscience* **1**:363–383.
- Hewitt MJ and Meddis R (1994) A computer model of amplitude-modulation sensitivity of single units in the inferior colliculus. *Journal of the Acoustical Society of America* **95**:2145–2159.
- Hillery CM (1984) Detection of amplitude-modulation tones by frogs: implications for temporal processing mechanisms. *Hearing Research* **14**:129–143.
- Hose B, Langner G, and Scheich H (1987) Topographic representation of periodicities in the forebrain of the mynah bird: one map for pitch and rhythm? *Brain Research* **422**:367–373.
- Huffman RF, Argeles PC, and Covey E (1998) Processing of sinusoidally frequency modulated signals in the nuclei of the lateral lemniscus of the big brown bat, *Eptesicus fuscus*. *Hearing Research* **126**:161–180.
- Hurley LM and Pollak GD (1999) Serotonin differentially modulates responses to tones and frequency-modulated sweeps in the inferior colliculus. *Journal of Neuroscience* **19**:8071–8082.
- Joris PX, Schreiner CE, and Rees A (2004) Neural processing of amplitude-modulated sounds. *Physiological Reviews* **84**:541–577.
- Kim DO, Sirianni JG, and Chang SO (1990) Responses of DCN-PVCN neurons and auditory nerve fibers in unanesthetized decerebrate cats to AM and pure tones: analysis with autocorrelation/power-spectrum. *Hearing Research* **45**:95–113.
- Koch U and Grothe B (1998) GABAergic and glycinergic inhibition sharpens tuning for frequency modulations in the inferior colliculus of the big brown bat. *Journal of Neurophysiology* **80**:71–82.
- Krishna BS and Semple MN (2000) Auditory temporal processing: responses to sinusoidally amplitude-modulated tones in the inferior colliculus. *Journal of Neurophysiology* **84**:255–273.
- Kuwada S, Batra R, Yin TCT, Oliver DL, Haberly LB, and Stanford TR (1997) Intracellular recordings in response to monaural and binaural stimulation of neurons in the inferior colliculus of the cat. *Journal of Neuroscience* **17**:7565–7581.
- Langner G (1981) Neuronal mechanisms for pitch analysis in the time domain. *Experimental Brain Research* **44**:450–454.
- Langner G (1983) Evidence for neuronal periodicity detection in the auditory system of the guinea fowl: implications for pitch analysis in the time domain. *Experimental Brain Research* **52**:333–355.
- Langner G (1988) Physiological properties of units in the cochlear nucleus are adequate for a model of periodicity analysis in the auditory midbrain. In: Syka J and Masterton RB (eds). *Auditory Pathway: Structure and Function*. Plenum Press, New York, pp. 207–212.
- Langner G (1992) Periodicity coding in the auditory system. *Hearing Research* **60**:115–142.
- Langner G (1997) Neural processing and representation of periodicity pitch. *Acta Otolaryngologica Supplement* **532**:68–76.
- Langner G (2004) Topographic representation of periodicity information: the 2nd neural axis of the auditory system. In: Syka J and Merzenich MM (eds). *Plasticity of the*

- Central Auditory System and Processing of Complex Acoustic Signals*. Springer-Verlag, New York, pp. 19–33.
- Langner G and Schreiner CE (1988) Periodicity coding in the inferior colliculus of the cat. I. Neuronal mechanisms. *Journal of Neurophysiology* **60**:1799–1822.
- Langner G and Schreiner CE (1996) Contributions of the auditory brainstem to periodicity pitch coding. *Advances in Speech, Hearing and Language Processing* **3**:447–461.
- Langner G, Schreiner CE, and Merzenich MM (1987) Covariation of latency and temporal resolution in the inferior colliculus of the cat. *Hearing Research* **31**:197–202.
- Langner G, Schreiner CE, and Albert M (1992) Tonotopy and periodotopy in the auditory midbrain of cat and guinea fowl. In: Cazals Y, Horner K, and Demany L (eds). *Auditory Physiology and Perception*. Pergamon Press, New York, pp. 241–248.
- Langner G, Sams M, Heil P, and Schulze H (1997) Frequency and periodicity are represented in orthogonal maps in the human auditory cortex: evidence from magnetoencephalography. *Journal of Comparative Physiology, A* **181**:665–676.
- Langner G, Albert M, and Briede T (2002) Temporal and spatial coding of periodicity information in the inferior colliculus of awake chinchilla (*Chinchilla laniger*). *Hearing Research* **168**:110–130.
- Large EW and Crawford JD (2002) Auditory temporal computation: interval selectivity based on post-inhibitory rebound. *Journal of Computational Neuroscience* **13**:125–142.
- Le Beau FEN, Rees A, and Malmierca MS (1996) Contribution of GABA- and glycine-mediated inhibition to the monaural temporal response properties of neurons in the inferior colliculus. *Journal of Neurophysiology* **75**:902–919.
- Lesser HD, Frisina RD, and O'Neill WE (1986) Responses to amplitude-modulated sounds in the inferior colliculus of the mustached bat. *Society for Neuroscience Abstracts* **12**:345.
- Lesser HD, O'Neill WE, Frisina RD, and Emerson RG (1990) ON-OFF units in the mustached bat inferior colliculus are selective for transients resembling acoustic glint from fluttering insect targets. *Experimental Brain Research* **82**:137–148.
- Licklider JCR (1951) A duplex theory of pitch perception. *Experientia* **7**:128–134.
- Misawa H and Suga N (2001) Multiple combination-sensitive neurons in the auditory cortex of the mustached bat. *Hearing Research* **151**:15–29.
- Mittmann DH and Wenstrup JJ (1995) Combination-sensitive neurons in the inferior colliculus. *Hearing Research* **90**:185–191.
- Møller AR (1974) Coding of amplitude and frequency modulated sounds in the cochlear nucleus. *Acustica* **31**:292–299.
- Møller AR and Rees A (1986) Dynamic properties of the responses of single neurons in the inferior colliculus of the rat. *Hearing Research* **24**:203–215.
- Moore BCJ (2003) *An Introduction to the Psychology of Hearing*. Academic Press, San Diego.
- Müller-Preuss P, Flachskamm C, and Bieser A (1994) Neural encoding of amplitude modulation within the auditory midbrain of squirrel monkeys. *Hearing Research* **80**:197–208.
- Nelson PG and Erulkar SD (1963) Synaptic mechanisms of excitation and inhibition in the central auditory pathway. *Journal of Neurophysiology* **26**:908–923.
- Nelson PG, Erulkar SD, and Bryan JS (1966) Response of units in the inferior colliculus to time-varying acoustic stimuli. *Journal of Neurophysiology* **29**:834–860.
- Neuert V, Pressnitzer D, Patterson RD, and Winter IM (2001) The responses of single units in the inferior colliculus of the guinea pig to damped and ramped sinusoids. *Hearing Research* **159**:36–52.



- Palombi PS, Backoff PM, and Caspary DM (2001) Responses of young and aged rat inferior colliculus neurons to sinusoidally amplitude modulated stimuli. *Hearing Research* **153**:174–180.
- Pedemonte M, Torterolo P, and Velluti RA (1997) In vivo intracellular characteristics of inferior colliculus neurons in guinea pigs. *Brain Research* **759**:24–31.
- Pinheiro AD, Jen PH-S, and Wu M (1991) Encoding repetition rate and duration in the inferior colliculus of the big brown bat, *Eptesicus fuscus*. *Journal of Comparative Physiology A* **169**:69–85.
- Plomp R (1983) The role of modulation in hearing. In: Klinke R and Hartmann R (eds). *Hearing—Physiological Bases and Psychophysics*. Springer-Verlag, Berlin, pp. 270–276.
- Poon PWF and Chiu TW (1997) Single cell responses to AM tones of different envelopes at the auditory midbrain. In: Syka J (ed). *Acoustical Signal Processing in the Central Auditory System*. Plenum Press, New York, pp. 253–261.
- Poon PWF, Chen XY, and Cheung YM (1992) Differences in FM response correlate with morphology of neurons in the rat inferior colliculus. *Experimental Brain Research* **91**: 94–104.
- Portfors CV and Wenstrup JJ (2001) Topographical distribution of delay-tuned responses in the mustached bat inferior colliculus. *Hearing Research* **151**:95–105.
- Pressnitzer D, Winter IM, and Patterson RD (2000) The responses of single units in the ventral cochlear nucleus of the guinea pig to damped and ramped sinusoids. *Hearing Research* **149**:155–166.
- Rall W (1962) Theory of physiological properties of dendrites. *Annals of the New York Academy of Sciences* **96**:1071–1092.
- Rees A and Møller AR (1983) Responses of neurons in the inferior colliculus of the rat to AM and FM tones. *Hearing Research* **10**:301–330.
- Rees A and Møller AR (1987) Stimulus properties influencing the responses of inferior colliculus neurons to amplitude-modulated sounds. *Hearing Research* **27**:129–143.
- Rees A and Palmer AR (1988) Rate-intensity functions and their modification by broad-band noise for neurons in the guinea-pig inferior colliculus. *Journal of the Acoustical Society of America* **83**:1488–1498.
- Rees A and Palmer AR (1989) Neuronal responses to amplitude-modulated and pure-tone stimuli in the guinea-pig inferior colliculus, and their modification by broad-band noise. *Journal of the Acoustical Society of America* **85**:1978–1994.
- Rees A and Sarbaz A (1997) The influence of intrinsic oscillations on the encoding of amplitude modulation by neurons in the inferior colliculus. In: Syka J (ed). *Acoustical Signal Processing in the Central Auditory System*, Plenum Press, New York, pp. 239–252.
- Rees A, Sarbaz A, Malmierca MS, and Le Beau FEN (1997) Regularity of firing of neurons in the inferior colliculus. *Journal of Neurophysiology* **77**:2945–2965.
- Reimer K (1987) Coding of sinusoidally amplitude modulated acoustic stimuli in the inferior colliculus of the rufous horseshoe bat, *Rhinolophus rouxi*. *Journal of Comparative Physiology A* **161**:305–313.
- Riquelme R, Saldaña E, Osen KK, Ottersen OP, and Merchán MA (2001) Colocalization of GABA and glycine in the ventral nucleus of the lateral lemniscus in rat: an in situ hybridization and semiquantitative immunocytochemical study. *Journal of Comparative Neurology* **432**:409–424.
- Rockel AJ and Jones EG (1973) The neuronal organization of the inferior colliculus of the adult cat. I. The central nucleus. *Journal Comparative Neurology* **147**:11–60.

- Rose GJ and Capranica RR (1984) Processing of amplitude-modulated sounds by the auditory midbrain of two species of toads: matched temporal filters. *Journal of Comparative Physiology A* **154**:211–219.
- Rose GJ and Capranica RR (1985) Sensitivity to amplitude modulated sounds in the anuran auditory nervous system. *Journal of Neurophysiology* **53**:446–465.
- Rose JE, Greenwood DD, Goldberg JM, and Hind JE (1963) Some discharge characteristics of single neurons in the inferior colliculus of the cat. I. Tonotopical organization, relation of spike-counts to tone intensity, and firing patterns of single elements. *Journal of Neurophysiology* **26**:294–320.
- Rosen S (1992) Temporal information in speech: acoustic, auditory and linguistic aspects. *Philosophical Transactions of the Royal Society of London, series B, Biological Sciences* **336**:367–373.
- Ryan A and Miller J (1978) Single unit responses in the inferior colliculus of the awake and performing rhesus monkey. *Experimental Brain Research* **32**:389–407.
- Saitoh I and Suga N (1995) Long delay lines for ranging are created by inhibition in the inferior colliculus of the mustached bat. *Journal of Neurophysiology* **74**:1–11.
- Schouten JF (1940) The perception of pitch. *Philips Technical Review* **5**:286–294.
- Schouten JF (1970) The residue revisited. In: Plomp R and Smoorenburg GF (eds). *Frequency Analysis and Periodicity Detection in Hearing*. Sijthoff, Leiden, pp. 41–54.
- Schouten JF, Ritsma RJ, and Cardozo BL (1962) Pitch of the residue. *Journal of the Acoustical Society of America* **34**:1418–1424.
- Schreiner CE and Langner G (1988) Periodicity coding in the inferior colliculus of the cat. I. Topographical organization. *Journal of Neurophysiology* **60**:1823–1840.
- Schreiner CE and Langner G (1997) Laminar fine structure of frequency organization in auditory midbrain. *Nature* **388**:383–386.
- Schreiner CE and Snyder RL (1987) Modulation transfer characteristics of neurons in the dorsal cochlear nucleus of the cat. *Society for Neuroscience Abstracts* **13**:1258.
- Schroeder MR (1981) Modulation transfer functions: definition and measurement. *Acustica* **49**:179–182.
- Schuller G (1979) Coding of small sinusoidal frequency and amplitude modulations in the inferior colliculus of the “CF-FM” bat, *Rhinolophus ferrumequinum*. *Experimental Brain Research* **34**:117–132.
- Schuller G (1984) Natural ultrasonic echos from wing beating insects are encoded by collicular neurons in the CF-FM bat, *Rhinolophus ferrumequinum*. *Journal of Comparative Physiology A* **155**:121–128.
- Semal C and Demany L (1990) The upper limit of musical pitch. *Music Perception* **8**:165–175.
- Shannon RV, Zeng FG, Kamath V, Wygonski J, and Ekelid M (1995) Speech recognition with primarily temporal cues. *Science* **270**:303–304.
- Shofner WP (2000) Comparison of frequency discrimination thresholds for complex and single tones in chinchillas. *Hearing Research* **149**:106–114.
- Shofner WP and Yost WA (1997) Discrimination of rippled-spectrum noise from flat-spectrum noise by chinchillas: evidence for a spectral dominance region. *Hearing Research* **110**:15–24.
- Smith ZM, Delgutte B, and Oxenham AJ (2002) Chimaeric sounds reveal dichotomies in auditory perception. *Nature* **416**:87–90.
- Stiebler I and Ehret G (1985) Inferior colliculus of the house mouse. I. A quantitative study of tonotopic organization, frequency representation, and tone-threshold representation. *Journal of Comparative Neurology* **238**:65–76.

- Suga N (1965) Analysis of frequency-modulated sounds by auditory neurons of echolocating bats. *Journal of Physiology (London)* **179**:26–53.
- Suga N (1969) Classification of inferior collicular neurons of bats in terms of responses to pure tones, FM sounds and noise bursts. *Journal of Physiology(London)* **200**:555–574.
- Suga N and O'Neill WE (1979) Neural axis representing target range in the auditory cortex of the mustache bat. *Science* **206**:351–353.
- Syka J, Popelář J, Kvašňák E, and Astl J (2000) Response properties of neurons in the central nucleus and external and dorsal cortices of the inferior colliculus in guinea pig. *Experimental Brain Research* **133**:254–266.
- Terhardt E (1970) Frequency analysis and periodicity detection in the sensation of roughness and periodicity pitch. In: Plomp R and Smoorenburg GF (eds). *Frequency Analysis and Periodicity Detection in Hearing*. Sijthoff, Leiden, pp. 278–290.
- Thornton S and Rees A (2001) Across-frequency interactions in the inferior colliculus. *Association for Research in Otolaryngology Abstracts* **24**:54.
- Tomlinson RWW and Schwarz DWF (1988) Perception of the missing fundamental in nonhuman primates. *Journal of the Acoustical Society of America* **84**:560–565.
- Treurniet WC and Boucher DR (2001) A masking level difference due to harmonicity. *Journal of the Acoustical Society of America* **109**:306–320.
- Van Stokkum IHM (1987) Sensitivity of neurons in the dorsal medullary nucleus of the grassfrog to spectral and temporal characteristics of sound. *Hearing Research* **29**:223–235.
- Van Stokkum IHM (1989) A model for the auditory midbrain of the grassfrog for monaural stimuli. *Hearing Research* **43**:231–250.
- Vater M, Habbicht H, Kössl M, and Grothe B (1992) The functional role of GABA and glycine in monaural and binaural processing in the inferior colliculus of horseshoe bats. *Journal of Comparative Physiology A* **171**:541–553.
- Vater M, Covey E, and Casseday JH (1997) The columnar region of the ventral nucleus of the lateral lemniscus in the big brown bat (*Eptesicus fuscus*): synaptic arrangements and structural correlates of feedforward inhibitory function. *Cell and Tissue Research* **289**:223–233.
- Viemeister NF and Plack CJ (1993) Time analysis. In: Yost WA, Popper AN, and Fay RR (eds). *Springer Handbook of Auditory Research, Volume 3: Human Psychophysics*. Springer-Verlag, New York, pp. 116–154.
- Walkowiak W (1984) Neuronal correlates of the recognition of pulsed sound signals in the grass frog. *Journal of Comparative Physiology A* **155**:57–66.
- Walton JP, Frisina RD, Ison JR, and O'Neill WE (1997) Neural correlates of behavioral gap detection in the inferior colliculus of the young CBA mouse. *Journal of Comparative Physiology A* **181**:161–176.
- Walton JP, Frisina RD, and O'Neill WE (1998) Age-related alteration in processing of temporal sound features in the auditory midbrain of the CBA mouse. *Journal of Neuroscience* **18**:2764–2776.
- Wenstrup JJ and Grose CD (1995) Inputs to combination-sensitive neurons in the medial geniculate body of the mustached bat: the missing fundamental. *Journal of Neuroscience* **15**:4693–4711.
- Wenstrup JJ and Leroy SA (2001) Spectral integration in the inferior colliculus: role of glycinergic inhibition in response facilitation. *Journal of Neuroscience* **21** RC124:1–6.
- Wever EG (1949) *Theory of Hearing*. John Wiley & Sons, New York.

- Willott JF and Urban GP (1978) Response properties of neurons in nuclei of the mouse inferior colliculus. *Journal of Comparative Physiology A* **127**:175–184.
- Wilson WW and Walton JP (2002) Background noise improves gap detection in tonically inhibited inferior colliculus neurons. *Journal of Neurophysiology* **87**:240–249.
- Wright AA, Rivera JJ, Hulse SH, Shyan M, and Neiwirth JJ (2000) Music perception and octave generalization in rhesus monkeys. *Journal of Experimental Psychology* **129**: 291–307.
- Yan J and Suga N (1996) The midbrain creates and the thalamus sharpens echo-delay tuning for the cortical representation of target-distance information in the mustached bat. *Hearing Research* **93**:102–110.
- Yang LC and Pollak GD (1997) Differential response properties to amplitude modulated signals in the dorsal nucleus of the lateral lemniscus of the mustache bat and the roles of GABAergic inhibition. *Journal of Neurophysiology* **77**:324–340.
- Yost WA and Sheft S (1989) Across-critical-band processing of amplitude-modulated tones. *Journal of the Acoustical Society of America* **85**:848–857.
- Yost WA and Sheft S (1993) Auditory perception. In: Yost WA, Popper AN, and Fay RR (eds). *Springer Handbook of Auditory Research*, Volume 3: *Human Psychophysics*. Springer-Verlag, New York, pp. 193–236.

# Chapter 13

## Binaural and Spatial Coding in the Inferior Colliculus

ALAN R. PALMER AND SHIGEYUKI KUWADA

### 1. INTRODUCTION

#### *1.1. THE IMPORTANCE OF THE INFERIOR COLLICULUS IN BINAURAL PROCESSING*

The inferior colliculus (IC) is a critical center for binaural processing because the outputs of the primary binaural processors, the medial and lateral superior olives (MSO and LSO), terminate in the IC. The MSO and LSO also project to the nuclei of the lateral lemniscus, which in turn project to the IC. Direct monaural inputs from both cochlear nuclei, as well as second-order monaural inputs from the ventral nucleus of the lateral lemniscus, can also converge with the binaural inputs in the IC. Thus, there is considerable opportunity for binaural as well as monaural convergence. Because of this potential for multiple-source interaction, the responses of IC neurons cannot be a simple reflection of a single source (see Chapter 2).

#### *1.2. BINAURAL CUES AND FUNCTIONS OF BINAURAL HEARING*

There are generally considered to be three major cues for the localization of sound: interaural time (ITDs) and level differences (ILDs) and spectral cues. Although often treated as independent, because in some circumstances only one cue is available, for any spectrally rich sound all of these cues are simultaneously present and consistent. Interaural time differences are created by the differences in arrival time of the sound at the two ears. The largest ITD depends on head size and in humans it is approximately 700  $\mu$ s (Nordlund 1962; Shaw 1974; Blauert 1983). The upper frequency limit for ITD cues is limited by head size, that is, when the wavelength of the sound is shorter than the head width. In humans, this limit is about 1300 Hz. Above this frequency the effect of the head in "shadowing" the sound reaching the ear further from the sound source creates ILDs as well as multiple peaks (ambiguity) in the ITD function. In low-frequency sounds ILDs are small because their wavelengths are much larger than the head width. However, even at low frequencies sound sources very near

the ear (in the near field) can generate considerable ILDs. In humans, the ILDs at high frequencies can exceed 20 dB (Shaw 1974; Blauert 1983). The utility of this cue is maximized in animals with small head widths by extending their high-frequency hearing (Masterton et al. 1969). Spatially dependent spectral cues are generated by interference at the eardrum between sound waves traveling directly from the source with those reflected from the torso and pinna. The effect of the interfering waves is to produce peaks and notches in the spectra of wide-band sounds, the frequency position of which depends on the spatial position (Blauert 1983). The filtering effect of the head and body in the sound field can be measured in the ear canal as the head-related transfer function (HRTF). For any sound source, the sound reaching the two ears will be filtered by the HRTF, appropriate for that position in space at that ear, to produce interaural spectra that contain ITDs, ILDs, and spectral information specific to the sound position.

Many advantages accrue from listening with both ears rather than only one ear, for example, improved detection and increased loudness of sounds. Other advantages stem directly from the ability to localize the sound source, such as the detection of moving sound sources. Whether this is a simple process of sequentially evaluating spatial position or involves specialized motion detectors is still in debate (Grantham 1997). Finally, there are benefits in source isolation and segregation. In our everyday environment we experience not only direct sound, but also many echoes that usually do not interfere with our listening to the primary source. This reduced perceptual salience of echoes depends on processing the differences at the two ears and identifying sources that arrive earliest (the precedence effect; Wallach et al. 1949). Binaural hearing also helps in segregating target sounds from interfering backgrounds that derive from spatially separated sources. A well investigated surrogate for this ability is the binaural masking level difference in which spatially separating masker sounds from target sounds improves target detection (Hirsh 1948a,b; Licklider 1948).

## 2. PROCESSING OF INTERAURAL TIME DIFFERENCES FOR LOW-FREQUENCY SOUNDS

### *2.1. TYPES OF INTERAURAL TIME DIFFERENCE SENSITIVITY AND THEIR POSSIBLE ORIGIN*

It is important to distinguish between two types of ITD, onset and ongoing. The onset ITD is the initial difference produced by the sound reaching the ears at different times, while the ongoing ITD is produced by ongoing differences in the pattern of pressure variations at the two ears. In humans, the ongoing ITDs are much more potent than onset ITDs in determining the position of a lateralized image (Tobias and Schubert 1959; Buell et al. 1991). Consistent with this observation, neurons sensitive to ongoing ITD (those showing a cyclic ITD function with a period equal to that of the stimulating tone; Fig. 13.1) generally display no sensitivity to onset ITDs (Kuwada and Yin 1983). Although the ev-




Figure 13.1. Types of ITD functions. Examples of types of response to changes in ITD across frequency in IC neurons sensitive to low-frequency sounds. *Left column*: each ITD function is the response to a particular stimulus frequency. *Middle column*: “composite curves” obtained by averaging the ITD functions (*left column*). The peak of the composite curve is fit by a parabola and the center of the parabola is an estimate of the neuron’s best ITD (BD). *Right column*: plots of the mean phase of the response vs. the stimulating frequency. The slope of the best fit line represents the characteristic delay (CD), and the intercept is the characteristic phase (CP). (A) Peak-type response: the CP is near 0, the CD occurs near the peak of each ITD function, and the BD of the composite curve resembles the CD. (B) Trough-type response: CP is near  $\pm 0.5$ , the CD occurs near the trough of each ITD function, and the BD differs substantially from the CD. (C) Intermediate-type response: the CP is near  $\pm 0.25$ , the CD occurs between the peaks and troughs of the ITD function, and the BD differs from the CD, but not as much as for the trough-type neuron. (From Kuwada et al. 1987.) *Contra*, Contralateral; *Ipsi*, ipsilateral.

idence for neural sensitivity to ongoing ITDs is abundant and has been observed at all levels of the auditory pathway (Kuwada et al. 1997), sensitivity to onset ITDs has been studied by few investigators (e.g., Harnischfeger 1980; Caird and Klinke 1983, 1987; Kelly and Phillips 1991). Such investigations are complicated by the fact that even transient stimuli can give rise to cyclic ITD functions (Carney and Yin 1989). This occurs because brief sounds can “ring” the basilar membrane and evoke a spike discharge that is synchronized to the fiber’s best frequency (Kiang et al. 1965; Carney and Yin 1989).

Rose and colleagues (1966) described three types of ongoing ITD-sensitive neurons in the IC that are currently referred to as peak-, trough-, and intermediate-type. Quantitative criteria were established for these three types (Yin and Kuwada 1983; Fig. 13.1) that require the neuron's sensitivity to ITDs be measured at several frequencies. For peak-type neurons, the peaks (maxima) of the ITD functions for different stimulus frequencies align at or near a particular ITD (Fig. 13.1, top row), for trough-type neurons the alignment is at or near the trough (minima; Fig. 13.1, middle row), and for intermediate-type neurons the alignment is intermediate (Fig. 13.1, bottom row). The alignment of ITD functions is determined by a linear fit to the plot of the mean interaural phase vs. stimulating frequency (Fig. 13.1, right column). The slope of the fit is the characteristic delay (CD) and the phase intercept at 0 Hz is the characteristic phase (CP). In the Jeffress scheme (see Section 2.2), the CD would correspond to the difference in conduction times of the neural path lengths from each ear to the binaural coincidence detector, that is, the ITD required for coincident arrival of the inputs. The CP is a measure of whether the alignment occurred at the peak (CPs near 0 or 1 cycle), trough (CPs near 0.5 cycles), or at an intermediate position (CP near 0.25 or 0.75 cycles). The most common ITD functions display a peak with flanking slopes (Figs. 13.1A and 13.2D), and can be evoked by tones or more complex stimuli such as amplitude-modulated tones, noise, clicks, and vowels (Kuwada and Yin 1983; Yin et al. 1986a; Carney and Yin 1989; Palmer et al. 1990; D'Angelo et al. 2003; Sterbing et al. 2003a).

Averaging the ITD functions (Fig. 13.1, left column) for all the stimulating frequencies to which the neuron was ITD sensitive generates the composite curve, which is representative of the neuron's ITD sensitivity to a broad-band sound (Yin et al. 1986a). The neuron's best ITD (BD) is estimated by fitting a parabola to the peak of the composite curve (Fig. 13.1, middle column). A neuron's BD can be predicted by the equation: best ITD = CD + CP/best frequency of ITD sensitivity (Fitzpatrick et al. 2000, 2002).

In a Jeffress type scheme, peak-type ITD sensitivity arises from excitatory inputs from each side, whereas trough-type sensitivity is created by excitatory inputs from one side and inhibitory inputs from the other (Fig. 13.2). Thus, it is tempting to assume that peak-type neurons in the IC reflect inputs from the MSO, because MSO neurons receive excitatory input from the cochlear nucleus neurons of each side. Similarly, trough-type IC neurons could reflect LSO inputs because LSO neurons receive excitatory inputs from the ipsilateral cochlear nucleus, and inhibitory inputs from the contralateral cochlear nucleus via the medial nucleus of the trapezoid body. Intermediate-type neurons are not compatible with a Jeffress type scheme and yet constitute a sizeable portion of ITD-sensitive IC neurons. Their phase-vs.-frequency plots are often complex and often of the intermediate type. There is compelling empirical and modeling evidence (McAlpine et al. 1998; Shackleton et al. 2000) that intermediate-type IC neurons are created by the convergence of inputs from different coincidence detectors from the same nucleus (e.g., MSO) or different nuclei (MSO and LSO). Because of the high likelihood that IC neurons receive ITD-sensitive inputs from many






Figure 13.2. Jeffress model for peak ITD sensitivity and its variant for trough units. (A) The classic Jeffress network has excitatory inputs arranged in a delay line and that feed coincidence detectors (circles labeled A–E) to convert interaural time differences to a place code for position (output axons labeled a–e). Activation is shown for a spatial position on the right as in (B). (C) A putative Jeffress network with inhibition from one side giving rise to suppression when the inputs are coincident (labels as in A). (D) A typical “peak” unit response from the Jeffress network. (E) A typical “trough” response from the variant of the Jeffress network.

different nuclei, the type of ITD sensitivity observed is not easily attributable to a single input, even for peak- and trough-type neurons.

It has been suggested on the basis of frequency tuning and ITD and ILD sensitivity that IC neuronal responses are dominated by parallel inputs from different lower level brain stem nuclei (Davis et al. 1999, 2003; Ramachandran et al. 1999; Ramachandran and May 2002). Thus, cells with V-shaped responses areas (“type V”) tend to be found at low frequencies, have excitatory input from both ears, and have “peak” type ITD functions. The dominant input to these cells is hypothesized to be the MSO. Cells with level tolerant narrow tuning (“type I”) tend to have trough- or intermediate-type carrier and envelope ITD functions, receive excitatory input from the contralateral ear and inhibitory input from the ipsilateral ear, and are sensitive to ILDs. These are hypothesized to be dominated by inputs from the LSO. Finally, units with an island of low-level excitation caused by strong best frequency inhibition at higher levels (“type O”)

are a more diverse group that often show very weak binaural sensitivities, but could show ITD or ILD sensitivity and were hypothesized to be dominated by input from the dorsal cochlear nucleus.

## 2.2. *THE JEFFRESS MODEL FOR INTERAURAL TIME DELAY SENSITIVITY*

The human ear is exquisitely sensitive to ITDs. For noise and clicks the just noticeable difference in ITD is 5 to 12  $\mu$ s and for tones it is 20 to 30  $\mu$ s, dependent on frequency and duration (Tobias and Zerlin 1959; Durlach 1972). An obvious question is, What aspects of the neural ITD function carry the salient information to code the ITD in an external sound source? The dominant model for the way in which ITDs are processed in the brain was proposed by Jeffress (1948) (Fig. 13.2A), who postulated a network of neurons that acted as coincidence detectors, receiving excitatory input from the two ears. Originally he suggested the medial geniculate body (Jeffress 1948), but later correctly suggested that it likely took place in the MSO (Jeffress 1958). Input to the coincidence detector from both sides occurring within a short time window leads to summation of the excitatory postsynaptic potentials and generation of an action potential. A variant of this scheme (Fig. 13.2C) has inhibitory input from one ear. When excitation and inhibition are coincident the cell firing is prevented. With this variant (Fig. 13.2C), responses of ITD-sensitive neurons are consistent with Jeffress's notion of coincidence detectors. To enable such cells to respond maximally (or minimally, Fig. 13.2C) to specific spatial locations, Jeffress proposed that the cells were connected to each ear by axons of differing length to provide a delay line. Only when the internal delay compensated exactly for the later arrival of the sound at one ear would a specific coincidence detector fire. Thus, the specific coincidence detector activated provides a code for a specific spatial position and the array of coincidence detectors converts ITDs into a place code representing all azimuthal sound locations.

A sound source would optimally activate neurons with ITD functions whose peaks correspond to the ITD created by the sound source, as well as other neurons whose ITD functions overlapped with the optimally activated functions, all contributing to increasing the activity at a locus in the neural ITD array. This viewpoint is supported in a study in the unanesthetized rabbit showing that such arrays in the SOC, IC, and auditory thalamus could achieve high acuity in ITD discriminability (Fitzpatrick et al. 1997).

## 2.3. *DISTRIBUTION OF BEST INTERAURAL TIME DELAY SENSITIVITIES*

Subsequent work focussed on the value of the ITD evoking the maximum output of the coincidence detectors. Whether in response to best frequency tones or to broadband noise, the numbers of IC neurons with the various best delays (BD) tended to peak at relatively long ITDs, representing sensitivity preferentially to

sounds in the contralateral hemifield, and this applies when different animals were examined (Fig. 13.3A, B; Palmer et al. 1990; McAlpine et al. 1996b). Later studies suggested that within such plots, at least for the gerbil and guinea pig, there was a frequency dependence: BDs near 0 ITD were found in neurons with high best frequencies, while longer BDs were found in neurons with lower best frequencies (Fig. 13.3C). This was confirmed when CD to tones (Fig. 13.3C; McAlpine et al. 1996b) or BD in response to noise (McAlpine et al. 2001) was measured as a function of best frequency.

Despite large differences in head width and therefore the physiological ITDs available to the animal, the shapes of the ITD functions and the distribution of their BDs are similar across many mammals. Furthermore, even the sharpest ITD function can span several hundred microseconds and the bulk of their BDs are off the midline. So, for animals with small head widths, an external sound source would not activate the peak of the ITD functions, and even if the peak was located near 0 ITD, the modulation of the response would be negligible across the physiological ITDs available to such animals.

In guinea pigs, even at high best frequencies, BDs were on the edge of the estimated physiological range, whereas the medial slopes of the ITD functions were well within. Their physiological range was estimated to be  $\pm 150 \mu\text{s}$ , al-




Figure 13.3. Distributions of best delays from different animals and the frequency dependence of BD. (A) The distribution of BD to BF tone and noise stimulation. (From Palmer et al. 1990.) (B) The distribution of noise BDs in the cat and guinea pig. (From Yin et al. 1986a; Palmer et al. 1990.) (C) The variation of BD and CD for BF tones with BF. *Open circles* show BD, *crosses* show CD. (From McAlpine et al. 1996b.) (D) The distribution of BDs and steepest slopes of noise ITD functions. (From McAlpine et al. 2001.) (E) The distribution of BDs as a function of stimulus frequency in rabbit.

though another report measured it to be about  $\pm 330 \mu\text{s}$  at frequencies  $< 1600$  Hz (Sterbing et al. 2003b). Taking this larger range, guinea pig neurons with best frequencies  $> 400$  Hz have their BDs within the physiological range (McAlpine et al. 2001), albeit near its extreme end.

A consequence of the frequency dependence of ITD sensitivity is that the steepest slopes of the delay functions at any frequency pass through the physiological range of the animal (Fig. 13.3D). Several other reports, however, show BD distributions (from composite curves) that, although still displaced toward the contralateral hemifield, have modes between 90 and 200  $\mu\text{s}$  (cat: Yin and Kuwada 1983, 1984; rabbit: Kuwada et al. 1987; Stanford et al. 1992) and that contain many neurons with BDs at and near 0 ITD. A sampling bias or perhaps even species difference may contribute to the differences between the cat/rabbit BDs and those of the guinea pig. The best frequency dependence of the BD is marked when frequencies  $< 300$  Hz are plotted, which are more prevalent in the guinea pig than in the cat and rabbit. Above 300 Hz, the data sets are much more similar with BDs closer to 0 ITD. For example, a plot of BD vs. dominant frequency (highly correlated to best frequency) (Joris et al. 2005) resembles Fig. 13.3C but extends only down to about  $\pm 300$  Hz and has a greater representation of neurons with BDs near 0 ITD. Furthermore, when BDs distributions are viewed at individual stimulus frequencies many neurons have BDs near 0 delay at all frequencies (Fig. 13.3E). Differences between studies may be methodological, and, if so, may shed light on ITD processing under normal listening conditions. The data in Fig. 13.3E are plotted as a function of stimulus frequency, which usually was not the best frequency of the neuron. In Jeffress' original network formulation this would make no difference (i.e., all BDs are available at all best frequencies). However, if the BD dependence on best frequency is general, the distribution of best frequencies will be germane, as a preponderance of high best frequencies will lead to BD distributions closer to 0 ITD. Certainly, some rabbit near-0 BDs could be from the low-frequency tails of high best-frequency neurons. Thus, the plot of BD at each stimulating frequency would encompass the distribution across best frequency for all neurons sampled that responded to that frequency. The importance of this (developed later) is that, under normal sound stimulation levels, the population of neurons will be activated at their tail region or at their best frequency. Under these conditions, depending on the best frequency distribution, the population response may indeed be dominated by neurons with BDs near 0 ITD.

From a Jeffress perspective, the full range of ITD tuning should be present in every frequency channel, but this was not the case in the guinea pig. However, McAlpine et al. (2001) used a low stimulus level (10 to 20 dB above the neurons' threshold) to reveal the shifts in BD with frequency. Even at moderate levels (60 to 70 dB), low-frequency ITD functions can be generated in high-frequency neurons, presumably by stimulating their low-frequency tails. In the rabbit, ITD functions evoked at 70 dB SPL show tuning to a range of BDs within this animal's physiological range at all frequencies (Fig. 13.3E). So, using near-threshold levels may not capture the population of ITD-sensitive neurons evoked at higher stimulus levels. However, the high-frequency neurons recruited at su-

prathreshold levels likely have short BDs (near 0), but their ITD tuning width will largely be determined by the stimulus period (Fitzpatrick and Kuwada 2001). Thus, although the population at high levels will contain many neurons responding maximally near 0 to low-frequency stimuli, many of these will not provide much variation in discharge as ITDs change within the physiological range. However, even under these circumstances neurons with BDs remote from 0 ITD and, hence, with steep slopes through 0 ITD, should still provide good resolution.

A distribution of BDs that is densest off the midline becomes a problem for the Jeffress model if its centroid is far from 0 ITD and/or the ITD functions are extremely sharp. In mammals, neither is the case as shown by averaging the delay functions from many neurons and plotting them as well as their mirror images (Fig. 13.4A). Averaging the two population ITD functions reveals a central peak at 0 ITD and is a simple consequence of the medial tails of each side overlapping. If the peaks were widely separated and the ITD functions were extremely sharp, then this overlap would be minimal and no activity could be evoked at 0 ITD.

#### 2.4. ALTERNATIVES TO THE JEFFRESS MODEL

Stillman (1971) first proposed that the steeply sloping part of the ITD functions in the kangaroo rat might be an important feature because the peaks of the ITD functions occurred outside the natural range of ITDs in this species. Consistent with this suggestion, Skottun (1998) and Shackleton et al. (2003) showed empirically that the slope of the ITD function has sufficient information to discriminate changes in ITD with high acuity. The steepest part of the slopes, independent of best frequency, fell near 0 ITD (McAlpine et al. 2001) because of the best-frequency dependent shift in BD (see Section 3.2.5). This finding, the paucity of BDs near 0 in the guinea pig, and the finding that inhibition can systematically shift the peaks of ITD functions, questions the validity of the Jeffress delay line model and has led to reconsiderations of alternative coding schemes that compare activity in two neural populations (possibly on either side of the brain) evoked by a sound with a particular ITD (McAlpine et al. 2001). Such a view is like the count-comparison models (von Békésy 1960) and refined within an anatomical and physiological context (van Bergeijk 1962).

McAlpine et al. (2001) recognized that their count-comparison scheme, in its simplest form, is incompatible with lesion/localization studies. Unilateral lesions of the IC, brachium of the IC, and auditory cortex have shown that the deficits in localizing sounds are confined to the contralateral sound field and localization appears normal in the ipsilateral sound field (Jenkins and Masterton 1982). Certainly, hemisphere comparisons remove the inherent ambiguities generated by other factors, such as sound level, that change the neural activity. However, direct access in one hemisphere to a monaural level dependent signal could achieve the same disambiguation.

A further caveat in making use of changes on the slopes occurs when the peak of the ITD function lies within the physiological range. In the cat and




Figure 13.4. A count comparison scheme applied to ITD-sensitive IC neurons in the unanesthetized rabbit. (A) The ITD functions from 300 to 1500 Hz were averaged to depict the activity to a sound containing all these frequencies (i.e., noise). This averaged function (*black line*) was also flipped to depict the activity in the contralateral IC (*coarse dashed line*). The difference (count comparison) between the activity of the two sides at ipsilateral delays of 100, 200, and 300  $\mu\text{s}$  is expressed as a percentage. The two ITD functions were averaged to show that the resultant function (*fine dashed line*) had maximal activity near 0 ITD. If these peaks were further displaced from 0 ITD, such averaging would lead to a function with minimal activity at 0 ITD. (B, C) The ITD functions only at 500 Hz or 1300 Hz were averaged. In all cases, the peak of the ITD function is within the rabbits' physiological range. In such cases, a count comparison scheme provides an ambiguous code for ITD that degrades as frequency increases.

rabbit and likely in human, most IC neurons display BDs within their physiological ranges. As the ITD increases, activity along the upslope and the downslope of the ITD function can produce the same count comparison difference, at very different ITDs. This is a situation (Fig. 13.4) in which the ITD functions from the IC of the unanesthetized rabbit for all frequencies between 300 and 1500 Hz are averaged (Fig. 13.4A), or only those at 500 Hz (Fig. 13.4B) or 1300 Hz (Fig. 13.4C). The averaged ITD functions in one IC are flipped to display also the averaged ITD function in the opposite IC. The activity evoked

at 0-, 100-, 200-, and 300- $\mu$ s delays appears as a percentage difference between the activities in the two. An ITD of 100 and 200  $\mu$ s (Fig. 13.4A) increases neural activity by 16%, whereas the increase between ITDs at 200 and 300  $\mu$ s is only 1%. At 1300 Hz, the count-comparison scheme becomes more ambiguous, that is, activity increases by 7% between a 100- and 200- $\mu$ s ITD, but decreases by 11% between a 200- and 300- $\mu$ s ITD. For these animals, under such stimulation conditions, a slope/count comparison scheme leads to an ambiguous code.

Although the measured ITD functions can produce ITD discriminability (Fitzpatrick et al. 1997; Skottun 1998; Shackleton et al. 2003) that matches or approaches that seen in humans, under similar behavioral testing conditions rabbits, gerbils, kangaroo rats, and guinea pigs show poorer localization ability than humans and the cat (Heffner and Heffner 1985, 1987, 1988a,b; Heffner 1997). This brings into question the processing capabilities of different animals, our inability to assess properly their ITD acuity, or our focus on neural codes that do not correlate with their behavioral acuity.

The controversy over whether the system uses the slopes or peaks of the ITD function centers around peak-type responses, as these features are within or close to the physiological delays an animal would experience. In contrast, the peaks of trough-type neurons almost invariably occur outside the animal's physiological range and their troughs show little, if any, modulation over this range. These features suggest that troughs are unlikely to play a role in sound localization. A proposed role for trough-type neurons rests on the idea that the maxima of peak-type and peaks of trough-type neurons formed a continuous ITD axis (Fig. 13.3) (Fitzpatrick et al. 2000, 2002). The ITDs of peak-type neurons fell within the physiological range and therefore may participate in sound localization, whereas the peaks of trough-type neurons occurred outside and could be used for analysis of features such as the spaciousness of the acoustic environment (Batra et al. 1997b).

## 2.5. MECHANISMS TO GENERATE INTERAURAL DELAYS

The Jeffress circuit model for ITD sensitivity accounts for many aspects of the psychophysics, and several of its tenets such as coincidence detection and characteristic delay are well supported by the physiological data. In the owl and chick, there is excellent evidence that the anatomical substrate for a delay line exists and, although apparently different in these two birds, that it operates much as Jeffress envisaged (Konishi et al. 1988; Rubel and Parks 1988; Carr and Konishi 1990; Carr 1993). However, the evidence for a delay-line system in mammals is more equivocal. This issue has been addressed by reconstructing the paths of axons to the MSO; however, the difficulty of these confirmatory experiments means that even those few reports (Smith et al. 1993; Beckius et al. 1999) are neither comprehensive nor, unfortunately, entirely convincing. There are several ways in which the necessary delays could arise such as graded length collaterals on both sides, graded axon lengths on both sides, or graded

collaterals or axons on only one side (Beckius et al. 1999). Both studies show evidence that axons that could provide the necessary graded delays and both conclude that the anatomical substrate for a delay line system is present in the cat.

Another method of generating interaural delays that does not involve axonal length variation was proposed by Schroeder (1977), expanded and formally modeled by Shamma et al. (1989), then by Bonham and Lewis (1999) and Shackleton et al. (2000), and investigated physiologically (Joris et al. 2005). Phase changes created by the conduction time of the activity along the basilar membrane to different frequencies are used directly by mismatching the best frequencies of the inputs to the coincidence detectors from each ear. Cross-correlation functions between auditory nerve fibers with different best frequencies have delays that vary in appropriate ways and over suitable ranges to account for ITD sensitivity measured in the IC (Joris et al. 2005).

Another way of generating interaural delay exists (Brand et al. 2002). Blocking inhibitory inputs to MSO neurons shifted the BD toward 0 ITD. They proposed that inhibition rather than axonal delay accounts for the empirically observed CD. The phase-locked, but ITD-insensitive inhibition shifts the peak of the ITD curve from 0 to about 1/8 of the period of the best frequency tone, thus providing the dependence of BD on best frequency. In support of this role for inhibition, in neonatal animals all BDs are near 0 ITD and they move away from 0 with development, unless deprived of spatial information (Seidl and Grothe 2003). Together, these studies suggest that in some species ITDs at low frequency may not be processed by a delay-line system, but rather an inhibitory input to MSO tunes the BD away from 0 ITD and optimally positions the slope in the physiological range.

## 2.6. TUNING TO INTERAURAL TIME DELAY

The ITD tuning widths systematically narrow at progressively higher stations along the auditory pathway. This frequency-by-frequency narrowing (Fig. 13.5A) appears in the composite curves (Fig. 13.5B) for the SOC, IC, and auditory thalamic neurons in the unanesthetized rabbit, all at 70 dB SPL. Equating for frequency range across structures (300 to 1000 Hz), the mean widths decreased 25% between the SOC and IC, and the total sharpening between the SOC and auditory thalamus was 50%.

For peak-type neurons, the widths of the peak of their composite curves decreases between the SOC and auditory thalamus, but for trough-type neurons, their widths remain relatively constant above the SOC (Fig. 13.5B). The widths in the SOC are probably overestimated because ITD sensitivity in this sample was not seen >1000 Hz, whereas in the samples at higher structures, many neurons had ITD sensitivity to 1500 Hz and more (Fig. 13.5A). With this caveat the decrease in peak width was 35% between the SOC and IC and, like the peak-widths by frequency (Fig. 13.5A), the total extent of sharpening in peak width from SOC to thalamus was 50%.






Figure 13.5. Peak width of ITD functions for neurons in the SOC, IC, and auditory thalamus as measured 50% from maximal discharge. (A) Peak width in cycles of interaural phase as a function of stimulus frequency. (B) Peak width ( $\mu\text{s}$ ) of composite curves of peak- and trough-type neurons.

Such sharpening of ITD functions could be due to many mechanisms (Kuwada et al. 1997). For example, inhibitory, peak-type inputs from the ipsilateral dorsal nucleus of the lateral lemniscus (DNLL) could interact with excitatory peak-type MSO inputs to create sharper peak-type IC responses, or inhibitory, peak-type contralateral DNLL inputs interacting with excitatory peak-type inputs from the opposite MSO could create an asymmetrical, sharper ITD function

(Fig. 13.1C). Similar sharpening could be created by peak-type inputs from the contralateral IC. Although this projection appears to be excitatory, it could sharpen if it acted through inhibitory interneurons. Other candidates are inhibitory, trough-type LSO or DNLL inputs interacting with excitatory, peak-type MSO inputs. IC sharpening may also be due to projections within the IC. Most, if not all, IC neurons display axon collaterals (Oliver et al. 1991) and 19% use  $\gamma$ -aminobutyric acid (GABA) (Oliver et al. 1994). Finally, sharpening may involve coincidence between similarly tuned ITD inputs converging onto an IC neuron.

Even in the auditory thalamus ITD functions are still broad and encompass about half of the physiological range of the rabbit (Fitzpatrick et al. 1997; Skottun 1998; Shackleton et al. 2003). Perhaps the auditory system uses neural ensembles, that is, a population code, to localize sound in space. A position-variable type model used to test the efficacy of the population code found that fewer neurons are required to detect ITD changes. The auditory thalamic sharpening appears optimal as further artificial sharpening degraded acuity (Fitzpatrick et al. 1997).

## 2.7. DYNAMIC VS. STATIC INTERAURAL TIME DELAY

Many studies varied ITD in discrete steps (i.e., statically) to measure neural ITD sensitivity and early comparisons of cat static and dynamic ITD sensitivity (generated by binaural beat stimuli) revealed only small differences (Yin and Kuwada 1983). However, in gerbil, when dynamic ITD stimuli were presented that corresponded to azimuthal movement of low-frequency sound sources through restricted arcs (e.g.,  $90^\circ$  centered at various azimuthal positions), sensitivities different from those with static ITDs emerged (Spitzer and Semple 1991, 1993). Such responses (Fig. 13.6A) show histograms locked to the cycle of repeating interaural phase modulations with the responses to the two directions shaded differently. The responses from such histograms as a function of the interaural phase difference (IPD: the ITD transformed into the phase difference for the frequency presented; Fig. 13.6B, dashed line) show the static IPD sensitivity. Some IPD values give static responses that are quite different from the same IPD presented dynamically. Furthermore, the same IPD in two different dynamic stimuli can vary from near minimum to near maximum discharge. In the extreme, IPD sensitivity curves could peak at very different IPD values when the IPD was static or part of a dynamic stimulus (Fig. 13.6C). A component of the sensitivity to the direction of dynamic IPD stimuli was consistent with adaptation (McAlpine et al. 2000). However, the static/dynamic differences (Spitzer and Semple 1991, 1993) greatly exceed those found by other authors, even using similar paradigms and this may reflect species and/or anesthesia differences manifest as different strengths of GABAergic inhibition. Larger effects were seen when tonic GABAergic inhibition to ITD-sensitive cells in the guinea pig was blocked, consistent with the adaptation of excitation hypothesis (McAlpine and Palmer 2002). Thus, adapting inhibitory inputs seem to be also required.




Figure 13.6. Dynamic and static ITD sensitivity compared. IC neural responses to static and dynamic variation in interaural phase difference (IPD). (A) The *curved arrow* shows the simulated movements generated by the phase modulations. The *unfilled bins* in the histograms show the response during clockwise motion and the *filled bins* the response during the motion in the opposite direction (center IPD values shown). (B) Response as a function of IPDs computed from the histograms in (A). *Dashed lines* show responses to static presentation and the *solid lines* that from the different presentations of dynamic IPD stimuli. (From Spitzer and Semple 1993). (C) Complete separation of the sensitivity curves to IPD when the response to specific IPDs are computed from dynamic stimuli or from static stimuli. (From M. Semple, personal communication.)

### 3. PROCESSING INTERAURAL TIME DELAYS IN ENVELOPES OF HIGH-FREQUENCY SOUNDS

Humans can resolve small differences in ITDs conveyed by the fine structure of sinusoids for frequencies to about 1200 Hz. Beyond this, ITD sensitivity decreases rapidly and disappears  $>1400$  Hz (Zwislocki and Feldman 1956). However, if the sound is amplitude modulated, ITDs are detected at much higher frequencies by using the ITDs in the envelopes of complex, high-frequency sounds (Leakey et al. 1958; David et al. 1959; McFadden and Pasanen 1976; Nuetzel and Hafter 1976; Bernstein and Trahiotis 1985a,b, 1994). Neurons sen-

sitive to ITDs in envelopes exist at many auditory stations (SOC: Yin et al. 1984; Yin and Chan 1990; Joris and Yin 1995; Joris 1996; Batra et al. 1997a,b; IC: Crow et al. 1980; Batra et al. 1989, 1993; auditory cortex: Fitzpatrick et al. 2000). The mechanism underlying envelope ITD sensitivity is similar in principle to that for low-frequency tones, beginning with envelope coding (discharges phase- or temporally locked to the envelope frequency) in the auditory nerve (Joris and Yin 1992) that is transmitted to cochlear nucleus bushy cells. These inputs converge onto the binaural SOC neurons that use a coincidence mechanism to become optimally tuned to a particular ITD.

### 3.1. TYPES OF ENVELOPE INTERAURAL TIME DELAY SENSITIVITY AND THEIR FREQUENCY DISTRIBUTION

The types of ITD sensitivity for low-frequency tones (peak-, trough-, and intermediate-type; Figs. 13.1 and 13.2) are also seen in the envelopes of amplitude-modulated, high-frequency tones (Batra et al. 1997a,b; Fitzpatrick et al. 2000, 2002), although ITD sensitivity in envelopes differs from that to tones: its lower and upper frequency limits are much lower (Fig. 13.7). In the rabbit, ITD envelope sensitivity begins at around 25 Hz (the lowest frequency tested) and the optimal frequency is approximately 150 Hz. In contrast, tone sensitivity begins at about 200 Hz (in the guinea pig <100 Hz) and the optimal frequency is nearly 900 Hz (Fig. 13.3). The frequency region where this encoding declines corresponds to the point where most neurons respond to envelopes. Combining tones and envelopes, ITD sensitivity extends from 25 to 2000 Hz.




Figure 13.7. The ranges of carrier and modulation frequencies over which rabbit IC neurons were sensitive to ITDs. Each point is the percentage of cells that showed ITD sensitivity at a tonal or modulation frequency. By using both tones and envelopes the frequency range is continuous from 25 Hz (the lowest modulation frequency tested) to about 2000 Hz.

Most neural and psychophysical studies used higher carrier frequencies in ITD envelope studies than those used to study ITD sensitivity to sound fine structure. Neurons with ITD sensitivity to fine structure and those preferring envelopes overlapped over a common range of best frequencies (Joris 2003). These findings are in accord with a model using normalized interaural correlation and with human data showing that envelope sensitivity occurs at a far lower carrier frequency than the best tonal ITD sensitivity (around 1300 Hz). The transition of ITD sensitivity in the fine structure to that of envelopes approximates a low-pass filter with an approximately 425 Hz cutoff (Bernstein and Trahiotis 1996a,b).

### 3.2. CODING OF ENVELOPE INTERAURAL TIME DELAYS

This broad ITD tuning suggests that at low envelope frequencies response modulation is nearly negligible over the physiological ITD range, leading to the speculation that envelope ITD sensitivity is a manifestation of dynamic interaural level sensitivity (Joris and Yin 1998). Envelope sensitive neurons, especially those that are trough-type, can have BDs far beyond the animal's physiological range. Such BDs could be activated by low interaural correlations from sound reflections in a reverberant environment. These neurons may not be involved in sound location per se, but in analyzing the properties of the auditory space. Based on the broadness of ITD functions evoked by envelopes, behavioral ITD sensitivity is predicted to be poor. Surprisingly, human listeners detect changes in the ITDs in envelopes  $<200 \mu\text{s}$  for a sinusoidal envelope of 64 Hz and even less for a 64-Hz "transposed" envelope that provides the high-frequency channels with information similar to that available in low-frequency channels (Bernstein and Trahiotis 2002, 2003). Models using normalized interaural correlations are in accord with such behavioral acuity as are neural studies showing high sensitivity to changes in interaural correlation (Yin et al. 1986b).

## 4. PROCESSING OF INTERAURAL LEVEL DIFFERENCES

### 4.1. TYPES OF INTERAURAL LEVEL DIFFERENCE SENSITIVITY

The first major site of binaural interaction takes place in the superior olive in the brain stem. LSO neurons receiving excitatory input from one ear and inhibitory input from the other are sensitive to ILDs (Goldberg and Brown 1969). As the inhibition (I) is from the contralateral ear and the excitation (E) is ipsilateral, such cells are IE. Owing to the crossed projection to the IC from the high-frequency part of the LSO, at higher levels of the auditory pathway the cells responsive to ILDs are designated as EI (excited contralaterally and inhibited ipsilaterally). Two different and largely complementary stimulus paradigms are used to investigate ILD sensitivity. In one, the contralateral stimulus is held at a constant level while the ipsilateral stimulus level is varied (excitatory mon-

aural intensity constant: EMI). For EI cells this method reveals the inhibitory effects of ipsilateral stimulation. A second paradigm varies the levels at the two ears symmetrically in opposite directions to provide ILD variations about an average binaural intensity level (ABI-constant). This better mimics ILD variation as a sound source traverses azimuthal positions, increasing level at one ear as it decreases at the other (Irvine 1986, 1991).

Most EI cells in the IC with monotonic rate-level functions to monaural, contralateral sounds have a stereotypical sigmoid-shaped sensitivity to ILDs with EMI or ABI (Fig. 13.8A). The response is maximal over an ILD range at which




Figure 13.8. Types of ILD curves and their distribution. (A) Normalized spike rate as a function of ILD for 6 IC neurons for BF tones showing the variation in the position of the half-maximum ILD value. (Modified from Irvine 1986.) (B, C) Normalized spike rate as a function of ILD for two cells measured using a constant BF tone at the contralateral ear (response shown as *dashed line*). (D, E) As for (B, C) for the same two neurons but measured with the average binaural intensity constant. (B, E modified from Irvine 1991.) (F, G) Normalized spike rate as a function of ITD for three cells. (F) A monotonic neuron whose ILD sensitivity is invariant with ABI. (G) A monotonic cell with a shift to the right as the ABI is increased. (H) A nonmonotonic unit shifts to the left with increased ABI. (F-H modified from Irvine and Gago 1990.) (I) Distribution of the half-maximal ILD values. (From Irvine and Gago 1990.)

the contralateral excitatory sound level is greatest, and it is partially or totally suppressed over ILDs at which the ipsilateral inhibitory ear is loudest. Between these extremes neural output varies directly with the ILD.

For neurons with nonmonotonic rate-level functions to monaural contralateral stimuli the ILD functions using EMI and ABI methods are usually different. The ILD function with the fixed contralateral tone may be sigmoid (Fig. 13.8B, C) whereas the ABI-derived ILD function for the same neurons is highly nonmonotonic (Fig. 13.8D, E) (Irvine and Gago 1990). Because the ABI paradigm moves from left to right in the plot (+40 to -40 dB ILD) the contralateral stimulus is declining while the ipsilateral stimulus grows, so that the ILD function is the net result of level dependent ipsilateral and contralateral influences. The responses to the contralateral stimulus (Fig. 13.8, dashed lines) show that ILD functions reflect facilitatory action of the ipsilateral input at positive ILDs, and the converse at negative ILDs, when the ipsilateral tone was louder (Irvine 1987). Mixed responses to stimulation of both ears are also reported (Irvine 1991).

Analyses of sound level variations on ILD sensitivity are available in gerbil, mustached bat, cat, and other species (Semple and Kitzes 1987; Wenstrup et al. 1988; Irvine and Gago 1990) and show that level affects all but a few neurons (Fig. 13.8F). The commonest variation as ABI increased was for the sigmoid-shaped ILD function to shift to the right, that is, for activity to encroach on the region where ipsilateral inhibition dominated the ILD function at lower ABIs (or into the ipsilateral spatial hemifield: Fig. 13.8G). More dramatic changes occur in neurons with nonmonotonic ILD functions (Fig. 13.8H) and a few cells have nonmonotonicity in their responses to ILD and to ABI (Semple and Kitzes 1987; Irvine and Gago 1990). For such neurons the maximal discharge rate is often evoked only by a limited ILD and ABI range.

#### 4.2. DISTRIBUTION OF INTERAURAL LEVEL DIFFERENCES

We do not know what aspect of ILD sensitive neural activity is important for ILD and spatial position discrimination. Perhaps ILD is encoded by a population response. As the azimuthal position of a sound source moves from the ipsilateral to the contralateral hemifield, IC neurons will first be strongly inhibited by the ipsilateral ear input, then progressively activated, and eventually saturate. A growing population of cells is activated by the variation in the position of the ILD range over which the neuron is sensitive (Fig. 13.8A). The transition region ranges between domination by the different ear inputs and is quantified as the ILD value at half-maximum output. Its distribution for cat central nucleus cells (Fig. 13.8I) spans the range of ILDs that this animal naturally experiences (Irvine and Gago 1990). Such a population code might depend only on those cells whose medial ILD borders are invariant with ABI or on a more complex representation using all information, including that on changes with ABI.

An alternative to population coding suggests that neurons have both a preferred ILD and ABI (Semple and Kitzes 1987). The position at which a cell is

maximally activated might provide an ILD place code. Although the data are limited there is evidence for a systematic variation in the position of the half-maximum ILD, moving from rostral to caudal along a single IC frequency lamina (Irvine and Gago 1990), and the ILD functions more rostrally show sharper transitions and stronger inhibition.

### *4.3. MECHANISMS OF INTERAURAL LEVEL DIFFERENCE SENSITIVITY*

Undoubtedly much of the initial ILD information processing takes place in the LSO, and the excitation from the ipsilateral ear and contralateral ears endows many high-frequency LSO cells with their ILD sensitivity (Goldberg and Brown 1969; Boudreau and Tsuchitani 1970; Tollin and Yin 2002a,b). However, other mechanisms clearly give rise to, or modify, the ILD sensitivity in the IC. Evidence includes deactivation of the superior olivary complex (even bilaterally) or DNLL, which does not abolish ILD sensitivity (Li and Kelly 1992a,b; Sally and Kelly 1992), and deactivation within the IC of local inhibition or of inhibitory inputs to the IC leaves some ILD sensitivity intact but can modify or abolish the ILD of many cells (Pollak et al. 2002). It is suggested that, owing to the level dependency of first spike latency, that ILD sensitivity ensues from brain stem coincidence detectors operating like the ITD-sensitive MSO cells (Jeffress 1948). Testing this model by counteracting the level dependent latency shift abolished the ILD sensitivity only in some IC and LSO cells (Irvine et al. 1995, 2001). Using virtual space stimuli, the spatial receptive fields of LSO cells are generated by their sensitivity to the ILDs in a band of frequencies around the cells' characteristic frequency; however, LSO spatial tuning accounts for only 75% of that in IC (Tollin and Yin 2002a,b). These data and the mixed facilitation and inhibition in ILD functions at the IC suggest that diverse supraolivary mechanisms shape midbrain ILD sensitivity.

## **5. ROLE OF THE INFERIOR COLLICULUS IN PERCEPTUAL PHENOMENA**

### *5.1. BINAURAL MASKING LEVEL DIFFERENCES*

The ability to detect a signal masked by a noise is strongly influenced by the relative interaural similarity between signal and masker. When a signal, identical at the two ears, is masked by a noise identical at the two ears, inverting either can achieve a 12- to 15-dB improvement in signal detection (Hirsh 1948a,b; Licklider 1948). For unmodulated tonal signals the mechanism for this binaural masking level difference (BMLD) might reside in low-level ITD-sensitive cells, because large BMLDs occur only at low frequency and BMLDs depend on the relative phase at the ears. Large BMLDs occur at higher frequencies when the signal envelope fluctuates from mechanisms analogous to those at low frequency,



but involves cells sensitive to envelope ITDs (Bernstein and Trahiotis 1992; van de Par and Kohlrausch 1997).

Early attempts to find a neural mechanism for this unmasking by comparing activity to identical noise (N0) and tone (S0) at the two ears with that to identical noise (N0) but inverted signal ( $S\pi$ ) were not successful, although reduced MSO discharge to  $N0S\pi$  stimuli was consistent with coincidence detection models (Langford 1984). Subsequent studies tracking masked thresholds while varying masker or signal ITD showed greatest ITD masking for the masking noise evoking most activity (Caird et al. 1991; McAlpine et al. 1996a). Tracking methods were, however, limited to discharge rate increases, and only when full rate-level functions to the signals and fixed levels of masker noise were employed with signal detection methods did a more consistent picture emerge. IC responses to BMLD signals are entirely consistent with their ITD sensitivity to noise and tones (Jiang et al. 1997a,b; Palmer et al. 2000). In most neurons with peak or intermediate ITD functions and near-0 BD, S0 tones can be detected by increased discharge and  $S\pi$  tones by a decrease (Fig. 13.9A). These BMLDs are consistent with the psychophysics. Trough units show the inverse. Identical noises (N0) and noise inverted at one ear ( $N\pi$ ) drive neurons according to their respective ITD functions and S0 tones are again detected by an increased discharge above this rate (Fig. 13.9B). Masked thresholds are obtained from rate-level functions using signal detection standard separation methods (Sakitt 1973) (Fig. 13.9A, B: arrows). Such masked thresholds for populations of neurons responding to  $N0S0$ ,  $N0S\pi$ , and  $N\pi S0$  signals at 500 Hz show the lowest thresholds near the signal frequency and  $N0S0$  masked thresholds tend to be  $>N0S\pi$  (Fig. 14.9C) or the  $N\pi S0$  (Fig. 14.9D). This is further emphasized by taking the average masked thresholds for different BMLD stimuli (Fig. 13.9E, F), which is smaller than that in human psychophysics. However, that  $N0S0$  vs.  $N0S\pi$  BMLD is larger than  $N0S0$  vs.  $N\pi S0$  BMLD is consistent with the hierarchy in humans (Hirsh 1948a,b). Different neural populations contribute to detection of the tonal signal in different binaural configurations (Jiang et al. 1997a,b). The empirical neural responses to BMLD stimuli are consistent with both the type of ITD sensitivities and with the predictions from interaural cross-correlation models.

## 5.2. THE PRECEDENCE EFFECT

This term signifies that the auditory system can discern the actual location of sound sources despite the mass of reflections and echoes that arrive soon after the direct waves strike the ears (Wallach et al. 1949). The precedence effect has been studied by presenting two brief, successive sounds through two spatially separated speakers or through headphones. The leading sound, the “conditioner,” simulates the direct wavefront whereas the second, or delayed sound, the “probe,” simulates a reflection or an echo. Speakers are usually placed in front of the listener and on either side of the midline, and the sound from the probe speaker is systematically delayed. Under headphones, ITDs commonly provide




Figure 13.9. Responses of IC units to BMLD stimuli. **(A)** Rate level functions to S0 and  $S\pi$  tones at 500 Hz in the presence of NO noise masker. The arrows show the masked thresholds determined by signal detection methods. The inset shows the value of the MLD. **(B)** As for **(A)**: rate-level functions to S0 tones at 500 Hz in the presence of NO and N $\pi$  noise masker. **(C)** Masked threshold for NOS0 (filled circles) and NOS $\pi$  (open squares) for IC neurons responding to a 500-Hz tone. Vertical line, 500 Hz; dashed lines, 300 and 800 Hz. **(D)** As for **(C)** for NOS0 (filled circles) and N $\pi$ S0 (open squares). **(E)** Average masked thresholds across the 300–800-Hz neurons from **(C)**. **(F)** Average masked thresholds across 300- to 800-Hz neurons from **(D)**. (Modified from Jiang et al. 1997a; Palmer et al. 2000.)

different lateralities of the conditioner and probe. Behaviorally, a sequence of perceptual events occurs as the interval between two identical sounds from different locations increases (Zurek 1987; Blauert 1983; Litovsky et al. 1999). When the “conditioner-probe interval” (CPI) is approximately 1 ms, the listener hears a single, fused auditory event and judges its location to be between the conditioner and probe. This is the window of “summing localization” and the perception follows the physical cues present from the combination of the sources (Blauert 1983). When the CPI exceeds 1 ms, the listener still hears a fused auditory perception but the location is assigned to the conditioner. This is the start of the precedence (or localization dominance) window. The end of the precedence window occurs at “echo threshold,” or the CPI at which the probe is heard as a separate sound in its own location. For clicks, echo threshold occurs at CPIs ranging from 2 to 10 ms, and headphone studies have shorter values (from 2 to 4 ms) than free-field studies. Far beyond echo threshold, the probe is heard more faintly than the conditioner (Blauert 1983). Although many data have been obtained from humans, experiments in rats (Kelly 1974) and cats (Cranford and Oberholzer 1976; Populin and Yin 1998; Tollin 2003) indicate similar phenomena.

The neural basis for the precedence effect has been studied in several species and structures (Yin 1994; Fitzpatrick et al. 1995, 1999; Keller and Takahashi 1996; Parham et al. 1996, 1998; Wickesberg 1996; Litovsky and Yin 1998a,b; Wickesberg and Stevens 1998). The CPI at which the neural response to the probe is 50% of that to the conditioner (the half-maximal delay) has been used as a measure of the neuron’s recovery time. Within the summing localization window (CPI < 1 ms), IC neurons sensitive to sound-source location in cats and owls respond to “phantom images” in accord with the physical cues present during this window (Yin 1994; Keller and Takahashi 1996).

The recovery functions for neurons in the auditory nerve, cochlear nucleus, superior olivary complex, IC, and auditory cortex in unanesthetized animals show median recovery times in the first three stations that are within the precedence window (< 3 ms). In contrast, in the IC and auditory cortex they increase to 6 and 20 ms, respectively (Fig. 13.10). Anesthesia increases the recovery times of IC and auditory cortex neurons. In the unanesthetized rabbit and behaving cat IC (Fitzpatrick et al. 1995; Tollin 2003) the median 50% recovery was near or within the precedence window (< 10 ms) while in anesthetized cats median recovery times extend beyond (17 to 28 ms; Yin 1994; Litovsky and Yin 1998a,b). In the unanesthetized rabbit auditory cortex median recovery times are about 20 ms (Fitzpatrick et al. 1999) and those in the anesthetized cat are approximately 100 ms (Reale et al. 1995).

While the precedence effect was long thought to involve inhibitory mechanisms (Lindemann 1986) modeling studies (Hartung and Trahiotis 2001; Trahiotis and Hartung 2002) indicate that cochlear filter interactions and hair cell-based compression and adaptation account for the results of many precedence experiments, including behavioral data (Wallach et al. 1949) and neural responses (Fitzpatrick et al. 1999).




Figure 13.10. Population recovery functions to click pairs at different levels along the auditory pathway.  $n$  = Number of cells. Those from the auditory nerve (AN), AVCN, and SOC were all similar and short, with 50% recovery times of about 2 ms. By echo threshold (4 ms CPI) the recovery was near 70%. For the IC, recovery was much slower, with 50% at approximately 7 ms and nearly 40% recovery at echo threshold. In the auditory cortex the recovery was slower yet, with 50% recovery at 20 ms and 25% recovery at echo threshold. Neurons in the AN and AVCN were from an unanesthetized, decerebrate cat, and the others from an unanesthetized rabbit.

The IC and cortical neurons with long recovery times have behavioral correlates, including a reduction in the discriminability of the ITD in the probe (Shinn-Cunningham et al. 1993; Tollin and Henning 1998), and reduced perceived loudness of the probe compared to the conditioner (Blauert 1983) for a period well beyond echo threshold. Such neurons may participate in coding features unrelated to the precedence effect, for example, spaciousness or timbre. Inhibition may modulate such neurons since inactivating glycinergic (M. Paterson, personal communication) or GABAergic inhibition (Zhou and Jen 2003) can reduce IC recovery time in guinea pig and big brown bat neurons, respectively.

### 5.3. BINAURAL PITCH

Pitch not only is important for musical perception, but also is probably the strongest cue for segregation of sounds (Darwin 1997). Some pitches are created solely from the binaural cues that result when narrow frequency bands within binaurally identical noise (correlated), have a reduced correlation (i.e., decorrelation, Culling et al. 1998). Huggins pitch is a binaural pitch associated with the frequency at which decorrelation is caused by a rapid phase difference between two dichotic noise sources (Cramer and Huggins 1958). This pitch resem-

bles a binaural unmasking effect resulting from the decorrelation of activity within the frequency channels at which the phase difference occurs (Culling et al. 1998). Physiological and modeling experiments show that IC neural responses are consistent with this interpretation, showing firing rate changes and decorrelation effects consistent with the cross correlation model of binaural hearing (Hancock and Delgutte 2002). The neural responses to stimuli generating binaural pitches are entirely consistent with models of binaural cross correlation.

## 6. FUTURE DIRECTIONS

Detailed descriptions are now available that account for many elements of the auditory signal that are used in specific binaural contexts represented in the auditory midbrain. However, there has been much emphasis on abstracted binaural cues presented via earphones as a tool for understanding underlying mechanisms. This would seem to be a propitious point to advocate the inclusion of more ecological and realistic listening situations. The effort to link perception to the responses recorded in anesthetized animals can be taken only so far and recordings in the unanesthetized animal engaged in behaviors such as localizing may provide new and more relevant information on the brain processes underlying these behaviors. Remarkable abilities such as sound source segregation and the binding of their components into a single auditory object ultimately depend on auditory midbrain processing and require consideration of the animal's reactions to the sounds. Virtual auditory environments may represent an enabling technology to allow new stimulus paradigms and for the study of IC spatial receptive fields (Sterbing et al. 2003b). The use of individualized HRTFs is critical for accurate assessment of spatial receptive fields, which are smaller than those reported when using nonindividualized HRTFs.

Some binaural abilities will no doubt be reflected only in population measures of activation or even synchronized or correlated population activity. If so, such representations will require simultaneous monitoring of many neurons, approaches that are becoming routinely available. Finally, we encourage the use of naturally occurring and meaningful sounds and analysis of the necessarily complicated responses that are evoked. This approach will probably entail study of the auditory ecology of animal models that have proved so successful in the case of the bat and barn owl.

## Abbreviations

ABI	average binaural intensity
AN	auditory nerve
AVCN	anteroventral cochlear nucleus
BD	best delay
BMLD	binaural masking level difference
CD	characteristic delay

CP	characteristic phase
CPI	conditioner-probe interval
DNLL	dorsal nucleus of lateral lemniscus
E	excitation
EMI	excitatory monaural intensity constant
GABA	$\gamma$ -aminobutyric acid
HRTF	head-related transfer function
I	inhibition
IC	inferior colliculus
ILD	interaural level difference
IPD	interaural phase difference
ITD	interaural time difference
LSO	lateral superior olive
MSO	medial superior olive
$N\pi$	inverted noise
$N0$	noise
$S0$	signal
SOC	superior olivary complex
$S\pi$	inverted signal

## References

- Batra R, Kuwada S, and Stanford TR (1989) Temporal coding of envelopes and their interaural delays in the inferior colliculus of the unanesthetized rabbit. *Journal of Neurophysiology* **61**:257–268.
- Batra R, Kuwada S, and Stanford TR (1993) High-frequency neurons in the inferior colliculus that are sensitive to interaural delays of amplitude-modulated tones: evidence for dual binaural influences. *Journal of Neurophysiology* **70**:64–80.
- Batra R, Kuwada S, and Fitzpatrick DC (1997a) Sensitivity to interaural temporal disparities of low- and high-frequency neurons in the superior olivary complex. I. Heterogeneity of responses. *Journal of Neurophysiology* **78**:1222–1236.
- Batra R, Kuwada S, and Fitzpatrick DC (1997b) Sensitivity to interaural temporal disparities of low- and high-frequency neurons in the superior olivary complex. II. Coincidence detection. *Journal of Neurophysiology* **78**:1237–1247.
- Beckius GE, Batra R, and Oliver DL (1999) Axons from anteroventral cochlear nucleus that terminate in medial superior olive of the cat: observations related to delay lines. *Journal of Neuroscience* **19**:3146–3161.
- Bernstein LR and Trahiotis C (1985a) Lateralization of low-frequency, complex waveforms: the use of envelope-based temporal disparities. *Journal of the Acoustical Society of America* **77**:1868–1880.
- Bernstein LR and Trahiotis C (1985b) Lateralization of sinusoidally amplitude-modulated tones: effects of spectral locus and temporal variation. *Journal of the Acoustical Society of America* **78**:514–523.
- Bernstein LR and Trahiotis C (1992) Discrimination of interaural envelope correlation and its relation to binaural unmasking at high-frequencies. *Journal of the Acoustical Society of America* **91**:306–316.

- Bernstein LR and Trahiotis C (1994) Detection of interaural delay in high-frequency sinusoidally amplitude-modulated tones, two-tone complexes, and bands of noise. *Journal of the Acoustical Society of America* **95**:3561–3567.
- Bernstein LR and Trahiotis C (1996a) The normalized correlation: accounting for bin-aural detection across center frequency. *Journal of the Acoustical Society of America* **100**:3774–3784.
- Bernstein LR and Trahiotis C (1996b) On the use of the normalized correlation as an index of interaural envelope correlation. *Journal of the Acoustical Society of America* **100**:1754–1763.
- Bernstein LR and Trahiotis C (2002) Enhancing sensitivity to interaural delays at high frequencies by using “transposed stimuli.” *Journal of the Acoustical Society of America* **112**:1026–1036.
- Bernstein LR and Trahiotis C (2003) Enhancing interaural-delay-based extents of laterality at high frequencies by using “transposed stimuli”. *Journal of the Acoustical Society of America* **113**:3335–3347.
- Blauert J (1983) *Spatial Hearing. The Psychophysics of Human Sound Localization*. MIT Press, Cambridge.
- Bonham BH and Lewis ER (1999) Localization by interaural time difference (ITD): effects of interaural frequency mismatch. *Journal of the Acoustical Society of America* **106**:281–290.
- Boudreau JC and Tsuchitani C (1970) Cat superior olive S-segment cell discharge to tonal stimulation. In: Neff WD (ed). *Contributions to Sensory Physiology*. Academic Press, New York, pp. 143–213.
- Brand A, Behrend O, Marquardt T, McAlpine D, and Grothe B (2002) Precise inhibition is essential for microsecond interaural time difference coding. *Nature* **417**:543–547.
- Buell TN, Trahiotis C, and Bernstein LR (1991) Lateralization of low-frequency tones: relative potency of gating and ongoing interaural delays. *Journal of the Acoustical Society of America* **90**:3077–3085.
- Caird D and Klinke R (1983) Processing of binaural stimuli by cat superior olivary complex neurons. *Experimental Brain Research* **52**:385–399.
- Caird D and Klinke R (1987) Processing of interaural time and intensity differences in the cat inferior colliculus. *Experimental Brain Research* **68**:379–392.
- Caird DM, Palmer AR, and Rees A (1991) Binaural masking level difference effects in single units of the guinea pig inferior colliculus. *Hearing Research* **57**:91–106.
- Carney LH and Yin TCT (1989) Responses of low-frequency cells in the inferior colliculus to interaural time differences of clicks: excitatory and inhibitory components. *Journal of Neurophysiology* **62**:144–161.
- Carr CE (1993) Delay line models of sound localization in the barn owl. *American Zoologist* **33**:79–85.
- Carr CE and Konishi M (1990) A circuit for detection of interaural time differences in the brain stem of the barn owl. *Journal of Neuroscience* **10**:3227–3246.
- Cramer EM and Huggins WH (1958) Creation of pitch through binaural interaction. *Journal of the Acoustical Society of America* **30**:858–866.
- Cranford JL and Oberholzer M (1976) Role of neocortex in binaural hearing in the cat. II. The ‘precedence effect’ in sound localization. *Brain Research* **111**:225–239.
- Crow G, Langford TL, and Moushegian G (1980) Coding of interaural time differences by some high-frequency neurons of the inferior colliculus: responses to noise bands and two-tone complexes. *Hearing Research* **3**:147–153.

- Culling JF, Summerfield AQ, and Marshall DH (1998) Dichotic pitches as illusions of binaural unmasking. I. Huggin's pitch and the "binaural edge pitch." *Journal of the Acoustical Society of America* **103**:3509–3526.
- D'Angelo WR, Sterbing SJ, Ostapoff E-M, and Kuwada S (2003) Effects of amplitude modulation on the coding of interaural time differences of low-frequency sounds in the inferior colliculus. II. Neural mechanisms. *Journal of Neurophysiology* **90**:2827–2836.
- Darwin CJ (1997) Auditory grouping. *Trends in Cognitive Sciences* **1**:327–333.
- David EE, Guttman N, and van Bergeijk WA (1959) Binaural interaction of high-frequency complex stimuli. *Journal of the Acoustical Society of America* **31**:774–782.
- Davis KA, Ramachandran R, and May BJ (1999) Single-unit responses in the inferior colliculus of decerebrate cats II. Sensitivity to interaural level differences. *Journal of Neurophysiology* **82**:164–175.
- Davis KA, Ramachandran R, and May BJ (2003) Auditory processing of spectral cues for sound localization in the inferior colliculus. *Journal of the Association for Research in Otolaryngology* **4**:148–163.
- Durlach NI (1972) Binaural signal detection: equalization and cancellation theory. In: Tobias JV (ed). *Foundations of Modern Auditory Theory*, Volume II. Academic Press, New York, pp. 369–462.
- Fitzpatrick DC and Kuwada S (2001) Tuning to interaural time differences across frequency. *Journal of Neuroscience* **21**:4844–4851.
- Fitzpatrick DC, Kuwada S, Batra R, and Trahiotis C (1995) Neural responses to simple simulated echoes in the auditory brain stem of the unanesthetized rabbit. *Journal of Neurophysiology* **74**:2469–2486.
- Fitzpatrick DC, Batra R, Stanford TR, and Kuwada S (1997) A neuronal population code for sound localization. *Nature* **388**:871–874.
- Fitzpatrick DC, Kuwada S, Kim DO, Parham K, and Batra R (1999) Responses of neurons to click-pairs as simulated echoes: auditory nerve to auditory cortex. *Journal of the Acoustical Society of America* **106**:3460–3472.
- Fitzpatrick DC, Kuwada S, and Batra R (2000) Neural sensitivity to interaural time differences: beyond the Jeffress model. *Journal of Neuroscience* **20**:1605–1615.
- Fitzpatrick DC, Kuwada S, and Batra R (2002) Transformations in processing interaural time differences between the superior olivary complex and inferior colliculus: beyond the Jeffress model. *Hearing Research* **168**:79–89.
- Goldberg JM and Brown PB (1969) Response of binaural neurons of dog superior olivary complex to dichotic tonal stimuli: some physiological mechanisms of sound localization. *Journal of Neurophysiology* **32**:613–636.
- Grantham DW (1997) Auditory motion perception: snapshots revisited. In: Gilkey RH and Anderson TR (eds). *Binaural Hearing in Real and Virtual Environments*. Lawrence Erlbaum, Mahwah, pp. 295–314.
- Hancock KE and Delgutte B (2002) Neural correlates of the Huggins dichotic pitch. *Association for Research in Otolaryngology Abstracts* **25**:153.
- Harnischfeger G (1980) Brainstem units of echolocating bats code binaural time differences in the microsecond range. *Naturwissenschaften* **67**:314–316.
- Hartung K and Trahiotis C (2001) Peripheral auditory processing and investigations of the "precedence effect" which utilize successive transient stimuli. *Journal of the Acoustical Society of America* **110**:1505–1513.
- Heffner HE and Heffner RS (1985) Sound localization in wild Norway rats (*Rattus norvegicus*). *Hearing Research* **19**:151–155.



- Heffner RS (1997) Comparative study of sound localization and its anatomical correlates in mammals. *Acta Otolaryngologica (Stockholm)*. Supplement **532**:46–53.
- Heffner RS and Heffner HE (1987) Localization of noise, use of binaural cues, and a description of the superior olivary complex in the smallest carnivore, the least weasel (*Mustela nivalis*). *Behavioral Neuroscience* **101**:701–708, 744–745.
- Heffner RS and Heffner HE (1988a) Sound localization acuity in the cat: effect of azimuth, signal duration, and test procedure. *Hearing Research* **36**:221–232.
- Heffner RS and Heffner HE (1988b) Sound localization and use of binaural cues by the gerbil (*Meriones unguiculatus*). *Behavioral Neuroscience* **102**:422–428.
- Hirsh IJ (1948a) Binaural summation and interaural inhibition as a function of the noise level of the masking noise. *American Journal of Psychology* **56**:205–213.
- Hirsh IJ (1948b) The influence of interaural phase on interaural summation and inhibition. *Journal of the Acoustical Society of America* **20**:536–544.
- Irvine DRF (1986) The auditory brainstem. A review of the structure and function of auditory brainstem processing mechanisms. In: Autrum H, Ottoson D, Perl ER, Schmidt RF, Shimazu H, and Willis WD (eds). *Progress in Sensory Physiology* **7**:1–279.
- Irvine DRF (1987) A comparison of two methods for the measurement of neural sensitivity to interaural intensity differences. *Hearing Research* **30**:169–180.
- Irvine DRF (1991) Physiology of the auditory brainstem. In: Popper AN and Fay RR (eds). *Springer Handbook of Auditory Research*, Volume 2: *The Mammalian Auditory Pathway: Neurophysiology*. Springer-Verlag, New York, pp. 153–231.
- Irvine DRF and Gago G (1990) Binaural interaction in high-frequency neurons in inferior colliculus of the cat: effects of variations in sound pressure levels on sensitivity to interaural intensity differences. *Journal of Neurophysiology* **63**:570–591.
- Irvine DRF, Park VN, and Mattingley JB (1995) Responses of neurons in the inferior colliculus of the rat to interaural time and intensity differences in transient stimuli: implications for the latency hypothesis. *Hearing Research* **85**:127–141.
- Irvine DRF, Park VN, and McCormick L (2001) Mechanisms underlying the sensitivity of neurons in the lateral superior olive to interaural intensity differences. *Journal of Neurophysiology* **86**:2647–2666.
- Jeffress LA (1948) A place code theory of sound localization. *Journal of Comparative and Physiological Psychology* **44**:35–39.
- Jeffress LA (1958) Medial geniculate body: a disavowal. *Journal of the Acoustical Society of America* **30**:802–803.
- Jenkins WM and Masterton RB (1982) Sound localization: effects of unilateral lesions in central auditory system. *Journal of Neurophysiology* **47**:987–1016.
- Jiang D, McAlpine D, and Palmer AR (1997a) Detectability index measures of binaural masking level difference across populations of inferior colliculus neurons. *Journal of Neuroscience* **17**:9331–9339.
- Jiang D, McAlpine D, and Palmer AR (1997b) Responses of neurons in the guinea pig inferior colliculus to binaural masking level difference stimuli measured by rate-versus-level functions. *Journal of Neurophysiology* **77**:3085–3106.
- Joris PX (1996) Envelope coding in the lateral superior olive. II. Characteristic delays and comparison with responses in the medial superior olive. *Journal of Neurophysiology* **76**:2137–2156.
- Joris PX (2003) Interaural time sensitivity dominated by cochlea-induced envelope patterns. *Journal of Neuroscience* **23**:6345–6350.
- Joris PX and Yin TCT (1992) Responses to amplitude-modulated tones in the auditory nerve of the cat. *Journal of the Acoustical Society of America* **91**:215–232.

- Joris PX and Yin TCT (1995) Envelope coding in the lateral superior olive. I. Sensitivity to interaural time differences. *Journal of Neurophysiology* **73**:1043–1062.
- Joris PX and Yin TCT (1998) Envelope coding in the lateral superior olive. III. Comparison with afferent pathways. *Journal of Neurophysiology* **79**:253–269.
- Joris PX, van der Heijden M, Louage D, van de Sande B, and van Kerckhoven C (2005) Dependence of binaural and cochlear “best delays” on characteristic frequency. In: Pressnitzer D, de Cheveigné A, McAdams S, and Collet L (eds). *Auditory Signal Processing: Physiology, Psychophysics, and Models*. Springer-Verlag, New York, pp. 396–402.
- Keller CH and Takahashi TT (1996) Responses to simulated echoes by neurons in the barn owl’s auditory space map. *Journal of Comparative Physiology A* **178**:499–512.
- Kelly JB (1974) Localization of paired sound sources in the rat: small time differences. *Journal of the Acoustical Society of America* **55**:1277–1284.
- Kelly JB and Phillips DP (1991) Coding of interaural time differences of transients in auditory cortex of *Rattus norvegicus*: implications for the evolution of mammalian sound localization. *Hearing Research* **55**:39–44.
- Kiang NY-S, Watanabe T, Thomas EC, and Clark LF (1965) *Discharge Patterns of Single Fibers in the Cat’s Auditory Nerve*. MIT Press, Cambridge.
- Konishi M, Takahashi TT, Wagner H, Sullivan WE, and Carr CE (1988) Neurophysiological and anatomical substrates of sound localization in the owl. In: Edelman GM, Gall WE, and Cowan WM (eds). *Auditory Function. Neurobiological Bases of Hearing*. John Wiley & Sons, New York, pp. 721–746.
- Kuwada S, Stanford TR, and Batra R (1987) Interaural phase-sensitive units in the inferior colliculus of the unanesthetized rabbit: effects of changing frequency. *Journal of Neurophysiology* **57**:1338–1360.
- Kuwada S and Yin TCT (1983) Binaural interaction in low-frequency neurons in inferior colliculus of the cat. I. Effects of long interaural delays, intensity, and repetition rate on interaural delay function. *Journal of Neurophysiology* **50**:981–999.
- Kuwada S, Batra R, and Fitzpatrick DC (1997) Neural processing of binaural temporal cues. In: Gilkey RH and Anderson TR (eds). *Binaural and Spatial Hearing in Real and Virtual Environments*. Lawrence Erlbaum Associates, Mahwah, pp. 399–425.
- Langford TL (1984) Responses elicited from medial superior olivary neurons by stimuli associated with binaural masking and unmasking. *Hearing Research* **15**:39–50.
- Leakey DM, Sayers BM, and Cherry C (1958) Binaural fusion of low- and high-frequency sounds. *Journal of the Acoustical Society of America* **30**:222.
- Li L and Kelly JB (1992a) Binaural responses in rat inferior colliculus following kainic acid lesions of the superior olive: interaural intensity difference functions. *Hearing Research* **61**:73–85.
- Li L and Kelly JB (1992b) Inhibitory influence of the dorsal nucleus of the lateral lemniscus on binaural responses in the rat’s inferior colliculus. *Journal of Neuroscience* **12**:4530–4539.
- Licklider JCR (1948) The influence of interaural phase relations upon the masking of speech by white noise. *Journal of the Acoustical Society of America* **20**:150–159.
- Lindemann W (1986) Extension of a binaural cross-correlation model by contralateral inhibition. Part II: the law of the first wavefront. *Journal of the Acoustical Society of America* **80**:1623–1630.
- Litovsky RY and Yin TCT (1998a) Physiological studies of the precedence effect in the inferior colliculus of the cat. I. Correlates of psychophysics. *Journal of Neurophysiology* **80**:1285–1301.

- Litovsky RY and Yin TCT (1998b) Physiological studies of the precedence effect in the inferior colliculus of the cat. II. Neural mechanisms. *Journal of Neurophysiology* **80**: 1302–1316.
- Litovsky RY, Colburn HS, Yost WA, and Guzman SJ (1999) The precedence effect. *Journal of the Acoustical Society of America* **106**:1633–1654.
- Masterton B, Heffner H, and Ravizza R (1969) The evolution of human hearing. *Journal of the Acoustical Society of America* **45**:966–985.
- McAlpine D and Palmer AR (2002) Blocking GABAergic inhibition increases sensitivity to sound motion cues in the inferior colliculus. *Journal of Neuroscience* **22**:1443–1453.
- McAlpine D, Jiang D, and Palmer AR (1996a) Binaural masking level differences in the inferior colliculus of the guinea pig. *Journal of the Acoustical Society of America* **100**: 490–503.
- McAlpine D, Jiang D, and Palmer AR (1996b) Interaural delay sensitivity and the classification of low best-frequency binaural responses in the inferior colliculus of the guinea pig. *Hearing Research* **97**:136–152.
- McAlpine D, Jiang D, Shackleton TM, and Palmer AR (1998) Convergent input from brainstem coincidence detectors onto delay-sensitive neurons in the inferior colliculus. *Journal of Neuroscience* **18**:6026–6039.
- McAlpine D, Jiang D, Shackleton TM, and Palmer AR (2000) Responses of neurons in the inferior colliculus to dynamic interaural phase cues: evidence for a mechanism of binaural adaptation. *Journal of Neurophysiology* **83**:1356–1365.
- McAlpine D, Jiang D, and Palmer AR (2001) A neural code for low-frequency sound localization in mammals. *Nature Neuroscience* **4**:396–401.
- McFadden D and Pasanen E (1976) Lateralization at high frequencies based on ITDs. *Journal of the Acoustical Society of America* **59**:634–639.
- Nordlund B (1962) Physical factors in angular localization. *Acta Oto-Laryngologica* **54**: 75–93.
- Nuetzel JM and Hafter ER (1976) Lateralization of complex waveforms: effect of fine structure, amplitude, and duration. *Journal of the Acoustical Society of America* **60**: 1339–1346.
- Oliver DL, Kuwada S, Yin TCT, Haberly L, and Henkel CK (1991) Dendritic and axonal morphology of HRP-injected neurons in the inferior colliculus of the cat. *Journal of Comparative Neurology* **303**:75–100.
- Oliver DL, Winer JA, Beckius GE, and Saint Marie RL (1994) Morphology of GABAergic neurons in the inferior colliculus of the cat. *Journal of Comparative Neurology* **340**:27–42.
- Palmer AR, Rees A, and Caird D (1990) Interaural delay sensitivity to tone and broad band signals in the guinea-pig inferior colliculus. *Hearing Research* **50**:71–86.
- Palmer AR, Jiang D, and McAlpine D (2000) Neural responses in the inferior colliculus to binaural masking level differences created by inverting the noise in one ear. *Journal of Neurophysiology* **84**:844–852.
- Parham K, Zhao HB, and Kim DO (1996) Responses of auditory nerve fibers of the unanesthetized decerebrate cat to click pairs as simulated echoes. *Journal of Neurophysiology* **76**:17–29.
- Parham K, Zhao HB, Ye Y, and Kim DO (1998) Responses of anteroventral cochlear nucleus neurons of the unanesthetized decerebrate cat to click pairs as simulated echoes. *Hearing Research* **125**:131–146.
- Pollak GD, Burger RM, Park TJ, Klug A, and Bauer EE (2002) Roles of inhibition for transforming binaural properties in the brainstem auditory system. *Hearing Research* **168**:60–78.

- Populin LC and Yin TCT (1998) Behavioral studies of sound localization in the cat. *Journal of Neuroscience* **18**:2147–2160.
- Ramachandran R and May BJ (2002) Functional segregation of ITD sensitivity in the inferior colliculus of decerebrate cats. *Journal of Neurophysiology* **88**:2251–2261.
- Ramachandran R, Davis KA, and May BJ (1999) Single-unit responses in the inferior colliculus of decerebrate cats I. Classification based on frequency response maps. *Journal of Neurophysiology* **82**:152–163.
- Reale RA, Brugge JF, and Hind JE (1995) Encoding of stimulus direction by A1 neurons in the cat: effects of sound arriving at different times and from different directions. *Society for Neuroscience Abstracts* **21**:667.
- Rose JE, Gross NB, Geisler CD, and Hind JE (1966) Some neural mechanisms in the inferior colliculus of the cat which may be relevant to localization of a sound source. *Journal of Neurophysiology* **26**:288–314.
- Rubel EW and Parks TN (1988) Organization and development of the avian brainstem auditory system. In: Edelman GM, Gall WE, and Cowan WM (eds). *Auditory Function. Neurobiological Bases of Hearing*. John Wiley & Sons, New York, pp. 3–92.
- Sakitt B (1973) Indices of discriminability. *Nature* **241**:133–134.
- Sally SL and Kelly JB (1992) Effects of superior olivary complex lesions on binaural responses in rat inferior colliculus. *Brain Research* **572**:5–18.
- Schroeder MR (1977) New viewpoints in binaural interactions. In: Evans EF and Wilson JP (eds). *Psychophysics and Physiology of Hearing*. Academic Press, New York, pp. 455–467.
- Seidl AH and Grothe B (2003) Acoustic experience is essential for the development of normal ITD processing. *Association for Research in Otolaryngology Abstracts* **26**:930.
- Semple MN and Kitzes LM (1987) Binaural processing of sound pressure level in the inferior colliculus. *Journal of Neurophysiology* **57**:1130–1147.
- Shackleton TM, McAlpine D, and Palmer AR (2000) Modelling convergent input onto interaural-delay sensitive inferior colliculus neurones. *Hearing Research* **149**:199–215.
- Shackleton TM, Skottun BC, Arnott RH, and Palmer AR (2003) Interaural time difference discrimination thresholds for single neurons in the inferior colliculus of guinea pigs. *Journal of Neuroscience* **23**:716–724.
- Shamma SA, Shen NM, and Gopalaswamy P (1989) Stereausis: binaural processing without neural delays. *Journal of the Acoustical Society of America* **86**:989–1006.
- Shaw EAG (1974) The external ear. In: Keidel WD and Neff WD (eds). *Handbook of Sensory Physiology, Volume V, Part 1: Auditory System, Anatomy, Physiology (Ear)*. Springer-Verlag, Berlin, pp. 455–490.
- Shinn-Cunningham BG, Zurek PM, and Durlach NI (1993) Adjustment and discrimination measurements of the precedence effect. *Journal of the Acoustical Society of America* **93**:2923–2932.
- Skottun BC (1998) Sound localization and neurons. *Nature* **393**:531.
- Smith PH, Joris PX, and Yin TCT (1993) Projections of physiologically characterized spherical bushy cell axons from the cochlear nucleus of the cat: evidence for delay lines to the medial superior olive. *Journal of Comparative Neurology* **331**:245–260.
- Spitzer MW and Semple MN (1991) Interaural phase coding in auditory midbrain: influence of dynamic stimulus features. *Science* **254**:721–724.
- Spitzer MW and Semple MN (1993) Responses of inferior colliculus neurons to time-varying interaural phase disparity: effects of shifting the locus of virtual motion. *Journal of Neurophysiology* **69**:1245–1263.

- Stanford TR, Kuwada S, and Batra R (1992) A comparison of the interaural time sensitivity of neurons in the inferior colliculus and thalamus of the unanesthetized rabbit. *Journal of Neuroscience* **12**:3200–3216.
- Sterbing SJ, D'Angelo WR, Ostapoff E-M, and Kuwada S (2003a) Effects of amplitude modulation on the coding of interaural time differences of low-frequency sounds in the inferior colliculus. I. Response properties. *Journal of Neurophysiology* **90**:2818–2826.
- Sterbing SJ, Hartung K, and Hoffmann K-P (2003b) Spatial tuning to virtual sounds in the inferior colliculus of the guinea pig. *Journal of Neurophysiology* **90**:2648–2659.
- Stillman RD (1971) Characteristic delay neurons in the inferior colliculus of the kangaroo rat. *Experimental Neurology* **32**:404–412.
- Tobias JV and Schubert ED (1959) Effective onset duration of auditory stimuli. *Journal of the Acoustical Society of America* **31**:1595–1605.
- Tobias JV and Zerlin S (1959) Lateralization threshold as a function of stimulus duration. *Journal of the Acoustical Society of America* **31**:1591–1594.
- Tollin DJ (2003) The lateral superior olive: a functional role in sound source localization. *Neuroscientist* **9**:127–143.
- Tollin DJ and Henning GB (1998) Some aspects of the lateralization of echoed sound in man. I. The classical interaural-delay based precedence effect. *Journal of the Acoustical Society of America* **104**:3030–3038.
- Tollin DJ and Yin TCT (2002a) The coding of spatial location by single units in the lateral superior olive of the cat. I. Spatial receptive fields in azimuth. *Journal of Neuroscience* **22**:1454–1467.
- Tollin DJ and Yin TCT (2002b) The coding of spatial location by single units in the lateral superior olive of the cat. II. The determinants of spatial receptive fields in azimuth. *Journal of Neuroscience* **22**:1468–1479.
- Trahiotis C and Hartung K (2002) Peripheral auditory processing, the precedence effect and responses of single units in the inferior colliculus. *Hearing Research* **168**:55–59.
- van Bergeijk WA (1962) Variation on a theme of Békésy: a model of binaural interaction. *Journal of the Acoustical Society of America* **34**:1431–1437.
- van de Par S and Kohlrausch A (1997) A new approach to comparing binaural masking level differences at low and high frequencies. *Journal of the Acoustical Society of America* **101**:1671–1680.
- von Békésy G (1960) *Experiments in Hearing*. McGraw-Hill, New York.
- Wallach H, Newman EB, and Rozenzweig MR (1949) The precedence effect in sound localization. *American Journal of Psychology* **62**:315–336.
- Wenstrup JJ, Fuzessery ZM, and Pollak GD (1988) Binaural neurons in the mustache bat's inferior colliculus. I. Responses of 60-kHz EI units to dichotic sound stimulation. *Journal of Neurophysiology* **60**:1369–1383.
- Wickesberg RE (1996) Rapid inhibition in the cochlear nuclear complex of the chinchilla. *Journal of the Acoustical Society of America* **100**:1691–1702.
- Wickesberg RE and Stevens HE (1998) Responses of auditory nerve fibers to trains of clicks. *Journal of the Acoustical Society of America* **103**:1990–1999.
- Yin TCT (1994) Physiological correlates of the precedence effect and summing localization in the inferior colliculus of the cat. *Journal of Neuroscience* **14**:5170–5186.
- Yin TCT and Chan JCK (1990) Interaural time sensitivity in medial superior olive of cat. *Journal of Neurophysiology* **64**:465–488.

- Yin TCT and Kuwada S (1983) Binaural interaction in low-frequency neurons in inferior colliculus of the cat. III. Effects of changing frequency. *Journal of Neurophysiology* **50**:1020–1042.
- Yin TCT and Kuwada S (1984) Neuronal mechanisms of binaural interaction. In: Edelman GM, Cowan WM, and Gall WE (eds). *Dynamic Aspects of Neocortical Function*. John Wiley & Sons, New York, pp. 263–313.
- Yin TCT, Kuwada S, and Sujaku Y (1984) Interaural time sensitivity of high-frequency neurons in the inferior colliculus. *Journal of the Acoustical Society of America* **76**: 1401–1410.
- Yin TCT, Chan JCK, and Irvine DRF (1986a) Effects of interaural time delays of noise stimuli on low-frequency cells in the cat's inferior colliculus. I. Responses to wideband noise. *Journal of Neurophysiology* **55**:280–300.
- Yin TCT, Chan JCK, and Carney LH (1986b) Effects of interaural time delays of noise stimuli on low-frequency cells in the cat's inferior colliculus. III. Evidence for cross-correlation. *Journal of Neurophysiology* **58**:562–583.
- Zhou X and Jen PH-S (2003) The effect of bicuculline application on azimuth-dependent recovery cycle of inferior collicular neurons of the big brown bat, *Eptesicus fuscus*. *Brain Research* **973**:131–141.
- Zurek PM (1987) The precedence effect. In: Yost WA and Gourevitch G (eds). *Directional Hearing*. Springer-Verlag, New York, pp. 85–105.
- Zwislocki JJ and Feldman RS (1956) Just noticeable dichotic phase difference. *Journal of the Acoustical Society of America* **28**:152–154.

# Chapter 14

## Coding of Communication Sounds in the Inferior Colliculus

CHRISTINE V. PORTFORS AND DONAL G. SINEX

### 1. INTRODUCTION

Many animals, including humans, use species-specific vocalizations to convey biologically relevant information to members of the same species. These communication sounds (referred to as speech sounds in humans and vocalizations in nonhuman animals) are often complex in frequency and time. A fundamental function of the auditory system is to process these complex sounds and impart specific meanings to each sound that are important for communication behaviors. It is unclear how communication sounds are processed and represented in the auditory system, and in particular, it is unclear how these sounds are differentiated from other types of complex sounds. The majority of previous studies have examined processing of communication sounds in the auditory forebrain (Margoliash and Fortune 1992; Ohlemiller et al. 1996; Wang 2000), but the inferior colliculus (IC) likely plays an important role in the neural analysis of communication sounds. We consider the representation of communication sounds in the IC and discuss current evidence for neural specializations in the IC for encoding communication sounds.

The degree to which auditory neurons are specialized in encoding vocalizations has been debated since about 1980 (Symmes 1981). In the auditory cortex, some neurons exhibit specializations that affect the representation of vocalizations so that they are encoded differently than other types of complex sounds. These neural specializations include selectivity for the species' own vocalizations (Wang and Kadia 2001), selectivity for particular types of vocalizations (Wang 2000), reduced responsiveness to spectrally or temporally modified vocalizations (Wang et al. 1995) and experience-dependent plasticity (Wang et al. 1995). Although some specializations are apparent, it is still uncertain whether the information in vocalizations is represented in the auditory cortex by the pattern of spatially distributed activity, by subsets of highly selective, highly specialized neurons ("call detectors"), or by a combination of these mechanisms (Wang 2000; Rauschecker and Tian 2000).

In contrast, peripheral neurons do not exhibit specializations that lead to different representations of speech or vocalizations compared to other types of

complex sounds. Instead, their responses to communication sounds are largely a function of their frequency selectivity, as determined by cochlear mechanics (Robles and Ruggero 2001). For example, auditory nerve fibers respond to complex sounds with an overall temporal pattern that resembles a band-pass-filtered version of the stimulus (Sinex and McDonald 1988).

Consequently, the question arises as to where in the ascending auditory system specializations for encoding communication sounds emerge. Most studies have focused on the representation of communication sounds in the auditory cortex because the forebrain is thought to be inherently more complex than brain stem auditory nuclei (Schreiner 1998; Rauschecker and Tian 2000) and therefore a likely site for the emergence of specializations for encoding complex sounds (Rauschecker et al. 1997).

Few studies have examined responses of IC neurons in mammals to species-specific communication sounds (Aitkin et al. 1994; Klug et al. 2002; Portfors et al. 2002; Šuta et al. 2003). However, the IC, with its ascending inputs from nearly every nucleus in the caudal brain stem (see Chapters 3 to 5), and its descending (see Chapter 8) and intrinsic (see Chapters 2 and 6) connections, is a rich substrate for the integration of spectral information. This convergence of information makes the IC a plausible candidate in the representation and transformation of communication sounds. Understanding the midbrain's role will provide useful information about hierarchical processing of communication sounds in the auditory system, and clarify the species-specific neural mechanisms involved in processing of vocalizations. Moreover, studies of IC neural responses to speech sounds in nonhuman animals will increase understanding of how speech sounds are processed in the absence of linguistic experience.

A brief summary of the production and acoustic characteristics of communication sounds is followed by a review of studies of the representation of communication sounds in the IC. Contemporary studies in mammals are emphasized and the representation of human speech is considered. Finally, the prospects for future studies are discussed.

## 2. ACOUSTIC PROPERTIES OF SPEECH AND VOCALIZATIONS

For simplicity, the term *vocalizations* is used to refer to the communication sounds produced by nonhuman animals, and *speech* is reserved for human sounds. The distinction is made on acoustic grounds alone. Questions about how animals use vocalizations, and the extent to which the use of vocalizations resembles humans' use of speech, are beyond the scope of this chapter and are treated elsewhere (Hauser et al. 2002).

The "source-filter theory" (Fant 1960) provides a useful framework for understanding the production and acoustic characteristics of speech and vocaliza-



tions. When these sounds are produced, air from the lungs is forced through the larynx, providing a source of energy, and resonances of the supraglottal vocal tract filter the source spectrum and shape the sound spectrum. Vocal tract resonant frequencies are determined by its physical characteristics, and by the locations and elasticity of constrictions produced primarily by the tongue, teeth, and lips.

Forcing air through vocal folds that are held close together causes them to open and close repetitively, with each opening producing a “glottal pulse.” A sequence of glottal pulses produces a quasi-periodic waveform, and the rate at which glottal pulses occur determines the fundamental frequency ( $F_0$ ) of the sound. In speech,  $F_0$  varies between 100 and 250 Hz (Peterson and Barney 1952) whereas  $F_0$  varies much more in vocalizations, within and across species. Values range from several hundred Hertz (guinea pig: Šuta et al. 2003; macaque: Rendall et al. 1998) to several kHz (marmoset: Wang et al. 1995) to several tens of kiloHertz (mustached bat: Kanwal et al. 1994; mouse: Portfors 2003). Varying subglottal air pressure and the position and tension of the vocal folds causes changes in  $F_0$  (Borden et al. 2003). Modulations of  $F_0$  can be minimal (Fig. 14.1A) or extensive (Fig. 14.1B). The line spectrum of the glottal waveform is shaped by the transfer function of the upper vocal tract, so that the sound that is radiated from the lips may have a distinctive pattern of spectral peaks and valleys. The peaks are produced by resonances of the vocal tract, called “formants.” In speech, the locations of formants vary (Fig. 14.1A) and provide powerful cues that contribute to phonetic judgments.

Nonhuman animals have much less control over the configuration of the vocal tract (Schön Ybarra 1996), so formants in their vocalizations are relatively static. However, nonhuman animals may still use the information in formants to identify individuals and body size of conspecifics (Fitch 1997).

If the vocal folds are held partially open and air is forced through a constricted vocal tract, the resulting sound will be aperiodic and may be perceived as noisy. Aperiodic segments occur in speech (Fig. 14.1A) and also in vocalizations (Kanwal et al. 1994). As with voiced sounds, the shape of the emitted spectrum reflects the spectrum of the aperiodic source and the transfer function of the vocal tract.

Speech and vocalizations also vary over time. In speech, temporal information arises because the configuration of the vocal tract rarely reaches a steady state, and one consequence is that the amplitude envelope fluctuates at a rate typically between 5 and 50 Hz (Rosen 1992). During vocalizations, vocal tract configuration is more stable, but  $F_0$  may change dramatically. In some vocalizations, short spectrally similar segments are repeated in bursts (Kanwal et al. 1994) so the temporal envelope may fluctuate at a rate similar to that of speech. Spectral and temporal changes are linked because both are natural consequences of the same movements of the articulators and changes in voicing.




Figure 14.1. Spectrograms of speech and of nonhuman vocalizations. **(A)** Spectrogram of the sentence “Gus saw pine trees and redwoods.” Acoustic properties of representative consonants and vowels can be seen. For example, the fricative consonant, /s/, produced with aperiodic excitation of the vocal tract, is characterized by a broad peak at about 5 kHz (150 to 250 ms). During vowels, formant frequencies occur at lower frequencies, and formant transitions are usually visible between segments (for example, F2, F3, and F4 rise between 800 and 900 ms, preceding the vowel /i/ in “tree”). During voiced segments, individual glottal pulses appear as vertical striations. In speech, formant frequencies vary but F0 is relatively stable, as indicated by the approximately equal spacing between glottal pulses. In vocalizations, the opposite is usually true. The sentence material is from the DARPA TIMIT Acoustic-Phonetic Continuous Speech Corpus. **(B)** Two mustached bat communication calls. The frequencies in the calls are much higher than human speech, but the calls are complex in frequency and time.

### 3. REPRESENTATION OF VOCALIZATIONS IN THE INFERIOR COLLICULUS

#### 3.1. RESPONSES TO COMPLEX STIMULI THAT MAY UNDERLIE PROCESSING OF VOCALIZATIONS

Very few studies have explicitly examined neural responses to communication sounds in the IC of mammals, but there are many examples of responses to

complex acoustic stimuli in the IC that may underlie processing of vocalizations. Single neurons demonstrate complex response properties to spectrally and temporally complex acoustic stimuli that involve amplitude modulation (Rees and Møller 1987; Sinex et al. 2002), frequency modulation (Schuller 1979), duration (Casseday et al. 1994), and the temporal interval between stimuli (Walton et al. 1998). More than half of the IC neurons of the mustached bat (*Pteronotus p. parnellii*) show nonlinear facilitation or inhibition to the combination of two sounds in different frequency bands (Portfors and Wenstrup 1999). These combination-sensitive neurons are a distinct class (Portfors and Wenstrup 1999) and were initially described as specialized for echolocation behaviors. However, they are now thought to play a more general role in processing of spectrally and temporally complex sounds (Leroy and Wenstrup 2000).

### 3.2. RESPONSES TO VOCALIZATIONS IN THE CENTRAL NUCLEUS OF THE INFERIOR COLLICULUS

Two studies have suggested that some IC neurons in bats respond selectively to species-specific vocalizations. It is useful to distinguish between “selectivity” and “specialization” when discussing neural responses to communication sounds. Selectivity implies that the neuronal response can be reasonably well predicted based on the same cell’s response to simpler sounds, whether presented alone or in combination with other sounds. Specialized responses, in contrast, are not well predicted. In the IC, neural responses to vocalizations can be fairly well predicted based on excitatory and inhibitory frequency response areas.

Excitatory frequency tuning curves of neurons in the IC of the Mexican free-tailed bat (*Tadarida brasiliensis mexicana*) were measured and their responses to 10 different vocalizations assessed (Klug et al. 2002). These responses were not well predicted by the excitatory tuning curves because most neurons responded to a subset of the vocalizations even though most of those to which the neurons were unresponsive had energy in the neuron’s excitatory frequency response area. However, responses to vocalizations were much better predicted when inhibition was blocked by iontophoretic application of bicuculline or bicuculline and strychnine, which caused the neurons to respond to many vocalizations and eliminated their selectivity. These data suggest that IC neurons show selectivity to vocalizations, and that one mechanism underlying this selectivity is inhibition.

Selective responses in the IC of the mustached bat IC were also found with a two-tone paradigm to assess excitatory and inhibitory frequency response areas of single neurons, and then predict their responses to fourteen different mustached bat communication calls (Portfors et al. 2002). The benefit of a two-tone paradigm is that the sound frequencies that elicit inhibition are determined, and the neural response to a communication call can be predicted, based on whether the call has energy in the frequency range that elicits an inhibitory response. To test the frequencies of inhibition, one sound is presented at the frequency that elicits the best response from the neuron while a second sound that varies in frequency is presented (Portfors and Wenstrup 2002). Neurons with inhibitory

frequency response areas were selective to certain communication calls whereas those without inhibitory frequency response areas responded to many of the communication calls.

Neurons with inhibitory response areas too remote from the excitatory frequency area to be considered as side-band inhibition present an interesting mechanism for suppressing responses to certain vocalizations. For example, consider a neuron with a narrow excitatory frequency-tuning curve centered at 50 kHz, and with two inhibitory frequency regions that flank the excitatory response area, and a third inhibitory region centered at 19 kHz (Portfors et al. 2002). Without knowing the areas of inhibition, we would predict that this neuron would respond to all calls that had energy in the 50-kHz regions. However, this neuron actually responded to only 1 of 14 vocalizations even though other calls had energy in the cell's excitatory tuning curve. The vocalization to which the neuron responded was a three harmonic, frequency-modulated call with the second harmonic passing through the 50-kHz range. The fundamental frequency of the call never entered the low-frequency inhibitory region of the neuron. The other vocalizations either did not have energy in the excitatory region of the neuron or had energy in the low-frequency inhibitory region (18- to 23-kHz range) that suppressed the excitatory response. Therefore, inhibitory frequency response areas are effective in providing for selectivity to vocalizations in the mustached bat IC. To understand the mechanisms underlying neural selectivity to communication calls, then, it is necessary to understand how both the excitatory and inhibitory frequency response areas of neurons are organized and interact.

However, excitatory and inhibitory frequency tuning cannot explain selectivity in all neurons. Another mechanism that may provide for selectivity to vocalizations is combination-sensitive facilitation (Portfors et al. 2002) in which a neuron responded only to one call (a multiharmonic, upward frequency modulated sweep) regardless of intensity (Fig. 14.2A, B). The high selectivity could be explained by the neuron's combination-sensitive facilitatory response to pairs of tones. The neuron responded weakly to tones near 57 kHz, was unresponsive to any low-frequency tones, and responded well to the combination of a 57-kHz and a 29-kHz tone presented simultaneously (Fig. 14.2B). The vocalization to which this neuron selectively responded contained energy in both the 57-kHz and 29-kHz bandwidths and these frequency components overlapped in time. The cardinal features of this response were the weak or nonexistent response to the individual components (tones or call elements) and the robust facilitatory response to the combination of the frequency components. These features caused the neuron to respond poorly or not at all to vocalizations that had energy in only one of the frequency bands and to respond well to the vocalizations with energy in at least two of the frequency bands, as long as the spectral components of the vocalization were in the proper temporal relationship. Thus, the combination-sensitive properties of some mustached bat IC neurons provided for selectivity to particular vocalizations.

Interestingly, mouse IC neurons also show inhibitory and facilitatory combination sensitivity (Portfors 2003) suggesting that neural selectivity to species-specific vocalizations may occur in species other than bats. In contrast, however,




Figure 14.2. Responses of an IC neuron illustrating that combination sensitivity can explain some neuronal selectivity to social vocalizations. **(A)** Raster plot of the neuron's response to its preferred vocalization (a three-harmonic, upward frequency modulated sweep as shown in the figure). **(B)** Raster plots show that such response selectivity may be explained by the neuron's response to single tones and combinations of tones. The cell responded weakly to a 57.1-kHz tone alone, did not respond to a 29 kHz tone, and was strongly facilitated by the combination of the tones. The facilitatory frequency tuning explains the neuron's selectivity to vocalizations because only one vocalization had sufficient energy in both of the facilitatory frequency bands in the appropriate temporal relationship to elicit a strong response. *Black bars* under raster plots indicate onset time and duration of stimulus.

are other findings that guinea pig IC neurons showed little selectivity to four vocalizations (Šuta et al. 2003). Most neurons responded to all the vocalizations, but they did show some selectivity in that they preferred vocalizations to pure tones or noise bursts. The temporal firing patterns matched the sound envelope of the vocalization, suggesting that the neurons encoded the spectrotemporal aspects of the vocalizations. The reasons for different findings between species are unknown.

An important aspect of encoding of vocalizations in the IC is that individual neurons showed different selectivity to vocalizations both in terms of how many

vocalizations elicited a response from each neuron and the particular vocalizations eliciting a response. It has been suggested that because neurons show different types of selectivity, each vocalization should evoke a unique pattern of activity across the population of IC neurons, and this activation should be greater than if the individual IC neurons showed no selectivity (Klug et al. 2002).

### *3.3. RESPONSES TO VOCALIZATIONS IN OTHER INFERIOR COLLICULUS SUBDIVISIONS*

When the responses of neurons in all IC subdivisions of the cat IC to four different vocalizations were recorded, many cells responded more vigorously to vocalizations than to tones, in particular neurons in the external nucleus (EN), but they were not selective for individual vocalizations. Instead, the increased response to vocalizations appeared to reflect a general preference for broadband sounds (Aitkin et al. 1994). The authors proposed that the EN neurons strongly preferring vocalizations might be part of a premotor pathway involved in the initiation of vocalization. Further studies will be required to substantiate this idea.

## 4. REPRESENTATION OF SPEECH IN THE INFERIOR COLLICULUS

Few reports of the representation of selected speech sounds by IC neurons in the anesthetized cat are available (Watanabe and Sakai 1978; Delgutte et al. 1998). In both studies, envelope information was prominently represented by IC neurons. In one study the discharge rate of auditory nerve fibers and cochlear nucleus neurons generally followed the envelope of the speech in the frequency band near the neurons' characteristic frequencies (CFs). In contrast, IC neurons responded phasically to the same stimulus, with activity restricted to the onsets of syllables and the release bursts of stop consonants. Quantitative comparisons of spectral properties of the speech to discharge patterns were not reported, but the findings suggest that IC neurons provide a good representation of temporal cues (Delgutte et al. 1998).

A series of studies of the representation of consonant–vowel syllables differing in voice-onset time (VOT) has been carried out in the chinchilla IC (Chen et al. 1996; Chen and Sinex 1999; Sinex and Chen 2000). VOT is the delay between the release of a stop consonant and the onset of voicing, and it determines the identity of stop consonants such as /d/ and /t/ that are produced at the same place of articulation (Lisker and Abramson 1964). Small changes in VOT near the /d/-/t/ perceptual boundary are highly audible, but larger changes away from the boundary are not (Abramson and Lisker 1970). This finding, coupled with the observation that chinchillas exhibit psychophysical acuity for VOT that is quantitatively similar to that of humans (Kuhl 1981), led to the hypothesis that certain sounds are advantageous for vocal communication be-

cause they naturally elicit distinctive responses in the mammalian auditory system (Kuhl 1986). Studies of the neural representation of VOT continua were undertaken in the chinchilla to determine how these sounds are processed in the absence of confounding linguistic experience. That is, the goal was explicitly to examine nonspecialized processing of acoustic cues for consonant voicing.

Examples of the representation of syllables differing in VOT are shown in Fig. 14.3. The top row shows waveforms of syllables with VOTs of 30 and 70 ms, heard by humans as /da/ and /ta/, respectively. The responses of an IC neuron to two syllables (Fig. 14.3, bottom row) can be compared to those of an auditory nerve fiber with the same CF, 0.8 kHz, to the same syllables (Fig. 14.3, middle row). The IC neuron responded phasically at the onset of voicing in each syllable and also exhibited a less obvious response to the steady-state vowel. In contrast,




Figure 14.3. Responses to syllables differing in VOT. VOT is indicated in each panel by a vertical dashed line. (A) Waveform of a synthesized syllable with VOT of 30 ms. The syllable is heard as /da/. (B) Discharge pattern of a chinchilla auditory nerve fiber to the syllable shown in A. Fiber 8713-66; CF: 0.8 kHz; threshold: 11 dB SPL. (C) Discharge pattern of a chinchilla IC neuron to the syllable shown in A. Neuron 9311-78; CF: 0.8 kHz; threshold: 25 dB SPL. (D) Waveform of a synthesized syllable with VOT = 70 ms. The syllable is heard as /ta/. (E) Discharge pattern of the same auditory nerve fiber shown in (B), for the syllable shown in (D). (F) Discharge pattern of the same IC neuron shown in (C), for the syllable shown in (D).

the auditory nerve fiber discharged additional spikes at the release of the syllable, and higher sustained discharge rates throughout the vowel. Clearly, the representation of the onset of voicing was less precisely marked by the peripheral neuron.

The characteristic discharge patterns of individual IC neurons to VOT syllables reflect the acoustic properties of the syllable but are strongly affected by characteristics of individual neurons such as frequency selectivity, threshold, the temporal pattern of responses to steady-state tones, and binaural sensitivity (Chen et al. 1996; Chen and Sinex 1999). Because neurons with similar CFs differ in these other characteristics, the same syllable may elicit a range of discharge patterns, and IC neurons do respond in complex ways to changes in VOT. For example, some low-CF neurons represent VOT nonmonotonically, showing higher discharge rates for the syllables with intermediate VOT for which psychophysical acuity would be highest (Chen and Sinex 1999). Discharge rates of individual auditory nerve fibers do not vary as strongly with VOT (Sinex and McDonald 1988), suggesting that this difference is a consequence of more complex spectral integration at the level of the IC.

## 5. CONCLUSIONS

The studies reviewed here are consistent with the view that the processing of communication sounds is organized in a hierarchical fashion (Rauschecker et al. 1997). Responses observed in the IC are generally more complex than those in the peripheral auditory system (Sinex and McDonald 1988; Chen et al. 1996; Delgutte et al. 1998) but appear less complex than the cortical representation (Ohlemiller et al. 1996; Esser et al. 1997). There is also a suggestion that, in the bat and perhaps in other species with stereotypical vocalizations, some IC neurons have intricate patterns of spectral and temporal integration that are especially well matched to the acoustic properties of the species-specific vocalizations (Portfors et al. 2002; Portfors 2003). The generality and significance of combination sensitivity for the processing of vocalizations in other species should be evaluated in future studies to elucidate the extent of common neural mechanisms across nonhuman species.

In nonhuman vocalizations, the range of sounds that may occur is far more restricted than is the case with speech. When individual vocalizations are somewhat stereotyped, and there are limits to the number of different sounds that must be processed, strategies such as combination sensitivity may be very efficient. Although there have been suggestions that combination sensitivity may contribute to the processing of speech sounds (Suga 1996; Sussman et al. 1998), it is not clear that the same strategy would be as effective for representing the phonetic elements of speech in a sound-selective way. This is because the range of sounds that occur in speech is enormous. There are only modest constraints on the possible relationships between elements such as formant frequencies for



an individual talker, and even greater variation exists across talkers and languages. Thus, it seems unlikely that combination-sensitive neurons that can encode each potential variation exist, at least at the level of the IC. If some form of combination sensitivity does exist in higher structures, at or beyond the primary auditory cortex, it seems likely that the properties of those neurons might be language dependent. That is, experience with one's native language may alter the selectivity of neurons (cf. Wang et al. 1995), such that some sounds are encoded more effectively and some less effectively. Previously we noted that studies of the representation of speech in nonhuman animals are most useful for addressing the role of mechanisms that are not species-specific and not language dependent. Thus, if combination-sensitive processing does occur in the human, imaging studies of extraordinary resolution will likely be required to reveal them.

## 6. GENERAL ISSUES AND FUTURE CONSIDERATIONS

### 6.1. SPECIES-SPECIFIC DIFFERENCES AND SIMILARITIES

Selectivity to species-specific vocalizations in the mammalian IC has been found only in bats. Responses in the cat IC to vocalizations are more complex than responses to tones, but there is little evidence of specific neuronal selectivity to particular vocalizations (Aitkin et al. 1994). Similarly, responses in the guinea pig IC show little selectivity to vocalizations (Šuta et al. 2003). This may be a valid species-specific difference or may reflect the choice of vocalizations presented or the experimental paradigm. Only further experiments in other species will resolve these questions. An important consideration, however, is the choice of vocalizations used as stimuli. To dissect critically how the IC encodes species-specific vocalizations and determine the extent of neural selectivity and/or specializations, a full array of vocalizations should be explored as stimuli. This includes all the different categories of calls in a species' repertoire and also sufficient calls per category to counterbalance the variability seen in the repertoire of calls emitted, both within and between individual animals. Presenting only a few "exemplar" calls as representative stimuli may provide a misleading impression of both neural selectivity and specializations in the IC.

As shown in both species of bats, excitatory and inhibitory frequency response areas are important in providing for selectivity to vocalizations, and it is expected that these features of frequency tuning will also be important in other species (Portfors 2003). The importance of combination sensitivity as a mechanism for selectivity in species other than the mustached bat is unclear. Although there is evidence of nonlinear interactions in the response properties of cat (Escabí and Schreiner 2002) and Mexican free-tailed bat (Klug et al. 2002) IC neurons, the appropriate tests to investigate combination sensitivity have not been done. In the mouse IC some neurons are combination sensitive (Egorova et al. 2001;

Portfors 2003), but their role in selectivity for vocalizations is unclear. Certainly assessing the excitatory, inhibitory, and facilitatory frequency response areas of IC neurons on a more secure comparative basis will increase our understanding of the mechanisms underlying midbrain selectivity to species-specific vocalizations.

## 6.2. SELECTIVITY FOR VOCALIZATIONS VS. SPECIALIZATION FOR VOCALIZATIONS

The evidence supporting the idea of neural selectivity to communication calls in the IC suggests that the neural responses to communication calls could be predicted reasonably well from their discharges to simpler stimuli. This implies that the responses are selective, but not specialized, for encoding species-specific vocalizations.

There is little consensus about the degree to which central auditory neurons are specialized for vocalizations, and this is partly a reflection of the difficulty in determining what a nonspecialized response should look like. Neurons in the IC certainly exhibit selectivity for vocalizations, but to claim that the neuron is specialized, it is first necessary to establish which response patterns could have been predicted given adequate information about the same neuron's responses to simpler sounds. Thus, "adequate" information should be obtained when assessing responses to complex sounds. This could include measures of (at least) the form of the temporal discharge pattern elicited by a pure tone, the distribution of responses to tone pairs, and temporal interactions such as the spectrotemporal receptive field (STRF, Escabí and Schreiner 2002). For example, some IC neurons show excitatory or inhibitory combination sensitivity that affects their responses to vocalizations. If combination sensitivity had not been documented, the response of such neurons might mistakenly have appeared to express specialization for vocalizations rather than selectivity. Similar misinterpretations could potentially be made for neurons with excitatory–inhibitory temporal interactions of the type revealed by the STRF. The current perspective suggests that IC neurons are selective for vocalizations, but not specialized. If stronger claims for specialization are to be made in the future, they should be supported with correspondingly extensive and rigorous measurements with simpler sounds.

## Abbreviations

CF	characteristic frequency
F0	fundamental frequency
IC	inferior colliculus
STRF	spectrotemporal receptive field
VOT	voice-onset time

## Acknowledgments

Preparation of this chapter was supported by NIDCD Grants DC04733 (C.V.P.) and DC00341 (D.G.S.). The authors thank Drs. Susan Nittrouer and Andrew Lotto for comments on a preliminary version of the manuscript.

## References

- Abramson AS and Lisker L (1970) Discriminability along the voicing continuum: cross-language tests. *Proceedings of the Sixth International Congress of Phonetic Sciences*. Academia Publishing House of the Czechoslovak Academy of Sciences, Prague, pp. 70.
- Aitkin L, Tran L, and Syka J (1994) The responses of neurons in subdivisions of the inferior colliculus of cats to tonal, noise and vocal stimuli. *Experimental Brain Research* **98**:53–64.
- Borden GJ, Harris KS, and Raphael LJ (2003) *Speech Science Primer: Physiology, Acoustics, and Perception of Speech*. Lippincott Williams & Wilkins, Philadelphia.
- Casseday JH, Ehrlich D, and Covey DE (1994) Neural tuning for sound duration: role of inhibitory mechanisms in the inferior colliculus. *Science* **264**:847–850.
- Chen GD and Sinex DG (1999) Effects of interaural time differences on the responses of chinchilla inferior colliculus neurons to consonant-vowel syllables. *Hearing Research* **138**:29–44.
- Chen GD, Nuding SC, Narayan SS, and Sinex DG (1996) Responses of single neurons in the chinchilla inferior colliculus to consonant-vowel syllables differing in voice onset time. *Auditory Neuroscience* **3**:179–198.
- Delgutte B, Hammond BM, and Cariani PA (1998) Neural coding of the temporal envelope of speech: relation to modulation transfer functions. In: Palmer AR, Rees A, and Summerfield Q (eds). *Psychophysical and Physiological Advances in Hearing*. Whurr, London, pp. 595–603.
- Egorova M, Ehret G, Vartanian I, and Esser K-H (2001) Frequency response areas of neurons in the mouse inferior colliculus. I. Threshold and tuning characteristics. *Experimental Brain Research* **140**:145–161.
- Escabí MA and Schreiner CE (2002) Nonlinear spectrotemporal sound analysis by neurons in the auditory midbrain. *Journal of Neuroscience* **22**:4114–4131.
- Esser K-H, Condon CJ, Suga N, and Kanwal JS (1997) Syntax processing by auditory cortical neurons in the FM-FM area of the mustached bat *Pteronotus parnellii*. *Proceedings of the National Academy of Sciences of the United States of America* **94**:14019–14024.
- Fant G (1960) *Acoustic Theory of Speech Production*. Mouton, The Hague.
- Fitch WT (1997) Vocal tract length and formant frequency dispersion correlate with body size in rhesus macaques. *Journal of the Acoustical Society of America* **102**:1213–1222.
- Hauser MD, Chomsky N, and Fitch WT (2002) The faculty of language: what is it, who has it, and how did it evolve? *Science* **298**:1569–1579.
- Kanwal JS, Matsumura S, Ohlemiller K, and Suga N (1994) Analysis of acoustic elements and syntax in communication sounds emitted by mustached bats. *Journal of the Acoustical Society of America* **96**:1229–1254.

- Klug A, Bauer EE, Hanson JT, Hurley L, Meitzen J, and Pollak GD (2002) Response selectivity for species-specific calls in the inferior colliculus of Mexican free-tailed bats is generated by inhibition. *Journal of Neurophysiology* **88**:1941–1954.
- Kuhl PK (1981) Discrimination of speech by nonhuman animals: basic auditory sensitivities conducive to the perception of speech-sound categories. *Journal of the Acoustical Society of America* **70**:340–349.
- Kuhl PK (1986) Theoretical contributions of tests on animals to the special-mechanisms debate in speech. *Experimental Biology* **45**:233–265.
- Leroy SA and Wenstrup JJ (2000) Spectral integration in the inferior colliculus of the mustached bat. *Journal of Neuroscience* **20**:8533–8541.
- Lisker L and Abramson AS (1964) A cross-language study in initial stops: acoustical measurements. *Word* **20**:384–422.
- Margoliash D and Fortune ES (1992) Temporal and harmonic combination-sensitive neurons in the zebra finch's HVC. *Journal of Neuroscience* **12**:4309–4326.
- Ohlemiller KK, Kanwal JS, and Suga N (1996) Facilitative responses to species-specific calls in cortical FM-FM neurons of the mustached bat. *NeuroReport* **7**:1749–1755.
- Peterson GE and Barney HL (1952) Control methods used in a study of the vowels. *Journal of the Acoustical Society of America* **24**:175–184.
- Portfors CV (2003) Spectral and temporal response properties of single neurons in the inferior colliculus of the mouse. *Association for Research in Otolaryngology Abstracts* **26**:1540.
- Portfors CV and Wenstrup JJ (1999) Delay-tuned neurons in the inferior colliculus of the mustached bat: implications for analyses of target distance. *Journal of Neurophysiology* **82**:1326–1338.
- Portfors CV and Wenstrup JJ (2002) Excitatory and facilitatory frequency response areas in the inferior colliculus of the mustached bat. *Hearing Research* **168**:131–138.
- Portfors CV, Gans DP, and Wenstrup JJ (2002) Responses to communication calls in the inferior colliculus of the mustached bat. *Association for Research in Otolaryngology Abstracts* **25**:143.
- Rauschecker JP and Tian B (2000) Mechanisms and streams for processing of “what” and “where” in auditory cortex. *Proceedings of the National Academy of Sciences of the United States of America* **97**:11800–11806.
- Rauschecker JP, Tian B, Pons T, and Mishkin M (1997) Serial and parallel processing in rhesus monkey auditory cortex. *Journal of Comparative Neurology* **382**:89–103.
- Rees A and Møller AR (1987) Stimulus properties influencing the responses of inferior colliculus neurons to amplitude-modulated sounds. *Hearing Research* **27**:129–143.
- Rendall D, Owren MJ, and Rodman PS (1998) The role of vocal tract filtering in identity cueing in rhesus monkey (*Macaca mulatta*) vocalizations. *Journal of the Acoustical Society of America* **103**:602–614.
- Robles L and Ruggero MA (2001) Mechanics of the mammalian cochlea. *Physiological Reviews* **81**:1305–1352.
- Rosen S (1992) Temporal information in speech: acoustic, auditory and linguistic aspects. *Philosophical Transactions of the Royal Society of London, series B, Biological Sciences* **336**:367–373.
- Schön Ybarra MA (1996) A comparative approach to the non-human primate vocal tract: implications for sound production. In: Zimmermann E, Newman JD, and Jürgens U (eds). *Current Topics in Primate Vocal Communication*. Plenum Press, New York, pp. 185–198.

- Schreiner CE (1998) Spatial distribution of responses to simple and complex sounds in the primary auditory cortex. *Audiology and Neurootology* **3**:104–122.
- Schuller G (1979) Coding of small sinusoidal frequency and amplitude modulations in the inferior colliculus of the CF-FM bat, *Rhinolophus ferrumequinum*. *Experimental Brain Research* **34**:117–132.
- Sinex DG and Chen GD (2000) Neural responses to the onset of voicing are unrelated to other measures of temporal resolution. *Journal of the Acoustical Society of America* **107**:486–495.
- Sinex DG and McDonald LP (1988) Average discharge rate representation of voice onset time in the chinchilla auditory nerve. *Journal of the Acoustical Society of America* **83**:1817–1827.
- Sinex DG, Henderson J, Li H, and Chen GD (2002) Responses of chinchilla inferior colliculus neurons to amplitude-modulated tones with different envelopes. *Journal of the Association for Research in Otolaryngology* **3**:390–402.
- Suga N (1996) Basic acoustic patterns and neural mechanisms shared by humans and animals for auditory perception: a neuroethologists's view. In: Ainsworth A and Greenberg S (eds). *Auditory Basis of Speech Perception*. Keele University, Staffordshire, pp. 31–38.
- Sussman HM, Fruchter D, Hilbert J, and Sirosh J (1998) Linear correlates in the speech signal: the orderly output constraint. *Behavioral and Brain Sciences* **21**:241–299.
- Šuta D, Kvašňák E, Popelář J, and Syka J (2003) Representation of species-specific vocalizations in the inferior colliculus of the guinea pig. *Journal of Neurophysiology* **90**:3794–3808.
- Symmes D (1981) On the use of natural stimuli in neurophysiological studies of audition. *Hearing Research* **4**:203–214.
- Walton JP, Frisina RD, and O'Neill WE (1998) Age-related alteration in processing of temporal sound features in the auditory midbrain of the CBA mouse. *Journal of Neuroscience* **18**:2764–2776.
- Wang X (2000) On cortical coding of vocal communication sounds in primates. *Proceedings of the National Academy of Sciences of the United States of America* **97**:11843–11849.
- Wang X and Kadia SC (2001) Differential representation of species-specific primate vocalizations in the auditory cortices of marmoset and cat. *Journal of Neurophysiology* **86**:2616–2620.
- Wang X, Merzenich MM, Beitel R, and Schreiner CE (1995) Representation of a species-specific vocalization in the primary auditory cortex of the common marmoset: temporal and spectral characteristics. *Journal of Neurophysiology* **74**:2685–2706.
- Watanabe T and Sakai H (1978) Responses of the cat's collicular auditory neuron to human speech. *Journal of the Acoustical Society of America* **64**:333–337.

# Chapter 15

## Acoustic Behavior and Midbrain Function

TOM C.T. YIN AND BRADFORD J. MAY

### 1. INTRODUCTION

This chapter provides an overview of the role of the central nucleus of the inferior colliculus (ICC) in auditory behavioral tasks. This question must be addressed to understand the function of this large structure. The substantial literature on the anatomy, physiology, and pharmacology reflect its importance as a nexus for ascending and descending auditory information processing.

A primary goal is to relate these anatomical and physiological properties of the ICC to acoustically dependent animal behavior. For this, studies of the ICC that incorporate behavior are critical and must address the relationship between neuronal activity and behavior. Given the difficulty in quantifying behavior in animals and recording neural activity in humans, many early studies related human psychophysics to recordings from anesthetized animals (Siebert 1970). This is not an ideal combination, as the behavioral “function” of a neuron in the awake brain and anesthetized brain is different. These ambiguities are compounded when the animal’s auditory behavioral ability is unknown.

Traditionally, the behavioral function of neural tissue has been inferred from the effects of activation (electrical stimulation in intact animals) or inactivation (lesions or cooling) on neurons. Such manipulations, although useful, are constrained by the virtual impossibility of electrically stimulating neurons with naturalistic stimuli and because lesions cause loss of neurons but also elicit compensatory responses in the brain.

The development of techniques to record from neurons in awake, behaving animals, particularly in the visual system, has enabled striking advances in relating neural activity with behavior. Major advances in understanding include the neural basis for saccadic eye movements (Wurtz 1996; Sparks 2002), the linkage between single-cell responses and visual motion perception (Parker and Newsome 1998), the neuronal substrate for directed visual attention (Maunsell and Cook 2002), and neuronal processing of intended movements (Andersen and Buneo 2002). Studies in the somatosensory system have related single-neuron discharges to the perception of vibration (LaMotte and Mountcastle 1975; Vallbo and Johansson 1984). Deciphering the relationship

between perceptual state and neural activity is a central goal of modern neurobiology.

Unfortunately, the auditory community has lagged behind the visual and the somatosensory communities in relating neural activity and behavior in animals. It was noted that “Remarkably, we have been unable to identify studies of the auditory pathways in which neural signals have been measured at the same time as the subject is performing at near-threshold levels in a detection or discrimination task” (Parker and Newsome 1998, p. 260). The few behavioral studies of the ICC and its input systems from a functional point of view limit our understanding of its function.

This chapter summarizes experiments that have been directed toward the functional organization and processing in the ICC with respect to behavior, sound localization, and on the coding of localization cues. Classically, the duplex theory posits that the important cues for sound localization are interaural time and level disparities (Rayleigh 1907). Since about 1980 it has become clear that additional cues are necessary for localization in elevation, as can be readily appreciated by considering localization of sources that vary in elevation along the midsagittal plane, where interaural cues are minimal. The spectral filtering by the head, shoulders, and external ears is now considered important for localization in elevation (May 2000) while the interaural cues are critical for azimuthal localization (Yin 2002).

## 2. SOUND LOCALIZATION AND THE INFERIOR COLLICULUS

### 2.1. CONTRALATERAL ENCODING OF SPACE

Most mammalian sensory and motor systems are organized in a contralateral fashion: the left brain receives input from the right side of the body and extrapersonal space, and controls the muscles on the right side of the body. This contralateral representation is documented in the visual, somatosensory and skeletal motor systems, for example, the precise partial decussation of fibers at the optic chiasm, such that the right side of visual space is projected to the left lateral geniculate body and visual cortex, is well known. That the auditory system follows a similar contralateral representation is less obvious but not unexpected.

However, the auditory system differs markedly from the visual and somatosensory systems in the manner in which space is encoded. In vision and touch the spatial location of the stimulus is naturally encoded by receptor location: the retinal map of the visual field and the somatosensory receptors in the skin are naturally topographically arranged. Retinotopic and somatotopic organization is preserved in both systems by the anatomical projections at successive nuclei to the sensory cortices, and in both systems there is a precise projection of the appropriate fibers across the midline such that the cortex has a topographical representation of the contralateral visual field and body.

On the other hand the auditory system initially encodes frequency, not spatial location. The auditory system must compute the location of the stimulus based on the interaural cues of time, intensity, as well as spectral cues. The result of this neuronal computation also provides for a contralateral representation of auditory space.

## 2.2. CODING OF INTERAURAL LOCALIZATION CUES

### 2.2.1. Interaural Time Disparities

It has been known since about 1900 that the interaural time (ITD) and interaural level differences (ILD) of sounds in free field provide the critical cues for localization of sound sources along the horizontal dimension (Rayleigh 1907). Numerous studies have shown that ITDs are first encoded by cells in the medial superior olive (MSO; Goldberg and Brown 1969; Yin and Chan 1990), which receives bilateral input from the spherical bushy cells of the anteroventral cochlear nucleus (AVCN; Fig. 15.1A; Warr 1966; Tolbert et al. 1982; Cant and Casseday 1986; Smith et al. 1993). Cells in the MSO project to the ICC so that ITD sensitivity is also a common feature of low-frequency ICC cells (Rose et al. 1966; Kuwada and Yin 1983). The physiological properties of these cells are described in more detail elsewhere (see Chapter 13) (Yin 2002).

For pure tone stimuli, which were used by most of the early studies, auditory nerve fibers preserve timing information in the stimulus by discharging over a limited range of phase angles of the input sinusoid, a feature usually called phase locking. Phase locking in mammals is limited to low frequencies <3 to 4 kHz (Johnson 1980) and it is preserved, indeed, enhanced, by AVCN spherical bushy cells (Joris et al. 1994) so that the bilateral inputs arriving at the MSO are highly phase-locked. A fundamental feature of binaural ITD-sensitive MSO cells is that they behave like coincidence detectors in accord with the original model (Jeffress 1948): they respond maximally when the inputs from the left and right sides coincide in time (Goldberg and Brown 1969; Yin and Chan 1990; Batra et al. 1997). Thus, when both ears are stimulated at the same frequency and the delay between the onset of the tones is varied, an MSO cell will respond maximally at a particular interaural phase angle, which corresponds to the condition when the arrival of the left and right inputs is in coincidence. For each ear a time delay results from the angular position of the stimulus with respect to the midsagittal plane, cochlear delays, and the path length between the AVCN on that side to the MSO cell. When the time delays on the two sides are equal, then the inputs arrive in phase and the MSO cell will discharge maximally.

A fundamental concept, first proposed on the basis of observations in only four cells, was characteristic delay (CD) (Rose et al. 1966). CD refers to what happens in a cell as the frequency of the tone is varied since changes in frequency also result in changes in phase. Thus, the interaural phase difference of the stimulus at which the maximal response is obtained may vary with the stimulus frequency for any given MSO cell. A cell that shows CD is one in






Figure 15.1. Parallel circuits in the superior olive for encoding interaural time (A) and interaural level (B) disparities in the medial and lateral superior olive, respectively. AN, Auditory nerve; AVCN, anteroventral cochlear nucleus; GBC, globular bushy cell; ICC, central nucleus of the inferior colliculus; LSO, lateral superior olive; MNTB, medial nucleus of the trapezoid body; MSO, medial superior olive; SBC, spherical bushy cell. Inset diagrams indicate how the neuron would be expected to respond to sound sources delivered in free field.

which the same relative discharge rate (e.g., maximum) is reached at the same ITD at all stimulus frequencies (Rose et al. 1966). This concept was quantified by showing that cells with CD had a linear relationship between interaural phase and stimulus frequency (Yin and Kuwada 1983). The slope of the phase-frequency plot corresponds to the CD while the phase intercept is the characteristic phase (CP) and specifies whether the point of common relative discharge is at the peak (CP = 0.0 or 1.0), trough (CP = 0.5), or in-between. Cells sensitive to ITD and exhibiting CD have been described in the MSO, ICC, dorsal nucleus of the lateral lemniscus (DNLL), medial geniculate body, and primary auditory cortex (Kuwada et al. 1987; Reale and Brugge 1990; Yin and Chan 1990; Stanford et al. 1992).

For stimuli with multiple spectral components, such as noise, a composite curve can be calculated by averaging all the ITD curves at the different frequencies. This composite curve resembles the ITD obtained using the broadband stimulus (Yin et al. 1986; Palmer et al. 1990). This relationship suggests that the individual frequency components of the stimulus are summed in near-linear fashion. An important feature of these responses to broadband stimuli is that the maxima of the composite curves of MSO cells tend to fall in the contralateral sound field, that is, with the ipsilateral ear delayed. This is expected, as the bushy cell axons from the contralateral AVCN must travel further to reach the ipsilateral MSO and hence coincidence would be expected if the contralateral stimulus is given a lead in time. Because the MSO projects to the ipsilateral ICC, we predict that ICC cells would also respond best to sound stimuli in the contralateral sound field, which is indeed the case. Thus, in ICC there is a representation of the contralateral sound field in terms of ITD (Fig. 15.1A).

### 2.2.2. Interaural Level Disparities

The other important cue for localization in the horizontal plane is created by the reflection and diffraction of sound by the head, shoulders, and external ears. Because the degree of reflection depends on sound wavelength, and hence on its frequency, the ILD cue is highly frequency dependent, with significant ILDs only at high frequencies where the wavelength is short (Blauert 1997). The neural circuit thought to be important for encoding ILDs involves lateral superior olive cells (LSO) which receive excitatory input from the ipsilateral spherical bushy cells and inhibitory input from the medial nucleus of the trapezoid body (MNTB), which is in turn driven by globular bushy cells of the contralateral AVCN with a large calyx of Held synapse on the MNTB cells (Fig. 15.1B; Warr 1966, 1972; Cant and Casseday 1986; Guinan and Li 1990; Smith et al. 1991, 1993). Thus, LSO cells are excited by a sound source in the ipsilateral sound field, where the excitation exceeds the inhibition, and inhibited when the source is in the contralateral sound field (Fig. 15.1B; Tsuchitani and Boudreau 1966; Tollin and Yin 2002a,b; Tollin 2003).

The projection of the LSO to the ICC is unusual and complex. LSO axons project bilaterally to the ICC, but some axons are glutamatergic and excitatory

while others are glycinergic and inhibitory (Glendenning and Masterton 1983; Saint Marie et al. 1989). In the cat the crossed projection originates primarily from the medial, high-frequency limb and is excitatory while the uncrossed projection from the lateral, low-frequency limb is both excitatory and inhibitory. Because LSO cells are excited by sound sources in the ipsilateral sound field, an excitatory crossed projection or an inhibitory uncrossed projection will result in ICC neurons that are excited by sound sources in the contralateral sound field (Fig. 15.1B; Glendenning et al. 1992). Note two caveats to this scheme. First, the inhibitory uncrossed projection will not result in an ICC cell that responds to sounds in the contralateral field alone: it needs to act on spontaneous activity or other driven excitation. Second, the excitatory uncrossed projection (not shown in Fig. 15.1B) provides inputs to the ICC that are excited by sounds in the ipsilateral sound field. Thus, most, but not all, ICC cells receive inputs that are responsive to stimuli in the contralateral, but not the ipsilateral, sound field.

Most binaural ICC cells respond to ITDs and ILDs that represent the contralateral sound field, like their visual and somatosensory counterparts. To ensure a unified perception of space, it is not surprising that all modalities follow the contralateral representation of space. What is remarkable is that the auditory system achieves this representation using a computational scheme that involves a massive, but precisely ordered, array of bilateral cochlear nucleus projections to the LSO, and in turn from the LSO to the ICC. This representation is important behaviorally because it predicts that a lesion above the level of the superior olive should affect localization in the contralateral sound field with little ipsilateral effect.

### *2.3. PERCEPTION AND PROCESSING OF SPECTRAL CUES FOR SOUND LOCALIZATION*

Anatomical projections for the binaural processing of ITD and ILD cues converge on the superior olive from the two cochlear nuclei (Ramón y Cajal 1909; Cant and Gaston 1982; Glendenning et al. 1985; Cant and Casseday 1986). By contrast, sound localization pathways with a selectivity for spectral information do not have overt anatomical features and have been inferred from the unusual spectral integration properties of populations in dorsal cochlear nucleus neurons (Nelken and Young 1997; Spirou et al. 1999; Imig et al. 2000) and ICC (Ehret and Merzenich 1988; Leroy and Wenstrup 2000; Davis et al. 2003). The anatomical specializations that subservise spectral processing may appear less differentiated than those for the binaural brain stem, but the functional enhancements are equally important in natural behavioral contexts where survival mandates the accurate localization of complex sounds.

#### **2.3.1 The Head-Related Transfer Function of the Cat**

Our description of spectral cues for sound localization emphasizes the large body of behavioral and physiological evidence from domestic cats. These directional

cues are described by the head-related transfer function (HRTF), which is the filter function characterizing relative spectral changes in complex sound propagating from the free field to the ear drum (Wightman and Kistler 1989a,b; Musicant et al. 1990). With ITD and ILD cues that are inherent in binaural comparisons of the HRTF, directionally dependent spectral features encode sound source location (Kulkarni and Colburn 1998). These spectral cues are critical for monaural sound localization, but binaural comparisons of HRTF spectral shapes also provides spatial information (Rice et al. 1992).

The HRTFs of a representative adult cat (Rice et al. 1992) were measured by recording sound energy near the ear drum with a miniature implanted microphone (Fig. 15.2). Rather than focusing on the specific directional properties of individual functions, several functions from the horizontal and median planes are juxtaposed to show generalized localization cues. One transfer function (Fig. 15.2, bold line) in both data sets indicates the ear's filtering effects for a broadband sound directly in front of the head ( $0^\circ$  AZ,  $0^\circ$  EL). By convention, a 0-dB gain indicates the HRTF has neither amplified nor attenuated the sound relative to its free-field sound pressure level.

The HRTFs suggest three domains of spectral information. Frequencies  $<5$  kHz are amplified, rising to an energy peak at about 4 kHz. Middle frequencies (5 to 20 kHz) usually have one deep and narrow energy trough, or spectral notch. Frequencies  $>20$  kHz manifest a complex spectral shape with multiple peaks and notches. HRTF gain falls with increasing frequency, implying a reduced salience for the high-frequency spectral cues of broadband sounds.

Directional features of the HRTF are revealed by comparing transfer functions across sound locations. The functions represent different azimuthal locations in the horizontal plane ( $\pm 75^\circ$  in  $15^\circ$  increments) (Fig. 15.2A) as well as different elevations in the median plane ( $-30^\circ$  to  $90^\circ$  in  $7.5^\circ$  increments) (Fig. 15.2B).

The HRTFs (Fig. 15.2) suggest that its low-frequency components are either amplified by the pinna when the sound source azimuth is in the near field, or attenuated by the head's "sound shadow" when it is in the far field. The difference in gain between the ears is equivalent to the ILD cue. This frequency region is virtually unaffected by elevation changes in the median plane.

The mid-frequency HRTF spectrum provides rich directional information for both azimuth and elevation. As a sound source moves in the horizontal or median plane, changes in location are reflected in notch frequency. This feature is critical for encoding sound source elevation, but it also may contribute to the identification of azimuth because horizontal plane spatial acuity improves with increasing stimulus bandwidth (Heffner and Heffner 1988).

The high-frequency HRTF displays complex and less systematic filtering properties. Consequently, directionally dependent relationships between energy peaks and notches have not been established. Nevertheless, high-frequency sounds tend to impart the perception of high elevation (Pratt 1930; Butler and Belendiuk 1977; Blauert 1997), presumably because they are less attenuated by the HRTF at such elevations (Wightman and Kistler 1989a; Rice et al. 1992).




Figure 15.2. Head-related transfer functions (HRTFs) of an adult cat for sound locations in the interaural horizontal plane (A) and median vertical plane (B). The transfer function for a sound source directly in front of the subject is repeated in both panels (*bold line*). Filtering effects fall into three frequency domains. At low frequencies, the HRTF gain increases as the sound source moves from contralateral to ipsilateral locations in the horizontal plane. These gain changes create interaural level differences (ILDs). At mid-frequencies, the frequency of a prominent spectral notch increases as the sound rises from low to high elevations in the median plane. At high frequencies, spectral peaks and notches display a less orderly relationship to the direction of the sound source. (Adapted from Rice et al. 1992.)

### 2.3.2. Role of High-Frequency Spectral Information in Spatial Discrimination

The spectral notches of the HRTF are conveyed to the brain by the discharge rates of auditory-nerve fibers (Poon and Brugge 1993). Computational models suggest a dichotomous role for spectral information in the high- vs. mid-frequency HRTF domains (Nandy and Ben-Arie 1996; May and Huang 1997). Although the high-frequency HRTF components are robust, individual spectral features are not unambiguously associated with discrete sound locations and thus are most effective for the simple detection of a change from one location to another. Mid-frequency spectral notches, in contrast, influence fewer auditory neurons but create more spatially localized response patterns, dictating performance in behavioral tasks that demand the absolute identification of sound source location.

The most common metric for evaluating the perceptual significance of HRTF-borne directional information is the minimum audible angle (MAA; Mills 1958), the threshold for detecting directional changes between paired sound sources. Relative localization tasks are often used in animal behavioral studies because they require less training and are easily related to psychophysical thresholds.

MAA tasks have been used extensively in cats (Martin and Webster 1987; Heffner and Heffner 1988; May et al. 1995). Behavioral results using noise bursts with different frequency domains were used to isolate the frequency dependence of HRTF filtering effects (Fig. 15.3) (Huang and May 1996a). Broadband noise contained spectral cues  $>5$  kHz. Mid-frequency noise was limited to the spectral notch region from 5 to 20 kHz. High-frequency noise conveyed only complex spectral cues  $>20$  kHz. For each stimulus, the MAA was defined as the smallest elevation change that elicited correct responses with a probability corresponding to the signal detection criterion  $d' = 1$ .

Correlations between directional acuity and the spectral domain of complex sounds have been demonstrated in cats (Martin and Webster 1987; Populin and Yin 1998a) and humans (Musicant and Butler 1984; Carlile et al. 1999; van




Figure 15.3. Effects of spectral frequency on minimum audible angles (MAAs) in the horizontal (A) and median planes (B). MAA is the smallest change in location relative to a reference speaker at  $0^\circ$  AZ,  $0^\circ$  EL that elicited threshold performance ( $d' = 1$ ). Histograms indicate the average thresholds ( $\pm$ SD) of three cats. Largest performance deficits were noted for tests in the median plane with mid-frequency noise (arrow). BBN, Broadband noise ( $>5$  kHz); HFN, High-frequency noise ( $>20$  kHz); MFN, mid-frequency noise (5 to 20 kHz). (Adapted from Huang and May 1996a.)

Schaik et al. 1999) and are replicated by responses to mid- and high-frequency noise bursts (Fig. 15.3). As predicted by the quality of directional information inherent in the HRTF (May and Huang 1997), high-frequency noise bursts contain many spectral cues for directional change. Their thresholds were similar to broadband performance. Cats showed deficits when tested with noise that conveyed the more singular mid-frequency notch cue. The relationship between behavioral performance and sound spectra was clearer for sound sources varying in elevation in the median plane than those varying in horizontal plane azimuth.

### 2.3.3. Role of Mid-Frequency Spectral Cues in Sound Orientation

Natural sound localization behaviors are concerned with the determination of actual locations, not the detection of spectral differences between paired sound sources. The unnatural aspects of the MAA task are circumvented in behavioral paradigms that require the subject to indicate the apparent location of an auditory stimulus by indicating or approaching the sound source. Such procedures are especially useful when directional percepts are predicted to change with modifications of acoustic parameters or lesions. Thus, limiting spectral content to high frequencies provides enough information to convey relative positional changes in the MAA task (Fig. 15.3), but does this manipulation preserve the absolute directionality of an auditory stimulus?

Early studies of cat directional hearing combined an array of sound sources with food or water reward. Sound presentations signaled the active goal box and the subject obtained the reward by approaching the proper speaker (Casseday and Neff 1973). When combined with surgical lesions, this procedure revealed valuable insight into the information processing roles of major auditory centers (Moore et al. 1974; Casseday and Neff 1975; Neff and Casseday 1977).

Unilateral ICC lesions induce only contralateral sound localization deficits in the goal-box task (Strominger and Oesterreich 1970; Goreva and Shcherbakov 1978; Jenkins and Masterton 1982). Robust perceptual deficits are only observed if the subject is forced to make an absolute localization among multiple sound sources (Jenkins and Masterton 1982; Jenkins and Merzenich 1984). Normal performance is achieved if the localization task is a simple spatial discrimination between paired sound sources (Butler and Musicant 1993; Kelly and Kavanagh 1994) and in nonspatial tasks such as the pure-tone discrimination (Neff et al. 1975). In these behavioral contexts, only the multiple speaker goal-box procedure specifically addresses the effects of the lesion on absolute directional hearing.

The dominant role of spectral information in directional hearing exists beyond the horizontal plane. Since flightless animals cannot approach an elevated stimulus, behavioral studies rely on a response in which the subject points to the perceived location of the sound source. In animals, the response is usually based on changes in position of the head (Thompson and Masterton 1978) or eyes

(Populin and Yin 1998a). This reflexive behavior may be used briefly in naive subjects (Sutherland et al. 1998a) or shaped into a food-reinforced operant paradigm for more detailed psychophysical analyses (May and Huang 1996; Populin and Yin 1998a,b; Tollin and Yin 2003a,b).

The effects of spectral content on the accuracy of sound orientation behavior was assessed in cats trained to orient to broadband noise (Fig. 15.4A) (Huang and May 1996b), then probing behavioral performance with occasional bursts of mid-frequency (Fig. 15.4B) or high-frequency noise (Fig. 15.4C). Band-limited stimuli elicited less accurate orientation than broadband noise, but spatially organized responses were maintained when the mid-frequency notches were available. High-frequency probes evoked inconsistent and inaccurate head movements. These deficits confirm that the high-frequency filtering effects of the cat's HRTF fail to impart an absolute directional identity, even though this domain is an excellent source of information for relative directional changes (Fig. 15.3).

The directionally dependent HRTF spectral shapes explain illusory elevation effects seen when paired sounds are presented from two locations in close temporal proximity (Tollin and Yin 2003a). Under summing localization conditions (see Section 3.1.1.), cats unexpectedly localize sounds at higher elevations than predicted by actual locations, a striking effect when both sound sources are located in the horizontal plane. These response errors are correlated with mid-frequency spectral notch alterations arising from the time-delayed summation of energy from the two locations.

Illusions of spatial azimuth and elevation also can be created by synthesizing HRTF-based spectral and binaural cues over headphones (Wightman and Kistler 1989b; Pralong 1996; Langendijk and Bronkhorst 2000). The perceptual realism of the resulting virtual space stimuli implies that the auditory system cannot isolate HRTF directional properties from acoustic source spectrum features. Nevertheless, to function effectively in the biological world, the auditory system must be able to derive the HRTF shape without prior knowledge of the stimulus. This processing dilemma is simplified under normal listening conditions because the amplitude spectra of most natural sounds lack sharp spectral features that may be confused with HRTF-filtering effects (Zakarauskas and Cynader 1993). The source spectrum can be disambiguated further by comparing the multiple directional perspectives provided by the two ears (Rice et al. 1992), head movement (Goossens and van Opstal 1999; Wightman and Kistler 1999), and mobile pinna (Young et al. 1996; Populin and Yin 1998b). Despite these heuristics, optimal localization is achieved when the auditory stimulus is a familiar sound (McGregor et al. 1985; Blauert 1997) and it is filtered by one's own ears (Wenzel et al. 1993; Hofman et al. 1998; Middlebrooks 1999).

#### 2.3.4. Coding of Spectral Cues

Electrophysiological studies of the ICC suggest specializations for processing spectral cues for sound localization. The existence of this discrete information






Figure 15.4. Effects of frequency domain on sound orientation behavior. Results from one cat summarized across orientation sessions in which discrete noise bursts were presented from randomly selected speaker locations. *Lines* connect individual responses (*circles*) to stimulus locations (*plus symbols*). Presentations of broadband noise comprised 85% of the trials (**A**). Noisebands with mid-frequency spectral notches (**B**) or high-frequency complex cues (**C**) were presented during the remaining probe trials. In this behavioral context, large performance deficits were observed for high-frequency stimuli (*arrow*). *Numerical labels* indicate stimulus azimuth and elevation in double-pole coordinates. (Adapted from Huang and May 1996b.)

processing pathway is based on the physiological classification of inhibitory patterns in the ICC of decerebrate cats (Ramachandran et al. 1999). Type V units have a V-shaped receptive field without strong inhibition. Type I units show a narrow I-shaped excitatory field bounded by lateral inhibitory areas. Type O units may be inhibited by any combination of frequency and level except for an O-shaped excitatory area near threshold.

Similarities in the inhibitory features of frequency response maps have linked type O responses to ascending projections from the dorsal cochlear nucleus (DCN; Davis et al. 1999, 2003; Ramachandran et al. 1999; Ramachandran and May 2002). The response map of a representative type O unit is compared with that of its hypothesized input, a type IV DCN neuron (Fig. 15.5A, B) (Young et al. 1992). The conservation of pure-tone responses between the putative target and source neurons suggests that the DCN spectral processing pathways remain functionally segregated in the ICC (Davis 2002).

The largely inhibitory pure-tone type O responses give way to excitation when the neurons are tested with more natural stimuli (Davis et al. 2003). The response class is particularly sensitive to broadband sounds containing HRTF-filtering effects. The spectral cues influencing these responses have been characterized with parametric stimuli that reduce the complex HRTF to a rectangular notch in broadband noise. A tuned excitatory response is elicited when the notch is swept in frequency across the receptive field of the type O unit (Fig. 15.5C). Because notch frequency is a directionally dependent feature of the cat's HRTF, notch-selective neurons can encode sound location (Young et al. 1992; Imig et al. 2000).

The directional sensitivity of type O responses has been demonstrated under closed-field conditions with noise bursts that provide more realistic simulations of binaural and spectral properties of the HRTF (Fig. 15.5E) (Delgutte et al. 1999; Davis et al. 2003). As predicted by the notch frequency sweep (Fig. 15.5C), maximum discharge rates align with a spatial contour that is defined by near-BF spectral notches. Interestingly, the unit exhibits a preferred location along the iso-notch contour that cannot be explained purely by notch directional selectivity. This enhanced spatial tuning may derive from wideband spectral integration properties allowing type O units to analyze the broader spectral context of mid-frequency notches, or a sensitivity to covarying binaural cues in some type O units (Ramachandran and May 2002).

Most type O units are silenced by surgical or pharmacological manipulations that disrupt neural transmission in the dorsal acoustic stria (DAS; Davis 2002), the principal DCN output pathway to the ICC. Although type IV units are distinguished by nonlinear spectral integration properties that endow sensitivity to HRTF-based spectral notches, type O units do not simply recapitulate their DCN inputs. Type IV units (Fig. 15.5D) display a frequency tuned inhibitory response to spectral notches (Young et al. 1992; Imig et al. 2000). Additional excitatory, inhibitory, and binaural inputs are required to transform this DCN pattern into the directionally selective type O unit (Davis et al. 2003).




Figure 15.5. Physiological properties of type O units in the ICC and type IV units in the DCN. Each row of the response maps shows driven discharge rates at a different level of attenuation. Inhibitory responses of the two unit types are similar for pure-tone frequency sweeps (**A**, **B**), but the type O unit shows a tuned excitatory response for notch frequency that is absent in the type IV unit (**C**, **D**). This notch selectivity creates a spatial receptive field when type O units are tested with virtual-space stimuli (**E**). Responses along the preferred iso-notch contour exhibit a “best location” that implies directional processing beyond notch frequency (*arrow*). (Adapted from Davis et al. 2003.)

### 2.3.5. Effects of Dorsal Cochlear Nucleus and Central Nucleus Lesions on Localization Behavior

The functions of DCN projections to the ICC have been investigated by measuring the effects of DAS lesions on localization behavior (Sutherland et al. 1998a,b; May 2000). Unlike trapezoid body lesions that induce deafness, this surgical procedure has little effect on hearing sensitivity in quiet and in background noise (Masterton and Granger 1988; Masterton et al. 1994). Consequently, it is possible to explore the nature of auditory deficits in subjects with bilateral DAS lesions.

A cat with DAS lesions was tested with bandpass noise to restrict spectral cues to HRTF-based notches between 5 and 20 kHz (Fig. 15.6). The subject oriented correctly to random source locations throughout the frontal sound field prior to the lesions (Fig. 15.6A), but had large errors after transecting DCN pathways (Fig. 15.6B) (May 2000). These deficits confirm that DCN/ICC pathways fulfill a critical spectral processing role in feline sound localization (Davis et al. 2003).

Elevation errors are pervasive (Fig. 15.6B). Regardless of the actual elevation of randomly selected sound sources, most responses fell within  $\pm 30^\circ$  of the horizontal plane. The under-estimation of sound source elevations exceeded  $60^\circ$  for extreme locations. These results reproduce the elevation-dependent errors induced when spectral cues are constrained in narrowband stimuli (Middlebrooks 1992). In both situations, the perceptual consequences of impaired spectral information are minimized for testing in the horizontal plane by the normal function of the binaural pathways. The remaining deficits suggest that monaural spectral representations within the DCN/ICC pathway also contribute to the perception of azimuth. This dual role is supported by the observation that most monaural ICC units are sensitive to both azimuth and elevation of virtual-space stimuli (Delgutte et al. 1999).

The functional specificity of DAS sound localization deficits has been evaluated by testing cats with MAA procedures (Sutherland et al. 1998a; May 2000) (Fig. 15.6C). Although these three cats had orientation deficits after the DAS lesion, none showed impaired spatial acuity when tested with mid-frequency noise bursts in the MAA procedure, confirming that DAS lesions do not produce general hearing deficits (Masterton and Granger 1988). These results further substantiate the view that relative spatial acuity and absolute directional hearing utilize different cues and processing pathways (Jenkins and Masterton 1982; May and Huang 1997).

### 2.3.6. The Neurobiology of Spectral Processing

Behavioral and electrophysiological studies suggest that type O units represent a specialized pathway for processing of spectral cues for sound localization. Although type O response patterns are common in decerebrate cats, they are rare in the ICC of some other laboratory species. For example, V-shaped fre-




Figure 15.6. Effects of dorsal acoustic stria (DAS) lesions on directional hearing. Orientation responses of a representative cat before (A) and after (B) bilateral DAS lesions. Elimination of DCN inputs to the ICC disrupted absolute orientation behavior, especially sound source elevation (*arrow*). Three cats with bilateral DAS lesions had excellent post-lesion performance in MAA procedures (C). Psychometric functions plot the percentage of correct responses relative to changes of elevation in the median plane. False alarms to catch trials without directional changes (*xts*) confirm the low probability of false positive responses in well-trained subjects. (Adapted from May 2000.)

quency response areas comprise most ICC units in anesthetized guinea pigs (Syka et al. 2000; Le Beau et al. 2001). (See Fig. 15.7A.) The remaining non-V-shaped units tend to have narrow tuning like the type I units in decerebrate cats.

Species differences in the prevalence of type O units cannot be explained by the use of anesthesia in guinea pigs, or the surgical elimination of descending projections to the ICC in decerebrate cats. After decerebration, guinea pigs show the same inhibitory patterns described in anesthetized preparations, including a conspicuous absence of type O units in frequency response maps (Fig. 15.7B).

An alternative explanation is that type O units reflect an auditory specialization that is restricted to species that require accurate sound localization, a biological need often associated with predatory lifestyles. The sound localization abilities of laboratory mice support this neuroethological premise. Besides poor directional acuity (Ehret and Dreyer 1984), single-unit recordings suggest an under-representation of type O responses in anesthetized mice (Egorova et al. 2001; Ehret et al. 2003). These species differences do not simply reflect the confounding influences of anesthesia and decerebration on the inhibitory prop-




Figure 15.7. Frequency response maps in nonpredatory species. Non-V-shape and V-shape responses are common in anesthetized guinea pigs (A), decerebrate guinea pigs (B), and awake mice (C). Type O units are rare in these species. (A: Adapted from Le Beau et al. 2001.) (B, C: Unpublished data from BJ May.)

erties of ICC neurons. V-shaped units also represent the most common response type in the ICC of awake, intact mice (Fig. 15.7C; May 2003).

This review relies heavily on research in domestic cats to describe the perception and processing of spectral cues for sound localization. Though regarded as a general model of mammalian hearing, the cat auditory system has evolved to meet the demands of nocturnal predation. These adaptations for prey localization may be species-specific.

Natural behavior provides insights into the functional significance of the common neural pathways and species specializations that remain to be addressed by laboratory studies. An emphasis on comparative behavior, particularly in terms of predator and prey relationships, has contributed greatly to investigations of the brain stem binaural sound localization system (Neff et al. 1975; Heffner and Heffner 1986, 1987, 1989; Moore 1987). A neurobiological perspective ought to prove equally important for understanding the spectral processing pathways of the ICC.

#### *2.4. EFFECTS OF LESIONS OF THE CENTRAL NUCLEUS OF THE INFERIOR COLLICULUS ON LOCALIZATION*

Physiological studies of the encoding of interaural localization cues suggest that the contralateral sound field is represented in the ICC even though the superior olive has both ipsilaterally and contralaterally driven cells. This dominant innervation has been confirmed by ICC lesions. A contralateral localization deficit followed a unilateral lesion of its output pathway, the brachium of the ICC (Strominger and Oesterreich 1970), and a unilateral lesion that affected LSO output in the lateral lemniscus or ICC caused a deficit in localizing sounds in the field contralateral to the lesion (Jenkins and Masterton 1982) and after large unilateral ICC lesions in the ferret (Kelly and Kavanagh 1994).

Since the ascending projections from the ICC to the medial geniculate body and then to the primary auditory cortex are predominantly uncrossed (see Chapter 7), we predict the same functional contralateral representation in the physiology of thalamic and cortical cells as well as a contralateral deficit in localization following a unilateral lesion of these higher centers. Indeed, most medial geniculate and primary auditory cortex cells have receptive fields in the contralateral sound field (Imig et al. 1990; Brugge et al. 1996; Irvine et al. 1996; Samson et al. 2000) and unilateral auditory cortex lesions create a contralateral localization deficit (Jenkins and Merzenich 1984).

#### *2.5 EFFECTS OF ELECTRICAL STIMULATION OF THE INFERIOR COLLICULUS*

There are few studies of behavioral results of electrical stimulation of the ICC. By 1970 (Syka and Straschill 1970; Syka and Radil-Weiss 1971) it was known

that head and eye movements could be evoked by stimulation of the ICC and that these effects used connections to the deep layers of the superior colliculus, a well-known sensorimotor integration site (Wurtz 1996; Sparks 2002).

In rodents the ICC has been implicated as one of the sites for abnormal brain activity in the form of seizures. Stimulation of the ICC by acoustic, electrical, or pharmacological means can evoke seizure activity in particular seizure-prone strains of rodents, the genetically epilepsy prone rat (GEPR; Ribak and Morin 1995; Ross and Coleman 2000). This system has been used as a model for studying the interaction between excitatory and inhibitory neurotransmitters in inducing epileptic activity (see Chapter 21).

### 3. PHYSIOLOGY IN AWAKE AND BEHAVING ANIMALS

In all of the experiments reviewed in the preceding, the physiological responses to localization cues were recorded in anesthetized or decerebrated animals. How anesthesia affects the responses is not understood and undoubtedly depends on the dosage, type, and the site of recording as well as the species. A major advance since 1960 is the development of techniques to record from awake, behaving animals, which avoids the contaminating effects of anesthesia and allows one to study the neural bases of higher-order cognitive effects.

There are few studies since 1980 in awake behaving preparations, even though several earlier studies used this technique. Responses of monkey ICC cells in an auditory reaction time task were compared to responses to the same stimuli when the monkey was quiescent. Discharge patterns resembled those in acute experiments and rates significantly increased during task performance (Ryan and Miller 1977, 1978). Similar comparisons were also made from the cochlear nucleus to auditory cortex (Ryan et al. 1984). Moreover, in behaving animals there are multisensory interactions as eye position affects some ICC neuron responses as much as sound source location in monkeys (Groh et al. 2001).

#### 3.1. THE PRECEDENCE EFFECT

##### 3.1.1. Psychophysics

There has been considerable interest in an auditory illusion known as the precedence effect (PE), or law of the first wavefront (Wallach et al. 1949), shown by placing a subject between two speakers (Fig. 15.8A) and delivering the same transient sound to both speakers with a variable interstimulus delay (ISD) between the sounds. An idealized plot (Blauert 1997) of the perceived location of the sound as a function of ISD (Fig. 15.8B) shows that, if both speakers are activated simultaneously, the observer will localize the sound to a single phantom source midway between them. As ISD gradually lengthens the perceived source moves towards the leading speaker until, at approximately 1 ms ISD the sound is perceived near the leading source. The interval between about  $\pm 1$  ms






Figure 15.8. The precedence effect. (A) Diagram of the stimulus configuration in free field with the subject midway between two speakers (A, B) (top) and (B) the perceived azimuthal location of the sound as a function of the interclick delay. By convention negative delays correspond to speaker B leading. (Adapted from Blauert 1997 and Yin 1994.)

has been termed summing localization. ISDs between 1 and 5 to 10 ms represent the period of the precedence effect, or localization dominance: the sound is localized to the position of the leading source even though the lagging source is readily localized when presented alone. At longer ISDs the subject hears both sounds at their respective positions, and the shortest ISD at which this occurs is the echo threshold. In practice the echo threshold value is variable and depends on the instructions given to the subject and on the duration and type of acoustic stimulus. Sustained and complex sounds (music, speech) usually have longer echo thresholds.

The term, precedence effect, refers to the way in which the leading sound appears to take precedence over the lagging one in judgments of sound location, as if the lagging sound's location were suppressed by the auditory system. However, the lagging source still plays a role in perception other than localization: for delays in the range of localization dominance, the fused sound can be readily discriminated from the leading sound by a change in timbre, although it is localized to the position of the leading sound. It is thought that the precedence effect is critical for accurately localizing sound in a reverberant environment, such as a room, where any sound will produce time-delayed echoes from many different spatial locations.

An important question regarding the neural bases of PE is how it is experienced by animals. PE has been demonstrated in cats (Populin and Yin 1998a; Tollin and Yin 2003b), birds (Keller and Takahashi 1996; Dent and Dooling 2003), dogs (Ashmead et al. 1986), rats (Kelly 1974), and crickets (Wytenbach and Hoy 1993). Mean horizontal eye position as a function of ISD of a cat trained to fixate sound sources when presented with pairs of short noise bursts to simulate the PE (Tollin and Yin 2003b) demonstrates this effect (Fig. 15.9). With the head held, all cats consistently undershoot the target (Fig. 15.9A, B: leftward arrows).

The cat (Fig. 15.9) and human (Fig. 15.8B) responses are similar during the localization summing period except that the period is only  $\pm 400 \mu\text{s}$  in the cat because of its smaller head width. For delays between 1 and 10 ms, the cat showed localization dominance to the position of the leading speaker and at delays  $>10$  ms it often made saccades to the location of the lagging sound or double saccades to both the leading and lagging sounds, resulting in mean responses near 0 and large variability. We infer from this behavior, never seen for




Figure 15.9. Responses of a cat to the PE stimuli. Azimuthal final eye position as a function of interstimulus delay (ISD). The locations of the speakers A and B are indicated by the arrows on the right and the horizontal lines ( $\pm$  standard deviation) show the mean eye position for the two speakers when delivered in isolation. (Adapted from Tollin and Yin 2003b, used with permission.)

shorter delays, that the cat was perceiving both sounds and was unsure which to look at. Thus, the cat echo threshold is between 10 and 15 ms.

### 3.1.2. Physiology

The first studies of the physiological responses to PE stimuli were in anesthetized animals (Yin 1994). ICC cells respond to click pairs delivered dichotically (Fig. 15.10). A +400  $\mu$ s ITD, an effective stimulus for this cell, was imposed on both pairs of clicks while the interclick delay (ICD) varied from 1 to 101 ms. At long ICDs the response to both pairs of clicks is clear, but as the ICD decreases gradually the response to the lagging click declines until it disappears at ICDs <7 ms, while there is little change in the response to the leading one. A recovery curve plot shows the response to the lagging click normalized by the response to the same click when delivered in isolation (Fig. 15.9B). The delay at 50% recovery is often used as a metric for half-maximal suppression, and in this cell is 11 ms.

Suppression of the response to the lagging click was common in the ICC and has been studied in other species (rabbit: Fitzpatrick et al. 1995; barn owl: Keller and Takahashi 1996) and extends from the auditory nerve to the cortex (Parham et al. 1996, 1998; Fitzpatrick et al. 1999). Spatial location, ITD, level, duration, and binaurality of the leading sound can each affect suppression (Litovsky and Yin 1998a,b; Litovsky and Delgutte 2002). The half-maximal delay values in different cells were variable, with significant differences in the degree of suppression in the anesthetized cat and unanesthetized rabbit (Yin 1994; Fitzpatrick et al. 1995, 1999; Litovsky and Yin 1998a,b).

To investigate whether the differences between the cat and rabbit results were due to the species difference or to the anesthesia, recordings were also made in the ICC while the cat was awake and performing a sound localization paradigm. The distribution of half-maximal delays for cells in the anesthetized cat (Litovsky and Yin 1998b; Fig. 15.11, gray bars), for unanesthetized, quiescent rabbit (Fitzpatrick et al. 1995; Fig. 15.11, white bars) and in awake, behaving cats (Fig. 15.11, black bars) are available. The half-maximal delays in the behaving cat (circles representing mean and s.d.) were significantly different than in the anesthetized cat but similar in the awake rabbit and suggest that the differences reflect anesthesia. Mean recovery curves for the population of ICC cells in the awake animal were computed separately for either ipsilateral side or contralateral side-leading speaker and show significantly stronger contralateral leading suppression (Fig. 15.10B). The PE period (Fig. 15.10, shaded region) shows considerable suppression in the response of the population of ICC cells when PE is behaviorally effective.

## 4. FUTURE DIRECTIONS

We have focused on the relationship between sound localization behavior and neural response patterns in the ascending pathways of the auditory midbrain.




Figure 15.10. Echo suppression in the ICC under dichotic stimulation. **(A)** Dot rasters show the responses to 40 repetitions of ICDs ranging from 1 to 101 ms. Each dichotic click had an ITD of +400  $\mu$ s. **(B)** Recovery curve of the data in **(A)**. The response to the lagging click was normalized by the response to the same click without the leading click. Half-maximal delay was 12 ms. (Adapted from Yin 1994.)

This emphasis reflects the prominent role the ICC has played in current behavioral and physiological assessments of the inferior colliculus. It is almost certain that the midbrain is involved in other auditory behaviors and that these processes reflect descending influences from higher auditory centers.

Physiological studies find that the auditory processing of a meaningful sound is enhanced by corticofugal pathways that terminate in the dorsal cortex of the inferior colliculus (Zhang et al. 1997; Jen et al. 1998; Yan and Ehret 2001, 2002). When repeating tones are paired with noxious shocks, neurons encoding the fear-conditioned stimulus show an expanded reorganization in the inferior colliculus of big brown bats (Gao and Suga 1998). Analogous changes in frequency tuning are evoked by electrical stimulation of auditory cortex (Chowdhury and Suga 2000), or abolished by pharmacological inactivation of the same locations (Gao and Suga 2000). By contrast, specialized echolocation areas in the inferior colliculus of mustached bats show increased activity and sharper frequency tuning when matching cortical frequency regions are stimulated (Zhang and Suga




Figure 15.11. (A) Histogram of the half-maximal delays in anesthetized cats (Litovsky and Yin 1998b; *gray bars*), awake rabbits (Fitzpatrick et al. 1995; *white bars*), and awake, behaving cats (black bars). The means and standard error of the mean of the distributions are shown by the *circles above*. (B) Mean normalized recovery curves of all cells as a function of ISD with the ipsilateral (*filled circles*) or contralateral (*open squares*) speaker leading. (Adapted from Tollin et al. 2005.)

2000), and those of unmatched neurons are suppressed. This compressive reorganization creates a more selective representation of biosonar signals (Suga et al. 2002).

The physiological correlates of attentional filtering were predicted by the classic behavioral paradigms (Jane et al. 1965), in which reciprocal functional roles for the central nucleus and dorsal cortex were proposed. Unlike the deafening effects of ICC lesions, surgical manipulations of the dorsal cortex do not impair sound localization but diminish auditory attention. When redundant auditory and visual cues signal an impending electrical shock, intact cats tend to base their avoidance behaviors on auditory information. This natural “prepotency” of the auditory stimulus is supplanted by visual information in cats with lesions of the dorsal cortex.

Cognitive influences such as learning, attention, and compensation represent exciting new directions for future studies of the physiological underpinnings of acoustic behavior. Within the auditory midbrain, it is becoming increasingly clear that while these complex processes are driven by ascending sensory representations, a set of descending perceptual filters that are suppressed by anesthesia, silenced by decerebration, and altered by experience operate in parallel. From this perspective, acoustic behavior provides an essential context for understanding midbrain function.

## Abbreviations

AN	auditory nerve
AVCN	anteroventral cochlear nucleus
CD	characteristic delay
CP	characteristic phase
DAS	dorsal acoustic stria
DCN	dorsal cochlear nucleus
DNLL	dorsal nucleus of lateral lemniscus
GBC	globular bushy cell
HRTF	head-related transfer function
IC	inferior colliculus
ICC	central nucleus of inferior colliculus
ILD	interaural level difference
ISD	interstimulus delay
ITD	interaural time difference
LSO	lateral superior olive
MAA	minimum audible angle
MSO	medial superior olive
PE	precedence effect
SBC	spherical bushy cell

## References

- Andersen RA and Buneo CA (2002) Intentional maps in posterior parietal cortex. *Annual Review of Neuroscience* **25**:189–220.
- Ashmead DH, Clifton RK, and Reese EP (1986) Development of auditory localization in dogs: single source and precedence effect sounds. *Developmental Psychobiology* **19**: 91–103.
- Batra R, Kuwada S, and Fitzpatrick DC (1997) Sensitivity to interaural temporal disparities of low- and high-frequency neurons in the superior olivary complex. II. Coincidence detection. *Journal of Neurophysiology* **78**:1237–1247.
- Blauert J (1997) *Spatial Hearing. The Psychophysics of Human Sound Localization*. MIT Press, Cambridge.
- Brugge JF, Reale RA, and Hind JE (1996) The structure of spatial receptive fields of neurons in primary auditory cortex of the cat. *Journal of Neuroscience* **16**:4420–4437.
- Butler RA and Belendiuk K (1977) Spectral cues utilized in the localization of sound in the median sagittal plane. *Journal of the Acoustical Society of America* **61**:1264–1269.
- Butler RA and Musicant AD (1993) Binaural localization: influence of stimulus frequency and the linkage to covert peak areas. *Hearing Research* **67**:220–229.
- Cant NB and Gaston KC (1982) Pathways connecting the right and left cochlear nuclei. *Journal of Comparative Neurology* **212**:313–326.
- Cant NB and Casseday JH (1986) Projections from the anteroventral cochlear nucleus to

- the lateral and medial superior olivary nuclei. *Journal of Comparative Neurology* **247**:457–476.
- Carlile S, Delaney S, and Corderoy A (1999) The localisation of spectrally restricted sounds by human listeners. *Hearing Research* **128**:175–189.
- Casseday JH and Neff WD (1973) Localization of pure tones. *Journal of the Acoustical Society of America* **54**:365–372.
- Casseday JH and Neff WD (1975) Auditory localization: role of auditory pathways in brain stem of the cat. *Journal of Neurophysiology* **38**:842–858.
- Chowdhury SA and Suga N (2000) Reorganization of the frequency map of the auditory cortex evoked by cortical electrical stimulation in the big brown bat. *Journal of Neurophysiology* **83**:1856–1863.
- Davis KA (2002) Evidence of a functionally segregated pathway from dorsal cochlear nucleus to inferior colliculus. *Journal of Neurophysiology* **87**:1824–1835.
- Davis KA, Ramachandran R, and May BJ (1999) Single-unit responses in the inferior colliculus of decerebrate cats II. Sensitivity to interaural level differences. *Journal of Neurophysiology* **82**:164–175.
- Davis KA, Ramachandran R, and May BJ (2003) Auditory processing of spectral cues for sound localization in the inferior colliculus. *Journal of the Association for Research in Otolaryngology* **4**:148–163.
- Delgutte B, Joris PX, Litovsky RY, and Yin TCT (1999) Receptive fields and binaural interactions for virtual-space stimuli in the cat inferior colliculus. *Journal of Neurophysiology* **81**:2833–2851.
- Dent ML and Dooling RJ (2003) Investigations of the precedence effect in budgerigars: the perceived location of auditory images. *Journal of the Acoustical Society of America* **113**:2159–2169.
- Egorova M, Ehret G, Vartanian I, and Esser K-H (2001) Frequency response areas of neurons in the mouse inferior colliculus. I. Threshold and tuning characteristics. *Experimental Brain Research* **140**:145–161.
- Ehret G and Dreyer A (1984) Localization of tones and noise in the horizontal plane by unrestrained house mice (*Mus musculus*). *Journal of Experimental Biology* **109**:163–174.
- Ehret G and Merzenich MM (1988) Complex sound analysis (frequency resolution, filtering and spectral integration) by single units of the inferior colliculus of the cat. *Brain Research* **472**:139–163.
- Ehret G, Egorova M, Hage SR, and Mueller BA (2003) Spatial map of frequency tuning-curve shapes in the mouse inferior colliculus. *NeuroReport* **14**:1365–1369.
- Fitzpatrick DC, Kuwada S, Batra R, and Trahiotis C (1995) Neural responses to simple simulated echoes in the auditory brain stem of the unanesthetized rabbit. *Journal of Neurophysiology* **74**:2469–2486.
- Fitzpatrick DC, Kuwada S, Kim DO, Parham K, and Batra R (1999) Responses of neurons to click-pairs as simulated echoes: auditory nerve to auditory cortex. *Journal of the Acoustical Society of America* **106**:3460–3472.
- Gao E and Suga N (1998) Experience-dependent corticofugal adjustment of midbrain frequency map in bat auditory system. *Proceedings of the National Academy of Sciences of the United States of America* **95**:12663–12670.
- Gao E and Suga N (2000) Experience-dependent plasticity in the auditory cortex and the inferior colliculus of bats: role of the corticofugal system. *Proceedings of the National Academy of Sciences of the United States of America* **97**:8081–8086.
- Glendenning KK and Masterton RB (1983) Acoustic chiasm: efferent projections of the lateral superior olive. *Journal of Neuroscience* **3**:1521–1537.

- Glendenning KK, Hutson KA, Nudo RJ, and Masterton RB (1985) Acoustic chiasm II: anatomical basis of binaurality in lateral superior olive of cat. *Journal of Comparative Neurology* **232**:261–285.
- Glendenning KK, Baker BN, Hutson KA, and Masterton RB (1992) Acoustic chiasm V: inhibition and excitation in the ipsilateral and contralateral projections of LSO. *Journal of Comparative Neurology* **319**:100–122.
- Goldberg JM and Brown PB (1969) Response of binaural neurons of dog superior olivary complex to dichotic tonal stimuli: some physiological mechanisms of sound localization. *Journal of Neurophysiology* **32**:613–636.
- Goossens HH and van Opstal AJ (1999) Influence of head position on the spatial representation of acoustic targets. *Journal of Neurophysiology* **81**:2720–2736.
- Goreva OA and Shcherbakov VI (1978) Behavior of cats after unilateral section of the brachii of the inferior colliculi. *Zhurnal Vysshei Nervnoi Deiatelnosti Imeni Ivan Pavlovich Pavlova* **28**:727–734 (in Russian).
- Groh JM, Trause AS, Underhill AM, Clark KR, and Inati S (2001) Eye position influences auditory responses in primate inferior colliculus. *Neuron* **29**:509–518.
- Guinan JJ Jr and Li RY (1990) Signal processing in brainstem auditory neurons which receive giant endings (calyces of Held) in the medial nucleus of the trapezoid body of the cat. *Hearing Research* **49**:321–334.
- Heffner RS and Heffner HE (1986) Localization of tones by horses: use of binaural cues and the role of the superior olivary complex. *Behavioral Neuroscience* **100**:93–103.
- Heffner RS and Heffner HE (1987) Localization of noise, use of binaural cues, and a description of the superior olivary complex in the smallest carnivore, the least weasel (*Mustela nivalis*). *Behavioral Neuroscience* **101**:701–708, 744–745.
- Heffner RS and Heffner HE (1988) Sound localization acuity in the cat: effect of azimuth, signal duration, and test procedure. *Hearing Research* **36**:221–232.
- Heffner RS and Heffner HE (1989) Sound localization, use of binaural cues and the superior olivary complex in pigs. *Brain, Behavior and Evolution* **33**:248–258.
- Hofman PM, Van Riswick JG, and Van Opstal AJ (1998) Relearning sound localization with new ears. *Nature Neuroscience* **1**:417–421.
- Huang AY and May BJ (1996a) Spectral cues for sound localization in cats: effects of frequency domain on minimum audible angles in the median and horizontal planes. *Journal of the Acoustical Society of America* **100**:2341–2348.
- Huang AY and May BJ (1996b) Sound orientation behavior in cats. II. Mid-frequency spectral cues for sound localization. *Journal of the Acoustical Society of America* **100**:1070–1080.
- Imig TJ, Irons WA, and Samson FR (1990) Single-unit selectivity to azimuthal direction and sound pressure level of noise bursts in cat high-frequency primary auditory cortex. *Journal of Neurophysiology* **63**:1448–1466.
- Imig TJ, Bibikov NG, Poirier P, and Samson FK (2000) Directionality derived from pinna-cue spectral notches in cat dorsal cochlear nucleus. *Journal of Neurophysiology* **83**:907–925.
- Irvine DRF, Rajan R, and Aitkin LM (1996) Sensitivity to interaural intensity differences of neurons in primary auditory cortex of the cat. I. Types of sensitivity and effects of variations in sound pressure level. *Journal of Neurophysiology* **75**:75–96.
- Jane JA, Masterton RB, and Diamond IT (1965) The function of the tectum for attention to auditory stimuli in the cat. *Journal of Comparative Neurology* **125**:165–191.
- Jeffress L (1948) A place theory of sound localization. *Journal of Comparative and Physiological Psychology* **41**:35–39.



- Jen PH-S, Chen QC, and Sun XD (1998) Corticofugal regulation of auditory sensitivity in the bat inferior colliculus. *Journal of Comparative Physiology A* **183**:683–697.
- Jenkins WM and Masterton RB (1982) Sound localization: effects of unilateral lesions in central auditory system. *Journal of Neurophysiology* **47**:987–1016.
- Jenkins WM and Merzenich MM (1984) Role of cat primary auditory cortex for sound-localization behavior. *Journal of Neurophysiology* **52**:819–847.
- Johnson DH (1980) The relationship between spike rate and synchrony in responses of auditory-nerve fibers to single tones. *Journal of the Acoustical Society of America* **68**:1115–1122.
- Joris PX, Carney LH, Smith PH, and Yin TCT (1994) Enhancement of neural synchronization in the anteroventral cochlear nucleus. I. Responses to tones at the characteristic frequency. *Journal of Neurophysiology* **71**:1022–1036.
- Keller CH and Takahashi TT (1996) Responses to simulated echoes by neurons in the barn owl's auditory space map. *Journal of Comparative Physiology A* **178**:499–512.
- Kelly JB (1974) Localization of paired sound sources in the rat: small time differences. *Journal of the Acoustical Society of America* **55**:1277–1284.
- Kelly JB and Kavanagh GL (1994) Sound localization after unilateral lesions of inferior colliculus in the ferret (*Mustela putorius*). *Journal of Neurophysiology* **71**:1078–1087.
- Kulkarni A and Colburn HS (1998) Role of spectral detail in sound-source localization. *Nature* **396**:747–749.
- Kuwada S and Yin TCT (1983) Binaural interaction in low-frequency neurons in inferior colliculus of the cat. I. Effects of long interaural delays, intensity, and repetition rate on interaural delay function. *Journal of Neurophysiology* **50**:981–999.
- Kuwada S, Stanford TR, and Batra R (1987) Interaural phase-sensitive units in the inferior colliculus of the unanesthetized rabbit: effects of changing frequency. *Journal of Neurophysiology* **57**:1338–1360.
- LaMotte RH and Mountcastle VB (1975) Capacities of humans and monkeys to discriminate vibratory stimuli of different frequency and amplitude: a correlation between neural events and psychological measurements. *Journal of Neurophysiology* **38**:539–559.
- Langendijk EH and Bronkhorst AW (2000) Fidelity of three-dimensional-sound reproduction using a virtual auditory display. *Journal of the Acoustical Society of America* **107**:528–537.
- Le Beau FEN, Malmierca MS, and Rees A (2001) Iontophoresis in vivo demonstrates a key role for GABA<sub>A</sub> and glycinergic inhibition in shaping frequency response areas in the inferior colliculus of guinea pig. *Journal of Neuroscience* **21**:7303–7312.
- Leroy SA and Wenstrup JJ (2000) Spectral integration in the inferior colliculus of the mustached bat. *Journal of Neuroscience* **20**:8533–8541.
- Litovsky RY and Delgutte B (2002) Neural correlates of the precedence effect in the inferior colliculus: effect of localization cues. *Journal of Neurophysiology* **87**:976–994.
- Litovsky RY and Yin TCT (1998a) Physiological studies of the precedence effect in the inferior colliculus of the cat. I. Correlates of psychophysics. *Journal of Neurophysiology* **80**:1285–1301.
- Litovsky RY and Yin TCT (1998b) Physiological studies of the precedence effect in the inferior colliculus of the cat. II. Neural mechanisms. *Journal of Neurophysiology* **80**:1302–1316.
- Martin RL and Webster WR (1987) The auditory spatial acuity of the domestic cat in the interaural horizontal and median vertical planes. *Hearing Research* **30**:239–252.

- Masterton RB and Granger EM (1988) Role of the acoustic striae in hearing: contribution of dorsal and intermediate striae to detection of noises and tones. *Journal of Neurophysiology* **60**:1841–1860.
- Masterton RB, Granger EM, and Glendenning KK (1994) Role of acoustic striae in hearing: mechanism for enhancement of sound detection in cats. *Hearing Research* **73**:209–222.
- Maunsell JHR and Cook EP (2002) The role of attention in visual processing. *Philosophical Transactions of the Royal Society of London, series B, Biological Sciences* **357**:1063–1072.
- May BJ (2000) Role of the dorsal cochlear nucleus in the sound localization behavior of cats. *Hearing Research* **148**:74–87.
- May BJ (2003) Species differences in the spectral processing pathways of the inferior colliculus. *Association for Research in Otolaryngology Abstracts* **26**:173.
- May BJ and Huang AY (1996) Sound orientation behavior in cats. I. Localization of broadband noise. *Journal of the Acoustical Society of America* **100**:1059–1069.
- May BJ and Huang AY (1997) Spectral cues for sound localization in cats: a model for discharge rate representations in the auditory nerve. *Journal of the Acoustical Society of America* **101**:2705–2719.
- May BJ, Huang AY, Aleszczyk CM, and Hienz RD (1995) Design and conduct of sensory experiments for domestic cats. In: Klumpf G, Dooling RJ, Fay RR, and Stebbins WC (eds). *Methods in Comparative Psychoacoustics*. Birkhäuser Verlag, Basel, pp. 95–108.
- McGregor P, Horn AG, and Todd MA (1985) Are familiar sounds ranged more accurately? *Perceptual and Motor Skills* **61**:1082.
- Middlebrooks JC (1992) Narrow-band sound localization related to external ear acoustics. *Journal of the Acoustical Society of America* **92**:2607–2624.
- Middlebrooks JC (1999) Individual differences in external-ear transfer functions reduced by scaling in frequency. *Journal of the Acoustical Society of America* **106**:1480–1492.
- Mills AW (1958) On the minimum audible angle. *Journal of the Acoustical Society of America* **30**:237–246.
- Moore CN, Casseday JH, and Neff WD (1974) Sound localization: the role of the commissural pathways of the auditory system of the cat. *Brain Research* **82**:13–26.
- Moore JK (1987) The human auditory brain stem: a comparative view. *Hearing Research* **29**:1–32.
- Musicant AD and Butler RA (1984) The influence of pinnae-based spectral cues on sound localization. *Journal of the Acoustical Society of America* **75**:1195–1200.
- Musicant AD, Chan JCK, and Hind JE (1990) Direction-dependent spectral properties of cat external ear: new data and cross-species comparisons. *Journal of the Acoustical Society of America* **87**:757–781.
- Nandy D and Ben-Arie J (1996) An auditory localization model based on high-frequency spectral cues. *Annals of Biomedical Engineering* **24**:621–638.
- Neff WD and Casseday JH (1977) Effects of unilateral ablation of auditory cortex on monaural cat's ability to localize sound. *Journal of Neurophysiology* **40**:44–52.
- Neff WD, Diamond IT, and Casseday JH (1975) Behavioral studies of auditory discrimination: central nervous system. In: Keidel WD and Neff WD (eds). *Handbook of Sensory Physiology*, Volume V, Part 2: *Auditory System, Anatomy, Physiology (Ear)*. Springer-Verlag, Berlin, pp. 307–400.
- Nelken I and Young ED (1997) Linear and nonlinear spectral integration in type IV neurons of the dorsal cochlear nucleus. I. Regions of linear interaction. *Journal of Neurophysiology* **78**:790–799.

- Palmer AR, Rees A, and Caird D (1990) Interaural delay sensitivity to tones and broadband signals in the guinea-pig inferior colliculus. *Hearing Research* **50**:71–86.
- Parham K, Zhao HB, and Kim DO (1996) Responses of auditory nerve fibers of the unanesthetized decerebrate cat to click pairs as simulated echoes. *Journal of Neurophysiology* **76**:17–29.
- Parham K, Zhao HB, Ye Y, and Kim DO (1998) Responses of anteroventral cochlear nucleus neurons of the unanesthetized decerebrate cat to click pairs as simulated echoes. *Hearing Research* **125**:131–146.
- Parker AJ and Newsome WT (1998) Sense and the single neuron: probing the physiology of perception. *Annual Review of Neuroscience* **21**:227–277.
- Poon PWF and Brugge JF (1993) Virtual-space receptive fields of single auditory nerve fibers. *Journal of Neurophysiology* **70**:667–676.
- Populin LC and Yin TCT (1998a) Behavioral studies of sound localization in the cat. *Journal of Neuroscience* **18**:2147–2160.
- Populin LC and Yin TCT (1998b) Pinna movements of the cat during sound localization. *Journal of Neuroscience* **18**:4233–4243.
- Pralong D (1996) The role of individualized headphone calibration for the generation of high fidelity virtual auditory space. *Journal of the Acoustical Society of America* **100**:3785–3793.
- Pratt CC (1930) The spatial character of high and low tones. *Journal of Experimental Psychology* **13**:278–285.
- Ramachandran R and May BJ (2002) Functional segregation of ITD sensitivity in the inferior colliculus of decerebrate cats. *Journal of Neurophysiology* **88**:2251–2261.
- Ramachandran R, Davis KA, and May BJ (1999) Single-unit responses in the inferior colliculus of decerebrate cats I. Classification based on frequency response maps. *Journal of Neurophysiology* **82**:152–163.
- Ramón y Cajal S (1909) *Histologie du Système Nerveux de l'Homme et des Vertébrés*. Maloine, Paris.
- Rayleigh L (1907) On our perception of sound direction. *Philosophical Magazine* **13**:214–232.
- Reale RA and Brugge JF (1990) Auditory cortical neurons are sensitive to static and continuously changing interaural phase cues. *Journal of Neurophysiology* **64**:1247–1260.
- Ribak CE and Morin CL (1995) The role of the inferior colliculus in a genetic model of audiogenic seizures. *Anatomy and Embryology* **191**:279–295.
- Rice JJ, May BJ, Spirou GA, and Young ED (1992) Pinna-based spectral cues for sound localization in cat. *Hearing Research* **58**:132–152.
- Rose JE, Gross NB, Geisler CD, and Hind JE (1966) Some neural mechanisms in the inferior colliculus of the cat which may be relevant to localization of a sound source. *Journal of Neurophysiology* **29**:288–314.
- Ross KC and Coleman JR (2000) Developmental and genetic audiogenic seizure models: behavior and biological substrates. *Neuroscience and Biobehavioral Reviews* **24**:639–653.
- Ryan A and Miller J (1977) Effects of behavioral performance on single-unit firing patterns in inferior colliculus of the rhesus monkey. *Journal of Neurophysiology* **40**:943–956.
- Ryan A and Miller J (1978) Single unit responses in the inferior colliculus of the awake and performing rhesus monkey. *Experimental Brain Research* **32**:389–407.

- Ryan AF, Miller JM, Pflugst BE, and Martin GK (1984) Effects of reaction time performance on single-unit activity in the central auditory pathway of the rhesus macaque. *Journal of Neuroscience* **4**:298–308.
- Saint Marie RL, Ostapoff E-M, Morest DK, and Wenthold RJ (1989) Glycine-immunoreactive projection of the cat lateral superior olive: possible role in midbrain ear dominance. *Journal of Comparative Neurology* **279**:382–396.
- Samson FK, Barone P, Irons WA, Clarey JC, Poirier P, and Imig TJ (2000) Directionality derived from differential sensitivity to monaural and binaural cues in the cat's medial geniculate body. *Journal of Neurophysiology* **84**:1330–1345.
- Siebert WM (1970) Frequency discrimination in the auditory system: place or periodicity mechanism. *Proceedings of the Institute of Electrical and Electronics Engineers* **58**:723–730.
- Smith PH, Joris PX, Carney LH, and Yin TCT (1991) Projections of physiologically characterized globular bushy cell axons from the cochlear nucleus of the cat. *Journal of Comparative Neurology* **304**:387–407.
- Smith PH, Joris PX, and Yin TCT (1993) Projections of physiologically characterized spherical bushy cell axons from the cochlear nucleus of the cat: evidence for delay lines to the medial superior olive. *Journal of Comparative Neurology* **331**:245–260.
- Sparks DL (2002) The brainstem control of saccadic eye movements. *Nature Reviews Neuroscience* **3**:952–964.
- Spirou GA, Davis KA, Nelken I, and Young ED (1999) Spectral integration by type II interneurons in dorsal cochlear nucleus. *Journal of Neurophysiology* **82**:648–663.
- Stanford TR, Kuwada S, and Batra R (1992) A comparison of the interaural time sensitivity of neurons in the inferior colliculus and thalamus of the unanesthetized rabbit. *Journal of Neuroscience* **12**:3200–3216.
- Strominger NL and Oesterreich RE (1970) Localization of sound after section of the brachium of the inferior colliculus. *Journal of Comparative Neurology* **138**:1–18.
- Suga N, Xiao Z, Ma X, and Ji W (2002) Plasticity and corticofugal modulation for hearing in adult animals. *Neuron* **36**:9–18.
- Sutherland DP, Glendenning KK, and Masterton RB (1998a) Role of acoustic striae in hearing: discrimination of sound-source elevation. *Hearing Research* **120**:86–108.
- Sutherland DP, Masterton RB, and Glendenning KK (1998b) Role of acoustic striae in hearing: reflexive responses to elevated sound-sources. *Behavioural Brain Research* **97**:1–12.
- Syka J and Radil-Weiss T (1971) Electrical stimulation of the tectum in freely moving cats. *Brain Research* **28**:567–572.
- Syka J and Straschill M (1970) Activation of superior colliculus neurons and motor responses after electrical stimulation of the inferior colliculus. *Experimental Neurology* **28**:384–392.
- Syka J, Popelář J, Kvašňák E, and Astl J (2000) Response properties of neurons in the central nucleus and external and dorsal cortices of the inferior colliculus in guinea pig. *Experimental Brain Research* **133**:254–266.
- Thompson GC and Masterton RB (1978) Brain stem auditory pathways involved in reflexive head orientation to sound. *Journal of Neurophysiology* **41**:1183–1202.
- Tolbert LP, Morest DK, and Yurgelun-Todd DA (1982) The neuronal architecture of the anteroventral cochlear nucleus of the cat in the region of the cochlear nerve root: horseradish peroxidase labelling of identified cell types. *Neuroscience* **7**:3031–3052.
- Tollin DJ (2003) The lateral superior olive: a functional role in sound source localization. *The Neuroscientist* **9**:127–143.

- Tollin DJ and Yin TCT (2002a) The coding of spatial location by single units in the lateral superior olive of the cat. I. Spatial receptive fields in azimuth. *Journal of Neuroscience* **22**:1454–1467.
- Tollin DJ and Yin TCT (2002b) The coding of spatial location by single units in the lateral superior olive of the cat. II. The determinants of spatial receptive fields in azimuth. *Journal of Neuroscience* **22**:1468–1479.
- Tollin DJ and Yin TCT (2003a) Spectral cues explain illusory elevation effects with stereo sounds in cats. *Journal of Neurophysiology* **90**:525–530.
- Tollin DJ and Yin TCT (2003b) Psychophysical investigation of an auditory spatial illusion in cats: the precedence effect. *Journal of Neurophysiology* **90**:2149–2162.
- Tollin DJ, Dent ML, and Yin TCT (2005) Psychophysical and physiological studies of the precedence effect and echo threshold in the behaving cat. In: Pressnitzer D, de Cheveigné A, McAdams S, and Collet L (eds). *Auditory Signal Processing: Physiology, Psychophysics, and Models*. Springer-Verlag, New York, pp. 429–435.
- Tsuchitani C and Boudreau JC (1966) Single unit analysis of cat superior olive S segment with tonal stimuli. *Journal of Neurophysiology* **29**:684–697.
- Vallbo ÅB and Johansson RS (1984) Properties of cutaneous mechanoreceptors in the human hand related to touch sensation. *Human Neurobiology* **3**:3–14.
- van Schaik A, Jin C, and Carlile S (1999) Human localisation of band-pass filtered noise. *International Journal of Neural Systems* **9**:441–446.
- Wallach H, Newman EB, and Rosenzweig MR (1949) The precedence effect in sound localization. *American Journal of Psychology* **57**:315–336.
- Warr WB (1966) Fiber degeneration following lesions in the anterior ventral cochlear nucleus of the cat. *Experimental Neurology* **14**:453–474.
- Warr WB (1972) Fiber degeneration following lesions in the multipolar and globular cell areas in the ventral cochlear nucleus of the cat. *Brain Research* **40**:247–270.
- Wenzel EM, Arruda M, Kistler DJ, and Wightman FL (1993) Localization using non-individualized head-related transfer functions. *Journal of the Acoustical Society of America* **94**:111–123.
- Wightman FL and Kistler DJ (1989a) Headphone simulation of free-field listening. I: stimulus synthesis. *Journal of the Acoustical Society of America* **85**:858–867.
- Wightman FL and Kistler DJ (1989b) Headphone simulation of free-field listening. II: psychophysical validation. *Journal of the Acoustical Society of America* **85**:868–878.
- Wightman FL and Kistler DJ (1999) Resolution of front-back ambiguity in spatial hearing by listener and source movement. *Journal of the Acoustical Society of America* **105**:2841–2853.
- Wurtz RH (1996) Vision for the control of movement. The Friedenwald Lecture. *Investigative Ophthalmology and Visual Science* **37**:2130–2145.
- Wytenbach RA and Hoy RR (1993) Demonstration of the precedence effect in an insect. *Journal of the Acoustical Society of America* **94**:777–784.
- Yan J and Ehret G (2001) Corticofugal reorganization of the midbrain tonotopic map in mice. *NeuroReport* **12**:3313–3316.
- Yan J and Ehret G (2002) Corticofugal modulation of midbrain sound processing in the house mouse. *European Journal of Neuroscience* **16**:119–128.
- Yin TCT (1994) Physiological correlates of the precedence effect and summing localization in the inferior colliculus of the cat. *Journal of Neuroscience* **14**:5170–5186.
- Yin TCT (2002) Neural mechanisms of encoding binaural localization cues in the auditory brainstem. In: Oertel D, Fay RR, and Popper AN (eds). *Springer Handbook of*

- Auditory Research*, Volume 15: *Integrative Functions in the Mammalian Auditory Pathway*. Springer-Verlag, New York, pp. 99–159.
- Yin TCT and Chan JCK (1990) Interaural time sensitivity in medial superior olive of cat. *Journal of Neurophysiology* **64**:465–488.
- Yin TCT and Kuwada S (1983) Binaural interaction in low-frequency neurons in inferior colliculus of the cat. III. Effects of changing frequency. *Journal of Neurophysiology* **50**:1020–1042.
- Yin TCT, Chan JCK, and Irvine DRF (1986) Effects of interaural time delays of noise stimuli on low-frequency cells in the cat's inferior colliculus. I. Responses to wideband noise. *Journal of Neurophysiology* **55**:280–300.
- Young ED, Spirou GA, Rice JJ, and Voigt HF (1992) Neural organization and responses to complex stimuli in the dorsal cochlear nucleus. *Philosophical Transactions of the Royal Society of London, series B, Biological Sciences* **336**:407–413.
- Young ED, Rice JJ, and Tong SC (1996) Effects of pinna position on head-related transfer functions in the cat. *Journal of the Acoustical Society of America* **99**:3064–3076.
- Zakarauskas P and Cynader MS (1993) A computational theory of spectral cue localization. *Journal of the Acoustical Society of America* **94**:1323–1331.
- Zhang Y and Suga N (2000) Modulation of responses and frequency tuning of thalamic and collicular neurons by cortical activation in mustached bats. *Journal of Neurophysiology* **84**:325–333.
- Zhang Y, Suga N, and Yan J (1997) Corticofugal modulation of frequency processing in bat auditory system. *Nature* **387**:900–903.

# Chapter 16

## Auditory Midbrain of Fish, Amphibians, and Reptiles: Model Systems for Understanding Auditory Function

ANDREW H. BASS, GARY J. ROSE, AND  
MICHAEL B. PRITZ

### 1. INTRODUCTION

We review studies that identify midbrain mechanisms in fish, amphibians, and reptiles that solve acoustic problems common to all vertebrates, including humans. The homologue of the inferior colliculus (IC) in fish, amphibians, and reptiles is the torus semicircularis (TS) (Nieuwenhuys et al. 1998). The TS, like the IC, “is a nexus of the auditory system because it processes and integrates almost all ascending acoustic information from lower centers, and it determines the form in which information is conveyed to higher regions in the forebrain” (Pollak et al. 2003). To remind the readers of this homology, we will use the term TS/IC.

Limited space precludes full review of the studies available on fish and amphibian auditory midbrain (Fay and Popper 1999). We therefore concentrate on neurophysiological studies of the encoding of vocalizations. Studies of nonmammalian audition have long focused on anuran amphibians (frogs and toads), showing how their auditory system is exquisitely adapted to detecting acoustic signals that diverge in spectral and temporal properties both within and between species. Studies of fish have concentrated on sound producing teleosts, whose detection and interpretation of vocal signals is essential for survival and reproduction.

We consider midbrain auditory circuitry devoted to encoding the temporal structure of vocalizations. “Temporal structure” refers to the modulations of the envelope of the acoustic waveform, either its amplitude (AM) and/or frequency (FM) over time. Temporal “fine structure” is the time course of the waveform beneath the envelope, and is most important when substantial energy is concentrated <500 Hz. We emphasize temporal codes because “the production and physical reality of animal signals lie in the time domain . . . the time varying acoustic waveform is the physical signal that is actually produced by the temporally patterned action of the motor system under ongoing control of the central

nervous system. . . . The two systems [vocal and auditory] co-evolved and we should expect them to share the same underlying code for signal generation and recognition" (Capranica 1992). Temporal processing has received increased attention in studies of humans (van Tasell et al. 1987; Shannon et al. 1995) and other mammals (Wang 2000; Klug et al. 2002; Nagarajan et al. 2002). We propose that studies of fish and anurans can reveal the most fundamental mechanisms underlying temporal coding in the more complex vocal communication systems of mammals as well as birds.

We also integrate a review of the reptilian auditory system, which is challenging because of the few sonic reptiles and the paucity of central neurophysiological studies for sonic or nonsonic species. The cardinal features of nonmammalian organization are conserved in mammals.

## 2. VOCALIZING FISH

Sound production is widespread among teleost fishes (Fine et al. 1977), the largest group of living fishes and of vertebrates (Pough et al. 2002). Among sonic/vocal fish, we concentrate on two distantly related teleosts, the marine batrachoidids and freshwater mormyrids, species whose midbrain TS/IC has received the most attention among teleosts (Bass and McKibben 2003). Batrachoidids are a single order and family of teleosts of 19 genera that include midshipman fish and toadfish (Nelson 1984). Mormyrids are a single family of 16 genera (Nelson 1984) known for their electrosensory and electrogenic abilities (Bullock and Heiligenberg 1986).

### 2.1. VOCAL COMMUNICATION SIGNALS OF TELEOST FISH

The acoustic communication signals of fish, like those of anurans (see Section 3.1), are diverse in their spectral (broadband or tonal harmonic series) and temporal (amplitude or frequency modulations) properties. Signals of fish are perhaps the simplest because the potential for diversity is constrained by the contraction rates of sonic muscles that vibrate or move one body part against another and the absence of a vocal tract and other resonators, although a resonant, gas-filled swimbladder may be involved in some species (Bradbury and Vehrencamp 1998). The simplicity of fish vocalizations can be mimicked with computer-synthesized signals to identify auditory sensitivities with either behavioral or neurophysiological methods. The acoustic signals of midshipman fish (*Porichthys notatus*) exemplify this simplicity. Males court females with long duration (seconds to >1 hour) advertisement calls, "hums," whose fundamental frequency (F0) is 90 to 100 Hz and which have prominent harmonics and essentially a flat envelope shape (Bass et al. 1999; Fig. 16.1). The temporal structure of hums differs from that of "grunts" which are brief (50 to 100 ms), broadband signals that midshipman males produce during agonistic encounters at rates of only 1 to 2 Hz (grunt "trains," Fig. 16.1). The pulse repetition rate



(PRR) in a grunt is similar to the F0 of hums. The temporal attributes of concurrent hums (acoustic beats) and AM-like grunt trains have shaped the design of behavioral and neurophysiological studies of auditory processing (Bass and McKibben 2003).

The temporal structure of mormyrid sounds has strongly influenced psychophysical and neurophysiological studies, where PRR is the vocal parameter of interest (Fig. 16.1). Mormyrids generate species-specific “moans,” “grunts,” and “growls” (Crawford et al. 1997). Moans are multiharmonic signals with differences in F0 of 200 to 400 Hz and 100- to 200-ms durations. Grunts are broadband, pulsatile signals with durations of 100 to 400 ms and PRRs of 30 to 60 Hz. Growls extend for seconds with PRRs of about 25 Hz.


## 2.2. *TORUS SEMICIRCULARIS AND INFERIOR COLLICULUS CIRCUITRY OF TELEOST FISH*

Like all terrestrial vertebrates, the teleost inner ear has three semicircular canals and two otolithic end organs, a sacculus and a utriculus (Popper and Fay 1999). Teleosts also have an otolithic lagena, a trait shared with anuran and urodele amphibians, reptiles, and birds. One or more of the otolithic organs may detect acoustic (and/or vestibular) stimuli (Popper and Fay 1999). The sacculus is the main auditory end organ in most teleosts (Popper and Fay 1999) and it encodes acoustic stimuli in amphibians (Lewis et al. 1982) and mammals (McCue and Guinan 1994). The macula neglecta is a nonotolithic end organ of unknown function in teleosts. As it serves an auditory function in cartilaginous fishes (Corwin 1981), it may also in teleosts. Some groups, including mormyrids, are categorized as “hearing specialists” because of accessory structures that enhance the detection of a propagating pressure wave. Although “hearing nonspecialists” or “generalists” such as batrachoidids lack these adaptations, their hearing sensitivity is comparable to that of specialists in the relevant frequency range (McKibben and Bass 1999).

While comparisons of hindbrain and forebrain auditory nuclei in teleost fish to those in tetrapods are not entirely resolved, the homology of TS to the IC is widely accepted (McCormick 1999). Studies of both toadfish and midshipman identify corresponding auditory centers at hindbrain, midbrain, and forebrain levels (Highstein et al. 1992; Bass et al. 1994, 2000, 2001; Edds-Walton et al. 1999; Goodson and Bass 2002). Comparable circuitry is found in vocal mormyrids (Kozloski and Crawford 1998; Precht et al. 1998; von der Emde and Precht 1999) and nonvocal teleosts in general (McCormick 1999) (Fig. 16.2).

The midbrain’s TS/IC has separate auditory and lateral line divisions. Although these divisions are often not obvious in Nissl- and Bodian-stained material, they are apparent using neurophysiological and tract-tracing methods. Medullary auditory and lateral line afferents enter the TS/IC via medial and lateral components, respectively, of the lateral lemniscus. The auditory region, nucleus centralis (TSnc, Fig. 16.2A; Knudsen 1977) occupies medial and dorsal TS/IC regions and caps the lateral line-recipient nucleus ventrolateralis (TSvl,

### A Midshipman



### B Mormyrids




Figure 16.1. Communication signals of vocal fish. (A) Oscillograms of a midshipman hum and grunt train on two time scales. (From Bass et al. 1999.) (B) Oscillograms of mormyrid grunts, moans and growls on two time scales. (From Crawford 1997.)

Fig. 16.2A). As in anurans and reptiles (see later), TSnc receives input from two medullary auditory nuclei (Fig. 16.2B), one an eighth nerve recipient nucleus, the descending octaval nucleus (DO), and a nearby nucleus, the secondary octaval nucleus (SO) that is DO's main target (Highstein et al. 1992; Bass et al. 1994; Mensinger et al. 1997; Edds-Walton et al. 1999). TSnc is reciprocally connected with multiple hindbrain and midbrain sites including DO, SO, the contralateral TSnc, and several sites involved in vocal production (see Section 2.3). Other brain stem targets of TSnc include a nucleus of the lateral lemniscus, while forebrain targets are the hypothalamus, the posterior tuberculum, and a dorsal thalamic nucleus projecting to the telencephalon (Fig. 16.2B). The precise role of these sites in auditory processing is unknown.

### 2.3. AUDITORY–VOCAL INTEGRATION


The pattern of auditory circuitry is similar in nonvocal and vocal teleosts with one important exception: auditory–vocal integration sites have been identified in batrachoidids. Midshipman and toadfish have an expansive vocal pacemaker–motoneuron circuit near the spinal–hindbrain junction that generates rhythmic, oscillatory-like activity similar in frequency and duration to that of natural vocalizations (Bass and Baker 1990, 1991). Each of a pair of midline sonic motor nuclei innervates the ipsilateral sonic muscle attached to the swimbladder walls, a gas-filled chamber involved in buoyancy control (Pough et al. 2002). Vocal pacemaker neurons cluster ventrolateral to the motoneurons and innervate both motor nuclei; their synchronous activation leads to the simultaneous contraction of both sonic muscles (Bass and Baker 1990).

Tracing studies coupled with electrical brain stimulation identify auditory–vocal integration sites at forebrain, midbrain and hindbrain levels (Bass et al. 1994; Goodson and Bass 2000, 2002). The TS/IC projects to low threshold midbrain tegmental activation sites where electrical stimuli evoke fictive vocalizations as monitored by intracranial recordings from the sonic nerve roots (Bass and Baker 1990; Goodson and Bass 2000). These sites include the periaqueductal gray (PAG) and the paralemniscal tegmentum (PL), both of which receive TS/IC input (Fig. 16.2B). As in mammals, the PAG and PL receive auditory input from the IC and are sites of electrically evoked vocalizations (Suga et al. 1973; Metzner 1996; Jürgens 2002). While neurophysiological studies have identified audio–vocal mechanisms in bats (Smotherman et al. 2003), such studies are just beginning in midshipman (Weeg and Bass 2003).

### 2.4. NEURAL CODING OF TEMPORAL PARAMETERS

Like those in other teleosts (Feng and Schellart 1999; Popper and Fay 1999), eighth nerve afferents in batrachoidids and mormyrids provide a periodicity code of frequency by phase locking to temporal modulations of the acoustic waveform's fine structure. As with anuran amphibians (see Section 3.4), the range of

**A**



**B**




Figure 16.2. TS/IC auditory circuitry in teleost fish. **(A)** A line drawing (*left side*) of a transverse section through the midbrain of midshipman fish (*Porichthys notatus*) showing the position of the auditory division, nucleus centralis (TSnc). Also shown is the lateral line division, nucleus ventrolateralis (TSvl). To the right is a line drawing of a dorsal view of the brain; the *vertical arrow* indicates the approximate midbrain level of the transverse section. **(B)** Schematic diagram of the major components of TS/IC circuitry. Following labeling of one eighth nerve with neurobiotin or biocytin, there is robust filling

frequency sensitivity is consistent with the energy spectrum of each species' calls (Fay and Edds-Walton 1997; McKibben and Bass 1999; Suzuki et al. 2002; Weeg et al. 2002; Sisneros and Bass 2003). Species differences in frequency sensitivity, like the F0s of vocalizations, often reflect differences in recording temperature (Bass et al. 2001). However, other studies of midshipman fish show seasonal changes in frequency sensitivity that are not temperature dependent (Sisneros and Bass 2003). Thus, the onset of male advertisement calling during the breeding season shows upward shifts among females in the frequency sensitivity of their saccular afferents such that it overlaps both the F0 and the most prominent harmonics of the male's hum; afferent sensitivity mainly overlaps only the F0 at other times. Midbrain neurons are more often sharply tuned than primary afferents in both nonvocal and vocal species (Crawford 1993; Lu and Fay 1996; Bodnar and Bass 1997). Seasonal shifts in midbrain frequency sensitivity among vocal species may yet be found.

Studies of midbrain auditory neurons in midshipman have focused on the encoding of acoustic beats because of their prominence in this species' acoustic environment (Bodnar and Bass 1997; Bass et al. 1999). Neighboring males that concurrently hum, generate acoustic beats with a modulation rate determined by the difference frequency (dF) between their F0s (Fig. 16.3). Beats are imitated by two concurrent tones with dFs like the natural range ( $<10$  Hz). Playback studies show that beat dF and modulation depth are important parameters for discriminating hums from beats and other vocalizations with low modulation rates (grunt trains, Fig. 16.1) (McKibben and Bass 1998, 2001a).

There is a peripheral-to-midbrain transformation in the temporal coding of a beat's dF and modulation depth. Saccular afferents exhibit robust phase locking to a beat's component tones, and code its modulation rate dF in their temporal modulations of spike rate and synchronization (McKibben and Bass 2001b). By contrast, TS/IC neurons show poor phase locking to tones, but robust phase locking to dFs over their natural range, which is  $\pm 10$  Hz (Bodnar and Bass 1997, 1999, 2001a). About 35% of midbrain units are broadly tuned across the behaviorally relevant range of  $\pm 10$  Hz (broad dF-selective, Fig. 16.3B). The remaining units show symmetrical peaks in dF selectivity (narrow dF-selective, Fig. 16.3C); 25% of these units exhibit asymmetrical tuning (narrow dF sign-selective, Fig. 16.3D). This synchronization code cannot be explained by either spike rate profiles (Fig. 16.3E–G) or frequency tuning. Spike synchronization increases with increasing modulation depth; primary afferents increase in both spike rate and spike synchronization (McKibben and Bass 2001b).

---

Figure 16.2. *Continued*

of medullary (descending octaval; DO) neurons bilaterally, likely due to transneuronal transport (Bass et al. 1994, 2000). Bilateral projections are reported in several teleost taxa (McCormick 1997; Kozloski and Crawford 1998; H. Straka and R. Baker, unpublished observations). ?, evidence for a projection that has not been demonstrated conclusively. For abbreviations please see the list.

# Acoustic Beat




Figure 16.3. Encoding of acoustic beats. **(A)** The temporal waveform of two concurrent hums on two time scales. **(B–G)** Single-unit responses in the auditory midbrain (TSnc) of midshipman fish to beat stimuli. Each stimulus was formed by an F1 of 90 Hz and an F2 of  $90 \pm 2$  to 20 Hz (see legend in lower right). **(B–D)** Plots of vector strength of synchronization (vs.  $\pm$ SE) over dF stimulus range. Units show a range of dF selectivities, but exhibit low synchronization to the individual components (F1, F2) of a beat. **(E–G)** Corresponding average spike rate ( $\pm$ SE) curves for the same units shown in **(B–D)**. **(B–G)** from Bodnar and Bass 1997.)

Other experiments show two midbrain populations of AM-encoding neurons. About 65% of TS/IC neurons have a similar tuning profile for the modulation rate of either beats or AM signals produced by the modulation of a single acoustic waveform's amplitude over time (Bodnar and Bass 1997). Thus, despite differences in fine structure between these signals (beats are formed by the interaction of two waveforms), their envelope shape is often encoded in a similar way. Importantly, 35% of TS/IC neurons distinguish AM-like signals from beats that could contribute to a behavioral distinction between beats and grunts (Fig. 16.1).

In contrast to eighth nerve afferents that show little if any adaptation with increasing sound duration (McKibben and Bass 1999), midbrain neurons show a range of adaptation rates (Bodnar and Bass 2001b). While the lack of evidence for duration selective units is consistent with the behavioral evidence showing that a wide range of tone durations are recognized as hums (McKibben and Bass 1998, 2001a), spike rate variability across midbrain units may contribute to a population code (Bodnar and Bass 2001b).

Mormyrid psychophysical experiments show interclick interval discrimination consistent with the natural variation of PRRs (Marvit and Crawford 2000; Fletcher and Crawford 2001; Fig. 16.1). The click interval sensitivities of midbrain neurons overlap the PRRs of mormyrid moans (5-ms intervals/200 Hz), grunts (20-ms intervals/50 Hz), and growls (40-ms intervals/25 Hz) (Crawford 1997). This temporal code is apparently formed in the midbrain, since medullary neurons have a broad range of interclick interval sensitivities (Kozloski and Crawford 2000). Thus, like midshipman, mormyrid studies provide evidence for midbrain temporal coding.

### 3. AUDITORY MIDBRAIN OF ANURANS

Acoustic communication is vital in the reproductive biology of most anurans. Hence, several robust behaviors can be used to address questions of auditory processing. Phonotaxis and evoked calling paradigms have been particularly important for investigating the auditory system's analytic ability to code the spectral and temporal structure of anuran calls. We emphasize behavioral work that relates to our understanding of the function of the anuran TS/IC (Fay and Popper 1999).

#### 3.1. VOCAL COMMUNICATION SIGNALS OF ANURANS

Spectrally, most anuran communication signals are narrowband "tonal," broadband, dual-peaked, a harmonic series, or some combination of these. Dual-peaked signals resemble the formant structure of human speech sounds, particularly vowels, which can generally be identified based on their first two formants. In such anuran vocalizations, each spectral peak is within the frequency sensitivity range of one of the two peripheral auditory organs that function primarily for detecting communication signals, the amphibian and basilar

papillae; typically, the higher frequency peak is close to the tuning of the basilar papilla. The bullfrog (*Rana catesbeiana*), for example, produces calls with broad spectral “formants” centered at about 200 Hz and 1400 Hz (Capranica 1965, 1966). The precise positions of these spectral maxima differ among the calls of the various species within this genus. Whereas most interspecific calls differ in spectral and temporal structure, intraspecific call types often are spectrally highly similar and differ chiefly in temporal structure; calls of closely related cryptic species may differ exclusively in their temporal structure.

The basic temporal features of a vocalization are its duration and rise/fall characteristics; the temporal structure of a tonal call can be described by these parameters. Many communication signals of anurans, however, show distinctive, more complex, AM patterns. Sometimes AM reflects the interactions of harmonically related spectral components of the vocalization. In most cases, however, AM results from passive and/or active mechanical processes such as the contractions of thoracic musculature and muscle-related vibrations of the cartilaginous laryngeal elements (Martin 1971; Schneider 1988). The diverse AM patterns probably result from complex active control and passive properties of laryngeal structures. Further involvement of skeletal and respiratory-related musculature in vocalization is absent in teleosts, but enhances potential anuran call diversity and complexity (and in reptiles, birds, and mammals; Bradbury and Vehrencamp 1998). Amplitude modulation ranges from a nearly sinusoidal periodic pattern to discrete sound pulses. The rise and fall characteristics of individual pulses can vary even between the calls of closely related species, and, with PRR, constitute the primary temporal cues that enable discrimination between conspecific and heterospecific calls (Gerhardt 1982, 1988). As discussed later, many intraspecific call types differ virtually exclusively in PRR; individual pulses have similar shapes and spectral composition. Where pulse number is similar between call types, call duration is inversely related to PRR. Anuran PRR is physiologically equivalent to PRR in sonic fish, the number of sound pulses per second. The latter is also the case for F0 in teleosts, but F0 among anurans is determined by vocal cord vibratory properties (Martin 1971). PRR and F0 in teleosts, like PRR in anurans, are established by a central pattern generator (anurans: Schmidt 1976; teleosts: Bass and Baker 1990).

### 3.2. BEHAVIORAL STUDIES

Evoked calling experiments on bullfrogs demonstrate that energy must be present in the low- and high-, but not in the middle-, frequency regions for stimuli to maximally elicit responses from males (Capranica 1965, 1966). These experiments and others support the concept of a neural AND gate that detects simultaneous low- and high-frequency energy peaks.

Many anuran communication sounds have nearly identical spectral composition but differ in temporal structure. This disjunction occurs in the calls of some closely related species, but is found more commonly in the intraspecific call repertoire of particular species. The intraspecific call types of the Pacific treefrog (*Hyla regilla*) are differentiated virtually completely by temporal structure. Pa-



cific treefrogs produce an “encounter” (aggressive) call and two types of advertisement calls (Allen 1973). These calls consist of pulses, each 10 to 12 ms in duration, repeated at about 25 pulses/s (pps) and 100 pps, respectively (Fig. 16.4A); the exact value depends on temperature. To alter PRR, they change the pulse duty cycle (ratio of pulse duration to interpulse interval). Because the advertisement and encounter pulses are nearly identical in spectral composition, shape, and duration, the calls differ primarily in PRR; calls also differ in duration but this is quite variable (Rose and Brenowitz 2002). Behavioral studies define two discrete sensory “channels” for processing the two call types (Brenowitz and Rose 1994; Rose and Brenowitz 1997). These findings suggest neural filters that are highly selective for PRR. Other behavioral studies (Rose and Brenowitz 2002) indicate that temporal filters tuned to the fast PRRs, as found in the advertisement calls of *H. regilla*, respond only after several consecutive pulses, delivered with 10-ms interpulse intervals (time between onsets of adjacent pulses), have occurred.

An excellent example of temporal differentiation in the calls of closely related anurans is the advertisement calls of two species of gray treefrogs, *H. versicolor* and *H. chrysoscelis*, which differ only by chromosome number and call structure (Gerhardt 1982). Their advertisement calls are spectrally essentially identical, but differ in temporal structure (Fig. 16.4B). The *versicolor* advertisement pulses have slower rise time, longer duration and slower PRR. Female gray treefrogs use both pulse shape and PRR to select a conspecific’s advertisement call over that of the other species (Gerhardt 1988; Diekamp and Gerhardt 1995).

Frequency modulation (FM) is especially common among certain anuran genera (e.g., *Leptodactylus* and *Physalaemus*). The call direction and rate of frequency change is species-specific, with some using upsweeps and others downsweeps. The FM frequency range in these calls corresponds to that represented by the sensitivity range of the amphibian papilla. Interestingly, species such as *L. labialis* have calls with concurrent AM and FM (Straughan and Heyer 1976). In the advertisement calls of the Tungara frog, *P. pustulosus*, frequency sweeps downward exponentially from 900 Hz to 300 Hz (Fig. 16.4C) as signal amplitude decreases smoothly. In phonotaxis studies with *P. pustulosus*, females chose the forward call over a time-reversed version (Ryan 1983). Behavioral (evoked calling) studies of males showed that they can discriminate, in an intensity-independent fashion, between synthetic signals that differ only in FM direction (Rose et al. 1988).

### 3.3. ANURAN TORUS SEMICIRCULARIS/INFERIOR COLLICULUS CIRCUITRY

Anurans possess amphibian and basilar papillae and there is evidence for an acoustic function for the sacculus. The amphibian papilla is unique to amphibians, while the basilar papilla may be homologous to that of reptiles, birds, and mammals (Lewis and Narins 1999; McCormick 1999). Anuran brain stem auditory regions are homologous to the major auditory areas in the mammalian brain stem (Wilczynski 1988; Feng and Lin 1991; ten Donkelaar 1998a; Mc-




Figure 16.4. Anuran communication signals. (A) Oscillograms of encounter and advertisement calls of *Hyla regilla*; advertisement calls consist of diphasic and monophasic types. (B) Oscillograms of advertisement calls of *H. versicolor* and *H. chrysoscelis*. (C) Spectrograms of advertisement calls of *Physalaemus pustulosus*; the frequency-modulated (whine) portion can be followed by one or more harmonically rich “chucks.”

Cormick 1999). The TS/IC has laminar, principal, and magnocellular subdivisions (Fig. 16.5A). Afferents include the dorsal lateral nucleus (DLN, first-order auditory region), the superior olivary nucleus (SO, second-order target of auditory input from the DLN), and the lateral lemniscal nucleus (nLL, also termed the superficial reticular nucleus) near the reticular formation. Portions of the TS/IC also receive afferents from the pretectum, preoptic area and hypothalamus. The TS/IC has reciprocal connections with SO, the nLL and the contralateral TS/IC. The TS/IC projects to three thalamic nuclei (anterior, central, and posterior), some of which also receive SO input (Fig. 16.5B).

### 3.4. SPECTRAL PROCESSING IN ANURAN TORUS SEMICIRCULARIS/ INFERIOR COLLICULUS

The range of spectral sensitivity of units in the anuran TS/IC parallels that seen in the auditory nerve. This range is represented by the frequency sensitivities of



**A****B**

Figure 16.5. TS/IC auditory circuitry in anurans. **(A)** A line drawing (*to the left*) of a transverse section through the midbrain of the bullfrog, *Rana catesbeiana*, showing the position of the laminar (TSnl) and principal (TSpr) nuclei. A magnocellular division has no clearly defined boundaries but includes large cells that lie ventral and lateral to TSnl and TSpr. To the right is a line drawing of a dorsal view of the brain; the *vertical arrow* indicates the approximate midbrain level of the transverse section. **(B)** Schematic diagram of the major components of TS/IC circuitry as described in the text. The TS/IC also receives contralateral SO and nLL input and has bilateral outputs to SO and nLL (not shown for simplicity, but see Feng and Lin 1991). [Line drawings in **A** and **B** based, respectively, on Northcutt (1974) and Wilczynski and Northcutt (1983).]

the amphibian and basilar papillae (Lewis and Lombard 1988). Basilar papilla units are tuned to higher frequencies and excitatory responses cannot be suppressed by higher frequency tones. The tonotopically organized amphibian papilla represents the low- and mid-frequency range, with only the low-frequency sensitive fibers, those innervating rostral segments, showing two-tone suppression. Like auditory-nerve fibers, many TS/IC units have a single region of frequency sensitivity; some cells even show V-shaped excitatory frequency tuning curves. Unlike eighth-nerve fibers, however, many TS/IC neurons have well developed inhibitory receptive fields that are not attributable to peripheral two-tone suppression (Walkowiak 1980; Fuzessery 1988; Hall 1994). Thus, units most sensitive to middle frequencies can be inhibited by frequencies below and/or above their excitatory regions. As a result, a major transformation in coding is that many individual TS/IC neurons in anurans show “level-tolerant” frequency sensitivity in which the excitatory frequency band is a narrow region independent of stimulus intensity; in the most extreme cases, excitatory receptive fields consist of a discrete region in frequency-amplitude “space.” Blocking  $\gamma$ -aminobutyric acid-A ( $GABA_A$ ) receptors with bicuculline showed that neural inhibition underlies these complex receptive fields (Hall 1994).

Approximately 3% of TS/IC neurons have two regions of frequency sensitivity (W-shaped tuning curves); in some cases this W function is shallow, indicating that the neuron simply shows very broad frequency tuning (Walkowiak 1980). Although most neurons with W-shaped tuning curves respond when energy is present in either of their excitatory receptive field regions, a few show properties characteristic of an “AND” logical operation, responding only, or in a facilitated manner, when energy is in both frequency bands. Units that require energy in both low- and high-frequency bands represent the neural correlate of the AND gate proposed as the filter matched to the spectral structure of the advertisement call (bullfrogs and leopard frogs). This spectral selectivity can be more generally classified as “formant selectivity.” Such selectivity is rare in the anuran TS/IC, but robust in the posterior thalamic nucleus (Fuzessery and Feng 1983).

### 3.5. TEMPORAL PROCESSING IN ANURAN *TORUS SEMICIRCULARIS/INFERIOR COLLICULUS*

As discussed previously, the temporal structure of anuran communication signals is important in their reproductive biology. What neural mechanisms underlie the abilities of anurans to discriminate between calls that differ in temporal structure? We next consider how patterns of AM are represented and processed in the auditory system.

Constituting early evidence suggesting midbrain specializations for processing AM signals, some neurons in *R. ridibunda* preferred amplitude-modulated tones to pure tones, and maximally synchronized their spikes to a particular phase of the modulation cycle at rates of 10 to 20 Hz AM (Bibikov and Gorodetskaya 1980). Also, TS/IC units were recorded that responded to a sequence of sound pulses, but not to pure tones (Walkowiak 1980). Sinusoidally amplitude-

modulated (SAM) white noise stimuli were used to further investigate how AM is represented in anurans. With white noise as the carrier, long-term spectral properties of the stimulus do not change with AM rate; selective neural responses, therefore, reflect temporal, not spectral, stimulus features. While auditory-nerve fibers coded AM rate in their discharge periodicities, their mean spike rate was independent of the AM rate; this result is perhaps not surprising because stimulus energy and spectral properties were constant across AM rate. In the anuran TS/IC, however, the firing rate of most neurons depended strongly on the AM rate (Rose and Capranica 1983, 1985). This conclusion is supported by many studies in which AM rate or sound pulse repetition rate was varied (Walkowiak 1988; Diekamp and Gerhardt 1995). For sinusoidal AM, the temporal selectivity of cells is level-tolerant and is classed as low-pass, high-pass, band-suppression, or band-pass. About one third of the cells are band-pass types, responding best over a narrow range of modulation rates (“AM-tuned”). Using tonal carriers as stimuli, almost 70% of TS/IC cells are band-pass to the AM rate (Alder and Rose 2000). Similarly, band-pass selectivity for the AM rate is seen in avian (Albert et al. 1989) and mammalian midbrain cells (Langner 1992; Krishna and Semple 2000; Langner et al. 2002).

Remarkably, anuran TS/IC neurons most sharply tuned to AM poorly encode the modulation rate in their discharge periodicity (quantified by calculating a synchronization coefficient) (Rose and Capranica 1984; Rose 1995). Similarly, in mammals, there is a significant reduction in spike synchronization to the AM stimulus at the midbrain (Langner 1992; Eggermont 2001; Frisina 2001). The transformation from a peripheral periodicity code of AM rate to a temporal filter representation in the midbrain has been observed for fish (see Section 2.4), frogs (Rose and Capranica 1983, 1985), birds (Langner 1981), and mammals (Langner and Schreiner 1988; Pinheiro et al. 1991; Condon et al. 1996). The distribution of AM tuning values for anuran TS/IC neurons is species-specific and nicely related to the range of PRRs in their calls (Rose and Capranica 1984; Rose et al. 1985).

The mechanisms that underlie AM selectivity are only beginning to be understood. Theoretically, band-pass selectivity for SAM waveforms might arise from sensitivity to stimulus rise time and duration. Although some anuran TS/IC neurons are sensitive to stimulus rise time and/or duration, these properties contribute little to their AM selectivity (Gooler and Feng 1992; Alder and Rose 2000). For example, band-pass selectivity exists for square-wave AM, where pulse rise time is constant across AM rates.

Rise-time sensitivity accounts only for the slight differences in the shapes of band-pass functions for square-wave AM vs. sinusoidal AM; at slow AM rates, responses are weaker for sinusoidal AM because cells prefer fast rise times. This preference appears to originate in auditory regions caudal to the anuran TS/IC. Neurons in DLN and SO that respond phasically to tone bursts prefer rise times <25 ms (Hall and Feng 1988, 1991; Condon et al. 1991). Their phasic response properties may account for their diminished responses to slow SAM rates (Hall 1994). Bicuculline injections, presumably reversing their phasic properties, can

transform neurons from band-pass to low-pass, and from high-pass to all-pass (Hall 1994).

Processes other than just rise time and duration sensitivity must therefore underlie the band-pass selectivity of TS/IC neurons. This conclusion is perhaps not surprising considering that the intraspecific calls of many anurans differ primarily in PRR, not pulse shape or duration, for example, *H. regilla* (see Section 3.2). As predicted from the behavioral studies described previously, extracellular single-unit recordings confirm that one class of TS/IC neurons responds highly selectively to fast PRRs, as in the advertisement calls, but not to PRRs seen in the encounter call (25 pps) (Alder and Rose 1998, 2000). These neurons respond only after a threshold number of pulses, each separated by a cell-specific interpulse interval (time between the onsets of consecutive pulses) (Fig. 16.6A). The salient temporal feature for eliciting responses is the number of consecutive “correct” intervals rather than mean pulse rate, suggesting an interval-counting process (Edwards et al. 2002). This integration process may account for the impressive selectivity of these units for AM or PRR. Remarkably, a single interval that falls outside a particular range can reset the integration process (Fig. 16.6B–D). This integration process bears some similarity to “sequence sensitivity” in cat auditory cortex (Brosch and Schreiner 2000) and to combination sensitivity in bat auditory cortex (O’Neill and Suga 1979) and in songbird forebrain (Margoliash 1983; Margoliash and Fortune 1992). An integration process of this nature may contribute to AM rate discrimination in humans (Lee 1994).

Surprisingly, the enigmatic “band-suppression” neurons, so called because they respond to slow and fast, but not intermediate, AM rates, are also interval-integrating types (Edwards et al. 2002). Relative to band-pass neurons, these cells require few (median: 2) intervals and have broad interval tolerance. Thus, mechanistically, and perhaps functionally, band-suppression neurons and band-pass interval-integrating neurons may be one physiological class.

Another class, the “recovery neurons,” respond well to slow PRRs, but only weakly at the onset of faster PRR stimuli (Alder and Rose 2000). At slow PRRs and AM rates, they respond phasically to each pulse and, therefore, are low-pass when pulse shape, duration and number are held constant, and band-pass to SAM (Fig. 16.7). These findings suggest that recovery processes contribute to their PRR selectivity; after the excitation from a stimulus pulse, a recovery period is required before the next pulse can excite the cell. The phasic nature of these responses may reflect inhibition, as bicuculline can transform neurons from band-pass to low-pass (Hall 1994).

Most TS/IC neurons selective for AM rate are either recovery- or interval-integrating types. Thus, we suggest there are primarily two physiologically distinct classes of AM selective neurons. The mechanisms that underlie the integration process in the “interval-counting” neurons and the recovery process of the “recovery-type” cells are unknown. Temporally selective TS/IC neurons are hypothesized to project to the central thalamic nucleus (Hall and Feng 1987). Thus, parallel spectral and temporal processing continues at the posterior and




Figure 16.6. Properties of integration-type midbrain neurons that respond selectively to fast PRRs. **(A)** Raster plots and histograms of responses of a unit to multiple repetitions of stimuli having three (*top*) or four (*bottom*) pulses. **(B)** Histograms of responses to stimuli consisting of nine consecutive intervals of 10 ms each (*top*), or alternating 5-ms or 15-ms intervals (*bottom*). **(C)** Effects of a single long (30-ms) interval in resetting the interval-integrating process; this unit had an interval number threshold of 8. **(D)** Normalized response levels of two units as a function of the duration of a single interpulse interval embedded in a series of optimal intervals. (Adapted from Edwards et al. 2002.)

central thalamic nuclei, respectively (Hall and Feng 1987). Further spectral processing involves generating “AND-type” selectivity, where neurons respond only if energy in two spectral bands occurs simultaneously. Central nucleus neurons show temporal selectivity; however, further work is needed to delineate how it differs from temporally selective TS/IC neurons.

## 4. REPTILIAN AUDITORY MIDBRAIN

Residing in the reptilian tectal plate are the modalities of vision, somatic sensation, and audition. Although each modality is represented in an individual target area, sensory channels are not entirely separate and multimodal neurons exist (Kachunts 1982; Belekhova et al. 1985). Vision is represented in the optic tectum (Ulinski et al. 1992) and somatic sensation is located in the caudolateral midbrain roof (Ebbesson 1967; Pritz and Stritzel 1989). As with fish and amphibians, the TS is the midbrain auditory area and the reptilian homologue of the mammalian IC (ten Donkelaar 1998b).

### 4.1. MORPHOLOGY AND SUBDIVISIONS OF REPTILIAN TORUS SEMICIRCULARIS/INFERIOR COLLICULUS

The morphology and complexity of the TS/IC varies among species, reflecting the role audition plays in their behavior. The optic tectum rings the TS/IC on all but its ventral surface where it is bounded by the midbrain tegmentum (Díaz et al. 2000). The tectal ventricle separates the optic tectum from the TS/IC. Internal subdivisions exist in the TS/IC. The main auditory portion is the central nucleus (TSnc, Fig. 16.8A). The deep layers of the optic tectum are continuous with TSnc. This feature has been used to distinguish an external or lateral nucleus (also known as nucleus laminaris), continuous with the deep tectal layers, from TSnc (Pritz 1974; Browner and Rubinson 1977; Browner and Baruch 1984) (Fig. 16.8A). In other reptiles an additional subdivision termed cortical (Browner and Rubinson 1977) or superficial (Browner and Baruch 1984) has been described, and in some reptiles (Tegu lizards) the central nucleus is subdivided into two regions (Browner et al. 1981). The topographic relations and individual cellular morphology of each of these subdivisions have been detailed in several species. Selected features of neuronal morphology (Table 16.1) and immunocytochemical/histochemical properties (Table 16.2) are summarized.

### 4.2. REPTILIAN TORUS SEMICIRCULARIS/INFERIOR COLLICULUS CIRCUITRY

The basilar papilla is the main auditory organ in reptiles; its morphology varies widely especially among lizards (Manley 2000). Many brain stem subdivisions project to the central nucleus of the torus, which is part of a larger scheme (Fig.






Figure 16.7. “Recovery-type” neuron. (A) This unit showed band-pass properties to sinusoidal AM (SAM) rate, but was low-pass for stimuli when only pulse repetition rate was varied (variable duty cycle [VDC], pulse number, shape, and duration were constant). (B) Phasic responses of this unit to stimuli of these two types. (From Alder and Rose 2000.)

Table 16.1. Cell types within the TS/IC in several reptiles.

Subdivision	Somatic shape	Somatic diameter ( $\mu\text{m}$ )	Spines	Species <sup>a</sup>
Central nucleus	Round	6–25	Few	Turtle
		6–12	Many	Skink
		13–21	Some	Lizard
	Triangular	15–27	Rare	Turtle
		14–28	Few	Skink
		21–27	Moderate	Lizard
		Fusiform	10–26	None
	20–30		Few	Skink
	Laminar nucleus	Ovoid	19–21	—
9–15			Few	Turtle
8–12			Many	Skink
Triangular		14–20	Few	Lizard
		20–40	None	Turtle
		18–28	Few	Skink
		19–36	None	Lizard
Fusiform		20–40	None	Turtle
		20–40	None	Skink
	18–33	Few	Lizard	

<sup>a</sup>Based on data on turtles (Browner et al. 1981), skinks (Browner and Baruch 1984), and lizards (Browner and Rubinson 1977).

16.8B). Brain stem projections to the central nucleus arise bilaterally from the cochlear nuclei, with a contralateral predominance (Foster and Hall 1978; Belekova et al. 1985; Künzle 1986), bilaterally from the superior olive (Foster and Hall 1978; Belekova et al. 1985) and from the nucleus of the lateral lemniscus (ipsilateral in iguana, Foster and Hall 1978 and bilaterally in turtles, Belekova et al. 1985). Ascending connections of TS terminate bilaterally, with an ipsilateral dominance in a caudomedial thalamic nucleus, nucleus reuniens (Pritz 1974; Browner 1983; Belekova et al. 1985) or medialis (Foster and Hall 1978). Additional ascending terminations target a nucleus embedded in this tectoreuniens fiber tract (Pritz 1974) and a separate, lateral portion of nucleus reuniens, the pars diffusa, as opposed to the heavier terminations in the pars centralis (Pritz 1974).

Descending connections of the TS/IC have not received much attention experimentally. Limited data describe descending efferents as terminating primarily in the ipsilateral midbrain and medullary reticular formation and nucleus of the lateral lemniscus (Foster and Hall 1978). In addition to commissural projections to the opposite central nucleus, axons terminate in the ipsilateral deep layers of the optic tectum (Foster and Hall 1978; Belekova et al. 1985) (see Chapter 6).

Experimental analyses of nucleus laminaris neural circuitry are likewise limited. Efferents have been traced to the spinal cord (ten Donkelaar and de Boer-van Huizen 1978; Butler and Bruce 1981; Follett 1989), the inferior raphe (ten




Figure 16.8. TS/IC auditory centers in reptiles. **(A)** A line drawing (*to the left*) of a transverse section through midbrain of the crocodile, *Caiman crocodilus*, showing the position of the central (TSnc) and laminar (TSnl) nuclei. To the right is a line drawing of a dorsal view of the brain; the *vertical arrow* indicates the approximate midbrain level of the transverse section. **(B)** Schematic diagram of the main components of TS/IC circuitry as described in the text. [Line drawings based on Pritz (1974).]

Table 16.2. Immunocytochemical properties of the torus semicircularis of turtles<sup>a</sup>

Marker	Subdivision			
	Central Nucleus		Laminar Nucleus	
	Fibers/Boutons	Cells	Fibers/Boutons	Cells
5-HT	+	—	+++	—
TH	+	—	+++	—
SP	+	—	+++	+
<i>m</i> ENK	+	—	+++	+
NPY	+++	+	+++	+
NADPH-d	++	++	+++	+++

Density: +, Light; ++, moderate; +++, heavy; —, not described.

*m*ENK, met-enkephalin; 5-HT, serotonin (5-hydroxytryptamine); NADPH-d, nicotinamide adenine dinucleotide-diaphorase; NPY, neuropeptide Y; SP, substance P; TH, tyrosine hydroxylase.

<sup>a</sup>Based on data of Belekhoval et al. (2002).

Donkelaar and de Boer-van Huizen 1987), and the anterior dorsal ventricular ridge (ten Donkelaar and de Boer-van Huizen 1988), while afferents come from the spinal cord (Künzle and Woodson 1982).

#### 4.3. NEUROPHYSIOLOGICAL STUDIES OF AUDITORY TORUS SEMICIRCULARIS/INFERIOR COLLICULUS

Electrophysiological studies have investigated response to sound in several reptilian groups. Auditory units have been recorded in lizard central nucleus (Kennedy 1974; Foster and Hall 1978) and in *Caiman crocodilus*, where the most detailed information is available (Manley 1971). In *Caiman*, TS/IC is organized tonotopically. Most units have phasic responses with the characteristic frequency of individual units varying from 70 to 1850 Hz. Tuning curves often were V-shaped. Although unimodal auditory units were identified in the torus of turtles, some cells were bimodal, responding to both audition as well as somatic sensation (Kachunts 1982). Characteristic frequencies ranged from 40 to 6000 Hz with optimal frequencies between 100 and 400 Hz. Three types of discharge patterns were observed: phasic, burst, and sustained (Kachunts 1982).

#### 4.4. BRAIN STIMULATION AND BEHAVIOR STUDIES

Regardless of whether some central nucleus units are bimodally responsive to both auditory and somatic sensory stimuli, audition is the dominant modality. However, vocalization, as elicited by electrical stimulation, is localized to the lateral nucleus (Kennedy 1975). Not surprisingly, the lateral nucleus rather than the central nucleus contains androgen receptors (Tang et al. 2001). This is potentially significant because geckos are the only lizards capable of producing

complex sounds involved in social behavior (Carr 1992). Although not yet examined in geckos or other lizards, ultrastructural analyses in turtles, which do vocalize (Carr 1992), revealed lipid-like droplets and agranular endoplasmic reticulum in certain nucleus laminaris neurons. These are features of steroid secreting cells (Drakontides and Browner 1986).

#### *4.5. COMMON FEATURES OF TORUS SEMICIRCULARIS/INFERIOR COLLICULUS ORGANIZATION IN REPTILES*

This brief review of relevant anatomical, physiological, and behavioral features of the TS/IC notes significant areas where data are either limited or lacking for both the central nucleus as well as the external or laminar nucleus. Nevertheless, certain key features are salient. First, much of the midbrain neural circuitry for audition among reptiles is similar although not identical. Second, nucleus laminaris is probably involved in vocalization and this behavioral feature is likely to be influenced by steroid hormones. Third, unlike in mammals (see Chapter 8), no long efferent tracts of telencephalic origin (either cortex, dorsal ventricular ridge, or striatum) to the torus have yet been described in any reptilian group, or in fish and amphibians.

### 5. CONCLUDING COMMENTS AND DIRECTIONS OF FUTURE WORK

Fish model systems present the simplest and perhaps most ancient extant example of how a vertebrate auditory system encodes context-dependent vocal signals. In some teleosts, like the species discussed here, the physical attributes of the acoustic waveform are essentially a direct translation of the temporal attributes of a central pattern generator (CPG). This work has demonstrated a vocal-acoustic network that closely resembles the organizational pattern in mammals, while showing how that circuitry can modulate the output of the vocal CPG. Behavioral studies demonstrate that the recognition of temporal structure is an essential acoustic task: studies of midshipman fish emphasize the importance of recognizing perturbations of the acoustic waveform's envelope shape for distinguishing advertisement calls (hums) that essentially lack any envelope modulation from AM-like signals (beats and grunt trains), while studies of mormyrids provide evidence for the recognition of interclick intervals that compare closely to the PRRs of their calls. As in other sonic vertebrates, the range of frequency sensitivity of both eighth nerve afferents and midbrain neurons is consistent with the energy spectrum of each species' calls. Midbrain auditory neurons in midshipman and mormyrids are well adapted to encode the temporal structure of acoustic waveforms: the midshipman's acute behavioral sensitivity to dF, the modulation rate of acoustic beats, is matched by dF-selective neurons; the mormyrid's sensitivity to PRR is paralleled by interval-selective neurons.

The biophysical mechanisms underlying midbrain temporal sensitivity are unknown but studies of the electrosensory system of other teleosts (Fortune and Rose 2001) suggest that fish model systems hold great promise in this regard. Given the evolutionary emergence of the most fundamental mechanisms of vertebrate audition in fishes, such studies will influence both our basic understanding of the evolution and development of auditory and audio–vocal mechanisms and the functional organization of genetic model systems (zebrafish).

The temporal structure of anuran vocalizations are more complex than those of fish, in part because of the specialized coupling between vocal organs, respiratory-related musculature and resonant chambers that enhance the coupling of the vibrations of their vocal organs to terrestrial habitats (Bradbury and Vehrencamp 1998). Intraspecific call types often are spectrally similar and differ only in temporal structure; moreover, the calls of closely related species may only differ in their temporal structure. Studies in anurans first showed that a peripheral periodicity code of AM is transformed in the midbrain into a temporal filter representation. The distribution of AM tuning values for anuran TS/IC neurons are species-specific and match the range of call PRRs. This resembles the peripheral-to-midbrain transformations for the encoding of dF and PRR in fish. Further studies in anurans have revealed two populations of AM neurons encoding interpulse intervals. Theoretical studies suggest that the interplay between excitation and inhibition might underlie temporal selectivity, particularly interval analysis, in hearing (Buonomano 2000; Large and Crawford 2002). In these models, interval analysis stems from differences in the timing, time course and plasticity (Buonomano 2000) of excitatory and inhibitory inputs. GABAergic inhibition is present in the anuran TS/IC (Hall 1994, 1999), but its role in temporal processing is incompletely understood. Intracellular studies should be pivotal in testing these models and revealing the mechanisms that underlie temporal selectivity.

As noted earlier, the temporal structure of acoustic signals is characterized primarily by the changes in amplitude and frequency over time. Anurans offer the opportunity to study the neural representations and processing of these two forms of temporal information in isolation because particular species specialize in the use of AM or FM. Although there have not yet been any neurophysiological investigations of FM processing in anurans, they provide tractable systems in which to study how FM information is represented and the mechanisms that underlie FM selective behavioral responses.

Studies of reptiles remain essential to a more complete understanding of the auditory system of birds and mammals because of their pivotal position in the phylogeny of amniotic vertebrates. The general organization of the central auditory system of reptiles resembles that of birds and, to a lesser extent, that of mammals (Manley 2000). Besides those differences mentioned earlier (such as a lack of direct telencephalic input to the midbrain), multiple forebrain representations and their associated neural circuitry seem to be the rule rather than the exception in birds and mammals (Nieuwenhuys et al. 1998). Similar features have not yet been documented in reptiles with the exception of a second tel-

encephalic auditory area in crocodylians (Pritz and Stritzel 1992). Such anatomical data have prompted the view that the central nervous system of reptiles is dominated by the midbrain (Kappers et al. 1967) but such generalizations for these and other nonmammalian groups must be tempered by the sparse behavioral and physiological evidence available. In keeping with the themes described for fish and amphibians, important questions to be answered in the future include the following. If neural circuitry is similar among lizards, then how does the auditory circuitry of sonic geckos interface with central vocal circuitry? Furthermore, why are so few reptiles, with the exception of geckos, some turtles and crocodylians (Carr 1992), capable of vocalization? For reptiles that do vocalize, what is the role of steroid hormones and how do these relate to behavior? Lastly, what are the auditory specializations for animals that are aquatic and terrestrial as compared with those species that are solely terrestrial?

The nonmammalian auditory systems discussed here and elsewhere (Fay and Popper 1999; Simmons et al. 2003) have just begun to articulate the experiments among fish, amphibians, and reptiles that will reveal neural mechanisms for hearing that are common to all vertebrates. Such model systems will provide clues in infrahuman species about how the human auditory system has evolved to meet environmental challenges common to all vertebrates.

## Abbreviations

AM	amplitude modulated
Cb	cerebellum
CN	cochlear nuclei
D	diencephalon
dF	difference frequency
DLN	dorsolateral medullary nucleus
DO	descending octaval nucleus
DVR	dorsal ventricular ridge
F <sub>0</sub>	fundamental frequency
FM	frequency modulated
IC	inferior colliculus
Mid	midbrain
MO	medulla oblongata
ms	milliseconds
nLL	nucleus of the lateral lemniscus
OB	olfactory bulb
PAG	periaqueductal gray
PL	paralemniscal tegmentum
POA	preoptic area
pps	pulses per second
PRR	pulse repetition rate
Rc	nucleus reuniens pars centralis

Rd	nucleus reuniens pars diffusa
RF	reticular formation
RI	inferior raphe
SAM	sinusoidal amplitude modulation
SC	spinal cord
SO	secondary octaval nucleus of teleost fish <i>or</i> superior olive of anurans and reptiles
Tel	telencephalon
TeO	optic tectum
TS	torus semicircularis
TSnc	central nucleus of the torus semicircularis
TSnl	laminar nucleus of the torus semicircularis
TSpr	principal nucleus of the torus semicircularis
TSvl	ventrolateral nucleus of the torus semicircularis
TV	tectal ventricle
VS	vector strength of synchronization
III	third ventricle

## Acknowledgments

We thank C. McCormick and the editors for many helpful comments on the text, and C. Brown, J. Corbitt, M. Marchaterre, T. Natoli, J. Sisneros, and W. Wilczynski for assistance with the illustrations. This work was supported in part by NIH-NIDCD Grants DC00092 (A.H.B.) and DC003788 (G.J.R.).

## References

- Albert M, Hose B, and Langner G (1989) Modulation transfer functions in the auditory midbrain (MLD) of the guinea fowl (*Numida meleagris*). In: Elsner N and Singer W (eds). *Dynamics and Plasticity in Neuronal Systems*. Thieme Verlag, Stuttgart, pp. 280.
- Alder TB and Rose GJ (1998) Long-term temporal integration in the anuran auditory system. *Nature Neuroscience* **1**:519–522.
- Alder TB and Rose GJ (2000) Integration and recovery processes contribute to the temporal selectivity of neurons in the midbrain of the northern leopard frog, *Rana pipiens*. *Journal of Comparative Physiology A* **186**:923–937.
- Allen DM (1973) Some relationships of vocalization to behavior in the Pacific treefrog, *Hyla regilla*. *Herpetologica* **29**:366–371.
- Bass AH and Baker R (1990) Sexual dimorphisms in the vocal control system of a teleost fish: morphology of physiologically identified neurons. *Journal of Neurobiology* **21**: 1155–1168.
- Bass AH and Baker R (1991) Adaptive modification of homologous vocal control traits in teleost fishes. *Brain, Behavior and Evolution* **38**:240–254.
- Bass AH and McKibben JR (2003) Neural mechanisms and behaviors for acoustic communication in teleost fish. *Progress in Neurobiology* **69**:1–26.



- Bass AH, Marchaterre MA, and Baker R (1994) Vocal-acoustic pathways in a teleost fish. *Journal of Neuroscience* **14**:4025–4039.
- Bass AH, Bodnar DA, and Marchaterre MA (1999) Complementary explanations for existing phenotypes in an acoustic communication system. In: Hauser M and Konishi M (eds) *Neural Mechanisms of Communication*. MIT Press, Cambridge, pp. 493–514.
- Bass AH, Bodnar DA, and Marchaterre MA (2000) Midbrain acoustic circuitry in a vocalizing fish. *Journal of Comparative Neurology* **419**:505–531.
- Bass AH, Bodnar DA, and Marchaterre MA (2001) Acoustic nuclei in the medulla and midbrain of the vocalizing gulf toadfish, *Opsanus beta*. *Brain, Behavior and Evolution* **57**:63–79.
- Belekhova MG, Zharskaja VD, Kachunts AS, Gaidaenko GV, and Tumanova NL (1985) Connections of the mesencephalic, thalamic and telencephalic auditory centers in turtles. Some structural bases for audiosomatic interrelations. *Journal für Hirnforschung* **26**:127–152.
- Belekhova MG, Kenigfest-Rio NB, Vesselkin NP, Rio J-P, Repérant J, and Ward R (2002) Evolutionary significance of different neurochemical organisation of the internal and external regions of auditory centres in the reptilian brain: an immunocytochemical and reduced NADPH-diaphorase histochemical study in turtles. *Brain Research* **925**:100–106.
- Bibikov NG and Gorodetskaya ON (1980) Single unit responses in the auditory center of the frog mesencephalon to amplitude-modulated tones. *Neirofiziologija* **12**:264–271.
- Bodnar DA and Bass AH (1997) Temporal coding of concurrent acoustic signals in auditory midbrain. *Journal of Neuroscience* **17**:7553–7564.
- Bodnar DA and Bass AH (1999) A midbrain combinatorial code for temporal and spectral information in concurrent acoustic signals. *Journal of Neurophysiology* **81**:552–563.
- Bodnar DA and Bass AH (2001a) The coding of concurrent vocal signals by the auditory midbrain: the effects of stimulus level and depth of modulation. *Journal of the Acoustical Society of America* **109**:809–825.
- Bodnar DA and Bass AH (2001b) Coding of concurrent signals by the auditory midbrain: effects of duration. *Journal of Comparative Physiology A* **187**:381–391.
- Bradbury JW and Vehrencamp SL (1998) *Principles of Animal Communication*. Sinauer Associates, Sunderland.
- Brenowitz EA and Rose GJ (1994) Behavioural plasticity mediates aggression in choruses of the Pacific treefrog. *Animal Behavior* **47**:633–641.
- Brosch M and Schreiner CE (2000) Sequence sensitivity of neurons in cat primary auditory cortex. *Cerebral Cortex* **10**:1155–1167.
- Browner RH (1983) The ascending connections of the torus semicircularis central nucleus in *Chrysemys scripta elegans*. *Hearing Research* **12**:139–143.
- Browner RH and Baruch AM (1984) The cytoarchitecture of the torus semicircularis in the golden skink, *Mabuya multifasciata*. *Journal of Morphology* **180**:223–242.
- Browner RH and Rubinson K (1977) The cytoarchitecture of the torus semicircularis in the Tegu lizard, *Tupinambis nigropunctatus*. *Journal of Comparative Neurology* **176**:539–558.
- Browner RH, Kennedy MC, and Facelle T (1981) The cytoarchitecture of the torus semicircularis in the red-eared turtle. *Journal of Morphology* **169**:207–223.
- Bullock TH and Heiligenberg W (eds.) (1986) *Electroreception*. John Wiley & Sons, New York.
- Buonomano DV (2000) Decoding temporal information: a model based on short-term synaptic plasticity. *Journal of Neuroscience* **20**:1129–1141.

- Butler AB and Bruce LL (1981) Nucleus laminaris of the torus semicircularis: projection to spinal cord in reptiles. *Neuroscience Letters* **25**:221–225.
- Capranica RR (1965) The evoked vocal response of the bullfrog: a study of communication by sound. *MIT Research Monographs* **33**, MIT Press, Cambridge.
- Capranica RR (1966) Vocal response of the bullfrog to natural and synthetic mating calls. *Journal of the Acoustical Society of America* **40**:1131–1139.
- Capranica RR (1992) The untuning of the tuning curve: is it time? *Seminars in the Neurosciences* **4**:401–408.
- Carr CE (1992) Evolution of the central auditory system in reptiles and birds. In: Webster DB, Fay RR, and Popper AN (eds). *The Evolutionary Biology of Hearing*. Springer-Verlag, New York, pp. 511–543.
- Condon CJ, Chang SH, and Feng AS (1991) Processing of behaviorally relevant temporal parameters of acoustic stimuli by single neurons in the superior olivary nucleus of the leopard frog. *Journal of Comparative Physiology A* **168**:709–725.
- Condon CJ, White KR, and Feng AS (1996) Neurons with different temporal firing patterns in the inferior colliculus of the little brown bat differentially process sinusoidal amplitude-modulated signals. *Journal of Comparative Physiology A* **178**:147–157.
- Corwin JT (1981) Peripheral auditory physiology in the lemon shark: evidence of parallel otolithic and non-otolithic sound detection. *Journal of Comparative Physiology A* **142**:379–390.
- Crawford JD (1993) Central auditory neurophysiology of a sound producing fish: the mesencephalon of *Pollimyrus isidori* (Mormyridae). *Journal of Comparative Physiology A* **172**:139–152.
- Crawford JD (1997) Feature-detecting auditory neurons in the brain of a sound-producing fish. *Journal of Comparative Physiology A* **180**:439–450.
- Crawford JD, Cook AP, and Heberlain AS (1997) Bioacoustic behavior of African fishes (Mormyridae): potential cues for species and individual recognition in *Pollimyrus*. *Journal of the Acoustical Society of America* **102**:1200–1212.
- Díaz C, Yanes C, Trujillo C-M, and Puelles L (2000) Cytoarchitectonic subdivisions in the subtectal midbrain of the lizard *Gallotia galloti*. *Journal of Neurocytology* **29**:569–593.
- Diekamp B and Gerhardt HC (1995) Selective phonotaxis to advertisement calls in the gray treefrog, *Hyla versicolor*: behavioral experiments and neurophysiological correlates. *Journal of Comparative Physiology A* **177**:173–190.
- Drakontides AB and Browner RB (1986) Ultrastructural features of the ventromedial region of the laminar nucleus in the red-eared turtle *Chrysemys scripta elegans*. *Journal of Comparative Neurology* **248**:555–572.
- Ebbesson SOE (1967) Ascending axon degeneration following hemisection of the spinal cord in the Tegu lizard (*Tupinambis nigropunctatus*). *Brain Research* **5**:178–206.
- Edds-Walton PL, Fay RR, and Highstein SM (1999) Dendritic arbors and central projections of physiologically characterized auditory fibers from the saccule of the toadfish, *Opsanus tau*. *Journal of Comparative Neurology* **411**:212–238.
- Edwards CJ, Alder TB, and Rose GJ (2002) Auditory midbrain neurons that count. *Nature Neuroscience* **5**:934–936.
- Eggermont JJ (2001) Between sound and perception: reviewing the search for a neural code. *Hearing Research* **157**:1–42.
- Fay RR and Edds-Walton PL (1997) Diversity in frequency response properties of saccular afferents of the toadfish, *Opsanus tau*. *Hearing Research* **113**:235–246.
- Fay RR and Popper AN (eds.) (1999) *Springer Handbook of Auditory Research*, Volume 11: *Comparative Hearing: Fish and Amphibians*. Springer-Verlag, New York.

- Feng AS and Lin WY (1991) Differential innervation patterns of three divisions of frog auditory midbrain (*torus semicircularis*). *Journal of Comparative Neurology* **306**:613–630.
- Feng AS and Schellart NAM (1999) Central auditory processing in fish and amphibians. In: Fay RR and Popper AN (eds). *Springer Handbook of Auditory Research*, Volume 11: *Comparative Hearing: Fish and Amphibians*. Springer-Verlag, New York, pp. 218–268.
- Fine ML, Winn HE, and Olla BL (1977) Communication in fishes. In: Sebeok TA (ed). *How Animals Communicate*. Indiana University Press, Bloomington, pp. 472–518.
- Fletcher LB and Crawford JD (2001) Acoustic detection by sound-producing fishes (Mormyridae): the role of gas-filled tympanic bladders. *Journal of Experimental Biology* **204**:175–183.
- Follett KA (1989) A telencephalospinal projection in the Tegu lizard (*Tupinambis tequixin*). *Brain Research* **496**:89–97.
- Fortune ES and Rose GJ (2001) Short-term synaptic plasticity as a temporal filter. *Trends in Neurosciences* **24**:381–385.
- Foster RE and Hall WC (1978) The organization of central auditory pathways in a reptile, *Iguana iguana*. *Journal of Comparative Neurology* **178**:783–832.
- Frisina RD (2001) Subcortical neural coding mechanisms for auditory temporal processing. *Hearing Research* **158**:1–27.
- Fuzessery ZM (1988) Frequency tuning in the anuran central auditory system. In: Fritzsche B, Wilczynski W, Ryan MJ, and Walkowiak W (eds). *The Evolution of the Amphibian Auditory System*. John Wiley & Sons, New York, pp. 253–273.
- Fuzessery ZM and Feng AS (1983) Mating call selectivity in the thalamus and midbrain of the leopard frog (*Rana p. pipiens*): single and multiunit activity. *Journal of Comparative Physiology A* **150**:333–344.
- Gerhardt HC (1982) Sound pattern recognition in some North American treefrogs (Anura: Hylidae): implications for mate choice. *American Zoologist* **22**:581–595.
- Gerhardt HC (1988) Acoustic properties used in call recognition by frogs and toads. In: Fritzsche B, Wilczynski W, Ryan MJ, and Walkowiak W (eds). *The Evolution of the Amphibian Auditory System*. John Wiley & Sons, New York, pp. 275–294.
- Goodson JL and Bass AH (2000) Forebrain peptide modulation of sexually polymorphic vocal motor circuitry. *Nature* **403**:769–772.
- Goodson JL and Bass AH (2002) Vocal-acoustic circuitry and descending vocal motor pathways in teleost fish: convergence with terrestrial vertebrates reveals conserved traits. *Journal of Comparative Neurology* **448**:298–321.
- Gooler DM and Feng AS (1992) Temporal coding in the frog midbrain: the influence of duration and rise-fall time on the processing of complex amplitude-modulated stimuli. *Journal of Neurophysiology* **67**:1–22.
- Hall JC (1994) Central processing of communication sounds in the anuran auditory system. *American Zoologist* **34**:670–684.
- Hall JC (1999) GABAergic inhibition shapes frequency tuning and modifies response properties in the auditory midbrain of the leopard frog. *Journal of Comparative Physiology A* **185**:479–491.
- Hall JC and Feng AS (1987) Evidence for parallel processing in the frog's auditory thalamus. *Journal of Comparative Neurology* **258**:407–419.
- Hall JC and Feng AS (1988) Influence of envelope rise time on neural responses in the auditory system of anurans. *Hearing Research* **36**:261–276.
- Hall JC and Feng AS (1991) Temporal processing in the dorsal medullary nucleus of the northern leopard frog (*Rana pipiens pipiens*). *Journal of Neurophysiology* **66**:955–973.

- Highstein SM, Kitch R, Carey J, and Baker R (1992) Anatomical organization of the brainstem octavolateralis area of the oyster toadfish, *Opsanus tau*. *Journal of Comparative Neurology* **319**:501–518.
- Jürgens U (2002) Neural pathways underlying vocal control. *Neuroscience and Biobehavioral Reviews* **26**:235–258.
- Kachunts AS (1982) Electrophysiologic characteristics of representation of auditory and somatosensory systems in the turtle midbrain. *Neurofiziologiya* **14**:260–269.
- Kappers CUA, Huber GC, and Crosby EC (1967) *The Comparative Anatomy of the Nervous System of Vertebrates, Including Man*. Hafner, New York.
- Kennedy MC (1974) Auditory multiple-unit activity in the midbrain of the Tokay Gecko (*Gekko gekko*, L.). *Brain, Behavior and Evolution* **10**:257–264.
- Kennedy MC (1975) Vocalization elicited in a lizard by electrical stimulation of the midbrain. *Brain Research* **91**:321–325.
- Klug A, Bauer EE, Hanson JT, Hurley L, Meitzen J, and Pollak GD (2002) Response selectivity for species-specific calls in the inferior colliculus of Mexican free-tailed bats is generated by inhibition. *Journal of Neurophysiology* **88**:1941–1954.
- Knudsen EI (1977) Distinct auditory and lateral line nuclei in the midbrain of catfishes. *Journal of Comparative Neurology* **173**:417–432.
- Kozloski J and Crawford JD (1998) Functional neuroanatomy of auditory pathways in the sound-producing fish *Pollimyrus*. *Journal of Comparative Neurology* **401**:227–252.
- Kozloski J and Crawford JD (2000) Transformations of an auditory temporal code in the medulla of a sound-producing fish. *Journal of Neuroscience* **20**:2400–2408.
- Krishna BS and Semple MN (2000) Auditory temporal processing: responses to sinusoidally amplitude-modulated tones in the inferior colliculus. *Journal of Neurophysiology* **84**:255–273.
- Künzle H (1986) Projections from the cochlear nuclear complex to rhombencephalic auditory centers and torus semicircularis in the turtle. *Brain Research* **379**:307–319.
- Künzle H and Woodson W (1982) Mesodiencephalic and other target regions of ascending spinal projections in the turtle, *Pseudemys scripta elegans*. *Journal of Comparative Neurology* **212**:349–364.
- Langner G (1981) Neuronal mechanisms for periodicity analysis in the time domain. In: Klinke R and Hartmann R (eds). *Hearing—Physiological Bases and Psychoacoustics*. Springer-Verlag, Berlin, pp. 334–343.
- Langner G (1992) Periodicity coding in the auditory system. *Hearing Research* **60**:115–142.
- Langner G and Schreiner CE (1988) Periodicity coding in the inferior colliculus of the cat. I. Neuronal mechanisms. *Journal of Neurophysiology* **60**:1799–1822.
- Langner G, Albert M, and Briede T (2002) Temporal and spatial coding of periodicity information in the inferior colliculus of awake chinchilla (*Chinchilla laniger*). *Hearing Research* **168**:110–130.
- Large EW and Crawford JD (2002) Auditory temporal computation: interval selectivity based on post-inhibitory rebound. *Journal of Computational Neuroscience* **13**:135–142.
- Lee J (1994) Amplitude modulation rate discrimination with sinusoidal carriers. *Journal of the Acoustical Society of America* **96**:2140–2147.
- Lewis ER and Lombard RE (1988) The amphibian inner ear. In: Fritzsche B, Wilczynski W, Ryan MJ, and Walkowiak W (eds). *The Evolution of the Amphibian Auditory System*. John Wiley & Sons, New York, pp. 93–123.

- Lewis ER and Narins PM (1999) The acoustic periphery of amphibians: anatomy and physiology. In: Popper AN and Fay RR (eds). *Springer Handbook of Auditory Research*, Volume 11: *Comparative Hearing: Fish and Amphibians*. Springer, New York, pp. 101–154.
- Lewis ER, Baird RA, Leverenz EL, and Koyama H (1982) Inner ear: dye injection reveals peripheral origins of specific sensitivities. *Science* **215**:1641–1643.
- Lu Z and Fay RR (1996) Two-tone interaction in primary afferents and midbrain neurons of the goldfish, *Carassius auratus*. *Auditory Neuroscience* **2**:257–273.
- Manley JA (1971) Single unit studies in the midbrain auditory area of *Caiman*. *Zeitschrift für vergleichende Physiologie* **71**:255–261.
- Manley JA (2000) Cochlear mechanisms from a phylogenetic viewpoint. *Proceedings of the National Academy of Sciences of the United States of America* **97**:11736–11743.
- Margoliash D (1983) Acoustic parameters underlying the responses of song-specific neurons in the white-crowned sparrow. *Journal of Neuroscience* **3**:1039–1057.
- Margoliash D and Fortune ES (1992) Temporal and harmonic combination-sensitive neurons in the zebra finch's HVC. *Journal of Neuroscience* **12**:4309–4326.
- Martin WF (1971) Mechanics of sound production in toads of the genus *Bufo*: passive elements. *Journal of Experimental Biology* **176**:273–294.
- Marvit P and Crawford JD (2000) Auditory discrimination in a sound-producing electric fish (*Pollimyrus*): tone frequency and click-rate difference detection. *Journal of the Acoustical Society of America* **108**:1819–1825.
- McCormick CA (1997) Organization and connections of octaval and lateral line centers in the medulla of a clupeid, *Dorosoma cepedianum*. *Hearing Research* **110**:39–60.
- McCormick CA (1999) Anatomy of the central auditory pathways of fish and amphibians. In: Popper AN and Fay RR (eds.) *Springer Handbook of Auditory Research*, Volume 11: *Comparative Hearing: Fish and Amphibians*. Springer-Verlag, New York, pp. 155–217.
- McCue MP and Guinan JJ Jr (1994) Acoustically responsive fibers in the vestibular nerve of the cat. *Journal of Neuroscience* **14**:6058–6070.
- McKibben JR and Bass AH (1998) Behavioral assessment of acoustic parameters relevant to signal recognition and preference in a vocal fish. *Journal of the Acoustical Society of America* **104**:3520–3533.
- McKibben JR and Bass AH (1999) Peripheral encoding of behaviorally relevant acoustic signals in a vocal fish: single tones. *Journal of Comparative Physiology A* **184**:563–576.
- McKibben JR and Bass AH (2001a) Effects of temporal envelope modulation on acoustic signal recognition in a vocal fish. *Journal of the Acoustical Society of America* **109**:2934–2943.
- McKibben JR and Bass AH (2001b) Peripheral encoding of behaviorally relevant acoustic signals in a vocal fish: harmonic and beat stimuli. *Journal of Comparative Physiology A* **187**:271–285.
- Mensing AF, Carey JP, Boyle R, and Highstein SM (1997) Differential central projections of physiologically characterized horizontal semicircular canal vestibular nerve afferents in the toadfish, *Opsanus tau*. *Journal of Comparative Neurology* **384**:71–85.
- Metzner W (1996) Anatomical basis for audio-vocal integration in echolocating horseshoe bats. *Journal of Comparative Neurology* **368**:252–269.
- Nagarajan SS, Cheung SW, Bedenbaugh P, Beitel RE, Schreiner CE, and Merzenich MM (2002) Representation of spectral and temporal envelope of twitter vocalizations in

- common marmoset primary auditory cortex. *Journal of Neurophysiology* **87**:1723–1737.
- Nelson JS (1984) *Fishes of the World*. John Wiley & Sons, New York.
- Nieuwenhuys R, ten Donkelaar HJ, and Nicholson C (eds.) (1998) *The Central Nervous System of Vertebrates*. Springer-Verlag, New York.
- Northcutt RG (1974) Some histochemical observations on the telencephalon of the bullfrog, *Rana catesbeiana* Shaw. *Journal of Comparative Neurology* **157**:379–390.
- O'Neill WE and Suga N (1979) Target range-sensitive neurons in the auditory cortex of the mustache bat. *Science* **203**:69–72.
- Pinheiro AD, Wu M, and Jen PH-S (1991) Encoding repetition rate and duration in the inferior colliculus of the big brown bat *Eptesicus fuscus*. *Journal of Comparative Physiology A* **169**:69–85.
- Pollak GD, Burger RM, and Klug A (2003) Dissecting the circuitry of the auditory system. *Trends in Neurosciences* **26**:33–39.
- Popper AN and Fay RR (1999) The auditory periphery in fishes. In: Fay RR and Popper AN (eds). *Springer Handbook of Auditory Research*, Volume 11: *Comparative Hearing: Fish and Amphibian*. Springer-Verlag, New York, pp. 43–100.
- Pough FH, Janis CM, and Heiser JB (2002) *Vertebrate Life*. Prentice-Hall, Upper Saddle River.
- Prechtl JC, von der Emde G, Wolfart J, Karamürsel S, Akoev GN, Andrianov YN, and Bullock TH (1998) Sensory processing in the pallium of a mormyrid fish. *Journal of Neuroscience* **18**:7381–7393.
- Pritz MB (1974) Ascending connections of a midbrain auditory area in a crocodile, *Caiman crocodilus*. *Journal of Comparative Neurology* **153**:179–198.
- Pritz MB and Stritzel ME (1989) Reptilian somatosensory midbrain: identification based on input from the spinal cord and dorsal column nucleus. *Brain, Behavior and Evolution* **33**:1–14.
- Pritz MB and Stritzel ME (1992) A second auditory area in the non-cortical telencephalon of a reptile. *Brain Research* **569**:146–151.
- Rose GJ (1995) Representation of temporal patterns of amplitude modulation in the anuran auditory system and electrosensory system. In: Covey E, Hawkins HL, and Port RF (eds). *Representation of Temporal Patterns*. Plenum Press, New York, pp. 1–23.
- Rose GJ and Brenowitz EA (1997) Plasticity of aggressive thresholds in *Hyla regilla*: discrete accommodation to encounter calls. *Animal Behavior* **53**:353–361.
- Rose GJ and Brenowitz EA (2002) Pacific treefrogs use temporal integration to differentiate advertisement from encounter calls. *Animal Behavior* **63**:1183–1190.
- Rose G and Capranica RR (1983) Temporal selectivity in the central auditory system of the leopard frog *Rana pipiens*. *Science* **219**:1087–1089.
- Rose GJ and Capranica RR (1984) Processing amplitude-modulated sounds by the auditory midbrain of two species of toads: matched temporal filters. *Journal of Comparative Physiology A* **154**:211–219.
- Rose G and Capranica RR (1985) Sensitivity to amplitude modulated sounds in the anuran auditory nervous system. *Journal of Neurophysiology* **53**:446–465.
- Rose GJ, Brenowitz EA, and Capranica RR (1985) Species specificity and temperature dependency of temporal processing by the auditory midbrain of two species of treefrogs. *Journal of Comparative Physiology A* **157**:763–769.
- Rose GJ, Zellick R, and Rand SA (1988) Auditory processing of temporal information in a neotropical frog is independent of signal intensity. *Ethology* **77**:330–336.

- Ryan MJ (1983) Frequency modulated calls and species recognition in a neotropical frog. *Journal of Comparative Physiology A* **150**:217–221.
- Schmidt RS (1976) Neural correlates of frog calling: isolated brainstem. *Journal of Comparative Physiology A* **108**:99–113.
- Schneider H (1988) Peripheral and central mechanisms of vocalization. In: Fritzsche B, Wilczynski W, Ryan MJ, and Walkowiak W (eds). *The Evolution of the Amphibian Auditory System*. John Wiley & Sons, New York, pp. 275–294.
- Shannon RV, Zeng F-G, Kamath V, Wygonski J, and Ekelid M (1995) Speech recognition with primarily temporal cues. *Science* **270**:303–304.
- Simmons AM, Popper AN, and Fay RR (eds.) (2003) *Springer Handbook of Auditory Research, Volume 16: Acoustic Communication*. Springer-Verlag, New York.
- Sisneros J and Bass AH (2003) Seasonal plasticity of peripheral auditory frequency sensitivity. *Journal of Neuroscience* **23**:1049–1058.
- Smotherman M, Zhang S, and Metzner W (2003) A neural basis for auditory feedback control of vocal pitch. *Journal of Neuroscience* **23**:1464–1477.
- Straughan IR and Heyer WR (1976) A functional analysis of the mating calls of the neotropical frog genera of the *Leptodactylus* complex (Amphibia, Leptodactylidae). *Papeis Avulsos Zoologica Sao Paulo* **29**:221–245.
- Suga N, Schlegel P, Shimozawa T, and Simmons J (1973) Orientation sounds evoked from echolocating bats by electrical stimulation of the brain. *Journal of the Acoustical Society of America* **54**:793–797.
- Suzuki A, Kozloski J, and Crawford JD (2002) Temporal encoding for auditory computation: physiology of primary afferent neurons in sound-producing fish. *Journal of Neuroscience* **22**:6290–6301.
- Tang Y-Z, Piao Y-S, Zhuang L-Z, and Wang Z-W (2001) Expression of androgen receptor mRNA in the brain of *Gekko gekko*: implications for understanding the role of androgens in controlling auditory and vocal processes. *Journal of Comparative Neurology* **438**:136–147.
- ten Donkelaar HJ (1998a) Anurans. In: Nieuwenhuys R, ten Donkelaar HJ, and Nicholson C (eds). *The Central Nervous System of Vertebrates*. Springer-Verlag, New York, pp. 1151–1314.
- ten Donkelaar HJ (1998b) Reptiles. In: Nieuwenhuys R, ten Donkelaar HJ, and Nicholson C (eds). *The Central Nervous System of Vertebrates*. Springer-Verlag, New York, pp. 1315–1524.
- ten Donkelaar HJ and de Boer van-Huizen R (1978) Cells of origin of pathways descending to the spinal cord in a lizard (*Lacerta galloti*). *Neuroscience Letters* **9**:123–128.
- ten Donkelaar HJ and de Boer-van Huizen R (1987) A possible pain control system in a non-mammalian vertebrate (a lizard, *Gekko gekko*). *Neuroscience Letters* **83**:65–70.
- ten Donkelaar HJ and de Boer-van Huizen R (1988) Brain stem afferents to the anterior dorsal ventricular ridge in a lizard (*Varanus exanthematicus*). *Anatomy and Embryology* **177**:465–475.
- Ulinksí PS, Dacey DM, and Sereno MI (1992) Optic tectum. In: Gans C and Ulinksí PS (eds). *Biology of the Reptilia, Volume 17: Neurology C, Sensorimotor Integration*. University of Chicago Press, Chicago, pp. 241–366.
- Van Tasell DJ, Soli SD, Kirby VM, and Widin GP (1987) Speech waveform envelope cues for consonant recognition. *Journal of the Acoustical Society of America* **82**:1152–1161.

- Von der Emde G and Prechtl JC (1999) Anatomical connections of auditory and lateral line areas of the dorsal telencephalon (Dm) in the osteoglossomorph teleost, *Gnathopeternus petersii*. *Brain Research* **818**:355–367.
- Walkowiak W (1980) The coding of auditory signals in the torus semicircularis of the fire-bellied toad and grass frog: responses to simple stimuli and to conspecific calls. *Journal of Comparative Physiology A* **138**:131–148.
- Walkowiak W (1988) Central temporal encoding. In: Fritzsche B, Wilczynski W, Ryan MJ, and Walkowiak W (eds). *The Evolution of the Amphibian Auditory System*. John Wiley & Sons, New York, pp. 275–294.
- Wang X (2000) On cortical coding of vocal communication sounds in primates. *Proceedings of the National Academy of Sciences of the United States of America* **97**: 11843–11849.
- Weeg, MS and Bass AH (2000) Midbrain lateral line circuitry in a vocalizing fish. *Journal of Comparative Neurology* **418**:841–864.
- Weeg MS and Bass AH (2003) Modulation of auditory and lateral line efferents during fictive vocalization. *Society for Neuroscience Abstracts* **28**:93.10.
- Weeg MS, Fay RR, and Bass AH (2002) Directionality and frequency tuning of primary saccular afferents of a vocal fish, the plainfin midshipman (*Porichthys notatus*). *Journal of Comparative Physiology A* **188**:631–641.
- Wilczynski W (1988) Brainstem auditory pathways in anurans. In: Fritzsche B, Wilczynski W, Ryan MJ, and Walkowiak W (eds). *The Evolution of the Amphibian Auditory System*. John Wiley & Sons, New York, pp. 209–231.
- Wilczynski W and Northcutt RG (1983) Connections of the bullfrog striatum: efferent projections. *Journal of Comparative Neurology* **214**:333–343.



# Chapter 17

## The Auditory Midbrain in Bats and Birds

ELLEN COVEY AND CATHERINE E. CARR

### 1. INTRODUCTION

#### *1.1. EVOLUTION AND PHYLOGENY*

Both bats and birds are flying vertebrates that diverged early in evolution. Their flight mechanisms are different, and so are their auditory systems. They share a common ancestor in the Carboniferous era, but their auditory systems evolved in parallel and are not homologous (Clack 1997). However, other ancestral mammalian traits, such as moveable ears, multiple ossicles, and echolocation have modified the ear and the auditory system of bats. We discuss both species because they reveal neuroethological principles that relate neural activity to behavior.

#### *1.2. BEHAVIORAL CONTEXT*

Bats and birds use hearing to localize and analyze sound sources passively, and to communicate with conspecifics (Kanwal et al. 1993; Konishi 2000). Some birds learn song and have forebrain areas to control vocal pathways. Others, like the barn owl, are nocturnal predators with superb sound localization skills. Bats use their extraordinary hearing in concert with stereotyped vocalizations and listen to the echoes from objects. By analyzing how echoes have been modified from the original sound by reflective objects, bats perceive obstacles and capture flying insects in darkness.

##### 1.2.1. Bat Hearing and Echolocation

Bats are Chiroptera, an order closely related to insectivores. Because bats use their sense of hearing to perform tasks that in other mammals are guided by vision, their auditory system has undergone a corresponding species-specific hypertrophy and specialization (Neuweiler 1990). The inferior colliculus (IC) is a structure critical for hearing that has evolved to fulfill unique functional needs for different species of bats.

There are approximately 1000 bat species and each uses a different pattern of vocalization for echolocation (Neuweiler 2000). Bats that hunt for insects in open spaces (e.g., the Mexican free-tailed bat, *Tadarida brasiliensis*) use short-duration frequency-modulated (FM) calls. Bats that hunt in foliage use calls with a long constant-frequency (CF) component and a short FM component (CF-FM calls). The best studied CF-FM species are the mustached bat, *Pteronotus parnellii*, and the horseshoe bats, *Rhinolophus rouxi* and *R. ferrumequinum*. Other species, like the big brown bat, *Eptesicus fuscus*, use either CF- or FM-type calls (Fig. 17.1) depending on circumstances (Simmons 1989; Grinnell 1995).

Bat echolocation calls are modified in predictable ways by objects. The interval between call and echo is proportional to the distance of the reflective object. The attenuation of the echo depends on the object's size and distance. Three-dimensional structures create characteristic interference patterns in the echoes (Simmons 1989). The IC contains neurons sensitive to all these features of echolocation signals.

All bat species listen to sounds passively and most echolocate. Species that glean from surfaces such as the pallid bat, *Antrozous pallidus*, listen to sounds made by prey (Fuzessery et al. 1993). Many bat species have a rich repertoire of communication sounds that differ from echolocation calls in temporal structure (Kanwal et al. 1993).

Because there is so much information on the patterns of sound that are behaviorally relevant, bats have become an important model system in which to study temporal aspects of sound processing. Although bats are specialized mammals, much of what has been learned about their auditory systems can be generalized to other mammals (Covey 2003).

### 1.2.2. Avian Hearing

Psychophysical tests have measured auditory sensitivity, loudness, and temporal resolving power in several species of birds. These data resemble those for other vertebrates, including humans (Dooling et al. 2000). Birdsong has become a model for complex sound processing, especially in the auditory midbrain (Theunissen et al. 2000). Studies of the owl, a sound-localization specialist, have implications for spatial hearing, while the development of the map of auditory space has become a paradigm for studies of experience-dependent plasticity (Knudsen 2002).

### 1.2.3. The Auditory Midbrain: A Hub of the Central Auditory System

In birds and mammals, the midbrain occupies a central position in the auditory system, receiving input from many ascending and descending sensory pathways as well as motor and modulatory projections. The IC projects to the thalamus and has abundant motor and premotor targets (Covey and Casseday 1995; Carr and Code 2000). Besides developing selectivity for biologically relevant sounds, the IC likely transforms the high rate of auditory input into a slower rate of




Figure 17.1. Echolocation calls of *Eptesicus* during foraging, pursuit, and insect capture. Calls are 100 dB SPL or more. While hunting the bat emits narrowband quasi-constant frequency calls (#1, #2) between 23 and 29 kHz and up to 20 ms or longer. After finding an insect, it emits broadband FM calls (#4 to #25) that decrease in duration and increase in rate until the terminal buzz (#25 to end) when durations are <1 ms and the repetition rate is up to 150/s. (From Simmons 1989; redrawn from Casseday and Covey 1995.)

output that matches the speed of motor performance (Casseday and Covey 1996).

## 2. INFERIOR COLLICULUS STRUCTURE

### 2.1. STRUCTURE AND CHEMOARCHITECTURE OF THE BAT INFERIOR COLLICULUS

The bat IC is enormous and protrudes between the cerebellum and the cortex. In Nissl-stained material it does not have obvious subdivisions (Fig. 17.2), although most species have a “pericentral” area that contains large neurons and




Figure 17.2. Nissl preparations showing the relative size and location of the IC in an echolocating bat, *Eptesicus fuscus*. (A) Parasagittal section. (B) Frontal section through the caudal one-third of the IC; the caudorostral level is indicated by the arrow (A).

probably corresponds to the external nucleus in other mammals (see Chapter 22).

### 2.1.1. Organization of the Bat Inferior Colliculus

Like that of other mammals, the bat IC contains various cell types, including small disc-shaped cells and large multipolar neurons (Zook et al. 1985; see Chapter 22). *Eptesicus* and other bats with a flexible repertoire of echolocation calls and behavior have a homogeneous IC. *Pteronotus*, a CF-FM bat, has an expanded region of the basilar membrane, the “auditory fovea” (Pollak et al. 1972; Bruns and Schmieszek 1980). There is a corresponding IC “foveal” frequency presentation that is distinguished cytoarchitecturally, the dorsomedial (or dorsoposterior) division (Zook and Casseday 1982; Zook et al. 1985).

### 2.1.2. Biochemical Gradients in the Bat Midbrain

Chemoarchitectural markers reveal regional differences or gradients within the IC. Immunoreactive synaptic endings (puncta) stained for the inhibitory neurotransmitter  $\gamma$ -aminobutyric acid (GABA) are virtually absent in a crescent-shaped area in the ventrolateral IC, but this same area is immunostained for glycine (Vater et al. 1992b; Winer et al. 1995). Local gradients in GABA and glycine immunoreactivity are organized so that each area is devoted to a single frequency range. This specific isofrequency lamina contains subregions in which neuroactive compounds are abundant or sparse. The distribution of GABA<sub>A</sub> and glycine receptors follows the distribution of GABA and glycine puncta; however, GABA<sub>B</sub> receptors are concentrated in the dorsomedial IC (Fubara et al. 1996). Because GABA<sub>A</sub> receptors provide rapid inhibition and GABA<sub>B</sub> receptors subserve longer inhibition, the same neurotransmitter could have different effects in different IC subregions.

The IC receives glutamatergic excitatory input from many sources in the cochlear nucleus (CN) and superior olivary complex (SOC) and modulatory input from neurons containing acetylcholine (Habbicht and Vater 1996) or serotonin (Kaiser and Covey 1997; Hurley and Thompson 2001). Serotonin concentrates in the dorsomedial half of the IC and at the lateral edge.

Other biochemical markers include the Ca<sup>2+</sup> binding proteins, parvalbumin, calbindin-D28 and calretinin (Zettel et al. 1991; Vater and Braun 1994), and the voltage-gated potassium channel subunit, *Kv1.1* (Rosenberger et al. 2003). Parvalbumin is present in many neurons of all types throughout the IC. Calbindin- and calretinin-positive neurons are concentrated in the dorsomedial IC and external nucleus. Calretinin-positive fibers are densest in an arc above the glycine-rich area, probably from heavy input from calretinin-positive cells in the ventral nucleus of the lateral lemniscus (VLL), which are also glycinergic (Vater et al. 1997).

The *Kv1.1* potassium channel subunit is related to ion channel kinetics implicated in precise temporal processing (Brew and Forsythe 1995). Neurons immunoreactive for *Kv1.1* in *Eptesicus* are mostly in the ventral two thirds of the

IC, decreasing from ventrolateral to dorsomedial (Rosenberger et al. 2003). Many immunolabeled cells are multipolar, indicating that disc-shaped cells have few or no *Kv1.1* ion channels. This regional concentration suggests that multipolar neurons in the ventrolateral IC are distinct and concerned with preserving precise timing.

## 2.2. STRUCTURE OF THE AUDITORY MIDBRAIN IN BIRDS

The avian auditory midbrain forms when a caudodorsal portion of the mesencephalic vesicle buckles and thickens to encroach upon the tectal ventricle (Puelles et al. 1994). This massive invagination protrudes beneath the tectal ventricle as the torus semicircularis in nonmammalian species. The torus contains three main parts: the intercollicular area, the toral nucleus, and the preisthmic superficial area (Puelles et al. 1994; Fig. 17.3). The toral nucleus includes the auditory region. It is surrounded rostrally and laterally by the intercollicular area (Wild et al. 1993) and caudally by the caudal preisthmic superficial area (Puelles et al. 1994).

Auditory midbrain nomenclature varies by author and custom. The main recipient of ascending auditory information, Puelles' toral nucleus, has been referred to as the mesencephalic lateral nucleus, pars dorsalis (MLd) (Karten 1967) and as the inferior colliculus (Knudsen 1983). We use the term inferior colliculus, as a major premise of this account is that the functional organization of the avian auditory midbrain parallels that of the mammalian IC.

IC borders and connections have been reevaluated and an intercollicular area between the IC and the tectum identified (Puelles et al. 1994; Fig. 17.3). The intercollicular core, also called DM in songbirds, mediates avian call production and has robust cholinergic immunoreactivity and steroid binding sites (Balthazart 1992; Gahr et al. 1993). Consistent with its function in call production, the intercollicular core receives descending input from the lateral hypothalamus and forebrain archistriatum, specifically the robust archistriatal nucleus, which provides descending motor control of the songbird vocal organ (Wild et al. 1993).

### 2.2.1. Organization of the Avian Inferior Colliculus

The avian IC contains a large tonotopically organized central nucleus (ICC) and two surrounding nuclei. The central nucleus contains a central core (ICCc) surrounded by lateral (ICCl) and medial (ICCM) shells. Lateral to the central nucleus is the external nucleus (ICX). Surrounding ICC and ICX dorsally is a periventricular lamina, the superficial nucleus (ICS; Wagner et al. 2003). Medial to the ICC is the caudomedial shell and paracentral nucleus, both intercollicular structures (Fig. 17.3B; Puelles et al. 1994).

### 2.2.2. Neurons and Internal Organization

The bird IC, like that of the bat, contains many large, loosely packed cells with fusiform, stellate, or round somata (Knudsen 1983). In Golgi preparations many




Figure 17.3. Avian auditory midbrain organization. The *inset* (upper left) partially removes the mesencephalic roof. (A) Schema of the right torus protruding into the optic lobe ventricle, showing its topographic relationships with tectum (*T*), tectal gray (*GT*), and isthmic structures: isthmo-optic nucleus (*IO*) and trochlear nerve (*iv*). The three toral subdivisions (*IC*, *To*, *S*) are shaded dotted contours; black arrow denotes subdivisions (cf. C). (B) Subdivisions of the right torus from a rostral perspective as projected in a transverse plane (directional arrows: dorsal and medial) and related to the paracentral toral nucleus (*PC*), external toral nucleus (*E*), toral periventricular lamina (*PV*), intercollicular area core (*Ic*) and shell (*Is*), periaqueductal gray (*PAG*), preisthmic superficial area shell and core (*Ss*, *Sc*), and preisthmic dorsal brain surface (right). (C) Toral nucleus subdivisions (as in A) showing the posterior location of the hilar nucleus (*H*) and the orientation of the central core (*Cc*). (Modified from Puelles et al. 1994.)

cells are multipolar with stellate or elongate somata (Fig. 17.4). Bipolar and unipolar cells are rare and lie near the margins of the nucleus. Unlike most mammals (see Chapter 2), there is neither fibrodendritic alignment nor preferred orientation. Dendritic fields may be polarized in any dimension and the lemniscal input does not enter or ramify in a laminar fashion (Knudsen 1983). External nucleus cells are smaller than those in the central nucleus, but the same cell types occur.




Figure 17.4. IC cell types and subdivisions in Golgi preparations. *Fine lines*, fibers; *ICC*, IC central core, dorsal and ventral; *ICS*, IC superficial nucleus; *ICX*, IC external nucleus; *OT*, optic tectum. (A) ICX stellate cell. (B) ICX elongate cell. (C) ICS bipolar cell. (Modified from Knudsen 1983.)



### 2.2.3. Biochemical Gradients in the Bird Inferior Colliculus

Biochemical features delineate and support the scheme of architectonic divisions (Fig. 17.5). The differential distributions of these markers identify nuclear subdivisions, which are more apparent than in bats.

In the barn owl and chicken, calretinin immunoreactive fibers from the nucleus laminaris demarcate the laminaris terminal field in the ICCc and thus the pathway used for interaural time difference (ITD) computations (Puelles et al. 1994; Takahashi et al. 1987; Kubke et al. 1999). Acetylcholinesterase defines the medial and lateral shell, and the interaural level difference (ILD) pathway (Adolphs 1993; Puelles et al. 1994). Other markers label other IC subdivisions, including the ICX and ICS (Puelles et al. 1994; Wagner et al. 2003).

GABAergic inhibition is a hallmark of the IC in all vertebrates, including birds (Carr et al. 1989; Fujita and Konishi 1991; Müller 1988; Zheng and Knudsen 1999). ICC contains both the largest number and the largest GABAergic neurons. Cytochrome oxidase distinguishes chicken ICCc from a moderately active ICX (Dezso et al. 1993) and the lateral shell had stronger neuropil reactivity than the central core.

## 3. CONNECTIONS OF THE INFERIOR COLLICULUS

### 3.1. CONNECTIONS OF THE INFERIOR COLLICULUS IN BATS

#### 3.1.1. Afferent Inputs

An enormous variety of projections converge in the IC (Fig. 17.6) (Covey and Casseday 1995; Casseday et al. 2002) and brain stem input extends nearly to the IC surface laterally and dorsally. This section focuses on characteristic features in the context of the physiology of the inputs (see Chapters 2 to 5).

Monaural projections (Fig. 17.6A) arise in the contralateral cochlear nucleus (CN), and indirectly from the ventral (VLL) and intermediate (ILL) nuclei of the lateral lemniscus. Binaural projections come from the medial superior olive (MSO), lateral superior olive (LSO), and dorsal nucleus of the lateral lemniscus (DLL). These projections converge throughout the IC with varying degrees of overlap. Dorsal cochlear nucleus (DCN) input extends through almost all of the IC, and is densest in the dorsomedial one third. Posteroventral cochlear nucleus (PVCN) projections are more prominent than in other mammals, and are uniformly distributed. Input from the anteroventral cochlear nucleus (AVCN) is dense ventrolaterally, declining dorsomedially (Zook and Casseday 1985, 1987).

ILL and VLL are hypertrophied and highly specialized in all species of bats (Covey 1993). The VLL of echolocating bats contains a segregated, functionally specialized population of small neurons, the columnar nucleus (VLLc) (Covey and Casseday 1986, 1991), whose neurons are segregated, permitting their inputs to be readily examined. Each sheet receives convergent input from many CN




Figure 17.5. Chick IC chemoarchitecture in sagittal sections through the central nucleus core (*Cc*). Dorsal is up, rostral to the right. (A) Acetylcholinesterase (*AchE*). (B) Nissl stained section. (C) Choline acetyltransferase (*ChAT*). (D) L-Enkephalin (*L-Enk*). (E) Calcium binding protein and/or calretinin. *Cc*, IC central core; *Cs*, IC shell; *E*, external nucleus; *H*, hilar toral nucleus; *Ic*, intercollicular area core; *Im*, nucleus isthmi, pars magnocellularis; *lp*, nucleus isthmi, pars parvicellularis; *Is*, intercollicular area shell; *PV*, toral periventricular lamina; *Sc*, preisthmic superficial area core; *T*, optic tectum; *ttb*, tectobulbar tract. (Modified from Puelles et al. 1994.)

cells, which represent a broad range of frequencies and project to an equally broad IC territory (Covey and Casseday 1986). The VLLc projection is the most dense and widespread of IC inputs. VLLc neurons are glycinergic, so presumably provide widespread inhibition (Vater et al. 1997). The projections of ILL and the remainder of VLL extend throughout the IC and overlap those of VLLc (Covey and Casseday 1986).

Binaural projections originate from LSO and DLL bilaterally, and MSO ipsilaterally (Fig. 17.6B) (Casseday et al. 1988; Vater et al. 1995) except in *Tadarida*, whose MSO input is bilateral (Grothe et al. 1994). The MSO in most bats probably serves a different function than MSO in species with prominent low frequency hearing (Casseday et al. 1988; Covey et al. 1991; Grothe et al. 1992). Instead of processing interaural time differences, which in bats are probably too small to be a useful cue for sound location, the MSO may represent the amplitude envelope of a high frequency carrier (Grothe et al. 1992). The MSO, LSO, and DLL projections extend throughout IC and overlap with each other and with monaural input (Zook and Casseday 1987; Covey et al. 1991; Vater et al. 1995). Although inputs converge, those from each source form gradients: DCN predominates in the dorsomedial part of each isofrequency lamina, while AVCN, LSO, and NLL predominate ventrolaterally.

### 3.1.2. Other Afferents

Mammalian IC receives projections arising from auditory cortex (Herbert et al. 1991; Winer et al. 1998), parts of the medial geniculate body (MGB) and posterior thalamus (Senatorov and Hu 2002), and the substantia nigra pars reticulata, globus pallidus, and central gray (Winer et al. 2002). Most of these pathways have not been studied in bats. In *Pteronotus*, layer V cells in auditory cortex project to the dorsomedial IC but not to the ventrolateral part. Input from the basal amygdaloid nucleus in *Pteronotus* and *Antrozous* (Marsh et al. 2002) is distributed throughout most of the IC.

### 3.1.3. Projections of the Inferior Colliculus

The IC projects to the MGB (Wenstrup et al. 1994; see Chapter 7) and to the superior colliculus (Covey et al. 1987) and pontine gray (Frisina et al. 1989; Schuller et al. 1991; Wenstrup et al. 1994). Input to the pontine gray is comparable in size to the tectothalamic path, suggesting that auditory input to the cerebellum is crucial for aerial species that echolocate.

## 3.2. CONNECTIONS OF THE AVIAN INFERIOR COLLICULUS

Avian IC projections follow the same pattern as other vertebrates, with brain stem input terminating topographically in the contralateral IC (Fig. 17.7; Carr and Code 2000). The surrounding intercollicular area receives descending input from the forebrain archistriatum (Puelles et al. 1994; Wild 1995).




Figure 17.6. Ascending projections to the bat IC. **(A)** Monaural direct and indirect CN pathways to contralateral IC. **(B)** Binaural input from both CN via the superior olivary complex and DLL. *AVCN*, Anteroventral cochlear nucleus; *DCN*, dorsal cochlear nucleus; *DLL*, dorsal nucleus of the lateral lemniscus; *IC*, inferior colliculus; *ILL*, intermediate nucleus of the lateral lemniscus; *LSO*, lateral superior olive; *MSO*, medial superior olive; *MTB*, medial nucleus of the trapezoid body; *PVCN*, posteroventral cochlear nucleus; *VLLc*, ventral nucleus of the lateral lemniscus, columnar division; *VLLm*, ventral nucleus of the lateral lemniscus, multipolar cell division (modified from Woolley and Casseday 2004; see also Karten 1968; Schwartz et al. 1995).

### 3.2.1. Afferent Inputs

There are two lemniscal projections to the ICC, one from the cochlear nucleus angularis (NA) and another from the cochlear nucleus magnocellularis (NM). The NA projection is direct. The NM projects solely and bilaterally to the brain stem nucleus laminaris (NL) where ITD sensitivity first arises, and the NL projects to the IC. Thus, the NL input is analogous to the mammalian MSO projection. Selective, reversible ablation of each cochlear nucleus shows that the two ascending streams form separate neural channels to encode time and intensity cues for sound localization. Injection of local anesthetic in NM alters the space-specific cells' selectivity for ITD, leaving their ILD selectivity intact. Anesthetizing NA has the opposite effect (Takahashi et al. 1984).

Indirect pathways from the avian SO and lateral lemniscal (LL) nuclei also project to the ICC (Fig. 17.7; Leibler 1975; Conlee and Parks 1986; Takahashi and Konishi 1988a,b; Takahashi et al. 1989; Adolphs 1993). The NL and NA projections are segregated within the ICC, perhaps more so in the barn owl than in the chicken or pigeon (Conlee and Parks 1986; Wild 1995). In the barn owl, the ICC core receives ITD input from the contralateral NL and anterior DLL (Takahashi et al. 1987; Takahashi and Konishi 1988a; Adolphs 1993; Carr and Code 2000). Contralateral NA projects to the lateral and medial shell, conveying both ILD and other information. The lateral shell also receives ITD input from the contralateral core (Takahashi et al. 1989). These ITD and ILD signals converge in the lateral shell, which in turn projects to the ICX (see Section 5). Most of the avian auditory system to the level of the midbrain is binaural and without the specialized monaural pathways found in mammals.

### 3.2.2. Projections of the Avian Inferior Colliculus

The ICC projects to both the ICX and the thalamic nucleus ovoidalis (Fig. 17.7; Proctor and Konishi 1997; Carr and Code 2000). The ICX projects to the optic tectum where the auditory and visual maps are in register (Knudsen 1983). In the barn owl, the ICX also receives a topographic, visually based instructive signal from the tectum which calibrates the auditory space map (Luksch et al. 2000; Hyde and Knudsen 2002; see Section 6).

Auditory–motor connections are essential for bird vocalizations. The intercollicular area receives auditory input from the IC in the dove (Akesson et al. 1987) and is part of the descending telencephalic archistriatal pathway for vocal control (Wild 1993; see also Jarvis et al. 2000). The ICX projects to the deep layers of the tectum, which projects to the pontine tegmentum, enabling orientation to sound in space (Knudsen and Knudsen 1983; Knudsen et al. 1993). In birds, the connections of the IC to the tectum are probably concerned with sound localization, and in generating the nontotopic space map in the ICX. The tonotopically organized representation that is transferred to the thalamic pathway in both birds and bats may be concerned with other aspects of sound processing (Bonke et al. 1979; Durand et al. 1992).




Figure 17.7. Connections of the avian IC. Typical avian auditory system and the passerine song control nuclei. Ascending auditory circuits, *black*; *dashed gray lines*, main descending song control circuit; *solid gray lines*, descending input from the song nuclei (HVC shelf and RA cup) to MLd and Ov. *cHV*, Caudal hyperstriatum ventrale; *DM*, dorsal mesencephalic nucleus and part of ICo; *HVC*, hyperstriatum ventrocaudal (used as a proper name); *IC*, inferior colliculus; *ICo*, nucleus intercollicularis; field L, forebrain auditory area; *LL nuclei*, lateral lemniscal nuclei; *NA*, nucleus angularis; *NCM*, caudo-medial neostriatum; *Nif*, nucleus interfacialis; *NL*, nucleus laminaris; *NM*, nucleus magnocellularis; *Ov*, nucleus ovoidalis; *RA*, robust nucleus of the archistriatum; *SON*, superior olivary nucleus; *Uva*, nucleus uvaeformis of the thalamus. IC receives parallel inputs from lower brain stem and projects to the auditory thalamus (*Ov*), as does the ventral nucleus of the lateral lemniscus. (Modified from Woolley and Casseday 2004; see also Karten 1968; Wild 1987; Schwarz et al. 1995; Wild et al. 1997, 2001.)

## 4. TONOTOPIC ORGANIZATION OF THE INFERIOR COLLICULUS

### 4.1. TONOTOPIC ORGANIZATION IN BATS

The bat IC, like that of other mammals, is tonotopically organized with low-frequency afferents terminating dorsolaterally and high-frequency input ventromedially. Many bat species have expanded representations of specific frequency ranges that are critical in echolocation (Fig. 17.8).

#### 4.1.1. Tonotopy in Frequency Modulating Bats

*Eptesicus* is a species without cochlear specializations. The largest area in the ICC is the 20- to 40-kHz representation, with a continuous low-to-high sequence from dorsolateral to ventromedial (Casseday and Covey 1992). *Antrozous* uses echolocation and passive listening to detect and locate prey. IC tonotopy resembles that in *Eptesicus* except that there is a larger representation below 20 kHz and no expanded isofrequency contour. IC regions devoted to echolocation (30 to 60 kHz) and passive listening (<30 kHz) are segregated, with the dorsal and lateral regions serving passive listening and the ventral region serving echolocation (Fuzessery and Hall 1999).




Figure 17.8. Tonotopic organization in the left IC of three species of bats. *White areas*, frequencies <60 kHz; *gray areas*, >60 kHz. In *Pteronotus*, the light gray area is the expanded 60-63 kHz region and the dark gray area is 64 kHz and higher. *Arrows*, tonotopic axis from low-to-high frequency. In *Antrozous*: *ICd*, dorsal IC region; *ICl*, lateral IC region; *ICv*, ventral IC region. In *Pteronotus*: *AL*, Anterolateral region IC; *DP*, dorsal posterior IC region; *M*, medial IC region. Orientation: *d*, dorsal; *l*, lateral; *m*, medial; *v*, ventral.

#### 4.1.2. Specialized Tonotopy

The cochlear specializations in *Pteronotus* bias the auditory system at every level toward the second harmonic of the CF echolocation signal, at about 61 kHz (Kössl and Vater 1985; Henson and Henson 1991; Kössl et al. 1999). Thus, 61 kHz occupies its expected place in subcollicular tonotopic sites (Vater et al. 1985; Zook and Leake 1989). However, in the IC, this region is displaced to form the dorsoposterior region (DP), and frequencies >63 kHz are expanded concomitantly, forming a large medial division (M) while those <60 kHz are greatly reduced (Pollak and Casseday 1989). To accommodate this expansion, the sequence is convoluted and semicircular. However, in *Rhinolophus*, another CF-FM bat with an “acoustic fovea” created by cochlear specializations, ICC frequency organization resembles that in FM bats and other mammals (Rübsamen and Schäfer 1990).

#### 4.2. TONOTOPY IN THE AVIAN INFERIOR COLLICULUS

Birds have an IC tonotopic orientation like that in mammals (Carr and Code 2000) with low best frequencies dorsally and higher frequencies ventrally (chicken: Coles and Aitkin 1979; guinea fowl: Heil and Scheich 1986; owl: Wagner et al. 1987, 2002; zebra finch: Woolley and Casseday 2004). There is a monotonic relationship between best frequency and recording depth, although best frequency changes more slowly in the low-frequency region (Wagner et al. 2002).

Owl ICCc frequency tuning curves are typically single peaked and symmetrical, with steep slopes on both flanks (Fig. 17.9A–C). Tuning curve widths are related to best frequency (BF), such that the ratio of tuning width to BF decreases as BF increases (Wagner et al. 2002). They are also related to IC subdivisions: lateral shell neurons have broader tuning curves and ICX cells prefer noise (Knudsen and Konishi 1978; Wagner et al. 1987; Mazer 1997). In the chicken and zebra finch, BFs range from 1 to 6 kHz and tuning curve widths are variable (Coles and Aitkin 1979; Woolley and Casseday 2004; Fig 17.9D–F). Most zebra finch IC neurons have V-shaped tuning curves, with a quarter having multiple peaks, noncontiguous excitatory regions, or narrow tilted or columnar shapes. Double peaked tuning curves occur in about 12% of chicken neurons (Coles and Aitkin 1979), and single- and multi-peaked tuning curves are found in awake guinea fowl (Scheich et al. 1977). Temporal response patterns in the chicken and zebra finch revealed several categories (Coles and Aitkin 1979; Woolley and Casseday 2004). In anesthetized finches, 49% of IC neurons had onset responses, 20% primary-like, 19% sustained, and 12% primary-like with notch (Woolley and Casseday 2004).






Figure 17.9. Avian IC tuning curves. (A–C) The barn owl ICC is tonotopic with best frequency increasing from dorsal (A) to ventral (C). *Left column*, isointensity frequency response functions with pure tone stimuli. *Right column*, dot raster plots for the same response functions. (D–F) Excitatory tuning curves for chicken and zebra finch. (D) Mid-frequency tuning curves in chicken. Narrow (E) and broad (F) tuning curves in zebra finch. (From Coles and Aiken 1979; Wagner et al. 2002; and Woolley and Casseday 2004.)




Figure 17.10. (A) Response areas in *Eptesicus* IC. The range between 15 and 35 kHz (lower panel) is expanded to better illustrate narrow, level-tolerant frequency response areas. Gray shading, a V-shaped frequency response area for comparison. (B) Neurons with upper thresholds and closed frequency response areas. (From Casseday and Covey 1992.)

## 5. RESPONSE PROPERTIES OF INFERIOR COLLICULUS NEURONS

In all vertebrates the auditory midbrain receives excitatory, inhibitory, and modulatory input from many sources, arranged as overlapping and intersecting gradients. The inputs terminate in various ways on different neuronal types, each with specific membrane properties, creating heterogeneous response properties (see Chapters 2, 10, and 11). It is often possible to relate neuronal responses to sound attributes such as frequency, amplitude, duration, the rate and/or depth of modulation in both amplitude and frequency, and the interval separating sounds. It has been possible to localize specialized response properties to particular IC regions in both bats and birds.

### 5.1. FUNCTIONAL CHARACTERISTICS OF BAT INFERIOR COLLICULUS CELLS

#### 5.1.1. Frequency and Amplitude Tuning

Single neurons have many types of frequency response areas, including simple V-shaped tuning curves, multi-peaked areas, closed response areas with lower and upper thresholds, and narrow, level-tolerant “filter” types (Casseday and Covey 1992). (See Fig. 17.10.)

In CF-FM bats with cochlear specializations, frequency response areas of “foveal” neurons are largely determined by cochlear mechanical properties (Suga et al. 1975). These neurons typically respond to a 1- to 2-kHz frequency band, with their tuning breadth constant across amplitude. This “filter-type” response area (Fig. 17.10A) contrasts with the V-shaped frequency response areas of neurons tuned to other frequencies.

Although *Eptesicus* has no cochlear specializations, it too has neurons with filter-type frequency response areas, tuned to a 1- to 2-kHz band near the bat’s quasi-CF calls. Filter-type response areas have not been found below the level of the IC in *Eptesicus* (Haplea et al. 1994). Frequency tuning can be narrowed considerably by synaptic inhibition (Yang et al. 1992; Pollak and Park 1993; Fuzessery and Hall 1996). In FM bats without cochlear specializations, inhibition produces the same filter property that is mechanically derived in other species, an example of convergent evolution. Perhaps narrow, level-tolerant frequency response areas are essential for processing CF or quasi-CF calls and echoes.

Many IC neurons have nonmonotonic rate-level functions: as amplitude grows their activity increases up to a point and declines or saturates (Suga 1968; Casseday and Covey 1992). Because neurons with nonmonotonic rate-level functions have a variety of “best amplitudes” (Fig. 17.10B), some respond only to the loud echolocation call, others only to the faint echoes. Amplitude tuning





Figure 17.11. (A) A combination-sensitive delay tuned neuron in *Pteronotus* IC. *a, b*: Responses to 83-kHz and 27-kHz tones alone. *c, d*: Responses when the 27-kHz tone was followed by the 83-kHz tone at two different interstimulus intervals (ISI). *e*: Mean spikes/tone pair as a function of ISI. *f*: The neuron had one excitatory response area near the fundamental (first) harmonic of the echolocation call and another near the third harmonic. (From Portfors and Wenstrup 1999.) (B) Paradoxical latency shift (PLS; left) in *Eptesicus* IC. Latency to an 85-dB tone was almost 15 ms longer than to a 45-dB tone. Diagram (top right) of a circuit that could produce delay-tuning if neurons A and B have a similar PLS. Because FM1 is loud in the emitted call, neuron A's latency would be 30 ms, while B's response to the attenuated nth echo harmonic would be 15 ms. The combined EPSP evoked by A in response to the call and in B in response to the echo would be maximal at neuron C when there is a 15-msec echo delay.

might provide a mechanism for segregating responses to calls and echoes for parallel processing (Rübsamen et al. 1988).

### 5.1.2. Combination Sensitivity and Delay-Tuning



All bat species have IC neurons with multi-peaked frequency response areas (Portfors and Wenstrup 2002). Many neurons are facilitated by combinations of sounds at specific frequencies, with the response to both tones significantly larger than the sum of either alone (Fig. 17.11A). Such facilitation is combination sensitivity and was first described in the intercollicular area of *Eptesicus* (Feng et al. 1978) but is common in the auditory cortex (AC) (Suga et al. 1979; O'Neill and Suga 1979), thalamus (Olsen and Suga 1991), and IC (Mittmann and Wenstrup 1995; Dear and Suga 1995; Portfors and Wenstrup 2001). In one form of combination sensitivity, two frequencies facilitate each other only when presented sequentially in a specific temporal relationship (Fig. 17.11A). Because facilitation depends on the interstimulus interval, such neurons are "delay tuned."

In *Pteronotus*, delay-tuned neurons require tones of different frequencies to produce facilitation. The initial stimulus must often be within the frequency range of the first (fundamental) harmonic of the FM portion of the echolocation call (FM1). Because the FM1 amplitude is weak relative to that of the other harmonics, it likely will be below threshold in the echoes. The second stimulus must be within the frequency range of the second, third, or fourth FM harmonic (FM2, FM3, or FM4, collectively designated FMn). In *Pteronotus* AC, neurons tuned to different delays between FM1 and FMn are organized systematically (O'Neill and Suga 1982). However, there is no topographic organization of best delays in the IC, nor any systematic relation between response latency and best delay (Portfors and Wenstrup 2001), suggesting that "mapping" of best delay occurs at the AC.

The delay between the emitted call and its echo is proportional to the distance of the reflective object; the best delays in IC, MGB, and AC neurons correspond to the range of values for realistic object distances. Therefore, combination-sensitive delay-tuned neurons may be a specialization for echolocation. Other combination-sensitive delay-tuned neurons are facilitated by sounds in frequency combinations that are harmonically unrelated and/or that are outside the frequency range of echolocation calls (Leroy and Wenstrup 2000), and may be involved in processing communication sounds.

## 5.2. MECHANISM FOR DELAY-TUNING

Most work on delay-tuning mechanisms has used *Pteronotus*. One hypothesis that flowed from Jeffress's work (1948) was that delay-tuned neurons act as coincidence detectors, although on a much longer time scale than for sound localization. In this model, the delay-tuned neuron receives delayed excitatory input from FM1 neurons and rapid excitatory input from FMn-tuned neurons, firing only when the FM1-FMn delay compensates for the delay in the FM1



pathway, making the inputs coincident. Delay-tuned neurons respond to certain interpulse intervals with a peak “best delay” (O’Neill and Suga 1982), consistent with partial coincidence of excitatory postsynaptic potentials (EPSPs) at some delays and complete coincidence at the peak of the delay-tuning function. However, glycine antagonists abolish delay-tuning in IC neurons that show facilitation (Wenstrup and Leroy 2001), suggesting that inhibition is involved.

One mechanism might be paradoxical latency shift (PLS) (Sullivan 1982), presumably using neural inhibition. For most cells, latency decreases with increasing sound level, but paradoxical latency neurons respond with longer latencies as sound level grows (Fig. 17.11B). Some *Eptesicus* IC neurons show PLS due to increasing strength and duration of inhibition at higher sound levels (Covey et al. 1996). PLS cells would respond to the loud echolocation call with a longer latency than to the faint echo. The amount of latency shift would determine best delay (Fig. 17.11B). Since the latency shift varies according to sound level, this may be a mechanism for PLS neurons that changes best delay as a function of amplitude, and it might explain why blocking glycinergic inhibition abolishes delay-tuning.

### 5.3. DURATION TUNING

Duration sensitive neurons (Fig. 17.12A) are common in the bat IC. Duration tuned neurons typically have a peak response at some best duration (Pinheiro et al. 1991; Casseday et al. 1994; Ehrlich et al. 1997; Fuzessery and Hall 1999) with best durations near the range of call durations used by bats while echolocating. In *Eptesicus* one third of IC cells are duration tuned. They are mainly in the caudal half of the IC, and have not been found below the IC. The underlying mechanisms involve convergence of excitatory and inhibitory inputs with different latencies and discharge patterns (Casseday et al. 1994, 2000). Blocking inhibition abolishes duration tuning in nearly all such neurons (Casseday et al. 2000) and in a subset of these cells in *Antrozous* (Fuzessery and Hall 1999).

Figure 17.12. (A) Duration sensitive *Eptesicus* IC cells. Composite dot rasters (*left*) from band-pass (*top*) and long-pass (*bottom*) cells with 10 presentations/duration (*horizontal bars*). Spike count vs. duration (*right*) at several sound levels. Duration sensitivity is level tolerant. (From Faure et al. 2003.) (B) Model for duration sensitivity created by the interaction of excitatory and inhibitory inputs with different time courses and latencies. The band-pass neuron (*top traces*) receives a transient subthreshold EPSP with a 10-ms onset latency and a sustained 4-ms latency IPSP, followed by an excitatory rebound. The neuron does not reach threshold unless sound duration is such that the EPSP coincides with the rebound from inhibition. Best duration reflects latency differences between the EPSP and IPSP. A short-pass neuron would result when the difference is very small. The long-pass neuron (*bottom traces*) receives a sustained EPSP with a 5-ms latency which, alone, is suprathreshold. The neuron also receives a transient IPSP with similar latency. At short durations, the EPSP is cancelled by the IPSP, but at longer durations the later part of the EPSP is unaffected and elicits a response.

All *Eptesicus* duration tuned neurons receive inhibition for the stimulus duration and have a shorter latency than for excitation. At the end of the inhibitory period, corresponding to stimulus offset, an excitatory rebound occurs that, alone, is subthreshold (Covey et al. 1996). If the sound duration allows the rebound to coincide with the transient excitation evoked by sound onset, facilitation occurs and the neuron responds (Fig. 17.12B). Thus, the cell's best duration reflects the difference between the latencies of inhibition and excitation. Consistent with this model, duration tuned neurons respond at the stimulus offset.

Long-pass neurons' duration sensitivity results when a neuron receives sustained excitation and transient inhibition at about the same latency (Faure et al. 2003). At short durations the excitation is sufficiently brief to be nullified by inhibition, and at longer durations it outlasts the inhibition and drives the neuron (Fig. 17.12B).

#### 5.4. TUNING TO FREQUENCY AND AMPLITUDE MODULATIONS

All bats use FM sweeps in their echolocation call repertoire. Some IC neurons respond only to frequency sweeps and many of these are selective for the direction of frequency change (Suga 1969; Fuzessery 1994). IC neurons preferring either downward or upward FM (Fuzessery 1994) may process communication calls and echolocation signals.

For bats that use CF or quasi-CF calls, insect wing beats can produce periodic amplitude modulations and Doppler shifts in the echo frequency. Echoes of FM calls may contain significant amplitude modulations from interference patterns of reflections from multiple surfaces. Sinusoidal amplitude modulations (SAM) and sinusoidal frequency modulations (SFM) are artificial stimuli that mimic these patterns of amplitude and frequency change. Half of IC neurons respond to sinusoidally modulated stimuli with periodic, phase-locked discharges (Schuller 1979; Casseday et al. 1997). Some *Rhinolophus* foveal cells respond to SAM of as little as 3% depth and to SFMs with modulation depths as small as 0.025%.

IC neurons' discharge patterns in response to pure tones are related to their responses to modulated stimuli. *Pteronotus* ON-OFF neurons are unresponsive to SAM (Lesser et al. 1990) and receive an initial inhibitory input with a shorter latency than that of excitation. When presented with pairs of tones separated by a short interval, both populations experience a phenomenon similar to backward masking. These cells do not code periodic amplitude modulations because the response to the earlier stimulus is suppressed by the short-latency inhibition evoked by the second stimulus. Because many IC neurons receive inhibition before excitation (Covey et al. 1996), backward masking may set the upper limit for modulation rate. IC neurons respond to SAM only up to 100 to 200 Hz (Reimer 1987) even though the presynaptic cells respond to higher modulation rates. Because blocking GABAergic and glycinergic inhibition (Burger and Polak 1998) has little effect on SAM tuning, intrinsic properties may be more important than inhibition for tuning to AM rate.



Some *Eptesicus* IC neurons respond only to SFM (Casseday et al. 1997), requiring several cycles to become primed. Perhaps each cycle of frequency modulation elicits an inhibitory event followed by a subthreshold excitatory event. If the modulation rate for the excitatory rebound that follows the inhibition from one half-cycle coincides with the excitation evoked by the next half-cycle, the neuron reaches threshold and fires. Blocking inhibition broadens SFM tuning (Koch and Grothe 1998), suggesting that neural inhibition may contribute to tuning to SFM rate.

### 5.5. ROLE OF THE BAT INFERIOR COLLICULUS IN BINAURAL HEARING AND SOUND LOCALIZATION

Bat IC neurons, like those of other mammals, have many binaural interaction patterns, but there is no evidence of a space map like that in the barn owl. Because most mammals move their eyes, the relationship between the visual and auditory fields is variable. Many mammals also have mobile pinnae, complicating the relation between the head and the auditory field.

Many bats have large, complex, and mobile pinnae that are highly directional and accentuate ILDs. They also produce characteristic spectral patterns that could aid in localization in both azimuth and elevation (Fuzessery and Pollak 1984; Wotton et al. 1995). The response patterns of all IC neurons likely reflect the pinna's characteristics, imparting some spatial tuning, especially at sound levels just above threshold (Grothe et al. 1996).

IC binaural properties are not simply inherited from LSO or MSO, but instead reflect integration of excitatory and inhibitory synaptic input. Both GABAergic and glycinergic inhibition actively shapes binaural response properties (Vater et al. 1992a; Park and Pollak 1993, 1994; Klug et al. 1995; Pollak 1997). Some cells experience inhibition when sound is presented at one ear and excitation when it is presented at the other, and the latency and time course of each is different (Covey et al. 1996). Because binaural processing does not construct a fixed auditory space representation, it may have other roles such as enhancing the detection of signals in noise, reducing echo clutter, or coding sound source motion.

Specific binaural subclasses are excited by inputs from the contralateral ear and inhibited by the ipsilateral ear (EI) or excited by inputs from the contralateral and ipsilateral ear (EE). Cells representing these classes are segregated spatially and there are progressive shifts in the excitatory–inhibitory balance of inputs from the ears that form gradients across the IC. When ILD function 50% points (or cutoffs) are measured across an isofrequency contour, the points create a gradient in which ILD functions extend into ipsilateral space (Wenstrup et al. 1986). However, spatial receptive field (RF) properties interact with sound amplitude, frequency spectrum, modulation pattern, sound source movement, and other stimulus aspects (Grothe et al. 1996), so individual neurons, or even populations, are unlikely sources for unambiguous spatial information.

### 5.6. INTERACTIVE PROCESSING

Convergent inputs differentially tuned to many sound parameters suggest that neural responses are interactive. Stimulus amplitude or spectrotemporal changes affect spatial RF size and shape (Grothe et al. 1996; Wu and Jen 1996) and stimulus repetition rate can influence latency, rate-level functions, thresholds, frequency tuning, and duration tuning (Jen and Chen 1998; Jen et al. 2001; Zhou and Jen 2001). Some interactions depend on GABAergic inhibition (Jen et al. 2002). The sound-evoked sequence of synaptic currents can exceed the sound's duration, imposing context-dependent alterations of the subsequent responses (Covey et al. 1996).

### 5.7. PLASTICITY OF RESPONSES

IC neural response properties show long-term plastic changes caused by corticocollicular input (Suga et al. 2002; Suga and Ma 2003). Tuning changes can reflect behavioral conditioning (Gao and Suga 1998) or experimental manipulations of AC, including electrical stimulation (Yan and Suga 1996) or inactivation (Yan and Suga 1999). Effects on frequency tuning (Zhang et al. 1997), interstimulus interval (Yan and Suga 1996), and sound duration (Ma and Suga 2001) are seen. Conditioned changes in responses require 15 to 30 minutes of stimulation under constant conditions and last for hours (Ji et al. 2001). Cortical manipulations have similar effects on all parameters tested. In *Pteronotus*, AC electrical stimulation in an area where neurons are tuned to a particular parameter value augments IC responses of cells tuned to that parameter value and decreases that of neurons tuned to different values of the same parameter, perhaps enhancing the representation of a common sound.

## 6. FUNCTIONAL CHARACTERISTICS OF AVIAN MIDBRAIN NEURONS

Save for barn owls and songbirds, we do not know what acoustic properties are behaviorally salient to birds (Konishi 1985). Nevertheless, IC responses in barn owl, chicken, and zebra finch IC are similar, and principles learned from studies of the barn owl IC may apply to other birds (Knudsen et al. 1979).

Barn owls can localize sound precisely, and their midbrain responses are dominated by binaural computations subserving localization (Keller and Takahashi 1996; Konishi 2000). Their IC neurons are EE or EI (Moiseff and Konishi 1981) and sensitive to changes in ILD and ITD. Similarly, most chicken neurons are binaural, 46% being EI, 21% EE, 8% II (inhibited by either ear), and only 17% are monaural (Coles and Aitkin 1979).

Physiological properties are subdivision specific. The ICCc contains ITD-sensitive neurons and projects to the contralateral lateral shell, where ITD in-

formation is combined with ILD information (Fig. 17.13; Coles and Aitkin 1979; Takahashi et al. 1987a; Wagner et al. 1987; Volman and Konishi 1989; Spezio and Takahashi 2003). Thus, the progression from ICC to ICX is associated with the abstraction of stimulus features required to construct the ICX auditory space map (Mazer 1997).

### *6.1. INTERAURAL TIME DIFFERENCE TUNING*

ITD is the principal cue for auditory azimuth (Moiseff and Konishi 1981) and is computed in two stages: first, brain stem NL neurons act as coincidence detectors to encode interaural phase difference, firing maximally when simultaneously receiving inputs from both ears (Carr and Konishi 1990; Peña and Konishi 2001). At this stage, ambiguities exist about the correspondence between the NL response and the actual ITD in auditory space. The second stage of ITD computation occurs in the IC, where across-frequency integration filters phase-ambiguous side peaks, creating neurons coding the true ITD (Takahashi and Konishi 1986; Peña and Konishi, 2000).

The ICCc receives NL input and contains sharply frequency-tuned ITD sensitive neurons with primary-like responses (Fig. 17.13; Coles and Aitkin 1979; Wagner et al. 1987). In the ICCc one ITD activates a neural array tuned to many different frequencies (Fig. 17.13F; Wagner et al. 1987). This array code is transformed into unambiguous space-specific neuron responses as follows. Each array projects to ITD- and ILD-sensitive neurons in the contralateral lateral shell (ICCLs), which project to space-specific neurons in the contralateral ICX, endowing them with ITD selectivity and therefore azimuth coding (Takahashi and Konishi 1986; Mazer 1997; Peña and Konishi 2000).

### *6.2. TUNING TO LEVEL AND INTERAURAL LEVEL DIFFERENCE PROCESSING*

Less is known about ILD processing. In the barn owl, the vertical asymmetry in ear directionality makes ILD a cue for the vertical target axis at high frequencies. Level is encoded by CN neurons, most having monotonic rate level functions (Köppl and Carr 2003). ILD sensitivity first emerges in the dorsal nucleus of the lateral lemniscus, posterior part (LLDp) whose neurons are excited by contralateral stimulation and inhibited by ipsilateral stimulation (Takahashi and Keller 1992; Takahashi et al. 1995) and are EI cells coding ILD (Manley et al. 1988; Mogdans and Knudsen 1994). LLDp therefore is analogous to LSO (Tsuchitani 1977; Takahashi et al. 1995). LLDp projects bilaterally to the ICCLs, endowing these cells with ILD sensitivity (Adolphs 1993).

LLDp neurons do not unambiguously encode ILD. Although they prefer sound at the contralateral ear, they are also sensitive to changes in average binaural level. ILD tuning gradually emerges in the ICCLs (Mazer 1998). Response latencies are about 5 ms near the central core, and >10 ms at the ICX




Figure 17.13. (A) ITD tuning curves in the barn owl ICCc. *Left*, ITD tuning curves; *right*, raster plots. Positive ITDs are right-ear leading. Tuning curves (*a*, *b*, *d*) from low- and high-frequency neurons. (B) The stimulus is an 800-Hz tone; all others are broadband noise. Note the cyclical tuning, even in response to noise stimuli, and the inhibition of spontaneous activity at stimulus termination. (D, E) Calretinin immunohistochemistry labels the ICCc; *arrow*, best frequency axis. (F) The ensemble code for ITD in the ICC. The array is sensitive to interaural phase differences, but each response shows phase ambiguity. (Modified from Wagner et al. 1987, 2002; Kubke et al. 1999.)

border. Frequency tuning widths broaden with increasing lateral position, while ITD and ILD tuning widths sharpen. Almost all ICCls neurons are sensitive to both time and intensity cues (Mazer 1997).

Unlike the ITD coding array in the ICCc, a topographical representation of ILD is never observed in the ICCls (Mazer 1998). Recordings from the space specific neurons show instead that ILD varies as a function of frequency in a

complex manner for any given location (Fig. 17.14A). In ICX, ILD-alone RFs are generally horizontal swaths of activity at the elevation of the cell's normal spatial RF (Fig. 17.14A). An ITD-alone RF forms a vertical swath at the azimuth of the cell's normal RF, which thus lies at the intersection of the ITD- and ILD-alone RFs (Euston and Takahashi 2002). ILD sensitive neurons in the chicken are EI and encode a 10- to 15-dB range (Coles and Aitkin 1979), with responses like those in the barn owl.

### 6.3. THE MAP OF AUDITORY SPACE IN THE EXTERNAL NUCLEUS

The ICX space-specific neurons respond to sound only from a particular spatial locus and, when spontaneously active, are inhibited by stimulation of either ear (II) (Fig. 17.15; Knudsen and Konishi 1978; Knudsen and Knudsen 1983; Takahashi and Konishi 1986) and they are selective for combinations of ITD and ILD. Driven by noise, they do not show phase ambiguity, and thus differ from the ICCc ITD sensitive cells that provide their input (Peña and Konishi 2000). The phase-unambiguous response of space-specific neurons has been explained as follows. They receive inputs via the ICCIs from many ICCc isofrequency laminae (Knudsen 1983), presumably from the ITD-specific arrays (Wagner et al. 1987). These inputs interact at the postsynaptic cell: peaks signaling the correct ITD superimpose and add, while secondary, ambiguous peaks cancel by interacting with inhibitory sidebands from other or ambiguous frequencies (Takahashi and Konishi 1986; Keller and Takahashi 1996, 2000). The inhibitory interactions between different frequency channels may originate from GABAergic ICX neurons (Fujita and Konishi 1991).

ICX neurons act like analog AND gates for ITD and ILD, such that the two inputs are multiplied (Fig. 17.14B–E). Multiplication of separate postsynaptic potentials (PSPs) tuned to ITD and ILD, rather than an additive process, explains their subthreshold responses to ITD-ILD pairs. Comparing subthreshold PSPs and spike output for the same ICX neurons show that RFs measured in PSPs exceed those measured in spikes in both ITD and ILD dimensions. Thus, a spike threshold mechanism mediates formation of the restricted space specific RF, the stimulus-induced first spike having a lower threshold than subsequent or spontaneous spikes (Peña and Konishi 2001).

The spatially restricted ICX RFs are still much larger than the minimum detectable change in sound source location, which can be 3°. However, changes in neuronal activity across the space map show that most neurons can reliably signal source location changes smaller than the behavioral threshold. Each source is represented in the space map by a focus of activity in a neural population, and source displacement changes their activity pattern (Takahashi et al. 2003). This map of contralateral auditory space projects topographically to the optic tectum (OT), whose visual and auditory space maps are in register (Knudsen and Knudsen 1983). OT activity directs the owl's rapid head movements to auditory and visual stimuli (du Lac and Knudsen 1990).




Figure 17.14. Formation of space-specific responses in the barn owl ICX. (A) ITD and ILD contributions to the spatial RF of an ICX cell. Diamond, spikes (gray) evoked by a source at frontal hemisphere loci. *First column*: ILD-alone RFs for three cells obtained by filtering broadband noise with HRTFs altered to present the ILD spectra for each spatial location while holding the ITD constant at a cell's preferred value. *Second column*:

#### 6.4. THE ROLE OF EXPERIENCE IN FORMATION OF THE SPACE MAP

The auditory space map has become a paradigm for studies of experience-dependent plasticity (Knudsen 2002). Manipulating sensory experience reorganizes ICX in a manner congruent with behavioral learning, as accurate auditory orienting behavior is a measure for adaptive adjustment in sound localization after peripheral changes in sensory experience such as disrupting auditory cue values for spatial loci by ear plugging (Knudsen et al. 1984; Brainard and Knudsen 1995; Knudsen 1999). Owls with an earplug first err toward the open ear, then recover. On earplug removal, their orienting errors are in the opposite direction, and these resolve with experience. A second manipulation alters visual and auditory correspondence with prismatic spectacles that displace the visual field (Brainard and Knudsen 1998). Such owls learn new associations between auditory and visual cues to recalibrate both worlds.

Learning changes forebrain and midbrain neuron tuning to sound localization cues. In the midbrain space map, the assay for plasticity is precise and can be quantified for each neuron (Fig. 17.15; Knudsen 1982). Adaptive changes are centered in the ICX and depend on changes in axonal projections and adjustments in synaptic strength (Knudsen 2002; Nieder et al. 2003). In young owls with either earplugs or prismatic displacement, ICX tuning to sound localization cues alters adaptively to coordinate the auditory and the visual RFs (Fig. 17.15; Mogdans and Knudsen 1994). Gradually, these new responses strengthen, while those to the prior ITD range disappear. These changes are correlated with axonal remodeling of the topographic projection from the ICC to the ICX. Prism experience appears to induce the formation of modified circuitry in the ICX at least in part through axonal sprouting and synaptogenesis. Normal circuitry also persists, showing that alternative plastic and normal circuits coexist in this network.

Both *N*-methyl-D-aspartate (NMDA) and GABA<sub>A</sub> receptor changes are implicated in plasticity. NMDA receptors regulate expression of newly learned responses (Feldman and Knudsen 1998). In an ICX expressing a maximally shifted

---

Figure 17.14. *Continued*

the ITD-alone RFs. *Third column*: combining the plots in these columns yields the predicted spatial RFs. *Fourth column*: spatial RFs measured directly from the cell. Negative azimuths and elevations denote loci on the left of the midline and below eye level, respectively. (Modified from Takahashi et al. 2003.) **(B)** Space specific neuron showing PSPs to different ITD–ILD pairs. *Dotted lines*, mean resting potential. **(C)** Spiking responses of the same neuron to different ITD–ILD pairs. The large peak is the excitatory center and the flat area around it is the inhibitory surround (compare **B**, **C**). Negative (–) ITD and negative (–) ILD mean, respectively, sound in ipsilateral ear leading and louder. **(D)** An ICX neuron and its axon projecting to the OT. **(E)** The same neuron labeled with neurobiotin. (From Peña and Konishi 2001.)




Figure 17.15. The auditory space map in the ICX before (A) and after (B) shifts produced by early prism experience. (A) The ICC-ICX (*black arrows*) projection conveys ITD and other information and converges across frequency channels, creating spatially restricted auditory RFs and a space map. The auditory and visual maps of space merge in OT and receives a reciprocal instructive signal from it. (B) The schematic representation of the anatomical change (*dark arrows*) in the ICC-ICX projection resulting from early prism experience. For each axis of prismatic displacement, an abnormal rostralward projection of ICC axons (*dark gray arrows*) appears on one side of the brain and a caudalward projection appears contralaterally (data not shown), although the normal projection persists (*light gray arrows*). (Modified from Knudsen 2002.)

ITD map, focal blockade of GABA elicited the immediate appearance of normal responses. Thus, in a shifted-ITD map, synapses that support normal responses remain patent and coexist with synapses that support learned responses, but responses to the normal synapses are selectively nullified by GABAergic inhibition (Zheng and Knudsen 1999).

Changes in the ICX auditory space map are directed by a topographic map of visual space whose dominance is plausible since the pathway's primary function is to trigger gaze towards auditory targets (Knudsen et al. 1993; Wagner 1993). The power and precision of the visual instructive signal is shown when a small lesion in the tectum eliminates adaptive adjustments (Hyde and Knudsen 2002).

## 7. COMPARING BIRDS AND BATS

The auditory midbrain structures of birds and bats are probably homologous, and serve many similar purposes. The IC integrates information converging on it from many parallel pathways that will ultimately contribute to sound source localization and decoding of the auditory signals. The IC is an important source of output to motor pathways for orientation, vocalization, and other behaviors. It is part of a massive system that extends to the forebrain, and it may participate



in adaptive changes from learning and peripheral modifications. However, the evolutionary paths of birds and bats diverged so long ago that many of the substrates through which these tasks are accomplished are very different. The avian cochlea, central auditory system, and forebrain are organized differently from those of mammals so that the patterns of connections differ considerably between the two groups.

Midbrain organization is more hierarchical in owls, especially with regard to the emergence of an invariant map of auditory space in the different IC subdivisions. The barn owl midbrain is dominated by binaural processing and may have more in common with mammals that do not echolocate. Finally, the streams of processing for sound location and temporal patterns are more sharply segregated in owls. Future work should reveal whether birds and bats share some of the same neural circuits and cellular mechanisms for information processing despite the fact that many structures and pathways are different.

## Abbreviations

AVCN	anteroventral cochlear nucleus
CF	constant-frequency
CN	cochlear nucleus
DCN	dorsal cochlear nucleus
DLL	dorsal nucleus of the lateral lemniscus
EE	neuron excited by contralateral and ipsilateral sound
EI	neuron excited by contralateral sound and inhibited by ipsilateral sound
EPSP	excitatory postsynaptic potential
FM	frequency-modulated
GABA	$\gamma$ -aminobutyric acid
GC	tectal gray
IC	inferior colliculus
ICC	central nucleus of the inferior colliculus
ICCc	central core of the central nucleus of the inferior colliculus
ICCl <sub>s</sub>	lateral shell of the central core of the inferior colliculus
ICCm <sub>s</sub>	medial shell of the central core of the inferior colliculus
ICX	external nucleus of the central core of the inferior colliculus
III	oculomotor nucleus
ILD	interaural level difference
ILL	intermediate nucleus of lateral lemniscus
IO	isthmo-optic nucleus
IPSP	inhibitory postsynaptic potential
ITD	interaural time difference
IV	trochlear nucleus
iv	trochlear nerve
LL	lateral lemniscus

LLDp	dorsal nucleus of lateral lemniscus, posterior part
LSO	lateral superior olive
MGB	medial geniculate body
MLd	mesencephalic nucleus, pars dorsalis
MSO	medial superior olive
NA	nucleus angularis
NL	nucleus laminaris
NM	nucleus magnocellularis
NMDA	<i>N</i> -methyl-D-aspartate
PLS	paradoxical latency shift
PVCN	posteroventral cochlear nucleus
RF	receptive field
SAM	sinusoidal amplitude modulation
SFM	sinusoidal frequency modulation
SOC	superior olivary complex
VLL	ventral nucleus of lateral lemniscus
VLLc	columnar nucleus of VLL

## Acknowledgments

This work was supported by NIH/NIDCD Grants P30 DC04661 and R01 DC00607 (E.C.) and P30 DC04664 and R01 DCD00436 (C.E.C.).

## References

- Adolphs R (1993) Bilateral inhibition generates neuronal responses tuned to interaural level differences in the auditory brainstem of the barn owl. *Journal of Neuroscience* **13**:3647–3668.
- Akesson TR, de Lanerolle NC, and Cheng MF (1987) Ascending vocalization pathways in the female ring dove: projections of the nucleus intercollicularis. *Experimental Neurology* **95**:34–43.
- Balthazart J, Foidart A, Wilson EM, and Ball GF (1992) Immunocytochemical localization of androgen receptors in the male songbird and quail brain. *Journal of Comparative Neurology* **317**:407–420.
- Bonke B, Bonke D, and Scheich H (1979) Connectivity of the auditory forebrain nuclei in the guinea fowl (*Numida meleagris*). *Cell and Tissue Research* **200**:101–121.
- Brainard MS and Knudsen EI (1995) Dynamics of visually guided auditory plasticity in the optic tectum of the barn owl. *Journal of Neurophysiology* **73**:595–614.
- Brainard MS and Knudsen EI (1998) Sensitive periods for visual calibration of the auditory space map in the barn owl optic tectum. *Journal of Neuroscience* **18**:3929–3942.
- Brew HM and Forsythe ID (1995) Two voltage-dependent K<sup>+</sup> conductances with complementary functions in postsynaptic integration at a central auditory synapse. *Journal of Neuroscience* **15**:8011–8022.
- Bruns V and Schmieszek E (1980) Cochlear innervation in the greater horseshoe bat: demonstration of an acoustic fovea. *Hearing Research* **3**:27–43.

- Burger RM and Pollak GD (1998) Analysis of the role of inhibition in shaping responses to sinusoidally amplitude-modulated signals in the inferior colliculus. *Journal of Neurophysiology* **80**:1686–1701.
- Carr CE and Code RA (2000) The central auditory system of reptiles and birds. In: Dooling RJ, Fay RR, and Popper AN (eds). *Springer Handbook of Auditory Research*, Volume 13: *Comparative Hearing: Birds and Reptiles*. Springer-Verlag, New York, pp. 197–248.
- Carr CE and Konishi M (1990) A circuit for detection of interaural time differences in the brainstem of the barn owl. *Journal of Neuroscience* **10**:3227–3246.
- Carr CE, Fujita I, and Konishi M (1989) Distribution of GABAergic neurons and terminals in the auditory system of the barn owl. *Journal of Comparative Neurology* **286**: 190–207.
- Casseday JH and Covey E (1992) Frequency tuning properties of neurons in the inferior colliculus of an FM bat. *Journal of Comparative Neurology* **319**:34–50.
- Casseday JH and Covey E (1996) A neuroethological theory of the operation of the inferior colliculus. *Brain, Behavior and Evolution* **47**:311–336.
- Casseday JH, Covey E, and Vater M (1988) Connections of the superior olive in the rufous horseshoe bat, *Rhinolophus rouxi*. *Journal of Comparative Neurology* **278**:313–329.
- Casseday JH, Ehrlich D, and Covey E (1994) Neural tuning for sound duration: role of inhibitory mechanisms in the inferior colliculus. *Science* **264**:847–850.
- Casseday JH, Covey E, and Grothe B (1997) Neurons specialized for periodic frequency modulations in the inferior colliculus of the big brown bat, *Eptesicus fuscus*. *Journal of Neurophysiology* **77**:1595–1605.
- Casseday JH, Ehrlich D, and Covey E (2000) Neural measurement of sound duration: control by excitatory-inhibitory interactions in the inferior colliculus. *Journal of Neurophysiology* **84**:1475–1487.
- Casseday JH, Fremouw T, and Covey E (2002) The inferior colliculus: hub of the auditory system. In: Oertel D, Popper AN, and Fay RR (eds). *Springer Handbook of Auditory Research*, Volume 15: *Integrative Functions in the Mammalian Auditory Pathway*. Springer-Verlag, New York, pp. 238–318.
- Clack JA (1997) The evolution of tetrapod ears and the fossil record. *Brain, Behavior and Evolution* **50**:198–212.
- Cohen YE and Knudsen EI (1999) Maps versus clusters: different representations of auditory space in the midbrain and forebrain. *Trends in Neurosciences* **22**:128–135.
- Coles RB and Aitkin LM (1979) The response properties of auditory neurones in the midbrain of the domestic fowl (*Gallus gallus*) to monaural and binaural stimuli. *Journal of Comparative Physiology A* **134**:241–251.
- Conlee JW and Parks TN (1986) Origin of ascending auditory projections to the nucleus mesencephalicus lateralis pars dorsalis in the chicken. *Brain Research* **367**:96–113.
- Covey E (1993) The monaural nuclei of the lateral lemniscus: parallel pathways from cochlear nucleus to midbrain. In: Merchán MA, Juiz JM, and Godfrey DA (eds). *The Mammalian Cochlear Nuclei: Organization and Function*. Plenum Press, New York, pp. 321–334.
- Covey E (2003) Brainstem mechanisms for analyzing temporal patterns of echolocation sounds: a model for understanding early stages of speech processing? *Speech Communication* **41**:151–163.
- Covey E and Casseday JH (1986) Connectional basis for frequency representation in the nuclei of the lateral lemniscus of the bat, *Eptesicus fuscus*. *Journal of Neuroscience* **6**:2926–2940.

- Covey E and Casseday JH (1991) The monaural nuclei of the lateral lemniscus in an echolocating bat: parallel pathways for analyzing temporal features of sound. *Journal of Neuroscience* **11**:3456–3470.
- Covey E and Casseday JH (1995) The lower brainstem auditory pathways. In: Popper AN and Fay RR (eds). *Springer Handbook of Auditory Research*, Volume 5: *Hearing by Bats*. Springer-Verlag, New York, pp. 235–295.
- Covey E, Hall WC, and Kobler JB (1987) Subcortical connections of the superior colliculus in the mustache bat, *Pteronotus parnellii*. *Journal of Comparative Neurology* **263**:179–197.
- Covey E, Vater M, and Casseday JH (1991) Binaural properties of single units in the superior olivary complex of the mustache bat. *Journal of Neurophysiology* **66**:1080–1094.
- Covey E, Kauer JA, and Casseday JH (1996) Whole-cell patch clamp recording reveals subthreshold sound-evoked postsynaptic currents in the inferior colliculus of awake bats. *Journal of Neuroscience* **6**:3009–3018.
- Dear SP and Suga N (1995) Delay-tuned neurons in the midbrain of the big brown bat. *Journal of Neurophysiology* **73**:1084–1100.
- Dezso A, Schwarz DWF, and Schwarz IE (1993) A survey of auditory brainstem nuclei in the chicken (*Gallus domesticus*) with cytochrome oxidase histochemistry. *Journal of Otolaryngology* **22**:385–390.
- Dooling RJ, Lohr B, and Dent ML (2000) Hearing in birds and reptiles. In: Dooling RJ, Popper AN, and Fay RR (eds). *Springer Handbook of Auditory Research*, Volume 13: *Comparative Hearing: Birds and Reptiles*. Springer-Verlag, New York, pp. 308–359.
- du Lac S and Knudsen EI (1990) Neural maps of head movement vector and speed in the optic tectum of the barn owl. *Journal of Neurophysiology* **63**:131–146.
- Durand SE, Tepper JM, and Cheng M-F (1992) The shell region of the nucleus ovoidalis: a subdivision of the avian auditory thalamus. *Journal of Comparative Neurology* **323**:495–518.
- Ehrlich D, Casseday JH, and Covey E (1997) Neural tuning to sound duration in the inferior colliculus of the big brown bat, *Eptesicus fuscus*. *Journal of Neurophysiology* **77**:2360–2372.
- Euston DR and Takahashi TT (2002) From spectrum to space: the contribution of level difference cues to spatial receptive fields in the barn owl inferior colliculus. *Journal of Neuroscience* **22**:284–293.
- Faure PA, Fremouw T, Casseday JH, and Covey E (2003) Neural manifestations of temporal masking reveal properties of sound-evoked inhibition in duration tuned neurons of the inferior colliculus. *Journal of Neuroscience* **23**:3052–3065.
- Feldman DE and Knudsen EI (1998) Experience-dependent plasticity and the maturation of glutamatergic synapses. *Neuron* **20**:1067–1071.
- Feng AS, Simmons JA, and Kick SA (1978) Echo detection and target-ranging neurons in the auditory system of the bat *Eptesicus fuscus*. *Science* **202**:645–648.
- Frisina RD, O'Neill WE, and Zettel ML (1989) Functional organization of mustached bat inferior colliculus: II. Connections of the FM<sub>2</sub> region. *Journal of Comparative Neurology* **284**:85–107.
- Fubara BM, Casseday JH, Covey E, and Schwartz-Bloom RD (1996) Distribution of GABA<sub>A</sub>, GABA<sub>B</sub> and glycine receptors in the central auditory system of the big brown bat, *Eptesicus fuscus*. *Journal of Comparative Neurology* **369**:83–92.
- Fujita I and Konishi M (1991) The role of GABAergic inhibition in processing of interaural time difference in the owl's auditory system. *Journal of Neuroscience* **11**:722–739.

- Fuzessery ZM (1994) Response selectivity for multiple dimensions of frequency sweeps in the pallid bat inferior colliculus. *Journal of Neurophysiology* **72**:1061–1079.
- Fuzessery ZM, Buitenhoff P, Andrews B, Kennedy JM (1993) Passive sound localization of prey by the pallid bat (*Antrozous p. pallidus*). *Journal of Comparative Physiology, A* **171**:767–777.
- Fuzessery ZM and Hall JC (1996) Role of GABA in shaping frequency tuning and creating FM sweep selectivity in the inferior colliculus. *Journal of Neurophysiology* **76**:1059–1073.
- Fuzessery ZM and Hall JC (1999) Sound duration selectivity in the pallid bat inferior colliculus. *Hearing Research* **137**:137–154.
- Fuzessery ZM and Pollak GD (1984) Neural mechanisms of sound localization in an echolocating bat. *Science* **225**:725–728.
- Gahr M, Guttinger HR, and Kroodsma DE (1993) Estrogen receptors in the avian brain: survey reveals general distribution and forebrain areas unique to songbirds. *Journal of Comparative Neurology* **327**:112–122.
- Gao E and Suga N (1998) Experience-dependent corticofugal adjustment of midbrain frequency map in bat auditory system. *Proceedings of the National Academy of Sciences of the United States of America* **95**:12663–12770.
- Grinnell AD (1995) Hearing in bats: an overview. In: Popper AN and Fay RR (eds). *Springer Handbook of Auditory Research, Volume 5: Hearing by Bats*. Springer-Verlag, New York, pp. 1–36.
- Grothe B, Vater M, Casseday JH, and Covey E (1992) Monaural interaction of excitation and inhibition in the medial superior olive of the mustached bat: an adaptation for biosonar. *Proceedings of the National Academy of Sciences of the United States of America* **89**:5108–5112.
- Grothe B, Schweizer H, Pollak GD, Schuller G, and Rosemann C (1994) Anatomy and projection patterns of the superior olivary complex in the Mexican free-tailed bat, *Tadarida brasiliensis mexicana*. *Journal of Comparative Neurology* **343**:630–646.
- Grothe B, Covey E, and Casseday JH (1996) Spatial tuning of neurons in the inferior colliculus of the big brown bat: effects of sound level, stimulus type and multiple sound sources. *Journal of Comparative Physiology A* **179**:89–102.
- Habbicht H and Vater M (1996) A microiontophoretic study of acetylcholine effects in the inferior colliculus of horseshoe bats: implications for a modulatory role. *Brain Research* **724**:169–179.
- Haplea S, Covey E, and Casseday JH (1994) Frequency tuning and response latencies at three levels in the brainstem of the echolocating bat, *Eptesicus fuscus*. *Journal of Comparative Physiology A* **174**:671–683.
- Heil P and Scheich H (1986) Effects of unilateral and bilateral cochlea removal on 2-deoxyglucose patterns in the chick auditory system. *Journal of Comparative Neurology* **252**:279–301.
- Henson MM and Henson OW Jr (1991) Specializations for sharp tuning in the mustached bat: the tectorial membrane and spiral limbus. *Hearing Research* **56**:122–132.
- Herbert H, Aschoff A, and Ostwald J (1991) Topography of projections from the auditory cortex to the inferior colliculus in the rat. *Journal of Comparative Neurology* **304**:103–122.
- Hurley LM and Thompson AM (2001) Serotonergic innervation of the auditory brainstem of the Mexican free-tailed bat, *Tadarida brasiliensis*. *Journal of Comparative Neurology* **435**:78–88.
- Hyde PS and Knudsen EI (2002) The optic tectum controls visually guided adaptive plasticity in the owl's auditory space map. *Nature* **415**:73–76.

- Jarvis ED, Ribeiro S, da Silva ML, Ventura D, Vielliard J, and Mello CV (2000) Behaviourally driven gene expression reveals song nuclei in hummingbird brain. *Nature* **406**:628–632.
- Jeffress LA (1948) A place theory of sound localization. *Journal of Comparative and Physiological Psychology* **41**:359.
- Jen PH-S and Chen QC (1998) The effect of pulse repetition rate, pulse intensity, and bicuculline on the minimum threshold and latency of bat inferior collicular neurons. *Journal of Comparative Physiology A* **182**:455–465.
- Jen PH-S, Zhou X, and Wu CH (2001) Temporally patterned sound pulse trains affect intensity and frequency sensitivity of inferior collicular neurons of the big brown bat, *Eptesicus fuscus*. *Journal of Comparative Physiology A* **187**:605–616.
- Jen PH-S, Wu CH, Luan RH, and Zhou X (2002) GABAergic inhibition contributes to pulse repetition rate-dependent frequency selectivity in the inferior colliculus of the big brown bat, *Eptesicus fuscus*. *Brain Research* **948**:159–164.
- Ji W, Gao E, and Suga N (2001) Effects of acetylcholine and atropine on plasticity of central auditory neurons caused by conditioning in bats. *Journal of Neurophysiology* **86**:211–225.
- Kaiser A and Covey E (1997) 5-HT innervation of the auditory pathway in birds and bats. In: Syka J (ed). *Acoustical Signal Processing in the Central Auditory System*. Plenum Press, New York, pp. 71–78.
- Kanwal JS, Matsumura S, Ohlemiller K, and Suga N (1993) Analysis of acoustic elements and syntax in communication sounds emitted by mustached bats. *Journal of the Acoustical Society of America* **96**:1229–1254.
- Karten HJ (1967) The organization of the ascending auditory pathway in the pigeon (*Columba livia*). I. Diencephalic projections of the inferior colliculus (nucleus mesencephali lateralis, pars dorsalis). *Brain Research* **6**:409–427.
- Karten HJ (1968) The ascending auditory pathway in the pigeon (*Columba livia*) II. Telencephalic projections of the nucleus ovoidalis thalami. *Brain Research* **11**:134–153.
- Keller CH and Takahashi TT (1996) Binaural cross-correlation predicts the responses of neurons in the owl's auditory space map under conditions simulating summing localization. *Journal of Neuroscience* **16**:4300–4309.
- Keller CH and Takahashi TT (2000) Representation of temporal features of complex sounds by the discharge patterns of neurons in the owl's inferior colliculus. *Journal of Neurophysiology* **84**:2638–2650.
- Klug A, Park TJ, and Pollak GD (1995) Glycine and GABA influence binaural processing in the inferior colliculus of the mustache bat. *Journal of Neurophysiology* **74**:1701–1713.
- Knudsen EI (1982) Auditory and visual maps of space in the optic tectum of the owl. *Journal of Neuroscience* **9**:1177–1194.
- Knudsen EI (1983) Subdivisions of the inferior colliculus in the barn owl (*Tyto alba*). *Journal of Comparative Neurology* **218**:174–186.
- Knudsen EI (1999) Mechanisms of experience-dependent plasticity in the auditory localization pathway of the barn owl. *Journal of Comparative Physiology A* **185**:305–321.
- Knudsen EI (2002) Instructed learning in the auditory localization pathway of the barn owl. *Nature* **417**:322–328.
- Knudsen EI and Brainard MS (1991) Visual instruction of the neural map of auditory space in the developing optic tectum. *Science* **253**:85–87.

- Knudsen EI and Knudsen PF (1983) Space-mapped auditory projections from the inferior colliculus to the optic tectum in the barn owl (*Tyto alba*). *Journal of Comparative Neurology* **218**:187–196.
- Knudsen EI and Konishi M (1978) A neural map of auditory space in the owl. *Science* **200**:795–797.
- Knudsen EI, Blasdel GG, and Konishi M (1979) Sound localization by the barn owl (*Tyto alba*) measured with the search coil technique. *Journal of Comparative Physiology A* **133**:1–11.
- Knudsen EI, Esterly SD, Knudsen PF (1984) Monaural occlusion alters sound localization during a sensitive period in the barn owl. *Journal of Neurophysiology* **4**:1001–1011.
- Knudsen EI, Knudsen PF, and Masino T (1993) Parallel pathways mediating both sound localization and gaze control in the forebrain and midbrain of the barn owl. *Journal of Neuroscience* **13**:2837–2852.
- Koch U and Grothe B (1998) GABAergic and glycinergic inhibition sharpens tuning for frequency modulations in the inferior colliculus of the big brown bat. *Journal of Neurophysiology* **80**:71–82.
- Konishi M (1985) Birdsong: from behavior to neuron. *Annual Reviews of Neuroscience* **8**:125–170.
- Konishi M (2000) Study of sound localization by owls and its relevance to humans. *Comparative Biochemistry and Physiology A Molecular Integrative Physiology* **126**:459–469.
- Köppl C and Carr CE (2003) Computational diversity in the cochlear nucleus angularis of the barn owl. *Journal of Neurophysiology* **89**:2313–2329.
- Kössl M and Vater M (1985) The cochlear frequency map of the mustache bat, *Pteronotus parnellii*. *Journal of Comparative Physiology A* **157**:687–697.
- Kössl M, Mayer F, Frank G, Faulstich M, and Russell IJ (1999) Evolutionary adaptations of cochlear function in Jamaican mormoopid bats. *Journal of Comparative Physiology A* **185**:217–228.
- Kubke MF, Gauger B, Basu L, Wagner H, and Carr CE (1999) Development of calretinin immunoreactivity in the brainstem auditory nuclei of the barn owl (*Tyto alba*). *Journal of Comparative Neurology* **415**:189–203.
- Leibler LM (1975) *Monaural and Binaural Pathways in the Ascending Auditory System of the Pigeon*. Thesis. Massachusetts Institute of Technology, Cambridge.
- Leroy SA and Wenstrup JJ (2000) Spectral integration in the inferior colliculus of the mustached bat. *Journal of Neuroscience* **20**:8533–8541.
- Lesser HD, O'Neill WE, Frisina RD, and Emerson RC (1990) ON-OFF units in the mustached bat inferior colliculus are selective for transients resembling “acoustic glint” from fluttering insect targets. *Experimental Brain Research* **82**:137–148.
- Luksch H, Gauger B, and Wagner H (2000) A candidate pathway for a visual instructional signal to the barn owl’s auditory system. *Journal of Neuroscience* **20** RC70:1–4.
- Ma X and Suga N (2001) Corticofugal modulation of duration-tuned neurons in the midbrain auditory nucleus in bats. *Proceedings of the National Academy of Sciences of the United States of America* **98**:14060–14065.
- Manley GA, Köppl C, and Konishi M (1988) A neural map of interaural intensity difference in the brainstem of the barn owl. *Journal of Neuroscience* **8**:2665–2677.
- Marsh RA, Fuzessery ZM, Grose CD, and Wenstrup JJ (2002) Projection to the inferior colliculus from the basal nucleus of the amygdala. *Journal of Neuroscience* **22**:10449–10460.

- Mazer JA (1997) The integration of parallel processing streams in the sound localization system of the barn owl. *Computational Neuroscience: Trends in Research* **5**:735–739.
- Mazer JA (1998) How the owl resolves auditory coding ambiguity. *Proceedings of the National Academy of Sciences of the United States of America* **95**:10932–10937.
- Mittmann DH and Wenstrup JJ (1995) Combination-sensitive neurons in the inferior colliculus. *Hearing Research* **90**:185–191.
- Mogdans J and Knudsen EI (1994) Representation of interaural level difference in the VLVp, the first site of binaural comparison in the barn owl's auditory system. *Hearing Research* **74**:148–164.
- Moiseff A and Konishi M (1981) Neuronal and behavioral sensitivity to binaural time differences in the owl. *Journal of Neuroscience* **1**:40–48.
- Müller CM (1988) Distribution of GABAergic perikarya and terminals in the centers of the higher auditory pathway of the chicken. *Cell and Tissue Research* **252**:99–106.
- Neuweiler G (1990) Auditory adaptations for prey capture in echolocating bats. *Physiological Reviews* **70**:615–641.
- Neuweiler G (2000) *The Biology of Bats*. Oxford University Press, New York.
- Nieder B, Wagner H, and Luksch H (2003) Development of output connections from the inferior colliculus to the optic tectum in barn owls. *Journal of Comparative Neurology* **464**:511–524.
- Olsen JF and Suga N (1991) Combination-sensitive neurons in the medial geniculate body of the mustached bat: encoding of target range information. *Journal of Neurophysiology* **65**:1275–1296.
- O'Neill WE and Suga N (1979) Target range-sensitive neurons in the auditory cortex of the mustache bat. *Science* **203**:69–73.
- O'Neill WE and Suga N (1982) Encoding of target range and its representation in the auditory cortex of the mustached bat. *Journal of Neuroscience* **2**:17–31.
- Park TJ and Pollak GD (1993) GABA shapes sensitivity to interaural intensity disparities in the mustache bat's inferior colliculus: implications for encoding sound location. *Journal of Neuroscience* **13**:2050–2067.
- Park TJ and Pollak GD (1994) Azimuthal receptive fields are shaped by GABAergic inhibition in the inferior colliculus of the mustache bat. *Journal of Neurophysiology* **72**:1080–1102.
- Peña JL and Konishi M (2000) Cellular mechanisms for resolving phase ambiguity in the owl's inferior colliculus. *Proceedings of the National Academy of Sciences of the United States of America* **97**:11787–11792.
- Peña JL and Konishi M (2001) Auditory spatial receptive fields created by multiplication. *Science* **292**:249–252.
- Pinheiro AD, Wu M, and Jen PH-S (1991) Encoding repetition rate and duration in the inferior colliculus of the big brown bat, *Eptesicus fuscus*. *Journal of Comparative Physiology A* **169**:69–85.
- Pollak GD (1997) Roles of GABAergic inhibition for the binaural processing of multiple sound sources in the inferior colliculus. *Annals of Otolaryngology, Rhinology and Laryngology Supplement* **168**:44–54.
- Pollak GD and Casseday JH (1989) *The Neural Basis of Echolocation in Bats*. Springer-Verlag, Berlin.
- Pollak GD and Park TJ (1993) The effects of GABAergic inhibition on monaural response properties of neurons in the mustache bat's inferior colliculus. *Hearing Research* **65**:99–117.



- Pollak GD, Henson OW Jr, and Novick A (1972) Cochlear microphonic audiograms in the “pure tone” bat, *Chilonycteris parnellii parnellii*. *Science* **176**:66–68.
- Portfors CV and Wenstrup JJ (1999) Delay-tuned neurons in the inferior colliculus of the mustached bat: implications for analysis of target distance. *Journal of Neurophysiology* **82**:1326–1338.
- Portfors CV and Wenstrup JJ (2001) Topographical distribution of delay-tuned responses in the mustached bat inferior colliculus. *Hearing Research* **151**:95–105.
- Portfors CV and Wenstrup JJ (2002) Excitatory and facilitatory frequency response areas in the inferior colliculus of the mustached bat. *Hearing Research* **168**:131–138.
- Proctor L and Konishi M (1997) Representation of sound localization cues in the auditory thalamus of the barn owl. *Proceedings of the National Academy of Sciences of the United States of America* **94**:10421–10425.
- Puelles L, Robles C, and Martinez-de-la-Torre M (1994) New subdivision schema for the avian torus semicircularis: neurochemical maps in the chick. *Journal of Comparative Neurology* **340**:98–125.
- Reimer K (1987) Coding of sinusoidally amplitude modulated acoustic stimuli in the inferior colliculus of the rufous horseshoe bat, *Rhinolophus rouxi*. *Journal of Comparative Physiology A* **161**:305–313.
- Rosenberger MH, Fremouw T, Casseday JH, and Covey E (2003) Expression of the Kv1.1 ion channel subunit in the auditory system of the big brown bat, *Eptesicus fuscus*. *Journal of Comparative Neurology* **462**:101–120.
- Rübsamen R and Schäfer M (1990) Ontogenesis of auditory fovea representation in the inferior colliculus of the Sri Lankan rufous horseshoe bat, *Rhinolophus rouxi*. *Journal of Comparative Physiology A* **167**:757–769.
- Rübsamen R, Neuweiler G, and Sripathy K (1988) Comparative collicular tonotopy in two bat species adapted to movement detection, *Hipposideros speoris* and *Megaderma lyra*. *Journal of Comparative Physiology A* **163**:271–285.
- Scheich H, Langner G, and Koch R (1977) Coding of narrow-band and wide-band vocalizations in the auditory midbrain nucleus (MLD) of the guinea fowl (*Numida meleagris*). *Journal of Comparative Physiology A* **117**:245–266.
- Schuller G (1979) Coding of small sinusoidal frequency and amplitude modulations in the inferior colliculus of ‘CF-FM’ bat, *Rhinolophus ferrumequinum*. *Experimental Brain Research* **34**:117–132.
- Schuller G, Covey E, and Casseday JH (1991) Auditory pontine grey: connections and response properties in the horseshoe bat. *European Journal of Neuroscience* **3**:648–662.
- Schwarz DWF, Schwarz IE, and Dezso A (1995) Retrograde labelling of auditory brainstem neurons following tritiated glycine injection into the inferior colliculus of the chicken, *Gallus domesticus*. *Journal of Otolaryngology* **24**:31–41.
- Senatorov VV and Hu B (2002) Extracortical descending projections to the rat inferior colliculus. *Neuroscience* **115**:243–250.
- Simmons JA (1989) A view of the world through the bat’s ear: the formation of acoustic images in echolocation. *Cognition* **33**:155–199.
- Spezio ML and Takahashi TT (2003) Frequency-specific interaural level difference tuning predicts spatial response patterns of space-specific neurons in the barn owl inferior colliculus. *Journal of Neuroscience* **23**:4677–4688.
- Suga N (1968) Analysis of frequency modulated and complex sounds by single auditory neurones of bats. *Journal of Physiology (London)* **198**:51–80.
- Suga N (1969) Classification of inferior colliculus neurons of bats in terms of responses to pure tones, FM sounds, and noise bursts. *Journal of Physiology (London)* **200**:555–574.

- Suga N and Ma X (2003) Multiparametric corticofugal modulation and plasticity in the auditory system. *Nature Reviews Neuroscience* **4**:783–794.
- Suga N, Simmons JA, and Jen PH-S (1975) Peripheral specialization for fine analysis of Doppler-shifted echoes in the auditory system of the “CF-FM” bat, *Pteronotus parnellii*. *Journal of Experimental Biology* **63**:161–192.
- Suga N, O’Neill WE, and Manabe T (1979) Harmonic-sensitive neurons in the auditory cortex of the mustache bat. *Science*. **203**:270–274.
- Suga N, Xiao Z, Ma X, and Ji W (2002) Plasticity and corticofugal modulation for hearing in adult animals. *Neuron* **36**:9–18.
- Sullivan WE (1982) Possible neural mechanisms of target distance encoding in auditory system of the echolocating bat, *Myotis lucifugus*. *Journal of Neurophysiology* **48**:1033–1047.
- Takahashi TT and Keller CH (1992) Commissural connections mediate inhibition for the computation of interaural level difference in the barn owl. *Journal of Comparative Physiology A* **170**:161–169.
- Takahashi TT and Konishi M (1986) Selectivity for interaural time difference in the owl’s midbrain. *Journal of Neuroscience* **6**:3413–3422.
- Takahashi TT and Konishi M (1988a) Projections of nucleus angularis and nucleus laminaris to the lateral lemniscal nuclear complex of the barn owl. *Journal of Comparative Neurology* **274**:212–238.
- Takahashi TT and Konishi M (1988b) Projections of the cochlear nuclei and nucleus laminaris to the inferior colliculus of the barn owl. *Journal of Comparative Neurology* **274**:190–211.
- Takahashi TT, Barberini CL, and Keller CH (1995) An anatomical substrate for the inhibitory gradient in the VLVp of the owl. *Journal of Comparative Neurology* **358**:294–304.
- Takahashi TT, Moiseff A, and Konishi M (1984) Time and intensity cues are processed independently in the auditory system of the owl. *Journal of Neuroscience* **4**:1781–1786.
- Takahashi TT, Carr CE, Brecha N, and Konishi M (1987) Calcium binding protein-like immunoreactivity labels the terminal field of nucleus laminaris of the barn owl. *Journal of Neuroscience* **7**:1843–1856.
- Takahashi TT, Wagner H, and Konishi M (1989) Role of commissural projections in the representation of bilateral auditory space in the barn owl’s inferior colliculus. *Journal of Comparative Neurology* **281**:545–554.
- Takahashi TT, Bala AD, Spitzer MW, Euston DR, Spezio ML, and Keller CH (2003) The synthesis and use of the owl’s auditory space map. *Biological Cybernetics* **89**:378–387.
- Theunissen FE, Sen K, and Doupe AJ (2000) Spectral-temporal receptive fields of non-linear auditory neurons obtained using natural sounds. *Journal of Neuroscience* **20**:2315–2331.
- Tsutchitani C (1977) Functional organization of lateral cell groups of cat superior olivary complex. *Journal of Neurophysiology* **40**:296–318.
- Vater M and Braun K (1994) Parvalbumin, calbindin D-28k, and calretinin immunoreactivity in the ascending auditory pathway of horseshoe bats. *Journal of Comparative Neurology* **341**:534–558.
- Vater M, Feng AS, and Betz M (1985) An HRP-study of the frequency-place map of the horseshoe bat cochlea: morphological correlates of the sharp tuning to a narrow frequency band. *Journal of Comparative Physiology A* **157**:671–686.
- Vater M, Habbicht H, Kössl M, and Grothe B (1992a) The functional role of GABA and glycine in monaural and binaural processing in the inferior colliculus of horseshoe bats. *Journal of Comparative Physiology A* **171**:541–553.

- Vater M, Kössl M, and Horn AKE (1992b) GAD- and GABA-immunoreactivity in the ascending auditory pathway of horseshoe and mustached bats. *Journal of Comparative Neurology* **325**:183–206.
- Vater M, Casseday JH, and Covey E (1995) Convergence and divergence of ascending binaural and monaural pathways from the superior olives of the mustached bat. *Journal of Comparative Neurology* **351**:632–646.
- Vater M, Covey E, and Casseday JH (1997) The columnar region of the ventral nucleus of the lateral lemniscus in the big brown bat: synaptic arrangements and structural correlates of feedforward inhibitory function. *Cell and Tissue Research* **289**:223–233.
- Volman SF and Konishi M (1989) Spatial selectivity and binaural responses in the inferior colliculus of the great horned owl. *Journal of Neuroscience* **9**:3083–3096.
- Wagner H (1993) Sound-localization deficits induced by lesions in the barn owl's auditory space map. *Journal of Neuroscience* **13**:371–386.
- Wagner H, Takahashi T, and Konishi M (1987) Representation of interaural time difference in the central nucleus of the barn owl's inferior colliculus. *Journal of Neuroscience* **7**:3105–3116.
- Wagner H, Mazer JA, and von Campenhausen M (2002) Response properties of neurons in the core of the central nucleus of the inferior colliculus of the barn owl. *European Journal of Neuroscience* **15**:1343–1352.
- Wagner H, Günturkun O, and Nieder B (2003) Anatomical markers for the subdivisions of the barn owl's inferior-collicular complex and adjacent peri- and subventricular structures. *Journal of Comparative Neurology* **465**:145–159.
- Wenstrup JJ and Leroy SA (2001) Spectral integration in the inferior colliculus: role of glycinergic inhibition in response facilitation. *Journal of Neuroscience* **21** RC124:1–6.
- Wenstrup JJ, Ross LS, and Pollak GD (1986) Binaural response organization within a frequency-band representation of the inferior colliculus: implications for sound localization. *Journal of Neuroscience* **6**:962–973.
- Wenstrup JJ, Larue DT, and Winer JA (1994) Projections of physiologically defined subdivisions of the inferior colliculus in the mustached bat: targets in the medial geniculate body and extrathalamic nuclei. *Journal of Comparative Neurology* **346**:207–236.
- Wild JM (1987) Nuclei of the lateral lemniscus project directly to the thalamic auditory nuclei in the pigeon. *Brain Research* **408**:303–307.
- Wild JM (1993) Descending projections of the songbird nucleus robustus archistriatalis. *Journal of Comparative Neurology* **338**:225–241.
- Wild JM (1995) Convergence of somatosensory and auditory projections in the avian torus semicircularis, including the central auditory nucleus. *Journal of Comparative Neurology* **358**:465–486.
- Wild JM, Karten HJ, and Frost BJ (1993) Connections of the auditory forebrain in the pigeon (*Columba livia*). *Journal of Comparative Neurology* **337**:32–62.
- Wild JM, Li D, and Eagleton C (1997) Projections of the dorsomedial nucleus of the intercollicular complex (DM) in relation to respiratory-vocal nuclei in the brainstem of pigeon (*Columba livia*) and zebra finch (*Taeniopygia guttata*). *Journal of Comparative Neurology* **377**:392–413.
- Wild JM, Kubke MF, and Carr CE (2001) Tonotopic and somatotopic representation in the nucleus basalis of the barn owl, *Tyto alba*. *Brain, Behavior and Evolution* **57**:39–62.
- Winer JA, Larue DT, and Pollak GD (1995) GABA and glycine in the central auditory system of the mustache bat: structural substrates for inhibitory neuronal organization. *Journal of Comparative Neurology* **355**:317–353.

- Winer JA, Larue DT, Diehl JJ, and Hefti BJ (1998) Auditory cortical projections to the cat inferior colliculus. *Journal of Comparative Neurology* **400**:147–174.
- Winer JA, Chernock ML, Larue DT, and Cheung SW (2002) Descending projections to the inferior colliculus from the posterior thalamus and the auditory cortex in rat, cat, and monkey. *Hearing Research* **168**:181–195.
- Woolley SMN and Casseday JH (2004) Response properties of single neurons in the zebra finch auditory midbrain: response patterns, frequency coding, intensity coding and spike latencies. *Journal of Neurophysiology* **91**:136–151.
- Wotton JM, Haresign T, and Simmons JA (1995) Spatially dependent acoustic cues generated by the external ear of the big brown bat, *Eptesicus fuscus*. *Journal of the Acoustical Society of America* **98**:1423–1445.
- Wu MI and Jen PH-S (1996) Temporally patterned pulse trains affect directional sensitivity of inferior collicular neurons of the big brown bat, *Eptesicus fuscus*. *Journal of Comparative Physiology A* **179**:385–393.
- Yan J and Suga N (1996) Corticofugal modulation of time-domain processing of biosonar information in bats. *Science* **273**:1100–1103.
- Yan J and Suga N (1999) Corticofugal amplification of facilitative auditory responses of subcortical combination-sensitive neurons in the mustached bat. *Journal of Neurophysiology* **81**:817–824.
- Yang L, Pollak GD, and Resler C (1992) GABAergic circuits sharpen tuning curves and modify response properties in the mustache bat inferior colliculus. *Journal of Neurophysiology* **68**:1760–1774.
- Zettel ML, Carr CE, and O'Neill WE (1991) Calbindin-like immunoreactivity in the central auditory system of the mustached bat, *Pteronotus parnellii*. *Journal of Comparative Neurology* **313**:1–16.
- Zhang Y, Suga N, and Yan J (1997) Corticofugal modulation of frequency processing in bat auditory system. *Nature* **387**:900–903.
- Zheng W and Knudsen EI (1999) Functional selection of adaptive auditory space map by GABA<sub>A</sub>-mediated inhibition. *Science* **284**:962–965.
- Zhou X and Jen PH-S (2001) The effect of sound intensity on duration-tuning characteristics of bat inferior collicular neurons. *Journal of Comparative Physiology A* **187**:63–73.
- Zook JM and Casseday JH (1982) Cytoarchitecture of auditory system in lower brainstem of the mustache bat, *Pteronotus parnellii*. *Journal of Comparative Neurology* **207**:1–13.
- Zook JM and Casseday JH (1985) Projections from the cochlear nuclei of the mustache bat, *Pteronotus parnellii*. *Journal of Comparative Neurology* **237**:307–324.
- Zook JM and Casseday JH (1987) Convergence of ascending pathways at the inferior colliculus of the mustache bat, *Pteronotus parnellii*. *Journal of Comparative Neurology* **261**:347–361.
- Zook JM and Leake PA (1989) Connections and frequency representation in the auditory brainstem of the mustache bat, *Pteronotus parnellii*. *Journal of Comparative Neurology* **290**:243–261.
- Zook JM, Winer JA, Pollak GD, and Bodenhamer RD (1985) Topology of the central nucleus of the mustache bat's inferior colliculus: correlation of single unit properties and neuronal architecture. *Journal of Comparative Neurology* **231**:530–546.

# Chapter 18

## Development of Auditory Afferents to the Central Nucleus of the Inferior Colliculus

JUDY K. BRUNSO-BECHTOLD AND CRAIG K. HENKEL

### 1. INTRODUCTION

The inferior colliculus (IC) is a requisite site of convergent input for much of the ascending auditory system (Fig. 18.1). Although some ascending fibers bypass this structure, the IC is certainly a synaptic target for the vast majority of parallel auditory pathways ascending from the brain stem (see Chapters 2 and 4). In addition, the IC receives a heavy descending projection from cerebral cortex (see Chapter 8). Consequently, during development an immense population of actively growing axons converges on the IC. Of necessity, this heterogeneous and dynamic input must rapidly and efficiently develop local intrinsic networks to integrate and preserve ascending monaural and binaural frequency and timing cues extracted during neural processing in brain stem nuclei.

The IC is composed of a large central nucleus (CNIC) surrounded by dorsal (DCIC) and external (ECIC) cortical nuclei. The CNIC receives crossed and uncrossed ascending pathways (Fig. 18.1A), each converging in cochleotopic order within a series of fibrodendritic layers. A hallmark of the structural organization of the CNIC is the mosaic of bands and patches of afferent input from different sources projecting to somata and dendrites oriented parallel to the layers (Fig. 18.1B, C) (Geniec and Morest 1971; Rockel and Jones 1973; Fitzpatrick 1975; Morest and Oliver 1984; Oliver and Morest 1984; Faye-Lund and Osen 1985; Zook et al. 1985; Shneiderman and Henkel 1987; Oliver and Shneiderman 1989; Malmierca et al. 1993; Oliver 2000). How this elaborate and precise organization is created from a dynamic population of rapidly growing neurites is unknown but it must be governed by some of the classic ontogenetic principles prevailing elsewhere in the neuraxis (Fig. 18.2). This population of neurites includes axons arriving from a wide range of cell types in various nuclei on both sides of the brain, as well as intrinsic axons and developing dendrites. Once the basic axonal organization is established within the CNIC, their local distribution is refined and they begin to express mature features such as neurotransmitters and their receptors and transporters as well as calcium binding proteins. The development of these features appears to be influenced significantly by neural activity, often demonstrated by the effects of hearing loss.




Figure 18.1. Schematic overview of the architecture and afferent projections to the CNIC (simplified to include only ascending projections from principal nuclei). (A) Locations of the CNIC, DCIC and ECIC are shown at the *upper left* in an outline of a frontal section. *Diagonal lines* depict the orientation of fibrodendritic layers characteristic of the CNIC. The major excitatory (*open circles*) and inhibitory (*closed circles*) inputs to the layers are shown. (B) A portion of CNIC is enlarged to show four layers. Afferents from multiple sources converge in cochleotopic order in layers, forming distinct synaptic compartments among functional sets of inputs. (C) A single disc-shaped cell is illustrated in the expanded view of a layer. Cells with flattened dendritic fields are aligned with layered afferent inputs to form input–output circuits that preserve frequency processing. *CN*, cochlear nuclei; *CNIC*, central nucleus of inferior colliculus; *DCIC*, dorsal cortex of inferior colliculus; *DNLL*, dorsal nucleus of the lateral lemniscus; *ECIC*, external cortex of the inferior colliculus; *LSO*, lateral superior olivary nucleus; *MSO*, medial superior olivary nucleus; *VNLL*, ventral nucleus of the lateral lemniscus.

Although the precise mechanisms by which ear removal or hearing loss influence afferent organization and axonal identity remain to be determined, there is an increased awareness of the trophic impact of activity-driven neural events.

## 2. DEVELOPMENT OF THE CENTRAL NUCLEUS

### 2.1. AXONS AND SYNAPSES

Cells within the CNIC are born very early in development (Fig. 18.2), beginning at embryonic day 14 (E14) in rat, 3 weeks before hearing begins (Altman and




Figure 18.2. Summary of the time course of the developmental events in the IC relative to each other and in relation to onset of hearing.

Bayer 1981). By E13 some cochlear nucleus pioneer axons have extended along the perimeter of the hindbrain (Niblock et al. 1995), reaching the IC by E17 or E18 in rat (Kandler and Friauf 1993). Axons from the dorsal nucleus of the lateral lemniscus (DNLL) extend to the contralateral IC by E19 in ferret (Gabriele et al. 2000a). The medial superior olive (MSO) projection to the IC is present at birth in rat and gradually increases until adulthood (Okoyama et al. 1995). In carnivores as well as in the rat, there is a rich distribution of axons within the IC well before auditory experience (González-Hernández et al. 1989; Kandler and Friauf 1993; Gabriele et al. 2000a; Keiger et al. 2003). Initially, projections from DNLL distribute diffusely in the CNIC as linear axons with few side branches or protrusions but oriented parallel to the frequency layers (Fig. 18.3A). Second, still before the onset of hearing, axons from DNLL as well as those from lateral superior olive (LSO) begin to form a periodic distribution of fiber-rich bands or layers with intervening axon-sparse interband spaces or sublayers (Fig. 18.3B) (Gabriele et al. 2000a). Similarly the LSO projection in ferret distributes diffusely in the IC in the first postnatal week (Fig. 18.4A, B). One week later, and two weeks before hearing begins, a banded segregation of this projection is apparent (Fig. 18.4C, D) (Keiger et al. 2003).

In the only two afferent systems in which band formation has been studied, the DNLL and the LSO projections to IC, the projections are bilateral. Both the DNLL and LSO projections are banded within the adult IC (Kudo 1981; Shneiderman and Henkel 1987; Shneiderman et al. 1988; Oliver 2000). This pattern is in contrast to that of other projections such as those from the MSO and the ventral nucleus of the lateral lemniscus, which are almost exclusively ipsilateral and have little banding within the CNIC (Henkel and Spangler 1983; Whitley




Figure 18.3. A working model of steps in the developmental refinement of layered fibrodendritic circuits in the CNIC. **(A)** Afferent and dendritic elements of IC circuits initially establish their approximate distribution (*hatched area*) and then subsequently are pruned (*arrow*) to match cochleotopic position within fibrodendritic layers. **(B)** Afferent inputs from different ascending channels initially converge on the same layer and then segregate (*arrow*) in the vertical dimension to form sublayers (*black vs. white*). **(C)** Functional sets of inputs segregate (*arrow*) to form multiple synaptic domains (*black, overlapping, and white zones*) within fibrodendritic layers or sublayers.

and Henkel 1984; Oliver 2000). These observations are consistent with the possibility that competition between converging ipsilateral and contralateral input drives the development of banded projection patterns from the DNLL and the LSO. Although there is a banded pattern of input to CNIC from the dorsal cochlear nucleus (DCN) (Oliver 1984; Oliver et al. 1997), which also projects bilaterally, the contralateral component is significantly larger. This asymmetry of projection raises the issue of whether a projection from another source might overlap with that from ipsilateral cochlear nucleus (CN), presumably firing coincidentally with it and thus participating in a competition for synaptic space potentially occupied by fibers from the contralateral CN.

Rearrangement in the laterality of IC afferents prior to hearing onset is evident in the gradual decrease in the proportion of LSO cells projecting to the contralateral IC in ferret during the first postnatal month (Henkel and Brunso-Bechtold 1993). Initially the DNLL and LSO projections distribute diffusely across the width of the frequency layers and extend broadly along the length of the frequency layers (Fig. 18.3C: left). However, once the segregation of axons into bands and interband spaces has begun, these projections are confined to the length of the frequency layers (Fig. 18.3C: right). In the adult, projections from different sources such as DCN and LSO occupy distinct domains within the layers, consistent with a role for competition among afferent sources in the refinement of intralaminar synaptic domains. Although refinement of the banded






Figure 18.4. DiI labeling of projections from the LSO to the IC in ferret prior to hearing onset at the end of the first postnatal month. DiI-coated cactus needles that were placed in the LSO in lightly fixed ferret brains at P7 (A) and P14 (C) labeled axons that enter the CNIC from the lateral lemniscus and extend parallel to the isofrequency planes. Axons at P7 (B) distribute throughout the ventrolateral portion of the CNIC with early periodicity emerging. By P14 (D), LSO axons within the CNIC form bands (arrows). Dorsal is toward the *top* and medial is to the *right*. CNIC, Central nucleus of the inferior colliculus; DCIC, dorsal cortex of the inferior colliculus; ECIC, external cortex of the inferior colliculus; LL, lateral lemniscus; LSO, lateral superior olive; MNTB, medial nucleus of the trapezoid body; MSO, medial superior olive. The scale bars represent 150  $\mu\text{m}$ .

distribution of ascending auditory projections undoubtedly continues after ear opening, the basic banded pattern of DNLL and LSO input to the CNIC is evident several days before hearing onset in the rat (Gabriele et al. 2000a) and several weeks before hearing onset in the ferret (Keiger et al. 2003).

As input segregation along afferent bands is taking place, numerous varicosities and side branches emerge along the preterminal axons (Gabriele et al. 2000a). Whether these axonal events are accompanied by a synchronous formation of synaptic connections is currently unclear as little information is available on synaptogenesis within the IC. In fact, the only quantitative studies of synapse development pertain to marsupials for which data are not available about the development or absence of a banded distribution pattern (see Chapter 22, Fig. 22.2B). In the Northern quoll, a marsupial cat, synapses first can be detected

in the IC nearly 2 months prior to the onset of hearing. The ratio of synapses to neurons gradually increases until several days before hearing onset, and then rapidly accelerates to adult levels about 10 days later (Aitkin et al. 1996). Similarly, in the short-tail Brazilian opossum, synapses are apparent at the first age studied, about 1 week before hearing begins, and do not quantitatively change until hearing onset, after which synaptogenesis rapidly accelerates (Aitkin et al. 1997). In the rat, the available information suggests some synapses are present before hearing onset (Pysh 1969).

In the ferret, evidence from our laboratory indicates that synapses in the IC neuropil are present during the first postnatal week and increase steadily in number throughout the month before hearing begins. At postnatal day 4 (P4) IC synapses are distributed sparsely through the neuropil, but are small and immature, without glial encapsulation, and myelin is not present (Fig. 18.5A). In comparison, P28 synapses (Fig. 18.5B) are larger and more likely to be perforated, and have multiple targets. Moreover, myelin is now apparent and glial encapsulation of dendrites and synapses is in progress. Quantitative studies will be required to relate synaptogenesis to the beginning of afferent band formation within the ferret CNIC (Keiger et al. 2003). However, ferret MSO synapses are apparent in the neuropil during the first 2 weeks after birth with rapid somatic synaptogenesis beginning during the third postnatal week and a relatively mature synaptic organization present by ear opening (Brunso-Bechtold et al. 1992). Although quantitative data on synaptogenesis within the CNIC are not




Figure 18.5. Ultrastructure of ferret IC prior to hearing onset. At P4 (A), simple synapses (*arrowheads*) are sparsely distributed throughout the CNIC neuropil. Growth cones (*arrow*) are present, myelin has not yet begun to form, and glial processes only rarely extend along the surfaces of somata and dendrites. At P28 (B), just before hearing begins, synapses (*arrowheads*) are more complex and glial encapsulation (*arrows*) of synapses and dendrites is evident. Myelin formation also is underway. The scale bars represent 5  $\mu\text{m}$  (A) and 1  $\mu\text{m}$  (B).

currently available for any mammal, it is reasonable to extrapolate that a basal level of synapses is present during the segregation of afferent input and that increased synaptogenesis coincides with the refinement of afferent projections.

## 2.2. *DENDRITES*

Lamination of dendritic arbors as well as lemniscal axons is a characteristic feature of the CNIC (Fig. 18.1). This arrangement of afferent inputs and post-synaptic cells underlies the orderly coding of frequency in IC layers, with low frequencies represented dorsolaterally and higher frequencies more ventromedially. In rat Golgi material, dendritic features distinguishing disc-shaped and multipolar (stellate) cells in the CNIC and multipolar and giant cells in the ECIC and DCIC are present at birth (González-Hernández et al. 1989). Quantitative analysis of CNIC cells reveals a pronounced flattening of dendritic trees in the plane of these layers after the onset of hearing in kittens and rats (Meininger and Baudrimont 1981; Dardennes et al. 1984). Dendritic trunks parallel to the layers elongate by adding secondary processes while those extending beyond the layer are pruned. Dendritic spines, present on various CNIC cell types, gradually increase in density until one month of age then decline to adult values (Dardennes et al. 1984). Dense layers of lemniscal axons are present at birth in kittens (González-Hernández et al. 1989) and banding of afferent projections in rat (Gabrielle et al. 2000a) and ferret (Keiger et al. 2003) emerges well before hearing begins. Thus, the final dendritic maturation in the CNIC, including growth, branching, and spine elimination that occurs after hearing onset, may depend on functional maturation of afferent input (Fig. 18.2).

## 2.3. *FUNCTIONAL CORRELATES*

Almost as soon as responses to sound can be recorded in the IC, response properties such as frequency tuning, binaural response patterns, and response types emerge and resemble those in adult animals (Aitkin and Moore 1975; Aitkin and Reynolds 1975; Moore and Irvine 1980, 1981a,b; Blatchley and Brugge 1990; Romand and Ehret 1990). These reflect the early establishment and postnatal maturation described for the presynaptic and postsynaptic structural organization in the IC. Although higher thresholds, lower discharge rates, and broader tuning are evident initially, and in part are secondary to protracted development of the ear, further maturation of threshold and discharge rates undoubtedly depends on subsequent postnatal events that affect the balance of excitatory and inhibitory synaptic circuits and their respective receptor composition. Moreover, refinement of frequency tuning may involve interaction and matching of dendritic compartments with afferent input.

In contrast to the morphological sequence of cochlear maturation in which development of the high-frequency, basal cochlea precedes that of the low-frequency, apical portion, functional studies suggest that low-frequency hearing develops first. It has been proposed that this apparent paradox reflects a devel-

opmental shift in cochlear best frequency, and such shifts in sensitivity of hair cells from low- to high-frequency sounds are reported in many species (Lippe and Rubel 1983; Harris and Dallos 1984; Rubel et al. 1984; Echteler et al. 1989; Sanes et al. 1989; Friauf 1992; Schweitzer et al. 1996; Romand 1997; Mills and Rubel 1998). Evidence in the gerbil supports a developmental shift in the tonotopic map within the LSO and IC in a manner consistent with such frequency shifts in the developing cochlea (Ryan and Woolf 1988; Sanes et al. 1989). However, at a comparable time point in the rat, high- and low-frequency neural activity demonstrated by *c-fos* immunoreactivity is present in the IC as well as in brain stem auditory nuclei (Friauf 1992). No subsequent shift in the position of frequency responsiveness was observed, suggesting that any such frequency shift takes place before ear opening. Reconciling this observation with reports of frequency shifts in the rat cochlea taking place near hearing onset (Hyson and Rudy 1987) or even later (Müller 1991) remains an open issue.

### 2.3.1. Calcium-Binding Proteins

Calcium-binding proteins in neurons have multiple roles and their presence has been correlated with different functional properties. In immature neurons, calcium-binding proteins may identify the site of important calcium-dependent mechanisms and serve as simple markers of various components of developing auditory circuits (Baimbridge et al. 1992; Andressen et al. 1993). In adult rat, various calcium-binding proteins are distributed differentially in the IC and have unique developmental patterns (Friauf 1994; Lohmann and Friauf 1996). For instance, parvalbumin-immunoreactive neurons are distributed throughout the IC subnuclei in adult rat, but are not present until P8. Parvalbumin-immunopositive cells increase from P8 until P28 when adult levels are reached (Lohman and Friauf 1996). In contrast, calretinin- and calbindin-immunoreactive neurons are present in the IC around birth. Immunostaining for these calcium-binding proteins, as opposed to parvalbumin, declines thereafter and is absent shortly after hearing onset in the CNIC, but persists in other subnuclei (Friauf 1994; Lohman and Friauf 1996).

In addition to the distribution of calcium-binding proteins in neuronal somata within the IC, calretinin and calbindin concentrate selectively in axons afferent to the CNIC (McHaffie et al. 2000; Fuentes-Santamaria et al. 2003). The early distribution pattern of calbindin-immunoreactive endings in kitten (McHaffie et al. 2000) resembles the alternating bands described for lemniscal inputs to the CNIC (Fig. 18.6A). This pattern gradually becomes fainter but more sharply defined during the 2 postnatal months as calbindin-immunoreactive endings are restricted to the central region of the IC layers (Fig. 18.6B), suggesting that the segregation of those projections involves calcium-mediated mechanisms (Fig. 18.2).

### 2.3.2. Neurotransmitters

Glutamate, glycine, and  $\gamma$ -aminobutyric acid (GABA) are the dominant neurotransmitter systems in the brain stem auditory nuclei afferent to the IC. Iono-




Figure 18.6. Distribution of calbindin immunoreactivity in cat IC during postnatal development. Calbindin immunoreactivity in the CNIC at P1 (**A**) is localized densely in axons forming bands (*arrows*) oriented parallel to the isofrequency planes. By P42 (**B**), the bands of calbindin immunoreactive axons are sharper but less intense. Dorsal is toward the *top* and medial is to the *right* in (**A**) and to the *left* in (**B**). *CNIC*, Central nucleus of the inferior colliculus; *ECIC*, external cortex of the inferior colliculus. The scale bars represent 100  $\mu\text{m}$ .

tropic receptors for these amino acid neurotransmitters have in common that they are heteromeric complexes comprised of different subunits surrounding an ion channel. The specific receptor subunit composition modulates channel kinetics. Receptors for these neurotransmitters have complex individual subunit profiles and distributions in the IC (Sato et al. 2000). These receptor complexes undergo postnatal modification with slight variations in peak expression times of the various components (Fig. 18.2).

The major class of excitatory receptors in the mature IC is the ionotropic glutamate receptors. Both  $\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazole propionic acid (AMPA) and *N*-methyl-D-aspartate (NMDA) glutamate receptors are present. The functional complements of AMPA receptor subunits (GluR1-4 or A-D) and two of their splice variants (flip or “a” and flop or “b”) as well as the NMDA receptor subunits (NR1 and NR2) in the CNIC are developmentally regulated. There is a decrease in the immature, flip isoforms of the AMPA subunits in the IC by hearing onset while the mature, flop isoforms are expressed more highly at or after hearing onset. In contrast, NMDA subunit expression in the CNIC is moderate and relatively constant in postnatal and young adult rats (Caicedo and Eybalin 1999). For the AMPA receptors, there are significantly higher mRNA levels for the immature flip isoform of the GluR2 subunit in the IC compared to LSO and CN (Schmid et al. 2001). Higher GluR2 subunit levels in mature receptors may be related to slower kinetics and lower permeability to calcium ions of these neurons (Hollman et al. 1991; Hume et al. 1991). This calcium permeability of IC glutamate receptors appears to be developmentally regulated and is highest in postnatal rat around hearing onset (Caicedo et al. 1998).

Both GABA and glycine are prominent IC inhibitory neurotransmitters. The composition of GABA<sub>A</sub> and glycine receptor subunits changes postnatally, with the expression of mRNA for immature forms of both receptor complexes de-

clining by hearing onset (Laurie et al 1992; Kraushaar and Backus 2002). Although it is often the case that the glycine  $\alpha$ -2 receptor subunit is the neonatal form and  $\alpha$ -1 the adult form, the  $\alpha$ -2 message is weakly expressed at birth in the rat IC and remains low until P21. In contrast, the  $\alpha$ -1 subunits of the glycine receptor are absent at birth, expressed weakly at about P8, increase from P12-P20, and reach adult distribution by P28 (Friauf et al. 1997; Piechotta et al. 2001). As the glycine transporter protein GLYT2 is highly expressed in the CNIC at birth, before the mature  $\alpha$ -1 subunit appears, and declines to lower levels by P22, this transporter may participate in synapse maturation (Friauf et al. 1999).

Although its function has not been fully explored, a substantial cholinergic input to the CNIC appears postnatally and may be of basal forebrain origin (Gao and Suga 2000; Ma and Suga 2003). This input is mediated primarily by a subclass of nicotinic cholinergic receptors (nAChR) (Morley and Happe 2000). nAChRs in the IC are concentrated on presynaptic endings and may have modulatory effects on neurotransmitter release. In development, nAChRs have important roles in activity-dependent plasticity of synaptic connections (Radcliffe and Dani 1998; Dani 2001; McGehee 2002). Perhaps the developmental regulation of nAChRs in IC is similar to that in auditory cortex where they peak late in postnatal development (Aramakis et al. 2000) during the plastic period when auditory experience shapes functional and structural refinement of synaptic circuits.

Maturation of the neurotransmitter receptor subunit profiles, regardless of type, is a dynamic process and may reflect the organization of heterogeneous inputs to the CNIC. Moreover, cotransmitters such as somatostatin and leu-enkephalin, which modulate activity, are expressed transiently in the IC (Sekitani et al. 1990; Kungel and Friauf 1995; Thoss et al. 1996). The maturation of these profiles occurs well after initial circuits are formed and even after banded patterns in IC emerge (Fig. 18.2). Thus, the early functional role of receptor subunits may be to participate in the formation of afferent bands and early emergence of synaptic domains (see Chapter 2).

### 3. ROLE OF ACTIVITY IN INFERIOR COLLICULUS DEVELOPMENT

Activity-mediated mechanisms that shape IC circuits may depend on either spontaneous activity in the ascending auditory pathways before hearing onset or patterned sound-evoked activity during early hearing (Fig. 18.2). Ascending projections reach the auditory midbrain and elaborate diffuse axonal arbors near their ultimate targets well before hearing onset and undergo subsequent adjustment. The electrophysiological and biochemical environment plays an essential role in influencing the formation of a mature afferent projection pattern in the CNIC. The principal manipulations used to assay activity-specific effects include

cochlear removal and antibiotic infusion, which physically interrupt the link from sensory receptor to auditory targets and cause a sensorineural hearing loss. In contrast, ossicle removal and ear plugging leave the auditory pathway intact, but reduce the afferent signal, causing a conductive hearing loss. Here, we consider only studies that affect the formation of afferent bands within the developing IC.

### *3.1. EFFECT OF EAR REMOVAL ON INFERIOR COLLICULUS AFFERENTS*

Considerable information is available on the effects of ear removal on projections to, and morphometry of, mammalian CN, LSO, and MSO neurons (Webster and Webster 1977; Trune 1982a,b; Moore and Kowalchuk 1988; Moore 1990; Kitzes et al. 1995; Russell and Moore 1995; Tierney et al. 1997; Hardie and Shepherd 1999). The impact of sensorineural hearing loss on these structures is most dramatic when it occurs prior to hearing onset. This suggests that the effect of early ear removal on the input to the IC extends beyond a change in levels or patterns of activity to physically affect the neuronal populations contributing the axonal input. Thus, unilateral cochlear ablations made before ear opening in ferret significantly decrease cochlear nucleus volume ipsilateral to the lesion in adults and increase the projection from the contralateral CN to the IC on the same side (Moore and Kowalchuk 1988). However, conductive hearing loss does not change CN volume, yet still increases the number of CN neurons contralateral to the lesion that projected to the IC on the same side (Moore et al. 1989). This demonstrates forcefully that early changes in auditory activity can affect the projection pattern to the IC independent of cell loss.

In the gerbil, as in the ferret, early cochlear ablation increases the projection originating in the CN contralateral to the lesion and terminating in the IC on the same side (Nordeen et al. 1981). Moreover, the distribution of the ipsilateral CN projection within the IC is expanded (Moore and Kitzes 1985). In contrast, bilateral cochlear ablations at ear opening cause no significant differences in the projection from the CN to the IC (Moore 1990). Taken together, the effects of unilateral vs. bilateral cochlear ablation suggest the importance of competition between event-driven auditory activity from both ears in establishing the projection of ascending auditory afferents to the IC.

### *3.2. EFFECT OF EAR REMOVAL ON STRUCTURAL FEATURES OF THE INFERIOR COLLICULUS*

Few studies have evaluated structural changes within the IC ensuing from cochlear ablation. In the cat unilateral cochlear ablation near hearing onset did not affect the size of neurons within the IC, while bilateral ablation produced a small but significant decrease (Nishiyama et al. 2000). These authors suggest that the profound changes in subcollicular auditory nuclei are not evident in the

IC following unilateral ear removal because the massive afferent convergence insures sufficient levels of afferent activity to maintain somatic size. Similarly, synaptic density in mature cats and ferrets is unchanged by unilateral cochlear removal at the time of ear opening (Hardie et al. 1998; Fuentes-Santamaria et al. 2003). Interestingly, synapse density was significantly lower in the mature cat IC after bilateral neonatal cochlear ablation, which was interpreted as a global decrease of auditory activity and a consequent effect on trophic support (Hardie et al. 1998).

### *3.3. EFFECT OF EAR REMOVAL ON AFFERENT BAND FORMATION IN THE INFERIOR COLLICULUS*

Even less is known about the impact of cochlear ablation on the development of IC bands and patches of afferents. Changes in the banded projection pattern of DNLL fibers, however, have been studied following unilateral neonatal cochlear removal on P2 in rats, before afferent bands develop (Gabriele et al. 2000b). The rats survived until P12, around onset of hearing, when DNLL input to the CNIC is banded (Fig. 18.7A, B) (Gabriele et al. 2000a). The projection pattern of crossed DNLL axons was visualized with a fluorescent carbocyanine dye tracer, diI (1,1'-dioctadecyl-3,3',3'- tetramethylindocarbocyanine perchlorate), which showed a sparse, but banded, distribution of crossed DNLL input in the contralateral IC (Fig. 18.7C). Ipsilateral to the ablation, the crossed DNLL input is considerably heavier; however, no banding pattern is evident (Fig. 18.7D). Thus, before ear opening, cochlear ablation profoundly affects the refinement of afferent projection patterns within the IC. Precisely how cochlear ablation leads to the changes in the banded DNLL input to the IC is unknown. Because cochlear ablation reduces spontaneous activity, it is tempting to ascribe the effect to activity per se or to a difference in the activity-regulated release of neurotrophic factors that regulate the symmetry of afferent patterns.

### *3.4. EFFECT OF EAR REMOVAL ON INFERIOR COLLICULUS FUNCTION*

Functional changes in the IC after early postnatal deafening demonstrate the considerable plasticity latent in the central auditory pathways in young animals and that can persist to a degree even in adults. Thus, frequency-specific IC circuits are altered by partial hearing loss after focal damage to the spiral ganglion (Snyder and Sinex 2002). The several prospective contributions to this process may remain plastic even in adults (Harrison et al. 1998; Snyder et al. 2000) and include plasticity from structural rearrangement, sprouting of afferents in adjacent layers, unmasking of silent synapses, or alterations in synaptic efficacy. These various consequences have just begun to be elucidated. Neonatal lesions restricted to a portion of the gerbil cochlea reorganize the IC tonotopic map (Harrison et al. 1998) and there is evidence that experience dependent changes may be adaptive or restorative (Silverman and Clopton 1977; Knudsen






Figure 18.7. DiI-labeled axons in IC of P12 rats following a midline diI pin placement in the decussation of DNLL fibers. Axons from the contralateral DNLL in unoperated, control animals are banded (*arrows*, **A** and **B**). Following cochlear ablation at P2, DNLL axons projecting to the IC contralateral to the ablation distribute densely in an unbanding pattern (**C**). In contrast, labeled axons projecting to the IC ipsilateral to the ablation are sparser, but a banded distribution (*arrows*) is still evident (**D**). Dorsal is toward the *top* and medial is to the *right* in (**A**) and (**C**) and to the *left* in (**B**) and (**D**). The scale bars represent 100  $\mu\text{m}$ .

1999). The mechanisms underlying these changes may include synaptic reorganization as well as adjustments in the efficacy of synaptic function.

Changes in the composition and distribution of IC neurotransmitter receptors that may mediate plasticity of functional properties after either monaural or binaural hearing loss are variable. For instance, whereas GABA or glutamic acid decarboxylase levels in IC decrease following bilateral hearing loss in rats, GABA<sub>A</sub> receptor expression (Marianowski et al. 2000) and receptor binding (Milbrandt et al. 2000) are up-regulated. Moreover, in bilaterally deafened rat pups, decreased expression of mature, and increased expression of immature, forms of AMPA subunit mRNA occur, as well as decreased NMDA receptor subunit mRNA (Marianowski et al. 2000). Furthermore, unilateral deafening in adult guinea pig results in modest but transient elevation in the number and activity of AMPA receptors (Suneja et al. 2000).

Unilateral deafening results in a paradoxical increase in excitability of the IC ipsilateral to stimulation of the intact ear (McAlpine 1997). This excitability likely reflects plasticity of lower auditory connections as well as expanded ipsilateral excitatory CN inputs to the IC (Moore and Kitzes 1985; Moore et al. 1989). Moreover, the unmasking of responses in deafened animals, whether immature or adult, may be attributable to down-regulation of inhibitory circuits, particularly intrinsic GABAergic mechanisms (Mossop et al. 2000).

Such activity-dependent changes in receptor distribution and composition affect the dynamics of neurotransmission and could serve to maintain a balance of excitation and inhibition in IC circuits, compensating for peripheral changes in hearing sensitivity. Similarly, prior to hearing onset and during assembly and refinement of IC circuitry, bilateral cochlear ablation increases excitatory postsynaptic currents and decreases inhibitory postsynaptic currents in neonatal gerbil IC (Vale and Sanes 2002). These changes may contribute to altered levels and localization patterns of intracellular calcium that mediate functional and structural adjustments. How such changes in activity or calcium levels contribute to alteration of afferent patterns (Gabriele et al. 2000b) or levels of calcium-binding proteins (Fuentes-Santamaria et al. 2003) in the IC is unknown.

#### 4. CONCLUSION AND SYNTHESIS

We have reviewed several events in the development of pre- and postsynaptic elements of banded circuits in the IC that may contribute ultimately to their formation, refinement, and plasticity (Fig. 18.2). The development of the presynaptic component of these circuits begins with the growth of axons from multiple brain stem auditory sources into the IC. En route to the IC, these growing neurites must make important decisions at decussation points to establish the appropriate laterality and binaural interactions. These decisions may be self-regulated, programmed or the result of either permissive or inhibitory cues in the local microenvironment through which they grow. Once neurites invade the IC, intrinsic cues likely guide the initial establishment of a coarse cochleo-

topic order (Figs. 18.2 and 18.3A). The ascending projections are simple in morphology and distribution, only minimal synaptogenesis has taken place, and the constellation of neurotransmitter receptors is immature. Thus, even before hearing begins, ascending projections segregate into discrete bands of aural dominance (Figs. 18.2 and 18.3B) in an activity-dependent manner, perhaps driven by competition between spontaneously active pathways from each ear. These projections then proliferate within IC in a manner consistent with trophic support mediated by activation of the target neurons. These afferents also must compete within the fibrodendritic framework with similar inputs to establish precise synaptic domains and dendritic alignment (Figs. 18.2 and 18.3C). Thus, local circuits are established by activity-dependent, calcium-mediated mechanisms. Once the basic pattern of afferent bands is established, the many neurotransmitter receptor systems begin their elaboration and auditory-driven activity begins to sculpt individual projection pathways and their synaptic domains via gene expression and cell-cell interactions. Finally, as the head grows or the sensitivity of the two ears changes over time, adaptive changes in circuits and efficacy of synapses maintain or restore relative synaptic throughput within IC synaptic domains.

## 5. FUTURE STUDIES

This chapter has emphasized development of ascending afferent projections to the CNIC. However, many basic issues of the development of projections to the other subdivisions of the IC are less well understood. Tracing studies should document the development of descending cortical and intrinsic commissural projections in relation to inputs ascending to the IC. In addition, the question whether the projections to different IC subdivisions develop independently should be investigated.

Further experimental studies are required to understand the role of competition in the development of layered inputs and the segregation of synaptic domains. Moreover, the role of spontaneous and patterned activity in the development of the complex circuitry of afferent projections remains to be explored more fully. Similarly, no information is currently available on the role of growth factors in the early development of IC circuits. Future studies of this issue are critical because activity-based regulation of growth factor expression may be the mechanism through which layered inputs and synaptic domains are established.

Future studies also should address the development and plasticity of synaptic domains within different functional inputs to the IC. Histochemical or immunohistochemical identification of discrete synaptic domains of related input-output elements should aid these studies. Once identified, multiple anatomical methods could assay the spatial relationships of discrete sets of functional inputs. For instance, the banded profiles of calbindin immunostained plexuses that are pronounced in kitten IC provide borders for some type of functional domain

within IC layers. Their boundaries might constitute a template that defines the distribution of receptor expression and maturation, dendritic rearrangement, or spine modification. Such features could be assayed in conditions of altered overall activity, afferent competition, or growth factor expression in order to study their effect on circuit refinement.

## Abbreviations

AMPA	$\alpha$ -amino-3-hydroxy-5-methyl-isoxazole-4-propionate
CN	cochlear nuclei
CNIC	central nucleus of the inferior colliculus
DCIC	dorsal cortex of the inferior colliculus
DiI	1-1'-dioctadecyl-3,3',3'-tetramethylindocarbocyanine dye
DNLL	dorsal nucleus of the lateral lemniscus
E	embryonic day
ECIC	external cortex of the inferior colliculus
GABA	$\gamma$ -aminobutyric acid
GluR	glutamate receptor
GLYT	glycine transporter
HRP	horseradish peroxidase
IC	inferior colliculus
LL	lateral lemniscus
LSO	lateral superior olivary nucleus
MNTB	medial nucleus of the trapezoid body
MSO	medial superior olivary nucleus
NMDA	<i>N</i> -methyl-D-aspartate
nAChR	nicotinic cholinergic receptors
P	postnatal day
VNLL	ventral nucleus of the lateral lemniscus

## Acknowledgments

This research was supported by United States Public Health Service Grants R01 DC 04412 (C.K.H. and J.B.B.) and R01 DC00813 (C.K.H.).

## References

- Aitkin LM and Moore DR (1975) Inferior colliculus. II. Development of tuning characteristics and tonotopic organization in central nucleus of the neonatal cat. *Journal of Neurophysiology* **38**:1208–1216.
- Aitkin LM and Reynolds A (1975) Development of binaural responses in the kitten inferior colliculus. *Neuroscience Letters* **1**:315–319.

- Aitkin L, Nelson J, Martsi-McClintock A, and Swann S (1996) Features of the structural development of the inferior colliculus in relation to the onset of hearing in a marsupial: the Northern Quoll, *Dasyurus hallucatus*. *Journal of Comparative Neurology* **375**:77–88.
- Aitkin L, Cochran S, Frost S, Martsi-McClintock A, and Masterton B (1997) Features of the auditory development of the short-tailed Brazilian opossum, *Monodelphis domestica*: evoked responses, neonatal vocalizations and synapses in the inferior colliculus. *Hearing Research* **113**:69–75.
- Altman JA and Bayer SA (1981) Time of origin of neurons of the rat inferior colliculus and the relations between cytogenesis and tonotopic order in the auditory pathway. *Experimental Brain Research* **42**:411–423.
- Andressen C, Blümcke I, and Celio MR (1993) Calcium-binding proteins: selective markers of nerve cells. *Cell and Tissue Research* **271**:181–208.
- Aramakis VB, Hsieh CY, Leslie FM, and Metherate R (2000) A critical period for nicotine-induced disruption of synaptic development in rat auditory cortex. *Journal of Neuroscience* **20**:1606–1616.
- Baimbridge KE, Celio MR, and Rogers JH (1992) Calcium-binding proteins in the nervous system. *Trends in Neurosciences* **15**:303–308.
- Blatchley BJ and Brugge JF (1990) Sensitivity to binaural intensity and phase difference cues in kitten inferior colliculus. *Journal of Neurophysiology* **64**:582–597.
- Brunso-Bechtold JK, Henkel CK, and Linville C (1992) Ultrastructural development of the medial superior olive (MSO) in the ferret. *Journal of Comparative Neurology* **324**:539–556.
- Caicedo A and Eybalin M (1999) Glutamate receptor phenotypes in the auditory brainstem and mid-brain of the developing rat. *European Journal of Neuroscience* **11**:51–74.
- Caicedo A, Kungel M, Pujol R, and Friauf E (1998) Glutamate-induced  $\text{Ca}^{2+}$  uptake in rat auditory brainstem neurons reveals developmental changes in  $\text{Ca}^{2+}$  permeability of glutamate receptors. *European Journal of Neuroscience* **10**:941–954.
- Dani JA (2001) Overview of nicotinic receptors and their roles in the central nervous system. *Biological Psychiatry* **49**:166–174.
- Dardennes R, Jarreau PH, and Meininger V (1984) A quantitative Golgi analysis of the postnatal maturation of dendrites in the central nucleus of the inferior colliculus of the rat. *Brain Research* **318**:159–169.
- Echteler SM, Arjmand E, and Dallos P (1989) Developmental alterations in the frequency map of the mammalian cochlea. *Nature* **341**:147–149.
- Faye-Lund H and Osen KK (1985) Anatomy of the inferior colliculus in rat. *Anatomy and Embryology* **171**:1–20.
- Fitzpatrick KA (1975) Cellular architecture and topographic organization of the inferior colliculus of the squirrel monkey. *Journal of Comparative Neurology* **164**:185–208.
- Friauf E (1992) Tonotopic order in the adult and developing auditory system of the rat as shown by c-fos immunocytochemistry. *European Journal of Neuroscience* **4**:798–812.
- Friauf E (1994) Distribution of calcium-binding protein calbindin-D28k in the auditory system of adult and developing rats. *Journal of Comparative Neurology* **349**:193–211.
- Friauf E, Hammerschmidt B, and Kirsch J (1997) Development of adult-type inhibitory glycine receptors in the central auditory system of rats. *Journal of Comparative Neurology* **385**:117–134.

- Friauf E, Aragon C, Löhrike S, Westenfelder B, and Zafra F (1999) Developmental expression of the glycine transporter GLYT2 in the auditory system of rats suggests involvement in synapse maturation. *Journal of Comparative Neurology* **412**:17–37.
- Fuentes-Santamaria V, Alvarado JC, Brunso-Bechtold JK, and Henkel CK (2003) Upregulation of calretinin immunostaining in the ferret inferior colliculus after cochlear ablation. *Journal of Comparative Neurology* **460**:585–596.
- Gabriele ML, Brunso-Bechtold JK, and Henkel CK (2000a) Development of afferent patterns in the inferior colliculus of the rat: projection from the dorsal nucleus of the lateral lemniscus. *Journal of Comparative Neurology* **416**:368–382.
- Gabriele ML, Brunso-Bechtold JK, and Henkel CK (2000b) Plasticity in the development of afferent patterns in the inferior colliculus of the rat after unilateral cochlear ablation. *Journal of Neuroscience* **20**:6939–6949.
- Gao E and Suga N (2000) Experience-dependent plasticity in the auditory cortex and the inferior colliculus of bats: role of the corticofugal system. *Proceedings of the National Academy of Sciences of the United States of America* **97**:8081–8086.
- Geniec P and Morest DK (1971) The neuronal architecture of the human posterior colliculus. *Acta Oto-Laryngologica Supplement* **295**:1–33.
- González-Hernández TH, Meyer G, and Ferres-Torres R (1989) Development of neuronal types and laminar organization in the central nucleus of the inferior colliculus in the cat. *Neuroscience* **30**:127–141.
- Hardie NA and Shepherd RK (1999) Sensorineural hearing loss during development: morphological and physiological response of the cochlea and auditory brainstem. *Hearing Research* **128**:147–165.
- Hardie NA, Martsi-McClintock A, Aitkin LM, and Shepherd RK (1998) Neonatal sensorineural hearing loss affects synaptic density in the auditory midbrain. *NeuroReport* **9**:2019–2022.
- Harris DM and Dallos P (1984) Ontogenetic changes in frequency mapping of a mammalian ear. *Science* **225**:741–743.
- Harrison RV, Ibrahim D, and Mount RJ (1998) Plasticity of tonotopic maps in auditory midbrain following partial cochlear damage in the developing chinchilla. *Experimental Brain Research* **123**:449–460.
- Henkel CK and Brunso-Bechtold JK (1993) Laterality of superior olive projections to the inferior colliculus in adult and developing ferret. *Journal of Comparative Neurology* **331**:458–468.
- Henkel CK and Spangler KM (1983) Organization of the efferent projections of the medial superior olivary nucleus in the cat as revealed by HRP and autoradiographic tracing methods. *Journal of Comparative Neurology* **221**:416–428.
- Hollmann M, Hartley M, and Heinemann S (1991) Ca<sup>2+</sup> permeability of KA-AMPA-gated glutamate receptor channels depends on subunit composition. *Science* **252**:851–853.
- Hume RI, Dingledine R, and Heinemann SF (1991) Identification of a site in glutamate receptor subunits that controls calcium permeability. *Science* **253**:1028–1031.
- Hyson RL and Rudy JW (1987) Ontogenetic changes in the analysis of sound frequency in the infant rat. *Developmental Psychobiology* **20**:189–207.
- Kandler K and Friauf E (1993) Pre- and postnatal development of efferent connections of the cochlear nucleus in the rat. *Journal of Comparative Neurology* **328**:161–184.
- Keiger CJ, Henkel CK, and Brunso-Bechtold JK (2003) Normal postnatal development of banded afferent projections to the inferior colliculus of the ferret from the superior olivary complex. *Association for Research in Otolaryngology Abstracts* **26**:145–146.

- Kitzes LM, Kageyama GH, Semple MN, and Kil J (1995) Development of ectopic projections from the ventral cochlear nucleus to the superior olivary complex induced by neonatal ablation of the contralateral cochlea. *Journal of Comparative Neurology* **353**:341–363.
- Knudsen EI (1999) Mechanisms of experience-dependent plasticity in the auditory localization pathway of the barn owl. *Journal of Comparative Physiology A* **185**:305–321.
- Kraushaar U and Backus KH (2002) Characterization of GABA<sub>A</sub> and glycine receptors in neurons of the developing rat inferior colliculus. *Pflügers Archiv European Journal of Physiology* **445**:279–288.
- Kudo M (1981) Projections of the nuclei of the lateral lemniscus in the cat: an autoradiographic study. *Brain Research* **221**:57–69.
- Kungel M and Friauf E (1995) Somatostatin and leu-enkephalin in the rat auditory brainstem during fetal and postnatal development. *Anatomy and Embryology* **191**:425–443.
- Laurie DJ, Wisden W, and Seeburg PH (1992) The distributions of thirteen GABA<sub>A</sub> receptor subunit mRNAs in the rat brain. III. Embryonic and postnatal development. *Journal of Neuroscience* **12**:4151–4172.
- Lippe W and Rubel EW (1983) Development of the place principle: tonotopic organization. *Science* **219**:514–516.
- Lohmann C and Friauf E (1996) Distribution of the calcium-binding proteins parvalbumin and calretinin in the auditory brainstem of adult and developing rats. *Journal of Comparative Neurology* **367**:90–109.
- Ma X and Suga N (2003) Augmentation of plasticity of the central auditory system by the basal forebrain and/or somatosensory cortex. *Journal of Neurophysiology* **89**:90–103.
- Malmierca MS, Blackstad TW, Osen KK, Karagülle T, and Molowny RL (1993) The central nucleus of the inferior colliculus in rat: a Golgi and computer reconstruction study of neuronal and laminar structure. *Journal of Comparative Neurology* **333**:1–27.
- Marianowski R, Liao W-H, Van Den Abbeele T, Fillit P, Herman P, Frachet B, and Huy PTB (2000) Expression of NMDA, AMPA and GABA<sub>A</sub> receptor subunit mRNAs in the rat auditory brainstem. I. Influence of early auditory deprivation. *Hearing Research* **150**:1–11.
- McAlpine D, Martin RL, Mossop JE, and Moore DR (1997) Response properties of neurons in the inferior colliculus of the monaurally deafened ferret to acoustic stimulation of the intact ear. *Journal of Neurophysiology* **78**:767–779.
- McGehee DS (2002) Nicotinic receptors and hippocampal synaptic plasticity . . . it's all in the timing. *Trends in Neurosciences* **25**:171–172.
- McHaffie JG, Gabriele ML, Stein BE, Wallace MT, and Henkel CK (2000) Banded distribution of calbindin D-28k immunoreactivity in the auditory brainstem of neonatal and adult cats. *Society for Neuroscience Abstracts* **26**:676.
- Meininger V and Baudrimont M (1981) Postnatal modifications of the dendritic tree of cells in the inferior colliculus of the cat. *Journal of Comparative Neurology* **200**:339–355.
- Milbrandt JC, Holder TM, Wilson MC, Salvi RJ, and Caspary DM (2000) GAD levels and muscimol binding in rat inferior colliculus following acoustic trauma. *Hearing Research* **147**:251–260.
- Mills DM and Rubel EW (1998) Development of the base of the cochlea: place code shift in the gerbil. *Hearing Research* **122**:82–96.

- Moore DR (1990) Auditory brainstem of the ferret: bilateral cochlear lesions in infancy do not affect the number of neurons projecting from the cochlear nucleus to the inferior colliculus. *Developmental Brain Research* **54**:125–130.
- Moore DR and Irvine DRF (1980) Development of binaural input, response patterns and discharge rate in single units of the cat inferior colliculus. *Experimental Brain Research* **38**:103–108.
- Moore DR and Irvine DRF (1981a) Development of responses to acoustic interaural intensity differences in the cat inferior colliculus. *Experimental Brain Research* **41**:301–309.
- Moore DR and Irvine DRF (1981b) Plasticity of binaural interaction in the cat inferior colliculus. *Brain Research* **208**:198–202.
- Moore DR and Kitzes LM (1985) Projections from the cochlear nucleus to the inferior colliculus in normal and neonatally cochlea-ablated gerbils. *Journal of Comparative Neurology* **240**:180–195.
- Moore DR and Kowalchuk NE (1988) Auditory brainstem of the ferret: effects of unilateral cochlear lesions on cochlear nucleus volume and projections to the inferior colliculus. *Journal of Comparative Neurology* **272**:503–515.
- Moore DR, Hutchings ME, King AJ, and Kowalchuk NE (1989) Auditory brain stem of the ferret: some effects of rearing with a unilateral ear plug on the cochlea, cochlear nucleus, and projections to the inferior colliculus. *Journal of Neuroscience* **9**:1213–1222.
- Morest DK and Oliver DL (1984) The neuronal architecture of the inferior colliculus in the cat: defining the functional anatomy of the auditory midbrain. *Journal of Comparative Neurology* **222**:209–236.
- Morley BJ and Happe HK (2000) Cholinergic receptors: dual roles in transduction and plasticity. *Hearing Research* **147**:104–112.
- Mossop JE, Wilson MJ, Caspary DM, and Moore DR (2000) Down-regulation of inhibition following unilateral deafening. *Hearing Research* **147**:183–187.
- Müller M (1991) Developmental changes of frequency representation in the rat cochlea. *Hearing Research* **56**:1–7.
- Niblock MM, Brunso-Bechtold JK, and Henkel CK (1995) Fiber outgrowth and path-finding in the developing auditory brainstem. *Developmental Brain Research* **85**:288–292.
- Nishiyama N, Hardie NA, and Shepherd RK (2000) Neonatal sensorineural hearing loss affects neurone size in cat auditory midbrain. *Hearing Research* **140**:18–22.
- Nordeen KW, Killackey HP, and Kitzes LM (1981) Ascending projections to the inferior colliculus following unilateral cochlear ablation in the neonatal gerbil, *Meriones unguiculatus*. *Journal of Comparative Neurology* **214**:144–153.
- Okoyama S, Morizumi T, Kitao Y, Kawano J, and Kudo M (1995) Postnatal development of the projection from the medial superior olive to the inferior colliculus in the rat. *Hearing Research* **88**:65–70.
- Oliver DL (1984) Dorsal cochlear nucleus projections to the inferior colliculus in the cat. A light and electron microscopic study. *Journal of Comparative Neurology* **224**:155–172.
- Oliver DL (2000) Ascending efferent projections of the superior olivary complex. *Microscopic Research Techniques* **51**:355–363.
- Oliver DL and Morest DK (1984) The central nucleus of the inferior colliculus in the cat. *Journal of Comparative Neurology* **222**:237–264.



- Oliver DL and Shneiderman A (1989) The anatomy of the inferior colliculus: a cellular basis for integration of monaural and binaural information. In: Altschuler RA, Bobbin RP, Clopton BM, and Hoffman DW (eds). *Neurobiology of Hearing: The Central Auditory System*. Raven Press, New York, pp. 195–222.
- Oliver DL, Beckius GE, Bishop DC, and Kuwada S (1997) Simultaneous anterograde labeling of axonal layers from lateral superior olive and dorsal cochlear nucleus in the inferior colliculus of cat. *Journal of Comparative Neurology* **382**:215–229.
- Piechotta K, Weth E, Harvey RJ, and Friauf E (2001) Localization of rat glycine receptor  $\alpha 1$  and  $\alpha 2$  subunit transcripts in the developing auditory brainstem. *Journal of Comparative Neurology* **438**:336–352.
- Pysh JJ (1969) The development of the extracellular space in neonatal rat inferior colliculus. *American Journal of Anatomy* **124**:411–430.
- Radcliffe KA and Dani JA (1998) Nicotinic stimulation produces multiple forms of increased glutamatergic synaptic transmission. *Journal of Neuroscience* **18**:7075–7083.
- Rockel AJ and Jones EG (1973) The neuronal organization of the inferior colliculus of the adult cat. I. The central nucleus. *Journal of Comparative Neurology* **147**:11–60.
- Romand R (1997) Modification of tonotopic representation in the auditory system during development. *Progress in Neurobiology* **51**:1–17.
- Romand R and Ehret G (1990) Development of tonotopy in the inferior colliculus. I. Electrophysiological mapping in house mice. *Developmental Brain Research* **54**:221–234.
- Rubel EW, Lippe WR, and Ryals BM (1984) Development of the place principle. *Annals of Otolaryngology, Rhinology and Laryngology* **93**:609–615.
- Russell FA and Moore DR (1995) Afferent reorganization within the superior olivary complex of the gerbil: development and induction by neonatal, unilateral cochlear removal. *Journal of Comparative Neurology* **352**:607–625.
- Ryan AF and Woolf NK (1988) Development of tonotopic representation in the Mongolian gerbil: a 2-deoxyglucose study. *Brain Research* **469**:61–70.
- Sanes DH, Merickel M, and Rubel EW (1989) Evidence for an alteration of the tonotopic map in the gerbil cochlea during development. *Journal of Comparative Neurology* **279**:436–444.
- Sato K, Shiraishi S, Nakagawa H, Kuriyama H, and Altschuler RA (2000) Diversity and plasticity in amino acid receptor subunits in the rat auditory brain stem. *Hearing Research* **147**:137–144.
- Schmid S, Guthmann A, Ruppertsberg JP, and Herbert H (2001) Expression of AMPA receptor subunit flip/flop splice variants in the rat auditory brainstem and inferior colliculus. *Journal of Comparative Neurology* **430**:160–171.
- Schweitzer L, Lutz C, Hobbs M, and Weaver SP (1996) Anatomical correlates of the passive properties underlying the developmental shift in the frequency map of the mammalian cochlea. *Hearing Research* **97**:84–94.
- Sekitani M, Shiosaka S, Kuriyama H, Lee Y, Ikeda M, and Tohyama M (1990) Transient expression of somatostatin mRNA in the auditory system of neonatal rat. *Molecular Brain Research* **7**:177–181.
- Shneiderman A and Henkel CK (1987) Banding of lateral superior olivary nucleus afferents in the inferior colliculus: a possible substrate for sensory integration. *Journal of Comparative Neurology* **266**:519–534.

- Shneiderman A, Oliver DL, and Henkel CK (1988) Connections of the dorsal nucleus of the lateral lemniscus: an inhibitory parallel pathway in the ascending auditory system? *Journal of Comparative Neurology* **276**:188–208.
- Silverman MS and Clopton BM (1977) Plasticity of binaural interaction. I. Effect of early auditory deprivation. *Journal of Neurophysiology* **40**:1266–1274.
- Snyder RL and Sinex DG (2002) Immediate changes in tuning of inferior colliculus neurons following acute lesions of cat spiral ganglion. *Journal of Neurophysiology* **87**:434–452.
- Snyder RL, Sinex DG, McGee JD, and Walsh EW (2000) Acute spiral ganglion lesions change the tuning and tonotopic organization of cat inferior colliculus neurons. *Hearing Research* **147**:200–220.
- Suneja SK, Potashner SJ, and Benson CG (2000) AMPA receptor binding in adult guinea pig brain stem auditory nuclei after unilateral cochlear ablation. *Experimental Neurology* **165**:355–369.
- Thoss VS, Kungel M, Friauf E, and Hoyer D (1996) Presence of somatostatin sst<sub>2</sub> receptors in the developing rat auditory system. *Developmental Brain Research* **97**:269–278.
- Tierney TS, Russell FA, and Moore DR (1997) Susceptibility of developing cochlear nucleus neurons to deafferentation-induced death abruptly ends just before the onset of hearing. *Journal of Comparative Neurology* **378**:295–306.
- Trune DR (1982a) Influence of neonatal cochlear removal on the development of mouse cochlear nucleus: I. Number, size, and density of its neurons. *Journal of Comparative Neurology* **209**:409–424.
- Trune DR (1982b) Influence of neonatal cochlear removal on the development of mouse cochlear nucleus: II. Dendritic morphometry of its neurons. *Journal of Comparative Neurology* **209**:425–434.
- Vale C and Sanes DH (2002) The effect of bilateral deafness on excitatory and inhibitory synaptic strength in the inferior colliculus. *European Journal of Neuroscience* **16**:2394–2404.
- Webster DB and Webster M (1977) Neonatal sound deprivation affects brain stem auditory nuclei. *Archives of Otolaryngology* **103**:392–396.
- Whitley JM and Henkel CK (1984) Topographical organization of the inferior collicular projection and other connections of the ventral nucleus of the lateral lemniscus in the cat. *Journal of Comparative Neurology* **229**:257–270.
- Zook JM, Winer JA, Pollak GD, and Bodenhamer RD (1985) Topology of the central nucleus of the mustache bat's inferior colliculus: correlation of single unit properties and neuronal architecture. *Journal of Comparative Neurology* **231**:530–546.

# Chapter 19

## Inferior Colliculus: Aging and Plasticity

ROBERT D. FRISINA AND RAMESH RAJAN

### 1. INTRODUCTION

Classic views and theories of the central nervous system (CNS) emphasized that mammalian neural circuitry formed during development remained essentially invariant in adulthood. No new neurons are added, and CNS damage entails neuron loss, resulting in permanent anatomical damage and enduring functional impairments. However, advances in the study of plasticity and aging in the CNS, including the central auditory system, reveal that neural systems can reconfigure themselves with age and in response to the loss of peripheral inputs from sensory end-organs. For instance, peripheral abnormalities induce unmasking, rewiring, or changes in functional responses in the central auditory system, sometimes termed “peripherally induced central changes” (Frisina et al. 2001). Alternatively, in the case of age-related changes in the central auditory system, deficits can occur independently of age-dependent alterations of the inner ear. Instances of these plastic and aging phenomena are presented in this chapter, focusing on those manifested in the inferior colliculus (IC).

### 2. AGE-RELATED FUNCTIONAL CHANGES

#### 2.1. *TONOTOPIC ORGANIZATION*

Neuroanatomical and neurophysiological investigations revealed age-related tonotopic plasticity in the auditory midbrain in response to the loss of high-frequency input of cochlear origin (Willott 1984, 1986). The C57Bl/6 mouse strain suffers from accelerated peripheral age-related hearing loss, starting in the base of the cochlea. Like human age-related hearing loss (presbycusis), the C57 syndrome begins with an impairment of the high frequencies, which then spreads to the lower frequencies with age. The C57 high-frequency hearing loss is much more rapid than that of human presbycusis, even when correcting for the different absolute lifespans of mice and men. Specifically, young adult C57s about 6 months old have severe-to-profound high-frequency hearing losses and de-

velop profound deficits at all frequencies in their second year. Because a 6-month-old C57 mouse has an “old” ear but a young brain, it permits the dissection of aspects of presbycusis inherent in the inner ear and largely independent of the aging brain.

The IC tonotopic reorganization was documented by neurophysiological mapping in the central nucleus (ICC) across time in C57 mice (Willott 1984, 1986). In young adults, as in most mammals, there is a dorsoventral gradient of single-unit characteristic frequencies (CFs) orthogonal to the well defined neuroanatomical ICC laminae, with lower CFs dorsally and higher CFs ventrally. With age, high-CF units shift to lower frequencies, although the tuning curve tip sensitivity remains surprisingly good, indicative of a true CF shift and not merely a loss of the sensitive tips of high CF units (Fig. 19.1). Apparently, a development, strengthening, or enhancement process involving new synaptic connections occurs in the formerly high-frequency IC regions and/or its brain stem input nuclei, such that the low CFs become overrepresented in the tonotopic map.

In contrast to the C57, the CBA mouse strain has a slow, progressive age-related hearing loss that is similar (flat) across the auditory frequency range. This strain has been useful for studying age-related changes in the central auditory system when only moderate changes take place in the auditory periphery. Not surprisingly, in contrast to the C57 ICC reorganization, a tonotopic reorganization of the IC in aged CBA mice does not occur (Willott 1991a; Walton et al. 1998).

## 2.2. TEMPORAL PROCESSING

Sound temporal features are essential for effective processing and perception of biologically relevant acoustic stimuli such as speech, animal vocalizations, and music (see Chapters 12 and 14). In presbycusis, aged listeners often have auditory temporal processing deficits that contribute to their impairment in speech comprehension in the presence of background noise (Snell and Frisina 2000; Frisina et al. 2001). Temporal processing deficits may be due to a sloping hearing loss of peripheral origin. When aged persons have good peripheral sensitivity, speech perception problems may involve temporal processing deficits caused by brain stem auditory pathology, including the IC (Frisina and Frisina 1997; Frisina et al. 2001).

Gap stimuli and amplitude-modulated stimuli are two of the most commonly utilized sounds for investigating auditory temporal processing deficits in cases of hearing loss, including presbycusis (Frisina 2001a). Gap detection paradigms involve measuring the minimum detectable silent interval between two sounds. These sounds are sometimes referred to as “markers,” or the “masker” and the “target,” and may be pure tones or wideband noise bursts. Gap detection experiments assess the smallest gap that a human listener, behaving animal, or single neuron can perceive, encode, or process. Temporal envelope features are represented by amplitude-modulated (AM) sounds. The most common AM stimulus




Figure 19.1. The tonotopic (cochleotopic) map of CFs in the C57 mouse IC reorganizes after an age-related decline in inputs from the high-frequency region (basal turn) of the cochlea. Group average frequency-threshold tuning curves (MTCs) for units, as a function of age and dorsoventral ICC depth, show a statistically significant CF shift from high (25 kHz) to middle (10 to 12 kHz) frequencies at the three lowest depths for the 7- to 12-month old subjects relative to the young adults. The number of single-unit tuning curves comprising each of the group average MTCs is: 1 month,  $n = 90$ ; 7 months,  $n = 74$ ; 9 months,  $n = 50$ ; 12 months,  $n = 56$ . (From Willott 1986.)

invokes a periodic, usually sinusoidal, fluctuation in a tone's or wideband noise's envelope. Here, the threshold for the minimal depth of modulation can be measured or, more commonly in physiological experiments, the strength of the response of a single-unit or multiunit cluster is determined, either in terms of number of action potentials (rate) or timing of the action potentials (synchrony) in response to the AM (see Chapter 12). Both gap and AM coding can change with age in the IC.



### 2.2.1. Sound Gaps

Single-unit gap encoding in the ICC of unanesthetized CBA mice is consistent with behavioral gap detection, both of which degrade with age, starting in middle age (Walton et al. 1997, 1998). Specifically, units with short gap detection thresholds decline, and the strength of their gap responses decreases in the ICC of aged CBA mice (Fig. 19.2). It is unknown what portion of this age effect occurs within the ICC and in brain stem nuclei that project to the ICC. It is likely that much of this age-related temporal processing decline occurs at ICC, as the coding of gaps there is of a different nature than that at subcollicular sites. For instance, at the auditory nerve and cochlear nucleus, gaps are encoded by a decrease in single-unit firing rate during the gap. In contrast, ICC neurons respond to gaps with an increase in neural firing at the end of the gap, and it is this response that declines with age in CBA mice with relatively good peripheral hearing sensitivity.

### 2.2.2. Amplitude Modulation

The neural encoding of ongoing auditory temporal features is investigated in AM experiments. Using monaural stimuli in anesthetized rats, there were no dramatic changes in ICC sinusoidal AM coding with age, no significant declines in the upper cutoff frequencies of the modulation transfer functions, or any marked changes in other temporal and rate measures of AM coding. However,

Figure 19.2. The proportion of single units in the unanesthetized CBA mouse IC with short gap thresholds decreases in old animals. The percentage of units in young adult and old CBA mice with minimum gap thresholds (MGTs) from 1 to >11 ms for young adult (*hatched*,  $n = 78$ ) and aged mice (*solid*,  $n = 108$ ). The distribution shows longer gap thresholds for aged animals. IC cells in young adult units encode gaps with higher firing rates. Neural recovery functions for 30 phasic units in young adult (*top*) and aged (*bottom*) animals. Recovery (*vertical axis*) was measured by computing the number of spikes elicited by the noise burst following the gap, divided by those to the noise burst preceding it,  $\times 100$ . This calculation included all gap durations for each unit. A 100% recovery (*horizontal dashed line*) designates equal discharges to both bursts. Gap response recovery to the post-gap noise burst is 75% by <10 to 15 ms for almost all young adult units, whereas most aged units do not reach this criterion at even the longest gaps. (From Walton et al. 1998.)



there was a significant shift in the shape distribution of modulation transfer functions, with a decline in band-pass, and an increase in the low-pass shape, with age, in both the ICC and the external nucleus of the IC. This was interpreted as consistent with the hypothesis that IC inhibition contributes to the formation of band-pass AM transfer functions in young adult animals, and if this inhibition declines with age (see later), then band-pass specificity of AM coding will also degrade (Shaddock-Palombi et al. 2001).

Subsequent sinusoidal AM studies in the unanesthetized mouse ICC used free-field stimuli and also demonstrated the age-related shift from band-pass to low-pass modulation transfer function shapes, originally found in the aged rat (Walton et al. 2002). The various shapes of modulation transfer functions characteristic of the mammalian IC (Fig. 19.3A) shift from a preponderance of band-pass transfer functions in young CBA mice to more low-pass in old mice (Fig. 19.3B). Other aspects of AM processing change with age, such as declining upper cutoff and best modulation frequencies (Fig. 19.3C) of rate modulation transfer functions, and the increased spike rate for pure tone stimulation was amplified in old animals in response to AM stimuli, further supporting a loss of IC inhibition with age.




### 2.3. SPATIAL PROCESSING

The auditory system uses different neural mechanisms for separating signals of interest, such as speech, from interfering acoustic clutter, such as ambient noise. Spatial localization of a speaker of interest in the context of noise from other speakers is a critical auditory adaptation for enhancing the signal-to-noise ratio in complex acoustic environments. It has been postulated, from clinical research, that this ability degrades with age (Willott 1991b). Consistent with the initial behavioral experiments with human subjects and patients, animal experiments found that sound localization abilities decline with age in C57 mice, and in a manner independent of their high-frequency hearing loss that occurs between young adulthood and middle age (Heffner et al. 2001).

What is the neural basis for age-related declines in the ability of the brain stem auditory system to separate signals from noise spatially? Single IC units in middle-aged C57 mice did not benefit from moving background noise sources away from signal locations, as do cells in young adult C57 mice. At a given

Figure 19.3. Age-related effects on rate encoding of envelope periodicities for IC single units. (A) Four different single-unit amplitude modulation transfer functions (MTF) types showing the scheme for MTF filter shape classification. Five categories occur: low-pass with a maximum synchronization index (SI<sub>m</sub>) of 0.889; band-pass, SI<sub>m</sub> 0.939; high-pass, SI<sub>m</sub> 0.946; all-pass, SI<sub>m</sub> 0.703; band-reject, example not shown. (B) Young adult (*open bars*) and aged (*filled bars*) units with each rate MTF filter shape. (C) Young adult (*open bars*) and old (*closed bars*) mice units with rate BMFs at each of the nine AM frequencies tested. (From Walton et al. 2002 with permission.)





azimuthal location, the amount of masking from the background noise was greater in the older mice, a change that peripherally induced threshold shifts with age could not account for and that implicates age-related declines in the mammalian brain stem auditory spatial localization analysis system (McFadden and Willott 1994a,b).

### 3. NEUROANATOMICAL AND CHEMICAL CHANGES WITH AGE

#### 3.1. CHANGES IN INPUT PATHWAYS

Do IC brain stem connections also change in aged mice and, if so, is altered connectivity implicated in this syndrome? A neuroanatomical retrograde tracer, horseradish peroxidase, was injected into the centers of the IC regions from which recordings were obtained in temporal processing experiments (Frisina et al. 1998; Frisina and Walton 2001a,b). Inputs from all three divisions of the contralateral cochlear nucleus, the ipsilateral anterolateral periolivary nucleus, and portions of the ventral nucleus of the lateral lemniscus, each decline with age. These inputs likely contribute to the normal IC temporal processing abilities in young adult animals. As inputs age and decline, the balance of excitatory and inhibitory inputs to IC principal cells may become disrupted and precise temporal processing is diminished, leading to degradations in gap coding. These declines in brain stem inputs to the IC begin in middle-aged mice who have relatively normal auditory sensitivity as measured with auditory brain stem responses (ABRs), a finding consistent with the human mid-life psychophysical declines in temporal processing, even in subjects with otherwise normal audiograms (Snell and Frisina 2000).

#### 3.2. CHANGES IN INHIBITORY SYSTEM

GABA ( $\gamma$ -aminobutyric acid) is one of the main auditory brain stem neurotransmitters, and it is prominent in ICC neurons and synaptic endings where it has been implicated in many aspects of sound processing, including temporal and spatial coding. A study of IC GABAergic systems found that many aspects decline with age (Caspary et al. 1995). Quantitative analyses of GABA immunolabeling in IC perikarya in young and aged Fischer 344 rats found a 36% reduction with age (Caspary et al. 1990) and a reduced basal and  $K^+$ -evoked GABA release, in contrast to the normal values in control excitatory neurotransmitter levels and in acetylcholine release. Using quantitative receptor autoradiography, it was shown that declining GABA<sub>B</sub> receptor binding occurs in all three divisions of the aged rat IC (Milbrandt et al. 1994) while cerebellar tissue was unchanged. In contrast, GABA<sub>A</sub> receptors increased with age, perhaps in compensation for the GABA<sub>B</sub> decline (Milbrandt et al. 1996). Immunogold electron microscopy demonstrates that rat IC excitatory and inhibitory synapses

decrease in density with age (Helfert et al. 1999), just as does synaptic density in the IC of C57 mice (Kazee et al. 1995), although not in aged CBA mice (Kazee and West 1999). In conclusion, in rats, most aspects of GABA neurotransmitter biochemistry degrade with age, perhaps causing an imbalance of excitation and inhibition in the IC and concomitant functional impairments such as deficits in auditory signal extraction in background noise.

### 3.3. ACTIVITY-DEPENDENT CHANGES IN CALCIUM-BINDING PROTEINS

Several competing theories and models address age-related neural degeneration and cell death in the mammalian CNS. These include processes such as calcium excitotoxicity; apoptosis; and damage caused by the harmful by-products of oxidative phosphorylation such as free radicals, stress/inflammatory responses in the brain, and others. Calcium excitotoxicity can result from concentration imbalances in intracellular calcium-binding proteins such as calbindin, calretinin, or parvalbumin. Age-related changes in calcium-binding proteins have been found in the IC (Zettel et al. 1997; Frisina 2001b,c). Antibodies for calbindin and calretinin reveal an age-related decline in IC calbindin immunostaining of CBA and C57 strains. For calretinin, up regulation occurs in the aged CBA IC, but not in aged C57 mice. Perhaps the age-related up regulation of calretinin in CBA mice reflects sound-evoked IC activity, as this up regulation was absent in aged, deaf C57s. This hypothesis was confirmed in CBA mice deafened as young adults and compared to the aged controls (Zettel et al. 2001). Calretinin up regulation in aged, hearing CBAs was absent in the mice deafened as juveniles (Fig. 19.4).

## 4. PLASTICITY IN THE ADULT MAMMALIAN INFERIOR COLLICULUS CAUSED BY ALTERED COCHLEAR OUTPUTS

Plasticity in the IC has been studied at many time points in the lives of experimental animals. This section examines studies in the adult, but not aged, mammalian IC.

The mature IC possesses the neural machinery that, at other brain sites, plays important roles in neural plasticity, including the *N*-methyl-D-aspartate (NMDA) glutamate receptors, the  $\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazole propionic acid (AMPA) glutamate receptor, and GABA<sub>A</sub> receptors (Casparly et al. 2002; Kelly and Zhang 2002; Ma et al. 2002; Pollak et al. 2002). Many neuromodulators such as acetylcholine, serotonin, and noradrenaline (Habbicht and Vater 1996; Hurley et al. 2002) are present and can influence neuronal response plasticity in specific brain areas. IC cells express long-term potentiation (LTP), a synaptic phenomenon that is an important form of neural plasticity (Buonomano and Merzenich 1998). Various chemical levels in IC show immediate




Figure 19.4. Mean cell density of calretinin-stained (CR+) cells per 100  $\mu\text{m}^2$  in the dorsomedial IC of aged, deafened CBA mice, aged hearing CBA, young adult hearing CBA, and young adult hearing C57 mice. Aged deafened CBA mice do not show the statistically significant increase in CR+ cells discovered in the aged hearing CBA mice. There was no significant difference between the cross-sectional areas of the dorsomedial IC in any of the ages or strains tested (not shown). Error bars denote standard errors of the mean. (From Zettel et al. 2001.)

changes as well as long-term changes after major alterations in afferent input. It is therefore surprising that, except in aging, there are very few demonstrations of physiological response plasticity in the adult mammalian IC.

An important distinction is made between plasticity and “unmasking” of latent inputs. As noted in the somatosensory CNS after damage or other manipulations to the receptor surface, CNS plasticity cannot simply mean the expression of latent, functionally dormant inputs once these have been released from inhibition (i.e., “unmasked”) after removal of a dominant input (Calford and Tweedale 1988, 1991). Instead, plasticity must involve the conversion of implicit but unexpressed inputs into explicit responses expressed as part of the cell’s normal suprathreshold repertoire, whereas unmasking entails the expression of explicit inputs that are otherwise blocked by other dominant (often inhibitory) inputs. Plasticity must involve changes in synaptic weights (Calford and Tweedale 1988, 1991; Buonomano and Merzenich 1998) that would not occur in unmasking. This distinction between plasticity and unmasking has been made in auditory cortex (Rajan 2001) and in the IC (Snyder et al. 2000) and will guide this exposition. It does not mean that the IC only has a stereotyped repertoire of hard-wired responses unaltered by major alterations of afferent or efferent input, or that unmasking does not produce functionally important response changes. However, it does provide a conceptual context for thinking about these changes and their causes. Finally, studies of IC plasticity in adults are confined to ICC unless otherwise stated.

#### 4.1. INHIBITORY NEUROCHEMISTRY IN THE INFERIOR COLLICULUS AFTER COCHLEAR DAMAGE

IC neurochemistry has been examined often after cochlear ablation or total inactivation, and for focal cochlear damage with acoustic trauma. Although the former type of manipulation is not representative of most clinical conditions, it allows assessment of IC neurochemical reorganization after large changes in afferent input.

Many studies examined the effects on GABA<sub>A</sub> receptors, as hyperactivity or increased spread of activity often follows cochlear damage (Ryan et al. 1992; Bledsoe et al. 1995; Nagase et al. 2000; Salvi et al. 2000). IC cells express LTP only after blocking of some, if not most, inhibitory inputs such as GABA<sub>A</sub> receptors (Hosomi et al. 1995). LTP required NMDA and GABA<sub>B</sub> receptor activation—the latter likely acting presynaptically on GABAergic neurons to reduce postsynaptic inhibitory potentials mediated through GABA<sub>A</sub> receptors—and blocking of glycinergic receptors (Zhang and Wu 2002). LTP also occurs in brain slices in which GABAergic and glycinergic inhibition was routinely suppressed pharmacologically (Wu et al. 2002). These studies provide the rationale for examining changes in inhibition-related parameters in IC after cochlear damage.

There are considerable changes in GABAergic indices in IC after alterations of cochlear outflow. An immediate (Milbrandt et al. 2000; immediately after, and 42 hours after focal acoustic trauma) or rapid (Mossoop et al. 2000; 24 hours and 7 days post-cochleotomy, but not 4 hours post) decrease in levels of glutamic acid decarboxylase (GAD), the rate-limiting enzyme in GABA synthesis, occurs in both membrane and cytosolic fractions (Milbrandt et al. 2000). Return to normal GAD levels follows 30-day (Milbrandt et al. 2000) or 1-year survivals (Mossoop et al. 2000). However, GAD levels (primarily GAD67, the GAD isoform of molecular mass 67,000 kDa) in the entire IC increased immediately after focal acoustic trauma (Abbott et al. 1999), returned to near normal levels 2 days post-trauma, then dropped significantly below control levels at 30 days with a permanent hearing loss of 20 to 25 dB in auditory brain stem response thresholds. A similar trend occurs for GAD-immunoreactive cells in all IC divisions, although these changes were rarely significant and 30 days post-trauma the number was similar to that of normal animals. When salicylate was chronically administered in drinking water for 4 weeks or 4 months to produce a very small elevation in rat auditory brain stem evoked response (ABR) thresholds (Bauer et al. 2000), there was a marginal increase in GAD65 levels at 4 weeks treatment and a significant increase at 4 months.

Variability occurs in other indices of GABAergic function. A significant decrease in evoked GABA release occurs 21 days after ototoxic drug-induced bilateral deafness (Bledsoe et al. 1995). Evoked GABA release exceeded control levels in the contralateral IC 2 to 5 days after unilateral cochleotomy or ossicular removal, fell to normal at 59 days, and increased again at 145 days, with no

changes ipsilaterally. There were small decreases in GABA uptake in both ICs after cochleotomy only after the 145-day survival, in contrast with an immediate increase followed by a return to normal (ipsilateral) or subnormal (contralateral) levels after ossicle removal (Suneja et al. 1998). GABA<sub>A</sub> receptor function shows similar variability: increased binding when tested with muscimol (Milbrandt et al. 2000) and a small decrease in binding sites but increased receptor affinity to muscimol and not other GABA<sub>A</sub> ligands (Bauer et al. 2000). Thus, these studies do not show that a decrease in cochlear outflow consistently reduces IC GABAergic levels or expression.

Glycine is present in low amounts in the IC compared to GABA and has received correspondingly less attention. Little change in glycine release was seen 21 days after ototoxic drug-induced bilateral deafness in rats (Bledsoe et al. 1995). In guinea pigs a small and transient increase in glycine receptor binding occurred only in dorsal cortex (ICD) and the lateral nucleus (LIC) 2 days after unilateral cochleotomy and became normal 31 days later, decreased slightly in all three IC divisions at 60 days, and returned to normal at 147 days. The pattern was similar bilaterally (Suneja et al. 2000a).

#### *4.2. EXCITATORY TRANSMITTERS AND CALCIUM-RELATED INFERIOR COLLICULUS PROTEINS AFTER COCHLEAR DAMAGE*

Changes in glutamate after unilateral cochleotomy in adult guinea pigs were assessed by measuring the number and/or activity of intracellular AMPA-type glutamate receptors. There was a small, significant decrease in AMPA receptor binding in contralateral IC 2 days post-lesion, a gradual increase to above average 60 days later, and a return to normal at 147 days. The ipsilateral pattern showed a significant decrease starting at 31 days, then followed the contralateral pattern. These changes were predominantly in the ICC and the ICD but not in LIC (Suneja et al. 2000b).

Given the established role of high calcium levels in excitotoxic neuronal cell death, a few studies have examined changes in calcium-related proteins; different effects have been reported for different proteins. Parvalbumin was unchanged from immediately after focal acoustic trauma to 30 days later (Abbott et al. 1999). After unilateral cochleotomy in adult rats, increased immunostaining was found for calbindin, which regulates intracellular calcium and may act as a cytoplasmic calcium buffer and thereby exert a neuroprotective effect. Changes were found only in the IC contralateral to the ablated cochlea, with immunoreactivity increasing from about 23 days post-lesion to a plateau only by 18 weeks (Förster and Illing 2000). Both cochleotomy and focal acoustic trauma also upregulate GAP 43, a growth and plasticity-associated phosphoprotein that binds to calmodulin and acts a substrate of protein kinase C, in IC neurons (Illing et al. 1997; Michler and Illing 2002). GAP 43 influences early synaptogenetic processes and decreases to near-zero levels with maturation; thus, GAP 43 expression after cochlear damage represents its trauma-induced reemergence

and occurs only from day 10 post-lesion and increases to day 30; the ipsilateral increase is greatest until day 30 when contralateral expression predominates. Immediately after acoustic trauma there is a reactive transient decrease in the phosphorylated form of the cyclic adenosine monophosphate (cAMP) response element binding protein (phospho-CREB) in ipsilateral IC and a contralateral increase that persists for up to 239 days and exceeded any subsequent ipsilateral changes. These changes in calcium-related proteins and in CREB are indicative of reactive changes but not plasticity, unless the latter is loosely defined as any change in the parameter under measurement (Michler and Illing 2003).

#### *4.3. CHANGES IN INFERIOR COLLICULUS PHYSIOLOGY AFTER COCHLEAR DAMAGE*

Focal cochlear damage, quantified by changes in the compound action potential (CAP) audiogram, was created with acoustic trauma (Popelář and Syka 1982; Syka and Popelář 1982) or by spiral ganglion cell lesions (Snyder et al. 2000; Snyder and Sinex 2002), or lesions to the base of the cochlear partition (Irvine et al. 2003). Several indices, from potentials evoked at the IC surface to the responses of single neurons, before and after lesions, were assessed (Salvi et al. 1992; Wang et al. 1996). Responses were recorded from multineuron clusters across the dorsolateral-to-ventromedial tonotopic IC axis before and after cochlear lesions at similar depths (Snyder et al. 2000). Chronic implanted IC electrodes were used to assess neural responses from multineuron clusters pre- and post-lesion (Snyder and Sinex 2002). Multineuron responses several months after cochlear lesions were compared to normal animals (Irvine et al. 2003). The results from these studies are presented below according to the type of IC response measured.

#### *4.4. CHANGES IN EVOKED POTENTIALS*

In the IC contralateral to a damaged cochlea, changes in evoked potential (EP) amplitudes and thresholds to stimulation of the undamaged ipsilateral cochlea and to stimulation of undamaged parts of the contralateral test cochlea were obtained, although not in the same study. After unilateral cochlear deactivation with an ototoxic drug, the IC ipsilateral to the intact ear was now as sensitive and responsive to stimulation as the contralateral IC, whereas normally the contralateral IC is more sensitive. The decreased laterality differences were specifically due to changes in the IC responsiveness ipsilateral to the intact cochlea and occurred with a lag: the difference persisted at 1 to 2 days post-lesion and decreased at 7 days (Popelář et al. 1994).

A second effect was elicited by acoustic trauma (Salvi et al. 1992). Thirty days post-trauma a focal permanent hearing loss was seen with a peak of a 20- to 30-dB loss in CAP thresholds, and EP amplitude enhancement to frequencies below the peak of the cochlear hearing loss occurred, even at nearby frequen-

cies with deficits resembling the peak loss. Enhanced EP amplitude occurred at threshold and at mid-to-high stimulation intensities (>50 dB SPL). At frequencies above and below that of peak cochlear hearing loss, there was a decrease in EP amplitude in the IC at 8 hours post-lesion but not after 1 hour. Deficits could occur at frequencies at which CAP amplitudes (and EP amplitudes from cochlear nucleus) were below normal, or when CAP amplitudes had recovered but the amplitude of the cochlear nucleus EP was still depressed. These results were interpreted as due to relatively rapid, but not instantaneous, changes local to the IC or at least proximal to the cochlear nucleus (Salvi et al. 1992).

In contrast to these EP enhancement effects, other studies (Popelář et al. 1987) found none in awake guinea pigs after white noise trauma, even when the trauma caused a permanent hearing loss. Instead, EP amplitude was depressed and gradually recovered. That study (Popelář et al. 1987) also found EP enhancements in the auditory cortex, but not in the IC, when acoustic trauma was administered when the animals were awake but not when anesthetized. The absence of EP enhancement in the IC cannot simply reflect the use of a click stimulus, as the enhancement was found in the auditory cortex in the same animals. No EP enhancement occurred in the IC after acute noise trauma although the decrease of EP amplitude with increasing tone burst duration was slower in the lesioned animals, an effect mimicked by application of a GABA<sub>A</sub> blocker in controls (Szczepaniak and Møller 1995).

#### *4.5. CHANGES IN THE RESPONSE PROPERTIES OF SINGLE NEURONS*

The reduced ipsilateral–contralateral differences after deactivation of one cochlea (Popelář et al., 1994) indicate recruitment of more IC neurons ipsilateral to the intact ear to the ipsilateral stimulus, possibly through a decreased inhibition on binaural input. In keeping with this hypothesis, increased IC neurons responsive to ipsilateral stimulation occur in adult cats with partial high-frequency contralateral cochlear damage (Snyder and Sinex 2002; Irvine et al. 2003) and in adult ferrets after unilateral cochleotomy (Moore et al. 1993). The change was larger after cochlear lesions causing broad high-frequency losses extending over the entire frequency range with 26% to 100% increase in ipsilateral responsive sites in IC (Irvine et al. 2003) than the changes ensuing after more focal damage and <50% increase (Snyder and Sinex 2002).

Any increase in EP amplitudes at frequencies below the peak of cochlear damage in a focally damaged cochlea (Salvi et al. 1992) could indicate a recruitment of more neurons, and a plausible mechanism for such recruitment is decreased monaurally acting inhibition evoked by stimulating a partially damaged cochlea. Discounting for the moment the varied effects of cochlear damage on GABA function in the IC noted earlier, a decrease in such function should affect the frequency-bandwidth of the response areas (tuning curves) because inhibition can shape this property of central auditory neurons (Vater et al. 1992;



Yang et al. 1992; Suga 1995; Palombi and Caspary 1996; Rajan 1998). Consistent with the idea that cochlear damage reduced inhibition in IC cells and could thereby unmask responses in them, 21 days after ototoxicity-induced cochlear deafness, there was decreased IC GABA release and fewer IC cells were inhibited by electrical stimulation of the contralateral cochlea (Bledsoe et al. 1995). Another study, using the same technique and duration of cochlear deafening (Nagase et al. 2000), found more Fos-immunoreactive cells activated by an intense electrical stimulus to the basal cochlea (and fewer activated by a near-threshold stimulus). These results predict that, after cochlear damage, there should be changes in the profile of the response area and/or other response properties of IC neurons that do not simply mimic the peripheral desensitization changes seen in auditory nerve responses. Analyses of response properties (Wang et al. 1996; Snyder et al. 2000; Snyder and Sinex 2002) found changes in response area profiles consistent with a decrease in inhibition after cochlear damage. Acoustic trauma at frequencies above the characteristic frequency (CF) of individual neurons created cochlear hearing losses above the high-frequency edge of the IC neuron's response area (Wang et al. 1996). In essence, rapidly after acoustic trauma there was addition of excitation to the low-frequency side of the response area in about 40% of cells, and this was more likely to occur in mustached bat IC cells whose response areas are shaped by GABAergic inhibition than in neurons whose response areas were less influenced or uninfluenced by GABA. Most often the change was manifest as decreased thresholds in the low-frequency tails of the neuron's response area and occasionally as a broadening of bandwidth along the low-frequency slope between the CF tip and the low-frequency tail of the response area, with no change in the neuron's CF (Yang et al. 1992; Fig. 19.5).

The posttrauma effects (Wang et al. 1996) resemble those seen on tuning curves after applying GABA<sub>A</sub> blockers to mustached bat IC neurons (Yang et al. 1992). Following the earlier model (Yang et al. 1992), tuning curve changes were interpreted as an unmasking of low-frequency excitatory inputs normally suppressed by inhibitory inputs tuned to a higher CF than that of the IC neuron under study (Wang et al. 1996). They postulated that the cochlear hearing loss at frequencies higher than the response area of an IC neuron under study, desensitized inhibitory neurons with CFs above that of the IC neuron, thereby unmasking lower threshold responses on the low-frequency side (mainly in the tail) of the IC cell's response area (Fig. 19.5).

Unmasking responses and changes in the response area profile also occurred after focal cochlear damage in IC neurons that previously had (Snyder and Sinex 2002), or would have had (Snyder et al. 2000), a CF at or near frequencies sustaining cochlear hearing losses. A withdrawal of lesion-induced afferent drive at the appropriate frequencies in the response area and addition of excitation at other frequencies, could shift the tuning curve CF. The rapidity of unmasking shows that the IC physiology can change soon after alteration in afferent outflow and argues that the effect cannot be a form of adaptive plasticity (Snyder et al. 2000).




Figure 19.5. Rapid unmasking in the CNIC. **(A)** Prototypical changes in the tuning curves of level-tolerant CNIC neurons after a brief high-frequency loud tone (H.f. exposure) at a frequency above the high-frequency boundary of the IC neuron's tuning curve. (From Wang et al. 1996.) *Pre* is the response area boundary before the loud sound. The two *Post* curves show schematically two common changes seen immediately after the loud tone. In *Post (i)* the major change was the "addition" of a low-frequency "tail" at high intensities, whereas in *Post (ii)* the low-frequency "tail" occurred at lower intensities. *Post (ii)* shows another feature sometimes seen: a broadened response area width toward low frequencies at intensities nearer threshold. There is no change at the characteristic frequency (CF). **(B)** Schematic representation of the model (Wang et al. 1996). The final response area (*Final RA*) of the level-tolerant IC neuron is modeled as consisting of excitatory and inhibitory inputs; for clarity, only one of each input is shown. Both inputs have the same general profile but the inhibitory input is tuned to a higher frequency. In the hatched regions, the inhibitory input has a lower threshold than the excitatory input, and is therefore modeled to be a stronger input ( $I > E$ ) at suprathreshold levels, thus suppressing excitatory input at these frequencies. In the area *Final RA*, excitation is stronger than excitation as is observed with extracellular recording. If a high-frequency loud tone exposure elevated the inhibitory input thresholds, and if this effect was greater on the low-frequency side, low-frequency excitation would be unmasked to produce the *Post* curves in **(A)**.

The physiological unmasking is consistent with a reduction in inhibition; however, studies of IC neurochemistry demonstrate that cochlear damage can decrease or increase indices of GABAergic inhibition (see Section 4.1). The difference does not appear to be methodological: focal acoustic trauma in a single neuron study (Wang et al. 1996) caused immediate physiological unmasking, probably from reduced inhibition, while similar trauma increased GAD levels across the entire IC (Abbott et al. 1999). Perhaps small decreases in inhibitory gain suffice to unmask physiological responses, but these may not be detectable neurochemically, and certainly physiological unmasking was not seen in all cases (Wang et al. 1996; Snyder and Sinex 2002), nor was the pattern of unmasking related to the locus of cochlear damage (Snyder and Sinex 2002).

Other studies (Popelář et al. 1978; Popelář and Syka 1982) found no unmasking of new components in IC neural tuning curves after acoustic trauma, but only threshold elevation which, like the data in studies of unmasking, were obtained in the same neurons before and after acoustic trauma.

Changes in the maximum firing rate of single IC neurons have also been reported after cochlear damage. Acoustic trauma, targeted to produce hearing losses at frequencies in the response area of IC neurons, found a few neurons with an increased maximum firing rate at the CF (Lonsbury-Martin and Martin 1981). Even with acoustic trauma targeted at frequencies above the tuning curve of IC neurons, 70% increased their maximum response rate at CF, possibly through decreased suprathreshold inhibition. Because the same study reported response area changes in about 40% of neurons after acoustic trauma, in at least 30% of neurons, GABA-mediated IC inhibition does not play a significant role in shaping the threshold response area (Wang et al. 1996), although it may have effects at high intensities (Yang et al. 1992; Rajan 2001).

#### *4.6. RELATIONSHIP OF SINGLE-UNIT CHANGES TO EVOKED POTENTIAL SHIFTS*

Does the unmasking of new components in single neuron response areas account for the EP amplitude enhancement after cochlear trauma? Prior work did not provide a definitive answer because the principal unmasking took place at tail frequencies in neurons with CF below the frequency range with hearing loss (Wang et al. 1996). Because EP enhancement occurs (Wang et al. 1996) even for frequencies with significant hearing loss (frequencies near, but below, the peak hearing loss), the unmasking effects (Wang et al. 1996) are at frequencies too far from the damaged frequencies in the EP studies to provide the single-neuron basis for EP enhancement. Unmasking provides a more likely candidate, as it was found in neurons with CF at a frequency with hearing loss. Further, unmasking could often be at low frequencies very near to the pre-lesion CF of the IC neuron (Snyder and Sinex 2002).

Despite this, unmasking of new parts of the response area is unlikely to account for the EP amplitude enhancement because it does not occur as reliably or predictably as does the EP amplitude enhancement (Salvi et al. 1992). The

other factor likely to play a role—and probably a more important role since it appears to be a more common effect than unmasking in the response area profile—is the increase in IC neural maximum firing rates after cochlear damage (Salvi et al. 1992; Snyder and Sinex 2002).

#### 4.7. COCHLEOTOPIC MAP CHANGES AFTER COCHLEAR DAMAGE

This issue has been examined using paradigms described earlier to create focal cochlear hearing losses, assessed subsequently by changes in the CF progression across the dorsolateral-to-ventromedial tonotopic IC axis, examined by multi-neuron mapping before, and within hours after, a small focal lesion in the basal cochlea. Post-lesion there was a normal CF progression in dorsal IC regions receiving CF input from cochlear frequency regions below the lesion-affected region. Further ventrally there were discontinuities in the CF progression within and adjacent to the region receiving CF input from cochlear frequencies with hearing losses (Snyder et al. 2000). The term *lesion projection zone* (LPZ) denominates the region of a CNS structure that once received input from a receptor region that is now lesioned (Schmid et al. 1996). Despite the awkwardness of the term, because a lesion cannot project anywhere, it suffices as a short and easily remembered descriptor of such a region.

Two types of discontinuities were observed (Fig. 19.6). In one type, starting near the ventral edge of the LPZ, successive points had CFs at the low-frequency edge of the cochlear lesion, indicating an expanded CF map from the low-frequency margin of the cochlear lesion. Next, a succession of points occurred with CF at a frequency at the high-frequency edge of the cochlear lesion, representing an expanded CF map at the high-frequency margin of the cochlear lesion. Further ventrally in IC, there was a normal CF progression. The second type of discontinuity had an LPZ with an expanded CF frequency map at the low-frequency edge of the cochlear lesion and this enlarged map persisted throughout the depth of the IC even when exiting the LPZ and entering regions that, pre-lesion, had CF at frequencies without hearing losses. Thresholds at the new CFs in the expanded CF representations (in the map discontinuities) resembled, and were even more sensitive than, thresholds at the same CF(s) in neurons in the normal parts of the map.

Others have concluded (Snyder et al. 2000) that the rapid changes embody unmasking of suppressed inputs (see Section 4.5) and cannot represent adaptive use-dependent plasticity consequent on sensory deprivation. In the mapping study unmasking resulted in more homogeneous CF representation in the LPZ than would be predicted (see Section 4.4) from the wide scattering of CF with unmasking (Snyder and Sinex 2002). This may relate to the fact that the latter study produced much larger CAP losses (up to 60 dB); it would be interesting to determine whether rapid map changes of the sort seen in the first study, with small CAP losses, would have occurred in the second study.

This issue is particularly germane because further study found that only rarely could an expanded CF map of a lesion-edge frequency be interpreted as not




Figure 19.6. Changes in the frequency gradient curves (Snyder et al. 2000) in the CNIC after cochlear lesions. **(A, B)** Schematics of the frequency progression in CNIC when an electrode passes from dorsolateral to ventromedial and characteristic frequencies (CFs) are recorded at regular intervals. The *light line* (*Normal* in **B**) is the normal CF progression across the CNIC frequency axis. The *dashed line* shows the CF progression seen after a cochlear lesion produced a focal hearing loss (shown in **C** as elevations in cochlear compound action potential thresholds). In **(A)**, two discontinuities in CF progression are observed, and in both CFs remain about the same for some distance. Discontinuities include an expanded CF at the low-frequency edge of the cochlear lesioned range, and a CF over-representation at the high-frequency edge. **(B)** A single CF discontinuity at the low-frequency edge of the cochlear lesioned range extended across the remainder of the IC depth.

simply the residue of pre-lesion inputs to the LPZ (Irvine et al. 2003). In this residue argument (Rajan et al. 1993; Rajan and Irvine 1998) thresholds should increase across the expanded CF map in the LPZ (Fig. 19.6). Twenty sequences of recordings in the IC of 8 chronically lesioned animals found that the residue argument accounted for effects in almost every case when CFs were defined from the late component of multilineuron responses to tones (time window from 35 ms after tone onset to the end of the 150-ms tone) (Irvine et al. 2003). CFs defined from the onset component (time window 5 to 35 ms after tone onset) were comparable to the 50-ms time window in other studies (Snyder and Sinex 2002; Snyder et al. 2000) for 50-ms duration tones. For two thirds of the sequences the effects were consistent with responses in the LPZ being simply the residue of pre-lesion responses at those sites. Reservations as to why even the remaining one third may not represent plasticity of the type seen in studies in auditory cortex in the same experimental paradigm led to the conclusion that, if IC tonotopic map plasticity occurs, it is rare, unrelated to intersubject variability, and limited to the early responses to tonal stimuli (Irvine et al. 2003).

## 5. SUMMARY AND CONCLUSIONS

Despite differences in techniques used to alter cochlear outflow and to record from the IC, and in physiological responses measured in the IC, similar conclusions follow from most such studies: the changes in IC physiology after altering cochlear outflow are best explained as a post-lesion expression of preexisting inputs, either through unmasking of previously suppressed inputs or as the residue of pre-lesion inputs. These studies suggest that there is little native capacity for plasticity in IC physiology in the adult mammal in response to large changes in cochlear outflow, at least in the limited context of the few experimental paradigms available. This conclusion is at odds with the effects seen in IC physiology with presbycusis, and with the dramatic changes in IC physiology seen after neonatal cochleotomy (Moore et al. 1993), or following large and extensive perinatal cochlear lesions with ototoxic drugs (Harrison et al. 1998). The plasticity in the latter two instances may reflect developmental phenomena, as massive cochlear lesions made with ototoxic drugs in adult life do not produce the IC plasticity seen after similar neonatal lesions (Harrison 2001).

## Abbreviations

AI	primary auditory cortex
AM	amplitude modulation of sound envelopes
AP	all pass
BF	best frequency
BP	band-pass
C57	C57Bl/6 mouse strain

cAMP	cyclic adenosine monophosphate
CAP	compound action potential
CBA	CBA mouse strain
CF	characteristic frequency
CNS	central nervous system
CR+	calretinin-stained cells
CREB	(phospho-CREB) cyclic adenosine monophosphate response element binding protein
EP	evoked potential
GABA	$\gamma$ -aminobutyric acid
GAD	glutamic acid decarboxylase
GAP 43	growth associated protein 43
HP	high pass
IC	inferior colliculus
ICC	central nucleus of inferior colliculus
LIC	lateral nucleus of the inferior colliculus
LPZ	lesion projection zone
LTP	long-term potentiation
LP	low-pass
MGT	minimum gap threshold
MTC	frequency-threshold tuning curve
MTF	modulation transfer function
NMDA	<i>N</i> -methyl-D-aspartate
SI <sub>m</sub>	synchronization index maximum

## Acknowledgments

This report was funded by USPHS/NIH Grants P01-AG9524 from the National Institute on Aging, P30-DC05409 from the National Institute on Deafness and Communication Disorders, and the International Center for Hearing and Speech Research, Rochester, NY, USA (R.D.F.), and by the National Health and Medical Research Council of Australia (R.R.).

## References

- Abbott SD, Hughes LF, Bauer CA, Salvi R, and Caspary DM (1999) Detection of glutamate decarboxylase isoforms in rat inferior colliculus following acoustic exposure. *Neuroscience* **93**:1375–1381.
- Bauer CA, Brozoski TJ, Holder TM, and Caspary DM (2000) Effects of chronic salicylate on GABAergic activity in rat inferior colliculus. *Hearing Research* **147**:175–182.
- Bledsoe SC Jr, Nagase S, Miller JM, and Altschuler RA (1995) Deafness-induced plasticity in the mature central auditory system. *NeuroReport* **7**:225–229.
- Buonomano DV and Merzenich MM (1998) Cortical plasticity: from synapses to maps. *Annual Review of Neuroscience* **21**:149–186.

- Calford MB and Tweedale R (1988) Immediate and chronic changes in responses of somatosensory cortex in adult flying-fox after digit amputation. *Nature* **332**:446–448.
- Calford MB and Tweedale R (1991) Acute changes in cutaneous receptive fields in primary somatosensory cortex after digit denervation in adult flying fox. *Journal of Neurophysiology* **65**:178–187.
- Caspary DM, Raza A, Armour BAL, Pippin J, and Arnerić SP (1990) Immunocytochemical and neurochemical evidence for age-related loss of GABA in the inferior colliculus: implications for neural presbycusis. *Journal of Neuroscience* **10**:2363–2372.
- Caspary DM, Milbrandt JC, and Helfert RH (1995) Central auditory aging: GABA changes in the inferior colliculus. *Experimental Gerontology* **30**:349–360.
- Caspary DM, Palombi PS, and Hughes L (2002) GABAergic inputs shape responses to amplitude modulated stimuli in the inferior colliculus. *Hearing Research* **168**:163–171.
- Förster CR and Illing R-B (2000) Plasticity of the auditory brainstem: cochleotomy-induced changes in calbindin-D28K expression in the rat. *Journal of Comparative Neurology* **416**:173–187.
- Frisina DR (2001a) Subcortical neural coding mechanisms for auditory temporal processing. *Hearing Research* **158**:1–27.
- Frisina RD (2001b) Possible neurochemical and neuroanatomical bases of age-related hearing loss—presbycusis. *Seminars In Hearing: Innovations in Aging Auditory Research* **22**:213–225.
- Frisina RD (2001c) Anatomical and neurochemical bases of presbycusis. In: Hof PR and Mobbs CV (eds). *Functional Neurobiology of Aging*. Academic Press, San Diego, pp. 531–547.
- Frisina DR and Frisina RD (1997) Speech recognition in noise and presbycusis: relations to possible neural sites. *Hearing Research* **106**:95–104.
- Frisina RD and Walton JP (2001a) Neuroanatomy of the mouse central auditory system. In: Willott JP (ed). *Handbook of Mouse Auditory Research: From Behavior to Molecular Biology*. CRC Press, Boca Raton, pp. 243–278.
- Frisina RD and Walton JP (2001b) Aging of the mouse central auditory system. In: Willott JP (ed). *Handbook of Mouse Auditory Research: From Behavior to Molecular Biology*. CRC Press, Boca Raton, pp. 339–379.
- Frisina RD, Walton JP, Lynch-Armour MA, and Byrd JD (1998) Inputs to a physiologically characterized region of the inferior colliculus of the young adult CBA mouse. *Hearing Research* **115**:61–81.
- Frisina DR, Frisina RD, Snell KB, Burkard R, Walton JP, and Ison JR (2001) Auditory temporal processing during aging. In: Hof PR and Mobbs CV (eds). *Functional Neurobiology of Aging*. Academic Press, San Diego, pp. 565–579.
- Habbicht H and Vater M (1996) A microiontophoretic study of acetylcholine effects in the inferior colliculus of horseshoe bats: implications for a modulatory role. *Brain Research* **724**:169–179.
- Harrison RV (2001) Age-related tonotopic map plasticity in the central auditory pathways. *Scandinavian Audiology Supplement* **53**:8–14.
- Harrison RV, Ibrahim D, and Mount RJ (1998) Plasticity of tonotopic maps in auditory midbrain following partial cochlear damage in the developing chinchilla. *Experimental Brain Research* **123**:449–460.
- Heffner RS, Koay G, and Heffner H (2001) Focus: sound-localization acuity changes with age in C57Bl/6j mice. In: Willott JP (ed). *Handbook of Mouse Auditory Research: From Behavior to Molecular Biology*. CRC Press, Boca Raton, pp. 31–35.



- Helfert RH, Sommer TJ, Meeks J, Hofstetter P, and Hughes LF (1999) Age-related synaptic changes in the central nucleus of the inferior colliculus of Fischer-344 rats. *Journal of Comparative Neurology* **406**:285–298.
- Hosomi H, Hirai H, Okoda Y, and Amatsu M (1995) Long-term potentiation of neurotransmission in the inferior colliculus of the rat. *Neuroscience Letters* **195**:175–178.
- Hurley LM, Thompson AM, and Pollak GD (2002) Serotonin in the inferior colliculus. *Hearing Research* **168**:1–11.
- Illing R-B, Horvath M, and Laszig R (1997) Plasticity of the auditory brainstem: effects of cochlear ablation on GAP-43 immunoreactivity in the rat. *Journal of Comparative Neurology* **382**:116–138.
- Irvine DRF, Rajan R, and Smith S (2003) Effects of restricted cochlear lesions in adult cats on the frequency organization of the inferior colliculus. *Journal of Comparative Neurology* **467**:354–374.
- Kazee AM and West NR (1999) Preservation of synapses on principal cells of the central nucleus of the inferior colliculus with aging in the CBA mouse. *Hearing Research* **133**:98–106.
- Kazee AM, Han LY, Spongr VP, Walton JP, Salvi RJ, and Flood DG (1995) Synaptic loss in the central nucleus of the inferior colliculus correlates with sensorineural hearing loss in the C57Bl/6 mouse model of presbycusis. *Hearing Research* **89**:109–120.
- Kelly JB and Zhang H (2002) Contribution of AMPA and NMDA receptors to excitatory responses in the inferior colliculus. *Hearing Research* **168**:35–42.
- Lonsbury-Martin BL and Martin GK (1981) Effects of moderately intense sound on auditory sensitivity in rhesus monkeys: behavioral and neural observations. *Journal of Neurophysiology* **46**:563–586.
- Ma CL, Kelly JB, and Wu SH (2002) AMPA and NMDA receptors mediate synaptic excitation in the rat's inferior colliculus. *Hearing Research* **168**:25–34.
- McFadden SL and Willott JF (1994a) Responses of inferior colliculus neurons in C57Bl/6j mice with and without sensorineural hearing loss: effects of changing the azimuthal location of a continuous noise masker on responses to contralateral tones. *Hearing Research* **78**:132–148.
- McFadden SL and Willott JF (1994b) Responses of inferior colliculus neurons in C57Bl/6j mice with and without sensorineural hearing loss: effects of changing the azimuthal location of an unmasked pure-tone stimulus. *Hearing Research* **78**:115–131.
- Michler SA and Illing R-B (2002) Acoustic trauma induces a reemergence of the growth- and plasticity-associated protein GAP-43 in the rat auditory brainstem. *Journal of Comparative Neurology* **451**:250–266.
- Michler SA and Illing R-B (2003) Molecular plasticity in the rat auditory brainstem: modulation of expression and distribution of phosphoserine, phospho-CREB and TrkB after noise. *Audiology and Neurootology* **8**:190–206.
- Milbrandt JC, Albin RL, and Caspary DM (1994) Age-related decrease in GABA<sub>B</sub> receptor binding in the Fischer 344 rat inferior colliculus. *Neurobiology of Aging* **15**:699–703.
- Milbrandt JC, Albin RL, Turgeon SM, and Caspary DM (1996) GABA<sub>A</sub> receptor binding in the aging rat inferior colliculus. *Neuroscience* **73**:449–458.
- Milbrandt JC, Holder TM, Wilson MC, Salvi RJ, and Caspary DM (2000) GAD levels and muscimol binding in rat inferior colliculus following acoustic trauma. *Hearing Research* **147**:251–260.

- Moore DR, King AJ, McAlpine D, Martin RL, and Hutchings ME (1993) Functional consequences of neonatal unilateral cochlear removal. *Progress in Brain Research* **97**: 127–133.
- Mossop JE, Wilson MJ, Caspary DM, and Moore DR (2000) Down-regulation of inhibition following unilateral deafening. *Hearing Research* **147**:183–187.
- Nagase S, Miller JM, Dupont J, Lim HH, Sato K, and Altschuler RA (2000) Changes in cochlear electrical stimulation induced Fos expression in the rat inferior colliculus following deafness. *Hearing Research* **147**:242–250.
- Palombi PS and Caspary DM (1996) GABA inputs control discharge rate primarily within frequency receptive fields of inferior colliculus neurons. *Journal of Neurophysiology* **75**:2211–2219.
- Pollak GM, Burger RM, Park TJ, Klug A, and Bauer EE (2002) Roles of inhibition for transforming binaural properties in the brainstem auditory system. *Hearing Research* **168**:60–78.
- Popelář J and Syka J (1982) Noise impairment in the guinea pig. II. Changes of single unit responses in the inferior colliculus. *Hearing Research* **8**:273–283.
- Popelář J, Syka J, and Ulehlova L (1978) Effect of high-intensity sound on cochlear microphonics and activity of inferior colliculus neurons in the guinea pig. *Archives of Otorhinolaryngology* **221**:115–122.
- Popelář J, Syka J, and Berndt H (1987) Effect of noise on auditory evoked responses in awake guinea pigs. *Hearing Research* **26**:239–247.
- Popelář J, Erre JP, Aran JM, and Cazals Y (1994) Plastic changes in ipsi-contralateral differences of auditory cortex and inferior colliculus evoked potentials after injury to one ear in the adult guinea pig. *Hearing Research* **72**:125–134.
- Rajan R (1998) Receptor organ damage causes loss of cortical surround inhibition without topographic map plasticity. *Nature Neuroscience* **1**:138–143.
- Rajan R (2001) Plasticity of excitation and inhibition in the receptive field of primary auditory cortical neurons after limited receptor organ damage. *Cerebral Cortex* **11**: 171–182.
- Rajan R and Irvine DRF (1998) Neuronal responses across cortical field A1 in plasticity induced by peripheral auditory damage. *Audiology and Neurootology* **3**:123–144.
- Rajan R, Irvine DRF, Wise LZ, and Heil P (1993) Effect of unilateral partial cochlear lesions in adult cats on the representation of lesioned and unlesioned cochleas in primary auditory cortex. *Journal of Comparative Neurology* **338**:17–49.
- Ryan AF, Axelsson GA, and Woolf NK (1992) Central auditory metabolic activity induced by intense noise exposure. *Hearing Research* **61**:24–30.
- Salvi RJ, Powers NL, Saunders SS, Boettcher F, and Clock AE (1992) Enhancement of evoked response amplitude and single unit activity after noise exposure. In: Dancer A, Henderson D, Salvi RJ, and Hamernik RP (eds). *Noise-Induced Hearing Loss*. Mosby Year Book, St Louis, pp. 156–171.
- Salvi RJ, Huang J, and Ding D (2000) Auditory plasticity and hyperactivity following cochlear damage. *Hearing Research* **147**:261–274.
- Schmid LM, Rosa MG, Calford MB, and Ambler JS (1996) Visuotopic reorganization in the primary visual cortex of adult cats following monocular and binocular retinal lesions. *Cerebral Cortex* **6**:388–405.
- Shaddock-Palombi P, Backoff PM, and Caspary DM (2001) Responses of young and aged rat inferior colliculus neurons to sinusoidally amplitude modulated stimuli. *Hearing Research* **153**:174–180.

- Snell KB and Frisina DR (2000) Relationships among age-related differences in gap detection and word recognition. *Journal of the Acoustical Society of America* **107**: 1615–1626.
- Snyder RL and Sinex DG (2002) Immediate changes in tuning of inferior colliculus neurons following acute lesions of cat spiral ganglion. *Journal of Neurophysiology* **87**: 434–452.
- Snyder RL, Sinex DG, McGee AD, and Walsh EW (2000) Acute spiral ganglion lesions change the tuning and tonotopic organization of cat inferior colliculus neurons. *Hearing Research* **147**:200–220.
- Suga N (1995) Sharpening of frequency tuning by inhibition in the central auditory system: tribute to Yasuji Katsuki. *Neuroscience Research* **21**:287–299.
- Suneja SK, Benson CG, and Potashner SJ (2000a) Glycine receptors in adult guinea pig brain stem auditory nuclei: regulation after unilateral cochlear ablation. *Experimental Neurology* **154**:473–488.
- Suneja SK, Potashner SJ, and Benson CG (2000b) AMPA receptor binding in adult guinea pig brain stem auditory nuclei after unilateral cochlear ablation. *Experimental Neurology* **165**:355–369.
- Suneja SK, Potashner SJ, and Benson CG (1998) Plastic changes in glycine and GABA release and uptake in adult brain stem auditory nuclei after unilateral middle ear ossicle removal and cochlea ablation. *Experimental Neurology* **151**:273–288.
- Syka J and Popelář J (1982) Noise impairment in the guinea pig. I. Changes in electrical evoked activity along the auditory pathway. *Hearing Research* **8**:263–272.
- Szczepaniak WS and Möller AR (1995) Evidence of decreased GABAergic influence on temporal integration in the inferior colliculus following acute noise exposure: a study of evoked potentials in the rat. *Neuroscience Letters* **196**:77–80.
- Vater M, Habbicht H, Kössl M, and Grothe B (1992) The functional role of GABA and glycine in monaural and binaural processing in the inferior colliculus of horseshoe bats. *Journal of Comparative Physiology A* **171**:541–553.
- Walton JP, Frisina RD, Ison JR, and O'Neill WE (1997) Neural correlates of behavioral gap detection in the inferior colliculus of the young CBA mouse. *Journal of Comparative Physiology A* **181**:61–176.
- Walton JP, Frisina RD, and O'Neill WE (1998) Age-related alteration in processing of temporal sound features in the auditory midbrain of the CBA mouse. *Journal of Neuroscience* **18**:2764–2776.
- Walton JP, Simon H, and Frisina RD (2002) Age-related alterations in the neural coding of envelope periodicities. *Journal of Neurophysiology* **88**:565–578.
- Wang J, Salvi RJ, and Powers N (1996) Plasticity of response properties of inferior colliculus neurons following acute cochlear damage. *Journal of Neurophysiology* **75**: 171–183.
- Willott JF (1984) Changes in frequency representation in the auditory system of mice with age-related hearing impairment. *Brain Research* **309**:159–162.
- Willott JF (1986) Effects of aging, hearing loss, and anatomical location on thresholds of inferior colliculus neurons in C57BL/6 and CBA mice. *Journal of Neurophysiology* **56**:391–408.
- Willott JF (1991a) Central physiological correlates of ageing and presbycusis in mice. *Acta Otolaryngologica* **476**:153–156.
- Willott JF (1991b) *Aging and the Auditory System: Anatomy, Physiology and Psychophysics*. Singular, San Diego.

- Wu SH, Ma CL, Sivaramakrishnan S, and Oliver DL (2002) Synaptic modification in neurons of the central nucleus of the inferior colliculus. *Hearing Research* **168**:43–54.
- Yang L, Pollak GD, and Resler C (1992) GABAergic circuits sharpen tuning curves and modify response properties in the mustache bat inferior colliculus. *Journal of Neurophysiology* **68**:1760–1774.
- Zettel ML, Frisina, RD, Haider S, and O'Neill WE (1997) Age-related changes in calbindin D-28k and calretinin immunoreactivity in the inferior colliculus of CBA/CaJ and C57Bl/6 mice. *Journal of Comparative Neurology* **386**:92–110.
- Zettel ML, O'Neill WE, Trang TT, and Frisina RD (2001) Early bilateral deafening prevents calretinin up-regulation in the dorsal cortex of the inferior colliculus of aged CBA/CaJ mice. *Hearing Research* **158**:131–138.
- Zhang Y and Wu SH (2000) Long-term potentiation in the inferior colliculus studied in rat brain slice. *Hearing Research* **147**:92–103.

# Chapter 20

## Hearing Loss and the Inferior Colliculus

JAMES F. WILLOTT

### 1. HEARING LOSS AND THE INFERIOR COLLICULUS

It is a given that peripheral hearing loss will dictate which sounds inferior colliculus (IC) neurons can respond to; if the ear does not respond to a sound, neither will the IC. Of interest, then, is how the loss of peripheral sensitivity affects responses to still-audible sounds, and the physiological, histological, metabolic, and neurochemical effects this has upon IC neurons. This chapter addresses these topics with regard to the types of hearing loss occurring in adolescents and adults. Related issues are the effects of cochlear damage on the IC during development (see Chapter 18) and in aging or after peripheral trauma (see Chapter 19), which complement this exposition and explore themes in which peripheral challenge elicits reorganization.

Because the IC does not receive direct input from the auditory nerve, effects of hearing loss on IC physiology and anatomy must be centrally mediated, a function of altered properties of lower-order auditory pathways afferent to the IC. Spatiotemporal patterns of auditory-evoked activity in these pathways, altered by hearing loss, likely affect IC neuron responses and perhaps trigger neural plasticity. Denervation and transneuronal changes resulting from cochlear pathology and hearing loss in adults can also have lasting effects on the cochlear nucleus (CN) and other subcollicular auditory nuclei, including shrinkage of neurons, neuropil reduction, changes in neurotransmitter activity, and other sequelae. These effects depend in turn on whether the hearing loss is sensorineural (and which tonotopic regions of the cochlea are damaged), acute noise-induced, unilateral, conductive, or bilaterally profound. From the perspective of the central auditory system, each type of hearing loss has different consequences. Sensorineural cochlear trauma eliminates and/or diminishes evoked activity in damaged auditory nerve fibers, whereas other “channels” may remain intact. Moreover, a population of deafferented CN neurons is deprived of synaptic input and trophic support, possibly causing degenerative or compensatory changes. Acute noise-induced hearing loss results in a sudden attenuation of auditory input from some or most of the cochlea; however, the transsynaptic effects on the CN are presumably minimal in the short term. Unilateral damage dramati-

cally alters the pattern of neural activity received by binaural neurons of the IC and elsewhere, while depriving one CN of nearly all ascending input. Conductive hearing loss is typically “flat” across frequencies and results in no direct denervation of CN neurons. Severe bilateral sensorineural damage eliminates auditory input to the CN and causes massive denervation without the relative sparing of certain channels seen in less severe sensorineural or unilateral loss. Effects on the IC of each of these types of hearing loss are now addressed.

## 2. SENSORINEURAL HEARING LOSS

Sensorineural cochlear damage causes most hearing disorders, including those associated with some diseases, genetic conditions, aging (presbycusis), ototoxicity, and noise trauma (Schuknecht 1974; Willott 1991). The basal cochlea is often vulnerable and compromised, resulting in high-frequency hearing loss, and this has been the focus of most of the work reviewed here. Of course, this means that high-frequency tonotopic components of auditory pathways afferent to the IC are essentially inactivated, whereas those supporting lower frequencies are not. This imbalance of afferent input has a number of important consequences falling under the rubric hearing-loss-induced (HLI) plasticity—a variety of changes in neural response properties and frequency organization. Several processes may be involved in HLI plasticity including “unmasking” of responses supported by the still-intact circuit components by the removal of inhibitory or occlusive inputs from the now-defunct components, or synaptogenesis whereby the functional axons sprout branches or up-regulate synaptic events at sites “vacated” by the quieted inputs (see Chapter 19).

Two basic types of animal models have been used to study the effects of sensorineural cochlear damage on the IC and other central auditory nuclei: animals in which circumscribed cochlear lesions have been induced by frequency-specific noise trauma or surgery and inbred strains of mice that exhibit specific genetic cochlear pathology. We focus on murine models because the forms and consequences of pathology are considered elsewhere in this volume (see Chapters 19 and 21).

### 2.1. MOUSE MODELS

Inbred mice of the strains C57BL/6J (B6) and DBA/2J (D2), in particular, have provided much information on the effects of hearing loss on the IC and other central auditory nuclei. Both strains undergo genetically determined, progressive sensorineural hearing loss (Ralls 1967; Mikaelian 1979; Henry and Chole 1980; Willott 1981, 1986; Hunter and Willott 1987; Li and Borg 1991; Erway et al. 1993; Parham et al. 1997; Willott et al. 1995, 1997). B6 mice have hearing equal to that of young adults at 2 months of age and show a gradual change in IC properties as high-frequency hearing loss slowly ensues over the next few months. D2 mice, however, have poor sensitivity to high frequencies (>16 kHz)

even at 3 weeks, when abnormal IC properties are already observed (Willott 1981). At 2 months of age, auditory brain stem response (ABR) tone thresholds of D2 mice are roughly equivalent to those of 12- to 16-month-old B6 mice. Tones >20 kHz are virtually inaudible and thresholds of lower frequencies are elevated as well. Thus, D2 mice provide an animal model for adolescent-onset, rapid hearing loss, whereas B6 mice have adult-onset, gradual hearing loss.

## 2.2. FREQUENCY MAPS

Tonotopic organization of the IC is greatly altered in both B6 and D2 strains (Willott 1981, 1984, 1986, 1996). In normal-hearing mice (as in other mammals), neurons in the dorsolateral IC have sensitive thresholds and vigorous suprathreshold responses to low-frequency sounds. At increasingly ventromedial locations the best frequencies increase. After basal cochlear degeneration has occurred in B6 mice, this is no longer possible because severe high-frequency hearing loss removes the normal input to ventrally located neurons. However, the ventral neurons are not silent; rather, their thresholds for low- and middle-frequency sounds become similar to those in more dorsal IC regions (Fig. 20.1). Thus, neurons throughout the IC tend to respond best to low–middle frequencies.




Figure 20.1. Prototypical tuning curves of B6 mouse IC neurons (Willott 1986). The tuning curve (*filled circles*) is typical of that seen in the ventral IC of a young, normal-hearing B6 mouse. The neuron's best frequency (BF) is about 32 kHz and its low-frequency "tail" has high thresholds for frequencies <10 kHz. The same tuning curve would look very different (*unfilled circles*) after high-frequency hearing loss has occurred due to expression of the *Ahl* gene. This neuron no longer responds to its normal BF because of sensorineural damage to the basal cochlea; however, thresholds for lower frequencies are now lower, and the experimentally altered BF is near 8 kHz. The 50-dB BF threshold is comparable to 8-kHz thresholds observed in lower-frequency (more dorsal) tonotopic regions of the IC.

Indeed, more neurons now respond to the lower frequencies than is the case in normal-hearing mice. In D2 mice, tonotopic organization is also absent at a young age in both the ventral cochlear nucleus (Willott et al. 1982) and IC (Willott and Turner 2000). Tuning curves are similar throughout the IC, responding best to middle frequencies, between 12 and 15 kHz. As in hearing impaired B6 mice, ventral IC neurons have lower thresholds for low frequencies than would normally be the case. The tonotopic alteration of frequency threshold maps is one aspect of HLI plasticity, and occurs in the IC of several species when cochlear lesions have been experimentally induced during adulthood (see Chapter 19).

### 2.3. SUPRATHRESHOLD RESPONSES

Suprathreshold responses of IC neurons were evaluated in normal-hearing young adult B6 mice and middle-aged B6 mice with high-frequency hearing loss (Willott et al. 1988b). Single units were given tones (2 to 64 kHz) and noise stimuli at intensity levels from 80 dB SPL, decreasing by 10-dB steps to threshold. A small age-related increase (2% to 11%) in “sluggish” neurons (driven poorly by sound) occurred, but most neurons were well driven by suprathreshold stimuli in middle-aged mice. Indeed, in the ventral IC of middle-aged B6 mice, robust suprathreshold discharges were evoked by lower frequencies than normal. Similar changes were not observed in normal-hearing, middle-aged CBA mice.

Rate-level (intensity) functions have various shapes including “flat” (minimally affected by increased intensity), “monotonic” (increasing discharge rate as intensity increases), and “nonmonotonic” (decreased discharge rate at some suprathreshold intensities) (Willott et al. 1977). Nonmonotonicity presumably results when inhibition is evoked by stimuli at higher SPLs. IC neurons of both B6 and D2 mice with hearing loss exhibit fewer nonmonotonic intensity functions (Willott 1981; Willott et al. 1988b).

A significant proportion of IC neurons in adolescent D2 mice exhibit hyperactivity in the form of after-discharges (ADs) seen in a post-stimulus time histogram (PSTH) that is representative of those for IC neurons of a normal (Fig. 20.2, left) and a 21-day-old D2 mouse (Fig. 20.2, right). Normally, neurons responding to tones in a sustained fashion cease firing when the tone ends (Fig. 20.2, left). ADs occur when the discharges continue beyond tone termination, sometimes with a brief offset pause (Fig. 20.2, right). Interestingly, D2 mice exhibit violent, often lethal, audiogenic seizures (AGS) at the age when ADs occur (21 to 23 days); by 28 days both ADs and AGS subside. AGS susceptibility is likely to result from hearing loss; moreover, the IC plays a key role in AGS (Fig. 20.2; see Chapter 21).

### 2.4. DIRECTIONAL HEARING

Free-field (binaural) stimuli have been used to evaluate the effects of sensori-neural hearing loss on azimuth functions (discharge rates evoked by tone bursts






Figure 20.2. Characteristic post-stimulus time histograms (PSTHs) of mouse IC neurons (Willott 1981). The PSTHs depict the number and temporal pattern of action potentials evoked by a 100-ms tone burst (*silhouette above box*) summed for 15 trials. The PSTH (*left side*) is a typical sustained response in a normal-hearing mouse: the neuron fires and continues to do so for the duration of the tone, with little adaptation after onset. When the tone ceases, so does the neuron's responding. The PSTH (*right side*) resembles that of many IC cells in adolescent D2 mice with high-frequency hearing loss: the sustained firing does not stop immediately when the stimulus ceases, but continues as after-discharges. These hyperexcitable IC responses are most prevalent when D2 mice are susceptible to AGS and disappear as hearing loss progresses in severity.

as a function of stimulus azimuth) of IC neurons of B6 mice aged 2, 7, and 12 months (McFadden and Willott 1994a). Irrespective of age, nearly all neurons in the central area of the IC were sensitive to the azimuth of a best frequency (BF) stimulus, as revealed by azimuth functions in which firing rates varied by more than 50% from maximum to minimum at one or more intensities (the criterion used to designate directional sensitivity of an azimuth function). Much like IC neurons in other species, stimuli at azimuths contralateral to the IC recording site evoked excitatory responses, whereas ipsilateral stimuli evoked few discharges or inhibited the neuron. However, in hearing-impaired 7- and 12-month-old mice, more IC neurons had their strongest excitatory response evoked by ipsilateral stimulation, azimuth function "borders" (separating angles evoking high vs. low discharge rates) were more likely to have the high rates being evoked by the more ipsilateral angle, and azimuth functions often met the criterion for direction sensitivity only minimally.

## 2.5. MASKING

A single-unit study evaluated the effects of hearing loss on IC responses to a signal (BF tone) in a fixed position, when a continuous broadband masker was presented from different azimuth angles (McFadden and Willott 1994b). For both young and middle-aged B6 mice, BF tone thresholds were significantly elevated in the presence of noise at the three locations employed ( $-90^\circ$ ,  $0^\circ$ , and  $+90^\circ$ ).

Separating the signal (fixed at  $-90^\circ$ ) and masker sources improved masked tone thresholds of young, but not middle-aged, mice with hearing loss. Whereas the differences between normal-hearing and hearing-impaired B6 mice might have been influenced by peripheral or central monaural effects, the data indicated that improvement in thresholds as signal and masker are separated depends on binaural interactions. This implies that altered binaural excitatory–inhibitory interactions (within the IC and/or elsewhere) could contribute to the changes in binaural masking changes described here.

## *2.6. COMPARISON OF INFERIOR COLLICULUS, AUDITORY CORTEX, AND LOWER BRAIN STEM*

The representation of frequency in the primary auditory cortex (AI) of B6 mice was mapped (Willott et al. 1993) and compared with that obtained from single- and multiple-unit tuning curves from the ventral cochlear nucleus (VCN) and dorsal cochlear nucleus (DCN) (Willott et al. 1991). Reorganization of frequency maps was rapid and striking in cortex, but did not occur in VCN. The sensitization of tuning curve tails, seen in high-frequency IC neurons (Fig. 20.1), was not observed in VCN neurons. Indeed, as thresholds in the tuning curve tails of ventral IC neurons decreased with age, thresholds of VCN tuning curve tails in comparable tonotopic regions became elevated. Although the loss of high-frequency portions of VCN tuning curves inevitably results in lowering the BFs of some high-frequency neurons, that is, a portion of tuning curve tail became the “new” BF, the new BF thresholds of VCN neurons were elevated. In other words, the shift in the high-frequency IC to lower BFs, while maintaining low thresholds, did not occur in the VCN. Thus, VCN neurons appear to exhibit “pseudoplasticity” (Kaltenbach et al. 1996). These studies suggest that HLI plasticity in the IC is intermediate between that of VCN and auditory cortex. Moreover, sensitization of tuning curve tails of IC neurons is not accounted for by similar changes occurring in auditory nerve fibers or VCN neurons that are being conveyed to the IC. Rather, central neural processes appear to be responsible for the changes in the IC.

## *2.7. INTERPRETATION*

Sensorineural hearing loss can have significant effects on the IC of adult mice. The imbalance of neural traffic in high- vs. low-frequency pathways probably contributes to altered tonotopic organization. For example, pathways served by healthy cochlear regions may “out-compete” pathways served by impaired regions for synaptic expression or synaptogenesis on IC neurons. Alternatively, impaired pathways may not engage inhibitory circuits that normally “mask” excitatory responses evoked by lower frequencies. Degenerative, compensatory, or complex transsynaptic changes in the cochlear nucleus, secondary to denervation, may also come into play. For example, (inhibitory) glycinergic neurons in the CN are degraded in older, hearing-impaired B6 mice (Willott et al. 1997),

and this could alter the activity in neural circuits projecting to the IC. Indeed, observations outlined here suggest that inhibition may be diminished in the IC. These changes include fewer nonmonotonic rate-intensity functions, weaker ipsilaterally evoked free field inhibition, and ADs and AGS in D2 mice. In addition, IC neurons of older B6 mice exhibit increased spontaneous activity, which might be expected if tonic inhibition of neurons were reduced (Willott et al. 1988b).

### 3. ACUTE NOISE-INDUCED HEARING LOSS

Acute exposure to sufficiently intense noise causes immediate elevation of thresholds, and these typically recover partially or completely. Acute effects on the IC are likely to be secondary to impaired cochlear function and resulting changes in input via the auditory nerve. Two topics not addressed here are chronic exposure to intense noise in workplace or military occupations which cause permanent sensorineural hearing loss akin to that discussed in Section 2.7, and exposures to tones or narrow-band noises to demonstrate plasticity (see Chapter 19). The effects on IC responses can be evaluated using a within-subjects design by recording auditory activity before and after noise exposure, or using a between-groups design comparing exposed and control groups.

#### 3.1. WITHIN-SUBJECT, BEFORE-AND-AFTER DESIGN

Several studies have obtained responses of IC neurons before and after threshold elevations were induced acutely with noise exposure (Willott and Lu 1982). They recorded PSTHs and tuning curves in normal-hearing mice, exposed them to a threshold-elevating noise, and recorded again from the same neuron (only one neuron per mouse was studied). The neuron responded to tones with a transient onset response: one or two discharges at the beginning of the tone (left) as shown in typical PSTHs (Fig. 20.3). After noise exposure, about one third of the neurons showed evoked discharges after the initial onset response (Fig. 20.3, right side). It was hypothesized that the inhibition which normally suppresses the sustained discharges was diminished by the hearing loss, resulting in the increased discharges. Subsequent studies of this kind have supported the notion that inhibition is reduced (Salvi et al. 2000; Wang et al. 2002). This is confirmed by experiments in which IC responses of chinchillas were studied before and after intense tone exposure designed to attenuate inputs to the neural circuits responsible for activating the inhibitory sidebands above BF. The noise exposure caused a widening of some tuning curves at high sound intensities but had little or no effect on the low-threshold tip of the tuning curves. Among the neurons with strongly nonmonotonic discharge rate level functions, 93% showed a significant increase in discharge rate. Pauser PSTHs often significantly decreased in the inhibitory pause duration and increased in the sustained discharge rate after exposure (Wang et al. 1996).




Figure 20.3. A typical acute effect of noise-induced hearing loss on the auditory response of a mouse IC neuron. (From Willott and Lu 1982.) The PSTH (*left side*) depicts a typical onset response in a normal-hearing mouse. Unlike the sustained response (Fig. 20.2, *left side*), the neuron fires once or twice at the 100-ms tone onset, then ceases firing for the duration of the tone. A likely mechanism for producing an onset response is inhibition of sustained discharges. Immediately after the mouse was exposed to a threshold-elevating noise, the same neuron now responds with a more sustained pattern (*right side*) when noise-induced hearing loss resulted in stronger excitation/weaker inhibition.

### 3.2. BETWEEN-GROUPS DESIGN

Studies comparing noise-exposed and control groups also provide evidence of diminished inhibition in association with noise-induced hearing loss. This was demonstrated by a significant decrease in glutamic acid decarboxylase immunoreactivity in the IC after exposure of adult rats to high-intensity sound for 10 hours. Complete recovery occurred by 30 days post-exposure. These data suggest that transient changes in  $\gamma$ -aminobutyric acid (GABA) neurotransmission occur in the IC of animals exposed to intense sound (Milbrandt et al. 2000).

Several studies produced noise-induced hearing loss in adolescent mice and monitored the IC responses over several days. This procedure is particularly interesting because such mice become susceptible to AGS within 3 to 5 days, a propensity dubbed as “acoustic priming” for AGS (Henry 1967; Chen and Willott 1983). Thus, the changes in IC responses have a behavioral correlate, as is the case for D2 mice mentioned earlier. Measures of auditory evoked potentials from the IC of adolescent mice, made one day after noise exposure, show thresholds were elevated and suprathreshold amplitudes were reduced (Fig. 20.4). By 4 days, however, evoked potential amplitudes were enhanced compared to controls (Willott and Henry 1974). Extracellular single-unit IC responses in D2 mice revealed abnormal ADs such as those seen in adolescents (Fig. 20.3), and prolonged bursts of action potentials to click stimuli. These studies suggest that, after a period of partial threshold recovery from noise exposure, some IC neu-




Figure 20.4. Auditory evoked potentials from the IC in three groups of adolescent mice. (From Willott and Henry 1974.) A large electrode recorded evoked potentials in response to noise pips of various SPLs. Three groups were used: nonexposed control (*filled circles*); mice exposed to an intense, threshold-elevating noise and tested 1 day later (*unfilled squares*) or 4 days later (*open circles*). In mice tested 1 day after noise exposure, thresholds were greatly elevated and evoked potential amplitudes were reduced from control values. Mice tested 4 days after noise exposure had less severe threshold elevations (now responding to 40-dB stimuli). Interestingly, evoked potential amplitudes were larger than in controls at higher SPLs.

rons become hyperexcitable (perhaps because of diminished inhibition), and this results in susceptibility to AGS (Urban and Willott 1979).

### 3.3. INTERPRETATION

Acute noise-exposed hearing loss appears to have some effects on responses of IC neurons resembling those of sensorineural cochlear damage. Correlates of threshold elevations such as the emergence of ADs, increased excitatory driving, and altered tuning curves suggest diminution of inhibitory influences on the IC, similar to sensorineural damage. Because acute noise-induced hearing loss presumably causes little or no central trauma, the response changes probably result from weaker auditory activation of inhibitory pathways by affected regions of the cochlea.

## 4. UNILATERAL DEAFNESS

Unilateral deafness can result from trauma or disease in the ear or auditory nerve. Given the importance of binaural input to the IC, this is likely to have significant effects. For example, an imbalance between left and right ear input

might trigger competition for the “functionally vacated” synapses, whereby input supported by the intact ear would prevail. Such effects are especially dramatic when the unilateral deafferentation occurs during early development (see Chapter 19). Neonatal unilateral hearing loss in various species leads to substantial changes in organization of bilateral projections to the IC and bilateral neural responses (Trune 1983; Kitzes and Semple 1985; Moore 1985; Moore et al. 1989).

Unilateral deafening during adulthood has consequences as well. For example, unilateral cochlear removals in adult ferrets led to a rapid increase in the proportion of neurons in the IC ipsilateral to the intact ear that were excited by acoustic stimulation of that ear. Responses in the IC contralateral to the intact ear were largely unchanged (McAlpine et al. 1997). In an activity-sensitive measure, *c-fos* mRNA expression using in situ hybridization was assessed in the adult rat brain following cochlear ablations. After unilateral ablation, acoustically driven *c-fos* expression was eliminated or decreased in the contralateral IC and elsewhere. Recovery was not evident even after 6-month survivals (Luo et al. 1999). A physiological study concluded that unilateral deafening reduced inhibition in the adult gerbil IC as shown by multiple-unit recordings made from electrode penetrations before and after contralateral cochlear ablation. Up to 60% increases in the proportion of recording sites at which neural activity was elicited by ipsilateral stimulation were observed after the ablation. Novel excitatory responses were evident within minutes of ablation. Western blotting for glutamic acid decarboxylase, the precursor to GABA, showed significant decreases in the IC contralateral to cochlear ablation, relative to those in the ipsilateral IC. Their results suggested decreased inhibitory influence of two types: a rapid, stimulus-related, functional unmasking and a delayed reduction in the capacity of GABA synthesis in the IC (Mossop et al. 2000).

## 5. CONDUCTIVE HEARING LOSS

Occlusion of the outer or middle ears causes threshold elevations across all frequencies, so the most obvious effect would be reduced auditory evoked activity in the IC (assuming there is not amplification of stimuli). Thus, in adults, one might not expect strong effects on IC neurons, such as those brought about by sensorineural or unilateral deafness. Indeed, no effect of conductive hearing loss initiated during young adulthood was found on IC neural somatic size neurons or in other auditory brain stem structures (Webster 1983). Conductive hearing loss decreased 2-deoxy-D-glucose (2-DG) uptake, an activity sensitive metabolic marker studied in the IC of adult gerbils (Tucci et al. 1999, 2001), whereas others found minimal effects (Woolf et al. 1983). No effect of unilateral conductive hearing loss on cytochrome oxidase staining was observed in the adult gerbil IC (Tucci et al. 2002). Even when a conduction deficit was caused by neonatal ossicle removal, there was no detectable difference in relative IC cytochrome oxidase activity (Paterson and Hosea 1993).

Might more interesting, subtle changes occur following conductive hearing loss? Perhaps the elevation of thresholds is not completely linear or differential attenuation of low and high frequencies or changes in resonant frequencies from reduced volume of the ear canal cause small changes in the relative activity related to tonopcity, albeit much less than those occurring with sensorineural damage to cochlear regions. Moreover, complex excitatory–inhibitory responses of IC neurons might be affected, as in this hypothetical example. Sound levels in the ambient acoustic environment range from 0 to 80 dB. An IC neuron with a “nonmonotonic” intensity function responds with excitation to sounds in the 0- to 40-dB SPL range and is inhibited from 40 to 80 dB. After a 40-dB conductive loss, 0- to 40-dB sounds are eliminated and the 40- to 80-dB sounds are effectively attenuated to 0 to 40 dB. This manipulation creates an IC neuron that can respond only with excitation.

## 6. COMPLETE BILATERAL HEARING LOSS

The effects of bilateral cochlea ablation or auditory nerve section obviously result in a loss of auditory function in the IC and throughout the central auditory system. Because responses to all frequencies in all monaural and binaural circuits are equally affected, bilateral deafening does not cause the imbalance and synaptic competition in different neural circuit components that follow sensorineural and unilateral hearing loss. Thus, it is not surprising that effects on the IC can be minimal. Ferrets reared with bilateral cochlear removals from before the normal hearing onset were injected with wheat germ agglutinin conjugated to horseradish peroxidase in one IC. Neither the absolute number nor the bilateral symmetry of the labeled neurons differed significantly from normal adult ferrets (Moore 1990). Severe bilateral hearing loss also occurs in B6 mice older than 18 months of age (see section 2). In aged B6 mice, IC labeling with either 2-DG (Willott et al. 1988a) or cytochrome oxidase (Willott 2001) showed little, if any, change. Retrograde transport of wheat germ agglutinin–horseradish peroxidase from IC to CN of B6 mice revealed no evidence of altered projections or change in the number of labeled neurons (Willott et al. 1985). Little effect was observed on the size or the packing density of IC neurons, and there was no age-related neuron loss (Willott et al. 1994a). Another study observed a small but significant decrease in the size of principal neurons in the central nucleus of older B6 mice (Kazee et al. 1995).

This is not to say that bilateral deafness has no effect on the IC, as a study on aged B6 mice found fewer synapses of all morphologic types on principal neuronal somas, and the percentage of somatic membrane covered by synapses decreased by 67% (Kazee et al. 1995). After bilateral deafness of 21 days in guinea pigs there was an increase in evoked *fos*-immunoreactive neurons in IC to contralateral cochlear electrical stimulation, a reduction in IC neurons responding with suppression of activity to electrical stimulation, and decreased GABA release assayed by in vivo microdialysis (Bledsoe et al. 1995). Taken

together, these studies suggest that severe hearing loss may affect synaptic density or organization and inhibition, but does not have dramatic effects on morphology or metabolism. This implies that a uniform reduction in activity is less injurious to IC cells than unilateral contralateral deafferentation; perhaps this reflects upper and lower limits on IC capacity for reorganization after partial and complete reductions in activity.

## 7. BEHAVIORAL CORRELATES OF INFERIOR COLLICULUS CHANGES: PREPULSE INHIBITION

An issue often overlooked is whether neural plasticity in the IC and elsewhere has significant effects on auditory perception. What, if any, are the effects of altered tonotopicity or suprathreshold excitability? The correlation between IC neural properties and AGS, mentioned earlier, strongly suggest a behavioral outcome of altered IC physiology. Another auditory behavior that is localized to the IC is prepulse inhibition (PPI), a normal behavior exhibited by humans and other species.

In PPI, an audible “prepulse” stimulus (S1) is presented approximately 100 ms before an intense, startle-eliciting stimulus (S2). Whether the S1 is a tone burst, a gap in background noise, or some other acoustic change, it ultimately activates other components of the PPI neural circuitry, including pathways that descend to the reticular formation and inhibit neurons that trigger the startle reflex (Davis 1984; Willott et al. 1994b; Carlson and Willott 1998; Koch 1999; Ison 2001). Activation of the PPI circuit inhibits the startle pathway for several hundred milliseconds, producing an “inhibited” startle reflex with a reduced amplitude. Studies on humans find that PPI magnitude reflects the behavioral salience of the prepulse tones (Hoffman and Ison 1980; Ison 2001).

The IC plays a crucial role in the processing of prepulses and their ability to inhibit the startle response circuit. For example, IC lesions reduce or abolish PPI, whereas electrical stimulation of the IC produces PPI (Leitner and Cohen 1985; Li et al. 1998a,b). It follows, therefore, that changes in the responses of IC neurons to tones have the potential to modulate PPI, when those tones are used as prepulses.

PPI can be expressed by the simple ratio of startle reflex amplitude when a prepulse has been presented (S1 to S2) to startle reflex amplitude without the prepulse (S2-only). A stronger PPI is indicated by a smaller ratio in this computation. Studies of B6 mice have shown that the PPI magnitude changes are correlated with hearing loss and HLI plasticity (Willott et al. 1994b; Willott and Carlson 1995; Carlson and Willott 1996). As high-frequency sensitivity declines in aging B6 mice, the salience of a 24-kHz S1 in PPI also wanes (Fig. 20.5). However, at 5 months of age enhanced PPI is produced by a 12-kHz S1, which has become over-represented in the IC because of HLI plasticity. As B6 mice age from 5 to 12 months and hearing loss extends to 12 kHz and above, the






Figure 20.5. Prepulse inhibition in B6 mice (Willott et al. 1994b) aged 1, 5, or 12 months. In normal-hearing 1-month-olds, PPI ranged between 50% and 67% for 70-dB SPL with S1 stimuli of 4, 12, and 24 kHz. At 5 months (when thresholds >20 kHz have become elevated), the 24 kHz S1 is less effective, whereas the 12- and 4-kHz S1s produce stronger PPI. At 12 months, after high-frequency hearing loss has progressed, the 24-kHz S1 is ineffective, the 12-kHz S1 is less effective than it was a 5 months, but the 4 kHz S1 is more effective. The salience of S1 tones parallels changes in IC neuron frequency responses associated with HLI plasticity. At 5 months, more IC neurons respond to 70-dB SPL 12-kHz tones than at 1 month (Fig. 20.1), and 12-kHz tones elicit stronger PPI. At 12 months of age, even more IC neurons respond to 70-dB 4-kHz tones because the changes in frequency representation continue to favor even lower frequencies and 4-kHz tones elicit stronger PPI. (From Willott 1984, 1986.)

improvement in lower-frequency PPI continues to develop (Fig. 20.5). With a 4-kHz S1, PPI is even better in 12-month-olds than in 1- or 5-month-olds. Age-related changes in PPI do not occur in CBA mice, which have minimal hearing loss. Because the IC “drives” the inhibiting modulation of the startle reflex, improved PPI parallels HLI plasticity for the middle and low frequencies.

## 8. DIMINISHED INHIBITION

Much of the evidence presented here suggests that hearing loss results in diminished inhibition in the IC. This may be especially marked when the balance between excitatory and inhibitory processes becomes biased against inhibition: inhibitory processes become impaired and/or excitatory processes become augmented as a function of cochlear damage/hearing loss. Such events may be involved in HLI plasticity, development of susceptibility to AGS, and strengthening of PPI.

Several hypotheses have been offered to explain diminished inhibition (Willott 1996). One possibility is that diminished inhibition occurs in the CN or other structures projecting to the IC. Another hypothesis is that inhibitory portions of neuronal response areas are selectively impaired because of the topographic pattern of cochlear damage (Willott 1981). For example, the response areas of many IC neurons have inhibitory subregions activated by high frequencies. A high-frequency deficit (with relatively intact middle-to-low-frequency hearing) from noise or genetic manipulation would have an especially deleterious effect on activation of these inhibitory regions. When intense or broad band stimuli are presented to normal-hearing animals, both excitatory and inhibitory regions would be engaged simultaneously, affecting discharge rates; however, the same stimuli would activate only the excitatory region in hearing-impaired mice because the inhibitory region would be “lost,” and discharge rates would increase. A variant of this hypothesis is that the neurons would be inhibited less than normally by high frequencies in the ambient acoustic environment, which could have long-term effects on the neurons’ inhibitory response properties (a use- or activity-dependent phenomenon). It is also feasible that altered auditory activity affects inhibitory circuits directly.

## 9. IMPLICATIONS

Whatever the mechanisms of diminished inhibition and/or HLI plasticity prove to be, it is clear that the organization and response properties of IC neurons and circuits are altered with hearing loss in adults. We have seen the likely consequences in the form of AGS and altered PPI in mice, and there is no reason to believe that effects of hearing loss on the IC in human adults would not affect auditory perception. The general effect of HLI plasticity/diminished inhibition is to enhance the strength of still-audible sounds. Whether this is advantageous for a person with hearing loss is not yet clear. If the enhanced response is to a signal, this could be positive; if, on the other hand, the response to background noise is enhanced, it may be deleterious. Thus, it seems vital to better understand the processes involved in hearing-loss-induced changes in IC responses as a prelude to developing interventions to modulate or manage them clinically.

## Abbreviations

ABR	auditory brain stem response
AD	after-discharge
AGS	audiogenic seizures
AI	primary auditory cortex
BF	best frequency
CN	cochlear nucleus

DCN	dorsal cochlear nucleus
2-DG	2-deoxyglucose
GABA	$\gamma$ -aminobutyric acid
HLI	hearing-loss-induced
IC	inferior colliculus
PPI	prepulse inhibition
PSTH	post-stimulus-time histogram
VCN	ventral cochlear nucleus

## Acknowledgment

This research was supported by NIH Grant R37 AG07554.

## References

- Bledsoe SC Jr, Nagase S, Miller JM, and Altschuler RA (1995) Deafness-induced plasticity in the mature central auditory system. *NeuroReport* **7**:225–229.
- Carlson SC and Willott JF (1996) The behavioral salience of tones as indicated by prepulse inhibition of the startle response: relationship to hearing loss and central neural plasticity in C57BL/6J mice. *Hearing Research* **99**:168–175.
- Carlson S and Willott JF (1998) Caudal pontine reticular formation of C57BL/6J mice: responses to startle stimuli, inhibition by tones, and plasticity. *Journal of Neurophysiology* **79**:2603–2614.
- Chen CS and Willott JF (1983) Developmental plasticity: acoustic priming for audiogenic seizures. In: Willott JF (ed). *The Auditory Psychobiology of the Mouse*. Charles C Thomas, Springfield, pp. 426–469.
- Davis M (1984) The mammalian startle response. In: Eaton RC (ed). *Neural Mechanisms of Startle Behavior*. Plenum Press, New York, pp. 287–351.
- Erway LC, Willott JF, Archer JR, and Harrison D (1993) Genetics of age-related hearing loss in mice: I. Inbred and F1 hybrid strains. *Hearing Research* **65**:125–132.
- Henry KR (1967) Audiogenic seizure susceptibility induced in C57BL/6J mice by prior auditory exposure. *Science* **158**:938–940.
- Henry KR and Chole RA (1980) Genotypic differences in behavioral, physiological and anatomical expressions of age-related hearing loss in the laboratory mouse. *Audiology* **1**:369–383.
- Hoffman HS and Ison JR (1980) Reflex modification in the domain of startle. I. Some empirical findings and their implications for how the nervous system processes sensory input. *Psychological Bulletin* **87**:175–189.
- Hunter KP and Willott JF (1987) Aging and the auditory brainstem response in mice with severe or minimal presbycusis. *Hearing Research* **30**:207–218.
- Ison JR (2001) The acoustic startle response: reflex elicitation and reflex modification by preliminary stimuli. In: Willott JF (ed). *Handbook of Mouse Auditory Research: From Behavior to Molecular Biology*. CRC Press, Boca Raton, pp. 59–82.
- Kaltenbach JA, Meleca RJ, and Falzarano PR (1996) Alterations in tonotopic map of the cochlear nucleus following cochlear damage. In: Salvi RJ, Henderson DH, Colletti V,

- and Fiorino F (eds). *Auditory Plasticity and Regeneration*. Thieme, New York, pp. 317–332.
- Kazee AM, Han LY, Spongr VP, Walton JP, Salvi RJ, and Flood DG (1995) Synaptic loss in the central nucleus of the inferior colliculus correlates with sensorineural hearing loss in the C57BL/6 mouse model of presbycusis. *Hearing Research* **89**:109–120.
- Kitzes LM and Semple MN (1985) Single-unit responses in the inferior colliculus: effects of neonatal unilateral cochlear ablation. *Journal of Neurophysiology* **53**:1483–1500.
- Koch M (1999) The neurobiology of startle. *Progress in Neurobiology* **59**:107–128.
- Leitner DS and Cohen ME (1985) Role of the inferior colliculus in the inhibition of acoustic startle in the rat. *Physiology & Behavior* **34**:65–70.
- Li HS and Borg E (1991) Age-related loss of auditory sensitivity in two mouse genotypes. *Acta Otolaryngologica* **111**:827–834.
- Li L, Priebe RP, and Yeomans JS (1998a) Prepulse inhibition of acoustic or trigeminal startle of rats by unilateral electrical stimulation of the inferior colliculus. *Behavioral Neuroscience* **112**:1187–1198.
- Li L, Korngut LM, Frost BJ, and Beninger RJ (1998b) Prepulse inhibition following lesions of the inferior colliculus: prepulse intensity functions. *Physiology & Behavior* **65**:133–139.
- Luo L, Ryan AF, and Saint Marie RL (1999) Cochlear ablation alters acoustically induced c-fos mRNA expression in the adult rat auditory brainstem. *Journal of Comparative Neurology* **404**:271–283.
- McAlpine D, Martin RL, Mossop JE, and Moore DR (1997) Response properties of neurons in the inferior colliculus of the monaurally deafened ferret to acoustic stimulation of the intact ear. *Journal of Neurophysiology* **78**:767–779.
- McFadden SL and Willott JF (1994a) Responses of inferior colliculus neurons in C57BL/6J mice with and without sensorineural hearing loss: effects of changing the azimuthal location of an unmasked pure-tone stimulus. *Hearing Research* **78**:115–131.
- McFadden SL and Willott JF (1994b) Responses of inferior colliculus neurons in C57BL/6J mice with and without sensorineural hearing loss: effects of changing the azimuthal location of a continuous noise masker on responses to contralateral tones. *Hearing Research* **78**:132–148.
- Mikaelian DO (1979) Development and degeneration of hearing in the C57/b16 mouse: relation of electrophysiologic responses from the round window and cochlear nucleus to cochlear anatomy and behavioral responses. *Laryngoscope* **89**:1–15.
- Milbrandt JC, Holder TM, Wilson MC, Salvi RJ, and Caspary DM (2000) GAD levels and muscimol binding in rat inferior colliculus following acoustic trauma. *Hearing Research* **147**:251–260.
- Moore DR (1985) Postnatal development of the mammalian central auditory system and the neural consequences of auditory deprivation. *Acta Otolaryngologica Supplement* **421**:19–30.
- Moore DR (1990) Auditory brainstem of the ferret: bilateral cochlear lesions in infancy do not affect the number of neurons projecting from the cochlear nucleus to the inferior colliculus. *Developmental Brain Research* **54**:125–130.
- Moore DR, Hutchings ME, King AJ, and Kowalchuk NE (1989) Auditory brain stem of the ferret: some effects of rearing with a unilateral ear plug on the cochlea, cochlear nucleus, and projections to the inferior colliculus. *Journal of Neuroscience* **9**:1213–1222.
- Mossop JE, Wilson MJ, Caspary DM, and Moore DR (2000) Down-regulation of inhibition following unilateral deafening. *Hearing Research* **147**:183–187.

- Parham K, Gerber A, McKowen A, Sun X-M, and Kim DO (1997) Distortion product otoacoustic emissions of the DBA/2J mouse with early onset progressive hearing loss. *Association for Research in Otolaryngology Abstracts* **20**:194.
- Paterson JA and Hosea EW (1993) Auditory behaviour and brainstem histochemistry in adult rats with characterized ear damage after neonatal ossicle ablation or cochlear disruption. *Behavioural Brain Research* **53**:73–89.
- Ralls K (1967) Auditory sensitivity in mice: *Peromyscus* and *Mus musculus*. *Animal Behavior* **15**:123–128.
- Salvi RJ, Wang J, and Ding D (2000) Auditory plasticity and hyperactivity following cochlear damage. *Hearing Research* **147**:261–274.
- Schuknecht HF (1974) *Pathology of the Ear*. Harvard University Press, Cambridge.
- Trune DR (1983) Influence of neonatal cochlear removal on the development of mouse cochlear nucleus. III. Its efferent projections to inferior colliculus. *Brain Research* **285**: 1–12.
- Tucci DL, Cant NB, and Durham D (1999) Conductive hearing loss results in a decrease in central auditory system activity in the young gerbil. *Laryngoscope* **109**:1359–1371.
- Tucci DL, Cant NB, and Durham D (2001) Effects of conductive hearing loss on gerbil central auditory system activity in silence. *Hearing Research* **155**:124–132.
- Tucci DL, Cant NB, and Durham D (2002) Conductive hearing loss results in changes in cytochrome oxidase activity in gerbil central auditory system. *Journal of the Association for Research in Otolaryngology* **3**:89–106.
- Urban GP and Willott JF (1979) Response properties of neurons in the inferior colliculi of mice made susceptible to audiogenic seizures by acoustic priming. *Experimental Neurology* **63**:229–243.
- Wang J, Salvi RJ, and Powers N (1996) Plasticity of response properties of inferior colliculus neurons following acute cochlear damage. *Journal of Neurophysiology* **75**: 171–183.
- Wang J, Ding D, and Salvi RJ (2002) Functional reorganization in chinchilla inferior colliculus associated with chronic and acute cochlear damage. *Hearing Research* **168**: 238–249.
- Webster DB (1983) Late onset of auditory deprivation does not affect brainstem auditory neuron soma size. *Hearing Research* **12**:145–147.
- Webster DB and Webster M (1979) Effects of neonatal conductive hearing loss on brain stem auditory nuclei. *Annals of Otolaryngology, Rhinology and Laryngology* **88**:684–688.
- Willott JF (1981) Comparison of response properties of inferior colliculus neurons of two inbred mouse strains differing in susceptibility to audiogenic seizures. *Journal of Neurophysiology* **45**:35–47.
- Willott JF (1984) Changes in frequency representation in the auditory system of mice with age-related hearing impairment. *Brain Research* **309**:159–162.
- Willott JF (1986) Effects of aging, hearing loss, and anatomical location on thresholds of inferior colliculus neurons in C57BL/6 and CBA mice. *Journal of Neurophysiology* **56**:391–408.
- Willott JF (1991) *Aging and the Auditory System: Anatomy, Physiology, and Psychophysics*. Singular, San Diego.
- Willott JF (1996) Auditory system plasticity in the adult C57BL/6J mouse. In: Salvi RJ, Henderson DH, Colletti V, and Fiorino F (eds.) *Auditory Plasticity and Regeneration*. Thieme, New York, pp. 297–316.
- Willott JF (2001) Diversity of the mouse central auditory system In: Willott JF (ed). *Handbook of Mouse Auditory Research: From Behavior to Molecular Biology*. CRC Press, Boca Raton, pp. 239–242.

- Willott JF and Carlson S (1995) Modification of the acoustic startle response in hearing-impaired C57BL/6J mice: prepulse augmentation and prolongation of prepulse inhibition. *Behavioral Neuroscience* **109**:396–403.
- Willott JF and Henry KR (1974) Auditory evoked potentials: developmental changes of threshold and amplitude following early acoustic trauma. *Journal of Comparative and Physiological Psychology* **86**:1–7.
- Willott JF and Lu S (1982) Noise-induced hearing loss can alter neural coding and increase excitability in the central nervous system. *Science* **216**:1331–1332.
- Willott JF and Turner JG (2000) Neural plasticity in the mouse inferior colliculus: relationship to hearing loss, augmented acoustic stimulation, and prepulse inhibition. *Hearing Research* **147**:275–281.
- Willott JF, Chalupa LM, and Henry KR (1977) Responses of single units of the inferior colliculus of the mouse (*Mus musculus*) as a function of tone intensity. *Experimental Brain Research* **28**:443–448.
- Willott JF, Demuth RM, Lu SM, and Van Bergem P (1982) Abnormal tonotopic organization in the ventral cochlear nucleus of the hearing-impaired DBA/2 mouse. *Neuroscience Letters* **34**:13–17.
- Willott JF, Pankow D, Hunter KP, and Kordyban M (1985) Projections from the anterior ventral cochlear nucleus to the central nucleus of the inferior colliculus in young and aging C57BL/6 mice. *Journal of Comparative Neurology* **237**:545–551.
- Willott JF, Jackson LM, and Hunter KP (1987) Morphometric study of the anteroventral cochlear nucleus of two mouse models of presbycusis. *Journal of Comparative Neurology* **260**:472–480.
- Willott JF, Hunter KP, and Coleman JR (1988a) Aging and presbycusis: effects on 2-deoxy-D-glucose uptake in the mouse auditory brainstem in quiet. *Experimental Neurology* **99**:615–621.
- Willott JF, Parham K, and Hunter KP (1988b) Response properties of inferior colliculus neurons in middle-aged C57BL/6J mice with presbycusis. *Hearing Research* **37**:15–28.
- Willott JF, Parham K, and Hunter KP (1991) Comparison of the auditory sensitivity of neurons in the cochlear nucleus and inferior colliculus of young and aging C57BL/6J and CBA/J mice. *Hearing Research* **53**:78–94.
- Willott JF, Aitkin LM, and McFadden SM (1993) Plasticity of auditory cortex associated with sensorineural hearing loss in adult C57BL/6J mice. *Journal of Comparative Neurology* **329**:402–411.
- Willott JF, Bross LS, and McFadden SL (1994a) Morphology of the inferior colliculus in C57BL/6J and CBA/J mice across the life span. *Neurobiology of Aging* **15**:175–183.
- Willott JF, Carlson S, and Chen H (1994b) Prepulse inhibition of the startle response in mice: relationship to hearing loss and auditory system plasticity. *Behavioral Neuroscience* **108**:703–713.
- Willott JF, Erway LC, Archer JR, and Harrison D (1995) Genetics of age-related hearing loss in mice: II. Strain differences and effects of caloric restriction on cochlear pathology and evoked response thresholds. *Hearing Research* **88**:143–155.
- Willott JF, Milbrandt JC, Bross LS, and Caspary DM (1997) Glycine immunoreactivity and receptor binding in the cochlear nucleus of C57BL/6J and CBA/CaJ mice: effects of cochlear impairment and aging. *Journal of Comparative Neurology* **385**:405–414.
- Wolf NK, Sharp FR, Davidson TM, and Ryan AF (1983) Cochlear and middle ear effects on metabolism in the central auditory pathway during silence: a 2-deoxyglucose study. *Brain Research* **274**:119–127.

# Chapter 21

## The Midbrain and Audiogenic Seizures

CARL L. FAINGOLD

### 1. SEIZURES AND THE INFERIOR COLLICULUS

High-intensity acoustic stimuli can elicit motor convulsions (audiogenic seizures, AGS) in several animal species and are common in rodents (Hohenboken and Nellhaus 1970; Jobe et al. 1973; Beck et al. 1987; Gil Verona et al. 1990). Extensive experimentation has established that the inferior colliculus (IC) plays the key role in AGS initiation (Millan et al. 1986; Browning 1994; Nehlig et al. 1995; Faingold et al. 1998; Faingold 1999, 2002, 2004; Garcia-Cairasco 2002). Many techniques demonstrate the primacy of IC in AGS, including *c-fos* expression (McCown et al. 1991; Snyder-Keller and Pierson 1992; Clough et al. 1997; Kwon and Pierson 1997), lesion (Browning 1994; Ribak et al. 1994), focal microinjection (Frye et al. 1983, 1986; Millan et al. 1986; Faingold et al. 1989b,c, 1998; Riaz and Faingold 1994), 2-deoxyglucose (Eckardt et al. 1986), electrical stimulation (Frye et al. 1986; McCown et al. 1995), and neurophysiological studies (Li 1997). High-intensity acoustic stimuli evoke abnormal neuronal firing increases in the IC central nucleus (ICc) of rats with genetic or induced AGS susceptibility (Faingold et al. 1986b; Faingold and Boersma Anderson 1991; Faingold and Riaz 1995). This enhanced output from the IC initiates AGS. Most of the evidence indicates the ICc is the most critical site within the IC, but the other IC subnuclei are also important network components (McCown et al. 1984; Chakravarty and Faingold 1996, 1997; Faingold 2002).

AGS studies in rats began in 1924 (Jobe 1981). Several rodent strains inherit AGS susceptibility, and susceptibility is inducible in rodents by several experimental treatments (Tecott et al. 1995; Ross et al. 2001; Faingold 2002). Seizure mechanisms in rats were extensively studied in inherited forms of AGS (Jobe et al. 1973, 1992; Faingold and Naritoku 1992; Nehlig et al. 1995; Faingold 1999; Garcia-Cairasco 2002). Sustained AGS susceptibility is also produced in young rodents by inducing a hearing deficit during the critical period for hearing development (Ross et al. 2001), through thyroid deficiency, intense acoustic stimulation (priming), and ototoxic drugs (Faingold 2002). Normal adult rodents can be rendered susceptible to AGS by chemical stimulation in the IC (Millan et al. 1986; Faingold 1999) or after sustained focal perfusion of agents that inhibit

neuronal firing (Yang et al. 2001). AGS-like seizures can be evoked by IC electrical stimulation (McCown et al. 1984, 1991, 1995; McCown and Breese 1993), and acoustic priming lowers the threshold for evoking electrically induced seizures (Sakamoto and Niki 2001).

## 2. HEARING LOSS AND AUDIOGENIC SEIZURE SUSCEPTIBILITY

Hearing loss is associated with susceptibility to AGS in many rodents, such as DBA/2 mice (Willott and Turner 2000). Two strains of genetically epilepsy-prone rats (GEPR-3s, GEPR-9s) exhibit elevated hearing thresholds (Faingold et al. 1986a,b, 1990; Penny et al. 1986). Not all genetic forms of AGS involve a hearing deficit. Black Swiss mice exhibit AGS susceptibility without hearing deficits (Misawa et al. 2002). Treatments producing significant hearing loss during the critical period for hearing development render rodents susceptible to AGS (Van Middlesworth and Norris 1980; Chen and Aberdeen 1981; Henry 1984; Pierson and Swann 1988; Kwon and Pierson 1997; Iida et al. 1998; Ross and Coleman 1999; Willott and Turner 2000; Sakamoto and Niki 2001; Kai and Niki 2002). The degree of hearing loss does not correlate well with the intensity of AGS in thyroid-deficient rats and GEPRs (Patrick and Faingold 1989; Faingold et al. 1990). When hearing deficits rise above a certain level, AGS severity declines (Faingold et al. 1990). Acute AGS forms, such as ethanol withdrawal, do not involve hearing deficits (Wecker and Ison 1984). In AGS forms with early-onset hearing loss, the neurotransmitter changes in IC may occur as compensatory mechanisms for decreased acoustic input during the critical period (Faingold 2002). IC neurotransmitter changes brought about by several different mechanisms play critical roles in drug-induced and withdrawal-induced AGS.

## 3. GENETICALLY BASED AUDIOGENIC SEIZURES

Genetically epilepsy-prone rats (GEPRs) were derived by selective breeding from the Sprague–Dawley strain (Reigel et al. 1986). The two substrains, GEPR-9 and GEPR-3, exhibit consistently different severities of convulsive behavior during AGS (Jobe et al. 1973, 1992; Faingold and Naritoku 1992; Faingold 1999; Raisinghani et al. 2003). AGS mechanisms in GEPRs and other AGS-susceptible rats have been investigated worldwide (Garcia-Cairasco et al. 1998; Simler et al. 1999; Yechikhov et al. 2001). Many mouse strains also exhibit AGS (Faingold 2002). Thus far, genetic abnormalities have been identified only in DBA/2 and Frings mice (Neumann and Collins 1991; Skradski et al. 2001). Several “gene-knockout” mouse strains also exhibit AGS susceptibility (Faingold 2002; Upton and Stratton 2003), but how these abnormalities produce AGS susceptibility is unknown.



Some forms of AGS consistently exhibit generalized clonic seizures (such as GEPR-3s), while others consistently exhibit tonic seizures (such as GEPR-9s) (Reigel et al. 1986; Marescaux et al. 1987). Many similarities in the AGS networks of these closely related seizure forms exist, particularly the critical role of the IC in seizure initiation. However, there are also surprising differences with regard to the involvement of the amygdala (Browning et al. 1985; Garcia-Cairasco et al. 1996; Hirsch et al. 1997; Faingold 1999; Feng and Faingold 2002; Yang and Faingold 2002; Raisinghani and Faingold 2003).

## 4. INDUCED AUDIOGENIC SEIZURES

Long-lasting AGS susceptibility can be induced in normal rodents, commonly in young rodents during the critical period. However, AGS susceptibility induced in mature rodents is typically short lived.

### 4.1. DEVELOPMENTALLY RELATED AUDIOGENIC SEIZURES

In young rodents, AGS susceptibility can result from a hearing deficit during the critical period within the first 3 weeks of life (Ross et al. 2001). Approaches that induce AGS during development include thyroid deficiency, intense acoustic stimulation (priming), and ototoxic drugs (Reid and Collins 1986; Ross et al. 2001; Sakamoto and Niki 2001).

### 4.2. AUDIOGENIC SEIZURES INDUCED IN MATURE ANIMALS

Transient AGS susceptibility can be induced in adult rodents without detectable hearing deficit, including a post-ischemic syndrome (Kawai et al. 1995) and by drug administration (Wada 1970; Chung and Johnson 1983; Bac et al. 1998). AGS also occurs during withdrawal after continued administration of depressant drugs (Faingold 2002). AGS that occur during ethanol withdrawal (ETX) is well studied (Faingold et al. 1998; Atkins et al. 2000). If ethanol intake is prolonged or intense, most species, including humans, will develop convulsive seizures. In rodents such seizures are triggered by acoustic stimuli (Majchrowicz 1975; Faingold and Riaz 1994; Faingold et al. 1998; Koob 2000; Veatch and Gonzalez 2000; Devaud and Chadda 2001; Yang et al. 2003).

In normal rats AGS susceptibility is induced by focal microinjection of agents that enhance excitation or reduce inhibition in IC. The same agents injected into other brain sites do not induce AGS susceptibility (Millan et al. 1986; Bagri et al. 1989, 1999; Faingold et al. 1992). AGS are a major manifestation of withdrawal after chronic administration of alcohol, barbiturates, or benzodiazepines in rodents (Faingold et al. 1998; Oh and Ho 1999). These agents enhance the inhibitory action of  $\gamma$ -aminobutyric acid (GABA), and withdrawal results in AGS susceptibility involving GABA<sub>A</sub> receptor (GABA<sub>A</sub>R) down-regulation and/or de-




Figure 21.1. Inhibitory effects of iontophoretic application of GABA and a benzodiazepine, flurazepam (FLUR) on the responses of an ICc neuron to acoustic stimuli and the interaction of the agents applied simultaneously. The drug time course appears in the upper panel. The PSTHs are shown below. (A, C, E) The control response before drug application. (B) FLUR (200 nA, 2 min) alone is relatively ineffective. (D) The effect of GABA (40 nA, 1 min). (F) Combined application of GABA and FLUR is superadditive. The line below (A) is the stimulus duration. *N* is the total number of action potentials in the post-stimulus time histogram (PSTH). Stimulus: 100-ms tone bursts, 60 dB contralateral, 10 kHz. CF threshold: 40 dB SPL. (From Faingold et al. 1991a,b, Lippincott Williams & Wilkins.)

sensitization, which diminishes the effectiveness of GABA-mediated inhibition (Bianchi and Macdonald 2002; Chang et al. 2002; Mozrzymas et al. 2003).

## 5. NEUROTRANSMITTER MECHANISMS OF AUDIOGENIC SEIZURES

Several forms of AGS are associated with neurotransmitter abnormalities, most prominently in IC. GABA- and glutamate (GLU)-mediated neurotransmission are implicated in AGS. Monoaminergic abnormalities (norepinephrine, 5-hydroxytryptamine, histamine) are also implicated in AGS (Onodera et al. 1992; Jobe et al. 1994; Yan et al. 1998; Feng and Faingold 2000; Feng et al. 2001), but the IC may not be the most critical site of involvement of these transmitters.

### 5.1. ROLE OF $\gamma$ -AMINO BUTYRIC ACID

Reduced GABA effectiveness is a critical AGS mechanism (Roberts et al. 1985; Faingold et al. 1986b, 1994a,b; Frye et al. 1986; Ribak et al. 1993). The GABA<sub>A</sub>R hyperpolarizes most neurons by fluxing chloride (Sieghart 2000; Kitterler et al. 2002). GABA plays a key role in acoustic processing in IC neurons (Faingold et al. 1989b) based on anatomical, neurochemical, neurophysiological, and pharmacological studies (Faingold 2002; see Chapter 9). Direct application of GABA and agents that alter the GABA receptor activation modify IC inhibition (Faingold et al. 1989a, 1991a,b, 2000; Wagner 1996; Hosomoi et al. 1997) (Fig. 21.1).

GABA mediates several forms of inhibition in ICc. Exogenous GABA-mediated inhibition and endogenous acoustically evoked inhibition are increased by agents that enhance the action of GABA. These include a GABA uptake inhibitor and a benzodiazepine (Fig. 21.1). Acoustically evoked GABA-mediated neural inhibition is blocked by a GABA<sub>A</sub>R antagonist in vivo (Faingold et al. 1989a, 1991a,b; Casseday et al. 2000; Jen and Zhang 2000). The acoustically evoked inhibitory responses of ICc neurons include binaural inhibition (Fig. 21.2), nonmonotonic rate intensity response, offset-inhibition (Fig. 21.3), and pauser responses (Caird 1991; Covey and Casseday 1999). These inhibitory response patterns are subserved, in part, by local GABAergic inhibitory circuits within the IC and exogenous GABAergic inhibitory projections to IC (Faingold et al. 1989a, 1991a,b, 1993a; Li and Kelly 1992; González-Hernández et al. 1996; Burger and Pollak 2001). Agents that enhance or block the action of GABA will increase or reduce, respectively, inhibition in ICc neurons (Faingold et al. 1989a, 1991a,b; Fuzessery and Hall 1996; Le Beau et al. 1996; Palombi and Caspary 1996; Jen et al. 2001).

An inherent or induced deficiency of GABA-mediated ICc inhibition is a key mechanism subserving AGS susceptibility (Faingold et al. 1986a; N'Gouemo et al. 1996; Faingold 1999; Ishida et al. 2002). GABA transport mechanisms are also abnormal in genetic forms of AGS (Faingold et al. 1994b; Akbar et al.




Figure 21.2. The effects of the GABA<sub>A</sub> antagonist bicuculline (*BIC*) on the responses of an ICc neuron to monaural and binaural acoustic stimuli at CF. For drug time course and PSTHs see Fig. 21.1. (A, D) The control response. Binaural inhibition reduced firing >40% (A vs. D). The darkened line at the bottom of histogram (A) shows stimulus duration. After *BIC* application (100 nA) the inhibitory effect of binaural stimulation is almost completely blocked (B vs. D), but the monaural effect at the same time point is nearly normal (E). (C) Recovery of the binaural inhibition after ending *BIC* application. *N* is the total number of action potentials in the PSTH. Stimulus: 100-ms tone bursts, 50 dB contralateral or bilateral, 19.9 kHz. CF threshold: 30 dB SPL. (From Faingold et al. 1991a,b, Lippincott Williams & Wilkins.)

1998). The effectiveness of both acoustically evoked GABA-mediated inhibition and exogenously applied GABA is significantly decreased in both induced and genetically based AGS (Faingold et al. 1986a; Faingold and Boersma Anderson 1991; Li et al. 1994; N'Gouemo et al. 1996). Reduced GABA effectiveness also occurs in other brain sites of GEPRs (Gould et al. 1991, 1995; Evans et al. 1994; Molnar et al. 2000). Application of GABA onto ICc neurons induces dose-related inhibition (Fig. 21.1) that is enhanced by agents that increase GABA<sub>A</sub>R activation and diminished by agents that block GABA<sub>A</sub>R. The effectiveness of iontophoretically applied GABA is reduced in GEPRs and ETX (Faingold et al. 1986a; N'Gouemo et al. 1996). The nonmonotonic rate-intensity function is mediated, in part, by GABA (Faingold et al. 1989a, 1991a,b), and this form of inhibition is diminished in GEPR-9s (Faingold and Boersma Anderson 1991). The failure of the nonmonotonic rate-intensity function commonly seen in GEPR IC neurons mimics the effect of iontophoretic application of bicuculline in normal IC neurons (Faingold 2002).

Binaural inhibition is mediated partly by the action of GABA, and its effectiveness is also diminished in GEPRs (Faingold et al. 1986a; Faingold 2002). When a tone burst is presented a few normal ICc neurons exhibit a response peak at the offset of the stimulus, but an elevated incidence of offset responses occurs in GEPR-9s (Faingold et al. 1986b). The offset response is also induced by iontophoretic application of a GABA<sub>A</sub>R antagonist in normal rats (Faingold et al. 1986a) (Fig. 21.3). The offset response also involves diminished GABA-mediated offset inhibition (Faingold et al. 1986a). The source of the offset inhibition is an ipsilateral GABAergic projection to ICc from the superior paraolivary nucleus (SPON) (Kulesza and Berrebi 2000; Berrebi and Saldaña 2001). SPON neurons respond to acoustic stimuli with excitation at stimulus offset (Kulesza et al. 2002). The GABA projection from SPON appears to mediate offset inhibition. Diminished GABA effectiveness, genetically or pharmacologically induced, contributes to the emergence of the offset peak. A defect in GABA-mediated tonic inhibition, arising in the dorsal nucleus of the lateral lemniscus, which normally suppresses the offset response in ICc neurons, could contribute to the altered response pattern (Faingold et al. 1993a; Burger and Pollak 2001). A form of GABA-mediated inhibition evoked by paired electrical pulses *in vitro* is also diminished in IC and several other brain sites in AGS-susceptible rats (Dalkara 1986; Li 1997; Evans et al. 2000).

Agents focally microinjected into ICc that increase GABA<sub>A</sub>R activation will block AGS in GEPR-9s or during ETX. Microinjection of GABA<sub>A</sub>R or GABA<sub>B</sub>R agonists blocks AGS. Blocking GABA uptake or breakdown, or enhancing GABA synthesis focally within ICc, also blocks AGS susceptibility (Frye et al. 1983, 1986; Faingold et al. 1994a,b; Riaz and Faingold 1994). GABA uptake is increased in GEPR-9s as compared to controls (Faingold et al. 1994b). These data support the critical importance of a defect in GABA-mediated inhibition in the IC for susceptibility to AGS in GEPR-9s and ETX. A transient susceptibility to AGS is induced in normal rats after focal microinjection of bicuculline (Millan et al. 1986). AGS susceptibility was induced in normal rats on withdrawal after




Figure 21.3. PSTHs showing the induction of the onset–offset response in ICc neurons. The responses to increasing sound intensity in the GEPR are shown in the top row. (A) At 60 dB an onset response pattern was observed. (B) At 80 dB the pattern becomes an onset–offset response, a result that occurred significantly more often in GEPR IC neurons. The bottom row shows the induction of an onset–offset response by *BIC* in a control IC neuron. (C) Before *BIC* only an onset response was seen. (D) After *BIC* (100 nA for 2 min) the response is onset–offset as above. Typical acoustic stimulus appears in the middle row. *N* is the number of action potentials/histogram. PSTH parameters: 50 stimulus presentations, 1-ms bin width. Stimulus parameters: 63 dB binaural, 6.4 kHz; threshold: 33 dB SPL (*top row*), 7.2 kHz; threshold: 30 dB SPL (*bottom row*). (From Yang et al. 2001, with permission from Elsevier.)

continuous activation of GABA receptors by GABA infusion in ICc for 7 days (Yang et al. 2001). GABA receptor desensitization and down-regulation occur after prolonged exposure to GABA, which reduces GABA-mediated inhibition (Russek et al. 2000). These findings further support diminished GABAergic function in the IC as a critical mechanism for AGS initiation.

Withdrawal after continued systemic treatment with agents that enhance the action of GABA induces seizures in normal rats. Systemic administration of ethanol, barbiturates, or benzodiazepines for sufficient periods results in AGS susceptibility on withdrawal due partly to GABA-related mechanisms (Faingold et al. 1998; Harris 1999; Grobin et al. 2000; Loh and Ball 2000; Davis and Wu 2001; Ueno et al. 2001). GABA<sub>A</sub>R activation is strongly implicated in the depressive effects of ethanol (Loh and Ball 2000; Davis and Wu 2001; Criswell et al. 2003; Roberto et al. 2003). The regionally selective effects of ethanol on GABA<sub>A</sub>Rs may be lost when the neurons are isolated from their normal in vivo neuronal network (Criswell et al. 2003). The inhibitory effect of GABA directly applied onto IC neurons in vivo is enhanced by systemically administered ethanol (Simson et al. 1991). GABA<sub>A</sub>R subunit alterations are associated with altered responses to GABA (Buck et al. 1991; Mhatre and Ticku 1992; Devaud et al. 1997; Mihic et al. 1997; Faingold et al. 1998; Loh and Ball 2000).

The importance of GABA in the IC in AGS mechanisms is confirmed further by the ability to block GABA-mediated inhibition with significantly lower doses of bicuculline in genetic and induced AGS-susceptible rats than in normal rats (Faingold et al. 1986a, 2000).

## 5.2. THE ROLE OF GLUTAMATE

Glutamate (GLU) is the major excitatory brain neurotransmitter (Hardingham and Bading 2003), and GLU receptors (GLURs) play a critical role in modulating AGS. GLUR antagonists block AGS (Croucher et al. 1984; Faingold et al. 1992; Chapman et al. 2001; De Sarro et al. 2003).

Excitatory neurotransmission is critical in the responses of IC neurons to sound, and considerable data support GLU as a candidate (Feliciano and Potashner 1995; Goldsmith et al. 1995; Suneja et al. 1995, 2000; Saint Marie 1996; Gaza and Ribak 1997; Caicedo and Eybalin 1999). Application of GLUR agonists excite, and antagonists suppress, respectively, IC neuronal firing (Faingold et al. 1989b, 1991a,b, 2000; Smith 1992; Wagner 1996; Li et al. 1999; Jen et al. 2001; Zhang and Kelly 2001, 2003; Goldstein-Daruech et al. 2002) (Fig. 21.4).

GLU and aspartate levels in IC and certain other sites of AGS-susceptible rodents are abnormal (Chapman et al. 1986; Ribak et al. 1988; Lasley 1991; Marianowski et al. 1995). AGS susceptibility results from microinjection of agents that enhance GLUR activation in the IC (Millan et al. 1986; Faingold et al. 1989c; Browning et al. 1991). AGS susceptibility is also induced on withdrawal after continued systemic treatment or focal infusion of GLUR antagonists (Faingold and Riaz 1995; Tsai and Coyle 1998; Dahchour and De Witte 1999; Yang et al. 2001). Significantly greater than normal doses of GLUR antagonists




Figure 21.4. The effects of NMDA and the NMDA receptor antagonist, 2-APV, and their interactive effects on the responses of an IC neuron to acoustic stimuli at CF. The *top graph* is the time course, and the bottom PSTHs show the effects on the cell's response pattern. (A) The control response. The line below (A–E) is stimulus duration. (B) The excitatory effect of NMDA (30 nA). (C) Recovery from drug effects. (D) The same dose of NMDA was repeated with simultaneous application of APV (80 nA), reducing the NMDA effect. *N* is the total number of action potentials in the PSTH. Stimulus parameters: 100-ms tone burst; 30 dB SPL contralateral; 36.0 kHz; 0 dB SPL threshold. (From Faingold et al. 1989, with permission from Elsevier.)



are needed to block GLU-mediated excitation in IC dorsal cortex (ICd) neuron (in vitro) from AGS-susceptible animals (Faingold et al. 2000). (See Fig. 21.5.)

Systemic or focal microinjection into IC of agents that reduce GLUR activation will block AGS in several types of AGS-susceptible rats (Faingold et al. 1988, 1991a,b, 1992, 1993b; Patrick and Faingold 1989; Riaz and Faingold 1994; Higashiyama et al. 1998; Raisinghani and Faingold 2003). Focal microinjection of agents acting on metabotropic GLURs also blocks AGS (Chapman et al. 2001; Yip et al. 2001).

Focal IC microinjection of *N*-methyl-D-aspartate (NMDA) induces temporary susceptibility to AGS in normal rats (Millan et al. 1986; Faingold et al. 1989c; Browning et al. 1991). AGS susceptibility in normal rats is induced on termination of continuous focal perfusion into IC for 1 week with a NMDA receptor antagonist, presumably due to upregulation of NMDA receptors (Yang et al. 2001).

## 6. CONCLUSIONS

IC neurons and the ICc play a critical role in AGS initiation. Chronic susceptibility to AGS is a common rodent neurological disorder that can be genetically mediated (GEPRs) or induced by a hearing deficit produced during the critical period. Transient AGS susceptibility is readily induced by withdrawal from ethanol and other depressant drugs. Most forms of chronic AGS susceptibility are associated with early-onset hearing deficits. Several neurotransmitter abnormalities that occur in the IC involve GABA and GLU. GABA, acting largely at GABA<sub>A</sub>Rs, plays a major role in various types of normal acoustically evoked inhibition in IC. The effectiveness of these forms of GABA<sub>A</sub>R-mediated inhibition is reduced in AGS, which can be blocked by enhancing the action of GABA. AGS susceptibility in normal rodents is also induced by decreases in GABA receptor function in IC, and AGS susceptibility in normal rats can be induced by reducing the action of GABA. GLU likely mediates normal IC excitatory neurotransmission, and evidence for increased GLU availability is seen in AGS. AGS susceptibility can be temporarily blocked by GLUR antagonists, while it can be induced in normal animals by enhancing GLUR activation in IC. AGS susceptibility also follows GLUR antagonist withdrawal after continuous IC perfusion. These changes in amino acid neurotransmitters in the IC are proposed to occur either as compensatory mechanisms for hearing loss in many AGS forms, or are induced transiently in adulthood, by treatments such as withdrawal after prolonged treatment with agents such as ethanol.

## 7. FUTURE DIRECTIONS

Many questions about the role of the IC in AGS remain unanswered. The genetic defect responsible for AGS susceptibility in GEPRs is unknown. In the two forms of mouse AGS in which the abnormal gene has been identified, the mo-




Figure 21.5. (A) Inhibition *in vitro* of ICd neurons' excitatory postsynaptic potentials (EPSPs) by bath application of the non-NMDA excitatory amino acid receptor antagonist, CNQX. CNQX blocks the EPSP during ethanol withdrawal (ETX) in the presence of a competitive NMDA receptor antagonist, AP5 (10 to 70  $\mu$ M), and bicuculline (0.1 to 30  $\mu$ M). Membrane potential was held at  $-90$  mV. (B) Concentration-response (area, mV  $\times$  ms) curves for CNQX. The curves for ETX vs. control were significantly different. All data are means  $\pm$  SEM. (From Faingold et al. 2000, with permission from Elsevier.)

lecular cascade by which the gene abnormality leads to AGS susceptibility is unknown. Understanding how the genetic defects lead to the abnormalities in neurotransmission seen in AGS remains a challenge. Finally, development of experimental approaches to reverse the genetic deficit or prevent its development in AGS-susceptible animals is a major future research goal. Results of these investigations may have important therapeutic implications for genetic diseases of the human nervous system.

## Abbreviations

AGS	audiogenic seizures
AMPA	$\alpha$ -3-hydroxy-5-methyl-4-isoxazolepropionic acid
AP7	DL-2-amino-7-phosphonoheptanoic acid
2-APV(AP5)	2-amino-5-phosphonovalerate
BIC	bicuculline
CF	characteristic frequency

CNQX	6-cyano-7-nitroquinoxaline-2,3-dione
CONT	microinjection control
EPSPs	excitatory postsynaptic potentials
ETX	ethanol withdrawal
FLUR	flurazepam
GABA	$\gamma$ -aminobutyric acid
GABA <sub>A</sub> R	GABA <sub>A</sub> receptor subtype
GEPRs	genetically epilepsy-prone rats
GEPR-3s	moderate seizure strain of genetically epilepsy-prone rats
GEPR-9s	severe seizure strain of genetically epilepsy-prone rats
GLU	glutamate
GLUR	glutamate receptors
IC	inferior colliculus
ICc	inferior colliculus, central nucleus
ICd	inferior colliculus, dorsal cortex
nA	nanoamperes
NMDA	<i>N</i> -methyl-D-aspartate
PSTH	post-stimulus time histogram
SPON	superior paraolivary nucleus

## Acknowledgments

This work was supported by grants from the NIH, NIAAA, NS21281, and AA11628.

## References

- Akbar MT, Rattray M, Williams RJ, Chong NW, and Meldrum BS (1998) Reduction of GABA and glutamate transporter messenger RNAs in the severe-seizure genetically epilepsy-prone rat. *Neuroscience* **85**:1235–1251.
- Atkins AL, Rustay NR, and Crabbe JC (2000) Anxiety and sensitivity to ethanol and pentobarbital in alcohol withdrawal seizure-prone and withdrawal seizure-resistant mice. *Alcoholism Clinical Experimental Research* **24**:1743–1749.
- Bac P, Maurois P, Dupont C, Pages N, Stables JP, Gressens P, Evrard P, and Vamecq J (1998) Magnesium deficiency-dependent audiogenic seizures (MDDASs) in adult mice: a nutritional model for discriminatory screening of anticonvulsant drugs and original assessment of neuroprotection properties. *Journal of Neuroscience* **18**:4363–4373.
- Bagri A, Tomaz C, Brandão ML, and Carrive P (1989) Increased susceptibility of delectencephalated rats to audiogenic seizures induced by microinjection of bicuculline into the inferior colliculus. *Brazilian Journal of Medical and Biological Research* **22**: 1361–1370.
- Bagri A, Di Scala G, and Sandner G (1999) Myoclonic and tonic seizures elicited by microinjection of cholinergic drugs into the inferior colliculus. *Thérapie* **54**:589–594.

- Beck MM, Brown-Borg HM, and Jones TA (1987) Peripheral and brainstem auditory function in paroxysmal (*px*) White Leghorn chicks. *Brain Research* **406**:93–98.
- Berrebi AS and Saldaña E (2001) The superior paraolivary nucleus is a major source of GABAergic projections to the inferior colliculus of rats. *International Symposium on the Central Auditory System Abstracts*. Salamanca, Spain, pp. 19.
- Bianchi MT and Macdonald RL (2002) Slow phases of GABA<sub>A</sub> receptor desensitization: structural determinants and possible relevance for synaptic function. *Journal of Physiology (London)* **544**:3–18.
- Browning RA (1994) Anatomy of generalized convulsive seizures. In: Malafosse A, Genten P, Hirsch E, Marescaux C, Broglin D, and Bernasconi R (eds). *Idiopathic Generalized Epilepsies: Clinical, Experimental and Genetic Aspects*. John Libbey, London, pp. 399–413.
- Browning RA, Nelson DK, Mogharreban N, Jobe PC, and Laird HE (1985) Effect of midbrain and pontine tegmental lesions on audiogenic seizures in genetically epilepsy-prone rats. *Epilepsia* **26**:175–183.
- Browning RA, Wang C, and Faingold CL (1991) Effect of norepinephrine depletion on audiogenic-like seizures elicited by microinfusion of an excitant amino acid into the inferior colliculus of normal rats. *Experimental Neurology* **112**:200–205.
- Buck KJ, Hahner L, Sikela J, and Harris RA (1991) Chronic ethanol treatment alters brain levels of gamma-aminobutyric acid<sub>A</sub> receptor subunit mRNAs: relationship to genetic differences in ethanol withdrawal seizure severity. *Journal of Neurochemistry* **57**:1452–1455.
- Burger RM and Pollak GD (2001) Reversible inactivation of the dorsal nucleus of the lateral lemniscus reveals its role in the processing of multiple sound sources in the inferior colliculus of bats. *Journal of Neuroscience* **21**:4830–4843.
- Caicedo A and Eybalin M (1999) Glutamate receptor phenotypes in the auditory brainstem and mid-brain of the developing rat. *European Journal of Neuroscience* **11**:51–74.
- Caird D (1991) Processing in the colliculi. In: Altschuler RA, Bobbin RP, Clopton BM, and Hoffman DW (eds). *Neurobiology of Hearing: The Central Auditory System*. Raven Press, New York, pp. 253–292.
- Casseday JH, Ehrlich D, and Covey E (2000) Neural measurement of sound duration: control by excitatory-inhibitory interactions in the inferior colliculus. *Journal of Neurophysiology* **84**:1475–1487.
- Chakravarty DN and Faingold CL (1996) Increased responsiveness and failure of habituation in neurons of the external nucleus of inferior colliculus associated with audiogenic seizures of the genetically epilepsy-prone rat. *Experimental Neurology* **141**:280–286.
- Chakravarty DN and Faingold CL (1997) Aberrant neuronal responsiveness in the genetically epilepsy-prone rat: acoustic responses and influences of the central nucleus upon the external nucleus of inferior colliculus. *Brain Research* **761**:263–270.
- Chang Y, Ghansah E, Chen Y, Ye J, Weiss DS, and Chang Y (2002) Desensitization mechanism of GABA receptors revealed by single oocyte binding and receptor function. *Journal of Neuroscience* **22**:7982–7990.
- Chapman AG, Halsey MJ, Hart GP, Luff NP, Meldrum BS, and Wardley-Smith B (1986) Regional amino acid concentration in the brains of rats exposed to high pressures. *Journal of Neurochemistry* **47**:314–317.
- Chapman AG, Talebi A, Yip PK, and Meldrum BS (2001) Anticonvulsant activity of a mGlu<sub>4α</sub> receptor selective agonist, (1*S*,3*R*,4*S*)-1-aminocyclopentane-1,2,4-tricarboxylic acid. *European Journal of Pharmacology* **424**:107–113.

- Chen CS and Aberdeen GC (1981) The sensitive period for induction of susceptibility to audiogenic seizures by kanamycin in mice. *Archives of Oto-rhinolaryngology* **232**: 215–220.
- Chung SH and Johnson MS (1983) Experimentally induced susceptibility to audiogenic seizure. *Experimental Neurology* **82**:89–107.
- Clough RW, Eells JB, Browning RA, and Jobe PC (1997) Seizures and proto-oncogene expression of fos in the brain of adult genetically epilepsy-prone rats. *Experimental Neurology* **146**:341–353.
- Covey E and Casseday JH (1999) Timing in the auditory system of the bat. *Annual Reviews of Physiology* **61**:457–476.
- Criswell HE, Ming Z, Griffith BL, and Breese GR (2003) Comparison of effect of ethanol on N-methyl-D-aspartate- and GABA-gated currents from acutely dissociated neurons: absence of regional differences in sensitivity to ethanol. *Journal of Pharmacology and Experimental Therapeutics* **304**:192–199.
- Croucher MJ, Meldrum BS, Jones AW, and Watkins JC (1984) gamma-D-Glutamylaminomethylsulphonic acid (GAMS), a kainate and quisqualate antagonist, prevents sound-induced seizures in DBA/2 mice. *Brain Research* **322**:111–114.
- Dahchour A and De Witte P (1999) Effect of repeated ethanol withdrawal on glutamate microdialysate in the hippocampus. *Alcoholism Clinical Experimental Research* **23**: 1698–1703.
- Dalkara T (1986) Nipecotic acid, an uptake blocker, prevents fading of the  $\gamma$ -aminobutyric acid effect. *Brain Research* **366**:314–319.
- Davis KM and Wu JY (2001) Role of glutamatergic and GABAergic systems in alcoholism. *Journal of Biomedical Sciences* **8**:7–19.
- De Sarro G, Ferreri G, Gareri P, Russo E, De Sarro A, Gitto R, and Chimirri A (2003) Comparative anticonvulsant activity of some 2,3-benzodiazepine derivatives in rodents. *Pharmacology Biochemistry and Behavior* **74**:595–602.
- Devaud LL and Chadda R (2001) Sex differences in rats in the development of and recovery from ethanol dependence assessed by changes in seizure susceptibility. *Alcoholism Clinical Experimental Research* **25**:1689–1696.
- Devaud LL, Fritschy JM, Sieghart W, and Morrow AL (1997) Bidirectional alterations of GABA<sub>A</sub> receptor subunit peptide levels in rat cortex during chronic ethanol consumption and withdrawal. *Journal of Neurochemistry* **69**:126–130.
- Eckardt MJ, Campbell GA, Marietta CA, Majchrowicz E, Wixon HN, and Weight FF (1986) Cerebral 2-deoxyglucose uptake in rats during ethanol withdrawal and post-withdrawal. *Brain Research* **366**:1–9.
- Evans MS, Viola-McCabe KE, Caspary DM, and Faingold CL (1994) Loss of synaptic inhibition during repetitive stimulation in genetically epilepsy-prone rats (GEPR). *Epilepsy Research* **18**:97–105.
- Evans MS, Li Y, and Faingold C (2000) Inferior colliculus intracellular response abnormalities in vitro associated with susceptibility to ethanol withdrawal seizures. *Alcoholism Clinical Experimental Research* **24**:1180–1186.
- Faingold CL (1999) Neuronal networks in the genetically epilepsy-prone rat. *Advances in Neurology* **79**:311–321.
- Faingold CL (2002) Role of GABA abnormalities in the inferior colliculus pathophysiology–audiogenic seizures. *Hearing Research* **168**:223–237.
- Faingold CL (2004) Emergent properties of CNS neuronal networks as targets for pharmacology: application to anticonvulsant drug action. *Progress in Neurobiology* **72**:55–85.

- Faingold CL and Boersma Anderson CA (1991) Loss of intensity-induced inhibition in inferior colliculus neurons leads to audiogenic seizure susceptibility in behaving genetically epilepsy-prone rats. *Experimental Neurology* **113**:354–363.
- Faingold CL and Naritoku DK (1992) The genetically epilepsy-prone rat: neuronal networks and actions of amino acid neurotransmitters. In: Faingold CL and Fromm GH (eds). *Drugs for Control of Epilepsy: Actions on Neuronal Networks Involved in Seizure Disorders*. CRC Press, Boca Raton, pp. 277–308.
- Faingold CL and Riaz A (1994) Increased responsiveness of pontine reticular formation neurons associated with audiogenic seizure susceptibility during ethanol withdrawal. *Brain Research* **663**:69–76.
- Faingold CL and Riaz A (1995) Ethanol withdrawal induces increased firing in inferior colliculus neurons associated with audiogenic seizure susceptibility. *Experimental Neurology* **132**:91–98.
- Faingold CL, Gehlbach G, and Caspary DM (1986a) Decreased effectiveness of GABA-mediated inhibition in the inferior colliculus of the genetically epilepsy-prone rat. *Experimental Neurology* **93**:145–159.
- Faingold CL, Travis MA, Gehlbach G, Hoffmann WE, Jobe PC, Laird HE, and Caspary DM (1986b) Neuronal response abnormalities in the inferior colliculus of the genetically epilepsy-prone rat. *Electroencephalography and Clinical Neurophysiology* **63**:296–305.
- Faingold CL, Millan MH, Boersma CA, and Meldrum BS (1988) Excitant amino acids and audiogenic seizures in the genetically epilepsy-prone rat. I. Afferent seizure initiation pathway. *Experimental Neurology* **99**:678–686.
- Faingold CL, Gehlbach G, and Caspary DM (1989a) On the role of GABA as an inhibitory neurotransmitter in inferior colliculus neurons: iontophoretic studies. *Brain Research* **500**:302–312.
- Faingold CL, Hoffmann WE, and Caspary DM (1989b) Effects of excitant amino acids on acoustic responses of inferior colliculus neurons. *Hearing Research* **40**:127–136.
- Faingold CL, Millan MH, Boersma Anderson CA, and Meldrum BS (1989c) Induction of audiogenic seizures in normal and genetically epilepsy-prone rats following focal microinjection of an excitant amino acid into reticular formation and auditory nuclei. *Epilepsy Research* **3**:199–205.
- Faingold CL, Walsh EJ, Maxwell JK, and Randall ME (1990) Audiogenic seizure severity and hearing deficits in the genetically epilepsy-prone rat. *Experimental Neurology* **108**:55–60.
- Faingold CL, Boersma Anderson CA, and Caspary DM (1991a) Involvement of GABA in acoustically-evoked inhibition in inferior colliculus neurons. *Hearing Research* **52**:201–216.
- Faingold CL, Gehlbach G, and Caspary DM (1991b) Functional pharmacology of inferior colliculus neurons. In: Altschuler RA, Bobbin RP, Clopton BM, and Hoffmann DW (eds). *Neurobiology of Hearing: The Central Auditory System*. Raven Press, New York, pp. 223–251.
- Faingold CL, Naritoku DK, Copley CA, Randall ME, Riaz A, Anderson CA, and Arnerić SP (1992) Glutamate in the inferior colliculus plays a critical role in audiogenic seizure initiation. *Epilepsy Research* **13**:95–105.
- Faingold CL, Anderson CA, and Randall ME (1993a) Stimulation or blockade of the dorsal nucleus of the lateral lemniscus alters binaural and tonic inhibition in contralateral inferior colliculus neurons. *Hearing Research* **69**:98–106.

- Faingold CL, Randall ME, Naritoku DK, and Boersma Anderson CA (1993b) Noncompetitive and competitive NMDA antagonists exert anticonvulsant effects by actions on different sites within the neuronal network for audiogenic seizures. *Experimental Neurology* **119**:198–204.
- Faingold CL, Randall ME, and Anderson CA (1994a) Blockade of GABA uptake with tiagabine inhibits audiogenic seizures and reduces neuronal firing in the inferior colliculus of the genetically epilepsy-prone rat. *Experimental Neurology* **126**:225–232.
- Faingold CL, Marcinczyk MJ, Casebeer DJ, Randall ME, Arneric SP, and Browning RA (1994b) GABA in the inferior colliculus plays a critical role in control of audiogenic seizures. *Brain Research* **640**:40–47.
- Faingold CL, N’Gouemo P, and Riaz A (1998) Ethanol and neurotransmitter interactions—from molecular to integrative effects. *Progress in Neurobiology* **55**:509–535.
- Faingold CL, Li Y, and Evans MS (2000) Decreased GABA and increased glutamate receptor-mediated activity on inferior colliculus neurons in vitro are associated with susceptibility to ethanol withdrawal seizures. *Brain Research* **868**:287–295.
- Feliciano M and Potashner SJ (1995) Evidence for a glutamatergic pathway from the guinea pig auditory cortex to the inferior colliculus. *Journal of Neurochemistry* **65**:1348–1357.
- Feng H-J and Faingold CL (2000) Modulation of audiogenic seizures by histamine and adenosine receptors in the inferior colliculus. *Experimental Neurology* **163**:264–270.
- Feng H-J and Faingold CL (2002) Repeated generalized audiogenic seizures induce plastic changes in acoustically evoked neuronal firing in the amygdala. *Brain Research* **932**:61–69.
- Feng H-J, Naritoku DK, Randall ME, and Faingold CL (2001) Modulation of audiogenically kindled seizures by  $\gamma$ -aminobutyric acid-related mechanisms in the amygdala. *Experimental Neurology* **172**:477–481.
- Frye GD, McCown TJ, and Breese GR (1983) Characterization of susceptibility to audiogenic seizures in ethanol-dependent rats after microinjection of  $\gamma$ -aminobutyric acid (GABA) agonists into the inferior colliculus, substantia nigra or medial septum. *Journal of Pharmacology and Experimental Therapeutics* **227**:663–670.
- Frye GD, McCown TJ, Breese GR, and Peterson SL (1986) GABAergic modulation of inferior colliculus excitability: role in the ethanol withdrawal audiogenic seizures. *Journal of Pharmacology and Experimental Therapeutics* **237**:478–485.
- Fuzessery ZM and Hall JC (1996) Role of GABA in shaping frequency tuning and creating FM sweep selectivity in the inferior colliculus. *Journal of Neurophysiology* **76**:1059–1073.
- Garcia-Cairasco N (2002) A critical review on the participation of inferior colliculus in acoustic-motor and acoustic-limbic networks involved in the expression of acute and kindled audiogenic seizures. *Hearing Research* **168**:208–222.
- Garcia-Cairasco N, Wakamatsu H, Oliveira JA, Gomes EL, Del Bel EA, and Mello LE (1996) Neuroethological and morphological (Neo-Timm staining) correlates of limbic recruitment during the development of audiogenic kindling in seizure susceptible Wistar rats. *Epilepsy Research* **26**:177–192.
- Garcia-Cairasco N, Oliveira JAC, Wakamatsu H, Bueno STC, and Guimaraes FS (1998) Reduced exploratory activity of audiogenic seizures (*sic*) susceptible (*sic*) Wistar rats. *Physiology & Behavior* **64**:671–674.
- Gaza WC and Ribak CE (1997) Immunocytochemical localization of AMPA receptors in the rat inferior colliculus. *Brain Research* **774**:175–183.

- Gil Verona JA, Gomez Carretero ME, Manso ME, Gomez BP, and Garcia AN (1990) Morphology of the posterior colliculus in epileptic hamsters. *Archives of Neurobiology (Madrid)* **53**:112–118.
- Goldsmith JD, Kujawa SG, McLaren JD, and Bledsoe SC Jr (1995) In vivo release of neuroactive amino acids from the inferior colliculus of the guinea pig using brain microdialysis. *Hearing Research* **83**:80–88.
- Goldstein-Daruech N, Pedemonte M, Inderkum A, and Velluti RA (2002) Effects of excitatory amino acid antagonists on the activity of inferior colliculus neurons during sleep and wakefulness. *Hearing Research* **168**:174–180.
- González-Hernández T, Mantolán-Sarmiento B, González-González B, and Pérez-González H (1996) Sources of GABAergic input to the inferior colliculus of the rat. *Journal of Comparative Neurology* **372**:309–326.
- Gould EM, Craig CR, Fleming WW, and Taylor DA (1991) Sensitivity of cerebellar Purkinje neurons to neurotransmitters in genetically epileptic rats. *Journal of Pharmacology and Experimental Therapeutics* **259**:1008–1012.
- Gould EM, Curto KA, Craig CR, Fleming WW, and Taylor DA (1995) The role of GABA<sub>A</sub> receptors in the subsensitivity of Purkinje neurons to GABA in genetic epilepsy prone rats. *Brain Research* **698**:62–68.
- Grobin AC, Papadeas ST, and Morrow AL (2000) Regional variations in the effects of chronic ethanol administration on GABA<sub>A</sub> receptor expression: potential mechanisms. *Neurochemistry International* **37**:453–461.
- Hardingham GE and Bading H (2003) The Yin and Yang of NMDA receptor signalling. *Trends in Neurosciences* **26**:81–89.
- Harris RA (1999) Ethanol actions on multiple ion channels: which are important? *Alcoholism Clinical Experimental Research* **23**:1563–1570.
- Henry KR (1984) Noise and the young mouse: genotype modifies the sensitive period for effects on cochlear physiology and audiogenic seizures. *Behavioral Neuroscience* **98**:1073–1082.
- Higashiyama A, Ishida N, Nishimura T, Yasuda S, Kuroda Y, McEwen BS, and Kato N (1998) NMDA receptors in the inferior colliculus are critically involved in audiogenic seizures in the adult rats with neonatal hypothyroidism. *Experimental Neurology* **153**:94–101.
- Hirsch E, Danober L, Simler S, Pereira de Vasconcelos A, Maton B, Nehlig A, Marescaux C, and Vergnes M (1997) The amygdala is critical for seizure propagation from brainstem to forebrain. *Neuroscience* **77**:975–984.
- Hohenboken WD and Nellhaus G (1970) Inheritance of audiogenic seizures in the rabbit. *Journal of Heredity* **61**:107–112.
- Hosomi H, Mori M, Amatsu M, and Okada Y (1997) GABA-activated conductance in cultured rat inferior colliculus neurons. *Journal of Neurophysiology* **77**:994–1002.
- Iida K, Sasa M, Serikawa T, Noda A, Ishihara K, Akimitsu T, Hanaya R, Arita K, and Kurisu K (1998) Induction of convulsive seizures by acoustic priming in a new genetically defined model of epilepsy (Noda epileptic rat: NER). *Epilepsy Research* **30**:115–126.
- Ishida Y, Nakahara D, Hashiguchi H, Nakamura M, Ebihara K, Takeda R, Nishimori T, and Niki H (2002) Fos expression in GABAergic cells and cells immunopositive for NMDA receptors in the inferior and superior colliculi following audiogenic seizures in rats. *Synapse* **46**:100–107.
- Jen PH-S and Zhang J (2000) The role of GABAergic inhibition on direction-dependent sharpening of frequency tuning in bat inferior collicular neurons. *Brain Research* **862**:127–137.



- Jen PH-S, Sun X, and Chen QC (2001) An electrophysiological study of neural pathways for corticofugally inhibited neurons in the central nucleus of the inferior colliculus of the big brown bat, *Eptesicus fuscus*. *Experimental Brain Research* **137**:292–302.
- Jobe PC (1981) Pharmacology of audiogenic seizures. In: Brown RD (ed). *Pharmacology of Hearing: Experimental Clinical Bases*. John Wiley & Sons, New York, pp. 271–304.
- Jobe PC, Picchioni AL, and Chin L (1973) Role of brain norepinephrine in audiogenic seizure in the rat. *Journal of Pharmacology and Experimental Therapeutics* **184**: 1–10.
- Jobe PC, Mishra PK, and Dailey JW (1992) Genetically epilepsy-prone rats: actions of antiepileptic drugs and monoaminergic neurotransmitters. In: Faingold CL and Fromm GH (eds). *Drugs for Control of Epilepsy: Actions on Neuronal Networks Involved in Seizure Disorders*. CRC Press, Boca Raton, pp. 253–275.
- Jobe PC, Mishra PK, Browning RA, Wang C, Adams-Curtis LE, Ko KH, and Dailey JW (1994) Noradrenergic abnormalities in the genetically epilepsy-prone rat. *Brain Research Bulletin* **35**:493–504.
- Kai N and Niki H (2002) Altered tone-induced Fos expression in the mouse inferior colliculus after early exposure to intense noise. *Neuroscience Research* **44**:305–313.
- Kawai K, Penix LP, Kawahara N, Ruetzler CA, and Klatzo I (1995) Development of susceptibility to audiogenic seizures following cardiac arrest cerebral ischemia in rats. *Journal of Cerebral Blood Flow and Metabolism* **15**:248–258.
- Kittler JT, McAinsh K, and Moss SJ (2002) Mechanisms of GABA<sub>A</sub> receptor assembly and trafficking: implications for the modulation of inhibitory neurotransmission. *Molecular Neurobiology* **26**:251–268.
- Koob GF (2000) Animal models of craving for ethanol. *Addiction* **95** Supplement **2**:S73–S81.
- Kulesza RJ Jr and Berrebi AS (2000) Superior paraolivary nucleus of the rat is a GABAergic nucleus. *Journal of the Association for Research in Otolaryngology* **1**:255–269.
- Kulesza RJ, Spirou GA, and Berrebi AS (2002) Response properties of neurons in the superior paraolivary nucleus of the rat. *Association for Research in Otolaryngology Abstracts* **25**:111.
- Kwon J and Pierson M (1997) Fos-immunoreactive responses in inferior colliculi of rats with experimental audiogenic seizure susceptibility. *Epilepsy Research* **27**:89–99.
- Lasley SM (1991) Roles of neurotransmitter amino acids in seizure severity and experience in the genetically epilepsy-prone rat. *Brain Research* **560**:63–70.
- Le Beau FEN, Rees A, and Malmierca MS (1996) Contribution of GABA- and glycine-mediated inhibition to the monaural temporal response properties of neurons in the inferior colliculus. *Journal of Neurophysiology* **75**:902–919.
- Li Y (1997) In vitro electrophysiological properties of inferior colliculus cortex neurons in genetically epilepsy prone rats and rats undergoing ethanol withdrawal (seizures). *Dissertation Abstracts International* **59-02, Section B**:0548.
- Li L and Kelly JB (1992) Inhibitory influence of the dorsal nucleus of the lateral lemniscus on binaural responses in the rat's inferior colliculus. *Journal of Neuroscience* **12**:4530–4539.
- Li Y, Evans MS, and Faingold CL (1994) Inferior colliculus neuronal membrane and synaptic properties in genetically epilepsy-prone rats. *Brain Research* **660**:232–240.
- Li Y, Evans MS, and Faingold CL (1999) Synaptic response patterns of neurons in the cortex of rat inferior colliculus. *Hearing Research* **137**:15–28.

- Loh EW and Ball D (2000) Role of the GABA<sub>A</sub>  $\beta$ 2, GABA<sub>A</sub>  $\alpha$ 2, GABA<sub>A</sub>  $\alpha$ 1 and GABA<sub>A</sub>  $\gamma$ 2 receptor subunit genes cluster in drug responses and the development of alcohol dependence. *Neurochemistry International* **37**:413–423.
- Majchrowicz E (1975) Induction of physical dependence upon ethanol and the associated behavioral changes in rats. *Psychopharmacologia* **43**:245–254.
- Marescaux C, Vergnes M, Kiesmann M, Depaulis A, Micheletti G, and Warter JM (1987) Kindling of audiogenic seizures in Wistar rats: an EEG study. *Experimental Neurology* **97**:160–168.
- Marianowski R, Pollard H, Moreau J, Despres G, Ben Ari Y, Tran Ba HP, and Romand R (1995) *N*-methyl-D-aspartate receptor subunits NR1 and NR2C are over-expressed in the inferior colliculus of audiogenic mice. *Neuroscience Letters* **189**:190–194.
- McCown TJ and Breese GR (1993) A potential contribution to ethanol withdrawal kindling: reduced GABA function in the inferior collicular cortex. *Alcoholism Clinical Experimental Research* **17**:1290–1294.
- McCown TJ, Greenwood RS, Frye GD, and Breese GR (1984) Electrically elicited seizures from the inferior colliculus: a potential site for the genesis of epilepsy? *Experimental Neurology* **86**:527–542.
- McCown TJ, Duncan GE, and Breese GR (1991) Neuroanatomical characterization of inferior collicular seizure genesis: 2-deoxyglucose and stimulation mapping. *Brain Research* **567**:25–32.
- McCown TJ, Duncan GE, Johnson KB, and Breese GR (1995) Metabolic and functional mapping of the neural network subserving inferior collicular seizure generalization. *Brain Research* **701**:117–128.
- Mhatre MC and Ticku MK (1992) Chronic ethanol administration alters gamma-aminobutyric acid A receptor gene expression. *Molecular Pharmacology* **42**:415–422.
- Mihic SJ, Ye Q, Wick MJ, Koltchine VV, Krasowski MD, Finn SE, Mascia MP, Valenzuela CF, Hanson KK, Greenblatt EP, Harris RA, and Harrison NL (1997) Sites of alcohol and volatile anaesthetic action on GABA<sub>A</sub> and glycine receptors. *Nature* **389**:385–389.
- Millan MH, Meldrum BS, and Faingold CL (1986) Induction of audiogenic seizure susceptibility by focal infusion of excitant amino acid or bicuculline into the inferior colliculus of normal rats. *Experimental Neurology* **91**:634–639.
- Misawa H, Sherr EH, Lee DJ, Chetkovich DM, Tan A, Schreiner CE, and Brecht DS (2002) Identification of a monogenic locus (*jams1*) causing juvenile audiogenic seizures in mice. *Journal of Neuroscience* **22**:10088–10093.
- Molnar LR, Fleming WW, and Taylor DA (2000) Alterations in neuronal  $\gamma$ -aminobutyric acid<sub>A</sub> receptor responsiveness in genetic models of seizure susceptibility with different expression patterns. *Journal of Pharmacology and Experimental Therapeutics* **295**:1258–1266.
- Mozzrymas JW, Barberis A, Mercik K, and Zarnowska ED (2003) Binding sites, singly bound states, and conformation coupling shape GABA-evoked currents. *Journal of Neurophysiology* **89**:871–883.
- N'Gouemo P, Caspary DM, and Faingold CL (1996) Decreased GABA effectiveness in the inferior colliculus neurons during ethanol withdrawal in rats susceptible to audiogenic seizures. *Brain Research* **724**:200–204.
- Nehlig A, Vergnes M, Hirsch E, Boyet S, Koziel V, and Marescaux C (1995) Mapping of cerebral blood flow changes during audiogenic seizures in Wistar rats: effect of kindling. *Journal of Cerebral Blood Flow and Metabolism* **15**:259–269.

- Neumann PE and Collins RL (1991) Genetic dissection of susceptibility to audiogenic seizures in inbred mice. *Proceedings of the National Academy of Sciences of the United States of America* **88**:5408–5412.
- Oh S and Ho IK (1999) Changes of [<sup>3</sup>H]muscimol binding and GABA<sub>A</sub> receptor β2-subunit mRNA level by tolerance to and withdrawal from pentobarbital in rats. *Neurochemical Research* **24**:1603–1609.
- Onodera K, Tuomisto L, Tacke U, and Airaksinen M (1992) Strain differences in regional brain histamine levels between genetically epilepsy-prone and resistant rats. *Methods and Findings in Experimental and Clinical Pharmacology* **14**:13–16.
- Palombi PS and Caspary DM (1996) GABA inputs control discharge rate primarily within frequency receptive fields of inferior colliculus neurons. *Journal of Neurophysiology* **75**:2211–2219.
- Patrick DL and Faingold CL (1989) Auditory brainstem responses (ABRs) and effects of microinjection into inferior colliculus (IC) of an excitant amino acid (EAA) antagonist or GABA agonist on audiogenic seizures (AGS) in thyroid deficient (THX) or genetically epilepsy-prone rats (GEPRs). *Society for Neuroscience Abstracts* **15**:46.
- Penny JE, Brown RD, Wallace MS, and Henley CM (1986) Auditory aspects of seizure in the genetically epilepsy prone rat. *Life Sciences* **39**:887–895.
- Pierson MG and Swann JW (1988) The sensitive period and optimum dosage for induction of audiogenic seizure susceptibility by kanamycin in the Wistar rat. *Hearing Research* **32**:1–10.
- Raisinghani M and Faingold CL (2003) Identification of the requisite brain sites in the neuronal network subserving generalized clonic audiogenic seizures. *Brain Research* **967**:113–122.
- Raisinghani M, Feng H-J, and Faingold CL (2003) Glutamatergic activation of the amygdala differentially mimics the effects of audiogenic seizure kindling in two sub-strains of genetically epilepsy-prone rats. *Experimental Neurology* **183**:516–522.
- Reid HM and Collins RL (1986) Lateralized audiogenic seizure: motor asymmetries exhibited and the effects of interrupted stimulation. *Behavioral Neural Biology* **46**:424–431.
- Reigel CE, Dailey JW, and Jobe PC (1986) The genetically epilepsy-prone rat: an overview of seizure-prone characteristics and responsiveness to anticonvulsant drugs. *Life Sciences* **39**:763–774.
- Riaz A and Faingold CL (1994) Seizures during ethanol withdrawal are blocked by focal microinjection of excitant amino acid antagonists into the inferior colliculus and pontine reticular formation. *Alcoholism Clinical Experimental Research* **18**:1456–1462.
- Ribak CE, Byun MY, Ruiz GT, and Reiffenstein RJ (1988) Increased levels of amino acid neurotransmitters in the inferior colliculus of the genetically epilepsy-prone rat. *Epilepsy Research* **2**:9–13.
- Ribak CE, Lauterborn JC, Navetta MS, and Gall CM (1993) The inferior colliculus of GEPRs contains greater numbers of cells that express glutamate decarboxylase (GAD67) mRNA. *Epilepsy Research* **14**:105–113.
- Ribak CE, Khurana V, and Lien NT (1994) The effect of midbrain collicular knife cuts on audiogenic seizure severity in the genetically epilepsy-prone rat. *Journal für Hirnforschung* **35**:303–311.
- Roberto M, Madamba SG, Moore SD, Tallent MK, and Siggins GR (2003) Ethanol increases GABAergic transmission at both pre- and postsynaptic sites in rat central amygdala neurons. *Proceedings of the National Academy of Sciences of the United States of America* **100**:2053–2058.

- Roberts RC, Ribak CE, and Oertel WH (1985) Increased numbers of GABAergic neurons occur in the inferior colliculus of an audiogenic model of genetic epilepsy. *Brain Research* **361**:324–338.
- Ross KC and Coleman JR (1999) Audiogenic seizures in the developmentally primed Long-Evans rat. *Developmental Psychobiology* **34**:303–313.
- Ross KC, Coleman JR, and Jones LS (2001) Anti-epileptiform effects of audiogenic seizure priming on in vitro kindling in rat hippocampus. *Neuroscience Letters* **299**:234–238.
- Russek SJ, Bandyopadhyay S, and Farb DH (2000) An initiator element mediates autologous down regulation of the human type A  $\alpha$ -aminobutyric acid receptor  $\beta$ 1 subunit gene. *Proceedings of the National Academy of Sciences of the United States of America* **97**:8600–8605.
- Saint Marie RL (1996) Glutamatergic connections of the auditory midbrain: selective uptake and axonal transport of D-[ $^3$ H]aspartate. *Journal of Comparative Neurology* **373**:255–270.
- Sakamoto T and Niki H (2001) Acoustic priming lowers the threshold for electrically induced seizures in mice inferior colliculus, but not in the deep layers of superior colliculus. *Brain Research* **898**:358–363.
- Sieghart W (2000) Unraveling the function of GABA<sub>A</sub> receptor subtypes. *Trends in Pharmacological Sciences* **21**:411–413.
- Simler S, Vergnes M, and Marescaux C (1999) Spatial and temporal relationships between c-Fos expression and kindling of audiogenic seizures in Wistar rats. *Experimental Neurology* **157**:106–119.
- Simson PE, Criswell HE, and Breese GR (1991) Ethanol potentiates  $\gamma$ -aminobutyric acid-mediated inhibition in the inferior colliculus: evidence for local ethanol/ $\gamma$ -aminobutyric acid interactions. *Journal of Pharmacology and Experimental Therapeutics* **259**:1288–1293.
- Skradski SL, Clark AM, Jiang H, White HS, Fu H, and Ptáček LJ (2001) A novel gene causing a mendelian audiogenic mouse epilepsy. *Neuron* **31**:537–544.
- Smith PH (1992) Anatomy and physiology of multipolar cells in the rat inferior collicular cortex using the in vitro brain slice technique. *Journal of Neuroscience* **12**:3700–3715.
- Snyder-Keller AM and Pierson MG (1992) Audiogenic seizures induce c-fos in a model of developmental epilepsy. *Neuroscience Letters* **135**:108–112.
- Suneja SK, Benson CG, Gross J, and Potashner SJ (1995) Evidence for glutamatergic projections from the cochlear nucleus to the superior olive and the ventral nucleus of the lateral lemniscus. *Journal of Neurochemistry* **64**:161–171.
- Suneja SK, Potashner SJ, and Benson CG (2000) AMPA receptor binding in adult guinea pig brain stem auditory nuclei after unilateral cochlear ablation. *Experimental Neurology* **165**:355–369.
- Tecott LH, Sun LM, Akana SF, Strack AM, Lowenstein DH, Dallman MF, and Julius D (1995) Eating disorder and epilepsy in mice lacking 5-HT<sub>2C</sub> serotonin receptors. *Nature* **374**:542–546.
- Tsai G and Coyle JT (1998) The role of glutamatergic neurotransmission in the pathophysiology of alcoholism. *Annual Review of Medicine* **49**:173–184.
- Ueno S, Harris RA, Messing RW, Sanchez-Perez AM, Hodge CW, McMahon T, Wang D, Mehmert KK, Kelley SP, Haywood A, Olive MF, Buck KJ, Hood HM, Blednov Y, Findlay G, and Mascia MP (2001) Alcohol actions on GABA<sub>A</sub> receptors: from protein structure to mouse behavior. *Alcoholism Clinical Experimental Research* **25** Supplement **5**:76S–81S.

- Upton N and Stratton S (2003) Recent developments from genetic mouse models of seizures. *Current Opinion in Pharmacology* **3**:19–26.
- Van Middlesworth L and Norris CH (1980) Audiogenic seizures and cochlear damage in rats after perinatal antithyroid treatment. *Endocrinology* **106**:1686–1690.
- Veatch LM and Gonzalez LP (2000) Nifedipine alleviates alterations in hippocampal kindling after repeated ethanol withdrawal. *Alcoholism Clinical Experimental Research* **24**:484–491.
- Wada JA (1970) Susceptibility to audiogenic seizure induced by thiosemicarbazide. *Electroencephalography and Clinical Neurophysiology* **28**:100.
- Wagner T (1996) Lemniscal input to identified neurons of the central nucleus of mouse inferior colliculus: an intracellular brain slice study. *European Journal of Neuroscience* **8**:1231–1239.
- Wecker JR and Ison JR (1984) Acute exposure to methyl or ethyl alcohol alters auditory function in the rat. *Toxicology and Applied Pharmacology* **74**:258–266.
- Willott JF and Turner JG (2000) Neural plasticity in the mouse inferior colliculus: relationship to hearing loss, augmented acoustic stimulation, and prepulse inhibition. *Hearing Research* **147**:275–281.
- Yan QS, Dailey JW, Steenbergen JL, and Jobe PC (1998) Anticonvulsant effect of enhancement of noradrenergic transmission in the superior colliculus in genetically epilepsy-prone rats (GEPRs): a microinjection study. *Brain Research* **780**:199–209.
- Yang L and Faingold CL (2002) The amygdala is a requisite nucleus in the neuronal network for ethanol withdrawal seizures. *Society for Neuroscience Abstracts* **28**:607.1.
- Yang L, Long C, and Faingold CL (2001) Audiogenic seizure susceptibility is induced by termination of continuous infusion of  $\gamma$ -aminobutyric acid or an *N*-methyl-D-aspartic acid antagonist into the inferior colliculus. *Experimental Neurology* **171**:147–152.
- Yang L, Long C, Randall ME, and Faingold CL (2003) Neurons in the periaqueductal gray are critically involved in the neuronal network for audiogenic seizures during ethanol withdrawal. *Neuropharmacology* **44**: 275–281.
- Yechikhov S, Morenkov E, Chulanova T, Godukhin O, and Shchipakina T (2001) Involvement of cAMP- and  $\text{Ca}^{2+}$ /calmodulin-dependent neuronal protein phosphorylation in mechanisms underlying genetic predisposition to audiogenic seizures in rats. *Epilepsy Research* **46**:15–25.
- Yip PK, Meldrum BS, and Rattray M (2001) Elevated levels of group-III metabotropic glutamate receptors in the inferior colliculus of genetically epilepsy-prone rats following intracollicular administration of L-serine-*O*-phosphate. *Journal of Neurochemistry* **78**:13–23.
- Zhang H and Kelly JB (2001) AMPA and NMDA receptors regulate responses of neurons in the rat's inferior colliculus. *Journal of Neurophysiology* **86**:871–880.
- Zhang H and Kelly JB (2003) Glutamatergic and GABAergic regulation of neural responses in inferior colliculus to amplitude modulated sounds. *Journal of Neurophysiology* **90**:477–490.

# Chapter 22

## The Inferior Colliculus: Past, Present, and Future

JOHN H. CASSEDAY, CHRISTOPH E. SCHREINER, AND  
JEFFERY A. WINER

### 1. RATIONALE

From an historical perspective, the last few decades have seen a spectacular increase in our knowledge of the inferior colliculus (IC). The topics of the previous chapters cover the range of neurobiology, thus reflecting the complexity of the issues concerning the IC. However, the wide range makes it virtually impossible to provide a synthesis that does justice to all the chapters. Therefore, we address a few salient themes that, through our view, seem to merit analysis or that are otherwise underrepresented in the text. We hope this idiosyncratic view will be a polemic to encourage new ways of thinking about the auditory midbrain.

### 2. HISTORICAL OVERVIEW

Much of the early thought on function of the auditory midbrain (Whitfield 1967; Aitkin 1986) was influenced by Ramón y Cajal's concepts of the mammalian auditory midbrain. He described the principal component of lateral lemniscal ascending fibers as a "reflex bundle terminating in the IC" (Ramón y Cajal 1911). The second and large component consisted of fibers that bifurcated, sending a branch to the IC and another to the thalamus. Fibers continuing to the medial geniculate nucleus, the so-called "central acoustic tract," were considered a major constituent (one third or more) of the entire ascending body (Papez 1929). In short, the ascending auditory pathway to the telencephalon bypassed the IC, and the ascending auditory pathways to the IC served as the auditory input necessary for midbrain reflex functions. Thus, the IC was seen as serving hearing in much the same way as the superior colliculus served vision.

However, by the 1960s a different view began to emerge, one that is relatively close to the current view that the IC is an auditory center that is an obligatory synapse for nearly all ascending input to the medial geniculate nucleus (Whitfield 1967). The evidence came from several early tract tracing studies showing

that nearly all fibers in the lateral lemniscus terminated in the IC (Woollard and Harpman 1940; Barnes et al. 1943; Rasmussen 1946). The “central acoustic tract,” as first defined, is a quite minor component of the ascending auditory pathway (Casseday et al. 1989).

A critical step came as retrograde transport methods became available in the 1970s and 1980s (Adams 1979). From these studies an appreciation grew for the extent of convergence at the IC, from both ascending and descending auditory pathways (Aitkin 1986; Irvine 1986; Pollak and Casseday 1989; Casseday et al. 2002). Thus, the IC was construed as the “nexus” (Aitkin 1986) or “shunting-yard” (Ehret 1997) of the auditory pathway.

The same neuronal tracing methods called attention to the nonauditory inputs to the IC, some of which had been known from older degeneration techniques, such as the somatosensory inputs to the IC (RoBards et al. 1976), and some of which were newly identified such as projections from the substantia nigra (Ola-zábal and Moore 1989) and projections from the amygdala (Marsh et al. 2002).

Physiological techniques to block inhibitory transmitters revealed a crucial functional aspect of the convergence, namely, that most, if not all, IC neurons receive auditory evoked inhibition as well as excitation (see Chapters 9 to 13). Further, the timing of the excitatory and inhibitory inputs can be offset from one another, providing a possible mechanism for processing complex temporal information.

Thus, the IC truly merits a descriptive term such as Aitkin’s (1986) “nexus” or “hub” for its broad patterns of afferent convergence and for the active reshaping of sound encoding (Casseday et al. 2002). Although the IC is clearly the main route to auditory thalamus and cortex, it also serves motor functions that have to be taken into account when considering its manifold roles (Casseday et al. 1976). Ramón y Cajal’s idea of the IC as an auditory reflex pathway was simply incomplete in the important sense that the IC both transmits information to and receives information from “higher” auditory centers. Furthermore, the IC has functions that cannot be viewed as “reflex” in the classic sense, yet must certainly be construed as motor, in that it serves as a route to enable motor action via telencephalic auditory centers or via deep superior colliculus or pontine gray/cerebellar circuits. Naturally, the widespread intrinsic operations of the IC both at a cellular and a local circuit level make their own contributions in sculpting the many functions of this critical hub. These operations are just beginning to be understood (see Chapters 2 to 5).

Pioneering physiological studies (Thurlow et al. 1951) were followed by experiments revealing narrow-band, level-tolerant frequency tuning and inhibitory sidebands in IC cells (Katsuki et al. 1958; Erulkar 1959). Studies by Rose and Hind and colleagues were crucial in establishing tonotopy, binaural response types, and temporal response patterns (Rose et al. 1963). The first intracellular IC recordings provided direct evidence of long-lasting temporal sequences of excitation and inhibition in response to sound (Nelson and Erulkar 1963) showing that “. . . changes in membrane potential are specific to particular tonal frequencies and may outlast the duration of the tone. They serve to set the levels

of excitability of the neurons and to regulate the patterns of firing in response to long-lasting tonal stimuli” (Nelson and Erulkar 1963, p. 921). These observations led to interest in how IC neurons process time varying sounds (Nelson et al. 1966). Likewise, the IC became an early focus of neurophysiological studies of the mechanisms for processing temporal signals by echolocating bats (Grinnell 1963; Suga 1964).

Behavioral-ablation studies drew attention to subcortical mediation of auditory discrimination abilities. Cats with bilateral ablation of the entire auditory cortex had surprisingly little or no difficulty relearning frequency discrimination (Neff et al. 1975). For example, even after bilateral transection of the brachium of the inferior colliculus (BIC) cats could still relearn a frequency discrimination. These early anatomical, physiological, and behavioral studies already revealed the essence of the major themes that define our current understanding of IC structure and function as summarized in this volume.

### 3. COMPARATIVE NEUROBIOLOGY OF THE AUDITORY MIDBRAIN

Most studies of the auditory midbrain are concentrated in a few mammals, the rat, cat, bat, and on one bird, the barn owl. There are many investigations on the amphibian auditory midbrain, virtually all on anuran amphibians (Wilczynski 1988). There is a small but growing literature on fish, but the reptilian auditory midbrain has received far less attention (see Chapter 16). Moreover, the species examined are often chosen for their specializations, for example, the barn owl for sound localization and the bat for temporal processing (see Chapter 17). How can we make meaningful comparisons based on so limited a sample? If the auditory midbrain were a stereotyped structure with a common architectonic, neuroanatomical, neurochemical, neuroethological, neurophysiological, and ontogenetic arrangement, insularity with respect to a few aquatic, terrestrial, and aerial species would be a defensible position.

Nevertheless, the preceding chapters have revealed some features that appear to be common among vertebrates so far examined and might embody general principles of organization that appeared early in evolution (Grothe et al. 2004). One is the organization into a core that is the primary target of ascending auditory pathways and surrounding areas that receive other inputs, some of which may be descending and not all of which are auditory. Another is the presence of multiple ascending auditory pathways, some limbs of which ascend directly while others ascend via a synapse, providing different synaptic delays at their target in the IC (Casseday and Covey 1996). In fish, the pathways are few; in birds and mammals they are many; however, all vertebrates so far examined seem to have direct and indirect pathways. In this context, it is important to note the recent discovery that in the auditory midbrain of fish, specializations emerge for processing complex temporal features of sound, as is the case in



amphibians and mammals (see Chapter 16) and probably also in songbirds (Woolley and Casseday 2004). A third common feature is the close connection of the auditory midbrain with motor systems, especially those concerned with vocalization. Many of these points have been noted in prior work (Wilczynski 1988), and supporting evidence has grown considerably since then. The issue of how the auditory midbrain participates in processing and influencing species-specific vocalizations is an important venue for further comparative studies in all vertebrates (see Chapter 14).

#### 4. THE MAMMALIAN AUDITORY MIDBRAIN

Even considering only the mammalian IC in more detail, there are many unanswered questions. One might expect to see gross structural differences in the IC of different mammals in view of the enormous comparative differences in the cochlear nucleus (Moore 1980), trapezoid body (Richter et al. 1983), lateral lemniscal nuclei (Covey and Casseday 1995; Merchán et al. 1997), medial geniculate body (MGB) (Morest and Winer 1986), and auditory cortex (Winer 1992). However, at a macroscopic level, the IC in different species looks remarkably similar (Fig. 22.1): each species has a substantial central nucleus in which disc-shaped neurons form conspicuous fibrodendritic laminae (Fig. 22.2) much like those in the cat (Oliver and Morest 1984) and human (Geniec and Morest 1971). In addition, in each species the IC has an external cortex (Fig. 22.1A: bat; 22.1B: owl; 22.1C: rat; 22.1E: macaque), lateral nucleus (Fig. 22.1D: cat), or lateral zone (Fig. 22.1F: human).

Are these subdivisions homologous? With respect to architectonic, neuroanatomical, neurochemical, neuroethological, neurophysiological, and ontogenetic frames of reference—all pertinent to establishing such correspondences (Butler and Hodos 1996)—the evidence is too fragmentary to permit a conclusion. The cytoarchitecture of the lateral rind of the inferior colliculus region—denominated variously as external cortex/lateral nucleus/lateral zone/pericentral nucleus—has been described in the rat (Faye-Lund and Osen 1985). Its neurons have been classified to some extent (Malmierca 1991); certain neurochemical aspects have received attention (Chernock et al. 2004); and a limited amount of physiology (Syka and Popelář 1984; Smith 1992) and a developmental profile have been described (Altman and Bayer 1981). Many of the features in these studies suggest that it differs from the central nucleus, although we do not yet have a sufficiently large or secure database to propose that any given structure is homologous across species.

Three conclusions follow from this analysis: (1) It might be possible to assemble the elements for such a list of definitive criteria for the rat, but the task of looking at many species including bats, owls, cats, primates, and human would be daunting, and inspection of the contemporary literature shows few signs of providing the essential data. (2) Even if such a survey were available, there is little evidence to establish functional correspondences between species,




Figure 22.1. Cytoarchitectonic views of inferior colliculus organization in six common experimental species in Nissl preparations. Protocol: 30  $\mu\text{m}$  thick celloidin-embedded or frozen sections counterstained with cresyl violet acetate. (A) In the mustached bat, *Pteronotus p. parnellii*, the dorsoposterior division (DpD) is devoted to the approximately 61-kHz second harmonic, and the external cortex (Ex) is a slender rind (Zook et al. 1985). (B) The barn owl (*Tyto alba*) central nucleus (ICc) is massive, but has no apparent laminar organization (Knudsen 1983; Takahashi and Konishi 1988). (C) Rat external cortex (EC) and central nucleus both expand as a consequence of reduction of the dorsal cortex (DC) (Faye-Lund and Osen 1985). (D) The cat central nucleus (CN) shows regional architectonic variation and the dorsal cortex (DC) is massive (Oliver 1984; Winer et al. 1998). (E) The rhesus monkey (*Macaca mulatta*) has a large central nucleus (CN). (J.A. Winer and D.T. Larue, unpublished observations). (F) The human inferior colliculus has a fiber rich lateral zone (LZ) and a reduced dorsal cortex (DC) relative to the cat (D).

for example, whether the same types of neurons subserved central nucleus bin-aural processing. (3) No synaptic picture exists for serial, parallel, hierarchical, and multimodal (Aitkin et al. 1994) processing for external cortex/lateral nucleus/lateral zone/pericentral nucleus, and that would distinguish these structures from the central nucleus. The rigorous statement of the comparative position is that all possible comparisons would be required to establish homologies; a more liberal interpretation is that a subset of criteria would suffice (Morest and Winer 1986), much as cladistic studies rest on critical evidence (Eisenberg 1981). Resolution of these questions will require more knowledge about the canonical IC circuits as a prelude to specifying the nature of species differences.

## 5. THE CENTRAL NUCLEUS: *PRIMUS INTER PARES*

Much of the text in this volume is devoted to the central nucleus, which is a defensible emphasis in view of its primacy in hearing and its apparent homology among vertebrates (Grothe et al. 2004). Ultimately, however, the dorsal cortex, lateral nucleus, caudal cortex, rostral pole, and intercollicular tegmentum must each receive attention commensurate with their importance, as an example will show.

Physiological experiments suggest that the auditory cortex can modulate frequency-specific plasticity in the central nucleus of the inferior colliculus in bats (Jen et al. 1998; Suga et al. 2000) and mice (Yan and Ehret 2001). However, the present route for this effect in the central nucleus is unknown, and it may be indirect, requiring the participation of the external cortex/lateral nucleus (but see Chapter 5). The reason is that the most sensitive anterograde tract tracing techniques show only sparse input to the central nucleus in the cat after large cortical tracer deposits, although there are massive projections to the external cortex from many fields in the cat (Winer et al. 1998). In rats there is a difference of opinion in regard to the density of auditory cortex input to the central nucleus, with either a substantial (Saldaña et al. 1996) or a far smaller input (Herbert et al. 1991) comparable to that reported in the cat. A related issue is how the central nucleus is defined. In any case, the circuitry for expressing central feedback either depends on the external cortex/lateral nucleus because abundant intrinsic connections link it to the central nucleus (Saldaña and Merchán 1992), or there may be species differences in cortical input. In principle, the other inferior colliculus subdivisions are no less important with regard to their functional affiliations: for example, the dorsal cortex is implicated in attention (Jane et al. 1965), the intercollicular tegmentum in commissural and multimodal roles (RoBards et al. 1976), the lateral nucleus in the processing of complex sounds (Aitkin et al. 1994), and the rostral pole has significant projections to the superior colliculus (Harting and Van Lieshout 2000), presumably for spatial orientation.




Figure 22.2. The constraints inherent in the Nissl method as an architectonic tool (cf. Fig. 22.1) are readily apparent in Golgi preparations from the central nucleus in four species. Protocol: Golgi-Cox preparations, 140  $\mu\text{m}$  thick celloidin-embedded sections. Planapochromat, numerical aperture 0.65,  $\times 500$ . (A) In the mustached bat (see Fig. 22.1A) a laminar organization is readily apparent in DpD (cf. *black inset*) that is masked in Nissl preparations, including disc-shaped cells (1) as well as neurons with dendrites orthogonal to the laminar axis (2). Such neurons are common in the opossum (B), rat (C), and cat (D) inferior colliculus. (B) In the opossum, many disc-shaped cells have dendritic fields more constrained and flattened than those in *Pteronotus*. (C) In the rat there is an especially wide range of sizes and shapes in central nucleus cells that is matched by the cat (D). (D) Some cat disc-shaped cells have spiny dendrites (1).

## 6. THE QUESTION OF INHIBITORY INTERNEURONS IN THE INFERIOR COLLICULUS

This question requires first a definition of interneurons. The classic view assigns neurons to either type I (long axon) or type II (short axon) categories (Golgi 1873), a scheme that applies to all but a few cells, for example, axonless amacrine cells (Masland 2001). Classic type II cells (Ramón y Cajal 1911) have these features: a short, often unmyelinated, axon (Winer and Morest 1983) whose plexus and main sphere of influence is limited to perhaps 1 or 2 mm (Kisvárdy et al. 1993); smooth dendrites with a few long, slender, and pedunculated appendages (Winer and Larue 1988); a soma about 10  $\mu\text{m}$  in diameter (Winer 1986); and a primary transmitter of  $\gamma$ -aminobutyric acid (GABA) or glycine (Todd 1996).

With regard to the IC, this dichotomous arrangement encounters several difficulties. Thus, some 20% of central nucleus neurons are GABAergic, and these have subtypes that include both the smallest and the largest neurons (Oliver et al. 1994). Moreover, about the same proportion of cells that are GABAergic also project to the MGB (Winer et al. 1996), and intracellular labeling demonstrates that most neurons injected have a local axonal plexus, reminiscent of type II cells, as well as presumptive tectofugal long axonal projections, which is not characteristic of interneurons (Oliver et al. 1991). Thus, the central nucleus either has no classic type II cells, unlike the cochlear nucleus (Adams and Mugnaini 1987), MGB (Huang et al. 1999), and auditory cortex (Prieto et al. 1994), or it more closely resembles the superior olivary complex and lateral lemniscal nuclei in having GABAergic neurons with long projections (Saint Marie et al. 1989). An IC with no classic interneurons necessarily has a different functional organization than one with GABAergic neurons that have both local and remote projections. For example, the synaptic glomeruli that are a cardinal feature of MGB intrinsic organization and in which Golgi type II cells have an important role (Morest 1975) do not seem to occur in the IC (Rockel and Jones 1973).

## 7. CONTEXT DEPENDENCY OF INFERIOR COLLICULUS PROCESSING

Audition must capture events in time, auditory objects that are fleeting disturbances in air pressure, and reconstruct the nature of the sound source and the information it contains. This task is complicated by the fact that a given auditory object, such as a phoneme, may be embedded in different acoustical and informational conditions, or context, that must be considered when performing the analysis.

The context of an acoustic signal activating neurons in the IC is determined by the intensity distribution across the spectral profile of the signal at every

instant in time, by the time course of intensity changes across the spectrum, including dynamic changes in spectral components, and by the phase relationships and their changes among the spectral components, all of which are influenced by the location of the sound source in the auditory space. That is, under natural hearing conditions, sounds have a dynamic stimulation effect on the IC. Because of intrinsic mechanisms of adaptation and synaptic mechanisms of habituation, the instantaneous response of an IC neuron reflects not only the sound composition at the related instant in time but also the sound history, that is, the recent state of its own activity and that of neurons that provide input to it. For example, IC neurons may respond to bursts of a tone differently depending on whether the amplitude is variable or constant (Neuert et al. 2001), or whether a certain tone is reached by a frequency sweep or a frequency step (Malone and Semple 2001). Further, direction and speed of a frequency sweep may affect the response to a following tone (Suga 1969; Fuzessery 1994; Hage and Ehret 2003). Finally, transitions between sounds with different amplitude histograms (or contrasts) can elicit release from adaptation, thus substantially increasing the salience in the representation of a newly introduced amplitude statistics (Kvale and Schreiner 2004).

Accordingly, time-critical statistical properties of the acoustic environment can have a marked influence on the stimulus/object properties and the dynamics of neural responses. Consequences of these influences emerge when characterizing receptive fields with different stimulus statistics (Escabí and Schreiner 2002) and following a context change (Kvale and Schreiner 2004).

The ability of IC neurons to adapt their performance to the local statistics of the acoustic environment provides several advantages for information processing, including gain control, and enhanced detection, discrimination, and classification abilities. This context sensitivity may hold the key for understanding the transition from a peripheral analysis based on sound properties to a central one oriented to auditory objects.

## 8. STREAMS OF FUNCTION AND FUNCTIONS OF STREAMS

The inferior colliculus is located at a crucial transition in the flow of auditory information processing. Peripheral to the IC, incoming information can be viewed as being segregated into several processing streams that either emphasize specific stimulus aspects, including frequency composition, phase-locking, onsets and other amplitude variations, or compute specific stimulus aspects such as those related to the comparison of inputs to the two ears or the frequency filtering at the pinnae. Thus, the auditory stimulus is decomposed and undergoes parallel analysis. These multiple processing streams can be considered stimulus-based, and the description of the neuronal function is well served by reference to elemental sound attributes.

Rostral to the IC, the main principle of processing appears to be increasingly object-, task-, and perhaps action-related. Examples of identifiable task-specific central processing streams are reflected in the Doppler-shift and echo-delay maps in bats (O'Neill 1995), and the proposed "what" and "where" pathways in primates (Romanski et al. 1999). Other necessary central processing tasks include representational robustness in background noise, object equivalencies across sound source (speaker) variability, and enabling categorical signal classification. Even at the midbrain level, specific stimulus evaluations regarding urgent action-related tasks must be established or at least prepared (Casseday and Covey 1996). Examples include reflexive and other motor behaviors such as pinna movements, acoustic startle reflexes, responses to aversive stimuli, and vocalization (see Chapter 7). Each of these ascending (see Chapters 3 and 4) and descending (see Chapters 6 and 8) functional streams likely relies on specific superpositions of convergent feedforward and feedback information that shape and refine the task-specific processing. Potential segregations of different functional streams may be expressed in the many subdivisions in the midbrain and other nuclei (see Chapter 1), or in the different synaptic domains in the central nucleus of the inferior colliculus (see Chapter 2). It remains for future work to identify the origin, termination, and function of different streams, to establish whether incoming and outgoing projections are sufficiently compartmentalized that they can be unequivocally related to a specific functional stream, and, finally, whether is it even useful to refer to specific functional streams if the task-oriented refinement requires cooperation of several convergent projections at each node of the auditory pathway.

## Abbreviations

AID	anterolateral division of the inferior colliculus
Cb	cerebellum
CG	central gray
DC	dorsal cortex of the inferior colliculus
DM	dorsomedial part of central nucleus
DNLL	dorsal nucleus of lateral lemniscus
DpD	dorsoposterior division of the inferior colliculus
Ex	external nucleus of the inferior colliculus
GABA	$\gamma$ -aminobutyric acid
ICc	central nucleus of the inferior colliculus
IcT	intercollicular tegmentum
ICx	external nucleus of the inferior colliculus
Imc	nucleus isthmi, <i>pars magnocellularis</i>
Ipc	nucleus isthmi, <i>pars parvicellularis</i>
LL	lateral lemniscus
LN	lateral nucleus
LZ	lateral zone

MD	medial dorsal division of the inferior colliculus
MGB	medial geniculate body
SC	superior colliculus
Vln	ventrolateral nucleus

## References

- Adams JC (1979) Ascending projections to the inferior colliculus. *Journal of Comparative Neurology* **183**:519–538.
- Adams JC and Mugnaini E (1987) Patterns of glutamate decarboxylase immunostaining in the feline cochlear nucleus complex studied with silver enhancement and electron microscopy. *Journal of Comparative Neurology* **262**:375–401.
- Aitkin L (1986) *The Auditory Midbrain. Structure and Function in the Central Auditory Pathway*. Humana Press, Clifton.
- Aitkin L, Tran L, and Syka J (1994) The responses of neurons in subdivisions of the inferior colliculus of cats to tonal, noise and vocal stimuli. *Experimental Brain Research* **98**:53–64.
- Altman J and Bayer SA (1981) Time of origin of neurons of the rat inferior colliculus and the relations between cytogenesis and tonotopic order in the auditory pathway. *Experimental Brain Research* **42**:411–423.
- Barnes WT, Magoun HW, and Ranson SW (1943) Ascending auditory pathway in the brain stem of the monkey. *Journal of Comparative Neurology* **79**:129–152.
- Butler AB and Hodos W (1996) *Comparative Vertebrate Neuroanatomy: Evolution and Adaptation*. John Wiley & Sons, New York.
- Casseday JH and Covey E (1996) A neuroethological theory of the operation of the inferior colliculus. *Brain, Behavior and Evolution* **47**:311–336.
- Casseday JH, Diamond IT, and Harting JK (1976) Auditory pathways to the cortex in *Tupaia glis*. *Journal of Comparative Neurology* **166**:303–340.
- Casseday JH, Kobler JB, Isbey SF, and Covey E (1989) Central acoustic tract in an echolocating bat: an extralemniscal auditory pathway to the thalamus. *Journal of Comparative Neurology* **287**:247–259.
- Casseday JH, Fremouw T, and Covey E (2002) The inferior colliculus: hub of the auditory system. In: Oertel D, Popper AN, and Fay RR (eds). *Springer Handbook of Auditory Research, Volume 15: Integrative Functions in the Mammalian Auditory Pathway*. Springer, New York, pp. 238–318.
- Chernock ML, Larue DT, and Winer JA (2004) A periodic network of neurochemical modules in the inferior colliculus. *Hearing Research* **188**:12–20.
- Covey E and Casseday JH (1995) The lower brainstem auditory pathways. In: Popper AR and Fay RR (eds). *Springer Handbook of Auditory Research, Volume 5: Hearing by Bats*. Springer-Verlag, New York, pp. 235–295.
- Ehret G (1997) The auditory midbrain, a ‘shunting-yard’ of acoustical information processing. In: Ehret G and Romand R (eds). *The Central Auditory System*. Oxford University Press, New York, pp. 259–316.
- Eisenberg JF (1981) *The Mammalian Radiations*. University of Chicago Press, Chicago.
- Eruikar SD (1959) The responses of single units of the inferior colliculus of the cat to acoustic stimulation. *Proceedings of the Royal Society of London, Series B, Biological Sciences*, **150**:336–355.



- Escabi MA and Schreiner CE (2002) Nonlinear spectrotemporal sound analysis by neurons in the auditory midbrain. *Journal of Neuroscience* **22**:4114–4131.
- Faye-Lund H and Osen KK (1985) Anatomy of the inferior colliculus in the rat. *Anatomy and Embryology* **171**:1–20.
- Fuzessery ZM (1994) Response selectivity for multiple dimensions of frequency sweeps in the pallid rat inferior colliculus. *Journal of Neurophysiology* **72**:1061–1079.
- Geniec P and Morest DK (1971) The neuronal architecture of the human posterior colliculus: a study with the Golgi method. *Acta Oto-Laryngologica (Stockholm)* Supplement 295:1–33.
- Golgi C (1873) Sulla struttura della sostanza grigia del cervello. *Gazzetta Medicina Italiana Lombarda* **33**:244–246.
- Grinnell AD (1963) The neurophysiology of audition in bats: directional localization and binaural interaction. *Journal of Physiology (London)* **167**:97–113.
- Grothe B, Carr CE, Casseday JH, Fritzsche B, and Köppl C (2004) The evolution of central pathways and their neural processing patterns. In: Manley GA, Popper AN, and Fay RR (eds). *Springer Handbook of Auditory Research*, Volume 22: *Evolution of the Vertebrate Auditory System*. Springer-Verlag, New York, pp. 289–359.
- Hage SR and Ehret G (2003) Mapping responses to frequency sweeps and tones in the inferior colliculus of house mice. *European Journal of Neuroscience* **18**:2301–2312.
- Harting JK and Van Lieshout DP (2000) Projections from the rostral pole of the inferior colliculus to the cat superior colliculus. *Brain Research* **881**:244–247.
- Herbert H, Aschoff A, and Ostwald J (1991) Topography of projections from the auditory cortex to the inferior colliculus in the rat. *Journal of Comparative Neurology* **304**:103–122.
- Huang CL, Larue DT, and Winer JA (1999) GABAergic organization of the cat medial geniculate body. *Journal of Comparative Neurology* **415**:368–392.
- Irvine DRF (1986) The auditory brainstem. A review of the structure and function of auditory brainstem processing mechanisms. In: Aitman H, Ottoson D, Perl ER, Schmidt RF, Shimazu H, and Willis WD (eds). *Progress in Sensory Physiology* **7**:1–279.
- Jane JA, Masterton RB, and Diamond IT (1965) The function of the tectum for attention to auditory stimuli in the cat. *Journal of Comparative Neurology* **125**:165–192.
- Jen PH-S, Chen QC, and Sun XD (1998) Corticofugal regulation of auditory sensitivity in the bat inferior colliculus. *Journal of Comparative Physiology A* **183**:683–697.
- Katsuki Y, Sumi T, Uchiyama H, and Watanabe T (1958) Electric responses of auditory neurons in cat to sound stimulation. *Journal of Neurophysiology* **21**:569–588.
- Kisvárdy Z, Beaulieu C, and Eysel U (1993) Network of GABAergic large basket cells in cat visual cortex (area 18): implication for lateral disinhibition. *Journal of Comparative Neurology* **327**:398–415.
- Knudsen EI (1983) Subdivisions of the inferior colliculus in the barn owl (*Tyto alba*). *Journal of Comparative Neurology* **218**:174–186.
- Kvale M and Schreiner CE (2004) Adaptation to dynamic stimulus aspects and its effects on central auditory temporal receptive fields. *Journal of Neurophysiology* **91**:604–612.
- Malmierca MS (1991) *Reconstrucción Tridimensional por Ordenador de Neuronas Impregnadas con el Método de Golgi en el Colículo Inferior de la Rata*. Doctoral thesis, Universidad de Salamanca, Oslo and Salamanca.
- Malone BJ and Semple MN (2001) Effects of auditory stimulus context on the representation of frequency in the gerbil inferior colliculus. *Journal of Neurophysiology* **86**:1113–1130.

- Marsh RA, Fuzessery ZM, Grose CD, and Wenstrup JJ (2002) Projection to the inferior colliculus from the basal nucleus of the amygdala. *Journal of Neuroscience* **22**:10449–10460.
- Masland RH (2001) The fundamental plan of the retina. *Nature Neuroscience* **4**:877–886.
- Merchán MA, Malmierca MS, Bajo VM, and Bjaalig JG (1997) The nuclei of the lateral lemniscus. In: Syka J (ed). *Acoustical Signal Processing in the Central Auditory System*. Plenum Press, New York, pp. 211–226.
- Moore JK (1980) The primate cochlear nuclei: loss of lamination as a phylogenetic process. *Journal of Comparative Neurology* **193**:609–629.
- Morest DK (1975) Synaptic relations of Golgi type II cells in the medial geniculate body of the cat. *Journal of Comparative Neurology* **162**:157–194.
- Morest DK and Winer JA (1986) The comparative anatomy of neurons: homologous neurons in the medial geniculate body of the opossum and the cat. *Advances in Anatomy, Embryology and Cell Biology* **97**:1–96.
- Neff WD, Diamond IT, and Casseday JH (1975) Behavioral studies of auditory discrimination: central nervous system. In: Keidel WD and Neff WD (eds). *Handbook of Sensory Physiology, Volume V, Part 2: Auditory System, Anatomy, Physiology (Ear)*. Springer-Verlag, Berlin, pp. 307–400.
- Nelson PG and Erulkar SD (1963) Synaptic mechanisms of excitation and inhibition in the central auditory pathway. *Journal of Neurophysiology* **26**:908–923.
- Nelson PG, Erulkar SD, and Bryan JS (1966) Responses of units of the inferior colliculus to time-varying acoustic stimuli. *Journal of Neurophysiology* **29**:834–860.
- Neuert V, Pressnitzer D, Patterson RD, and Winter IM (2001) The responses of single units in the inferior colliculus of the guinea pig to damped and ramped sinusoids. *Hearing Research* **159**:36–52.
- Olazábal UE and Moore JK (1989) Nigrotectal projection to the inferior colliculus: horseradish peroxidase, transport and tyrosine hydroxylase immunohistochemical studies in rats, cats, and bats. *Journal of Comparative Neurology* **282**:98–118.
- Oliver DL (1984) Neuron types in the central nucleus of the inferior colliculus that project to the medial geniculate body. *Neuroscience* **11**:409–424.
- Oliver DL and Morest DK (1984) The central nucleus of the inferior colliculus in the cat. *Journal of Comparative Neurology* **222**:237–264.
- Oliver DL, Kuwada S, Yin TCT, Haberly LB, and Henkel CK (1991) Dendritic and axonal morphology of HRP-injected neurons in the inferior colliculus of the cat. *Journal of Comparative Neurology* **303**:75–100.
- Oliver DL, Winer JA, Beckius GE, and Saint Marie RL (1994) Morphology of GABAergic cells and axon terminals in the cat inferior colliculus. *Journal of Comparative Neurology* **340**:27–42.
- O'Neill WE (1995) The bat auditory cortex. In: Popper AN and Fay RR (eds). *Springer Handbook of Auditory Research, Volume 5: Hearing by Bats*. Springer-Verlag, New York, pp. 416–480.
- Papez JW (1929) Central acoustic tract in cat and man. *Anatomical Record* **42**:60.
- Pollak GD and Casseday JH (1989) *The Neural Basis of Echolocation in Bats*. Springer-Verlag, Berlin.
- Prieto JJ, Peterson BA, and Winer JA (1994) Morphology and spatial distribution of GABAergic neurons in cat primary auditory cortex (AI). *Journal of Comparative Neurology* **344**:349–382.

- Ramón y Cajal S (1911) *Histologie du Système Nerveux de l'Homme et des Vertébrés*. Maloine (original French edition). Consejo Superior de Investigaciones Científicas (1972 reprint) Paris.
- Rasmussen GL (1946) The olivary peduncle and other fiber projections of the superior olivary complex. *Journal of Comparative Neurology* **84**:141–219.
- Richter EA, Norris BE, Fullerton BC, Levine RA, and Kiang NY-S (1983) Is there a medial nucleus of the trapezoid body in humans? *American Journal of Anatomy* **68**:157–166.
- RoBards MJ, Watkins III DW, and Masterton RB (1976) An anatomical study of some somesthetic afferents to the intercollicular terminal zone of the midbrain of the opossum. *Journal of Comparative Neurology* **170**:499–524.
- Rockel AJ and Jones EG (1973) Observations on the fine structure of the central nucleus of the inferior colliculus of the cat. *Journal of Comparative Neurology* **147**:61–92.
- Romanski LM, Tian B, Fritz J, Mishkin M, Goldman-Rakic PS, and Rauschecker JP (1999) Dual streams of auditory afferents target multiple domains in the primate prefrontal cortex. *Nature Neuroscience* **2**:1131–1136.
- Rose JE, Greenwood DD, Goldberg JM, and Hind JE (1963) Some discharge characteristics of single neurons in the inferior colliculus of the cat. I. Tonotopical organization, relation of spike-counts to tone intensity, and firing patterns of single elements. *Journal of Neurophysiology* **26**:294–320.
- Saint Marie RL, Ostapoff E-M, Morest DK, and Wenthold RJ (1989) Glycine-immunoreactive projections of the cat lateral superior olive: possible role in midbrain ear dominance. *Journal of Comparative Neurology* **279**:382–396.
- Saldaña E and Merchán MA (1992) Intrinsic and commissural connections of the rat inferior colliculus. *Journal of Comparative Neurology* **319**:417–437.
- Saldaña E, Feliciano M, and Mugnaini E (1996) Distribution of descending projections from primary auditory neocortex to inferior colliculus mimics the topography of intracollicular projections. *Journal of Comparative Neurology* **371**:15–40.
- Smith PH (1992) Anatomy and physiology of multipolar cells in the rat inferior collicular cortex using the in vitro brain slice technique. *Journal of Neuroscience* **12**:3700–3715.
- Suga N (1964) Single unit activity in cochlear nucleus and inferior colliculus of echo-locating bats. *Journal of Physiology (London)* **172**:449–474.
- Suga N (1969) Echo-location and evoked potentials of bats after ablation of inferior colliculus. *Journal of Physiology (London)* **203**:707–728.
- Suga N, Gao E, Zhang Y, Ma X, and Olsen JF (2000) The corticofugal system for hearing: recent progress. *Proceedings of the National Academy of Sciences of the United States of America* **97**:11807–11814.
- Syka J and Popelář J (1984) Inferior colliculus in the rat: neuronal responses to stimulation of the auditory cortex. *Neuroscience Letters* **51**:235–240.
- Takahashi TT and Konishi M (1988) Projections of the nucleus angularis and nucleus laminaris to the lateral lemniscal nuclear complex of the barn owl. *Journal of Comparative Neurology* **274**:212–238.
- Thurlow WR, Gross NB, Kemp EH, and Lowy K (1951) Microelectrode studies of neural activity of cat. I. Inferior colliculus. *Journal of Neurophysiology* **14**:289–304.
- Todd AJ (1996) GABA and glycine in synaptic glomeruli of the rat spinal dorsal horn. *European Journal of Neuroscience* **8**:2492–2498.
- Whitfield IC (1967) *The Auditory Pathway*. Arnold, London.

- Wilczynski W (1988) Brainstem auditory pathways in anuran amphibians. In: Fritsch B, Ryan MJ, Wilczynski W, Hetherington TE, and Walkowiak W (eds). *The Evolution of the Amphibian Auditory System*. John Wiley & Sons, New York, pp. 209–232.
- Winer JA (1986) Neurons accumulating [<sup>3</sup>H]gamma-aminobutyric acid (GABA) in supragranular layers of cat primary auditory cortex (AI). *Neuroscience* **19**:771–793.
- Winer JA (1992) The functional architecture of the medial geniculate body and the primary auditory cortex. In: Webster DB, Popper AN, and Fay RR (eds). *Springer Handbook of Auditory Research, Volume 1: The Mammalian Auditory Pathway: Neuroanatomy*. Springer-Verlag, New York, pp. 222–409.
- Winer JA and Larue DT (1988) Anatomy of glutamic acid decarboxylase (GAD) immunoreactive neurons and axons in the rat medial geniculate body. *Journal of Comparative Neurology* **278**:47–68.
- Winer JA and Morest DK (1983) The neuronal architecture of the dorsal division of the medial geniculate body of the cat. A study with the rapid Golgi method. *Journal of Comparative Neurology* **221**:1–30.
- Winer JA, Saint Marie RL, Larue DT, and Oliver DL (1996) GABAergic feedforward projections from the inferior colliculus to the medial geniculate body. *Proceedings of the National Academy of Sciences of the United States of America* **93**:8005–8010.
- Winer JA, Larue DT, Diehl JJ, and Hefti BJ (1998) Auditory cortical projections to the cat inferior colliculus. *Journal of Comparative Neurology* **400**:147–174.
- Woollard HH and Harpman JA (1940) The connexions of the inferior colliculus and of the dorsal nucleus of the lateral lemniscus. *Journal of Anatomy (London)* **71**:441–458.
- Woolley SMN and Casseday JH (2004) Response properties of single neurons in the zebra finch auditory midbrain: response patterns, frequency coding, intensity coding, and spike latencies. *Journal of Neurophysiology* **91**:136–151.
- Yan J and Ehret G (2001) Corticofugal reorganization of the midbrain tonotopic map in mice. *NeuroReport* **12**:3313–3316.
- Zook JM, Winer JA, Pollak GD, and Bodenhamer RD (1985) Topology of the central nucleus of the mustache bat's inferior colliculus: correlation of single unit response properties and neuronal architecture. *Journal of Comparative Neurology* **231**:530–546.

# Subject Index

Species names appear after common names.

- ABR.** *See* Auditory brain stem response
- Acetylcholine**  
inferior colliculus, 258, 301, 302, 497–498, 567
- Acetylcholinesterase**, 501
- Acoustic chiasm**, 117
- Acoustic environments, complex**, 564
- Acoustic fovea.** *See* Fovea, auditory
- Acoustic nerve**, 268  
*See also* Auditory nerve
- Acoustic striae.** *See* Held, stria of; Monakow, stria of von
- Acoustic trauma**  
and cAMP, 571  
and white noise, 572  
decreased monaural inhibition after, 572  
effect on GABA-mediated inhibition, 575  
effect on tuning curve tails, 573  
preservation of characteristic frequency after trauma, 574  
*See also* Hearing loss
- Action potential, compound (CAP)**, 571, 576
- Acuity, directional**, 435
- Adaptation**, 390, 634
- Adaptation-of-excitation**, 262
- Adult plasticity**, 568
- After-discharge (AD)**, 588–589
- Afterhyperpolarization**, 287
- Aging**  
acetylcholine and, 567  
AMPA receptors and, 567  
changes in connectivity in, 566  
changes in neurotransmitter status, 568  
changes in synaptic organization, 567  
decline in GABA release, 268–269, 566–567  
effects on compound action potential (CAP), 571  
effects on GABA<sub>A</sub> receptors in inferior colliculus subdivisions, 566  
effects on inferior colliculus, 560  
effects on rate-intensity functions, 269  
gap thresholds in, 562  
NMDA receptors in, 567  
plasticity, 559  
temporal processing in, 566  
tonotopic organization, and, 559–560
- AI.** *See* Primary auditory cortex
- All-pass**, 474, 564
- AM (amplitude modulation)**  
aged rats, in, 562  
chemical mechanisms, 258  
coding, 349–361, 562  
inferior colliculus, 349–360  
intensity dependence, 353  
modulation transfer function, 350–356  
nonsinusoidal, 356–357  
noise effects, 353–354  
mechanisms, 359–361, 380  
primary auditory cortex, 38  
sinusoidal, 264, 352–354, 562  
speech, 347  
stimuli, 560  
torus semicircularis 467–468, 472–476  
wing beat frequencies, 516

- AMPA ( $\alpha$ -amino-3-hydroxy-4-isoxazoeppionic acid), 208, 248, 255, 258–259, 263, 294, 545, 550, 567
- AMPA antagonists, 254
- CNQX (6-cyano-7-nitroquinoxaline-2,3-dione), 253–254, 294–295
- NBQX (1,2,3,4-tetrahydro-6-nitro-2,3-dioxobenzol[f]quinoxaline-7-sulfonamide), 254
- AMPA-NMDA receptor-mediated excitation, proportion of, 263, 294
- Amphibian, 467–477
- Amplitude
- best, 510
- coding, 313, 333, 510
- tuning, 510
- Amygdala, 4, 36, 192, 201, 605
- Amygdalocollicular system, 231, 240, 242
- Anesthesia, 211, 217, 349, 562
- Angle, minimum audible, 434
- Angularis, nucleus, 506
- Anterior ventral cochlear nucleus
- anatomy, 5–14
- auditory nerve, 7
- cell types, 10–15
- multipolar cells, 10
- neuron types, 10
- phase locking in, 428
- projections to the bat inferior colliculus, 502
- projections to lateral nucleus of trapezoid body, 9
- projections to medial nucleus of trapezoid body, 12
- projections to medial superior olivary nucleus, 12
- projections to superior olive, 12
- spontaneous rate, 10
- See also* Cochlear nucleus
- Archistriatal nucleus, robust, 498
- Archistriatum, 498
- Aspartate
- auditory nerve, 7
- commissural neurotransmitter candidate, 174
- APV ( $\pm$ )-2-amino-phosphonovaleric acid, 209, 254
- Arousal, 193, 268
- Array code, 519
- papilla, 469
- Association cortex, projections of medial geniculate body, 42
- Attention, auditory, 36
- Audible angle, minimum (MAA), 434
- Audiogenic seizures (AGS)
- abnormal firing and, 588, 603
- acoustic priming, 592
- alcohol in, 605
- amygdala, role in, 605
- barbiturates in, 605
- benzodiazepines in, 605
- blockade of, 610, 613
- causes, 603
- chemical stimulation, 603
- c-fos* expression in, 603
- clonic, 605
- early-onset hearing loss, 604
- ethanol effects compared in vitro and in vivo, 611
- ethanol-induced audiogenic seizures, 604, 611
- focal microinjection and, 603, 610
- focal perfusion and, 603
- gene-knockouts and susceptibility to, 604
- genetic abnormalities in, 604
- genetically epilepsy-prone rats (GEPRs), 604
- GABA, 606
- GABA application and local inhibition, 606
- GABA enhancement increases seizure likelihood, 611
- GABA<sub>A</sub> receptor, 605, 611
- GABA<sub>A</sub> receptor subunit alterations, 611
- GABA<sub>B</sub> receptor, 605
- genetic and induced audiogenic-seizure-susceptibility, 611
- GEPR-3s, 604
- GEPR-9s, 604
- glutamate, 606, 611–613
- glutamate agonists enhance audiogenic seizure susceptibility, 611
- glutamate antagonists and enhancement of seizure susceptibility, 611
- hearing loss and, 604
- histamine, 606

- induction and initiation of, 603
- induction by electrical stimulation, 604
- induction in normal rodents, 605
- intense acoustic stimuli and, 603
- lesions and, 603
- metabotropic glutamate receptors and seizure blockade, 613
- neurotransmitter abnormalities in, 606
- norepinephrine, 606
- ototoxic drugs and, 603
- priming of, 603
- receptor desensitization, 611
- reduced effectiveness of GABAergic inhibition, 606
- rodents, 603–604
- seizure susceptibility and GABA, 610
- tonic, 605
- triggered by acoustic stimuli, 605
- withdrawal and enhancement of susceptibility, 605
- 2-deoxyglucose labeling, 603
- 5-hydroxytryptamine, 606
- Auditory and visual frames of reference, calibration of, 523
- Auditory brain stem response, 2
- Auditory cortex
  - binaural organization, 39
  - connections, 2–3, 35, 38–39, 42–43
  - human, 356
  - interaural time differences, cells sensitive to, 430
  - intrinsic connections, 14
  - layers, 37
  - neurochemistry, 39
  - projections to the cochlear nucleus, 42
  - projections to inferior colliculus, 41–42, 123
  - projections to the pontine nuclei, 42
  - projections to the superior olive, 42
  - pyramidal cells, 37
  - role in contralateral sound localization, 443
- Auditory cortical fields
  - cat, 35, 39, 42
- Auditory evoked potentials
  - human, 2
  - mouse, 593
  - noise-induced hearing loss, 593
- Auditory hemifield, 383
- Auditory localization. *See* Sound localization
- Auditory neocortex. *See* Auditory cortex
- Auditory nerve
  - ascending and descending branches, 5
  - bandwidth, 322
  - cat, 5
  - central projections, 5, 7–8
  - characteristic frequency
  - class I, class II, class III, class IV, 317
  - cross correlation functions, 388
  - descending branch, 8
  - effects of damage, 268
  - fiber bandwidth, 322
  - length, 5
  - phase locking, 428
  - projections to cochlear nucleus, 5
  - relationship to spiral ganglion fibers, 8
  - sensitivity, 317
  - synapse, 330
  - tuning, 317
  - See also* Acoustic nerve
- Auditory pathway
  - amphibian, 471–475
  - cat, 2
  - bird, 505
  - fish, 463
  - reptile, 478
- Auditory thalamocortical projections, 2–3
- Auditory-visual integration, 444
- Aurality, 315
- Average binaural level (ABI), 394, 519
  - See also* Excitatory monaural intensity constant (EMI)
- Avian. *See* Bird
- Axoplasmic transport, 118–119
- Azimuth, 145, 432, 445, 564, 589
- Azimuth, and interaural time difference processing, 519
- Azimuth perception, monaural spectral representation, contribution to, 440
- Background noise, masking from, 564**
- Background sources, moving, 564**
- Backward masking, effect on modulation rate, 516**
- Baclofen, 255, 267**
- Banding, 538–541, 548–550**

- Band-pass, 322, 512, 563  
 inhibitory contribution to amplitude-modulated transfer functions, 564
- Band-reject, 265, 564
- Band-width and level dependency, 327
- Bandwidth, 315
- Bandwidth, critical, 324–326
- Basilar membrane, 329
- Basilar membrane, and synchronization of spike discharge, 379
- Basilar papilla, 469, 477
- Bat
- auditory cortex, 504, 513
  - cochlear nucleus, 502, 504
  - connections of cochlear nucleus with inferior colliculus, 116, 118–119
  - expanded isofrequency representation, 506
  - GABA, 516–517
  - inferior colliculus, 510–517
  - medial geniculate body, 202–204, 206–207
  - pinnae, complexity of, 517
  - space map, absence in the bat mid-brain, 517
  - species-specific frequency representation, 507
  - superior olive, 503
  - transmitters, 497, 516, 523
  - trapezoid body, 503
  - ventral nucleus of lateral lemniscus, 503
  - vocalizations, 493–494
- Bat species
- Antrozous pallidus* (pallid bat), 494
  - Eptesicus fuscus* (big brown bat), 266, 315, 319, 323, 507, 509
  - Pteronotus p. parnellii* (mustached bat), 124, 205–206, 494
  - Rhinolophus rouxi* (horseshoe bat), 494
  - Tadarida brasiliensis* (Mexican free-tailed bat), 494
  - See also* Species
- Behavioral salience, 518
- Behavioral studies
- absence of studies of auditory performance in awake animals, 427
  - behavioral deficits and task specificity, 436
  - importance of, 426
  - in awake animals, 426, 444
  - natural, and functional perspective on localization, 443
  - precedence effect, 397–401, 444–450
  - sound localization, 628
- Belt auditory pathways, 167
- Best amplitude, 510
- Best duration, 512, 514
- Best frequency
- changes, 508
  - comparative, 508
  - inferior colliculus, 508
- Bicuculline, 261–267, 299–300, 349, 415, 608–610
- Bilateral projections, olivocochlear bundle, 185
- Binaural headphone cues and illusions, 436
- Binaural hearing
- biological significance, 377
  - cross correlation models, 401
  - inferior colliculus, 94
  - inhibition, 606
  - inhibition, release from, 262
  - level, average, sensitivity to, 519
  - masking level differences, 396–398
  - pathways for, 132, 142, 502
  - processing, 8, 90, 96, 142, 146, 396, 608
  - sensitivities, 381
  - target segregation in, 378
  - unmasking, 401
- Binaural input
- conditioner-probe interval, 399–400
  - lateral superior olivary nucleus, 15–19, 94–95, 502
  - masking level differences, 396
  - medial superior olivary nucleus, 91–94, 502
  - medial geniculate body, 217
  - primary auditory cortex, 39
  - trauma and decreased inhibition on, 572
- Biological coding, independence from intensity coding, 334
- Biophysical properties, 282
- Biosonar signals, 512–513, 516
- Biosonar sounds. *See* Biosonar signals
- Bipolar cells, in avian inferior colliculus, 500



- Bird
- auditory cortex, 505
  - caudomedial shell of the intercollicular region of the inferior colliculus, 498
  - cell types in inferior colliculus, 500
  - central core of the central nucleus of the inferior colliculus, 498
  - central nucleus of the inferior colliculus, 498
  - connections of the inferior colliculus, 505
  - dendritic fields of inferior colliculus neurons, 500
  - descending auditory-vocal circuits, 505
  - field L, 505
  - GABA, 523–524
  - hearing, 494
  - hyperstriatum, 505
  - inferior colliculus tuning curves, 515
  - intercollicular region of the midbrain, 498
  - lateral shell of the inferior colliculus, 498
  - medial geniculate body (nucleus ovoidalis), 505
  - medial shell of the inferior colliculus, 498
  - midbrain, 494, 499
  - owl sound localization, 494
  - paracentral nucleus of the intercollicular region of the inferior colliculus, 498
  - primary auditory cortex (field L), 505
  - projections from laminaris and magnocellular nuclei, 505
  - song nuclei, brain stem, 505
  - subdivisions of the midbrain, 498
  - superficial nucleus of the inferior colliculus, 498
  - thalamus, 505
  - tonotopy in central nucleus of the inferior colliculus, 508
  - torus semicircularis, 499
  - ventral nucleus of lateral lemniscus
  - vocal circuits, 505
- Brachium of, inferior colliculus, 84, 156, 169, 238, 628
- Brain slices, 253, 569
- Brain stem
- evoked response, 569
  - projections to medial geniculate body, 2–3
  - projections to cochlear nucleus, 186–187
  - projections to the inferior colliculus, 118–121
- Bursting firing mode, 233
- Bushy cells
- cochlear nucleus, 12, 392
  - projection to medial nucleus of the trapezoid body, 142
- See also* Spherical bushy cells
- Calbindin
- age and, 567
  - cytoplasmic calcium buffer, role as, 570
  - inferior colliculus, 497, 544–545, 567
- Calcium
- binding protein changes with aging, 567
  - binding proteins, 497, 544–545, 550
  - effects on GABA release, 251
  - intracellular, 570
  - levels after acoustic trauma, 570
- Calretinin, in inferior colliculus, 497, 501–502, 520, 544–545, 550, 567–568
- Carboniferous, 493
- Carnivore
- auditory pathway, 2
  - medial geniculate body, 32–37, 42
- Cartwheel cell, 12
- Cat (*Felis catus*)
- auditory nerve, 2
  - cochlear nucleus, 5–15
  - connections of cochlear nucleus with inferior colliculus, 116, 118–119
  - decerebrate, 319
  - frequency map of auditory neurons, 219
  - inferior colliculus, 206
  - lateral lemniscus, 19–21, 23
  - medial geniculate body, 32–37, 42
  - medial superior olivary nucleus, 15–19
  - olivocochlear system, 185–186
  - periolivary nuclei, 16, 183
  - primary auditory cortex, 2
  - superior olive, 119, 183
  - thalamocortical connections, 2, 35

- Cat (*Felis catus*) (*continued*)  
 ventral nucleus of lateral lemniscus, 119  
 visual system, 19  
*See also* Species
- CBA mouse, 560
- Cell death, excitotoxicity induced, 570
- Cell types (morphological)  
 cochlear nucleus, 10, 12  
 inferior colliculus, 75–78, 167, 169, 498, 631
- Cell types (physiological)  
 buildup, 11  
 chopper, 10  
 Class I to IV, 318, 328, 330, 332  
 excitatory-excitatory (EE), 517–518  
 excitatory-inhibitory (EI), 394, 517–519  
 I-type, 319, 438, 442  
 inhibitory-excitatory (IE), 393  
 O-type, 319, 332, 438–439, 442  
 pauser, 10, 606  
 primary-like, 10  
 V-type, 319, 324, 381, 438  
 tuning curves, frequency-response, 251  
*See also* Binaural, Monaural
- Central acoustic tract, 626–627  
 nucleus of, 204  
*See also* Medial geniculate body, subgeni-  
 culatate nucleus of
- Central nucleus of inferior colliculus  
 aging, 559–568  
 amplitude modulation, 349–361  
 behavioral function compared with the  
 dorsal nucleus of the inferior col-  
 liculus, 449  
 cell classes I to IV, 315, 318, 323, 326–  
 328, 330, 438  
 cell types, 167, 169, 319  
 I-type, 319, 330, 438  
 O-type, 319, 328, 330, 381, 438–  
 439, 443  
 V-type, 319, 330, 438, 443  
 central feedback, 632  
 changes in frequency gradient after  
 cochlear lesions, 577  
 characteristic frequency, 319  
 characteristic frequency, preservation  
 after acoustic trauma, 574  
 clustering of responses, 124  
 communication sounds, 415–418
- connections  
 banding of input, 91–93, 123, 538–  
 541, 548–550  
 cell-specific promoters, 126  
 cochlear nucleus, 2–3, 78, 115–116,  
 122, 183, 186  
 combinations of input, 125  
 convergence of input, significance of,  
 123, 125  
 dorsal nucleus of the lateral lemnis-  
 culus, 24, 79, 182  
 homogeneity of afferent terminal  
 fields, 125  
 intrinsic connections, 255  
 lateral superior olive, 28, 79, 94–95  
 medial superior olive, 28, 79, 91–94  
 ventral nucleus of the lateral lemnis-  
 culus, 79, 182
- core of central nucleus (ICCc) in owl,  
 520
- deficits, and the lateral superior olive,  
 443
- development, 537–552
- disc-shaped projection cells, 205, 538,  
 543
- effects of trauma, 575  
 excitatory input to, 249  
 flurazepam, effects of, 607
- frequency-band lamina, 320, 325, 330,  
 396, 510
- frequency gradient, 319–320, 324
- frequency tuning, 319
- functional modules, 86–88, 125
- glycinergic projections to, 255
- homology, 632
- importance in hearing, 627
- interaural time differences, cells sensi-  
 tive to, 430, 519
- intensity tuning in, 330
- lamina, fibrodendritic, 74, 81, 87, 89,  
 91–93, 124, 324, 328, 538–540
- lateral shell, ascending projections to,  
 in bird, 506
- medial shell, ascending projections to,  
 in bird, 506
- molecular biological approaches to dis-  
 secting function, 126
- neuron types, 74–78, 81, 87, 98, 205,  
 325
- neurotransmitters, 25

- nucleotopic organization, 124  
 overlap of ascending input, 123  
 owl, 519  
 periodicity coding, 352–361  
 physiology  
   binaural processing, 25, 96  
   eye movements, resulting from stimulation of, 444  
   facilitation, 326  
   monaural processing, 96  
   spectral processing, 25, 314, 325  
   stimulation of elicits seizures, 444  
   temporal processing, 104  
   tonotopic organization, 25, 88–89, 314, 320  
 projections from  
   contralateral inferior colliculus, 255  
   lateral lemniscus, 538  
   lateral superior olive, 538, 541  
   nucleus angularis in owl, 506  
   nucleus laminaris in owl, 506  
 projections to  
   cochlear nucleus, 186–188  
   contralateral medial geniculate body, 443  
   superior colliculus, 188–190  
 role in epileptic seizures, 603–613  
 role in localization of complex sounds, 432, 440  
 saturation in intensity coding, 330  
 segregation of ascending input, 78–82, 123–124  
 space map in the barn owl, participation in the construction of, 519  
 specificity of ascending input, 80–82  
 synaptic domains, 86–88, 91, 124  
 synaptic inhibition and frequency band laminae, 510  
 tonotopic organization of central nucleus in mouse, 560  
 tonotopy after cochlear lesions, 577  
 unmasking of responses after trauma, 574  
*See also* Inferior colliculus  
 Central pattern generator, 481  
 c-fos (or *c-fos*), 320, 573, 594  
 Characteristic delay (CD), 380, 383, 428  
 Characteristic frequency (CF), 313, 319–322, 494  
   cochlear nucleus, 6  
   cochlear root neurons, 190  
   discontinuities in after cochlear lesions, 577  
   expansion after cochlear damage, 577  
   focal damage and, 576  
   hair cell afferent neurons, 185  
   progression, 576  
   scatter and unmasking, 576  
   *See also* Isofrequency contours  
 Characteristic phase (CP), 380  
 Chemoarchitecture  
   acetylcholine, 501  
   aspartate, 16, 35, 33, 39  
   choline acetyltransferase, 16, 25, 501  
   enkephalins, 33  
   GABA ( $\gamma$ -aminobutyric acid), 16, 25, 33, 39  
   glutamate, 16, 33, 39  
   glycine, 16  
   serotonin (5-hydroxytryptamine), 39  
   somatostatin, 16  
 Chinchilla, connections of cochlear nucleus with inferior colliculus, 116, 118–119  
 Chiroptera, 493  
 Chopper, 16  
 Circuitry, cochlear nucleus, 12, 14  
 Claustrum, 43  
 CNQX (6-cyano-7-nitroquinoxaline-2,3-dione), 209, 253–254, 258, 294–296  
 Cochlea  
   ablation, 547–549, 595  
   central changes after ablation or inactivation, 569  
   changes in central firing rate after damage, 576  
   damage and effects on adult cats, 572  
   damage and effects on adult ferrets, 572  
   filtering, 327  
   lesions, basal cochlear, 571  
   lesions and maps of characteristic frequency, 576  
   olivocochlear input, 186  
   projections from olivocochlear system, 186  
   removal, effects of, 268, 547–549  
   specialization, in bats, 508, 510

- Cochlea (*continued*)  
 spiral ganglion cell lesions, 571  
 ventral cochlear nucleus, 183
- Cochlear nuclear complex. *See* Cochlear nucleus
- Cochlear nucleus, 5–15, 380  
 axons from, 81  
 bat, 503  
 brain stem projections, 117–123  
 branched axons, 120  
 bushy cell, 392  
 cartwheel cell, 12, 186  
 cat, 2, 6  
 changes in connections with age, 566  
 characteristic frequencies, 14  
 collaterals from olivocochlear neurons, 185  
 connections from cochlea, 13  
 contralateral input, 8–9, 20  
 convergence of input to, 431  
 divisions, 6  
 dorsal cochlear nucleus, 319  
 projections, 120, 382  
 end bulbs of Held, 13  
 fine structure, 13  
 fusiform cell, 120, 186  
 GABA, 7  
 GAD. *See* GABA  
 giant cell, 120  
 globular bushy cells, 12  
 glutamate, 7  
 glycine, 7  
 granule cell area, 12, 186  
 innervation, 12–13  
 interactions between ascending and descending input, 187, 319  
 intrinsic connections, 12, 14  
 isofrequency contours, 6–8  
 molecular layer, 186  
 monaural projections to inferior colliculus, 502  
 multipolar cell, 120  
 type I, 120  
 neuron types, 10  
 neuronal classes, 10  
 octopus cells, 10, 12, 120  
 olivocochlear innervation, 186  
 olivocochlear projections, 186–187  
 pauser/buildup, 10–11  
 primary synapses, 11  
 projections from inferior colliculus, 186  
 projections from lateral lemniscus, 20  
 projections from olivocochlear neurons, 186  
 projections from outer hair cells, 18  
 projections from periolivary cell groups, 17  
 projections to, 253  
 projections to inferior colliculus, 12, 20, 120, 380, 502  
 projections to lateral lemniscus, 12  
 projections to medial olivocochlear neurons, 185, 187  
 projections to superior olive, 12, 20  
 projections to ventral nucleus of lateral lemniscus, 12  
 spontaneous rate, 14  
 synapses, 13  
 topography of projections to inferior colliculus, 122  
 transmitters, 5  
 type II, 14  
 type IV, 14  
*See also* Anterior ventral cochlear nucleus, Dorsal cochlear nucleus, Ventral cochlear nucleus
- Cochleotopic map, 561
- Code, array, 519
- Coding, average-rate, likelihood of, 334
- Coding, independence of biological and intensity coding, 334
- Coincidence model, 267, 380, 382
- Coincidence and sinusoidal frequency modulation tuning, 517
- Columnar organization of lateral lemniscal nuclei, 503
- Combination sensitivity, 352, 416–417, 420–422, 511–512  
 echolocation and, 514  
 harmonically unrelated tones and, 514
- Commissure of inferior colliculus, 156  
 anatomy of, 157  
 axonal plexus, 161  
 comparative studies of the commissural axonal plexus, 159  
 comparative studies of commissural input, 156

- congruence with morphological and neurochemical diversity, 169  
 contribution to fibrodendritic laminae, 166  
 distribution of commissural axons, 159–161  
 effect of GABAergic cells on commissural transmission, 250  
 excitatory, 174, 176  
 fibers in, 157–158, 162  
 function, 172  
 homotopic projections, 166  
 hypothesis on function of, 166  
 inhibitory, 174  
 internal organization, 156–158  
 laminar organization, 166  
 location, 156  
 morphological diversity in form of cells of origin, 169  
 neurotransmitter, 174  
 other targets of commissural cells, 169  
 polysynaptic influences on, 174  
 Probst, commissure of, 262, 289  
 projections, 173, 255  
 richness of commissural projections, 169  
 sources of commissural axons, 156  
 spatial distribution of commissural cells, 172–173  
 tectal commissural column, 170  
 topography of commissural axons, 157, 159, 166  
 transmitters, 172  
 types of commissural neurons, 169  
 Commissural column, tectal, 170  
 Communication sounds, 411–422, 460–461, 494  
 Comparative studies  
   bats and birds, 524–525  
   behavior, 524–525  
   best delays, 386  
   brain stem afferents, 118–119  
   cytoarchitectonics, 630–631  
   filter properties among bats, 510  
   inferior colliculus, structure, 159, 628–632  
   physiology, medial geniculate body, 208–211  
   sound localization, 384–386, 390, 510  
   subdivisions, 629–632  
   superior olive, 118–119  
   torus semicircularis, 459–483, 498  
 Composite curves. *See* Interaural differences  
 Compound action potential (CAP), 571  
   amplitude after lesions, 572  
   decreases in aging, 571  
   decreases after lesions, 576  
   unmasking and, 576  
 Conditioned changes, establishment and duration of, 518  
 Conditioned responses, plasticity of, 449  
   influenced by electrical stimulation, 449, 518  
   influenced by pharmacologic inactivation, 449  
 Computational models in spectral processing, 433  
 Connectivity, age-related changes in, 566  
 Consonance to dissonance transition, 324  
 Constant frequency-frequency modulated (CF-FM), 494, 497, 510  
   quasi-constant frequency (CF) calls and echoes, 510  
 Context-dependency  
   affecting spatial receptive fields, 518  
   affecting tone responses, 634  
   stimulus statistics, 634  
 Contrast sensitivity, 333  
 Convergence, 328, 332, 377, 504, 510  
 Convergence of excitation and inhibition with different latencies and discharge patterns, 514  
 Corpora quadrigemina. *See* Inferior and Superior colliculus, Midbrain  
 Corpus callosum, 4  
 Corpus striatum, 42–43, 45  
 Corticocollicular system, 165, 232–234, 236–237, 250, 504  
   axons, 165, 233, 235  
   function, 236–237  
   neurons, 232, 239  
 Corticocortical connections, 174  
 Corticofugal projections, 39, 41–43, 174, 234, 236–237  
 Corticothalamic projections, 207, 236  
 Count-comparison schemes. *See* Interaural differences

- CPP ( $\pm$ )-3-(2-carboxypiperazin-4-yl)-propyl-1-phosphonic acid, 254, 250
- Critical bands, 29, 91, 324, 327–328, 330
- Cross correlation functions, 387–388
- Current-clamp, 253
- Current-pulse stimulation, 253
- Cutoff, upper, 564
- Cyclic AMP, (cAMP), cyclic adenosine monophosphate, 571
- Cytoarchitecture  
 inferior colliculus 72, 496  
 lack of obvious subdivisions in inferior colliculus, 496, 628–632
- Cytochrome oxidase, 502, 594–595
- Damage**  
 central aspects of, 573  
 competing peripheral and central effects of, 573  
 unmasking of low-frequency input, 573
- Deafferentation, 585
- Deafening, juvenile, 567
- Deafness. *See* Hearing loss
- Decerebration, 319, 400, 440, 450
- Delay, half maximal, 447
- Delay functions, 384
- Delay lines, 382  
 and axons to the medial superior olive, 387  
 graded delay lines, 388  
 phase changes, role in generating delay lines, 387–388  
 tuning and coincidence detectors, 514  
 tuning and combination-sensitive neurons, 511–512
- Delays, best, 382–383, 512, 514
- Descending control, 237
- Descending projections,  
 amygdalocollicular connections, 240, 241  
 corticocollicular connections, 234, 241, 334, 449  
 feedback, 236  
 from inferior colliculus, 182, 193  
 thalamotectal connections, 241, 334
- Desensitization, peripheral, 573
- Deoxyglucose, [ $^{14}$ C]-deoxyglucose, 73, 320, 356, 603
- Direction-dependent tuning, 263
- Detection criterion, signal,  $d'$ , 435
- Development. *See* Ontogeny
- Directional acuity, 435
- Directional hearing, 441, 588–589
- Disc-shaped cells, 74–78, 87, 98, 289, 497, 538, 540  
*Kv1.1* potassium channels on, 498  
 synapses on, 121
- Discrimination of tones in noise, 324
- Dorsal cochlear nucleus, 6–7  
 anatomy, 10, 12  
 contribution to ITD and ILD sensitivity, 382  
 concentration within inferior colliculus, 504  
 fusiform cells, 249  
 buildup, 249  
 nonmonotonicity, 249  
 pauser-buildup, 249  
 rate-intensity function, 249  
 temporal locking, 249
- GABA, 6  
 giant cells, 10, 120, 125  
 inferior colliculus projections, 85, 117–120  
 innervation, 382  
 lateral lemniscus, 118  
 neuron types, 7–11  
 physiology, 328, 438  
 projections from superior olive, 184  
 projections from inferior colliculus, 186, 188  
 projections to inferior colliculus, 266, 504  
 role in localization of complex sounds, 432, 438, 440–441  
 spectral integration in, 431  
*See also* Cochlear nucleus
- Dorsal cortex, of the inferior colliculus, 84–85, 122, 257, 320–321, 330, 449
- Dorsal nucleus of lateral lemniscus, 19–21, 139–142  
 defect in GABAergic inhibition, 610  
 GABA, 19, 139, 141, 184

- interaural time differences, cells sensitive to, 430
- ipsilateral and contralateral projections, 163
- mechanisms for sharpening interaural time difference functions, 389
- posterior part (LLDp) and comparison with lateral superior olive, 519
- projections from, 20–21, 139, 163, 183, 502
- projections to, 19, 21, 139
- Dorsomedial periolivary nucleus (DMPO)
  - comparative, 138
  - comparison with superior paraolivary nucleus, 138
  - projection to inferior colliculus, 138
- Drugs, ototoxic, 571
- Drugs, ototoxic, and laterality differences, 571
- Duplex theory, 427
- Duration selectivity, 266, 316
- Duration sensitivity, 512
- Duration tuned neurons, localization of, 514
- Duration tuning, 140, 514
  - abolition of, 514
  - latency and, 515
  - stimulus offset and, 516
- Dynamic range, 258, 330
- Ear-plugging, effects on azimuthal sensitivity of binaural cells, 217, 523
- Echo clutter, signal detection in, 517
- Echo suppression, 448
- Echo threshold, 446
- Echoes and paradoxical latency shift (PLS), 514
- Echoes, role in perceptual salience, 378
- Echolocation
  - bat, 219, 449, 494, 506
  - calls, 495, 497
  - compressive reorganization of biosonar signals, 449
  - delays, 364–366
- Efferent innervation
  - organ of Corti, 185, 187
  - cochlear nucleus, 183
- Efferent synapse
  - outer hair cells, 185
- Efferent system. *See* Olivocochlear system
- Eighth nerve. *See* Auditory nerve
- Elevation effects, illusory, 436
- Endbulbs of Held, 11, 13, 45, 430
- Enhancement of response, 517
- Envelope
  - coding, 104, 249, 258, 361, 392–393, 397
  - fluctuations, 562
  - periodicities, coding with aging, 564
  - sinusoidal fluctuations, 258, 562
- Epilepsy, 176, 444
  - genetically epilepsy-prone rats (GEPR) 444, 610
- Evoked potential, 2, 571
  - amplitude after hearing loss, 571–572
  - enhancement, 572, 575
  - unmasking and evoked potential amplitude, 575
- Evolution, 46, 482, 493, 628
- Excitatory-inhibitory interactions, models of, 261–262
- Excitatory monaural intensity (EMI), 393
  - See also* Average binaural level (ABI)
- Excitatory postsynaptic current (EPSC), 103, 253, 296, 300
- Excitatory postsynaptic potential (EPSP), 174, 253, 382, 512, 514
- Excitatory rebound, 512, 515
- Excitatory spectrotemporal receptive fields (sSRF), 313, 315, 319
  - among brain stem neurons, 328
  - diversity of shapes, 328
- External cortex, 82
- External nucleus of the inferior colliculus, adaptive changes in, and plasticity of, 523
  - computation of interaural time difference sensitivity, 519
  - construction of the space map in the barn owl, participation in, 519
  - dominance of visual map within, 524
  - frequency representation, 320
  - frequency tuning, 316, 322
  - intrinsic projections, 167–168

- External nucleus of the inferior colliculus  
*(continued)*  
 methodological issues in interpreting projections, 167  
 projection from the cochlear nucleus, 122  
 projection to central nucleus, 167–168  
 projection to superior colliculus (optic tectum) in barn owl, 522  
 relation to pericentral nucleus, 497  
 role in startle response (lateral nucleus), 190  
 spatial plasticity, and 424  
 spatially restricted receptive fields in, 522  
 ventral region, specialization for upward sweeps, 216  
 visual instructive signal, 524  
 vocalization, 418
- Extraauditory forebrain centers, 241–242
- Facial motor nuclei, 189–190
- Facilitation, 104, 214, 315, 396, 512, 513
- Feedback control, 231–235
- Ferret, connections of cochlear nucleus with inferior colliculus, 116, 118–119
- Field L  
 bird, 505
- Filter bandwidth, 313
- Filtering, multitask, 324
- Filtering, spectral, 312, 328
- Filtering of space-ambiguous side peaks, 519
- Filters  
 band-pass, 322, 474, 564  
 filter-type frequency response areas, 510
- First (fundamental) harmonic, 512
- Fish  
 hearing, 460–467  
 forebrain, 463  
 lateral lemniscal nuclei, 463  
 thalamus 463–464
- Flat cells, 76–78, 102, 289
- Flight mechanisms, of birds and bats compared, 493
- Flunitrazepam, 251–252, 270
- FM. *See* Frequency modulation
- fMRI (functional magnetic resonance imaging), 214
- Forebrain  
 archistriatum, 498  
 avian, 505  
 vocal pathways in, 493
- Formants, 322, 413
- Fos protein, 73, 89, 320, 573  
*See also* c-fos, c-fos
- Fourier analysis, 312
- Fovea, auditory, 7, 497, 508, 510
- Free field, 445
- Free radicals, 567
- Frequency  
 best, 313  
 characteristic, 313, 319–322, 494  
 dependence of best delay upon, 384  
 selectivity, 313  
 spectrum, 517
- Frequency analysis  
 phase versus frequency plots, 380
- Frequency-band  
 laminae, 320  
 processing, across, 325  
 processing, within, 325
- Frequency discrimination, 324
- Frequency domain, effects on sound localization, 437
- Frequency gradient, 218, 322
- Frequency modulation (FM), 361–364, 469, 495, 508, 511  
 amplitude, 512  
 bat, 266  
 directional, 266, 362  
 FM1–FMn, 514  
 FMn, 512  
 mechanisms, 266, 363  
 sweep vector influences, 516  
 sweeps, 516
- Frequency perception and sound source elevation, 433
- Frequency representation  
 complete, 322  
 physiological ambiguity, 320
- Frequency resolution gradients, 324
- Frequency-response map, 262  
 tuning curves, 260
- Frequency tuned inhibitory response, 440



- Frequency tuning, 267  
 development, 543–544
- Frequency tuning curve, excitatory, 313
- Fundamental frequency, 348, 413, 416, 446
- Fusiform cells  
 avian inferior colliculus, in, 498  
 axonal projections in dorsal acoustic stria, 120  
 comparison with bushy cell, 13  
 dendrites as synaptic targets, 186  
 dorsal cochlear nucleus, 11  
 role in spectral and temporal transformation of auditory nerve activity, 13  
 synaptic inputs, 13  
 targets of inferior colliculus descending input, 186
- GABA**, 17, 28, 87, 97, 205, 260, 263, 297–300, 390, 482, 502, 516, 544–550, 566, 569, 573, 592, 594, 606–610  
 absence of effects on an inferior colliculus subpopulation, 575  
 auditory cortex, 40, 633  
 bat, 34, 497  
 benzodiazepine, interactions with, 607  
 bicuculline, 255, 261–267, 299–300, 349, 608–609  
 binaural stimulation and, 608  
 bird, 522  
 blockade of, and normal responses, 524  
 cochlear nucleus, 6  
 cochleotomy and effects on, 569  
 commissural projection, 172  
 damage and, 573  
 decreased release after damage, 573  
 dorsal cochlear nucleus, 6  
 effects of adapting GABA input on static and dynamic differences, 390  
 effects of age, 268  
 effects of cochlear removal on, 569  
 effects of peripheral trauma, 575  
 flurazepam, effects of, 607  
 gradients in inferior colliculus, 497  
 inferior colliculus, 206, 250, 297–300, 497, 522, 544–550  
 lateral superior olivary nucleus, 133  
 levels in inferior colliculus, 250, 570  
 mechanisms for sharpening interaural time difference functions, 389  
 medial geniculate body, 208  
 medial nucleus of trapezoid body, 145, 380  
 medial superior olivary nucleus, 134, 249, 396, 504  
 mediates inhibitory responses, 257, 318  
 membrane and cytosolic fractions, effects of peripheral damage on, 569  
 primary auditory cortex, 39  
 projections to the inferior colliculus, 254, 606  
 receptors, 103, 297–300  
 recovery times and, 400  
 release, Ca<sup>2+</sup> dependent, 251, 298  
 release, K<sup>+</sup>-evoked, 566  
 release, presynaptic, 257  
 transaminase, 251  
 transport mechanisms, 606  
 thalamus, 34  
 uptake after peripheral trauma, 570
- GABA<sub>A</sub>**, 208, 251, 260, 262, 266–267, 269, 297–298, 497, 523, 566, 569  
 antagonist, 572–573, 610  
 postsynaptic receptors, 255, 269, 297, 497  
 subunit, 251, 269, 298
- GABA<sub>B</sub>**, 208, 253, 255–257, 267, 269, 298–300, 568–569  
 presynaptic receptor, 267, 270, 298–300, 497
- GAD**, 250, 268, 569, 594  
*See also* GABA
- GAD65** (glutamic acid decarboxylase), 569
- GAP43** (growth associated protein), influence on synaptogenesis, 570
- Gaps**  
 coding of, 566  
 detection thresholds, 562  
 post-gap noise burst, 562  
 single-unit firing rate, 562  
 stimuli, 560  
 thresholds in aging, 562

- Gates, AND, for interaural time and level difference processing, 522
- Gerbil, connections of cochlear nucleus with inferior colliculus, 116, 118–119
- Giant cell, dorsal cochlear nucleus, 12, 120, 186–187
- Globalular bushy cell, cochlear nucleus, 12
- Glutamate  
 agonists, 611  
 agonists enhance audiogenic seizure susceptibility, 611  
 antagonists, 613  
 antagonists and enhancement of seizure susceptibility, 611  
 changes in inferior colliculus subdivisions with trauma, 570  
 cochlear nucleus projections to the inferior colliculus, 497  
 effects of cochlear removal on, 570  
 inferior colliculus, 172, 208, 248, 256–257, 294–297, 431, 497, 544–546  
 medial geniculate body, 208  
 receptors, 103, 256
- Glutamic acid decarboxylase (GAD), 250, 268, 569, 594  
*See also* GABA
- Glycine  
 and binaural processing, 517  
 and long term potentiation, 569  
 cochlear nucleus, 6, 13  
 concentration in inferior colliculus, 570  
 delay tuning, relation to, 514, 516  
 inferior colliculus, 300, 544–546  
 lateral superior olivary nucleus, 133  
 medial nucleus of trapezoid body, 16  
 medial superior olivary nucleus, 133  
 periolivary cell groups, 16  
 periolivary nuclei, 16  
 projections to central nucleus of the inferior colliculus, 255, 431  
 receptors, 570
- Golgi method, 167, 500, 629, 631
- Golgi type II cell, 12, 99–100, 633
- Granule cells, 12
- Growth cone, 542
- Guinea pig (*Cavia porcellus*)  
 hearing loss in, 572  
 lateral nucleus, 30  
 sound localization in, 383–385  
*See also* Rodent
- Hair cell**  
 inner, 330  
 outer, 185
- Head orientation, 436
- Head-related transfer function (HRTF), 378, 432–433, 436, 438  
 directional information for azimuth and elevation, 432  
 domains of spectral information, 432  
 owl, 521
- Head width and best delays, 383
- Hearing loss  
 bilateral, 585–586, 595–596  
 c-fos, 594  
 compensatory mechanisms for, 613  
 conductive, 585–586, 594–595  
 evoked potential amplitude in, 571  
 flat, 586  
 frequency map, 587–588  
 GABA, effects on, 592, 594–595  
 high-frequency, 572  
 high-frequency and aging, 559  
 inhibition, diminished, 590–593, 597–598  
 inferior colliculus, 585–598  
 masking, 589–560  
 noise-induced, 585, 591–593  
 ototoxic-induced, 570, 573  
 permanent, 569, 571  
 plasticity, 586, 597–598  
 pre-pulse inhibition, 596–597  
 profound, 585  
 pseudo-plasticity, 590  
 rate-level functions, 588  
 sensorineural, 585–591  
 sloping, 560  
 transsynaptic changes, 590  
 unilateral, 593–594
- Hearing onset, 534–544, 547
- Held, stria of, 4, 9, 439–440
- Hierarchical midbrain organization in owls, 525

- High-frequency edge, effects of damage on, 573
- High frequency spectral contrast, and minimum audible angle (MAA), 435–436
- Homology, 459, 493, 626–631, 659
- Horseradish peroxidase, 75, 119, 164, 566
- Huggins pitch, 400–401
- Human
  - auditory cortical fields, 41, 43
  - hypothalamus, lateral, 498
  - inferior colliculus, 630
  - interaural time difference processing, 377
  - lateral lemniscus, 630
  - medial nucleus of trapezoid body, 17
  - superior colliculus, 630
- $I_h$  current, 210
- Inactivation and lesion studies, limitations of, 426
- Inferior colliculus
  - association projections, 160–161
  - axons from cochlear nucleus, 86–88
  - bands, 147, 161–162, 321
  - bat, 21, 27, 496
  - best frequency, 508
  - binaural input, 30, 504
  - binaural pathways and masking level differences, 398
  - cat, 630–631
  - cellular organization, 21–31, 74–83, 630–631
  - central nucleus, 2–3, 24, 74, 155, 283, 449, 560, 631
  - combination sensitivity, compared in inferior colliculus and medial geniculate body, 216
  - commissure, 24, 27, 156–158, 250
  - comparative, 21, 207, 626–631
  - comparison with the medial geniculate body, 200, 213, 215
  - connections, 28, 117
  - convergence of projections, 123, 253, 503–504
  - cortex, 25
  - crossed ascending projection from, 117
  - descending connections, 182–188
  - development, 537–552
  - dorsal cortex, 25, 30–31, 84–85, 163
  - dorsomedial subdivision, 497
  - dorsoposterior subdivision, 497
  - echolocation, 21
  - frequency bands, 24, 26, 87, 321
  - frequency band-specific tectothalamic projections, 87, 202
  - GABA, 24, 27, 29, 99–100, 297–300
    - role in frequency tuning, 215
  - GABA, sources to inferior colliculus, 251
  - GABAergic neurons, 250
  - GABAergic projections to the medial geniculate body, 205
  - glycine, 300
    - development, 544–546
  - human, 630
  - inhibitory bands, 100
  - inhibitory processing, 29, 100, 205, 318, 390, 544–546
  - intercollicular area, 501
  - interneurons, 633
  - intrinsic projections, 160, 163–165, 169, 266, 390
  - lamina, 74, 81, 87, 89, 122, 161, 320–321, 325
  - lateral cortex, 82–83
  - lateral nucleus (external cortex), 25, 30, 163–164
  - local connections, 29, 163
  - major divisions, 24
  - monaural cells and sideband inhibition, 215
  - monaural pathways, 30, 215
  - monotonic, 332
  - mouse, 320, 323
  - natural amplitude statistics, 333
  - neural processing, 26, 30
  - neuron types, 29, 74–78, 87–89, 167, 169
  - nexus, 627
  - non-human primate, 630
  - nonmonotonic cells, 333, 393–395
  - number of neurons, 155
  - obligatory synaptic station, 155
  - operating range, 333

Inferior colliculus (*continued*)

- pericentral area, 496
- projections from anterior ventral cochlear nucleus, 27
- projections from auditory cortex, 24, 30, 165
- projections from the amygdala, 504
- projections from brain stem, 115
- projections from GABAergic sources, 254
- projections from the globus pallidus, 504
- projections from inferior colliculus, 24, 35, 205
- projections from lateral lemniscus, 24, 117, 142–144
- projections from medial superior olive (MSO), 377
- projections from midbrain, 173
- projections from superior olivary nucleus, 24, 28
- projections from substantia nigra, 501
- projections from superior olive, 15–19, 24, 28, 91–95, 142, 377
- projections from superior paraolivary nucleus, 165
- projections from telencephalon, 232–241
- projections from ventral cochlear nucleus, 9, 12, 27, 118–120
- projections to cochlear nucleus, 20–21, 186
- projections to lateral lemniscus, 182–183
- projections to lateral (external) nucleus, 122
- projections to medial geniculate body, 27–28, 35, 200, 202–203, 524
- projections to medial geniculate body, crossed, 202
- projections to the superior colliculus, 504
- projections to superior olivary complex, 184
- projections to thalamus, 202
- ripple spectra, 333
- sharpness of tuning, 215, 323
- somatosensory projection to, 30

- spectral features, 215, 315
- spectral integration and two-tone stimuli, 215
- speech encoding, 418–420
- steroid binding sites, 498
- subdivisions, 70–86, 155
- synapses, 79–80
- synaptic domains, 86–88, 91, 148, 334
- temporal coding, 346–368
- thalamocortical auditory system, 96
- tonotopic organization, 29–30
- tonotopic organization, age-related changes in, 560
- topographic projections, 117
- transformation from inferior colliculus to cerebral cortex, 214
- transmitters, 25
- vocalization encoding, 411–422
- See also* Central nucleus of inferior colliculus,
  - Torus semicircularis
- Inflammatory response, and stress, 567
- Inheritance, physiological, 328, 517
- Inhibition
  - absence of effects on an inferior colliculus subpopulation, 575
  - age-related decline in, 564, 606
  - binaural, 610
  - contribution to modulation transfer functions, 564
  - decreased suprathreshold, 575
  - delay tuning, 514
  - dose-related, 610
  - one-tone, 315
  - release from, 265, 568
  - sculpturing effects on eSRFs, 328
  - sinusoidal frequency modulation tuning, 517
  - synaptic, 510
  - two-tone, 315, 318, 522
- Inhibitory amplification, 255
- Inhibitory areas, 313
- Inhibitory postsynaptic potential (IPSP), 174, 254, 298
- Inhibitory projections to the inferior colliculus, changes with age, 566
- Inhibitory responses, frequency tuned, 440

- Inhibitory sidebands, 315, 522, 591, 627
- Input-output transformations, 11, 211, 215, 312, 346, 473
- Insectivores, relation to Chiroptera, 493
- Intensity coding, 313, 329–334, 428  
 cues, importance of, 428  
 development, 543  
 dynamic range, 329  
 independence from biological coding, 334  
 loudness, 324, 329, 378, 400, 494  
 models of, 334  
 neural coding of intensity, 329  
 nonmonotonic rate-level coding, 334  
 onset envelope, 329  
 population coding, 329  
 pressure envelope, 329  
 psychoacoustical tests, 329  
 response sensitivity, 329–330  
 selectivity, 330  
 small group coding, 334  
 spread of excitation, 329  
 threshold, response sensitivity, 329
- Interactions of ascending inputs in bat inferior colliculus, 517
- Interactions sculpted from excitatory spectrotemporal receptive fields (eSRFs) by inhibition, 328
- Interaural click delays (ICDs), 448
- Interaural correlations  
 changes in, 393  
 normalized, 393
- Interaural level differences (ILDs), 377–378, 393, 396, 427–428, 502, 519  
 AND gates for interaural time and level difference processing, 522  
 birds, 502, 522  
 circuits in, 429  
 coding of, 522  
 coding with excitatory-inhibitory (EI) cells, 519  
 effects of lesions of superior olivary complex, 396  
 frequency and, 520  
 functions of, 395  
 gradients, 517  
 local inhibition and, 396  
 negative, 522  
 population coding, relation to, 395  
 sensitivity of inferior colliculus central core neurons to, 518  
 sensitivity to changes in, 518  
 shaped by diverse supraolivary mechanisms, 396  
 spikes and postsynaptic potentials, and interaural time and level processing, 522  
 topography of, 520  
 transition region, 395  
 types of interaural level difference curves, 394
- Interaural level disparities (ILDs), 430
- Interaural phase differences (IPDs), 390, 428, 519
- Interaural time delays, static and dynamic, 390
- Interaural time differences (ITDs), 377–379, 427–428, 504, 521  
 ambiguous peaks, 522  
 amplitude envelope, 504  
 at different levels of the auditory system, 388, 430  
 at low frequencies, 384  
 azimuth, 519  
 best, 380, 382  
 best delays at zero ITD, 379  
 binaural masking level differences, relation to ITD, 396  
 broad interaural time difference functions evoked by envelopes, 393  
 cancellation and filtering of ambiguous peaks, 522  
 characteristic phase, 430  
 circuits in, 429  
 coding, 382  
 comparative studies, 392  
 compatibility with behavioral studies, 385  
 composite curves, 384  
 computations, 522  
 count-comparison models, 385–387  
 created by inhibition, 388  
 cyclic functions, 378  
 dynamic and static ITD sensitivity, 391  
 effects of anesthesia upon, 390  
 ensemble coding of, 520

- Interaural time differences (ITDs) (*cont.*)  
   envelopes and dynamic interaural level  
     sensitivity, 392–393, 397  
   firing rate variations and, 385  
   functions, 383, 389  
   GABAergic influence on, 263, 388  
   head width, 382  
   high frequency carrier, 504  
   long, and relation to best delays, 383  
   low-pass filters, parallels with, 393  
   negative, 522  
   neural code of, 387  
   nonphysiological delays, trough-type  
     neurons with, 393  
   optimal sharpening for ITD in the me-  
     dial geniculate body, 390  
   overlap of functions, 382  
   peak, 381, 428  
   peaks, ambiguous, and filtering, 522  
   peaks within the physiological range,  
     385  
   perceptual ambiguity and, 377  
   prism experience, 523  
   processing, 506  
   processing in high frequency enve-  
     lopes, 391  
   range of ITD sensitivity, 392  
   remodeling of, 523  
   sensitivity to sound fine structure, 392  
   slope, 385, 429  
   spikes and postsynaptic potentials, and  
     interaural time and level process-  
     ing, 522  
   species differences in, 290, 392  
   species-specific sensitivity, 506  
   types of composite curves, 379  
     intermediate type, 379–380  
     peak and trough compared across  
       auditory nuclei, 388–389  
     peak-type, 379–380, 389  
     trough type, 379–380, 389, 393  
     vertical asymmetry, aural, 519
- Interclick delay (ICD), 447  
 Intercollicular area, of the inferior colli-  
   culus, 84, 501, 512  
 Intercollicular tegmentum, projections  
   from the archistriatum, 504  
 Interaural intensity differences, 250  
 Interaural phase differences, 250
- Intermediate nucleus of lateral lemniscus,  
   139, 141  
   bat, 21  
   cat, 19  
   projections, 141  
 Interneuron (Golgi type II cell), 27, 82,  
   98–100, 205, 207, 250, 390, 633  
 Intracellular filling, 27, 81, 169, 252  
 in vitro brain slices, 253  
 in vivo recordings, 257  
 Ion channels, 290–292  
   Ca<sup>2+</sup>, 293  
   Ca<sup>2+</sup>-activated K<sup>+</sup>, 285, 288, 292  
   K<sup>+</sup>, 283, 290–292  
   Kv channels, 290–292  
 Isofrequency contours, 81, 86–90, 164,  
   320, 325, 360, 507, 541, 631  
   cochlear nucleus, 6  
   internal organization, 123–125  
   sheets, 320  
 Isthmo-optic nucleus, avian, 499
- Jeffress model**, 16, 380–381, 384, 428  
**Jeffress model**, alternatives to, 385, 388  
**Just noticeable difference (JND)**, 382
- K<sup>+</sup>-GABA release**, 566  
   Kv, calcium channel subunit, 497  
**Koniocortex**. *See* Primary auditory cortex  
**Kynurenic acid**, 174–175
- Laminar nature of extrinsic projections**,  
   166  
   primary auditory cortex, 37, 39  
**Laminar organization**  
   central nucleus of the IC, 81, 87–89,  
     122  
   dorsal cortex of the IC, 84  
**Language areas of brain**, 44, 47  
**Latency**, effects of amplitude or spectro-  
   temporal changes on, 518  
**Latent input**, unmasking of, 568  
**Lateral cortex**, 70, 73, 82–83  
**Lateral geniculate body**, 43  
**Lateral inhibition**, 90, 260, 262, 395  
**Lateral lemniscus and lateral lemniscal  
   nuclei**, 20, 138–142  
   axons of, 85

- cat, 19–21, 23
- complex signal processing, 21
- GABA, 138–140
- glutamate, 140
- glycine, 138–140, 504
- major functions, 21
- participating in timing, 140
- physiologic organization, 23
- projections from cochlear nucleus, 139
- projections from superior olivary nuclei, 4, 139, 145
- projections in birds supporting interaural time differences, 506
- projections to cochlear nucleus, 4, 117
- projections to inferior colliculus, 140–142, 506
- projections to medial geniculate body, 204
- ventral nucleus of the lateral lemniscus, 205
- ventral nucleus of the lateral lemniscus, subdivisions, 138
- Lateral line, 461
- Lateral nucleus of inferior colliculus. *See* External nucleus
- Lateral nucleus of trapezoid body
  - projections to cochlear nucleus, 18
  - See also* Superior olive
- Lateral olivocochlear system
  - definition, 16
  - projections to organ of Corti, 16–17
- Lateral shell of owl inferior colliculus, 502
  - computation of interaural time difference sensitivity, 519
  - integration of interaural time and level information, 518
- Lateral superior olivary nucleus, 134, 249
  - aspartate, 15
  - bat, 503
  - binaural information, 250, 428
  - binaural input, 15, 94–95
  - calyx of Held, 430
  - cat, 16–19
  - choline acetyltransferase, 16
  - GABA, 16, 135
  - glutamate, 431
  - glycine, 15–16, 18, 431
  - Jeffress model, 16, 428
  - lateral lemniscus, 15
  - location, 16
  - low-frequency neurons, 16
  - mice, 18
  - neuron types, 15
  - physiology, 15–19
  - projections from anterior ventral cochlear nucleus, 15
  - projections from medial nucleus of trapezoid body, 430
  - projections from ventral cochlear nucleus, 8–10, 12, 115
  - projections to, 2–4
  - projections to inferior colliculus, 18, 22, 28, 121, 134
  - projections to lateral lemniscus, 22
  - sound localization, 17, 428, 430, 506
  - spatial tuning, 396
  - tonotopic organization, 16
  - transmitters, 16, 431
  - trough-type cells, 380
  - See also* Superior olive
- Lateral suppression, 317, 327
- Laterality differences, and drugs, 571
- Lateral tegmental system, 204
- Layers
  - central nucleus of inferior colliculus, 26, 29, 161, 538, 631
  - dorsal cortex of the inferior colliculus, 84, 161, 234–235, 242, 320
  - primary auditory cortex, 37, 39
- Learning
  - descending control of plasticity, 41
  - effects on midbrain function, 450, 518
  - medial geniculate body, 237
  - primary auditory cortex, 30
  - regulation of, 523
- L-enkephalin, 34, 35, 136, 174, 480, 502, 546
- Lesion projection zone, 576
  - discontinuities within, 576
  - plasticity and, 578
  - residue of pre-lesion input, 578
- Lesions, acoustic striae, and localization, 440
- Less-flat cells, 76–78, 102, 289
- Level tolerant tuning, 381, 472, 509–510, 512, 574

- Listening, passive, 494
- Location, conditions for optimal, 43
- Localization dominance, 445
- Long pass, 513
- Long-term potentiation (LTP), 257, 567, 569
- Low-frequency edge, effects of damage on, 573
- Low-pass, 564
- LTP. *See* Long-term potentiation
- Magnesium**, 254
- Magnocellularis, nucleus, 506
- Map**
- auditory, 506
  - calibration of, 506
  - cochleotopic, 561
  - creation of by axonal divergence, 5
  - registration of auditory and visual, 506
  - role in sound localization, 506
  - space, 506
  - visual, 506
- Marsupial**
- connections of cochlear nucleus with inferior colliculus, 116
  - northern native cat (*Dasyurus hallucatus*), 541–542
  - opossum (*Didelphys marsupialis*), 542, 631
- Masking**, 324, 564
- level differences, binaural, 396
- Masker**, narrow band, 324
- Medial geniculate body**
- anesthesia effects on neural discharge patterns, 211–212
  - APV (aminophosphonovalerate), 209
  - avian homologue, 506
  - bat, 34, 204
  - bat sonar signals, 204
  - best frequency, 201
  - binaural processing, 32–33, 36, 382
  - brachium of the inferior colliculus, 208
  - brain stem projections to, 200–205
  - burst firing mode, 209
  - burst-to-tonic shifts, 210
  - carnivore, 238, 242
  - cat, 32–36, 238, 242
  - characteristic frequency (CF) representation, 218–219
  - combination sensitivity, compared in inferior colliculus and medial geniculate body, 216
  - comparative connections, 204
  - comparative cytoarchitectonics, 32
  - comparative mammal, 202–205
  - comparative physiology, 213–220
  - comparative structure, 24
  - compared to inferior colliculus, 32, 200, 204, 214, 218
  - cytoarchitectonic subdivisions, 32–36, 201
  - cytoarchitecture, 200–203
  - delays, axonal and synaptic, 212
  - discharges during sleep, 211
  - dorsal division, 33, 36, 201, 204–205, 210
  - dorsal division, presence of higher harmonics, 219
  - dorsal division, response organization, 219
  - dorsal division, systematic organization of delay tuning, 219
  - EX/O neurons, 214
  - facilitation, compared in inferior colliculus and medial geniculate body, 216
  - frequency modulation, 216–217, 219
  - GABA ( $\gamma$ -aminobutyric acid), 34
  - GABA, influences on latencies, 212
  - GABA<sub>A</sub>, 208, 214
  - GABA<sub>B</sub>, 208
  - glomerulus, 34
  - glutamate, 33, 214
  - Golgi type II cell, 34, 207
  - gradients within an isofrequency band, 219
  - $I_h$  current, 210
  - independence of characteristic frequency gradients, 219
  - inferior colliculus projections to, 28
  - interaural phase disparities sensitivity comparisons, 214
  - interaural time differences, cells sensitive to, 430
  - in vitro and in vivo firing modes compared, 210
  - in vitro preparations, 208
  - innervation, 205–208



laminae, 32, 201  
 latencies, 212–213  
 lemniscal and nonlemniscal inputs, 200–205  
 location, 33  
 medial division, 33, 36, 201, 204–205, 220  
 medial division, tonotopic organization in, 220  
 medial division, tuning properties in, orthogonal to tonotopic axis, 220  
 modulation, effects on responsiveness, 214  
 modulation rate transfer functions, 213  
 monaural processing, 217  
 multisensory responses, 220  
 neuron intrinsic properties, 210  
 neuron tuning, 215  
 neuron types, 32–33, 36, 201, 208  
 nonauditory projections to, 204–205  
 NMDA receptors, 214  
 nonhuman primate, 34  
 offset (OFF), 212–213  
 onset (ON), 212–213  
 ovoidalis, nucleus, 505–506  
 paired-pulse depression mediated pre-synaptically, 214  
 paradoxical sleep, 211  
 physiology, 36, 201  
 possible role in ITD computations and optimal sharpening, 390  
 precision of information transfer in burst mode, 211  
 processing, comparing the effects of input and those of computation, 217  
 projections from contralateral inferior colliculus, 443  
 projections from inferior colliculus, 28, 122, 203, 205  
 projections to amygdala, 201  
 projections to auditory cortex, 32, 35–36  
 projections to inferior colliculus, 250  
 projections to nonauditory forebrain, 36  
 projections to primary auditory cortex, 32, 36  
 $Q_{10}$  dB, 215, 315, 323, 324  
 rabbit, 206

rat, 32  
 relation to limbic system, 220  
 response segregation, 218  
 role in coincidence detection, proposed, 382  
 role in contralateral sound localization, 443  
 rostral and medial parts of medial geniculate body, 220  
 serial processing and emergence of new representations, 218, 221  
 sound localization, 32  
 structure, 32–34  
 suprageniculate nucleus, 204, 216  
 sustained response properties, 213  
 synchronization, 213  
 synapses, 34  
 synaptic vesicles, 207  
 tectothalamic transformation, 204, 218  
 temporal discharge properties  
   independence from changes in discharge mode, 213  
   processing, 221  
 temporal sensitivity, compared in inferior colliculus and medial geniculate body, 216  
 thresholds, 208–210  
 tonic discharge, 208, 210, 213  
 tonic firing mode, 209  
 tonotopic organization, 219  
 topography of projections to medial geniculate body, 203  
 transmitters, 32  
 tuning, 215  
 ventral division, 28, 201, 203, 210  
**Medial nucleus of trapezoid body, 145, 380**  
 cat, 16  
   glycine, 22  
   guinea pig, 30  
   human, 17  
   location, 16  
   neuron types, 145  
   projections from, 22  
   projections from cochlear nucleus, 12  
   projections to cochlear nucleus, 18, 145  
   projections to inferior colliculus, 22  
   transmitters, 16

- Medial olivocochlear efferent fibers  
 bilateral projections, 17  
 distribution, 17  
 distribution along organ of Corti, 185  
 innervation of cochlear nucleus, 185  
 innervation of outer hair cells, 185  
 innervation of spiral afferent neurons,  
 185  
 lateral olivocochlear system, 16  
 number of neurons, 17  
 origin, 16–17  
 pathway, 185  
 projections, 16–17
- Medial olivocochlear system  
 definition, 16–17
- Medial superior olivary nucleus, 134,  
 249, 396, 504  
 axons to, and delay lines, 387, 428  
 banded input to inferior colliculus, 134  
 bilateral input, 134  
 cat, 15–19  
 coincidence detection in, 428  
 cytoarchitectonic features, 91–93  
 GABA, 134  
 glycine, 133–134  
 inhibitory input and creation of delay  
 lines, 388, 428  
 location, 2  
 mechanisms for sharpening interaural  
 time difference functions, 389–390  
 peak-type cells, 387  
 phase analysis, 428  
 physiology, 134  
 projections from, 134  
 projections from anterior ventral coch-  
 lear nucleus, 16–17  
 projections to cochlear nucleus, 18  
 projections to inferior colliculus, 22,  
 91–93, 134, 377, 388  
 role in coincidence detection, 382, 428  
 sound localization, 382  
 tonotopic organization, 16  
 transmitters, 133–134
- Membrane properties 282–300, 332  
 active, 282  
 input resistance, 288–289  
 passive, 282, 288  
 time constants, 288–289
- Mesencephalic lateral nucleus, pars dor-  
 salis, 498
- Mesencephalon, in birds, 498
- Metabotropic, 256
- Midbrain  
 auditory, 626–635  
 avian, 493–495  
 comparative perspective, 628–629  
 comparative projections to, 117–121  
 historical perspective, 626–628  
 mammalian, 629–632  
 projections from periolivary cell  
 groups, 117–121  
*See also* Inferior and Superior  
 colliculi, Mesencephalon
- Middle latency response, 325
- Minimum audible angle (MAA), 434–  
 435, 440
- Models of sound intensity coding, 334
- Modulated sounds, 263
- Modulation  
 frequencies, 263  
 frequencies, best, 564
- Modulation transfer function (MTF)  
 best modulation frequency, 350–353  
 inhibitory contribution to band-pass  
 amplitude-modulated transfer  
 functions, 564  
 firing rate (MTF<sub>FR</sub>), 258–259, 263, 351–  
 356  
 shapes in aged rats, 563  
 vector strength (MTF<sub>vs</sub>), 258–259, 350–  
 356
- Modulatory input, 510
- Monakow, stria of von, 4, 9
- Monaural  
 pathways, 132, 143–144, 146, 217, 502  
 specialized, and absence in birds, 506  
 response, medial geniculate body, 217  
 spectral representation, contribution to  
 azimuth perception, 440
- Mormyrids, 460–462
- Motion detection, 378
- Motor, 7, 43, 84, 232, 427, 603  
 pathways, 188  
 system, auditory relations with, 496
- Mouse (*Mus musculus*)  
 CBA, 560

- C57Bl/6, 559, 586  
 cochlear nucleus, 288  
 DBA/2J, 586  
 directional acuity in, 443  
 hearing loss, 586–598  
*See also* Rodent
- Multineuron responses after damage, 571
- Multipolar cells  
   cochlear nucleus, 10  
   in inferior colliculus, 500, 543  
   physiology, 10  
   ventral cochlear nucleus, 10
- Multisensory interactions, 444
- Muscimol, 251, 255, 270
- Music, 348, 400, 435, 445, 560
- Natural sounds, 325, 333, 361, 436
- NBQX (1,2,3,4-tetrahydro-6-nitro-2,3-dioxobenzol[*f*]quinoxaline-7-sulfonamide), 254, 258–259
- Neocortex, 37–44
- Neostriatum, bird 505
- Neural processing, inferior colliculus. *See* Inferior colliculus
- Neurochemistry  
   changes in aging, 567  
   comparative, 481, 502  
   inferior colliculus, 71, 260–268, 294–302, 366–367  
   primary auditory cortex, 39, 41  
   thalamus, 33–34
- Neuroethology, 411, 493
- Neuron types  
   cochlear nucleus, 10–12  
   disc-shaped, 75–76, 87, 98, 289, 497, 538, 540  
   dorsal cochlear nucleus, 10–12  
   ion channels, 101–102  
   medial geniculate body, 32, 34  
   membrane properties, 100  
   primary auditory cortex, 37–38  
   response types, 101  
   stellate, 12, 75, 100, 187, 249, 500
- Neuronal organization  
   cochlear nucleus, 12–16  
   cortex, 37–38  
   inferior colliculus, 69
- Neurotransmitters. *See* Transmitters
- Nissl preparation, 496, 630
- NMDA (*N*-methyl-D-aspartate), 208, 248, 255–256, 258, 263, 294–295, 523, 612–613
- NMDA receptor antagonists, 254
- APV ( $\pm$ )-2-amino-phosphonovaleric acid, 254, 294–295, 612
- CPP ( $\pm$ )-3-(2-carboxypiperazin-4-yl)-propyl-1-phosphonic acid, 254
- Noise, background, 560  
   broadband, 434, 436  
   discrimination of tones in, 324  
   efferent neurons, 184–185  
   high frequency and spectral cues, 435
- Noise bursts, 434
- Noise trauma, 268, 572, 591
- Noradrenaline, inferior colliculus, 567
- Norepinephrine, and inferior colliculus, 258
- Notch selective cells and localization, 438–439
- Nuclei of lateral lemniscus, 19–21, 86, 169, 295, 497, 609
- Nucleotopic organization, 124, 146
- Nucleus angularis, 506
- Nucleus basalis, 41
- Nucleus of central acoustic tract (NCAT), 216  
   projection to the supragenulate nucleus, 216
- Nucleus laminaris, 519
- Nucleus magnocellularis, 506
- O**ctaval nuclei, 464
- Octopus cells  
   cochlear nucleus, 10  
   connections, 140  
   physiology, 10
- Olivocochlear bundle  
   bilateral projections, 185  
   cat, 16–17  
   comparative number, 16–17  
   innervations of ear, 185  
   location in superior olive, 16–17  
   origin in CNS, 16–17, 184–185  
   pathway, 184–185  
   projections to cochlear nucleus, 185

- Olivocochlear bundle (*continued*)  
 to cochlear nucleus, 16–17  
*See also* Olivocochlear system
- Olivocochlear system  
 cat, 17  
 guinea pig, 17  
 mouse, 17  
 neuron immunocytochemistry, 185  
 neuron number, 17  
 parts, 16–17  
 superior olive, 184–186  
 transmitters, 185  
*See also* Olivocochlear bundle
- ON-OFF cells, unresponsiveness to sinusoidal amplitude modulation, 516
- Offset response, 611
- Onset response, 334, 378, 609
- Onset-offset responses, 609
- Ontogeny, 301  
 activity-dependence, 546  
 auditory evoked potentials, 2, 593  
 axons, 538–541  
 banding, 539–540, 548, 550  
 calbindin, 544–545, 550  
 calcium-binding proteins, 544–545, 550  
 calretinin, 544–545, 550  
 cholinergic receptors, 546  
 cochlear ablation, 547–549  
 dendrites, 539, 542–543  
 GABA, 544–546, 550  
 GAP43 influence on, 570  
 glutamate, 544–546  
 glycine, 544–546  
 inferior colliculus, 537–552  
 layers, 538–540  
 myelin, 542  
 neuronal birth, 539  
 onset of hearing, 539–544, 547  
 sensorineural hearing loss, 547–549  
 synapses, 541–543  
 synaptic domains, 551  
 synaptogenesis, 539–547, 570  
 time course, 539  
 tonotopy, 539  
 transmitters, 544–546
- Optic tectum, 506  
 projection of external nucleus of owl inferior colliculus to, 522  
 role in direction of head movements, 523
- Ossicles, removal of, 268
- Ossicles, removal of, and effects on evoked GABA release, 569
- Ototoxic drugs, 571
- Owl, 258, 263, 494, 515, 518–525
- Paradoxical latency shift (PLS), 511, 514
- Paragigantocellular nucleus, 193
- Paralemniscal tegmentum, 464
- Paralemniscal zone, 189
- Parallel pathways  
 auditory, 15  
 thalamus, 27–28  
 visual system, 29
- Parallel processing, 27–28, 41, 44
- Parvalbumin, 497, 570
- Passive listening, 494
- Patch clamp, 253, 267, 288, 294, 365
- Perceptual changes after lesions, 435
- Perceptual state and neural activity, relation of, 427
- Periaqueductal gray, 192, 464
- Pericentral area, 84, 323, 496, 629  
 relation to external nucleus, 497
- Pericentral nucleus. *See* Dorsal cortex
- Peripheral changes  
 desensitization, 573  
 effects on neurotransmitter status, 570
- Peripheral sensitivity, 560, 562
- Periodicity, 147, 352, 367, 463, 541  
 coding, 348, 360  
 pitch, 348
- Periodotopy, 356–358
- Periolivary nuclei (PON), 9, 135, 138  
 acetylcholine, 16  
 binaural processing, 137  
 cat, 16  
 glycine, 135  
 guinea pig, 133–138  
 neuron types, 135  
 projections to midbrain, 9, 15–19, 136  
 projections from cochlear nucleus, 135, 137

- projections from inferior colliculus,
  - 184, 188
  - size, 16, 135
  - superior periolivary nucleus, 137
  - transmitters, 18
- Phase ambiguity
  - filtering of side peaks, 519
  - responses, 522
- Phase locking, 313, 349, 392, 634
- Phase versus frequency plots, 380
- phospho-CREB, cyclic adenosine mono-  
phosphate response element bind-  
ing protein, 571
- Phylogenetic divergence of birds and  
bats, 525
- Physiology
  - primary auditory cortex, 38–40
  - superior olive, 15–18
- Picrotoxin, 251, 269
- Pinna
  - filtering. *See* Head-related transfer  
function
  - movements, 188
  - premotor centers, 189
  - role in sound localization, 378
- Place code, 330
- Plane, median, 434
- Planum temporale, 43
- Plasticity
  - adaptive, 573
  - adult, 568
  - aging and, 559, 567
  - compared with other forms of change,  
571
  - compared with unmasking, 568
  - developmental, 548
  - experience dependent, 494, 523
  - neural, 567
  - response, 249
- Plateau, rate level function, 332
- Pontine nuclei, 191
- Pontine tegmentum, 42
- Population coding, in spatial localization,  
390
- Postsynaptic potentials (PSPs), 255–256
  - multiplication of, 522
  - response, 254
  - spikes and postsynaptic potentials, and  
interaural time and level process-  
ing, 522
- Posterior ventral cochlear nucleus
  - anatomy, 12
  - and lateral lemniscus, 20
  - multipolar cells, 10
  - neuron types, 10
  - projections to the bat inferior collicu-  
lus, 502
  - spontaneous rate, 10
- Posterior intralaminar nucleus, 201
- Potassium channel subunits, Kv, 102,  
290, 291, 497–498, 697–698
- Potassium currents, 283–284, 290–293
  - A-type, 286, 292
  - apamin-sensitive, 285, 287
  - buildup-pauser, 286–289
  - Ca<sup>2+</sup>-activated, 285, 287
  - CTX-sensitive, 285, 287
  - delayed-rectifier, 285–286, 291
  - high-threshold, 287, 291
  - inward rectifier, 293
  - low-threshold, 286, 290
  - onset, 286–288
  - rebound, 285, 287
  - sustained-regular, 283–284, 287–288
  - TEA-sensitive, 283, 285
- Precedence effect, 397–400, 444–447
  - anteroventral cochlear nucleus, 400
  - auditory cortex, 400
  - auditory nerve, 400
  - cochlear filter interaction and, 399
  - conditioner and probe, 397
  - eye position, and, 446
  - hair cell-based compression and adap-  
tation, 399
  - inferior colliculus, 400
  - inhibition and, 399
  - interstimulus delay, 444
  - neural basis for, 399
  - perceptual uncertainty, and, 447
  - populations in central nucleus of the  
inferior colliculus during behavior,  
447
  - responses at different levels of the au-  
ditory system, 447
  - role of anesthesia in, 447
  - saccades, and, 446

- Precedence effect (*continued*)  
 saccades, double, 446  
 spaciousness, role in the creation of, 400  
 superior olivary complex, 400  
 timbre, role in, 400
- Preisthmic superficial nucleus, 498
- Premotor processing, 494
- Pre-pulse inhibition, 596–597
- Presbycusis, 559–560  
*See also* Hearing loss
- Primary auditory cortex  
 afferent input, 38  
 best frequency, 39–43  
 binaural processing, 38, 40  
 bird, 505  
 cat, 38–39  
 commissural pathways, 38  
 connectivity, 39, 42  
 corticocollicular cells of origin, 39  
 corticocortical connections, 38–39  
 corticofugal projections, 38, 41–42  
 corticothalamic cell origin, 38  
 functional organization, 38–43  
 GABA, 39  
 glutamate, 39  
 laminar organization, 39  
 layers, 37–44  
 learning, 41  
 memory, 41  
 neurochemistry, 39  
 neuron types, 39–41  
 neuronal organization, 37–41  
 phase detection, 40  
 physiology, 38–39  
 projections from medial geniculate body, 38  
 projections from thalamus, 38, 40  
 projections to thalamus, 38, 42  
 $Q_{10}$  dB, 40–41  
 sound localization, 40  
 tonotopic organization, 38  
 tonotopy, 39  
 transmitters, 39–40  
 tuning, 38  
*See also* Auditory cortex
- Probst, commissure of, 4
- Processing mechanisms, nonlinear mechanisms, 325
- Processing streams, 634–635
- Psychometric functions, 441
- Psychophysics, 494
- Puncta, 497  
*See also* Synaptic endings
- Pyramidal cells  
 cerebral cortex, 232, 239  
 cochlear nucleus, 10. *See also* Fusiform cell  
 origin of corticocollicular system, 232
- Q**, quality factor, 315, 322
- $Q_{10}$  dB  
 inferior colliculus, 262, 322–323  
 medial geniculate body, 215  
 primary auditory cortex, 38, 40  
 sharpness of, 215, 322–323
- Rabbit**  
 auditory cortex, 400  
 inferior colliculus, 392, 400
- Radiations, auditory, 2
- Rat  
 corticofugal system, 42  
 Fischer 344 strain, 566  
 inferior colliculus, 76–77, 155, 630–631  
*See also* Rodent
- Rate coding, 334, 350–351
- Rate, discharge, 575
- Rate-intensity curves. *See* Rate-level functions
- Rate-level functions, 33, 302, 313, 332, 353, 511  
 best level 330  
 bicuculline and, 610  
 complex, 332  
 confounding onset and sustained responses, 332  
 corticofugal influences, 332  
 different assignment criteria, 332  
 effects of aging on, 269  
 EI cells and, 394  
 factors influencing, 518  
 hearing loss, 588  
 limited sound range, 332  
 mediated by GABA, 610  
 monotonic, 260, 331–332, 393, 588  
 nonmonotonic, 260, 331–332, 510, 588, 595, 610

- plateau, 332
- thresholds 329–330
- Rebound, excitatory, 512
- Receptive field, 260, 313, 517
  - bandwidth, 322
  - excitatory, 313–314
  - influences upon, 518
  - interactions among, 517
  - subregions of, 314
- Receptor binding, 252
- Receptors
  - acetylcholine, 301–302
  - AMPA, 294–297
  - GABA<sub>A</sub>, 214, 297–300
  - GABA<sub>A</sub>, effects of aging, 269
  - GABA<sub>B</sub>, 297–300
  - glutamate, 294–297
  - glycine, 254, 300
  - NMDA, 174, 214, 294–295
  - serotonin, 301
  - See also* specific receptors
- Recovery time,
  - anesthesia, effects of on, 399
  - auditory cortex, 399
  - auditory nerve, 399
  - cochlear nucleus, 399
  - functions, 562
  - inferior colliculus, 399
  - superior olivary complex, 399
- Regimes, tonic, 332
- Repetition rate, 477, 495
  - stimulus, 518
- Reptile
  - hearing, 477–481
  - lateral lemniscus, 478–479
  - superior olive, 478
  - thalamus, 478–479
- Response
  - area, 509
  - closed, 510
  - multipeaked, 510
  - repertoire, 568
  - synchrony, 316
- Response plasticity, 518
  - effects of electrical stimulation, 518
  - effects of inactivation, 518
  - enhancement of, 518
- Responses
  - multineuron, after damage, 571
  - nonmonotonic, best level, 330
- Reverse correlation, 325
- Rewiring, 559
- Ripple density, 327
- Rodent
  - medial geniculate body, 239
  - olivocochlear system, 17
  - superior olive, 138
  - See also* specific species
- Rostral pole, 83, 188
- Sagulum, 23, 184, 204–205
- Saliency, behavioral, 518
- Secondary auditory cortex, 39, 42
- Seizures, motor, 603
  - See also* Audiogenic seizures
- Semantic information, 334
- Sensory-to-motor transformation, 494
- Serial processing, 41, 44
- Serotonin, inferior colliculus, 258, 268, 301
- Shell, lateral and medial in bird, projections to, from laminaris and magnocellular nuclei, 506
- Sidebands, inhibitory, 522
- Side peaks, filtering of space ambiguous, 519
- Signal detection criterion,  $d'$ , 434–435
  - in echo clutter, 517
- Signal, instructive, 524
- Signal-to-noise ratio, 11, 564
- Single neuron discharge patterns, importance of, 426
- Sinusoidal amplitude modulation (SAM), 516
  - depth, 516
  - See also* (AM) Amplitude modulation
- Sinusoidal frequency modulation (SFM), 516
  - coincidence and sinusoidal frequency modulation tuning, 517
  - selectivity of some inferior colliculus neurons to, 517
  - See also* Frequency modulation (FM)
- Slope(s), 315, 318
- Somatosensory system
  - projections to inferior colliculus, 30–31
  - projections to superior colliculus, 188–190

- Sound  
 biologically meaningful, 448, 494  
 broadband, 438
- Sound detection, superior olive 380–382
- Sound localization  
 adaptive adjustment in barn owls, 523  
 at low frequencies, 377  
 bat, 493  
 contralateral encoding of, 427, 430  
 deficits, and central nucleus of the inferior colliculus and the lateral superior olive, 443  
 errors in elevation, 440  
 head related transfer function, 378  
 importance of cues in, 427  
 lateral superior olive, 381  
 mechanisms of, 518  
 medial geniculate body, 32  
 medial superior olivary nucleus, 382  
 natural, and functional perspective on localization, 443  
 near field and interaural level differences, 378  
 nonpredatory lifestyle, and, 442  
 owl, 494  
 predatory lifestyle, and, 442  
 rat, 32  
 role of pinna, 378  
 specialists in, 494  
 vertical, 187
- Sound pressure level, 313
- Sound shadow, 432
- Sound source  
 location, detectable changes in, 522  
 movement, 378, 517  
 phantom, 445  
 segregation, 378  
 spectrum, disambiguation of, 436
- Sounds  
 artificial, 325  
 natural, 333
- Source-filter theory, 412–413
- Space map  
 absence in the bat midbrain, 517  
 experience-dependent plasticity, 523  
 midbrain in the owl, 517, 519, 524  
 multimodal, in owl inferior colliculus, 524  
 space-specific neuron responses, 519, 521–523
- Space-specific receptive fields, contributions of a spike threshold mechanism to, 522
- Spaciousness, acoustic and the environment, 387
- Spatial  
 acuity in the horizontal plane, 432  
 information, ambiguity of, 517, 519  
 processing, 564
- Species  
 bat, big brown (*Eptesicus fuscus*), 266, 315, 319, 323, 507, 509  
 bat, horseshoe (*Rhinolophus hipposideros*), 116  
 bat, Mexican free-tailed (*Tadarida brasiliensis mexicana*), 268, 415  
 bat, mustache or mustached (*Pteronotus p. parnellii*), 116, 124, 201–202, 205, 219, 260, 266–267, 395, 507, 517, 573, 630  
 bat, pallid (*Antrozous pallidus*), 266, 507  
 bullfrog (*Rana catesbeiana*), 468, 471  
 cat (*Felis catus*), 116, 119, 202, 219, 312, 315–316, 318–319, 322–324, 327, 332, 384–385, 387, 395, 400, 433, 436, 446, 630  
 chicken (*Gallus gallus*), 387, 508, 515, 521  
 chimpanzee (*Pan troglodytes*), 117  
 chinchilla (*Chinchilla lanigera*), 116, 120, 136, 159, 260, 266, 312  
 crocodile (*Caiman crocodilus*), 479  
 dog (*Felis canus*), 446  
 ferret (*Mustela putorius*), 116, 138, 205  
 gerbil (*Meriones unguiculatus*), 116, 119, 124, 137, 159, 255, 312, 322, 387, 395  
 guinea fowl (*Numida meleagris*), 508  
 guinea pig (*Cavia porcellus*), 116, 119, 202, 260, 312, 318–319, 322, 332, 383–385, 392, 442, 572  
 human (*Homo sapiens*), 356, 386–387, 630  
 marsupial (*Didelphis marsupialis*), 116, 630–631  
 midshipman fish (*Porichthys notatus*), 460–463  
 mole (*Mogera tokudae*), 116, 119



- monkey rhesus (*Macaca mulatta*), 312, 630
- mouse (*Mus musculus*), 116, 137, 159, 169–170, 202, 266, 315, 318, 322, 324, 332, 442, 559, 604
- opossum (*Didelphis marsupialis*), 137, 630–631
- owl, barn (*Tyto alba*), 387, 447, 508, 517–521, 630
- Pacific treefrog (*Hyla regilla*), 468, 471, 474
- rabbit (*Lepus americanus*), 382, 384, 386–387, 392, 400, 447
- rat (*Rattus norvegicus*), 116, 119, 155, 322, 332, 446, 603–604, 630
- rat kangaroo (*Dipodomys merriami*), 387
- squirrel monkey (*Saimiri sciureus*), 218, 238
- zebra finch (*Taeniopygia guttata*), 508, 515
- Species differences, 332
- Species-specific features, 15, 76, 301, 324, 421, 493
- Spectra
  - complex, 325
  - ripple, 327
  - tone, 327
- Spectral
  - cues, complex, 435, 438
  - decomposition, 312
  - envelopes, sinusoidal, 325
  - filters, 312
- Spectral cues, 378, 432, 435
- Spectral information
  - auditory cortex, 39
  - cues, 428
  - inferior colliculus, 90–91
  - medial geniculate body, 215, 217–220
- Spectral integration, 325, 327–328, 440
- Spectral minima and maxima, 313
- Spectral models and computational models, 433
- Spectral notch, 433–434, 440
- Spectral receptive field (SRF), 313–330
  - classes I–IV, 314–319, 323–324, 328–330
  - facilitation, 315
  - inferior colliculus subdivisions, 316
  - types I, O, V, 314–319, 324, 328, 330
  - W-shaped, 472
- Spectral representation, monaural, contribution to azimuth perception, 440
- Spectral tuning, 313–330
- Spectrotemporal
  - envelopes, 325
  - feature combinations, 327
  - receptive fields (STRFs), 325
  - receptive fields, composite, 329
- Speech, 347, 412–414, 560
- Speech encoding, 346, 418–420
- Spherical bushy cells, 142, 428
  - ventral cochlear nucleus, 10, 12
  - See also* Bushy cells
- Spherical cells. *See* Spherical bushy cells
- Spontaneous activity
  - inferior colliculus, 314
- Startle response, 190, 191
  - cochlear root neurons, 190
  - motor pathway, 190
- Stellate cell
  - dorsal cochlear nucleus, 12, 120, 187
  - in avian inferior colliculus, 498
  - inferior colliculus, 74–78, 289, 543
  - projection to the cochlear nucleus, 187
  - synapses on, 121
- Steroid binding sites, midbrain in avians, 498
- Stimulation, electrical and c-Fos immunoreactivity, 573
- Stimuli
  - broadband, 325
  - complex, 380
  - gap, 560
- Stimulus repetition rate, 518
- Streaming, 346
- Stress and inflammatory responses, 567
- Structure-function relations, 334
- Strychnine, 267, 299–300, 415
- Substantia nigra, 192, 238, 503, 627
  - projection to inferior colliculus, 504
- Summation of excitatory postsynaptic potentials (EPSPs), 382
- Summing localization, 445
- Summing window, 400
- Superior colliculus
  - auditory input, 204
  - carnivores, 630
  - cat, 630
  - connections, 188

- Superior colliculus (*continued*)  
 deep layers, 189  
 functional organization, 189  
 location, 70  
 projections from inferior colliculus, 188  
 role in startle response, 190
- Superior olive nucleus, medial superior olivary nucleus  
 anatomy, 133  
 bat, 503  
 bird, 505  
 cat, 22  
 comparative studies, 503–505  
 complex signal processing, 382  
 component nuclei, 15–19, 133  
 effects of lesion on interaural level differences, 396  
 location of olivocochlear cells, 185  
 major functions, 380–385  
 medial olivocochlear neurons, 16, 185  
 nuclei, 16, 133  
 olivocochlear system, 18, 133  
 physiology, 15, 382  
 projections from anterior ventral cochlear nucleus, 117–122  
 projections from cochlear nucleus, 117–122  
 projections to inferior colliculus, 22, 117, 497  
 rabbit, 382  
 reptile, 478  
 size, 16  
 tonotopic organization, 16  
 transmitters, 15–16  
*See also* Lateral superior olivary
- Superior paraolivary nucleus, 137–138, 254, 610
- Superior temporal gyrus, 43, 56
- Sustained response, 140, 334
- Synapse, 99, 102, 214, 266, 524, 543, 626  
 axodendritic endings, 207  
 axosomatic endings, 207  
 cochlear nucleus terminals in inferior colliculus, 121  
 GABAergic axon terminals, 207  
 inferior colliculus, 78–80  
 medial geniculate body, 206–207  
 triads, 207  
 vesicles from anteroventral cochlear nucleus, 121  
 vesicles from dorsal cochlear nucleus, 121  
 vesicles from lateral superior olive, 121  
 vesicles from medial superior olive, 121
- Synaptic  
 changes with age, 566–567  
 connections, new, 560  
 domain 86–88, 91, 124, 148, 334, 551  
 endings, 497  
 function, 268  
 organization, 103  
 response, early, 254
- Synaptic vesicle  
 inferior colliculus, 121  
 medial geniculate body, 207  
 round, 121
- Synaptogenesis, 539–543
- Synaptosomes, 269–270
- Synchronization code, 349–350, 465
- Synchrony, 562
- TBOB (*t*-[<sup>3</sup>H]butylbicycloorthobenzoate), 252, 270
- Tectothalamic transmitters, 25, 33, 205–207
- Tectum, 70, 506
- Tectum, optic, 506
- Tegmentum, 70, 83
- Telencephalon  
 auditory subdivisions, 32–44  
 projections to inferior colliculus, 230–242
- Teleost fish, 460–467
- Temporal coding  
 amphibians, 472–477  
 amplitude modulation, 349–361  
 birds, 515–516  
 degradation in the ascending auditory system, 214  
 duration, 364–366  
 inferior colliculus, 104, 346–351, 366–368, 377  
 fish, 465–467  
 frequency modulation, 361–364  
 gaps, 364–365  
 pitch 348,

- torus semicircularis, 465–467, 472–477
- Temporal lobe, 41, 43
- Temporal locking, 258, 392
- Temporal processing, 140, 560
- Temporal processing, effects of aging, 566
- Temporal response
  - buildup, 349
  - chopper, 349
  - medial geniculate body and inferior colliculus, 348–349
  - onset, 349, 508
  - patterns, 508
  - pauser, 349
  - primary-like, 508
  - primary-like with notch, 508
  - sustained, 508
- Temporal selectivity, 266
- Thalamic reticular nucleus, projections to medial geniculate body, 207
- Thalamocortical auditory system, 32–44
- Thalamocortical connections, cat, 32–42
- Thalamotectal system, 231, 238–239, 242, 504
- Thalamus
  - amphibian, 479
  - bat, 201–212
  - bird, 504–505
  - connections, 36, 201–212
  - fish, 463
  - GABA, 32–33
  - neurochemistry, 33
  - ovoidalis, nucleus, 506
  - physiology, 32–36
  - posterior, 238
  - projections from inferior colliculus, 200–208
  - projections from primary auditory cortex, 242
  - projections to primary auditory cortex, 36
  - reptile, 478
  - tonotopic organization, 33, 201–202
  - transmitters, 32–33
  - See also* Medial geniculate body
- Threshold, 73, 224, 260, 291, 356, 562
- Threshold, behavioral decision, 330
- Threshold, minimum, 313, 330
- Thresholds
  - hearing, 254
  - inferior colliculus, 329–330
  - medial geniculate body, 209–212
  - upper, 509
- Timbre, 348
- Timbre, and localization, 446
- Tone discrimination, 435
- Tonic regimes, 332
- Tonotopic organization
  - auditory cortex, 39
  - auditory pathway, 6, 16, 23, 25, 33, 39
  - cochlea, 312
  - cochlear nucleus, 6
  - hearing loss, 587
  - inferior colliculus, 25, 88–89, 319–321, 507
  - lateral superior olivary nucleus, 16
  - medial geniculate body, 33, 36
  - medial superior olivary nucleus, 16
  - primary auditory cortex, 39
  - specialized, 508
  - species-specific, 507
  - superior olive, 16
  - tectothalamic projections, 157
  - thalamus, 33
- Topographic map, 57, 319
  - frequency, 319–322, 352, 587
  - periodicity, 94, 356–360
  - prevalence of in the brain and across modalities, 427
  - projections, 161
  - sharpness of tuning, 323
  - space, 505, 521–524
  - threshold, 321
- Toral nucleus, 498
- Torus semicircularis
  - amphibians, 467–477
  - avian, 498
  - central nucleus, 477–480
  - comparative neuroscience, 21, 459–483
  - fish, 461–467
  - immunocytochemistry, 480
  - interval-counting neuron, 474
  - laminar nucleus, 477–480
  - projections, 463, 471, 478,
  - recovery-type neuron, 474–476
  - reptiles, 477–481

- Torus semicircularis (*continued*)  
 spectral processing, 470–472  
 subdivisions in birds, 498  
 temporal processing, 465–467, 482
- Tracers, neuroanatomical  
 axonal degeneration, 120  
 biotinylated dextran amines (BDA)  
 163, 173  
 double labeling for connectivity and  
 neurotransmitter, 172  
 horseradish peroxidase (HRP), 75, 119,  
 164  
 horseradish peroxidase conjugated to  
 wheat germ agglutinin (HRP-  
 WGA), 116  
*Phaseolus vulgaris*-leucoagglutinin  
 (PHA-L), 159, 163
- Trapezoid body, 9
- Transmitters and neuromodulators  
 acetylcholine (or ChAT), 16, 25, 33,  
 258, 498. *See also* choline acetyl-  
 transferase  
 age-related changes in, 568  
 aspartate, 20, 22, 25, 33, 174  
 bat, 205–207, 516  
 bird, 501  
 calbindin-D28, 497  
 calretinin, 497  
 cat, 6, 16, 22, 25, 33  
 cochlear nucleus, 20  
 enkephalins, 33, 136  
 excitatory, 174  
 excitatory and inhibitory interactions,  
 444  
 GABA, 6, 16, 25, 27, 33, 39, 45, 250–  
 251, 497  
 GAD. *See* GABA  
 glutamate, 20, 22, 25, 27, 33, 39  
 glycine, 6, 16, 22, 25, 251, 497  
 inferior colliculus, 20, 25  
 inhibitory, 174  
 lateral superior olive, 16  
 medial geniculate body, 25  
 medial nucleus of trapezoid body, 16  
 medial superior olive, 16  
 norepinephrine, 258  
 olivocochlear neurons, 185, 187  
 parvalbumin, 497  
 periolivary nuclei, 134, 136  
 primary auditory cortex, 39  
 serotonin, 39, 187, 258, 268  
 somatostatin, 136  
 tectothalamic, 25, 33, 205–207  
 thalamus, 33  
 ventral nucleus of lateral lemniscus,  
 23  
*See also* specific transmitters
- Transneuronal, 240, 242, 465, 585
- Transsynaptic changes, 584, 590
- Trapezoid body  
 cat, 4  
 projections from anterior ventral coch-  
 lear nucleus, 121
- Trauma  
 acoustic, and CAMP, 571  
 acute noise, 572  
 comparing neurophysiological and neu-  
 rochemical effects, 575  
 decreased inhibition on binaural input,  
 and, 572–573  
 discontinuities in characteristic fre-  
 quency following, 577  
 firing rate increase after cochlear dam-  
 age, 576  
 recruitment of neurons and, 572  
 residue of pre-lesion input after, 578
- Trochlear nerve, 499
- Tuning  
 broadband in owl, 520  
 cat auditory nerve, 317  
 class I-IV, 314–330  
 compared at the superior olive, inferior  
 colliculus, and medial geniculate  
 body, 388  
 curve tails, effects of damage upon,  
 573  
 curve tips of high-frequency units and  
 aging, 560  
 curves of response areas, effects of  
 damage upon, 572  
 curves  
 effects of inhibition on, 572  
 frequency-response, 251, 322  
 frequency-threshold, 561  
 level tolerant, 381  
 tail region, 384  
 lateral shell of inferior colliculus,  
 508

- multipeaked in birds, 508
  - V-shaped in birds, 508
  - V-shaped in anurans, 473
  - W-shaped in anurans, 473
- medial geniculate body, 211
- medial geniculate body neurons, 205–208
- owl central nucleus of inferior colliculus core (ICCc), 520
- primary auditory cortex, 39
- pure-tone, 322
- types I, O, V, 260, 319–330
- Type I spiral ganglion neurons. *See* Spiral ganglion
- Type II spiral ganglion neurons. *See* Spiral ganglion
- (*t*-[<sup>3</sup>H]butylbicycloorthobenzoate). *See* TBOB
- Two-tone inhibition. *See* Inhibition, two tone
  
- Unipolar cells, in avian inferior colliculus, 500
- Units, single and multiunits, 319
- Unmasking
  - hearing loss, 586
  - physiological, 575
  - plasticity and, 559, 568, 575
  - scatter of characteristic frequency and, 576
  - suppressed input, 576
  
- Ventral cochlear nucleus
  - amphibians, 471
  - anatomy, 12
  - effects of trauma, 268
  - multipolar cells, fine structure, 120, 121–122
  - neuron types, 12
  - physiology, 328
  - projections to inferior colliculus, 249
  - pseudoplasticity, 590
  - spherical bushy cells, 10, 12
  - See also* Cochlear nucleus
  
- Ventral nucleus of lateral lemniscus, 20, 139–141, 502
  - bat, 502
  - birds, 505
  - cat, 121
  - changes in connections with age, 566
  - columnar subdivision of, in bats, 140, 502
  - divisions, 140
  - function, 140
  - GABA, 23, 140
  - glycine, 23, 139
  - location, 23
  - monaural hearing, role in, 377
  - projections from, 277
  - projections to, 9, 140
  - tonotopic organization, 23
  - transmitters, 23
- Ventrolateral nucleus, 82
- Vestibular system, 36
- Virtual motion, 262
- Visual and auditory frames of reference, calibration of, 523
- Visual cortex, 38
- Vocalization
  - amphibians, 468–469
  - animal, 560
  - bat, 414–418, 493–494
  - circuit, 192
  - fish, 460–465
  - formants, 413
  - fundamental frequency, 413
  - natural, 325
  - processing, 411–422
  - selectivity, 415–417
  - specialization, 415, 422
  - vocal tract, 413
- Voice-onset time, 418–420
- Voltage-clamp, 253
  
- Waveform, 349, 355, 413, 466, 481
  
- Zone
  - lateral, 13, 629
  - paralemniscal, 189, 194

# Author Index

- Aas JE, 191, 195  
Abbas PJ, 329, 342  
Abbott SD, 268, 272, 569, 570, 575, 579  
Aberdeen GC, 604, 617  
Aboitiz F, 43, 50  
Abraham LD, 52, 223  
Abramson AS, 418, 423  
Adams JC, 5, 7, 11, 17, 19, 27, 50, 116, 119, 120, 121, 122, 126, 135, 140, 141, 150, 164, 169, 178, 238, 244, 248, 249, 250, 251, 254, 262, 272, 361, 368, 406, 457, 627, 633, 636  
Adams-Curtis LE, 621  
Adamson L, 53  
Adelman G, 53  
Adelman JP, 310  
Adolphs R, 501, 505, 519, 526  
Adrià HO, 38, 57  
Aertsen AMHJ, 337, 343  
Airaksinen M, 623  
Aitkin LM, vii, 3, 4, 12, 19, 21, 23, 26, 27, 29, 30, 31, 33, 35, 36, 38, 50, 51, 52, 64, 73, 85, 90, 91, 97, 106, 112, 116, 122, 123, 124, 126, 127, 128, 129, 130, 147, 150, 153, 157, 172, 178, 193, 195, 202, 203, 204, 207, 208, 213, 223, 231, 233, 238, 244, 312, 316, 319, 322, 323, 328, 330, 332, 336, 342, 343, 412, 418, 421, 423, 452, 509, 518, 519, 521, 527, 542, 543, 552, 553, 554, 602, 626, 627, 632, 636  
Akagi H, 310  
Akana SF, 624  
Akbar MT, 615  
Akesson TR, 505, 526  
Akimitsu T, 620  
Akiyama H, 114  
Akoev GN, 490  
Albert M, 44, 51, 372, 473, 484, 488  
Albin RL, 277, 581  
Alder TB, 473, 474, 476, 484, 486  
Aleszezyk CM, 454  
Alibardi L, 116, 120, 121, 127, 186, 195  
Allen DM, 469, 484  
Allon N, 215, 223  
Altman JA, 338, 538, 553, 629, 636  
Altschuler RA, 7, 51, 61, 62, 66, 67, 68, 110, 113, 151, 198, 272, 274, 275, 279, 280, 305, 309, 338, 342, 557, 579, 582, 599, 616, 618  
Alvarado JC, 554  
Alvarado-Mallart RM, 70, 112  
Amarillo Y, 304, 309  
Amatsu M, 306, 581, 619, 620  
Ambler JS, 582  
Amoroso J, 63  
Andersen RA, 23, 30, 51, 64, 112, 116, 129, 153, 157, 164, 176, 178, 201, 203, 204, 223, 342, 426, 450  
Anderson CA, 305  
Anderson CAB, 274, 305, 618, 619  
Anderson DJ, 50, 106, 225  
Anderson TR, 109, 404, 406  
Andressen C, 544, 553  
Andrews B, 245  
Andrejik JA, 193, 195  
Andrianov YN, 490  
Antanavage J, 307  
Aoki E, 112  
Appell PP, 188, 195  
Aragon C, 554

- Aramakis VB, 546, 553  
 Aran JM, 582  
 Araujo DM, 309  
 Archer JR, 599, 602  
 Arehole S, 279  
 Argeles PC, 371  
 Arita K, 620  
 Arjmand E, 553  
 Armour BAL, 196, 225, 273, 580  
 Arnerić SP, 273, 278, 305, 580, 618, 619  
 Arnott RH, 408  
 Arruda M, 457  
 Aschoff A, 132, 134, 135, 141, 142, 143, 151, 529, 637  
 Ashburner J, 225  
 Ashmead DH, 446, 450  
 Ashmore JF, 13, 51  
 Astl J, 229, 375, 456  
 Aston-Jones G, 199  
 Atkins AL, 605, 615  
 Atsoggiu T, 273  
 Attias H, 333, 336  
 Avan P, 249, 278, 328, 342  
 Axelsson GA, 582  
 Azerado WJ, 198  
 Azizi SA, 195
- Baba K, 180**  
 Bac P, 605, 615  
 Bachelard HS, 250, 251, 273  
 Backoff PM, 373, 582  
 Backus KH, ix, 298, 300, 301, 305, 307, 546, 555  
 Bading H, 611, 620  
 Bagri A, 605, 615  
 Baimbridge KE, 544, 553  
 Bain CJ, 280  
 Baird IL, 5, 45, 51, 489  
 Bajo VM, 19, 51, 59, 110, 157, 178, 180, 198, 224, 251, 272, 638  
 Baker BN, 55, 108, 250, 251, 270, 274, 452  
 Baker H, 310  
 Baker R, 80, 112, 136, 137, 138, 140, 153, 197, 231, 246, 465, 468, 484, 485, 487  
 Bal R, 287, 288, 289, 304, 332, 336  
 Bala AD, 534  
 Ball D, 611, 622
- Ball GF, 526  
 Balthazart J, 498, 526  
 Banay-Schwartz M, 250, 251, 269, 272  
 Bandyopadhyay S, 624  
 Banks MI, 130  
 Baranauskas G, 304  
 Barberis A, 622  
 Barker JL, 278  
 Barker MC, 305  
 Barnard EA, 293, 304  
 Barnes WT, 115, 127, 627, 636  
 Barnes-Davies M, 103, 106  
 Barney HL, 413, 424  
 Baron J, 107, 245  
 Barone P, 52, 224, 229, 456  
 Barruel P, 341  
 Barsz K, 364, 368  
 Bartlett E, 100, 106, 111, 205, 206, 207, 208, 210, 212, 213, 214, 223, 228  
 Baruch AM, 477, 479, 485  
 Basbaum AI, 127  
 Bass AH, 460, 461, 462, 464, 465, 466, 467, 468, 484, 485, 487, 489, 491, 492  
 Basta D, 286, 304  
 Basu L, 531  
 Batra R, 51, 91, 97, 104, 106, 107, 109, 111, 113, 135, 137, 140, 150, 152, 285, 307, 371, 387, 392, 402, 404, 406, 428, 450, 451, 453, 456  
 Baudrimont M, 543, 555  
 Bauer CA, 272, 569, 570, 579  
 Bauer EE, 109, 226, 407, 423, 488, 582  
 Bausch SB, 293, 304  
 Baxter D, 71, 111  
 Bayliss DA, 309  
 Bear D, 44, 51  
 Bear MF, 257, 272  
 Beaulieu C, 637  
 Beaver CJ, 339  
 Bechtold JK, 59  
 Beck MM, 603, 616  
 Beckius GE, ix, 17, 51, 61, 111, 116, 124, 129, 152, 227, 277, 341, 387, 388, 402, 407, 557, 638  
 Bedenbaugh PH, 52, 489  
 Behan M, 188, 195  
 Behrend O, 137, 150, 403  
 Beitel R, 425, 489  
 Belekhoa MG, 477, 479, 480, 485

- Belendiuk K, 432, 450  
 Belin P, 43, 51  
 Ben-Ari Y, 308, 622  
 Ben-Arie J, 433, 454  
 Beneyto M, 23, 24, 25, 38, 43, 51, 184,  
 195, 205, 223, 232, 244  
 Beninger RJ, 600  
 Benson CG, 113, 120, 124, 127, 150,  
 277, 279, 558, 583, 624  
 Benson DF, 43, 51  
 Benson PK, 368  
 Berbel P, 79, 97, 110  
 Berger AJ, 309  
 Bergman M, 248, 249, 272  
 Berman AL, 17, 18, 19, 51, 54, 58, 71,  
 82, 84, 107, 316, 329, 340  
 Bernstein LR, 391, 393, 397, 402, 403  
 Berrebi AS, 9, 11, 15, 22, 51, 66, 128,  
 136, 137, 152, 153, 178, 180, 276,  
 610, 616, 621  
 Berridge CW, 193, 195  
 Betz M, 534  
 Beyerl BD, 116, 127  
 Bianchi MT, 606, 616  
 Bibikov NG, 352, 353, 368, 452, 472,  
 485  
 Biebel UW, 352, 353, 359, 368  
 Bieser A, 372  
 Biggio G, 273  
 Bijak M, 277  
 Binns KE, 23, 29, 51, 316, 336  
 Bischofberger J, 57, 306  
 Bishop DC, 61, 109, 111, 116, 129, 557  
 Bjaalie JG, 59, 110, 180, 246, 272, 638  
 Bjordahl TS, 13, 51  
 Blackstad TW, 109, 340, 555  
 Blaine JT, 290, 304  
 Blakemore C, xiii, ix  
 Blasdel GG, 55, 531  
 Blatchley B, 369, 543, 553  
 Blauert J, 377, 378, 399, 400, 403, 430,  
 433, 438, 444, 445, 450  
 Blednov Y, 625  
 Bledsoe SC Jr, 15, 22, 51, 198, 268, 272,  
 569, 570, 573, 579, 599, 620  
 Bliss TVP, 257, 272  
 Blomqvist A, 114  
 Blum PS, 35, 52, 205, 223  
 Blümcke I, 553  
 Bock GR, 193, 199, 213, 223  
 Bodenhamer RD, 114, 230, 536, 558, 640  
 Bodnar DA, 465, 466, 467, 485  
 Boersma Anderson CA, 54, 274, 603,  
 610, 618, 619  
 Boettcher F, 582  
 Boettcher FA, 274  
 Bogdanovic N, 108  
 Bonham BH, 388, 403  
 Bonhoeffer T, 56  
 Bonke BA, 505, 526  
 Bonke D, 505, 526  
 Bonneau JM, 151, 275  
 Borden GJ, 413, 423  
 Bordi F, 211, 215, 218, 220, 223, 246  
 Borg E, 586, 600, 616  
 Borisyuk A, viii, ix  
 Bormann J, 253, 272  
 Borst M, 361, 369  
 Boucher DR, 351, 375  
 Boudreau JC, 8, 66, 396, 403, 430, 457  
 Bourk TR, 7, 11, 27, 51  
 Bowman EM, 41, 52  
 Boyet S, 622  
 Boyle R, 489  
 Bradbury JW, 460, 468, 482, 485  
 Braff DL, 198  
 Brainard MS, 274, 523, 526, 530  
 Brand A, 150, 266, 272, 285, 304, 366,  
 369, 388, 403  
 Brandner ML, 218, 219, 228  
 Braun AR, 55  
 Braun K, 85, 113, 497, 526, 534  
 Braun M, 185, 195  
 Braun S, 369  
 Brawer JR, 6, 9, 11, 14, 52  
 Braz JM, 126, 127  
 Brazier MAB, 37, 52  
 Brecha N, 534  
 Bredt DS, 306, 622  
 Breese GR, 604, 617, 619, 622, 624  
 Bregman AS, 369  
 Brenowitz EA, 469, 485, 490  
 Brenowitz S, 255, 273  
 Brew HM, 290, 304, 305, 497, 526  
 Briede T, 372, 488  
 Briggs F, 38, 52, 233, 244  
 Brimijoin WO, 212, 216, 217, 227  
 Britton BH, 310



- Broadfield DC, 55  
 Brodal A, 3, 52  
 Brodal P, 43, 52, 232, 244  
 Brödel M, 5, 52  
 Bronkhorst AW, 436, 453  
 Brosch M, 474, 485  
 Bross LS, 602  
 Brown AM, 305  
 Brown M, 88, 107, 122, 127, 320, 336  
 Brown MC, 13, 18, 52, 59  
 Brown PB, 328, 338, 370, 393, 404, 428, 452  
 Brown PE, 240, 244  
 Brown-Borg HM, 616  
 Brownell WE, 4, 52  
 Browner RB, 481, 486  
 Browner RH, 115, 127, 477, 479, 485  
 Browning RA, 305, 603, 605, 611, 613, 616, 617, 619, 621  
 Brozoski TJ, 579  
 Bruce LL, 479, 486  
 Brückner S, 3, 52, 124, 27, 319, 336  
 Bruges JF, 38, 39, 43, 50, 52, 56, 59, 62, 106, 353, 369, 408, 430, 433, 443, 450, 455, 543, 553  
 Bruns V, 497, 526  
 Brunso-Bechtold JK, 4, 52, 79, 107, 116, 124, 127, 134, 140, 146, 150, 151, 162, 178, 233, 244, 540, 542, 553, 554, 556  
 Bryan JS, 271, 372, 638  
 Buchwald J, 35, 52, 65  
 Buck KJ, 498, 611, 616, 625  
 Buell TN, 378, 403  
 Buhrow SA, 308  
 Bullier J, 245  
 Bullock TH, 460, 485, 490  
 Buneo CA, 426, 450  
 Buonomano DV, 482, 485, 567, 568, 580  
 Burger RM, 152, 226, 263, 266, 267, 273, 297, 304, 407, 490, 516, 527, 582, 606, 610, 616  
 Burkard R, 580  
 Burne RA, 191, 195, 225  
 Burnstock G, 304  
 Burt DR, 57, 269, 273  
 Burton H, 57  
 Butler AB, 45, 52, 479, 486, 629, 636  
 Butler RA, 433, 435, 450, 454  
 Buttenhoff P, 245  
 Büttner C, 356, 369  
 Buunen TJF, 109  
 Byers M, 116, 127  
 Byrd JD, 116, 128, 178, 580  
 Byun MY, 623  
 Cai H, viii, ix  
 Caicedo A, 85, 107, 182, 183, 184, 186, 187, 188, 191, 193, 195, 545, 553, 611, 616  
 Caird D, 379, 397, 403, 407, 455, 606, 616  
 Calford MB, 27, 29, 30, 33, 34, 35, 52, 65, 193, 195, 203, 204, 207, 211, 212, 213, 215, 217, 218, 223, 224, 233, 244, 343, 568, 580, 582  
 Calhoun BM, 325, 327, 336, 343  
 Callaway EM, 38, 52, 233, 244  
 Campbell GA, 617  
 Canlon B, 108  
 Cant NB, 4, 5, 8, 9, 11, 15, 16, 52, 64, 66, 116, 117, 119, 120, 124, 127, 130, 136, 137, 140, 143, 149, 150, 153, 159, 178, 184, 185, 198, 428, 431, 451, 601  
 Capranica RR, 350, 352, 373, 374, 460, 468, 473, 486, 490  
 Capt M, 64  
 Cardozo BL, 374  
 Carey C, 184, 195  
 Carey J, 487, 489  
 Cariani PA, 369, 423  
 Carlile S, 435, 451, 457  
 Carlyon RP, 348, 369  
 Carney LH, ix, 379, 380, 403, 410, 453, 456  
 Carr CE, 16, 52, 250, 251, 273, 387, 403, 406, 481, 483, 486, 494, 501, 503, 505, 508, 519, 527, 531, 535, 536  
 Carrive P, 615  
 Carterette EC, 65, 342  
 Caruncho HJ, 274  
 Casebeer DJ, 305, 619  
 Caspary DM, 159, 172, 178, 249, 250, 251, 260, 262, 263, 266, 269, 270, 272, 273, 274, 277, 278, 279, 297, 305, 308, 328, 330, 332, 338, 341, 370, 373, 555, 556, 566, 567, 572,

- 579, 580, 581, 582, 600, 602, 606,  
617, 618, 622, 623
- Cassady JM, 57
- Casseday JH, 8, 15, 19, 21, 52, 53, 60,  
62, 66, 104, 107, 116, 122, 131, 132,  
140, 141, 142, 146, 150, 151, 152,  
154, 164, 180, 181, 192, 193, 195,  
198, 204, 224, 238, 244, 250, 266,  
267, 273, 274, 280, 285, 286, 297,  
300, 304, 305, 309, 316, 319, 320,  
336, 361, 362, 363, 365, 369, 370,  
375, 414, 423, 428, 430, 431, 435,  
451, 454, 494, 495, 496, 497, 501,  
503, 504, 506, 507, 508, 509, 510,  
511, 515, 516, 517, 527, 528, 529,  
532, 533, 535, 536, 606, 616, 617,  
627, 628, 629, 635, 636, 637, 638,  
640
- Castañeyra-Perdomo A, 55, 179, 225
- Cazals Y, 342, 372, 582
- Celesia GG, 43, 52
- Celio MR, 553
- Cesselin F, 53
- Cetas JS, 32, 52, 204, 205, 218, 219, 224
- Chadda R, 605, 617
- Chakravarty DN, 174, 178, 603, 616
- Chalupa LM, 602
- Chan JCK, 18, 91, 68, 114, 392, 409,  
410, 412, 428, 430, 432, 454, 458, 570
- Chang SH, 486
- Chang SO, 276, 371
- Chang Y, 606, 616
- Chan-Palay V, 195
- Chapman AG, 50, 611, 613, 616
- Chase MB, 113, 279
- Chatterjee M, 329, 336
- Chavkin C, 304
- Chen CS, 593, 599, 617
- Chen GD, 266, 273, 285, 304, 366, 369,  
418, 420, 423, 425
- Chen L, 80, 107, 273
- Chen QC, 57, 179, 245, 247, 275, 339,  
453, 530, 621, 637
- Chen XH, 189, 195, 198, 373
- Chen Y, 604, 616
- Cheng MF, 518, 526, 528
- Cheramy A, 53
- Chernock ML, 166, 178, 244, 247, 536,  
629, 636
- Cherry C, 406
- Cherry EC, 348, 369
- Chetkovich DM, 622
- Cheung SW, 39, 52, 247, 373, 489, 536
- Chimirri A, 617
- Chin L, 621
- Chiry O, 113
- Chiu J, 304, 355, 356, 373
- Chole RA, 66, 167, 199, 586, 599
- Chomsky N, 423
- Chong NW, 615
- Chow A, 304, 309
- Chowdhury SA, 449, 451
- Chulanova T, 625
- Chung DY, 348, 369
- Chung SH, 605, 617
- Chusid JG, 3, 52
- Clack JA, 493, 527
- Clarey JC, 32, 39, 52, 200, 217, 224,  
229, 456
- Clark GM, 340
- Clark KR, 452
- Clark LF, 58, 324, 339, 406
- Clarke S, 41, 43, 53, 66, 72, 113
- Clerici WJ, 116, 122, 127, 137, 150, 164,  
166, 169, 178
- Clifton RK, 49, 52, 66, 336, 450, 636
- Clock AE, 391, 582
- Clopton BM, 61, 62, 66, 67, 68, 83, 107,  
110, 274, 275, 280, 305, 322, 336,  
338, 342, 362, 369, 550, 557, 558, 616
- Clough RW, 603, 617
- Cochran S, 553
- Code RA, 38, 53, 232, 245, 350, 494,  
503, 505, 508, 527
- Coetzee WA, 290, 304
- Cohen YE, 527, 596, 600
- Colavita FB, 35, 53, 348, 369
- Colburn HS, ix, 407, 432, 453
- Coleman JR, 85, 107, 116, 122, 127,  
137, 150, 164, 166, 169, 178, 280,  
444, 455, 602, 604, 624
- Coles RB, 509, 515, 518, 519, 521, 527
- Collett L, 457
- Collingridge GL, 257, 272
- Colwell S, 5, 53
- Concas A, 270, 273
- Condon CJ, 214, 224, 356, 359, 361,  
369, 423, 473, 486

- Conlee JW, 5, 21, 30, 53, 57, 186, 187, 188, 196, 197, 505, 527  
 Conley RA, 68  
 Connor JA, 286, 304, 325, 341  
 Conrath M, 53  
 Contreras NEIR, 250, 251, 273  
 Cook AP, 486  
 Copley CA, 618  
 Corderoy A, 451  
 Correa J, 179  
 Cortés R, 301, 304  
 Cortez AM, 154, 279  
 Corwin JT, 461, 486  
 Costa E, 274  
 Coveñas R, 35, 53  
 Covey E, 21, 39, 53, 66, 97, 107, 140, 141, 150, 151, 154, 188, 189, 192, 195, 196, 198, 224, 238, 244, 273, 274, 280, 285, 304, 305, 309, 316, 319, 320, 336, 349, 361, 365, 369, 370, 371, 375, 423, 490, 494, 495, 496, 497, 501, 503, 507, 509, 510, 511, 515, 516, 517, 518, 527, 528, 529, 530, 533, 535, 606, 616, 617, 628, 629, 635, 636  
 Cowan WM, 47, 53, 54  
 Cowey A, 39, 53  
 Coyle JT, 611, 624  
 Crabbe JC, 615  
 Crabtree JW, 236, 245  
 Craig CR, 537, 538, 540, 542, 544, 546, 548, 550, 552, 554, 556, 558, 620  
 Cramer EM, 400, 403  
 Cranford JL, 399, 403  
 Crawford JD, 360, 372, 461, 462, 464, 465, 467, 482, 486, 487, 488, 489, 491  
 Creutzfeldt O, 39, 54  
 Crewther D, 344  
 Crewther S, 344  
 Criswell HE, 611, 617, 624  
 Critz SD, 291, 305  
 Crook DK, 228  
 Crosby DC, 106, 336  
 Crosby EC, 488  
 Croucher MJ, 611, 617  
 Crow G, 392, 403, 489  
 Crowe JJ, 224  
 Culling JF, 400, 401, 403  
 Cunningham AM, 306, 400, 408  
 Curró Dossi RC, 229  
 Curto KA, 620  
 Cynader MS, 40, 64, 233, 246, 436, 458  
 Cynx J, 348, 369  
 Da Silva ML, 530  
 d'Aldin C, 107  
 Dailey JW, 621, 623, 625  
 Dalkara T, 610, 617  
 Dallman MF, 624  
 Dallos P, 341, 544, 553, 554  
 Damasio H, 56  
 Dan Y, 66  
 D'Angelo WR, 380, 404, 409  
 Dani JA, 546, 553, 557  
 Danober L, 620  
 Dardennes R, 543, 553  
 Darwin CJ, 348, 369, 400, 404  
 Dascal N, 306  
 Datta S, 229  
 David EE, 391, 404  
 David J, 273  
 Davidson N, 304, 306  
 Davidson TM, 602  
 Davies WE, 103, 250, 251, 274  
 Davis KA, 91, 107, 111, 125, 127, 129, 147, 151, 199, 249, 273, 328, 336, 341, 381, 404, 408, 431, 438, 439, 440, 451, 455, 456  
 Davis KM, 611, 617  
 Davis M, 187, 197, 599  
 De Biasi S, 224  
 De Blas AL, 275  
 de Boer-van Huizen R, 478, 480, 491  
 de Cheveigné A, 360, 369, 406, 457  
 De Lanerolle NC, 526  
 de Ribaupierre F, 53, 60, 64, 200, 202, 203, 204, 224, 225, 227, 228, 345  
 de Ribaupierre Y, 53, 60, 64, 225, 227, 228  
 De Sarro A, 617  
 De Sarro G, 617  
 Dear SP, 513, 528  
 Dehmel S, 137, 151  
 Deitmer JW, 305  
 Del Bel EA, 619  
 Del Mar Perez Delgado MM, 179  
 Delaney S, 451  
 Delgutte B, 217, 224, 328, 337, 355, 369,

- 374, 401, 404, 418, 420, 423, 438,  
440, 447, 451, 453
- Demuth RM, 602
- Dent ML, 446, 451, 454, 457, 528, 529
- Depaulis A, 622
- Depireux DA, 339
- Derer P, 110, 340
- Deschênes M, 39, 54, 236, 245
- Després G, 308, 622
- Devaud LL, 605, 611, 617
- Dezso A, 71, 85, 107, 501, 528, 533
- Diehl JJ, 68
- Di Scala G, 615
- Diamond IT, 33, 39, 40, 42, 54, 57, 60,  
247, 452, 454, 521, 636, 637, 638
- Díaz C, 476, 486
- DiCaprio RA, 152
- Dickerson L, 52
- Dickhaus H, 50, 336
- Diehl JJ, 43, 54, 68, 114, 247, 535, 640
- Diekamp B, 469, 473, 486
- Dimyan MA, 51
- Ding D, 601
- Dingledine R, 554
- Dinse HE, 44, 53
- Doble A, 269, 274
- Dodson PD, 290, 305
- Dooling RJ, 337, 446, 451, 454, 494,  
527, 528
- Doroshenko NZ, 159, 178
- Dorrscheidt GJ, 151
- Doubell TP, 84, 107, 238, 245
- Doucet JR, 38, 43, 54, 134, 144, 151,  
197, 231, 245
- Dougherty WJ, 130
- Doupe AJ, 344, 534
- Drakontides AB, 481, 486
- Dreyer A, 443, 451
- Druga R, 116, 119, 127, 164, 166, 169,  
178, 344
- du Lac S, 521, 528
- Ducic I, 269, 274
- Duncan GE, 622
- Dunlop CW, 35, 50, 208, 213, 223
- Dupont J, 582, 615
- Durand SE, 505
- Durlach NI, 382, 404, 408
- Duval D, 306
- Dykes RW, 57, 60
- Eagleton C, 535
- Ebbesson SOE, 477, 486
- Ebihara K, 620
- Echteler SM, 544, 553
- Eckardt MJ, 603, 617
- Edds-Walton PL, 461, 464, 465, 486
- Edeline J-M, 211, 212, 213, 215, 218,  
224, 226
- Edelman GM, 3, 54, 337, 344, 345, 406,  
408
- Edwards CJ, 474, 475, 486
- Edwards SB, 189, 196, 198
- Eells JB, 617
- Eggermont JJ, 39, 46, 47, 54, 325, 337,  
350, 351, 370, 473, 486
- Egorova M, 315, 316, 317, 318, 319,  
322, 323, 324, 328, 330, 337, 338,  
344, 421, 423, 443, 451
- Ehrenguber MU, 304
- Ehret G, 25, 27, 30, 38, 41, 54, 66, 90,  
107, 151, 169, 170, 172, 174, 180,  
184, 194, 196, 213, 224, 236, 247,  
253, 254, 278, 283, 287, 288, 289,  
309, 315, 316, 317, 318, 319, 320,  
322, 323, 324, 325, 327, 328, 329,  
330, 331, 332, 333, 334, 335, 337,  
338, 339, 342, 343, 344, 345, 349,  
359, 362, 363, 370, 374, 423, 431,  
443, 448, 451, 457, 543, 557, 627,  
632, 634, 636, 637, 640
- Ehrlich D, 107, 273, 285, 300, 304, 305,  
365, 366, 369, 370, 423, 527, 528,  
616
- Ehrlich I, 305, 515
- Eisenberg JF, 632, 636
- Ekelid M, 374, 491
- Elverland HH, 18, 54
- Emerson RG, 372, 531
- Emson PC, 112
- Endepols H, 238, 245
- Endo T, 1, 51, 78, 85, 110, 244, 268,  
396, 440, 481, 519, 606, 625
- Engeland B, 307
- Epping WJM, 350, 351, 370
- Erisir A, 309
- Erre JP, 582
- Erulkar SD, 27, 60, 208, 227, 349, 363,  
366, 372, 627, 628, 636, 638
- Erway LC, 542, 586, 599, 602

- Escabi MA, 60, 325, 327, 333, 338, 340, 341, 421, 422, 423, 634, 637
- Eshete F, 107
- Esser K-H, 420, 423, 451
- Euston DR, 521, 528, 534
- Evans EF, 13, 17, 38, 54, 338, 408
- Evans MS, 109, 276, 307, 610, 617, 619, 621, 622
- Evrard P, 615
- Eybalin M, 107, 545, 553, 611, 616
- Eysel UT, 214, 224, 637
- Facelle T, 485
- Faingold CL, 31, 54, 109, 174, 178, 247, 248, 249, 250, 251, 254, 257, 258, 260, 262, 267, 274, 276, 297, 300, 301, 305, 307, 308, 335, 338, 349, 370, 603, 604, 605, 606, 607, 608, 609, 610, 611, 612, 613, 614, 616, 617, 618, 619, 620, 621, 622, 623, 625
- Falconi A, 180
- Falzarano PR, 599
- Fant G, 412, 423, 554
- Farb DH, 278, 624
- Farley GR, 66, 301, 305
- Fastl H, 337, 354
- Faulstich M, 531
- Faure PA, 366, 370, 512, 515, 516, 528
- Fay RR, 3, 17, 52, 53, 54, 60, 62, 63, 64, 67, 71, 75, 82, 107, 108, 110, 129, 150, 152, 153, 154, 155, 156, 178, 195, 196, 197, 199, 224, 229, 230, 247, 273, 339, 341, 375, 376, 405, 454, 458, 459, 461, 465, 467, 483, 486, 487, 488, 489, 490, 491, 492, 527, 528, 529, 636, 638, 640
- Faye-Lund H, 71, 75, 82, 107, 155, 157, 178, 537, 553, 629, 630, 637
- Feeser HR, 208, 227
- Fekete DM, 5, 54, 116, 129
- Feldman DE, 249, 250, 258, 274, 297, 305, 523, 528
- Feldman ML, 16, 56
- Feldman RS, 391, 410, 523
- Feldtkeller R, 329, 345
- Feliciano M, 176, 178, 181, 232, 245, 611, 619, 639
- Felsheim C, 363, 370
- Feng AS, 3, 7, 23, 56, 116, 130, 162, 181, 186, 199, 212, 213, 214, 216, 224, 225, 226, 281, 311, 315, 338, 364, 366, 369, 370, 465, 469, 471, 472, 473, 474, 477, 486, 487, 513, 528, 534, 605, 606, 619, 623
- Feng H-J, 619, 623
- Fenzl T, 192, 196
- Fernandes MA, 339, 370
- Fernandez C, 115, 127
- Ferragamo MJ, 305
- Ferreri G, 617
- Ferres-Torres R, 55, 179, 225, 554
- Ferster D, 18, 55
- Fex J, 51, 320, 339
- Fields RD, 73, 89, 107, 113, 125
- Fillit P, 276, 555
- Findlay G, 625
- Fine ML, 487
- Finlayson PG, 135, 137, 151
- Finn SE, 622
- Fischer R, 337
- Fischer S, 198
- Fitch WT, 413, 423
- Fitzpatrick D, 38, 55, 97, 104, 106, 107, 109, 140, 150, 220, 224, 240, 245, 380, 382, 385, 387, 390, 392, 399, 402, 404, 406, 447, 449, 450, 451
- Fitzpatrick KA, 71, 107, 319, 338, 537, 553
- Flachskamm C, 372
- Flanagan JL, 325, 338
- Fleming WW, 620, 622
- Fletcher H, 322, 324, 338
- Fletcher KA, 467, 487
- Flood DG, 581, 600
- Flottorp G, 345
- Foeller E, 257, 274
- Foidart A, 526
- Folander K, 308
- Foley BJ, 305
- Forest R, 226
- Förster CR, 570, 580
- Forsythe ID, 103, 106, 290, 304, 305, 310, 497, 526
- Fortune ES, 216, 226, 411, 424, 474, 482, 487, 489
- Foster RE, 21, 55, 479, 480, 487
- Fotuhi M, 306
- Frachet B, 276, 555
- Frackowiak R, 225
- Frank G, 31, 68, 199, 324, 337, 531

- Frankenreiter M, 324, 337  
 Frankland PW, 31, 68, 199  
 Franzen L, 310  
 Frech MJ, 298, 305  
 Fremouw T, 150, 195, 273, 309, 370,  
 527, 528, 533, 636  
 Friauf E, 4, 55, 85, 109, 305, 320, 338,  
 539, 544, 546, 553, 554, 555, 557, 558  
 Friedman MA, 16, 52, 342  
 Frigyesi T, 236, 245  
 Frisina DR, 559, 560, 566, 580, 582  
 Frisina RD, 56, 116, 128, 159, 164, 166,  
 169, 178, 188, 191, 192, 196, 201,  
 202, 203, 204, 218, 224, 225, 228,  
 280, 372, 375, 425, 473, 487, 503,  
 528, 531, 560, 567, 578, 580, 583, 584  
 Fritschy JM, 277, 617  
 Fritz J, 63, 637, 639  
 Fritzsich B, 487–488, 637, 640  
 Frost BJ, 535, 600  
 Frost S, 553  
 Fruchter D, 425  
 Frye GD, 603, 606, 610, 619, 622  
 Fubara BM, 253, 274, 300, 305, 497, 528  
 Fuentes V, 157, 164, 178, 544, 548, 550,  
 554  
 Fuentes-Santamaria V, 544, 548, 550, 554  
 Fujimoto Y, 195  
 Fujino K, 188, 196  
 Fujita I, 263, 273, 274, 501, 521, 527,  
 528  
 Fukui Y, 198  
 Fullerton BC, 63, 639  
 Fuzessery ZM, 216, 225, 240, 245, 246,  
 266, 267, 274, 297, 305, 315, 316,  
 328, 338, 362, 363, 364, 365, 370,  
 409, 472, 487, 494, 507, 511, 515,  
 516, 517, 529, 531, 606, 619, 634,  
 637, 638  
**Gabriele ML**, 539, 541, 543, 548, 550,  
 554, 555  
 Gago G, 217, 225, 330, 332, 339, 394,  
 395, 396, 405  
 Gahr M, 498, 529  
 Gaidaenko GV, 485  
 Galambos R, 64  
 Galazyuk AV, 224, 332, 338  
 Gales RS, 342  
 Galindo-Mireles D, 55, 179, 225  
 Gall CM, 623  
 Gall WE, 54, 337, 344, 345, 406, 408  
 Games KD, 42, 55, 232, 245  
 Gannon PJ, 43, 55  
 Gans C, 491  
 Gans DP, 424, 461, 467, 482, 485, 486,  
 488, 491, 559  
 Gao E, 236, 245, 246, 344, 449, 451,  
 518, 529, 530, 546, 554, 639  
 Garcia AN, 603, 604, 605, 619, 620  
 García VR, 43, 50  
 Garcia-Cairasco N, 603, 604, 619  
 Gardner SM, 103, 107  
 Garell PC, 56  
 Gareri P, 617  
 Garey LJ, 13, 17, 67  
 Garraghty PE, 60  
 Garrett KM, 310  
 Gaston KC, 4, 52, 431, 450  
 Gauger B, 531  
 Gaza WC, 611, 620  
 Gazzaniga MS, 43  
 Gehi A, 58  
 Gehlbach G, 274, 305, 338, 370, 618  
 Geisler CD, 64, 408, 455  
 Geniec P, 75, 82, 108, 537, 554, 629, 637  
 Gerber A, 600  
 Gerhardt HC, 468, 469, 473, 486, 487  
 Gerken GM, 268, 274  
 Gerren RA, 35, 55, 238, 245  
 Gersuni GV, 316, 338  
 Gescheider GA, 56  
 Geschwind N, 43, 54, 55  
 Geyer MA, 198  
 Ghansah E, 616  
 Gibson A, 197  
 Gil Verona JA, 603, 620  
 Gilbert CD, 38, 59  
 Gilman S, 52, 223  
 Girard P, 245  
 Giraud AL, 214, 225  
 Gitto R, 617  
 Gleason AT, 113  
 Gleich O, 364, 370, 489  
 Glendenning KK, 18, 19, 22, 23, 30, 45,  
 54, 55, 56, 80, 91, 108, 134, 140, 144,  
 146, 151, 179, 225, 250, 251, 270,  
 274, 431, 452, 454, 456  
 Glenn L, 339  
 Glezer II, 85, 108

- Glickstein M, 197  
 Godfrey DA, 58, 107, 129, 130, 198, 280, 527  
 Godukhin O, 625  
 Goldberg JM, 54, 157, 180, 328, 338, 342, 349, 370, 374, 393, 396, 404, 428, 452, 639  
 Golding NL, 290, 305  
 Goldman-Rakic PS, 63, 639  
 Goldsmith JD, 611, 620  
 Goldstein NA, 64  
 Goldstein-Daruech N, 611, 620  
 Gomes EL, 619  
 Gomez BP, 620  
 Gomez Carretero ME, 620  
 Gonzalez LP, 605, 625  
 González-González B, 55, 108, 151, 179, 275, 620  
 González-Hernández TH, 3, 23, 55, 99, 108, 135, 137, 151, 157, 159, 164, 169, 170, 172, 179, 202, 225, 250, 275, 539, 543, 555, 606, 620  
 González-Lima F, 71, 85, 108, 111  
 Goodson JL, 461, 464, 487  
 Gooler DM, 228, 473, 487  
 Goossens HH, 436, 452  
 Gopalswamy P, 408  
 Gordon M, 204, 212, 216, 225, 362, 363, 370  
 Goreva OA, 435, 452  
 Gorga MP, 305  
 Gorodetskaya MM, 352, 368, 472, 485  
 Gould EM, 610, 620  
 Granger EM, 440, 454  
 Granner M, 56  
 Grant L, 343  
 Grant S, 51, 336  
 Grantham DW, 378, 404  
 Grasse KL, 59  
 Gratton MA, 279  
 Green EJ, 229  
 Green GGR, 304, 336  
 Greenberg S, 5, 31, 47, 63, 125, 129, 348, 370, 425  
 Greenblatt EP, 622  
 Greenwood DD, 324, 338, 339, 342, 374, 622, 639  
 Greenwood RS, 622  
 Gregory KM, 5, 55  
 Gressens P, 615  
 Griffith BL, 617  
 Grigg JJ, 290, 291, 305  
 Grinnell AD, 244, 494, 529, 628, 637  
 Grinvald A, 56  
 Grobin AC, 611, 620  
 Groh JM, 444, 452  
 Grose CD, 116, 130, 201, 202, 204, 215, 218, 230, 246, 344, 352, 375, 531, 638  
 Gross NB, 41, 64, 70, 71, 96, 218, 225, 408, 455, 624, 629, 639  
 Grothe B, 66, 150, 226, 263, 266, 267, 272, 275, 280, 297, 300, 304, 307, 310, 362, 363, 369, 371, 375, 388, 403, 408, 503, 517, 518, 527, 529, 531, 534, 583, 628, 632, 637  
 Guillery RW, 37, 65, 106, 223  
 Guimaraes FS, 619  
 Guinan JJ Jr, 15, 16, 23, 56, 58, 91, 108, 135, 151, 319, 328, 339, 430, 452, 461, 489  
 Guinan SS, 56, 108, 151, 339  
 Gulick WL, 56  
 Günturkun O, 535  
 Gupta A, 103, 108  
 Guthmann A, 112, 557  
 Guthrie PB, 308  
 Gutiérrez A, 269, 275  
 Guttinger HR, 529  
 Guttman N, 404  
 Guzman SJ, 407  
**Haack B**, 325, 337  
 Habbicht H, 66, 280, 302, 305, 310, 375, 497, 529, 534, 567, 580, 583  
 Haberly LB, ix, 61, 111, 152, 180, 307, 341, 371, 407, 638  
 Hackett TA, 39, 43, 56, 57  
 Hafter ER, 391, 407  
 Hage SR, 131, 324, 338, 339, 362, 363, 370, 451, 634, 637  
 Haggard MP, 339, 370  
 Hahner L, 616  
 Hajalilou R, 341  
 Hall JC, 23, 56, 266, 267, 274, 297, 305, 328, 338, 363, 364, 365, 370, 472, 473, 474, 477, 482, 487, 507, 511, 515, 529, 606, 619  
 Hall JW, 325, 339, 348, 370  
 Hall WC, 21, 55, 196, 203, 204, 227, 479, 480, 487, 528

- Hallanger AE, 59  
 Halsey MJ, 616  
 Halvorsen MB, 226  
 Hamann I, 370  
 Hammerschmidt B, 553  
 Hammond BM, 369, 423  
 Han LY, 581, 600  
 Hanaya R, 620  
 Hancock KE, 401, 404  
 Hanson JT, 109, 423, 488, 622  
 Haplea S, 511, 529  
 Happe HK, 301, 308, 428, 446, 546, 556  
 Hardie NA, 547, 548, 554, 556  
 Hardingham GE, 611, 620  
 Haresign T, 536  
 Harmison GG, 51  
 Harms MP, 214, 225  
 Harnischfeger G, 379, 404  
 Harpman JA, 115, 131, 627, 640  
 Harris DM, 554  
 Harris KS, 423  
 Harris RA, 611, 616, 620, 622, 624  
 Harrison JB, 244  
 Harrison JM, 16, 17, 52, 56  
 Harrison RV, 548, 554, 578, 580  
 Hart GP, 616  
 Harting JK, 83, 108, 188, 196, 238, 245, 632, 636, 637  
 Hartley M, 554  
 Hartmann R, 373, 488  
 Hartmann WM, 488  
 Hartung K, 399, 404, 409  
 Harty TP, 300, 306  
 Harvey AR, 281  
 Harvey RJ, 557  
 Hashiguchi H, 620  
 Hashikawa T, 56  
 Hattori T, 359, 370  
 Hattox AM, 190, 196  
 Hauser M, 228, 485  
 Hauser MD, 423  
 Häusser M, 66  
 He J, 40, 56, 210, 213, 218, 225,  
 Heberlain AS, 486  
 Heffner HE, 43, 56, 348, 371, 387, 404, 405, 407, 432, 434, 452, 580  
 Heffner RS, 43, 56, 387, 404, 405, 432, 434, 452, 564, 580  
 Hefti BJ, 68, 114, 233, 245, 247, 535, 640  
 Heil P, 39, 56, 329, 330, 339, 356, 371, 372, 460, 485, 508, 529, 582  
 Heiligenberg W, 460, 485  
 Heinemann S, 554  
 Heinemann SF, 554  
 Heinemann SH, 293, 308, 310  
 Heiser JB, 490  
 Heizmann CW, 112, 526  
 Held H, 4, 9, 13, 45, 55, 128, 157, 179, 393, 413, 430, 446, 452, 474, 614  
 Helfert RH, 51, 132, 134, 135, 137, 141, 142, 143, 151, 162, 172, 178, 198, 249, 250, 251, 269, 275, 567, 580, 581  
 Hellweg F-C, 54  
 Henderson J, 425, 582, 599, 601  
 Hendry SHC, 39, 56  
 Henkel CK, ix, 30, 61, 66, 82, 97, 111, 112, 113, 123, 128, 130, 134, 141, 147, 151, 153, 154, 155, 157, 179, 180, 189, 190, 196, 204, 205, 225, 226, 279, 341, 407, 537, 539, 540, 553, 554, 555, 556, 557, 558, 638  
 Hennevin E, 224  
 Henning GB, 400, 409  
 Henry KR, 586, 592, 593, 599, 602, 604, 620  
 Henson MM, 529  
 Henson OW Jr, 23, 56, 194, 196, 250, 275, 508, 529, 533  
 Herbert H, 112, 182, 183, 184, 186, 187, 188, 189, 191, 192, 193, 195, 196, 197, 503, 529, 557, 632, 637  
 Herman P, 37, 65, 208, 211, 229, 276, 464, 491, 555  
 Hernandez-Cruz A, 309  
 Hernandez-Pineda R, 309  
 Herrera M, 164, 166, 179  
 Herrnberger B, 334, 339  
 Hestrin S, 294, 306  
 Hewitt MJ, xiii, ix, 361, 371  
 Heyer WR, 469, 491  
 Hienz RD, 454  
 Higashiyama A, 311, 613, 620  
 Highstein SM, 461, 464, 486, 487, 489  
 Hilbert J, 425  
 Hillery CM, 350, 371  
 Hillman DE, 64, 310  
 Hind JE, 56, 64, 277, 342, 374, 408, 427, 450, 454, 455, 461, 464, 539, 627, 639  
 Hinman C, 52



- Hinrichsen CF, 196  
Hirai H, 581  
Hirai K, 310  
Hirsch JA, 288, 306, 605, 616, 620, 622  
Hirsh IJ, 378, 396, 397, 405  
Hishinuma F, 310  
Ho IK, 605, 623  
Hobbs M, 557  
Hockfield S, 111, 308  
Hodge CW, 624  
Hodos W, 45, 52, 629, 636  
Hof PR, 108, 580  
Hoffman HS, 61, 62, 66, 67, 68, 110, 198, 274, 275, 280, 305, 338, 342, 409, 557, 596, 599, 616, 618  
Hoffmann K-P, 198  
Hoffmann WE, 274, 618  
Hofman PM, 436, 452  
Hofstetter P, 275, 581  
Hohenboken WD, 603, 620  
Holder TM, 273, 277, 555, 579, 581, 600  
Hollmann M, 554  
Holloway RL, 55  
Honda CN, 61  
Hood HM, 295, 380, 539, 559, 564, 588, 594, 613, 625  
Horn AG, 454  
Horn AKE, 229, 280, 344, 534  
Horner K, 342, 372  
Hornung JP, 64  
Horvath M, 581  
Hose B, 371, 384  
Hosea EW, 594, 601  
Hosomi H, 298, 306, 569, 581, 620  
Houtani T, 180  
Howard MA, 43, 56  
Hoy RR, 63, 446, 457  
Hoyer D, 558  
Hsieh CY, 553  
Hu B, 225, 227, 503, 533  
Huang AY, 434, 435, 436, 437, 440, 452, 454  
Huang C, 320, 339  
Huang CL, 33, 35, 37, 41, 56, 68, 230, 232, 238, 245, 491, 633, 637  
Hübener M, 38, 56  
Huber GC, 378, 409, 488  
Huerta MF, 23, 43, 61, 70, 82, 87, 110, 124, 129, 188, 197, 238, 245  
Huffman RF, 23, 56, 194, 196, 250, 275, 362, 371  
Huggins WH, 400, 401, 403, 404  
Hughes JR, 64  
Hughes LF, 272, 273, 275, 279, 579, 581  
Hugues M, 306  
Huie D, 67  
Hume RI, 545, 554  
Hunter C, 9, 67, 277, 280  
Hunter KP, 280, 281, 599, 602  
Hupé J-M, 237, 245  
Hurley LM, 109, 186, 196, 226, 268, 275, 301, 306, 364, 371, 423, 488, 497, 529, 567, 581  
Huston JP, 195  
Hutchings ME, 556, 581, 600  
Hutsler JJ, 43, 56  
Hutson KA, 4, 23, 55, 56, 80, 108, 140, 146, 151, 157, 172, 179, 204, 225, 274, 452  
Huy PT, 276, 308, 555  
Hwang PM, 292, 306  
Hyde PS, 96, 157, 158, 505, 524, 529  
Ibrahim D, 554, 580  
Ichiyama M, 195  
Idrizbegovic E, 85, 108  
Iida K, 604, 620  
Ikeda M, 180, 557  
Ikeuchi Y, 293, 306  
Illing R-B, 29, 81, 118, 169, 238, 464, 570, 571, 580, 581  
Imai H, 277  
Imig TJ, 32, 34, 35, 36, 38, 39, 40, 42, 52, 57, 60, 63, 215, 217, 219, 224, 225, 228, 229, 232, 233, 245, 246, 431, 438, 440, 443, 452, 456  
Inati S, 452  
Inderkum A, 620  
Irons WA, 225, 229, 452, 456  
Irvine DRF, 10, 19, 20, 22, 26, 27, 28, 30, 57, 90, 91, 108, 135, 151, 188, 196, 217, 220, 225, 228, 330, 332, 334, 339, 394, 395, 396, 405, 410, 443, 452, 458, 543, 556, 571, 572, 578, 581, 582, 627, 637  
Irving R, 56  
Isbey SF, 224, 636  
Ishida N, 311, 606, 620

- Ishihara K, 620  
 Ison JR, 375, 580, 583, 599, 625  
 Ito M, 151  
 Itoh K, 60, 107, 226  
 Itoh M, 195, 198  
 Ivanov DP, 111  
 Ivarsson C, 217, 225  
 Iwata J, 59  
  
**Jack** JJB, 9, 57  
 Jackson BS, 329, 339  
 Jackson LM, 602  
 Jahnsen H, 208, 225  
 Jähn-Siebert TK, 369  
 James AC, 245  
 James N, 341  
 Jane JA, 31, 57, 449, 452, 632, 637  
 Janis CM, 490  
 Jarreau PH, 553  
 Jarolimek W, 277  
 Jarvis ED, 505, 530  
 Javel E, 305  
 Jeffress LA, 17, 57, 380, 381, 382, 384, 385, 387, 396, 404, 405, 428, 453, 514, 530  
 Jen PH-S, 7, 27, 29, 57, 66, 177, 179, 199, 236, 245, 247, 250, 251, 275, 300, 307, 315, 316, 319, 323, 328, 332, 339, 340, 341, 345, 373, 400, 405, 410, 448, 490, 518, 530, 532, 533, 536, 606, 611, 621, 632, 637  
 Jenkins WM, 23, 31, 32, 57, 385, 405, 435, 440, 443, 453  
 Ji W, 456, 530, 534  
 Jiang D, 276, 341, 405, 407  
 Jiang ZD, 109, 196, 197  
 Jin C, 188, 196, 457  
 Jishage K, 131  
 Jobe PC, 603, 604, 606, 616, 617, 618, 621, 623, 625  
 Johannesma PIM, 337, 343  
 Johansson RS, 426, 457  
 Johnson DH, 8, 15, 66, 428, 453  
 Johnson MS, 605, 617  
 Johnson KB, 622  
 Johnsrude I, 225  
 Johnston GA, 251, 273  
 Jonas P, 7, 57, 300, 306  
 Jones AW, 617  
 Jones DR, 61  
 Jones EG, 23, 25, 26, 29, 32, 33, 35, 39, 41, 52, 54, 56, 57, 61, 63, 67, 112, 166, 180, 226, 246, 373, 557, 639  
 Jones LS, 624  
 Jones TA, 616  
 Joris PX, 91, 108, 130, 153, 224, 346, 371, 384, 388, 392, 393, 405, 406, 408, 428, 451, 453, 456  
 Josephson EM, 116, 120, 128  
 Jubelier EM, 228  
 Judge PW, 58  
 Julius D, 624  
 Jürgens U, 192, 196, 424, 464, 488  
  
**Kaas** JH, 7, 27, 39, 56, 57, 60  
 Kachunts AS, 477, 480, 485, 488  
 Kaczmarek LK, 111, 291, 292, 307, 308, 309, 310  
 Kadia SC, 411, 425  
 Kagamiyama H, 131  
 Kageyama GH, 555  
 Kai N, 604, 621  
 Kaiser A, 497, 530  
 Kajikawa Y, 310  
 Kaltenbach JA, 590, 599  
 Kamada T, 341  
 Kamatchi GL, 269, 273  
 Kamath V, 374, 491  
 Kandler K, 191, 192, 197, 539, 554  
 Kane EC, 58  
 Kane ES, 5, 21, 30, 52, 53, 58, 186, 187, 188, 196, 197  
 Kaneko T, 60  
 Kanold PO, 292, 306  
 Kanwal JS, 227, 423, 424, 494, 530  
 Kapfer C, 150  
 Kappers CUA, 483, 488  
 Karagülle T, 109, 340, 555  
 Karamürsel S, 490  
 Karapas F, 115, 127  
 Karschin A, 306  
 Karschin C, 293, 306, 307  
 Karten HJ, 498, 504, 506, 530, 535  
 Kato N, 311, 620  
 Katsuki Y, 583, 627, 637  
 Kauer JA, 304, 369, 528  
 Kavanagh GL, 172, 179, 435, 443, 453  
 Kawahara N, 621

- Kawai J, 605, 621  
Kawamura K, 41, 58, 226  
Kawano J, 556  
Kawasaki M, 131  
Kazee AM, 567, 581, 595, 600  
Keating MJ, 51, 336  
Keiger CJ, 539, 541, 542, 543, 554  
Keipert C, ix, 7  
Kellar KJ, 309  
Keller A, 13, 37, 67, 196, 399, 446, 447, 518, 521, 603, 624  
Keller CH, 406, 453, 530, 534  
Kelly JB, 32, 35, 37, 38, 42, 58, 68, 90, 107, 108, 114, 137, 151, 172, 179, 181, 249, 250, 251, 254, 258, 259, 262, 263, 264, 265, 266, 268, 273, 275, 276, 279, 280, 281, 288, 290, 291, 297, 300, 303, 306, 308, 310, 311, 322, 330, 339, 379, 396, 399, 406, 408, 435, 443, 446, 453, 567, 581, 606, 611, 621, 625  
Kemp EH, 639  
Kemp JA, 280  
Kempf S, 339  
Kennedy JM, 245  
Kennedy MC, 485, 488  
Kentros C, 310  
Kenyon CE, 50, 116, 126, 127  
Kerckhoven C, 406  
Khan A, 226  
Khan ZU, 275  
Khurana V, 623  
Kiang NY-S, 9, 11, 58, 63, 339, 342, 406, 639  
Kidd SA, 262, 275, 280  
Kiesmann M, 622  
Kil J, 555  
Kilgard MP, 13, 43, 58  
Killackey HP, 116, 128, 152, 556  
Kim CS, 276  
Kim DO, 13, 58, 351, 352, 371, 404, 407, 451, 455, 600  
King AJ, 107, 109, 196, 197, 245, 246, 556, 581, 600  
Kinoshita N, 126, 128  
Kinouchi Y, 56  
Kirsch J, 553  
Kirson ED, 295, 306  
Kiss A, 207, 226  
Kistler DJ, 432, 433, 436, 457  
Kisvárdy Z, 633, 637  
Kitao Y, 116, 128, 151, 556  
Kitch R, 487  
Kittle M, 606, 621  
Kittler JT, 606, 621  
Kitzes LM, 38, 58, 62, 67, 116, 128, 152, 213, 229, 322, 332, 339, 343, 395, 408, 547, 550, 555, 556, 594, 600  
Klatzo I, 621  
Klein DJ, 327, 339  
Kleinschmidt A, 225  
Klepper A, 196  
Klinke R, 373, 379, 403, 488  
Klotz DA, 178, 196, 225  
Klug A, 104, 109, 152, 210, 212, 226, 262, 267, 275, 297, 300, 307, 407, 412, 415, 418, 421, 423, 460, 488, 490, 517, 530, 582  
Klump GM, 337, 341, 370, 454  
Knight PL, 59  
Knudsen EI, 29, 58, 84, 109, 249, 250, 258, 263, 274, 281, 297, 305, 461, 488, 494, 498, 500, 501, 505, 508, 519, 521, 523, 524, 526, 527, 528, 529, 530, 531, 532, 536, 550, 555, 630, 637  
Ko KH, 621  
Koay G, 580  
Kobler JB, 196, 224, 528, 636  
Koch R, 533  
Koch U, 275, 307, 371, 531  
Koh DS, 397, 409  
Kohlrausch A, 397, 409  
Koltchine VV, 622  
Kondoh A, 180  
Konishi M, 29, 58, 263, 273, 274, 387, 403, 406, 485, 493, 501, 505, 508, 518, 519, 521, 523, 527, 528, 531, 532, 533, 534, 535, 630, 639  
Koob GF, 605, 621  
Köppl C, 337, 341, 519, 531, 637  
Kopp-Scheinpflug C, 151  
Kordyban M, 602  
Korngut LM, 600  
Koskinas GN, 37, 67  
Kössl M, 66, 229, 274, 280, 310, 344, 508, 531, 534, 583

- Kotak VC, 180, 277, 300, 307, 308, 309  
 Kowalchuk NE, 547, 556, 600  
 Kowalski N, 325, 339, 343  
 Koyama H, 489, 539, 556  
 Koziel V, 622  
 Kozloski J, 461, 464, 467, 488, 491  
 Krasowski MD, 622  
 Kraushaar U, 298, 300, 301, 307, 546, 555  
 Krettek JE, 242, 246  
 Krishna BS, 265, 276, 351, 352, 353, 355, 371, 473, 488  
 Kroodsma DE, 529  
 Krout KE, 238, 246  
 Krubitzer LA, 59  
 Kubke MF, 501, 520, 531, 535  
 Kudo M, 36, 58, 60, 84, 109, 116, 119, 128, 132, 138, 142, 147, 151, 157, 166, 172, 179, 192, 197, 201, 203, 226, 328, 340, 369, 539, 555, 556  
 Kudoh M, 369  
 Kuhar MJ, 281  
 Kujawa SG, 620  
 Kujirai K, 229  
 Kulesza RJ Jr, 125, 128, 136, 137, 138, 152, 155, 180, 251, 254, 255, 276, 610, 621  
 Kulkarni A, 432, 453  
 Kumoi K, 134, 152  
 Kuner T, 295, 307  
 Kungel M, 546, 553, 555, 558  
 Künzle H, 479, 480, 488  
 Kurisu K, 620  
 Kuriyama H, 309, 557  
 Kuriyama K, 250, 251, 279  
 Kuroda Y, 620  
 Kuwabara N, 137, 138, 143, 152, 231, 238, 246  
 Kuwada S, ix, 61, 75, 76, 91, 97, 106, 107, 109, 110, 111, 113, 114, 116, 129, 135, 137, 147, 152, 180, 217, 226, 285, 307, 341, 349, 371, 378, 379, 380, 384, 385, 389, 390, 402, 404, 406, 407, 409, 410, 428, 430, 450, 451, 453, 456, 458, 557, 638  
 Kvale M, 634, 637  
 Kvašňák E, 229, 344, 425, 456  
 Kwon J, 603, 604, 621  
 Lafaille P, 51  
 Laird HE, 616, 618  
 Lajtha A, 272  
 Lam DM-K, 154, 279  
 LaMotte RH, 426, 453  
 Langendijk EH, 436, 453  
 Langford TL, 397, 403, 406  
 Langner G, 25, 27, 30, 46, 58, 65, 88, 90, 112, 176, 181, 213, 214, 226, 319, 320, 323, 324, 325, 328, 343, 350, 351, 352, 353, 354, 356, 359, 360, 361, 367, 368, 369, 371, 372, 374, 473, 484, 488, 533  
 Large EW, 372, 488  
 Larue DT, 5, 33, 47, 51, 56, 68, 114, 154, 195, 199, 207, 223, 230, 244, 247, 281, 345, 535, 536, 630, 633, 636, 637, 640  
 Lasley SM, 611, 621  
 Laszig R, 581  
 Latorre R, 310  
 Lau D, 309  
 Laurie DJ, 281, 546, 555  
 Lauterborn JC, 623  
 Lazdunski M, 306  
 Le Beau FEN, 91, 110, 180, 197, 226, 246, 260, 262, 263, 267, 276, 297, 300, 307, 316, 318, 319, 340, 349, 372, 442, 453, 606, 621  
 Leake PA, 6, 58, 391, 406, 508, 536  
 Leake-Jones PA, 6, 58  
 Leakey DM, 391, 406  
 LeDoux JE, 35, 38, 59, 63, 201, 203, 204, 205, 211, 215, 218, 220, 223, 226, 246  
 Ledwith III JV, 13, 52  
 Lee DJ, 622  
 Lee J, 474, 488  
 Lee Y, 190, 197, 557  
 Leergaard TB, 59, 110, 180  
 Leibler LM, 505, 531  
 Leitner DS, 596, 600  
 Lennartz RC, 211, 213, 226  
 Leonard C, 309  
 Leroy SA, 216, 226, 230, 319, 340, 352, 375, 415, 424, 431, 453, 513, 515, 531, 535  
 Leslie FM, 553  
 Lesser HD, 125, 204, 250, 350, 363, 372, 482, 516, 531

- Lester HA, 304  
Leumann L, 197  
LeVay S, 18, 38, 55, 59  
Levelt WJM, 324, 341  
Leverenz EL, 489  
Levey AI, 35, 59  
Levine RA, 63, 639  
Levitan ES, 307  
Levitsky W, 43, 55  
Lewis ER, 388, 403, 461, 469, 472, 488, 489, 619  
Li D, 535  
Li FF, 66  
Li H, 425  
Li HS, 586, 600  
Li L, 105, 114, 181, 199, 250, 251, 254, 262, 275, 276, 281, 396, 406, 530, 596, 600, 606, 621  
Li RY, 55, 452  
Li RY-S, 55  
Li W, 291, 307  
Li Y, 109, 257, 276, 283, 287, 288, 289, 292, 294, 297, 298, 307, 610, 611, 617, 619, 621, 622  
Liao WH, 276, 555  
Liberman MC, 17, 18, 54, 59, 316, 329, 340  
Licklider JCR, 360, 372, 378, 396, 406  
Lien NT, 623  
Lifschitz WS, 225  
Lim D, 255, 276  
Lim HH, 582  
Lin WY, 487  
Lindemann W, 399, 406  
Lingenhöhl K, 310  
Linke R, 201, 204, 226  
Linville C, 553  
Lippe W, 544, 555, 557  
Liscum A, 151  
Lisker L, 418, 423  
Litovsky RY, 224, 399, 406, 407, 447, 449, 451, 453  
Liu Q, 68  
Liu SQ, 291, 307  
Llano D, 212, 213, 214, 226, 338  
Llinás R, 208, 225, 229, 238, 246  
Lo Y-J, 298, 307  
Loewy AD, 246  
Loftus WC, 80, 83, 94, 95, 109, 111  
Loh EW, 611, 622  
Lohmann C, 85, 109, 544, 555  
Lohr B, 528  
Löhrke S, 305, 554  
Lolait SJ, 278  
Lombard RE, 472, 488, 637  
Lomber SG, 245  
Long C, 247, 625  
Lonsbury-Martin BL, 575, 581  
López DE, 51, 84, 109, 178, 190, 197, 198  
López-Colomé AM, 251, 276  
López-Poveda EA, 180  
Lorente de Nó R, 8, 11, 59  
Lorenzi C, 225  
Louage D, 406  
Lowenstein DH, 624  
Lowy K, 639  
Lu S, 592, 602  
Lu SM, 268, 280, 602  
Lu Y, 300, 307, 316, 319, 340  
Lu Z, 465, 489  
Luan RH, 530  
Luddens H, 269, 276  
Ludewig K, 197  
Ludwig J, 292, 307, 310  
Luethke LE, 38, 59  
Luff NP, 616  
Luksch H, 505, 531, 532  
Lund JS, 49, 55, 61, 71, 75, 82, 107, 155, 156, 178, 277, 377, 407, 537, 553, 629, 630, 637  
Luneau CJ, 291, 307  
Luo L, 112, 118, 174, 548, 594, 600  
Lutz C, 557  
Lynch-Armour MA, 116, 128, 178, 196, 225, 580  
**Ma** CL, 114, 249, 253, 254, 255, 267, 276, 281, 294, 295, 296, 298, 299, 300, 308, 310, 567, 581, 583  
**Ma** X, 174, 181, 246, 456, 518, 531, 533, 534, 546, 555, 639  
Macdonald RL, 269, 276, 606, 616  
Madamba SG, 623  
Maeda T, 277  
Maffi CL, 116, 123, 124, 128  
Magistretti PJ, 113  
Magoun HW, 127, 636

- Mahan LC, 278  
 Maiwald D, 329, 340  
 Majchrowicz E, 605, 617, 622  
 Majorossy K, 205, 207, 226  
 Malenka RC, 257, 272  
 Malherbe P, 278  
 Malmierca MS, 23, 59, 75, 76, 77, 78,  
     79, 81, 83, 88, 97, 110, 155, 162, 163,  
     164, 172, 174, 175, 180, 182, 183,  
     184, 186, 188, 197, 198, 202, 203,  
     204, 205, 226, 236, 246, 251, 272,  
     276, 307, 325, 340, 372, 373, 453,  
     537, 555, 621, 629, 637, 638  
 Malone BJ, 279, 634, 637  
 Manabe T, 229, 334, 344, 534  
 Manis PB, 113, 290, 292, 300, 306, 308  
 Manley JA, 337, 341, 477, 480, 482, 489,  
     519, 531, 637  
 Manso ME, 620  
 Mantolán-Sarmiento B, 55, 108, 151,  
     179, 275, 620  
 Manunta Y, 224  
 Marchaterre MA, 484, 485  
 Marcinczyk MJ, 305, 619  
 Marescaux C, 605, 616, 620, 622, 624  
 Margoliash D, 47, 59, 216, 226, 411,  
     424, 474, 489  
 Margolis RH, 324, 340  
 Marianowski R, 249, 276, 297, 308, 550,  
     555, 611, 622  
 Marietta CA, 617  
 Markovitz NS, 21, 25, 59  
 Markram H, 108  
 Marquardt T, 403  
 Marrion NV, 310  
 Marsh RA, 231, 240, 242, 246, 503, 531,  
     627, 638  
 Marshall DH, 403  
 Marshall J, 111, 307, 308  
 Martin GF, 116, 120, 121, 122, 131, 137,  
     138, 154, 164, 166, 169, 181  
 Martin GK, 456, 575, 581  
 Martin IL, 269, 274  
 Martin KA, 111, 308  
 Martin RL, 107, 127, 320, 336, 340, 434,  
     435, 454, 555, 581, 600  
 Martin WF, 468, 489  
 Martinez-de-la-Torre M, 533  
 Martsis-McClintock A, 553, 554  
 Maruseva EA, 338  
 Marvit P, 467, 489  
 Marx SO, 290, 308  
 Mascia MP, 622, 625  
 Masino T, 531  
 Massaux A, 211, 213, 226  
 Masterton RB, 23, 30, 52, 55, 57, 59, 61,  
     107, 108, 110, 112, 127, 150, 151,  
     178, 179, 225, 244, 274, 340, 344,  
     371, 405, 407, 452, 453, 454, 456,  
     553, 637, 639  
 Maton B, 620  
 Matos-Collazo S, 108  
 Matsubara JA, 38, 59  
 Matsumura S, 423, 530  
 Matsuoka H, 3, 35, 36, 40, 61  
 Matthews PBC, 47, 59  
 Mattingley JB, 405  
 Maurois P, 615  
 Maxwell JK, 618  
 May BJ, 91, 111, 113, 127, 129, 273,  
     341, 381, 404, 408, 433, 434, 435,  
     436, 437, 438, 440, 441, 442, 443,  
     451, 452, 454, 455  
 Mayer F, 531  
 Mayer ML, 294, 308  
 Mazer J, 508, 519, 520, 532, 535  
 McAdams S, 406, 457  
 McAinsh K, 621  
 McAlpine D, ix, 262, 276, 380, 383, 384,  
     385, 390, 397, 403, 405, 407, 408,  
     550, 555, 581, 594, 600  
 McCabe PM, 229, 617  
 McCormack T, 304  
 McCormick DA, 208, 227, 246, 405, 461,  
     464, 469, 484, 489  
 McCown TJ, 603, 604, 619, 622  
 McKee MP, 461, 489  
 McDonald AJ, 107, 412, 420, 425  
 McEwen BS, 620  
 McFadden SL, 269, 277, 391, 407, 566,  
     581, 589, 600, 602  
 McGee JD, 343, 558, 583  
 McGeer EG, 114, 277  
 McGeer PL, 114, 277  
 McGeer R, 309  
 McGehee DS, 546, 555  
 McGregor P, 438, 454  
 McHaffie JG, 544, 555

- McKernan RM, 269, 273, 277  
McKibben JR, 460, 461, 465, 467, 484, 489  
McKowen A, 600  
McLaren JD, 620  
McMahon T, 624  
McMullen NT, 52, 224  
Meddis R, viii, ix, 361, 370, 371  
Meeks J, 275, 581  
Mehmert KK, 625  
Meininger V, 71, 74, 110, 312, 322, 325, 340, 543, 553, 555  
Meitzen J, 109, 423, 488  
Melcher JR, 214, 225  
Meldrum BS, 54, 308, 615, 616, 617, 618, 622, 625  
Meleca RJ, 599  
Mello CV, 530  
Mello LE, 619  
Melo LL, 301, 308  
Meloni EG, 197  
Mendelson JR, 39, 41, 59, 65  
Mensing AF, 464, 489  
Merchán MA, 19, 51, 59, 79, 81, 97, 107, 110, 111, 112, 122, 129, 130, 155, 159, 160, 162, 163, 172, 178, 180, 181, 205, 226, 228, 236, 246, 251, 272, 277, 278, 279, 288, 373, 527, 629, 632, 638, 639  
Mercik K, 622  
Merzenich MM, 13, 23, 27, 38, 39, 41, 43, 51, 57, 58, 60, 64, 88, 90, 107, 110, 112, 116, 129, 153, 178, 223, 315, 316, 318, 319, 320, 324, 327, 328, 330, 331, 332, 333, 334, 337, 340, 342, 371, 372, 425, 431, 435, 443, 451, 453, 489, 567, 568, 580  
Messing RW, 624  
Metherate R, 553  
Metzner W, 189, 192, 197, 464, 489, 491  
Meyer G, 179, 554  
Meyer R, 293, 308  
Mhatre MC, 611, 622  
Micheletti G, 622  
Michler SA, 570, 571, 581  
Middlebrooks JC, 33, 35, 38, 39, 41, 60, 204, 227, 438, 440, 454  
Mielcarz JP, 52  
Mihailoff GA, 195  
Mihic SJ, 611, 622  
Mikaelian DO, 586, 600  
Miki T, 195, 198  
Milbrandt JC, 251, 253, 267, 268, 269, 270, 273, 277, 278, 550, 555, 566, 569, 570, 580, 581, 592, 600, 602  
Millan MH, 54, 297, 308, 603, 605, 610, 611, 613, 618, 622  
Miller J, 332, 342, 349, 374, 455, 456  
Miller JM, 272, 456, 579, 582, 599  
Miller LM, 37, 39, 60, 338, 340  
Mills AW, 434, 454  
Mills DM, 554, 555  
Ming Z, 617  
Mirsky R, 56  
Misawa H, 352, 372, 604, 622  
Misgeld U, 255, 277  
Mishima K, 214, 227  
Mishkin M, 63, 424, 639  
Mishra PK, 621  
Mitani A, 38, 60, 233, 246  
Mittman DH, 116, 131, 216, 227, 315, 319, 340, 344, 352, 366, 372, 513, 532  
Miyoshi S, 198  
Mizuno N, 60, 65, 198, 226, 246  
Mizuno T, 128, 131  
Moffat AJM, 213, 224, 316, 330, 332, 337, 338, 349, 370  
Mogdans J, 519, 523, 532  
Mogharreban N, 616  
Mohler H, 269, 277  
Moiseff A, 518, 519, 532, 534  
Møller AR, 263, 268, 309, 350, 352, 353, 362, 372, 373, 415, 424, 572  
Molnár C, viii, ix  
Molnar CE, 13, 58  
Molnar LR, 610, 622  
Molowny RL, 109, 340, 555  
Momiyama A, 310  
Monyer H, 281  
Mooney D, 210, 225, 227  
Moore BCJ, 324, 325, 340, 369, 372  
Moore CN, 172, 180, 435, 454  
Moore DR, 23, 50, 109, 116, 119, 120, 128, 138, 152, 169, 180, 196, 197, 253, 254, 255, 267, 277, 294, 297, 308, 543, 547, 550, 552, 555, 556, 557, 558, 578, 581, 582, 594, 595, 600

- Moore JK, 15, 60, 197, 198, 250, 251, 277, 443, 454, 627, 629
- Moore RY, 15, 60, 157, 180, 250, 251, 277
- Moore SD, 638
- Moreau J, 308, 622
- Morel A, 32, 33, 34, 36, 39, 57, 60, 215, 217, 219, 225, 227, 228
- Morenkov E, 625
- Moreno H, 304, 309, 310
- Morest DK, 4, 7, 11, 23, 26, 31, 32, 33, 34, 35, 40, 51, 52, 56, 58, 61, 64, 68, 71, 75, 82, 83, 84, 86, 88, 108, 110, 112, 116, 120, 122, 128, 129, 138, 152, 153, 197, 201, 204, 205, 207, 227, 247, 278, 312, 320, 325, 340, 341, 456, 537, 554, 556, 629, 632, 633, 637, 638, 639, 640
- Morgane PJ, 108
- Mori K, 131
- Mori M, 306, 620
- Moriiizumi T, 556
- Morinobu S, 311
- Morley BJ, 27, 65, 301, 308, 546, 556
- Morris SJ, 275
- Morrow AL, 617, 620
- Moss SJ, 621
- Mossop JE, 268, 277, 550, 555, 556, 569, 582, 594, 600
- Mount RJ, 554, 580
- Mountcastle VB, 25, 29, 60, 426, 453
- Moushegian G, 403
- Mower G, 191, 197
- Mozzrymas JW, 606, 622
- Mugnaini E, 5, 7, 9, 11, 15, 17, 18, 19, 27, 50, 51, 67, 107, 129, 130, 141, 150, 174, 181, 199, 250, 251, 254, 262, 272, 277, 280, 633, 636, 639
- Mulders WHAM, 185, 197
- Müller BA, 338
- Müller CM, 501, 532
- Müller M, 544, 556
- Müller-Preuss P, 350, 372
- Mulligan KA, 41, 60
- Musicant AD, 62, 432, 435, 450, 454
- Nadal MS, 304, 309
- Nagai T, 250, 251, 277
- Nagarajan SS, 52, 460, 489
- Nagase S, 272, 569, 573, 579, 582, 599
- Nakagawa H, 174, 180, 309, 557
- Nakahira M, 131
- Nakamura K, 132, 142, 147, 151
- Nakamura M, 620
- Nakamura Y, 116, 119, 128, 151, 328, 340
- Nandy D, 433, 454
- Narayan SS, 423
- Narins PM, 469, 488
- Naritoku DK, 273, 603, 604, 609, 618, 619
- Nauta WJH, 115, 128
- Navetta MS, 623
- Neff WD, 40, 54, 55, 60, 66, 180, 247, 403, 408, 435, 443, 451, 454, 597, 607, 628, 638
- Nehlig A, 603, 620, 622
- Neises GR, 51
- Nelken I, 68, 325, 340, 431, 455, 456
- Nellhaus G, 603, 620
- Nelson DK, 616
- Nelson J, 553
- Nelson JE, 127
- Nelson JS, 460, 489
- Nelson LR, 130
- Nelson PG, 27, 60, 113, 227, 349, 355, 361, 366, 372, 627, 628, 638
- Nelson RJ, 57
- Netzer R, 307
- Neubauer H, 329, 330, 339
- Neuert V, 355, 372, 634, 638
- Neumann PE, 604, 623
- Neuweiler G, 493, 494, 532, 533
- Newman EB, 65, 409, 424, 457
- Newsome WT, 426, 427, 455
- Niblock MM, 539, 556
- Nicholson C, 490, 491
- Nicoll RA, 306
- Nieder B, 532, 535
- Nienhuys TGW, 324, 340
- Nieuwenhuys R, 459, 482, 490, 491
- Niimi K, 3, 36, 40, 58, 61, 84, 109, 157, 166, 172, 179, 192, 197, 203, 226
- Niki H, 604, 605, 620, 621, 624
- Nishimori T, 620
- Nishimura T, 620
- Nishiyama N, 547, 556
- Nishizaki T, 293, 306



- Nizamov SV, 353, 368  
 Noble D, 57  
 Nobre AC, 47, 61  
 Noda A, 620  
 Noda T, 131  
 Nodal FR, 109, 190, 197, 224, 272  
 Noh MD, 56, 114, 141, 152, 180, 195,  
 197, 207, 277, 306, 520, 551, 614, 638  
 Nomura S, 60, 246  
 Nordeen KW, 116, 117, 119, 128, 137,  
 152, 547, 556  
 Nordlund B, 377, 407  
 Norris BE, 52, 56, 63, 108, 151, 339,  
 604, 625, 639  
 Norris CH, 625  
 Northcutt RG, 471, 490, 492  
 Novick A, 533  
 Nuding SC, 423  
 Nudo RJ, 55, 73, 88, 110, 452  
 Nuetzel JM, 391, 407  
  
**O**  
 Oakson G, 229  
 O'Carrol AM, 278  
 O'Connor KN, 325, 341  
 Oeckinghaus H, 337, 341  
 Oertel D, 13, 18, 63, 67, 107, 132, 138,  
 140, 141, 146, 150, 152, 153, 154,  
 195, 196, 199, 273, 305, 306, 308,  
 310, 458, 527, 636  
 Oertel WH, 250, 251, 277, 624  
 Oesterreich RE, 435, 443, 456  
 Oh S, 276, 605, 623  
 Ohl FW, 43, 61  
 Ohlemiller KK, 216, 227, 411, 420, 423,  
 424, 530  
 Ohlrogge M, 186, 197  
 Ohshima T, 85, 110, 112  
 Ojima H, 38, 56, 61  
 Okada Y, 306, 620  
 Okano H, 131  
 Okoda Y, 581  
 Okoyama S, 539, 556  
 Olazábal UE, 192, 197, 627, 638  
 Oliva C, 307  
 Olive MF, 625  
 Oliveira JA, 619  
 Oliver DL, vii, ix, 9, 13, 15, 17, 19, 20,  
 23, 25, 26, 27, 28, 30, 31, 33, 43, 51,  
 60, 61, 65, 68, 70, 71, 74, 75, 78, 79,  
 80, 81, 82, 83, 84, 85, 86, 87, 88, 92,  
 93, 96, 98, 99, 100, 101, 102, 109,  
 110, 111, 112, 113, 114, 116, 117,  
 120, 121, 122, 123, 124, 128, 129,  
 132, 133, 134, 135, 152, 153, 157,  
 164, 169, 170, 171, 180, 188, 197,  
 203, 204, 205, 226, 227, 228, 230,  
 247, 250, 277, 279, 281, 283, 284,  
 286, 287, 288, 290, 291, 292, 293,  
 307, 309, 310, 312, 318, 320, 325,  
 332, 340, 341, 343, 371, 390, 402,  
 407, 537, 539, 540, 556, 557, 583,  
 629, 630, 633, 638, 640  
 Olla BL, 487  
 Ollo C, 137, 152  
 Olsen JF, 204, 212, 224, 227, 245, 246,  
 344, 513, 532, 639  
 Olsen RW, 269, 276  
 Olson CR, 41, 52, 490, 491  
 Olszewski J, 71, 111  
 Onaya T, 110  
 O'Neill WE, 116, 128, 178, 196, 204,  
 212, 216, 217, 218, 220, 224, 225,  
 227, 228, 229, 280, 356, 362, 363,  
 370, 372, 375, 425, 474, 490, 513,  
 515, 528, 531, 532, 534, 536, 583,  
 584, 635, 638  
 Onodera K, 606, 623  
 Orman SS, 62  
 Osen KK, 5, 6, 7, 15, 61, 66, 71, 75, 82,  
 107, 109, 112, 115, 117, 120, 121,  
 122, 129, 155, 157, 178, 278, 340,  
 373, 537, 553, 555, 629, 630, 637  
 Ostad M, 64  
 Ostapoff E-M, 64, 111, 112, 116, 129,  
 153, 187, 197, 227, 251, 277, 278,  
 404, 409, 456, 639  
 Ostwald J, 4, 55, 196, 363, 370, 529, 637  
 Otani K, 116, 130, 181  
 Ottersen OP, 61, 63, 67, 112, 113, 129,  
 278, 373  
 Owren MJ, 424  
 Oxenham AJ, 374  
 Ozaita A, 304, 309  
  
**Pages N**, 615  
 Palacios JM, 301, 304  
 Palay SL, 62, 195  
 Palkovits M, 272

- Pallas SL, 205, 228  
 Palm G, 369  
 Palmer AR, ix, 262, 276, 328, 332, 333, 338, 341, 342, 345, 350, 351, 352, 353, 354, 356, 370, 373, 380, 383, 390, 397, 398, 403, 405, 407, 408, 423, 430, 455  
 Palmer LA, 66  
 Paloff AM, 78, 111  
 Palombi PS, 260, 262, 269, 273, 277, 278, 297, 308, 328, 330, 332, 341, 356, 373, 564, 572, 580, 582, 606, 623  
 Pandossio JE, 193, 198  
 Pandya PK, 3, 58, 65  
 Papadeas ST, 620  
 Papez JW, 15, 62, 626, 638  
 Paré D, 229, 238, 246  
 Parent A, 65  
 Parham K, 280, 281, 399, 404, 407, 447, 451, 455, 586, 600, 602  
 Park TJ, 226, 250, 262, 275, 278, 307, 308, 309, 341, 407, 511, 530, 532, 582  
 Park VN, 405  
 Parks TN, 249, 278, 387, 408, 426, 444, 456, 505, 527  
 Pasanen E, 391, 407  
 Paterson JA, 594, 601  
 Paterson M, 400  
 Patterson RD, 341, 372, 373, 638  
 Patterson TA, 304  
 Patuzzi R, 312, 341  
 Paxinos G, 66, 71, 82, 85, 111  
 Payne BR, 245  
 Pearl D, 59  
 Pedemonte M, 349, 373, 620  
 Peña JL, 519, 521, 523, 532  
 Penix LP, 621  
 Penny JE, 604, 623  
 Pepitoni S, 273  
 Pérez-González H, 55, 108, 151, 179, 275, 620  
 Perkel DJ, 306  
 Perney TM, 102, 111, 291, 307, 308  
 Peroutka SJ, 301, 308  
 Persohn E, 269, 278  
 Peruzzi D, 99, 100, 101, 102, 104, 111, 169, 180, 205, 206, 208, 228, 283, 287, 288, 289, 291, 309, 332, 341  
 Peters A, 7, 52, 62  
 Peterson BA, 62, 638  
 Peterson GE, 424  
 Petralia RS, 113, 280  
 Pettigrew JD, 39, 59  
 Phillips DP, 38, 44, 59, 62, 228, 406  
 Phillips SC, 23, 29, 50, 157, 178, 223, 244  
 Philpott P, 50, 116, 127  
 Piao Y-S, 491  
 Picchioni AL, 621  
 Pickles JO, 324, 327, 341  
 Piechotta K, 546, 557  
 Pieribone VA, 199  
 Pierre A, 51, 53, 59, 63, 200, 202, 203, 204, 224, 225, 227, 228, 345  
 Pierson M, 603, 604, 621  
 Pierson MG, 257, 278, 297, 309, 604, 623, 624  
 Pike B, 51  
 Pinek B, 107  
 Pinheiro AD, 351, 364, 373, 473, 490, 515, 532  
 Pippin J, 580  
 Plaza I, 59, 110, 277  
 Plomp R, 324, 341, 347, 373, 374, 375  
 Plunkett K, 47, 61  
 Poirier P, 215, 217, 225, 228, 229, 452, 456  
 Pol D, 110, 340  
 Pollak GD, 19, 21, 25, 59, 62, 64, 68, 109, 114, 116, 123, 124, 129, 130, 132, 141, 142, 146, 147, 152, 154, 164, 180, 230, 247, 250, 262, 263, 266, 267, 268, 273, 275, 278, 281, 297, 301, 304, 306, 307, 308, 309, 311, 328, 341, 345, 362, 364, 371, 376, 396, 407, 423, 459, 488, 490, 497, 508, 511, 516, 517, 527, 529, 530, 533, 535, 536, 558, 567, 581, 582, 583, 606, 610, 616, 627, 638, 640  
 Pollard H, 308, 622  
 Pongs O, 307  
 Pons T, 424  
 Pontes C, 42, 62  
 Poon PWF, 341, 373, 455  
 Popelář J, 229, 233, 247, 344, 375, 425, 456, 571, 572, 575, 582, 583, 629, 639

- Popper AN, 3, 52, 60, 62, 63, 64, 66, 67, 108, 110, 129, 150, 152, 153, 154, 195, 196, 197, 199, 224, 229, 230, 247, 273, 339, 341, 375, 376, 405, 458, 459, 461, 465, 467, 483, 486, 487, 488, 489, 490, 491, 527, 528, 529, 636, 637, 638, 640
- Populin LC, 399, 408, 435, 436, 446, 455
- Poremba A, 71, 85, 111
- Portfors CV, 212, 216, 219, 220, 228, 315, 319, 341, 359, 373, 412, 413, 415, 416, 420, 421, 424, 512, 513, 532, 533
- Potashner SJ, 61, 113, 176, 178, 197, 232, 245, 255, 272, 278, 279, 558, 583, 611, 619, 624
- Pough FH, 460, 464, 490
- Poulter MO, 269, 278
- Pountney D, 304
- Pow DV, 205, 228
- Powell TPS, 35, 43, 54, 57
- Powers N, 279, 344, 582, 583, 601
- Pozza MF, 310
- Pralong D, 436, 455
- Prasad V, 51
- Pratt CC, 433, 455
- Prechtl JC, 461, 490, 491
- Pressnitzer D, 355, 372, 373, 406, 457, 638
- Preuss TM, 56, 350, 372
- Price JL, 242, 246
- Price RO, 52, 224
- Priebe RP, 600
- Prieto JJ, 3, 13, 33, 34, 38, 39, 41, 42, 43, 51, 62, 195, 223, 232, 244, 246, 247, 633, 638
- Priets CA, 196
- Pritz MB, 23, 62, 459, 477, 478, 479, 483, 490
- Probst A, 4, 107, 108, 113, 141, 262, 273, 280, 289
- Proctor L, 505, 533
- Przybyszewski AW, 39, 62, 237, 246
- Práček LJ, 624
- Puchades A, 179
- Puel JL, 107
- Puelles L, 486, 498, 499, 501, 502, 503, 533
- Puil E, 229
- Pujol R, 553
- Pysh JJ, 542, 557
- Qiu A, 325, 327, 341**
- Quirion R, 309
- Rabow LE, 269, 278**
- Radcliffe KA, 546, 557
- Radil-Weiss T, 443, 456
- Radionova EA, 338
- Radtke-Schuller S, 31, 65
- Raisinghani M, 604, 605, 613, 623
- Rajan R, 56, 339, 452, 568, 572, 575, 578, 581, 582
- Rakic P, 33, 62, 63, 639
- Rall W, 373
- Ralls K, 586, 601
- Ramachandran R, 91, 111, 113, 125, 127, 129, 273, 315, 316, 319, 324, 328, 330, 341, 381, 404, 408, 438, 451, 455
- Ramón y Cajal S, 70, 71, 111, 155, 157, 180, 431, 455, 626, 627, 633, 639
- Randall ME, 54, 274, 305, 618, 619, 625
- Ranson SW, 127, 636
- Rao SC, 307
- Raphael LJ, 423
- Rasch RA, 444, 456
- Ratnikova GI, 338
- Ratray M, 615, 625
- Rauschecker JP, 39, 58, 62, 63, 216, 228, 411, 412, 420, 424, 639
- Ravizza R, 407
- Rayleigh L, 427, 428, 455
- Raza A, 251, 269, 273, 278, 580
- Read HL, 38, 41, 60, 63, 324, 338, 340, 341
- Reale RA, 35, 38, 39, 40, 42, 52, 57, 63, 232, 233, 245, 246, 399, 408, 430, 450, 455
- Recanzone GH, 41, 62
- Redgrave P, 246
- Redies H, 218, 219, 228
- Reeks KA, 67
- Rees A, 110, 180, 197, 226, 246, 263, 276, 278, 304, 307, 332, 336, 340, 342, 349, 350, 351, 352, 353, 354, 355, 356, 361, 362, 370, 371, 372, 373, 375, 403, 407, 414, 423, 424, 453, 455, 621
- Reese EP, 450, 604, 617, 619

- Reetz G, 169, 170, 172, 174, 180, 253, 254, 278, 283, 287, 288, 289, 309, 325, 332, 342
- Regenold W, 301, 309
- Reid HM, 605, 623
- Reid MD, 23, 27, 59, 88, 110, 319, 320, 340
- Reiffenstein RJ, 623
- Reigel CE, 604, 605, 623
- Reimer K, 320, 342, 352, 373, 516, 533
- Reis DJ, 59, 226
- Reis FF, 62
- Relkin EM, 329, 339
- Rendall D, 413, 424
- Resler C, 281, 311, 345, 536, 583
- Réthelyi M, 205, 226
- Reynolds A, 23, 50, 543, 552
- Rhode WS, 5, 9, 11, 31, 63, 121, 125, 129, 130, 319, 322, 328, 342
- Riaz A, 603, 605, 610, 611, 613, 618, 619, 623
- Ribak CE, 78, 112, 250, 278, 444, 455, 603, 606, 611, 620, 623, 624
- Ribeiro S, 530
- Ribera AB, 290, 304
- Rice JJ, 52, 63, 84, 106, 112, 178, 180, 223, 224, 242, 246, 323, 336, 432, 433, 436, 455, 458, 496, 629, 632
- Richards JG, 278, 291, 292, 309
- Richardson FC, 291, 292, 309
- Richter EA, 17, 63, 629, 639
- Rico B, 127, 204, 541
- Rinvik E, 34, 63, 245
- Rinzel J, ix, 309
- Riquelme R, 80, 112, 121, 129, 254, 278, 361, 373
- Ritsma RJ, 374
- Rittenhouse AR, 293, 308
- RoBards MJ, 59, 84, 112, 627, 632, 639
- Robert D, 47, 63, 611
- Robert JM, 345
- Roberto M, 611, 623
- Roberts RC, 78, 112, 250, 278, 606, 624
- Robertson D, 185, 197, 281
- Robles C, 533
- Robles L, 424
- Rockel AJ, 23, 26, 29, 33, 63, 71, 84, 112, 166, 180, 207, 226, 373, 537, 557, 633, 639
- Rockwood JM, 113, 279
- Rodman PS, 424
- Rodrigues-Dagaëff C, 33, 36, 64, 204, 212, 213, 218, 228
- Rogers JH, 553
- Romand R, 151, 196, 224, 249, 278, 308, 319, 320, 322, 328, 337, 338, 342, 543, 544, 557, 622, 636
- Romanski LM, 38, 39, 63, 231, 246, 635, 639
- Romey G, 306
- Romo R, 53, 126, 136, 248, 270, 281, 297, 364, 469, 567
- Rosa MG, 269, 582
- Rose GJ, 350, 352, 373, 374, 469, 473, 474, 476, 482, 484, 485, 486, 487, 490
- Rose JE, vi, 5, 8, 26, 34, 39, 40, 42, 64, 316, 322, 342, 349, 371, 380, 408, 428, 430, 455, 627, 639
- Rose L, 54, 245
- Rosemann C, 529
- Rosen S, 346, 347, 374, 413, 424
- Rosenberger MH, 290, 309, 497, 498, 533
- Rosenquist AC, 66
- Rosenzweig MR, 457
- Ross KC, 444, 455
- Ross LS, 64, 67, 116, 119, 129, 130, 147, 152, 156, 164, 172, 180, 535
- Roth GL, 4, 51, 59, 64, 91, 112, 116, 119, 123, 124, 129, 146, 153, 178, 223, 226, 263, 319, 322, 342
- Rotman Y, 340
- Rouiller EM, 33, 35, 39, 41, 51, 53, 54, 60, 64, 178, 228, 345
- Rubel EW, 309, 387, 408, 544, 555, 557
- Rubinson K, 477, 479, 485
- Rübsamen R, 3, 52, 124, 127, 151, 198, 319, 336, 508, 513, 533
- Rudolph U, 277
- Rudy B, 304, 309, 310
- Rudy JW, 554
- Rudy R, 309
- Ruetzler CA, 621
- Ruggiero MA, 412, 424
- Ruggiero DA, 58, 226
- Ruiz GT, 179, 623
- Ruppersberg JP, 112, 557
- Russek SJ, 278, 611, 624
- Russell FA, 547, 557, 558
- Russell IJ, 531

- Russo E, 617  
 Rustay NR, 615  
 Rustioni A, 53, 224  
 Ryals BM, 557  
 Ryan A, 112, 332, 342, 349, 374, 444, 455, 456, 544, 557, 582, 600, 602  
 Rybalko N, 268, 279  
 Ryugo DK, 5, 21, 38, 43, 54, 64, 67, 113, 116, 120, 122, 129, 134, 137, 138, 144, 151, 154, 166, 181, 186, 197, 199, 245
- Sachs MB, 318, 327, 329, 342  
 Saganich M, 304, 309  
 Sah P, 306, 616  
 Saint Marie RL, ix, 7, 18, 19, 22, 23, 25, 61, 64, 68, 80, 89, 109, 111, 112, 114, 135, 136, 137, 138, 140, 153, 162, 169, 174, 181, 184, 186, 188, 189, 191, 198, 205, 207, 228, 230, 231, 246, 247, 249, 250, 251, 277, 278, 341, 407, 431, 456, 600, 611, 624, 633, 638, 639, 640  
 Saito N, 152, 366, 374  
 Saitoh I, 366, 374  
 Sakai H, 418, 425  
 Sakamoto T, 604, 605, 624  
 Sakitt B, 397, 408  
 Sakmann B, 66  
 Saldaña E, 59, 67, 81, 109, 110, 112, 122, 128, 129, 130, 137, 153, 155, 157, 159, 160, 161, 162, 165, 167, 170, 172, 176, 178, 180, 181, 188, 198, 199, 205, 228, 236, 246, 251, 277, 278, 279, 373, 610, 616, 632, 639  
 Saleda R, 276  
 Sally SL, 83, 90, 108, 155, 159, 161, 162, 163, 166, 172, 186, 250, 279, 300, 396, 408, 498, 502, 508, 560  
 Salvi R, 272, 277, 279, 280, 344, 555, 579, 581, 582, 583, 599, 600, 601  
 Samson FR, 217, 225, 228, 229, 443, 452, 456  
 Sánchez del Campo F, 179  
 Sanchez-Perez AM, 624  
 Sanders DJ, 304, 336  
 Sandkühler J, 57, 306  
 Sandner G, 615  
 Sanes DH, 18, 64, 180, 262, 277, 279, 300, 307, 308, 309, 544, 550, 557, 558  
 Saper CB, 57  
 Sarbaz A, 350, 356, 361, 373  
 Sasa M, 620  
 Sato K, 270, 298, 309, 426, 545, 557, 582, 585, 590, 604, 613  
 Saunders SS, 51, 279, 582  
 Sawamoto K, 131  
 Sayers BM, 348, 369, 406  
 Scates KW, 185, 198  
 Schachner M, 526  
 Schafer M, 508, 533  
 Schafer TH, 324, 342  
 Scheich H, 43, 61, 371, 508, 526, 529, 533  
 Schellart NAM, 465, 487  
 Schirra C, 306  
 Schlegel P, 491  
 Schmid LM, 582  
 Schmid S, 112, 557  
 Schmiedt RA, 328, 343  
 Schmieszek E, 497, 526  
 Schneider GE, 45, 56, 64, 468, 491  
 Schneiderman N, 229  
 Schoepfer R, 295, 307  
 Schofield BR, 15, 16, 64, 116, 119, 120, 130, 132, 135, 136, 137, 140, 143, 148, 150, 153, 154, 184, 185, 186, 187, 188, 198  
 Schön Ybarra MA, 413, 424  
 Schouten JF, 348, 360, 374  
 Schreibmayer W, 306  
 Schreiner CE, 25, 30, 37, 38, 39, 40, 41, 44, 54, 57, 58, 60, 63, 64, 65, 88, 90, 112, 176, 181, 233, 246, 319, 320, 323, 324, 325, 327, 328, 330, 333, 336, 338, 340, 341, 343, 350, 351, 352, 353, 356, 359, 360, 361, 371, 372, 374, 412, 421, 422, 423, 424, 425, 473, 474, 485, 488, 489, 622, 634, 637  
 Schroeder MR, 352, 374, 388, 408  
 Schubert ED, 378, 409  
 Schuck D, 116, 123, 126, 147, 150, 328, 336  
 Schuknecht HF, 324, 343, 586, 601  
 Schuller G, 31, 65, 189, 191, 192, 196, 198, 350, 353, 374, 414, 424, 503, 516, 529, 533

- Schult W, 50, 336  
 Schulze H, 371, 372  
 Schwartz IR, 15, 19, 65, 132, 135, 137, 138, 152, 154, 506  
 Schwartz RD, 301, 309  
 Schwartz-Bloom RD, 274, 305, 528  
 Schwarz DWF, 107, 229, 375, 528, 533  
 Schwarz IE, 107, 528, 533  
 Schweitzer L, 544, 557  
 Schweizer H, 130, 164, 169, 181, 198, 529  
 Scott LL, 199  
 Seeburg PH, 281, 555  
 Séguéla P, 65  
 Seidl AH, 388, 408  
 Sekitani M, 546, 557  
 Seltzer B, 3, 65  
 Semal C, 348, 374  
 Semba R, 112  
 Semple MN, ix, 62, 90, 91, 104, 112, 113, 124, 130, 147, 153, 213, 229, 262, 265, 276, 279, 319, 332, 343, 351, 352, 353, 355, 371, 390, 391, 395, 408, 473, 488, 555, 594, 600, 634, 637  
 Sen K, 344  
 Senatorov V, 225, 227, 503, 533  
 Senut MC, 70, 112  
 Serikawa T, 620  
 Serôdio P, 292, 309  
 Servièrre J, 3, 65, 320, 340, 343, 344  
 Seto-Ohshima A, 85, 112  
 Shackleton TM, viii, ix, 276, 380, 385, 387, 388, 390, 407, 408  
 Shaddock-Palombi P, 564, 582  
 Shaia WT, 113  
 Shamma SA, 325, 339, 343, 344, 388, 408  
 Shannon RV, 346, 374, 460, 491  
 Shapiro M, 348, 369  
 Shapiro SM, 113  
 Sharp FR, 602  
 Shatz CJ, 59  
 Shaw EAG, 310, 377, 378, 408, 490  
 Shcherbakov VI, 435, 452  
 Shchipakina T, 625  
 Sheft S, 348, 376  
 Shen NM, 159, 178, 408  
 Sheng HZ, 73, 89, 113  
 Shepherd RK, 547, 554, 556  
 Sherman SM, 37, 65, 208, 211, 229  
 Sherr EH, 622  
 Shewmaker CA, 342  
 Shimokouchi M, 38, 60, 246  
 Shimozawa T, 491  
 Shinn-Cunningham BG, 400, 408  
 Shinonaga Y, 35, 65, 193, 198, 231, 246  
 Shiosaka S, 557  
 Shipley C, 35, 65  
 Shiraishi S, 103, 113, 253, 279, 309, 557  
 Shiraishi Y, 113, 279  
 Shneiderman A, 23, 30, 61, 64, 65, 70, 79, 80, 82, 85, 111, 112, 113, 123, 129, 130, 141, 147, 152, 153, 157, 179, 250, 251, 254, 279, 537, 539, 556, 557  
 Shofner WP, 345, 348, 374  
 Shoham D, 56  
 Shore SE, 186, 187, 198  
 Siebert WM, 369, 426, 456  
 Sieghart W, 269, 279, 606, 617, 624  
 Siggins GR, 623  
 Sikela J, 616  
 Silverman MS, 550, 558  
 Simada Z, 260, 280  
 Simler S, 604, 620, 624  
 Simm G, 64, 228  
 Simmons AM, 483, 491  
 Simmons JA, 370, 491, 494, 495, 528, 533, 536  
 Simon JZ, 167, 339, 583  
 Simson PE, 611, 624  
 Sinclair D, 13, 65  
 Sindberg RM, 40, 65  
 Sinex DG, 52, 224, 327, 343, 411, 412, 414, 416, 418, 420, 422, 423, 424, 425, 548, 558, 571, 572, 573, 575, 576, 578, 582, 583  
 Singer JH, 301, 309, 464, 484, 489  
 Sirianni JG, 371  
 Sirinathsinghji DJS, 281  
 Sirosh J, 425  
 Sismanis A, 113  
 Sisneros J, 465, 484, 491  
 Sivaramakrishnan S, 86, 100, 101, 102, 111, 114, 180, 281, 283, 284, 286, 287, 288, 290, 291, 292, 293, 309, 310, 332, 341, 343, 583

- Skaliora I, 107, 245  
Skottun BC, 385, 387, 390, 408  
Skradski SL, 604, 624  
Small AM, 340  
Smith JS, 307  
Smith KL, 278, 309  
Smith PH, 63, 106, 111, 113, 130, 143,  
153, 181, 223, 228, 245, 279, 309,  
342, 343, 408, 453, 456, 624, 639  
Smith S, 581  
Smith Y, 33, 65  
Smith ZM, 374  
Smolders JWT, 325, 343  
Smotherman M, 464, 491  
Snead CR, 51, 275  
Snell KB, 560, 566, 580, 582  
Snyder RL, 58, 178, 343, 558, 582, 583  
Snyder SH, 306  
Snyder-Keller AM, 603, 624  
Solecki JM, 274  
Sommer TJ, 275, 581  
Song WJ, 290, 309  
Sousa-Pinto A, 3, 62, 65  
Spangler KM, 8, 18, 27, 66, 123, 128,  
143, 153, 539, 554  
Sparks DL, 426, 444, 456  
Spencer RF, 85, 113, 198  
Spezio ML, 533, 534  
Spirou GA, 22, 66, 91, 113, 132, 152,  
153, 276, 319, 343, 431, 455, 456,  
458, 621  
Spitzer MN, 104, 113, 262, 279, 390,  
391, 408, 534  
Spongr VP, 581, 600  
Spruston N, 66  
Sripathy K, 533  
Staatz-Benson C, 272  
Stables JP, 615  
Stanford TR, 91, 106, 107, 109, 113,  
152, 307, 371, 384, 402, 404, 430,  
453, 456  
Stanforth DA, 64, 112, 228  
Stanley GB, 66  
Stansfeld C, 307  
Stark JM, 106, 223  
Steenbergen JL, 625  
Steffan R, 310  
Stein BE, 198, 555  
Stein RB, 308  
Steinschneider M, 56  
Sterbing SJ, 380, 383, 401, 404, 409  
Sterchi D, 197  
Steriade M, 208, 210, 229, 246  
Stevens B, 107  
Stevens CF, 304  
Stevens HE, 409  
Stevens SS, 345  
Stiebler I, 25, 27, 30, 66, 319, 320, 322,  
324, 330, 334, 343, 359, 374  
Stillman RD, 385, 409  
Storm-Mathisen J, 61, 63, 67  
Stornetta R, 226  
Stotler WA, 115, 130  
Strack AM, 624  
Straka H, 465  
Straschill M, 444, 456  
Stratton S, 604, 625  
Straughan IR, 469, 491  
Streit P, 64  
Stritzel ME, 477, 483, 490  
Strominger AI, 130  
Strominger NL, 130, 456  
Strutz J, 370  
Stryker MP, 59  
Stuart G, 9, 66  
Stühmer W, 310  
Suga N, 7, 31, 39, 66, 68, 174, 181, 204,  
212, 215, 216, 220, 224, 227, 229,  
230, 236, 240, 245, 246, 247, 315,  
316, 332, 334, 344, 345, 352, 356,  
359, 362, 363, 364, 366, 370, 372,  
374, 375, 376, 420, 423, 424, 425,  
449, 451, 456, 458, 464, 474, 490,  
491, 511, 513, 515, 516, 518, 528,  
529, 530, 531, 532, 533, 534, 536,  
546, 554, 555, 572, 583, 628, 632,  
634, 639  
Sugimoto T, 180  
Sugita S, 116, 130, 181  
Sullivan WE, 406, 515, 534  
Sumi T, 594, 637  
Summerfield AQ, 370, 403, 423  
Sun LM, 624  
Sun W, 195  
Sun X, 236, 247, 275, 341, 621  
Sun XD, 57, 179, 199, 245, 339, 453,  
637  
Sun X-M, 600  
Suneja SK, 249, 250, 251, 268, 279, 550,  
558, 570, 583, 611, 624

- Sur M, 57, 228  
 Surmeier DJ, 304  
 Sussman HM, 420, 425  
 Šuta D, 412, 413, 416, 421, 425  
 Sutherland DP, 436, 440, 456  
 Sutter ML, 38, 41, 59, 65, 330, 341, 344  
 Suzuki H, 131, 465, 491  
 Svirskis G, 290, 309  
 Swann JW, 278, 309, 623  
 Swann S, 553  
 Swanson LW, 49, 57  
 Swanson R, 308  
 Swerdlow NR, 191, 198  
 Syka J, 51, 52, 61, 106, 109, 116, 119,  
     127, 151, 153, 164, 166, 169, 178,  
     212, 218, 223, 226, 229, 233, 247,  
     268, 279, 316, 319, 322, 323, 330,  
     332, 336, 340, 344, 349, 371, 373,  
     375, 423, 425, 442, 444, 456, 530,  
     571, 575, 582, 583, 629, 636, 638, 639  
 Symmes D, 411, 425  
 Szczepaniak WS, 268, 279, 300, 309,  
     572, 583  
 Szeligo FV, 53  
  
 Tabuchi K, 131  
 Tachibana M, 250, 251, 279  
 Tacke U, 76, 623  
 Takada M, 65, 198, 246  
 Takahashi TT, 406, 453, 501, 505, 521,  
     523, 528, 530, 533, 534, 639  
 Takeda R, 620  
 Takeuchi Y, 189, 190, 195, 198  
 Talebi A, 616  
 Tallent MK, 623  
 Talley EM, 309  
 Tan A, 622  
 Tanaka C, 152  
 Tang Y-Z, 438, 480, 491  
 Tapia R, 276  
 Tardif E, 43, 66, 85, 113  
 Taylor DA, 53, 620, 622  
 Tecott LH, 603, 624  
 Tempel BL, 305  
 ten Donkelaar HJ, 469, 477, 479, 480,  
     490, 491  
 Tennigkeit F, 208, 210, 211, 213, 229  
 Tepper JM, 528  
 Terhardt E, 353, 375  
 Terlau H, 292, 310  
  
 Theunissen FE, 325, 344, 494, 534  
 Thiele A, 188, 198  
 Thomas EC, 58, 67, 154, 181, 337, 339,  
     343, 406, 599  
 Thompson AM, 4, 66, 132, 135, 148,  
     154, 182, 183, 184, 185, 186, 192,  
     196, 198, 199, 275, 306, 310, 529, 581  
 Thompson GC, 52, 59, 107, 116, 127,  
     140, 150, 151, 154, 178, 182, 183,  
     184, 185, 199, 244, 250, 251, 279,  
     310, 436, 456  
 Thompson PO, 342  
 Thompson R, 32, 40, 66  
 Thornton S, 352, 375  
 Thoss VS, 546, 558  
 Thurlow WR, 627, 639  
 Tian B, 39, 62, 63, 228, 411, 412, 424,  
     639  
 Ticku MK, 611, 622  
 Tierney TS, 547, 558  
 Tkatch T, 304  
 Tobias JV, 342, 378, 382, 404, 409  
 Todd AJ, 639  
 Todd MA, 454  
 Toffano G, 273  
 Tohyama M, 557  
 Tokunaga A, 116, 119, 130, 164, 166,  
     169, 181  
 Tokuno H, 116, 128  
 Tolbert LP, 428, 456  
 Tollin DJ, 396, 399, 400, 409, 430, 436,  
     446, 449, 457  
 Tomaz C, 615  
 Tomlinson RWW, 348, 368, 375  
 Türk I, 41  
 Toros-Morel A, 228  
 Torterolo P, 373  
 Trahiotis C, 391, 393, 397, 399, 402,  
     403, 404, 409, 451  
 Tran Ba Huy P, 308, 622  
 Tran L, 51, 106, 223, 336, 343, 423, 636  
 Trang TT, 584  
 Trause AS, 452  
 Travis MA, 618  
 Treurniet WC, 351, 375  
 Troncoso AC, 195  
 Trujillo C-M, 486  
 Trune DR, 547, 558, 594, 601  
 Trussell L, 107, 140, 154, 273  
 Tsai G, 611, 624



- Tsien RW, 57  
 Tsuchitani C, 8, 15, 18, 66, 396, 403,  
 430, 457, 519, 534  
 Tsujimoto T, 310  
 Tucci DL, 594, 601  
 Tumanova NL, 485  
 Tuomisto L, 623  
 Turgeon SM, 277, 581  
 Turner JG, 250, 279, 588, 602, 604, 625  
 Tusa RJ, 7, 39, 66  
 Tweedale R, 568, 580
- Uchiyama H, 637  
 Ueda O, 131  
 Ueno S, 611, 619, 624  
 Ueyama T, 180  
 Ulehlova L, 582  
 Underhill AM, 452  
 Upton N, 604, 625  
 Urban A, 369, 592  
 Urban GP, 344, 376, 601  
 Urban R, 272, 304  
 Usunoff KG, 111
- Vale C, 550, 558  
 Valenzuela CF, 622  
 Valla J, 108  
 Vallbo ÅB, 426, 457  
 Vamecq J, 615  
 van Adel BA, 151, 262, 280  
 Van Bergeijk WA, 385, 404, 409  
 Van Bergem P, 602  
 Van Bockstaele EJ, 193, 199  
 van de Par S, 397, 409  
 van der Heijden M, 406  
 Van de Sande B, 406  
 Van Den Abbeele T, 276, 555  
 Van Lieshout DP, 83, 108, 188, 196, 632,  
 637  
 Van Middlesworth L, 604, 625  
 Van Noort J, 115, 120, 130, 157, 172, 181  
 Van Opstal AJ, 436, 452  
 Van Riswick JG, 452  
 Van Schaik A, 435, 457  
 Van Stokkum IHM, 360, 375  
 Vartanian I, 324, 337, 338, 344, 423, 451  
 Vasquez J, 58  
 Vater M, 3, 7, 8, 33, 55, 66, 85, 113,  
 116, 130, 147, 154, 162, 181, 186,  
 199, 207, 215, 229, 250, 251, 254,  
 260, 262, 267, 273, 274, 280, 286,  
 297, 300, 302, 304, 305, 310, 316,  
 344, 349, 361, 375, 497, 503, 508,  
 517, 527, 528, 529, 531, 534, 535,  
 567, 572, 580, 583  
 Vaughn MD, 300, 310  
 Veale JL, 336  
 Veatch LM, 605, 625  
 Vega-Saenz de Miera E, 304, 309, 310  
 Vehrencamp SL, 460, 468, 482, 485  
 Veinante P, 54, 245  
 Velenovsky DS, 52, 224  
 Velluti RA, 373, 620  
 Ventura D, 530  
 Vergara C, 285, 310  
 Vergnes M, 620, 622, 624  
 Versnel H, 325, 343, 344  
 Vetter DE, 17, 18, 67, 174, 181, 184,  
 185, 199  
 Vicini S, 125, 204, 274  
 Vidal PP, 197  
 Vielliard J, 530  
 Viemeister NF, 333, 344, 364, 375  
 Villa A, 63, 228, 345  
 Villa AEP, 64, 345  
 Vincent JP, 306  
 Viñuela A, 128, 170, 180, 181  
 Viola-McCabe KE, 617  
 Voigt HF, 13, 68, 345, 458  
 Volkov IO, 56  
 Vollenweider F, 197  
 Volman SF, 535  
 von Békésy G, 324, 344, 385, 409  
 Von Campenhausen M, 535  
 Von der Emde G, 461, 490, 491  
 von Economo C, 37, 67
- Wable J, 225  
 Wada JA, 605, 625  
 Wafford KA, 269, 280  
 Wagner H, 406, 498, 501, 509, 519, 520,  
 521, 524, 531, 532, 534, 535  
 Wagner T, 169, 172, 181, 253, 280, 283,  
 288, 294, 297, 298, 310, 325, 332,  
 344, 606, 611, 625  
 Wainer BH, 59  
 Waitzman DM, 70, 84, 113  
 Walkowiak W, 238, 245, 352, 375, 472,  
 473, 487, 488, 491, 492, 640  
 Wallace MN, 40, 67

- Wallace MS, 623  
Wallace MT, 555  
Wallach H, 378, 397, 399, 409, 444, 457  
Walsh EJ, 618  
Walsh EW, 343, 558, 583  
Walton JP, 116, 128, 178, 196, 225, 269,  
280, 349, 359, 364, 365, 368, 375,  
376, 414, 425, 461, 464, 465, 486,  
560, 562, 564, 566, 580, 581, 583, 600  
Wamsley JK, 281  
Wang C, 616, 621  
Wang D, 624  
Wang J, 268, 280, 315, 316, 344, 571,  
573, 574, 575, 591, 601  
Wang LY, 291, 310  
Wang X, 411, 413, 421, 425, 460, 492  
Wang Y, 108, 113  
Wang YX, 103, 113  
Wang Z-W, 491  
Wardley-Smith B, 616  
Warr WB, 8, 10, 14, 17, 18, 28, 67, 109,  
115, 120, 121, 130, 153, 185, 199,  
231, 233, 247, 312, 428, 430, 457  
Warter JM, 622  
Watanabe T, 58, 260, 280, 339, 406, 418,  
425, 637  
Watanabe Y, 131  
Waterhouse BD, 193, 195  
Watkins DW III, 112  
Watkins JC, 617  
Watson C, 78, 82, 111, 196  
Watson CD, 190, 196  
Weaver SP, 557  
Webb TE, 304  
Webber TJ, 208, 229  
Webster D, 195  
Webster DB, 3, 52, 60, 63, 64, 67, 110,  
127, 129, 153, 197, 199, 230, 247,  
486, 558, 601, 640  
Webster HdeF, 62  
Webster M, 558, 601  
Webster WR, 3, 13, 17, 65, 67, 107, 127,  
223, 224, 336, 340, 343, 344, 454  
Wecker JR, 604, 625  
Weedman DL, 21, 38, 43, 67  
Weeg MS, 464, 465, 492  
Weight FF, 328, 568, 617  
Weinberg RJ, 37, 67  
Weinberger NM, 35, 43, 51, 55, 67, 211,  
213, 226, 238, 245  
Weiser M, 291, 310  
Weiss DS, 444, 456, 616  
Wenstrup JJ, 29, 30, 67, 116, 123, 124,  
130, 131, 182, 191, 193, 199, 200,  
201, 202, 203, 204, 205, 207, 215,  
216, 218, 219, 220, 226, 227, 228,  
229, 230, 246, 315, 319, 340, 341,  
344, 352, 359, 366, 372, 373, 375,  
395, 409, 415, 424, 431, 453, 503,  
511, 513, 515, 517, 531, 532, 533,  
535, 638  
Wenthold RJ, 7, 9, 19, 51, 64, 67, 112,  
113, 151, 153, 248, 249, 250, 251,  
272, 275, 278, 280, 456, 639  
Wenzel EM, 438, 457  
Wepsic JG, 220, 230  
Weseloh R, 307  
West NR, 567, 581  
Westbook GL, 308  
Westby GWM, 246  
Westenfelder B, 554  
Westman J, 114  
Weth E, 557  
Wever EG, 361, 375  
White EL, 13, 37, 67, 232, 247  
White HS, 624  
White KR, 224, 369, 486  
Whitfield IC, 38, 54, 348, 371, 626, 639  
Whiting PJ, 269, 277, 280  
Whitley JM, 97, 113, 141, 154, 558  
Wiberg M, 84, 114  
Wible BA, 305  
Wickesberg RE, 13, 18, 67, 109, 132,  
138, 140, 141, 146, 152, 399, 409  
Wightman FL, 432, 433, 436, 457  
Wilczynski W, 469, 471, 487, 488, 491,  
492, 628, 629, 639, 640  
Wild JM, 498, 503, 505, 506, 535  
Willard FH, 116, 120, 121, 122, 129,  
131, 137, 138, 154, 164, 166, 169, 181  
Williams JB, 307  
Williams JT, 301, 304  
Williams RJ, 615  
Willott JF, 154, 181, 268, 269, 277, 280,  
281, 315, 316, 337, 344, 349, 376,  
559, 560, 561, 564, 566, 580, 581,  
583, 586, 587, 588, 589, 590, 591,  
592, 593, 595, 596, 597, 598, 599,  
600, 601, 602, 604, 625  
Wilson EM, 526

- Wilson GF, 62  
 Wilson JP, 408  
 Wilson MC, 277, 555, 581, 600  
 Wilson MJ, 277, 556, 582, 600  
 Wilson WW, 364, 376  
 Winer JA, vii, ix, 3, 5, 17, 19, 21, 23, 26, 27, 29, 30, 31, 33, 35, 37, 38, 39, 40, 41, 42, 43, 51, 54, 55, 61, 63, 68, 84, 99, 111, 113, 114, 134, 140, 154, 166, 178, 195, 199, 200, 205, 206, 207, 215, 223, 230, 231, 232, 233, 234, 238, 242, 245, 246, 247, 248, 250, 277, 281, 303, 318, 341, 345, 407, 497, 503, 535, 536, 558, 629, 630, 632, 633, 636, 637, 638, 640  
 Winfield JA, 83, 107, 322, 336, 362, 369  
 Winn HE, 54, 487  
 Winter IM, 229, 328, 341, 345, 372, 373, 638  
 Winters RW, 229  
 Wisden W, 269, 276, 281, 555  
 Withington DJ, 51, 336  
 Wixon HN, 617  
 Wolfart J, 490  
 Wollberg Z, 223  
 Wong D, 37, 42, 58, 68  
 Wong ROL, 228  
 Woods CI, 198, 414, 480, 488  
 Woodson W, 480, 488  
 Woodward DJ, 195  
 Woolf NK, 544, 557, 582, 594, 602  
 Woollard HH, 115, 131, 627, 640  
 Woolley SMN, 504, 506, 509, 536, 629, 640  
 Woolsey CN, 34, 52, 63, 66  
 Wooten GF, 309  
 Wotton JM, 517, 536  
 Wrege KS, 58  
 Wright DD, 113, 186, 199, 348, 376  
 Wu CH, 530  
 Wu FJ, 339  
 Wu JY, 617  
 Wu M, 373, 490, 532, 536  
 Wu SH, 103, 107, 114, 181, 273, 276, 281, 283, 288, 290, 297, 298, 300, 308, 310, 311, 581, 584  
 Wurtz RH, 426, 444, 457  
 Wygonski J, 374, 491  
 Wynne B, 269, 281  
 Wyttenbach RA, 446, 457  
 Xiao Z, 456, 534  
 Xu J, 281, 311  
 Yaari Y, 295, 306  
 Yahr MD, 245  
 Yajima Y, 31, 66  
 Yamashita T, 180  
 Yan J, 39, 48, 68, 216, 229, 230, 236, 247, 332, 345, 364, 376, 448, 457, 458, 536, 640  
 Yan QS, 625  
 Yanes C, 486  
 Yang L, 25, 68, 226, 238, 247, 260, 281, 297, 311, 328, 345, 511, 536, 572, 573, 575, 583, 604, 605, 613, 625  
 Yang LC, 362, 376  
 Yang Y, 89, 90, 114  
 Yasuda S, 297, 311, 620  
 Yasuhara O, 85, 114  
 Ye J, 616  
 Ye Q, 622  
 Ye Y, 407, 455  
 Yechikhov S, 604, 625  
 Yeomans JS, 31, 68, 191, 199, 600  
 Yeshurun Y, 223  
 Yigit Y, vii, ix, 7  
 Yin TCT, 15, 61, 68, 91, 108, 109, 111, 114, 130, 152, 153, 180, 217, 224, 226, 307, 341, 371, 378, 379, 380, 383, 384, 390, 392, 393, 396, 399, 403, 405, 406, 407, 408, 409, 410, 427, 428, 430, 435, 436, 445, 446, 447, 448, 449, 451, 453, 455, 456, 457, 458, 638  
 Yip PK, 613, 616, 625  
 Yosef OB, 340  
 Yoshihara Y, 126, 128, 131  
 Yost WA, 348, 374, 375, 376, 407, 410  
 Young ED, 9, 13, 31, 68, 186, 199, 319, 328, 343, 345, 431, 436, 438, 440, 455, 456, 458  
 Yurgelun-Todd DA, 456  
 Zafra F, 554  
 Zakarauskas P, 436, 458

- Zarbin MA, 267, 281  
Zarnowska ED, 622  
Zatorre RJ, 51  
Zellick R, 490  
Zeng F-G, 374, 491  
Zerlin S, 325, 345, 382, 409  
Zettel ML, 116, 128, 178, 196, 224, 497,  
528, 536, 567, 568, 583, 584  
Zhang DX, 114, 164, 181, 281  
Zhang H, 275, 281, 306, 311, 581, 625  
Zhang S, 199, 491  
Zhang Y, 29, 39, 68, 103, 114, 229, 281,  
311, 344, 458, 536, 584  
Zhang Z-W, 54, 245  
Zhao HB, 407, 455  
Zharskaja VD, 485  
Zheng W, 263, 281, 501, 524, 536  
Zhou X, 332, 345, 400, 410, 518, 530,  
536  
Zhu WJ, 274, 452, 491  
Zhuang L-Z, 491  
Zimmer SD, 50, 53  
Zimmerman E, 424  
Zimmerman M, 50, 336, 337, 424  
Zook JM, 33, 35, 60, 64, 75, 114,  
116, 122, 129, 131, 137, 141,  
143, 152, 154, 164, 180, 181, 204,  
218, 227, 230, 231, 238, 246, 497,  
501, 503, 508, 536, 537, 558, 630,  
640  
Zrull MC, 107  
Zurek PM, 399, 408, 410  
Zurita P, 330, 345  
Zwicker E, 324, 329, 345