



Commutative C^* -Algebras Generated by Toeplitz Operators on the Super Unit Ball

R. Quiroga-Barranco* and A. Sánchez-Nungaray

Abstract. We extend known results about commutative C^* -algebras generated Toeplitz operators over the unit ball to the supermanifold setup. This is obtained by constructing commutative C^* -algebras of super Toeplitz operators over the super ball $\mathbb{B}^{p|q}$ and the super Siegel domain $\mathbb{U}^{p|q}$ that naturally generalize the previous results for the unit ball and the Siegel domain. In particular, we obtain one such commutative C^* -algebra for each even maximal Abelian subgroup of automorphisms of the super ball.

Mathematics Subject Classification. Primary 47B35; Secondary 58C50, 32M15.

Keywords. Toeplitz operator, Super unit ball, Commutative C^* -algebras.

1. Introduction

In [4] it was proved, under mild conditions, that a C^* -algebra generated by Toeplitz operators is commutative on each weighted Bergman space of the unit disk if and only if there is a pencil of hyperbolic geodesics of the unit disk such that the symbols of the Toeplitz operators are constant on the cycles of this pencil. In fact, the cycles are the orbits of a one-parameter subgroup of isometries for the hyperbolic geometry on the unit disk. We note that there are three different non-conjugate model classes of such subgroups: elliptic, parabolic and hyperbolic. This provides us with the following scheme: the C^* -algebra generated by Toeplitz operators is commutative on each weighted Bergman space on the unit disk if and only if there is a maximal Abelian subgroup of Möbius transformations such that the symbols of the Toeplitz operators are invariant under the action of this subgroup.

A generalization of this scheme was given in [8–10]. The generalization is obtained by considering a maximal Abelian subgroup of biholomorphisms of the unit ball, then the C^* -algebra generated by the Toeplitz operators

The authors were supported by SNI and Conacyt grants.

*Corresponding author.

whose symbols are invariant under the action of such subgroup is commutative on each weighted Bergman space. It was also noted that there are five different non-conjugate model classes of such subgroups: quasi-elliptic, quasi-parabolic, quasi-hyperbolic, nilpotent, and quasi-nilpotent. We refer to the above mentioned works for further details.

On the other hand, the general theory of non-perturbative quantization for a class of Hermitian symmetric supermanifolds (a particular case is the super ball) was developed in [2, 3]. Such quantization is based on the notion of super Toeplitz operator defined on a suitable \mathbb{Z}_2 -graded Hilbert space of superholomorphic functions that effectively defines super Bergman spaces. These quantized supermanifolds yield the C^* -algebra generated by such super Toeplitz operators. Along these lines, in [5] it is given an exhaustive description of the super Toeplitz operators over the super ball using classical Toeplitz-type operators.

Recently and quite unexpectedly it was observed in [6, 7, 11] that there are five different non-conjugate classes of maximal abelian supergroups of isomorphisms of the super disk labeled by the names super-elliptic, quasi-elliptic, super-parabolic, quasi-parabolic and quasi-hyperbolic. In these works it is proved that the C^* -algebra of super Toeplitz operators whose symbols are invariant under the action of one of these subgroups is commutative on each weighted super Bergman space.

The main goal of this work is to extend the previous results and theory to the case of the super unit ball $\mathbb{B}^{p|q}$ and its unbounded realization $\mathbb{U}^{p|q}$, the super Siegel domain.

Hence, we introduce in Sect. 2 the super Bergman space for the super Siegel domain and prove its unitary equivalence with the super Bergman space of the super ball. This allows us to use the known theory to compute the super Bergman projection and define the super Toeplitz operators on the super Siegel domain.

With the above setup, we obtain the full list of even maximal Abelian subgroups of the group of automorphisms of the super unit ball. Our classification is based on the analysis of the corresponding maximal Abelian subalgebras that are described in Theorem 3.2. This is used in Sect. 3 to give an explicit description of the actions of the even maximal Abelian subgroups on the super unit ball. Most of these are easier to present in the super Siegel domain. There exists 5 non-equivalent types of even maximal Abelian subgroups for our supermanifold setup, one of them depending on a parameter for a total of $n+2$ different conjugacy classes. We label the five types with the names quasi-elliptic, quasi-parabolic, nilpotent, quasi-hyperbolic, and quasi-nilpotent.

Section 4 introduces a super Bargmann transform corresponding to each one of the conjugacy classes of even maximal Abelian subgroups mentioned above. These super Bargmann transforms generalize those presented in [9]. At the same time, our transforms allow us to prove that the C^* -algebra generated by the Toeplitz operators whose symbols are invariant by one of the even maximal Abelian subgroups is commutative. This is obtained in Sect. 6 and is thus the core of this work. The relevant results, for each type

of even maximal Abelian subgroups, are Theorems 6.1, 6.2, 6.3, 6.4, 6.5. We note that our results have the same strength of those for the classical case in that our approach using a super Bargmann transform allows us to realize the commuting Toeplitz operators in each case as multiplication operators.

2. Weighted Super Bergman Spaces and Projections

For $p \geq 1$, let $\mathcal{O}(\mathbb{B}^p)$ denote the algebra of holomorphic functions $\psi(z_1, \dots, z_p)$ on the open unit ball

$$\mathbb{B}^p = \{z = (z_1, \dots, z_p) : |z|^2 = |z_1|^2 + \dots + |z_p|^2 < 1\}$$

in \mathbb{C}^p .

For $p \geq 1$, let $\mathcal{O}(\mathbb{U}^p)$ denote the algebra of all functions $\psi(w_1, \dots, w_p)$ that are holomorphic on the Siegel domain

$$\mathbb{U}^p = \{w = (w', w_p) = (w_1, \dots, w_p) \in \mathbb{C}^p : \text{Im}(w_p) - |w'|^2 > 0\},$$

where $w' \in \mathbb{C}^{p-1}$.

Definition 2.1. For $\nu > p$, the weighted Bergman space

$$H_\nu^2(\mathbb{B}^p) = \mathcal{O}(\mathbb{B}^p) \cap L^2(\mathbb{B}^p, d\mu_\nu)$$

consists of all holomorphic functions on \mathbb{B}^p which are square-integrable for the probability measure

$$d\mu_\nu = c_\nu(1 - z\bar{z})^{\nu-p-1}dz, \quad \nu > p,$$

where the normalizing constant is given by

$$c_\nu = \frac{\Gamma(\nu)}{\pi^p \Gamma(\nu - p)}$$

with $z\bar{w} = z_1\bar{w}_1 + \dots + z_p\bar{w}_p$ on \mathbb{C}^p and dz is the Lebesgue measure. Correspondingly, the weighted Bergman space

$$H_\nu^2(\mathbb{U}^p) = \mathcal{O}(\mathbb{U}^p) \cap L^2(\mathbb{U}^p, d\tilde{\mu}_\nu)$$

consists of all holomorphic functions on \mathbb{U}^p which are square-integrable for the probability measure

$$d\tilde{\mu}_\nu = \frac{c_\nu}{4}(\text{Im}(w_p) - w'\bar{w}')^{\nu-p-1}dz, \quad \nu > p,$$

where dz is the Lebesgue measure.

It is well-known (see [12]) that $H_\nu^2(\mathbb{B}^p)$ has the reproducing kernel

$$K_{\mathbb{B}^p, \nu}(z, w) = (1 - z\bar{w})^{-\nu},$$

for all $z, w \in \mathbb{B}^p$. And that $H_\nu^2(\mathbb{U}^p)$ has the reproducing kernel

$$K_{\mathbb{U}^p, \nu}(z, w) = \left(\frac{z_n - \bar{w}_n}{2i} - z'\bar{w}' \right)^{-\nu},$$

for all $z, w \in \mathbb{U}^p$.

Let Λ_q denote the complex Grassmann algebra with the generators ξ_1, \dots, ξ_q satisfying the relations

$$\xi_i \xi_j + \xi_j \xi_i = 0,$$

for $1 \leq i, j \leq q$. If we take $Q := \{1, \dots, q\}$, then we have

$$\Lambda_q = \mathbb{C}\langle \xi_I : I \subset Q \rangle,$$

where $\xi_I = \xi_{i_1} \cdots \xi_{i_k}$ if $I = \{i_1 < \dots < i_k\}$. For disjoint subsets I, J we have $\xi_I \xi_J = \varepsilon_{I,J} \xi_{I \cup J}$ for some $\varepsilon_{I,J} = \pm 1$ whose value depends on the pair I, J .

The tensor product algebra

$$\mathcal{O}(\mathbb{B}^{p|q}) := \mathcal{O}(\mathbb{B}^p) \otimes \Lambda_q$$

consists of all super-holomorphic functions

$$\Psi(z, \xi) = \sum_{I \subset Q} \Psi_I \xi_I,$$

where $\Psi_I \in \mathcal{O}(\mathbb{B}^p)$ for all $I \subset Q$. In a similar way, the tensor product algebra

$$\mathcal{O}(\mathbb{U}^{p|q}) := \mathcal{O}(\mathbb{U}^p) \otimes \Lambda_q$$

consists of all super-holomorphic functions. In this case we have a similar expression for the super-holomorphic functions.

Let $\Lambda_q^{\mathbb{C}}$ denote the complex Grassmann algebra with the generators $\xi_1, \dots, \xi_q, \bar{\xi}_1, \dots, \bar{\xi}_q$ satisfying the relations

$$\begin{aligned} \xi_i \xi_j + \xi_j \xi_i &= 0, \\ \bar{\xi}_i \bar{\xi}_j + \bar{\xi}_j \bar{\xi}_i &= 0, \\ \xi_i \bar{\xi}_j + \xi_j \bar{\xi}_i &= 0, \end{aligned}$$

for $1 \leq i, j \leq q$.

Thus we have

$$\Lambda_q^{\mathbb{C}} = \mathbb{C}\langle \xi_I \xi_J^* : I, J \subset Q \rangle,$$

where $\xi_I = \xi_{i_1} \cdots \xi_{i_k}$ and $\xi_I^* = \bar{\xi}_{j_1} \cdots \bar{\xi}_{j_l}$ if $I = \{i_1 < \dots < i_k\}$ and $J = \{j_1 < \dots < j_l\}$.

The tensor product algebra

$$\mathcal{C}(\mathbb{B}^{p|q}) := \mathcal{C}(\mathbb{B}^p) \otimes \Lambda_q^{\mathbb{C}}$$

consists of all continuous super functions

$$\Psi = \sum_{I, J \subset Q} \Psi_{I,J} \xi_I \xi_J^*,$$

where $\Psi_{I,J} \in \mathcal{C}(\mathbb{B}^p)$ for all $I, J \subset Q$. Similarly

$$\mathcal{C}(\mathbb{U}^{p|q}) := \mathcal{C}(\mathbb{U}^p) \otimes \Lambda_q^{\mathbb{C}}$$

consists of all continuous super functions

$$\Psi = \sum_{I, J \subset Q} \Psi_{I,J} \xi_I \xi_J^*,$$

where $\Psi_{I,J} \in \mathcal{C}(\mathbb{U}^p)$ for all $I, J \subset Q$.

There is a natural involution $\Psi \mapsto \Psi^*$ on these spaces of super functions, which is defined by

$$\Psi^* = \sum_{I, J \subset Q} \bar{\Psi}_{I, J} \xi_J \xi_I^*,$$

for Ψ as above.

The Berezin integral on $\mathbb{V}^{p|q}$ is defined by

$$\int_{\mathbb{V}^{p|q}} dz d\xi F(z, \xi) = \int_{\mathbb{V}^p} f_{Q, Q}(z) dz$$

for $F \in \mathcal{C}(\mathbb{V}^{p|q})$, where the normalization is given by

$$\int_{\mathbb{V}^{p|q}} dz d\xi \xi_Q^* \xi_Q = 1,$$

where \mathbb{V}^p is either \mathbb{B}^p or \mathbb{U}^p .

For any given morphism γ between super domains we define

$$\gamma'(Z) = \text{Ber} \begin{pmatrix} \frac{\partial w}{\partial z} & \frac{\partial \omega}{\partial z} \\ \frac{\partial w}{\partial \xi} & \frac{\partial \omega}{\partial \xi} \end{pmatrix} = \text{Ber} \frac{\partial W}{\partial Z}, \tag{2.1}$$

where $Z = (z_1, \dots, z_p, \xi_1, \dots, \xi_q)$, $W = (w_1, \dots, w_p, \omega_1, \dots, \omega_q)$ and where the Berezinian is defined as follows

$$\text{Ber} \begin{pmatrix} A & B \\ C & D \end{pmatrix} = \det(A - BD^{-1}C) \det(D)^{-1}. \tag{2.2}$$

We refer to [1] for more details.

We now recall a natural biholomorphism of supermanifolds between the super p -ball $\mathbb{B}^{p|q}$ and the super Siegel domain $\mathbb{U}^{p|q}$.

We define the super Cayley transform from $\mathbb{B}^{p|q}$ to $\mathbb{U}^{p|q}$ in local coordinates by

$$\psi(z_1, \dots, z_p, \xi_1, \dots, \xi_q) = (w_1, \dots, w_p, \omega_1, \dots, \omega_q), \tag{2.3}$$

where

$$\begin{aligned} w_k &= \frac{iz_k}{1 + z_p}, & \text{for } k = 1, \dots, p - 1, \\ w_p &= i \frac{1 - z_p}{1 + z_p}, \\ \omega_k &= \frac{i\xi_k}{1 + z_p}, & \text{for } k = 1, \dots, q. \end{aligned}$$

The inverse transform is given by

$$\begin{aligned} z_k &= \frac{-2iw_k}{1 - iw_p}, & \text{for } k = 1, \dots, p - 1, \\ z_p &= i \frac{1 + iw_p}{1 - iw_p}, \\ \xi_k &= \frac{-2i\omega_k}{1 - iw_p}, & \text{for } k = 1, \dots, q. \end{aligned}$$

Lemma 2.2. *Let $Z = (z, \xi)$, $U = (u, \eta)$, $\psi(Z) = (w, \omega) = W$ and $\psi(U) = (v, \zeta) = V$ where ψ is given by (2.3). Then, we have*

$$(1 + z\bar{u} - \xi\bar{\eta})(1 + z_p)^{-1}(1 + u_p)^{-1} = \left(\frac{w_p - \bar{v}_p}{2i} - w'\bar{v}' - \omega\bar{\zeta} \right), \tag{2.4}$$

$$(1 + z\bar{u} - \xi\bar{\eta}) = \left(\frac{w_p - \bar{v}_p}{2i} - w'\bar{v}' - \omega\bar{\zeta} \right) 4(1 - iw_p)^{-1}(1 - iv_p)^{-1}, \tag{2.5}$$

where $\xi\bar{\theta} = \xi_1\bar{\theta}_1 + \dots + \xi_q\bar{\theta}_q$ and $z\bar{w} = z_1\bar{w}_1 + \dots + z_p\bar{w}_p$.

Proof. Using the above and substituting $\psi(Z) = (w, \omega) = W$ and $\psi(U) = (v, \zeta) = V$ on the left hand side of Eq. (2.4) it follows that

$$\begin{aligned} & \left(\frac{w_p - \bar{v}_p}{2i} - w'\bar{v}' - \omega\bar{\zeta} \right) \\ &= \left(\frac{1}{2i} \left[i \frac{1 - z_p}{1 + z_p} - i \frac{1 - u_p}{1 + u_p} \right] - \sum_{k=1}^{p-1} \frac{iz_k}{1 + z_p} \frac{\overline{i u_k}}{1 + u_p} - \sum_{k=1}^q \frac{i\xi_k}{1 + z_p} \frac{\overline{i\eta_k}}{1 + u_p} \right) \\ &= \left(\frac{1}{2} [(1 - z_p)(1 + \bar{u}_p) + (1 - \bar{u}_p)(1 + z_p)] - \sum_{k=1}^{p-1} z_k\bar{u}_k - \sum_{k=1}^q \xi_k\bar{\eta}_k \right) \\ &\quad \times \frac{1}{(1 + z_p)} \frac{1}{(1 + \bar{u}_p)} \\ &= \frac{1}{(1 + z_p)} \frac{1}{(1 + \bar{u}_p)} \left(1 - z_p\bar{u}_p - \sum_{k=1}^{p-1} z_k\bar{v}_k - \sum_{k=1}^q \xi_k\bar{\eta}_k \right) \\ &= (1 + z_p)^{-1}(1 + u_p)^{-1}(1 + z\bar{u} - \xi\bar{\eta}). \end{aligned}$$

Its clear that $1 + z_p = 2(1 - iw_p)^{-1}$, then we obtain (2.5). □

Definition 2.3. For any parameter $\nu > p - q + 1$, the (weighted) super-Bergman space

$$H^2_\nu(\mathbb{B}^{p|q}) \subset \mathcal{O}(\mathbb{B}^{p|q})$$

consists of all super-holomorphic functions $\Psi(z, \xi)$ which satisfy the square-integrability condition

$$\begin{aligned} & (\Psi|\Psi)_{\mathbb{B}^{p|q}, \nu} \\ &= \frac{\Gamma(\nu)}{\pi^p \Gamma(\nu + q - p)} \int_{\mathbb{B}^{p|q}} dzd\xi (1 - z\bar{z} - \xi\bar{\xi})^{\nu + q - p - 1} \Psi(z, \xi)^* \Psi(z, \xi) < \infty, \end{aligned}$$

where $\xi\bar{\theta} = \xi_1\bar{\theta}_1 + \dots + \xi_q\bar{\theta}_q$ and $z\bar{w} = z_1\bar{w}_1 + \dots + z_p\bar{w}_p$. Moreover

$$\frac{(1 - z\bar{z} - \xi\bar{\xi})^{\nu + q - p - 1}}{\pi^p \Gamma(\nu + q - p)} = \sum_{J \subset Q} \frac{(1 - z\bar{z})^{\nu + |J| - p - 1}}{\pi^p \Gamma(\nu + |J| - p)} \xi_{Q \setminus J}^* \xi_{Q \setminus J}$$

In [5] the authors proved that the super Bergman space has a decomposition in direct sum of classical Bergman space and gave an explicit expression for the super Bergman projection. If we take $\Psi = \sum_{M \subset Q} \psi_M \xi_M \in$

$H_\nu^2(\mathbb{B}^{p|q}) \subset \mathcal{O}(\mathbb{B}^{p|q})$ then the inner product has the form

$$\begin{aligned} & \frac{\Gamma(\nu)}{\pi^p \Gamma(\nu + q - p)} \int_{\mathbb{B}^{p|q}} dz d\xi (1 - z\bar{z} - \xi\bar{\xi})^{\nu+p-p-1} \Psi(z, \xi)^* \Psi(z, \xi) \\ &= \sum_{m=1}^q \frac{\Gamma(\nu)}{\Gamma(\nu + m)} \sum_{M \subset Q, |M|=m} \|\psi_M(z)\|_{\mathbb{B}^{p, \nu+m}}^2. \end{aligned} \tag{2.6}$$

Thus the super Bergman space over the super ball has an orthogonal decomposition

$$H_\nu^2(\mathbb{B}^{p|q}) = \sum_{m=0}^q H_{\nu+m}^2(\mathbb{B}^p) \otimes \Lambda^m(\mathbb{C}^q)$$

into a sum of weighted Bergman spaces for $0 \leq m \leq q$, with multiplicity $\binom{m}{q}$. Moreover, this super Bergman space has the reproducing kernel property

$$\begin{aligned} P_{\mathbb{B}^{p|q}, \nu} \Psi(z, \xi) &= \frac{\Gamma(\nu)}{\pi^p \Gamma(\nu + q - p)} \int_{\mathbb{B}^{p|q}} dw d\omega (1 - w\bar{w} - \omega\bar{\omega})^{\nu+q-p-1} \\ &\quad \times (1 - z\bar{w} - \xi\bar{\omega})^{-\nu} \Psi(w, \omega) = \Psi(w, \omega). \end{aligned}$$

In other words, $H_\nu^2(\mathbb{B}^{p|q})$ has the reproducing kernel

$$K_{\mathbb{B}^{p|q}, \nu}(z, \xi, w, \omega) = (1 - z\bar{w} - \xi\bar{\omega})^{-\nu}.$$

Definition 2.4. For any parameter $\nu > p - q + 1$, the (weighted) super-Bergman space

$$H_\nu^2(\mathbb{U}^{p|q}) \subset \mathcal{O}(\mathbb{U}^{p|q})$$

consists of all super-holomorphic functions $\Psi(w, \omega)$ which satisfy the square-integrability condition

$$\begin{aligned} (\Psi|\Psi)_{\mathbb{U}^{p|q}, \nu} &= \frac{\Gamma(\nu)}{4\pi^p \Gamma(\nu + q - p)} \int_{\mathbb{U}^{p|q}} dw d\omega (\text{Im}(w_p) - w'\bar{w}' - \omega\bar{\omega})^{\nu+q-p-1} \\ &\quad \times \Psi(z, \omega)^* \Psi(w, \omega) < \infty, \end{aligned}$$

where $\xi\bar{\omega} = \xi_1\bar{\omega}_1 + \dots + \xi_q\bar{\omega}_q$ and $z'\bar{w}' = z_1\bar{w}_1 + \dots + z_{p-1}\bar{w}_{p-1}$. Where one can prove that

$$\begin{aligned} & \frac{1}{4\pi^p \Gamma(\nu + q - p)} \left(\frac{w_p - \bar{w}_p}{2i} - w'\bar{w}' - \omega\bar{\omega} \right)^{\nu+q-p-1} \\ &= \sum_{J \subset Q} \frac{1}{4\pi^p \Gamma(\nu + |J| - p)} \left(\frac{w_p - \bar{w}_p}{2i} - w'\bar{w}' \right)^{\nu+|J|-p-1} \omega_{Q \setminus J}^* \omega_{Q \setminus J}. \end{aligned}$$

We now observe that the Berezinian of the Jacobian matrix of the transformation ψ given by (2.3) satisfies

$$\psi'(Z) = \text{Ber} \begin{pmatrix} A & B \\ 0 & C \end{pmatrix} = \det(A) \cdot \det(C^{-1}),$$

where

$$A = \text{Diagonal}(i(1 + z_p)^{-1}, \dots, i(1 + z_p)^{-1}, -2i(1 + z_p)^{-2})$$

and

$$C = \text{Diagonal}(i(1 + z_p)^{-1}, \dots, i(1 + z_p)^{-1}).$$

As a consequence we obtain that $\psi'(Z) = -2i^{p-q+1}(1 + z_p)^{q-p-1}$ and analogously $(\psi^{-1})'(W) = -2^{p-q}i^{q-p-1}(1 - iw_p)^{q-p-1}$.

Definition 2.5. We define the operator $U_\nu : H_\nu^2(\mathbb{B}^{p|q}) \rightarrow H_\nu^2(\mathbb{U}^{p|q})$ given by

$$U_\nu(\Psi)(W) = \Psi(\psi^{-1}(W)) \left(\frac{2}{1 - iw_p} \right)^\nu,$$

and its adjoint $U_\nu^* : H_\nu^2(\mathbb{U}^{p|q}) \rightarrow H_\nu^2(\mathbb{B}^{p|q})$ given by

$$U_\nu^*(\Psi)(Z) = \Psi(\psi(Z)) \left(\frac{1}{1 + z_p} \right)^\nu.$$

Theorem 2.6. *The operator $U_\nu : H_\nu^2(\mathbb{B}^{p|q}) \rightarrow H_\nu^2(\mathbb{U}^{p|q})$ is unitary.*

Proof. Consider $\Psi \in H_\nu^2(\mathbb{B}^{p|q})$. Then we have

$$(\Psi|\Psi)_{\mathbb{B}^{p|q}, \nu} = \frac{\Gamma(\nu)}{\pi^p \Gamma(\nu + q - p)} \int_{\mathbb{B}^{p|q}} dz d\xi (1 - z\bar{z} - \xi\bar{\xi})^{\nu+q-p-1} \Psi(z, \xi)^* \Psi(z, \xi).$$

Using Lemma 2.2 and the change of variable given by the super Cayley transform (2.3) on the right side of the above equation, we obtain

$$\begin{aligned} & \frac{\Gamma(\nu)}{\pi^p \Gamma(\nu + q - p)} \int_{\mathbb{U}^{p|q}} dw d\omega (\psi^{-1})'(W) \overline{(\psi^{-1})'(W)} \\ & \times (\text{Im}(w_p) - w'\bar{w}' - \omega\bar{\omega})^{\nu+q-p-1} 4^{\nu+q-p-1} ((1 - iw_p)^{-1})^{\nu+q-p-1} \\ & \times \overline{(1 - iw_p)^{-1}}^{\nu+q-p-1} \Psi(\psi^{-1}(W))^* \Psi(\psi^{-1}(W)) \\ & = \frac{\Gamma(\nu)}{\pi^p \Gamma(\nu + q - p)} \int_{\mathbb{U}^{p|q}} dw d\omega 4^{p-q} (1 - iw_p)^{q-p-1} \overline{(1 - iw_p)^{q-p-1}} \\ & \times \left(\frac{w_p - \bar{w}_p}{2i} - w'\bar{z}' - \omega\bar{\omega} \right)^{\nu+q-p-1} 4^{\nu+q-p-1} ((1 - iw_p)^{-1})^{\nu+q-p-1} \\ & \times \overline{(1 - iw_p)^{-1}}^{\nu+q-p-1} \Psi(\psi^{-1}(W))^* \Psi(\psi^{-1}(W)) \\ & = \frac{\Gamma(\nu)}{4\pi^p \Gamma(\nu + q - p)} \int_{\mathbb{U}^{p|q}} dw d\omega \left(\frac{w_p - \bar{w}_p}{2i} - w'\bar{z}' - \omega\bar{\omega} \right)^{\nu+q-p-1} \\ & \times \left[\Psi(\psi^{-1}(W)) \left(\frac{2}{1 - iw_p} \right)^\nu \right]^* \left[\Psi(\psi^{-1}(W)) \left(\frac{2}{1 - iw_p} \right)^\nu \right] \\ & = (U_\nu(\Psi)|U_\nu(\Psi))_{\mathbb{U}^{p|q}, \nu}. \end{aligned}$$

□

Corollary 2.7. *If $\Psi = \sum_{M \subset Q} \psi_M \xi_M \in \mathcal{O}_\nu(\mathbb{U}^{p|q})$, then we have*

$$\frac{\Gamma(\nu)}{\pi^p \Gamma(\nu + q - p)} \int_{\mathbb{U}^{p|q}} dw d\omega \left(\frac{w_p - \bar{w}_p}{2i} - w' \bar{w}' - \omega \bar{\omega} \right)^{\nu+q-p-1} \\ \times \Psi(w, \omega)^* \Psi(w, \omega) = \sum_{m=1}^q \frac{\Gamma(\nu)}{\Gamma(\nu + m)} \sum_{M \subset Q, |M|=m} \|\psi_M(z)\|_{\mathbb{U}^p, \nu+m}^2.$$

In other words, there is an orthogonal decomposition

$$H_\nu^2(\mathbb{U}^{p|q}) = \sum_{m=0}^q H_{\nu+m}^2(\mathbb{U}^p) \otimes \Lambda^m(\mathbb{C}^q)$$

into a sum of weighted Bergman spaces for $0 \leq m \leq q$, whose corresponding multiplicities are $\binom{m}{q}$.

Proof. We rewrite the operator U_ν as follows

$$U_\nu(\Psi)(W) = \sum_{M \subset Q} (-i)^{|M|} \psi_M(\psi_0^{-1}(w)) \left(\frac{2}{1 - iw_p} \right)^{\nu+|M|} \omega_M \\ = \sum_{M \subset Q} (-i)^{|M|} V_{\nu+|M|}(\psi_M)(w) \omega_M,$$

where $V_{\nu+m} : H_{\nu+m}^2(\mathbb{B}^p) \rightarrow H_{\nu+m}^2(\mathbb{U}^p)$ is defined by

$$V_{\nu+m}(f) = f(\psi_0^{-1}(w)) \left(\frac{2}{1 - iw_p} \right)^{\nu+|M|}.$$

Since we know that $V_{\nu+m}$ is a unitary operator, the result follows. □

Proposition 2.8. *For $\nu > p$ and $\Psi = \sum_{M \subset Q} \psi_M \xi_M \in \mathcal{O}_\nu(\mathbb{U}^{p|q})$, we have the reproducing kernel property*

$$P_{\mathbb{U}^{p|q}, \nu} \Psi(w, \omega) = \frac{\Gamma(\nu)}{\pi^p \Gamma(\nu + q - p)} \int_{\mathbb{U}^{p|q}} dv d\zeta \left(\frac{v_p - \bar{v}_p}{2i} - v' \bar{v}' - \zeta \bar{\zeta} \right)^{\nu+q-p-1} \\ \times \left(\frac{w_p - \bar{w}_p}{2i} - w' \bar{w}' - \omega \bar{\omega} \right)^{-\nu} \Psi(v, \zeta) = \Psi(w, \omega).$$

In particular, $H_\nu^2(\mathbb{U}^{p|q})$ has the reproducing kernel

$$K_{\mathbb{U}^{p|q}, \nu}(w, \omega, v, \zeta) = \left(\frac{w_p - \bar{w}_p}{2i} - w' \bar{w}' - \omega \bar{\omega} \right)^{-\nu}.$$

Proof. We know that U_ν is unitary, therefore the Bergman projection over $H_\nu^2(\mathbb{U}^{p|q})$ is given by

$$P_{\nu, \mathbb{U}^{p|q}} = U_\nu P_{\nu, \mathbb{B}^{p|q}} U_\nu^*,$$

thus $P_{\mathbb{U}^{p|q}, \nu}(\Psi)$ is given by

$$U_\nu \left[\frac{\Gamma(\nu)}{\pi^p \Gamma(\nu + q - p)} \int_{\mathbb{B}^{p|q}} dud\eta (1 - u\bar{u} - \eta\bar{\eta})^{\nu+q-p-1} (1 - z\bar{u} - \xi\bar{\eta})^{-\nu} U_\nu^*(\Psi(V)) \right].$$

Using the change of variable (2.3), taking $W = (w, \omega) = \psi(z, \xi) = \psi(Z)$, $V = (v, \zeta) = \psi(u, \eta) = \psi(U)$ and Lemma 2.2, we obtain

$$\begin{aligned}
 & P_{\mathbb{U}^{p|q}, \nu}(\Psi) \\
 &= \left(\frac{2}{1 - iw_p} \right)^\nu \frac{\Gamma(\nu)}{4\pi^p \Gamma(\nu + q - p)} \int_{\mathbb{U}^{p|q}} dvd\zeta \left(\frac{v_p - \bar{v}_p}{2i} - v' \bar{v}' - \zeta \bar{\zeta} \right)^{\nu + q - p - 1} \\
 &\quad \times (2(1 - iv_p)^{-1})^\nu (2(1 - i\bar{v}_p)^{-1})^\nu \Psi(V) \left(\frac{1}{1 + u_p} \right)^\nu \\
 &\quad \times \left[\left(\frac{w_p - \bar{v}_p}{2i} - w' \bar{v}' - \omega \bar{\zeta} \right) 4(1 - iw_p)^{-1} (1 - i\bar{v}_p)^{-1} \right]^{-\nu} \\
 &= \frac{\Gamma(\nu)}{4\pi^p \Gamma(\nu + q - p)} \int_{\mathbb{U}^{p|q}} dvd\zeta \left(\frac{v_p - \bar{v}_p}{2i} - v' \bar{v}' - \zeta \bar{\zeta} \right)^{\nu + q - p - 1} \\
 &\quad \times \left(\frac{w_p - \bar{v}_p}{2i} - w' \bar{v}' - \omega \bar{\zeta} \right)^{-\nu} \Psi(V) \left(\frac{1}{1 + u_p} \right)^\nu \left(\frac{2}{1 - iv_p} \right)^\nu \\
 &= \frac{\Gamma(\nu)}{4\pi^p \Gamma(\nu + q - p)} \int_{\mathbb{U}^{p|q}} dvd\zeta \left(\frac{v_p - \bar{v}_p}{2i} - v' \bar{v}' - \zeta \bar{\zeta} \right)^{\nu + q - p - 1} \\
 &\quad \times \left(\frac{w_p - \bar{v}_p}{2i} - w' \bar{v}' - \omega \bar{\zeta} \right)^{-\nu} \Psi(V).
 \end{aligned}$$

□

3. The Super-Group $SU(p, 1|q)$ and Its Even MASG's

There is a super Lie group denoted by $SU(p, 1|q)$ that is naturally associated to the super unit ball $B^{p,q}$ and whose definition we now recall (see [2]). The base manifold of $SU(p, 1|q)$ is the Lie group $SU(p, 1) \times SU(q)$. For the structure sheaf we use the Grassmann algebra $\Lambda(M_{p+q+1}(\mathbb{C}))$, where $M_{p+q+1}(\mathbb{C})$ is the space of complex square matrices of size $(p + q + 1) \times (p + q + 1)$, and we consider the tensor product

$$C^\infty(SU(p, 1)) \otimes \Lambda(M_{p+q+1}(\mathbb{C})).$$

The variables corresponding to the matrix entries are given the following parity assignments

$$p(\gamma_{jk}) = p(\bar{\gamma}_{jk}) = \begin{cases} 0, & \text{if } 1 \leq j, k \leq p + 1 \text{ or } p + 1 < j, k \leq p + q + 1, \\ 1, & \text{otherwise.} \end{cases}$$

Thus we have that the super-matrix has a natural block decomposition as follows

$$\gamma = \begin{pmatrix} A & C \\ D & B \end{pmatrix},$$

where A and B are even square matrices with sizes $(p + 1) \times (p + 1)$ and $q \times q$, respectively, and C and D are odd matrices with sizes $q \times (p + 1)$

and $(p + 1) \times q$, respectively. The structure sheaf of $SU(p, 1|q)$ is obtained by considering the set of matrices γ as above that satisfy

$$\text{Ber}\gamma = 1, \quad \gamma^* J_{p,1|q} \gamma = J_{p,1|q},$$

where

$$J_{p,1|q} = \begin{pmatrix} I_p & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -I_q \end{pmatrix}.$$

In particular, the Lie super algebra $\mathfrak{su}(p, 1|q)$ of $SU(p, 1|q)$ is given by the set of matrices γ that satisfy the conditions

$$\text{str}(\gamma) = 0, \quad \gamma^* J_{p,1|q} + J_{p,1|q} \gamma^* = 0,$$

where $\text{str}(\gamma) = \text{tr}(A) - \text{tr}(B)$ and $\text{tr}(A)$ are the usual supertrace and trace, respectively. Since the parity of the entries for such matrices is defined as above, we conclude that the Lie algebra of even elements of $\mathfrak{su}(p, 1|q)$ is given by

$$\mathfrak{su}(p, 1|q)_0 = \left\{ \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix} : A \in \mathfrak{u}(p, 1), B \in \mathfrak{u}(q), \text{tr}(A) = \text{tr}(B) \right\}.$$

On the other hand, the super Siegel domain realization $\mathbb{U}^{p|q}$ of the super unit ball $\mathbb{B}^{p|q}$ together with the super Cayley transform introduced before yield another realization of the super Lie group $SU(p, 1|q)$. More precisely, we obtain the super Lie group $SU(K_{p|q})$ whose base Lie group is

$$SU(K_p) \times SU(q),$$

where $SU(K_p)$ is the Lie group of unitary transformation of the pseudo-Hermitian form on \mathbb{C}^{p+1} whose matrix is the following (see [10] for comparison)

$$K_p = \begin{pmatrix} 2I_{p-1} & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix}.$$

The structure sheaf of $SU(K_{p|q})$ is given by the set of matrices γ as above that now satisfy

$$\text{Ber}(\gamma) = 1, \quad \gamma^* K_{p|q} \gamma = K_{p|q},$$

where

$$K_{p|q} = \begin{pmatrix} K_p & 0 \\ 0 & -I_q \end{pmatrix}.$$

In particular, the super Lie algebra of $SU(K_{p|q})$, denoted by $\mathfrak{su}(K_{p|q})$, has the following Lie algebra as space of even elements

$$\mathfrak{su}(K_{p|q})_0 = \left\{ \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix} : A \in \mathfrak{u}(K_p), B \in \mathfrak{u}(q), \text{tr}(A) = \text{tr}(B) \right\}.$$

Here, we have used the notation where $\mathfrak{u}(K_p)$ denotes the Lie algebra of the Lie group $U(K_p)$ of unitary transformations for the pseudo-Hermitian product with matrix K_p .

We note that $SU(p, 1|q)$, $SU(K_{p|q})$ and their Lie algebras are conjugated through the super Cayley transform considered before. At the base manifold level this implies that $\mathfrak{su}(p, 1)$ and $\mathfrak{su}(K_p)$ are conjugate.

Definition 3.1. An even maximal Abelian subalgebra of $\mathfrak{su}(p, 1|q)$, or an even MASA for short, is a maximal Abelian subalgebra of the Lie algebra $\mathfrak{su}(p, 1|q)_0$. An even maximal Abelian subgroup of $SU(p, 1|q)$, or an even MASG, is a connected super Lie subgroup of $SU(p, 1|q)$ whose Lie super algebra is an even MASA.

We now list the collection of all even MASA of $\mathfrak{su}(p, 1|q)$ up to conjugacy. Note that since $\mathfrak{su}(p, 1|q)$ and $\mathfrak{su}(K_{p|q})$ are conjugated the even MASA of both super Lie algebras correspond to each other. In particular, the conjugacy classes of even MASA's can be described in terms of either one of these super Lie algebras. Also note that it follows from the above remarks that the maximal Abelian subalgebras of $\mathfrak{su}(p, 1)$ and $\mathfrak{su}(K_p)$ correspond to each other as well.

Theorem 3.2. For every even MASA \mathfrak{h} of $\mathfrak{su}(p, 1|q)$ there exist maximal Abelian subalgebras $\mathfrak{h}_1 \subset \mathfrak{u}(p, 1)$ and $\mathfrak{h}_2 \subset \mathfrak{u}(q)$ such that \mathfrak{h} is conjugate to the even MASA

$$\mathfrak{h}_1 \boxtimes_0 \mathfrak{h}_2 = \left\{ \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix} : A \in \mathfrak{h}_1, B \in \mathfrak{h}_2, \text{tr}(A) = \text{tr}(B) \right\}.$$

Furthermore, we can assume that \mathfrak{h}_2 is the Lie subalgebra of diagonal matrices in $\mathfrak{u}(q)$ and that \mathfrak{h}_1 is given by one of the following where $D(k)$ denotes the Lie algebra of $k \times k$ diagonal matrices with pure imaginary entries.

1. *Quasi-elliptic:* The Lie subalgebra of diagonal matrices in $\mathfrak{u}(p, 1)$.
2. *Quasi-parabolic:* The Lie subalgebra of $\mathfrak{su}(K_p)$ that consists of the matrices of the form

$$\begin{pmatrix} D & 0 & 0 \\ 0 & z & 0 \\ 0 & 0 & \bar{z} \end{pmatrix},$$

where $D \in D(p - 1)$, $z \in \mathbb{C}$ and $\text{tr}(D) + 2i\text{Im}(z) = 0$.

3. *Quasi-hyperbolic:* The Lie subalgebra of $\mathfrak{su}(K_p)$ that consists of the matrices of the form

$$\begin{pmatrix} D & 0 & 0 \\ 0 & iy & a \\ 0 & 0 & iy \end{pmatrix},$$

where $D \in D(p - 1)$, $a, y \in \mathbb{R}$ and $\text{tr}(D) + 2iy = 0$.

4. *Nilpotent:* The Lie subalgebra of $\mathfrak{su}(K_p)$ that consists of the matrices of the form

$$\begin{pmatrix} 0 & 0 & b^t \\ 2ib & 0 & a \\ 0 & 0 & 0 \end{pmatrix},$$

where $a \in \mathbb{R}$ and $b \in \mathbb{R}^{p-1}$.

5. *Quasi-nilpotent: For some k such that $1 \leq k \leq p - 2$, the Lie subalgebra of $\mathfrak{su}(K_p)$ that consists of the matrices of the form*

$$\begin{pmatrix} D & 0 & 0 & 0 \\ 0 & iyI_{p-k-1} & 0 & b^t \\ 0 & 2ib & iy & a \\ 0 & 0 & 0 & iy \end{pmatrix},$$

where $a, y \in \mathbb{R}$, $b \in \mathbb{R}^{p-k-1}$, $D \in D(k)$ and $\text{tr}(D) + iy(p - k + 1) = 0$.

Proof. Let \mathfrak{h} be an even MASA subalgebra of $\mathfrak{su}(p, 1|q)$. In particular, \mathfrak{h} is a MASA (maximal Abelian subalgebra) of $\mathfrak{su}(p, 1|q)_0$. We first note that

$$\mathfrak{su}(p, 1|q)_0 \simeq \mathfrak{su}(p, 1) \times \mathfrak{u}(q)$$

as Lie algebras where the isomorphism is given by the assignment

$$\begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix} \mapsto \left(A - \frac{1}{p+1} \text{tr}(A)I_{p+1}, B \right).$$

And so, we can consider \mathfrak{h} as a MASA of $\mathfrak{su}(p, 1) \times \mathfrak{u}(q)$.

Let \mathfrak{h}_1 and \mathfrak{h}_2 be the projections of \mathfrak{h} into the first and second factors, respectively. Hence, \mathfrak{h}_1 and \mathfrak{h}_2 are both Abelian Lie algebras. It is clear that $\mathfrak{h} \subset \mathfrak{h}_1 \times \mathfrak{h}_2$ and the maximality of \mathfrak{h} implies that $\mathfrak{h} = \mathfrak{h}_1 \times \mathfrak{h}_2$. Furthermore, the same argument shows that \mathfrak{h}_1 and \mathfrak{h}_2 are MASA's of $\mathfrak{su}(p, 1)$ and $\mathfrak{u}(q)$, respectively. It is well known that there is a single conjugacy class of MASA's of $\mathfrak{u}(q)$ with a representative given by $D(q)$. The conjugacy classes of MASA's of $\mathfrak{su}(p, 1)$ are also known and they are listed in [10]. From this and the above isomorphism of Lie algebras the result now follows directly. It is also important to note that the MASA's of $\mathfrak{u}(p, 1)$ are of the form $\mathfrak{h}_1 \times \mathbb{R}$ where \mathfrak{h}_1 is a MASA of $\mathfrak{su}(p, 1)$. □

After exponentiating the even MASA's listed above we obtain the following even MASG's viewed through their actions on either $\mathbb{B}^{p|q}$ or $\mathbb{U}^{p|q}$. The content of Theorem 3.2 is that, up to conjugacy, these are the only MASG's of $\text{SU}(p, 1|q)$.

Quasi-elliptic group of super biholomorphisms of the super unit ball $\mathbb{B}^{p|q}$ is isomorphic to \mathbb{T}^{p+q} with the following group action:

$$(t, s) : (z, \theta) \in \mathbb{B}^{p|q} \mapsto (tz, s\theta) = (t_1z_1, \dots, t_pz_p, s_1\theta_1, \dots, s_q\theta_q) \in \mathbb{B}^{p|q},$$

for each $t = (t_1, \dots, t_p, s_1, \dots, s_q)$.

Note that if the super function F is invariant under the action of the quasi-elliptic group, then

$$F(z, \xi) = \sum_{I \subset Q} f_I(r) \xi_I \xi_I^*,$$

where $r = (r_1, \dots, r_p) = (|z_1|, \dots, |z_p|)$.

Quasi-parabolic group of biholomorphisms of the super Siegel domain $\mathbb{U}^{p|q}$ is isomorphic to $\mathbb{T}^{p-1} \times \mathbb{R} \times \mathbb{T}^q$ with the following group action:

$$(t, h, s) : (z', z_p, \theta) \in \mathbb{U}^{p|q} \mapsto (tz', z_p + h, s\theta) \in \mathbb{U}^{p|q},$$

for each $(t, h, s) \in \mathbb{T}^{p-1} \times \mathbb{R} \times \mathbb{T}^q$.

In this case, if the super function F is invariant under the action of the quasi-parabolic group, then

$$F(z, \xi) = \sum_{I \subset Q} f_I(r', \text{Im}(z_n)) \xi_I \xi_I^*,$$

where $r' = (r_1, \dots, r_{p-1}) = (|z_1|, \dots, |z_{p-1}|)$.

Quasi-hyperbolic group of biholomorphisms of the Siegel domain $\mathbb{U}^{p|q}$ is isomorphic to $\mathbb{T}^{p-1} \times \mathbb{R}_+ \times \mathbb{T}^q$ with the following group action:

$$(t, r, s) : (z', z_p, \theta) \in \mathbb{U}^{p|q} \mapsto (r^{1/2}tz', rz_p, r^{1/2}s\theta) \in \mathbb{U}^{p|q},$$

for each $(t, r, s) \in \mathbb{T}^{p-1} \times \mathbb{R}_+ \times \mathbb{T}^q$.

We now have that, if the super function F is invariant under the action of the quasi-hyperbolic group, then

$$F(z, \xi) = \sum_{I \subset Q} f_I(\rho_1, \dots, \rho_{n-1}, \arg(z_n - i|z'|^2)) |z_n - i|z'|^2|^{-|I|} \xi_I \xi_I^*,$$

where $z' = (z_1, \dots, z_{p-1})$ and

$$\rho_k = \frac{|z_k|}{\sqrt{|z'|^2 + |z_n - i|z'|^2|}},$$

for $k = 1, \dots, n - 1$.

Nilpotent group of biholomorphisms of the Siegel domain $\mathbb{U}^{p|q}$ is isomorphic to $\mathbb{R}^{p-1} \times \mathbb{R} \times \mathbb{T}^q$ with the following group action:

$$(b, h, s) : (z', z_p, \theta) \in \mathbb{U}^{p|q} \mapsto (z' + b, z_p + h + 2iz' \cdot b + i|b|^2, s\theta) \in \mathbb{U}^{p|q},$$

for each $(b, h, s) \in \mathbb{R}^{p-1} \times \mathbb{R} \times \mathbb{T}^q$.

In this case, we have that if the super function F is invariant under the action of the nilpotent group, then

$$F(z, \xi) = \sum_{I \subset Q} f_I(\text{Im } z', \text{Im } z_n - |z'|^2) \xi_I \xi_I^*,$$

where $z' = (z_1, \dots, z_{p-1})$.

Quasi-nilpotent group of biholomorphisms of the Siegel domain $\mathbb{U}^{p|q}$ is isomorphic to $\mathbb{T}^k \times \mathbb{R}^{p-k-1} \times \mathbb{R} \times \mathbb{T}^q$ where $0 < k < p - 1$, with the following group action:

$$(t, b, h, s) : (z', z'', z_p, \theta) \in \mathbb{U}^{p|q} \mapsto (tz', z'' + b, z_p + h + 2iz' \cdot b + i|b|^2, s\theta) \in \mathbb{U}^{p|q},$$

for each $(t, b, h, s) \in \mathbb{T}^k \times \mathbb{R}^{p-k-1} \times \mathbb{R} \times \mathbb{T}^q$.

And for this case, we have that if the super function F is invariant under the action of the quasi-nilpotent group, then

$$F(z, \xi) = \sum_{I \subset Q} f_I(r, y', \text{Im } z_n - |z'|^2) \xi_I \xi_I^*,$$

where $r = (|z_1|, \dots, |z_k|)$, $y' = \text{Im } w'$, and $w' = (z_{k+1}, \dots, z_{p-1})$.

4. Super Bargmann Transform

In [9] it was introduced a Bargmann type transform for each of the five cases on the unit ball. These Bargmann transforms are used in [9] to provide very useful descriptions of the Bergman spaces in terms of coordinates corresponding to the actions of maximal abelian subgroups. In this section we define Bargmann type transforms corresponding to the even MASA's considered above. These transforms are natural analogues of those defined in [9]

4.1. Quasi-Elliptic Case

Denote by $\tau(\mathbb{B}^p)$ the base of the unit ball \mathbb{B}^p , considered as a Reinhardt domain. In other words, we have

$$\tau(\mathbb{B}^p) = \{r = (r_1, \dots, r_p) = (|z_1|, \dots, |z_p|) : r^2 = r_1^2 + \dots + r_n^2 \in [0, 1]\},$$

which is contained in \mathbb{R}_+^p . Consider in \mathbb{C}^p the polar coordinates $z_k = t_k r_k$, where $r_k \in \mathbb{R}_+$ and $t_k \in \mathbb{T}$, for $k = 1, \dots, p$. Then, with respect to the identification

$$z = (z_1, \dots, z_p) = (t_1 r_1, \dots, t_p r_p) = (t, r),$$

where $t = (t_1, \dots, t_p) \in \mathbb{T}^p, r = (r_1, \dots, r_p) \in \tau(\mathbb{B}^p)$, we have $\mathbb{B}^p = \mathbb{T}^p \times \tau(\mathbb{B}^p)$, and

$$L_2(\mathbb{B}^p, \mu_\nu) = L_2(\mathbb{T}^p) \otimes L_2(\tau(\mathbb{B}^p), \mu_\nu),$$

where

$$L_2(\mathbb{T}^p) = \bigotimes_{k=1}^p L_2\left(\mathbb{T}, \frac{dt_k}{it_k}\right),$$

and the measure $d\mu_\nu(r)$ in $L_2(\tau(\mathbb{B}^p), \mu_\nu)$ is given by

$$d\mu_\nu(r) = \frac{\Gamma(\nu)}{\pi^p \Gamma(\nu - p)} (1 - r^2)^{\nu - p - 1} r dr.$$

We denote the discrete Fourier transform $\mathcal{F} : L_2(\mathbb{T}) \mapsto l_2 = l_2(\mathbb{Z})$ by

$$\mathcal{F} : f \mapsto c_n = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{T}} f(t) t^{-n} \frac{dt}{it}.$$

Of course, the operator \mathcal{F} is unitary and

$$\mathcal{F}^{-1} = \mathcal{F}^* : \{c_n\}_{n \in \mathbb{Z}} \mapsto f = \frac{1}{\sqrt{2\pi}} \sum_{n \in \mathbb{Z}} c_n t^n.$$

We consider the operator

$$U = \mathcal{F}_{(p)} \otimes I : L_2(\mathbb{T}^p) \otimes L_2(\mathbb{B}^p, \mu_\nu) \mapsto l_2(\mathbb{Z}^p) \otimes L_2(\mathbb{B}^p, \mu_\nu),$$

where $\mathcal{F}_{(p)} = \mathcal{F} \otimes \dots \otimes \mathcal{F}$. As in [9], we use the isometric embedding

$$R_{0,\nu} : l_2(\mathbb{Z}_+^p) \longrightarrow l_2(\mathbb{Z}^p) \otimes L_2(\mathbb{B}^p, \mu_\nu),$$

defined by

$$R_{0,\nu} : \{c_n\}_{n \in \mathbb{Z}_+^p} \mapsto c_n(r) = \begin{cases} \left(\frac{(2\pi)^p \Gamma(|n| + \nu)}{n! \Gamma(\nu)}\right)^{1/2} c_n r^n, & \text{for } n \in \mathbb{Z}_+^p \\ 0, & \text{for } n \in \mathbb{Z}^p \setminus \mathbb{Z}_+^p \end{cases}.$$

Hence, it is easily seen that the map

$$R_{0,\nu}^* : l_2(\mathbb{Z}^p) \otimes L_2(\mathbb{B}^p, \mu_\nu) \longrightarrow l_2(\mathbb{Z}_+^p)$$

satisfies

$$R_{0,\nu}^* : \{f_n(r)\}_{n \in \mathbb{Z}_+^p} \mapsto \left\{ \left(\frac{(2\pi)^p \Gamma(|n| + \nu)}{n! \Gamma(\nu)}\right)^{1/2} \int_{\tau(\mathbb{B}^p)} f_n(r) r^n d\mu_\nu \right\}_{\mathbb{Z}_+^p}.$$

In [9] the authors introduced the operator $R_\nu = R_{0,\nu}^* U$ from $L_2(\mathbb{B}^p, \mu_\nu)$ onto $l_2(\mathbb{Z}_+^p)$ and the adjoint operator $R_\nu^* = U^* R_{0,\nu}$ from $l_2(\mathbb{Z}_+^p)$ onto $H_\nu^2(\mathbb{B}^p)$. They proved that R_ν^* is the isometric isomorphism from $l_2(\mathbb{Z}_+^p)$ onto the subspace $H_\nu^2(\mathbb{B}^p)$.

Furthermore, we have

$$(4.1) \quad \begin{aligned} R_\nu R_\nu^* &= I : l_2(\mathbb{Z}_+^p) \rightarrow l_2(\mathbb{Z}_+^p), \\ R_\nu^* R_\nu &= P_{\mathbb{B}^p, \nu} : L_2(\mathbb{B}^p, \mu_\nu) \rightarrow H_\nu^2(\mathbb{B}^p), \end{aligned}$$

where $P_{\mathbb{B}^p, \nu}$ is the Bergman projection.

The inner product in $(l_2(\mathbb{Z}_+^p))^{2^q}$ is given by

$$\begin{aligned} &\langle \{a_{M,n}\}_{n \in \mathbb{Z}_+^p}\rangle_{M \subset Q}, \langle \{b_{M,n}\}_{n \in \mathbb{Z}_+^p}\rangle_{M \subset Q} \\ &= \sum_{M \subset Q} \frac{\Gamma(\nu)}{\Gamma(\nu + |M|)} \left(\sum_{n \in \mathbb{Z}_+^p} a_{M,n} \overline{b_{M,n}} \right). \end{aligned}$$

Definition 4.1. Consider the operator

$$R_{\nu, (p|q)} : H_\nu^2(\mathbb{B}^{p|q}) \rightarrow (l_2(\mathbb{Z}_+^p))^{2^q},$$

defined by

$$R_{\nu, (p|q)} \left(\sum_{M \subset Q} \psi_M \xi_M \right) = (R_{\nu+|M|}(\psi_M))_{M \subset Q},$$

whose explicit expression is given by

$$\begin{aligned} &R_{\nu, (p|q)} \left(\sum_{M \subset Q} \psi_M \xi_M \right) \\ &= \left(\left\{ (2\pi)^{-\frac{p}{2}} \left(\frac{(2\pi)^p \Gamma(|n| + \nu + |M|)}{n! \Gamma(\nu + |M|)}\right)^{1/2} \int_{\mathbb{B}^p} \psi_M \overline{z^n} d\mu_{\nu+|M|}(z) \right\}_{\mathbb{Z}_+^p} \right)_{M \subset Q}. \end{aligned}$$

The adjoint operator

$$R_{\nu, (p|q)}^* : (l_2(\mathbb{Z}_+^p))^{2^q} \rightarrow H_\nu^2(\mathbb{B}^{p|q})$$

is defined by

$$R_{\nu, (p|q)}^* ((\{c_{M,n}\}_{n \in \mathbb{Z}_+^p})_{M \subset Q}) = \sum_{M \subset Q} R_{\nu+|M|}^* (\{c_{M,n}\}_{n \in \mathbb{Z}_+^p}) \xi_M,$$

whose explicit expression is now

$$R_{\nu, (p|q)}^* ((\{c_{M,n}\}_{n \in \mathbb{Z}_+^p})_{M \subset Q}) = \sum_{M \subset Q} \left((2\pi)^{-\frac{p}{2}} \sum_{n \in \mathbb{Z}_+^p} \left(\frac{(2\pi)^p \Gamma(|n| + \nu + |M|)}{n! \Gamma(\nu)} \right)^{1/2} c_{M,n} z^n \xi_M \right).$$

Theorem 4.2. *The operators*

$$R_{\nu, (p|q)} : H_\nu^2(\mathbb{B}^{p|q}) \rightarrow (l_2(\mathbb{Z}_+^p))^{2^q}$$

and

$$R_{\nu, (p|q)}^* : (l_2(\mathbb{Z}_+^p))^{2^q} \rightarrow H_\nu^2(\mathbb{B}^{p|q})$$

are isometric isomorphisms. Furthermore, we have

$$R_{\nu, (p|q)} R_{\nu, (p|q)}^* = I : (l_2(\mathbb{Z}_+^p))^{2^q} \rightarrow (l_2(\mathbb{Z}_+^p))^{2^q},$$

$$R_{\nu, (p|q)}^* R_{\nu, (p|q)} = I : H_\nu^2(\mathbb{B}^{p|q}) \rightarrow H_\nu^2(\mathbb{B}^{p|q}).$$

Proof. First, we will prove that $R_{\nu, (p|q)}$ is a unitary operator. We first take an element $(\{c_{M,n}\}_{n \in \mathbb{Z}_+^p})_{M \subset Q}$ in $(l_2(\mathbb{Z}_+^p))^{2^q}$, then

$$\begin{aligned} I &= \|R_{\nu, (p|q)}^* ((\{c_{M,n}\}_{n \in \mathbb{Z}_+^p})_{M \subset Q})\|_{\mathbb{B}^{p|q}, \nu}^2 \\ &= \left(\sum_{M \subset Q} R_{\nu+|M|} (\{c_{M,n}\}_{n \in \mathbb{Z}_+^p}) \xi_M, \sum_{M \subset Q} R_{\nu+|M|} (\{c_{M,n}\}_{n \in \mathbb{Z}_+^p}) \xi_M \right)_{\mathbb{B}^{p|q}, \nu} \\ &= \sum_{m=1}^q \frac{\Gamma(\nu)}{\Gamma(\nu+m)} \sum_{M \subset Q, |M|=m} \|R_{\nu+|M|} (\{c_{M,n}\}_{n \in \mathbb{Z}_+^p})\|_{\mathbb{U}^{\nu+m}}^2. \end{aligned}$$

By (4.1) we know that R_ν is an isometric isomorphism, then

$$I = \sum_{M \subset Q} \frac{\Gamma(\nu)}{\Gamma(\nu+|M|)} \sum_{n \in \mathbb{Z}_+^p} |c_{M,n}|^2 = \|(\{c_{M,n}\}_{n \in \mathbb{Z}_+^p})_{M \subset Q}\|_{(l_2(\mathbb{Z}_+^p))^{2^q}}^2,$$

which proves our claim. □

4.2. Quasi-Parabolic Case

We define $\mathbb{D}^p = \mathbb{C}^{p-1} \times \mathbb{R} \times \mathbb{R}_+$ whose points we denote by (z', u, v) . Consider the space $L_2(\mathbb{D}^p, \eta_\nu)$, where η_ν is given by

$$\eta_\nu(z', u, v) = \frac{\Gamma(\nu)}{4\pi^p \Gamma(\nu-p)} v^{\nu-p-1}.$$

Consider the operator $U_0 : L_2(\mathbb{B}^p, \mu_\nu) \rightarrow L_2(\mathbb{D}^p, \eta_\nu)$ defined by

$$U_0(f)(z', u, v) = f(\kappa(z', u, v)),$$

where $\kappa(z', u, v) = (z', u + iv + i|w'|)$. It is clear that the operator is unitary and the inverse operator is given by

$$U_0^{-1}(f)(z) = f(\kappa^{-1}(z)).$$

We represent the space $L_2(\mathbb{D}^p, \eta_\nu)$ as the following tensor product

$$L_2(\mathbb{D}^p, \eta_\nu) = L_2(\mathbb{C}^{p-1}) \otimes L_2(\mathbb{R}) \otimes L_2(\mathbb{R}_+, \eta_\nu),$$

and consider the unitary operator $U_1 = I \otimes F \otimes I$ acting on it. Here F is the standard Fourier transform on $L_2(\mathbb{R})$ and $f(z', u, v) \mapsto U_1 f(z', \xi, v)$.

On the other hand, we have the decomposition

$$L_2(\mathbb{D}^p, \eta_\nu) = L_2(\mathbb{R}_+^{p-1}) \otimes L_2(\mathbb{T}^{p-1}) \otimes L_2(\mathbb{R}) \otimes L_2(\mathbb{R}_+, \eta_\nu),$$

where

$$L_2(\mathbb{T}^{p-1}) = \bigotimes_{k=1}^{p-1} L_2\left(\mathbb{T}, \frac{dt_k}{it_k}\right).$$

As before, we consider the discrete Fourier transform $\mathcal{F} : L_2(\mathbb{T}^p) \mapsto l_2 = l_2(\mathbb{Z})$, and consider the unitary operator $U_2 = I \otimes \mathcal{F}_{(p-1)} \otimes I \otimes I$ acting from

$$L_2(\mathbb{R}_+^{p-1}) \otimes L_2(\mathbb{T}^{p-1}) \otimes L_2(\mathbb{R}) \otimes L_2(\mathbb{R}_+, \eta_\nu)$$

onto

$$\begin{aligned} &L_2(\mathbb{R}_+^{p-1}) \otimes l_2(\mathbb{Z}^{p-1}) \otimes L_2(\mathbb{R}) \otimes L_2(\mathbb{R}_+, \eta_\nu) \\ &= l_2(\mathbb{Z}^{p-1}, L_2(\mathbb{R}_+^{p-1})) \otimes L_2(\mathbb{R}) \otimes L_2(\mathbb{R}_+, \eta_\nu), \end{aligned}$$

where $\mathcal{F}_{(p-1)} = \mathcal{F} \otimes \dots \otimes \mathcal{F}$ and $f(z', \xi, v) \mapsto \{U_2 f(r, n, \xi, v)\}_{n \in \mathbb{Z}^{p-1}}$

We now consider isometric embedding

$$R_{\nu,0} : l_2(\mathbb{Z}_+^{p-1}, L_2(\mathbb{R}_+)) \rightarrow l_2(\mathbb{Z}^{p-1}, L_2(\mathbb{R}_+^{p-1})) \otimes L_2(\mathbb{R}) \otimes L_2(\mathbb{R}_+, \eta_\nu)$$

defined by the assignment

$$\begin{aligned} &R_{\nu,0} \{b_n(\xi)\}_{n \in \mathbb{Z}_+^{p-1}} \\ &= \left\{ \chi_{\mathbb{Z}_+^{p-1}}(n) \chi_{\mathbb{R}_+}(\xi) \left(\frac{2(2\pi)^p (2\xi)^{|n|+\nu-1}}{n! \Gamma(\nu)} \right)^{\frac{1}{2}} r^n e^{\xi(|z'|^2+v)} b_n(\xi) \right\}_{n \in \mathbb{Z}_+^{p-1}}, \end{aligned}$$

where the function $b_n(\xi)$ is extended by zero for $\xi \in \mathbb{R} \setminus \mathbb{R}_+$ and $n \in \mathbb{Z}^{p-1} \setminus \mathbb{Z}_+^{p-1}$. Its adjoint

$$R_{\nu,0}^* : l_2(\mathbb{Z}^{p-1}, L_2(\mathbb{R}_+^{p-1})) \otimes L_2(\mathbb{R}) \otimes L_2(\mathbb{R}_+, \eta_\nu) \rightarrow l_2(\mathbb{Z}_+^{p-1}, L_2(\mathbb{R}_+))$$

is given by the expression

$$R_{\nu,0}^* : \{d_n(r, \xi, v)\}_{n \in \mathbb{Z}^{p-1}}$$

$$= \left\{ \chi_{\mathbb{R}_+}(\xi) \left(\frac{2(2\pi)^p (2\xi)^{|n|+\nu-1}}{n! \Gamma(\nu)} \right)^{\frac{1}{2}} \int_{\mathbb{R}_+^n} r^n e^{\xi(|z'|^2+v)} d_n(r, \xi, v) r dr \frac{c_\nu v^{\nu-p-1}}{4} dv \right\}_{n \in \mathbb{Z}_+^{p-1}}.$$

In [9] the authors introduced the operator $R_\nu = R_{0,\nu}^* U$ from $L_2(\mathbb{U}^p, \mu_\nu)$ onto $l_2(\mathbb{Z}_+^{p-1}, L_2(\mathbb{R}_+))$, and the adjoint operator $R_\nu^* = U^* R_{0,\nu}$ from $l_2(\mathbb{Z}_+^{p-1},$

$L_2(\mathbb{R}_+)$ onto $H_\nu^2(\mathbb{U}^p)$ where $U = U_2U_1U_0$. They proved that R_ν^* is the isometric isomorphism of $l_2(\mathbb{Z}_+^{p-1}, L_2(\mathbb{R}_+))$ onto the subspace $H_\nu^2(\mathbb{U}^p)$. Furthermore, we have

$$R_\nu R_\nu^* = I : l_2(\mathbb{Z}_+^{p-1}, L_2(\mathbb{R}_+)) \rightarrow l_2(\mathbb{Z}_+^{p-1}, L_2(\mathbb{R}_+))$$

$$R_\nu^* R_\nu = P_{\mathbb{U}^p, \nu} : L_2(\mathbb{U}^p, \mu_\nu) \rightarrow H_\nu^2(\mathbb{U}^p),$$

where $P_{\mathbb{U}^p, \nu}$ is the Bergman projection of $L_2(\mathbb{U}^p, \mu_\nu)$ to $H_\nu^2(\mathbb{U}^p)$.

The inner product in $(l_2(\mathbb{Z}_+^{p-1}, L_2(\mathbb{R}_+)))^{2^q}$ is given by

$$\langle (\{a_{M,n}(\xi)\}_{n \in \mathbb{Z}_+^{p-1}})_{M \subset Q}, (\{b_{M,n}(\xi)\}_{n \in \mathbb{Z}_+^{p-1}})_{M \subset Q} \rangle$$

$$= \sum_{M \subset Q} \frac{\Gamma(\nu)}{\Gamma(\nu + |M|)} \left(\sum_{n \in \mathbb{Z}_+^{p-1}} ((a_{M,n}(\xi), b_{M,n}(\xi))_{L_2(\mathbb{R}_+)}) \right).$$

Definition 4.3. Consider the operator

$$R_{\nu, (p|q)} : H_\nu^2(\mathbb{U}^{p|q}) \rightarrow (l_2(\mathbb{Z}_+^{p-1}, L_2(\mathbb{R}_+)))^{2^q}$$

defined by

$$R_{\nu, (p|q)} \left(\sum_{M \subset Q} \psi_M \xi_M \right) = (R_{\nu+|M|}(\psi_M))_{M \subset Q},$$

whose explicit expression is given by

$$R_{\nu, (p|q)} \left(\sum_{M \subset Q} \psi_M \xi_M \right)$$

$$= \left(\left\{ (2\pi)^{-\frac{p}{2}} \left(\frac{2(2\pi)^p (2\xi)^{|n|+\nu+|M|-1}}{n! \Gamma(\nu + |M|)} \right)^{\frac{1}{2}} \right. \right.$$

$$\left. \times \int_{\mathbb{U}^p} \psi_M(z) (\bar{z}')^n e^{-i\xi z_n} (\text{Im} z_n - |z'|)^{\nu+|M|-p-1} dv(z) \right\}_{\mathbb{Z}_+^{p-1}} \right)_{M \subset Q}.$$

Its adjoint operator is

$$R_{\nu, (p|q)}^* : (l_2(\mathbb{Z}_+^{p-1}, L_2(\mathbb{R}_+)))^{2^q} \rightarrow H_\nu^2(\mathbb{U}^{p|q})$$

defined by

$$R_{\nu, (p|q)}^* ((\{b_{M,n}(\xi)\}_{n \in \mathbb{Z}_+^{p-1}})_{M \subset Q}) = \sum_{M \subset Q} R_{\nu+|M|}(\{b_{M,n}(\xi)\}_{n \in \mathbb{Z}_+^{p-1}}) \xi_M,$$

whose explicit expression is now given by

$$R_{\nu, (p|q)}^* ((\{b_{M,n}(\xi)\}_{n \in \mathbb{Z}_+^{p-1}})_{M \subset Q})$$

$$= \sum_{M \subset Q} \left((2\pi)^{-\frac{p}{2}} \sum_{n \in \mathbb{Z}_+^{p-1}} \int_{\mathbb{R}_+} \left(\frac{2(2\pi)^p (2\xi)^{|n|+\nu+|M|-1}}{n! \Gamma(\nu + |M|)} \right)^{\frac{1}{2}} b_{n, M}(\xi) (z')^n e^{i\xi z_n} d\xi \right) \xi_M.$$

Theorem 4.4. *The operators*

$$R_{\nu,(p|q)} : H_{\nu}^2(\mathbb{U}^{p|q}) \rightarrow (l_2(\mathbb{Z}_+^{p-1}, L_2(\mathbb{R}_+)))^{2^q}$$

and

$$R_{\nu,(p|q)}^* : (l_2(\mathbb{Z}_+^{p-1}, L_2(\mathbb{R}_+)))^{2^q} \rightarrow H_{\nu}^2(\mathbb{U}^{p|q})$$

are isometric isomorphisms. Furthermore, we have

$$R_{\nu,(p|q)} R_{\nu,(p|q)}^* = I : (l_2(\mathbb{Z}_+^{p-1}, L_2(\mathbb{R}_+)))^{2^q} \rightarrow (l_2(\mathbb{Z}_+^{p-1}, L_2(\mathbb{R}_+)))^{2^q}$$

$$R_{\nu,(p|q)}^* R_{\nu,(p|q)} = I : H_{\nu}^2(\mathbb{U}^{p|q}) \rightarrow H_{\nu}^2(\mathbb{U}^{p|q}).$$

The proof of this result is similar to that of Theorem 4.2.

4.3. Nilpotent Case

We consider the space

$$L_2(\mathbb{D}^p, \eta_{\nu}) = L_2(\mathbb{C}^{p-1}) \otimes L_2(\mathbb{R}) \otimes L_2(\mathbb{R}_+, \eta_{\nu}),$$

and the unitary operator $U_1 = I \otimes F \otimes I$ acting on it, where F is the Fourier transform on $L_2(\mathbb{R})$.

Using the standard Cartesian coordinates $x' = (x_1, \dots, x_{p-1})$ and $y' = (y_1, \dots, y_{p-1})$, where $z_k = x_k + iy_k$, in $\mathbb{C}^{p-1} = \mathbb{R}^{p-1} \times \mathbb{R}^{p-1}$, we have

$$L_2(\mathbb{D}^p, \eta_{\nu}) = L_2(\mathbb{R}^{p-1}) \otimes L_2(\mathbb{R}^{p-1}) \otimes L_2(\mathbb{R}) \otimes L_2(\mathbb{R}_+, \eta_{\nu}).$$

Consider the unitary operator $U_2 = F_{(p-1)} \otimes I \otimes I \otimes I$, where $F_{(p-1)} = F_1 \otimes \dots \otimes F_{p-1}$ is $(p-1)$ -dimensional Fourier transform, acting on this tensor decomposition.

Consider the change of variables

$$u_k = \frac{1}{2\sqrt{\xi}} \xi_k - \sqrt{\xi} y_k, \quad v_k = \frac{1}{2\sqrt{\xi}} \xi_k + \sqrt{\xi} y_k, \quad k = 1, \dots, p-1,$$

which is equivalent to

$$\xi_k = \sqrt{\xi} (u_k + v_k), \quad y_k = \frac{1}{2\sqrt{\xi}} (-u_k + v_k), \quad k = 1, \dots, p-1,$$

and the corresponding unitary operator U_3 acting on $L_2(\mathbb{R}^{p-1}) \otimes L_2(\mathbb{R}^{p-1}) \otimes L_2(\mathbb{R}) \otimes L_2(\mathbb{R}_+, \eta_{\nu})$ by the rule

$$(U_3 \varphi)(u', v', \xi, v) = \varphi \left(\sqrt{\xi} (u' + v'), \frac{1}{2\sqrt{\xi}} (-u' + v'), \xi, v \right),$$

where $u' = (u_1, \dots, u_{p-1})$ and $v' = (v_1, \dots, v_{p-1})$.

In this case, we consider the isometric embedding

$$R_{0,\nu} : L_2(\mathbb{R}^{p-1} \times \mathbb{R}_+) \longrightarrow L_2(\mathbb{R}^{p-1}) \otimes L_2(\mathbb{R}^{p-1}) \otimes L_2(\mathbb{R}) \otimes L_2(\mathbb{R}_+, \eta_{\nu})$$

given by the assignment

$$R_{0,\nu}(\psi(u', \xi)) = \pi^{-\frac{p-1}{4}} e^{-\xi v - \frac{|v'|^2}{2}} \chi_{\mathbb{R}_+}(\xi) \left(\frac{4(2\xi)^{\nu-p}}{c_{\nu} \Gamma(\nu-p)} \right)^{\frac{1}{2}} \psi(u', \xi),$$

where the function $\psi(u', \xi)$ is extended by zero for $\xi \in \mathbb{R} \setminus \mathbb{R}_+$ for each $u' \in \mathbb{R}^{p-1}$. The adjoint operator

$$R_{0,\nu}^* : L_2(\mathbb{R}^{p-1}) \otimes L_2(\mathbb{R}^{p-1}) \otimes L_2(\mathbb{R}) \otimes L_2(\mathbb{R}_+, \eta_{\nu}) \longrightarrow L_2(\mathbb{R}^{p-1} \times \mathbb{R}_+)$$

obviously has the form

$$R_{0,\nu}^*(\varphi(u', v', \xi, v)) = \pi^{-\frac{p-1}{4}} \int_{\mathbb{R}^{p-1} \times \mathbb{R}_+} e^{-\xi v - \frac{|v'|^2}{2}} \left(\frac{4(2\xi)^{\nu-p}}{c_\nu \Gamma(\nu-p)} \right)^{\frac{1}{2}} f(u', v', \xi, v) dv' \frac{c_\nu}{4} v^{\nu-p-1} dv.$$

In [9] the authors introduced the operator $R_\nu = R_{0,\nu}^*U$ from $L_2(\mathbb{D}^p, \tilde{\mu}_\nu)$ onto $L_2(\mathbb{R}^{p-1} \times \mathbb{R}_+)$ and the adjoint operator R_ν^* from $L_2(\mathbb{R}^{p-1} \times \mathbb{R}_+)$ onto the subspace $\mathcal{A}_\nu^2(\mathbb{D}^p)$. They proved that R_ν^* is the isometric isomorphism of $L_2(\mathbb{R}^{p-1} \times \mathbb{R}_+)$ onto the subspace $H_\nu^2(\mathbb{U}^p)$. Furthermore

$$R_\nu R_\nu^* = I : L_2(\mathbb{R}^{p-1} \times \mathbb{R}_+) \longrightarrow L_2(\mathbb{R}^{p-1} \times \mathbb{R}_+),$$

$$R_\nu^* R_\nu = B_{\mathbb{D}^p, \nu} : L_2(\mathbb{D}^p, \tilde{\mu}_\nu) \longrightarrow \mathcal{A}_\nu^2(\mathbb{D}^p),$$

where $B_{\mathbb{D}^p, \nu}$ is the Bergman projection.

Now we define the analogue of the Bargmann transform for the super case.

Definition 4.5. Consider the operator

$$R_{\nu, (p|q)} : H_\nu^2(\mathbb{U}^{p|q}) \rightarrow (L_2(\mathbb{R}^{p-1} \times \mathbb{R}_+))^{2^q}$$

defined by

$$R_{\nu, (p|q)} \left(\sum_{M \subset Q} \psi_M \xi_M \right) = (R_{\nu+|M|}(\psi_M))_{M \subset Q}.$$

Whose adjoint operator

$$R_{\nu, (p|q)}^* : (L_2(\mathbb{R}^{p-1} \times \mathbb{R}_+))^{2^q} \rightarrow H_\nu^2(\mathbb{U}^{p|q})$$

is defined by

$$R_{\nu, (p|q)}^*(\psi_M(u', \xi))_{M \subset Q} = \sum_{M \subset Q} R_{\nu+|M|}(\psi_M(u', \xi)) \xi_M.$$

The inner product in $(L_2(\mathbb{R}^{p-1} \times \mathbb{R}_+))^{2^q}$ is given by

$$\begin{aligned} &\langle (\psi_M(u', \xi))_{M \subset Q}, (\phi_M(u', \xi))_{M \subset Q} \rangle \\ &= \sum_{M \subset Q} \frac{\Gamma(\nu)}{\Gamma(\nu + |M|)} (\psi_M(u', \xi), \phi_M(u', \xi))_{L_2(\mathbb{R}_+)}. \end{aligned}$$

Theorem 4.6. *The operators*

$$R_{\nu, (p|q)} : H_\nu^2(\mathbb{U}^{p|q}) \rightarrow (L_2(\mathbb{R}^{p-1} \times \mathbb{R}_+))^{2^q}$$

and

$$R_{\nu, (p|q)}^* : (L_2(\mathbb{R}^{p-1} \times \mathbb{R}_+))^{2^q} \rightarrow H_\nu^2(\mathbb{U}^{p|q})$$

are isometric isomorphisms. Furthermore

$$R_{\nu, (p|q)} R_{\nu, (p|q)}^* = I : (L_2(\mathbb{R}^{p-1} \times \mathbb{R}_+))^{2^q} \rightarrow (L_2(\mathbb{R}^{p-1} \times \mathbb{R}_+))^{2^q}$$

$$R_{\nu, (p|q)}^* R_{\nu, (p|q)} = I : H_\nu^2(\mathbb{U}^{p|q}) \rightarrow H_\nu^2(\mathbb{U}^{p|q}).$$

The proof is similar to that of Theorem 4.2.

4.4. Quasi-Nilpotent Case

Given an integer $1 \leq k \leq n - 2$, we will write points of \mathbb{D}^p as $z = (z', w', z_n)$, where $z' \in \mathbb{C}^k$ and $w' \in \mathbb{C}^{p-k-1}$, and points of \mathbb{D}^p as (z', w', ζ) , respectively.

According to this notation we represent

$$L_2(\mathbb{D}^p, \eta_\nu) = L_2(\mathbb{C}^k) \otimes L_2(\mathbb{C}^{p-k-1}) \otimes L_2(\mathbb{R}) \otimes L_2(\mathbb{R}_+, \eta_\nu).$$

Applying, as in the previous two cases, the unitary operator $U_1 = I \otimes I \otimes F \otimes I$, we have that the image $\mathcal{A}_1(\mathbb{D}^p) = U_1(\mathcal{A}_0(\mathbb{D}^p))$.

Now introducing in \mathbb{C}^k the polar coordinates, $z_l = r_l t_l$, where $r_l \in \mathbb{R}_+$, $t_l \in S^1 = \mathbb{T}$, $l = 1, \dots, k$, and Cartesian coordinates in \mathbb{C}^{p-k-1} , $x' = (x_1, \dots, x_{p-k-1})$, $y' = (y_1, \dots, y_{p-k-1})$, where $w_m = x_m + iy_m$, $m = 1, \dots, p - k - 1$, we have that $L_2(\mathbb{D}^p, \eta_\nu)$ decomposes as the tensor product

$$L_2(\mathbb{R}_+^k, r dr) \otimes L_2(\mathbb{T}^k) \otimes L_2(\mathbb{R}^{p-k-1}) \otimes L_2(\mathbb{R}^{p-k-1}) \otimes L_2(\mathbb{R}) \otimes L_2(\mathbb{R}_+, \eta_\nu).$$

Consider the unitary operator $U_2 = I \otimes \mathcal{F}_{(k)} \otimes F_{(p-k-1)} \otimes I \otimes I \otimes I$ acting from $L_2(\mathbb{D}^p, \eta_\nu)$ onto

$$\begin{aligned} &L_2(\mathbb{R}_+^k, r dr) \otimes l_2(\mathbb{Z}^k \otimes L_2(\mathbb{R}^{p-k-1}) \otimes L_2(\mathbb{R}^{p-k-1}) \otimes L_2(\mathbb{R}) \otimes L_2(\mathbb{R}_+, \eta_\nu)) \\ &= l_2(\mathbb{Z}^k, L_2(\mathbb{R}_+^k, r dr) \otimes L_2(\mathbb{R}^{p-k-1}) \otimes L_2(\mathbb{R}^{p-k-1}) \otimes L_2(\mathbb{R}) \otimes L_2(\mathbb{R}_+, \eta_\nu)), \end{aligned}$$

where $\mathcal{F}_{(k)} = \mathcal{F} \otimes \dots \otimes \mathcal{F}$ is the k -dimensional discrete Fourier transform and $F_{(p-k-1)} = F \otimes \dots \otimes F$ is the $(p - k - 1)$ -dimensional Fourier transform.

Next consider the change of variables

$$u_m = \frac{1}{2\sqrt{\xi}} \xi_m - \sqrt{\xi} y_m, \quad v_m = \frac{1}{2\sqrt{\xi}} \xi_m + \sqrt{\xi} y_m, \quad m = 1, \dots, p - k - 1,$$

which is equivalent to

$$\xi_m = \sqrt{\xi} (u_m + v_m), \quad y_m = \frac{1}{2\sqrt{\xi}} (-u_m + v_m), \quad m = 1, \dots, p - k - 1.$$

Then, there is a corresponding unitary operator U_3 acting on

$$l_2(\mathbb{Z}^k, L_2(\mathbb{R}_+^k, r dr) \otimes L_2(\mathbb{R}^{p-k-1}) \otimes L_2(\mathbb{R}^{p-k-1}) \otimes L_2(\mathbb{R}) \otimes L_2(\mathbb{R}_+, \eta_\nu))$$

by the assignment

$$U_3 \{d_p(r, \xi', y', \xi, v)\}_{p \in \mathbb{Z}^k} = \left\{ d_p \left(r, \sqrt{\xi} (u' + v'), \frac{1}{2\sqrt{\xi}} (-u' + v'), \xi, v \right) \right\}_{p \in \mathbb{Z}^k},$$

where $u' = (u_1, \dots, u_{p-k-1})$ and $v' = (v_1, \dots, v_{p-k-1})$.

In [9] it is introduced the isometric imbedding $R_{0,\nu}$ from the Hilbert space $l_2(\mathbb{Z}_+^k, L_2(\mathbb{R}^{p-k-1} \times \mathbb{R}_+))$ into

$$l_2(\mathbb{Z}^k, (L_2(\mathbb{R}_+^k, r dr) \otimes L_2(\mathbb{R}^{p-k-1}) \otimes L_2(\mathbb{R}^{p-k-1}) \otimes L_2(\mathbb{R}) \otimes L_2(\mathbb{R}_+, \eta_\nu)),$$

which maps $\{c_n(u', \xi)\}_{n \in \mathbb{Z}_+^k}$ to

$$\left\{ \pi^{-\frac{p-k-1}{4}} \chi_{\mathbb{Z}_+^k}(n) \chi_{\mathbb{R}_+}(\xi) \left(\frac{2^{k+2} (2\xi)^{|n|+\nu-p+k}}{c_\nu n! \Gamma(\nu-p)} \right)^{\frac{1}{2}} r^\nu e^{-\xi(|r|^2+v)} - \frac{|v'|^2}{2} c_n(u', \xi) \right\}_{n \in \mathbb{Z}^k},$$

where the functions $c_n(u', \xi)$ are extended by zero for $\xi \in \mathbb{R} \setminus \mathbb{R}_+$ for each $u' \in \mathbb{R}^{p-k-1}$ and each $n \in \mathbb{Z}^k$.

The adjoint operator $R_{0,\nu}^*$ acts from

$l_2(\mathbb{Z}^k, (L_2(\mathbb{R}_+^k, r dr) \otimes L_2(\mathbb{R}^{p-k-1}) \otimes L_2(\mathbb{R}^{p-k-1}) \otimes L_2(\mathbb{R}) \otimes L_2(\mathbb{R}_+, \eta_\nu))$
 onto $l_2(\mathbb{Z}_+^k, L_2(\mathbb{R}^{p-k-1} \times \mathbb{R}_+))$ and maps a sequence $\{d_n(r, u', v', \xi, v)\}_{n \in \mathbb{Z}^k}$
 into

$$R_{0,\nu}^*(\{d_n(r, u', v', \xi, v)\}_{n \in \mathbb{Z}^k}) = \left\{ \pi^{-\frac{p-k-1}{4}} \left(\frac{2^{k+2}}{c_\nu} \frac{(2\xi)^{|n|+\nu-p+k}}{n! \Gamma(\nu-p)} \right)^{\frac{1}{2}} \int_{\mathbb{R}_+^k \times \mathbb{R}^{p-k-1} \times \mathbb{R}_+} r^n e^{-\xi(|r|^2+v)-\frac{|v'|^2}{2}} \times d_n(r, u', v', \xi, v) r dr dv' \frac{c_\nu v^{\nu-p-1}}{4} dv \right\}_{n \in \mathbb{Z}_+^k}.$$

In [9] the authors introduce the operator $R_\nu = R_{0,\nu}^*U$ from $L_2(\mathbb{D}^p, \tilde{\mu}_\nu)$ onto $l_2(\mathbb{Z}_+^k, L_2(\mathbb{R}^{p-k-1} \times \mathbb{R}_+))$, and the adjoint operator R_ν^* being the isometric isomorphism of $l_2(\mathbb{Z}_+^k, L_2(\mathbb{R}^{p-k-1} \times \mathbb{R}_+))$ onto the subspace $\mathcal{A}_\nu^2(\mathbb{D}^p)$ of $L_2(\mathbb{D}^p, \tilde{\mu}_\nu)$. Furthermore

$$R_\nu R_\nu^* = I : l_2(\mathbb{Z}_+^k, L_2(\mathbb{R}^{p-k-1} \times \mathbb{R}_+)) \longrightarrow l_2(\mathbb{Z}_+^k, L_2(\mathbb{R}^{p-k-1} \times \mathbb{R}_+)),$$

$$R_\nu^* R_\nu = B_{\mathbb{D}^p, \nu} : L_2(\mathbb{D}^p, \tilde{\mu}_\nu) \longrightarrow \mathcal{A}_\nu^2(\mathbb{D}^p),$$

where $B_{\mathbb{D}^p, \nu}$ is the Bergman projection.

Definition 4.7. Consider the operator

$$R_{\nu, (p|q)} : H_\nu^2(\mathbb{U}^{p|q}) \rightarrow (l_2(\mathbb{Z}_+^k, L_2(\mathbb{R}^{p-k-1} \times \mathbb{R}_+)))^{2^q}$$

defined by

$$R_{\nu, (p|q)} \left(\sum_{M \subset Q} \psi_M \xi_M \right) = (R_{\nu+|M|}(\psi_M))_{M \subset Q}.$$

The adjoint operator

$$R_{\nu, (p|q)}^* : (l_2(\mathbb{Z}_+^k, L_2(\mathbb{R}^{p-k-1} \times \mathbb{R}_+)))^{2^q} \rightarrow H_\nu^2(\mathbb{U}^{p|q})$$

is defined by

$$R_{\nu, (p|q)}^*(\{c_{M,n}(u', \xi)\}_{n \in \mathbb{Z}_+^k})_{M \subset Q} = \sum_{M \subset Q} R_{\nu+|M|}(\{c_{M,n}(u', \xi)\}_{n \in \mathbb{Z}_+^k})_{\xi_M},$$

where the inner product in $(l_2(\mathbb{Z}_+^k, L_2(\mathbb{R}^{p-k-1} \times \mathbb{R}_+)))^{2^q}$ is given by

$$\langle (\{a_{M,n}(u', \xi)\}_{n \in \mathbb{Z}_+^k})_{M \subset Q}, (\{b_{M,n}(u', \xi)\}_{n \in \mathbb{Z}_+^k})_{M \subset Q} \rangle = \sum_{M \subset Q} \frac{\Gamma(\nu)}{\Gamma(\nu+|M|)} \sum_{n \in \mathbb{Z}_+^k} (a_{M,n}(u', \xi), b_{M,n}(u', \xi))_{L_2(\mathbb{R}^{p-k-1} \times \mathbb{R}_+)}.$$

Theorem 4.8. The operators

$$R_{\nu, (p|q)} : H_\nu^2(\mathbb{U}^{p|q}) \rightarrow (l_2(\mathbb{Z}_+^k, L_2(\mathbb{R}^{p-k-1} \times \mathbb{R}_+)))^{2^q}$$

and

$$R_{\nu, (p|q)}^* : (l_2(\mathbb{Z}_+^k, L_2(\mathbb{R}^{p-k-1} \times \mathbb{R}_+)))^{2^q} \rightarrow H_\nu^2(\mathbb{U}^{p|q})$$

are isometric isomorphisms. Furthermore

$$\begin{aligned} R_{\nu,(p|q)}R_{\nu,(p|q)}^* &= I : (l_2(\mathbb{Z}_+^k, L_2(\mathbb{R}^{p-k-1} \times \mathbb{R}_+)))^{2q} \\ &\quad \rightarrow (l_2(\mathbb{Z}_+^k, L_2(\mathbb{R}^{p-k-1} \times \mathbb{R}_+)))^{2q} \\ R_{\nu,(p|q)}^*R_{\nu,(p|q)} &= I : H_\nu^2(\mathbb{U}^{p|q}) \rightarrow H_\nu^2(\mathbb{U}^{p|q}). \end{aligned}$$

The proof is similar to that of Theorem 4.2.

4.5. Quasi-Hyperbolic Case

We represent $\mathbb{D}^p = \mathbb{C}^{p-1} \times \mathbb{R} \times \mathbb{R}_+$ in the form $\mathbb{C}^{p-1} \times \Pi$, where Π is the upper half-plane, and introduce in \mathbb{D}^p the “non-isotropic” upper semi-sphere

$$\Omega = \{(z', \zeta) \in \mathbb{C}^{p-1} \times \Pi : |z'|^2 + |\zeta| = 1\}.$$

The points of Ω admit the natural parameterization

$$\begin{aligned} z_k &= s_k t_k, \quad \text{where } s_k \in [0, 1), \quad t_k \in S^1, \quad k = 1, \dots, p-1, \\ \zeta &= \rho e^{i\theta}, \quad \text{where } \rho \in (0, 1], \quad \theta \in (0, \pi), \end{aligned}$$

and

$$\sum_{k=1}^{p-1} s_k^2 + \rho = 1.$$

This in turn induces the following representation of the points $(z', \zeta) \in \mathbb{D}^p = \mathbb{C}^{p-1} \times \Pi$

$$z_k = r^{\frac{1}{2}} s_k t_k, \quad k = 1, \dots, p-1, \quad \zeta = r \rho e^{i\theta},$$

where $r \in \mathbb{R}_+$.

We represent now $\mathbb{D}^p = \tau(\mathbb{B}^{p-1}) \times \mathbb{T}^{p-1} \times \mathbb{R}_+ \times (0, \pi)$, where

$$\tau(\mathbb{B}^{p-1}) = \left\{ s = (s_1, \dots, s_{p-1}) \in \mathbb{R}_+^{p-1} : \sum_{k=1}^{p-1} s_k^2 < 1 \right\}$$

is the base (in the sense of a Reinhardt domain) of the unit ball \mathbb{B}^{p-1} , and $\mathbb{T}^{p-1} = S^1 \times \dots \times S^1$ is the $p-1$ dimensional torus.

Introduce the new coordinate system (s, t, r, θ) in \mathbb{D}^p , where we have $s = (s_1, \dots, s_{p-1}) \in \tau(\mathbb{B}^{p-1})$, $t = (t_1, \dots, t_{p-1}) \in \mathbb{T}^{p-1}$, $r \in \mathbb{R}_+$, and $\theta \in (0, \pi)$, which is connected with the old one (z', ζ) by the formulas

$$s_k = \frac{|z_k|}{\sqrt{|z'|^2 + |\rho|}}, \quad t_k = \frac{z_k}{|z_k|}, \quad r = |z'|^2 + |\rho|, \quad \theta = \arg \zeta, \quad (4.2)$$

or

$$z_k = r^{\frac{1}{2}} s_k t_k, \quad \zeta = r(1 - |s|^2)e^{i\theta},$$

where $k = 1, \dots, p-1$.

A direct computation shows that under the change of variables (4.2) we have

$$dv(z', \zeta) = r^p (1 - |s|^2) \prod_{k=1}^{p-1} s_k ds_k \prod_{k=1}^{p-1} \frac{dt_k}{it_k} dr d\theta,$$

and

$$\eta_\nu = \frac{C_\nu}{4} r^{\nu-p-1} (1 - |s|^2)^{\nu-p-1} \frac{C_\nu}{4} \sin^{\nu-p-1} \theta.$$

Introduce the unitary operator $U_1 = I \otimes \mathcal{F}_{(p-1)} \otimes M_\nu \otimes I$ which acts from the space

$$L_2(\tau(\mathbb{B}^{p-1}), (1 - |s|^2)^{\nu-p} s ds) \otimes L_2(\mathbb{T}^{p-1}) \otimes L_2(\mathbb{R}_+, r^{\nu-1} dr) \otimes L_2\left((0, \pi), \frac{c_\nu}{4} \sin^{\nu-p-1} \theta d\theta\right)$$

onto the space

$$l_2\left(\mathbb{Z}^{p-1}, L_2(\tau(\mathbb{B}^{p-1}), (1 - |s|^2)^{\nu-p} s ds) \otimes L_2(\mathbb{R}) \otimes L_2\left((0, \pi), \frac{c_\nu}{4} \sin^{\nu-p-1} \theta d\theta\right)\right),$$

where the Mellin transform $M_\nu : L_2(\mathbb{R}_+, r^{\nu-1} dr) \rightarrow L_2(\mathbb{R})$ is given by

$$(M_\nu \psi)(\xi) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}_+} r^{-i\xi + \frac{\nu}{2} - 1} \psi(r) dr,$$

and the inverse is as follows

$$(M_\nu^{-1} \psi)(r) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} r^{-i\xi - \frac{\nu}{2}} \psi(\xi) d\xi,$$

and $\mathcal{F}_{(p-1)} = \mathcal{F} \otimes \dots \otimes \mathcal{F}$ is the $(p-1)$ -dimensional discrete Fourier transform.

Introduce the isometric imbedding $R_{0,\nu}$ of the space $l_2(\mathbb{Z}_+^{p-1}, L_2(\mathbb{R}))$ into the space

$$l_2\left(\mathbb{Z}_+^{p-1}, L_2(\tau(\mathbb{B}^{p-1}), (1 - |s|^2)^{\nu-p} s ds) \otimes L_2(\mathbb{R}) \otimes L_2\left((0, \pi), \frac{c_\nu}{4} \sin^{\nu-p-1} \theta d\theta\right)\right)$$

by the rule

$$R_{0,\nu} : \{c_n(\xi)\}_{n \in \mathbb{Z}_+^{p-1}} \mapsto \{c_n(\xi) \alpha_{n,\nu}(\xi) \beta_{n,\nu}(s, \xi, \theta)\}_{n \in \mathbb{Z}^{p-1}},$$

where the functions $\beta_{n,\nu} = \beta_{n,\nu}(s, \xi, \theta)$ and $\alpha_{n,\nu}(\xi)$ are given by

$$\beta_{n,\nu} = s^n [1 - (1 + i)|s|^2]^{-\frac{\nu+|n|}{2} + i\xi} e^{-2(\xi + i\frac{\nu+|n|}{2}) \arctan[(1 - i\frac{|s|^2}{1-|s|^2}) \tan \frac{\theta}{2} + \frac{|s|^2}{1-|s|^2}]} \tag{4.3}$$

and

$$\alpha_{n,\nu}(\xi) = \left(\int_{\tau(\mathbb{B}^{p-1}) \times (0, \pi)} |\beta_{n,\nu}(s, \xi, \theta)|^2 (1 - |s|^2)^{\nu-p} \frac{c_\nu}{4} \sin^{\nu-p-1} \theta s ds d\theta \right)^{-\frac{1}{2}} \tag{4.4}$$

The adjoint operator $R_{0,\nu}^*$ which acts from

$$l_2\left(\mathbb{Z}_+^{p-1}, L_2(\tau(\mathbb{B}^{p-1}), (1 - |s|^2)^{\nu-p} s ds) \otimes L_2(\mathbb{R}) \otimes L_2\left((0, \pi), \frac{c_\nu}{4} \sin^{\nu-p-1} \theta d\theta\right)\right)$$

onto the space $l_2(\mathbb{Z}_+^{p-1}, L_2(\mathbb{R}))$ has obviously the form

$$R_{0,\nu}^* (\{d_n(s, \xi, \theta)\}_{n \in \mathbb{Z}^{p-1}}) = \left\{ \alpha_{n,\nu}(\xi) \int_{\tau(\mathbb{B}^{p-1}) \times (0, \pi)} \overline{\beta_{n,\nu}(s, \xi, \theta)} d_n(s, \xi, \theta) (1 - |s|^2)^{\nu-p} \times \frac{c_\nu}{4} \sin^{\nu-p-1} \theta s ds d\theta \right\}_{n \in \mathbb{Z}_+^{p-1}}.$$

In [9] the authors introduced the operator $R_\nu = R_{0,\nu}^* U$ from $L_2(\mathbb{D}^p, \tilde{\mu}_\nu)$ onto $l_2(\mathbb{Z}_+^{p-1}, L_2(\mathbb{R}))$, and the adjoint operator R_ν^* being an isometric isomorphism from $l_2(\mathbb{Z}_+^{p-1}, L_2(\mathbb{R}))$ onto the subspace $\mathcal{A}_\nu^2(\mathbb{D}^p)$. Furthermore

$$R_\nu R_\nu^* = I : l_2(\mathbb{Z}_+^{p-1}, L_2(\mathbb{R})) \longrightarrow l_2(\mathbb{Z}_+^{p-1}, L_2(\mathbb{R})),$$

$$R_\nu^* R_\nu = B_{\mathbb{U}^p, \nu} : L_2(\mathbb{U}^p, \tilde{\mu}_\nu) \longrightarrow \mathcal{A}_\nu^2(\mathbb{U}^p),$$

where $B_{\mathbb{U}^p, \nu}$ is the Bergman projection.

Now we define the analogue operator for the super case

Definition 4.9. Consider the operator

$$R_{\nu, (p|q)} : H_\nu^2(\mathbb{U}^{p|q}) \rightarrow (l_2(\mathbb{Z}_+^{p-1}, L_2(\mathbb{R})))^{2^q}$$

defined by

$$R_{\nu, (p|q)} \left(\sum_{M \subset Q} \psi_M \xi_M \right) = (R_{\nu+|M|}(\psi_M))_{M \subset Q}.$$

The adjoint operator

$$R_{\nu, (p|q)}^* : (l_2(\mathbb{Z}_+^{p-1}, L_2(\mathbb{R})))^{2^q} \rightarrow H_\nu^2(\mathbb{U}^{p|q})$$

is defined by

$$R_{\nu, (p|q)}^* (\{c_{M,n}(\xi)\}_{n \in \mathbb{Z}_+^{p-1}})_{M \subset Q} = \sum_{M \subset Q} R_{\nu+|M|} (\{c_{M,n}(\xi)\}_{n \in \mathbb{Z}_+^{p-1}}) \xi_M,$$

where the inner product in $(l_2(\mathbb{Z}_+^{p-1}, L_2(\mathbb{R})))^{2^q}$ is given by

$$\langle (\{a_{M,n}(\xi)\}_{n \in \mathbb{Z}_+^{p-1}})_{M \subset Q}, (\{b_{M,n}(\xi)\}_{n \in \mathbb{Z}_+^{p-1}})_{M \subset Q} \rangle = \sum_{M \subset Q} \frac{\Gamma(\nu)}{\Gamma(\nu + |M|)} \sum_{n \in \mathbb{Z}_+^{p-1}} (a_{M,n}(\xi), b_{M,n}(\xi))_{L_2(\mathbb{R}_+)}.$$

Theorem 4.10. *The operators*

$$R_{\nu, (p|q)} : H_\nu^2(\mathbb{U}^{p|q}) \rightarrow (l_2(\mathbb{Z}_+^{p-1}, L_2(\mathbb{R})))^{2^q}$$

and

$$R_{\nu, (p|q)}^* : (l_2(\mathbb{Z}_+^{p-1}, L_2(\mathbb{R})))^{2^q} \rightarrow H_\nu^2(\mathbb{U}^{p|q})$$

are isometric isomorphisms. Furthermore, we have

$$R_{\nu,(p|q)}R_{\nu,(p|q)}^* = I : (l_2(\mathbb{Z}_+^{p-1}, L_2(\mathbb{R}))^{2^q} \rightarrow (l_2(\mathbb{Z}_+^{p-1}, L_2(\mathbb{R}))^{2^q}$$

$$R_{\nu,(p|q)}^*R_{\nu,(p|q)} = I : H_\nu^2(\mathbb{U}^{p|q}) \rightarrow H_\nu^2(\mathbb{U}^{p|q}).$$

The proof is similar to that of Theorem 4.2.

5. Toeplitz Operators

Definition 5.1. For F an element of $\mathcal{C}(\mathbb{B}^{p|q})$ (an element of $\mathcal{C}(\mathbb{U}^{p|q})$), the super-Toeplitz operator T_F^ν on $H_\nu^2(\mathbb{B}^{p|q})$ (on $H_\nu^2(\mathbb{U}^{p|q})$, respectively) is defined by

$$T_F^\nu \Psi = P^\nu(F\Psi),$$

where P^ν denotes the orthogonal projection onto $H_\nu^2(\mathbb{B}^{p|q})$ (onto $H_\nu^2(\mathbb{U}^{p|q})$, respectively).

In [5] the authors proved that every super Toeplitz operator T_F^ν on $H_\nu^2(\mathbb{B}^{p|q})$ is given by the $2^q \times 2^q$ -matrix. With respect to the decomposition

$$\Psi = \sum_{M \subset Q} \psi_M \xi_M H_\nu^2(\mathbb{B}^{p|q}),$$

the super Toeplitz operator has the form

$$(T_F^\nu)_{I,J} = \sum_{I \cup J \subset K \subset Q} \left[\varepsilon_{K \setminus I, I} \varepsilon_{K \setminus J, J} \frac{\Gamma(\nu + |I| - p)}{\Gamma(\nu + |J| - p)} \right. \tag{5.1}$$

$$\left. \times T_{\nu+|I|}^{\nu+|J|}(F_{K \setminus I, K \setminus J}(w)(1 - ww)^{|K|-|I|}) \right]. \tag{5.2}$$

Here, for $0 \leq i, j \leq q$, $T_{\nu+i}^{\nu+j}$ denotes the Bergman-type Toeplitz operator

$$T_{\nu+i}^{\nu+j}(f) = P_{\nu+i} f P_{\nu+j} : H_{\nu+j}^2(\mathbb{B}^p) \rightarrow H_{\nu+i}^2(\mathbb{B}^p)$$

from $H_{\nu+j}^2(\mathbb{B}^p)$ to $H_{\nu+i}^2(\mathbb{B}^p)$.

The following proposition is a consequence of the above result and the fact that $H_\nu^2(\mathbb{B}^{p|q})$ is isometric to $H_\nu^2(\mathbb{U}^{p|q})$, which is given by Theorem 2.6.

Proposition 5.2. *With respect to the decomposition*

$$\Psi = \sum_{M \subset Q} \psi_M \xi_M H_\nu^2(\mathbb{U}^{p|q}),$$

the super Toeplitz operator T_F^ν on $H_\nu^2(\mathbb{U}^{p|q})$ is given by the $2^q \times 2^q$ -matrix

$$(T_F^\nu)_{I,J} = \sum_{I \cup J \subset K \subset Q} \left[\varepsilon_{K \setminus I, I} \varepsilon_{K \setminus J, J} \frac{\Gamma(\nu + |I| - p)}{\Gamma(\nu + |J| - p)} \right.$$

$$\left. \times T_{\nu+|I|}^{\nu+|J|} \left(F_{K \setminus I, K \setminus J}(w) \left(\frac{w_p - \bar{w}_p}{2i} - w' \bar{w}' \right)^{|K|-|I|} \right) \right].$$

Here, for $0 \leq i, j \leq q$, $T_{\nu+i}^{\nu+j}$ denotes the Bergman-type Toeplitz operator

$$T_{\nu+i}^{\nu+j}(f) = P_{\nu+i} f P_{\nu+j} : H_{\nu+j}^2(\mathbb{U}^p) \rightarrow H_{\nu+i}^2(\mathbb{U}^p)$$

from $H_{\nu+j}^2(\mathbb{U}^p)$ to $H_{\nu+i}^2(\mathbb{U}^p)$.

Corollary 5.3. *Let F be a super function of the form*

$$F(z, \xi) = \sum_{M \subset Q} F_M(z) \xi_M \xi_M^*.$$

Then the super Toeplitz operator T_F^ν on $H_\nu^2(\mathbb{B}^{p|q})$ (or $H_\nu^2(\mathbb{B}^{p|q})$) is given by the $2^q \times 2^q$ diagonal-matrix

$$(T_F^\nu)_{I,I} = \sum_{I \subset K \subset Q} T_{\nu+|I|}^{\nu+|I|}(F_{K \setminus I}(w)(1 - w\bar{w})^{|K|-|I|}), \tag{5.3}$$

or

$$(T_F^\nu)_{I,I} = \sum_{I \subset K \subset Q} T_{\nu+|I|}^{\nu+|I|} \left(F_{K \setminus I}(w) \left(\frac{w_p - \bar{w}_p}{2i} - w' \bar{w}' \right)^{|K|-|I|} \right).$$

6. Super Toeplitz Operators with Special Symbols

6.1. Quasi-Elliptic

We will call a super function F quasi-elliptic if it is invariant under the action of the quasi-elliptic group, in other words, when F has the following form

$$F(z, \xi) = \sum_{I \subset Q} F_I(r) \xi_I \xi_I^*,$$

where $r = (r_1, \dots, r_p) = (|z_1|, \dots, |z_p|)$.

Theorem 6.1. *Let F be a bounded measurable quasi-elliptic super function. Then the Toeplitz operator T_F^ν acting on $H_\nu^2(\mathbb{B}^{p|q})$ is unitarily equivalent to the multiplication operator $\gamma_{F,\nu} I = R_{\nu,(p|q)} T_F^\nu R_{\nu,(p|q)}^*$ acting on $(l_2(\mathbb{Z}_+^p))^{2^q}$, where $R_{\nu,(p|q)}$ and $R_{\nu,(p|q)}^*$ are given in Definition 4.1. The sequence*

$$\gamma_{F,\nu} = (\{\gamma_{F,\nu}(n, M)\}_{n \in \mathbb{Z}_+^p})_{M \subset Q}$$

is given by

$$\gamma_{F,\nu}(n, M) = \frac{2^p \Gamma(|n| + \nu + |M|)}{n! \Gamma(\nu + |M| - p)} \sum_{K \in \mathcal{I}_{M \setminus \tau}(\mathbb{B}^p)} \int F_{K \setminus M}(r) r^{2n} (1 - r^2)^{\nu + |K| - p - 1} r dr,$$

where $\mathcal{I}_M = \{K \subset Q : M \subset K\}$.

Proof. For every super function F the Toeplitz operator is unitarily equivalent to

$$R_{\nu,(p|q)} T_F^\nu R_{\nu,(p|q)}^* : (l_2(\mathbb{Z}_+^p))^{2^q} \rightarrow (l_2(\mathbb{Z}_+^p))^{2^q},$$

where the components of the operator are

$$(R_{\nu,(p|q)} T_F^\nu R_{\nu,(p|q)}^*)_{I,J} = R_{\nu+|I|} (T_F^\nu)_{I,J} R_{\nu+|J|}^*. \tag{6.1}$$

Now, we consider a super function of the form $F = \sum_{I \subset Q} F_I(r) \xi_I \xi_I^*$ and using Corollary 5.3 we have that the Toeplitz operator has the form

$$(T_F^\nu)_{I,J} = \begin{cases} 0 & I \neq J \\ \sum_{K \in \mathcal{I}_I} T_{\nu+|I|}^{\nu+|I|}(F_{K \setminus I}(r)(1 - |r|^2)^{|K|-|I|}) & I = J \end{cases} \tag{6.2}$$

From the above equation and 6.1 we have

$$(R_{\nu,(p|q)}T_F^\nu R_{\nu,(p|q)}^*)_{I,J} = \begin{cases} 0 & I \neq J \\ \sum_{K \in \mathcal{I}_I} R_{\nu+|I} P_{\nu+|I} (F_{K \setminus I}(r)(1 - |r|^2)^{|K|-|I|}) P_{\nu+|I} R_{\nu+|I}^* & I = J \end{cases}.$$

By Theorem 10.1 in [9] the above operators are multiplication operators then

$$\begin{aligned} & \sum_{K \in \mathcal{I}_I} R_{\nu+|I} P_{\nu+|I} (F_{K \setminus I}(r)(1 - |r|^2)^{|K|-|I|}) P_{\nu+|I} R_{\nu+|I}^* (\{c_{n,I}\}_{n \in \mathbb{Z}_+^p}) \\ &= \left\{ \left[l_{\nu+|I}(n) \sum_{K \in \mathcal{I}_{I \setminus \tau}(\mathbb{B}^p)} \int (F_{K \setminus I}(r)(1 - r^2)^{|K|-|I|}) \right. \right. \\ & \quad \left. \left. \times r^{2n}(1 - r^2)^{\nu+|I|-p-1} r dr \right] c_{n,I} \right\}_{n \in \mathbb{Z}_+^p} \\ &= \left\{ \left[l_{\nu+|I}(n) \sum_{K \in \mathcal{I}_{I \setminus \tau}(\mathbb{B}^p)} \int F_{K \setminus I}(r) r^{2n} (1 - r^2)^{\nu+|K|-p-1} r dr \right] c_{n,I} \right\}_{n \in \mathbb{Z}_+^p}, \end{aligned}$$

where

$$l_{\nu+|I}(n) = \frac{2^p \Gamma(|n| + \nu + |I|)}{n! \Gamma(\nu + |I| - p)}.$$

□

6.2. Quasi-Parabolic Case

We will call a super function F quasi-parabolic if it is invariant under the action of the quasi-parabolic group, in other words, when F has the following form

$$F(z, \xi) = \sum_{I \subset Q} f_I(r, \text{Im}(z_n)) \xi_I \xi_I^*,$$

where $r = (r_1, \dots, r_{p-1}) = (|z_1|, \dots, |z_{p-1}|)$.

Theorem 6.2. *Let F be a bounded measurable quasi-parabolic super function. Then the Toeplitz operator T_F^ν acting on $H_\nu^2(\mathbb{D}^{p|q})$ is unitarily equivalent to the multiplication operator $\gamma_{F,\nu} I = R_{\nu,(p|q)} T_F^\nu R_{\nu,(p|q)}^*$ acting on $(l_2(\mathbb{Z}_+^{p-1}, L_2(\mathbb{R}_+)))^{2^q}$, where $R_{\nu,(p|q)}$ and $R_{\nu,(p|q)}^*$ are given in Definition 4.3. The sequence*

$$\gamma_{F,\nu} = (\{\gamma_{F,\nu}(n, \xi, M)\}_{n \in \mathbb{Z}_+^{p-1}})_{M \subset Q}$$

is given by

$$\begin{aligned} \gamma_{F,\nu}(n, \xi, M) &= \frac{(2\xi)^{|n|+\nu+|M|-1}}{n! \Gamma(\nu + |M| - p)} \\ &\times \sum_{K \in \mathcal{I}_M \mathbb{R}_+^p} \int F_{K \setminus M}(\sqrt{r'}, v + \hat{r}) r^n e^{-2\xi(v+\hat{r})} v^{\nu+|K|-p-1} dr dv, \end{aligned}$$

where $r = (r_1, \dots, r_{p-1})$, $\sqrt{r} = (\sqrt{r_1}, \dots, \sqrt{r_{p-1}})$, $\hat{r} = r_1 + \dots + r_{p-1}$ and $\mathcal{I}_M = \{K \subset Q : M \subset K\}$.

Proof. For every super function F the Toeplitz operator is unitarily equivalent to

$$R_{\nu, (p|q)} T_F^\nu R_{\nu, (p|q)}^* : (l_2(\mathbb{Z}_+^{p-1}, L_2(\mathbb{R}_+)))^{2^q} \rightarrow (l_2(\mathbb{Z}_+^{p-1}, L_2(\mathbb{R}_+)))^{2^q},$$

where the components of the operator are

$$(R_{\nu, (p|q)} T_F^\nu R_{\nu, (p|q)}^*)_{I, J} = R_{\nu+|I|} (T_F^\nu)_{I, J} R_{\nu+|I|}^*. \tag{6.3}$$

Now, we consider a super function $F = \sum_{I \subset Q} F_I(r, \text{Im}(z_n)) \xi_I \xi_I^*$ and using Corollary 5.3 we obtain

$$(T_F^\nu)_{I, J} = \begin{cases} 0 & I \neq J \\ \sum_{K \in \mathcal{I}_I} T_{\nu+|I|}^{\nu+|I|} (F_{K \setminus I}(r, v + \hat{r}) v^{|K|-|I|}) & I = J \end{cases}.$$

From the above equation and 6.3 we have

$$\begin{aligned} & (R_{\nu, (p|q)} T_F^\nu R_{\nu, (p|q)}^*)_{I, J} \\ &= \begin{cases} 0 & I \neq J \\ \sum_{K \in \mathcal{I}_I} R_{\nu+|I|} P_{\nu+|I|} (F_{K \setminus I}(r, v + \hat{r}) (v)^{|K|-|I|}) P_{\nu+|I|} R_{\nu+|I|}^* & I = J \end{cases}. \end{aligned}$$

By Theorem 10.2 in [9] the above operators are multiplication operators

$$\begin{aligned} & \sum_{K \in \mathcal{I}_I} R_{\nu+|I|} P_{\nu+|I|} (F_{K \setminus I}(r', v + \hat{r}) (v)^{|K|-|I|}) \\ & \quad \times P_{\nu+|I|} R_{\nu+|I|}^* (\{c_{n, I}(\xi)\}_{n \in \mathbb{Z}_+^{p-1}}) \\ &= \left\{ \left[\sum_{K \in \mathcal{I}_I} \frac{(2\xi)^{|n|+\nu+|I|-1}}{n! \Gamma(\nu + |M| - p)} \int_{\mathbb{R}_+^p} (F_{K \setminus I}(\sqrt{r'}, v + \hat{r}) v^{|K|-|I|}) \right. \right. \\ & \quad \left. \left. \times r^n e^{-2\xi(v+\hat{r})} v^{\nu+|I|-p-1} dr dv \right] c_{n, I}(\xi) \right\}_{n \in \mathbb{Z}_+^{p-1}} \\ &= \left\{ \left[\frac{(2\xi)^{|n|+\nu+|I|-1}}{n! \Gamma(\nu + |M| - p)} \sum_{K \in \mathcal{I}_I} \int_{\mathbb{R}_+^p} F_{K \setminus I}(\sqrt{r'}, v + \hat{r}) \right. \right. \\ & \quad \left. \left. \times r^n e^{-2\xi(v+\hat{r})} v^{\nu+|K|-p-1} dr dv \right] c_{n, I}(\xi) \right\}_{n \in \mathbb{Z}_+^{p-1}}. \end{aligned}$$

□

6.3. Nilpotent Case

We will call a super function F nilpotent if it is invariant under the action of the Nilpotent group, in other words, when F has the following form

$$F(z, \xi) = \sum_{I \subset Q} f_I(\text{Im } z', \text{Im } z_n - |z'|^2) \xi_I \xi_I^*,$$

where $z' = (z_1, \dots, z_{p-1})$.

Theorem 6.3. *Let F be a bounded measurable nilpotent super function. Then the Toeplitz operator T_F^ν acting on $H_\nu^2(\mathbb{U}^{p|q})$ is unitarily equivalent to the multiplication operator $\gamma_{F,\nu}I = R_{\nu,(p|q)}T_F^\nu R_{\nu,(p|q)}^*$ acting on $(L_2(\mathbb{R}^{p-1} \times \mathbb{R}_+))^{2^q}$, where $R_{\nu,(p|q)}$ and $R_{\nu,(p|q)}^*$ are given in Definition 4.5. The function*

$$\gamma_{F,\nu} = (\gamma_{F,\nu}(u', \xi, M))_{M \subset Q}$$

is given by

$$\begin{aligned} \gamma_{F,\nu}(u', \xi, M) &= \frac{(2\xi)^{\nu+|M|-p}}{\pi^{\frac{p-1}{2}} \Gamma(\nu + |M| - p)} \sum_{K \in \mathcal{I}_M} \int_{\mathbb{R}^{n-1} \times \mathbb{R}_+} \\ &\times F_{K \setminus M} \left(\frac{1}{2\sqrt{\xi}}(-u' + v'), v \right) e^{-2\xi v - |v'|^2} v^{\nu+|K|-p-1} dv' dv, \end{aligned}$$

where $\mathcal{I}_M = \{K \subset Q : M \subset K\}$, $u' \in \mathbb{R}^{n-1}$ and $\xi \in \mathbb{R}_+$.

Proof. For every super function F the Toeplitz operator is unitarily equivalent to

$$R_{\nu,(p|q)}T_F^\nu R_{\nu,(p|q)}^* : (L_2(\mathbb{R}^{p-1} \times \mathbb{R}_+))^{2^q} \rightarrow (L_2(\mathbb{R}^{p-1} \times \mathbb{R}_+))^{2^q},$$

where the components of the operator are

$$(R_{\nu,(p|q)}T_F^\nu R_{\nu,(p|q)}^*)_{I,J} = R_{\nu+|I|}(T_F^\nu)_{I,J}R_{\nu+|J|}^*. \tag{6.4}$$

Now, we consider a super function of the form

$$F = \sum_{I \subset Q} F_I(\text{Im } z', \text{Im } z_n - |z'|^2)\xi_I \xi_I^*.$$

Using (5.3) we obtain

$$(T_F^\nu)_{I,J} = \begin{cases} 0 & I \neq J \\ \sum_{K \in \mathcal{I}_I} T_{\nu+|I|}^{\nu+|I|}(F_{K \setminus I}(\text{Im } z', \text{Im } z_n - |z'|^2)v^{|K|-|I|}) & I = J \end{cases} \tag{6.5}$$

From the above equation and 6.4 we have

$$\begin{aligned} &(R_{\nu,(p|q)}T_F^\nu R_{\nu,(p|q)}^*)_{I,J} \\ &= \begin{cases} 0 & I \neq J \\ \sum_{K \in \mathcal{I}_I} R_{\nu+|I|}P_{\nu+|I|}(F_{K \setminus I}(y', v)(v)^{|K|-|I|})P_{\nu+|I|}R_{\nu+|I|}^* & I = J \end{cases} \end{aligned}$$

By Theorem 10.3 in [9] the above operators are the multiplication operators

$$\begin{aligned} &\sum_{K \in \mathcal{I}_I} R_{\nu+|I|}P_{\nu+|I|}(F_{K \setminus I}(y', v)(v)^{|K|-|I|})P_{\nu+|I|}R_{\nu+|I|}^*(c_I(u', \xi)) \\ &= \left[\sum_{K \in \mathcal{I}_I} \frac{(2\xi)^{\nu+|M|-p}}{\pi^{\frac{p-1}{2}} \Gamma(\nu + |M| - p)} \int_{\mathbb{R}^{n-1} \times \mathbb{R}_+} F_{K \setminus M} \left(\frac{1}{2\sqrt{\xi}}(-u' + v'), v \right) \right. \\ &\quad \left. \times e^{-2\xi v - |v'|^2} v^{\nu+|K|-p-1} dv' dv \right] c_I(u', \xi). \end{aligned}$$

□

6.4. Quasi-Nilpotent Case

We will call a super function F quasi-nilpotent if it is invariant under the action of the quasi-nilpotent group, in other words, when F has the following form

$$F(z, \xi) = \sum_{I \subset Q} f_I(r, y', \text{Im } z_n - |z'|^2) \xi_I \xi_I^*,$$

where $r = (|z_1|, \dots, |z_k|)$, $y' = \text{Im } w'$, and $w' = (z_{k+1}, \dots, z_{p-1})$.

Theorem 6.4. *Let F be a bounded measurable quasi-nilpotent super function. Then the Toeplitz operator T_F^ν acting on $H_\nu^2(\mathbb{U}^{p|q})$ is unitarily equivalent to the multiplication operator $\gamma_{F,\nu} I = R_{\nu,(p|q)} T_F^\nu R_{\nu,(p|q)}^*$ acting on $(l_2(\mathbb{Z}_+^k, L_2(\mathbb{R}^{p-k-1} \times \mathbb{R}_+)))^{2^q}$, where $R_{\nu,(p|q)}$ and $R_{\nu,(p|q)}^*$ are given in Definition 4.7. The function*

$$\gamma_{F,\nu} = (\{\gamma_{F,\nu}(n, u', \xi, M)\}_{n \in \mathbb{Z}_+^k})_{M \subset Q}$$

is given by

$$\begin{aligned} \gamma_{F,\nu}(n, u', \xi, M) &= \pi^{-\frac{p-k-1}{2}} \frac{(2\xi)^{|n|+\nu+|M|-p+k}}{n! \Gamma(\nu + |M| - p)} \sum_{K \in \mathcal{I}_M} \int_{\mathbb{R}_+^k \times \mathbb{R}^{n-k-1} \times \mathbb{R}_+} \\ &\quad \times F_{K \setminus M} \left(\sqrt{r}, \frac{1}{2\sqrt{\xi}}(-u' + v'), v + r_1 + \dots + r_k \right) r^p \\ &\quad \times e^{-2\xi(v+r_1+\dots+r_k)-|v'|^2} v^{\nu+|K|-p-1} dr dv' dv, \end{aligned}$$

where $\mathcal{I}_M = \{K \subset Q : M \subset K\}$, $\sqrt{r} = (\sqrt{r_1}, \dots, \sqrt{r_k})$, $n \in \mathbb{Z}_+^k$, $u' \in \mathbb{R}^{n-k-1}$ and $\xi \in \mathbb{R}_+$.

Proof. For every super function F the Toeplitz operator is unitarily equivalent to

$$\begin{aligned} R_{\nu,(p|q)} T_F^\nu R_{\nu,(p|q)}^* &: (l_2(\mathbb{Z}_+^k, L_2(\mathbb{R}^{p-k-1} \times \mathbb{R}_+)))^{2^q} \\ &\rightarrow (l_2(\mathbb{Z}_+^k, L_2(\mathbb{R}^{p-k-1} \times \mathbb{R}_+)))^{2^q}, \end{aligned}$$

where the components of the operator are

$$(R_{\nu,(p|q)} T_F^\nu R_{\nu,(p|q)}^*)_{I,J} = R_{\nu+|I|} (T_F^\nu)_{I,J} R_{\nu+|J|}^*. \tag{6.6}$$

Now, we consider a super function of the form $F = \sum_{I \subset Q} F_I(r, y', \text{Im } z_n - |z'|^2) \xi_I \xi_I^*$ and using Corollary 5.3 we obtain

$$(T_F^\nu)_{I,J} = \begin{cases} 0 & I \neq J \\ \sum_{K \in \mathcal{I}_I} T_{\nu+|I|}^{\nu+|I|} (F_{K \setminus I}(r, y', \text{Im } z_n - |z'|^2) v^{|K|-|I|}) & I = J \end{cases} \tag{6.7}$$

From the above equation and 6.6 we have

$$\begin{aligned} &(R_{\nu,(p|q)} T_F^\nu R_{\nu,(p|q)}^*)_{I,J} \\ &= \begin{cases} 0 & I \neq J \\ \sum_{K \in \mathcal{I}_I} R_{\nu+|I|} P_{\nu+|I|} (F_{K \setminus I}(r, y', v) (v)^{|K|-|I|}) P_{\nu+|I|} R_{\nu+|I|}^* & I = J \end{cases} \end{aligned}$$

In particular by Theorem 10.4 in [9] the following operator is a multiplication operator

$$\begin{aligned} & \sum_{K \in \mathcal{I}_I} R_{\nu+|I|} P_{\nu+|I|}(F_{K \setminus I}(r, y', v)(v)^{|K|-|I|}) P_{\nu+|I|} R_{\nu+|I|}^*(c_{n,I}(u', \xi)) \\ &= \sum_{K \in \mathcal{I}_I} \pi^{-\frac{p-k-1}{2}} \frac{(2\xi)^{|n|+\nu+|M|-p+k}}{n! \Gamma(\nu+|M|-p)} \int_{\mathbb{R}_+^k \times \mathbb{R}^{n-k-1} \times \mathbb{R}_+} \\ & \quad \times F_{K \setminus M}(\sqrt{r}, \frac{1}{2\sqrt{\xi}}(-u' + v'), v + r_1 + \dots + r_k) \\ & \quad \times r^n e^{-2\xi(v+r_1+\dots+r_k)-|v'|^2} v^{\nu+|K|-p-1} dr dv' dv \cdot [c_{n,I}(u', \xi)]. \end{aligned}$$

□

6.5. Quasi-Hyperbolic Case

We will call a super function F quasi-nilpotent if it is invariant under the action of the Quasi-nilpotent group, in other words, when F has the following form

$$F(z, \xi) = \sum_{I \subset Q} f_I(s_1, \dots, s_{n-1}, \arg(z_n - i|z'|^2)) |z_n - i|z'|^2|^{-|I|} \xi_I \xi_I^*,$$

where

$$s_k = \frac{|z_k|}{\sqrt{|z'|^2 + |z_n - i|z'|^2|}},$$

$k = 1, \dots, n - 1$ and $z' = (z_1, \dots, z_{p-1})$.

Theorem 6.5. *Let F be a bounded measurable quasi-nilpotent super function. Then the Toeplitz operator T_F^ν acting on $H_\nu^2(\mathbb{U}^{p|q})$ is unitarily equivalent to the multiplication operator $\gamma_{F,\nu} I = R_{\nu,(p|q)} T_F^\nu R_{\nu,(p|q)}^*$ acting on $(l_2(\mathbb{Z}_+^{p-1}, L_2(\mathbb{R})))^{2^q}$, where $R_{\nu,(p|q)}$ and $R_{\nu,(p|q)}^*$ are given in Definition 4.9. The function*

$$\gamma_{F,\nu} = (\{\gamma_{F,\nu}(n, \xi, M)\}_{n \in \mathbb{Z}_+^{p-1}})_{M \subset Q}$$

is given by

$$\begin{aligned} \gamma_{F,\nu}(n, \xi, M) &= \sum_{K \in \mathcal{I}_M} \alpha_{n,\nu+|M|}^2(\xi) \int_{\tau(\mathbb{B}^{n-1}) \times (0,\pi)} F_{K \setminus M}(s, \theta) |\beta_{n,\nu+|M|}(s, \xi, \theta)|^2 \\ & \quad \times (1 - |s|^2)^{\nu+|M|-p} \frac{C_{\nu+|M|}}{4} \sin^{\nu+|K|-p-1} \theta \, ds d\theta, \end{aligned}$$

where $\mathcal{I}_M = \{K \subset Q : M \subset K\}$, and the functions $\alpha_p(\xi)$ and $\beta_p(s, \xi, \theta)$ are given by (4.4) and (4.3), respectively.

Proof. For every super function F the Toeplitz operator is unitarily equivalent to

$$R_{\nu,(p|q)} T_F^\nu R_{\nu,(p|q)}^* : (l_2(\mathbb{Z}_+^{p-1}, L_2(\mathbb{R})))^{2^q} \rightarrow (l_2(\mathbb{Z}_+^{p-1}, L_2(\mathbb{R})))^{2^q},$$

where the components of the operator are

$$(R_{\nu,(p|q)} T_F^\nu R_{\nu,(p|q)}^*)_{I,J} = R_{\nu+|I|} (T_F^\nu)_{I,J} R_{\nu+|J|}^*. \tag{6.8}$$

Now, we consider a super function of the form $F = \sum_{I \subset Q} F_I(s, \theta) r^{-|I|} \xi_I \xi_I^*$ and using Corollary 5.3 we obtain

$$(T_F^\nu)_{I,J} = \begin{cases} 0 & I \neq J \\ \sum_{K \in \mathcal{I}_I} T_{\nu+|I|}^{\nu+|I|} (F_{K \setminus I}(s, \theta) r^{-|K \setminus I|} v^{|K|-|I|}) & I = J \end{cases} \quad (6.9)$$

From the above equation and 6.8 we have

$$(R_{\nu, (p|q)} T_F^\nu R_{\nu, (p|q)}^*)_{I,J} = \begin{cases} 0 & I \neq J \\ \sum_{K \in \mathcal{I}_I} R_{\nu+|I|} P_{\nu+|I|} (F_{K \setminus I}(s, \theta) \sin^{|K|-|I|} \theta) P_{\nu+|I|} R_{\nu+|I|}^* & I = J \end{cases}$$

By Theorem 10.5 in [9] the above operators are multiplication operators

$$\begin{aligned} & \sum_{K \in \mathcal{I}_I} R_{\nu+|I|} P_{\nu+|I|} F_{K \setminus I}(s, \theta) \sin^{|K|-|I|} \theta P_{\nu+|I|} R_{\nu+|I|}^*(c_{n,I}(\xi)) \\ &= \sum_{K \in \mathcal{I}_I} \alpha_{n, \nu+|I|}^2(\xi) \int_{\tau(\mathbb{B}^{n-1}) \times (0, \pi)} F_{K \setminus I}(s, \theta) \sin^{|K|-|I|} \theta |\beta_{n, \nu+|I|}(s, \xi, \theta)|^2 \\ & \quad \times (1 - |s|^2)^{\nu+|I|-p} \frac{C_{\nu+|I|}}{4} \sin^{\nu+|I|-p-1} \theta s ds d\theta \cdot [c_{n,I}(u', \xi)] \\ &= \sum_{K \in \mathcal{I}_I} \alpha_{n, \nu+|I|}^2(\xi) \int_{\tau(\mathbb{B}^{n-1}) \times (0, \pi)} F_{K \setminus I}(s, \theta) |\beta_{n, \nu+|I|}(s, \xi, \theta)|^2 \\ & \quad \times (1 - |s|^2)^{\nu+|I|-p} \frac{C_{\nu+|I|}}{4} \sin^{\nu+|K|-p-1} \theta s ds d\theta \cdot [c_{n,I}(u', \xi)]. \end{aligned}$$

□

References

- [1] Berezin, F.A.: Introduction to Superanalysis. Reidel, Dordrecht (1987)
- [2] Borthwick, D., Klimek, S., Lesniewski, A., Rinaldi, M.: Super Toeplitz operators and non-perturbative deformation quantization of supermanifolds. Commun. Math. Phys. **153**, 49–76 (1993)
- [3] Borthwick, D., Klimek, S., Lesniewski, A., Rinaldi, M.: Matrix Cartan superdomains, super upper operators, and quantization. J. Funct. Anal. **127**(2), 456–510 (1995)
- [4] Grudsky, S., Quiroga, R., Vasilevski, N.: Commutative C^* -algebras of Toeplitz operators and quantization on the unit disk. J. Funct. Anal. **234**, 1–44 (2006)
- [5] Loaiza, M., Upmeyer, H.: Toeplitz C^* -algebras on super-Cartan domains. Rev. Mat. Complut. **21**(2), 489–518 (2008)
- [6] Loaiza, M., Sánchez-Nungaray, A.: On C^* -algebras of super Toeplitz operators with radial symbols. Oper. Theory Adv. Appl. **210**, 175–188 (2010)
- [7] Loaiza, M., Sánchez-Nungaray, A.: Commutative algebras of Toeplitz operators on the super upper half-plane: quasi-hyperbolic and quasi-parabolic cases. Oper. Theory Adv. Appl. **236**, 317–336 (2014)
- [8] Quiroga-Barranco, R., Vasilevski, N.: Commutative algebras of Toeplitz operators on Reinhardt domains. Integral Equ. Oper. Theory **59**, 67–98 (2007)

- [9] Quiroga-Barranco, R., Vasilevski, N.: Commutative C^* -algebra of Toeplitz operators on the unit ball, I. Bargmann type transform and spectral representations of Toeplitz operators. *Integral Equ. Oper. Theory* **59**(3), 379–419 (2007)
- [10] Quiroga-Barranco, R., Vasilevski, N.: Commutative C^* -algebra of Toeplitz operators on the unit ball, II. Geometry of the level sets of symbols. *Integral Equ. Oper. Theory* **59**(1), 89–132 (2008)
- [11] Sánchez-Nungaray, A.: Commutative algebras of Toeplitz operators on the super upper half-plane: super parabolic case. *Oper. Theory Adv. Appl.* **220**, 263–280 (2012)
- [12] Zhu, K.: *Spaces of Holomorphic Functions in the Unit Ball*. Springer, New York (2005)

R. Quiroga-Barranco
Centro de Investigación en Matemáticas
Guanajuato
Mexico
e-mail: quiroga@cimat.mx

A. Sánchez-Nungaray
Facultad de Matemáticas
Universidad Veracruzana
Veracruz
Mexico
e-mail: armsanchez@uv.mx

Received: April 17, 2015.

Accepted: August 12, 2015.