

A new look at the Jordan-Hölder theorem for semimodular lattices

G. GRÄTZER AND J.B. NATION

ABSTRACT. We show that in a semimodular lattice L of finite length, from any prime interval we can reach any maximal chain C by an up- and a down-perspectivity. Therefore, C is a congruence-determining sublattice of L .

A classical theorem of R. Dedekind [2] (see also R. Dedekind [3]) states that the factors of any chief series (maximal chain of normal subgroups) of a finite group are invariant. C. Jordan, O. Hölder, and H. Wielandt generalized this result to the factors of any composition series (maximal chain of subnormal subgroups).

Formulations of this result for modular lattices are well-known; they are called the *Jordan-Hölder Theorem*. For semimodular lattices of finite length, most books (see, for instance, [1, page 40] and [4, page 226]) contain only a much weaker result: any two maximal chains in a semimodular lattice have the same length.

However, there is a stronger version for semimodular lattices that requires only a slight extension of Dedekind's original argument. This can be found in [6, page 444] or [7, Chapter 9]. Recall that the intervals $[a, b]$ and $[c, d]$ are *perspective* if $b \vee c = d$ and $b \wedge c = a$ or *vice versa*. *Projectivity* is the transitive closure of perspectivity.

Theorem 1. *In a in a semimodular lattice, let*

$$\begin{aligned} C: 0 = c_0 \prec c_1 \prec \cdots \prec c_n = 1, \\ D: 0 = d_0 \prec d_1 \prec \cdots \prec d_n = 1. \end{aligned}$$

Then there is a permutation π of the set $\{1, \dots, n\}$ such that $[c_{i-1}, c_i]$ is projective to $[d_{\pi(i)-1}, d_{\pi(i)}]$, for all i .

In this note, we improve on Theorem 1 by replacing projectivity with *two* perspectivities.

Presented by G. Czedli.

Received June 8, 2009; accepted in final form February 10, 2010.

2000 *Mathematics Subject Classification*: Primary: 06C10; Secondary: 06B05.

Key words and phrases: semimodular lattice, maximal chain, finite length, congruence-determining sublattice.

Theorem 2. *In a semimodular lattice L , let*

$$\begin{aligned} C: 0 = c_0 \prec c_1 \prec \cdots \prec c_n = 1, \\ D: 0 = d_0 \prec d_1 \prec \cdots \prec d_n = 1. \end{aligned}$$

Then there is a permutation π of the set $\{1, \dots, n\}$ with the following property: for each i , there exists a (prime) interval \mathfrak{p}_i such that $[c_{i-1}, c_i]$ is up-perspective to \mathfrak{p}_i and \mathfrak{p}_i is down-perspective to $[d_{\pi(i)-1}, d_{\pi(i)}]$.

Remark. Recall that the lattice of subnormal subgroups of a finite group is lower semimodular, so the dual of this theorem applies to yield the full Jordan-Hölder theorem for groups.

Proof. By induction on the length $\text{len}(L)$ of the lattice L . The statement is obvious for $\text{len}(L) \leq 2$, so let $\text{len}(L) > 2$.

Let k be the largest integer with $c_1 \not\leq d_k$; note that $k < n$. If $k = 0$, then $c_1 = d_1$ and the statement follows by the induction hypothesis. So we can assume that $k > 0$.

For $0 \leq j \leq n$, let $e_j = c_1 \vee d_j$. Note that $e_0 = c_1$ and $e_k = e_{k+1} = d_{k+1}$, and indeed $e_j = d_j$, for $j \geq k+1$. Now

$$c_1 = e_0 \prec e_1 \prec \cdots \prec e_k = e_{k+1} \prec e_{k+2} \prec \cdots \prec e_n = 1$$

is a maximal chain in the interval $[c_1, 1]$. By induction, there is a bijective map $\sigma: \{2, \dots, n\} \rightarrow \{1, \dots, k, k+2, \dots, n\}$ such that, for $i > 1$, each interval $[c_{i-1}, c_i]$ is up-perspective to some prime interval \mathfrak{p}_i in L , which in turn is down-perspective to $[e_{\sigma(i)-1}, e_{\sigma(i)}]$. For $j \leq k$, $[e_{j-1}, e_j]$ is down-perspective to $[d_{j-1}, d_j]$, while for $j > k+1$ we have $[e_{j-1}, e_j] = [d_{j-1}, d_j]$. Meanwhile, $[0, c_1]$ is up-perspective to $[d_k, d_{k+1}]$. So we may take π to be the permutation with $\pi(i) = \sigma(i)$, for $i \neq 1$, and $\pi(1) = k+1$. \square

Given a sublattice S of a lattice L , we say that S is a *congruence-determining sublattice* of L if any congruence relation on L is uniquely determined by its restriction to S ; see [5]. The reader should have little difficulty in deriving the following statement.

Corollary 3. *Let L be a semimodular lattice and C be a finite maximal chain in L . Then C is a congruence-determining sublattice of L .*

REFERENCES

- [1] Birkhoff, G.: Lattice Theory, 3rd edition. Amer. Math. Soc., Providence (1967)
- [2] Dedekind, R.: Über die von drei Moduln erzeugte Dualgruppe. Math. Annalen **(53)** (1900), 371–403. In: Gesammelte mathematische Werke, Vol. 2, pp. 236–271, Chelsea (1968)
- [3] Dedekind, R.: Gesammelte mathematische Werke. Bände I–III. (German) Herausgegeben von R. Fricke, E. Noether und O. Ore. Chelsea Publishing Co., New York (1968)

- [4] Grätzer, G.: General Lattice Theory, 2nd edition. New appendices by the author with B.A. Davey, R. Freese, B. Ganter, M. Greferath, P. Jipsen, H.A. Priestley, H. Rose, E.T. Schmidt, S.E. Schmidt, F. Wehrung, and R. Wille. Birkhäuser Verlag, Basel (1998)
- [5] Grätzer, G.: The Congruences of a Finite Lattice. A *Proof-by-Picture* Approach. Birkhäuser Boston (2006)
- [6] Jacobson, N.: *Basic algebra. I.* 2nd edn. W.H. Freeman and Company, New York (1985)
- [7] Nation, J.B.: Notes on Lattice Theory.
<http://www.math.hawaii.edu/~jb/books.html>
- [8] Wielandt, H.: Eine Verallgemeinerung der invarianten Untergruppen. *Math. Zeit.* **45**, 209–244 (1939)

G. GRÄTZER

Department of Mathematics, University of Manitoba, Winnipeg, MB R3T 2N2, Canada

e-mail, G. Grätzer: gratzer@ms.umanitoba.ca

URL, G. Grätzer: <http://server.math.umanitoba.ca/homepages/gratzer/>

J.B. NATION

University of Hawaii, Honolulu, HI, USA

e-mail, J.B. Nation: jb@math.hawaii.edu