

Equidistribution of signs for modular eigenforms of half integral weight

ILKER INAM AND GABOR WIESE

Abstract. Let f be a cusp form of weight $k + 1/2$ and at most quadratic nebentype character whose Fourier coefficients $a(n)$ are all real. We study an equidistribution conjecture of Bruinier and Kohnen for the signs of $a(n)$. We prove this conjecture for certain subfamilies of coefficients that are accessible via the Shimura lift by using the Sato–Tate equidistribution theorem for integral weight modular forms. Firstly, an unconditional proof is given for the family $\{a(tp^2)\}_p$, where t is a squarefree number and p runs through the primes. In this case, the result is in terms of natural density. To prove it for the family $\{a(tn^2)\}_n$ where t is a squarefree number and n runs through all natural numbers, we assume the existence of a suitable error term for the convergence of the Sato–Tate distribution, which is weaker than one conjectured by Akiyama and Tanigawa. In this case, the results are in terms of Dedekind–Dirichlet density.

Mathematics Subject Classification (2010). Primary 11F37;
Secondary 11F30.

Keywords. Forms of half-integer weight, Shimura lift, Fourier coefficients of automorphic forms, Sato–Tate equidistribution.

1. Introduction. The signs of the coefficients of half-integral weight modular forms have attracted some recent attention. In particular, Bruinier and Kohnen conjectured an equidistribution of signs. Using the Shimura lift and the Sato–Tate equidistribution theorem, we obtain results towards this conjecture.

I.I. was supported by The Scientific and Technological Research Council of Turkey (TUBITAK). I.I. would also like to thank the University of Luxembourg for having hosted him as a postdoctoral fellow of TUBITAK. G.W. acknowledges partial support by the priority program 1489 of the Deutsche Forschungsgemeinschaft (DFG) and by the Fonds National de la Recherche Luxembourg (INTER/DFG/12/10).

The authors would like to thank Winfried Kohnen and Jan Bruinier for having suggested the problem and encouraged them to write this article.

Throughout this paper, the notation is as follows. Let $k \geq 2$, $4 \mid N$ be integers, χ a Dirichlet character modulo N s.t. $\chi^2 = 1$, and let $f = \sum_{n=1}^{\infty} a(n)q^n \in S_{k+1/2}(N, \chi)$ be a non-zero cuspidal Hecke eigenform (for operators T_{p^2} for primes $p \nmid N$) of weight $k + \frac{1}{2}$ (see [15], p. 458) with real coefficients. Let $F_t = \sum_{n=1}^{\infty} A_t(n)q^n \in S_{2k}(N/2, \chi^2)$ be the Hecke eigenform (for operators T_p for primes $p \nmid N$) of weight $2k$ corresponding to f under the Shimura lift for a fixed squarefree t such that $a(t) \neq 0$ (see Section 3). We work under the assumption that F_t does not have complex multiplication.

The question about the sign changes of coefficients of half integral weight modular forms has been asked by Bruinier and Kohnen [4], and there it was shown that the sequence $\{a(tn^2)\}_{n \in \mathbb{N}}$ of coefficients of half integral weight modular forms has infinitely many sign changes under the additional hypothesis that a certain L -function has no zeros in the interval $(0, 1)$ (later this hypothesis has been removed in [7]). In addition to this partly conditional result, the authors came up with a suggestion supported by some numerical experiments. They claimed that half of the coefficients are positive among all non-zero coefficients due to observations on computations made for one example of weight $11/2$ given in [10] and for one example of weight $3/2$ given in [5] for indices up to 10^6 . This claim is also supported by five different examples of weight $3/2$ in computations done in the preparation process of [6] for indices up to 10^7 . Kohnen et al. [8] also study the sign change problem on specific sets of integers. They establish lower bounds of the best possible order of magnitude for the number of those coefficients that have the same signs. These give an improvement on some results in [4] and [7]. Although the problem of the equidistribution of signs has been mentioned informally in [4], the conjecture was only given in [8] formally.

In this paper we improve the previously known results towards the Bruinier–Kohnen conjecture by proving the equidistribution of the signs of $\{a(tn^2)\}_n$ for n running through the set of primes (see Theorem 4.1) and through the set of all positive integers (see Corollary 5.2). The former result is formulated in terms of natural density, the latter for Dedekind–Dirichlet density. For the latter result, we have to assume a certain error term for the convergence of the Sato–Tate distribution for integral weight modular forms, which is weaker than an error term conjectured by Akiyama and Tanigawa [1]. Relying on the Shimura lift as in the papers cited above, the techniques of the present article also do not extend to study the equidistribution of the signs of $\{a(n)\}_n$, when n runs through the squarefree integers.

The ideas and techniques of the present paper have been adapted by Kumar [9] to q -exponents of generalised modular functions. In a sequel to this work [2], written together with Sara Arias-de-Reyna, we extend the main theorem to the case when F_t has complex multiplication and we also weaken the assumption on the error bound.

2. Density and regular sets of primes. Let \mathbb{P} denote the set of all prime numbers. We first recall some definitions. Let $S \subseteq \mathbb{P}$ be a set of primes. It is said to have *natural density* $d(S)$ if the limit $\lim_{x \rightarrow \infty} \frac{\pi_S(x)}{\pi(x)}$ exists and is equal to $d(S)$,

where $\pi(x) = \#\{p \leq x \mid p \in \mathbb{P}\}$ is the prime number counting function and $\pi_S(x) := \#\{p \leq x \mid p \in S\}$. The set S is said to have *Dirichlet density* $\delta(S)$ if the limit $\lim_{z \rightarrow 1^+} \frac{\sum_{p \in S} \frac{1}{p^z}}{\log(\frac{1}{z-1})}$ exists and is equal to $\delta(S)$ (the limit $\lim_{z \rightarrow 1^+}$ is defined via sequences of real numbers tending to 1 from the right). If S has a natural density, then it also has a Dirichlet density and the two coincide. As in [12], p. 343f, we call S *regular* if there is a function $g(z)$ holomorphic on $\text{Re}(z) \geq 1$ such that $\sum_{p \in S} \frac{1}{p^z} = \delta(S) \log\left(\frac{1}{z-1}\right) + g(z)$. Note that \mathbb{P} is a regular set of primes of Dirichlet density 1. We collect some more straight forward properties of regular sets of primes in the following lemma.

- Lemma 2.1.**
1. *Let S be any set of primes such that the series $\sum_{p \in S} \frac{1}{p}$ converges to a finite value. Then S has a Dirichlet density equal to 0.*
 2. *Let S be a regular set of primes. Then the Dirichlet density of S is 0 if and only if the series $\sum_{p \in S} \frac{1}{p}$ converges to a finite value.*
 3. *Let S_1, S_2 be two regular sets of primes having the same Dirichlet density $\delta(S_1) = \delta(S_2)$. Then the function $\sum_{p \in S_1} \frac{1}{p^z} - \sum_{q \in S_2} \frac{1}{q^z}$ is analytic on $\text{Re}(z) \geq 1$.*

For our application in Section 5, we need regular sets of primes. Hence, we now include a proposition showing that if a set of primes S has a natural density and the convergence satisfies a certain error term, then it is regular. This may be known, but, we are not aware of any reference; so we include a full proof.

Proposition 2.2. *Let $S \subseteq \mathbb{P}$ have natural density $d(S)$. Let $E(x) := \frac{\pi_S(x)}{\pi(x)} - d(S)$ be the error function. Suppose that there are $\alpha > 0$ and $C > 0$ such that for all sufficiently large x we have $|E(x)| \leq Cx^{-\alpha}$.*

Then S is a regular set of primes.

Proof. We will use the notation $D_S(z) := \sum_{p \in S} \frac{1}{p^z}$ and $D(z) := \sum_{p \in \mathbb{P}} \frac{1}{p^z}$. We abbreviate $d := d(S)$, put $g(x) := E(x)\pi(x)$ and $f(z) := \sum_{n=2}^{\infty} g(n) \left(\frac{1}{n^z} - \frac{1}{(n+1)^z}\right)$. As $g(x)$ is a step function with jumps only at integers, we have

$$f(z) = z \cdot \sum_{n=2}^{\infty} g(n) \int_n^{n+1} \frac{1}{x^{z+1}} dx = z \cdot \int_2^{\infty} \frac{g(x)}{x^{z+1}} dx.$$

From Theorem 29 of [14], we know that $\pi(x) < \frac{x}{\log(x)-4}$ for $x > 55$, yielding $|g(x)| \leq C \cdot \pi(x) \cdot x^{-\alpha} \leq C \cdot \frac{x^{1-\alpha}}{\log(x)-4}$. Thus for $\text{Re}(z) > 1 - \frac{\alpha}{2}$ we get

$$\left| \int_{55}^{\infty} \frac{g(x) dx}{x^{z+1}} \right| \leq \int_{55}^{\infty} \frac{|g(x)|}{x^{\text{Re}(z)+1}} dx \leq C \int_{55}^{\infty} \frac{1}{x^{1+\frac{\alpha}{2}}} dx.$$

As the last integral is convergent, we conclude that $f(z)$ is an analytic function on $\text{Re}(z) \geq 1$. Since for $\text{Re}(z) > 1$ we have

$$D_S(z) = \sum_{n=2}^{\infty} \underbrace{(d\pi(n) + g(n))}_{=\pi_S(n)} \left(\frac{1}{n^z} - \frac{1}{(n+1)^z} \right) = dD(z) + f(z)$$

the proposition follows as \mathbb{P} is regular of density 1. □

Finally we recall a notion of density on sets of natural numbers analogous to that of Dirichlet density. A subset $A \subseteq \mathbb{N}$ is said to have *Dedekind–Dirichlet density* $\delta(A)$ if the limit $\lim_{z \rightarrow 1^+} (z - 1) \sum_{n=1, n \in A}^{\infty} \frac{1}{n^z}$ exists and is equal to $\delta(A)$. If $A \subseteq \mathbb{N}$ has a natural density (defined in the evident way), then it also has a Dedekind–Dirichlet density, and they are the same.

3. Shimura lift and Sato–Tate equidistribution. We continue to use the notation described in the introduction. We now summarise some properties of the Shimura lift that are proved in [15] and Chapter 2 of [13]. There is the direct relation between the Fourier coefficients of f and those of its lift F_t , namely

$$A_t(n) := \sum_{d|n} \chi_{t,N}(d) d^{k-1} a \left(\frac{tn^2}{d^2} \right), \tag{3.1}$$

where $\chi_{t,N}$ denotes the character $\chi_{t,N}(d) := \chi(d) \left(\frac{(-1)^k N^2 t}{d} \right)$. As we assume f to be a Hecke eigenform for the Hecke operator T_{p^2} , F_t is an eigenform for the Hecke operator T_p , for all primes $p \nmid N$. In fact, in this case $F_t = a(t)F$, where F is a normalised Hecke eigenform independent of t . Moreover, from the Euler product formula for the Fourier coefficients of half integral weight modular forms, one obtains the multiplicativity relation for $(m, n) = 1$

$$a(tm^2)a(tn^2) = a(t)a(tm^2n^2). \tag{3.2}$$

Note that the assumption that χ be (at most) quadratic implies that F_t has real coefficients if f does. Furthermore, the coefficients of F_t satisfy the Ramanujan–Pettersson bound $\left| \frac{A_t(p)}{a(t)} \right| \leq 2p^{k-1/2}$. We normalise them by letting

$$B_t(p) := \frac{A_t(p)}{2a(t)p^{k-1/2}} \in [-1, 1].$$

Recall now that we are assuming F_t without complex multiplication. One defines the *Sato–Tate measure* μ to be the probability measure on the interval $[-1, 1]$ given by $\frac{2}{\pi} \sqrt{1-t^2} dt$.

Theorem B of [3], case 3 with $\zeta = 1$, gives the important *Sato–Tate equidistribution theorem* for $\Gamma_0(N)$, which applies in particular to $F = \frac{F_t}{a(t)}$.

Theorem 3.1. (Barnet-Lamb, Geraghty, Harris, Taylor) *Let $k \geq 1$, and let $F = \sum_{n \geq 1} A(n)q^n$ be a normalised cuspidal Hecke eigenform of weight $2k$ for $\Gamma_0(N)$ without complex multiplication. Then the numbers $B(p) = \frac{A_t(p)}{2p^{k-1/2}}$ are μ -equidistributed in $[-1, 1]$ when p runs through the primes not dividing N .*

This has the following simple corollary, which we formulate for the Shimura lift F_t .

Corollary 3.2. *Let $[a, b] \subseteq [-1, 1]$ be a subinterval and $S_{[a,b]} := \{p \text{ prime} \mid p \nmid N, B_t(p) \in [a, b]\}$. Then $S_{[a,b]}$ has natural density equal to $\frac{2}{\pi} \int_a^b \sqrt{1-t^2} dt$.*

4. Equidistribution of signs for $\{a(tp^2)\}_{p \text{ prime}}$. Our main unconditional result is the following theorem.

Theorem 4.1. *We use the assumptions from the introduction, in particular, F_t has no complex multiplication. Define the set of primes*

$$\mathbb{P}_{>0} := \{p \in \mathbb{P} \mid a(tp^2) > 0\}$$

and similarly $\mathbb{P}_{<0}$, $\mathbb{P}_{\geq 0}$, $\mathbb{P}_{\leq 0}$, and $\mathbb{P}_{=0}$ (depending on f and t).

Then the sets $\mathbb{P}_{>0}$, $\mathbb{P}_{<0}$, $\mathbb{P}_{\geq 0}$, $\mathbb{P}_{\leq 0}$ have natural density $1/2$ and the set $\mathbb{P}_{=0}$ has natural density 0 .

Proof. Denote by $\pi_{>0}(x) := \#\{p \leq x \mid p \in \mathbb{P}_{>0}\}$ and similarly $\pi_{<0}(x)$, $\pi_{\geq 0}(x)$, $\pi_{\leq 0}(x)$, and $\pi_{=0}(x)$. Since dividing f by $a(t)$ does not affect the assertions of the theorem, we may and do assume $a(t) = 1$. In that case F_t is a normalised eigenform. Equation (3.1) specialises to $a(tp^2) = A_t(p) - \chi_{t,N}(p)p^{k-1}$ for all primes p , implying the equivalence:

$$a(tp^2) > 0 \Leftrightarrow B_t(p) > \frac{\chi_{t,N}(p)}{2\sqrt{p}}.$$

The idea is to use the Sato–Tate equidistribution to show that $|A_t(p)|$ dominates the term $\chi_{t,N}(p)p^{k-1}$ for ‘most’ primes. Let $\epsilon > 0$. Since for all $p > \frac{1}{4\epsilon^2}$ one has $|\frac{\chi_{t,N}(p)}{2\sqrt{p}}| = \frac{1}{2\sqrt{p}} < \epsilon$, we obtain

$$\pi_{>0}(x) + \pi\left(\frac{1}{4\epsilon^2}\right) \geq \#\{p \leq x \text{ prime} \mid B_t(p) > \epsilon\}. \tag{4.1}$$

By Corollary 3.2, we have $\lim_{x \rightarrow \infty} \frac{\#\{p \leq x \text{ prime} \mid B_t(p) > \epsilon\}}{\pi(x)} = \mu([\epsilon, 1])$. This implies that $\liminf_{x \rightarrow \infty} \frac{\pi_{>0}(x)}{\pi(x)} \geq \mu([\epsilon, 1])$ for all $\epsilon > 0$, whence we can conclude $\liminf_{x \rightarrow \infty} \frac{\pi_{>0}(x)}{\pi(x)} \geq \mu([0, 1]) = \frac{1}{2}$.

A similar argument yields $\liminf_{x \rightarrow \infty} \frac{\pi_{\leq 0}(x)}{\pi(x)} \geq \mu([0, 1]) = \frac{1}{2}$. Using $\pi_{\leq 0}(x) = \pi(x) - \pi_{>0}(x)$ gives $\limsup_{x \rightarrow \infty} \frac{\pi_{>0}(x)}{\pi(x)} \leq \mu([0, 1]) = \frac{1}{2}$, thus showing that $\lim_{x \rightarrow \infty} \frac{\pi_{>0}(x)}{\pi(x)}$ exists and is equal to $\frac{1}{2}$, whence by definition $\mathbb{P}_{>0}$ has natural density $\frac{1}{2}$. The arguments for $\mathbb{P}_{<0}(x)$, $\mathbb{P}_{\geq 0}(x)$, $\mathbb{P}_{\leq 0}(x)$ are exactly the same, and the conclusion for $\mathbb{P}_{=0}$ immediately follows. \square

We next show that the sets of primes in Theorem 4.1 are regular if the Sato–Tate equidistribution converges with a certain error term. A stronger error term was conjectured by Akiyama and Tanigawa (see [11], Conjecture 2.2, and Conjecture 1, on p. 1204 in [1]).

Theorem 4.2. *We make the same assumptions as in Theorem 4.1. We additionally assume that there are $C > 0$ and $\alpha > 0$ such that for all subintervals $[a, b] \subseteq [-1, 1]$ and all sufficiently large x one has*

$$\left| \frac{\#\{p \leq x \text{ prime} \mid \frac{A_t(p)}{a(t)2p^{k-1/2}} \in [a, b]\}}{\pi(x)} - \mu([a, b]) \right| \leq \frac{C}{x^\alpha}.$$

Then the sets $\mathbb{P}_{>0}$, $\mathbb{P}_{<0}$, $\mathbb{P}_{\geq 0}$, $\mathbb{P}_{\leq 0}$, and $\mathbb{P}_{=0}$ are regular sets of primes.

Proof. We start as in the proof of Theorem 4.1 up to Eq. (4.1) and plug in the error term to get for all $\epsilon > 0$ and all sufficiently large x

$$\begin{aligned} \pi_{>0}(x) + \pi\left(\frac{1}{4\epsilon^2}\right) &\geq \#\{p \leq x \text{ prime} \mid B_t(p) > \epsilon\} \\ &\geq -C\pi(x)x^{-\alpha} + \mu([\epsilon, 1])\pi(x) \geq -C\pi(x)x^{-\alpha} + \left(\frac{1}{2} - \mu([0, \epsilon])\right)\pi(x). \end{aligned}$$

Using the estimates $\mu([0, \epsilon]) = \int_0^\epsilon \sqrt{1-t^2} dt \leq \epsilon$ and $\frac{x}{\log(x)+2} \leq \pi(x) \leq \frac{x}{\log(x)-4}$, which is valid for $x > 55$ (see [14] Theorem 29 A on p. 211), we get

$$\frac{\pi_{>0}(x)}{\pi(x)} - \frac{1}{2} \geq -\left(Cx^{-\alpha} + \epsilon + \frac{(\log(x) + 2)}{-4x\epsilon^2(\log(4\epsilon^2) + 4)}\right) \geq -C_1x^{-\alpha}$$

for some $C_1 > 0$ and all x big enough, where for the last inequality we set $\epsilon = x^{-2\alpha}$. The same argument with $\pi_{<0}(x)$ yields

$$\frac{\pi_{<0}(x)}{\pi(x)} - \frac{1}{2} \geq -C_1x^{-\alpha}.$$

Using $\pi_{\geq 0}(x) = \pi(x) - \pi_{<0}(x)$, we get $\frac{\pi_{>0}(x)}{\pi(x)} - \frac{1}{2} \leq \frac{\pi_{\geq 0}(x)}{\pi(x)} - \frac{1}{2} \leq C_1x^{-\alpha}$. Thus, one has

$$\left| \frac{\pi_{>0}(x)}{\pi(x)} - \frac{1}{2} \right| \leq \frac{C_1}{x^\alpha}.$$

Proposition 2.2 now implies that $\mathbb{P}_{>0}$ is a regular set of primes. The regularity of $\mathbb{P}_{<0}$ is obtained in the same way, implying also the regularity of $\mathbb{P}_{=0}$, $\mathbb{P}_{\geq 0}$, and $\mathbb{P}_{\leq 0}$. □

Remark 4.3. Assume the setup of Theorem 4.1. Let $[a, b] \subseteq [-1, 1]$ be a subinterval. Then using the same arguments as in the proof of Theorem 4.1, one can show that the set of primes $\{p \mid \frac{a(tp^2)}{2a(t)p^{k-1/2}} \in [a, b]\}$ has a natural density equal to $\mu([a, b])$. Similarly, a more general version of Theorem 4.2 also holds.

5. Equidistribution of signs for $\{a(tn^2)\}_{n \in \mathbb{N}}$. In this section we prove our equidistribution result for the signs of the coefficients $a(tn^2)$, when n runs through the natural numbers. The same arguments also work for n running through k -free positive integers for any $k \geq 1$.

Theorem 5.1. *We make the same assumptions as in Theorem 4.2. We also assume $a(t) > 0$. We define the multiplicative (but, not completely multiplicative) function*

$$s(n) = \begin{cases} 1 & \text{if } a(tn^2) > 0, \\ -1 & \text{if } a(tn^2) < 0, \\ 0 & \text{if } a(tn^2) = 0. \end{cases}$$

Let $S(z) := \sum_{n=1}^{\infty} \frac{s(n)}{n^z}$ be the Dirichlet series of $s(n)$.

Then $S(z)$ is holomorphic for $\text{Re}(z) \geq 1$.

Proof. Since $s(n)$ is multiplicative because of Eq. (3.2) and the assumption $a(t) > 0$, we have the Euler product $S(z) = \prod_{p \in \mathbb{P}} \sum_{k=0}^{\infty} s(p^k)p^{-kz}$. Taking the logarithm of $S(z)$, we obtain

$$\begin{aligned} \log S(z) &= \sum_{p \in \mathbb{P}_{>0}} \log \left(1 + \frac{1}{p^z} + g(z, p) \right) + \sum_{p \in \mathbb{P}_{<0}} \log \left(1 - \frac{1}{p^z} + g(z, p) \right) \\ &\quad + \sum_{p \in \mathbb{P}=0} \log(1 + g(z, p)), \end{aligned}$$

where $g(z, p) := \sum_{m=2}^{\infty} \frac{s(p^m)}{p^{mz}}$ is a holomorphic function on $\text{Re}(z) \geq 1$ in the variable z for fixed p . Note that for all p , we have $|g(z, p)| \leq \frac{1}{p^z(p^z-1)}$. Using this, we conclude that

$$\log S(z) = \sum_{p \in \mathbb{P}_{>0}} \frac{1}{p^z} - \sum_{p \in \mathbb{P}_{<0}} \frac{1}{p^z} + k(z),$$

where $k(z)$ is a holomorphic function on $\text{Re}(z) \geq 1$.

Since $\mathbb{P}_{>0}$ and $\mathbb{P}_{<0}$ are regular sets of primes having the same density $\frac{1}{2}$ by Theorem 4.2, we can conclude that $\log S(z)$ is analytic on $\text{Re}(z) \geq 1$ by Lemma 2.1. Since the exponential function is holomorphic, we find that $S(z)$ is analytic on $\text{Re}(z) \geq 1$ by taking the exponential of $\log S(z)$, as was to be shown. □

We now deduce our main density statement from Theorem 5.1.

Corollary 5.2. *Assume the setting of Theorem 4.2. Then the sets*

$$\{n \in \mathbb{N} \mid a(tn^2) > 0\} \text{ and } \{n \in \mathbb{N} \mid a(tn^2) < 0\}$$

have equal positive Dedekind–Dirichlet densities, that is, both are precisely half of the Dedekind–Dirichlet density of the set $\{n \in \mathbb{N} \mid a(tn^2) \neq 0\}$.

Proof. We assume without loss of generality $a(t) > 0$ since the statement is invariant under replacing f by $-f$. We use the notation of Theorem 5.1. Since the set of all natural numbers has a Dedekind–Dirichlet density of 1 (since the Riemann-zeta function has a pole of order 1 and residue 1 at 1), we can write

$$\lim_{z \rightarrow 1^+} (z - 1) \sum_{n \in \mathbb{N}} \frac{1}{n^z} = 1.$$

Since $S(z) = \sum_{a(tn^2)>0} \frac{1}{n^z} - \sum_{a(tn^2)<0} \frac{1}{n^z}$ is holomorphic for $\text{Re}(z) \geq 1$ by Theorem 5.1 (in fact, much less suffices), we deduce

$$\lim_{z \rightarrow 1^+} (z - 1) \left[2 \sum_{a(tn^2)>0} \frac{1}{n^z} + \sum_{a(tn^2)=0} \frac{1}{n^z} \right] = 1.$$

Define $t(n) := [s(n)]^2$ and $T(z) := \sum_{n=1}^\infty \frac{t(n)}{n^z}$. Since $t(n)$ is multiplicative, $T(z)$ has an Euler product, which we use now. Put $A(z) := \frac{T(z)}{\zeta(z)}$, then

$$\begin{aligned} A(z) &= \prod_p \left(1 - \frac{1}{p^z} \right) \cdot \prod_p \left(1 + \sum_{n=1, a(tp^{2n}) \neq 0}^\infty \frac{1}{p^z} \right) \\ &= \prod_{p, a(tp^2) \neq 0} \left(\left(1 - \frac{1}{p^z} \right) \left(1 + \frac{1}{p^z} + \sum_{n=2, a(tp^{2n}) \neq 0}^\infty \frac{1}{p^{nz}} \right) \right) \\ &\quad \cdot \prod_{p, a(tp^2) = 0} \left(\left(1 - \frac{1}{p^z} \right) \left(1 + \sum_{n=2}^\infty \frac{1}{p^{nz}} \right) \right) \\ &= \prod_{p, a(tp^2) \neq 0} \left(1 - \frac{1}{p^{2z}} + r_1(z, p) \right) \cdot \prod_{p, a(tp^2) = 0} \left(1 - \frac{1}{p^z} + r_2(z, p) \right), \end{aligned}$$

where $r_1(z, p)$ and $r_2(z, p)$ are the remaining terms. Taking the logarithm of $A(z)$, we conclude that $\sum_{p, a(tp^2) \neq 0} \log \left(1 - \frac{1}{p^z} + r_2(z, p) \right)$ is holomorphic on $\text{Re}(z) \geq 1$ since $\mathbb{P}_{=0}$ is a regular set of primes of density 0, by Theorem 4.2. Moreover, $\sum_{p, a(tp^2) \neq 0} \log \left(1 - \frac{1}{p^{2z}} + r_1(z, p) \right)$ is also holomorphic on $\text{Re}(z) \geq 1$. Taking the exponential shows that $A(z) = \frac{T(z)}{\zeta(z)}$ is holomorphic for $\text{Re}(z) \geq 1$. We obtain $A(1) > 0$ and $\lim_{z \rightarrow 1^+} (z - 1)T(z) = A(1)$. Therefore we conclude that the set $\{n \in \mathbb{N} \mid a(tn^2) \neq 0\}$ has a Dedekind–Dirichlet density equal to $A(1)$. Hence, the limit $\lim_{z \rightarrow 1^+} (z - 1) \sum_{a(tn^2)=0} \frac{1}{n^z} = 1 - A(1)$ exists. So we conclude that

$$\lim_{z \rightarrow 1^+} (z - 1) \sum_{a(tn^2)>0} \frac{1}{n^z} = \frac{A(1)}{2}.$$

This implies that the Dedekind–Dirichlet densities of the two sets in the statement are equal and this completes the proof. □

References

- [1] S. AKIYAMA AND Y. TANIGAWA, Calculation of Values of L - Functions Associated to Elliptic Curves, *Mathematics of Computation* **68** (1999), 1201–1231.
- [2] S. ARIAS-DE-REYNA, I. INAM, AND G. WIESE, On Conjectures of Sato–Tate and Bruinier–Kohnen, preprint (2013), arXiv:1305.5443.
- [3] T. BARNET-LAMB ET AL., A Family of Calabi–Yau varieties and Potential Automorphy II, *Pub. Res. Inst. Math. Sci.* **47** (2011), 29–98.

- [4] J. H. BRUINIER AND W. KOHNEN, Sign Changes of Coefficients of Half Integral Weight Modular Forms, *Modular forms on Schiermonnikoog*, Eds.: B. Edixhoven, G. van der Geer, and B. Moonen, Cambridge University Press (2008), 57–66.
- [5] N. DUMMIGAN, Congruences of Modular Forms and Selmer Groups, *Math. Res. Lett.* **8** (2001), 479–494.
- [6] I. INAM, Selmer Groups in Twist Families of Elliptic Curves, *Quaestiones Mathematicae* **35** (2012), 471–487.
- [7] W. KOHNEN, A Short Note on Fourier Coefficients of Half-Integral Weight Modular Forms, *Int. J. of Number Theory* **06** (2010), 1255–1259.
- [8] W. KOHNEN, Y.-K. LAU, AND J. WU, Fourier Coefficients of Cusp Forms of Half-Integral Weight, *Math. Zeitschrift* **273** (2013), 29–41.
- [9] N. KUMAR, On sign changes of q -exponents of generalized modular functions, *J. Number Theory* **133** (2013), 3589–3597.
- [10] W. KOHNEN AND D. ZAGIER, Values of L -series of Modular Forms at the Center of the Critical Strip, *Invent. Math.* **64** (1981), 175–198.
- [11] B. MAZUR, Finding Meaning in Error Terms, *Bulletin of the Amer. Math. Soc.* **45** (2008), 185–228.
- [12] W. NARKIEWICZ, *Elementary and Analytic Theory of Algebraic Numbers*, Third Edition, Springer, 708 pp., (1990).
- [13] S. NIWA, Modular Forms of Half Integral Weight and the Integral of Certain Theta-Functions, *Nagoya Math. J.* **56** (1974), 147–161.
- [14] B. ROSSER, Explicit Bounds for Some Functions of Prime Numbers, *Amer. Journal of Mathematics* **63** (1941), 211–232.
- [15] G. SHIMURA, On Modular Forms of Half-Integral Weight, *Annals of Math.* **97** (1973), 440–481.

ILKER INAM

Department of Mathematics, Faculty of Arts and Sciences,
Uludag University,
16059 Gorukle/Bursa,
Turkey
e-mail: ilker.inam@gmail.com

GABOR WIESE

Faculté des Sciences, de la Technologie et de la Communication,
Université du Luxembourg,
6, rue Richard Coudenhove-Kalergi,
1359 Luxembourg,
Luxembourg
e-mail: gabor.wiese@uni.lu

Received: 15 May 2013