

Visions & Reflections (Minireview)

The semenogelins: proteins with functions beyond reproduction?

M. Jonsson*, Å. Lundwall and J. Malm

Department of Laboratory Medicine, Section for Clinical Chemistry, Lund University, Malmö University Hospital, 205 02 Malmö (Sweden), Fax: +46 40 33 62 86

Received 21 June 2006; received after revision 16 August 2006; accepted 28 September 2006
Online First 20 November 2006

Abstract. The coagulum proteins of human semen, semenogelins I and II, are secreted in abundance by the seminal vesicles. Their function in reproduction is poorly understood as they are rapidly degraded in ejaculated se-

men. However, more recent results indicate that it is time to put the semenogelins in a broader physiological perspective that goes beyond reproduction and fertility.

Keywords. Semenogelin, zinc, fertility, seminal vesicle, kallikrein, reproduction.

In the 1980s, semenogelin I and semenogelin II were – as their names imply – recognized as the major structural and gel-forming proteins in human semen [1–4]. Studies conducted at that time also showed that the semenogelins are secreted mainly by the seminal vesicles, and they were identified as the major substrates for the kallikrein-like protease called prostate-specific antigen (PSA) [5–7]. Semenogelin I has a molecular mass of 50 kDa, and semenogelin II is a 63-kDa protein that also exists in a glycosylated form [8]. The semenogelins exhibit 78% similarity in their primary structure, and they are composed of repetitive units [9]. The genes that encode these proteins are located 11.5 kbp apart on the long arm of chromosome 20 [10]. The two gene products belong to a family of proteins that are encoded by three exons: the first gives rise to the signal peptide, the second encodes the secreted protein and the third contains the 3' untranslated nucleotides. The first and third exons are conserved between family members, whereas the second is not. Thus, the expressed proteins in this family show wide variation in primary structure. Accordingly, it seems that they have evolved rapidly [11, 12], which would explain why semenogelins I and II are apparently found solely in

primates. Even within that taxonomic group, there are obvious interspecies differences in the molecules [13, 14]. Rats and mice also express seminal vesicle proteins that belong to the same family and show high sequence similarity in the products encoded by exons one and three, whereas the translated middle exon exhibits little similarity to the primate counterpart [12].

It is common with a history of rapid evolution for proteins to be involved in reproduction [15]. Naturally, as major constituents of semen, the semenogelins are closely associated with reproduction, and a large proportion of our knowledge about these proteins consists of results and hypotheses related to fertility [16]. A correlation has been found between the degree of female promiscuity within a species and the evolution of the two genes encoding semenogelins I and II, and thus it has been suggested that post-copulatory sperm competition drives the evolution of these genes [13, 17, 18]. In addition to having coagulum-forming functions and being susceptible to cleavage by kallikrein-like peptidases, the semenogelins are assumed to activate sperm hyaluronidase [19], affect sperm motility [20], possess anti-microbial activity [21], serve as substrates for transglutaminase [22] and have amyloid properties [23]. In short, it is suggested these two proteins participate in several steps that affect the ability of a sper-

* Corresponding author.

matozoon to reach and fuse with an ovum. In the human population, differences in the two semenogelin genes are poorly mapped, and the only studies of genetic anomalies that have been performed have concerned individuals carrying a common variant semenogelin I allele encoding a truncated protein. No homozygotes for that allele have been found, and heterozygosity does not seem to affect fertility [24, 25].

Studies have shown that semenogelins I and II are expressed in tissues unrelated to fertility, e.g. breast, salivary glands, trachea, kidney and intestine [26], and in lung cancer cell lines [27], which indicates that these proteins are involved in functions beyond reproduction. Notably, it was subsequently discovered that the semenogelins represent the major binders of the zinc ions that are found in abundance in semen [28]. Even more recently, it was also reported that the semenogelins are expressed in retina, a tissue known to contain high levels of zinc [29]. The high capacity of the semenogelins for extracellular binding of Zn^{2+} is a function that may have implications throughout the body, and hence it has been proposed that they play a physiological role as modulators of zinc-dependent serine proteases [28]. PSA is the major protease in seminal plasma. It is a member of the tissue kallikrein family and has a unique specificity [30]. The proteolytic activity of PSA is inhibited by zinc ions [30], and experiments *in vitro* have shown that adding semenogelins to a zinc-rich environment such as semen can decrease the amount of free zinc to a level low enough to block the inhibition and thereby enable activation of PSA [28]. However, since the semenogelins are continually broken down into smaller fragments, it seems likely that the number of zinc-binding sites gradually decreases, which would in turn increase the number of zinc ions available for binding to PSA and eventually more or less inhibit the protease. This is an example of a self-limited system in which PSA is maintained in an active state by its substrates [28]. Human kallikrein related peptidases 2 and 5 (KLK2 and KLK5) are other proteins in same family as PSA. Interestingly, their activity is also inhibited by zinc [31, 32], suggesting that these peptidases may interact with Zn^{2+} and the semenogelins in a way similar to PSA. Furthermore, the semenogelins are co-localized with PSA, KLK2 and KLK5 in several tissues outside the reproductive organs, such as breast, salivary gland, trachea and the intestine [26, 33–35]

The two semenogelin genes are surrounded by a locus containing 14 genes that are structurally similar in that they exhibit conservation of the first exon encoding the signal peptide and the last exon containing the 3' untranslated nucleotides. The locus produces small proteins that have been suggested to inhibit serine proteases and that have been found to contain a characteristic structural motif, which was first discovered in whey acidic protein and is now known to consist of four conserved disulfide bridges. The locus includes the genes encoding elafin and

secretory leukocyte protease inhibitor [36, 37], and both those genes have been observed to show homology to exons one and three of the semenogelin genes [38]. The expression patterns of the small serine protease inhibitors are not restricted to the reproductive organs, and it has been suggested that these molecules play a role in the defense against a wide spectrum of microbes. Antimicrobial activity has also been proposed for the semenogelins [21]. Furthermore, the protein eppin, which is encoded in the whey-acidic-protein-type locus, has been identified in complex with semenogelin I both in seminal plasma and on the surface of spermatozoa [39]. Thus it seems that the proteins encoded in this locus are evolutionary and physiologically connected with the semenogelins.

It is plausible that semenogelins I and II, like many other proteins, have several functions. Some of the effects that have already been assigned to the semenogelins, and the role of these proteins in semen may be both essential for and specific to reproduction. Nonetheless, the semenogelins are expressed in non-reproductive organs, they have a high capacity for binding zinc and they may also modulate zinc-dependent proteases outside the reproduction area. It seems that we need to take a wider view of these proteins. Therefore, if we assume that a positive selection pressure underlies the rapid evolution of the semenogelin genes, the findings mentioned might motivate going beyond reproductive biology to search for the factor that provokes such selection. One such function might be as extracellular zinc regulators.

- Lilja, H. and Laurell, C. B. (1985) The predominant protein in human seminal coagulate. *Scand. J. Clin. Lab. Invest.* 45, 635–641.
- Lilja, H., Abrahamsson, P. A. and Lundwall, A. (1989) Semenogelin, the predominant protein in human semen. Primary structure and identification of closely related proteins in the male accessory sex glands and on the spermatozoa. *J. Biol. Chem.* 264, 1894–1900.
- Lilja, H., Laurell, C. B. and Jeppsson, J. O. (1984) Characterization of the predominant basic protein in human seminal plasma, one cleavage product of the major seminal vesicle protein. *Scand. J. Clin. Lab. Invest.* 44, 439–446.
- McGee, R. S. and Herr, J. C. (1987) Human seminal vesicle-specific antigen during semen liquefaction. *Biol. Reprod.* 37, 431–439.
- Lilja, H. and Laurell, C. B. (1984) Liquefaction of coagulated human semen. *Scand. J. Clin. Lab. Invest.* 44, 447–452.
- McGee, R. S. and Herr, J. C. (1988) Human seminal vesicle-specific antigen is a substrate for prostate-specific antigen (or P-30). *Biol. Reprod.* 39, 499–510.
- Lilja, H. (1985) A kallikrein-like serine protease in prostatic fluid cleaves the predominant seminal vesicle protein. *J. Clin. Invest.* 76, 1899–1903.
- Malm, J., Hellman, J., Magnusson, H., Laurell, C. B. and Lilja, H. (1996) Isolation and characterization of the major gel proteins in human semen, semenogelin I. and semenogelin II. *Eur J. Biochem.* 238, 48–53.
- Lilja, H. and Lundwall, A. (1992) Molecular cloning of epididymal and seminal vesicular transcripts encoding a semenogelin-related protein. *Proc. Natl. Acad. Sci. USA* 89, 4559–4563.
- Ulvback, M., Lazure, C., Lilja, H., Spurr, N. K., Rao, V. V., Loffler, C., Hansmann, I. and Lundwall, A. (1992) Gene struc-

- ture of semenogelin I. and II. The predominant proteins in human semen are encoded by two homologous genes on chromosome 20. *J. Biol. Chem.* 267, 18080–18084.
- 11 Lundwall, A. and Lazure, C. (1995) A novel gene family encoding proteins with highly differing structure because of a rapidly evolving exon. *FEBS Lett.* 374, 53–56.
 - 12 Lundwall, A. (1996) The cloning of a rapidly evolving seminal-vesicle-transcribed gene encoding the major clot-forming protein of mouse semen. *Eur. J. Biochem.* 235, 424–430.
 - 13 Jensen-Seaman, M. I. and Li, W. H. (2003) Evolution of the hominoid semenogelin genes, the major proteins of ejaculated semen. *J. Mol. Evol.* 57, 261–270.
 - 14 Ulvsback, M. and Lundwall, A. (1997) Cloning of the semenogelin I. I. gene of the rhesus monkey. Duplications of 360 bp extend the coding region in man, rhesus monkey and baboon. *Eur. J. Biochem.* 245, 25–31.
 - 15 Birkhead, T. R. and Pizzari, T. (2002) Postcopulatory sexual selection. *Nat. Rev. Genet.* 3, 262–273.
 - 16 Robert, M. and Gagnon, C. (1999) Semenogelin I: a coagulum forming, multifunctional seminal vesicle protein. *Cell. Mol. Life Sci.* 55, 944–960.
 - 17 Dorus, S., Evans, P. D., Wyckoff, G. J., Choi, S. S. and Lahn, B. T. (2004) Rate of molecular evolution of the seminal protein gene SEMG2 correlates with levels of female promiscuity. *Nat. Genet.* 36, 1326–1329.
 - 18 Kingan, S. B., Tatar, M. and Rand, D. M. (2003) Reduced polymorphism in the chimpanzee semen coagulating protein, semenogelin I. *J. Mol. Evol.* 57, 159–169.
 - 19 Mandal, A. and Bhattacharyya, A. K. (1995) Sperm hyaluronidase activation by purified predominant and major basic human seminal coagulum proteins. *Hum. Reprod.* 10, 1745–1750.
 - 20 Robert, M. and Gagnon, C. (1996) Purification and characterization of the active precursor of a human sperm motility inhibitor secreted by the seminal vesicles: identity with semenogelin. *Biol. Reprod.* 55, 813–821.
 - 21 Bourgeon, F., Evrard, B., Brillard-Bourdet, M., Colleu, D., Jegou, B. and Pineau, C. (2004) Involvement of semenogelin-derived peptides in the antibacterial activity of human seminal plasma. *Biol. Reprod.* 70, 768–774.
 - 22 Peter, A., Lilja, H., Lundwall, A. and Malm, J. (1998) Semenogelin I. and semenogelin I. I., the major gel-forming proteins in human semen, are substrates for transglutaminase. *Eur. J. Biochem.* 252, 216–221.
 - 23 Linke, R. P., Joswig, R., Murphy, C. L., Wang, S., Zhou, H., Gross, U., Rocken, C., Westermarck, P., Weiss, D. T. and Solomon, A. (2005) Senile seminal vesicle amyloid is derived from semenogelin I. *J. Lab. Clin. Med.* 145, 187–193.
 - 24 Lundwall, A., Giwercman, A., Ruhayel, Y., Giwercman, Y., Lilja, H., Hallden, C. and Malm, J. (2003) A frequent allele codes for a truncated variant of semenogelin I., the major protein component of human semen coagulum. *Mol. Hum. Reprod.* 9, 345–350.
 - 25 Miyano, S., Yoshida, K., Yoshiike, M., Miyamoto, C., Furuichi, Y. and Iwamoto, T. (2003) A large deletion of the repeat site in semenogelin I. is not involved in male infertility. *Int. J. Mol. Med.* 11, 435–440.
 - 26 Lundwall, A., Bjartell, A., Olsson, A. Y. and Malm, J. (2002) Semenogelin I. and I. I., the predominant human seminal plasma proteins, are also expressed in non-genital tissues. *Mol. Hum. Reprod.* 8, 805–810.
 - 27 Rodrigues, R. G., Panizo-Santos, A., Cashel, J. A., Krutzsch, H. C., Merino, M. J. and Roberts, D. D. (2001) Semenogelins are ectopically expressed in small cell lung carcinoma. *Clin. Cancer Res.* 7, 854–860.
 - 28 Jonsson, M., Linse, S., Frohm, B., Lundwall, A. and Malm, J. (2005) Semenogelins I. and I. I. bind zinc and regulate the activity of prostate-specific antigen. *Biochem. J.* 387, 447–453.
 - 29 Bonilha, V. L., Rayborn, M. E., Shadrach, K., Ake, L., Malm, J., Bhattacharya, S. K., Crabb, J. W. and Hollyfield, J. G. (2006) Characterization of semenogelin proteins in the human retina. *Exp. Eye Res.* 83, 120–127.
 - 30 Malm, J., Hellman, J., Hogg, P. and Lilja, H. (2000) Enzymatic action of prostate-specific antigen (PSA or hK3): substrate specificity and regulation by Zn(2+), a tight-binding inhibitor. *Prostate* 45, 132–139.
 - 31 Lovgren, J., Airas, K. and Lilja, H. (1999) Enzymatic action of human glandular kallikrein 2 (hK2). Substrate specificity and regulation by Zn2+ and extracellular protease inhibitors. *Eur. J. Biochem.* 262, 781–789.
 - 32 Michael, I. P., Pampalakis, G., Mikolajczyk, S. D., Malm, J., Sotiropoulou, G. and Diamandis, E. P. (2006) Human tissue kallikrein 5 is a member of a proteolytic cascade pathway involved in seminal clot liquefaction and potentially in prostate cancer progression. *J. Biol. Chem.* 281, 12743–12750.
 - 33 Yousef, G. M. and Diamandis, E. P. (1999) The new kallikrein-like gene, KLK-L2. Molecular characterization, mapping, tissue expression, and hormonal regulation. *J. Biol. Chem.* 274, 37511–37516.
 - 34 Brattsand, M. and Egelrud, T. (1999) Purification, molecular cloning, and expression of a human stratum corneum trypsin-like serine protease with possible function in desquamation. *J. Biol. Chem.* 274, 30033–30040.
 - 35 Olsson, A. Y., Bjartell, A., Lilja, H. and Lundwall, A. (2005) Expression of prostate-specific antigen (PSA) and human glandular kallikrein 2 (hK2) in ileum and other extraprostatic tissues. *Int. J. Cancer* 113, 290–297.
 - 36 Clauss, A., Lilja, H. and Lundwall, A. (2005) The evolution of a genetic locus encoding small serine proteinase inhibitors. *Biochem. Biophys. Res. Commun.* 333, 383–389.
 - 37 Clauss, A., Lilja, H. and Lundwall, A. (2002) A locus on human chromosome 20 contains several genes expressing protease inhibitor domains with homology to whey acidic protein. *Biochem. J.* 368, 233–242.
 - 38 Lundwall, A. and Ulvsback, M. (1996) The gene of the protease inhibitor SKALP/elafin is a member of the REST gene family. *Biochem. Biophys. Res. Commun.* 221, 323–327.
 - 39 Wang, Z., Widgren, E. E., Sivashanmugam, P., O’Rand, M. G. and Richardson, R. T. (2005) Association of eppin with semenogelin on human spermatozoa. *Biol. Reprod.* 72, 1064–1070.

