



Structural disorder in plant proteins: where plasticity meets sessility

Alejandra A. Covarrubias¹ · Cesar L. Cuevas-Velazquez¹ ·
Paulette S. Romero-Pérez¹ · David F. Rendón-Luna¹ ·
Caspar C. C. Chater¹

Received: 17 May 2017 / Accepted: 1 June 2017 / Published online: 22 June 2017
© Springer International Publishing AG 2017

Abstract Plants are sessile organisms. This intriguing nature provokes the question of how they survive despite the continual perturbations caused by their constantly changing environment. The large amount of knowledge accumulated to date demonstrates the fascinating dynamic and plastic mechanisms, which underpin the diverse strategies selected in plants in response to the fluctuating environment. This phenotypic plasticity requires an efficient integration of external cues to their growth and developmental programs that can only be achieved through the dynamic and interactive coordination of various signaling networks. Given the versatility of intrinsic structural disorder within proteins, this feature appears as one of the leading characters of such complex functional circuits, critical for plant adaptation and survival in their wild habitats. In this review, we present information of those intrinsically disordered proteins (IDPs) from plants for which their high level of predicted structural disorder has been correlated with a particular function, or where there is experimental evidence linking this structural feature with its protein function. Using examples of plant IDPs involved in the control of cell cycle, metabolism, hormonal signaling and regulation of gene expression, development and responses to stress, we demonstrate the critical importance of IDPs throughout the life of the plant.

Keywords Intrinsically disordered proteins · Plant development · Plant metabolism · Plant stress responses · Plant signaling · LEA proteins · Transcription factors

Introduction

Throughout evolution plants have developed an extraordinary ability to overcome fluctuating and drastic environmental changes. Their sessile nature has imposed the selection of particular defense strategies allowing them efficient and effective adjustment or acclimation responses to these conditions, as well as skilled mechanisms to tolerate and survive them. The different endurance strategies selected in these organisms are the result of complex structural and interconnected regulatory networks, which have evolved in an intimate relationship with developmental programs. For instance, in many plant species, the reproductive stage waits for favorable climatic conditions to instrument a crucial set of processes for their perpetuation; root architecture modifies according to the availability of water, phosphorus and other nutrients; and orthodox seeds once desiccated can remain dormant for many years without significant loss in viability until they find sufficient water to germinate [1, 2]. This outstanding resourcefulness suggests mechanisms that make them capable of detecting diverse changes in the plant cell milieu, imposed by the external environment or by developmental programs.

Many molecular response mechanisms are efficiently adapted for rapid detection of subtle environmental fluctuations, as can be observed in mechano-sensitivity, ion channels, proton pumps and post-translational protein modifications. The modification of protein structure also seems to be an efficient and effective transducer of a great diversity of signals. This phenomenon is commonly

✉ Alejandra A. Covarrubias
crobles@ibt.unam.mx

¹ Departamento de Biología Molecular de Plantas, Instituto de Biotecnología, Universidad Nacional Autónoma de México, 62250 Cuernavaca, Mexico

associated with changes in protein conformation produced by phosphorylation or acetylation, or by interactions between proteins or other partners such as nucleic acids or other small molecules acting as substrates, cofactors, and allosteric regulators [3, 4]. However, an adaptation that has received less attention is that related to the intrinsic plasticity found in those proteins that have the ability to present different transient structures depending on the nature of their surroundings.

During the last decade, we have witnessed significant advances in the identification and characterization of many proteins showing intrinsic structural disorder. This has increased our knowledge of their functional relevance, structural properties and dynamics, as well as mechanisms of action (For review see [5–7]). Intrinsically disordered proteins (IDPs) are widely distributed in all domains of life. Although only a few complete proteomes from the different domains are currently available, various bioinformatic studies agree that Eukaryota proteomes show a higher average of disorder, compared to those of Bacteria, which in turn present higher disorder than those of Archaea. Interestingly, the predicted disorder in eukaryote proteomes spans a broad range of score values, with both very low and very high disorder [8, 9]. Overall, current information indicates that the level of disorder is higher in eukaryotic organisms than in prokaryotes. Even more important is the observation that protein superfamilies, which have undergone massive diversification during evolution present more structural disorder than other families. These data also correlate with the expansion of the number of cell types in an organism, revealing a positive relationship between proteome disorder and organism complexity [10, 11].

The accumulated knowledge on IDPs has revealed their functional versatility resulting from their peculiar properties. For example, IDPs can form ensembles with different structural conformations, allowing variability in the exposed surfaces [7, 12–14]. This structural plasticity confers to IDPs the ability to differentially exhibit different post-translational modification sites and/or recognition motifs, depending on specific conditions to interact transiently, but specifically with proteins or nucleic acids. With this in mind, it is not surprising the central roles that IDPs play in cellular functions, achieving regulatory and signaling roles as well as acting as scaffold or assembly proteins.

In this review, we present a general panorama of the available knowledge on protein disorder in plants. We have put together this information in the context of fundamental biological processes such as development, metabolism and stress responses, which in spite of the limited number of studies unveil the functional relevance of these proteins in the life of plants. The different IDPs referred to in this work are compiled in Tables 1 and 2.

IDPs distribution in plants

In recent years, the discovery and characterization of proteins with different amounts of structural disorder has revealed their high representation in plants [15–18]. Large-scale analysis of IDPs and intrinsically disordered regions (IDRs) in *Arabidopsis thaliana*, a widely used experimental model in plant biology, has shown that approximately 30% of its proteome is mostly disordered [10, 16], whereas *Zea mays* and *Glycine max* proteomes contain an even higher proportion of disorder (~50%) [19]. Interestingly, the chloroplast and mitochondrial proteomes show a significantly lower occurrence of disorder (between 2 and 19%) when compared to nuclear proteomes of different plant species. The abundance of disorder in these organellar proteomes is comparable to that of *Archaea* and bacteria, in accordance with the bacterial origin of the genes encoding their proteins [20]. The IDPs encoded in these organellar genomes are mostly involved in translation, transcription or RNA biosynthesis, and some are structural constituents of ribosomes, having in common the ability to form large complexes or to interact with numerous partners as expected from their intrinsic structural flexibility [5, 20]. It is interesting to note that for those proteins with paralogues of nuclear origin, both copies tend to show similarly low levels of disorder, suggesting again a common extra-nuclear origin or functional constraints [20]. Furthermore, the recent data obtained from the examination of the distribution of genes encoding IDPs in the genomes of *A. thaliana* and *Oryza sativa* indicate that they are not randomly arranged and that their organization may result from high recombination rates and chromosomal rearrangements. These observations are in accordance with the location of genes for proteins with highly disordered content within recombination hotspots and possessing high G + C content; this codon usage related to the over-representation of specific amino acid residues in IDPs (e.g., Arg, Gly, Ala and Pro) [19].

In silico analyses of the *Arabidopsis* proteome and of proteins from other plant species have found that IDPs are highly represented in functions related to cell cycle, nucleic acid metabolism, protein synthesis, hormone signaling and regulation of gene expression, development and responses to stress [16, 17, 19, 21–23]. This last functional category seems to be particularly associated with plant IDPs, including proteins involved in detection and signaling of external stimuli, chaperone activities and secondary metabolism; all essential functions for the phenotypic plasticity needed for plant adaptation and survival, as will be further discussed in this review. It should be noted that the examples we present in this review cannot be unequivocally classified and they may belong to several functional classes, which alludes to their functional promiscuity.

IDPs in plant development

The study of plant development and the characterization of the mechanisms involved have identified many proteins playing major control roles in this process. Further detailed analyses have revealed the presence of IDRs in some of these proteins. Germination and early seedling development [24], adventitious shoot formation [25], xylem development [26], photomorphogenesis [27], phytohormone signaling and response [28], flowering [29], and vegetative and reproductive growth [30] are some of the processes where IDR-containing proteins appear as key players. Interestingly, the structural plasticity arising from IDRs of several of these IDPs has been shown to be essential for proper function.

TCP (TB1-CYC-PCF1) transcription factors

The appropriate development and function of vegetative (leaves, shoot and roots) and reproductive (flowers) organs is orchestrated by several proteins, which are subjected to adaptable but precise spatio-temporal control, resulting in a timely fine-tuning of cell proliferation, expansion and differentiation [31]. Many of these proteins are transcription factors, some of which contain IDRs of significant length, that by interacting with other proteins and/or binding to DNA decode a specific signal in the activation or repression of gene expression. The TCP [from TEOSINTE BRANCHED1 (TB1), CYCLOIDEA (CYC) and PROLIFERATING CELL NUCLEAR ANTIGEN FACTOR1 (PCF1)] protein family consists of plant-specific transcription factors involved in plant shape developmental control. Bioinformatic analyses have shown that these transcription factors are IDPs [30, 32]. TCPs are classified as class I or class II according to the characteristics of their conserved and non-canonical basic helix-loop-helix (bHLH) DNA-binding domain [30, 32]. Class I TCP transcription factors participate in organ shape and growth, pollen development, germination, and inflorescence and flower development [33]. Class II TCPs, in addition to their redundant function in the regulation of lateral organ morphogenesis, also participate in endosperm, cotyledon, leaf, petal and stamen development, as well as other aspects of plant development and other processes [33]. Some of the functions assigned to TCP transcription in plant growth and development are a consequence of their involvement in the biosynthesis of some phytohormones, such as brassinosteroids and jasmonic acid, and other metabolites with biological activity such as flavonoids [21]. Analysis of the 24 *Arabidopsis* TCP protein sequences has shown a differential structural disorder content between the two TCP classes; with class I being more disordered than class II

[30]. Biochemical analysis of TCP8, a class I TCP shows three IDRs of more than 50 residues in length containing a cluster of serine residues, at least one of which is phosphorylated [30]. In addition, the IDR located in the TCP8 C-terminal region corresponds to a trans-activation domain (TAD), which is required for the formation of high-order TCP8 homo-oligomers [30]. The identification of molecular recognition features (MoRFs) in the TCP8 TAD [30] and evidence of its requirement to bind TCP15 and PNM1, a pentatricopeptide repeat protein [34], are consistent with TCPs' function as mediators of different stimuli or signals (Fig. 1a). Furthermore, they demonstrate the importance of IDRs as protein domains able to confer the ability to recognize various different partners, a feature needed for precise and flexible control.

NAC (NAM-ATA-CUC2) transcription factors

Another fundamental aspect of plant development is the maintenance of the shoot apical meristem (SAM). NAC (NAM/ATAF/CUC2) transcription factors constitute one of the largest families described in plants that, in addition to their involvement in other processes, control key aspects of SAM maintenance [35]. A conserved and folded DNA-binding domain defines these transcription factors; however, an additional feature of some NAC transcription factors is the presence of a variable and disordered TAD [24]. This characteristic has been experimentally confirmed for several NAC TADs [21, 36], such as ANAC019, involved in germination and early seedling development; HvNAC005 and HvNAC013, in senescence; NTL8, ANAC013, NAP, ANAC046 and SOG1 in germination and senescence; CUC1 in adventitious shoot formation and ANAC012 in xylem fiber development [24, 25, 37, 38]. It is known that TADs from HvNAC013 and ANAC046 interact with the RST (RCD1-SRO-TAF4) multi-binding domain of the hub protein RCD1 (RADICAL-INDUCED CELL DEATH1), a regulator of developmental, hormonal and stress responses [37]. Differing from the folding-upon-binding phenomenon, no structural rearrangement of the two disordered TADs occur upon binding to RCD1 [37], indicating that these ensembles might function as fuzzy complexes.

Elongated hypocotyl (HY5), bZIP transcription factor

Light is absolutely required for plant life. The presence or absence of light causes developmental reprogramming. The light-dependent modulation of plant development is known as photomorphogenesis. This developmental program leads to cotyledon expansion, hypocotyl shortening and chloroplast development [27]. HY5 (Elongated hypocotyl) is a

Table 1 Plant intrinsically disordered proteins involved in development, metabolism and stress response

Biological Process	Protein	Functional role	Organism	Reference
Development	TCP8	TAD and formation of high order oligomers	<i>Arabidopsis thaliana</i>	Valsecchi et al. [30]
	HvNAC005	Senescence	<i>Hordeum vulgare</i>	Kjaersgaard et al. [37]
	HvNAC013			
	NTL8	Germination and Senescence	<i>Arabidopsis thaliana</i>	O'Shea et al. [38]
	ANAC013			
	NAP			
	ANAC046			
	SOG1			
	ANAC019	Germination and earlyseedling development		
	HDC1	Flowering and petiole development	<i>Arabidopsis thaliana</i>	Perrella et al. [53]
	HY5	Photomorphogenesis	<i>Arabidopsis thaliana</i>	Yoon et al. [27]
	BRI1	Brassinosteroid signaling	<i>Arabidopsis thaliana</i>	Hothorn et al. [60]
	BKI1	Brassinosteroid signaling	<i>Arabidopsis thaliana</i>	Jiang et al. [61]
	LD	Flowering	<i>Arabidopsis thaliana</i>	Chakrabortee et al. [29]
	AtRGL1n	GA signaling	<i>Arabidopsis thaliana</i>	Sun et al. [70]
	AtRGL2n			
	AtRGL3n			
	AtGAI			
	AtRGA			
	RHT1n	GA signaling	<i>Triticum aestivum</i>	Sun et al. [70]
CRY1	Blue light signaling	<i>Arabidopsis thaliana</i>	Brautigam et al. [47]	
CRY2		<i>Arabidopsis thaliana</i>	Partch et al. [48]	
Metabolism	CP12	Regulation of GAPDH and PRK activities	<i>Chlamydomonas reinhardtii</i> <i>Arabidopsis thaliana</i> <i>Chlamydomonas reinhardtii</i> <i>Chlamydomonas reinhardtii</i> <i>Arabidopsis thaliana</i> <i>Chlamydomonas reinhardtii</i>	Erales et al. [92] Fermani et al. [96] Mileo et al. [97] Marri et al. [99] Moparthi et al. [98]
	MSP	Regulation and stability of cofactors needed for correct functioning of PSII	<i>Spinacia oleracea</i> <i>Spinacia oleracea</i> <i>Spinacia oleracea</i>	Lydakiss-Simantiris et al. [118] Shutova et al. [116] Popelkova et al. [119]
	BTPC	Catalyzes the addition of bicarbonate to phosphoenolpyruvate	<i>Ricinus communis</i>	Dalziel et al. [126]
	Alb3	Recruits cpSRP to the thylakoid membrane	<i>Arabidopsis thaliana</i>	Falk et al. [120]
	UreG	Chaperone needed to activate urease	<i>Glycine max</i>	Real-Guerra et al. [130]
	Jaburetox	Insecticidal popypeptide	<i>Canavalia ensiformis</i>	Lopes et al. [136]
	RuBisCO	Activates RuBisCO	<i>Arabidopsis thaliana</i> <i>Nicotiana tabacum</i> <i>Arabidopsis thaliana</i> <i>Nicotiana tabacum</i> <i>Camelina sativa</i> <i>Arabidopsis thaliana</i>	Zhang & Portis [108] Stotz et al. [109] Carmo-Silva et al. [112] Wang & Portis [111]

Table 1 continued

Biological Process	Protein	Functional role	Organism	Reference
Stress Response	GR-RBP	RNA chaperone	<i>Arabidopsis thaliana</i>	Kwak et al. [237]
			<i>Nicotiana tabacum</i>	Khan et al. [246]
	VIPPI	Plastid vesicle formation and thylakoid membrane biogenesis	<i>Arabidopsis thaliana</i>	McDonald et al. [259]
			<i>Arabidopsis thaliana</i>	Zhang et al. [260]
	DREB2A	Transcription factor	<i>Arabidopsis thaliana</i>	Kragelund et al. [21]
			<i>Arabidopsis thaliana</i>	Blomberg et al. [262]
			<i>Arabidopsis thaliana</i>	O'Shea et al. [264]
	bZIP28	Transcription factor in the unfolded protein response	<i>Arabidopsis thaliana</i>	Srivastava et al. [270]
	OsWAK14	Positively regulates resistance to <i>Magnaporthe oryzae</i>	<i>Oryza sativa</i>	Delteil et al. [273]
	OsWAK92			
OsWAK91	Positively regulates resistance to <i>Magnaporthe oryzae</i> . Required for H ₂ O ₂ production			
OsWAK112d	Negatively regulates resistance to <i>Magnaporthe oryzae</i>			
Hsp90	Helps to form the Hsp90-Sgt1-CHORD complex involved in viral resistance	<i>Hordeum vulgare</i>	Zhang et al. [281]	

bZIP transcription factor that positively regulates photomorphogenesis [39]. Disorder within the N-terminal region of HY5, responsible for the interaction with its negative regulator COP1, a multifunctional E3 ubiquitin ligase, has been demonstrated by various biophysical methods including limited proteolysis, mass spectrometry, circular dichroism (CD) and nuclear magnetic resonance (NMR) [27]. It is proposed that this disordered character might modulate the interaction with its partners, although functional characterization is still needed.

Cryptochromes (CRYs), blue light receptors

Plants are able to sense light quality (or wavelength) using different proteins such as phytochromes, phototropins and cryptochromes (CRY). Cryptochromes are blue light receptors that control developmental processes such as seedling de-etiolation, growth by elongation and initiation of flowering [40, 41]. CRYs consist of two domains: a conserved light-sensing N-photolyase-homologous region (PHR) of about 500 residues, and a C-terminal tail of variable sequence and length (CRY C-terminal Extension, CCE) [42, 43]. The CCE tail interacts with the PHR domain in a globular well-defined structure. Light activation of the *Arabidopsis* receptors CRY1 and CRY2 releases the CCE tail from the PHR, inducing the unfolding of the

tail and allowing the interaction of both the PHR and the CCE with other proteins (e.g., COP1 and SPA1, a suppressor of phytochrome A1) to promote the blue light signal transduction pathway [44–46]. The light-induced disordered state of CRY receptors has been characterized by several biophysical methods such as limited proteolysis, CD, NMR and X-ray crystallography [47–49]. It is possible that plant CRYs use their disordered CCE region to efficiently recognize diverse binding partners through high-specificity/low-affinity interactions, potentially expanding the repertoire of plant signaling pathways coordinated by light [17].

HDC1 (histone deacetylase complex 1)

Regulation of chromatin accessibility is an important event of gene expression control, fundamental in developmental processes to fulfill the cell requirements within its organismal context. This process depends on the action of multiprotein complexes that control different modifications in DNA and histones [50]. One of these complexes is the histone deacetylase complex (HDAC), which in plants consists of histone deacetylases, co-repressors and histone-binding proteins [51]. HDC1 (HISTONE DEACETYLASE COMPLEX1) is a protein component of *Arabidopsis* HDAC containing a disordered N-terminal region [52, 53].




Fig. 1 Schematic representation of two examples of plant proteins containing IDRs that participate in developmental and metabolic processes. **a** TCP8 is a plant-specific transcription factor involved in plant shape developmental control. TCP8 contains three IDRs (represented by *curved lines*). In these IDRs, there are conserved serine residues, from which at least one is phosphorylated (fill blue *small circle* in the *middle IDR*). The IDR at the C-terminal region corresponds to a trans-activation domain (TAD) required for the formation of TCP8 homo-oligomers. This TAD is also required to bind different partners, such as TCP15 or PNM (*red irregular oval*). The IDR at the amino-terminal region (*purple irregular line*) is part of the TCP8 DNA-binding domain; this disordered region gains structure when TCP8 binds to DNA. **b** CP12 plays a key role in the regulation of the Calvin cycle by translating changes in light

availability into the modulation of GAPDH and PRK enzyme activities. CP12 is a scaffold protein (represented by *curved lines* at the *top* of this panel) that forms a ternary complex with GAPDH (*blue and red irregular ovals*) and PRK (*brown irregular oval*) (GAPDH-CP12-PRK) (represented by the association of the three components at the *bottom* of the panel). During the formation of the GAPDH-CP12-PRK complex, GAPDH associates with CP12 by conformational selection. Upon this interaction, the CP12 N-terminal region remains in a fuzzy state, serving as a linker that facilitates the interaction with PRK. Once the complex is formed, it dimerizes to form a native complex in which there are two dimers of PRK, two tetramers of GAPDH and two monomers of CP12 (figure at the *bottom right* of this panel). Using this mechanism, it seems that CP12 is able to modulate GAPDH and PRK activities

Interestingly, an HDC1 knockout mutant shows impaired leaf growth and delayed flowering, demonstrating its participation in plant development [52]. As expected for an IDP, HDC1 interacts with a wide variety of partners (HDA6, HDA19, SNL3, SNL2, SAP18, ING2 and MSI1) [53]. Deletion of the N-terminal disordered region considerably weakens HDC1 interaction with those proteins. This result together with evidence obtained from complementation experiments shows that the HDC N-terminal IDR plays a significant role in the coordination of flowering and petiole development [53].

BRI1 and BKI1, brassinosteroids signaling proteins

Brassinosteroids (BRs) are plant hormones that control a variety of growth and developmental processes, such as vascular differentiation, leaf development, stem elongation, flowering, senescence, stomatal development and male fertility [54–56]. BRs are perceived at the cell surface by

BRI1 (BRASSINOSTEROID INSENSITIVE 1), a leucine-rich repeat receptor-like kinase (LRR-RLK) and its co-receptor BAK1 (BRI1-ASSOCIATED RECEPTOR KINASE 1) [57]. In the absence of BRs, the cytosolic kinase activity of BRI1 is maintained at low levels by auto-inhibition through its C terminus and by interacting with the repressor protein BKI1 (BRI1 KINASE INHIBITOR 1) [58]. When BRs bind to the extracellular domain of BRI1, the intracellular kinase domain is activated through auto- and trans-phosphorylation. BKI1 is then phosphorylated by BRI1 and released to the cytosol [59]. In contrast to animal LRR toll-like receptors, the extracellular region of the BR receptor contains a superhelix of twenty-five twisted LRRs; moreover, a ~70 amino acid ‘island’ domain has been localized between LRRs 21 and 22, which together constitute a hormone binding region. BR binding causes a conformational change in the BRI1 receptor that leads to its auto-phosphorylation. Remarkably, the ‘island’ domain connects to the LRR core through two long-disordered

loops that become fully ordered upon binding to the steroid ligand. This makes the receptor competent to interact with other proteins, a conversion that may be necessary for receptor activation. It has been proposed that the BRI1 IDR may be an LRR receptor adaptation for efficient detection of small ligands [60]. Further participation of protein structural disorder is evident in this BR sensing protein ensemble, as the BKI1 C-terminal region presents high levels of disorder, particularly, at the BRI1 interacting motif (BIM). It is interesting to note that even though angiosperm BKI1 orthologues are highly diverse, the BIM IDR shows a high degree of conservation [61]. This, together with the finding that the absence of the IDR leads to increased BR sensitivity, establishes its relevance in BR signaling in plants [61].

Luminidependens (LD), a plant prion

The most diverse group of plants corresponds to the flowering plants (angiosperms). Flowering needs to be precisely controlled to generate flowers in an optimal time frame, where environmental conditions match with the presence of pollinators to promote fertilization and reproduction processes [62]. Flowering often follows vernalization, a process achieved after a prolonged period of cold (winter), which ensures flowering in the spring [63]. Interestingly, Chakrabortee and collaborators found that a high proportion of proteins related to flowering in *Arabidopsis* are predicted to contain prion-like domains (PrDs) [29]. Some of these proteins are involved in transcription or regulation of RNA stability in the autonomous flowering pathway: Luminidependens (LD), Flowering Locus PA (FPA), Flowering Locus Y (FY) and Flowering Locus CA (FCA) [29]. Prions are proteins that retain the molecular memory of the cell because they are able to adopt different conformations and can be self-perpetuating [64]. PrDs are enriched in glutamine, asparagine, glycine, proline, serine and tyrosine and it has been shown that they are intrinsically disordered [65, 66]. LD is the first protein reported to have prion-like properties in plants, and can fully complement the activity of the Sup35 PrD, a well-characterized yeast prion [29]. As expected for a prion-like protein, LD protein shows a high level of structural disorder (64.6% according to PONDR, this work) [67], indicating that it is an IDP, even though this property has not been experimentally tested. Notably, LD orthologues from different plant species (*Z. mays*, *O. sativa*, *Phaseolus vulgaris* and *Physcomitrella patens*) also show a high percentage of disorder (51–66%, this work) [67]. As mentioned above, LD, along with a substantial percentage of *Arabidopsis* PrD-containing proteins, participates in flowering processes. This suggests that these proteins may play adaptive roles in the plant environmental memory required for fast

responses to changing conditions, fine-tuning reproductive functions and consequently plant species preservation.

GRAS (GAI-RGA-SCR) transcription factors

The plant-specific GRAS [GIBBERELIC ACID INSENSITIVE (GAI), REPRESSOR OF GAI (RGA), SCARECROW (SCR)] protein family is essential in diverse developmental processes, acting as integrators of signals from different plant growth regulatory inputs (for an extensive review refer to 68). GRAS proteins modulate gene expression through interaction with different transcription factors, thereby controlling their activities. Along with the conserved and folded GRAS domain, GRAS proteins are characterized by a disordered N-domain enriched in MoRFs [69]. Remarkably, the predicted MoRFs exclusively reside in the N-domain conserved motifs that define each subfamily, suggesting that structural disorder permits interactions with different proteins [17]. As has been established for other unstructured proteins, GRAS IDRs containing MoRFs experience disorder-to-order transitions when interacting with their ligands [17, 68–70]. GRAS proteins are classified in ten subfamilies. One of these subfamilies, composed of DELLA (Asp-Glu-Leu-Leu-Ala) proteins, is particularly important for hormonal regulation because DELLA proteins participate as negative regulators of gibberellic acid (GA)-induced plant growth. These are negatively regulated under increasing GA, as GA binds to its receptor (GID), prompting the interaction of the GID-GA complex with the disordered N-domain of DELLAs. This, in turn, promotes the degradation of the DELLA proteins through the ubiquitin–proteasome pathway, resulting in derepression of plant growth [71]. This interaction is mediated by the conserved DELLA and VHYNP motifs localized in an IDR that upon binding to the GID1/GA complex, experiences a disorder-to-order transition [70]. The participation of GRAS IDRs in this signaling pathway highlights their prevalence and function among hub network proteins, operating as integrators of environmental and developmental cues in plants.

MAP65-1, a microtubule associated protein

MAP65-1 is a microtubule (MT)-bundling protein implicated in central spindle formation and cytokinesis in animals, yeast and plants [72]. The *Arabidopsis* genome has nine genes encoding MAP65 proteins [73]. All these proteins have an N-terminal dimerization domain and an MT-binding domain. The MT-binding domain is localized at the second half of the MAP65-1 protein. The N-terminal region of this part of the MAP65-1 protein contains a conserved sequence responsible for MT binding, whereas its C-terminal region is more variable and predicted to be disordered

[74, 75]. It was recently shown that *Arabidopsis* MAP65-1 is phosphorylated by Aurora α -kinases at two amino acid residues within its C-terminal disordered tail. The phosphorylation of these residues renders its detachment from MTs, leading to cell cycle progression, suggesting that the unfolded structure in MAP65-1 is required to modulate the accessibility of the two phosphorylatable residues to Aurora kinases, hence ensuring appropriate cell proliferation during plant development [75].

NRPE1, the largest subunit of Pol V

The RNA-directed DNA methylation (RdDM) pathway may act to repress the transcription of transposable elements to maintain genome integrity, mostly during critical plant development stages [76]. In *A. thaliana*, the canonical RdDM pathway is characterized by the participation of heterochromatic 24 nt small RNAs (hc-siRNAs) which are mainly produced by the interplay between RNA POLYMERASE IV (POLIV) and RNA-DEPENDENT RNA POLYMERASE 2 (RDR2). These enzymes generate a double stranded RNA that is subsequently trimmed into a 24 nt duplex by a type III ribonuclease, DICER-LIKE 3 (DCL3) [77, 78]. The generated hc-siRNAs are then methylated by HEN1 at the 3' end of each strand [79] to be exported to the cytoplasm where one strand associates with the ARGONAUTE 4 (AGO4) complex [80]. The complex is then imported to the nucleus where hc-siRNA pairs may bind by base complementarity to a scaffold long non-coding RNA produced by RNA POLYMERASE V (POLV) [81]. The association of AGO4 in the silencing complex allows a physical interaction between this protein and POLV carboxy-terminal domain (CTD) via AGO hooks (described below) aided by the function of KTF1/SPT5L (Suppressor of Ty insertion 5—such as a homologue of SPT5 Pol II-associated elongation factor) [82]. This triggers the recruitment of a plethora of proteins which remove active chromatin marks and establish repressive ones, such as DNA methylation, DNA and histone modifications and chromatin remodeling features (reviewed extensively in [83]).

A peculiarity of the RdDM pathway in plants is the participation of two plant-exclusive RNA polymerases, POLIV and POLV. The catalytic domain of these polymerases is highly conserved, but their specific activities are conferred by their largest subunits; NRPD1 for POLIV, and NRPE1 for POLV [76, 84]. These subunits possess a characteristic carboxy-terminal domain which, in the case of NRPE1, contains a region rich in GW, WG and GWG amino acid residue arrangements, known as AGO hooks [84, 85]. This region constitutes an AGO-binding platform necessary for the interaction between NRPE1 and AGO4 and the consequent small RNA-directed DNA methylation

[86]. Besides NRPE1, AGO hooks are also present in other AGO-binding proteins with up to 45 repeats. Along with their repetitive character, AGO-binding platforms have been predicted to be IDRs [87]. Interestingly, whereas the AGO-binding platform of NRPE1 orthologues is highly divergent in the primary sequence, the intrinsic disorder and the presence of AGO hooks are hallmarks of AGO-binding platforms across NRPE1 s. These characteristics are also extended to other AGO-binding proteins such as SPT5L, suggesting that this repetitive disordered structure is required to interact with a broad repertoire of targets, presumably regardless of sequence conservation [84]. Moreover, the evolutionary analyses reported by Trujillo et al. [84] suggest that this repetitive disordered array has been conserved to allow rapid sequence divergence while maintaining key functions in these proteins.

Protein disorder in plant metabolism

Large-scale computational approaches have found that IDP functions seem to be more common in signaling and regulation processes, whereas structural order is more frequent in proteins involved in catalysis, in binding of small ligands and in membrane proteins (channels or transporters) [88]. However, this dichotomy contrasts with the description of some enzymes containing IDRs in loops or tails, which participate in the modification of protein conformation upon substrate binding, and thus expose catalytic residues and contribute to catalysis [89–91]. Furthermore, one must consider the role of some IDRs as sites for post-translational modifications, acting as switches of activation/inactivation or as modulators of their own activity. Many of these IDR-containing proteins are involved in the fundamental housekeeping of the plant.

In this section, we will describe those IDPs known to participate in different aspects of plant metabolism; some of them involved in photosynthesis, in metal binding or in antioxidant mechanisms.

Chloroplast protein 12 (CP12)

Few studies have investigated the role of protein structural disorder in the plant photosynthetic machinery. However, with the advancement of the characterization of proteins implicated in this process, more data are emerging showing the impact of intrinsic disorder in this essential plant function. An example of this is the chloroplast protein 12 (CP12), a well-characterized scaffold protein that forms a ternary complex with glyceraldehyde-3-phosphate dehydrogenase (GAPDH) and phosphoribulokinase (PRK), named the GAPDH-CP12-PRK complex. CP12, present in

most photosynthetic organisms, also regulates GAPDH and PRK activities [92, 93]. CP12 is a small protein (8.5 kDa) encoded in the nuclear genome and translocated to chloroplasts; although it contains cysteine residues, it has been shown to have all the properties of an IDP. Because its degree of disorder is higher in vascular plant orthologues than in eukaryotic algae, it has been proposed that CP12 has evolved to become more flexible, which correlates with its increased multifunctionality [94, 95]. In the plant kingdom, CP12 proteins share common features; however, their N termini, in addition to being highly disordered, show high sequence variability [95, 96]. During the formation of the GAPDH-CP12 or PRK-CP12 binary complexes, CP12 structural disorder remains, in particular in its N-terminal region, indicating that these are fuzzy complexes. These observations have suggested that the fuzziness of this association could facilitate the binding of either GAPDH or PRK [97, 98]. The integration of the different lines of evidence suggests a model for the formation of the GAPDH-CP12-PRK complex, where GAPDH associates with CP12 by conformational selection; first recognizing specific conformation(s) in CP12 to establish the binding. Upon this interaction event, the CP12N-terminal remains in a fuzzy state acting as a linker to facilitate the association with PRK. Once the GAPDH-CP12-PRK complex is formed, it dimerizes to form the native complex, composed finally of two dimers of PRKs, two tetramers of GAPDH, and probably, two monomers of CP12 [93, 96, 97] (Fig. 1b).

CP12 plays a key role in the regulation of the Calvin cycle, transducing changes in light availability such as those occurring during the day–night transition. This event leads to the generation of a hyperoxidant state, which is detected by the two cysteine residues in the CP12 C terminus forming a disulfide bridge. This leads to a conformational change in CP12, resulting in its N-terminal region folding into α -helix [96], which subsequently prevents the entrance of the NADPH cofactor in the GAPDH catalytic site. In the night-to-day transition, the conformation is reversed; the disulfide bridge is reduced by thioredoxin permitting NADPH entry and resulting in GAPDH activation. This inhibiting effect exerted by the CP12 also occurs on the PRK enzyme, as part of the complex. Interestingly, accumulating evidence indicates that CP12 assembles in larger supramolecular complexes, as happens in *Chlamydomonas reinhardtii*, where the GAPDH-CP12-PRK complex associates with aldolase [92], thus suggesting additional roles in other metabolic processes [23]. From the differential lines of evidence, it can be concluded that CP12, as with some other IDPs, has a moonlighting activity, being able to act as a scaffold for GAPDH and PRK [93], as a regulator of these enzyme

activities, and as a protective shield against oxidative damage [23, 99, 100].

Glyceraldehyde-3-phosphate dehydrogenase (GAPDH)

GAPDH plays a central role in glycolysis and gluconeogenesis. In vascular plants, GAPDH can exist as heterotetramers of two GapA and two GapB (A_2B_2) subunits, as homotetramers of four GapA subunits (A_4) or as hexadecamer of eight GapA and eight GapB subunits (A_8B_8). Interestingly, the GapB subunit also contains a C terminus highly similar to the CP12 C-terminal IDR [101]. The presence of two cysteine residues in this region permits photosynthetic NADPH-dependent GAPDH containing the GapB subunits to detect redox changes. Oxidative conditions induce the formation of a disulfide bridge in its CP12-like C terminus, promoting the NAD-dependent arrangement of higher homo-oligomers that result in auto-inhibition of its NADPH-dependent catalytic activity. This conformational change and complex formation is needed for the reduction of 1,3-bisphosphoglycerate to produce glyceraldehyde-3-phosphate [101–103]. This intrinsically disordered feature of GapB confers on A_2B_2 GAPDH a CP12-autonomous regulation by the redox status of the cell.

Ribulose-1,5-bisphosphate carboxylase/oxygenase (Rubisco) activase

Rubisco (Ribulose-1,5-bisphosphate carboxylase/oxygenase), the most abundant protein on Earth [104, 105], is an enzyme responsible for fixing atmospheric CO_2 into RuBP (ribulose 1,5-bisphosphate) to produce two phosphoglycerate molecules. The activity of this enzyme depends on the binding of Mg^{2+} ions and the carbamylation of a lysine residue located in its active site; however, the binding of RuBP can reduce the efficiency of carbamylation, and consequently the activation of the enzyme [106]. Nature has solved this limitation through proteins known as Rubisco activases that because of their ATPase and chaperone activity, allow Rubisco carbamylation by removing RuBP from the active site and giving access to CO_2 molecules. Photosynthetic organisms present two Rubisco activase isoforms (α and β) [107] containing a C-terminal extension (20–50 amino acid residues) which is predicted as an intrinsically disordered region [23]. As is the case for CP12, this IDR contains two highly conserved cysteine residues in α isoforms [108, 109], responsible for the light regulation of Rubisco activase. This control is achieved by the action of thioredoxin *f* on the two cysteine residues, such that upon oxidation, inhibition of α isoform activity

by light is abolished [110, 111]. The Rubisco activase function can be recovered by the reduction of the C-terminal disulfide bridge by thioredoxin *f*, depending on the redox status of the chloroplasts [112]. Interestingly, in spite of the functional or structural differences among Rubisco activases in diverse photosynthetic organisms, their C-terminal IDRs have been conserved; for example, in the case of cyanobacteria, they are involved in carboxysome targeting [23, 113]. Overall, intrinsic disorder in Rubisco activase strongly suggests that it is a conserved feature responsible for its functional versatility as an ATPase, a chaperone and as a fine-tuning regulator that has contributed to the broad adaptability of the photosynthetic process.

Manganese stabilizing protein (MSP)

Plants capture sunlight through the light-harvesting complex (LHC) or antenna complex as part of Photosystem II (PSII). This complex of proteins and pigments is embedded in thylakoid membranes and connects the antenna to the chlorophylls in the reaction center. The photons captured by PSII initiate a chain of redox states through electron transfer reactions needed for the oxidation of two water molecules to O₂. This photolysis reaction takes place in the oxygen-evolving complex (OEC), one of the PSII subunits. The different polypeptides of PSII are needed for an efficient O₂ evolution; in particular, three extrinsic proteins of 17, 23, and 33 kDa, which are located on the luminal side of PSII. This last protein, also termed manganese stabilizing protein (MSP), is required to maintain stability and an efficient cycling of the four oxidizing manganese atoms [114–117]. MSP lacks a compact structure and is composed of 55% turns and random coils. These properties, together with its amino acid composition and other features, establish its intrinsic structural disorder. In vitro experiments suggest that the structural flexibility of this protein is required for its function, possibly by facilitating effective protein–protein interactions as an integral member of PSII [118]. Moreover, it has been shown that conserved charged amino acid residues in MSP are important for the retention of Cl[−] ions, to maintain their concentration at the levels needed for the effective redox reactions of the manganese cluster [119]. Again, MSP exemplifies the participation of protein structural disorder as an essential attribute to achieve precise and opportune roles in a complex system able to adjust to the changing environment.

Alb3, a thylakoid membrane protein

The membrane invertase protein Alb3 controls the insertion, folding and assembly of a diverse group of proteins

into the thylakoid membrane of chloroplasts. Alb3 interacts with chloroplast signal recognition particles (cpSRP) in the thylakoid membrane through its C-terminal intrinsically disordered region. This IDR has two conserved positively charged motifs needed for the association with cpSRPs that follows a coupled binding and folding mechanism [120]. Once the Alb3-cpSRP complex is formed, it participates in the post-translational insertion of the light-harvesting chlorophyll a/b-binding protein (LHCP), a highly abundant protein in thylakoid membranes. The insertion of LHCP into these membranes strictly requires the involvement of cpSRP and Alb3. Alb3 is also needed for the targeting and insertion of cytochrome *b₆* into the thylakoid membrane [121]. Cytochrome *b₆* is a largely disordered protein in aqueous solution, but by interaction with lipids from the membrane it folds into an α -helical structure just before its membrane insertion [122]. An additional function assigned to the Alb3 C-terminal IDR is the light-dependent modulation of Alb3 stability [123].

Polyphenol oxidases (PPOs)

Tyrosinases and catecholases from plants and fungi are generally named polyphenol oxidases (PPOs). In plants, PPOs mediate the production of melanin, responsible for the brown color in fruits when they suffer damage. They are nuclear-encoded and are transported to the chloroplast thylakoid lumen, where they can be in a soluble form or in a weak association to the thylakoid membranes. They are activated by the proteolytic cleavage of their C-terminal region. Using bioinformatic approaches to analyze multiple plant PPO sequences, it was found that the region between the N-terminal and C-terminal corresponds to a disordered linker essential to establish those conditions in which the PPOs are processed and may be activated [124]. This prediction suggests that the PPO IDR may acquire certain levels of order depending on the environment. Although experimental data are needed, the presence of a conserved phosphorylation site within this IDR suggests auto-regulation of PPO activities and/or that PPOs have roles as signaling molecules [124].

Phosphoenolpyruvate carboxylase (PEPC)

Carbon assimilation is not only accomplished by the activity of Rubisco, but also by PEPC (Phosphoenolpyruvate carboxylase), a ubiquitous enzyme in plants. PEPC also plays a critical role in plants with C₄ photosynthesis and crassulacean acid metabolism (CAM), by producing oxaloacetate from HCO₃[−] [125]. Two types of PEPC enzymes have been described in plants, known as plant-type PEPC (PTPC) and a distantly related bacterial-type PEPC (BTPC). The BTPC enzymes show low sequence identity with PTPCs, they lack the typical serine

phosphorylation motif located in the PTPC N-terminal region and they are encoded in all plant genomes sequenced to date. Of particular interest is the fact that BTPCs contain an insertion of approximately 142 amino acid residues predicted as a structurally disordered region. This IDR seems to be highly divergent and a distinctive characteristic of BTPCs. PEPC enzymes are organized as oligomers, from which two classes have been identified: class-1 oligomers consisting of PTPC homotetramers, and class-2 complexes corresponding to heterotetramers composed of three PTPC and one BTPC subunits [125]. Recently, it was demonstrated that the BTPC IDR from the castor oil plant (*Ricinus communis*) is needed for its association with the PTPC subunit in a class-2 PEPC complex. Furthermore, even though the N-terminal region conserved in PTPCs is not conserved in BTPCs, it was thought that these enzymes were non-phosphorylatable. However, it has been shown that RcBTPC is phosphorylated in vivo at least at two serine residues. One of these modifications occurs at serine-451, a highly conserved target residue located within the IDR of these proteins. This event exerts a regulatory role, causing the inhibition of the catalytic activity of the enzyme within the class-2 PEPC complex [126].

UreG G protein

GTP-binding proteins (G proteins) are GTPases that catalyze the hydrolysis of GTP to yield GDP and inorganic phosphate. The structure of the catalytic domain of this enzyme is usually a β -sheet delimited by flexible regions of α -helices and loops. The binding of GTP or GDP activates (in the case of GTP) or inactivates (in the case of GDP) these GTPases, associations stabilized by the binding of specific protein regulators that promote conformational modifications. UreG is a bacterial-type G protein involved in urease maturation that has been demonstrated to belong to the class of intrinsically disordered enzymes. The structural disorder in UreG is mostly concentrated in a region of ~ 50 residues localized in the center of its protein sequence, which seems to influence the structure of the GTP-binding pocket [127–129]. In plants, one gene shows sequence similarity with bacterial *UreG* GTPase, which functions as an urease accessory protein, promoting optimal urease activation by allowing nickel or zinc incorporation in its active site and the GTP-dependent CO_2 transfer required for lysine carbamylation. This protein has been characterized in soybean (*G. max*), where it has shown a differential binding affinity to Ni^{2+} and Zn^{2+} . Furthermore, it has the highest affinity for Zn^{2+} described to date for any UreG protein. This observation suggests a role for UreG as a Zn^{2+} accumulator protein that may modulate the available levels of this metal in the cell. Analysis of its

quaternary structure indicates that UreG is monomeric in solution and that dimers can be formed and stabilized upon Zn^{2+} binding, due to conformational rearrangements in the protein. The association with Zn^{2+} decreases the levels of secondary structure, but perhaps stabilizes the subsequent dimerization by facilitating the folding of the active site domain. However, this binding alone is not enough to yield a high UreG activity, suggesting that additional factors are needed to achieve its optimal GTPase activity [130]. UreG further illustrates the functional versatility conferred by intrinsic disorder to proteins with catalytic and regulatory roles in plant metabolism.

Jaburetox, an intrinsically disordered insecticidal polypeptide

Ureases are nickel-dependent metallo-enzymes that catalyze the hydrolysis of urea into ammonia and CO_2 [131]. It was discovered that canatoxin, considered an isoform of a jack bean urease (from seeds of *Canavalia ensiformis*), corresponds to a 10-kDa peptide (JBU) produced from urease hydrolysis by cathepsin-like enzymes. This JBU peptide is toxic to mammals, fungi and insects. One of the major urease isoforms from jack bean seeds shows toxicity to hemiptera insects independent of its ureolytic activity, and instead its effect is produced by the action of digestive enzymes present in the insect gut [132, 133]. This entomotoxic activity is caused by an internal peptide (pepcanatox) product of this hydrolysis. Jaburetox, a recombinant version of the in vivo generated peptide, is derived from the N-terminal sequence of the *C. ensiformis* urease isoform and possesses a potent insecticidal effect on crop pests [133, 134]. A motif present on this peptide is also found in pore-forming and neurotoxic peptides which present membrane-disturbing activities [135]. A large hydrodynamic radius, together with light scattering, CD and NMR spectroscopic data, shows that Jaburetox is a monomeric disordered peptide with an α -helix motif by its N terminus and two turn-like structures in the central region and by the C terminus of the peptide. It is suggested that the Jaburetox IDP might act as a membrane protein or as a scaffold protein, but evidence for this is still lacking, therefore a comprehensive view of its insecticidal activity remains elusive [136].

Protein structural disorder in plant abiotic stress responses

Prediction of structural intrinsic disorder from plant proteomes reveals a noteworthy participation of IDPs in plant responses to their environment and to stress conditions.

Table 2 Intrinsically disordered LEA proteins

Group	Protein	Specie	Function*	Reference
LEA 1	AtEM6	<i>Arabidopsis thaliana</i>	Protein protection activity	Campos et al. [155]
LEA 2	CrCOR15	<i>Citrus clementina x Citrus reticulata</i>	Protein protection activity	Sanchez-Ballesta et al. [151]
	DHN-5	<i>Triticum durum</i>	Protein protection activity	Drira & Antonny [258] Drira et al. [154]
	YSK2	<i>Vitis riparia</i>	Protein protection activity	Hughes et al. [160]
	PCA60	<i>Prunus persica</i>		
	DHN5	<i>Hordeum vulgare</i>		
	TsDHN-1	<i>Thellungiella salsuginea</i>	Vesicle stabilization	Rahman et al. [177]
	TsDHN-2	<i>Thellungiella salsuginea</i>	Protection activityVesicle stabilization	Hughes et al. [160] Rahman et al. [177]
	OpsDHN-1	<i>Opuntia streptacantha</i>	Protein protection activity	Hughes et al. [160]
	COR85	<i>Spinacia oleracea</i>	Protein protection activity	Kazuoka & Oeda [162]
	WCS120	<i>Triticum aestivum</i>	Protein protection activity	Houde et al. [163]
	HvP-80/ Dhn5	<i>Hordeum vulgare</i>	Protein protection activity	Bravo et al. [164]
	Cor15am	<i>Arabidopsis thaliana</i>	Protein protection activity	Nakayama et al. [165]
	ERD10	<i>Arabidopsis thaliana</i>	Protein protection activity, Calcium-binding activity	Kovacs et al. [166] Reyes et al. [158] Alsheikh et al. [207]
	ERD14	<i>Arabidopsis thaliana</i>	Protection activity, Calcium-binding activity	Kovacs et al. [166] Tantos et al. [152] Alsheikh et al. [206]
	35KD	<i>Vigna unguiculata</i>	Lipid vesicles binding	Ismail et al. [172]
	DHN1	<i>Zea mays</i>	Stabilization of vesicles	Koag et al. [175]
	rGmDHN1	<i>Glycine max</i>	Lipid membrane binding	Soulaiges et al. [176]
	Lti30	<i>Arabidopsis thaliana</i>	Stabilization of vesicles, Antioxidant activity	Ericksson et al. [178] Hara et al. [204]
	K2	<i>Vitis riparia</i>	Stabilization of vesicles	Clarke et al. [179]
	VcaB45	<i>Apium graveolens</i>	Calcium-binding activity	Heyen et al. [202]
	BjDHN2	<i>Brassica juncea</i>	Antioxidant activity (<i>in vivo</i>)	Xu et al. [208]
	BjDHN3			
	ITP	<i>Ricinus communis</i>	Fe transport in the phloem	Krüger et al. [205]
	CuCOR15	<i>Citrus unshiu</i>	RNA and DNA binding	Hara et al. [209]
	WCI16	<i>Triticum aestivum</i>	DNA binding	Sasaki et al. [210]
	DHN1a	<i>Vitis vinifera</i>	Protection activity, DNA binding	Rosales et al. [211]
	COR47	<i>Arabidopsis thaliana</i>	Calcium-binding activity	Alsheikh et al. [207]
	HIRD11	<i>Arabidopsis thaliana</i>	Antioxidant activity, Cu binding	Hara et al. [204]
LEA 3	LEA76	<i>Arabidopsis thaliana</i>	Protection activity	Reyes et al. [149]
	AavLEA3	<i>Aphelenchus avenae</i>	Protection activity	Goyal et al. [150]
	PsLEAm	<i>Pisum sativum</i>	Protection activity	Grelet et al. [156]
	Cor15am	<i>Arabidopsis thaliana</i>	Protection activity	Nakayama et al. [165]
	Cor15a	<i>Arabidopsis thaliana</i>	Protection activity, Vesicle stabilization	Thalhammer et al. [169]
	Cor15b			

Table 2 continued

Group	Protein	Specie	Function*	Reference
LEA 4	AtLEA4-2	<i>Arabidopsis thaliana</i>	Chaperone-like	Cuevas-Velazquez et al. [139]
	AtLEA4-5			
	GmPM1	<i>Glycine max</i>	Fe ³⁺ binding, Antioxidant activity	Liu et al. [200]
	GmPM9			
LEA 7	GmASR1	<i>Glycine max</i>	Zn ²⁺ , Fe ³⁺ binding	Liu et al. [194]
	VvMSA	<i>Vitis vinifera</i>	Protection activity, Transcription factor	Ricardi et al. [212]
	ZmASR1	<i>Zea mays</i>	Transcription factor	Virilouvet et al. [214]
	ASR1	<i>Solanum tuberosum</i>	Glucose content modulator	Frankel et al. [215]

* Function is inferred from phenotypes as described in the text

However, not many abiotic stress response proteins have been confirmed as IDPs and there is limited information about their function. Here, we compile those stress-responsive IDPs for which there is evidence of function and structural organization.

Late embryogenesis abundant (LEA) proteins

Late embryogenesis abundant (LEA) proteins belong to an emblematic group of IDPs distinctively involved in plant stress responses, notably, in adverse conditions of low water availability. LEA proteins can be classified into seven groups or families based on amino acid sequence similarity, although nomenclature can vary. In this review, we will follow that proposed by Battaglia et al. [137], who also report the presence of distinctive motifs for each family, some of which correspond to MoRFs [138, 139]. LEA proteins do not show significant sequence similarity with any other proteins of known function, making their characterization a challenging task. LEA proteins are considered ubiquitous in the *Viridiplantae* kingdom because they have been found in angiosperms, gymnosperms, non-vascular plants and algae [137, 140]. Although for some time they were considered exclusive to plants, interestingly they have also been detected in other organisms including insects, nematodes, crustaceans, rotifers and bacteria [141–145]. In all cases, their abundance is related to water deficit, but some also respond to other stress conditions. In general, LEA proteins are highly hydrophilic with a high content of glycine residues or other small amino acids, and they are usually deficient in tryptophan and cysteine residues; all characteristics of IDPs [137, 146, 147]. These properties are conserved in a wider group of water deficit response proteins, the ‘hydrophilins’, which are conserved across all domains of life [148]. As is documented for other IDPs, LEA proteins possess key qualities that enable them to perform more than one function; this ‘moonlighting’ characteristic will be

described below. As in the case of IDPs involved in development and metabolism, the plasticity and molecular flexibility of LEA proteins appear to be central to their function (Fig. 2A).

One of the most general functions across the LEA group is an ability to protect the integrity of other enzymes. This has been demonstrated using several non-plant reporter enzymes with in vitro partial dehydration and freeze–thaw treatments, whereby the presence of LEAs prevents inactivation, denaturation and consequent aggregation of enzymes such as lactate (LDH) and malate dehydrogenases (MDH), citrate synthase (CS), β -glucosidase G (β glG), and glucose oxidase/peroxidase (GOD/POD) [137, 146, 149–155]. In the case of group 3 LEA proteins from *Pisum sativum* (PsLEAm), this protective effect has been demonstrated on plant proteins such as mitochondrial rhodanase and fumarase [156]. The protective activities resemble that of small heat shock proteins (sHSPs), which circumvent protein aggregation upon heat shock treatments in the absence of ATP [157]. Hence, it appears that LEA proteins may function as chaperones during water deficit stress. These observations suggest a protective role specifically against protein damage caused when water limitation inhibits cellular functions. Furthermore, it appears that this unique function cannot be provided by other types of chaperones [149, 158] (Cuevas-Velazquez et al. unpublished).

The different lines of evidence from in vitro enzyme assays have led to two main hypotheses to explain the LEA protein protecting activity. Because high concentrations of LEA proteins are able to prevent inactivation and aggregation of other proteins, it has been proposed that they may act as ‘molecular shields’. Given their large hydrodynamic radius in aqueous solution, they may create a protein molecular net, thereby promoting the alignment of their hydrophilic amino acid residues around the surface of a target protein, and in this way, prevent the loss of its bulk water and consequent changes in its native structure [159, 160]. However, there is also evidence




Fig. 2 Schematic representation of two examples of plant IDPs that participate in abiotic and biotic stress responses. **A** LEA proteins (represented as *purple curved lines*) belong to a representative group of plant IDPs involved in plant abiotic stress responses. LEA proteins are able to prevent the inactivation of reporter enzymes under in vitro partial dehydration and freeze–thaw treatments. One action mechanism supported by different lines of evidence indicates that LEA proteins function as chaperones during water deficit *a* by interacting with their protein target(s) (*green irregular ovals*) and avoiding the damage (denaturation represented by *green irregular lines* emerging from the *green ovals*) caused by the effects of low water availability in the cell. The possibility that LEA proteins may bind and recognize their targets by conformational selection under water deficit has been suggested by in vitro data. In addition, there is evidence indicating that LEA proteins are able to stabilize membrane (*double blue circles*) integrity *b* during water deficit, by interaction through the amphipathic regions present in some LEA proteins. It has been suggested that LEA proteins might achieve more stable conformations upon membrane association. It has been proposed that this interaction induces LEA protein folding. *b* An additional attribute of at least some LEA proteins is their ability to bind metal ions (Fe^{3+} , Ni^{2+} , Cu^{2+} , Co^{2+} and Zn^{2+}) (*small gray fill circles*), which in some cases,

showing that small amounts of LEA proteins (down to 1:1–1:5 ratios of LEA:reporter enzyme) are also capable of protecting target proteins to a similar degree [149, 161–166]. This indicates that LEA proteins may function in a ‘chaperone-like’ mode, where interaction is required to select and protect their targets, binding as monomers or oligomers [149, 161, 167, 168] (Fig. 2A(a)). This hypothesis is supported by evidence indicating that these disordered proteins can fold in α -helix under high osmolarity or high macromolecular crowding, prevalent

by these means scavenge reactive oxygen species (*c*). For some LEA proteins, metal binding promotes a reduction in the content of structural disorder; however, this is not a common observation. In this panel, continue arrows represent the protective effect of LEA proteins, whereas discontinuous arrows indicate the consequent damage produced by stress in the absence of these proteins. **B** Biotic stress produced by plant pathogens has led to the selection of refined mechanisms to detect their presence and to mount complex inducible responses to efficiently counteract their attack. The participation of IDPs along the different steps of pathogen invasion, from their perception to the plant defense response has been documented. The RbohD protein (*green curved lines*), which belongs to the NADPH oxidase family, represents an example of this. This protein, partially integrated in the membrane, is responsible for the early generation of ROS, upstream of calcium and phosphorylation signaling. The RbohD cytoplasmic N terminus possesses an IDR, which contains EF-hand motifs involved in calcium binding. The malleable nature of this region results in extended conformational changes induced by the synergistic effect of calcium binding and its phosphorylation, which in turn modulates the interaction with small GTPase proteins (*orange irregular oval*); a process needed to set up the plant protection response against pathogens

conditions under water deficit, which would lead to a decrease in their hydrodynamic radius [138, 139, 147, 169]. Crucially, this property seems to be associated with their chaperone-like activity [139]. With this in mind, and considering the role of conformational plasticity [138, 139, 147, 169–171], it is possible that LEA proteins may bind and recognize their targets following a mechanism that resembles conformational selection under water deficit, the natural conditions under which they accumulate in the cell. Although similar lines

of evidence have been obtained for LEA proteins from different groups (LEA2, LEA3, LEA4, LEA7), further experimentation is needed to support these alternative mechanisms. In particular, it is imperative to establish strategies to obtain *in vivo* data that could help provide a more comprehensive view of their action mechanisms.

The intrinsic disorder of LEA proteins may also function in the stabilization of membrane integrity under stress (Fig. 2A(b)). Some LEA proteins, mainly those from groups 2–4, are able to bind *in vitro* to lipid vesicles [172–176]. In some cases, these vesicles have been produced using phospholipids and galactolipids common to plant chloroplast and mitochondrial envelopes [169, 177], or they have been obtained from thylakoid membrane fractions of spinach leaf tissue [178]. Interestingly, α -helix folding upon vesicle binding has also been shown for some LEA proteins [179, 180]. For group 2 LEA proteins (dehydrins), it has been found that the K-segment, a distinctive motif of this family is necessary for liposome binding, which is consistent with its amphipathic nature [178, 180, 181] (Fig. 2A(b)). Distinctive motifs of LEA3 and LEA4 proteins also present amphipathic properties, which help explain their ability to bind to lipid vesicles surfaces [137, 182, 183].

Unfortunately, to date there remains no evidence of any of these activities *in vivo*. Despite this, *in vitro* functions correlate with the accumulation of LEA proteins in plant tissues under low water potentials induced by dehydration or by cold or freezing temperatures; conditions in which enzymes can be inactivated and membrane injuries are produced. Interestingly, some *Arabidopsis* LEA proteins are required for plant optimal adjustment to cold, water deficit and/or salinity, as can be inferred from the phenotypes produced by mutants lacking genes encoding LEA proteins from group 1 [184], group 2 [185], group 3 [169] and group 4 [186]. Additionally, the acquisition of tolerance to water limitation or low temperatures by the overexpression of several LEA proteins (from groups 2–4 and 7) in different plant species strongly supports their role as protector molecules under these stress conditions [187–197].

As can be seen with IDPs involved in plant development and metabolism, dynamic structural order can be attained by interaction with metal ions. LEA proteins have also been shown to bind metal ions (Fe^{3+} , Ni^{2+} , Cu^{2+} , Co^{2+} and Zn^{2+}) and scavenge reactive oxygen species (reviewed in [198, 199]). LEA proteins showing high affinity to these metals include LEA2 or dehydrins, LEA3 and LEA4 [191, 198–200]. Acid dehydrins (RAB17 and VCaB45) are able to bind calcium, possibly to modulate intracellular calcium levels, thereby acting as ionic buffers during water deficit: a hypothesis that still needs to be tested [201, 202]. The metal binding properties in these proteins have been attributed to the abundance of histidine residues or to the

presence of metal binding motifs (HX₃H or HH) [203]. Importantly, for some LEA proteins, it is known that metal binding can promote the gain of an ordered conformation [204].

Remarkably, a group 2 LEA protein (ITP, iron transport protein) has been shown to carry iron *in vivo* and bind iron *in vitro* (Fig. 2A(c)). This protein was found associated with iron in phloem exudate from *R. communis* L. [205]. ITP also binds Ni^{2+} , Cu^{2+} , Zn^{2+} and Mn^{2+} *in vitro*, preferentially binding to Fe^{3+} but not to Fe^{2+} . This indicates that this LEA protein may function as a phloem micronutrient transport protein [205], opening up the possibility that this novel function may exist for other LEA proteins or IDPs able to bind iron or other micronutrients.

For some group 2 proteins, it has been shown that their phosphorylation is required for the metal association to occur [202, 206, 207]. Group 2 and group 4 proteins can also circumvent the production of reactive oxygen species (ROS), given their capacity to bind metals able to promote ROS generation (Fig. 2A(c)). Evidence for this activity has been obtained *in vitro* and *in vivo* [200, 204, 208]. This mechanism could be advantageous under abiotic stress conditions such as water deficit, when ROS production and sensitivity to secondary stresses are exacerbated.

The multifunctionality of LEA proteins and the role of metal ions are reinforced by data indicating that group 2 and group 7 LEA proteins can also bind nucleic acids. Group 2 LEA proteins (CuCOR15, VvDHN1a and WC116) have been shown to associate with DNA and RNA. In the case of CuCOR15 and WC116, this occurs in the presence of physiological concentrations of Zn^{2+} [209–211]. This evidence suggests that nucleic acids need similar protection from the effects of water limitation.

DNA binding has also been demonstrated for a group 7 LEA protein (ASR1, ABA [Abscisic acid] stress ripening 1), a widely occurring plant LEA protein that does not exist in *A. thaliana*. Strikingly, in addition to its protein protective role, ASR1 can also function as a transcription factor. It has been shown that ASR1 is able to bind to the regulatory regions of genes related to cell wall synthesis and remodeling, as well as genes encoding membrane channels implicated in water and solute trafficking [212]. Grape ASR1, VvMSA, recognizes specific sites in the regulatory region of the *hexose transport 1 (Ht1)* gene [213], and ASR orthologues are also involved in sugar and amino acid accumulation in species such as maize and potato [214, 215].

Phosphorylation of IDRs in some LEA groups may also play a role in LEA protein function. Members of group 2, 4, 6 contain phosphorylatable motifs and *in vivo* and *in vitro* phosphorylation has been verified for group 2 LEA proteins (dehydrins/DHNs) in *Arabidopsis*, wheat, maize, and other plants [201, 207, 216–222]. Although the role of

this modification is not well understood, for group 2 LEA proteins it may be needed to modulate membrane interaction and lipid phase transition [178, 180], as well as nuclear localization [223, 224]. However, although phosphorylation pattern correlates with tolerance to water limitation, it is unknown whether this modification is required to modulate LEA protein protective activity and/or target selectivity by allowing the display of different binding motifs and/or MoRFs.

The multifunctionality found *in vitro* for the different LEA proteins is often compatible with their *in vivo* intracellular localization, suggesting that there may be both subcellular specialization as well as redundancy. LEA proteins from all groups have been localized to cytosol, nucleus, mitochondria and chloroplast [137, 199, 225]. However, at least in the case of group 3 LEA proteins, the most diverse LEA family, not all its members show the same localization. Some are found in cytosol or nucleus, others in the chloroplast and some others only in mitochondria [226]. This implies a requirement and possible functional specificity of LEA proteins during the plant stress response.

Further evidence of their deep functional divergence, as well as their ubiquitousness, can be seen in the high conservation of most LEA families throughout the *Plantae* kingdom's evolution. LEA proteins from group 1–4 can be detected in genomes from the most recent angiosperms through to the bryophytes, including the liverwort *Marchantia*, the most basal plant model described to date. Group 6 and 7 LEA proteins have been found only in seed plants, and in the case of group 7 LEA proteins, do not seem to be present in all phyla [137, 155, 186, 227, 228]. The broad distribution and conservation of these plant IDPs throughout evolution illustrate not only the relevance of these proteins for the organisms in this kingdom, but also the importance of disorder for the various functions they achieve.

The ubiquity of LEA proteins across land plants is testament to their versatility. Even though LEA protein action mechanisms remain elusive, the intrinsically disordered nature of these proteins matches their apparent 'moonlighting' character, as exhibited by diverse data where the same LEA protein is able to protect proteins and membranes, and bind metals and/or nucleic acids (see Table 2). These characteristics are compatible with their ability to utilize the same or overlapping regions to exert distinct effects and to switch functions by adopting different conformations upon binding [229].

Small heat shock proteins (sHSPs)

Small heat shock proteins (sHSPs) are ubiquitous molecular chaperones, which play important roles in protein

homeostasis and in plant responses to stress. sHSPs are classified in 11 subfamilies, six localized to cytoplasm/nucleus and five to organelles. These chaperones bind diverse partially unfolded polypeptides maintaining their refolding capacity until they can return to their native structure with the help of other chaperone proteins, such as HSP70. In this way, sHSPs protect cells from the loss of essential proteins and from the penalties caused by protein aggregation. Commonly these proteins respond to high temperatures, but also to other stress conditions, and some may also be produced even under optimal growth conditions. In contrast to other molecular chaperones, sHSPs form large and dynamic oligomers with different stoichiometry. All sHSPs contain a core α -crystallin domain bordered by a short C-terminal region and an N-terminal extension of variable length and sequence (for review see [157, 230]). Both regions participate in the recognition of—and binding to—clients and in the formation of their oligomers (for review see [230, 231]). It has been proposed that during heat stress the oligomeric sHSPs undergo conformational rearrangements leading to their dissociation. These structural changes enable the interaction of these chaperones with hydrophobic patches in the partially denatured clients, subsequently forming large soluble complexes, protecting protein clients from further damage. Biochemical and biophysical evidence indicate that the intrinsically disordered N-terminal arm is able to present different interaction sites revealing a mechanism to efficiently protect the integrity of many different substrates in the cell [157, 232]. Although many questions still remain unanswered regarding mechanistic details and *in vivo* evidence is required, sHSPs offer a view of the need for structural plasticity and promiscuity to maintain cell functions during stress.

Glycine-rich RNA-binding proteins (GR-RBPs)

Although LEA proteins are important to the plant cold stress response, other IDPs are known to play a protective role. Plants exposed to low temperatures experience a slowing down or even a pause of their metabolic processes and this may result, directly or indirectly, in damage to macromolecules and cellular structures [233]. Among the proteins synthesized to overcome the impairment that cold and other abiotic and biotic stresses cause to macromolecules are the so-called glycine-rich RNA-binding proteins (GR-RBPs) [234, 235]. Some of the functions characterized for GR-RBPs are the facilitation of mRNA transport and participation in splicing and translation: roles mediated by their RNA chaperone activity [236, 237]. GR-RBPs contain an RNA recognition motif (RRM) in the N-terminal region and a disordered glycine-rich region (GR) at their C-terminal end, and they can be classified into

eight groups, each one with apparently different roles [238, 239]. In *Arabidopsis*, AtGR-RBP7, in addition to being a circadian regulator and promoter of flowering and mRNA splicing, accumulates in response to cold stress [236, 237, 240–244]. Deletion of *AtGR-RBP7* leads to low-temperature sensitive phenotypes, highlighting its role in optimal plant adjustment to cold stress [245]. NMR analysis confirms the structural disorder of the GR domain for NtGR-RBP1, an AtGR-RBP7 orthologue from *Nicotiana tabacum* [246]. As expected, NtGR-RBP1 is shown to bind RNA and single stranded DNA through the RRM. Furthermore, the NtGR-RBP1 GR interacts transiently with its RRM domain, promoting self-association to effectively increase its local concentration, and hence its affinity for nucleic acids. These findings suggest a mechanism for the unfolding of non-native structures in RNA by NtGR-RBP1, which may be involved in enhancing its RNA chaperone activity [246].

Vesicle-inducing protein in plastid 1 (VIPP1)

The integrity of thylakoid membranes is crucially important for photosynthesis and chloroplast functions. Multiple reports have shown the participation of a protein called VIPP1 (Vesicle inducing protein in plastid 1) in thylakoid membrane biogenesis and thylakoid membrane maintenance during drought, heat and osmotic stress [247, 248], not only in cyanobacteria and green algae, but also in vascular plants [249–251]. The evolutionary emergence of this protein seems to be specific to oxygenic photosynthetic organisms [251]. Recent evidence suggests that while VIPP1 may have multiple roles in plastids, it strongly protects the chloroplast envelope [252]. The N-terminal region of VIPP1 presents high sequence similarity to its bacterial orthologue PspA (Phage shock protein A) [253], which plays a central role in the well-characterized bacterial system Psp, involved in the protection of membrane integrity under various stresses [254]. During membrane damage, PspA and VIPP1 bind to membranes forming high-order oligomeric effector complexes able to repair the inner membrane and conserve its integrity [255, 256]. Interestingly, this occurs despite the absence of trans-membrane domains in these proteins [252, 253]. CD spectroscopy studies show that PspA and VIPP1 N-terminal peptides are disordered in solution and fold upon membrane association, as occurs in a typical membrane amphipathic helix [257, 258]. The membrane binding of these proteins depends on differences in stored curvature elastic stress, a feature of damaged membranes [259]. These observations suggest that the folding transition associated with PspA and VIPP N-terminal membrane binding might act as a stress-sensing mechanism controlling the effector function of these proteins.

During the evolution of photosynthetic organisms, the PspA orthologue VIPP1 has acquired an additional C-terminal tail (Vc) that also presents the characteristics of an intrinsically disordered region [260]. Using live imaging experiments performed in vivo in *Arabidopsis*, with GFP (green fluorescent protein) translational fusions of VIPP1 or VIPP1 lacking Vc (VIPP1 Δ Vc), it was shown that Vc enables VIPP1 to form oligomeric effector complexes along cell envelopes, whereas VIPP1 Δ Vc leads to the formation of irregular aggregates of VIPP1 particles. The expression of VIPP1 Δ Vc complemented the *vipp1* knock out mutation in *Arabidopsis*, but exhibited sensitivity to heat shock. Furthermore, transgenic plants over-expressing wild-type *VIPP1* showed enhanced tolerance against heat shock. *Vipp1* knockout *Arabidopsis* mutants show reduced content and other structural defects of thylakoid membranes, as well as reduced photosynthetic activity. In addition to its role in membrane biogenesis, it has been proposed that VIPP1 may also function as a lipid transfer protein, delivering structural lipids into thylakoid or envelope membranes [253]. Overall, these data suggest that the involvement of the Vc disordered region in the formation of the oligomeric effector complexes might be relevant for the control of VIPP1 association/dissociation states. Under conditions of membrane stress, this IDR may permit the insertion of their amphipathic helix into the lipid bilayer to relax the curvature elastic stress in membranes [259].

Dehydration-responsive element binding protein 2A (DREB2A)

Dehydration-responsive element binding protein 2A (DREB2A) is a key transcription factor for drought and heat stress tolerance in *Arabidopsis*. DREB2A induces the expression of dehydration and heat stress responsive genes [261]. This transcription factor contains several IDRs allowing it to interact with multiple proteins, a characteristic consistent with interactome data showing that DREB2A is a hub protein with 26 nodes [21]. DREB2A may interact with its negative regulators DRIP1 and DRIP2 (DREB2A-interacting protein1 and 2), with ribosomal proteins such as RPL15 (ribosomal protein L15), other transcription factors such as RCD1 (Radical cell death 1), and the transcription co-regulator MED25 (Mediator 25), among others [21]. It has been shown that MED25 binds to one of the DREB2A IDRs and that this interaction results in a gain of ordered structure in this region. Similarly, the binding of DREB2A to its canonical DNA sequence also leads to an increase in the secondary structure of the protein. Data also show that DREB2A conformational changes induced by DNA binding reduce its interaction with the

MED25 acid domain, which does not exclude the possibility that this modification may promote its association to another Mediator subunit close by [262]. RCD1 controls DREB2A function, and is itself rapidly removed during abiotic stress [263]. O'Shea et al. [264] showed by NMR spectroscopy that DREB2A undergoes coupled folding and binding with α -helix formation upon interaction with RCD1.

bZIP28, a transcription factor in the unfolded protein response

Under adverse conditions such as heat stress, pathogenesis and by inhibition of protein glycosylation [265–267], the demand for protein folding can exceed the capacity of protein homeostasis systems. This results in the increase of misfolded or unfolded proteins in the endoplasmic reticulum (ER) lumen. This series of events leads to ER stress that subsequently induces the unfolded protein response (UPR) to fulfill the requirement of protein folding and degradation [268]. Two branches of the UPR signaling pathway have been described in plants: one involving the membrane-associated basic leucine zipper (bZIP) transcription factors and the other involving a bifunctional protein, with kinase and ribonuclease activities, known as inositol-requiring enzyme 1 (IRE1), which functions as an RNA splicing factor [269]. In *Arabidopsis*, bZIP28 is an ER membrane-associated transcription factor; its N-terminal region contains a transcriptional activation domain oriented towards the cytoplasm, while its disordered C-terminal tail localizes to the ER lumen [270]. It is proposed that bZIP28 senses ER stress through its interaction with the ER chaperone BiP (binding immunoglobulin protein), a master regulator of the ER stress sensor. Under non-stress conditions, BiP binds to bZIP28 IDRs present in its lumen-facing tail and retains it in the ER. Upon stress, BiP is competed away from bZIP28 by the accumulation of misfolded proteins in the ER, releasing bZIP28 and allowing its exit from the ER, to move towards the Golgi apparatus [271]. Then, bZIP28 is cleaved by proteases, releasing its transcriptional activation domain that will be translocated to the nucleus to up-regulate stress response genes [271]. The bZIP28 IDR represents one additional example of the role of IDRs in controlling signaling in plant stress responses.

Protein structural disorder in plant biotic stress responses

From germination to reproduction, plants confront a large diversity of parasitic organisms that can cause disease. These pathogens include viruses, bacteria, fungi, nematodes and insects that exploit resources and replication

systems in plants [272]. Infection by these organisms has driven plants to evolve refined mechanisms to detect their presence and to mount complex inducible responses to efficiently counteract their attack. As in other plant processes, plant defense systems are tightly regulated, many of them through the participation of kinases and phosphatases that modulate the phosphorylation status of key control proteins [272–274]. Marín and Ott [22] have reported the prediction and extensive compilation of different IDPs involved in this process. Because this information has been recently published, in the present work we include only a summary of the material for which functional and structural evidence is available.

Plants are able to specifically recognize their aggressors through receptors localized at the cell membrane. These receptors include LRR-RLKs, a common class of receptors in plants, where intrinsic disorder is present. An example of this is the aforementioned BAK1, an RLK that in this process functions as a co-receptor of the two of the best-characterized pathogen LRR-RLK receptors, FLS2 (Flagellin-sensing 2) and EFR (EF-Tu receptor) [275]. The relevance of the BAK1 IDR C-region resides in its ability to discriminate between two signal transduction pathways, even though the same phosphorylation site (Tyr-610) inside this region is involved in both brassinosteroid sensing and in the pathogen defense response [276]. It is well established that plant perception of pathogens is accompanied by an oxidative burst, where RbohD plays a central role (Fig. 2b). This protein belongs to the NADPH oxidase family, responsible for the early generation of ROS, upstream of calcium and protein phosphorylation signaling. Different experimental evidence supports the presence of an IDR in the RbohD cytoplasmic N terminus, a region that contains an EF-hand motif involved in calcium binding. The malleable nature of this region results in extended conformational changes induced by the synergistic effect of calcium binding and its phosphorylation, which in turn modulates the interaction with small GTPase proteins [277, 278]; fundamental events to set up protection responses to pathogenic agents (Fig. 2b). Following perception at the cell envelope, the signaling process continues in the cytosol, where different molecules play relevant roles. One of these protein molecules is the HSP90 molecular chaperone that given its refolding capacity, is an essential participant in many signaling pathways in plants and animals [279]. The structural organization of this chaperone shows an N-terminal region with an ATPase domain and a linker region composed of charged residues that connect its middle domain with the dimerization region localized at the C terminus. Interestingly, the N-terminal domain undergoes consecutive conformational changes upon ATP binding, leading to the formation of a transient dimer with different co-chaperone partners. The

association of HSP90 with the RAR1 co-chaperone results in an order-to-disorder transition of this ATP domain, which enables its movement to allow the entrance of the catalytic loop localized at the middle HSP90 region [280]. These interaction events are essential for the competence of RAR1 function, which together with the SGT1 co-chaperone, is needed to activate the majority of R proteins, detectors of pathogen effector molecules, by mediating NLR (nucleotide binding leucine-rich repeat receptor) function [281]. This signal pathway flows towards a MAP kinase cascade, whose activation ends in the phosphorylation of transcription factors (e.g., WRKY33) that induce the expression of defense genes. Two of these MAP kinases, MEK and MEKK1, show long disordered regions in their N termini that, in the case of MEKK1, have been shown to play a regulatory role; their removal results in a constitutively active kinase [22, 282]. The reprogramming of those genes encoding the proteins that will counteract pathogen incursion needs the action of transcription factors (TFs). Various TF families are involved in this process including MYC, MYB, TGA, WRKY and ERF. Among the TFs known to have a role in the plant pathogen response are MYC2, MYB30, TGA3, WRKY1, WRKY4, WRKY52, WRKY53 and ERF. All these proteins contain, in addition to their DNA-binding domains, IDR-containing linker domains with regulatory functions [21]. Some of these linker domain IDRs have been shown to interact with co-transcription factors that might contribute to the modulation of the spatio-temporal expression of target genes and to the selectivity required to distinguish the identity of particular pathogens (for review see [283]).

Computational analyses using available plant genome sequences predict the presence of significant structural disorder in many more proteins implicated in plant pathogen responses. However, as yet there is no experimental support for their structure or function. Hence, new discoveries await our curiosity and creativity.

Conclusions and future directions

Plants provide a clear picture of the importance of intrinsic disorder in eukaryote protein function. The structural flexibility and molecular promiscuity afforded to a wealth of plant proteins with intrinsically disordered domains have ensured pivotal and multifunctional roles in core processes, including development and metabolism as well as biotic and abiotic stress responses. Technical and experimental barriers to the study of IDPs have limited IDP research *in planta*, and up to now, there has been a strong reliance on interpretation and extrapolation from *in vitro* analyses; in particular, for those which are highly disordered and function under stress. It is hoped that the recent explosion

in molecular genetic technologies will pave the way for further exploration of the *in vivo* mechanisms and interactions of plant IDPs. We are only beginning to understand their place in the story of plant evolution and their essential functions in life as a whole.

Acknowledgements We thank Jose L. Reyes for critical reviewing this manuscript and for his kindness in helping us with the design and drawing of the figures in this manuscript. This work was partially supported by a Grant from CONACyT-México (221448). CLC-V and CC are supported by post-doctoral fellowships from CONACyT (221448) and Newton Fund-Mexican Academy of Sciences-CONACyT, respectively, and PSR-P and DFR-L by CONACyT PhD fellowships.

Appendix

Amphipathic helix A type of α -helix in which one face along the long axis of the helix is polar and the opposing face is nonpolar [284].

Chaperone A type of protein that ensures the proper folding of nascent polypeptidic chains by binding to them, stabilizing one structural conformation and releasing from the folded protein [285, 286].

Chloroplast signal recognition particle (CSRPs) A protein complex formed by cpSRP43 and cpSRP54 involved in the transport and insertion of proteins of the LHC into the membrane of thylakoids [287, 288].

Conformational selection A protein binding mechanism in which the binding protein structure shifts among different conformations and the ligand “selects” one such conformation, stabilizing the structure of the binding protein [289, 290].

Crassulacean acid metabolism (CAM) A photosynthetic pathway in which CO₂ is “stored” in the form of malate in the vacuole during the night and released during the day for RuBisCO-mediated carbon fixation [291].

Folding-upon-binding A mechanism in which IDPs or IDRs fold upon binding to their targets [292, 293].

Fuzzy complex A mechanism in which the parts of the proteins involved in the formation of macromolecular complexes do not fold, but remain in a disordered state [294].

Hydrodynamic radius The radius required for a perfect solid sphere to achieve the same frictional coefficient as the protein analyzed [295].

Hydrophilins Any protein that has a high glycine and/or small amino acid residues content (>6%) and a high hydrophilicity index (>1.0) [148, 167].

Intrinsically disordered protein (IDP) These proteins lack a fixed three-dimensional structure. These proteins fail to form a stable conformation and cannot be adequately described by a single equilibrium 3D structure, yet they exhibit biological activity [7, 296].

Intrinsically disordered region (IDR) Significant disordered sequences contained in a protein with structured globular domains [7].

Light-harvesting complex (LHC) In green plants, these complexes contain multiple proteins and molecules such as chlorophylls a and b and carotenoids, which increase the capacity for photon capture during the photosynthesis [1].

Molecular recognition feature (MORF) Protein structural element or feature that mediates the binding events of disordered regions with other proteins or nucleic acids. This element undergoes coupled binding and folding within a disorder region [296].

Moonlight activity The ability of a protein to fulfill several, apparently unrelated functions [229].

Orthodox seed Seeds that acquire desiccation tolerance during development maintain dormancy and might be stored under dry environments [297].

Store curvature elasticity stress Physical torque stress that occurs within a bilayer when lipids in the constituent monolayers are forced to adopt an unfavorable packing conformation. This stress results in hydrophobic cavities in the membrane, which are known as lipid-packing defects [298].

Oxygen-evolving complex (OEC) A Mn_4O_5Ca cluster embedded in the Photosystem II (PSII) protein complex. It is considered the active site of water oxidation where O_2 is produced and protons are released into the lumen during the light reactions of photosynthesis [299].

Unfolded protein response (UPR) Signaling pathways triggered to maintain a productive endoplasmic reticulum protein-folding environment [300].

References

- Jones R, Ougham H, Thomas H, Waaland S (2013) The molecular life of plants. Wiley-Blackwell, Hoboken
- Raven PH, Evert RF, Eichhorn SE (1999) Biology of plants. W.H. Freeman, Worth Publishers, New York
- Bögre L (2001) Cell signaling mechanisms in plants. In: eLS. Wiley, New York. doi:10.1002/9780470015902.a0020134
- Hetherington AM, Bardwell L (2011) Plant signaling pathways: a comparative evolutionary overview. *Curr Biol* 21(9):R317–R319. doi:10.1016/j.cub.2011.04.013
- Calabretta S, Richard S (2015) Emerging roles of disordered sequences in RNA-binding proteins. *Trends Biochem Sci* 40(11):662–672. doi:10.1016/j.tibs.2015.08.012
- Oldfield CJ, Dunker AK (2014) Intrinsically disordered proteins and intrinsically disordered protein regions. *Annu Rev Biochem* 83(1):553–584. doi:10.1146/annurev-biochem-072711-164947
- Wright PE, Dyson HJ (2015) Intrinsically disordered proteins in cellular signaling and regulation. *Nat Rev Mol Cell Biol* 16(1):18–29. doi:10.1038/Nrm3920
- Ward JJ, Sodhi JS, McGuffin LJ, Buxton BF, Jones DT (2004) Prediction and functional analysis of native disorder in proteins from the three kingdoms of life. *J Mol Biol* 337(3):635–645. doi:10.1016/j.jmb.2004.02.002
- Vucetic S, Brown CJ, Dunker AK, Obradovic Z (2003) Flavors of protein disorder. *Proteins* 52(4):573–584. doi:10.1002/prot.10437
- Schad E, Tompa P, Hegyi H (2011) The relationship between proteome size, structural disorder and organism complexity. *Genome Biol* 12(12):R120. doi:10.1186/gb-2011-12-12-r120
- Vogel C, Chothia C (2006) Protein family expansions and biological complexity. *PLoS Comput Biol* 2(5):e48. doi:10.1371/journal.pcbi.0020048
- Dosztanyi Z, Chen J, Dunker AK, Simon I, Tompa P (2006) Disorder and sequence repeats in hub proteins and their implications for network evolution. *J Proteome Res* 5(11):2985–2995. doi:10.1021/pr060171o
- Fuxreiter M, Tompa P, Simon I, Uversky VN, Hansen JC, Asturias FJ (2008) Malleable machines take shape in eukaryotic transcriptional regulation. *Nat Chem Biol* 4(12):728–737. doi:10.1038/nchembio.127
- Mittag T, Forman-Kay JD (2007) Atomic-level characterization of disordered protein ensembles. *Curr Opin Struct Biol* 17(1):3–14. doi:10.1016/j.sbi.2007.01.009
- Peng Z, Yan J, Fan X, Mizianty MJ, Xue B, Wang K, Hu G, Uversky VN, Kurgan L (2015) Exceptionally abundant exceptions: comprehensive characterization of intrinsic disorder in all domains of life. *Cell Mol Life Sci* 72(1):137–151. doi:10.1007/s00018-014-1661-9
- Pietrosemoli N, García-Martín JA, Solano R, Pazos F (2013) Genome-wide analysis of protein disorder in *Arabidopsis thaliana*: implications for plant environmental adaptation. *PLoS One* 8(2):e55524. doi:10.1371/journal.pone.0055524
- Sun X, Rikkerink EHA, Jones WT, Uversky VN (2013) Multifarious roles of intrinsic disorder in proteins illustrate its broad impact on plant biology. *Plant Cell* 25:38–55. doi:10.1105/tpc.112.106062
- Dunker AK, Bondos SE, Huang F, Oldfield CJ (2015) Intrinsically disordered proteins and multicellular organisms. *Semin Cell Dev Biol* 37:44–55. doi:10.1016/j.semcdb.2014.09.025
- Yruela I, Contreras-Moreira B (2013) Genetic recombination is associated with intrinsic disorder in plant proteomes. *BMC Genom* 14(1):772. doi:10.1186/1471-2164-14-772
- Yruela I, Contreras-Moreira B (2012) Protein disorder in plants: a view from the chloroplast. *BMC Plant Biol* 12:165. doi:10.1186/1471-2229-12-165
- Kragelund BB, Jensen MK, Skriver K (2012) Order by disorder in plant signaling. *Trends Plant Sci* 17(11):625–632. doi:10.1016/j.tplants.2012.06.010
- Marín M, Ott T (2014) Intrinsic disorder in plant proteins and phytopathogenic bacterial effectors. *Chem Rev* 114:6912–6932. doi:10.1021/cr400488d
- Thieulin-Pardo G, Avilan L, Kojadinovic M, Gontero B (2015) Fairy “tails”: flexibility and function of intrinsically disordered extensions in the photosynthetic world. *Front Mol Biosci* 2:23. doi:10.3389/fmolb.2015.00023
- Jensen MK, Kjaersgaard T, Nielsen MM, Galberg P, Petersen K, O’Shea C, Skriver K (2010) The *Arabidopsis thaliana* NAC transcription factor family: structure-function relationships and determinants of ANAC019 stress signaling. *Biochem J* 426:183–196. doi:10.1042/BJ20091234
- Taoka K, Yanagimoto Y, Daimon Y, Hibara K, Aida M, Tasaka M (2004) The NAC domain mediates functional specificity of CUP-SHAPED COTYLEDON proteins. *Plant J* 40(4):462–473. doi:10.1111/j.1365-3113X.2004.02238.x
- Ko JH, Yang SH, Park AH, Lerouxel O, Han KH (2007) ANAC012, a member of the plant-specific NAC transcription factor family, negatively regulates xylary fiber development in *Arabidopsis thaliana*. *Plant J* 50(6):1035–1048. doi:10.1111/j.1365-3113X.2007.03109.x

27. Yoon MK, Shin J, Choi G, Choi BS (2006) Intrinsically unstructured N-terminal domain of bZIP transcription factor HY5. *Proteins Struct Funct Bioinf* 65(4):856–866. doi:[10.1002/prot.21089](https://doi.org/10.1002/prot.21089)
28. Wang J, Jiang JJ, Wang J, Chen L, Fan SL, Wu JW, Wang XL, Wang ZX (2014) Structural insights into the negative regulation of BRI1 signaling by BRI1-interacting protein BKI1. *Cell Res* 24(11):1328–1341. doi:[10.1038/cr.2014.132](https://doi.org/10.1038/cr.2014.132)
29. Chakrabortee S, Kayatekin C, Newby GA, Mendillo ML, Lancaster A, Lindquist S (2016) Luminidependens (LD) is an *Arabidopsis* protein with prion behavior. *Proc Natl Acad Sci USA* 113(21):6065–6070. doi:[10.1073/pnas.1604478113](https://doi.org/10.1073/pnas.1604478113)
30. Valsecchi I, Guittard-Crilat E, Maldiney R, Habricot Y, Lignon S, Lebrun R, Miginiac E, Ruelland E, Jeannette E, Lebreton S (2013) The intrinsically disordered C-terminal region of *Arabidopsis thaliana* TCP8 transcription factor acts both as a transactivation and self-assembly domain. *Mol Biosyst* 9(9):2282–2295. doi:[10.1039/c3mb70128j](https://doi.org/10.1039/c3mb70128j)
31. Ingram GC, Waites R (2006) Keeping it together: co-ordinating plant growth. *Curr Opin Plant Biol* 9(1):12–20. doi:[10.1016/j.pbi.2005.11.007](https://doi.org/10.1016/j.pbi.2005.11.007)
32. Viola IL, Guttlein LN, Gonzalez DH (2013) Redox modulation of plant developmental regulators from the class I TCP transcription factor family. *Plant Physiol* 162(3):1434–1447. doi:[10.1104/pp.113.216416](https://doi.org/10.1104/pp.113.216416)
33. Aguilar-Martinez JA, Sinha N (2013) Analysis of the role of *Arabidopsis* class I TCP genes AtTCP7, AtTCP8, AtTCP22, and AtTCP23 in leaf development. *Front Plant Sci*. doi:[10.3389/fpls.2013.00406](https://doi.org/10.3389/fpls.2013.00406)
34. Hammani K, Gobert A, Hleibieh K, Choulier L, Small I, Giege P (2011) An *Arabidopsis* dual-localized pentatricopeptide repeat protein interacts with nuclear proteins involved in gene expression regulation. *Plant Cell* 23(2):730–740. doi:[10.1105/tpc.110.081638](https://doi.org/10.1105/tpc.110.081638)
35. Olsen AN, Ernst HA, Lo Leggio L, Skriver K (2005) NAC transcription factors: structurally distinct, functionally diverse. *Trends Plant Sci* 10(2):79–87. doi:[10.1016/j.tplants.2004.12.010](https://doi.org/10.1016/j.tplants.2004.12.010)
36. Sun XL, Rikkerink EHA, Jones WT, Uversky VN (2013) Multifarious roles of intrinsic disorder in proteins illustrate its broad impact on plant biology. *Plant Cell* 25(1):38–55. doi:[10.1105/tpc.112.106062](https://doi.org/10.1105/tpc.112.106062)
37. Kjaersgaard T, Jensen MK, Christiansen MW, Gregersen P, Kragelund BB, Skriver K (2011) Senescence-associated barley NAC (NAM, ATAF1,2, CUC) transcription factor interacts with radical-induced cell death 1 through a disordered regulatory domain. *J Biol Chem* 286(41):35418–35429. doi:[10.1074/jbc.M111.247221](https://doi.org/10.1074/jbc.M111.247221)
38. O'shea C, Kryger M, Stender EGP, Kragelund BB, Willemoes M, Skriver K (2015) Protein intrinsic disorder in *Arabidopsis* NAC transcription factors: transcriptional activation by ANAC013 and ANAC046 and their interactions with RCD1. *Biochem J* 465:281–294. doi:[10.1042/BJ20141045](https://doi.org/10.1042/BJ20141045)
39. Oyama T, Shimura Y, Okada K (1997) The *Arabidopsis* HY5 gene encodes a bZIP protein that regulates stimulus-induced development of root and hypocotyl. *Gene Dev* 11(22):2983–2995. doi:[10.1101/Gad.11.22.2983](https://doi.org/10.1101/Gad.11.22.2983)
40. Ahmad M (2016) Photocycle and signaling mechanisms of plant cryptochromes. *Curr Opin Plant Biol* 33:108–115. doi:[10.1016/j.pbi.2016.06.013](https://doi.org/10.1016/j.pbi.2016.06.013)
41. Chaves I, Pokorny R, Byrdin M, Hoang N, Ritz T, Brettel K, Essen LO, van der Horst GTJ, Batschauer A, Ahmad M (2011) The cryptochromes: blue light photoreceptors in plants and animals. *Annu Rev Plant Biol* 62:335–364. doi:[10.1146/annurev-arplant-042110-103759](https://doi.org/10.1146/annurev-arplant-042110-103759)
42. Cashmore AR, Jarillo JA, Wu YJ, Liu D (1999) Cryptochromes: blue light receptors for plants and animals. *Science* 284(5415):760–765
43. Lin C, Shalitin D (2003) Cryptochrome structure and signal transduction. *Annu Rev Plant Biol* 54:469–496. doi:[10.1146/annurev.arplant.54.110901.160901](https://doi.org/10.1146/annurev.arplant.54.110901.160901)
44. Liu B, Zuo Z, Liu H, Liu X, Lin C (2011) *Arabidopsis* cryptochrome 1 interacts with SPA1 to suppress COP1 activity in response to blue light. *Genes Dev* 25(10):1029–1034. doi:[10.1101/gad.2025011](https://doi.org/10.1101/gad.2025011)
45. Wang H, Ma L-G, Li J-M, Zhao H-Y, Deng XW (2001) Direct interaction of *Arabidopsis* cryptochromes with COP1 in light control development. *Science* 294(5540):154–158. doi:[10.1126/science.1063630](https://doi.org/10.1126/science.1063630)
46. Wang Q, Barshop WD, Bian M, Vashisht AA, He R, Yu X, Liu B, Nguyen P, Liu X, Zhao X, Wohlschlegel JA, Lin C (2015) The blue light-dependent phosphorylation of the CCE domain determines the photosensitivity of *Arabidopsis* CRY2. *Mol Plant* 8(4):631–643. doi:[10.1016/j.molp.2015.03.005](https://doi.org/10.1016/j.molp.2015.03.005)
47. Brautigam CA, Smith BS, Ma ZQ, Palnitkar M, Tomchick DR, Machius M, Deisenhofer J (2004) Structure of the photolyase-like domain of cryptochrome 1 from *Arabidopsis thaliana*. *Proc Natl Acad Sci USA* 101(33):12142–12147. doi:[10.1073/pnas.0404851101](https://doi.org/10.1073/pnas.0404851101)
48. Partch CL, Clarkson MW, Ozgur S, Lee AL, Sancar A (2005) Role of structural plasticity in signal transduction by the cryptochrome blue-light photoreceptor. *Biochem* 44(10):3795–3805. doi:[10.1021/bi047545g](https://doi.org/10.1021/bi047545g)
49. Kondoh M, Shiraishi C, Muller P, Ahmad M, Hitomi K, Getzoff ED, Terazima M (2011) Light-induced conformational changes in full-length *Arabidopsis thaliana* cryptochrome. *J Mol Biol* 413(1):128–137. doi:[10.1016/j.jmb.2011.08.031](https://doi.org/10.1016/j.jmb.2011.08.031)
50. Pfluger J, Wagner D (2007) Histone modifications and dynamic regulation of genome accessibility in plants. *Curr Opin Plant Biol* 10(6):645–652. doi:[10.1016/j.pbi.2007.07.013](https://doi.org/10.1016/j.pbi.2007.07.013)
51. Mehdi S, Derkacheva M, Ramstrom M, Kralemann L, Bergquist J, Hennig L (2016) The WD40 domain protein MSII functions in a histone deacetylase complex to fine-tune abscisic acid signaling. *Plant Cell* 28(1):42–54. doi:[10.1105/tpc.15.00763](https://doi.org/10.1105/tpc.15.00763)
52. Perrella G, Lopez-Vernaza MA, Carr C, Sani E, Gossele V, Verduyn C, Kellermeier F, Hannah MA, Amtmann A (2013) Histone deacetylase complex1 expression level titrates plant growth and abscisic acid sensitivity in *Arabidopsis*. *Plant Cell* 25(9):3491–3505. doi:[10.1105/tpc.113.114835](https://doi.org/10.1105/tpc.113.114835)
53. Perrella G, Carr C, Asensi-Fabado MA, Donald NA, Paldi K, Hannah MA, Amtmann A (2016) The histone deacetylase complex 1 protein of *Arabidopsis* has the capacity to interact with multiple proteins including histone 3-binding proteins and histone 1 variants. *Plant Physiol* 171(1):62–70. doi:[10.1104/pp.15.01760](https://doi.org/10.1104/pp.15.01760)
54. Clouse SD (2011) Brassinosteroid signal transduction: from receptor kinase activation to transcriptional networks regulating plant development. *Plant Cell* 23(4):1219–1230. doi:[10.1105/tpc.111.084475](https://doi.org/10.1105/tpc.111.084475)
55. Kim TW, Wang ZY (2010) Brassinosteroid signal transduction from receptor kinases to transcription factors. *Annu Rev Plant Biol* 61:681–704. doi:[10.1146/annurev.arplant.043008.092057](https://doi.org/10.1146/annurev.arplant.043008.092057)
56. Xie HB, Vucetic S, Iakoucheva LM, Oldfield CJ, Dunker AK, Uversky VN, Obradovic Z (2007) Functional anthology of intrinsic disorder. 1. Biological processes and functions of proteins with long disordered regions. *J Proteome Res* 6(5):1882–1898. doi:[10.1021/pr060392u](https://doi.org/10.1021/pr060392u)
57. Nam KH, Li JM (2002) BRI1/BAK1, a receptor kinase pair mediating brassinosteroid signaling. *Cell* 110(2):203–212. doi:[10.1016/S0092-8674\(02\)00814-0](https://doi.org/10.1016/S0092-8674(02)00814-0)

58. Wang XL, Chory J (2006) Brassinosteroids regulate dissociation of BKII, a negative regulator of BRI1 signaling, from the plasma membrane. *Science* 313(5790):1118–1122. doi:[10.1126/science.1127593](https://doi.org/10.1126/science.1127593)
59. Wang XF, Kota U, He K, Blackburn K, Li J, Goshe MB, Huber SC, Clouse SD (2008) Sequential transphosphorylation of the BRI1/BAK1 receptor kinase complex impacts early events in brassinosteroid signaling. *Dev Cell* 15(2):220–235. doi:[10.1016/j.devcel.2008.06.011](https://doi.org/10.1016/j.devcel.2008.06.011)
60. Hothorn M, Belkhadir Y, Dreux M, Dabi T, Noel JP, Wilson IA, Chory J (2011) Structural basis of steroid hormone perception by the receptor kinase BRI1. *Nature* 474(7352):467–471. doi:[10.1038/nature10153](https://doi.org/10.1038/nature10153)
61. Jiang JJ, Wang T, Wu ZH, Wang J, Zhang C, Wang HJ, Wang ZX, Wang XL (2015) The intrinsically disordered protein BKII is essential for inhibiting BRI1 signaling in plants. *Mol Plant* 8(11):1675–1678. doi:[10.1016/j.molp.2015.07.012](https://doi.org/10.1016/j.molp.2015.07.012)
62. Song YH, Ito S, Imaizumi T (2013) Flowering time regulation: photoperiod- and temperature-sensing in leaves. *Trends Plant Sci* 18(10):575–583. doi:[10.1016/j.tplants.2013.05.003](https://doi.org/10.1016/j.tplants.2013.05.003)
63. Schmitz RJ, Amasino RM (2007) Vernalization: a model for investigating epigenetics and eukaryotic gene regulation in plants. *BBA Gene Struct Expr* 1769(5–6):269–275. doi:[10.1016/j.bbaexp.2007.02.003](https://doi.org/10.1016/j.bbaexp.2007.02.003)
64. Shorter J, Lindquist S (2005) Prions as adaptive conduits of memory and inheritance. *Nat Rev Genet* 6(6):435–450. doi:[10.1038/nrg1616](https://doi.org/10.1038/nrg1616)
65. Mukhopadhyay S, Krishnan R, Lemke EA, Lindquist S, Deniz AA (2007) A natively unfolded yeast prion monomer adopts an ensemble of collapsed and rapidly fluctuating structures. *Proc Natl Acad Sci USA* 104(8):2649–2654. doi:[10.1073/pnas.0611503104](https://doi.org/10.1073/pnas.0611503104)
66. Malinowska L, Kroschwald S, Alberti S (2013) Protein disorder, prion propensities, and self-organizing macromolecular collectives. *BBA Proteins Proteom* 1834(5):918–931. doi:[10.1016/j.bbapap.2013.01.003](https://doi.org/10.1016/j.bbapap.2013.01.003)
67. Romero P, Obradovic Z, Li XH, Garner EC, Brown CJ, Dunker AK (2001) Sequence complexity of disordered protein. *Proteins Struct Funct Genet* 42(1):38–48. doi:[10.1002/1097-0134\(20010101\)42:1<38:Aid-Prot50>3.0.Co;2-3](https://doi.org/10.1002/1097-0134(20010101)42:1<38:Aid-Prot50>3.0.Co;2-3)
68. Sun XL, Jones WT, Rikkerink EHA (2012) GRAS proteins: the versatile roles of intrinsically disordered proteins in plant signaling. *Biochem J* 442:1–12. doi:[10.1042/BJ20111766](https://doi.org/10.1042/BJ20111766)
69. Sun X, Xue B, Jones WT, Rikkerink E, Dunker AK, Uversky VN (2011) A functionally required unfoldome from the plant kingdom: intrinsically disordered N-terminal domains of GRAS proteins are involved in molecular recognition during plant development. *Plant Mol Biol* 77:205–223. doi:[10.1007/s11103-011-9803-z](https://doi.org/10.1007/s11103-011-9803-z)
70. Sun XL, Jones WT, Harvey D, Edwards PJB, Pascal SM, Kirk C, Considine T, Sheerin DJ, Rakonjac J, Oldfield CJ, Xue B, Dunker AK, Uversky VN (2010) N-terminal domains of DELLA proteins are intrinsically unstructured in the absence of interaction with GID1/gibberellic acid receptors. *J Biol Chem* 285(15):11557–11571. doi:[10.1074/jbc.M109.027011](https://doi.org/10.1074/jbc.M109.027011)
71. Daviere JM, Achard P (2016) A pivotal role of DELLAs in regulating multiple hormone signals. *Mol Plant* 9(1):10–20. doi:[10.1016/j.molp.2015.09.011](https://doi.org/10.1016/j.molp.2015.09.011)
72. Smertenko AP, Chang H-Y, Wagner V, Kaloriti D, Fenyk S, Sonobe S, Lloyd C, Hauser M-T, Hussey PJ (2004) The *Arabidopsis* microtubule-associated protein AtMAP65-1: molecular analysis of its microtubule bundling activity. *Plant Cell* 16(8):2035–2047. doi:[10.1105/tpc.104.023937](https://doi.org/10.1105/tpc.104.023937)
73. Hussey PJ, Hawkins TJ, Igarashi H, Kaloriti D, Smertenko A (2002) The plant cytoskeleton: recent advances in the study of the plant microtubule-associated proteins MAP-65, MAP-190 and the *Xenopus* MAP215-like protein, MOR1. *Plant Mol Biol* 50(6):915–924. doi:[10.1023/A:1021236307508](https://doi.org/10.1023/A:1021236307508)
74. Smertenko AP, Kaloriti D, Chang H-Y, Fiserova J, Opatrný Z, Hussey PJ (2008) The C-terminal variable region specifies the dynamic properties of *Arabidopsis* microtubule-associated protein MAP65 isoforms. *Plant Cell* 20(12):3346–3358. doi:[10.1105/tpc.108.063362](https://doi.org/10.1105/tpc.108.063362)
75. Boruc J, Weimer A, Stoppin-Mellet V, Mylle E, Kosetsu K, Cedeno C, Jaquinod M, Njo M, De Milde L, Tompa P, Gonzalez N, Inze D, Beeckman T, Vantard M, van Damme D (2016) Phosphorylation of MAP65-1 by *Arabidopsis* Aurora kinases is required for efficient cell cycle progression. *Plant Physiol.* doi:[10.1104/pp.16.01602](https://doi.org/10.1104/pp.16.01602)
76. Matzke MA, Mosher RA (2014) RNA-directed DNA methylation: an epigenetic pathway of increasing complexity. *Nat Rev Genet* 15(6):394–408. doi:[10.1038/nrg3683](https://doi.org/10.1038/nrg3683)
77. Blevins T, Podicheti R, Mishra V, Marasco M, Wang J, Rusch D, Tang H, Pikaard CS (2015) Identification of Pol IV and RDR2-dependent precursors of 24 nt siRNAs guiding de novo DNA methylation in *Arabidopsis*. *Elife* 4:e09591. doi:[10.7554/eLife.09591](https://doi.org/10.7554/eLife.09591)
78. Henderson IR, Zhang X, Lu C, Johnson L, Meyers BC, Green PJ, Jacobsen SE (2006) Dissecting *Arabidopsis* thaliana DICER function in small RNA processing, gene silencing and DNA methylation patterning. *Nat Genet* 38(6):721–725. doi:[10.1038/ng1804](https://doi.org/10.1038/ng1804)
79. Yu B, Bi L, Zhai J, Agarwal M, Li S, Wu Q, Ding SW, Meyers BC, Vaucheret H, Chen X (2010) siRNAs compete with miRNAs for methylation by HEN1 in *Arabidopsis*. *Nucleic Acids Res* 38(17):5844–5850. doi:[10.1093/nar/gkq348](https://doi.org/10.1093/nar/gkq348)
80. Ye R, Wang W, Iki T, Liu C, Wu Y, Ishikawa M, Zhou X, Qi Y (2012) Cytoplasmic assembly and selective nuclear import of *Arabidopsis* Argonaute4/siRNA complexes. *Mol Cell* 46(6):859–870. doi:[10.1016/j.molcel.2012.04.013](https://doi.org/10.1016/j.molcel.2012.04.013)
81. Wierzbicki AT, Ream TS, Haag JR, Pikaard CS (2009) RNA polymerase V transcription guides ARGONAUTE4 to chromatin. *Nat Genet* 41(5):630–634. doi:[10.1038/ng.365](https://doi.org/10.1038/ng.365)
82. Rowley MJ, Avrutsky MI, Sifuentes CJ, Pereira L, Wierzbicki AT (2011) Independent chromatin binding of ARGONAUTE4 and SPT5L/KTF1 mediates transcriptional gene silencing. *PLoS Genet* 7(6):e1002120. doi:[10.1371/journal.pgen.1002120](https://doi.org/10.1371/journal.pgen.1002120)
83. Matzke MA, Kanno T, Matzke AJ (2015) RNA-directed DNA methylation: the evolution of a complex epigenetic pathway in flowering plants. *Annu Rev Plant Biol* 66:243–267. doi:[10.1146/annurev-arplant-043014-114633](https://doi.org/10.1146/annurev-arplant-043014-114633)
84. Trujillo JT, Beilstein MA, Mosher RA (2016) The Argonaute-binding platform of NRPE1 evolves through modulation of intrinsically disordered repeats. *New Phytol* 212(4):1094–1105. doi:[10.1111/nph.14089](https://doi.org/10.1111/nph.14089)
85. Till S, Lejeune E, Thermann R, Bortfeld M, Hothorn M, Enderle D, Heinrich C, Hentze MW, Ladurner AG (2007) A conserved motif in Argonaute-interacting proteins mediates functional interactions through the Argonaute PIWI domain. *Nat Struct Mol Biol* 14(10):897–903. doi:[10.1038/nsmb1302](https://doi.org/10.1038/nsmb1302)
86. El-Shami M, Pontier D, Lahmy S, Braun L, Picart C, Vega D, Hakimi MA, Jacobsen SE, Cooke R, Lagrange T (2007) Reiterated WG/GW motifs form functionally and evolutionarily conserved ARGONAUTE-binding platforms in RNAi-related components. *Genes Dev* 21(20):2539–2544. doi:[10.1101/gad.451207](https://doi.org/10.1101/gad.451207)
87. Pfaff J, Meister G (2013) Argonaute and GW182 proteins: an effective alliance in gene silencing. *Biochem Soc Trans* 41(4):855–860. doi:[10.1042/bst20130047](https://doi.org/10.1042/bst20130047)
88. Radivojac P, Clark WT, Oron TR, Schnoes AM, Wittkop T, Sokolov A, Graim K, Funk C, Verspoor K, Ben-Hur A, Pandey G, Yunes JM, Talwalkar AS, Repo S, Souza ML, Piovesan D,

- Casadio R, Wang Z, Cheng J, Fang H, Gough J, Koskinen P, Toronen P, Nokso-Koivisto J, Holm L, Cozzetto D, Buchan DW, Bryson K, Jones DT, Limaye B, Inamdar H, Datta A, Manjari SK, Joshi R, Chitale M, Kihara D, Lisewski AM, Erdin S, Venner E, Lichtarge O, Rentzsch R, Yang H, Romero AE, Bhat P, Paccanaro A, Hamp T, Kassner R, Seemayer S, Vicedo E, Schaefer C, Achten D, Auer F, Boehm A, Braun T, Hecht M, Heron M, Honigschmid P, Hopf TA, Kaufmann S, Kiening M, Krompass D, Landerer C, Mahlich Y, Roos M, Bjorne J, Salakoski T, Wong A, Shatka H, Gatzmann F, Sommer I, Wass MN, Sternberg MJ, Skunca N, Supek F, Bosnjak M, Panov P, Dzeroski S, Smuc T, Kourmpetis YA, van Dijk AD, ter Braak CJ, Zhou Y, Gong Q, Dong X, Tian W, Falda M, Fontana P, Lavezzo E, Di Camillo B, Toppo S, Lan L, Djuric N, Guo Y, Vucetic S, Bairoch A, Linial M, Babbitt PC, Brenner SE, Orengo C, Rost B, Mooney SD, Friedberg I (2013) A large-scale evaluation of computational protein function prediction. *Nat Methods* 10(3):221–227. doi:10.1038/nmeth.2340
89. Fersht AR, Knill-Jones JW, Bedouelle H, Winter G (1988) Reconstruction by site-directed mutagenesis of the transition state for the activation of tyrosine by the tyrosyl-tRNA synthetase: a mobile loop envelops the transition state in an induced-fit mechanism. *Biochemistry* 27(5):1581–1587. doi:10.1021/bi00405a028
90. McElheny D, Schnell JR, Lansing JC, Dyson HJ, Wright PE (2005) Defining the role of active-site loop fluctuations in dihydrofolate reductase catalysis. *Proc Natl Acad Sci USA* 102(14):5032–5037. doi:10.1073/pnas.0500699102
91. Schulenburg C, Hilvert D (2013) Protein conformational disorder and enzyme catalysis. *Top Curr Chem* 337:41–67. doi:10.1007/128_2012_411
92. Erales J, Lorenzi M, Lebrun R, Fournel A, Etienne E, Courcelle C, Guigliarelli B, Gontero B, Belle V (2009) A new function of GAPDH from *Chlamydomonas reinhardtii*: a thiol-disulfide exchange reaction with CP12. *Biochemistry* 48:6034–6040. doi:10.1021/bi900569h
93. Graciet E, Gans P, Wedel N, Lebreton S, Camadro JM, Gontero B (2003) The small protein CP12: a protein linker for supramolecular complex assembly. *Biochemistry* 42:8163–8170. doi:10.1021/bi034474x
94. Groben R, Kaloudas D, Raines CA, Offmann B, Maberly SC, Gontero B (2010) Comparative sequence analysis of CP12, a small protein involved in the formation of a Calvin cycle complex in photosynthetic organisms. *Photosynth Res* 103:183–194. doi:10.1007/s11120-010-9542-z
95. Stanley DN, Raines CA, Kerfeld CA (2013) Comparative analysis of 126 cyanobacterial genomes reveals evidence of functional diversity among homologs of the redox-regulated CP12 protein. *Plant Physiol* 161(2):824–835. doi:10.1104/pp.112.210542
96. Fermani S, Trivelli X, Sparla F, Thumiger A, Calvaresi M, Marri L, Falini G, Zerbetto F, Trost P (2012) Conformational selection and folding-upon-binding of intrinsically disordered protein CP12 regulate photosynthetic enzymes assembly. *J Biol Chem* 287:21372–21383. doi:10.1074/jbc.M112.350355
97. Mileo E, Lorenzi M, Erales J, Lignon S, Puppo C, Le Breton N, Etienne E, Marque SRA, Guigliarelli B, Gontero B, Belle V (2013) Dynamics of the intrinsically disordered protein CP12 in its association with GAPDH in the green alga *Chlamydomonas reinhardtii*: a fuzzy complex. *Mol Biosyst* 9:2869–2876. doi:10.1039/c3mb70190e
98. Moparthi SB, Thieulin-Pardo G, Mansuelle P, Rigneault H, Gontero B, Wenger J (2014) Conformational modulation and hydrodynamic radii of CP12 protein and its complexes probed by fluorescence correlation spectroscopy. *FEBS J* 281:3206–3217. doi:10.1111/febs.12854
99. Marri L, Thieulin-Pardo G, Lebrun R, Puppo R, Zaffagnini M, Trost P, Gontero B, Sparla F (2014) CP12-mediated protection of Calvin-Benson cycle enzymes from oxidative stress. *Biochimie* 97:228–237. doi:10.1016/j.biochi.2013.10.018
100. Lopez-Calcagno PE, Howard TP, Raines CA (2014) The CP12 protein family: a thioredoxin-mediated metabolic switch? *Front Plant Sci* 5:9. doi:10.3389/fpls.2014.00009
101. Pohlmeier K, Paap BK, Soll J, Wedel N (1996) CP12: a small nuclear-encoded chloroplast protein provides novel insights into higher-plant GAPDH evolution. *Plant Mol Biol* 32:969–978. doi:10.1007/BF00020493
102. Matsumura H, Kai A, Maeda T, Tamoi M, Satoh A, Tamura H, Hirose M, Ogawa T, Kizu N, Wadano A, Inoue T, Shigeoka S (2011) Structure basis for the regulation of glyceraldehyde-3-phosphate dehydrogenase activity via the intrinsically disordered protein CP12. *Structure* 19:1846–1854. doi:10.1016/j.str.2011.08.016
103. Sparla F, Pupillo P, Trost P (2002) The C-terminal extension of glyceraldehyde-3-phosphate dehydrogenase subunit B acts as an autoinhibitory domain regulated by thioredoxins and nicotinamide adenine dinucleotide. *J Biol Chem* 277(47):44946–44952. doi:10.1074/jbc.M206873200
104. Ellis RJ (1979) The most abundant protein in the world. *Trends Biochem Sci* 4:241–244. doi:10.1016/0968-0004(79)90212-3
105. Raven JA (2013) RuBisCO: still the most abundant protein of Earth? *New Phytol* 198:1–3. doi:10.1111/nph.12197
106. Cleland WW, Andrews TJ, Gutteridge S, Hartman FC, Lorimer GH (1998) Mechanism of RuBisCO: the carbamate as general base. *Chem Rev* 98:549–562. doi:10.1021/cr970010r
107. Gontero B, Salvucci ME (2014) Regulation of photosynthetic carbon metabolism in aquatic and terrestrial organisms by RuBisCO activase, redox-modulation and CP12. *Aquat Bot* 118:14–23. doi:10.1016/j.aquabot.2014.05.011
108. Zhang N, Portis AR (1999) Mechanism of light regulation of RuBisCO: a specific role for the larger RuBisCO activase isoform involving reductive activation by Thioredoxin-f. *Proc Natl Acad Sci USA* 96:9438–9443. doi:10.1073/pnas.96.16.9438
109. Stotz M, Mueller-Cajar O, Ciniawsky S, Wendler P, Hartl FU, Bracher A, Hayer-Hartl M (2011) Structure of green-type RuBisCO activase from tobacco. *Nat Struct Mol Biol* 18:1366–1370. doi:10.1038/nsmb.2171
110. Shen JB, Ogren WL (1992) Alteration of spinach ribulose-1,5-bisphosphate carboxylase/oxygenase activase activities by site-directed mutagenesis. *Plant Physiol* 99(3):1201–1207. doi:10.1104/pp.99.3.1201
111. Wang D, Portis AR (2006) Increased sensitivity of oxidized large isoform of ribulose-1,5-bisphosphate carboxylase/oxygenase (RuBisCO) activase to ADP inhibition is due to an interaction between its carboxyl extension and nucleotide-binding pocket. *J Biol Chem* 281:25241–25249. doi:10.1074/jbc.M604756200
112. Carmo-Silva AE, Salvucci ME (2013) The regulatory properties of RuBisCO activase differ among species and affect photosynthetic induction during light transitions. *Plant Physiol* 161:1645–1655. doi:10.1104/pp.112.213348
113. Zarzycki J, Axen SD, Kinney JN, Kerfeld CA (2013) Cyanobacterial-based approaches to improving photosynthesis in plants. *J Exp Bot* 64(3):787–798. doi:10.1093/jxb/ers294
114. Seidler A (1996) The extrinsic polypeptides of photosystem II. *Biochim Biophys Acta Bioenerg* 1277:35–60. doi:10.1016/S0005-2728(96)00102-8
115. Lydakis-Simantiris N, Betts SD, Yocum CF (1999) Leucine 245 is a critical residue for folding and function of the manganese stabilizing protein of photosystem II. *Biochemistry* 38:15528–15535. doi:10.1021/bi991599m

116. Shutova T, Irrgang KD, Shubin V, Klimov VV, Renger G (1997) Analysis of pH-induced structural changes of the isolated extrinsic 33 kilodalton protein of photosystem II. *Biochemistry* 36:6350–6358. doi:[10.1021/bi963115h](https://doi.org/10.1021/bi963115h)
117. Yi X, McChargue M, Laborde S, Frankel LK, Bricker TM (2005) The manganese-stabilizing protein is required for photosystem II assembly/stability and photoautotrophy in higher plants. *J Biol Chem* 280(16):16170–16174. doi:[10.1074/jbc.M501550200](https://doi.org/10.1074/jbc.M501550200)
118. Lydakakis-Simantiris N, Hutchison RS, Betts SD, Barry BA, Yocum CF (1999) Manganese stabilizing protein of photosystem II is a thermostable, natively unfolded polypeptide. *Biochemistry* 38:404–414. doi:[10.1021/bi981847z](https://doi.org/10.1021/bi981847z)
119. Popelkova H, Betts SD, Lydakakis-Symantiris N, Im MM, Swenson E, Yocum CF (2006) Mutagenesis of basic residues R151 and R161 in manganese-stabilizing protein of photosystem II causes inefficient binding of chloride to the oxygen-evolving complex. *Biochemistry* 45:3107–3115. doi:[10.1021/bi0523759](https://doi.org/10.1021/bi0523759)
120. Falk S, Ravaud S, Koch J, Sinning I (2010) The C terminus of the Alb3 membrane insertase recruits cpSRP43 to the thylakoid membrane. *J Biol Chem* 285:5954–5962. doi:[10.1074/jbc.M109.084996](https://doi.org/10.1074/jbc.M109.084996)
121. Krolczewski J, Piskozub M, Bartoszewski R, Krolczewska B (2016) ALB3 insertase mediates cytochrome b6 co-translational import into the thylakoid membrane. *Sci Rep* 6:34557. doi:[10.1038/srep34557](https://doi.org/10.1038/srep34557)
122. Brynion EA, Rankin SE, Carey M, Watts A, Pinheiro TJ (1999) Folding of apocytochrome c in lipid micelles: formation of alpha-helix precedes membrane insertion. *Biochemistry* 38(30):9758–9767. doi:[10.1021/bi990119o](https://doi.org/10.1021/bi990119o)
123. Urbiscek M, Nick von Braun S, Brylok T, Gügel IL, Richter A, Koskela M, Grimm B, Mulo P, Bölter B, Soll J, Ankele E, Schwenkert S (2015) The extreme Albino3 (Alb3) C terminus is required for Alb3 stability and function in *Arabidopsis thaliana*. *Planta* 242(3):733–746. doi:[10.1007/s00425-015-2352-y](https://doi.org/10.1007/s00425-015-2352-y)
124. Marusek CM, Trobaugh NM, Flurkey WH, Inlow JK (2006) Comparative analysis of polyphenol oxidase from plant and fungal species. *J Inorg Biochem* 100:108–123. doi:[10.1016/j.jinorgbio.2005.10.008](https://doi.org/10.1016/j.jinorgbio.2005.10.008)
125. O'Leary B, Park J, Plaxton WC (2011) The remarkable diversity of plant PEPC (phosphoenolpyruvate carboxylase): recent insights into the physiological functions and post-translational controls of non-photosynthetic PEPCs. *Biochem J* 436:15–34. doi:[10.1042/BJ20110078](https://doi.org/10.1042/BJ20110078)
126. Dalziel KJ, O'Leary B, Brikis C, Rao SK, She Y-M, Cyr T, Plaxton WC (2012) The bacterial-type phosphoenolpyruvate carboxylase isozyme from developing castor oil seeds is subject to in vivo regulatory phosphorylation at serine-451. *FEBS Lett* 586:1049–1054. doi:[10.1016/j.febslet.2012.02.054](https://doi.org/10.1016/j.febslet.2012.02.054)
127. Musiani F, Ippoliti E, Micheletti C, Carloni P, Ciurli S (2013) Conformational fluctuations of UreG, an intrinsically disordered enzyme. *Biochemistry* 52:2949–2954. doi:[10.1021/bi4001744](https://doi.org/10.1021/bi4001744)
128. Neyroz P, Zambelli B, Ciurli S (2006) Intrinsically disordered structure of *Bacillus pasteurii* UreG as revealed by steady-state and time-resolved fluorescence spectroscopy. *Biochemistry* 45:8918–8930. doi:[10.1021/bi060227s](https://doi.org/10.1021/bi060227s)
129. Zambelli B, Cremades N, Neyroz P, Turano P, Uversky VN, Ciurli S (2012) Insights in the (un)structural organization of *Bacillus pasteurii* UreG, an intrinsically disordered GTPase enzyme. *Mol BioSyst* 8:220–228. doi:[10.1039/c1mb05227f](https://doi.org/10.1039/c1mb05227f)
130. Real-Guerra R, Staniscuaski F, Zambelli B, Musiani F, Ciurli S, Carlini CR (2012) Biochemical and structural studies on native and recombinant *Glycine max* UreG: a detailed characterization of a plant urease accessory protein. *Plant Mol Biol* 78(4–5):461–475. doi:[10.1007/s11103-012-9878-1](https://doi.org/10.1007/s11103-012-9878-1)
131. Karplus PA, Pearson MA, Hausinger RP (1997) 70 years of crystalline urease: what have we learned? *Acc Chem Res* 30:330–337. doi:[10.1021/ar960022j](https://doi.org/10.1021/ar960022j)
132. Carlini CR, Oliveira AE, Azambuja P, Xavier-Filho J, Wells MA (1997) Biological effects of canatoxin in different insect models: evidence for a proteolytic activation of the toxin by insect cathepsin-like enzymes. *J Econ Entomol* 90:340–348. doi:[10.1093/jee/90.2.340](https://doi.org/10.1093/jee/90.2.340)
133. Staniscuaski F, Ferreira-Dasilva CT, Mulinari F, Pires-Alves M, Carlini CR (2005) Insecticidal effects of canatoxin on the cotton stainer bug *Dysdercus peruvianus* (Hemiptera: Pyrrhocoridae). *Toxicon* 45(6):753–760. doi:[10.1016/j.toxicon.2005.01.014](https://doi.org/10.1016/j.toxicon.2005.01.014)
134. Mulinari F, Staniscuaski F, Bertholdo-Vargas LR, Postal M, Oliveira-Neto OB, Rigden DJ, Grossi-de-Sá MF, Carlini CR (2007) Jaburetox-2Ec: an insecticidal peptide derived from an isoform of urease from the plant *Canavalia ensiformis*. *Peptides* 28:2042–2050. doi:[10.1016/j.peptides.2007.08.009](https://doi.org/10.1016/j.peptides.2007.08.009)
135. Menez A (1998) Functional architectures of animal toxins: a clue to drug design? *Toxicon* 36(11):1557–1572. doi:[10.1016/S0041-0101\(98\)00148-2](https://doi.org/10.1016/S0041-0101(98)00148-2)
136. Lopes FC, Dobrovolska O, Real-Guerra R, Broll V, Zambelli B, Musiani F, Uversky VN, Carlini CR, Ciurli S (2015) Pliable natural biocide: Jaburetox is an intrinsically disordered insecticidal and fungicidal polypeptide derived from jack bean urease. *FEBS J* 282:1043–1064. doi:[10.1111/febs.13201](https://doi.org/10.1111/febs.13201)
137. Battaglia M, Olvera-Carrillo Y, Garcarrubio A, Campos F, Covarrubias AA (2008) The enigmatic LEA proteins and other hydrophilins. *Plant Physiol* 148(1):6–24. doi:[10.1104/pp.108.120725](https://doi.org/10.1104/pp.108.120725)
138. Rivera-Najera LY, Saab-Rincon G, Battaglia M, Amero C, Pulido NO, Garcia-Hernandez E, Solorzano RM, Reyes JL, Covarrubias AA (2014) A group 6 late embryogenesis abundant protein from common bean is a disordered protein with extended helical structure and oligomer-forming properties. *J Biol Chem* 289(46):31995–32009. doi:[10.1074/jbc.M114.583369](https://doi.org/10.1074/jbc.M114.583369)
139. Cuevas-Velazquez CL, Saab-Rincon G, Reyes JL, Covarrubias AA (2016) The unstructured N-terminal region of *Arabidopsis* group 4 late embryogenesis abundant (LEA) proteins is required for folding and for chaperone-like activity under water deficit. *J Biol Chem* 291(20):10893–10903. doi:[10.1074/jbc.M116.720318](https://doi.org/10.1074/jbc.M116.720318)
140. Shih M-D, Hoekstra FA, Hsing Y-IC (2008) Late embryogenesis abundant proteins. *Adv Bot Res* 48:211–255. doi:[10.1016/S0065-2296\(08\)00404-7](https://doi.org/10.1016/S0065-2296(08)00404-7)
141. Hand SC, Jones D, Menze MA, Witt TL (2007) Life without water: expression of plant LEA genes by an anhydrobiotic arthropod. *J Exp Zool A Ecol Genet Physiol* 307(1):62–66. doi:[10.1002/jez.a.343](https://doi.org/10.1002/jez.a.343)
142. Stacy RAP, Aalen RB (1998) Identification of sequence homology between the internal hydrophilic repeated motifs of Group 1 late-embryogenesis-abundant proteins in plants and hydrophilic repeats of the general stress protein GsiB of *Bacillus subtilis*. *Planta* 206(3):476–478. doi:[10.1007/S004250050424](https://doi.org/10.1007/S004250050424)
143. Tunnacliffe A, Lapinski J, McGee B (2005) A putative LEA protein, but no trehalose, is present in anhydrobiotic bdelloid rotifers. *Hydrobiologia* 546:315–321. doi:[10.1007/s10750-005-4239-6](https://doi.org/10.1007/s10750-005-4239-6)
144. Browne J, Tunnacliffe A, Burnell A (2002) Anhydrobiosis: plant desiccation gene found in a nematode. *Nature* 416(6876):38. doi:[10.1038/416038a](https://doi.org/10.1038/416038a)
145. Hand SC, Menze MA, Toner M, Boswell L, Moore D (2011) LEA proteins during water stress: not just for plants anymore. *Annu Rev Physiol* 73:115–134. doi:[10.1146/annurev-physiol-012110-142203](https://doi.org/10.1146/annurev-physiol-012110-142203)

146. Hinch DK, Thalhammer A (2012) LEA proteins: IDPs with versatile functions in cellular dehydration tolerance. *Biochem Soc Trans* 40:1000–1003. doi:10.1042/Bst20120109
147. Hundertmark M, Popova AV, Rausch S, Seckler R, Hinch DK (2012) Influence of drying on the secondary structure of intrinsically disordered and globular proteins. *Biochem Biophys Res Commun* 417(1):122–128. doi:10.1016/J.Bbrc.2011.11.067
148. Garay-Arroyo A, Colmenero-Flores JM, Garcarrubio A, Covarrubias AA (2000) Highly hydrophilic proteins in prokaryotes and eukaryotes are common during conditions of water deficit. *J Biol Chem* 275(8):5668–5674. doi:10.1074/jbc.275.8.5668
149. Reyes JL, Rodrigo MJ, Colmenero-Flores JM, Gil JV, Garay-Arroyo A, Campos F, Salamini F, Bartels D, Covarrubias AA (2005) Hydrophilins from distant organisms can protect enzymatic activities from water limitation effects in vitro. *Plant Cell Environ* 28(6):709–718. doi:10.1111/J.1365-3040.2005.01317.X
150. Goyal K, Walton LJ, Tunnacliffe A (2005) LEA proteins prevent protein aggregation due to water stress. *Biochem J* 388:151–157. doi:10.1042/BJ20041931
151. Sanchez-Ballesta MT, Rodrigo MJ, LaFuente MT, Granell A, Zacarias L (2004) Dehydrin from citrus, which confers in vitro dehydration and freezing protection activity, is constitutive and highly expressed in the flavedo of fruit but responsive to cold and water stress in leaves. *J Agr Food Chem* 52(7):1950–1957. doi:10.1021/Jf035216
152. Tantos A, Friedrich P, Tompa P (2009) Cold stability of intrinsically disordered proteins. *FEBS Lett* 583(2):465–469. doi:10.1016/J.Febslet.2008.12.054
153. Brini F, Saibi W, Amara I, Gargouri A, Masmoudi K, Hanin M (2010) Wheat dehydrin DHN-5 exerts a heat-protective effect on beta-glucosidase and glucose oxidase activities. *Biosci Biotech Biochem* 74(5):1050–1054. doi:10.1271/Bbb.90949
154. Drira M, Saibi W, Brini F, Gargouri A, Masmoudi K, Hanin M (2013) The K-segments of the wheat dehydrin DHN-5 are essential for the protection of lactate dehydrogenase and beta-glucosidase activities in vitro. *Mol Biotechnol* 54(2):643–650. doi:10.1007/s12033-012-9606-8
155. Campos F, Cuevas-Velazquez C, Fares MA, Reyes JL, Covarrubias AA (2013) Group 1 LEA proteins, an ancestral plant protein group, are also present in other eukaryotes, and in the archaea and bacteria domains. *Mol Genet Genom* 288(10):503–517. doi:10.1007/s00438-013-0768-2
156. Grelet J, Benamar A, Teyssier E, Avelange-Macherel MH, Grunwald D, Macherel D (2005) Identification in pea seed mitochondria of a late-embryogenesis abundant protein able to protect enzymes from drying. *Plant Physiol* 137(1):157–167. doi:10.1104/pp.104.052480
157. Santhanagopalan I, Basha E, Ballard KN, Bopp NE, Vierling E (2015) Model chaperones: small heat shock proteins from plants. In: Tanguay RM, Hightower LE (eds) *The big book on small heat shock proteins*. Springer International Publishing, Cham, pp 119–153. doi:10.1007/978-3-319-16077-1_5
158. Reyes JL, Campos F, Wei H, Arora R, Yang YI, Karlson DT, Covarrubias AA (2008) Functional dissection of hydrophilins during in vitro freeze protection. *Plant Cell Environ* 31(12):1781–1790. doi:10.1111/J.1365-3040.2008.01879.X
159. Tunnacliffe A, Wise MJ (2007) The continuing conundrum of the LEA proteins. *Naturwissenschaften* 94(10):791–812. doi:10.1007/S00114-007-0254-Y
160. Hughes SL, Schart V, Malcolmson J, Hogarth KA, Martynowicz DM, Tralman-Baker E, Patel SN, Graether SP (2013) The importance of size and disorder in the cryoprotective effects of dehydrins. *Plant Physiol* 163(3):1376–1386. doi:10.1104/pp.113.226803
161. Cuevas-Velazquez CL, Rendon-Luna DF, Covarrubias AA (2014) Dissecting the cryoprotection mechanisms for dehydrins. *Front Plant Sci* 5:583. doi:10.3389/fpls.2014.00583
162. Kazuoka T, Oeda K (1994) Purification and characterization of Cor85-oligomeric complex from cold-acclimated spinach. *Plant Cell Physiol* 35(4):601–611
163. Houde M, Daniel C, Lachapelle M, Allard F, Laliberte S, Sarhan F (1995) Immunolocalization of freezing-tolerance-associated proteins in the cytoplasm and nucleoplasm of wheat crown tissues. *Plant J* 8(4):583–593. doi:10.1046/J.1365-313x.1995.8040583.X
164. Bravo LA, Gallardo J, Navarrete A, Olave N, Martinez J, Alberdi M, Close TJ, Corcuera LJ (2003) Cryoprotective activity of a cold-induced dehydrin purified from barley. *Physiol Plant* 118(2):262–269. doi:10.1034/J.1399-3054.2003.00060.X
165. Nakayama K, Okawa K, Kakizaki T, Honma T, Itoh H, Inaba T (2007) *Arabidopsis* Cor15am is a chloroplast stromal protein that has cryoprotective activity and forms oligomers. *Plant Physiol* 144(1):513–523. doi:10.1104/pp.106.094581
166. Kovacs D, Kalmar E, Torok Z, Tompa P (2008) Chaperone activity of ERD10 and ERD14, two disordered stress-related plant proteins. *Plant Physiol*. doi:10.1104/pp.108.118208
167. Olvera-Carrillo Y, Luis Reyes J, Covarrubias AA (2011) Late embryogenesis abundant proteins: versatile players in the plant adaptation to water limiting environments. *Plant Signal Behav* 6(4):586–589. doi:10.4161/psb.6.4.15042
168. Kovacs D, Agoston B, Tompa P (2008) Disordered plant LEA proteins as molecular chaperones. *Plant Signal Behav*. doi:10.4161/psb.3.9.6434
169. Thalhammer A, Bryant G, Sulpice R, Hinch DK (2014) Disordered cold regulated15 proteins protect chloroplast membranes during freezing through binding and folding, but do not stabilize chloroplast enzymes in vivo. *Plant Physiol* 166(1):190–201. doi:10.1104/Pp.114.245399
170. Szalaine Agoston B, Kovacs D, Tompa P, Perczel A (2011) Full backbone assignment and dynamics of the intrinsically disordered dehydrin ERD14. *Biomol NMR Assign* 5(2):189–193. doi:10.1007/s12104-011-9297-2
171. Shih MD, Hsieh TY, Lin TP, Hsing YIC, Hoekstra FA (2010) Characterization of two soybean (*Glycine max* L.) LEA IV proteins by circular dichroism and Fourier transform infrared spectrometry. *Plant Cell Physiol* 51(3):395–407. doi:10.1093/Pcp/Pcq005
172. Ismail AM, Hall AE, Close TJ (1999) Purification and partial characterization of a dehydrin involved in chilling tolerance during seedling emergence of cowpea. *Plant Physiol* 120(1):237–244. doi:10.1104/Pp.120.1.237
173. Hara M, Terashima S, Kuboi T (2001) Characterization and cryoprotective activity of cold-responsive dehydrin from *Citrus unshiu*. *J Plant Physiol* 158(10):1333–1339. doi:10.1078/0176-1617-00600
174. Koag MC, Wilkens S, Fenton RD, Resnik J, Vo E, Close TJ (2009) The K-segment of maize DHN1 mediates binding to anionic phospholipid vesicles and concomitant structural changes. *Plant Physiol* 150(3):1503–1514. doi:10.1104/pp.109.136697
175. Koag MC, Fenton RD, Wilkens S, Close TJ (2003) The binding of maize DHN1 to lipid vesicles. Gain of structure and lipid specificity. *Plant Physiol* 131(1):309–316. doi:10.1104/Pp.011171
176. Soulages JL, Kim K, Arrese EL, Walters C, Cushman JC (2003) Conformation of a group 2 late embryogenesis abundant protein from soybean. Evidence of poly (L-proline)-type II structure. *Plant Physiol* 131(3):963–975. doi:10.1104/Pp.015891
177. Rahman LN, McKay F, Giuliani M, Quirk A, Moffatt BA, Harauz G, Dutcher JR (2013) Interactions of *Thellungiella salziginea* dehydrins TsDHN-1 and TsDHN-2 with membranes at cold and ambient temperatures-surface morphology and single-molecule force measurements show phase separation, and reveal tertiary and quaternary associations. *Biochim Biophys Acta* 1828(3):967–980. doi:10.1016/j.bbame.2012.11.031

178. Eriksson SK, Kutzer M, Procek J, Grobner G, Harryson P (2011) Tunable membrane binding of the intrinsically disordered dehydrin Lti30, a cold-induced plant stress protein. *Plant Cell* 23(6):2391–2404. doi:10.1105/TPC.111.085183
179. Clarke MW, Boddington KF, Warnica JM, Atkinson J, McKenna S, Madge J, Barker CH, Graether SP (2015) Structural and functional insights into the cryoprotection of membranes by the intrinsically disordered dehydrins. *J Biol Chem* 290(45):26900–26913. doi:10.1074/jbc.M115.678219
180. Eriksson S, Eremina N, Barth A, Danielsson J, Harryson P (2016) Membrane-induced folding of the plant stress dehydrin Lti30. *Plant Physiol* 171(2):932–943. doi:10.1104/pp.15.01531
181. Dure L (1993) Structural motifs in lea proteins. *Struct Motifs Lea Proteins* 10:91–103. doi:10.1104/pp.105.072967
182. Tolleter D, Jaquinod M, Mangavel C, Passirani C, Saulnier P, Manon S, Teyssier E, Payet N, Avelange-Macherel MH, Macherel D (2007) Structure and function of a mitochondrial late embryogenesis abundant protein are revealed by desiccation. *Plant Cell* 19(5):1580–1589. doi:10.1105/TPC.107.050104
183. Hundertmark M, Dimova R, Lengefeld J, Seckler R, Hinch DK (2011) The intrinsically disordered late embryogenesis abundant protein LEA18 from *Arabidopsis thaliana* modulates membrane stability through binding and folding. *BBA Biomembr* 1808(1):446–453. doi:10.1016/j.Bbamem.2010.09.010
184. Manfre AJ, Lanni LM, Marcotte WR (2006) The *Arabidopsis* group 1 LATE EMBRYOGENESIS ABUNDANT protein AtEM6 is required for normal seed development. *Plant Physiol* 140(1):140–149. doi:10.1104/PP.105.072967
185. Saavedra L, Svensson J, Carballo V, Izemendi D, Welin B, Vidal S (2006) A dehydrin gene in *Physcomitrella patens* is required for salt and osmotic stress tolerance. *Plant J* 45(2):237–249. doi:10.1111/J.1365-313X.2005.02603.X
186. Olvera-Carrillo Y, Campos F, Reyes JL, Garcarrubio A, Covarrubias AA (2010) Functional analysis of the group 4 late embryogenesis abundant proteins reveals their relevance in the adaptive response during water deficit in *Arabidopsis*. *Plant Physiol* 154(1):373–390. doi:10.1104/pp.110.158964
187. Cheng Z, Targolli J, Huang X, Wu R (2002) Wheat LEA genes, PMA80 and PMA1959, enhance dehydration tolerance of transgenic rice (*Oryza sativa* L.). *Mol Breed* 10(1):71–82. doi:10.1023/a:1020329401191
188. Figueras M, Pujal J, Saleh A, Save R, Pages M, Goday A (2004) Maize Rab17 overexpression in *Arabidopsis* plants promotes osmotic stress tolerance. *Ann Appl Biol* 144(3):251–257. doi:10.1111/j.1744-7348.2004.tb00341.x
189. Xu D, Duan X, Wang B, Hong B, Ho THD, Wu R (1996) Expression of a late embryogenesis abundant protein gene, HVA1, from barley confers tolerance to water deficit and salt stress in transgenic rice. *Plant Physiol* 110(1):249–257. doi:10.1104/pp.110.1.249
190. Park B-J, Liu Z, Kanno A, Kameya T (2005) Genetic improvement of Chinese cabbage for salt and drought tolerance by constitutive expression of a *B. napus* LEA gene. *Plant Sci* 169(3):553–558. doi:10.1016/j.plantsci.2005.05.008
191. Liu Y, Wang L, Xing X, Sun L, Pan J, Kong X, Zhang M, Li D (2013) ZmLEA3, a multifunctional group 3 LEA protein from maize (*Zea mays* L.), is involved in biotic and abiotic stresses. *Plant Cell Physiol* 54(6):944–959. doi:10.1093/pcp/pct047
192. Jha B, Lal S, Tiwari V, Yadav SK, Agarwal PK (2012) The SbASR-1 gene cloned from an extreme halophyte *Salicornia brachiata* enhances salt tolerance in transgenic tobacco. *Mar Biotechnol (N Y)* 14(6):782–792. doi:10.1007/s10126-012-9442-7
193. Hu W, Huang C, Deng X, Zhou S, Chen L, Li Y, Wang C, Ma Z, Yuan Q, Wang Y, Cai R, Liang X, Yang G, He G (2013) TaASR1, a transcription factor gene in wheat, confers drought stress tolerance in transgenic tobacco. *Plant Cell Environ* 36(8):1449–1464. doi:10.1111/pce.12074
194. Liu J, Jia C, Dong F, Wang J, Zhang J, Xu Y, Xu B, Jin Z (2013) Isolation of an abscisic acid senescence and ripening inducible gene from litchi and functional characterization under water stress. *Planta* 237(4):1025–1036. doi:10.1007/s00425-012-1820-x
195. Jeanneau M, Gerentes D, Foueillassar X, Zivy M, Vidal J, Toppan A, Perez P (2002) Improvement of drought tolerance in maize: towards the functional validation of the Zm-Asr1 gene and increase of water use efficiency by over-expressing C4-PEPC. *Biochimie* 84(11):1127–1135
196. Kalifa Y, Gilad A, Konrad Z, Zaccari M, Scolnik PA, Bar-Zvi D (2004) The water- and salt-stress-regulated Asr1 (abscisic acid stress ripening) gene encodes a zinc-dependent DNA-binding protein. *Biochem J* 381(Pt 2):373–378. doi:10.1042/bj20031800
197. Yang C-Y, Chen Y-C, Jauh GY, Wang C-S (2005) A lily ASR protein involves abscisic acid signaling and confers drought and salt resistance in *Arabidopsis*. *Plant Physiol* 139(2):836–846. doi:10.1104/pp.105.065458
198. Hara M, Shinoda Y, Kubo M, Kashima D, Takahashi I, Kato T, Horiike T, Kuboi T (2011) Biochemical characterization of the *Arabidopsis* KS-type dehydrin protein, whose gene expression is constitutively abundant rather than stress dependent. *Acta Physiol Plant* 33(6):2103–2116. doi:10.1007/s11738-011-0749-1
199. Graether SP, Boddington KF (2014) Disorder and function: a review of the dehydrin protein family. *Front Plant Sci* 5:576. doi:10.3389/fpls.2014.00576
200. Liu G, Xu H, Zhang L, Zheng Y (2011) Fe binding properties of two soybean (*Glycine max* L.) LEA4 proteins associated with antioxidant activity. *Plant Cell Physiol* 52(6):994–1002. doi:10.1093/pcp/pcr052
201. Plana M, Itarte E, Eritja R, Goday A, Pages M, Martinez MC (1991) Phosphorylation of maize RAB-17 protein by casein kinase 2. *J Biol Chem* 266(33):22510–22514
202. Heyen BJ, Alsheikh MK, Smith EA, Torvik CF, Seals DF, Randall SK (2002) The calcium-binding activity of a vacuole-associated, dehydrin-like protein is regulated by phosphorylation. *Plant Physiol* 130(2):675–687. doi:10.1104/pp.002550
203. Hara M, Fujinaga M, Kuboi T (2005) Metal binding by citrus dehydrin with histidine-rich domains. *J Exp Bot* 56(420):2695–2703. doi:10.1093/Jxb/Eri262
204. Hara M, Kondo M, Kato T (2013) A KS-type dehydrin and its related domains reduce Cu-promoted radical generation and the histidine residues contribute to the radical-reducing activities. *J Exp Bot* 64(6):1615–1624. doi:10.1093/jxb/ert016
205. Kruger C, Berkowitz O, Stephan UW, Hell R (2002) A metal-binding member of the late embryogenesis abundant protein family transports iron in the phloem of *Ricinus communis* L. *J Biol Chem* 277(28):25062–25069. doi:10.1074/Jbc.M201896200
206. Alsheikh MK, Heyen BJ, Randall SK (2003) Ion binding properties of the dehydrin ERD14 are dependent upon phosphorylation. *J Biol Chem* 278(42):40882–40889. doi:10.1074/jbc.M307151200
207. Alsheikh MK, Svensson JT, Randall SK (2005) Phosphorylation regulated ion-binding is a property shared by the acidic subclass dehydrins. *Plant Cell Environ* 28(9):1114–1122. doi:10.1111/J.1365-3040.2005.01348.X
208. Xu J, Zhang YX, Wei W, Han L, Guan ZQ, Wang Z, Chai TY (2008) BjDHNs confer heavy-metal tolerance in plants. *Mol Biotechnol* 38(2):91–98. doi:10.1007/s12033-007-9005-8
209. Hara M, Shinoda Y, Tanaka Y, Kuboi T (2009) DNA binding of citrus dehydrin promoted by zinc ion. *Plant Cell Environ* 32(5):532–541. doi:10.1111/J.1365-3040.2009.01947.X

210. Sasaki K, Christov NK, Tsuda S, Imai R (2014) Identification of a novel LEA protein involved in freezing tolerance in wheat. *Plant Cell Physiol* 55(1):136–147. doi:10.1093/pcp/pct164
211. Rosales R, Romero I, Escribano MI, Merodio C, Sanchez-Ballesta MT (2014) The crucial role of Φ - and K-segments in the in vitro functionality of *Vitis vinifera* dehydrin DHN1a. *Phytochemistry* 108:17–25. doi:10.1016/j.phytochem.2014.10.006
212. Ricardi MM, Gonzalez RM, Zhong S, Dominguez PG, Duffy T, Turjanski PG, Salgado Salter JD, Alleva K, Carrari F, Giovanoni JJ, Estevez JM, Iusem ND (2014) Genome-wide data (ChIP-seq) enabled identification of cell wall-related and aquaporin genes as targets of tomato ASR1, a drought stress-responsive transcription factor. *BMC Plant Biol* 14:29. doi:10.1186/1471-2229-14-29
213. Hayes MA, Davies C, Dry IB (2007) Isolation, functional characterization, and expression analysis of grapevine (*Vitis vinifera* L.) hexose transporters: differential roles in sink and sourcettissues. *J Exp Bot* 58(8):1985–1997. doi:10.1093/jxb/erm061
214. Virlouvet L, Jacquemot MP, Gerentes D, Corti H, Bouton S, Gilard F, Valot B, Trouverie J, Tcherkez G, Falque M, Damerval C, Rogowsky P, Perez P, Noctor G, Zivy M, Coursol S (2011) The ZmASR1 protein influences branched-chain amino acid biosynthesis and maintains kernel yield in maize under water-limited conditions. *Plant Physiol* 157(2):917–936. doi:10.1104/pp.111.176818
215. Frankel N, Nunes-Nesi A, Balbo I, Mazuch J, Centeno D, Iusem ND, Fernie AR, Carrari F (2007) ci21A/Asr1 expression influences glucose accumulation in potato tubers. *Plant Mol Biol* 63(5):719–730. doi:10.1007/s11103-006-9120-0
216. Brini F, Hanin M, Lumbreras V, Irar S, Pages M, Masmoudi K (2007) Functional characterization of DHN-5, a dehydrin showing a differential phosphorylation pattern in two Tunisian durum wheat (*Triticum durum* Desf.) varieties with marked differences in salt and drought tolerance. *Plant Sci* 172(1):20–28. doi:10.1016/j.plantsci.2006.07.011
217. Bonhomme L, Valot B, Tardieu F, Zivy M (2012) Phosphoproteome dynamics upon changes in plant water status reveal early events associated with rapid growth adjustment in maize leaves. *Mol Cell Proteomics* 11(10):957–972. doi:10.1074/mcp.M111.015867
218. Mayank P, Grossman J, Wuest S, Boisson-Dernier A, Roschitzki B, Nanni P, Nuhse T, Grossniklaus U (2012) Characterization of the phosphoproteome of mature *Arabidopsis* pollen. *Plant J* 72(1):89–101. doi:10.1111/j.1365-313X.2012.05061.x
219. Yao Q, Bollinger C, Gao J, Xu D, Thelen JJ (2012) P(3)DB: an integrated database for plant protein phosphorylation. *Front Plant Sci* 3:206. doi:10.3389/fpls.2012.00206
220. Jiang X, Wang Y (2004) Beta-elimination coupled with tandem mass spectrometry for the identification of in vivo and in vitro phosphorylation sites in maize dehydrin DHN1 protein. *Biochemistry* 43(49):15567–15576. doi:10.1021/bi0483965
221. Röhrig H, Schmidt J, Colby T, Brautigam A, Hufnagel P, Bartels D (2006) Desiccation of the resurrection plant *Craterostigma plantagineum* induces dynamic changes in protein phosphorylation. *Plant Cell Environ* 29(8):1606–1617. doi:10.1111/j.1365-3040.2006.01537.x
222. Gao J, Xu D (2012) Correlation between posttranslational modification and intrinsic disorder in protein. *Pac Symp Biocomput*. doi:10.1142/9789814366496_0010
223. Riera M, Figueras M, López C, Goday A, Pagès M (2004) Protein kinase CK2 modulates developmental functions of the abscisic acid responsive protein Rab17 from maize. *Proc Natl Acad Sci USA* 101(26):9879–9884. doi:10.1073/pnas.0306154101
224. Goday A, Jensen AB, Culiñez-Macia FA, Mar Alba M, Figueras M, Serratos J, Torrent M, Pages M (1994) The maize abscisic acid-responsive protein Rab17 is located in the nucleus and interacts with nuclear localization signals. *Plant cell* 6(3):351–360. doi:10.1105/tpc.6.3.351
225. Hara M (2010) The multifunctionality of dehydrins: an overview. *Plant Signal Behav* 5(5):503–508. doi:10.4161/psb.11085
226. Candat A, Paszkiewicz G, Neveu M, Gautier R, Logan DC, Avelange-Macherel MH, Macherel D (2014) The ubiquitous distribution of late embryogenesis abundant proteins across cell compartments in *Arabidopsis* offers tailored protection against abiotic stress. *Plant Cell* 26(7):3148–3166. doi:10.1105/tpc.114.127316
227. Dominguez PG, Carrari F (2015) ASR1 transcription factor and its role in metabolism. *Plant Signal Behav* 10(4):e992751. doi:10.4161/15592324.2014.992751
228. Gao J, Lan T (2016) Functional characterization of the late embryogenesis abundant (LEA) protein gene family from *Pinus tabulaeformis* (Pinaceae) in *Escherichia coli*. *Sci Rep* 6:19467. doi:10.1038/srep19467
229. Tompa P, Szasz C, Buday L (2005) Structural disorder throws new light on moonlighting. *Trends Biochem Sci* 30(9):484–489. doi:10.1016/j.tibs.2005.07.008
230. Basha E, O'Neill H, Vierling E (2012) Small heat shock proteins and alpha-crystallins: dynamic proteins with flexible functions. *Trends Biochem Sci* 37(3):106–117. doi:10.1016/j.tibs.2011.11.005
231. Waters ER (2013) The evolution, function, structure, and expression of the plant sHSPs. *J Exp Bot* 64(2):391–403. doi:10.1093/jxb/ers355
232. Jaya N, Garcia V, Vierling E (2009) Substrate binding site flexibility of the small heat shock protein molecular chaperones. *Proc Natl Acad Sci USA* 106(37):15604–15609. doi:10.1073/pnas.0902177106
233. Graham D, Patterson BD (1982) Responses of plants to low, nonfreezing temperatures: proteins, metabolism, and acclimation. *Annu Rev Plant Physiol* 33(1):347–372. doi:10.1146/annurev.pp.33.060182.002023
234. Mangeon A, Junqueira RM, Sachetto-Martins G (2010) Functional diversity of the plant glycine-rich proteins superfamily. *Plant Signal Behav* 5(2):99–104. doi:10.4161/psb.5.2.10336
235. Khan F, Sultana T, Deeba F, Naqvi S (2013) Dynamics of mRNA of glycine-rich RNA-binding protein during wounding, cold and salt stresses in *Nicotiana tabacum*. *Pak J Bot* 45:297–300
236. Kim JY, Kim WY, Kwak KJ, Oh SH, Han YS, Kang H (2010) Glycine-rich RNA-binding proteins are functionally conserved in *Arabidopsis thaliana* and *Oryza sativa* during cold adaptation process. *J Exp Bot* 61(9):2317–2325. doi:10.1093/jxb/erq058
237. Kwak KJ, Park SJ, Han JH, Kim MK, Oh SH, Han YS, Kang H (2011) Structural determinants crucial to the RNA chaperone activity of glycine-rich RNA-binding proteins 4 and 7 in *Arabidopsis thaliana* during the cold adaptation process. *J Exp Bot* 62(11):4003–4011. doi:10.1093/jxb/err101
238. Freire MA (2012) The *Zea mays* glycine-rich RNA-binding protein MA16 is bound to a ribonucleotide(s) by a stable linkage. *J Plant Res* 125(5):653–660. doi:10.1007/s10265-012-0476-8
239. Lorkovic ZJ (2009) Role of plant RNA-binding proteins in development, stress response and genome organization. *Trends Plant Sci* 14(4):229–236. doi:10.1016/j.tplants.2009.01.007
240. Heintzen C, Nater M, Apel K, Staiger D (1997) AtGRP7, a nuclear RNA-binding protein as a component of a circadian-regulated negative feedback loop in *Arabidopsis thaliana*. *Proc Natl Acad Sci* 94(16):8515–8520. doi:10.1073/pnas.94.16.8515
241. Staiger D, Zecca L, Wiczeorek Kirk DA, Apel K, Eckstein L (2003) The circadian clock regulated RNA-binding protein AtGRP7 autoregulates its expression by influencing alternative

- splicing of its own pre-mRNA. *Plant J* 33(2):361–371. doi:[10.1046/j.1365-313X.2003.01629.x](https://doi.org/10.1046/j.1365-313X.2003.01629.x)
242. Streitner C, Danisman S, Wehrle F, Schoning JC, Alfano JR, Staiger D (2008) The small glycine-rich RNA binding protein AtGRP7 promotes floral transition in *Arabidopsis thaliana*. *Plant J* 56(2):239–250. doi:[10.1111/j.1365-313X.2008.03591.x](https://doi.org/10.1111/j.1365-313X.2008.03591.x)
 243. Kim JS, Park SJ, Kwak KJ, Kim YO, Kim JY, Song J, Jang B, Jung CH, Kang H (2007) Cold shock domain proteins and glycine-rich RNA-binding proteins from *Arabidopsis thaliana* can promote the cold adaptation process in *Escherichia coli*. *Nucleic Acids Res* 35(2):506–516. doi:[10.1093/nar/gkl1076](https://doi.org/10.1093/nar/gkl1076)
 244. Streitner C, Koster T, Simpson CG, Shaw P, Danisman S, Brown JW, Staiger D (2012) An hnRNP-like RNA-binding protein affects alternative splicing by in vivo interaction with transcripts in *Arabidopsis thaliana*. *Nucleic Acids Res* 40(22):11240–11255. doi:[10.1093/nar/gks873](https://doi.org/10.1093/nar/gks873)
 245. Kim JS, Jung HJ, Lee HJ, Kim KA, Goh CH, Woo Y, Oh SH, Han YS, Kang H (2008) Glycine-rich RNA-binding protein 7 affects abiotic stress responses by regulating stomata opening and closing in *Arabidopsis thaliana*. *Plant J* 55(3):455–466. doi:[10.1111/j.1365-313X.2008.03518.x](https://doi.org/10.1111/j.1365-313X.2008.03518.x)
 246. Khan F, Daniels MA, Folkers GE, Boelens R, Saqlan Naqvi SM, van Ingen H (2014) Structural basis of nucleic acid binding by *Nicotiana tabacum* glycine-rich RNA-binding protein: implications for its RNA chaperone function. *Nucleic Acids Res* 42(13):8705–8718. doi:[10.1093/nar/gku468](https://doi.org/10.1093/nar/gku468)
 247. McCain DC, Croxdale J, Markley JL (1989) Thermal damage to chloroplast envelope membranes. *Plant Physiol* 90(2):606–609. doi:[10.1104/pp.90.2.606](https://doi.org/10.1104/pp.90.2.606)
 248. Dekov I, Tsonev T, Yordanov I (2000) Effects of water stress and high-temperature stress on the structure and activity of photosynthetic apparatus of *Zea mays* and *Helianthus annuus*. *Photosynthetica* 38(3):361–366. doi:[10.1023/a:1010961218145](https://doi.org/10.1023/a:1010961218145)
 249. Vothknecht UC, Otters S, Hennig R, Schneider D (2012) Vipp1: a very important protein in plastids? *J Exp Bot* 63(4):1699–1712. doi:[10.1093/jxb/err357](https://doi.org/10.1093/jxb/err357)
 250. Zhang L, Sakamoto W (2013) Possible function of VIPP1 in thylakoids: protection but not formation? *Plant Signal Behav* 8(2):e22860. doi:[10.4161/psb.22860](https://doi.org/10.4161/psb.22860)
 251. Zhang L, Sakamoto W (2015) Possible function of VIPP1 in maintaining chloroplast membranes. *Biochim Biophys Acta* 1847(9):831–837. doi:[10.1016/j.bbabi.2015.02.013](https://doi.org/10.1016/j.bbabi.2015.02.013)
 252. Zhang L, Kato Y, Otters S, Vothknecht UC, Sakamoto W (2012) Essential role of VIPP1 in chloroplast envelope maintenance in *Arabidopsis*. *Plant Cell* 24(9):3695–3707. doi:[10.1105/tpc.112.103606](https://doi.org/10.1105/tpc.112.103606)
 253. Kroll D, Meierhoff K, Bechtold N, Kinoshita M, Westphal S, Vothknecht UC, Soll J, Westhoff P (2001) VIPP1, a nuclear gene of *Arabidopsis thaliana* essential for thylakoid membrane formation. *Proc Natl Acad Sci USA* 98(7):4238–4242. doi:[10.1073/pnas.061500998](https://doi.org/10.1073/pnas.061500998)
 254. Flores-Kim J, Darwin AJ (2016) The phage shock protein response. *Annu Rev Microbiol* 70:83–101. doi:[10.1146/annurev-micro-102215-095359](https://doi.org/10.1146/annurev-micro-102215-095359)
 255. Hankamer BD, Elderkin SL, Buck M, Nield J (2004) Organization of the AAA+ adaptor protein PspA is an oligomeric ring. *J Biol Chem* 279(10):8862–8866. doi:[10.1074/jbc.M307889200](https://doi.org/10.1074/jbc.M307889200)
 256. Lenn T, Gkekas CN, Bernard L, Engl C, Jovanovic G, Buck M, Ying L (2011) Measuring the stoichiometry of functional PspA complexes in living bacterial cells by single molecule photobleaching. *Chem Commun* 47(1):400–402. doi:[10.1039/C0CC01707H](https://doi.org/10.1039/C0CC01707H)
 257. Drin G, Casella J-F, Gautier R, Boehmer T, Schwartz TU, Antony B (2007) A general amphipathic [alpha]-helical motif for sensing membrane curvature. *Nat Struct Mol Biol* 14(2):138–146. doi:[10.1038/nsmb1194](https://doi.org/10.1038/nsmb1194)
 258. Drin G, Antony B (2010) Amphipathic helices and membrane curvature. *FEBS Lett* 584(9):1840–1847. doi:[10.1016/j.febslet.2009.10.022](https://doi.org/10.1016/j.febslet.2009.10.022)
 259. McDonald C, Jovanovic G, Wallace BA, Ces O, Buck M (2016) Structure and function of PspA and Vipp1 N-terminal peptides: insights into the membrane stress sensing and mitigation. *Biochim Biophys Acta* 1859(1):28–39. doi:[10.1016/j.bbamem.2016.10.018](https://doi.org/10.1016/j.bbamem.2016.10.018)
 260. Zhang L, Kondo H, Kamikubo H (2016) VIPP1 has a disordered C-terminal tail necessary for protecting photosynthetic membranes against stress. *Plant Physiol* 171(3):1983–1995. doi:[10.1104/pp.16.00532](https://doi.org/10.1104/pp.16.00532)
 261. Sato H, Mizoi J, Tanaka H, Maruyama K, Qin F, Osakabe Y, Morimoto K, Ohori T, Kusakabe K, Nagata M, Shinozaki K, Yamaguchi-Shinozaki K (2014) *Arabidopsis* DPB3-1, a DREB2A interactor, specifically enhances heat stress-induced gene expression by forming a heat stress-specific transcriptional complex with NF-Y subunits. *Plant Cell* 26(12):4954–4973. doi:[10.1105/tpc.114.132928](https://doi.org/10.1105/tpc.114.132928)
 262. Blomberg J, Aguilar X, Brännström K, Rautio L, Olofsson A, Wittung-Stafshede P, Björklund S (2012) Interactions between DNA, transcriptional regulator Dreb2a and the Med25 mediator subunit from *Arabidopsis thaliana* involve conformational changes. *Nucleic Acids Res* 40(13):5938–5950. doi:[10.1093/nar/gks265](https://doi.org/10.1093/nar/gks265)
 263. Vainonen JP, Jaspers P, Wrzaczek M, Lamminmaki A, Reddy RA, Vaahtera L, Brosche M, Kangasjarvi J (2012) RCD1-DREB2A interaction in leaf senescence and stress responses in *Arabidopsis thaliana*. *Biochem J* 442:573–581. doi:[10.1042/BJ20111739](https://doi.org/10.1042/BJ20111739)
 264. O’Shea C, Staby L, Bendsen SK, Tidemand FG, Redsted A, Willemoës M, Kragelund BB, Skriver K (2016) Structures and short linear motif of disordered transcription factor regions provide clues to the interactome of the cellular hub radical-induced cell death1. *J Biol Chem*. doi:[10.1074/jbc.M116.753426](https://doi.org/10.1074/jbc.M116.753426)
 265. Che P, Bussell JD, Zhou W, Estavillo GM, Pogson BJ, Smith SM (2010) Signaling from the endoplasmic reticulum activates brassinosteroid signaling and promotes acclimation to stress in *Arabidopsis*. *Sci Signal* 3(141):ra69. doi:[10.1126/scisignal.2001140](https://doi.org/10.1126/scisignal.2001140)
 266. Gao H, Brandizzi F, Benning C, Larkin RM (2008) A membrane-tethered transcription factor defines a branch of the heat stress response in *Arabidopsis thaliana*. *Proc Natl Acad Sci USA* 105(42):16398–16403. doi:[10.1073/pnas.0808463105](https://doi.org/10.1073/pnas.0808463105)
 267. Urade R (2009) The endoplasmic reticulum stress signaling pathways in plants. *BioFactors* 35(4):326–331. doi:[10.1002/biof.45](https://doi.org/10.1002/biof.45)
 268. Howell SH (2013) Endoplasmic reticulum stress responses in plants. *Annu Rev Plant Biol* 64:477–499. doi:[10.1146/annurev-arplant-050312-120053](https://doi.org/10.1146/annurev-arplant-050312-120053)
 269. Deng Y, Humbert S, Liu JX, Srivastava R, Rothstein SJ, Howell SH (2011) Heat induces the splicing by IRE1 of a mRNA encoding a transcription factor involved in the unfolded protein response in *Arabidopsis*. *Proc Natl Acad Sci USA* 108(17):7247–7252. doi:[10.1073/pnas.1102117108](https://doi.org/10.1073/pnas.1102117108)
 270. Srivastava R, Deng Y, Shah S, Rao AG, Howell SH (2013) BINDING PROTEIN is a master regulator of the endoplasmic reticulum stress sensor/transducer bZIP28 in *Arabidopsis*. *Plant Cell* 25(4):1416–1429. doi:[10.1105/tpc.113.110684](https://doi.org/10.1105/tpc.113.110684)
 271. Srivastava R, Deng Y, Howell SH (2014) Stress sensing in plants by an ER stress sensor/transducer, bZIP28. *Front Plant Sci* 5:59. doi:[10.3389/fpls.2014.00059](https://doi.org/10.3389/fpls.2014.00059)
 272. Dangel JL, Jones JD (2001) Plant pathogens and integrated defence responses to infection. *Nature* 411(6839):826–833. doi:[10.1038/35081161](https://doi.org/10.1038/35081161)

273. Delteil A, Gobbato E, Cayrol B, Estevan J, Michel-Romiti C, Dievart A, Kroj T, Morel JB (2016) Several wall-associated kinases participate positively and negatively in basal defense against rice blast fungus. *BMC Plant Biol* 16:17. doi:[10.1186/s12870-016-0711-x](https://doi.org/10.1186/s12870-016-0711-x)
274. Rasmussen MW, Roux M, Petersen M, Mundy J (2012) MAP kinase cascades in *Arabidopsis* innate immunity. *Front Plant Sci* 3:169. doi:[10.3389/fpls.2012.00169](https://doi.org/10.3389/fpls.2012.00169)
275. Shiu SH, Bleecker AB (2001) Receptor-like kinases from *Arabidopsis* form a monophyletic gene family related to animal receptor kinases. *Proc Natl Acad Sci USA* 98(19):10763–10768. doi:[10.1073/Pnas.181141598](https://doi.org/10.1073/Pnas.181141598)
276. Kemmerling B, Schwedt A, Rodriguez P, Mazzotta S, Frank M, Qamar SA, Mengiste T, Betsuyaku S, Parker JE, Mussig C, Thomma BP, Albrecht C, de Vries SC, Hirt H, Nurnberger T (2007) The BRI1-associated kinase 1, BAK1, has a brassinolide-independent role in plant cell-death control. *Curr Biol* 17(13):1116–1122. doi:[10.1016/j.cub.2007.05.046](https://doi.org/10.1016/j.cub.2007.05.046)
277. Ogasawara Y, Kaya H, Hiraoka G, Yumoto F, Kimura S, Kadota Y, Hishinuma H, Senzaki E, Yamagoe S, Nagata K, Nara M, Suzuki K, Tanokura M, Kuchitsu K (2008) Synergistic activation of the *Arabidopsis* NADPH oxidase AtrbohD by Ca²⁺ and phosphorylation. *J Biol Chem* 283(14):8885–8892. doi:[10.1074/jbc.M708106200](https://doi.org/10.1074/jbc.M708106200)
278. Oda T, Hashimoto H, Kuwabara N, Akashi S, Hayashi K, Kojima C, Wong HL, Kawasaki T, Shimamoto K, Sato M, Shimizu T (2010) Structure of the N-terminal regulatory domain of a plant NADPH oxidase and its functional implications. *J Biol Chem* 285(2):1435–1445. doi:[10.1074/jbc.M109.058909](https://doi.org/10.1074/jbc.M109.058909)
279. Sangster TA, Queitsch C (2005) The HSP90 chaperone complex, an emerging force in plant development and phenotypic plasticity. *Curr Opin Plant Biol* 8(1):86–92. doi:[10.1016/j.pbi.2004.11.012](https://doi.org/10.1016/j.pbi.2004.11.012)
280. Shirasu K, Schulze-Lefert P (2003) Complex formation, promiscuity and multi-functionality: protein interactions in disease-resistance pathways. *Trends Plant Sci* 8(6):252–258. doi:[10.1016/S1360-1385\(03\)00104-3](https://doi.org/10.1016/S1360-1385(03)00104-3)
281. Zhang MH, Kadota Y, Prodromou C, Shirasu K, Pearl LH (2010) Structural basis for assembly of Hsp90-Sgt1-CHORD protein complexes: implications for chaperoning of NLR innate immunity receptors. *Mol Cell* 39(2):269–281. doi:[10.1016/j.molcel.2010.05.010](https://doi.org/10.1016/j.molcel.2010.05.010)
282. Kong Q, Qu N, Gao MH, Zhang ZB, Ding XJ, Yang F, Li YZ, Dong OX, Chen S, Li X, Zhang YL (2012) The MEKK1-MKK1/MKK2-MPK4 kinase cascade negatively regulates immunity mediated by a mitogen-activated protein kinase kinase in *Arabidopsis*. *Plant Cell* 24(5):2225–2236. doi:[10.1105/tpc.112.097253](https://doi.org/10.1105/tpc.112.097253)
283. Jones JDG, Dangl JL (2006) The plant immune system. *Nature* 444(7117):323–329. doi:[10.1038/nature05286](https://doi.org/10.1038/nature05286)
284. Segrest JP, De Loof H, Dohlman JG, Brouillette CG, Anantharamaiah GM (1990) Amphipathic helix motif: classes and properties. *Protein Struct Funct Bioinform* 8(2):103–117. doi:[10.1002/prot.340080202](https://doi.org/10.1002/prot.340080202)
285. Ellis J (1987) Proteins as molecular chaperones. *Nature* 328(6129):378–379. doi:[10.1038/328378a0](https://doi.org/10.1038/328378a0)
286. Hendrick JP, Hartl FU (1995) The role of molecular chaperones in protein folding. *FASEB J* 9(15):1559–1569
287. Schunemann D (2004) Structure and function of the chloroplast signal recognition particle. *Curr Genet* 44(6):295–304. doi:[10.1007/s00294-003-0450-z](https://doi.org/10.1007/s00294-003-0450-z)
288. Rosenblad MA, Samuelsson T (2004) Identification of chloroplast signal recognition particle RNA genes. *Plant Cell Physiol* 45(11):1633–1639. doi:[10.1093/pcp/pch185](https://doi.org/10.1093/pcp/pch185)
289. Weikl TR, Paul F (2014) Conformational selection in protein binding and function. *Protein Sci* 23(11):1508–1518. doi:[10.1002/pro.2539](https://doi.org/10.1002/pro.2539)
290. Csermely P, Palotai R, Nussinov R (2010) Induced fit, conformational selection and independent dynamic segments: an extended view of binding events. *Trends Biochem Sci* 35(10):539–546. doi:[10.1016/j.tibs.2010.04.009](https://doi.org/10.1016/j.tibs.2010.04.009)
291. Cushman JC (2001) Crassulacean acid metabolism. A plastic photosynthetic adaptation to arid environments. *Plant Physiol* 127(4):1439–1448. doi:[10.1104/pp.010818](https://doi.org/10.1104/pp.010818)
292. Wright PE, Dyson HJ (1999) Intrinsically unstructured proteins: re-assessing the protein structure-function paradigm. *J Mol Biol* 293(2):321–331. doi:[10.1006/Jmbi.1999.3110](https://doi.org/10.1006/Jmbi.1999.3110)
293. Wright PE, Dyson HJ (2009) Linking folding and binding. *Curr Opin Struct Biol* 19(1):31–38. doi:[10.1016/j.sbi.2008.12.003](https://doi.org/10.1016/j.sbi.2008.12.003)
294. Tompa P, Fuxreiter M (2008) Fuzzy complexes: polymorphism and structural disorder in protein–protein interactions. *Trends Biochem Sci* 33(1):2–8. doi:[10.1016/J.Tibs.2007.10.003](https://doi.org/10.1016/J.Tibs.2007.10.003)
295. Janson J-C (2011) Protein purification: principles, high resolution methods, and applications, 3rd edn. Wiley, New York. doi:[10.1002/9780470939932](https://doi.org/10.1002/9780470939932)
296. Oldfield CJ, Cheng YG, Cortese MS, Romero P, Uversky VN, Dunker AK (2005) Coupled folding and binding with alpha-helix-forming molecular recognition elements. *Biochemistry* 44(37):12454–12470. doi:[10.1021/Bi050736e](https://doi.org/10.1021/Bi050736e)
297. Roberts EH (1973) Predicting the storage life of seeds. *Seed Sci Technol* 1:499–514
298. McDonald C, Jovanovic G, Ces O, Buck M (2015) Membrane stored curvature elastic stress modulates recruitment of maintenance proteins PspA and Vippl1. *MBio* 6(5):e01188–e01467. doi:[10.1128/mBio.01188-15](https://doi.org/10.1128/mBio.01188-15)
299. Vogt L, Vinyard DJ, Khan S, Brudvig GW (2015) Oxygen-evolving complex of Photosystem II: an analysis of second-shell residues and hydrogen-bonding networks. *Curr Opin Chem Biol* 25:152–158. doi:[10.1016/j.cbpa.2014.12.040](https://doi.org/10.1016/j.cbpa.2014.12.040)
300. Wang M, Kaufman RJ (2014) The impact of the endoplasmic reticulum protein-folding environment on cancer development. *Nat Rev Cancer* 14(9):581–597. doi:[10.1038/nrc3800](https://doi.org/10.1038/nrc3800)