



# Positive and Generalized Positive Real Lemma for Slice Hyperholomorphic Functions

Daniel Alpay, Fabrizio Colombo, Izchak Lewkowicz  
and Irene Sabadini 

**Abstract.** In this paper we prove a quaternionic positive real lemma as well as its generalized version, in case the associated kernel has negative squares for slice hyperholomorphic functions. We consider the case of functions with positive real part in the half space of quaternions with positive real part, as well as the case of (generalized) Schur functions in the open unit ball.

**Mathematics Subject Classification.** Primary 30C40, Secondary 30G35, 93B15.

**Keywords.** Positive real functions, Kalman–Yakubovich–Popov lemma, Slice hyperholomorphic functions, Unit ball, Half space, Negative squares.

## 1. Introduction

Scalar rational function which analytically map the open right half of the complex plane to its closure are called in system and control engineering positive real and we shall denote them by  $\mathcal{PR}$ . More generally, matrix-valued functions analytic in the open unit disk and having a positive real part there play an important role in engineering, network theory and operator theory. They can be characterized in a number of ways, and we mention in particular Herglotz integral representations. They admit numerous generalizations, for instance to quaternions (see e.g. [1, 6]) and to the case of negative squares. The latter was initiated and studied in much details by Krein and Langer; see e.g. [34, 35].

The scalar rational case is of special interest. Indeed, for nearly a century it has been recognized that the driving point impedance of an electrical

---

Daniel Alpay thanks the Foster G. and Mary McGaw Professorship in Mathematical Sciences, which supported this research.

circuit comprised of resistors, inductors and capacitors is a  $\mathcal{PR}$  function and conversely every  $\mathcal{PR}$  function may be realized as a driving point impedance of such a circuit; see [20, 21]. A classical electro-mechanical duality reveals that mechanical systems comprised of mass-damper-spring (strictly speaking, mass should be replaced by inerter; see [38] for the latter) are described by  $\mathcal{PR}$  functions. Thus positive real functions are perceived as a mathematical model of linear passive systems. The positive real lemma (also called Kalman–Yakubovich–Popov lemma) offers an easy-to-check characterization of matrix-valued  $\mathcal{PR}$  functions with no pole at infinity. There are a few hundreds of papers on this subject. The purpose of this work is to introduce an extension of the positive real lemma in the setting of quaternionic analysis and rational slice hyperholomorphic functions. To set the framework, we first discuss the complex setting, and recall that an important question in the theory of matrix-valued rational functions is to relate metric properties of the function to those of one of its minimal realization. The case of functions taking unitary values on the boundary is considered in [14, 32]. The positive real lemma characterizes minimal realizations of functions having a positive real part in a half-plane and reads (see [16, 24, 29, 30]):

**Theorem 1.1.** *Let  $R$  be a  $\mathbb{C}^{n \times n}$ -valued rational function, analytic at infinity, and with minimal realization*

$$R(z) = D + C(zI_N - A)^{-1}B, \tag{1.1}$$

with  $D = R(\infty) \in \mathbb{C}^{n \times n}$  and  $(C, A, B) \in \mathbb{C}^{n \times N} \times \mathbb{C}^{N \times N} \times \mathbb{C}^{N \times n}$ . Then  $R$  has a real positive real part in the open right half-plane if and only if there exists a negative definite matrix  $H \in \mathbb{C}^{N \times N}$  satisfying

$$\begin{pmatrix} H & 0 \\ 0 & I_n \end{pmatrix} \begin{pmatrix} A & B \\ C & D \end{pmatrix} + \begin{pmatrix} A & B \\ C & D \end{pmatrix}^* \begin{pmatrix} H & 0 \\ 0 & I_n \end{pmatrix} \geq 0. \tag{1.2}$$

*Remark 1.2.* The matrix  $H$  is not unique and the set of all matrices  $H$  satisfying (1.2) is convex. Assuming  $D + D^*$  positive definite, the Schur complement formula shows that (1.2) holds if and only if  $H$  satisfies the Riccati inequality

$$HA + A^*H - (HB + C^*)(D + D^*)^{-1}(C + B^*H) \leq 0, \tag{1.3}$$

see [15, Propositions 3.1 and 3.2], and it can be proved (see [29]) that this set has a minimal and a maximal solution. The minimal solution correspond to the spectral factorization. To see the link with factorizations, write the left side of (1.2) as

$$\begin{pmatrix} L \\ M \end{pmatrix} (L^* \quad M^*),$$

with  $L \in \mathbb{C}^{N \times \ell}$  and  $M \in \mathbb{C}^{n \times \ell}$ , and where  $\ell$  is equal to the rank of the left hand side of (1.2).

We have for  $z \in \mathbb{C}$  (see [29, p. 26])

$$R(z) + R(-\bar{z})^* = (M + C(zI - A)^{-1}H^{-1}L)(M + C(-\bar{z}I - A)^{-1}H^{-1}L)^* \tag{1.4}$$

and so, for  $x \in \mathbb{R}$

$$R(x) + R(-x)^* = (M + C(xI - A)^{-1}H^{-1}L)(M + C(-xI - A)^{-1}H^{-1}L)^*. \tag{1.5}$$

Note that (1.5) (and not (1.4) because of the noncommutativity) makes sense in the quaternionic setting, and then can be extended from  $x \in \mathbb{R}$  to general quaternions  $q \in \mathbb{H}$  using the  $\star$ -product see formulas (4.13) and (4.11) and next section.

*Remark 1.3.* Theorem 1.1 has been extended in [24] to the case of rational functions which are only positive on the imaginary axis. The matrix  $H$  in (1.2) is then only Hermitian non-singular.

Closely related to the above right-half plane Kalman–Yakubovich–Popov lemma in (1.2) (and its generalization to generalized positive functions) is the analogous result for rational matrix-valued functions analytic in the open unit disk and contractive on the unit circle (i.e. Schur functions), or possibly having a (finite) number of poles in the open unit disk (i.e. generalized Schur functions). See [24, (16), p. 604] for the following theorem.

**Theorem 1.4.** *Let  $S$  be a  $\mathbb{C}^{n \times m}$ -valued rational function, analytic at infinity, with minimal realization  $S(z) = D + C(zI_N - A)^{-1}B$ . Then,  $S$  takes contractive values on the unit circle if and only if there is an invertible Hermitian matrix  $H$  such that*

$$\begin{pmatrix} H & 0 \\ 0 & I_n \end{pmatrix} - \begin{pmatrix} A & B \\ C & D \end{pmatrix}^* \begin{pmatrix} H & 0 \\ 0 & I_n \end{pmatrix} \begin{pmatrix} A & B \\ C & D \end{pmatrix} \geq 0. \tag{1.6}$$

Furthermore,  $S$  is a Schur function if and only if  $H$  is negative.

Generalized Schur functions were characterized in [34] as quotient of Schur functions, the denominator being a finite matrix Blaschke product. This result allows to reduce the study of realizations of generalized Schur functions to the study of Schur functions. They are been studied in a number of works; see e.g. [13, 17, 19, 33], [39, Section II].

When leaving the setting of the complex numbers and going to quaternionic slice hyperholomorphic functions, a number of questions have to be addressed, the first one being what are the objects to be studied, that is, what are now Schur and generalized Schur functions, and positive real and generalized positive real Schur functions. Since the spectral theorem holds for Hermitian matrices with quaternionic entries, one can defined positive definite kernels, and kernels having a finite number of negative squares; see [9] and Definition 5.1 below. We will define the classes of functions to be studied in terms of the number of negative squares of an associated kernel; in the complex setting the two definitions are equivalent, but this is not the case anymore in the quaternionic setting. The following definitions first appeared in [7, 8]. In the statement, and in this paper in general,  $\mathbb{H}$  denotes the skew-field of the quaternions.

**Definition 1.5.** A  $\mathbb{H}^{n \times n}$ -valued function  $\phi$ , rational slice hyperholomorphic in a neighborhood  $\Omega$  of the origin in  $\mathbb{H}$  is called positive (resp. generalized positive) if the kernel

$$K_\phi(p, q) = (\phi(p) + \phi(q)^*) \star (p + \bar{q})^{-\star} \tag{1.7}$$

is positive definite (resp. has a finite number of negative squares) in  $\Omega$ .

**Definition 1.6.** A  $\mathbb{H}^{n \times m}$ -valued function  $S$ , rational slice hyperholomorphic in a neighborhood  $\Omega$  of the origin in  $\mathbb{H}$  is called a Schur function (resp. a generalized Schur function) if the kernel

$$K_S(p, q) = \sum_{u=0}^{\infty} p^u (I_n - S(p)S(q)^*) \bar{q}^u = (I_n - S(p)S(q)^*) \star (1 - p\bar{q})^{-\star} \tag{1.8}$$

is positive definite (resp. has a finite number of negative squares) in  $\Omega$ .

*Remark 1.7.* Realizations of operator-valued slice hyperholomorphic functions with real positive part have been considered in [7,9]. There the main tools were reproducing kernel Hilbert spaces and the theory of relations in Hilbert and Pontryagin spaces. Here the strategy is different (although a bit overlapping). The strategy used to prove a quaternionic positive real lemma is to start from a positive definite kernel in the quaternions, restrict it to the real line and apply the map  $\chi$  (see (2.1)) to obtain a condition on a complex matrix-valued positive definite on an open interval of the real line. An extension theorem (called Fitzgerald’s theorem in [28, Theorem p. 144]; see [31]) allows to extend it to the complex plane and so apply the classical positive real lemma on the new kernel. The case of negative squares requires an extension of this result. The underlying structure allows to go back to the quaternionic setting.

*Remark 1.8.* When the kernel  $K_\Phi$  is not assumed to have a finite number of negative squares, one can still obtain realizations with an interesting structure using Krein spaces; see [25,26] and, in the quaternionic setting, [10].

The paper contains eight sections besides the introduction, and we now proceed to describe their contents. Section 2 contains some preliminaries on quaternions and slice hyperholomorphic functions. Section 3 provides some characterizations of the image of quaternionic matrices under a suitable map denoted by  $\chi$  and also of the minimality of a triple of matrices. In Sect. 4 we prove a quaternionic version of the positive real lemma in the case of the half space and unit ball. Section 5 contains the extension result in the case of kernels with negative squares mentioned in the previous paragraph. Finally, in Sect. 6 we extend the results in Sect. 4 to the case of negative squares.

## 2. Slice Hyperholomorphic Functions

In this section we recall some basic facts about quaternions and slice hyperholomorphic functions which may be useful in the sequel. We refer the interested reader to the book [9] for a more complete treatment of these topics.

The skew field  $\mathbb{H}$  of quaternions consists of elements of the form  $p = x_0 + x_1i + x_2j + x_3k$ ,  $x_i \in \mathbb{R}$ ,  $i = 0, 1, 2, 3$ , where the imaginary units  $i, j, k$  satisfy

$$i^2 = j^2 = k^2 = -1, \quad ij = -ji = k, \quad jk = -kj = i, \quad ki = -ik = j.$$

The real number  $x_0$  also denoted by  $\text{Re}(p)$  is called real part of  $p$  while  $x_1i + x_2j + x_3k$  is called vector part or imaginary part of  $p$ . A quaternion  $p = x_0 + x_1i + x_2j + x_3k$  may be identified with a vector in  $\mathbb{R}^4$  and its norm is the Euclidean norm in  $\mathbb{R}^4$ , i.e.  $|p| = \sqrt{x_0^2 + x_1^2 + x_2^2 + x_3^2}$ . For any  $p \in \mathbb{H}$  as above its conjugate is defined by  $\bar{p} = x_0 - x_1i - x_2j - x_3k$ ; note that  $p\bar{p} = \bar{p}p = |p|^2$ .

By selecting one of the imaginary units of the basis, e.g. the unit  $i$ , a quaternion  $p$  can be written in terms of two complex numbers  $z_1, z_2$  belonging to the complex plane containing elements of the form  $x + iy$ , namely  $p = z_1 + z_2j$  with  $z_1 = x_0 + ix_1, z_2 = x_2 + ix_3$ . Moreover, one may identify the quaternion  $p$  with a  $2 \times 2$  complex valued matrix via the map  $\chi_i$  defined by (see [40])

$$\chi_i(p) = \begin{pmatrix} z_1 & z_2 \\ -\bar{z}_2 & \bar{z}_1 \end{pmatrix}. \tag{2.1}$$

In the sequel, in order to ease the notation the subscript will be omitted and we will write  $\chi$  instead of  $\chi_i$ . The map  $\chi : \mathbb{H} \rightarrow \mathbb{C}^{2 \times 2}$  is an injective homomorphism of rings, i.e.  $\chi(p + q) = \chi(p) + \chi(q)$  and  $\chi(pq) = \chi(p)\chi(q)$ , for any  $p, q \in \mathbb{H}$ . Further properties of  $\chi$  may be found in [40].

It is also useful to introduce the set  $\mathbb{S}$  of purely imaginary quaternions with square equal  $-1$ . This set coincides with the set of purely imaginary quaternions with norm 1 and it is a 2-dimensional sphere in  $\mathbb{H}$  identified with  $\mathbb{R}^4$ . Any nonreal quaternion  $q$  uniquely identifies the element  $I \in \mathbb{S}$  defined by  $I = x_1i + x_2j + x_3k / |x_1i + x_2j + x_3k|$ , so that  $p = x + Iy$  with  $x = x_0$  and  $y = |x_1i + x_2j + x_3k|$ . Moreover  $p$  defines a 2-sphere  $[p]$  given by  $[p] = \{x + Jy, J \in \mathbb{S}\}$ .

We now introduce the notion of slice hyperholomorphic function with values in a quaternionic Banach space. These functions are naturally defined on open sets  $\Omega$  which are axially symmetric, i.e. such that  $p \in \Omega$  implies that  $[p] \subset \Omega$ . It is immediate that the quaternionic unit ball  $\mathbb{B}$  and the half space  $\mathbb{H}_+$  of quaternions with positive real part are axially symmetric.

**Definition 2.1.** Let  $\Omega \subseteq \mathbb{H}$  be an axially symmetric set and let  $\mathcal{X}$  be a two sided quaternionic Banach space. A function  $f : \Omega \rightarrow \mathcal{X}$  of the form  $f(p) = f(x + Iy) = \alpha(x, y) + I\beta(x, y)$  where  $\alpha, \beta : \Omega \rightarrow \mathcal{X}$  depend only on  $x, y$ , are real differentiable, satisfy the Cauchy-Riemann equations

$$\begin{cases} \partial_x \alpha - \partial_y \beta = 0 \\ \partial_y \alpha + \partial_x \beta = 0, \end{cases} \tag{2.2}$$

and

$$\alpha(x, -y) = \alpha(x, y), \quad \beta(x, -y) = -\beta(x, y) \tag{2.3}$$

is said to be (left) slice hyperholomorphic.

Observe that, in the definition, if  $p = x$  is a real quaternion, then  $I$  is not uniquely defined but the hypothesis that  $\beta$  is odd in the variable  $y$  implies  $\beta(x, 0) = 0$ .

One can also introduce the definition of right slice hyperholomorphic function: the only change in the previous definition is that  $f(p) = f(x + Iy) = \alpha(x, y) + \beta(x, y)I$ .

The definition holds, in particular, when  $\mathcal{X} = \mathbb{H}^{n \times m}$  the quaternionic two sided Banach space of  $n \times m$  matrices with quaternionic entries and in the scalar valued case.

When we consider two slice hyperholomorphic functions  $f, g$  on  $\Omega$  with values in  $\mathcal{X}, \mathcal{X}_1$ , respectively, and it makes sense to define the product  $AB$  of  $A \in \mathcal{X}, B \in \mathcal{X}_1$  we can define the  $\star$ -product of  $f$  and  $g$  as follows: let  $f(p) = f(x + Iy) = \alpha(x, y) + I\beta(x, y), g(p) = g(x + Iy) = \gamma(x, y) + I\delta(x, y)$ , then

$$f \star g = (\alpha\gamma - \beta\delta) + I(\alpha\delta + \beta\gamma).$$

In the case a function  $f$  is scalar valued, it makes sense to define its  $\star$ -reciprocal. We refer the reader to [9, 22] for more details. Here it is sufficient to recall how to construct the  $\star$ -inverses of the functions  $\varphi(p) = p + q, \psi(p) = 1 - p\bar{q}$  which are left hyperholomorphic in  $p$  (and right slice hyperholomorphic in  $\bar{q}$ ):

$$\begin{aligned} (1 - p\bar{q})^{-\star} &= (1 - 2\operatorname{Re}(q)p + |q|^2p^2)^{-1}(1 - pq) \\ (p + \bar{q})^{-\star} &= (|q|^2 + 2\operatorname{Re}(q)p + p^2)^{-1}(p + q). \end{aligned}$$

### 3. Minimality

The map  $\chi = \chi_i$  that has been introduced in the previous section can be extended to quaternionic matrices, in fact if  $M = (q_{\ell s}) \in \mathbb{H}^{n \times m}, q_{\ell s} = z_{1, \ell s} + z_{2, \ell s}j, z_{1, \ell s}, z_{2, \ell s} \in \mathbb{C}$  we set  $M = A + Bj$  where  $A = (z_{1, \ell s}), B = (z_{2, \ell s}) \in \mathbb{C}^{n \times m}$ . Then we define

$$\chi(M) = \chi(A + Bj) = \begin{pmatrix} A & B \\ -\bar{B} & \bar{A} \end{pmatrix}.$$

In this section, we study the connections between the matrices associated to a quaternionic realization and their images under the map  $\chi$ . Our first goal is to characterize the range of  $\chi$ .

**Proposition 3.1.** *Let  $(A, B) \in \mathbb{C}^{n \times m} \times \mathbb{C}^{n \times m}$  and let  $\xi_1, \xi_2 \in \mathbb{C}^m$  be such that*

$$\begin{pmatrix} A & B \\ -\bar{B} & \bar{A} \end{pmatrix} \begin{pmatrix} \xi_1 \\ -\bar{\xi}_2 \end{pmatrix} = \begin{pmatrix} 0_n \\ 0_n \end{pmatrix}. \tag{3.1}$$

Then

$$\begin{pmatrix} A & B \\ -\bar{B} & \bar{A} \end{pmatrix} \begin{pmatrix} \xi_2 \\ \bar{\xi}_1 \end{pmatrix} = \begin{pmatrix} 0_n \\ 0_n \end{pmatrix}. \tag{3.2}$$

*Proof.* Applying conjugation to both sides of (3.1) we get

$$\begin{pmatrix} \bar{A} & \bar{B} \\ -B & A \end{pmatrix} \begin{pmatrix} \bar{\xi}_1 \\ -\xi_2 \end{pmatrix} = \begin{pmatrix} 0_n \\ 0_n \end{pmatrix}, \tag{3.3}$$

and so

$$\underbrace{\begin{pmatrix} 0 & -I_m \\ I_m & 0 \end{pmatrix} \begin{pmatrix} \bar{A} & \bar{B} \\ -B & A \end{pmatrix} \begin{pmatrix} 0 & I_m \\ -I_m & 0 \end{pmatrix}}_{\begin{pmatrix} A & B \\ -\bar{B} & \bar{A} \end{pmatrix}} \begin{pmatrix} 0 & -I_m \\ I_m & 0 \end{pmatrix} \begin{pmatrix} \bar{\xi}_1 \\ -\xi_2 \end{pmatrix} = \begin{pmatrix} 0_n \\ 0_n \end{pmatrix}, \tag{3.4}$$

which is (3.2). □

As a corollary we get:

**Corollary 3.2.** *With  $A, B, \xi_1$  and  $\xi_2$  as above,*

$$\chi(A + Bj) \begin{pmatrix} \bar{\xi}_1 \\ -\xi_2 \end{pmatrix} = \begin{pmatrix} 0_n \\ 0_n \end{pmatrix} \iff (A + Bj)(\xi_1 + \xi_2 j) = 0. \tag{3.5}$$

**Lemma 3.3.** *A matrix  $M \in (\mathbb{C}^{n \times m})^{2 \times 2}$  is of the form*

$$\begin{pmatrix} A & B \\ -\bar{B} & \bar{A} \end{pmatrix}$$

*if and only if it satisfies*

$$E_n^{-1} \bar{M} E_m = M, \tag{3.6}$$

*where*

$$E_m = \begin{pmatrix} 0 & I_m \\ -I_m & 0 \end{pmatrix}. \tag{3.7}$$

*Proof.* Let

$$M = \begin{pmatrix} M_{11} & M_{12} \\ M_{21} & M_{22} \end{pmatrix}$$

be the decomposition of  $M$  into  $\mathbb{C}^{n \times m}$  matrices. We have

$$E_n^{-1} \bar{M} E_m = \begin{pmatrix} \overline{M_{22}} & -\overline{M_{21}} \\ -\overline{M_{12}} & \overline{M_{11}} \end{pmatrix},$$

which will be equal to  $M$  if and only if  $M_{11} = \overline{M_{22}}$  and  $M_{12} = -\overline{M_{21}}$ , that is, if and only if  $M$  is of the form (3.6). □

*Remark 3.4.* We note that

$$E_m^{-1} = -E_m = E_m^* \tag{3.8}$$

and that the map  $a$  defined by

$$a(M) = E_m^{-1} \bar{M} E_m \tag{3.9}$$

is an involution and is real-linear. Moreover,  $a$  maps positive matrices into positive matrices since  $E_m^{-1} = E_m^*$ .

The notion of minimality in this section is according to the following definition:

**Definition 3.5.** We say that the triple of matrices  $(C, A, B) \in \mathbb{H}^{n \times N} \times \mathbb{H}^{N \times N} \times \mathbb{H}^{N \times m}$  is minimal if the following two conditions hold: The pair  $(C, A)$  is observable, that is

$$\bigcap_{t=0}^{\infty} \ker CA^t = \{0\} \tag{3.10}$$

and the pair  $(A, B)$  is controllable, that is

$$\bigcap_{t=0}^{\infty} \ker B^*A^{*t} = \{0\}. \tag{3.11}$$

Note that these intersections are finite (see [9], Section 9). The map  $\chi$  allows to reduce the study of minimality to the case of matrices with complex entries.

**Theorem 3.6.** *Let  $(C, A, B) \in \mathbb{H}^{n \times N} \times \mathbb{H}^{N \times N} \times \mathbb{H}^{N \times m}$ .*

- (1) *The pair  $(C, A) \in \mathbb{H}^{n \times N} \times \mathbb{H}^{N \times N}$  is controllable if and only if the corresponding pair  $(\chi(C), \chi(A))$  is controllable.*
- (2) *The pair  $(A, B) \in \mathbb{H}^{N \times N} \times \mathbb{H}^{N \times n}$  is observable if and only if the corresponding pair  $(\chi(A), \chi(B))$  is controllable.*
- (3) *The triple  $(C, A, B) \in \mathbb{H}^{n \times N} \times \mathbb{H}^{N \times N} \times \mathbb{H}^{N \times m}$  is minimal if and only if the corresponding triple  $(\chi(C), \chi(A), \chi(B))$  is minimal.*

*Proof.* Using (3.5) we have

$$\xi_1 + \xi_2 j \in \bigcap_{u=0}^{\infty} \ker CA^u \iff \begin{pmatrix} \xi_1 \\ -\xi_2 \end{pmatrix} \in \bigcap_{u=0}^{\infty} \ker \chi(C)\chi(A)^u,$$

from which we get the first claim; the other claims follow easily. □

We remark that other kind of realizations are possible. Assuming that  $R$  is slice hyperholomorphic at  $\infty$  (that is, in a set of the form  $|p| > r$ ) one can consider realizations of the form

$$R(p) = D + C \star (pI - A)^{-*} B.$$

In the setting of complex numbers and analytic functions, a matrix-valued analytic function is contractive in the open unit disk if and only if the kernel

$$K_S(z, w) = \frac{I_n - S(z)S(w)^*}{1 - z\bar{w}} \tag{3.12}$$

is positive definite in the open unit disk. We now mention an extension result (see [4, Théorème 2.6.5, p. 45], which roughly speaking, states that, in the positive case, positivity of the kernel  $K_S$  implies analyticity of the function  $S$ . Such results (for Carathéodory functions rather than Schur functions) originate in the work of Loewner, see [28, 36], when the positivity is assumed on the boundary. Here we use a different result, where the positivity is not on the boundary. For completeness we give a proof.



**Theorem 3.7.** *Let  $S$  be a  $\mathbb{C}^{n \times m}$ -valued function defined on a subset  $\Omega \subset \mathbb{D}$  and such that the kernel  $K_S$  is positive definite in  $\Omega$ . Assume that  $\Omega$  has an accumulation point inside  $\mathbb{D}$ . Then,  $S$  is the restriction of a uniquely defined Schur function in  $\mathbb{D}$ , and conversely.*

*Outline of the proof.* To ease the notation we consider the scalar case and write  $s(z)$  rather than  $S(z)$ . Only one direction is non trivial and will be proved. Let  $\mathbf{H}_2$  denote the Hardy space of the disk. Define a linear relation  $\mathcal{R}$  in  $\mathbf{H}_2 \times \mathbf{H}_2$  as the linear span of the pairs

$$\left( \frac{1}{1 - z\bar{w}}, \frac{\overline{s(w)}}{1 - z\bar{w}} \right), \quad w \in \Omega.$$

$\mathcal{R}$  is densely defined since  $\Omega$  has an accumulation point, and is contractive because the kernel  $K_S$  is positive definite on  $\Omega$ . It follows that  $\mathcal{R}$  extends to the graph of an everywhere defined contraction, say  $T$ . Its adjoint satisfies

$$(T^*1)(w) = s(w), \quad w \in \Omega$$

and is analytic, and hence  $s$  extends to a uniquely defined analytic function. To see that this function is a contraction, first check that for every integer  $n$ , and with  $M_z$  denoting the operator of multiplication by  $z$ ,

$$(T^*M_z^n 1)(w) = w^n s(w), \quad w \in \Omega.$$

Thus  $T^*$  is the multiplication operator by  $s$ , and so  $s$  is contractive since  $T^*$  is contractive. □

*Remark 3.8.* An alternative proof consists of solving the Nevanlinna-Pick theorem for an increasing family of finite subsets of  $\Omega$  and use Montel's theorem, see also Corollary 3.4 in [5] for the quaternionic setting. An earlier result, also for the quaternionic setting and for a subclass of Schur functions, is given in [8, Theorem 4.9, p. 98].

In the quaternionic setting, the kernel  $K_S$  (defined by (3.12)) takes now the form

$$(I_n - S(p)S(q)^*) \star (1 - p\bar{q})^{-*}. \tag{3.13}$$

The equivalence of the positivity of (3.13) with the fact that  $S$  is slice hyperholomorphic in the quaternionic unit ball and bounded by 1 in norm there holds for  $\mathbb{H}$ -valued functions (see [5]), but not for matrix-valued functions, as illustrated by the example (see [9, §8.4, p. 213])

$$U(p) = \frac{1}{\sqrt{2}} \begin{pmatrix} p & i \\ pi & 1 \end{pmatrix}.$$

By Cayley transform a similar remark holds for Herglotz functions.

### 4. The Quaternionic Positive Real Lemma

We now give the quaternionic version of the positive real lemma. As in the case of complex numbers, the set of Hermitian matrices  $H$  appearing in the statement form a convex set.

**Theorem 4.1.** *Let  $\phi$  be a  $\mathbb{H}^{n \times n}$ -valued slice hyperholomorphic rational function, slice hyperholomorphic at infinity, with minimal realization*

$$\phi(p) = D + C \star (pI_N - A)^{-\star} B. \tag{4.1}$$

Then,

$$(\phi(p) + \phi(q)^*) \star (p + \bar{q})^{-\star} \geq 0, \quad p, q \in \mathbb{H}_+ \tag{4.2}$$

if and only if there exists a negative definite matrix  $H \in \mathbb{H}^{N \times N}$  such that

$$\begin{pmatrix} H & 0 \\ 0 & I_n \end{pmatrix} \begin{pmatrix} A & B \\ C & D \end{pmatrix} + \begin{pmatrix} A & B \\ C & D \end{pmatrix}^* \begin{pmatrix} H & 0 \\ 0 & I_n \end{pmatrix} \geq 0. \tag{4.3}$$

*Proof.* Restricting to the positive real line and using the map  $\chi$ , we get that the kernel

$$\frac{\chi(\phi)(x) + \chi(\phi)(y)^*}{x + y} \tag{4.4}$$

is positive definite for  $x, y$  on the positive real line. By Theorem 3.6 the realization

$$\chi(\phi)(x) = \chi(D) + \chi(C) (xI_{2N} - \chi(A))^{-1} \chi(B)$$

is minimal. It follows from Theorem 3.7 (using Cayley transforms both on the variable and on  $\chi(\phi)$ ) that the kernel is positive definite in the whole right half-plane. We can therefore apply the positive real lemma which asserts that there exists a negative definite matrix  $H \in \mathbb{C}^{2N \times 2N}$  such that

$$\begin{pmatrix} H & 0 \\ 0 & I_{2n} \end{pmatrix} \begin{pmatrix} \chi(A) & \chi(B) \\ \chi(C) & \chi(D) \end{pmatrix} + \begin{pmatrix} \chi(A) & \chi(B) \\ \chi(C) & \chi(D) \end{pmatrix}^* \begin{pmatrix} H & 0 \\ 0 & I_{2n} \end{pmatrix} \geq 0. \tag{4.5}$$

We now show that there exists a solution  $H_1$  to (4.5) such that  $H_1 = a(H)$ , where  $a$  is defined by (3.9). By Lemma 3.3 and Remark 3.4, we deduce that there exists a positive quaternionic matrix  $X$  such that  $H_1 = \chi(X)$ . To that purpose, we show that if  $H$  satisfies (4.5) so does  $a(H)$ . Then,  $\frac{H+a(H)}{2}$  satisfies the symmetry (3.6). Using (3.8) we first note that

$$\begin{pmatrix} E_N^{-1} & 0 \\ 0 & E_n^{-1} \end{pmatrix} = \begin{pmatrix} E_N & 0 \\ 0 & E_n \end{pmatrix}^*.$$

Thus, we have from (4.5)

$$\begin{aligned} & \begin{pmatrix} E_N^{-1} & 0 \\ 0 & E_n^{-1} \end{pmatrix} \begin{pmatrix} \overline{H} & 0 \\ 0 & I_{2n} \end{pmatrix} \begin{pmatrix} E_N & 0 \\ 0 & E_n \end{pmatrix} \begin{pmatrix} E_N^{-1} & 0 \\ 0 & E_n^{-1} \end{pmatrix} \\ & \times \begin{pmatrix} \overline{\chi(A)} & \overline{\chi(B)} \\ \chi(C) & \chi(D) \end{pmatrix} \begin{pmatrix} E_N & 0 \\ 0 & E_n \end{pmatrix} \\ & + \begin{pmatrix} E_N^{-1} & 0 \\ 0 & E_n^{-1} \end{pmatrix} \begin{pmatrix} \overline{\chi(A)} & \overline{\chi(B)} \\ \chi(C) & \chi(D) \end{pmatrix}^* \begin{pmatrix} E_N & 0 \\ 0 & E_n \end{pmatrix} \begin{pmatrix} E_N^{-1} & 0 \\ 0 & E_n^{-1} \end{pmatrix} \\ & \times \begin{pmatrix} \overline{H} & 0 \\ 0 & I \end{pmatrix} \begin{pmatrix} E_N & 0 \\ 0 & E_n \end{pmatrix} \geq 0. \end{aligned} \tag{4.6}$$

Using Lemma 3.3 we have

$$\begin{aligned} & \begin{pmatrix} E_N^{-1} & 0 \\ 0 & E_n^{-1} \end{pmatrix} \begin{pmatrix} \overline{H} & 0 \\ 0 & I_{2n} \end{pmatrix} \begin{pmatrix} E_N & 0 \\ 0 & E_n \end{pmatrix} \begin{pmatrix} \chi(A) & \chi(B) \\ \chi(C) & \chi(D) \end{pmatrix} + \\ & + \begin{pmatrix} \chi(A) & \chi(B) \\ \chi(C) & \chi(D) \end{pmatrix}^* \begin{pmatrix} E_N^{-1} & 0 \\ 0 & E_n^{-1} \end{pmatrix} \begin{pmatrix} \overline{H} & 0 \\ 0 & I_n \end{pmatrix} \begin{pmatrix} E_N & 0 \\ 0 & E_n \end{pmatrix} \geq 0. \end{aligned} \quad (4.7)$$

Thus  $E_N^{-1}\overline{H}E_N$  satisfies (4.5), and so  $H_1 = \frac{H+a(H)}{2}$  is in the range of  $\chi$ . By Lemma 3.3 with  $H = \chi(X)$  it follows (4.3).

The converse follows the computations in [29, p. 26] and in [30, p. 129]. As in [29] we denote by

$$\begin{pmatrix} Q & S \\ S^* & R \end{pmatrix} \geq 0$$

the right handside of (4.3). Then

$$HA + A^*H = Q \quad (4.8)$$

$$HB + C^* = S \quad (4.9)$$

$$D + D^* = R. \quad (4.10)$$

For  $x, y \in \mathbb{R}$ , and replacing  $B$  by  $H^{-1}(S - C^*)$  we have

$$\begin{aligned} \phi(x) + \phi(y)^* &= D + C(xI - A)^{-1}B + D^* + B^*(yI - A^*)^{-1}C^* \\ &= D + C(xI - A)^{-1}H^{-1}(S - C^*) + D^* \\ &\quad + (S^* - C)H^{-1}(yI - A^*)^{-1}C^* \\ &= R + C(xI - A)^{-1}H^{-1}S + S^*H^{-1}(yI - A^*)^{-1} - \\ &\quad - \{C(xI - A)^{-1}H^{-1}C^* + CH^{-1}(yI - A^*)^{-1}C^*\} \\ &= R + C(xI - A)^{-1}H^{-1}S + S^*H^{-1}(yI - A^*)^{-1} - \\ &\quad - (x + y) \{C(xI - A)^{-1}H^{-1}(yI - A^*)^{-1}C^*\} + \\ &\quad + C(xI - A)^{-1}H^{-1}QH^{-1}(yI - A^*)^{-1}C^*. \end{aligned}$$

Hence

$$\begin{aligned} K_\phi(x, y) &= \frac{\phi(x) + \phi(y)^*}{x + y} = -C(xI - A)^{-1}H^{-1}(yI - A^*)^{-1}C^* + \\ &\quad + \frac{(C(xI - A)^{-1}H^{-1} \quad I) \begin{pmatrix} Q & S \\ S^* & R \end{pmatrix} \begin{pmatrix} H^{-1}(yI - A^*)^{-1}C^* \\ I \end{pmatrix}}{x + y}. \end{aligned} \quad (4.11)$$

The kernel (4.11) admits an extension slice hyperholomorphic in  $p$  and right slice hyperholomorphic in  $\bar{q}$  given by

$$\begin{aligned} K_\phi(p, q) &= (\phi(p) + \phi(q)^*)(p + \bar{q})^{-*} \\ &= -C \star (pI - A)^{-*}H^{-1}(\bar{q} - A^*)^{-*} \star_r C^* + \\ &\quad + \left( (C \star (pI - A)^{-*}H^{-1} \quad I) \begin{pmatrix} Q & S \\ S^* & R \end{pmatrix} \begin{pmatrix} H^{-1}(\bar{q}I - A^*)^{-*} \star_r C^* \\ I \end{pmatrix} \right) \star (p + \bar{q})^{-*} \end{aligned} \quad (4.12)$$

(where all the  $\star$ -product are computed in the variable  $p$ ) and so  $K_\phi$  is positive definite in the right half-space.  $\square$

*Remark 4.2.* Alternatively, and replacing now  $C$  by  $S^* - HB^*$  we also have

$$\begin{aligned} \phi(x) + \phi(y)^* &= D + C(xI - A)^{-1}B + D^* + B^*(yI - A^*)^{-1}C^* \\ &= R + (S - B^*H)(xI - A)^{-1}B + B^*(yI - A^*)^{-1}(S^* - HB) \\ &= R + S(xI - A)^{-1}B + B^*(yI - A)^{-*}S^* - \\ &\quad - \{B^*H(xI - A)^{-1}B + B^*(yI - A^*)^{-1}HB\} \\ &= R + S(xI - A)^{-1}B + B^*(yI - A)^{-*}S^* - \\ &\quad - \{B^*(yI - A^*)^{-1}(H(xI - A) + (yI - A^*)H)B \\ &\quad + B^*(xI - A)^{-1}B\} \\ &= R + S(xI - A)^{-1}B + B^*(yI - A)^{-*}S^* - \\ &\quad - (x + y)B^*(yI - A^*)^{-1}H(xI - A)^{-1}B + \\ &\quad + B^*(yI - A^*)^{-1}Q(xI - A)^{-1}B. \end{aligned}$$

Hence

$$\begin{aligned} \frac{\phi(x) + \phi(y)^*}{x + y} &= -B^*(yI - A^*)^{-1}H(xI - A)^{-1}B + \\ &\quad + \frac{\begin{pmatrix} B^*(yI - A^*)^{-1} & I \end{pmatrix} \begin{pmatrix} Q & S \\ S^* & R \end{pmatrix} \begin{pmatrix} (xI - A)^{-1}B \\ I \end{pmatrix}}{x + y}. \end{aligned} \tag{4.13}$$

We now consider the case of the quaternionic unit ball.

**Theorem 4.3.** *A  $\mathbb{H}^{n \times n}$ -valued rational function slice hyperholomorphic at  $\infty$  and with minimal realization (4.1) is contractive in the quaternionic unit ball if and only if there exists a negative definite matrix  $H \in \mathbb{H}^{N \times N}$  such that*

$$\begin{pmatrix} H & 0 \\ 0 & I_n \end{pmatrix} - \begin{pmatrix} A & B \\ C & D \end{pmatrix}^* \begin{pmatrix} H & 0 \\ 0 & I_n \end{pmatrix} \begin{pmatrix} A & B \\ C & D \end{pmatrix} \geq 0. \tag{4.14}$$

*Proof.* As in the previous, one reduces the claim to the complex setting using the map  $\chi$ , and use [24, Theorem 2]. We then go back to the quaternionic setting as in the proof of the previous theorem.  $\square$

We also remark that the set of Hermitian matrices  $H$  satisfying (4.14) form a convex set.

### 5. The Case of Negative Squares: An Extension Theorem

We start this section by recalling a definition which, as explained in the introduction, makes sense since the spectral theorem holds for quaternionic Hermitian matrices (see also [9]).

**Definition 5.1.** Let  $K(u, v)$  be a  $\mathbb{H}^{n \times n}$ -valued function defined for  $u, v$  in some set  $\Omega$  and assume that  $K(u, v) = K(v, u)^*$  for all  $u, v \in \Omega$ . The function  $K(u, v)$  is said to have  $\kappa$  negative squares if the following condition holds: for every  $N \in \mathbb{N}$ , and every choice of points  $u_1, \dots, u_N \in \Omega$  and vectors  $\xi_1, \dots, \xi_N \in \mathbb{H}^n$  the  $N \times N$  Hermitian matrix with  $(k, j)$  entry  $\xi_k^* K(u_k, u_j) \xi_j$  has at most  $\kappa$  strictly negative eigenvalues, and exactly  $\kappa$  strictly negative eigenvalues for some choice of  $N, u_1, \dots, u_N, \xi_1, \dots, \xi_N$ .

**Theorem 5.2.** Let  $F$  be a  $\mathbb{C}^{n \times n}$ -valued rational function, analytic at infinity. Then the following are equivalent:

- (1)  $\operatorname{Re} F(iy) \geq 0, \quad y \in \mathbb{R}.$
- (2) The kernel

$$K_F(z, w) = \frac{F(z) + F(w)^*}{z + \bar{w}}$$

has a finite number of negative squares in the right half-plane.

*Proof.* Assume (1). By maybe adding to  $F$  a term of the form  $i\epsilon I_n$  we do not change the first condition but may assume that the Cayley transform  $S = (I_n + F)^{-1}(I_n - F)$  is well defined. We make another Cayley transform this time on the variable, to get to a function  $S_1$  contractive on the unit circle. By multiplying  $S_1$  by a finite scalar Blaschke factor  $b$  we get a function  $bS_1$  which is analytic in  $\mathbb{D}$  and contractive on the unit circle. It is therefore a Schur function  $S_0$  and so  $S_1 = \frac{S_0}{b}$  is a generalized Schur function, and the corresponding kernel

$$K_{S_1}(z, w) = \frac{I_n - S_1(z)S_1(w)^*}{1 - z\bar{w}}$$

has a finite number of negative squares, see [34]. Since  $K_F$  and  $K_{S_1}$  are related by Cayley transforms, we get that  $K_F$  has a finite number of negative squares.

The converse is proved by reading backwards the above arguments.  $\square$

We now remark that Theorem 3.7 will not hold in general in the case of negative squares. As recalled in [13, p. 82], the function defined by

$$S(z) = \begin{cases} 1, & z \in \mathbb{D} \setminus \{0\} \\ 0, & z = 0, \end{cases}$$

is not meromorphic in the open unit disk, while the corresponding kernel  $K_S$  has one negative square. A claim on a counterpart of the extension theorem was made in [11, Theorem 3.2], but the proof turned out to be flawed; see [12]. We now prove a weaker result, which is enough for our purposes, when one knows ahead of time that the function  $K_S$  is meromorphic in the open unit disk, and has a finite number of negative squares in a subset  $\Omega$  which has an accumulation point. We focus on the matrix-valued rational case for simplicity, and which is what is needed in the present paper. The result itself can be seen as an extension of [28, p. 144] to the setting of negative squares (although beyond the scope of the current work, note that the case of analytic functions can be similarly treated).

**Theorem 5.3.** *Let  $S$  be a  $\mathbb{C}^{n \times n}$ -valued rational function such that the kernel  $K_S(z, w)$  has a finite number of negative squares in  $(-1, 1)$ , from which have been removed possibly a finite number of points  $P$ . Then, the kernel  $K_S(z, w)$  has the same number of negative squares in the open unit disk, from which have been removed possibly a finite number of points  $P_1$ .*

Before proving the theorem we recall a result on positive definite kernels.

**Theorem 5.4.** *Let  $\Lambda_1(z, w)$  and  $\Lambda_2(z, w)$  be two  $\mathbb{C}^{n \times n}$ -valued kernels, positive definite on the set  $\Omega$ , and let  $\mathcal{H}(\Lambda_1), \mathcal{H}(\Lambda_2)$  and  $\mathcal{H}(\Lambda_1 + \Lambda_2)$  denote the reproducing kernel Hilbert spaces associated to  $\Lambda_1, \Lambda_2$  and  $\Lambda_1 + \Lambda_2$  respectively. Then*

$$\mathcal{H}(\Lambda_1 + \Lambda_2) = \mathcal{H}(\Lambda_1) + \mathcal{H}(\Lambda_2). \tag{5.1}$$

*More precisely, every element in  $\mathcal{H}(\Lambda_1 + \Lambda_2)$  can be written as a (in general non-unique) sum  $f = f_1 + f_2$ , where  $f_j \in \mathcal{H}(\Lambda_j), j = 1, 2$ . For every such decomposition*

$$\|f\|_{\mathcal{H}(\Lambda_1 + \Lambda_2)}^2 \leq \|f_1\|_{\mathcal{H}(\Lambda_1)}^2 + \|f_2\|_{\mathcal{H}(\Lambda_2)}^2, \tag{5.2}$$

*and there is a unique decomposition for which equality holds. For the function*

$$f(z) = \sum_{u=1}^U (\Lambda_1(z, w_u) + \Lambda_2(z, w_u))c_u, \tag{5.3}$$

*the minimal (unique) decomposition for which equality holds in (5.2) is given by*

$$f_1(z) = \sum_{u=1}^U \Lambda_1(z, w_u)c_u, \quad \text{and} \quad f_2(z) = \sum_{u=1}^U \Lambda_2(z, w_u)c_u. \tag{5.4}$$

The proof of this result can be found in [18]. The fact that the equality in (5.2) holds for (5.4) when  $f$  is of the form (5.3) is crucial in the proof.

*Proof of Theorem 5.3.* The strategy of the proof is as follows.

$$K_S(z, w) = \frac{I_n}{1 - z\bar{w}} - \frac{S(z)S(w)^*}{1 - z\bar{w}}$$

is a difference of two positive definite functions, and hence is the reproducing kernel of a (in general not uniquely defined) reproducing kernel Krein space; see [37]. We study the restriction of the elements of this space to  $(-1, 1) \setminus P$  to see that it is a Pontryagin space.

We proceed in a number of steps.

STEP 1: *The function*

$$M_S(z, w) = \frac{I_n}{1 - z\bar{w}} + \frac{S(z)S(w)^*}{1 - z\bar{w}}$$

*is positive definite in  $\mathbb{D} \setminus P_1$ . It is the reproducing kernel of a (uniquely defined) Hilbert space of  $\mathbb{C}^n$ -valued functions analytic in  $\mathbb{D} \setminus P_1$ .*

The first part of the claim is just Aronszajn-Moore result; the second part follows from the joint analyticity of  $M_S(z, w)$  in  $z$  and  $\bar{w}$ . Such kernels are called Bergman kernels in [28]. See e.g. [28, Theorem p. 92] or [4, Théorème 2.3.5, p.31] for the latter claim on analyticity. We will denote the reproducing kernel Hilbert space with reproducing kernel  $M_S(z, w)$  by  $\mathcal{H}(M_S)$ .

STEP 2: *The linear relation  $R$  of  $\mathcal{H}(M_S) \times \mathcal{H}(M_S)$  spanned by the functions*

$$(M_S(\cdot, w)c, K_S(\cdot, w)c) \tag{5.5}$$

*extends to the graph of a contractive self-adjoint map from  $\mathcal{H}(M_S)$  into itself.*

The proof follows, with slight modifications, the arguments in [3, Proof of Lemma 4, p. 177]. It is convenient to set

$$\Lambda_1(z, w) = \frac{I_n}{1 - z\bar{w}} \quad \text{and} \quad \Lambda_2(z, w) = \frac{S(z)S(w)^*}{1 - z\bar{w}}. \tag{5.6}$$

Let first  $f$  be a finite linear combination of kernels:

$$f(z) = \sum_{u=1}^U M_S(z, w_u)c_u = f_1(z) + f_2(z),$$

with

$$f_1(z) = \sum_{u=1}^U \frac{c_u}{1 - z\bar{w}_u} \in \mathcal{H}(\Lambda_1) \tag{5.7}$$

$$f_2(z) = \sum_{u=1}^U \frac{S(z)S(w_u)^*c_u}{1 - z\bar{w}_u} \in \mathcal{H}(\Lambda_2), \tag{5.8}$$

and assume  $f(z) \equiv 0$ . We have  $(0, g) \in R$  with  $g(z) = \sum_{u=1}^U K_S(z, w_u)c_u$  and we wish to show that  $g(z) \equiv 0$ . We have

$$\sum_{u=1}^U c_v^* M_S(w_v, w_u)c_u = 0, \quad v = 1, \dots, U$$

and so

$$\sum_{u,v=1}^U c_v^* M_S(w_v, w_u)c_u = 0,$$

which implies in particular that

$$\sum_{u,v=1}^U \frac{c_v^*c_u}{1 - w_v\bar{w}_v} = 0 \quad \sum_{u,v=1}^U \frac{c_v^*S(w_v)S(w_u)^*c_u}{1 - w_v\bar{w}_v} = 0$$

and so, by positivity of the functions  $\frac{I_n}{1 - z\bar{w}}$  and  $\frac{S(z)S(w)^*}{1 - z\bar{w}}$  we get

$$f_1(z) \equiv 0 \quad \text{and} \quad f_2(z) \equiv 0,$$

that is,  $g(z) \equiv 0$ , and  $R$  is the graph of a densely defined operator  $T$ . We note that

$$\|f\|_{\mathcal{H}(M_S)}^2 = \min_{\substack{f=h_1+h_2 \\ h_j \in \mathcal{H}(\Lambda_j) \ j=1,2}} \|h_1\|^2 + \|h_2\|^2 = \|f_1\|_{\mathcal{H}(\Lambda_1)}^2 + \|f_2\|_{\mathcal{H}(\Lambda_2)}^2. \tag{5.9}$$

STEP 3: We show now that the operator  $T$  is a self-adjoint contraction.

We have with  $f, f_1, f_2$  as above,

$$f = f_1 + f_2 \quad \text{and} \quad T(f_1 + f_2) = f_1 - f_2 \stackrel{\text{def.}}{=} g.$$

Note that Theorem 5.4 insures that  $Tf \in \mathcal{H}(M_S)$ . Moreover

$$\|g\|_{\mathcal{H}(M_S)}^2 = \min \left( \|g_1\|_{\mathcal{H}(\Lambda_1)}^2 + \|g_2\|_{\mathcal{H}(\Lambda_2)}^2 \right),$$

where the infimum, which in fact is a minimum, is computed over all decomposition  $g = g_1 - g_2$  with  $g_1 \in \mathcal{H}(\Lambda_1)$  and  $g_2 \in \mathcal{H}(\Lambda_2)$ . Thus, by (5.9)

$$\|Tf\|^2 = \|g\|^2 \leq \|f_1\|_{\mathcal{H}(\Lambda_1)}^2 + \|f_2\|_{\mathcal{H}(\Lambda_2)}^2 = \|f\|^2. \tag{5.10}$$

We now show that  $T$  is self-adjoint. We prove it on the kernels and will show that

$$\begin{aligned} \langle T((\Lambda_1(\cdot, w) + \Lambda_2(\cdot, w))c), (\Lambda_1(\cdot, v) + \Lambda_2(\cdot, v))d \rangle_{\mathcal{H}(M_S)} &= \\ &= \langle (\Lambda_1(\cdot, w) + \Lambda_2(\cdot, w))c, T((\Lambda_1(\cdot, v) + \Lambda_2(\cdot, v))d) \rangle_{\mathcal{H}(M_S)}, \end{aligned} \tag{5.11}$$

or equivalently

$$\begin{aligned} \langle (\Lambda_1(\cdot, w) - \Lambda_2(\cdot, w))c, (\Lambda_1(\cdot, v) + \Lambda_2(\cdot, v))d \rangle_{\mathcal{H}(M_S)} &= \\ &= \langle (\Lambda_1(\cdot, w) + \Lambda_2(\cdot, w))c, (\Lambda_1(\cdot, v) - \Lambda_2(\cdot, v))d \rangle_{\mathcal{H}(M_S)}. \end{aligned} \tag{5.12}$$

The reproducing kernel property shows that both sides of (5.12) coincide and are equal to

$$d^*(\Lambda_1(v, w) - \Lambda_2(v, w))c.$$

Being contractive and densely defined,  $T$  extends to a unique everywhere defined self-adjoint contraction from  $\mathcal{H}(M_S)$  into itself, which we still denote by  $T$ .

We set  $T = \sigma P$ , where  $P$  is the unique positive squareroot of  $T^2$  and  $\sigma$  is the sign of  $T$ . Recall that

$$P = \int_{\mathbb{R}} |t| dE_t \quad \text{and} \quad \sigma = \int_{\mathbb{R} \setminus \{0\}} \frac{t}{|t|} dE_t$$

where  $T = \int_{\mathbb{R}} t dE_t$  is the spectral decomposition of  $T$  (and where the integral is in fact on a bounded integral since  $T$  is bounded), and that  $\sigma$  and  $P$  commute.

STEP 4: The space

$$\mathcal{K} = \left\{ F = \sqrt{P}u, u \in \mathcal{H}(M_S); \|F\| = \|(I - \pi)u\|_{\mathcal{H}(M_S)} \right\} \tag{5.13}$$

where  $\pi$  is the orthogonal projection onto  $\ker P$ , is a Hilbert space.

STEP 5: Define

$$[Tu, Tv]_{\mathcal{K}} = \langle Tu, v \rangle_{\mathcal{H}(M_S)} \tag{5.14}$$

and

$$\langle Tu, Tv \rangle_{\mathcal{K}} = \langle Pu, v \rangle_{\mathcal{H}(M_S)}. \tag{5.15}$$



Then,  $(\mathcal{K}, \langle \cdot, \cdot \rangle_{\mathcal{K}})$  is a Hilbert space, and there exists a bounded and boundedly invertible self-adjoint operator  $G$  such that

$$[Tu, Tv]_{\mathcal{K}} = \langle Tu, GTv \rangle_{\mathcal{K}}. \quad (5.16)$$

We have

$$\begin{aligned} [\sigma Tu, \sigma Tv]_{\mathcal{K}} &= [Tu, Tv]_{\mathcal{K}} \\ [\sigma Tu, Tv]_{\mathcal{K}} &= [Tu, \sigma Tv]_{\mathcal{K}} \end{aligned}$$

so that  $\sigma$  satisfies the equations  $\sigma^2 = I$  and  $\sigma = \sigma^*$  in  $\mathcal{K}$ . We take  $G = \sigma$ .

STEP 6: By its definition, the space  $\mathcal{K}$  with the indefinite inner product (5.14) is a reproducing kernel Krein space the inner product with reproducing kernel  $K_S$ .

By (5.5) we have that  $z \mapsto K_S(z, w)c$  belongs to  $\mathcal{K}$ . Furthermore, we have:

$$\begin{aligned} \kappa &= [Tu, T(M_S(\cdot, w)c)]_{\mathcal{K}} \\ &= \langle Tu, (M_S(\cdot, w)c) \rangle_{\mathcal{H}(M_S)} \\ &= c^*(Tu)(w). \end{aligned} \quad (5.17)$$

STEP 7: The linear span  $\mathcal{M}$  of the functions  $K_S(\cdot, x)c$  with  $x \in (-1, 1) \setminus P$  is dense in  $\mathcal{K}$ .

Indeed, by the reproducing kernel property, any element  $f$  orthogonal to the linear span vanishes on  $(-1, 1) \setminus P$ . In view of the analyticity of the element of  $\mathcal{K}$ , and since  $\mathbb{D} \setminus P_1$  is connected (since  $P_1$  is finite), we get  $f \equiv 0$ .

At this stage of the proof we define  $\mathcal{P}(K_S)$  to be the (uniquely defined) reproducing kernel Pontryagin space with reproducing kernel  $K_S(z, w)$  with  $z, w$  restricted to  $(-1, 1) \setminus P$ .

STEP 8: The restriction map  $f \mapsto f|_{(-1, 1) \setminus P}$  is one to one from  $\mathcal{M}$  into  $\mathcal{P}(K_S)$ .

The proof of this step follows directly from the analyticity of the elements of  $\mathcal{M}$ .

STEP 9: The inner products of the elements of  $\mathcal{M}$  (and restricted to  $(-1, 1) \setminus P$ ) coincide in  $\mathcal{K}$  and  $\mathcal{P}(K_S)$  respectively.

Step 9 follows from the reproducing kernel property.

STEP 10:  $\mathcal{K}$  is in fact a Pontryagin space, made of the analytic extensions of the functions of  $\mathcal{P}(K_S)$  to  $\mathbb{D} \setminus P_1$ .

The linear span  $\mathcal{M}'$  of the functions  $K_S(\cdot, y)c$  with  $y \in (-1, 1) \setminus P$  and  $c \in \mathbb{C}^n$  (i.e. of the restrictions of the elements of  $\mathcal{M}$  to  $(-1, 1) \setminus P$ ) is dense in the Pontryagin space  $\mathcal{P}(K_S)$  because of the reproducing kernel property. It contains therefore a maximal strictly negative subspace  $\mathcal{P}_-$ . Write

$$\mathcal{P}(K_S) = \mathcal{P}_+ [+] \mathcal{P}_-$$

and for  $f \in \mathcal{M}'$  write

$$f = f_+ + f_-.$$

Both  $f_+$  and  $f_-$  are by definition finite linear combinations of elements of the form  $K(\cdot, y)c$ . Denote by  $I$  the restriction map appearing in Step 8. The spaces

$$\{I^{-1}(f_+); f = f_+ + f_-; f \in \mathcal{M}'\} \quad \text{and} \quad \{I^{-1}(f_-); f = f_+ + f_-; f \in \mathcal{M}'\} \tag{5.18}$$

are orthogonal in  $\mathcal{K}$ , and respectively positive and negative in  $\mathcal{K}$  thanks to Step 9. Since the second one is finite dimensional, (5.18) is a fundamental decomposition of  $\mathcal{K}$  with a finite dimensional negative part, and  $\mathcal{K}$  is a Pontryagin space.  $\square$

*Remark 5.5.* One can also build a reproducing kernel Krein space with reproducing kernel  $K_S$  of functions analytic in  $|z| < r < 1$  by multiplying  $K_S(z, w)$  by a scalar function  $f(z)\overline{f(w)}$  and apply [2, Theorem 3.1 p. 1198].

Let  $S$  be a rational slice function, hyperholomorphic at the origin, and with realization  $S(p) = D + pC \star (I - pA)^{-\star}B$ . The map  $x \mapsto \chi(S)(x)$  is the restriction of the rational function of a complex variable

$$\chi(S)(z) = D + z\chi(C)(I - z\chi(A))^{-1}\chi(B)$$

to the interval  $(-1, 1) \setminus P$  (at the possible exception of a finite set  $P_1$  in  $(-1, 1)$  where  $\det(I - xA)$  vanishes). By the above theorem, the kernel  $K_S$  has  $\kappa$  negative squares in  $(-1, 1)$  if and only if it has  $\kappa$  negative squares in  $\mathbb{D} \setminus P_1$ , where  $P_1$  are the zeros of  $\det(zI - A)$  in  $\mathbb{D}$ .

### 6. Generalized Quaternionic Positive Real Lemma

In this section we prove a generalized version of Theorem 4.1 in case the kernel has negative squares.

**Theorem 6.1.** *Let  $\phi$  be a  $\mathbb{H}^{n \times n}$ -valued slice hyperholomorphic rational function, slice hyperholomorphic at infinity, with minimal realization (4.1) Then, the kernel  $K_\Phi$  (defined by (4.2)) has a finite number of negative squares in the open right half-space  $\mathbb{H}_+$ , from which is possibly removed a finite set of points, if and only if there exists an Hermitian non-singular matrix  $H \in \mathbb{H}^{N \times N}$  such that (4.3) holds.*

*Proof.* Assume first that  $K_\Phi$  has a finite number of negative squares in  $\mathbb{H}_+ \setminus Q$ , where  $Q$  is a finite set of points (or is the empty set). Restricting  $p$  and  $q$  to  $\mathbb{R}_+ \setminus Q$  and applying the map  $\chi$  we see that the kernel (4.4) has a finite number of negative squares on  $\mathbb{R}_+ \setminus Q$ . This kernel is the restriction to  $(\mathbb{R}_+ \setminus Q)^2$  of the kernel

$$\frac{(\chi(\phi))(z) + ((\chi(\phi))(w))^*}{z + \bar{w}}. \tag{6.1}$$

It follows from the representation theorem of Krein-Langer for generalized Schur function, and using a Cayley transform, or by the representation theorem of generalized positive real functions due to [23, 27] (both in the scalar case; see also [15]) we see that  $\text{Re } \chi(\phi)(x) \geq 0$  on the imaginary line. We can now resort to the result of [24] to assert that there is an Hermitian matrix

$H$  satisfying (4.5). The rest of this side of the proof goes as in the proof of Theorem 4.1.

The converse is as in the converse of Theorem 4.1, since formula (4.11) still holds for any invertible Hermitian matrix  $H$ . In (4.12) the kernel

$$\begin{aligned}
 & - C \star (pI - A)^{-\star} H^{-1} (\bar{q} - A^*)^{-\star_r} \star_r C^* + \\
 & + \left( (C \star (pI - A)^{-\star} H^{-1} \quad I) \begin{pmatrix} Q & S \\ S^* & R \end{pmatrix} \begin{pmatrix} H^{-1} (\bar{q}I - A^*)^{-\star_r} \star_r C^* \\ I \end{pmatrix} \right) \\
 & \star (p + \bar{q})^{-\star}
 \end{aligned}$$

has now (possibly) a finite number of negative squares and the kernel

$$(C \star (pI - A)^{-\star} H^{-1} \quad I) \begin{pmatrix} Q & S \\ S^* & R \end{pmatrix} \begin{pmatrix} H^{-1} (\bar{q}I - A^*)^{-\star_r} \star_r C^* \\ I \end{pmatrix} (p + \bar{q})^{-\star}$$

is still positive negative, for real  $x, y$  in both cases and then in the open right half-space by slice hyperholomorphic extension.  $\square$

We now consider the counterpart of Theorem 1.4 in the quaternionic setting for generalized Schur functions. Here we omit the proof since it makes use of the map  $\chi$  and of the proof of Theorem 6.1. We also remark that the set of Hermitian matrices which satisfy (6.2) form a convex set.

**Theorem 6.2.** *A  $\mathbb{H}^{n \times n}$ -valued rational function slice hyperholomorphic at  $\infty$  and with minimal realization (4.1) is contractive on the quaternionic unit sphere if and only if there exists an Hermitian non-singular matrix  $H \in \mathbb{H}^{N \times N}$  such that*

$$\begin{pmatrix} H & 0 \\ 0 & I_n \end{pmatrix} - \begin{pmatrix} A & B \\ C & D \end{pmatrix}^* \begin{pmatrix} H & 0 \\ 0 & I_n \end{pmatrix} \begin{pmatrix} A & B \\ C & D \end{pmatrix} \geq 0. \tag{6.2}$$

**Acknowledgements**

It is a pleasure to thank Prof. Rovnyak for comments on Theorem 5.2.

**Publisher’s Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**References**

- [1] Abu-Ghanem, Kh, Alpay, D., Colombo, F., Lewkowicz, I., Sabadini, I.: Herglotz functions of several quaternionic variables. *J. Math. Anal. Appl.* **466**(1), 169–182 (2018)
- [2] Alpay, D.: Some remarks on reproducing kernel Krein spaces. *Rocky Mt. J. Math.* **21**, 1189–1205 (1991)
- [3] Alpay, D.: On linear combination of positive functions, associated reproducing kernel spaces and a nonhermitian Schur algorithm. *Arch. Math. (Basel)* **58**, 174–182 (1992)
- [4] Alpay, D.: Algorithmes de Schur, espaces à noyau reproduisant et théorie des systèmes. *Panoramas et Synthèses [Panoramas and Syntheses]*, vol. 6. Société Mathématique de France, Paris (1998)

- [5] Alpay, D., Bolotnikov, V., Colombo, F., Sabadini, I.: Self-mappings of the quaternionic unit ball: multiplier properties, Schwarz–Pick inequality, and Nevanlinna–Pick interpolation problem. *Indiana Univ. Math. J.* **64**, 151–180 (2015)
- [6] Alpay, D., Colombo, F., Kimsey, D.P., Sabadini, I.: An extension of Herglotz’s theorem to the quaternions. *J. Math. Anal. Appl.* **421**(1), 754–778 (2015)
- [7] Alpay, D., Colombo, F., Lewkowicz, I., Sabadini, I.: Realizations of slice hyperholomorphic generalized contractive and positive functions. *Milan J. Math.* **83**, 91–144 (2015)
- [8] Alpay, D., Colombo, F., Sabadini, I.: Pontryagin-de Branges-Rovnyak spaces of slice hyperholomorphic functions. *J. Anal. Math.* **121**, 87–125 (2013)
- [9] Alpay, D., Colombo, F., Sabadini, I.: Slice Hyperholomorphic Schur Analysis, volume 256 of *Operator Theory: Advances and Applications*. Springer, Basel (2016)
- [10] Alpay, D., Colombo, F., Sabadini, I.: Realizations of holomorphic and slice hyperholomorphic functions: the Krein space case. *ArXiv e-prints* (2018)
- [11] Alpay, D., Constantinescu, T., Dijksma, A., Rovnyak, J.: Notes on interpolation in the generalized Schur class II. Nudel’man’s problem. *Trans. Am. Math. Soc.* **355**(2), 813–836 (2003)
- [12] Alpay, D., Dijksma, A., Rovnyak, J.: Corrigendum to “Notes on interpolation in the generalized Schur class. II. Nudel’man’s problem”. *Trans. Am. Math. Soc.* **371**(5), 3743–3745 (2019)
- [13] Alpay, D., Dijksma, A., Rovnyak, J., de Snoo, H.: *Schur Functions, Operator Colligations, and Reproducing Kernel Pontryagin Spaces*, volume 96 of *Operator Theory: Advances and Applications*. Birkhäuser Verlag, Basel (1997)
- [14] Alpay, D., Gohberg, I.: Unitary rational matrix functions. In: Gohberg, I. (ed.) *Topics in Interpolation Theory of Rational Matrix-Valued Functions*. *Operator Theory: Advances and Applications*, vol. 33, pp. 175–222. Birkhäuser Verlag, Basel (1988)
- [15] Alpay, D., Lewkowicz, I.: The positive real lemma and construction of all realizations of generalized positive rational functions. *Syst. Control Lett.* **60**, 985–993 (2011)
- [16] Anderson, B.D.O., Moore, J.B.: Algebraic structure of generalized positive real matrices. *SIAM J. Control* **6**, 615–624 (1968)
- [17] Arlinskii, Y.: The Kalman–Yakubovich–Popov inequality for passive discrete time-invariant systems. *Oper. Matrices* **2**(1), 15–51 (2008)
- [18] Aronszajn, N.: Theory of reproducing kernels. *Trans. Am. Math. Soc.* **68**, 337–404 (1950)
- [19] Bolotnikov, V., Rodman, L.: Krein–Langer factorizations via pole triples. *Integral Equ. Oper. Theory* **47**(2), 169–195 (2003)
- [20] Brune, O.: Synthesis of a finite two-terminal network whose driving-point impedance is a prescribed function of frequency. *J. Math. Phys.* **10**, 191–236 (1931)
- [21] Cauer, W.: The realization of impedances of prescribed frequency dependence (in German). *Arch. Elektrotech.* **17**, 355–388 (1926)
- [22] Colombo, F., Sabadini, I., Struppa, D.C.: *Noncommutative Functional calculus*, volume 289 of *Progress in Mathematics*. Theory and applications of slice hyperholomorphic functions. Birkhäuser, Basel (2011)

- [23] Derkach, V., Hassi, S., de Snoo, H.: Operator models associated with Kac subclasses of generalized Nevanlinna functions. *Methods Funct. Anal. Topol.* **5**(1), 65–87 (1999)
- [24] Dickinson, B., Delsarte, Ph, Genin, Y., Kamp, Y.: Minimal realizations of pseudo-positive and pseudo-bounded rational matrices. *IEEE Trans. Circuits Syst.* **32**, 603–605 (1985)
- [25] Dijksma, A., Langer, H., de Snoo, H.: Unitary Colligations in Kreĭn Spaces and Their Role in the Extension Theory of Isometries and Symmetric Linear Relations in Hilbert Spaces. *Functional analysis. II (Dubrovnik, 1985)*, pp. 1–42. Springer, Berlin (1987)
- [26] Dijksma, A., Langer, H., de Snoo, H.S.V.: Representations of holomorphic operator functions by means of resolvents of unitary or selfadjoint operators in Kreĭn spaces. In: Helson, H., Arsene, G. (eds.) *Operators in Indefinite Metric Spaces, Scattering Theory and Other Topics (Bucharest, 1985)*, volume 24 of *Oper. Theory Adv. Appl.*, pp. 123–143. Birkhäuser, Basel (1987)
- [27] Dijksma, A., Langer, H., Luger, A., Shondin, Yu.: A factorization result for generalized Nevanlinna functions of the class  $\mathcal{N}_\kappa$ . *Integral Equ. Oper. Theory* **36**(1), 121–125 (2000)
- [28] Donoghue, W.F.: *Monotone Matrix Functions and Analytic Continuation*, volume 207 of *Die Grundlehren der mathematischen Wissenschaften*. Springer, London (1974)
- [29] Faurre, P.: *Réalisations markoviennes de processus stationnaires*. Ph.D. thesis, INRIA (1973)
- [30] Faurre, P., Clerget, M., Germain, F.: *Opérateurs rationnels positifs*, volume 8 of *Méthodes Mathématiques de l’Informatique [Mathematical Methods of Information Science]*. Dunod, Paris, (1979). Application à l’hyperstabilité et aux processus aléatoires
- [31] Fitzgerald, C.H.: On analytic continuation to a schlicht function. *Proc. Am. Math. Soc.* **18**, 788–792 (1967)
- [32] Genin, Y., Van Dooren, P., Kailath, T., Delosme, J.M., Morf, M.: On  $\Sigma$ -lossless transfer functions and related questions. *Linear Algebra Appl.* **50**, 251–275 (1983)
- [33] Hassibi, B., Sayed, A.H., Kailath, T.: *Indefinite-quadratic estimation and control*, volume 16 of *SIAM Studies in Applied Mathematics*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, (1999). A unified approach to  $H^2$  and  $H^\infty$  theories
- [34] Kreĭn, M.G., Langer, H.: Über die verallgemeinerten Resolventen und die charakteristische Funktion eines isometrischen Operators im Raume  $\Pi_k$ . In *Hilbertspace operators and operator algebras (Proc. Int. Conf. Tihany, 1970)*, pages 353–399. North-Holland, Amsterdam, *Colloquia Math. Soc. János Bolyai* (1972)
- [35] Kreĭn, M.G., Langer, H.: Über die  $Q$ -Funktion eines  $\pi$ -hermiteschen Operators im Raume  $\pi_\kappa$ . *Acta Sci. Math. (Szeged)* **34**, 191–230 (1973)
- [36] Loewner, K.: Über monotone matrixfunktionen. *Math. Z.* **38**, 177–216 (1934)
- [37] Schwartz, L.: Sous espaces hilbertiens d’espaces vectoriels topologiques et noyaux associés (noyaux reproduisants). *J. Analyse Math.* **13**, 115–256 (1964)
- [38] Smith, M.C.: Synthesis of mechanical networks: the inerter. *IEEE Trans. Automat. Control* **47**(10), 1648–1662 (2002)

- [39] Vaidyanathan, P.P.: The discrete-time bounded-real lemma in digital filtering. *IEEE Trans. Circuits Syst.* **32**, 918–924 (1985)
- [40] Zhang, F.: Quaternions and matrices of quaternions. *Linear Algebra Appl.* **251**, 21–57 (1997)

Daniel Alpay  
Faculty of Mathematics, Physics, and Computation,  
Schmid College of Science and Technology  
Chapman University  
One University Drive  
Orange CA92866  
USA  
e-mail: [alpay@chapman.edu](mailto:alpay@chapman.edu)

Fabrizio Colombo and Irene Sabadini (✉)  
Dipartimento di Matematica  
Politecnico di Milano  
Via E. Bonardi, 9  
20133 Milan  
Italy  
e-mail: [irene.sabadini@polimi.it](mailto:irene.sabadini@polimi.it)

Fabrizio Colombo  
e-mail: [fabrizio.colombo@polimi.it](mailto:fabrizio.colombo@polimi.it)

Izchak Lewkowicz  
Department of Electrical Engineering  
Ben-Gurion University of the Negev  
P.O.B. 65384105 Beer-Sheva  
Israel  
e-mail: [izchak@ee.bgu.ac.il](mailto:izchak@ee.bgu.ac.il)

Received: October 12, 2018.