



Dyadic models for ideal MHD

Mimi Dai  and Susan Friedlander

Communicated by G. P. Galdi

Abstract. We study two dyadic models for incompressible ideal magnetohydrodynamics, one with a uni-directional energy cascade and the other one with both forward and backward energy cascades. Global existence of weak solutions and local well-posedness are established for both models. In addition, solutions to the model with uni-directional energy cascade associated with positive initial data are shown to develop blow-up at a finite time. Moreover, a set of fixed points is found for each model. Linear instability about some particular fixed points is proved.

Mathematics Subject Classification. 35Q35, 76B03, 76W05.

Keywords. Magnetohydrodynamics, Dyadic model, Energy cascade, Blow-up, Fixed points.

1. Introduction

1.1. Dyadic Euler Model

Dyadic models for hydrodynamics have a long history traced back to the first such models introduced by Desnyanskiy and Novikov [22], Gledzer [28], and Ohkitani and Yamada [36], usually referred as DN model and GOY model respectively. Following these early papers, various models [10, 18, 23, 35] were developed by simplifying the nonlinearity in different ways. Among them, the Katz–Pavlovic (KP) model [32] was derived applying Littlewood–Paley techniques to the Euler equation

$$u_t + (u \cdot \nabla)u + \nabla P = 0.$$

The KP model with external forcing is the following nonlinearly coupled infinite system of ordinary differential equations (ODEs)

$$\frac{d}{dt}a_j + \lambda_j^\theta a_j a_{j+1} - \lambda_{j-1}^\theta a_{j-1}^2 = f_j, \quad (1.1)$$

for $j \geq 0$ and $a_{-1} = 0$, with $f = (f_0, f_1, f_2, \dots)$. The quantity $\frac{1}{2}a_j^2$ stands for the energy of the j th dyadic shell with wavenumber $\lambda_j = \lambda^j$ for a constant $\lambda > 1$. The parameter θ takes the form $\theta = \frac{5-\delta}{2}$ with δ being the intermittency dimension for the turbulent fluid (cf. [13]). For 3D fluids, $\delta \in [0, 3]$ and hence $\theta \in [1, \frac{5}{2}]$. Evidently smaller δ corresponds to larger θ . Conceptually a turbulent flow with smaller intermittency dimension δ is more singular and hence exhibits stronger nonlinear effects.

The main features of (1.1) include: (i) it formally conserves the energy when $f = 0$; (ii) it only takes into account the nonlinear interactions of nearest neighboring shells through a Littlewood–Paley decomposition treatment; (iii) it is equipped with a forward energy cascade mechanism; (iv) spatial structure is oversimplified. Although model (1.1) has a seemingly simple form, analyzing it is highly non-trivial. Fortunately, two important properties play a vital role in the analysis: positivity of solutions starting from positive data and monotonicity of the rescaled quantity $\lambda_j^{\frac{1}{3}\theta} a_j$.

There is an extensive literature for the study of model (1.1), cf. [16] for a thorough description. We highlight some results from the literature on the Euler model without the intention of providing a complete list. Finite time blow-up was first shown in [32] and later sharpened in [33]. For a closely related vector model, the authors of [27] also discovered finite time blow-up. A model of a combination of DN type and Obukhov type [35] was studied in [30] and shown to exhibit finite time blow-up when the Obukhov type nonlinearity is sufficiently weak. Anomalous energy dissipation and self-similar solutions for the unforced model were studied in [6]. The problems of uniqueness, well-posedness and regularity for the inviscid model were addressed in [4, 8]. The authors of [17] proved regularizing properties of the nonlinear term of (1.1) due to the forward energy cascade mechanism. The models with stochastic forcing were also investigated in many articles, for instance, see [5, 7, 26, 38].

The results of Cheskidov, Friedlander and Pavlovic given in [14, 15] are relevant to our current paper. In [14], the authors studied (1.1) with $\theta = 1$ and positive forcing. They showed the existence of a unique positive fixed point for the dyadic Euler model (1.1) with positive initial data. They also established linear stability of the fixed point in the sense that there is no positive eigenvalue for the linearized system about the fixed point and that there exist solutions to the linearized system in H^s with $s < \frac{1}{3}$ which decay exponentially fast in time. Moreover, they proved that every solution with bounded $H^{1/3}$ norm approaches the fixed point in the energy space l^2 . Hence solutions with positive initial l^2 data blow up in finite time in $H^{1/3}$ norm. In [15] the authors further examined the properties of the unique positive fixed point and showed that the fixed point is an exponential global attractor. Via this property the authors provided another proof of finite time blowup of solutions in the norm H^s with $s \geq \frac{1}{3}$. We note that persistence of positivity of the solutions plays a crucial role in the analysis of [14, 15].

1.2. Dyadic Models for Magnetohydrodynamics

The equations for incompressible ideal (inviscid and non-resistive) magnetohydrodynamics (MHD) are the set of partial differential equations

$$u_t + (u \cdot \nabla)u - (B \cdot \nabla)B + \nabla P = 0, \tag{1.2a}$$

$$B_t + (u \cdot \nabla)B - (B \cdot \nabla)u = 0, \tag{1.2b}$$


$$\nabla \cdot u = 0, \quad \nabla \cdot B = 0. \tag{1.2c}$$

System (1.2a)–(1.2c) describes electrically conducting fluids in geophysics and astrophysics when the underlying length scales are very large, and hence the kinetic viscosity and magnetic diffusivity are negligible. In (1.2a)–(1.2c) the unknown vector fields u and B and the scalar function P denote the fluid velocity, magnetic field and pressure.

Inspired by results described above for the dyadic model (1.1) of the Euler equation, we study two particular dyadic models (suggested in [19]) associated with the PDEs (1.2a)–(1.2c) of MHD. One such model has a uni-directional energy cascade. The other model has both forward and backward energy cascade mechanisms. The model with uni-directional energy cascade is presented as

$$\begin{aligned} \frac{d}{dt}a_j &= -(\lambda_j^\theta a_j a_{j+1} - \lambda_{j-1}^\theta a_{j-1}^2) - (\lambda_j^\theta b_j b_{j+1} - \lambda_{j-1}^\theta b_{j-1}^2) + f_j, \\ \frac{d}{dt}b_j &= \lambda_j^\theta a_j b_{j+1} - \lambda_j^\theta b_j a_{j+1}, \end{aligned} \tag{1.3}$$

for $j \geq 0$ with $a_{-1} = b_{-1} = 0$ and $f = (f_0, f_1, f_2, \dots)$. For positive solutions, the forward energy cascade within the dynamics can be illustrated below



The model with both forward and backward energy cascades is

$$\begin{aligned}\frac{d}{dt}a_j &= -(\lambda_j^\theta a_j a_{j+1} - \lambda_{j-1}^\theta a_{j-1}^2) + (\lambda_j^\theta b_j b_{j+1} - \lambda_{j-1}^\theta b_{j-1}^2) + f_j, \\ \frac{d}{dt}b_j &= -(\lambda_j^\theta a_j b_{j+1} - \lambda_j^\theta b_j a_{j+1}),\end{aligned}\tag{1.4}$$

with the bi-directional energy transfer illustrated as,

$$\begin{array}{ccccccc} \cdots & \longrightarrow & a_{j-1} & \longrightarrow & a_j & \longrightarrow & a_{j+1} & \longrightarrow & \cdots \\ & & & & \uparrow & \swarrow & \uparrow & \swarrow & \uparrow \\ \cdots & \longrightarrow & b_{j-1} & \longrightarrow & b_j & \longrightarrow & b_{j+1} & \longrightarrow & \cdots \end{array}$$

The quantities $\frac{1}{2}a_j^2$ and $\frac{1}{2}b_j^2$ in (1.3) and (1.4) represent the kinetic energy and magnetic energy in the j th shell, respectively. One can see that only interactions with the nearest neighbor shells are taken into account in the modeling. Both models (1.3) and (1.4) preserve the most essential feature of the original dynamics of (1.2a)–(1.2c), namely the total energy

$$E(t) = \frac{1}{2} \sum_{j \geq 1} (a_j^2(t) + b_j^2(t))$$

is conserved formally. In addition, the cross helicity defined by

$$H^c(t) = \sum_{j=0}^{\infty} a_j b_j$$

is invariant for strong solutions of the model (1.4), but not for model (1.3).

It is obvious that eliminating b (i.e. $b_j = 0$ for $j \geq 0$) reduces systems (1.3) and (1.4) to the fluid model (1.1). Beside sharing all of the similar difficulties arising from the nonlinearity in (1.1), the MHD models (1.3) and (1.4) exhibit extra obstacles due to the interactions between the fluid and magnetic fields. In particular, it is unlikely that solutions with positive initial data stay positive for all the time in a general setting. It is also not clear whether the quantities $\lambda_j^{\frac{1}{3}\theta} a_j$ and $\lambda_j^{\frac{1}{3}\theta} b_j$ exhibit certain monotonicity. On the other hand, the conservation of positivity is not a natural property, especially in the context of MHD models. Numerical simulations suggest solutions with positive initial data can become negative.

We take a brief look at the energy transfer mechanisms and get some basic sense of the difficulties. Denote the flux by $\Pi_j = \lambda_j^\theta (a_j^2 + b_j^2) a_{j+1}$ for $j \geq 0$. The rate of change of the total energy of the j th shell for (1.3) is given by

$$\begin{aligned}\frac{1}{2} \frac{d}{dt} (a_j^2 + b_j^2) &= \lambda_{j-1}^\theta (a_{j-1}^2 + b_{j-1}^2) a_j - \lambda_j^\theta (a_j^2 + b_j^2) a_{j+1} + f_j a_j \\ &= \Pi_{j-1} - \Pi_j + f_j a_j,\end{aligned}$$

where Π_{j-1} is the flux coming from the previous shell and Π_j is the flux escaping to the next shell provided $a_j, a_{j+1} \geq 0$, and vice versa if $a_j, a_{j+1} \leq 0$. It thus represents a uni-directional energy cascade mechanism. Nevertheless, since the conservation of positivity is not valid in general setting, analyzing the energy transfer is not trivial. Moreover, there are obstacles to understand the energy flux $b_j^2 a_{j+1}$ involved with interactions between the velocity and the magnetic fields. The situation is even more subtle for system (1.4). Denote $\tilde{\Pi}_j = \lambda_j^\theta (a_j^2 - b_j^2) a_{j+1}$ for $j \geq 0$. Then for (1.4) the energy change rate of the j th shell is

$$\frac{1}{2} \frac{d}{dt} (a_j^2 + b_j^2) = \tilde{\Pi}_{j-1} - \tilde{\Pi}_j + f_j a_j.$$

We note that the sign of the flux $\tilde{\Pi}_j$ does not only depend on a_{j+1} but also on the size of $|a_j|$ and $|b_j|$. Hence it is challenging to obtain a rather comprehensive picture of the energy transfer for (1.4).

Despite the obstacles described above, the authors [21] were able to obtain results on uniqueness and non-uniqueness for the models (1.3) and (1.4) with viscous linear terms $\lambda_j^2 a_j$ and $\lambda_j^2 b_j$. More precisely,

uniqueness of Leray-Hopf solution was established for $\theta \leq 2$; while non-unique Leray-Hopf solutions were constructed for $\theta > 2$. We remark that positivity is not known to hold for the viscous dyadic MHD models. The techniques used in [21] are independent of the positivity property.

We note that dyadic models for the MHD turbulence were introduced a few decades ago in the physics community, see [11, 25, 29]. A majority of contributions, for instance in [1–3, 9], concern numerical studies of various dyadic models. In fact, one of the motivations to develop dyadic models for the MHD is that direct numerical simulations for the original dynamics face serious computing limitations. More complex dyadic models have been proposed to include non-local interactions [24] and anisotropy [34] to understand the MHD turbulence. Although the dyadic models do not preserve geometric features due to the lack of spatial structure, numerics for these models show intermittency statistics and chaotic behaviour, which are in agreement with experimental MHD turbulence. For a full background in this regard, the reader may consult the recent review article [37] and references therein.

The main objective of the present article is to examine the inviscid systems (1.3) and (1.4). Global weak solutions and local well-posedness are obtained by employing standard arguments. In the context of the dyadic MHD model (1.3), we show finite time blow-up for (1.3) by constructing an appropriate Lyapunov function. We note that it remains an open question whether a solution of (1.4) with bi-directional energy cascade develops blow-up in finite time.

A set of fixed points for (1.3) and (1.4) are found with an explicit form. Specifically, assuming f takes the form

$$f_0 > 0, \quad \text{and} \quad f_j = 0, \quad \forall \quad j \geq 1,$$

we find infinitely many steady states (\bar{a}, \bar{b}) of (1.3) in $l^2 \times l^2$ which satisfy

$$\bar{a}_j^2 + \bar{b}_j^2 = \lambda^{\frac{1}{3}\theta} f_0 \lambda_j^{-\frac{2}{3}\theta}, \quad \forall \quad j \geq 0,$$

and infinitely many steady states (\bar{a}, \bar{b}) of (1.4) satisfying

$$\bar{a}_j^2 - \bar{b}_j^2 = \lambda^{\frac{1}{3}\theta} f_0 \lambda_j^{-\frac{2}{3}\theta}, \quad \forall \quad j \geq 0.$$

In stark contrast with the dyadic Euler model studied in [14, 15], it is much harder to analyze the properties of the fixed points of systems (1.3) and (1.4). One obvious reason is that the fixed points are not unique. The more sophisticated difficulty comes from the interactions between the velocity and magnetic fields. Our understanding of the energy transfer from the j th shell of velocity to the j th shell of magnetic field (a_j to b_j) is limited at the moment. Although it is challenging to show stability or instability of the fixed points in general setting, we are able to establish linear instability results for some special fixed points. In particular, for the fixed point with zero magnetic field $(\bar{a}, 0)$, we observe that the linearized system is decoupled between the velocity and magnetic field and we prove that it is linearly unstable. More specifically, if \bar{a} is positive, the velocity component is stable while the magnetic field component is unstable; if \bar{a} is negative, both of the velocity and magnetic field are unstable.

1.3. Organisation of the Paper

- Section 2 consists of notations and definitions of solutions for the dyadic systems (1.3) and (1.4).
- Section 3 gives a proof of global existence of weak solutions for any initial data with finite energy. This result holds for both systems (1.3) and (1.4).
- Section 4 gives a proof of local well-posedness in H^s for $s \geq \theta$. This result holds for both systems (1.3) and (1.4).
- In Sect. 5 we show that the system (1.3) starting from positive initial data blows up in finite time in H^s for $s > \frac{1}{3}\theta$.
- Section 6 describes the fixed point sets for systems (1.3) and (1.4).
- In Sect. 7 we examine the linear instability for perturbations about two specific fixed points.

2. Preliminaries

2.1. Notation and Definition of Solutions

We will often use C_0 and c, c_0, c_1, \dots to denote constants which may vary from line to line. They are universal constants unless specified otherwise. We say $a = (a_0, a_1, a_2, \dots)$ is positive if all of the components are positive.

Parallel to the energy space L^2 and Sobolev space H^s for functions of space and time, we need to refer an analogue of such spaces for sequences by l^2 and H^s (the same notation), with l^2 endowed with the standard scalar product and norm

$$(u, v) := \sum_{n=1}^{\infty} u_n v_n, \quad |u| := \sqrt{(u, u)},$$

and H^s equipped with the scalar product

$$(u, v)_s := \sum_{n=1}^{\infty} \lambda_n^{2s} u_n v_n \quad \text{with } \lambda_n = \lambda^n$$

and the norm

$$\|u\|_{H^s} = \|u\|_s := \sqrt{(u, u)_s}.$$

We define the strong distance d_s and weak distance d_w on l^2 as,

$$d_s(u, v) := |u - v|, \quad d_w(u, v) := \sum_{n=1}^{\infty} \frac{1}{\lambda^{n^2}} \frac{|u_n - v_n|}{1 + |u_n - v_n|}, \quad u, v \in l^2.$$

Naturally, d_w generates a weak topology on any bounded subset of l^2 . Weak convergence of a bounded sequence $\{u^k\} \subset l^2$ to $u \in l^2$ is understood in the usual way,

$$d_w(u^k, u) \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

The functional spaces $C([0, T]; l_w^2)$ and $C([0, \infty); l_w^2)$ are defined as

$$C([0, T]; l_w^2) := \{u(\cdot) : [0, T] \rightarrow l^2, u_n(t) \text{ is continuous for all } n\}$$

endowed with the distance

$$d_{C([0, T]; l_w^2)}(u, v) := \sup_{t \in [0, T]} d_w(u(t), v(t)),$$

and

$$C([0, \infty); l_w^2) := \{u(\cdot) : [0, \infty) \rightarrow l^2, u_n(t) \text{ is continuous for all } n\}$$

with

$$d_{C([0, \infty); l_w^2)} := \sum_{T \in \mathbb{N}} \frac{1}{2^T} \frac{\sup\{d_w(u(t), v(t)) : 0 \leq t \leq T\}}{1 + \sup\{d_w(u(t), v(t)) : 0 \leq t \leq T\}}.$$

Definition 2.1. A pair of l^2 -valued functions $(a(t), b(t))$ defined on $[t_0, \infty)$ is said to be a weak solution of (1.3) or (1.4) if a_j and b_j satisfy (1.3) or (1.4) and $a_j, b_j \in C^1([t_0, \infty))$ for all $j \geq 0$.

Definition 2.2. A solution $(a(t), b(t))$ of (1.3) or (1.4) is strong on $[T_1, T_2]$ if $\|a\|_{H^1}$ and $\|b\|_{H^1}$ are bounded on $[T_1, T_2]$. A solution is strong on $[T_1, \infty)$ if it is strong on every interval $[T_1, T_2]$ for any $T_2 > T_1$.

3. Weak Solutions

In this part, we show the global existence of weak solutions for any data with finite energy by the Galerkin approximating method.

Theorem 3.1. *For any initial data $(a^0, b^0) \in l^2 \times l^2$, there exists a weak solution $(a(t), b(t))$ of system (1.3) and (1.4) on $[0, \infty)$ with $a(0) = a^0$ and $b(0) = b^0$.*

Proof. We only need to show the existence of a weak solution to system (1.3), since minor modification of the proof works for system (1.4). For any fixed integer $k \geq 1$, we consider the truncated variables $\{(a^k(t), b^k(t))\}$ given by

$$\begin{aligned} a^k(t) &= (a_0^k(t), a_1^k(t), a_2^k(t), \dots, a_k^k(t), 0, 0, \dots), \\ b^k(t) &= (b_0^k(t), b_1^k(t), b_2^k(t), \dots, b_k^k(t), 0, 0, \dots), \end{aligned}$$

with $a_j^k(0) = a_j^0$ and $b_j^k(0) = b_j^0$ for $0 \leq j \leq k$, and $\{(a^k(t), b^k(t))\}$ satisfies the finite system

$$\begin{aligned} \frac{d}{dt} a_j^k &= -\lambda_j^\theta a_j^k a_{j+1}^k + \lambda_{j-1}^\theta (a_{j-1}^k)^2 - \lambda_j^\theta b_j^k b_{j+1}^k \\ &\quad + \lambda_{j-1}^\theta (b_{j-1}^k)^2 + f_j, \quad j \leq k-1 \\ \frac{d}{dt} b_j^k &= \lambda_j^\theta a_j^k b_{j+1}^k - \lambda_j^\theta b_j^k a_{j+1}^k, \quad j \leq k-1 \\ \frac{d}{dt} a_k^k &= \lambda_{k-1}^\theta (a_{k-1}^k)^2 + \lambda_{k-1}^\theta (b_{k-1}^k)^2 + f_k, \\ \frac{d}{dt} b_k^k &= 0. \end{aligned} \tag{3.1}$$

We notice that the functions on the right hand side of (3.1) are continuous in a^k and b^k . Hence, there exists a unique solution $(a^k(t), b^k(t))$ to (3.1) on $[0, T]$ for arbitrary $T > 0$. In order to pass a subsequence of $\{(a^k(t), b^k(t))\}$ to a limit, we will apply the Ascoli-Arzelà theorem. It is sufficient to show that the sequence $\{(a^k(t), b^k(t))\}$ is weakly equicontinuous. For any fixed $k \geq 1$, there exists a constant C_0 independent of k such that

$$|a_j^k(t)| \leq C_0, \quad |b_j^k(t)| \leq C_0 \quad \forall t \in [0, T] \quad \text{and} \quad 0 \leq j \leq k. \tag{3.2}$$

It follows from (3.1) and (3.2) that

$$\begin{aligned} &|a_j^k(t) - a_j^k(s)| \\ &\leq \int_s^t |-\lambda_j^\theta a_j^k(\tau) a_{j+1}^k(\tau) + \lambda_{j-1}^\theta (a_{j-1}^k(\tau))^2 \\ &\quad - \lambda_j^\theta b_j^k(\tau) b_{j+1}^k(\tau) + \lambda_{j-1}^\theta (b_{j-1}^k(\tau))^2 + f_j| \, d\tau \\ &\leq (2\lambda_j^\theta C_0^2 + 2\lambda_{j-1}^\theta C_0^2 + f_j) |t - s| \end{aligned} \tag{3.3}$$

for $0 \leq j \leq k$ and $0 \leq t \leq s \leq T$, and similarly

$$\begin{aligned} &|b_j^k(t) - b_j^k(s)| \\ &\leq \int_s^t |\lambda_j^\theta a_j^k(\tau) b_{j+1}^k(\tau) - \lambda_j^\theta b_j^k(\tau) a_{j+1}^k(\tau)| \, d\tau \\ &\leq 2\lambda_j^\theta C_0^2 |t - s|. \end{aligned} \tag{3.4}$$

Therefore, we have from (3.3) and (3.4)

$$\begin{aligned} d_w(a^k(t), a^k(s)) &= \sum_{j=0}^{\infty} \frac{1}{\lambda^{j^2}} \frac{|a_j^k(t) - a_j^k(s)|}{1 + |a_j^k(t) - a_j^k(s)|} \leq c|t - s|, \\ d_w(b^k(t), b^k(s)) &= \sum_{j=0}^{\infty} \frac{1}{\lambda^{j^2}} \frac{|b_j^k(t) - b_j^k(s)|}{1 + |b_j^k(t) - b_j^k(s)|} \leq c|t - s|, \end{aligned}$$

for an absolute constant c . Obviously, $\{a^k\}$ and $\{b^k\}$ are weakly equicontinuous in $C([0, T]; l^2)$, and hence are relatively compact in $C([0, T]; l_w^2)$ by the Ascoli-Arzelà theorem. Therefore, there exists a subsequence $\{(a^{k_n}, b^{k_n})\}$ and a weakly continuous pair of functions $(a(t), b(t))$ in $l^2 \times l^2$ such that

$$a^{k_n} \rightharpoonup a, \quad b^{k_n} \rightharpoonup b \quad \text{as } k_n \rightarrow \infty \quad \text{in } C([0, T]; l_w^2).$$

Hence,

$$a_j^{k_n}(t) \rightarrow a_j(t), \quad b_j^{k_n}(t) \rightarrow b_j(t) \quad \text{as } k_n \rightarrow \infty \quad \text{for all } j \geq 0 \quad \text{and } t \in [0, T].$$

In particular, we have $a(0) = a^0$ and $b(0) = b^0$.

It is left to show that the limit (a, b) is a solution of (1.3). By (3.1), we have

$$\begin{aligned} a_j^{k_n}(t) &= a_j^{k_n}(0) + \int_0^t \left(-\lambda_j^\theta a_j^{k_n}(\tau) a_{j+1}^{k_n}(\tau) + \lambda_{j-1}^\theta (a_{j-1}^{k_n}(\tau))^2 \right) d\tau \\ &\quad + \int_0^t \left(-\lambda_j^\theta b_j^{k_n}(\tau) b_{j+1}^{k_n}(\tau) + \lambda_{j-1}^\theta (b_{j-1}^{k_n}(\tau))^2 + f_j \right) d\tau, \\ b_j^{k_n}(t) &= b_j^{k_n}(0) + \int_0^t \left(\lambda_j^\theta a_j^{k_n}(\tau) b_{j+1}^{k_n}(\tau) - \lambda_j^\theta b_j^{k_n}(\tau) a_{j+1}^{k_n}(\tau) \right) d\tau, \end{aligned}$$

for $0 \leq j \leq k_n - 1$. After taking the limit $k_n \rightarrow \infty$ we get

$$\begin{aligned} a_j(t) &= a_j(0) + \int_0^t \left(-\lambda_j^\theta a_j(\tau) a_{j+1}(\tau) + \lambda_{j-1}^\theta (a_{j-1}(\tau))^2 \right) d\tau \\ &\quad + \int_0^t \left(-\lambda_j^\theta b_j(\tau) b_{j+1}(\tau) + \lambda_{j-1}^\theta (b_{j-1}(\tau))^2 + f_j \right) d\tau, \\ b_j(t) &= b_j(0) + \int_0^t \left(\lambda_j^\theta a_j(\tau) b_{j+1}(\tau) - \lambda_j^\theta b_j(\tau) a_{j+1}(\tau) \right) d\tau. \end{aligned}$$

Furthermore, since a_j and b_j are continuous for all $j \geq 0$, the integral form implies that $a_j \in C^1[0, T]$ and $b_j \in C^1[0, T]$. Thus (a, b) solves (1.3). \square

4. Local Well-Posedness

System (1.3) and (1.4) are locally well-posed in H^s with $s \geq \theta$. Namely, we prove:

Theorem 4.1. *Let $s \geq \theta$. Assume $(a^0, b^0) \in H^s \times H^s$ and $f \in H^s$. There exists a time $T > 0$ such that system (1.3) with initial data (a^0, b^0) has a unique solution $(a(t), b(t))$ in $H^s \times H^s$ on $[0, T]$.*

Proof. The local existence follows from a standard argument and a priori estimate for solutions in H^s . We only show the a priori estimate here. Multiplying the a_j equation of (1.3) by $\lambda_j^{2s} a_j$, the b_j equation by $\lambda_j^{2s} b_j$, and taking the sum over $j \geq 0$, we obtain

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \sum_{j=0}^{\infty} (\lambda_j^{2s} a_j^2 + \lambda_j^{2s} b_j^2) &= (\lambda^{2s} - 1) \sum_{j=0}^{\infty} \lambda_j^{\theta+2s} a_j^2 a_{j+1} \\ &\quad + (\lambda^{2s} - 1) \sum_{j=0}^{\infty} \lambda_j^{\theta+2s} b_j^2 a_{j+1} + \sum_{j=0}^{\infty} \lambda_j^{2s} a_j f_j. \end{aligned}$$

Applying Cauchy-Schwartz's inequality gives

$$\begin{aligned} \sum_{j=0}^{\infty} \lambda_j^{\theta+2s} a_j^2 a_{j+1} &= \lambda^{-\frac{1}{3}(\theta+2s)} \sum_{j=0}^{\infty} \lambda_j^{\frac{2}{3}(\theta+2s)} a_j^2 \cdot \lambda_j^{\frac{1}{3}(\theta+2s)} a_{j+1} \\ &\leq \frac{2}{3} \lambda^{-\frac{1}{3}(\theta+2s)} \sum_{j=0}^{\infty} \lambda_j^{\theta+2s} a_j^3 + \frac{1}{3} \lambda^{-\frac{1}{3}(\theta+2s)} \sum_{j=0}^{\infty} \lambda_{j+1}^{\theta+2s} a_{j+1}^3 \\ &\leq \lambda^{-\frac{1}{3}(\theta+2s)} \left(\sum_{j=0}^{\infty} \lambda_j^{\frac{2}{3}(\theta+2s)} a_j^2 \right)^{\frac{3}{2}}, \end{aligned}$$

and similarly

$$\begin{aligned} \sum_{j=0}^{\infty} \lambda_j^{\theta+2s} b_j^2 a_{j+1} &= \lambda^{-\frac{1}{3}(\theta+2s)} \sum_{j=0}^{\infty} \lambda_j^{\frac{2}{3}(\theta+2s)} b_j^2 \cdot \lambda_j^{\frac{1}{3}(\theta+2s)} a_{j+1} \\ &\leq \frac{2}{3} \lambda^{-\frac{1}{3}(\theta+2s)} \sum_{j=0}^{\infty} \lambda_j^{\theta+2s} b_j^3 + \frac{1}{3} \lambda^{-\frac{1}{3}(\theta+2s)} \sum_{j=0}^{\infty} \lambda_{j+1}^{\theta+2s} a_{j+1}^3 \\ &\leq \frac{2}{3} \lambda^{-\frac{1}{3}(\theta+2s)} \left(\sum_{j=0}^{\infty} \lambda_j^{\frac{2}{3}(\theta+2s)} b_j^2 \right)^{\frac{3}{2}} \\ &\quad + \frac{1}{3} \lambda^{-\frac{1}{3}(\theta+2s)} \left(\sum_{j=0}^{\infty} \lambda_j^{\frac{2}{3}(\theta+2s)} a_j^2 \right)^{\frac{3}{2}}. \end{aligned}$$

We also have

$$\sum_{j=0}^{\infty} \lambda_j^{2s} a_j f_j \leq \sum_{j=0}^{\infty} \lambda_j^{2s} a_j^2 + \sum_{j=0}^{\infty} \lambda_j^{2s} f_j^2.$$

Therefore, putting the last four estimates together yields

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \sum_{j=0}^{\infty} (\lambda_j^{2s} a_j^2 + \lambda_j^{2s} b_j^2) &\leq \frac{4}{3} (\lambda^{2s} - 1) \lambda^{-\frac{1}{3}(\theta+2s)} \left(\sum_{j=0}^{\infty} \lambda_j^{\frac{2}{3}(\theta+2s)} a_j^2 \right)^{\frac{3}{2}} \\ &\quad + \frac{2}{3} (\lambda^{2s} - 1) \lambda^{-\frac{1}{3}(\theta+2s)} \left(\sum_{j=0}^{\infty} \lambda_j^{\frac{2}{3}(\theta+2s)} b_j^2 \right)^{\frac{3}{2}} \\ &\quad + \sum_{j=0}^{\infty} \lambda_j^{2s} a_j^2 + \sum_{j=0}^{\infty} \lambda_j^{2s} f_j^2. \end{aligned}$$

Since $s \geq \theta$, we have $\frac{2}{3}(\theta + 2s) \leq 2s$ and hence

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \sum_{j=0}^{\infty} (\lambda_j^{2s} a_j^2 + \lambda_j^{2s} b_j^2) &\leq C(\lambda, s, \theta) \left(\sum_{j=0}^{\infty} (\lambda_j^{2s} a_j^2 + \lambda_j^{2s} b_j^2) \right)^{\frac{3}{2}} \\ &\quad + \sum_{j=0}^{\infty} \lambda_j^{2s} a_j^2 + \sum_{j=0}^{\infty} \lambda_j^{2s} f_j^2 \end{aligned}$$

for a constant $C(\lambda, s, \theta)$. Thus, for $f \in H^s$, there exists a time $T > 0$ and another constant $C(\lambda, s, \theta, f)$ such that

$$\|a(t)\|_{H^s}^2 + \|b(t)\|_{H^s}^2 \leq C(\lambda, s, \theta, f) (\|a(0)\|_{H^s}^2 + \|b(0)\|_{H^s}^2), \quad t \in [0, T].$$

The uniqueness can be obtained by the classical approach, i.e. establishing a Gröwall's inequality for the difference of two solutions starting from the same data. We omit the details here. \square

Theorem 4.2. *Let $s \geq \theta$. Assume $(a^0, b^0) \in H^s \times H^s$ and $f \in H^s$. There exists a time $T > 0$ such that system (1.4) with initial data (a^0, b^0) has a unique solution $(a(t), b(t))$ in $H^s \times H^s$ on $[0, T]$.*

The proof follows similarly as that of Theorem 4.1.

Remark 4.3. Recall $\theta = \frac{5-\delta}{2}$ and $\delta \in [0, 3]$. When $\theta = \frac{5}{2}$ ($\delta = 0$) the local well-posedness of (1.3) and (1.4) holds in H^s with $s \geq \frac{5}{2}$ which is consistent with the well-posedness result for the ideal MHD system. On the other hand, if $\theta = 1$ ($\delta = 3$), (1.3) and (1.4) are locally well-posed in H^s with $s \geq 1$.

5. Finite Time Blow-up

In this section we show that a solution of the dyadic ideal MHD model (1.3) starting from positive initial data develops blow-up at a finite time in the space H^s with $s > \frac{1}{3}\theta$. For dyadic models, we are aware of two approaches of proving finite time blow-up. For the forced dyadic Euler system the authors of [15] made use of properties of the fixed point and provided a dynamical approach to show the development of blow-up. Another method is to construct a suitable Lyapunov function for the underlying system that blows up at finite time, for instance, see [20, 30]. In particular, the model studied in [30] contains both the DN type and Obukov type of nonlinearities, and exhibits both forward and backward energy cascade mechanisms. Consequently, the solutions considered there are not known to be positive. However, the authors demonstrated blow-up by developing a special Lyapunov function approach. Their approach has certain flexibility and is robust. In the context of model (1.3), the interactions of the fluid and magnetic field make it harder to explore the energy transfer from shell to shell; as a result, a dynamical approach to show blow-up is out of reach. Therefore we choose to take the path of a Lyapunov function. Nevertheless, it seems quite challenging to construct finite time blow-up solution for system (1.4) by applying the aforementioned approaches. In fact, the mechanism of both forward and backward energy cascades may prevent finite time blow-up.

The result of blow-up for (1.3) is stated below.

Theorem 5.1. *Let $\theta > 0$ and $f_0 \geq 0$. The solution $(a(t), b(t))$ of (1.3) with positive initial data develops blow-up at a finite time in the H^s norm with $s > \frac{1}{3}\theta$.*

Proof. The plan is to show that the quantity

$$E_s(t) := \|a(t)\|_{H^s}^2 + \|b(t)\|_{H^s}^2$$

with $s > \frac{1}{3}\theta$ is not locally integrable, which will be achieved through a contradiction argument.

We introduce the rescaled variables $w_j = \lambda_j^\theta a_j$ and $z_j = \lambda_j^\theta b_j$. It follows from (1.3) that $w = (w_0, w_1, w_2, \dots)$ and $z = (z_0, z_1, z_2, \dots)$ satisfy

$$\begin{aligned} w'_j &= -\lambda^{-\theta} w_j w_{j+1} + \lambda^\theta w_{j-1}^2 - \lambda^{-\theta} z_j z_{j+1} + \lambda^\theta z_{j-1}^2 + \lambda_j^\theta f_j, \\ z'_j &= \lambda^{-\theta} w_j z_{j+1} - \lambda^{-\theta} z_j w_{j+1}, \end{aligned} \tag{5.1}$$

for $j \geq 0$, with $w_{-1} = z_{-1} = 0$, $f_0 > 0$ and $f_j = 0$ for $j \geq 1$. Consider the quantities

$$\begin{aligned} \phi(t) &= \sum_{j=0}^{\infty} (\lambda_j^{-\gamma} w_j^2(t) + \lambda_j^{-\gamma} z_j^2(t)), \\ \psi(t) &= \sum_{j=0}^{\infty} (\lambda_j^{-\gamma} w_j(t) + c_0 \lambda_j^{-\gamma} z_j(t)), \end{aligned}$$

for appropriate constants $\gamma > 0$ and $0 < c_0 < 1$ to be specified later.

Assume $E_s(t)$ is locally integrable for $s > \frac{1}{3}\theta$. Notice that

$$0 < \phi(t) = \sum_{j=0}^{\infty} (\lambda_j^{2\theta-\gamma} a_j^2(t) + \lambda_j^{2\theta-\gamma} b_j^2(t)) \leq E_s(t)$$

provided

$$2\theta - \gamma \leq 2s, \tag{5.2}$$

and hence $\phi(t)$ is locally integrable. We will proceed to show that $\psi(t)$ is locally integrable as well and in the same time it satisfies a Riccati type of inequality for an appropriate value of γ , which leads to an obvious contradiction.

First of all, by Cauchy-Schwarz's inequality, we have

$$\begin{aligned} \psi^2(t) &= \left(\sum_{j=0}^{\infty} (\lambda_j^{-\gamma} w_j(t) + c_0 \lambda_j^{-\gamma} z_j(t)) \right)^2 \\ &\leq \left(\sum_{j=0}^{\infty} \lambda_j^{-\gamma} \right) \sum_{j=0}^{\infty} \lambda_j^{-\gamma} (w_j(t) + c_0 z_j(t))^2 \\ &\leq \frac{2}{1 - \lambda^{-\gamma}} \sum_{j=0}^{\infty} \lambda_j^{-\gamma} (w_j^2(t) + c_0^2 z_j^2(t)) \\ &\leq \frac{2c_0^2 \phi(t)}{1 - \lambda^{-\gamma}}. \end{aligned} \tag{5.3}$$

As an immediate consequence, $\psi(t)$ is locally integrable.

On the other hand, straightforward computation based on (5.1) shows

$$\begin{aligned} \frac{d}{dt} \psi(t) &= - \sum_{j=0}^{\infty} \lambda^{-\theta} \lambda_j^{-\gamma} w_j w_{j+1} + \sum_{j=0}^{\infty} \lambda^\theta \lambda_j^{-\gamma} w_{j-1}^2 \\ &\quad - \sum_{j=0}^{\infty} \lambda^{-\theta} \lambda_j^{-\gamma} z_j z_{j+1} + \sum_{j=0}^{\infty} \lambda^\theta \lambda_j^{-\gamma} z_{j-1}^2 \\ &\quad + c_0 \sum_{j=0}^{\infty} \lambda^{-\theta} \lambda_j^{-\gamma} w_j z_{j+1} - c_0 \sum_{j=0}^{\infty} \lambda^{-\theta} \lambda_j^{-\gamma} z_j w_{j+1} + f_0. \end{aligned} \tag{5.4}$$

Since $\phi(t)$ is locally integrable, it is finite almost everywhere. Notice that the infinite sums on the right hand side of (5.4) are in the order of $\phi(t)$ and hence are defined almost everywhere. It is clear that

$$\sum_{j=0}^{\infty} \lambda^{\theta} \lambda_j^{-\gamma} w_{j-1}^2 + \sum_{j=0}^{\infty} \lambda^{\theta} \lambda_j^{-\gamma} z_{j-1}^2 = \lambda^{\theta-\gamma} \phi(t). \tag{5.5}$$

Applying Cauchy-Schwarz's inequality, we estimate the infinite sums as follows

$$\begin{aligned} \sum_{j=0}^{\infty} \lambda^{-\theta} \lambda_j^{-\gamma} w_j w_{j+1} &= \lambda^{\frac{\gamma}{2}-\theta} \sum_{j=0}^{\infty} \left(\lambda_j^{-\frac{\gamma}{2}} w_j \right) \left(\lambda_{j+1}^{-\frac{\gamma}{2}} w_{j+1} \right) \\ &\leq \lambda^{\frac{\gamma}{2}-\theta} \left(\sum_{j=0}^{\infty} \lambda_j^{-\gamma} w_j^2 \right)^{\frac{1}{2}} \left(\sum_{j=0}^{\infty} \lambda_{j+1}^{-\gamma} w_{j+1}^2 \right)^{\frac{1}{2}}, \end{aligned}$$

and similarly

$$\begin{aligned} \sum_{j=0}^{\infty} \lambda^{-\theta} \lambda_j^{-\gamma} z_j z_{j+1} &\leq \lambda^{\frac{\gamma}{2}-\theta} \left(\sum_{j=0}^{\infty} \lambda_j^{-\gamma} z_j^2 \right)^{\frac{1}{2}} \left(\sum_{j=0}^{\infty} \lambda_{j+1}^{-\gamma} z_{j+1}^2 \right)^{\frac{1}{2}}, \\ \sum_{j=0}^{\infty} \lambda^{-\theta} \lambda_j^{-\gamma} z_j w_{j+1} &\leq \lambda^{\frac{\gamma}{2}-\theta} \left(\sum_{j=0}^{\infty} \lambda_j^{-\gamma} z_j^2 \right)^{\frac{1}{2}} \left(\sum_{j=0}^{\infty} \lambda_{j+1}^{-\gamma} w_{j+1}^2 \right)^{\frac{1}{2}}, \\ - \sum_{j=0}^{\infty} \lambda^{-\theta} \lambda_j^{-\gamma} w_j z_{j+1} &\leq \lambda^{\frac{\gamma}{2}-\theta} \left(\sum_{j=0}^{\infty} \lambda_j^{-\gamma} w_j^2 \right)^{\frac{1}{2}} \left(\sum_{j=0}^{\infty} \lambda_{j+1}^{-\gamma} z_{j+1}^2 \right)^{\frac{1}{2}}. \end{aligned}$$

Combining the last four inequalities yields

$$\begin{aligned} &\sum_{j=0}^{\infty} \lambda^{-\theta} \lambda_j^{-\gamma} w_j w_{j+1} + \sum_{j=0}^{\infty} \lambda^{-\theta} \lambda_j^{-\gamma} z_j z_{j+1} \\ &\quad - c_0 \sum_{j=0}^{\infty} \lambda^{-\theta} \lambda_j^{-\gamma} w_j z_{j+1} + c_0 \sum_{j=0}^{\infty} \lambda^{-\theta} \lambda_j^{-\gamma} z_j w_{j+1} \\ &\leq (1 + 2c_0) \lambda^{\frac{\gamma}{2}-\theta} \phi(t). \end{aligned} \tag{5.6}$$

Since $a_j(t) > 0$ and $b_j(t) > 0$ for all $j \geq 0$ and $t \geq 0$, it is also true that $w_j(t) > 0$ and $z_j(t) > 0$ for all $j \geq 0$ and $t \geq 0$. Therefore, we deduce from (5.4), (5.5), (5.6) and (5.3) that

$$\begin{aligned} \frac{d}{dt} \psi(t) &\geq \left(\lambda^{\theta-\gamma} - (1 + 2c_0) \lambda^{\frac{\gamma}{2}-\theta} \right) \phi(t) + f_0 \\ &\geq \frac{1 - \lambda^{-\gamma}}{2c_0^2} \left(\lambda^{\theta-\gamma} - (1 + 2c_0) \lambda^{\frac{\gamma}{2}-\theta} \right) \psi^2(t) + f_0 \end{aligned}$$

provided $\lambda^{\theta-\gamma} - (1 + 2c_0) \lambda^{\frac{\gamma}{2}-\theta} > 0$. It follows from the Riccati type of inequality that $\psi(t)$ becomes infinity at a finite time if

$$\frac{1 - \lambda^{-\gamma}}{2c_0^2} \left(\lambda^{\theta-\gamma} - (1 + 2c_0) \lambda^{\frac{\gamma}{2}-\theta} \right) \psi^2(0) + f_0 > 0. \tag{5.7}$$

Therefore, the contradiction is achieved.

In the end, we summarize all the conditions on the parameters

$$2\theta - \gamma \leq 2s, \quad \lambda^{\theta-\gamma} - (1 + 2c_0) \lambda^{\frac{\gamma}{2}-\theta} > 0.$$

In order to have the latter one satisfied, we can choose

$$\theta - \gamma > \frac{\gamma}{2} - \theta, \quad 0 < c_0 < \frac{1}{2}\lambda^{2\theta - \frac{3}{2}\gamma} - \frac{1}{2}.$$

Thus, γ can be chosen as

$$2\theta - 2s \leq \gamma < \frac{4}{3}\theta.$$

Notice that $[2\theta - 2s, \frac{4}{3}\theta)$ is not empty given the assumption $s > \frac{1}{3}\theta$. We also note condition (5.7) is satisfied for any positive initial data and forcing $f_0 \geq 0$. \square

We note the importance of the critical threshold index $\frac{1}{3}\theta$ in terms of the energy equality. For the original ideal MHD (1.2a)–(1.2c), it was proved in [12] and [31] that solutions conserving energy need to have the minimum regularity of $\frac{1}{3}$, which is analogous to Onsager’s critical regularity index for the Euler equation. We note that $\frac{1}{3}\theta = \frac{5}{6}$ for $\theta = \frac{5}{2}$ (and $\delta = 0$); and the space $H^{\frac{5}{6}}$ has the same scaling as $B_{3,\infty}^{\frac{1}{3}}$ and $B_{3,c(\mathbb{N})}^{\frac{1}{3}}$ in three dimensions, which have Onsager’s critical scaling proved in [12] and [31].

6. The Set of Fixed Points

We consider (1.3) and (1.4) with the forcing $f = (f_0, f_1, f_2, \dots)$ satisfying

$$f_0 > 0; \quad f_j = 0, \quad j \geq 1.$$

We will show the existence of fixed points of both systems (1.3) and (1.4); moreover, we provide an explicit form of them.

In order to simplify notations, we will rescale the stationary system of (1.3) and (1.4). Define

$$A_j = \lambda^{-\frac{1}{6}\theta} f_0^{-\frac{1}{2}} \lambda_j^{\frac{1}{3}\theta} a_j, \quad B_j = \lambda^{-\frac{1}{6}\theta} f_0^{-\frac{1}{2}} \lambda_j^{\frac{1}{3}\theta} b_j.$$

Let $\frac{d}{dt}a_j = \frac{d}{dt}b_j = 0$ for all $j \geq 0$. System (1.3) reduces to

$$A_{j-1}^2 - A_j A_{j+1} + B_{j-1}^2 - B_j B_{j+1} = 0, \quad j \geq 1, \tag{6.1a}$$

$$A_j B_{j+1} - B_j A_{j+1} = 0, \quad j \geq 1, \tag{6.1b}$$

$$-A_0 A_1 - B_0 B_1 = -1, \tag{6.1c}$$

$$A_0 B_1 - B_0 A_1 = 0. \tag{6.1d}$$

Lemma 6.1. *The solutions $\{(A_j, B_j)\}$ of (6.1a)–(6.1d) satisfy*

$$A_j = A_0, \quad B_j = B_0, \quad \forall j \geq 1,$$

for some constants A_0 and B_0 with the constraint

$$A_0^2 + B_0^2 = 1.$$

Proof. The two Eqs. (6.1b) and (6.1d) imply

$$\frac{A_{j+1}}{A_j} = \frac{B_{j+1}}{B_j} = c_j, \quad \forall j \geq 0, \tag{6.2}$$

for certain constant c_j . It follows that $A_{j+1} = c_j A_j$ and $B_{j+1} = c_j B_j$ for all $j \geq 0$. Then (6.1a) can be rewritten as

$$(A_j^2 + B_j^2) (c_{j-1}^{-2} - c_j) = 0, \quad \forall j \geq 1. \tag{6.3}$$

It follows

$$c_j = c_{j-1}^{-2}, \quad \forall j \geq 1.$$

As a consequence of the recursive relation, we have

$$c_j = c_0^{(-2)^j}, \quad j \geq 0. \quad (6.4)$$

Therefore, (6.2) together with (6.4) gives rise to

$$\begin{aligned} A_{j+1} &= c_j A_j = \cdots = c_j c_{j-1} \cdots c_1 c_0 A_0 \\ &= c_0^{(-2)^j + (-2)^{j-1} + \cdots + (-2) + 1} A_0 \\ &= c_0^{(1 - (-2)^{j+1})/3} A_0, \end{aligned}$$

and

$$B_{j+1} = c_0^{(1 - (-2)^{j+1})/3} B_0.$$

We further simplify the previous form and obtain

$$A_{j+1} = \begin{cases} c_0^{\frac{1+2^{j+1}}{3}} A_0, & j \text{ is even,} \\ c_0^{\frac{1-2^{j+1}}{3}} A_0, & j \text{ is odd,} \end{cases} \quad B_{j+1} = \begin{cases} c_0^{\frac{1+2^{j+1}}{3}} B_0, & j \text{ is even,} \\ c_0^{\frac{1-2^{j+1}}{3}} B_0, & j \text{ is odd.} \end{cases} \quad (6.5)$$

Consequently, we claim that c_0 has to be 1, since the sequences $\{a_j\}$ and $\{b_j\}$ are bounded. Thus, (6.4) implies that $c_j = 1$ for all $j \geq 1$. Hence, $A_j = A_0$ and $B_j = B_0$ for all $j \geq 1$. In addition, it follows from (6.1c) that

$$A_0^2 + B_0^2 = 1.$$

It concludes the proof of the lemma. □

With the same rescaling as above, the stationary system of (1.4) can be written as

$$A_{j-1}^2 - A_j A_{j+1} - B_{j-1}^2 + B_j B_{j+1} = 0, \quad j \geq 1, \quad (6.6a)$$

$$-A_j B_{j+1} + B_j A_{j+1} = 0, \quad j \geq 1, \quad (6.6b)$$

$$-A_0 A_1 + B_0 B_1 = -1, \quad (6.6c)$$

$$-A_0 B_1 + B_0 A_1 = 0. \quad (6.6d)$$

Lemma 6.2. *The solutions $\{(A_j, B_j)\}$ of (6.6a)–(6.6d) satisfy*

$$A_j = A_0, \quad B_j = B_0, \quad \forall j \geq 1,$$

for some constants A_0 and B_0 with the constraint

$$A_0^2 - B_0^2 = 1.$$

Proof. The proof is analogous as that of Lemma 6.1. It follows from Eqs. (6.6b) and (6.6d) that

$$A_{j+1} = c_j A_j, \quad B_{j+1} = c_j B_j$$

for certain constant c_j and all $j \geq 0$. Thus we obtain from (6.6a) that

$$(A_j^2 - B_j^2) (c_{j-1}^{-2} - c_j) = 0, \quad \forall j \geq 1. \quad (6.7)$$

Therefore,

$$c_j = c_{j-1}^{-2}, \quad \text{or} \quad A_j^2 - B_j^2 = 0 \quad \forall j \geq 1.$$

Thanks to (6.6b) and (6.6d), we observe if $A_j^2 - B_j^2 = 0$ for one j , it has to be true for all $j \geq 0$. However, (6.6c) implies $c_0(A_0^2 - B_0^2) = 1$. Consequently, $A_0^2 - B_0^2 \neq 0$ and hence $A_j^2 - B_j^2 \neq 0$ for all $j \geq 0$. We conclude the only possibility is $c_j = c_{j-1}^{-2}$ for all $j \geq 0$. It then follows from the same argument from the proof of Lemma 6.1 that $c_j = 1$ for all $j \geq 1$. Thus we have $A_j = A_0$ and $B_j = B_0$ for all $j \geq 1$. In the end, (6.6c) implies that

$$A_0^2 - B_0^2 = 1. \quad \square$$

7. Linear Instability

This section concerns the long time behavior of solutions, specifically, whether a solution of (1.3) converges to a fixed point or not. Evidently, from Lemmas 6.1 and 6.2, we know that a fixed point of (1.3) or (1.4) is not unique. Without loss of generality, we choose $f_0 = \lambda^{-\frac{1}{3}\theta}$ such that $\lambda^{\frac{1}{6}\theta} f_0^{\frac{1}{2}} = 1$; thus, the steady state (\bar{a}, \bar{b}) can be written as

$$\bar{a}_j = A_0 \lambda_j^{-\frac{1}{3}\theta}, \quad \bar{b}_j = B_0 \lambda_j^{-\frac{1}{3}\theta}, \quad \forall j \geq 0. \tag{7.1}$$

Consider the perturbation of the steady state,

$$\begin{aligned} a_j(t) &= A_0 \lambda_j^{-\frac{1}{3}\theta} + \epsilon \omega_j(t), \quad j \geq 0, \\ b_j(t) &= B_0 \lambda_j^{-\frac{1}{3}\theta} + \epsilon \zeta_j(t), \quad j \geq 0. \end{aligned} \tag{7.2}$$

Based on (1.3), we derive the equations for ω_j and ζ_j

$$\begin{aligned} \omega'_j &= A_0 \lambda_j^{\frac{2}{3}\theta} \left(2\lambda^{-\frac{2}{3}\theta} \omega_{j-1} - \lambda^{-\frac{1}{3}\theta} \omega_j - \omega_{j+1} \right) \\ &\quad + B_0 \lambda_j^{\frac{2}{3}\theta} \left(2\lambda^{-\frac{2}{3}\theta} \zeta_{j-1} - \lambda^{-\frac{1}{3}\theta} \zeta_j - \zeta_{j+1} \right) \\ &\quad - \epsilon \lambda_j^\theta \omega_j \omega_{j+1} + \epsilon \lambda_{j-1}^\theta \omega_{j-1}^2 - \epsilon \lambda_j^\theta \zeta_j \zeta_{j+1} + \epsilon \lambda_{j-1}^\theta \zeta_{j-1}^2, \\ \zeta'_j &= B_0 \lambda_j^{\frac{2}{3}\theta} \left(\lambda^{-\frac{1}{3}\theta} \omega_j - \omega_{j+1} \right) - A_0 \lambda_j^{\frac{2}{3}\theta} \left(\lambda^{-\frac{1}{3}\theta} \zeta_j - \zeta_{j+1} \right) \\ &\quad + \epsilon \lambda_j^\theta \omega_j \zeta_{j+1} - \epsilon \lambda_j^\theta \zeta_j \omega_{j+1} \end{aligned} \tag{7.3}$$

for $j \geq 0$ and $\omega_{-1} = \zeta_{-1} = 0$. We focus on the two special cases:

- (i) $A_0 = 1, B_0 = 0$: the fixed point corresponds to the fixed point of the dyadic Euler equation. We show linear instability with stable velocity component and unstable magnetic field.
- (ii) $A_0 = -1, B_0 = 0$: the fixed point also corresponds to a steady state of the dyadic Euler equation. We show linear instability with both unstable velocity and unstable magnetic field.

The main results described in (i) and (ii) are presented below.

Theorem 7.1. *Let (\bar{a}, \bar{b}) be the fixed point of (1.3) with $A_0 = 1$ and $B_0 = 0$ (hence $\bar{b} = 0$). For the linearized system about (\bar{a}, \bar{b}) , there are no positive eigenvalues corresponding to the velocity component; while every real number is an eigenvalue for the magnetic field linearized equation.*

Proof. When $A_0 = 1$ and $B_0 = 0$, it follows from (7.3) that the linearized system is

$$\omega'_j = \lambda_j^{\frac{2}{3}\theta} \left(2\lambda^{-\frac{2}{3}\theta} \omega_{j-1} - \lambda^{-\frac{1}{3}\theta} \omega_j - \omega_{j+1} \right) \tag{7.4a}$$

$$\zeta'_j = -\lambda_j^{\frac{2}{3}\theta} \left(\lambda^{-\frac{1}{3}\theta} \zeta_j - \zeta_{j+1} \right) \tag{7.4b}$$

for $j \geq 0$ and $\omega_{-1} = \zeta_{-1} = 0$. Notice that ω_j and ζ_j are decoupled in the linearized system, and (7.4a) is the linearized equation of dyadic Euler model about the fixed point $\bar{a}_j = \lambda_j^{-\frac{1}{3}\theta}$. We will apply the continued fraction approach from [14] to (7.4a)–(7.4b). We look for a solution to (7.4a) and (7.4b) in the form

$$\omega_j = c_j e^{pt}, \quad \zeta_j = d_j e^{qt}$$

with real values of p and q . Inserting the form $\omega_j = c_j e^{pt}$ to (7.4a) yields

$$c_{j+1} + \left(p \lambda_j^{-\frac{2}{3}\theta} + \lambda^{-\frac{1}{3}\theta} \right) c_j - 2\lambda^{-\frac{2}{3}\theta} c_{j-1} = 0 \tag{7.5}$$

for $j \geq 0$, with $c_{-1} = 0$. Inserting $\zeta_j = d_j e^{qt}$ to (7.4b) gives

$$d_{j+1} = \left(q \lambda_j^{-\frac{2}{3}\theta} + \lambda^{-\frac{1}{3}\theta} \right) d_j, \quad j \geq 0 \tag{7.6}$$

with $d_{-1} = 0$.

Following the arguments of [14], one can deduce from the iterative equation (7.5) that it is impossible to have $p \geq 0$. Therefore, for a real and negative p , we can choose $c_0 = 1$, and hence

$$c_j = \left(p\lambda_{j-1}^{-\frac{2}{3}\theta} + \lambda^{-\frac{1}{3}\theta} \right) \left(p\lambda_{j-2}^{-\frac{2}{3}\theta} + \lambda^{-\frac{1}{3}\theta} \right) \cdots \left(p\lambda_1^{-\frac{2}{3}\theta} + \lambda^{-\frac{1}{3}\theta} \right) \lambda^{-\frac{1}{3}\theta}.$$

It implies that

$$c_j \sim \lambda_j^{-\frac{1}{3}\theta}, \quad \forall j \geq J \text{ for some } J > 0.$$

Therefore, the sequence $\{c_j\}$ has finite H^s norm for $s < \frac{1}{3}\theta$, and in particular has finite l^2 norm.

On the other hand, it follows from (7.6) that

$$d_{j+1} = \alpha_j \alpha_{j-1} \alpha_0 d_0, \quad \text{with } \alpha_j = q\lambda_j^{-\frac{2}{3}\theta} + \lambda^{-\frac{1}{3}\theta}.$$

As a consequence, we know

$$\lim_{j \rightarrow \infty} \alpha_j = \lambda^{-\frac{1}{3}\theta}, \quad \lim_{j \rightarrow \infty} d_j = 0, \quad (7.7)$$

which hold for any $q \in \mathbb{R}$. In view of (7.12), we also have

$$d_j \sim \lambda_j^{-\frac{1}{3}\theta}, \quad \text{for large } j.$$

Thus the sequence $\{c_j\}$ has finite H^s norm for $s < \frac{1}{3}\theta$. Hence the H^s norm of $\{\zeta_j\}$ is finite for all the time if $q \leq 0$ and grows exponentially if $q > 0$. \square

Theorem 7.2. *Let (\bar{a}, \bar{b}) be the fixed point of (1.3) with $A_0 = -1$ and $B_0 = 0$. There are no negative eigenvalue to the velocity equation in the linearized system about such (\bar{a}, \bar{b}) . Every real number is an eigenvalue for the magnetic field equation in the linearized system. Thus the linearized system is unstable.*

Proof. Taking $A_0 = -1$ and $B_0 = 0$ in (7.3) gives the linearized system

$$\begin{aligned} \omega'_j &= -\lambda_j^{\frac{2}{3}\theta} \left(2\lambda^{-\frac{2}{3}\theta} \omega_{j-1} - \lambda^{-\frac{1}{3}\theta} \omega_j - \omega_{j+1} \right) \\ \zeta'_j &= \lambda_j^{\frac{2}{3}\theta} \left(\lambda^{-\frac{1}{3}\theta} \zeta_j - \zeta_{j+1} \right) \end{aligned} \quad (7.8)$$

Similarly, looking for solutions of the linearized system in the form

$$\omega_j = c_j e^{pt}, \quad \zeta_j = d_j e^{qt},$$

we have

$$c_{j+1} + \left(\lambda^{-\frac{1}{3}\theta} - p\lambda_j^{-\frac{2}{3}\theta} \right) c_j - 2\lambda^{-\frac{2}{3}\theta} c_{j-1} = 0, \quad (7.9)$$

$$d_{j+1} = \left(\lambda^{-\frac{1}{3}\theta} - q\lambda_j^{-\frac{2}{3}\theta} \right) d_j, \quad (7.10)$$

for $j \geq 0$. Now denote $\alpha_j = \lambda^{-\frac{1}{3}\theta} - p\lambda_j^{-\frac{2}{3}\theta}$. Following the analysis of [14], we infer from (7.9) that for $j \geq 1$

$$p\lambda_j^{-\frac{2}{3}\theta} - \lambda^{-\frac{1}{3}\theta} = -\alpha_j = [\alpha_{j+1}, \alpha_{j+2}, \dots] \quad (7.11)$$

with the continued fraction defined as

$$[\alpha_{j+1}, \alpha_{j+2}, \dots] = \frac{1}{\lambda^{\frac{3}{2}\theta} \alpha_{j+1} + \frac{\lambda^{\frac{3}{2}\theta}}{\lambda^{\frac{3}{2}\theta} \alpha_{j+2} + \frac{\lambda^{\frac{3}{2}\theta}}{\dots}}}$$

We note that $\alpha_j > 0$ for all $j \geq J$ for some large enough J , and hence the right hand side of (7.11) is positive. That implies p needs to be positive. Therefore, we know that the velocity component is unstable about the steady state with $A_0 = -1$ and $B_0 = 0$.

It follows from (7.10) that

$$\lim_{j \rightarrow \infty} \alpha_j = \lambda^{-\frac{1}{3}\theta}, \quad \lim_{j \rightarrow \infty} d_j = 0, \tag{7.12}$$

for any $q \in \mathbb{R}$. Thus the magnetic field is unstable due to a similar analysis as in the proof of Theorem 7.1. \square

Theorem 7.3. *Let (\bar{a}, \bar{b}) be the fixed point of (1.4) with $A_0 = -1$ and $B_0 = 0$. There are no negative eigenvalue to the velocity equation in the linearized system about such (\bar{a}, \bar{b}) . Every real number is an eigenvalue for the magnetic field equation in the linearized system. Thus the linearized system is unstable.*

Proof. In analogy with previous analysis for system (1.3), considering perturbation

$$a_j(t) = A_0 \lambda_j^{-\frac{1}{3}\theta} + \epsilon \omega_j(t), \quad b_j(t) = B_0 \lambda_j^{-\frac{1}{3}\theta} + \epsilon \zeta_j(t), \quad j \geq 0$$

for system (1.4), the pair (ω_j, ζ_j) satisfies

$$\begin{aligned} \omega'_j &= A_0 \lambda_j^{\frac{2}{3}\theta} \left(2\lambda^{-\frac{2}{3}\theta} \omega_{j-1} - \lambda^{-\frac{1}{3}\theta} \omega_j - \omega_{j+1} \right) \\ &\quad - B_0 \lambda_j^{\frac{2}{3}\theta} \left(2\lambda^{-\frac{2}{3}\theta} \zeta_{j-1} - \lambda^{-\frac{1}{3}\theta} \zeta_j - \zeta_{j+1} \right) \\ &\quad - \epsilon \lambda_j^\theta \omega_j \omega_{j+1} + \epsilon \lambda_{j-1}^\theta \omega_{j-1}^2 + \epsilon \lambda_j^\theta \zeta_j \zeta_{j+1} - \epsilon \lambda_{j-1}^\theta \zeta_{j-1}^2, \\ \zeta'_j &= -B_0 \lambda_j^{\frac{2}{3}\theta} \left(\lambda^{-\frac{1}{3}\theta} \omega_j - \omega_{j+1} \right) + A_0 \lambda_j^{\frac{2}{3}\theta} \left(\lambda^{-\frac{1}{3}\theta} \zeta_j - \zeta_{j+1} \right) \\ &\quad - \epsilon \lambda_j^\theta \omega_j \zeta_{j+1} + \epsilon \lambda_j^\theta \zeta_j \omega_{j+1} \end{aligned}$$

for $j \geq 0$ and $\omega_{-1} = \zeta_{-1} = 0$. If $A_0 = 1$ and $B_0 = 0$ the linearized system with $\epsilon = 0$ is

$$\begin{aligned} \omega'_j &= \lambda_j^{\frac{2}{3}\theta} \left(2\lambda^{-\frac{2}{3}\theta} \omega_{j-1} - \lambda^{-\frac{1}{3}\theta} \omega_j - \omega_{j+1} \right), \\ \zeta'_j &= \lambda_j^{\frac{2}{3}\theta} \left(\lambda^{-\frac{1}{3}\theta} \zeta_j - \zeta_{j+1} \right), \end{aligned} \tag{7.13}$$

while if $A_0 = -1$ and $B_0 = 0$ the linearized system is

$$\begin{aligned} \omega'_j &= -\lambda_j^{\frac{2}{3}\theta} \left(2\lambda^{-\frac{2}{3}\theta} \omega_{j-1} - \lambda^{-\frac{1}{3}\theta} \omega_j - \omega_{j+1} \right), \\ \zeta'_j &= -\lambda_j^{\frac{2}{3}\theta} \left(\lambda^{-\frac{1}{3}\theta} \zeta_j - \zeta_{j+1} \right). \end{aligned} \tag{7.14}$$

In view of the form of the linearized systems (7.4a)–(7.4b), (7.8), (7.13) and (7.14), we note that the instability results stated in Theorem 7.1 and Theorem 7.2 hold as well for the linearized systems of (1.4) about the fixed point with $A_0 = 1$ and $B_0 = 0$ and the fixed point with $A_0 = -1$ and $B_0 = 0$ respectively. \square

Acknowledgements. The work of M. Dai is partially supported by NSF Grants DMS-1815069 and DMS-2009422; the work of S. Friedlander is partially supported by NSF Grant DMS-1613135. S. Friedlander is grateful to IAS for its hospitality in 2020-2021.

Declarations

Conflict of interest The authors certify that there is no conflict of interest.

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

- [1] Alexakis, A., Mininni, P.D., Pouquet, A.: Shell-to-shell energy transfer in magnetohydrodynamics. I. Steady state turbulence. *Phys. Rev. E* **72**(4), 046301 (2005)
- [2] Antonov, T., Frick, P.: Cascade and scaling in a class of shell models for MHD-turbulence. *Math. Model. Syst. Processes* **8**, 4–10 (2000)
- [3] Antonov, T., Lozhkin, S., Frick, P., Sokoloff, D.: A shell model for free decaying MHD-turbulence and the role of the magnetic Prandtl number. *Magnetohydrodynamics* **37**, 87–92 (2001)
- [4] Barbato, D., Flandoli, F., Morandin, F.: A theorem of uniqueness for an inviscid dyadic model. *C.R. Math. Acad. Sci. Paris* **348**(9–10), 525–528 (2010)
- [5] Barbato, D., Flandoli, F., Morandin, F.: Anomalous dissipation in a stochastic inviscid dyadic model. *Ann. Appl. Probab.* **21**(6), 2424–2446 (2011)
- [6] Barbato, D., Flandoli, F., Morandin, F.: Energy dissipation and self-similar solutions for an unforced inviscid dyadic model. *Trans. Am. Math. Soc.* **363**(4), 1925–1946 (2011)
- [7] Barbato, D., Flandoli, F., Morandin, F.: Uniqueness for a stochastic inviscid dyadic model. *Proc. Am. Math. Soc.* **138**(7), 2607–2617 (2010)
- [8] Barbato, D., Morandin, F.: Positive and non-positive solutions for an inviscid dyadic model: well-posedness and regularity. *Nonlinear Differ. Equ. Appl.* **20**(3), 1105–1123 (2013)
- [9] Basu, A., Sain, A., Dhar, S.K., Pandit, R.: Multiscaling in models of magnetohydrodynamic turbulence. *Phys. Rev. Lett.* **81**, 2687–2690 (1998)
- [10] Biferale, L.: Shell models of energy cascade in turbulence. *Annu. Rev. Fluid Mech.* **35**, 441468 (2003)
- [11] Biskamp, D.: Cascade models for magnetohydrodynamic turbulence. *Phys. Rev. E* **50**, 2702–2711 (1994)
- [12] Caffisch, R.E., Klapper, I., Steele, G.: Remarks on singularities, dimension and energy dissipation for ideal hydrodynamics and MHD. *Commun. Math. Phys.* **184**, 443–455 (1997)
- [13] Cheskidov, A., Dai, M.: Kolmogorov’s dissipation number and the number of degrees of freedom for the 3D Navier–Stokes equations. *Proc. R. Soc. Edinb. Sect. A* **149**(2), 429–446 (2019)
- [14] Cheskidov, A., Friedlander, S., Pavlović, N.: Inviscid dyadic model of turbulence: the fixed point and Onsager’s conjecture. *J. Math. Phys.* **48**(6), 065503 (2007)
- [15] Cheskidov, A., Friedlander, S., Pavlović, N.: An inviscid dyadic model of turbulence: the global attractor. *Discret. Contin. Dyn. Syst.* **26**(3), 781–794 (2010)
- [16] Cheskidov, A., Glatt-Holtz, N., Pavlović, N., Shvydkoy, R., Vicol, V.: Susan Friedlander’s contributions in mathematical fluid dynamics. *Not. Am. Math. Soc.* **68**(3), 331–343 (2021)
- [17] Cheskidov, A., Zaya, K.: Regularizing effect of the forward energy cascade in the inviscid dyadic model. *Proc. Am. Math. Soc.* **144**, 73–85 (2016)
- [18] Constantin, P., Levant, B., Titi, E.: Analytic study of the shell model of turbulence. *Physica D* **219**(2), 120–141 (2006)
- [19] Dai, M.: Blow-up of a dyadic model with intermittency dependence for the Hall MHD. *Phys. D Nonlinear Phenom.* (2021) (to appear)
- [20] Dai, M.: Blow-up of dyadic MHD models with forward energy cascade. [arXiv: 2102.03498](https://arxiv.org/abs/2102.03498) (2021)
- [21] Dai, M., Friedlander, S.: Uniqueness and non-uniqueness results for dyadic MHD models. [arXiv:2107.04073](https://arxiv.org/abs/2107.04073) (2021)
- [22] Desnyanskiy, V.N., Novikov, E.A.: Evolution of turbulence spectra toward a similarity regime. *Izv. Akad. Nauk SSSR Fiz. Atmos. Okeana* **10**, 127–136 (1974)
- [23] Dinaburg, E.I., Sinai, Y.G.: A quasi-linear approximation of three-dimensional Navier–Stokes system. *Moscow Math. J.* **1**, 381–388 (2001)
- [24] Frick, P., Stepanov, R., Sokoloff, D.: Large- and small-scale interactions and quenching in an α^2 -dynamo. *Phys. Rev. E* **74**(6), 066310 (2006)
- [25] Friik, P.G.: Two-dimensional MHD turbulence: a hierarchical model. *Magnetohydrodynamics* **20**(3), 262–267 (1984)
- [26] Friedlander, S., Glatt-Holtz, N., Vicol, V.: Inviscid limits for a stochastically forced shell model of turbulent flow. *Ann. Inst. Henri Poincaré Probab. Stat.* **52**(3), 1217–1247 (2016)
- [27] Friedlander, S., Pavlović, N.: Blowup in a three-dimensional vector model for the Euler equations. *Commun. Pure Appl. Math.* **57**(6), 705–725 (2004)
- [28] Gledzer, E.B.: System of hydrodynamic type admitting two quadratic integrals of motion. *Sov. Phys. Dokl.* **18**, 216–217 (1973)
- [29] Gloaguen, C., Léorat, J., Pouquet, A., Grappin, R.: A scalar model for MHD turbulence. *Physica D* **17**(2), 154–182 (1985)
- [30] Jeong, I., Li, D.: A blow-up result for dyadic models of the Euler equations. *Commun. Math. Phys.* **337**, 1027–1034 (2015)
- [31] Kang, E., Lee, J.: Remarks on the magnetic helicity and energy conservation for ideal magneto-hydrodynamics. *Nonlinearity* **20**(11), 2681–2689 (2007)
- [32] Katz, N., Pavlović, N.: Finite time blow-up for a dyadic model of the Euler equations. *Trans. Am. Math. Soc.* **357**(2), 695–708 (2005)
- [33] Kiselev, A., Zlatoš, A.: On discrete models of the Euler equation. *Int. Math. Res. Not.* **38**, 2315–2339 (2005)
- [34] Nigro, G., Malara, F., Carbone, V., Veltri, P.: Nanoflares and MHD turbulence in coronal loops: a hybrid shell model. *Phys. Rev. Lett.* **92**(19), 194501 (2004)

- [35] Obukhov, A.M.: Some general properties of equations describing the dynamics of the atmosphere. *Izv. Akad. Nauk. SSSR Ser. Fiz. Atmos. Okeana* **7**, 695–704 (1971)
- [36] Ohkitani, K., Yamada, M.: Temporal intermittency in the energy cascade process and local Lyapunov analysis in fully-developed model of turbulence. *Prog. Theor. Phys.* **81**, 329–341 (1989)
- [37] Plunian, F., Stepanov, R., Frick, P.: Shell models of magnetohydrodynamic turbulence. *Phys. Rep.* **523**, 1–60 (2013)
- [38] Romito, M.: Uniqueness and blow-up for a stochastic viscous dyadic model. *Probab. Theor. Relat. Fields* **158**, 895–924 (2014)

Mimi Dai
Department of Mathematics, Statistics and
Computer Science
University of Illinois at Chicago
Chicago IL60607
USA
e-mail: mdai@uic.edu

Susan Friedlander
Department of Mathematics
University of Southern California
Los Angeles CA90089
USA
e-mail: susanfri@usc.edu

(accepted: October 19, 2021; published online: January 31, 2022)