



# Index Theory and Topological Phases of Aperiodic Lattices

C. Bourne and B. Mesland

**Abstract.** We examine the non-commutative index theory associated with the dynamics of a Delone set and the corresponding transversal groupoid. Our main motivation comes from the application to topological phases of aperiodic lattices and materials and applies to invariants from tilings as well. Our discussion concerns semifinite index pairings, factorisation properties of Kasparov modules and the construction of unbounded Fredholm modules for lattices with finite local complexity.

## Contents

Introduction	1970
1. Preliminaries on Groupoids and Kasparov Theory	1973
1.1. Kasparov Modules and Semifinite Spectral Triples	1973
1.2. Étale Groupoids, Twisted Algebras and $C^*$ -Modules	1975
1.3. The $C^*$ -Module of a Groupoid and the Reduced Twisted $C^*$ -Algebra	1976
1.4. Morita Equivalence of Twisted Groupoid $C^*$ -Algebras	1978
1.5. Exact Cocycles and Unbounded $KK$ -Cycles	1980
1.6. The Extension Class of an $\mathbb{R}$ -Valued Cocycle	1982
2. Delone Sets and the Transversal Groupoid	1985
2.1. Delone Dynamical Systems	1985
2.2. The Transversal Groupoid	1987
2.3. The Twisted Groupoid Algebra and Its $K$ -theory	1991
2.3.1. The Bulk $KK$ -Cycle	1993
2.4. One-Dimensional Delone Sets as Cuntz–Pimsner Algebras	1993
2.4.1. The Cuntz–Pimsner Extension Class	1995
3. Factorisation and the Bulk–Boundary Correspondence	1997
3.1. The Edge Groupoid	1997
3.2. The Bulk–Boundary Extension	1999
3.3. Factorisation	2001

3.3.1. A Remark on More General Boundaries	2001
3.4. $KK$ -Cycles with Higher Codimension	2002
3.4.1. Another Factorisation	2005
4. Spectral Triple Constructions	2006
4.1. The Evaluation Spectral Triple	2006
4.2. Invariant Measures and the Semifinite Spectral Triple	2007
5. Unbounded Fredholm Modules for Lattices with Finite Local Complexity	2010
5.1. The Pearson–Bellissard Spectral Triple	2011
5.2. The Product Operator	2013
5.3. The Bounded Transform as a Fredholm Module	2016
6. Index Pairings and Topological Phases	2023
6.1. Complex Pairings	2025
6.1.1. Weak Chern Numbers	2026
6.2. Real Pairings and Analytic Indices	2026
6.2.1. Extending the Pairings	2028
6.2.2. Weak Indices	2029
6.3. Pairings for Lattices with Finite Local Complexity	2029
6.4. The Bulk–Boundary Correspondence	2030
6.5. Examples From Materials Science and Meta-Materials	2031
Acknowledgements	2032
References	2033

## Introduction

Models of systems in solid-state physics that do not make reference to a Bloch decomposition or Fourier transform are essential if one wishes to understand topological phases of disordered or aperiodic systems. A description of disordered media using crossed product  $C^*$ -algebras has successfully adapted many important properties of periodic topological insulators to the disordered setting, see [13, 42, 77] for example.

Recently, newer proposed models of topological materials and meta-materials have emerged whose underlying lattice has a quasicrystalline [8] or amorphous configuration [68]. In the case of amorphous lattice configurations, because there is no canonical labelling of the lattice points, the Hamiltonians of interest cannot be described by a crossed product  $C^*$ -algebra. Hence, the techniques and results on the bulk–boundary correspondence in [20] cannot be applied to recent results on edge states and transport in topological amorphous (meta-)materials [68]. Such amorphous systems are instead modelled by the transversal (étale) groupoid  $\mathcal{G}$  associated with a Delone set developed in [10, 14, 47, 48]. One of the key results of the paper is the extension of the  $K$ -theoretic framework for topological phases to such algebras and aperiodic media.

Quasicrystalline materials often display finite local complexity, meaning that up to translation the lattice is determined by a finite number of patterns or polyhedra (cf. Definition 2.2). If the lattice has finite local complexity, the aperiodic but ordered pattern configurations can be described using tiling spaces.

By [85, Theorem 2], the tiling space of such lattices is homeomorphic to the  $d$ -fold suspension of a  $\mathbb{Z}^d$ -subshift space, though not necessarily topologically conjugate. This result implies that, adding some extra sites to quasicrystalline lattice configurations if necessary, there is a  $\mathbb{Z}^d$ -labelling of points and the system can be described by a discrete crossed product  $C(Z) \rtimes \mathbb{Z}^d$ .

The advantage of the transversal groupoid approach is that it does not require finite local complexity nor a  $\mathbb{Z}^d$ -labelling. In particular, the framework can accommodate non-periodic  $\mathbb{R}^d$ -actions. Thus, the modifications needed to obtain a  $\mathbb{Z}^d$ -labelling (which may not be physically reasonable) can be avoided. Furthermore, the transversal groupoid covers a broader range of examples which are not covered by the results in [85] such as the pinwheel tiling.

Given a Hamiltonian on an aperiodic system, computational techniques are currently in development to determine its spectrum [9]. If the Hamiltonian contains a spectral gap, we can associate a topological phase with the system by modelling its dynamics via the transversal groupoid  $\mathcal{G}$ . In particular, topological indices and  $K$ -theoretic properties of such Hamiltonians are determined using the groupoid  $C^*$ -algebra.

In previous work, this groupoid description was used to describe bulk topological phases [21]. In this paper, we show that a (gapped) Hamiltonian acting on a Delone set  $\mathcal{L} \subset \mathbb{R}^d$  is enough to define strong and weak topological phases as well as show the bulk–edge correspondence of Hamiltonians acting on the lattice  $\ell^2(\mathcal{L})$ . Furthermore, if the unit space  $\Omega_0$  of the transversal groupoid  $\mathcal{G}$  has an invariant measure, then Chern number formulas can also be defined for complex topological phases.

Because of the generality of Delone sets, they are able to model materials that go beyond what is normally considered when discussing topological phases of matter. These include quasicrystal structures but also other materials such as glasses and some liquids, see [11] for example. This tells us, at least from a mathematical perspective, that our constructions and results are potentially applicable to a broader range of materials and meta-materials in addition to the applications to (possibly disordered) crystals.

In the present paper, our central object of study is an unbounded  $KK$ -cycle for the transversal groupoid  $C^*$ -algebra which gives rise to a class in  $KK^d(C_r^*(\mathcal{G}, \sigma), C(\Omega_0))$  (real or complex) with  $d$  the dimension of the underlying space,  $\sigma$  a magnetic twisting and  $\Omega_0$  the transversal space. When this  $KK$ -cycle is coupled with the  $K$ -theoretic phase of a gapped free-fermionic Hamiltonian [which gives a class in  $K_n(C_r^*(\mathcal{G}, \sigma))$ ], the corresponding index pairing gives analytic indices that encode the strong topological phase. When the transversal  $\Omega_0$  has an invariant probability measure, we can construct a semifinite spectral triple and measure this (disorder-averaged) pairing using the semifinite local index formula (considered for ergodic measures in [21]).

The factorisation properties of the unbounded  $KK$ -cycle also allow us to express the index pairing as a pairing over a closed subgroupoid  $\Upsilon$  that encodes the dynamics of the transversal in  $(d - 1)$ -dimensions and models an edge system. Namely, we can link these systems explicitly via a short exact sequence

$$0 \rightarrow C_r^*(\Upsilon, \sigma) \otimes \mathbb{K} \rightarrow \mathcal{T} \rightarrow C_r^*(\mathcal{G}, \sigma) \rightarrow 0$$

with  $\mathcal{T}$  modelling a half-space system. When the lattices we consider have a canonical  $\mathbb{Z}^d$ -labelling, then this short exact sequence is the usual Toeplitz extension of a crossed product considered in [77]. Our result, analogous to the crossed product case, [20, 77], is that our  $d$ -dimensional pairing with  $C_r^*(\mathcal{G}, \sigma)$  (as an element in the  $K$ -theory of the configuration space or a numerical phase label of this pairing) is equal to or in the same  $K$ -theory class as the  $(d - 1)$ -dimensional pairing with  $C_r^*(\Upsilon, \sigma)$  up to a possible sign.

For aperiodic lattices with finite local complexity, the transversal  $\Omega_0$  is totally disconnected. In this case, a general construction due to Pearson and Bellissard [74] gives a family of spectral triples on  $C(\Omega_0)$ . Coupling the unbounded  $KK$ -cycle for  $(C_r^*(\mathcal{G}), C(\Omega_0))$  to such a spectral triple for  $C(\Omega_0)$  using the unbounded Kasparov product gives us  $K$ -homology representatives for  $C_r^*(\mathcal{G})$ . The construction of the product operator employs the techniques developed in [66], but the commutators with  $C_r^*(\mathcal{G})$  turn out to be unbounded. Nonetheless, using arguments similar to [36] and recent results in [63], we are able to show that the operator represents the Kasparov product of the given classes via the bounded transform. The analytic difficulties with the commutators can be directly attributed to the disorder, that is, the non-periodicity of the Delone set.

Let us remark that the unbounded Fredholm module constructed from quasicrystalline lattice configurations allows us to consider new topological phases that cannot be defined in periodic systems or disordered systems with a contractible disorder space of configurations. Indeed, the totally disconnected structure of the transversal  $\Omega_0$  is a crucial ingredient in defining these new phases.

Some of our results show parallels with those of Kubota and of Ewert–Meyer, who study topological phases associated with Delone sets and the corresponding Roe algebra [31, 54]. Briefly, the Roe algebra, by its universal nature, provides a means to compare topological phases from different lattice configurations (see [54, Lemma 2.19]). Conversely, the transversal groupoid algebra is used to determine the topological phase of Hamiltonians associated with a fixed lattice configuration. Because the groupoid algebra is separable (while the Roe algebra is not), it is more susceptible to the use of  $KK$ -theoretic machinery, which is a central theme of this paper. In particular, it is generally easier to both define and compute the pairings with  $KK$ -cycles or cyclic cocycles that characterise the numerical phase labels; see [21, Section 3] for numerical simulations.

Lastly, the groupoid of a transversal is typically used to study the dynamics of aperiodic tilings and related dynamical systems. We have not emphasised the application to tilings in this manuscript, though our constructions and results may have broader interest.

**Outline.** Because our paper draws from aspects of dynamical systems, operator algebras, Kasparov theory and physics, we aim to give a systematic and largely self-contained exposition of our results.

We first give a brief overview of the mathematical tools we require in Sect. 1, which include Kasparov theory, semifinite index theory and  $C^*$ -algebras of étale groupoids twisted by a 2-cocycle. We consider  $C^*$ -modules constructed from étale groupoids and review how groupoid equivalences can be naturally expressed in terms of  $C^*$ -modules. In particular, we consider groupoids with a normalised 2-cocycle, where groupoid equivalence for compatible twists gives rise to a Morita equivalence of the twisted groupoid  $C^*$ -algebras. We also provide a higher-dimensional extension of the result in [65], where if one has a continuous 1-cocycle  $c : \mathcal{G} \rightarrow \mathbb{R}^n$  that is exact in the sense of [65], then this cocycle gives rise to a Dirac-like operator and unbounded Kasparov module over the twisted  $C^*$ -algebra of  $\mathcal{G}$  relative to that of a closed subgroupoid  $\mathcal{H} = \text{Ker}(c)$ . We further provide a condition on  $\mathbb{R}$ -valued cocycles that guarantees injectivity of the Busby invariant directly. This condition is satisfied in all examples considered in the paper.

In Sect. 2, we review the construction of the transversal groupoid following [10, 14, 47, 50] and show how it fits into our general  $KK$ -theoretic framework. In the case of dimension 1, we give an alternative description of the groupoid  $C^*$ -algebra and unbounded  $KK$ -cycle using Cuntz–Pimsner algebras and results from [37, 80, 81].

In Sect. 3, we show how the unbounded  $KK$ -cycle we build factorises into the product of an ‘edge’  $KK$ -cycle modelling a system of codimension 1 with a linking  $KK$ -cycle that relates the two systems. This can also be extended to higher codimension and is related to weak topological insulators.

We then consider spectral triple constructions in Sects. 4 and 5. We construct spectral triples using the evaluation map of the transversal, an invariant measure (which gives a semifinite spectral triple) and the product with a Pearson–Bellissard spectral triple. The latter construction yields an unbounded Fredholm module with mildly unbounded commutators as in [36], so that the bounded transform represents the Kasparov product.

Lastly, we apply our results to topological phases in Sect. 6, where the physical invariants of interest naturally arise as index pairings of classes in  $K_n(C_r^*(\mathcal{G}, \sigma))$  with our unbounded  $KK$ -cycles (or spectral triples). Here we prove Chern number formulas for complex phases, analytic strong and weak indices for systems with anti-linear symmetries and the bulk–boundary correspondence. Much like the crossed product setting, our bulk indices are also well defined for a much larger algebra that can be constructed using non-commutative  $L^p$ -spaces. A connection of these extended indices to regions of dynamical or spectral localisation remains an open problem.

## 1. Preliminaries on Groupoids and Kasparov Theory

### 1.1. Kasparov Modules and Semifinite Spectral Triples

In this section, we establish basic results and notation that we will need for this paper. Because we are motivated by topological phases whose relation to

real  $K$ -theory is now well-established [34, 40, 49], we will work in both real and complex vector spaces and algebras.

Given a real or complex right- $B$   $C^*$ -module  $E_B$ , we denote the right action by  $e \cdot b$  and the  $B$ -valued inner product  $(\cdot | \cdot)_B$ . The set of adjointable endomorphisms on  $E_B$  with respect to this inner product is denoted by  $\text{End}^*(E_B)$ . The rank-1 operators  $\Theta_{e,f}$ ,  $e, f \in E_B$ , are defined such that

$$\Theta_{e,f}(g) = e \cdot (f | g)_B, \quad e, f, g \in E_B.$$

The norm closure of the algebraic span of the set of such rank-1 operators is the set of compact operators on  $E_B$ , and we denote this set by  $\mathbb{K}(E_B)$ . We will often work with  $\mathbb{Z}_2$ -graded algebras and spaces and denote by  $\hat{\otimes}$  the graded tensor product (see [44, Section 2] and [17, Section 14]). A densely defined closed symmetric operator  $T : \text{Dom } T \rightarrow E_B$  is *self-adjoint and regular* if the operators  $T \pm i : \text{Dom } T \rightarrow E_B$  have dense range. See [60, Chapter 9–10] for the basic theory of unbounded operators on  $C^*$ -modules.

**Definition 1.1.** Let  $A$  and  $B$  be  $\mathbb{Z}_2$ -graded real (resp. complex)  $C^*$ -algebras. A real (resp. complex) unbounded Kasparov module  $(\mathcal{A}, \pi E_B, D)$  (also called an unbounded  $KK$ -cycle) for  $(A, B)$  consists of

- (1) a  $\mathbb{Z}_2$ -graded real (resp. complex)  $C^*$ -module  $E_B$ ,
- (2) a graded  $*$ -homomorphism  $\pi : A \rightarrow \text{End}^*(E_B)$ ,
- (3) an unbounded self-adjoint, regular and odd operator  $D$  and a dense  $*$ -subalgebra  $\mathcal{A} \subset A$  such that for all  $a \in \mathcal{A} \subset A$ ,

$$[D, \pi(a)]_{\pm} \in \text{End}^*(E_B), \quad \pi(a)(1 + D^2)^{-1} \in \mathbb{K}(E_B).$$

For complex algebras and spaces, one can also remove the gradings, in which case the Kasparov module is called odd (otherwise even).

We will often omit the representation  $\pi$  when the left action is unambiguous. Unbounded Kasparov modules represent classes in the  $KK$ -group  $KK(A, B)$  or  $KKO(A, B)$  [7]. We note that an unbounded  $A$ - $\mathbb{C}$  or  $A$ - $\mathbb{R}$  Kasparov module is precisely the definition of a complex or real spectral triple.

Another non-commutative extension of index theory and closely related to unbounded Kasparov theory are semifinite spectral triples [25, 26]. Let  $\tau$  be a fixed faithful, normal, semifinite trace on a von Neumann algebra  $\mathcal{N}$ . We denote by  $\mathcal{K}_{\mathcal{N}}$  the  $\tau$ -compact operators in  $\mathcal{N}$ , that is, the norm closed ideal generated by the projections  $P \in \mathcal{N}$  with  $\tau(P) < \infty$ .

**Definition 1.2.** Let  $\mathcal{N} \subset \mathcal{B}(\mathcal{H})$  be a graded semifinite von Neumann algebra with trace  $\tau$ . A semifinite spectral triple  $(\mathcal{A}, \mathcal{H}, D)$  is given by a  $\mathbb{Z}_2$ -graded Hilbert space  $\mathcal{H}$ , a graded  $*$ -algebra  $\mathcal{A} \subset \mathcal{N}$  with  $C^*$ -closure  $A$  and a graded representation on  $\mathcal{H}$ , together with a densely defined odd unbounded self-adjoint operator  $D$  affiliated to  $\mathcal{N}$  such that

- (1)  $[D, a]_{\pm}$  is well defined on  $\text{Dom}(D)$  and extends to a bounded operator on  $\mathcal{H}$  for all  $a \in \mathcal{A}$ ,
- (2)  $a(1 + D^2)^{-1} \in \mathcal{K}_{\mathcal{N}}$  for all  $a \in A$ .

For  $\mathcal{N} = \mathcal{B}(\mathcal{H})$  and  $\tau = \text{Tr}$ , one recovers the usual definition of a spectral triple. A semifinite spectral triple relative to  $(\mathcal{N}, \tau)$  with  $\mathcal{A}$  unital is called  $p$ -summable if  $(1 + D^2)^{-s/2}$  is  $\tau$ -trace class for all  $s > p$ . We also call a semifinite spectral triple  $QC^\infty$  if  $a, [D, a] \in \text{Dom}(\delta^k)$  for all  $k \in \mathbb{N}$  with  $\delta(T) = [|D|, T]$  being the partial derivation.

Semifinite spectral triples can be paired with  $K$ -theory classes in  $K_*(\mathcal{A})$  via a semifinite Fredholm index [16]. An operator  $T \in \mathcal{N}$  that is invertible modulo  $\mathcal{K}_{\mathcal{N}}$  has a semifinite Fredholm index

$$\text{Index}_\tau(T) = \tau(P_{\text{Ker}(T)}) - \tau(P_{\text{Ker}(T^*)}).$$

If the semifinite spectral triples are  $p$ -summable and  $QC^\infty$ , the complex index pairing can be computed using the resolvent cocycle and the semifinite local index formula [25, 26]. By writing the index pairing as a pairing with cyclic cohomology, the topological invariants of interest can more easily be connected to physics [77]. See [16, 26, 32] for further details on semifinite index theory and [76] for results concerning numerical implementation.

Suppose  $(\mathcal{A}, E_B, D)$  is an unbounded Kasparov module for a separable  $C^*$ -algebra  $A$  and the right-hand algebra  $B$  has a faithful, semifinite and norm lower semicontinuous trace  $\tau_B$ . We work with faithful traces as we can always pass to a quotient algebra  $B/\text{Ker}(\tau_B)$  if necessary. Assuming such a trace, one can often construct a semifinite spectral triple via a dual trace construction [58]. We follow this approach in Sect. 4.2. By constructing a semifinite spectral triple from a Kasparov module, we obtain a  $KK$ -theoretic interpretation of the semifinite index pairing, which can be expressed via the map

$$K_*(A) \times KK^*(A, B) \rightarrow K_0(B) \xrightarrow{(\tau_B)^*} \mathbb{R}, \tag{1}$$

with  $(\mathcal{A}, E_B, D)$  representing the class in  $KK^*(A, B)$ . Equation (1) allows us to more explicitly characterise the range of the semifinite index pairing (which is in general  $\mathbb{R}$ -valued). The local index formula then gives us a computable expression for the  $KK$ -theoretic composition in Eq. (1).

### 1.2. Étale Groupoids, Twisted Algebras and $C^*$ -Modules

We start with some basic definitions for convenience. Our standard reference for groupoid  $C^*$ -algebras is [79].

**Definition 1.3.** A groupoid is a set  $\mathcal{G}$  with a subset  $\mathcal{G}^{(2)} \subset \mathcal{G} \times \mathcal{G}$ , a multiplication map  $\mathcal{G}^{(2)} \rightarrow \mathcal{G}$ ,  $(\gamma, \xi) \mapsto \gamma\xi$  and an inverse  $\mathcal{G} \rightarrow \mathcal{G}$   $\gamma \mapsto \gamma^{-1}$  such that

- (1)  $(\gamma^{-1})^{-1} = \gamma$  for all  $\gamma \in \mathcal{G}$ ,
- (2) if  $(\gamma, \xi), (\xi, \eta) \in \mathcal{G}^{(2)}$ , then  $(\gamma\xi, \eta), (\gamma, \xi\eta) \in \mathcal{G}^{(2)}$ ,
- (3)  $(\gamma, \gamma^{-1}) \in \mathcal{G}^{(2)}$  for all  $\gamma \in \mathcal{G}$ ,
- (4) for all  $(\gamma, \xi) \in \mathcal{G}^{(2)}$ ,  $(\gamma\xi)\xi^{-1} = \gamma$  and  $\gamma^{-1}(\gamma\xi) = \xi$ .

Given a groupoid, we denote by  $\mathcal{G}^{(0)} = \{\gamma\gamma^{-1} : \gamma \in \mathcal{G}\}$  the space of units and define the source and range maps  $r, s : \mathcal{G} \rightarrow \mathcal{G}^{(0)}$  by the equations

$$r(\gamma) = \gamma\gamma^{-1}, \quad s(\gamma) = \gamma^{-1}\gamma$$

for all  $\gamma \in \mathcal{G}$ . The source and range maps allow us to characterise

$$\mathcal{G}^{(2)} = \{(\gamma, \xi) \in \mathcal{G} \times \mathcal{G} : s(\gamma) = r(\xi)\}.$$

We furthermore assume that  $\mathcal{G}$  has a locally compact topology such that  $\mathcal{G}^{(0)} \subset \mathcal{G}$  is Hausdorff in the relative topology and multiplication, inversion, source and range maps all continuous. In this work, we restrict ourselves to groupoids that are both Hausdorff and étale.

**Definition 1.4.** A topological groupoid  $\mathcal{G}$  is called étale if the range map  $r : \mathcal{G} \rightarrow \mathcal{G}^{(0)}$  is a local homeomorphism.

**Definition 1.5.** Let  $\mathcal{G}$  be a locally compact and Hausdorff groupoid. A continuous map  $\sigma : \mathcal{G}^{(2)} \rightarrow \mathbb{T} \simeq U(1)$  is a 2-cocycle if

$$\sigma(\gamma, \xi)\sigma(\gamma\xi, \eta) = \sigma(\gamma, \xi\eta)\sigma(\xi, \eta)$$

for any  $(\gamma, \xi), (\xi, \eta) \in \mathcal{G}^{(2)}$ , and

$$\sigma(\gamma, s(\gamma)) = 1 = \sigma(r(\gamma), \gamma)$$

for all  $\gamma \in \mathcal{G}$ . We will call a groupoid 2-cocycle normalised if  $\sigma(\gamma, \gamma^{-1}) = 1$  for all  $\gamma \in \mathcal{G}$ .

*Remark 1.6.* We can also define  $O(1) \simeq \mathbb{Z}_2$ -valued groupoid 2-cocycles whose cocycle relation is the same as the  $U(1)$  case. Generally speaking, if we are working in the category of complex spaces and algebras, we will use  $U(1)$ -valued 2-cocycles. If we are in the real category, we work with  $O(1)$ -valued 2-cocycles.

Because the algebraic structure is the same in either setting, we will abuse notation slightly and denote by  $\sigma$  a generic groupoid 2-cocycle, where the range of this 2-cocycle will be clear from the context.

Given an étale groupoid  $\mathcal{G}$  and 2-cocycle  $\sigma$ , we define  $C_c(\mathcal{G}, \sigma)$  to be the  $*$ -algebra of compactly supported functions on  $\mathcal{G}$  with twisted convolution and involution

$$(f_1 * f_2)(\gamma) = \sum_{\gamma=\xi\eta} f_1(\xi)f_2(\eta)\sigma(\xi, \eta), \quad f^*(\gamma) = \sigma(\gamma, \gamma^{-1})\overline{f(\gamma^{-1})}.$$

The 2-cocycle condition ensures that  $C_c(\mathcal{G}, \sigma)$  is an associative  $*$ -algebra. In the present paper, we restrict ourselves to considering normalised cocycles, which covers all examples of interest to us. Our definition of the groupoid 2-cocycle and twisted convolution algebra comes from Renault [79]. For a broader version of twisted groupoid algebra, see [56].

### 1.3. The $C^*$ -Module of a Groupoid and the Reduced Twisted $C^*$ -Algebra

Take an étale groupoid  $\mathcal{G}$  with a normalised 2-cocycle  $\sigma$ . The space  $C_c(\mathcal{G}, \sigma)$  is a right module over  $C_0(\mathcal{G}^{(0)})$  via  $(f \cdot g)(\xi) = f(\xi)g(s(\xi))$ . Since  $\mathcal{G}^{(0)} \subset \mathcal{G}$  is closed, we can consider the restriction map  $\rho : C_c(\mathcal{G}) \rightarrow C_0(\mathcal{G}^{(0)})$ . This defines a  $C_0(\mathcal{G}^{(0)})$  valued inner product on the right module  $C_c(\mathcal{G}, \sigma)$  via



$$\begin{aligned} (f_1 \mid f_2)_{C_0(\mathcal{G}^{(0)})}(x) &:= \rho(f_1^* * f_2)(x) \\ &= \sum_{\xi \in s^{-1}(x)} \overline{f_1(\xi^{-1})} f_2(\xi^{-1}) \sigma(\xi, \xi^{-1}) = \sum_{\xi \in r^{-1}(x)} \overline{f_1(\xi)} f_2(\xi) \end{aligned}$$

as  $\sigma$  is normalised. Denote by  $E_{C_0(\mathcal{G}^{(0)})}$  the  $C^*$ -module completion of  $C_c(\mathcal{G})$  in this inner product. There is an action of the  $*$ -algebra  $C_c(\mathcal{G}, \sigma)$  on the  $C^*$ -module  $E_{C_0(\mathcal{G}^{(0)})}$  by bounded adjointable endomorphisms, extending the action of  $C_c(\mathcal{G}, \sigma)$  on itself by left multiplication.

**Definition 1.7** (cf. [52]). The reduced groupoid  $C^*$ -algebra  $C_r^*(\mathcal{G}, \sigma)$  is the completion of  $C_c(\mathcal{G}, \sigma)$  in the norm inherited from the embedding  $C_c(\mathcal{G}, \sigma) \hookrightarrow \text{End}^*(E_{C_0(\mathcal{G}^{(0)})})$ .

**Definition 1.8.** Let  $\mathcal{G}$  be an étale groupoid over  $\mathcal{G}^{(0)}$ . An  $s$ -cover of  $\mathcal{G}$  is a locally finite countable open cover  $\mathcal{V} := \{V_i\}_{i \in \mathbb{N}}$  consisting of pre-compact sets, such that  $s : V_i \rightarrow \mathcal{G}^{(0)}$  is a homeomorphism onto its image.

**Lemma 1.9.** Let  $\mathcal{V} := \{V_i\}_{i \in \mathbb{N}}$  be an  $s$ -cover of  $\mathcal{G}$  and  $\chi_i : V_i \rightarrow \mathbb{R}$  a partition of unity subordinate to  $\mathcal{V}$ , that is,  $\sum_i \chi_i(\eta)^2 = 1$  for all  $\eta \in \mathcal{G}$ . Write  $u_m := \sum_{i \leq m} \Theta_{\chi_i, \chi_i}$ . Then for all  $f \in C_c(\mathcal{G}, \sigma)$  there exists  $N \in \mathbb{N}$  such that for all  $n \geq N$

$$f(\eta) = u_n f(\eta) = \sum_{i \leq n} \chi_i * \rho(\chi_i^* * f)(\eta).$$

In particular,  $u_m f$  converges to  $f$  in the norm of  $E_{C_0(\mathcal{G}^{(0)})}$ .

*Proof.* The above, together with the fact that we have an  $s$ -cover gives

$$\begin{aligned} \sum_i (\chi_i \cdot \rho(\chi_i^* * f))(\eta) &= \sum_i \chi_i(\eta) \rho(\chi_i^* * f)(s(\eta)) \\ &= \sum_i \sum_{\xi \in r^{-1}(s(\eta))} \chi_i(\eta) \chi_i^*(\xi) f(\xi^{-1}) \sigma(\xi, \xi^{-1}) \\ &= \sum_i \sum_{\xi \in r^{-1}(s(\eta))} \chi_i(\eta) \chi_i(\xi^{-1}) f(\xi^{-1}) \\ &= \sum_i \sum_{\xi \in s^{-1}(s(\eta))} \chi_i(\eta) \chi_i(\xi) f(\xi) \\ &= \sum_{\{i: \eta \in V_i\}} \chi_i^2(\eta) f(\eta) = f(\eta). \end{aligned}$$

Since  $f$  has compact support, there exists  $N = N_f$  such that  $\chi_n|_{\text{supp} f} = 0$  for all  $n \geq N$ . Thus, the sum above is uniformly finite and hence convergent in the  $\rho$ -norm. □

Note that the above result implies that  $u_n$  is a sequence of local units for  $C_c(\mathcal{G}^{(2)}) \subset \mathbb{K}(E_{C_0(\mathcal{G}^{(0)})})$ .

**Lemma 1.10.** We have  $\sup_n \|u_n\|_{\text{End}^*(E_{C_0(\mathcal{G}^{(0)})})} \leq 1$ .

*Proof.* We compute the operator norm of the  $u_n$  directly. Let  $f \in C_c(\mathcal{G}, \sigma)$ :

$$\begin{aligned} (u_n f \mid u_n f)_{C(\mathcal{G}^{(0)})}(x) &= \sum_{\xi \in r^{-1}(x)} |u_n f(\xi)|^2 \\ &= \sum_{\xi \in r^{-1}(x)} \left| \sum_{i \leq n} \chi_i(\xi)^2 f(\xi) \right|^2 \leq \sum_{\xi \in r^{-1}(x)} \left( \sum_{i \leq n} \chi_i(\xi)^2 |f(\xi)| \right)^2 \\ &\leq \sum_{\xi \in r^{-1}(x)} |f(\xi)|^2 = (f \mid f)_{C(\mathcal{G}^{(0)})}(x) \end{aligned}$$

Thus, it follows that

$$\begin{aligned} \|u_n f\|_{C_0(\mathcal{G}^{(0)})}^2 &= \sup_{x \in \mathcal{G}^{(0)}} (u_n f \mid u_n f)_{C(\mathcal{G}^{(0)})}(x) \\ &\leq \sup_{x \in \mathcal{G}^{(0)}} (f \mid f)_{C(\mathcal{G}^{(0)})}(x) = \|f\|_{C_0(\mathcal{G}^{(0)})}^2, \end{aligned}$$

and we find that  $\sup \|u_n\|_{\text{End}^*(E_{C_0(\mathcal{G}^{(0)})})} \leq 1$  as claimed. □

**Proposition 1.11.** *The sequence  $u_n$  forms an approximate unit for  $\mathbb{K}(E_{C_0(\mathcal{G}^{(0)})})$ . In other words, the ordered set of elements  $\chi_i \in E_{C_0(\mathcal{G}^{(0)})}$  forms a frame for  $E_{C_0(\mathcal{G}^{(0)})}$ .*

*Proof.* The sequence  $u_n$  is uniformly bounded in operator norm and converges strongly to 1 on a dense subset. This implies that it converges strongly to 1 on all of  $E_{C_0(\mathcal{G}^{(0)})}$ , which is equivalent to being an approximate unit for  $\mathbb{K}(E_{C_0(\mathcal{G}^{(0)})})$ . □

### 1.4. Morita Equivalence of Twisted Groupoid $C^*$ -Algebras

In this section we work with an arbitrary étale groupoid  $\mathcal{G}$  with closed subgroupoid  $\mathcal{H}$  that admits a Haar system and a normalised 2-cocycle  $\sigma : \mathcal{G}^{(2)} \rightarrow \mathbb{T}$  or  $\{\pm 1\}$ . The map  $\sigma$  restricts to a 2-cocycle on the subgroupoid  $\mathcal{H}$ . Denote by

$$\rho_{\mathcal{H}} : C_c(\mathcal{G}, \sigma) \rightarrow C_c(\mathcal{H}, \sigma),$$

the restriction map. This map is a generalised conditional expectation by [79, Proposition 2.9]. It gives rise to a  $C_c(\mathcal{H}, \sigma)$ -valued inner product, where

$$\begin{aligned} (f_1 \mid f_2)_{C_c(\mathcal{H}, \sigma)}(\eta) &= \rho_{\mathcal{H}}(f_1^* * f_2)(\eta) \\ &= \sum_{\xi \in r_{\mathcal{G}}^{-1}(r_{\mathcal{H}}(\eta))} f_1^*(\eta^{-1}\xi) f_2(\xi^{-1}) \sigma(\eta^{-1}\xi, \xi^{-1}). \end{aligned}$$

This map is compatible with the right action,

$$(f \cdot h)(\gamma) = \sum_{\eta \in r_{\mathcal{H}}^{-1}(s_{\mathcal{G}}(\gamma))} f(\gamma\eta) h(\eta^{-1}) \sigma(\gamma\eta, \eta^{-1}), \quad f \in C_c(\mathcal{G}, \sigma), \quad h \in C_c(\mathcal{H}, \sigma).$$

We again take the completion of  $C_c(\mathcal{G}, \sigma)$  in the  $C_r^*(\mathcal{H}, \sigma)$ -valued inner product to obtain a right  $C^*$ -module  $E_{C_r^*(\mathcal{H}, \sigma)}$ . The left action of  $C_c(\mathcal{G}, \sigma)$  on itself makes  $E_{C_r^*(\mathcal{H}, \sigma)}$  into a  $(C_r^*(\mathcal{G}, \sigma), C_r^*(\mathcal{H}, \sigma))$ -bimodule by [67, Theorem

1.4]. These bimodules often support a natural operator making them into  $KK$ -cycles, as we will discuss in Sect. 1.5.

At present we wish to describe the compact operators on  $E_{C_r^*(\mathcal{H},\sigma)}$ . To this end, we first define

$$\mathcal{G}/\mathcal{H} \cong \{[\xi] : \xi \in \mathcal{G}, [\gamma] = [\xi] \iff \text{there exists } \eta \in \mathcal{H} \text{ with } \gamma\eta = \xi\}.$$

We can define a new groupoid by considering a left action of  $\mathcal{G}$  on this quotient space. Namely, we take

$$\mathcal{G} \ltimes \mathcal{G}/\mathcal{H} := \{(\xi, [\gamma]) \in \mathcal{G} \times \mathcal{G}/\mathcal{H} : s_{\mathcal{G}}(\xi) = r_{\mathcal{G}}(\gamma)\},$$

where we have  $(\mathcal{G} \ltimes \mathcal{G}/\mathcal{H})^{(0)} = \mathcal{G}/\mathcal{H}$  and

$$\begin{aligned} r(\xi, [\gamma]) &= [\xi\gamma], & s(\xi, [\gamma]) &= [\gamma], \\ (\xi, [\gamma])^{-1} &= (\xi^{-1}, [\xi\gamma]), & (\xi, [\gamma]) \circ (\eta, [\eta^{-1}\gamma]) &= (\xi\eta, [\eta^{-1}\gamma]). \end{aligned}$$

Furthermore, we can again use the 2-cocycle  $\sigma$  on  $\mathcal{G}$  to define a 2-cocycle on  $\mathcal{G} \ltimes \mathcal{G}/\mathcal{H}$ ,

$$\sigma((\xi, [\gamma]), (\eta, [\eta^{-1}\gamma])) = \sigma(\xi, \eta).$$

The groupoid  $\mathcal{G}$  naturally implements an equivalence between  $\mathcal{G} \ltimes \mathcal{G}/\mathcal{H}$  and  $\mathcal{H}$  in the sense of [71]. Namely  $\mathcal{G}$  is a free and proper left  $(\mathcal{G} \ltimes \mathcal{G}/\mathcal{H})$ -space and a free and proper right  $\mathcal{H}$ -space via the groupoid actions,

$$(\xi, [\gamma]) \cdot \eta = \xi\eta, \quad s(\xi) = r(\gamma) = r(\eta), \quad \gamma \cdot \eta = \gamma\eta, \quad s(\gamma) = r(\eta).$$

In particular,  $C_c(\mathcal{G})$  can be completed into Morita equivalence bimodules for both the full and reduced  $C^*$ -algebras of  $\mathcal{H}$  and  $\mathcal{G} \ltimes \mathcal{G}/\mathcal{H}$  [71, 90]. If the 2-cocycles on  $\mathcal{H}$  and  $\mathcal{G} \ltimes \mathcal{G}/\mathcal{H}$  are compatible (e.g. if both are inherited from a fixed 2-cocycle on  $\mathcal{G}$ ), then the full twisted groupoid  $C^*$ -algebras are Morita equivalent by [30, Theorem 9.1]. Morita equivalence was extended to the reduced  $C^*$ -algebras of Fell bundles in [69, 70, 91], which includes twisted reduced groupoid  $C^*$ -algebras (see [70, Proposition 6.2]). We briefly review this construction for the special case in which we are working.

We define a left action of  $C_c(\mathcal{G} \ltimes \mathcal{G}/\mathcal{H}, \sigma)$  on  $C_c(\mathcal{G}, \sigma)$  (seen as a right  $C_c(\mathcal{H}, \sigma)$ -module) by the formula

$$\begin{aligned} (\pi(g)f)(\gamma) &= \sum_{\xi \in r^{-1}(r(\gamma))} g(\xi, [\xi^{-1}\gamma])f(\xi^{-1}\gamma)\sigma(\xi, \xi^{-1}\gamma), \\ g &\in C_c(\mathcal{G} \ltimes \mathcal{G}/\mathcal{H}, \sigma), \quad f \in C_c(\mathcal{G}, \sigma). \end{aligned}$$

As we argue below, this action extends to an isomorphism

$$C_r^*(\mathcal{G} \ltimes \mathcal{G}/\mathcal{H}, \sigma) \xrightarrow{\cong} \mathbb{K}(E_{C_r^*(\mathcal{H},\sigma)}),$$

to obtain the following result.

**Proposition 1.12** ([70], Theorem 5.5, [90], Theorem 4.1, [91], Theorem 14). *The  $C^*$ -module  $E_{C_r^*(\mathcal{H},\sigma)}$  is a Morita equivalence bimodule between the  $C^*$ -algebras  $C_r^*(\mathcal{H}, \sigma)$  and  $C_r^*(\mathcal{G} \ltimes \mathcal{G}/\mathcal{H}, \sigma)$ .*

This statement is derived from the proof in [90] with fairly minor alterations. The more general Fell bundle setting requires more machinery, see [70, 91]. We define the linking groupoid as the topological disjoint union,

$$L = (\mathcal{G} \ltimes \mathcal{G}/\mathcal{H}) \sqcup \mathcal{G} \sqcup \mathcal{G}^{\text{op}} \sqcup \mathcal{H}$$

where  $\mathcal{G}^{\text{op}}$  is the opposite groupoid  $\mathcal{G}^{\text{op}} = \{\bar{\gamma} : \gamma \in \mathcal{G}\}$ , which we can equip with a 2-cocycle  $\sigma^{\text{op}}(\bar{\gamma}_1, \bar{\gamma}_2) = \sigma(\gamma_2, \gamma_1)$ . As the name suggests,  $L$  is a groupoid with unit space  $\mathcal{G}/\mathcal{H} \sqcup \mathcal{H}$  and source and range maps inherited from the groupoid structure on its parts [90, Lemma 2.1]. We can consider the twisted convolution algebra of  $L$ , with respect to the cocycle  $\hat{\sigma} : L \rightarrow \mathbb{T}$  which coincides with the given cocycles on each of the components of the disjoint union.

The algebraic machinery used in [72, 90] also works in the twisted case (see [79, Chapter II, Lemma 2.5] or [69, Chapter 5]), and the argument in [90] follows through to obtain the result.

### 1.5. Exact Cocycles and Unbounded $KK$ -Cycles

We now discuss the construction of  $KK$ -cycles from the data of a continuous 1-cocycle  $c : \mathcal{G} \rightarrow \mathbb{R}^n$  which is *exact* in the sense of [65, Definition 3.3]. We will assume  $\mathcal{G}$  is étale, so that in this higher-dimensional setting exactness entails that  $\text{Ker}(c)$  admits a Haar system and the map

$$r \times c : \mathcal{G} \rightarrow \mathcal{G}^{(0)} \times \mathbb{R}^n, \quad \xi \mapsto (r(\xi), c(\xi)),$$

is a quotient map onto its image.

Given  $\mathcal{H} = \text{Ker}(c)$  a closed subgroupoid of  $\mathcal{G}$ , we will construct a  $KK$ -cycle from  $c$  supported on the module  $E_{C_r^*(\mathcal{H}, \sigma)}$  constructed in the previous section. We use the representation of  $C_c(\mathcal{G}, \sigma)$  on  $E_{C_r^*(\mathcal{H}, \sigma)}$  by left multiplication,  $\pi(f_1)f_2 = f_1 * f_2$  for  $f_2 \in C_c(\mathcal{G}, \sigma) \subset E_{C_r^*(\mathcal{H}, \sigma)}$ . Again by [67, Theorem 1.4] this action extends to a representation of  $C_r^*(\mathcal{G}, \sigma)$ .

The components of the exact cocycle  $c : \mathcal{G} \rightarrow \mathbb{R}^n$  give  $n$  real cocycles  $c_k(\xi) := (\pi_k \circ c)(\xi)$  by composition with the  $k$ th coordinate projection

$$\pi_k : \mathbb{R}^n \rightarrow \mathbb{R}, \quad x = (x_1, \dots, x_n) \mapsto x_k.$$

Following [79],  $C_r^*(\mathcal{G}, \sigma)$  has  $n$  mutually commuting one-parameter groups of automorphisms  $\{u_t^{(k)}\}_{k=1}^n$ , which on  $C_c(\mathcal{G}, \sigma)$  are given by

$$(u_t^{(k)} f)(\xi) = e^{itc_k(\xi)} f(\xi), \quad t \in \mathbb{R}$$

with  $c_k = \pi_k \circ c$  as above. The generators of these automorphisms are derivations  $\{\partial_j\}_{j=1}^n$  on  $C_c(\mathcal{G}, \sigma)$ , where  $(\partial_j f)(\xi) = c_j(\xi)f(\xi)$  (pointwise multiplication). We denote by  $D_{c_j}$  the extension of these derivations to an unbounded operator on  $E_{C_r^*(\mathcal{H}, \sigma)}$ .

We use this differential structure to define an unbounded operator that plays the role of an elliptic differential operator. Our construction mimics the construction of the elements  $\alpha$  and  $\beta$  in [44, Section 5] and, as such, uses the exterior algebra  $\bigwedge^* \mathbb{R}^n$ . We briefly establish our Clifford algebra notation,

where  $Cl_{r,s}$  is the (real)  $\mathbb{Z}_2$ -graded  $C^*$ -algebra generated by the mutually anti-commuting odd elements  $\{\gamma^j\}_{j=1}^r, \{\rho^k\}_{k=1}^s$  such that

$$(\gamma^j)^2 = 1, \quad (\gamma^j)^* = \gamma^j, \quad (\rho^k)^2 = -1, \quad (\rho^k)^* = -\rho^k.$$

The exterior algebra  $\bigwedge^* \mathbb{R}^n$  has representations of  $Cl_{0,n}$  and  $Cl_{n,0}$  with generators

$$\rho^j(\omega) = e_j \wedge \omega - \iota(e_j)\omega, \quad \gamma^j(\omega) = e_j \wedge \omega + \iota(e_j)\omega,$$

where  $\{e_j\}_{j=1}^n$  is the standard basis of  $\mathbb{R}^n$  and  $\iota(\nu)\omega$  is the contraction of  $\omega$  along  $\nu$ . One readily checks that  $\rho^j$  and  $\gamma^j$  mutually anti-commute and generate representations of  $Cl_{0,n}$  and  $Cl_{n,0}$ , respectively. An analogous construction holds in the complex case where  $\text{End}_{\mathbb{C}}(\bigwedge^* \mathbb{C}^n) \cong Cl_n \hat{\otimes} Cl_n$ , where the two representations graded-commute.

**Proposition 1.13.** *Let  $\mathcal{G}$  be an étale groupoid and  $c : \mathcal{G} \rightarrow \mathbb{R}^n$  an exact cocycle with kernel  $\mathcal{H}$ . The triple*

$${}_n\lambda_{\mathcal{H}}^c = \left( C_c(\mathcal{G}, \sigma) \hat{\otimes} Cl_{0,n}, E_{C_r^*(\mathcal{H}, \sigma)} \hat{\otimes} \bigwedge^* \mathbb{R}^n, D_c = \sum_{j=1}^n D_{c_j} \hat{\otimes} \gamma^j \right)$$

is an unbounded real Kasparov module for  $(C_r^*(\mathcal{G}, \sigma), C_r^*(\mathcal{H}, \sigma))$ . If we use complex algebras and  $\bigwedge^* \mathbb{C}^n$ , the Kasparov module is complex.

*Proof.* The essential self-adjointness and regularity of  $D$  follow since the subset

$$C_c(\mathcal{G}, \sigma) \hat{\otimes} \bigwedge^* \mathbb{R}^n \subset E_{C_r^*(\mathcal{H}, \sigma)} \hat{\otimes} \bigwedge^* \mathbb{R}^n,$$

is a core for  $D$  and  $D^2 = c^2 \hat{\otimes} 1_{\bigwedge^* \mathbb{R}^n}$  with  $(c^2 f)(\xi) = (c_1(\xi)^2 + \dots + c_n(\xi)^2) f(\xi)$ . Therefore,  $1 + D^2$  has dense range. We note that in particular

$$(1 + D^2)^{-1} = (1 + c^2)^{-1} \hat{\otimes} 1_{\bigwedge^* \mathbb{R}^n}.$$

Using exactness of  $c$ , the same argument as [65, Theorem 3.9] can now be applied to show that  $(1 + D^2)^{-1}$  is compact in  $E_{C_r^*(\mathcal{H})}$ . For  $f \in C_c(\mathcal{G}, \sigma)$ , a simple computation using the regular representation gives that

$$[D_c, \pi(f)] = \sum_{j=1}^n [D_{c_j}, \pi(f)] \hat{\otimes} \gamma^j = \sum_{j=1}^n \pi(\partial_j f) \hat{\otimes} \gamma^j$$

which is adjointable as  $C_c(\mathcal{G}, \sigma)$  is invariant under the derivations  $\{\partial_j\}_{j=1}^n$ .  $\square$

We remark that there is additional structure on the  $KK$ -cycles constructed in Proposition 1.13. Namely, using the action of  $\text{Spin}_{n,0}$  or  $\text{Spin}_{0,n}$  on  $\bigwedge^* \mathbb{R}^n$  defined in [44, §2.18] and using the notation from [44, §5], the unbounded  $KK$ -cycle  ${}_n\lambda_{\mathcal{H}}^c$  determines a class in the equivariant Kasparov group  $KKO_{\text{Spin}_n}^{\mathbb{R}^n}(C_r^*(\mathcal{G}, \sigma), C_r^*(\mathcal{H}, \sigma))$  or  $KK_{\text{Spin}_n}^{\mathbb{C}^n}(C_r^*(\mathcal{G}, \sigma), C_r^*(\mathcal{H}, \sigma))$ . We can then restrict the  $C^*$ -module  $E_{C_r^*(\mathcal{H}, \sigma)} \hat{\otimes} \bigwedge^* \mathbb{R}^n$  to the irreducible spinor representation space. By [44, §5, Lemma 1], this restriction gives an isomorphism of  $KK$ -groups.

For concreteness, we write out the unbounded representatives of the spinor Kasparov modules explicitly. Given  $n$ , we fix an irreducible complex

or real representation space  $S_n^{\mathbb{C}}$  and  $S_n$  of  $\mathbb{C}l_n$  and  $Cl_{n,0}$  respectively. We also set  $K_n = \mathbb{R}, \mathbb{C}$  or  $\mathbb{H}$  to be the maximal commuting subalgebra for the irreducible real representation of  $Cl_{n,0}$  on  $S_n$  (see [61, Chapter I, §5]).

**Proposition 1.14.** *Let  $\mathcal{G}$  be an étale groupoid and  $c : \mathcal{G} \rightarrow \mathbb{R}^n$  an exact cocycle and  $\mathcal{H} := \text{Ker}(c)$ . Then the triple*

$${}_n\lambda_{\mathcal{H}}^{S_c^{\mathbb{C}}} = \left( C_c(\mathcal{G}, \sigma), E_{C_r^*(\mathcal{H}, \sigma)} \hat{\otimes} S_n^{\mathbb{C}}, \sum_{j=1}^n D_{c_j} \hat{\otimes} \gamma^j \right)$$

is a complex Kasparov module of parity  $n \bmod 2$ .

Let  $S_n$  be an irreducible representation space of  $Cl_{n,0}$ . If  $n \not\equiv 1 \bmod 4$ , then

$${}_n\lambda_{\mathcal{H}}^S = \left( C_c(\mathcal{G}, \sigma), (E \hat{\otimes} S_n)_{C_r^*(\mathcal{H}, \sigma) \hat{\otimes} K_n}, \sum_{j=1}^n D_{c_j} \hat{\otimes} \gamma^j \right),$$

$$K_n = \begin{cases} \mathbb{R}, & n = 0, 2 \bmod 8, \\ \mathbb{C}, & n = 3 \bmod 4, \\ \mathbb{H}, & n = 4, 6 \bmod 8 \end{cases}$$

is a real graded unbounded Kasparov module. If  $n \equiv 1 \bmod 4$ , then

$${}_n\lambda_{\mathcal{H}}^S = \left( C_c(\mathcal{G}, \sigma) \hat{\otimes} Cl_{0,1}, \begin{pmatrix} E_{C_r^*(\mathcal{H}, \sigma)} \otimes S_n \\ E_{C_r^*(\mathcal{H}, \sigma)} \otimes S_n \end{pmatrix}_{K_n}, \left( \sum_{j=1}^n D_{c_j} \otimes \gamma^j \right) \hat{\otimes} \sigma_1 \right),$$

$$K_n = \begin{cases} \mathbb{R}, & n = 1 \bmod 8, \\ \mathbb{H}, & n = 5 \bmod 8 \end{cases}$$

is an unbounded Kasparov module, where the left action of  $Cl_{0,1}$  is generated by  $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ .

The spinor Kasparov modules have the advantage that the left algebra is no longer graded, which is useful if we wish to apply the local index formula (for complex semifinite spectral triples constructed from  ${}_n\lambda_{\mathcal{H}}^{S_c^{\mathbb{C}}}$ ). We will predominantly work with the ‘oriented’ Kasparov module  ${}_n\lambda_{\mathcal{H}}^c$  and class  $[{}_n\lambda_{\mathcal{H}}^c] \in KK^n(C_r^*(\mathcal{G}, \sigma), C_r^*(\mathcal{H}, \sigma))$  (real or complex) as the representations are more tractable and we can work in the real or complex category interchangeably. Though we emphasise that at the level of  $K$ -groups (and up to a possible normalisation), there is no loss of information working with either the spin or oriented  $KK$ -cycles.

### 1.6. The Extension Class of an $\mathbb{R}$ -Valued Cocycle

Consider the Kasparov module from Proposition 1.13. In case  $n = 1$  we obtain an ungraded Kasparov module to which we can associate an extension of  $C^*$ -algebras with positive semisplitting.

In this section we fix  $\varepsilon > 0$  and a continuous non-decreasing function  $\chi_+ : \mathbb{R} \rightarrow \mathbb{R}$  satisfying

$$\chi_+(x) := \begin{cases} 0 & \text{if } x \leq -\varepsilon \\ 1 & \text{if } x \geq 0. \end{cases}$$

**Lemma 1.15.** *The operator*

$$\Pi_c : C_c(\mathcal{G}, \sigma) \rightarrow C_c(\mathcal{G}, \sigma), \quad \Pi_c f(\eta) := \chi_+(c(\eta))f(\eta),$$

extends to a self-adjoint operator  $\Pi_c \in \text{End}_{C_r^*(\mathcal{H}, \sigma)}^*(E)$  with  $\|\Pi_c\| \leq 1$ . For all  $f \in C_c(\mathcal{G}, \sigma)$  it holds that  $\pi(f)(\Pi_c^2 - \Pi_c) \in \mathbb{K}(E)$ .

*Proof.* The operator  $D_c$  is self-adjoint and regular in  $E_{C_r^*(\mathcal{H}, \sigma)}$  and  $\Pi_c := \chi_+(D)$  as defined by the continuous functional calculus. It follows that  $\Pi_c$  is a self-adjoint operator on  $E_{C_r^*(\mathcal{H}, \sigma)}$ .

The action of  $\Pi_c^2 - \Pi_c$  is implemented by the function  $K_c \in C_c(\mathcal{G} \times \mathcal{G}/\mathcal{H}, \sigma)$

$$K_c(\xi, [\eta]) = (\chi_c(c(\eta))^2 - \chi_c(c(\eta)))f(\xi),$$

and thus defines a compact operator. □

The same argument as the proof of Lemma 1.15 shows that the (ungraded) Kasparov module  $(C_r^*(\mathcal{G}, \sigma), E_{C_r^*(\mathcal{H}, \sigma)}, 2\Pi_c - 1)$  represents the same class in  $KKO^1(C_r^*(\mathcal{G}, \sigma), C_r^*(\mathcal{H}, \sigma))$  (or complex) as the bounded representative of  $(C_c(\mathcal{G}, \sigma) \hat{\otimes} Cl_{0,1}, E_{C_r^*(\mathcal{H}, \sigma)} \hat{\otimes} \wedge^* \mathbb{R}, D_c \hat{\otimes} \gamma^1)$ .

In order to construct an extension of  $C_r^*(\mathcal{G}, \sigma)$  by  $C_r^*(\mathcal{G} \times \mathcal{G}/\mathcal{H})$ , which is Morita equivalent to  $C_r^*(\mathcal{H}, \sigma)$ , we need to consider the Busby invariant. To this end, we first note the following result on étale groupoids.

**Proposition 1.16** ([79], Chapter II, Proposition 4.2 and [89], Proposition 3.3.3). *Let  $\mathcal{G}$  be an étale groupoid with a fixed 2-cocycle  $\sigma$ . The identity map  $C_c(\mathcal{G}, \sigma) \rightarrow C_c(\mathcal{G}, \sigma)$  extends to a continuous injection  $j : C_r^*(\mathcal{G}, \sigma) \rightarrow C_0(\mathcal{G})$ . For  $a \in C_r^*(\mathcal{G})$  the map  $j$  is given by*

$$j_{\mathcal{G}}(a)(\eta) := (\pi(a)\delta_{s(\eta)} \mid \delta_{\eta})_{C_0(\mathcal{G}^{(0)})}(r(\eta)), \tag{2}$$

where  $\delta_{\eta} \in C_c(\mathcal{G}, \sigma)$  is any function for which

$$r : \text{supp}(\delta_{\eta}) \rightarrow r(\text{supp}(\delta_{\eta})), \quad s : \text{supp}(\delta_{\eta}) \rightarrow s(\text{supp}(\delta_{\eta})),$$

are homeomorphisms and  $\delta_{\eta}(\eta) = 1$  on a neighbourhood of  $\eta$ .

**Definition 1.17.** Let  $c : \mathcal{G} \rightarrow \mathbb{R}$  be a continuous cocycle. We say that  $c$  is *r-unbounded* if for all  $x \in \mathcal{G}^{(0)}$  and all  $M > 0$  there exists  $\eta \in r^{-1}(x)$  for which  $c(\eta) > M$ .

Recall that the *Calkin algebra* of a  $C^*$ -module  $E$  over a  $C^*$ -algebra  $B$  is the quotient  $\mathcal{Q}(E_B) := \text{End}_B^*(E)/\mathbb{K}(E_B)$ . We denote by  $q : \text{End}_B^*(E) \rightarrow \mathcal{Q}(E_B)$  the quotient map. Lastly, we denote by  $\mathcal{M}(B)$  the multiplier algebra of  $B$ .

**Proposition 1.18.** *Let  $c : \mathcal{G} \rightarrow \mathbb{R}$  be an exact cocycle on a Hausdorff étale groupoid  $\mathcal{G}$ ,  $\mathcal{H} = \text{Ker}(c)$  and  $\sigma$  a 2-cocycle on  $\mathcal{G}$ . If  $c$  is  $r$ -unbounded, then the  $*$ -homomorphism*

$$\varphi : C_r^*(\mathcal{G}, \sigma) \rightarrow \mathcal{Q}(E_{C_r^*(\mathcal{H}, \sigma)}), \quad \varphi(a) = q(\Pi_c a \Pi_c)$$

is injective.

*Proof.* Using Proposition 1.16, we can view elements of  $C_r^*(\mathcal{G} \times \mathcal{G}/\mathcal{H}, \sigma)$  as  $C_0$ -functions on  $\mathcal{G} \times \mathcal{G}/\mathcal{H}$ . Let  $E = E_{C_r^*(\mathcal{H}, \sigma)}$  and  $F = F_{\mathcal{G}/\mathcal{H}}$  the  $C^*$ -module over the unit space of  $\mathcal{G} \times \mathcal{G}/\mathcal{H}$ . Then

$$\text{End}_{C_r^*(\mathcal{H}, \sigma)}^*(E) = \mathcal{M}(\mathbb{K}(E_{C_r^*(\mathcal{H}, \sigma)})) = \mathcal{M}(C_r^*(\mathcal{G} \times \mathcal{G}/\mathcal{H}, \sigma)).$$

Since the representation of  $C_r^*(\mathcal{G} \times \mathcal{G}/\mathcal{H}, \sigma)$  on  $F$  is essential, we see that  $\text{End}_{C_r^*(\mathcal{H}, \sigma)}^*(E)$  acts on  $F$ . Thus, if  $j = j_{\mathcal{G} \times \mathcal{G}/\mathcal{H}}$  and  $T \in \text{End}_{C_r^*(\mathcal{H}, \sigma)}^*(E)$ , then formula (2) defines a continuous function  $j(T) : \mathcal{G} \times \mathcal{G}/\mathcal{H} \rightarrow \mathbb{C}$ . The functions  $\delta_{(\xi, [\eta])}$  can be chosen so that  $\|\delta_{(\xi, [\eta])}\|_F \leq 1$ , so we obtain the pointwise estimate

$$\begin{aligned} |j(T)(\xi, [\eta])| &= |(\pi(T)\delta_{[\eta]} | \delta_{(\xi, [\eta])})_{C_0(\mathcal{G}/\mathcal{H})}([\xi\eta])| \leq \|(\pi(T)\delta_{[\eta]} | \delta_{(\xi, [\eta])})\|_{C_0(\mathcal{G}/\mathcal{H})} \\ &\leq \|T\|_{\text{End}_{C_r^*(\mathcal{H}, \sigma)}^*(E)} \|\delta_{[\eta]}\|_F \|\delta_{(\xi, [\eta])}\|_F \leq \|T\|. \end{aligned}$$

In particular, if  $T_n \rightarrow T$  in norm in  $\text{End}_{C_r^*(\mathcal{H}, \sigma)}^*(E)$  then  $j(T_n) \rightarrow j(T)$  pointwise on  $\mathcal{G} \times \mathcal{G}/\mathcal{H}$ .

Suppose that  $a \neq 0 \in C_r^*(\mathcal{G}, \sigma)$  and choose  $\xi \in \mathcal{G}$  with  $|j(a)(\xi)| \geq 3\delta > 0$ . Choose  $f \in C_c(\mathcal{G}, \sigma)$  with  $\|f - a\|_{C_r^*(\mathcal{G}, \sigma)} < \delta$  and  $|f(\xi)| \geq 2\delta$ . Then for every  $(\xi, [\eta]) \in \mathcal{G} \times \mathcal{G}/\mathcal{H}$  it holds that

$$|j(\Pi_c(f - a)\Pi_c)(\xi, [\eta])| \leq \|\Pi_c(f - a)\Pi_c\| \leq \|f - a\| < \delta.$$

For  $f \in C_c(\mathcal{G})$  it holds that

$$j(\Pi_c f \Pi_c)(\xi, [\eta]) = \chi_+(c(\xi\eta))\chi_+(c(\eta))f(\xi).$$

Thus, for all  $[\eta] = (r(\eta), c(\eta))$  satisfying  $c(\eta) \geq \max\{0, -c(\xi)\}$  we estimate

$$\begin{aligned} |j(\Pi_c a \Pi_c)(\xi, [\eta])| &\geq |j(\Pi_c f \Pi_c)(\xi, \eta)| - |j(\Pi_c(f - a)\Pi_c)(\xi, [\eta])| \\ &= |f(\xi)| - |j(\Pi_c(f - a)\Pi_c)(\xi, [\eta])| \\ &\geq |f(\xi)| - \|\Pi_c(f - a)\Pi_c\| \\ &\geq |f(\xi)| - \|f - a\| > \delta. \end{aligned}$$

Since  $c$  is exact, there is a homeomorphism

$$\mathcal{G}/\mathcal{H} \rightarrow \{(r(\xi), c(\xi)) : \xi \in \mathcal{G}\} \subset \mathcal{G}^{(0)} \times \mathbb{R},$$

where the latter set carries the relative topology. Since  $c$  is  $r$ -unbounded, for fixed  $\xi$  there is a non-compact set of pairs  $(\xi, [\eta]) \in \mathcal{G} \times \mathcal{G}/\mathcal{H}$  with  $|j(\Pi_c a \Pi_c)(\xi, [\eta])| > \delta$ . Therefore,  $j(\Pi_c a \Pi_c) \notin C_0(\mathcal{G} \times \mathcal{G}/\mathcal{H})$  and

$$\Pi_c a \Pi_c \notin C_r^*(\mathcal{G} \times \mathcal{G}/\mathcal{H}, \sigma) = \mathbb{K}_{C_r^*(\mathcal{H}, \sigma)}(E).$$

This is equivalent to the statement that the map

$$\varphi : C_r^*(\mathcal{G}, \sigma) \rightarrow \mathcal{Q}(E_{C_r^*(\mathcal{H}, \sigma)}), \quad \varphi(a) = q(\Pi_c a \Pi_c),$$

is injective. □



Using the isomorphism  $\mathbb{K}(E_{C_r^*(\mathcal{H},\sigma)}) \cong C_r^*(\mathcal{G} \ltimes \mathcal{G}/\mathcal{H}, \sigma)$  and the injectivity of  $\varphi$ , we construct the generalised Toeplitz extension

$$0 \rightarrow C_r^*(\mathcal{G} \ltimes \mathcal{G}/\mathcal{H}, \sigma) \rightarrow C^*(\Pi_c C_r^*(\mathcal{G}, \sigma) \Pi_c, C_r^*(\mathcal{G} \ltimes \mathcal{G}/\mathcal{H}, \sigma)) \rightarrow C_r^*(\mathcal{G}, \sigma) \rightarrow 0$$

with completely positive semisplitting  $a \mapsto \Pi_c a \Pi_c$  and Busby invariant  $\varphi$ . The algebra

$$\mathcal{T} = C^*(\Pi_c C_r^*(\mathcal{G}, \sigma) \Pi_c, C_r^*(\mathcal{G} \ltimes \mathcal{G}/\mathcal{H}, \sigma))$$

is represented on  $\Pi_c E_{C_r^*(\mathcal{H},\sigma)}$ .

## 2. Delone Sets and the Transversal Groupoid

We briefly summarise the construction of a groupoid of an aperiodic hull. Results and further details can be found in [2, 12, 14, 47, 48, 50]. We most closely follow the perspective of [12, 14] and construct a dynamical system and transversal groupoid from the topology of point measures in  $\mathbb{R}^d$ .

### 2.1. Delone Dynamical Systems

**Definition 2.1.** Let  $\mathcal{L} \subset \mathbb{R}^d$  be discrete and infinite and fix  $0 < r < R$ .

- (1)  $\mathcal{L}$  is  $r$ -uniformly discrete if  $|B(x; r) \cap \mathcal{L}| \leq 1$  for all  $x \in \mathbb{R}^d$ .
- (2)  $\mathcal{L}$  is  $R$ -relatively dense if  $|B(x; R) \cap \mathcal{L}| \geq 1$  for all  $x \in \mathbb{R}^d$ .

An  $r$ -uniformly discrete and  $R$ -relatively dense set  $\mathcal{L}$  is called an  $(r, R)$ -Delone set.

We will occasionally want extra structure on our Delone set.

**Definition 2.2.** Let  $\mathcal{L} \subset \mathbb{R}^d$  be discrete and infinite.

- (1) A patch of radius  $R > 0$  of  $\mathcal{L}$  is a subset of  $\mathbb{R}^d$  of the form  $(\mathcal{L} - x) \cap B(0; R)$ , for some  $x \in \mathcal{L}$ . If for all  $R > 0$  the set of its patches of radius  $R$  is finite, then  $\mathcal{L}$  has finite local complexity.
- (2) We call  $\mathcal{L}$  repetitive if given any finite subset  $p \subset \mathcal{L}$  and  $\varepsilon > 0$ , there is an  $R > 0$  such that in any ball  $B(x; R)$  there is a subset  $p' \subset \mathcal{L} \cap B(x; R)$  that is a translation of  $p$  within the distance  $\varepsilon$ ; that is, there is an  $a \in \mathbb{R}^d$  such that the Hausdorff distance between  $p'$  and  $p + a$  is less than  $\varepsilon$ .
- (3) We call  $\mathcal{L}$  aperiodic if there is no  $x \neq 0 \in \mathbb{R}^d$  such that  $\mathcal{L} - x = \mathcal{L}$ .

There is an equivalence between discrete sets and point measures in  $\mathbb{R}^d$ . Let  $\mathcal{M}(\mathbb{R}^d)$  denote the space of measures on  $\mathbb{R}^d$  and consider

$$\begin{aligned}
 QD(\mathbb{R}^d) &= \{ \nu \in \mathcal{M}(\mathbb{R}^d) : \forall x \in \mathbb{R}^d, \nu \text{ is pure point and } \nu(\{x\}) \in \mathbb{N} \}, \\
 UD_r(\mathbb{R}^d) &= \{ \nu \in QD(\mathbb{R}^d) : \forall x \in \mathbb{R}^d, \nu(B(x; r)) \leq 1 \}.
 \end{aligned}$$

For  $\nu \in QD(\mathbb{R}^d)$ ,  $\mathcal{L}^{(\nu)} = \text{supp}(\nu)$  is discrete. Similarly for a discrete set  $\mathcal{L}$ , we can define a measure  $\delta_{\mathcal{L}} = \sum_{x \in \mathcal{L}} \delta_x \in QD(\mathbb{R}^d)$ , where  $\delta_x$  is the point measure. We can also relate measures and Delone sets.

**Proposition 2.3.** Let  $\nu \in UD_r(\mathbb{R}^d)$  be a measure such that for all  $x \in \mathbb{R}^d$ ,  $\nu(B(x; R)) \geq 1$ . Then  $\mathcal{L}^{(\nu)}$  is an  $(r, R)$ -Delone set.

As  $\mathcal{M}(\mathbb{R}^d)$  is a subspace of  $C_c(\mathbb{R}^d)^*$ , it can be given the weak  $*$ -topology.

**Proposition 2.4** ([14], Theorem 1.5). *The set  $UD_r(\mathbb{R}^d)$  is a compact subspace of  $\mathcal{M}(\mathbb{R}^d)$ .*

**Proposition 2.5** (cf. [59], Sect. 3, [33], Chapter 1). *The set of  $(r, R)$ -Delone sets,  $\text{Del}_{(r,R)}$ , is a compact and metrizable space. Let  $d_H$  denote the Hausdorff distance between sets. A neighbourhood base at  $\omega \in \text{Del}_{(r,R)}$  is given by the sets*

$$U_{\epsilon, M}(\omega) = \{ \eta \in \text{Del}_{(r,R)} : d_H(\mathcal{L}^{(\omega)} \cap B(0; M), \mathcal{L}^{(\eta)} \cap B(0; M)) < \epsilon \}$$

with  $M, \epsilon > 0$ .

The translation action on  $\mathbb{R}^d$  gives an action on  $C_c(\mathbb{R}^d)$  and thus an action on  $UD_r(\mathbb{R}^d)$ , where

$$(T_a \nu)(f) = \nu(T_{-a} f), \quad (T_{-a} f)(x) = f(x - a), \quad f \in C_c(\mathbb{R}^d).$$

As expected, the  $\mathbb{R}^d$ -action on  $UD_r(\mathbb{R}^d)$  induces an  $\mathbb{R}^d$ -action on the discrete lattices  $\mathcal{L}^{(\nu)}$  by translation,  $T_a(\mathcal{L}^{(\nu)}) = \mathcal{L}^{(\nu)} + a$ .

**Definition 2.6** (cf. [10], Sect. 2, [14], Definition 1.7). Let  $\mathcal{L}$  be a uniformly discrete subset of  $\mathbb{R}^d$ . The *continuous hull* of  $\mathcal{L}$  is the dynamical system  $(\Omega_{\mathcal{L}}, \mathbb{R}^d, T)$ , where  $\Omega_{\mathcal{L}}$  is the closure of the orbit of  $\nu \in UD_r(\mathbb{R}^d)$  such that  $\text{supp}(\nu) = \mathcal{L}$ .

We note that  $\Omega_{\mathcal{L}}$  is compact by Proposition 2.4. The translation action on  $UD_r(\mathbb{R}^d)$  gives the family of homeomorphisms  $\{T_a\}_{a \in \mathbb{R}^d}$  on  $\Omega_{\mathcal{L}}$ . Thus, starting from a Delone set  $\mathcal{L}$ , we may associate with it a continuous topological dynamical system  $(\Omega_{\mathcal{L}}, T, \mathbb{R}^d)$ . This dynamical system is minimal if and only if the lattice  $\mathcal{L}$  is repetitive [14, Theorem 2.13].

*Example 2.7.* Let  $\mathcal{L}$  be a periodic and cocompact group  $G$ , then it is immediate that  $\Omega_{\mathcal{L}} \cong \mathbb{R}^d/G$ . This is the classical picture with no aperiodicity or disorder on our lattice. We can use Rieffel induction on the  $C^*$ -dynamical system to simplify the crossed product algebra

$$C(\Omega_{\mathcal{L}}) \rtimes \mathbb{R}^d \cong C(\mathbb{R}^d/G) \rtimes \mathbb{R}^d \cong C^*(G) \otimes \mathbb{K},$$

which then implies that, for  $\mathcal{L} = \mathbb{Z}^d$ ,  $K_*(C(\Omega_{\mathcal{L}}) \rtimes \mathbb{R}^d) \cong K^*(\mathbb{T}^d)$ . Considering applications to topological phases, we see that for periodic lattices the dynamics of the hull reproduces the  $K$ -theoretic topological phase of the Bloch bundle over the Brillouin torus.

There is a loose equivalence between Delone sets and tilings of  $\mathbb{R}^d$ , where much of the terminology we use was originally formulated [2, 33, 47, 50].

**Definition 2.8.** A tile of  $\mathbb{R}^d$  is a compact subset of  $\mathbb{R}^d$  that is homeomorphic to the closed unit ball. A tiling of  $\mathbb{R}^d$  is a covering of  $\mathbb{R}^d$  by a family of tiles whose interiors are pairwise disjoint.

Given a tiling  $\mathcal{T}$  with a uniform minimum and maximum bound on the radius of each tile, we can choose a point from the interior of every tile to

obtain a Delone set  $\mathcal{L}_{\mathcal{T}}$ . There is also an explicit passage from Delone sets to tilings via the Voronoi tiling.

**Definition 2.9.** Let  $\mathcal{L}$  be an  $(r, R)$ -Delone set in  $\mathbb{R}^d$ . The Voronoi tile around a point  $x \in \mathcal{L}$  is the set

$$V_x = \{y \in \mathbb{R}^d : \|y - x\| \leq \|y - x'\| \text{ for all } x' \in \mathcal{L}\}.$$

The Voronoi tiling  $\mathcal{V}$  associated with  $\mathcal{L}$  is the family  $\{V_x\}_{x \in \mathcal{L}}$ .

*Remark 2.10* (A note on topologies). Given a Delone set, one may instead consider the corresponding Voronoi tiling. If each tile in the Voronoi tiling comes from a finite collection of prototiles, there is a canonical tiling space with tiling metric (cf. [84, Chapter 1]). The topology of the tiling space is strictly finer than the topology coming from the weak-\* topology on the space of Delone sets. However, if the Delone set is repetitive and has finite local complexity, then the topologies are equivalent, see [12, Section 2] and [50].

We will mostly work under the assumption that  $\mathcal{L}$  is  $(r, R)$ -Delone only. Therefore, if one wishes to apply our work to tilings, one should also assume that  $\mathcal{L}$  is repetitive and has finite local complexity.

### 2.2. The Transversal Groupoid

The notion of an abstract transversal in a groupoid allows one to replace a topological groupoid by a smaller subgroupoid, up to Morita equivalence.

**Definition 2.11.** A topological groupoid  $\mathcal{F}$  admits an *abstract transversal* if there is a closed subset  $X \subset \mathcal{F}^{(0)}$  such that

- (1)  $X$  meets every orbit of the  $\mathcal{F}$ -action on  $\mathcal{F}^{(0)}$ ;
- (2) for the relative topologies on  $X$  and

$$\mathcal{F}_X := \{\xi \in \mathcal{F} : r(\xi) \in X\} \subset \mathcal{F},$$

the restrictions  $r : \mathcal{F}_X \rightarrow X$  and  $s : \mathcal{F}_X \rightarrow \mathcal{F}^{(0)}$  are open maps for the relative topologies on  $\mathcal{F}_X$  and  $X$ .

The set  $\mathcal{G} := \mathcal{F}_X \cap \mathcal{F}_X^{-1}$  is a closed subgroupoid and  $\mathcal{F}_X$  is a groupoid equivalence between  $\mathcal{F}$  and  $\mathcal{G}$  (with its relative topology), see [71, Example 2.7]. Abstract transversals were studied more generally in [75, Section 3]. We will describe an abstract transversal  $\mathcal{G} \subset \Omega_{\mathcal{L}} \rtimes \mathbb{R}^d$  which is Hausdorff and étale in the relative topology.

**Definition 2.12.** The transversal of a lattice  $\mathcal{L}$  is given by the set

$$\Omega_0 = \{\omega \in \Omega_{\mathcal{L}} : 0 \in \mathcal{L}^{(\omega)}\},$$

We see that  $\Omega_0$  is a closed subset of  $\Omega_{\mathcal{L}}$  and so is compact by Proposition 2.4.

**Proposition 2.13** ([14], Proposition 2.3, [12], Proposition 2.24). *Let  $\mathcal{L}$  be a Delone set.*

- (1) *If  $\mathcal{L}$  has finite local complexity, then  $\Omega_0$  is totally disconnected.*
- (2) *If  $\mathcal{L}$  is repetitive, aperiodic and of finite local complexity, then  $\Omega_0$  is a Cantor set (totally disconnected with no isolated points).*

The passage from the continuous hull  $\Omega_{\mathcal{L}}$  to the transversal  $\Omega_0$  discretises the  $\mathbb{R}^d$ -action at the cost that we no longer have a group action, but only a groupoid structure.

**Proposition 2.14** ([10], Sect. 3, [48], Lemma 2). *Given a Delone set  $\mathcal{L}$  with transversal  $\Omega_0$ , define the set*

$$\mathcal{G} := \{(\omega, x) \in \Omega_0 \times \mathbb{R}^d : T_{-x}\omega \in \Omega_0\} = \{(\omega, x) \in \Omega_0 \times \mathbb{R}^d : x \in \mathcal{L}^{(\omega)}\}.$$

Then  $\mathcal{G}$  is a Hausdorff étale groupoid with maps

$$\begin{aligned} (\omega, x)^{-1} &= (T_{-x}\omega, -x), & (\omega, x) \cdot (T_{-x}\omega, y) &= (\omega, x + y), \\ s(\omega, x) &= T_{-x}\omega, & r(\omega, x) &= \omega \end{aligned} \tag{3}$$

and unit space  $\mathcal{G}^{(0)} = \Omega_0$ .

The transversal groupoid  $\mathcal{G}$  and its corresponding (twisted)  $C^*$ -algebra will be our central object of study. The space  $\Omega_0$  is an abstract transversal in the sense of Definition 2.11, so that  $\mathcal{G} \subset \Omega_0 \times \mathbb{R}^d$  with its subspace topology is Morita equivalent to  $\Omega_{\mathcal{L}} \rtimes \mathbb{R}^d$ . This result is well known to experts, see [33, Chapter 2, Section 2] for the case of tilings. We find it worthwhile to give a detailed proof in the Delone lattice setting. To this end, we first make the following observation.

**Lemma 2.15.** *Let  $0 < \varepsilon < r/2$ . For any  $\omega \in \Omega_0$ , the intersection  $\mathcal{L}^{(\omega)} \cap B(y; \varepsilon)$  contains at most one point.*

*Proof.* Suppose that the intersection is non-empty and  $x_1, x_2 \in \mathcal{L}^{(\omega)} \cap B(y; \varepsilon)$ . Then  $d(x_1, x_2) < 2\varepsilon < r$  so it must hold that  $x_1 = x_2$ .  $\square$

For  $\mu \in \mathbb{R}_{>0}$  we denote by

$$P_\mu := \{\mathcal{L}^{(\omega)} \cap B(0; \mu) : \omega \in \Omega_0\},$$

the set of patterns of radius  $\mu$ . The sets

$$U_{p,\mu} := \{\omega \in \Omega_{\mathcal{L}} : 0 \in \mathcal{L}^{(\omega)}, \mathcal{L}^{(\omega)} \cap B(0; \mu) = p\} \subset \Omega_0, \quad \mu \in \mathbb{R}_{>0}, p \in P_\mu,$$

define the relative topology on the closed subset  $\Omega_0 := \{\omega \in \Omega_{\mathcal{L}} : 0 \in \mathcal{L}^{(\omega)}\}$ . In case  $\mathcal{L}$  has finite local complexity each set  $P_\mu$  is finite and the clopen sets  $U_{p,\mu}$  determine the totally disconnected topology on  $\Omega_0$ . We now provide the proof that  $\Omega_0$  is indeed an abstract transversal.

**Proposition 2.16.** *Let  $\mathcal{L} \subset \mathbb{R}^d$  be a uniformly  $r$ -discrete subset with transversal  $\Omega_0$  and associated groupoid  $\mathcal{G}$ . For  $U \subset \Omega_0$  an open set, the sets*

$$\begin{aligned} V_{(U,y,\varepsilon)} &:= (U \times B(y; \varepsilon)) \cap \mathcal{G} \\ &= \{(\omega, x) \in \Omega_0 \times \mathbb{R}^d : \omega \in U, x \in \mathcal{L}^{(\omega)} \cap B(y; \varepsilon)\}, \end{aligned}$$

form a base for the topology on  $\mathcal{G}$ . For  $0 < \varepsilon < r/2$ , the restriction  $s : V_{(U,y,\varepsilon)} \rightarrow \Omega_0$  is a homeomorphism onto its image. Moreover, the restrictions

$$s : \Omega_{\mathcal{L}} \rtimes \mathbb{R}^d \cap r^{-1}(\Omega_0) \rightarrow \Omega_{\mathcal{L}}, \quad r : \Omega_{\mathcal{L}} \rtimes \mathbb{R}^d \cap r^{-1}(\Omega_0) \rightarrow \Omega_0,$$

are open maps. Therefore, the set  $\Omega_0$  is an abstract transversal and the groupoid  $\mathcal{G} \subset \Omega_{\mathcal{L}} \rtimes \mathbb{R}^d$ , with the subspace topology, is Morita equivalent to  $\Omega_{\mathcal{L}} \rtimes \mathbb{R}^d$ .

*Proof.* The sets  $V_{(U,y,\varepsilon)}$  generate the relative topology on  $\mathcal{G}$  as a subset of the crossed product groupoid  $\Omega_{\mathcal{L}} \rtimes \mathbb{R}^d$  of the hull of  $\mathcal{L}$ . To see that each of the basic sets is an  $s$ -set, we adapt the proof of [12, Lemma 2.10]. The map  $s$  is injective on  $V_{(U,y,\varepsilon)}$ , for  $(\omega, x), (\eta, z) \in V_{(p,\mu,y,\varepsilon)}$  the equality  $T_{-x}\omega = T_{-z}\eta$  implies that  $\omega = T_{x-z}\eta$ , and  $x - z \in \mathcal{L}^{(\omega)}$ . Now  $x, z \in B(y; \varepsilon)$  so  $d(x, z) < 2\varepsilon < r$ , and thus,  $x = z$  because  $\mathcal{L}^{(\omega)}$  is  $r$ -discrete. It then follows that  $\omega = \eta$  as well. Now consider  $s : (\omega, x) \mapsto T_{-x}\omega$  and the image

$$\begin{aligned} s(V_{(U,y,\varepsilon)}) &= \{\omega \in \Omega_0 : \exists x \in B(y; \varepsilon), T_x\omega \in U\} \\ &= \{\omega \in \Omega_0 : \exists x \in B(0; \varepsilon), T_{x+y}\omega \in U\} \\ &= \Omega_0 \cap T_{-y}(\{\omega \in \Omega_0 : \exists x \in B(0; \varepsilon), T_x\omega \in U\}) \\ &= \Omega_0 \cap T_{-y}(s(U \times B(0; \varepsilon))), \end{aligned}$$

with  $s(\omega, x) = \phi(\omega, -x)$  and  $\phi$  as in [12, Lemma 2.10], and by that result the map  $s$  is a homeomorphism onto its image. Thus, since  $y$  is fixed, the set  $s(V_{(U,y,\varepsilon)})$  is open in  $\Omega_0$ . Now  $\omega \in s(V_{(U,y,\varepsilon)})$  implies that  $B(-y; \varepsilon) \cap \mathcal{L}^{(\omega)} \neq \emptyset$  and thus contains a unique point  $x_\omega^{-y}$ . The map

$$t_y : s(V_{(U,y,\varepsilon)}) \rightarrow V_{(U,y,\varepsilon)}, \quad \omega \mapsto (T_{-x_\omega^{-y}}\omega, -x_\omega^{-y}),$$

is an inverse for  $s$ : If  $\omega = T_{-x}\eta$  with  $\{x\} = B(y; \varepsilon) \cap \mathcal{L}^{(\eta)}$  then

$$x_\omega^{-y} = x_{T_{-x}\eta}^{-y} = B(-y; \varepsilon) \cap \mathcal{L}^{(T_{-x}\eta)} = -x,$$

and so indeed

$$t_y \circ s(\eta, x) = s_y(\omega) = (T_x\omega, x) = (\eta, x).$$

The points  $x_\omega^y$  satisfy the equality  $x_\omega^y = y + x_{T_{-y}\omega}^0$ , and thus, the map  $t_y$  can be written

$$t_y(\omega) = (T_{-x_\omega^{-y}}\omega, -x_\omega^{-y}) = (T_y T_{-x_{T_y\omega}^0}\omega, y - x_{T_y\omega}^0) = (T_y \times T_y) \circ t_0 \circ T_y(\omega).$$

The map  $t_0$  is continuous by [12, Lemma 2.10] and  $y$  is fixed, proving continuity of  $t_y$ .

We now proceed to show the maps  $s, r$  are open when restricted to  $r^{-1}(\Omega_0)$ . As above we have

$$s(U \times B(y; \varepsilon) \cap r^{-1}(\Omega_0)) = \{T_{-x}(\omega) : \omega \in U \cap \Omega_0, x \in B(y; \varepsilon) \cap \mathcal{L}^{(\eta)}\},$$

and to prove that the map  $s|_{r^{-1}(\Omega_0)}$  is open we may restrict ourselves to sets  $U = U_{\delta,M}(\omega) \cap \Omega_0$  and  $M$  sufficiently large,  $\delta$  sufficiently small. It then suffices to show that the set  $s(U \times B(y; \varepsilon) \cap r^{-1}(\Omega_0))$  contains a basic open neighbourhood of any of its elements  $T_{-x}(\omega)$ . Let  $\delta < \varepsilon < r/2$  and  $M > \delta$ . Then if  $\eta \in \Omega_{\mathcal{L}}$  is such that

$$d_H(B(0; M + \|y\| + r) \cap \mathcal{L}^{(T_{-x}\omega)}, B(0; M + \|y\| + r) \cap \mathcal{L}^{(\eta)}) < \delta/2,$$

we have that  $-x \in \mathcal{L}^{(T_{-x}\omega)} \cap B(0; M + \|y\| + r)$ . By definition of the Hausdorff distance, we have

$$\inf_{w \in B(0; M + \|y\|) \cap \mathcal{L}^{(\eta)}} \|w + x\| < \delta/2,$$

and since the sets involved are discrete, there exists a point  $w \in \mathcal{L}^{(\eta)}$  with  $\|w + x\| \leq \delta/2$ . Moreover, if  $\delta < r$  then this point  $w$  is unique because  $\mathcal{L}^{(\eta)}$  is  $r$ -discrete. Then for  $z \in B(0; M) \cap \mathcal{L}^{(\omega)}$  and  $v \in B(0; M) \cap \mathcal{L}^{(T^{-x}\eta)}$  we have  $(z - w) \in B(0; M + \|y\| + r) \cap \mathcal{L}^{(T^{-x}\omega)}$ ,  $(v + x) \in B(0; M + \|y\| + r) \cap \mathcal{L}^{(\eta)}$ , from which we deduce

$$\|z - v\| \leq \|(v + x) - (z - w)\| + \|x + w\| < \delta,$$

and therefore it follows that

$$d_H(B(0; M) \cap \mathcal{L}^{(\omega)}, B(0; M) \cap \mathcal{L}^{(T^{-w}\eta)}) < \delta.$$

Since  $\|w + y\| \leq \|w + x\| + \|x - y\| < \delta < \varepsilon$ , it holds that  $(T_{-w}\eta, -w) \in U \times B(y, \varepsilon)$  and  $0 \in T_{-w}\eta$ . Therefore,  $\eta \in s((U \times B(y, \varepsilon)) \cap r^{-1}(\Omega_0))$  and  $s : r^{-1}(\Omega_0) \rightarrow \Omega_{\mathcal{L}}$  is an open map. The statement that  $r$  is an open map is immediate because  $\Omega_0$  carries the relative topology inherited from  $\Omega_{\mathcal{L}}$ . This completes the proof.  $\square$

From this, we derive several structure statements for the groupoid  $\mathcal{G}$ .

**Proposition 2.17.** *For any  $1 \leq k \leq d$  the groupoid cocycles*

$$\hat{c}_k := (c_1, \dots, c_k) : (\omega, x) \mapsto (x_1, \dots, x_k),$$

*are exact in the sense of [65, Definition 3.3].*

*Proof.* The subspace topology on  $\mathcal{G}$  has a base consisting of the sets

$$(U_{(p,\mu)} \times B(y; \varepsilon)) \cap \mathcal{G} = \{(\omega, x) \in \Omega_0 \times \mathbb{R}^d : \mathcal{L}^{(\omega)} \cap B(0; \mu) = p, \\ x \in \mathcal{L}^{(\omega)} \cap B(y; \varepsilon)\},$$

with  $\mu \in [0, \infty)$ ,  $p \in P_\mu$ ,  $y \in \mathbb{R}^d$  and  $0 < \varepsilon < r/2$ . For  $(\omega, x) \in \mathcal{G}$ , choose  $\mu > \|x\| + r/2$  and let  $p := \mathcal{L}^{(\omega)} \cap B(0; \mu)$ . Consider  $(\eta, z) \in (U_{p,\mu} \times B(x; \varepsilon)) \cap \mathcal{G}$ . Then it holds that  $\|z - x\| < \varepsilon < r/2$  and

$$z, x \in \mathcal{L}^{(\eta)} \cap B(0, \mu) = \mathcal{L}^{(\omega)} \cap B(0, \mu),$$

from which we conclude that  $z = x$ . In particular, each  $\hat{c}_k$  is locally constant and  $\hat{c}_k^{-1}(0)$  is a clopen subgroupoid. Since  $\mathcal{G}$  is étale, counting measures define a Haar system on  $\hat{c}_k^{-1}(0)$ . Exactness of the cocycles  $\hat{c}_k$  entails that the map  $(\omega, x) \mapsto (\omega, \hat{c}_k(x)) = (\omega, x_1, \dots, x_k)$  is a complete quotient map onto its image. This map is equal to the restriction of the map  $\text{id} \times \pi_k$  to  $\mathcal{G}$ , with  $\pi_k : \mathbb{R}^d \rightarrow \mathbb{R}^k$  the projection onto the first  $k$  coordinates, which is a complete quotient map.  $\square$

Note that the above proof applies to any cocycle  $c : \mathcal{G} \rightarrow \mathbb{R}^k$  that factors through the cocycle  $\hat{c}_d : \mathcal{G} \rightarrow \mathbb{R}^d$ . Now that we have characterised the étale topology on  $\mathcal{G}$ , we recall the constructions in Sect. 1 and consider an  $s$ -cover for  $\mathcal{G}$  (Definition 1.8), which will then give a frame for the  $C^*$ -module over the unit space, which we denote  $E_{C(\Omega_0)}$ . We fix a choice of  $0 < \varepsilon < r/2$  and a countable set of points  $Y \subset \mathbb{R}^d$  for which  $B(y; \varepsilon)$  form an open cover of  $\mathbb{R}^d$ . Note that we can choose the set  $Y = \lambda\mathbb{Z}^d$  with  $\lambda > 0$  sufficiently small, which is convenient but not necessary.

**Proposition 2.18.** *Let  $\mathcal{L} \subset \mathbb{R}^d$  be a uniformly discrete subset,  $\mathcal{G}$  the associated groupoid and  $E_{C(\Omega_0)}$  the  $C^*$ -module over the unit space. For any  $0 < \varepsilon < r/2$  and any countable cover  $\{B(y; \varepsilon)\}_{y \in Y}$  the open sets*

$$V_y := V_{(0,0,y,\varepsilon)} = \{(\omega, x) \in \Omega_0 \times \mathbb{R}^d : x \in \mathcal{L}^{(\omega)} \cap B(y; \varepsilon)\},$$

*form an  $s$ -cover for  $\mathcal{G}$ . Any partition of unity  $\chi_y$  subordinate to the cover  $\{B(y; \varepsilon)\}_{y \in Y}$  of  $\mathbb{R}^d$  can be lifted to a partition of unity subordinate to the cover  $V_y$  of  $\mathcal{G}$  via  $\chi_y(\omega, x) = \chi_y(x)$ . Consequently, the functions  $\chi_y : \mathcal{G} \rightarrow \mathbb{R}$  define a frame for  $E_{C(\Omega_0)}$ .*

*Proof.* The sets  $V_y$  form an open cover of  $\mathcal{G}$  because each  $(\omega, x) \in \mathcal{G}$  is an element of  $V_y$  whenever  $x \in B(y; \varepsilon)$  and such  $y$  exists because  $B(y; \varepsilon)$  form an open cover. Moreover, each of the  $V_y$  is an  $s$ -set by Lemma 2.16. The functions  $\chi_y$  define a frame by Proposition 1.11. □

**2.3. The Twisted Groupoid Algebra and Its  $K$ -theory**

Given our transversal groupoid, we fix a normalised 2-cocycle  $\sigma : \mathcal{G}^{(2)} \rightarrow \mathbb{T}$  (or  $\{\pm 1\}$  in the real case). Our central motivation for working with twisted groupoid algebras comes from the following example.

*Example 2.19* (Magnetic twists). For the transversal groupoid, we can encode the action of a magnetic field that twists the translation action of the lattice. Working first with the continuous hull  $\Omega_{\mathcal{L}} \rtimes \mathbb{R}^d$ , we follow [15, Section 2.2] and define a 2-cocycle,

$$\sigma : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathcal{U}(C(\Omega_{\mathcal{L}})), \quad \sigma(x, y) = \exp(-i\Gamma\langle 0, x, x + y \rangle)$$

where  $\Gamma\langle 0, x, x + y \rangle$  is the magnetic flux through the triangle defined by the points  $0, x, x + y \in \mathbb{R}^d$ . The magnetic field need not be constant over  $C(\Omega_{\mathcal{L}})$  and can generally be described by a continuous map  $B : \Omega_{\mathcal{L}} \rightarrow \wedge^2 \mathbb{R}^d$ , where  $\Gamma\langle x, y, z \rangle = \int_{\langle x,y,z \rangle} B_\omega$  and  $\langle x, y, z \rangle \subset \mathbb{R}^{2d}$  is the triangle with corners  $x, y, z \in \mathbb{R}^d$ . If the magnetic field is constant over  $\Omega_{\mathcal{L}}$ , then our general flux equation can be simplified by a skew-symmetric matrix  $B$  with

$$\sigma(x, y) = \exp(-i\langle x, B(x + y) \rangle) = \exp(-i\langle x, By \rangle).$$

Our choice of 2-cocycle on the crossed product  $C(\Omega_{\mathcal{L}}) \rtimes_{\sigma} \mathbb{R}^d$  restricts to a 2-cocycle on the transversal groupoid, which we also denote by  $\sigma$ . Namely, we define

$$\sigma((\omega, x), (T_{-x}\omega, y)) = \exp(-i\Gamma_{\mathcal{L}^{(\omega)}}\langle 0, x, x + y \rangle)$$

where  $\Gamma_{\mathcal{L}^{(\omega)}}\langle 0, x, x + y \rangle$  is the magnetic flux through the triangle defined by the points  $0, x, x + y \in \mathcal{L}^{(\omega)}$ . We note that our twist will always be trivial for  $d = 1$  and is normalised because

$$\sigma((\omega, x), (T_{-x}, -x)) = \exp(-i\Gamma_{\mathcal{L}^{(\omega)}}\langle 0, x, 0 \rangle) = 1.$$

The cocycle condition on  $\sigma$  translates into the condition that for  $x, x + y, x + y + z \in \mathcal{L}^{(\omega)}$ ,

$$\begin{aligned} & \Gamma_{\mathcal{L}(\omega)} \langle 0, x, x + y \rangle + \Gamma_{\mathcal{L}(\omega)} \langle 0, x + y, x + y + z \rangle \\ &= \Gamma_{\mathcal{L}(\omega)} \langle 0, x, x + y + z \rangle + \Gamma_{\mathcal{L}(\tau_{-x}\omega)} \langle 0, y, y + z \rangle, \end{aligned}$$

which follows from Stokes' Theorem and the observation that

$$\Gamma_{\mathcal{L}(\tau_{-x}\omega)} \langle 0, y, y + z \rangle = \Gamma_{\mathcal{L}(\omega)} \langle x, x + y, x + y + z \rangle.$$

Given our groupoid  $\mathcal{G}$  and cocycle  $\sigma$ , we can construct the groupoid  $C^*$ -algebra by the method given in Sect. 1.3, acting on the  $C^*$ -module over the unit space. The  $K$ -theory of the twisted groupoid algebra is used to describe topological phases of gapped Hamiltonians, which we will then pair with  $KK$ -cycles to obtain numerical labels for these phases. In the absence of a 2-cocycle twist, the continuous dynamical system  $(\Omega_{\mathcal{L}}, T, \mathbb{R}^d)$  can be described via the crossed product groupoid  $\Omega_{\mathcal{L}} \rtimes \mathbb{R}^d$ , which is then groupoid equivalent to  $\mathcal{G}$ . Applying the equivalence theorem of [71, 90] and the Connes–Thom isomorphism [27],

$$K_*(C_r^*(\mathcal{G})) \cong K_*(C(\Omega_{\mathcal{L}}) \rtimes \mathbb{R}^d) \cong K_{*-d}(C(\Omega_{\mathcal{L}})) \cong K^{d-*}(\Omega_{\mathcal{L}}),$$

in both real and complex  $K$ -theory. This result remains true for twists by 2-cocycles.

**Proposition 2.20.** *Let  $\mathcal{L}$  be a Delone set and  $\sigma : (\Omega_{\mathcal{L}} \rtimes \mathbb{R}^d)^{(2)} \rightarrow \mathbb{T}$  (or  $\{\pm 1\}$  in the real case) a continuous 2-cocycle. Then the twisted groupoid  $C^*$ -algebra  $C^*(\mathcal{G}, \sigma)$  is Morita equivalent to the twisted crossed product  $C(\Omega_{\mathcal{L}}) \rtimes_{\sigma} \mathbb{R}^d$  and there is an isomorphism  $K_*(C_r^*(\mathcal{G}, \sigma)) \rightarrow K^{d-*}(\Omega_{\mathcal{L}})$ .*

*Proof.* As the 2-cocycle on  $\mathcal{G}$  comes from the restriction of a 2-cocycle on  $\Omega_{\mathcal{L}} \rtimes \mathbb{R}^d$ , we can apply [30, Theorem 9.1], which gives that  $C_r^*(\mathcal{G}, \sigma)$  is Morita equivalent to the twisted crossed product  $C(\Omega_{\mathcal{L}}) \rtimes_{\sigma} \mathbb{R}^d$ . Then, by Packer–Raeburn stabilisation, [73, Section 3], and the Connes–Thom isomorphism, we obtain that

$$\begin{aligned} K_*(C_r^*(\mathcal{G}, \sigma)) &\cong K_*(C(\Omega_{\mathcal{L}}) \rtimes_{\sigma} \mathbb{R}^d) \cong K_*((C(\Omega_{\mathcal{L}}) \otimes \mathbb{K}) \rtimes \mathbb{R}^d) \\ &\cong K_{*-d}(C(\Omega_{\mathcal{L}}) \otimes \mathbb{K}) \cong K^{d-*}(\Omega_{\mathcal{L}}). \end{aligned}$$

Hence, the  $K$ -theory of the twisted groupoid  $C^*$ -algebra reduces to that of the continuous hull  $\Omega_{\mathcal{L}}$ . □

Let us emphasise that the computation of the  $K$ -theory of  $\Omega_{\mathcal{L}}$  is highly non-trivial. A homological description of the  $K$ -theory of  $\Omega_{\mathcal{L}}$  for a large class of tilings with finite local complexity is given in [33] as well as computational techniques. See also the review [43]. In the case that  $\mathcal{L}$  is repetitive, aperiodic and has finite local complexity, one can characterise  $\Omega_{\mathcal{L}}$  as a projective limit [2, 12, 50] and compute its  $K$ -theory using the Pimsner–Voiculescu spectral sequence [87] (adapted from the spectral sequence used by Kasparov [45, §6.10]), whose  $E_2$ -page is isomorphic to the Čech cohomology of  $\Omega_{\mathcal{L}}$  with integer coefficients. In the case of low-dimensional substitution tilings with finite local complexity and a primitive and injective substitution map, Gonçalves–Ramirez–Solano relate the Čech cohomology of  $\Omega_{\mathcal{L}}$  to the  $K$ -theory of the groupoid  $C^*$ -algebra of the unstable equivalence relation on  $\Omega_{\mathcal{L}}$  (note that



this groupoid  $C^*$ -algebra is Morita equivalent to  $C_r^*(\mathcal{G})$  [39, Theorem 2.3]. See [39] for a detailed exposition on these (and other) matters.

In contrast to  $\Omega_{\mathcal{L}}$ , the  $K$ -theory of the transversal  $\Omega_0$  is often very simple to compute. If  $\mathcal{L}$  is a Delone set with finite local complexity, then by Proposition 2.13  $\Omega_0$  is totally disconnected and, by continuity of the  $K$ -functor,  $K_*(C(\Omega_0)) \cong C(\Omega_0, K_*(\mathbb{F}))$ , where  $\mathbb{F} = \mathbb{C}$  or  $\mathbb{R}$ .

**2.3.1. The Bulk  $KK$ -Cycle.** We now introduce our main tool to extract numerical invariants from  $K_*(C_r^*(\mathcal{G}, \sigma))$  (see Sect. 6). The transversal groupoid  $\mathcal{G}$  is étale, and the cocycles  $\hat{c}_k : \mathcal{G} \rightarrow \mathbb{R}^k$ ,  $\hat{c}_k(\omega, x) = (x_1, \dots, x_k)$  are exact by Proposition 2.17. Hence, we can construct a family of unbounded  $KK$ -cycles for  $\mathcal{G}$  by Proposition 1.13.

We call the special case  $c(\omega, x) := \hat{c}_d(\omega, x) = x$ , where  $\text{Ker}(c) \cong \mathcal{G}^{(0)} \cong \Omega_0$ , the bulk  $KK$ -cycle as it spans all dimensions of the lattice, where the terminology is taken from topological phases. Explicitly,

$${}^d\lambda_{\Omega_0} = \left( C_c(\mathcal{G}, \sigma) \hat{\otimes} Cl_{0,d}, E_{C(\Omega_0)} \hat{\otimes} \bigwedge^* \mathbb{R}^d, \sum_{j=1}^d X_j \hat{\otimes} \gamma^j \right), \tag{4}$$

is an unbounded Kasparov module, with  $X_j$  the self-adjoint regular operator  $(X_j f)(\omega, x) = x_j f(\omega, x)$  on  $E_{C(\Omega_0)}$ . We will consider other unbounded  $KK$ -cycles from cocycles on  $\mathcal{G}$  and their properties in Sect. 3.

**2.4. One-Dimensional Delone Sets as Cuntz–Pimsner Algebras**

Given an  $(r, R)$ -Delone set  $\mathcal{L} \subset \mathbb{R}^d$ , we have constructed the groupoid  $\mathcal{G}$  and a class in  $KK^d(C_r^*(\mathcal{G}, \sigma), C(\Omega_0))$  that encodes the translation action on the transversal. For the case  $d = 1$  and trivial cocycle  $\sigma = 1$ , we now give an equivalent description of  $C_r^*(\mathcal{G})$  as a Cuntz–Pimsner algebra. We also find that the Kasparov cycle from Eq. (4) is equivalent to the class of the defining extension of the Cuntz–Pimsner algebra. We remark that a similar construction is done in [95] that includes higher dimensions but for more restrictive substitution tilings. Here we leave open the question of higher dimensions where, in analogy with crossed products by  $\mathbb{Z}^d$ , a description of  $C^*(\mathcal{G}, \sigma)$  as an iterated Cuntz–Pimsner algebra or  $C^*$ -algebra of a product system [92] is a natural aim.

In the case  $d = 1$ , recall the cocycle  $c(\omega, x) = x \in \mathbb{R}$  and write

$$\mathcal{G}^{(0)} = \mathcal{G}_0 := c^{-1}(0), \quad \mathcal{G}_1 := c^{-1}(r, R), \quad \mathcal{G}_{-1} := c^{-1}(-R, -r).$$

**Lemma 2.21.** *Let  $(\omega, x) \in \mathcal{G}$  and  $x > 0$ . There exist  $(\omega_j, x_j) \in \mathcal{G}_1$ ,  $j = 1, \dots, n$  such that*

$$(\omega, x) = \prod_{j=1}^n (\omega_j, x_j),$$

and this decomposition is unique. A similar statement holds for  $(\omega, x)$  with  $x < 0$  where we replace  $\mathcal{G}_1$  with  $\mathcal{G}_{-1}$ .

*Proof.* The lattice  $\mathcal{L}^{(\omega)} \subset \mathbb{R}$  is discrete, so we can order it as

$$\mathcal{L}^{(\omega)} = \{y_n\}_{n \in \mathbb{Z}}, \quad y_0 = 0, \quad y_j < y_{j+1}, \quad r < y_{j+1} - y_j < R. \quad (5)$$

Then  $(\omega, x) = (\omega, y_n)$  for some  $n$  and we set

$$\omega_j := T_{-x_{j-1}}\omega, \quad x_j := y_j - y_{j-1}.$$

It follows that

$$\begin{aligned} (\omega, x) &= (\omega, y_n) = (\omega, y_1) \cdot (T_{-y_1}\omega, y_2 - y_1) \cdots (T_{-y_{n-1}}\omega, y_n - y_{n-1}) \\ &= \prod_{j=1}^n (\omega_j, x_j), \end{aligned}$$

as claimed. Suppose that

$$(\omega, x) = \prod_{j=1}^m (\eta_j, z_j),$$

is another such decomposition and assume without loss of generality that  $m \geq n$ . Then  $\eta_1 = \omega_1 = \omega$ . Since  $z_1, x_1 \in \mathcal{L}^{(\omega)} \cap (r, R)$ , it follows that  $z_1 = x_1$ . This argument can be repeated to find  $\eta_j = \omega_j$  and  $x_j = z_j$  for  $1 \leq j \leq n$ , so the decompositions are the same if  $m = n$ . If  $m > n$ , then

$$(\eta_{n+1}, 0) = (\eta_{n+1}, z_{n+1}) \cdots (\eta_m, z_m) = (\eta_{n+1}, z_{n+1} + \cdots + z_m),$$

so  $0 < z_{n+1} + \cdots + z_m = 0$ , a contradiction. □

The previous result indicates that the one-dimensional transversal groupoid is in some sense generated by  $\mathcal{G}_1 = c^{-1}(r, R)$ . This then gives us a pathway to recharacterise  $C_r^*(\mathcal{G})$  as a Cuntz–Pimsner algebra. The following result comes from standard arguments.

**Lemma 2.22.** *Suppose  $d = 1$  and let  $E_{C(\Omega_0)}^{(r,R)}$  be the completion of  $C_c(\mathcal{G}_1)$  in  $C_r^*(\mathcal{G})$ . Then  $E_{C(\Omega_0)}^{(r,R)}$  is a  $C^*$ -bimodule over  $C(\Omega_0)$  with structure*

$$\begin{aligned} (f_1 \mid f_2)_{C(\Omega_0)}(\omega) &= (f_1^* * f_2)(\omega, 0), & C(\Omega_0)(f_1 \mid f_2)(\omega) &= (f_1 * f_2^*)(\omega, 0), \\ (g_1 \cdot f \cdot g_2)(\omega, x) &= g_1(\omega)f(\omega, x)g_2(T_{-x}\omega). \end{aligned}$$

An analogous result also holds for the completion of  $C_c(\mathcal{G}_{-1})$ .

Denote by  $d : \mathcal{G} \rightarrow \mathbb{Z}$  the map that associates with an element  $(\omega, x)$  the integer  $n$  for which  $x = y_n$  with  $\mathcal{L}^{(\omega)} = \{y_n\}_{n \in \mathbb{Z}}$  as in Eq. (5) on page 1994. We call  $d(\omega, x)$  the *degree* of  $(\omega, x)$ .

**Proposition 2.23.** *The map  $d : \mathcal{G} \rightarrow \mathbb{Z}$  is a continuous 1-cocycle that is unperforated in the sense of [81]. Consequently,  $C_r^*(\mathcal{G})$  is isomorphic to the Cuntz–Pimsner algebra  $\mathcal{O}_{E^{(r,R)}}$  and for  $n > 0$  the sets*

$$\mathcal{G}_{\pm n} := \{\xi_1 \cdots \xi_n : \xi_i \in \mathcal{G}_{\pm 1}\},$$

define a decomposition  $\mathcal{G} = \bigcup_{n \in \mathbb{Z}} \mathcal{G}_n$  into clopen subsets.

*Proof.* We prove that  $d$  is locally constant. Let  $(\omega, x) \in \mathcal{G}$  and choose  $\mu, y$  such that  $x \in B(y; \varepsilon) \subset B(0; \mu)$  with  $\varepsilon < r/2$ . Then  $(\omega, x) \in V_{(p, \mu, y, \varepsilon)}$  for  $p = \mathcal{L}^{(\omega)} \cap B(0; \mu)$  and consider  $(\eta, z) \in V_{(p, \mu, y, \varepsilon)}$ . Since

$$x, z \in B(y; \varepsilon) \subset \mathcal{L}^{(\omega)} \cap B(0; \mu) = \mathcal{L}^{(\eta)} \cap B(0; \mu),$$

and  $\varepsilon < r/2$ , it follows that  $x = z$ . Then since

$$\mathcal{L}^{(\omega)} \cap B(0; \mu) = \mathcal{L}^{(\eta)} \cap B(0; \mu)$$

it follows that  $d(\omega, x) = d(\eta, z)$ . Thus, the degree is locally constant on  $\mathcal{G}$ . By Lemma 2.21 the degree is additive, and it thus defines a continuous 1-cocycle with

$$d^{-1}(n) = \mathcal{G}_n := (\mathcal{G}_{\frac{n}{|n|}})^{|n|},$$

and each  $\mathcal{G}_n$  is clopen. We thus satisfy the hypothesis of [81, Proposition 10], which gives the isomorphism  $\mathcal{O}_{E^{(r,R)}} \rightarrow C_r^*(\mathcal{G})$ .  $\square$

**2.4.1. The Cuntz–Pimsner Extension Class.** We extend the equivalence of the one-dimensional transversal groupoid with a Cuntz–Pimsner algebra to a compatibility of the bulk  $KK$ -cycle from Eq. (4) on page 1993 with the class in  $KK^1(\mathcal{O}_{E^{(r,R)}}, C(\Omega_0))$  that comes from the defining extension of  $\mathcal{O}_{E^{(r,R)}}$ .

**Lemma 2.24.** *The  $C^*$ -module  $E_{C(\Omega_0)}^{(r,R)}$  is a self-Morita equivalence bimodule (SMEB).*

*Proof.* Given  $\omega \in \Omega_0$  with ordering  $\mathcal{L}^{(\omega)} = \{x_n\}_{n \in \mathbb{Z}}$  with  $x_0 = 0$  and  $x_n - x_{n-1} \in (r, R)$ , a generic element in  $c^{-1}(r, R)$  can be written as  $(T_{-x_n}\omega, x_{n+1} - x_n)$ . We first compute

$$\begin{aligned} ({}_{C(\Omega_0)}(f_1 \mid f_2) \cdot f_3)(T_{-x_n}\omega, x_{n+1} - x_n) &= (f_1 * f_2^*)(T_{x_n}\omega, 0)f_3(T_{-x_n}\omega, x_{n+1} - x_n) \\ &= f_1(T_{-x_n}\omega, x_{n+1} - x_n)f_2^*(T_{x_{n+1}}\omega, x_n - x_{n+1})f_3(T_{-x_n}\omega, x_{n+1} - x_n) \\ &= f_1(T_{-x_n}\omega, x_{n+1} - x_n)\overline{f_2(T_{-x_n}\omega, x_{n+1} - x_n)}f_3(T_{-x_n}\omega, x_{n+1} - x_n) \end{aligned}$$

and then compare to

$$\begin{aligned} (f_1 \cdot (f_2 \mid f_3)_{C(\Omega_0)})(T_{-x_n}\omega, x_{n+1} - x_n) &= f_1(T_{-x_n}\omega, x_{n+1} - x_n)(f_2^* * f_3)(T_{x_{n+1}}\omega, 0) \\ &= f_1(T_{-x_n}\omega, x_{n+1} - x_n)f_2^*(T_{x_{n+1}}\omega, x_n - x_{n+1})f_3(T_{x_n}\omega, x_{n+1} - x_n) \\ &= f_1(T_{-x_n}\omega, x_{n+1} - x_n)\overline{f_2(T_{-x_n}\omega, x_{n+1} - x_n)}f_3(T_{-x_n}\omega, x_{n+1} - x_n) \end{aligned}$$

as required. Lastly the bimodule is full as by the compactness of  $\Omega_0$ , any  $g \in C(\Omega_0)$  can be written

$$\begin{aligned} g(\omega) &= f_1(\omega, x_1)\overline{f_2(\omega, x_1)} = {}_{C(\Omega)}(f_1 \mid f_2)(\omega) \\ &= \overline{\tilde{f}_1(T_{-x_1}\omega, -x_1)}\tilde{f}_2(T_{-x_1}\omega, -x_1) = (\tilde{f}_1 \mid \tilde{f}_2)_{C(\Omega_0)}(\omega) \end{aligned}$$

for some  $f_1, f_2, \tilde{f}_1, \tilde{f}_2 \in C_c(c^{-1}(r, R))$ .  $\square$

Given the bimodule  $E_{C(\Omega_0)}^{(r,R)}$ , the Cuntz–Pimsner algebra  $\mathcal{O}_{E^{(r,R)}}$  is defined by a short exact sequence

$$0 \rightarrow \mathbb{K}((F_{E^{(r,R)}})_{C(\Omega_0)}) \rightarrow \mathcal{T}_{E^{(r,R)}} \rightarrow \mathcal{O}_{E^{(r,R)}} \rightarrow 0, \tag{6}$$

where  $\mathcal{T}_{E^{(r,R)}}$  is generated by creation and annihilation operators on the Fock module  $F_{E^{(r,R)}} = \bigoplus_{n \geq 0} (E^{(r,R)})_{C(\Omega_0)}^{\otimes n}$  with  $(E^{(r,R)})_{C(\Omega_0)}^{\otimes 0} := C(\Omega_0)$ .

The extension Eq. (6) gives a  $KK^1$ -class [ext] which can be composed with the natural Morita equivalence between  $\mathbb{K}((F_{E^{(r,R)}})_{C(\Omega_0)})$  and  $C(\Omega_0)$ . Thus, the Cuntz–Pimsner algebra gives an element  $[\text{ext}] \hat{\otimes}_{\mathbb{K}} [F_E] \in KK^1(\mathcal{O}_{E^{(r,R)}}, C(\Omega_0))$ . We can use [80, Section 3.1] to construct an unbounded Kasparov module representing this class.

Using the conjugate module  $\overline{E}_{C(\Omega)}$ , we define for  $n < 0$ ,  $(E^{(r,R)})^{\otimes n} = (\overline{E}^{(r,R)})^{\otimes |n|}$ . We can then consider the bi-infinite Fock module

$$F_{E,\mathbb{Z}} := \bigoplus_{n \in \mathbb{Z}} (E^{(r,R)})_{C(\Omega_0)}^{\otimes n},$$

which carries a natural representation of  $\mathcal{O}_E$  and an operator making it into a  $KK$ -cycle.

**Proposition 2.25** ([80], Theorem 3.1). *Define an operator  $N$  on the algebraic direct sum  $\bigoplus_{m \in \mathbb{Z}}^{\text{alg}} E^{\otimes m}$  by  $N\xi = n\xi$  for  $\xi \in E^{\otimes n}$ . There is a  $*$ -homomorphism  $\mathcal{O}_{E^{(r,R)}} \rightarrow \text{End}^*((F_{E,\mathbb{Z}})_{C(\Omega_0)})$  such that  $S_f \cdot \xi := f \otimes \xi$  for all  $f \in E^{(r,R)}$  and  $\xi \in (E^{(r,R)})^{\otimes n}$ . The triple  $(\mathcal{O}_{E^{(r,R)}}, (F_{E,\mathbb{Z}})_{C(\Omega_0)}, N)$  is an unbounded odd Kasparov module that represents the class  $[\text{ext}] \hat{\otimes}_{\mathbb{K}} [F_E] \in KK^1(\mathcal{O}_{E^{(r,R)}}, C(\Omega_0))$ .*

**Corollary 2.26.** *The odd Kasparov module from Proposition 2.25 defines the same class in  $KK^1(C_r^*(\mathcal{G}), C(\Omega_0))$  as the bulk Kasparov module  ${}_d\lambda_{\Omega_0}$  from Eq. (4) on page 1993 with  $d = 1$ .*

*Proof.* The  $C^*$ -algebras are isomorphic by Proposition 2.23. Furthermore, the positive semisplitting from both the groupoid and Cuntz–Pimsner Kasparov modules is the projection onto elements with nonnegative cocycle values. Hence, the extensions are equivalent, which also gives equivalence within  $KK^1$ .  $\square$

By the presentation of  $C_r^*(\mathcal{G})$  as a Cuntz–Pimsner algebra, we can use the long (cyclic) exact sequence as a tool for the computation of  $K_*(C_r^*(\mathcal{G}))$ . Namely, for complex algebras,

$$\begin{array}{ccccc} K_0(C(\Omega_0)) & \xrightarrow{\otimes([C(\Omega_0)] - [E^{(r,R)}])} & K_0(C(\Omega_0)) & \xrightarrow{\iota_*} & K_0(C_r^*(\mathcal{G})) \\ \uparrow \partial & & & & \downarrow \partial \\ K_1(C_r^*(\mathcal{G})) & \xleftarrow{\iota_*} & K_1(C(\Omega_0)) & \xleftarrow{\otimes([C(\Omega_0)] - [E^{(r,R)}])} & K_1(C(\Omega_0)) \end{array}$$

where the map  $K_*(C(\Omega_0)) \xrightarrow{\otimes[E^{(r,R)}]} K_*(C(\Omega_0))$  comes from the internal product of the  $K$ -theory class with the element  $[E^{(r,R)}] \in KK(C(\Omega_0), C(\Omega_0))$ . There is an analogous but longer exact sequence for real  $C^*$ -algebras,

$$\begin{aligned} \dots &\rightarrow KO_j(C(\Omega_0)) \xrightarrow{\otimes([C(\Omega_0)] - [E^{(r,R)}])} KO_j(C(\Omega_0)) \\ &\xrightarrow{\iota_*} KO_j(C_r^*(\mathcal{G})) \xrightarrow{\partial} KO_{j-1}(C(\Omega_0)) \rightarrow \dots \end{aligned}$$

By the Morita equivalence of  $C_r^*(\mathcal{G})$  and  $C(\Omega_{\mathcal{L}}) \rtimes \mathbb{R}$ , we know that  $K_*(C_r^*(\mathcal{G})) \cong K_{*-1}(C(\Omega_{\mathcal{L}}))$  by the Connes–Thom isomorphism. As the  $K$ -theory of  $C(\Omega_{\mathcal{L}})$  is generally quite difficult to compute, the Pimsner exact sequence for  $C_r^*(\mathcal{G})$  is a helpful tool for such  $K$ -theory computations. For example, if  $K_1(C(\Omega_0)) = 0$  (e.g. if  $\mathcal{L}$  has finite local complexity), then we immediately obtain that

$$K_0(C_r^*(\mathcal{G})) \cong \text{Coker}(1 - [E^{(r,R)}]), \quad K_1(C_r^*(\mathcal{G})) \cong \text{Ker}(1 - [E^{(r,R)}]).$$

Hence, for a one-dimensional lattice  $\mathcal{L}$  with finite local complexity,

$$K_0(C(\Omega_{\mathcal{L}})) \cong \text{Ker}(1 - [E^{(r,R)}]), \quad K_1(C(\Omega_{\mathcal{L}})) \cong \text{Coker}(1 - [E^{(r,R)}]).$$

Of course, this result is restricted to one-dimensional lattices or tilings. A description of  $C_r^*(\mathcal{G})$  for higher dimensions using the  $C^*$ -algebra of a product system or as an iterated Cuntz–Pimsner algebra may be possible. We leave this analysis to future research.

*Remark 2.27.* As a brief cautionary remark, we note that our bimodule  $C(\Omega_0)E_{C(\Omega_0)}^{(r,R)}$  looks quite similar but is different from the crossed product bimodule  ${}_{\alpha}A_A$  with  $\alpha : \mathbb{Z} \rightarrow \text{Aut}(A)$  and such that  $\mathcal{O}_{\alpha} \cong A \rtimes_{\alpha} \mathbb{Z}$ . Indeed, given  $\omega \in \Omega_0$  and  $x_1 \in \mathcal{L}^{(\omega)} \cap (r, R)$ , there is no guarantee that  $T_{-2x_1}\omega \in \Omega_0$  as would be the case for a  $\mathbb{Z}$ -action.

### 3. Factorisation and the Bulk–Boundary Correspondence

A key attribute of the operator algebra approach to topological phases via crossed products is that both bulk and boundary systems can be treated under the same general framework with an extension of  $C^*$ -algebras linking the two systems. Namely, up to stabilisation the edge algebra can be described via  $C(\Omega) \rtimes_{\sigma} \mathbb{Z}^{d-1}$  and, we can recover the bulk algebra by the iterated crossed product  $(C(\Omega) \rtimes_{\sigma} \mathbb{Z}^{d-1}) \rtimes \mathbb{Z} \cong C(\Omega) \rtimes_{\sigma} \mathbb{Z}^d$  for normalised twists.

In this section we use the groupoid cocycle  $c_d : \mathcal{G} \rightarrow \mathbb{R}$  to consider the closed subgroupoid  $\Upsilon = \text{Ker}(c_d)$ . This subgroupoid is too small to completely model an edge system but is groupoid equivalent to one that we argue encodes the translation dynamics on the transversal in  $(d-1)$ -directions. Furthermore, we show that the subgroupoid  $\Upsilon$  gives rise to a canonical *bulk–boundary extension* of reduced  $C^*$ -algebras that generalises the Toeplitz extension for crossed products. In particular, we use this extension to factorise the groupoid  $KK$ -cycle into a product of a  $(d-1)$ -dimensional system and the bulk–boundary extension that recovers the bulk system.

#### 3.1. The Edge Groupoid

We now apply our results on twisted groupoid equivalences to the transversal groupoid and the bulk–boundary short exact sequence.

Recall the groupoid cocycle  $c_d : \mathcal{G} \rightarrow \mathbb{R}$ ,  $c_d(\omega, x) = x_d$ . Because  $c_d$  is exact, we can apply the results from Sect. 1.5 and construct an unbounded  $KK$ -cycle. We consider the closed subgroupoid  $\Upsilon = \text{Ker}(c_d)$ , namely

$$\Upsilon = \{(\omega, y) \in \Omega_0 \times \mathbb{R}^{d-1} : T_{(-y,0)}\omega \in \Omega_0\}.$$

with multiplication, range and source maps inherited from  $\mathcal{G}$ . Furthermore, the restriction of  $\sigma$  to  $\Upsilon$  gives a well-defined 2-cocycle for  $\Upsilon$ . Recalling Sect. 1.4, the restriction map

$$\rho_\Upsilon : C_c(\mathcal{G}, \sigma) \rightarrow C_c(\Upsilon, \sigma)$$

defines a  $(C_r^*(\mathcal{G}, \sigma), C_r^*(\Upsilon, \sigma))$ -bimodule  $E_{C_r^*(\Upsilon, \sigma)}$ . Applying Proposition 1.13 to the cocycle  $c_d : \mathcal{G} \rightarrow \mathbb{R}$  and writing  $X_d := D_{c_d}$  gives us the following.

**Proposition 3.1** ([65], Theorem 3.9). *The triple*

$$d\lambda_{d-1} = \left( C_c(\mathcal{G}, \sigma) \hat{\otimes} Cl_{0,1}, E_{C_r^*(\Upsilon, \sigma)} \hat{\otimes} \bigwedge^* \mathbb{R}, X_d \hat{\otimes} \gamma \right)$$

is a real or complex unbounded Kasparov module.

The groupoid  $\Upsilon$  is too small to be thought of as representing an edge system. Instead, we will consider the groupoid  $\mathcal{G} \times \mathcal{G}/\Upsilon$  whose twisted reduced  $C^*$ -algebra is Morita equivalent to  $C_r^*(\Upsilon, \sigma)$ , cf. Sect. 1.4.

The cocycle  $c_d$  determines the subset  $\text{Ran}(c_d) \subset \mathbb{R}^d$  (which need not be a subgroup). Having fixed this set, the groupoid  $\mathcal{G} \times \mathcal{G}/\Upsilon$  allows us to put a groupoid structure back into our system with the translation action in  $(d-1)$ -directions.

The space  $\mathcal{G}/\Upsilon$  is given by equivalence classes of elements  $[(\omega, x)] \in \mathcal{G}$  under the relation

$$(\omega, x) \sim (\omega', x') \Leftrightarrow \exists (T_{-x}\omega, (y, 0)) \in \Upsilon \quad (\omega, x + (y, 0)) = (\omega', x').$$

Hence, the quotient  $\mathcal{G}/\Upsilon$  can be described by equivalence classes of pairs  $[(\omega, x_d)]$  with  $(\omega, x_d) \in \Omega_0 \times \text{Ran}(c_d)$ . We have the presentation of  $\mathcal{G} \times \mathcal{G}/\Upsilon$  by pairs

$$\begin{aligned} \mathcal{G} \times \mathcal{G}/\Upsilon &\cong \{((\omega, x), [(\omega', y_d)]) : r_{\mathcal{G}}(c_d^{-1}(y_d)) = T_{-x}\omega\} \\ &\cong \{((\omega, x), [(T_{-x}\omega, y_d)])\} \subset \mathcal{G} \times \mathcal{G}/\Upsilon. \end{aligned}$$

Recall that  $(\omega, x) \in \mathcal{G}$  if  $x \in \mathcal{L}^{(\omega)}$ . Our presentation says that  $((\omega, x), [(T_{-x}\omega, y_d)]) \in \mathcal{G} \times \mathcal{G}/\Upsilon$  if there is some  $u \in \mathbb{R}^{d-1}$  such that  $x, x + (u, y_d) \in \mathcal{L}^{(\omega)}$ . The unit space is given by

$$(\mathcal{G} \times \mathcal{G}/\Upsilon)^{(0)} = \mathcal{G}/\Upsilon,$$

and the groupoid structure is determined by

$$\begin{aligned} s((\omega, x), [(T_{-x}\omega, y_d)]) &= [(T_{-x}\omega, y_d)], \quad r((\omega, x), [(T_{-x}\omega, y_d)]) = [(\omega, x_d + y_d)], \\ ((\omega, x), [(T_{-x}\omega, y_d)])^{-1} &= ((T_{-x}\omega, -x), [(\omega, x_d + y_d)]), \\ ((\omega, x), [(T_{-x}\omega, y_d)]) \cdot ((T_{-x}\omega, z), [(T_{-x-z}\omega, y_d - z_d)]) & \\ &= ((\omega, x + z), [(T_{-x-z}\omega, y_d - z_d)]). \end{aligned}$$

We note that for  $((T_{-x}\omega, z), [(T_{-x-z}\omega, y_d - z_d)])$  to be in  $\mathcal{G} \times \mathcal{G}/\Upsilon$ , there must be some  $v \in \mathbb{R}^{d-1}$  such that  $x + (v, y_d) \in \mathcal{L}^{(\omega)}$ . Because  $((\omega, x), [(T_{-x}\omega, y_d)]) \in \mathcal{G} \times \mathcal{G}/\Upsilon$  implies  $x + (u, y_d) \in \mathcal{L}^{(\omega)}$  for some  $u \in \mathbb{R}^{d-1}$ , the groupoid multiplication involves a translation in  $(d-1)$ -dimensions only. Thus, we see that the groupoid

$\mathcal{G} \ltimes \mathcal{G}/\Upsilon$  models the dynamics of the transversal  $\Omega_0$  relative to the fixed set  $\text{Ran}(c_d)$ . We use the 2-cocycle  $\sigma$  on  $\mathcal{G}$  to define a 2-cocycle on  $\mathcal{G} \ltimes \mathcal{G}/\Upsilon$  via

$$\sigma(((\omega, x), [(T_{-x}\omega, y_d)]), ((T_{-x}\omega, z), [(T_{-x-z}\omega, y_d - z_d)])) = \sigma((\omega, x), (T_{-x}\omega, z)).$$

Applying Proposition 1.12, we obtain the following.

**Proposition 3.2.** *The  $C^*$ -module  $E_{C_r^*(\Upsilon, \sigma)}$  is a Morita equivalence bimodule between  $C_r^*(\mathcal{G} \ltimes \mathcal{G}/\Upsilon, \sigma)$  and  $C_r^*(\Upsilon, \sigma)$ . In particular, there is an isomorphism  $C_r^*(\mathcal{G} \ltimes \mathcal{G}/\Upsilon, \sigma) \cong \mathbb{K}(E_{C_r^*(\Upsilon, \sigma)})$ .*

The Morita equivalence bimodule gives an invertible element in  $KK(C_r^*(\mathcal{G} \ltimes \mathcal{G}/\Upsilon, \sigma), C_r^*(\Upsilon, \sigma))$ . From the perspective of index theory, we can work with either  $\Upsilon$  or  $\mathcal{G} \ltimes \mathcal{G}/\Upsilon$ . While we consider  $\mathcal{G} \ltimes \mathcal{G}/\Upsilon$  to be our edge groupoid, the subgroupoid  $\Upsilon \subset \mathcal{G}$  will be easier to work with for some of our mathematical arguments.

### 3.2. The Bulk–Boundary Extension

Because the Kasparov module from Proposition 3.1 comes from an exact  $\mathbb{R}$ -valued cocycle, we can construct an extension of  $C^*$ -algebras. As in Lemma 1.15 in Sect. 1.6, we fix an  $\varepsilon > 0$  and a function  $\chi_+$  to define a self-adjoint operator  $\Pi_d := \Pi_{c_d} \in \text{End}_{C_r^*(\Upsilon, \sigma)}^*(E)$  on the  $C^*$ -module  $E_{C_r^*(\Upsilon, \sigma)}$  satisfying  $\Pi_d^2 - \Pi_d \in \mathbb{K}_{C_r^*(\Upsilon, \sigma)}(E)$ . Since the Delone set  $\mathcal{L}$  is relatively dense, the cocycle  $c_d$  takes arbitrarily large values in each  $r$ -fibre and is  $r$ -unbounded. Therefore, the map

$$\varphi : C_r^*(\mathcal{G}, \sigma) \rightarrow \mathcal{Q}(E_{C_r^*(\Upsilon, \sigma)}), \quad \varphi(a) = q(\Pi_d a \Pi_d),$$

is injective by Proposition 1.18. Hence, we can construct the generalised Toeplitz extension

$$0 \rightarrow C_r^*(\mathcal{G} \ltimes \mathcal{G}/\Upsilon, \sigma) \rightarrow C^*(\Pi_d C_r^*(\mathcal{G}, \sigma) \Pi_d, C_r^*(\mathcal{G} \ltimes \mathcal{G}/\Upsilon, \sigma)) \rightarrow C_r^*(\mathcal{G}, \sigma) \rightarrow 0 \tag{7}$$

with completely positive semisplitting  $a \mapsto \Pi_d a \Pi_d$  and Busby invariant  $\varphi$ .

In the case that our lattices have a canonical  $\mathbb{Z}^d$ -labelling, then this extension reduces to the usual bulk–boundary short exact sequence considered in [77]. In the case  $d = 1$ , we have  $\Upsilon \cong \mathcal{G}^{(0)} \cong \Omega_0$ , and the extension (7) is equivalent to the Toeplitz–Cuntz–Pimsner extension for  $C_r^*(\mathcal{G}, \sigma)$  of Corollary 2.26.

For a fixed  $\omega \in \Omega_0$ , the Toeplitz algebra can be represented on the space  $\Pi_d^\omega \ell^2(\mathcal{L}^{(\omega)})$ , which we can interpret as a half-infinite system with boundary. Because we work with general Delone sets, 0 need not be an isolated point in  $\text{Ran}(c_d)$ . As such, our boundary operator  $\Pi_d$  is not a projection in general, but if we have a  $\mathbb{Z}^d$ -labelling, the above construction yields a genuine projection.

*Remark 3.3* (Integer-valued cocycles and the Pimsner–Voiculescu extension) It is shown in [65, Proposition 3.22] that if the an exact cocycle  $c_d$  is integer-valued, then the associated  $KK$ -class  $[D_{c_d}] \in KK^1(C_r^*(\mathcal{G}, \sigma), C_r^*(\Upsilon, \sigma))$  coincides with the  $KK$ -class defined from the circle action

$$\alpha^c : \mathbb{T} \rightarrow \text{Aut}(C_r^*(\mathcal{G}, \sigma)), \quad \alpha_t^c(f)(\xi) := e^{itc_d(\xi)} f(\xi),$$

via the construction in [24]. For crossed products by  $\mathbb{Z}$ , the Kasparov module of a circle action is the same as the Kasparov module constructed from the Toeplitz extension of the crossed product (constructed in, for example, [20])

$$0 \rightarrow C_r^*(\Upsilon, \sigma) \otimes \mathbb{K} \rightarrow \mathcal{T} \rightarrow C_r^*(\Upsilon, \sigma) \rtimes \mathbb{Z} \rightarrow 0$$

with  $C_r^*(\Upsilon, \sigma) \rtimes \mathbb{Z} \cong C_r^*(\mathcal{G}, \sigma)$ . A similar result holds for semisaturated circle actions (see [3, Section 3.3] or [4, Section 3]). Hence, we recover the ‘usual’ bulk–boundary extension considered in [77] for special cases of integer-valued cocycles  $c_d$ . This applies in particular if  $c_d$  is unperforated.

*Remark 3.4* (The Connes–Thom class). Let us now consider the relation between the Kasparov module of Proposition 3.1 and its Toeplitz extension with the Connes–Thom isomorphism and the Wiener–Hopf extension of [51], when  $d \geq 2$ .

The transversal groupoid  $\mathcal{G}$  is Morita equivalent to the crossed product groupoid  $\Omega_{\mathcal{L}} \rtimes \mathbb{R}^d$ , and for  $d \geq 2$  the boundary groupoid  $\Upsilon$  is equivalent to  $\Omega_{\mathcal{L}} \rtimes \mathbb{R}^{d-1}$ . Given a normalised 2-cocycle  $\sigma : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathcal{U}(C(\Omega_{\mathcal{L}}))$ , there is an isomorphism  $C(\Omega_{\mathcal{L}}) \rtimes_{\sigma} \mathbb{R}^d \cong (C(\Omega_{\mathcal{L}}) \rtimes_{\sigma} \mathbb{R}^{d-1}) \rtimes \mathbb{R}$  and a Wiener–Hopf extension

$$0 \rightarrow (C(\Omega_{\mathcal{L}}) \rtimes_{\sigma} \mathbb{R}^{d-1}) \otimes \mathbb{K}[L^2(\mathbb{R})] \rightarrow \mathcal{W} \rightarrow C(\Omega_{\mathcal{L}}) \rtimes_{\sigma} \mathbb{R}^d \rightarrow 0,$$

see [51]. In [22, Section 6], it was shown that the Wiener–Hopf extension can be represented by the unbounded Kasparov module

$$\left( C_c(\mathbb{R}, C(\Omega_{\mathcal{L}}) \rtimes_{\sigma} \mathbb{R}^{d-1}) \hat{\otimes} Cl_{0,1}, F_{C(\Omega_{\mathcal{L}}) \rtimes_{\sigma} \mathbb{R}^{d-1}} \hat{\otimes} \bigwedge^* \mathbb{R}, X \hat{\otimes} \gamma \right), \quad (8)$$

where  $F_{C(\Omega_{\mathcal{L}}) \rtimes_{\sigma} \mathbb{R}^{d-1}}$  is the bimodule obtained from the conditional expectation induced from the restriction to the closed subgroupoid  $\Omega_{\mathcal{L}} \rtimes \mathbb{R}^{d-1} \subset \Omega_{\mathcal{L}} \rtimes \mathbb{R}^d$ ,

$$\begin{aligned} \rho : C_c(\Omega_{\mathcal{L}} \rtimes_{\sigma} \mathbb{R}^d) &\rightarrow C_c(\Omega_{\mathcal{L}} \rtimes_{\sigma} \mathbb{R}^{d-1}), \\ \rho(f)(x_1, \dots, x_{d-1}) &= f(x_1, \dots, x_{d-1}, 0). \end{aligned}$$

We consider the composition of  $KK$ -classes

$$\begin{aligned} &\left( C(\Omega_{\mathcal{L}}) \rtimes_{\sigma} \mathbb{R}^d, F_{C_r^*(\mathcal{G}, \sigma)}^d, 0 \right) \hat{\otimes}_{C_r^*(\mathcal{G}, \sigma)} [d\lambda_{d-1}] \\ &\hat{\otimes}_{C_r^*(\Upsilon, \sigma)} \left( C_r^*(\Upsilon, \sigma), (F^{d-1})_{C(\Omega_{\mathcal{L}}) \rtimes_{\sigma} \mathbb{R}^{d-1}}^* \right), \end{aligned}$$

where the left and right Kasparov modules represent the Morita equivalence (resp. dual Morita equivalence) of the groupoid algebras and crossed products. The end result of this triple product is a Kasparov module representing a class in  $KK^1(C(\Omega_{\mathcal{L}}) \rtimes_{\sigma} \mathbb{R}^d, C(\Omega_{\mathcal{L}}) \rtimes_{\sigma} \mathbb{R}^{d-1})$ . Its relation to the Kasparov module in Eq. (8) is as follows. By Definition 2.11 and Proposition 2.16, the Morita equivalence bimodule  $F_{C_r^*(\mathcal{G}, \sigma)}^d$  is obtained from restriction of the crossed product dynamics to the transversal  $\Omega_0$ . The  $C^*$ -module  $E_{C_r^*(\Upsilon, \sigma)}$  from  $d\lambda_{d-1}$  in Proposition 3.1 is defined by a restriction  $C_c(\mathcal{G}, \sigma) \rightarrow C_c(\Upsilon, \sigma)$ . Lastly the dual Morita equivalence bimodule  $(F^{d-1})_{C(\Omega_{\mathcal{L}}) \rtimes_{\sigma} \mathbb{R}^{d-1}}^*$  is induced by the inclusion



of  $\Upsilon$  into  $\Omega_{\mathcal{L}} \rtimes \mathbb{R}^{d-1}$ . Hence, the inner product on the balanced tensor product can be considered as coming from a generalised conditional expectation  $C_c(\Omega_{\mathcal{L}} \rtimes \mathbb{R}^d, \sigma) \rightarrow C_c(\Omega_{\mathcal{L}} \rtimes \mathbb{R}^{d-1}, \sigma)$  and there is a natural identification

$$F^d \otimes_{C_r^*(\mathcal{G}, \sigma)} E \otimes_{C_r^*(\Upsilon, \sigma)} (F^{d-1})^*_{C(\Omega_{\mathcal{L}}) \rtimes_{\sigma} \mathbb{R}^{d-1}} \xrightarrow{\sim} F_{C(\Omega_{\mathcal{L}}) \rtimes_{\sigma} \mathbb{R}^{d-1}}.$$

An argument similar to that in the proof of Theorem 3.6 shows that the operator  $X$  in  $F_{C(\Omega_{\mathcal{L}}) \rtimes_{\sigma} \mathbb{R}^{d-1}}$  satisfies the connection condition with respect to the operator  $X \otimes 1$  in  $E \otimes_{C_r^*(\Upsilon, \sigma)} (F^{d-1})^*_{C(\Omega_{\mathcal{L}}) \rtimes_{\sigma} \mathbb{R}^{d-1}}$ . As these maps are also compatible with the Clifford actions, we recover the unbounded representative of the Wiener–Hopf extension from Eq. (8) on page 2000. The boundary maps in  $K$ -theory and  $K$ -homology from the Wiener–Hopf extension, i.e. the product with the unbounded Kasparov module from Eq. (8), implement the inverse of the Connes–Thom isomorphism [82]. Hence, these maps are represented by our Toeplitz extension up to groupoid/Morita equivalence.

### 3.3. Factorisation

By the same basic argument as the bulk algebra, we can build a  $KK$ -cycle for  $C_r^*(\Upsilon, \sigma)$  which is stably isomorphic to the edge algebra  $C_r^*(\mathcal{G} \rtimes \Upsilon, \sigma)$ . We denote by  $F_{C(\Omega_0)}$  the  $(C_r^*(\Upsilon, \sigma), C(\Omega_0))$ - $C^*$ -bimodule coming from the restriction of  $C_r^*(\Upsilon, \sigma)$  to the unit space. The notation  $F_{C(\Omega_0)}$  distinguishes it from the  $C^*$ -module  $E_{C(\Omega_0)}$  constructed from  $C_r^*(\mathcal{G}, \sigma)$ . Specifically,

$${}_{d-1}\lambda_{\Omega_0} = \left( C_c(\Upsilon, \sigma) \hat{\otimes} Cl_{0,d-1}, F_{C(\Omega_0)} \hat{\otimes} \bigwedge^* \mathbb{R}^{d-1}, \sum_{j=1}^{d-1} X_j \hat{\otimes} \gamma^j \right)$$

is an unbounded Kasparov module and gives rise to a class in  $KK^{d-1}(C_r^*(\Upsilon, \sigma), C(\Omega_0))$  (real or complex). We have constructed a class representing our bulk–boundary extension and a  $KK$ -cycle for the edge algebra. The key  $K$ -theoretic result that drives the bulk–boundary correspondence is that these two  $KK$ -cycles can be put together using the unbounded Kasparov product to reconstruct the bulk  $KK$ -cycle.

**Theorem 3.5.** *Under the boundary map coming from the extension of Eq. (7) on page 1999,*

$$\partial[{}_{d-1}\lambda_{\Omega_0}] = [{}_{d-1}\lambda_{\Omega_0}] \hat{\otimes}_{C_r^*(\Upsilon, \sigma)} [{}_{d-1}\lambda_{\Omega_0}] = (-1)^{d-1} [{}_{d-1}\lambda_{\Omega_0}],$$

with  ${}_{d-1}\lambda_{\Omega_0}$  the bulk  $KK$ -cycle from Eq. (4) on page 1993 and where  $-[x]$  denotes the inverse in the  $KK$ -group. Furthermore, the equality is an unbounded equivalence up to a permutation of the Clifford algebra basis.

Theorem 3.5 is a special case of Theorem 3.6 with  $k = d - 1$ . Hence, we delay the proof until Sect. 3.4.

**3.3.1. A Remark on More General Boundaries.** Our edge groupoid  $\Upsilon$  can be thought of as the result of a cut of the Delone sets  $\mathcal{L}^{(\omega)} \in \Omega_0$  along the plane defined by  $\text{Ker}(c_d) \cong \mathbb{R}^{d-1} \times \{0\}$ . This choice of cut or boundary is somewhat arbitrary. Let us briefly consider more general boundary choices though, as we will show, our  $KK$ -theoretic factorisation still applies.

Let  $b : \mathbb{R}^d \rightarrow \mathbb{R}$  be a continuous homomorphism such that as a vector space  $\dim(\text{Ran}(b)) = 1$ . The plane defined by  $\text{Ker}(b)$  defines a new  $(d - 1)$ -dimensional plane in  $\mathbb{R}^d$  which we can cut along to make a new boundary. It is easy to check that the corresponding map  $c_b : \mathcal{G} \rightarrow \mathbb{R}$ ,  $c_b(\omega, x) = b(x)$  is an exact groupoid cocycle. Hence, we can study this boundary via the closed subgroupoid  $\Upsilon_b = \text{Ker}(c_b)$  and equivalent groupoid  $\mathcal{G} \times \mathcal{G}/\Upsilon_b$ , which models the  $(d - 1)$ -dimensional dynamics of the transversal relative to  $\text{Ran}(b)$ . Because  $c_b$  is exact, we can construct an ungraded and unbounded Kasparov module  $(C_c(\mathcal{G}, \sigma), E_{C_r^*(\Upsilon_b, \sigma)}, D_b)$  that gives a class  $[\text{ext}_b] \in KK^1(C_r^*(\mathcal{G}, \sigma), C_r^*(\Upsilon_b, \sigma))$  and a bulk–boundary short exact sequence

$$0 \rightarrow C_r^*(\mathcal{G} \times \mathcal{G}/\Upsilon_b, \sigma) \rightarrow \mathcal{T}_b \rightarrow C_r^*(\mathcal{G}, \sigma) \rightarrow 0.$$

As a vector space,  $\text{Ker}(b)$  is  $(d - 1)$ -dimensional and so fix an orthonormal basis  $\{z_1, \dots, z_{d-1}\}$ . These basis vectors give rise to an exact  $\mathbb{R}^{d-1}$ -valued cocycle on the groupoid  $\Upsilon_b$ , which we use to build an unbounded Kasparov module and a class  $[\lambda_{\Omega_0}^b] \in KK^{d-1}(C_r^*(\Upsilon_b), C(\Omega_0))$ .

Following the proof of Theorem 3.6 with  $k = d - 1$ , the product of the class of the Kasparov modules  $[\text{ext}_b]$  and  $[\lambda_{\Omega_0}^b]$  is represented by the unbounded Kasparov module

$$\left( C_c(\mathcal{G}, \sigma) \hat{\otimes} Cl_{0,d}, E_{C(\Omega_0)} \hat{\otimes} \bigwedge^* \mathbb{R}^d, \sum_{j=1}^{d-1} Z_j \hat{\otimes} \gamma^{j+1} + D_b \hat{\otimes} \gamma^1 \right).$$

At this point, we can take transformation from the basis  $\{z_1, \dots, z_{d-1}, z_d\}$  to the standard basis of  $\mathbb{R}^d$ . This transformation recovers the bulk  $K$ -cycle  $d\lambda_{\Omega_0}$  up to a Clifford basis reordering. Fixing the Clifford basis reordering, we have that

$$[\text{ext}_b] \hat{\otimes}_{C_r^*(\Upsilon_b, \sigma)} [\lambda_{\Omega_0}^b] = (-1)^{d-1} [d\lambda_{\Omega_0}]$$

and our factorisation result extends.

Let us briefly note that while any crystallographic group  $G \subset \mathbb{R}^d$  is a Delone set and our choice of boundary is quite general, the factorisation and bulk–boundary result in Theorem 3.5 is too coarse to detect boundary indices derived from the crystalline structure as in [38].

### 3.4. $KK$ -Cycles with Higher Codimension

Let us now generalise the constructions and ideas from the previous section to consider subinvariants of arbitrary codimension. Such invariants are linked to so-called weak topological phases which are characterised by elements in  $K_*(C_r^*(\mathcal{G}, \sigma))$  that are not detected by the ‘top degree form’ that comes via a pairing with  $d\lambda_{\Omega_0}$ .

Once again we use a groupoid homomorphism  $\check{c}_k : \mathcal{G} \rightarrow \mathbb{R}^{d-k}$  via  $\check{c}_k(\omega, x) = (x_{k+1}, \dots, x_d)$  and define  $\Upsilon_k = \text{Ker}(\check{c}_k)$ , where we characterise

$$\Upsilon_k = \left\{ (\omega, x_1, \dots, x_k) \in \Omega_0 \times \mathbb{R}^k : (x_1, \dots, x_k, 0, \dots, 0) \in \mathcal{L}^{(\omega)} \right\}.$$

As in the case of  $k = d - 1$ ,  $\Upsilon_k$  is a closed subgroupoid of  $\mathcal{G}$  and is equivalent to  $\mathcal{G} \times \mathcal{G}/\Upsilon_k$ . By Proposition 1.13, we can build a Kasparov module

$${}_d\lambda_k = \left( C_c(\mathcal{G}, \sigma) \hat{\otimes} Cl_{0,d-k}, E_{C_r^*(\Upsilon_k, \sigma)}^{d-k} \hat{\otimes} \bigwedge^* \mathbb{R}^{d-k}, \sum_{j=k+1}^d X_j \hat{\otimes} \gamma^{j-k} \right).$$

In the case  $k = d - 1$ ,  ${}_d\lambda_{d-1}$  is the unbounded  $KK$ -cycle representing the bulk-boundary extension considered in Sect. 3.1. We will be interested in pairings of  ${}_d\lambda_k$  with the  $K$ -theory of  $C_r^*(\mathcal{G}, \sigma)$ . Such pairings naturally take values in the  $K$ -theory of  $C_r^*(\Upsilon_k, \sigma)$ .

As in the case of  $k = d - 1$ , we can construct an unbounded  $KK$ -cycle for the subgroupoid  $\Upsilon_k$ ,

$${}_k\lambda_{\Omega_0} = \left( C_c(\Upsilon_k, \sigma) \hat{\otimes} Cl_{0,k}, F_{C(\Omega_0)}^k \hat{\otimes} \bigwedge^* \mathbb{R}^k, \sum_{j=1}^k X_j \hat{\otimes} \gamma^j \right),$$

$$F_{C(\Omega_0)}^k := \overline{C_c(\Upsilon_k, \sigma)}_{C(\Omega_0)}, \tag{9}$$

which represents the class  $[{}_k\lambda_{\Omega_0}] \in KK(C_r^*(\Upsilon_k, \sigma) \hat{\otimes} Cl_{0,k}, C(\Omega_0))$ .

We now present our main factorisation result, which allows us to decompose the bulk Kasparov module  ${}_d\lambda_{\Omega_0}$  as the product of  ${}_d\lambda_k$  with  ${}_k\lambda_{\Omega_0}$  (up to a sign related to the orientation of Clifford algebras).

**Theorem 3.6.** *Taking the Kasparov product,*

$$[{}_d\lambda_k] \hat{\otimes}_{C_r^*(\Upsilon_k, \sigma)} [{}_k\lambda_{\Omega_0}] = (-1)^{k(d-k)} [{}_d\lambda_{\Omega_0}].$$

*Furthermore, our equivalence is at the unbounded level up to a permutation of the Clifford basis.*

*Proof.* Much of this proof is bookkeeping and is very similar to the proof in [20, Theorem 3.4]. To take the product of the  $C_r^*(\mathcal{G}, \sigma) \hat{\otimes} Cl_{0,d-k} - C_r^*(\Upsilon_k, \sigma)$  Kasparov module with a  $C_r^*(\Upsilon_k, \sigma) \hat{\otimes} Cl_{0,k} - C(\Omega_0)$  Kasparov module, we first take the external product of  ${}_d\lambda_k$  with a  $KK$ -cycle representing the identity in  $KK(Cl_{0,k}, Cl_{0,k})$ . This identity class can be represented by  $(Cl_{0,k}, Cl_{0,k} Cl_{0,k}, 0)$  with right and left actions by multiplication. We then take the product of a  $(C_r^*(\mathcal{G}, \sigma) \hat{\otimes} Cl_{0,d}, C_r^*(\Upsilon_k, \sigma) \hat{\otimes} Cl_{0,k})$  Kasparov module with a  $(C_r^*(\Upsilon_k, \sigma) \hat{\otimes} Cl_{0,k}, C(\Omega_0))$  Kasparov module. First, the balanced tensor product gives the  $C^*$ -module

$$\begin{aligned} & \left( E^{d-k} \hat{\otimes} \bigwedge^* \mathbb{R}^{d-k} \hat{\otimes} Cl_{0,k} \right) \hat{\otimes}_{C_r^*(\Upsilon_k, \sigma) \hat{\otimes} Cl_{0,k}} \left( F^k \hat{\otimes} \bigwedge^* \mathbb{R}^k \right)_{C(\Omega_0)} \\ & \cong \left( E^{d-k} \otimes_{C_r^*(\Upsilon_k, \sigma)} F_{C(\Omega_0)}^k \right) \hat{\otimes} \bigwedge^* \mathbb{R}^{d-k} \hat{\otimes} \left( Cl_{0,k} \hat{\otimes}_{Cl_{0,k}} \bigwedge^* \mathbb{R}^k \right) \\ & \cong \left( E^{d-k} \otimes_{C_r^*(\Upsilon_k, \sigma)} F^k \right)_{C(\Omega_0)} \hat{\otimes} \bigwedge^* \mathbb{R}^{d-k} \hat{\otimes} \bigwedge^* \mathbb{R}^k \end{aligned}$$

as  $Cl_{0,d-1}$  acts on  $\bigwedge^* \mathbb{R}^{d-1}$  non-degenerately. Next we define a unitary isomorphism

$$E^{d-k} \otimes_{C_r^*(\Upsilon_k, \sigma)} F_{C(\Omega_0)}^k \rightarrow E_{C(\Omega_0)},$$

by first considering defining a map on dense submodules,

$$v : C_c(\mathcal{G}, \sigma) \otimes_{C_c(\Upsilon_k, \sigma)} C_c(\Upsilon_k, \sigma)_{C(\Omega_0)} \ni f \otimes h \mapsto f \cdot h \in C_c(\mathcal{G}, \sigma)_{C(\Omega_0)}.$$

This map preserves the inner product structures, is thus uniformly bounded and, hence, extends to an isomorphism of  $C^*$ -modules. Furthermore, the map commutes with the representation of  $C_r^*(\mathcal{G}, \sigma)$  as elements in  $\text{End}^*(E_{C_r^*(\Upsilon_k, \sigma)}^{d-k})$  commute with the right action of  $C^*(\Upsilon_k, \sigma)$ . Similarly,  $\{X_l\}_{l=k+1}^d$  also commutes with this map as  $X_l$  is right  $C_r^*(\Upsilon_k, \sigma)$ -linear on  $E_{C_r^*(\Upsilon_k, \sigma)}^{d-k}$ . The operators  $\{X_j\}_{j=1}^k$  satisfy the connection condition under the unitary isomorphism  $v$ . Let  $f \in C_c(\mathcal{G}, \sigma)$  and consider the map

$$v \circ |f\rangle : C_c(\Upsilon_k, \sigma) \rightarrow C_c(\mathcal{G}, \sigma), \quad h \mapsto f \cdot h.$$

Then we need to check that  $X_j \circ v \circ |f\rangle - v \circ |f\rangle \circ X_j$  defines a bounded operator  $F_{C(\Omega_0)}^k \rightarrow E_{C(\Omega_0)}$ . This follows since each  $X_j$  acts as a derivation of  $C_c(\mathcal{G}, \sigma)$ :

$$\begin{aligned} & ((X_j f) \cdot h + f \cdot (X_j h))(\omega, x) \\ &= \sum_{(y, 0_{d-k}) \in \mathcal{L}^{(\omega)} - x} \left( (x_j + y_j) f(\omega, x + y) h(T_{-x-y}\omega, -y) \right. \\ & \quad \left. + f(\omega, x + y)(-y_j) h(T_{-x-y}\omega, -y) \right) \sigma((\omega, x + y), (T_{-x-y}\omega, -y)) \\ &= x_j \left( \sum_{(y, 0_{d-k}) \in \mathcal{L}^{(\omega)} - x} f(\omega, x + y) h(T_{-x-y}\omega, -y) \sigma((\omega, x + y), (T_{-x-y}\omega, -y)) \right) \\ &= X_j(f \cdot h)(\omega, x). \end{aligned}$$

It follows that  $X_j \circ v \circ |f\rangle - v \circ |f\rangle \circ X_j = v \circ |X_j f\rangle$ , which is a bounded adjointable operator. The left and right Clifford actions on  $\bigwedge^* \mathbb{R}^{d-k} \hat{\otimes} \bigwedge^* \mathbb{R}^k$  are given by

$$\begin{aligned} \rho^l \hat{\otimes} 1(\omega_1 \hat{\otimes} \omega_2) &= (e_l \wedge \omega_1 - \iota(e_l)\omega_1) \hat{\otimes} \omega_2, \\ 1 \hat{\otimes} \rho^j(\omega_1 \hat{\otimes} \omega_2) &= (-1)^{|\omega_1|} \omega_1 \hat{\otimes} (e_j \wedge \omega_2 - \iota(e_j)\omega_2), \\ \gamma^l \hat{\otimes} 1(\omega_1 \hat{\otimes} \omega_2) &= (e_l \wedge \omega_1 + \iota(e_l)\omega_1) \hat{\otimes} \omega_2, \\ 1 \hat{\otimes} \gamma^j(\omega_1 \hat{\otimes} \omega_2) &= (-1)^{|\omega_1|} \omega_1 \hat{\otimes} (e_j \wedge \omega_2 + \iota(e_j)\omega_2), \end{aligned}$$

with  $|\omega|$  the degree of the form and  $\{e_l\}_{l=1}^{d-k}$  and  $\{e_j\}_{j=1}^k$  the standard bases of  $\mathbb{R}^{d-k}$  and  $\mathbb{R}^k$ , respectively.

We relate  $\bigwedge^* \mathbb{R}^{d-k} \hat{\otimes} \bigwedge^* \mathbb{R}^k \cong \bigwedge^* \mathbb{R}^d$ , which sends the  $Cl_{0, d-k} \hat{\otimes} Cl_{0, k} \rightarrow Cl_{0, d}$  by the map on generators,

$$\rho^l \hat{\otimes} 1 \mapsto \rho^l, \quad 1 \hat{\otimes} \rho^j \mapsto \rho^{d-k+j}$$

with  $l \in \{1, \dots, d-k\}$  and  $j \in \{1, \dots, k\}$  (see [44, §2.16]). There is an analogous map for the right action of  $Cl_{d-k, 0} \hat{\otimes} Cl_{k, 0}$ .

This leads us to conclude that the unbounded Kasparov module

$$\left( C_c(\mathcal{G}, \sigma) \hat{\otimes} Cl_{0,d}, E_{C(\Omega_0)} \hat{\otimes} \bigwedge^* \mathbb{R}^d, \sum_{l=k+1}^d X_l \hat{\otimes} \gamma^{l-k} + \sum_{j=1}^k X_j \hat{\otimes} \gamma^{d-k+j} \right), \tag{10}$$

represents the Kasparov product  $[{}_d\lambda_k] \hat{\otimes}_{C_r^*(\Upsilon_k, \sigma)} [{}_k\lambda_{\Omega_0}]$ , because it satisfies the hypotheses of [55, Theorem 13]. Its operator satisfies the connection condition as shown above, the domain of the operator is included in the domain of  $\sum_{l=k+1}^d X_l \hat{\otimes} \gamma^{l-k}$ , and since the  $X_j \hat{\otimes} \gamma^j$  mutually anti-commute, the positivity condition is satisfied as well.

The Kasparov module (10) recovers the bulk module  ${}_d\lambda_{\Omega_0}$  up to a reordering of the Clifford basis, as we now show. We consider the map  $\eta_{d-k}(\gamma^j) = \gamma^{j-(d-k)}$  on  $Cl_{d,0}$  where we identify  $\gamma^l = \gamma^{d-l}$  if  $l \leq 0$ . We define the same map on  $\rho^j$  and  $Cl_{0,d}$ . The map  $\eta$  is an automorphism of Clifford algebras but may reverse the canonical orientation, namely

$$\begin{aligned} \eta_{d-k}(\omega_{Cl_{d,0}}) &= \eta_{d-k}(\gamma^1 \dots \gamma^d) = \gamma^{k+1} \dots \gamma^d \gamma^1 \dots \gamma^k \\ &= (-1)^{k(d-k)} \gamma^1 \gamma^2 \dots \gamma^d = (-1)^{k(d-k)} \omega_{Cl_{d,0}}, \end{aligned}$$

with the same result for the orientation of  $Cl_{0,d}$ . We can apply the map  $\eta_{d-k}$  to obtain the bulk cycle  ${}_d\lambda_{\Omega_0}$  but at the expense that at the level of  $KK$ -classes  $[x] \mapsto (-1)^{k(d-k)}[x]$  [44, §5, Theorem 3]. This finishes the proof.  $\square$

**3.4.1. Another Factorisation.** Let us also show another way our Kasparov modules can be factorised using a different short exact sequence. Starting with  $\Upsilon_k, \Upsilon_{k-1}$  is a closed subgroupoid and we can build the  $C^*$ -bimodule  $F_{C_r^*(\Upsilon_{k-1}, \sigma)}$  via the restriction  $C_c(\Upsilon_k, \sigma) \rightarrow C_c(\Upsilon_{k-1}, \sigma)$ . Applying Proposition 1.13, we obtain the unbounded Kasparov module

$${}_k\lambda_{k-1} = \left( C_c(\Upsilon_k, \sigma) \hat{\otimes} Cl_{0,1}, F_{C_r^*(\Upsilon_{k-1}, \sigma)} \hat{\otimes} \bigwedge^* \mathbb{R}, X_k \hat{\otimes} \gamma \right)$$

and for  $(\Pi_k f)(\omega, y) = \chi_+(y_k) f(\omega, y)$  with  $\chi_+$  as in Sect. 1.6, we have an extension

$$\begin{aligned} 0 \rightarrow C_r^*(\Upsilon_k \times \Upsilon_k / \Upsilon_{k-1}, \sigma) \rightarrow C^*(\Pi_k C_r^*(\Upsilon_k, \sigma) \Pi_k, \\ C_r^*(\Upsilon_k \times \Upsilon_k / \Upsilon_{k-1}, \sigma)) \rightarrow C_r^*(\Upsilon_k, \sigma) \rightarrow 0. \end{aligned}$$

**Theorem 3.7.** *Taking the Kasparov product,*

$$[{}_d\lambda_k] \hat{\otimes}_{C_r^*(\Upsilon_k, \sigma)} [{}_k\lambda_{k-1}] = (-1)^{d-k} [{}_d\lambda_{k-1}].$$

*Furthermore, our equivalence is at the unbounded level up to a permutation of the Clifford basis.*

*Proof.* The proof follows a very similar argument as the proof of Theorem 3.6. First, we define a map

$$v : C_c(\mathcal{G}, \sigma) \otimes_{C_c(\Upsilon_k, \sigma)} C_c(\Upsilon_k, \sigma)_{C_c(\Upsilon_{k-1}, \sigma)} \rightarrow E_{C_r^*(\Upsilon_{k-1}, \sigma)}^{d-(k-1)}, \quad f \otimes h_k \mapsto f \cdot h_k$$

where we consider  $f \cdot h_k$  as an element in  $E_{C_r^*(\Upsilon_{k-1}, \sigma)}^{d-(k-1)}$ . One can check analogously to the proof of Theorem 3.5 that this map extends to a unitary isomorphism of  $C^*$ -modules

$$v : E^{d-k} \otimes_{C_r^*(\Upsilon_k, \sigma)} F_{C_r^*(\Upsilon_{k-1}, \sigma)} \rightarrow E_{C_r^*(\Upsilon_{k-1}, \sigma)}^{d-(k-1)}.$$

Similarly, we check that

$$(X_k \circ v \circ |f\rangle - v \circ |f\rangle \circ X_k)h = (X_k f) \cdot h = v \circ |X_j f\rangle(h)$$

defines a bounded operator and for  $j \in \{k + 1, \dots, d\}$ ,  $(X_j \otimes 1) \mapsto X_j$  as  $X_j$  is right  $C_r^*(\Upsilon_k)$ -linear.

Applying the isomorphism and grouping together the Clifford actions, we obtain the unbounded Kasparov module

$$\left( C_c(\mathcal{G}, \sigma) \hat{\otimes} Cl_{0, d-(k-1)}, E_{C_r^*(\Upsilon_k)}^{d-(k-1)} \hat{\otimes} \bigwedge^* \mathbb{R}^{d-(k-1)}, X_k \hat{\otimes} \gamma^{d-k+1} + \sum_{j=k+1}^d X_j \hat{\otimes} \gamma^{j-k} \right),$$

which as before satisfies [55, Theorem 13] and thus represents the Kasparov product. To relate this  $KK$ -cycle to  ${}_d\lambda_{k-1}$ , we correct the Clifford labelling by the map  $\gamma^j \mapsto \gamma^{j+1}$  for  $1 \leq j \leq d - k$  and  $\gamma^{d-k+1} \mapsto \gamma^1$ . Such a map will change the orientation of  $Cl_{0, d-(k-1)}$  and  $Cl_{d-(k-1), 0}$  by a factor of  $(-1)^{d-k}$ . The result follows. □

### 4. Spectral Triple Constructions

We now present two constructions of (semifinite) spectral triples obtained from localising the bulk  $KK$ -cycle for  $(C_r^*(\mathcal{G}, \sigma), C(\Omega_0))$  over a state of  $C(\Omega_0)$ . Their index-theoretic properties are discussed in Sect. 6.

#### 4.1. The Evaluation Spectral Triple

We can directly construct a spectral triple on  $\ell^2(\mathcal{L}(\omega))$  by considering the internal product of the Kasparov module  ${}_d\lambda_{\Omega_0}$  with the trivially graded Kasparov module  $ev_\omega = (C(\Omega_0), ev_\omega \mathbb{R}_{\mathbb{R}}, 0)$  coming from the evaluation map on  $C(\Omega_0) \rightarrow \mathbb{R}$  (or  $\mathbb{C}$ ). This spectral triple was considered in [21] for complex algebras. The Kasparov module  $ev_\omega$  gives a class in  $KKO(C(\Omega_0), \mathbb{R})$  or  $KK(C(\Omega_0), \mathbb{C})$  if the algebra and space is complex. If we take the internal product, then

$$\begin{aligned} & \left( C_c(\mathcal{G}, \sigma) \hat{\otimes} Cl_{0, d}, E_{C(\Omega_0)} \hat{\otimes} \bigwedge^* \mathbb{R}^d, X = \sum_{j=1}^d X_j \hat{\otimes} \gamma^j \right) \hat{\otimes}_{C(\Omega_0)} (C(\Omega_0), ev_\omega \mathbb{R}_{\mathbb{R}}, 0) \\ & \cong \left( C_c(\mathcal{G}, \sigma) \hat{\otimes} Cl_{0, d}, (E_{C(\Omega_0)} \otimes_{ev_\omega} \mathbb{R}) \hat{\otimes} \bigwedge^* \mathbb{R}^d, \sum_{j=1}^d X_j \otimes 1 \hat{\otimes} \gamma^j \right). \end{aligned}$$

There is an isometric isomorphism  $E_{C(\Omega_0)} \otimes_{ev_\omega} \mathbb{R} \rightarrow \ell^2(s^{-1}(\omega))$  (see for instance [52, p. 50]). Since

$$s^{-1}(\omega) = \{(T_{-x}\omega, -x) : x \in \mathcal{L}^{(\omega)}\} \simeq \mathcal{L}^{(\omega)}, \quad (T_{-x}\omega, -x) \mapsto x,$$

the Hilbert space  $\ell^2(s^{-1}(\omega))$  can be canonically identified with  $\ell^2(\mathcal{L}^{(\omega)})$ . This gives a map

$$\rho_\omega : E_{C(\Omega_0)} \otimes_{\text{ev}_\omega} \mathbb{R} \rightarrow \ell^2(\mathcal{L}^{(\omega)}), \quad \rho_\omega(f \otimes t)(x) = tf(T_{-x}\omega, -x),$$

and the action of  $C_c(\mathcal{G}, \sigma)$  is then computed to be

$$\begin{aligned} \rho_\omega(\pi(f_1)f_2)(x) &= (f_1 * f_2)(T_{-x}\omega, -x) \\ &= \sum_{y \in \mathcal{L}^{(\omega)} - x} \sigma((T_{-x}\omega, y), (T_{-x-y}\omega, -x - y))f_1(T_{-x}\omega, y)f_2(T_{-x-y}\omega, -x - y) \\ &= \sum_{u \in \mathcal{L}^{(\omega)}} \sigma((T_{-x}\omega, u - x), (T_{-u}\omega, -u))f_1(T_{-x}\omega, u - x)f_2(T_{-u}\omega, -u) \\ &= \sum_{u \in \mathcal{L}^{(\omega)}} \sigma((T_{-x}\omega, u - x), (T_{-u}\omega, -u))f_1(T_{-x}\omega, u - x)(\rho_\omega f_2)(u). \end{aligned}$$

Hence, for  $f \in C_c(\mathcal{G}, \sigma)$  the representation of  $C_r^*(\mathcal{G}, \sigma)$  on  $\ell^2(\mathcal{L}^{(\omega)})$  is given by

$$(\pi_\omega(f)\psi)(x) = \sum_{y \in \mathcal{L}^{(\omega)}} \sigma((T_{-x}\omega, y - x), (T_{-y}\omega, -y))f(T_{-x}\omega, y - x)\psi(y).$$

**Proposition 4.1** ([21], Proposition 5.1). *The triple*

$${}_d\lambda_\omega = \left( C_c(\mathcal{G}, \sigma) \hat{\otimes} Cl_{0,d}, \pi_\omega \ell^2(\mathcal{L}^{(\omega)}) \hat{\otimes} \bigwedge^* \mathbb{R}^d, \sum_{j=1}^d X_j \hat{\otimes} \gamma^j \right)$$

is a  $QC^\infty$  and  $d$ -summable spectral triple. If  $\omega, \omega' \in \Omega_0$  are such that  $\omega' = T_{-a}\omega$ . then the spectral triples  ${}_d\lambda_\omega$  and  ${}_d\lambda_{\omega'}$  define the same class in the  $K$ -homology of  $C_r^*(\mathcal{G}, \sigma)$ .

### 4.2. Invariant Measures and the Semifinite Spectral Triple

Measure theoretic properties of the continuous hull  $\Omega_{\mathcal{L}}$  have been extensively studied. We note a useful result below.

**Proposition 4.2** ([12, 86]). *There is a one-to-one correspondence between measures on  $\Omega_{\mathcal{L}}$  invariant under the  $\mathbb{R}^d$ -action and measures on the unit space  $\Omega_0$  invariant under the groupoid action. Furthermore, if  $\mathcal{L}$  is repetitive, aperiodic and has finite local complexity, then there is a one-to-one correspondence between the invariant measures on  $\Omega_{\mathcal{L}}$  and a canonical positive cone in  $H_d(\Omega_{\mathcal{L}}, \mathbb{R})$ .*

Hence, under additional hypotheses, invariant measure theory on the transversal  $\Omega_0$  can be reduced to a homological condition on the continuous hull  $\Omega_{\mathcal{L}}$ . We will now assume that the unit space  $\Omega_0$  has a probability measure  $\mathbf{P}$  that is invariant under the groupoid action with  $\text{supp}(\mathbf{P}) = \Omega_0$ . Using [58, Theorem 1.1], given the trace

$$\tau_{\mathbf{P}} : C(\Omega_0) \rightarrow \mathbb{C}, \quad f \mapsto \int f(\omega) d\mathbf{P}(\omega)$$

on  $C(\Omega_0)$  we can define the dual trace on finite-rank endomorphisms  $\text{Fin}(E_{C(\Omega_0)}) \subset \mathbb{K}(E_{C(\Omega_0)})$  by the formula

$$\text{Tr}_\tau(\Theta_{e_1, e_2}) = \tau_{\mathbf{P}}((e_2 \mid e_1)_{C(\Omega_0)}),$$

which then extends to a faithful, semifinite and norm lower semicontinuous trace on the von Neumann algebra  $\mathcal{N} = \text{Fin}(E_{C(\Omega_0)})'' \subset \mathcal{B}(\mathcal{H}_\tau)$ , with  $\mathcal{H}_\tau$  the completion of  $C_c(\mathcal{G}, \sigma)$  under the inner product

$$\langle f_1, f_2 \rangle = \int_{\Omega_0} (f_1 \mid f_2)_{C(\Omega_0)}(\omega) \, d\mathbf{P}(\omega) = \int_{\Omega_0} (f_1^* * f_2)(\omega, 0) \, d\mathbf{P}(\omega).$$

We note that for  $f \in C_c(\mathcal{G}, \sigma)$ , the dual trace  $\text{Tr}_\tau$  can also be written by the simple formula

$$\text{Tr}_\tau(f) = \int_{\Omega_0} f(\omega, 0) \, d\mathbf{P}(\omega). \tag{11}$$

The semifinite trace we use is quite abstract but can be related to the so-called trace per unit volume if we also assume ergodicity.

**Proposition 4.3** ([21], Proposition 4.23). *If the measure on  $\Omega_{\mathcal{L}}$  is ergodic under the translation action, then for almost all  $\omega \in \Omega_0$  and any  $f \in C_c(\mathcal{G}, \sigma)$ ,*

$$\text{Tr}_\tau(f) = \text{Tr}_{\text{Vol}}(\pi_\omega(f)) := \lim_{\Lambda \nearrow \mathcal{L}^{(\omega)}} \frac{1}{|\Lambda|} \text{Tr}_{\ell^2(\mathcal{L}^{(\omega)})}(P_\Lambda \pi_\omega(f)),$$

$$P_\Lambda : \ell^2(\mathcal{L}^{(\omega)}) \rightarrow \ell^2(\Lambda),$$

where the limit  $\Lambda \nearrow \mathcal{L}^{(\omega)}$  is an increasing sequence of finite sets approximating  $\mathcal{L}^{(\omega)}$ .

The following result does not require an ergodicity assumption.

**Proposition 4.4.** *The triple*

$${}_d\lambda_\tau = \left( C_c(\mathcal{G}, \sigma) \hat{\otimes} Cl_{0,d}, \mathcal{H}_\tau \hat{\otimes} \bigwedge^* \mathbb{R}^d, \sum_{j=1}^d X_j \otimes \gamma^j \right)$$

is a  $QC^\infty$  and  $d$ -summable semifinite spectral triple relative to  $(\mathcal{N}, \text{Tr}_\tau)$ . Furthermore, for  $f \in C_c(\mathcal{G}, \sigma)$ , the identity

$$\text{res}_{z=d} \text{Tr}_\tau(\pi(f)(1 + |X|^2)^{-s/2}) = \text{Vol}_{d-1}(S^{d-1}) \text{Tr}_\tau(f),$$

holds true.

*Proof.* The representation of  $C_r^*(\mathcal{G}, \sigma)$  on  $E_{C(\Omega_0)}$  gives a representation  $\pi : C_r^*(\mathcal{G}, \sigma) \rightarrow \mathcal{B}(\mathcal{H}_\tau)$  as  $\mathcal{H}_\tau \cong E \otimes_{C(\Omega_0)} L^2(\Omega_0, \mathbf{P})$ . This representation retains the property that  $[X_j, \pi(f)] = \pi(\partial_j f)$  and, as such,  $[|X|^k, \pi(f)]$  is well defined and bounded for all  $k \in \mathbb{N}$ . To consider the summability, we first note that  $(1 + X^2)^{-s/2} = (1 + |X|^2)^{-s/2} \hat{\otimes} 1_{\bigwedge^* \mathbb{R}^d}$  and so it suffices to prove the summability of  $(1 + |X|^2)^{-s/2}$ . We then observe that the space of trace class elements under the dual trace  $\mathcal{L}^1(\mathcal{N}, \text{Tr}_\tau)$  contains the trace class operators on the



space  $\int_{\Omega_0}^{\oplus} \ell^2(\mathcal{L}(\omega)) \, d\mathbf{P}(\omega)$  and, on this subalgebra, the dual trace acts as the usual trace on the direct integral. With this in mind, we first compute

$$\begin{aligned} & (\pi(f)(1 + |X|^2)^{-s/4}\psi)(T_{-x}\omega, -x) \\ &= \sum_{y \in \mathcal{L}(\omega)} \sigma((T_{-x}\omega, y - x), (T_{-y}\omega, -y))f(T_{-x}\omega, y - x) \\ & \quad (1 + |y|^2)^{-s/4}\psi(T_{-y}\omega, -y). \end{aligned}$$

Hence, the ‘integral kernel’ of this operator is

$$k_f(\omega; x, y) = \sigma((T_{-x}\omega, y - x), (T_{-y}\omega, -y))f(T_{-x}\omega, y - x)(1 + |y|^2)^{-s/4}.$$

Similarly, one can compute that the integral kernel of  $(1 + |X|^2)^{-s/4}\pi(f^*)$  is

$$k_{f^*}(\omega; x, y) = \sigma((T_{-x}\omega, y - x), (T_{-y}\omega, -y))f^*(T_{-x}\omega, y - x)(1 + |x|^2)^{-s/4}.$$

Then we can estimate the  $\text{Tr}_\tau$ -Hilbert–Schmidt norm

$$\begin{aligned} & \|\pi(f)(1 + |X|^2)^{-s/4}\|_2^2 = \int_{\Omega_0} \sum_{x, y \in \mathcal{L}(\omega)} k_{f^*}(\omega; x, y)k_f(\omega; y, x) \, d\mathbf{P}(\omega) \\ &= \int_{\Omega_0} \sum_{x, y \in \mathcal{L}(\omega)} \sigma((T_{-x}\omega, y - x), (T_{-y}\omega, -y))\sigma(T_{-y}\omega, x - y), (T_{-x}\omega, -x)) \\ & \quad \times f^*(T_{-x}\omega, y - x)f(T_{-y}\omega, x - y)(1 + |x|^2)^{-s/2} \, d\mathbf{P}(\omega) \\ &= \int_{\Omega_0} \sum_{x, y \in \mathcal{L}(\omega)} \sigma((T_{-x}\omega, y - x), (T_{-y}\omega, x - y))\sigma((T_{-x}\omega, 0), (T_{-x}\omega, -x)) \\ & \quad \times |f(T_{-y}\omega, x - y)|^2(1 + |x|^2)^{-s/2} \, d\mathbf{P}(\omega) \\ &= \int_{\Omega_0} \sum_{x, y \in \mathcal{L}(\omega)} |f(T_{-y}\omega, x - y)|^2(1 + |x|^2)^{-s/2} \, d\mathbf{P}(\omega) \\ &= \int_{\Omega_0} \sum_{x \in \mathcal{L}(\omega)} \sum_{u \in \mathcal{L}(\omega) - x} |f(T_{u-x}\omega, u)|^2(1 + |x|^2)^{-s/2} \, d\mathbf{P}(\omega) \\ &\leq C \int_{\Omega_0} \sum_{x \in \mathcal{L}(\omega)} (1 + |x|^2)^{-s/2} \, d\mathbf{P}(\omega) = C \int_{\Omega_0} C_s(\omega) \, d\mathbf{P}(\omega), \end{aligned}$$

where in the third line we have used the cocycle identity, where we then note that

$$\begin{aligned} & \sigma((T_{-x}\omega, y - x), (T_{-y}\omega, x - y))\sigma((T_{-x}\omega, 0), (T_{-x}\omega, -x)) \\ &= \sigma(\xi, \xi^{-1})\sigma(r(\eta), \eta) = 1. \end{aligned}$$

Because Delone subsets of  $\mathbb{R}^d$  display the same summability asymptotics as  $\mathbb{Z}^d$ , we see that  $C_s(\omega)$  is bounded for all  $\omega \in \Omega_0$  and  $s > d$ . Hence, we have that  $\pi(f)(1 + |X|^2)^{-s/4}$  is  $\text{Tr}_\tau$ -Hilbert–Schmidt. Therefore,  $(1 + |X|^2)^{-s/4}\pi(f^*f)(1 + |X|^2)^{-s/4}$  is  $\text{Tr}_\tau$ -trace class for all  $f \in C_c(\mathcal{G}, \sigma)$  and  $s > d$ . In particular,  $(1 + |X|^2)^{-s/2}$  is  $\text{Tr}_\tau$ -trace class for  $s > d$ .

Let us now consider the residue trace of  $\pi(f)(1 + |X|^2)^{-z/2}$  for  $\Re(z) < d$ . By the properties of the dual trace, we can compute the trace by summing along the diagonal of this integral kernel.

$$\begin{aligned} \text{Tr}_\tau (\pi(f)(1 + |X|^2)^{-z/2}) &= \int_{\Omega_0} \sum_{x \in \mathcal{L}(\omega)} k(x, x) \, d\mathbf{P}(\omega) \\ &= \int_{\Omega_0} \sum_{x \in \mathcal{L}(\omega)} \sigma((T_{-x}\omega, 0), (T_{-x}, -x)) f(T_{-x}\omega, 0) (1 + |x|^2)^{-z/2} \, d\mathbf{P}(\omega) \\ &= \int_{\Omega_0} f(\omega, 0) \sum_{x \in \mathcal{L}(\omega)} (1 + |x|^2)^{-z/2} \, d\mathbf{P}(\omega) \\ &= C(z) \int_{\Omega_0} f(\omega, 0) \, d\mathbf{P}(\omega), \end{aligned}$$

where we have used that  $\sigma(r(\xi), \xi) = 1$  for all  $\xi \in \mathcal{G}$  and the invariance of the measure  $\mathbf{P}$  under the groupoid action. For any Delone set  $\omega \in \Omega_0$ , we use an integral approximation to compute that

$$C(z) = \sum_{x \in \mathcal{L}(\omega)} (1 + |x|^2)^{-z/2} = \text{Vol}_{d-1}(S^{d-1}) \frac{\Gamma(\frac{d}{2}) \Gamma(\frac{z-d}{2})}{2\Gamma(\frac{d}{2})} + h(z)$$

with  $h$  a function holomorphic in a neighbourhood of  $\Re(z) = d$ . The function  $C(z)$  has a meromorphic extension to the complex plane with a simple pole at  $z = d$  with  $\text{res}_{z=d} C(z) = \text{Vol}_{d-1}(S^{d-1})$ . The result follows.  $\square$

Also of use to us for complex algebras is the semifinite spectral triple from the spin<sup>c</sup>  $KK$ -cycle in Proposition 1.14. That is,

$${}_d\lambda_\tau^{Sc} = \left( C_c(\mathcal{G}, \sigma), \mathcal{H}_\tau \hat{\otimes} \mathbb{C}^{2^{\lfloor \frac{d}{2} \rfloor}}, \sum_{j=1}^d X_j \hat{\otimes} \gamma^j \right) \tag{12}$$

is a  $QC^\infty$  and  $d$ -summable semifinite spectral triple that is even or odd depending on the parity of  $d$ . We recall that, as the spin and oriented Kasparov modules are equivalent at the level of  $KK$ -theory (up to a renormalisation), we can equivalently consider pairings with the spin semifinite spectral triple.

### 5. Unbounded Fredholm Modules for Lattices with Finite Local Complexity

We will now assume that our lattice  $\mathcal{L}$  has finite local complexity. Recall from Proposition 2.13 that this implies that the transversal  $\Omega_0$  is totally disconnected. In particular, we have an explicit description of the basis of the topology of  $\Omega_0$  by closed and open sets. Namely, for some  $n \in \mathbb{N}$  and  $P \subset B(0; n)$  discrete, the sets  $U_{P,n} = \{\omega \in \Omega_0 : B(0; n) \cap \mathcal{L}(\omega) = P\}$  give a basis of the topology of  $\Omega_0$ , see [47]. We will use these sets to characterise  $\Omega_0$  as the boundary of a rooted tree. This then allows us to use the Pearson–Bellissard construction to obtain a spectral triple and corresponding class in  $KK(C(\Omega_0), \mathbb{C})$ .

We compose this spectral triple with our bulk  $KK$ -cycle via the unbounded Kasparov product. As in [36, Section 6], the resulting operator exhibits mildly unbounded commutators with the algebra  $C_c(\mathcal{G})$  and its bounded transform is a Fredholm module.

Spectral triple constructions for  $C_r^*(\mathcal{G})$  building from the Pearson–Bellissard framework have already appeared in the tiling literature [57, 64]. While the setting of each construction is quite different, it would be interesting to better understand the relationship between these spectral triples and our unbounded Fredholm module.

**5.1. The Pearson–Bellissard Spectral Triple**

In the case that  $\mathcal{L}$  has finite local complexity,  $\Omega_0$  is totally disconnected and can be conveniently described as the boundary of a rooted tree  $\mathcal{T} = \mathcal{T}_{\mathcal{L}}$  using the local patterns  $p \in P_{\mu}$ . The set of vertices of  $\mathcal{T}_{\mathcal{L}}$  is denoted  $\mathcal{V}_{\mathcal{L}}$  and the set of edges by  $\mathcal{E}_{\mathcal{L}}$ . They are given explicitly by

$$\mathcal{V}_{\mathcal{L}} := \{p \in P_{nR} : n \in \mathbb{N}\}, \quad \mathcal{E}_{\mathcal{L}} := \{(p, q) \in P_{nR} \times P_{(n+1)R} : p \subset q\}.$$

Thus, the vertices are the patterns seen at all levels  $nR$  and there is an edge from  $p \in P_{nR}$  to  $q \in P_{(n+1)R}$  if and only if  $p \subset q$ . The root of this tree is the unique element  $\{0\} \in P_0$ . The vertex set  $\mathcal{V}$  is naturally  $\mathbb{N}$ -graded by

$$\mathcal{V}_n := \{p \in \mathcal{V} : p \in P_{nR}\},$$

and we denote the degree of  $v \in \mathcal{V}$  by  $|v|$ . The boundary  $\partial\mathcal{T}$  is defined to be the set of infinite paths  $\alpha = p_0 \cdots p_n \cdots$  with

$$\{0\} = p_0 \subset p_1 \subset \cdots \subset p_n \subset p_{n+1} \subset \cdots$$

Such a boundary point determines a unique set  $\mathcal{L}^{(\alpha)} := \bigcup_{n=0}^{\infty} p_n \subset \mathbb{R}^d$ , and since  $0 \in \mathcal{L}^{(\alpha)}$ , we have  $\mathcal{L}^{(\alpha)} \in \Omega_0$ . Conversely, any element  $\mathcal{L} \in \Omega_0$  defines a boundary point by setting  $p_n := \mathcal{L} \cap B(0; nR)$ .

The topology on the boundary of a tree is defined by the so-called *cylinder sets* associated to vertices

$$\mathcal{C}_p := \{\alpha \in \partial\mathcal{T} : \alpha_{|p|} = p\} \simeq \{\omega \in \Omega_0 : \mathcal{L}^{(\omega)} \cap B(0; nR) = p\} = U_{(nR,p)},$$

where the latter identification is given by sending a boundary point to its associated set. Thus, the topology on  $\partial\mathcal{T}$  matches that on  $\Omega_0$  and the two spaces are homeomorphic. Equivalently, the topology on  $\partial\mathcal{T}$  can be defined through the ultrametric

$$\rho(\alpha, \omega) = \min\{e^{-nR} : \exists p \in P_{nR} \quad \alpha, \omega \in \mathcal{C}_p\}.$$

By a *choice function*, we mean a map  $\tau : \mathcal{V} \rightarrow \partial\mathcal{T}$  such that  $\tau(v) \in \mathcal{C}_v$ . A choice function defines a representation

$$\pi_{\tau} : C(\Omega_0) \rightarrow B(\ell^2(\mathcal{V})), \quad \pi(f)\phi(v) := f(\tau(v))\phi(v).$$

It is straightforward to verify that for any pair of choice functions  $(\tau_+, \tau_-)$  the pair  $(\pi_{\tau_+}, \pi_{\tau_-})$  defines a quasi-homomorphism  $C(\Omega_0) \rightarrow \mathbb{K}(\ell^2(\mathcal{V}))$  and hence a class in  $KK(C(\Omega_0), \mathbb{C})$  [29]. We associate a spectral triple to this data in the spirit of Pearson–Bellissard [74], with some extra flexibility for reasons

similar to those in [36], related to the pathologies of the unbounded Kasparov product.

**Proposition 5.1.** *Let  $(\tau_+, \tau_-)$  be a pair of choice functions,  $\rho$  an ultrametric on  $\Omega_0$  and let  $\zeta : \mathbb{N} \rightarrow \mathbb{R}_{\geq 0}$  be a sequence with  $\zeta_n \rightarrow \infty$  and for which there exists  $C > 0$  such that  $\zeta_n \leq C (\sup_{p \in \mathcal{V}_n} \text{diam}_\rho \mathcal{C}_p)^{-1}$ . The representation*

$$\pi(f) \begin{pmatrix} \phi_+ \\ \phi_- \end{pmatrix} (v) := \begin{pmatrix} f(\tau_+(v))\phi_+(v) \\ f(\tau_-(v))\phi_-(v) \end{pmatrix},$$

and self-adjoint operator

$$D \begin{pmatrix} \phi_+ \\ \phi_- \end{pmatrix} (v) = \begin{pmatrix} 0 & D_- \\ D_+ & 0 \end{pmatrix} \begin{pmatrix} \phi_+ \\ \phi_- \end{pmatrix} (v) := \begin{pmatrix} \zeta_{|v|}\phi_-(v) \\ \zeta_{|v|}\phi_+(v) \end{pmatrix},$$

define a spectral triple  $(\text{Lip}(\Omega_0), \ell^2(\mathcal{V}, \mathbb{C}^2), D)$  for  $C(\Omega_0)$  whose  $K$ -homology class coincides with that of the quasi-homomorphism  $(\pi_{\tau_+}, \pi_{\tau_-})$ .

*Proof.* The only thing to check is that the Lipschitz functions for the metric  $\rho$  have bounded commutators with each such  $D$ . This follows since

$$\|[D, f]\phi(v)\| = \zeta_{|v|} \|f(\tau_+(v)) - f(\tau_-(v))\| \|\phi(v)\|,$$

and by assumption the sequence  $\zeta_{|v|}$  satisfies  $\zeta_{|v|} \leq C \rho(\tau_+(v), \tau_-(v))^{-1}$ . □

The spectral triple constructed in [74, Proposition 8] corresponds to choosing the sequence  $\zeta_n := e^{nR}$ . Here we choose the sequence  $\zeta_n := \log(1+n)$ . Before we proceed, we record the following observation which serves as the main technical tool in our arguments below.

**Lemma 5.2.** *Let  $x, y \in B(0; nR)$  and  $\|x - y\| < r$ . If  $\alpha, \omega \in \Omega_0$  are such that  $x \in \mathcal{L}^{(\omega)}$ ,  $y \in \mathcal{L}^{(\alpha)}$  and  $\rho(T_{-x}\omega, T_{-y}\alpha) \leq e^{-nR}$ , then  $x = y$  and  $\rho(\alpha, \omega) \leq e^{-nR + \|x\|}$ .*

*Proof.* Since  $\rho(T_{-x}\omega, T_{-y}\alpha) \leq e^{-nR}$ , it holds that

$$\mathcal{L}^{(T_{-x}\omega)} \cap B(0; nR) = \mathcal{L}^{(T_{-y}\alpha)} \cap B(0; nR),$$

and  $x, y \in \mathbb{R}^d \cap B(0; nR)$  gives

$$-x, -y \in \mathcal{L}^{(T_{-x}\omega)} \cap B(0; nR) = \mathcal{L}^{(T_{-y}\alpha)} \cap B(0; nR),$$

and since  $\|x - y\| < r$ , it follows that  $x = y$ . Then because

$$B(-x; nR - \|x\|) \subset B(0; nR), \quad T_x(B(-x; nR - \|x\|)) = B(0; nR - \|x\|)$$

it follows that

$$\mathcal{L}^{(\omega)} \cap B(0; nR - \|x\|) = \mathcal{L}^{(\alpha)} \cap B(0; nR - \|x\|).$$

This means that  $\rho(\alpha, \omega) \leq e^{-(nR - \|x\|)} = e^{-nR + \|x\|}$ . □

### 5.2. The Product Operator

We now proceed to describe the product operator (in the sense of [66]) defined from the unbounded Kasparov module of Proposition 1.13 and the Pearson–Bellissard spectral triples of Proposition 5.1. Because the formulas that appear in this section are somewhat involved, we condense our notation for the groupoid  $\mathcal{G}$ . Namely, let  $\xi = (\omega, x) \in \mathcal{G}$  be a generic groupoid element and let  $x(\xi) \in \mathbb{R}^d$  be the image of the cocycle  $(\omega, x) \mapsto x \in \mathbb{R}^d$  with  $x_k(\xi)$  the  $k$ th component,  $x_k$ . Furthermore, to reduce computational complexity, for the remainder of this section we set  $\sigma = 1$ . The case of a non-trivial 2-cocycle twist requires a separate treatment and involves even longer computations, though we expect the analytic subtleties to be similar.

Given a choice function  $\tau : \mathcal{V} \rightarrow \partial\mathcal{T} = \Omega_0$ , consider the fibre product

$$\mathcal{G}_s \times_\tau \mathcal{V} := \{(\xi, v) \in \mathcal{G} \times \mathcal{V} : s(\xi) = \tau(v)\}.$$

Denote by  $L^2(\mathcal{G}_s \times_\tau \mathcal{V})$  the Hilbert space completion of  $C_c(\mathcal{G}_s \times_\tau \mathcal{V})$  in the inner product

$$\langle \phi, \psi \rangle = \sum_{v \in \mathcal{V}} \sum_{\xi, s(\xi) = \tau(v)} \overline{\phi(\xi, v)} \psi(\xi, v).$$

The following lemma is a straightforward verification.

**Lemma 5.3.** *Let  $\tau : \mathcal{V} \rightarrow \Omega_0$  be a choice function. The map*

$$\alpha : C_c(\mathcal{G}) \otimes_{\pi_\tau}^{\text{alg}} C_c(\mathcal{V}) \rightarrow C_c(\mathcal{G}_s \times_\tau \mathcal{V}), \quad \alpha(f \otimes \psi)(\xi, v) := f(\xi)\psi(v),$$

*extends to a unitary isomorphism  $E_{C(\Omega_0)} \otimes_{\pi_\tau} L^2(\mathcal{V}) \xrightarrow{\sim} L^2(\mathcal{G}_s \times_\tau \mathcal{V})$ . The left representation of  $C_c(\mathcal{G})$  is concretely expressed as*

$$f * \phi(\eta, w) = \sum_{s(\xi) = r(\eta)} f(\xi)\phi(\xi^{-1}\eta, v).$$

Using this lemma, we can decompose the tensor product Hilbert space via the choice function,

$$\mathcal{H} = \mathcal{H}_+ \oplus \mathcal{H}_- = L^2(\mathcal{G}_s \times_{\tau_+} \mathcal{V}) \hat{\otimes} \bigwedge^* \mathbb{R}^d \oplus L^2(\mathcal{G}_s \times_{\tau_-} \mathcal{V}) \hat{\otimes} \bigwedge^* \mathbb{R}^d,$$

though we note that  $\mathcal{H}_\pm$  is *not* the decomposition of the tensor product Hilbert space due to the grading, which also takes into account the  $\mathbb{Z}_2$ -graded structure of  $\bigwedge^* \mathbb{R}^d$ . On this Hilbert space, the operator  $X$  from the bulk  $KK$ -cycle in Eq. (4) on page 1993 is mapped to the operator

$$X = \sum_{k=1}^d X_k \hat{\otimes} \gamma^k : \mathcal{H}_\pm \rightarrow \mathcal{H}_\pm, \quad X(\phi \otimes w)(\xi, v) = \sum_{k=1}^d x_k(\xi)\phi(\xi, v) \hat{\otimes} \gamma^k w.$$

We fix  $\varepsilon$  with  $0 < \varepsilon < \frac{r}{2}$ , a discrete lattice  $Y \subset \mathbb{R}^d$  and a uniformly locally finite cover for  $\mathbb{R}^d$  with subordinate partition of unity

$$\mathcal{Y} := \{B(y; \varepsilon)\}_{y \in Y}, \quad \chi_y : B(y; \varepsilon) \rightarrow [0, 1], \quad \sum_{y \in Y} \chi_y^2 = 1,$$

Recalling Proposition 2.18, from  $\mathcal{Y}$ , we consider the sets  $\{V_y\}_{y \in Y}$ ,

$$V_y = \{\xi = (\omega, x) \in \Omega_0 \times \mathbb{R}^d : x \in \mathcal{L}^{(\omega)} \cap B(y; \varepsilon)\},$$

which form an  $s$ -cover of  $\mathcal{G}$ . Consequently, the functions  $\chi_y : \mathcal{G} \rightarrow \mathbb{R}$  define a frame for  $E_{C(\Omega_0)}$ . In order to construct the connection operator, we wish to describe the maps

$$\langle \chi_y^\pm | : L^2(\mathcal{G} \times_{\tau_\pm} \mathcal{V}) \rightarrow \ell^2(\mathcal{V}), \quad |\chi_y^\pm \rangle : \ell^2(\mathcal{V}) \rightarrow L^2(\mathcal{G}_s \times_{\tau_\pm} \mathcal{V}).$$

Since the support  $\chi_y$  is a compact subset of  $B(y; \varepsilon)$ , the convolution product takes the form

$$\begin{aligned} \chi_y^* * f(\eta, v) &= \sum_{\xi \in r^{-1}(r(\eta))} \chi_y^*(\xi) f(\xi^{-1}\eta, v) = \sum_{\xi \in r^{-1}(r(\eta))} \chi_y(\xi^{-1}) f(\xi^{-1}\eta, v) \\ &= \sum_{\{\xi \in s^{-1}(r(\eta)) \cap V_y\}} \chi_y(\xi) f(\xi\eta, v) \\ &= \chi_y(\xi) f(\xi\eta, v), \quad \text{with } \xi \in s^{-1}(r(\eta)) \cap V_y, \end{aligned}$$

and 0 when the latter set is empty. This shows that the maps become

$$\langle \chi_y^\pm | : L^2(\mathcal{G} \times_{\tau_\pm} \mathcal{V}) \rightarrow \ell^2(\mathcal{V}), \quad \langle \chi_y^\pm | \phi(v) := \chi_y(\xi_\pm^y(v)) \phi(\xi_\pm^y(v), v),$$

whenever  $V_y \cap s^{-1}(\tau_\pm(v)) \neq \emptyset$  and  $\xi_\pm^y(v)$  is the unique point in  $V_y \cap s^{-1}(\tau_\pm(v))$ . In case  $V_y \cap s^{-1}(\tau_\pm(v)) = \emptyset$  we have  $\langle \chi_y^\pm | \phi(v) = 0$ . We can now define the operators

$$T_\pm : C_c(\mathcal{G} \times_{\tau_\pm} \mathcal{V}) \rightarrow C_c(\mathcal{G} \times_{\tau_\mp} \mathcal{V}),$$

by

$$T_+ \phi_+(\eta, v) = \sum_y \sum_{\xi \in s^{-1}(\tau_+(v))} \zeta_{|v|} \chi_y(\eta) \chi_y(\xi) \phi_+(\xi, v), \quad s(\eta) = \tau_-(v).$$

The above sum is in fact finite for each  $(\eta, v) \in \mathcal{G} \times_{\tau_-} \mathcal{V}$ , since the summands are nonzero only for those  $y$  with  $\eta \in V_y$  and  $V_y \cap s^{-1}(\tau_+(v)) \neq \emptyset$ . For  $T_-$  we have an analogous formula.

The operators  $T_\pm$  can be viewed as being constructed from the Grassmann connection associated with the frame  $\{\chi_y\}$  as in [66, Section 3.4]. We use the methods developed there to address self-adjointness properties of these operators.

**Lemma 5.4.** *The operator*

$$T := \begin{pmatrix} 0 & T_- \\ T_+ & 0 \end{pmatrix} : C_c(\mathcal{G} \times_{\tau_+} \mathcal{V}) \oplus C_c(\mathcal{G}_s \times_{\tau_-} \mathcal{V}) \rightarrow L^2(\mathcal{G} \times_{\tau_+} \mathcal{V}) \oplus L^2(\mathcal{G}_s \times_{\tau_-} \mathcal{V}),$$

*is essentially self-adjoint.*

*Proof.* For fixed  $z$  the continuous functions

$$(\chi_y | \chi_z)_{C(\Omega_0)}(\omega) = \sum_{\xi \in s^{-1}(\omega)} \chi_y(\xi) \chi_z(\xi),$$

are possibly nonzero only for those  $y$  with  $B(y; \varepsilon) \cap B(z; \varepsilon) \neq \emptyset$ . There are only finitely many such  $y$  since the cover  $\mathcal{Y}$  has finite intersection number.

Moreover, they are locally constant since for  $\rho(\alpha, \omega) < e^{-nR}$  sufficiently small we have

$$(\chi_y | \chi_z)_{C(\Omega_0)}(\alpha) = (\chi_y | \chi_z)_{C(\Omega_0)}(\omega),$$

by Lemma 5.2. Thus, the frame  $\{\chi_y\}_{y \in Y}$  is column finite in the sense of [66, Proposition 3.2], the operators  $\Theta_{z,z} := \Theta_{\chi_z, \chi_z}$  preserve a core for  $T$  by [66, Lemma 3.15] and the commutators  $[T, \Theta_{z,z}]$  extend to bounded operators by [66, Lemma 3.8].

It remains to show that there exists an approximate unit  $u_n$  in the convex hull of the  $\Theta_{z,z}$  that satisfies [66, Definition 3.9]. For a fixed  $n$ , consider the set

$$I_n := \bigcup_{v \subset B(0;nR)} \{y \in Y : s^{-1}(\tau_{\pm}(v)) \cap V_y \neq \emptyset\},$$

and consider the operator  $u_n := \sum_{y \in I_n} \Theta_{y,y}$ . Since

$$[T, \Theta_{y,y}]\phi(\eta, \nu) = \sum_{z, \xi} (\chi_y(\xi)^2 - \chi_y(\eta)^2) \zeta_{|\nu|} \chi_z(\xi) \chi_z(\eta) \phi(\xi, \nu),$$

we see that for  $|v| \geq nR$ , Lemma 5.2 gives that  $x(\xi) = x(\eta)$  and thus  $\chi_y(\xi) = \chi_y(\eta)$ , so we have  $[T, u_n]\phi(\eta, \nu) = 0$ . For  $|v| \leq nR$ , we find

$$\sum_{y \in I_n} (\chi_y(r(\xi))^2 - \chi_y(r(\eta))^2) \zeta_{|v|} \chi_z(\xi) \chi_z(\eta) \phi(\xi, \nu) = 0,$$

because  $\xi, \eta \in s^{-1}(\tau_{\pm}(v))$  and  $\nu \subset B(0;nR)$  so  $\sum_{y \in I_n} \chi_y^2(\xi) = \sum_{y \in I_n} \chi_y^2(\eta) = 1$ . This proves that  $[T, u_n] = 0$ , so  $\{\chi_y\}_{y \in Y}$  form a complete frame and  $T$  is essentially self-adjoint by [66, Theorem 3.18].  $\square$

Denote by

$$\mathcal{C} = \mathcal{C}_+ \oplus \mathcal{C}_- := C_c(\mathcal{G} \times_{\tau_+} \mathcal{V}) \hat{\otimes} \bigwedge^* \mathbb{R}^d \oplus C_c(\mathcal{G} \times_{\tau_-} \mathcal{V}) \hat{\otimes} \bigwedge^* \mathbb{R}^d,$$

the common core for  $X$  and  $T$  and by  $\kappa$  the grading operator on  $\bigwedge^* \mathbb{R}^d$ . Then we have  $X\kappa = -\kappa X$  and  $T\kappa = \kappa T$ . We now address self-adjointness of the densely defined symmetric Hilbert space operator  $D = X + T\kappa$ , using the methods of [63].

**Proposition 5.5.** *The resolvent  $(X \pm i)^{-1}$  maps the core  $\mathcal{C}$  bijectively onto itself. For  $\phi \in \mathcal{C}_{\pm}$  and  $w \in \bigwedge^* \mathbb{R}^d$  we have the estimate*

$$\langle (XT\kappa + T\kappa X)(\phi \otimes w), (XT\kappa + T\kappa X)(\phi \otimes w) \rangle \leq r^2 \|T\kappa(\phi \otimes w)\|^2.$$

*Consequently, the sum operator  $D := X + T\kappa$  is essentially self-adjoint with compact resolvent and the bounded transforms of  $X$  and  $D$  satisfy the Connes–Skandalis positivity and connection conditions (see [28, Appendix A]).*

*Proof.* The statement about the resolvent is immediate since  $X$  is given by Clifford multiplication by a real valued function. Thus, the anti-commutator

$XT\kappa + T\kappa X = (XT - TX)\kappa$  is defined on  $\mathcal{C}$ . The commutator  $XT - TX$  can be explicitly computed as

$$\begin{aligned} (TX - XT)(\phi(\eta, v) \otimes w) &= \sum_{k=1}^d \sum_{y, \xi} (x_k(\xi) - x_k(\eta)) \zeta_{|v|} \chi_y(\xi) \chi_y(\eta) \phi(\xi, v) \otimes \gamma^k w \\ &= \sum_{k=1}^d (x_k(\xi) - x_k(\eta)) (T\phi)(\eta, v) \otimes \gamma^k w, \end{aligned}$$

and since the  $\gamma^k$  are Clifford matrices, it holds that

$$\left\| \sum_{k=1}^d (x_k(\xi) - x_k(\eta)) (T\phi)(\eta, v) \otimes \gamma^k w \right\|^2 = \sum_k \|x_k(\xi) - x_k(\eta)\|^2 \|T\kappa(\phi \otimes w)\|^2.$$

Since  $x(\xi), x(\eta) \in B(y; \varepsilon)$ , it follows that  $\sum_k \|x_k(\xi) - x_k(\eta)\|^2 < 4\varepsilon^2 \leq r^2$ , which gives us the desired estimate. Self-adjointness, compact resolvent and positivity follow from [63, Theorem 4.5, Theorem 7.4, Proposition 7.12], and the connection condition follows from [66, Theorem 4.4].  $\square$

*Remark 5.6.* Note that we have not yet shown that  $D$  has bounded commutators with  $C_c(\mathcal{G})$  and that this is the only obstruction to  $D$  representing the unbounded Kasparov product via [63, Theorem 7.4]. In fact, the operator  $X$  has bounded commutators with all  $f \in C_c(\mathcal{G})$ , but the operator  $T$  does not. In the next section we will show that whenever  $\zeta_{|v|}$  is chosen so that it grows sufficiently slowly, the bounded transform of  $D$  will be a Fredholm module. This Fredholm module will satisfy the Connes–Skandalis connection and positivity conditions by the previous proposition and thus represents the Kasparov product.

### 5.3. The Bounded Transform as a Fredholm Module

Recall that a continuous function  $b : \mathbb{R} \rightarrow [-1, 1]$  is a *normalising function* if it is odd and  $\lim_{x \rightarrow \pm\infty} b(x) = \pm 1$ . To prove that for the right choice of  $\zeta_{|v|}$  we obtain a Fredholm module, we use the following lemma.

**Lemma 5.7.** *Let  $(S, T)$  be a weakly anti-commuting pair of self-adjoint regular operators on the Hilbert  $C^*$ -module  $E_B$ ,  $a \in \text{End}^*(E_B)$ ,  $b : \mathbb{R} \rightarrow [-1, 1]$  normalising function and  $0 < \delta < \frac{1}{2}$ . Write  $D = S + T$  and suppose that  $a(D \pm i)^{-1}$  is compact and the operators*

$$[S, a], \quad (1 + S^2)^{-\delta} [T, a], \quad [T, a] (1 + S^2)^{-\delta},$$

*extend boundedly to all of  $E_B$ . Then  $[b(D), a]$  is a compact operator.*

*Proof.* We need only show that  $[T, a](1 + D^2)^{-\delta}$  and its adjoint extend to bounded operators. Then [36, Theorem A.6] applies to reach the conclusion. Since we have the factorisation

$$[T, a](1 + D^2)^{-\delta} = [T, a](1 + S^2)^{-\delta} (1 + S^2)^\delta (1 + D^2)^{-\delta},$$

it suffices to show that  $(1 + S^2)^\delta (1 + D^2)^{-\delta}$  is bounded. Now if  $R$  is a densely defined operator on  $E_B$  with bounded adjoint, then for  $e_1 \in \text{Dom } R$ ,  $e_2 \in E_B$ ,



$$\|(Re_1 | e_2)_B\| = \|(e_1 | R^*e_2)_B\| \leq C\|e_1\|\|e_2\|$$

so  $R$  is bounded on its domain. Since  $D = S + T$  is self-adjoint and regular on  $\text{Dom } S \cap \text{Dom } T$ , the operators  $(S \pm i)(D \pm i)^{-1}$  are everywhere defined and closed, hence bounded. Their adjoints are the extension of  $(D \pm i)^{-1}(S \pm i)$ . Hence,

$$\begin{aligned} & ((1 + D^2)^{-1}e | e)_B \\ &= ((D + i)^{-1}e | (D + i)^{-1}e)_B \\ &= ((D + i)^{-1}(S + i)(S + i)^{-1}e | (D + i)^{-1}(S + i)(S + i)^{-1}e)_B \\ &\leq C((S + i)^{-1}e | (S + i)^{-1}e)_B = C((1 + S^2)^{-1}e | e)_B. \end{aligned}$$

Thus, it holds that

$$(1 + D^2)^{-1} \leq C(1 + S^2)^{-1}.$$

For  $0 < \delta < \frac{1}{2}$  we have the form estimate

$$((1 + D^2)^{-\delta}(1 + S^2)^\delta e | (1 + D^2)^{-\delta}(1 + S^2)^\delta e)_B \leq C^\delta(e | e)_B,$$

which proves that the adjoint is bounded. □

By Proposition 1.11 every  $f \in C_c(\mathcal{G})$  can be expressed as a finite sum  $f = \sum_y \chi_y f_y$  with  $f_y \in C(\Omega_0)$ . It follows that  $f^* = \sum f_y^* \chi_y^*$ , and since  $C_c(\mathcal{G})$  is closed under the adjoint operation, it suffices to show that for all  $f \in \text{Lip}(\Omega_0)$  and all  $\chi_k^*$  Lemma 5.7 is satisfied for certain choices of  $\zeta_{|v|}$ . From now on, we fix the choice  $\zeta_{|v|} := \log(1 + |v|)$ .

**Lemma 5.8.** *Let  $f \in \text{Lip}(\Omega_0)$ . Then*

$$\|f\|_{\log} := \sup_{\alpha \neq \omega} \frac{|f(\alpha) - f(\omega)|}{\log(1 - \log(\rho(\alpha, \omega)))} < \infty,$$

and so  $\|f(\alpha) - f(\omega)\| \leq \|f\|_{\log} \log(1 - \log(\rho(\alpha, \omega)))$ .

*Proof.* Since  $0 \leq \rho(\alpha, \omega) \leq 1$ , it follows that

$$\rho(\alpha, \omega) \leq \log(e - \log(\rho(\alpha, \omega))) \leq \log(1 - \log(\rho(\alpha, \omega))) + \log(e - 1).$$

So for  $\rho(\alpha, \omega)$  small there is a uniform constant with

$$\rho(x, y) \leq C \log(1 - \log(\rho(\alpha, \omega))),$$

and  $\|f\|_{\log} \leq \frac{\|f\|_{\text{Lip}}}{C}$ . The statement follows. □

**Lemma 5.9.** *Let  $\mathcal{V} := \{V_y\}_{y \in Y}$  be an  $s$ -cover of a groupoid  $\mathcal{G}$  with intersection number  $N$  and  $\chi_y$  a subordinate partition of unity. For  $\eta \in \mathcal{G}$  and  $\omega \in \mathcal{G}^{(0)}$  fixed, the set*

$$Y_{\eta, \omega} := \{(\xi, y) \in s^{-1}(\omega) \times Y : \chi_y(\xi)\chi_y(\eta) \neq 0\},$$

*contains at most  $N$  elements.*

*Proof.* First of all observe that  $(\xi, y) \in Y_{\eta, \omega}$  only if  $\xi, \eta \in V_y$  and there can be at most  $N$  distinct indices  $y$  for which  $\eta \in V_y$ . Secondly if  $(\xi, y), (\xi', y) \in Y_{\eta, \omega}$ , then since  $V_y$  is an  $s$ -cover, it follows that  $\xi = \xi'$ . Thus, there are at most  $N$  distinct pairs  $(\xi, y) \in Y_{\eta, \omega}$ . □

**Proposition 5.10.** *For  $f \in \text{Lip}(\Omega_0)$  and  $\delta > 0$  the operators  $(1 + X^2)^{-\delta}[T, f]$  and  $[T, f](1 + X^2)^{-\delta}$  extend to bounded operators.*

*Proof.* Because  $X^2$  and  $T$  act diagonally on the finite-dimensional Clifford representation space  $\bigwedge^* \mathbb{R}^d$ , it suffices to prove boundedness on  $L^2(\mathcal{G}_s \times_{\tau_+} \mathcal{V}) \oplus L^2(\mathcal{G}_s \times_{\tau_-} \mathcal{V})$ . For  $f \in \text{Lip}(\Omega_0)$  the commutator takes the simple form

$$\begin{aligned}
 & [T_+, f](1 + X^2)^{-\delta} \phi_+(\eta, v) \\
 &= \sum_y \sum_{\xi \in s^{-1}(\tau_+(v))} \zeta_{|v|} (1 + \|x(\xi)\|^2)^{-\delta} (f(r(\xi)) - f(r(\eta))) \chi_y(\eta) \chi_y(\xi) \phi_+(\xi, v),
 \end{aligned}$$

with  $s(\eta) = \tau_-(v)$ . We consider the two cases namely

$$\xi \in J_v := x^{-1}(B(0; |v|R - r)), \quad \xi \notin J_v.$$

In the first case, we have  $\chi_y(\xi), \chi_y(\eta) \neq 0$  only if  $x(\eta) \in B(0; |v|R)$ , and Lemma 5.2 gives that  $x(\xi) = x(\eta)$ . Then applying Lemma 5.8 yields the estimate

$$\|f(r(\xi)) - f(r(\eta))\| \leq \|f\|_{\log} \log(1 + |v|R - \|x(\xi)\|).$$

Since

$$\sup_v \sup_{\xi \in J_v} \log(1 + |v|R - \|x(\xi)\|) (1 + \|x(\xi)\|^2)^{-\delta} \log(1 + |v|) < \infty,$$

and denoting by  $J_v^c$  the complement of  $J_v$ ,

$$\sup_v \sup_{\xi \in J_v^c} \zeta_{|v|} (1 + \|x(\xi)\|^2)^{-\delta} < \infty,$$

so we have the following norm estimates (with  $C$  denoting a generic constant):

$$\begin{aligned}
 & \|[T_+, f](1 + X^2)^{-\delta} \phi_+(\eta, v)\|^2 \\
 &= \sum_{\substack{v \in \mathcal{V}, \\ \eta \in s^{-1}(\tau_-(v))}} \left\| \sum_{\xi \in s^{-1}(\tau_+(v))} \sum_y \zeta_{|v|} (1 + \|x(\xi)\|^2)^{-\delta} \right. \\
 & \quad \left. \times (f(r(\xi)) - f(r(\eta))) \chi_y(\eta) \chi_y(\xi) \phi_+(\xi, v) \right\|^2 \\
 & \leq C \sum_{v, \eta} \left( \sum_{\xi \in s^{-1}(\tau_+(v))} \left\| \sum_y \chi_y(\eta) \chi_y(\xi) \phi_+(\xi, v) \right\| \right)^2 \\
 & \leq C \sum_{v, \eta} \left( \sum_{\xi \in s^{-1}(\tau_+(v))} \sum_y \chi_y(\eta) \chi_y(\xi) \|\phi_+(\xi, v)\| \right)^2 \\
 & \leq CN \sum_{v, \eta} \sum_{\xi \in s^{-1}(\tau_+(v))} \sum_y \chi_y(\eta)^2 \chi_y(\xi)^2 \|\phi_+(\xi, v)\|^2,
 \end{aligned}$$

where we have used Lemma 5.9 and the estimate  $(a_1 + \dots + a_N)^2 \leq N(a_1^2 + \dots + a_N^2)$ . Now we use that for a fixed  $\xi \in s^{-1}(\tau_+(v))$  and  $y \in Y$  with  $\chi_y(\xi) \neq 0$  there is at most one  $\eta \in s^{-1}(\tau_-(v))$  with  $0 \neq \chi_y(\eta) \leq 1$  so

$$\begin{aligned} \|[T_+, f](1 + X^2)^{-\delta} \phi_+(\eta, v)\|^2 &\leq CN \sum_{v \in \mathcal{V}} \sum_{\xi \in s^{-1}(\tau_+(v))} \sum_y \chi_y(\xi)^2 \|\phi_+(\xi, v)\|^2 \\ &\leq CN \sum_{v \in \mathcal{V}} \sum_{\xi \in s^{-1}(\tau_+(v))} \|\phi_+(\xi, v)\|^2 = N \|\phi\|^2, \end{aligned}$$

it follows that  $[T, f](1 + X^2)^{-\delta}$  defines a bounded operator for all  $\delta > 0$ .  $\square$

Next we consider the commutator  $[T, \chi_z^*]$  with  $\chi_z$  the frame elements. We obtain the same statement for them.

**Lemma 5.11.** *For  $\phi \in C_c(\mathcal{G}_s \times_{\tau_+} \mathcal{V})$  we have*

$$\begin{aligned} &\langle [T_+, \chi_z^*] \phi, [T_+, \chi_z^*] \phi \rangle \\ &= \sum_{v \in \mathcal{V}} \sum_{\eta \in s^{-1}(\tau_-(v))} \zeta_{|v|}^2 \\ &\times \left\| \sum_{\xi \in s^{-1}(\tau_+(v))} \sum_{y, \alpha, \beta} (\chi_z(\alpha) \chi_y(\beta \eta) \chi_y(\alpha \xi) - \chi_z(\beta) \chi_y(\eta) \chi_y(\xi)) \phi(\xi, v) \right\|^2, \end{aligned} \tag{13}$$

where we used the shorthand notation  $\sum_{y, \alpha, \beta} := \sum_{\alpha \in s^{-1}(r(\xi))} \sum_{\beta \in s^{-1}(r(\eta))} \sum_{y \in Y}$ .

*Proof.* The formula is obtained by direct calculation. First, we compute the commutator acting on a function  $\phi \in C_c(\mathcal{G}_s \times_{\tau_+} \mathcal{V})$ :

$$\begin{aligned} [T_+, \chi_z] \phi(\eta, v) &= \sum_y \sum_{\xi \in s^{-1}(\tau_+(v))} \sum_{\alpha \in r^{-1}(r(\xi))} \zeta_{|v|} \chi_y(\eta) \chi_y(\xi) \chi_z(\alpha) \phi(\alpha^{-1} \xi, v) \\ &\quad - \sum_y \sum_{\beta \in r^{-1}(r(\eta))} \sum_{\xi \in s^{-1}(\tau_+(v))} \zeta_{|v|} \chi_z(\beta) \chi_y(\beta^{-1} \eta) \chi_y(\xi) \phi(\xi, v) \\ &= \sum_y \sum_{\xi \in s^{-1}(\tau_+(v))} \sum_{\alpha \in s^{-1}(r(\xi))} \zeta_{|v|} \chi_z(\alpha) \chi_y(\eta) \chi_y(\alpha \xi) \phi(\xi, v) \\ &\quad - \sum_y \sum_{\beta \in r^{-1}(r(\eta))} \sum_{\xi \in s^{-1}(\tau_+(v))} \zeta_{|v|} \chi_z(\beta) \chi_y(\beta^{-1} \eta) \chi_y(\xi) \phi(\xi, v) \\ &= \sum_{y, \xi} \zeta_{|v|} \left( \sum_{\alpha \in s^{-1}(r(\xi))} \chi_z(\alpha) \chi_y(\eta) \chi_y(\alpha \xi) \right. \\ &\quad \left. - \sum_{\beta \in r^{-1}(r(\eta))} \chi_z(\beta) \chi_y(\beta^{-1} \eta) \chi_y(\xi) \right) \phi(\xi, v). \end{aligned}$$

The  $L^2$ -norm of the vector  $[T_+, \chi_z^*]\phi$  is thus computed as

$$\begin{aligned} \langle [T_+, \chi_z^*]\phi, [T_+, \chi_z^*]\phi \rangle &= \sum_{v \in \mathcal{V}} \sum_{\eta \in s^{-1}(\tau_-(v))} \|[T_+, \chi_k^*]\phi(\eta, v)\|^2 \\ &= \sum_{v, \eta} \zeta_{|v|}^2 \left\| \sum_{y, \xi} \left( \sum_{\alpha \in s^{-1}(r(\xi))} \chi_z(\alpha)\chi_y(\eta)\chi_y(\alpha\xi) \right. \right. \\ &\quad \left. \left. - \sum_{\beta \in r^{-1}(r(\eta))} \chi_z(\beta)\chi_y(\beta^{-1}\eta)\chi_y(\xi) \right) \phi(\xi, v) \right\|^2 \\ &= \sum_{v, \eta} \zeta_{|v|}^2 \left\| \sum_{y, \xi} \left( \sum_{\alpha \in s^{-1}(r(\xi))} \sum_{\beta \in s^{-1}(r(\eta))} \chi_z(\alpha)\chi_y(\beta\eta)\chi_y(\alpha\xi) \right. \right. \\ &\quad \left. \left. - \chi_z(\beta)\chi_y(\eta)\chi_y(\xi) \right) \phi(\xi, v) \right\|^2 \\ &= \sum_{v, \eta} \zeta_{|v|}^2 \left\| \sum_{\xi} \sum_{y, \alpha, \beta} (\chi_z(\alpha)\chi_y(\beta\eta)\chi_y(\alpha\xi) - \chi_z(\beta)\chi_y(\eta)\chi_y(\xi)) \phi(\xi, v) \right\|^2. \end{aligned}$$

This is the desired formula. □

In the inner product expression (13) on page 2019, we split the sum over  $\mathcal{V}$  into a sum over

$$\mathcal{V}_z := \{v \in \mathcal{V} : z \in B(0; |v|R - r)\}, \quad \text{and} \quad \mathcal{V} \setminus \mathcal{V}_z = \{v \in \mathcal{V} : z \notin B(0; |v|R - r)\}.$$

The sum over  $\mathcal{V} \setminus \mathcal{V}_z$  is easily seen to define a bounded operator  $B$ . We further examine the expression that occurs inside the norm bars in (13), namely

$$\sum_{\xi \in s^{-1}(\tau_+(v))} \sum_{\alpha \in s^{-1}(r(\xi))} \sum_{\beta \in s^{-1}(r(\eta))} \sum_y (\chi_z(\alpha)\chi_y(\beta\eta)\chi_y(\alpha\xi) - \chi_z(\beta)\chi_y(\eta)\chi_y(\xi)) \phi(\xi, v), \tag{14}$$

for  $v \in \mathcal{V}_z$  and  $\eta \in s^{-1}(\tau_-(v))$  fixed. We need to further distinguish between two cases for  $\xi \in s^{-1}(\tau_+(v))$  with  $\xi, \eta \in V_y$ . For the fixed function  $\chi_z$ , we split the sum over  $\xi \in s^{-1}(\tau_+(v))$  into a sum over

$$J_{(z,v)} := \{\xi \in s^{-1}(\tau_+(v)) : x(\xi) \notin B(0; |v|R - \|z\| - r)\}, \quad \text{and} \\ s^{-1}(\tau_+(v)) \setminus J_{(z,v)},$$

for which we obtain the following expression.

**Lemma 5.12.** *We have an equality*

$$\begin{aligned} &\sum_{\substack{v \in \mathcal{V}, \\ \eta \in s^{-1}(\tau_-(v))}} \zeta_{|v|}^2 \left\| \sum_{\xi \in s^{-1}(\tau_+(v))} \sum_{y, \alpha, \beta} (\chi_z(\alpha)\chi_k(\beta\eta)\chi_y(\alpha\xi) - \chi_z(\beta)\chi_y(\eta)\chi_y(\xi)) \phi(\xi, v) \right\|^2 \\ &= \langle B\phi, B\phi \rangle \end{aligned}$$

$$+ \sum_{\substack{v \in \mathcal{V}_z, \\ \eta \in s^{-1}(\tau_-(v))}} \zeta_{|v|}^2 \left\| \sum_{\xi \in J_{(z,v)}} \sum_{y, \alpha, \beta} (\chi_z(\alpha)\chi_y(\beta)\eta)\chi_y(\alpha\xi) - \chi_z(\beta)\chi_y(\eta)\chi_y(\xi) \right\| \phi(\xi, v) \Big\|^2,$$

where  $B$  is a bounded operator.

*Proof.* We need to show that the sum over

$$\xi \in s^{-1}(\tau_+(v)) \setminus J_{(z,v)} = \{\xi \in s^{-1}(\tau_+(v)) : x(\xi) \in B(0; |v|R - \|z\| - r)\}$$

vanishes. To this end, we prove the implication

$$\xi \in s^{-1}(\tau_+(v)) \setminus J_{(z,v)} \Rightarrow x(\xi) = x(\eta), \quad x(\alpha) = x(\beta), \quad x(\alpha\xi) = x(\beta\eta). \quad (15)$$

Using (15) we deduce that the sum over  $J_{(z,v)}^c := s^{-1}(\tau_+(v)) \setminus J_{(z,v)}$  vanishes, because  $\chi_z(\eta)$  depends only on  $x(\eta)$ :

$$\begin{aligned} & \sum_y \sum_{\xi \in J_{(z,v)}^c} \sum_{\alpha \in s^{-1}(r(\xi))} \sum_{\beta \in s^{-1}(r(\eta))} (\chi_z(\alpha)\chi_y(\beta)\eta)\chi_y(\alpha\xi) \\ & \quad - \chi_z(\beta)\chi_y(\eta)\chi_y(\xi) \phi(\xi, v) \\ & = \sum_{\xi \in J_{(z,v)}^c} \sum_{\alpha \in s^{-1}(r(\xi))} \sum_{\beta \in s^{-1}(r(\eta))} \sum_y \chi_z(\beta) (\chi_y(\beta\eta)^2 - \chi_y(\eta)^2) \phi(\xi, v) = 0, \end{aligned}$$

and we are left with the sum over the complement  $J_{(z,v)}$ . It thus remains to show (15) holds true.

Let  $\eta \in s^{-1}(\tau_-(v))$  and  $\xi \in s^{-1}(\tau_+(v))$  with  $x(\xi), x(\eta) \in B(y; \varepsilon)$  as well as  $\xi \in s^{-1}(\tau_+(v)) \setminus J_{(z,v)}$ . First, observe that we have

$$\mathcal{L}^{(\tau_+(v))} \cap B(0; |v|R) = \mathcal{L}^{(\tau_-(v))} \cap B(0; |v|R),$$

since  $\rho(\tau_+(v), \tau_-(v)) \leq e^{-|v|R}$ . Then since

$$x(\xi) \in B(0; |v|R - \|z\| - r) \subset B(0; |v|R - r)$$

and  $\|x(\xi) - x(\eta)\| < r$  it must hold that  $x := x(\xi) = x(\eta)$ . We also conclude that

$$\rho(T_x \tau_+(v), T_x \tau_-(v)) \leq e^{-|v|R + \|x(\xi)\|},$$

by Lemma 5.2, and thus,

$$\mathcal{L}^{(T_x \tau_+(v))} \cap B(0; |v|R - \|x(\xi)\|) = \mathcal{L}^{(T_x \tau_-(v))} \cap B(0; |v|R - \|x(\xi)\|).$$

For any two elements

$$\alpha = (T_{x(\alpha)} r(\xi), x(\alpha)) \in s^{-1}(r(\xi)) \cap V_z, \quad \beta = (T_{x(\beta)} r(\eta), x(\beta)) \in s^{-1}(r(\eta)) \cap V_z,$$

we have

$$-x(\alpha) \in \mathcal{L}^{(T_x \tau_+(v))} \cap B(z; \varepsilon), \quad -x(\beta) \in \mathcal{L}^{(T_x \tau_-(v))} \cap B(z; \varepsilon).$$

Now for  $x = x(\xi) = x(\eta) \in B(0; |v|R - \|z\| - r)$  we have  $B(z; \varepsilon) \subset B(0; |v|R - \|x(\xi)\|)$ . Using Lemma 5.2, we find

$$\begin{aligned} & -x(\alpha), -x(\beta) \in \mathcal{L}^{(T_x \tau_+(v))} \cap B(0; |v|R - \|x(\xi)\|) \\ & = \mathcal{L}^{(T_x \tau_-(v))} \cap B(0; |v|R - \|x(\xi)\|), \end{aligned}$$

and since  $\|x(\alpha) - x(\beta)\| < r$ , it must hold that  $z := x(\alpha) = x(\beta)$ . Thus,

$$\alpha = (T_{x+w}(\tau_+(v)), w), \quad \beta = (T_{x+w}(\tau_-(v)), w)$$

where  $-w$  is the unique point in  $\mathcal{L}^{(T_x\tau_+(v))} \cap B(z; \varepsilon) = \mathcal{L}^{(T_x\tau_-(v))} \cap B(z; \varepsilon)$ . We then have

$$\alpha\xi = (T_{w+x}(\tau_+(v)), w + x), \quad \beta\eta = (T_{w+x}(\tau_-(v)), w + x),$$

that is  $x(\alpha\xi) = x(\beta\eta)$ . This proves (15) on page 2021. □

**Proposition 5.13.** *Let  $\chi_k$  be the partition of unity elements associated with the  $s$ -cover  $\{V_y\}_{y \in Y}$ . The operators*

$$[T, \chi_k^*](1 + X^2)^{-\delta}, \quad (1 + X^2)^{-\delta}[T, \chi_k^*]$$

*extend boundedly to all of  $L^2(\mathcal{G} \times_{\tau_+} \mathcal{V}) \hat{\otimes} \Lambda^* \mathbb{R}^d \oplus L^2(\mathcal{G} \times_{\tau_-} \mathcal{V}) \hat{\otimes} \Lambda^* \mathbb{R}^d$ .*

*Proof.* We can again ignore the finite-dimensional space  $\Lambda^* \mathbb{R}^d$ , where  $X^2$  and  $T$  act diagonally. Consider

$$\begin{aligned} & \langle [T, \chi_k^*](1 + X^2)^{-\delta} \phi, [T, \chi_k^*](1 + X^2)^{-\delta} \phi \rangle - \langle B\phi, B\phi \rangle \\ &= \sum_{\substack{v \in \mathcal{V}_z, \\ \eta \in s^{-1}(\tau_-(v))}} \zeta_{|v|}^2 \left\| \sum_{\xi \in J(z, v)} (1 + \|x(\xi)\|^2)^{-\delta} \sum_{y, \alpha, \beta} (\chi_k(\alpha)\chi_y(\beta\eta)\chi_y(\alpha\xi) \right. \\ & \quad \left. - \chi_k(\beta)\chi_y(\eta)\chi_y(\xi)) \phi(\xi, v) \right\|^2 \\ &\leq \sum_{\substack{v \in \mathcal{V}_z, \\ \eta \in s^{-1}(\tau_-(v))}} \left( \sum_{\xi \in J(z, v)} \left\| \sum_{y, \alpha, \beta} (\chi_k(\alpha)\chi_y(\beta\eta)\chi_y(\alpha\xi) - \chi_z(\beta)\chi_y(\eta)\chi_y(\xi)) \phi(\xi, v) \right\| \right)^2 \\ &\leq \sum_{\substack{v \in \mathcal{V}_z, \\ \eta \in s^{-1}(\tau_-(v))}} \sum_{\xi \in J(z, v)} N \left\| \sum_{y, \alpha, \beta} (\chi_z(\alpha)\chi_y(\beta\eta)\chi_y(\alpha\xi) - \chi_z(\beta)\chi_y(\eta)\chi_y(\xi)) \right\|^2 \|\phi(\xi, v)\|^2, \end{aligned}$$

where the last inequality follows from Lemma 5.9. We proceed

$$\begin{aligned} & \langle [T, \chi_k^*](1 + X^2)^{-\delta} \phi, [T, \chi_k^*](1 + X^2)^{-\delta} \phi \rangle - \langle B\phi, B\phi \rangle \\ &\leq 2N \sum_{\substack{v \in \mathcal{V}_z, \\ \eta \in s^{-1}(\tau_-(v))}} \sum_{\xi \in J(z, v)} \sum_{y, \alpha, \beta} (\chi_z(\alpha)^2 \chi_y(\alpha\xi)^2 \chi_y(\beta\eta)^2 \\ & \quad + \chi_z(\beta)^2 \chi_y(\xi)^2 \chi_y(\eta)^2) \|\phi(\xi, v)\|^2 \\ &\leq 2N \sum_{v \in \mathcal{V}_z} \sum_{\xi \in J(z, v)} \sum_{\alpha, \beta} (\chi_z(\alpha)^2 + \chi_z(\beta)^2) \|\phi(\xi, v)\|^2 \\ &\leq 4N \sum_{v \in \mathcal{V}_z} \sum_{\xi \in J(z, v)} \|\phi(\xi, v)\|^2 \leq 4N \|\phi\|^2, \end{aligned}$$

and we conclude that  $[T, \chi_k^*](1 + X^2)^{-\delta}$  is bounded for all  $\delta$ . The statement for  $(1 + X^2)^{-\delta}[T, \chi_k^*]$  follows in a similar manner. □

**Theorem 5.14.** *The triple*

$$\left( C_c(\mathcal{G}) \hat{\otimes} Cl_{0,d}, L^2(\mathcal{G}_s \times_{\tau_+} \mathcal{V}) \hat{\otimes} \bigwedge^* \mathbb{R}^d \oplus L^2(\mathcal{G}_s \times_{\tau_-} \mathcal{V}) \hat{\otimes} \bigwedge^* \mathbb{R}^d, X + T\kappa \right)$$

is an  $\varepsilon$ -unbounded Fredholm module for all  $0 < \varepsilon < 1$  in the sense of [36, Definition A.1]. It represents the Kasparov product of the class  $[_d\lambda_{\Omega_0}] \in KK(C_r^*(\mathcal{G}) \hat{\otimes} Cl_{0,d}, C(\Omega_0))$  of Eq. (4) on page 1993 and the quasi-homomorphism  $[(\pi_{\tau_+}, \pi_{\tau_-})] \in KK(C(\Omega_0), \mathbb{C})$ .

*Proof.* By [36, Theorem A.6] the bounded transform of  $X + T\kappa$  is a Fredholm module. By Proposition 5.5 and [28, Theorem A.3], this Fredholm module represents the Kasparov product of the indicated classes.  $\square$

As previously mentioned, it would be interesting to compare the  $K$ -homology class of the  $\varepsilon$ -unbounded Fredholm module from Theorem 5.14 with similar constructions in the tiling literature [57, 64].

We have concretely represented a  $K$ -homology class containing information of both the transversal dynamics and internal structure of the unit space. In Sect. 6.3 we will briefly consider its potential applications to topological phases of lattices or tilings with finite local complexity (e.g. quasicrystals) via the index pairing.

## 6. Index Pairings and Topological Phases

Up to now we have largely been concerned with the  $K$ -homology and  $KK$ -theory of  $C_r^*(\mathcal{G}, \sigma)$ . In this section, we use these constructions and properties to consider homomorphisms on  $K$ -theory. That is, we are interested in product pairings in real or complex  $K$ -theory

$$K_n(C_r^*(\mathcal{G}, \sigma)) \times KK^d(C_r^*(\mathcal{G}, \sigma), C(\Omega_0)) \rightarrow K_{n-d}(C(\Omega_0)). \tag{16}$$

Our motivation for studying such pairings comes from applications to topological phases of Hamiltonians on aperiodic lattices, which we briefly introduce. A low-energy quantum mechanical system with negligible interactions between particles is modelled via a self-adjoint Hamiltonian acting on a complex separable Hilbert space  $\mathcal{H}$ . This Hamiltonian is often an element of or affiliated to a  $C^*$ -algebra of observables. One can then consider underlying symmetries of the Hamiltonian, where Wigner’s theorem implies that such symmetries arise on  $\mathcal{H}$  as projective unitary or anti-unitary representations of a symmetry group [94]. In the case of an anti-unitary representation of  $\mathbb{Z}_2$  (e.g. a time-reversal symmetry), conjugation by the generator of this representation often gives a *Real structure*  $\tau$  on the  $C^*$ -algebra of observables. That is, an anti-linear order-2 automorphism that commutes with the  $*$ -operation.

While other symmetry groups can be considered, for free-fermionic topological phases, one generally considers a symmetry group  $G \subset \mathbb{Z}_2 \times \mathbb{Z}_2$ . The symmetries that generate this group are chiral symmetry (unitary, anti-commutes with Hamiltonian), time-reversal symmetry (anti-unitary, commutes with Hamiltonian) and particle-hole symmetry (anti-unitary,

anti-commutes with Hamiltonian). A key reason for studying such symmetries comes from the following result.

**Proposition 6.1** ([49]). *Let  $h$  be a self-adjoint element in the complex  $C^*$ -algebra  $A$  with a spectral gap at 0 (taking a shift  $h - \mu$  if necessary).*

- (1) *The spectral projection  $\chi_{(-\infty,0]}(h)$  gives a class in  $K_0(A)$ .*
- (2) *If  $h$  has a chiral symmetry, then  $h$  determines an element in  $K_1(A)$ .*
- (3) *If  $h$  has a time-reversal symmetry and/or particle–hole symmetry, then  $h$  determines an element in  $KO_n(A^\tau)$ . The degree  $n$  of the  $KO$ -theory group and the Real structure  $\tau$  is determined by the symmetry of the Hamiltonian (cf. [49, Section 6]).*

*Remarks 6.2.* (1) Proposition 6.1 has appeared in numerous forms, see [19, 34, 53, 93] for example.

- (2) We wish to apply Proposition 6.1 to the case  $A = C_r^*(\mathcal{G}, \sigma)$  (complex  $C^*$ -algebras) so that we can then apply our  $KK$ -theoretic machinery to the case of invertible Hamiltonians  $h \in C_r^*(\mathcal{G}, \sigma)$ . For systems with no anti-linear symmetries, this is no problem. For systems with time-reversal symmetry or particle–hole symmetry, we require the corresponding real subalgebra  $C_r^*(\mathcal{G}, \sigma)^\tau$  to have a presentation as a twisted real groupoid  $C^*$ -algebra,  $C_r^*(\mathcal{G}, \sigma_{\mathbb{R}})_{\mathbb{R}}$  with  $\sigma_{\mathbb{R}}$  a  $O(1)$ -valued twist. This places a restriction on the  $U(1)$ -valued twist  $\sigma$ , but is immediate if the twist is trivial. Such assumptions are to be expected as, for example, the case of magnetic twists from Example 2.19 should in general not be compatible with a time-reversal symmetry.

Hence, we shall from now on assume that our algebra of observables is given by the real or complex transversal groupoid  $C^*$ -algebra and that we have a class in  $K_n(C_r^*(\mathcal{G}, \sigma))$  (real or complex) to take pairings with.

- (3) Because we have an unbounded Kasparov module

$$d\lambda_{\Omega_0} = \left( C_c(\mathcal{G}, \sigma) \hat{\otimes} Cl_{0,d}, E_{C(\Omega_0)} \hat{\otimes} \bigwedge^* \mathbb{R}^d, X = \sum_{j=1}^d X_j \hat{\otimes} \gamma^j \right),$$

the completion  $\mathcal{A} = \overline{C_c(\mathcal{G}, \sigma)}$  of  $C_c(\mathcal{G}, \sigma)$  in the norm  $\|f\| + \|[X, f]\|$  is a Banach  $*$ -algebra that is stable under the holomorphic functional calculus [18]. Having fixed such an algebra  $\mathcal{A}$ , the spectral gap assumption on  $h \in C_r^*(\mathcal{G}, \sigma)$ , means that we can improve Proposition 6.1 and obtain an element of the group  $K_n(\mathcal{A}) \cong K_n(C_r^*(\mathcal{G}, \sigma))$ .

Given a  $K$ -theory class from Proposition 6.1, we can thus consider pairings such as Eq. (16) on page 2023. In general, the pairing in Eq. (16) can be described using a Clifford index similar to Atiyah–Bott–Shapiro [5]. This index then serves as an explicit phase label of the  $K$ -theory class from Proposition 6.1.

Also of importance are numerical pairings, which can be defined in a few ways. One is by point evaluation  $C(\Omega_0) \rightarrow \mathbb{C}$  or  $\mathbb{R}$ , which leads to  $\mathbb{Z}$  or  $\mathbb{Z}_2$ -valued invariants. Alternatively, we fix a faithful and invariant measure on



$\Omega_{\mathcal{L}}$ , which gives an invariant measure on  $\Omega_0$ . This defines a trace on  $C(\Omega_0)$  and a homomorphism  $K_0(C(\Omega_0)) \rightarrow \mathbb{R}$ . In particular, for complex algebras, the composition

$$K_*(C_r^*(\mathcal{G}, \sigma)) \times KK^*(C_r^*(\mathcal{G}, \sigma), C(\Omega_0)) \rightarrow K_0(C(\Omega_0)) \xrightarrow{f} \mathbb{R} \quad (17)$$

can be computed using the semifinite local index formula. The cyclic formula that we obtain from the local index formula is then more amenable to physical interpretation and numerical approximation.

### 6.1. Complex Pairings

For complex algebras, we use the semifinite local index formula to pair complex  $K$ -theory classes in  $K_*(C_r^*(\mathcal{G}, \sigma))$  with the  $\text{spin}^c$  semifinite spectral triple from Eq. (12) on page 2010. Algebraic manipulation of the Dirac operator means that only the top degree term survives as in [16, Appendix]. Then we can evaluate the resolvent cocycle, which uses the residue trace computation from Proposition 4.4. We will simply state the result as the proof follows the same argument as analogous results in [22, 23].

**Proposition 6.3.** *Let  $u$  be a complex unitary in  $M_q(\mathcal{A})$  and  ${}_d\lambda_\tau^{S^c}$  the complex semifinite spectral triple from Eq. (12) on page 2010 with  $d$  odd. Then the semifinite index pairing is given by the formula*

$$\langle [u], [{}_d\lambda_\tau^{S^c}] \rangle = \tilde{C}_d \sum_{\rho \in S_d} (-1)^\rho (\text{Tr}_{\mathbb{C}^q} \otimes \text{Tr}_\tau) \left( \prod_{j=1}^d u^* \partial_{\rho(j)} u \right),$$

$$\tilde{C}_{2n+1} = \frac{2(2\pi i)^n n!}{(2n+1)!},$$

where  $\text{Tr}_{\mathbb{C}^q}$  is the matrix trace on  $\mathbb{C}^q$  and  $S_d$  is the permutation group on  $d$  letters.

If  $p$  is a projection in  $M_q(\mathcal{A})$ , then the pairing with  ${}_d\lambda_\tau^{S^c}$  with  $d$  even is given by

$$\langle [p], [{}_d\lambda_\tau^{S^c}] \rangle = C_d \sum_{\rho \in S_d} (-1)^\rho (\text{Tr}_{\mathbb{C}^q} \otimes \text{Tr}_\tau) \left( p \prod_{j=1}^d \partial_{\rho(j)} p \right), \quad C_{2n} = \frac{(-2\pi i)^n}{n!}.$$

If the measure on the unit space is ergodic, then we can almost surely describe the semifinite index pairing via the usual  $\mathbb{Z}$ -valued index pairing with the evaluation spectral triple from Proposition 4.1. Namely, setting  $F_X = X(1 + X^2)^{-1/2}$  and  $\Pi_q = \frac{1}{2}(1 + F_X) \otimes 1_q$ , we have for almost all  $\omega \in \Omega_0$ ,

$$\begin{aligned} & \text{Index} (\Pi_q \pi_\omega(u) \Pi_q + (1 - \Pi_q)) \\ &= \tilde{C}_d \sum_{\rho \in S_d} (-1)^\rho (\text{Tr}_{\mathbb{C}^q} \otimes \text{Tr}_{\text{Vol}}) \left( \prod_{j=1}^d \pi_\omega(u)^* [X_{\rho(j)}, \pi_\omega(u)] \right), \\ & \text{Index} (\pi_\omega(p) (F_X \otimes 1_q)_+ \pi_\omega(p)) \\ &= C_d \sum_{\rho \in S_d} (-1)^\rho (\text{Tr}_{\mathbb{C}^q} \otimes \text{Tr}_{\text{Vol}}) \left( \pi_\omega(p) \prod_{j=1}^d [X_{\rho(j)}, \pi_\omega(p)] \right), \end{aligned}$$

which was proved by slightly different means in [21].

**6.1.1. Weak Chern Numbers.** Analogous to the construction in Sect. 4.2, we can construct a semifinite spectral triple  ${}_d\lambda_{\tau_k}^{S_c}$  from the Kasparov module  ${}_d\lambda_k$  via the dual trace  $\text{Tr}_{\tau_k}$  constructed from the trace  $\tau_k$  on  $C_r^*(\Upsilon_k, \sigma)$ . This semifinite spectral triple is  $QC^\infty$  and  $(d - k)$ -summable with a residue trace evaluation analogous to Proposition 4.4. The semifinite pairing with  ${}_d\lambda_{\tau_k}^{S_c}$  represents the composition

$$K_{d-k}(C_r^*(\mathcal{G}, \sigma)) \times KK^{d-k}(C_r^*(\mathcal{G}, \sigma), C_r^*(\Upsilon_k, \sigma)) \rightarrow K_0(C_r^*(\Upsilon_k, \sigma)) \xrightarrow{\tau_k} \mathbb{R}. \tag{18}$$

We again use the local index formula to compute this pairing; the interested reader can consult [23], where the proof transfers to this setting without issue.

**Proposition 6.4.** *The composition from Eq. (18) is computed by, for  $d - k$  even and  $p \in M_q(\mathcal{A})$  a projection,*

$$\langle [p], [{}_d\lambda_{\tau_k}^{S_c}] \rangle = C_{d-k} \sum_{\rho \in S_{d-k}} (-1)^\rho (\text{Tr}_\tau \otimes \text{Tr}_{\mathbb{C}^q}) \left( p \prod_{j=k+1}^d \partial_{\rho(j)}(p) \right),$$

$$C_{2n} = \frac{(-2\pi i)^n}{n!}.$$

If  $d - k$  is odd and  $u \in M_q(\mathcal{A})$  is unitary, then

$$\langle [u], [{}_d\lambda_{\tau_k}^{S_c}] \rangle = \tilde{C}_{d-k} \sum_{\rho \in S_{d-k}} (-1)^\rho (\text{Tr}_\tau \otimes \text{Tr}_{\mathbb{C}^q}) \left( \prod_{j=k+1}^d u^* \partial_{\rho(j)}(u) \right),$$

$$\tilde{C}_{2n+1} = \frac{2(2\pi i)^n n!}{(2n + 1)!}.$$

Hence, we recover and extend results from [23, 78].

**6.2. Real Pairings and Analytic Indices**

Our aim for this section is to define an analytic index representing the map

$$KO_n(C_r^*(\mathcal{G}, \sigma)) \times KK O^d(C_r^*(\mathcal{G}, \sigma), C(\Omega_0)) \rightarrow KO_{n-d}(C(\Omega_0))$$

Suppose we are given a gapped Hamiltonian  $h = h^*$  in a  $C^*$ -algebra  $A$  such that  $h$  is compatible with the  $CT$ -symmetry group  $G \subset \{1, T, C, CT\}$ . Then, following the construction in [19, Section 3.3], one is able to construct, up to Morita equivalence, a finitely generated and projective module  $pA_A^{\oplus N}$  with a representation  $Cl_{n,0} \rightarrow \text{End}^*(pA_A^{\oplus N})$  constructed from the symmetry group  $G$ . Note that if the Hamiltonian is particle-hole symmetric, then the projection  $p \in M_N(A)$  is closely related, but not equal, to the Fermi projection  $\chi_{(-\infty, E_F]}(h)$ .

When we apply this construction to the transversal groupoid, we obtain the projective module  $pC_r^*(\mathcal{G}, \sigma)_{C_r^*(\mathcal{G}, \sigma)}^{\oplus N}$  which, with its left  $Cl_{n,0}$  action, is a representative of the class  $[h] \in KO_n(C_r^*(\mathcal{G}, \sigma))$  from Proposition 6.1. The fact that the Hamiltonian is gapped implies that this class can be represented by a projective module over the smooth dense subalgebra  $\mathcal{A} \subset C_r^*(\mathcal{G}, \sigma)$ .

The perspective outlined in [19, 40] is that topological phases are measured via a pairing of this  $K$ -theory class  $[h] \in KO_n(C_r^*(\mathcal{G}, \sigma))$  with a dual element. In our case, this element is precisely the bulk  $KK$ -cycle from Eq. (4) on page 1993. Hence, we compute the product

$$\begin{aligned} & \left( Cl_{n,0}, pC_r^*(\mathcal{G}, \sigma)_{C_r^*(\mathcal{G}, \sigma)}^{\oplus N}, 0 \right) \hat{\otimes}_{C_r^*(\mathcal{G}, \sigma)} d\lambda_{\Omega_0} \\ &= \left( Cl_{n,d}, pE_{C(\Omega_0)}^{\oplus N} \hat{\otimes} \mathbb{R}^{2^d}, p(X \otimes 1_N)p \right) \end{aligned}$$

Making small adjustments (that do not change the  $KK$ -class) if necessary, we can ensure that the product  $pXp$  graded-commutes with the left  $Cl_{n,d}$ -action. We denote by  $F_{pXp}$  the bounded transform of  $pXp$ . If the operator  $F_{pXp}$  is a regular Fredholm operator (as characterised in [41, Section 4.3]), then  $\text{Ker}(F_{pXp})_{C(\Omega_0)}$  is a complemented  $C^*$ -submodule of  $pE_{C(\Omega_0)}^{\oplus N} \hat{\otimes} \wedge^* \mathbb{R}^d$  with a graded left action of  $Cl_{n,d}$ . Furthermore, all index-theoretic information of the Kasparov product is contained in the Clifford module  $\text{Ker}(F_{pXp})_{C(\Omega_0)}$ , see [19, Appendix B]. If  $F_{pXp}$  is not regular, then we can amplify  $F_{pXp}$  to a regular Fredholm operator at the expense that this changes the supporting model  $pE^{\oplus N} \oplus C(\Omega_0)^K$  for some  $K$ . The physical significance of this amplification is not always clear and, as such, needs to be considered on a case by case basis.

We briefly summarise our argument.

**Proposition 6.5.** *If  $F_{pXp}$  is regular, then the  $C^*$ -module  $\text{Ker}(F_{pXp})_{C(\Omega_0)}$  with left  $Cl_{n,d}$ -action represents the Kasparov product of the class  $[h] \in KO_n(C_r^*(\mathcal{G}, \sigma))$  with the bulk  $KK$ -cycle from Eq. (4) on page 1993.*

Let us now associate an analytic index representing the Kasparov product.

**Definition 6.6.** We let  ${}_{r,s}\mathfrak{M}_{C(\Omega_0)}$  be the Grothendieck group of equivalence classes of real  $\mathbb{Z}_2$ -graded right- $C(\Omega_0)$   $C^*$ -modules carrying a graded left representation of  $Cl_{r,s}$ .

Provided  $F_{pXp}$  is regular, the product  $\text{Ker}(F_{pXp})$  determines a class in the quotient group  ${}_{n,d}\mathfrak{M}_{C(\Omega_0)}/i^*({}_{n+1,d}\mathfrak{M}_{C(\Omega_0)})$ , where  $i^*$  comes from restricting a Clifford action of  $Cl_{n+1,d}$  to  $Cl_{n,d}$ . Next, we use an extension of the Atiyah–Bott–Shapiro isomorphism, see [88, §2.3], to make the identification

$${}_{n,d}\mathfrak{M}_{C(\Omega_0)}/i^*{}_{n+1,d}\mathfrak{M}_{C(\Omega_0)} \cong KO_{n-d}(C(\Omega_0)).$$

**Definition 6.7.** If  $F_{pXp}$  is regular, the Clifford index of  $F_{pXp}$  is given by the class

$$\text{Index}_{n-d}(F_{pXp}) = [\text{Ker}(F_{pXp})] \in {}_{n,d}\mathfrak{M}_{C(\Omega_0)}/i^*{}_{n+1,d}\mathfrak{M}_{C(\Omega_0)} \cong KO_{n-d}(C(\Omega_0)).$$

*Remark 6.8* (Range of the pairing). In general it is a difficult task to compute  $KO_{n-d}(C(\Omega_0))$  for a transversal set  $\Omega_0$  that comes from a generic Delone set. However, if our original Delone lattice has finite local complexity, then  $\Omega_0$  is totally disconnected (Proposition 2.13), so by the continuity of the  $K$ -functor,

$$KO_j(C(\Omega_0)) \cong C(\Omega_0, KO_j(\mathbb{R})) = \begin{cases} C(\Omega_0, \mathbb{Z}), & j = 0 \pmod{4}, \\ C(\Omega_0, \mathbb{Z}_2), & j = 1, 2 \pmod{8}, \\ 0, & \text{otherwise.} \end{cases}$$

*Example 6.9* (Spectral triple pairings). By the evaluation map  $ev_\omega : C(\Omega_0) \rightarrow \mathbb{R}$ , we can also pair our  $K$ -theory classes with the evaluation spectral triple  ${}_d\lambda_\omega$  from Proposition 4.1,

$$KO_n(C_r^*(\mathcal{G})) \times KO^d(C_r^*(\mathcal{G})) \rightarrow KO_{n-d}(\mathbb{R}).$$

The  $\mathbb{Z}$  or  $\mathbb{Z}_2$ -valued indices can be measured using results from [6, 40, 46]. Writing these pairings explicitly,

$$[h] \hat{\otimes} [{}_d\lambda_\omega] = \begin{cases} \dim_{\mathbb{R}} \text{Ker}((F_{p_\omega} X_{p_\omega})_+) - \dim_{\mathbb{R}} \text{Ker}((F_{p_\omega} X_{p_\omega})_-), & n - d = 0 \pmod{8} \\ \dim_{\mathbb{R}} \text{Ker}((F_{p_\omega} X_{p_\omega})_+) \pmod{2}, & n - d = 1 \pmod{8} \\ \dim_{\mathbb{C}} \text{Ker}((F_{p_\omega} X_{p_\omega})_+) \pmod{2}, & n - d = 2 \pmod{8} \\ \dim_{\mathbb{H}} \text{Ker}((F_{p_\omega} X_{p_\omega})_+) - \dim_{\mathbb{H}} \text{Ker}((F_{p_\omega} X_{p_\omega})_-), & n - d = 4 \pmod{8} \\ 0, & \text{otherwise} \end{cases}$$

under the decomposition of  $F = \begin{pmatrix} 0 & F_- \\ F_+ & 0 \end{pmatrix}$  by the grading. By considering  $\mathbb{H}$  as an even-dimensional complex space, the quaternionic index naturally takes values in  $2\mathbb{Z}$ .

Let us also briefly remark that complex topological phase labels can also be defined via a Clifford index, though generally indices defined via cyclic cocycles can be more easily related to measurable physical phenomena.

**6.2.1. Extending the Pairings.** In [21], complex bulk indices are extended to a larger algebra constructed from the non-commutative Sobolev spaces  $\mathcal{W}_{r,p}$ , obtained as the closure of  $C_c(\mathcal{G}, \sigma)$  in the norms

$$\|f\|_{r,p} = \sum_{|\alpha| \leq r} \text{Tr}_\tau(|\partial^\alpha f|^p)^{1/p}, \quad r \in \mathbb{N}, \quad p \in [1, \infty), \quad \alpha \in \mathbb{N}^d,$$

$$\partial^\alpha = \partial_1^{\alpha_1} \cdots \partial_d^{\alpha_d}, \quad |\alpha| = \sum_{j=1}^d \alpha_j.$$

We also consider the von Neumann algebra generated by the GNS representation of  $C_r^*(\mathcal{G}, \sigma)$  with respect to the dual trace  $\text{Tr}_\tau$ , denoted by  $L^\infty(\mathcal{G}, \text{Tr}_\tau)$ . Following [21], we define  $\mathcal{A}_{\text{Sob}}$  as the intersection of  $\mathcal{W}_{r,p}$  for  $r, p \in \mathbb{N}$  with  $L^\infty(\mathcal{G}, \text{Tr}_\tau)$ , but emphasise that the topology of  $\mathcal{A}_{\text{Sob}}$  comes only from the Sobolev norms  $\|\cdot\|_{r,p}$  and not the von Neumann norm (see also [22, Section 5]).

If the measure on the continuous hull  $\Omega_{\mathcal{L}}$  is ergodic under the translation action, then  $\mathbb{Z}$  and  $\mathbb{Z}_2$ -valued bulk topological phases can be defined over  $\mathcal{A}_{\text{Sob}}$ . For complex pairings, the Hochschild cocycle from the semifinite spectral triple is also well defined for the Sobolev algebra and, as this cocycle represents the Chern character (because the lower-order terms vanish), the cyclic formulas

for the index also extend to the Sobolev algebra. For real pairings with an ergodic measure, the analytic indices considered in Example 6.9 are almost surely well defined and constant over  $\Omega_0$  in the Sobolev setting. See [22, 40, 46] for a more comprehensive treatment.

For Hamiltonians acting on  $\ell^2(\mathbb{Z}^d) \otimes \mathbb{C}^n$  without a spectral gap but instead a region  $\Delta \subset \sigma(h)$  of dynamical localisation [which implies, amongst other properties, that  $\Delta \cap \sigma_{pp}(h)$  is dense in  $\Delta$ ], the pairings with  $\mathcal{A}_{\text{Sob}}$  are connected to these localised regions via the Aizenman–Molchanov bound [1, 77]. For the case of a general Delone set, the Hamiltonian  $h \in \mathcal{A}_{\text{Sob}}$  acts on the family  $\{\ell^2(\mathcal{L}^{(\omega)})\}_{\omega \in \Omega_0}$ . In the general Delone setting, spectral properties of the Hamiltonian are more difficult to determine. See [35, 62, 83] for more information.

**6.2.2. Weak Indices.** Our  $KK$ -theoretic pairings can also be used to define analytic indices for the pairing with the higher codimension Kasparov modules constructed in Sect. 3.4. Namely, using the  $KK$ -cycles  ${}_d\lambda_k$  from Sect. 3.4, we have a well-defined map,

$$KO_n(C_r^*(\mathcal{G}, \sigma)) \times KK O^{d-k}(C_r^*(\mathcal{G}, \sigma), C_r^*(\Upsilon_k, \sigma)) \rightarrow KO_{n-(d-k)}(C_r^*(\Upsilon_k, \sigma)).$$

Once again this index can be described using Clifford modules.

**6.3. Pairings for Lattices with Finite Local Complexity**

The complex and real pairings from the previous section can be defined for general Delone sets. When the underlying lattice  $\mathcal{L}$  used to construct the continuous hull  $\Omega_{\mathcal{L}}$  has finite local complexity, we can define new numerical phase labels via the  $\varepsilon$ -unbounded Fredholm module from Theorem 5.14.

Recall the unbounded operator  $X + T\kappa$  on  $(L^2(\mathcal{G}_s \times_{\tau_+} \mathcal{V}) \oplus L^2(\mathcal{G}_s \times_{\tau_-} \mathcal{V})) \hat{\otimes} \bigwedge^* \mathbb{R}^d$  whose bounded transform  $b(X + T\kappa)$  is Fredholm and has compact commutators with the representation of  $C_c(\mathcal{G})$  (cf. Lemma 5.7).

There are well-defined index pairings for the  $K$ -theoretic phase of the Hamiltonian  $[h] \in K_n(C_r^*(\mathcal{G}))$  with the  $K$ -homology class  $[b(X + T\kappa)] \in K^d(C_r^*(\mathcal{G}))$  via a Fredholm index for complex phases and a skew-adjoint Fredholm index for real phases,

$$K_n(C_r^*(\mathcal{G})) \times K^d(C_r^*(\mathcal{G})) \rightarrow K_{n-d}(\mathbb{F}), \quad \mathbb{F} = \mathbb{R}, \mathbb{C}.$$

We emphasise that unlike the cases of  $\mathbb{Z}$  or  $\mathbb{Z}_2$ -valued indices that can be defined by the evaluation map  $\text{ev}_{\omega} : \Omega_0 \rightarrow \mathbb{F}$ , these indices depend on the ultrametric structure of the transversal. To more explicitly show this, we note the following result, which is an immediate consequence of the associativity of the Kasparov product.

**Proposition 6.10.** *The index pairing of the  $K$ -theoretic Hamiltonian phase  $[h] \in K_n(C_r^*(\mathcal{G}))$  with the class  $[b(X + T\kappa)] \in K^d(C_r^*(\mathcal{G}))$  is the same as the pairing of the class of the Clifford module  $[\text{Ker}(F_p X_p)] \in K_{n-d}(C(\Omega_0))$  from Proposition 6.5 with the Pearson–Bellissard spectral triple  $[(\pi_{\tau_+}, \pi_{\tau_-})] \in K^0(C(\Omega_0))$  from Proposition 5.1.*

It is worth noting that the index pairing of any fixed class  $\alpha \in K_{n-d}(C(\Omega_0))$  with  $[(\pi_{\tau_+}, \pi_{\tau_-})]$  depends on only finitely many of the values  $\tau_{\pm}(v)$ , viewed as a pair of point evaluations. This follows from the fact that  $K^0(C(\Omega_0))$  is generated by the classes of indicator functions  $\chi_p$  of the cylinder sets  $\mathcal{C}_p$ . For  $|v| > |p|$ , it holds that  $\tau_+(v) \in \mathcal{C}_p$  if and only if  $\tau_-(v) \in \mathcal{C}_p$ , and thus,

$$[\chi_p] \otimes [(\pi_{\tau_+}, \pi_{\tau_-})] = \sum_{|v| \leq |p|} [\chi_p] \otimes [(\pi_{\tau_+}(v), \pi_{\tau_-}(v))].$$

This generic observation was used in [36, Theorem 6.3.1] to determine the rational  $K$ -homology class of an analogous operator. This does not seem to be possible in the present context.

The physical distinction between the indices defined via  $b(X + T\kappa)$  and the more standard bulk index pairings in Sects. 6.1 and 6.2 is currently unclear to us as well. Another question is whether the class of the  $\varepsilon$ -unbounded Fredholm module has a finitely summable representative and, if so, whether the corresponding Chern character gives additional physical information.

**6.4. The Bulk–Boundary Correspondence**

Because our topological phases arise as pairings with the bulk  $KK$ -cycle, the results from Sect. 3 can be used to relate pairings of differing dimension. Recall that we have the extension,

$$0 \rightarrow C_r^*(\mathcal{G} \times \mathcal{G}/\Upsilon, \sigma) \rightarrow \mathcal{T} \rightarrow C_r^*(\mathcal{G}, \sigma) \rightarrow 0,$$

with  $\mathcal{G} \times \mathcal{G}/\Upsilon$  the edge groupoid and  $\mathcal{T}$  acting on a half-infinite space. Suppose that  $[h] \in K_n(C_r^*(\mathcal{G}, \sigma))$  (real or complex) and consider the product with  ${}_d\lambda_{\Omega_0}$ . Then by Theorem 3.5,

$$\begin{aligned} [h] \hat{\otimes}_{C_r^*(\mathcal{G}, \sigma)} [{}_d\lambda_{\Omega_0}] &= (-1)^{d-1} [h] \hat{\otimes}_{C_r^*(\mathcal{G}, \sigma)} ([{}_d\lambda_{d-1}] \hat{\otimes}_{C_r^*(\Upsilon, \sigma)} [{}_{d-1}\lambda_{\Omega_0}]) \\ &= (-1)^{d-1} ([h] \hat{\otimes}_{C_r^*(\mathcal{G}, \sigma)} [{}_d\lambda_{d-1}]) \hat{\otimes}_{C_r^*(\Upsilon, \sigma)} [{}_{d-1}\lambda_{\Omega_0}] \\ &= (-1)^{d-1} \partial[h] \hat{\otimes}_{C_r^*(\Upsilon, \sigma)} [{}_{d-1}\lambda_{\Omega_0}] \end{aligned}$$

with  $\partial[h] \in KO_{n-1}(C_r^*(\mathcal{G} \times \mathcal{G}/\Upsilon, \sigma))$  the image under the boundary map in  $K$ -theory. That is, the pairing with respect to the bulk algebra  $C_r^*(\mathcal{G}, \sigma)$  is non-trivial if and only if the pairing  $\partial[h] \hat{\otimes}_{C_r^*(\Upsilon, \sigma)} [{}_{d-1}\lambda_{\Omega_0}]$  over the edge algebra  $C_r^*(\Upsilon, \sigma)$  [or  $C_r^*(\mathcal{G} \times \mathcal{G}/\Upsilon, \sigma)$ ] is non-trivial.

Furthermore, because the semifinite pairings involve the Kasparov product and then the trace, the bulk–edge correspondence also holds for the Chern number formulas (using the Morita equivalence between  $\text{spin}^c$  and oriented structures). Using the notation from Proposition 6.3,

$$\langle [p], [{}_d\lambda_{\tau}^{Sc}] \rangle = -\langle \partial[p], [{}_{d-1}\lambda_{\tau}^{Sc}] \rangle, \quad \langle [u], [{}_d\lambda_{\tau}^{Sc}] \rangle = \langle \partial[u], [{}_{d-1}\lambda_{\tau}^{Sc}] \rangle.$$

Similarly, our weak or higher codimension pairings also factorise by Theorem 3.7. Namely, via the short exact sequence,

$$0 \rightarrow C_r^*(\Upsilon_k \times \Upsilon_k/\Upsilon_{k-1}, \sigma) \rightarrow \mathcal{I}_k \rightarrow C_r^*(\Upsilon_k, \sigma) \rightarrow 0, \tag{19}$$

we have the equality of pairings,

$$[h] \hat{\otimes}_{C_r^*(\mathcal{G}, \sigma)} [{}_d\lambda_{k-1}] = (-1)^{d-k} [h] \hat{\otimes}_{C_r^*(\mathcal{G}, \sigma)} ([{}_d\lambda_k] \hat{\otimes}_{C_r^*(\Upsilon_k, \sigma)} [{}_k\lambda_{k-1}])$$

$$\begin{aligned}
 &= (-1)^{d-k} ([h] \hat{\otimes}_{C_r^*(\mathcal{G}, \sigma)} [d\lambda_k]) \hat{\otimes}_{C_r^*(\Upsilon_k, \sigma)} [k\lambda_{k-1}] \\
 &= (-1)^{d-k} \partial([h] \hat{\otimes}_{C_r^*(\mathcal{G}, \sigma)} [d\lambda_k]).
 \end{aligned}$$

That is, our weak pairing  $[h] \hat{\otimes}_{C_r^*(\mathcal{G}, \sigma)} [d\lambda_k]$  takes values in  $K_{n-d+k}(C_r^*(\Upsilon_k, \sigma))$  and if we apply the boundary map coming from the short exact sequence in Eq. (19) on page 2030, then up to a sign we obtain the weak pairing  $[h] \hat{\otimes}_{C_r^*(\mathcal{G}, \sigma)} [d\lambda_{k-1}] \in K_{n-d+k-1}(C_r^*(\Upsilon_{k-1}, \sigma))$ . Of course this equality is not necessarily related to the presence of a boundary and is more a property of the Kasparov modules that we use to define the weak topological phases.

For lattices with finite local complexity, we also obtain a factorisation of index pairings of the class  $[h] \in K_n(C_r^*(\mathcal{G}, \sigma))$  with the  $\varepsilon$ -unbounded Fredholm module from Theorem 5.14,

$$\begin{aligned}
 &[h] \hat{\otimes}_{C_r^*(\mathcal{G}, \sigma)} ([d\lambda_{\Omega_0}] \hat{\otimes}_{C(\Omega_0)} [(\pi_{\tau_+}, \pi_{\tau_-})]) \\
 &= (-1)^{d-1} \partial([h] \hat{\otimes}_{C_r^*(\Upsilon, \sigma)} ([d-1\lambda_{\Omega_0}] \hat{\otimes}_{C(\Omega_0)} [(\pi_{\tau_+}, \pi_{\tau_-})])),
 \end{aligned}$$

where the right-hand side is a pairing  $K_{n-1}(C_r^*(\Upsilon)) \times K^{d-1}(C_r^*(\Upsilon)) \rightarrow K_{n-d}(\mathbb{R})$  (or complex).

**6.5. Examples From Materials Science and Meta-Materials**

Constructing model Hamiltonians for generic Delone sets is in general a difficult task, particularly if the underlying lattice is amorphous. However, given  $\omega \in \Omega_0$ , we can write down a basic Hamiltonian by coupling lattice sites with exponential decay and twisting by a magnetic flux,

$$(H_\omega \psi)(x) = \sum_{y \in \mathcal{L}(\omega)} e^{-i\Gamma_{\mathcal{L}(\omega)}(0,x,y)} e^{-\beta|x-y|} \psi(y), \quad \beta > 0, \quad \psi \in \ell^2(\mathcal{L}(\omega)).$$

There is some element  $h \in C_r^*(\mathcal{G}, \sigma)$  such that  $\pi_\omega(h) = H_\omega$  using the pointwise representation. If  $\Delta \subset \mathbb{R}$  is a spectral gap of  $h$ , then the spectral projection  $P_E = \chi_{(-\infty, E]}(h)$  is in the smooth \*-subalgebra  $\mathcal{A} \subset C_r^*(\mathcal{G}, \sigma)$  for any  $E \in \Delta$ . One of the advantages of introducing a magnetic flux into the Hamiltonian is that it can potentially open gaps in the spectrum of  $h$ , as is required by our assumptions. Let us also remark that our choice of Hamiltonian can also be used to model mechanical or gyroscopic meta-materials provided the energies are low, see [68] for example.

In the setting of a spectral gap, our results give that for  $d$  even,

$$\begin{aligned}
 &C_d \sum_{\mu \in S_d} (-1)^\mu \text{Tr}_\tau \left( P_E \prod_{j=1}^d \partial_{\mu(j)}(P_E) \right) \\
 &= -C_{d-1} \sum_{\nu \in S_{d-1}} (-1)^\nu (\text{Tr}_\tau^\Upsilon \otimes \text{Tr}_{\mathcal{H}}) \left( \prod_{j=1}^{d-1} \hat{u}_h^* \partial_{\nu(j)}(\hat{u}_h) \right) \quad (20)
 \end{aligned}$$

where  $\hat{u}_h = e^{2\pi i f(\Pi_d h \Pi_d)}$  with  $\Pi_d$  the projection onto a half-space and  $f$  a function that smoothly goes from 0 to 1 inside the spectral gap  $\Delta$ . We also use that  $C_r^*(\mathcal{G} \rtimes \mathcal{G}/\Upsilon, \sigma) \cong C_r^*(\Upsilon, \sigma) \otimes \mathbb{K}(\mathcal{H})$  and so our boundary semifinite

pairing can be written using the semifinite spectral triple over  $C_r^*(\Upsilon, \sigma) \otimes \mathbb{K}(\mathcal{H})$  relative to  $\text{Tr}_\tau^\Upsilon \otimes \text{Tr}_{\mathcal{H}}$ .

If  $0 \notin \sigma(h)$  (shifting by a constant term if necessary) and  $R_C h R_C = -h$  for some self-adjoint unitary  $R_C \in \mathcal{A}$ , then we can define the Fermi unitary  $U_F = \frac{1}{2}(1 - R_C)(1 - 2P_F)\frac{1}{2}(1 + R_C)$  with  $[U_F] \in K_1(\mathcal{A})$ . Then, for  $d$  odd

$$\begin{aligned}
 & C_d \sum_{\mu \in S_d} (-1)^\mu \text{Tr}_\tau \left( \prod_{j=1}^d U_F^* \partial_{\mu(j)} U_F \right) \\
 &= C_{d-1} \sum_{\nu \in S_{d-1}} (-1)^\nu (\text{Tr}_\tau^\Upsilon \otimes \text{Tr}_{\mathcal{H}}) \left( \text{Ind}(U_F) \prod_{j=1}^{d-1} \partial_{\nu(j)} \text{Ind}(U_F) \right) \quad (21)
 \end{aligned}$$

with  $\text{Ind}(U_F)$  the index map in complex  $K$ -theory.

If the measure on  $\Omega_{\mathcal{L}}$  is ergodic under the translation action, we can replace the dual trace in the left-hand-side Eqs. (20) and (21) with the trace per unit volume on  $\ell^2(\mathcal{L}^{(\omega)})$  for almost all  $\omega \in \Omega_0$ . In this setting, the bulk cyclic formulas continue to be well defined and integer-valued if the assumption on  $\Delta$  is relaxed to a mobility gap (as characterised in [21, Section 6.2]).

We can implement other  $CT$ -symmetries on  $h$  by a choice of Real structure on  $C_r^*(\mathcal{G}, \sigma)$ . Because the equation for  $h$  is quite generic, such symmetries and invariants can be implemented by passing to matrices over  $C_r^*(\mathcal{G}, \sigma)$ . The corresponding bulk and boundary pairings are described in Sect. 6.2, though let us note that if there is no magnetic flux (such as in time-reversal symmetric Hamiltonians), then a model Hamiltonian with spectral gap may be difficult to construct for a generic Delone lattice. However, gaps in the spectrum without a magnetic field may be possible by considering more ordered (but still aperiodic) lattices coming from quasicrystals or substitution tilings.

### Acknowledgements

We thank Jean Bellissard, Magnus Goffeng, Johannes Kellendonk, Aidan Sims and Makoto Yamashita for helpful discussions. We thank the anonymous referees for their careful reading of the manuscript and valuable feedback. CB was supported by a postdoctoral fellowship for overseas researchers from The Japan Society for the Promotion of Science (No. P16728), and both authors were supported by a KAKENHI Grant-in-Aid for JSPS fellows (No. 16F16728). This work is also supported by World Premier International Research Center Initiative (WPI), MEXT, Japan. BM gratefully acknowledges support from the Hausdorff Center for Mathematics and the Max Planck Institute for Mathematics in Bonn, Germany, as well as Tohoku University, Sendai, Japan, for its hospitality. Part of this work was carried out during the Lorentz Center



programme *KK-theory, Gauge Theory and Topological Phases* held in Leiden, Netherlands, in March 2017. We also thank the Leibniz Universität Hannover, Germany, the Radboud University Nijmegen, Netherlands, and the Erwin Schrödinger Institute, University of Vienna, Austria, for hospitality.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

## References

- [1] Aizenman, M., Molchanov, S.: Localization at large disorder and at extreme energies: an elementary derivation. *Commun. Math. Phys.* **157**(2), 245–278 (1993)
- [2] Anderson, J.E., Putnam, I.F.: Topological invariants for substitution tilings and their associated  $C^*$ -algebras. *Ergodic Theory Dyn. Syst.* **18**(3), 509–537 (1998)
- [3] Arici, F., D'Andrea, F., Landi, G.: Pimsner algebras and circle bundles. In: Alpay, D., Cipriani, F., Colombo, F., Guido, D., Sabadini, I., Sauvageot, J.-L. (eds.) *Noncommutative Analysis, Operator Theory and Applications*, pp. 1–25. Springer International Publishing, Cham (2016)
- [4] Arici, F., Kaad, J., Landi, G.: Pimsner algebras and Gysin sequences from principal circle actions. *J. Noncommut. Geom.* **10**(1), 29–64 (2016)
- [5] Atiyah, M.F., Bott, R., Shapiro, A.: Clifford modules. *Topology* **3**(suppl. 1), 3–38 (1964)
- [6] Atiyah, M.F., Singer, I.M.: Index theory for skew-adjoint Fredholm operators. *Inst. Hautes Études Sci. Publ. Math.* **37**, 5–26 (1969)
- [7] Baaj, S., Julg, P.: Théorie bivariante de Kasparov et opérateurs non bornés dans les  $C^*$ -modules hilbertiens. *C. R. Acad. Sci. Paris Sér. I Math.* **296**(21), 875–878 (1983)
- [8] Bandres, M.A., Rechtsman, M.C., Segev, M.: Topological photonic quasicrystals: fractal topological spectrum and protected transport. *Phys. Rev. X* **6**, 011016 (2016)
- [9] Beckus, S., Bellissard, J., De Nittis, G.: Spectral continuity for aperiodic quantum systems I. *General Theory J. Funct. Anal.* **275**(11), 2917–2977 (2018)
- [10] Bellissard, J.:  $K$ -theory of  $C^*$ -algebras in solid state physics. In: *Statistical Mechanics and Field Theory: Mathematical Aspects* (Groningen, 1985), volume 257 of *Lecture Notes in Phys.*, Springer, Berlin pp. 99–156 (1986)
- [11] Bellissard, J.: Delone sets and materials science: a program. In: *Mathematics of Aperiodic Order*, volume 309 of *Progr. Math.*, Birkhäuser/Springer, Basel, pp. 405–428 (2015)
- [12] Bellissard, J., Benedetti, R., Gambaudo, J.-M.: Spaces of tilings, finite telescopic approximations and gap-labeling. *Commun. Math. Phys.* **261**(1), 1–41 (2006)
- [13] Bellissard, J., van Elst, A., Schulz-Baldes, H.: The non-commutative geometry of the quantum Hall-effect. *J. Math. Phys.* **35**, 5373–5451 (1994)
- [14] Bellissard, J., Herrmann, D.J.L., Zarrouati, M.: Hulls of aperiodic solids and gap labelling theorems. *Directions in Mathematical Quasicrystals*. Volume 13 of *CIRM Monograph Series*, pp. 207–259 (2000)
- [15] Belmonte, F., Lein, M., Măntoiu, M.: Magnetic twisted actions on general abelian  $C^*$ -algebras. *J. Operator Theory* **69**(1), 33–58 (2013)

- [16] Benameur, M., Carey, A.L., Phillips, J., Rennie, A., Sukochev, F.A., Wojciechowski, K.P.: An analytic approach to spectral flow in von Neumann algebras. In: Booß-Bavnbek, B., Klimek, S., Lesch, M., Zhang, W. (eds.) *Analysis, Geometry and Topology of Elliptic Operators*, pp. 297–352. World Scientific Publishing, Singapore (2006)
- [17] Blackadar, B.: *K-Theory for Operator Algebras*. Volume 5 of *Mathematical Sciences Research Institute Publications*, Cambridge University Press, Cambridge (1998)
- [18] Blackadar, B., Cuntz, J.: Differential Banach algebra norms and smooth subalgebras of  $C^*$ -algebras. *J. Operator Theory* **26**, 255–282 (1991)
- [19] Bourne, C., Carey, A.L., Rennie, A.: A non-commutative framework for topological insulators. *Rev. Math. Phys.* **28**(2), 1650004 (2016)
- [20] Bourne, C., Kellendonk, J., Rennie, A.: The  $K$ -theoretic bulk-edge correspondence for topological insulators. *Ann. Henri Poincaré* **18**(5), 1833–1866 (2017)
- [21] Bourne, C., Prodan, E.: Non-commutative Chern numbers for generic aperiodic discrete systems. *J. Phys. A* **51**(23), 235202 (2018)
- [22] Bourne, C., Rennie, A.: Chern numbers, localisation and the bulk-edge correspondence for continuous models of topological phases. *Math. Phys. Anal. Geom.* **21**(3), 16 (2018)
- [23] Bourne, C., Schulz-Baldes, H.: Applications of semifinite index theory to weak topological phases. In: Wood, D., de Gier, J., Praeger, C., Tao, T. (eds.) *2016 Matrix Annals*. Springer, Cham (2018)
- [24] Carey, A.L., Neshveyev, S., Nest, R., Rennie, A.: Twisted cyclic theory, equivariant  $KK$ -theory and KMS states. *J. Reine Angew. Math.* **650**, 161–191 (2011)
- [25] Carey, A.L., Phillips, J., Rennie, A., Sukochev, F.A.: The local index formula in semifinite von Neumann algebras i: spectral flow. *Adv. Math.* **202**(2), 451–516 (2006)
- [26] Carey, A.L., Phillips, J., Rennie, A., Sukochev, F.A.: The local index formula in semifinite von Neumann algebras ii: the even case. *Adv. Math.* **202**(2), 517–554 (2006)
- [27] Connes, A.: An analogue of the Thom isomorphism for crossed products of a  $C^*$ -algebra by an action of  $R$ . *Adv. Math.* **39**(1), 31–55 (1981)
- [28] Connes, A., Skandalis, G.: The longitudinal index theorem for foliations. *Publ. RIMS, Kyoto Univ.* **20**, 1139–1183 (1984)
- [29] Cuntz, J.: A new look at  $KK$ -theory. *K-theory* **1**, 31–51 (1987)
- [30] Daenzer, C.: A groupoid approach to noncommutative  $T$ -duality. *Commun. Math. Phys.* **288**(1), 55–96 (2009)
- [31] Ewert, E.E., Meyer, R.: Coarse geometry and topological phases. (2018). [arXiv:1802.05579](https://arxiv.org/abs/1802.05579)
- [32] Fack, T., Kosaki, H.: Generalised  $s$ -numbers of  $\tau$ -measurable operators. *Pac. J. Math.* **123**(2), 269–300 (1986)
- [33] Forrest, A., Hunton, J., Kellendonk, J.: Topological invariants for projection method patterns. *Mem. Am. Math. Soc.* **159**(758), x+120 (2002)
- [34] Freed, D.S., Moore, G.W.: Twisted equivariant matter. *Ann. Henri Poincaré* **14**(8), 1927–2023 (2013)

- [35] Germinet, F., Müller, P., Rojas-Molina, C.: Ergodicity and dynamical localization for Delone–Anderson operators. *Rev. Math. Phys.* **27**(9), 1550020 (2015)
- [36] Goffeng, M., Mesland, B.: Spectral triples and finite summability on Cuntz–Krieger algebras. *Doc. Math.* **20**, 89–170 (2015)
- [37] Goffeng, M., Mesland, B., Rennie, A.: Shift-tail equivalence and an unbounded representative of the Cuntz–Pimsner extension. *Ergodic Theory Dyn. Syst.* **38**(4), 1389–1421 (2018)
- [38] Gomi, K., Thiang, G.C.: Crystallographic bulk-edge correspondence: glide reflections and twisted mod 2 indices. *Lett. Math. Phys.*, online first (2018). <https://doi.org/10.1007/s11005-018-1129-1>
- [39] Gonçalves, D., Ramirez-Solano, M.: On the  $K$ -theory of  $C^*$ -algebras for substitution tilings (a pedestrian version). (2017). [arXiv:1712.09551](https://arxiv.org/abs/1712.09551)
- [40] Grossmann, J., Schulz-Baldes, H.: Index pairings in presence of symmetries with applications to topological insulators. *Commun. Math. Phys.* **343**(2), 477–513 (2016)
- [41] Gracia-Bondía, J.M., Várilly, J.C., Figueroa, H.: *Elements of Noncommutative Geometry*. Birkhäuser Advanced Texts Basler Lehrbücher. Birkhäuser, Boston (2001)
- [42] Hannabuss, K., Mathai, V., Thiang, G.C.: T-duality simplifies bulk-boundary correspondence: the noncommutative case. *Lett. Math. Phys.* **108**(5), 1163–1201 (2018)
- [43] Hunton, J.: Spaces of projection method patterns and their cohomology. In: *Mathematics of Aperiodic Order*, volume 309 of *Progr. Math.*, Birkhäuser/Springer, Basel, pp. 105–135 (2015)
- [44] Kasparov, G.G.: The operator  $K$ -functor and extensions of  $C^*$ -algebras. *Math. USSR Izv.* **16**, 513–572 (1981)
- [45] Kasparov, G.G.: Equivariant  $KK$ -theory and the Novikov conjecture. *Invent. Math.* **91**(1), 147–201 (1988)
- [46] Katsura, H., Koma, T.: The noncommutative index theorem and the periodic table for disordered topological insulators and superconductors. *J. Math. Phys.* **58**(3), 031903 (2018)
- [47] Kellendonk, J.: Noncommutative geometry of tilings and gap labelling. *Rev. Math. Phys.* **7**, 1133–1180 (1995)
- [48] Kellendonk, J.: The local structure of tilings and their integer group of coinvariants. *Commun. Math. Phys.* **187**, 115–157 (1997)
- [49] Kellendonk, J.: On the  $C^*$ -algebraic approach to topological phases for insulators. *Ann. Henri Poincaré* **18**(7), 2251–2300 (2017)
- [50] Kellendonk, J., Putnam, I.: Tilings,  $C^*$ -algebras, and  $K$ -theory. In: *Directions in Mathematical Quasicrystals*, Volume 13 of *CRM Monogr. Ser.*, pp. 177–206. Amer. Math. Soc., Providence, RI (2000)
- [51] Kellendonk, J., Richard, S.: Topological boundary maps in physics. In: Boca, F., Purice, R., Strătilă, Ș. (eds), *Perspectives in Operator Algebras and Mathematical Physics*. Theta Ser. Adv. Math., volume 8, pp. 105–121 Theta, Bucharest (2008). [arXiv:math-ph/0605048](https://arxiv.org/abs/math-ph/0605048)
- [52] Khoshkam, M., Skandalis, G.: Regular representation of groupoid  $C^*$ -algebras and applications to inverse semigroups. *J. Reine Angew. Math.* **546**, 47–72 (2002)

- [53] Kubota, Y.: Notes on twisted equivariant  $K$ -theory for  $C^*$ -algebras. *Int. J. Math.* **27**, 1650058 (2016)
- [54] Kubota, Y.: Controlled topological phases and bulk-edge correspondence. *Commun. Math. Phys.* **349**(2), 493–525 (2017)
- [55] Kucerovsky, D.: The  $KK$ -product of unbounded modules.  *$K$ -Theory* **11**, 17–34 (1997)
- [56] Kumjian, A.: On  $C^*$ -diagonals. *Canad. J. Math.* **38**, 969–1008 (1986)
- [57] Julien, A., Kellendonk, J., Savinien, J.: On the noncommutative geometry of tilings. In: *Mathematics of Aperiodic Order*, volume 309 of *Progr. Math.*, Birkhäuser/Springer, Basel, pp. 259–306 (2015)
- [58] Laca, M., Neshveyev, S.: KMS states of quasi-free dynamics on Pimsner algebras. *J. Funct. Anal.* **211**(2), 457–482 (2004)
- [59] Lagarias, J.C., Pleasants, P.A.B.: Repetitive Delone sets and quasicrystals. *Ergodic Theory Dyn. Syst.* **23**(3), 831–867 (2003)
- [60] Lance, E.C.: *Hilbert  $C^*$ -Modules: A Toolkit for Operator Algebraists*. Volume 210 of *London Mathematical Society Lecture Note Series*, Cambridge University Press, Cambridge (1995)
- [61] Lawson, H.B., Michelsohn, M.-L.: *Spin geometry*. Volume 38 of *Princeton Mathematical Series*, Princeton University Press, Princeton (1989)
- [62] Lenz, D., Peyerimhoff, N., Veselić, I.: Groupoids, von Neumann algebras and the integrated density of states. *Math. Phys. Anal. Geom.* **10**(1), 1–41 (2007)
- [63] Lesch, M., Mesland, B.: Sums of regular self-adjoint operators in Hilbert- $C^*$ -modules. *J. Math. Anal. Appl.* **472**(1), 947–980 (2019)
- [64] Mampusti, M., Whittaker, M.: Fractal spectral triples on Kellendonk’s  $C^*$ -algebra of a substitution tiling. *J. Geom. Phys.* **112**, 224–239 (2017)
- [65] Mesland, B.: Groupoid cocycles and  $K$ -theory. *Münster J. Math.* **4**, 227–250 (2011)
- [66] Mesland, B., Rennie, A.: Nonunital spectral triples and metric completeness in unbounded  $KK$ -theory. *J. Funct. Anal.* **271**(9), 2460–2538 (2016)
- [67] Macho Stadler, M., O’Uchi, M.: Correspondence of groupoid  $C^*$ -algebras. *J. Operator Theory* **42**, 103–119 (1999)
- [68] Mitchell, N.P., Nash, L.M., Hexner, D., Turner, A.M., Irvine, W.T.M.: Amorphous topological insulators constructed from random point sets. *Nat. Phys.* **14**, 380–385 (2018)
- [69] Moutuou, E.M.: *Twisted groupoid  $KR$ -theory*. Ph.D. thesis, Université de Lorraine, Universität Paderborn (2012)
- [70] Moutuou, E.M., Tu, J.-L.: Equivalence of fell systems and their reduced groupoid  $C^*$ -algebras (2011). [arXiv:1101.1235](https://arxiv.org/abs/1101.1235)
- [71] Muhly, P.S., Renault, J., Williams, D.P.: Equivalence and isomorphism for groupoid  $C^*$ -algebras. *J. Operator Theory* **17**, 3–22 (1987)
- [72] Muhly, P.S., Williams, D.P.: Renault’s equivalence theorem for groupoid crossed products. Volume 3 of *NYJM Monographs*, State University of New York, University at Albany, Albany (2008)
- [73] Packer, J.A., Raeburn, I.: Twisted crossed products of  $C^*$ -algebras. *Math. Proc. Cambridge Philos. Soc.* **106**, 293–311 (1989)

- [74] Pearson, J., Bellissard, J.: Noncommutative Riemannian geometry and diffusion on ultrametric Cantor sets. *J. Noncommut. Geom.* **3**(3), 447–480 (2009)
- [75] Putnam, I.F., Spielberg, J.: The structure of  $C^*$ -algebras associated with hyperbolic dynamical systems. *J. Funct. Anal.* **163**(2), 279–299 (1999)
- [76] Prodan, E.: *A Computational Non-commutative Geometry Program for Disordered Topological Insulators*. Springer, Cham (2017)
- [77] Prodan, E., Schulz-Baldes, H.: *Bulk and Boundary Invariants for Complex Topological Insulators: From  $K$ -Theory to Physics*. Springer, Berlin (2016)
- [78] Prodan, E., Schulz-Baldes, H.: Generalized Connes–Chern characters in  $KK$ -theory with an application to weak invariants of topological insulators. *Rev. Math. Phys.* **28**, 1650024 (2016)
- [79] Renault, J.: *A groupoid approach to  $C^*$ -algebras*. Lecture Notes in Mathematics, vol. 793, Springer (1980)
- [80] Rennie, A., Robertson, D., Sims, A.: The extension class and KMS states for Cuntz–Pimsner algebras of some bi-Hilbertian bimodules. *J. Topol. Anal.* **9**(2), 297–327 (2017)
- [81] Rennie, A., Robertson, D., Sims, A.: Groupoid algebras as Cuntz–Pimsner algebras. *Math. Scand.* **120**(1), 115–123 (2017)
- [82] Rieffel, M.: Connes’ analogue for crossed products of the Thom isomorphism. *Contemp. Math.* **10**, 143–154 (1982)
- [83] Rojas-Molina, C.: *Random Schrödinger Operators on discrete structures* (2017). [arXiv:1710.02293](https://arxiv.org/abs/1710.02293)
- [84] Sadun, L.: *Topology of Tiling Spaces*. Volume 46 of University Lecture Series, American Mathematical Society, Providence (2008)
- [85] Sadun, L., Williams, R.W.: Tiling spaces are Cantor set fiber bundles. *Ergodic Theory Dyn. Syst.* **23**(1), 307–316 (2003)
- [86] Savinien, J.: *Cohomology and  $K$ -theory of aperiodic tilings*. Ph.D. thesis, Georgia Institute of Technology (2008)
- [87] Savinien, J., Bellissard, J.: A spectral sequence for the  $K$ -theory of tiling spaces. *Ergodic Theory Dyn. Syst.* **29**(3), 997–1031 (2009)
- [88] Schröder, H.:  *$K$ -Theory for Real  $C^*$ -Algebras and Applications*. Longman Scientific & Technical, Harlow (1993). Copublished in the United States with Wiley, New York
- [89] Sims, A.: Étale groupoids and their  $C^*$ -algebras. To appear In: Szabo, G., Williams, D., Sims, A. (eds), *Operator Algebras and Dynamics: Groupoids, Crossed Products and Rokhlin Dimension*, Birkhäuser (2017). [arXiv:1710.10897](https://arxiv.org/abs/1710.10897)
- [90] Sims, A., Williams, D.P.: Renault equivalence theorem for reduced groupoid  $C^*$ -algebras. *J. Operator Theory* **68**(1), 223–239 (2012)
- [91] Sims, A., Williams, D.P.: An equivalence theorem for reduced Fell bundle  $C^*$ -algebras. *New York J. Math.* **19**, 159–178 (2013)
- [92] Sims, A., Yeend, T.:  $C^*$ -algebras associated to product systems of Hilbert bimodules. *J. Operator Theory* **64**(2), 349–376 (2010)
- [93] Thiang, G.C.: On the  $K$ -theoretic classification of topological phases of matter. *Ann. Henri Poincaré* **17**(4), 757–794 (2016)
- [94] Wigner, E.P.: *Group theory: and its application to the quantum mechanics of atomic spectra*. Expanded and improved ed. Translated from the German by J.

J. Griffin. Volume 5 of Pure and Applied Physics, Academic Press, New York (1959)

- [95] Williamson, P.: Cuntz–Pimsner algebras associated with substitution tilings. Ph.D thesis., University of Victoria (2016)

C. Bourne

WPI-Advanced Institute for Materials Research (WPI-AIMR)

Tohoku University

2-1-1 Katahira, Aoba-ku

Sendai 980-8577

Japan

e-mail: [chris.bourne@tohoku.ac.jp](mailto:chris.bourne@tohoku.ac.jp)

and

RIKEN iTHEMS

2-1 Hirosawa, Wako

Saitama 351-0198

Japan

B. Mesland

Max Planck Institut für Mathematik

Vivatsgasse 7

53111 Bonn

Germany

e-mail: [brammesland@gmail.com](mailto:brammesland@gmail.com)

Communicated by Jean Bellissard.

Received: July 31, 2018.

Accepted: January 12, 2019.