

TWISTED FORMS OF TORIC VARIETIES

ALEXANDER DUNCAN*

Department of Mathematics
University of South Carolina
Columbia, SC 29208, USA
duncan@math.sc.edu

Abstract. We consider the set of forms of a toric variety over an arbitrary field: those varieties which become isomorphic to a toric variety after base field extension. In contrast to most previous work, we also consider arbitrary isomorphisms rather than just those that respect a torus action. We define an injective map from the set of forms of a toric variety to a non-abelian second cohomology set, which generalizes the usual Brauer class of a Severi–Brauer variety. Additionally, we define a map from the set of forms of a toric variety to the set of forms of a separable algebra along similar lines to a construction of A. Merkurjev and I. Panin. This generalizes both a result of M. Blunk for del Pezzo surfaces of degree 6, and the standard bijection between Severi–Brauer varieties and central simple algebras.

1. Introduction

Let k be an arbitrary field. A *toric variety* is a normal variety X over k with a faithful action of a torus T which has a dense open orbit.

When the field k is the complex numbers \mathbb{C} , toric varieties have been extensively studied (see, for example, [Ful93] or [Cox95]). Among the simplest examples is the projective line \mathbb{P}^1 where the torus \mathbb{G}_m acts via

$$\lambda \cdot (x : y) = (\lambda x : y)$$

for $\lambda \in \mathbb{C}^\times$ with open orbit $x, y \neq 0$. In most literature, as part of the definition of “toric variety” the torus is identified with the open orbit on which it acts. This identification is always possible over \mathbb{C} .

However, when the ground field k is not separably closed, the torus T may not be the standard split torus \mathbb{G}_m^n . Indeed, when k is the real numbers \mathbb{R} , the projective line also has an action of the circle group S^1 via multiplication by rotation matrices. Thus, $\mathbb{P}_{\mathbb{R}}^1$ has the structure of a toric variety for the two non-isomorphic tori S^1 and \mathbb{G}_m . (Rotation matrices are diagonalizable over \mathbb{C} so they are isomorphic in that situation.)

DOI: 10.1007/s00031-016-9394-5

*The author was partially supported by National Science Foundation RTG grants DMS 0838697 and DMS 0943832.

Received November 27, 2014. Accepted March 12, 2016.

Published online June 8, 2016.

Corresponding Author: Alexander Duncan, e-mail: duncan@math.sc.edu.

Moreover, for general base fields the open orbit may not even have a rational point. Indeed, when $k = \mathbb{R}$, the conic

$$C = \{x^2 + y^2 + z^2 = 0\} \subset \mathbb{P}^1(\mathbb{R})$$

has an action under the circle group S^1 via rotation matrices on the coordinates x, y . Thus the conic C is a toric variety, but has no \mathbb{R} -points. (Over \mathbb{C} , we obtain $\mathbb{P}^1_{\mathbb{C}}$ by projecting away from the point $(1 : i : 0)$.)

Let us compare our description of the real forms of \mathbb{P}^1 to that of Section 3.1 of [ELFST14]. Suppose $\mathbb{C}(t)$ is the function field of $\mathbb{P}^1_{\mathbb{C}}$ and $\lambda \in \mathbb{C}^\times$ acts by $t \mapsto \lambda t$. We have three anti-holomorphic involutions which normalize the torus action given by

$$t \mapsto \bar{t}, \quad t \mapsto \bar{t}^{-1}, \quad \text{and} \quad t \mapsto -\bar{t}^{-1}$$

where \bar{t} denotes the complex conjugate. In the first two cases, the corresponding real variety is $\mathbb{P}^1_{\mathbb{R}}$ since there are fixed points; while in the third, we obtain C since there are no fixed points. For the torus, we obtain $\lambda \mapsto \bar{\lambda}$ for the first case, so the real torus is \mathbb{R}^\times . However, in the second two cases $\lambda \mapsto \bar{\lambda}^{-1}$ and we obtain the circle group S^1 with $|\lambda| = 1$.

When the torus T is split, we call X a *split* toric variety. Most of the literature on toric varieties is about the split case since this is the only case when $k = \mathbb{C}$. In [Vos82], V. Voskresenskiĭ studies toric varieties for general tori T , but assumes that the open orbit has a rational point; we will call these *neutral* toric varieties.

Recall that a k -form of a k -variety X is a k -variety X' such that

$$X \times_{\text{Spec}(k)} \text{Spec}(K) \simeq X' \times_{\text{Spec}(k)} \text{Spec}(K)$$

for some field extension K/k . For example, $\mathbb{P}^1_{\mathbb{R}}$ and the conic C both become isomorphic to $\mathbb{P}^1_{\mathbb{C}}$ after base extension to \mathbb{C} . All toric varieties are k -forms of a split toric variety. The main goal of this paper is to study the set of isomorphism classes of forms of toric varieties.

Note that there are several different natural notions of isomorphism that one can use. We consider three different categories of toric varieties (made precise in Section 3), each of which has a different notion of isomorphism, and thus, of k -form.

When the toric variety is *neutral* we can identify the torus with its open orbit and consider the category \mathcal{W} with *toric morphisms* which restrict to group homomorphisms of the tori; this is the notion of isomorphism studied in [Vos82]. The category \mathcal{N} consists of toric varieties with torus-equivariant morphisms; this category is implicit in, for example, [VK84], [MP97], and [ELFST14]. Finally, the category \mathcal{R} consists of toric varieties with arbitrary morphisms which completely ignore the toric structure. To see how the different categories differ, note that the two different torus actions on $\mathbb{P}^1_{\mathbb{R}}$ discussed above give rise to two distinct isomorphism classes in \mathcal{N} , but they are not distinct in \mathcal{R} since the underlying varieties are the same.

Our approach is a direct generalization of standard techniques for studying forms of projective space. We review this now. Recall that a *Severi–Brauer variety*

is a k -form of projective space. The set of isomorphism classes of k -forms of a (suitably nice) algebraic object is in bijection with the étale cohomology set $H^1(k, \text{Aut}(X))$, which is simply Galois cohomology since k is a field. Thus the set of isomorphism classes of Severi–Brauer varieties is in bijection with the set $H^1(k, \text{PGL}_n)$.

Recall that the projective space $X = \mathbb{P}^{n-1}$ has a homogeneous coordinate ring whose spectrum is a vector space $V \simeq k^n$. The space X is a quotient of an open subset of the vector space V by the group \mathbb{G}_m . This construction gives rise to an exact sequence

$$1 \rightarrow \mathbb{G}_m \rightarrow \text{GL}_n \rightarrow \text{PGL}_n \rightarrow 1 \tag{1.1}$$

where GL_n acts on V and PGL_n is the automorphism group of \mathbb{P}^{n-1} .

From the exact sequence (1.1), the long exact sequence in non-abelian Galois cohomology produces a well-known injection

$$H^1(k, \text{PGL}_n) \hookrightarrow H^2(k, \mathbb{G}_m) \tag{1.2}$$

where $H^1(k, \text{PGL}_n)$ is in bijection with the set of isomorphism classes of Severi–Brauer varieties of dimension $n - 1$ and the group $H^2(k, \mathbb{G}_m)$ is the Brauer group $\text{Br}(k)$.

For a complete split toric variety X , the *Cox ring* of X , introduced in [Cox95], generalizes the usual homogeneous coordinate ring for projective space. Again, the spectrum of the Cox ring is a vector space V that has an open subset from which X can be reconstructed as a quotient by a diagonalizable group S . Here S is the Cartier dual of the class group of X , which generalizes \mathbb{G}_m from the case where X is \mathbb{P}^{n-1} .

There exists a linear algebraic group $\widetilde{\text{Aut}}(X)$ acting on the Cox ring, generalizing GL_n from the case of projective space, which has a more convenient description than the automorphism group scheme $\text{Aut}(X)$. Under certain assumptions on X , the standard sequence (1.1) is generalized by the bottom row of the following commutative diagram with exact rows

$$\begin{array}{ccccccc}
 1 & \longrightarrow & S & \longrightarrow & \widetilde{T} \times W & \longrightarrow & T \times W \longrightarrow 1 \\
 & & \parallel & & \downarrow & & \downarrow \\
 1 & \longrightarrow & S & \longrightarrow & \widetilde{\text{Aut}}(X) & \longrightarrow & \text{Aut}(X) \longrightarrow 1
 \end{array} \tag{1.3}$$

where T is a maximal torus of $\text{Aut}(X)$, the group \widetilde{T} is a maximal torus of $\widetilde{\text{Aut}}(X)$, and W is the group of toric automorphisms of X . The diagram (1.3) is essentially due to D. Cox; it follows from Theorem 4.5 below.

Galois cohomology can be viewed as a functor $H^1(k, -)$ which behaves well in exact sequences. In Theorem 5.1, we show how the sets of isomorphism classes for the three categories, as well as their relationships to each other and to their subsets of neutral forms, can be readily obtained from the Galois cohomology sets associated to (1.3). Applying $H^1(k, -)$ to the rightmost square we may interpret

each set as the k -forms of X (up to isomorphism) in an appropriate category:

$$\begin{array}{ccc}
 \begin{array}{|c|} \hline \text{Forms in } \mathcal{W} \\ \hline \text{neutral } X \\ \text{specified } T \\ \hline \end{array} & \begin{array}{c} \leftarrow \text{---} \\ \hookrightarrow \\ \leftarrow \text{---} \\ \hookrightarrow \end{array} & \begin{array}{|c|} \hline \text{Forms in } \mathcal{N} \\ \hline \text{arbitrary } X \\ \text{specified } T \\ \hline \end{array} \\
 \downarrow \quad \uparrow \text{---} & & \downarrow \\
 \begin{array}{|c|} \hline - \\ \hline \text{neutral } X \\ \text{unspecified } T \\ \hline \end{array} & \begin{array}{c} \leftarrow \text{---} \\ \hookrightarrow \\ \leftarrow \text{---} \\ \hookrightarrow \end{array} & \begin{array}{|c|} \hline \text{Forms in } \mathcal{R} \\ \hline \text{arbitrary } X \\ \text{unspecified } T \\ \hline \end{array}
 \end{array} \tag{1.4}$$

Here the vertical maps are surjections, the horizontal maps are injections and the dashed arrows are canonical sections or retracts. We see that the isomorphism classes are naturally partitioned into *neutralization classes* each of which contains exactly one neutral toric variety.

Note that $\text{Aut}(X)$ has a natural induced action on $\text{Cl}(X)$ and thus on its Cartier dual S . If J is the image of the action of $\text{Aut}(X)$ on $\text{Cl}(X)$, then the group J is finite but non-trivial in general. Thus, unlike the sequence (1.1) for projective space, S is not necessarily central in (1.3) and one should not expect a map $H^1(k, \text{Aut}(X)) \rightarrow H^2(k, S)$ for a general toric variety. Nevertheless, using the theory of non-abelian H^2 from [Spr66] and [Gir71], we prove in Theorem 7.4 that there is an injection

$$H^1(k, \text{Aut}(X)) \hookrightarrow H^2(k, S \rightarrow J) \tag{1.5}$$

where $H^2(k, S \rightarrow J)$ is a structured set defined below which is a natural analog of the Brauer group. One may view the map (1.5) as a refinement of the *elementary obstruction* from [CTS87].

Finally, given a split smooth projective toric variety X , we construct a canonical k -algebra B in a similar vein as a construction of A. Merkurjev and I. Panin in [MP97]. To each k -form of X we associate a k -form of B . This construction is a common generalization of the usual association of a Severi–Brauer variety to a central simple algebra and a construction of M. Blunk [Blu10] for del Pezzo surfaces of degree 6. We investigate when the isomorphism class of the k -forms of X can be recovered from the isomorphism classes of B ; it turns out to be closely related to retract rationality of the k -forms of X .

The paper is structured as follows. In Sections 2, 3, and 4, we fix notation, state definitions, and review basic facts that will be needed later in the paper. In Section 5, we use Galois cohomology and the structure theory of split toric varieties to classify their forms. In Section 6, we extend the structure theory of the split toric variety to the general case. In Section 7, we define the set $H^2(k, S \rightarrow J)$ and prove the injection (1.5). In Sections 8, 9, and 10, we use the preceding theory to investigate when Blunk’s construction can be generalized to other toric varieties.

Acknowledgements. The author would like to thank B. Antieau, M. Borovoi, M. Brown, D. Cox, A. Merkurjev, Z. Reichstein, A. Ruoizzi, and anonymous referees for helpful comments and discussions.

2. Preliminaries

Let k be a field. We will denote by k_s a separable closure of k . We denote by Γ_k the absolute Galois group of the field k , which is a profinite group.

A *variety* X is a geometrically integral separated scheme of finite type over a field k . A *group scheme* G will always be a group scheme of finite type over a field k . An *algebraic group* G is a smooth group scheme of finite type over a field k .

Given a field extension K/k we denote by

$$X_K := X \times_{\text{Spec}(k)} \text{Spec}(K)$$

the pullback, which is a variety defined over K . For the separable closure, we use the shorthand $\bar{X} := X_{k_s}$. A *k-form* of X is a variety X' defined over k such that $X'_K \simeq X_K$ for some field extension K/k .

We assume that reader is familiar with Galois cohomology (see, e.g., [Ser02]). For an algebraic group G , $H^i(k, G)$ will denote the i th Galois cohomology set $H^i(\Gamma_k, G(k_s))$. This is an abelian group when G is abelian, a group when $i = 0$, and a pointed set when G is non-abelian and $i = 1$.

Of fundamental importance to this paper is the well-known functorial bijection of pointed sets

$$H^1(k, \text{Aut}(X)) \simeq \left\{ \begin{array}{l} \text{isomorphism classes} \\ \text{of } k\text{-forms of } X \end{array} \right\}$$

which holds when X is quasiprojective and $\text{Aut}(X)$ is an algebraic group.

For an algebraic group G , an element $\gamma \in H^1(k, G)$ can be represented by a *cocycle* c or by a G -torsor T (also called a principal homogeneous space). If X is an algebraic variety (or algebraic group, or k -algebra) with a G -action, then we can construct the twisted variety ${}_cX$ or ${}^T X$ as in §5.3 of [Ser02].

2.1. Algebras

We refer the reader to §1, §18, and §23 of [KMRT98] for many of the results that follow. We assume throughout that all algebras are associative and unital. Given an algebra A , we denote its opposite algebra by A^{op} .

An *étale k-algebra* E is a direct product

$$E = F_1 \times \cdots \times F_r$$

where F_1, \dots, F_r are separable field extensions of k . An étale algebra E is *split* if every field F_i in the decomposition is isomorphic to k . The *degree* of an étale algebra E is its dimension as a vector space over k .

A *central simple algebra over k* is a k -algebra A such that there exists a field K for which $A_K \simeq M_n(K)$ where $M_n(K)$ is the algebra of $n \times n$ -matrices over K . The algebra A is *split* if $A \simeq M_n(k)$ over the original field.

A *separable algebra* A is a finite-dimensional k -algebra which is a finite product

$$A = A_1 \times \cdots \times A_r$$

where each A_i is a central simple algebra over a finite separable field extension F_i of k . The algebra A is *neutral* if every algebra A_i is split as a central simple

algebra over F_i . The algebra A is *split* if $F_1 = \cdots = F_r = k$ and every central simple algebra A_i is split over k . Note that the number of simple algebras r may increase after a base field extension.

Every separable algebra has a *split form* A_{split} which is the unique split k -form of A . Every separable algebra A has a *neutralization* A_{neut} where we replace each A_i in the product with its split form as an F_i -algebra. We define a *neutralization class* of a separable k -algebra A to be the set of k -forms of A which have the same neutralization. The center $Z(A)$ of a separable k -algebra is an étale k -algebra. The center $Z(A)$ is invariant within neutralization classes.

2.2. Automorphisms

Given a k -algebra A , let $\text{Aut}(A)$ denote the group scheme of automorphisms of A .

If E is an étale k -algebra of degree n , then $\text{Aut}(E)$ is a form of the symmetric group S_n ; thus, $\text{Aut}(E)$ is finite étale over k .

Given a finite-dimensional k -algebra A , we denote by $\text{GL}_1(A)$ the algebraic group representing the functor

$$R \rightarrow (A_R)^\times$$

on commutative k -algebras R (see §20 of [KMRT98]).

Let A be a separable k -algebra with center $Z(A)$. By §23 of [KMRT98], the connected component of $\text{Aut}(A)$ is given by

$$\text{Aut}(A)^\circ \simeq \text{GL}_1(A) / \text{GL}_1(Z(A))$$

and $\pi_0(\text{Aut}(A))$, the quotient by the connected component, is a subgroup of $\text{Aut}(Z(A))$.

Given an algebraic subgroup I of $\pi_0(\text{Aut}(A))$, the *I-restricted automorphism group of A*, denoted $\text{Aut}_I(A)$, is the preimage of I in $\text{Aut}(A)$. Note, in particular, when I is trivial, $\text{Aut}_1(A) = \text{Aut}(A)^\circ$.

2.3. Groups of multiplicative type

Most of the material here can be found in, e.g., [Vos98].

Let $\mathbb{G}_m = \text{GL}_1(k)$. A group scheme S of finite type is *diagonalizable* if S is a closed subgroup of $(\mathbb{G}_m)^n$ for some positive integer n . A *group of multiplicative type* is a group scheme S such that \overline{S} is diagonalizable. A group of multiplicative type is *split* if it is diagonalizable. An algebraic group S is a *torus* if $\overline{S} \simeq (\mathbb{G}_m)^n$ for some non-negative integer n .

Let Γ be a profinite group. A Γ -*module* L is a finitely generated abelian group L with a continuous action of Γ where L is endowed with the discrete topology. A Γ -*lattice* is a torsion-free Γ -module.

There is an exact anti-equivalence between the category of groups of multiplicative type and the category of Γ_k -modules which we will call “duality.” Given a group S of multiplicative type, the *character group*, \widehat{S} , is the corresponding Γ_k -module. Conversely, given a Γ_k -module L , the corresponding group of multiplicative type will be denoted $D(L)$. Under this equivalence, tori correspond to Γ -lattices.

We remark that if the torsion subgroup of the underlying abelian group of a Γ_k -module L has order relatively prime to the characteristic, then $D(L)$ is an algebraic group (in other words, it is smooth).

The image of the map $\Gamma_k \rightarrow \text{Aut}(\widehat{S})$ is a finite group which we call the *decomposition group of S* . One can also define the *cocharacter group of S* as the Γ_k -lattice $\text{Hom}(\widehat{S}, \mathbb{Z})$. One can recover the original group scheme S from its cocharacter lattice if and only if S is a torus.

2.4. Weil restrictions and Galois cohomology

Let E be an étale k -algebra and let \mathcal{F} be a functor from E -algebras to sets. We define the *Weil restriction* $R_{E/k} \mathcal{F}$ of \mathcal{F} as the functor from commutative k -algebras to sets given by

$$R_{E/k} \mathcal{F}(R) = \mathcal{F}(R \otimes_k E)$$

for each k -algebra R . The Weil restriction of an algebraic group (resp. variety) is also an algebraic group (resp. variety).

Lemma 2.1. *Let E be an étale k -algebra and G be an algebraic group over E . There is a natural isomorphism $H^i(k, R_{E/k} G) \simeq H^i(E, G)$ of groups (or pointed sets).*

Proof. See Lemma 29.6 of [KMRT98]. □

Proposition 2.2 (Hilbert 90). *For any separable k -algebra A , the cohomology set $H^1(k, \text{GL}_1(A))$ is trivial.*

Proof. See Theorem 29.2 of [KMRT98]. □

Proposition 2.3. *For any separable k -algebra A and for any cocycle c representing an element in $H^1(k, \text{Aut}(A))$, we have*

$${}_c\text{GL}_1(A) = \text{GL}_1({}_cA)$$

where ${}_cA$ (resp. ${}_c\text{GL}_1(A)$) denotes the form of A (resp. $\text{GL}_1(A)$) twisted by the cocycle c .

Proof. The embedding $\text{GL}_1(A)(k_s) \rightarrow A_{k_s}$ is $\text{Aut}(A_{k_s})(k_s)$ -equivariant. □

2.5. Permutation lattices and quasi-split tori

A Γ -lattice L is *permutation* if L has a basis which is permuted by Γ . A torus S is *quasi-split* if \widehat{S} is permutation.

There is an antiequivalence of categories between the category of étale algebras and the category of Γ_k -sets where Γ_k -orbits correspond to the subfields F_i of E .

Given an étale k -algebra E , the group $\text{GL}_1(E)$ is a torus. Indeed,

$$T = \text{GL}_1(E) \simeq R_{E/k} \mathbb{G}_{m,E}$$

is a Weil restriction. The character lattice \widehat{T} is a permutation Γ_k -lattice with a basis indexed by the Γ_k -set corresponding to E . The torus T is always a quasi-split torus and any quasi-split torus can arise in this way. Note, however, that non-isomorphic étale algebras may give rise to isomorphic tori since the choice of basis is not canonical.

3. Preliminaries on general toric varieties

There are several reasonable notions of a “toric variety” over a general field. Here we fix the definitions for the remainder of the paper.

Definition 3.1. Let T be a torus. A *toric T -variety* X is a normal k -variety with a faithful T -action and a dense open T -orbit X_0 . A toric T -variety X is *neutral* if there exists a T -equivariant isomorphism $T \rightarrow X_0$. A toric T -variety X is *split* if T is a split torus.

Over an algebraically closed field, the notions of neutral and split are vacuous. It is often desirable to keep track of a specific isomorphism $T \rightarrow X_0$ when X is neutral, but this is not part of our definition. Note that there are no non-trivial torsors under a split torus, so a split toric T -variety is always neutral.

Definition 3.2. We say X is a *toric variety* if there exists a torus T with an action on X giving X the structure of a toric T -variety. We say X is *neutral* if one can choose T such that X is neutral as a toric T -variety. We say X is *split* if one can choose T such that X is split as a toric T -variety.

The difference between Definitions 3.1 and 3.2 is whether a specific torus is fixed a priori or not. Note that a toric variety X is neutral for any choice of torus if it is neutral for any one choice. However, beware that a split toric variety may also be a toric T -variety for a different torus T that is *not* split.

We introduce 3 different categories of toric varieties to emphasize the different kind of morphisms one might consider in light of the above considerations. In Section 5, these categories will be used as natural settings for the machinery of descent.

Definition 3.3. The category \mathcal{R} :

- (1) objects are toric varieties,
- (2) morphisms are morphisms of varieties.

Definition 3.4. The category \mathcal{N} :

- (1) objects are pairs (T, X) where T is a torus and X is a toric T -variety,
- (2) morphisms from (T, X) to (T', X') are pairs (g, f) where $g : T \rightarrow T'$ is a group homomorphism and $f : X \rightarrow X'$ is a morphism of varieties which is T -equivariant via g .

The automorphisms in the category \mathcal{N} amount to automorphisms of the subvariety X_0 which extend to all of X .

Definition 3.5. The category \mathcal{W} :

- (1) objects are triples (T, X, ι) where T is a torus, X is a neutral toric T -variety, and $\iota : T \hookrightarrow X$ is an isomorphism with the dense open orbit,
- (2) morphisms from (T, X, ι) to (T', X', ι') are pairs (g, f) where $g : T \rightarrow T'$ is a group homomorphism and $f : X \rightarrow X'$ is a morphism of varieties such

that the diagram

$$\begin{array}{ccc}
 X & \xrightarrow{f} & X' \\
 \uparrow \iota & & \uparrow \iota' \\
 T & \xrightarrow{g} & T'
 \end{array}$$

commutes.

The morphisms of \mathcal{W} are called *toric morphisms* in the literature. Observe that multiplication by a non-trivial element of the torus is an automorphism in \mathcal{N} , but not in \mathcal{W} .

We mention two useful results regarding general toric varieties.

Proposition 3.6. *If T is a torus then there exists a smooth projective T -variety X .*

Proof. See [CTHS05]. \square

Proposition 3.7. *A smooth projective toric variety X is neutral if and only if X has a rational k -point.*

Proof. See Proposition 4 of [VK84]. \square

4. Structure of split toric varieties

Throughout this section, X is a proper split toric T -variety with a specified embedding $T \hookrightarrow X$.

Hypothesis 4.1. Throughout the paper we will make the following additional technical assumptions:

- the automorphisms functor $\text{Aut}(X)$ is a linear algebraic group, and
- the order of the torsion subgroup of the class group $\text{Cl}(X)$ is relatively prime to the characteristic of k .

We shall see that both these assumptions hold when X is smooth or X has characteristic 0.

Here we outline the well-known structure theory of a split toric variety to fix notation. We assume the reader is familiar with standard references on toric varieties (for example, [Ful93] or [CLS11]). Many references only consider the base field \mathbb{C} , but much of the theory goes through unchanged in the split case.

Let $M = \widehat{T}$ be the character lattice of T , and let N be the cocharacter lattice of T . Of course, since T is split, both lattices have trivial Γ_k -action.

Split toric T -varieties X are in bijective correspondence with *fans* in their cocharacter lattices N . From the data of a fan Σ , one can determine whether X is smooth, projective, or proper over k . We denote by $\Sigma(k)$ the set of cones of dimension k in Σ ; in particular, $\Sigma(1)$ is the set of rays. We denote by \widetilde{M} the free abelian group with basis indexed by $\Sigma(1)$. The dual lattice \widetilde{N} is canonically isomorphic to \widetilde{M} by using this basis. The lattice \widetilde{M} is isomorphic to the group of T -invariant Weil divisors of X .

For a split toric variety, we have canonical isomorphisms $\text{Cl}(X) = \text{Cl}(\overline{X})$ and $\text{Pic}(X) = \text{Pic}(\overline{X})$ (this follows from their descriptions via torus-invariant divisors

as in, e.g., Theorem 4.1.3 of [CLS11]). The divisor class group, $\text{Cl}(\overline{X})$, has a natural structure as a (trivial) Γ_k -module. Let S be the group of multiplicative type $D(\text{Cl}(\overline{X}))$. Note that by our assumption above, S is a linear algebraic group.

There is an exact sequence

$$1 \rightarrow M \rightarrow \widetilde{M} \rightarrow \text{Cl}(\overline{X}) \rightarrow 1 \tag{4.1}$$

when $k_s[\overline{X}]^\times \simeq k_s$. When X is smooth, $\text{Cl}(\overline{X})$ is canonically isomorphic to $\text{Pic}(\overline{X})$ and is torsion free.

4.1. Cox rings

We now review the theory of Cox rings (see [Cox95]). For non-split toric varieties this can be subtle (see Section 6 and [DP14]), but here we assume X is split and proper.

The *Cox ring* $\text{Cox}(X)$ of X is the polynomial ring

$$\text{Cox}(X) := k[x_1, \dots, x_r]$$

where the monomials x_1, \dots, x_r correspond to the rays ρ_1, \dots, ρ_r in $\Sigma(1)$. Let \widetilde{T} be the torus dual to \widetilde{M} . Note that $\text{Cox}(X)$ has a canonical embedding into $k[\widetilde{T}]$ where monomials can be identified with elements of \widetilde{M} . The ring $\text{Cox}(X)$ has a natural $\text{Cl}(\overline{X})$ grading via the morphism $\widetilde{M} \rightarrow \text{Cl}(\overline{X})$ from (4.1).

For a Weil divisor D , we denote the graded component of $\text{Cox}(X)$ corresponding to $[D] \in \text{Cl}(\overline{X})$ by $\text{Cox}(X)_{[D]}$ or $\text{Cox}(X)_D$. Denoting $\mathcal{O}_X(D)$ as the reflexive sheaf associated to D , there are isomorphisms

$$\text{Cox}(X)_D \simeq H^0(X, \mathcal{O}_X(D)) = \{f \in k(X)^\times : \text{div}(f) + D \geq 0\} \cup \{0\}$$

for every Weil divisor D .

We define the *irrelevant ideal* B of $\text{Cox}(X)$ as the monomial ideal generated by products $x_{i_1} \cdots x_{i_r}$ corresponding to subsets of rays $\{\rho_{i_1}, \dots, \rho_{i_r}\}$ which are the complement of a cone in Σ . Note that $V = \text{Spec}(\text{Cox}(X))$ is an affine variety with a natural vector space structure. The ideal B cuts out a closed subvariety $Z \subset V$ whose complement \widetilde{X} we call the *characteristic space* of X .

Since S is the Cartier dual of $\text{Cl}(\overline{X})$, the $\text{Cl}(\overline{X})$ -grading on $\text{Cox}(X)$ corresponds to generically-free actions of S on V and on \widetilde{X} (obtained via restrictions of the action of the torus \widetilde{T}). We may recover X as the categorical quotient of \widetilde{X} by S ; over \mathbb{C} this is Theorem 2.1 of [Cox95], but the proof works in general with minor modifications. In the case that X is smooth, S is a torus and, thus, the action of S on \widetilde{X} is free and the quotient

$$\psi : \widetilde{X} \rightarrow X$$

is an S -torsor (in fact, a *universal torsor* in the sense of [CTS87]).

4.2. Automorphisms

The automorphism group scheme of a smooth proper split toric variety was determined in [Dem70]. Our exposition is heavily inspired by [Cox95] where the automorphism group of a split proper simplicial toric variety over \mathbb{C} is determined indirectly via the Cox ring (see also [Cox14]). The simplicial hypothesis is removed in [Büh96]. The automorphism group of a projective split toric variety is determined in [BG99] over an arbitrary field.

Recall that we assume that $\text{Aut}(X)$ is a linear algebraic group. The author knows no counterexample to this assumption for an arbitrary proper toric variety and it holds many cases of interest:

Lemma 4.2. *Let X be a proper split toric variety and suppose that either*

- (1) X is smooth, or
- (2) k has characteristic 0 and X is projective.

Then the functor $\text{Aut}(X)$ of automorphisms of X is a linear algebraic group.

Proof. Note that since X is proper, $\text{Aut}(X)$ is a group scheme of locally finite type over k (see Theorem 3.7 of [MO67]). When X is smooth, $\text{Aut}(X)$ is a linear algebraic group by Proposition 11 of [Dem70]. In characteristic 0, the neutral component $\text{Aut}(X)^\circ$ must be smooth. From [BG99], we have a description of $\text{Aut}(X)(K)$ for any field K when X is projective. From this description we see that $\text{Aut}(X)$ has finitely many connected components and thus $\text{Aut}(X)$ is a linear algebraic group. \square

Let W be the group of toric automorphisms of X (the subgroup of $\text{GL}(N) \simeq \text{GL}_n(\mathbb{Z})$ which takes cones to cones). Note that W has induced actions on \widetilde{T} and T .

Let V_λ be the weight subspace of V corresponding to λ in $\text{Cl}(\overline{X})$ and let n_λ be its dimension. Let Λ be the subset of $\text{Cl}(\overline{X})$ corresponding to the non-trivial weight subspaces V_λ of V .

Definition 4.3. The *Cox endomorphism algebra of X* is the split separable k -algebra

$$A := \prod_{\lambda \in \Lambda} \text{End}(V_\lambda).$$

We define $W^\circ = W \cap \text{GL}_1(A)$ and note that

$$W^\circ \simeq \prod_{\lambda \in \Lambda} S_{n_\lambda}$$

where each S_{n_λ} is the symmetric group on n_λ letters. The group W° is isomorphic to the Weyl group of $\text{GL}_1(A)$.

Definition 4.4. The *group of class group automorphisms of X* , denoted J , is the image of the map $\text{Aut}(X) \rightarrow \text{Aut}(S)$ (recall that we obtain an induced action of $\text{Aut}(X)$ on S since it is the dual of the class group $\text{Cl}(X)$). The group J is a finite constant group which will be of fundamental importance for the remainder of the paper.

Let $\widetilde{\text{Aut}}(X)$ be the normalizer of S in the automorphism group functor of \widetilde{X} . We will see that, in fact, $\widetilde{\text{Aut}}(X)$ is a linear algebraic group.

Theorem 4.5. *Let X be a projective split toric variety with $\text{Aut}(X)$ smooth. Diagram (1.3):*

$$\begin{array}{ccccccc}
 1 & \longrightarrow & S & \longrightarrow & \widetilde{T} \rtimes W & \longrightarrow & T \rtimes W \longrightarrow 1 \\
 & & \parallel & & \downarrow & & \downarrow \\
 1 & \longrightarrow & S & \longrightarrow & \widetilde{\text{Aut}}(X) & \longrightarrow & \text{Aut}(X) \longrightarrow 1
 \end{array}$$

commutes and has exact rows. Moreover, $T \rtimes W$ (resp. $\widetilde{T} \rtimes W$) is the normalizer of a maximal torus in $\text{Aut}(X)$ (resp. $\widetilde{\text{Aut}}(X)$). There is a unipotent subgroup U such that

$$\widetilde{\text{Aut}}(X) \simeq U \rtimes \text{GL}_1(A) \rtimes J,$$

and an isomorphism

$$W \simeq W^\circ \rtimes J$$

where the splitting is unique up to conjugacy.

Proof. When $k = \mathbb{C}$ and X is simplicial, the commutative diagram (1.3) is essentially the main theorem of §4 of [Cox95]. From [BG99], one can obtain the commutativity and exactness of the diagram on the level of k -points where k is an arbitrary field. The group $\widetilde{\text{Aut}}(X)$ can be explicitly constructed as a closed subgroup of a general linear group acting on a subspace of $k[V]$ (the exposition in [Cox14] is especially clear). Using this description, one checks that its Lie algebra has the appropriate dimension and thus it is a linear algebraic group.

Since $\text{Aut}(X)$ is also a linear algebraic group, we may establish the remaining statements by first assuming k is algebraically closed and then showing that all of the structural maps are actually defined over the original field k . We need to establish the splittings of $\widetilde{\text{Aut}}(X)$ and W . By the references above, the group $\widetilde{\text{Aut}}(X) \cap \text{GL}(V)$ is isomorphic to the product $\text{GL}_1(A)W$. Moreover, there is a surjective homomorphism $\widetilde{\text{Aut}}(X) \rightarrow \text{GL}_1(A)W$ with kernel a unipotent subgroup U .

Note that V is the vector space dual to the subspace spanned by the generators x_1, \dots, x_r of $\text{Cox}(X)$ and thus is spanned by the dual basis x_1^*, \dots, x_r^* which may be identified with rays of Σ . The group W permutes these basis vectors. The group W° consists of all permutations of x_1^*, \dots, x_r^* which preserve the decomposition into weight subspaces V_λ .

Since $\text{GL}_1(A)$ is connected and J is a constant finite group, we obtain an exact sequence

$$1 \rightarrow W^\circ \rightarrow W \rightarrow J \rightarrow 1$$

which we want to show is split. The group J acts faithfully on the set Λ . Note that if W splits as $W^\circ \rtimes J$, then $\text{GL}_1(A)W$ splits as $\text{GL}_1(A) \rtimes J$.

Choose orderings for x_1^*, \dots, x_r^* within each subspace V_λ . Pick a set-theoretic section $s : J \rightarrow W$. For each $j \in J$, we have $s(j)(V_\lambda) = V_{j(\lambda)}$. By comparing the ordered bases of V_λ and $V_{j(\lambda)}$, the element j gives rise to an element $w_\lambda \in S_{n_\lambda}$. Taking the product of the w_λ 's for each $\lambda \in \Lambda$, we obtain an element $w_j \in W^\circ$ such that $w_j^{-1} \circ s(j) : V_\lambda \rightarrow V_{j(\lambda)}$ is an isomorphism of vector spaces with ordered

bases. Since such isomorphisms are unique, the set-theoretic section $\tilde{s} : J \rightarrow W$ given by $\tilde{s}(j) = w_j^{-1} \circ s(j)$ is a group homomorphism as desired.

Note that the section constructed only depends on the choice of orderings for the bases within each V_λ . These are permuted by W° , so the section is unique up to conjugacy.

We have established the theorem when k is algebraically closed. However, $U, \tilde{T}, T, \text{GL}_1(A), W, W^\circ, J$ and the splittings can be all be defined over a general field k . All the remaining statements follow since they are true over an algebraic closure. \square

Remark 4.6. As remarked above, it is unclear if $\text{Aut}(X)$ is ever not a smooth group scheme. Regardless, there is a group scheme homomorphism $\widetilde{\text{Aut}}(X)/S \rightarrow \text{Aut}(X)$ defined via the universal property of the categorical quotient $\tilde{X} \rightarrow X$, which is an isomorphism on the level of K -points for any field K . The theorem still holds without the smoothness assumption if we replace $\text{Aut}(X)$ with $\widetilde{\text{Aut}}(X)/S$.

Remark 4.7. Note that in [Cox95], it is erroneously stated that *all* graded endomorphisms of $\text{Cox}(X)$ form a (not necessarily separable) algebra B . This would lead to a description $\widetilde{\text{Aut}}(X) \simeq \text{GL}_1(B) \times J$ above. This error was first addressed in [Cox14]. In fact, the graded endomorphisms form an algebra if and only if the unipotent radical U is trivial. Indeed, if U is trivial then $B = A$. However, if U is non-trivial then there exists an element $g \in B$ which acts as $g(x_i^*) = x_i^*$ for all $i \neq j$ and $g(x_j^*) = x_j^* + M$ where j is an integer in $1, \dots, r$, and M is some monomial in x_1^*, \dots, x_r^* of degree ≥ 2 . If B were an algebra then g would commute with scalar multiplication, but this fails since $\text{deg}(M) \neq 1$.

Remark 4.8. Let V be a vector space of dimension n . If X is $\mathbb{P}(V)$, the set of 1-dimensional subspaces of V , then the Cox endomorphism algebra A is simply $\text{End}(V)$. As a special case of the Severi–Brauer construction, we may recover X as the variety of right ideals of A (see [Sal99] or [KMRT98]).

5. Twists of the split toric variety

Throughout this section, we assume that X is a split projective toric T -variety with a fixed torus embedding $T \hookrightarrow X$ as in the previous section. The projectivity assumption is to ensure that descent is effective. This assumption can be often be weakened (see [Hur11] for a complete description of when descent is effective for toric T -varieties.) Moreover, we also assume Hypothesis 4.1 holds.

Note that the Galois cohomology set $H^1(k, G)$ is functorial in the group G . Thus, we may apply the functor $H^1(k, -)$ to the commutative diagram (1.3). The following theorem investigates the result of this operation:

Theorem 5.1. *The commutative square*

$$\begin{array}{ccc}
 H^1(k, \tilde{T} \rtimes W) & \longrightarrow & H^1(k, T \rtimes W) \\
 \downarrow & & \downarrow \\
 H^1(k, \widetilde{\text{Aut}}(X)) & \longrightarrow & H^1(k, \text{Aut}(X))
 \end{array} \tag{5.1}$$

obtained from the commutative diagram (1.3) is canonically isomorphic to the square

$$\begin{array}{ccc}
 H^1(k, W) & \xrightarrow{\quad} & H^1(k, T \rtimes W) \\
 \downarrow & \nearrow & \downarrow \\
 H^1(k, J) & \xrightarrow{\quad} & H^1(k, \text{Aut}(X))
 \end{array} \tag{5.2}$$

where the downward maps are surjective and the rightward maps are injective. Moreover, the rightward maps have canonical retracts and the left downward map has a canonical section.

Before proving the theorem, we make some remarks. Recalling the categories from Section 3 we find:

$$\text{Aut}_{\mathcal{R}}(X) = \text{Aut}(X), \quad \text{Aut}_{\mathcal{N}}(X) = T \rtimes W, \quad \text{Aut}_{\mathcal{W}}(X) = W.$$

Thus, three of the coefficient groups appearing in (5.2) are simply the automorphism groups of toric varieties within these categories. The Galois cohomology sets represent the forms of X within each category. As \mathcal{W} contains only neutral toric varieties, we recover the interpretation from (1.4).

From this diagram, we recover the well-known fact that there is a unique isomorphism class of a split toric variety (resp. split toric T -variety) among all the possible k -forms. For a given toric variety X we call the unique split variety *the associated split toric variety* and denote it by X_{split} .

From the canonical retracts, we see that every toric variety X has an *associated neutral toric variety*, or *neutralization*, which we denote by X_{neut} . We call the set of forms which have a common neutralization a *neutralization class* and remark that these partition the isomorphism classes of forms of X .

The canonical section provides every neutral toric variety with a canonical isomorphism class of torus among the many which may act on the variety. When the toric variety is split, the canonical section recovers the split torus.

Remark 5.2. An investigation of the top row of Theorem 4.5 using Galois cohomology was also carried out in [ELFST14].

Remark 5.3. In [VK84] and [MP97], the neutralization is defined when T is fixed and is called the “associated toric T -model.” The theorem above shows that one can define the neutralization independently of the particular torus action chosen.

We now prove Theorem 5.1. We begin with some technical lemmas:

Lemma 5.4. *Let A and C be algebraic groups where C acts on A . The map $H^1(k, A \rtimes C) \rightarrow H^1(k, C)$ is surjective and the map $H^1(k, C) \rightarrow H^1(k, A \rtimes C)$ is injective.*

Proof. Apply the functor $H^1(k, -)$ to the composition $C \rightarrow A \rtimes C \rightarrow C$. \square

Lemma 5.5. *Let A and C be algebraic groups where C acts on A . Let ξ be a set of cocycle representatives for the set $H^1(k, C)$. There is a canonical bijection*

$$H^1(k, A \rtimes C) \simeq \coprod_{c \in \xi} H^1(k, {}_cA) / H^0(k, {}_cC),$$

functorial in A , where each component of the disjoint union is a fibre of the canonical map to $H^1(k, C)$. In particular, $H^1(k, {}_cA)$ is trivial for every $c \in \xi$ if and only if the map $H^1(k, A \rtimes C) \rightarrow H^1(k, C)$ is a bijection.

Proof. The last statement follows from the previous statements since the indexing set ξ is bijective with $H^1(k, C)$.

By Lemma 5.4, there is a canonical choice of preimage in $H^1(k, A \rtimes C)$ for elements in $H^1(k, C)$. By Corollary 2 of §5.5 of [Ser02] we obtain the desired bijection. More precisely, for $c \in \xi$, we obtain a map $H^1(k, {}_cA) \rightarrow H^1(k, {}_c(A \rtimes C))$ whose image is $H^1(k, {}_cA)/H^0(k, {}_cC)$; then we use the bijection $\tau_c : H^1(k, A \rtimes C) \rightarrow H^1(k, {}_c(A \rtimes C))$ as in §5.4 of [Ser02]. (Note that this bijection is canonical up to the choice of cocycle representatives ξ .)

To prove functoriality, consider another algebraic group B with an action of C and a C -equivariant homomorphism $A \rightarrow B$. The morphism $f \rtimes C : A \rtimes C \rightarrow B \rtimes C$ gives rise to a commutative diagram

$$\begin{CD} H^1(k, A \rtimes C) @>>> H^1(k, B \rtimes C) \\ @VVV @VVV \\ H^1(k, C) @= H^1(k, C) \end{CD}$$

Thus, the map takes fibres to fibres. Note that twisting by $c \in \xi$ is a functorial operation, so we can twist f by c to obtain ${}_cf : {}_cA \rightarrow {}_cB$. The map

$$\coprod_{c \in \xi} H^1(k, {}_cA)/H^0(k, {}_cC) \rightarrow \coprod_{c \in \xi} H^1(k, {}_cB)/H^0(k, {}_cC)$$

is obtained by taking a disjoint union of quotients by $H^0(k, {}_cC)$. This process preserves compositions and the identity. \square

Lemma 5.6. *Let A be a separable k -algebra with center $Z(A)$. Let S be a closed subgroup of $\text{GL}_1(Z(A))$ and suppose J is an algebraic subgroup of $\text{Aut}(A)$ which stabilizes S . Then the induced morphism of Galois cohomology sets*

$$H^1(k, \text{GL}_1(A) \rtimes J) \rightarrow H^1(k, \text{GL}_1(A)/S \rtimes J)$$

is canonically isomorphic to the injection

$$H^1(k, J) \hookrightarrow H^1(k, \text{GL}_1(A)/S \rtimes J)$$

and has a canonical retract.

Proof. Since J acts on the algebra A by automorphisms, for any cocycle $c \in Z^1(k, J)$ we have ${}_c\text{GL}_1(A) = \text{GL}_1({}_cA)$ by Proposition 2.3. Thus, by Hilbert 90, we see that $H^1(k, {}_c\text{GL}(A))$ is trivial for any cocycle $c \in Z^1(k, J)$.

By Lemma 5.5, we conclude that

$$H^1(k, \text{GL}_1(A) \rtimes J) \simeq H^1(k, J).$$

We obtain injectivity and the canonical retract from Lemma 5.4. \square

We now observe that, from the perspective of Galois cohomology, the unipotent radical is irrelevant:

Proposition 5.7. *There are canonical bijections*

$$H^1(k, \widetilde{\text{Aut}}(X)) \simeq H^1(k, \text{GL}_1(A) \rtimes J)$$

and

$$H^1(k, \text{Aut}(X)) \simeq H^1(k, (\text{GL}_1(A)/S) \rtimes J).$$

Proof. Since X is split, X can be defined over the prime field k_0 of k ; in other words, $X = Y_k$ for a split toric variety Y over k_0 . All prime fields are perfect, thus the unipotent radicals of $\widetilde{\text{Aut}}(Y)$ and $\text{Aut}(Y)$ are k_0 -split. As a consequence, the unipotent radicals of $\widetilde{\text{Aut}}(X)$ and $\text{Aut}(X)$ are k -split and, thus, we may apply Lemma 7.20 of [GMB13]. \square

Proof of Theorem 5.1. First we consider the top row. Recall that \widetilde{T} is a split torus and thus is $\text{GL}_1(E)$ for a split étale k -algebra E . The group W permutes a basis for V and thus acts on E by algebra automorphisms. From Lemma 5.6, we obtain injectivity with a canonical retract.

By Proposition 5.7, we may assume the bottom row in (5.1) is

$$H^1(k, \text{GL}_1(A) \rtimes J) \rightarrow H^1(k, (\text{GL}_1(A)/S) \rtimes J).$$

Again, Lemma 5.6 provides injectivity with a canonical retract.

Now, we show that the vertical maps are surjections. For any cocycle $c \in Z^1(k, J)$ the map

$${}_c(\widetilde{T} \rtimes W^\circ) \rightarrow {}_c\text{GL}_1(A)$$

is the inclusion of the normalizer of a maximal torus into a connected algebraic group and thus the induced map

$$H^1(k, {}_c(\widetilde{T} \rtimes W^\circ)) \rightarrow H^1(k, {}_c\text{GL}_1(A))$$

is surjective by Corollary 5.3 of [CGR08] (Lemma III.4.3.6 of [Ser02] when k is perfect). Via Lemma 5.5, we conclude that the left vertical map in (5.1) is surjective. Similarly, we conclude the right vertical map is surjective.

Since J and W are finite constant groups, the elements of $H^1(k, J)$ and $H^1(k, W)$ are simply homomorphisms from Γ_k to J or W up to conjugacy. By Theorem 4.5, there exists a section of the map $W \rightarrow J$ which is unique up to conjugacy. Thus, the induced map $H^1(k, J) \rightarrow H^1(k, W)$ is independent of the choice of section $J \rightarrow W$ and, thus, is canonical. \square

Example 5.8 ($\mathbb{P}^1_{\mathbb{R}} \times \mathbb{P}^1_{\mathbb{R}}$). Consider $X = \mathbb{P}^1 \times \mathbb{P}^1$ over the real numbers \mathbb{R} . There is precisely one other \mathbb{R} -form C of \mathbb{P}^1 over \mathbb{R} corresponding to the subvariety

$$x^2 + y^2 + z^2 = 0$$

cut out of \mathbb{P}^2 . There are only two isomorphism classes of 1-dimensional tori over \mathbb{R} which we denote by \mathbb{R}^\times and S^1 .

Here $W \simeq D_8$, and $T \simeq \mathbb{G}_m^2$. The group $J \simeq C_2$ can be thought of as the group which interchanges the two fibrations $X \rightarrow \mathbb{P}^1$. In this case, one can compute all the cohomology groups in Theorem 5.1 explicitly. We summarize the conclusions of this computation.

There are two neutralization classes. The first contains:

- (a) $\mathbb{P}^1 \times \mathbb{P}^1$ with possible tori $\mathbb{R}^\times \times \mathbb{R}^\times$, $\mathbb{R}^\times \times S^1$ and $S^1 \times S^1$,
- (b) $\mathbb{P}^1 \times C$ with possible tori $\mathbb{R}^\times \times S^1$ and $S^1 \times S^1$,
- (c) $C \times C$ with torus $S^1 \times S^1$.

And the second class contains:

- (d) $\mathbb{R}_{\mathbb{C}/\mathbb{R}}\mathbb{P}_{\mathbb{C}}^1$ with torus $\mathbb{R}_{\mathbb{C}/\mathbb{R}}\mathbb{C}^\times$.

We point out the role of J in these computations using Lemma 5.4. First, the action of $H^0(k, {}_cJ)$ is necessary to establish that $\mathbb{P}^1 \times C$ and $C \times \mathbb{P}^1$ are isomorphic. Second, the necessity of non-trivial cocycles $c \in Z^1(k, J)$ is witnessed by the existence of $\mathbb{R}_{\mathbb{C}/\mathbb{R}}\mathbb{P}_{\mathbb{C}}^1$.

6. Properties of twisted forms

Throughout this section, X is a projective toric variety which is *not necessarily* split. However, we assume the split form X_{split} of X satisfies Hypothesis 4.1 (which then also holds for X).

Clearly, one may define the automorphism group $\text{Aut}(X)$ and its unipotent radical U . As before, we define the algebraic group S as the dual of the Γ_k -module $\text{Cl}(\overline{X})$. We define the *group of class group automorphisms* J as the image of the morphism $\text{Aut}(X) \rightarrow \text{Aut}(S)$ as before. The group J is a finite algebraic group, but may not be constant. If we do specify a torus T , then we define W as the Weyl group of T in $\text{Aut}(X)$; in other words $W := N_{\text{Aut}(X)}(T)/T$. All of these objects agree with the corresponding objects of the split form over a separable closure.

The theory of Cox rings is more complex for non-split toric varieties. Depending on one’s definitions, they may not exist or they may not be polynomial rings (see [DP14]). Despite this, one may still define a Cox endomorphism algebra for a general toric variety.

Proposition 6.1. *There exists an algebra A and a morphism $\text{GL}_1(A) \rightarrow \text{Aut}(X)$ which coincides with the split case over the separable closure. The isomorphism class of the center $Z(A)$ of A is determined only by the neutralization class of X .*

We call the algebra A in the proposition above the *Cox endomorphism algebra of X* , generalizing Definition 4.3.

Proof. We use the fact that X is a k -form of X_{split} .

Recall that the algebra A_{split} associated to X_{split} comes with a map $\text{GL}_1(A_{\text{split}}) \rightarrow \text{Aut}(X_{\text{split}})$. From Theorem 4.5, we have a morphism

$$\text{Aut}(X_{\text{split}}) \rightarrow \text{Aut}(A_{\text{split}})$$

since S_{split} is a subgroup of the group $\text{GL}_1(Z(A_{\text{split}}))$. Thus the morphism $\text{GL}_1(A_{\text{split}}) \rightarrow \text{Aut}(X_{\text{split}})$ is $\text{Aut}(X_{\text{split}})$ -equivariant, and we obtain the desired algebra A and map by twisting.

By composition, there is also a map $\text{Aut}(X_{\text{split}}) \rightarrow \text{Aut}(Z(A_{\text{split}}))$ which factors through J_{split} since $\text{GL}_1(A_{\text{split}})$ centralizes $Z(A_{\text{split}})$. Thus, the induced morphism

$$H^1(k, \text{Aut}(X_{\text{split}})) \rightarrow H^1(k, \text{Aut}(Z(A_{\text{split}})))$$

then factors through $H^1(k, J_{\text{split}})$. Thus, the isomorphism class of $Z(A)$ depends only on the neutralization class of X . \square

Example 6.2 (Severi–Brauer varieties). Given a central simple algebra A , one can construct the Severi–Brauer variety X associated to A . Proposition 6.1 simply reverses this process. In the case of Severi–Brauer varieties, the isomorphism class of the toric variety X is completely determined by the isomorphism class of its Cox algebra A .

Example 6.3 (del Pezzo of degree 6). Let X be the split del Pezzo surface of degree 6. As a toric variety, its effective T -invariant divisors are spanned by 6 elements which correspond to the blow-ups of three points on \mathbb{P}^2 and the strict transforms of the lines between them.

There is a common convention for the coordinates of the rays of this surface. Using this convention, the map $\tilde{N} \rightarrow N$ is given by

$$\begin{pmatrix} 1 & 0 & -1 & -1 & 0 & 1 \\ 0 & 1 & -1 & 0 & -1 & 1 \end{pmatrix}.$$

By duality we obtain a map $\tilde{M} \rightarrow \text{Pic}(X)$ given by

$$\begin{pmatrix} 0 & 0 & 0 & 1 & 1 & 1 \\ 1 & 0 & 0 & 0 & -1 & -1 \\ 0 & 1 & 0 & -1 & 0 & -1 \\ 0 & 0 & 1 & -1 & -1 & 0 \end{pmatrix}$$

where we denote the ordered basis of $\text{Cl}(\tilde{X}) \simeq \text{Pic}(X)$ by H, E_1, E_2, E_3 .

Note that as all 6 rays have different images in $\text{Cl}(\tilde{X})$, all the spaces V_λ are 1-dimensional. Thus, the Cox endomorphism algebra A is an étale algebra and thus equal to its center. By Proposition 6.1, the algebras associated to the k -forms of X are determined only by the neutralization class.

However, not all forms of X are neutral. Thus, the Cox endomorphism algebras A do not suffice to distinguish k -forms of X in this case. However, in [Blu10], a different pair of separable algebras is associated to each k -form of X which do distinguish all isomorphism classes of X . We investigate this phenomenon in Section 10 below; in particular, see Example 10.2.

Remark 6.4. (Maximal étale algebras) For Severi–Brauer varieties, there is a bijective correspondence between maximal étale subalgebras of A and maximal tori of $\text{Aut}(X)$. This holds for toric varieties when the unipotent radical U is trivial. Conditions for when the hypothesis $U = 0$ holds are investigated in [Nil06].

In Cox’s original paper, the Cox ring was only defined for split toric varieties (over \mathbb{C}). Cox rings can be extended to more general varieties as a ring structure on the direct sum of the vector spaces $H^0(X, \mathcal{O}_X(D))$ for all $D \in \text{Cl}(X)$. This definition makes sense when the base field is not algebraically closed, but we lose the connection with universal torsors. An alternate definition was proposed in [DP14], which is better suited to applications over non-closed fields. However, like universal torsors, these Cox rings do not always exist, nor are they necessarily unique. The following is in the spirit of the latter definition.

Recall that in the split case, the *characteristic space* of a toric variety X is the S -invariant open subset \tilde{X} of V along with the categorical quotient map $\psi : \tilde{X} \rightarrow X$.

We define a *characteristic space* of a (not necessarily split) toric variety X as a subvariety \widetilde{X} of an affine space V along with a map $\psi : \widetilde{X} \rightarrow X$ which coincides with the characteristic space of the split toric variety over the separable closure. When \widetilde{X} does exist, we define $\widetilde{\text{Aut}}(X)$ to be normalizer of S in $\text{Aut}(\widetilde{X})$ as before.

Proposition 6.5. *There exists a characteristic space $\psi : \widetilde{X} \rightarrow X$ if and only if X is neutral. When \widetilde{X} exists, we have a decomposition*

$$\widetilde{\text{Aut}}(X) \simeq U \times \text{GL}_1(A) \rtimes J$$

as in the split case.

Proof. Assume a characteristic space exists. Then we have a dominant rational map $V \rightarrow X$ where V is an affine space. Thus X has a Zariski-dense set of k -points. We conclude that X is neutral since any open T -orbit for any T -action must contain a k -point.

Now, assume that X is neutral. By Theorem 5.1, we may choose a cocycle c in $Z^1(K, J_{\text{split}})$ such that $X \simeq_c(X_{\text{split}})$. Since J_{split} maps to $\text{Aut}(X_{\text{split}})$ and leaves S_{split} stable, we may define an affine space $V := {}_c(V_{\text{split}})$ containing a characteristic space $\widetilde{X} := {}_c(\widetilde{X}_{\text{split}})$ both with S -actions. We define the map $\psi : \widetilde{X} \rightarrow X$ as the twist ${}_c(\psi_{\text{split}})$.

The splitting $J_{\text{split}} \xrightarrow{\sim} \widetilde{\text{Aut}}(X_{\text{split}})$ is J_{split} -equivariant and thus, when we twist by c , the morphism $\widetilde{\text{Aut}}(X) \rightarrow J$ splits and we have the desired decomposition. \square

Remark 6.6. Note that the map $\psi : \widetilde{X} \rightarrow X$ is not unique as an S -scheme over X — even up to isomorphism. Indeed, when X is smooth, ψ is a universal torsor of X ; their isomorphism classes are in bijection with $H^1(k, S)$ (see §2 of [CTS87]).

Remark 6.7. (Canonical torus) When X is neutral, we can define \widetilde{T} with an action on \widetilde{X} as in the split case. In this case, there is a canonical choice for the isomorphism class of the torus \widetilde{T} (and thus T). This follows from the fact that the morphism $H^1(k, W_{\text{split}}) \rightarrow H^1(k, J_{\text{split}})$ has a canonical section. Specifically, as A is a neutral separable k -algebra, it is a product of matrix algebras

$$A = M_{n_1}(F_1) \times \cdots \times M_{n_r}(F_r)$$

where F_1, \dots, F_r are separable field extensions of k and n_1, \dots, n_r are positive integers. There is a maximal étale subalgebra of A of the form

$$E = (F_1)^{n_1} \times \cdots \times (F_r)^{n_r}.$$

The canonical torus \widetilde{T} is then $\text{GL}_1(E)$.

Remark 6.8. (Restricted automorphism groups) Let X be a neutral toric variety. Given a finite algebraic subgroup I of J , the I -restricted automorphism group of X , denoted $\text{Aut}_I(X)$, is the preimage of I in $\text{Aut}(X)$. Note that, in particular, $\text{Aut}_1(X) = \text{Aut}(X)^\circ$. The Galois cohomology set $H^1(k, \text{Aut}_1(X))$ can be interpreted as the set of isomorphism classes of toric varieties Y isomorphic to X along

with a fixed isomorphism $\text{Pic}(Y) \simeq \text{Pic}(X)$. Note that if we drop the explicit isomorphism, we obtain the set

$$H^1(k, \text{Aut}_1(X))/H^1(k, J)$$

which is the set of isomorphism classes of toric varieties Y with a neutralization isomorphic to X .

Example 6.9 (Products of projective spaces). Let $X_{\text{split}} = (\mathbb{P}^{q-1})^n$ be a product of projective spaces given two integers $q, n > 1$. Here $\text{Pic}(X_{\text{split}}) \simeq \mathbb{Z}^n$, $S \simeq \mathbb{G}_m^n$ and $J \simeq S_n$. In this case, (1.3) is as follows:

$$\begin{array}{ccccccc} 1 & \longrightarrow & \mathbb{G}_m^n & \longrightarrow & (\mathbb{G}_m^q)^n \rtimes (S_q \wr S_n) & \longrightarrow & (\mathbb{G}_m^q/\mathbb{G}_m)^n \rtimes (S_q \wr S_n) \longrightarrow 1 \\ & & \parallel & & \downarrow & & \downarrow \\ 1 & \longrightarrow & \mathbb{G}_m^n & \longrightarrow & \text{GL}_m \wr S_n & \longrightarrow & \text{PGL}_m \wr S_n \longrightarrow 1 \end{array}$$

where $G \wr S_n$ is the wreath product $G^n \rtimes S_n$ for a group G .

Here the algebra A_{split} is simply $M_q(k)^n$ and thus the automorphism group $\text{Aut}(X_{\text{split}}) \simeq \text{PGL}_q \wr S_n$ of the variety is isomorphic to $\text{Aut}(A_{\text{split}})$. Thus, forms X of X_{split} correspond to forms A of A_{split} . The neutral forms X correspond to neutral forms A , which, in turn, correspond to étale k -algebras of degree n .

More geometrically, k -forms X are products of Weil restrictions

$$\text{R}_{F_1/k} X_1 \times \cdots \times \text{R}_{F_r/k} X_r$$

where each X_i is a Severi–Brauer variety of dimension $q-1$ over the field F_i . When X is neutral, each X_i is a projective space.

7. Non-abelian H^2

In this section, we extend the long exact sequences in Galois cohomology from Theorem 4.5 to H^2 . Unlike the situation for Severi–Brauer varieties, the original sequences do not correspond to central extensions, so we cannot use the ordinary abelian cohomology group $H^2(k, S)$.

In [Spr66] and §IV.4.2 of [Gir71], the ordinary sequence from Galois cohomology is extended to a non-abelian version of H^2 (see also more recent work in [Bor93], [FSS98] and [Flo04]). Our applications are less ambitious, so we have the luxury of a simpler exposition. Rather than consider the abelian group $H^2(k, S)$, we consider $H^2(k, S \rightarrow J)$ which has the structure of a set with a distinguished subset. Here the notation for the “coefficients” is meant to suggest a *crossed module* as in, for example, [Bre10].

Let S be a smooth split k -group of multiplicative type and let J be a finite subgroup of $\text{Aut}(S)$. Note that J is a constant group.

We have a natural action of J on each cocycle $c \in Z^1(k, J)$ via $j(c)_\sigma := jc_\sigma j^{-1}$ for all $\sigma \in \Gamma_k$. Two cycles are cohomologous if and only if they are in the same orbit under this action (essentially by definition).

Given a cocycle $c \in Z^1(k, J)$ and an element $j \in J$ we have an isomorphism

$$j_* : H^2(k, {}_cS) \rightarrow H^2(k, {}_{j(c)}S)$$

which is defined on cocycles $s \in Z^2(k, {}_cS)$ via

$$j_*(s)_{\sigma, \tau} := j(s_{\sigma, \tau}) \text{ for all } \sigma, \tau \in \Gamma_k.$$

One checks that the image cocycle sits in $Z^2(k, {}_{j(c)}S)$ as expected and that cohomology classes are preserved.

We now define the main object of study in this section:

Definition 7.1. We define the following set:

$$H^2(k, S \rightarrow J) := \left(\coprod_{c \in Z^1(k, J)} H^2(k, {}_cS) \right) / J.$$

We define the set of *neutral elements* as the subset of $H^2(k, S \rightarrow J)$ containing the trivial elements in each component $H^2(k, {}_cS)$. This endows the set $H^2(k, S \rightarrow J)$ with the structure of a set with a distinguished subset. The image of the trivial element from $H^2(k, S)$ will be referred to as the *trivial element* of $H^2(k, S \rightarrow J)$.

Now we define the connecting homomorphism. Suppose \tilde{G} and G are algebraic groups sitting in an exact sequence

$$1 \rightarrow S \rightarrow \tilde{G} \rightarrow G \rightarrow 1 \tag{7.1}$$

such that the conjugation action of G on S induces a surjection $\pi : G \rightarrow J$.

Let a be a cocycle in $Z^1(k, G)$. Let $\tilde{a} : \Gamma_k \rightarrow \tilde{G}(k_s)$ be a continuous function lifting a (which always exists since Γ_k is profinite and $G(k_s), \tilde{G}(k_s)$ have the discrete topology). Define a function $\Delta\tilde{a} : \Gamma_k \times \Gamma_k \rightarrow \tilde{G}(k_s)$ via

$$(\Delta\tilde{a})_{\sigma, \tau} := \tilde{a}_\sigma(\sigma\tilde{a}_\tau)(\tilde{a}_{\sigma\tau})^{-1}$$

for all $\sigma, \tau \in \Gamma_k$. We will see that $\Delta\tilde{a}$ is a cocycle in $Z^2(k, \pi(a)S)$, and that this gives rise to a well-defined map

$$\delta : H^1(k, G) \rightarrow H^2(k, S \rightarrow J)$$

which we call the *connecting map*.

Lemma 7.2. *The connecting map*

$$\delta : H^1(k, G) \rightarrow H^2(k, S \rightarrow J)$$

is well defined and canonically isomorphic as sets with distinguished subsets to the map

$$\delta^1 : H^1(k, \tilde{G}, S) \rightarrow H^2(k, S \text{ rel } \tilde{G})$$

from 1.20 of [Spr66].

Proof. We will prove the isomorphism with Springer’s map; the fact that our construction is well defined then follows for free.

Let a be a cocycle in $Z^1(k, G)$ and let $\tilde{a} : \Gamma_k \rightarrow \tilde{G}(k_s)$ be a continuous function lifting a . In Springer’s notation, the set $Z^1(k, \tilde{G}, S)$ is simply the set of continuous functions $b : \Gamma_k \rightarrow \tilde{G}(k_s)$ lifting cocycles in $Z^1(k, G)$. We may simply assume $\tilde{a} = b$. To each cocycle b there is an associated 2-cocycle (f, g) for $H^2(k, S \text{ rel } \tilde{G})$ where

$$f_\sigma(s) = b_\sigma(\sigma s)(b_\sigma)^{-1},$$

$$g_{\sigma, \tau} = b_\sigma(\sigma b_\tau)(b_{\sigma\tau})^{-1}$$

for $s \in S(k_s)$ and $\sigma, \tau \in \Gamma_k$. Comparing this to Definition 7.1, we will see that f corresponds to the choice of cocycle c and g corresponds to a cocycle in $Z^2(k, {}_cS)$.

This cocycle (f, g) sits naturally inside $Z^2(k, S, \lambda_a)$ where λ_a is the Γ_k -kernel induced by the automorphisms f_σ of S . Since S is abelian, the kernels λ_a are honest continuous maps $\Gamma_k \rightarrow \text{Aut}(S(k_s))$ (where $\text{Aut}(S(k_s))$ has the discrete topology). Thus, the kernel λ_a coincides with the cocycle $c = \pi_*(a)$ in $Z^1(k, J)$. The cocycle g is equal to Δb as functions and they are then 2-cocycles in $Z^2(k, {}_cS)$.

The normalizer N of S in \tilde{G} is \tilde{G} itself. Given an element $n \in N$ we have the image $j = \pi(n) \in J$. The induced action of n on the set of Γ_k -kernels of S coincides with the induced action of j on $Z^1(k, J)$ defined above, and the induced action of n on the pairs (f, g) coincides with the action of j on 2-cocycles in the sets $Z^2(k, {}_cS)$. Thus the constructions agree. \square

Now, suppose that $\tilde{G} = R \rtimes J$ where R is an algebraic group containing S as a J -stable central closed subgroup. We have a J -equivariant exact sequence

$$1 \rightarrow S \rightarrow R \rightarrow R/S \rightarrow 1$$

corresponding to a central extension.

Lemma 7.3. *Let ξ be a set of cocycle representatives for $H^1(k, J)$. The connecting map*

$$\delta : H^1(k, G) \rightarrow H^2(k, S \rightarrow J)$$

is canonically isomorphic to the map

$$\coprod_{c \in \xi} \left(\frac{H^1(k, {}_c(R/S))}{H^0(k, {}_cJ)} \right) \rightarrow \coprod_{c \in \xi} \left(\frac{H^2(k, {}_cS)}{H^0(k, {}_cJ)} \right)$$

induced from the usual connecting maps $H^1(k, {}_c(R/S)) \rightarrow H^2(k, {}_cS)$.

Proof. We have the decomposition of $H^1(k, G)$ from Lemma 5.5.

The orbits of J in $Z^1(k, J)$ are precisely the cohomology classes. Thus, the images of the maps $H^2(k, {}_cS) \rightarrow H^2(k, S \rightarrow J)$ over all cocycles $c \in \xi$ are mutually disjoint and jointly surjective. For a particular cocycle $c \in \xi$, an element $j \in J$ fixes c if and only if j is in $H^0(k, {}_cJ)$. Thus, the image of the map

$H^2(k, {}_cS) \rightarrow H^2(k, S \rightarrow J)$ is simply the quotient by $H^0(k, {}_cJ)$. This establishes the decomposition of $H^2(k, S \rightarrow J)$.

It remains to prove that the connecting maps agree. Fix a cocycle $c \in \xi$. Let u be a cocycle in $Z^1(k, {}_c(R/S))$ and let $a_\sigma = u_\sigma c_\sigma$ be a corresponding cocycle in $Z^1(k, G)$. The cocycles u and a map to the same element of $H^1(k, G)$ in the decomposition from Lemma 5.5. Select a lift $v : \Gamma_k \rightarrow {}_cR(k_s)$ of u and note that $b_\sigma = v_\sigma c_\sigma : \Gamma_k \rightarrow \tilde{G}(k_s)$ is a lift of a_σ .

The cocycle Δv in $Z^2(k, {}_cS)$ is constructed via

$$(\Delta v)_{\sigma,\tau} := (v_\sigma)(\sigma'v_\tau)(v_{\sigma\tau})^{-1}$$

using the twisted action $\sigma's = (c_\sigma)(\sigma s)(c_\sigma)^{-1}$ for $s \in S(k_s)$ and $\sigma \in \Gamma_k$. The computation

$$\begin{aligned} (\Delta v)_{\sigma,\tau} &= (v_\sigma)[(c_\sigma)(\sigma v_\tau)(c_\sigma)^{-1}](v_{\sigma\tau})^{-1} \\ &= (v_\sigma)(c_\sigma)(\sigma v_\tau)[(\sigma c_\tau)(c_{\sigma\tau})^{-1}](v_{\sigma\tau})^{-1} \\ &= (b_\sigma)(\sigma b_\tau)(b_{\sigma\tau})^{-1} = (\Delta b)_{\sigma,\tau} \end{aligned}$$

shows that this has the desired image in $H^2(k, S \rightarrow J)$. \square

The above shows that there is a surjective map $H^2(k, S \rightarrow J) \rightarrow H^1(k, J)$ with a unique neutral element in each fiber. Given an element α in the set $H^2(k, S \rightarrow J)$, we define its *neutral class* as the fiber of this map and we define the *neutralization* of α as the unique neutral element therein.

Let A, B, C respectively be sets with distinguished subsets A', B', C' respectively. We say that the composition $g \circ f$ of functions $f : A \rightarrow B, g : B \rightarrow C$ is *exact* if $\text{im}(f) = g^{-1}(C')$. We can now state the main theorem of this section.

Theorem 7.4. *Let X be a split projective toric variety satisfying Hypothesis 4.1. Applying Galois cohomology to the sequence (1.3), we extend (5.2) to the commutative diagram*

$$\begin{CD} H^1(k, W) @<<< H^1(k, T \times W) @>>> H^2(k, S \rightarrow J) \\ @VVV @VVV @| \\ H^1(k, J) @<<< H^1(k, \text{Aut}(X)) @<<< H^2(k, S \rightarrow J) \end{CD} \tag{7.2}$$

where the rows are exact sequences of sets with a distinguished subset and the bottom right horizontal map is injective.

Note that exact sequences for sets with distinguished subsets behave differently from exact sequences of abelian groups. Thus it is perfectly reasonable to have an exact sequence consisting of two injections. In the bottom row of the above diagram, the exactness and injectivity imply that the distinguished subset of $H^2(k, S \rightarrow J)$ is in bijection with $H^1(k, J)$.

Proof. The top row is the case where $R = \widetilde{T} \rtimes W^\circ$; the bottom, where $R = \mathrm{GL}_1(A)$. The exactness follows from Proposition 1.28 of [Spr66]. This can also be checked directly using Lemmas 5.5 and 7.3. The only statement that remains to be proved is the injectivity of the map $H^1(k, \mathrm{Aut}(X)) \rightarrow H^2(k, S \rightarrow J)$.

Let c be a cocycle representative of an element in $H^1(k, J)$. Consider the central extension

$$1 \rightarrow {}_cS \rightarrow {}_c\mathrm{GL}_1(A) \rightarrow {}_c(\mathrm{GL}_1(A)/S) \rightarrow 1.$$

Since this is a central extension, we obtain an exact sequence of pointed sets

$$H^1(k, {}_c\mathrm{GL}_1(A)) \rightarrow H^1(k, {}_c(\mathrm{GL}_1(A)/S)) \rightarrow H^2(k, {}_cS).$$

Note that ${}_c(\mathrm{GL}_1(A)/S)$ acts on ${}_c\mathrm{GL}_1(A) \simeq \mathrm{GL}_1({}_cA)$ by algebra automorphisms of ${}_cA$. Thus, for any cocycle d in $Z^1(k, {}_c(\mathrm{GL}_1(A)/S))$, we see that

$$H^1(k, {}_d({}_c\mathrm{GL}_1(A))) \simeq H^1(k, \mathrm{GL}_1({}_d({}_cA)))$$

is trivial by Hilbert 90. By the Corollary in §5.7 of [Ser02], we conclude the map $H^1(k, {}_c(\mathrm{GL}_1(A)/S)) \rightarrow H^2(k, {}_cS)$ is injective. Injectivity is preserved after taking quotients by $H^0(k, {}_cJ)$, and the result follows from Lemma 7.3. \square

Remark 7.5. Suppose X and X' are toric varieties with fixed torus actions. Theorem 7.4 allows one to determine whether the underlying varieties are isomorphic by considering the map $H^1(k, T \rtimes W) \rightarrow H^2(k, S \rightarrow J)$ instead of $H^1(k, T \rtimes W) \rightarrow H^1(k, \mathrm{Aut}(X))$. The former involves only the cohomology of abelian and finite constant groups, which may be easier to use in some applications. This generalizes the common trick where one distinguishes Severi–Brauer varieties (or central simple algebras) by their Brauer classes rather than by their classes in $H^1(k, \mathrm{PGL}_n)$.

Remark 7.6. (Torsors) Note that Lemma 7.3 allows us to easily describe the set $H^2(k, S \rightarrow J)$ using torsors instead of 1-cocycles. Indeed, if ξ is a collection of torsors representing elements of $H^1(k, J)$ then we have

$$H^2(k, S \rightarrow J) \simeq \coprod_{T \in \xi} \left(\frac{H^2(k, {}^T S)}{H^0(k, {}^T J)} \right).$$

Note that this agrees with Proposition 4.2.5(ii) of [Gir71].

Remark 7.7. (Restriction) One can define a restriction map

$$\mathrm{Res}_k^K : H^2(k, S \rightarrow J) \rightarrow H^2(K, S_K \rightarrow J_K)$$

for any field extension K/k . For each J -torsor T , we have a restriction map

$$H^2(k, {}^T S) \rightarrow H^2(K, {}^{T_K} S_K)$$

which is equivariant with respect to the homomorphism

$$H^0(k, {}^T J) \rightarrow H^0(K, {}^{T_K} J_K).$$

Noting that $T \mapsto T_K$ is the restriction morphism

$$H^1(k, J) \rightarrow H^1(K, J)$$

of the indexing sets, we obtain a restriction map via Remark 7.6. One checks that the restriction maps are compatible and we may think of $H^2(-, S \rightarrow J)$ as being a functor from the category of field extensions K/k to the category of sets with a distinguished subset.

Remark 7.8. (Period) The set $H^2(k, S \rightarrow J)$ is not a group in general. However, we may still consider powers of elements. Indeed, any element $\alpha \in H^2(k, S \rightarrow J)$ is represented by an element β in some group $H^2(k, {}_cS)$ for some cocycle $c \in Z^1(k, J)$. For any integer n , we define α^n as the image of β^n . Since J acts on each component of the disjoint union by group automorphisms, this is well defined. In particular, the notion of “period” still makes sense: the *period* of $\alpha \in H^2(k, S \rightarrow J)$ is the minimal positive integer n such that α^n is neutral.

Remark 7.9. (Index) There are at least two natural notions of the “index” of an element $\alpha \in H^2(k, S \rightarrow J)$. We define the *neutralizing index* (resp. *splitting index*) of α as the greatest common divisor of the degrees of all finite extensions L/k such that the restriction $\text{Res}_k^L(\alpha)$ is neutral (resp. trivial).

Example 7.10. Let k be the real numbers \mathbb{R} . Consider $X = \mathbb{P}^1 \times \mathbb{P}^1$ where $S = \mathbb{G}_m^2$ and $J = C_2$. By Lemma 7.3 we find that

$$H^2(\mathbb{R}, S \rightarrow J) \simeq (\text{Br}(\mathbb{R})^2/C_2) \sqcup \text{Br}(\mathbb{C})$$

and see that every Brauer class corresponds to a form of X from Example 5.8.

In contrast, consider $X = \mathbb{P}^1 \times \mathbb{P}^3$ where $S = \mathbb{G}_m^2$ and J is trivial. Here

$$H^2(\mathbb{R}, S \rightarrow J) \simeq \text{Br}(\mathbb{R})^2$$

and the k -forms are

$$\mathbb{P}^1 \times \mathbb{P}^3, C \times \mathbb{P}^3, \mathbb{P}^1 \times C', C \times C'$$

where C and C' , respectively, are the non-split forms of \mathbb{P}^1 and \mathbb{P}^3 , respectively.

Note that in both these examples S is the same. However, the finite group J is different in each case.

Example 7.11 (Permutation lattices). Let \widehat{P} be an S_n -lattice with a basis $\{p_1, \dots, p_n\}$ permuted by S_n . Let P be the dual torus $D(\widehat{P})$. We would like to understand the set $H^2(k, P \rightarrow S_n)$.

Recall that elements of $H^1(k, S_n)$ are in bijection with Γ -actions on $\{p_i\}$ and with étale k -algebras of degree n . Thus, for any cocycle $c \in Z^1(k, S_n)$, the twisted torus ${}_cP$ is simply a Weil restriction $\text{R}_{E/k} \mathbb{G}_{m,E}$ where E is the corresponding étale k -algebra.

Writing $E = F_1 \times \dots \times F_r$ as a decomposition of field extensions of k we have

$$\begin{aligned} H^2(k, {}_cP) &\simeq H^2(k, \text{R}_{E/k} \mathbb{G}_m) \simeq \prod_{i=1}^r H^2(k, \text{R}_{F_i/k} \mathbb{G}_m) \\ &\simeq \prod_{i=1}^r H^2(F_i, \mathbb{G}_m) \simeq \prod_{i=1}^r \text{Br}(F_i) \end{aligned}$$

for each cocycle $c \in Z^1(k, S_n)$. Note that the action of $H^0(k, {}_cS_n)$ permutes isomorphic field extensions in the decomposition of E .

By the decomposition in Lemma 7.3, we conclude that $H^2(k, P \rightarrow S_n)$ consists of all sets

$$\{(\alpha_1, F_1), \dots, (\alpha_r, F_r)\}$$

of pairs, where each F_i is a separable field extension of k and each α_i is an element of $\text{Br}(F_i)$, satisfying the condition $\sum_{i=1}^r [F_i : k] = n$.

Remark 7.12. (Elementary obstruction) The elements in $H^2(k, S \rightarrow J)$ are closely related to the “elementary obstructions” from [CTS87]. Indeed, given a k -form of split smooth projective toric variety, its class in $H^2(k, S \rightarrow J)$ is neutral if and only if the elementary obstruction is trivial. In Proposition 1.3 of [Sko09], A. Skorobogatov shows that the images of the elementary obstruction and the classes defined above actually coincide in the set $H^2(k, S \rightarrow \text{Aut}(S))$ where one must extend the definition above to consider the infinite group $\text{Aut}(S)$ (in this reference, the type of a universal torsor is only well-defined up to an isomorphism of the Picard group).

8. Comparison to ordinary Brauer groups

There is a natural bijection between the isomorphism classes of Severi–Brauer varieties and their associated central simple algebras. This is a stronger statement than merely identifying the classes of these algebras in $H^2(k, \mathbb{G}_m)$. In [Blu10], M. Blunk constructs a pair of separable algebras B and Q and exhibits a bijection with the k -forms of a del Pezzo surface of degree 6 and a subset of the k -forms of the algebra $B \times Q$.

Both of these constructions are well-behaved under extension of the base field. In both cases we have a toric variety X and a separable k -algebra B along with an injection

$$\left\{ \begin{array}{c} \text{isomorphism classes} \\ \text{of } k\text{-forms of } X \end{array} \right\} \hookrightarrow \left\{ \begin{array}{c} \text{isomorphism classes} \\ \text{of } k\text{-forms of } B \end{array} \right\} \tag{8.1}$$

which is natural in the field k . The remainder of the paper focuses on a partial answer to this question.

Recall that the automorphism group $\text{Aut}(B)$ of a separable algebra B is a linear algebraic group with connected component $\text{Aut}(B)^\circ$. Given a finite closed subgroup J of the quotient $\text{Aut}(B)/\text{Aut}(B)^\circ$, the J -restricted automorphism group of B , denoted $\text{Aut}_J(B)$, is the preimage of J in $\text{Aut}(B)$.

For a finite group J , a finitely generated J -module is *permutation* if it is free as a \mathbb{Z} -module with basis permuted by J ; a J -module is *invertible* if it is a direct summand of a permutation J -module. Note that, for an algebraic group G over a field k , one can view $H^1(-, G)$ as a functor from the category of field extensions K/k to the category of sets.

The following theorem is our main goal for the rest of the paper and will be proved in Section 10.

Theorem 8.1. *Let X be a smooth projective split toric variety over a field k . Let J be the image of the homomorphism $\text{Aut}(X) \rightarrow \text{Aut}(\text{Pic}(X))$. Suppose that $\text{Pic}(X)$ is an invertible J -lattice. There exists a canonical separable algebra B over k , and a natural transformation*

$$H^1(-, \text{Aut}(X)) \rightarrow H^1(-, \text{Aut}_J(B)), \tag{8.2}$$

which is injective for every field K/k .

Note that $\text{Pic}(X)$ is an invertible Γ -lattice for all smooth projective toric surfaces by [Vos67]. Thus, the conclusion of Theorem 8.1 holds for all smooth projective split toric surfaces.

When $\text{Aut}_J(B) = \text{Aut}(B)$, as is the case for the central simple algebras associated to Severi–Brauer varieties and Blunk’s algebra B , the set $H^1(k, \text{Aut}_J(B))$ can be interpreted directly as isomorphism classes of k -forms of B . When J is trivial, the set $H^1(k, \text{Aut}_1(B))$ is simply the set of isomorphism classes of k -forms of B which preserve a fixed labelling of each central simple algebra in the product. In the intermediate cases, $\text{Aut}_J(B)$ preserves this labelling up to certain symmetries. Thus, Theorem 8.1 can be interpreted as a variant of our desired injection (8.1) for a mildly weaker notion of isomorphism.

In [MP97], A. Merkurjev and I. Panin construct a “motivic category” which contains both toric varieties and separable algebras. In that paper, which was a starting point for Blunk’s work, they also associate forms of separable algebras to forms of toric varieties. In Theorem 9.1, we show that their method is, up to Brauer equivalence, essentially the same as our own.

8.1. Morphisms of non-abelian H^2

Throughout this section, X will be a split smooth projective toric variety. Recall that this implies that $\text{Cl}(\overline{X}) \simeq \text{Pic}(\overline{X})$ is a lattice and thus that S is a torus. Note that $\text{Pic}(\overline{X})$ has a trivial Γ_k -action, but a non-trivial J -action; it will be convenient to view $\widehat{S} = \text{Pic}(\overline{X})$ as a J -lattice.

In this section we make precise to what degree the information contained in $H^2(k, S \rightarrow J)$ can be captured by “ordinary Brauer groups.” To make this precise, we need to define a notion of morphisms between non-abelian H^2 sets. Unfortunately, defining “functoriality” for non-abelian H^2 is a rather delicate problem (see, e.g., [AN09]). However, in our restricted context there is a natural induced morphism which is well-behaved enough for our applications.

Definition 8.2. Let S and P be split groups of multiplicative type with actions of finite groups J and I respectively. Suppose (m, g) is a pair of group homomorphisms $m : S \rightarrow P$ and $g : J \rightarrow I$ such that m is J -equivariant where the J -action on P is given via g . Define the *induced map*

$$m_* : H^2(k, S \rightarrow J) \rightarrow H^2(k, P \rightarrow I)$$

via the decomposition from Lemma 7.3 and the induced maps

$$H^2(k, {}_cS) \rightarrow H^2(k, {}_{g_*(c)}P)$$

for every cocycle c representing a cohomology class in $H^1(k, J)$. One checks that m_* is compatible with restriction, and thus may be viewed as a natural transformation

$$m_* : H^2(-, S \rightarrow J) \rightarrow H^2(-, P \rightarrow I)$$

of functors from field extensions K/k to the category of sets with distinguished subsets.

A G -lattice M is *invertible* if M is a direct summand of a permutation G -module. Recall that a k -variety Y is *retract rational* if there exists an affine space

V , a dominant rational map $\psi : V \dashrightarrow X$ and a rational map $\eta : X \dashrightarrow V$ such that the composition $\psi \circ \eta$ is defined and equivalent to the identity as a rational map.

Theorem 8.3. *Let X be a smooth projective split toric variety. The following are equivalent:*

- (a) *There exists a morphism $\widehat{m} : \widehat{P} \rightarrow \widehat{S}$ of J -lattices, where \widehat{P} is permutation, such that the morphism of functors*

$$m_* : H^2(-, S \rightarrow J) \rightarrow H^2(-, P \rightarrow J)$$

is injective,

- (b) *\widehat{S} is an invertible J -lattice,*
- (c) *for every field extension K/k , every neutral K -form of X_K is retract rational.*

Remark 8.4. Note that retract rationality is a birational invariant; thus, the retract rationality of a toric variety can be determined by consideration of only the open orbit. Retract rational tori have been completely classified in small dimensions. See [Vos67], [Kun87], and [HY12]. In fact, all 2-dimensional tori are rational; thus all toric surfaces satisfy the equivalent conditions of Theorem 8.3.

We may interpret Theorem 8.3 as a statement about ordinary Brauer groups in the following way. Fix a choice of basis ω for \widehat{P} . Then twisting by a cocycle $c \in Z^1(k, J)$ gives a Γ_k -set ${}_c\omega$ as basis for ${}_c\widehat{P}$. Each Γ_k -set corresponds to an étale algebra ${}_cL$. Thus $H^2(k, {}_cP) \simeq H^2({}_cL, \mathbb{G}_m) \simeq \text{Br}({}_cL)$.

By the injection of Theorem 7.4, we have a composite map

$$H^1(k, \text{Aut}(X)) \rightarrow H^2(k, S \rightarrow J) \rightarrow H^2(k, P \rightarrow J)$$

which is injective if and only if the map m_* is injective. Given a fixed cocycle $c \in H^1(k, J)$, we have a morphism

$$H^1(k, \text{Aut}_1({}_cX)) \rightarrow \text{Br}({}_cL)$$

for each neutralization class (recall the definition of $\text{Aut}_1({}_cX)$ from Remark 6.8).

We recall some preliminaries on flasque and coflasque tori from, for example, [CTS77]. Let G be a finite group. A G -lattice M is *flasque* (resp. *coflasque*) if the Tate cohomology group $\widehat{H}^{-1}(H, M)$ (resp. $\widehat{H}^1(H, M)$) is trivial for every subgroup $H \subset G$. Every G -lattice M has a *coflasque resolution*, which is an exact sequence

$$1 \rightarrow A \rightarrow B \rightarrow M \rightarrow 1$$

where A is a coflasque G -lattice and B is a permutation G -lattice.

For any torus T defined over a field k , the action of Γ_k on the character lattice M factors through a finite group G . A torus T is defined to be *flasque*, *coflasque*, or *invertible* if the corresponding property is true of the G -module M .

Lemma 8.5. *Consider an exact sequence of J -lattices*

$$1 \rightarrow \widehat{Q} \xrightarrow{\widehat{m}} \widehat{S} \rightarrow 1$$

where \widehat{Q} is coflasque, \widehat{P} is permutation, and \widehat{S} is flasque. Let $Q, P,$ and S be the corresponding dual tori and let $m : S \rightarrow P$ the dual morphism obtained from \widehat{m} . Let $m_* : H^2(-, S \rightarrow J) \rightarrow H^2(-, P \rightarrow J)$ be the natural transformation induced from m as in Definition 8.2. Let $m'_* : H^2(-, S) \rightarrow H^2(-, P)$ be the usual induced map on abelian Galois cohomology obtained from m . Then the following are equivalent:

- (a) $m_* : H^2(-, S \rightarrow J) \rightarrow H^2(-, P \rightarrow J)$ is injective,
- (b) ${}_c(m'_*) : H^2(K, {}_cS) \rightarrow H^2(K, {}_cP)$ is injective for every field extension K/k and every cocycle $c \in Z^1(K, J)$,
- (c) $H^1(K, {}_cQ)$ is trivial for every field extension K/k and every cocycle $c \in Z^1(K, J)$,
- (d) \widehat{Q} is invertible,
- (e) \widehat{S} is invertible.

Proof. First, we prove the equivalence of (a) and (b). By the decomposition from Lemma 7.3, it suffices to check whether each constituent map

$$H^2(K, {}_cS)/H^0(K, {}_cJ) \rightarrow H^2(k, {}_cP)/H^0(K, {}_cJ)$$

is injective for each $c \in Z^1(K, J)$. If each ${}_c(m'_*)$ is injective, then so must be each constituent map. Conversely, since each ${}_c(m'_*)$ is a morphism of groups and $H^0(K, {}_cJ)$ acts by group automorphisms, if the preimage of the trivial element is trivial then ${}_c(m'_*)$ is injective.

The equivalence of (b) and (c) follow from the triviality of $H^1(K, {}_cP)$ for all cocycles $c \in Z^1(K, J)$ since \widehat{P} is a permutation J -lattice.

The implication (d) \implies (c) follows since there is a factorization $\widehat{Q} \rightarrow \widehat{M} \rightarrow \widehat{Q}$ of the identity morphism for some permutation lattice M . This means that the identity morphism on $H^1(K, {}_cQ)$ factors through the trivial group $H^1(K, {}_cM)$ for every cocycle c .

The implication (c) \implies (d) follows from Theorem 3.2 of [Mer10]. Indeed, we know that, in particular, the generic ${}_cQ$ -torsor is trivial. The theorem tells us that the class of the extension corresponding to a flasque resolution of \widehat{Q} is trivial. We conclude that \widehat{Q} is a direct summand of a permutation module as desired.

Finally, the equivalence of (d) and (e) follows from Lemma 6 of [CTS77]. Indeed, \widehat{S} is obtained via a flasque resolution of \widehat{Q} and, conversely, \widehat{Q} is obtained via a coflasque resolution of \widehat{S} . \square

Proposition 8.6 (Versality). *Suppose \widehat{Q} is coflasque. Let R be a torus with a J -action such that the dual \widehat{R} is a permutation J -lattice, and let $q : S \rightarrow R$ be a J -equivariant morphism of tori. Then*

$$q_* : H^2(k, S \rightarrow J) \rightarrow H^2(k, R \rightarrow J)$$

factors through

$$m_* : H^2(k, S \rightarrow J) \rightarrow H^2(k, P \rightarrow J).$$

Proof. Fix a cocycle $c \in Z^1(k, J)$ and note that it may be viewed as a morphism $c : \Gamma_k \rightarrow J$ since J is a constant group. The action of Γ_k on the character lattices of ${}_cS, {}_cR, {}_cP,$ and ${}_cQ$ then factor through J . We may conclude that ${}_cR$ is quasi-split and

$$1 \rightarrow {}_cS \rightarrow {}_cP \rightarrow {}_cQ \rightarrow 1$$

is a coflasque resolution of tori. Thus, the morphism $cq : {}_cS \rightarrow {}_cR$ factors through ${}_cm : {}_cS \rightarrow {}_cP$ by Lemma 4 of [CTS77]. We conclude that q_* factors through m_* via Definition 8.2. \square

We point out the following fact which is essentially due to Colliot-Thélène and Sansuc.

Proposition 8.7. *The J -lattice \widehat{S} is flasque.*

Proof. We may assume that k is of characteristic 0 since the statement only depends on data associated to the fan. Furthermore, by possibly taking a base extension of k , we may assume that there exists a surjection $c : \Gamma_k \rightarrow J$. The twisted torus ${}_cS$ has decomposition group J . Proposition 6 of [CTS77], shows that the Picard group $\text{Pic}(\overline{X})$ of a smooth compactification of a torus is flasque as a Γ_k -module. We conclude that ${}_cS$ is flasque as a Γ_k -module and thus $\text{Pic}(\overline{X})$ is flasque as a J -module. \square

Proof of Theorem 8.3. There exists a coflasque resolution

$$1 \rightarrow \widehat{Q} \rightarrow \widehat{P} \rightarrow \widehat{S} \rightarrow 1$$

of \widehat{S} . By Lemma 8.6, any morphism as in (a) factors through the one obtained from this resolution. Therefore, it suffices to consider only this morphism. Since S is flasque by Proposition 8.7, we are in the situation of Lemma 8.5. Thus (a) and (b) are equivalent.

By Theorem 3.14 of [Sal84], a neutral toric variety X is retract rational if and only if $\text{Pic}(\overline{X})$ is invertible as a Γ_k lattice. Thus (b) and (c) are equivalent. \square

9. Comparison to a construction of Merkurjev–Panin

In §7 of [MP97], A. Merkurjev and I. Panin describe a construction which assigns a separable algebra to each toric variety. Up to Brauer equivalence, their construction is essentially equivalent to ours. In this section, we make this precise.

In their construction, they fix a smooth projective toric variety Y and an action of a specific torus T on Y . Their notion of isomorphism amounts to T -equivariant isomorphisms of toric varieties. The set of isomorphism classes is thus in bijection with $H^1(k, T)$ and corresponds to the class of the open orbit Y_0 viewed as a T -torsor. All the forms of Y belong to the same neutralization class since T acts trivially on $\text{Pic}(\overline{Y})$. The neutralization X of Y corresponds to the trivial element of $H^1(k, T)$ and there is an injection $H^1(k, T) \rightarrow H^1(k, \text{Aut}_1(X))$ where $\text{Aut}_1(X)$ is the group of restricted automorphisms as in Remark 6.8.

First, we review the construction of Section 8 in a manner which is more directly comparable. Note that $\widehat{S} = \text{Pic}(\overline{X})$ is a Γ_k -lattice. Choose a morphism $\widehat{m} : \widehat{P} \rightarrow \widehat{S}$

of Γ_k -lattices where \widehat{P} is permutation with basis ω . Our choice of ω gives rise to a quasitrivial torus P and an étale algebra L and we obtain a homomorphism

$$\alpha_\omega : H^1(k, T) \rightarrow H^1(k, \text{Aut}_1(X)) \rightarrow \text{Br}(L)$$

which takes the isomorphism class $[Y_0]$ to a Brauer class in $\text{Br}(L)$.

Now we review the construction of Merkurjev–Panin. For the moment, assume that ω is a single Γ_k -orbit and thus that L is a field extension of k . The choice of ω gives us a morphism

$$S \rightarrow P \simeq R_{L/k}(\mathbb{G}_{m,L})$$

which is equivalent to a morphism $S_L \rightarrow \mathbb{G}_{m,L}$, which, by duality, gives a morphism $\mathbb{Z} \rightarrow S_L^*$. This last morphism can be viewed as an element Ω of $\text{Pic}(X_L)$.

Fix a Galois splitting field K/k for T which contains L . We obtain an element $[Y_0] \in H^1(\Gamma_{K/k}, T(K))$. From the exact sequence

$$1 \rightarrow T^* \rightarrow \widetilde{T}^* \rightarrow S^* \rightarrow 1$$

and the isomorphism $\text{Pic}(X_L) \simeq H^0(\Gamma_{K/L}, S^*)$, we obtain an element $\partial[\Omega] \in H^1(\Gamma_{K/L}, T^*)$ via the connecting homomorphism. We may take the cup product $[(Y_0)_L] \cup \partial[\Omega]$ in $H^2(\Gamma_{K/L}, K^\times)$ via the standard pairing $T(K) \otimes T^* \rightarrow K^\times$. Thus, we obtain a map

$$\beta_\omega : H^1(k, T) \rightarrow \text{Br}(L)$$

as desired. When ω is not a single Γ_k -orbit, we simply take products and again obtain an element Ω in $\text{Pic}(X_L)$ and a morphism β_ω .

Theorem 9.1. *For any element $[Y_0] \in H^1(k, T)$, our construction α_ω and the Merkurjev–Panin construction β_ω satisfy the relation*

$$\alpha_\omega([Y_0]) = -\beta_\omega([Y_0])$$

in the group $\text{Br}(L)$.

Proof. It suffices to prove the theorem for ω a single Γ_k -orbit. First, we claim that the following diagram commutes

$$\begin{CD} H^2(k, S) @>{m_*}>> H^2(k, P) \\ @VVV @VV{\simeq}V \\ H^2(L, S_L) @>{\Omega^*}>> H^2(L, \mathbb{G}_m) \end{CD}$$

where $P = R_{L/k}(\mathbb{G}_m)$. Note that the morphisms

$$m : S \rightarrow R_{L/k}(\mathbb{G}_m) \text{ and } D(\Omega) : S_L \rightarrow \mathbb{G}_m$$

arise from the adjunction of the functors $(-)_L$ and $R_{L/k}(-)$. Thus the morphism m factors as $R_{L/k}(D(\Omega)) \circ \eta(S)$ where $\eta(S) : S \rightarrow R_{L/k}(S_L)$ is the unit of adjunction. The morphism $H^2(k, S) \rightarrow H^2(L, S_L)$ factors through $H^2(k, S) \rightarrow$

$H^2(k, R_{L/k}(S_L))$ by the functorial construction of the restriction map (see §2.5 of [Ser02]). Thus, the diagram above is the same as

$$\begin{CD}
 H^2(k, S) @>>> H^2(k, R_{L/k}(S_L)) @>>> H^2(k, R_{L/k}(\mathbb{G}_m)) \\
 @. @VV \simeq V @VV \simeq V \\
 @. H^2(L, S_L) @>>> H^2(L, \mathbb{G}_m)
 \end{CD}$$

which commutes since $R_{L/k}(-)(k) \rightarrow (-)(L)$ is a natural transformation of functors from algebraic groups over k to abstract groups.

Recall that $\alpha_\omega([Y_0]) = m_*(\partial[Y_0])$; thus, by the above, we may instead write $\alpha_\omega([Y_0]) = \Omega^*(\partial[Y_0]_L) = \partial[(Y_0)_L] \cup \Omega$. Note that $\beta_\omega([Y_0]) = [(Y_0)_L] \cup \partial\Omega$. Choosing a Galois splitting field K for L with Galois group G , we need to compare two different methods of evaluating the map

$$H^1(G, T(K)) \otimes H^0(G, S^*) \rightarrow H^2(G, K^\times).$$

The theorem follows from Lemma 9.2 below. \square

Lemma 9.2. *Let G be a finite group. Let A, B, C be G -lattices and I be a G -module. Suppose there is an exact sequence*

$$1 \rightarrow A \rightarrow B \rightarrow C \rightarrow 1.$$

Up to the sign $(-1)^i$, the following square commutes

$$\begin{CD}
 H^i(G, \text{Hom}(A, I)) \otimes H^j(G, C) @>{\partial \otimes \text{id}}>> H^{i+1}(G, \text{Hom}(C, I)) \otimes H^j(G, C) \\
 @V \text{id} \otimes \partial VV @VV \cup V \\
 H^i(G, \text{Hom}(A, I)) \otimes H^{j+1}(G, A) @>{\cup}>> H^{i+j+1}(G, I)
 \end{CD}$$

for any integers $i \geq 0, j \geq 0$.

Proof. First, we show that, for any G -lattice M , the cup product and evaluation morphism

$$a : H^i(G, \text{Hom}(M, I)) \otimes H^j(G, M) \rightarrow H^{i+j}(G, I)$$

is isomorphic to the composition morphism

$$b : \text{Ext}_G^i(M, I) \otimes \text{Ext}_G^j(\mathbb{Z}, M) \rightarrow \text{Ext}_G^{i+j}(\mathbb{Z}, I).$$

The isomorphisms between Ext groups and cohomology groups follow from Proposition III.2.2 of [Bro82] since M is a free \mathbb{Z} -module. Picking a free resolution F_\bullet of the G -module \mathbb{Z} , the maps a and b can be represented by maps of chain complexes

$$\text{Hom}_G(F_\bullet, \text{Hom}(M, I)) \otimes \text{Hom}_G(F_\bullet, F_\bullet \otimes M) \rightarrow \text{Hom}_G(F_\bullet, I)$$

as in V.4.2 of [Bro82]. The map a takes elements $u \otimes v$ to $eval_*((u \otimes id_M) \circ v)$. The map b is defined as $\psi(u) \circ v$ where

$$\psi : \mathcal{H}om_G(F_\bullet, \text{Hom}(M, I)) \simeq \mathcal{H}om_G(F_\bullet \otimes M, I)$$

is the canonical isomorphism. If $v(f) = \sum_i f_i \otimes m_i$ for some $f \in F_j$ then we obtain

$$eval_*((u \otimes id_M) \circ v)(f) = eval_*\left(\sum_i u(f_i) \otimes m_i\right) = \sum_i u(f_i)(m_i) = (\psi(u) \circ v)(f)$$

for all $u \in \mathcal{H}om(F_\bullet, \text{Hom}(M, I))$ and $v \in \mathcal{H}om(F_\bullet, F_\bullet \otimes M)$ as desired.

Consider arbitrary elements $\alpha \in \text{Ext}_G^i(A, I)$ and $\gamma \in \text{Ext}_G^j(\mathbb{Z}, C)$. In view of the above, the statement of the lemma is equivalent to showing that $(\partial\alpha) \circ \gamma = (-1)^i \alpha \circ (\partial\gamma)$ as elements of $\text{Ext}_G^{i+j+1}(\mathbb{Z}, I)$. The exact sequence is an extension of C by A and thus it may be considered as an element β of $\text{Ext}_G^1(C, A)$. From Theorem III.9.1 of [ML63] we find that $\partial(\alpha) = (-1)^i \alpha \circ \beta$ and $\partial(\gamma) = \beta \circ \gamma$. Thus, the equality follows from the associativity of the composition $(\alpha \circ \beta) \circ \gamma = \alpha \circ (\beta \circ \gamma)$ as in Theorem III.5.4 of [ML63]. \square

Remark 9.3. As an immediate corollary of Theorem 9.1 we see that the Merkurjev–Panin construction is independent of the choice of torus T . Additionally, their construction can be extended to arbitrary forms of X rather than a fixed neutralization class by properly accounting for the group of class group automorphisms J .

Remark 9.4. The construction of Merkurjev–Panin as stated in [MP97] actually produces *algebras* rather than simply elements of the Brauer group. However, these algebras ultimately depend on a specific choice of torus T and a choice of splitting field. We discuss a more intrinsic method of producing algebras in the next section.

Remark 9.5. Note that $\Omega \in \text{Pic}(X_L)$ is denoted by “ Q ” in [MP97]. One cannot in general recover ω from Ω , as a corresponding set ω would only be Γ_k -invariant rather than J -invariant. However, this is not a serious shortcoming as one can always simply expand ω to be J -stable. Indeed, this is necessary if one wants to extend the Merkurjev–Panin construction to *all* forms of X rather than a single fixed neutralization class.

10. Associated separable algebras

In this section, we sharpen the results of Section 8 and prove Theorem 8.1. As discussed in Example 6.9, products of projective spaces are in bijective correspondence with their Cox endomorphism algebras. Thus, studying the isomorphism classes of separable algebras is equivalent to studying products of projective spaces.

Theorem 10.1. *Let X be a split smooth projective toric variety. Let J be the group of class group automorphisms of X . There exists a morphism*

$$f : X \rightarrow Y := \prod_{i=1}^N \mathbb{P}^{m_i}$$

where n_1, \dots, n_N are positive integers, and a morphism of functors

$$f_* : H^1(-, \text{Aut}(X)) \rightarrow H^1(-, \text{Aut}_J(Y))$$

such that, if X' is a form of X and Y' is a form of Y representing the class $f_*([X'])$, then there is a map

$$f' : X' \rightarrow Y'$$

which coincides with f over some field extension.

Proof. Given a morphism

$$f : X \rightarrow Y := \prod_{i=1}^N \mathbb{P}^{m_i}$$

as above, we obtain an induced map $\text{Pic}(Y) \rightarrow \text{Pic}(X)$. Thus, such morphisms give rise to a permutation basis ω of a permutation J -lattice $\widehat{P} := \text{Pic}(Y)$ as in Section 8. Conversely, if every element of ω corresponds to a sheaf which is generated by global sections then we obtain such a morphism.

Since X is projective, the ample cone spans $\text{Pic}(\overline{X}) \otimes \mathbb{R}$. In addition, the fixed locus of J must be a proper closed subset of $\text{Pic}(\overline{X}) \otimes \mathbb{R}$. Thus, we may assume that there exists a subset ω of $\text{Pic}(\overline{X})$ whose corresponding sheaves are generated by their global sections and such that J acts faithfully on ω . This provides the space Y and the desired map $f : X \rightarrow Y$.

The space Y is also a smooth split projective toric variety. We write \widetilde{Y} as its characteristic space sitting inside a vector space W and note that $\widehat{m} : \widehat{P} \rightarrow \text{Pic}(\overline{X})$ is just the morphism $\text{Pic}(Y) \rightarrow \text{Pic}(X)$ induced by f .

Labeling the basis of \widehat{P} as $\{E_1, \dots, E_r\}$ and labeling $\omega = \{D_1, \dots, D_r\}$, we have $\widehat{m}(E_i) = D_i$ for every $i = 1, \dots, r$. By duality, we obtain a morphism $m : S \rightarrow P$ of tori.

The construction of f is equivalent to an isomorphism

$$W^\vee = \bigoplus_{i=1}^r H^0(k, \mathcal{O}_Y(E_i)) \rightarrow \bigoplus_{i=1}^r H^0(k, \mathcal{O}_X(D_i))$$

which gives rise to a ring homomorphism $F^* : \text{Cox}(Y) \rightarrow \text{Cox}(X)$ since W contains all the generators of $\text{Cox}(Y)$. The ring homomorphism F^* is equivalent to a morphism $F : V \rightarrow W$. Since each line bundle $\mathcal{O}_X(D_i)$ is generated by global sections, the image of \widetilde{X} is contained in \widetilde{Y} so we have a restricted morphism $\widetilde{f} : \widetilde{X} \rightarrow \widetilde{Y}$. This descends to the morphism $f : X \rightarrow Y$ since the map \widetilde{f} is S -equivariant via $m : S \rightarrow P$.

Let B be the Cox endomorphism algebra of Y . We have the isomorphism

$$\text{GL}_1(B) \simeq \prod_{i=1}^r \text{GL}(H^0(k, \mathcal{O}(E_i))).$$

Note that $\widetilde{\text{Aut}}(Y) \simeq \text{GL}_1(B) \rtimes I$ where I is a finite group containing all possible permutations of the subgroups in the product. The group $\text{GL}_1(B) \rtimes J$ is the preimage of the restricted automorphism group $\text{Aut}_J(Y)$.

The group $\text{GL}_1(A)$ embeds in $\text{GL}_1(B)$ since it has a linear action on each component of $\text{Cox}(X)_{D_i}$. The group J embeds in $\text{GL}_1(B) \rtimes J$ by permuting monomials in the generators of $\text{Cox}(Y)$ since ω is J -stable. We obtain an S -equivariant homomorphism $\text{GL}_1(A) \rtimes J \rightarrow \text{GL}_1(B) \rtimes J$.

This descends to a morphism $\text{GL}_1(A)/S \rtimes J \rightarrow \text{Aut}_J(Y)$ after taking the quotient by S . We conclude that the morphism f is $\text{GL}_1(A)/S \rtimes J$ -equivariant. Using Proposition 5.7, we obtain $f' : X' \rightarrow Y'$ by descent. \square

Example 10.2 (Blunk’s algebras). Let X be a split del Pezzo surface of degree 6. In the notation of Example 6.3, consider the elements

$$\begin{aligned} a_1 &= H, & a_2 &= 2H - E_1 - E_2 - E_3, \\ b_1 &= H - E_1, & b_2 &= H - E_2, & b_3 &= H - E_3 \end{aligned}$$

in $\text{Pic}(X)$. The elements a_1, a_2 correspond to morphisms $X \rightarrow \mathbb{P}^2$ while b_1, b_2, b_3 correspond to morphisms $X \rightarrow \mathbb{P}^1$. Here, J is isomorphic to $S_3 \times S_2$ and the J -orbits are $\omega_1 = \{a_1, a_2\}$ and $\omega_2 = \{b_1, b_2, b_3\}$.

Let $\omega = \omega_1 \cup \omega_2$. In this special case, $\text{Aut}_J(Y) = \text{Aut}(Y)$. Thus Theorem 10.1 produces a natural transformation

$$f_* : H^1(-, \text{Aut}(X)) \rightarrow H^1(-, \text{Aut}(Y)).$$

Taking B to be the Cox endomorphism algebra of Y , we see that ω_1 produces an Azumaya algebra B_1 of rank 3^2 over an étale k -algebra of degree 2; ω_2 , produces an Azumaya algebra B_2 of rank 2^2 over an étale k -algebra of degree 3. Theorem 3.4 of [Blu10] shows f_* is injective in this case.

(Note that Blunk also specifies embeddings of an étale subalgebra into the algebras B_1 and B_2 . This is necessary to characterize the image of f_* , but we are only concerned with whether the map is injective. Blunk’s Theorem 3.4 uses a stronger notion of isomorphism on the algebras B_1, B_2 which respects the subalgebra embeddings. However, we show that the stronger notion of isomorphism is equivalent to the usual one; one can also see this directly.)

As our goal is to show that the natural transformation f_* is injective under suitable conditions, the following example demonstrates why we want to consider $\text{Aut}_J(Y)$ rather than $\text{Aut}(Y)$ in general.

Example 10.3. Continuing Example 7.10, consider $X = \mathbb{P}^1 \times \mathbb{P}^3$ over \mathbb{R} and let

$$f : X \rightarrow \mathbb{P}^3 \times \mathbb{P}^3$$

be the product of an inclusion $\mathbb{P}^1 \rightarrow \mathbb{P}^3$ as a linear subspace and the isomorphism $\mathbb{P}^3 \rightarrow \mathbb{P}^3$. The induced functor

$$H^1(-, \text{Aut}(X)) \rightarrow H^1(-, \text{Aut}(Y))$$

is *not injective* since the classes of $C \times \mathbb{P}^3$ and $\mathbb{P}^1 \times C'$ map to the same element. However, if we consider the functor

$$H^1(-, \text{Aut}(X)) \rightarrow H^1(-, \text{Aut}_J(Y))$$

then we do have injectivity. In this case injectivity can be fixed by a different choice of ω , but more generally the group J will not be a product of symmetric groups and such a fix will not exist.

Theorem 10.4. *Suppose we are in the situation of Theorem 10.1. If $\text{Pic}(\overline{X})$ is invertible as a J -lattice there is a canonical choice of maps such that f_* is injective.*

Proof. The construction of ω is as follows. Let $\text{Nef}(X)$ be the Nef cone of X . By Theorem 3.1 of [Mus02], we see that the divisors in the Nef cone of X are precisely those generated by their global sections; moreover, $\text{Nef}(X)$ can be extracted from the fan so it suffices to assume we are working over \mathbb{C} . We may identify $\text{Nef}(X)$ with a strongly convex rational polyhedral cone of full dimension in $\text{Pic}(X) \otimes \mathbb{R} = S^* \otimes \mathbb{R}$ (see Theorems 6.3.20 and 6.3.22 of [CLS11]). The intersection $\text{Nef}(X) \cap S^*$ is precisely the set of line bundles which are generated by their global sections.

For every subgroup G of J , consider the intersection $M_G = \text{Pic}(\overline{X})^G \cap \text{Nef}(X)$. Since the intersection of a rational polyhedral cone and a subspace cut out by rational linear equations is again a rational polyhedral cone, the monoid M_G is finitely generated. Indeed, since the cones are strongly convex, each monoid M_G has a canonical minimal generating set C_G . Take ω to be the union of the C_G for each $G \subset J$. Note that ω is J -stable since $j(C_G) = C_{j(G)}$ for each $j \in J$.

We claim that C_G spans $\text{Pic}(\overline{X})^G$ for every $G \subset J$. Indeed, we may find a field K and a cocycle $c \in Z^1(K, G)$ such that $\text{Pic}(cX) = \text{Pic}(\overline{X})^G$. Since projectivity is a geometric property, the ample cone (and thus the Nef cone) is of full dimension in $\text{Pic}(cX)$. Thus, M_G is of full dimension in $\text{Pic}(\overline{X})^G$. Thus C_G spans $\text{Pic}(\overline{X})^G$ as desired.

As in Lemma 3 of [CTS77], since we have chosen ω such that $\widehat{P}^G \rightarrow \text{Pic}(\overline{X})^G$ is surjective for all subgroups $G \subset J$, we conclude that the kernel \widehat{Q} is coflasque. In particular, the action of J is faithful. Furthermore, every element of ω corresponds to a sheaf which is generated by global sections. The result now follows from Lemma 8.5, Theorem 10.1, and Lemma 10.5 below. \square

Lemma 10.5. *In the situation of Theorem 10.1 we have the following commutative diagram*

$$\begin{CD} H^1(k, \text{Aut}(X)) @<\delta<< H^2(k, S \rightarrow J) \\ @V f_* VV @VV m_* V \\ H^1(k, \text{Aut}_J(Y)) @<\delta<< H^2(k, P \rightarrow J) \end{CD}$$

where the horizontal maps are as in Theorem 7.4.

Proof. From the proof of Theorem 10.1, there is a morphism

$$s : \text{GL}_1(A) \rtimes J \rightarrow \text{GL}_1(B) \rtimes J$$

which descends to the group homomorphism

$$r : \mathrm{GL}_1(A)/S \rtimes J \rightarrow \mathrm{Aut}_J(Y)$$

making f equivariant.

Let a be a cocycle in $H^1(k, \mathrm{GL}_1(A) \rtimes J) \simeq H^1(k, \mathrm{Aut}(X))$. To map a to $H^2(k, S \rightarrow J)$, we lift to a continuous function $b : \Gamma_k \rightarrow (\mathrm{GL}_1(A) \rtimes J)(k_s)$ and form the 2-cocycle Δb in $Z^2(k, {}_cS)$ for an appropriate cocycle $c \in Z^1(k, J)$. To map this to $H^2(k, P \rightarrow J)$, we simply take the 2-cocycle $s(\Delta b)$ in $Z^2(k, {}_cP)$.

Going the other way, map a to $H^1(k, \mathrm{Aut}_J(Y))$ by the morphism r . To map $r(a)$ to $H^2(k, P \rightarrow J)$, we may take the function $s(b)$ as a lift and then take $\Delta(s(b))$; again, this cocycle sits in $Z^2(k, {}_cP)$.

Since $s(\Delta b) = \Delta(s(b))$, the diagram commutes. \square

Remark 10.6. Note that the construction used in the proof of Theorem 10.4 has the advantage of being canonical. However, it may not be very economical: in Example 10.2, ω is simply the minimal generating set for $\mathrm{Nef}(X)$ while our canonical construction produces a larger set.

Finally, we are in position to prove Theorem 8.1.

Proof of Theorem 8.1. By Theorem 10.1 we may construct a product of projective spaces $Y = \prod \mathbb{P}^{n_i}$ along with a natural transformation

$$H^1(-, \mathrm{Aut}(X)) \rightarrow H^1(-, \mathrm{Aut}_J(Y)).$$

Moreover, since $\mathrm{Pic}(\overline{X})$ is an invertible J -lattice, we may select Y such that the natural transformation is injective by Theorem 10.4. Following Example 6.9 there is a canonical identification between the forms of Y and the forms of its Cox endomorphism algebra B . Thus we have a canonical separable algebra B with an injective natural transformation

$$H^1(-, \mathrm{Aut}(X)) \rightarrow H^1(-, \mathrm{Aut}_J(B))$$

as desired. \square

References

[AN09] E. Aldrovandi, B. Noohi, *Butterflies I. Morphisms of 2-group stacks*, Adv. Math. **221** (2009), no. 3, 687–773.

[BG99] W. Bruns, J. Gubeladze, *Polytopal linear groups*, J. Algebra **218** (1999), no. 2, 715–737.

[Blu10] M. Blunk, *Del Pezzo surfaces of degree 6 over an arbitrary field*, J. Algebra **323** (2010), no. 1, 42–58.

[Bor93] M. V. Borovoi, *Abelianization of the second nonabelian Galois cohomology*, Duke Math. J. **72** (1993), no. 1, 217–239.

[Bre10] L. Breen, *Notes on 1- and 2-gerbes*, in: *Towards Higher Categories*, IMA Vol. Math. Appl., Vol. 152, Springer, New York, 2010, pp. 193–235.

- [Bro82] K. S. Brown, *Cohomology of Groups*, Graduate Texts in Mathematics, Vol. 87, Springer-Verlag, New York, 1982. Russian transl.: К. С. Браун, *Когомология групп*, Наука, М., 1987.
- [Büh96] D. Bühler, *Homogener Koordinatenring und Automorphismengruppe vollständiger torischer Varietäten*, PhD thesis, University of Basel, 1996.
- [CGR08] V. Chernousov, P. Gille, Z. Reichstein, *Reduction of structure for torsors over semilocal rings*, Manuscripta Math. **126** (2008), no. 4, 465–480.
- [CLS11] D. A. Cox, J. B. Little, H. K. Schenck, *Toric Varieties*, Graduate Studies in Mathematics, Vol. 124, American Mathematical Society, Providence, RI, 2011.
- [Cox95] D. A. Cox, *The homogeneous coordinate ring of a toric variety*, J. Algebraic Geom. **4** (1995), no. 1, 17–50.
- [Cox14] D. A. Cox, *Erratum to “The homogeneous coordinate ring of a toric variety”*, J. Algebraic Geom. **23** (2014), no. 2, 393–398.
- [CTHS05] J.-L. Colliot-Thélène, D. Harari, A. N. Skorobogatov, *Compactification équivariante d’un tore (d’après Brylinski et Künnemann)*, Expo. Math. **23** (2005), no. 2, 161–170.
- [CTS77] J.-L. Colliot-Thélène, J.-J. Sansuc, *La R-équivalence sur les tores*, Ann. Sci. École Norm. Sup. (4) **10** (1977), no. 2, 175–229.
- [CTS87] J.-L. Colliot-Thélène, J.-J. Sansuc, *La descente sur les variétés rationnelles II*, Duke Math. J. **54** (1987), no. 2, 375–492.
- [Dem70] M. Demazure, *Sous-groupes algébriques de rang maximum du groupe de Cremona*, Ann. Sci. École Norm. Sup. (4) **3** (1970), no. 4, 507–588.
- [DP14] U. Derenthal, M. Pieropan, *Cox rings over nonclosed fields*, arXiv:1408.5358 (2014).
- [ELFST14] E. Javier Elizondo, P. Lima-Filho, F. Sottile, Z. Teitler, *Arithmetic toric varieties*, Math. Nachr. **287** (2014), no. 2–3, 216–241.
- [Flo04] M. Florence, *Zéro-cycles de degré un sur les espaces homogènes*, Int. Math. Res. Not. **54** (2004), 2897–2914.
- [FSS98] Y. Z. Flicker, C. Scheiderer, R. Sujatha, *Grothendieck’s theorem on non-abelian H^2 and local-global principles*, J. Amer. Math. Soc. **11** (1998), no. 3, 731–750.
- [Ful93] W. Fulton, *Introduction to Toric Varieties*, Annals of Mathematics Studies, Vol. 131, Princeton University Press, Princeton, NJ, 1993.
- [Gir71] J. Giraud, *Cohomologie Non Abélienne*, Die Grundlehren der mathematischen Wissenschaften, Band 179, Springer-Verlag, Berlin, 1971.
- [GMB13] P. Gille, L. Moret-Bailly, *Actions algébriques de groupes arithmétiques*, in: *Torsors, Étale Homotopy and Applications to Rational Points*, London Math. Soc. Lecture Note Ser., Vol. 405, Cambridge Univ. Press, Cambridge, 2013, pp. 231–249.
- [Hur11] M. Huruguen, *Toric varieties and spherical embeddings over an arbitrary field*, J. Algebra **342** (2011), 212–234.
- [HY12] A. Hoshi, A. Yamasaki, *Rationality problem for algebraic tori*, arXiv:1210.4525v4 (2012).

- [KMRT98] M.-A. Knus, A. Merkurjev, M. Rost, J.-P. Tignol, *The Book of Involutions*, American Mathematical Society Colloquium Publications, Vol. 44, American Mathematical Society, Providence, RI, 1998, with a preface in French by J. Tits.
- [Kun87] Б. Э. Кунявский, *Трёхмерные алгебраические торы*, в сб. *Исследования по теории чисел*, Саратовский гос. унив., Саратов, 1987, 90–111. Engl. transl.: В. Ё. Kunyavskii, *Three-dimensional algebraic tori*, Selecta Math. Soviet. **9** (1990), no. 1, 1–21.
- [ML63] S. Mac Lane, *Homology*, Die Grundlehren der mathematischen Wissenschaften, Bd. 114, Academic Press, New York, Springer-Verlag, Berlin, 1963. Russian transl.: С. Маклейн, *Гомология*, Мир, М., 1966.
- [MO67] H. Matsumura, F. Oort *Representability of group functors, and automorphisms of algebraic schemes*, Invent. Math. **4** (1967), 1–25.
- [Mer10] A. Merkurjev, *Periods of principal homogeneous spaces of algebraic tori*, Vestnik St. Petersburg Univ.: Math. **43** (2010), no. 1, 39–43.
- [MP97] A. Merkurjev, I. Panin, *K-theory of algebraic tori and toric varieties*, K-Theory **12** (1997), no. 2, 101–143.
- [Mus02] M. Mustaŭă, *Vanishing theorems on toric varieties*, Tohoku Math. J. (2) **54** (2002), no. 3, 451–470.
- [Nil06] B. Nill, *Complete toric varieties with reductive automorphism group*, Math. Z. **252** (2006), no. 4, 767–786.
- [Sal84] D. J. Saltman, *Retract rational fields and cyclic Galois extensions*, Israel J. Math. **47** (1984), no. 2–3, 165–215.
- [Sal99] D. J. Saltman, *Lectures on Division Algebras*, CBMS Regional Conference Series in Mathematics, Vol. 94, American Mathematical Society, Providence, RI, 1999.
- [Ser02] J.-P. Serre, *Galois Cohomology*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2002.
- [Sko09] А. Н. Скоробогатов, *Автоморфизмы и формы торических факторов однородных пространств*, Матем. сб. **200** (2009), ном. 10, 107–122. Engl. transl.: А. N. Skorobogatov, *Automorphisms and forms of torus quotients of homogeneous spaces*, Sbornik: Math. **200** (2009), no. 10, 1521–1536.
- [Spr66] Т. А. Springer, *Nonabelian H^2 in Galois cohomology*, in: *Algebraic Groups and Discontinuous Subgroups*, Proc. Sympos. Pure Math., Boulder, Colo., 1965, Amer. Math. Soc., Providence, RI, 1966, pp. 164–182.
- [VK84] В. Е. Воскресенский, А. А. Клячко, *Торические многообразия Фано и системы корней*, Изв. АН СССР, Сер. матем. **48** (1984), вып. 2, 237–263. Engl. transl.: V. E. Voskresenskii, A. A. Klyachko, *Toroidal Fano varieties and root systems*, Math. USSR-Izv. **24** (1985), no. 2, 221–244.
- [Vos67] В. Е. Воскресенский, *О двумерных алгебраических торах II*, Изв. АН СССР, Сер. матем. **31** (1967), вып. 3, 711–716. Engl. transl.: V. E. Voskresenskii, *On two-dimensional algebraic tori II*, Math. USSR-Izv. **1** (1967), no. 3, 691–696.
- [Vos82] В. Е. Воскресенский, *Проективные инвариантные модели Демажюра*, Изв. АН СССР, Сер. матем. **46** (1982), вып. 2, 195–210. Engl. transl.:

- V. E. Voskresenskii, *Projective invariant Demazure models*, Math. USSR-Izv. **20** (1983), no. 2, 189–202.
- [Vos98] V. E. Voskresenskii, *Algebraic Groups and Their Birational Invariants, Translations of Mathematical Monographs*, Vol. 179, American Mathematical Society, Providence, RI, 1998.
- [Wat79] W. C. Waterhouse, *Introduction to Affine Group Schemes*, Graduate Texts in Mathematics, Vol. 66, Springer-Verlag, New York, 1979.