

KÄHLER–EINSTEIN METRICS ON STABLE VARIETIES AND LOG CANONICAL PAIRS

ROBERT J. BERMAN AND HENRI GUENANCIA

Abstract. Let X be a canonically polarized variety, i.e. a complex projective variety such that its canonical class K_X defines an ample \mathbb{Q} -line bundle, and satisfying the conditions G_1 and S_2 . Our main result says that X admits a Kähler–Einstein metric iff X has semi-log canonical singularities i.e. iff X is a stable variety in the sense of Kollár–Shepherd-Barron and Alexeev (whose moduli spaces are known to be compact). By definition a Kähler–Einstein metric in this singular context simply means a Kähler–Einstein on the regular locus of X with volume equal to the algebraic volume of K_X , i.e. the top intersection number of K_X . We also show that such a metric is uniquely determined and extends to define a canonical positive current in $c_1(K_X)$. Combined with recent results of Odaka our main result shows that X admits a Kähler–Einstein metric iff X is K -stable, which thus confirms the Yau–Tian–Donaldson conjecture in this general setting of (possibly singular) canonically polarized varieties. More generally, our results are shown to hold in the setting of log minimal varieties and they also generalize some prior results concerning Kähler–Einstein metrics on quasi-projective varieties.

Contents

1	Introduction	1683
2	Preliminaries	1690
3	Singular Kähler–Einstein metrics	1694
4	Variational solutions	1701
5	Smoothness of the Kähler–Einstein metric	1708
6	Applications	1721
7	Outlook	1724
	References	1726

1 Introduction

According to the seminal works of Aubin [Aub78] and Yau [Yau78] any *canonically polarized* compact complex manifold X (i.e. X is a non-singular projective algebraic variety such that the canonical line bundle K_X is ample) admits a unique Kähler–Einstein metric ω in the first Chern class $c_1(K_X)$. One of the main goals of the present paper is to extend this result to the case when X is singular or more precisely when

X has *semi-log canonical* singularities. A major motivation comes from the fact that such singular varieties are used to compactify the moduli space of canonically polarized manifolds—a subject where there has been great progress in the last years in connection to the (log) Minimal Model Program (MMP) in birational algebraic geometry [Kol, Kov12]. The varieties in question are usually referred to as *stable varieties* (or *canonical models*) as they are the higher dimensional generalization of the classical notion of stable curves of genus $g > 1$, which form the Deligne–Mumford compactification of the moduli space of non-singular genus g curves [Kol, Kov12]. It is a classical fact that any stable curve admits a unique Kähler–Einstein metric on its regular part, whose total area is equal to the (arithmetic) degree of the curve X (see the section on stable curves further in this Introduction for more details). Our first (and main) result gives a generalization of this fact to the higher dimensional setting:

Theorem A. *Let X be a projective complex algebraic variety with semi-log canonical singularities such that K_X is ample. Then there exists a Kähler metric ω on the regular locus X_{reg} , satisfying*

$$\text{Ric } \omega = -\omega$$

and such that the volume of (X_{reg}, ω) coincides with the volume of K_X , i.e. $\int_{X_{\text{reg}}} \omega^n = c_1(K_X)^n$. Moreover, the metric extends to define a current ω in $c_1(K_X)$ which is uniquely determined by X .

We will refer to the current ω in the previous theorem as the (singular) Kähler–Einstein metric on X . Moreover, the current ω will be shown to be of *finite energy*, in the sense of [GZ07, BBGZ09] and as discussed in the last section of the present paper this allows one to define a canonical (singular) Weil–Petersson metric on the compact moduli space in terms of Deligne pairings. The notion of semi-log canonical singularities of a variety X —which is the most general class of singularities appearing in the (log) Minimal Model Program—will be recalled below. For the moment let us just point out that the definition involves two ingredients: first a condition which makes sure that the canonical divisor K_X is defined as a \mathbb{Q} -Cartier divisor (i.e. \mathbb{Q} -line bundle) which is in particular needed to make so sense of the notion of ampleness of K_X and secondly, the definition of semi-log canonical singularities involves a bound on the discrepancies of X on any resolution of singularities.

In fact, we will conversely show that if K_X is ample and the variety X admits a Kähler–Einstein metric then X has semi-log canonical singularities and this brings us to our second motivation for studying Kähler–Einstein metrics in the the singular setting, namely the Yau–Tian–Donaldson conjecture. Recall that this conjecture concerns polarized algebraic manifolds (X, L) , i.e. algebraic manifolds together with an ample line bundle $L \rightarrow X$ and it says that the first Chern class $c_1(L)$ of L contains a Kähler metric ω with constant scalar curvature if and only if (X, L) is K -stable. The latter notion of stability is of algebro-geometric nature and can be seen as an asymptotic form of the classical notions of Chow and Hilbert stability appearing in

Geometric Invariant Theory (GIT). However, while the notion of K -stability makes equal sense when X is singular it is less clear how to give a proper definition of a constant scalar curvature metric for a singular polarized variety (X, L) . But, as it turns out, the situation becomes more transparent in the case when L is equal to K_X or its dual, the anti-canonical bundle $-K_X$. The starting point is the basic fact that, when X is smooth, a Kähler metric in ω in $c_1(\pm K_X)$ has constant scalar curvature on all of X precisely when it has constant Ricci curvature, i.e. when ω is a Kähler–Einstein metric. Various generalizations of Kähler–Einstein metrics to the singular setting have been proposed in the literature, see e.g. [EGZ09, BEGZ10, BBEGZ11] etc. In this paper we will adopt the definition which appears in the formulation of the previous theorem (see also Section 3), i.e. a positive current in $c_1(\pm K_X)$ is said to define a (singular) Kähler–Einstein metric if defines a bona fide Kähler–Einstein metric on the regular locus X_{reg} and if its total volume there coincides with the algebraic top intersection number of $c_1(\pm K_X)$. This definition, first used in the Fano case in [BBEGZ11], has the virtue of generalizing all previously proposed definitions, regardless of the sign of the canonical line bundle. Combing our results with recent results of Odaka [Oda13b, Oda11], which say that a canonically polarized variety has semi-log canonical singularities precisely when (X, K_X) is K -stable, gives the following theorem, which can be seen as a confirmation of the generalized form of the Yau–Tian–Donaldson conjecture for canonically polarized varieties (satisfying the conditions G_1 and S_2 , cf. Theorem 3.10 for a more precise statement):

Theorem B. *Let X be a complex projective variety such that K_X is ample. Then X admits a Kähler–Einstein metric if and only if (X, K_X) is K -stable.*

It may also be illuminating to compare this result with the case when $L := -K_X$ is ample (i.e. X is Fano). Then it was shown in [Ber12], in the general singular setting, that the existence of a Kähler–Einstein metric indeed implies K -(poly)stability. As for the converse it was finally settled very recently in the deep works by Chen–Donaldson–Sun [CDS12a, CDS12b, CDS13] and Tian [Tia13], independently, in the case when X is smooth. The existence problem in the singular case is still open in general, except for the toric case [BB12]; cf. also [OSS12] for a related problem in the case of singular del Pezzo surfaces.

Coming back to the present setting we point out that the starting point of our approach is that, after passing to a suitable resolution of singularities, we may as well assume that the variety X is smooth if we work in the setting of *log pairs* (X, D) , where D is a \mathbb{Q} -divisor on X with simple normal crossings (SNC) and where the role of the canonical line bundle is played by the *log canonical line bundle* $K_X + D$ (which appears as the pull-back to the resolution of the original canonical line bundle). In this notation the original variety has semi-log canonical singularities precisely when the log pair (X, D) is *log canonical (lc)* in the usual sense of the Minimal Model Program, i.e. the coefficients of D are at most equal to one (but negative coefficients are allowed). However, it should be stressed that for this gain in regularity we have, of course, to pay a loss of positivity: even if the original canonical line bundle is

ample, the corresponding log canonical line bundle is only *semi-ample* (and big) on the resolution, since it is trivial along the exceptional divisors of the corresponding resolution.

The upshot is that the natural setting for our results is the setting of log smooth log canonical pairs (X, D) such that the log canonical line bundle $K_X + D$ is semi-ample and big. To any such pair we will associate a canonical Kähler–Einstein metric ω in the sense that ω is a current in the first Chern class $c_1(K_X + D)$ such that ω restricts to a bona fide Kähler–Einstein metric on a Zariski open set of X and such that, globally on X , the current defined by the divisor D gives a singular contribution to the Ricci curvature of ω .

Theorem C. *Let X be a Kähler manifold and D a simple normal crossings \mathbb{R} -divisor on X with coefficients in $] -\infty, 1]$ such that $K_X + D$ is semi-positive and big (i.e. $(K_X + D)^n > 0$). Then there exists a unique current ω in $c_1(K_X + D)$ which is smooth on a Zariski open set U of X and such that*

$$\text{Ric } \omega = -\omega + [D]$$

holds on X in the weak sense and $\int_U \omega^n = (K_X + D)^n$. More precisely, U can be taken to be the complement of D in the ample locus of $K_X + D$. Moreover, any such current ω on X automatically has finite energy.

Recall that the *ample locus* of a big line bundle L may be defined as the Zariski open set whose complement is the *augmented base locus* of L , i.e. the intersection of all effective \mathbb{Q} -divisors E such that $L - E$ is ample. In particular, if Y is a variety with semi-log canonical singularities and $\pi : (X, D) \rightarrow Y$ is a log resolution of the normalization (endowed with its conductor), then the exceptional locus of π is contained in the augmented base locus of $K_X + D$ and hence Theorem C above indeed implies Theorem A.

The existence proof of Theorem C (and its generalizations described below) will be divided into two parts: in the first part we construct a *variational solution* with *finite energy*, by adapting the variational techniques developed in [BBGZ09] to the present setting. Then, in the second part, we show that the variational solutions have appropriate regularity using a priori Laplacian estimates, building on the works of Aubin [Aub78] and Yau [Yau78] and ramifications of their work by Kobayashi [Kob84] and Tian–Yau [TY87] to the setting of quasi-projective varieties—in particular we will be relying on Yau’s maximum principle. For the second part we will need to perturb the line bundle $L := K_X + D$ (to make it ample) and regularize the klt part of the divisor D (to make the divisor purely log canonical).

It is interesting to note that, so far, we are not able of proving Theorem C without using the variational method, i.e. relying only on a priori estimates. The reason is that our estimates on the potential of the solution near $\text{Supp}(D)$ are not good enough to extend it directly as a current with full Monge–Ampère mass, so as to get the Kähler–Einstein metric of (X, D) .

Let us also point out that the variational part of our proof only requires $K_X + D$ to be *big* and thus produces a unique singular Kähler–Einstein metric of finite energy on any variety of log general type. As for the regularity part it applies as long as the corresponding log canonical ring is *finitely generated*. In fact, according to one of the fundamental conjectures of the general log minimal model program the latter finiteness property always holds. Note that for log canonical pairs, the finite generation is known to hold for $n \leq 4$, cf. [K+92, Fuj10] and the references therein. We also recall that in the case of varieties of log general type with log terminal singularities (which in our notation means that D_{lc} vanishes) the finite generation in question was established in the seminal work [BCHM10] and Kähler–Einstein metrics were first constructed in [EGZ09].

In the last section of the paper some applications of Theorem A are given. First, we explain the link with Yau–Tian–Donaldson as we indicated above in Theorem B. Then, we give a short analytic proof of the fact that the automorphism group of a normal stable variety is finite (see [FG12] or [BHPS12] for algebro-geometric proofs of more general results in this spirit, cf. also [Miy83]). We also discuss the problem of deducing Miyaoka–Yau type inequalities from Theorem A.

Further comparison with previous results

Stable curves. A stable curve, as defined by [DM69], is a reduced one equidimensional projective scheme over \mathbb{C} with only nodes as singularities and with finite automorphism group. The latter finiteness assumption can be replaced with various equivalent conditions, for example that the canonical sheaf of X is ample or in differential geometric terms: every connected component of $X_{\text{reg}} = X \setminus \{\text{nodes}\}$ is covered by the disk. In turn, this is equivalent to asking that every connected component of X_{reg} admits a *complete* hyperbolic metric.

As above, the higher dimensional analogue of Deligne–Mumford stable curves are the so-called stable varieties which are reduced equidimensional complex projective schemes with semi-log canonical singularities (i.e. double normal crossing singularities in codimension one, and log canonical singularities in higher codimension) and ample canonical bundle. In the light of the discussion above one could be tempted to believe that the regular locus of a stable variety can always be endowed with a complete Kähler–Einstein metric. However, this is not the case, mainly because of the singularities in codimension ≥ 2 (the relationship between the completeness of the metric and the singularities of the variety will be analyzed in [GW], and has also been largely investigated in the recent work of Song [Son14] that appeared after the first version of this article). Indeed, our main result (cf. Theorem 3.10) says that if we are given a scheme X (reduced, equidimensional, complex and projective) whose only singularities in codimension one are double normal crossings and such that K_X is ample, then X has semi-log canonical singularities (i.e. X is stable) if and only if X_{reg} admits a Kähler–Einstein metric ω with negative curvature such

that $\text{Vol}_\omega(X_{\text{reg}}) = c_1(K_X)^n$. This volume condition replaces in higher dimension the completeness condition, which does not hold in general. Further, this condition was known for a long time to be equivalent to the other ones in the one dimensional case already, see e.g. [HK13, Section 1].

Quasi-projective varieties. Theorem C also extends some of the results of Wu in [Wu08, Wu09], concerning the setting of Kähler–Einstein metrics on quasi-projective projective varieties of the form $X_0 := X - D$, where X is smooth and D is reduced SNC divisor. We recall that the case when $K_X + D$ is ample was independently settled by Kobayashi [Kob84] and Tian–Yau [TY87]. The case when X is an orbifold and $K_X + D$ is semi-ample and big was considered by Tian–Yau in [TY87] and as later shown by Yau [Yau93] the corresponding Kähler–Einstein metric is then complete on X_0 . (in the orbifold sense). However, in our general setting the metric will typically not be complete on the regular locus. This is only partly due to the klt singularities (which generalize orbifold singularities)—there is also a complication coming from the presence of negative coefficients on a resolution.

To illustrate this we recall that a standard example of log canonical pairs (X, D) is given by the Borel–Baily compactification $X := X_0 \cup D$ of an arithmetic quotient, i.e. $X_0 = B/\Gamma$, where B is a bounded symmetric domain and Γ is discrete subgroup of the automorphism group of B . In this case any toroidal resolution X' has the property that the corresponding divisor D' on the resolution is reduced (and hence purely log canonical) if Γ is neat, i.e. if there are no fixed points. The corresponding Kähler–Einstein metric on X_0 is the complete one induced from the corresponding metric on B , constructed in [CY80, MY83]. When Γ has fixed points these give rise to an additional fractional klt part D'_{klt} in D' so that the corresponding Kähler–Einstein metric is only complete in the orbifold sense [TY87]. However, for general log canonical singularity (X, D) the klt part D'_{klt} of D' may not be fractional or more seriously: it may contain negative coefficients and the main novelty of the present paper is to show how to deal with this problem by combining a variational approach with a priori estimates.

Behaviour at the boundary. It is also interesting to compare with the case when the pair (X, D) is log smooth with $K_X + D$ ample and with D effective and klt (i.e. with coefficients in $[0, 1]$), where very precise regularity results have been obtained recently. For example, in [Bre11, CGP13, JMR11, GP13] it is shown that the corresponding Kähler–Einstein metric ω has conical singularities along D (sometimes also called edge singularities in the literature), thus confirming a previous conjecture of Tian. As for the mixed case when the coefficient 1 is also allowed in D it was studied in [Gue12, GP13], where it was shown that ω has mixed cone and Poincaré type singularities. A common theme in these results is that singularities of the metric ω are encoded by a suitable local model (with cone or Poincaré type singularities) determined by D . However, the difficulty in the situation studied in the present paper is the presence of negative coefficients in D and the associated loss of positivity

which appears when we pass to a log resolution of a singular variety X . It would be very interesting if one could associate local models to this situation as well, but this seems very challenging even in the case when X has canonical singularities.

We should mention that the behaviour of the Kähler–Einstein metric of a log canonical pair (X, D) such that $K_X + D$ is ample and D is effective will be investigated in [GW]. As a consequence of the results therein, the Kähler–Einstein metric of a stable variety is equivalent to the cusp metric near the ordinary double points.

Organization of the paper

- Section 2: We introduce the preliminary material that we will need, concerning the pluripotential theoretic setting of singular metrics on line bundles over varieties which are not necessarily normal.
- Section 3: Here we give the definition of a Kähler–Einstein metric on a canonically polarized variety X and more generally on a log pair (X, D) . As we explain a purely differential-geometric definition can be given which only involves the regular locus X_{reg} of X . But, as we show, the corresponding metric automatically extends in a unique manner to define a singular current on X (which will allow us to prove the uniqueness of the Kähler–Einstein metric, later on in Section 4). We first treat the case when X has log canonical (and hence normal singularities) and then the general case of a variety X with semi-log singularities. Anyway, as we recall, the latter case reduces to the former (if one works in the setting of pairs) if one passes to the normalization.
- Section 4: We prove the uniqueness and existence of a weak Kähler–Einstein metric in the general setting of varieties of log general type. The existence is proved by adapting the variational approach to complex Monge–Ampère equations introduced in [BBGZ09] to the present setting. This method produces a singular Kähler–Einstein metric with finite energy (the new feature here compared to [BBGZ09] is that the reference measure does not have an L^1 density). We also use the variational approach to establish a stability result for the solutions to the equations induced from an (ample) perturbation of the log canonical line bundle on a resolution.
- Section 5: Here we establish the smoothness of the Kähler–Einstein metric, produced by the variational approach, on the regular locus of the variety X (or more generally, the pair (X, D)). The proof uses a perturbation argument in order to reduce the problem to the original setting of Kobayashi and Tian–Yau, combined with a priori estimates. But it should be stressed that in order to control the \mathcal{C}^0 norms we need to invoke the variational stability result proved in the previous section.
- Section 6: We give some applications to automorphism groups and show how to deduce the Yau–Tian–Donaldson conjecture for canonically polarized varieties from our results.
- Section 7: The paper is concluded with a brief outlook on possible applications to Miyaoka–Yau types inequalities, as well as the Weil–Peterson geometry of the

moduli space of stable varieties. These applications will require a more detailed regularity analysis of the Kähler–Einstein metrics that we leave for the future.

2 Preliminaries

We collect here some useful tools or notions that we are going to work with in this paper. We start with a compact Kähler manifold X of dimension n , and we consider a class $\alpha \in H^{1,1}(X, \mathbb{R})$ which is big. By definition, this means that α lies in the interior of the pseudo-effective cone, so that there exists a Kähler current $T \in \alpha$, that is a current which dominates some smooth positive form ω on X . We fix θ , a smooth representative of α .

The ample locus. An important invariant attached to α is the *ample locus* of α , denoted $\text{Amp}(\alpha)$, and introduced in [Bou04, Section 3.5]. This is the largest Zariski open subset U of X such that for all $x \in U$, there exists a Kähler current $T_x \in \alpha$ with analytic singularities such that T_x is smooth in an (analytic) neighbourhood of x . Its complement, called the augmented base locus, is usually denoted by $\mathbb{B}_+(\alpha)$. In the case when $\alpha = c_1(L)$ is the Chern class of a line bundle, it is known (see e.g. [BBP10]) that:

$$\mathbb{B}_+(L) = \bigcap_{\substack{L=A+E \\ A \text{ ample}, E \geq 0}} \text{Supp}(E)$$

Currents with minimal singularities. We will be very brief about this well-known notion, and refer e.g. to [Bou04, Section 2.8], [BBGZ09, Section 1], [Ber09] or [BD12] for more details and recent results.

By definition, if T, T' are two positive closed currents in the same cohomology class α , we say that T is less singular than T' if the local potentials φ, φ' of these currents satisfy $\varphi' \leq \varphi + O(1)$. It is clear that this definition does not depend on the choice of the local potentials, so that the definition is consistent. In each (pseudo-effective) cohomology class α , one can find a positive closed current T_{\min} which will be less singular than all the other ones; this current is not unique in general; only its class of singularities is. Such a current will be called current with minimal singularities.

One way to find such a current is to pick $\theta \in \alpha$ a smooth representative, and define then, following Demailly, the upper envelope

$$V_\theta := \sup\{\varphi \text{ } \theta\text{-psh}, \varphi \leq 0 \text{ on } X\}$$

Once observed that V_θ is θ -psh (in particular upper semi-continuous), it becomes clear that $\theta + dd^c V_\theta$ has minimal singularities.

Non-pluripolar Monge–Ampère operator. In the paper [BEGZ10], the authors define the non-pluripolar product $T \mapsto \langle T^n \rangle$ of any closed positive $(1, 1)$ -current $T \in \alpha$, which is shown to be a well-defined measure on X putting no mass on pluripolar sets, and extending the usual Monge–Ampère operator for Kähler forms (or having merely bounded potentials, cf. [BT87]). Let us note that when T is a smooth positive form ω on a Zariski dense open subset $\Omega \subset X$, then its Monge–Ampère $\langle T^n \rangle$ is simply the extension by 0 of the measure ω^n defined on Ω .

Given now a θ -psh function φ , one defines its non-pluripolar Monge–Ampère by $\text{MA}(\varphi) := \langle (\theta + dd^c \varphi)^n \rangle$. Then one can check easily from the construction that the total mass of $\text{MA}(\varphi)$ is less than or equal to the volume $\text{vol}(\alpha)$ of the class α (cf. [Bou02]):

$$\int_X \text{MA}(\varphi) \leq \text{vol}(\alpha)$$

A particular class of θ -psh functions that appears naturally is the one for which the last inequality is an equality. We will say that such functions (or the associated currents) have *full Monge–Ampère mass*. For example, θ -psh functions with minimal singularities have full Monge–Ampère mass (cf. [BEGZ10, Theorem 1.16]).

Plurisubharmonic functions on complex spaces. Here again, we just intend to give a short overview of the extension of the pluripotential theory to (reduced) complex Kähler spaces. A very good reference is [Dem85], or [EGZ09, Section 5] which is written in relation to singular Kähler–Einstein metric. We also refer to the preliminary parts of [Var89] or [FS90].

The data of a reduced complex space X includes the data of the sheaves of continuous and holomorphic functions. So the first object we would like to give a sense to is the sheaf \mathcal{C}_X^∞ of smooth functions. It may be defined as the restriction of smooth functions in some local embeddings of X in some \mathbb{C}^n . One defines similarly the sheaves of smooth (p, q) -forms $\mathcal{A}_X^{p,q}$ which carry the differentials $d, \partial, \bar{\partial}$ satisfying the usual rules; the space of currents is by definition the dual of the space of differential forms as in the smooth case. The sheaves complexes that are induced (Dolbeault, de Rham, etc.) are however not exact in general.

Another important sheaf is the one of pluriharmonic functions. They are defined to be smooth functions locally equal to the imaginary part of some holomorphic functions. One can show (see e.g. [FS90]) that a continuous function which is pluriharmonic on X_{reg} in the usual sense is automatically pluriharmonic on X . We denote by \mathcal{PH}_X the sheaf of real-valued pluriharmonic functions on X .

Let us move on to psh functions now. There are actually two possible definitions which extend the usual one for complex manifolds. The first one, introduced by Grauert and Remmert, mimics the one in the smooth case: we will say that a function $\varphi : X \rightarrow \mathbb{R} \cup \{-\infty\}$ is plurisubharmonic if it is upper semi-continuous and if for all holomorphic map $f : \Delta \rightarrow X$ from the unit disc in \mathbb{C} , the function $\varphi \circ f$ is subharmonic.

We could also introduce a more local definition: a function $\varphi : X \rightarrow \mathbb{R} \cup \{-\infty\}$ is strongly plurisubharmonic if in any local embeddings $i_\alpha : X \supset U_\alpha \hookrightarrow \mathbb{C}^n$, φ is the restriction of a psh function defined on an open set $\Omega_i \subset \mathbb{C}^n$ containing $i_\alpha(U_\alpha)$, if $X = \cup_\alpha U_\alpha$ is an open covering.

Clearly, a strongly psh function is also psh. Actually, Fornaess and Narasimhan [FN80] showed that these notions coincide: a function on X is psh if and only if it is strongly psh. On *normal* spaces one still has a Riemann extension theorem for psh functions, thanks to [GR56]. More precisely, if X is normal, $Y \subsetneq X$ is any proper analytic subspace, and $\varphi : X \setminus Y \rightarrow \mathbb{R} \cup \{-\infty\}$ is psh, then φ extends to a (unique) psh function on X if and only if it is locally bounded above near the points of Y , condition which is always realized if Y has codimension at least two in X . In particular, if X is normal, the data of a psh function on X is equivalent to the data of a psh function on X_{reg} .

Moreover, one can show (cf [BEG13, Lemma 3.6.1]) that a pluriharmonic function on X_{reg} automatically extends to a pluriharmonic function on X .

On non-normal spaces, one has to be more cautious, and it is convenient to introduce the notion of weakly psh function. Let X be a reduced complex space, and $\nu : X^\nu \rightarrow X$ its normalization. We say that a function $\varphi : X \rightarrow \mathbb{R} \cup \{-\infty\}$ is weakly psh if $\nu^*\varphi = \varphi \circ \nu$ is psh. It is not hard to see that a weakly psh function φ induces a bona fide psh function on X_{reg} which is locally bounded from above near the points of X_{sing} . Conversely, any psh function on X_{reg} which is locally upper bounded extends to a weakly psh function on X . On a normal space, a weakly psh function is of course psh, but in general these notions are different: consider $X = \{zw = 0\} \subset \mathbb{C}^2$, and $\varphi(x) = 0$ or 1 according to the connected component of $x \in X$. We refer to [Dem85, Théorème 1.10] for equivalent characterizations of weakly psh functions and conditions on a weakly psh function that ensure that it is already psh.

Finally, one can check that a (strongly) psh function φ on a complex space X is always locally integrable with respect to the area measure induced by any local embedding of X in \mathbb{C}^n (note that this is stronger than saying that φ is locally integrable on X_{reg} with respect to some volume form). Moreover, a locally integrable function φ is (almost everywhere) weakly psh if and only if it is locally bounded from above and $dd^c\varphi$ is a positive current.

Weights and Chern classes. From now on, X will be a *normal* complex space unless stated otherwise.

The definition of a (smooth) Kähler form is rather natural: it is a smooth real $(1, 1)$ -form written locally as $dd^c\psi$ for some (smooth) strictly psh function ψ ; equivalently this is locally the restriction of a Kähler form in an embedding in \mathbb{C}^n . Note that we could interpret this definition in terms of hermitian metrics on the Zariski tangent bundle of X , cf. [Var89].

Let us now consider a line bundle L on X . A smooth hermitian metric h on L is defined as in the smooth case: using trivialisations $\tau_\alpha : L|_{U_\alpha} \xrightarrow{\cong} U_\alpha \times \mathbb{C}$, we just

ask h to be written as $h(v) = |\tau_\alpha(v)|^2 e^{-\varphi_\alpha(z)}$ where φ_α is a smooth function on U_α . We say that the data $\phi := \{(U_\alpha, \varphi_\alpha)\}$ is a weight on L , so that it is equivalent to consider a weight or an hermitian metric.

Observe that if $(g_{\alpha\beta} : U_\alpha \cap U_\beta \rightarrow \mathbb{C}^*)$ is the cocycle in $H^1(X, \mathcal{O}_X^*)$ determined by the τ_α 's (more precisely $\tau_\alpha \circ \tau_\beta^{-1}(z, v) = (z, g_{\alpha\beta}v)$), then we have necessarily $\varphi_\beta - \varphi_\alpha = \log |g_{\alpha\beta}|^2$. In particular, the forms $dd^c \varphi_\alpha$ glue to a global smooth $(1, 1)$ -form on X called curvature of (L, h) and denoted by $c_1(L, h)$. This forms lives naturally in the space $H^0(X, \mathcal{C}_X^\infty / \mathcal{PH}_X)$, and using the exact sequence

$$0 \longrightarrow \mathcal{PH}_X \longrightarrow \mathcal{C}_X^\infty \longrightarrow C_X^\infty / \mathcal{PH}_X \longrightarrow 0$$

one may attach to (L, h) a class $\hat{c}_1(L, h) \in H^1(X, \mathcal{PH}_X)$. It is then easy to see that this class actually does not depend on the choice of h , so we will denote it by $\hat{c}_1(L)$. If X is smooth, $H^1(X, \mathcal{PH}_X) \simeq H^{1,1}(X, \mathbb{R})$, and it is well-known that $\hat{c}_1(L)$ coincides with the image of $L \in H^1(X, \mathcal{O}_X^*)$ in $H^2(X, \mathbb{Z})$ via the connecting morphism induced by the exponential exact sequence

$$0 \longrightarrow \mathbb{Z} \longrightarrow \mathcal{O}_X \xrightarrow{e^{2i\pi \cdot}} \mathcal{O}_X^* \longrightarrow 0.$$

This sequence also exists on any (even non-reduced) complex space, so that $c_1(L) \in H^2(X, \mathbb{Z})$ is well-defined; it will be more convenient for us to look at the image of $c_1(L)$ in $H^2(X, \mathbb{R})$ however. To relate it to $\hat{c}_1(L)$, we may use the following exact sequence:

$$0 \longrightarrow \mathbb{R} \longrightarrow \mathcal{O}_X \xrightarrow{-2\text{Im}(\cdot)} \mathcal{PH}_X \longrightarrow 0. \quad (2.1)$$

It is not hard to check that the connecting morphism $H^1(X, \mathcal{PH}_X) \rightarrow H^2(X, \mathbb{R})$ sends $\hat{c}_1(L)$ to $c_1(L)$ as expected.

We will also have to consider singular weights, which are by definition couples $\phi := \{(U_\alpha, \varphi_\alpha)\}$ where U_α is covering of X trivializing L , and φ_α are locally integrable on U_α , satisfying $\varphi_\beta - \varphi_\alpha = \log |g_{\alpha\beta}|^2$ on $U_\alpha \cap U_\beta$. The associated curvature current, denoted by $dd^c \phi$, is well-defined on X . The weight is said psh if the φ_α are, in which case $dd^c \phi$ is a positive current. Moreover, we can proceed as in the smooth case to attach to $dd^c \phi$ a class $\hat{c}_1(L) \in H^1(X, \mathcal{PH}_X)$ (consider ϕ as a section of the sheaf $L_{\text{loc}}^1 / \mathcal{PH}_X$ and use the natural exact sequence), whose image in $H^2(X, \mathbb{R})$ via the long exact sequence in cohomology induced by (2.1) is $c_1(L)$. Therefore when ϕ is a singular weight on L , we may say that $dd^c \phi$ is a current in $c_1(L)$.

We claim that a (possibly singular) psh weight ϕ on $L|_{X_{\text{reg}}}$ —and thus a psh weight in the usual sense—automatically extends to a (unique) psh weight $\tilde{\phi}$ on L . Indeed, by Grauert and Remmert's theorem, the φ_α 's defined on $U_\alpha \cap X_{\text{reg}}$ extend to a psh function $\tilde{\varphi}_\alpha$ on the whole U_α , which is moreover defined by $\tilde{\varphi}_\alpha(z_0) = \limsup_{X_{\text{reg}} \ni z \rightarrow z_0} \varphi_\alpha(z)$. Therefore, the relation $\tilde{\varphi}_\beta = \tilde{\varphi}_\alpha + \log |g_{\alpha\beta}|^2$ is immediately satisfied on the whole $U_\alpha \cap U_\beta$, which proves the claim.

If we get back to the non-normal case, we can define psh weights using weakly psh functions instead of psh functions. So the general philosophy is that we always

pull-back our objects to the normalization where things behave better, and the notions downstairs are defined and studied upstairs. For example, we can define on a normal variety the analogue of the non-pluripolar product and consider Monge–Ampère equations as in the smooth case (cf. [BBEGZ11, Section 1.1–1.2]). Then, if we write them on a non-normal variety, they have to be thought as pulled-back to the normalization.

Log canonical pairs. Following the by now common terminology of Mori theory and the minimal model program (cf. e.g. [KM98]), a pair (X, D) is by definition a complex normal projective variety X carrying a Weil \mathbb{Q} -divisor D (not necessarily effective). We will say that the pair (X, D) is a log canonical pair if $K_X + D$ (which is a priori defined as a Weil divisor) is \mathbb{Q} -Cartier, and if for some (or equivalently any) log resolution $\pi : X' \rightarrow X$, we have:

$$K_{X'} = \pi^*(K_X + D) + \sum a_i E_i$$

where E_i are either exceptional divisors or components of the strict transform of D , and the coefficients a_i satisfy the inequality $a_i \geq -1$.

3 Singular Kähler–Einstein metrics

3.1 Kähler–Einstein metrics on pairs. In this section, we will consider log pairs (X, D) where X is a complex normal projective variety, D is a Weil divisor, and $K_X + D$ is assumed to be \mathbb{Q} -Cartier. On X_{reg} , $D = D|_{X_{\text{reg}}}$ is a Cartier divisor, therefore there exists a psh weight ϕ_D on X_{reg} satisfying $dd^c \phi_D = [D|_{X_{\text{reg}}}]$, where $[D|_{X_{\text{reg}}}]$ stands for the current of integration along $D|_{X_{\text{reg}}}$. The first definition concerns the Ricci curvature of currents:

DEFINITION 3.1. *Let ω be a positive current on X_{reg} ; we say that ω is admissible if it satisfies:*

1. *Its non-pluripolar product $\langle \omega^n \rangle$ defines a (locally) absolutely continuous measure on X_{reg} with respect to $d\mathbf{z} \wedge d\bar{\mathbf{z}}$, where $\mathbf{z} = (z_i)$ are local holomorphic coordinates.*
2. *The function $\log(\langle \omega^n \rangle / d\mathbf{z} \wedge d\bar{\mathbf{z}})$ belongs to $L^1_{\text{loc}}(X_{\text{reg}})$.*

In that case, we define (on X_{reg}) the Ricci curvature of ω by setting $\text{Ric } \omega := -dd^c \log \langle \omega^n \rangle$.

Another way of thinking of this is to interpret the positive measure $\langle \omega^n \rangle|_{X_{\text{reg}}}$ as a singular metric on $-K_{X_{\text{reg}}}$ whose curvature is $\text{Ric } \omega$ by definition.

The measure e^ϕ for ϕ a weight on K_X . In the same spirit, we will use the convenient but somehow abusive notation e^ϕ for ϕ a weight on K_X (whenever the latter is defined as a \mathbb{Q} -line bundle) to refer to the positive measure $e^{\varphi z} d\mathbf{z} \wedge d\bar{\mathbf{z}}$ defined on

X_{reg} and extended by 0 to X ; where $\varphi_{\mathbf{z}}$ is the expression on some trivializing chart of X_{reg} (and hence of $K_{X_{\text{reg}}}$ too) of ϕ . In particular, for ϕ a psh weight on $K_X + D$, the measure $e^{\phi - \phi_D}$ can easily be pulled-back to any log resolution (π, X', D') of (X, D) (we pull it back over $X_{\text{reg}} \setminus \text{Supp}(D)$ and then extend it by 0), where it become $e^{\phi \circ \pi - \phi_{D'}}$.

We may now introduce the notion of (negatively curved) Kähler–Einstein metric attached to a pair (X, D) :

DEFINITION 3.2. *Let (X, D) be a log pair; we say that a positive admissible current ω is a Kähler–Einstein metric with negative curvature for (X, D) if:*

1. $\text{Ric}\omega = -\omega + [D]$ on X_{reg} ,
2. $\int_{X_{\text{reg}}} \langle \omega^n \rangle = c_1(K_X + D)^n$.

These conditions are sufficient to show that a Kähler–Einstein metric is a global solution of a Monge–Ampère equation. More precisely, we have the following:

PROPOSITION 3.3. *Let (X, D) be a log pair, and ω be a Kähler–Einstein metric for (X, D) . Then $\phi := \log \langle \omega^n \rangle + \phi_D$ extends to X as a psh weight with full Monge–Ampère mass on $K_X + D$, solution of*

$$\langle (dd^c \phi)^n \rangle = e^{\phi - \phi_D}.$$

Conversely, any psh weight ϕ on $K_X + D$ with full Monge–Ampère mass solution of the equation above induces a Kähler–Einstein metric $\omega := dd^c \phi$ for (X, D) .

Proof. On X_{reg} , we have $dd^c \phi = \omega$ thus ϕ is a psh weight on $(K_X + D)|_{X_{\text{reg}}}$, and thanks to a theorem of Grauert and Remmert, it extends through X_{sing} which has codimension at least 2. Clearly, we have $\omega = dd^c \phi$ on X , and by condition 2. in the definition of a Kähler–Einstein metric, ϕ has full Monge–Ampère mass. Then by definition, the two (non-pluripolar) measures $\langle (dd^c \phi)^n \rangle$ and $e^{\phi - \phi_D}$ coincide.

For the converse, let $\omega := dd^c \phi$; clearly condition 2. is satisfied. Moreover, ϕ and ϕ_D are locally integrable, so that ω is admissible and $\text{Ric}\omega = -dd^c(\phi - \phi_D) = -\omega + [D]$. \square

This proposition shows that the different definitions of what should be a singular Kähler–Einstein metric, appearing e.g. in [Ber11, BEGZ10, CGP13, EGZ09] etc. coincide. Moreover, one could equally define positively curved Kähler–Einstein metrics in an equivalent way as in [BBEGZ11]. In particular this object, intrinsically defined on X , can also be seen on any log resolution in the usual way; in practice, we will most of the time work on log resolutions when dealing with existence or smoothness questions.

Note also that we could have chosen to define a Kähler–Einstein metric attached to a pair (X, D) (say satisfying $K_X + D$ ample) to be a *smooth* Kähler metric ω on $X_{\text{reg}} \setminus \text{Supp}(D)$ which extends to an admissible current on X_{reg} satisfying

there $\text{Ric}\omega = -\omega + [D]$ and the mass condition $\left(\int_{X_{\text{reg}}}\langle\omega^n\rangle =\right)\int_{X_{\text{reg}}\setminus\text{Supp}(D)}\omega^n = c_1(K_X + D)^n$.

Then, our regularity Theorem (say combined with Proposition 6.1) shows a posteriori that this definition would have coincided with Definition 3.2.

Let us also mention that in the case of a log smooth log canonical pair (X, D) , the same proof as [Gue12, Proposition 2.5] combined with [GW] will show that the data of a negatively curved Kähler–Einstein on (X, D) is equivalent to giving an admissible current ω on $X \setminus \text{Supp}(D)$ such that:

- $\text{Ric}\omega = -\omega$ on $X \setminus \text{Supp}(D)$,
- There exists $C > 0$ such that

$$C^{-1}dV \leq \prod_{a_j < 1} |s_j|^{2a_j} \cdot \prod_{a_k = 1} (|s_k|^2 \log^2 |s_k|^2) \omega^n \leq CdV$$

for some volume form dV on X , and where $D = \sum a_i D_i$, s_i is a defining section of D_i .

3.2 Kähler–Einstein metrics on stable varieties. Stable varieties, as considered e.g. in [KSB88, Ale96, Kar00, Kol, Kov12] are the appropriate singular varieties to look at if one wants to compactify the moduli space of canonically polarized projective varieties (cf. also [Vie95]). Before giving the precise definition of a stable variety, we explain very briefly that notion and give the connection with Kähler–Einstein theory. In the next section, we will give a more detailed account of the type of singularities involved.

So first of all, we will consider complex varieties that are Gorenstein in codimension 1 (this condition replaces regularity in codimension 1 for normal varieties) and satisfy the condition S_2 of Serre. Basically, the singularities in codimension 1 of our varieties are those of the union of two coordinate hyperplanes (“double crossing”), so it is important to be aware that such varieties are in general not irreducible, and hence their normalization will not be connected.

Now we want to recast them in the context given by the singularities of the minimal model program (MMP); so we consider such a variety X and its normalization $\nu : X^\nu \rightarrow X$. One can write $\nu^*K_X = K_{X^\nu} + D$ for some reduced divisor D called the conductor of ν ; it sits above the codimension 1 component of the singular locus of X . We then say that X has semi log canonical singularities if the pair (X^ν, D) is log canonical in the usual sense. The generalization of the notion of stable curve is given by the following definition:

DEFINITION 3.4. *A projective variety X is called stable if X has semi-log canonical singularities, and K_X is an ample \mathbb{Q} -line bundle.*

There is a subtlety for the definition of K_X , but we refer to Section 3.3 for appropriate explanations. It is actually possible to define the notion of Kähler–Einstein metric for a stable variety:

DEFINITION 3.5. Let X be a stable variety. A Kähler–Einstein metric on X is a positive admissible current ω on X_{reg} such that:

1. $\text{Ric } \omega = -\omega$ on X_{reg} ,
2. $\int_{X_{\text{reg}}} \langle \omega^n \rangle = c_1(K_X)^n$.

In the non-normal case however, psh weight do not automatically extend across the singularities, so that it is not clear that the Kähler–Einstein metric will extend as a positive current on K_X satisfying the usual Monge–Ampère equation *globally*. Actually, this is the case as shows the following proposition:

PROPOSITION 3.6. Let X be a stable variety, and ω a Kähler–Einstein metric on X . Then the weight $\phi := \log \omega^n$ extends to X as a weakly psh weight on K_X solution of the Monge–Ampère equation $\langle (dd^c \phi)^n \rangle = e^\phi$.

Proof. Taking the dd^c of each side in the definition of ϕ and using the Ricci equation, we find $\omega = dd^c \phi$ on X_{reg} , and therefore ϕ satisfies $\langle (dd^c \phi)^n \rangle = \langle \omega^n \rangle = e^\phi$. Pulling back this equation to normalization X^ν , we find a psh weight $\phi' = \nu^* \phi$ on $c_1(\nu^* K_X)|_{\nu^{-1}(X_{\text{reg}})}$ solution of $\langle (dd^c \phi')^n \rangle = e^{\phi' - \phi_D}$ where D is the conductor of the normalization. As we work inside X_{reg}^ν and the integral $\int_{\nu^{-1}(X_{\text{reg}})} e^{\phi' - \phi_D}$ is finite, we infer from Lemma 3.7 below that ϕ' extends (as a psh weight) across D_{reg} . So ϕ' induces a psh weight on $c_1(\nu^* K_X)|_{X_{\text{reg}}^\nu \setminus D_{\text{sing}}}$, and by normality of X^ν , it extends to the whole X^ν , which means precisely that ϕ extends as a weakly psh weight on K_X . The expected Monge–Ampère equation holds automatically on X (or equivalently on X') since both measures $\langle (dd^c \phi)^n \rangle$ and e^ϕ put no mass on X_{sing} by definition. \square

In the previous proof, we used the following extension result:

LEMMA 3.7. Let U be a neighbourhood of $0 \in \mathbb{C}^n$, $H = \{z_1 = 0\} \subset \mathbb{C}^n$, and φ be a psh function defined on $U \setminus H$. We assume that the integral

$$\int_{U \setminus H} \frac{e^\varphi}{|z_1|^2} dV$$

is finite. Then φ extends across H , and more precisely φ tends to $-\infty$ near H .

Proof. (thanks to Bo Berndtsson for providing us with this elegant proof) Assume, to get a contradiction, that φ does not tend to $-\infty$ near H , and let $V := U \setminus H$. Then we can find a sequence (x_k) of points in V converging to H such that $\varphi(x_k) \geq -C$ for some constant C . We write $x_k = (x_{1,k}, \dots, x_{n,k})$, and we set $r_k = |x_{1,k}|/2$; the sequence (r_k) converges to 0, and if D_k denotes the polydisk centered at x_k with polyradius $(r_k, \delta, \dots, \delta)$ for some fixed $\delta > 0$, then we have $D_k \subset V$.

Using the mean value inequality for φ at x_k , we find:

$$-C \leq \frac{1}{\text{vol}(D_k)} \int_{D_k} \varphi dV.$$

Therefore, using Jensen’s inequality, we obtain, up to modifying C by a normalization factor depending only on the dimension n :

$$e^{-C} \leq \int_{D_k} \frac{e^\varphi dV}{r_k^2 \delta^{2(n-1)}}$$

but on D_k , $|z_1| \leq 3r_k$ so

$$e^{-C'} \leq \int_{D_k} \frac{e^\varphi dV}{|z_1|^2}$$

for $C' = C + \log 9 - 2(n - 1) \log \delta$. As the measure of D_k goes to zero when $k \rightarrow +\infty$, it shows that the integral $\int_{U \setminus H} \frac{e^\varphi}{|z_1|^2} dV$ is infinite, which is absurd. \square

3.3 Singularities of stable varieties. In this paragraph, we intend to give a more precise overview of the notion of semi-log canonical singularities. As we will just touch on this topic, we refer to the nice survey [Kov12] for a broader study. Other good references are [Kol, KSB88].

In the following, X will always be a reduced and equidimensional scheme of finite type over \mathbb{C} , and we set $n := \dim X$. We emphasize again on the fact that X will not be irreducible in general.

The conditions G_1 and S_2 . As we saw earlier, we need a canonical sheaf. The condition G_1 will guarantee its existence, and the condition S_2 will (among other things) ensure its uniqueness.

If X is Cohen–Macaulay (for every $x \in X$, the depth of $\mathcal{O}_{X,x}$ is equal to its Krull dimension), then X admits a dualizing sheaf ω_X . We say that X is Gorenstein if X is Cohen–Macaulay and ω_X is a line bundle. We say that X is G_1 if X is Gorenstein in codimension 1, which means that there is an open subset $U \subset X$ which is Gorenstein and satisfies $\text{codim}_X(X \setminus U) \geq 2$.

We say that X satisfies the condition S_2 of Serre if for all $x \in X$, we have $\text{depth}(\mathcal{O}_{X,x}) \geq \min\{\text{ht}(\mathfrak{m}_{X,x}), 2\}$, where $\text{ht}(\mathfrak{m}_{X,x}) = \text{codim}(\bar{x})$ denotes the height of the maximal ideal $\mathfrak{m}_{X,x}$ of $\mathcal{O}_{X,x}$. This condition is equivalent to saying that for each closed subset $i : Z \hookrightarrow X$ of codimension at least two, the natural map $\mathcal{O}_X \rightarrow i_* \mathcal{O}_{X \setminus Z}$ is an isomorphism.

If X is G_1 and S_2 , and $U \subset X$ is a Gorenstein open subset whose complement has codimension at least 2, one can then define the canonical sheaf ω_X by $\omega_X := j_* \omega_U$ where $j : U \hookrightarrow X$ is the open embedding, and ω_U is the dualizing sheaf of U . By definition, this is a rank one reflexive sheaf. When X is projective, we know that it admits a dualizing sheaf; as it is reflexive, it coincides with ω_X by the S_2 condition.

We would like to have an interpretation of ω_X , or at least ω_U in terms of Weil divisors as in the normal case where we define the Weil divisor K_X as the closure of some Weil divisor representing the line bundle $K_{X_{\text{reg}}}$. But we have to be more cautious in the non normal case as it is not clear how we should extend a Weil

divisor given on X_{reg} . Actually, this is where the G_1 conditions appears: as ω_U is a line bundle, or equivalently a Cartier divisor, we may choose a Weil divisor K_U whose support does not contain any component of X_{sing} of codimension 1 and represent ω_U (write ω_U as the difference of two very ample line bundles). Then we define K_X to be the closure of K_U . Clearly, the divisorial sheaf $\mathcal{O}_X(K_X)$ is reflexive, and coincides with $\omega_U = \omega_X|_U$ on U , so that by the S_2 condition, we get:

$$\omega_X \simeq \mathcal{O}_X(K_X).$$

In fact, if $\omega_X^{[m]}$ denotes the m th reflexive power of ω_X , the same arguments yield $\omega_X^{[m]} \simeq \mathcal{O}_X(mK_X)$. Therefore, the Weil divisor K_X is \mathbb{Q} -Cartier if and only if ω_X is a \mathbb{Q} -line bundle, i.e. $\omega_X^{[m]}$ is a line bundle for some $m > 0$.

Conductors and slc singularities. Let now X be a (reduced) scheme, and $\nu : X^\nu \rightarrow X$ its normalization. We recall that if X is not irreducible, its normalization is defined to be the disjoint union of the normalization of its irreducible components. The *conductor ideal*

$$\text{cond}_X := \mathcal{H}om_{\mathcal{O}_X}(\nu_*\mathcal{O}_{X^\nu}, \mathcal{O}_X)$$

is the largest ideal sheaf on X that is also an ideal sheaf on X^ν . If we think of the case where B is the integral closure of some integral ring A , then we can easily see that $\text{Hom}_A(B, A)$ injects in A (via the evaluation at 1), and the image of this map is the annihilator $\text{Ann}_A(B/A) = \{f \in A; fB \subset A\}$, or equivalently the largest ideal $\mathcal{I} \subset A$ that is also an ideal in B .

Coming back to the case of varieties, we will denote by cond_{X^ν} the conductor seen as an ideal sheaf on X^ν , and we define the conductor subschemes as $C_X := \text{Spec}_X(\mathcal{O}_X/\text{cond}_X)$ and $C_{X^\nu} := \text{Spec}_{X^\nu}(\mathcal{O}_{X^\nu}/\text{cond}_{X^\nu})$. If X is S_2 , then one can show that these schemes have pure codimension 1 (and hence define Weil divisors) but they are in general not reduced (e.g. the cusp $y^2 = x^3$).

If K_X is \mathbb{Q} -Cartier and X is demi-normal (i.e. X is S_2 and has only double crossing singularities in codimension 1, cf. [Kol]), we have the following relation:

$$\nu^*K_X = K_{X^\nu} + C_{X^\nu}. \quad (3.1)$$

The proof of this identity goes as follows: first, using the demi-normality assumption, we may assume that the only singularities of X are double normal crossings. Then, using the universal property of the dualizing sheaf (which coincide with the canonical sheaf as we observed above) and the projection formula, we have $\nu_*\omega_{X^\nu} = \omega_X(-C_X)$. We pull-back this relation to X^ν using the fact that the sheaf $\mathcal{O}_X(-C_X)$ becomes precisely $\mathcal{O}_{X^\nu}(-C_{X^\nu})$. By the assumptions on the singularities, the latter sheaf is actually an invertible sheaf so that we get the expected identity (cf. point 8 in [Kol]). As we will explain below, we do not want to assume a priori that our varieties are demi-normal. Therefore, it may happen that C_{X^ν} is not Cartier, and the formula

(3.1) may not be true anymore. So whenever we will deal with Kähler–Einstein on those varieties, we will have to apply the arguments on a log-resolution of the normalization instead of the normalization itself. Anyway, this will not cause any trouble.

An important point is that whenever the conductor is reduced, then necessarily X is seminormal (i.e. every finite morphism $X' \rightarrow X$ (with X' reduced) that is a bijection on points is an isomorphism); moreover, a seminormal scheme which is G_1 and S_2 has only double crossing singularities in codimension 1, i.e. it is deminormal. We refer to [Tra70, GT80, KSS10] for the previous assertions. This leads to the following definition:

DEFINITION 3.8. *We will say that X has semi-log canonical singularities if:*

1. X is G_1 and S_2 ,
2. K_X is \mathbb{Q} -Cartier,
3. The pair (X^ν, C_{X^ν}) is log-canonical.

If X has semi-log canonical singularities (slc), then C_{X^ν} is necessarily reduced, and therefore the codimension 1 singularities of X are only double crossing as we explained above. This assumption is usually added in the definitions (cf. [Kol, Kov12]), but we may keep it or not without any change. This justifies the seemingly different definition given in the previous section. Finally, we can give the definition of a stable variety:

DEFINITION 3.9. *We say that X is stable if*

1. X is projective,
2. X has semi-log canonical singularities,
3. K_X is \mathbb{Q} -ample.

Singularities and Kähler–Einstein metrics: a summary. If we take a closer look at the proof of Proposition 3.6, we see that we did not use all of the properties of a stable variety to see that a Kähler–Einstein metric always extend. Actually, we just used the fact that the conductor was a divisor. Therefore, using the existence and regularity results that we are going to prove in the next sections, and the restriction on the singularities of a pair carrying a Kähler–Einstein metric (cf. Proposition 6.1), we can summarize the problem of the existence of a Kähler–Einstein metric on a stable variety in the following statement:

Theorem 3.10. *Let X be a reduced n -equidimensional projective scheme over \mathbb{C} , satisfying the conditions G_1 and S_2 , and such that K_X is an ample \mathbb{Q} -line bundle. Then the following are equivalent:*

1. There exists a Kähler form ω on X_{reg} such that $\text{Ric } \omega = -\omega$ and $\int_{X_{\text{reg}}} \omega^n = c_1(K_X)^n$,
2. There exists ω as above which extends to define a positive current in $c_1(K_X)$,

3. X has semi-log canonical singularities, i.e. X is stable.
 Moreover, by the results of Odaka [Oda13b, Oda11], the latter condition is equivalent to:
4. The pair (X, K_X) is K -stable.

Proof. The proofs of these results lie at various places in this paper:

$1 \Leftrightarrow 2$ is a consequence of Proposition 3.6 as we mentioned above.

$1 \Rightarrow 3$ is the content of Proposition 6.1.

$3 \Rightarrow 1$ is the main result of this paper, obtained as a combination of the results in Sections 4 and 5. \square

4 Variational solutions

4.1 General setting. Consider the following general setting: X is a compact Kähler manifold and $[\omega]$ a big class, with ω smooth (but not necessarily positive). We say that a function $u \in PSH(X, \omega)$ has *full Monge–Ampère mass*, and we will write $u \in \mathcal{E}(X, \omega)$, if the total mass of $\text{MA}(u)$ is equal to the volume of the class $[\omega]$, where the volume in question may be defined by $V := \text{vol}([\omega]) := \int_X \text{MA}(u_{\min})$, for u_{\min} any element in $PSH(X, \omega)$ with minimal singularities, cf. Section 2.

We now recall an important subspace of $\mathcal{E}(X, \omega)$ denoted by $\mathcal{E}^1(X, \omega)$, and consisting of functions with finite energy. The energy $\mathcal{E}(u)$ of an ω -psh function u (not necessarily in $\mathcal{E}(X, \omega)$) is defined in the following way (cf. [GZ07, BEGZ10, BBGZ09] for more details—the energy is sometimes denoted by E in the aforementioned papers).

First, if $u \in PSH(X, \omega)$ has minimal singularities, we set

$$\mathcal{E}(u) := \frac{1}{(n+1)V} \sum_{j=0}^n \int_X (u - V_\theta) \text{MA}\left(u^{(j)}, V_\theta^{(n-j)}\right)$$

where MA is the mixed non-pluripolar Monge–Ampère operator. If now u is any ω -psh function, we defined

$$\mathcal{E}(u) := \inf\{\mathcal{E}(v) \mid v \in PSH(X, \omega) \text{ with minimal singularities, } v \geq u\}.$$

Then we set $\mathcal{E}^1(X, \omega) := \{u \in PSH(X, \omega), \mathcal{E}(u) > -\infty\}$. Actually, [BEGZ10, Proposition 2.11] gives another characterization of this last space: a function $u \in PSH(X, \omega)$ belongs to $\mathcal{E}^1(X, \omega)$ if and only if $u \in \mathcal{E}(X, \omega)$ and $\int_X (u - V_\theta) \text{MA}(u) < +\infty$ (and for any $u \in \mathcal{E}(X, \omega)$, the explicit integral formula for $\mathcal{E}(u)$ above is still valid). Using this result, it becomes clear that $\mathcal{E}^1(X, \omega) \subset \mathcal{E}(X, \omega)$ as announced.

We should finally add that \mathcal{E} is an upper-semicontinuous (usc) concave functional on $PSH(X, \omega)$, and that it is the normalized primitive of the Monge–Ampère operator, i.e.

$$(d\mathcal{E})_u = \frac{1}{V} \text{MA}(u)^n. \quad (4.1)$$

4.2 Uniqueness. Given a measure μ on X (possibly non-finite) we consider the following MA-equation for $u \in PSH(X, \omega)$ attached to the pair (ω, μ) :

$$\omega_u^n = e^u \mu, \tag{4.2}$$

where $\omega_u^n := MA(u)$ is the non-pluripolar Monge–Ampère operator as before. This equation is equivalent to the following *normalized* MA-equation on $\mathcal{E}(X, \omega)/\mathbb{R}$:

$$\frac{\omega_u^n}{V} = \frac{e^u \mu}{\int e^u \mu}, \tag{4.3}$$

The equivalence follows immediately from the \mathbb{R} -invariance of the latter equation and the substitution $u \mapsto u - \log \int e^u \mu$ which maps solutions of Equation (4.2) to solutions of the Equation (4.3).

PROPOSITION 4.1. *Any two solutions u and v of the MA-equation (4.2) such that u and v are in $\mathcal{E}(X)$ coincide.*

Proof. This is an immediate consequence of the comparison principle [BEGZ10, Corollary 2.3]: if u and v are in $\mathcal{E}(X)$ then

$$\int_{\{u < v\}} MA(v) \leq \int_{\{u < v\}} MA(u).$$

But the MA above then forces $u = v$ a.e w.r.t. the measure μ . Since μ cannot charge pluripolar sets (as $MA(u)$ does not) it follows that $u = v$ away from a pluripolar set and hence everywhere, by basic properties of psh functions. \square

4.3 Existence results for log canonical pairs. Let (X, D) be a log canonical pair such that the log canonical divisor $K_X + D$ is big. Assume that (X, D) is a log smooth, i.e. X is smooth and

$$D = \sum_i c_i D_i$$

is a normal crossing divisor with $c_i \in]-\infty, 1]$. To the pair (X, D) we can associate the following Kähler–Einstein type equation for a metric ϕ on $L := K_X + D$:

$$(dd^c \phi)^n = e^{\phi - \phi_D}, \tag{4.4}$$

where $\phi_D = \sum_i c_i \log |s_i|^2$ and s_i are sections cutting out the divisors D_i above.

Theorem 4.2. *There is a unique finite energy solution ϕ to the equation above.*

Proof. The proof is a modification of the variational approach in [BBGZ09] (concerning the case when D is trivial). To explain this we fix a smooth form $\omega \in c_1(K_X + D)$. Then the equation above is equivalent to a Monge–Ampère equation for an ω -psh function u :

$$\omega_u^n = e^u \mu, \tag{4.5}$$

where the measure μ is of the form $\mu = \rho dV$ for a function ρ in $L^{1-\delta}(X)$ (but ρ is not in $L^1(X)$!). We let

$$\mathcal{L}(u) := -\log \int e^u \mu.$$

Then, at least formally, solutions of Equation (4.5) are critical points of the functional

$$\mathcal{G}(u) := \mathcal{E}(u) + \mathcal{L}(u).$$

in view of the Equation (4.1) satisfied by \mathcal{E} . \mathcal{L} also defines an usc concave functional on $PSH(X, \omega)$ and we let $\mathcal{L}(X, \omega) := \{\mathcal{L} > -\infty\}$ (the upper semi-continuity follows from Fatou's lemma).

Note that Lemma 4.3 below guarantees that the intersection $\mathcal{E}^1(X, \omega) \cap \mathcal{L}(X, \omega)$ is non-empty. Hence, $\mathcal{G}(u)$ is not identically equal to $-\infty$ on its domain of definition that we will take to be $\mathcal{E}^1(X, \omega)$ (equipped with the usual $L^1(X)$ -topology).

Next, we observe that

$$\mathcal{G}(u) \leq \mathcal{E}(u) - \int u \mu_0 + C'' \tag{4.6}$$

Indeed, since $\mu \geq C\mu_0$, where μ_0 is finite measure on X integrating all quasi-psh functions on X (in our case we may take $\mu_0 = \|s'\|^2 dV$ for some holomorphic section s' defined by the negative coefficients of D):

$$\int e^u \mu \geq C \int e^u \mu_0$$

and hence

$$\mathcal{L}(u) \leq C' - \log \int e^u \mu_0 \leq C'' - \int u \mu_0$$

using Jensen's inequality, which proves (4.6). In particular, $\mathcal{G}(u)$ is bounded from above. Indeed, by scaling invariance we may assume that $\sup_X u = 0$ and then use that, by basic compactness properties of ω -psh functions, $\sup u \leq \int u \mu_0 + C$.

Let now $u_j \in \mathcal{E}^1(X, \omega)$ be a sequence such that

$$\mathcal{G}(u_j) \rightarrow \sup_{\mathcal{E}^1(X, \omega)} \mathcal{G} := S < \infty.$$

Again, by the scale invariance of \mathcal{G} we may assume that $\sup_X u_j = 0$. In particular,

$$\mathcal{L}(u_j) \geq S/2 - \mathcal{E}(u_j)$$

for $j > j_0$. But, by (4.6), $\mathcal{E}(u_j)$ is bounded from below and hence there is a constant C such that

$$\mathcal{E}(u_j) \geq -C, \quad \mathcal{L}(u_j) \geq -C.$$

Let now u_* be a limit point of u_j . By upper-semicontinuity we have that

$$\mathcal{E}(u) \geq -C, \quad \mathcal{L}(u) \geq -C.$$

Finally, we note that u_* satisfies the Equation (4.5) by applying the projection argument from [BBGZ09] as follows. Fixing $v \in C^\infty(X)$ let $f(t) := \mathcal{E}_\omega(P_\omega(u_* + tv)) + \mathcal{L}(u_* + tv)$, where

$$P_\omega(u)(x) := \sup\{v(x) : v \leq u, v \in PSH(X, \omega)\}$$

(note that $f(t)$ is finite for any t). The functional $\mathcal{L}(u)$ is decreasing in u and hence the sup of $f(t)$ on \mathbb{R} is attained for $t = 0$. Now $\mathcal{E}_\omega \circ P_\omega$ is differentiable with differential $MA(P_\omega u)$ at u [BBGZ09]. Hence, the condition $df/dt = 0$ for $t = 0$ gives that the variational equation (4.5) holds when integrated against any $v \in C^\infty(X)$. \square

Let us now prove the following result, that we used in the proof:

LEMMA 4.3. *Let (X, D) be a log smooth pair and L a big line bundle. Let θ be a smooth $(1, 1)$ form whose cohomology class is $c_1(L)$. Let s_0 be a section of D , and $|\cdot|$ a smooth hermitian metric on $\mathcal{O}_X(D)$. Then there exists a θ -psh function $u \in \mathcal{E}^1(X, \theta)$ such that $e^u/|s_0|^2$ is integrable.*

Proof. As L is big, there exists m big enough such that $mL - D$ is effective. We choose t a holomorphic section of $mL - D$, and consider $s := s_0 t$ which is a section of mL vanishing along D . Let h_0 be a smooth hermitian metric on L with curvature form θ , and let V_θ be the upper envelope of all (normalized) θ -psh functions. We define on mL the hermitian metric $h := h_0^{\otimes m} e^{-mV_\theta}$. For $0 < \alpha < 1$ small enough we claim that the function

$$u := V_\theta - \left(-\frac{1}{m} \log |s|_{h_0}^2 \right)^\alpha$$

suits our requirements.

First of all, it is θ -psh because of the following general fact: if ψ is θ -psh and $\chi : \mathbb{R} \rightarrow \mathbb{R}$ is convex and non-decreasing satisfying $\chi' \leq 1$, then $V_\theta + \chi(\psi - V_\theta)$ is θ -psh. Indeed, $dd^c(V_\theta + \chi(\psi - V_\theta)) = (1 - \chi')(\theta + dd^c V_\theta) + \chi'(\theta + dd^c \psi) + \chi''|d(\psi - V_\theta)|^2$ where χ' and χ'' are evaluated at $\psi - V_\theta$. Now we apply this to $\psi = 1/m \log |s|_{h_0}^2$.

For the integrability property, we use the following inequality for x a real number (big enough): $x^\alpha \geq (n + 1) \log x - C$ for some $C > 0$ depending only on α . Now we observe that $V_\theta + \chi(\psi - V_\theta) \leq \chi(\psi)$: indeed, $\chi(\psi - V_\theta) - \chi(\psi) \leq \sup \chi' \cdot (-V_\theta) \leq -V_\theta$, so that in our case, $u \leq \left(-\frac{1}{m} \log |s|_{h_0}^2 \right)^\alpha$. If we apply the basic inequality stated above to $x = -\frac{1}{m} \log |s|_{h_0}^2$ which can be made big enough by multiplying h_0 by a big constant (this does not change the curvature), we get

$$e^u \leq C \left(-\frac{1}{m} \log |s|_{h_0}^2 \right)^{-(n+1)}.$$

As D has snc support, and $|t|$ is bounded from above, we are left to check that the integral

$$\int_D \frac{dV}{\prod_{i \leq n} |z_i|^2 \cdot \log^{n+1}(\prod_{i \leq n} |z_i|^2)}$$

over the unit polydisc D in \mathbb{C}^n converges. But after a polar change of coordinate, we are led to estimate $\int_{[0,1]^n} \frac{dx_1 \cdots dx_n}{\prod_{i \leq n} x_i \cdot \log^{n+1}(\prod_{i \leq n} x_i^2)}$, which equals $\frac{1}{2^{n+1}n} \int_{[0,1]^{n-1}} \frac{dx_1 \cdots dx_{n-1}}{\prod_{i \leq n-1} x_i \cdot \log^n(\prod_{i \leq n-1} x_i)}$. By induction, and using the Poincaré case, it concludes.

Finally, one has to check that $u \in \mathcal{E}^1(X, \theta)$. We compute the capacity $\text{Cap}_\theta(u < V_\theta - t)$ for t big. But $(u < V_\theta - t) = (\frac{1}{m} \log |s|_h^2 < -t^{1/\alpha}) \subset ((\frac{1}{m} \log |s|_{h_0}^2 < -t^{1/\alpha}))$ and thus $\text{Cap}_\theta(u < V_\theta - t) \leq \frac{C}{t^{1/\alpha}}$ because for every θ -psh function ψ , one has $\text{Cap}_\theta(\psi < -t) \leq \frac{C_\psi}{t}$ (this is an easy generalization of [GZ05, Proposition 2.6]). Therefore, if $\alpha < \frac{1}{n+1}$, one has

$$\int_0^{+\infty} t^n \text{Cap}_\theta(u < V_\theta - t) dt < +\infty$$

which, using the characterization given in [BBGZ09, Lemma 2.9], ends the proof of the lemma. \square

REMARK 4.4. The proof of the preceding lemma yields actually a stronger result. If $\sum a_i \text{div}(s_i)$ is an effective divisor with snc support meeting D transversally and such that $a_i < 1$ for all i , then the function u obtained above satisfies $e^u / \prod |s_i|^{2a_i} |s_0|^2 \in L^1(dV)$, and more generally this is still true for $e^{\varepsilon u}$ for all $\varepsilon > 0$ (use the inequality $\varepsilon x^\alpha \geq (n+1) \log x - C$ for $x = -\frac{1}{m} \log |s|_{h_0}^2$ this time).

4.4 Stability under perturbations. Let now L be a semipositive and big line bundle, and consider the perturbed ample line bundles $L_j := L + \varepsilon_j A$, for ε_j a sequence of positive numbers tending to 0 and A a fixed ample line bundle. Fixing also a Kähler form $\omega_A \in c_1(A)$ and a smooth semipositive form $\omega \in c_1(L)$, we write $\omega_j := \omega + \varepsilon_j \omega_A$. Let μ_j be the sequence of measures on X given by

$$\mu_j = \prod_\alpha (|s_\alpha|^2 + \varepsilon_j)^{e_\alpha} \frac{dV}{\prod_\beta |s_\beta|^2}$$

where $e_\alpha > -1$ for all α , and the divisor $\sum_\alpha \text{div}(s_\alpha) + \sum_\beta \text{div}(s_\beta)$ is a reduced normal crossing divisor. This is precisely the sequence of approximations we are going to use to solve our Kähler–Einstein equation.

Consider now the following Monge–Ampère equations for $u_j \in \mathcal{E}(X, \omega_j)$ (and sup-normalized):

$$\omega_{u_j}^n / V = \frac{e^{u_j} \mu_j}{\int_X e^{u_j} \mu_j}$$

and similarly

$$\omega_u^n/V = \frac{e^u \mu}{\int_X e^u \mu}$$

for $u \in \mathcal{E}(X, \omega)$.

Theorem 4.5. *The unique sup-normalized solution u_j of the first equation above converges, in the $L^1(X)$ -topology, to the unique sup-normalized solution u to the latter equation. Equivalently, the solutions v_j of the corresponding non-normalized equations converge in $L^1(X)$ to v solving the corresponding limiting non-normalized equation.*

Proof. We denote by \mathcal{G}_j (resp. \mathcal{L}_j) the functional determined by the pair (ω_j, μ_j) (resp. μ_j), and by u_j the sup-normalized maximizer of \mathcal{G}_j . We also denote by u_0 the sup-normalized fixed ω -psh function given by Lemma 4.3. Let us add that in the course of the proof, the precise value of the constant C may, as usual, change from line to line. We split the proof into four steps.

Step 1. We first show that

$$-C \leq \mathcal{G}_j(u_j) \leq C. \tag{4.7}$$

As u_0 is ω -psh, it is also ω_j -psh. Moreover, the capacity computation of Lemma 4.3 shows that the energy of u_0 with respect to ω_j is finite, and as $\mathcal{E}_{\omega_j}(u_0)$ increases with j , we obtain

$$\mathcal{E}_{\omega_j}(u_0) \geq -C.$$

Besides, by dominated convergence, we have $\lim_{j \rightarrow +\infty} \mathcal{L}_j(u_0) = \mathcal{L}(u_0)$ and therefore we get $\mathcal{L}_j(u_0) \geq -C$. Consequently, $\mathcal{G}_j(u_j) \geq \mathcal{G}_j(u_0) \geq -C$ which gives a first bound (recall that u_j maximize \mathcal{G}_j by Theorem 4.2 and the translation invariance of \mathcal{G}_j).

Choose now a probability measure μ_0 satisfying $\mu_j \geq e^{-C} \mu_0$ for all j (its existence is clear given the precise form of μ_j). Then Jensen's inequality gives

$$\mathcal{L}_j(u_j) \leq - \int_X u_j d\mu_0 - C$$

but the compactness properties of quasi-psh functions (all u_j 's are $C\omega_A$ -psh) also gives the inequality

$$\sup u_j = 0 \leq \int_X u_j d\mu_0 + C.$$

Combining the two previous inequalities, we get

$$\mathcal{G}_j(u_j) \leq \mathcal{E}_{\omega_j}(u_j) + C$$

which gives both the uniform upper bound for $\mathcal{G}_j(u_j)$ (as \mathcal{E}_{ω_j} is always non-positive) and a uniform lower bound $\mathcal{E}_{\omega_j}(u_j) \geq -C$.

Let u be an L^1 -limit point in $PSH(X, \omega)$ of the sequence (u_j) .

Step 2. We next show that

$$\mathcal{G}(u) \geq \limsup \mathcal{G}_j(u_j). \quad (4.8)$$

First by Fatou's lemma, we have

$$\mathcal{L}(u) \geq \limsup \mathcal{L}_j(u_j).$$

Moreover

$$\mathcal{E}(u) \geq \limsup \mathcal{E}_{\omega_j}(u_j)$$

as follows from Lemma 4.6 below. Putting these two inequalities together gives the desired bound.

Step 3. u is a sup-normalized maximizer of \mathcal{G} .

For any given sup-normalized ω -psh function v , we need to show that

$$\mathcal{G}(u) \geq \mathcal{G}(v). \quad (4.9)$$

Of course, one can assume that $\mathcal{G}(v)$ is finite. Thanks to step 2, it is enough to show that $\limsup \mathcal{G}_j(v) \geq \mathcal{G}(v)$. But this inequality is far from clear as we cannot directly apply the dominated convergence theorem here. Indeed, for the energy part, it could happen that $\mathcal{E}_\omega(v)$ is finite though $\mathcal{E}_{\omega_j}(v) = -\infty$ for all j . As for the other part, despite $e^v \in L^1(\mu)$, it is not obvious that $e^v \in L^1(\mu_j)$ (because of the “zeroes” of μ which do not appear in μ_j).

To bypass these difficulties we will use a regularization/perturbation argument. More precisely, we pick a family of smooth ω -psh functions $(v_\delta)_{\delta>0}$ which decrease to v , and we set for all positive δ, ε :

$$v_{\delta,\varepsilon} := (1 - \varepsilon)v_\delta + \varepsilon u_0$$

where we recall that u_0 denotes the particular (sup-normalized) ω -psh function constructed in Lemma 4.3.

As v_δ is smooth and $u_0 \in \mathcal{E}^1(X, \omega_j)$ for all j , $v_{\delta,\varepsilon}$ has finite ω_j -energy, the dominated convergence theorem shows that

$$\lim_{j \rightarrow +\infty} \mathcal{E}_{\omega_j}(v_{\delta,\varepsilon}) = \mathcal{E}_\omega(v_{\delta,\varepsilon}). \quad (4.10)$$

Moreover, as we observed in Remark 4.4, the function $e^{\varepsilon u_0}$ is in $L^1(\mu)$ for all $\varepsilon > 0$, and $e^{v_{\delta,\varepsilon}} \leq e^{\varepsilon u_0}$. Therefore, by dominated convergence, we get

$$\lim_{j \rightarrow +\infty} \mathcal{L}_j(v_{\delta,\varepsilon}) = \mathcal{L}(v_{\delta,\varepsilon}). \quad (4.11)$$

Combining (4.8) with (4.10) and (4.11), we get $\mathcal{G}(u) \geq \mathcal{G}(v_{\delta,\varepsilon})$ for all $\delta, \varepsilon > 0$. Set $v_\varepsilon := (1 - \varepsilon)v + \varepsilon u_0$. By monotonicity of \mathcal{E}_ω , we have $\mathcal{E}_\omega(v_{\delta,\varepsilon}) \geq \mathcal{E}_\omega(v_\varepsilon)$. Using the

dominated convergence theorem, we also see that $\mathcal{L}(v_{\delta,\varepsilon}) \rightarrow \mathcal{L}(v_\varepsilon)$. Therefore, we have $\mathcal{G}(u) \geq \mathcal{G}(v_\varepsilon)$. Finally, using the concavity of \mathcal{G} , we get $\mathcal{G}(u) \geq (1 - \varepsilon)\mathcal{G}(v) + \varepsilon\mathcal{G}(u_0)$, and we get (4.9) by letting ε go to zero.

Step 4. Back to the non-normalized equation. We have $v_j = u_j + \mathcal{L}_j(u_j)$ and as shown above in (4.7), $\mathcal{G}_j(u_j)$ is a bounded sequence (more precisely, it converges to the maximal value S of \mathcal{G}) and $0 \leq -\mathcal{E}(u_j) \leq C$, which implies that $\mathcal{L}_j(u_j)$ is also a bounded sequence. After passing to a subsequence we may thus assume that $\mathcal{L}_j(u_j) \rightarrow l \in \mathbb{R}$ so that $v_j \rightarrow v := u + l$, solving the desired equation [and $v \in \mathcal{E}^1(X, \omega)$]. By the uniqueness of solutions of the latter equation this means that the whole sequence u_j converges to v , which concludes the proof. \square

Let us now give the proof of the following result which was essential for step 2:

LEMMA 4.6. *Let $[\omega_j]$ and $[\omega]$ be semi-positive big classes such that $\omega_j \rightarrow \omega$ in the C^∞ -topology of smooth (semipositive) forms. If $u_j \in \mathcal{E}(X, \omega_j)$ (and $u \in \mathcal{E}(X, \omega)$) such that $u_j \rightarrow u$ in $L^1(X)$, then*

$$\mathcal{E}_\omega(u) \geq \limsup_j \mathcal{E}_{\omega_j}(u_j).$$

Proof. When $\omega_j = \omega$ the lemma amounts to the well-known fact that \mathcal{E}_ω is usc. We may as well assume that $\sup u_j = \sup u = 0$.

First of all, we modify the sequence (u_j) to make it non-increasing. More precisely, we set $\tilde{u}_j := (\sup_{k \geq j} u_k)^*$, which defines an ω_j -psh function. Then $\tilde{u}_j \geq u_j$ and the sequence $(\tilde{u}_j)_j$ is non-increasing. Given v an ω -psh function and $c \in \mathbb{R}$, we will write $v^c := \max\{v, c\}$. By construction \tilde{u}_j^c decreases to u^c , and all these functions are ω_0 -psh. By the local convergence result of Bedford–Taylor for mixed Monge–Ampère expressions and the smooth convergence of ω_j to ω , we see that

$$\mathcal{E}_\omega(u^c) = \lim_{j \rightarrow +\infty} \mathcal{E}_{\omega_j}(\tilde{u}_j^c).$$

As $u_j \leq \tilde{u}_j \leq \tilde{u}_j^c$, the monotonicity of \mathcal{E}_ω ensures that

$$\mathcal{E}_\omega(u^c) = \limsup_{j \rightarrow +\infty} \mathcal{E}_{\omega_j}(u_j).$$

Taking the infimum over all c and using the definition of the functional \mathcal{E}_ω , we obtain the desired inequality. \square

5 Smoothness of the Kähler–Einstein metric

Before we go into the details of the proof of the regularity theorem, we would like to give an overview of previous related results and underline the main differences that will appear in our specific case, namely the case of general log canonical pairs. As we will rely on the so called logarithmic case (i.e. (X, D) is log smooth, D is reduced, and $K_X + D$ is ample), the next section will be devoted to recall some of the main tools appearing in this setting. Then, we will give the proof of the main regularity theorem, which will constitute the core of this section.

5.1 Special features in the log canonical case. We should first mention the case of varieties with log terminal singularities, or more generally klt pairs, which correspond to the pairs where the discrepancies a_i defined earlier satisfy the strict inequalities $a_i > -1$. Then the situation is relatively well understood: In the non-positively curved case, we know that the Kähler–Einstein metric exists, is unique, has bounded potential, and induces on the regular locus a genuine Kähler–Einstein metric (see e.g. [BEGZ10,EGZ08,EGZ09,DP10]). As for the case of positive curvature, or log Fano manifolds, then there exist criteria (like the properness of the Mabuchi functional) to guarantee the existence and uniqueness (modulo automorphisms of X) of a Kähler–Einstein metric [BBEGZ11]; this metric is also known to have bounded potential and to be smooth on the regular locus of X (see again [BBEGZ11,Pău08]).

However, the behavior of the Kähler–Einstein metric near the singularities of X is mostly unknown (except if the singularities are orbifold). In the case of a klt pair, we know that the metric will not be smooth along the divisor, but its singularities can sometimes be understood outside of the singular part of (X, D) . For example, a recent result in this direction states that the Kähler–Einstein metric has cone singularities near each point where (X, D) is log smooth, i.e. X is smooth and D has simple normal crossing support (cf. [Gue13,GP13]).

When (X, D) is now a log smooth pair, the situation gets easier because there is no more loss of positivity coming from the resolution of singularities. For example, if the coefficients of D are in $[0, 1)$ (the pair is then klt), the Kähler–Einstein metric is known to have cone singularities along D , as it was proved by [GP13] in full generality, and by [Bre11,CGP13,JMR11] under some assumptions on D .

When now every coefficient of D is equal to 1, and $K_X + D$ is ample, then we know from the work of Kobayashi [Kob84] and Tian–Yau [TY87] that there exists a unique complete Kähler–Einstein metric having Poincaré singularities along D . The situation where the coefficients of D are in $]0, 1]$ behaves like a product of cone and Poincaré geometries and was studied in [Gue12,GP13].

In a slightly different direction, Tsuji [Tsu88a] has considered the case of a singular variety with ample canonical line bundle such that only one divisor appears in its resolution, with discrepancy equal to -1 . Finally, Wu [Wu08,Wu09] has worked out the case of a quasi-projective manifold compactified by a snc divisor $\sum D_i$ such that $K_X + \sum a_i D_i$ is ample for some coefficients $a_i \geq -1$. In our case however, such a strong positivity assumption will never happen as soon as we have to perform a non-trivial resolution.

As one can already observe in the log smooth case studied by Kobayashi and Tian–Yau, the log canonical case is very different from the klt case. Let us mention some striking divergences: first of all, the potentials are no more bounded even in the ample case so that the solution does not have minimal singularities. Moreover, the Kähler–Einstein equation in this setting is closely related to a negative curvature geometry. Indeed, if we first consider the Ricci-flat case, then it is impossible to

write the equation on the whole X . Indeed, the current solution obtained on X_{reg} will not have finite mass near the singularities, and hence it will not extend as a global positive current on X . This phenomenon already happens in [TY90]. Finally, it has been proven in [BBEGZ11, Proposition 3.8] that any pair (X, D) with X normal and $-(K_X + D)$ ample admitting a Kähler metric ω on X_{reg} with continuous potentials solution of $\text{Ric}\omega = \omega + [D]$ is necessarily klt. Therefore it is pointless to look for positively curved Kähler–Einstein metric in the general setting of log canonical pairs instead of klt pairs.

To finish this discussion, let us stress the fact that the class of varieties with semi-log canonical singularities can be realized as a subclass of log canonical pairs (cf. definition 3.8). This is the largest “reasonable” class to look for Kähler–Einstein metrics: for example, if X is a smooth Fano manifold carrying a smooth divisor $D \in |-K_X|$, then for any $\varepsilon > 0$, one has $K_X + (1 + \varepsilon)D > 0$; however, there is no smooth Kähler–Einstein metric with negative scalar curvature on $X \setminus D$. Indeed, its existence would contradict the Yau–Schwarz lemma applied with the complete Ricci-flat Kähler metric constructed in [TY90].

Moreover, we will see that the existence of a negatively curved Kähler–Einstein metric on the regular part of a normal projective variety with maximal volume forces the singularities to be at worst log canonical, cf. Proposition 6.1.

5.2 The logarithmic case. In this section, we will briefly recall the Theorem of Kobayashi and Tian–Yau constructing negatively-curved Kähler–Einstein metrics on quasi-projective varieties $X \setminus D$ where D is a reduced divisor with simple normal crossings, and $K_X + D$ is ample. In the course of the proof of Theorem 5.6, we will use in an essential manner the functional spaces introduced by these authors, namely the “quasi-coordinates” version of the usual Hölder spaces $\mathcal{C}^{k,\alpha}$. For now, X_0 will denote $X \setminus D$.

DEFINITION 5.1. *We say that a Kähler metric ω on X_0 is of Carlson–Griffiths type if there exists a Kähler form ω_0 on X such that $\omega = \omega_0 - \sum_K dd^c \log \log \frac{1}{|s_k|^2}$.*

In [CG72], Carlson and Griffiths introduced such a metric for some $\omega_0 \in c_1(K_X + D)$, but one can easily see that such a metric always exists on a Kähler manifold without assumptions on the bundle $K_X + D$. One can also observe that the existence of such a metric ω forces the cohomology class $\{\omega\}$ to be Kähler by Demailly’s regularization theorem [Dem82, Dem92].

The reason why we exhibit this particular class of Kähler metrics on X_0 having Poincaré singularities along D is that we have an exact knowledge on its behaviour along D , which is much more precise than knowing its membership in the aforementioned class. This is precisely the class in which one will look for a Kähler–Einstein metric, so that one needs to define the appropriate analogue of the usual Hölder spaces $\mathcal{C}^{k,\alpha}$. And to do so, one may (almost) boil down to the usual euclidean situation.

The key point is that (X_0, ω) has bounded geometry at any order. Let us get a bit more into the details. To simplify the notations, we will assume that D is irreducible, so that locally near a point of D , X_0 is biholomorphic to $\mathbb{D}^* \times \mathbb{D}^{n-1}$, where \mathbb{D} (resp. \mathbb{D}^*) is the unit disc (resp. punctured disc) of \mathbb{C} . We want to show that, roughly speaking, the components of ω in some appropriate coordinates have bounded derivatives at any order. The right way to formalize it consists in introducing quasi-coordinates: they are maps from an open subset $V \subset \mathbb{C}^n$ to X_0 having maximal rank everywhere. So they are just locally invertible, but these maps are not injective in general.

To construct such quasi-coordinates on X_0 , we start from the universal covering map $\pi : \mathbb{D} \rightarrow \mathbb{D}^*$, given by $\pi(w) = e^{\frac{w+1}{w-1}}$. Formally, it sends 1 to 0. The idea is to restrict π to some fixed ball $B(0, R)$ with $1/2 < R < 1$, and compose it (at the source) with a biholomorphism Φ_η of \mathbb{D} sending 0 to η , where η is a real parameter which we will take close to 1. If one wants to write a formula, we set $\Phi_\eta(w) = \frac{w+\eta}{1+\eta w}$, so that the quasi-coordinate maps are given by $\Psi_\eta = \pi \circ \Phi_\eta \times \text{Id}_{\mathbb{D}^{n-1}} : V = B(0, R) \times \mathbb{D}^{n-1} \rightarrow \mathbb{D}^*$, with $\Psi_\eta(v, v_2, \dots, v_n) = \left(e^{\frac{1+\eta}{1-\eta} \frac{v+1}{v-1}}, v_2, \dots, v_n \right)$.

Once we have said this, it is easy to see that X_0 is covered by the images $\Psi_\eta(V)$ when η goes to 1, and for all the trivializing charts for X , which are in finite number. Now, an easy computation shows that the derivatives of the components of ω with respect to the v_i 's are bounded uniformly in η . This can be thought as a consequence of the fact that the Poincaré metric is invariant by any biholomorphism of the disc.

At this point, it is natural to recall the definition of the Hölder space of $\mathcal{C}_{qc}^{k,\alpha}$ -functions on X_0 introduced by Cheng–Yau [CY80]:

DEFINITION 5.2. *For a non-negative integer k , a real number $\alpha \in]0, 1[$, we define:*

$$\mathcal{C}_{qc}^{k,\alpha}(X_0) = \left\{ u \in \mathcal{C}^k(X_0); \sup_{V,\eta} \|u \circ \Psi_\eta\|_{k,\alpha} < +\infty \right\}$$

where the supremum is taken over all our quasi-coordinate maps V (which cover X_0). Here $\|\cdot\|_{k,\alpha}$ denotes the standard $\mathcal{C}_{qc}^{k,\alpha}$ -norm for functions defined on a open subset of \mathbb{C}^n .

The following fact, though easy, is very important (see e.g. [Kob84] or [Gue12, Lemma 1.6] for a detailed proof):

LEMMA 5.3. *Let ω be a Carlson–Griffiths type metric on X_0 , and ω_0 some Kähler metric on X . Then*

$$F_0 := \log \left(\prod |s_k|^2 \log^2 |s_k|^2 \cdot \omega^n / \omega_0^n \right)$$

belongs to the space $\mathcal{C}_{qc}^{k,\alpha}(X_0)$ for every k and α .

Finding the Kähler–Einstein metric consists then in showing that the Monge–Ampère equation $(\omega + dd^c \varphi)^n = e^{\varphi+f} \omega^n$ has a unique solution $\varphi \in \mathcal{C}_{qc}^{k,\alpha}(X_0)$

for all functions $f \in \mathcal{C}_{qc}^{k,\alpha}(X_0)$ with $k \geq 3$. This can be done using the continuity method in the quasi-coordinates. In particular, applying this to $f = F := -\log(\prod |s_k|^2 \log^2 |s_k|^2 \cdot \omega^n / \omega_0^n) + (\text{smooth terms on } X)$, which the previous lemma allows to do, this will prove the existence of a negatively curved Kähler–Einstein metric, which is equivalent to ω (in the *strong* sense: $\varphi \in \mathcal{C}_{qc}^{k,\alpha}(X_0)$ for all k, α).

In this continuity method, one needs to obtain first uniform estimates; they follow from a consequence of Yau’s maximum principle for complete manifolds which we recall here (see [CY80, Proposition 4.1]):

PROPOSITION 5.4. *Let (X, ω) be a n -dimensional complete Kähler manifold, and $F \in \mathcal{C}^2(X)$ a bounded function. We assume that we are given $u \in \mathcal{C}^2(X)$ satisfying $\omega + dd^c u > 0$ and*

$$(\omega + dd^c u)^n = e^{u+F} \omega^n.$$

Suppose that the bisectional curvature of (X, ω) is bounded below by some constant, and that u is bounded. Then

$$\inf_X u \geq -\sup_X F \quad \text{and} \quad \sup_X u \leq -\inf_X F.$$

There are similar results for the Laplacian estimates, but as we will not use them directly, we do not state them here. To summarize the discussion, one obtains:

Theorem 5.5. (Kobayashi [Kob84], Tian–Yau [TY87]). *Let X be a compact Kähler manifold, D a reduced divisor with simple normal crossings, ω a Kähler form of Carlson–Griffiths type on $X \setminus D$, and $F \in \mathcal{C}_{qc}^{k,\alpha}(X \setminus D)$ for some $k \geq 3$. Then there exists $\varphi \in \mathcal{C}_{qc}^{k,\alpha}(X \setminus D)$ solution to the following equation:*

$$(\omega + dd^c \varphi)^n = e^{\varphi+F} \omega^n.$$

In particular if $K_X + D$ is ample, then there exists a (unique) Kähler–Einstein metric of curvature -1 equivalent to ω .

5.3 Statement of the regularity theorem. In this section, we prove that the Kähler–Einstein metric attached to a log canonical pair (X, D) (satisfying $K_X + D$ ample) by Theorem 4.2 is smooth on $X_0 = X_{\text{reg}} \setminus \text{Supp}(D)$. As usual, we will work on a log resolution $\pi : X' \rightarrow X$, where:

$$K_{X'} = \pi^*(K_X + D) + \sum a_i E_i,$$

E_i being either an exceptional divisor or a component of the strict transform of D , and the coefficients a_i (called discrepancies) satisfy the inequality $a_i \geq -1$.

The Kähler–Einstein metric is given on X' by a (singular) psh weight ϕ on $\pi^*(K_X + D)$ satisfying

$$(dd^c \phi)^n = e^{\phi + \sum a_i \phi_{E_i}}$$

where ϕ_{E_i} is a psh weight on $\mathcal{O}_{X'}(E_i)$ such that $dd^c\phi_{E_i} = [E_i]$. So if in local coordinates, E_i is given by $\{z_n = 0\}$, then $\phi_{E_i} = \log|z_n|^2$.

Our aim is to obtain regularity properties for the solutions of degenerate Monge–Ampère equations like the previous one; this is the content of the following theorem:

Theorem 5.6. *Let X be a compact Kähler manifold of dimension n , dV some volume form, $D = \sum a_i D_i$ a \mathbb{R} -divisor with coefficients in $(-\infty, 1]$ and defining sections s_i , $E = \sum c_\alpha E_\alpha$ an effective \mathbb{R} -divisor such that $D_{red} + E$ has snc support, and θ a semipositive form with $\int_X \theta^n > 0$ such that $\{\theta\} - c_1(E)$ is a Kähler class. Then the θ -psh function φ with full Monge–Ampère mass, which is a solution of*

$$\langle (\theta + dd^c\phi)^n \rangle = \frac{e^\varphi dV}{\prod_i |s_i|^{2a_j}}$$

is smooth outside of $\text{Supp}(D) \cup \text{Supp}(E)$.

Note that although φ has full Monge–Ampère mass, it is in general far from having minimal singularities as soon as some coefficient a_i of D equals 1. Think for example of the logarithmic case (a log smooth pair (X, D) where $K_X + D$ is ample; then the potential of the Kähler–Einstein metric is not bounded whereas the class is ample.

Let us go back to the general Kähler–Einstein case. We would like to apply the previous results with E being some positive combination of the E_i 's. The problem is that there might be no such divisors; for example if π happens to be a small resolution, its exceptional locus has codimension at least 2. Therefore we need to perform another modification.

On X' , $\pi^*(K_X + D)$ is no more ample, and by [BBP10, Proposition 1.5], its augmented base locus is $\mathbb{B}_+(\pi^*(K_X + D)) = \pi^{-1}(\mathbb{B}_+(K_X + D)) \cup \text{Exc}(\pi) = \text{Exc}(\pi)$, and lies above $X_{\text{sing}} \cup \text{Supp}(D)$. It is well-known that one can find a log resolution $\mu : X'' \rightarrow X'$ of $(X', \mathbb{B}_+(\pi^*(K_X + D)))$, and an effective \mathbb{Q} -divisor F with snc support lying above $\mathbb{B}_+(\pi^*(K_X + D))$ and such that $\mu^*\pi^*(K_X + D) - F$ is ample. Moreover one can also assume that $F + \sum E'_i$ has snc support, where E'_i denotes the strict transform of E_i by μ .

Let us recall the argument. We start by resolving the singularities of a Kähler current $T \geq \omega$ (ω a Kähler form on X') in $\pi^*(K_X + D)$ computing $\mathbb{B}_+(\pi^*(K_X + D))$, then we write Siu's decomposition $\mu^*T = \theta + [D]$ with θ semi-positive dominating $\mu^*\omega$, and D lying above $\mathbb{B}_+(\pi^*(K_X + D))$. Finally, we choose a μ -exceptional \mathbb{Q} -divisor G such that $-G$ is μ -ample; it exists because μ is a finite composition of blow-ups with smooth centers. Then it becomes clear that for $\varepsilon > 0$ small enough, $\{\mu^*\theta\} - \varepsilon G$ is a Kähler class, and we have $\mu^*\pi^*(K_X + D) = (\{\mu^*\theta\} - \varepsilon G) + (\varepsilon G + D)$, with $\varepsilon G + D$ lying above $\mathbb{B}_+(\pi^*(K_X + D))$ and having simple normal crossing support. If one had chosen a log resolution of the ideal sheaf generated by the augmented base locus of $\pi^*(K_X + D)$ and the $\mathcal{O}'_X(E_i)$, we would have obtained the refined result that $F + E'$ has snc support.

Set $\nu := \pi \circ \mu : X'' \rightarrow X$, and write $K_{X''} = \nu^*(K_X + D) + E_\nu$. We know that E_ν is a divisor with snc support and coefficients ≥ -1 , and by the construction above, there exists a snc divisor F on X'' lying above $X_{\text{sing}} \cup \text{Supp}(D)$ such that $F + (E_\nu)_{\text{red}}$ has snc support and $\nu^*(K_X + D) - F$ is ample. Applying Theorem 5.6, we get:

Corollary 5.7. *Let (X, D) be a log canonical pair such that $K_X + D$ is ample. Then the Kähler–Einstein metric ω_{KE} on (X, D) is smooth on $X_{\text{reg}} \setminus \text{Supp}(D)$.*

As we shall see in the course of the proof (cf. Section 5.5.2), we do not obtain very precise estimates on the potential of the solution, even at order zero. However, it is tempting to believe that the potential ϕ_{KE} of the Kähler–Einstein metric should be locally bounded outside of the non-klt locus of (X, D) defined as the support of the sheaf $\mathcal{O}_X/\mathcal{I}(X, D)$ where $\mathcal{I}(X, D)$ is the multiplier ideal sheaf of (X, D) (cf. e.g. [Laz04]). However, as this locus cannot be read easily on some log resolution, it does not seem obvious how one should tackle this question.

5.4 Preliminaries: the regularized equation. We now borrow the notations of Theorem 5.6, and we let ω_0 be a Kähler form on X ; it will be our reference metric in the following. Recall that we want to solve the equation

$$\text{MA}(\varphi) = \frac{e^\varphi dV}{\prod_i |s_i|^{2a_i}}$$

where the unknown function is φ a θ -psh function, s_i are non-zero sections of $\mathcal{O}_X(D_i)$, $|\cdot|_i$ are smooth hermitian metrics on $\mathcal{O}_X(D_i)$, $f \in \mathcal{C}^\infty(X)$ and dV is a smooth volume form on X . Moreover, the expression $\text{MA}(\varphi)$ has to be understood as the non-pluripolar Monge–Ampère operator. It will be convenient for the following to differentiate the “klt part” of D from its “lc part”, so we introduce the following notation:

$$D = \underbrace{\sum_{a_j < 1} a_j D_j}_{D_{\text{klt}}} + \underbrace{\sum_{a_k = 1} D_k}_{D_{\text{lc}}}.$$

By Theorem 4.5, we know that the solution is the limit of any sequence of solutions of some appropriate regularized equations. The regularization process we are going to use concerns both the a priori non-Kähler class $\{\theta\}$ and the “klt part” in the volume form: $\prod_{a_j < 1} |s_j|^{-2a_j}$. More concretely, we will be studying the following equation:

$$\langle (\theta + t\omega_0 + dd^c \varphi_{t,\varepsilon})^n \rangle = \frac{e^{\varphi_{t,\varepsilon} + f} dV}{\prod_{a_j < 1} (|s_j|^2 + \varepsilon^2)^{a_j} \prod_{a_k = 1} |s_k|^2}. \tag{5.1}$$

Smoothness of the regularized solution. At this point, it is still not completely clear that the solution $\varphi_{t,\varepsilon}$ of Equation (5.1) is smooth on $X \setminus D_{\text{lc}}$. So we translate our equation into the logarithmic setting : we set

$$\omega_{t,lc} := \theta + t\omega_0 - \sum_{a_k=1} dd^c \log(\log |s_k|^2)^2.$$

We may choose the hermitian metrics $|\cdot|_k$ such that $|s_k| < 1$ and such that $\omega_{t,lc}$ defines a Kähler metric on $X \setminus D_{lc}$ (cf. [CG72, Gri76] e.g.). Of course this rescaling will depend on t , but we will explain how to bypass this problem later.

So using this new reference metric, one may rewrite Equation (5.1) in the following form:

$$(\omega_{t,lc} + dd^c \psi_{t,\varepsilon})^n = \frac{e^{\psi_{t,\varepsilon} + f_t} \omega_{t,lc}^n}{\prod_{a_j < 1} (|s_j|^2 + \varepsilon^2)^{a_j}}$$

where $\psi_{t,\varepsilon} = \varphi_{t,\varepsilon} + \sum_{a_k=1} \log(\log |s_k|^2)^2$ and $f_t = -\log\left(\frac{\prod_k |s_k|^2 \log^2 |s_k|^2 \omega_{t,lc}^n}{dV}\right)$. Clearly, f_t is bounded (but only the lower bound is uniform in t) and smooth on $X \setminus D_{lc}$, but we know by Lemma 5.3 that f_t is smooth when read in the quasi-coordinates adapted to the pair (X, D_{lc}) . Therefore, using the Theorem of Kobayashi and Tian–Yau (see Theorem 5.5), we know that the solution $\psi_{t,\varepsilon}$ is bounded on $X \setminus D_{lc}$: there exists $C_{t,\varepsilon} > 0$ such that

$$-C_{t,\varepsilon} - \sum_{a_k=1} \log(\log |s_k|^2)^2 \leq \varphi_{t,\varepsilon} \leq C_{t,\varepsilon} - \sum_{a_k=1} \log(\log |s_k|^2)^2. \quad (5.2)$$

Moreover, $\psi_{t,\varepsilon}$ is smooth in the quasi-coordinates. In particular, $\omega_{t,lc} + dd^c \psi_{t,\varepsilon}$ is a Kähler metric with bounded geometry on $X \setminus D_{lc}$ and with Poincaré type growth along D_{lc} . Therefore it is complete and has a bounded curvature tensor. To prove the regularity theorem, we will thus have to obtain on each compact subset of X_0 estimates on the potential φ_ε at any order.

A first attempt at the uniform estimate. The previous observation allows us to apply Yau’s maximum principle (cf. Proposition 5.4), and obtain that

$$\sup_{X \setminus D_{lc}} \psi_{t,\varepsilon} \leq \sup_{X \setminus D_{lc}} \left(\sum a_i \log(|s_i|^2 + \varepsilon^2) - f_t \right)$$

and similarly $\inf_{X \setminus D_{lc}} \psi_{t,\varepsilon} \leq \inf_{X \setminus D_{lc}} (\sum a_i \log(|s_i|^2 + \varepsilon^2) - f_t)$. If some coefficient a_i is negative, then we cannot obtain a bound for $\sup \psi_{t,\varepsilon}$. As for the lower bound, $-f_t$ is not uniformly bounded from below because $\omega_{t,lc}$ degenerates at $t = 0$ (and if some a_i is positive, $a_i \log(|s_i|^2 + \varepsilon^2)$ is not uniformly bounded below neither), so we cannot expect to find a lower bound for $\psi_{t,\varepsilon}$ using this strategy. Therefore we need another method to obtain a zero-order estimate on the potential of the solution. In fact, we will need to add some barrier function to gain positivity, in the spirit of Tsuji’s trick [Tsu88b] for the Laplacian estimate of a degenerate class; the novelty in our situation is that this is also needed for the zero-order estimates (as opposed to the klt case).

5.5 Uniform estimate. Before going any further, let us fix some notations.

5.5.1 A new framework. We will denote by $s_i, i \in I$ (non-zero) sections of the (reduced) components of the divisor $D_{red}+E$, and by $s_\alpha, \alpha \in A$ (non-zero) sections of the (reduced) components of E ; we endow all these line bundles with suitable smooth hermitian metrics (see below). Finally, we set $X_0 := X \setminus (\text{Supp}(D) \cup \text{Supp}(E))$, and define $F := (D_{red} + E)_{red}$ as the reduced divisor $X \setminus X_0$.

The idea is to work on X_0 . Of course, if we endow the last space with the Kähler metric $\omega_{t,lc}$, it will not be complete (near D_{klt} e.g.), so we won't be able to use Yau's maximum principle. Instead, we will rather use the following metric:

$$\omega_\chi := \theta + t\omega_0 - \sum_{i \in I} dd^c \log(\log |s_i|^2) + dd^c \chi$$

where $\chi := \sum_\alpha c_\alpha \log |s_\alpha|^2$ (recall that the c_α 's are the coefficients of E).

We do here a slight abuse of notation because ω_χ depends on t . However, the following lemma shows that the dependence is harmless:

LEMMA 5.8. *Up to changing the previously chosen hermitian metrics, the (1,1)-form ω_χ defines on X_0 a smooth Kähler metric with Poincaré growth along F having bounded geometry, all of those properties being satisfied uniformly in t .*

What we mean by this statement is that there exists a Poincaré-type metric ω_P on X_0 and a constant $C > 0$ independent of t such that $C^{-1}\omega_P \leq \omega_\chi \leq C\omega_P$, and that in the appropriate quasi-coordinates attached to the pair (X, F) , the coefficients $g_{i\bar{j}}$ of ω_χ satisfy $\left| \frac{\partial^{|\alpha|+|\beta|} g_{i\bar{j}}}{\partial z^\alpha \partial \bar{z}^\beta} \right| \leq C_{\alpha,\beta}$ for some constants $C_{\alpha,\beta} > 0$ independent of t . In particular, ω_χ has a uniformly (in t) bounded curvature tensor.

Proof of Lemma 5.8. We know that $\{\theta\} - c_1(E)$ is ample. Therefore, up to changing the hermitian metrics h_α on $\mathcal{O}_X(E_\alpha)$, we may suppose that $\eta := \theta - \sum c_\alpha \Theta_{h_\alpha}(E_\alpha)$ defines a smooth Kähler form on X [we designated by $\Theta_{h_\alpha}(E_\alpha)$ the curvature form of the hermitian line bundle (E_α, h_α)]. Therefore, on X_0 , we have:

$$\omega_\chi = \eta + t\omega_0 - \sum_{i \in I} dd^c \log(\log |s_i|^2)$$

and the statement follows easily from the computations of [CG72, Proposition 2.1] and [Kob84, Lemma 2] or [TY87]. □

5.5.2 Getting the lower bound. First of all, we will use the crucial information that $\varphi_{t,\varepsilon}$ converges (in the weak sense of distributions) to some θ -psh function (cf. first paragraph). By the elementary properties of psh (or quasi-psh) functions, we know that $(\varphi_{t,\varepsilon})$ is uniformly bounded above on the compact set X (see e.g. [Hör94, Theorem 3.2.13]). Therefore, we obtain some uniform constant C such that

$$\varphi_{t,\varepsilon} \leq C. \tag{5.3}$$

Now, recall that we chose ω_χ to be the new reference metric, so our equation becomes

$$(\omega_\chi + dd^c u_{t,\varepsilon})^n = e^{u_{t,\varepsilon} + G_\varepsilon} \omega_\chi^n \quad (5.4)$$

where

$$u_{t,\varepsilon} := \varphi_{t,\varepsilon} + \sum_{i \in I} \log(\log |s_i|^2)^2 - \chi$$

and

$$G_\varepsilon := \chi + f + \sum_{a_j < 1} \log \left(\frac{|s_j|^2}{(|s_j|^2 + \varepsilon^2)^{a_j}} \right) - \log \left(\frac{\prod_{i \in I} |s_i|^2 \log^2 |s_i|^2 \omega_\chi^n}{dV} \right).$$

Here again we should mention that G_ε also depends on t through the last term involving ω_χ . For our purpose, we can ignore this dependence in order to simplify the notations.

We can see from (5.3) and Lemma 5.8 that G_ε has a uniform (in t and ε) upper bound on X_0 :

$$\sup_{X_0} G_\varepsilon \leq C.$$

Moreover, we know from (5.2) that $\varphi_{t,\varepsilon} + \sum_{a_k=1} \log(\log |s_k|^2)^2$ is bounded. Therefore, it follows immediately that $u_{t,\varepsilon}$ is bounded *from below* (but a priori non uniformly). Applying Yau's maximum principle (cf. Proposition 5.4) to the smooth function $u_{t,\varepsilon}$ on the complete Kähler manifold (X_0, ω_χ) ensures that $\inf_{X_0} u_{t,\varepsilon} \geq -\sup_{X_0} G_\varepsilon \geq -C$. In terms of $\varphi_{t,\varepsilon}$, and recalling inequality (5.3) we get:

$$C \geq \varphi_{t,\varepsilon} \geq \chi - C - \sum_{i \in I} \log(\log |s_i|^2)^2.$$

5.6 Laplacian estimate. For the Laplacian estimate, we still work on the open manifold X_0 . We endow it with the complete Kähler metric ω_χ , and we recall from Lemma 5.8 that ω_χ has uniformly bounded (bisectional) curvature.

As usual when one wants to compare to Kähler metrics ω and ω' , the strategy is to use an inequality of the form $\Delta F \geq G$, where F, G involve terms like $\text{tr}_\omega \omega'$, $\text{tr}_{\omega'} \omega$ or the local potentials of $\omega' - \omega$. There exist several variants of such inequalities, due e.g. to Chen-Lu, Yau, Siu, etc. involving different assumptions on the curvature of the metrics at stake. In our case, as we have a control on the bisectional curvature of the reference metric ω_χ , on the Ricci curvature of the "unknown metric" $\omega_\chi + dd^c u_{t,\varepsilon}$, and on the Laplacian $\Delta_{\omega_\chi} G_\varepsilon$, we could use any of these formulas.

We have chosen to use a variant of Siu's inequality [Siu87, p. 99], which can be found in [CGP13, Proposition 2.1] (see also [Pău08, BBEGZ11]); notice the important feature allowing the factor e^{-F_-} for F_- quasi-psh which is crucial for us since the RHS of our Monge–Ampère equation has poles:

PROPOSITION 5.9. *Let X be a Kähler manifold of dimension n , ω, ω' two cohomologous Kähler metrics on X . We assume that $\omega' = \omega + dd^c u$ with $\omega'^n = e^{u+F_+-F_-}\omega^n$. and that we have a constant $C > 0$ satisfying:*

- (i) $dd^c F_{\pm} \geq -C\omega$,
- (ii) $\Theta_{\omega}(T_X) \geq -C\omega \otimes \text{Id}_{T_X}$.

Then there exists some constant $A > 0$ depending only on n and C such that

$$\Delta_{\omega'}(\log \text{tr}_{\omega}\omega' - Au + F_-) \geq \text{tr}_{\omega'}\omega - nA.$$

Moreover, if one assumes that $\sup F_+ \leq C, u \geq -C$ and that $\log \text{tr}_{\omega}\omega' - Au + F_-$ attains its maximum on X , then there exists $M > 0$ depending on n and C only such that:

$$\omega' \leq Me^{Au-F_-}\omega.$$

Here Δ (resp. Δ') is the Laplacian with respect to ω (resp. ω'), and $\Theta_{\omega}(T_X)$ is the Chern curvature tensor of the hermitian holomorphic vector bundle (T_X, ω) (which may be identified with the tensor of holomorphic bisectional curvatures, usually denoted by the letter R).

Sketch of proof. Siu’s inequality applied to $\omega = \sum g_{i\bar{j}}dz_i \wedge d\bar{z}_k$ and $\omega' = \sum g'_{i\bar{j}}dz_i \wedge d\bar{z}_k$ yields:

$$\Delta'(\log \text{tr}_{\omega}\omega') \geq \frac{1}{\text{tr}_{\omega}\omega'} \left(-g^{j\bar{i}}R_{j\bar{i}} + \Delta(u + F_+ - F_-) + g'^{k\bar{l}}R_{k\bar{l}}^{j\bar{i}}g'_{j\bar{i}} \right).$$

Recollecting terms coming (with different signs) from the scalar and the Ricci curvature, we will obtain a similar inequality involving only a lower bound for the holomorphic bisectional curvature, namely

$$\Delta' \log \text{tr}_{\omega}\omega' \geq \frac{\Delta(u + F_+ - F_-)}{\text{tr}_{\omega}(\omega')} - B \text{tr}_{\omega'}\omega \tag{5.5}$$

where B is a lower bound for the bisectional curvature of ω : this is the content of [CGP13, Lemma 2.2].

Clearly, $\Delta u = \text{tr}_{\omega}\omega' - n$ so that $\Delta(u + F_+) \geq -n(C + 1)$. As $\text{tr}_{\omega}\omega' \text{tr}_{\omega'}\omega \geq n$, we get

$$\frac{\Delta(u + F_+)}{\text{tr}_{\omega}\omega'} \geq -(1 + C)\text{tr}_{\omega'}\omega. \tag{5.6}$$

As for the second Laplacian, we write

$$0 \leq C\omega + dd^c F_- \leq \text{tr}_{\omega'}(C\omega + dd^c F_-)\omega'$$

and we take the trace with respect to ω :

$$\frac{nC + \Delta_{\omega}F_-}{\text{tr}_{\omega}\omega'} \leq C\text{tr}_{\omega'}\omega + \Delta_{\omega'}F_-$$

so that

$$\Delta_{\omega'} F_- \geq \frac{\Delta_{\omega} F_-}{\text{tr}_{\omega'} \omega} - C \text{tr}_{\omega'} \omega. \tag{5.7}$$

Plugging (5.6) and (5.7) into (5.5), we get:

$$\Delta'(\log \text{tr}_{\omega} \omega' + F_-) \geq -C_1 \text{tr}_{\omega'} \omega$$

for $C_1 = 1 + B + 2C$. Finally, using $\Delta' u = n - \text{tr}_{\omega'} \omega$, we see that

$$\Delta'(\log \text{tr}_{\omega} \omega' - (C_1 + 1)u + F_-) \geq \text{tr}_{\omega'} \omega - n(C_1 + 1)$$

which shows the first assertion by choosing $A := 1 + C_1$.

As for the second part, if we denote by p the point where the maximum is attained, then one has $(\text{tr}_{\omega'} \omega)(p) \leq C_2$. Using the basic inequality $\text{tr}_{\omega} \omega' \leq e^{u+F_+-F_-} (\text{tr}_{\omega'} \omega)^{n-1}$, one gets

$$\begin{aligned} \log(\text{tr}_{\omega} \omega') &= (\log \text{tr}_{\omega} \omega' - Au + F_-) + Au - F_- \\ &\leq (u(p) + F_+(p) - F_-(p)) + (n - 1) \log(nA) - Au(p) + F_-(p) + Au - F_- \\ &\leq C_2 + Au - F_- \end{aligned}$$

where $C_2 = \sup F_+ + (n - 1) \log(nA) - (A - 1) \inf u$ (recall that A can be chosen to be positive). This concludes the proof of the proposition. \square

Recall that we are interested in Equation (5.4) given by

$$(\omega_{\chi} + dd^c u_{t,\varepsilon})^n = e^{u_{t,\varepsilon} + G_{\varepsilon}} \omega_{\chi}^n.$$

We obtained the zero-order estimate on $u_{t,\varepsilon}$ in the last section, and now we want a Laplacian estimate. In order to use the previous proposition we first have to decompose G_{ε} as a difference of $C\omega_{\chi}$ -psh functions in order to use the result above. Recall that

$$G_{\varepsilon} := \chi + f + \sum_{a_j < 1} \log \left(\frac{|s_j|^2}{(|s_j|^2 + \varepsilon^2)^{a_j}} \right) - \log \left(\frac{\prod_{i \in I} |s_i|^2 \log^2 |s_i|^2 \omega_{\chi}^n}{dV} \right).$$

By [Kob84] or [Gue12, Lemma 1.6], the last term is already known to be smooth in the quasi-coordinates (and it depends neither on t nor on ε).

We claim that

$$G_{\varepsilon} = \underbrace{\left[\chi + f + \sum_{a_j < 1} \log |s_j|^2 + \sum_{a_j < 0} \log (|s_j|^2 + \varepsilon^2)^{-a_j} \right]}_{F_+} - \underbrace{\left[\sum_{0 < a_j < 1} \log (|s_j|^2 + \varepsilon^2)^{a_j} \right]}_{F_-}$$

gives the desired decomposition $G_{\varepsilon} = F_+ - F_-$ in the notations of Proposition 5.9. Indeed, $\chi, f, \log |s_i|^2$ are quasi-psh, thus $C\omega_{\chi}$ -psh for some uniform $C > 0$ as ω_{χ}

dominates some fixed Kähler form, cf. Lemma 5.8. Moreover, a simple computation leads to the identity:

$$dd^c \log(|s|^2 + \varepsilon^2) = \frac{\varepsilon^2}{(|s|^2 + \varepsilon^2)^2} \cdot \langle D's, D's \rangle - \frac{|s|^2}{|s|^2 + \varepsilon^2} \cdot \Theta_h$$

where Θ_h is the curvature of the hermitian metric h implicit in the term $|s|^2$, and $D's$ is the $(1, 0)$ -part of Ds where D is the Chern connection attached to $(\mathcal{O}_X(\text{div}(s)), h)$. It follows that F_{\pm} are $C\omega_{\chi}$ -psh for some uniform $C > 0$.

We can now apply Proposition 5.9 to the setting: $\omega = \omega_{\chi}, \omega' = \omega_{\chi} + dd^c u_{t,\varepsilon}$, $F_+ - F_- = G_{\varepsilon}$. Indeed, it is clear that F_+ is uniformly upper bounded, we just saw that F_{\pm} are $C\omega_{\chi}$ -psh, and we know from the previous section that $u_{t,\varepsilon}$ has a uniform lower bound. Furthermore, $\log \text{tr}_{\omega} \omega' - Au + F_-$ attains its maximum on X_0 : indeed, $-Au$ tends to $-\infty$ near the boundary of X_0 , F_- is bounded (it is even smooth on X), and $\text{tr}_{\omega} \omega' = \Delta_{\omega_{\chi}}(\varphi_{t,\varepsilon} + \sum_{i \in I} \log(\log |s_i|^2)^2 - \chi)$ is bounded on X_0 (we know it for the term $\Delta_{\omega_{\chi}}(\varphi_{t,\varepsilon} - \chi)$ and it is an elementary computation for the other term).

In conclusion, we may use Proposition 5.9 to obtain the following estimate:

$$\theta + t\omega_0 + dd^c \varphi_{t,\varepsilon} \leq M \left(\prod_{i \in I} \log^2 |s_i|^2 \right)^C \cdot \prod_{\alpha \in A} |s_{\alpha}|^{-c_{\alpha} \cdot C} \prod_{a_j > 0} |s_j|^{-2a_j} \omega_{\chi}.$$

For the “reverse inequality”, we use the identity

$$(\omega_{\chi} + dd^c u_{t,\varepsilon})^n = e^{u_{t,\varepsilon} + G_{\varepsilon}} \omega_{\chi}^n$$

which leads to the inequality

$$\text{tr}_{\omega_{\chi} + dd^c u_{t,\varepsilon}} \omega_{\chi} \leq e^{-(u_{t,\varepsilon} + G_{\varepsilon})} (\text{tr}_{\omega_{\chi}} (\omega_{\chi} + dd^c u_{t,\varepsilon}))^{n-1}$$

and therefore

$$\begin{aligned} & \theta + t\omega_0 + dd^c \varphi_{t,\varepsilon} \\ & \geq M^{-1} \prod_{a_j < 1} \frac{|s_j|^2}{(|s_j|^2 + \varepsilon^2)^{a_j}} \cdot \left(\prod_{i \in I} \log^2 |s_i|^2 \right)^{-C} \cdot \prod_{\alpha \in A} |s_{\alpha}|^{c_{\alpha} \cdot C} \prod_{a_j > 0} |s_j|^{2a_j} \omega_{\chi} \end{aligned}$$

for some uniform $C, M > 0$ (different from the previous ones).

In particular, for any compact set $K \Subset X_0$, there exists a constant $C_K > 0$ satisfying

$$C_K^{-1} \omega_0 \leq \theta + t\omega_0 + dd^c \varphi_{t,\varepsilon} \leq C_K \omega_0.$$

Using Evans–Krylov theorem and the classical elliptic theory shows that the potential $\varphi_{t,\varepsilon}$ satisfies uniform $\mathcal{C}^{k,\alpha}$ estimates on any $\Omega \Subset K$ for each k, α . Thus the theorem is proved.

REMARK 5.10. One can easily obtain somewhat more precise estimates. Indeed, if α is a nef and big class and E an effective \mathbb{R} -divisor such that $\alpha - E$ is ample, then we have in fact that for every $\delta > 0$, $\alpha - \delta E$ is ample (write $\alpha - \delta E = (1 - \delta)\alpha + \delta(\alpha - E)$). Applying this observation to Theorem 5.6, we see that for every $\delta > 0$, there exists $C_\delta > 0$ such that:

$$\varphi_{\text{KE}} \geq \delta \sum c_\alpha \log |s_\alpha|^2 - \sum_{i \in I} \log(\log |s_i|^2)^2 - C_\delta$$

for every $\delta > 0$. One could apply the same argument to the Laplacian estimates.

About uniqueness of the Kähler–Einstein metrics. First of all, in the case where X is smooth and K_X is ample, then uniqueness of the Kähler–Einstein metric constructed by Aubin and Yau is a straightforward consequence of the maximum principle. Generalizing this principle to some complete Kähler manifolds in [Yau75], Yau could prove that on a Kähler manifold, there can be only one *complete* Kähler metric ω satisfying $\text{Ric} \omega = -\omega$. In particular, this result has been applied by Kobayashi and Tian–Yau to show the uniqueness of the Kähler–Einstein metric for a log smooth pair (X, D) satisfying $K_X + D$ ample, cf. [Kob84, TY87]. Using Yau–Schwarz lemma in a more subtle way (through the notion of almost complete metric), they also show uniqueness when $K_X + D$ is only assumed nef, big and ample modulo D , which means that $K_X + D$ intersects positively every curve not contained in D . For example, if (X, D) is a log resolution of some canonically polarized singular variety, these assumptions are not satisfied.

In our situation we proceed in a different manner: we first use the volume assumption (as a replacement for completeness) to show that the Kähler–Einstein metric, originally defined on the (log) regular locus, extends to define a positive current on X whose local potentials glue to define a solution with full Monge–Ampère mass of a global Monge–Ampère equation to which we can apply the comparison principle to finally deduce the uniqueness.

6 Applications

6.1 Yau–Tian–Donaldson conjecture for singular varieties. Let us start with the following converse of Theorem A stated in the Section 1.

PROPOSITION 6.1. *Let X be projective variety satisfying the conditions G_1 and S_2 , and such that K_X is \mathbb{Q} -ample. If X admits a Kähler–Einstein metric ω in the following sense: ω is a K–E metric on X_{reg} and its total volume there is equal to K_X^n , then X has semi-log canonical singularities.*

Proof. We begin with the case where X is normal.

Let us first show that $\phi := \log \omega^n$ on X_{reg} extends to an element in $\mathcal{E}(X, K_X)$. By the K–E equation $dd^c \phi := -\text{Ric} \omega = \omega \geq 0$ and hence, since X is normal, ϕ

extends to a positively curved singular metric on K_X over all of X . Thus, writing $\omega = dd^c\psi$ for some other such metric ψ the compactness of X forces the K–E equation $\text{MA}(\phi) = e^\phi$ (up to shifting ϕ by a constant) globally on X (since the non-pluripolar MA does not charge the singular locus of X). Moreover, by the volume assumption we have that $\int_X \text{MA}(\phi) = K_X^n$ and hence $\phi \in \mathcal{E}(X, K_X)$, as desired.

Next, fix a resolution $\pi : X' \rightarrow X$ and assume, to get a contradiction, that $p^*K_X = K_{X'} + D$ where D is a snc \mathbb{Q} -divisor such that $D = D' + (1 + \delta)E$ for some $\delta > 0$, where D' is a \mathbb{Q} -divisor and E is a smooth irreducible divisor transversal to the support of D' . Since ϕ has maximal MA-mass it follows, as shown in [BBEGZ11] using an Izumi type estimate, that $\pi^*\phi$ has no Lelong numbers. In particular, it follows from the characterization of Lelong numbers that there exists a neighbourhood U of E such that $\pi^*\phi \geq \frac{1}{2}\delta \log |s_E|^2 - C$ in local trivializations. Moreover, we may take U such that D does not intersect U . But then it follows from the K–E equation that

$$K_X^n \geq C' \int_U e^{\pi^*\phi - (1+\delta)\log |s_E|^2} \geq C'' \int_U e^{-(1+\delta/2)\log |s_E|^2} = \infty,$$

which gives the desired contradiction.

We move on to the general case when X is only assumed to be G_1 and S_2 . As we observed in Section 3.3, the result of Proposition 3.6 holds actually in the general G_1 and S_2 case (we did not use at all that the singularities were slc); however we should be careful and work instead on a log-resolution of (X^ν, C_{X^ν}) because the formula $\nu^*K_X = K_{X^\nu} + C_{X^\nu}$ could not be satisfied anymore if C_{X^ν} is not Cartier, cf. Section 3.3 and the remarks following the identity (3.1). So the first conclusion is that the weight $\phi := \log \omega^n$ on X_{reg} extends on the normalization X^ν to a psh weight in $\mathcal{E}(X^\nu, \nu^*K_X)$. Then we take a log-resolution $\pi : X' \rightarrow X^\nu$ of the pair (X^ν, C_{X^ν}) where C_{X^ν} is the conductor of the normalization; a priori, this is just an effective divisor, possibly non-reduced. We write (X', D') for the new pair that we obtain on X' . Then same arguments as earlier show that D' has coefficients less than or equal to 1, which amounts to saying that (X^ν, C_{X^ν}) is log canonical, or equivalently that X has semi-log canonical singularities. \square

To relate this to K -stability we recall that Odaka [Oda13b] has shown that, if X is K -semistable, then X has semi-log canonical singularities (recall that we assume that X is G_1 and S_2 and that $K_X > 0$). Conversely, if X is semi-log canonical, then X is K -stable [Oda11]. Hence, combining our results with Odaka’s results gives the following confirmation of the Yau–Tian–Donaldson conjecture for varieties being G_1 and S_2 , with K_X ample:

Theorem 6.2. *Let X be a G_1 and S_2 projective variety such that K_X is ample. Then X admits a Kähler–Einstein metric iff X is K -stable.*

It would be interesting to have a direct analytical proof of the implication “Kähler–Einstein implies K -stable” as shown in [Ber12] the (log) Fano case (where

K -stability has to be replaced by K -polystability in the presence of holomorphic vector fields).

6.2 Automorphism groups of canonically polarized varieties. The existence and uniqueness of Kähler–Einstein metrics established in Theorem A allows us to give an analytical proof of the following result, that is already shown in [FG12, Corollary 3.13] (for lc pairs of general type) and [BHPS12] (in the stable case, with a geometric and a cohomological proof). Also, Kollár informed us that the result below was already well-known for some time, cf. e.g. [Miy83].

Theorem 6.3. *Let X be a normal stable variety. Then $\text{Aut}(X)$ is finite.*

Proof. By general results on automorphism groups of normal varieties (see [BBEGZ11, Lemma 5.2] and references therein) it is equivalent to show that any holomorphic vector field V on X_{reg} vanishes identically. To prove this vanishing we first observe that, by normality, V is the infinitesimal generator of a complex one-parameter family of automorphism F of X and in particular of X_{reg} . Fix a Kähler–Einstein current ω on X . By the naturality of the KE-equation it follows that $F^*\omega$ is also a KE-current and hence by uniqueness $F^*\omega = \omega$ on X_{reg} . Let us denote by V_r and V_i the real and imaginary parts of V , which are infinitesimal generators of real one parameter families of automorphisms that we will denote by F_r and F_i respectively, which, by the previous argument, also preserve ω . Next, note that any automorphism automatically lifts to the line bundle K_X over X_{reg} and thus it follows from general principles that the real part V_r of V is a Hamiltonian vector field, i.e. $i_{V_r}\omega = dh$ for some smooth function h on X_{reg} . But then, by Cartan’s formula, the Lie derivative $L_{V_i}\omega$ is given by $d(i_{V_i}\omega) = dJi_{V_i}\omega = dJdh = dd^c h$. Since the flow F_i defined by V_i also preserves ω it thus follows that $dd^c h = 0$. But by normality it follows that $h = 0$ (indeed, by normality h is bounded and we can thus apply the maximum principle on a resolution). Hence $i_{V_r}\omega = 0$ on X_{reg} , which forces $V_r = 0$ on X_{reg} , since ω is Kähler there and in particular pointwise non-degenerate. Finally, by the same argument $V_i = 0$ (for example replacing V with JV) and hence $V = 0$ as desired. \square

REMARK 6.4. In the case when X is smooth there is a simple cohomological proof of the previous proposition: by Serre duality $H^0(X, TX)$ is isomorphic to $H^{n-1}(X, -K_X)$, which is trivial by Kodaira vanishing (since K_X is ample). In the case when X is log canonical a similar cohomological argument can be used [BHPS12], relying on the Bogomolov–Sommese vanishing result for log canonical singularities, established in [GKKP11, Theorem 7.2]. Indeed, if V does not vanish identically then contracting with V on X_{reg} maps K_X to a rank one reflexive sheaf in $\text{Hom}(K_X, \Omega_X^{[n-1]})$, where $\Omega_X^{[n-1]}$ is the sheaf of reflexive $(n-1)$ -forms on X and hence by the Bogomolov–Sommese vanishing result in [GKKP11] the Kodaira dimension of K_X is at most $n-1$, which contradict the ampleness of K_X .

7 Outlook

7.1 Towards Miyaoka–Yau type inequalities. For simplicity we will only consider the case $n = 2$ (but a similar discussion applies in the general case). We set $E := \Omega_X^1$, the cotangent bundle of X . The classical case is when X is smooth with K_X ample, where the Miyaoka–Yau inequality says

$$c_1(E)^2 \leq 3c_2(E).$$

Let us briefly recall Yau’s differential-geometric proof. We equip E with the Hermitian metric induced by ω and denote by (E, ω) the corresponding Hermitian vector bundle. Then, if ω is Kähler–Einstein a direct local calculation gives the *point-wise* inequality

$$c_1(E, \omega)^2 \leq 3c_2(E, \omega)$$

formulated in terms of the Chern–Weil representatives $c_i(E, \omega)$ of the corresponding Chern classes. Hence, integrating immediately gives the Miyaoka–Yau inequality. Repeating this argument in the singular case when X a stable surface and using Theorem A gives the following

PROPOSITION 7.1. *The following inequality holds for a stable surface equipped with the canonical Kähler–Einstein metric ω on its regular part:*

$$c_1(K_X)^2 \leq 3 \int_{X_{\text{reg}}} c_2(E, \omega)$$

with equality iff ω has constant holomorphic sectional curvature, i.e. (X_{reg}, ω) is locally isometric to a ball.

Proof. Since the point-wise inequality above still holds, by the KE-condition, we can simply integrate it over X_{reg} and use that, by Theorem A, $c_1(K_X)^2 = \int_{X_{\text{reg}}} c_1(E, \omega)^2$. The conditions for equality are well-known in the point-wise inequality. \square

Since ω is canonically attached to X one could simply define the rhs appearing in the inequality above as the “analytical second Chern number” $c_{2,\text{an}}(X)$ of X . However, it should be stressed that it is not even a priori clear that $c_{2,\text{an}}(X)$ is finite, even though we expect that this is the case. More precisely, we expect that $c_{2,\text{an}}(X)$ can be identified with (or at least bounded from above) by a suitable algebraically defined second Chern class number $c_2(X)$. Various definitions of such Chern numbers have been proposed in the literature and we refer the reader to the paper of Langer [Lan00] where very general algebraic Miyaoka–Yau type inequalities are obtained, which in particular apply to stable surfaces. More generally, as before, our arguments apply to log canonical pairs.

7.2 The Weil–Peterson geometry of the moduli space of stable varieties.

In this section we will briefly explain how the finite energy property of the Kähler–Einstein metric on a stable variety naturally appears in the geometric study of the moduli space \mathcal{M} of all stable varieties. In a nutshell, the Kähler–Einstein metrics on stable varieties induces a metric on the \mathbb{Q} -line bundle $\mathcal{L} \rightarrow \mathcal{M}$ over the moduli space

defined by the top Deligne pairing of K_X and the finite energy condition is precisely the condition which makes sure that the metric is point-wise finite. The relation to Weil–Petersson geometry comes from the well-known fact that the curvature form of the corresponding metric over the moduli space \mathcal{M}_0 of all *smooth* stable varieties (i.e. all canonically polarized n -dimensional manifolds, with a fixed oriented smooth structure) coincides with the Weil–Petersson metric Ω_{WP} on \mathcal{M}_0 [FS90, Sch12].

To be a bit more precise we first recall that given a line bundle $L \rightarrow X$ over a (complex) n -dimensional algebraic variety X its top Deligne pairing i.e. the $(n+1)$ -fold Deligne pairing of L with itself is a complex line that we will denote by $\langle L \rangle$ [Elk89, Elk90]. Equipping $\langle L \rangle$ with an Hermitian metric ϕ (using additive notation as before) induces a Hermitian metric $\langle \phi \rangle$ on $\langle L \rangle$, satisfying the change of metric formula: $\langle \phi \rangle - \langle \psi \rangle = (\mathcal{E}(\phi) - \mathcal{E}(\psi))$ (up to a multiplicative normalization constant), where \mathcal{E} is the energy functional appearing in Section 4 (compare [PS04]). Fixing a smooth reference metric ϕ_0 on L one can use the latter transformation formula to define the metric $\langle \phi \rangle$ as long as ϕ has finite energy. The resulting metric $\langle \phi \rangle$ is then independent of the choice of reference metric ϕ_0 . More generally, in the relative case of a flat morphism $\mathcal{X} \rightarrow B$ between integral schemes of relative dimension n and an Hermitian line bundle $L \rightarrow \mathcal{X}$ this construction produces an Hermitian line bundle $\langle L \rangle \rightarrow B$ over the base B .

In particular, taking X to be an n -dimensional stable variety $L := K_X$ one obtains a canonical metric on the complex line $\langle K_X \rangle$, induced by the finite energy metric on K_X determined by the volume form of the Kähler–Einstein metric on the regular locus of X .

Let now \mathcal{M} denote the moduli space of all n -dimensional stable varieties with a fixed Hilbert polynomial [Kol10, Kol]. Using the existence of a universal stable family \mathcal{X} (in the sense of Kollar) over a finite cover of each irreducible component of the moduli space one obtains a \mathbb{Q} -line bundle \mathcal{L} over \mathcal{M} , induced by the fiber-wise top Deligne pairings $\langle K_X \rangle$. We conjecture that the metric on \mathcal{L} induced by the fiber-wise Kähler–Einstein metrics is continuous (as in the case of stable curves [Fre12]). Confirming this conjecture would require a more detailed analysis of the dependence of the Kähler–Einstein metric on the complex structure that we leave for the future.

As is well-known the curvature of the corresponding metric over \mathcal{M}_0 coincides (up to a numerical factor) with the Weil–Petersson metric Ω_{WP} . In particular, it is strictly positive as a form (in the orbifold sense). Under the validity of the previous conjecture one thus obtains a canonical extension of the induced Weil–Petersson metric on \mathcal{M}_0 to its compactification in \mathcal{M} as a positive current with continuous potentials. It would also be very interesting to know under which assumptions the extension is *strictly* positive in a suitable sense, for example if it is, locally, the restriction of a Kähler metric? These problems are (e.g. by Grauert’s generalization of Kodaira’s embedding theorem to singular varieties) closely related to the problem of showing that the corresponding line bundle \mathcal{L} over the moduli space \mathcal{M} is ample (on each irreducible component) and it should be compared with the recent work of Schumacher

[Sch12], where an analytic proof of the quasi-projectivity of \mathcal{M}_0 is given. As shown by Schumacher the Weil–Petersson metric Ω_{WP} on \mathcal{M}_0 admits a (non-canonical) extension as a positive current with analytic singularities to Artin’s Moishezon compactification of \mathcal{M}_0 . But the conjecture above is closely related to the problem of obtaining a *canonical* extension of Ω_{WP} to \mathcal{M} as a positive current with continuous potentials.

Finally, it should be pointed out that the top Deligne pairing used above essentially coincides with Tian’s CM-line bundle in this setting (by the Knudson–Mumford expansion and Zhang’s isomorphism realizing the Chow divisor as a top Deligne pairing). The ampleness of the induced CM-line bundle over general moduli spaces of K -stable polarized varieties was recently speculated on by Odaka [Oda13a].

Acknowledgments

It is a pleasure to thank Bo Berndtsson and Sébastien Boucksom for helpful discussions and Sándor Kovács for answering our questions concerning the singularities of stable varieties. The joint work started when the second author visited Chalmers University, which he would like to thank for the warm hospitality and the inspiring environment.

References

- [Ale96] V. ALEXEEV. Log Canonical Singularities and Complete Moduli of Stable Pairs. [arXiv:alg-geom/9608013](https://arxiv.org/abs/alg-geom/9608013) (1996).
- [Aub78] T. AUBIN. Équations du type Monge–Ampère sur les variétés Kähleriennes compactes. *Bull. Sci. Math.*, **102** (1978)
- [BB12] R. J. BERMAN and B. BERNDTSSON. Real Monge–Ampère Equations and Kähler–Ricci Solitons on Toric Log Fano Varieties. [arXiv:1207.6128](https://arxiv.org/abs/1207.6128) (2012).
- [BBEGZ11] R. J. BERMAN, S. BOUCKSOM, P. EYSSIDIEUX, V. GUEDJ and A. ZERIAHI. Kähler–Einstein Metrics and the Kähler–Ricci Flow on Log-Fano Varieties. [arXiv:1111.7158v2](https://arxiv.org/abs/1111.7158v2) (2011).
- [BBGZ09] R. J. BERMAN, S. BOUCKSOM, V. GUEDJ and A. ZERIAHI. A variational approach to complex Monge–Ampère equations (2009) (to appear in Publ. IHES, [arXiv:0907.4490](https://arxiv.org/abs/0907.4490)).
- [BBP10] S. BOUCKSOM, A. BROUSTET and G. PACIENZA. Uniruledness of stable base loci of adjoint linear systems with and without Mori theory (2010) (to appear in *Math. Zeit.*, [arXiv:0902.1142](https://arxiv.org/abs/0902.1142)).
- [BCHM10] C. BIRKAR, P. CASCINI, C. HACON and J. MCKERNAN. Existence of minimal models for varieties of log general type. *J. Am. Math. Soc.*, **23** (2010), 405468
- [BD12] R. J. BERMAN and J.-P. DEMAILLY. Regularity of plurisubharmonic upper envelopes in big cohomology classes, Perspectives in analysis, geometry, and topology. In: *Progress in Mathematics*, Vol. 296. Birkhäuser/Springer, New York (2012), p. 3966.
- [BEG13] S. BOUCKSOM, P. EYSSIDIEUX and V. GUEDJ. Introduction to the Kähler–Ricci flow. In: *Lecture Notes in Mathematics* (to appear). Springer, New York (2013).

- [BEGZ10] S. BOUCKSOM, P. EYSSIDIEUX, V. GUEDJ and A. ZERIAHI. Monge–Ampère equations in big cohomology classes. *Acta Math.*, (2)**205** (2010), 199262
- [Ber09] R. J. BERMAN. Bergman kernels and equilibrium measures for line bundles over projective manifolds. *Am. J. Math.* (5)**131** (2009), 14851524
- [Ber11] R. J. BERMAN. A thermodynamical formalism for Monge–Ampère equations, Moser–Trudinger inequalities and Kähler–Einstein metrics. [arXiv:1011.3976](https://arxiv.org/abs/1011.3976) (2011).
- [Ber12] R. J. BERMAN. K-polystability of \mathbb{Q} -Fano varieties admitting Kähler–Einstein metrics. [arXiv:1205.6214](https://arxiv.org/abs/1205.6214) (2012).
- [BHPS12] B. BHATT, W. HO, Z. PATAKFALVI and C. SCHNELL. Moduli of products of stable varieties. [arXiv:1206.0438](https://arxiv.org/abs/1206.0438) (2012).
- [Bou02] S. BOUCKSOM. On the volume of a line bundle. *Int. J. Math.*, (10)**13** (2002), 10431063
- [Bou04] S. BOUCKSOM. Divisorial Zariski decompositions on compact complex manifolds. *Ann. Sci. École Norm. Sup. (4)*, (1)**37** (2004), 4576
- [Bre11] S. BRENDLE. Ricci flat Kähler metrics with edge singularities (2011) (to appear in IMRN, [arXiv:1103.5454](https://arxiv.org/abs/1103.5454))
- [BT87] E. BEDFORD and B. TAYLOR. Fine topology, Silov boundary, and $(dd^c)^n$. *J. Funct. Anal.*, (2)**72** (1987), 225251
- [CDS12a] X. CHEN, S. DONALDSON and S. SUN. Kähler–Einstein Metrics on Fano Manifolds, I: Approximation of Metrics with Cone Singularities. [arXiv:1211.4566](https://arxiv.org/abs/1211.4566) (2012).
- [CDS12b] X. CHEN, S. DONALDSON and S. SUN. Kähler–Einstein Metrics on Fano Manifolds, II: Limits with Cone Angle Less Than 2π . [arXiv:1212.4714](https://arxiv.org/abs/1212.4714) (2012).
- [CDS13] X. CHEN, S. DONALDSON and S. SUN. Kähler–Einstein Metrics on Fano Manifolds, III: Limits as Cone Angle Approaches 2π and Completion of the Main Proof. [arXiv:1302.0282](https://arxiv.org/abs/1302.0282) (2013).
- [CG72] J. CARLSON and P. GRIFFITHS. A defect relation for equidimensional holomorphic mappings between algebraic varieties. *Ann. Math.* **95** (1972), 557584
- [CGP13] F. CAMPANA, H. GUENANCIA and M. PĂUN. Metrics with cone singularities along normal crossing divisors and holomorphic tensor fields. *Ann. Sci. École Norm. Sup.*, **46** (2013), 879916
- [CY80] S.-Y. CHENG and S.-T. YAU. On the existence of a complete Kähler metric on non-compact complex manifolds and the regularity of Fefferman’s equation. *Commun. Pure Appl. Math.*, **33** (1980), 507544
- [Dem82] J.-P. DEMAILLY. Estimations L^2 pour l’opérateur $\bar{\partial}$ d’un fibré vectoriel holomorphe semi-positif au-dessus d’une variété kählérienne complète. *Ann. Sci. École Norm. Sup. (4)*, (3)**15** (1982), 457511
- [Dem85] J.-P. DEMAILLY. Mesures de Monge–Ampère et caractérisation géométrique des variétés algébriques affines. *Mém. Soc. Math. France (N.S.)*, **19** (1985), 124
- [Dem92] J.-P. DEMAILLY. Regularization of closed positive currents and intersection theory. *J. Algebraic Geom.*, (3)**1** (1992), 361409
- [DM69] P. DELIGNE and D. MUMFORD. The irreducibility of the space of curves of given genus. *Inst. Hautes Études Sci. Publ. Math.*, **36** (1969), 75109
- [DP10] J.-P. DEMAILLY and N. PALI. Degenerate complex Monge–Ampère equations over compact Kähler manifolds. *Int. J. Math.*, **21** (2010)

- [EGZ08] P. EYSSIDIEUX, V. GUEDJ and A. ZERIAHI. A priori L^∞ -estimates for degenerate complex Monge–Ampère equations. *Int. Math. Res. Notes* (2008), Art. ID rnn 070, 8
- [EGZ09] P. EYSSIDIEUX, V. GUEDJ and A. ZERIAHI. Singular Kähler–Einstein metrics. *J. Am. Math. Soc.*, **22** (2009), 607639
- [Elk89] R. ELKIK. Fibrés d’intersections et intégrales de classes de Chern. *Ann. Sci. École Norm. Sup. (4)*, (2)**22** (1989), 195226
- [Elk90] R. ELKIK. Métriques sur les fibrés d’intersection. *Duke Math. J.*, (1)**61** (1990), 303328
- [FG12] O. FUJINO and Y. GONGYO. Log Pluricanonical Representations and Abundance Conjecture. [arXiv:1104.0361](https://arxiv.org/abs/1104.0361) (2012).
- [FN80] J. E. FORNÆSS and R. NARASIMHAN. The Levi problem on complex spaces with singularities. *Math. Ann.*, (1)**248** (1980), 4772
- [Fre12] G. FREIXAS I MONTPLET. An arithmetic Hilbert–Samuel theorem for pointed stable curves. *J. Eur. Math. Soc. (JEMS)*, (2)**14** (2012), 321351
- [FS90] A. FUJIKI and G. SCHUMACHER. The moduli space of extremal compact Kähler manifolds and generalized Weil–Petersson metrics. *Publ. Res. Inst. Math. Sci.*, (1)**26** (1990), 101183
- [Fuj10] O. FUJINO. Finite generation of the log canonical ring in dimension four. *Kyoto J. Math.*, (4)**50** (2010), 671684
- [GKKP11] D. GREB, S. KEBEKUS, S. J. KOVÁCS and T. PETERNELL. Differential forms on log canonical spaces. *Publ. Math. Inst. Hautes Études Sci.* **114** (2011), 87169
- [GP13] H. GUENANCIA and M. PĂUN. Conic Singularities Metrics with Prescribed Ricci Curvature: The Case of General Cone Angles Along Normal Crossing Divisors. [arXiv:1307.6375](https://arxiv.org/abs/1307.6375) (2013).
- [GR56] H. GRAUERT and R. REMMERT. Plurisubharmonische Funktionen in komplexen Räumen. *Math. Z.*, **65** (1956), 175194
- [Gri76] P. A. GRIFFITHS. Entire Holomorphic Mappings in One and Several Complex Variables (A. of Mathematics Studies, ed.). Princeton University Press, Princeton (1976).
- [GT80] S. GRECO and C. TRAVERSO. On seminormal schemes. *Compos. Math.*, (3)**40** (1980), 325365
- [Gue12] H. GUENANCIA. Kähler–Einstein metrics with mixed Poincaré and cone singularities along a normal crossing divisor (2012) (to appear in *Ann. Inst. Fourier*, [arXiv:1201.0952](https://arxiv.org/abs/1201.0952))
- [Gue13] H. GUENANCIA. Kähler–Einstein metrics with cone singularities on klt pairs. *Int. J. Math.*, **24** (2013)
- [GW] H. GUENANCIA and D. WU. On the boundary behaviour of Kähler–Einstein metrics of log canonical pairs (in preparation)
- [GZ05] V. GUEDJ and A. ZERIAHI. Intrinsic capacities on compact Kähler manifolds. *J. Geom. Anal.*, (4)**15** (2005), 607639
- [GZ07] V. GUEDJ and A. ZERIAHI. The weighted Monge–Ampère energy of quasi plurisubharmonic functions. *J. Funct. Anal.*, **250** (2007), 442482
- [HK13] J. H. HUBBARD and S. KOCH. An analytic construction of the Deligne–Mumford compactification of the moduli space of curves. [arXiv:1301.0062](https://arxiv.org/abs/1301.0062) (2013).
- [Hör94] L. HÖRMANDER. *Notions of Convexity*. Birkhäuser, Basel (1994).
- [JMR11] T. JEFFRES, R. MAZZEO and Y. RUBINSTEIN. Kähler–Einstein metrics with

- edge singularities (2011) (to appear in *Ann. Math.*, [arXiv:1105.5216](#), with an appendix by C. Li and Y. Rubinstein)
- [K+92] J. KOLLÁR et al. Flips and Abundance for Algebraic Threefolds. Société Mathématique de France, Paris. In: *Papers from the Second Summer Seminar on Algebraic Geometry held at the University of Utah, Salt Lake City, August 1991*, Astérisque No. 211 (1992).
- [Kar00] K. KARU. Minimal models and boundedness of stable varieties. *J. Algebraic Geom.*, (1)**9** (2000), 93109
- [KM98] J. KOLLÁR and S. MORI. Birational geometry of algebraic varieties. In: *Cambridge Tracts in Mathematics*, Vol. 134. Cambridge University Press, Cambridge (1998) (with the collaboration of C. H. Clemens and A. Corti, translated from the 1998 Japanese original).
- [Kob84] R. KOBAYASHI. Kähler–Einstein metric on an open algebraic manifolds. *Osaka 1. Math.*, **21** (1984), 399418
- [Kol] J. KOLLÁR. *Book on Moduli of Surfaces* (ongoing project, available at the author’s webpage <https://web.math.princeton.edu/kollar/book/chap3.pdf>).
- [Kol10] J. KOLLÁR. Moduli of Varieties of General Type. [arXiv:1008.0621](#) (2010).
- [Kov12] S. J. KOVÁCS. Singularities of stable varieties (2012) (to appear in *Handbook of Moduli*, [arXiv:1102.1240](#)).
- [KSB88] J. KOLLÁR and N. I. SHEPHERD-BARRON. Threefolds and deformations of surface singularities. *Invent. Math.*, (2)**91** (1988), 299338
- [KSS10] S. J. KOVÁCS, K. SCHWEDE and K. E. SMITH. The canonical sheaf of Du Bois singularities. *Adv. Math.*, (4)**224** (2010), 16181640
- [Lan00] A. LANGER. Logarithmic Orbifold Euler Numbers of Surfaces with Applications. [arXiv:0012180](#) (2000).
- [Laz04] R. LAZARSFELD. *Positivity in Algebraic Geometry II*. Springer, New York (2004).
- [Miy83] M. MIYANISHI. On the affine-ruledness of algebraic varieties. Algebraic geometry. Proceedings of the Japan–France Conference, Tokyo and Kyoto 1982. In: *Lecture Notes in Mathematics*, Vol. 1016 (1983), pp. 449–485.
- [MY83] N. MOK and S.-T. YAU. Completeness of the Kähler–Einstein metric on bounded domains and the characterization of domains of holomorphy by curvature conditions. In: *The Mathematical Heritage of Henri Poincaré, Part 1 (Bloomington, Ind., 1980)*. Proceedings of the Symposium on Pure Mathematics, Vol. 39. American Mathematical Society, Providence, RI (1983), p. 4159.
- [Oda11] Y. ODAKA. The Calabi Conjecture and K-stability. [arXiv:1010.3597](#) (2011).
- [Oda13a] Y. ODAKA. On the Moduli of Kahler–Einstein Fano Manifolds. [arXiv:1211.4833](#) (2013).
- [Oda13b] Y. ODAKA. The GIT-stability of polarised varieties via discrepancy. *Ann. Math.*, (2)**177** (2013), 645661
- [OSS12] Y. ODAKA, C. SPOTTI and S. SUN. Compact Moduli Spaces of Del Pezzo Surfaces and Kähler–Einstein Metrics. [arXiv:1210.0858](#) (2012).
- [Pău08] M. PĂUN. Regularity properties of the degenerate Monge–Ampère equations on compact Kähler manifolds. *Chin. Ann. Math. Ser. B*, (6)**29** (2008), 623630
- [PS04] D. H. PHONG and J. STURM. Scalar curvature, moment maps, and the Deligne pairing. *Am. J. Math.*, (3)**126** (2004), 693712
- [Sch12] G. SCHUMACHER. Positivity of Relative Canonical Bundles and Applications. [arXiv:1201.2930](#) (2012).

- [Siu87] Y.-T. SIU. *Lectures on Hermitian–Einstein Metrics for Stable Bundles and Kähler–Einstein Metrics*. Birkhäuser, Basel (1987).
- [Son14] J. SONG. Riemannian Geometry of Kähler–Einstein currents. [arXiv:1404.0445](https://arxiv.org/abs/1404.0445) (2014).
- [Tia13] G. TIAN. K-stability and Kähler–Einstein Metrics. [arXiv:1211.4669](https://arxiv.org/abs/1211.4669) (2013).
- [Tra70] C. TRAVERSO. Seminormality and Picard group. *Ann. Scuola Norm. Sup. Pisa* (3), **24** (1970), 585595
- [Tsu88a] H. TSUJI. A characterization of ball quotients with smooth boundary. *Duke Math. J.*, (2)**57** (1988), 537553
- [Tsu88b] H. TSUJI. Existence and degeneration of Kähler–Einstein metrics on minimal algebraic varieties of general type. *Math. Ann.*, (1)**281** (1988), 123133
- [TY87] G. TIAN and S.-T. YAU. Existence of Kähler–Einstein metrics on complete Kähler manifolds and their applications to algebraic geometry. *Adv. Ser. Math. Phys.* 1, **1** (1987), 574628 (mathematical aspects of string theory, San Diego, California, 1986)
- [TY90] G. TIAN and S.-T. YAU. Complete Kähler manifolds with zero Ricci curvature. *I. J. Am. Math. Soc.*, (3)**3** (1990), 579609
- [Var89] J. VAROUCHAS. Kähler spaces and proper open morphisms. *Math. Ann.*, (1)**283** (1989), 1352
- [Vie95] E. VIEHWEG. *Quasi-projective moduli for polarized manifolds*. In: *Ergebnisse der Mathematik und ihrer Grenzgebiete* (3) [*Results in Mathematics and Related Areas* (3)], Vol. 30. Springer-Verlag, Berlin (1995).
- [Wu08] D. WU. Kähler–Einstein metrics of negative Ricci curvature on general quasi-projective manifolds. *Commun. Anal. Geom.*, (2)**16** (2008), 395435
- [Wu09] D. WU. Good Kähler metrics with prescribed singularities. *Asian J. Math.*, (1)**13** (2009), 131150
- [Yau75] S.-T. YAU. Harmonic functions on complete Riemannian manifolds. *Commun. Pure Appl. Math.*, **28** (1975), 201228
- [Yau78] S.-T. YAU. On the Ricci curvature of a compact Kähler manifold and the complex Monge–Ampère equation, I. *Commun. Pure Appl. Math.*, **31** (1978), 339411
- [Yau93] S.-T. YAU. A splitting theorem and an algebraic geometric characterization of locally Hermitian symmetric spaces. *Commun. Anal. Geom.*, (3–4)**1** (1993), 473486

ROBERT J. BERMAN, Chalmers Techniska Hogskola, Göteborg, Sweden

robertb@math.chalmers.se

HENRI GUENANCIA, Institut de Mathématiques de Jussieu, Université Pierre et Marie Curie, Paris, France

guenancia@math.jussieu.fr

and

Département de Mathématiques et Applications, École Normale Supérieure, Paris, France

guenanci@dma.ens.fr

Received: March 21, 2014

Revised: July 7, 2014

Accepted: July 13, 2014