

Spectral Measures Associated with the Factorization of the Lebesgue Measure on a Set via Convolution

Jean-Pierre Gabardo · Chun-Kit Lai

Received: 25 September 2013 / Revised 20 February 2014 / Published online: 8 April 2014
© Springer Science+Business Media New York 2014

Abstract Let Q be a fundamental domain of some full-rank lattice in \mathbb{R}^d and let μ and ν be two positive Borel measures on \mathbb{R}^d such that the convolution $\mu * \nu$ is a multiple of χ_Q . We consider the problem as to whether or not both measures must be spectral (i.e. each of their respective associated L^2 space admits an orthogonal basis of exponentials) and we show that this is the case when $Q = [0, 1]^d$. This theorem yields a large class of examples of spectral measures which are either absolutely continuous, singularly continuous or purely discrete spectral measures. In addition, we propose a generalized Fuglede's Conjecture for spectral measures on \mathbb{R}^1 and we show that it implies the classical Fuglede's Conjecture on \mathbb{R}^1 .

Keywords Convolutions · Fuglede's Conjecture · Lebesgue measures · Spectral measures · Spectra

Mathematics Subject Classification Primary 42B05 · 42A85 · 28A25.

1 Introduction

Let μ be a compactly supported Borel probability measure on \mathbb{R}^d . We say that μ is a *spectral measure* if there exists a countable set $\Lambda \subset \mathbb{R}^d$ called *spectrum* such

Communicated by Yang Wang.

The first named author was supported by an NSERC grant.

J.-P. Gabardo · C.-K. Lai (✉)
Department of Mathematics and Statistics, McMaster University, Hamilton, ON L8S 4K1, Canada
e-mail: cklai@math.mcmaster.ca

J.-P. Gabardo
e-mail: gabardo@mcmaster.ca

that $E(\Lambda) := \{e^{2\pi i(\lambda \cdot x)} : \lambda \in \Lambda\}$ is an orthonormal basis for $L^2(\mu)$. If $\Omega \subset \mathbb{R}^d$ is measurable with finite positive Lebesgue measure and $d\mu(x) = \chi_\Omega(x)dx$ is a spectral measure, then we say that Ω is a *spectral set*. Spectral sets were first introduced by Fuglede [7] and have a very delicate and mysterious relationship with translational tiling because of the *spectral set conjecture* (known also as *Fuglede’s Conjecture*) proposed by Fuglede.

Conjecture (Fuglede’s Conjecture): *A bounded measurable set Ω on \mathbb{R}^d of positive Lebesgue measure is a spectral set if and only if Ω is a translational tile.*

We say that Ω is a translational tile if there exists a discrete set \mathcal{J} such that $\bigcup_{t \in \mathcal{J}} (\Omega + t) = \mathbb{R}^d$, and the Lebesgue measure of $(\Omega + t) \cap (\Omega + t')$ is zero for any distinct t and t' in \mathcal{J} . Although this conjecture was eventually disproved in dimension $d \geq 3$ [11, 12, 22], most of the known examples of spectral sets are constructed from translational tiles. An important class of examples of spectral sets constructed in [20] consists of sets of the form $A + [0, 1]$ tiling $[0, N]$ for some N , where $A \subset \mathbb{Z}$. In fact, in this case, the corresponding equally weighted discrete measure on A is a spectral measure.

The first singular spectral measure was constructed by Jorgensen and Pedersen [8]. They showed that the standard Cantor measures are spectral measures if the contraction is $\frac{1}{2n}$, while there are at most two orthogonal exponentials when the contraction is $\frac{1}{2n+1}$. Following this discovery, more spectral self-similar/self-affine measures were also found [4, 15, 21]. In these investigations, the tiling conditions on the digit sets play an important role. An interesting question arises naturally:

Question: *What kind of measures are spectral measures and how are they related to translational tilings?*

This question seems to be out of reach using our current knowledge. In this paper, we aim to describe a unifying framework bridging the gap between singular spectral measures and spectral sets. Let us introduce some simple notations. Denote by \mathcal{L} the Lebesgue measure in \mathbb{R}^d and by \mathcal{L}_E the normalized Lebesgue measure restricted to the measurable set E of finite positive Lebesgue measure (i.e. $\mathcal{L}_E(F) = \mathcal{L}(E \cap F) / \mathcal{L}(E)$). For a finite set A , we denote by $|A|$ the cardinality of A and by δ_A the measure $\sum_{a \in A} \delta_a$, where δ_a is the Dirac mass at a . We also write $A \oplus B = C$ if every element in C can be uniquely expressed as a sum $a + b$ with $a \in A$ and $b \in B$. We now make some observations about specific examples of spectral measures known in the literature.

- (1) According to [20], if $A \subset \mathbb{Z}$ and the set $\Omega = A + [0, 1)$ tiles $[0, N)$, then Ω is a spectral set. We can thus find a set B such that $A \oplus B = \{0, 1, \dots, N - 1\}$. This means that $(\frac{1}{|B|} \delta_B) * \mathcal{L}_\Omega = \mathcal{L}_{[0, N]}$.
- (2) Let μ be the standard one-fourth Cantor (probability) measure defined by the self-similar identity

$$\mu(\cdot) = \frac{1}{2} \mu(4 \cdot) + \frac{1}{2} \mu(4 \cdot - 2).$$

It is known that μ is a spectral measure [8]. At the same time, we observe that if we define ν to be the one-fourth Cantor measure obeying the equation

$$\nu(\cdot) = \frac{1}{2} \nu(4 \cdot) + \frac{1}{2} \nu(4 \cdot - 1),$$

then $\mu * \nu = \mathcal{L}_{[0,1]}$. This can be seen directly by computing the Fourier transform of both measures.

In fact, we may view the operation of convolution with a positive measure as certain kind of generalized translation. The above examples suggest the following question. Let Q be a fundamental domain of some full-rank lattice on \mathbb{R}^d .

$\mathcal{F}(Q)$: Any positive Borel measures μ and ν such that $\mu * \nu = \mathcal{L}_Q$ are spectral measures.

Unfortunately, we cannot expect the above statement to be true for all Q . In fact, if $\mu = \mathcal{L}_E$ with E is the translational tile without a spectrum constructed in [11], then $\mu * \nu = \mathcal{L}_Q$ for some fundamental domain Q as seen directly from the construction of this counterexample. However, in order to understand which measures are spectral, it is useful to know to what extent the statement $\mathcal{F}(Q)$ is true for some specific Q . Our first main result unifies the examples of discrete spectral measures, spectral sets and the singular spectral measures given in (1) and (2) above.

Theorem 1.1 For any $d \geq 1$, the statement $\mathcal{F}([0, 1]^d)$ is true. Moreover, for any positive Borel measures μ and ν such that $\mu * \nu = \mathcal{L}_{[0,1]^d}$, we can find spectra Λ_μ and Λ_ν for μ and ν respectively satisfying the property that

$$\Lambda_\mu \oplus \Lambda_\nu = \mathbb{Z}.$$

We now give a brief explanation of the proof of Theorem 1.1. We first focus on \mathbb{R}^1 where the proof involves two main steps. The first step is a complete characterization of the Borel probability measures μ and ν satisfying the identity $\mu * \nu = \mathcal{L}_{[0,1]}$. This characterization is actually a known result in probability due to Lewis [16]. In particular, Lewis proved that only two cases could occur: either one measure is absolutely continuous and the other one is purely discrete or they are both singular. To prove our theorem, we will express the measures μ and ν as weak limits of convolutions of some discrete measures using the result of Lewis (see Sect. 2). The second step is to construct spectra for μ and ν . This is done by observing that the discrete measures obtained at each level are spectral measures. We then show that the spectral property carries over by passing to the weak limit. This argument is a generalization of the proof in [2] (see Sect. 3). After the dimension one case is established, we characterize the Borel probability measures μ and ν satisfying $\mu * \nu = \mathcal{L}_{[0,1]^d}$ as Cartesian products of one-dimensional Borel probability measures σ_i and τ_i , $i = 1, \dots, d$, on \mathbb{R}^1 satisfying $\sigma_i * \tau_i = \mathcal{L}_{[0,1]}$ and also prove the spectral property for those (see Sect. 5).

The results in [11] show that the statement $\mathcal{F}(Q)$ might not be true when Q is a general fundamental domain of a lattice. On the other hand, we also know that it can be true for some non-cube fundamental domains. For instance, the set $Q = [0; 1] + \{0, 3\}$ is not a cube but it is a fundamental domain of the lattice $2\mathbb{Z}$ and it tiles a cube, so the statement $\mathcal{F}(Q)$ holds in this case using Theorem 1.1. Of course, we don't know what happens for the vast majority of the fundamental domains. We will focus our attention on \mathbb{R}^1 in which Fuglede's Conjecture remains open. We propose the following generalized Fuglede's Conjecture for spectral measures on \mathbb{R}^1 and it is immediate to see that a full generality of $\mathcal{F}(Q)$ on \mathbb{R}^1 will imply one direction of this generalized conjecture.

Conjecture (Generalized Fuglede’s Conjecture): *A compactly supported Borel probability measure μ on \mathbb{R}^1 is spectral if and only if there exists a Borel probability measure ν and a fundamental domain Q of some lattice on \mathbb{R}^1 such that $\mu * \nu = \mathcal{L}_Q$.*

This is an open conjecture on \mathbb{R}^1 and we will prove that it extends the classical Fuglede’s Conjecture.

Theorem 1.2 *The generalized Fuglede’s Conjecture implies Fuglede’s Conjecture on \mathbb{R}^1 .*

Let us make some remarks on the classical Fuglede’s Conjecture on \mathbb{R}^1 . There is some evidence that the conjecture may be true on \mathbb{R}^1 . In particular, the known fact that all tiling sets of a tile and all spectra of a spectral set are periodic offers some credibility to the conjecture [10, 13]. Moreover, some algebraic conditions, if satisfied, are sufficient to settle the conjecture on \mathbb{R} , although these conditions are not easy to check [6].

As our focus is the one-dimensional case, we organize our paper as follows: In Sect. 2, we describe the factorization of the Lebesgue measure on $[0, 1]$ given by Lewis and, for the reader’s convenience, we provide a somewhat different proof of the factorization theorem that avoids some of the complications of the original ones stemming from the use of probabilistic tools. We then prove the spectral property in Sect. 3 and discuss the **generalized Fuglede’s Conjecture** on \mathbb{R}^1 in Sect. 4. We will finally prove Theorem 1.1 in higher dimension in Sect. 5. As this piece of work offers us several new directions for further research, we end this paper with some remarks and open question in Sect. 6.

Note: During the preparation of the manuscript, we were made aware that Professor Xinggang He and his student [1] discovered independently a new class of one-dimensional spectral measures obtained via a Moran construction of fractals. These one-dimensional spectral measures turn out to coincide exactly with those we consider in this paper.

2 Factorization of Lebesgue Measures

Let $\mathcal{L}_{[0,1]}$ be the Lebesgue measure supported on $[0, 1]$ and let μ and ν be two Borel probability measures supported on $[0, 1]$. We say that (μ, ν) is a *complementary pair* of measures with respect to $\mathcal{L}_{[0,1]}$ if

$$\mu * \nu = \mathcal{L}_{[0,1]}.$$

Let $\mathcal{N} = \{N_k\}_{k=1}^\infty$ be a sequence of positive integers greater than or equal to 2. We associate with \mathcal{N} the discrete measures

$$\nu_k = \frac{1}{N_k} \sum_{j=0}^{N_k-1} \delta_{\frac{j}{N_1 \cdots N_k}}, \quad k \geq 1. \quad (2.1)$$

For a given Borel set E of finite positive Lebesgue measure, recall that \mathcal{L}_E is the normalized Lebesgue measure supported on E . We now observe that the Lebesgue measure supported on $[0, 1]$ admits a natural decomposition as convolution products.

$$\begin{aligned} \mathcal{L}_{[0,1]} &= \nu_1 * \left(\mathcal{L}_{\left[0, \frac{1}{N_1}\right]} \right) \\ &= \nu_1 * \nu_2 * \left(\mathcal{L}_{\left[0, \frac{1}{N_1 N_2}\right]} \right) \\ &= \dots \\ &= \nu_1 * \nu_2 * \dots * \nu_k * \left(\mathcal{L}_{\left[0, \frac{1}{N_1 \dots N_k}\right]} \right). \end{aligned}$$

The sequence of measures $\nu_1 * \nu_2 * \dots * \nu_k$ converges weakly to $\mathcal{L}_{[0,1]}$. Therefore, one can write the Lebesgue measure as an infinite convolution of discrete measures.

$$\mathcal{L}_{[0,1]} = \nu_1 * \nu_2 * \dots \tag{2.2}$$

Given a set \mathcal{N} as above, we will consider two types of factorization (type I and type II) of $\mathcal{L}_{[0,1]}$ as the convolution of two measures obtained from the infinite factorization obtained in (2.2).

Type I. There exists a finite positive integer k such that we have either

$$\mu_{\mathcal{N}} = \nu_1 * \nu_3 * \dots * \nu_{2k-1} \text{ and } \nu_{\mathcal{N}} = \nu_2 * \nu_4 * \dots * \nu_{2k} * \left(\mathcal{L}_{\left[0, \frac{1}{N_1 N_2 \dots N_{2k}}\right]} \right)$$

or

$$\mu_{\mathcal{N}} = \nu_1 * \nu_3 * \dots * \nu_{2k-1} * \left(\mathcal{L}_{\left[0, \frac{1}{N_1 N_2 \dots N_{2k}}\right]} \right) \text{ and } \nu_{\mathcal{N}} = \nu_2 * \nu_4 * \dots * \nu_{2k}.$$

Type II

$$\mu_{\mathcal{N}} = \nu_1 * \nu_3 * \dots * \nu_{2k-1} * \dots \tag{2.3}$$

$$\nu_{\mathcal{N}} = \nu_2 * \nu_4 * \dots * \nu_{2k} \dots \tag{2.4}$$

Remark 2.1 The reader might want to construct more general decompositions obtained by choosing other factorizations of (2.2), but note that if convolution product of two consecutive factors of (2.2) belong to the same factor in the factorization, say ν_k and ν_{k+1} , then we have

$$\nu_k * \nu_{k+1} = \frac{1}{N_k N_{k+1}} \sum_{j=0}^{N_k N_{k+1} - 1} \delta_{j/N_1 N_2 \dots (N_k N_{k+1})}$$

and we would then be able to write the given convolution product as one of type I or type II associated with a different \mathcal{N} .

Note in both cases that $\mu_{\mathcal{N}} * \nu_{\mathcal{N}} = \mathcal{L}_{[0,1]}$ by (2.2). Therefore, they are $\mu_{\mathcal{N}}$ and $\nu_{\mathcal{N}}$ form a complementary pair with respect to $\mathcal{L}_{[0,1]}$. In the case of the type I decomposition, one is purely discrete and one is absolutely continuous while in the type II

decomposition, both factors are singularly continuous measures. We say that a complementary pair (μ, ν) is *natural* if we can find a sequence \mathcal{N} of positive integers such that $(\mu, \nu) = (\mu_{\mathcal{N}}, \nu_{\mathcal{N}})$.

Theorem 2.2 *If μ and ν are positive Borel probability measures supported on $[0, 1]$ and $\mu * \nu = \mathcal{L}_{[0,1]}$, then μ and ν are natural complementary pair.*

This theorem is essentially due to Lewis [16] who considered the problem in probability consisting in characterizing the type of the distributions of pairs of independent random variables X and Y whose sum $X + Y$ is a uniform random variable on $[-\pi, \pi]$. For the reader’s convenience, we will give here another proof based on his ideas as his result is not widely known. Moreover, the proof we give here is more analytical in flavor and avoids some of the complications arising in the original proof from the use of probability tools. The main important step of the proof is to show that if two probability measures μ and ν satisfy $\mu * \nu = \mathcal{L}_{[0,1]}$, then one of them, say μ , must be “ $1/N$ periodic” in the sense that $\mu = \left(1/N \sum_{j=0}^{N-1} \delta_{j/N}\right) * \mu_1$ for some integer $N \geq 2$ and $\mu_1 * \nu = \mathcal{L}_{[0,1/N]}$. This is done by analyzing the structure of the zeros of the Fourier transform of μ and ν (Lemma 2.5).

We now define the (complex) Fourier transform of a compactly supported probability measure μ by the formula

$$\widehat{\mu}(\xi) = \int e^{-2\pi i \xi x} d\mu(x), \quad \xi \in \mathbb{C}.$$

We will consider convolution products yielding the Lebesgue measure supported on $[-1/2, 1/2]$ instead of $[0, 1]$ to exploit some symmetric properties of the solutions (as explained below). Note that $\mu * \nu = \mathcal{L}_{[-1/2, 1/2]}$ is equivalent to

$$\widehat{\mu}(\xi)\widehat{\nu}(\xi) = \widehat{\mathcal{L}_{[-1/2, 1/2]}}(\xi) = \frac{\sin \pi \xi}{\pi \xi}. \tag{2.5}$$

The zero set of the Fourier transform $\widehat{\mu}$ in the complex plane will be denoted by

$$\mathcal{Z}(\widehat{\mu}) = \{\xi \in \mathbb{C} : \widehat{\mu}(\xi) = 0\}.$$

Since $(\delta_x * \mu) * (\delta_{-x} * \nu) = \mathcal{L}_{[-1/2, 1/2]}$ for any real numbers x , we may assume the smallest closed interval containing the support of μ is given by $[-a, a]$. Denote by $\text{supp } \mu$ the closed support of μ . Given a probability measure ρ , we also define the measure $\check{\rho}$ to be the measure satisfying $\check{\rho}(B) = \rho(-B)$ for any Borel set $B \subset \mathbb{R}$.

Lemma 2.3 *Let μ and ν be two probability measures such that $\mu * \nu = \mathcal{L}_{[-1/2, 1/2]}$ and assume that the smallest closed interval containing $\text{supp } \mu$ is of the form $[-a, a]$, $a > 0$. Then we have*

$$\mathbb{Z} \setminus \{0\} = \mathcal{Z}(\widehat{\mu}) \cup \mathcal{Z}(\widehat{\nu}) \text{ (as a disjoint union)}. \tag{2.6}$$

Moreover, the smallest closed interval containing $\text{supp } \nu$ is given by $[-b, b]$ where $b = 1/2 - a$ and both μ and ν have symmetric distributions around the origin (i.e. $\check{\mu} = \mu$ and $\check{\nu} = \nu$).

Proof It is well-known that $\widehat{\mu}$ is a non-zero entire analytic function, so its zero set is a discrete set in the complex plane. Furthermore, since the zeros of $\chi_{[-1/2, 1/2]}$ are simple, (2.6) follows from (2.5). Let $[c, b]$ be the smallest closed interval containing the support of ν . Then $a + b = 1/2$ and $-a + c = -1/2$ showing that $c = -b$ and $b = 1/2 - a$.

Finally, note that, since μ is a positive measure, $\mathcal{Z}((\check{\mu})^\wedge) = \mathcal{Z}(\widehat{\mu})$. Therefore, $\mathcal{Z}((\check{\mu})^\wedge) \cup \mathcal{Z}(\widehat{\nu}) = \mathbb{Z} \setminus \{0\}$. Consider the tempered distribution $\rho := \check{\mu} * \nu * \delta_{\mathbb{Z}}$. Then $\widehat{\rho} = (\check{\mu})^\wedge \cdot \widehat{\nu} \cdot \delta_{\mathbb{Z}} = \delta_0$. Hence, ρ is the Lebesgue measure on \mathbb{R} and the restriction of ρ to the interval $[-1/2, 1/2]$ is $\check{\mu} * \nu$. This shows that $\check{\mu} * \nu = \mathcal{L}_{[-1/2, 1/2]}$, which means that $\check{\mu} * \nu = \mu * \nu$. Taking Fourier transform, we obtain $\check{\mu} = \mu$. The proof of the symmetry of ν is similar. □

Note that Lewis used the Hadamard factorization theorem to prove the symmetry property of μ and ν in Lemma 2.3. The ideas of the following two lemmas are due to Lewis and form the crucial parts of the argument.

Lemma 2.4 *Let $r \geq 1$ be the smallest positive zero of $\widehat{\mu}$. Then*

$$\frac{1}{4r} \leq a \leq \frac{1}{2r} \text{ and } \frac{1}{2} - \frac{1}{2r} \leq b \leq \frac{1}{2} - \frac{1}{4r}.$$

Proof We just need to prove the lower estimates for both a and b as the upper ones will follow from these and the fact that $a + b = 1/2$. Since r is a zero of $\widehat{\mu}$, then $-r$ is also a zero and we must have $\int \cos(2\pi r x) d\mu(x) = 0$. This implies that $2\pi r a \geq \frac{\pi}{2}$ and thus $a \geq \frac{1}{4r}$. In particular, the claim is true for $r = 1$.

For the upper bound, we consider the following functions for different r .

$$h(x) := \begin{cases} \cos(2\pi x), & r = 2; \\ \cos(2\pi x) - \cos(2\pi 2x), & r = 3; \\ \cos\left(\frac{\pi r x}{2}\right) \prod_{j=1}^{k-1} \left(\cos(2\pi x) - \cos\frac{2(2j-1)\pi}{r}\right), & r > 2, r = 2k; \\ \cos\left(\frac{\pi(r-1)x}{2}\right) - \cos\left(\frac{\pi(r+1)x}{2}\right) \prod_{j=1}^{k-2} \left(\cos(2\pi x) - \cos\frac{(2\pi)(2j)}{r}\right), & r > 2, r = 2k - 1. \end{cases}$$

By expanding $h(x)$, we see that $h(x)$ is a linear combination of $\cos(2\pi k x)$, for $k = 1, \dots, r - 1$. Hence $\int h(x) d\nu(x) = 0$ as $1, \dots, r - 1$ are zeros of $\widehat{\nu}$. By checking the sign of each factor, we see that if $2\pi x \leq \pi(r - 1)/r$, then $h(x) \geq 0$.

Consider the case where $r > 2$ is even. We have either $2\pi b \geq \pi(r - 1)/r$ (i.e. $b \geq 1/2 - 1/2r$) or ν is supported on the atoms $\pm(1/r), \dots, \pm(r - 3)/r$. However, ν cannot be supported on those atoms since $\widehat{\nu}$ would be a polynomial in $\cos(2\pi x/r)$ of degree at most $r - 3$, but there are $r - 1$ zeros for $\widehat{\nu}$, a contradiction. Therefore, we must have $b \geq 1/2 - 1/(2r)$. The proof for the other cases follows from a similar argument. □

Lemma 2.5 *Let $N > 0$ be a positive integer and let μ and ν be two probability measures on \mathbb{R} such that $\mu * \nu = \mathcal{L}_{[0,1/N]}$ with neither $\widehat{\mu}$ nor $\widehat{\nu}$ being identically one. Suppose that $N \in \mathcal{Z}(\widehat{\nu})$ and let Nr with $r > 1$ be the smallest positive zero of $\widehat{\mu}$. Then*

$$\mathcal{Z}(\widehat{\mu}) \subset Nr\mathbb{Z}.$$

Proof By rescaling the measures by a factor of N , it is easy to see that it suffices to consider the case $N = 1$. By translating the measure (i.e. $\mu * (\delta_{-1/2} * \nu) = \mathcal{L}_{[-1/2,1/2]}$), it suffices to prove the lemma for the case $\mu * \nu = \mathcal{L}_{[-1/2,1/2]}$, where $\check{\mu} = \mu$ and $\check{\nu} = \nu$.

Let $\rho(E) = \nu(\{0\})\delta_0(E) + 2\nu(E \cap (0, 1/2])$ and $\check{\rho}(E) = \rho(-E)$ for E Borel. Then, the fact that $\nu(E) = \nu(-E)$ implies that $\rho + \check{\rho} = 2\nu$. Therefore,

$$\mu * \rho + \mu * \check{\rho} = 2\mathcal{L}_{[-1/2,1/2]}. \tag{2.7}$$

This implies, in particular, that $\mu * \rho$ is absolutely continuous with respect to the Lebesgue measure and we can let $g(x) \geq 0$ be its density. Then $g(-x)$ is the density of $(\mu * \check{\rho}) = \mu * \check{\rho}$. By (2.7), we have a.e.

$$g(x) + g(-x) = 2.$$

As $\text{supp}(\mu * \check{\rho})$ (and hence $\text{supp} g(-x)$) is contained in $[-1/2, a]$, $g(x) = 2$ on $[a, 1/2]$. We may therefore write

$$\begin{aligned} g &= 2\chi_{[a,1/2]} + g\chi_{[-a,a]} = 2\chi_{[a,1/2]} + g\chi_{[-a,0]} + (2 - g(-x))\chi_{[0,a]} \\ &= 2\chi_{[0,1/2]} + (g\chi_{[-a,0]} - g(-x)\chi_{[0,a]}). \end{aligned}$$

Note that $2\chi_{[0,1/2]}$ is the density of the measure $\mathcal{L}_{[0,1/2]}$. Taking Fourier transform, we have

$$\widehat{\mu}(\xi)\widehat{\rho}(\xi) = \widehat{g}(\xi) = \widehat{\mathcal{L}_{[0,1/2]}}(\xi) + 2i \int_0^a g(-x) \sin(2\pi\xi x) dx \tag{2.8}$$

Suppose that r is even. As $\widehat{\mu}(r) = 0$, we must have

$$\int_0^a g(-x) \sin(2\pi r x) dx = 0.$$

Since $a \leq 1/2r$ by Lemma 2.4, we have $\sin(2\pi r x) \geq 0$ on $[0, a]$ and thus $g(-x) = 0$ there. Thus, (2.8) implies that

$$\widehat{\mu}(\xi)\widehat{\rho}(\xi) = \widehat{\mathcal{L}_{[0,1/2]}}(\xi). \tag{2.9}$$

Hence, $\mathcal{Z}(\widehat{\mu}) \subset 2\mathbb{Z}$.

Writing $r = 2^n m$ where m is odd, we deduce from the above argument that $\mathcal{Z}(\widehat{\mu}) \subset 2\mathbb{Z}$. Consider the measure $\mu_1(E) = \mu(E/2)$ and $\rho_1(E) = \rho(E/2)$ we have $\widehat{\mu}_1(\xi) = \widehat{\mu}(2\xi)$ and $\widehat{\rho}_1(\xi) = \widehat{\rho}(2\xi)$. By (2.9), we have $\widehat{\mu}_1(\xi)\widehat{\rho}_1(\xi) = \widehat{\mathcal{L}_{[0,1]}}(\xi)$ (i.e. $\mu_1 * (\delta_{-1/2} * \rho_1) = \mathcal{L}_{[-1/2, 1/2]}$). Moreover, $\mathcal{Z}(\widehat{\mu}_1) = \frac{1}{2}\mathcal{Z}(\widehat{\mu})$. In this case, the smallest positive zero of $\widehat{\mu}_1$ will be $2^{n-1}m$. Therefore, repeating the above argument, we have $\mathcal{Z}(\widehat{\mu}) \subset 2^n\mathbb{Z}$ and the proof will be finished if we can prove our claim if r is odd.

Suppose now that r is odd. We consider the measures $\nu_1(E) = \nu(E \cap [-a, b])$ and $\nu_2(E) = \nu(E \cap [-b, -a])$ (Here, it is more convenient not to normalize ν_1 and ν_2 as probability measures). We have then $\nu = \nu_1 + \nu_2$ and $\mathcal{L}_{[-1/2, 1/2]} = \mu * \nu_1 + \mu * \nu_2$. Let g_1 and g_2 be the density of $\mu * \nu_1$ and $\mu * \nu_2$ respectively. The above implies that

$$g_1(x) + g_2(x) = 1 \text{ a.e. on } [-1/2, 1/2].$$

Note that the $\text{supp } g_1$ is contained in $[-2a, 1/2]$ and $\text{supp } g_2$ is contained in $[-1/2, 0]$. It follows that $g_1 = 1$ almost everywhere on $[0, 1/2]$. We may therefore write

$$g_1 = \chi_{[0, 1/2]} + g_1\chi_{[-2a, 0]}.$$

Taking Fourier transforms and noting that $\widehat{g_1}(\xi) = \widehat{\mu}(\xi)\widehat{\nu_1}(\xi)$, we obtain

$$\widehat{\mu}(\xi)\widehat{\nu_1}(\xi) = \widehat{\chi_{[0, 1/2]}}(\xi) + \int_0^{2a} g_1(-x)e^{2\pi i \xi x} dx. \tag{2.10}$$

As $\widehat{\mu}(r) = 0$, by substituting $\xi = r$ and equating the imaginary parts, we have

$$\frac{1}{\pi r} = \int_0^{2a} g_1(-x) \sin(2\pi r x) dx.$$

By Lemma 2.4, $2a \geq 1/2r$ and therefore,

$$\begin{aligned} \frac{1}{\pi r} &= \int_0^{1/2r} g_1(-x) \sin(2\pi r x) dx + \int_{1/2r}^{2a} g_1(-x) \sin(2\pi r x) dx \\ &\leq \int_0^{1/2r} g_1(-x) \sin(2\pi r x) dx \quad (\text{as } \sin(2\pi r x) \leq 0 \text{ on } [1/2r, 2a]) \\ &\leq \int_0^{1/2r} \sin(2\pi r x) dx = \frac{1}{\pi r} \quad (\text{as } g_1(-x) \leq 1). \end{aligned}$$

Hence, we must have $g_1(-x) = 1$ on $[0, 1/2r]$ and $\int_{1/2r}^{2a} g_1(-x) \sin(2\pi r x) dx = 0$, which implies that $g_1(-x) = 0$ on $[1/2r, 2a]$. Considering the real part of the Eq. (2.10) and noting that $\widehat{\mu}(\xi)$ is real-valued (as $\check{\mu} = \mu$), we have

$$\widehat{\mu}(\xi)\text{Re}(\widehat{\nu}_2(\xi)) = \frac{\sin \pi \xi}{2\pi \xi} + \int_0^{1/2r} \cos(2\pi \xi x) dx = \frac{1}{2\pi \xi} \left(\sin \pi \xi + \sin \frac{\pi \xi}{r} \right).$$

Since $\mathcal{Z}(\widehat{\mu}) \subset \mathbb{Z}$, the previous equation shows that in fact $\mathcal{Z}(\widehat{\mu}) \subset r\mathbb{Z}$, completing the proof. □

Proof of Theorem 2.2 Let (μ, ν) be a complementary pair with respect to $\mathcal{L}_{[0,1]}$. We may assume that $\widehat{\nu}(1) \neq 0$ and we let $N_1 > 1$ be the smallest positive zero of $\widehat{\nu}$. We have $\mathcal{Z}(\widehat{\nu}) \subset N_1\mathbb{Z}$ by Lemma 2.5. As the zero sets of $\widehat{\mu}$ and $\widehat{\nu}$ are disjoint (see (2.6)), the set $\{k \in \mathbb{Z} : \widehat{\mu}(k) \neq 0\}$ is contained in $N_1\mathbb{Z}$.

Consider the periodization of the measure μ defined by $\mu_p = \mu * \delta_{\mathbb{Z}}$. Its distributional Fourier transform (as a tempered distribution) is given by

$$\widehat{\mu}_p = \widehat{\mu} \cdot \delta_{\mathbb{Z}} = \widehat{\mu} \cdot \delta_{N_1\mathbb{Z}}$$

Hence, μ_p is indeed $1/N_1$ -periodic. It follows immediately that

$$\mu = \nu_1 * \alpha_1 \quad \text{and} \quad \nu * \alpha_1 = \mathcal{L}_{[0,1/N_1]} \tag{2.11}$$

where $\nu_1 = \frac{1}{N_1} \sum_{j=0}^{N_1-1} \delta_{j/N_1}$ and $\alpha_1(E) = N_1\mu(E \cap [0, 1/N_1])$ for any Borel set E . The case where α_1 is the Dirac measure at the origin immediately yields a type I decomposition. Otherwise, we apply Lemma 2.5 on the pair (ν, α_1) . Since $\widehat{\nu}(N_1) = 0$, we have $\widehat{\alpha}_1(N_1) \neq 0$ and we can let N_2 be the smallest positive integer such that $\widehat{\alpha}_1(N_1N_2) = 0$. By Lemma 2.5, we have $\mathcal{Z}(\widehat{\alpha}_1) \subset N_1N_2\mathbb{Z}$. We obtain

$$\mu = \nu_1 * \alpha_1, \quad \nu = \nu_2 * \alpha_2 \quad \alpha_1 * \alpha_2 = \mathcal{L}_{[0,1/N_1N_2]}$$

where $\nu_2 = \frac{1}{N_2} \sum_{j=0}^{N_2-1} \delta_{j/N_1N_2}$. The case where α_2 is a Dirac measure at the origin yields again a type I decomposition. Otherwise, we continue this inductive process and define recursively the probability measures $\alpha_k, k \geq 1$. If $\alpha_k = \delta_0$ for some k , the process stops and we have arrived at a type I decomposition. If $\alpha_k \neq \delta_0$ for all k , we have then expressed both measures μ and ν at the infinite convolution products $\mu = \nu_1 * \nu_3 * \dots, \quad \nu = \nu_2 * \nu_4 * \dots$, which yields a type II decomposition. □

Theorem 2.2 also gives us a new proof of classification of the set A and B such that $A \oplus B = \{0, \dots, n - 1\}$ which was proved in [17] and [20] using a theorem of De Bruijn.

Corollary 2.6 *Let $\mathcal{E}_n = \{0, 1, \dots, n - 1\}$ and let \mathcal{A} and \mathcal{B} be two finite set of integers such that $\mathcal{A} \oplus \mathcal{B} = \{0, \dots, n - 1\}$. Suppose that $1 \in \mathcal{A}$. Then there exist integers N_1, \dots, N_{2k} such that $N_1 \dots N_{2k} = n$ and*

$$\begin{aligned} A &= \mathcal{E}_{N_0} \oplus N_0N_1\mathcal{E}_{N_2} \oplus \dots \oplus N_0N_1 \dots N_{2k-1}\mathcal{E}_{2k}, \\ B &= N_0\mathcal{E}_{N_1} \oplus N_0N_1N_2\mathcal{E}_{N_3} \oplus \dots \oplus N_0N_1 \dots N_{2k-2}\mathcal{E}_{2k-1}. \end{aligned}$$

Proof As $\mathcal{A} \oplus \mathcal{B} = \{0, \dots, n - 1\}$, we have

$$\left(\frac{1}{|\mathcal{A}|} \delta_{\frac{1}{n}\mathcal{A}}\right) * \left(\frac{1}{|\mathcal{B}|} \delta_{\frac{1}{n}\mathcal{B}}\right) * \mathcal{L}_{[0,1/n]} = \mathcal{L}_{[0,1]}.$$

By Theorem 2.2, the measures $\mu = \left(\frac{1}{|\mathcal{A}|} \delta_{\frac{1}{n}\mathcal{A}}\right)$ and $\nu = \left(\frac{1}{|\mathcal{B}|} \delta_{\frac{1}{n}\mathcal{B}}\right) * \mathcal{L}_{[0,1/n]}$ are natural complementary pair. As one of them is discrete and the other is absolutely continuous, they correspond to a type I decomposition. Since $1 \in \mathcal{A}$, we have thus $1/n \in \frac{1}{n}\mathcal{A}$. By comparing the support of the measures, we obtain the existence of integers N'_1, N'_2, \dots such that

$$\begin{aligned} \frac{1}{n}\mathcal{A} &= \frac{1}{N'_1} \mathcal{E}_{N'_1} \oplus \frac{1}{N'_1 N'_2 N'_3} \mathcal{E}_{N'_3} \oplus \dots \oplus \frac{1}{N'_1 N'_2 \dots N'_{2k-1}} \mathcal{E}_{N'_{2k-1}}, \\ \frac{1}{n}\mathcal{B} &= \frac{1}{N'_1 N'_2} \mathcal{E}_{N'_2} \oplus \frac{1}{N'_1 N'_2 N'_3 N'_4} \mathcal{E}_{N'_4} \oplus \dots \oplus \frac{1}{N'_1 N'_2 \dots N'_{2k}} \mathcal{E}_{N'_{2k}}, \end{aligned}$$

and $n = N'_1 \dots N'_{2k}$. Letting $N_r = N'_{2k-r}$, we obtain the desired factorization. □

3 The Spectral Property

In this section, we show that all measures appearing in natural complementary pairs are spectral measures. Recall that a Borel probability measure μ is called a *spectral measure* with associated *spectrum* Λ if the collection of exponentials $E(\Lambda) = \{e^{2\pi i \lambda x}\}_{\lambda \in \Lambda}$ forms an orthonormal basis for $L^2(\mu)$. It is easy to see that $E(\Lambda)$ is an orthonormal set in $L^2(\mu)$ if and only if

$$\Lambda - \Lambda \subset \mathcal{Z}(\widehat{\mu}) \cup \{0\}.$$

By a well-known result in [8], Λ is a spectrum of μ if and only if

$$Q(\xi) := \sum_{\lambda \in \Lambda} |\widehat{\mu}(\xi + \lambda)|^2 \equiv 1. \tag{3.1}$$

In fact, if $E(\Lambda)$ is an orthonormal set, $Q(\xi) \leq 1$ and Q is an entire function of exponential type ([8], see also [2]). Let $\mathcal{N} = \{N_i\}_{i=1}^\infty$ be a collection of positive integers and consider the type I and II decomposition as in the previous section. Let

$$\mu^{(k)} = \nu_1 * \nu_3 * \dots * \nu_{2k-1}, \quad \nu^{(k)} = \nu_2 * \nu_4 * \dots * \nu_{2k}$$

and, for a given \mathcal{N} , we let $A_1 = \{0, \dots, N_1 - 1\}$ and $A_n = N_1 \dots N_{n-1} \cdot \{0, \dots, N_n - 1\}$ for $n \geq 2$. We start with a simple observation.

Proposition 3.1 *Each ν_n is a spectral measure with spectrum A_n . For all $k \geq 1$, $\mu^{(k)}$ is a spectral measure with spectrum given by*

$$\Lambda_k = \bigoplus_{j=1}^k A_{2j-1}. \tag{3.2}$$

In particular, the type I natural complementary pair $\mu_{\mathcal{N}}$ and $\nu_{\mathcal{N}}$ defined in the previous section are spectral measures.

Proof It is immediate to see that the measure $\frac{1}{N_n} \sum_{j=0}^{N_n-1} \delta_{j/N_n}$ is a spectral measure with spectrum $\{0, \dots, N_n - 1\}$. Therefore, $\nu_n = \frac{1}{N_n} \sum_{j=0}^{N_n-1} \delta_{j/(N_1 \dots N_n)}$ is a spectral measure with spectrum $N_1 \dots N_{n-1} \cdot \{0, \dots, N_n - 1\} = A_n$.

Note that $\mathcal{Z}(\widehat{\nu_n}) = N_1 N_2 \dots N_n \mathbb{Z} \setminus N_1 N_2 \dots N_{n-1} \mathbb{Z}$ and

$$\widehat{\mu^{(k)}}(\xi) = \prod_{j=1}^k \widehat{\nu_{2j-1}}(\xi).$$

For notational convenience, we define $N_0 = 1$. Taking distinct $\lambda_1, \lambda_2 \in \Lambda_k$ and writing $\lambda_\ell = \sum_{j=1}^k r_{\ell,j} N_1 N_2 \dots N_{2j-2}$, for $\ell = 1, 2$, we have

$$\lambda_1 - \lambda_2 = \sum_{j=1}^k (r_{1,j} - r_{2,j}) N_1 N_2 \dots N_{2j-2} = \sum_{j=J}^k s_j N_1 N_2 \dots N_{2j-2},$$

where J is the first index such that $r_{1,j} \neq r_{2,j}$ and $-(N_{2J-1} - 1) \leq s_J \leq N_{2J-1} - 1$. so $\widehat{\nu_{2J-1}}(\lambda_1 - \lambda_2) = \widehat{\nu_{2J-1}}(N_1 \dots N_{2J-2} s_J) = 0$. Therefore, $\widehat{\mu^{(k)}}(\lambda_1 - \lambda_2) = 0$. This proves the orthogonality of $E(\Lambda_k)$ in $L^2(\mu^{(k)})$. As $L^2(\mu^{(k)})$ is a finite dimensional vector space of dimension $N_1 N_3 \dots N_{2k-1} = \text{card}(E(\Lambda))$, the collection $E(\Lambda)$ must be complete in $L^2(\mu^{(k)})$.

To prove the last statement, we just consider the case where $\mu_{\mathcal{N}} = \mu^{(k)}$ and $\nu_{\mathcal{N}} = \nu^{(k)} * (\mathcal{L}_{[0, \frac{1}{N_1 \dots N_{2k}}]})$, as the case $\mu_{\mathcal{N}} = \nu_1 * \nu_3 * \dots * \nu_{2k-1} * (\mathcal{L}_{[0, \frac{1}{N_1 N_2 \dots N_{2k}}]})$ and $\nu_{\mathcal{N}} = \nu_2 * \nu_4 * \dots * \nu_{2k}$ is similar.

It is easily seen, as before, that $\nu^{(k)}$ is also a discrete spectral measure with spectrum

$$\widetilde{\Lambda}_k = \bigoplus_{j=1}^k A_{2j}.$$

Moreover, $\widehat{\nu^{(k)}}$ is $N_1 \dots N_{2k}$ -periodic. Let α denote the measure $\mathcal{L}_{[0, \frac{1}{N_1 N_2 \dots N_{2k}}]}$. Then α has $N_1 N_2 \dots N_{2k} \mathbb{Z}$ as a spectrum. It follows that

$$\begin{aligned} \sum_{\lambda \in \widetilde{\Lambda}_k + N_1 \dots N_{2k} \mathbb{Z}} |\widehat{\nu_{\mathcal{N}}}(\xi + \lambda)|^2 &= \sum_{\lambda \in \widetilde{\Lambda}_k, m \in \mathbb{Z}} \left| \widehat{\nu^{(k)}}(\xi + \lambda + N_1 \dots N_{2k} m) \right|^2 |\widehat{\alpha}(\xi + \lambda + N_1 \dots N_{2k} m)|^2 \\ &= \sum_{\lambda \in \widetilde{\Lambda}_k} \left| \widehat{\nu^{(k)}}(\xi + \lambda) \right|^2 \cdot \sum_{m \in \mathbb{Z}} |\widehat{\alpha}(\xi + \lambda + N_1 \dots N_{2k} m)|^2 \equiv 1. \end{aligned}$$

Hence, $\nu_{\mathcal{N}}$ a spectral measure with spectrum $\widetilde{\Lambda}_k + N_1 \cdots N_{2k}\mathbb{Z}$. □

It remains to deal with the spectral property for complementary pairs $\mu_{\mathcal{N}}$ and $\nu_{\mathcal{N}}$ of type II. Since these two measures have essentially the same form, we will discuss only the case $\mu := \mu_{\mathcal{N}}$. Note that the measure μ will be the weak limit of the measures $\mu^{(k)}$ and

$$\widehat{\mu}(\xi) = \prod_{j=1}^{\infty} \widehat{\nu_{2j-1}}(\xi) = \widehat{\mu^{(k)}}(\xi) \cdot \prod_{j=k+1}^{\infty} \widehat{\nu_{2j-1}}(\xi). \tag{3.3}$$

Here we recall that $\nu_{2j-1} = \frac{1}{N_{2j-1}} \sum_{r=0}^{N_{2j-1}-1} \delta_{\frac{r}{N_1 \cdots N_{2j-1}}}$ and its Fourier transform is given by

$$\widehat{\nu_{2j-1}}(\xi) = e^{-\pi i(N_{2j-1}-1)\xi/(N_1 \cdots N_{2j-1})} \frac{\sin(\pi\xi/(N_1 \cdots N_{2j-2}))}{N_{2j-1} \sin(\pi\xi/(N_1 \cdots N_{2j-1}))}. \tag{3.4}$$

Let

$$\Lambda_{\mu} = \bigoplus_{j=1}^{\infty} A_{2j-1} = \bigcup_{k=1}^{\infty} \Lambda_k,$$

(with only finite sums of elements of A_{2j-1} , $j \geq 1$, appearing in Λ_{μ}). The exponentials $\{e^{2\pi i\lambda x}\}_{\lambda \in \Lambda_{\mu}}$ are mutually orthogonal in $L^2(\mu)$ by Proposition 3.1. Our goal is verify (3.1). To do this, we note that, as Q is an entire function, we just need to show that $Q(\xi) \equiv 1$ on a neighborhood of 0. Let

$$Q_k(\xi) = \sum_{\lambda \in \Lambda_k} |\widehat{\mu}(\xi + \lambda)|^2.$$

Now, we fix two positive integers n and p . By (3.3) and the fact that $\{\Lambda_k\}_{k \geq 1}$ is an increasing sequence of sets,

$$\begin{aligned} Q_{n+p}(\xi) &= Q_n(\xi) + \sum_{\lambda \in \Lambda_{n+p} \setminus \Lambda_n} |\widehat{\mu}(\xi + \lambda)|^2 \\ &= Q_n(\xi) + \sum_{\lambda \in \Lambda_{n+p} \setminus \Lambda_n} \left| \widehat{\mu^{(n+p)}}(\xi + \lambda) \right|^2 \cdot \left| \prod_{j=n+p+1}^{\infty} \widehat{\nu_{2j-1}}(\xi + \lambda) \right|^2. \end{aligned} \tag{3.5}$$

We need the following proposition which provides a crucial estimate for the last term in the previous expression in order to establish the spectral property.

Proposition 3.2 *There exists $c > 0$ such that*

$$\inf_{k \geq 1} \inf_{\lambda \in \Lambda_k} \left| \prod_{j=k+1}^{\infty} \widehat{\nu_{2j-1}}(\xi + \lambda) \right|^2 \geq c$$

for all $|\xi| < 1/2$, where Λ_k is given in (3.2).

Proof Let $\lambda \in \Lambda_k$ and $x_{k,\lambda} = \frac{\xi + \lambda}{N_1 N_2 \dots N_k}$. We first note that, by (3.4),

$$\begin{aligned} \left| \prod_{j=k+1}^{\infty} \widehat{v_{2^{j-1}}}(\xi + \lambda) \right|^2 &= \prod_{j=k+1}^{\infty} \frac{\sin^2(\pi(\xi + \lambda)/(N_1 \dots N_{2^{j-2}}))}{N_{2^{j-1}}^2 \sin^2(\pi(\xi + \lambda)/(N_1 \dots N_{2^{j-1}}))} \\ &= \prod_{j=k+1}^{\infty} \frac{\sin^2(\pi x_{2^{j-2}, \lambda})}{N_{2^{j-1}}^2 \sin^2(\pi x_{2^{j-1}, \lambda})}. \end{aligned} \tag{3.6}$$

Writing $\lambda = \sum_{j=1}^k r_j N_1 N_2 \dots N_{2^{j-2}}$ with $0 \leq r_j \leq N_{2^{j-1}} - 1$, we see immediately that $\lambda \leq N_1 \dots N_{2^{k-1}} - 1$. Hence, we have

$$\frac{\lambda}{N_1 \dots N_{2^k}} \leq \frac{N_1 \dots N_{2^{k-1}} - 1}{N_1 \dots N_{2^k}} \leq \frac{1}{N_{2^k}} \leq \frac{1}{2}.$$

Therefore, for all $|\xi| < 1/2$, we have

$$C := \sup_{k \geq 1} \sup_{\lambda \in \Lambda_k} x_{2^k, \lambda} = \sup_{k \geq 1} \sup_{\lambda \in \Lambda_k} \frac{\xi + \lambda}{N_1 \dots N_{2^k}} < \frac{3}{4}$$

as all $N_j \geq 2$. Note that $N_k x_{k,\lambda} = x_{k-1,\lambda}$ and using two elementary inequalities $\sin x \leq x$ and $\sin x \geq x - \frac{x^3}{3!}$, we have the following estimation for the product in (3.6),

$$\begin{aligned} \prod_{j=k+1}^{\infty} \frac{\sin^2(\pi x_{2^{j-2}, \lambda})}{N_{2^{j-1}}^2 \sin^2(\pi x_{2^{j-1}, \lambda})} &\geq \prod_{j=k+1}^{\infty} \left(1 - \frac{\pi^2}{6} x_{2^{j-2}, \lambda}^2 \right)^2 \\ &= \prod_{j=k+1}^{\infty} \left(1 - \frac{\pi^2}{6} \left(\frac{x_{2^k, \lambda}}{N_{2^{k+1}} \dots N_{2^{j-2}}} \right)^2 \right)^2 \\ &\geq \prod_{j=k+1}^{\infty} \left(1 - \frac{\pi^2}{6} \left(\frac{C}{2^{2(j-k)-2}} \right)^2 \right)^2 \\ &= \prod_{j=1}^{\infty} \left(1 - \frac{3\pi^2}{32} \left(\frac{1}{2^{2j-2}} \right)^2 \right)^2 := c. \end{aligned}$$

As $\sum_{j=1}^{\infty} 1/2^{2j-2} < \infty$ and all factors are positive, $c > 0$ and hence the proof is complete. □

Proof of Theorem 1.1 on \mathbb{R}^1 . In view of Theorem 2.2, we just need to show that all natural complementary pairs are spectral measures. Let \mathcal{N} be a sequence of positive integers greater than or equal to 2. If the pair is of type I, then Proposition 3.1 shows that both factors are spectral measures.

It remains to consider the type II case. Let $\mu_{\mathcal{N}}$ and $\nu_{\mathcal{N}}$ be defined in (2.3) and (2.4). As mentioned before, we only need to prove that $\mu = \mu_{\mathcal{N}}$ is a spectral measure. Let c be the positive number determined in Proposition 3.2. By Proposition 3.1 and (3.1), we have

$$\sum_{\lambda \in \Lambda_{n+p} \setminus \Lambda_n} \left| \widehat{\mu^{(n+p)}}(\xi + \lambda) \right|^2 = 1 - \sum_{\lambda \in \Lambda_n} \left| \widehat{\mu^{(n+p)}}(\xi + \lambda) \right|^2.$$

Using this fact and Proposition 3.2, we obtain from (3.5) that

$$Q_{n+p}(\xi) \geq Q_n(\xi) + c \cdot \left(1 - \sum_{\lambda \in \Lambda_n} \left| \widehat{\mu^{(n+p)}}(\xi + \lambda) \right|^2 \right).$$

Fixing n and letting p go to infinity, it follows that

$$Q(\xi) \geq Q_n(\xi) + c \left(1 - \sum_{\lambda \in \Lambda_n} |\widehat{\mu}(\xi + \lambda)|^2 \right) = Q_n(\xi) + c(1 - Q_n(\xi)).$$

Finally, taking n to infinity, we obtain that $c(1 - Q(\xi)) \leq 0$. But $c > 0$ and $Q(\xi) \leq 1$ because $\{e^{2\pi i \lambda x}\}_{\lambda \in \Lambda}$ is an orthogonal set in $L^2(\mu)$. This show that $Q(\xi) = 1$ for $|\xi| \leq 1/2$ and thus for all $\xi \in \mathbb{R}$ by analyticity, completing the proof.

We now establish the tiling property of the spectra. Suppose that we are given a type I decomposition. Then Proposition 3.1 implies that $\mu_{\mathcal{N}}$ and $\nu_{\mathcal{N}}$ have the following spectra:

$$\Lambda_{\mu} = \bigoplus_{j=1}^k A_{2j-1}, \quad \Lambda_{\nu} = \bigoplus_{j=1}^{k-1} A_{2j} \oplus N_1 \cdots N_{2k-1} \mathbb{Z}.$$

It can be seen immediately that $\Lambda_{\mu} \oplus \Lambda_{\nu} = \{0, 1, \dots, N_{2k-1} - 1\} \oplus N_{2k-1} \mathbb{Z} = \mathbb{Z}$.

Suppose now the decomposition is of type II. Note that the complementary measures have the following spectra using the above notations.

$$\Lambda_{\mu} = \bigoplus_{j=1}^{\infty} A_{2j-1}, \quad \Lambda_{\nu} = \bigoplus_{j=1}^{\infty} A_{2j}.$$

Note that $-\Lambda_{\nu}$ is also spectrum of ν . We now claim that $\Lambda_{\mu} \oplus (-\Lambda_{\nu}) = \mathbb{Z}$. Observe that

$$\begin{aligned} A_1 \oplus (-A_2) &= \{-N_1 N_2 + N_1, \dots, N_1 - 1\}. \\ A_1 \oplus (-A_2) \oplus A_3 &= \{-N_1 N_2 + N_1, \dots, N_1 N_2 N_3 - N_1 N_2 + N_1 - 1\}. \end{aligned}$$

Inductively, the sets $A_1 \oplus (-A_2) \oplus \dots \oplus (-1)^{k-1} A_k$ cover an increasing sequence of consecutive integers. showing that $\Lambda_{\mu} \oplus (-\Lambda_{\nu}) = \mathbb{Z}$. This proves our claim. \square

4 Generalized Fuglede’s Conjecture

In this section, we will formulate a generalization of Fuglede’s conjecture and prove that it implies the original one. Recall the conjecture we are interested in:

Conjecture (Generalized Fuglede’s Conjecture): A compactly supported Borel probability measure μ on \mathbb{R}^1 is spectral if and only if there exists a Borel probability measure ν and a fundamental domain Q of some lattice on \mathbb{R}^1 such that $\mu * \nu = \mathcal{L}_Q$.

We first prove the following proposition.

Proposition 4.1 *Let Ω and Q be bounded measurable sets of positive Lebesgue measure on \mathbb{R}^1 . Suppose that $\mathcal{L}_\Omega * \nu = \mathcal{L}_Q$, for some Borel probability measure ν . Then*

$$\nu = \sum_{k=1}^N \frac{1}{N} \delta_{a_k}, \quad Q = \bigcup_{k=1}^N (\Omega + a_k)$$

and $\mathcal{L}((\Omega + a_k) \cap (\Omega + a_\ell)) = 0$ for all $k \neq \ell$.

Proof We first note that $\mathcal{L}_\Omega * \nu = \mathcal{L}_Q$ if and only if $(\mathcal{L}_\Omega * \delta_y) * (\nu * \delta_x * \delta_{-y}) = (\mathcal{L}_Q * \delta_x)$ for any real numbers x and y . Therefore, there is no loss of generality to assume that the smallest closed intervals containing Ω and Q are respectively $[0, a]$ and $[0, b]$. As $\overline{Q} = \text{supp}(\mathcal{L}_\Omega * \nu) = \overline{\Omega} + \text{supp} \nu$, The support of ν has to be contained in the non-negative part of the real line.

Let $\epsilon > 0$ and consider the interval $E_\epsilon = [0, \epsilon]$. Let $\eta_\epsilon \in E_\epsilon$ be a Lebesgue point of χ_Q . Then, using $\mathcal{L}_\Omega * \nu = \mathcal{L}_Q$,

$$\begin{aligned} \frac{1}{\mathcal{L}(Q)} \mathcal{L}(Q \cap [\eta_\epsilon, \eta_\epsilon + h]) &= \frac{1}{\mathcal{L}(\Omega)} \int_0^{\eta_\epsilon + h} \mathcal{L}(\Omega \cap ([\eta_\epsilon, \eta_\epsilon + h] - y)) d\nu(y) \\ &= \frac{1}{\mathcal{L}(\Omega)} \int_0^{\eta_\epsilon + h} \mathcal{L}((\Omega + y) \cap [\eta_\epsilon, \eta_\epsilon + h]) d\nu(y), \end{aligned}$$

since Ω and $\text{supp} \nu$ are contained in $[0, \infty)$. This implies that

$$\frac{\mathcal{L}(\Omega)}{\mathcal{L}(Q)} \mathcal{L}(Q \cap [\eta_\epsilon, \eta_\epsilon + h]) \leq \mathcal{L}([\eta_\epsilon, \eta_\epsilon + h])\nu([0, \eta_\epsilon + h]) = h\nu([0, \eta_\epsilon + h]).$$

Since η_ϵ is a Lebesgue point of χ_Q , we have $\lim_{h \rightarrow 0} \frac{\mathcal{L}(Q \cap [\eta_\epsilon, \eta_\epsilon + h])}{h} = 1$. Therefore, by taking $h \rightarrow 0$, we deduce that $\frac{\mathcal{L}(\Omega)}{\mathcal{L}(Q)} \leq \nu([0, \eta_\epsilon])$. Letting ϵ approach zero, we obtain the inequality

$$\frac{\mathcal{L}(\Omega)}{\mathcal{L}(Q)} \leq \nu(\{0\}). \tag{4.1}$$

Since $\mathcal{L}(\Omega) > 0$, ν has an atom at 0 and we can write

$$\nu = p_0\delta_0 + (1 - p_0)\nu_1, \quad p_0 = \nu(\{0\}) \text{ and } \nu_1(\{0\}) = 0. \tag{4.2}$$

The equation $\mathcal{L}_\Omega * \nu = \mathcal{L}_Q$ can thus be rewritten as

$$(1 - p_0)\mathcal{L}_\Omega * \nu_1 = \mathcal{L}_Q - p_0\mathcal{L}_\Omega. \tag{4.3}$$

Since the left hand side of (4.3) is still a positive measure, this implies that

$$0 \leq (\mathcal{L}_Q - p_0\mathcal{L}_\Omega)(\Omega) \leq \frac{\mathcal{L}(\Omega)}{\mathcal{L}(Q)} - p_0.$$

Combining this with (4.1), we conclude that $p_0 = \frac{\mathcal{L}(\Omega)}{\mathcal{L}(Q)}$ and, using (4.3), we obtain

$$\mathcal{L}_\Omega * \nu_1 = \mathcal{L}_{Q \setminus \Omega}.$$

If $p_0 = 1$, then $Q = \Omega$ and $\nu = \delta_0$, so we are done. If not, we then repeat the argument with Q replaced by $Q \setminus \Omega$. We can find $\Omega + a_1 \subset Q \setminus \Omega$ such that $p_1 := \nu_1(\{a_1\}) > 0$ and $\nu_1 = p_1\delta_{a_1} + (1 - p_1)\nu_2$. Moreover, $p_1 = \mathcal{L}(\Omega)/\mathcal{L}(Q \setminus \Omega)$. By (4.2),

$$\nu = \frac{\mathcal{L}(\Omega)}{\mathcal{L}(Q)} (\delta_0 + \delta_{a_1}) + (1 - p_1)\nu_2.$$

The theorem will be proved if $p_1 = 1$. Otherwise, we continue this process to obtain a maximal number N of measure disjoint translates of Ω , $\Omega + a_1, \dots, \Omega + a_{N-1}$ such that $Q \supset \bigcup_{k=0}^{N-1} (\Omega + a_k)$. Since $\mathcal{L}(\Omega) > 0$ and $\mathcal{L}(Q) \geq N\mathcal{L}(\Omega)$, N is the largest integer such that $\mathcal{L}(Q) \geq N\mathcal{L}(\Omega)$. We can then write

$$\nu = \frac{\mathcal{L}(\Omega)}{\mathcal{L}(Q)} (\delta_0 + \dots + \delta_{a_{N-1}}) + (1 - p_{N-1})\nu_N.$$

If $p_{N-1} < 1$, we could iterate this process to obtain one more disjoint translate of Ω contained in Q , which is certainly impossible by this choice of N . Hence, $p_{N-1} = 1$. As ν is a probability measure, we must have $\mathcal{L}(\Omega)/\mathcal{L}(Q) = 1/N$. Therefore, the proposition is proved. □

Theorem 4.2 *The validity of generalized Fuglede’s Conjecture implies that of the original Fuglede’s Conjecture on \mathbb{R}^1 .*

Proof Suppose that Ω is a bounded spectral set, then \mathcal{L}_Ω is a spectral measure. By the **generalized Fuglede’s Conjecture**, we can find a probability measure ν and a fundamental domain Q of some lattice Γ such that

$$\mathcal{L}_\Omega * \nu = \mathcal{L}_Q.$$

By Proposition 4.1, ν is a purely discrete measure that can be written as $\nu = \frac{1}{\#\mathcal{A}} \delta_{\mathcal{A}}$ for some finite discrete subset \mathcal{A} and

$$Q = \bigcup_{a \in \mathcal{A}} (\Omega + a).$$

As Q is a fundamental domain Q of the lattice Γ , Ω is a translational tile with tiling set given by $\mathcal{A} + \Gamma$.

Conversely, suppose that Ω is a bounded translational tile with tiling set \mathcal{J} . By the result of Lagarias and Wang [13], all tiling sets on \mathbb{R}^1 are periodic. This implies that we can find a finite set $A \subset \mathbb{R}$ and a lattice Γ such that $\mathcal{J} = \mathcal{A} + \Gamma$. This means that the set $Q = \Omega + \mathcal{A}$ is a fundamental domain of Γ . Letting $\nu = \frac{1}{\#\mathcal{A}}\delta_{\mathcal{A}}$, $\mathcal{L}_{\Omega} * \nu = \mathcal{L}_Q$. By the **generalized Fuglede’s Conjecture**, \mathcal{L}_{Ω} is a spectral measure and Ω is a spectral set. □

5 The Higher Dimensional Case

Let μ_1, \dots, μ_d be Borel probability measures on \mathbb{R}^1 . The Cartesian product of these measures is the unique Borel probability measure $\mu_1 \otimes \dots \otimes \mu_d$ on \mathbb{R}^d such that

$$(\mu_1 \otimes \dots \otimes \mu_d)(E_1 \times \dots \times E_d) = \prod_{i=1}^d \mu_i(E_i),$$

for any Borel sets E_i , $1 \leq i \leq d$, on \mathbb{R}^1 . In this section, we characterize the measures μ and ν on \mathbb{R}^d which are solutions of the equation

$$\mu * \nu = \mathcal{L}_{[0,1]^d}. \tag{5.1}$$

as Cartesian products of the measures satisfying the corresponding one-dimensional equation.

Theorem 5.1 *Let μ and ν be compactly supported probability measures on \mathbb{R}^d . Then μ and ν are solutions to (5.1) if and only if there exists compactly supported Borel probability measures $\{\sigma_i\}_{i=1}^d$ and $\{\tau_i\}_{i=1}^d$ on \mathbb{R}^1 such that*

$$\mu = \sigma_1 \otimes \dots \otimes \sigma_d, \nu = \tau_1 \otimes \dots \otimes \tau_d \tag{5.2}$$

and $\sigma_i * \tau_i = \mathcal{L}_{[0,1]}$ for all $i = 1, \dots, d$.

Note that the sufficiency part of the theorem follows by a direct computation. We only need to establish the necessity part of the theorem. Denote by P the orthogonal projection of the first coordinate on \mathbb{R}^d and Q the orthogonal projection of the corresponding orthogonal complement. If μ is a positive Borel measure on \mathbb{R}^d , we denote by μP^{-1} the positive Borel measure on \mathbb{R}^1 defined by $\mu P^{-1}(E) = \mu(P^{-1}(E))$ for any Borel set $E \subset \mathbb{R}$ and the measure μQ^{-1} is similarly defined. We will need the following lemmas.

Lemma 5.2 *Let μ and ν be two probability measures on \mathbb{R}^d . Then*

$$(\mu * \nu)P^{-1} = (\mu P^{-1}) * (\nu P^{-1}), \text{ and } (\mu * \nu)Q^{-1} = (\mu Q^{-1}) * (\nu Q^{-1}).$$

In particular, if μ and ν are two Borel probability measures satisfying (5.1), then we have

$$(\mu P^{-1}) * (\nu P^{-1}) = \mathcal{L}_{[0,1]} \text{ and } (\mu Q^{-1}) * (\nu Q^{-1}) = \mathcal{L}_{[0,1]^{d-1}}.$$

Proof The proof follows easily from the fact that

$$(\mu P^{-1})^\wedge(\xi) = \widehat{\mu}(\xi, 0, \dots, 0), \text{ and } (\mu Q^{-1})^\wedge(\xi_2, \dots, \xi_d) = \widehat{\mu}(0, \xi_2, \dots, \xi_d).$$

□

Lemma 5.3 *Let ν be a Borel probability measure on \mathbb{R}^d . Then, there is at most one probability measure μ on \mathbb{R}^d satisfying $\mu * \nu = \mathcal{L}_{[0,1]^d}$.*

Proof If μ is as above, we have

$$\widehat{\mu}(\xi)\widehat{\nu}(\xi) = (\mathcal{L}_{[0,1]^d})^\wedge(\xi), \quad \xi \in \mathbb{R}^d. \tag{5.3}$$

Therefore, $\widehat{\mu}(\xi)$ is thus determined on the set

$$F = \left\{ \xi \in \mathbb{R}^d : \xi_i \notin \mathbb{Z}^d, i = 1, \dots, d \right\},$$

Since $\overline{F} = \mathbb{R}^d$ and $\widehat{\mu}$ is continuous (as μ is compactly supported), $\widehat{\mu}$ and thus μ is completely determined by (5.3). □

The previous lemma is also valid if $[0, 1]^d$ is replaced by a d -dimensional rectangular box. Now, we proceed to the proof of Theorem 5.1.

Proof of Theorem 5.1 We prove the necessity part of the theorem by induction on the dimension. The statement is proved when $d = 1$ in Theorem 2.2. Assuming that the statement is true for $d - 1$, we now establish it on \mathbb{R}^d .

Let μ and ν be two Borel probability measures satisfying $\mu * \nu = \mathcal{L}_{[0,1]^d}$. By Lemma 5.2 and Theorem 2.2 [see also Eq. (2.11)], we can find an integer $N_1 \geq 2$ such that μP^{-1} and νP^{-1} can be decomposed (after possibly interchanging these two measures) as

$$\mu P^{-1} = \nu_1 * \alpha_1, \text{ and } \alpha_1 * (\nu P^{-1}) = \mathcal{L}_{[0,1/N_1]} \tag{5.4}$$

where $\nu_1 = 1/N_1 \sum_{j=0}^{N_1-1} \delta_{j/N_1}$ and $\alpha_1(E) = N_1(\mu P^{-1})(E \cap [0, 1/N_1])$ for any Borel set E . Let C_{N_1} be the d -dimensional rectangular box $\left[0, \frac{1}{N_1}\right) \times [0, 1]^{d-1}$. Then $[0, 1]^d \setminus C_{N_1} = \left[\frac{1}{N_1}, 1\right] \times [0, 1]^{d-1}$ and

$$\mu(C_{N_1}) = \mu P^{-1} \left(\left[0, \frac{1}{N_1}\right) \right) = \frac{1}{N_1}.$$

Hence, we can define two Borel probability measures on \mathbb{R}^d , ρ_1 and $\tilde{\rho}_1$, satisfying

$$\rho_1(E) = N_1 \mu(E \cap C_{N_1}), \quad \tilde{\rho}_1(E) = \frac{N_1}{N_1 - 1} \mu\left(E \cap ([0, 1]^d \setminus C_{N_1})\right)$$

for any Borel sets E . Then $\mu = \frac{1}{N_1} \rho_1 + (1 - \frac{1}{N_1}) \tilde{\rho}_1$. Since $\text{supp } \tilde{\rho} \subset [0, 1]^d \setminus C_{N_1}$ and $\text{supp } \nu \subset [0, 1]^d$, we have $\nu * \tilde{\rho} = 0$ on the rectangular box C_{N_1} . Hence,

$$\rho_1 * \nu = N_1(\mu * \nu) = \mathcal{L}_{C_{N_1}} \text{ on } C_{N_1}.$$

We can thus write $\rho_1 * \nu = \mathcal{L}_{C_{N_1}} + \eta$ where η is a positive measure. However, $\eta = 0$ as $\rho_1 * \nu$ and $\mathcal{L}_{C_{N_1}}$ are probability measures. Hence,

$$(\nu_1 \otimes \delta_{0_{d-1}}) * \rho_1 * \nu = \left(\frac{1}{N_1} \sum_{j=0}^{N_1-1} \delta_{(j/N_1, 0, \dots, 0)} \right) * \rho_1 * \nu = \mathcal{L}_{[0, 1]^d}$$

where $0_{d-1} = (0, \dots, 0) \in \mathbb{R}^{d-1}$. By Lemma 5.3, we have that

$$\mu = (\nu_1 \otimes \delta_{0_{d-1}}) * \rho_1, \text{ and } \rho_1 * \nu = \mathcal{L}_{[0, 1/N_1] \times [0, 1]^{d-1}}. \tag{5.5}$$

Furthermore, $\rho_1 P^{-1} = \alpha_1$ where α_1 is defined in (5.4).

We now consider two cases depending on whether μP^{-1} and νP^{-1} correspond to a type I or type II decomposition (as defined in Sect. 2).

Case 1 (Type I decomposition): Using the notations introduced in Sect. 2, we have then, without loss of generality, that

$$\mu P^{-1} = \nu_1 * \dots * \nu_{2k-1}, \quad \nu P^{-1} = \nu_2 * \dots * \nu_{2k} * \mathcal{L}_{\left[0, \frac{1}{N_1 \dots N_{2k}}\right]}.$$

By the previous steps, the identities in (5.5) hold. A similar argument, shows the existence of a probability measure ρ_2 such that

$$\nu = (\nu_2 \otimes \delta_{0_{d-1}}) * \rho_2 \text{ and } \rho_1 * \rho_2 = \mathcal{L}_{\left[0, \frac{1}{N_1 N_2}\right] \times [0, 1]^{d-1}}.$$

Continuing this procedure $2k$ -times, we deduce the existence of probability measures ρ_{2k-1} and ρ_{2k} such that

$$\mu = ((\nu_1 * \nu_3 * \dots * \nu_{2k-1}) \otimes \delta_{0_{d-1}}) * \rho_{2k-1}, \tag{5.6}$$

$$\nu = ((\nu_2 * \nu_4 * \dots * \nu_{2k}) \otimes \delta_{0_{d-1}}) * \rho_{2k}, \tag{5.7}$$

and

$$\rho_{2k-1} * \rho_{2k} = \mathcal{L}_{[0, 1/N_1 N_2 \dots N_{2k}] \times [0, 1]^{d-1}}. \tag{5.8}$$

By (5.6) and Lemma 5.2, $\mu P^{-1} = \nu_1 * \dots * \nu_{2k-1} * \rho_{2k-1} P^{-1}$, showing that $\rho_{2k-1} P^{-1} = \delta_0$. Hence, we can write $\rho_{2k-1} = \delta_0 \otimes \sigma$ for some positive measure σ on \mathbb{R}^{d-1} . Using (5.8) and Lemma 5.2 again, we obtain that $\sigma * (\rho_{2k} Q^{-1}) = \mathcal{L}_{[0,1]^{d-1}}$. Hence,

$$\begin{aligned} \rho_{2k-1} * \rho_{2k} &= \mathcal{L}_{[0,1/N_1 N_2 \dots N_{2k}]} \otimes \mathcal{L}_{[0,1]^{d-1}} \\ &= \mathcal{L}_{[0,1/N_1 N_2 \dots N_{2k}]} \otimes \left(\sigma * \left(\rho_{2k} Q^{-1} \right) \right) \\ &= (\delta_0 \otimes \sigma) * \left(\mathcal{L}_{[0,1/N_1 N_2 \dots N_{2k}]} \otimes \left(\rho_{2k} Q^{-1} \right) \right) \\ &= \rho_{2k-1} * \left(\mathcal{L}_{[0,1/N_1 N_2 \dots N_{2k}]} \otimes \left(\rho_{2k} Q^{-1} \right) \right). \end{aligned}$$

Lemma 5.3 shows that $\rho_{2k} = \mathcal{L}_{[0,1/N_1 N_2 \dots N_{2k}]} \otimes (\rho_{2k} Q^{-1})$ and (5.7) implies that $\nu = \nu P^{-1} \otimes \rho_{2k} Q^{-1}$. Finally, applying the induction hypothesis to the identity $\sigma * (\rho_{2k} Q^{-1}) = \mathcal{L}_{[0,1]^{d-1}}$, we can write $\sigma = \sigma_2 \otimes \dots \otimes \sigma_d$ and $\rho_{2k-1} Q^{-1} = \tau_2 \otimes \dots \otimes \tau_d$ with $\sigma_i * \tau_i = \mathcal{L}_{[0,1]}$ and Theorem 5.1 for dimension d follows.

Case 2 (Type II decomposition). In this case, we can without loss of generality assume that

$$\mu P^{-1} = \nu_1 * \nu_3 * \dots, \quad \nu P^{-1} = \nu_2 * \nu_4 * \dots$$

and we still have (5.6), (5.7) and (5.8) for all $k = 1, 2, \dots$ with $\rho_n P^{-1} \neq \delta_0$ for any integer n . As ρ_n are all probability measures, we can assume, by passing to subsequences if necessary, that the sequences $\{\rho_{2k-1}\}$ and $\{\rho_{2k}\}$ converge weakly to some probability measures that we denote by σ and τ , respectively. From (5.8), it is immediate to see that the supports of σ and τ are both contained in $\{0\} \times [0, 1]^{d-1}$. We can write $\sigma = \delta_0 \otimes \sigma'$ and $\tau = \delta_0 \otimes \tau'$. By passing to weak limit in (5.6) and (5.7), we have

$$\mu = \left(\mu P^{-1} \otimes \delta_{0_{d-1}} \right) * (\delta_0 \otimes \sigma'), \quad \nu = \left(\nu P^{-1} \otimes \delta_{0_{d-1}} \right) * (\delta_0 \otimes \tau'). \tag{5.9}$$

As $\mu * \nu = \mathcal{L}_{[0,1]^d}$ and $\left(\mu P_1^{-1} \otimes \delta_{0_{d-1}} \right) * \left(\nu P_1^{-1} \otimes \delta_{0_{d-1}} \right) = \mathcal{L}_{[0,1]} \otimes \delta_{0_{d-1}}$, we have

$$\sigma' * \tau' = \mathcal{L}_{[0,1]^{d-1}},$$

The conclusion follows immediately by (5.9) using the induction hypothesis. □

Proof of Theorem 1.1 on \mathbb{R}^d . The proof follows from the result on \mathbb{R}^1 . By Theorem 5.1, we can write $\mu = \sigma_1 \otimes \dots \otimes \sigma_d$ and $\nu = \tau_1 \otimes \dots \otimes \tau_d$ with $\sigma_i * \tau_i = \mathcal{L}_{[0,1]}$. Therefore, our conclusion on \mathbb{R}^1 implies that σ_i and τ_i are spectral measures on \mathbb{R}^1 with spectrum Λ_{σ_i} and Λ_{τ_i} respectively. Moreover, they satisfies $\Lambda_{\sigma_i} \oplus \Lambda_{\tau_i} = \mathbb{Z}$. Now we define

$$\Lambda_\mu = \bigotimes_{i=1}^d \Lambda_{\sigma_i}, \quad \Lambda_\nu = \bigotimes_{i=1}^d \Lambda_{\tau_i},$$

where $\bigotimes_{i=1}^d A_i := \{(a_1, \dots, a_d) : a_i \in A_i\}$ for sets $A_i \subset \mathbb{R}^1$. We claim that Λ_μ is a spectrum for μ (the proof that Λ_ν is a spectrum for ν is similar).

Note that $\widehat{\mu}(\xi) = \prod_{i=1}^d \widehat{\sigma}_i(\xi_i)$. From this, it follows easily that

$$\sum_{\lambda \in \Lambda_\mu} |\widehat{\mu}(\xi + \lambda)|^2 = \prod_{i=1}^d \left(\sum_{\lambda_i \in \Lambda_{\sigma_i}} |\widehat{\sigma}_i(\xi_i + \lambda_i)|^2 \right) = 1.$$

Hence, Λ_μ is a spectrum for μ . That the tiling property of the spectra (i.e. $\Lambda_\mu \oplus \Lambda_\nu = \mathbb{Z}^d$) follows immediately from the tiling property of Λ_{σ_i} and Λ_{τ_i} . \square

6 Remarks and Open Questions

As indicated in the introduction, the statement $\mathcal{F}(Q)$ is false in general. Nonetheless, this statement suggests many related questions that may help us understand the relationship among convolutions, translational tilings and spectral measures. Motivated by the **generalized Fuglede's Conjecture**, one of the main questions we would like to ask is:

(Q1): For which Q is the statement $\mathcal{F}(Q)$ true?

This question seems to be hard if we go beyond cubes as the methods of this paper would be difficult to extend. An easier, but still interesting question concerns the decomposition of the Lebesgue measure on sets as convolution product of singular measures:

(Q2): For what kind of measurable (resp. spectral) sets Q can \mathcal{L}_Q be decomposed into the convolution of two singularly continuous (resp. spectral) measure?

One natural type of such sets will be the self-affine tiles [14]. These tiles can be described as infinite convolution product of discrete measures and can therefore be decomposed into two singular measures using methods similar to those in Sect. 2.

Fourier frames and exponential Riesz bases are natural generalization of exponential orthonormal bases. It has been an interesting question to produce singular measures with Fourier frames but not exponential orthonormal bases. By now we only know we can produce such measures by considering measures which are absolutely continuous with respect to a spectral measure with density bounded above and away from 0 or convolving a spectral measure with some discrete measures [5, 9]. These methods are rather restrictive. As absolutely continuous (w.r.t. Lebesgue) measures with Fourier frames were completely classified in [19], we ask

(Q3): Can we produce new singular measures admitting Fourier frames by decomposing an absolutely continuous (w.r.t. Lebesgue) measures with Fourier frames? Conversely, is it true that all measures admitting Fourier frames are constructed in this way?

Given a spectral measure μ , another important issue is to classify its spectrum. This question has been studied for Lebesgue measures and some Cantor measures in [2, 3, 18]. However, there is no satisfactory answer when the measure is singular. The tiling statement of Theorem 1.1, suggests a possible answer.

(Q4): Let μ and ν be a natural complementary pair of $\mathcal{L}_{[0,1]}$. Let also Λ_μ be a spectrum for $L^2(\mu)$, does there exist a spectrum Λ_ν for $L^2(\nu)$ such that $\Lambda_\mu \oplus \Lambda_\nu = \mathbb{Z}$?

It is not difficult to prove that **(Q4)** actually holds for type I decompositions. The remaining challenge is to answer the question for type II decompositions.

Acknowledgments The authors would like to thank the anonymous referee for his/her valuable comments and suggestions.

References

1. An, L.-X., He, X.-G.: A class of spectral Moran measures. *J. Funct. Anal.* **266**, 343–354 (2014)
2. Dai, X.-R., He, X.-G., Lai, C.-K.: Spectral property of Cantor measures with consecutive digits. *Adv. Math.* **242**, 187–208 (2013)
3. Dutkay, D., Han, D., Sun, Q.: On spectra of a Cantor measure. *Adv. Math.* **221**, 251–276 (2009)
4. Dutkay, D., Jorgensen, P.: Fourier frequencies in affine iterated function systems. *J. Funct. Anal.* **247**, 110–137 (2007)
5. Dutkay, D., Lai, C.-K.: Uniformity of measures with Fourier frames. *Adv. Math.* **252**, 684–707 (2014)
6. Dutkay, D., Lai, C.-K.: Some reductions of the spectral set conjecture to integers. *Math. Proc. Cambridge Phil. Soc.* **156**, 123–135 (2014)
7. Fuglede, B.: Commuting self-adjoint partial differential operators and a group theoretic problem. *J. Funct. Anal.* **16**, 101–121 (1974)
8. Jorgensen, P., Pedersen, S.: Dense analytic subspaces in fractal L^2 spaces. *J. Anal. Math.* **75**, 185–228 (1998)
9. He, X.-G., Lai, C.-K., Lau, K.-S.: Exponential spectra in $L^2(\mu)$. *Appl. Comput. Harmon. Anal.* **34**, 327–338 (2013)
10. Iosevich, A., Kolountzakis, M.: Periodicity of the spectrum in dimension one. *Anal. PDE* **6**, 819–827 (2013)
11. Kolountzakis, M., Matolcsi, M.: Tiles with no spectra. *Forum Math.* **18**, 519–528 (2006)
12. Kolountzakis, M., Matolcsi, M.: Complex Hadamard matrices and the spectral set conjecture. [arXiv: math/0411512](https://arxiv.org/abs/math/0411512) [math.CA]
13. Lagarias, J., Wang, Y.: Tiling the line by translates of one tile. *Invent. Math.* **124**, 341–365 (1996)
14. Lagarias, J.: Self-Affine tiles in \mathbb{R}^n . *Adv. Math.* **121**, 21–49 (1996)
15. Łaba, I., Wang, Y.: On spectral Cantor measures. *J. Funct. Anal.* **193**, 409–420 (2002)
16. Lewis, T.: The factorization of the rectangular distribution. *J. Appl. Probab.* **4**, 529–542 (1967)
17. Long, C.: Addition theorems for sets of integers. *Pacific J. Math.* **23**, 107–112 (1967)
18. Lagarias, J., Reeds, J., Wang, Y.: Orthonormal bases of exponentials for the n -cubes. *Duke Math. J.* **103**, 25–37 (2000)
19. Lai, C.-K.: On Fourier frame of absolutely continuous measures. *J. Funct. Anal.* **261**, 2877–2889 (2011)
20. Pedersen, S., Wang, Y.: Universal spectra, universal tiling sets and the spectral set conjecture. *Math Scand.* **88**, 246–256 (2001)
21. Strichartz, R.: Mock Fourier series and transforms associated with certain Cantor measures. *J. Anal. Math.* **81**, 209–238 (2000)
22. Tao, T.: Fuglede’s conjecture is false in 5 or higher dimensions. *Math. Res. Lett.* **11**, 251–258 (2004)