



Extrapolation of Compactness on Banach Function Spaces

Emiel Lorist¹ · Zoe Nieraeth²

Received: 8 November 2023 / Revised: 5 April 2024 / Accepted: 6 April 2024 /
Published online: 2 May 2024
© The Author(s) 2024

Abstract

We prove an extrapolation of compactness theorem for operators on Banach function spaces satisfying certain convexity and concavity conditions. In particular, we show that the boundedness of an operator T in the weighted Lebesgue scale and the compactness of T in the unweighted Lebesgue scale yields compactness of T on a very general class of Banach function spaces. As our main new tool, we prove various characterizations of the boundedness of the Hardy-Littlewood maximal operator on such spaces and their associate spaces, using a novel sparse self-improvement technique. We apply our main results to prove compactness of the commutators of singular integral operators and pointwise multiplication by functions of vanishing mean oscillation on, for example, weighted variable Lebesgue spaces.

Keywords Banach function space · Compact operator · Extrapolation · Muckenhoupt weight

Mathematics Subject Classification Primary 46E30 · Secondary 46B50 · 42B25

1 Introduction

The classical Rubio de Francia extrapolation theorem [23, 44] is one of the most powerful tools in the theory of weighted norm inequalities. In its simplest form, it states that if an operator T is bounded on the weighted Lebesgue space $L_w^p(\mathbf{R}^d)$ some

Communicated by Oscar Blasco.

✉ Emiel Lorist
e.lorist@tudelft.nl

Zoe Nieraeth
zoe.nieraeth@gmail.com

¹ Delft Institute of Applied Mathematics, Delft University of Technology, P.O. Box 5031, 2600 GA Delft, The Netherlands

² BCAM-Basque Center for Applied Mathematics, Bilbao, Spain

$p \in (1, \infty)$ and all weights $w \in A_p$, then T is automatically bounded on $L_w^p(\mathbf{R}^d)$ for all $p \in (1, \infty)$ and $w \in A_p$. Here we call a positive function w a weight, we let $L_w^p(\mathbf{R}^d)$ denote the space of all functions f such that $fw \in L^p(\mathbf{R}^d)$, and write $w \in A_p$ if

$$[w]_p := \sup_Q \left(\frac{1}{|Q|} \int_Q w^p \right)^{\frac{1}{p}} \left(\frac{1}{|Q|} \int_Q w^{-p'} \right)^{\frac{1}{p'}} < \infty,$$

where the supremum is taken over all cubes $Q \subseteq \mathbf{R}^d$.

In recent years, Rubio de Francia’s extrapolation theorem has been extended to compact operators. Again in its simplest form, Hytönen and Lappas [29] showed that if

- T is bounded on $L_w^p(\mathbf{R}^d)$ for some $p \in (1, \infty)$ and all weights $w \in A_p$;
- T is compact on $L_w^p(\mathbf{R}^d)$ for some $p \in (1, \infty)$ and some weight $w \in A_p$;

then T is compact on $L_w^p(\mathbf{R}^d)$ for all $p \in (1, \infty)$ and $w \in A_p$. Note that in typical applications, one checks the compactness assumption for $p = 2$ and $w = 1$, i.e. on the Hilbert space $L^2(\mathbf{R}^d)$.

In order to fix ideas, let us briefly sketch the proof of Hytönen and Lappas [29] in the simplest case stated above. In essence, the argument has three main ingredients:

- (1) The Rubio de Francia extrapolation theorem;
- (2) Interpolation of compactness: For Banach function spaces X_0 and X_1 and an operator T which is compact on X_0 and bounded on X_1 , T is compact on the product space $X_0^{1-\theta} \cdot X_1^\theta$ for $\theta \in (0, 1)$;
- (3) The self-improvement property of Muckenhoupt classes: For $w \in A_p$ there is an $1 < r < p$ such that $w^r \in A_{p/r}$.

Using these ingredients, one starts by observing that T is bounded on $L_w^p(\mathbf{R}^d)$ for all $p \in (1, \infty)$ and $w \in A_p$ by (1) and hence, by (2), the compactness of the operator T on $L_w^p(\mathbf{R}^d)$ can be used to deduce the compactness of T on

$$L^2(\mathbf{R}^d) = L_w^p(\mathbf{R}^d)^{\frac{1}{2}} \cdot L_{w^{-1}}^{p'}(\mathbf{R}^d)^{\frac{1}{2}}.$$

Next, fix $p \in (1, \infty)$ and $w \in A_p$. Using (3) on both $w \in A_p$ and $w^{-1} \in A_{p'}$, one finds $1 < r < p < s < \infty$ such that $w^r \in A_{\frac{p}{r}}$ and $w^{-s'} \in A_{\frac{p'}{s'}}$ and $1 - \frac{1}{r} = \frac{1}{s}$. One readily checks that this is equivalent to the condition $w_{r,s} \in A_{p_{r,s}}$, where

$$\frac{1}{p_{r,s}} := \frac{\frac{1}{p} - \frac{1}{s}}{\frac{1}{r} - \frac{1}{s}}, \quad \text{and} \quad w_{r,s} := w^{\frac{1}{\frac{1}{r} - \frac{1}{s}}}.$$

Note that the affine transformation that maps $\frac{1}{p} \rightarrow \frac{1}{p_{r,s}}$ is the one that maps the interval $(\frac{1}{s}, \frac{1}{r})$ to the interval $(0, 1)$ through a translation by $-\frac{1}{s}$ and a scaling by a factor of

$\frac{1}{r} - \frac{1}{s}$. Similarly, the space $X_{r,s} := L_{w_{r,s}}^{p_{r,s}}(\mathbf{R}^d)$ can be scaled and translated back to the space $X := L_w^p(\mathbf{R}^d)$ through the factorization

$$X = (X_{r,s})^{\frac{1}{r} - \frac{1}{s}} \cdot L^s(\mathbf{R}^d) = (X_{r,s})^{1-\theta} \cdot L^2(\mathbf{R}^d)^\theta, \quad \theta = \frac{2}{s}. \tag{1.1}$$

Therefore, since T is bounded on $X_{r,s}$ and compact on $L^2(\mathbf{R}^d)$, this means it is also compact on $L_w^p(\mathbf{R}^d)$ by (2), proving the result.

The extrapolation theorem of Rubio de Francia has been generalized to a general class of Banach function spaces. This was first done for rearrangement invariant spaces by Curbera, García-Cuerva, Martell, and Pérez in [18]. In the recent work of Cao, Márin and Martell [9] weighted Banach function spaces under weighted analogues of the conditions of [4] were considered. In [43] by the second author this was generalized to the class of saturated spaces, which is also the class of Banach function spaces we consider in this work, see Sect. 2. We refer the reader to [43, Sect. 4.7] for a direct comparison of the assumptions with those of [9]. In particular, it was shown in [43] that if T is bounded on $L_w^p(\mathbf{R}^d)$ for some $p \in (1, \infty)$ and all weights $w \in A_p$, then T is bounded on any Banach function space X for which

$$M : X \rightarrow X, \quad M : X' \rightarrow X',$$

where M denotes the Hardy–Littlewood maximal operator and X' the associate space of X . This, of course, includes the weighted Lebesgue spaces $X = L_w^p(\mathbf{R}^d)$ for $w \in A_p$, as it is well-known that M is bounded on X and $X' = L_{w^{-1}}^{p'}(\mathbf{R}^d)$.

Thus, we observe that (1) is available in the setting of Banach function spaces. Moreover, (2) is already phrased in this setting. Therefore to extend the extrapolation of compactness theorem of Hytönen and Lappas [29] to the setting of Banach function spaces, one only needs to find a suitable replacement for (3), which will be the main contribution of this paper.

In the above proof sketch, (3) was used to deduce the factorization in Eq. (1.1). Thus, we are looking for a self-improvement property of the form: If X is a Banach function space such that

$$M : X \rightarrow X, \quad M : X' \rightarrow X',$$

then there are $1 < r < s < \infty$ with $1 - \frac{1}{r} = \frac{1}{s}$ such that

$$M : X_{r,s} \rightarrow X_{r,s}, \quad M : (X_{r,s})' \rightarrow (X_{r,s})' \tag{1.2}$$

for some suitable space $X_{r,s}$ satisfying Eq. (1.1). In [43] it was shown that a space $X_{r,s}$ such that Eq. (1.1) holds exists if and only if X is r -convex and s -concave, i.e.,

$$\begin{aligned} \|(|f|^r + |g|^r)^{\frac{1}{r}} \|_X &\leq (\|f\|_X^r + \|g\|_X^r)^{\frac{1}{r}}, & f, g \in X \\ (\|f\|_X^s + \|g\|_X^s)^{\frac{1}{s}} &\leq \| (|f|^s + |g|^s)^{\frac{1}{s}} \|_X, & f, g \in X. \end{aligned}$$

We note that in this case $X_{r,s}$ is given by the formula

$$X_{r,s} := \left[[(X^r)']^{(\frac{s}{r})'} \right]'$$

Combined with the boundedness of M on $X_{r,s}$ for r small enough and s large enough, which we will discuss below, we have sketched the proof of our first main result.

Theorem 1.1 *Let*

$$T : \bigcup_{\substack{p \in (1, \infty) \\ w \in A_p}} L_w^p(\mathbf{R}^d) \rightarrow L^0(\mathbf{R}^d)$$

be a linear operator such that

- *T is bounded on $L_w^p(\mathbf{R}^d)$ for some $p \in (1, \infty)$ and all weights $w \in A_p$;*
- *T is compact on $L_w^p(\mathbf{R}^d)$ for some $p \in (1, \infty)$ and some weight $w \in A_p$.*

Let X be a Banach function space over \mathbf{R}^d such that

$$M : X \rightarrow X, \quad M : X' \rightarrow X',$$

and assume X is r -convex and s -concave for some $1 < r < s < \infty$. Then $T : X \rightarrow X$ is compact.

We prove Theorem 1.1 as Theorem 4.1 below. We note that the r -convexity and s -concavity conditions in the case of the Lebesgue space $X = L_w^p(\mathbf{R}^d)$ are satisfied with $r = s = p$. Theorem 1.1 is also applicable to, for example, weighted variable Lebesgue spaces $X = L_w^{p(\cdot)}(\mathbf{R}^d)$. Here the function $p : \mathbf{R}^d \rightarrow (0, \infty)$ has to satisfy

$$1 < \text{ess inf } p \leq \text{ess sup } p < \infty$$

so that X is r -convex and s -concave with $r = \text{ess inf } p, s = \text{ess sup } p$, and the weight and the exponent have to satisfy some additional condition ensuring the boundedness of M on X (see, e.g., [21, 34] for the unweighted setting and [17] for the weighted setting).

As we shall see in Sect. 6, Theorem 1.1 can be used to deduce the compactness of commutators of singular integral operators and multiplication by functions with vanishing mean oscillation. We refer the reader to [29, Sect. 5–8, p. 10] for further examples of operators to which Theorem 1.1 is applicable.

Remark 1.2 It is clear that the proof strategy of Theorem 1.1 cannot work without the convexity and concavity assumptions, since, as mentioned before, the existence of the factorization Eq. (1.1) implies that X is r -concave and s -concave. This means that Theorem 1.1 is, in particular, not applicable to Morrey spaces, as Morrey spaces are not s -concave for any $s < \infty$. In [33], Lappas proved that extrapolation of compactness in the Morrey scale is possible if one replaces the compactness assumption on $L_w^p(\mathbf{R}^n)$

by a compactness assumption on a Morrey space. Since there are factorization results in the spirit of Eq. (1.1), but with $L^s(\mathbf{R}^d)$ replaced by this Morrey space, this allows one to follow the same lines of reasoning as before. For a general, non-convex or non-concave Banach function space X , it is not clear what a suitable replacement for $L^s(\mathbf{R}^d)$ would be.

Let us return to the analogue of the self-improvement property of the Muckenhoupt classes needed to prove Theorem 1.1, which was stated in Eq. (1.2). For a Banach function space X , it was shown by Lerner and Ombrosi [36] that if $M : X \rightarrow X$, then there is an $r > 1$ such that also $M : X^r \rightarrow X^r$ and hence, if

$$M : X \rightarrow X, \quad M : X' \rightarrow X'.$$

as in Theorem 1.1, we can find $1 < r < s < \infty$ so that

$$M : X^r \rightarrow X^r, \quad M : (X')^{s'} \rightarrow (X')^{s'}. \tag{1.3}$$

Unfortunately, it is not clear if this implies the bounds

$$M : X_{r,s} \rightarrow X_{r,s}, \quad M : (X_{r,s})' \rightarrow (X_{r,s})'. \tag{1.4}$$

In fact, it was shown in [43, Theorem 2.36] that Eq. (1.4) implies Eq. (1.3) and the converse is an open problem, see [43, Conjecture 2.39].

Instead of using the self-improvement result of [36] to the spaces X and X' separately, we will prove a simultaneous self-improvement result to show that if M is bounded on X and X' , and the space X is r^* -concave and s^* -concave for some $1 < r^* \leq s^* < \infty$, then Eq. (1.4) holds for all $1 < r \leq r^*$ small enough and $s^* \leq s < \infty$ large enough. This is a direct consequence of our second result.

Proposition 1.3 *Let $r^* \in (1, \infty)$ and let X be an r^* -convex Banach function space over \mathbf{R}^d . Then the following are equivalent:*

- (i) *We have $M : X \rightarrow X$ and $M : X' \rightarrow X'$;*
- (ii) *There is an $r_0 \in (1, r^*]$ so that for all $r \in (1, r_0]$ we have*

$$M : X^r \rightarrow X^r, \quad M : (X^r)' \rightarrow (X^r)';$$

- (iii) *There is an $r \in (1, r^*]$ so that $M : (X^r)' \rightarrow (X^r)'$.*

Proposition 1.3 is proved as Theorem 3.1 below. It relies on a sparse characterization of the boundedness of M on X and X' , followed by a sparse self-improvement result based on a novel use of the classical reverse Hölder inequality of Muckenhoupt weights. Applying Theorem 1.3 first to X and then to the resulting space $(X^r)'$ yields Eq. (1.4) for some $1 < r < s < \infty$, see Corollary 3.3. We note that Proposition 1.3 is also of independent interest, as various works use (3) as an assumption, often not realizing that it is equivalent to (1.3).

Rubio de Francia’s extrapolation theorem for $L_w^p(\mathbf{R}^d)$ has been generalized to the off-diagonal setting by Harboure, Macías and Segovia [26] and to the limited range

setting by Auscher and Martell [3]. Both settings were extended to general (quasi)-Banach function spaces X in [43]. Using [43, Theorem A], we also obtain a limited range, off-diagonal extrapolation of compactness theorem for Banach function spaces, which is our final main result. We refer the reader to Sect. 5 for the definition of $A_{\vec{p},(\vec{r},\vec{s})}$.

Theorem 1.4 *Let $\alpha \in \mathbf{R}$ and let $r_1, r_2 \in [1, \infty)$, $s_1, s_2 \in (1, \infty]$ satisfy $r_j < s_j$ for $j \in \{1, 2\}$ and*

$$\frac{1}{r_1} - \frac{1}{r_2} = \frac{1}{s_1} - \frac{1}{s_2} = \alpha.$$

Define

$$\mathcal{P} := \{(p_1, p_2) \in (0, \infty]^2 : \frac{1}{p_j} \in [\frac{1}{s_j}, \frac{1}{r_j}], j \in \{1, 2\}, \frac{1}{p_1} - \frac{1}{p_2} = \alpha\}$$

and let

$$T : \bigcup_{\substack{(p_1, p_2) \in \mathcal{P} \\ w \in A_{\vec{p},(\vec{r},\vec{s})}}} L_w^{p_1}(\mathbf{R}^d) \rightarrow L^0(\mathbf{R}^d)$$

be a linear operator such that

- *T is bounded from $L_w^{p_1}(\mathbf{R}^d)$ to $L_w^{p_2}(\mathbf{R}^d)$ for some $(p_1, p_2) \in \mathcal{P}$ and all $w \in A_{\vec{p},(\vec{r},\vec{s})}$;*
- *T is compact from $L_w^{p_1}(\mathbf{R}^d)$ to $L_w^{p_2}(\mathbf{R}^d)$ for some $(p_1, p_2) \in \mathcal{P}$ and some $w \in A_{\vec{p},(\vec{r},\vec{s})}$.*

Let $r_j < r_j^ < s_j^* < s_j$ and let X_j be an r_j^* -convex and s_j^* -concave Banach function space for $j \in \{1, 2\}$ satisfying*

$$(X_1)_{r_1, s_1} = (X_2)_{r_2, s_2}$$

and

$$M : (X_1)_{r_1, s_1} \rightarrow (X_1)_{r_1, s_1}, \quad M : ((X_1)_{r_1, s_1})' \rightarrow ((X_1)_{r_1, s_1})'$$

Then $T : X_1 \rightarrow X_2$ is compact.

We prove Theorem 1.4 as Theorem 5.1 below. Note that Theorem 1.4 recovers [29, Theorems 2.3 and 2.4] for X_1 and X_2 weighted Lebesgue spaces in a unified result.

Remark 1.5 For weighted Lebesgue spaces, the result in [29] was further generalized to multilinear operators by Cao, Olivo and Yabuta [8] (see also [28] by Hytönen and Lappas). Currently, multilinear extrapolation in quasi-Banach function spaces has not yet been proven in its full expected generality (see [43, Sect. 6]). Moreover, the compactness of multilinear operators on product spaces seems to be only available for bilinear operators (see [14, Sect. 3]) and, although appearing naturally in the multilinear setting, quasi-Banach function spaces are not allowed in this result. Because of

these limitations, we will not develop multilinear versions of our results in the current paper. We do point out that a bilinear compact extrapolation for products of weights classes extending [8, Theorem 1.2] can easily be obtained using [43, Theorem B], but since the proof is just an application of the linear case presented in this paper on each component, we do not consider this an important contribution to the literature, and therefore omit it.

Finally, we would like to note that for weighted Lebesgue spaces, an extrapolation of compactness result has also been obtained in the two-weight setting by Liu, Wu and Yang [39].

The plan for this paper is as follows. We start in Sect. 2 by defining Banach function spaces and all their relevant properties. Afterwards, in Sect. 3, we prove the self-improvement property of the maximal operator on Banach function spaces stated in Theorem 1.3. Sections 4 and 5 are devoted to proving the extrapolation of compactness results in the full range case (Theorem 1.1) and limited range, off-diagonal case (Theorem 1.4) respectively. Finally, in Sect. 6 we apply Theorem 1.1 to deduce the compactness of commutators of both Calderón–Zygmund and rough homogeneous singular integral operators with pointwise multiplication by a function with vanishing mean oscillation.

2 Banach Function Spaces

Let (Ω, μ) be a σ -finite measure space. Let $L^0(\Omega)$ denote the space of measurable functions on (Ω, μ) . A vector space $X \subseteq L^0(\Omega)$ equipped with a norm $\|\cdot\|_X$ is called a *Banach function space over Ω* if it satisfies the following properties:

- *Ideal property:* If $f \in X$ and $g \in L^0(\Omega)$ with $|g| \leq |f|$, then $g \in X$ with $\|g\|_X \leq \|f\|_X$.
- *Fatou property:* If $0 \leq f_n \uparrow f$ for $(f_n)_{n \geq 1}$ in X and $\sup_{n \geq 1} \|f_n\|_X < \infty$, then $f \in X$ and $\|f\|_X = \sup_{n \geq 1} \|f_n\|_X$.
- *Saturation property:* For every measurable $E \subseteq \Omega$ of positive measure, there exists a measurable $F \subseteq E$ of positive measure with $\mathbf{1}_F \in X$.

We note that the saturation property is equivalent to the assumption there is an $f \in X$ such that $f > 0$ a.e. Moreover, the Fatou property ensures that X is complete.

We define the associate space X' as the space of all $g \in L^0(\mathbf{R}^n)$ such that

$$\|g\|_{X'} := \sup_{\|f\|_X \leq 1} \int_{\mathbf{R}^n} |fg| < \infty,$$

which is again a Banach function space. By the Lorentz–Luxembourg theorem we have $X'' = X$ with equal norms.

For proofs and a further elaboration on these properties we refer the reader to the book of Zaanan [49] and the recent survey [40].

Remark 2.1 Throughout the literature, following the book by Bennet and Sharpley [4], in the definition of a Banach function space X it is often in addition assumed that for

all measurable $E \subseteq \Omega$ with $\mu(E) < \infty$ one has

$$\mathbf{1}_E \in X \quad \text{and} \quad \mathbf{1}_E \in X'. \tag{2.1}$$

Note that this implies the saturation property. However, Eq. (2.1) is too restrictive to study weighted norm inequalities in harmonic analysis. Indeed, there are examples of weighted Lebesgue spaces $L_w^p(\mathbf{R}^d)$ for $p \in (1, \infty)$ and $w \in A_p$ that do not satisfy Eq. (2.1), see [45, Sect. 7.1].

2.1 Convexity Properties

Let X be a Banach function space over Ω and $1 \leq p \leq q \leq \infty$. We call X p -convex if

$$\|(|f|^p + |g|^p)^{\frac{1}{p}}\|_X \leq (\|f\|_X^p + \|g\|_X^p)^{\frac{1}{p}}, \quad f, g \in X,$$

and we call X q -concave if

$$(\|f\|_X^q + \|g\|_X^q)^{\frac{1}{q}} \leq \|(|f|^q + |g|^q)^{\frac{1}{q}}\|_X, \quad f, g \in X.$$

Note that any Banach function space is 1-convex by the triangle inequality and ∞ -concave by the ideal property. One often defines p -convexity and q -concavity using finite sums of elements from X and a constant in the defining inequalities, but, by [38, Theorem 1.d.8], one can always renorm X such that these constants are equal to one, yielding our definition.

We note that if X is p -convex and q -concave, then X is also p_0 convex and q_0 -concave for all $p_0 \in [1, p]$ and $q_0 \in [q, \infty]$ and X' is q' -convex and p' -concave (see, e.g., [38, Sect. 1.d]).

For $p \in (0, \infty)$ we define the p -concavification of X by

$$X^p := \{f \in L^0(\Omega) : |f|^{\frac{1}{p}} \in X\},$$

i.e. for a positive $f \in L^0(\Omega)$ we have $f \in X$ if and only if $f^p \in X^p$. We equip X^p with the quasi-norm

$$\|f\|_{X^p} := \| |f|^{\frac{1}{p}} \|_X^p, \quad f \in X^p.$$

Note that X is a Banach function space if and only if X is $p \vee 1$ -convex.

Definition 2.2 Let $1 \leq r < s \leq \infty$ and let X be an r -convex and s -concave Banach function space. We define the (r, s) -rescaled Banach function space of X by

$$X_{r,s} := \left[[(X^r)']^{\left(\frac{s}{r}\right)'} \right]',$$

which is again a Banach function space.

2.2 Calderón–Lozanovskii Products

Let X_0 and X_1 be Banach function spaces over Ω and $\theta \in (0, 1)$. We define the *Calderón–Lozanovskii product* (see [7, 41]) $X_\theta := X_0^{1-\theta} \cdot X_1^\theta$ as the space of those $h \in L^0(\Omega)$ for which there exist $0 \leq f \in X_0$, $0 \leq g \in X_1$ such that $|h| \leq f^{1-\theta} g^\theta$. We equip this space with the norm

$$\|h\|_{X_\theta} := \inf \|f\|_{X_0}^{1-\theta} \|g\|_{X_1}^\theta,$$

where the infimum is taken over all $0 \leq f \in X_0$, $0 \leq g \in X_1$ for which $|h| \leq f^{1-\theta} g^\theta$. We note that X_θ is a Banach function space, see e.g. [19, Appendix 7].

The following proposition will play a key role in the proof of Theorem 1.1 and Theorem 1.4. Note that one can interpret the appearing products as Calderón–Lozanovskii products since

$$L^s(\Omega) = L^{1+\frac{s}{r}}(\Omega)^{1-(\frac{1}{r}-\frac{1}{s})}.$$

Proposition 2.3 *Let $1 \leq r < s \leq \infty$ and let X Banach function space over Ω .*

(i) *If X is r -convex and s -concave, then*

$$X = (X_{r,s})^{\frac{1}{r}-\frac{1}{s}} \cdot L^s(\Omega).$$

(ii) *If there is a Banach function space Y such that*

$$X = Y^{\frac{1}{r}-\frac{1}{s}} \cdot L^s(\Omega),$$

then X is r -convex and s -concave and $Y = X_{r,s}$.

Proof For 2.3 we refer the reader to [43, Corollary 2.12], and 2.3 is proven analogously to [43, Theorem 2.13], substituting $X_{r,s}$ by Y . \square

3 Sparse Self-Improvement for the Maximal Operator

As we will need Proposition 1.3 in the proof of Theorem 1.1, we start in this section by proving Proposition 1.3 and its consequences. First we introduce some notation. For $r \in (0, \infty)$, a cube $Q \subseteq \mathbf{R}^d$ and function $f \in L^0(\mathbf{R}^d)$ we define the r -average of f by $\langle f \rangle_{r,Q} := (\frac{1}{|Q|} \int_Q |f|^r)^{1/r}$, and we define the Hardy–Littlewood maximal operator by

$$Mf := \sup_Q \langle f \rangle_{1,Q} \mathbf{1}_Q,$$

where the supremum is taken over all cubes Q in \mathbf{R}^d .

Theorem 3.1 *Let $r^* \in (1, \infty)$ and let X be an r^* -convex Banach function space over \mathbb{R}^d . Then the following are equivalent:*

- (i) *We have $M : X \rightarrow X$ and $M : X' \rightarrow X'$;*
- (ii) *There is an $r_0 \in (1, r^*]$ so that for all $r \in (1, r_0]$ we have*

$$M : X^r \rightarrow X^r, \quad M : (X^r)' \rightarrow (X^r)';$$

- (iii) *There is an $r \in (1, r^*]$ so that $M : (X^r)' \rightarrow (X^r)'$.*

Before we turn to the proof, we provide two useful corollaries. First of all, we obtain the following equivalent formulations of the bounds

$$M : X \rightarrow X, \quad M : X' \rightarrow X',$$

assumed in Theorem 1.1.

Corollary 3.2 *Let X be a Banach function space for which there exist $1 < r^* < s^* < \infty$ such that X is r^* -convex and s^* -concave. Then the following are equivalent:*

- (i) *We have $M : X \rightarrow X$ and $M : X' \rightarrow X'$;*
- (ii) *There is an $r \in (1, r^*]$ so that $M : (X^r)' \rightarrow (X^r)'$;*
- (iii) *There is an $s \in [s^*, \infty)$ so that $M : X_{1,s} \rightarrow X_{1,s}$.*

Proof The equivalence of 3.2 and 3.2 is contained in Theorem 3.1 whereas the equivalence of 3.2 and 3.2 follows from applying the first equivalence with X replaced by X' , which is $(s^*)'$ -convex, to find an $s' \in (1, (s^*)']$ for which M is bounded on $[(X')^{s'}]' = X_{1,s}$. □

Our second corollary is the self-improvement property in Banach function spaces discussed in the introduction, which will replace the self-improvement property of the Muckenhoupt classes in the proof of Theorem 1.1 and Theorem 1.4:

Corollary 3.3 *Let X be a Banach function space for which there exist $1 < r^* < s^* < \infty$ such that X is r^* -convex, s^* -concave, and*

$$M : X \rightarrow X, \quad M : X' \rightarrow X'.$$

Then there exist $r_0 \in (1, r^]$, $s_0 \in [s^*, \infty)$ such that for all $r \in (1, r_0]$ and $s \in [s_0, \infty)$ we have*

$$M : X_{r,s} \rightarrow X_{r,s}, \quad M : (X_{r,s})' \rightarrow (X_{r,s})'.$$

Proof By Theorem 3.1 there is a $r_0 \in (1, r^*]$ for which

$$M : X^{r_0} \rightarrow X^{r_0}, \quad M : (X^{r_0})' \rightarrow (X^{r_0})'.$$

Note that $(X^{r_0})'$ is $(\frac{s^*}{r_0})'$ -convex. By applying Theorem 3.1 to $(X^{r_0})'$, we find a $t_0 \in (1, (\frac{s^*}{r_0})']$ such that M is bounded on $[(X^{r_0})']^{t_0}$ and $[(X^{r_0})']^{t_0}'$.

Now define $s_0 := r_0 t'_0 > s^*$. Then we have $t_0 = (\frac{s_0}{r_0})'$, so that

$$M : X_{r_0, s_0} \rightarrow X_{r_0, s_0}, \quad M : (X_{r_0, s_0})' \rightarrow (X_{r_0, s_0})'.$$

Letting $r \in (1, r_0)$, $s \in (s_0, \infty)$ and noting that $(X_{r_0, s_0})' = (X')_{s'_0, r'_0}$ by [43, Proposition 2.14], it follows from [43, Proposition 2.30] that also

$$M : X_{r, s} \rightarrow X_{r, s}, \quad M : (X_{r, s})' \rightarrow (X_{r, s})'.$$

The assertion follows. □

We now turn to the proof of Theorem 3.1. As a final preparation, we require a lemma on sparse operators. A collection of cubes \mathcal{S} in \mathbf{R}^d is called *sparse* if for each $Q \in \mathcal{S}$ there is a measurable set $E_Q \subseteq Q$ for which $|E_Q| \geq \frac{1}{2}|Q|$ and, furthermore, the collection $(E_Q)_{Q \in \mathcal{S}}$ is pairwise disjoint. For a sparse collection of cubes \mathcal{S} and $f \in L^0(\mathbf{R}^d)$ we define

$$T_{\mathcal{S}} f := \sum_{Q \in \mathcal{S}} \langle f \rangle_{1, Q} \mathbf{1}_Q.$$

We have the following characterization of when M is bounded on X and X' in terms of $T_{\mathcal{S}}$.

Lemma 3.4 *Let X be a Banach function space over \mathbf{R}^d . Then*

$$M : X \rightarrow X, \quad M : X' \rightarrow X'$$

if and only if we have $T_{\mathcal{S}} : X \rightarrow X$ for all sparse collections of cubes \mathcal{S} with

$$\sup_{\mathcal{S} \text{ is sparse}} \|T_{\mathcal{S}}\|_{X \rightarrow X} < \infty.$$

Proof For $f, g \in L^0(\mathbf{R}^d)$ we define

$$M_{1,1}(f, g) := \sup_Q \langle f \rangle_{1, Q} \langle g \rangle_{1, Q} \mathbf{1}_Q.$$

By [43, Proposition 6.1] it suffices to show that $M_{1,1} : X \times X' \rightarrow L^1(\mathbf{R}^d)$ if and only if for all sparse collections \mathcal{S} we have $T_{\mathcal{S}} : X \rightarrow X$ uniformly in \mathcal{S} with

$$\sup_{\mathcal{S} \text{ is sparse}} \|T_{\mathcal{S}}\|_{X \rightarrow X} \approx_d \|M_{1,1}\|_{X \times X' \rightarrow L^1(\mathbf{R}^d)}.$$

Assume first that $M_{1,1} : X \times X' \rightarrow L^1(\mathbf{R}^d)$. If \mathcal{S} is sparse, then for $f \in X$ and $g \in X'$

$$\begin{aligned} \|(T_S f)g\|_{L^1(\mathbf{R}^d)} &= \sum_{Q \in \mathcal{S}} \langle f \rangle_{1,Q} \langle g \rangle_{1,Q} |Q| \\ &\leq 2 \sum_{Q \in \mathcal{S}} \int_{E_Q} M_{1,1}(f, g) \, dx \\ &\leq 2 \|M_{1,1}(f, g)\|_{L^1(\mathbf{R}^d)} \end{aligned}$$

Hence,

$$\sup_{\mathcal{S} \text{ is sparse}} \|T_S\|_{X \rightarrow X} = \sup_{\mathcal{S} \text{ is sparse}} \sup_{\substack{\|f\|_X=1 \\ \|g\|_{X'}=1}} \|(T_S f)g\|_{L^1(\mathbf{R}^d)} \leq 2 \|M_{1,1}\|_{X \times X' \rightarrow L^1(\mathbf{R}^d)}.$$

Conversely, using [42, Proposition 3.2.10], for each $f, g \in L^0(\mathbf{R}^d)$, each dyadic lattice \mathcal{D} in \mathbf{R}^d , and each finite collection $\mathcal{F} \subseteq \mathcal{D}$, there exists a sparse collection $\mathcal{S} \subseteq \mathcal{F}$ such that

$$M_{1,1}^{\mathcal{F}}(f, g) \leq 4 \sum_{Q \in \mathcal{S}} \langle f \rangle_{1,Q} \langle g \rangle_{1,Q} \mathbf{1}_Q,$$

where the superscript \mathcal{F} indicates that the defining supremum in the definition of $M_{1,1}$ is only taken over $Q \in \mathcal{F}$. Hence,

$$\begin{aligned} \|M_{1,1}^{\mathcal{F}}(f, g)\|_{L^1(\mathbf{R}^d)} &\leq 4 \sum_{Q \in \mathcal{S}} \langle f \rangle_{1,Q} \langle g \rangle_{1,Q} |Q| = 4 \|(T_S f)g\|_{L^1(\mathbf{R}^d)} \\ &\leq 4 \sup_{\mathcal{S} \text{ is sparse}} \|T_S\|_{X \rightarrow X} \|f\|_X \|g\|_{X'}. \end{aligned}$$

The assertion now follows from the monotone convergence theorem and the 3^d lattice theorem. □

Proof of Theorem 3.1 For 3.1 \Rightarrow 3.1, we first note that the sharp reverse Hölder inequality [30, Theorem 2.3] states that there is a dimensional constants $c_d \geq 1$ so that for all weights w such that

$$[w]_{A_\infty} := \sup_Q \frac{1}{w(Q)} \int_Q M(w \mathbf{1}_Q) < \infty$$

and all $r \in (1, \infty)$ satisfying $r' \geq c_d [w]_{A_\infty}$ we have

$$\langle w \rangle_{r,Q} \leq 2 \langle w \rangle_{1,Q}$$

for all cubes Q in \mathbf{R}^d .

Fix an $r_0 \in (1, r^*]$ with $r'_0 \geq 2c_d \|M\|_{X \rightarrow X}$, and let $r \in (1, r_0]$. For $f \in X^r$ we define

$$w := \sum_{k=0}^{\infty} \frac{M^k(|f|^{\frac{1}{r}})}{2^k \|M\|_{X \rightarrow X}^k},$$

where M^k denotes the k -th iterate of M . Then we have

$$c_d[w]_{A_\infty} \leq 2c_d \|M\|_{X \rightarrow X} \leq r',$$

and therefore $\langle w \rangle_{r,Q} \leq 2\langle w \rangle_{1,Q}$ for all cubes Q .

Let \mathcal{S} be a sparse collection. Since $|f|^{\frac{1}{r}} \leq w$ and $\|w\|_X \leq 2\|f\|_{X^r}^{\frac{1}{r}}$, we obtain

$$\begin{aligned} \|T_{\mathcal{S}}f\|_{X^r} &= \left\| \left(\sum_{Q \in \mathcal{S}} \langle |f|^{\frac{1}{r}} \rangle_{r,Q}^r \mathbf{1}_Q \right)^{\frac{1}{r}} \right\|_X^r \leq \left\| \sum_{Q \in \mathcal{S}} \langle w \rangle_{r,Q} \mathbf{1}_Q \right\|_X^r \\ &\leq 2^r \|T_{\mathcal{S}}w\|_X^r \leq 2^r \|T_{\mathcal{S}}\|_{X \rightarrow X}^r \|w\|_X^r \leq 4^r \|T_{\mathcal{S}}\|_{X \rightarrow X}^r \|f\|_{X^r}. \end{aligned}$$

Thus, we have

$$\sup_{\mathcal{S} \text{ is sparse}} \|T_{\mathcal{S}}\|_{X^r \rightarrow X^r} \leq 4^r \left(\sup_{\mathcal{S} \text{ is sparse}} \|T_{\mathcal{S}}\|_{X \rightarrow X} \right)^r.$$

The result now follows from Lemma 3.4.

The implication 3.1 \Rightarrow 3.1 is immediate. To prove 3.1 \Rightarrow 3.1, we apply [43, Theorem 4.16 and Remark 4.17] to the classical bound

$$\|M\|_{L_w^p(\mathbf{R}^d) \rightarrow L_w^p(\mathbf{R}^d)} \lesssim_d p' [w]_p^{p'}$$

by Buckley [6] to conclude that

$$\|M\|_{X \rightarrow X} \lesssim_d r' 2^{\frac{1}{r-1}} \|M\|_{(X^r)' \rightarrow (X^r)'}, \quad \|M\|_{X' \rightarrow X'} \lesssim_d r \|M\|_{(X^r)' \rightarrow (X^r)'}$$

This proves the result. □

Remark 3.5 Theorem 3.1 can easily be extended to Banach function spaces X over a space of homogeneous type (S, d, μ) in the sense of Coifman and Weiss [15]. Maximal operator bounds and sparse domination estimates in this setting are available through the use of e.g. Hytönen–Kairema cubes [27] (see also Christ [11]) and the sharp reverse Hölder inequality needs to be replaced by the weak sharp reverse Hölder inequality by Hytönen, Perez and Rela [31].

4 Full Range Compact Extrapolation

This section is dedicated to the proof of the full range extrapolation of compactness theorem, Theorem 1.1:

Theorem 4.1 *Let*

$$T : \bigcup_{\substack{p \in (1, \infty) \\ w \in A_p}} L_w^p(\mathbf{R}^d) \rightarrow L^0(\mathbf{R}^d)$$

be a linear operator such that

- *T is bounded on $L_w^p(\mathbf{R}^d)$ for some $p \in (1, \infty)$ and all weights $w \in A_p$;*
- *T is compact on $L_w^p(\mathbf{R}^d)$ for some $p \in (1, \infty)$ and some weight $w \in A_p$.*

Let X be a Banach function space over \mathbf{R}^d such that

$$M : X \rightarrow X, \quad M : X' \rightarrow X',$$

and assume X is r^ -convex and s^* -concave for some $1 < r^* < s^* < \infty$. Then $T : X \rightarrow X$ is compact.*

As noted in the introduction, we will need three main ingredients to prove Theorem 4.1. We will need the Rubio de Francia extrapolation result from [43], the self-improvement result from Sect. 3 and a result on the compactness of operators on product spaces. The latter is a special case of a result of Cobos, Kühn and Schonbek [13, Theorem 3.1] with the function parameter $\rho(t) = t^\theta$, which we formulate next.

Proposition 4.2 *[[13, Theorem 3.1]] Let (Ω, μ) be a σ -finite measure space and let X_0, X_1, Y_0, Y_1 be Banach function spaces over Ω . Let $T : X_0 + X_1 \rightarrow Y_0 + Y_1$ be a linear operator such that*

- *T is bounded from X_0 to Y_0 ;*
- *T is compact from X_1 to Y_1 .*

Then

$$T : X_0^{1-\theta} \cdot X_1^\theta \rightarrow Y_0^{1-\theta} \cdot Y_1^\theta$$

is compact for all $\theta \in (0, 1)$.

Having all main ingredients at our disposal, the proof of Theorem 4.1 is rather short.

Proof of Theorem 4.1 Note that, by the classical Rubio de Francia extrapolation theorem, T is bounded on $L_w^p(\mathbf{R}^d)$ for all $p \in (1, \infty)$ and $w \in A_p$. Moreover, since

$$L^2(\mathbf{R}^d) = L_w^p(\mathbf{R}^d)^{\frac{1}{2}} \cdot L_{w^{-1}}^{p'}(\mathbf{R}^d)^{\frac{1}{2}}$$

and $w \in A_p$ if and only if $w^{-1} \in A_{p'}$, Proposition 4.2 implies that T is compact on $L^2(\mathbf{R}^d)$.

By Corollary 3.3, there are $r_0 \in (1, r^*]$ and $s_0 \in [s^*, \infty)$ such that M is bounded on $X_{r,s}$ and $(X_{r,s})'$ for all $r \in (1, r_0]$ and $s \in [s_0, \infty)$. Hence, by Rubio de Francia extrapolation in Banach function spaces as in [43, Theorem A], T is bounded on $X_{r,s}$ for all $r \in (1, r_0]$ and $s \in [s_0, \infty)$.

By Proposition 2.3, we have

$$X = X_{r,s}^{1-\theta} \cdot L^p(\mathbf{R}^d)^\theta.$$

with $\theta = 1 - (\frac{1}{r} - \frac{1}{s}) \in (0, 1)$ and $p = 1 + \frac{s}{r'}$. Choosing $\frac{1}{r'} = \frac{1}{s}$ small enough, we have $p = 2$ and T is bounded on $X_{r,s}$. Since T is compact on $L^2(\mathbf{R}^d)$, T is compact on X by Proposition 4.2. This proves the result. □

5 Limited Range, Off-diagonal Compact Extrapolation

In this section, we prove Theorem 1.4. Essentially, the steps are the same as in the proof of Theorem 4.1, and the new difficulties lie mainly in unwinding the definitions while incorporating the additional parameters.

For $1 \leq r < p < s \leq \infty$, we say that a weight w belongs to the limited range Muckenhoupt class $A_{p,(r,s)}$ if

$$[w]_{p,(r,s)} := \sup_Q \langle w \rangle_{\frac{1}{p} - \frac{1}{s}, Q} \langle w^{-1} \rangle_{\frac{1}{r} - \frac{1}{p}, Q} < \infty.$$

For $\alpha \in \mathbf{R}$, $r_1, r_2 \in [1, \infty)$, $s_1, s_2 \in (1, \infty]$ for which $\frac{1}{s_j} < \frac{1}{r_j}$ for $j \in \{1, 2\}$ and

$$\frac{1}{r_1} - \frac{1}{r_2} = \frac{1}{s_1} - \frac{1}{s_2} = \alpha,$$

we note that we have $A_{p_1,(r_1,s_1)} = A_{p_2,(r_2,s_2)}$. For a weight w in this class we will write $w \in A_{\vec{p},(\vec{r},\vec{s})}$. Recall that our limited range, off-diagonal extrapolation of compactness theorem reads as follows:

Theorem 5.1 *Let $\alpha \in \mathbf{R}$ and let $r_1, r_2 \in [1, \infty)$, $s_1, s_2 \in (1, \infty]$ satisfy $\frac{1}{s_j} < \frac{1}{r_j}$ for $j \in \{1, 2\}$ and*

$$\frac{1}{r_1} - \frac{1}{r_2} = \frac{1}{s_1} - \frac{1}{s_2} = \alpha.$$

Define

$$\mathcal{P} := \{(p_1, p_2) \in (0, \infty]^2 : \frac{1}{p_j} \in [\frac{1}{s_j}, \frac{1}{r_j}], j \in \{1, 2\}, \frac{1}{p_1} - \frac{1}{p_2} = \alpha\}$$

and let

$$T : \bigcup_{\substack{(p_1, p_2) \in \mathcal{P} \\ w \in A_{\vec{p}, (\vec{r}, \vec{s})}}} L_w^{p_1}(\mathbf{R}^d) \rightarrow L^0(\mathbf{R}^d)$$

be a linear operator such that

- T is bounded from $L_w^{p_1}(\mathbf{R}^d)$ to $L_w^{p_2}(\mathbf{R}^d)$ for some $(p_1, p_2) \in \mathcal{P}$ and all $w \in A_{\vec{p}, (\vec{r}, \vec{s})}$;
- T is compact from $L_w^{p_1}(\mathbf{R}^d)$ to $L_w^{p_2}(\mathbf{R}^d)$ for some $(p_1, p_2) \in \mathcal{P}$ and some $w \in A_{\vec{p}, (\vec{r}, \vec{s})}$.

Let $r_j < r_j^* < s_j^* < s_j$ and let X_j be an r_j^* -convex and s_j^* -concave Banach function space for $j \in \{1, 2\}$ satisfying

$$(X_1)_{r_1, s_1} = (X_2)_{r_2, s_2}$$

and

$$M : (X_1)_{r_1, s_1} \rightarrow (X_1)_{r_1, s_1}, \quad M : ((X_1)_{r_1, s_1})' \rightarrow ((X_1)_{r_1, s_1})'$$

Then $T : X_1 \rightarrow X_2$ is compact.

Remark 5.2 The limited range, off-diagonal Rubio de Francia extrapolation theorem in [43] is phrased for quasi-Banach function spaces. However, one of our other main ingredients, the compactness result in Proposition 4.2, does not seem to be available in the quasi-setting. Therefore, we have stated Theorem 5.1 in the Banach function space setting and leave its extension to the quasi-Banach function space setting with $r_1, r_2 \in (0, \infty)$ as an open problem. Note that quasi-Banach function spaces typically only show up in harmonic analysis in multilinear or endpoint settings.

As said, the main challenge in the proof of Theorem 5.1 is to unpack all definitions and to keep track of the additional parameters. In order to do so, we prove a couple of technical lemmata. We start with a limited range version of the self-improvement property for the maximal operator.

Lemma 5.3 *Let $1 \leq r < r^* < s^* < s \leq \infty$ and let X be an r^* -convex and s^* -concave Banach function space. If*

$$M : X_{r, s} \rightarrow X_{r, s}, \quad M : (X_{r, s})' \rightarrow (X_{r, s})',$$

then there are $r_0 \in (r, r^*]$, $s_0 \in [s^*, s)$ so that for all $\tilde{r} \in (r, r_0)$, $\tilde{s} \in (s_0, s)$ we have

$$M : X_{\tilde{r}, \tilde{s}} \rightarrow X_{\tilde{r}, \tilde{s}}, \quad M : (X_{\tilde{r}, \tilde{s}})' \rightarrow (X_{\tilde{r}, \tilde{s}})'$$

Proof By Theorem 3.1 there is a $q_0 \in (1, r^*]$ so that for all $q \in (1, q_0]$ we have

$$M : X_{r, s}^q \rightarrow X_{r, s}^q, \quad M : (X_{r, s}^q)' \rightarrow (X_{r, s}^q)'$$

Defining

$$\frac{1}{r_0} := \frac{1}{q} \left(\frac{1}{r} - \frac{1}{s} \right) + \frac{1}{s} \in \left(\frac{1}{s}, \frac{1}{r} \right),$$

it follows from [43, Lemma 2.32] that for $q > 1$ chosen small enough so that $r_0 \leq r^*$, we have

$$X_{r,s}^q = X_{r_0,s}.$$

Note that $(X_{r_0,s})' = [(X^{r_0})']^{(\frac{s}{r_0})'}$ is t^* -convex, where

$$t^* := \frac{\frac{1}{r_0} - \frac{1}{s}}{\frac{1}{r_0} - \frac{1}{s^*}}.$$

Thus, by Theorem 3.1, we find a $t_0 \in (1, t^*)$ so that for all $t \in (1, t_0]$ we have that M is bounded on $[(X^{r_0})']^{t(\frac{s}{r_0})'}$ and $[(X^{r_0})']^{t(\frac{s}{r_0})'}$.

Now define

$$\frac{1}{s_0} := \frac{1}{r_0} - \frac{1}{t} \left(\frac{1}{r_0} - \frac{1}{s} \right) = \frac{1}{q} \frac{1}{t} \left(\frac{1}{r} - \frac{1}{s} \right) + \frac{1}{s} \in \left(\frac{1}{s}, \frac{1}{r_0} \right)$$

which, if $t > 1$ is chosen small enough, satisfies $s_0 \geq s^*$. Then we have

$$t \left(\frac{s}{r_0} \right)' = \left(\frac{s_0}{r_0} \right)'$$

so that

$$M : X_{r_0,s_0} \rightarrow X_{r_0,s_0}, \quad M : (X_{r_0,s_0})' \rightarrow (X_{r_0,s_0})'.$$

Now let $\tilde{r} \in (r, r_0)$, $\tilde{s} \in (s_0, s)$. Noting that $(X_{r_0,s_0})' = ((X^r)')_{(\frac{s_0}{r}, \frac{r_0}{r})'}$ by [43, Proposition 2.14], it follows from [43, Proposition 2.30] that also

$$M : X_{\tilde{r},\tilde{s}} \rightarrow X_{\tilde{r},\tilde{s}}, \quad M : (X_{\tilde{r},\tilde{s}})' \rightarrow (X_{\tilde{r},\tilde{s}})'.$$

The assertion follows. □

Next, we prove a rescaling lemma.

Lemma 5.4 *Let $1 \leq r < \tilde{r} < \tilde{s} < s \leq \infty$ and let X be an \tilde{r} -convex and \tilde{s} -concave Banach function space. Define*

$$\frac{1}{t} := \frac{\frac{1}{r} \frac{1}{\tilde{s}} - \frac{1}{\tilde{r}} \frac{1}{s}}{\frac{1}{r} - \frac{1}{s}}$$

Then X is t -concave, and $[X_{\tilde{r},t}]^{\frac{1}{\tilde{r}}}$ is r -convex and s -concave with

$$([X_{\tilde{r},t}]^{\frac{1}{\tilde{r}}})_{r,s} = X_{\tilde{r},\tilde{s}}.$$

Proof We have

$$\frac{1}{t} = \frac{\frac{1}{r}(\frac{1}{\tilde{s}} - \frac{1}{s}) + (\frac{1}{r} - \frac{1}{\tilde{r}})\frac{1}{s}}{\frac{1}{r} - \frac{1}{s}} > 0$$

and

$$\frac{1}{\tilde{s}} - \frac{1}{t} = \frac{1}{s} \frac{\frac{1}{\tilde{r}} - \frac{1}{\tilde{s}}}{\frac{1}{r} - \frac{1}{s}} \geq 0.$$

Thus, $t \geq \tilde{s}$ and, hence, X is t -concave. Therefore $([X_{\tilde{r},t}]^{\frac{1}{\tilde{r}}})' = X_{\tilde{r},t}$ is a Banach function space, and hence, $[X_{\tilde{r},t}]^{\frac{1}{\tilde{r}}}$ is r -convex. Moreover, since

$$\left(\frac{t}{\tilde{r}}\right)' \left(\frac{s}{\tilde{r}}\right)' = \left(\frac{\tilde{s}}{\tilde{r}}\right)'$$

we have

$$([([X_{\tilde{r},t}]^{\frac{1}{\tilde{r}}})')^{(\frac{t}{\tilde{r}})'})^{(\frac{s}{\tilde{r}})'} = [(X^{\tilde{r}})']^{(\frac{\tilde{s}}{\tilde{r}})'} = (X_{\tilde{r},\tilde{s}})'$$

which is a Banach function space, since X is \tilde{r} -convex and \tilde{s} -concave. This proves that $[X_{\tilde{r},t}]^{\frac{1}{\tilde{r}}}$ is s -concave. Moreover, taking associate spaces in this equality, the final assertion follows. \square

We finish our preparation for the proof of Theorem 5.1 with a factorization lemma in the limited range setting.

Lemma 5.5 *Let $1 \leq r < \tilde{r} < \tilde{s} < s \leq \infty$ and let X be an \tilde{r} -convex and \tilde{s} -concave Banach function space. Define*

$$\frac{1}{t} := \frac{\frac{1}{r}\frac{1}{\tilde{s}} - \frac{1}{\tilde{r}}\frac{1}{s}}{\frac{1}{r} - \frac{1}{s}}$$

and

$$\theta := 1 - \frac{\frac{1}{\tilde{r}} - \frac{1}{\tilde{s}}}{\frac{1}{r} - \frac{1}{s}} \in (0, 1), \quad p = \frac{\frac{1}{\tilde{s}} - \frac{1}{s} + \frac{1}{r} - \frac{1}{\tilde{r}}}{\frac{1}{r}(\frac{1}{\tilde{s}} - \frac{1}{s}) + (\frac{1}{r} - \frac{1}{\tilde{r}})\frac{1}{s}} \in (1, \infty).$$

Then

$$X = ([X_{\tilde{r},t}]^{\frac{1}{\tilde{r}}})^{1-\theta} \cdot L^p(\mathbf{R}^d)^\theta.$$

Proof Since

$$\frac{1}{r}(1 - \theta) = \frac{1}{\tilde{r}} - \frac{1}{t}$$

and $t = \frac{p}{\theta}$, we have by Proposition 2.3 and Lemma 5.4

$$([X_{\tilde{r},t}]^{\frac{1}{r}})^{1-\theta} \cdot L^p(\mathbf{R}^d)^\theta = (X_{\tilde{r},t})^{\frac{1}{r}-\frac{1}{t}} \cdot L^t(\mathbf{R}^d) = X,$$

as asserted. □

Proof of Theorem 5.1 Let $(p_1, p_2) \in \mathcal{P}$ and $w \in A_{\vec{p},(\vec{r},\vec{s})}$ so that $T : L_w^{p_1}(\mathbf{R}^d) \rightarrow L_w^{p_2}(\mathbf{R}^d)$ is compact. Note that, by limited range, off-diagonal Rubio de Francia extrapolation for weighted Lebesgue spaces as in [16, Theorem 1.8] (which is also a special case of [43, Theorem A]), we actually obtain the boundedness assumption on T for all $(q_1, q_2) \in \mathcal{P}$ and weights in $A_{\vec{q},(\vec{r},\vec{s})}$. In particular, this is the case for $(q_1, q_2) \in \mathcal{P}$ defined by

$$\frac{1}{q_j} := \frac{1}{s_j} + \frac{1}{r_j} - \frac{1}{p_j}, \quad j \in \{1, 2\},$$

and the weight $w^{-1} \in A_{\vec{q},(\vec{r},\vec{s})}$, which one can directly verify using the definition of the weight constant. Since

$$L^{\frac{1}{\frac{1}{2}(\frac{1}{r_j} + \frac{1}{s_j})}}(\mathbf{R}^d) = L_w^{p_j}(\mathbf{R}^d)^{\frac{1}{2}} \cdot L_{w^{-1}}^{q_j}(\mathbf{R}^d)^{\frac{1}{2}},$$

it follows Proposition 4.2 that $T : L^{\frac{1}{\frac{1}{2}(\frac{1}{r_1} + \frac{1}{s_1})}}(\mathbf{R}^d) \rightarrow L^{\frac{1}{\frac{1}{2}(\frac{1}{r_2} + \frac{1}{s_2})}}(\mathbf{R}^d)$ is compact. Thus, we have reduced the compactness assumption to the case $\frac{1}{p_j} = \frac{1}{2}(\frac{1}{r_j} + \frac{1}{s_j})$ for $j \in \{1, 2\}$ and $w = 1$.

Next, by Lemma 5.3 we have

$$M : (X_1)_{\tilde{r}_1, \tilde{s}_1} \rightarrow (X_1)_{\tilde{r}_1, \tilde{s}_1}, \quad M : ((X_1)_{\tilde{r}_1, \tilde{s}_1})' \rightarrow ((X_1)_{\tilde{r}_1, \tilde{s}_1})'$$

for all $\tilde{r}_1 \in (r_1, r_1^*]$, $\tilde{s}_1 \in [s_1^*, s_1)$ with

$$\frac{1}{\tilde{s}_1} - \frac{1}{s_1} = \frac{1}{r_1} - \frac{1}{\tilde{r}_1} = \varepsilon$$

for $\varepsilon > 0$ small enough.

Defining $\frac{1}{\tilde{r}_2} := \frac{1}{r_1} - \alpha$ and $\frac{1}{\tilde{s}_2} := \frac{1}{s_1} - \alpha$, it follows from [43, Lemma 2.32] that for

$$\beta := \frac{\frac{1}{r_1} - \frac{1}{s_1}}{\frac{1}{\tilde{r}_1} - \frac{1}{\tilde{s}_1}} = \frac{\frac{1}{r_2} - \frac{1}{s_2}}{\frac{1}{\tilde{r}_2} - \frac{1}{\tilde{s}_2}}, \quad \gamma := \frac{\frac{1}{\tilde{r}_1} - \frac{1}{r_1}}{\frac{1}{\tilde{r}_1} - \frac{1}{r_1} + \frac{1}{s_1} - \frac{1}{\tilde{s}_1}} = \frac{\frac{1}{\tilde{r}_2} - \frac{1}{r_2}}{\frac{1}{\tilde{r}_2} - \frac{1}{r_2} + \frac{1}{s_2} - \frac{1}{\tilde{s}_2}}$$

we have

$$\begin{aligned} (X_1)_{\tilde{r}_1, \tilde{s}_1} &= (X_1)_{r_1, s_1}^\beta \cdot L^{\frac{1}{1-\gamma}}(\mathbf{R}^d)^{1-\beta} \\ &= (X_2)_{r_2, s_2}^\beta \cdot L^{\frac{1}{1-\gamma}}(\mathbf{R}^d)^{1-\beta} = (X_2)_{\tilde{r}_2, \tilde{s}_2}. \end{aligned}$$

Hence, by Lemma 5.4 we conclude that

$$\left([(X_1)_{\tilde{r}_1, t_1}]^{\frac{1}{r_1}} \right)_{r_1, s_1} = (X_1)_{\tilde{r}_1, \tilde{s}_1} = (X_2)_{\tilde{r}_2, \tilde{s}_2} = \left([(X_2)_{\tilde{r}_2, t_2}]^{\frac{1}{r_2}} \right)_{r_2, s_2},$$

where

$$\frac{1}{t_j} = \frac{\frac{1}{r_j} \frac{1}{s_j} - \frac{1}{\tilde{r}_j} \frac{1}{s_j}}{\frac{1}{r_j} - \frac{1}{s_j}}, \quad j \in \{1, 2\}.$$

Thus, by [43, Theorem A] it follows that

$$T : [(X_1)_{\tilde{r}_1, t_1}]^{\frac{1}{r_1}} \rightarrow [(X_2)_{\tilde{r}_2, t_2}]^{\frac{1}{r_2}}.$$

Moreover, by Lemma 5.5 we have

$$\begin{aligned} X_1 &= \left([(X_1)_{\tilde{r}_1, t_1}]^{\frac{1}{r_1}} \right)^{1-\theta} \cdot L^{p_1}(\mathbf{R}^d)^\theta, \\ X_2 &= \left([(X_2)_{\tilde{r}_2, t_2}]^{\frac{1}{r_2}} \right)^{1-\theta} \cdot L^{p_2}(\mathbf{R}^d)^\theta \end{aligned}$$

with

$$\theta = 1 - \frac{\frac{1}{\tilde{r}_1} - \frac{1}{\tilde{s}_1}}{\frac{1}{r_1} - \frac{1}{s_1}} = 1 - \frac{\frac{1}{r_2} - \frac{1}{s_2}}{\frac{1}{r_2} - \frac{1}{s_2}},$$

and

$$\frac{1}{p_j} = \frac{1}{2} \left(\frac{1}{r_j} + \frac{1}{s_j} \right), \quad j \in \{1, 2\}.$$

Hence, it follows from Proposition 4.2 that $T : X_1 \rightarrow X_2$ is compact. This proves the assertion. □

6 Applications

In this final section, we will briefly outline applications of Theorem 1.1 and Theorem 1.4 to the compactness of commutators of singular integral operators and multiplication by functions with vanishing mean oscillation. We refer the reader to [29, Sect. 5–10] for further examples of operators to which Theorem 1.1 or Theorem 1.4 are applicable.

We start with an application to Calderón–Zygmund singular integral operators. Let $T : L^2(\mathbf{R}^d) \rightarrow L^2(\mathbf{R}^d)$ be a bounded linear operator and suppose that, for any $f \in C_c^\infty(\mathbf{R}^d)$, T has the representation

$$Tf(x) = \int_{\mathbf{R}^d} K(x, y)f(y) \, dy, \quad x \in \mathbf{R}^d \setminus \text{supp}(f),$$

where the kernel satisfies the estimates

$$\begin{aligned}
 |K(x, y) - K(x, y')| &\leq \omega\left(\frac{|x-x'|}{|x-y|}\right), & 0 < |x-x'| < \frac{1}{2}|x-y|, \\
 |K(x, y) - K(x, y')| &\leq \omega\left(\frac{|x-x'|}{|x-y|}\right), & 0 < |y-y'| < \frac{1}{2}|x-y|,
 \end{aligned}$$

for some increasing, subadditive $\omega : [0, 1] \rightarrow [0, \infty)$. If $\int_0^1 \omega(t) \frac{dt}{t} < \infty$, we call T a Calderón–Zygmund operator with Dini-continuous kernel.

We will be concerned with the commutators

$$[b, T]f := bT(f) - T(bf)$$

for pointwise multipliers $b \in L^1_{\text{loc}}(\mathbf{R}^d)$ belonging to the space

$$\text{CMO} := \overline{C_c^\infty(\mathbf{R}^d)}^{\|\cdot\|_{\text{BMO}}},$$

where BMO denotes the classical space of functions with bounded mean oscillation. Note that the space CMO is the space of all functions with vanishing mean oscillation and is therefore also denoted by VMO in the literature, see, e.g., [47, Lemma 3].

Theorem 6.1 *Let T be a Calderón–Zygmund operator with Dini-continuous kernel. Let $1 < r < s < \infty$ and suppose X is an r -convex and s -concave Banach function space over \mathbf{R}^d such that*

$$M : X \rightarrow X, \quad M : X' \rightarrow X'.$$

For $b \in \text{CMO}$, the commutator $[b, T] : X \rightarrow X$ is a compact operator.

Proof In view of Theorem 1.1, it suffices to check that $[b, T]$ is bounded on $L^2_w(\mathbf{R}^d)$ for all $w \in A_2$ and that $[b, T]$ is compact on $L^2(\mathbf{R}^d)$. For the weighted boundedness it suffices to check that T is bounded on $L^2_w(\mathbf{R}^d)$ for all $w \in A_2$ by [1, Theorem 2.13] (see [5] for a modern approach). The boundedness of T on $L^2_w(\mathbf{R}^d)$ for all $w \in A_2$ is classical (see [32, 37] for a modern approach). Finally, the compactness of $[b, T]$ on $L^2(\mathbf{R}^d)$ was shown by Uchiyama [47]. \square

Theorem 6.1 in the specific case where X is a weighted Lebesgue space was previously obtained by Clop and Cruz [12] and subsequently proven using an extrapolation of compactness argument by Hytönen and Lappas [29]. Theorem 6.1 yields many examples of Banach function spaces X for which the compactness of $[b, T]$ was previously unknown.

For example, one can consider weighted variable Lebesgue spaces $X = L^{p(\cdot)}_w(\mathbf{R}^d)$. Here the function $p : \mathbf{R}^d \rightarrow (0, \infty)$ has to satisfy

$$1 < \text{ess inf } p \leq \text{ess sup } p < \infty \tag{6.1}$$

so that we can take $r = \text{ess inf } p, s = \text{ess sup } p$ in Theorem 1.1, and some additional condition ensuring the boundedness of M on X (see, e.g., [21, 34] for the unweighted

setting and [17] for the weighted setting). As a matter of fact, it was originally shown by Diening in [20, Theorem 8.1] that in the case of unweighted variable Lebesgue spaces satisfying Eq. (6.1) we have $M : X \rightarrow X$ if and only if $M : X' \rightarrow X'$. We also refer the reader to [34] for an explicit characterization of when M is bounded on X in terms of the exponent function p . Diening's duality result was extended to weighted variable Lebesgue spaces $X = L_w^{p(\cdot)}(\mathbf{R}^d)$ by Lerner in [35]. In conclusion, in the setting of weighted variable Lebesgue spaces, the boundedness condition of M in Theorem 1.1 only needs to be checked on X , as the boundedness on X' is then guaranteed.

We refer the reader to [43] for further examples of Banach function spaces satisfying the conditions in Theorem 6.1. We note that for the case that X is a Morrey space, similar results were obtained by Arai and Nakai [2] and Lappas [33], which lie beyond the scope of our general framework as explained in the introduction.

Next, let us consider rough singular integral operators

$$T_{\Omega} f := \text{p.v.} \int_{\mathbf{R}^d} \frac{\Omega(x-y)}{|x-y|^d} f(y) dy, \quad x \in \mathbf{R}^d,$$

where $\Omega \in L^r(\mathbf{S}^{d-1})$ for some $r \in [1, \infty]$ is homogeneous of order zero and has mean value zero.

Theorem 6.2 *Let $1 \leq r < r^* < s^* < \infty$ and let $\Omega \in L^{r'}(\Omega)$. Suppose X is an r^* -convex and s^* -concave Banach function space over \mathbf{R}^d such that*

$$M : X^r \rightarrow X^r, \quad M : (X^r)' \rightarrow (X^r)'.$$

For $b \in \text{CMO}$, the commutator $[b, T_{\Omega}] : X \rightarrow X$ is a compact operator.

Proof In view of Theorem 1.4, it suffices to check that $[b, T]$ is bounded on $L_w^p(\mathbf{R}^d)$ for some $p \in (r, \infty)$ and all $w \in A_{p, (r, \infty)}$ and that $[b, T]$ is compact on $L^p(\mathbf{R}^d)$. As in the proof of Theorem 6.1, for the weighted boundedness it suffices note that T_{Ω} is bounded on $L_w^p(\mathbf{R}^d)$ for all $w \in A_{p, (r, \infty)}$, which was shown independently by Watson [48] and Duoandikoetxea [22]. The compactness of $[b, T_{\Omega}]$ on $L^p(\mathbf{R}^d)$ was shown by Chen and Hu [10]. \square

Theorem 6.2 in the specific case where X is a weighted Lebesgue space was previously obtained by Guo and Hu [24] and subsequently proven using an extrapolation of compactness argument by Hytönen and Lappas [29]. In the case that X is a Morrey space, similar results were also obtained in [24] and [33].

To conclude our applications section, we note that in the recent work by Tao, Yang, Yuan and Zhang [46], compactness of $[b, T_{\Omega}]$ for $b \in \text{CMO}$ was studied in the setting of Banach function spaces as well. Let us give a comparison between Theorem 6.2 in the specific case $r = 1$ and [46, Theorem 3.1]

- We work with Banach function spaces as in Sect. 2, whereas [46] uses ball Banach function spaces, the former being a more general in the sense that any ball Banach function space is also a Banach function space in the sense of Sect. 2. However,

since we assume M to be bounded on X and X' , so in particular $\mathbf{1}_B \in X$ and $\mathbf{1}_B \in X'$ for all balls $B \subseteq \mathbf{R}^d$, we also have that any Banach function space satisfying the assumptions of Theorem 6.2 is a ball Banach function space.

- In [46] it is assumed that there is an $r^* > 1$ such that X^{r^*} is a Banach function space (i.e. X is r^* -convex) and M is bounded on $(X^{r^*})'$. This is equivalent to M being bounded on X and X' by Theorem 1.3. Thus, the only difference in the assumptions on X between Theorem 6.2 with $r = 1$ and [46, Theorem 3.1] is that Theorem 6.2 in addition assumes X to be s -concave for some $s < \infty$. This, as discussed before, is a necessary assumption in our general approach.
- In Theorem 6.2 with $r = 1$ it is assumed that $\Omega \in L^\infty(\mathbb{S}^{d-1})$, whereas in [46, Theorem 3.1] a much stronger assumption on Ω is imposed. Indeed, Ω is assumed to satisfy a Dini continuity condition, see [46, Definition 2.15].
- The proof of Theorem 6.2 uses “soft” techniques, whereas [46] takes a more hands-on and technical approach, developing and using a Frechet–Kolmogorov compactness criterion in ball Banach function spaces (see also [25]).

Funding Z. N. is supported by the grant Juan de la Cierva formación 2021 FJC2021-046837-I, the Basque Government through the BERC 2022–2025 program, by the Spanish State Research Agency project PID2020-113156GB-I00/AEI/10.13039/501100011033 and through BCAM Severo Ochoa excellence accreditation SEV-2023-2026.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Álvarez, J., Bagby, R.J., Kurtz, D.S., Pérez, C.: Weighted estimates for commutators of linear operators. *Stud. Math.* **104**(2), 195–209 (1993)
2. Arai, R., Nakai, E.: Compact commutators of Calderón-Zygmund and generalized fractional integral operators with a function in generalized Campanato spaces on generalized Morrey spaces. *Tokyo J. Math.* **42**(2), 471–496 (2019)
3. Auscher, P., Martell, J.M.: Weighted norm inequalities, off-diagonal estimates and elliptic operators. I. General operator theory and weights. *Adv. Math.* **212**(1), 225–276 (2007)
4. Bennett, C., Sharpley, R.: Interpolation of operators. In: *Pure and Applied Mathematics*, vol. 129. Academic Press Inc, Boston (1988)
5. Bényi, Á., Martell, J.M., Moen, K., Stachura, E., Torres, R.H.: Boundedness results for commutators with BMO functions via weighted estimates: a comprehensive approach. *Math. Ann.* **376**(1–2), 61–102 (2020)
6. Buckley, S.: Estimates for operator norms on weighted spaces and reverse Jensen inequalities. *Trans. Amer. Math. Soc.* **340**(1), 253–272 (1993)
7. Calderón, A.P.: Intermediate spaces and interpolation, the complex method. *Stud. Math.* **24**, 113–190 (1964)
8. Cao, M., Olivo, A., Yabuta, K.: Extrapolation for multilinear compact operators and applications. *Trans. Amer. Math. Soc.* **375**(7), 5011–5070 (2022)

9. Cao, M., Marín, J.J., Martell, J.M.: Extrapolation on function and modular spaces, and applications. *Adv. Math.* **406**, 108520 (2022)
10. Chen, J., Hu, G.: Compact commutators of rough singular integral operators. *Canad. Math. Bull.* **58**(1), 19–29 (2015)
11. Christ, M.: Lectures on singular integral operators. In: *CBMS Regional Conference Series in Mathematics*, vol. 77. American Mathematical Society, Providence (1990)
12. Clop, A., Cruz, V.: Weighted estimates for Beltrami equations. *Ann. Acad. Sci. Fenn. Math.* **38**(1), 91–113 (2013)
13. Cobos, F., Kühn, T., Schonbek, T.: One-sided compactness results for Aronszajn-Gagliardo functors. *J. Funct. Anal.* **106**(2), 274–313 (1992)
14. Cobos, F., Fernández-Cabrera, L.M., Martínez, A.: On compactness results of Lions-Peetre type for bilinear operators. *Nonlinear Anal.* **199**, 111951 (2020)
15. Coifman, R.R., Weiss, G.: Analyse harmonique non-commutative sur certains espaces homogènes. In: *Lecture Notes in Mathematics*, vol. 242. Springer, Berlin (1971)
16. Cruz-Uribe, D., Martell, J.M.: Limited range multilinear extrapolation with applications to the bilinear Hilbert transform. *Math. Ann.* **371**(1–2), 615–653 (2018)
17. Cruz-Uribe, D., Fiorenza, A., Neugebauer, C.J.: Weighted norm inequalities for the maximal operator on variable Lebesgue spaces. *J. Math. Anal. Appl.* **394**(2), 744–760 (2012)
18. Curbera, G.P., García-Cuerva, J., Martell, J.M., Pérez, C.: Extrapolation with weights, rearrangement-invariant function spaces, modular inequalities and applications to singular integrals. *Adv. Math.* **203**(1), 256–318 (2006)
19. Cwikel, M., Nilsson, P.G., Schechtman, G.: Interpolation of weighted Banach lattices: a characterization of relatively decomposable Banach lattices. *Mem. Amer. Math. Soc.* **165**(787), 127 (2003)
20. Diening, L.: Maximal function on Musielak-Orlicz spaces and generalized Lebesgue spaces. *Bull. Sci. Math.* **129**(8), 657–700 (2005)
21. Diening, L., Harjulehto, P., Hästö, P., Ružička, M.: Lebesgue and Sobolev spaces with variable exponents. In: *Lecture Notes in Mathematics*, vol. 2017. Springer, Heidelberg (2011)
22. Duoandikoetxea, J.: Weighted norm inequalities for homogeneous singular integrals. *Trans. Amer. Math. Soc.* **336**(2), 869–880 (1993)
23. García-Cuerva, J., de Francia, J.L.R.: Weighted norm inequalities and related topics. In: *North-Holland Mathematics Studies*. North-Holland Publishing, Amsterdam (1985)
24. Guo, X., Hu, G.: On the commutators of singular integral operators with rough convolution kernels. *Canad. J. Math.* **68**(4), 816–840 (2016)
25. Guo, W., Zhao, G.: On relatively compact sets in quasi-Banach function spaces. *Proc. Amer. Math. Soc.* **148**(8), 3359–3373 (2020)
26. Harboure, E., Macías, R., Segovia, C.: Extrapolation results for classes of weights. *Amer. J. Math.* **110**(3), 383–397 (1988)
27. Hytönen, T.P., Kairema, A.: Systems of dyadic cubes in a doubling metric space. *Colloq. Math.* **126**(1), 1–33 (2012)
28. Hytönen, T.P., Lappas, S.: Extrapolation of compactness on weighted spaces: bilinear operators. *Indag. Math.* **33**(2), 397–420 (2022)
29. Hytönen, T.P., Lappas, S.: Extrapolation of compactness on weighted spaces. *Rev. Mat. Iberoam.* **39**(1), 91–122 (2023)
30. Hytönen, T.P., Pérez, C.: Sharp weighted bounds involving A_∞ . *Anal. PDE* **6**(4), 777–818 (2013)
31. Hytönen, T.P., Pérez, C., Rela, E.: Sharp reverse Hölder property for A_∞ weights on spaces of homogeneous type. *J. Funct. Anal.* **263**(12), 3883–3899 (2012)
32. Lacey, M.T.: An elementary proof of the A_2 bound. *Israel J. Math.* **217**(1), 181–195 (2017)
33. Lappas, S.: Extrapolation of compactness on weighted Morrey spaces. *Stud. Math.* **265**(2), 177–195 (2022)
34. Lerner, A.K.: A boundedness criterion for the maximal operator on variable Lebesgue spaces. <https://arxiv.org/abs/2302.02475> (2023)
35. Lerner, A.K.: On a dual property of the maximal operator on weighted variable L^p spaces. In: *Functional Analysis, Harmonic Analysis, and Image Processing: A Collection of Papers in Honor of Björn Jawerth*, pp. 288–300. American Mathematical Society, Providence (2017)
36. Lerner, A.K., Ombrosi, S.: A boundedness criterion for general maximal operators. *Publ. Mat.* **54**(1), 53–71 (2010)

37. Lerner, A.K., Ombrosi, S.: Some remarks on the pointwise sparse domination. *J. Geom. Anal.* **30**(1), 1011–1027 (2020)
38. Lindenstrauss, J., Tzafriri, L.: *Classical Banach Spaces*, vol. 97. Springer, Berlin (1979)
39. Liu, S., Wu, H., Yang, D.: A note on extrapolation of compactness. *Collect. Math.* **74**(2), 375–390 (2023)
40. Lorist, E., Nieraeth, Z.: Banach function spaces done right. *Indag. Math.* **35**(2), 247–268 (2024)
41. Lozanovskii, G., Ya.: On some Banach lattices. *Sib. Math. J.* **10**(3), 419–431 (1969)
42. Nieraeth, Z.: *Sharp Estimates and Extrapolation for Multilinear Weight Classes*. Delft University of Technology, Delft (2020)
43. Nieraeth, Z.: Extrapolation in general quasi-Banach function spaces. *J. Funct. Anal.* **285**(10), 110130 (2023)
44. Rubio de Francia, J.L.: Factorization and extrapolation of weights. *Bull. Amer. Math. Soc.* **7**(2), 393–395 (1982)
45. Sawano, Y., Ho, K.P., Yang, D., Yang, S.: Hardy spaces for ball quasi-Banach function spaces. *Dissertationes Math.* **525**, 102 (2017)
46. Tao, J., Yang, D., Yuan, W., Zhang, Y.: Compactness characterizations of commutators on ball Banach function spaces. *Potential Anal.* **58**(4), 645–679 (2023)
47. Uchiyama, A.: On the compactness of operators of Hankel type. *Tohoku Math. J.* **30**(1), 163–171 (1978)
48. Watson, D.K.: Weighted estimates for singular integrals via Fourier transform estimates. *Duke Math. J.* **60**(2), 389–399 (1990)
49. Zaanen, A.C.: *Integration*. North-Holland Publishing Co., Amsterdam (1967)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.