

# Analysis of 6-methoxy podophyllotoxin and podophyllotoxin in hairy root cultures of *Linum album* Kotschy ex Boiss

Najmeh Ahmadian Chashmi · Mozafar Sharifi ·  
Morteza Yousefzadi · Mehrdad Behmanesh ·  
Hassan Rezadoost · Alejandra Cardillo · Javier Palazon

Received: 23 January 2012 / Accepted: 18 April 2012 / Published online: 4 May 2012  
© Springer Science+Business Media, LLC 2012

**Abstract** *Linum album* is a herbaceous medicinal plant that contains some lignans with antiviral and anticancer properties such as podophyllotoxin (PTOX) and 6-methoxy podophyllotoxin (MPTOX). In this research, hairy root cultures of *L. album* were established by transformation with *Agrobacterium rhizogenes* strains LBA9402, A4, AR15834 and *Agrobacterium tumefaciens* strain C58C1 (pRiA4). The presence of PTOX and MPTOX in the hairy

roots was verified by ESI/MS in positive ion mode. MPTOX was confirmed and its enantiomer determined by nuclear magnetic resonance spectroscopy and circular dichroism spectroscopy, respectively. PTOX and MPTOX production was determined by HPLC, in different lines of hairy roots. The results showed that all obtained hairy root lines produced higher yield of lignans than mother plant roots. In addition, the lignan content in the roots derived from *A. rhizogenes* strain LBA9402 was higher than in those obtained from *A. tumefaciens* strain C58C1.

**Electronic supplementary material** The online version of this article (doi:10.1007/s00044-012-0067-1) contains supplementary material, which is available to authorized users.

**Keywords** *Agrobacterium* · Hairy roots · *Linum album* · Podophyllotoxin (PTOX) · 6-methoxy podophyllotoxin (MPTOX)

N. A. Chashmi · M. Sharifi (✉)  
Department of Plant Biology, Faculty of Biological Sciences,  
Tarbiat Modares University, P.O. Box: 14115-154, Tehran, Iran  
e-mail: msharifi@modares.ac.ir

## Abbreviations

Ags	Agropine synthase
CD	Circular dichroism
LC–ESI–MS/MS	Liquid chromatography–electrospray–mass spectrometry/mass spectrometry
MPTOX	6-methoxy podophyllotoxin
NMR	Nuclear magnetic resonance
PTOX	Podophyllotoxin

M. Yousefzadi  
Department of Biology, Faculty of Sciences, Hormozgan  
University, Bandar Abbas, Iran

M. Behmanesh  
Department of Genetics, Faculty of Biological Sciences,  
Tarbiat Modares University, Tehran, Iran

H. Rezadoost  
Department of Phytochemistry, Medicinal Plants and Drugs  
Research Institute, Shahid Beheshti University, Tehran, Iran

A. Cardillo  
Catedra de Microbiología Industrial y Biotecnología,  
Facultad de Farmacia y Bioquímica, Universidad de Buenos  
Aires, Buenos Aires, Argentina

J. Palazon  
Laboratori de Fisiologia Vegetal, Facultat de Farmacia,  
Universitat de Barcelona, Barcelona, Spain

## Introduction

Lignans constitute a large group of secondary metabolites synthesized by many plants. These compounds are usually formed from two phenylpropanoid units and manifest considerable biological activity (Smolny *et al.*, 1998). Podophyllotoxin (PTOX), a lignan with antiviral and antineoplastic activities, is used today primarily as a

precursor for the semi-synthesis of established cancer therapeutics such as etoposide, teniposide and etopophos (Gordaliza *et al.*, 2004). To date, extracting PTOX and 6-methoxy podophyllotoxin (MPTOX) from the subterrrestrial parts of *Podophyllum* species is still preferred to their chemical synthesis due to low product yields and the complexity of the enantioselective reactions (Botta *et al.*, 2001). Plants from the genus *Linum* (the family Linaceae), with their ability to synthesize and accumulate lignans in callus and hairy root cultures, provide an additional source of PTOX and MPTOX (Farkya and Bisaria, 2008).

Biotechnological methods are being increasingly studied for their application as alternative ways of obtaining these valuable metabolites (Ramachandra and Ravishankar, 2002), although only a few have so far been produced at an industrial level owing to certain limitations (Chattopadhyay *et al.*, 2002; Zhong, 2002). As suspension cultures are not genetically stable (Bais *et al.*, 2001), new methods have been sought to improve lignan production. Hairy roots are transformed roots derived from the infection of wounded plant tissue by the soil bacterium, *Agrobacterium rhizogenes*. Such hairy roots often grow as fast as plant cell cultures or faster, and do not require hormones in the growth medium (Giri and Narasu, 2000). The greatest advantage of using hairy roots is that their biosynthetic capacity for metabolite production is similar to that of the mother plants or greater. Most of the research on the production of PTOX and related lignans has used undifferentiated cell cultures derived from non-transformed explants of naturally growing plants or aseptically growing seedlings (Smolny *et al.*, 1998; Farkya *et al.*, 2004; Kuhlmann *et al.*, 2002; Nikolay and Ionkova, 2005). The *in vitro* production of PTOX in *Podophyllum hexandrum* has been reported at the bioreactor level under batch, fed-batch and continuous modes with optimized medium conditions only in non-transformed undifferentiated cell suspension cultures (Chattopadhyay *et al.*, 2002, 2003a, b, c). Transformed calli of *P. hexandrum* obtained by infecting its embryos with *A. rhizogenes*, in the presence of phytohormones, showed a threefold increase in PTOX content compared with non-transformed cell cultures (Giri *et al.*, 2001).

In this study, hairy root cultures were established by infecting derived leaf segments of *L. album* with *A. rhizogenes* strains LBA9402, A4, AR15834 and *A. tumefaciens* strain C58C1 (pRiA4). PTOX and MPTOX production in two lines of hairy roots was compared. The presence of PTOX and MPTOX in the samples was verified by ESI/MS in positive ion mode. In order to confirm MPTOX and determine its enantiomer, nuclear magnetic resonance (NMR) spectroscopy and circular dichroism (CD) spectroscopy were used, respectively.

## Experimental

### Plant material

*Linum album* seeds were collected from Sohanak, (35° 48' N, 51° 32' E and altitude of 1,900 m) Iran. The plant material was kindly identified by Dr. Shahrokh Kazempour (Tarbiat Modares University, Tehran, Iran) and a voucher specimen was deposited at the herbarium of the Biological Sciences Faculty, Tarbiat Modares University, Tehran, Iran. The seeds were surface-sterilized with 70 % (v/v) ethanol for 70 s, and 2 % (v/v) sodium hypochlorite for 20 min, and then were washed with sterile distilled water more than three times. Sterile seeds were treated with GA<sub>3</sub> (500 mg L<sup>-1</sup>) overnight, then cultured on hormone-free MS medium supplemented with sucrose (3 %) (Murashige and Skoog, 1962) and incubated at 25 °C with a 16-h photoperiod. Sterile grown seedlings were used to initiate cultures.

### Agrobacterium culture

*Agrobacterium rhizogenes* strains LBA9402, A4, AR15834 and *A. tumefaciens* C58C1 (pRiA4) were used for infections. The bacteria from glycerol stock (stored at -80 °C) were cultured on LB solid medium (Park and Facchini, 2000) containing 50 mg L<sup>-1</sup> (w/v) rifampicin for screening and selecting the desired colony. Bacterial suspensions were incubated in liquid LB medium for 2 days at 28 °C.

### Plant transformation

Two procedures were tested for initiation of *L. album* hairy roots: the syringe needle and the leaf disk methods (Lin *et al.*, 2003). In the first procedure, separate syringe needles were contaminated with *Agrobacteria* colonies strains LBA9402, A4, AR15834 and C58C1 (pRiA4) and used to infect 4-week-old *L. album* seedlings by pricking the leaves and stems, and then placed on solid MS medium as described above. Also, seedlings pricked with a sterile syringe needle, without infection, were used as the control. After incubation in the dark at 25 °C for 2 days, the infected seedlings were placed on solid MS medium containing 200 mg L<sup>-1</sup> cefotaxime antibiotic and incubated at 25 °C.

In the leaf disk method, leaves of *L. album* seedlings were excised and cut in half transversally. The leaf disks were immersed in the *A. rhizogenes* suspension for 10 min, blotted to remove excess bacteria, and placed on solid MS medium as described above. Uninfected sterile leaf explants were grown on the MS medium culture as the control. The obtained hairy roots were cultured in MS medium by serial sub-culturing every 2 weeks.

## PCR analysis

DNA was extracted from non-transformed root, obtained from seedlings, and transformed hairy roots using the acetyl trimethyl ammonium bromide (CTAB) method (Khan *et al.*, 2007). The transgenic nature of hairy roots was confirmed by PCR *rolC*, *Ags* (agropine synthase) and *VirD* specific primers. Sequences of these primers are shown in Table 1. Amplification conditions for *rolC*, *Ags* and *VirD* were 1 cycle at 94 °C for 2 min followed by 35 cycles of amplification (45 s at 94 °C, 40 s at 55 °C and 45 s at 72 °C) and 1 cycle at 72 °C for 10 min. PCR products were analyzed by electrophoresis in a 1 % (w/v) agarose–ethidium bromide gel. Meanwhile, the Ri-plasmid of *A. rhizogenes* was used as a positive control (Wang *et al.*, 2006).

## Growth kinetics and biomass measurements

The growth rate was measured as hairy root growth kinetics at the first of the experiment. The experimental conditions were as follows. Hairy root lines with 2.4 g of root mass (on fresh weight basis) were transferred to a 250-ml Erlenmeyer flask containing 50 ml of hormone-free MS liquid medium at 25 °C for period of 4 weeks. Flasks were harvested with three replications each week (up to day 28 after inoculation). The growth of hairy root lines was measured every week. Hairy roots were dried by blotting paper to remove all external moisture and weighed to determine the final fresh weight (FW) before being frozen for lignan extraction.

## Extraction of lignans

Powdered hairy roots (2 g) were extracted by sonication in methanol (80 %) during 1 h. Dichloromethane and water (1:1 v/v) were added to obtain a partition of compounds between two layers. The dichloromethane fractions were collected, dried and dissolved in 500 µl of HPLC grade methanol and then injected into the HPLC (Van Uden *et al.*, 1989) (Philips, UV/Vis detector; Pu 41110). The

presence of PTOX and MPTOX in the hairy roots was determined by ESI/MS in positive ion mode and 0.05 % formic acid was used.

## MPTOX purification and confirmation

Analytical separation was carried out to confirm MPTOX and determine its enantiomer. Separations were performed on a C18-S50DS3 (250 × 4.6 mm) RP column with a flow rate of 1.0 ml min<sup>-1</sup>. A fraction containing MPTOX was collected, lyophilized and dissolved in DMSO and subjected to 500 MHz NMR spectroscopy. A NMR spectrum was recorded on Bruker DRX500 Avance running XWIN-NMR 2.6 software using 500 µl samples. The chemical shifts (delta) are reported in parts per million (ppm). Also, FT-RI was carried out with a Bruker tensor 27 spectrometer.

The CD spectrum was obtained with a JASCO J-715 spectropolarimeter using cuvettes with optical lengths of 1 cm and detection at 250–350 nm. The final data were obtained by subtracting the buffer contribution (DMSO) from the original spectrum.

Optical rotation ( $[\alpha]_D$  values) of MPTOX was measured on a Perkin Elmer 241 automatic polarimeter (Überlingen, Germany) at 20 °C. The specific rotation ( $[\alpha]_D$ ) at the sodium D line ( $\lambda$  589 Å) was calculated as follows:

$$[\alpha]_D = (100 \times \alpha) / (l \times c), \alpha = \text{observed angle of rotation, } l = \text{length of the path of light through the solution in decimeters (dm), } c = \text{concentration of the solution in g/100 ml.}$$

## Analysis of lignans

PTOX and MPTOX quantification was analyzed using HPLC as described previously by Yousefzadi *et al.* (2010b). The stationary phase was a C18-ODS3, 5 µm (250–4.6 mm) column. The elution solvent was composed of water and acetonitrile with a gradient system. This system with acetonitrile (A%) and water (B%) was used as follows: (0 min, flow rate 0.8): A (40), B (33); (10 min, flow rate 1): A (67), B (33); (17 min, flow rate 1): A (40), B


**Table 1** Sequences of used primers

Genes	Sequences	Amplicon size (bp)
<i>rolC</i>	Forward primer 5'-TAACATGGCTGAAGACGACC-3'	534
	Reverse primer 5'-AAACTTGCACTCGCCATGCC-3'	
<i>Ags</i>	Forward primer 5'-GGCGTGAGCACCTCATATCCG-3'	347
	Reverse primer 5'-TTCGAAGCCTTTGCCTGCAAA-3'	
<i>virD</i>	Forward primer 5'-ATGTCGCAAGGCAGTAAGCCC-3'	438
	Reverse primer 5'-GGAGTCTTTCAGCATGGAGCAA-3'	

(60); (21 min, flow rate 0.8): A (40), B (60). A spectrophotometric UV detection was used (290 nm). The calibration was performed with standard PTOX ( $RT = 9.5 \pm 0.1$  min) and MPTOX ( $RT = 11.5 \pm 0.1$  min) (Fig. 1). Every sample was assayed with three replications. The lignan production was determined using a calibration curve compared with standards and a co-chromatogram of the standards and samples.


#### Statistical analysis

All data were analyzed using MSTAT-C software. Duncan's multiple range tests were used to show statistical differences between treatment methods and controls.  $P \leq 0.05$  or  $\leq 0.01$  was considered significantly different.



**Fig. 1** 6-Methoxy podophyllotoxin (a) and podophyllotoxin (b) calibration curve using HPLC. Each value is the mean of three replications


**Fig. 3** Electrophoretogram showing the presence of *rolC* (a), *Ags* (b) genes and absence of *VirD* (c) gene (534, 347, 483 bp bands, respectively) in transformed hairy roots; M molecular weight marker, PC positive control, R1 and R2: transformed root lines LBA9402 and C58C1, respectively, R<sup>-</sup>: the non-transformed root



## Results and Discussion

### *L. album* hairy root induction

Although all four *Agrobacterium* strains applied in this work are among the most virulent in common usage (Lin *et al.*, 2003), despite several attempts using different bacterial strains and infection sites, no hairy roots were obtained using the leaf disk infection technique. Infection by syringe needle was more successful, with strains LBA9402 and C58C1 (pRiA4) showing their rooting ability, the former more readily. LBA9402 hairy roots of *L. album* were induced after 4 weeks of infection and had a higher transformed percentage (20 %), while C58C1 (pRiA4) hairy roots were induced after 5 weeks with a lower percentage of transformation (10 %) (data are not shown). Some transgenic roots turned brown and aged considerably faster than others. These brown roots were discarded and only those showing a good growth capacity were maintained for further characterization. Although various explant sources can be used to produce hairy roots *in vitro*, in some cases specific tissues perform better than others (Mishra *et al.*, 2011). Regarding *L. album*, young stems produced more primary cultures than other tissues.




**Fig. 2** Growth pattern of hairy roots induced by *A. rhizogenes* strains LBA9402 and *A. tumefaciens* C58C1 (pRiA4). Data represent average values from four separate experiments  $\pm$  SD

Efficiency of transformation differed among the *Agrobacterium* strains, as reported by other authors (Giri *et al.*, 2001; Mishra *et al.*, 2011).


A time course experiment (up to 4 weeks at defined time intervals) was performed to monitor root growth and biomass production and also to estimate lignan content. The growth kinetics of *L. album* hairy roots showed three different phases: the lag phase (first week) when the specific growth rate is essentially slow, the exponential growth

phase (second and third weeks) when the specific growth rate is constant and the stationary phase (fourth week) with the emergence of at least one limiting parameter. The highest hairy root growth was recorded with strain LBA9402 (12.455 g FW) after 4 weeks of culture (Fig. 2). Hairy roots induced by C58C1 (pRiA4) showed comparatively slow growth (11.083 g FW). However, transformed hairy roots induced by both strains exhibited negative geotropism. The transformed roots with *A. rhizogenes* have

**Fig. 4** CD spectrum of MPTOX. Detected solvents: methanol of chromatographic grade, detected temperature: 25 °C. UV detected wavelength at 250–350 nm



**Fig. 5** Selected molecular ion (445 m/z) in TIC (up) and mass spectrum of 6-MPTOX (down)



an altered phenotype, showing profuse lateral growth, lack of geotropism and fast growth in the culture (Mishra *et al.*, 2011).

#### Confirmation of transformation

In order to assess the genetic status of the hairy roots, we used a PCR-based analysis that targeted *rolC*, *Ags* (opine synthase) and *virD* genes. The *rolC* and *Ags* genes are diagnostic for T-DNA integration into the host genome. The *virD* gene, located outside the T-DNA, is diagnostic for the presence of any remaining *Agrobacterium* in the root tissue. PCR analysis revealed the presence of the *rolC* and *Ags* genes in hairy roots, indicating that the T-DNA was retained in hairy roots (Fig. 3). Confirmation of contamination was also done by PCR analysis using *virD* (483 bp) gene-specific primers. It was not found in the hairy root lines, thus indicating the absence of contamination with *Agrobacterium* strains LBA9402 and C58C1. Also, *rolC*, *Ags* and *virD* genes were not detected in the non-transformed root.

#### Confirmation of MPTOX and PTOX

##### Identification of MPTOX in transformed root cultures

Lignan (dimers of monolignols) formation and lignin (polymers of monolignols) biosynthesis are catalyzed by

different enzymes, and a consequence is that natural lignans are normally enantiomerically pure because they arise from stereochemically controlled coupling (Dewik, 2002). For example, the lignan pinoresinol, and its derived matairesinol, was found only as their (-)-isomers in suspension-cultured *Pinus taeda* (Eberhardt *et al.* 1993). Data acquired using a polarimeter (optical rotation =  $[\alpha]_D - 60^\circ$ ) and CD spectrum confirmed the presence of the (-)-isomer of MPTOX (Fig. 4).


Moreover, this less hydrophilic compound was identified by NMR and IR spectroscopy. Significant  $^1\text{H-NMR}$  data are shown in the following order: chemical shift ( $\delta$ ) expressed in ppm, multiplicity (s, singlet; d, doublet; m, multiplet and ABq, AB quartet).

$^1\text{H-NMR}$  (500 MHz,  $(\text{CD}_3)_2\text{SO}$ ):  $\delta = 6.39$  (2H, s), 6.29 (1H, s), 5.97 (2H, d), 5.04 (1H, d), 4.93 (1H, m), 4.45 (2 H, ABq), 4.08 (1H, dd), 3.95 (3H, s,  $\text{OCH}_3$ ), 3.62 (6H, s,  $(\text{OCH}_3)_2$ ), 3.6 (3H, s,  $\text{OCH}_3$ ), 3.0 (1H, dd). Melting point: 80–83. IR: 3449.20, 2924.41, 2862.45, 1775.34, 1634.25, 1465.59, 1247.14, 1124.43, 1050.77.

##### MS analyses of MPTOX

For identification, liquid chromatography–electrospray–mass spectrometry/mass spectrometry (LC–ESI–MS/MS) analysis was carried out. Several reports have been cited in the literature demonstrating the applicability of MS and MS/MS coupled with liquid chromatography in the

**Fig. 6** The fragmentation of MPTOX (according to Li *et al.* 2007)



analyses of PTOX and its derivatives (Li *et al.*, 2007; Lin *et al.*, 2008). In this work,  $[M + H]^+$  and  $[M + Na]^+$  ions of 6-MPTOX were determined. According to Fig. 5, the observed mass at 467 Da corresponds to the  $[M + Na]^+$  ion of 6-MPTOX and the observed mass of 445 Da is related to the  $[M + H]^+$  ion of 6-MPTOX. The other peaks at mass of 427 and 343 Da correspond to fragments illustrated in Fig. 6.

#### Confirmation of PTOX

The presence of PTOX in the hairy roots was verified by ESI/MS in positive ion mode, as previously described (Yousefzadi *et al.*, 2010a).

#### Quantification of lignans

Among the examined samples, hairy roots induced by strain LBA9402 were found to be the best in terms of biomass as well as lignan production. The maximum concentrations of PTOX ( $105 \mu\text{g g}^{-1}$  DW), and 6-MPTOX ( $48 \text{ mg g}^{-1}$  DW) were recorded in hairy roots induced by strain LBA9402 (Fig. 7). In the roots of mother plants which were collected from its natural habitat, concentrations of PTOX and 6-MPTOX were 14.11 and  $1.13 \text{ mg g}^{-1}$  DW, respectively. When compared to the roots of


mother plant, the hairy root cultures were found to contain higher concentrations of lignans. This confirms that the insertion sites of Ri-T-DNA interfere with the secondary metabolite biosynthetic pathway of lignans. Similar results have been reported by several groups (Mishra *et al.*, 2011; Verma *et al.*, 2007).

#### Conclusion

In general, the amount of anti-cancer lignans (PTOX and MPTOX) in plants is considerably less than in transformed hairy roots. In addition, the hairy roots exhibited more rapid growth than plants. This study may open possibilities for bioreactor-based production of the pharmaceutically important metabolites PTOX and 6-MPTOX.

#### References

- Bais HP, Sudha G, George J, Ravishankar GA (2001) Influence of exogenous hormones on growth and secondary metabolite production in hairy root cultures of *Cichorium intybus* L. Cv. Lucknow local. *In Vitro Cell Dev Biol Plant* 37:293–299
- Botta B, Delle Monache G, Misiti D, Vitali A, Zappia G (2001) Aryltetralin lignans: chemistry, pharmacology and biotransformation. *Curr Med Chem* 8:1363–1381
- Chattopadhyay S, Srivastava AK, Bisaria VS (2002) Optimization of culture parameters for production of podophyllotoxin in suspension culture of *Podophyllum hexandrum*. *Appl Biochem Biotechnol* 102:381–393
- Chattopadhyay S, Bisaria VS, Srivastava AK (2003a) Enhanced production of podophyllotoxin by *Podophyllum hexandrum* using in situ cell retention bioreactor. *Biotechnol Prog* 19:1026–1028
- Chattopadhyay S, Bisaria VS, Bhojwani SS, Srivastava AK (2003b) Enhanced production of podophyllotoxin by fed-batch cultivation of *Podophyllum hexandrum*. *Can J Chem Eng* 81:1–8
- Chattopadhyay S, Farkya S, Srivastava AK, Bisaria VS (2003c) Bioprocess considerations for production of secondary metabolites by plant cell suspension cultures. *Biotechnol Bioprocess Eng* 7:138–149
- Dewik PM (2002) Medicinal natural products, a biosynthetic approach. Wiley, Chichester, pp 121–136
- Eberhardt TL, Bernards MA, He L, Davin LB, Wooten JB, Lewis NG (1993) Lignification in cell suspension cultures of *Pinus taeda*: in situ characterization of a Gymnosperm lignin. *J Biol Chem* 268:21088–21096
- Farkya S, Bisaria VS (2008) Exogenous hormones affecting morphology and biosynthetic potential of hairy root line (LYR2i) of *Linum album*. *J Biosci Bioeng* 105:140–146
- Farkya S, Bisaria VS, Srivastava AK (2004) Biotechnological aspects of the production of the anticancer drug podophyllotoxin. *Appl Microbiol Biotechnol* 65:504–519
- Giri A, Narasu ML (2000) Transgenic hairy roots: recent trends and applications. *Biotechnol Adv* 18:1–22
- Giri A, Giri CC, Dhingra V, Narasu ML (2001) Enhanced podophyllotoxin production from *Agrobacterium rhizogenes* transformed cultures of *Podophyllum hexandrum*. *Nat Prod Lett* 15:229–235



**Fig. 7** PTOX and MPTOX production in a 4-week period. PTOX: podophyllotoxin. MPTOX: 6-methoxy podophyllotoxin. Data represent average values from three separate experiments  $\pm$  SD

- Gordaliza M, Garc PA, del Corral M, Castro MA, Gomez-Zurita MA (2004) Podophyllotoxin: distribution, sources, applications and new cytotoxic derivatives. *Toxicol* 44:441–459
- Khan S, Irfan Qureshi M, Kamaluddin Alam T, Abdin MZ (2007) Protocol for isolation of genomic DNA from dry and fresh roots of medicinal plants suitable for RAPD and restriction digestion. *Afr J Biotechnol* 6:175–178
- Kuhlmann S, Kranz K, Lücking B, Alfermann AW, Petersen M (2002) Aspects of cytotoxic lignan biosynthesis in suspension cultures of *Linum nodiflorum*. *Phytochem Rev* 1:37–43
- Li Q, Yan G, Ge T (2007) A fragmentation study of podophyllotoxin and its 40-demethyl-4b-substituted derivatives by electrospray ionization ion-trap time-of-flight tandem mass spectrometry. *Rapid Commun Mass Spectrom* 21:2843–2852
- Lin H, Kwok KH, Doran PM (2003) Development of *Linum flavum* hairy root cultures for production of Coniferin. *Biotechnol Lett* 25:521–525
- Lin MC, Lin JH, Chen SK, Cheng YW, Cheng HW (2008) Simultaneous determination of podophyllotoxin, quercetin and kaempferol in podophyllin by liquid chromatography tandem mass Spectrometry. *J Food Drug Anal* 16:29–40
- Mishra J, Bhandari H, Singh M, Rawat S, Agnihotri RK, Mishra S, Purohit S (2011) Hairy root culture of *Picrorhiza kurroa* Royle ex Benth.: a promising approach for the production of picrotin and picrotoxinin. *Acta Physiol Plant* 33(5):1841–1846
- Murashige T, Skoog F (1962) A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Plant Physiol* 15:473–497
- Nikolay V, Ionkova I (2005) Lignan production by cell cultures of *Linum setaceum* and *Linum campanulatum*. *Pharm Biol* 43:509–511
- Park S, Facchini P (2000) *Agrobacterium rhizogenes*-mediated transformation of opium poppy, *Papaver somniferum* L., and California poppy, *Escholizia californica* cham., root cultures. *J Exp Bot* 51:1005–1016
- Ramachandra RS, Ravishankar GA (2002) Plant cell cultures: chemical factories of secondary metabolites. *Biotechnol Adv* 20:101–153
- Smollny T, Wichers H, Kalenberg S, Shahsavari A, Petersen M, Alfermann WA (1998) Accumulation of podophyllotoxin and related lignans in cell suspension cultures of *Linum album*. *Phytochemistry* 48:975–979
- Van Uden W, Pras N, Visser JF, Malingre TM (1989) Detection and identification of podophyllotoxin produced by cell cultures derived from *Podophyllum hexandrum* Royle. *Plant Cell Rep* 8:165–168
- Verma PC, Rahman L, Negi AS, Jain DC, Khanuja SPS, Banerjee S (2007) *Agrobacterium rhizogenes* mediated transformation of *Picrorhiza kurroa* Royle ex Benth.: establishment and selection of superior hairy root clone. *Plant Biotechnol Rep* 1:169–174
- Wang B, Zhang G, Zhu L, Chen L, Zhang Y (2006) Genetic transformation of *Echinacea purpurea* with *Agrobacterium rhizogenes* and bioactive ingredient analysis in transformed cultures. *Colloid Surface B* 53:101–104
- Yousefzadi M, Sharifi M, Behmanesh M, Ghasempour A, Moyano E, Palazon J (2010a) Salicylic acid improves podophyllotoxin production in cell cultures of *Linum album* by increasing the expression of genes related with its biosynthesis. *Biotechnol Lett* 32:1739–1743
- Yousefzadi M, Sharifi M, Ahmadian Chashmi N, Behmanesh M, Ghasempour A (2010b) Optimization of podophyllotoxin extraction method from *Linum album* cell cultures. *Pharm Biol* 48:1421–1425
- Zhong JJ (2002) Plant cell culture for production of paclitaxel and other taxanes. *J Biosci Bioeng* 94:591–599