

coloproctology 2019 · 41:110
<https://doi.org/10.1007/s00053-019-0355-z>
 Published online: 4 April 2019
 © Springer Medizin Verlag GmbH, ein Teil von
 Springer Nature 2019



A. P. Wysocki^{1,2}

¹ Department of Surgery, Logan Hospital, Meadowbrook, Australia

² Griffith Health Centre—G40, Gold Coast Campus, Griffith University Medical School, Gold Coast, Australia

Erratum to: Primary open treatment of sacrococcygeal pilonidal disease

Erratum to:
coloproctology 2019
<https://doi.org/10.1007/s00053-019-0346-0>

In **Fig. 2** the color correlation was not clearly given. You can see the corrected figure below.

Corresponding address

A. P. Wysocki, MBBS, FRACS
 Department of Surgery, Logan Hospital
 Corner Meadowbrook and Loganlea Roads,
 4133 Meadowbrook, Queensland, Australia
 arek_p@ecn.net.au

References

12. Stauffer VK et al (2018) Common surgical procedures in pilonidal sinus disease: a meta-analysis, merged data analysis, and comprehensive study on recurrence. *Nature Sci Rep* 8(1):1–27




Fig. 2 ▲ Procedure-specific recurrence rates. (Re-drawn with data from Reference [1])

The online version of the original article can be found under <https://doi.org/10.1007/s00053-019-0346-0>.