

Strahlenther Onkol 2012 · 188:696–701
 DOI 10.1007/s00066-012-0122-3
 Received: 22 January 2012
 Accepted: 26 March 2012
 Published online: 23. Juni 2012
 © Springer-Verlag 2012

R. Wiggendaad¹ · A. Verbeek-de Kanter¹ · M. Mast¹ · R. Molenaar² · H.B. Kal³ · G. Lycklama à Nijeholt⁴ · C. Vecht⁵ · H. Struikmans^{1,6}

¹ Radiotherapy Centre West, The Hague

² Department of Neurology, Diaconessenhuis, Leiden

³ Maassluis

⁴ Department of Radiology, Medical Centre Haaglanden, The Hague

⁵ Department of Neurology, Medical Centre Haaglanden, The Hague

⁶ Department of Radiotherapy, Leiden University Medical Centre, Leiden

Local progression and pseudo progression after single fraction or fractionated stereotactic radiotherapy for large brain metastases

A single centre study

Stereotactic radiotherapy (SRT) is an established treatment modality for patients with brain metastases [1]. Local control of the metastases is the aim of treatment as progressive tumor growth may lead to new neurologic symptoms [2]. The dose that can be safely administered depends upon the size of the metastasis [3]. In Radiation Therapy Oncology Group (RTOG) study 90–05 the maximum tolerated single fraction dose for metastases with a diameter >3 cm was found to be 15 Gy, as a higher dose of 18 Gy was associated with an unacceptably high rate of grade 3–5 neurotoxicity. Progression after radiotherapy may be caused by proliferation of tumor cells but may also be a manifestation of radiation toxicity (pseudo progression). The distinction between these two types of progression is difficult to make on magnetic resonance imaging (MRI), but perfusion MRI may be helpful in this respect [4].

The 1-year local control of metastases >3 cm is reported to be 37–62% after 15 Gy [3, 5, 6]. With FSRT 1-year local control rates >70% were reported [7, 8]. However, a comparative study of SRT and FSRT for large brain metastases has

not yet been published. Furthermore, only scarce data are available about the rates of pseudo progression after SRT or FSRT. We observed a disappointing 12-month local control rate of 37% after a single-fraction dose of 15 Gy in our patients with large brain metastases [6]. In an attempt to improve local control rates we embarked on an FSRT protocol in September 2007, treating this category of patients with 3 fractions of 8 Gy. The purpose of the present study is to compare the local control rates as well as the rates of pseudo progression between these two treatment protocols.

Materials and methods

Patients

Two patient cohorts received SRT for brain metastases in two consecutive periods. In both cohorts we included patients with metastases with a planning target volume (PTV) of >13 cm³ or metastases in or close to the brainstem. The prescribed dose was 15 Gy between June 2004 and January 2007 (group A) and 24 Gy in 3 fractions of 8 Gy (in 8 days) between

September 2007 and September 2009 (group B). To be able to study the effect of SRT dose on local control, we excluded the patients who had SRT as a boost after whole brain irradiation (WBI). Karnofsky performance scores (KPS) were determined prospectively and the RTOG recursive partitioning analysis (RPA) scores were determined retrospectively in all patients.

Treatment

Patients had a CT scan with 2-mm slice thickness while fixed in a relocatable stereotactic head frame (Brainlab AG Feldkirchen, Germany) [9, 10]. All patients also had an MRI planning scan (T1-weighted 3D MPRAGE after gadolinium administration; voxel size 1.1×1.1×1.3 mm³). Co-registration of CT and MRI, contouring and treatment planning were done on Brainscan 5.31 or iPlan 4.0 (Brainlab AG Feldkirchen, Germany). The gross tumor volume (GTV) was defined as the volume of the contrast-enhancing tumor as visualized on the MRI scan. The PTV was created by 3D expansion of the GTV with 2 mm. All patients were treated on

Tab. 1 Patient and treatment characteristics in both groups			
	Group A^a n (%)	Group B^b n (%)	Group A vs Group B
Patients (n)	41	51	
Sex			
Male	15 (37)	19 (37)	p=0.9
Female	26 (63)	32 (63)	
Age (years)			
<65	22 (54)	29 (57)	p=0.8
≥65	19 (46)	22 (43)	
KPS			
<90	21 (51)	29 (57)	p=0.6
≥90	20 (49)	22 (43)	
RPA class			
1	7 (17)	8 (16)	p=0.7
2	30 (73)	43 (84)	
3	4 (10)	0	
Primary tumor			
Lung	19 (46)	24 (47)	p=0.2
Breast	10 (24)	8 (16)	
Melanoma	2 (6)	9 (18)	
Other	10 (24)	10 (19)	
Brain metastases (n)			
1	20 (49)	28 (55)	p=0.6
>1	21 (51)	23 (45)	
Treatment			
No WBI	29 (70)	32 (63)	p=0.4
SRT as salvage after WBI	6 (15)	13 (25)	
WBI as salvage after SRT	6 (15)	6 (12)	

^a Group A received 1 fraction of 15 Gy ^b Group B received 3 fractions of 8 Gy KPS Karnofsky performance score, RPA recursive partitioning analysis, WBI whole brain irradiation, SRT stereotactic radiotherapy.

Tab. 2 Tumor characteristics in both groups			
	Group A^a n (%)	Group B^b n (%)	Group A vs Group B
Metastases (n)	46	65	
Primary tumor			
Lung	22 (48)	26 (40)	p=0.3
Breast	11 (24)	13 (20)	
Melanoma	2 (4)	10 (15)	
Other	11 (24)	16 (25)	
PTV			
<13 cm ³	11 (24)	18 (28)	p=0.4
13–20 cm ³	17 (37)	16 (25)	
>20 cm ³	18 (39)	31 (47)	
Treatment			
No WBI	33 (72)	42 (65)	p=0.6
SRT as salvage after WBI	7 (15)	15 (23)	
WBI as salvage after SRT	6 (13)	8 (12)	

^a Group A received 1 fraction of 15 Gy ^b Group B received 3 fractions of 8 Gy PTV planning target volume, WBI whole brain irradiation, SRT stereotactic radiotherapy.

the Novalis, a dedicated linear accelerator (Brainlab AG Feldkirchen, Germany). Dynamic conformal arc was used as treatment technique for all metastases. Doses were prescribed to the 80% isodose. We allowed a maximum dose of 8 Gy (SRT) and 15 Gy (FSRT) to the optic system and a maximum dose of 15 Gy (SRT) and 24 Gy (FSRT) to the brainstem.

Follow-up

All patients were followed-up at 3-month intervals at the outpatient clinic as long as their condition allowed them to come. These follow-up visits were combined with MRI scans at 1.5 T (Siemens, Erlangen, Germany). The imaging protocol consisted of T1-weighted images without and with gadolinium, T2 images and diffusion-weighted imaging. MR perfusion imaging, using a SE-EPI sequence, was performed when increase or recurrence of gadolinium enhancement was observed in lesions. Analysis of the MR perfusion data was performed by calculating the relative cerebral blood volume (r-CBV) maps and by comparing the r-CBV maps with the post-gadolinium T1-weighted images. R-CBV maps were considered to be suggestive for viable tumor tissue when r-CBV in the enhancing part of the tumor was equal to or higher than cerebral gray matter (based on visual assessment by a neuroradiologist).

Telephonic follow-up was done if patients were not able to visit the hospital anymore; however, information thus acquired was only used for survival analysis. Local control was calculated from the first day of (F)SRT. All MRI scans were reviewed and tumors were measured in three dimensions. Response to treatment was classified according to the Macdonald criteria [11]. The date of the first MRI showing any local progression was used as the date of progression. Pseudo progression was diagnosed when perfusion MRI showed no signs of viable tissue [4]. Tumor progression was defined as tumor proliferation not caused by pseudo progression, i.e., with MR perfusion imaging characteristics compatible with viable tissue. Tumor progression could be preceded by pseudoprogression. Endpoints were

any local progression, tumor progression and pseudo progression.

Analyses

Statistical analysis was performed using SPSS 16.0 (SPSS Inc., Chicago, IL, USA). The Pearson χ^2 test was used to analyze whether the characteristics of both cohorts were equally divided. Overall survival and local progression-free survival (LPFS) curves were calculated using the Kaplan–Meier method. For the calculation of the actuarial freedom from any progression only the first progression of a metastasis was used as an event. The log-rank test was used for the univariate analyses.

Results

Patients, metastases and treatment

Group A consisted of 41 patients with 46 brain metastases and group B consisted of 51 patients with 65 brain metastases. The median follow-up of all patients was 5.3 months and the minimum follow-up of patients still alive was 22 months. Patient and treatment characteristics were equally divided between both groups (■ **Tab. 1**). The tumor characteristics of both cohorts are shown in ■ **Tab. 2**.

Local progression-free survival (LPFS)

The actuarial rates of freedom from any local progression in group A and B are shown in ■ **Fig. 1**. The 6- and 12-month local control rates were 89% and 54% with 1 fraction of 15 Gy and 84% and 61% with 3 fractions of 8 Gy. LPFS rates were not significantly different between the two groups ($p=0.93$). In two brain metastases tumor progression was preceded by pseudo progression. Only one resection of a progressive lesion was performed (in a patient from the SRT cohort). The histological diagnosis was radiation necrosis.

The actuarial freedom from tumor progression in groups A and B are shown in ■ **Fig. 2**. Pseudo progression was not considered an event in this figure. The 6- and 12-month rates of freedom from tumor progression were 89% and 67%

Strahlenther Onkol 2012 · 188:696–701 DOI 10.1007/s00066-012-0122-3
© Springer-Verlag 2012

R. Wiggeraad · A. Verbeek-de Kanter · M. Mast · R. Molenaar · H.B. Kal · G. Lycklama à Nijeholt · C. Vecht · H. Struikmans

Local progression and pseudo progression after single fraction or fractionated stereotactic radiotherapy for large brain metastases. A single centre study

Abstract

Purpose. The 1-year local control rates after single-fraction stereotactic radiotherapy (SRT) for brain metastases >3 cm diameter are less than 70%, but with fractionated SRT (FSRT) higher local control rates have been reported. The purpose of this study was to compare our treatment results with SRT and FSRT for large brain metastases.

Materials and methods. In two consecutive periods, 41 patients with 46 brain metastases received SRT with 1 fraction of 15 Gy, while 51 patients with 65 brain metastases received FSRT with 3 fractions of 8 Gy. We included patients with brain metastases with a planning target volume of >13 cm³ or metastases in the brainstem.

Results. The minimum follow-up of patients still alive was 22 months. Comparing 1 frac-

tion of 15 Gy with 3 fractions of 8 Gy, the 1-year rates of freedom from any local progression (54% and 61%, $p=0.93$) and pseudo progression (85% and 75%, $p=0.25$) were not significantly different. Overall survival rates were also not different.

Conclusion. The 1-year local progression and pseudo progression rates after 1 fraction of 15 Gy or 3 fractions of 8 Gy for large brain metastases and metastases in the brainstem are similar. For better local control rates, FSRT schemes with a higher biological equivalent dose may be necessary.

Keywords

Brain metastasis · Stereotactic radiotherapy · Fractionation · Progression · Pseudo progression

Lokale Progression und Pseudoprogression bei großen Hirnmetastasen nach stereotaktischer Radiotherapie mit einer oder mehreren Fraktionen. Eine monoinstitutionelle Studie

Zusammenfassung

Hintergrund. Die einjährige lokale Tumorkontrollrate von Hirnmetastasen mit einem Durchmesser von mind. 3 cm beträgt nach stereotaktischer Bestrahlung mit einer Fraktion (SRT) weniger als 70%. Höhere lokale Tumorkontrollraten sind nach fraktionierter stereotaktischer Bestrahlung (FSRT) beschrieben worden. In dieser Studie werden die Behandlungsergebnisse der SRT und der FSRT für größere Hirnmetastasen verglichen.

Material und Methoden. In 2 aufeinanderfolgenden Perioden wurden 41 Patienten mit 46 Hirnmetastasen mit 1-mal 15 Gy SRT und 51 Patienten mit 65 Hirnmetastasen mit 3-mal 8 Gy FSRT behandelt. Alle inkludierten Patienten hatten einen PTV von mindestens 13 cm³ oder Metastasen im Hirnstamm-bereich.

Ergebnisse. Das minimale Follow-up von überlebenden Patienten betrug 22 Monate.

Beim Vergleich der Resultate von 1-mal 15 Gy mit 3-mal 8 Gy ergaben sich nach einem Jahr keine signifikanten Unterschiede bei den Raten ohne lokale Progression (54% bzw. 61%; $p=0,93$), der Pseudoprogression (85% bzw. 75%; $p=0,25$) und im Gesamtüberleben.

Schlussfolgerung. Die lokale Progression und Pseudoprogression von großen Hirnmetastasen oder Hirnstammmetastasen sind ein Jahr nach Behandlung mit 1-mal 15 Gy oder 3-mal 8 Gy vergleichbar. Für eine bessere lokale Kontrolle sind vielleicht FSRT-Regime mit höherer biologischer Äquivalentdosis notwendig.

Schlüsselwörter

Hirnmetastase · Stereotaktische Radiotherapie · Fraktionierung · Progression · Pseudoprogression

with 1 fraction of 15 Gy and 92% and 75% with 3 fractions of 8 Gy. There was no significant difference between both groups ($p=0.27$).

The actuarial rates of freedom from pseudo progression in group A and B are

shown in ■ **Fig. 3**. The 6- and 12-month rates of freedom from pseudo progression were 93% and 85% with 1 fraction of 15 Gy and 91% and 75% with 3 fractions of 8 Gy. There was no significant difference between the two groups ($p=0.25$).




Fig. 1 ◀ The actuarial rate of freedom from all tumor progression in both groups of metastases. Tumor progression and pseudo progression were considered as events. $p=0.93$




Fig. 2 ◀ The actuarial rate of freedom from tumor progression in both groups of metastases. Only tumor progressions, but not pseudo progressions were considered as events. $p=0.27$

In a univariate analysis dose (15 or 24 Gy), KPS, PTV, previous WBI only and all WBI (previous or later) were correlated with the occurrence of all tumor progressions, pseudo progression and tumor progression (■ Tab. 3). We found no relation between any type of progression and dose, KPS or tumor volume. However, after previous WBI a significantly higher rate of pseudo progression was found

($p=0.02$). Moreover, a higher rate of tumor progression was found with previous or later WBI. As this was the only significant relation, a multivariate analysis was not performed.

Survival

The median survival of all patients was 5.3 months. The 6- and 12-month over-

all survival rates were 41% and 23%, respectively. In the univariate analysis the only prognostic factor for survival was KPS ($p=0.02$). We found no difference in survival rates between group A and B ($p=0.58$).

Discussion

This is a retrospective comparison of SRT (1 fraction of 15 Gy) and FSRT (3 fractions of 8 Gy) used in two consecutive cohorts of patients with large-sized brain metastases. Actuarial survival rates and rates of freedom from progression or pseudo progression were found not to differ significantly between 1 fraction of 15 Gy and 3 fractions of 8 Gy. Therefore, FSRT with 3 fractions of 8 Gy does not seem to be an improvement over 1 fraction of 15 Gy for large brain metastases and metastases in the brainstem.

Surgery may be the treatment of choice for large brain metastases, if feasible. SRT however is also an attractive option for patients with large metastases, although symptomatic improvement and local control are not optimal after single fraction treatment [6, 12, 13, 14]. There is no agreement on the optimal SRT dose for these tumors. Recently we performed a systematic literature search to summarize the evidence about the relation between SRT dose and local control [13]. We found that 12-month local control after SRT was highly dependent upon dose and was high after >21 Gy, but low after <15 Gy [13]. However, it is known from RTOG 90-05 that unacceptably high neurotoxicity rates are found after treating larger recurrent brain metastases (diameter >3 cm) with single doses >15 Gy [3]. Higher local control rates were reported after FSRT in larger metastases with acceptable rates of radiation toxicity (■ Tab. 4, [8, 15, 16, 17, 18]). Therefore, to improve the results of SRT of large brain metastases, FSRT is a logical step, enabling a higher tumor dose with a lower risk of neurotoxicity.

We decided to treat these patients with 3 fractions of 8 Gy, after we had observed the disappointing efficacy of 1 fraction 15 Gy. The rationale for this new scheme was the better biologically effective dose (BED) “profile” of the fractionated scheme [6]. The BED model describes the re-




Fig. 3 ◀ The actuarial rate of freedom from pseudo progression in both groups of metastases. $p=0.25$

sponses to ionizing radiation well at doses up to about 18 Gy [19, 20, 21]. The BED_2 values (for normal tissue, $\alpha/\beta=2$ Gy) for 1 fraction of 15 Gy and 3 fractions of 8 Gy are 127.5 Gy and 120 Gy respectively and the BED_{12} values (for tumor, $\alpha/\beta=12$ Gy) 33.8 Gy and 40 Gy respectively. Based on the BED model we initially expected an improved local control with less late toxicity. In hindsight these expected differences are probably too small to detect in the relatively small numbers studied here. To improve local control for brain metastases with a PTV >13 cm³ it would be logical to use FSRT with a higher BED_{12} , keeping in mind that the rate of adverse treatment effects may also increase. In our department we decided to change the protocol for these large metastases to 3 fractions of 8.5 Gy, following the conclusions from our literature search [13].

As this is not a randomized comparison, conclusions from this study have to be viewed with caution. Local control was found to be independent upon PTV and all other factors that may influence local control are well balanced between the two cohorts. Therefore we think that it is justified to conclude that local control rates are similar with both dose schemes, with all well-known restrictions of a retrospective study.

An enlargement of the treated volume after radiotherapy may be caused by an increased proliferation rate of tumor cells but may also be a manifestation of radiation toxicity (pseudo progression). We prefer to use the term pseudo progression instead of radiation necrosis like it is used in gliomas, where real progression can also be preceded by pseudo progression [22]. The histology of this radiation effect usually is a chronic inflammatory reaction of brain tissue combined with necrosis of normal brain tissue and tumor tissue [23]. The distinction between real tumor progression and pseudo progression is difficult to make using standard morphologic MR imaging, but modern MR imaging techniques, especially perfusion MRI, may be helpful to differentiate viable tumor tissue from tissue with radiation effects [4, 24]. In our patients a substantial proportion of all progressions were diagnosed as pseudo progression. As the rate of pseudo progression was not significantly different between group A and B (15% versus 25% at 1 year), whereas the metastases in group B were slightly larger than those in group A, 3 fractions of 8 Gy is certainly feasible for the larger metastases.

Not much is known about ways to reduce the rate of pseudo progression. A relation has been reported between the rate

of radiation necrosis and the V_{12} (volume of tissue that received a single dose of ≥ 12 Gy) [25, 26]. A lower BED to normal brain tissue and at the same time a higher BED to tumor tissue would be needed to reduce the rate of pseudo progression without compromising local control. To this end FSRT may be used, but more research is needed to find the optimal scheme.

An interesting finding in our study is the higher rate of pseudo progression in patients with prior WBI. This would imply that it would be safe to give higher biological doses than advised based on RTOG 90–05 if no prior WBI has been given. Until now, only a few and somewhat conflicting data are available on this issue. Chao et al. [5] retreated 111 patients with SRT for recurrence after WBI and found only two cases of radiation necrosis. Yang et al. [14] treated 70 patients with metastases >3 cm diameter with SRT, 33 of whom had previous WBI. After 2 months patients with previous WBI had slightly worse edema response and symptom relief than those without. Fourteen of 29 patients with imaging assessment >6 months after SRT had adverse radiation effects (48%), but the authors did not mention a relation with previous WBI.

We conclude that local control rates with 1 fraction of 15 Gy or 3 fractions of 8 Gy for large brain metastases are similar. Large brain metastases can be safely treated with a hypofractionated scheme, but the optimal dose remains to be determined.

Corresponding address

Dr. R. Wiggeraad
Radiotherapy Centre West
Lijnbaan 32, 2512 VA The Hague
The Netherlands
r.wiggeraad@mchaaglanden.nl

Conflict of interest. R.W. was a former speaker for Brainlab. No conflicts of interest for the additional authors.

Tab. 3 Univariate analysis of prognostic factors for local progression

		All local progressions	Tumor progression	Pseudo progression
Dose	1 fraction of 15 Gy vs. 3 fractions of 8 Gy	p=0.93	p=0.27	p=0.25
KPS	<90 vs. ≥90	p=0.50	p=0.17	p=0.50
PTV	<13 cm ³ vs. 13–20 cm ³ vs. >20 cm ³	p=0.72	p=0.60	p=0.36
Previous and later WBI	Yes vs. no	p=0.07	p=0.04 ^a	p=0.02 ^b
Only previous WBI	Yes vs no	p=0.02	p=0.63	p=0.02 ^c

Log rank: p value ^aSix of 75 metastases without WBI developed tumor progression, 8 of 36 metastases with WBI developed tumor progression ^bThree of 75 metastases without WBI developed pseudo progression, 6 of 36 metastases with WBI developed pseudo progression ^cSix of 22 metastases with previous WBI developed pseudo progression, 3 of 89 metastases without previous WBI developed pseudo progression, no metastasis with WBI after SRT developed pseudo progression **KPS** Karnofsky performance score, **WBI** whole brain irradiation, **SRT** stereotactic radiotherapy.

Tab. 4 Results from the literature on SRT for large brain metastases

Author	Diameter/volume	Dose	BED ₁₂ (Gy)	12-month local control (%)	Radiation toxicity (%)
Vogelbaum [16]	3–4.5 cm	1 fraction of 15 Gy	33.8	45	na
Ernst-Stecken [18]	1–5 cm	5–6 fractions of 7 Gy	45–55.4	76	14% (V _{4Gy} <23 ml) ^c 70% (V _{4Gy} >23 ml)
Narayana [17]	2–5 cm	5 fractions of 6 Gy	45	70	na
Higuchi [8]	3–4.5 cm	3 fractions of 10 Gy	55	76	0 ^b
Marchetti [15]	0.3–48.2 cm ³	3 fractions of 8 Gy	40	59	2 ^b
This series	76% >3 cm	1 fraction of 15 Gy	33.8	54	15 ^a
	76% >3 cm	3 fractions of 8 Gy	40	61	25 ^a

na not available V_{4Gy} volume of tissue receiving at least 4 Gy per fraction ^apseudoprogression ^blate radiation necrosis ^cnew or increasing necrotic lesions.

References

- Linskey ME, Andrews DW, Asher AL et al (2010) The role of stereotactic radiosurgery in the management of patients with newly diagnosed brain metastases: a systematic review and evidence-based clinical practice guideline. *J Neurooncol* 96:45–68
- Regine WF, Huhn JL, Patchell RA et al (2002) Risk of symptomatic brain tumor recurrence and neurologic deficit after radiosurgery alone in patients with newly diagnosed brain metastases: results and implications. *Int J Radiat Oncol Biol Phys* 52:333–338
- Shaw E, Scott C, Souhami L et al (2000) Single dose radiosurgical treatment of recurrent previously irradiated primary brain tumors and brain metastases: final report of RTOG protocol 90–05. *Int J Radiat Oncol Biol Phys* 47:291–298
- Hoefnagels FW, Lagerwaard FJ, Sanchez E et al (2009) Radiological progression of cerebral metastases after radiosurgery: assessment of perfusion MRI for differentiating between necrosis and recurrence. *J Neurol* 256:878–887
- Chao ST, Barnett GH, Vogelbaum MA et al (2008) Salvage stereotactic radiosurgery effectively treats recurrences from whole-brain radiation therapy. *Cancer* 113:2198–2204
- Molenaar R, Wiggendaad R, Verbeek-de KA et al (2009) Relationship between volume, dose and local control in stereotactic radiosurgery of brain metastasis. *Br J Neurosurg* 23:170–178
- Fahrig A, Ganslandt O, Lambrecht U et al (2007) Hypofractionated stereotactic radiotherapy for brain metastases—results from three different dose concepts. *Strahlenther Onkol* 183:625–630
- Higuchi Y, Serizawa T, Nagano O et al (2009) Three-Stage stereotactic radiotherapy without whole brain irradiation for large metastatic brain tumors. *Int J Radiat Oncol Biol Phys* 74(5):1543–1548
- Theelen A, Martens J, Bosmans G et al (2012) Relocatable fixation systems in intracranial stereotactic radiotherapy. Accuracy of serial CT scans and patient acceptance in a randomized design. *Strahlenther Onkol* 188:84–90
- Santvoort J van, Wiggendaad R, Bos P (2008) Positioning accuracy in stereotactic radiotherapy using a mask system with added vacuum mouth piece and stereoscopic X-ray positioning. *Int J Radiat Oncol Biol Phys* 72:261–267
- Macdonald DR, Cascino TL, Schold SC Jr, Cairncross JG (1990) Response criteria for phase II studies of supratentorial malignant glioma. *J Clin Oncol* 8:1277–1280
- Herfarth KK, Izwekowa O, Thilmann C et al (2003) Linear-accelerated radiosurgery of cerebral melanoma metastases. Analysis of 122 metastases treated in 64 patients. *Strahlenther Onkol* 179:366–371
- Wiggendaad R, Verbeek-de KA, Kal HB et al (2011) Dose-effect relation in stereotactic radiotherapy for brain metastases. A systematic review. *Radiother Oncol* 98:292–297
- Yang HC, Kano H, Lunsford LD et al (2011) What factors predict the response of larger brain metastases to radiosurgery? *Neurosurgery* 68:682–690
- Marchetti M, Milanesi I, Falcone C et al (2011) Hypofractionated stereotactic radiotherapy for oligometastases in the brain: a single-institution experience. *Neurol Sci* 32:393–399
- Vogelbaum MA, Angelov L, Lee SY et al (2006) Local control of brain metastases by stereotactic radiosurgery in relation to dose to the tumor margin. *J Neurosurg* 104:907–912
- Narayana A, Chang J, Yenice K et al (2007) Hypofractionated stereotactic radiotherapy using intensity-modulated radiotherapy in patients with one or two brain metastases. *Stereotact Funct Neurosurg* 85:82–87
- Ernst-Stecken A, Ganslandt O, Lambrecht U et al (2006) Phase II trial of hypofractionated stereotactic radiotherapy for brain metastases: results and toxicity. *Radiother Oncol* 81:18–24
- Barendsen GW (1982) Dose fractionation, dose rate and iso-effect relationships for normal tissue responses. *Int J Radiat Oncol Biol Phys* 8:1981–1997
- Thames HD Jr, Withers HR, Peters LJ, Fletcher GH (1982) Changes in early and late radiation responses with altered dose fractionation: implications for dose-survival relationships. *Int J Radiat Oncol Biol Phys* 8:219–226
- Brenner DJ (2008) The linear-quadratic model is an appropriate methodology for determining isoeffective doses at large doses per fraction. *Semin Radiat Oncol* 18:234–239
- Brandtsma D, Stalpers L, Taal W et al (2008) Clinical features, mechanisms, and management of pseudoprogression in malignant gliomas. *Lancet Oncol* 9:453–461
- Yoshii Y (2008) Pathological review of late cerebral radionecrosis. *Brain Tumor Pathol* 25:51–58
- Jain R, Narang J, Sundgren PM et al (2010) Treatment induced necrosis versus recurrent/progressing brain tumor: going beyond the boundaries of conventional morphologic imaging. *J Neurooncol* 100:17–29
- Korytko T, Radivoyevitch T, Colussi V et al (2006) 12 Gy gamma knife radiosurgical volume is a predictor for radiation necrosis in non-AVM intracranial tumors. *Int J Radiat Oncol Biol Phys* 64:419–424
- Minniti G, Clarke E, Lanzetta G et al (2011) Stereotactic radiosurgery for brain metastases: analysis of outcome and risk of brain radionecrosis. *Radiat Oncol* 6:48