

Adult stem cells and their trans-differentiation potential—perspectives and therapeutic applications

Sabine Hombach-Klonisch · Soumya Panigrahi ·
Iran Rashedi · Anja Seifert · Esteban Alberti ·
Paola Pocar · Maciej Kurpisz · Klaus Schulze-Osthoff ·
Andrzej Mackiewicz · Marek Los

Received: 26 April 2008 / Revised: 16 June 2008 / Accepted: 18 June 2008 / Published online: 16 July 2008
© Springer-Verlag 2008

Abstract Stem cells are self-renewing multipotent progenitors with the broadest developmental potential in a given tissue at a given time. Normal stem cells in the adult organism are responsible for renewal and repair of aged or damaged tissue. Adult stem cells are present in virtually all tissues and during most stages of development. In this review, we introduce the reader to the basic information about the field. We describe selected stem cell isolation techniques and stem cell markers for various stem cell

populations. These include makers for endothelial progenitor cells (CD146/MCAM/MUC18/S-endo-1, CD34, CD133/prominin, Tie-2, Flk1/KD/VEGFR2), hematopoietic stem cells (CD34, CD117/c-Kit, Sca1), mesenchymal stem cells (CD146/MCAM/MUC18/S-endo-1, STRO-1, Thy-1), neural stem cells (CD133/prominin, nestin, NCAM), mammary stem cells (CD24, CD29, Sca1), and intestinal stem cells (NCAM, CD34, Thy-1, CD117/c-Kit, Flt-3). Separate section provides a concise summary of recent clinical trials involving

Sabine Hombach-Klonisch, Soumya Panigrahi, and Iran Rashedi contributed equally to this review manuscript.

Klaus Schulze-Osthoff and Marek Los share senior authorship.

S. Hombach-Klonisch · A. Seifert
Department of Human Anatomy and Cell Science,
University of Manitoba,
Winnipeg, Canada

S. Panigrahi
Department of Physiology, University of Manitoba,
Winnipeg, Canada

S. Panigrahi · I. Rashedi
Manitoba Institute of Cell Biology, CancerCare Manitoba,
University of Manitoba,
Winnipeg, Manitoba, Canada R3E 0V9

I. Rashedi
Department of Biochemistry and Medical Genetics,
University of Manitoba,
Winnipeg, Canada

E. Alberti
Department of Neurobiology,
International Center of Neurological Restoration, CIREN,
Havana, Cuba

P. Pocar
Department of Animal Science, Faculty of Veterinary Medicine,
University of Milan,
Milan, Italy

M. Kurpisz
Institute of Human Genetics, Polish Academy of Science,
Poznan, Poland

K. Schulze-Osthoff
Institute of Molecular Medicine, University of Duesseldorf,
Duesseldorf, Germany

A. Mackiewicz
Department of Cancer Immunology,
Poznan University of Medical Sciences,
and Great-Poland Cancer Center,
Poznan, Poland

M. Los (✉)
BioApplications Enterprises,
Winnipeg, Manitoba, Canada
e-mail: mjelos@gmail.com

stem cells directed towards improvement of a damaged myocardium. In the last part of the review, we reflect on the field and on future developments.

Keywords Autoimmune disease · G-CSF · Graft vs. host reaction · Stem/progenitor cell · Trans-differentiation

Abbreviations

AGM	aorta–gonad–mesonephros
BCRP1	breast cancer resistance protein1
BM	bone marrow
CABG	coronary artery bypass graft
CNS	central nervous system
CSC	cardiac stem cell
ESC	embryonic stem cells
Flk1	fetal liver kinase-1
G-CSF	granulocyte-colony stimulating factor
GM-CSF	granulocyte-macrophage-colony stimulating factor
HLA	human leukocyte antigen
HSC	hematopoietic stem cell
LVAD	left ventricular assist device
LVEF	left ventricular ejection fraction
MAPC	multipotent adult progenitor cells
MRF	myogenic regulatory factor
MSC	mesenchymal stromal cell
NCAM	neural cell adhesion molecule
NSC	neuronal stem cells
NYHA	New York Heart Association
PB	peripheral blood
Sca1	stem cell antigen 1
SP	side population
UC	umbilical cord
VEGFR2	vascular endothelial growth factor receptor 2

Introduction—adult stem cells and their surface markers

Adult stem cells are clonogenic, self-renewing, and pluripotent cells with a plasticity to differentiate into cell types of the particular tissue in which they reside and often to trans-differentiate into different types of tissues [1]. The tremendous proliferative potential of these cells may lead to the development of cancer if the control of their differentiation, and/or proliferation, and/or apoptotic program is impaired [2]. Stem cell activity has been demonstrated in many tissues/organs, but the exact location of these adult stem cells is not always clear because of a current lack of well-defined organ-specific stem cell markers. Adult stem cells are usually located in a specific cellular niche, and niche microenvironment determine the

status of stem cell activation, thus ensuring a balance between maintenance of the stem cell pool and production of progenitor cells engaged in tissue differentiation [3]. The identification and selection of stem cells within a given tissue/organ largely relies on the presence of specific cell surface markers (Table 1). Other identification methods include the ability of certain stem cells to exclude fluorescent dyes (rhodamine 123, Hoechst 33342) and DNA label retention as well as their ability to form colonies and differentiate into certain lineages as seen in mesenchymal/stromal stem cells. Dye exclusion techniques permit selection of a “side population,” and this is facilitated by the ABCG2 gene product, also named breast-cancer-resistant protein (BCRP1) [4]. This member of the family of ABC transporters has been shown to be a positive selection marker of pluripotent cells from various adult tissue sources. However, ABCG2 is neither unique to nor ubiquitously expressed in all stem/progenitor cells [5]. Another way to identify slowly cycling adult stem cells is DNA label retention. Quiescent stem cells retain the DNA label for much longer than dividing cells where the DNA label is diluted with each cell division [6]. In the following paragraphs, we will characterize at greater detail the various stem cells populations, their therapeutic potential, and existing isolation techniques.

Growth and isolation of stem cells and tests to confirm stem cell presence

The trans-differentiation potential of adult stem cells and their capacity for tissue renewal and damage repair has attracted much attention from biotechnologists and clinicians [7], and the isolation and in vitro maintenance of stem cells have immense importance in applied biology. Although flow cytometric separations of stem cells or positive and negative selections using magnetic beads tagged with antibodies targeting specific markers on the surface of stem cells are used routinely now in many applications, alternative approaches for stem cell identification have been proposed based on the specific behavior of individual stem cells. One of these approaches exploits stem cell homing characteristics [8]. These and other principles were proven to be effective in isolating stem cells for research and biological applications. Taking hematopoietic stem cells (HSCs) as an example, we will provide an overview of some of these important isolation techniques, which could also be applied to various other stem cells.

Isolation of stem cells by flow cytometry In flow cytometry, a mixture of cells tagged by appropriate fluorochrome-labeled stem cell markers is passed through a laser beam.

Table 1 Commonly used markers to identify adult stem cells in different tissues

Marker	Synonym	Significance	Literature
Endothelial progenitor cells			
CD146	MCAM, MUC18, S-endo-1	Integral membrane protein expressed on endothelial precursor cells and circulating endothelial cells	[115]
CD34		Cell surface protein on bone marrow cells, HSCs, endothelial progenitor and muscle stem cell	[115]
CD133	Prominin		[116]
Tie-2		Receptor for angiopoietin on endothelial progenitor cells and HCS	[115]
Fetal liver kinase-1	Flk1, KDR VEGFR2	Endothelial cell surface receptor protein that identifies endothelial cell progenitor	[117]
Hematopoietic stem cells			
CD34		Cell surface protein on bone marrow cells, HSCs, endothelial progenitor and muscle stem cell	[24, 25]
c-Kit	CD117, YB5.B8	Cell surface receptor on bone marrow cell types that identifies HSC, hematopoietic progenitor cells and mesenchymal stem cell (MSC)	[15, 25]
Stem cell antigen	Sca1, Ly-6A/E	Cell surface protein on bone marrow (BM) cell, indicative of HSC and MSC	[118]
Mesenchymal stem cells			
CD146	MCAM, MUC18, S-endo-1	Cell surface protein found on bone marrow fibroblasts	[118]
STRO-1 antigen		Cell surface glycoprotein on subsets of bone marrow stromal (mesenchymal) cells	[119]
Thy-1		Cell surface protein on HSC, MSC	[120]
Neural stem cells			
CD133	Prominin	Cell surface protein that identifies neuronal stem cells	[21]
Nestin		Marker for NSC in the CNS and in culture	[121]
Neuronal cell adhesion molecule	NCAM	Cell surface molecule that promotes cell–cell interaction; NSC migration; primitive neuroectoderm formation	[122]
Breast stem cells			
CD24 CD29		Cell surface proteins on mammary repopulating units in mice	[123]
Stem cell antigen	Sca1, Ly-6A/E	Cell surface protein on human mammary repopulating units	[124]
Intestinal stem cells			
NCAM		Marker for liver and pancreas stem cells	[125]
CD34		Marker for liver and pancreas stem cells	[126]
Thy-1		Marker for liver and pancreas stem cells	[126]
c-Kit		Marker for liver and pancreas stem cells	[126]
Flt-3		Marker for liver and pancreas stem cells	[126]

The cells scatter the fluorescence which provides information on the cell morphology, composition of surface proteins, DNA content, and cytoplasmic processes. The advantage of flow cytometry is the speed of the flow that allows quick processing of large population of cells. Cytometric analysis goes one step further and allows cell sorting [9], a process that breaks the fluid stream containing the cells into droplets by piezoelectric perturbation. It is then possible to deflect a selected droplet with precise timing on a charge given to the stream as it passes through an electric field. Droplets containing the (stem) cells of interest are deflected in the electric field and collected [10].

Immunomagnetic-beads-based isolation method Immunomagnetic beads coated with specific antibodies are used either for isolation or depletion of various types of cells. Positive or negative cell isolation can be performed

depending on the nature of the cell surface markers and its specific application. Positive cell isolation amends itself to any downstream application after removal of the beads. Negative cell isolation is the method of choice to ensure that cells of interest remain unaffected. Cells with multiple cell surface markers can be isolated by the combination of negative and positive cell isolation [11]. Such technologies based on immunomagnetic beads are currently in use in two major clinical applications: (1) CD34-positive stem cell isolation and (2) in vitro T cell isolation and expansion for clinical trials in novel adoptive immunotherapy [12].

Other techniques Various other stem cell isolation techniques have been proposed and tested in recent years. Stem cell isolation by acoustic standing waves (acoustophoresis) in microfluidic channels is one of such methods where controlled acoustic waves were used within a fluid flow

column with antinodes and nodes maintained in different flow layers of the channel, allowing fluid components to differentially migrate to areas of preferred acoustic interaction [13]. Acoustophoresis allows particles and cells to be driven towards the node or anti-node of a standing wave that depends on the characteristics of density and compressibility [14].

Isolation of HSCs HSCs are highly enriched in the population characterized by low or undetectable levels of the lineage markers found on mature hematopoietic cells (B220, CD3, CD4, CD5, CD8, Mac-1, GR-1, Ter119, and NK1.1) and by high levels of c-Kit and Sca-1 [15]. Furthermore, differential expression of SLAM receptors (CD150, CD244, and CD48) on HSC and more restricted multipotent hematopoietic progenitors allows further, more distinct selection. Highly purified HSCs are CD150⁺, CD244⁻, and CD48⁻, while multipotent hematopoietic progenitors are CD244⁺, CD150⁻, and CD48⁻. In the course of further differentiation/commitment, they become CD48⁺, CD244⁺, and CD150⁻ [16, 17]. Thus, sorting bone-marrow-derived cells according to the above criteria is a relatively simple and practical way of stem cell enrichment. Another approach to isolating stem-cell-enriched populations is based on the ‘dye-efflux’ properties of proliferating HSCs. Due to high expression of ABCG2 transporter, HSCs rapidly efflux the DNA dye Hoechst 33342. Cells maintaining an efficient efflux of the dye can be identified by flow cytometry, and cells that efflux the dye are referred to as ‘side population’ or SP cells when visualized on dot plot [4].

Growth and validation for stem cell presence Like other mammalian cells, stem cells are also grown and maintained at 37°C in humidified cell culture incubators under a 5% CO₂ atmosphere. The media requirements vary among individual stem cell types, and it is essential not to induce differentiation of the cultured stem cells. HSCs are obtained from bone marrow aspirates, placental, or umbilical cord blood, and their high growth rate makes them prone to differentiation in culture. Bone marrow origin stem cells are grown in culture media supplemented with serum [18], while HSCs usually require a co-culture system containing fibroblast feeder layers that support growth and differentiation. The change of cytokine conditions in such culture systems affects the differentiation of stem cells [19]. The effects of specific culture conditions and speed of differentiation varies among different stem cell populations. Bone marrow stromal cells attach to culture dishes and continue to grow slowly for weeks before differentiation [20]. Neural stem cells from fetal or adult brain tissue can grow suspended in culture medium without any additional serum supplements [21].

Validation of the presence of stem cells in a culture system with histochemical methods, antigenic markers, and morphogenetic studies are primarily dependent on the surface morphology and immuno-phenotype of the stem cells of interest. For in vitro studies, cytogenetic analysis or RT-PCR-based methods are commonly applied.

Hematopoietic stem cells

HSCs are among the best characterized adult stem cells and the only stem cells being routinely used in the clinics. HSCs are able to renew themselves or differentiate into precursors which produce specialized hematopoietic cells, including lymphocytes, dendritic and natural killer cells, megakaryocytes, erythrocytes, granulocytes, and macrophages [22]. Cells in the hematopoietic hierarchy have different multi-potentiality, differentiation and self-renewal capacities, and are able to cope with the high demand for continuously producing large numbers of blood cells. The first progeny of HSCs are multipotent progenitors which retain the ability to differentiate into all hematopoietic lineages but show a lower capacity to proliferate [23]. Multipotent progenitors are more abundant than HSCs and differentiate into oligopotent progenitors, which in turn give rise to more lineage-committed precursor cells.

During sequential differentiation, stem cells present various antigenic characteristics which are associated with their properties and function. These antigens allow for the definition of stem cell subpopulations and permit clinicians to improve the outcome of HSCs transplantation by increasing the purity of HSCs product used in allo-engraftments. Human HSCs express CD34 surface antigen (CD34⁺), which is commonly used as a marker in clinical settings to identify and quantify the population of progenitor cells to be infused [24, 25]. Scientists have been trying to narrow the subset of progenitor cells by defining a set of markers that are more consistently expressed on these cells. Human HSCs are known to exhibit CD34⁺, Thy1⁺, CD38^{lo/-}, C-kit^{-/lo}, CD105⁺, Lin⁻ phenotype [25, 26]. However, there is not a general agreement on the association between any combination of these antigenic properties and function of stem cells.

Sources of HSCs HSCs can be isolated from bone marrow (BM). BM also accommodates stromal cells, mesenchymal stem cells, and variably mature blood cells and their progenitors. HSCs constitute only a small fraction of BM population (1 in 10⁴ to 1 in 10⁸ of BM nucleated cells) [27]. HSCs can also be isolated from peripheral blood (PB) where they can be found in small numbers [27, 28]. Stimulation with mobilizing agents, including cytokines such as G-CSF alone or in a combination with GM-CSF

and/or other agents, dramatically increases the release of HSCs from BM to PB [28]. This allows collecting a high number of progenitor cell types, which constitutes an important improvement in engraftment success and effective repopulation with neutrophils and platelets [24]. Another important source of HSCs for clinical purposes is umbilical cord (UC)/placenta blood. Its therapeutic use has become popular after successful applications of UC-derived HSCs in children with Fanconi anemia [29]. A practical constraint with HSCs collected from UC blood is their limited quantity, which is predominantly due to the low blood volume obtained from these tissues. The low UC stem cell dosage of approximately 10% of the amount of marrow transplants is adequate for transplants in children and low-weight patients but delivers too few stem cells for use in most adults with a recommended dose of more than 2.5×10^6 CD34⁺ cells per kilogram body weight [24]. In addition to their quantity, HSCs collected from UC and PB differ in some other characteristics (Table 2). In general, HSCs obtained from tissues at earlier developmental stages have greater capacity for self-replication and long-term growth in vitro [30] and show different homing and surface properties [31]. Nonetheless, it is currently unclear whether these variations have any clinical significance. In animal models, HSCs can also be obtained from the fetal hematopoietic system including aorta–gonad–mesonephros (AGM), liver and spleen, which provide useful sources of stem cells for experimental purposes.

Heart and muscle stem cells and potential for regeneration of the heart muscle

Satellite cells, the muscle stem cells, were first described by Mauro [32] in 1961 as committed precursor cells residing in skeletal muscle. Upon an injury, the basic role of these cells is to restore skeletal muscle function. An inflicted stress may stimulate these cells to transform into myoblasts by the

activation of myogenic regulatory (transcription) factors (MRF) [33]. Quiescent (satellite) cells appear to be negative for MRF, but during activation, their expression is upregulated (MyoD, Myf5, myogenin, MRF4) [33]. Moreover, some transcripts transform to alternatively spliced isoforms when satellite cells start to proliferate (CD34⁺ from truncated to a full-length form, MNFβ to MNFα) [34]. MRF are first expressed during early embryogenesis when myoblasts are formed (prior to cardiac-specific cells), and in case of muscle injury, they appear about 6 h after injury [33]. Following proliferation, differentiation and multi-fusion, myoblast cells form new myotubes that become finally differentiated muscle fibers [35]. This potential of myoblasts and satellite cells for regeneration of heart muscle was first explored in a preclinical trial using a dog heart model of cryoinjury [36]. Myoblasts or satellite cells may give rise to both self-renewing muscle-type-specific populations of myogenic stem cells and to progenitor muscle cells.

In rodents, a stem cell population was initially identified as muscle SP that could reconstitute hematopoiesis [37]. The analysis of the transcriptional profile of this SP population obtained from skeletal muscle revealed that these cells shared transcripts with embryonic stem cells (ESC) and overlapped with bone marrow SP in almost half of all transcripts [38]. Recent studies have shown that the majority of obtained SP cells were positive for Sca1 (stem cell antigen 1), CD31, and CD45 [39]. FACS sorting of these rodent cells resulted in the identification of three subpopulations with different myogenic and hematopoietic potential and distinct proliferative and regenerative capacities.

Since the heart has a low regenerative capacity, efforts have been undertaken to restore its function after, e.g., ischemic or non-ischemic damage. Its own modest regenerative potential (possibly only by ~1% renewal per year) can be either provided by circulating multipotential stem cells [40] or cardiac progenitor stem cells residing in specific heart niches [41]. Putative human cardiac stem and progenitor cells (hCSCs) are now distinguished into four

Table 2 Comparison of BM with two other sources of HSC

	PB/G-CSF	UC blood
Advantage	Easier, less aggressive collection High number of progenitor cells Faster engraftment Faster neutrophil, platelet, immune system recovery Less frequency of donation-associated risks	Easy to collect without any risk for the donor Less restriction for HLA compatibility Lower rate of GVHD Better accessibility from UC blood banks for unrelated transplantations Can be stored for years
Disadvantage	Administration of mobilizing agents is required Higher risk of chronic GVHD in allografts	Limited supply Low number of progenitor cells, inadequate for transplant in adults Slower engraftment Unable to provide additional cells in case of need for second transplant

PB/G-CSF Peripheral-blood-derived HSC treated with G-CSF

main resident cardiac cell types characterized by c-Kit, Sc α 1 (rodents), MDR-1, and Isl-1 markers [42]. Divided into primary and secondary heart fields (according to the heart development), these cells may reconstitute main structural elements of the myocardium such as cardiomyocytes, endothelial and conductive cells [43], and/or post-natal cardiac progenitors. First and second heart field progenitors express common mesodermal markers: Oct4, Mesp, Bry, Nkx2.5, while Isl-1 belongs exclusively to the second field characteristics giving rise to all previously mentioned cells, while the first heart field progenitors may differentiate only into cardiac conductive and cardiac muscle cells. Attempts to multiply in vitro hCSCs for possible heart regeneration are ongoing [44]. The most ambitious aim would be to practically utilize totipotent stem cells, and preclinical studies in rodents are well advanced with application of in vitro differentiated cardiomyocytes and/or genetically pre-programmed pluripotent cells [45]. Yet it seems that at the present stage, the numbers of cells obtained in this way are low and inadequate for clinical trials [46].

Despite advances in pharmacological treatment, organ transplantation and cardiac devices (LVAD), data from the Framingham Heart Study indicate that the survival of patients who were diagnosed with heart failure is still poor [47]. Stem cells, including myoblasts, seem to ideally suit for autologous transplantation, and more than ten clinical trials involving myoblast transplantation have so far been performed (Table 3). Several features favor myoblasts as the first target for treatment of an ischemic myocardium: (1) Autologous origin excludes adverse immune responses; (2) relatively low plasticity due to their progenitor commitment reduces the risk of teratoma formation as observed with totipotent cells; (3) myoblasts have a relatively high proliferation potential, although there could still be better

candidates among muscle-derived-stem cells; and (4) myoblasts are relatively resistant to ischemic conditions. Table 3 summarizes current clinical trials involving myoblast transplantation. The majority of the trials are at stage I/II; no proper control groups were assigned (except for the MAGIC trial). A vast majority of the trials were conducted at the opportunity of other surgical interventions (CABG or LVAD implantations), which makes it difficult to evaluate the beneficial effect of myoblasts alone and which resulted in the premature termination of the expensive MAGIC trial, with statistical power in over a hundred patients being compromised by multifactorial assessment of CABG operations. Nonetheless, at least three major issues overcome a general pessimistic view over the current stage of stem cells interventions in the failing heart: (a) heart hemodynamic parameters improved and correlated with higher numbers of myoblasts implanted; (b) autologous myoblasts appeared to be safe and feasible candidate cells, although the arrhythmia episodes associated in a majority of the trials required the concurrent cardioverter/defibrillator implantation [48].

Cells with myogenic (contractile) potential can be considered as the better alternative to bone-marrow-derived stem cells, which can only be harvested in low quantities, do not possess contractile properties, and only have transitory benefits [49]. Although myoblasts do not remain in the myocardium infinitely, their beneficial effect on the ejection fraction (LVEF), which begins 3–6 months after implantation and may last until almost 2 years (Kurpisz, unpublished), has been observed in most of the trials. This holds a promise for repetitive dosing of myoblasts using a low invasive percutaneous approach. Such data are already available from preclinical trials in rats and pigs [50]. The cell homing inside the myocardium seems to be sufficient when using either a percutaneous or an intramyocardial

Table 3 Clinical trials involving myoblast transplantation

Approach	Patient No.	Results	Reference
Intramyocardial, adjunct to CABG	10 (no controls)	LVEF increased, improved NYHA class	[127]
Intramyocardial, adjunct to LVAD implantation	5 (no controls)	Myofibers parallel to host myocardial fibers, increased blood vessel density	[128]
Intramyocardial, adjunct to CABG	10 (no controls)	LVEF and regional wall motion increased	[129]
Percutaneous	9 (no controls)	LVEF improved in 6 out of 9 cases	[130]
Intramyocardial, adjunct to CABG	18 (no controls)	LVEF regional wall motion and viability increased improved NYHA class, reduction of scar size	[131]
Intramyocardial, adjunct to CABG	11 (no controls)	LVEF, regional wall motion and viability increased	[132]
Percutaneous	5 (no controls)	LVEF and regional wall motion increased	[133]
Intramyocardial, adjunct to CABG or LVAD implantation	18 (no controls)	LVEF and viability increased, engraftment of myoblasts	[134]
Percutaneous	6 (plus controls)	LVEF increased, improved NYHA class	[135]
Intramyocardial, adjunct to CABG, allogeneic	2 (no controls)	LVEF increased	[136]
Intramyocardial, adjunct to CABG	120 (plus control)	Reduction of LV remodeling, LVEF increased at high dose	[48]

approach [51]. A long-term survival of myoblasts within the myocardium has been recently demonstrated with pro-angiogenic gene modifications, for example, with VEGF-transduced myoblasts in rats [52]. Their functionality may also be improved by connexin 43 overexpression [53]. Finally, different types of stem cells can be combined at progenitor-committed stages, thus greatly enhancing the therapeutic outcome [54] and ultimately leading to the rejuvenation of the whole organ.

Adult stem cell plasticity and the implications for regenerative medicine


Cell-based therapy may in the nearer future represent a new strategy to treat a wide array of clinical conditions. The use of adult stem cells as opposed to human embryonic stem cells for therapy avoids ethical problems and has two additional advantages: (a) Adult stem cells can be isolated from patients, and this overcomes the problem of immunological rejection and (b) the risk of tumor formation is greatly reduced as compared to the use of embryonic stem cells [55].

While pluripotency and plasticity are considered properties of early ESC, adult stem cells are traditionally thought to be restricted in their differentiation potential to the progeny of the tissue in which they reside. When parts of an organ or tissue are transplanted to a new site, the transplanted tissue maintains its original character. Similarly, when dissociated cells from an organ or tissue are cultured, they also tend to maintain their original phenotype. Despite losing some of their differentiation properties, they do not acquire differentiated characteristics of a different cell lineage. However, a remarkable plasticity in differentiation potential of stem cells derived from adult tissues was

recently suggested [56] (Fig. 1). In 1998, Ferrari et al. [57] first reported that mouse bone-marrow-derived cells give rise to skeletal muscle cells when transplanted into damaged mouse muscle. Thereafter, transplanted bone marrow cells were reported to generate a wide spectrum of different cell types, including hepatocytes [58], endothelial, myocardial [59, 60], neuronal, and glial cells [61]. Moreover, HSC can produce cardiac myocytes and endothelial cells [62], functional hepatocytes [63], and epithelial cells of the liver, gut, lung, and skin [64]. Mesenchymal stromal cells (MSC) of the bone marrow can generate brain astrocytes [65]. Enriched stem cells from adult mouse skeletal muscle were shown to produce blood cells [66, 67]. In most of these plasticity studies, genetically marked cells from one organ of an adult mouse apparently gave rise to cell type characteristics of other organs following transplantation, suggesting that even cell types once thought to be terminally differentiated are far more plastic in their developmental potential than previously thought. A critical aspect of the observation of adult stem cell plasticity is that in order for plasticity to occur, cell injury is necessary [68]. This suggests that microenvironmental exposure to the products of injured cells may play a key role in determining the differentiated expression of marrow stem cells [69].

The events underlying stem cells plasticity could relate to a variety of mechanisms such as dedifferentiation, transdifferentiation, epigenetic changes, and/or cell fusion. Re-routing of cell fate may result from the multistep process known as dedifferentiation where cells revert to an earlier, more primitive phenotype characterized by alterations in gene expression pattern which confer an extended differentiation potential (Fig. 2). In urodele amphibians, cell dedifferentiation is a common mechanism resulting in the functional regeneration of complex body structures throughout life, including limbs, tail, and even spinal cord

Fig. 1 Diagram illustrating plasticity of bone-marrow-derived cells






Fig. 2 Diagram illustrating the different mechanisms of cell fate switching in adult stem cells

[70]. Recent studies on the plasticity of murine myotubes [71] and other cells derived from adult tissues suggest that dedifferentiation may also be possible in mammals [72]. At the molecular level, *MSX1* has been identified as a possible factor involved in dedifferentiation processes in both urodele and human cells [71, 73]. The small molecule, reversine, can induce murine myogenic lineage-committed cells to become multipotent mesenchymal progenitor cells that can proliferate and re-differentiate into bone and fat cells [74]. Epigenetic cell changes are probably involved and may be mediated by signals received from the injured cells. The identification of signals that induce dedifferentiation of somatic cells are key to elucidating the molecular mechanism of this phenomenon and may ultimately provide effective tools for the *in vivo* regeneration of mammalian tissues. Recent studies favor a model of continuous stem cell differentiation at the level of progenitor cells with dynamic transcriptional regulation of stem cell cycle phases and chromatin alterations associated with cell cycle transit [69]. In this model, stem cells represent a highly flexible ever-changing cell system in which the potential and characteristics of the stem cell are continually and reversibly changing with the cell cycle until a terminal differentiating stimulus is encountered at a cycle-susceptible time [69]. In this asynchronous stem cell population, there would always be a small percentage of cells receptive or primed for a specific differentiation (or de-differentiation) induction at any particular time.

Another mechanism put forward to explain stem cell switch to a novel phenotype is a process known as trans-differentiation (Fig. 2). Cells may differentiate from one cell type into another within the same tissue or develop into a completely different tissue without acquiring an intermediate recognizable, undifferentiated progenitor state. Recent

studies show clearly that bone-marrow-derived cells can colonize a wide variety of tissues in the body of a host [75, 76]. Although derived from the embryonic mesoderm, the developmental potential of bone marrow cells is not restricted to this germ layer, but these cells have also been shown to populate tissues of ectodermal and endodermal origin [77]. Both mesenchymal stem cells and bone-marrow-derived cells can give rise to a wide array of non-hematopoietic cell types such as astrocytes and neurons in the brain [61, 78], cardiac myocytes in models of infarction [60], skeletal muscle [57], and hepatocytes [79]. However, the reported frequencies of colonization are low, and it is unlikely that there is much repair of organ damage by bone marrow in the normal individual.

Despite examples of trans-differentiation events of adult stem cells being reported, these findings are still controversial. Most of the reports could not be confirmed in subsequent investigations [80], and to date, trans-differentiation has never been conclusively demonstrated in any experimental setting. In every case, differentiation from a rare population of stem cells has never been excluded, or “trans-differentiation” events turned out to be misinterpretations caused by cell fusion resulting in nuclear reprogramming and changes in cell fate [81, 82] (Fig. 2). It is now recognized that adult stem cells from bone marrow may fuse with cell of the target organ. So far, bone-marrow-derived cells were shown to form fusion heterokaryons with liver, skeletal muscle, cardiac muscle, and neurons [83]. There is evidence that such fused cells become mononucleated again, either by nuclear fusion or by elimination of supernumerary nuclei [82, 84]. Fusion and nuclear transfer experiments demonstrated that genes previously silenced during development could be reactivated by cytoplasmic factors modulating the epigenetic mechanisms responsible

for the maintenance of a specific state of cell differentiation. However, nuclear transfer experiments demonstrate that the capacity of cells to successfully reprogram diminishes with increasing developmental progression of the donor nuclei. Despite this limitation and the low frequency, cell fusion may be considered as a potential avenue for tissue repair.

In addition to the aforementioned phenomena of cell fate switching, the presence of a rare population of pluripotent primitive stem cells may also explain the acquisition of an unexpected phenotype. Recently, non-hematopoietic cell populations from bone marrow and umbilical cord blood were enriched by *in vitro* culture and demonstrate to have the potential to differentiate into derivatives of all three germ line layers with meso-, endo-, and ectodermal characteristics [85, 86]. Known as multipotent adult progenitor cells (MAPC), these cells contribute to most, if not all, somatic cell lineages, including brain, when injected into a mouse blastocyst [87]. Interestingly, while MAPC express Oct4, a transcription factor required for undifferentiated embryonic stem cells maintenance [88] at levels approaching those of ESC, MAPC do not express two other transcription factors known to play a major role in ESC pluripotency, Nanog and Sox2 [89]. This particular expression profile may contribute to the fact that the use of ESC, but not MAPC, carries the risk of generating tumors. Thus, MAPC are a promising source of autologous stem cells in regenerative medicine. Their low tumorigenicity, high regenerative plasticity, and optimal immunological compatibility are essential assets for the successful transplantation of MAPC-derived tissue-committed cells without immune-mediated rejection [90].

Bone marrow transplantation as a clinical example of regenerative medicine HSCs were primarily used in the treatment of patients with hematological malignancies. During the course of treatment, patient's cancerous cells are first destroyed by chemo/radiotherapy and subsequently replaced with BM or PB/G-CSF transplant from a human leukocyte antigen (HLA)-matched donor [91]. Allogeneic marrow transplants have also been used in the treatment of hereditary blood disorders including aplastic anemia, β -thalassemia, Wiskott–Aldrich syndrome and SCID, as well as inborn errors of metabolism disorders such as Hunter's syndrome and Hurler's syndrome [92–96].

During the therapy of hematological malignancies, autologous PB/G-CSF HSCs are collected prior to the treatment and reinfused into the patients after the course of the aggressive chemotherapy. With a similar approach, autologous HSCs may be used to reprogram immune system and reconstitute non-autoreactive immune cells as a treatment for autoimmune disorders. The problem using auto-engrafts to rescue HSC population in cancer treatment is that patient's cancerous cells may be inadvertently

collected and reinfused back into the patients along with the stem cells.

HSC transplants are also used as a therapeutic strategy against various types of solid tumors [97]. Graft-versus-tumor effect of allogeneic HSC transplants seems to be a result of an immune reaction between donor cytotoxic T cells and patient's malignant cells [98].

HSCs have the ability to generate cell types other than blood cells. Circulating HSCs are able to reside in distant tissues and participate in regeneration process by trans-differentiating into non-hematopoietic cells such as hepatocytes [15], skeletal muscle and cardiac myocytes [59, 60], neurons [99], and epithelial cells [64]. The existence of pluripotent stem cells, which are reprogrammed in the new microenvironment and differentiate into the cell types of the tissue to which they were recruited, has significant implications for regenerative medicine. However, the potential plasticity of HSCs are disputed by several reports of failure to show trans-differentiation [56] as well as other perplexing possibilities such as technical problems and the possible existence of several types of non-hematopoietic stem cells in BM [100].

Over the past two decades, HSCs have also been targeted for *ex vivo* gene therapy as a vehicle to transfer a modified gene in autologous settings [101]. This approach provides a promising alternative treatment for various inherent and acquired human diseases and may provide an alternative to currently still risky allogeneic HSC transplants.

Closing remarks

Since the beginning of the basic stem cell research in 1960s, scientists have been facing serious challenges with identifying true stem cells and proliferating and maintaining them in culture. For example, the scarcity of HSCs along with their morphological resemblance to other PB or marrow cell types in culture makes isolation and purification of stem cells difficult. Surface biomarkers currently used to define HSCs do not appear to be exclusively expressed on stem cells. This leads to the isolation of a heterogeneous population of cells that were mistakenly assumed as true stem cells. Self-renewal capacity of stem cells is an important property of these cells. Stem cells that maintain this property and do not differentiate into their progeny can provide an unlimited source of cells for both therapeutic and research applications.

Self-renewal divisions of stem cells are rare events in BM; HSCs replicate themselves slowly with an average turnover time of 30 days in adult BM [102]. This self-replication is also hard to induce *in vitro*, which altogether hinders the study of self-renewal and differentiation of stem cells, influencing factors and signaling involved in these

processes and, subsequently, further targeted manipulation of these pathways. In addition to these technical difficulties, various functional assays being used by researchers make the comparison and meta-analysis of data obtained from different studies inapplicable. Furthermore, many of the studies on stem cells including *in vivo* experiments have been conducted in animal models, and the true relevance and reliability of the results in the human remains to be determined.

One of the major challenges with HSC transplants is failure to engraft, which is mediated by donor T cells as a result of graft-versus-host disease. The lack of assays to selectively deplete the anti-host alloreactive T cells *ex vivo* and to separate graft versus host disease from beneficial effect of ‘graft versus tumor’ results in increased morbidity and mortality associated with HSC transplants. In addition, variability of HSC migration by mobilizing agents in different patients and the serious side effects and toxicity caused by some of them in donors require further development of methods to overcome cell dose barrier in HSC transplants [103]. Moreover, the absence of unrelated HLA-matched donors for many patients in need of HSC transplants and the high incidence of relapse of underlying diseases in transplant recipients are important challenges remained to be addressed in future attempts to improve the clinical outcome of HSCs-based therapies.

HSC/BM transplantation has been the lifesaver and a critical element of anticancer therapies, particularly in leukemias. Nowadays, however, other less toxic therapeutic options emerged. Tremendous progress has been made in the area of therapeutic antibodies [104, 105]. A more integrated approach is being taken to detect new targets for the treatment of cancer and other diseases [106, 107]. New proteins and peptides have recently been discovered that have cancer (semi-)selective properties [108–110]. Entirely new concepts that connect cell death, cell survival, and cell proliferation have been developed that warrant new targets [111]. Targeted therapies, with the epidermal growth factor receptor pathway as the best example gain importance [112], and finally, “traditional” chemotherapy approaches are being further developed so that new generations of chemotherapeutic drugs would become available [113].

Nevertheless, the therapeutic potential of adult stem cells as powerful tools in tissue regeneration and engineering has been recognized, and intense efforts are ongoing to harness and direct adult stem cell plasticity. Understanding the basic molecular mechanisms underlying cell fate switching of adult stem cells will be an essential contribution to ensuring their safe use in regenerative medicine. In the near future, it will most likely be possible to transplant genetically modified stem cells that carry a set of genes critical for, e.g., trans-differentiation, that are under externally regulated promoters [114] and, depending on the therapeutic require-

ments, direct their differentiation into desired cell populations. Even more “clinically friendly” systems may be developed where pharmacologic “small molecules” will be used to directly influence the trans- or re-differentiation potential of therapeutically applied adult stem cells both prior and after their administration into patient’s body.

Acknowledgments The authors (SHK, ML) would like to thank the Manitoba Health Research Council (MHRC) for their generous support. We apologize to those authors whose research was not cited in this review due to the reference number limitations.

References

- Filip S, English D, Mokry J (2004) Issues in stem cell plasticity. *J Cell Mol Med* 8:572–577
- Hombach-Klonisch S, Paranjothy T, Wiehac E, Pocar P, Mustafa T, Seifert A, Zahl C, Gerlach KL, Biermann K, Steger K, Hoang-Vu C, Schulze-Osthoff K, Los M (2008) Cancer stem cells as targets for cancer therapy: selected cancers as examples. *Arch Immunol Ther Exp* 56:165–180
- Kindler V (2005) Postnatal stem cell survival: does the niche, a rare harbor where to resist the ebb tide of differentiation, also provide lineage-specific instructions? *J Leukoc Biol* 78:836–844
- Scharenberg CW, Harkey MA, Torok-Storb B (2002) The ABCG2 transporter is an efficient Hoechst 33342 efflux pump and is preferentially expressed by immature human hematopoietic progenitors. *Blood* 99:507–512
- van Herwaarden AE, Schinkel AH (2006) The function of breast cancer resistance protein in epithelial barriers, stem cells and milk secretion of drugs and xenotoxins. *Trends Pharmacol Sci* 27:10–16
- Potten CS, Booth C, Pritchard DM (1997) The intestinal epithelial stem cell: the mucosal governor. *Int J Exp Pathol* 78:219–243
- Mimeault M, Hauke R, Batra SK (2007) Stem cells: a revolution in therapeutics—recent advances in stem cell biology and their therapeutic applications in regenerative medicine and cancer therapies. *Clin Pharmacol Ther* 82:252–264
- Juopperi TA, Schuler W, Yuan X, Collector MI, Dang CV, Sharkis SJ (2007) Isolation of bone marrow-derived stem cells using density-gradient separation. *Exp Hematol* 35:335–341
- Jayasinghe SM, Wunderlich J, McKee A, Newkirk H, Pope S, Zhang J, Staehling-Hampton K, Li L, Haug JS (2006) Sterile and disposable fluidic subsystem suitable for clinical high speed fluorescence-activated cell sorting. *Cytometry B Clin Cytom* 70:344–354
- Kamentsky LA (1979) Future directions for flow cytometry. *J Histochem Cytochem* 27:1649–1654
- Neurauter AA, Bonyhadi M, Lien E, Nokleby L, Ruud E, Camacho S, Aarvak T (2007) Cell isolation and expansion using dynabeads (R). *Adv Biochem Eng Biotechnol* 106:41–73
- Chen HW, Liao CH, Ying C, Chang CJ, Lin CM (2006) *Ex vivo* expansion of dendritic-cell-activated antigen-specific CD4⁺ T cells with anti-CD3/CD28, interleukin-7, and interleukin-15: potential for adoptive T cell immunotherapy. *Clin Immunol* 119:21–31
- Das CM, Becker F, Vernon S, Noshari J, Joyce C, Gascoyne PR (2005) Dielectrophoretic segregation of different human cell types on microscope slides. *Anal Chem* 77:2708–2719
- Petersson F, Nilsson A, Holm C, Jonsson H, Laurell T (2005) Continuous separation of lipid particles from erythrocytes by

- means of laminar flow and acoustic standing wave forces. *Lab Chip* 5:20–22
15. Pearce DJ, Ridler CM, Simpson C, Bonnet D (2004) Multi-parameter analysis of murine bone marrow side population cells. *Blood* 103:2541–2546
 16. Kiel MJ, Yilmaz OH, Iwashita T, Yilmaz OH, Terhorst C, Morrison SJ (2005) SLAM family receptors distinguish hematopoietic stem and progenitor cells and reveal endothelial niches for stem cells. *Cell* 121:1109–1121
 17. Wang N, Morra M, Wu C, Gullo C, Howie D, Coyle T, Engel P, Terhorst C (2001) CD150 is a member of a family of genes that encode glycoproteins on the surface of hematopoietic cells. *Immunogenetics* 53:382–394
 18. Landau T, Sachs L (1971) Characterization of the inducer required for the development of macrophage and granulocyte colonies. *Proc Natl Acad Sci USA* 68:2540–2544
 19. Ailles LE, Gerhard B, Hogge DE (1997) Detection and characterization of primitive malignant and normal progenitors in patients with acute myelogenous leukemia using long-term coculture with supportive feeder layers and cytokines. *Blood* 90:2555–2564
 20. Knospe WH, Hinrichs B, Fried W, Robinson W, Trobaugh FE Jr (1976) Normal colony stimulating factor (CSF) production by bone marrow stromal cells and abnormal granulopoiesis with decreased CFUc in S1/S1d mice. *Exp Hematol* 4:125–130
 21. Uchida N, Buck DW, He D, Reitsma MJ, Masek M, Phan TV, Tsukamoto AS, Gage FH, Weissman IL (2000) Direct isolation of human central nervous system stem cells. *Proc Natl Acad Sci USA* 97:14720–14725
 22. Rashedi I, Panigrahi S, Ezzati P, Ghavami S, Los M (2007) Autoimmunity and apoptosis—therapeutic implications. *Curr Med Chem* 14:3139–3159
 23. Bryder D, Rossi DJ, Weissman IL (2006) Hematopoietic stem cells: the paradigmatic tissue-specific stem cell. *Am J Pathol* 169:338–346
 24. Menichella G, Lai M, Serafini R, Pierelli L, Vittori M, Ciarli M, Rumi C, Puggioni P, Scambia G, Sica S, Leone G (1999) Large volume leukapheresis for collecting hemopoietic progenitors: role of CD 34+ precursors in predicting successful collection. *Int J Artif Organs* 22:334–341
 25. Murray L, Chen B, Galy A, Chen S, Tushinski R, Uchida N, Negrin R, Tricot G, Jagannath S, Vesole D et al (1995) Enrichment of human hematopoietic stem cell activity in the CD34+ Thy-1+Lin- subpopulation from mobilized peripheral blood. *Blood* 85:368–378
 26. Pierelli L, Scambia G, Bonanno G, Rutella S, Puggioni P, Battaglia A, Mozzetti S, Marone M, Menichella G, Rumi C, Mancuso S, Leone G (2000) CD34+/CD105+ cells are enriched in primitive circulating progenitors residing in the G0 phase of the cell cycle and contain all bone marrow and cord blood CD34+/CD38low/- precursors. *Br J Haematol* 108:610–620
 27. Group SCTC (2005) Allogeneic peripheral blood stem-cell compared with bone marrow transplantation in the management of hematologic malignancies: an individual patient data meta-analysis of nine randomized trials. *J Clin Oncol* 23:5074–5087
 28. Flomenberg N, Devine SM, Dippersio JF, Liesveld JL, McCarty JM, Rowley SD, Vesole DH, Badel K, Calandra G (2005) The use of AMD3100 plus G-CSF for autologous hematopoietic progenitor cell mobilization is superior to G-CSF alone. *Blood* 106:1867–1874
 29. Gluckman E, Broxmeyer HA, Auerbach AD, Friedman HS, Douglas GW, Devergie A, Esperou H, Thierry D, Socie G, Lehn P et al (1989) Hematopoietic reconstitution in a patient with Fanconi's anemia by means of umbilical-cord blood from an HLA-identical sibling. *N Engl J Med* 321:1174–1178
 30. Kim DK, Fujiki Y, Fukushima T, Ema H, Shibuya A, Nakauchi H (1999) Comparison of hematopoietic activities of human bone marrow and umbilical cord blood CD34 positive and negative cells. *Stem Cells* 17:286–294
 31. Surbek DV, Steinmann C, Burk M, Hahn S, Tichelli A, Holzgreve W (2000) Developmental changes in adhesion molecule expressions in umbilical cord blood CD34 hematopoietic progenitor and stem cells. *Am J Obstet Gynecol* 183:1152–1157
 32. Mauro A (1961) Satellite cell of skeletal muscle fibers. *J Biophys Biochem Cytol* 9:493–495
 33. Dhawan J, Rando TA (2005) Stem cells in postnatal myogenesis: molecular mechanisms of satellite cell quiescence, activation and replenishment. *Trends Cell Biol* 15:666–673
 34. Beauchamp JR, Heslop L, Yu DS, Tajbakhsh S, Kelly RG, Wernig A, Buckingham ME, Partridge TA, Zammit PS (2000) Expression of CD34 and Myf5 defines the majority of quiescent adult skeletal muscle satellite cells. *J Cell Biol* 151:1221–1234
 35. Grounds MD, White JD, Rosenthal N, Bogoyevitch MA (2002) The role of stem cells in skeletal and cardiac muscle repair. *J Histochem Cytochem* 50:589–610
 36. Marelli D, Ma F, Chiu RC (1992) Satellite cell implantation for neomyocardial regeneration. *Transplant Proc* 24:2995
 37. Meeson AP, Hawke TJ, Graham S, Jiang N, Elterman J, Hutcheson K, Dimairo JM, Gallardo TD, Garry DJ (2004) Cellular and molecular regulation of skeletal muscle side population cells. *Stem Cells* 22:1305–1320
 38. Seidel M, Rozwadowska N, Tomczak K, Kurpisz M (2006) Myoblast preparation into injured myocardium. *Eur Heart J* 8 (Suppl H):H8–15
 39. Uezumi A, Ojima K, Fukada S, Ikemoto M, Masuda S, Miyagoe-Suzuki Y, Takeda S (2006) Functional heterogeneity of side population cells in skeletal muscle. *Biochem Biophys Res Commun* 341:864–873
 40. Fiszer D, Seidel M, Siminiak T, Kurpisz M (2007) Future trends: cell engineering for cardiac repair. *EuroIntervention* 2:889–894
 41. Leri A, Kajstura J, Anversa P (2005) Cardiac stem cells and mechanisms of myocardial regeneration. *Physiol Rev* 85:1373–1416
 42. Torella D, Ellison GM, Mendez-Ferrer S, Ibanez B, Nadal-Ginard B (2006) Resident human cardiac stem cells: role in cardiac cellular homeostasis and potential for myocardial regeneration. *Nat Clin Pract Cardiovasc Med* 3(Suppl 1):S8–13
 43. Srivastava D, Ivey KN (2006) Potential of stem-cell-based therapies for heart disease. *Nature* 441:1097–1099
 44. Moretti A, Lam J, Evans SM, Laugwitz KL (2007) Biology of Isl1 + cardiac progenitor cells in development and disease. *Cell Mol Life Sci* 64:674–682
 45. Guan K, Nayernia K, Maier LS, Wagner S, Dressel R, Lee JH, Nolte J, Wolf F, Li M, Engel W, Hasenfuss G (2006) Pluripotency of spermatogonial stem cells from adult mouse testis. *Nature* 440:1199–1203
 46. Passier R, Oostwaard DW, Snapper J, Kloots J, Hassink RJ, Kuijk E, Roelen B, de la Riviere AB, Mummery C (2005) Increased cardiomyocyte differentiation from human embryonic stem cells in serum-free cultures. *Stem Cells* 23:772–780
 47. Levy D, Kenchaiah S, Larson MG, Benjamin EJ, Kupka MJ, Ho KK, Murabito JM, Vasan RS (2002) Long-term trends in the incidence of and survival with heart failure. *N Engl J Med* 347:1397–1402
 48. Menasche P (2006) Randomized, placebo-controlled Myoblast Autologous Grafting in Ischemic Cardiomyopathy (MAGIC) Trial. *AHA Meeting, Chicago, November 11–15*
 49. Laflamme MA, Murry CE (2005) Regenerating the heart. *Nat Biotechnol* 23:845–856
 50. Tambara K, Tabata Y, Komeda M (2004) Factors related to the efficacy of skeletal muscle cell transplantation and future

- approaches with control-released cell growth factors and minimally invasive surgery. *Int J Cardiol* 95(Suppl 1):S13–15
51. Kurpisz M, Czepczynski R, Grygielska B, Majewski M, Fiszler D, Jerzykowska O, Sowinski J, Siminiak T (2007) Bone marrow stem cell imaging after intracoronary administration. *Int J Cardiol* 121:194–195
 52. Askari A, Unzek S, Goldman CK, Ellis SG, Thomas JD, DiCorleto PE, Topol EJ, Penn MS (2004) Cellular, but not direct, adenoviral delivery of vascular endothelial growth factor results in improved left ventricular function and neovascularization in dilated ischemic cardiomyopathy. *J Am Coll Cardiol* 43:1908–1914
 53. Reinecke H, Minami E, Virag JI, Murry CE (2004) Gene transfer of connexin43 into skeletal muscle. *Hum Gene Ther* 15:627–636
 54. Ott HC, Bonaros N, Marksteiner R, Wolf D, Margreiter E, Schachner T, Laufer G, Hering S (2004) Combined transplantation of skeletal myoblasts and bone marrow stem cells for myocardial repair in rats. *Eur J Cardiothorac Surg* 25:627–634
 55. Smith AG (2001) Embryo-derived stem cells: of mice and men. *Annu Rev Cell Dev Biol* 17:435–462
 56. Wagers AJ, Weissman IL (2004) Plasticity of adult stem cells. *Cell* 116:639–648
 57. Ferrari G, Cusella-De Angelis G, Coletta M, Paolucci E, Stornaiuolo A, Cossu G, Mavilio F (1998) Muscle regeneration by bone marrow-derived myogenic progenitors. *Science* 279:1528–1530
 58. Petersen BE, Bowen WC, Patrene KD, Mars WM, Sullivan AK, Murase N, Boggs SS, Greenberger JS, Goff JP (1999) Bone marrow as a potential source of hepatic oval cells. *Science* 284:1168–1170
 59. Lin Y, Weisdorf DJ, Solovey A, Hebbel RP (2000) Origins of circulating endothelial cells and endothelial outgrowth from blood. *J Clin Invest* 105:71–77
 60. Orlic D, Kajstura J, Chimenti S, Limana F, Jakoniuk I, Quaini F, Nadal-Ginard B, Bodine DM, Leri A, Anversa P (2001) Mobilized bone marrow cells repair the infarcted heart, improving function and survival. *Proc Natl Acad Sci USA* 98:10344–10349
 61. Brazelton TR, Rossi FM, Keshet GI, Blau HM (2000) From marrow to brain: expression of neuronal phenotypes in adult mice. *Science* 290:1775–1779
 62. Jackson KA, Majka SM, Wang H, Pocius J, Hartley CJ, Majesky MW, Entman ML, Michael LH, Hirschi KK, Goodell MA (2001) Regeneration of ischemic cardiac muscle and vascular endothelium by adult stem cells. *J Clin Invest* 107:1395–1402
 63. Lagasse E, Connors H, Al-Dhalimy M, Reitsma M, Dohse M, Osborne L, Wang X, Finegold M, Weissman IL, Grompe M (2000) Purified hematopoietic stem cells can differentiate into hepatocytes in vivo. *Nat Med* 6:1229–1234
 64. Krause DS, Theise ND, Collector MI, Henegariu O, Hwang S, Gardner R, Neutzel S, Sharkis SJ (2001) Multi-organ, multi-lineage engraftment by a single bone marrow-derived stem cell. *Cell* 105:369–377
 65. Kopen GC, Prockop DJ, Phinney DG (1999) Marrow stromal cells migrate throughout forebrain and cerebellum, and they differentiate into astrocytes after injection into neonatal mouse brains. *Proc Natl Acad Sci USA* 96:10711–10716
 66. Gussoni E, Soneoka Y, Strickland CD, Buzney EA, Khan MK, Flint AF, Kunkel LM, Mulligan RC (1999) Dystrophin expression in the mdx mouse restored by stem cell transplantation. *Nature* 401:390–394
 67. Pang W (2000) Role of muscle-derived cells in hematopoietic reconstitution of irradiated mice. *Blood* 95:1106–1108
 68. Abedi M, Greer DA, Colvin GA, Demers DA, Dooner MS, Harpel JA, Pimentel J, Menon MK, Quesenberry PJ (2004) Tissue injury in marrow transdifferentiation. *Blood Cells Mol Diseases* 32:42–46
 69. Quesenberry PJ, Colvin G, Dooner G, Dooner M, Aliotta JM, Johnson K (2007) The stem cell continuum: cell cycle, injury, and phenotype lability. *Ann N Y Acad Sci* 1106:20–29
 70. Tanaka EM (2003) Cell differentiation and cell fate during urodele tail and limb regeneration. *Curr Opin Genet Dev* 13:497–501
 71. Odelberg SJ, Kollhoff A, Keating MT (2000) Dedifferentiation of mammalian myotubes induced by *msx1*. *Cell* 103:1099–1109
 72. Tsai RY, Kittappa R, McKay RD (2002) Plasticity, niches, and the use of stem cells. *Dev Cell* 2:707–712
 73. Schnapp E, Tanaka EM (2005) Quantitative evaluation of morpholino-mediated protein knockdown of GFP, *MSX1*, and *PAX7* during tail regeneration in *Ambystoma mexicanum*. *Dev Dyn* 232:162–170
 74. Chen S, Zhang Q, Wu X, Schultz PG, Ding S (2004) Dedifferentiation of lineage-committed cells by a small molecule. *J Am Chem Soc* 126:410–411
 75. Ianus A, Holz GG, Theise ND, Hussain MA (2003) In vivo derivation of glucose-competent pancreatic endocrine cells from bone marrow without evidence of cell fusion. *J Clin Invest* 111:843–850
 76. Jang YY, Sharkis SJ (2004) Metamorphosis from bone marrow derived primitive stem cells to functional liver cells. *Cell Cycle* 3:980–982
 77. Direkze NC, Forbes SJ, Brittan M, Hunt T, Jeffery R, Preston SL, Poulos R, Hodivala-Dilke K, Alison MR, Wright NA (2003) Multiple organ engraftment by bone-marrow-derived myofibroblasts and fibroblasts in bone-marrow-transplanted mice. *Stem Cells* 21:514–520
 78. Corti S, Locatelli F, Donadoni C, Guglieri M, Papadimitriou D, Strazzer S, Del Bo R, Comi GP (2004) Wild-type bone marrow cells ameliorate the phenotype of SOD1-G93A ALS mice and contribute to CNS, heart and skeletal muscle tissues. *Brain* 127:2518–2532
 79. Theise ND, Nimmakayalu M, Gardner R, Illei PB, Morgan G, Teperman L, Henegariu O, Krause DS (2000) Liver from bone marrow in humans. *Hepatology* 32:11–16
 80. Morshead CM, Benveniste P, Iscove NN, van der Kooy D (2002) Hematopoietic competence is a rare property of neural stem cells that may depend on genetic and epigenetic alterations. *Nat Med* 8:268–273
 81. Vassilopoulos G, Russell DW (2003) Cell fusion: an alternative to stem cell plasticity and its therapeutic implications. *Curr Opin Genet Dev* 13:480–485
 82. Wang X, Willenbring H, Akkari Y, Torimaru Y, Foster M, Al-Dhalimy M, Lagasse E, Finegold M, Olson S, Grompe M (2003) Cell fusion is the principal source of bone-marrow-derived hepatocytes. *Nature* 422:897–901
 83. Pomerantz J, Blau HM (2004) Nuclear reprogramming: a key to stem cell function in regenerative medicine. *Nat Cell Biol* 6:810–816
 84. Alvarez-Dolado M, Pardal R, Garcia-Verdugo JM, Fike JR, Lee HO, Pfeffer K, Lois C, Morrison SJ, Alvarez-Buylla A (2003) Fusion of bone-marrow-derived cells with Purkinje neurons, cardiomyocytes and hepatocytes. *Nature* 425:968–973
 85. D'Ippolito G, Diabira S, Howard GA, Menei P, Roos BA, Schiller PC (2004) Marrow-isolated adult multilineage inducible (MIAMI) cells, a unique population of postnatal young and old human cells with extensive expansion and differentiation potential. *J Cell Sci* 117:2971–2981
 86. Kogler G, Sensken S, Airey JA, Trapp T, Muschen M, Feldhahn N, Liedtke S, Sorg RV, Fischer J, Rosenbaum C, Greschat S, Knipper A, Bender J, Degistirici O, Gao J, Caplan AI, Colletti EJ, Almeida-Porada G, Muller HW, Zanjani E, Wernet P (2004) A new human somatic stem cell from placental cord blood with

- intrinsic pluripotent differentiation potential. *J Exp Med* 200:123–135
87. Jiang Y, Jahagirdar BN, Reinhardt RL, Schwartz RE, Keene CD, Ortiz-Gonzalez XR, Reyes M, Lenvik T, Lund T, Blackstad M, Du J, Aldrich S, Lisberg A, Low WC, Largaespada DA, Verfaillie CM (2002) Pluripotency of mesenchymal stem cells derived from adult marrow. *Nature* 418:41–49
 88. Nichols J, Zevnik B, Anastasiadis K, Niwa H, Klewe-Nebenius D, Chambers I, Scholer H, Smith A (1998) Formation of pluripotent stem cells in the mammalian embryo depends on the POU transcription factor Oct4. *Cell* 95:379–391
 89. Boyer LA, Lee TI, Cole MF, Johnstone SE, Levine SS, Zucker JP, Guenther MG, Kumar RM, Murray HL, Jenner RG, Gifford DK, Melton DA, Jaenisch R, Young RA (2005) Core transcriptional regulatory circuitry in human embryonic stem cells. *Cell* 122:947–956
 90. Shizuru JA, Negrin RS, Weissman IL (2005) Hematopoietic stem and progenitor cells: clinical and preclinical regeneration of the hematolymphoid system. *Annu Rev Med* 56:509–538
 91. Tabata M, Satake A, Okura N, Yamazaki Y, Toda A, Nishioka K, Tanaka H, Chin M, Itsukuma T, Yamaguchi M, Misawa M, Kai S, Hara H (2002) Long-term outcome after allogeneic bone marrow transplantation for hematological malignancies with non-remission status. Results of a single-center study of 24 patients. *Ann Hematol* 81:582–587
 92. Iannone R, Casella JF, Fuchs EJ, Chen AR, Jones RJ, Woolfrey A, Amylon M, Sullivan KM, Storb RF, Walters MC (2003) Results of minimally toxic nonmyeloablative transplantation in patients with sickle cell anemia and beta-thalassemia. *Biol Blood Marrow Transplant* 9:519–528
 93. Pai SY, DeMartini D, Forino C, Cavagnini S, Lanfranchi A, Giliiani S, Moratto D, Mazza C, Porta F, Imberti L, Notarangelo LD, Mazzolari E (2006) Stem cell transplantation for the Wiskott–Aldrich syndrome: a single-center experience confirms efficacy of matched unrelated donor transplantation. *Bone Marrow Transplant* 38:671–679
 94. Peters C, Krivit W (2000) Hematopoietic cell transplantation for mucopolysaccharidosis IIB (Hunter syndrome). *Bone Marrow Transplant* 25:1097–1099
 95. Salerno M, Busiello R, Esposito V, Cosentini E, Adriani M, Selleri C, Rotoli B, Pignata C (2004) Allogeneic bone marrow transplantation restores IGF-I production and linear growth in a gamma-SCID patient with abnormal growth hormone receptor signaling. *Bone Marrow Transplant* 33:773–775
 96. Staba SL, Escolar ML, Poe M, Kim Y, Martin PL, Szabolcs P, Allison-Thacker J, Wood S, Wenger DA, Rubinstein P, Hopwood JJ, Krivit W, Kurtzberg J (2004) Cord-blood transplants from unrelated donors in patients with Hurler's syndrome. *N Engl J Med* 350:1960–1969
 97. Ueno NT, Cheng YC, Rondon G, Tannir NM, Gajewski JL, Couriel DR, Hosing C, de Lima MJ, Anderlini P, Khouri IF, Booser DJ, Hortobagyi GN, Pagliaro LC, Jonasch E, Giral SA, Champlin RE (2003) Rapid induction of complete donor chimerism by the use of a reduced-intensity conditioning regimen composed of fludarabine and melphalan in allogeneic stem cell transplantation for metastatic solid tumors. *Blood* 102:3829–3836
 98. Kurokawa T, Fischer K, Bertz H, Hoegerle S, Finke J, Mackensen A (2002) In vitro and in vivo characterization of graft-versus-tumor responses in melanoma patients after allogeneic peripheral blood stem cell transplantation. *Int J Cancer* 101:52–60
 99. Mezey E, Chandross KJ, Harta G, Maki RA, McKercher SR (2000) Turning blood into brain: cells bearing neuronal antigens generated in vivo from bone marrow. *Science* 290:1779–1782
 100. Kucia M, Reza R, Jala VR, Dawn B, Ratajczak J, Ratajczak MZ (2005) Bone marrow as a home of heterogeneous populations of nonhematopoietic stem cells. *Leukemia* 19:1118–1127
 101. Lubovny M, McCune S, Dong JY, Prchal JF, Townes TM, Prchal JT (1996) Stable transduction of recombinant adeno-associated virus into hematopoietic stem cells from normal and sickle cell patients. *Biol Blood Marrow Transplant* 2:24–30
 102. Bradford GB, Williams B, Rossi R, Bertonecello I (1997) Quiescence, cycling, and turnover in the primitive hematopoietic stem cell compartment. *Exp Hematol* 25:445–453
 103. Stiff P, Gingrich R, Luger S, Wyres MR, Brown RA, LeMaistre CF, Perry J, Schenkein DP, List A, Mason JR, Bensinger W, Wheeler C, Freter C, Parker WRL, Emmanouilides C (2000) A randomized phase 2 study of PBPC mobilization by stem cell factor and filgrastim in heavily pretreated patients with Hodgkin's disease or non-Hodgkin's lymphoma. *Bone Marrow Transplant* 26:471–481
 104. Booy EP, Johar D, Maddika S, Pirzada H, Sahib MM, Gehrke I, Loewen SD, Louis SD, Kadkhoda K, Mowat M, Los M (2006) Monoclonal and bispecific antibodies as novel therapeutics. *Arch Immunol Ther Exp* 54:1–17
 105. Krzemieniecki K, Szpyt E, Rashedi I, Gawron K, Los M (2006) Targeting of solid tumors and blood malignancies by antibody-based therapies. *Centr Eur J Biol* 1:167–182
 106. Anderson JE, Hansen LL, Mooren FC, Post M, Hug H, Zuse A, Los M (2006) Methods and biomarkers for the diagnosis and prognosis of cancer and other diseases: towards personalized medicine. *Drug Resist Updat* 9:198–210
 107. Krocak TJ, Baran J, Pryjma J, Siedlar M, Rashedi I, Hernandez E, Alberti EMS, Los M (2006) The emerging importance of DNA mapping and other comprehensive screening techniques as tools to identify new drug targets and as a mean of (cancer) therapy personalization. *Expert Opin Ther Targets* 10:289–302
 108. Ghavami S, Asoodeh A, Klonisch T, Halayko AJ, Kadkhoda K, Krocak TJ, Gibson SB, Booy EP, Naderi-Manesh H, Los M (2008) Brevinin-2R(1) semi-selectively kills cancer cells by a distinct mechanism, which involves the lysosomal-mitochondrial death pathway. *J Cell Mol Med* 12:1005–1022
 109. Maddika S, Bay GH, Krocak TJ, Ande SR, Maddika S, Wiechec E, Gibson SB, Los M (2007) Akt is transferred to the nucleus of cells treated with apoptin, and it participates in apoptin-induced cell death. *Cell Prolif* 40:835–848
 110. Maddika S, Mendoza FJ, Hauff K, Zamzow CR, Paranjothy T, Los M (2006) Cancer-selective therapy of the future: apoptin and its mechanism of action. *Cancer Biol Ther* 5:10–19
 111. Maddika S, Ande SR, Panigrahi S, Paranjothy T, Weglarczyk K, Zuse A, Eshraghi M, Manda KD, Wiechec E, Los M (2007) Cell survival, cell death and cell cycle pathways are interconnected: implications for cancer therapy. *Drug Resist Updat* 10:13–29
 112. Johnston JB, Navaratnam S, Pitz MW, Maniate JM, Wiechec E, Baust H, Gingerich J, Skliris GP, Murphy LC, Los M (2006) Targeting the EGFR pathway for cancer therapy. *Curr Med Chem* 13:3483–3492
 113. Zuse A, Prinz H, Muller K, Schmidt P, Gunther EG, Schweizer F, Prehn JH, Los M (2007) 9-Benzylidene-naphtho[2,3-b]thiophen-4-ones and benzylidene-9(10H)-anthracenones as novel tubulin interacting agents with high apoptosis-inducing activity. *Eur J Pharmacol* 575:34–45
 114. Alexander HK, Booy EP, Xiao W, Ezzati P, Baust H, Los M (2007) Selected technologies to control genes and their products for experimental and clinical purposes. *Arch Immunol Ther Exp* 55:139–149
 115. Rafii S, Lyden D (2003) Therapeutic stem and progenitor cell transplantation for organ vascularization and regeneration. *Nat Med* 9:702–712

116. Peichev M, Naiyer AJ, Pereira D, Zhu Z, Lane WJ, Williams M, Oz MC, Hicklin DJ, Witte L, Moore MA, Rafii S (2000) Expression of VEGFR-2 and AC133 by circulating human CD34 (+) cells identifies a population of functional endothelial precursors. *Blood* 95:952–958
117. Takahashi H, Shibuya M (2005) The vascular endothelial growth factor (VEGF)/VEGF receptor system and its role under physiological and pathological conditions. *Clin Sci (Lond)* 109:227–241
118. Yamamoto Y, Yasumizu R, Amou Y, Watanabe N, Nishio N, Toki J, Fukuhara S, Ikehara S (1996) Characterization of peripheral blood stem cells in mice. *Blood* 88:445–454
119. Pittenger MF, Mackay AM, Beck SC, Jaiswal RK, Douglas R, Mosca JD, Moorman MA, Simonetti DW, Craig S, Marshak DR (1999) Multilineage potential of adult human mesenchymal stem cells. *Science* 284:143–147
120. Rege TA, Hagood JS (2006) Thy-1, a versatile modulator of signaling affecting cellular adhesion, proliferation, survival, and cytokine/growth factor responses. *Biochim Biophys Acta* 1763:991–999
121. Rietze RL, Valcanis H, Brooker GF, Thomas T, Voss AK, Bartlett PF (2001) Purification of a pluripotent neural stem cell from the adult mouse brain. *Nature* 412:736–739
122. Bonfanti L (2006) PSA-NCAM in mammalian structural plasticity and neurogenesis. *Prog Neurobiol* 80:129–164
123. Shackleton M, Vaillant F, Simpson KJ, Stingl J, Smyth GK, Asselin-Labat ML, Wu L, Lindeman GJ, Visvader JE (2006) Generation of a functional mammary gland from a single stem cell. *Nature* 439:84–88
124. Welm BE, Tepera SB, Venezia T, Graubert TA, Rosen JM, Goodell MA (2002) Sca-1(pos) cells in the mouse mammary gland represent an enriched progenitor cell population. *Dev Biol* 245:42–56
125. Roskams T (2006) Liver stem cells and their implication in hepatocellular and cholangiocarcinoma. *Oncogene* 25:3818–3822
126. Burke ZD, Thowfeeu S, Peran M, Tosh D (2007) Stem cells in the adult pancreas and liver. *Biochem J* 404:169–178
127. Menasche P, Hagege AA, Scorsin M, Pouzet B, Desnos M, Duboc D, Schwartz K, Vilquin JT, Marolleau JP (2001) Myoblast transplantation for heart failure. *Lancet* 357:279–280
128. Pagani FD, DerSimonian H, Zawadzka A, Wetzel K, Edge AS, Jacoby DB, Dinsmore JH, Wright S, Aretz TH, Eisen HJ, Aaronson KD (2003) Autologous skeletal myoblasts transplanted to ischemia-damaged myocardium in humans. Histological analysis of cell survival and differentiation. *J Am Coll Cardiol* 41:879–888
129. Siminiak T, Kalawski R, Fiszer D, Jerzykowska O, Rzezniczak J, Rozwadowska N, Kurpisz M (2004) Autologous skeletal myoblast transplantation for the treatment of postinfarction myocardial injury: phase I clinical study with 12 months of follow-up. *Am Heart J* 148:531–537
130. Siminiak T, Fiszer D, Jerzykowska O, Grygielska B, Rozwadowska N, Kalmucki P, Kurpisz M (2005) Percutaneous trans-coronary-venous transplantation of autologous skeletal myoblasts in the treatment of post-infarction myocardial contractility impairment: the POZNAN trial. *Eur Heart J* 26:1188–1195
131. Chachques JC, Herreros J, Trainini J, Juffe A, Rendal E, Prosper F, Genovese J (2004) Autologous human serum for cell culture avoids the implantation of cardioverter-defibrillators in cellular cardiomyoplasty. *Int J Cardiol* 95(Suppl 1):S29–33
132. Herreros J, Prosper F, Perez A, Gavira JJ, Garcia-Velloso MJ, Barba J, Sanchez PL, Canizo C, Rabago G, Marti-Climent JM, Hernandez M, Lopez-Holgado N, Gonzalez-Santos JM, Martin-Luengo C, Alegria E (2003) Autologous intramyocardial injection of cultured skeletal muscle-derived stem cells in patients with non-acute myocardial infarction. *Eur Heart J* 24:2012–2020
133. Smits PC, van Geuns RJ, Poldermans D, Bountiokos M, Onderwater EE, Lee CH, Maat AP, Serruys PW (2003) Catheter-based intramyocardial injection of autologous skeletal myoblasts as a primary treatment of ischemic heart failure: clinical experience with six-month follow-up. *J Am Coll Cardiol* 42:2063–2069
134. Dib N, McCarthy P, Campbell A, Yeager M, Pagani FD, Wright S, MacLellan WR, Fonarow G, Eisen HJ, Michler RE, Binkley P, Buchele D, Korn R, Ghazoul M, Dinsmore J, Opie SR, Diethrich E (2005) Feasibility and safety of autologous myoblast transplantation in patients with ischemic cardiomyopathy. *Cell Transplant* 14:11–19
135. Ince H, Petzsch M, Rehders TC, Chatterjee T, Nienaber CA (2004) Transcatheter transplantation of autologous skeletal myoblasts in postinfarction patients with severe left ventricular dysfunction. *J Endovasc Ther* 11:695–704
136. Law PK, Fang G, Chua F (2003) First-in man myoblast allografts for heart degeneration. *Int J Med Implants Devices* 1:100–155