

Rapid linkage disequilibrium decay in the *Lr10* gene in wild emmer wheat (*Triticum dicoccoides*) populations

Hanan Sela · Caroline Loutre · Beat Keller ·
Alan Schulman · Eviatar Nevo · Abraham Korol ·
Tzion Fahima

Received: 11 February 2010 / Accepted: 25 August 2010 / Published online: 22 September 2010
© Springer-Verlag 2010

Abstract

Introduction Recombination is a key evolutionary factor enhancing diversity. However, the effect of recombination on diversity in inbreeding species is expected to be low. To estimate this effect, recombination and diversity patterns of *Lr10* gene were studied in natural populations of the inbreeder species, wild emmer wheat (*Triticum dicoccoides*). Wild emmer wheat is the progenitor of most cultivated wheats and it harbors rich genetic resources for disease resistance. *Lr10* is a leaf rust resistance gene encoding three domains: a coiled-coil, nucleotide-binding site, and leucine-rich repeat (CC-NBS-LRR).

Results *Lr10* was sequenced from 58 accessions representing 12 diverse habitats in Israel. Diversity analysis

revealed a high rate of synonymous and non-synonymous substitutions ($d_S = 0.029$, $d_N = 0.018$, respectively) in the NBS-LRR domains. Moreover, in contrast to other resistance genes, in *Lr10* the CC domain was more diverse than the NBS-LRR domains ($d_S = 0.069$ vs. 0.029 , $d_N = 0.094$ vs. 0.018) and was subjected to positive selection in some of the populations. Seventeen recombination events were detected between haplotypes, especially in the CC domain. Linkage disequilibrium (LD) analysis has shown a rapid decay from $r^2 = 0.5$ to $r^2 = 0.1$ within a 2-kb span.

Conclusion These results suggest that recombination is a diversifying force for the *R*-gene, *Lr10*, in the selfing species *T. dicoccoides*. This is the first report of a short-range LD decay in wild emmer wheat.

Communicated by R. Waugh.

Electronic supplementary material The online version of this article (doi:10.1007/s00122-010-1434-2) contains supplementary material, which is available to authorized users.

H. Sela · E. Nevo · A. Korol · T. Fahima (✉)
Department of Evolutionary and Environmental Biology,
Institute of Evolution, Faculty of Natural Sciences,
University of Haifa, Mt. Carmel, Haifa 31905, Israel
e-mail: fahima@research.haifa.ac.il

C. Loutre · B. Keller
Institute of Plant Biology, University of Zürich,
Zollikerstrasse, 107, 8008 Zürich, Switzerland

A. Schulman
MTT/BI Plant Genomics Laboratory,
Institute of Biotechnology, University of Helsinki,
P.O. Box 65, 00014 Helsinki, Finland

A. Schulman
Biotechnology and Food Research,
MTT Agrifood Research Finland, Myllytie 1,
31600 Jokioinen, Finland

Introduction

Recombination is a major mechanism underlying genetic diversity (Korol et al. 1994; Gaut et al. 2007; Wicker et al. 2007; Morrell et al. 2006). Tracking recombination events can illuminate the evolution of genes and genomes. Recombination and linkage disequilibrium (LD) are basic tools in genetic mapping and positional cloning of genes in plant genomes (Ersoz et al. 2008; Waugh et al. 2009). Unlike linkage studies of bi-parental mapping populations, in wild populations LD patterns depend on a long history of recombination and therefore enable to obtain improved mapping resolution. On the other hand, LD is affected by population structure and selection, and hence linkage between markers may not necessarily reflect physical proximity (Flint-Garcia et al. 2003; Rostoks et al. 2006). The extent of LD in barley, a selfing plant and a relative of wheat, was found to be much lower in wild populations (*Hordeum spontaneum*) than in a cultivated collection (*H. vulgare*). LD rapidly

decays over the first kb in the wild forms but extends over more than 200 kb in elite cultivars (Caldwell et al. 2006). In durum wheat (*Triticum durum*), LD stretches for a long distance (Maccaferri et al. 2005), with high LD values found between loci located 50 cM apart. In bread wheat (*T. aestivum*) collections, the LD extent ranged from 1 cM to 5 cM (Chao et al. 2007; Horvath et al. 2009; Somers et al. 2007). In *Aegilops tauschii*, the wild donor of the D genome to *T. aestivum*, a 1,400-bp LD block was observed in the grain softness locus (Massa and Morris 2006). In wild emmer wheat (*T. dicoccoides*) populations from Amiad and Tabigha in northern Israel, high LD values were observed even between loci on different chromosomes. It seems that LD values in *T. dicoccoides* populations are inflated by the population structure (Li et al. 2000a, b). To date, only one study has targeted the intra-genic LD in *T. dicoccoides* populations and only two studies targeted LD within *R*-genes in selfers (Mauricio et al. 2003; Kuang et al. 2008; Haudry et al. 2007). Therefore, the current study may add insights into LD patterns in *T. dicoccoides* populations and LD patterns in *R*-genes.

Leaf rust resistance gene *Lr10* is one of the few cloned disease resistance genes (*R*-genes) in bread wheat. *Lr10* encodes a three domain protein: coiled-coil, nucleotide-binding site and leucine-rich repeats (CC-NBS-LRR) (Feuillet et al. 2003). *Lr10* is located 36 cM from the centromere on the short arm of chromosome 1A. Previous studies have shown that *Lr10* is a very variable (polymorphic) gene while the CC domain is its most diverse part. The high diversity in the CC domain is probably due to positive selection (Loutre et al. 2009). Many plant *R*-genes exhibit high diversity, positive selection, and sequence exchanges since the arms race with pathogens drives them to change recognition specificity frequently (Wicker et al. 2007; Bergelson et al. 2001; Mondragon-Palomino and Gaut 2005). In some *R*-genes, diversity is maintained by frequency-dependent selection where alleles are cycled between high and low frequency in a long co-evolution with pathogens as a form of “trench warfare”. The trench warfare model predicts that the polymorphism will be old and under balancing selection (Holub 2001; Stahl et al. 1999). Although many studies of *R*-gene diversity were conducted [reviewed in McDowell and Simon (2006)], most of them were limited to a few alleles or to paralogs and only a few studies focused on population genetics (Mauricio et al. 2003; Meaux and Neema 2003; Butterbach 2007; Yahiaoui et al. 2009; Kuang et al. 2008). These studies mainly targeted the LRR domain, which is considered the most diverse domain due to its interaction with pathogen effectors. However, in *Lr10*, the CC domain was found more diverse than the LRR domain (Loutre et al. 2009). Therefore, in the current study, the CC domain was analyzed separately from the NBS and LRR domains.

T. dicoccoides is the tetraploid (genome BBAA) progenitor of most cultivated wheats. The natural habitats of *T. dicoccoides* are located in the Middle East Fertile Crescent. *T. dicoccoides* populations harbor a valuable pool of resistance genes that can be transferred into cultivated wheat (Rong et al. 2000; Knott et al. 2005; Mergoum et al. 2005; Uauy et al. 2005; Marais et al. 2005; Gustafson et al. 2009). Numerous studies of *T. dicoccoides* genetic diversity have shown that the population in Israel is sub-divided. The selfing nature of emmer wheat and the large difference between *T. dicoccoides* habitats in Israel largely contribute to this pattern (Golenberg and Nevo 1987; Nevo et al. 2002; Peleg et al. 2008b; Sela et al. 2009).

Leaf rust, caused by *Puccinia triticina*, is the most common rust disease of wheat. It is highly specific and can spread thousands of kilometers by wind (Bolton et al. 2008). In Israel, the pathogen cannot complete its life cycle due to the absence of an alternate host and harsh summer conditions. It is assumed that new inocula of *P. triticina* arrive every year from regions adjacent to Israel (Y. Anikster, personal communication). Most of *T. dicoccoides* accessions from Israel are susceptible to leaf rust and only a small fraction of the accessions are resistant (Anikster et al. 2005; Moseman et al. 1985). Nevertheless, leaf rust resistance genes from *T. dicoccoides* were introgressed into cultivated wheat (Marais et al. 2005).

The main objectives of the present study were the following: (1) to estimate *Lr10* nucleotide diversity among emmer wheat populations originating from Israel; (2) test the differences in *Lr10* nucleotide diversity between and within *T. dicoccoides* populations; (3) estimate the extent of LD along the gene; (4) study the effect of recombination on sequence diversity.

Materials and methods

One hundred *T. dicoccoides* accessions from 12 populations in Israel and vicinity were used in the current study (Table 1; Fig. 1). A full description of their geographic origin and climatic conditions can be found in Nevo and Beiles (1989). DNA was extracted from 10-day-old seedlings using the ArchivePure DNA extraction kit (5 Prime). The whole *Lr10* gene (4 kb) was amplified using Pfu Ultra Fusion II HS polymerase (Stratagene) according to the supplier's protocol (annealing temperature was 53°C and elongation time 1 min) with the primers ThLR10_V (CGGA ACTATGGAGAGTGAAC) and ThLR10_U (GGGAAATGTAGACAGGTACAT) (Feuillet et al. 2003). PCR products were separated on agarose gels and extracted using MinElute gel extraction kits (Qiagen). The extracted DNA was cloned into the pSMART vector using a GC cloning kit (Lucigen). Samples were sequenced using Big

Table 1 List of wild emmer wheat accessions used in the study

Population	Population no ^a	No of Accessions	Accessions
Mt. Hermon	1	6	1-22, 1-24, 1-42, 1-51, 1-59, 1-64
Gamla	8	4	8-22, 8-28, 8-35, 8-40
Rosh Pinna	9	5	9-29, 9-3, 9-33, 9-45, 9-55
Amiad	10	1	10-85
Tabigha	15	3	13-53, 15-25, 15-6
Mt. Gilboa	16	4	16-28, 16-33, 16-34, 16-51
Gitit	18	8	18-1, 18-16, 18-20, 18-25, 18-27, 18-39, 18-4, 18-44
Kokhav-Hashahar	19	4	19-10, 19-13, 19-16, 19-5
Jaba	23	4	23-24, 23-3, 23-45, 23-6,
Amirim	24	5	24-12, 24-13, 24-36, 24-46, 24-47
Achihud	26	1	26-22
Bat-Shelomo	30	2	30-18, 30-4
Beit-Oren	28	5	28-14, 28-2, 28-30, 28-33, 28-53
Givat-Koach	33	4	33-58, 33-13, 33-19, 33-48
Burj el Maliech ^b		1	Be-1
Ma'ale Merav ^b		1	Me-2
<i>T. urartu</i> ^c		4	TU17119, TU17122, TU17159, TU6735
<i>T. aestivum</i> ^d		1	AY2701574

^a Numbers are according to Nevo and Beiles (1989)

^b *T. dicoccoides* accessions from Peleg et al. (2008b)

^c *T. urartu* accessions described in Loutre et al. (2009)

^d GenBank accession from Feuillet et al. (2003)




Fig. 1 Geographic distribution of the 12 populations of wild emmer wheat, *T. dicoccoides*, tested in this study. For names of the numbered populations see list in Table 1. The numbers of populations are according to Nevo and Beiles (1989)

Dye Terminator chemistry on ABI 3700 or ABI 3730 instruments (Applied Biosystems). Alternatively, eight PCR fragments covering the whole gene region were

extracted from the gel and sequenced directly (Table 2). Sequence reads were checked and assembled using Phred, Phrap, and Consed software provided by B. Ewing, P. Green & D. Gordon (available at <http://www.phrap.org/>). The sequences were aligned using MUSCLE (Edgar 2004) and manually corrected using BioEdit (Hall 2007).

Statistical analysis

DNA sequence alignments were analyzed for nucleotide diversity of synonymous (d_S) and non-synonymous substitutions (d_N) (Nei 1987) estimated by Kumar's method as implemented in MEGA software (Kumar et al. 2004). Z tests for selection based on the difference between synonymous and non-synonymous substitutions ($d_S - d_N$) were conducted using MEGA software as well (Nei and Kumar 2000). A consensus maximum parsimony (MP) tree from 500 bootstrapped trees was constructed using MEGA. Pairwise distances between all sequences were calculated using MEGA. Estimations of evolutionary distances were done using the maximum composite likelihood (Tamura and Kumar 2002). The rate of variation among sites was modeled with a gamma distribution (shape parameter = 1). The differences in the composition bias among sequences were considered in evolutionary comparisons (Tamura et al. 2004). The evolutionary distances were calculated separately for the CC domain and for the NBS-LRR domains. Based on the distances, a principal coordinates analysis (PCO) was conducted using GENALEX (Peakall and Smouse 2006). Raw precipitation data for each population

Table 2 Primers used for amplification and sequencing of overlapping fragments of *Lr10* sequence

Primer name		Size (bp)	Start position*
LR10A2F	ATGCATCTTTACCCGTCCAC	840	-100
LR10A4R	TGAGCGTGAATTGAGACAG		
LR10S1F	GACGACTCTTGCCAAACAGG	831	610
LR10S1R	TGTCAAATTCCTTTTCATCG		
LR10b2F	CTACGTCCGTTATTCAAAGAT	1,500	844
LR10b2R	ACCAAATACTCGGTTCAAAA		
LR10T1F	TGGGATGTTTTGAAATGTGC	800	2,024
LR10T2R	AGAAGCACCGAGAGATCGAG		
LR10U1F	GGCACGGTTCTACAGAGTCC	836	2,629
LR10U1R	AAGCAGCCATTTGTTTCATCC		
LR10V2F	ATGAGCTGGGCAAATTAACG	750	3,195
LR10V2R	GCCATTTTGTCACTCATCCTG		

*Start position is relative to the start codon

in the past 20 years was obtained from the Israeli Meteorology Service (Beit-Dagan, Israel) and summarized by SPSS (SPSS Inc.). Monthly and annual means and the variability of rain in the sites of collection of the populations (Table S1) were tested for correlations with diversity indices. Variability of rain was calculated as standard deviation/annual mean.

Linkage disequilibrium (LD) coefficients $-r^2$ and p values were calculated in TASSEL (Bradbury et al. 2007). For convenience, p values were transformed into $-\log p$ ($-\log P$) values. Data were filtered for SNPs with more than 20% frequency of the minor allele. In order to minimize the effect of population structure, a subset of 20 sequences that represents major branches on the MP tree was selected (Fig. S2). This subset was analyzed and compared with the whole data set. Regressions and correlations were analyzed in SPSS (SPSS Inc.). Recombination events in the sequence alignment were detected with RDP software that combines seven different recombination detection methods (Martin et al. 2005). Recombination events were confirmed when at least three methods showed $p < 0.01$ for the event. The population recombination parameter ρ (Hudson 2001) and the population mutation parameter θ (Watterson 1975) were estimated using “rhopmap” software in the LDhat package (Auton and McVean 2007). All the 33 haplotypes revealed in the alignment were used in the analysis. All polymorphic sites were included for the calculation of θ . Only sites with two alleles were used for the estimation of ρ . The run parameters were: 1,100,000 iterations, 100,000 iterations of burn-in period and sampling interval at every 100 iterations. Illegitimate recombination events were detected using Dotter (Sonnhammer and Durbin 1995) and Bioedit (Hall 2007) following the pattern described in Wicker et al. (2007). Illegitimate recombination is a process of asymmet-

ric pairing of two dispersed homologous sequences of only a few-bp-length followed by sequence exchange that can result in either duplications or deletions (Devos et al. 2002). Two tests were implemented to test for deviation from neutrality of the haplotypes that resulted from illegitimate recombination. The Slatkin exact test (Slatkin 1994) was conducted using Arlequin software (Excoffier et al. 2005), and Tajima D test (Tajima 1989) was carried out in DNAsp (Rozas et al. 2003).

Resistance tests

Ten-day-old seedling of all *T. dicoccoides* accessions were tested for leaf rust resistance as described by Schachermayr et al. (1995) using isolate BRW97512-19, which is avirulent to the *Lr10* resistance gene in bread wheat (*T. aestivum*). Seedlings were sprayed with the rust isolate suspended in Soltrol 170 oil (Chevron Phillips Chemical) and incubated overnight at high humidity (90%) and low temperature (16°C). Visual scoring of the phenotypes were performed 10 days after infection on a 0–4 scale (0 = resistant; 4 = susceptible) (McIntosh et al. 1995).

Results

A total of 100 accessions of *T. dicoccoides* from 12 populations were tested for the presence of *Lr10* (Table 1). In 95 accessions *Lr10* was present as determined by PCR amplification of at least three primer pairs listed in Table 2. The full-length 4 kb *Lr10* from 58 accessions was cloned and sequenced. First, reads were assembled into contigs for each accession and then DNA and protein sequence alignments were generated. The sequences were deposited in GenBank (accessions GU393247-GU393304). Four accessions, originating from two populations (Jaba and Givat-Koach), had a 1.2 kb deletion of the NBS domain and therefore were excluded from the diversity analysis. Three sequences obtained from Mt. Hermon accessions had a premature stop codon at position 130.

Sequence diversity

Out of 54 *Lr10* sequences, 33 different haplotypes were detected in the alignment; 332 sites out of 4,114 sites were polymorphic, and the nucleotide diversity (π) was 0.029. Within-population nucleotide diversity for coding regions was calculated for nine populations that were represented by at least three accessions in the alignment. Previous studies have shown that the CC domain is the most diverse region of the *Lr10* (Feuillet et al. 2003; Loutre et al. 2009). Therefore, the analysis was conducted on the

CC domain and on the NBS-LRR domains separately. Diversity values for synonymous and non-synonymous substitution rate (d_S and d_N , respectively) of the NBS-LRR domains differed significantly between populations ($p = 0.04$) (Table 3). The highest diversity was observed in the population from Kokhav-Hashahar ($d_S = 0.036$, $d_N = 0.021$) while the lowest diversity was observed in the population from Beit-Oren ($d_S = 0$, $d_N = 0$; Table 3). The overall d_S value was 0.0295 while d_N value was only 0.0182. The difference $d_S - d_N$ was mainly positive and varied between populations, ranging from -0.0003 in the Gamla population to 0.0154 in the Kokhav-Hashahar population. Z tests for purifying selection based on $d_S - d_N$ values revealed significant purifying selection in the populations of Tabigha, Mt. Gilboa, Gitit and Kokhav-Hashahar ($p < 0.05$) (Table 3). The mean d_S value of these populations was significantly higher than the mean d_S value of the rest of the populations ($d_S = 0.030$ and 0.008, respectively, $p = 0.016$), while no significant difference was observed for d_N values ($d_N = 0.017$ and 0.006, respectively; $p = 0.11$). The population diversity values, d_S and d_N , were tested for correlation with previously obtained simple sequence repeat marker (SSR) gene diversity (He) values of the same populations (Peleg et al. 2008b) (Table S1). d_S and d_N were positively correlated with He values ($r = 0.73$, $p = 0.027$ and $r = 0.76$, $p = 0.021$, respectively) (Fig. 2a, b). In the CC domain, mean population d_S and d_N values were positively correlated with the NBS-LRR d_S

and d_N values ($r = 0.81$, $p = 0.008$ and $r = 0.80$, $p = 0.048$, respectively). The overall d_S value of the CC domain was 0.069, which is more than twofold higher than the NBS-LRR d_S value (0.0295), and the overall d_N value for the CC domain was 0.094, which is fivefold higher than NBS-LRR d_N value (0.0182). In contrast to NBS-LRR values, $d_S - d_N$ values of the CC domain were negative. The overall $d_S - d_N$ value was -0.025 and ranged among the populations from -0.05 in Kokhav-Hashahar population to 0 in Beit-Oren population. In four populations (Mt. Hermon, Mt. Gilboa, Kokhav-Hashahar and Amirim) a significant positive selection was detected.

Climatic factors associated with diversity

The associations between climatic factors and selection have revealed that populations with putative purifying selection in the NBS-LRR domains are associated with low precipitation and high rain variability. In the sites where those populations were collected, the annual precipitation ranged from 375 mm to 432 mm and rain variation ranged from 0.34 to 0.46, while in the other sites the ranges were 535–1,300 mm and 0.29–0.32, respectively (Table S1). NBS-LRR d_S and $d_S - d_N$ values of the populations were positively correlated with rain variation in the collection sites (Spearman rank correlation $r_s = 0.79$, $p = 0.011$ and $r_s = 0.84$, $p = 0.005$, respectively) (Fig. 2c, d).

Fig. 2 Correlations of diversity indices and rain variability.

a Correlation of d_N values of the populations with SSR He (gene diversity) of the same populations (Peleg et al. 2008b).

b Correlation of d_S values of the populations with SSR He .

c Correlation of $d_S - d_N$ values of the populations with rain variability (SD of annual precipitation/mean annual precipitation).

d Correlation of d_S values of the populations with rain variability




Table 3 Nucleotide diversity values d_N and d_S of *T. dicoccoides* populations

Pop no ^a	Population	<i>N</i>	No of haplotypes	NBS-LRR				CC			
				d_S	d_N	d_S-d_N	Sig ^b	d_S	d_N	d_S-d_N	Sig ^c
1	Mt. Hermon	6	4	0.0262	0.0202	0.003		0.0400	0.0650	-0.030	*
8	Gamla	4	2	0.0000	0.0003	-0.0003		0.0000	0.0000	0.000	
9	Rosh-Pinna	5	2	0.0124	0.0111	0.001		0.0390	0.0450	-0.006	
15	Tabigha	3	3	0.0342	0.0189	0.015	*	0.0540	0.0630	-0.008	
16	Mt. Gilboa	4	2	0.0240	0.0137	0.010	*	0.0310	0.0650	-0.030	*
18	Gitit	8	4	0.0291	0.0173	0.011	*	0.0670	0.0820	-0.018	
19	Kokhav-Hashahar	4	3	0.0363	0.0210	0.015	*	0.0560	0.1040	-0.050	*
24	Amirim	5	2	0.0029	0.0015	0.001		0.0400	0.0740	-0.030	*
28	Beit-Oren	5	1	0.0000	0.0000	0.000		0.0000	0.0000	0.000	
	Mean			0.0230	0.0200	0.010		0.0400	0.0570	-0.017	
	Overall	44	15	0.0295	0.0182	0.010	*	0.0690	0.0940	-0.025	

^a Numbers of populations are according to Nevo and Belis (1989)

^b NBS-LRR Sig: populations with significant purifying selection in the NBS-LRR domains ($p < 0.05$)

^c CC Sig: populations with significant positive selection in the CC domain ($p < 0.05$)

Principal coordinates analysis PCO

Based on the sequence distances of the NBS-LRR domains and the CC domain, two PCO graphs were plotted (Fig. 3a, b, respectively). Sequences that had the same coordinates were considered as one haplotype. In the plots, many haplotypes were represented by two or more genotypes of the same population and/or genotypes from more than one population. Nevertheless, in most of the cases, populations represented by the same haplotype did not share similar geographic or climatic characteristics. Moreover, different haplotypes of the same population were scattered on the plot and not clustered together. Four *T. urartu* sequences

(Loutre et al. 2009) that were included in the plot did not cluster together. The PCO plots of the CC and the NBS-LRR domains showed different patterns. In the CC plot, haplotypes were scattered all over the plot with some clustering but with no obvious eco-geographic relations between the clustered populations. In the NBS-LRR plot, a clear division into two groups was observed. Seven populations and the *T. urartu* accessions were represented in both groups, while four populations were represented by only two haplotypes that were almost identical. The main difference in the two groups of sequences lies in the LRR domain where two well conserved, ~400-bp-long, haplotypes exist. These haplotypes diverge from one another. Tajima's test




Fig. 3 Principal coordinates analysis (PCO) based on the genetic distances between *Lr10* sequences of *T. dicoccoides* accessions from Israel. **a** PCO based on NBS-LRR domains. **b** PCO based on CC domain. Diamonds represent position of haplotypes. Numbers near the diamonds are the populations represented in each haplotype x number

of accessions represented. Population numbers are according to Table 1. TA is ThatcherLr10, the first cloned *Lr10* from *T. aestivum* cv. ThatcherLr10 (Feuillet et al. 2003). TU are *T. urartu* accessions (Table 1). Arrows connect the most frequent haplotypes to their position

(Tajima 1989) has shown that in a stretch of 114 bp sequence within these haplotypes there is a deviation from neutrality, suggesting that balancing selection was involved in shaping the diversity ($D = 2.46$, $p < 0.05$).

Linkage disequilibrium analysis

A previous analysis of population structure showed that *T. dicoccoides* populations are well structured (Peleg et al. 2008a). Therefore, in order to avoid the bias of the population structure on LD, a subset of 20 sequences was selected from the MP phylogenetic tree (Fig. S2). LD values ($-\log P$ and r^2) between all pairwise comparisons of polymorphic sites were plotted against the physical distance (bp) between sites in the whole set and in the selected subset (Fig. 4). For the whole set, locally weighted scatterplot smoothing (LOESS) curves of the association values, $-\log P$ and r^2 , declined within 3 and 2 kb, respectively (Fig. 4a, b). For the subset, the decline was much steeper with sharp LD decay found within only 1 kb. The differences in LD decay

patterns between the whole set and the subset were more profound for $-\log P$ values than for r^2 values. Curve estimation of regression models showed that the logarithmic model is the best one to describe the relationships between LD values and physical distance. LD- r^2 values of the whole set and the subset were correlated with distance ($R^2 = 0.40$ and $R^2 = 0.41$, respectively, $p < 0.001$), while $-\log P$ values were correlated for the subset ($R^2 = 0.40$, $p < 0.001$) but much less for the whole set ($R^2 = 0.16$, $p < 0.001$). LD plots along the gene showed the same pattern (Fig. 5). In the plot of the whole set, low p values were found away from the diagonal line, indicating associations that are not a result of physical proximity. This pattern was not noted either for p values of the subset or for r^2 values of the whole set and the subset. In these plots, small LD blocks (<250 bp) were observed along the diagonal with the exception of a larger block of ~ 500 bp at the 3' end of the gene. This block lies in the LRR domain in the region where the two well-conserved haplotypes mentioned earlier were observed.

Recombination detection analysis implemented in RDP revealed 17 highly significant recombination events





Fig. 4 LD values of pairwise comparisons between polymorphic sites plotted against their physical distances (bp). LOESS curves are in blue, logarithmic regression curves are in red. **a, b** $-\log P$ and r^2 values

calculated for the whole dataset (60 accessions), respectively. **c, d** $-\log P$ and r^2 values of the subset (20 selected accessions), respectively

Fig. 5 LD plot of p and r^2 values of pairwise comparisons between polymorphic sites. **a** Values for the whole set. **b** Values for the subset. Upper triangles are r^2 values; lower triangles are p values. Only polymorphic sites ($f > 0.2$) are represented



(confirmed by at least three detection methods ($p < 0.01$). These events resulted in 22 different combinations of recombinant fragments (Fig. 6). The distribution pattern of breakpoints was different between the CC domain and other domains. While in the CC domain most of the break points were found within the domain, in the NBS and LRR domains most of the breakpoints were located near the borders of the domain. To test the significance of this observation, the distance of the breakpoints to the nearest border was calculated and transformed to a relative distance as a fraction of the domain size. The distances in the CC domain were significantly higher than in the NBS-LRR domains (mean 0.6 and 0.25, respectively; $p = 0.004$). A significant difference ($p = 0.001$) was found in the length of the recombinant fragments between the domains. In the CC domain fragments were short, ranging from 23 to 361 bp (mean 177 bp) while in the NBS and LRR domains fragments were longer, ranging from 349 to 3,020 bp (mean 1,246 bp). Only one breakpoint was found in the intron region. All events occurred between sequences obtained from different populations and were detected in 2–30 pairs of sequences (mean 11). The estimation of population recombination parameter rho (ρ) was 0.012 per base pair and the estimation of population mutation parameter, theta (θ), was 0.021 per base pair. The ratio between recombination and mutation rates (ρ/θ) was 0.59.

Ten illegitimate recombination sites were detected in the alignment. Three events have resulted in deletions of 20–26 bp in the intron region and four more resulted in trinucleotide deletions that were scattered along the gene. These deletions were present in 2–32 of the sequences. One repeat of 81 bp in the CC domain was found only in one sequence. Another repeat of 32 bp was found in the intron region of all sequences. This repeat was about 40% degenerated and a subsequent deletion was nested within it indicating that this region is prone to illegitimate recombination (Fig. 7a, b). None of the illegitimate recombination events

observed in the alignment had caused frame shift. A region with high indel variation was revealed between the CC and the NBS domains (positions 380–460) where six indel haplotypes were observed (Fig. 7c). A repeat of 24 bp in the region, caused by illegitimate recombination, and subsequent recombination errors formed a 54-bp difference between the shortest and the longest sequences. However, none of these indels have caused any frame shift. This region is a linker between the CC and the NBS domains. Slatkin's exact test and Tajima's D test did not reveal a significant deviation from neutrality in this region.

Resistance tests

Resistance tests were conducted with the leaf rust isolate BRW97512-19, which is avirulent to some *Lr10* alleles from *T. durum* and *T. aestivum*. All 100 wild emmer wheat accessions tested in the current study showed high infection types (3–4) with this isolate.

Discussion

In the current study, the sequence diversity of *Lr10* within and between *T. dicoccoides* populations from Israel was estimated, the LD and recombination patterns along the *Lr10* gene were analysed, and the processes shaping the diversity and the LD were revealed.

Lr10 diversity within and between populations

The diversity revealed in the current paper is of the same magnitude and distribution of synonymous and nonsynonymous substitutions along the gene as observed by Loutre et al. (2009). *T. dicoccoides* populations showed large differences in diversity. These differences could be caused




Fig. 6 Schematic sequence display of recombination events detected by RDP (Martin et al. 2005). Full length bars are the acceptor (“daughter” in RDP terms) sequence. Short bars below each full length bar are recombinant fragments (“parent” in RDP terms). Different colors represent different representative accessions. Vertical lines are the domains borders

by the population demographic history, e.g., bottlenecks, or by selection pressure of the pathogen. The d_S and d_N values of the NBS-LRR domains obtained in this study were strongly correlated with the diversity index H_e obtained for

SSR markers in the same populations studied earlier (Peleg et al. 2008b). This correlation suggests that demographic processes have a large effect on *Lr10* nucleotide variation, since they act similarly on all loci. However, the negative d_S-d_N values in the CC domain obtained in all populations indicate that this domain is under positive selection. Such positive selection in the CC domain might indicate that this domain is interacting with pathogen effectors as suggested by Loutre et al. (2009) for *Lr10* and by Bergelson et al. (2001) in general. The d_S-d_N values differed several-fold between populations both in the CC domain and in the NBS-LRR domains. In the NBS-LRR domains, a significant $d_S-d_N > 0$ difference in four populations was observed, suggesting the involvement of purifying selection. However, the significant $d_S-d_N > 0$ values in the populations resulted from high d_S values and not from low d_N values. Therefore, the results may not reveal higher purifying selective pressure in the populations but rather higher diversity in the populations. Nevertheless, purifying selection may have acted on the NBS-LRR domains and its signature maintained in these populations while decaying in the other populations. The populations with significant $d_S-d_N > 0$ values originated in the transition zone between Mediterranean and steppe climate with annual precipitation of less than 450 mm and rain variability higher than 0.34. The high d_S values may reflect the overall high genomic variability in this climatic transition zone as suggested by Safriel et al. (1994), Volis et al. (2001), Kark et al. (2004), Kark et al. (2008) and Peleg et al. (2008b). In the CC domain, no clear pattern was observed for the distribution of populations with significant $d_S-d_N < 0$ values.

The PCO analysis used in the current study showed that the revealed diversity is old and probably existed before populations were separated to the current status, since haplotypes from the same populations did not form clusters. Furthermore, the haplotypes of *T. urartu*, the donor of the A genome to *T. dicoccoides*, are scattered and are not clustered. This pattern was also observed in Loutre et al. (2009) where different *Triticum* species shared the same *Lr10* haplotypes. The different pattern observed between the CC PCO and the NBS-LRR PCO could point to different selection pressures and/or a high recombination rate between the domains. The old origin of diversity is supported by the observation that most recombination events were detected between sequences obtained from different populations, implying that the resulting haplotypes were formed before populations separated. Significant gene flow could also contribute to low population differentiation but then one would expect geographical relationships between clustered accessions and this is not the case. Moreover, *T. dicoccoides* is a selfer with low gene flow rate (Golenberg and Nevo 1987). The genetic distance between *T. dicoccoides* populations is not correlated with geographic distance,




Fig. 7 Illegitimate recombination and indel diversity. **a** A typical deletion caused by illegitimate recombination. The upper sequence consists of a tri-nucleotide repeat (**bold**). The sequence between the tri-nucleotide repeats was deleted in the lower sequence. **b** An example of a degenerated repeat present in all sequences. A sequence aligned against itself. **Bold letters** are the short hepta-nucleotide repeat that is the cause of the illegitimate recombination. **Underlined letters** are the

tri-nucleotide repeat marked in **bold** in Fig. 7a. **c** High indel polymorphism at the linker region between CC and NBS domains (alignment of amino acid sequence). *Upper case* residues are identical or similar to the consensus. *Lower case* residues are dissimilar. *Lower case x* in the consensus sequence are non-consensus residues. **Bold** and **underlined** residues are part of a repeat

contrary to the expectations for situations where gene flow is substantial (Peleg et al. 2008b; Sela et al. 2009; Haudry et al. 2007). The ancient diversity observed in the current study supports a long evolutionary history of *Lr10*–pathogen effector interaction in the “trench warfare” model where polymorphism is maintained by alleles that are recycling between high and low frequencies shifted by frequency-dependent selection (Holub 2001; Stahl et al. 1999). The long coalescent time is also supported by the significant Tajima test of the two haplotypes in the LRR domain suggesting balancing selection. These findings of old polymorphism are in agreement with the observations of Loutre et al. (2009).

Resistance tests with a leaf rust isolate which is avirulent to *Lr10* of bread wheat did not detect any *Lr10*-dependent resistance in the collection of *T. dicoccoides* tested in the current study. Therefore, it was impossible to assess the direct impact of mutations of the different haplotypes on gene function and specificity. However, in most accessions, an intact open-reading frame was observed and the gene was expressed in all eight accessions that were tested for *Lr10* mRNA (data not shown). These findings suggest that *T. dicoccoides* *Lr10* haplotypes are functional against unknown leaf rust isolates present now, or that were present in the recent past.

Recombination and linkage disequilibrium

The alignment of *Lr10* sequences obtained from 54 accessions of *T. dicoccoides* revealed many recombination events. In the CC domain there were mostly very short recombinant sequences that are typical of gene conversion (Jeffrys and May 2004), while the NBS and LRR domains exhibited longer sequence exchanges that could result from crossing over events. These results are in agreement with LD analysis, which showed an LD block at the 3' end (Fig. 5). Recombination events were detected between *Lr10* sequences that were derived from different *T. dicoccoides* populations and not within populations. This may suggest that recombination events took place prior to the current separation between *T. dicoccoides* populations. Old recombination events imply that selective pressures may have an effect on the recombination outcome. Shuffling haplotypes at the CC domain enhances diversity and may have a selective advantage if this domain is involved in pathogen recognition as predicted. In the LRR domain, two well-conserved haplotypes were observed. These two haplotypes have a long evolutionary history and can be tracked back to *T. urartu*, the A genome donor of *T. dicoccoides* (Loutre et al. 2009). It seems that recombination is suppressed in this region. Apparently, there is a tendency not to shuffle

haplotypes within the NBS and LRR domains, since most of the breakpoints in the region are restricted to the domain borders. Michelmore and Meyers (1998) proposed that inter-allelic recombination enhances diversity in the LRR domain of *R*-genes. This expectation was validated by Kuang et al. (2004) and Yahiaoui et al. (2006). The data presented here about the LD block in the *Lr10* LRR domain do not match this expectation, but in the CC domain this expectation can be met. The LD block in the *Lr10* LRR domain is similar to the LD block that was observed in the *Rps2* LRR domain from *Arabidopsis thaliana*: both blocks are located around a hot spot of diversity. The *Rps2* hot spot of divergence differentiates between resistant and susceptible clades while the *Lr10* hot spot does not distinguish between resistant and susceptible haplotypes (Mauricio et al. 2003; Loutre et al. 2009). It could be because the *A. thaliana* *Rps2* block is located in the 5' region of the LRR domain while in the current *Lr10* study the LD block is located in the 3' region. The two parts of the LRR play a different role in the protein function (Lukasik and Takken 2009).

Understanding LD decay patterns is essential when conducting association studies or for estimating rates of recombination. Population structure can lead to spurious LD associations, especially in selfing species (Ewens and Spielman 1995). In the current study r^2 values were better predictors of physical distance than LD p values since they were less affected by population structure. Choosing a subset of sequences from a tree proved to be a good method to avoid population structure bias. This approach is somewhat similar to the method implemented by Breseghello and Sorrells (2006), even though they have removed only very similar haplotypes, and in the current study only distant haplotypes were selected. Taking into account that *T. dicoccoides* is a selfer, LD decay in *Lr10* was very rapid, within 1–2 kb. This LD extent was also found in wild barley, *Hordeum spontaneum*, a relative of wheat and a selfer as well (Morrell et al. 2006; Caldwell et al. 2006). Whole genome LD studies in the selfer *A. thaliana* have shown a longer extent of LD, up to 250 kb (Nordborg et al. 2002). LD in bread wheat collections extended 1–5 cM (Chao et al. 2007; Horvath et al. 2009; Somers et al. 2007). Longer LD extents may be due to longer stretches between the sampled loci that overlook local gene conversion events and only display crossing over events (Plagnol et al. 2006). This situation creates a paradox, where high-resolution analysis, on a base-pair scale, shows a steep decay in LD, while in low-resolution analysis, on the centiMorgan scale, the decay is far more moderate (Andolfatto and Nordborg 1998). From a practical point of view, attempts to associate very closely linked SNPs with a trait, especially in the highly diverse *R* genes, may be hampered by the short LD extent. The rapid decay may be a result of a high number of

point mutations or recombination events. In the current study the ratio ρ/θ was lower than 1 (0.59), which may imply that recombination events had a smaller effect than mutations on sequence diversity. Nevertheless, the ρ/θ ratio is fairly high for inbreeders. For reference, the average ρ/θ ratio in inbreeders ranges from 0.05 in *A. thaliana* to 1.5 in wild barley (Morrell et al. 2006; Nordborg et al. 2002). In the outbreeder wild maize, teosinte (*Zea mays ssp. parviglumis*), ρ/θ ratio is 4.5 (Wright et al. 2005).

Recombination is a major force enhancing indel diversity as revealed by the numerous events of illegitimate recombination, especially in the intron region of *Lr10*. One interesting example is the linker region between the NBS and the CC domains of *Lr10* where high length variability was observed between six haplotypes. It is not clear if the variability in this region affects the function of the protein. Several studies have shown that the CC domain in *R* genes can function even when expressed separately from the rest of the protein [e.g. Rairdan et al. (2008) and Moffett et al. (2002)]. Neutrality tests looking for old polymorphism (Slatkin 1994; Tajima 1989) did not reject the neutrality of indel variation of this region but the long and short haplotypes found in all ploidy levels (*T. urartu* AA, *T. dicoccoides* BBAA, *T. aestivum* BBAADD) may indicate that these haplotypes are old and stable (Loutre et al. 2009).

The results presented here highlight the role of recombination in enhancing diversity in wild populations of the selfing plant species *T. dicoccoides*. The results support the trench warfare model of old polymorphism in *R*-genes. The high diversity observed in *Lr10* in *T. dicoccoides* populations emphasizes the need to further explore and conserve this diverse gene pool.

Acknowledgments This work was supported by grants from the European Union's Sixth Framework Programme (FP6) in the BioExploit project (No. CT-2005-513949). The authors acknowledge The Israel Science Foundation equipment grants #048/99, #1478/04, #1719/08. The authors wish to thank Ortal Mergi and Anne-Mari Narvanto for technical assistance.

References

- Andolfatto P, Nordborg M (1998) The effect of gene conversion on intralocus associations. *Genetics* 148:1397–1399
- Anikster Y, Manisterski J, Long DL, Leonard KJ (2005) Leaf rust and stem rust resistance in triticum *dicoccoides* populations in Israel. *Plant Dis* 89:55–62
- Auton A, McVean G (2007) Recombination rate estimation in the presence of hotspots. *Genome Res* 17:1219
- Bergelson J, Kreitman M, Stahl EA, Tian D (2001) Evolutionary dynamics of plant *R*-Genes. *Science* 292:2281–2285
- Bolton MD, Kolmer JA, Garvin DF (2008) Wheat leaf rust caused by *Puccinia triticina*. *Mol Plant Pathol* 9:563–575
- Bradbury PJ, Zhang Z, Kroon DE, Casstevens TM, Ramdoss Y, Buckler ES (2007) TASSEL: software for association mapping of complex traits in diverse samples. *Bioinformatics* 23:2633–2635

- Breseghele F, Sorrells ME (2006) Association mapping of kernel size and milling quality in wheat (*Triticum aestivum* L.) cultivars. *Genetics* 172:1165–1177
- Butterbach P (2007) Molecular evolution of the disease resistance gene Rx in Solanum. PhD dissertation. Wageningen University, The Netherlands
- Caldwell KS, Russell J, Langridge P, Powell W (2006) Extreme population-dependent linkage disequilibrium detected in an inbreeding plant species, *Hordeum vulgare*. *Genetics* 172:557–567
- Chao S, Zhang W, Dubcovsky J, Sorrells M (2007) Evaluation of genetic diversity and genome-wide linkage disequilibrium among U.S. Wheat (*Triticum aestivum* L.) germplasm representing different market classes. *Crop Sci* 47:1018–1030
- Devos KM, Brown JKM, Bennetzen JL (2002) Genome size reduction through illegitimate recombination counteracts genome expansion in *Arabidopsis*. *Genome Res* 12:1075–1079
- Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* 32:1792
- Ersoz ES, Yu J, Buckler ES (2008) Applications of linkage disequilibrium and association mapping in crop plants. In: Varshney RK, Tuberosa R (eds) *Genomics-assisted crop improvement*. Springer, Netherlands, pp 97–119
- Ewens WJ, Spielman RS (1995) The transmission/disequilibrium test: history, subdivision, and admixture. *Am J Hum Genet* 57:455
- Excoffier L, Laval G, Schneider S (2005) Arlequin version 3.0: an integrated software package for population genetics data analysis. *Evol Bioinform Online* 1:47–50
- Feuillet C, Travella S, Stein N, Albar L, Nublait A, Keller B (2003) Map-based isolation of the leaf rust disease resistance gene *Lr10* from the hexaploid wheat (*Triticum aestivum* L.) genome. *Proc Natl Acad Sci USA* 100:15253–15258
- Flint-Garcia SA, Thornsberry JM, E S, IV B (2003) Structure of linkage disequilibrium in plants. *Annu Rev Plant Biol* 54:357–374
- Gaut BS, Wright SI, Rizzon C, Dvorak J, Anderson LK (2007) Opinion: recombination: an underappreciated factor in the evolution of plant genomes. *Nat Rev Genet* 8:77–84
- Golenberg EM, Nevo E (1987) Multilocus differentiation and population structure in a selfer, wild emmer wheat. *Triticum dicoccoides* *Heredity* 451–456
- Gustafson JP, Raskina O, Ma X, Nevo E (2009) Evolutionary genomics of wheat. In: Carver BF (ed) *Wheat: Science and Trade*. Wiley-Blackwell, NJ, pp 5–30
- Hall T (2007) BioEdit, version 7.0. 9. Computer program and documentation. Ibis Biosciences, Carlsbad
- Haudry A, Cenci A, Ravel C, Bataillon T, Brunel D, Poncet C, Hochu I, Poirier S, Santoni S, Glemin S (2007) Grinding up wheat: a massive loss of nucleotide diversity since domestication. *Mol Biol Evol* 24:1506–1517
- Holub EB (2001) The arms race is ancient history in *Arabidopsis*, the wildflower. *Nat Rev Genet* 2:516–527
- Horvath A, Didier A, Koenig J, Exbrayat F, Charmet G, Balfourier F (2009) Analysis of diversity and linkage disequilibrium along chromosome 3B of bread wheat (*Triticum aestivum* L.). *Theor Appl Genet* 119:1523–1537
- Hudson RR (2001) Two locus sampling distributions and their application. *Genetics* 159:1805–1817
- Jeffrys AJ, May CA (2004) Intense and highly localized gene conversion activity in human meiotic crossover hot spots. *Nat Genet* 36:151–156
- Kark S, Volis S, Novoplansky A (2004) Biodiversity along core-periphery distribution clines. In: Shachak M, Gosz JR, Pickett STA, Perevolotsky A (eds) *Biodiversity in drylands*. Oxford University Press, Oxford, pp 30–56
- Kark S, Hadany L, Safriel UN, Noy-Meir I, Eldredge N, Tabarroni C, Randi E (2008) How does genetic diversity change towards the range periphery? An empirical and theoretical test. *Evol Ecol Res* 10:391–414
- Knott DR, DaPeng B, Zale J (2005) The transfer of leaf and stem rust resistance from wild emmer wheats to durum and common wheat. *Can J Plant Sci* 85:49–57
- Korol AB, Preygel IA, Preygel SI (1994) *Recombination variability and evolution*. Chapman & Hall, London, p 361
- Kuang H, Woo SS, Meyers BC, Nevo E, Michelmore RW (2004) Multiple genetic processes result in heterogeneous rates of evolution within the major cluster disease resistance genes in lettuce. *Plant Cell* 16:2870–2894
- Kuang H, van Eck HJ, Sicard D, Michelmore R, Nevo E (2008) Evolution and genetic population structure of prickly lettuce (*Lactuca serriola*) and its RGC2 resistance gene cluster. *Genetics* 178:1547–1558
- Kumar S, Tamura K, Nei M (2004) MEGA3: integrated software for molecular evolutionary genetics analysis and sequence alignment. *Brief Bioinform* 5:150–163
- Li YC, Fahima T, Peng JH, Röder MS, Kirzhner VM, Beiles A, Korol AB, Nevo E (2000a) Edaphic microsatellite DNA divergence in wild emmer wheat, *Triticum dicoccoides*, at a microsite: Tabigha, Israel. *Theor Appl Genet* 101:1029–1038
- Li Y, Röder MS, Fahima T, Kirzhner VM, Beiles A, Korol AB, Nevo E (2000b) Natural selection causing microsatellite divergence in wild emmer wheat at the ecologically variable microsite at Ammiad, Israel. *Theor Appl Genet* 100:985–999
- Loutre C, Wicker T, Travella S, Galli P, Scofield S, Fahima T, Feuillet C, Keller B (2009) Two different CC-NBS-LRR genes are required for Lr10-mediated leaf rust resistance in tetraploid and hexaploid wheat. *Plant J* 60:1043–1054
- Lukasik E, Takken FLW (2009) STANDING strong, resistance proteins instigators of plant defence. *Curr Opin Plant Biol* 12:427–436
- Maccaferri M, Sanguineti MC, Noli E, Tuberosa R (2005) Population structure and long-range linkage disequilibrium in a durum wheat elite collection. *Mol Breed* 15:271–290
- Marais GF, Pretorius ZA, Wellings CR, McCallum B, Marais AS (2005) Leaf rust and stripe rust resistance genes transferred to common wheat from *Triticum dicoccoides*. *Euphytica* 143:115–123
- Martin DP, Williamson C, Posada D (2005) RDP2: recombination detection and analysis from sequence alignments. *Bioinformatics* 21:260–262
- Massa A, Morris C (2006) Molecular evolution of the puroindoline-a, puroindoline-b, and grain softness protein-1 genes in the tribe *Triticeae*. *J Mol Evol* 63:526–536
- Mauricio R, Stahl EA, Korves T, Tian D, Kreitman M, Bergelson J (2003) Natural selection for polymorphism in the disease resistance gene Rps2 of *Arabidopsis thaliana*. *Genetics* 163:735–746
- Mcdowell JM, Simon SA (2006) Recent insights into R gene evolution. *Mol Plant Pathol* 7:437–448
- McIntosh RA, Wellings CR, Park RF (1995) *Wheat rusts: an atlas of resistance genes*. CSIRO, East Melbourne
- Meaux J, Neema C (2003) Spatial patterns of diversity at the putative recognition domain of resistance gene candidates in wild bean populations. *J Mol Evol* 57:S90–S102
- Mergoum M, Froberg RC, Miller JD, Stack RW (2005) Registration of ‘Steele-ND’ wheat registration by CSSA. *Crop Sci* 45:1163–1164
- Michelmore RW, Meyers BC (1998) Clusters of resistance genes in plants evolve by divergent selection and a birth-and-death process. *Genome Res* 8:1113–1130
- Moffett P, Farnham G, Peart J, Baulcombe DC (2002) Interaction between domains of a plant NBS-LRR protein in disease resistance-related cell death. *EMBO J* 21:4511–4519
- Mondragon-Palomino M, Gaut BS (2005) Gene conversion and the evolution of three leucine-rich repeat gene families in *Arabidopsis thaliana*. *Mol Biol Evol* 22:2444–2456

- Morrell PL, Toleno DM, Lundy KE, Clegg MT (2006) Estimating the contribution of mutation, recombination and gene conversion in the generation of haplotypic diversity. *Genetics* 173:1705
- Moseman JG, Nevo E, Gerechter-Amitai ZK, El-Morshidy MA, Zohary D (1985) Resistance of *Triticum dicoccoides* collected in Israel to infection with *Puccinia recondita tritici*. *Crop Sci* 25:262–265
- Nei M (1987) Molecular evolutionary genetics. Columbia University Press, USA
- Nei M, Kumar S (2000) Molecular evolution and phylogenetics. Oxford University Press, USA
- Nevo E, Beiles A (1989) Genetic diversity of wild emmer wheat in Israel and Turkey—structure, evolution, and application in breeding. *Theor Appl Genet* 77:421–455
- Nevo E, Korol AB, Beiles A, Fahima T (2002) Evolution of wild emmer and wheat improvement: population genetics, genetic resources and genome organization of wheat's progenitor *Triticum dicoccoides*. Springer, Berlin
- Nordborg M, Borevitz JO, Bergelson J, Berry CC, Chory J, Hagenblad J, Kreitman M, Maloof JN, Noyes T, Oefner PJ (2002) The extent of linkage disequilibrium in *Arabidopsis thaliana*. *Nat Genet* 30:190–193
- Peakall R, Smouse PE (2006) GENALEX 6: genetic analysis in Excel Population genetic software for teaching and research. *Mol Ecol Notes* 6:288–295
- Peleg Z, Fahima T, Abbo S, Krugman T, Saranga Y (2008a) Genetic structure of wild emmer wheat populations as reflected by transcribed versus anonymous SSR markers. *Genome* 51:187–195
- Peleg Z, Saranga Y, Krugman T, Abbo S, Nevo E, Fahima T (2008b) Allelic diversity associated with aridity gradient in wild emmer wheat populations. *Plant Cell Environ* 31:39–49
- Plagnol V, Padhukasahasram B, Wall JD, Marjoram P, Nordborg M (2006) Relative influences of crossing over and gene conversion on the pattern of linkage disequilibrium in *Arabidopsis thaliana*. *Genetics* 172:2441–2448
- Rairdan GJ, Collier SM, Sacco MA, Baldwin TT, Boettrich T, Moffett P (2008) The coiled-coil and nucleotide binding domains of the potato rx disease resistance protein function in pathogen recognition and signaling. *Plant Cell* 20:739–751
- Rong J, Millet E, Manisterski J, Feldman M (2000) A new powdery mildew resistance gene: introgression from wild emmer into common wheat and RFLP-based mapping. *Euphytica* 115:121–126
- Rostoks N, Ramsay L, MacKenzie K, Cardle L, Bhat PR, Roose ML, Svensson JT, Stein N, Varshney RK, Marshall DF, Graner A, Close TJ, Waugh R (2006) Recent history of artificial outcrossing facilitates whole-genome association mapping in elite inbred crop varieties. *Proc Natl Acad Sci USA* 103:18656–18661
- Rozas J, Sanchez-DelBarrio JC, Messeguer X, Rozas R (2003) DnaSP, DNA polymorphism analyses by the coalescent and other methods. *Bioinformatics* 19:2496–2497
- Safriel UN, Volis S, Karr S (1994) Core and peripheral populations and global climate. *Isr J Plant Sci* 42:331–345
- Schachermayr GM, Messmer MM, Feuillet C, Winzeler H, Winzeler M, Keller B (1995) Identification of molecular markers linked to the *Agropyron elongatum*-derived leaf rust resistance gene *Lr24* in wheat. *TAG Theor Appl Genet* 90:982–990
- Sela H, Cheng J, Jun Y, Nevo E, Fahima T (2009) Divergent diversity patterns of NBS and LRR domains of resistance gene analogs in wild emmer wheat populations. *Genome* 52:557–565
- Slatkin M (1994) An exact test for neutrality based on the Ewens sampling distribution. *Genet Res* 64:71–74
- Somers DJ, Banks T, DePauw R, Fox S, Clarke J, Pozniak C, McCartney C (2007) Genome-wide linkage disequilibrium analysis in bread wheat and durum wheat. *Genome* 50:557–567
- Sonnhammer ELL, Durbin R (1995) A dot-matrix program with dynamic threshold control suited for genomic DNA and protein sequence analysis. *Gene* 167
- Stahl EA, Dwyer G, Mauricio R, Kreitman M, Bergelson J (1999) Dynamics of disease resistance polymorphism at the *Rpm1* locus of *Arabidopsis*. *Nature* 400:667–671
- Tajima F (1989) Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics* 123:585–595
- Tamura K, Kumar S (2002) Evolutionary distance estimation under heterogeneous substitution pattern among lineages. *Mol Biol Evol* 19:1727
- Tamura K, Nei M, Kumar S (2004) Prospects for inferring very large phylogenies by using the neighbor-joining method. *Proc Natl Acad Sci USA* 101:11030
- Uauy C, Brevis JC, Chen X, Khan I, Jackson L, Chicaiza O, Distelfeld A, Fahima T, Dubcovsky J (2005) High-temperature adult-plant (HTAP) stripe rust resistance gene *Yr36* from *Triticum turgidum* ssp. *dicoccoides* is closely linked to the grain protein content locus *Gpc-B1*. *Theor Appl Genet* 112:97–105
- Volis S, Yakubov B, Shulgina I, Ward D, Zur V, Mendlinger S (2001) Tests for adaptive RAPD variation in population genetic structure of wild barley, *Hordeum spontaneum* Koch. *Biol J Linn Soc* 74:289–303
- Watterson GA (1975) On the number of segregating sites in genetical models without recombination. *Theor Popul Biol* 7:256–276
- Waugh R, Jannink JL, Muehlbauer GJ, Ramsay L (2009) The emergence of whole genome association scans in barley. *Curr Opin Plant Biol* 12:218–222
- Wicker T, Yahiaoui N, Keller B (2007) Illegitimate recombination is a major evolutionary mechanism for initiating size variation in plant resistance genes. *Plant J* 51:631–641
- Wright SI, Bi IV, Schroeder SG, Yamasaki M, Doebley JF, McMullen MD, Gaut BS (2005) The effects of artificial selection on the maize genome. *Science* 308:1310
- Yahiaoui N, Brunner S, Keller B (2006) Rapid generation of new powdery mildew resistance genes after wheat domestication. *Plant J* 47:85–98
- Yahiaoui N, Kaur N, Keller B (2009) Independent evolution of functional *Pm3* resistance genes in wild tetraploid wheat and domesticated bread wheat. *Plant J* 57:846–856