

Exercise restores endothelial function independently of weight loss or hyperglycaemic status in db/db mice

F. Moien-Afshari · S. Ghosh · S. Elmi · M. M. Rahman ·
N. Sallam · M. Khazaei · T. J. Kieffer ·
R. W. Brownsey · I. Laher

Published online: 13 November 2008
© Springer-Verlag 2008

Erratum to: Diabetologia (2008) 51:1327–1337
DOI 10.1007/s00125-008-0996-x

The last author regrets the omission of three authors from this paper. The correct author list is shown above.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00125-008-0996-x>.

F. Moien-Afshari · S. Elmi · M. M. Rahman · N. Sallam ·
I. Laher (✉)
Department of Pharmacology and Therapeutics,
Faculty of Medicine, University of British Columbia,
2176 Health Sciences Mall,
Vancouver, BC V6T 1Z3, Canada
e-mail: ilaher@interchange.ubc.ca

S. Ghosh
Department of Pediatrics, Child and Family Research Institute,
Vancouver, BC, Canada

M. Khazaei
Department of Physiology,
Isfahan University of Medical Sciences,
Isfahan, Iran

T. J. Kieffer
Departments of Cellular & Physiological Sciences and Surgery,
University of British Columbia,
Vancouver, BC, Canada

R. W. Brownsey
Department of Biochemistry and Molecular Biology,
Faculty of Medicine, University of British Columbia,
Vancouver, BC, Canada