

# Arctigenin, a natural compound, activates AMP-activated protein kinase via inhibition of mitochondria complex I and ameliorates metabolic disorders in *ob/ob* mice

S.-L. Huang · R.-T. Yu · J. Gong · Y. Feng · Y.-L. Dai ·  
F. Hu · Y.-H. Hu · Y.-D. Tao · Y. Leng

Received: 9 September 2011 / Accepted: 6 October 2011 / Published online: 18 November 2011  
© Springer-Verlag 2011

## Abstract

**Aims/hypothesis** Arctigenin is a natural compound that had never been previously demonstrated to have a glucose-lowering effect. Here it was found to activate AMP-activated protein kinase (AMPK), and the mechanism by which this occurred, as well as the effects on glucose and lipid metabolism were investigated.

**Methods** 2-Deoxyglucose uptake and AMPK phosphorylation were examined in L6 myotubes and isolated skeletal muscle. Gluconeogenesis and lipid synthesis were evaluated in rat primary hepatocytes. The acute and chronic effects of arctigenin on metabolic abnormalities were observed in C57BL/6J and *ob/ob* mice. Changes in mitochondrial membrane potential were measured using the J-aggregate-forming dye, JC-1. Analysis of respiration of L6 myotubes or isolated mitochondria was conducted in a channel oxygen system.

**Results** Arctigenin increased AMPK phosphorylation and stimulated glucose uptake in L6 myotubes and isolated

skeletal muscles. In primary hepatocytes, it decreased gluconeogenesis and lipid synthesis. The enhancement of glucose uptake and suppression of hepatic gluconeogenesis and lipid synthesis by arctigenin were prevented by blockade of AMPK activation. The respiration of L6 myotubes or isolated mitochondria was inhibited by arctigenin with a specific effect on respiratory complex I. A single oral dose of arctigenin reduced gluconeogenesis in C57BL/6J mice. Chronic oral administration of arctigenin lowered blood glucose and improved lipid metabolism in *ob/ob* mice.

**Conclusions/interpretation** This study demonstrates a new role for arctigenin as a potent indirect activator of AMPK via inhibition of respiratory complex I, with beneficial effects on metabolic disorders in *ob/ob* mice. This highlights the potential value of arctigenin as a possible treatment of type 2 diabetes.

**Keywords** AMP-activated protein kinase · Gluconeogenesis · Glucose uptake · Lipid synthesis · Mitochondria · Natural compound · Respiratory complex I

**Electronic supplementary material** The online version of this article (doi:10.1007/s00125-011-2366-3) contains peer-reviewed but unedited supplementary material, which is available to authorised users.

S.-L. Huang · J. Gong · Y. Feng · Y.-L. Dai · F. Hu · Y.-H. Hu ·  
Y. Leng (✉)

State Key Laboratory of Drug Research,  
Shanghai Institute of Materia Medica,  
Chinese Academy of Sciences,  
Zu Chong Zhi Road 555,  
Shanghai 201203, People's Republic of China  
e-mail: yleng@mail.shnc.ac.cn

R.-T. Yu · Y.-D. Tao (✉)

Northwest Plateau of Biology Institute,  
Chinese Academy of Sciences,  
59 Xiguan Street,  
Xining 810008, People's Republic of China  
e-mail: taoyanduo@163.com

## Abbreviations

ACC	Acetyl coenzyme A synthetase
AICAR	5-Amino-imidazole carboxamide ribonucleotide
AMPK	AMP-activated protein kinase
CaMKK	Ca <sup>2+</sup> /calmodulin-dependent protein kinase kinase
EDL	Extensor digitorum longus
LKB1	Liver kinase B1
$\Delta\psi_m$	Mitochondrial membrane potential
NOS	Nitric oxide synthase
ONOO <sup>-</sup>	Peroxyntirite
PI3-kinase	Phosphoinositide-3 kinase

siRNA	Small interfering RNA
UPR	Unfolded protein response

## Introduction

Type 2 diabetes is characterised by abnormal glucose and lipid metabolism due to insulin resistance and impaired insulin secretion. Skeletal muscle is the primary tissue in glucose homeostasis, accounting for approximately 80% of whole-body glucose disposal under insulin-stimulated conditions [1]. Glucose transport is the first rate-limiting step in glucose metabolism. Reduced glucose transport in skeletal muscle leads to impaired whole-body glucose uptake and contributes to the pathogenesis of type 2 diabetes mellitus [2–4].

Glucose transport in skeletal muscle can be activated by two separate pathways (insulin-dependent and insulin-independent). In type 2 diabetic patients, insulin-mediated glucose transport in skeletal muscle is impaired due to defects in insulin signalling pathways [5]. Thus, therapies that target insulin-independent pathways in the regulation of glucose transport are likely to be more beneficial in the treatment of type 2 diabetes mellitus. Several factors, such as muscle contraction (exercise), hypoxia and cellular stress have been demonstrated to increase glucose transport independently of insulin in skeletal muscle, effects to which the activation of AMP-activated protein kinase (AMPK) has been suggested to contribute [6–8]. Moreover, it has been shown that activation of AMPK by known AMPK activators restored glucose transport in skeletal muscle from patients with type 2 diabetes mellitus, while insulin-induced glucose transport was impaired [9]. As such, activation of AMPK in skeletal muscle could be an attractive way to increase glucose transport independently of insulin, thus bypassing the defective insulin signalling in type 2 diabetic patients.

AMPK is a heterotrimeric protein composed of a catalytic  $\alpha$  subunit and regulatory  $\beta$  and  $\gamma$  subunits. It is activated in response to an increase of the intracellular AMP/ATP ratio due to several stimuli, e.g. hypoxia, glucose deprivation or the action of various hormones, cytokines and adipokines. AMPK thus acts as an efficient sensor of cellular energy states and plays an important role in the regulation of whole-body energy homeostasis [9]. Liver kinase B1 (LKB1) and  $\text{Ca}^{2+}$ /calmodulin-dependent protein kinase kinase (CaMKK) are two distinct upstream kinases that activate AMPK by phosphorylation of Thr172. LKB1 is activated by an increase of the AMP/ATP ratio, while CaMKK is activated by an increase in intracellular  $\text{Ca}^{2+}$  [10, 11]. Once activated, AMPK leads to a concomitant activation of energy-producing processes (e.g. fatty acid oxidation and glucose uptake) and inhibition of

energy-consuming biosynthetic processes (e.g. fatty acid and sterol synthesis) to restore energy balance [12]. Genetic and pharmacological studies have demonstrated that AMPK controls whole-body glucose homeostasis by regulating metabolism in various peripheral tissues, such as skeletal muscle, liver and adipose tissues [13]. Metformin, one of the most commonly used drugs for the treatment of type 2 diabetes, has been demonstrated to exert pleiotropic beneficial effects by activation of AMPK [14], suggesting that AMPK could be a very attractive drug target for the treatment of type 2 diabetes.

Recently, several small molecules have been reported to activate AMPK. Direct activators of AMPK, such as the thienopyridone family, act without increasing cellular AMP/ATP ratios [15], while various pharmacological agents have been reported to activate AMPK indirectly by modes of action that have not been fully delineated. Natural products have played a significant role in new drug discovery and development processes [16], with some reported to possess glucose-lowering effects [17, 18]. Moreover, natural products such as alkaloids, bitter melon extracts and epigallocatechin-3-gallate (EGCG) have been found to activate AMPK and improve metabolic disease [17, 19]. However, most of these agents displayed inadequate efficacy *in vivo*, leading to the demand to develop more effective AMPK activators.

Given the importance of skeletal muscle's capacity for glucose transport for whole-body glucose homeostasis and that glucose transport could be induced by activation of AMPK, we conducted a primary screening of our natural products library for 2-deoxyglucose uptake in L6 myotubes, followed by a secondary screening based on the evaluation of AMPK phosphorylation. During these screenings, arctigenin, a natural compound that had never been previously shown to have a glucose-lowering effect, was found to increase glucose uptake and AMPK phosphorylation in L6 myotubes. In the present study, the mechanisms involved in the activation of AMPK by arctigenin were explored, and the effects of arctigenin on glucose and lipid metabolism investigated *in vitro* and *in vivo*.

## Methods

### Animals

C57BL/6J male mice and Sprague–Dawley male rats were purchased from SLAC Laboratory Animals (Shanghai, China). B6.V-*Lep<sup>ob</sup>/Lep<sup>ob</sup>* mice (Jackson Laboratory, Bar Harbor, ME, USA) were bred at the Shanghai Institute of Materia Medica (Chinese Academy of Sciences, Shanghai, People's Republic of China). The animals were maintained

under a 12 h light–dark cycle with free access to water and food. Animal experiments were approved by the Animal Care and Use Committee, Shanghai Institute of Materia Medica.

#### 2-Deoxyglucose uptake, GLUT4 translocation and protein phosphorylation in L6 myotubes

L6 myoblasts were differentiated to myotubes and treated with or without arctigenin, followed by 2-deoxyglucose uptake measurement and analysis of GLUT4 translocation or protein phosphorylation by western blot (electronic supplementary material [ESM] [Methods](#)).

#### 2-Deoxyglucose uptake and protein phosphorylation in isolated muscle

Soleus and extensor digitorum longus (EDL) muscles were isolated for analysis of 2-deoxyglucose uptake and protein phosphorylation (see ESM [Methods](#)).

#### Gluconeogenesis, lipid synthesis and protein phosphorylation in primary cultured rat hepatocytes

Rat hepatocytes were isolated by collagenase digestion from rats that had been fasted for 24 h (gluconeogenesis analysis) or from fed rats (protein phosphorylation or lipid synthesis analysis). Gluconeogenesis, fatty acid and sterol synthesis, and protein phosphorylation were measured as described in ESM [Methods](#).

#### Transfection of small interfering RNA

L6 myotubes were transfected with small interfering RNA (siRNA) using siLentFect lipid (catalogue number 170–3360; Bio-Rad Laboratories, Hercules, CA, USA) as described in ESM [Methods](#).

#### Western blot analysis

Protein preparation and western blot analysis were performed as described in ESM [Methods](#).

#### Acute arctigenin administration in normal mice

The effects of acute administration of arctigenin on gluconeogenesis and AMPK $\alpha$  phosphorylation were observed in male C57BL/6J mice (see ESM [Methods](#)).

#### AMPK activity assay

AMPK activity was measured by monitoring phosphorylation of the AMARA peptide substrate (see ESM [Methods](#)).

#### Hela cells culture

Full details are provided in ESM [Methods](#).

#### Mitochondrial membrane potential assay

This assay was performed according to a previous report [20], with modifications (see ESM [Methods](#)).

#### Adenine nucleotide extraction and measurement

Adenine nucleotides were extracted from L6 myotubes and measured by HPLC as described in ESM [Methods](#).

#### Measurement of respiration in L6 myotubes and isolated mitochondria

Liver mitochondria were prepared from Sprague–Dawley male rats. Isolated liver was finely chopped, homogenised (Polytron Homogeniser; Kinematica, Lucerne, Switzerland) and centrifuged (300 g) as described in a previous report [21]. Mitochondria were resuspended in respiration medium at a concentration of 60 mg/ml. Respiration measurements in L6 myotubes and isolated mitochondria were performed in a 782 2-channel oxygen system (Strathkelvin Instruments, Motherwell, UK) (see ESM [Methods](#)).

#### Determination of lactate content

L6 myotubes were cultured in a 24 well plate and treated for 1 or 4 h with 3  $\mu$ mol/l arctigenin or 50  $\mu$ mol/l rosiglitazone (positive control) in FBS-free DMEM. Lactate in the medium was measured with a lactate assay kit (Nanjing Jiancheng Bioengineering Institute, Nanjing, People's Republic of China).

#### Arctigenin treatment in *ob/ob* mice

**Study 1** Female *ob/ob* mice (6–7 weeks old) were assigned to three groups ( $n=8$ ) based on blood glucose and body weight, and subjected to gavage treatment with vehicle (0.5% carboxymethyl cellulose, wt/vol.), 200 mg/kg arctigenin or 250 mg/kg metformin, respectively, for 23 days. The effects of arctigenin on metabolic disorders of *ob/ob* mice were investigated. HOMA-IR was calculated (see ESM [Methods](#)).

**Study 2** Female *ob/ob* mice (6–7 weeks) were assigned to two groups ( $n=18$ ) based on blood glucose and body weight, and subjected to twice daily intraperitoneal injection of 25 mg/kg arctigenin or vehicle (0.5% Tween 80, vol./vol.) for 22 days. AKT phosphorylation in skeletal muscle, liver and fat isolated from arctigenin-treated or

vehicle-treated (control) mice was evaluated under basal and insulin-stimulated conditions (see ESM Methods).

#### RNA isolation and real-time PCR

Gene expression was analysed by real-time PCR as described in ESM Methods. For details on primer sequences, see ESM Table 1.

#### Statistical analysis

Results are expressed as mean±SEM. The statistical analysis was performed with a two-tailed unpaired *t* test. Values of  $p < 0.05$  were considered to be statistically significant.

## Results

### Arctigenin increases glucose uptake and AMPK phosphorylation in L6 myotubes

To identify new glucose-lowering lead compounds from natural products, we screened our phytochemicals collection for 2-deoxyglucose uptake in L6 myotubes. Arctigenin, a natural product (Fig. 1a), was found to increase glucose uptake under basal and insulin-stimulated conditions. Incubation of L6 myotubes with 1 µmol/l arctigenin resulted in a 1.49-fold increase in glucose uptake, which is comparable with the effect of 100 nmol/l insulin (Fig. 1b). Arctigenin increased basal glucose uptake in a dose- and time-dependent manner. Glucose uptake increased to 1.83-fold of control after treatment with 3 µmol/l arctigenin for 2 h, whereas the same dose of berberine only caused a 1.49-fold enhancement (Fig. 1c, d). Moreover, arctigenin treatment increased GLUT4 protein levels in the plasma membrane, whereas no change of total GLUT4 abundance was observed, suggesting that arctigenin stimulated GLUT4 translocation in L6 myotubes (Fig. 1e). Arctigenin dose-dependently increased AMPK and acetyl coenzyme A synthetase (ACC) phosphorylation in L6 myotubes (Fig. 1f, g). Significant increases of AMPK and ACC phosphorylation were observed within 15 min of arctigenin treatment (3 µmol/l) and were maintained for up to 120 min (Fig. 1h, i).

### Arctigenin increases glucose uptake and AMPK phosphorylation in isolated muscles

Measurement of 2-deoxyglucose uptake was performed in isolated EDL and soleus muscles from C57BL/6J mice. Incubation of EDL or soleus muscles with 3 µmol/l arctigenin

resulted in an increase of glucose uptake by 31.7% ( $p = 0.1$ ) and 18% ( $p < 0.05$ ), or 42% ( $p < 0.05$ ) and 20.6% ( $p < 0.01$ ) under basal and insulin-stimulated conditions, respectively (Fig. 2a, b). Arctigenin also dose-dependently increased AMPK and ACC phosphorylation in isolated EDL and soleus muscles (Fig. 2c–f).

### Arctigenin-stimulated glucose uptake is AMPK-dependent

To determine whether arctigenin-stimulated glucose uptake involves either the phosphoinositide-3 kinase (PI3 kinase)/AKT pathway or activation of AMPK, the effect of the PI3 kinase inhibitor, wortmannin, or the AMPK inhibitor, compound C, on arctigenin-stimulated glucose uptake was observed in L6 myotubes. As shown in Fig. 3a, wortmannin blocked insulin-stimulated glucose uptake, but had no effect on arctigenin stimulation, whereas arctigenin-stimulated glucose uptake was fully blocked by compound C. Furthermore, arctigenin-stimulated glucose uptake was completely lost upon *Ampkα1* (also known as *Prkaa1*) and *Ampkα2* (also known as *Prkaa2*) silencing in L6 myotubes (Fig. 3c, d). These data indicate that arctigenin-stimulated glucose uptake was mediated by activation of AMPK.


### Arctigenin increases AMPK phosphorylation, and inhibits gluconeogenesis and lipid synthesis in primary hepatocytes

Primary rat hepatocytes were incubated for 1 h in the absence or presence of 2.5, 5, 10, 20 or 40 µmol/l arctigenin. AMPK and ACC phosphorylation were increased dose-dependently, suggesting that arctigenin activated the AMPK signalling pathway in hepatocytes (Fig. 4a). Since activation of AMPK is known to regulate glucose and lipid metabolism in liver, the effects of arctigenin on gluconeogenesis and lipid synthesis were evaluated in primary rat hepatocytes. As shown in Fig. 4b–d, glucagon-induced hepatic gluconeogenesis was significantly suppressed by arctigenin treatment in a dose-dependent manner; arctigenin also significantly decreased the insulin-stimulated synthesis of fatty acid and sterol. Moreover, the suppression of gluconeogenesis and lipid synthesis by arctigenin was completely blocked by compound C pretreatment (Fig. 4e–h), suggesting that arctigenin inhibited hepatic gluconeogenesis and lipid synthesis by activating AMPK.

### Arctigenin inhibits gluconeogenesis and increases AMPK phosphorylation in vivo

To assess the effect of arctigenin on gluconeogenesis in vivo, we performed a pyruvate tolerance test, as administration of the gluconeogenic substrate pyruvate increases blood glucose levels by promoting gluconeogenesis in liver.





**Fig. 1** Arctigenin (ARC-G) increases glucose uptake and AMPK phosphorylation in L6 myotubes. **a** Chemical structure of arctigenin. **b** Increased basal and insulin-stimulated (100 nmol/l) glucose uptake in L6 myotubes after 2 h of treatment with 1  $\mu$ mol/l arctigenin. **c** Dose-dependent increase in glucose uptake by L6 myotubes exposed to arctigenin at the indicated concentrations for 2 h. Berberine (3  $\mu$ mol/l) was set as a positive control. **d** Time-dependent stimulation of glucose uptake by 3  $\mu$ mol/l arctigenin. **e** GLUT4 translocation in L6 myotubes treated by arctigenin or insulin for 2 h. GAPDH, glyceraldehyde-3-

phosphate dehydrogenase; PM, plasma membrane. **f** Dose-dependent increase in AMPK and ACC phosphorylation (p-AMPK, p-ACC) in L6 myotubes by 2 h treatment with arctigenin at the indicated concentrations, quantified (**g**) as relative optical density. White bars, p-AMPK $\alpha$ /AMPK $\alpha$ ; black bars, p-ACC/GAPDH. **h** Time-dependent stimulation of AMPK and ACC phosphorylation by 3  $\mu$ mol/l arctigenin, quantified (**i**) as relative optical density. Key as above (**g**). Results are shown as means $\pm$ SEM from three independent experiments; \* $p$ <0.05 and \*\* $p$ <0.01 vs corresponding controls (as indicated)

As shown in Fig. 5a, b, administration of pyruvate significantly enhanced blood glucose levels in control mice, but this effect was attenuated in mice pre-treated with 300 mg/kg arctigenin by gavage, indicating reduced gluconeogenesis. In addition, the phosphorylation of AMPK and ACC in liver isolated from arctigenin-treated mice was significantly increased by 36% and 74%, respectively (Fig. 5c, d), suggesting that arctigenin activated AMPK in vivo.

Arctigenin activates AMPK indirectly, an action that is not dependent on LKB1, CaMKK or peroxynitrite


AMPK enzymatic activity measurement was performed by monitoring phosphorylation of the AMARA peptide. The results showed that arctigenin did not activate AMPK directly (ESM Fig. 1a). Moreover, arctigenin increased AMPK phosphorylation in HeLa cells lacking LKB1 (ESM

Fig. 1b, c). The increased AMPK phosphorylation and glucose uptake induced by arctigenin was also unaffected by the CaMKK inhibitor, STO-609, and by the nitric oxide synthase (NOS) inhibitor, L-NAME (ESM Fig. 1d–g). These data indicate that arctigenin activated AMPK independently of LKB1, CaMKK, and the peroxynitrite (ONOO<sup>-</sup>) or NOS pathways.

Arctigenin reduces mitochondrial respiration by inhibiting respiratory complex I

Several studies have linked mitochondrial function with AMPK activation [17, 18, 22, 23]. We observed effects of arctigenin on mitochondrial membrane potential ( $\Delta\psi_m$ ). As shown in Fig. 6a, arctigenin depolarised  $\Delta\psi_m$  in a dose-dependent manner, but with relatively mild effect (maximum effect lower than 30%). Arctigenin treatment also dose-dependently increased the AMP/ATP ratio in L6

**Fig. 2** Arctigenin (ARC-G) increases glucose uptake and AMPK phosphorylation in isolated muscles of C57BL/6J mice. **a** Increased basal and insulin-stimulated (12 nmol/l) glucose uptake in EDL and **(b)** soleus muscles by 2 h of treatment with 3  $\mu\text{mol/l}$  arctigenin. White bars, control; black bars, arctigenin. **c** Dose-dependent increase in AMPK and ACC phosphorylation (p-AMPK, p-ACC) in EDL muscles by 1 h of treatment with arctigenin at the indicated concentrations. AICAR (2 mmol/l) was set as a positive control. GAPDH, glyceraldehyde-3-phosphate dehydrogenase. **d** Blot findings **(c)**, quantified as relative optical density. White bars, p-AMPK $\alpha$ /AMPK $\alpha$ ; black bars, p-ACC/GAPDH. **e** Dose-dependent increase in AMPK and ACC phosphorylation in soleus muscles with treatment as above **(c, d)**, quantified **(f)** as relative optical density. Key as above **(d)**. Results are means $\pm$ SEM;  $n=10\text{--}14$  **(a, b)**,  $n=4\text{--}6$  **(c-f)**; \* $p<0.05$  and \*\* $p<0.01$  vs corresponding controls (as indicated)




myotubes (Fig. 6b). To determine whether the change in nucleotide ratio was due to an effect on cellular respiration, we examined oxygen consumption in L6 myotubes and observed a dose-dependent inhibition of respiration (Fig. 6d). Furthermore, the effects of arctigenin on ADP-stimulated respiration in the presence of complex I (glutamate + malate) or complex II (succinate) substrates were examined in isolated liver mitochondria. Arctigenin produced a dose-dependent inhibition of oxygen consumption with complex I-linked substrate, but not with complex II-linked respiration (Fig. 6e). In addition, arctigenin increased lactate production in L6 myotubes, as the reduction of aerobic respiration may lead to compensated elevation of anaerobic respiration (Fig. 6f).

#### Arctigenin ameliorates metabolic disorders in *ob/ob* mice

The acute and chronic glucose-lowering potential of arctigenin (200 mg/kg, orally) was evaluated in *ob/ob*

mice, with metformin (250 mg/kg, orally) included as positive control. A single administration of arctigenin resulted in a statistically significant decrease of blood glucose levels by 34.2% after 2 h, although this effect was less pronounced than that of metformin (decrease 62.3%) (Fig. 7a). Chronic administration of arctigenin for 23 days significantly decreased random and fasting blood glucose levels. During the whole treatment period, administration of 200 mg/kg arctigenin reduced random and fasting blood glucose levels, with an average reduction rate of 32.2% and 35.2%; administration of 250 mg/kg metformin caused average reductions of 22.2% and 46.0%, respectively (Fig. 7b, c). After 20 days of treatment with arctigenin, glucose tolerance improved, an effect similar to that seen with metformin (Fig. 7d). No changes in body weight were observed during arctigenin or metformin treatment (Table 1). However, arctigenin treatment caused a significant reduction of subcutaneous, but not of perirenal or mesenteric fat weight by 21% (Table 1). Arctigenin did






**Fig. 4** Effects of arctigenin (ARC-G) on AMPK phosphorylation (p-AMPK), gluconeogenesis and lipid synthesis in rat primary hepatocytes. **a** Dose-dependent increase in p-AMPK and ACC phosphorylation (p-ACC) by 1 h of treatment with arctigenin at the indicated concentrations. GAPDH, glyceraldehyde-3-phosphate dehydrogenase. **b** Dose-dependent inhibition of glucagon-stimulated (10 nmol/l) gluconeogenesis by 20 h incubation with arctigenin treatment at the indicated concentrations. **c** Dose-dependent inhibition of insulin-stimulated (10 nmol/l) fatty acid and sterol (**d**) synthesis by 24 h of exposure to arctigenin at the indicated concentrations. **e** Rat primary hepatocytes were preincubated for 30 min in the absence or presence of compound C (20 μmol/l) and then treated with

20 μmol/l arctigenin for 20 h, followed by the measurement of glucagon-stimulated (10 nmol/l) gluconeogenesis. **f** Rat primary hepatocytes were preincubated as above (**e**) and then treated with 20 μmol/l arctigenin for 24 h, followed by the measurement of insulin-stimulated (10 nmol/l) fatty acid and (**g**) sterol synthesis. **e–g** White bars, without compound C; black bars, with compound C. **h** Compound C (20 μmol/l) treatment inhibited the ACC phosphorylation induced by arctigenin in rat primary hepatocytes. Values (**b–g**) are means±SEM from three independent experiments; \* $p < 0.05$  and \*\* $p < 0.01$  vs corresponding controls (as indicated); † $p < 0.05$  and †† $p < 0.01$  vs ‘0’ group

Glucose transport in skeletal muscle is the major component of whole-body glucose uptake and plays a key role in maintaining whole-body glucose homeostasis [2]. Here, a screening of our natural compounds library led to an astonishing discovery, namely that arctigenin significantly stimulated glucose uptake in L6 myotubes, an effect observed to occur in addition to the increased glucose uptake induced by insulin. Since AMPK activation is implicated as an insulin-independent mechanism in stimulation of glucose uptake in skeletal muscle [29], we assessed the effect of arctigenin on AMPK phosphorylation in L6 myotubes. As expected, AMPK phosphorylation was significantly enhanced by arctigenin treatment, a finding that was associated with increased phosphorylation of

ACC, a downstream target of AMPK [30], suggesting that activation of AMPK had occurred. In agreement with our studies in L6 myotubes, arctigenin also stimulated glucose uptake, and increased AMPK and ACC phosphorylation in isolated mouse soleus and EDL muscle, providing further evidence of activation of AMPK by arctigenin treatment in skeletal muscle. To investigate the role of AMPK activation in mediating arctigenin-stimulated glucose uptake, inhibitor and siRNA studies were performed in L6 myotubes. Compound C, an AMPK inhibitor, fully blocked arctigenin-stimulated glucose uptake, whereas wortmannin, a PI3 kinase inhibitor, showed no effect. Furthermore, arctigenin-stimulated glucose uptake was completely lost upon silencing of *Ampkα1/α2*. Thus, arctigenin-stimulated






**Fig. 5** Acute effect of arctigenin (ARC-G) on gluconeogenesis and hepatic AMPK phosphorylation (p-AMPK) in C57BL/6J mice. Mice were treated as described in Methods. In the pyruvate tolerance test, blood glucose levels (**a**) were measured and the AUC (**b**) calculated. White circles, vehicle control; black squares, arctigenin. **c** Hepatic p-AMPK and ACC phosphorylation (p-ACC) were analysed by western blot and quantified (**d**) as relative optical density. White bars, vehicle control; black bars, arctigenin. GAPDH, glyceraldehyde-3-phosphate dehydrogenase. Values (**a**, **b**, **d**) are means $\pm$ SEM;  $n=8$ ; \* $p<0.05$  and \*\* $p<0.01$  vs vehicle

glucose uptake in skeletal muscle was mediated by activation of AMPK, which is similar to the effect of


metformin and 5-amino-imidazole carboxamide ribonucleotide (AICAR) [9, 14].

Liver is the major site for storage and release of carbohydrates, and for fatty acid synthesis. Hepatic metabolism plays a key role in the regulation of whole-body energy homeostasis. Elevated hepatic glucose production is a major cause of fasting hyperglycaemia in type 2 diabetic patients [31], while chronic hyperinsulinaemia in the insulin-resistant state results in increased lipid accumulation in the liver [32]. Activation of AMPK in liver leads to the stimulation of fatty acid oxidation, and inhibition of gluconeogenesis and lipogenesis, thus playing important roles in the regulation of glucose and lipid metabolism [15]. Evidence that the activation of AMPK in liver by metformin was required for metformin to exert its inhibitory effect on hepatocyte glucose production further reinforced the importance of AMPK activation in hepatic metabolism [33]. Since arctigenin has been shown to activate AMPK in skeletal muscle, we further studied its effect on liver. As expected, an increase in AMPK and ACC phosphorylation was observed in primary hepatocytes treated with arctigenin, suggesting that activation of AMPK in liver had occurred. Moreover, arctigenin dose-dependently inhibited glucagon-stimulated gluconeogenesis and insulin-stimulated lipid synthesis in rat primary hepatocytes. However, the latter two inhibitory effects of arctigenin were fully abrogated by treatment with com-



**Fig. 6** Effect of arctigenin (ARC-G) on the AMP/ATP ratio and respiration. **a** Dose-dependent depolarisation of  $\Delta\psi_m$  in L6 myotubes by 10 min of arctigenin treatment at the indicated concentrations. Berberine (3  $\mu\text{mol/l}$ ) and carbonyl cyanide *m*-chlorophenyl hydrazone (CCCP; 10  $\mu\text{mol/l}$ ) were set as positive control. **b** Dose-dependent increase of the AMP/ATP ratio in L6 myotubes exposed to arctigenin at the indicated concentrations for 2 h. **c** Time-dependent stimulation of the AMP/ATP ratio in L6 myotubes by 10  $\mu\text{mol/l}$  arctigenin. **d** Dose-dependent inhibition of the respiration of intact L6 myotubes by arctigenin treatment at the indicated concentrations. **e** Effect of

arctigenin on the respiration of isolated mitochondria from rat liver. Arctigenin dose-dependently inhibited mitochondrial respiration in ADP-stimulated respiration in the presence of complex I (glutamate+malate) substrates, but not of complex II (succinate) substrates. **f** Effect of arctigenin (3  $\mu\text{mol/l}$ ) on lactate release in L6 myotubes after 1 h (white bars) or 4 h (black bars) of treatment. Rosiglitazone (50  $\mu\text{mol/l}$ ) was set as a positive control. Values are means $\pm$ SEM;  $n=3$  (**a**),  $n=4$  (**b**, **c**, **f**),  $n=7$  (**d**),  $n=6$  (**e**); \* $p<0.05$ , \*\* $p<0.01$  vs corresponding controls (as indicated)



**Fig. 7** Effect of arctigenin (ARC-G) on metabolic abnormalities of *ob/ob* mice. **a** Arctigenin lowered fasting blood glucose values after a single oral administration. Random-fed blood glucose (**b**) and fasting blood glucose (**c**) were determined throughout the treatment. **d** Glucose tolerance was determined on day 20 of the treatment. White circles, vehicle; black squares, arctigenin (200 mg/kg); black triangles, metformin 250 (mg/kg). **e** AKT phosphorylation (p-AKT) in gastro-

cnemius muscle, liver (**f**) and perirenal fat (**g**) was evaluated. **h** Protein abundance of GLUT4 in gastrocnemius muscle was determined by western blot. **i** Hepatic *Pepck* and *G6pase* mRNA levels were determined by real-time PCR. White bars (**e–g**, **i**), vehicle control; black bars, arctigenin. Values are means $\pm$ SEM for  $n=8–9$  mice; \* $p<0.05$  and \*\* $p<0.01$  vs vehicle-treated mice

**Table 1** Chronic effects of arctigenin and metformin on metabolic variables in *ob/ob* mice

Variable	Treatment		
	Vehicle	Arctigenin	Metformin
Body weight gain (g)	11.2 $\pm$ 1.2	10.1 $\pm$ 0.9	11.7 $\pm$ 0.6
Perirenal fat (g)	0.65 $\pm$ 0.05	0.64 $\pm$ 0.06	0.73 $\pm$ 0.09
Mesenteric fat (g)	2.05 $\pm$ 0.16	2.03 $\pm$ 0.10	1.99 $\pm$ 0.13
Subcutaneous fat (g)	5.51 $\pm$ 0.32	4.53 $\pm$ 0.20 <sup>a</sup>	5.56 $\pm$ 0.27
Serum triacylglycerol (mmol/l)	0.82 $\pm$ 0.04	0.80 $\pm$ 0.07	0.79 $\pm$ 0.04
Serum cholesterol (mmol/l)	4.43 $\pm$ 0.22	3.85 $\pm$ 0.08 <sup>a</sup>	4.97 $\pm$ 0.18
Serum NEFA (mmol/l)	1.13 $\pm$ 0.11	1.16 $\pm$ 0.10	0.97 $\pm$ 0.16
Serum adiponectin ( $\mu$ g/ml)	9.24 $\pm$ 0.55	11.62 $\pm$ 1.45	9.88 $\pm$ 1.08
Serum insulin (pmol/l)	1,695.1 $\pm$ 246.5	2,118.8 $\pm$ 173.3	1,103.8 $\pm$ 182.3 <sup>b</sup>
HOMA-IR	228.3 $\pm$ 26.9	181.4 $\pm$ 29.5 <sup>c</sup>	61.2 $\pm$ 11.8 <sup>b</sup>
Muscle triacylglycerol ( $\mu$ mol/g)	81.01 $\pm$ 4.55	64.68 $\pm$ 3.64 <sup>a</sup>	58.07 $\pm$ 7.17 <sup>a</sup>
Hepatic triacylglycerol ( $\mu$ mol/g)	242.33 $\pm$ 20.20	176.67 $\pm$ 20.09 <sup>a</sup>	163.60 $\pm$ 10.39 <sup>b</sup>
Hepatic cholesterol ( $\mu$ mol/g)	1.09 $\pm$ 0.14	0.99 $\pm$ 0.07	1.01 $\pm$ 0.10

Data represent the mean $\pm$ SEM of eight mice

<sup>a</sup> $p<0.05$  (Student's *t* test for unpaired values with respect to vehicle)

<sup>b</sup> $p<0.01$  (Student's *t* test for unpaired values with respect to vehicle)

<sup>c</sup> $p=0.08$  (Student's *t* test for unpaired values with respect to vehicle)

pound C, demonstrating that AMPK activation was required for arctigenin-mediated effects to take place. In addition, the activation of hepatic AMPK and suppression of gluconeogenesis by arctigenin treatment was also demonstrated *in vivo*.

Although arctigenin activated AMPK in skeletal muscle and liver, it showed no effect on the activation of recombinant AMPK kinase, suggesting that arctigenin activated AMPK indirectly. LKB1 and CaMKK are two upstream kinases of AMPK [34], and a recent study showed that ONOO<sup>-</sup> played a critical role in AMPK activation by metformin in liver and that endothelial NOS was required for metformin action *in vitro* and *in vivo* [35]. However, our results showed that arctigenin activated AMPK independently of LKB1, CaMKK, and the ONOO<sup>-</sup> or NOS pathways.

Since mild perturbation of mitochondrial function has been shown to be associated with activation of AMPK [36–38], the effect of arctigenin on mitochondrial potential was examined. After arctigenin treatment in L6 myotubes, a relatively mild depolarisation of  $\Delta\psi_m$  was observed. Several factors may lead to a fall in  $\Delta\psi_m$ , including inhibition of mitochondrial respiration, induction of proton leakage (uncoupling) [39] or opening of the mitochondrial permeability transition pore. The action of arctigenin was not related to mitochondrial permeability transition pore opening, as no apoptosis was observed after arctigenin treatment (data not shown), whereas mitochondrial permeability transition pore opening was tightly correlated with the initiation of apoptosis [40]. We also ruled out the possibility that arctigenin acts directly as an uncoupling agent, as it did not accelerate the respiration rate in isolated mitochondria, as typical uncoupling agents would do [41]. Thus, decreased  $\Delta\psi_m$  caused by arctigenin is likely to be associated with inhibition of mitochondrial respiration. This hypothesis was further confirmed in L6 myotubes by the dose-dependently inhibited respiration and increase of the AMP/ATP ratio after arctigenin treatment. Moreover, in isolated mitochondria, arctigenin inhibited ADP-stimulated respiration with glutamate and malate as substrates, but had no effect on respiration with succinate as the substrate. This suggests that arctigenin inhibited respiration via a direct effect on mitochondrial respiratory complex I. Therefore, arctigenin activated AMPK by inhibiting mitochondrial respiratory complex I, further leading to an elevated AMP/ATP ratio. This action shares similar mechanisms with several clinically used glucose-lowering drugs such as rosiglitazone and metformin [38, 42, 43], highlighting the prospect of arctigenin as a lead compound for the treatment of type 2 diabetes mellitus.

Whole-body glucose homeostasis is maintained by a balance between glucose production and glucose uptake by peripheral tissues. Activation of AMPK increases glucose

uptake independently of insulin signalling [44] in skeletal muscle and suppresses hepatic glucose production, thus producing a glucose-lowering effect in type 2 diabetes [15]. In the present study, a single oral administration of arctigenin significantly decreased blood glucose levels after 2 h in *ob/ob* mice, but this effect was less pronounced than that of metformin. Our study of chronic administration of arctigenin showed that arctigenin significantly reduced random and fasting blood glucose levels in *ob/ob* mice. Moreover, unlike the weaker effect of arctigenin in the acute study, the glucose-lowering effect of arctigenin (200 mg/kg orally) on random-fed blood glucose was comparable with that of metformin (250 mg/kg), suggesting that arctigenin worked longer than metformin, as the random-fed blood glucose values were measured at 24 h post-dose. The glucose-lowering effect of arctigenin was not due to insulin secretion, since administration of arctigenin to *ob/ob* mice didn't cause insulin secretion upon glucose loading (ESM Fig. 3).

In addition to glucose-lowering, lipid metabolism in *ob/ob* mice was also improved by arctigenin treatment. A significant decrease in serum cholesterol levels was observed in *ob/ob* mice after 23 days of treatment with arctigenin. The triacylglycerol content in liver was also reduced, consistent with previous reports on AICAR, metformin or A-769662 [15, 33]. This reduced lipid accumulation in liver can be explained by arctigenin-induced activation of AMPK, and the subsequent inhibition of hepatic lipogenesis and increase of fatty acid oxidation [45]. In addition, chronic treatment with arctigenin in *ob/ob* mice resulted in a significant reduction of triacylglycerol content in skeletal muscle, possibly due to the activation of AMPK induced by arctigenin, followed by an increase of fatty acid oxidation in skeletal muscle [46]. Since excess lipid accumulation in muscle and liver is associated with insulin resistance in humans [47] and rodents [48], the decrease in hepatic and intramuscular lipid content will subsequently improve insulin sensitivity, further improving glucose metabolism. These findings are further supported by the increased insulin-stimulated AKT phosphorylation in gastrocnemius muscle and liver of *ob/ob* mice after chronic arctigenin treatment. Moreover, hepatic *G6pase* mRNA was significantly reduced in arctigenin-treated mice, suggesting that the decreased expression of gluconeogenic genes may also contribute to the glucose-lowering effect of arctigenin. Taken together, our results indicate that arctigenin improves overall glucose and lipid metabolism in *ob/ob* mice, an effect likely to be mediated by the activation of AMPK in liver and skeletal muscle.

In conclusion, our studies demonstrate the beneficial effects of arctigenin as a potent indirect activator of AMPK on the regulation of glucose and lipid homeostasis. Arctigenin caused inhibition of mitochondrial respiratory complex I, thus

leading to activation of AMPK and subsequent beneficial metabolic outcomes, such as enhanced glucose uptake in skeletal muscle, and suppression of hepatic gluconeogenesis and lipid synthesis. Administration of arctigenin decreased blood glucose, improved glucose tolerance and dyslipidaemia, decreased triacylglycerol accumulation and increased insulin-stimulated AKT phosphorylation in skeletal muscle and liver in *ob/ob* mice. These results highlight the potential value of arctigenin as a potent lead compound for the treatment of type 2 diabetes.

**Acknowledgements** This study was supported by the National Basic Research Program of China (973 Program, 2009CB522300) and National Nature Science Foundation of China (Grant 90813029).

**Contribution statement** S-LH, R-TY, JG, YF, Y-LD, FH, Y-HH, Y-DT and YL all contributed to the conception and design of the study, and contributed to analysis and interpretation of data. All the authors participated in drafting and revising the article and all have approved the final version of the article.

**Duality of interest** The authors declare that there is no duality of interest associated with this manuscript.

## References

- DeFronzo RA, Jacot E, Jequier E, Maeder E, Wahren J, Felber JP (1981) The effect of insulin on the disposal of intravenous glucose. Results from indirect calorimetry and hepatic and femoral venous catheterization. *Diabetes* 30:1000–1007
- Ren JM, Marshall BA, Gulve EA et al (1993) Evidence from transgenic mice that glucose transport is rate-limiting for glycogen deposition and glycolysis in skeletal muscle. *J Biol Chem* 268:16113–16115
- Hansell CA, Schiering C, Kinstry R et al (2011) Universal expression and dual function of the atypical chemokine receptor D6 on innate-like B cells in mice. *Blood* 117:5413–5424
- Kelley DE, Goodpaster B, Wing RR, Simoneau JA (1999) Skeletal muscle fatty acid metabolism in association with insulin resistance, obesity, and weight loss. *Am J Physiol* 277:E1130–E1141
- Leng Y, Steiler TL, Zierath JR (2004) Effects of insulin, contraction, and phorbol esters on mitogen-activated protein kinase signaling in skeletal muscle from lean and *ob/ob* mice. *Diabetes* 53:1436–1444
- Zierath JR, Krook A, Wallberg-Henriksson H (2000) Insulin action and insulin resistance in human skeletal muscle. *Diabetologia* 43:821–835
- Long YC, Zierath JR (2006) AMP-activated protein kinase signaling in metabolic regulation. *J Clin Invest* 116:1776–1783
- Winder WW (2001) Energy-sensing and signaling by AMP-activated protein kinase in skeletal muscle. *J Appl Physiol* 91:1017–1028
- Koistinen HA, Galuska D, Chibalin AV et al (2003) 5-aminoimidazole carboxamide riboside increases glucose transport and cell-surface GLUT4 content in skeletal muscle from subjects with type 2 diabetes. *Diabetes* 52:1066–1072
- Shaw RJ, Kosmatka M, Bardeesy N et al (2004) The tumor suppressor LKB1 kinase directly activates AMP-activated kinase and regulates apoptosis in response to energy stress. *Proc Natl Acad Sci U S A* 101:3329–3335
- Woods A, Dickerson K, Heath R et al (2005) Ca<sup>2+</sup>/calmodulin-dependent protein kinase kinase-beta acts upstream of AMP-activated protein kinase in mammalian cells. *Cell Metab* 2:21–33
- Towler MC, Hardie DG (2007) AMP-activated protein kinase in metabolic control and insulin signaling. *Circ Res* 100:328–341
- Viollet B, Lantier L, Devin-Leclerc J et al (2009) Targeting the AMPK pathway for the treatment of type 2 diabetes. *Front Biosci* 14:3380–3400
- Musi N, Hirshman MF, Nygren J et al (2002) Metformin increases AMP-activated protein kinase activity in skeletal muscle of subjects with type 2 diabetes. *Diabetes* 51:2074–2081
- Cool B, Zinker B, Chiou W et al (2006) Identification and characterization of a small molecule AMPK activator that treats key components of type 2 diabetes and the metabolic syndrome. *Cell Metab* 3:403–416
- Newman DJ, Cragg GM (2007) Natural products as sources of new drugs over the last 25 years. *J Nat Prod* 70:461–477
- Lee YS, Kim WS, Kim KH et al (2006) Berberine, a natural plant product, activates AMP-activated protein kinase with beneficial metabolic effects in diabetic and insulin-resistant states. *Diabetes* 55:2256–2264
- Baur JA, Pearson KJ, Price NL et al (2006) Resveratrol improves health and survival of mice on a high-calorie diet. *Nature* 444:337–342
- Collins QF, Liu HY, Pi J, Liu Z, Quon MJ, Cao W (2007) Epigallocatechin-3-gallate (EGCG), a green tea polyphenol, suppresses hepatic gluconeogenesis through 5'-AMP-activated protein kinase. *J Biol Chem* 282:30143–30149
- Wagner BK, Kitami T, Gilbert TJ et al (2008) Large-scale chemical dissection of mitochondrial function. *Nat Biotechnol* 26:343–351
- Turner N, Li JY, Gosby A et al (2008) Berberine and its more biologically available derivative, dihydroberberine, inhibit mitochondrial respiratory complex I: a mechanism for the action of berberine to activate AMP-activated protein kinase and improve insulin action. *Diabetes* 57:1414–1418
- Zou MH, Kirkpatrick SS, Davis BJ et al (2004) Activation of the AMP-activated protein kinase by the anti-diabetic drug metformin in vivo. Role of mitochondrial reactive nitrogen species. *J Biol Chem* 279:43940–43951
- Fryer LG, Parbu-Patel A, Carling D (2002) The anti-diabetic drugs rosiglitazone and metformin stimulate AMP-activated protein kinase through distinct signaling pathways. *J Biol Chem* 277:25226–25232
- Zhao F, Wang L, Liu K (2009) In vitro anti-inflammatory effects of arctigenin, a lignan from *Arctium lappa* L., through inhibition on iNOS pathway. *J Ethnopharmacol* 122:457–462
- Sun S, Wang X, Wang C et al (2011) Arctigenin suppresses unfolded protein response and sensitizes glucose deprivation-mediated cytotoxicity of cancer cells. *Planta Med* 77:141–145
- Hotamisligil GS (2010) Endoplasmic reticulum stress and the inflammatory basis of metabolic disease. *Cell* 140:900–917
- Deldicque L, Cani PD, Philp A et al (2010) The unfolded protein response is activated in skeletal muscle by high-fat feeding: potential role in the downregulation of protein synthesis. *Am J Physiol Endocrinol Metab* 299:E695–E705
- Ye R, Jung DY, Jun JY et al (2010) Grp78 heterozygosity promotes adaptive unfolded protein response and attenuates diet-induced obesity and insulin resistance. *Diabetes* 59:6–16
- Bergeron R, Russell RR 3rd, Young LH et al (1999) Effect of AMPK activation on muscle glucose metabolism in conscious rats. *Am J Physiol* 276:E938–E944
- Hardie DG (2003) Minireview: the AMP-activated protein kinase cascade: the key sensor of cellular energy status. *Endocrinology* 144:5179–5183
- DeFronzo RA, Ferrannini E, Simonson DC (1989) Fasting hyperglycemia in non-insulin-dependent diabetes mellitus: con-



- tributions of excessive hepatic glucose production and impaired tissue glucose uptake. *Metabolism* 38:387–395
32. Wolfrum C, Asilmaz E, Luca E, Friedman JM, Stoffel M (2004) Foxa2 regulates lipid metabolism and ketogenesis in the liver during fasting and in diabetes. *Nature* 432:1027–1032
  33. Zhou G, Myers R, Li Y et al (2001) Role of AMP-activated protein kinase in mechanism of metformin action. *J Clin Invest* 108:1167–1174
  34. Sanders MJ, Grondin PO, Hegarty BD, Snowden MA, Carling D (2007) Investigating the mechanism for AMP activation of the AMP-activated protein kinase cascade. *Biochem J* 403:139–148
  35. Fujita Y, Hosokawa M, Fujimoto S et al (2010) Metformin suppresses hepatic gluconeogenesis and lowers fasting blood glucose levels through reactive nitrogen species in mice. *Diabetologia* 53:1472–1481
  36. Hayashi T, Hirshman MF, Fujii N, Habinowski SA, Witters LA, Goodyear LJ (2000) Metabolic stress and altered glucose transport: activation of AMP-activated protein kinase as a unifying coupling mechanism. *Diabetes* 49:527–531
  37. Kosaka T, Okuyama R, Sun W et al (2005) Identification of molecular target of AMP-activated protein kinase activator by affinity purification and mass spectrometry. *Anal Chem* 77:2050–2055
  38. Konrad D, Rudich A, Bilan PJ et al (2005) Troglitazone causes acute mitochondrial membrane depolarisation and an AMPK-mediated increase in glucose phosphorylation in muscle cells. *Diabetologia* 48:954–966
  39. Hutchinson DS, Csikasz RI, Yamamoto DL et al (2007) Diphenylene iodonium stimulates glucose uptake in skeletal muscle cells through mitochondrial complex I inhibition and activation of AMP-activated protein kinase. *Cell Signal* 19:1610–1620
  40. Narita M, Shimizu S, Ito T et al (1998) Bax interacts with the permeability transition pore to induce permeability transition and cytochrome c release in isolated mitochondria. *Proc Natl Acad Sci U S A* 95:14681–14686
  41. Harper JA, Dickinson K, Brand MD (2001) Mitochondrial uncoupling as a target for drug development for the treatment of obesity. *Obes Rev* 2:255–265
  42. Brunmair B, Staniek K, Gras F et al (2004) Thiazolidinediones, like metformin, inhibit respiratory complex I: a common mechanism contributing to their antidiabetic actions? *Diabetes* 53:1052–1059
  43. Owen MR, Doran E, Halestrap AP (2000) Evidence that metformin exerts its anti-diabetic effects through inhibition of complex I of the mitochondrial respiratory chain. *Biochem J* 348:607–614
  44. Narkar VA, Downes M, Yu RT et al (2008) AMPK and PPARdelta agonists are exercise mimetics. *Cell* 134:405–415
  45. Viollet B, Guigas B, Leclerc J et al (2009) AMP-activated protein kinase in the regulation of hepatic energy metabolism: from physiology to therapeutic perspectives. *Acta Physiol (Oxf)* 196:81–98
  46. Winder WW, Hardie DG (1996) Inactivation of acetyl-CoA carboxylase and activation of AMP-activated protein kinase in muscle during exercise. *Am J Physiol* 270:E299–E304
  47. Pan DA, Lillioja S, Kriketos AD et al (1997) Skeletal muscle triglyceride levels are inversely related to insulin action. *Diabetes* 46:983–988
  48. Oakes ND, Bell KS, Furler SM et al (1997) Diet-induced muscle insulin resistance in rats is ameliorated by acute dietary lipid withdrawal or a single bout of exercise: parallel relationship between insulin stimulation of glucose uptake and suppression of long-chain fatty acyl-CoA. *Diabetes* 46:2022–2028