

Stéphane Legriél
Bruno Mourvillier
Nicolas Bele
Jose Amaro
Pierre Fouet
Philippe Manet
François Hilpert

Outcomes in 140 critically ill patients with status epilepticus

Published online: 16 November 2007
© Springer-Verlag 2007

The online version of the original can be found at
<http://dx.doi.org/10.1007/s00134-007-0915-5>.

S. Legriél (✉) · N. Bele
Hôpital Saint-Louis, Service de Réanimation Médicale,
1 Avenue Claude Vellefaux, 75010 Paris, France
e-mail: stlegriél@invivo.edu
Tel.: +33-1-42499421
Fax: +33-1-42499426

B. Mourvillier · J. Amaro · P. Fouet · P. Manet · F. Hilpert
Hôpital Robert Ballanger, Service de réanimation polyvalente,
Boulevard Robert Ballanger, 93602 Aulnay-sous-bois cedex, France

Intensive Care Med (2007)
DOI 10.1007/s00134-007-0915-5

Table 2 was published with a shift in the results of the six last lines. The correct version of this table is reproduced here.

Table 2 Seizure classification and causes of status epilepticus. SE, status epilepticus; CNS, central nervous system; ACD, anticonvulsant drug

	All patients	Patients without previous epilepsy	Patients with previous epilepsy
Seizure classification			
Nonconvulsive SE	16 (11)	8 (50)	8 (50)
Electrographic SE	16 (100)		
Convulsive SE	124 (88.6)	63 (88.7)	61 (88.4)
Tonic-clonic generalized	89 (64)	42 (59.1)	47 (68.1)
Partial	27 (19)	16 (22.5)	11 (15.9)
Myoclonic	7 (5)	5 (7)	2 (2.9)
Tonic	1 (1)	0	1 (1.4)
Causes of SE ^a			
Alcohol-related SE	10 (7.1)	4 (5.6)	6 (8.7)
Cerebrovascular disease	24 (17.1)	17 (23.9)	7 (10.1)
CNS infection	6 (4.3)	5 (7)	1 (1.4)
Metabolic disorder	13 (9.3)	12 (16.9)	1 (1.4)
Drug poisoning	7 (5)	6 (8.5)	1 (1.4)
Tumor	3 (2.1)	3 (4.2)	0
Undetermined	49 (35)	24 (33.8)	25 (36.2)
ACD withdrawal	28 (20)	0	28 (40.6)

^a Some patients had more than one diagnosis