



Correction to: Anterolateral ligament reconstruction improves the clinical and functional outcomes of anterior cruciate ligament reconstruction in athletes

Fawzy Hamido^{1,4} · Abdelrahman A. Habiba¹ · Yousef Marwan^{1,2} · Aymen S. I. Soliman³ · Tarek A. Elkhadrawe³ · Mohamed G. Morsy³ · Wael Shoaeb¹ · Ahmed Nagi¹

Published online: 20 August 2020

© European Society of Sports Traumatology, Knee Surgery, Arthroscopy (ESSKA) 2020

Correction to:

Knee Surgery, Sports Traumatology, Arthroscopy
<https://doi.org/10.1007/s00167-020-06119-w>

Authors would like correct the errors in Table 1 and typo in co-author name “Mohamed G. Morsy”. The author name is corrected in author group and the corrected version of Table 1 updated below.

The original article can be found online at <https://doi.org/10.1007/s00167-020-06119-w>.

✉ Fawzy Hamido
fhf19633@yahoo.com

Abdelrahman A. Habiba
aakramh@yahoo.com

Yousef Marwan
yousefmarwan@hotmail.com

Aymen S. I. Soliman
Sssoliman@hotmail.com

Tarek A. Elkhadrawe
elkhadrawe@yahoo.com

Mohamed G. Morsy
morsimoh@gmail.com

Wael Shoaeb
wael_shoaeb@yahoo.com

Ahmed Nagi
nagi_ot@hotmail.com

- ¹ Division of Sports Medicine, Al-Razi Orthopaedic Hospital, Ministry of Health, Kuwait, Kuwait
- ² Division of Orthopaedic Surgery, Department of Surgery, Faculty of Medicine, Health Sciences Center, Kuwait University, Kuwait, Kuwait
- ³ Department of Orthopaedic Surgery, Faculty of Medicine, Alexandria University, Alexandria, Egypt
- ⁴ Department of Orthopaedic Surgery, Al-Razi Orthopaedic Hospital, Safat, P.O. Box 24923, 13110 Kuwait, Kuwait

Table 1 Background characteristics of the patients

	Group A		Group B	
Characteristic	52 (100%)		50 (100%)	
Median age (range) years	(26) 18–40		(24) 18–33	
Sex male/female no	52/0		50/0	
Median duration between injury and operation (range) months	(3) 2.5–3.7		(3) 2.5–3.8	
Median follow-up time (range)	60 (55–65) months			
Patients with Meniscal injury	15 (28.8%)		13 (26.0%)	
Medial	10		8	
	P.M.	M.S.	P.M.	M.S.
	8	2	7	1
Lateral	6		5	
	P.M.	M.S.	P.M.	M.S.
	4	2	4	1
Both	1		0	
	P.M.	M.S.	P.M.	M.S.
	2	–	–	–
Patients available for final assessment	52		50	

ACL anterior cruciate ligament, *ALL* anterolateral ligament, *P.M.* partial meniscectomy *M.S.* meniscal sutures

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.