

Three hierarchies of simple games parameterized by “resource” parameters

Tatiana Gvozdeva · Lane A. Hemaspaandra · Arkadii Slinko

Accepted: 23 October 2011 / Published online: 23 November 2011
© Springer-Verlag 2011

Abstract This paper contributes to the program of numerical characterization and classification of simple games outlined in the classic monograph of von Neumann and Morgenstern. We suggest three possible ways to classify simple games beyond the classes of weighted and roughly weighted games. To this end we introduce three hierarchies of games and prove some relationships between their classes. We prove that our hierarchies are true (i.e., infinite) hierarchies. In particular, they are strict in the sense that more of the key “resource” (which may, for example, be the size or structure of the “tie-breaking” region where the weights of the different coalitions are considered so close that we are allowed to specify either winningness or nonwinningness of the coalition) yields the flexibility to capture strictly more games.

Keywords Simple game · Weighted majority game · Roughly weighted game · Hierarchy

1 Introduction

A simple game is a mathematical object that is used in economics and political science to describe the distribution of power among coalitions of players

T. Gvozdeva (✉) · A. Slinko
Department of Mathematics, University of Auckland, Private Bag 92019, Auckland, New Zealand
e-mail: t.gvozdeva@math.auckland.ac.nz

A. Slinko
e-mail: a.slinko@auckland.ac.nz

L. A. Hemaspaandra
Department of Computer Science, University of Rochester, Rochester, NY 14627-0226, USA
URL: <http://www.cs.rochester.edu/~lane>

(von Neumann and Morgenstern 1944; Shapley 1962). Recently, simple games have been studied as access structures of secret sharing schemes (Blakley 1979). They have also appeared, in some cases under other names, in a variety of mathematical and computer science contexts, e.g., in threshold logic (Muroga 1971). Simple games are closely related to hypergraphs, coherent structures, Sperner systems, clutters, and abstract simplicial complexes. The term “simple” was introduced by von Neumann and Morgenstern (1944), because in this type of games players strive not for monetary rewards but for power, and each coalition is either all-powerful or completely ineffectual. However these games are far from being simple.

An important class of simple games—well studied in economics—is the weighted majority game (von Neumann and Morgenstern 1944; Shapley 1962). In such a game every player is assigned a real number, his or her weight. The winning coalitions are the sets of players whose weights total at least q , a certain threshold. However, it is well known that not every simple game has a representation as a weighted majority game (von Neumann and Morgenstern 1944). The first step in attempting to characterize nonweighted games was the introduction of the class of roughly weighted games (Taylor and Zwicker 1999). Formally, a simple game G on the player set $P = [n] = \{1, 2, \dots, n\}$ is *roughly weighted* if there exist nonnegative real numbers w_1, \dots, w_n and a real number q , called the *quota*, not all equal to zero, such that for $X \in 2^P$ the condition $\sum_{i \in X} w_i > q$ implies X is winning, and $\sum_{i \in X} w_i < q$ implies X is losing. This concept realizes a very common idea in social choice that sometimes a rule needs an additional “tie-breaking” procedure that helps to decide the outcome if the result falls on a certain “threshold.” Taylor and Zwicker (1999) demonstrated the usefulness of this concept. Rough weightedness was studied by Gvozdeva and Slinko (2011), where it was characterized in terms of trading transforms, similar to the characterization of weightedness by Elgot (1961) and Taylor and Zwicker (1992).

Before moving on, it is worth mentioning in passing the notion of complete games. In a simple game player, i is said to be at least as desirable as player j (as a coalition partner) if replacing i in a winning coalition with j never makes that coalition losing. This desirability relation was introduced and studied by Isbell (1956). Weighted majority games have the property that players are totally ordered by the desirability relation. Thus another natural extension of the class of weighted majority games is the class of complete games, for which the desirability relation is a total order. This class is significantly larger than the class of weighted majority games since it contains simple games of any dimension (Freixas and Puente 2008) while, the dimension of a weighted game is always 1. Extensive theoretical and computational results on complete simple games have been obtained by Freixas and Molinero (2009). The strictness of our hierarchies for that important class of simple games is an interesting open question.

It might seem that nonweighted games and even games without rough weights are somewhat strange. However, an important observation of (von Neumann and Morgenstern, 1944, Sect. 53.2.6) states that they “correspond to a different organizational principle that deserves closer study.” In some of these games, as von Neumann and Morgenstern noted, all the minimal winning coalitions are minorities and at the same time “no player has any advantage over any other” (e.g., the Fano game introduced later). This is an attractive feature for secret sharing since in the case of a large number of users it is advantageous to keep minimal authorized coalitions relatively small. This

may be why weighted threshold secret sharing schemes were largely ignored and were characterized only recently (Beimel et al. 2005).

The parameter of the first of the three hierarchies we will discuss reflects the balance of power between small and large coalitions; the larger this parameter the more powerful some of the small coalitions are. Gvozdeva and Slinko (2011) proved that for a game G that is not roughly weighted there exists a certificate of nonweightedness (see the definition in Sect. 2) of the form

$$\mathcal{T} = (X_1, \dots, X_j, P; Y_1, \dots, Y_j, \emptyset), \quad (1)$$

where X_1, \dots, X_j are winning coalitions of G , P is the grand coalition, and Y_1, \dots, Y_j are losing coalitions. However, sometimes it is possible to have more than one grand coalition in the certificate. This may occur when coalitions X_1, \dots, X_j are small but nonetheless winning.

A certificate of nonweightedness of the form

$$\mathcal{T} = (X_1, \dots, X_j, P^\ell; Y_1, \dots, Y_j, \emptyset^\ell) \quad (2)$$

will be called ℓ -potent of length $j + \ell$. Each game that possesses such a certificate will be said to belong to the class of games \mathcal{A}_q , where $q = \ell / (j + \ell)$. The parameter q can take values in the open interval $(0, \frac{1}{2})$. We will show that $\mathcal{A}_p \supseteq \mathcal{A}_q$ for any p and q such that $0 < p \leq q < \frac{1}{2}$, and that if $0 < p < q < \frac{1}{2}$, then this inclusion is strict, i.e., we have the $\mathcal{A}_p \supsetneq \mathcal{A}_q$.

Another hierarchy emerges when we allow several thresholds instead of just one in the case of roughly weighted games. We say that a simple game G belongs to the class \mathcal{B}_k , $k \in \{1, 2, 3, \dots\}$, if there are k thresholds, $0 < q_1 \leq q_2 \leq \dots \leq q_k$, and any coalition with total weight of players smaller than q_1 is losing, any coalition with total weight greater than q_k is winning. We also impose an additional condition that, if a coalition X has total weight $w(X)$ that satisfies $q_1 \leq w(X) \leq q_k$, then $w(X) = q_i$ for some i . All games of the class \mathcal{B}_1 are roughly weighted. In fact, as we will prove in Section 4, almost all roughly weighted games belong to this class: \mathcal{B}_1 is exactly the class of roughly weighted games with nonzero quota. We will show that the Fano game (Gvozdeva and Slinko 2011) belongs to \mathcal{B}_2 but does not belong to \mathcal{B}_1 . We will prove that the \mathcal{B} -hierarchy is strict, that is,

$$\mathcal{B}_1 \subsetneq \mathcal{B}_2 \subsetneq \dots \subsetneq \mathcal{B}_\ell \subsetneq \dots,$$

with the union of these classes being the class of all simple games.

Yet another way to capture more games is by making the threshold “thicker.” We here will not use a point but rather an interval $[a, b]$ for the threshold, $a \leq b$. That is, all coalitions with total weight less than a will be losing and all coalitions whose total weight is greater than b will be winning. This time—in contrast with the k limit of \mathcal{B}_k —we do not care how many different values weights of coalitions falling in $[a, b]$ may take on. (A good example of this situation would be a faculty vote, where if neither side controls a 2/3 majority—calculated in faculty members or their grant dollars—then the Dean would decide the outcome as he wished.) We can keep weights

normalized so that the lower end of the interval is fixed at 1. Then the right end of the interval α becomes a “resource” parameter. Formally, a simple game G belongs to the class C_α if all coalitions in G with total weight less than 1 are losing and every coalition whose total weight is greater than α is winning. We show that the class of all simple games is split into a hierarchy of classes of games $\{C_\alpha\}_{\alpha \in [1, \infty)}$ defined by this parameter. We show that as α increases we get strictly greater descriptive power, i.e., strictly more games can be described, that is, if $\alpha < \beta$, then $C_\alpha \subsetneq C_\beta$. In this sense the hierarchy is strict. This strict hierarchy result, and our strict hierarchy results for hierarchies \mathcal{A} and \mathcal{B} , have very much the general flavor of hierarchy results found in computer science: more resources yield more power (whether computational power to accept languages as in a deterministic or nondeterministic time hierarchy theorem, or as is the case here, description flexibility to capture more games).

The strictness of the latter hierarchy was achieved because we allowed games with arbitrary (but finite) numbers of players. The situation will be different if we keep the number of players, n , fixed. Then there is an interval $[1, s(n)]$ such that all games with n players belong to $C_{s(n)}$ and $s(n)$ is minimal with this property. There will be also finitely many numbers $q \in [1, s(n)]$ such that the interval $[1, q]$ represents more n -player games than any interval $[1, q']$ with $q' < q$. We call the set of such numbers the n th *spectrum* and denote it $\text{Spec}(n)$. We also call a game with n players *critical* if it belongs to C_α with $\alpha \in \text{Spec}(n)$ but does not belong to any C_β with $\beta < \alpha$. We calculate the spectrum for $n < 7$ and also produce a set of critical games, one for each element of the spectrum. We in addition try to give a reasonably tight upper bound for $s(n)$.

All three of our hierarchies provide measures of how close a given game is to being a simple weighted voting game. That is, they each quantify the nearness to being a simple weighted voting game (e.g., hierarchies \mathcal{B} and \mathcal{C} quantify based on the extent and structure of a “flexible tie-breaking” region). And the main theme and contribution of this paper is that we prove for each of the three hierarchies that allowing more quantitative distance from simple weighted voting games yields *strictly more* games, i.e., the hierarchies are proper hierarchies.

2 Preliminaries

Definition 1 A simple game is a pair $G = (P, W)$, where W is a subset of the power set 2^P satisfying the monotonicity condition:

if $X \in W$ and $X \subsetneq Y \subseteq P$, then $Y \in W$,

and $W \not\subseteq \{\emptyset, 2^P\}$ (nontriviality assumption).

Elements of the set W are called *winning coalitions*. We also define the set $L = 2^P \setminus W$ and call elements of this set *losing coalitions*. A winning coalition is said to be *minimal* if each of its proper subsets is a losing coalition. Due to monotonicity, every simple game is fully determined by the set of its minimal winning coalitions. A player who does not belong to any minimal winning coalitions is called a *dummy* (or a *null player*).

For $X \subseteq P$ we will denote the complement of X , $P \setminus X$, as X^c .

Definition 2 A simple game is called proper if $X \in W$ implies that $X^c \in L$ and is called strong if $X \in L$ implies that $X^c \in W$. A simple game that is proper and strong is called a constant-sum game.

The following definition is framed the same way as it was in [Gvozdeva and Slinko \(2011\)](#).

Definition 3 A simple game $G = (P, W)$ is called roughly weighted if there exist nonnegative real numbers w_1, \dots, w_n and a nonnegative real number q , not all equal to zero, such that for $X \in 2^P$ the condition $\sum_{i \in X} w_i < q$ implies $X \in L$ and $\sum_{i \in X} w_i > q$ implies $X \in W$. We say that $[q; w_1, \dots, w_n]$ is a rough voting representation for G ; the number q is called the *quota*.

Example 1 (The Fano game) This important example first appeared in ([von Neumann and Morgenstern, 1944](#), Sect. 53.2.6). Let $P = [7]$ be identified with the set of seven points of the projective plane of order two, called the Fano plane. Let us take the seven lines of this projective plane as minimal winning coalitions:

$$\{1, 2, 3\}, \{3, 4, 5\}, \{1, 5, 6\}, \{1, 4, 7\}, \{2, 5, 7\}, \{3, 6, 7\}, \{2, 4, 6\}. \quad (3)$$

We will denote them by X_1, \dots, X_7 , respectively. This, as is easy to check, defines a constant-sum game. That game is called the *Fano game*. As we will see later, it has no rough voting representation. As we can see from the list of minimal winning coalitions they are all minorities, yet symmetry makes all players equal in this example.

We remind the reader that a sequence of coalitions

$$\mathcal{T} = (X_1, \dots, X_j; Y_1, \dots, Y_j) \quad (4)$$

is a trading transform ([Taylor and Zwicker 1999](#)) if the coalitions X_1, \dots, X_j can be converted into the coalitions Y_1, \dots, Y_j by rearranging players. This can also be expressed as

$$|\{i : a \in X_i\}| = |\{i : a \in Y_i\}| \quad \text{for all } a \in P.$$

We say that the length of \mathcal{T} is j .

Definition 4 A trading transform $(X_1, \dots, X_j; Y_1, \dots, Y_j)$ with all coalitions X_1, \dots, X_j winning and all coalitions Y_1, \dots, Y_j losing is called a certificate of non-weightedness. This certificate is said to be potent if the grand coalition P is among X_1, \dots, X_j and the empty coalition is among Y_1, \dots, Y_j .

Elgot proved (using a different terminology) that the existence of a certificate of nonweightedness implies that the game in question is not weighted and that every nonweighted game has a certificate of nonweightedness. [Taylor and Zwicker \(1999\)](#) showed that for a nonweighted game with n players a certificate of nonweightedness exists whose length is at most 2^{2^n} ; [Gvozdeva and Slinko \(2011\)](#) lowered this bound to $(n+1)2^{\frac{1}{2}n \log_2 n}$.

Theorem 1 (Criterion of rough weightedness (Gvozdeva and Slinko 2011)) *A simple game G with n players is roughly weighted if and only if for no positive integer $j \leq (n+1)2^{\frac{1}{2}n \log_2 n}$ does there exist a potent certificate of nonweightedness of length j .*

In Example 1 the following eight winning coalitions— X_1, \dots, X_7, P —of the Fano game can be transformed into the following eight losing coalitions: $X_1^c, \dots, X_7^c, \emptyset$. So the sequence

$$(X_1, \dots, X_7, P; X_1^c, \dots, X_7^c, \emptyset) \quad (5)$$

is a potent certificate of nonweightedness for this game. So the game is not roughly weighted, thanks to Theorem 1.

As one might expect, games with a very small number of players, or a slightly less small number of players but certain additional properties, are roughly weighted.

Theorem 2 (Gvozdeva and Slinko 2011)

- (a) *Each game with 4 or fewer players is roughly weighted.*
- (b) *Each strong or proper game with 5 or fewer players is roughly weighted.*
- (c) *Each constant sum game with 6 or fewer players is roughly weighted.*

Definition 5 (Taylor and Zwicker 1999, p. 6) We say that a player p in a game is a *dictator* if p belongs to every winning coalition and to no losing coalition. If all coalitions containing player p are winning, this player is called a *passer*. A player p is called a *vetoer* if p is contained in the intersection of all winning coalitions.

Proposition 1 (Gvozdeva and Slinko 2011) *Suppose G is a simple game with n players. Then G is roughly weighted if any of the following three conditions hold:*

- (a) *G has a passer.*
- (b) *G has a vetoer.*
- (c) *G has a losing coalition that consists of $n - 1$ players.*

Due to Proposition 1(a), there is a trivial way to make any game roughly weighted. This can be done by adding an additional player and making him or her a passer. Then we can introduce rough weights by assigning weight 1 to the passer and weight 0 to every other player and setting the quota equal to 0. Note that if the original game is not roughly weighted, then such rough representation is unique (up to multiplicative scaling). In our view, adding a passer trivializes the game but does not make it closer to a weighted majority game; this is why in definitions of our hierarchies \mathcal{B} and \mathcal{C} we do not allow 0 as a threshold value.

As in Gvozdeva and Slinko (2011) we would like to represent trading transforms algebraically. $\{-1, 0, 1\}^n$ denotes the Cartesian product of n copies of $\{-1, 0, 1\}$. For any pair (X, Y) of subsets $X, Y \in [n]$ we define

$$\mathbf{v}_{X,Y} = \chi(X) - \chi(Y) \in \{-1, 0, 1\}^n,$$

where $\chi(X)$ and $\chi(Y)$ are the characteristic vectors of subsets X and Y , respectively.

Let now $G = (P, W)$ be a simple game. We will associate an algebraic object with G . For any pair (X, Y) , where X is winning and Y is losing, we put the pair in correspondence with the vector $\mathbf{v}_{X,Y}$. The set of all such vectors will be denoted $I(G)$ and will be called the ideal of the game. Saying that $(X_1, \dots, X_j; Y_1, \dots, Y_j)$ is a certificate of nonweightedness is equivalent to saying that the following vector sum of the ideal is $\mathbf{0}$: $\mathbf{v}_{X_1,Y_1} + \mathbf{v}_{X_2,Y_2} + \dots + \mathbf{v}_{X_j,Y_j} = \mathbf{0}$. An ℓ -potent certificate $(X_1, \dots, X_j, P^\ell; Y_1, \dots, Y_j, \emptyset^\ell)$ will be represented as

$$\mathbf{v}_{X_1,Y_1} + \mathbf{v}_{X_2,Y_2} + \dots + \mathbf{v}_{X_j,Y_j} + \ell \cdot \mathbf{1} = \mathbf{0},$$

where $\mathbf{1}$ is a vector whose entries are each 1.

3 The \mathcal{A} -hierarchy

This hierarchy of classes \mathcal{A}_α tries to capture the richness of the class of games that do not have rough weights, and does so by introducing a parameter $\alpha \in (0, \frac{1}{2})$. Our method of classification is based on the existence of potent certificates of nonweightedness for such games (Gvozdeva and Slinko 2011). We will now show that potent certificates can be further classified. We will extract a very important parameter from this classification.

Definition 6 A certificate of nonweightedness

$$\mathcal{T} = (X_1, \dots, X_m; Y_1, \dots, Y_m)$$

is called an ℓ -potent certificate of length m if it contains at least ℓ grand coalitions among X_1, \dots, X_m and at least ℓ empty sets among Y_1, \dots, Y_m .

Obviously, every ℓ -potent certificate of length m is also an ℓ' -potent certificate of the same length for any $\ell' < \ell$.

Definition 7 Let q be a rational number. A game G belongs to the class \mathcal{A}_q of the \mathcal{A} -hierarchy if G possesses some ℓ -potent certificate of nonweightedness of length m , such that $q = \ell/m$. If α is irrational, we set $\mathcal{A}_\alpha = \bigcap_{\{q: q < \alpha \text{ and } q \text{ is rational}\}} \mathcal{A}_q$.

It is easy to see that, if $q \geq \frac{1}{2}$, then \mathcal{A}_q is empty. Why? Well, suppose $q \geq \frac{1}{2}$ and \mathcal{A}_q is not empty. Then there is a game G with a certificate of nonweightedness

$$\mathcal{T} = (X_1, \dots, X_k, P^m; Y_1, \dots, Y_k, \emptyset^m) \tag{6}$$

with $m \geq k$. This is not possible since m copies of P contain more elements than are contained in the sets Y_1, \dots, Y_k taken together and so (6) is not a trading transform. So our hierarchy consists of a family of classes $\{\mathcal{A}_\alpha\}_{\alpha \in (0, \frac{1}{2})}$. We would like to show that this hierarchy is strict, that is, a smaller parameter captures more games. The following proposition starts us on our path toward showing this, and after proving this proposition we will soon establish actual strictness when α is strictly smaller than β .

Proposition 2 *If $0 < \alpha \leq \beta < \frac{1}{2}$, then $\mathcal{A}_\alpha \supseteq \mathcal{A}_\beta$.*

Proof It is sufficient to prove this statement when α and β are rational. Suppose that we have a game G in \mathcal{A}_β that possesses a certificate of length n_1 with k_1 grand coalitions and $\beta = k_1/n_1$. Let $\alpha = k_2/n_2$. We can then represent these numbers as $\beta = m_1/n$ and $\alpha = m_2/n$, where $n = \text{lcm}(n_1, n_2)$. Since $\alpha \leq \beta$, we have $m_2 \leq m_1$. Since $n = n_1h$ and $m_1 = k_1h$ for some integer h , we can now combine h certificates for G to obtain one with length n and m_1 grand coalitions. As $m_1 \geq m_2$ we will get a certificate for G of length n with m_2 grand coalitions. So $G \in \mathcal{A}_\alpha$. \square

We say that a game G is *critical* for \mathcal{A}_α if it belongs to \mathcal{A}_α but does not belong to any \mathcal{A}_β with $\beta > \alpha$.

Theorem 3 *For every rational $\alpha \in (0, \frac{1}{2})$ there exists a critical game $G \in \mathcal{A}_\alpha$.*

Proof First, we will construct a two-parameter family of simple games. For any integers $a \geq 2$ and $b \geq 2$, let $G = ([a^2 + a + b + 1], W)$ be a simple game for which a coalition X is winning exactly if $|X| > a^2 + 1$ or X contains a subset whose characteristic vector is a cyclic permutation of $(\underbrace{1, \dots, 1}_{a+1}, \underbrace{0, \dots, 0}_{a^2+b})$.

Let $X_1, \dots, X_{a^2+a+b+1}$ be winning coalitions, whose characteristic vectors are cyclic permutations of $(\underbrace{1, \dots, 1}_{a+1}, \underbrace{0, \dots, 0}_{a^2+b})$. Also let $Y_1, \dots, Y_{a^2+a+b+1}$ be losing coalitions, whose characteristic vectors are cyclic permutations of

$$(\underbrace{1, \dots, 1}_a, 0, | \underbrace{1, \dots, 1}_a, 0, | \underbrace{1, \dots, 1}_a, 0, | \dots, | \underbrace{1, \dots, 1}_a, 0, | 0, 1, \underbrace{0, \dots, 0}_{b-1}),$$

where there are a groups of symbols $\underbrace{1, \dots, 1}_a, 0$. Regarding the $b - 1$ of the rightmost part, it is important to keep in mind that $\underbrace{a}_{b-1} \geq 1$.

This game possesses the following potent certificate of nonweightedness

$$\mathcal{T} = (X_1, \dots, X_{a^2+a+b+1}, P^{a^2-a}; Y_1, \dots, Y_{a^2+a+b+1}, \emptyset^{a^2-a}). \tag{7}$$

One can see that \mathcal{T} is a valid potent certificate. By symmetry losing coalitions in \mathcal{T} each contain $a^2 + 1$ copies of every player and winning coalitions $X_1, \dots, X_{a^2+a+b+1}$ have only $a + 1$ copies of every player. Hence we need to add $a^2 - a$ grand coalitions to make it a trading transform. Clearly the condition $a \geq 2$ is necessary, because otherwise the certificate \mathcal{T} will not be potent.

So $G \in \mathcal{A}_{\frac{a^2-a}{2a^2+b+1}}$. Let us prove that G is critical for this class, that is, it does not belong to any $\mathcal{A}_{q'}$ for $q' > q$. Note that the vectors $\mathbf{v}_i = \mathbf{v}_{X_i, Y_i}$ belong to the ideal of this game. Note also that the sum of all coefficients of \mathbf{v}_i is $\mathbf{v}_i \cdot \mathbf{1} = a - a^2$ and that for any other vector $\mathbf{v} \in I(G)$ from the ideal of this game we have $\mathbf{v} \cdot \mathbf{1} \geq a - a^2$.

Suppose G also has a potent certificate of nonweightedness

$$(A_1, \dots, A_s, P^t; B_1, \dots, B_s, \emptyset^t), \tag{8}$$

with $q' = \frac{t}{t+s} > \frac{a^2-a}{2a^2+b+1} = q$. The latter is equivalent to $\frac{a^2+q+b+1}{a^2-a} > \frac{s}{t}$. Let $\mathbf{u}_i = \mathbf{v}_{A_i, B_i} \in I(G)$; then (8) can be written as

$$\mathbf{u}_1 + \mathbf{u}_2 + \cdots + \mathbf{u}_s + t \cdot \mathbf{1} = \mathbf{0}.$$

Since $\mathbf{u}_i \cdot \mathbf{1} \geq a - a^2$, taking the dot product of both sides with $\mathbf{1}$ we get $t(a^2 + a + b + 1) \leq s(a^2 - a)$, which is equivalent to $\frac{a^2+q+b+1}{a^2-a} \leq \frac{s}{t}$, so we have reached a contradiction.

We will now show that any rational number between 0 and $\frac{1}{2}$ is representable as $\frac{a^2-a}{2a^2+1+b}$ for some positive integers $a \geq 2$ and $b \geq 2$. Let $\frac{p}{q} \in (0, \frac{1}{2})$. Then $q - 2p > 0$ and it is possible to choose a positive integer k such that $k^2 p(q - 2p) - kq - 3 > 0$. By the choice of k one can see that $kp > 1 + \frac{3+2pk}{k(q-2p)} \geq 2$. Take $a = kp$ and $b = k^2 p(q - 2p) - kq - 1$. Substituting these values we get $\frac{a^2-a}{2a^2+1+b} = \frac{p}{q}$. \square

Corollary 1 *If $0 < \alpha < \beta < \frac{1}{2}$, then $\mathcal{A}_\alpha \supsetneq \mathcal{A}_\beta$.*

Example 2 Let us illustrate this proof by an example. Suppose a game G is defined on the set of players $P = [10]$ with $a = 2$ and $b = 3$. Let us include in W all sets of cardinality greater than five and all coalitions with three consecutive players (we think of players as situated on the circle so that 10 and 1 are neighbors). The 3-player minimal winning coalitions, denoted X_1, \dots, X_{10} , are

$$\begin{aligned} &\{1, 2, 3\}, \{2, 3, 4\}, \{3, 4, 5\}, \{4, 5, 6\}, \{5, 6, 7\}, \\ &\{6, 7, 8\}, \{7, 8, 9\}, \{8, 9, 10\}, \{9, 10, 1\}, \{10, 1, 2\}. \end{aligned}$$

Let Y_1, \dots, Y_{10} be the losing coalitions, whose characteristic vectors are cyclic permutations of $(1, 1, 0, 1, 1, 0, 0, 1, 0, 0)$, respectively. For example, $Y_1 = \{1, 2, 4, 5, 8\}$, $Y_2 = \{2, 3, 5, 6, 9\}$, and $Y_{10} = \{10, 1, 3, 4, 7\}$.

Then the potent certificate of nonweightedness

$$\mathcal{T} = (X_1, \dots, X_{10}, P^2; Y_1, \dots, Y_{10}, \emptyset^2)$$

shows that this game belongs to $\mathcal{A}_{1/6}$. As we saw in the proof of Theorem 3, for no β satisfying $1/6 < \beta < 1/2$ does G belong to \mathcal{A}_β .

As was mentioned before, the larger parameter α is, the more relatively “small” winning coalitions and relatively “large” losing coalitions the game has. To see this, consider a simple game $G = ([n], W)$ with an ℓ -potent certificate of length $j + \ell$,

$$\mathcal{T} = (X_1, \dots, X_j, P^\ell; Y_1, \dots, Y_j, \emptyset^\ell).$$

Set $\alpha = \ell/(\ell + j)$. Then $G \in \mathcal{A}_\alpha$, and the average number of players in winning coalitions X_1, \dots, X_j is σ/j where $\sigma = \sum_{i \in [j]} |X_i|$. At the same time the average number of players in losing coalitions Y_1, \dots, Y_j is $(\sigma + n\ell)/j$. On average a losing coalition in \mathcal{T} contains $n\ell/j$ more players than a winning coalition in \mathcal{T} . From above

we know that $j > \ell$. The bigger $\ell/(\ell + j)$ is the bigger the ratio ℓ/j is and hence the bigger $n\ell/j$ is. This means that when α is increasing some winning coalitions become smaller and some losing coalitions become larger.

4 \mathcal{B} -hierarchy

The \mathcal{B} -hierarchy generalizes the idea behind rough weightedness to allow more “points of (decision) flexibility.”

Definition 8 A simple game $G = (P, W)$ belongs to \mathcal{B}_k if there exist real numbers $0 < q_1 \leq q_2 \leq \dots \leq q_k$, called thresholds, and a weight function $w: P \rightarrow \mathbb{R}^{\geq 0}$ such that

- (a) if $\sum_{i \in X} w(i) > q_k$, then X is winning,
- (b) if $\sum_{i \in X} w(i) < q_1$, then X is losing,
- (c) if $q_1 \leq \sum_{i \in X} w(i) \leq q_k$, then $w(X) = \sum_{i \in X} w(i) \in \{q_1, \dots, q_k\}$.

Games from \mathcal{B}_k will be sometimes called k -rough.

The condition $0 < q_1$ in Definition 8 is essential. If we allow the first threshold q_1 be zero, then every simple game can be represented as a 2-rough game. To do this we assign weight 1 to the first player and 0 to everyone else. It is also worthwhile to note that adding a passer does not change the class of the game, that is, a game G belongs to \mathcal{B}_k if and only if the game G' obtained from G by adding a passer belongs to \mathcal{B}_k . This is because a passer can be assigned a very large weight. Thus \mathcal{B}_1 consists of the roughly weighted simple games with nonzero quota.

Example 3 We know that the Fano game is not roughly weighted. Let us assign weight 1 to every player of this game and select two thresholds, $q_1 = 3$ and $q_2 = 4$. Then each coalition whose weight falls below the first threshold is in L , and each coalition whose total weight exceeds the second threshold is in W . If a coalition has total weight of three or four, i.e., its weight is equal to one of the thresholds, it can be either winning or losing. Thus the Fano is a 2-rough game.

Example 4 Let $n = 8$ and assume we have four types of players with players $2i - 1$ and $2i$ forming the i th type. Let us include in W all sets that contain two elements from the same type. Minimal winning coalitions for this game are $\{1, 2\}, \{3, 4\}, \{5, 6\}, \{7, 8\}$. The trading transform

$$\mathcal{T} = (\{1, 2\}^2, \{3, 4\}^2, \{5, 6\}^2, \{7, 8\}^2, P; \{1, 3, 5, 7\}^3, \{2, 4, 6, 8\}^3, \emptyset^3)$$

is the potent certificate of nonweightedness. So by Theorem 1, G is not roughly weighted.

On the other hand, if we assign weight 1 to every player, then G is a 3-rough game with thresholds 2, 3, and 4. Let us show that G is not 2-rough. Assume, to the contrary, that we have weights w_1, \dots, w_8 and two positive thresholds q_1 and q_2 that make this game 2-rough. Without loss of generality we can assume that $w_{2i-1} \geq w_{2i}$ for every type i . Two players of the same type form a winning coalition. This means that the

weight of this coalition is at least q_1 . Moreover the players w_1, w_3, w_5, w_7 with the biggest weight in each type have weight not smaller than $\frac{q_1}{2}$ each. Let us consider the following three losing coalitions with strictly increasing weights

$$\{1, 3\} \subsetneq \{1, 3, 5\} \subsetneq \{1, 3, 5, 7\}.$$

The coalition $\{1, 3\}$ has weight at least $\frac{q_1}{2} + \frac{q_1}{2} = q_1$. In the worst-case scenario this coalition lies exactly on the first threshold q_1 . So the weight of coalition $\{1, 3, 5\}$ is greater than or equal to q_2 . We can see that in every possible scenario the losing coalition $\{1, 3, 5, 7\}$ has weight strictly greater than q_2 , a contradiction.

Let us generalize the idea of Example 4.

Theorem 4 *For every natural number $k \in \mathbb{N}^+$, there exists a game in $\mathcal{B}_{k+1} \setminus \mathcal{B}_k$.*

Proof We will construct a simple game that is a $(k + 1)$ -rough but not k -rough. Let $G_{k+1,n} = ([n], W)$ be a simple game with $n = 2k + 4$ players. We have $k + 2$ types of players with the i th type consisting of two elements $2i - 1$ and $2i$. The set of minimal winning coalitions of this game is $W^m = \{2i - 1, 2i \mid i = 1, 2, \dots, k + 2\}$.

If we assign weight 1 to every player, then $G_{k+1,n}$ is $(k + 1)$ -rough game with thresholds $q_1 = 2, q_2 = 3, \dots, q_{k+1} = k + 2$. Let us assume that this game is j -rough for some $j < k + 1$, and let w be the new weight function. Without loss of generality we may assume that the players are ordered so that $w(2i - 1) \geq w(2i)$. Since the coalition $\{2i - 1, 2i\}$ is winning we have $w(2i - 1) \geq q_1/2 > 0$ for any $i = 1, 2, \dots, k + 2$. The coalitions $L_j = \{2i - 1 \mid 1 \leq i \leq j\}$ are losing, their weights are different, and each of them has weight at least $j q_1/2$ for all $2 \leq j \leq k + 2$. Thus at least $k + 1$ coalitions of different weights lie in the tie-breaking region. This is a contradiction. Thus $G_{k+1,n}$ is not j -rough for any $j < k + 1$. \square

An obvious upper bound on the number of thresholds is $K - k + 1$, where K is the cardinality of the largest losing coalition and k the cardinality of the smallest winning coalition. Indeed, it can be made $(K - k + 1)$ -rough by choosing weights $w(i) = 1$ for all $i \in [n]$ and setting thresholds $k, k + 1, \dots, K$. However this bound is not tight as is seen from the following example.

Example 5 Let $G = ([7], W)$ be a simple game with minimal winning coalitions $\{1, 2\}, \{6, 7\}, \{3, 4, 5\}$ and all coalitions of four players except $\{2, 3, 4, 6\}$. This game is not roughly weighted, because we have the following potent certificate of nonweightedness

$$\mathcal{T} = \left\{ \{1, 2\}^7, \{3, 4, 5\}^9, P; \{2, 3, 5\}^3, \{2, 3, 4\}^3, \right. \\ \left. \{2, 3, 6\}, \{2, 3, 7\}, \{1, 3, 4\}, \{1, 3, 5\}, \{1, 4, 5\}^6, \emptyset \right\}.$$

Let us assign weight 0 to the third player and $\frac{1}{2}$ to everyone else. Then the following four statements hold:

$$- \quad w(\{1, 2\}) = w(\{6, 7\}) = w(\{3, 4, 5\}) = 1 \text{ and } w(\{2, 3, 4, 6\}) = \frac{3}{2}.$$

- If X is winning coalition with four or more players, then $w(X) \geq \frac{3}{2}$.
- If X is losing coalition with three players, then $w(X) \in \{1, \frac{3}{2}\}$.
- If X is losing coalition with fewer than three players, then $w(X) \leq 1$.

Thus G is a 2-rough game with thresholds 1 and $\frac{3}{2}$. Note that the third player has weight zero but is not a dummy.

5 \mathcal{C} -hierarchy

Let us consider another extension of the idea of rough weightedness. This time we will use a threshold interval instead of a single threshold or (as in \mathcal{B} -hierarchy) a collection of threshold points. It is convenient to “normalize” the weights so that the left end of our threshold interval is 1. We do not lose any generality by doing this.

Definition 9 We say that a simple game $G = (P, W)$ is in the class C_α , $\alpha \in \mathbb{R}^{\geq 1}$, if there exists a weight function $w: P \rightarrow \mathbb{R}^{\geq 0}$ such that for $X \in 2^P$ the condition $w(X) > \alpha$ implies that X is winning, and $w(X) < 1$ implies X is losing. Games from C_α will sometimes be called rough $_\alpha$.

The roughly weighted games with nonzero quota form the class C_1 . From Example 3 we can conclude that the Fano game is in $C_{4/3}$ (by giving each player weight $1/3$). We also note that adding or deleting a passer does not change the class of the game.

Definition 10 We say that a game G is *critical* for C_α if it belongs to C_α but does not belong to any C_β with $\beta < \alpha$.

It is clear that if $\alpha \leq \beta$, then $C_\alpha \subseteq C_\beta$. However, we can show more.

Proposition 3 Let c and d be natural numbers with $1 < d < c$. Then there is a simple game G that is rough $_{c/d}$, but that for each $\alpha < c/d$ is not rough $_\alpha$.

Proof Define a game $G = (P, W)$, where $P = [cd]$. Similarly to the proof of Theorem 4, we have c types of players with d players in each type and the different types do not intersect. Winning coalitions are sets with at least $c + 1$ players and also sets having all d players from the same type. By i_j we will denote the i th player of j th type.

If we assign weight $1/d$ to each player, then the lightest winning coalition (d players from the same type) has weight 1 and the heaviest losing coalition has weight c/d . Thus G belongs to $C_{c/d}$.

Let us show that G is not rough $_\alpha$ for any $\alpha < c/d$. Suppose G is rough $_\alpha$ relative to a weight function w . Let $\max\{1_j, \dots, d_j\}$ be the element of the set $\{1_j, \dots, d_j\}$ that has the biggest weight relative to w .

For any type j we know that $w(\max\{1_j, 2_j, \dots, d_j\}) \geq \frac{1}{d}$. The coalition

$$Y = \{\max\{1_1, \dots, d_1\}, \dots, \max\{1_c, \dots, d_c\}\}$$

is losing by definition. Moreover, it has weight $w(Y) \geq c/d$. So c/d is the smallest number that can be taken as α so that G is rough $_\alpha$. \square

Theorem 5 For each $1 \leq \alpha < \beta$, it holds that $C_\alpha \subsetneq C_\beta$.

Proof We know that $C_\alpha \subseteq C_\beta$. If β is a rational number, then by Proposition 3 there exists a game G that is rough $_\beta$ but is not rough $_\alpha$. If β is an irrational number, then choose a rational number r , such that $\alpha < r < \beta$. By Proposition 3 there exists a game G that is rough $_r$ but is not rough $_\alpha$. So $C_\alpha \subsetneq C_r$. All that remains to notice is that $C_r \subseteq C_\beta$. □

Theorem 6 Let G be a simple game that is not roughly weighted and is critical for C_a . Suppose G also belongs to \mathcal{A}_q for some $0 < q < \frac{1}{2}$. Then

$$a \geq \frac{1 - q}{1 - 2q}.$$

Proof Obviously we can assume that q is rational. Since G is in \mathcal{A}_q , it possesses a certificate of nonweightedness \mathcal{T} of the kind

$$\mathcal{T} = (X_1, \dots, X_t, P^s; Y_1, \dots, Y_t, \emptyset^s).$$

Suppose we have a weight function $w: P \rightarrow \mathbb{R}^{\geq 0}$ instantiating $G \in C_a$. Then since $w(X_i) \geq 1$ and $w(P) \geq a$, we have

$$w(X_1) + \dots + w(X_t) + sw(P) \geq t + sa. \tag{9}$$

On the other hand, $w(Y_i) \leq a$ and

$$w(Y_1) + \dots + w(Y_t) \leq ta. \tag{10}$$

From these two inequalities we get $t + sa \leq ta$ or $a \geq \frac{t}{t-s}$. Since $q = \frac{s}{t+s}$ we obtain $a \geq \frac{1-q}{1-2q}$, which proves the theorem. □

6 Degrees of roughness of games with a small number of players

First, we will derive bounds on the largest number $s(n)$ of the spectra $\text{Spec}(n)$.

Theorem 7 For $n \geq 4$, $\frac{1}{2} \lfloor \frac{n}{2} \rfloor \leq s(n) \leq \frac{n-2}{2}$.

Proof Let G be a game with n players. Without loss of generality we can assume that G does not contain passers. Moreover the maximal value of $s(n)$ is achieved on games that are not roughly weighted. By Proposition 1 the biggest losing coalition contains at most $n - 2$ players and the smallest winning coalition has at least two players. If we assign weight $\frac{1}{2}$ to every player, then G is in $C_{(n-2)/2}$.

We can use a game similar to the one from Theorem 4 to prove the lower bound. Suppose our game has n players. If n is odd, then one player will be a dummy. The remaining $2 \lfloor \frac{n}{2} \rfloor$ players will be divided into $\lfloor \frac{n}{2} \rfloor$ pairs: $\{1, 2\}, \{2, 3\}, \dots, \{m - 1, m\}$,

where $m = \lfloor \frac{n}{2} \rfloor$. These pairs are declared minimal winning coalitions. Given any weight function w we have $w(\max\{2i - 1, 2i\}) \geq \frac{1}{2}$ for each i . Then

$$w(\{\max\{1, 2\}, \dots, \max\{m - 1, m\}\}) \geq \frac{m}{2},$$

while this coalition is losing. So $s(n) \geq m/2$ which proves the lower bound. □

Now let us calculate the spectra for $n \leq 6$. By Theorem 2 all games with four players are roughly weighted. Since we may assume that the game does not have passers we may assume that the quota is nonzero. Hence we have $\text{Spec}(4) = \{1\}$. So the first nontrivial case is $n = 5$.

Let $G = ([n], W)$ be a simple game. The problem of finding the smallest α such that $G \in C_\alpha$ holds is a linear programming question. Indeed, let W^{\min} and L^{\max} be the set of minimal winning coalitions and the set of maximal losing coalitions, respectively. We need to find the minimum α such that the following system of linear inequalities is consistent:

$$\begin{cases} w(X) \geq 1 & \text{for } X \in W^{\min}, \\ w(Y) \leq \alpha & \text{for } Y \in L^{\max}. \end{cases}$$

This is equivalent to the following optimization problem:

Minimize: α .
 Subject to: $\sum_{i \in X} w_i \geq 1, \sum_{i \in Y} w_i - \alpha \leq 0$, and $w_i \geq 0; X \in W^{\min}, Y \in L^{\max}$.

Theorem 8 $\text{Spec}(5) = \{1, \frac{6}{5}, \frac{7}{6}, \frac{8}{7}, \frac{9}{8}\}$.

Proof Let G be a critical game with five players. If G has a passer, then as was noted, the passer can be deleted without changing the class of G , hence $G \in C_1$. If G has no passers and does not belong to C_1 , then it is not roughly weighted. By Theorem 2 each game that is not roughly weighted is not strong and is not proper. Thus we have a winning coalition X such that X^c is also winning and a losing coalition Y such that Y^c is also losing.

By Proposition 1 we may assume that the cardinalities of both X and Y are two. Without loss of generality we assume that $X = \{1, 2\}$ and $X^c = \{3, 4, 5\}$. Note that Y cannot be contained in X^c as otherwise Y^c contains X and is not losing. So without loss of generality we assume that $Y = \{1, 5\}, Y^c = \{2, 3, 4\}$.

We have two levels of as yet unclassified coalitions, which can be set either losing or winning:

- level 1 : $\{1, 3, 4\}, \{1, 3, 5\}, \{1, 4, 5\}, \{2, 3, 5\}, \{2, 4, 5\}$,
- level 2 : $\{1, 3\}, \{1, 4\}, \{2, 5\}, \{3, 5\}, \{4, 5\}$.

We wrote Maple code using the ‘‘LPSolve’’ command. First we choose losing coalitions on level 1 and delete all subsets of them from level 2. We add every unclassified coalition from level 1 to winning coalitions. After that we choose losing coalitions

Table 1 Examples of critical simple games for every number of the 5th spectrum

α	Minimal winning coalitions and maximal losing coalitions	Weight representation
$\frac{9}{8}$	$W^{min} = \{\{1, 2\}, \{1, 3, 5\}, \{1, 4, 5\}, \{3, 4, 5\}\}$ $L^{max} = \{\{1, 5\}, \{1, 3, 4\}, \{2, 3, 4\}, \{2, 3, 5\}, \{2, 4, 5\}\}$	$w_1 = \frac{5}{8}, w_2 = \frac{3}{8}, w_5 = \frac{4}{8}$ $w_3 = w_4 = \frac{2}{8}$
$\frac{8}{7}$	$W^{min} = \{\{1, 2\}, \{2, 5\}, \{1, 3, 4\}, \{3, 4, 5\}\}$ $L^{max} = \{\{1, 3, 5\}, \{1, 4, 5\}, \{2, 3, 4\}\}$	$w_1 = w_5 = \frac{3}{7}, w_2 = \frac{4}{7}$ $w_3 = w_4 = \frac{2}{7}$
$\frac{7}{6}$	$W^{min} = \{\{1, 2\}, \{1, 4, 5\}, \{3, 4, 5\}\}$ $L^{max} = \{\{1, 3, 4\}, \{1, 3, 5\}, \{2, 3, 4\}, \{2, 3, 5\}, \{2, 4, 5\}\}$	$w_1 = w_2 = \frac{3}{6}$ $w_3 = w_4 = w_5 = \frac{2}{6}$
$\frac{6}{5}$	$W^{min} = \{\{1, 2\}, \{1, 3\}, \{1, 4\}, \{2, 5\}, \{3, 5\}, \{4, 5\}\}$ $L^{max} = \{\{1, 5\}, \{2, 3, 4\}\}$	$w_1 = w_5 = \frac{3}{5}$ $w_2 = w_3 = w_4 = \frac{2}{5}$

on level 2. We run through all possible combinations of losing coalitions on both levels and solve the respective linear programming problems. The results of these calculations are displayed in Table 1. □

Theorem 9 *The 6th spectrum $Spec(6)$ contains $Spec(5)$ and also the following fractions:*

$$\frac{3}{2}, \frac{4}{3}, \frac{5}{4}, \frac{9}{7}, \frac{10}{9}, \frac{11}{9}, \frac{11}{10}, \frac{12}{11}, \frac{13}{10}, \frac{13}{11}, \frac{13}{12}, \frac{14}{11}, \frac{14}{13}, \frac{15}{13}, \frac{15}{14}, \frac{16}{13}, \frac{16}{15}, \frac{17}{13}, \frac{17}{14}, \frac{17}{15}, \frac{17}{16}, \frac{18}{17}$$

Proof Let G be a critical game with six players. If G has a passer, then $G \in C_\alpha$ where $\alpha \in Spec(5)$. In the other words $Spec(5) \subseteq Spec(6)$. If G doesn't have a passer, then assume it is not roughly weighted. By Theorem 2 we know that every game with six players that is not roughly weighted is either not strong ($Y, Y^c \in L$ for some $Y \in 2^P$) or is not proper ($X, X^c \in W$ for some $X \in 2^P$). By Proposition 1 we can restrict ourselves to the consideration of games for which every coalition with less than two players is losing and every coalition with more that four players is winning. Since G is not roughly weighted there is a potent certificate of nonweightedness $\mathcal{T} = (X_1, \dots, X_k, P; Y_1, \dots, Y_k, \emptyset)$, where the coalitions X_1, \dots, X_k are winning and the coalitions Y_1, \dots, Y_k are losing. The latter absorb all players in X_1, \dots, X_k and the grand coalition. Then there exists a losing coalition Y_j among Y_1, \dots, Y_k with more players than in the smallest winning coalition X_i among X_1, \dots, X_k . If X_i consists of two players, then Y_j has at least three players. If X_i has three players, then Y_j has four players and any subset of it with three players is also losing. Clearly X_i cannot have four players or more. So in any case we have a losing coalition with three players and a winning coalition with three players. Without loss of generality we need to check only six possible cases:

- If G is not proper:
 1. $\{1, 2\}, \{3, 4, 5, 6\} \in W$ and $\{1, 3, 4\} \in L$;

2. $\{1, 2\}, \{3, 4, 5, 6\} \in W$ and $\{3, 4, 5\} \in L$;
 3. $\{1, 2, 3\}, \{4, 5, 6\} \in W$ and $\{1, 4, 5\} \in L$.
- If G is not strong:
4. $\{1, 2\}, \{3, 4, 5, 6\} \in L$ and $\{1, 3, 4\} \in W$;
 5. $\{1, 2\}, \{3, 4, 5, 6\} \in L$ and $\{1, 2, 3\} \in W$;
 6. $\{1, 2, 3\}, \{4, 5, 6\} \in L$ and $\{1, 2, 4\} \in W$.

For each case the algorithm considers all possible assignments of the attributes “winning” and “losing” to coalitions that are not yet classified. Let level 1 consists of all 4-element coalitions, level 2 consists of 3-element coalitions, and level 3 consists of 2-element coalitions. As in the code discussed in the proof of Proposition 8, first the algorithm selects losing coalitions at level 1 (everything else at level 1 will be winning) and classifies all subsets of these coalitions from levels 2 and 3 as losing. Next it selects losing coalitions among coalitions of level 2, which are not yet classified, and classifies all subsets of them from level 3 as losing. Finally, it selects losing coalitions among remaining coalitions of level 3 and solves the linear programming problem using “LPSolve” in Maple, which tries to assign weights to players consistent with the classification of coalitions. We repeat everything for each possible combination of losing coalitions at all levels. The code and the list of critical games are available from the authors. \square

7 Conclusion and further research

Economics has extensively studied weighted majority games. This class was previously extended to the class of roughly weighted games (Taylor and Zwicker 1999; Gvozdeva and Slinko 2011). However, many games are not even roughly weighted, and some of these games are important both for theory and applications.

In this paper we introduced three hierarchies, each of which partitions the class of games without rough weights according to some parameter that can be viewed as capturing some resource—either a measure of our flexibility on the size and structure of the tie-breaking region or allowing certain types of certificates of nonweightedness.

It is important to look for further connections between the classes of the three hierarchies, and we commend that direction to the interested reader.

Also, in this paper we studied only the \mathcal{C} -spectrum. Some interesting questions about this spectrum still remain; for example, the bounds for $s(n)$ are of considerable interest. It would be interesting to study both the \mathcal{A} -spectrum and \mathcal{B} -spectrum as well.

As we mentioned in the introduction, another important generalization of a class of weighted majority games is the class of complete games. All questions that we investigated in this article can be reformulated for games in this class: strictness of the three hierarchies, description of spectra, etc. This is an important direction for future research.

Acknowledgments During a talk on roughly weighted games at the March 2010 Dagstuhl Seminar on Computational Foundations of Social Choice, Bill Zwicker asked whether those games were merely the start of some hierarchy. That question motivated this paper, and we are deeply grateful to Bill both for that question and for encouragement and helpful conversations. We are also very grateful to Edith Elkind for helpful conversations and to the anonymous referees for their supportive reports and many helpful suggestions. Tatiana

Gvozdeva was supported by the University of Auckland’s Faculty of Science Research Development Fund 3624495/9844. Lane A. Hemaspaandra was supported by grants NSF-CCF-0426761, NSF-CCF-0915792, NSF-CCF-1101479, and ARC-DP110101792, and a Friedrich Wilhelm Bessel Research Award from the Alexander von Humboldt Foundation. Arkadii Slinko was supported by the University of Auckland’s Faculty of Science Research Development Fund 3624495/9844 and by grant ARC-DP110101792.

References

- Beimel A, Tassa T, Weinreb E (2005) Characterizing ideal weighted threshold secret sharing, theory of cryptography. In: Lecture notes in computer science vol 3378 Springer, Berlin, pp 600–619
- Blakley GR (1979) Safeguarding cryptographic keys. In: Proceedings of the national computer conference vol 48 AFIPS, Monval, pp 313–317
- Elgot CC (1961) Truth functions realizable by single threshold organs. In: Proceedings of the second annual symposium on switching circuit theory and logical design. Detroit, MI, pp 225–245
- Freixas J, Puente MA (2008) Dimension of complete simple games with minimum. *Eur J Oper Res* 188: 555–568
- Freixas J, Molinero X (2009) Simple games and weighted games: a theoretical and computational viewpoint. *Discr Appl Math* 157:1496–1508
- Gvozdeva T, Slinko A (2011) Weighted and roughly weighted simple games. *Math Soc Sci* 61:20–30
- Isbell JR (1956) A class of majority games. *Q J Math* 7:183–187
- Muroga S (1971) Threshold logic and its applications. Wiley, NY
- Richardson M (1956) On finite projective games. *Proc Am Math Soc* 7:458–465
- Shapley LS (1962) Simple games: an outline of the descriptive theory. *Behav Sci* 7:59–66
- Taylor AD, Zwicker WS (1992) A characterization of weighted voting. *Proc Am Math Soc* 115:1089–1094
- Taylor AD, Zwicker WS (1999) Simple games. Princeton University Press, Princeton
- von Neumann J, Morgenstern O (1944) Theory of games and economic behavior. Princeton University Press, Princeton