

## Erratum to: Analysis of daily teriparatide treatment for osteoporosis in men


R. Niimi<sup>1</sup> · T. Kono<sup>1</sup> · A. Nishihara<sup>1</sup> · M. Hasegawa<sup>2</sup> ·  
A. Matsumine<sup>2</sup> · T. Kono<sup>1</sup> · A. Sudo<sup>2</sup>

Published online: 20 June 2015

© International Osteoporosis Foundation and National Osteoporosis Foundation 2015

### Erratum to: Osteoporos Int DOI 10.1007/s00198-014-3001-1

Figure 3a in the original publication used a wrongly scaled y-axis. The corrected figure is given here:



**Fig. 3** The mean changes in procollagen type I N-terminal propeptide (PINP) (a) and urinary N-telopeptide (uNTX) (b) during teriparatide treatment (\* $p < 0.01$ , \*\* $p < 0.05$ , n.s.; not significant versus baseline; Wilcoxon signed-rank test, n.s. (upper); not significant between men and women; Mann–Whitney  $U$  test). Data are means+SDs

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00198-014-3001-1>.

✉ R. Niimi  
furikakefuri@hotmail.co.jp

<sup>1</sup> Department of Orthopaedic Surgery, Tomidahama Hospital, 26-14, Tomidahamacho, Yokkaichi, Mie 510-8008, Japan

<sup>2</sup> Department of Orthopaedic Surgery, Mie University Graduate School of Medicine, Tsu, Japan