

Systematic scoping review of patients' perceived needs of health services for osteoporosis

L. Chou¹ · P. Shamdasani¹ · A. M. Briggs^{2,3} · F. M. Cicuttini¹ · K. Sullivan⁴ ·
K. L. M. D. Seneviwickrama¹ · A. E. Wluka¹

Received: 10 February 2017 / Accepted: 17 July 2017 / Published online: 31 July 2017
© International Osteoporosis Foundation and National Osteoporosis Foundation 2017

Abstract Health service planners, administrators and providers need to understand the patients' perspective of health services related to osteoporosis to optimise health outcomes. The aims of this study were to systematically identify and review the literature regarding patients' perceived health service needs relating to osteoporosis and osteopenia. A systematic scoping review was performed of publications in MEDLINE, EMBASE, CINAHL and PsycINFO (1990–2016). Descriptive data regarding study design and methodology were extracted and risk of bias assessed. Aggregates of patients' perceived needs of osteoporosis health services were categorised. Thirty-three studies (19 quantitative and 14 qualitative) from 1027 were relevant. The following areas of perceived need emerged: (1) patients sought healthcare from doctors to obtain information and initiate management. They were dissatisfied with poor communication, lack of time and poor continuity of care. (2) Patients perceived a role for osteoporosis

pharmacotherapy but were concerned about medication administration and adverse effects. (3) Patients believed that exercise and vitamin supplementation were important, but there is a lack of data examining the needs for other non-pharmacological measures such as smoking cessation and alcohol. (4) Patients wanted diagnostic evaluation and ongoing surveillance of their bone health. This review identified patients' needs for better communication with their healthcare providers. It also showed that a number of important cornerstones of therapy for osteoporosis, such as pharmacotherapy and exercise, are identified as important by patients, as well as ongoing surveillance of bone health. Understanding patients' perceived needs and aligning them with responsive and evidence-informed service models are likely to optimise patient outcomes.

Keywords Needs assessment of health services · Osteoporosis · Patients

Louisa Chou and Pallavi Shamdasani contributed equally to the work.

Electronic supplementary material The online version of this article (doi:10.1007/s00198-017-4167-0) contains supplementary material, which is available to authorized users.

✉ A. E. Wluka
anita.wluka@monash.edu

¹ Department of Epidemiology and Preventative Medicine, School of Public Health and Preventative Medicine, Monash University, Commercial Road, Melbourne, Victoria, Australia

² School of Physiotherapy and Exercise Science, Curtin University, Bentley, Western Australia, Australia

³ Move: Muscle, Bone and Joint Health, Melbourne, Victoria, Australia

⁴ Monash University Library, Monash University, Melbourne, Victoria, Australia

Introduction

Osteoporosis is increasingly being recognised as an important public health concern due to an ageing population and rise in chronic diseases [1]. It is estimated that one in two women and one in five men over the age of 50 will sustain a fracture due to osteoporosis [2]. Fragility fractures related to osteoporosis are associated with significant morbidity and mortality. The direct medical costs of this global health burden are substantial, amounting to an estimated \$17 billion in the USA in 2005 [3], € 37 billion in the European Union in 2010 [4] and more than \$9 billion in China in 2010 [5]. This is projected to surpass \$25 billion by 2025 [3, 5, 6].

To close the evidence-practice gap in osteoporosis management and address the burden of osteoporosis [6, 7], several peak organisations have developed clinical practice guidelines

to guide clinicians in optimising bone health and managing osteoporosis [8–12]. Recent strategies have been implemented to improve the uptake of evidence-based recommendations, such as education programs, fracture-liaison services, orthogeriatric models of care and audits of healthcare services [13–15]. However, despite these measures, the management of osteoporosis and bone health following fragility fractures remains inadequate [16–18]. Previous studies have shown that just up to 25% of patients identified as high risk had further investigations for osteoporosis and less than 20% of patients with osteoporosis or a history of fragility fractures received treatment to prevent future fractures [15–17, 19, 20].

Optimal osteoporosis outcomes, for the patient and health service, depend on a variety of factors at multiple levels—from health policy through to patients’ self-management behaviours: all of these factors may affect the effective implementation of guidelines and models of care [21]. Understanding why management deviates from guidelines so frequently is important to improve bone health outcomes. A recent seminal report by the International Osteoporosis Foundation [6] has summarised current international gaps in quality service delivery for people with poor bone health and has suggested strategies from a health services and policy perspective for improvement. However, these issues are not considered through the lens of the consumer. As management requires the patient to access and use healthcare services, identifying their perceived needs may provide insight into why optimal management does not occur, or is not sustained (of particular relevance to osteoporosis management). It may also suggest more effective strategies for healthcare providers and policy makers for implementing consumer-centred strategies and promoting patient-centred care: taking the patients’ perceived needs into account may inform clinical decision making, helping doctors to optimise osteoporosis treatment. Although there are published systematic reviews that examine patients’ health beliefs relating to osteoporosis [22] or their experience of living with osteoporosis [23], these do not examine the patients’ perceived needs of health services. There have also been several studies that explore the patients’ perspective and perceived needs of health services for osteoporosis, either directly or indirectly, but no review has been performed to identify and summarise the existing literature. Therefore, we performed a systematic scoping review to identify the literature regarding patients’ perceived needs for health services for osteoporosis and osteopenia management.

Methods

A systematic scoping review was performed to identify what is known about patients’ perceived health service needs for osteoporosis and osteopenia within a larger project examining the patients’ perceived needs relating to musculoskeletal

health [24]. Throughout, we refer to ‘osteoporosis’, which is inclusive of osteopenia. Given the breadth of the topic, a systematic scoping review, based on the framework proposed by Arksey and O’Malley [25], was conducted to comprehensively explore of the patients’ perspective, map the existing literature and to identify gaps in the evidence [26, 27].

Search strategy and study selection

An electronic search of MEDLINE, EMBASE, CINAHL and PsycINFO was performed to identify studies examining patients’ perceived needs relating to osteoporosis health services between January 1990 and July 2016. This time period was chosen to include relevant studies examining the current patient perspective. The search strategy was developed iteratively by an academic librarian, clinical researchers (rheumatologists and physiotherapists) and a healthcare organisation representing consumers with osteoporosis and musculoskeletal disorders. It combined both text words and MeSH terms to capture information regarding the constructs of osteoporosis and bone health, patients’ perceived need(s) and factors related to health services. The term ‘patients’ perceived needs’ was used to broadly capture the patients’ perception of their capacity to benefit from services, including their expectations of satisfaction with and preferences for various services [28]. The term ‘health services’ includes ‘services relating to the diagnosis and treatment of disease, or the promotion, maintenance and restoration of health’, as described by the World Health Organisation [29]. The term ‘health service needs’ describes the patients’ perception of their capacity to benefit from services relating to the diagnosis and treatment of osteoporosis, or the promotion, maintenance and restoration of health, relating to osteoporosis. The detailed search strategy for MEDLINE is provided in the Supplementary Appendix.

Two investigators (LC and PS) independently assessed all the titles and abstracts of the studies identified by the initial search for relevance. The initial screening of manuscripts identified by the search strategy was designed to be as inclusive as possible to identify relevant studies, within the specific inclusion and exclusion criteria to capture the breadth of the literature. The reference lists of retrieved articles and review articles were also manually assessed for further studies for inclusion. To be included in the review, studies had to (1) concern patients older than 18 years and at risk of osteoporosis or having osteoporosis (either diagnosed by a physician, based on bone densitometry results, or individuals taking medications for osteoporosis); (2) report on patients’ perceived needs of health services; (3) concern osteoporosis (either primary or secondary), osteopenia or bone health; and (4) full-text articles. Both qualitative and quantitative studies were included to provide an in-depth review of the topic. Only studies in the English language were retained due to resource constraints. Studies that appeared to meet the inclusion criteria and

relevant reviews were retrieved, and the full text was assessed for relevance by two investigators (LC and PS). Any disagreements in the inclusion of studies were resolved through consensus or reviewed by a third investigator (AW).

Data extraction and analysis

Two investigators (LC and PS) independently extracted the data from relevant studies using a standardised data extraction form developed for this scoping review. The included studies were described and reported according to the following: (1) author and year of publication; (2) study population (patient age and gender, population source, population size and definition of osteoporosis); (3) primary study aim; and (4) description of the study methods. Two authors (LC and PS) independently reviewed and extracted relevant data from the included studies using the principles of meta-ethnography to synthesise qualitative data [30]. This involved a process of identifying key concepts from the included studies, and reciprocal translational analysis was undertaken to translate and compare the concepts from individual studies to other studies and gradually explore and develop overarching themes [31]. Importantly, reciprocal translational analysis allows for the development of a concept or theme by considering different viewpoints related to the same issue, described in different ways. In the first stage, one author (PS) initially developed a framework of concepts and underlying themes, based on primary data in the studies and any pertinent points raised by the authors in the discussion. In the second stage, another author (LC) independently reviewed the studies and further developed the framework of themes and concepts. In the third stage, two

senior authors (FC and AW) with over 10 years of clinical rheumatology consultant-level experience independently reviewed the framework of concepts and themes to ensure clinical meaningfulness and face validity.

Methodological quality assessment


To assess the methodological quality of the included studies, two reviewers independently assessed all of the included studies (LC and PS). For qualitative studies, the Critical Appraisal Skills Programme (CASP) tool was used [32]. The risk of bias tool was utilised to assess the external and internal validity of quantitative studies: low risk of bias of quantitative studies was defined as scoring 8 or more ‘yes’ answers, moderate risk of bias was defined as 6 to 7 ‘yes’ answers and high risk of bias was defined as 5 or fewer ‘yes’ answers [33]. The reviewers discussed and resolved disagreements through consensus. Any disagreements in scoring were reviewed by a third reviewer (AW).

Results

Overview of studies

The search strategy identified 1030 studies, of which 33 articles met the inclusion criteria for this review [34–67]. A PRISMA flowchart detailing the study selection is shown in Fig. 1. The descriptive characteristics of the included studies are shown in Table 1.

Fig. 1 PRISMA flow chart of study selection



Of the included studies, 20 were from North America [34, 35, 37–40, 43, 45–48, 52, 54–57, 59, 60, 64, 65], 6 from Europe [41, 42, 50, 53, 61, 67], 3 from the United Kingdom [36, 49, 51], 1 from South America [66] and 1 from the Middle-east [63]. There was one multi-centre study [44]. A total of 16,975 patients were included; the sample size of the quantitative studies ranged from 21 to 3438, with a median of 765 and the sample size of the qualitative studies ranged from 14 to 164, with a median of 25. Across the studies, 95% of the participants were female: 22 studies examined only female participants [34, 36, 38, 40, 41, 44–50, 52, 53, 58, 60, 61, 63–67] and the remaining 11 studies evaluated mainly women [35, 37, 39, 42, 43, 51, 54–57, 59]. The mean age of participants was 68 years. Eight studies recruited participants with a previous fragility fracture or at high risk of osteoporotic fractures and 6 studies included patients requiring prescription medications, with or without a previous history of fractures. Only 4 studies provided details regarding other co-morbidities: two studies reported that more than 50% of their participants had less than one co-morbidity [51, 61] and two studies had more than 70% of participants with more than two co-morbidities [42, 63].

Nineteen studies used quantitative methods [34, 37, 39, 40, 42, 44, 46, 47, 50–52, 58, 59, 61, 63–67], all of which were cross-sectional surveys; of these, 13 used questionnaires [37, 39, 42, 46, 47, 50, 51, 58, 61, 63, 65, 67], 5 used surveys [34, 40, 44, 59, 64] and 1 used interviews [52]. Fourteen used qualitative methods [35, 36, 38, 41, 43, 45, 48, 49, 53–57, 60]; of these, 10 used interviews [35, 36, 38, 41, 48, 49, 54–57], 4 used focus groups [38, 43, 45, 53] and 1 used video recordings [60]. There were no mixed method studies.

The inclusion criteria for study participants varied across studies. Patients were classified as having osteoporosis based on bone densitometry in seven studies [34, 41, 46, 48, 53, 65], requiring prescription medications in six studies [42, 45, 52, 59, 63, 66] or on the basis of previous fragility fractures or high risk of osteoporotic fractures in eight studies [37–39, 47, 54–56, 61]. The diagnosis of osteoporosis or osteopenia was unspecified in 13 studies [36, 40, 43, 44, 49–51, 57, 58, 60, 64, 67].

Quality of studies

Quality assessments of the included studies are presented in the Supplementary Appendix, Figs. 1 and 2. The quality of qualitative studies was poor, especially for CASP criteria 4 to 6 (Supplementary Appendix, Fig. 1). The quantitative studies were of low quality: 18 studies were at high risk of bias and 1 study was at moderate risk of bias (Supplementary appendix, Fig. 2). These scores for both qualitative and quantitative studies reflected potential biases with participant recruitment and data collection.

Results of review

Four main areas of patients' perceived needs of health services for osteoporosis emerged from this review.

Patients' perceived needs of healthcare providers in the management of their bone health and osteoporosis (Table 2)

Patient preference for consulting medical practitioners and their role

Eight studies identified patients' preference for seeing a medical practitioner for osteoporosis and their perceived role [35, 38, 41, 43, 45, 48, 49, 56]. Four studies found that patients sought care from a medical practitioner for their bone health [43, 45, 48, 49]. Two studies reported that patients believed and trusted medical specialists such as endocrinologists and rheumatologists more than their primary care physician, and they perceived their specialists as being more interested in their bone health than primary care providers [35, 43]. Feldstein found that patients who had sustained a fracture advocated for standardised protocols for integrating and involving medical specialists in the management of osteoporosis [38]. The role of the medical practitioner was perceived to perform a thorough examination [41], provide osteoporosis information and education [38, 41, 49, 56], initiate screening for osteoporosis [38, 56], prescribe and monitor treatment [38, 45, 48, 56] and provide support for optimal self-management [45].

Desirable characteristics of the medical practitioner

Four studies reported on the desired characteristics of medical practitioners in the management of osteoporosis [36, 41, 45, 52]. Besser found that patients wanted to be involved with decisions related to osteoporosis treatment [36]. Lau and Rizzoli reported that the patients wanted follow up from healthcare providers for support and monitoring of medications [45, 52]. Also, patients wanted their osteoporosis to be taken seriously by their practitioners [41] and to be able to discuss medication problems and concerns [45]. Lau reported that patients wanted non-judgemental care [45].

Dissatisfaction with, or concerns about, medical and non-medical practitioners

Six studies identified patients' dissatisfaction and concerns with medical practitioners relating to their osteoporosis management [35, 36, 43, 46, 48, 49]. Patients perceived poor communication, lack of an adequate explanation of the

Table 1 Descriptive characteristics of included studies

Author, year, country	Definition of osteoporosis	No. of participants	Source of participants	Age and gender	Primary study aims	Study design
Besser 2012 UK [36]	Diagnosed with osteoporosis/-osteopenia (criteria unspecified) for >6 months and prescribed osteoporosis medication.	14	Rheumatology clinic and osteoporosis screening unit at a teaching hospital	Mean age 69,100% female	To inform the development of a psychological intervention to increase adherence to treatment. The study aimed to investigate the osteoporosis patients' perceptions of their illness and medications to provide an evidence base for investigating adherence and how to improve it.	Qualitative: semi-structured interviews and drawings
Bogoch 2008 Canada [37]	Women aged ≥ 40 years or men ≥ 50 years who had sustained a fragility fracture of the wrist	166	Fracture clinic of a large teaching hospital	Mean age of men 65 (SD 10.1) and women 64.8 (SD 13.5) 81% female	To provide information for practitioners regarding osteoporosis-related needs of patients who present with low-trauma wrist fractures and are at high risk of subsequent hip fracture	Quantitative: questionnaires
Feldstein 2008 USA [38]	Patients who had had a fragility fracture with no osteoporosis management in prior 12 months	67	Health maintenance organisation. 10 patients.	Age greater >67 years 100% female	To gain perspective on an outreach program and barriers to care for osteoporosis treatment.	Qualitative: interviews and focus groups
Fraenkel 2006 USA [39]	Individuals at high risk for an osteoporotic hip fracture (Fracture Index Score > 7)	76	Patients who recently (within 2 weeks) had a DEXA scan were recruited from 6 centres performing bone densitometry	Mean age 78 (SD 5) 95% female	To determine older adults' treatment preferences for osteoporosis comparing bisphosphonates and hip protectors	Quantitative: questionnaires
Gold 2006 USA [40]	Women diagnosed with osteoporosis by a physician	617 in the preference study	Patients who were a part of the Risendronate Claims Study were sent an email invitation.	Mean age 67.3 (SD 9.4) 100% females	To determine how patients' preferences for weekly vs monthly bisphosphonate therapy is influenced by their knowledge of the medication's proven fracture efficacy.	Quantitative: online survey
Hansen 2014 Denmark [41]	Women with BMD T score < -2.5, no previous fracture with prescription for osteoporosis medication	15	Patients undergoing DXA scans at participating hospitals were recruited	Median age 72 (range 65–79) 100% female	To examine the experiences of women living with osteoporosis during the first 6 months after diagnosis.	Qualitative: interviews
Hiligsman 2014 Netherlands [42]	Patients with or at risk of osteoporosis to whom medication or lifestyle changes were proposed	257	Consecutive patients were recruited during outpatients' clinics in 2 Belgian osteoporosis centres.	Mean age 67.1 (SD 10.4) 83.3% female	To evaluate the preferences of patients with, or at risk of, osteoporosis for medication attributes, and to establish how patients	Quantitative: questionnaires

Table 1 (continued)

Author, year, country	Definition of osteoporosis	No. of participants	Source of participants	Age and gender	Primary study aims	Study design
Iversen 2011 USA [43]	Patients with diagnosed osteoporosis (criteria unspecified) on treatment for osteoporosis.	32	Participants recruited via advertisements in a tertiary hospital medical center newsletter.	Age range 65–85 93% female	trade between these attributes To determine factors influencing adherence to osteoporosis medications.	Qualitative: focus group discussions
Keen 2006 Multi-centre (UK, Germany, Spain, Italy and France) [44]	Physician diagnosis of postmenopausal osteoporosis	1248	Women from 5 European countries were recruited. Recruitment details unspecified.	Mean age 66, 100% female	To determine participant preference for weekly vs monthly bisphosphonate therapy for osteoporosis after being informed about differences in fracture efficacy	Quantitative: survey
Lau 2008 Canada [45]	Post-menopausal women taking prescription or over the counter medications for osteoporosis (definition unspecified)	37	Recruited by family physicians, geriatrician, rheumatologist and community pharmacists	Age distribution not specified. 100% female	To examine patients' perceptions of osteoporosis medications, reasons for non-adherence to therapy and effectiveness of strategies to improve adherence.	Qualitative: focus group discussion
Martin 1997 USA [46]	Clinical osteoporosis (BMD > T score – 2.5 with a history of fragility fracture)	465	Source of participants unspecified (222 participants met definition of clinical osteoporosis and 243 were defined as non-osteoporotic)	78% of osteoporotic women aged 70 or older 100% female	To quantify the effect of osteoporosis on quality of life of all women	Quantitative: questionnaires
Mauck 2002 USA [47]	Low impact fracture (i.e. fall from standing height or less).	21	Consecutive postmenopausal women >50 years who were hospitalised with a low-impact acute proximal femur fracture in May–August 2000, identified from the computerised admission records.	Mean age 81 (SD7) 100% female	To explore the process a woman negotiates when deciding to accept pharmacologic treatment for osteoporosis after a hip fracture.	Quantitative: questionnaires
Mazor 2010 USA [48]	Osteoporosis (BMD < T score – 2.5)	36	A multispeciality group practice in Massachusetts.	Age range > 65 years 100% female	To examine individuals' beliefs and experiences related to osteoporosis and treatment.	Qualitative: phone interviews
McKenna 2008 UK [49]	Diagnosed with osteoporosis (criteria unspecified)	21	Patients recruited through National Osteoporosis Society support groups, osteoporosis exercises classes and South Asian community centres	Age range 43–82 years 100% female	To compare the experiences of osteoporotic Caucasian women and South Asian women during their primary care physician consultations.	Qualitative: interviews
Payer 2009	Women with BMD diagnosed	2035	Participants recruited voluntarily for	Mean age 64 years 100% female	The aim of this VIVA II questionnaire-based study was to analysis	Quantitative: questionnaires

Table 1 (continued)

Author, year, country	Definition of osteoporosis	No. of participants	Source of participants	Age and gender	Primary study aims	Study design
Slovakia [50]	osteoporosis (criteria unspecified).		bisphosphonate treatment		the reasons for preferring once monthly bisphosphonates in patients with post-menopausal osteoporosis as a follow up to the published VIVA study.	
Richards 2007 UK [51]	Osteoporosis status not determined.	2485	Population-based healthy twin volunteers, > 55 years	Mean age 64.5 (SD 6.4) 90.3% female	To discern which therapeutic attributes would be most preferred by a population representative of the age and sex distribution of patients with osteoporosis	Quantitative: questionnaires
Ringe 2006 Germany [67]	No definition of osteoporosis, half of the participants selected as current bisphosphonate users	164	Postmenopausal women aged >55yo recruited from Germany or UK. Source of participants otherwise unspecified.	Mean age 69 (SD 8.8) 100% female	To evaluate whether the intake instructions and packaging of the new combination packaging with the once-weekly bisphosphonate risedronic acid and once-daily calcium tablets were better understood and preferred by postmenopausal women than if these women received separate packs of once-weekly bisphosphonate and calcium tablets.	Quantitative: semi-structured questionnaires
Rizzoli 2010 USA [52]	Post menopausal osteoporosis diagnosed by a physician and were currently or in the past 2 years prescribed medications	844 patients and 837 physicians	Source of participants and recruitment not specified.	Age range unspecified (post- menopausal women) 100% female	To investigate gaps between physician and patient knowledge on osteoporosis, understand barriers to patient adherence and improve communication	Quantitative: telephone interviews
Rothmann 2014 Denmark [53]	Women both with and without osteoporosis (DXA BMD T score < -2.5.)	31	Purposive sampling of participants from the ROSE study in Southern Denmark	Age range 65–80 100% female	To investigate women's perspectives and experiences with screening for osteoporosis.	Qualitative: focus group discussions
Sale 2010 Canada [54]	Patients >65 years old, with or without a history of osteoporosis treatment, who had a fragility fracture in the last 5 years and	21	Purposive sampling of patients identified from a fracture clinic osteoporosis screening program at an urban teaching hospital	Age range 65–88 71% female	To examine patients' experiences with the decision to take osteoporosis medication after they sustained a fracture	Qualitative: interviews

Table 1 (continued)

Author, year, country	Definition of osteoporosis	No. of participants	Source of participants	Age and gender	Primary study aims	Study design
Sale 2014 Canada [55]	deemed high risk for future fracture Patients >50 years, who had a fragility fracture (WHO definition)	25	Purposive sampling of patients presenting to a teaching hospital who experienced a fragility fracture and were candidates for fracture risk assessment.	Age range 50–79 88% female	To examine patients' experiences regarding BMD testing and bone health treatment after being screened through Ontario's Fracture Clinic Screening Program	Qualitative: interviews
Sale 2014 Canada [57]	Patients with osteoporosis-related fractures, but definition of osteoporosis not defined	25	Urban fracture clinic	Age range 50–79 88% female	To examine patients' self-management of bone health and fracture risk, particularly behaviours other than medication use and seeking diagnostic testing.	Qualitative: interviews
Sale 2014 Canada [56]	Patients who had a fragility fracture at >50 years and were not taking osteoporosis pharmacotherapy at the time of the fracture	28	Advertisement in a patient group newsletter	Age range 51–89 93% female	To examine experiences and behaviours with bone health management post-fracture among members of a national osteoporosis patient group	Qualitative: telephone interviews
Sale 2015 Canada [35]	Patients who had a fragility fracture at >50 years and were not taking osteoporosis pharmacotherapy at the time of the fracture	28	Advertisement in a patient group newsletter	Age range 51–89 93% female	To examine messages perceived by members of an osteoporosis patient group from various healthcare providers regarding bone health	Qualitative: telephone interviews
Saltman 2006 Australia [58]	3 participant categories were chosen: (a) patients diagnosed with a preventable condition ie osteoporotic fracture, taking bisphosphonates), (b) patients with other chronic conditions and (c) acute or no conditions	1096	Patients recruited by general practitioners (110 primary care physicians from research network databases held at the University of Sydney each recruited 10 patients)	Mean age of patients with preventable condition 74.7, mean age of patients with chronic illness 71.3, mean age of patients with acute/no illness 69.2 100% female	To explore whether various models that have described patient beliefs and motivations for medication taking applied to patient preferences and decision making across a range of patients with different types of conditions and varying experiences of medication frequencies, and whether there were differences in characteristics between these groups	Quantitative: questionnaires
Schousboe 2011 USA [59]	Patients with a prescription for an oral bisphosphonates	686	Patients recruited after reviewing the electronic medication record of Park Nicollet Clinic and	Mean age 66.3 (SD 10.1) 94% female	To estimate the associations of patients' perceived need of medication for fracture	Quantitative: surveys

Table 1 (continued)

Author, year, country	Definition of osteoporosis	No. of participants	Source of participants	Age and gender	Primary study aims	Study design
			who had a clinic visit within 6 months of the mailing date of the survey		prevention with objective indicators of fracture risk, patients' concerns about medications and the quality of the patient–physician relationship	
Scoville 2011 USA [60]	Postmenopausal women aged ≥ 50 years with osteopenia or osteoporosis and not already taking bisphosphonates or other prescription medications	18	10 academic primary care sites partaking in Osteoporosis Choice (randomised trial of a decision aid)	Patients mean age 70.6 (SD 9.4) 100% female patients	To determine the reasons women present when expressing hesitation about initiation of bisphosphonates during primary care consultations with clinics and how these clinicians react by studying video recordings of these consultations	Qualitative: video recordings of encounters
Turbi 2004 Spain [61]	Postmenopausal women >55 years of age and at risk for osteoporotic fractures (physician diagnosed)	909	Open label, prospective, observational, nonrandomized study conducted at 154 centres across Spain.	Mean age 64.4 (SD 6.9) 100% female	To assess the compliance of postmenopausal women at risk for osteoporotic fractures who were treated with raloxifene vs alendronate during a 12 month observation period in a routine clinical setting.	Quantitative: questionnaires
Weiss 2005 Israel [63]	Postmenopausal women, treated with alendronate daily for at least 1 month within the preceding year.	3438	Medical providers from 14 hospital and 150 primary care community clinics recruited subjects.	Mean age 66.7 (SD 8.9) 100% female	To measure compliance, convenience, tolerance and relative preference of alendronate oral weekly treatment among postmenopausal women with osteoporosis and physician satisfaction compared with previous treatment with alendronate oral daily.	Quantitative: questionnaires
Weiss 2007 USA [64]	Patients with a history of osteoporosis or at risk of osteoporosis (unspecified definition)	999	Women were surveyed via the Internet as part of the National Health and Wellness survey	Mean age 65.1 (SD 8.2) 100% female	To assess patient preferences for 2 osteoporosis medications	Quantitative: surveys
Yood 2008 USA [65]	Osteoporosis defined as BMD T score < -2.5 .	236	A multispecialty practice.	Age 35–33: 1.7%, 45–54: 10.6%, 55–64: 25.4%, 65–74: 28.4%, >75 : 33.9% 100% female	To evaluate the influence of patient characteristics, perceptions, knowledge and beliefs about osteoporosis on the decision to initiate	Quantitative: questionnaires

Table 1 (continued)

Author, year, country	Definition of osteoporosis	No. of participants	Source of participants	Age and gender	Primary study aims	Study design
Yu 2015 USA [34]	Osteoporosis defined as a diagnostic ICD code for osteoporosis and evidence of BMD test	430	Patients identified from Optum Research Database and a cross-sectional mail survey was conducted	Mean age 61 100% female	osteoporotic treatment To examine patients' reasons for not initiating osteoporosis treatment among women with osteoporosis	Quantitative: surveys
Zanchetta 2005 Argentina [66]	Postmenopausal women who had received prescription for raloxifene and had undergone BMD measurement	419	Patients identified from the Metabolic Research Institute database	Mean age 61.4 (SD 7.4) 100% female	To assess the raloxifene compliance and continuance rates and adverse effects over 24 months in clinical practice	Quantitative: telephone interviews

diagnosis and poor continuity of care to be barriers to a good relationship with their doctor [36, 46]. Patients were dissatisfied with the lack of time during consultations and felt that they were unable to ask questions or raise issues with medications with their physicians [35, 36, 43]. Furthermore, they felt that their primary care providers were dismissive of their concerns about osteoporosis [35]. Patients were disappointed with the strong focus on medications and expressed distrust when medical practitioners were too quick to recommend medications, rather than adopt a more holistic approach to care, inclusive of non-pharmacologic options [48, 49]. Moreover, patients reported inconsistent recommendations from different practitioners, and in particular, they found the advice from other disciplines of healthcare, such as nutritionists, physiotherapists and chiropractors to be contradictory, sporadic and not forthcoming [35].

Patients' needs related to pharmacotherapy for osteoporosis and bone health (Table 3)

Perceptions and roles of medications

Eleven studies examined the patients' preference for medications and the perceived role of pharmacotherapy [36, 37, 39, 45, 47, 48, 54, 56, 59, 61, 65]. While some studies found that patients had a preference for pharmacological management of osteoporosis [36, 37, 39, 45, 54, 56, 59], other studies did not [45, 48, 54, 56]. The patients who were more willing to take medication had been told of the diagnosis of osteoporosis [47, 65] and had previous bone mineral density (BMD) testing [47], believed they were susceptible to fractures [59], had a good relationship with their doctor or trusted their physicians [54, 59] and believed in the effectiveness of medications [65].

The role of pharmacotherapy was perceived to help eliminate symptoms, help avoid further deterioration in bone health, provide extra strength for the bone and improve bone density [48, 54]. A single study that compared patients' predilection for pharmacotherapy compared to hip protectors in high-risk patients found that although patients preferred bisphosphonates for the management of their osteoporosis, older patients were more likely to avoid prescription medications and preferred hip protectors [39]. In contrast, several studies reported that patients did not prefer pharmacotherapy for osteoporosis management [45, 48, 54, 56]. Mauck reported that most women who were admitted to a tertiary hospital after a fragility fracture were either unaware of osteoporosis or had never considered pharmacological treatment [47]. Some patients viewed osteoporosis as a consequence of ageing and did not perceive a need for medications [48] and some patients wanted a drug holiday from bisphosphonate treatment [56]. Also, some patients preferred lifestyle modifications rather than pharmacotherapy for osteoporosis management [45, 48, 56].

Concerns about medications

There were 12 studies that reported the patients' concerns with osteoporosis medications [34, 36, 41–43, 45, 48, 53, 54, 60, 65, 66]. Patients who believed they had good health were concerned about taking medications for a condition that was otherwise asymptomatic [53, 60]. Those with a family member who had osteoporosis with no complications were less likely to perceive a benefit with pharmacotherapy [53, 60]. Moreover, patients were unwilling to take medications if they had family members or friends who had experienced adverse events, or if they heard about side effects from the media [34, 45, 48]. Potential side effects from medications were a major

Table 2 Patients' perceived need of healthcare providers for osteoporosis

Author, year	Results
Patient preference for consulting medical practitioners and their role Feldstein 2008 [38]	<ul style="list-style-type: none"> • Patients advocated for standardised protocols for integrating and involving specialists in the management of osteoporosis at the time of fracture.
Hansen 2014 [41]	<ul style="list-style-type: none"> • Most patients thought that specialists should provide basic education in osteoporosis and initiate screening or treatment, with follow-up by a primary care provider or care manager
Iversen, 2011 [43]	<ul style="list-style-type: none"> • Patients expect a thorough examination and informative consultation and clarity of diagnosis
Lau, 2008 [45]	<ul style="list-style-type: none"> • Patients believed and trusted specialists such as endocrinologists and rheumatologists more than their primary care physician regarding osteoporosis management.
Mazor 2010 [48]	<ul style="list-style-type: none"> • Patients found follow up from health care providers for support and monitoring for medications very useful for improving adherence
McKenna, 2008 [49]	<ul style="list-style-type: none"> • Many women relied on their physicians' recommendation in deciding whether to take osteoporosis medications
Sale 2014 [56]	<ul style="list-style-type: none"> • Patients preferred attending consultations, expecting primary care physician dialogue on osteoporosis
Sale, 2015 [35]	<ul style="list-style-type: none"> • Patients believe the role of the doctor in their bone health was to prescribe medication that they requested, to do other routine activities such as annual exams and facilitate access to tests and provide current information
Desirable characteristics of the medical practitioner Besser 2012 [36] Hansen 2014 [41]	<ul style="list-style-type: none"> • Patients were satisfied with the care they received from specialists with many patients reporting that their specialists were more interested in their bone health than their primary care provider. • Specialists were perceived to be more knowledgeable about osteoporosis, and they took more time to discuss their bone.
Lau, 2008 [45]	<ul style="list-style-type: none"> • Patients wanted to be more involved with decisions related to treatment
Rizzoli, 2010 [52]	<ul style="list-style-type: none"> • Patients wanted their osteoporosis to be taken seriously by their physician, which promoted a feeling of care and trust
Dissatisfaction with, or concerns about, medical and non-medical practitioners Besser 2012 [36]	<ul style="list-style-type: none"> • A patient described how she felt taken seriously when her primary care physician referred to a specialist clinic for osteoporosis treatment • Patients found follow up from health care providers for support and monitoring for medications very useful for improving adherence • Patients wanted to be able to discuss their medication problems with their physicians • Patients wanted a non-judgmental service from their doctors • Patients found it helpful to have more frequent contact with their physicians regarding osteoporosis
	<ul style="list-style-type: none"> • Patients perceived lack of time during consultation, poor communication, lack of continuity of care as barriers to a good relationship with their doctor

Table 2 (continued)

Author, year	Results
Iversen, 2011 [43]	<ul style="list-style-type: none"> • Patients felt their health visits were not long enough to be able to discuss all their questions with their doctor. They also felt they were unable to bring up medication issues as their physicians were very rushed
Mazor 2010 [48]	<ul style="list-style-type: none"> • Some women expressed distrust and felt that doctors were too quick to recommend prescription medication
Martin, 1997 [46]	<ul style="list-style-type: none"> • Patients felt they had lack of definitive answers from their physician regarding osteoporosis
McKenna, 2008 [49]	<ul style="list-style-type: none"> • Patients were disappointed that consultations had a strong focus on medication and wanted to discuss other treatment options
Sale, 2015 [35]	<ul style="list-style-type: none"> • Patients perceived that their primary care providers were not interested in their bone health, and were dismissive of their concerns about osteoporosis. They also reported that the recommendations from different healthcare providers appeared to be inconsistent.
Preference for other healthcare providers	
Sale, 2015 [35]	<ul style="list-style-type: none"> • The messages received from other healthcare providers such as nutritionists, physiotherapists and chiropractors were perceived as sporadic, inconsistent and not forthcoming.

concern for many patients [34, 36, 41–43, 45, 48, 53, 54, 60, 65, 66], as well as possible drug interactions from polypharmacy [36, 66], the potential for addiction and overdosing [36]. In particular, some patients had specific concerns including the potential for jaw osteonecrosis, gastrointestinal side effects, breast and oesophageal cancer, thrombotic effects and cardiovascular events [34, 42, 45, 53, 66]. Patients also reported a dislike of chemicals [36, 45], distrust of medications [65] and of pharmaceutical companies [36]. Dissatisfaction with their doctor or the physician's attitude were other reasons for patients to not want to pursue pharmacotherapy for the management of osteoporosis [54, 66]. Furthermore, Iversen reported that patients found the method of medication administration and instructions difficult to understand and remember [43].

Preferable therapeutic attributes of medications

Patients' preferred therapeutic attributes of osteoporosis pharmacotherapy were also examined through this review [40, 42, 44, 45, 50–52, 58, 63, 64, 67]. Patients wanted osteoporosis medications to be effective [40, 44, 64], to not interact with other medications [52], have fewer side effects [52] and be easier to administer [44, 52, 64]. A single study evaluating combination packaging of bisphosphonates and calcium supplementation found that patients preferred the ease and convenience of combination packaging [67]. Some studies found that patients preferred weekly to daily or monthly dosing [40, 44, 58, 64]; however, other studies reported a preference for monthly administration [42, 66].

Patients' perceived needs of non-pharmacological management of osteoporosis (Table 4)

Four studies examined the patients' perceived needs of non-pharmacological management of osteoporosis [37, 38, 45, 57]. Patients' preference for calcium and vitamin D supplementation were examined by four articles [37, 38, 45, 57], which found that patients wanted these supplements for osteoporosis management. Patients expressed more willingness and comfort with taking supplements than prescription medication [38] and believed them to be more natural and safe [45]. Bogoch and Sale found that patients see a role for exercise for osteoporosis management [37, 57]. There were no studies identified that examined the patients' perceived needs of other non-pharmacological strategies such as smoking cessation, attitudes to interventions related to falls prevention and avoidance of excessive alcohol.

Table 3 Patients' perceived needs of pharmacotherapy for bone health and osteoporosis

Author, year	Results
Preference for medications and role of medications	
Besser 2012 [36]	<ul style="list-style-type: none"> • Half of the patients said medication use in general was positive
Bogoch 2008 [37]	<ul style="list-style-type: none"> • More than 85% of patients stated they would take medication for osteoporosis if their physician recommended such treatment
Fraenkel 2006 [39]	<ul style="list-style-type: none"> • Patients preferred bisphosphonates over hip protectors however older adults preferred to avoid taking prescription drugs for most health problems were more likely to prefer hip protectors
Lau 2008 [45]	<ul style="list-style-type: none"> • Improvement in BMD, not having a fracture and having a quicker recovery after a fall positively reinforced persistence in taking osteoporosis medications • Some patients believed that lifestyle modifications would be enough to prevent osteoporosis and that medications should be used as a last resort
Mauck 2002 [47]	<ul style="list-style-type: none"> • Most women (62%) who were admitted to a tertiary hospital were either unaware of osteoporosis or had never considered pharmacologic treatment • Previous BMD evaluation and a diagnosis of osteoporosis were associated with patients considering or currently taking medication
Mazor 2010 [48]	<ul style="list-style-type: none"> • Some women viewed osteoporosis as a common consequence of ageing and believed that medication was therefore not needed or not likely to be of benefit • Some women expressed confidence in the effectiveness of prescription osteoporosis medications and thought that it may be helpful to eliminate their symptoms, help them avoid further decline in health, replace something that they cannot obtain through diet alone • Some patients seemed confused about medications, thinking that it would reduce pain • Some women reported that they did not like medications and would avoid whenever possible • For some women it is seen as the last resort, only when calcium supplements and exercise had failed • A diagnosis of osteoporosis seemed to lead directly to the perception that medications were needed • Knowledge of others' experiences also affected views on medication • Patients have a preference for using lifestyle changes rather than prescription medications for osteoporosis treatment
Sale 2010 [54]	<ul style="list-style-type: none"> • One participant considered his bisphosphonate to be a minor medication, just more like supplements • Participants were more likely to take bisphosphonates if they had a good relationship with their health care provider or trusted their doctor • Some patients perceive the benefits of bisphosphonates to include keeping the bone from weakening, providing extra strength for the bone, preventing further bone loss or improving the bone density • Some patients did not understand what the medications were for • One patient described being convinced to take medications because her physician gave a detailed explanation of the condition and medication and she felt confident to take the medications
Sale 2014 [56]	<ul style="list-style-type: none"> • Some participants describe requesting prescriptions for anti-resorptive medications • Patients who were not taking medications gave a variety of reasons including refusal to initiate the first prescription, refusal to continue the prescription, deciding to take a drug holiday, wanting to try non-pharmacological strategies
Schousboe 2011 [59]	<ul style="list-style-type: none"> • Patients' belief in their susceptibility to and severity of fractures and trust that the prescribing physician is competent and willing to consider their interests are associated with the patients' perceived need for medications

Table 3 (continued)

Author, year	Results
Turbi 2004 [61]	<ul style="list-style-type: none"> • More patients reported being satisfied with raloxifene compared with alendronate
Yood 2008 [65]	<ul style="list-style-type: none"> • Patients who have been told they had osteoporosis were more likely to start prescription medications than those who had not been told they had osteoporosis • Belief in medication effectiveness was associated with initiation of medications
Concerns about medications	
Besser 2012 [36]	<ul style="list-style-type: none"> • Half of the patients listed various concerns about medications including side effects, harmfulness, over prescribing, addiction, suspicion of pharmaceutical companies, dislike of chemicals, drug interactions and overdosing
Hansen 2014 [41]	<ul style="list-style-type: none"> • Many participants expressed concerns about media reports of the link between bisphosphonates and oesophageal cancer
Hilgsmann 2014 [42]	<ul style="list-style-type: none"> • The comprehensive package leaflet in the medication package and possible side effects caused worries and anxiety about taking medications
Iversen, 2011 [43]	<ul style="list-style-type: none"> • Patients disliked being at risk of gastrointestinal disorders more than being at risk of skin reactions or flu-like symptoms
Lau 2008 [45]	<ul style="list-style-type: none"> • Patients expressed uncertainty about how to take their medications • Side effects of medications were a primary reason for lack of adherence • Method of medication administration and instructions were difficult for patients to remember • Patients were unwilling to take a medication if they heard that a family member or friend had a negative experience or if they heard negative publicity about the medication in the media • Some patients did not like to idea of taking any medications because they viewed medications as artificial and thought they had unpredictable effects. • Fear of breast cancer or cardiovascular events from hormone replacement therapy dominant patients' risk benefit assessments more than fear of other adverse effects, however, patients were willing to take hormone replacement therapy if they perceived their personal risk of these serious adverse events to be low • Some women expressed serious concerns about medications generally and fear of side effects in particular • Some declined medications due to their concerns about side effects • Some women discontinued medications after hearing reports of side effects through the media and other sources, even if they had not personally experienced side effects • The administration of some medications (e.g. sitting upright) was interpreted by some patients as evidence that the medication was dangerous
Mazor 2010 [48]	
Rothmann 2014 [53]	<ul style="list-style-type: none"> • A patient did not want treatment as she had a first hand experience of a serious side effect (osteonecrosis of the jaw) in a close relative • Patients were concerned about side-effects of taking a medication for a condition that otherwise was asymptomatic • Some participants described concerns with bisphosphonates regarding adverse effects (wanting to see their dentist before starting treatment, history of multiple allergies and concern with further medications) • Some patients were "turned off" medications by their physicians attitude
Sale 2010 [54]	
Scoville 2011 [60]	<ul style="list-style-type: none"> • Some reasons patients do not accept treatment include concern about side effects, history of adverse effects, distrust of medications, history of family member with no osteoporosis complications, good health without other treatments, perceived low value of potential benefits (too old to benefit, limited knowledge of osteoporosis, medications will not produce benefit)

Table 3 (continued)

Author, year	Results
Yood 2008 [65]	<ul style="list-style-type: none"> • Patients that had more distrust of medications and concern about side effects were more likely to not initiate medications
Yu 2015 [34]	<ul style="list-style-type: none"> • The primary reasons for not initiating osteoporosis medication were concern over side effects (77.3%), medication costs (34.1%) and pre-existing gastrointestinal concerns (25%).
Zanchetta 2004 [66]	<ul style="list-style-type: none"> • Patients' reasons for not starting treatment for raloxifene included fear of thrombolytic events, lack of interest in starting treatment, other physicians' advice, family problems, dissatisfaction with the prescribing physician, treatment cost, health problems unrelated to osteoporosis, mistrust in the prescription, advice from family and friends, fear of breast cancer, belief that raloxifene is hormonal and polypharmacy.
Preferable therapeutic attributes of medications	
Gold 2006 [40]	<ul style="list-style-type: none"> • Once patients were informed of the differences in fracture efficacy between the 2 therapies, more patients preferred weekly therapy over monthly therapy
Hilgsmann 2014 [42]	<ul style="list-style-type: none"> • Patients preferred either an oral monthly tablet or 6 monthly subcutaneous injection above weekly oral tablets, 3-month subcutaneously or yearly intravenous injections
Keen 2006 [44]	<ul style="list-style-type: none"> • Patients preferred weekly bisphosphonate therapy to monthly
Lau 2008 [45]	<ul style="list-style-type: none"> • In the UK, patients aged 55–59 and those over 70 preferred weekly compared to women in their 60s that preferred monthly
Payer 2009 [50]	<ul style="list-style-type: none"> • Patients preferred weekly therapy due to perceived efficacy, dosing and convenience • Patients who found rearranging their daily routines difficult preferred the once-daily dosing option of bisphosphonates • Those patients who had successfully integrated taking medication into their daily routines found it easier to take medication every day rather than once weekly
Richards 2007 [51]	<ul style="list-style-type: none"> • Patients prefer once monthly dosing due to the convenience and simplicity of treatment and the need to take fewer pills. Other reasons that a minority of patients reported included fewer reminders of the disease and independence.
Richards 2007 [51]	<ul style="list-style-type: none"> • 45% of participants preferred daily medications, 20% preferred weekly and 30% preferred monthly medications
Ringe 2006 [67]	<ul style="list-style-type: none"> • The least popular dosing frequency was twice per day • Participants that were not already taking anti-osteoporotic medications preferred daily therapy without having to remain fasting and upright after taking the medication compared with a weekly regime and monthly therapy
Ringe 2006 [67]	<ul style="list-style-type: none"> • Subjects already taking non-weekly anti-osteoporotic medications preferred continuing with this routine
Ringe 2006 [67]	<ul style="list-style-type: none"> • Subjects using weekly anti-osteoporotic therapy preferred weekly preparations
Rizzoli 2010 [52]	<ul style="list-style-type: none"> • Patients preferred combination packaging of bisphosphonates and calcium supplementation due to convenience, ease of understanding dosing instructions
Rizzoli 2010 [52]	<ul style="list-style-type: none"> • Patients desired osteoporosis treatments to not interact with other medications, have fewer side effects, require less frequent dosing, be easier to take and affect their regular routine less and have a less complicated dosing.
Saltman 2006 [58]	<ul style="list-style-type: none"> • Patients have a preference for weekly medication compared to monthly dosing

Table 3 (continued)

Author, year	Results
Weiss 2005 [63]	<ul style="list-style-type: none"> • 96% of women preferred alendronate weekly to the daily regime • Patients found weekly preparations more convenient • 77.6% of those who had previously stopped therapy with alendronate daily due to intolerance were willing to continue with weekly alendronate • Effectiveness was ranked as the most important determinant of preference. <p>Other less important reasons for a patient to prefer one drug over another included time on market, dosing procedure and dosing frequency.</p>
Weiss 2007 [64]	

Patients' perceived needs of investigations for osteoporosis (Table 5)

Three studies described patients' perceived need for investigations for the diagnosis of osteoporosis [48, 53, 56]. Patients saw a role for bone densitometry testing for diagnostic evaluation [48, 56]. Rothmann found that patients interpreted screening for osteoporosis as an opportunity to get reassurance about bone health and to optimise their own general health [53]. Three studies described patients' perceived need for investigations for ongoing surveillance of bone health [36, 48, 56]. Patients wanted feedback from bone density scans to evaluate the efficacy of pharmacotherapy [36, 48]. Sale reported that patients felt that had to 'nag' their physicians and follow up their own results [56].

Discussion

This systematic scoping review identified 33 studies that explored patients' perceived health service needs for osteoporosis. We identified specific health service needs among people with osteoporosis or osteopenia, highlighting opportunities for specific enhancement in models of service delivery for these conditions to ensure they continue to evolve in a patient-centred manner.

This review found that patients sought care from medical practitioners for the management of their osteoporosis [35, 43, 45, 48, 49]. In particular, patients tended to prefer management from specialists over primary care physicians. This is similar to other musculoskeletal conditions, such as low back pain [68, 69], and may reflect a lack of confidence or prioritisation by general practitioners in the management of bone health [70]. This may be attributed to limited knowledge of primary care providers [70] and suggests a need for future targeted education programs to bridge this gap, which have been shown to improve patient outcomes in osteoporosis as well as other chronic illnesses such as diabetes, asthma and congestive cardiac failure [71, 72]. Patients' expectation of healthcare providers was to perform a thorough examination, provide osteoporosis information and education, initiate screening for osteoporosis and to prescribe and monitor treatment [38, 41, 45, 48, 49, 56]. They wanted supportive and non-judgemental physicians [35, 45, 52], which enabled and promoted shared decision making. Indeed, this represents a key enabler to more effective self-management and sustainability to positive bone health behaviour change. They expressed dissatisfaction with the lack of time given by physicians, poor communication [35, 36, 43] and the inconsistent messages from different healthcare providers [35], again highlighting the need

Table 4 Patients' perceived needs of non-pharmacological management of osteoporosis

Author, year	Results
Calcium and vitamin D	
Bogoch 2008 [37]	• Patients generally agreed that regular exercise and calcium intake are beneficial in preventing osteoporosis
Feldstein 2008 [38]	• Patients expressed more willingness and comfort with taking supplements (calcium and vitamin D) than prescription medication for osteoporosis
Lau, 2008 [45]	• Calcium and vitamin D were perceived to be more “natural” than other osteoporosis medications and generally thought to be safe
Sale 2014 [57]	• Some participants watched their diet and/or taking supplements to improve their bone health • Patients exercise, have a healthy diet and take supplements to manage their bone health
Exercise therapy	
Bogoch 2008 [37]	• Patients generally agreed that regular exercise and calcium intake are beneficial in preventing osteoporosis
Sale 2014 [57]	• Patients exercise, have a healthy diet and take supplements to manage their bone health

for standardisation in cross-discipline education. Additionally, the dismissive approach, strong focus on pharmacotherapy and lack of continuity of care from healthcare providers were other areas of discontent among patients [35, 36, 43, 46, 48, 49]. It also underscores the patients' preference for patient-centred care and reinforces the need for clinicians to provide holistic care to improve the provider-patient relationship, which may facilitate improved uptake of osteoporosis clinical guidelines. This desire for improved communication from healthcare providers and holistic care is a common perceived need of patients with other chronic musculoskeletal conditions, including osteoarthritis, low back pain and inflammatory arthritides [24, 73].

Patients perceived a role for medications in the management of osteoporosis [36, 37, 39, 45, 54, 56, 59]. This is congruent with current clinical practice guidelines for osteoporosis which emphasise the use of pharmacotherapy [8–12], based on strong evidence for a number of effective medications in improving BMD and reducing fracture risk [74]. In

particular, this review found that individuals who were aware of the diagnosis of osteoporosis [47, 65], those who believed they were susceptible to future fractures [59], or had previous evaluation of their bone health [47] had a preference for medications. Furthermore, patients with a good relationship with their healthcare provider were more likely to have a preference for pharmacotherapy [54, 59], and this may reflect a more patient-centred approach to communication and shared therapeutic decision-making. Despite this perceived need for pharmacotherapy, there are high rates of treatment non-adherence for osteoporosis, with an estimated 50% of patients not taking medications by 12 months [75]. Educating patients regarding the benefits and rationale for effective pharmacotherapies for osteoporosis, a largely asymptomatic condition in the absence of fracture, may help to improve patient adherence with therapies and health outcomes, particularly a reduction in fracture risk [76, 77]. This contrasts with other chronic musculoskeletal conditions such as osteoarthritis, low back pain and inflammatory arthritis, where the perceived need for pharmacotherapy is often driven by a desire for symptom and pain control

Table 5 Patients' perceived needs of investigations for osteoporosis

Author, year	Results
Investigations for diagnosis	
Mazor 2010 [48]	• Patients noted the BMD test results at the time of diagnosis
Rothmann 2014 [53]	• Patients interpreted screening as an opportunity to get reassurance about bone status and take care of their own health.
Sale 2014 [56]	• Some participants reported persisting with the request to their family physician for a BMD test because of concern about their bones
Investigations for ongoing surveillance of bone health	
Besser 2012 [36]	• Patients wanted feedback from the DEXA scans to see if the medications were beneficial
Mazor 2010 [48]	• Patients thought the BMD results provided relevant feedback on the impact of their actions
Sale 2014 [56]	• Patients reported having to nag and follow up on their BMD test results

and maintenance of function and mobility [24, 73, 78–80]. Furthermore, addressing patients' concerns regarding pharmacotherapy, coupled with a broader approach to care that addresses lifestyle factors and support for effective self-management choices, may improve uptake of medications and health outcomes.

This review identified a number of patient beliefs regarding pharmacotherapy that may impact of adherence to osteoporosis pharmacotherapy. These included concerns regarding medication side effects, the potential for addiction and overdosing and the confusion and difficulty with the method of administration of medications [34, 36, 41–43, 45, 48, 53, 54, 60, 65, 66]. Furthermore, patients report a lack of knowledge about medications and they desire more health information [38, 43, 45, 48, 81, 82]. Medication non-adherence is also a growing concern in other chronic conditions, such as cardiovascular disease [83] and diabetes mellitus [84]. Poor adherence to medications is often multifactorial and may be due to patient, disease, medication, socioeconomic and healthcare system-related factors [85]. These areas of concern for osteoporosis pharmacotherapies may be addressed by multimodal interventions, including the provision of patient education and the development of novel systems to allow the mode of administration of medications to be more acceptable to patients and the use of technologies to prompt taking medications. Furthermore, the patients' beliefs and preferences for pharmacotherapy reported by the included studies need to be contextualised by healthcare providers. These findings demonstrate the breadth of patients' beliefs and preferences, and they may not apply to an individual patient. Clinicians should be cognisant of providing a tailored management approach to each specific patient, which may also improve the provider-patient relationship and foster a better therapeutic relationship.

Another finding from this review is that although some patients preferred medications [36, 37, 39, 48, 54, 56], they also perceived a need for lifestyle modifications and non-pharmacological therapies, such as exercise and vitamin supplementation to improve bone health [37, 38, 45, 57]. These non-pharmacological therapies were seen to be associated with lower-risk than prescription medications [38, 86]. Patients expressed dissatisfaction with the strong focus on pharmacotherapy from medical practitioners [48]. It appeared that driving the need for non-pharmacological therapies was the desire for a more holistic approach to healthcare management [36]. Despite exercise being a cornerstone therapy for the management of osteoporosis, a relatively smaller volume of literature was identified relating to patients' needs regarding exercise. This represents an important area for future exploration given the under-utilisation of exercise among people with osteoporosis. Capitalising on this need may also improve the relationship between providers and patients and improve osteoporosis outcomes. Integrating the patients' perceived needs of non-pharmacological management will improve

guideline adherence, especially as these recommend [8–12], based on evidence [74, 87–89] the use of physical therapy and vitamin D and calcium supplementation in osteoporosis management. However, there is a paucity of data regarding patients' perceived needs of other non-pharmacological lifestyle measures which may influence bone health, such as smoking cessation, attitudes to interventions related to falls prevention and avoidance of heavy alcohol: future research is required.

Clinical practice guidelines suggest the use of bone densitometry for the diagnosis of osteoporosis, to determine risk and need for therapy in people who have not sustained minimal trauma fractures [90]. This aligns with the findings of this review regarding the patients' perceived need of investigations for osteoporosis for diagnostic evaluation, and also for ongoing surveillance of the efficacy of pharmacotherapy [36, 48, 53, 56]. Yet, in spite of this, previous studies have found low rates of investigation of bone health in high-risk patients [18], thus, underscoring a lost window of opportunity to improve the uptake and adherence to pharmacotherapy. However, these studies included mainly older female participants, known to be at increased risk of osteoporosis: whether these results are generalizable to the perceived need for investigations in male patients with osteoporosis and younger women are unknown.

This review needs to be interpreted in light of a number of limitations. First, the results of this review have been inferred from heterogeneous studies that evaluated different study questions and had different inclusion criteria for participants. Furthermore, the majority of included studies were conducted in English-speaking, developed countries and examined elderly females. Thus, the results may not be generalizable to men, younger populations or people of different ethnicities and economies. Although our search strategy encompassed both primary and secondary osteoporosis, there were no studies identified that examined other high-risk groups such as those with long-term glucocorticoid use, end-stage renal failure and other secondary causes of osteoporosis. Moreover, many of the included studies were susceptible to bias, particularly regarding participant recruitment and data collection, as more interested patients may be inclined to participate in these studies. Also, some studies that evaluated pharmacotherapy for osteoporosis were funded by the pharmaceutical industry and many others did not acknowledge sources of funding or state the influence of funding on the study outcomes. These limitations in study quality highlight a need for future high-quality studies to confirm the findings in this review to better understand the patient's perceived needs for osteoporosis health services.

Despite these limitations, this review also has many strengths. A comprehensive scoping review was conducted across four complementary databases and included both qualitative and quantitative studies to capture the breadth of the existing literature. The rigorous and reproducible nature of our

methods therefore aligns with the intent of a systematic literature review, demonstrating a notable strength in our approach compared to narrative scoping reviews. The inclusion of qualitative studies provides invaluable insight into patient beliefs and attitudes and is particularly suitable for exploring biopsychosocial paradigms. Furthermore, several common themes emerged from the included studies, irrespective of study design or study quality; thus, this triangulation of data adds weight to the validity and credibility of the data. Additionally, participants were drawn from across care settings: from the community, from both primary care settings and hospital settings.

This systematic scoping review has identified patients' needs for improved health service delivery and better communication from healthcare professionals. Despite concerns regarding medication administration, side effects and compliance, patients have identified that osteoporosis pharmacotherapy is important. Patients also perceive a need for vitamin supplementation, exercise and ongoing surveillance of bone health. These findings may be unexpected given the low rates of screening and treatment for osteoporosis. Moving forward, the results from this review reinforce the need to improve the education provided not only to patients but also to cross-discipline healthcare practitioners regarding osteoporosis care. Workforce capacity building initiatives need to address the knowledge and skill deficits not only in pharmacologic management, including availability of different administration regimens for various therapies, but also important non-pharmacologic interventions like appropriate exercise and positive lifestyle choices. Given access limitations in many countries to medical specialists, capacity-building initiatives should be targeted in primary care settings. For consumers, education about the impact of osteoporosis and fractures remains critical to shift unhelpful nihilistic beliefs that the condition is an inevitable part of ageing and the risk-benefit balance of adherence of therapy. Their results confirm that clinicians need to provide patient-centred care through improved communication with patients, providing individualised information regarding the diagnosis and management of osteoporosis, encouraging multi-disciplinary shared care models and the use of decision aids to facilitate shared decision making. Moreover, given that poor treatment uptake is a significant practice gap in osteoporosis care, patient representatives should be involved in developing clinical practice guidelines and management initiatives to incorporate the patient perspective to develop patient-focused strategies, which may result in improved therapeutic relationships and compliance. The effects of this partnership will need to be evaluated to assess whether this ultimately translates into improved osteoporosis outcomes. These findings align well with the recent International Osteoporosis Foundation 2016 report [6], and together with the results from this review, provides important strategies for improving health services for people with bone

health impairments from multiple perspectives, which are critical to consider in any system-level reform initiatives.

Compliance with ethical standards

Conflict of interest None.

Funding This work was performed in partnership with Move: muscle, bone and joint health and was supported by a partnership grant awarded by the organisation. L.C is the recipient of an Australian Postgraduate Award and Arthritis Foundation Scholarship. A.E.W is the recipient of a NHMRC Career Development Fellowship (Clinical Level 2 #1063574). AMB is the recipient of a NHMRC TRIP Fellowship (#1132548).

References

1. Watts J, Abimanyi-Ochom J, Sanders K (2013) Osteoporosis costing all Australians: a new burden of disease analysis - 2012 to 2022. The University of Melbourne, Australia
2. Van Staa T, Dennison E, Leufkens H, Cooper C (2001) Epidemiology of fractures in England and Wales. *Bone* 29(6): 517–522
3. Burge R, Dawson-Hughes B, Solomon DH, Wong JB, King A, Tosteson A (2007) Incidence and economic burden of osteoporosis-related fractures in the United States, 2005–2025. *J Bone Miner Res* 22(3):465–475
4. Hernlund E, Svedbom A, Ivergard M, Compston J, Cooper C, Stenmark J et al (2013) Osteoporosis in the European Union: medical management, epidemiology and economic burden. A report prepared in collaboration with the International Osteoporosis Foundation (IOF) and the European Federation of Pharmaceutical Industry Associations (EFPIA). *Arch Osteoporos* 8:136
5. Si L, Winzenberg TM, Jiang Q, Chen M, Palmer AJ (2015) Projection of osteoporosis-related fractures and costs in China: 2010–2050. *Osteoporos Int* 26(7):1929–1937
6. Harvey NC, McCloskey EV (2016) Gaps and solutions in bone health. A global framework for improvement. International Osteoporosis Foundation, Nyon
7. Sanchez-Riera L, Carnahan E, Vos T, Veerman L, Norman R, Lim SS et al (2014) The global burden attributable to low bone mineral density. *Ann Rheum Dis* 73(9):1635–1645
8. Cosman F, de Beur SJ, LeBoff MS, Lewiecki EM, Tanner B, Randall S et al (2014) Clinician's guide to prevention and treatment of osteoporosis. *Osteoporos Int* 25(10):2359–2381
9. Kanis JA, McCloskey EV, Johansson H, Cooper C, Rizzoli R, Reginster JY (2013) European guidance for the diagnosis and management of osteoporosis in postmenopausal women. *Osteoporos Int* 24(1):23–57
10. Papaioannou A, Morin S, Cheung AM, Atkinson S, Brown JP, Feldman S et al (2010) Clinical practice guidelines for the diagnosis and management of osteoporosis in Canada: summary. *CMAJ: Canadian Medical Association journal = journal de l'Association medicale canadienne* 182(17):1864–1873
11. Wang M, Bolland M, Grey A (2016) Management recommendations for osteoporosis in clinical guidelines. *Clin Endocrinol* 84(5): 687–692
12. Watts NB, Bilezikian JP, Camacho PM, Greenspan SL, Harris ST, Hodgson SF et al (2010) American Association of Clinical Endocrinologists Medical Guidelines for clinical practice for the diagnosis and treatment of postmenopausal osteoporosis. *Endocr Pract* 16(Suppl 3):1–37

13. Mitchell P, Akesson K, Chandran M, Cooper C, Ganda K, Schneider M (2016) Implementation of models of care for secondary osteoporotic fracture prevention and orthogeriatric models of care for osteoporotic hip fracture. *Best Pract Res Clin Rheumatol* 30(3):536–558
14. Yates CJ, Chauchard MA, Liew D, Bucknill A, Wark JD (2015) Bridging the osteoporosis treatment gap: performance and cost-effectiveness of a fracture liaison service. *J Clin Densitom* 18(2):150–156
15. Sale JE, Beaton D, Posen J, Elliot-Gibson V, Bogoch E (2011) Systematic review on interventions to improve osteoporosis investigation and treatment in fragility fracture patients. *Osteoporos Int* 22(7):2067–2082
16. Kanis JA, Svedbom A, Harvey N, McCloskey EV (2014) The osteoporosis treatment gap. *J Bone Miner Res* 29(9):1926–1928
17. Giangregorio L, Papaioannou A, Cranney A, Zytaruk N, Adachi JD (2006) Fragility fractures and the osteoporosis care gap: an international phenomenon. *Semin Arthritis Rheum* 35(5):293–305
18. Elliot-Gibson V, Bogoch ER, Jamal SA, Beaton DE (2004) Practice patterns in the diagnosis and treatment of osteoporosis after a fragility fracture: a systematic review. *Osteoporos Int* 15(10):767–778
19. Díez-Pérez A, Hooven FH, Adachi JD, Adami S, Anderson FA, Boonen S et al (2011) Regional differences in treatment for osteoporosis. The global longitudinal study of osteoporosis in women (GLOW). *Bone* 49(3):493–498
20. Jennings LA, Auerbach AD, Maselli J, Pekow PS, Lindenauer PK, Lee SJ (2010) Missed opportunities for osteoporosis treatment in patients hospitalized for hip fracture. *J Am Geriatr Soc* 58(4):650–657
21. Briggs AM, Chan M, Slater H (2016) Models of care for musculoskeletal health: moving towards meaningful implementation and evaluation across conditions and care settings. *Best Pract Res Clin Rheumatol* 30(3):359–374
22. McLeod KM, Johnson CS (2011) A systematic review of osteoporosis health beliefs in adult men and women. *J Osteoporos* 2011:197454
23. Barker KL, Toye F, Lowe CJ (2016) A qualitative systematic review of patients' experience of osteoporosis using meta-ethnography. *Arch Osteoporos* 11(1):33
24. Wluka AE, Chou L, Briggs AM, FM C (2016) Understanding the needs of consumers with musculoskeletal conditions: consumers' perceived needs of health information, health services and other non-medical services: A systematic scoping review. *MOVE muscle, bone & joint health*
25. Arksey H, O'Malley L (2005) Scoping studies: towards a methodological framework. *Int J Soc Res Methodol* 8(1):19–32
26. Armstrong R, Hall BJ, Doyle J, Waters E (2011) 'Scoping the scope' of a cochrane review. *J Public Health* 33(1):147–150
27. Levac D, Colquhoun H, O'Brien KK (2010) Scoping studies: advancing the methodology. *Implement Sci* 5:69–69
28. Asadi-Lari M, Tamburini M, Gray D (2004) Patients' needs, satisfaction, and health related quality of life: towards a comprehensive model. *Health Qual Life Outcomes* 2:32–32
29. World Health Organisation, (WHO) http://www.who.int/topics/health_services/en/. Accessed 27 Oct 2016
30. Noblit G. HR. *Meta-ethnography: synthesising qualitative studies*. 1988 Sage Publications, California
31. Walsh D, Downe S (2005) Meta-synthesis method for qualitative research: a literature review. *J Adv Nurs* 50(2):204–211
32. Critical Appraisal Skills Programme (CASP) (2014) CASP Checklists (<http://www.casp-uk.net/casp-tools-checklists>) Oxford. CASP
33. Hoy D, Brooks P, Woolf A, Blyth F, March L, Bain C et al (2012) Assessing risk of bias in prevalence studies: modification of an existing tool and evidence of interrater agreement. *J Clin Epidemiol* 65:934–939
34. Yu J, Brennehan SK, Sazonov V, Modi A (2015) Reasons for not initiating osteoporosis therapy among a managed care population. *Patient Preference Adherence* 9:821–830
35. Sale JEM, Hawker G, Cameron C, Bogoch E, Jain R, Beaton D et al (2015) Perceived messages about bone health after a fracture are not consistent across healthcare providers. *Rheumatol Int* 35(1):97–103
36. Besser SJ, Anderson JE, Weinman J (2012) How do osteoporosis patients perceive their illness and treatment? Implications for clinical practice. *Arch Osteoporos* 7(1–2):115–124
37. Bogoch E, Elliot-Gibson V, Escott B, Beaton D (2008) The osteoporosis needs of patients with wrist fracture. *J Orthop Trauma* 22(8 Suppl):S73–S78
38. Feldstein AC, Schneider J, Smith DH, Vollmer WM, Rix M, Glauber H et al (2008) Harnessing stakeholder perspectives to improve the care of osteoporosis after a fracture. *Osteoporos Int* 19(11):1527–1540
39. Fraenkel L, Gulanski B, Wittink D (2006) Preference for hip protectors among older adults at high risk for osteoporotic fractures. *J Rheumatol* 33(10):2064–2068
40. Gold D, Safi W, Trinh H (2006) Patient preference and adherence: comparative US studies between two bisphosphonates, weekly risedronate and monthly ibandronate. *Curr Med Res Opin* 22(12):2383–2391
41. Hansen C, Konradsen H, Abrahamsen B, Pedersen BD (2014) Women's experiences of their osteoporosis diagnosis at the time of diagnosis and 6 months later: a phenomenological hermeneutic study. *Int J Qual Stud Health Well-being* 9:22438
42. Hiligsmann M, Dellaert B, Dirksen C, Van Der Qeijden T, Goemaere S, Reginster J et al (2014) Patients' preferences for osteoporosis drug treatment: a discrete-choice experiment. *Arthritis Res Ther* 16(1):R36
43. Iversen MD, Vora RR, Servi A, Solomon DH (2011) Factors affecting adherence to osteoporosis medications: a focus group approach examining viewpoints of patients and providers. *J Geriatr Phys Ther* 34(2):72–81
44. Keen R, Jodar E, Iolascon G, Kruse H, Varbanov A, Mann B et al (2006) European women's preference for osteoporosis treatment: influence of clinical effectiveness and dosing frequency. *Curr Med Res Opin* 22(12):2375–2381
45. Lau E, Papaioannou A, Dolovich L, Adachi J, Sawka AM, Burns S et al (2008) Patients' adherence to osteoporosis therapy: exploring the perceptions of postmenopausal women. *Can Fam Physician* 54(3):394–402
46. Martin A, Holmes R, Lydick E (1997) Fears, knowledge, and perceptions of osteoporosis among women. *Drug Inf J* 31(1):301–306
47. Mauck K, Cuddlhy M, Trousdale R, Pond G, Pankratz V, Melton LJ 3rd (2002) The decision to accept treatment for osteoporosis following hip fracture: exploring the woman's perspective using a stage-of-change model. *Osteoporos Int* 13(7):560–564
48. Mazar KM, Velten S, Andrade SE, Yood RA (2010) Older women's views about prescription osteoporosis medication: a cross-sectional, qualitative study. *Drugs Aging* 27(12):999–1008
49. McKenna J, Ludwig AF (2008) Osteoporotic Caucasian and South Asian women: a qualitative study of general practitioners' support. *J R Soc Promot Heal* 128(5):263–270 268p
50. Payer J, Ciemy D, Killinger Z, Sulkova I, Behuliac M, Celec P (2009) Preferences of patients with post-menopausal osteoporosis treated with bisphosphonates—the VIVA II study. *J Int Med Res* 37(4):1225–1229
51. Richards J, Cherkas L, Spector T (2007) An analysis of which anti-osteoporosis therapeutic regimen would improve compliance in a population of elderly adults. *Curr Med Res Opin* 23(2):293–299
52. Rizzoli R, Brandi M, Dreinhofer K, Thomas T, Wahl D, Cooper C (2010) The gaps between patient and physician understanding of the emotional and physical impact of osteoporosis. *Arch Osteoporos* 5(1–2):145–153

53. Rothmann MJ, Huniche L, Ammentorp J, Barkmann R, Gluer CC, Hermann AP (2014) Women's perspectives and experiences on screening for osteoporosis (risk-stratified osteoporosis strategy evaluation, ROSE). *Arch Osteoporos* 9:192
54. Sale J, Beaton D, Sujic R, Bogoch E (2010) 'If it was osteoporosis, I would have really hurt myself'. Ambiguity about osteoporosis and osteoporosis care despite a screening programme to educate fragility fracture patients. *J Eval Clin Pract* 16(3):590–596
55. Sale J, Bogoch E, Hawker G, Gignac M, Beaton D, Jaglal S et al (2014) Patient perceptions of provider barriers to post-fracture secondary prevention. *Osteoporos Int* 25(11):2581–2589
56. Sale J, Cameron C, Hawker G, Jaglal S, Funnell L, Jain R et al (2014) Strategies used by an osteoporosis patient group to navigate for bone health care after a fracture. *Arch Orthop Trauma Surg* 134(2):229–235
57. Sale J, Gignac M, Hawker G, Beaton D, Bogoch E, Webster F et al (2014) Non-pharmacological strategies used by patients at high risk for future fracture to manage fracture risk—a qualitative study. *Osteoporos Int* 25(1):281–288
58. Saltman D, Sayer G, O'Dea N (2006) Dosing frequencies in general practice—whose decision and why? *Aust Fam Physician* 35(11):915–919
59. Schousboe J, Davison M, Dowd B, Thiede Call K, Johnson P, Kane R (2011) Predictors of patients' perceived need for medication to prevent fracture. *Med Care* 49(3):273–280
60. Scoville E, Ponce de Leon Lovaton P, Shah N, Pencille L, Montori V (2011) Why do women reject bisphosphonates for osteoporosis? A videographic study. *PLoS ONE* 6(4):e18468
61. Turbi C, Herrero-Beaumont G, Acebes J, Torrijos A, Grana J, Miguelez R et al (2004) Compliance and satisfaction with raloxifene versus alendronate for the treatment of postmenopausal osteoporosis in clinical practice: an open-label, prospective, nonrandomized, observational study. *Clin Ther* 26(2):245–256
62. Walliser J, Bolge S, Sen S (2006) Patients' preference for osteoporosis medications: prefer-international. *J Clin Rheumatol* 12(4):S29
63. Weiss M, Vered I, Foldes A, Cohen Y, Shamir-Elron Y, Ish-Shalom S (2005) Treatment preference and tolerability with alendronate once weekly over a 3-month period: an Israeli multi-center study. *Aging Clin Exp Res* 17(2):143–149
64. Weiss T, McHorney C (2007) Osteoporosis medication profile preference: results from the PREFER-US study. *Health Expect* 10(3):211–223
65. Yood RA, Mazor KM, Andrade SE, Ermani S, Chan W, Kahler K (2008) Patient decision to initiate therapy for osteoporosis: the influence of knowledge and beliefs. *J Gen Intern Med* 23(11):1815–1821
66. Zanchetta J, Hakim C, Lombas C (2004) Observational study of compliance and continuance rates of raloxifene in the prevention and treatment of osteoporosis. *Curr Ther Res Clin Exp* 65(6):470–480
67. Ringe JD, van der Geest SA, Moller G (2006) Importance of calcium co-medication in bisphosphonate therapy of osteoporosis: an approach to improving correct intake and drug adherence. *Drugs Aging* 23(7):569–578
68. Amonkar SJ, Dunbar AM (2011) Do patients and general practitioners have different perceptions about the management of simple mechanical back pain? *Int Musculoskelet Med* 33(1):3–7
69. Carey TS, Garrett J, Jackman A, McLaughlin C, Fryer J, Smucker DR (1995) The outcomes and costs of care for acute low back pain among patients seen by primary care practitioners, chiropractors, and orthopedic surgeons. The North Carolina back pain project. *N Engl J Med* 333(14):913–917
70. Otmar R, Reventlow SD, Nicholson GC, Kotowicz MA, Pasco JA (2012) General medical practitioners' knowledge and beliefs about osteoporosis and its investigation and management. *Arch Osteoporos* 7:107–114
71. Forsetlund L, Bjorndal A, Rashidian A, Jamtvedt G, O'Brien MA, Wolf F et al (2009) Continuing education meetings and workshops: effects on professional practice and health care outcomes. *Cochrane Database Syst Rev* 2:CD003030
72. Weingarten SR, Henning JM, Badamgarav E, Knight K, Hasselblad V, Gano A Jr et al (2002) Interventions used in disease management programmes for patients with chronic illness—which ones work? Meta-analysis of published reports. *BMJ* 325(7370):925
73. Papandony MC, Chou L, Seneviwickrama M, Cicuttini FM, Lasserre K, Teichtahl AJ et al (2017) Patients' perceived health service needs for osteoarthritis (OA) care: a scoping systematic review. *Osteoarthr Cartil* 25:1010–1025
74. Body JJ, Bergmann P, Boonen S, Boutsens Y, Devogelaer JP, Goemaere S et al (2010) Evidence-based guidelines for the pharmacological treatment of postmenopausal osteoporosis: a consensus document by the Belgian bone Club. *Osteoporos Int* 21(10):1657–1680
75. Weycker D, Macarios D, Edelsberg J, Oster G (2006) Compliance with drug therapy for postmenopausal osteoporosis. *Osteoporos Int* 17(11):1645–1632
76. Kastner M, Straus SE (2008) Clinical decision support tools for osteoporosis disease management: a systematic review of randomized controlled trials. *J Gen Intern Med* 23(12):2095–2105
77. Winzenberg T, Oldenburg B, Frendin S, De Wit L, Riley M, Jones G (2006) The effect on behavior and bone mineral density of individualized bone mineral density feedback and educational interventions in premenopausal women: a randomized controlled trial [NCT00273260]. *BMC Public Health* 6:12
78. Jensen AL, Lomborg K, Wind G, Langdahl BL (2014) Effectiveness and characteristics of multifaceted osteoporosis group education—a systematic review. *Osteoporos Int* 25(4):1209–1224
79. Laliberte MC, Perreault S, Jouini G, Shea BJ, Lalonde L (2011) Effectiveness of interventions to improve the detection and treatment of osteoporosis in primary care settings: a systematic review and meta-analysis. *Osteoporos Int* 22(11):2743–2768
80. Smith CA (2010) A systematic review of healthcare professional-led education for patients with osteoporosis or those at high risk for the disease. *Orthop Nurs* 29(2):119–132
81. French MR, Moore K, Vernace-Inserra F, Hawker GA (2005) Factors that influence adherence to calcium recommendations. *Can J Diet Pract Res* 66(1):25–29
82. Sale JE, Beaton DE, Sujic R, Bogoch ER (2010) 'If it was osteoporosis, I would have really hurt myself'. Ambiguity about osteoporosis and osteoporosis care despite a screening programme to educate fragility fracture patients. *J Eval Clin Pract* 16(3):590–596
83. Ho PM, Bryson CL, Rumsfeld JS (2009) Medication adherence. Its importance in cardiovascular outcomes. *Circulation* 119(23):3028–3035
84. Krass I, Schieback P, Dhippayom T (2015) Adherence to diabetes medication: a systematic review. *Diabet Med* 32(6):725–737
85. World Health Organisation (2003) Adherence to long-term therapy: evidence for action Available at: http://www.who.int/chp/knowledge/publications/adherence_introduction.pdf. Accessed 27 Oct 2016
86. Lau E, Papaioannou A, Dolovich L, Adachi J, Sawka AM, Burns S et al (2008) Patients' adherence to osteoporosis therapy: exploring the perceptions of postmenopausal women. *Can Fam Physician* 54:394–402 399p
87. Body JJ, Bergmann P, Boonen S, Boutsens Y, Bruyere O, Devogelaer JP et al (2011) Non-pharmacological management of osteoporosis: a consensus of the Belgian bone Club. *Osteoporos Int* 22(11):2769–2788

88. Howe TE, Shea B, Dawson LJ, Downie F, Murray A, Ross C, et al. (2011) Exercise for preventing and treating osteoporosis in postmenopausal women. *Cochrane Database Syst Rev* (7): CD000333
89. Kemmler W, Haberle L, von Stengel S (2013) Effects of exercise on fracture reduction in older adults: a systematic review and meta-analysis. *Osteoporos Int* 24(7):1937–1950
90. Kanis J, McCloskey E, Johansson H, Cooper C, Rizzoli R, Reginster J (2012) European guidance for the diagnosis and management of osteoporosis in postmenopausal women. *Osteoporos Int* 24(1):23–57