



# The small RNA *STnc1480* contributes to the regulation of biofilm formation and pathogenicity in *Salmonella typhimurium*

Jing Li<sup>1</sup> · Chengcheng Ning<sup>1</sup> · Na Li<sup>1</sup> · Yun Guo<sup>1</sup> · Chunhui Ji<sup>1</sup> · Xiaozhen Zhu<sup>1</sup> · Xingxing Zhang<sup>2</sup> · Qingling Meng<sup>1</sup> · Xianzhu Xia<sup>1</sup> · Xuepeng Cai<sup>3</sup> · Jun Qiao<sup>1,4</sup>

Received: 14 April 2022 / Revised: 14 October 2022 / Accepted: 10 November 2022 / Published online: 19 November 2022  
© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2022

## Abstract

*Salmonella Typhimurium* (STM) is one of the most important food-borne bacteria that seriously harms livestock and human beings, which is capable of regulating the expression of its own genes in a variety of ways to adapt to a wide variety of adverse environmental stresses. To understand the regulatory roles of sRNA *STnc1480* on the capability of STM, the *STnc1480* gene-deficient strain  $\Delta$ *STnc1480* and its complement strain  $\Delta$ *STnc1480*/*STnc1480* were generated, and the impacts of *STnc1480* gene deficiency on the capability of responding to different environmental stresses, biofilm(BF)formation and pathogenicity were analyzed, respectively. Then the target genes that were regulated by *STnc1480* were also analyzed and explored. Compared with parent and complement strains, the deficiency of the *STnc1480* gene significantly reduced the BF formation. Moreover, the capacities of adhesion and invasiveness of the  $\Delta$ *STnc1480* strain to macrophages were also significantly reduced, while the LD<sub>50</sub> in mice was significantly increased. The bacterial loads in liver and spleen were significantly reduced, and the pathological damage was alleviated. It was confirmed that the *STnc1480* could be complementary to the 5'-UTR (-52 to -71 bases) region of *lpfA* mRNA. The bacterial dual-plasmid reporting system confirmed that *STnc1480* was capable of interacting with the mRNA of the *lpfA* gene, suggesting that *STnc1480* can regulate the 5'-UTR of the *lpfA* mRNA at post-transcription level to reduce the expression of the bacterial fimbria, thus reducing the BF formation and pathogenicity of STM.

**Keywords** *STnc1480* · *Salmonella typhimurium* · *lpfA* gene · Environmental stress · Biofilm · Pathogenicity

---

Communicated by Erko Stackebrandt.

---

Jing Li, Chengcheng Ning, and Na Li contributed equally to this work.

---

✉ Jun Qiao  
qj710625@shzu.edu.cn

Jing Li  
xiaoshouyii@163.com

Chengcheng Ning  
1214858343@qq.com

Na Li  
1505376491@qq.com

Yun Guo  
2722369104@qq.com

Chunhui Ji  
1922531568@qq.com

Xiaozhen Zhu  
1872200351@qq.com

Xingxing Zhang  
shzuzx@163.com

Qingling Meng  
2448986506@qq.com

Xianzhu Xia  
1012385302@qq.com

Xuepeng Cai  
caixp@vip.163.com

- <sup>1</sup> College of Animal Science and Technology, Shihezi University, Shihezi 832003, Xinjiang, China
- <sup>2</sup> Institute of Animal Science and Veterinary Research, Xinjiang Academy of Agricultural and Reclamation Science, Shihezi 832000, Xinjiang, China
- <sup>3</sup> State Key Lab of Veterinary Etiological Biology, Lanzhou Veterinary Research Institute, Chinese Academy of Agricultural Sciences, Lanzhou 730046, Gansu, China
- <sup>4</sup> College of Animal Science and Technology, Shihezi University, 280 N 4th road, Shihezi 832000, Xinjiang, China

## Introduction

*Salmonella Typhimurium* (STM) is a Gram-negative intracellular bacterium with a wide range of host, which is widely distributed in the digestive tracts of livestock, birds and human (Fàbrega and Vila 2013; Majowicz et al. 2010). As one of important food-borne pathogens, STM can infect human through the contaminated meat, milk, eggs, and other animal-derived foods, resulting in human acute gastroenteritis, which has posed a serious threat to global food hygiene and safety. In the recent years, food poisoning caused by STM has occurred frequently in many countries, especially in the developing countries, causing serious harm to human health (Senior 2009).

Small RNA (sRNA) is non-coding RNA with about 50–400 nucleotides (nt) in length (Fröhlich and Vogel 2009). They are widely found in the genomes of many bacterial species (Harfouche et al. 2015). The currently available studies have found that sRNA is involved in regulation of the expression of certain genes of bacteria in adverse stress environments, and play an important role in responding to the environmental stresses, community effect, and the pathogenicity of bacteria (Fröhlich and Vogel 2009; Wagner and Romby 2015). A number of studies have confirmed that bacterial sRNA can perform a variety of biological functions by binding to mRNAs of their target genes to adapt to the environmental changes by regulating them at the post-transcriptional level (Papenfert and Vogel 2009).

Recently, more than 100 non-coding-sRNAs have been found and identified in *Salmonella* (Kröger et al. 2012). However, the biological functions of most of these sRNAs

are not yet known. A few studies have revealed that the expression level of sRNA *STnc1480* with 395 nucleotides in length was significantly increased during the infection process of STM into macrophages (Kröger et al. 2012; Srikumar et al. 2015). However, the regulatory roles of *STnc1480* on the response of STM to environmental stress, biofilm (BF) formation, and pathogenicity of STM remain unclear. Therefore, the main purpose of this study was to understand the regulatory roles of *STnc1480* on environmental stress, BF formation, and pathogenicity in STM. Here, we constructed and analyzed the response to environmental stresses, BF formation, and pathogenicity of the *STnc1480* gene-deficient strain and complement strain, and identified the target genes regulated by sRNA *STnc1480*, aiming to provide new insights into the molecular mechanism of sRNA regulation in STM.

## Materials and methods

### Design of the primers

According to the STM SL1344 (accession number: FQ312003.1) sequence and pKD3 (AY048742.1) plasmid sequence deposited in the GeneBank, we designed the specific primers using Primer 5.0 software (Premier Inc., Canada), and synthesized by Beijing Genomics Institute (China). All the primers used in this study are listed in Table 1.

**Table 1** Primers used in this study

Primer	Sequences (5'–3')	Product size (bp)	Template plasmid/gene target
F1	TAAGTCGGATGCGTTAGTTA	760/1394/498	SL1344/ $\Delta$ STnc1480::cat/ $\Delta$ : STnc1480
R1	CAGACCTTATGGCGCGAGTATGA		
F2	CGGCATCGATTAACTAACGAAATAAAATAGCC TTTATTATTACGCGGTGTAGGCTGGAGCTGCTTC	1114	pKD3
R2	TATTCCGCGAATTGTTTCAGAAACAGAGCGCCAATC AAAGTGCATAACCATATGAATATCCTCCTTAGTT		
F3	AAGCTTTAAGTCGGATGCGTTAGTTA	760	SL1344/pBR322-STnc1480
R3	GGATCCCAGACCTTATGGCGCGAGTATGA		
16s-F	ATGCCGCGTGTATGAAGAAGG	132	SL1344
16s-R	GCGGCTGCTGGCACGGAGTTAG		
lpfA-F	GCCATGCGTCGTTTCTACTG	294	SL1344
lpfA-R	GTGGATTACCGGTGTTTCATTC		
F4	CGGGATCCATGCGTTAGTTTACCATAG	642	SL1344/ pUT18C-STnc1480
R4	GGGGTACCAAACAGAGCGCCAATCAAAGTGC		
F5	CCCAAGCTT AATTATTTTTATATATAC	285	SL1344/ pMR-LacZ-lpfA
R5	CGGGGTACCATTGACCCAGCACAACCTCC		

## Generation of STnc1480-deficient mutant strain and complement strains

The strains and plasmid used are listed in Table 2. STM Wild Type (WT) SL1344 strains were cultured with brain–heart immersion (BHI) broth (Hopebio, China) at 37 °C for 12–16 h. The gene sequence of STM STnc1480 was amplified with F1/R1 primers. The STnc1480-deficient mutant strain was constructed using  $\lambda$ -Red recombination technology (Datsenko and Wanner 2000). Briefly, chloramphenicol-containing gene products were first amplified by PCR with F2/R2 primer pair, which were used to construct recombinant strain  $\Delta$ : STnc1480:: cat. Then the STnc1480-deficient mutant strain was identified by PCR amplification with F1/R1 primer pair in combination with sequencing technology. The complement strain  $\Delta$ STnc1480/STnc1480 was constructed using F3/R3 primer pair.

## Environmental stresses assay

Briefly, the strains of SL1344,  $\Delta$ STnc1480, and  $\Delta$ STnc1480/STnc1480 were inoculated into 50 mL of BHI liquid medium, and cultured at 37 °C overnight; This was followed by inoculation of the bacteria on an LB agar plate overnight and observation of the individual colony sizes by continuous zoom microscopy. The bacterial optical density at 600 nm ( $OD_{600nm}$ ) value was adjusted to 0.5, the bacteria solution was inoculated at 1:100 ratio in the BHI liquid medium and cultured continuously at 37 °C, their  $OD_{600}$  values were measured every hour and their growth curves were plotted. At the same time and 37°C, the bacterial solutions were inoculated at a ratio of 1:100, respectively, to HCl-adjusted pH 4.8 BHI liquid medium, NaOH-adjusted pH 10 BHI liquid medium, and BHI liquid medium containing 2 M NaCl and 2 mM 30%  $H_2O_2$ . The  $OD_{600}$

values were measured every hour and their growth curves were plotted. The effects of *STnc1480* gene deficiency on the responses of STM to different environmental stress conditions were examined and analyzed.

## Biofilm formation assay

The overnight-cultured active bacterial solution was inoculated in 1:100 ratio into the BHI liquid medium and cultured up to its  $OD_{600nm}$  of 0.2. Then 100  $\mu$ L of bacteria solution was aspired and transferred to 96-well microplates. Each sample was divided into two groups and eight parallel repetitions were set for each group. BF of STM was measured according to methods reported in the literature (Kint et al. 2010). After being cultured for 22, 23, and 24 h, the differences in BF formation capability between SL1344,  $\Delta$ STnc1480, and  $\Delta$ STnc1480/STnc1480 strains were compared and the effects of *STnc1480* gene deficiency on STM BF formation capacity were analyzed.

## Cell adhesion, invasion, and intracellular survival assay

Briefly, mouse macrophages RAW264.7 were placed in culture incubator at 37 °C and 5% carbon dioxide ( $CO_2$ ) and cultured overnight to grow into a single layer with about  $2 \times 10^5$  cells per well. The bacterial adhesion and invasion tests were performed according to the methods described in the literature with a slight modification (Ryan et al. 2018). After infection of 1 h, for the cell invasion test, the bacterial cells that did not enter the macrophages were killed with Dulbecco's modified Eagle medium (DMEM) (Gibco, USA) containing gentamicin [(100 mg/mL) (BIOTOPPED, China)] at 37 °C. The remaining steps were the same as those described above. The adhesion rates and invasion rates

**Table 2** Strains and plasmids used in this study

Strains	Characteristic	Reference
<i>Salmonella enterica serovar Typhimurium</i> SL1344	Wild type	8
$\Delta$ STnc1480:: cat	STnc1480 deletion mutant carrying cat, $Cm^r$	This study
$\Delta$ : STnc1480	STnc1480 deletion mutant	This study
$\Delta$ : STnc1480/STnc1480	Complementation of $\Delta$ : STnc1480, $Amp^r$	This study
BTH101 <i>Escherichia coli</i>		This study
Plasmids		
pKD3	Contains FRT-flanked chloramphenicol resistance gene	11
pKD46	Expresses lambda red recombinase, $Amp^r$	11
pCP20	Expresses FLP recombinase, $Amp^rCm^r$	11
pBR322	Cloning vectors, $Amp^r$	This study
pMD19-T	Cloning vectors, $Amp^r$	Takara
PUT18C	$Amp^r$	This study
pMR-LacZ	$Kan^r$	This study

were calculated. Intracellular survival tests were conducted with reference to the literature (Hölzer et al. 2009). At 2, 4, 6, 8, 10, and 12 h after culture, cultured solution was aspirated, cleaned, and lysed. The bacterial colonies were counted. The test was repeated three times and three parallels were set each time.

### Animal infections

One hundred and fourteen 6-week-old BALB/c mice were randomly divided into one control group and six different dilution of bacterial infection groups  $2 \times 10^2$  cfu/mL,  $2 \times 10^3$  cfu/mL,  $2 \times 10^4$  cfu/mL,  $2 \times 10^5$  cfu/mL,  $2 \times 10^6$  cfu/mL, and  $2 \times 10^7$  cfu/mL, respectively. STM SL1344,  $\Delta$ STnc1480, and  $\Delta$ STnc1480/STnc1480 strains were cultured in BHI liquid medium, respectively, and used to infect mice when the logarithmic phase  $OD_{600}$  was 0.58. The mice in each subgroup were intraperitoneally injected with 0.5 mL of serially diluted cultures and the control group was treated with the same volume of phosphate buffered saline (PBS) and observed continuously for 10 days, and the  $LD_{50}$  of the mice was calculated by the modified Kurt method (Angalabiri-Owei and Isirima 2014). Six-week-old mice were infected with  $2 \times 10^4$  cfu/mL SL1344,  $\Delta$ STnc1480, and  $\Delta$ STnc1480/STnc1480 strains, respectively. Mice were sacrificed on days 1, 3, 5, 7, and 9 after infection, and the amounts of bacteria in these organs were simultaneously measured (Kumawat et al. 2016). Meanwhile, the liver and spleen were collected and pathological tissue sections were prepared 5 days after injection and the histopathological changes were observed by optical microscopy (Olympus, Japan).

### Prediction and screening of target genes regulated by STnc1480

TargetRNA2 online software was used to predict the target genes of *STnc1480* (Kery et al. 2014). The STM SL1344 genome was selected and the *STnc1480* RNA sequence was entered. The genes with longer bases being paired with *STnc1480* were selected as the target genes.

### RNA extraction and real-time quantitative PCR

Briefly, the bacteria were cultured to BF formation state, and then collected by centrifugation. Total RNA was extracted with TRIzol reagent (Invitrogen, USA). Then complementary DNA (cDNA) was synthesized using PrimeScript-RRT kit including gDNA Eraser (TaKaRa Bio, Inc., Japan). Briefly, using 16sF/16sR and *lpfAF/lpfAR* as

primers, the transcription levels of the target genes were determined by real-time quantitative PCR (RT-qPCR) using the ABI7500 instrument (Applied Biosystems, USA). Using the 16sRNA gene as the reference gene, the relative transcription levels of the target genes were calculated by the  $2^{-\Delta\Delta CT}$  method (Livak and Schmittgen 2001). Each test was repeated three times.

### The bacterial dual-plasmid reporting system experiment

The interaction between *STnc1480* and its target genes was analyzed using the bacterial dual-plasmid reporter system. In brief, F4/R4 and F5/R5 primers were used to amplify the operon DNA sequences of *STnc1480* and *lpfA*. Then the pUT18C-*STnc1480* and pMR-LacZ-*lpfA* plasmids were constructed; the constructed recombinant plasmids were co-transfected into BTH101 *E. coli* competent cells via electroporation. The positive clones on a Kan<sup>r</sup> (100 mg/mL) Amp<sup>r</sup> (100 mg/9 mL) (BIOTOPPED, China) double-resistant plate coated with X-gal (TaKaRa Bio, Inc., Japan) and isopropyl-beta-D-thiogalactopyranoside (IPTG) (Solarbio Science & Technology Co., Ltd., China) were screened out and were placed at 37 °C overnight, and the color change of the bacterial lawn was observed.


### Statistical analysis of data

GraphPad Prism 5.0 software (<https://www.graphpad.com/>, USA) was used for data analysis, and *t* test was used to conduct data significance analysis. The value of  $P < 0.05$  was considered significantly different, while  $P < 0.01$  was considered extremely significant different.

## Results

### STnc1480 does not affect the growth of STM in different environments

Through PCR and sequencing confirmation, the deletion strain  $\Delta$ STnc1480 and the complement strain  $\Delta$ STnc1480/STnc1480 were obtained (Supplementary Fig. S1 and Supplementary Fig. S2), respectively. The growth rates and bacterial single colony of the SL1344,  $\Delta$ STnc1480, and  $\Delta$ STnc1480/STnc1480 at 37 °C, were not significantly different ( $P > 0.05$ ) (Fig. 1A) and Supplementary Fig. S4).





**Fig. 1** Growth curves of SL1344,  $\Delta$ STnc1480, and  $\Delta$ STnc1480/STnc1480 strains under different conditions. A–E: Growth curves of SL1344,  $\Delta$ STnc1480, and  $\Delta$ STnc1480/STnc1480 strains at

37 °C, pH4.8, pH10, 2 M NaCl, 2 mM 30%  $H_2O_2$ , respectively.  $OD_{600}$  = optical density at 600 nm

When being cultured under the condition of pH4.8, pH10, 2 M NaCl, and 2 mM 30%  $H_2O_2$ , the growth of  $\Delta$ STnc1480 was not significantly different from those of the complement strain and the parent strain ( $P > 0.05$ ) (Fig. 1B, C, D and E), suggesting that *STnc1480* does not affect the growth of STM in different environments.


### STnc1480 affects biofilm formation of STM

Cultured for 22, 23, and 24 h, respectively, the SL1344,  $\Delta$ STnc1480, and  $\Delta$ STnc1480/STnc1480 were all capable of forming BF (Fig. 2A). However, the BF-forming ability of  $\Delta$ STnc1480 strain was significantly weakened



**Fig. 2** Assay of biofilm formation ability of SL1344,  $\Delta$ STnc1480, and  $\Delta$ STnc1480/STnc1480 strain, respectively. **A** Biofilm observation by microscopic (40 $\times$ 10), **B** Biofilm forming abilities

determined by OD<sub>600</sub>. Values represent the mean  $\pm$  SEM (\**P* < 0.05) OD<sub>600</sub> = optical density at 600 nm



**Fig. 3** Determination of cellular adhesion, invasion, and survival abilities in the RAW264.7 cell line. **A** cellular adhesion rate, **B** cellular invasion rate, **C** intracellular survival abilities. Values represent the mean  $\pm$  SEM (\* $P < 0.05$ , \*\* $P < 0.01$ ). CFU colony-forming unit

compared to the others at the indicated time ( $P < 0.05$ ) (Fig. 2B), suggesting that *STnc1480* had a potential influence on BF formation of STM.

### **STnc1480 affects adherence, invasion, and intracellular survival of STM to cells**

Compared with those of SL1344 and  $\Delta$ STnc1480/STnc1480, the adhesion of  $\Delta$ STnc1480 to RAW264.7 cells

was extremely significantly weakened ( $P < 0.01$ ) (Fig. 3A), and the invasion ability was also significantly weakened ( $P < 0.05$ ) (Fig. 3B). At 6, 8, 10, and 12 h after infection, the quantity of  $\Delta$ STnc1480 strains was significantly lower than those of SL1344 and  $\Delta$ STnc1480/STnc1480 strains ( $P < 0.01$ ) (Fig. 3C and Supplementary Fig. S5), indicating that *STnc1480* affects the intracellular survival of STM.

**Table 3** Determination of LD<sub>50</sub> of SL1344, ΔSTnc1480, and ΔSTnc1480/STnc1480 in 6-week-old mice after intraperitoneal injection

Strains	cfu/mL	Mice	Mouse death	LD <sub>50</sub> (cfu/mL)
SL1344	2 × 10 <sup>2</sup>	6	1	3.6 × 10 <sup>4</sup>
	2 × 10 <sup>3</sup>	6	1	
	2 × 10 <sup>4</sup>	6	2	
	2 × 10 <sup>5</sup>	6	5	
	2 × 10 <sup>6</sup>	6	6	
	2 × 10 <sup>7</sup>	6	6	
ΔSTnc1480	2 × 10 <sup>2</sup>	6	0	4 × 10 <sup>5</sup>
	2 × 10 <sup>3</sup>	6	0	
	2 × 10 <sup>4</sup>	6	1	
	2 × 10 <sup>5</sup>	6	2	
	2 × 10 <sup>6</sup>	6	6	
	2 × 10 <sup>7</sup>	6	6	
ΔSTnc1480/ STnc1480	2 × 10 <sup>2</sup>	6	0	8.5 × 10 <sup>4</sup>
	2 × 10 <sup>3</sup>	6	1	
	2 × 10 <sup>4</sup>	6	1	
	2 × 10 <sup>5</sup>	6	5	
	2 × 10 <sup>6</sup>	6	6	
	2 × 10 <sup>7</sup>	6	6	
Control	PBS	6	0	–


CFU colony-forming unit, LD<sub>50</sub> = 50% lethal dose

### Deleting STnc1480 attenuates virulence of STM in mice

The LD<sub>50</sub> of SL1344, ΔSTnc1480, and ΔSTnc1480/STnc1480 were 3.6 × 10<sup>4</sup> cfu/ mL, 4 × 10<sup>5</sup> cfu/ mL, and 8.5 × 10<sup>4</sup> cfu/ mL, respectively (Table 3). The survival curves indicated that the pathogenicity of ΔSTnc1480 group was significantly reduced (Fig. 4C). On the 5th, 7th, and 9th days after infection, as compared with those of SL1344 and ΔSTnc1480/STnc1480 groups, the bacterial loads of ΔSTnc1480 group in the liver and spleen of mice were significantly lower ( $P < 0.01$ ) (Fig. 4A, B). Compared with those of SL1344 and ΔSTnc1480/STnc1480 groups, the histopathological damages in liver, spleen, and small intestine of the mice infected with the ΔSTnc1480 were alleviated (Fig. 5).

### STnc1480 regulates *lpfA* expression

TargetRNA2 prediction of the target genes of *STnc1480* led to the finding that the *STnc1480* were complementarily paired with the 5'-UTR (– 52 to – 71 bases) of *lpfA* mRNA




**Fig. 4** Bacterial loads and survival curves in mice infected by SL1344, ΔSTnc1480, and ΔSTnc1480/STnc1480 strain, respectively. **A** Liver, **B** spleen, **C** survival curve. Values represent the mean ± SEM (\*\* $P < 0.01$ ). CFU = colony-forming unit

(Fig. 6A, B). In the state of BF formation, RT-qPCR analysis results showed that compared with SL1344 and ΔSTnc1480/STnc1480, the transcription level of *lpfA* gene in ΔSTnc1480 strain was significantly lower ( $P < 0.01$ ) (Fig. 7), suggesting that the expression of *lpfA* gene may be regulated by the *STnc1480*.

### STnc1480 can interact with target gene *lpfA*

The pMR-LacZ-*lpfA* and pUT18C-*STnc1480* plasmids were constructed (Supplementary Fig. S3). Compared with the transformed pUT18C, pMR-LacZ and the co-transformed pUT18C and pMR-LacZ empty plasmids, the *E. coli* moss transformed with pUT18C-*STnc1480* and pMR-LacZ-*lpfA* turned blue on the X-gal plate (Fig. 8), indicating that *STnc1480* plays a positively regulatory role on the expression of *lpfA* gene.





**Fig. 5** Histopathological changes in liver, spleen, and intestine of mice inoculated by PBS, SL1344,  $\Delta$ STnc1480, and  $\Delta$ STnc1480/STnc1480 strains, respectively. (magnification,  $\times 200$ , hematoxylin and eosin stain(HE) staining)

## Discussion

Many studies have shown that sRNA may regulate the expression of target genes in a variety of ways at the post-transcriptional levels. A large majority of sRNA regulatory factors block the entry of 30S ribosome by combining mRNA's Shine–Dalgarno (SD) sequences or AUG start codon, thus preventing the start of translation. Alternatively, sRNA can also be paired with the target gene base to protect it from cell RNA degradation, thereby stabilizing the protein expression of the target gene

(Bouvier et al. 2008; Opdyke et al. 2004). In addition, sRNA can be paired with the 5'-non-coding (UTR) regions of mRNA to open the secondary structures of the masking SD sequence and promote the binding of the SD sequence to the 30S ribosome, thus activating the expression of the target genes (Bobrovskyy and Vanderpool 2014).

In the recent years, many investigators have carried out research work on sRNAs of STM. To date, approximately 280 sRNAs expressed under various conditions have been detected in the STM LT2 strain (Kröger et al. 2013). Shabarinath et al. (Srikumar et al. 2015) found a significant increase in the expression level of 31 sRNAs




**Fig. 6** Complementary pairing position of sRNA *STnc1480* with mRNA of *lpfA* gene of STM. **A** The schematic represents the pairing region of *STnc1480* sRNA with *lpfA* gene, **B** The -35, -10

region of *lpfA* gene promoter were underlined. The complementary sequences for *STnc1480* targeting region are shown by black spots

through transcriptome studies of STM-infected mouse macrophages, while expression levels of some sRNAs were very low during extracellular growth, but were

significantly higher in macrophages, suggesting that these sRNAs may be related to STM's pathogenicity and intracellular survival (Sittka et al. 2008).




**Fig. 7** Determination of the transcription levels of *lpfA* gene in  $\Delta$ STnc1480,  $\Delta$ STnc1480/STnc1480, and SL1344 by RT-qPCR. Values represent the mean  $\pm$  SEM (\*\* $P < 0.01$ )

It has been found that expression level of sRNA *STnc1480* was significantly increased after STM-infected cells (Srikumar et al. 2015). However, the regulatory roles of *STnc1480* and its target genes have not yet been studied. Here, we confirmed that the cell invasion ability and pathogenicity of the *STnc1480*-deficient strain were significantly reduced, suggesting that the *STnc1480* was involved in regulation of pathogenicity in STM. Bioinformational predictions revealed that *lpfA* was one of the target genes regulated by *STnc1480*. It has been proved that *lpfA* was the main subunit of *lpf* fimbria, which helps STM form BF in the infected intestinal epithelia and increases ability of STM's intestinal colonization and pathogenicity to mice (Quan et al. 2019; Ledebor et al. 2006). RT-qPCR analysis further confirmed that expression level of *lpfA* in the  $\Delta$ STnc1480 was lower than that of SL1344, indicating that *STnc1480* contributes to the expression regulation of *lpfA* gene, thus resulting in the reduced capacities of BF formation, cell invasion, and pathogenicity of STM.

To confirm the interaction between *STnc1480* and 5'-UTR of *lpfA* mRNA, the bacterial dual-plasmid reporting system was employed in this study. By co-transformed the plasmid expressing sRNA and the plasmid containing their target mRNA-binding sites and the reporting gene, the effects of *STnc1480* and *lpfA* on LacZ gene expression using X-gal as color agent were analyzed. The results revealed that *STnc1480* could bind with the 5'UTR region of *lpfA* mRNA. Therefore,

**Fig. 8** Verification of interaction between *STnc1480* and target gene *lpfA* using two plasmids co-expression system. **A** Bacterial lawn of BTH101 transformed by pUT18C, **B** bacterial lawn of BTH101 transformed by pMR-LacZ, **C** bacterial lawn of BTH101 co-transformed by pUT18C and pMR-LacZ, **D** bacterial lawn of BTH101 co-transformed by pUT18C-*STnc1480* and pMR-LacZ-*lpfA*



we speculated that the molecular mechanism of *STnc1480* is to regulate the expression level of the *lpfA* gene by binding to mRNA of *lpfA* gene, protecting it from RNase degradation, and thus controlling the expression levels of the target genes, which possess similar regulatory mechanisms with the literature (Fröhlich et al. 2013).

In conclusion, this study revealed the regulatory roles of sRNA *STnc1480* on BF formation and pathogenicity in STM, which provided new insights into the regulatory mechanism of sRNAs in STM.

**Supplementary Information** The online version contains supplementary material available at <https://doi.org/10.1007/s00203-022-03331-9>.

**Acknowledgements** The authors thank the field staff for providing the materials for this study.

**Author contributions** LJ and QJ wrote the main manuscript text. LJ, Ning CC, LN, GY, JC, ZX, and ZX prepared figures. MQ, XX, CX, and QJ reviewed and edited. All authors reviewed the manuscript.

**Funding** This work was supported by grant from the national key research and development program (No. 2016YFD0500900), the Grant from Youth Science and Technology Innovation Leader of

Xinjiang Production and Construction Corps (No. 2016BC001), and the Key Scientific and Technological Project in Agriculture of Xinjiang Production and Construction Corps (No. 2019GG026).

## Declarations

**Conflict of interests** This manuscript has not been simultaneously submitted for publication in another journal and been approved by all co-authors. The authors declare that they do not have any conflict of interest.

**Ethical approval** The experiments were carried out in accordance with the guidelines issued by the Ethical Committee of Shihezi University.

## References

- Angalabiri-Owei BE, Isirima JC (2014) Evaluation of the lethal dose of the methanol extract of *Rhizophora racemosa* leaf using Karbers method. *Cell Pathol* 2:65–68. <https://doi.org/10.5897/AJCPA TH14.009>
- Bobrovskyy M, Vanderpool CK (2014) The small RNA SgrS: roles in metabolism and pathogenesis of enteric bacteria. *Front Cell Infect Microbiol* 4:61–68. <https://doi.org/10.3389/fcimb.2014.00061>

- Bouvier M, Sharma CM, Mika F, Nierhaus KH, Vogel J (2008) Small RNA binding to 5' mRNA coding region inhibits translational initiation. *Mol Cell* 32:827–837. <https://doi.org/10.1016/j.molcel.2008.10.027>
- Datsenko KA, Wanner BL (2000) One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc Natl Acad Sci USA* 97:6640–6645. <https://doi.org/10.1073/pnas.120163297>
- Fàbrega A, Vila J (2013) *Salmonella enterica* serovar *Typhimurium* skills to succeed in the host: virulence and regulation. *Clin Microbiol Rev* 26:308–341. <https://doi.org/10.1128/CMR.00066-12>
- Fröhlich KS, Vogel J (2009) Activation of gene expression by small RNA. *Curr Opin Microbiol* 12:674–682. <https://doi.org/10.1016/j.mib.2009.09.009>
- Fröhlich KS, Papenfort K, Fekete A, Vogel J (2013) A small RNA activates CFA synthase by isoform-specific mRNA stabilization. *EMBO J* 32:2963–2979. <https://doi.org/10.1038/emboj.2013.222>
- Harfouche L, Haichar FZ, Achouak W (2015) Small regulatory RNAs and the fine-tuning of plant-bacteria interactions. *New Phytol* 206:98–106. <https://doi.org/10.1111/nph.13195>
- Hölzer SU, Schlumberger MC, Jäckel D, Hensel M (2009) Effect of the O-antigen length of lipopolysaccharide on the functions of type III secretion systems in *Salmonella enterica*. *Infect Immun* 77:5458–5470. <https://doi.org/10.1128/IAI.00871-09>
- Kery MB, Feldman M, Livny J, Tjaden B (2014) TargetRNA2: identifying targets of small regulatory RNAs in bacteria. *Nucleic Acids Res* 42:124–129. <https://doi.org/10.1093/nar/gku317>
- Kint G, De Coster D, Marchal K, Vanderleyden J, De Keersmaecker SC (2010) The small regulatory RNA molecule *MicA* is involved in *Salmonella enterica* serovar *Typhimurium* biofilm formation. *BMC Microbiol* 10:276–284. <https://doi.org/10.1186/1471-2180-10-276>
- Kröger C, Dillon SC, Cameron AD, Papenfort K, Sivasankaran SK, Hokamp K, Chao Y, Sittka A, Hébrard M, Händler K, Colgan A, Leekitcharoenphon P, Langridge GC, Lohan AJ, Loftus B, Lucchini S, Ussery DW, Dorman CJ, Thomson NR, Vogel J, Hinton JC (2012) The transcriptional landscape and small RNAs of *Salmonella enterica* serovar *Typhimurium*. *Proc Natl Acad Sci USA* 109:E1277–1286. <https://doi.org/10.1073/pnas.1201061109>
- Kröger C, Colgan A, Srikumar S, Händler K, Sivasankaran SK, Hammarlöf DL, Canals R, Grissom JE, Conway T, Hokamp K, Hinton JC (2013) An infection-relevant transcriptomic compendium for *Salmonella enterica* Serovar *typhimurium*. *Cell Host Microbe* 14:683–695. <https://doi.org/10.1016/j.chom.2013.11.010>
- Kumawat M, Pesingi PK, Agarwal RK, Goswami TK, Mahawar M (2016) Contribution of protein isoaspartate methyl transferase (PIMT) in the survival of *Salmonella typhimurium* under oxidative stress and virulence. *Int J Med Microbiol* 306:222–230. <https://doi.org/10.1016/j.ijmm.2016.04.005>
- Ledeboer NA, Frye JG, McClelland M, Jones BD (2006) *Salmonella enterica* serovar *Typhimurium* requires the *Lpf*, *Pef*, and *Tafi* fimbriae for biofilm formation on HEp-2 tissue culture cells and chicken intestinal epithelium. *Infect Immun* 74:3156–3169. <https://doi.org/10.1128/IAI.01428-05>
- Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* 25:402–408. <https://doi.org/10.1006/meth.2001.1262>
- Majowicz SE, Musto J, Scallan E, Angulo FJ, Kirk M, O'Brien SJ, Jones TF, Fazil A, Hoekstra RM (2010) The global burden of nontyphoidal *Salmonella* gastroenteritis. *Clin Infect Dis* 50:882–889. <https://doi.org/10.1086/650733>
- Opdyke JA, Kang JG, Storz G (2004) GadY, a small-RNA regulator of acid response genes in *Escherichia coli*. *J Bacteriol* 186:6698–6705. <https://doi.org/10.1128/JB.186.20.6698-6705.2004>
- Papenfort K, Vogel J (2009) Multiple target regulation by small noncoding RNAs rewires gene expression at the post-transcriptional level. *Res Microbiol* 160:278–287. <https://doi.org/10.1016/j.resmic.2009.03.004>
- Quan X, Xia P, Zhao J, Zhu C, Meng X, Yang Y, Wang Y, Tian Y, Ding X, Zhu G (2019) Fimbriae and related receptors for *Salmonella enteritidis*. *Microb Pathog* 126:357–362. <https://doi.org/10.1016/j.micpath.2018.10.025>
- Ryan D, Mukherjee M, Nayak R, Dutta R, Suar M (2018) Biological and regulatory roles of acid-induced small RNA RyeC in *Salmonella typhimurium*. *Biochimie* 150:48–56. <https://doi.org/10.1016/j.biochi.2018.05.001>
- Senior K (2009) Estimating the global burden of foodborne disease. *Lancet Infect Dis* 9:80–81. [https://doi.org/10.1016/s1473-3099\(09\)70008-8](https://doi.org/10.1016/s1473-3099(09)70008-8)
- Sittka A, Lucchini S, Papenfort K, Sharma CM, Rolle K, Binnewies TT, Hinton JC, Vogel J (2008) Deep sequencing analysis of small noncoding RNA and mRNA targets of the global post-transcriptional regulator. Hfq *Plos Genet* 4:e1000163. <https://doi.org/10.1371/journal.pgen.1000163>
- Srikumar S, Kröger C, Hébrard M, Colgan A, Owen SV, Sivasankaran SK, Cameron AD, Hokamp K, Hinton JC (2015) RNA-seq brings new insights to the intra-macrophage transcriptome of *Salmonella typhimurium*. *PLoS Pathog* 11:e1005262. <https://doi.org/10.1371/journal.ppat.1005262>
- Wagner EGH, Romby P (2015) Small RNAs in bacteria and archaea: who they are, what they do, and how they do it. *Adv Genet* 90:133–208. <https://doi.org/10.1016/bs.adgen.2015.05.001>

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.