

L^1 Convergence to the Barenblatt Solution for Compressible Euler Equations with Damping

FEIMIN HUANG, RONGHUA PAN & ZHEN WANG

Communicated by A. BRESSAN

Abstract

We study the asymptotic behavior of compressible isentropic flow through a porous medium when the initial mass is finite. The model system is the compressible Euler equation with frictional damping. As $t \rightarrow \infty$, the density is conjectured to obey the well-known porous medium equation and the momentum is expected to be formulated by Darcy's law. In this paper, we prove that any L^∞ weak entropy solution to the Cauchy problem of damped Euler equations with finite initial mass converges strongly in the natural L^1 topology with decay rates to the Barenblatt profile of the porous medium equation. The density function tends to the Barenblatt solution of the porous medium equation while the momentum is described by Darcy's law. The results are achieved through a comprehensive entropy analysis, capturing the dissipative character of the problem.

1. Introduction

This paper is the continuation of the program toward the mathematical justification of Darcy's law as a long time asymptotic limit for compressible isentropic porous medium flow, modeled by the following Cauchy problem of the compressible Euler equation with frictional damping,

$$\begin{cases} \rho_t + (\rho u)_x = 0, \\ (\rho u)_t + (\rho u^2 + p(\rho))_x = -\alpha \rho u, \\ \rho(x, 0) = \rho_0(x), \quad u(x, 0) = u_0(x). \end{cases} \quad (1.1)$$

Here ρ , u and $p = \kappa \rho^\gamma$, $\kappa = \frac{(\gamma-1)^2}{4\gamma}$, ($1 < \gamma < 3$) denote density, velocity, momentum and pressure, respectively. $\alpha > 0$ is a given positive constant modeling frictional force induced by the medium. We also use momentum $m =: \rho u$ in what follows for convenience. For simplicity, we assume $\alpha = \kappa = \frac{(\gamma-1)^2}{4\gamma}$. Such choice

of constants κ and α is purely for convenience, which simplifies the form of the entropy functions we have employed below.

Because of the dissipative nature of frictional force, it is natural to expect the inertial terms in the momentum equation to decay to zero faster than other terms, so that the pressure gradient force is balanced by the frictional force, as stated by Darcy’s law. In other words, as $t \rightarrow \infty$, the density is conjectured to obey the well-known porous medium equation (PME) and the momentum is expected to be formulated by Darcy’s law, as observed in experiments. Therefore, taking time asymptotically, the system (1.1) is conjectured to be equivalent to the following decoupled system

$$\begin{cases} \bar{\rho}_t = (\bar{\rho}^\gamma)_{xx}, & \text{Porous Medium Equation} \\ \bar{m} = -(\bar{\rho}^\gamma)_x, & \text{Darcy’s law.} \end{cases} \tag{1.2}$$

The particular emphasis here is when the initial total mass is finite, allowing vacuum states in the solutions initially, namely,

$$\int_{-\infty}^{+\infty} \rho(x, t) \, dx = \int_{-\infty}^{+\infty} \bar{\rho}(x, t) \, dx = \int_{-\infty}^{+\infty} \rho_0(x) \, dx = M < \infty,$$

in view of mass conservation. We shall prove in this paper the L^1 convergence from any L^∞ entropy weak solution of the compressible Euler equations to the Barenblatt solution [2] of the corresponding porous medium equation carrying finite total mass. The first evidence toward this expectation was hinted at in the inspiring paper of LIU [26] through an interesting explicit solution which behaves as the Barenblatt solution of the porous medium equation. Recent evidence was provided by HUANG ET AL. [21] for L^∞ entropy weak solutions for $\gamma \in \left(\frac{1+\sqrt{5}}{2}, \sqrt{2} + 1\right)$ measured in the energy norm of the porous medium equation. The purpose of the current paper is to finally provide a complete treatment of the case when $\gamma \in (1, 3)$, with measurements in both L^1 (the mass norm) and $L^{\gamma+1}$ (the energy norm) and thus in L^p for $1 \leq p \leq \gamma + 1$.

Mathematical study of system (1.1) dates back to the 1970s. Following the pioneering work of NISHIDA [33], many contributions have been made to solving this problem. In the case away from vacuum, system (1.1) can be transferred to the damped p -system by changing to the Lagrangian coordinates; see [32,43]. Frictional damping prevents the breaking of waves with small amplitude, leading to the global existence of smooth solutions when the initial data are small and smooth [33]. However, waves break down in finite time when the initial derivatives of initial data exceed a certain threshold [45]. The global existence of weak solutions in L^p was established by the method of compensated compactness in [10,11,25,46]. The global BV solutions were proved in [7,9,27]. The conjecture mentioned above was first justified by HSIAO and LIU in [13,14], and further improved by many mathematicians for small smooth or piecewise smooth solutions away from vacuum based on the energy estimates for derivatives; see [12,15–18,30,34–36,44,47]. Recently, DAFERMOS and PAN [9] constructed global BV solutions to the damped p -system and proved the conjecture with sharp decay rates in L^2 . In these results, the solutions of the damped p -system were shown to converge to the self-similar solutions of the

corresponding porous medium equations constructed in [40], since the end-states of the initial density are away from vacuum.

When a vacuum occurs in the solution, the difficulty of the problem greatly increases, mainly due to the interaction of nonlinear convection, lower order dissipation of damping and the resonance due to the vacuum. It is known that non-linearity is the reason for shock formation in a hyperbolic system. For hyperbolic conservation laws, self-similarity is an important feature in constructing fundamental Riemann solutions and in describing the large time behaviors of solutions. Although damping provides weak dissipation, it breaks the self-similarity of the system, which is crucial for the large weak solutions. Another difficulty is due to the resonance near the vacuum which develops a new singularity; see [28, 29]. Due to this new singularity, it is very difficult to obtain the solutions with any degree of regularity. This makes (1.1) difficult to understand analytically and makes the construction of effective numerical methods for computing solutions a highly non-trivial problem. Indeed, the only global weak solution with vacuum is constructed in L^∞ space by using the method of compensated compactness; see DING ET AL. [10] for $1 < \gamma \leq \frac{5}{3}$ and HUANG AND PAN [19] for $1 \leq \gamma < 3$. Thus, to study the large time behavior of solutions of (1.1) with vacuum, it is suitable to consider the L^∞ weak solution.

Definition 1.1. For any $T > 0$, the bounded measurable functions $(\rho, m)(x, t) \in L^\infty(\mathbf{R} \times [0, T])$ are called entropy solutions of (1.1), if

$$\begin{cases} \rho_t + m_x = 0, \\ m_t + \left(\frac{m^2}{\rho} + \kappa\rho^\gamma\right)_x + \alpha m = 0, \\ \eta_t + q_x + \alpha\eta_m m \leq 0, \end{cases} \tag{1.3}$$

hold in the sense of distributions, where (η, q) is any weak convex entropy–flux pair $(\eta(\rho, m), q(\rho, m))$ satisfying

$$\nabla q = \nabla \eta \nabla f, \quad f = \left(m, \frac{m^2}{\rho} + \kappa\rho^\gamma\right)^t, \quad \eta(0, 0) = 0. \tag{1.4}$$

As the L^∞ weak solution does not have any degree of regularity, the methods for the case away from vacuum are not applicable here. Earlier attempts were made by HUANG and PAN [19], where the authors followed the rescaling argument due to SERRE and HSIAO [42] and obtained the first justification for the conjecture regarding the vacuum case. It showed that the density in the L^∞ weak entropy solutions of (1.1) converges to the similarity solution of the porous medium equation along the level curve of the diffusive similarity profiles, provided that one of the initial end-states is nonzero. The long time behavior of the momentum is not known, however. This is far from satisfactory. In [20], HUANG and PAN developed a new technique based on the conservation of mass and entropy analysis to attack this conjecture. They showed that the L^∞ weak entropy solutions with vacuum, selected by the physical entropy-flux pairs, converge strongly in $L^p(\mathbf{R})$ ($p \geq p_0$ for some $p_0 \geq 2$) with decay rates to the similarity solution of the porous medium equation determined uniquely by the end-states and the mass distribution of the

initial data, provided that the end-states are away from vacuum. This approach seems remarkable, since it does not need smallness assumptions on the solutions. Inspired by this result, HUANG ET AL. [21] further studied this problem with finite total mass, where the asymptotic profile is the celebrated Barenblatt solution of the porous medium equation. More precisely, this result is cited below:

Theorem. (HUANG ET AL., [21]) *Suppose $0 \leq \rho_0(x) \in L^1(\mathbf{R}) \cap L^\infty(\mathbf{R})$ and*

$$0 < M = \int_{-\infty}^{+\infty} \rho_0(x) \, dx < \infty.$$

Let (ρ, m) be an L^∞ entropy solution of the Cauchy problem (1.1), satisfying the following estimates

$$0 \leq \rho(x, t) \leq C, \quad |m(x, t)| \leq C\rho(x, t), \tag{1.5}$$

where the constant C is independent of t . Let $\bar{\rho}$ be the Barenblatt solution of the porous medium equation (1.2) with mass M and $\bar{m} = -(\bar{\rho}^\gamma)_x$ which satisfies

$$\begin{aligned} \|\bar{\rho}\|_{L^2}^2 &\leq C(1+t)^{-\frac{1}{\gamma+1}}, \\ \|\bar{\rho}\|_{L^\gamma}^\gamma &\leq C(1+t)^{-\frac{\gamma-1}{\gamma+1}}. \end{aligned} \tag{1.6}$$

Define

$$y = - \int_{-\infty}^x (\rho - \bar{\rho})(r, t) \, dr.$$

If $y(x, 0) \in L^2(\mathbf{R})$, then there exist positive constants $k_1 = \min \left\{ \frac{\gamma^2}{(\gamma+1)^2}, \frac{\gamma-1}{\gamma} \right\}$, $k_2 = \min \left\{ \frac{\gamma^2}{(\gamma+1)^2}, \frac{1}{\gamma} \right\}$ and C such that for any $\varepsilon > 0$,

$$\begin{aligned} \|(\rho - \bar{\rho})(x, t)\|_{L^2}^2 &\leq C(1+t)^{-k_1+\varepsilon}, \quad \text{if } 1 < \gamma \leq 2, \\ \|(\rho - \bar{\rho})(x, t)\|_{L^\gamma}^\gamma &\leq C(1+t)^{-k_2+\varepsilon}, \quad \text{if } \gamma \geq 2. \end{aligned} \tag{1.7}$$

Furthermore,

$$\begin{aligned} k_1 &> \frac{1}{\gamma+1}, \quad \text{if } \frac{1+\sqrt{5}}{2} < \gamma \leq 2, \\ k_2 &> \frac{\gamma-1}{\gamma+1}, \quad \text{if } 2 \leq \gamma < 1+\sqrt{2}. \end{aligned} \tag{1.8}$$

Remark 1.1. Since Barenblatt’s solution $\bar{\rho}$ decays itself, it is necessary to compare the decay rate of $\bar{\rho}$ with that of $\rho - \bar{\rho}$. This theorem shows that $\|\rho - \bar{\rho}\|_{L^2}$ decays faster than $\|\bar{\rho}\|_{L^2}$ when $\frac{1+\sqrt{5}}{2} < \gamma \leq 2$ and $\|\rho - \bar{\rho}\|_{L^\gamma}$ decays faster than $\|\bar{\rho}\|_{L^\gamma}$ when $2 \leq \gamma < 1 + \sqrt{2}$. Thus this theorem states that any L^∞ entropy weak solutions must converge to the Barenblatt solution of the PME with the same mass when $\gamma \in \left(\frac{1+\sqrt{5}}{2}, 1 + \sqrt{2} \right)$.

Remark 1.2. The entropy dissipation method introduced in [20,21] is an effective approach in proving the large time asymptotic behavior for L^∞ weak entropy solutions for hyperbolic conservation laws with dissipation. Further application of the approach can be found in [39] for an initial boundary value problem and in [22] for the Euler-Poisson system modeling semi-conductor devices.

Although this is the first result providing the convergence of L^∞ weak entropy solutions of (1.1) to the Barenblatt solution of (1.2), the result itself is not definitive for several reasons. One may thus ask the following questions:

- **Question 1:** Is it possible to remove assumption (1.5)? Assumption (1.5) is essential in the proof of [21]. This assumption is quite reasonable, since the solutions obtained in [10] and [19] satisfy this condition where invariant region theory is applied to the viscosity regularized system. On the other hand, there is no uniqueness theory available for the L^∞ weak entropy solutions of (1.1), it is not clear whether there are any uniformly bounded entropy weak solutions of (1.1) that do not satisfy (1.5) with uniform constant C . Therefore, it is natural to ask this question.
- **Question 2:** Is it possible to prove convergence for any $\gamma \in (1, 3)$, not only on the interval $\left(\frac{1+\sqrt{5}}{2}, 1 + \sqrt{2}\right)$, as in the last Theorem? Although the interval $\left(\frac{1+\sqrt{5}}{2}, 1 + \sqrt{2}\right)$ contains some physical cases, most physical gases live in the larger interval $(1, 3)$. This generalization is thus important from the physical point of view.
- **Question 3:** Is it possible to prove decay of $\|\rho - \bar{\rho}\|_{L^1}$? Since the compressible Euler equation is conserved, it is natural to measure the difference between ρ and the Barenblatt solution $\bar{\rho}$ in L^1 space. Furthermore, the L^1 norm does not decay for either ρ or $\bar{\rho}$, the L^1 decay is very convincing. Therefore, the last question is whether it is possible to obtain an L^1 convergence result.

In this paper, we will address the three questions listed above and give definite answers to them. After a quick review of some information on Barenblatt’s solution in Section 2, we will first prove in Section 3 the following invariant region theory for L^∞ weak entropy solution to (1.1).

Theorem 1.1. *Suppose that $(\rho_0, u_0)(x) \in L^\infty(\mathbf{R})$ satisfies*

$$0 \leq \rho_0(x) \leq C, \quad |m_0(x)| \leq C\rho_0(x).$$

Let $(\rho, u) \in L^\infty(R \times [0, T])$ be an L^∞ weak entropy solution of the system (1.1) with $\gamma > 1$. Then (ρ, m) satisfies

$$0 \leq \rho(x, t) \leq C, \quad |m(x, t)| \leq C\rho(x, t), \tag{1.9}$$

where the constant C depends solely on the initial data.

Remark 1.3. Theorem 1.1 is valid for L^∞ weak entropy solutions to the homogeneous compressible Euler systems; this is quite clear from the proof in Section 3 below. This theorem is an invariant region theorem for the weak solutions to compressible Euler equations with damping. In the invariant region theorem [6], the

derivative property of a solution is essential. But here it is impossible to use any derivative properties due to the lack of regularity. Theorem 1.1 is proved by choosing infinitely many convex weak entropies. This idea dates back to DA FERMOs [8] and SERRE [41]. A similar idea appears in [24, 38] for homogeneous two by two systems, including compressible Euler equations and elastodynamics. The distinct feature in our proof for this theorem is the choice of entropies designed for the dissipative source term.

In Section 4, we will pursue the sharper decay rates of the density to the Barenblatt solution, which will finally give the decay rates in the L^1 space. These are stated in the following Theorem.

Theorem 1.2. *Suppose $\rho_0(x) \in L^1(\mathbf{R}) \cap L^\infty(\mathbf{R})$, $u_0(x) \in L^\infty(\mathbf{R})$ and*

$$M = \int_{-\infty}^{\infty} \rho_0(x) \, dx > 0.$$

Let $1 < \gamma < 3$ and (ρ, m) be an L^∞ entropy solution of the Cauchy problem (1.1). Let $\bar{\rho}$ be the Barenblatt solution of porous medium equation (1.2) with mass M and $\bar{m} = -(\bar{\rho}^\gamma)_x$. Define

$$y = - \int_{-\infty}^x (\rho - \bar{\rho})(r, t) \, dr.$$

If $y(x, 0) \in L^2(\mathbf{R})$, then for any $\varepsilon > 0$ and $t > 0$,

$$\begin{aligned} \|(\rho - \bar{\rho})(\cdot, t)\|_{L^{\gamma+1}}^{\gamma+1} &\leq C(1+t)^{-1+\frac{1}{2(\gamma+1)}+\varepsilon}, \\ \|(\rho - \bar{\rho})(\cdot, t)\|_{L^1} &\leq C(1+t)^{-\frac{1}{4(\gamma+1)}+\varepsilon}. \end{aligned} \tag{1.10}$$

Remark 1.4. Due to the mass conservation law, it is natural to use L^1 norm to measure the difference between the entropy solution ρ and Barenblatt’s solution $\bar{\rho}$. (1.10) implies that any L^∞ entropy solutions must converge to the Barenblatt solution of PME’s with the same mass for $1 < \gamma < 3$. This is because the L^1 norm is conserved for both ρ and $\bar{\rho}$. Furthermore, we remark that the $L^{\gamma+1}$ norm of $\rho - \bar{\rho}$ decays faster than $\bar{\rho}$, since for the latter one has, from Lemma 2.2 below, that

$$\|\bar{\rho}(\cdot, t)\|_{L^{\gamma+1}}^{\gamma+1} \leq C(1+t)^{-\frac{\gamma}{\gamma+1}}$$

which decays slower than the rate in (1.10) by roughly $(1+t)^{\frac{1}{2(\gamma+1)}}$.

We now make some comments on the new ideas and approaches in this paper, based on intensive entropy analysis. In the proof of Theorem 1.1, infinitely many convex entropy functions are used. In the proof of Theorem 1.2, we first follow the method of [21] to pave the road. With the help of a sharp estimate on the degeneracy of $p(\rho)$ at vacuum proved in Lemma 3.1, a carefully chosen entropy is applied to obtain a much sharper decay in $L^{\gamma+1}$ norm of the density, which is missing in [21]. This much faster decay rate guaranteed that $\|\rho - \bar{\rho}\|_{L^{\gamma+1}}$ decays faster than $\|\bar{\rho}\|_{L^{\gamma+1}}$ for any $\gamma \in (1, 3)$. Finally, this refined estimate, together with a key observation

on the distribution of $\|\rho - \bar{\rho}\|_{L^1}$ over the support of $\bar{\rho}$, leads to the L^1 decay rates in Theorem 1.2.

In the last section, we further discuss a Barenblatt type solution of (1.1) with particular initial data, constructed by LIU in [26]. This interesting solution, which we call Liu’s solution, behaves just like Barenblatt’s solution and thus could serve as the large time asymptotic ansatz for solutions of (1.1) with finite total mass. Theorem 1.2, together with the results in [26], implies the L^1 decay to Liu’s solution as well. The reason for the choice of Barenblatt’s solution is simply because of its explicit form which is not available for Liu’s solution. Furthermore, the decay rates in Theorem 1.2 might not be optimal, some explanations will be presented there at the end of the paper.

2. Barenblatt’s solutions

According to [21], the solutions to (1.1) with finite total mass should converge in large time to the fundamental solutions of the porous media equation, that is, the Barenblatt solutions [2]. In this section, we provide some background information on the Barenblatt solutions.

Consider

$$\begin{cases} \bar{\rho}_t = (\bar{\rho}^\gamma)_{xx}, \\ \bar{\rho}(-1, x) = M\delta(x), \quad M > 0, \end{cases} \tag{2.1}$$

which admits a unique solution (see also [1,2]) given below

$$\bar{\rho}(x, t) = (t + 1)^{-\frac{1}{\gamma+1}} \{(A - B\xi^2)_+\}^{\frac{1}{\gamma-1}}. \tag{2.2}$$

Here $\xi = x(t + 1)^{-\frac{1}{\gamma+1}}$, $(f)_+ = \max\{0, f\}$, $B = \frac{\gamma-1}{2\gamma(\gamma+1)}$ and A is determined by

$$2A^{\frac{\gamma+1}{2(\gamma-1)}} B^{-\frac{1}{2}} \int_0^{\frac{\pi}{2}} (\cos \theta)^{\frac{\gamma+1}{\gamma-1}} d\theta = M. \tag{2.3}$$

Due to the degeneracy at vacuum, the derivative of $\bar{\rho}$ is not continuous across the interface between the gas and vacuum. Instead, $\bar{\rho}$ is a weak solution to (2.1) such that

$$\int_{-\infty}^{+\infty} \bar{\rho} dx = M, \tag{2.4}$$

and

$$\bar{\rho} = 0, \quad \text{if } |\xi| \geq \sqrt{\frac{A}{B}}. \tag{2.5}$$

Hence, for any finite time $T > 0$, $\bar{\rho}$ has compact support. This is the property of the finite speed of propagation for the porous medium equation. For the definition of the weak solution to (2.1), we refer to [1–3,23,31]. KAMIN proved in [23] that (2.1) admits at most one solution. Here, we addressed the initial data at $t = -1$ to avoid the singularity at $t = 0$. Thus, we have the following lemmas from (2.2)–(2.5).

Lemma 2.1. *If $M > 0$ is finite, then there is one and only one solution $\bar{\rho}(x, t)$ to (2.1). Furthermore, the following holds.*

- $\bar{\rho}(x, t)$ is continuous on \mathbf{R} .
- There is a number $b = \sqrt{\frac{A}{B}} > 0$, such that $\bar{\rho}(x, t) > 0$ if $|x| < bt^{\frac{1}{\gamma+1}}$; and $\bar{\rho}(x, t) = 0$ if $|x| \geq bt^{\frac{1}{\gamma+1}}$.
- $\bar{\rho}(x, t)$ is smooth if $|x| < bt^{\frac{1}{\gamma+1}}$.

In terms of the explicit form of $\bar{\rho}$, it is easy to check the following estimates.

Lemma 2.2. *For $\bar{\rho}$ defined in (2.2) and $t > 0$, it holds that*

$$\begin{cases} |\bar{\rho}| \leq C(1+t)^{-\frac{1}{\gamma+1}}, \\ |(\bar{\rho}^{\gamma-1})_x| \leq C(1+t)^{-\frac{\gamma}{\gamma+1}}, \quad |(\bar{\rho}^{\gamma-1})_t| \leq C(1+t)^{-\frac{2\gamma}{\gamma+1}}, \\ |(\bar{\rho}^\gamma)_x| \leq C(1+t)^{-1}, \quad |(\bar{\rho}^\gamma)_t| \leq C(1+t)^{-\frac{2\gamma+1}{\gamma+1}}, \end{cases} \quad (2.6)$$

and

$$\begin{cases} \int_{-\infty}^{+\infty} \bar{\rho}^p \, dx \leq C(1+t)^{-\frac{p-1}{\gamma+1}}, \quad \forall p \geq 1, \\ \int_{-\infty}^{+\infty} (\bar{\rho}^{\gamma-1})_x^2 \, dx \leq C(1+t)^{-\frac{2\gamma-1}{\gamma+1}}, \quad \int_{-\infty}^{+\infty} (\bar{\rho}^{\gamma-1})_t^2 \, dx \leq C(1+t)^{-\frac{4\gamma-1}{\gamma+1}}, \\ \int_{-\infty}^{+\infty} (\bar{\rho}^\gamma)_x^2 \, dx \leq C(1+t)^{-\frac{2\gamma+1}{\gamma+1}}, \quad \int_{-\infty}^{+\infty} (\bar{\rho}^\gamma)_t^2 \, dx \leq C(1+t)^{-\frac{4\gamma+1}{\gamma+1}}. \end{cases} \quad (2.7)$$

3. Invariant region for weak solutions

In this section, we shall show a proof to Theorem 1.1, which serves as an invariant region theory [6] for L^∞ weak entropy solutions to (1.1). It confirms that any L^∞ weak entropy solutions to (1.1) will stay inside the physical region

$$0 \leq \rho(x, t) \leq C, \quad |m(x, t)| \leq C\rho(x, t),$$

if the initial data do so.

To proceed, we first recall some results on the entropies available for (1.1). According to [24], all weak entropies of (1.1) are given by the following formula:

$$\begin{aligned} \eta(\rho, u) &= \int g(\xi)\chi(\xi; \rho, u) \, d\xi = \rho \int_{-1}^1 g(u + z\rho^\theta)(1 - z^2)^\lambda \, dz, \\ q(\rho, m) &= \int g(\xi)(\theta\xi + (1 - \theta)u)\chi(\xi; \rho, u) \, d\xi \\ &= \rho \int_{-1}^1 g(u + z\rho^\theta)(u + \theta z\rho^\theta)(1 - z^2)^\lambda \, dz \end{aligned} \quad (3.1)$$

where $\theta = \frac{\gamma-1}{2}$, $\lambda = \frac{3-\gamma}{2(\gamma-1)}$, $g(\xi)$ is any smooth function of ξ , and

$$\chi(\xi; \rho, u) = (\rho^{\gamma-1} - (\xi - u)^2)_+^\lambda. \tag{3.2}$$

This remarkable formula can be derived from the entropy equation (1.4) utilizing the kinetic formulation or by fundamental solution of the linear wave equation. We remark that when $g(\xi) = 1$, $\eta(\rho, m) = \rho$; when $g(\xi) = \xi$, $\eta(\rho, m) = m$; and when $g(\xi) = \frac{1}{2}\xi^2$, then $\eta = \frac{m^2}{2\rho} + \frac{\kappa}{\gamma-1}\rho^\gamma$ is mechanical energy.

As the convexity of the entropy function is crucial in the definition of admissible weak solutions, the characterization of convexity of entropy functions is important. In our case, the following lemma provides full details in this direction.

Lemma 3.1 (LIONS ET AL., [24]). *Weak entropy $\eta(\rho, m)$ defined in (3.1) is convex with respect to ρ and m if and only if $g(\xi)$ is a convex function.*

We now present the proof for Theorem 1.1.

3.1. Proof of Theorem 1.1

Choosing $g(\xi) = g_k(\xi) = e^{k\xi^2}$ in (3.1), for positive parameter $k > 0$, the corresponding entropy $\eta_k \geq 0$ is clearly convex. By the definition, if (ρ, m) is an L^∞ entropy weak solution, we have the following entropy inequality

$$\eta_{k,t} + q_{k,x} + \kappa\eta_{k,m}m \leq 0, \tag{3.3}$$

in the sense of distributions. Assuming temporarily that $\eta_{k,m}m \geq 0$, we define

$$M(T) = \|u\|_{L^\infty(\mathbb{R} \times [0, T])} + \theta \|\rho^\theta\|_{L^\infty(\mathbb{R} \times [0, T])},$$

which measures the largest possible amplitude of the characteristic speed of system (1.1) up to $T > 0$. It follows, by the L^∞ divergence-measure field theory [5], that for any $a > 0$ and $T > 0$,

$$\int_{-a}^a \eta_k(\rho, m)(x, T)dx \leq \int_{-a-M(T)T}^{a+M(T)T} \eta_k(\rho, m)(x, 0)dx, \tag{3.4}$$

In view of the positivity of η_k , one has

$$\left(\int_{-a}^a \eta_k(\rho, m)(x, T)dx \right)^{\frac{1}{k}} \leq \left(\int_{-a-M(T)T}^{a+M(T)T} \eta_k(\rho, m)(x, 0)dx \right)^{\frac{1}{k}}. \tag{3.5}$$

Noting that

$$\eta_k = \rho \int_{-1}^1 e^{k(u+z\rho^\theta)^2} (1 - z^2)^\lambda dz,$$

(3.5) implies, for $Q_a(T) = \{(x, t, z) : -a \leq x \leq a, t = T, -1 \leq z \leq 1\}$, that

$$\begin{aligned} \|\rho^{\frac{1}{k}} e^{(u+z\rho^\theta)^2} (1-z^2)^{\frac{\lambda}{k}}\|_{L^k(Q_a(T))} &\leq \left(\int_{-a}^a \eta_k(\rho, m)(x, T) dx \right)^{\frac{1}{k}} \\ &\leq \left(\int_{-a-M(T)T}^{a+M(T)T} \eta_k(\rho, m)(x, 0) dx \right)^{\frac{1}{k}}. \end{aligned} \tag{3.6}$$

Letting $k \rightarrow +\infty$ and then $a \rightarrow +\infty$ in (3.6), we have

$$\begin{aligned} &\max\{\|(u - \rho^\theta)(x, T)\|_{L^\infty}, \|(u + \rho^\theta)(x, T)\|_{L^\infty}\} \\ &\leq \max\{\|(u - \rho^\theta)(x, 0)\|_{L^\infty}, \|(u + \rho^\theta)(x, 0)\|_{L^\infty}\}, \end{aligned} \tag{3.7}$$

which implies, for any $T > 0$, that

$$0 \leq \rho(x, T) \leq C, \quad 0 \leq |u(x, T)| \leq C. \tag{3.8}$$

where C only depends on the initial data.

It remains to show that $\eta_{k,m}m \geq 0$. In fact,

$$\begin{aligned} m\eta_{k,m} &= 2km \int_{-1}^1 e^{k(u+z\rho^\theta)^2} (u+z\rho^\theta)(1-z^2)^\lambda dz \\ &= 2km \sum_{n=0}^\infty \int_{-1}^1 \frac{k^n}{n!} (u+z\rho^\theta)^{2n+1} (1-z^2)^\lambda dz \\ &= 2\rho u^2 \sum_{n=0}^\infty \frac{k^{n+1}}{n!} \sum_{j=0}^{2n+1} C_{2n+1}^j \int_{-1}^1 u^{2n-j} (z\rho^\theta)^j (1-z^2)^\lambda dz \\ &= 2\rho u^2 \sum_{n=0}^\infty \frac{k^{n+1}}{n!} \sum_{j=0}^n C_{2n+1}^{2j} \int_{-1}^1 u^{2(n-j)} (z\rho^\theta)^{2j} (1-z^2)^\lambda dz \\ &\geq 0. \end{aligned} \tag{3.9}$$

This concludes the proof of Theorem 1.1.

4. Decay estimates

We now proceed to the decay estimates. To begin, we first prove two important inequalities which give sharp information on the pressure near vacuum, where the theory is most lacking.

Lemma 4.1. *If $0 \leq \rho, \bar{\rho} \leq C$, there are two constants $c_1 > 0$ and $c_2 > 0$ such that*

$$\begin{cases} c_1 (\rho^{\gamma-1} + \bar{\rho}^{\gamma-1}) (\rho - \bar{\rho})^2 \leq \rho^{\gamma+1} - \bar{\rho}^{\gamma+1} - (\gamma + 1)\bar{\rho}^\gamma (\rho - \bar{\rho}) \\ \qquad \qquad \qquad \leq c_2 (\rho^{\gamma-1} + \bar{\rho}^{\gamma-1}) (\rho - \bar{\rho})^2 \\ c_1 (\rho^{\gamma-1} + \bar{\rho}^{\gamma-1}) (\rho - \bar{\rho})^2 \leq (\rho^\gamma - \bar{\rho}^\gamma)(\rho - \bar{\rho}) \\ \qquad \qquad \qquad \leq c_2 (\rho^{\gamma-1} + \bar{\rho}^{\gamma-1})(\rho - \bar{\rho})^2. \end{cases} \tag{4.1}$$

Proof. By Taylor’s theorem, we have

$$\begin{aligned} & \rho^{\gamma+1} - \bar{\rho}^{\gamma+1} - (\gamma + 1)\bar{\rho}^\gamma (\rho - \bar{\rho}) \\ &= \gamma(\gamma + 1)(\rho - \bar{\rho})^2 \left[\int_0^1 (1 - s)((1 - s)\bar{\rho} + s\rho)^{\gamma-1} ds \right], \\ & (\rho^\gamma - \bar{\rho}^\gamma)(\rho - \bar{\rho}) \\ &= (\rho - \bar{\rho})^2 \gamma \left[\int_0^1 ((1 - s)\bar{\rho} + s\rho)^{\gamma-1} ds \right]. \end{aligned} \tag{4.2}$$

Noting that for any $0 < s < 1$,

$$\max\{((1 - s)\bar{\rho})^{\gamma-1}, (s\rho)^{\gamma-1}\} \leq ((1 - s)\bar{\rho} + s\rho)^{\gamma-1} \leq \max\{\bar{\rho}^{\gamma-1}, \rho^{\gamma-1}\}, \tag{4.3}$$

(4.1) follows easily from (4.2) and (4.3).

We are now ready to derive the decay estimates. Suppose that (ρ, m) is a weak entropy solution of (1.1) satisfying conditions in Theorem 1.2, then (ρ, m) satisfies

$$\begin{cases} \rho_t + m_x = 0, \\ m_t + \left(\frac{m^2}{\rho} + \kappa(\rho^\gamma)\right)_x = -\kappa m. \end{cases} \tag{4.4}$$

Let $\bar{\rho}$ be the Barenblatt solution of the porous medium equation carrying the same total mass M as ρ , and $\bar{m} = -(\bar{\rho}^\gamma)_x$, then

$$\begin{cases} w = \rho - \bar{\rho}, \\ z = m - \bar{m}, \end{cases} \tag{4.5}$$

satisfying

$$\begin{cases} w_t + z_x = 0 \\ z_t + \left(\frac{m^2}{\rho}\right)_x + k(\rho^\gamma - \bar{\rho}^\gamma)_x + \kappa z = -\bar{m}_t. \end{cases} \tag{4.6}$$

Setting

$$y = - \int_{-\infty}^x w(r, t) dr, \tag{4.7}$$

we have

$$y_x = -w, \quad z = y_t. \tag{4.8}$$

Thus the equation (4.6) turns into a nonlinear wave equation with source terms, degenerate at vacuum:

$$y_{tt} + \left(\frac{m^2}{\rho}\right)_x + \kappa(\rho^\gamma - \bar{\rho}^\gamma)_x + \kappa y_t = -\bar{m}_t. \tag{4.9}$$

Multiplying y with (4.9), integrating over $[0, t] \times (-\infty, \infty)$, integrating by parts, one has

$$\begin{aligned} & \int_{-\infty}^{+\infty} \left(y_t y + \frac{\kappa}{2} y^2 \right) dx + \int_0^t \int_{-\infty}^{+\infty} \kappa (\rho^\gamma - \bar{\rho}^\gamma) (\rho - \bar{\rho}) dx d\tau \\ & \leq C + \int_0^t \int_{-\infty}^{+\infty} \left(y_t^2 + \frac{m^2}{\rho} y_x \right) dx d\tau + \left| \int_0^t \int_{-\infty}^{+\infty} \bar{m}_t y dx d\tau \right|. \end{aligned} \tag{4.10}$$

Since $\|y_x\|_{L^1} = \|\rho - \bar{\rho}\|_{L^1} \leq 2M$, we have, from (2.7), the following estimate

$$\begin{aligned} \left| \int_0^t \int_{-\infty}^{+\infty} \bar{m}_t y dx d\tau \right| &= \left| \int_0^t \int_{-\infty}^{+\infty} \bar{\rho}_t^\gamma y_x dx d\tau \right| \\ &\leq C \int_0^t (1 + \tau)^{-1 - \frac{\gamma}{\gamma+1}} d\tau \leq C. \end{aligned} \tag{4.11}$$

We thus obtain our first estimate.

Lemma 4.2. *Let the conditions of Theorem 1.2 be satisfied, it holds*

$$\begin{aligned} & \int_{-\infty}^{+\infty} \left(y_t y + \frac{\kappa}{2} y^2 \right) dx + \int_0^t \int_{-\infty}^{+\infty} \kappa (\rho^\gamma - \bar{\rho}^\gamma) (\rho - \bar{\rho}) dx d\tau \\ & \leq C + \int_0^t \int_{-\infty}^{+\infty} y_t^2 dx d\tau + \int_0^t \int_{-\infty}^{+\infty} \frac{m^2}{\rho} y_x dx d\tau. \end{aligned} \tag{4.12}$$

This estimate is very rough in its nature. In order to obtain a definite estimate, a higher order estimate is required. Due to the lack of regularity, we will need to close the estimate in the next run, which is achieved by entropy inequality.

Choosing $g(\xi) = \frac{1}{2} \xi^2$ in (3.1), the entropy η takes the form of mechanical energy

$$\eta_e = \frac{m^2}{2\rho} + \frac{\kappa}{\gamma - 1} \rho^\gamma,$$

and q_e the corresponding flux. We read from the entropy inequality that

$$\eta_{et} + q_{ex} + \kappa \frac{m^2}{\rho} \leq 0, \tag{4.13}$$

which implies with the help of divergence measure field theory of [5] that

$$\int_{-\infty}^{+\infty} \eta_e(x, t) dx + \kappa \int_0^t \int_{-\infty}^{+\infty} \frac{m^2}{\rho} dx d\tau \leq C. \tag{4.14}$$

Since $y_t = m - \bar{m}$, we thus have

$$\begin{aligned} \int_0^t \int_{-\infty}^{+\infty} y_t^2 dx d\tau &\leq 2 \int_0^t \int_{-\infty}^{+\infty} \bar{m}^2 dx d\tau + 2 \int_0^t \int_{-\infty}^{+\infty} m^2 dx d\tau \\ &\leq C \int_0^t (1 + \tau)^{-\frac{2\gamma+1}{\gamma+1}} d\tau + C \int_0^t \int_{-\infty}^{+\infty} \frac{m^2}{\rho} dx d\tau \\ &\leq C. \end{aligned} \tag{4.15}$$

and

$$\left| \int_0^t \int_{-\infty}^{+\infty} \frac{m^2}{\rho} y_x \, dx d\tau \right| \leq C \int_0^t \int_{-\infty}^{+\infty} \frac{m^2}{\rho} \, dx d\tau \leq C. \tag{4.16}$$

Substituting (4.14), (4.15) and (4.16) into (4.12), we arrive at the following uniform estimate.

Lemma 4.3. *Under the conditions of Theorem 1.2, for any $t > 0$, it holds that*

$$\begin{aligned} & \int_{-\infty}^{+\infty} \left(\rho^\gamma + y_t^2 + \frac{m^2}{\rho} + y^2 \right) dx + \int_0^t \int_{-\infty}^{+\infty} \left(\frac{m^2}{\rho} + y_t^2 \right) dx d\tau \\ & + \int_0^t \int_{-\infty}^{+\infty} (\rho^\gamma - \bar{\rho}^\gamma) (\rho - \bar{\rho}) \, dx d\tau \leq C. \end{aligned} \tag{4.17}$$

Remark 4.1. It holds for the Barenblatt solution that

$$\int_0^\infty \int_{-\infty}^\infty \bar{\rho}^{\gamma+1} dx dt = \infty. \tag{4.18}$$

On the other hand, one has

$$(\rho^\gamma - \bar{\rho}^\gamma)(\rho - \bar{\rho}) \geq |\rho - \bar{\rho}|^{\gamma+1}.$$

(4.17) thus hints ρ tends to the Barenblatt solution time asymptotically. In fact, a further study carried in [21] gives a rough decay rate on $\|\rho - \bar{\rho}\|_{L^\gamma}$. We remark that η_e measures ρ^γ while the last term on the left-hand side of (4.17) is in the form of $L^{\gamma+1}$. The mismatch of the exponent leads to essential difficulty for better decay rates.

For a sharper decay rate, we shall use the entropy inequality again, which measures the $L^{\gamma+1}$ norm in density. Choosing $g(\xi) = |\xi|^{\frac{2\gamma}{\gamma-1}}$ in (3.1), the entropy reads as

$$\tilde{\eta} = \rho \int_{-1}^1 |u + z\rho^\theta|^{\frac{2\gamma}{\gamma-1}} (1 - z^2)^\lambda dz. \tag{4.19}$$

We shall see below that such an entropy function measures the $L^{\gamma+1}$ norm of ρ , and thus matches the double integral term in (4.17).

By Taylor’s theorem, we have

$$\begin{aligned} \tilde{\eta} &= \rho \int_{-1}^1 g(u + z\rho^\theta) (1 - z^2)^\lambda dz \\ &= \rho \int_{-1}^1 \left[g(z\rho^\theta) + g'(z\rho^\theta)u + \frac{1}{2}g''(z\rho^\theta)u^2 \right] (1 - z^2)^\lambda dz \\ &\quad + \rho \int_{-1}^1 \left[g(u + z\rho^\theta) - g(z\rho^\theta) - g'(z\rho^\theta)u - \frac{1}{2}g''(z\rho^\theta)u^2 \right] (1 - z^2)^\lambda dz \\ &=: C_1\rho^{\gamma+1} + C_2m^2 + A(\rho, m), \end{aligned} \tag{4.20}$$

where

$$\begin{aligned}
 A(\rho, m) &= \rho \int_{-1}^1 \left[g(u + z\rho^\theta) - g(z\rho^\theta) - g'(z\rho^\theta)u - \frac{1}{2}g''(z\rho^\theta)u^2 \right] (1 - z^2)^\lambda dz \\
 &= \rho u^3 \int_{-1}^1 \int_0^1 \frac{(1 - s)^2}{2} g^{(3)}(su + z\rho^\theta) (1 - z^2)^\lambda ds dz, \tag{4.21}
 \end{aligned}$$

$$\begin{cases} C_1 = \int_{-1}^1 |z|^{\frac{2\gamma}{\gamma-1}} (1 - z^2)^\lambda dz = \frac{1}{2} B\left(\frac{\gamma + 1}{2(\gamma - 1)}, \frac{\gamma + 1}{2(\gamma - 1)}\right), \\ C_2 = \frac{\gamma(\gamma + 1)}{(\gamma - 1)^2} \int_{-1}^1 |z|^{\frac{2\gamma}{\gamma-1}} (1 - z^2)^\lambda dz = 2 \frac{\gamma(\gamma + 1)}{(\gamma - 1)^2} C_1 \end{cases} \tag{4.22}$$

and $B(p, q)$ is a Beta function defined by

$$B(p, q) = \int_0^1 x^{p-1} (1 - x)^{q-1} dx.$$

Here we have used the following fact

$$\int_{-1}^1 g'(z\rho^\theta) u (1 - z^2)^\lambda dz = \frac{2\gamma}{\gamma - 1} \rho^\theta u \int_{-1}^1 |z\rho^\theta|^{\frac{2}{\gamma-1}} z (1 - z^2)^\lambda dz = 0. \tag{4.23}$$

Further advantages of this entropy $\tilde{\eta}$ are summarized in the following lemma.

Lemma 4.4. *For $A(\rho, m)$ defined in (4.21), it holds that*

- 1) $|A| \leq C\rho|u|^3(|u|^{\frac{3-\gamma}{\gamma-1}} + \rho^{1-\theta})$,
- 2) $A(\rho, m) \geq 0$,
- 3) $A_m m \geq 0$.

Proof. From the formula (4.21) of $A(\rho, m)$, it is obvious that

$$|A(\rho, m)| \leq C\rho|u|^3 |g^{(3)}(|u| + \rho^\theta)| \leq C\rho|u|^3 \left(|u|^{\frac{3-\gamma}{\gamma-1}} + \rho^{1-\theta} \right).$$

We now prove 2) and 3). In terms of (4.21) and the fact

$$\int_{-1}^1 g^{(3)}(z\rho^\theta) (1 - z^2)^\lambda dz = 0,$$

we get

$$\begin{aligned}
 A &= \rho u^3 \int_{-1}^1 \int_0^1 \frac{(1 - s)^2}{2} \left[g^{(3)}(su + z\rho^\theta) - g^{(3)}(z\rho^\theta) \right] (1 - z^2)^\lambda ds dz \\
 &= \rho u^4 \int_{-1}^1 \int_0^1 \int_0^1 \frac{(1 - s_1)^2 s_2}{2} g^{(4)}(s_1 s_2 u + z\rho^\theta) (1 - z^2)^\lambda ds_1 ds_2 dz \geq 0 \tag{4.24}
 \end{aligned}$$

and

$$A_m m = 3A + \rho u^4 \int_{-1}^1 \int_0^1 \frac{s(1-s)^2}{2} g^{(4)}(su + z\rho^\theta) (1-z^2)^\lambda ds dz \geq 0 \tag{4.25}$$

where we have used $g^{(4)} \geq 0$. Thus Lemma 4.4 is proved.

Let \tilde{q} be the flux corresponding to entropy $\tilde{\eta}$. Define

$$\eta_* = \tilde{\eta} - C_1 \bar{\rho}^{\gamma+1} - C_1(\gamma + 1) \bar{\rho}^\gamma (\rho - \bar{\rho}) \tag{4.26}$$

where $\bar{\rho}$ is the Barenblatt solution defined in (2.2). The entropy inequality implies

$$\eta_{*t} + (C_1 \bar{\rho}^{\gamma+1} + C_1(\gamma + 1) \bar{\rho}^\gamma (\rho - \bar{\rho}))_t + \tilde{q}_x + 2\kappa C_2 m^2 + \kappa A_m m \leq 0. \tag{4.27}$$

Note that

$$C_1(\gamma + 1) = 2\kappa C_2 \tag{4.28}$$

due to (4.22), and

$$(\bar{\rho}^{\gamma+1})_t = (\gamma + 1) \bar{\rho}^\gamma \bar{\rho}_t = -(\gamma + 1) \bar{m}^2 + (\dots)_x, \tag{4.29}$$

where $(\dots)_x$ denotes terms which vanish after integrating over \mathbf{R} . We thus update (4.27) by

$$\begin{aligned} \eta_{*t} + (2\kappa C_2 \bar{\rho}^\gamma (\rho - \bar{\rho}))_t + 2\kappa C_2 (m - \bar{m})^2 \\ + 4\kappa C_2 \bar{m} (m - \bar{m}) + \kappa A_m m + (\dots)_x \leq 0. \end{aligned} \tag{4.30}$$

Since $\rho - \bar{\rho} = -y_x$ and

$$\bar{m} (m - \bar{m}) = -(\bar{\rho}^\gamma)_x y_t = -(\bar{\rho}_x^\gamma y)_t + (\bar{\rho}^\gamma)_t (\rho - \bar{\rho}) + (\dots)_x, \tag{4.31}$$

(4.30) is reduced into

$$\begin{aligned} \eta_{*t} - 2\kappa C_2 (\bar{\rho}_x^\gamma y)_t + 2\kappa C_2 (m - \bar{m})^2 + 4\kappa C_2 (\bar{\rho}^\gamma)_t (\rho - \bar{\rho}) \\ + \kappa A_m m + (\dots)_x \leq 0. \end{aligned} \tag{4.32}$$

For any small positive constant $\varepsilon > 0$, we define

$$\mu(\varepsilon) = 1 - \frac{1}{2(\gamma + 1)} - \varepsilon.$$

Multiplying (4.32) by $(1 + t)^{\mu(\varepsilon)}$, integrating the result on $\mathbf{R} \times [0, t]$ and using Lemma 4.4, we find

$$\begin{aligned}
 & (1 + t)^{\mu(\varepsilon)} \int_{-\infty}^{+\infty} \eta_* \, dx + \int_0^t \int_{-\infty}^{+\infty} (1 + \tau)^{\mu(\varepsilon)} [2\kappa C_2(m - \bar{m})^2 + \kappa A_m m] \, dx d\tau \\
 & \leq C + C \int_0^t \int_{-\infty}^{+\infty} (1 + \tau)^{\mu(\varepsilon)-1} \eta_* \, dx d\tau + C(1 + t)^{\mu(\varepsilon)} \int_{-\infty}^{+\infty} |\bar{\rho}_x^\gamma| |y| \, dx \\
 & \quad + C \int_0^t \int_{-\infty}^{+\infty} (1 + \tau)^{\mu(\varepsilon)-1} |\bar{\rho}_x^\gamma| |y| \, dx d\tau \\
 & \quad + C \int_0^t \int_{-\infty}^{+\infty} (1 + \tau)^{\mu(\varepsilon)} |\bar{\rho}_t^\gamma| |\rho - \bar{\rho}| \, dx d\tau \\
 & =: C + \sum_{i=1}^4 I_i.
 \end{aligned} \tag{4.33}$$

From (1.5), (4.20), (4.26), (4.1) and Lemma 4.4, we have

$$\eta_* \leq C(\rho^\gamma - \bar{\rho}^\gamma)(\rho - \bar{\rho}) + C \frac{m^2}{\rho}, \tag{4.34}$$

which, together with Lemma 4.3, implies

$$\begin{aligned}
 I_1 & \leq C \int_0^t \int_{-\infty}^{+\infty} \eta_* \, dx d\tau \\
 & \leq C \int_0^t \int_{-\infty}^{+\infty} \left((\rho^\gamma - \bar{\rho}^\gamma)(\rho - \bar{\rho}) + \frac{m^2}{\rho} \right) \, dx d\tau \leq C.
 \end{aligned} \tag{4.35}$$

By Lemma 4.3 again, we have

$$\begin{aligned}
 I_2 & \leq C \int_{-\infty}^{+\infty} y^2 \, dx + C(1 + t)^{2\mu(\varepsilon)} \int_{-\infty}^{+\infty} |\bar{\rho}_x^\gamma|^2 \, dx \\
 & \leq C + C(1 + t)^{2\mu(\varepsilon)-2+\frac{1}{\gamma+1}} \\
 & \leq C + C(1 + t)^{-2\varepsilon} \leq C.
 \end{aligned} \tag{4.36}$$

Similarly, we have

$$\begin{aligned}
 I_3 & \leq C \int_0^t \int_{-\infty}^{+\infty} (1 + \tau)^{\mu(\varepsilon)-1} |\bar{\rho}_x^\gamma| |y| \, dx d\tau \\
 & \leq \int_0^t \int_{-\infty}^{+\infty} (1 + \tau)^{-1-\varepsilon} y^2 \, dx d\tau + C \int_0^t \int_{-\infty}^{+\infty} (1 + \tau)^{2\mu(\varepsilon)-1+\varepsilon} |\bar{\rho}_x^\gamma|^2 \, dx d\tau \\
 & \leq \int_0^t (1 + \tau)^{-1-\varepsilon} \, d\tau + C \int_0^t (1 + \tau)^{2\mu(\varepsilon)-3+\varepsilon+\frac{1}{\gamma+1}} \, d\tau \\
 & \leq C + C \int_0^t (1 + \tau)^{-1-\varepsilon} \, d\tau \\
 & \leq C.
 \end{aligned}$$

It remains to deal with the term I_4 . From (2.2) and (4.1), we obtain, when $1 < \gamma < 3$,

$$\begin{aligned}
 I_4 &\leq C \int_0^t \int_{-\infty}^{+\infty} (1 + \tau)^{\mu(\varepsilon)} |\bar{\rho}_t^\gamma| |\rho - \bar{\rho}| \, dx d\tau \\
 &\leq C \int_0^t \int_{-\infty}^{+\infty} \bar{\rho}^{\gamma-1} (\rho - \bar{\rho})^2 \, dx d\tau \\
 &\quad + C \int_0^t \int_{-\infty}^{+\infty} (1 + \tau)^{2\mu(\varepsilon)} \bar{\rho}^{1-\gamma} |\bar{\rho}_t^\gamma|^2 \, dx d\tau \\
 &\leq C + C \int_0^t (1 + \tau)^{2\mu(\varepsilon)-2-\frac{\gamma}{\gamma+1}} \, d\tau \\
 &\leq C + C \int_0^t (1 + \tau)^{-1-2\varepsilon} \, d\tau \leq C.
 \end{aligned}
 \tag{4.37}$$

Thus, in view of (4.33), (4.35), (4.36) and (4.37), we get

$$\int_{-\infty}^{+\infty} \eta_* \, dx \leq C(1 + t)^{-\mu(\varepsilon)}.
 \tag{4.38}$$

Therefore we can prove the following lemma,

Lemma 4.5. *Under the conditions of Theorem 1.2, it holds for any $t > 0$ that*

$$\begin{aligned}
 &\| (m - \bar{m})(\cdot, t) \|_{L^2}^2 + \| (\rho - \bar{\rho})(\cdot, t) \|_{L^{\gamma+1}}^{\gamma+1} \\
 &\quad + \int_{-\infty}^{+\infty} (\rho^{\gamma-1} + \bar{\rho}^{\gamma-1}) (\rho - \bar{\rho})^2 \, dx \leq C(1 + t)^{\frac{1}{2(\gamma+1)} + \varepsilon - 1}, \\
 &\int_0^t (1 + \tau)^{1 - \frac{1}{2(\gamma+1)} - \varepsilon} \| (m - \bar{m})(\cdot, \tau) \|_{L^2}^2 \, d\tau \leq C,
 \end{aligned}
 \tag{4.39}$$

for any positive constant ε .

Proof. From Lemma 4.1 and the following fact

$$(\rho^{\gamma-1} + \bar{\rho}^{\gamma-1}) (\rho - \bar{\rho})^2 \geq C(\rho^\gamma - \bar{\rho}^\gamma) (\rho - \bar{\rho}) \geq C|\rho - \bar{\rho}|^{\gamma+1},$$

the estimates on ρ are directly from (4.39). It remains to show the estimate on m . For this purpose, we have

$$\begin{aligned}
 \int_{-\infty}^{+\infty} (m - \bar{m})^2 \, dx &\leq C \int_{-\infty}^{+\infty} (m^2 + \bar{m}^2) \, dx \\
 &\leq C \int_{-\infty}^{+\infty} \eta_* \, dx + \int_{-\infty}^{+\infty} (\bar{\rho}^\gamma)_x^2 \, dx \\
 &\leq C(1 + t)^{\frac{1}{2(\gamma+1)} + \varepsilon - 1} + C(1 + t)^{-\frac{2\gamma+1}{\gamma+1}} \\
 &\leq C(1 + t)^{\frac{1}{2(\gamma+1)} + \varepsilon - 1},
 \end{aligned}$$

where we have used the fact that

$$-\frac{2\gamma + 1}{\gamma + 1} < \frac{1}{2(\gamma + 1)} - 1 = -\frac{2\gamma + 1}{2(\gamma + 1)}.$$

The proof of Lemma 4.5 is complete.

Next, we shall use Lemma 4.5 to further obtain an L^1 convergence rate on density, which is based on the following key observation.

Lemma 4.6. *If $\rho \geq 0$ and $\bar{\rho} \geq 0$ have the same total mass M , then for any $t > 0$,*

$$\int_{-\infty}^{+\infty} |\rho - \bar{\rho}|(x, t) dx \leq 2 \int_{\bar{\rho}>0} |\rho - \bar{\rho}|(x, t) dx. \tag{4.40}$$

Proof. Because

$$\int_{-\infty}^{+\infty} \rho(x, t) dx = \int_{\bar{\rho}>0} \rho dx + \int_{\bar{\rho}=0} \rho dx = \int_{-\infty}^{+\infty} \bar{\rho} dx = \int_{\bar{\rho}>0} \bar{\rho} dx, \tag{4.41}$$

we have

$$\int_{\bar{\rho}=0} \rho dx = \int_{\bar{\rho}>0} (\bar{\rho} - \rho) dx \leq \int_{\bar{\rho}>0} |\bar{\rho} - \rho| dx \tag{4.42}$$

and

$$\int_{-\infty}^{+\infty} |\rho - \bar{\rho}| dx = \int_{\bar{\rho}>0} |\rho - \bar{\rho}| dx + \int_{\bar{\rho}=0} \rho dx \leq 2 \int_{\bar{\rho}>0} |\rho - \bar{\rho}| dx. \tag{4.43}$$

Therefore Lemma 4.6 is complete.

Remark 4.2. Lemma 4.6 discloses the fact that $|\rho - \bar{\rho}|$ on the support of the Barenblatt solution $\bar{\rho}$ plays a leading role in the L^1 estimate. It is noted that the support of $\bar{\rho}$ is a bounded domain for any fixed $t > 0$, thus it is possible to obtain an L^1 convergence rate from an L^p convergence rate.

The decay rates of the L^1 distance between ρ and $\bar{\rho}$ is given in the next lemma.

Lemma 4.7. *Assume the conditions in Theorem 1.2 are satisfied, then*

$$\|\rho - \bar{\rho}\|_{L^1} \leq C(1 + t)^{-\frac{1}{4(\gamma+1)} + \varepsilon}, \quad \forall t > 0,$$

for any $\varepsilon > 0$.

Proof. To take care of the singularity of $\bar{\rho}$, we divide the support of $\bar{\rho}$ into two parts:

$$\begin{aligned} \Omega_0 &= \left(-\sqrt{\frac{A}{B}}(1+t)^{\frac{1}{\gamma+1}} + (1+t)^{-\beta}, \sqrt{\frac{A}{B}}(1+t)^{\frac{1}{\gamma+1}} - (1+t)^{-\beta} \right), \\ \Omega_1 &= \left(-\sqrt{\frac{A}{B}}(1+t)^{\frac{1}{\gamma+1}}, -\sqrt{\frac{A}{B}}(1+t)^{\frac{1}{\gamma+1}} + (1+t)^{-\beta} \right) \\ &\cup \left(\sqrt{\frac{A}{B}}(1+t)^{\frac{1}{\gamma+1}} - (1+t)^{-\beta}, \sqrt{\frac{A}{B}}(1+t)^{\frac{1}{\gamma+1}} \right), \end{aligned} \tag{4.44}$$

where $\beta > 1$ is a constant. Then $\{\bar{\rho} > 0\} = \Omega_0 \cup \Omega_1$.

With the help of Lemma 4.5 and Lemma 4.6, we now compute

$$\begin{aligned} &\int_{\bar{\rho} > 0} |\rho - \bar{\rho}| dx \\ &= \int_{\Omega_0} |\rho - \bar{\rho}| dx + \int_{\Omega_1} |\rho - \bar{\rho}| dx \\ &\leq \left(\int_{\Omega_0} \bar{\rho}^{\gamma-1} |\rho - \bar{\rho}|^2 dx \right)^{\frac{1}{2}} \left(\int_{\Omega_0} \bar{\rho}^{1-\gamma} dx \right)^{\frac{1}{2}} + \int_{\Omega_1} |\rho - \bar{\rho}| dx \\ &\leq C(1+t)^{-\frac{1}{2} + \frac{1}{4(\gamma+1)} + \frac{\varepsilon}{2}} (1+t)^{\frac{\gamma-1}{2(\gamma+1)}} \\ &\quad \times \left(\int_{\Omega_0} \left(A - \frac{Bx^2}{(1+t)^{\frac{2}{\gamma+1}}} \right)^{-1} dx \right)^{\frac{1}{2}} + C(1+t)^{-\beta} \\ &\leq C(1+t)^{-\frac{1}{4(\gamma+1)} + \frac{\varepsilon}{2}} \sqrt{\ln(1+t)} \leq C(1+t)^{-\frac{1}{4(\gamma+1)} + \varepsilon}. \end{aligned} \tag{4.45}$$

Here we have used the fact that

$$\begin{aligned} &\left| \int_{\Omega_0} \left(A - \frac{Bx^2}{(1+t)^{\frac{2}{\gamma+1}}} \right)^{-1} dx \right| \\ &= B^2(1+t)^{\frac{1}{\gamma+1}} \int_{\xi = -\sqrt{\frac{A}{B} + (1+t)^{-\beta - \frac{1}{\gamma+1}}} }^{\xi = \sqrt{\frac{A}{B} - (1+t)^{-\beta - \frac{1}{\gamma+1}}} } \left(\sqrt{\frac{A}{B}} - \xi \right)^{-1} \left(\sqrt{\frac{A}{B}} + \xi \right)^{-1} d\xi \\ &\leq C(1+t)^{\frac{1}{\gamma+1}} \ln(1+t). \end{aligned}$$

This completes the proof of Lemma 4.7.

Theorem 1.2 follows from Lemmas 4.5 and 4.7.

5. Remarks and discussions

This section is devoted to some further remarks on two issues. The first is about alternative asymptotic profiles of (1.1) constructed by T. Liu. The other one is about the optimality of the rates in our Theorem 1.2.

5.1. Liu’s solution

In [26], LIU constructed an interesting solution to (1.1) mimic the Barenblatt solution to (1.2). In this section, we shall discuss Liu’s solution in its one dimensional version and the relation to our studies in this section.

Define $\phi = p'(\rho) = k\gamma\rho^{\gamma-1}$ be the square of the sound speed, one can rewrite (1.1) in terms of u and ϕ :

$$\begin{cases} \phi_t + u\phi_x + (\gamma - 1)\phi u_x = 0 \\ u_t + uu_x + \frac{1}{\gamma-1}\phi_x = -\alpha u. \end{cases} \tag{5.1}$$

Consider solutions with finite mass such that

$$\rho(x, t) \equiv 0, \text{ for } |x| \geq \left(\frac{e(t)}{b(t)}\right)^{\frac{1}{2}},$$

and set

$$\phi(x, t) = \left[e(t) - b(t)x^2\right]_+, \quad u(x, t) = a(t)x, \tag{5.2}$$

for some non-negative smooth functions $a(t)$, $b(t)$ and $e(t)$ to be determined. One thus obtains the following ordinary differential equations for the functions $a(t)$, $b(t)$ and $e(t)$:

$$\begin{cases} e' + (\gamma - 1)ae = 0 \\ b' + (\gamma + 1)ab = 0 \\ a' + a^2 + \alpha a - \frac{2}{\gamma-1}b = 0. \end{cases} \tag{5.3}$$

The first two equations imply that

$$\left[\ln\left(\frac{e^{\gamma+1}}{b^{\gamma-1}}\right)\right]' = 0, \tag{5.4}$$

which induces that

$$e^{\gamma+1} = e_1 b^{\gamma-1}, \tag{5.5}$$

for some constant $e_1 > 0$. Therefore, one only needs to solve the equations for a and b in (5.3). The existence of global solutions to (5.3) thus could be done by the phase-plane diagram analysis on the region

$$a \geq 0, \quad b \geq 0,$$

where $(0, 0)$ is the only stable equilibrium. Furthermore, $b' < 0$ inside the region while a' changes sign across the curve

$$\Gamma_1 : b = \frac{\gamma - 1}{2}(a^2 + \alpha a).$$

Indeed, $a' > 0$ above Γ_1 , and $a' < 0$ below Γ_1 . A further conclusion from this analysis leads to $(a, b)(t) \rightarrow 0$ as $t \rightarrow \infty$.

The above procedure can be performed for the porous medium equation to obtain the Barenblatt solution with the same ansatz as (5.2):

$$\bar{\phi} = \left[\bar{e}(t) - \bar{b}(t)x^2 \right]_+, \quad \bar{u} = \bar{a}(t)x. \tag{5.6}$$

Instead of (5.3), for the porous medium equation, one has

$$\begin{cases} \bar{e}' + (\gamma - 1)\bar{a}\bar{e} = 0 \\ \bar{b}' + (\gamma + 1)\bar{a}\bar{b} = 0 \\ \alpha\bar{a} - \frac{2}{\gamma-1}\bar{b} = 0, \end{cases} \tag{5.7}$$

which is solved explicitly as

$$\begin{cases} \bar{a}(t) = \frac{1}{\gamma+1}t^{-1} \\ \bar{b}(t) = \frac{\alpha(\gamma-1)}{2(\gamma+1)}t^{-1} \\ \bar{e}(t) = e_0t^{-\frac{\gamma-1}{\gamma+1}}, \end{cases} \tag{5.8}$$

if the initial density is chosen as a pointed mass located at the origin. From this solution, one has

$$\bar{\phi} = t^{-\frac{\gamma-1}{\gamma+1}} \left[e_0 - \frac{\alpha(\gamma-1)}{\gamma+1}\xi^2 \right]_+, \quad \bar{\rho} = \left(\frac{\bar{\phi}}{k\gamma} \right)^{\frac{1}{\gamma-1}} = t^{-\frac{1}{\gamma+1}} \left[A - B\xi^2 \right]_+^{\frac{1}{\gamma-1}}, \tag{5.9}$$

where e_0 is chosen as $k\gamma A$, while A, B and ξ were defined in Section 2.

When the initial data for $a(t), b(t)$ and $e(t)$ are chosen such that the initial density is also a pointed mass at the origin with the same total mass as the Barenblatt's defined through $(\bar{a}, \bar{b}, \bar{e})$, these two solutions share the very similar behavior. The main result of [26] is that (a, b, e) approaches to $(\bar{a}, \bar{b}, \bar{e})$ time asymptotically with very fast decay rates, which is restated in the following theorem.

Theorem 5.1. (LIU [26]) *If the solution $(\rho, u)(x, t)$ of (1.1) constructed in terms of a, b and e as solutions of (5.3) carries the same total mass M as $(\bar{\rho}, \bar{u})$ given by \bar{a}, \bar{b} and \bar{e} , then as $t \rightarrow +\infty$, it holds*

$$\frac{b(t)}{a(t)} \rightarrow \frac{\bar{b}}{\bar{a}} = \frac{\alpha(\gamma-1)}{2}, \tag{5.10}$$

and

$$(a(t), b(t), e(t)) = (\bar{a}(t), \bar{b}(t), \bar{e}(t)) \left(1 + O(1)\frac{\ln t}{t} \right), \tag{5.11}$$

where $O(1)$ is independent of $t \geq 1$, but varies with the trajectories of (5.3).

Remark 5.1. Theorem 5.1 is sharper than the original statement in [26], but (5.11) should be the true statement as it is clearly stated in the proof of [26]. On the other hand, there seems to be some misprint in the original statement, as both $\bar{a}(t)$ and $\bar{b}(t)$ decay faster than the remainder term $O(1)\frac{\ln t}{t}$ there.

The detailed information of Liu’s solution, denoted by $(\rho_L, u_L)(x, t)$ given in Theorem 5.1 indicates that

$$|(\rho_L - \bar{\rho})(x, t)| \leq C \left[\frac{\ln(t + 1)}{t + 1} \right]^{\frac{1}{\gamma-1}} \bar{\rho}(x, t), \tag{5.12}$$

which implies that

$$\|(\rho_L - \bar{\rho})(\cdot, t)\|_{L^1} \leq C \left[\frac{\ln(t + 1)}{t + 1} \right]^{\frac{1}{\gamma-1}}. \tag{5.13}$$

Now, one easily concludes from (5.13) and Theorem 1.2 that

$$\|(\rho - \rho_L)(\cdot, t)\|_{L^1} \leq C(1 + t)^{-\frac{1}{4(\gamma+1)} + \varepsilon}, \text{ for } \forall \varepsilon > 0. \tag{5.14}$$

5.2. Optimality of decay rates

It is not clear whether the decay rates we obtained in Theorem 1.2 are optimal or not. When first examining the rates in (5.12) and (5.13), one may feel that the rates in Theorem 1.2 are too slow. Indeed, since Liu’s solution is a particular solution with the same initial data as Barenblatt’s solution, the decay rates obtained in (5.12)–(5.13) are much faster than those stated in Theorem 1.2.

However, Theorem 1.2 is general in its nature. First of all, the results in Theorem 1.2 are valid for any L^∞ entropy weak solutions without any regularity, while Liu’s solution is Lipschitz almost everywhere and is Hölder at the vacuum boundary. Other reasons include the generic condition of the initial mass distribution in Theorem 1.2, without specification of the center of the mass. We note that a shift of the center of the mass could result in a large error compared to the specific choice made in Liu’s solution, which matches the Barenblatt solution in a perfect way. For instance, one could easily estimate the L^1 decay between Barenblatt’s solution and a shifted one as follows.

$$\begin{aligned} \|(\bar{\rho}(x, t) - \bar{\rho}(x + d, t))\|_{L^1} &= O(1)(1 + t)^{-\frac{1}{\gamma+1}} \int_0^{\sqrt{\frac{A}{B}}} \left(\sqrt{\frac{A}{B}} - \xi \right)^{\frac{1}{\gamma-1} - 1} d\xi \\ &= O(1)(1 + t)^{-\frac{1}{\gamma+1}}. \end{aligned} \tag{5.15}$$

This partially explains why the decay rate of Theorem 1.2 is much slower than (5.11).

We further remark that one should not expect the rate in (5.15) for the general case in our problem and for the porous medium equation. In fact, CARRILLO and TOSCANI [4] proved that the L^1 convergence rate from solutions of the porous medium equation with finite total mass and finite second moment to the Barenblatt solution is about $(1 + t)^{-\frac{1}{3\gamma-1}}$ using the relation between the porous medium equation and the Fokker-Planck equation. This idea is later adopted in the recent work of OGAWA [37] for the Keller-Segel system. This rate is much closer to our rates in Theorem 1.2. The difference between these two rates could be explained in many

ways. A distinct feature here is the lack of regularity in our L^∞ solutions. It is also remarkable that we did not question the conditions on the boundedness of the second moment. Furthermore, [4] is on the difference between solutions of the porous medium equation, while we compare the solutions of compressible Euler equations with damping, which is hyperbolic, with the Barenblatt solution. One thus expects slower decay rates in our case than in [4]. Finally, we note the constraints in our proof for better decay rates are from the terms of I_i ($i = 1, 2, 3, 4$) in (4.33)–(4.38). It is clear that the current rates are very hard to improve with current approach.

On the other hand, we did not find an argument on the optimality of our rates in Theorem 1.2. Therefore, the optimal decay rate of $\|\rho - \bar{\rho}\|_{L^1}$ remains as an interesting open problem.

Acknowledgments. FEIMIN HUANG's research is supported in part by NSFC Grant No. 10825102 for distinguished youth scholars, NSFC-NSAF Grant No. 10676037 and 973 project of China, Grant No. 2006CB805902. RONGHUA PAN's research is supported by the National Science Foundation under grant DMS-0807406. ZHEN WANG's research is supported in part by NSFC Grant No. 10871199.

References

1. ARONSON, D.G.: The porous media equations. "Nonlinear Diffusion Problem", Lecture Notes in Math., Vol. 1224 (Eds. Fasano A. and Primicerio M.) Springer, Berlin, 1986
2. BARENBLATT, G.I.: On one class of the one-dimensional problem of non-stationary filtration of a gas in a porous medium. *Prikl. Mat. i Mekh.* **17**, 739–742 (1953)
3. BREZIS, H., CRANDALL, M.: Uniqueness of solutions of the initial-value problem for $u_t - \Delta\phi(u) = 0$. *J. Math. pures et. appl.* **58**, 153–163 (1979)
4. CARRILLO, J.A., TOSCANI, G.: Asymptotic L^1 -decay of solutions of the Porous Medium Equation to self-similarity. *Indiana U. Math. J.* **49**, 113–142 (2000)
5. CHEN, G., FRID, H.: Divergence-measure fields and hyperbolic conservation laws. *Arch. Ration. Mech. Anal.* **147**, 89–118 (1999)
6. CHUEH, K., CONLEY, C., SMOLLER, J.: Positively invariant regions for systems of nonlinear diffusion equations. *Indiana U. Math. J.* **26**, 373–392 (1977)
7. DAFERMOS, C.M.: A system of hyperbolic conservation laws with frictional damping. *Z. Angew. Math. Phys.* **46**, 294–307 (1995)
8. DAFERMOS, C.M.: Estimates for conservation laws with little viscosity. *SIAM J. Math. Anal.* **18**, 409–421 (1987)
9. DAFERMOS, C.M., PAN, R.: Global BV solutions for the p -system with frictional damping. *SIAM J. Math. Anal.* **41**(3), 1190–1205 (2009)
10. DING, X., CHEN, G., LUO, P.: Convergence of the fractional step Lax-Friedrichs and Godunov scheme for isentropic system of gas dynamics. *Commun. Math. Phys.* **121**, 63–84 (1989)
11. DIPERNA, R.: Convergence of viscosity method for isentropic gas dynamics. *Commun. Math. Phys.* **91**, 1–30 (1983)
12. HSIAO, L.: *Quasilinear Hyperbolic Systems and Dissipative Mechanisms*. World Scientific, Singapore, 1997
13. HSIAO, L., LIU, T.: Convergence to nonlinear diffusion waves for solutions of a system of hyperbolic conservation laws with damping. *Commun. Math. Phys.* **143**, 599–605 (1992)
14. HSIAO, L., LIU, T.: Nonlinear diffusive phenomena of nonlinear hyperbolic systems. *Chin. Ann. Math.* **14B**(4), 465–480 (1993)

15. HSIAO, L., LUO, T.: Nonlinear diffusive phenomena of entropy weak solutions for a system of quasilinear hyperbolic conservation laws with damping. *Q. Appl. Math.* **56**(1), 173–198 (1998)
16. HSIAO, L., PAN, R.: The damped p -system with boundary effects. *Contemp. Math.* **255**, 109–123 (2000)
17. HSIAO, L., TANG, S.: Construction and qualitative behavior of solutions for a system of nonlinear hyperbolic conservation laws with damping. *Q. Appl. Math.* **LIII**(3), 487–505 (1995)
18. HSIAO, L., TANG, S.: Construction and qualitative behavior of solutions of perturbed Riemann problem for the system of one-dimensional isentropic flow with damping. *J. Differ. Equ.* **123**(2), 480–503 (1995)
19. HUANG, F., PAN, R.: Asymptotic behavior of the solutions to the damped compressible Euler equations with vacuum. *J. Differ. Equ.* **220**, 207–233 (2006)
20. HUANG, F., PAN, R.: Convergence rate for compressible Euler equations with damping and vacuum. *Arch. Ration. Mech. Anal.* **166**, 359–376 (2003)
21. HUANG, F., MARCATI, P., PAN, R.: Convergence to Barenblatt solution for the compressible Euler equations with damping and vacuum. *Arch. Ration. Mech. Anal.* **176**, 1–24 (2005)
22. HUANG, F., PAN, R., YU, H.: Large time behavior of Euler-Poisson system for semiconductor. *Sci. China Ser. A: Math.* **51**, 965–972 (2008)
23. KAMIN, S.: Source-type solutions for equations of nonstationary filtration. *J. Math. Anal. Appl.* **64**, 263–276 (1978)
24. LIONS, P.L., PERTHAME, B., TADMOR, E.: Kinetic formulation of the isentropic gas dynamics and p -systems. *Commun. Math. Phys.* **163**, 169–172 (1994)
25. LIONS, P.L., PERTHAME, B., SOUGANIDIS, P.: Existence and stability of entropy solutions for the hyperbolic systems of isentropic gas dynamics in Eulerian and Lagrangian coordinates. *Commun. Pure Appl. Math.* **49**, 599–638 (1996)
26. LIU, T.: Compressible flow with damping and vacuum. *Japan J. Appl. Math.* **13**(1), 25–32 (1996)
27. LUSKIN, M., TEMPLE, B.: The existence of a global weak solution to the nonlinear water-hammer problem. *Commun. Pure Appl. Math.* **35**, 697–735 (1982)
28. LIU, T., YANG, T.: Compressible Euler equations with vacuum. *J. Differ. Equ.* **140**, 223–237 (1997)
29. LIU, T., YANG, T.: Compressible flow with vacuum and physical singularity. *Methods Appl. Anal.* **7**, 495–509 (2000)
30. LUO, T., YANG, T.: Interaction of elementary waves for compressible Euler equations with frictional damping. *J. Differ. Equ.* **161**, 42–86 (2000)
31. MARCATI, P., MILANI, A.: The one dimensional Darcy’s law as the limit of a compressible Euler flow. *J. Differ. Equ.* **84**, 129–147 (1990)
32. MARCATI, P., RUBINO, B.: Hyperbolic to parabolic relaxation theory for quasilinear first order systems. *J. Differ. Equ.* **162**, 359–399 (2000)
33. NISHIDA, T.: Nonlinear hyperbolic equations and related topics in fluid dynamics. *Publ. Math. D’Orsay* 46–53 (1978)
34. NISHIHARA, K.: Convergence rates to nonlinear diffusion waves for solutions of system of hyperbolic conservation laws with damping. *J. Differ. Equ.* **131**, 171–188 (1996)
35. NISHIHARA, K., WANG, W., YANG, T.: L_p -convergence rate to nonlinear diffusion waves for p -system with damping. *J. Differ. Equ.* **161**, 191–218 (2000)
36. NISHIHARA, K., YANG, T.: Boundary effect on asymptotic behavior of solutions to the p -system with damping. *J. Differ. Equ.* **156**, 439–458 (1999)
37. OGAWA, T.: Asymptotic stability of a decaying solution to the Keller-Segel system of degenerate type. *Diff. Integr. Equ.* **21**, 1113–1154 (2008)
38. PERTHAME, B., TZAVARAS, A.: Kinetic formulation for system of two conservation laws and elastodynamics. *Arch. Ration. Mech. Anal.* **155**, 1–48 (2000)
39. PAN, R., ZHAO, K.: Initial boundary value problem for compressible Euler equations with damping. *Indiana Univ. Math. J.* **57**, 2257–2282 (2008)

40. PELETIER, L.A., VAN DUYN, C.J.: A class of similarity solutions of the nonlinear diffusion equations. *Nonlinear Anal. TMA* **1**, 223–233 (1977)
41. SERRE, D.: Domaines invariants pour les systèmes hyperboliques de lois de conservation. *J. Differ. Equ.* **69**, 46–62 (1987)
42. SERRE, D., XIAO, L.: Asymptotic behavior of large weak entropy solutions of the damped p -system. *J. Pure. Differ. Equ.* **10**, 355–368 (1997)
43. SMOLLER, J.A.: *Shock Waves and Reaction-Diffusion* equations. Springer, New York, 1980
44. ZHAO, H.: Convergence to strong nonlinear diffusion waves for solutions of p -system with damping. *J. Differ. Equ.* **174**, 200–236 (2001)
45. ZHENG, Y.: Global smooth solutions to the adiabatic gas dynamics system with dissipation terms. *Chinese Ann. Math.* **17A**, 155–162 (1996)
46. ZHU, C.J.: Convergence of viscosity solutions for the system Of nonlinear elasticity. *J. Math. Anal. Appl.* **209**, 585–604 (1997)
47. ZHU, C.J.: Convergence Rates to Nonlinear Diffusion Waves for Weak Entropy Solutions to p -System with Damping. *Sci. China Ser. A* **46**, 562–575 (2003)

Institute of Applied Mathematics,
AMSS, Academia Sinica,
Beijing, China.
e-mail: fhuang@amt.ac.cn

and

School of Mathematics,
Georgia Institute of Technology,
Atlanta, GA 30332, USA.
e-mail: panrh@math.gatech.edu

and

Institute of Physics and Mathematics,
Academia Sinica,
Wuhan, China.
e-mail: zhenwang@wipm.ac.cn

(Received February 5, 2010 / Accepted July 19, 2010)
Published online August 20, 2010 – © Springer-Verlag (2010)