



Control and Stabilization of the Benjamin-Ono Equation in $L^2(\mathbb{T})$

CAMILLE LAURENT, FELIPE LINARES & LIONEL ROSIER

Communicated by A. BRESSAN

Abstract

We study the control and stabilization of the Benjamin-Ono equation in $L^2(\mathbb{T})$, the lowest regularity where the initial value problem is well-posed. This problem was already initiated in Linares and Rosier (Trans Am Math Soc 367:4595–4626, 2015) where a stronger stabilization term was used (that makes the equation of parabolic type in the control zone). Here we employ a more natural stabilization term related to the L^2 -norm. Moreover, by proving a theorem of controllability in L^2 , we manage to prove the global controllability in large time. Our analysis relies strongly on the bilinear estimates proved in Molinet and Pilod (Anal PDE 5:365–395, 2012) and some new extension of these estimates established here.

1. Introduction

In this paper, we consider the BO equation posed on the periodic domain $\mathbb{T} = \mathbb{R}/(2\pi\mathbb{Z})$:

$$u_t + \mathcal{H}u_{xx} - uu_x = 0, \quad x \in \mathbb{T}, \quad t \in \mathbb{R}, \quad (1.1)$$

where u is real valued and the Hilbert transform \mathcal{H} is defined via the Fourier transform as

$$(\widehat{\mathcal{H}u})(\xi) = -i \operatorname{sgn}(\xi) \widehat{u}(\xi), \quad \xi \in \mathbb{Z}.$$

The BO equation, posed on the line, arises as a model in wave propagation in stratified fluids (see [6, 41]) and has widely been studied in many different contexts (see [1–3, 8, 10, 14–16, 18, 20, 23, 25, 36, 38–40, 50, 51]).

In the periodic setting the best known result to date regarding well-posedness for the Cauchy problem was obtained by MOLINET [34] (see also [37]), which guarantees the global well-posedness of the problem (1.1) for initial L^2 data. One of the main tools employed to obtain this result is the use of Tao's gauge transform [51].

The BO equation possesses an infinite number of conserved quantities (see [9]). The first two conserved quantities are

$$I_1(t) = \int_{\mathbb{T}} u(x, t) \, dx$$

and

$$I_2(t) = \int_{\mathbb{T}} u^2(x, t) \, dx.$$

Since the BO equation was derived to study the propagation of interfaces of stratified fluids, it is natural to think of I_1 and I_2 as expressing conservation of volume (or mass) and energy, respectively.

Here we will study the Equation (1.1) from a control point of view with a forcing term $f = f(x, t)$ added to the equation as a control input:

$$u_t + \mathcal{H}u_{xx} - uu_x = f(x, t), \quad x \in \mathbb{T}, \, t \in \mathbb{R}, \tag{1.2}$$

where f is assumed to be supported in a given open set $\omega \subset \mathbb{T}$. In control theory the following problems are essential:

Exact Control Problem: Given an initial state u_0 and a terminal state u_1 in a certain space, can one find an appropriate control input f so that the Equation (1.2) admits a solution u which satisfies $u(\cdot, 0) = u_0$ and $u(\cdot, T) = u_1$?

Stabilization Problem: Can one find a feedback law $f = Ku$ so that the resulting closed-loop system

$$u_t + \mathcal{H}u_{xx} - uu_x = Ku, \quad x \in \mathbb{T}, \, t \in \mathbb{R}^+$$

is asymptotically stable as $t \rightarrow +\infty$?

Those questions were first investigated by RUSSELL AND ZHANG in [49] for the Korteweg-de Vries equation, which serves as a model for propagation of surface waves along a channel:

$$u_t + u_{xxx} - uu_x = f, \quad x \in \mathbb{T}, \, t \in \mathbb{R}. \tag{1.3}$$

In their work, in order to keep the mass $I_1(t)$ conserved, the control input was chosen in the form

$$f(x, t) = (\mathcal{G}h)(x, t) := a(x) \left(h(x, t) - \int_{\mathbb{T}} a(y)h(y, t) \, dy \right) \tag{1.4}$$

where h is considered as a new control input, and $a(x)$ is a given nonnegative smooth function such that $\{x \in \mathbb{T}; a(x) > 0\} = \omega$ and

$$2\pi[a] = \int_{\mathbb{T}} a(x) \, dx = 1.$$

For the chosen a , it is easy to see that

$$\frac{d}{dt} \int_{\mathbb{T}} u(x, t) \, dx = \int_{\mathbb{T}} f(x, t) \, dx = 0 \quad \forall t \in \mathbb{R}$$

for any solution $u = u(x, t)$ of the system

$$u_t + u_{xxx} - uu_x = \mathcal{G}h, \quad x \in \mathbb{T}, \quad t \in \mathbb{R}. \quad (1.5)$$

Thus the *mass* of the system is indeed conserved.

The control of dispersive nonlinear wave equations on a periodic domain has been extensively studied in the last decade: see example [28, 46, 49] for the Korteweg-de Vries equation, [33] for the Boussinesq system, [48] for the Benjamin-Bona-Mahony equation, and [12, 26, 27, 44, 47] for the nonlinear Schrödinger equation. By contrast, the control theory of the BO equation is at its early stage. The linear problem was studied by LINARES AND ORTEGA [29]. They established the following results regarding control and stabilization in this case.

Theorem A. ([29]) *Let $s \geq 0$ and $T > 0$ be given. Then for any $u_0, u_1 \in H^s(\mathbb{T})$ with $[u_0] = [u_1]$ one can find a control input $h \in L^2(0, T, H^s(\mathbb{T}))$ such that the solution of the system*

$$u_t + \mathcal{H}u_{xx} = \mathcal{G}h, \quad u(x, 0) = u_0(x) \quad (1.6)$$

satisfies $u(x, T) = u_1(x)$.

In order to stabilize (1.6), they employed a simple control law

$$h(x, t) = -\mathcal{G}^*u(x, t) = -\mathcal{G}u(x, t).$$

The resulting closed-loop system reads

$$u_t + \mathcal{H}u_{xx} = -\mathcal{G}\mathcal{G}^*u.^1$$

Theorem B. ([29]) *Let $s \geq 0$ be given. Then there exist some constants $C > 0$ and $\lambda > 0$ such that for any $u_0 \in H^s(\mathbb{T})$, the solution of*

$$u_t + \mathcal{H}u_{xx} = -\mathcal{G}\mathcal{G}^*u, \quad u(x, 0) = u_0(x),$$

satisfies

$$\|u(\cdot, t) - [u_0]\|_{H^s(\mathbb{T})} \leq C e^{-\lambda t} \|u_0 - [u_0]\|_{H^s(\mathbb{T})} \quad \forall t \geq 0.$$

One of the main difficulties of extending the linear results to the nonlinear ones comes from the fact that one cannot use the contraction principle in its usual form to establish the local well-posedness of BO in $H_0^s(\mathbb{T}) = \{u \in H^s(\mathbb{T}), [u] = 0\}$ for $s \geq 0$ (see [39]). The method of proof in [34, 37] strongly used Tao's gauge transform, and it is not clear whether this approach can be followed when an additional control term is present in the equation. Nevertheless, LINARES AND ROSIER in [31] obtained the first results regarding stabilization and control for the BO equation. For completeness we will briefly describe these results.

¹ Actually, it is easily seen that \mathcal{G} is self-adjoint (that is $\mathcal{G}^* = \mathcal{G}$). However, we shall keep the usual notation for the feedback $h = -\mathcal{G}^*u$ throughout.

In [31] was considered the following feedback law

$$h = -D(\mathcal{G}u),$$

where $\widehat{D}u(\xi) = |\xi| \widehat{u}(\xi)$ is used to stabilize the BO equation. Thus scaling in (1.3) by u gives (at least formally)

$$\frac{1}{2} \|u(T)\|_{L^2(\mathbb{T})}^2 + \int_0^T \|D^{\frac{1}{2}}(\mathcal{G}u)\|_{L^2(\mathbb{T})}^2 dt = \frac{1}{2} \|u_0\|_{L^2(\mathbb{T})}^2, \tag{1.7}$$

which suggests that the energy is dissipated over time. On the other hand, (1.7) reveals a smoothing effect, at least in the region $\{a > 0\}$. Using a *propagation of regularity property* in the same vein as in [12,26–28], it was proved that the smoothing effect holds everywhere, that is

$$\|u\|_{L^2(0,T;H^{\frac{1}{2}}(\mathbb{T}))} \leq C(T, \|u_0\|). \tag{1.8}$$

Using this smoothing effect and the classical compactness/uniqueness argument, it was shown that the corresponding closed-loop equation is semi-globally exponentially stable. More precisely,

Theorem C. ([31]) *Let $R > 0$ be given. Then there exist some constants $C = C(R)$ and $\lambda = \lambda(R)$ such that for any $u_0 \in H_0^0(\mathbb{T})$ with $\|u_0\| \leq R$, the weak solutions in the sense of vanishing viscosity of*

$$u_t + \mathcal{H}u_{xx} - uu_x = -\mathcal{G}(D(\mathcal{G}u)), \quad u(x, 0) = u_0(x), \tag{1.9}$$

satisfy

$$\|u(t)\| \leq Ce^{-\lambda t} \|u_0\| \quad \forall t \geq 0.$$

Using again the smoothing effect (1.8), it was possible to extend (at least locally) the exponential stability from $H_0^0(\mathbb{T})$ to $H_0^s(\mathbb{T})$ for $s > 1/2$.

Theorem D. ([31]) *Let $s \in (\frac{1}{2}, 2]$. Then there exists $\rho > 0$ such that for any data $u_0 \in H_0^s(\mathbb{T})$ with $\|u_0\|_{H^s(\mathbb{T})} < \rho$, there exists for all $T > 0$ a unique solution $u(t)$ of (1.9) in the class $C([0, T], H_0^s(\mathbb{T})) \cap L^2(0, T, H_0^{s+\frac{1}{2}}(\mathbb{T}))$. Furthermore, there exist some constants $C > 0$ and $\lambda > 0$ such that*

$$\|u(t)\|_s \leq Ce^{-\lambda t} \|u_0\|_s \quad \forall t \geq 0.$$

Finally, including the same feedback law $h = -D(\mathcal{G}u)$ in the control input to obtain a smoothing effect, an exact controllability result for the full equation was derived as well. More precisely,

Theorem E. ([31]) *Let $s \in (\frac{1}{2}, 2]$ and $T > 0$ be given. Then there exists $\delta > 0$ such that for any $u_0, u_1 \in H_0^s(\mathbb{T})$ satisfying*

$$\|u_0\|_{H^s(\mathbb{T})} \leq \delta, \quad \|u_1\|_{H^s(\mathbb{T})} \leq \delta$$

one can find a control input $h \in L^2(0, T, H^{s-\frac{1}{2}}(\mathbb{T}))$ such that the system (1.5) admits a solution $u \in C([0, T], H_0^s(\mathbb{T})) \cap L^2(0, T, H_0^{s+\frac{1}{2}}(\mathbb{T}))$ satisfying

$$u(x, 0) = u_0(x), \quad u(x, T) = u_1(x).$$

Our main purpose here is to obtain the control and stabilization result in the largest space where local well-posedness is known for the BO equation, that is, for initial data in $L^2(\mathbb{T})$. As we commented above the main difficulty in treating the problem for data in L^2 is using the gauge transform $w = -\frac{i}{2}P_+(ue^{-\frac{i}{2}F})$ introduced in [51]. We first consider the Cauchy problem associated to the BO equation with a forcing term, that is,

$$\begin{cases} u_t + \mathcal{H}u_{xx} - uu_x = g, & x \in \mathbb{T}, t > 0, \\ u(x, 0) = u_0(x) \end{cases} \quad (1.10)$$

for data in $L^2(\mathbb{T})$ and $g \in L^2([0, T], L^2(\mathbb{T}))$. Then, we establish the well-posedness theory for (1.10). The main tool used is the bilinear estimates approach employed by MOLINET AND PILOD [37] to show the local well-posedness of the BO equation in both cases in the real line and in the periodic setting for data in $L^2(\mathbb{T})$ (see [18] for the first proof of this result in $L^2(\mathbb{R})$). The argument in [37] takes advantage of uniform estimates for small data for which the gauge transform is well defined and a scaled argument (subcritical) is used to consider any size data. In our case, we apply some Sobolev estimates which avoid the use of uniform estimates and we apply directly the argument in [37]. Then we have to analyze the gauge transform for large frequency data, for which this transformation has a bad behavior. To do so we extend previous estimates to overcome this difficulty. For instance we refine the bilinear estimates and obtain an extra time factor (see Lemma 2.8) as well as intermediate estimates to deal with large data (see the ‘‘Appendix’’).

Next we describe the main results in this paper.

We begin with the exact controllability result, more precisely:

Theorem 1.1. (i) *(Small data) For any $T > 0$, there exists some $\delta > 0$ such that for any $u_0, u_1 \in L^2(\mathbb{T})$ with*

$$\|u_0\| \leq \delta, \quad \|u_1\| \leq \delta \quad \text{and} \quad [u_0] = [u_1]$$

one can find a control input $h \in L^2([0, T], L^2(\mathbb{T}))$ such that the solution u of the system

$$\begin{cases} u_t + \mathcal{H}u_{xx} - uu_x = \mathcal{G}h, \\ u(x, 0) = u_0(x) \end{cases} \quad (1.11)$$

satisfies $u(x, T) = u_1(x)$ on \mathbb{T} .

(ii) *(Large data) For any $R > 0$, there exists a positive $T = T(R)$ such that the above property holds for any $u_0, u_1 \in L^2(\mathbb{T})$ with*

$$\|u_0\| \leq R, \quad \|u_1\| \leq R \quad \text{and} \quad [u_0] = [u_1].$$

Remark 1.1. We will give the proof of the theorem assuming that $[u_0] = [u_1] = 0$. The general case can be done after the change of variables $\tilde{u}(x, t) = u(x - t[u_0], t) - [u_0]$ is made in the equation in (1.11). The results in Theorem A remain valid for the new equation in \tilde{u} .

The argument of proof is as follows. The control result for large data (ii) will be a combination of a stabilization result presented below and of the result (i) for small data, as is usual in control theory (see for instance [12, 13, 26–28]). The local control result (i) will be proved using a perturbation argument from the linear result [29]. The difficulty comes from the fact that we need some estimates at higher order for the nonlinear equation, which are quite complicated due to the absence of a direct Duhamel formulation. Indeed, we need to have an expansion for all the elements used for the well-posedness theory. These estimates will come from the Benjamin-Ono equation verified by u , from the equation verified by the gauge transform w and from the “inversion” of the gauge transform. For example, we will need some estimates of the form $2iw = P_+u + o(\|(u, v)\|_X^2)$ and some Lipschitz type estimates of this fact.

In order to stabilize (1.11), we employ a simple control law

$$h(x, t) = -\mathcal{G}^*u(x, t),$$

where \mathcal{G}^* denotes the adjoint operator of \mathcal{G} . Thus the resulting closed-loop system is

$$u_t + \mathcal{H}u_{xx} + uu_x = -\mathcal{G}\mathcal{G}^*u.$$

Theorem 1.2. *There exist some functions $c, \lambda : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ with c nondecreasing and λ nonincreasing such that for any $u_0 \in L^2(\mathbb{T})$, the solution of*

$$\begin{cases} u_t + \mathcal{H}u_{xx} - uu_x = -\mathcal{G}\mathcal{G}^*u, \\ u(x, 0) = u_0(x) \end{cases} \tag{1.12}$$

satisfies

$$\|u(\cdot, t) - [u_0]\|_{L^2(\mathbb{T})} \leq C e^{-\lambda t} \|u_0 - [u_0]\|_{L^2(\mathbb{T})} \quad \forall t \geq 0$$

where $\lambda = \lambda(\|u_0\|)$, $C = C(\|u_0\|)$.

Remark 1.2. After the change of variables $\tilde{u}(x, t) = u(x - t[u_0], t) - [u_0]$ described in Remark 1.1, we are left with some solutions with $[\tilde{u}] = 0$ and with the same equation but with \mathcal{G} replaced by \mathcal{G}_μ defined by

$$(\mathcal{G}_\mu h)(x, t) := a(x - \mu t) \left(h(x - \mu t, t) - \int_{\mathbb{T}} a(y)h(y, t) dy \right) \tag{1.13}$$

with $\mu = [u_0]$.

We will give the proof of the theorem assuming that $\mu = [u_0] = 0$ and detail in some remarks the necessary modifications when \mathcal{G} is replaced by \mathcal{G}_μ .

The proof of the stabilization theorem will be obtained by an observability estimate proved by the compactness-uniqueness method. The difference from the case of the nonlinear Schrödinger equation (see for instance [12, 26, 27]) or other nonlinear conservative equations ([13, 28, 42, 43]) comes again from the use of the gauge transform. We need to propagate some information from the zone of damping to the whole space, yet the absence of a Duhamel formulation prevents us from doing that directly on the solution u . The idea is to transfer this information from

the solution u to the gauge transform w , then to apply some propagation results to w , and finally to obtain the expected result by returning to the original solution u .

As was already mentioned, one of the main ingredients in our analysis is the study of the well-posedness of the IVP

$$\begin{cases} u_t + \mathcal{H}u_{xx} - uu_x = g, & x \in \mathbb{T}, t > 0, \\ u(x, 0) = u_0(x), \end{cases} \quad (1.14)$$

for data $u_0 \in L^2(\mathbb{T})$ and where $g \in L^2([0, T], L^2(\mathbb{T}))$.

More precisely,

Theorem 1.3. *For any $u_0 \in L^2(\mathbb{T})$, $g \in L^2([0, T], L^2(\mathbb{T}))$ and $T > 0$, there exists a solution*

$$u \in C([0, T], L^2(\mathbb{T})) \cap X_T^{-1,1} \cap L_T^4 L^4(\mathbb{T}) \quad (1.15)$$

of (1.14) such that

$$w = \partial_x P_+ \left(e^{-\frac{i}{2} \partial_x^{-1} \tilde{u}} \right) \in X_2 \quad (1.16)$$

where

$$\tilde{u} = u(x - t[u_0], t) - [u_0] \quad \text{and} \quad \widehat{\partial_x^{-1}} := \frac{1}{i\xi}, \quad \xi \in \mathbb{Z}^*,$$

and $X_T^{-1,1}$ and X_2 are defined in (2.3) and (2.5) below, respectively.

The solution u is unique in the class

$$u \in L^\infty((0, T); L^2(\mathbb{T})) \cap L^4((0, T) \times \mathbb{T}) \quad \text{and} \quad w \in X_T^{0,1/2},$$

where $X_T^{0,1/2}$ is defined in (2.3) below.

Moreover, $u \in C([0, T]; L^2(\mathbb{T}))$ and the flow map data–solution: $(u_0, g) \rightarrow u$ is locally Lipschitz continuous from $L^2(\mathbb{T})$ into $C([0, T]; L^2(\mathbb{T}))$.

The proof of this theorem does not follow directly from the theory established by MOLINET AND PILOD [37]. We have to distinguish two cases, small and large data, instead. To apply the scale argument in [37] there is a need of some uniform estimates that do not hold when we introduce the source term g and consider any data. To deal with large data we have to go around the gauge transform since in this case it is not “invertible”. We prove the theorem when $[u_0] = 0$. The general case follows as commented above.

The plan of the paper is as follows: in Section 2 we introduce some notations and establish estimates needed in our analysis. General estimates which do not depend on the size of the data are derived in Section 3. In Section 4 we deduce estimates used to complete the proof of Theorem 1.3 for small data. Next, in Section 5, we establish Theorem 1.3 for large data. The control results are proved in Section 6. Section 7 contains preliminary tools to establish Theorem 1.2. The stabilization is demonstrated in Section 8, and finally we include an appendix containing some technical results.

2. Notations and Preliminary Estimates

We use standard notation in Partial Differential Equations. In addition, we will use C to denote various constants that may change from line to line. For any positive numbers a and b , we use the notation $a \lesssim b$ to mean that there exists a constant C such that $a \leq C b$.

For a 2π -periodic function f we define its Fourier transform on \mathbb{Z} by

$$\widehat{f}(\xi) = \int_{\mathbb{T}} e^{-ix\xi} f(x) dx, \quad \xi \in \mathbb{Z}.$$

The free group associated with the linearized Benjamin-Ono equation denoted by $\mathcal{U}(\cdot)$ is defined as

$$\widehat{\mathcal{U}(t)f}(\xi) = e^{-it|\xi|\xi} \widehat{f}(\xi).$$

The norm of Sobolev spaces $H^s(\mathbb{T})$ will be denoted by $\|\cdot\|_s$ and when $s = 0$ the notation $\|\cdot\|$ will be used.

We will use the following projection operators: for $N \in \mathbb{N}^*$ we define

$$\begin{aligned} \widehat{P_{\pm}f} &= \chi_{\mathbb{R}_{\pm}^*} \widehat{f}, & \widehat{P_{\leq 0}f} &= \chi_{\{\xi \leq 0\}} \widehat{f}, & \widehat{P_{\geq N}f} &= \chi_{\{\xi \geq N\}} \widehat{f} \\ \widehat{P_{\leq N}f} &= \chi_{\{\xi \leq N\}} \widehat{f}, & \widehat{P_0f} &= \widehat{f}(0), & \widehat{P_Nf} &= \chi_{\{|\xi| \leq N\}} \widehat{f}, \end{aligned} \tag{2.1}$$

and

$$\widehat{Q_Nf} = \chi_{\{|\xi| > N\}} \widehat{f}. \tag{2.2}$$

For $s, b \in \mathbb{R}$ we define the spaces $X^{s,b}$, $Z^{s,b}$ and $\widetilde{Z}^{s,b}$ via the norms

$$\|v\|_{X^{s,b}} = \left(\int \int \langle \tau + |\xi|\xi \rangle^{2b} \langle \xi \rangle^{2s} |\widehat{v}(\xi, \tau)|^2 d\xi d\tau \right)^{1/2} \tag{2.3}$$

$$\|v\|_{Z^{s,b}} = \left(\int \left(\int \langle \tau + |\xi|\xi \rangle^b \langle \xi \rangle^s |\widehat{v}(\xi, \tau)| d\xi \right)^2 d\tau \right)^{1/2}, \tag{2.4}$$

and

$$\|v\|_{\widetilde{Z}^{s,b}} = \|P_0v\|_{Z^{s,b}} + \left(\sum_N \|v_N\|_{Z^{s,b}}^2 \right)^{1/2},$$

where $d\xi$ denotes the counting measure in \mathbb{Z} , $\langle x \rangle = 1 + |x|$, and v_N corresponds to a classical Littlewood-Paley decomposition. More precisely, let $\phi \in C_0^\infty(\mathbb{R})$, even such that $\phi \equiv 1$ on $[-1, 1]$, $\text{Supp } \phi \subset (-2, 2)$ and $0 \leq \phi \leq 1$. Define a partition of unity ϕ_N on \mathbb{R} for dyadic $N \geq 1$

$$\phi_1 = \phi \quad \text{and} \quad \phi_N(\xi) = \phi\left(\frac{\xi}{N}\right) - \phi\left(\frac{2\xi}{N}\right) \quad \text{for } N \geq 2.$$

For $v : \mathbb{T} \mapsto \mathbb{C}$ we define v_N by

$$\widehat{v_N} = \phi_N \widehat{v}.$$

We also define the spaces X_2^s and $X_2^{s'}$ as

$$\|v\|_{X_2^s} = \|v\|_{X^{s,1/2}} + \|v\|_{\tilde{Z}^{s,0}} \quad (2.5)$$

and

$$\|v\|_{X_2^{s'}} = \|v\|_{X^{s,-1/2}} + \|v\|_{\tilde{Z}^{s,-1}}. \quad (2.6)$$

When $s = 0$ we will use X_2 and X_2' to simplify the notation.

Next we define the restriction in time spaces. More precisely, for any function space B and any $T > 0$, we denote by B_T the corresponding restriction in time space endowed with the norm

$$\|u\|_{B_T} = \inf_{v \in B} \{ \|v\|_B : v(\cdot) = u(\cdot) \text{ in } [0, T] \}. \quad (2.7)$$

We will still use X_2^s to denote X_{2T}^s for sake of clearness. Note that, as pointed out in [37], we have the embedding

$$X_2^s \subset \tilde{Z}_T^{s,0} \subset C([0, T], H^s(\mathbb{T})).$$

We end this section by introducing the functional space X with the norm

$$\|(u, w)\|_X = \|u\|_{L^\infty([0, T], L^2)} + \|u\|_{L^4([0, T] \times \mathbb{T})} + \|u\|_{X_T^{-1,1}} + \|w\|_{X_T^{0,1/2}} + \|w\|_{\tilde{Z}_T^{0,0}}$$

and we denote by X_1 the functional space corresponding to the norm for u and X_2 the functional space given by the norm corresponding to w as above. In some computations we will also use the notation $U = (u, w) \in X$ and the subscript i to denote different functions U_i .

For most of this work we consider T such that $0 \leq T < 1$.

2.1. Linear Estimates

We will begin presenting some estimates in the Bourgain spaces. We follow [17], where the reader can also find the proofs.

Let $\eta \in C_0^\infty(\mathbb{R})$ such that $0 \leq \eta \leq 1$, $\eta \equiv 1$ in $[-1, 1]$ and $\text{supp}(\eta) \subset [-2, 2]$.

Lemma 2.1. *Let $s \in \mathbb{R}$. Then*

$$\|\eta(t)\mathcal{U}(t)h\|_{X_2^s} \lesssim \|h\|_{H^s}.$$

Lemma 2.2. *Let $s \in \mathbb{R}$. Then for any $0 < \delta < 1/2$,*

$$\left\| \eta(t) \int_0^t \mathcal{U}(t-t') H(t') dt' \right\|_{X^{s, \frac{1}{2}+\delta}} \lesssim \|H\|_{X^{s, -\frac{1}{2}+\delta}} \quad (2.8)$$

and

$$\left\| \eta(t) \int_0^t \mathcal{U}(t-t') H(t') dt' \right\|_{X_2^s} \lesssim \|H\|_{X^{s, -\frac{1}{2}}} + \|H\|_{\tilde{Z}^{s, -1}}. \quad (2.9)$$

Lemma 2.3. *For any $T > 0$, $s \in \mathbb{R}$ and $-\frac{1}{2} < b' \leq b < \frac{1}{2}$, it holds*

$$\|u\|_{X_T^{0,b'}} \lesssim T^{b-b'} \|u\|_{X_T^{0,b}}. \tag{2.10}$$

The next estimate is a Strichartz type estimate (see [37]).

Lemma 2.4. *It holds that*

$$\|u\|_{L_{x,t}^4} \lesssim \|u\|_{X^{0,\frac{3}{8}}}$$

and for any $T > 0$ and $\frac{3}{8} \leq b \leq \frac{1}{2}$,

$$\|u\|_{L_{x,T}^4} \lesssim T^{b-\frac{3}{8}} \|u\|_{X_T^{0,b}}.$$

Recalling the notation $X'_2 = X^{0,-1/2} \cap \widetilde{Z}^{0,-1}$ we have

Lemma 2.5. *For $0 < \varepsilon \ll 1$, it holds*

$$\|z\|_{X'_2} \lesssim T^{1/2-\varepsilon} \|z\|_{L^2([0,T] \times \mathbb{T})} \tag{2.11}$$

where X'_2 is defined as in (2.6).

Proof. From the definition we have that

$$\begin{aligned} \|z\|_{Z^{0,-1}}^2 &= \int \left(\int \langle \tau + |\xi|\xi \rangle^{-1} |\widehat{z}(\tau, \xi)| d\tau \right)^2 d\xi \\ &= \int \left(\int \langle \sigma \rangle^{-1} |\widehat{z}(\sigma - |\xi|\xi, \xi)| d\sigma \right)^2 d\xi \\ &\lesssim \iint \langle \sigma \rangle^{-1+2\varepsilon} |\widehat{z}(\sigma - |\xi|\xi, \xi)|^2 d\sigma d\xi \\ &\lesssim \iint \langle \tau + |\xi|\xi \rangle^{-1+2\varepsilon} |\widehat{z}(\tau, \xi)|^2 d\tau d\xi \\ &= \|z\|_{X^{0,-1/2+\varepsilon}}^2 \end{aligned}$$

where we have used the Cauchy-Schwarz inequality and the integrability of $\langle \sigma \rangle^{-1-2\varepsilon}$. We can use Lemma 2.3 to finish the proof. \square

Define $F = \partial_x^{-1} u$ where $\widehat{\partial_x^{-1} u} = \xi^{-1} \widehat{u}$, $\xi \in \mathbb{Z}^*$. We denote by F_i the previous operator F defined for different functions u_i .

Lemma 2.6. *Let F_i , $i = 1, 2$, be defined as above. Then*

$$\left\| e^{\frac{i}{2}F_1} - e^{\frac{i}{2}F_2} \right\|_{L^\infty(\mathbb{T})} \lesssim \|u_1 - u_2\|_{L^2}, \tag{2.12}$$

$$\left\| P_+ \left(e^{\frac{i}{2}F_1} - e^{\frac{i}{2}F_2} \right) \right\|_{L^\infty(\mathbb{T})} \lesssim (1 + \|u_1\|_{L^2} + \|u_2\|_{L^2}) \|u_1 - u_2\|_{L^2}, \tag{2.13}$$

and

$$\left\| P_+ \left(e^{\frac{i}{2}F_1} - 1 \right) \right\|_{L^\infty(\mathbb{T})} \lesssim \|u_1\|_{L^2}. \tag{2.14}$$

Proof. Using the Sobolev embedding and the definition of F_i we have

$$\left\| e^{\frac{i}{2}F_1} - e^{\frac{i}{2}F_2} \right\|_{L^\infty(\mathbb{T})} \leq \|F_1 - F_2\|_{L^\infty} \leq C \|F_1 - F_2\|_{H^1} \leq C \|u_1 - u_2\|_{L^2}.$$

Next we prove (2.13). Since P_+ is not a continuous operator in L^∞ we use first the Sobolev embedding and then the L^2 continuity.

$$\begin{aligned} \left\| P_+ \left(e^{\frac{i}{2}F_1} - e^{\frac{i}{2}F_2} \right) \right\|_{L^\infty(\mathbb{T})} &\leq C \left\| e^{\frac{i}{2}F_1} - e^{\frac{i}{2}F_2} \right\|_{H^1(\mathbb{T})} \\ &\leq C (1 + \|F_1\|_{H^1} + \|F_2\|_{H^1}) \|F_1 - F_2\|_{H^1}. \end{aligned}$$

The definition of F_i yields the result.

Similarly, we obtain

$$\left\| P_+ \left(e^{\frac{i}{2}F_1} - 1 \right) \right\|_{L^\infty(\mathbb{T})} \leq C \left\| e^{\frac{i}{2}F_1} - 1 \right\|_{H^1(\mathbb{T})} \leq C \|F_1\|_{H^1} \leq C \|u_1\|_{L^2}.$$

□

The next result guarantees the existence of smooth solutions for the Cauchy problem (1.11).

Theorem 2.1. (Existence of smooth solutions)

- (Local) For a given $T > 0$ and $s > 3/2$, the initial value problem

$$\begin{cases} u_t + \mathcal{H}u_{xx} - uu_x = g, & x \in \mathbb{T}, t > 0, \\ u(x, 0) = u_0(x) \end{cases} \quad (2.15)$$

is locally well-posed for initial data $u_0 \in H^s(\mathbb{T})$ and $g \in L^2([0, T], H^s(\mathbb{T}))$. Moreover, the map $(u_0, g) \rightarrow u$ is continuous from $H^s(\mathbb{T}) \times L^2([0, T], H^s(\mathbb{T})) \rightarrow C([0, T], H^s(\mathbb{T}))$.

- (Global) Let $s \geq 2$, $T > 0$. Then for any $(u_0, g) \in H^s(\mathbb{T}) \times L^2([0, T], H^s(\mathbb{T}))$ the IVP (2.15) has a unique solution

$$u \in C([0, T], H^s(\mathbb{T})).$$

We notice that the result is also true with a damping term $-\mathcal{G}g^*u$ instead of g .

Proof. The local theory can be established using parabolic regularization (see for instance [21]) for the existence and uniqueness. The continuous dependence follows by using the Bona-Smith argument [4].

To extend the local theory globally we can employ the quantities conserved by the BO flow (see for instance [9,20]) to derive the appropriate *a priori* estimates.

□

Lemma 2.7. Let $u \in X$. Then, the function

$$\begin{cases} f : (0, T] \rightarrow \mathbb{R} \\ t \mapsto \|u\|_{X_t} \end{cases} \quad (2.16)$$

is continuous, where X_t is the norm of restriction on $[0, t)$. Moreover, there exists C such that

$$\lim_{t \rightarrow 0} f(t) \leq C \|(u(0), w(0))\|_{L^2}.$$

Proof. For a proof of this result see, for instance, [26] (Lemma 1.4) or [24] (Lemma 6.3). \square

In what follows, we will omit the dependence on T of the space X and we will denote $X = X_T$ for simplicity.

2.2. The Equation After Gauge Transform

Now we define the gauge transformation to obtain a new equation (see [37, 51]). Consider the IVP

$$\begin{cases} \partial_t u + \mathcal{H}\partial_x^2 u = u\partial_x u + g = \frac{1}{2}\partial_x(u^2) + g, \\ u(0) = u_0 \end{cases} \quad (2.17)$$

where u is a real valued function and u_0, g are real valued functions with zero mean. Set $w = \partial_x P_+(e^{-\frac{i}{2}F})$, $F = \partial_x^{-1}u$, and $G = \partial_x^{-1}g$, then (2.17) becomes

$$\begin{cases} \partial_t F + \mathcal{H}\partial_x^2 F = \frac{1}{2}(\partial_x F)^2 - \frac{1}{2}P_0(F_x^2) + G, \\ F(0) = \partial_x^{-1}u_0. \end{cases} \quad (2.18)$$

Denote by $W = P_+(e^{-\frac{i}{2}F})$, the solution of

$$\partial_t W - i\partial_x^2 W = -P_+[(\partial_x^2 P_- F)e^{-\frac{i}{2}F}] - \frac{i}{2}P_+\left[\left(-\frac{1}{2}P_0(F_x^2) + G\right)e^{-\frac{i}{2}F}\right].$$

Since $e^{-\frac{i}{2}F} = W + P_{\leq 0}e^{-\frac{i}{2}F}$ and the cancelation $P_+[(\partial_x^2 P_- F)P_{\leq 0}e^{-\frac{i}{2}F}] = 0$ we obtain

$$\partial_t W - i\partial_x^2 W = -P_+[(\partial_x^2 P_- F)W] - \frac{i}{2}P_+\left[\left(-\frac{1}{2}P_0(F_x^2) + G\right)e^{-\frac{i}{2}F}\right].$$

Denoting $w = \partial_x W = -\frac{i}{2}P_+(ue^{-\frac{i}{2}F})$, it follows that

$$\begin{aligned} \partial_t w - i\partial_x^2 w &= -\partial_x P_+[W(P_- \partial_x u)] - \frac{i}{2}\partial_x P_+\left[\left(-\frac{1}{2}P_0(F_x^2) + G\right)e^{-\frac{i}{2}F}\right] \\ &= -\partial_x P_+[W(P_- \partial_x u)] - \frac{i}{2}P_+[ge^{-\frac{i}{2}F}] \\ &\quad - \frac{1}{4}P_+\left[\left(-\frac{1}{2}P_0(u^2) + G\right)ue^{-\frac{i}{2}F}\right]. \end{aligned}$$

We want to prove that at order one, if $u(0) = u_0$, $w = -\frac{i}{2}P_+u$ with

$$u(t) = e^{t\mathcal{H}\partial_x^2}u_0 + \int_0^t e^{(t-\tau)\mathcal{H}\partial_x^2}g(\tau) \, d\tau + \mathcal{O}\left(\|u_0\|_{L^2}^2 + \|g\|_{L^2([0,T],L^2)}^2\right).$$

The main tool in our analysis is equation

$$\begin{cases} \partial_t w - i \partial_x^2 w = -\partial_x P_+ [W(P_- \partial_x u)] - \frac{i}{2} P_+ [g e^{-\frac{i}{2} F}] \\ -\frac{1}{4} P_+ \left[\left(-\frac{1}{2} P_0(u^2) + G \right) u e^{-\frac{i}{2} F} \right] \\ w(0) = w_0. \end{cases} \quad (2.19)$$

We denote the right hand side of (2.19) as

$$\begin{aligned} & -\partial_x P_+ [W(P_- \partial_x u)] - \frac{i}{2} P_+ [g e^{-\frac{i}{2} F}] - \frac{1}{4} P_+ \left[\left(-\frac{1}{2} P_0(u^2) + G \right) u e^{-\frac{i}{2} F} \right] \\ & \equiv \text{I} + \text{II} + \text{III}. \end{aligned} \quad (2.20)$$

In our arguments we use the integral equivalent form of the solution of (2.19), that is,

$$\begin{aligned} w(t) &= \mathcal{W}(t)w_0 + \int_0^t \mathcal{W}(t-t') (\text{I} + \text{II} + \text{III})(t') dt' \\ &\equiv \mathcal{W}(t)w_0 + \mathcal{J}\mathcal{W}(t), \end{aligned} \quad (2.21)$$

where $\mathcal{W}(t)$ denotes the unitary group associated to the linear Schrödinger equation.

Remark 2.1. We observe that the estimates in Lemmas 2.1–2.4 also hold for solution of the IVP (2.19).

We will end this section giving the statement of the key (main) bilinear estimate we will use in our analysis. It was proved by Molinet and Pilod (see Proposition 3.5 of [37]).

Lemma 2.8. *Let $\theta \in (0, 1/8)$. We have uniformly for $0 \leq T \leq 1$*

$$\|\partial_x P_+ [W(P_- \partial_x u)]\|_{X'_2} \leq CT^\theta \|(u, w)\|_X^2.$$

We shall remark that this estimate is a slight modification of the estimates in [37] since we also have a factor T^θ that will be very useful in making some bootstrap and absorption arguments. In the appendix, we sketch the modification of proof of MOLINET-PILOD [37] that allows us to obtain that term.

In what follows, we will fix $\theta \in (0, 1/8)$ such that Lemma 2.8 and (2.11) hold for $0 < T \leq 1$.

3. General Estimates

In this section we will derive estimates needed in our argument which are independent of the size of the initial data.

3.1. Estimates on w

Lemma 3.1. *It holds that*

$$\|w\|_{L^\infty([0, T], L^2)} \leq C \|w\|_{Z_T^{0,0}} \leq C \|(u, w)\|_X,$$

and

$$\|w\|_{L^4([0, T] \times \mathbb{T})} \leq C \|w\|_{X_T^{0,1/2}} \leq C \|(u, w)\|_X.$$

Proof. It readily follows from the space definition and Lemma 2.4. \square

Lemma 3.2. For w in (2.19) and $\theta \in (0, 1/8)$ the following estimates hold

$$\begin{aligned} \|w\|_{X_2} &\leq \|w_0\|_{L^2} + CT^\theta \left(\|g\|_{L^2([0,T],L^2)} + \|g\|_{L^2([0,T],L^2)}^2 \right. \\ &\quad \left. + \|(u, w)\|_X^2 + \|(u, w)\|_X^3 \right). \end{aligned} \tag{3.1}$$

Moreover,

$$\|w - w_L\|_{X_2} \leq CT^\theta \left(\|g\|_{L^2([0,T],L^2)} \|(u, w)\|_X + \|(u, w)\|_X^2 + \|(u, w)\|_X^3 \right) \tag{3.2}$$

where w_L is solution of

$$\begin{cases} \partial_t w_L - i\partial_x^2 w_L = -\frac{i}{2}P_+g \\ w_L(0) = w_0. \end{cases} \tag{3.3}$$

Proof. We first prove that the nonlinear term in (2.21) satisfies

$$\begin{aligned} \|\mathcal{JW}\|_{X_2} &\leq CT^\theta \left(\|(u, w)\|_X^2 + \|g\|_{L^2([0,T]\times\mathbb{T})} + \|(u, w)\|_X^3 \right. \\ &\quad \left. + \|g\|_{L^2([0,T],L^2)} \|(u, w)\|_X \right). \end{aligned} \tag{3.4}$$

Using Lemma 2.2 we obtain

$$\|\mathcal{JW}\|_{X_2} \lesssim \|\mathbb{I}\|_{X'_2} + \|\mathbb{II}\|_{X'_2} + \|\mathbb{III}\|_{X'_2} \tag{3.5}$$

where $\mathbb{I}, \mathbb{II}, \mathbb{III}$ are defined as in (2.20).

From Lemma 2.8 we have that

$$\|\mathbb{I}\|_{X'_2} \leq CT^\theta \|(u, w)\|_X^2.$$

The term \mathbb{II} will be crucial in our analysis. We rewrite it as

$$\begin{aligned} \mathbb{II} &= -\frac{i}{2}P_+ \left[g e^{-\frac{i}{2}F} \right] = -\frac{i}{2}P_+g - \frac{i}{2}P_+ \left[g \left(e^{-\frac{i}{2}F} - 1 \right) \right] \\ &= \mathbb{II}_L + \mathbb{II}_{NL}. \end{aligned} \tag{3.6}$$

Hence Lemma 2.5 yields

$$\begin{aligned} \|\mathbb{II}\|_{X'_2} &\leq CT^\theta \|\mathbb{II}_L\|_{L^2([0,T]\times\mathbb{T})} + \|\mathbb{II}_{NL}\|_{L^2([0,T]\times\mathbb{T})} \\ &\leq CT^\theta \left(\|g\|_{L^2([0,T]\times\mathbb{T})} + \|g\|_{L^2([0,T]\times\mathbb{T})} \left\| e^{-\frac{i}{2}F} - 1 \right\|_{L^\infty([0,T]\times\mathbb{T})} \right) \\ &\leq CT^\theta \left(\|g\|_{L^2([0,T]\times\mathbb{T})} + \|g\|_{L^2([0,T]\times\mathbb{T})} \|u\|_{L^\infty([0,T],L^2)} \right) \\ &\leq CT^\theta \left(\|g\|_{L^2([0,T]\times\mathbb{T})} + \|g\|_{L^2([0,T]\times\mathbb{T})} \|(u, w)\|_X \right). \end{aligned}$$

For \mathbb{III} , we also use Lemma 2.5

$$\begin{aligned} \|\mathbb{III}\|_{X_2'} &\leq C T^\theta \|\mathbb{III}\|_{L^2([0,T] \times \mathbb{T})} \leq C T^\theta \left\| -\frac{1}{2} P_0(u^2) + G \|_{L^2([0,T], L^\infty)} \|u\right\|_{L^\infty([0,T], L^2)} \\ &\leq C T^\theta \left(\|u\|_{L^\infty([0,T], L^2)}^2 + \|g\|_{L^2([0,T], L^2)} \right) \|u\|_{L^\infty([0,T], L^2)} \\ &\leq C T^\theta \left(\|u, w\|_X^3 + \|g\|_{L^2([0,T], L^2)} \|u, w\|_X \right). \end{aligned}$$

Combining the above estimates with (3.5) yields (3.4).

Finally, using Duhamel formulation (2.21), Lemmas 2.1, 2.2, and (3.5) we have

$$\begin{aligned} \|w\|_{X_2} &= \|w\|_{X^{0,1/2}} + \|w\|_{Z^{0,0}} \\ &\leq \|w_0\|_{L^2} + C T^\theta \left(\|g\|_{L^2([0,T], L^2)} + \|g\|_{L^2([0,T], L^2)} \|u, w\|_X + \|u, w\|_X^2 \right. \\ &\quad \left. + \|u, w\|_X^3 \right) \\ &\leq \|w_0\|_{L^2} + C T^\theta \left(\|g\|_{L^2([0,T], L^2)} + \|g\|_{L^2([0,T], L^2)}^2 + \|u, w\|_X^2 + \|u, w\|_X^3 \right). \end{aligned} \tag{3.7}$$

Similarly,

$$\begin{aligned} \|w - w_L\|_{X_2} &= \|w - w_L\|_{X^{0,1/2}} + \|w - w_L\|_{Z^{0,0}} \\ &\leq C T^\theta \left(\|g\|_{L^2([0,T], L^2)} \|u, w\|_X + \|u, w\|_X^2 + \|u, w\|_X^3 \right), \end{aligned} \tag{3.8}$$

where w_L is as in (3.3). \square

Next we derive the Lipschitz estimates corresponding to the X_2 norm.

Lemma 3.3. (Lipschitz estimates) *Let w_1 and w_2 be solutions of (2.19) and $\theta \in (0, 1/8)$. Then the following estimates hold:*

$$\begin{aligned} \|w_1 - w_2\|_{X_2} &\leq \|w_{0,1} - w_{0,2}\| + C T^\theta \|g_1 - g_2\|_{L^2([0,T] \times \mathbb{T})} (1 + \|U_1\|_X) \\ &\quad + C T^\theta \|U_1 - U_2\|_X \left(\|U_1\|_X + \|U_2\|_X + (\|U_1\|_X + \|U_2\|_X)^2 \right. \\ &\quad \left. + \|U_2\|_X^3 + \|g_2\|_{L^2([0,T] \times \mathbb{T})} (1 + \|U_2\|_X) \right) \end{aligned} \tag{3.9}$$

and

$$\begin{aligned} \|(w_1 - w_{L,1}) - (w_2 - w_{L,2})\|_{X_2} &\leq C T^\theta \|g_1 - g_2\|_{L^2([0,T] \times \mathbb{T})} \|U_1\|_X \\ &\quad + C T^\theta \|U_1 - U_2\|_X \left(\|U_1\|_X + \|U_2\|_X + (\|U_1\|_X + \|U_2\|_X)^2 \right. \\ &\quad \left. + \|U_2\|_X^3 + \|g_2\|_{L^2([0,T] \times \mathbb{T})} (1 + \|U_2\|_X) \right) \end{aligned} \tag{3.10}$$

where $w_{L,1}$ and $w_{L,2}$ are solutions of (3.3).

Proof. We first show that

$$\begin{aligned} \|\mathcal{J}\mathcal{W}_1 - \mathcal{J}\mathcal{W}_2\|_{X_2} &\leq CT^\theta \|g_1 - g_2\|_{L^2((0,T] \times \mathbb{T})} (1 + \|U_1\|_X) \\ &\quad + CT^\theta \|U_1 - U_2\|_X \left(\|U_1\|_X + \|U_2\|_X + (\|U_1\|_X + \|U_2\|_X)^2 \right. \\ &\quad \left. + \|U_2\|_X^3 + \|g_2\|_{L^2((0,T] \times \mathbb{T})} (1 + \|U_2\|_X) \right). \end{aligned} \tag{3.11}$$

Let w_1 and w_2 be two solutions of (2.19). We use the notation (2.20). From Lemma 2.2 we obtain

$$\|\mathcal{J}\mathcal{W}_1 - \mathcal{J}\mathcal{W}_2\|_{X_2} \lesssim \|\mathbf{I}\|_{X'_2} + \|\mathbf{II}\|_{X'_2} + \|\mathbf{III}\|_{X'_2}. \tag{3.12}$$

Since \mathbf{I} is bilinear, we get the estimates by writing

$$\begin{aligned} \mathbf{I}_1 - \mathbf{I}_2 &= -\partial_x P_+[W_1(P_-\partial_x u_1)] + \partial_x P_+[W_2(P_-\partial_x u_2)] \\ &= -\partial_x P_+[(W_1 - W_2)(P_-\partial_x u_1)] - \partial_x P_+[W_2(P_-\partial_x(u_1 - u_2))]. \end{aligned}$$

Thus

$$\begin{aligned} \|\mathbf{I}_1 - \mathbf{I}_2\|_{X'_2} &\leq CT^\theta \|w_1 - w_2\|_{X_2} \|u_1\|_{X_1} + \|w_2\|_{X_2} \|u_1 - u_2\|_{X_1} \\ &\leq CT^\theta \|U_1 - U_2\|_X (\|U_1\|_X + \|U_2\|_X). \end{aligned}$$

We write $\mathbf{II} = \mathbf{II}_L + \mathbf{II}_{NL}$ as in (3.6). Since \mathbf{II}_L is linear, we obtain

$$\|\mathbf{II}_{L,1} - \mathbf{II}_{L,2}\|_{X'_2} \leq CT^\theta \|\mathbf{II}_{L,1} - \mathbf{II}_{L,2}\|_{L^2((0,T] \times \mathbb{T})} \leq CT^\theta \|g_1 - g_2\|_{L^2((0,T] \times \mathbb{T})}.$$

For \mathbf{II}_{NL} , we write

$$\begin{aligned} \mathbf{II}_{NL,1} - \mathbf{II}_{NL,2} &= -\frac{i}{2} P_+ \left[\left[g_1 \left(e^{-\frac{i}{2} F_1} - 1 \right) \right] - \left[g_2 \left(e^{-\frac{i}{2} F_2} - 1 \right) \right] \right] \\ &= -\frac{i}{2} P_+ \left[\left[(g_1 - g_2) \left(e^{-\frac{i}{2} F_1} - 1 \right) \right] \right. \\ &\quad \left. + \left[g_2 \left(e^{-\frac{i}{2} F_1} - e^{-\frac{i}{2} F_2} \right) \right] \right] \\ \|\mathbf{II}_{NL,1} - \mathbf{II}_{NL,2}\|_{X'_2} &\leq CT^\theta \|\mathbf{II}_{NL,1} - \mathbf{II}_{NL,2}\|_{L^2((0,T] \times \mathbb{T})} \\ &\leq CT^\theta \left(\|g_1 - g_2\|_{L^2((0,T] \times \mathbb{T})} \|u_1\|_{L^\infty((0,T], L^2)} \right. \\ &\quad \left. + \|g_2\|_{L^2((0,T] \times \mathbb{T})} \|u_1 - u_2\|_{L^\infty((0,T], L^2)} \right) \\ &\leq CT^\theta \left(\|g_1 - g_2\|_{L^2((0,T] \times \mathbb{T})} \|U_1\|_X \right. \\ &\quad \left. + \|g_2\|_{L^2((0,T] \times \mathbb{T})} \|U_1 - U_2\|_X \right). \end{aligned}$$

For \mathbf{III} , we write

$$\begin{aligned} \mathbf{III}_1 - \mathbf{III}_2 &= P_+ \left(\left[\left(-\frac{1}{2} P_0(u_1^2) + G_1 \right) u_1 e^{-\frac{i}{2} F_1} \right] \right. \\ &\quad \left. - \left[\left(-\frac{1}{2} P_0(u_2^2) + G_2 \right) u_2 e^{-\frac{i}{2} F_2} \right] \right) \\ &= P_+ \left[\left(-\frac{1}{2} P_0((u_1 - u_2)(u_1 + u_2)) + G_1 - G_2 \right) u_1 e^{-\frac{i}{2} F_1} \right] \\ &\quad + P_+ \left[\left(-\frac{1}{2} P_0(u_2^2) + G_2 \right) \left((u_1 - u_2) e^{-\frac{i}{2} F_1} + u_2 \left(e^{-\frac{i}{2} F_1} - e^{-\frac{i}{2} F_2} \right) \right) \right]. \end{aligned}$$

Using the same estimates (the only difference is that we use (2.6) for the Lipschitz estimate of the term $e^{-\frac{i}{2}F}$), we then have

$$\begin{aligned}
& \|\mathbb{I}_1 - \mathbb{I}_2\|_{X'_2} \leq CT^\theta \|\mathbb{I}_1 - \mathbb{I}_2\|_{L^2([0,T] \times \mathbb{T})} \\
& \leq CT^\theta \left\| -\frac{1}{2}P_0((u_1 - u_2)(u_1 + u_2)) + G_1 - G_2 \right\|_{L^2([0,T], L^\infty)} \|u_1\|_{L^\infty([0,T], L^2)} \\
& \quad + CT^\theta \left\| -\frac{1}{2}P_0(u_2^2) + G_2 \right\|_{L^2([0,T], L^\infty)} \left(\|u_1 - u_2\|_{L^\infty([0,T], L^2)} \right. \\
& \quad \left. + \|u_2\|_{L^\infty([0,T], L^2)} \left\| e^{-\frac{i}{2}F_1} - e^{-\frac{i}{2}F_2} \right\|_{L^\infty([0,T] \times \mathbb{T})} \right) \\
& \leq CT^\theta \|u_1 - u_2\|_{L^4([0,T] \times \mathbb{T})} \left(\|u_1\|_{L^4([0,T] \times \mathbb{T})} + \|u_2\|_{L^4([0,T] \times \mathbb{T})} \right) \|u_1\|_{L^\infty([0,T], L^2)} \\
& \quad + CT^\theta \|g_1 - g_2\|_{L^2([0,T], L^2)} \|u_1\|_{L^\infty([0,T], L^2)} \\
& \quad + CT^\theta \left(\|u_2\|_{L^\infty([0,T], L^2)}^2 + \|g_2\|_{L^2([0,T], L^2)} \right) \\
& \quad \left(1 + \|u_2\|_{L^\infty([0,T], L^2)} \right) \|u_1 - u_2\|_{L^\infty([0,T], L^2)} \\
& \leq CT^\theta \left(\|U_1 - U_2\|_X (\|U_1\|_X + \|U_2\|_X)^2 + \|g_1 - g_2\|_{L^2([0,T], L^2)} \|U_2\|_X \right) \\
& \quad + CT^\theta \left(\|U_2\|_X^3 + \|g_2\|_{L^2([0,T], L^2)} (1 + \|U_2\|_X) \right) \|U_1 - U_2\|_X.
\end{aligned}$$

The estimates (3.9) and (3.10) follow by using Duhamel formulation, Lemmas 2.1, 2.2, (3.5) and standard estimates. \square

3.2. The Estimates Coming from the Original Benjamin-Ono Equation

We consider again the IVP

$$\begin{cases} \partial_t u + \mathcal{H}\partial_x^2 u = \frac{1}{2}\partial_x(u^2) + g \\ u(0) = u_0. \end{cases} \quad (3.13)$$

Lemma 3.4. *Let u be a solution of (3.13). Then the following estimate holds:*

$$\|u\|_{X_T^{-1,1}} \leq C \left(\|u_0\|_{L^2} + \|(u, w)\|_X^2 + \|g\|_{L^2([0,T], L^2)} \right). \quad (3.14)$$

Moreover, if we denote by u_L the solution of

$$\begin{cases} \partial_t u_L + \mathcal{H}\partial_x^2 u_L = g \\ u(0) = u_0, \end{cases}$$

we have

$$\|u - u_L\|_{X_T^{-1,1}} \leq \|(u, w)\|_X^2. \quad (3.15)$$

Proof. Using the definition of the space $X_T^{-1,1}$ we have the estimate

$$\|u\|_{X_T^{-1,1}} \leq \|u\|_{L^2([0,T], H^{-1})} + \left\| \frac{1}{2}\partial_x(u^2) + g \right\|_{L^2([0,T], H^{-1})}.$$

Using energy estimates we get

$$\|u\|_{L^2([0,T],H^{-1})} \leq C\|u_0\|_{L^2} + C\left\|\frac{1}{2}\partial_x(u^2) + g\right\|_{L^1([0,T],H^{-1})}.$$

Finally, we deduce that

$$\begin{aligned} \|u\|_{X_T^{-1,1}} &\leq C\|u_0\|_{L^2} + \left\|\frac{1}{2}\partial_x(u^2) + g\right\|_{L^2([0,T],H^{-1})} \\ &\leq C\left(\|u_0\|_{L^2} + \left\|u^2\right\|_{L^2([0,T],L^2)} + \|g\|_{L^2([0,T],L^2)}\right) \\ &\leq C\left(\|u_0\|_{L^2} + \|u\|_{L^4([0,T],L^4)}^2 + \|g\|_{L^2([0,T],L^2)}\right) \\ &\leq C\left(\|u_0\|_{L^2} + \|(u, w)\|_X^2 + \|g\|_{L^2([0,T],L^2)}\right). \end{aligned} \tag{3.16}$$

□

Using the previous lemma, it is not difficult to deduce the Lipschitz estimates in the $X_T^{-1,1}$ -norm. Indeed, we have:

Lemma 3.5. (Lipschitz estimates) *The next estimates hold*

$$\begin{aligned} \|u_1 - u_2\|_{X_T^{-1,1}} &\lesssim \|u_{0,1} - u_{0,2}\|_{L^2} + \|g_1 - g_2\|_{L^2([0,T]\times\mathbb{T})} \\ &\quad + \|u_1 - u_2\|_{L^4([0,T]\times\mathbb{T})}(\|(u_1, w_1)\|_X + \|(u_2, w_2)\|_X) \end{aligned} \tag{3.17}$$

and

$$\begin{aligned} \|u_1 - u_{L,1} - (u_2 - u_{L,2})\|_{X_T^{-1,1}} &\lesssim \|(u_1, w_1) - (u_2, w_2)\|_X \\ &\quad \times (\|(u_1, w_1)\|_X + \|(u_2, w_2)\|_X). \end{aligned} \tag{3.18}$$

Proof. The arguments in Lemma 3.4 with the required modifications yield the inequalities. □

3.3. Estimates Coming from the Gauge Transform

Since $w = -\frac{i}{2}P_+(ue^{-\frac{i}{2}F})$ we have that $ue^{-\frac{i}{2}F} = 2iw + P_{\leq 0}(ue^{-\frac{i}{2}F})$. Hence

$$u = 2iw e^{\frac{i}{2}F} + e^{\frac{i}{2}F} P_{\leq 0}(ue^{-\frac{i}{2}F}). \tag{3.19}$$

However, the second term is bad; it roughly says (at the first order in u) that for the negative frequencies, $u = u$, and so it does not allow for inversion. For positive frequencies however,

$$\begin{aligned} P_+u &= 2iP_+[we^{\frac{i}{2}F}] + P_+[P_+(e^{\frac{i}{2}F})P_{\leq 0}(ue^{-\frac{i}{2}F})] \\ &= 2iP_+w + 2iP_+[w(e^{\frac{i}{2}F} - 1)] + P_+[P_+(e^{\frac{i}{2}F})P_{\leq 0}(ue^{-\frac{i}{2}F})] \\ &= 2iw + A + B, \end{aligned} \tag{3.20}$$

where we have used $P_+[P_{\leq 0}(e^{\frac{i}{2}F})P_{\leq 0}(ue^{-\frac{i}{2}F})] = 0$. In this case, the third term is quadratic in u . Indeed,

$$\begin{aligned} B &= P_+ \left[P_+ \left(e^{\frac{i}{2}F} \right) P_{\leq 0} \left(ue^{-\frac{i}{2}F} \right) \right] = P_+ \left[P_+ \left(e^{\frac{i}{2}F} - 1 \right) P_{\leq 0} \left(ue^{-\frac{i}{2}F} \right) \right] \\ &\quad + P_+ \left[P_{\leq 0} \left(ue^{-\frac{i}{2}F} \right) \right] \\ &= P_+ \left[P_+ \left(e^{\frac{i}{2}F} - 1 \right) P_{\leq 0} \left(ue^{-\frac{i}{2}F} \right) \right]. \end{aligned} \tag{3.21}$$

Lemma 3.6. *It holds that*

$$\|P_+u\|_{L^\infty([0,T],L^2)} + \|P_+u\|_{L^4([0,T]\times\mathbb{T})} \leq \|w\|_{X_2} + \|(u, w)\|_{X'}^2 \tag{3.22}$$

and

$$\|P_+u - 2iw\|_{L^\infty([0,T],L^2)} + \|P_+u - 2iw\|_{L^4([0,T]\times\mathbb{T})} \leq \|(u, w)\|_{X'}^2. \tag{3.23}$$

Proof. We use the decomposition (3.20) to estimate P_+ . From Lemma 3.1 it follows that

$$\|2iw\|_{L^\infty([0,T],L^2)} + \|2iw\|_{L^4([0,T]\times\mathbb{T})} \leq \|w\|_{Z_T^{0,0}} + \|w\|_{X_T^{0,1/2}} \leq \|w\|_{X_2}. \tag{3.24}$$

Since P_+ is a pseudo-differential operator of order 0, it maps L^4 into itself. Hence

$$\begin{aligned} \|A\|_{L^\infty([0,T],L^2)} + \|A\|_{L^4([0,T]\times\mathbb{T})} &\lesssim (\|w\|_{L^\infty([0,T],L^2)} + \|w\|_{L^4([0,T]\times\mathbb{T})}) \left\| e^{\frac{i}{2}F} - 1 \right\|_{L^\infty([0,T]\times\mathbb{T})} \\ &\lesssim (\|w\|_{Z_T^{0,0}} + \|w\|_{X_T^{0,1/2}}) \|u\|_{L^\infty([0,T],L^2)} \leq \|(u, w)\|_{X'}^2 \end{aligned} \tag{3.25}$$

and

$$\begin{aligned} \|B\|_{L^\infty([0,T],L^2)} + \|B\|_{L^4([0,T]\times\mathbb{T})} &\lesssim \|P_+ \left(e^{\frac{i}{2}F} - 1 \right)\|_{L^\infty([0,T]\times\mathbb{T})} \left(\|u\|_{L^\infty([0,T],L^2)} \right. \\ &\quad \left. + \|u\|_{L^4([0,T]\times\mathbb{T})} \right) \\ &\lesssim \|(u, w)\|_{X'}^2. \end{aligned}$$

Combining these estimates we obtain the desired inequalities. \square

Next we obtain Lipschitz estimates for the terms involving the operator P_+ . More precisely,

Lemma 3.7. (Lipschitz Estimates) *It holds that*

$$\begin{aligned} & \|P_+u_1 - 2iw_1 - (P_+u_2 - 2iw_2)\|_{L^\infty([0,T],L^2)} + \|P_+u_1 - 2iw_1 \\ & \quad - (P_+u_1 - 2iw_2)\|_{L^4([0,T]\times\mathbb{T})} \\ & \leq \|(u_1, w_1) - (u_2, w_2)\|_X \left(\|(u_1, w_1)\|_X + \|(u_2, w_2)\|_X + \|(u_2, w_2)\|_X^2 \right) \end{aligned} \quad (3.26)$$

and

$$\begin{aligned} & \|P_+u_1 - P_+u_2\|_{L^\infty([0,T],L^2)} + \|P_+u_1 - P_+u_2\|_{L^4([0,T]\times\mathbb{T})} \\ & \leq \|w_1 - w_2\|_{X_2} + \|(u_1, w_1) - (u_2, w_2)\|_X \left(\|(u_1, w_1)\|_X \right. \\ & \quad \left. + \|(u_2, w_2)\|_X + \|(u_2, w_2)\|_X^2 \right). \end{aligned} \quad (3.27)$$

Proof. Recall the decomposition (3.20). To establish the corresponding Lipschitz estimates we write

$$\begin{aligned} A_1 - A_2 &= 2iP_+ \left[w_1 \left(e^{\frac{i}{2}F_1} - 1 \right) \right] - 2iP_+ \left[w_2 \left(e^{\frac{i}{2}F_2} - 1 \right) \right] \\ &= 2iP_+ \left[(w_1 - w_2) \left(e^{\frac{i}{2}F_1} - 1 \right) + w_2 \left(e^{\frac{i}{2}F_1} - e^{\frac{i}{2}F_2} \right) \right]. \end{aligned}$$

Hence

$$\begin{aligned} & \|A_1 - A_2\|_{L^\infty([0,T],L^2)} + \|A_1 - A_2\|_{L^4([0,T]\times\mathbb{T})} \\ & \leq (\|w_1 - w_2\|_{L^\infty([0,T],L^2)} + \|w_1 - w_2\|_{L^4([0,T]\times\mathbb{T})}) \|e^{\frac{i}{2}F_1} - 1\|_{L^\infty([0,T]\times\mathbb{T})} \\ & \quad + (\|w_2\|_{L^\infty([0,T],L^2)} + \|w_2\|_{L^4([0,T]\times\mathbb{T})}) \|e^{\frac{i}{2}F_1} - e^{\frac{i}{2}F_2}\|_{L^\infty([0,T]\times\mathbb{T})} \\ & \leq (\|w_1 - w_2\|_{Z_T^{0,0}} + \|w_1 - w_2\|_{X_T^{0,1/2}}) \|u_1\|_{L^\infty([0,T],L^2)} \\ & \quad + (\|w_2\|_{Z_T^{0,0}} + \|w_2\|_{X_T^{0,1/2}}) \|u_1 - u_2\|_{L^\infty([0,T],L^2)} \\ & \leq \|U_1 - U_2\|_X (\|U_1\|_X + \|U_2\|_X). \end{aligned}$$

On the other hand, we write

$$\begin{aligned} B_1 - B_2 &= P_+ \left[P_+ \left(e^{\frac{i}{2}F_1} - 1 \right) P_{\leq 0} (u_1 e^{-\frac{i}{2}F_1}) - P_+ \left(e^{\frac{i}{2}F_2} - 1 \right) P_{\leq 0} (u_2 e^{-\frac{i}{2}F_2}) \right] \\ &= P_+ \left[P_+ \left(e^{\frac{i}{2}F_1} - e^{\frac{i}{2}F_2} \right) P_{\leq 0} (u_1 e^{-\frac{i}{2}F_1}) \right. \\ & \quad \left. + P_+ \left(e^{\frac{i}{2}F_2} - 1 \right) P_{\leq 0} \left((u_1 - u_2) e^{-\frac{i}{2}F_1} + u_2 \left(e^{-\frac{i}{2}F_1} - e^{-\frac{i}{2}F_2} \right) \right) \right]. \end{aligned}$$

Thus

$$\begin{aligned} & \|B_1 - B_2\|_{L^\infty([0,T],L^2)} + \|B_1 - B_2\|_{L^4([0,T]\times\mathbb{T})} \\ & \leq C \left(\|U_1\|_X + \|U_2\|_X + (\|U_1\|_X + \|U_2\|_X)^2 \right) \|U_1 - U_2\|_X. \end{aligned}$$

These estimates lead to inequalities (3.26) and (3.27). \square

4. Small Data Estimates

The control results as well as the proof of Theorem 1.3 depend on the size data. In this section we aim to establish all the estimates related to small data needed in our arguments. We will also give the proof of Theorem 1.3 corresponding to small data.

Since we are just considering real valued functions we have $P_-u = \overline{P_+u}$, thus

$$\|u\|_{L^\infty([0,T],L^2)} + \|u\|_{L^4([0,T]\times\mathbb{T})} \approx \|P_+u\|_{L^\infty([0,T],L^2)} + \|P_+u\|_{L^4([0,T]\times\mathbb{T})}.$$

Gathering the information from Lemmas 3.2, 3.6 and 3.4 yields

$$\|w\|_{X_2} \leq \|w_0\|_{L^2} + C(\|g\|_{L^2([0,T],L^2)} + \|g\|_{L^2([0,T],L^2)}^2 + \|(u, w)\|_X^2 + \|(u, w)\|_X^3)$$

$$\|u\|_{L^\infty([0,T],L^2)} + \|u\|_{L^4([0,T]\times\mathbb{T})} \leq \|w\|_{X_2} + \|(u, w)\|_X^2$$

$$\|u\|_{X_T^{-1,1}} \leq \|u_0\|_{L^2} + \|(u, w)\|_X^2 + \|g\|_{L^2([0,T],L^2)},$$

and finally

$$\|(u, w)\|_X \lesssim \|u_0\|_{L^2} + \|g\|_{L^2([0,T],L^2)} + \|g\|_{L^2([0,T],L^2)}^2 + \|(u, w)\|_X^2 + \|(u, w)\|_X^3$$

If $\|g\|_{L^2([0,T],L^2)}$ is small enough, it follows that

$$\|(u, w)\|_X \lesssim \|u_0\|_{L^2} + \|g\|_{L^2([0,T],L^2)} + \|(u, w)\|_X^2 + \|(u, w)\|_X^3.$$

Then, since all these quantities are continuous in T and have value at $t = 0$ bounded by $C\|u_0\|_{L^2}$, see Lemma 2.7, we can apply a bootstrap argument to get for $\|u_0\|$ and $\|g\|_{L^2([0,T],L^2)}$, small enough, that

$$\|(u, w)\|_X \lesssim \|u_0\|_{L^2} + \|g\|_{L^2([0,T],L^2)}. \quad (4.1)$$

4.1. First Order Estimates

Using the Lipschitz bound in the previous section we will establish some estimates useful to prove the control results.

The estimates (3.23) and (3.10) yield

$$\|P_+u - 2iw\|_{L^\infty([0,T],L^2)} \leq \|(u, w)\|_X^2 \leq \|u_0\|_{L^2}^2 + \|g\|_{L^2([0,T],L^2)}^2$$

$$\|w - w_L\|_{X_2} = \|w - w_L\|_{X^{0,1/2}} + \|w - w_L\|_{Z^{0,0}}$$

$$\leq \|g\|_{L^2([0,T],L^2)} \|(u, w)\|_X + \|(u, w)\|_X^2.$$

Finally, we have by triangular inequality (noting that $Z^{0,0} \subset L^\infty([0, T], L^2)$)

$$\|P_+u - 2iw_L\|_{L^\infty([0,T],L^2)} \leq \|u_0\|_{L^2}^2 + \|g\|_{L^2([0,T],L^2)}^2$$

where

$$\begin{cases} \partial_t w_L - i\partial_x^2 w_L = \frac{i}{2} P_+g, \\ w_L(0) = w_0. \end{cases}$$

In particular, since u is real valued, $P_-u = \overline{P_+u}$, so

$$\|P_-u - \overline{2i w_L}\|_{L^\infty([0,T],L^2)} \leq \|u_0\|_{L^2}^2 + \|g\|_{L^2([0,T],L^2)}^2.$$

However, we notice that $u = P_+u + P_-u$ and $u_L = 2i w_L + \overline{2i w_L}$. In particular, it follows that

$$\|u - u_L\|_{L^\infty([0,T],L^2)} \lesssim \|u_0\|_{L^2}^2 + \|g\|_{L^2([0,T],L^2)}^2. \tag{4.2}$$

4.2. Proof of Theorem 1.3 (small data case)

In this section we will prove the Lipschitz estimates needed to complete the proof of Theorem 1.3 for small data. More precisely, we have:

Lemma 4.1. *It holds that*

$$\|(u_1, w_1) - (u_2, w_2)\|_X \lesssim \|u_{0,1} - u_{0,2}\| + \|g_1 - g_2\|_{L^2([0,T] \times \mathbb{T})} \tag{4.3}$$

and

$$\|u_1 - u_{L,1} - (u_2 - u_{L,2})\|_{L^\infty([0,T],L^2)} \lesssim \varepsilon (\|u_{0,1} - u_{0,2}\| + \|g_1 - g_2\|_{L^2([0,T] \times \mathbb{T})}). \tag{4.4}$$

Proof. If $\|u_{0,i}\|, \|g_i\|_{L^2([0,T],L^2)}, i = 1, 2$, are small enough, we still have

$$\|(u_i, w_i)\|_X \lesssim \|u_{0,i}\| + \|g_i\|_{L^2([0,T],L^2)}.$$

Assume that $\|u_{0,i}\| + \|g_i\|_{L^2([0,T],L^2)} \leq \varepsilon$, with ε small. Then (3.9) becomes

$$\|w_1 - w_2\|_{X_2} \lesssim \|u_{0,1} - u_{0,2}\| + \|g_1 - g_2\|_{L^2([0,T] \times \mathbb{T})} + \varepsilon \|(u_1, w_1) - (u_2, w_2)\|_X.$$

Inequality (3.17) becomes

$$\|u_1 - u_2\|_{X_T^{-1,1}} \lesssim \|u_{0,1} - u_{0,2}\| + \|g_1 - g_2\|_{L^2([0,T] \times \mathbb{T})} + \varepsilon \|(u_1, w_1) - (u_2, w_2)\|_X$$

and inequality (3.27) becomes

$$\begin{aligned} & \|P_+u_1 - P_+u_2\|_{L^\infty([0,T],L^2)} + \|P_+u_1 - P_+u_2\|_{L^4([0,T] \times \mathbb{T})} \\ & \lesssim \|w_1 - w_2\|_{X_2} + \varepsilon \|(u_1, w_1) - (u_2, w_2)\|_X. \end{aligned}$$

Thus gathering all these estimates of the X -norm components (again using that u is real valued and that the estimates about P_+u are sufficient for the $L^\infty([0, T], L^2)$ and $L^4([0, T] \times \mathbb{T})$ norms), we get

$$\begin{aligned} & \|(u_1, w_1) - (u_2, w_2)\|_X \lesssim \|u_{0,1} - u_{0,2}\| \\ & + \|g_1 - g_2\|_{L^2([0,T] \times \mathbb{T})} + \varepsilon \|(u_1, w_1) - (u_2, w_2)\|_X, \end{aligned}$$

which gives, by absorption, for ε small enough, the inequality (4.3).

For the estimates of the second order terms, we have from (3.10) and (4.3)

$$\|w_1 - w_{L,1} - (w_2 - w_{L,2})\|_{X_2} \lesssim \varepsilon (\|u_{0,1} - u_{0,2}\| + \|g_1 - g_2\|_{L^2([0,T] \times \mathbb{T})})$$

and from (3.26) that

$$\begin{aligned} & \|P_+u_1 - 2iw_1 - (P_+u_2 - 2iw_2)\|_{L^\infty([0,T],L^2)} \\ & \lesssim \varepsilon (\|u_{0,1} - u_{0,2}\| + \|g_1 - g_2\|_{L^2([0,T]\times\mathbb{T})}). \end{aligned}$$

The triangular inequality yields

$$\begin{aligned} & \|P_+u_1 - 2iw_{L,1} - (P_+u_2 - 2iw_{L,2})\|_{L^\infty([0,T],L^2)} \\ & \lesssim \varepsilon (\|u_{0,1} - u_{0,2}\| + \|g_1 - g_2\|_{L^2([0,T]\times\mathbb{T})}). \end{aligned}$$

Once again, we notice that since u_i, g_i are real valued, this implies (4.4). This finishes the proof of the lemma. \square

Proof (Proof of Theorem 1.3 (small data case)). The main ingredient is the Lipschitz inequality (4.3). Once this is established, the proof of the theorem follows from standard arguments. For a detailed and careful proof of it, see [37]. \square

5. Large Data Estimates

In this section we will complete the proof of Theorem 1.3 and establish some estimates for large data useful for further analysis.

5.1. Low-Frequency Estimate

In this subsection, we establish some estimates of the low frequency component of the solution for eventually large data. These estimates are necessary because the gauge transform is not invertible at low frequency for large data.

Let $N \in \mathbb{N}^*$ be fixed. We will prove some estimates that might depend on N .

Lemma 5.1. *For u solution of the IVP (3.13), we have the following uniform estimate for $0 \leq T \leq 1$*

$$\begin{aligned} & \|P_N u\|_{L^\infty([0,T],L^2)} + \|P_N u\|_{L^4([0,T]\times\mathbb{T})} \\ & \leq C(N)\|u_0\|_{L^2} + C(N)\|g\|_{L^2([0,T]\times\mathbb{T})} + C(N)T^{1/2}\|u\|_{L^4([0,T]\times\mathbb{T})}^2, \end{aligned} \quad (5.1)$$

and for two different solutions

$$\begin{aligned} & \|P_N(u_1 - u_2)\|_{L^\infty([0,T],L^2)} + \|P_N(u_1 - u_2)\|_{L^4([0,T]\times\mathbb{T})} \\ & \leq C(N)\|u_{0,1} - u_{0,2}\|_{L^2} + C(N)\|g_1 - g_2\|_{L^2([0,T]\times\mathbb{T})} \\ & \quad + C(N)T^{1/2} (\|u_1\|_{L^4([0,T]\times\mathbb{T})} + \|u_2\|_{L^4([0,T]\times\mathbb{T})}) \|u_1 - u_2\|_{L^4([0,T]\times\mathbb{T})}. \end{aligned} \quad (5.2)$$

Proof. We first link the L^4 estimate to the $L_t^\infty L_x^2$ one by using Sobolev estimates, that is,

$$\begin{aligned} \|P_N u\|_{L^4([0,T] \times \mathbb{T})} &\leq C T^{1/4} \|P_N u\|_{L^\infty([0,T],L^4)} \leq C \|P_N u\|_{L^\infty([0,T],H^1)} \\ &\leq C(N) \|P_N u\|_{L^\infty([0,T],L^2)}. \end{aligned}$$

To estimate the $L_t^\infty L_x^2$ norm we use the solution u_N of the equation satisfied by $u_N = P_N u$, that is

$$\begin{cases} \partial_t u_N + \mathcal{H}(\partial_x^2 u_N) = \frac{1}{2} P_N \partial_x (u^2) + P_N g, \\ u_N(0) = P_N u_0. \end{cases}$$

From semi-group estimates we obtain

$$\begin{aligned} \|P_N u\|_{L^\infty([0,T],L^2)} &\leq C \|u_0\|_{L^2} + C \|P_N \partial_x (u^2)\|_{L^1([0,T],L^2)} + C \|P_N g\|_{L^1([0,T],L^2)} \\ &\leq C \|u_0\|_{L^2} + C(N) T^{1/2} \|P_N (u^2)\|_{L^2([0,T],L^2)} + C \|g\|_{L^1([0,T],L^2)} \\ &\leq C \|u_0\|_{L^2} + C \|g\|_{L^2([0,T],L^2)} + C(N) T^{1/2} \|u\|_{L^4([0,T] \times \mathbb{T})}^2. \end{aligned}$$

The same argument leads to (5.2). \square

5.1.1. Global large data estimates We change a little the decomposition (3.19). The point is, roughly speaking, to make the gauge transform “invertible” at high frequency (the frequency will depend on the size of the data). For large F the second term of the gauge transform (3.19) is bad, and the gauge transform cannot be inverted for large F , but in some sense, it can be inverted at large positive frequencies. Note that another way of obtaining global estimates for large data used by MOLINET [34] was to use the scaling argument to get to some small data (because the equation is subcritical). However this approach requires us to have some estimates uniform on the lengths of the interval. Many of the estimates we used are indeed uniform, but this is not the case for the Sobolev embedding $H^1 \hookrightarrow L^\infty$. We recall that we use the Sobolev spaces H_λ^s for a $(2\pi\lambda)$ -periodic function defined by

$$\|f\|_{H_\lambda^s} = \left(\int_{\mathbb{R}/(2\pi\lambda)\mathbb{Z}} \left(1 + |\xi|^2\right)^s |\widehat{f}(\xi)|^2 (d\xi)_\lambda \right)^{1/2}, \quad \xi \in \lambda^{-1}\mathbb{Z}, \quad (5.3)$$

where $(d\xi)_\lambda$ stands for renormalized counting measure on $\lambda^{-1}\mathbb{Z}$ for large periods λ (see [34], p. 636). One **solution** found by MOLINET in [34] was to apply this estimate to P_1 and the estimate indeed becomes uniform. Our approach is much in the same spirit however we believe that it gives another point of view and may be more reliable in the case of a damped equation or source term.

Let $N \in \mathbb{N}^*$ be large, to be chosen later. Applying the operator $P_{\geq N}$ to both sides to (3.19) we obtain

$$\begin{aligned} P_{\geq N} u &= 2i P_{\geq N} \left[w e^{\frac{i}{2} F} \right] + P_{\geq N} \left[P_{\geq N} \left(e^{\frac{i}{2} F} \right) P_{\leq 0} \left(u e^{-\frac{i}{2} F} \right) \right] \\ &= A_N + B_N. \end{aligned}$$

We apply the same estimates as before to get (the action of operator $P_{\geq N}$ on L^4 can be easily seen to be uniform on N by noticing that $P_{\geq N} = e^{iNx} P_+ e^{-iNx}$)

$$\begin{aligned} \|A_N\|_{L^\infty([0,T],L^2)} + \|A_N\|_{L^4([0,T]\times\mathbb{T})} &\leq C (\|w\|_{L^\infty([0,T],L^2)} + \|w\|_{L^4([0,T]\times\mathbb{T})}) \\ &\leq C (\|w\|_{Z_T^{0,0}} + \|w\|_{X_T^{0,1/2}}) \leq C \|w\|_{X_2}. \end{aligned}$$

For B_N , we also use Lemma A.3 in the Appendix:

$$\begin{aligned} \|B_N\|_{L^\infty([0,T],L^2)} + \|B_N\|_{L^4([0,T]\times\mathbb{T})} \\ \leq \|P_{\geq N}\left(e^{\frac{i}{2}F}\right)\|_{L^\infty([0,T]\times\mathbb{T})} (\|u\|_{L^\infty([0,T],L^2)} + \|u\|_{L^4([0,T]\times\mathbb{T})}) \\ \leq \frac{C}{\sqrt{N}}\|(u, w)\|_X^2. \end{aligned}$$

The above estimates yield:

Lemma 5.2. *For $N \in \mathbb{N}^*$ large, it holds that*

$$\|P_{\geq N}u\|_{L^\infty([0,T],L^2)} + \|P_{\geq N}u\|_{L^4([0,T]\times\mathbb{T})} \leq \|w\|_{X_2} + \frac{C}{\sqrt{N}}\|(u, w)\|_X^2. \tag{5.4}$$

5.1.2. Large data Lipschitz estimates The previous decomposition is still not sufficient. This time, it is because of the first term of the gauge transform. The difficulty now comes from the low frequency of F which, *a priori*, do not allow us to make $P_{\geq N}[w_2(e^{\frac{i}{2}F_1} - e^{\frac{i}{2}F_2})]$ small.

We change a little bit the decomposition inspired by MOLINET [34] p. 663. Let $N \in \mathbb{N}^*$ be large, to be chosen later:

$$e^{-\frac{i}{2}F} = P_{\geq N}W + P_{\leq N}\left(e^{-\frac{i}{2}F}\right).$$

Then decomposing $F = Q_N F + P_N F$, we get

$$e^{-\frac{i}{2}Q_N F} = e^{\frac{i}{2}P_N F} \left(P_{\geq N}W + P_{\leq N}\left(e^{-\frac{i}{2}F}\right) \right).$$

Taking a derivative in x yields

$$\begin{aligned} (Q_N u) e^{-\frac{i}{2}Q_N F} &= -(P_N u) e^{\frac{i}{2}P_N F} \left(P_{\geq N}W + P_{\leq N}\left(e^{-\frac{i}{2}F}\right) \right) \\ &\quad + e^{\frac{i}{2}P_N F} \left(2i P_{\geq N}w + P_{\leq N}\left(ue^{-\frac{i}{2}F}\right) \right) \end{aligned}$$

or

$$\begin{aligned} Q_N u &= -(P_N u) e^{\frac{i}{2}P_N F} \left(P_{\geq N}W + P_{\leq N}\left(e^{-\frac{i}{2}F}\right) \right) \\ &\quad + e^{\frac{i}{2}P_N F} \left(2i P_{\geq N}w + P_{\leq N}\left(ue^{-\frac{i}{2}F}\right) \right) \\ &\quad + (Q_N u) \left(1 - e^{-\frac{i}{2}Q_N F} \right). \end{aligned}$$

We now apply $P_{\geq 3N}$ to the last identity to obtain

$$\begin{aligned} P_{\geq 3N}u &= -P_{\geq 3N} \left[(P_N u) e^{\frac{i}{2} P_N F} P_{\geq N} W \right] - P_{\geq 3N} \left[(P_N u) e^{\frac{i}{2} P_N F} P_{\leq N} \left(e^{-\frac{i}{2} F} \right) \right] \\ &\quad + 2i P_{\geq 3N} \left[e^{\frac{i}{2} P_N F} P_{\geq N} w \right] + P_{\geq 3N} \left[e^{\frac{i}{2} P_N F} P_{\leq N} \left(u e^{-\frac{i}{2} F} \right) \right] \\ &\quad + P_{\geq 3N} \left[(Q_N u) \left(1 - e^{-\frac{i}{2} Q_N F} \right) \right] \\ &= A_N + B_N + C_N + D_N + E_N. \end{aligned}$$

We write

$$\begin{aligned} A_{N,1} - A_{N,2} &= -P_{\geq 3N} \left[(P_N(u_1 - u_2)) e^{\frac{i}{2} P_N F_1} P_{\geq N} W_1 \right. \\ &\quad \left. + (P_N u_2) \left(e^{\frac{i}{2} P_N F_1} - e^{\frac{i}{2} P_N F_2} \right) P_{\geq N} W_1 \right. \\ &\quad \left. + (P_N u_2) e^{\frac{i}{2} P_N F_2} P_{\geq N} (W_1 - W_2) \right]. \end{aligned}$$

Then using the Sobolev estimate (A.3) of the Appendix on the terms with W , we get

$$\begin{aligned} &\|A_{N,1} - A_{N,2}\|_{L^\infty([0,T],L^2)} + \|A_{N,1} - A_{N,2}\|_{L^4([0,T] \times \mathbb{T})} \\ &\leq \frac{C}{\sqrt{N}} \|U_1 - U_2\|_X \left[\|U_1\|_X + \|U_1\|_X^2 \|U_1\|_X + \|U_1\|_X^2 \|U_2\|_X + \|U_2\|_X^2 \right]. \end{aligned} \tag{5.5}$$

To obtain estimates on B_N , we first notice that due to the frequency localization, we have

$$B_N = -P_{\geq 3N} \left[(P_N u) P_{\geq N} \left(e^{\frac{i}{2} P_N F} \right) P_{\leq N} \left(e^{-\frac{i}{2} F} \right) \right].$$

Thus we can write

$$\begin{aligned} B_{N,1} - B_{N,2} &= P_{\geq 3N} \left[(P_N(u_1 - u_2)) P_{\geq N} \left(e^{\frac{i}{2} P_N F_1} \right) P_{\leq N} \left(e^{-\frac{i}{2} F_1} \right) \right. \\ &\quad \left. + (P_N u_2) P_{\geq N} \left(e^{\frac{i}{2} P_N F_1} - e^{\frac{i}{2} P_N F_2} \right) P_{\leq N} \left(e^{-\frac{i}{2} F_1} \right) \right. \\ &\quad \left. + (P_N u_2) P_{\geq N} \left(e^{\frac{i}{2} P_N F_2} \right) P_{\leq N} \left(e^{-\frac{i}{2} F_1} - e^{-\frac{i}{2} F_2} \right) \right]. \end{aligned}$$

Using the estimates of Lemma A.3 for the term with $P_{\geq N}$, we obtain

$$\begin{aligned} &\|B_{N,1} - B_{N,2}\|_{L^\infty([0,T],L^2)} + \|B_{N,1} - B_{N,2}\|_{L^4([0,T] \times \mathbb{T})} \\ &\leq \frac{C}{\sqrt{N}} \|U_1 - U_2\|_X \left[\|U_1\|_X + \|U_1\|_X^2 + \|U_1\|_X^3 + \|U_2\|_X + \|U_2\|_X^2 + \|U_2\|_X^3 \right]. \end{aligned} \tag{5.6}$$

C_N is the crucial term that contains the information on w and we have to estimate it with respect to the norm of $P_N u$.

$$C_{N,1} - C_{N,2} = 2i P_{\geq 3N} \left[\left(e^{\frac{i}{2} P_N F_1} - e^{\frac{i}{2} P_N F_2} \right) P_{\geq N} w_1 - e^{\frac{i}{2} P_N F_2} P_{\geq N} (w_1 - w_2) \right].$$

Thus

$$\begin{aligned} & \|C_{N,1} - C_{N,2}\|_{L^\infty((0,T),L^2)} + \|C_{N,1} - C_{N,2}\|_{L^4((0,T)\times\mathbb{T})} \\ & \leq C \|P_N(u_n - u_2)\|_{L^\infty((0,T),L^2)} \|U_1\|_X + C \|w_1 - w_2\|_{X_2}. \end{aligned} \quad (5.7)$$

D_N can be treated similarly to B_N . By frequency localization, we also obtain

$$D_N = P_{\geq 3N} \left[P_{\geq 2N} \left(e^{\frac{i}{2}P_N F} \right) P_{\leq N} \left(u e^{-\frac{i}{2}F} \right) \right]$$

and then

$$\begin{aligned} & \|D_{N,1} - D_{N,2}\|_{L^\infty((0,T),L^2)} + \|D_{N,1} - D_{N,2}\|_{L^4((0,T)\times\mathbb{T})} \\ & \leq \frac{C}{\sqrt{N}} \|U_1 - U_2\|_X \left[\|U_1\|_X + \|U_1\|_X^2 + \|U_1\|_X^3 + \|U_2\|_X + \|U_2\|_X^2 + \|U_2\|_X^3 \right]. \end{aligned} \quad (5.8)$$

For E_N , we use the third estimate of Lemma A.3.

$$\begin{aligned} E_{N,1} - E_{N,2} &= P_{\geq 3N} \left[(Q_N(u_1 - u_2)) \left(1 - e^{-\frac{i}{2}Q_N F_1} \right) \right] \\ & \quad + P_{\geq 3N} \left[(Q_N u_2) \left(e^{-\frac{i}{2}Q_N F_2} - e^{-\frac{i}{2}Q_N F_1} \right) \right]. \end{aligned}$$

Hence,

$$\begin{aligned} & \|E_{N,1} - E_{N,2}\|_{L^\infty((0,T),L^2)} + \|E_{N,1} - E_{N,2}\|_{L^4((0,T)\times\mathbb{T})} \\ & \leq \frac{C}{\sqrt{N}} \|U_1 - U_2\|_X \left[\|U_1\|_X + \|U_1\|_X^2 + \|U_2\|_X + \|U_2\|_X^2 \right]. \end{aligned} \quad (5.9)$$

Gathering the estimates (5.5)–(5.9) results in:

Lemma 5.3.

$$\begin{aligned} & \|P_{\geq 3N}(u_1 - u_2)\|_{L^\infty((0,T),L^2)} + \|P_{\geq 3N}(u_1 - u_2)\|_{L^4((0,T)\times\mathbb{T})} \\ & \leq C \|P_N(u_n - u_2)\|_{L^\infty((0,T),L^2)} \|U_1\|_X + C \|w_1 - w_2\|_{X_2} \\ & \quad + \frac{C}{\sqrt{N}} \|U_1 - U_2\|_X \left[\|U_1\|_X + \|U_1\|_X^2 + \|U_1\|_X^3 + \|U_2\|_X + \|U_2\|_X^2 + \|U_2\|_X^3 \right]. \end{aligned} \quad (5.10)$$

5.2. Large Data Global Estimates

From Lemmas 3.2, 5.1, 5.2, and 3.16 we have

$$\begin{aligned} & \|w\|_{X_2} \leq \|w_0\|_{L^2} + CT^\theta \left(\|g\|_{L^2((0,T),L^2)} + \|g\|_{L^2((0,T),L^2)}^2 \right) \\ & \quad + \|(u, w)\|_{X'}^2 + \|(u, w)\|_{X'}^3, \\ & \|P_{\geq N}u\|_{L^\infty((0,T),L^2)} + \|P_{\geq N}u\|_{L^4((0,T)\times\mathbb{T})} \leq \|w\|_{X_2} + \frac{C}{\sqrt{N}} \|(u, w)\|_{X'}^2, \\ & \|u\|_{X_T^{1,1}} \leq C \|u_0\|_{L^2} + \|u\|_{L^4((0,T),L^4)}^2 + \|g\|_{L^2((0,T),L^2)}, \end{aligned}$$

and

$$\begin{aligned} & \|P_N u\|_{L^\infty([0,T],L^2)} + \|P_N u\|_{L^4([0,T]\times\mathbb{T})} \\ & \leq C(N)\|u_0\|_{L^2} + C(N)\|g\|_{L^2([0,T]\times\mathbb{T})} + C(N)T^{1/2}\|u\|_{L^4([0,T]\times\mathbb{T})}^2. \end{aligned}$$

Since the term $\|u\|_{L^4([0,T],L^4)}^2$ in the third inequality has no factor of T or $N^{-1/2}$ we need to combine the second and third estimates to obtain

$$\begin{aligned} \|u\|_{X_1} & \leq C(N) \left[\|u_0\|_{L^2} + \|g\|_{L^2([0,T],L^2)} \right] + \|w\|_{X_2} \\ & \quad + \frac{C}{\sqrt{N}} \|(u, w)\|_X^2 + C(N)T^{1/2}\|u\|_{L^4([0,T]\times\mathbb{T})}^2 \\ & \quad + \left(C(N) \left[\|u_0\|_{L^2} + \|g\|_{L^2([0,T],L^2)} \right] + \|w\|_{X_2} \right. \\ & \quad \left. + \frac{C}{\sqrt{N}} \|(u, w)\|_X^2 + C(N)T^{1/2}\|u\|_{L^4([0,T]\times\mathbb{T})}^2 \right)^2. \end{aligned}$$

Notice that $\|w_0\|_{L^2} \leq \|u_0\|_{L^2}$, and since we assume that $0 \leq T \leq 1, N \in N^*$, we have

$$\begin{aligned} \|(u, w)\|_X & \leq C(N) \left[\|u_0\|_{L^2} + \|g\|_{L^2([0,T],L^2)} + \|g\|_{L^2([0,T],L^2)}^2 \right] \\ & \quad + \left(CT^\theta + \frac{C}{\sqrt{N}} + C(N)T^{1/2} \right) \left(\|(u, w)\|_X^2 + \|(u, w)\|_X^3 \right) \\ & \quad + C(N) \left[\|u_0\|_{L^2} + \|g\|_{L^2([0,T],L^2)} + \|g\|_{L^2([0,T],L^2)}^2 \right] \\ & \quad + \left(CT^\theta + \frac{C}{\sqrt{N}} + C(N)T^{1/2} \right) \left(\|(u, w)\|_X^2 + \|(u, w)\|_X^3 \right)^2 \\ & \leq C(N) \left[\|u_0\|_{L^2} + \|u_0\|_{L^2}^2 + \|g\|_{L^2([0,T],L^2)} + \|g\|_{L^2([0,T],L^2)}^4 \right] \\ & \quad + \left(CT^\theta + \frac{C}{\sqrt{N}} + C(N)T^{1/2} \right) \left(\|(u, w)\|_X + \|(u, w)\|_X^6 \right). \end{aligned} \tag{5.11}$$

With this estimate at hand we proceed to establish a uniform Lipschitz bound for large data.

We can now use a bootstrap argument (see for instance Lemma 2.2 of [5]) using Lemma 2.7 for the continuity and limit in zero. Let $R > 0$ such that $\|u_0\|_{L^2} + \|g\|_{L^2([0,T],L^2)} \leq R$. We first pick N large enough and then T small enough (only depending on R and universal constants) to get by a boot strap

$$\|(u, w)\|_X \leq C(R) \left[\|u_0\|_{L^2} + \|g\|_{L^2([0,T],L^2)} \right].$$

This gives the expected result for times T small enough (the smallness only depending on R , that is $T \leq C(R)$). To globalize the result, we use the energy estimate (for t positive or negative), obtained by multiplying the equation by u and integration by parts, and which is valid for smooth solutions

$$\|u(t)\|_{L^2}^2 = \|u(0)\|_{L^2}^2 - 2 \int_0^t g u \, ds.$$

By Cauchy-Schwarz inequality,

$$\|u(t)\|_{L^2}^2 \leq \|u(0)\|_{L^2}^2 + \|g\|_{L^2([0,t],L^2)}^2 + \frac{1}{2} \int_0^t \|u(s)\|_{L^2}^2 ds.$$

We conclude by Gronwall's lemma that the L^2 -norm is bounded on every compact interval.

5.3. Proof of Theorem 1.3 (large data)

In this section we complete the proof of Theorem 1.3 by showing the Lipschitz estimates needed when large data is considered.

Lemma 5.4. *Let $u_0 \in L^2(\mathbb{T})$ and $g \in L^2([0, \tilde{T}], L^2(\mathbb{T}))$ such that $\|u_0\|_{L^2} + \|g\|_{L^2([0,T],L^2)} \leq R$. Let $N \in \mathbb{Z}^+$ be sufficiently large depending on R . Then there exists $T > 0$ small depending on N such that*

$$\|U_1 - U_2\|_X \leq C(N, R) (\|u_{0,1} - u_{0,2}\|_{L^2} + \|g_1 - g_2\|_{L^2([0,T] \times \mathbb{T})}) \quad (5.12)$$

holds.

Proof. Since we assume $\|u_0\|_{L^2} + \|g\|_{L^2([0,T],L^2)} \leq R$. From the previous section we have

$$\|U\|_X \leq C(R).$$

Thus the estimates (3.9) and (5.2) yield

$$\|w_1 - w_2\|_{X_2} \leq \|w_{0,1} - w_{0,2}\|_{L^2} + C(R) \|g_1 - g_2\|_{L^2([0,\tilde{T}] \times \mathbb{T})} + C(R) T^\theta \|U_1 - U_2\|_X \quad (5.13)$$

and

$$\begin{aligned} & \|P_N(u_1 - u_2)\|_{L^\infty([0,T],L^2)} + \|P_N(u_1 - u_2)\|_{L^4([0,T] \times \mathbb{T})} \\ & \leq C(N) \|u_{0,1} - u_{0,2}\|_{L^2} + C(N) \|g_1 - g_2\|_{L^2([0,T] \times \mathbb{T})} + C(N, R) T^{1/2} \|U_1 - U_2\|_X. \end{aligned} \quad (5.14)$$

The estimate (5.10) can be rewritten as

$$\begin{aligned} & \left\| P_{\geq N}(u_1 - u_2) \right\|_{L^\infty([0,T],L^2)} + \left\| P_{\geq N}(u_1 - u_2) \right\|_{L^4([0,T] \times \mathbb{T})} \\ & \leq C(R) \|P_N(u_1 - u_2)\|_{L^\infty([0,T],L^2)} + C \|w_1 - w_2\|_{X_2} + \frac{C(R)}{\sqrt{N}} \|U_1 - U_2\|_X, \end{aligned} \quad (5.15)$$

and from Lemma 3.5 it follows that

$$\|u_1 - u_2\|_{X_T^{-1,1}} \leq \|u_{0,1} - u_{0,2}\|_{L^2} + \|g_1 - g_2\|_{L^2([0,T] \times \mathbb{T})} + C(R) \|u_1 - u_2\|_{L^4([0,T] \times \mathbb{T})}. \quad (5.16)$$

Therefore, combining the estimates (5.13)–(5.16) leads to

$$\begin{aligned} \|U_1 - U_2\|_X &\leq C(N, R) (\|u_{0,1} - u_{0,2}\|_{L^2} + \|g_1 - g_2\|_{L^2([0, T] \times \mathbb{T})}) \\ &\quad + \left(C(N, R) T^{1/2} + \frac{C(R)}{\sqrt{N}} + C(R) T^\theta \right) \|U_1 - U_2\|_X. \end{aligned}$$

So, by choosing N large enough only depending on R and then T small enough (only depending on N), we get

$$\|U_1 - U_2\|_X \leq C(N, R) (\|u_{0,1} - u_{0,2}\|_{L^2} + \|g_1 - g_2\|_{L^2([0, T] \times \mathbb{T})}). \tag{5.17}$$

□

End of the proof of Theorem 1.3. The estimate (5.17) allows us to establish the existence and uniqueness of the solution of the problem (1.14). The continuity of the flow follows by Bona-Smith argument. This completes the proof of Theorem 1.3.

6. Control Results

In this section we will prove Theorem 1.1. First we will consider the damped equation and apply the theory established in the previous section to this case.

6.1. Damped Equation

We consider the equation

$$\begin{cases} \partial_t u + \mathcal{H} \partial_x^2 u = \frac{1}{2} \partial_x (u^2) - \mathcal{G} \mathcal{G}^* u \\ u(T) = 0. \end{cases} \tag{6.1}$$

Setting $g = -\mathcal{G} \mathcal{G}^* u$ in (2.15) we can see the damping term as a source term. Thus, we have

$$\|g\|_{L^2([0, T], L^2)} \leq CT^{1/2} \|u\|_{L^\infty([0, T], L^2)} \leq CT^{1/2} \|(u, w)\|_X.$$

Hence, the estimate (5.11) becomes

$$\begin{aligned} \|(u, w)\|_X &\leq C(N) \left[\|u_0\|_{L^2} + \|u_0\|_{L^2}^2 \right] \\ &\quad + \left(CT^\theta + \frac{C}{\sqrt{N}} + C(N) T^{1/2} \right) \left(\|(u, w)\|_X + \|(u, w)\|_X^6 \right). \end{aligned}$$

We can conclude similarly by a bootstrap for small times (only depending on the size). This gives the expected result for small times. For large times, we use the energy estimate (for t positive or negative)

$$\|u(t)\|_{L^2}^2 = \|u(0)\|_{L^2}^2 - 2 \int_0^t \|\mathcal{G}u(s)\|_{L^2}^2 ds.$$

For positive times, the energy is decreasing, for negative times, we conclude by Gronwall’s lemma.

We conclude similarly for the Lipschitz estimates.

Now we proceed to prove Theorem 1.1. We want to control the following problem

$$\begin{cases} \partial_t u + \mathcal{H}\partial_x^2 u = u\partial_x u - \mathcal{G}\mathcal{G}^*h, \\ u(T) = 0 \end{cases}$$

where \mathcal{G} is defined as in (1.4).

We seek control of the form of a solution of (h_0 is real valued with zero mean)

$$\begin{cases} \partial_t h + \mathcal{H}\partial_x^2 h = 0 \\ h(0) = h_0. \end{cases}$$

We denote by B the nonlinear operator defined by $Bh_0 = u(0)$. Let u_L be the solution of

$$\begin{cases} \partial_t u_L + \mathcal{H}\partial_x^2 u_L = -\mathcal{G}\mathcal{G}^*h \\ u_L(T) = 0. \end{cases}$$

The linear operator from $L^2(\mathbb{T})$ to itself defined by $Lh_0 = u_L(0)$ is the HUM operator (see [32]), which is a bijection of $L^2(\mathbb{T})$ by the observability inequality of LINARES-ORTEGA [29]. We note that $Wh_0 = u(0) = u_L(0) + Kh_0 = Lh_0 + Kh_0$, and that (4.2) can be written as

$$\|Kh_0\|_{L^2} \leq C\|h_0\|_{L^2}^2$$

for $\|h_0\|_{L^2}$ small enough. Let $u_0 \in L^2(\mathbb{T})$. So, the objective $Wh_0 = Lh_0 + Kh_0 = u_0$ is equivalent to $h_0 = -L^{-1}Kh_0 + L^{-1}u_0$, that is h_0 is a fixed point of B defined by $Bh_0 = -L^{-1}Kh_0 + L^{-1}u_0$.

However, we have

$$\|Bh_0\|_{L^2} \leq C\|Kh_0\|_{L^2} + C\|u_0\|_{L^2} \leq C\|h_0\|_{L^2}^2 + C\|u_0\|_{L^2}.$$

Thus, if we denote B_R the unit ball of L^2 of radius $R = 2C\|u_0\|_{L^2}$, B sends B_R into itself if $CR \leq 1/2$, that is, if $\|u_0\|_{L^2}$ is chosen small enough.

Moreover, estimate (4.4) shows that for two solutions of the nonhomogeneous problem u_1, u_2 coming from the controls $g_i = -\mathcal{G}\mathcal{G}^*h_i$ (recall that we have assumed $\|g_i\|_{L^2([0,T] \times \mathbb{T})} \leq \varepsilon$, which is equivalent to $\|h_{0,i}\|_{L^2} \lesssim \varepsilon$), we have the estimates

$$\begin{aligned} \|Kh_{0,1} - Kh_{0,2}\|_{L^2} &\leq \|u_1 - u_{L,1} - (u_2 - u_{L,2})\|_{L^\infty([0,T],L^2)} \\ &\lesssim \varepsilon \|g_1 - g_2\|_{L^2([0,T] \times \mathbb{T})}. \end{aligned}$$

Since $Bh_{0,1} - Bh_{0,2} = Kh_{0,1} - Kh_{0,2}$, this means that for ε small enough (that is $R = 2C\|u_0\|_{L^2}$ small enough), B is contracting and reproduces B_R . Therefore, it has a fixed point which is the expected control.

This ends the proof of Theorem 1.1 part (i) for small data.

7. Study of the Convergence of Some Bounded Sequences of Solutions of the Damped Equations

The aim of this section is to show the convergence of some bounded sequences $\{u_n\}$ of solutions to the damped equation such that $\mathcal{G}u_n$ converges to zero. The results here will be used to perform a contradiction argument which allows us to establish the stabilization theory claimed in Theorem 1.2. As we commented before we have to distinguish two cases: large and small data.

7.1. Large Data

We consider a sequence of solutions $\{u_n\}$ of the damped equation

$$\begin{aligned} \partial_t u_n + \mathcal{H}\partial_x^2 u_n &= u_n \partial_x u_n (-\mathcal{G}\mathcal{G}^* u_n) \\ u_n(0) &= u_{0,n}. \end{aligned}$$

We first suppose that (u_n, w_n) is bounded in X . We then show that $u_n \rightharpoonup 0$ in $L^2([0, T], L^2)$ and weakly-* in $L^\infty([0, T], L^2)$ (see Lemma 7.1 below). Finally we will prove strong convergence in L^2 (see Theorem 7.2 below). One of the main ingredients to obtain the latter result is the propagation compactness theory in [26]. We assume $[u_0] = 0$ to make the proofs more clear but at the end of each proof we will explain in a remark the required modifications for the general case. These modifications are in the same spirit as the ones noticed in Remark 1.2 for \tilde{u} and \mathcal{G}_μ .

Lemma 7.1. *Suppose $u_{n,0}$ is a bounded sequence in L^2 with $[u_{0,n}] = 0$ with associated solutions u_n*

$$\begin{aligned} \partial_t u_n + \mathcal{H}\partial_x^2 u_n &= u_n \partial_x u_n (-\mathcal{G}\mathcal{G}^* u_n) \\ u_n(0) &= u_{0,n}. \end{aligned}$$

Assume moreover that $\mathcal{G}u_n \rightarrow 0$ in $L^2([0, T] \times \mathbb{T})$. Then, $u_n \rightharpoonup 0$ in $L^2([0, T], L^2)$ and weakly- in $L^\infty([0, T], L^2)$.*

Remark 7.1. We could certainly get rid of the term $\mathcal{G}\mathcal{G}^* u_n$ (converging to 0 in $L^2([0, T] \times L^2)$) and work with solutions of the free equation, yet we have chosen to keep it since it does not greatly change the analysis and it seems easier than proving a general perturbation theorem.

Proof. We assume $u_n \rightharpoonup u$ in $L^2([0, T], L^2)$ up to subsequence. The only problem in using directly Proposition 2.8 of [31] is that, because of the nonlinear term, it is not clear that u is a solution of the Benjamin-Ono equation (see MOLINET [35,36] for some cases where the limit equation is a modified equation).

Denote the sequence $c_n(t) = \int_{\mathbb{T}} a(y)u_n(x, t)dy$, bounded in $L^2([0, T])$ and weakly convergent to $c(t)$. Denote $r_n = u_n - c_n$. We have $a(x)[r_n] \rightarrow 0$, in $L^2([0, T] \times \mathbb{T})$. In particular, $r_n \rightarrow 0$ in $L^2([0, T] \times \omega)$ and $r_n^2 \rightarrow 0$ in $L^1([0, T] \times \omega)$.

ω). Therefore, $r_n^2 \rightharpoonup 0$ in the distributional sense of $D'([0, T] \times \omega)$ and hence it is the same for $\partial_x r_n^2$. However, we have

$$\begin{aligned}\partial_x r_n^2 &= \partial_x u_n^2 - 2\partial_x(c_n u_n) + \partial_x c_n^2 = \partial_x u_n^2 - 2\partial_x(c_n u_n) \\ &= \partial_x u_n^2 - 2\partial_x(c_n r_n),\end{aligned}$$

So, we have $r_n \rightarrow 0$ in $L^2([0, T] \times \omega)$ so $c_n r_n \rightarrow 0$ in $L^1([0, T] \times \omega)$ and so in the distributional sense in $D'([0, T] \times \omega)$, and the same for $\partial_x(c_n r_n)$.

Thus, we conclude that $\partial_x u_n^2 \rightharpoonup 0$ in the distributional sense of $D'([0, T] \times \omega)$. In particular, the weak limit u is a solution in the sense of distribution $D'([0, T] \times \omega)$ of

$$u_t + \mathcal{H}u_{xx} = \dot{c}(t) + \mathcal{H}u_{xx} = 0.$$

Therefore, we have $\mathcal{H}u_{xxx} = 0$ in the distributional sense of $D'([0, T] \times \omega)$.

Moreover, since r_n converges to zero in the distributional sense of $D'([0, T] \times \omega)$, it is the same for $r_{n,xxx} = u_{n,xxx}$ and we have $u_{xxx} = 0$ in the distributional sense of $D'([0, T] \times \omega)$.

We conclude that $u \equiv 0$ as in LINARES-ROSIER [31] Proposition 2.8. \square

Remark 7.2. In the case of $[u_{0,n}] = \mu \neq 0$, we can, for instance, set $\underline{u}_n = u_n - \mu$ and obtain, similarly, $u_n \rightharpoonup \mu$, using that \underline{u}_n is a solution of $\partial_t \underline{u}_n + \mathcal{H}\partial_x^2 \underline{u}_n = \mu \partial_x \underline{u}_n + \underline{u}_n \partial_x \underline{u}_n - \mathcal{G}\mathcal{G}^* \underline{u}_n$.

Lemma 7.2. $u_n \rightharpoonup 0$ in $L^2([0, T], L^2)$ implies $w_n \rightharpoonup 0$ in $L^2([0, T], L^2)$ with $w_n = P_+(u_n e^{-\frac{i}{2}F_n})$ (for solutions). The same holds for $u_n e^{-\frac{i}{2}F_n}$.

Proof. F_n weakly-* converges to 0 in $L^\infty([0, T], H^1)$ and $\partial_t F_n$ is bounded in $L^\infty([0, T], H^{-1})$ and hence by Aubin-Lions lemma, F_n converges strongly to 0 in $L^\infty([0, T], L^\infty)$. By the mean value theorem, $e^{-\frac{i}{2}F_n}$ converges strongly to 1 in $L^\infty([0, T], L^\infty)$. Actually, $\partial_t F_n = -\mathcal{H}(\partial_x^2 F_n + \frac{1}{2}(\partial_x F_n)^2 - \frac{1}{2}P_0(F_{n,x}^2) + \partial_x^{-1}\mathcal{G}\mathcal{G}^* u_n)$ is bounded in $L^\infty([0, T], H^{-1})$ since $\partial_x^2 F_n = \partial_x u_n$ and $(\partial_x F_n)^2 = u_n^2$ is bounded in $L^\infty([0, T], L^1) \subset L^\infty([0, T], H^{-1})$.

Let $\varphi \in L^2([0, T], L^2)$. We write

$$\begin{aligned}\iint_{[0,T] \times \mathbb{T}} w_n \bar{\varphi} &= \iint_{[0,T] \times \mathbb{T}} u_n e^{-\frac{i}{2}F_n} P_+ \bar{\varphi} \\ &= \iint_{[0,T] \times \mathbb{T}} u_n \left(e^{-\frac{i}{2}F_n} - 1 \right) P_+ \bar{\varphi} + \iint_{[0,T] \times \mathbb{T}} u_n P_+ \bar{\varphi}.\end{aligned}$$

The first term converges to 0 by strong convergence of $e^{-\frac{i}{2}F_n} - 1$ in $L^\infty([0, T], L^\infty)$. The second converges to 0 by weak convergence. The same proof without P_+ gives the result for $u_n e^{-\frac{i}{2}F_n}$. \square

Lemma 7.3. $c_n(t) = \int a(y)u_n(y, t)dy$ is bounded in $H^1([0, T])$ and converges strongly to 0 in $L^2([0, T])$.

Proof. We compute

$$\begin{aligned} \dot{c}_n(t) &= \int_{\mathbb{T}} a(y) \left[\mathcal{H}\partial_y^2 u_n(t, y) + \frac{1}{2}\partial_y(u_n(t, y)^2) + \mathcal{G}\mathcal{G}^*u_n \right] dy \\ &= - \int_{\mathbb{T}} \mathcal{H}\partial_y^2 a(y)u_n(t, y)dy - \frac{1}{2} \int_{\mathbb{T}} \partial_y a(y)(u_n(t, y)^2)dy \\ &\quad + \int_{\mathbb{T}} a(y)\mathcal{G}\mathcal{G}^*u_n dy \\ |\dot{c}_n(t)| &\leq C\|u_n(t)\|_{L^2(\mathbb{T})} + C\|u_n(t)\|_{L^2(\mathbb{T})}^2. \end{aligned}$$

So $\|\dot{c}_n(t)\|_{L^2([0, T])} \leq C\|u_n\|_{L^2([0, T], L^2)} + C\|u_n\|_{L^\infty([0, T], L^2)}^2 \leq C$. Lemma 7.3 follows by Sobolev embedding and weak convergence to 0 of u_n . \square

Remark 7.3. For Lemmas 7.2 and 7.3 the same argument applies with \mathcal{G} replaced by \mathcal{G}_μ since we only use the boundedness as an operator of $L^\infty([0, T], L^2)$.

Lemma 7.4. $\mathcal{G}u_n \rightarrow 0$ in $L^2([0, T] \times \mathbb{T})$ implies that $a(x)w_n \rightarrow 0$ in $L^2([0, T] \times \mathbb{T})$.

Proof. Lemma 7.3 implies that we have in fact $a(x)u_n \rightarrow 0$ in $L^2([0, T] \times \mathbb{T})$. Hence $a(x)e^{-\frac{i}{2}F_n}u_n \rightarrow 0$ in $L^2([0, T] \times \mathbb{T})$. Similarly, $P_+(a(x)e^{-\frac{i}{2}F_n}u_n) \rightarrow 0$ in $L^2([0, T] \times \mathbb{T})$.

We write

$$a(x)w_n = a(x)P_+ \left(u_n e^{-\frac{i}{2}F_n} \right) = P_+ \left(a(x)u_n e^{-\frac{i}{2}F_n} \right) + [a(x), P_+] \left(u_n e^{-\frac{i}{2}F_n} \right).$$

Thus, it remains to prove that the second term in the right hand side is strongly convergent in $L^2([0, T] \times \mathbb{T})$.

From Lemma 2.5 of Linares-Rosier [31], we get

$$\left\| [a(x), P_+] \left(u_n e^{-\frac{i}{2}F_n} \right) \right\|_{L^2([0, T] \times \mathbb{T})} \leq \|u_n e^{-\frac{i}{2}F_n}\|_{L^2([0, T], H^{-1})}.$$

Using Lemma 7.2, we get that $u_n e^{-\frac{i}{2}F_n} \rightarrow 0$ in $L^2([0, T] \times \mathbb{T})$. To apply Aubin-Lions' lemma, we just need to prove that $\partial_t(u_n e^{-\frac{i}{2}F_n})$ is bounded in some space $L^p([0, T], H^{-s})$, $s \in \mathbb{R}$. We also notice that $u_n e^{-\frac{i}{2}F_n} = 2i\partial_x e^{-\frac{i}{2}F_n}$, so, we just have to prove that $\partial_t e^{-\frac{i}{2}F_n} = -\frac{i}{2}(\partial_t F_n)e^{-\frac{i}{2}F_n}$ is bounded in some $L^p([0, T], H^{-s})$. However, we have shown in the proof of Lemma 7.2 that $\partial_t F_n$ was bounded in $L^\infty([0, T], H^{-1})$ and we notice easily that $e^{-\frac{i}{2}F_n}$ is bounded in $L^\infty([0, T], H^1)$, since $\partial_x e^{-\frac{i}{2}F_n} = -\frac{i}{2}u_n e^{-\frac{i}{2}F_n}$ is bounded in $L^\infty([0, T], L^2)$. Thus $\partial_t e^{-\frac{i}{2}F_n}$ is bounded in $L^\infty([0, T], H^{-1})$. \square

Remark 7.4. Lemma 7.4 has to be modified when \mathcal{G} is replaced by \mathcal{G}_μ . Indeed, in that case, the damping zone is moving at speed μ , but, we can use the infinite speed of propagation to get a similar result as follows.

If $\mathcal{G}_\mu \tilde{u}_n \rightarrow 0$ in $L^2([0, T] \times \mathbb{T})$, we infer as before that $a(x - t\mu)\tilde{u}_n \rightarrow 0$ in $L^2([0, T] \times \mathbb{T})$. If, for instance, $[-c, c] \subset \omega$ with $c > 0$, we easily conclude that there exists an ε (a priori depending on μ) such that $\tilde{u}_n \rightarrow 0$ in $L^2([0, \varepsilon] \times [-\varepsilon, \varepsilon])$.

This result will be enough for what follows since the time $T > 0$ and the open set ω are arbitrary (ω non empty).

Now, we borrow a propagation theorem from [26].

Theorem 7.1. *Let w_n be a sequence of solutions of*

$$i \partial_t w_n + \partial_x^2 w_n = f_n$$

such that for one $0 \leq b \leq 1$, we have

$$\|w_n\|_{X_T^{0,b}} \leq C, \quad \|w_n\|_{X_T^{-1+b,-b}} \rightarrow 0 \quad \text{and} \quad \|f_n\|_{X_T^{-1+b,-b}} \rightarrow 0.$$

Moreover, we assume that there is a non empty open set ω such that $w_n \rightarrow 0$ in $L^2([0, T], L^2(\omega))$.

Then $w_n \rightarrow 0$ in $L^2_{loc}([0, T], L^2(\mathbb{T}))$.

We want to apply this lemma with $b = 1/2$ and

$$\begin{aligned} f_n &= -\partial_x P_+ [W_n(P_- \partial_x u_n)] - \frac{i}{2} P_+ [g_n e^{-\frac{i}{2} F_n}] \\ &\quad - \frac{1}{4} P_+ \left[\left(-\frac{1}{2} P_0(u_n^2) + G_n \right) u_n e^{-\frac{i}{2} F_n} \right] \end{aligned}$$

with $g_n = \mathcal{G}\mathcal{G}^* u_n$, $G_n = \partial_x^{-1} g_n$. That is, we need to prove

$$\|w_n\|_{X_T^{0,\frac{1}{2}}} \leq C, \quad \|w_n\|_{X_T^{-\frac{1}{2},-\frac{1}{2}}} \rightarrow 0 \quad \text{and} \quad \|f_n\|_{X_T^{-\frac{1}{2},-\frac{1}{2}}} \rightarrow 0.$$

We decompose $f_n = f_n^a + f_n^b$ with $f_n^a = -\partial_x P_+ [W_n(P_- \partial_x u_n)]$. We know that

- $\|w_n\|_{X^{0,1/2}} \leq C$,
- $\|f_n\|_{X^{0,-1/2}} \leq C$.

Lemma 7.5. *For any $\theta \in [0, 1]$, we have the bound*

$$\|f_n\|_{X_T^{-2\theta, -\frac{1-\theta}{2}}} \leq C.$$

Proof. If we prove $\|f_n\|_{X_T^{-2,0}} = \|f_n\|_{L^2([0,T], H^{-2})} \leq C$, by interpolation, this gives

$$\|f_n\|_{X_T^{-2\theta, -\frac{1-\theta}{2}}} \leq C.$$

So, to bound f_n^a , we need to prove that $W_n(P_- \partial_x u_n)$ is bounded in $L^2([0, T], H^{-1})$.

We write

$$\begin{aligned} \|W_n(P_- \partial_x u_n)\|_{L^2([0,T], H^{-1})} &\leq C \|W_n\|_{L^2([0,T], H^1)} \|\partial_x u_n\|_{L^\infty([0,T], H^{-1})} \\ &\leq C \|w_n\|_{L^2([0,T], L^2)} \|u_n\|_{L^\infty([0,T], L^2)} \leq C \end{aligned}$$

where we have used that H^1 is an algebra and so by duality, $H^1 * H^{-1} \hookrightarrow H^{-1}$.

The bound for f_n^b is easier (and corresponds actually to the estimates of *II* and *III* in Section 3.1)

$$\begin{aligned} \|f_n^b\|_{L^2([0,T],H^{-2})} &\leq \|f_n^b\|_{L^2([0,T],L^2)} \leq \|g_n\|_{L^2([0,T],L^2)} \\ &+ \left\| -\frac{1}{2}P_0(u_n^2) + G_n \right\|_{L^\infty([0,T]\times\mathbb{T})} \|u_n\|_{L^2([0,T],L^2)} \\ &\leq \|g_n\|_{L^2([0,T],L^2)} + \left(\|u_n\|_{L^\infty([0,T],L^2)}^2 + \|g_n\|_{L^2([0,T],L^2)} \right) \|u_n\|_{L^2([0,T],L^2)}. \end{aligned}$$

□

With for instance $\theta = \frac{1}{5}$,

$$\|f_n\|_{X_T^{-\frac{2}{5},-\frac{4}{10}}} \leq C.$$

So, since $-\frac{1}{2} < -\frac{2}{5}$ and $-\frac{1}{2} < -\frac{4}{10}$, the embedding of $X_T^{-\frac{2}{5},-\frac{4}{10}} \hookrightarrow X_T^{-\frac{1}{2},-\frac{1}{2}}$ is compact and we can extract a subsequence and pick a function f such that

$$\|f_n - f\|_{X_T^{-\frac{1}{2},-\frac{1}{2}}} \rightarrow 0.$$

We can get that $f = 0$ using the equation verified by w_n , f_n and use the fact that we already know that $w_n \rightarrow 0$ in the distributional sense.

$$\|f_n\|_{X_T^{-\frac{1}{2},-\frac{1}{2}}} \rightarrow 0.$$

This constitutes the case $b = \frac{1}{2}$. The same holds for w_n .

We can now state the following theorem.

Theorem 7.2. *Under the same assumptions as in Lemma 7.1, we have $u_n \rightarrow 0$ in $L^2_{loc}([0, T], L^2(\mathbb{T}))$*

Proof. Using Theorem 7.1 and the previous analysis, we get $w_n \rightarrow 0$ in $L^2_{loc}([0, T], L^2(\mathbb{T}))$. By definition of w_n , we have $w_n = -\frac{i}{2}P_+(u_n e^{-\frac{i}{2}F_n})$. However, we have proved in Lemma 7.2 that $e^{-\frac{i}{2}F_n}$ converges strongly to 1 in $L^\infty([0, T], L^\infty)$, so we can write

$$P_+u_n = 2i w_n - P_+ \left[u_n \left(e^{-\frac{i}{2}F_n} - 1 \right) \right].$$

This gives $P_+u_n \rightarrow 0$ in $L^2_{loc}([0, T], L^2(\mathbb{T}))$ and the same result for u_n since u_n is real valued. □

7.2. Small Data

Lemma 7.6. *For $u_0 \in L^2(\mathbb{T})$, let u denote the solution of*

$$\begin{cases} \partial_t u + \mathcal{H}\partial_x^2 u = u\partial_x u - \mathcal{G}\mathcal{G}^*u, \\ u(0) = u_0, \end{cases}$$

and let u_L denote the solution of

$$\begin{cases} \partial_t u_L + \mathcal{H}\partial_x^2 u_L = -\mathcal{G}\mathcal{G}^* u_L \\ u_L(0) = u_0. \end{cases} \quad (7.1)$$

Then, there exist some constants $T_0 > 0$, $\varepsilon > 0$ and $C > 0$ such that for $T < T_0$ and $\|u_0\|_{L^2} \leq \varepsilon$, we have

$$\|u - u_L\|_{L^\infty([0,T],L^2)} \leq C \|u_0\|_{L^2}^2.$$

Proof. Let v denote the solution of

$$\begin{cases} \partial_t v + \mathcal{H}\partial_x^2 v = -\mathcal{G}\mathcal{G}^* u \\ v(0) = u_0. \end{cases} \quad (7.2)$$

From (4.2) applied with $g = -\mathcal{G}\mathcal{G}^* u$ and $\|u_0\|_{L^2}$ small enough, we have

$$\begin{aligned} \|u - v\|_{L^\infty([0,T],L^2)} &\lesssim \|\mathcal{G}\mathcal{G}^* u\|_{L^2([0,T],L^2)}^2 + \|u_0\|_{L^2}^2 \\ &\lesssim \|u\|_{L^2([0,T],L^2)}^2 + \|u_0\|_{L^2}^2 \\ &\lesssim C_1(T+1) \|u_0\|_{L^2}^2 \end{aligned} \quad (7.3)$$

for some constant $C_1 > 0$ where we used the fact that $\|u(t)\|_{L^2} \leq \|u_0\|_{L^2}$ for any $t \geq 0$.

On the other hand, by classical semigroup estimates we have that

$$\begin{aligned} \|v - u_L\|_{L^\infty([0,T],L^2)} &\lesssim \|\mathcal{G}\mathcal{G}^*(u - u_L)\|_{L^1([0,T],L^2)} \\ &\lesssim \|u - u_L\|_{L^1([0,T],L^2)} \\ &\leq C_2 T \|u - u_L\|_{L^\infty([0,T],L^2)}. \end{aligned} \quad (7.4)$$

Combining (7.3) and (7.4) and the triangle inequality yield

$$\|u - u_L\|_{L^\infty([0,T],L^2)} \leq C_1 \|u_0\|_{L^2}^2 + C_2 T \|u - u_L\|_{L^\infty([0,T],L^2)}.$$

Thus

$$\|u - u_L\|_{L^\infty([0,T],L^2)} \leq \frac{C_1 T}{1 - C_2 T} \|u_0\|_{L^2}^2$$

whenever $\|u_0\|_{L^2} < \varepsilon$ and $T < T_0 = \frac{1}{2C_2}$. \square

Proposition 7.1. Suppose $u_{n,0}$ is a sequence of smooth functions strongly convergent to 0 in L^2 , with associated solutions u_n of the problem

$$\begin{cases} \partial_t u_n + \mathcal{H}\partial_x^2 u_n = u_n \partial_x u_n - \mathcal{G}\mathcal{G}^* u_n \\ u_n(0) = u_{0,n}, \end{cases}$$

and in addition assume that

$$\|\mathcal{G}u_n\|_{L^2([0,T] \times \mathbb{T})} \leq \frac{1}{n} \|u_{0,n}\|_{L^2}. \quad (7.5)$$

Then, $u_{0,n} = 0$ for n large enough.

Proof. We denote $u_{n,L}$ the solution of

$$\begin{cases} \partial_t u_{n,L} + \mathcal{H} \partial_x^2 u_{n,L} = -\mathcal{G} \mathcal{G}^* u_{n,L} \\ u_{n,L}(0) = u_{0,n}. \end{cases}$$

By the observability of the linear damped system, proved in [29] Theorem 1.2, we know that for some T that can be chosen $T < T_0$, we have

$$\|u_{0,n}\|_{L^2}^2 \leq C \int_0^T \|\mathcal{G} u_{n,L}\|_{L^2}^2. \tag{7.6}$$

But, we have from triangular inequality, (7.5) and Lemma 7.6, that

$$\begin{aligned} \|\mathcal{G} u_{n,L}\|_{L^2([0,T] \times \mathbb{T})}^2 &\leq C \|\mathcal{G} u_n\|_{L^2([0,T] \times \mathbb{T})}^2 + C \|\mathcal{G}(u_{n,L} - u_n)\|_{L^2([0,T] \times \mathbb{T})}^2 \\ &\leq \frac{C}{n} \|u_{0,n}\|_{L^2}^2 + C \|u_{0,n}\|_{L^2}^4. \end{aligned}$$

Combining the previous estimate with (7.6) gives

$$\|u_{0,n}\|_{L^2}^2 \leq C \left(\frac{1}{n} + \|u_{0,n}\|_{L^2}^2 \right) \|u_{0,n}\|_{L^2}^2.$$

Since $\|u_{0,n}\|_{L^2}$ converges to zero, this gives $u_{0,n} = 0$ for n large enough. \square

8. Stabilization

In this section we will prove Theorem 1.2. We will use the argument in [31]. However to complete the arguments we have to make some modifications as we did in the previous section.

Because of the identity

$$\frac{1}{2} \|u(t)\|^2 + \|\mathcal{G} u\|_{L^2([0,T] \times \mathbb{T})}^2 = \frac{1}{2} \|u_0\|^2 \tag{8.1}$$

we observe that $\|u(t)\|$ is nonincreasing, so that the exponential decay is guaranteed if

$$\|u((n + 1)T)\| \leq \kappa \|u(nT)\| \quad \text{for some } \kappa < 1.$$

To prove the theorem, it is sufficient to show the following observability inequality: for any $T > 0$ and any $R \gg 1$ there exists a constant $C(R, T)$ such that for any $u_0 \in H_0^0(\mathbb{T})$ with $\|u_0\| \leq R$ it holds

$$\|u_0\|^2 \leq C \int_0^T \|\mathcal{G} u(t)\|^2 dt \tag{8.2}$$

where u denotes the solution of (1.12).

Suppose there exists a sequence $u_{0,n} \in H_0^0(\mathbb{T})$ such that for each n we have $\|u_{0,n}\| \leq R$ (where $R \gg 1$) but

$$\|u_{0,n}\| > n \int_0^T \|\mathcal{G}u_n(t)\|^2 dt. \tag{8.3}$$

Then from Theorem 7.2, we have for a subsequence (still denoted by u_n) that

$$u_n \rightarrow 0 \text{ in } L^2_{\text{loc}}([0, T], L^2(\mathbb{T})).$$

Thus one can select some $t_0 \in (0, T)$ such that, extracting again a subsequence,

$$u_n(t_0) \rightarrow 0 \text{ in } L^2(\mathbb{T}).$$

Since

$$\frac{1}{2} \|u_n(t_0)\|^2 + \|\mathcal{G}u_n\|^2_{L^2([0,t_0] \times \mathbb{T})} = \frac{1}{2} \|u_{0,n}\|^2,$$

we conclude that

$$\|u_{0,n}\|_{L^2} \rightarrow 0.$$

The assumptions of Proposition 7.1 are fulfilled. We finally get $u_{0,n} = 0$ for n large enough, which is a contradiction to (8.3).

Appendix

In the following, we will establish a slight modification of the bilinear estimate of MOLINET-PILOD [37], Proposition 3.5, with a gain of a power of T . Such an estimate will allow us to avoid the dilation argument for small time and large data and it could also be interesting for other purposes. This type of gain was already obtained by BOURGAIN in [7] for the KdV equation and it relies mainly on the fact that the equation we consider is subcritical on L^2 with respect to the scaling.

For sake of completeness we will recall the notation used in [37].

Let

$$I = \int_{\mathcal{D}} \frac{\xi}{\langle x\sigma \rangle^{1/2}} \widehat{h}(\xi, \tau) \xi_1^{-1} \widehat{w}(\xi_1, \tau_1) \xi_2 \widehat{u}(\xi_2, \tau_2) dv \tag{A.1}$$

and

$$J = \sum_N \int_{\mathcal{D}} \frac{\xi}{\langle x\sigma \rangle^{1/2}} g_N(\xi, \tau) \phi_N(\xi) \xi_1^{-1} \widehat{w}(\xi_1, \tau_1) \xi_2 \widehat{u}(\xi_2, \tau_2) dv,$$

where $dv = d\xi d\xi_1 d\tau d\tau_1$, $\xi_2 = \xi - \xi_1$, $\tau_2 = \tau - \tau_1$, $\sigma = \tau - \xi|\xi|$ and $\sigma_i = \tau_i - \xi_i|\xi_i|$, $i = 1, 2$ and

$$\mathcal{D} = \{(\xi, \xi_1, \tau, \tau_1) \in \mathbb{R} : \xi > 1, \xi_1 \geq 1 \text{ and } \xi_2 \leq 0\}.$$

We recall that the measures $d\xi$ and $d\xi_1$ refer to the counting measure (see (5.3)).

By using a dyadic decomposition in space-frequency for the functions in the integrals (A.1), one can write

$$I = \sum_{N, N_1, N_2} I_{N, N_1, N_2} \quad \text{and} \quad J = \sum_{N, N_1, N_2} J_{N, N_1, N_2}$$

where

$$I_{N,N_1,N_2} = \int_{\mathcal{D}} \frac{\xi}{\langle x\sigma \rangle^{1/2}} \widehat{P_N h}(\xi, \tau) \xi_1^{-1} \widehat{P_{N_1} w}(\xi_1, \tau_1) \xi_2 \widehat{P_{N_2} u}(\xi_2, \tau_2) \, dv$$

and

$$J_{N,N_1,N_2} = \sum_N \int_{\mathcal{D}} \frac{\xi}{\langle x\sigma \rangle^{1/2}} g_N(\xi, \tau) \phi_N(\xi) \xi_1^{-1} \widehat{P_{N_1} w}(\xi_1, \tau_1) \xi_2 \widehat{P_{N_2} u}(\xi_2, \tau_2) \, dv.$$

Let us now consider

$$\begin{aligned} A_{N,N_2} &= \left\{ (\xi, \xi_1, \tau, \tau_1) \in \mathcal{D} : |\sigma| \geq \frac{1}{6} N N_2 \right\} \\ B_{N,N_2} &= \left\{ (\xi, \xi_1, \tau, \tau_1) \in \mathcal{D} : |\sigma_1| \geq \frac{1}{6} N N_2, |\sigma| \leq \frac{1}{6} N N_2 \right\} \\ C_{N,N_2} &= \left\{ (\xi, \xi_1, \tau, \tau_1) \in \mathcal{D} : |\sigma| < \frac{1}{6} N N_2, |\sigma_1| < \frac{1}{6} N N_2, |\sigma_2| \geq \frac{1}{6} N N_2 \right\}. \end{aligned} \tag{A.2}$$

Define by

$$\begin{aligned} I_A &= \sum_{N,N_1,N_2} I_{N,N_1,N_2}^{A,N,N_2}, \quad I_B = \sum_{N,N_1,N_2} I_{N,N_1,N_2}^{B,N,N_2}, \quad \text{and} \quad I_C = \sum_{N,N_1,N_2} I_{N,N_1,N_2}^{C,N,N_2}, \\ J_A &= \sum_{N,N_1,N_2} J_{N,N_1,N_2}^{A,N,N_2}, \quad J_B = \sum_{N,N_1,N_2} J_{N,N_1,N_2}^{B,N,N_2}, \quad \text{and} \quad J_C = \sum_{N,N_1,N_2} J_{N,N_1,N_2}^{C,N,N_2}, \end{aligned}$$

where I_{N,N_1,N_2}^{A,N,N_2} , I_{N,N_1,N_2}^{B,N,N_2} and I_{N,N_1,N_2}^{C,N,N_2} denote the restriction of I_{N,N_1,N_2} to the regions in (A.2). Similarly, J_{N,N_1,N_2}^{A,N,N_2} , J_{N,N_1,N_2}^{B,N,N_2} and J_{N,N_1,N_2}^{C,N,N_2} for J_{N,N_1,N_2} .

Now we are ready to establish the new estimates.

Lemma A.1. *We have for $0 \leq T \leq 1$ the estimate*

$$\left\| \partial_x P_+ [W(P_- \partial_x u)] \right\|_{X_T^{0,-1/2}} \leq CT^{1/8} \|(u, w)\|_X^2.$$

Proof. Let h, u, w be some extension (same notation by abuse). We use the same notation as in [37] and continue the computations started in [37]. We continue the estimates of I_A starting from estimate (3.32) in [37]:

$$|I_A| \leq \|h\|_{L^2} \|w\|_{X_T^{0,3/8}} \|u\|_{L^4} \leq CT^{1/2-3/8} \|h\|_{L^2} \|w\|_{X_T^{0,1/2}} \|u\|_{L^4}.$$

For I_B , we use (3.33) and (3.34) of [37] to get

$$\begin{aligned} |I_B| &\leq \left(\sum_{N_1} \left\| P_{N_1} \left(\frac{\widehat{h}}{\langle \sigma \rangle^{1/2}} \right)^\vee \right\|_{L^4}^2 \right)^{1/2} \|w\|_{X_T^{0,1/2}} \|u\|_{L^4} \\ &\leq CT^{1/2-3/8} \left\| \left(\frac{\widehat{h}}{\langle \sigma \rangle^{1/2}} \right)^\vee \right\|_{X_T^{0,1/2}} \|h\|_{L^2} \|w\|_{X_T^{0,3/8}} \|u\|_{L^4} \\ &\leq CT^{1/2-3/8} \|h\|_{L^2} \|w\|_{X_T^{0,3/8}} \|u\|_{L^4}, \end{aligned}$$

and using (3.38)

$$\begin{aligned} |I_C| &\leq \left(\sum_{N_1} \left\| P_{N_1} \left(\frac{\widehat{h}}{\langle \sigma \rangle^{1/2}} \right)^\vee \right\|_{L^4}^2 \right)^{1/2} \|w\|_{X_T^{0,3/8}} \|u\|_{X_T^{-1,1}} \\ &\leq CT^{1/2-3/8} \|h\|_{L^2} \|w\|_{X_T^{0,1/2}} \|u\|_{X_T^{-1,1}}. \end{aligned}$$

□

Lemma A.2. *Let $\varepsilon > 0$. The following estimate holds uniformly for $0 \leq T \leq 1$:*

$$\|\partial_x P_+[W(P_-\partial_x u)]\|_{\widetilde{Z}_T^{0,-1}} \leq CT^{1/8-\varepsilon} \|(u, w)\|_X^2.$$

Proof. We can see using (3.43) of [37] that

$$\begin{aligned} |J_A| &\leq \left(\sum_N \|g_N\|_{L_\xi^2 L_\tau^\infty}^2 \right)^{1/2} \|w\|_{X_T^{0,3/8}} \|u\|_{L^4} \\ &\leq CT^{1/2-3/8} \left(\sum_N \|g_N\|_{L_\xi^2 L_\tau^\infty}^2 \right)^{1/2} \|w\|_{X_T^{0,1/2}} \|u\|_{L^4}, \end{aligned}$$

and using estimate line 15 p 381 of [37] that

$$|J_B| + |J_C| \leq \left(\left\| \left(\frac{g}{\langle \sigma \rangle} \right)^\vee \right\|_{\widetilde{L}^4} + \left\| \left(\frac{|g|}{\langle \sigma \rangle} \right)^\vee \right\|_{\widetilde{L}^4} \right) \|w\|_{X_T^{0,1/2}} (\|u\|_{L^4} + \|u\|_{X_T^{-1,1}}).$$

However

$$\begin{aligned} \left\| \left(\frac{g}{\langle \sigma \rangle} \right)^\vee \right\|_{\widetilde{L}^4} + \left\| \left(\frac{|g|}{\langle \sigma \rangle} \right)^\vee \right\|_{\widetilde{L}^4} &\leq CT^{1/2-3/8-\varepsilon} \left\| \left(\frac{|g|}{\langle \sigma \rangle} \right)^\vee \right\|_{X_T^{0,1/2-\varepsilon}} \\ &\leq CT^{1/8-\varepsilon} \left\| \frac{|g|}{\langle \sigma \rangle^{1/2+\varepsilon}} \right\|_{L_\tau^2 L_\xi^2} \\ &\leq CT^{1/8-\varepsilon} \left(\sum_N \|\langle \sigma \rangle^{-1/2-\varepsilon} g_N\|_{L_\tau^2 L_\xi^2}^2 \right)^{1/2} \\ &\leq CT^{1/8-\varepsilon} \left(\sum_N \int_\xi \int_\tau \langle \tau + \xi |\xi| \rangle^{-1-2\varepsilon} |g_N(\tau, \xi)|^2 \right)^{1/2} \\ &\leq CT^{1/8-\varepsilon} \left(\sum_N \|g_N\|_{L_\xi^2 L_\tau^\infty}^2 \right)^{1/2}, \end{aligned}$$

which gives

$$|J_B| + |J_C| \leq CT^{1/8-\varepsilon} \left(\sum_N \|g_N\|_{L_\xi^2 L_\tau^\infty}^2 \right)^{1/2} \|w\|_{X_T^{0,1/2}} (\|u\|_{L^4} + \|u\|_{X_T^{-1,1}}).$$

The result follows by duality. □

Note that the two previous lemmas give Lemma 2.8.

Lemma A.3. *Let $N \in \mathbb{N}^*$. We have the following estimates, uniformly in N .*

$$\begin{aligned} \|P_{\geq N} \left(e^{i\frac{F}{2}} \right)\|_{L^\infty} &\leq \frac{C}{\sqrt{N}} \|F\|_{H^1}, \\ \|P_{\geq N} \left(e^{i\frac{F_1}{2}} - e^{i\frac{F_2}{2}} \right)\|_{L^\infty} &\leq \frac{C}{\sqrt{N}} [1 + \|F_1\|_{H^1} + \|F_2\|_{H^1}] \|F_1 - F_2\|_{H^1}, \end{aligned}$$

and

$$\|e^{i\frac{Q_N F_1}{2}} - e^{i\frac{Q_N F_2}{2}}\|_{L^\infty} \leq \frac{C}{\sqrt{N}} [1 + \|F_1\|_{H^1} + \|F_2\|_{H^1}] \|F_1 - F_2\|_{H^1}.$$

Remark A.1. Note that these estimates are very close to the estimate

$$\|P_{\geq 1} \left(e^{i\frac{F}{2}} \right)\|_{L^\infty((0,\lambda))} \leq C \|F_x\|_{L^2((0,\lambda))}$$

but uniform on the length λ , as used by MOLINET [34].

This gain comes from the fact that the inequality is subcritical. Roughly speaking, L^∞ scales like $H^{1/2}$ which gives a gain at high frequency.

Proof. We will mainly use the following estimate, which is only the Sobolev embedding at high frequency:

$$\|P_{\geq N}(f)\|_{L^\infty} \leq \sum_{n=N}^{+\infty} |\hat{f}(n)| \leq \left(\sum_{n=N}^{+\infty} \frac{1}{n^2} \right)^{1/2} \left(\sum_{n=N}^{+\infty} n^2 |\hat{f}(n)|^2 \right)^{1/2} \leq \frac{C}{\sqrt{N}} \|f\|_{H^1}. \tag{A.3}$$

We apply this estimate to $f = e^{i\frac{F_1}{2}} - e^{i\frac{F_2}{2}}$:

$$\begin{aligned} &\|P_{\geq N} \left(e^{i\frac{F_1}{2}} - e^{i\frac{F_2}{2}} \right)\|_{L^\infty} \\ &\leq \frac{C}{\sqrt{N}} \|e^{i\frac{F_1}{2}} - e^{i\frac{F_2}{2}}\|_{H^1} \leq \frac{C}{\sqrt{N}} \left(\|e^{i\frac{F_1}{2}} - e^{i\frac{F_2}{2}}\|_{L^2} + \|(\partial_x F_1)e^{i\frac{F_1}{2}} - (\partial_x F_2)e^{i\frac{F_2}{2}}\|_{L^2} \right) \\ &\leq \frac{C}{\sqrt{N}} (\|F_1 - F_2\|_{L^2} + \|F_1\|_{H^1} \min(1, \|F_1 - F_2\|_{H^1}) + \|F_2\|_{H^1} \|F_1 - F_2\|_{H^1}), \end{aligned}$$

where, for the last estimate, we have used estimate (2.6) together with

$$(\partial_x F_1)e^{i\frac{F_1}{2}} - (\partial_x F_2)e^{i\frac{F_2}{2}} = (\partial_x F_1) \left(e^{i\frac{F_1}{2}} - e^{i\frac{F_2}{2}} \right) + e^{i\frac{F_2}{2}} \partial_x (F_1 - F_2).$$

This proves the second inequality of the lemma. The first estimate of the lemma is a consequence of the previous inequality with $F_2 = 0$ using the fact that $P_{\geq N} \left(e^{i\frac{F}{2}} \right) = P_{\geq N} \left(e^{i\frac{F}{2}} - 1 \right)$.

The last inequality is proved using the Gagliardo-Nirenberg inequality on \mathbb{T} , that is

$$\left\| e^{i\frac{Q_N F_1}{2}} - e^{i\frac{Q_N F_2}{2}} \right\|_{L^\infty} \leq \left\| e^{i\frac{Q_N F_1}{2}} - e^{i\frac{Q_N F_2}{2}} \right\|_{L^2} + \left\| e^{i\frac{Q_N F_1}{2}} - e^{i\frac{Q_N F_2}{2}} \right\|_{H^1}^{1/2} \left\| e^{i\frac{Q_N F_1}{2}} - e^{i\frac{Q_N F_2}{2}} \right\|_{L^2}^{1/2}.$$

The previous computation shows that

$$\begin{aligned} \left\| e^{i\frac{Q_N F_1}{2}} - e^{i\frac{Q_N F_2}{2}} \right\|_{H^1} &\leq C \left[1 + \|Q_N F_1\|_{H^1} + \|Q_N F_2\|_{H^1} \right] \|Q_N F_1 - Q_N F_2\|_{H^1} \\ &\leq C \left[1 + \|F_1\|_{H^1} + \|F_2\|_{H^1} \right] \|F_1 - F_2\|_{H^1}, \end{aligned}$$

but, the mean value theorem gives

$$\left\| e^{i\frac{Q_N F_1}{2}} - e^{i\frac{Q_N F_2}{2}} \right\|_{L^2} \leq \|Q_N(F_1 - F_2)\|_{L^2} \leq \frac{C}{N} \|F_1 - F_2\|_{H^1}.$$

□

Acknowledgments. CL was partially supported by the Agence Nationale de la Recherche (ANR), Projet Blanc EMAQS number ANR-2011-BS01-017-01. CL would also like to thank IMPA for its hospitality. FL was partially supported by FAPERJ and CNPq Brazil. LR was partially supported by the Agence Nationale de la Recherche (ANR), Project CISIFS, grant ANR-09-BLAN-0213-02.

References

1. ABDELOUHAB, L., BONA, J.L., FELLAND, M., SAUT, J.-C.: Nonlocal models for nonlinear dispersive waves. *Phys. D* **40**, 360–392 (1989)
2. ABLOWITZ, M.J., FOKAS, A.S.: The inverse scattering transform for the Benjamin-Ono equation—a pivot to multidimensional problems. *Stud. Appl. Math.* **68**, 1–10 (1983)
3. AMICK, C., TOLAND, J.: Uniqueness and related analytic properties for the Benjamin-Ono equation—a nonlinear Neumann problem in the plane. *Acta Math.*, **167**, 107–126 (1991)
4. BONA, J.L., SMITH, R.: The initial value problem for the Korteweg-de Vries equation. *Philos. Trans. R. Soc. Lond., Ser. A* **278**, 555–601 (1975)
5. BAHOURI, H., GÉRARD, P.: High frequency approximation of critical nonlinear wave equations. *Am. J. Math.* **121**, 131–175 (1999)
6. BENJAMIN, T.B.: Internal waves of permanent form in fluids of great depth. *J. Fluid Mech.* **29**, 559–592 (1967)
7. BOURGAIN, J.: Fourier transform restriction phenomena for certain lattice subsets and applications to nonlinear evolution equations, Part II: the KdV equation. *Geom. Funct. Anal.*, **3**, 209–262 (1993)
8. BURQ, N., PLANCHON, F.: On the well-posedness for the Benjamin-Ono equation. *Math. Ann.* **340**, 497–542 (2008)
9. CASE, K.M.: Benjamin-Ono related equations and their solutions. *Proc. Nat. Acad. Sci. USA* **76**, 1–3 (1979)
10. COIFMAN, R., WICKERHAUSER, M.: The scattering transform for the Benjamin-Ono equation. *Inverse Probl.* **6**, 825–860 (1990)
11. DAWSON, L., MCGAHAGAN, H., PONCE, G.: On the decay properties of solutions to a class of Schrödinger equations. *Proc. Amer. Math. Soc.* **136**, 2081–2090 (2007)

12. DEHMAN, B., GÉRARD, P., LEBEAU, G.: Stabilization and control for the nonlinear Schrödinger equation on a compact surface. *Math. Z* **254**, 729–749 (2006)
13. DEHMAN, B., LEBEAU, G., ZUAZUA, E.: *Stabilization and control for the subcritical semilinear wave equation*. *Ann. Sci. de l'École Normale Supérieure* **36**, 525–551 (2003)
14. FONSECA, G., LINARES, F.: Benjamin-Ono equation with unbounded data. *J. Math. Anal. Appl.* **247**, 426–447 (2000)
15. FONSECA, G., PONCE, G.: The I.V.P for the Benjamin-Ono equation in weighted Sobolev spaces. *J. Funct. Anal.* **260**, 436–459 (2011)
16. FONSECA, G., LINARES, F., PONCE, G.: The I.V.P for the Benjamin-Ono equation in weighted Sobolev spaces II. *J. Funct. Anal.* **262**, 2031–2049 (2012)
17. GINIBRE, J., TSUTSUMI, Y., VELO, G.: On the Cauchy problem for the Zakharov system. *J. Funct. Anal.* **151**(2), 384–436 (1997)
18. IONESCU, A.D., KENIG, C.E.: Global well-posedness of the Benjamin-Ono equation on low-regularity spaces. *J. Amer. Math. Soc.* **20**, 753–798 (2007)
19. IONESCU, A.D., KENIG, C.E.: Local and global well-posedness of periodic KP-I equations, *Mathematical Aspects of Nonlinear Dispersive Equations*, pp. 181–212. Princeton University Press (2007)
20. IORIO, R.J.: On the Cauchy problem for the Benjamin-Ono equation. *Comm. Partial Differ. Equations* **11**, 1031–1081 (1986)
21. IORIO, R., IORIO, V.M.: *Fourier analysis and partial differential equations*. Cambridge Stud. Adv. Math., **70** (2001)
22. KATO, T., PONCE, G.: Commutator estimates and the Euler and Navier-Stokes equations. *Comm. Pure Appl. Math.* **41**, 891–907 (1988)
23. KENIG, C.E., KOENIG, K.D.: On the local well-posedness of the Benjamin-Ono and modified Benjamin-Ono equations. *Math. Res. Letters* **10**, 879–895 (2003)
24. KENIG, C.E., PILOD, D.: Well-posedness for the fifth-order KdV equation in the energy space, to appear in *transactions AMS*
25. KOCH, H., TZVETKOV, N.: On the local well-posedness of the Benjamin-Ono equation on $H^s(\mathbb{R})$. *Int. Math. Res. Not.* **26**, 1449–1464 (2003)
26. LAURENT, C.: Global controllability and stabilization for the nonlinear Schrödinger equation on an interval. *ESAIM Control Optim. Calc. Var.* **16**(2), 356–379 (2010)
27. LAURENT, C.: Global controllability and stabilization for the nonlinear Schrödinger equation on some compact manifolds of dimension 3. *SIAM J. Math. Anal.* **42**(2), 785–832 (2010)
28. LAURENT, C., ROSIER, L., ZHANG, B.Y.: Control and stabilization of the Korteweg-de Vries equation on a periodic domain. *Comm. Partial Differ. Equations* **35**(4), 707–744 (2010)
29. LINARES, F., ORTEGA, J.H.: On the controllability and stabilization of the linearized Benjamin-Ono equation. *ESAIM Control Optim. Calc. Var.* **11**, 204–218 (2005)
30. LINARES, F., PAZOTO, A.F.: On the exponential decay of the critical generalized Korteweg-de Vries equation with localized damping. *Proc. Amer. Math. Soc.* **135**(5), 1515–1522 (2007)
31. LINARES, F., ROSIER, L.: Control and Stabilization for the Benjamin-Ono equation. *Trans. Amer. Math. Soc.* **367**(7), 4595–4626 (2015)
32. LIONS, J.L.: *Contrôlabilité Exacte, perturbations et stabilisation de Systèmes Distribués. Tome 1,2*, RMA **8**(9) (1988)
33. MICU, S., ORTEGA, J.H., ROSIER, L., ZHANG, B.Y.: Control and stabilization of a family of Boussinesq systems. *Discrete Cont. Dyn. Syst.* **24**(2), 273–313 (2009)
34. MOLINET, L.: Global well-posedness in L^2 for the periodic Benjamin-Ono equation. *Amer. J. Math.* **130**(3), 635–683 (2008)
35. MOLINET, L.: On ill-posedness for the one-dimensional periodic cubic Schrödinger equation. *Math. Res. Lett.* **16**(1), 111–120 (2009)
36. MOLINET, L.: Sharp ill-posedness result for the periodic Benjamin-Ono equation. *J. Funct. Anal.* **257**(11), 3488–3516 (2009)

37. MOLINET, L., PILOD, D.: The Cauchy problem for the Benjamin-Ono equation in L^2 revisited. *Anal. PDE* **5**(2), 365–395 (2012)
38. MOLINET, L., RIBAUD, F.: Well-posedness in H^1 for generalized Benjamin-Ono equations on the circle. *Discrete Contin. Dyn. Syst.* **23**(4), 1295–1311 (2009)
39. MOLINET, L., SAUT, J.C., TZVETKOV, N.: Ill-posedness issues for the Benjamin-Ono and related equations. *SIAM J. Math. Anal.* **33**, 982–988 (2001)
40. PONCE, G.: On the global well-posedness of the Benjamin-Ono equation. *Diff. Int. Eqs.* **4**, 527–542 (1991)
41. ONO, H.: Algebraic solitary waves in stratified fluids. *J. Phys. Soc. Japan* **39**, 1082–1091 (1975)
42. PAZOTO, A.: Unique continuation and decay for the Korteweg-de Vries equation with localized damping. *ESAIM Control Optim. Calc. Var.* **11**(3), 473–486 (2005)
43. PERLA MENZALA, G., VASCONCELLOS, C.F., ZUAZUA, E.: Stabilization of the Korteweg-de Vries equation with localized damping. *Quart. Appl. Math.* **60**(1), 111–129 (2002)
44. ROSIER, L., ZHANG, B.Y.: Local exact controllability and stabilizability of the nonlinear Schrödinger equation on a bounded interval. *SIAM J. Control Optim.* **48**(2), 972–992 (2009)
45. ROSIER, L., ZHANG, B.Y.: Exact boundary controllability of the nonlinear Schrödinger equation. *J. Differ. Equations* **246**(10), 4129–4153 (2009)
46. ROSIER, L., ZHANG, B.Y.: Control and stabilization of the Korteweg-de Vries equation: recent progresses. *J. Syst. Sci. Complex.* **22**, 647–682 (2009)
47. ROSIER, L., ZHANG, B.Y.: Control and stabilization of the nonlinear Schrödinger equation on rectangles. *M3AS Math. Models Methods Appl. Sci.* **20**(12), 2293–2347 (2010)
48. ROSIER, L., ZHANG, B.-Y.: Unique continuation property and control for the Benjamin-Bona-Mahony equation on a periodic domain. *J. Differ. Equations* **254**(1), 141–178 (2013)
49. RUSSELL, D.L., ZHANG, B.Y.: Exact controllability and stabilizability of the Korteweg-de Vries equation. *Trans. Amer. Math. Soc.* **348**, 3643–3672 (1996)
50. SAUT, J.-C.: Sur quelques généralisations de l'équation de Korteweg-de Vries. *J. Math. Pures Appl.* **58**, 21–61 (1979)
51. TAO, T.: Global well-posedness of the Benjamin-Ono equation in $H^1(\mathbb{R})$. *J. Hyperbolic Differ. Equ.* **1**, 27–49 (2004)

Laboratoire Jacques-Louis Lions,
CNRS, Sorbonne Universités, UPMC Univ Paris 06, UMR 7598
Boîte courrier 187, 75252,
Paris Cedex 05, France.
e-mail: camille.laurent@upmc.fr

and

Instituto de Matematica Pura e Aplicada,
Estrada Dona Castorina 110,
Rio de Janeiro, 22460-320, Brazil.
e-mail: linares@impa.br

and

Centre Automatique et Systèmes,
MINES ParisTech, PSL Research University,
60 boulevard Saint-Michel, 75272,
Paris Cedex 06, France.
e-mail: lionel.rosier@mines-paristech.fr

(Received October 20, 2014 / Accepted May 6, 2015)

Published online May 20, 2015 – © Springer-Verlag Berlin Heidelberg (2015)