



Metastability of Kolmogorov Flows and Inviscid Damping of Shear Flows

ZHIWU LIN  & MING XU

Communicated by P. CONSTANTIN

Abstract

First, we consider Kolmogorov flow (a shear flow with a sinusoidal velocity profile) for 2D Navier–Stokes equation on a torus. Such flows, also called bar states, have been numerically observed as one type of metastable state in the study of 2D turbulence. For both rectangular and square tori, we prove that the non-shear part of perturbations near Kolmogorov flow decays in a time scale much shorter than the viscous time scale. The results are obtained for both the linearized NS equations with any initial vorticity in L^2 , and the nonlinear NS equation with initial L^2 norm of vorticity of the size of viscosity. In the proof, we use the Hamiltonian structure of the linearized Euler equation and the RAGE theorem to control the low frequency part of the perturbation. Second, we consider two classes of shear flows for which a sharp stability criterion is known. We show the inviscid damping in a time average sense for non-shear perturbations with initial vorticity in L^2 . For the unstable case, the inviscid damping is proved on the center space. Our proof again uses the Hamiltonian structure of the linearized Euler equation and an instability index theory recently developed by Lin and Zeng for Hamiltonian PDEs.

1. Introduction

Consider a 2D Navier–Stokes (NS) equation

$$\partial_t U + U \cdot \nabla U - \nu \Delta U = -\nabla P \quad (1.1)$$

on a torus

$$\mathbb{T}_\alpha = \left\{ 0 < y < 2\pi, 0 < x < \frac{2\pi}{\alpha} \right\}, \quad \alpha > 0,$$

with the incompressible condition $\nabla \cdot U = 0$, where $U = (u, v)$ is the fluid velocity and $\nu > 0$ is the viscosity. More precisely, we impose the periodic boundary conditions

$$U(0, y, t) = U(2\pi/\alpha, y, t), \quad U(x, 0, t) = U(x, 2\pi, t).$$

The vorticity form of NS equation (1.1) is

$$\omega_t + u\omega_x + v\omega_y - \nu \Delta \omega = 0, \quad \omega = v_x - u_y. \tag{1.2}$$

It is convenient to introduce the stream function ψ such that $\omega = -\Delta \psi$ and $U = \nabla^\perp \psi = (\psi_y, -\psi_x)$.

In the numerical and experimental study of 2D turbulence, it was often observed [6, 18, 23] that the solutions to the two-dimensional Navier–Stokes (NS) equation with small viscosity rapidly approach certain long-lived coherent structures. Evidence also suggested that these quasi-stationary, or metastable, solutions are closely related to stationary solutions of the inviscid Euler equations

$$\omega_t + u\omega_x + v\omega_y = 0, \quad \omega = v_x - u_y.$$

Since there is no forcing in (1.2), when $t \rightarrow \infty$, $\|\omega(t)\|_{L^2} \rightarrow 0$ in the viscous time scale $O(\frac{1}{\nu})$, where $\omega(t)$ is the solution of (1.2) with initial data $\omega(0) \in L^2$. We are interested in the dynamics of (1.2), particularly the appearance and persistence of coherent states in the intermediate time scale $(0, T)$, where $1 \ll T \ll O(\frac{1}{\nu})$. The first step is to prove that nearby solutions converge rapidly to these coherent states in a time scale $T \ll O(\frac{1}{\nu})$. Such a metastability problem is also called enhanced damping in the literature. Among the candidates of Euler steady solutions to explain the coherent structures, some authors (e.g. [6, 18, 23]) suggested that certain maximal entropy solutions of the inviscid Euler equation are the most probable quasi-stationary states that one would observe. The simplest of such maximal entropy solutions is the Kolmogorov flow (also called bar states in [23]), that is, $u_0 = (\sin y, 0)$ or $(\cos y, 0)$. The solution to (NS) with initial data u_0 is $u^v(t, y) = e^{-\nu t} (\sin y, 0)$. The linearized (NS) equation near u^v is

$$\partial_t \omega = \nu \Delta \omega - e^{-\nu t} \left[\sin y \partial_x \left(1 + \Delta^{-1} \right) \right] \omega = L(t) \omega, \tag{1.3}$$

where ν is the viscosity and ω is the vorticity perturbation. In [3], Beck and Wayne studied the following approximation of the linearized problem:

$$\partial_t \omega = \nu \Delta \omega - e^{-\nu t} \sin y \partial_x \omega = \tilde{L}(t) \omega, \tag{1.4}$$

by dropping the nonlocal term $e^{-\nu t} \sin y \partial_x \Delta^{-1} \omega$ in (1.3). Define the following weighed H^1 space for non-shear vorticity functions:

$$Z = \left\{ \sum_{k \neq 0} \omega = \omega_k(y) e^{ikx} \in L^2, \right. \tag{1.5}$$

$$\left. \|\omega\|_Z^2 := \sum_{k \neq 0} \left[\|\omega_k\|_2^2 + \sqrt{\frac{\nu}{|k|}} \|\partial_y \omega_k\|_2^2 + \frac{1}{\sqrt{\nu} |k|^{\frac{3}{2}}} \|C^k \omega_k\|_2^2 \right] < \infty \right\},$$

where $C^k \omega_k = -ik e^{\nu t} (\cos y) \omega_k$. It was proved in [3] that for any $\tau > 0$ and $T \in [0, \frac{\tau}{\nu}]$ there exist constants K, M such that if ν is small enough, then the solution to (1.4) with the initial data $\omega(0) \in Z$ satisfies the estimate

$$\|\omega(t)\|_Z^2 \leq K e^{-M\sqrt{\nu}t} \|\omega(0)\|_Z^2, \quad t \in [0, T].$$

The proof of this used Villani’s hypocoercivity method ([22]). For the full linearized NS equation, numerical evidence in [3] suggested the same decay rate $O(e^{-\sqrt{\nu}t})$.

In this paper, we study the full linearized equation (1.3) and the nonlinear equation (1.1) on a torus

$$\mathbb{T}_\alpha = \left\{ 0 < y < 2\pi, 0 < x < \frac{2\pi}{\alpha} \right\},$$

with $\alpha \geq 1$, which is the sharp stability condition of Kolmogorov flows for the 2D Euler equation (see Lemma 4.1 of [14]). To simplify notation, we use $\omega^\nu(t)$ for both solutions of the linearized Navier–Stokes equation (1.3) and the perturbation solutions of the nonlinear Navier–Stokes equation (3.1) near Kolmogorov flows. Our first result is about the enhanced damping for the linearized NS equation.

Theorem 1.1. *Consider the linearized NS equation (1.3) on \mathbb{T}_α with $\alpha \geq 1$. Define the non-shear vorticity space*

$$X = \left\{ \omega \in L^2 \mid \omega = \sum_{0 \neq k \in \mathbf{Z}} \omega_k(y) e^{ik\alpha x} \right\}. \tag{1.6}$$

(i) *Then (Rectangular torus) Consider $\alpha > 1$. For any $\tau > 0$ and $\delta > 0$, if ν is small enough, then the solution $\omega^\nu(t)$ to (1.3) with non-shear initial data $\omega^\nu(0) \in X$ satisfies $\|\omega^\nu(\frac{\tau}{\nu})\|_{L^2} < \delta \|\omega^\nu(0)\|_{L^2}$.*

(ii) *(Square torus) Consider $\alpha = 1$. Let P_a be the orthogonal projection from the non-shear space*

$$X = \left\{ \omega \in L^2 \mid \omega = \sum_{k \neq 0} \omega_k(y) e^{ikx} \right\}$$

to the ‘anomalous’ space W_a spanned by $\{\cos x, \sin x\}$. For any $\tau > 0$ and $\delta > 0$, if ν is small enough, then the solution $\omega^\nu(t)$ to (1.3) with initial data $\omega^\nu(0) \in X$ satisfies

$$\|(I - P_a) \omega^\nu\left(\frac{\tau}{\nu}\right)\|_{L^2} < \delta \|(I - P_a) \omega^\nu(0)\|_{L^2}. \tag{1.7}$$

Since τ can be arbitrarily small, the above result implies a much enhanced decay in the time scale $O(\frac{\tau}{\nu})$ compared with the viscous time scale $O(\frac{1}{\nu})$. For shear initial data $\omega(0) = \omega_0(y)$, the linearized NS equation (1.3) is reduced to the heat equation $\partial_t \omega = \nu \partial_{yy} \omega$ and there is no enhanced decay. On the square torus, there is a two-dimensional additional kernel space W_a of the operator $1 + \Delta^{-1}$ spanned by $\{\cos x, \sin x\}$, which corresponds to exact solutions $e^{-\nu t} \{\cos x, \sin x\}$ of the Navier–Stokes equations. These so called ‘anomalous modes’ (see [3]) need to be removed for the enhanced damping to hold true.

For the nonlinear Navier–Stokes equation, we have

Theorem 1.2. Consider the nonlinear NS equation (1.1) on \mathbb{T}_α with $\alpha \geq 1$. Denote $P_{\mathcal{K}}$ to be the projection of $L^2(\mathbb{T}_\alpha)$ to the subspace of Kolmogorov flows $W_{\mathcal{K}} = \text{span}\{\cos y, \sin y\}$. Then,

(i) (Rectangular torus) Suppose $\alpha > 1$. There exist $d > 0$, such that for any $\tau > 0$ and $\delta > 0$, if ν is small enough, then for any solution $\omega^\nu(t)$ to (1.1) with initial data $\omega^\nu(0) \in L^2$ satisfying

$$\|(I - P_{\mathcal{K}})\omega^\nu(0)\|_{L^2} \leq d\nu, \tag{1.8}$$

we have

$$\left\| P_{\neq 0}\omega^\nu\left(\frac{\tau}{\nu}\right) \right\|_{L^2} < \delta \|P_{\neq 0}\omega^\nu(0)\|_{L^2}.$$

Here, $P_{\neq 0}$ is the projection of L^2 to the non-shear space X , that is,

$$P_{\neq 0}\omega = \omega - \frac{\alpha}{2\pi} \int_0^{2\pi/\alpha} \omega dx.$$

(ii) (Square torus) Suppose $\alpha = 1$. There exist $d > 0$, such that: for any $M > 0$, $\tau > 0$ and $\delta > 0$, if ν is small enough, then for any solution $\omega^\nu(t)$ to (1.1) with initial data $\omega^\nu(0) \in L^2$ satisfying $\|(I - P_{\mathcal{K}})\omega^\nu(0)\|_{L^2} \leq d\nu$, either

$$\max_{0 \leq t \leq \frac{\tau}{\nu}} \|P_a\omega^\nu(t)\|_{L^2} \geq M \|P_{\neq 0}\omega^\nu(0)\|_{L^2} \tag{1.9}$$

or

$$\inf_{0 \leq t \leq \frac{\tau}{\nu}} \|(1 - P_a)P_{\neq 0}\omega^\nu(t)\|_{L^2} < \delta \|P_{\neq 0}\omega^\nu(0)\|_{L^2} \tag{1.10}$$

must hold true.

In the above theorem, the metastability of Kolmogorov flow is studied for perturbations of the size ν . On the rectangular torus, it is shown that the non-shear part of the perturbation is reduced to a factor δ of the initial size before the time scale $\frac{\tau}{\nu}$, which is much smaller than the viscous time scale $\frac{1}{\nu}$. Moreover, by taking the constant d to be smaller, we can ensure that $\|(I - P_{\mathcal{K}})\omega^\nu(t)\|_{L^2} \leq d\nu$ for all $t > 0$, thus we can repeatedly use Theorem 1.2 i) to get the rapid decay of the non-shear part before the viscous time scale.

The situation for the square torus is more subtle due to the existence of ‘anomalous modes’. By taking M to be large and τ, δ to be small, Theorem 1.2 i) implies that if the ‘anomalous modes’ do not get too amplified before the time scale $\frac{\tau}{\nu}$, then the non-shear part of the perturbation without ‘anomalous modes’ is rapidly reduced. If the ‘anomalous modes’ are indeed much amplified (i.e. 1.9) before the time scale $\frac{\tau}{\nu}$, it suggests that the energy of the perturbation is transferred to ‘anomalous modes’. Intuitively, this might be considered as a hint that the non-shear part without ‘anomalous modes’ should be reduced. A quantitative estimate of the enhanced damping for such case is under investigation.

Next, we discuss some key ideas in the proof of Theorems 1.1 and 1.2. Our work is partly motivated by the work of Constantin et al. [9] for the linear reaction diffusion equation

$$\partial_t \phi + v_0 \cdot \nabla \phi - \nu \Delta \phi = 0,$$

with an incompressible flow $v_0(x)$. In [9], the enhanced damping in the sense of Theorem 1.1 is proved under the assumption that the operator $v_0 \cdot \nabla$ has no non-constant eigenfunction in H^1 . Their proof is to consider the high and low frequency parts of the solution $\phi(t)$ separately. For the high frequency part (i.e. $\|\nabla \phi\|_{L^2} \approx N \|\phi\|_{L^2}$ for N large), the enhanced damping is ensured by the energy dissipation law

$$\partial_t \|\phi\|_{L^2}^2 = -\nu \|\nabla \phi\|_{L^2}^2. \tag{1.11}$$

The lower frequency part is shown to converge to zero in the time average sense, by using the following RAGE Theorem for the unitary group e^{itL} with the self-adjoint generator $L = i v_0 \cdot \nabla$:

Theorem (RAGE) [7] Let L be a self-adjoint operator in a Hilbert space H , P_c is the projection to the continuous spectrum space of L and B is any compact operator, then

$$\frac{1}{T} \int_0^T \|B e^{itL} P_c \psi\|_H^2 dt \rightarrow 0, \quad \text{when } T \rightarrow \infty.$$

In the proof of enhanced damping, B is taken to be the projection to the low frequency modes. Then the RAGE Theorem implies that the low frequency modes decay in the time average sense.

To apply these ideas to prove the enhanced damping for the linearized Navier–Stokes equation (1.3), there are a few difficulties to be overcome. First, for the Eq. (1.3), there is no obvious dissipation law as (1.11). We derive the identity

$$\frac{d}{dt} \int_{\mathbb{T}^\alpha} (|\omega^\nu|^2 - |\nabla \psi^\nu|^2) dx dy = -2\nu \int_{\mathbb{T}^\alpha} (|\nabla \omega^\nu|^2 - |\omega^\nu|^2) dx dy, \tag{1.12}$$

where $\psi^\nu = (-\Delta)^{-1} \omega^\nu$ is the stream function. When $\alpha > 1$, the quadratic forms on both sides of (1.12) are positive definite for non-shear vorticity (i.e. $\omega^\nu \in X$). When $\alpha = 1$, the positivity is still true in the space $X_1 = (I - P_a) X_0$. This provides a substitute of (1.11).

Second, even if we ignore the factor $e^{-\nu t}$ in (1.3), the linearized Euler operator $A = -\sin y \partial_x (1 + \Delta^{-1})$ is not anti-self-adjoint and the RAGE theorem cannot be applied directly to e^{tA} . An important observation is that A can be written in the Hamiltonian form $A = JL$, where

$$J = -\sin y \partial_x, \quad L = 1 + \Delta^{-1} \tag{1.13}$$

are anti-selfadjoint and selfadjoint operators in L^2 respectively. When $\alpha > 1$, since $L = 1 + \Delta^{-1} > 0$ on the non-shear space X , we can define a new inner product by $[\cdot, \cdot] = \langle L\cdot, \cdot \rangle$ on X , which is equivalent to the L^2 inner product. We observe that the operator A is anti-selfadjoint in the space $(X, [\cdot, \cdot])$. Moreover, on the space X ,

the operator A can be shown to have no embedded eigenvalues in the continuous spectra. Thus RAGE theorem can be applied to the semigroup e^{tA} to show the decay of the low frequency part in the time average sense. The linear enhanced damping for (1.3) then follows similarly as in [9]. For the square torus ($\alpha = 1$), there are additional anomalous modes $\{\cos x, \sin x\}$ lying in $\ker(1 + \Delta^{-1})$. For any initial data $\omega^v(0) \in X$, we note that $\omega_1^v = (I - P_a)\omega^v$ satisfies the equation

$$\partial_t \omega_1^v = \nu \Delta \omega_1^v - e^{-\nu t} \left[(I - P_a) \sin y \partial_x (1 + \Delta^{-1}) \right] \omega_1^v.$$

Let

$$A_1 = - (I - P_a) \sin y \partial_x (1 + \Delta^{-1}) = (I - P_a) J L,$$

then it can be checked that A_1 is anti-selfadjoint on the space $X_1 = (I - P_a)X$ in the inner product $[\cdot, \cdot] = \langle L \cdot, \cdot \rangle$, where the positivity of $L|_{X_1}$ is used. Thus applying the RAGE theorem to the semigroup e^{tA_1} on X_1 , we can again show that the low frequency part of $(I - P_a)e^{tJL}\omega^v(0)$ decays in the time average sense.

For the nonlinear NS equation (1.2), the evolution of the shear and non-shear parts are strongly coupled. For an initial perturbation $\omega^v(0)$ of size $O(\nu)$ in L^2 , the interaction terms are controllable and the nonlinear enhanced damping (metastability) still holds true. On the square torus, the analysis for the nonlinear problem is more involved due to the anomalous modes. We decompose the perturbation into four parts in: $W_a = \text{span}\{\cos x, \sin x\}$ and its complementary subspace W_a^\perp in the non-shear space, $W_{\mathcal{K}} = \text{span}\{\cos y, \sin y\}$ and its complementary subspace $W_{\mathcal{K}}^\perp$ in the shear space. By carefully analyzing the interaction of these four parts, we can show that the interaction terms in the nonlinear term $U^v \cdot \nabla \omega^v$ are under control when $\|\omega^v(0)\|_{L^2} = O(\nu)$. As a result, we can still split the non-shear vorticity into the low and high frequency parts and treat them separately as we do for the linearized equation. Then the nonlinear metastability can be proved. On the square torus, numerical evidences [5,6] suggested that the dipole states of the form $\omega_0 = \cos x + \cos y$ or $\sin x + \sin y$ appear more often in the long time dynamics of 2D Turbulence. The dipole flows are nonparallel and the enhanced damping problem is much more subtle to study. At the end of Section 3, we discuss some partial results and difficulties with the dipole states. In particular, in Proposition 3.1, we give a RAGE type theorem for the linearized Euler equation at dipoles.

In our proof of linear and nonlinear enhanced damping for Navier–Stokes equation, the Hamiltonian structures of the linearized Euler operator play an important role both in the derivation of the dissipation law (1.12) and in the control of the low frequency part. As a further application of these Hamiltonian structures, we consider the linear inviscid damping of more general shear flows $(U(y), 0)$. We study two classes of shear flows. One class is the flows without inflection points, which are spectrally stable by the classical Rayleigh criterion. The other class (called class \mathcal{K}^+) is the flows $U(y)$ with one inflection value U_s and $-\frac{U''}{U-U_s} > 0$. These two classes cover all the shear flows whose nonlinear stability in L^2 vorticity might be studied by the energy-Casimir method (see Remark 4.3). The flows in the first class are nonlinearly stable for any x period $2\pi/\alpha$ and are minimizers of

the energy-Casimir functional. The flows in the second class are stable only when $\alpha > \alpha_{\max}$ for some critical wave number α_{\max} , and are maximizers of the energy-Casimir functional. These shear flows often appear as long lived coherent states in 2D turbulence; for example, the Kolmogorov flows are in class \mathcal{K}^+ .

In Theorem 4.1, we give a RAGE theorem on the non-shear subspace X of L^2 for stable shear flows in the first class and in class \mathcal{K}^+ with $\alpha > \alpha_{\max}$. As a consequence, the decay of velocity (in the time average sense) is proved for any non-shear initial data with L^2 vorticity. Another consequence is the decay of low frequency modes in the L^2 norm of vorticity, which gives a justification of the dual cascade of 2D turbulence in a weak sense (see Remark 4.2). For the critical case $\alpha = \alpha_{\max}$, the linearized Euler operator JL (defined in (1.13)) has zero as an embedded eigenvalue due to the nontrivial $\ker L$. This case is very similar to the case of bar states on the square torus and can be treated similarly. The linear damping can be obtained by projecting out $\ker L$.

The flows in class \mathcal{K}^+ are unstable when $\alpha < \alpha_{\max}$ [10]. Moreover, by using an instability index theory recently developed in [15] for Hamiltonian PDEs, we give an exact counting formula (Proposition 4.1) for the dimension of unstable modes of the linearized Euler equation. A corollary of this formula is that $L|_{E^c} \geq 0$, where E^c is the center space corresponding to the spectra of the linearized Euler operator JL (defined in (1.13)) on the imaginary axis. Then the RAGE theorem, and as a consequence the damping of velocity, are obtained for the linearized Euler equation on E^c .

The inviscid damping was first known for the Couette flow in the 1907 work of Orr [19]. In recent years, the inviscid damping phenomena attracted new attention. In [13], it was showed that if we consider initial (vorticity) perturbation in the Sobolev space H^s ($s < \frac{3}{2}$) then the nonlinear damping is not true due to the existence of nonparallel steady flows of the form of Kelvin's cats eye near Couette. In [4], nonlinear inviscid damping was proved for perturbations near Couette in the Gevrey class (i.e. almost analytic).

The linear inviscid damping for more general shear flows was also recently studied by some authors. Monotone shears were considered in [27] for the case near Couette, and in [24] for more general cases. The optimal decay rates $O(1/t)$, $O(1/t^2)$ for the horizontal and vertical velocities were obtained for initial vorticity in H^1 and H^2 respectively. In [25], general shear flows satisfying some nondegeneracy conditions were considered, and certain space-time estimates for velocities of the linearized Euler equation were obtained for initial vorticity in H^1 . The optimal decay rates were also obtained in [25] for a special class of symmetric shear flows. The non-existence of embedded eigenvalues was assumed in above works.

We comment on some differences of our results on inviscid damping with the previous work. First, for the two classes of shear flows we considered, we do not need to assume the non-existence of embedded eigenvalues. This assumption is proved to be true for flows without inflection points and for flows in class \mathcal{K}^+ with $\alpha > \alpha_{\max}$, but for $\alpha = \alpha_{\max}$ and some $\alpha < \alpha_{\max}$, zero is indeed an embedded eigenvalue. For these cases, the inviscid damping can still be proved as in Corollary 4.1 ii) and Theorem 4.2 ii), as well as for the Kolmogorov flow on the square torus. Second, the inviscid damping results we obtained are for the initial vorticity

in L^2 . In [24,25], initial vorticity with higher regularity was considered and the linear damping for L^2 vorticity was not studied. In [24,25,27], the linearized Euler equation was studied in a channel. Here, we treat the cases of the channel and tori in a unified way. We note that RAGE theorem type results contain more information than the damping of velocities, for example the decay of low frequency part of the vorticity. Third, our approach, which exploits the Hamiltonian structures of the Euler equation, does not rely on ODE techniques as in other papers. Therefore, it could be used for nonparallel flows, see Proposition 3.1 for dipoles and Theorem 11.7 in [15] for general steady Euler flows. Moreover, more information on the damping could be derived from the regularity properties of the spectral measure of JL (see Remark 4.1). This might provide an alternative approach to study the inviscid damping in other models.

This paper is organized as follows: in Section 2, we study the linear enhanced damping for the linearized Navier–Stokes equation. In Section 3, the nonlinear enhanced damping (i.e. metastability of Kolmogorov flows) is proved for the nonlinear Navier–Stokes equation. We discuss the cases of rectangular and square tori separately in Sections 2 and 3. In Section 4, the linear inviscid damping is proved for both stable and unstable shear flows. Throughout the paper, we use C to denote a generic constant in the estimates and the dependence on parameters will be specified when necessary.

2. Linear Enhanced Damping

In this section, we prove Theorem 1.1 on the enhanced damping for the linearized Navier–Stokes equation (1.3). We consider the cases of rectangular and square tori separately.

2.1. Linearized Navier–Stokes on a Rectangular Torus

Consider the linearized equation (1.3) on a torus

$$\mathbb{T}_\alpha = \left\{ 0 < y < 2\pi, 0 < x < \frac{2\pi}{\alpha} \right\}, \alpha > 1.$$

We divide the proof of Theorem 1.1 i) into several steps. In the proof, we shall use C to denote a generic constant in the estimates. First, we prove the dissipation law (1.12).

Lemma 2.1. *Let $\omega^\nu(t)$ be a solution of (1.3) with the initial data $\omega^\nu(t) \in L^2(\mathbb{T}_\alpha)$. Then*

$$\frac{d}{dt} \int_{\mathbb{T}_\alpha} (|\omega^\nu|^2 - |\nabla \psi^\nu|^2) dx dy = -2\nu \int_{\mathbb{T}_\alpha} (|\nabla \omega^\nu|^2 - |\omega^\nu|^2) dx dy \quad (2.1)$$

for any $t > 0$.

Proof. The Eq. (1.3) can be written as

$$\partial_t \omega^\nu = \nu \Delta \omega + e^{-\nu t} J L \omega,$$

where J, L are defined in (1.13). Thus we have

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{T}_\alpha} (|\omega^\nu|^2 - |\nabla \psi^\nu|^2) dx dy &= \frac{d}{dt} \langle L \omega^\nu, \omega^\nu \rangle = 2 \langle L \omega^\nu, \omega_t^\nu \rangle \\ &= e^{-\nu t} \langle L \omega^\nu, J L \omega^\nu \rangle \\ &\quad + 2 \int_{\mathbb{T}_\alpha} \nu \Delta \omega^\nu (\omega^\nu - \psi^\nu) dx dy \\ &= -2\nu \int_{\mathbb{T}_\alpha} (|\nabla \omega^\nu|^2 - |\omega^\nu|^2) dx dy. \end{aligned}$$

In the last equality above, we use integration by parts and the fact that J is anti-selfadjoint. \square

In the next lemma, we show that the quadratic forms on both sides of (2.1) are positive definite for a non-shear vorticity.

Lemma 2.2. *Let $\alpha > 1$ and $\omega \in X \cap H^1(\mathbb{T}_\alpha)$. Then there exists a constant $c_0 > 0$ depending only on α such that*

$$\int_{\mathbb{T}_\alpha} (|\omega|^2 - |\nabla \psi|^2) dx dy \geq c_0 \|\omega\|_{L^2}^2, \tag{2.2}$$

and

$$\int_{\mathbb{T}_\alpha} (|\nabla \omega|^2 - |\omega|^2) dx dy \geq c_0 \|\omega\|_{H^1}^2. \tag{2.3}$$

Proof. For any

$$\omega = \sum_{k \neq 0} \omega_k(y) e^{ik\alpha x} \in H^1(\mathbb{T}_\alpha),$$

we have

$$\begin{aligned} \int_{\mathbb{T}_\alpha} (|\omega|^2 - |\nabla \psi|^2) dx dy &= \sum_{0 \neq k \in \mathbf{Z}} \left\langle \left(1 - \left(-\frac{d^2}{dy^2} + \alpha^2 k^2 \right)^{-1} \right) \omega_k, \omega_k \right\rangle \\ &\geq (1 - \alpha^{-2}) \sum_{0 \neq k \in \mathbf{Z}} (\omega_k, \omega_k) = (1 - \alpha^{-2}) \|\omega\|_{L^2}^2, \end{aligned}$$

and

$$\begin{aligned} \int_{\mathbb{T}_\alpha} (|\nabla \omega|^2 - |\omega|^2) dx dy &= \sum_{0 \neq k \in \mathbf{Z}} \left\langle \left(-\frac{d^2}{dy^2} + \alpha^2 k^2 - 1 \right) \omega_k, \omega_k \right\rangle \\ &\geq \sum_{0 \neq k \in \mathbf{Z}} \int |\omega'_k(y)|^2 dy + (\alpha^2 - 1) \int |\omega_k(y)|^2 dy \\ &\geq \min \{ 1, \alpha^2 - 1 \} \|\omega\|_{H^1}^2. \end{aligned}$$

\square

Next, we study the linearized Euler equation at the Kolmogorov flow

$$\partial_t \omega = -\sin y \partial_x \left(1 + \Delta^{-1}\right) \omega = JL\omega. \tag{2.4}$$

Lemma 2.3. *Let $\omega(t)$ be a solution of (2.4) with $\omega(0) \in X \cap H^1(\mathbb{T}_\alpha)$. Then*

$$\|\omega(t)\|_{H^1} \leq C(1+t)\|\omega(0)\|_{H^1}$$

for some constant C .

Proof. First, we note that

$$\langle L\omega, \omega \rangle = \int_{\mathbb{T}_\alpha} (|\omega|^2 - |\nabla \psi|^2) dx dy$$

is conserved for (1.3). Therefore, by the positivity estimate (2.2) we have

$$\|\omega(t)\|_{L^2} = \left\| e^{tJL} \omega(0) \right\|_{L^2} \leq C \|\omega(0)\|_{L^2} \tag{2.5}$$

for some constant C . Taking ∂_x of (2.4), we have

$$\partial_t \partial_x \omega = -\sin y \partial_x \left(1 + \Delta^{-1}\right) \partial_x \omega$$

and therefore

$$\|\partial_x \omega(t)\|_{L^2} = \left\| e^{tJL} \partial_x \omega(0) \right\|_{L^2} \leq C \|\partial_x \omega(0)\|_{L^2}.$$

Taking ∂_y of (2.4), we have

$$\partial_t \partial_y \omega = -\sin y \partial_x \left(1 + \Delta^{-1}\right) \partial_y \omega - \cos y \partial_x \left(1 + \Delta^{-1}\right) \omega,$$

and

$$\partial_y \omega(t) = e^{tJL} \partial_y \omega(0) - \int_0^t e^{(t-s)JL} \cos y \partial_x \left(1 + \Delta^{-1}\right) \omega(s) ds.$$

Therefore

$$\begin{aligned} \|\partial_y \omega(t)\|_{L^2} &\leq C \left(\|\partial_y \omega(0)\|_{L^2} + \int_0^t \left\| \partial_x \left(1 + \Delta^{-1}\right) \omega(s) \right\|_{L^2} ds \right) \\ &\leq C(1+t) \|\nabla \omega(0)\|_{L^2}. \end{aligned}$$

This finishes the proof of the lemma. \square

In the next lemma, we study the spectral properties of the linearized Euler operator $A = JL$.

Lemma 2.4. (i) *The operator $A : X \rightarrow X$ is anti-selfadjoint in the inner product $[\cdot, \cdot] = \langle L\cdot, \cdot \rangle$.*

(ii) *The spectrum of A lies on the imaginary axis and is purely continuous.*

Proof. (i) By the positivity of L on X , $[\cdot, \cdot] = \langle L\cdot, \cdot \rangle$ defines an equivalent inner product to the L^2 inner product. For any $\omega_1, \omega_2 \in X$, we have

$$\begin{aligned} [A\omega_1, \omega_2] &= \langle LJL\omega_1, \omega_2 \rangle = \langle JL\omega_1, L\omega_2 \rangle = -\langle L\omega_1, JL\omega_2 \rangle \\ &= -[\omega_1, A\omega_2], \end{aligned}$$

and thus A is anti-selfadjoint on $(X, [\cdot, \cdot])$.

(ii) By property i), the spectrum of A in L^2 is on the imaginary axis. Since $A = -\sin y \partial_x (1 + \Delta^{-1})$ is a compact perturbation of $D = -\sin y \partial_x$, by Weyl's Theorem the continuous spectrum of A is the same as that of D , which is clearly the whole imaginary axis. It remains to show that there is no embedded eigenvalue of A on the imaginary axis. Suppose $A\omega = \lambda\omega$, where $\lambda \in i\mathbf{R}$ and $0 \neq \omega \in X$. Let

$$\omega = \sum_{0 \neq k \in \mathbf{Z}} \omega_k(y) e^{ik\alpha x}, \quad \psi_k = \left(-\frac{d^2}{dy^2} + \alpha^2 k^2 \right)^{-1} \omega_k.$$

Then if $\omega_k \neq 0$, we have

$$ik\alpha \sin y (\omega_k - \psi_k) = \lambda\omega_k,$$

which is equivalent to the Rayleigh equation

$$\left(-\frac{d^2}{dy^2} + \alpha^2 k^2 - \frac{\sin y}{\sin y - c} \right) \psi_k = 0,$$

with $c = \frac{\lambda}{ik\alpha} \in \mathbf{R}$. Since $\omega_k \in L^2$ implies $\psi_k \in H^2$, by Lemma 4.3 in the Appendix, we must have $c = 0$ which is the only inflection value of $\sin y$. Thus

$$\left(-\frac{d^2}{dy^2} + \alpha^2 k^2 - 1 \right) \psi_k = 0,$$

which implies that $\psi_k = 0$ since $\alpha > 1$. This contradiction rules out the embedded imaginary eigenvalues of A in X . \square

By the above Lemma and the RAGE theorem, we have

Corollary 2.1. *Let B be any compact operator in $L^2(\mathbb{T}_\alpha)$. Then*

$$\frac{1}{T} \int_0^T \|B\omega(t)\|_{L^2}^2 dt \rightarrow 0, \quad \text{when } T \rightarrow \infty$$

for any solution $\omega(t)$ of (2.4) with $\omega(0) \in X$.

For the proof of the enhanced damping, we need a more quantitative version of RAGE theorem. Let $\alpha^2 \leq \lambda_1 \leq \lambda_2 \leq \dots$ be the eigenvalues of the operator $-\Delta$ on X and e_1, e_2, \dots be the corresponding orthonormal eigenvectors. Denote P_N to be the L^2 projection to the subspace spanned by the first N eigenvectors

e_1, e_2, \dots, e_N of $-\Delta$ and $S = \{\omega \in X : \|\omega\|_{L^2} = 1\}$ be the unit sphere in X . Denote the norms

$$\begin{aligned} \|\omega\|_X^2 &= \left\langle \left(1 + \Delta^{-1}\right) \omega, \omega \right\rangle = \int_{\mathbb{T}_\alpha} (|\omega|^2 - |\nabla \psi|^2) dx dy, \\ \|\omega\|_{X^1}^2 &= \langle (-\Delta - 1) \omega, \omega \rangle = \int_{\mathbb{T}_\alpha} (|\nabla \omega|^2 - |\omega|^2) dx dy, \end{aligned}$$

which are equivalent to L^2 and H^1 norms on the non-shear space X . Let $\omega = \sum_{k \geq 1} c_k e_k$, then

$$\|\omega\|_X^2 = \sum_{k \geq 1} \left(1 - \frac{1}{\lambda_k}\right) |c_k|^2,$$

and

$$\|\omega\|_{X^1}^2 = \sum_{k \geq 1} (\lambda_k - 1) |c_k|^2.$$

The following version of the RAGE theorem can be obtained as in [9]:

Lemma 2.5. *Let $K \subset S$ be a compact set and J, L are defined in (1.13). For any $N, \kappa > 0$, there exists $T_c(N, \kappa, K)$ such that for all $T \geq T_c$ and any $\omega(0) \in K$,*

$$\frac{1}{T} \int_0^T \|P_N e^{tJL} \omega(0)\|_X^2 dt \leq \kappa \|\omega(0)\|_X^2.$$

Now we estimate the difference of the solutions of the linearized NS and Euler equations.

Lemma 2.6. *Let ω^v, ω^0 be the solutions of the linearized NS equation (1.3) and Euler equation (2.4) with the initial data in $X \cap H^1$. Then there exists some constant $C_0 > 0$ such that*

$$\frac{d}{dt} \|\omega^v - \omega^0\|_X^2 \leq C_0 v (1 + t^2) \|\omega^0(t)\|_{H^1}^2 \tag{2.6}$$

for all $t \in (0, +\infty)$.

Proof. Let ψ^v, ψ^0 be the corresponding stream functions. Denote $\omega = \omega^v - \omega^0$ and $\psi = \psi^v - \psi^0$, then

$$\omega_t + e^{-vt} \sin y \partial_x (\omega - \psi) + (e^{-vt} - 1) \sin y \partial_x (\omega^0 - \psi^0) - v \Delta \omega^v = 0.$$

We have

$$\begin{aligned}
 \frac{d}{dt} \frac{1}{2} \|\omega^v - \omega^0\|_X^2 &= \frac{d}{dt} \frac{1}{2} \int_{\mathbb{T}_\alpha} (|\omega|^2 - |\nabla \psi|^2) dx dy \\
 &= \int_{\mathbb{T}_\alpha} \omega_t (\omega - \psi) dx dy \\
 &= - \int_{\mathbb{T}_\alpha} (e^{-\nu t} - 1) \sin y \partial_x (\omega^0 - \psi^0) (\omega - \psi) dx dy \\
 &\quad + \nu \int_{\mathbb{T}_\alpha} \Delta \omega^v (\omega^v - \psi^v) dx dy \\
 &\quad - \nu \int_{\mathbb{T}_\alpha} \Delta \omega^v (\omega^0 - \psi^0) dx dy \\
 &= I + II + III.
 \end{aligned}$$

Since $0 \leq 1 - e^{-\nu t} \leq \nu t$ when $t > 0$, we have

$$\begin{aligned}
 I &= (1 - e^{-\nu t}) \int_{\mathbb{T}_\alpha} \sin y \partial_x (\omega^0 - \psi^0) (\omega^v - \psi^v) dx dy \\
 &= -(1 - e^{-\nu t}) \int_{\mathbb{T}_\alpha} \sin y \partial_x (\omega^v - \psi^v) (\omega^0 - \psi^0) dx dy \\
 &\leq C \nu t \|\omega^v\|_{H^1} \|\omega^0\|_{L^2}.
 \end{aligned}$$

By integration by parts and (2.3),

$$II = -\nu \int_{\mathbb{T}_\alpha} (|\nabla \omega^v|^2 - |\omega^v|^2) dx dy \leq -c_0 \nu \|\omega^v\|_{H^1}^2.$$

For the last term, we have

$$III = \nu \int_{\mathbb{T}_\alpha} \nabla \omega^v \cdot \nabla (\omega^0 - \psi^0) dx dy \leq \nu \|\omega^v\|_{H^1} \|\omega^0\|_{H^1}.$$

Combining the above, we get

$$\begin{aligned}
 \frac{d}{dt} \frac{1}{2} \int_{\mathbb{T}_\alpha} (|\omega|^2 - |\nabla \psi|^2) dx dy \\
 \leq \nu \left(-c_0 \|\omega^v\|_{H^1}^2 + \|\omega^v\|_{H^1} \left(C t \|\omega^0\|_{L^2} + \|\omega^0\|_{H^1} \right) \right) \\
 \leq C \nu \left(1 + t^2 \right) \|\omega^0(t)\|_{H^1}^2.
 \end{aligned}$$

This proves (2.6). \square

As a corollary, combining (2.6), Lemma 2.3 and (2.3), we have

$$\|\omega^v(t) - \omega^0(t)\|_X^2 \leq \|\omega^v(0) - \omega^0(0)\|_X^2 + C_1 \nu \left(1 + t^5\right) \|\omega^0(0)\|_{X^1}^2 \quad (2.7)$$

for some constant $C_1 > 0$.

We are now ready to prove the linear enhanced damping.

Proof of Theorem 1.1 i). The proof follows the arguments in [9]. Fixing $\delta, \tau > 0$, we choose N such that

$$\exp(-2\lambda_N \tau) < c_0 \delta^2,$$

where c_0 is the constant in (2.2). Define a compact set

$$K = \text{span} \{e_1, \dots, e_N\} = R(P_N).$$

Denote $t_1 = T_c(N, \frac{1}{10}, K)$ as in Lemma 2.5 and let $\nu(\delta, \tau)$ be such that

$$\nu(\delta, \tau) C_1 \left(1 + t_1^5\right) < \frac{1}{10\lambda_N},$$

where C_1 is the constant in the estimate (2.7). For $0 < \nu < \nu(\delta, \tau)$, suppose that

$$\|\omega^\nu(t)\|_{X^1}^2 > \lambda_N \|\omega^\nu(t)\|_X^2$$

is true for t in some interval $(a, b) \subset (0, \tau/\nu)$. Then by (2.1), we have

$$\|\omega^\nu(b)\|_X^2 \leq \exp(-\nu\lambda_N(b-a)) \|\omega^\nu(a)\|_X^2. \tag{2.8}$$

Now consider any $t_0 \in (0, \tau/\nu)$ satisfying

$$\|\omega^\nu(t_0)\|_{X^1}^2 \leq \lambda_N \|\omega^\nu(t_0)\|_X^2.$$

Denote $\omega_0 = \omega^\nu(t_0)$ and let $\omega^0(t)$ ($t \in [t_0, t_0 + t_1]$) be the solution of (2.4) with $\omega^0(t_0) = \omega_0$. By the choice of $t_0, \nu(\delta, \tau)$ and (2.7), we have

$$\|\omega^\nu(t) - \omega^0(t)\|_X^2 \leq \frac{1}{10} \|\omega_0\|_X^2, \quad \forall t \in [t_0, t_0 + t_1]. \tag{2.9}$$

By the definition of t_1 , we have

$$\frac{1}{t_1} \int_{t_0}^{t_0+t_1} \|P_N \omega^0(t)\|_X^2 dt \leq \frac{1}{10} \|\omega_0\|_X^2.$$

Since $\|\omega^0(t)\|_X = \|\omega_0\|_X$ by the conservation of $\langle L\omega, \omega \rangle$ for the Eq.(2.4), it follows that

$$\frac{1}{t_1} \int_{t_0}^{t_0+t_1} \|(1 - P_N) \omega^0(t)\|_X^2 dt \geq \frac{9}{10} \|\omega_0\|_X^2.$$

Combined with (2.9), the above implies that

$$\frac{1}{t_1} \int_{t_0}^{t_0+t_1} \|(1 - P_N) \omega^\nu(t)\|_X^2 dt \geq \frac{1}{2} \|\omega_0\|_X^2.$$

For $\omega^\nu(t) = \sum_{k \geq 1} c_k e_k$, we have

$$\begin{aligned} \|\omega^\nu(t)\|_{X^1}^2 &= \sum_{k \geq 1} (\lambda_k - 1) |c_k|^2 \geq \frac{1}{\lambda_N} \sum_{k \geq N+1} \left(1 - \frac{1}{\lambda_k}\right) |c_k|^2 \\ &= \frac{1}{\lambda_N} \|(1 - P_N) \omega^\nu(t)\|_X^2, \end{aligned}$$

and thus

$$\int_{t_0}^{t_0+t_1} \|\omega^v(t)\|_{X^1}^2 dt \geq \frac{\lambda_N t_1}{2} \|\omega_0\|_X^2 = \frac{\lambda_N t_1}{2} \|\omega^v(t_0)\|_X^2.$$

Then (2.1) implies that

$$\begin{aligned} \|\omega^v(t_0 + t_1)\|_X^2 &\leq \|\omega^v(t_0)\|_X^2 - 2\nu \int_{t_0}^{t_0+t_1} \|\omega^v(t)\|_{X^1}^2 dt \\ &\leq (1 - \lambda_N \nu t_1) \|\omega^v(t_0)\|_X^2 \leq e^{-\lambda_N \nu t_1} \|\omega^v(t_0)\|_X^2. \end{aligned} \tag{2.10}$$

We can split the interval $[0, \frac{\tau}{\nu}]$ into a union of intervals such that either (2.8) or (2.10) holds true. Therefore we have

$$\|\omega^v\left(\frac{\tau}{\nu}\right)\|_X^2 \leq e^{-\lambda_N \tau} \|\omega^v(0)\|_X^2 < c_0 \delta^2 \|\omega^v(0)\|_X^2,$$

and by (2.2),

$$\|\omega^v\left(\frac{\tau}{\nu}\right)\|_{L^2}^2 \leq \frac{1}{c_0} \|\omega^v\left(\frac{\tau}{\nu}\right)\|_X^2 < \delta^2 \|\omega^v(0)\|_X^2 < \delta^2 \|\omega^v(0)\|_{L^2}^2.$$

This finishes the proof of Theorem 1.1 i). \square

2.2. Linearized Navier–Stokes on a Square Torus

Now we consider the linearized equation (1.3) on the square torus

$$\mathbb{T}_1 = \{0 < y < 2\pi, 0 < x < 2\pi\}.$$

In this case, there is a two dimensional kernel space W_a spanned by $\{\cos x, \sin x\}$ of the operator $L = 1 + \Delta^{-1}$ on the non-shear space X . We will sketch the changes induced by these anomalous modes, in the proof of Theorem 1.1 ii).

First, we note that L is positive on the space $X_1 = (I - P_a)X$, where P_a is the projection of X to W_a . Let $\omega^v(t)$ be the solution of (1.3) with any initial data $\omega^v(0) \in X$. Then $\omega_1^v = (I - P_a)\omega^v$ satisfies the equation

$$\partial_t \omega_1^v = \nu \Delta \omega_1^v - e^{-\nu t} \left[(I - P_a) \sin y \partial_x (1 + \Delta^{-1}) \right] \omega_1^v. \tag{2.11}$$

It is easy to check that the same dissipation law

$$\frac{d}{dt} \int_{\mathbb{T}_1} (|\omega_1^v|^2 - |\nabla \psi_1^v|^2) dx dy = -2\nu \int_{\mathbb{T}_1} (|\nabla \omega_1^v|^2 - |\omega_1^v|^2) dx dy,$$

holds true for (2.11). Moreover, there exists $c_0 > 0$ such that

$$\int_{\mathbb{T}_1} (|\omega|^2 - |\nabla \psi|^2) dx dy \geq c_0 \|\omega\|_{L^2}^2 \tag{2.12}$$

and

$$\int_{\mathbb{T}_1} (|\nabla \omega|^2 - |\omega|^2) dx dy \geq c_0 \|\omega\|_{H^1}^2$$

for $\omega \in X_1$. Define the operator $A_1 : X_1 \rightarrow X_1$ by

$$A_1 = - (I - P_a) \sin y \partial_x \left(1 + \Delta^{-1} \right) = (I - P_a) J L.$$

Since $L|_{X_1} > 0$, $[\cdot, \cdot] = \langle L \cdot, \cdot \rangle$ is again an equivalent inner product on X_1 and A_1 is anti-selfadjoint on $(X_1, [\cdot, \cdot])$. Indeed, for any $w_1, w_2 \in X_1$

$$\begin{aligned} [A_1 w_1, w_2] &= \langle L (I - P_a) J L w_1, w_2 \rangle = \langle L J L w_1, w_2 \rangle = - \langle L w_1, J L w_2 \rangle \\ &= - \langle L w_1, (I - P_a) J L w_2 \rangle = - [w_1, A_1 w_2]. \end{aligned}$$

Therefore the spectrum of A_1 lies on the imaginary axis. Next we show that A_1 has no embedded imaginary eigenvalues.

Lemma 2.7. *The spectrum of A_1 is purely continuous.*

Proof. Suppose A_1 has an eigenvalue $\lambda \in i\mathbf{R}$ and $A_1 \omega = \lambda \omega$, where $0 \neq \omega \in X_1$. Then

$$A \omega - \lambda \omega = J L \omega - \lambda \omega = \tilde{\omega} \in W_a.$$

If $\lambda \neq 0$, by noting that $A \tilde{\omega} = 0$, we get

$$A \left(\omega + \frac{1}{\lambda} \tilde{\omega} \right) = \lambda \left(\omega + \frac{1}{\lambda} \tilde{\omega} \right),$$

that is, λ is an eigenvalue of A . This is a contradiction, since by the proof of Lemma 2.4, the operator A has no nonzero eigenvalues. If $\lambda = 0$, then we must have $A \omega = \tilde{\omega}$ for some $0 \neq \tilde{\omega} \in W_a$, since $\omega \in X_1$ implies that $A \omega \neq 0$. Let $\tilde{\omega} = c_1 e^{ix} + c_{-1} e^{-ix} \in W_a$ and

$$\psi = \Delta^{-1} \omega = a_1(y) e^{ix} + a_{-1}(y) e^{-ix}.$$

From the equation $A \omega = \sin y \partial_x (\omega + \psi) = \tilde{\omega}$, we get

$$a_1''(y) = \frac{c_1}{i \sin y}, \quad a_{-1}''(y) = \frac{c_{-1}}{-i \sin y},$$

and thus $\psi \notin H^2$. This shows that 0 is not an eigenvalue of A_1 and the proof of the lemma is finished. \square

By the above lemma, we can use RAGE theorem for the semigroup e^{tA_1} on X_1 , which corresponds to solutions of the projected linearized Euler equation

$$\partial_t \omega = (I - P_a) \sin y \partial_x \left(1 + \Delta^{-1} \right) \omega. \tag{2.13}$$

In particular, let P_N be the projection of L^2 to the space spanned by the first N eigenfunction of $-\Delta$ on X_1 , then we have for any $N, \kappa > 0$ that there exists $T_c(N, \kappa, K)$ such that for all $T \geq T_c$ and any $\omega(0) \in R(P_N)$,

$$\frac{1}{T} \int_0^T \| P_N e^{tJL} \omega(0) \|_X^2 dt \leq \kappa \| \omega(0) \|_X^2. \tag{2.14}$$

In the next Lemma, we obtain the same estimates on the growth of solutions of (2.13) as we did in Lemma 2.3.

Lemma 2.8. *Let $\omega(t)$ be a solution of (2.13) with $\omega(0) \in X_1 \cap H^1(\mathbb{T}_1)$. Then*

$$\|\omega(t)\|_{H^1} \leq C(1+t)\|\omega(0)\|_{H^1}$$

for some constant C .

Proof. The proof is very similar to that of Lemma 2.3. We only sketch it. By the conservation of $\langle L\omega, \omega \rangle$ for the Eq. (2.13) and the positivity of $L|_{X_1}$, the L^2 norm of $\omega(t)$ is bounded by $\omega(0)$. Since P_a is the projector of X to $\ker L = \ker(1 + \Delta^{-1})$ and ∇ is commutable with $1 + \Delta^{-1}$, so P_a is also commutable with ∇ . Then the estimates of $\partial\omega(t)$ follow in the same way as in the proof of Lemma 2.3. \square

Similarly, we can estimate the difference of solutions of (2.13) and (2.11).

Lemma 2.9. *Let ω^v, ω^0 be the solutions of the projected linearized NS equation (1.3) and Euler equation (2.4) with the initial data $\omega^v(0) \in X_1$ and $\omega_0 \in X_1 \cap H^1$. Then there exists constants $C_0 > 0$ such that*

$$\frac{d}{dt} \|\omega^v - \omega^0\|_X^2 \leq C_0 v (1+t^2) \|\omega^0(t)\|_{H^1}^2$$

for $t \in (0, +\infty)$.

The proof is the same as that of Lemma 2.6 by using the fact that $L|_{X_1} > 0$ and $\langle L(I - P_a)\cdot, \cdot \rangle = \langle L\cdot, \cdot \rangle$.

Then by the same proof of Theorem 1.1 i), we can show the enhanced damping for the solution $\omega_1^v(t)$ of the projected Eq. (2.11). More precisely, for any $\tau > 0$ and $\delta > 0$, if v is small enough, then

$$\left\| \omega_1^v\left(\frac{\tau}{v}\right) \right\|_{L^2} < \delta \|\omega_1^v(0)\|_{L^2}.$$

Since $\omega_1^v(t) = (I - P_a)\omega^v(t)$, this proves Theorem 1.1 ii).

3. Nonlinear Enhanced Damping

In this section, we prove the metastability and enhanced damping of Kolmogorov flows for the nonlinear Navier–Stokes equation (1.1) on a torus \mathbb{T}_α ($\alpha \geq 1$). Let $P_{\mathcal{K}}$ to be the projection of $L^2(\mathbb{T}_\alpha)$ to the subspace $W_{\mathcal{K}} = \text{span}\{\cos y, \sin y\}$. For any initial data $\omega^v(0) \in L^2$, let

$$P_{\mathcal{K}}\omega^v(0) = d_1 \cos y + d_2 \sin y = D \sin(y + y_1),$$

where $D = \sqrt{d_1^2 + d_2^2}$ and $y_1 = \tan^{-1}(d_1/d_2)$. When $\|(I - P_{\mathcal{K}})\omega^v(0)\|_{L^2}$ is small, we can equivalently consider the perturbation near the shear flow $U(y) = D \sin(y + y_1)$ with initial data $\omega^v(0)$ satisfying $P_{\mathcal{K}}\omega^v(0) = 0$, for which the analysis is almost the same as for the shear flow $U(y) = \sin y$. For simplicity, in the proof of Theorem 1.2, we only consider the perturbations near $U(y) = \sin y$ with $P_{\mathcal{K}}\omega^v(0) = 0$. As in the proof of Theorem 1.1 for the linearized NS equation, we

will treat the high and low frequency parts of the non-shear perturbation separately. In particular, for the low frequency part, we will compare the solutions of the nonlinear NS equation and the linearized Euler equation, and then use the RAGE Theorem to control the time average. However, a significant difference with the linearized NS equation is that the shear and non-shear parts are strongly coupled for the nonlinear NS equation. Therefore, the main issue is to control the interaction terms. We will consider the equations on rectangular and square tori separately. On the square torus, the existence of anomalous modes makes the interactions terms considerably more subtle to handle.

3.1. The Case of Rectangular Torus

Consider the nonlinear Navier–Stokes equation near the Kolmogorov flow

$$\begin{aligned} \partial_t \omega^\nu &= \nu \Delta \omega^\nu - e^{-\nu t} \left[\sin y \partial_x \left(1 + \Delta^{-1} \right) \right] \omega^\nu + U^\nu \cdot \nabla \omega^\nu \\ &= L(t) \omega^\nu + U^\nu \cdot \nabla \omega^\nu, \end{aligned} \tag{3.1}$$

on \mathbb{T}_α ($\alpha > 1$), where ω^ν, U^ν are the perturbations of vorticity and velocity. We split ω^ν, U^ν into shear and non-shear components. More precisely, we write $\omega^\nu = \omega_s^\nu + \omega_n^\nu$, where the shear part is

$$\omega_s^\nu(t, y) = \frac{\alpha}{2\pi} \int_0^{\frac{2\pi}{\alpha}} \omega^\nu(t, x, y) dx = P_0 \omega^\nu,$$

and the non-shear part is

$$\omega_n^\nu(t, x, y) = (I - P_0) \omega^\nu = P_{\neq 0} \omega^\nu \in X.$$

Correspondingly, $U^\nu = U_s^\nu + U_n^\nu$ and $\psi^\nu = \psi_s^\nu + \psi_n^\nu$ where ψ^ν is the stream function. We also denote

$$U^\nu = (u^\nu, v^\nu), \quad U_s^\nu = (u_s^\nu(t, y), 0), \quad U_n^\nu = (u_n^\nu, v_n^\nu).$$

Then the Eq.(3.1) can be written as

$$\partial_t \omega_s^\nu = \nu \partial_{yy} \omega_s^\nu + P_0 (U_n^\nu \cdot \nabla \omega_n^\nu) = \nu \partial_{yy} \omega_s^\nu + \partial_y P_0 (v_n^\nu \omega_n^\nu), \tag{3.2}$$

and

$$\begin{aligned} \partial_t \omega_n^\nu &= L(t) \omega_n^\nu + u_s^\nu \partial_x \omega_n^\nu + v_n^\nu \partial_y \omega_s^\nu + P_{\neq 0} (U_n^\nu \cdot \nabla \omega_n^\nu) \\ &= L(t) \omega_n^\nu + u_s^\nu \partial_x \omega_n^\nu + v_n^\nu \partial_y \omega_s^\nu + \partial_x (u_n^\nu \omega_n^\nu) + \partial_y P_{\neq 0} (v_n^\nu \omega_n^\nu). \end{aligned} \tag{3.3}$$

First, we show that the dissipation law (1.12) also holds true for solutions of the nonlinear equation (3.1).

Lemma 3.1. *Let $\omega^\nu(t)$ be a solution of (3.1) with the initial data $\omega^\nu(0) \in L^2(\mathbb{T}_\alpha)$. Then*

$$\frac{d}{dt} \int_{\mathbb{T}_\alpha} (|\omega^\nu|^2 - |\nabla \psi^\nu|^2) dx dy = -2\nu \int_{\mathbb{T}_\alpha} (|\nabla \omega^\nu|^2 - |\omega^\nu|^2) dx dy \tag{3.4}$$

for any $t > 0$.

Proof. We have

$$\begin{aligned}
 \frac{d}{dt} \int_{\mathbb{T}_\alpha} (|\omega^\nu|^2 - |\nabla \psi^\nu|^2) dx dy &= \int_{\mathbb{T}_\alpha} \omega_t^\nu (\omega^\nu - \psi^\nu) dx dy \\
 &= \langle L(t) \omega^\nu, (\omega^\nu - \psi^\nu) \rangle \\
 &\quad + \int_{\mathbb{T}_\alpha} U^\nu \cdot \nabla \omega^\nu (\omega^\nu - \psi^\nu) dx dy \\
 &= -2\nu \int_{\mathbb{T}_\alpha} (|\nabla \omega^\nu|^2 - |\omega^\nu|^2) dx dy \\
 &\quad + \frac{1}{2} \int_{\mathbb{T}_\alpha} U^\nu \cdot \nabla \frac{1}{2} \omega^{\nu 2} dx dy \\
 &\quad - \int_{\mathbb{T}_\alpha} (U^\nu \cdot \nabla \psi^\nu) \omega^\nu dx dy \\
 &= -2\nu \int_{\mathbb{T}_\alpha} (|\nabla \omega^\nu|^2 - |\omega^\nu|^2) dx dy.
 \end{aligned}$$

In the above, we used the fact that $U^\nu \cdot \nabla \psi^\nu = 0$ and

$$\langle L(t) \omega^\nu, (\omega^\nu - \psi^\nu) \rangle = -2\nu \int_{\mathbb{T}_\alpha} (|\nabla \omega^\nu|^2 - |\omega^\nu|^2) dx dy,$$

as in the proof of Lemma 2.1. \square

Denote Y to be the space of mean zero functions in $L^2(\mathbb{T}_\alpha)$, depending only on y . Denote $Y_1 = (I - P_{\mathcal{K}})Y$. Then the operator $L = 1 + \Delta^{-1}$ is positive on Y_1 . There exists $c_0 > 0$ such that

$$\int_{\mathbb{T}_\alpha} (|\omega^\nu|^2 - |\partial_y \psi^\nu|^2) dx dy \geq c_0 \|\omega^\nu\|_{L^2}^2 \tag{3.5}$$

and

$$\int_{\mathbb{T}_\alpha} (|\partial_y \omega^\nu|^2 - |\omega^\nu|^2) dx dy \geq c_0 \|\omega^\nu\|_{H^1}^2 \tag{3.6}$$

for all $\omega^\nu \in Y_1$.

For a solution $\omega^\nu = \omega_s^\nu + \omega_n^\nu$ of the nonlinear NS equation (3.1), let $\omega_s^\nu = \omega_{s1}^\nu + \omega_{s2}^\nu$, where $\omega_{s1}^\nu = P_{\mathcal{K}} \omega_s^\nu = P_{\mathcal{K}} \omega^\nu$ and $\omega_{s2}^\nu \in Y_1$. Then (3.4) implies that

$$\begin{aligned}
 &\frac{d}{dt} \left(\int_{\mathbb{T}_\alpha} (|\omega_{s2}^\nu|^2 - |\partial_y \psi_{s2}^\nu|^2) dx dy + \|\omega_n^\nu\|_X^2 \right) \\
 &= -2\nu \left(\int_{\mathbb{T}_\alpha} (|\partial_y \omega_{s2}^\nu|^2 - |\omega_{s2}^\nu|^2) dx dy + \|\omega_n^\nu\|_{X^1}^2 \right). \tag{3.7}
 \end{aligned}$$

In particular, by the positivity estimates (2.2), (2.3), it follows from the above that there exists $C > 0$ such that

$$\|\omega_{s2}^\nu\|_{L^2}(t) + \|\omega_n^\nu\|_{L^2}(t) \leq C (\|\omega_{s2}^\nu\|_{L^2}(0) + \|\omega_n^\nu\|_{L^2}(0)) \leq C d\nu, \tag{3.8}$$

where we assume $\|\omega^v\|_{L^2}(0) \leq d v$ for a constant $d > 0$ to be determined later. To estimate

$$\omega_{s1}^v = a_1 \frac{\sqrt{\alpha}}{2\pi} \cos y + a_2 \frac{\sqrt{\alpha}}{2\pi} \sin y,$$

we project (3.2) to $\left\{ \frac{\sqrt{\alpha}}{\sqrt{2\pi}} \cos y, \frac{\sqrt{\alpha}}{\sqrt{2\pi}} \sin y \right\}$ to get

$$\begin{aligned} \frac{d}{dt} a_1 &= -v a_1 - \frac{\sqrt{\alpha}}{\sqrt{2\pi}} \int_{\mathbb{T}_\alpha} v_n^v \omega_n^v \sin y \, dx dy, \\ \frac{d}{dt} a_2 &= -v a_2 + \frac{\sqrt{\alpha}}{\sqrt{2\pi}} \int_{\mathbb{T}_\alpha} v_n^v \omega_n^v \cos y \, dx dy. \end{aligned} \tag{3.9}$$

Let

$$a(t) = \|\omega_{s1}^v\|_{L^2} = \sqrt{a_1^2 + a_2^2},$$

then, by (3.9), we have

$$\begin{aligned} \frac{da}{dt} &\leq -v a(t) + \frac{\sqrt{\alpha}}{2\sqrt{2\pi}} \sqrt{\left| \int_{\mathbb{T}_\alpha} v_n^v \omega_n^v \sin y \, dx dy \right|^2 + \left| \int_{\mathbb{T}_\alpha} v_n^v \omega_n^v \cos y \, dx dy \right|^2} \\ &\leq -v a(t) + C \|v_n^v\|_{L^2}(t) \|\omega_n^v\|_{L^2}(t). \end{aligned}$$

Since $a(0) = \|P_{\mathcal{K}} \omega^v(0)\|_{L^2} = 0$, the above implies that

$$\begin{aligned} \|\omega_{s1}^v\|_{L^2} = a(t) &\leq C d v \int_0^t e^{-v(t-s)} \|\omega_n^v\|_{L^2}(s) \, ds \\ &\leq C \|\omega_n^v\|_{L^2}(0) \leq C d v. \end{aligned} \tag{3.10}$$

Combined with (3.8), it follows from the above that

$$\|\omega_s^v\|_{L^2}(t) \leq C d v, \text{ for all } t > 0. \tag{3.11}$$

In the dissipation law (3.7), ω_{s2}^v and ω_n^v are coupled. In the next lemma, we show that when d is small, the dissipation law for ω_n^v can be ‘‘separated’’ from (3.7).

Lemma 3.2. *There exists a constant d depending only on α , such that when $\|\omega^v\|_{L^2}(0) \leq d v$, then*

$$\frac{d}{dt} \int_{\mathbb{T}_\alpha} \|\omega_n^v\|_X^2 \leq -v \|\omega_n^v\|_{X^1}^2. \tag{3.12}$$

Proof. By (3.2), we have

$$\begin{aligned} & \frac{d}{dt} \int_{\mathbb{T}_\alpha} (|\omega_{s2}^v|^2 - |\partial_y \psi_{s2}^v|^2) dx dy \\ &= 2 \int_{\mathbb{T}_\alpha} \partial_t \omega_{s2}^v (\omega_{s2}^v - \psi_{s2}^v) dx dy = 2 \int_{\mathbb{T}_\alpha} \partial_t \omega_s^v (\omega_{s2}^v - \psi_{s2}^v) dx dy \\ &= -2\nu \int_{\mathbb{T}_\alpha} (|\partial_y \omega_{s2}^v|^2 - |\omega_{s2}^v|^2) dx dy + 2 \int_{\mathbb{T}_\alpha} \partial_y (v_n^v \omega_n^v) (\omega_{s2}^v - \psi_{s2}^v) dx dy \\ &\geq -2\nu \int_{\mathbb{T}_\alpha} (|\partial_y \omega_{s2}^v|^2 - |\omega_{s2}^v|^2) dx dy - C \|\omega_n^v\|_{X^1}^2 \|\omega_{s2}^v\|_{L^2} \\ &\geq -2\nu \int_{\mathbb{T}_\alpha} (|\partial_y \omega_{s2}^v|^2 - |\omega_{s2}^v|^2) dx dy - Cd\nu \|\omega_n^v\|_{X^1}^2. \end{aligned}$$

In the first inequality above, we use the Sobolev embedding and (2.3). Combined with (3.7), this gives

$$\frac{d}{dt} \int_{\mathbb{T}_\alpha} \|\omega_n^v\|_X^2 \leq -\nu \|\omega_n^v\|_{X^1}^2 (2 - Cd) \leq -\nu \|\omega_n^v\|_{X^1}^2$$

when $d \leq 1/C$. \square

To use the RAGE theorem to control the low frequency part of ω_n^v , we need to estimate the difference of the solutions of the nonlinear NS equation (3.1) and the linearized Euler equation (2.4).

Lemma 3.3. *There exists $d > 0$ such that for any solution ω^v of Navier–Stokes equation (3.1) with initial data satisfying $\|\omega^v\|_{L^2}(0) \leq d\nu$, and any solution ω^0 of the linearized Euler equation (2.4) with initial data in $X \cap H^1$, we have*

$$\frac{d}{dt} \|\omega_n^v - \omega^0\|_X^2 \leq C_0\nu (1 + t^2) \|\omega^0(t)\|_{H^1}^2, \quad \forall t > 0$$

for some constant $C_0 > 0$. Here, $\omega_n^v = P_{\neq 0} \omega^v$ is the non-shear part of ω^v .

Proof. Let ψ_n^v, ψ^0 be the corresponding stream functions. Denote $\omega = \omega_n^v - \omega^0$ and $\psi = \psi_n^v - \psi^0$, then

$$\begin{aligned} \omega_t &= -e^{-\nu t} \sin y \partial_x (\omega - \psi) - (e^{-\nu t} - 1) \sin y \partial_x (\omega^0 - \psi^0) + \nu \Delta \omega_n^v \\ &\quad + u_s^v \partial_x \omega_n^v + v_n^v \partial_y \omega_s^v + P_{\neq 0} (U_n^v \cdot \nabla \omega_n^v). \end{aligned}$$

Thus

$$\begin{aligned}
 \frac{d}{dt} \frac{1}{2} \|\omega^\nu - \omega^0\|_X^2 &= \int_{\mathbb{T}_\alpha} \omega_t (\omega - \psi) \, dx dy \\
 &= \left[- \int_{\mathbb{T}_\alpha} (e^{-\nu t} - 1) \sin y \partial_x (\omega^0 - \psi^0) (\omega - \psi) \, dx dy \right. \\
 &\quad \left. + \nu \int_{\mathbb{T}_\alpha} \Delta \omega_n^\nu (\omega - \psi) \, dx dy \right] \\
 &\quad + \int_{\mathbb{T}_\alpha} u_s^\nu \partial_x \omega_n^\nu (\omega - \psi) \, dx dy \\
 &\quad + \int_{\mathbb{T}_\alpha} v_n^\nu \partial_y \omega_s^\nu (\omega - \psi) \, dx dy \\
 &\quad + \int_{\mathbb{T}_\alpha} U_n^\nu \cdot \nabla \omega_n^\nu (\omega - \psi) \, dx dy \\
 &= I + II + III + IV.
 \end{aligned}$$

Similar to the proof of Lemma 2.6, the first term can be estimated by

$$I \leq \nu \left(-c_0 \|\omega_n^\nu\|_{H^1}^2 + C(1+t) \|\omega^\nu\|_{H^1} \|\omega^0\|_{H^1} \right).$$

For the last term, noticing that, as in the proof of Lemma 3.1,

$$\int_{\mathbb{T}_\alpha} U_n^\nu \cdot \nabla \omega_n^\nu (\omega_n^\nu - \psi_n^\nu) \, dx dy = 0,$$

we have

$$\begin{aligned}
 IV &= - \int_{\mathbb{T}_\alpha} U_n^\nu \cdot \nabla \omega_n^\nu (\omega^0 - \psi^0) \, dx dy = \int_{\mathbb{T}_\alpha} U_n^\nu \cdot \nabla (\omega^0 - \psi^0) \omega_n^\nu \, dx dy \\
 &\leq C \|U_n^\nu\|_{L^4} \|\omega^0\|_{H^1} \|\omega_n^\nu\|_{L^4} \leq C \|\omega_n^\nu\|_{L^2} \|\omega^0\|_{H^1} \|\omega_n^\nu\|_{H^1} \\
 &\leq C d \nu \|\omega^0\|_{H^1} \|\omega_n^\nu\|_{H^1}.
 \end{aligned}$$

The second term is estimated by

$$\begin{aligned}
 II &= \int_{\mathbb{T}_\alpha} u_s^\nu \partial_x \omega_n^\nu (\psi_n^\nu - \omega^0 + \psi^0) \, dx dy \\
 &= - \int_{\mathbb{T}_\alpha} u_s^\nu \omega_n^\nu (\partial_x \psi_n^\nu - \partial_x (\omega^0 - \psi^0)) \, dx dy \\
 &\leq \|u_s^\nu\|_{L^\infty} \|\omega_n^\nu\|_{L^2} \|v_n^\nu\|_{L^2} + \|u_s^\nu\|_{L^\infty} \|\omega_n^\nu\|_{L^2} \|\omega^0\|_{H^1} \\
 &\leq C \left(\|\omega_s^\nu\|_{L^2} \|\omega_n^\nu\|_{H^1}^2 + \|\omega_s^\nu\|_{L^2} \|\omega_n^\nu\|_{H^1} \|\omega^0\|_{H^1} \right) \\
 &\leq C d \nu \left(\|\omega_n^\nu\|_{H^1}^2 + \|\omega_n^\nu\|_{H^1} \|\omega^0\|_{H^1} \right),
 \end{aligned}$$

where we use (3.11) in the last inequality. Similarly,

$$\begin{aligned} III &= - \int_{\mathbb{T}_\alpha} U_n^v \cdot \nabla (\omega - \psi) \omega_s^v dx dy \\ &\leq C \|U_n^v\|_{L^\infty} \left(\|\omega_n^v\|_{H^1} + \|\omega^0\|_{H^1} \right) \|\omega_s^v\|_{L^2} \\ &\leq C d v \left(\|\omega_n^v\|_{H^1}^2 + \|\omega^0\|_{H^1} \|\omega_n^v\|_{H^1} \right), \end{aligned}$$

where the embedding $\|U_n^v\|_{L^\infty} \leq C \|\omega_n^v\|_{H^1}$ and the bound (3.11) are used in the last inequality above. Combining the above estimates, we get

$$\begin{aligned} \frac{d}{dt} \frac{1}{2} \|\omega^v - \omega^0\|_X^2 &\leq v \left(-(c_0 - Cd) \|\omega_n^v\|_{H^1}^2 + C(1+t) \|\omega_n^v\|_{H^1} \|\omega^0\|_{H^1} \right) \\ &\leq v \left(-\frac{1}{2} c_0 \|\omega_n^v\|_{H^1}^2 + C(1+t) \|\omega_n^v\|_{H^1} \|\omega^0\|_{H^1} \right) \\ &\leq C v \left(1 + t^2 \right) \|\omega^0(t)\|_{H^1}^2, \end{aligned}$$

by choosing $d \leq \frac{c_0}{2C}$. \square

By using Lemmas 3.1 and 3.3, Theorem 1.2 i) follows by the same arguments as in the proof of Theorem 1.1.

Remark 3.1. By (3.8) and (3.10), we have the following Liapunov stability result:

$$\|\omega^v(t) - P_{\mathcal{K}}\omega^v(0)\|_{L^2} \leq C \|(I - P_{\mathcal{K}})\omega^v(0)\|_{L^2}, \quad \forall t > 0, \tag{3.13}$$

for some constant $C > 0$ and any solution $\omega^v(t)$ of the NS equation (3.1). Thus for initial data $\omega^v(0)$ satisfying

$$\|(I - P_{\mathcal{K}})\omega^v(0)\|_{L^2} \leq \frac{1}{C} d v, \tag{3.14}$$

we can repeatedly use Theorem 1.2 i) to get the rapid decay of the non-shear part, before the dissipation term takes over.

3.2. The Case of Square Torus

In this subsection, we prove Theorem 1.2 ii) for the nonlinear Navier–Stokes equation (3.1) on the square torus \mathbb{T}_1 . As in the rectangular torus case, we consider initial data satisfying $P_{\mathcal{K}}\omega(0) = 0$. Compared with the rectangular torus, the new difficulty is the existence of anomalous modes $\{\cos x, \sin x\}$ in the kernel of $L = 1 + \Delta^{-1}$. We decompose the vorticity perturbation as

$$\omega^v = \omega_s^v + \omega_n^v = \omega_{s1}^v + \omega_{s2}^v + \omega_{n1}^v + \omega_{n2}^v, \tag{3.15}$$

where the shear part ω_s^v is decomposed as in the rectangular case with $\omega_{s1}^v = P_{\mathcal{K}}\omega_s^v$ and $\omega_{s2}^v \in Y_1$, and the non-shear part ω_n^v is decomposed as $\omega_n^v = \omega_{n1}^v + \omega_{n2}^v$ with $\omega_{n1}^v = P_a\omega_n^v$ and $\omega_{n2}^v \in X_1$. Here, we recall that P_a is the projection

to the anomalous space W_a spanned by $\{\cos x, \sin x\}$ and X_1 is the orthogonal complement of W_a in X . Correspondingly, the velocity U^v and stream function ψ^v are decomposed into four parts. Then the nonlinear term can be written as

$$U^v \cdot \nabla \omega^v = (U_{s1}^v + U_{n1}^v) \cdot \nabla (\omega_{n2}^v - \psi_{n2}^v) + U_{n1}^v \cdot \nabla (\omega_{s2}^v - \psi_{s2}^v) + U_{s2}^v \cdot \nabla \omega_{n2}^v + U_{n2}^v \cdot \nabla \omega_{s2}^v + U_{n2}^v \cdot \nabla \omega_{n2}^v, \tag{3.16}$$

where we use the observation

$$U_{s1}^v \cdot \nabla \omega_n^v + U_n^v \cdot \nabla \omega_{s1}^v = U_{s1}^v \cdot \nabla (\omega_n^v - \psi_n^v) = U_{s1}^v \cdot \nabla (\omega_{n2}^v - \psi_{n2}^v);$$

we proceed similarly for other terms. The dissipation law (3.7) becomes

$$\begin{aligned} & \frac{d}{dt} \left(\int_{\mathbb{T}_1} (|\omega_{s2}^v|^2 - |\partial_y \psi_{s2}^v|^2) dx dy + \|\omega_{n2}^v\|_X^2 \right) \\ &= -2\nu \left(\int_{\mathbb{T}_1} (|\partial_y \omega_{s2}^v|^2 - |\omega_{s2}^v|^2) dx dy + \|\omega_{n2}^v\|_{X^1}^2 \right). \end{aligned} \tag{3.17}$$

By using the positivity of the above functional on Y_1 and X_1 , this implies that

$$\|\omega_{s2}^v\|_{L^2}(t) + \|\omega_{n2}^v\|_{L^2} \leq C (\|\omega_{s2}^v\|_{L^2}(0) + \|\omega_{n2}^v\|_{L^2}(0)) \leq Cdv. \tag{3.18}$$

In the following lemma, we separate the dissipation law for ω_{n2}^v from (3.17):

Lemma 3.4. *There exists a constant $d > 0$, such that when $\|\omega^v\|_{L^2}(0) \leq dv$, then*

$$\frac{d}{dt} \|\omega_{n2}^v\|_X^2 \leq -\nu \|\omega_{n2}^v\|_{X^1} (\|\omega_{n2}^v\|_{X^1} - \|\omega_{n1}^v\|_{L^2}). \tag{3.19}$$

Proof. By (3.24), we have

$$\begin{aligned} & \frac{d}{dt} \int_{\mathbb{T}_1} (|\omega_{s2}^v|^2 - |\partial_y \psi_{s2}^v|^2) dx dy \\ &= 2 \int_{\mathbb{T}_1} \partial_t \omega_{s2}^v (\omega_{s2}^v - \psi_{s2}^v) dx dy = 2 \int_{\mathbb{T}_1} \partial_t \omega_s^v (\omega_{s2}^v - \psi_{s2}^v) dx dy \\ &= -2\nu \int_{\mathbb{T}_1} (|\partial_y \omega_{s2}^v|^2 - |\omega_{s2}^v|^2) dx dy \\ & \quad + 2 \int_{\mathbb{T}_1} (U_{n1}^v \cdot \nabla (\omega_{n2}^v - \psi_{n2}^v) + U_{n2}^v \cdot \nabla \omega_{n2}^v) (\omega_{s2}^v - \psi_{s2}^v) dx dy \\ &\geq -2\nu \int_{\mathbb{T}_1} (|\partial_y \omega_{s2}^v|^2 - |\omega_{s2}^v|^2) dx dy \\ & \quad - C \left(\|\omega_{n1}^v\|_{L^2} \|\omega_{n2}^v\|_{X^1} + \|\omega_{n2}^v\|_{X^1}^2 \right) \|\omega_{s2}^v\|_{L^2} \\ &\geq -2\nu \int_{\mathbb{T}_1} (|\partial_y \omega_{s2}^v|^2 - |\omega_{s2}^v|^2) dx dy \\ & \quad - Cdv \left(\|\omega_{n1}^v\|_{L^2} \|\omega_{n2}^v\|_{X^1} + \|\omega_{n2}^v\|_{X^1}^2 \right). \end{aligned}$$

By choosing d such that $Cd \leq 1$, (3.19) follows from above and (3.17). \square

Compared to the dissipation law (3.12) for the rectangular torus, from (3.19) we cannot even infer that $\|\omega_{n2}^v\|_X^2$ is decreasing due to the interaction of ω_{n2}^v and ω_{n1}^v .

To prove Theorem 1.2 ii), it suffices to show that when $\|\omega^v(0)\|_{L^2} \leq d\nu$ and

$$\max_{0 \leq t \leq \frac{\tau}{\nu}} \|\omega_{n1}^v(t)\|_{L^2} \leq M \|\omega_n^v(0)\|_{L^2}, \tag{3.20}$$

we have

$$\inf_{0 \leq t \leq \frac{\tau}{\nu}} \|\omega_{n2}^v(t)\|_{L^2} < \delta \|\omega_n^v(0)\|_{L^2} \tag{3.21}$$

for ν small enough. In the next two lemmas, we derive some estimates to be used later based on the assumption (3.20).

Lemma 3.5. *Assume $\|\omega^v(0)\|_{L^2} \leq d\nu$ and (3.20), then*

$$\|\omega_{s1}^v\|_{L^2}(t) \leq C'\nu, \quad 0 \leq t \leq \frac{\tau}{\nu} \tag{3.22}$$

for some constant C' depending on d and M .

Proof. Denote

$$\omega_{s1}^v = a_1 \frac{1}{\sqrt{2\pi}} \cos y + a_2 \frac{1}{\sqrt{2\pi}} \sin y, \quad \omega_{n1}^v = b_1 \frac{1}{\sqrt{2\pi}} \cos x + b_2 \frac{1}{\sqrt{2\pi}} \sin x,$$

and

$$a(t) = \|\omega_{s1}^v\|_{L^2}(t) = \sqrt{a_1^2 + a_2^2}, \quad b(t) = \|\omega_{n1}^v\|_{L^2}(t) = \sqrt{b_1^2 + b_2^2}.$$

Then by (3.20),

$$b(t) \leq M \|\omega_n^v(0)\|_{L^2} \leq M d\nu, \quad 0 \leq t \leq \frac{\tau}{\nu}. \tag{3.23}$$

By (3.16), we have

$$\begin{aligned} \partial_t \omega_s^v &= \nu \partial_{yy} \omega_s^v + P_0(U_n^v \cdot \nabla \omega_n^v) \\ &= \nu \partial_{yy} \omega_s^v + P_0(U_{n1}^v \cdot \nabla (\omega_{n2}^v - \psi_{n2}^v) + U_{n2}^v \cdot \nabla \omega_{n2}^v). \end{aligned} \tag{3.24}$$

Projecting above to $\{\cos y, \sin y\}$ and using (3.18), we get

$$\frac{da_i}{dt} \leq -\nu a_i(t) + C d\nu (b(t) + \|\omega_{n2}^v\|_{L^2}), \quad i = 1, 2,$$

and thus

$$\frac{da}{dt} \leq -\nu a(t) + C d\nu (b(t) + \|\omega_{n2}^v\|_{L^2}(t)). \tag{3.25}$$

Since $a(0) = \|P_{\mathcal{K}}\omega(0)\|_{L^2} = 0$, we have

$$a(t) \leq C d\nu \int_0^t e^{-\nu(t-s)} (b(s) + \|\omega_{n2}^v\|_{L^2}(s)) ds \leq C'(d, M)\nu,$$

by (3.18) and (3.23). \square

Remark 3.2. In the rectangular torus case, $\|\omega^v(0)\|_{L^2} = O(v)$ implies that $\|\omega_{s1}^v\|_{L^2}(t) = O(v)$ (see (3.10)). However, for the square torus case, it is not clear that $\|\omega^v(0)\|_{L^2} = O(v)$ can imply

$$\|\omega_{s1}^v\|_{L^2}(t), \|\omega_{n1}^v\|_{L^2}(t) = O(v),$$

due to the existence of anomalous modes. Indeed, projecting (3.1) to $\{\cos x, \sin x\}$, we get

$$\begin{aligned} \frac{db_1}{dt} &= -vb_1 - \frac{1}{2\sqrt{2}\pi} e^{-vt} \int_{\mathbb{T}_1} \omega_{n2}^v \sin y \sin x \, dx dy \\ &\quad + \frac{1}{\sqrt{2}\pi} \int_{\mathbb{T}_1} (U^v \cdot \nabla \omega^v) \cos x \, dx dy \\ &\leq -vb_1 + C \|\omega_{n2}^v\|_{L^2} + Cdv(a(t) + \|\omega_{n2}^v\|_{L^2}) \\ &\leq -vb_1 + C \|\omega_{n2}^v\|_{L^2} + Cdv a(t). \end{aligned}$$

In the above, the second term is due to the identity

$$\begin{aligned} &\int_{\mathbb{T}_1} [\sin y \partial_x (1 + \Delta^{-1})] \omega^v \cos x \, dx dy \\ &= \int_{\mathbb{T}_1} [\sin y \partial_x (1 + \Delta^{-1})] \omega_{n2}^v \cos x \, dx dy \\ &= \int_{\mathbb{T}_1} \omega_{n2}^v [(1 + \Delta^{-1})(\sin y \sin x)] \, dx dy \\ &= \frac{1}{2} \int_{\mathbb{T}_1} \omega_{n2}^v \sin y \sin x \, dx dy, \end{aligned}$$

and the third term is estimated as

$$\left| \int_{\mathbb{T}_1} (U^v \cdot \nabla \omega^v) \cos x \, dx dy \right| \leq Cdv(a(t) + \|\omega_{n2}^v\|_{L^2}),$$

by using (3.16), integration by parts and the bound (3.18). Similarly,

$$\frac{db_2}{dt} \leq -vb_2 + C \|\omega_{n2}^v\|_{L^2} + Cdv a(t),$$

and thus

$$\frac{db}{dt} \leq -vb + C \|\omega_{n2}^v\|_{L^2} + Cdv a(t).$$

Combining the above with (3.25) and choosing d to be small, we have

$$\frac{de}{dt} \leq -\frac{1}{2}ve(t) + C \|\omega_{n2}^v\|_{L^2}, \quad e(t) = a(t) + b(t),$$

from which only the bound $e(t) \leq C$ can be derived. This also partly explains why the assumption (3.20) is needed in the proof of the enhanced damping.

Our next lemma is to estimate the difference of ω_n^v and the solution of the linearized Euler equation.

Lemma 3.6. *There exists $d > 0$ such that for any solution $\omega^v(t)$ of the Navier–Stokes equation (3.3) on \mathbb{T}_1 satisfying $\|\omega^v\|_{L^2}(0) \leq dv$ and (3.20), and any solution $\omega^0(t)$ of the linearized Euler equation (2.4) with initial data $\omega^0(0) \in X_1 \cap H^1$, we have*

$$\frac{d}{dt} \|\omega_{n2}^v - \omega^0\|_X^2 \leq C_0 v \left((1 + t^2) \|\omega^0(t)\|_{H^1}^2 + \|\omega_{n1}^v\|_{L^2}^2 + \|\omega_{n1}^v\|_{L^2} \|\omega^0\|_{H^1} \right), \tag{3.26}$$

$\forall t \in (0, \frac{t}{v})$, for some constant $C_0 > 0$ depending only on d and M . Here, the notations in the decomposition (3.15) for ω^v is used.

Proof. Denote $\omega = \omega_{n2}^v - \omega^0$ and $\psi = \psi_{n2}^v - \psi^0$, then

$$\begin{aligned} \omega_t = & -e^{-vt} (I - P_a) \sin y \partial_x (\omega - \psi) - (e^{-vt} - 1) (I - P_a) \sin y \partial_x (\omega^0 - \psi^0) \\ & + v \Delta \omega_{n2}^v \\ & + (I - P_a) P_{\neq 0} \left((U_{s1}^v + U_{n1}^v) \cdot \nabla (\omega_{n2}^v - \psi_{n2}^v) \right) \\ & + (I - P_a) (U_{n1}^v \cdot \nabla (\omega_{s2}^v - \psi_{s2}^v)) \\ & + (I - P_a) (U_{s2}^v \cdot \nabla \omega_{n2}^v + U_{n2}^v \cdot \nabla \omega_{s2}^v) \\ & + (I - P_a) P_{\neq 0} (U_{n2}^v \cdot \nabla \omega_{n2}^v). \end{aligned}$$

Thus we have

$$\begin{aligned} \frac{d}{dt} \frac{1}{2} \|\omega_{n2}^v - \omega^0\|_X^2 &= \int_{\mathbb{T}_1} \omega_t (\omega - \psi) \, dx dy \\ &= \left[- \int_{\mathbb{T}_1} (e^{-vt} - 1) \sin y \partial_x (\omega^0 - \psi^0) (\omega - \psi) \, dx dy \right. \\ &\quad \left. + v \int_{\mathbb{T}_1} \Delta \omega_{n2}^v (\omega - \psi) \, dx dy \right] \\ &\quad + \left[\int_{\mathbb{T}_1} (U_{s2}^v \cdot \nabla \omega_{n2}^v + U_{n2}^v \cdot \nabla \omega_{s2}^v) (\omega - \psi) \, dx dy \right. \\ &\quad \left. + \int_{\mathbb{T}_1} U_{n2}^v \cdot \nabla \omega_{n2}^v (\omega - \psi) \, dx dy \right] \\ &\quad - \left[\int_{\mathbb{T}_1} (U_{s1}^v + U_{n1}^v) \cdot \nabla (\omega_{n2}^v - \psi_{n2}^v) (\omega^0 - \psi^0) \, dx dy \right] \\ &\quad + \int_{\mathbb{T}_1} U_{n1}^v \cdot \nabla (\omega_{s2}^v - \psi_{s2}^v) (\omega - \psi) \, dx dy \\ &= I + II + III + IV. \end{aligned}$$

For the first three terms, by using the bounds (3.20) and (3.22), as in the proof of Lemma 3.3, we get

$$I + II + III \leq -\frac{1}{2} c_0 v \|\omega_{n2}^v\|_{H^1}^2 + C v (1 + t) \|\omega_{n2}^v\|_{H^1} \|\omega^0\|_{H^1}$$

by choosing $d > 0$ small enough. The last term is estimated by

$$\begin{aligned} IV &= - \int_{\mathbb{T}_1} U_{n1}^v \cdot \nabla \left[(\omega_{n2}^v - \psi_{n2}^v) - (\omega^0 - \psi^0) \right] (\omega_{s2}^v - \psi_{s2}^v) \, dx dy \\ &\leq C v \|\omega_{n1}^v\|_{L^2} \left(\|\omega_{n2}^v\|_{H^1} + \|\omega^0\|_{H^1} \right). \end{aligned}$$

Combining the above, we have

$$\frac{d}{dt} \|\omega_{n2}^v - \omega^0\|_X^2 \leq C_0 v \left((1 + t^2) \|\omega^0\|_{H^1}^2 + \|\omega_{n1}^v\|_{L^2}^2 + \|\omega_{n1}^v\|_{L^2} \|\omega^0\|_{H^1} \right)$$

for some constant C_0 depending on d and M . \square

We are now ready to prove Theorem 1.2 ii).

Proof of Theorem 1.2 ii). For any fixed $M, \delta, \tau > 0$, we will show that under the assumption (3.20), there exists $\nu(M, \delta, \tau) > 0$ such that (3.21) holds when $0 < \nu < \nu(M, \delta, \tau)$. Suppose that (3.21) is not true. Then

$$\|\omega_{n2}^v(t)\|_{L^2} \geq \delta \|\omega_n^v(0)\|_{L^2}, \quad \forall 0 \leq t \leq \frac{\tau}{\nu}, \tag{3.27}$$

where $\omega^v(t)$ is the solution of (3.1) satisfying $P_{\mathcal{K}}\omega^v(0) = 0, \|\omega^v(0)\|_{L^2} \leq d\nu$ and (3.20). Here, d is a constant chosen such that Lemmas 3.4 and 3.6 hold true. Then by (3.20) and (3.27), for any $t, t' \in (0, \frac{\tau}{\nu})$, we have

$$\begin{aligned} \|\omega_{n1}^v(t)\|_{L^2} &\leq M \|\omega_n^v(0)\|_{L^2}^2 \leq \frac{1}{\delta} M \|\omega_{n2}^v(t')\|_{L^2} \\ &\leq \frac{c_0}{\delta} M \|\omega_{n2}^v(t')\|_X. \end{aligned} \tag{3.28}$$

Choose λ_N big enough such that

$$\exp\left(-\left(\frac{\lambda_N}{2} - \sqrt{\frac{\lambda_N}{2}} \frac{c_0}{\delta} M\right) \tau\right) < c_0 \delta^2,$$

where c_0 is the constant in (2.12). We consider two cases below.

Case 1: $\omega_{n2}^v(t) \in R(I - P_N)$ is true for t in some interval $(a, b) \subset (0, \tau/\nu)$. That is,

$$\|\omega_{n2}^v(t)\|_{X^1}^2 > \lambda_N \|\omega_{n2}^v(t)\|_X^2, \quad t \in (a, b).$$

Then, by (3.19) and (3.28), we have

$$\|\omega^v(b)\|_X^2 \leq \exp\left(-\nu\sqrt{\lambda_N}\left(\sqrt{\lambda_N} - \frac{c_0}{\delta} M\right)(b-a)\right) \|\omega^v(a)\|_X^2. \tag{3.29}$$

Case 2: Let t_0 be any point in $(0, \tau/\nu)$ satisfying

$$\|\omega_{n2}^v(t_0)\|_{X^1}^2 \leq \lambda_N \|\omega_{n2}^v(t_0)\|_X^2.$$

Denote $t_1 = T_c(N, \frac{1}{10})$ to be such that the RAGE Lemma 2.14 is true for $\kappa = 1/10$ and all $T \geq T_c$. Let $\omega^0(t)$ ($t \in [t_0, t_0 + t_1]$) be the solution of (2.4) with $\omega^0(t_0) = \omega_{n_2}^v(t_0)$. By (3.26), (3.28) and Lemma 2.8, we have

$$\begin{aligned} \left\| \omega_{n_2}^v(t_0 + t) - \omega^0(t_0 + t) \right\|_X^2 &\leq C_2 v \left\{ \left(1 + t^5\right) \left\| \omega_{n_2}^v(t_0) \right\|_{X^1}^2 \right. \\ &\quad + \left(\frac{c_0}{\delta} \right)^2 M^2 \left\| \omega_{n_2}^v(t_0) \right\|_X^2 \\ &\quad \left. + \frac{c_0}{\delta} M \left(1 + t^2\right) \left\| \omega_{n_2}^v(t_0) \right\|_X \left\| \omega_{n_2}^v(t_0) \right\|_{X^1} \right\} \end{aligned} \tag{3.30}$$

for any $t \in [0, t_1]$ and some constant $C_2 > 0$ independent of τ, v . Let $\nu(M, \delta, \tau)$ be such that

$$C_2 \nu(\delta, \tau) \left\{ \left(1 + t_1^5\right) \lambda_N + \left(\frac{c_0}{\delta} \right)^2 M^2 + \frac{c_0}{\delta} M \left(1 + t_1^2\right) \sqrt{\lambda_N} \right\} < \frac{1}{10}.$$

By (3.30), when $0 < v < \nu(M, \delta, \tau)$, we have

$$\left\| \omega_{n_2}^v(t) - \omega^0(t) \right\|_X^2 \leq \frac{1}{10} \left\| \omega_{n_2}^v(t_0) \right\|_X^2, \quad \forall t \in [t_0, t_0 + t_1].$$

Then as in the proof of Theorem 1.1, by using the RAGE theorem for $\omega^0(t)$, we obtain

$$\int_{t_0}^{t_0+t_1} \left\| \omega_{n_2}^v(t) \right\|_{X^1}^2 dt \geq \frac{\lambda_N t_1}{2} \left\| \omega_{n_2}^v(t_0) \right\|_X^2.$$

Thus by (3.19) and (3.28), we have

$$\begin{aligned} \left\| \omega_{n_2}^v(t_0 + t_1) \right\|_X^2 &\leq \left\| \omega_{n_2}^v(t_0) \right\|_X^2 - \nu \int_{t_0}^{t_0+t_1} \left\| \omega_{n_2}^v \right\|_{X^1} \left(\left\| \omega_{n_2}^v \right\|_{X^1} - \left\| \omega_{n_1}^v \right\|_{L^2} \right) dt \\ &\leq \left\| \omega_{n_2}^v(t_0) \right\|_X^2 - \nu \int_{t_0}^{t_0+t_1} \left\| \omega_{n_2}^v \right\|_{X^1}^2 dt \\ &\quad + \nu \left(\int_{t_0}^{t_0+t_1} \left\| \omega_{n_2}^v \right\|_{X^1}^2 dt \right)^{\frac{1}{2}} \sqrt{t_1} \frac{c_0}{\delta} M \left\| \omega_{n_2}^v(t_0) \right\|_X \\ &\leq \left\| \omega_{n_2}^v(t_0) \right\|_X^2 \left(1 - \frac{\lambda_N}{2} + \sqrt{\frac{\lambda_N}{2}} \frac{c_0}{\delta} M \right) \nu t_1. \\ &\leq \left\| \omega_{n_2}^v(t_0) \right\|_X^2 \exp \left(- \left(\frac{\lambda_N}{2} - \sqrt{\frac{\lambda_N}{2}} \frac{c_0}{\delta} M \right) \nu t_1 \right). \end{aligned} \tag{3.31}$$

Splitting the interval $[0, \frac{\tau}{v}]$ into a union of intervals such that either Case 1 or Case 2 is true, then we have

$$\begin{aligned} \left\| \omega_{n_2}^v \left(\frac{\tau}{v} \right) \right\|_X^2 &\leq \exp \left(- \left(\frac{\lambda_N}{2} - \sqrt{\frac{\lambda_N}{2}} \frac{c_0}{\delta} M \right) \tau \right) \left\| \omega_{n_2}^v(0) \right\|_X^2 \\ &< c_0 \delta^2 \left\| \omega_{n_2}^v(0) \right\|_X^2. \end{aligned}$$

This implies that

$$\left\| \omega_{n2}^\nu \left(\frac{\tau}{\nu} \right) \right\|_{L^2} < \delta \left\| \omega_{n2}^\nu(0) \right\|_{L^2} \leq \delta \left\| \omega_n^\nu(0) \right\|_{L^2},$$

which is in contradiction to the assumption (3.27). This finishes the proof of Theorem 1.2 ii). \square

Remark 3.3. In the statement (1.10) of Theorem 1.2 ii), the non-shear part removing anomalous modes is reduced to a factor δ of the initial norm of the whole non-shear part. This is different from the result (1.7) for the linearized NS equation on a square torus, where the anomalous modes can be separated. The nonlinear coupling due to the anomalous modes can be seen from the term $U_{n1}^\nu \cdot \nabla (\omega_{s2}^\nu - \psi_{s2}^\nu)$ in (3.16), which reflects the nonlinear interaction of the anomalous modes and the shear part. For the rectangular torus, there is no such interaction term and the nonlinear enhanced damping result is consistent with that for the linearized NS equation.

3.3. Further Issues and Dipole States

We comment on some further issues. First, it would be interesting to enlarge the metastability basin from $O(\nu)$ in Theorem 1.2 to be $O(\nu^\alpha)$ ($0 < \alpha < 1$) or independent of ν if possible. Also, it is desirable to improve the decay time scale from $O(\tau/\nu)$ to $O(1/\sqrt{\nu})$ as given in [3] for the approximated linearized equation (1.4). This might require us to work on initial data of higher regularity. We note that the time scale $O(1/\sqrt{\nu})$ in [3] was obtained for initial data in H^1 .

Numerical simulations ([6]) suggested that on the rectangular torus the bar states (i.e. Kolmogorov flows) are usually observed. However, on the square torus [23], the dipole states of the form $\omega_0 = \cos x + \cos y$ or $\sin x + \sin y$ appear more often than the bar states. These dipole states are maximum entropy solutions of the 2D Euler equation, and hence are likely candidates for relevant quasi-stationary states according to the statistical approaches of 2D turbulence (e.g. [20]). The dipole states represent nonparallel flows with saddle points on the stream lines and therefore are more difficult to study. Consider the dipole with $\omega_0 = \cos x + \cos y$, then the quasi-stationary Navier–Stokes solution is $\omega^\nu(t, x, y) = e^{-\nu t} (\cos x + \cos y)$. The linearized NS equation around it becomes

$$\partial_t \omega = \nu \Delta \omega + e^{-\nu t} \left[(\sin y \partial_x - \sin x \partial_y) \left(1 + \Delta^{-1} \right) \right] \omega. \tag{3.32}$$

There are some similarities with the linearized equation (1.3) near bar states. First, the same dissipation law (1.12) holds true for (3.32). The linearized Euler operator is of the Hamiltonian form

$$(\sin y \partial_x - \sin x \partial_y) \left(1 + \Delta^{-1} \right) = JL, \tag{3.33}$$

with

$$J = \sin y \partial_x - \sin x \partial_y, \quad L = 1 + \Delta^{-1}.$$

Consider the energy space X to be the set of L^2 functions with zero mean. Define P to be the projection of L^2 to $\ker J$. It was shown in [11] that for any $\phi \in L^2$,

$$P\phi|_{\gamma_i(c)} = \frac{\oint_{\gamma_i(c)} \frac{\phi(x,y)}{|\nabla\psi_0|} dl}{\oint_{\gamma_i(c)} \frac{1}{|\nabla\psi_0|} dl}, \tag{3.34}$$

where c is in the range of $\psi_0 = \cos x + \cos y$ and $\gamma_i(c)$ is a branch of $\{\psi_0 = c\}$. Define the operator $A, A_0 : H^2 \cap X \rightarrow X$ by

$$A = -\Delta - 1 + P, \quad A_0 = -\Delta - 1.$$

We note that $A \geq A_0 \geq 0$ and

$$\ker A_0 = \ker L = \{\cos x, \sin x, \cos y, \sin y\}.$$

Therefore $\ker A \subset \ker A_0$ and by Proposition 2.8 and Lemma 11.3 of [15], we have the decomposition

$$L^2 = \ker(JL) + R(J). \tag{3.35}$$

Here, $\ker(JL) \cap R(J) \subset \ker L$ and $\ker(JL), R(J)$ are both invariant under JL . The space $\ker(JL)$ corresponds to the steady solution space of the linearized Euler equation $\partial_t \omega = JL\omega$. Different from the case of bar states where $\ker(JL)$ is the space of shear flows, for the dipole states the steady space $\ker(JL)$ has a more complicated structure. We can restrict the Euler semigroup e^{tJL} on the invariant subspace $R(J)$. Denote P_3 to be the orthogonal projection of $L^2(\mathbb{T}_1)$ to

$$\ker L = \{\cos x, \sin x, \cos y, \sin y\}.$$

We have the following RAGE type result for JL on $R(J)$:

Proposition 3.1. *Suppose that the operator JL defined in (3.33) has no nonzero purely imaginary eigenvalues. Let B be any compact operator in $L^2(\mathbb{T}_1)$. Then for any $\omega(0) \in R(J)$, we have*

$$\frac{1}{T} \int_0^T \left\| B(I - P_3) e^{itL} \omega(0) \right\|_{L^2}^2 dt \rightarrow 0, \quad \text{when } T \rightarrow \infty. \tag{3.36}$$

Proof. The proof is similar to Lemma 2.7 for bar states. We only sketch it. For any solution $\omega(t)$ of the equation $\partial_t \omega = JL\omega$ with $\omega(0) \in R(J)$, define $\omega_1(t) = (I - P_3)\omega(t)$ and let $X_1 = (I - P_3)R(J)$. Then $\omega_1(t)$ satisfies the equation $\partial_t \omega_1 = (I - P_3)JL\omega$ on the space X_1 . Since $L|_{X_1} > 0$, the operator $(I - P_3)JL$ is anti-selfadjoint on $(X_1, \langle L \cdot, \cdot \rangle)$. Our assumption on the spectrum of JL implies that $(I - P_3)JL$ has no nonzero purely imaginary eigenvalues. To show that the operator $(I - P_3)JL$ has purely continuous spectrum on X_1 , it remains to show that 0 is not an embedded eigenvalue of $(I - P_3)JL$. Supposing otherwise, there exists $0 \neq \omega \in X_1$ such that $(I - P_3)JL\omega = 0$. Let $\psi_1 = L\omega$, then $J\psi_1 = P_3JL\omega \in \ker L$. Denote

$$J\psi_1 = a_1 \cos x + a_2 \cos y + b_1 \sin x + b_2 \sin y. \tag{3.37}$$

Since $\cos y + \cos x \in \ker J \perp J\psi_1$, we have $a_1 + a_2 = 0$. Let $a_1 = -a_2 = a$. It is easy to see that for any $c \neq 0$ in the range of $\psi_0 = \cos x + \cos y$, each of the two branches of $\{\psi_0 = c\}$ is symmetric to x, y in the sense that both (x, y) and (y, x) are on the branch. Since any function ϕ in $\ker J$ takes the form (3.34), we conclude that $\cos x - \cos y \perp \ker J$, which implies that $\cos x - \cos y \in R(J)$, so there exists a double periodic function ψ_2 such that $J\psi_2 = \cos x - \cos y$. By noting that $Jx = \sin y$ and $Jy = -\sin x$, it follows from (3.37) that $J(\psi_1 - a\psi_2 + b_1y - b_2x) = 0$. Then by (3.34), the function $\psi_1 - a\psi_2 + b_1y - b_2x$ must take constant on each branch of the level set $\{\psi_0 = c\}$. Since ψ_0 is double periodic in x, y , this implies that $\psi_1 - a\psi_2 + b_1y - b_2x$ is also double periodic. This contradiction shows that 0 is not an embedded eigenvalue of $(I - P_3)JL\omega$. Then (3.36) follows from the standard RAGE theorem. \square

Even with the dissipation law (1.12) and above RAGE theorem, there are still significant differences with the bar states to get the linear enhanced damping for dipoles, besides the issue of proving the non-existence of imaginary eigenvalues of JL . The most important difference is that the decomposition (3.35) is no longer invariant when the viscosity is added. It is under investigation to find a subspace of initial data such that the enhanced damping is true for dipoles.

4. Linear Inviscid Damping of Shear Flows

Consider a shear flow $u_0 = (U(y), 0)$ in a channel $\{y_1 \leq y \leq y_2\}$ or on a torus. The linearized Euler equation can be written as

$$\omega_t + U(y) \partial_x \omega + U''(y) \partial_x \psi = 0, \tag{4.1}$$

where ω and $\psi = (-\Delta)^{-1} \omega$ are the vorticity and stream functions respectively.

4.1. Stable Case

We consider two classes of stable shear flows.

Class 1: $U'' \neq 0$, that is, U has no inflection points. This case is restricted to a channel, since such flows can not exist on a torus. By the classical Rayleigh inflection point theorem, $(U(y), 0)$ is linearly stable. Suppose $U'' > 0$, choose a constant $U_s < \min U$. Then in the frame $(x - U_s t, y, t)$, the Eq.(4.1) becomes

$$\omega_t + (U(y) - U_s) \partial_x \omega + U''(y) \partial_x \psi = 0. \tag{4.2}$$

Define

$$K_1(y) = \frac{U''(y)}{U(y) - U_s} > 0.$$

Let the x period be $2\pi/\alpha$ for any $\alpha > 0$. Define the non-shear space on the periodic channel $S_{2\pi/\alpha} \times [y_1, y_2]$ by

$$X = \left\{ \omega = \sum_{k \in \mathbf{Z}, k \neq 0} e^{ik\alpha x} \omega_k(y), \|\omega\|_X^2 = \left\| \frac{1}{\sqrt{K_1(y)}} \omega \right\|_{L^2}^2 < \infty \right\}.$$

Clearly, $X \subset L^2$ and $L^2 = X$ if $\min K_1 > 0$. The Eq.(4.2) can be written in a Hamiltonian form

$$\omega_t = -U''(y) \partial_x \left(\frac{\omega}{K_1(y)} + \psi \right) = JL\omega,$$

where

$$J = -U''(y) \partial_x : X^* \rightarrow X, \quad L = \frac{1}{K_1(y)} + (-\Delta)^{-1} : X \rightarrow X^*$$

are anti-selfadjoint and self-adjoint respectively. Moreover, L is uniformly positive on X and thus JL is anti-selfadjoint in the equivalent inner product $\langle L \cdot, \cdot \rangle$. Since $X \subset L^2$, by Lemma 4.3, JL has no purely imaginary eigenvalues in X and the entire spectrum of JL is continuous.

Case 2: Assume that there exists U_s in the range of U such that

$$K_2(y) = -\frac{U''}{U - U_s} > 0 \tag{4.3}$$

is bounded. We call these flows to be in class \mathcal{K}^+ , as used in [10]. The assumption (4.3) implies that U_s is the only inflection value of U . Examples include $U(y) = \sin y$, $\tanh y$, and more generally any $U(y)$ satisfying the ODE $U'' = g(U)$ with a decreasing g . Then (4.3) is satisfied with $U_s = g^{-1}(0)$. Let the x period be $2\pi/\alpha$. We can consider the class \mathcal{K}^+ flows in a periodic channel $S_{2\pi/\alpha} \times [y_1, y_2]$ or on a torus $S_{2\pi/\alpha} \times S_{y_2-y_1}$. The linearized Euler equation (4.2) with U_s being the inflection value can be written in the Hamiltonian form

$$\omega_t = U''(y) \partial_x \left(\frac{\omega}{K_2(y)} - \psi \right) = JL\omega, \tag{4.4}$$

where

$$J = U''(y) \partial_x, \quad L = \frac{1}{K_2(y)} - (-\Delta)^{-1}.$$

Define the non-shear space of vorticity

$$X = \left\{ \omega = \sum_{k \in \mathbf{Z}, k \neq 0} e^{ik\alpha x} \omega_k(y), \|\omega\|_X^2 = \left\| \frac{1}{\sqrt{K_2(y)}} \omega \right\|_{L^2}^2 < \infty \right\}. \tag{4.5}$$

Again, $X \subset L^2$ and $L^2 = X$ if $\min K_2 > 0$. Denote $n^-(L)$ ($n^0(L)$) to be the number of negative (zero) directions of L on X . Define the operator

$$A_0 = -\Delta - K_2(y) : H^2 \rightarrow L^2 \tag{4.6}$$

in the channel $S_{2\pi/\alpha} \times [y_1, y_2]$ or on the torus $S_{2\pi/\alpha} \times S_{y_2-y_1}$ and

$$L_0 = -\frac{d^2}{dy^2} - K_2(y) : H^2(y_1, y_2) \rightarrow L^2(y_1, y_2), \tag{4.7}$$

with the Dirichlet boundary conditions for the channel and periodic boundary conditions for the torus. Then, by Lemma 11.3 in [15], we have

$$n^0(L) = n^0(A_0) = 2 \sum_{k \geq 1} n^0(L_0 + k^2 \alpha^2),$$

and

$$n^-(L) = n^-(A_0) = 2 \sum_{k \geq 1} n^-(L_0 + k^2 \alpha^2).$$

If $n^-(L_0) \neq 0$, let $-\alpha_{\max}^2$ be the smallest eigenvalue of L_0 and ϕ_0 be an eigenfunction. When $L_0 \geq 0$, let $\alpha_{\max} = 0$. Then by the above relations, L is positive when $\alpha > \alpha_{\max}$. Again, by Lemma 4.3, the spectrum of JL is purely continuous in X . When $\alpha = \alpha_{\max}$, we have $n^-(L) = 0$ and

$$\ker L = \left\{ e^{\pm i \alpha x} \omega_0(y) \right\}, \quad \omega_0(y) = \left(-\frac{d^2}{dy^2} + \alpha^2 \right) \phi_0.$$

Let P_1 be the projection of X to $\ker L$ and $X_1 = (I - P_1)X$. Then $L|_{X_1} > 0$ and $A_1 = (I - P_1)JL$ is anti-selfadjoint on $(X_1, \langle L \cdot, \cdot \rangle)$.

Lemma 4.1. *A_1 has purely continuous spectrum on X_1 .*

Proof. By Lemma 3.5 of [10], $\phi_0 \neq 0$ at at least one of the points in the set $\{U = U_s\}$. By using this fact, the rest of the proof follows that of Lemma 2.7, so we skip the details. \square

By the above spectral properties, the following is a direct consequence of the RAGE Theorem:

Theorem 4.1. *If i) $U'' \neq 0$ or ii) U is in class \mathcal{K}^+ and $\alpha > \alpha_{\max}$, then for any compact operator B on X , we have*

$$\frac{1}{T} \int_0^T \left\| B e^{itJL} \omega \right\|_X^2 dt \rightarrow 0, \quad \text{when } T \rightarrow \infty \tag{4.8}$$

for any $\omega \in X$. If U is in class \mathcal{K}^+ and $\alpha = \alpha_{\max}$, then for any compact operator B on X , we have

$$\frac{1}{T} \int_0^T \left\| B (I - P_1) e^{itJL} \omega \right\|_X^2 dt \rightarrow 0, \quad \text{when } T \rightarrow \infty$$

for any $\omega \in X$.

By choosing

$$B\omega = \nabla^\perp (-\Delta)^{-1} \omega = u,$$

that is, the mapping from vorticity to velocity, we get

Corollary 4.1. (i) If a) $U'' \neq 0$ or b) U is in class \mathcal{K}^+ and $\alpha > \alpha_{\max}$, then

$$\frac{1}{T} \int_0^T \|u(t)\|_{L^2}^2 dt \rightarrow 0, \quad \text{when } T \rightarrow \infty$$

for any solution $\omega(t)$ of (4.1) with $\omega(0) \in X$.

(ii) If U is in class \mathcal{K}^+ and $\alpha = \alpha_{\max}$, then

$$\frac{1}{T} \int_0^T \|u_1(t)\|_{L^2}^2 dt \rightarrow 0, \quad \text{when } T \rightarrow \infty,$$

where $u_1(t)$ is the velocity corresponding to the vorticity $(I - P_1)\omega(t)$ with $\omega(0) \in X$.

Remark 4.1. More information on the decay of $\|u(t)\|_{L^2}$ could be obtained by studying the regularity of the spectral measure of the anti-selfadjoint operator JL on $(X, \langle L \cdot, \cdot \rangle)$. In particular, if the spectral measure of JL is absolutely continuous (i.e. absence of singular continuous spectrum), then when $t \rightarrow \infty$, $\omega(t) \rightarrow 0$ weakly in X . As a result, $\|u(t)\|_{L^2} \rightarrow 0$ when $t \rightarrow \infty$.

Remark 4.2. Let $B = P_N$ in (4.8), i.e., the projection operator to the first N Fourier modes (in x), then

$$\frac{1}{T} \int_0^T \|P_N \omega(t)\|_{L^2}^2 dt \rightarrow 0, \quad \text{when } T \rightarrow \infty. \quad (4.9)$$

This shows that in the time averaged sense, the low frequency part of ω tends to zero. As can be seen in the proof of Theorems 1.1 and 1.2, this property plays an important role in the proof of metastability of Kolmogorov flows. In the fluid literature (see e.g. [5, 21]), a dual cascade was known for 2D turbulence that energy moves to low frequency end and the enstrophy (i.e. $\|\omega\|_{L^2}^2$) moves to the high frequency end. The result (4.9) can be seen as a justification of such physical intuition in a weak sense.

Remark 4.3. The two classes of shear flows considered above are linearly stable in the L^2 norm of vorticity (assuming $\inf K_i > 0$), in the Liapunov sense. This follows directly from the conservation of $\langle L\omega, \omega \rangle$ for the linearized Euler equation (4.2) and the positivity of $L|_X$. Moreover, these two classes seem to exhaust all the possible shear flows for which nonlinear stability could be proved by the energy-Casimir method initiated by Arnold [1, 2] in 1960s. We briefly discuss it below and refer to [17] for more discussions on energy-Casimir method for 2D Euler equations. The energy-Casimir functional is of the form

$$H(\omega) = \int \left(G(\omega) + \frac{1}{2} |\nabla \psi|^2 \right) dx dy,$$

which is invariant for the nonlinear Euler equation. Suppose $\psi_0 = F(\omega_0)$, where $\psi_0 = \int (U - U_s) dy$ and $\omega_0 = -U''$. Choose G such that $G'(\omega_0) = -F(\omega_0)$, then $H'(\omega_0) = 0$ and the second order variation is given by

$$\langle H''(\omega) \delta\omega, \delta\omega \rangle = \frac{1}{2} \int \frac{(\delta\omega)^2}{K_1(y)} + \frac{1}{2} |\nabla \delta\psi|^2 = \frac{1}{2} \langle L\delta\omega, \delta\omega \rangle$$

for $U(y)$ in class 1 and

$$\langle H''(\omega) \delta\omega, \delta\omega \rangle = \frac{1}{2} \int -\frac{(\delta\omega)^2}{K_2(y)} + \frac{1}{2} |\nabla \delta\psi|^2 = -\frac{1}{2} \langle L\delta\omega, \delta\omega \rangle$$

for $U(y)$ in class 2. In the above, we use the relation

$$G''(\omega_0) = -F'(\omega_0) = \frac{U - U_s}{U''}.$$

Thus $\langle H''(\omega) \delta\omega, \delta\omega \rangle$ on X is positive definite for class 1 flows and negative definite for class 2 flows when $\alpha > \alpha_{\max}$. Then nonlinear stability (in L^2 vorticity) could be proved by properly handling the higher order terms. However, if $\frac{U-U_s}{U''}$ (equivalently K_1, K_2) changes sign, then the quadratic form $\langle H''(\omega) \delta\omega, \delta\omega \rangle$ is highly indefinite and the energy-Casimir method does not work. Despite above restrictions, the steady flows whose stability can be studied by energy-Casimir method do appear often as observable coherent states in 2D turbulence.

4.2. Unstable Case

The shear flows $U(y)$ in class \mathcal{K}^+ are proved to be linearly unstable when the horizontal wave number $\alpha < \alpha_{\max}$, see Theorem 1.2 in [10]. In this subsection, we will prove the inviscid damping on the center space which is complementary to the stable and unstable subspaces. The proof of linear instability in [10] is by studying the possible neutral limiting modes and the bifurcation of unstable modes near them. By using the Hamiltonian structure of (4.2) and the instability index formula in [15], we can recover this linear instability criterion. Moreover, we get more detailed information about the number of unstable modes which is important to study the inviscid damping on the center space.

Proposition 4.1. *Consider $U(y)$ in class \mathcal{K}^+ and $\alpha > 0$, where $2\pi/\alpha$ is the x period. Let k_u be the total algebraic multiplicities of unstable eigenvalues of the operator JL defined in (4.4). Then $k_u = n^-(A_0)$, where A_0 is defined in (4.6).*

Proof. It is easy to see that the unstable eigenfunctions of JL are in the space X defined in (4.5). Since on the energy space X , $n^-(L) = n^-(A_0) < \infty$, we can use Theorem 2.3 in [15] to get the index formula

$$k_r + 2k_c + 2k_i^{\leq 0} + k_0^{\leq 0} = n^-(L). \tag{4.10}$$

Here, k_r and k_c are the algebraic multiplicities of unstable eigenvalues of JL lying on the positive axis and the first quadrant respectively, $k_i^{\leq 0}$ is the number of non-positive directions of L restricted to the generalized eigenspace of imaginary eigenvalues on $i\mathbf{R}^+$, and $k_0^{\leq 0}$ is the number of non-positive directions of L on $E_0/\ker L$ where E_0 is the generalized zero eigenspace of JL . Since JL has no nonzero imaginary eigenvalue, $k_i^{\leq 0} = 0$. As in the proof of Lemmas 2.7 and 4.1, it can be shown that $E_0 = \ker L$ and thus $k_0^{\leq 0} = 0$. Therefore (4.10) implies that

$$k_u = k_r + 2k_c = n^-(L) = n^-(A_0).$$

□

The space X has an invariant decomposition $X = \cup_{l \in \mathbf{Z}, l \neq 0} X^l$, where

$$X^l = \left\{ e^{i\alpha l x} \omega_l(y), \omega_l \in L^2_{\frac{1}{K_2(y)}}(y_1, y_2) \right\}.$$

On the subspace X^l , the operator JL is reduced to an ODE operator $J_l L_l$ on the weighted space $L^2_{\frac{1}{K_2(y)}}(y_1, y_2)$, where

$$J_l = U''(y) i\alpha l, \quad L_l = \frac{1}{K_2(y)} - \left(-\frac{d^2}{dy^2} + \alpha^2 l^2 \right)^{-1}. \tag{4.11}$$

We have a similar counting formula for unstable eigenvalues of $J_l L_l$.

Proposition 4.2. *Consider $U(y)$ in class \mathcal{K}^+ and $\alpha > \alpha_{\max}$. Let $k_{u,l}$ be the total algebraic multiplicities of the unstable eigenvalues of the operator $J_l L_l$ ($0 \neq l \in \mathbf{Z}$) defined in (4.11). Then*

$$k_{u,l} = n^-(L_0 + l^2 \alpha^2), \tag{4.12}$$

where L_0 is defined in (4.7).

Proof. Since J_l is not a real operator, we can not directly apply the index Theorem 2.3 in [15]. Define the space

$$\begin{aligned} Y_l &= X^l \oplus X^{-l} \\ &= \left\{ \cos(\alpha l x) \omega_1(y) + \sin(\alpha l x) \omega_2(y), \omega_1, \omega_2 \in L^2_{\frac{1}{K_2(y)}}(y_1, y_2) \right\}, \end{aligned}$$

which is isomorphic to the space $Y = \left(L^2_{\frac{1}{K_2(y)}}(y_1, y_2) \right)^2$. For any

$$\omega = \cos(\alpha l x) \omega_1(y) + \sin(\alpha l x) \omega_2(y) \in Y_l,$$

we have

$$JL\omega = (\cos(\alpha l x), \sin(\alpha l x)) J^l L^l \begin{pmatrix} \omega_1 \\ \omega_2 \end{pmatrix},$$

where

$$J^l = \begin{pmatrix} 0 & \alpha l U'' \\ -\alpha l U'' & 0 \end{pmatrix}, \quad L^l = \begin{pmatrix} L_l & 0 \\ 0 & L_l \end{pmatrix}.$$

In the above, the operator L_l is defined in (4.11). Thus to study JL on Y_l , it is equivalent to study $J^l L^l$ on Y . Let k_u^l be the total algebraic multiplicities of the

unstable eigenvalues of the operator $J^l L^l$. Then by Theorem 2.3 in [15] and the same proof of Proposition 4.2, we have

$$k_u^l = n^-(L^l) = 2n^-(L_l) = 2n^-(L_0 + l^2\alpha^2).$$

Since the spectra of $J_l L_l$ is complex conjugate of that of $J_{-l} L_{-l}$, so

$$k_u^l = k_{u,l} + k_{u,-l} = 2k_{u,l},$$

and this finishes the proof of the proposition. \square

Remark 4.4. Let $\lambda = i\alpha l c$ be an eigenvalue of $J_l L_l$ and $J_l L_l \omega = \lambda \omega$ for some $0 \neq \omega \in X^l$. Then the stream function

$$\psi(y) = \left(-\frac{d^2}{dy^2} + \alpha^2 l^2\right)^{-1} \omega(y)$$

satisfies the classical Rayleigh equation

$$\left(-\frac{d^2}{dy^2} + \alpha^2 l^2 + \frac{U''}{U - c}\right) \psi = 0 \tag{4.13}$$

with Dirichlet or periodic boundary conditions. It was shown in [8] that for $U(y)$ in class \mathcal{K}^+ , the total number (i.e. geometric multiplicities) of unstable eigenvalues (i.e. $\text{Im } c > 0$) can not exceed $n^-(L_0 + l^2\alpha^2)$. In [10], it was shown that $k_{u,l} \geq 1$ when $n^-(L_0 + l^2\alpha^2) \neq 0$. The precise index formula (4.12) not only gives an improvement over previous results, but also is important for studying the dynamics on the center space (see below).

Denote $E^s (E^u) \subset X$ to be the stable (unstable) eigenspace of JL , then

$$\dim E^s = \dim E^u = k_u = n^-(L).$$

Moreover, by Corollary 6.1 in [15], $L|_{E^s \oplus E^u}$ is non-degenerate and

$$n^-(E^s \oplus E^u) = \dim E^u = n^-(L). \tag{4.14}$$

Define the center space E^c to be the orthogonal (in the inner product $\langle L \cdot, \cdot \rangle$) complement of $E^s \oplus E^u$ in X , that is,

$$E^c = \{ \omega \in X \mid \langle L\omega, \omega_1 \rangle = 0, \quad \forall \omega_1 \in E^s \oplus E^u \}. \tag{4.15}$$

Then we have

Lemma 4.2. *The decomposition $X = E^s \oplus E^u \oplus E^c$ is invariant under JL . Moreover, $n^-(L|_{E^c}) = 0$ and as a consequence $L|_{E^c / \ker L} > 0$.*

Proof. The invariance of the decomposition follows from the invariance of $\langle L \cdot, \cdot \rangle$ under JL . By (4.14), we have

$$n^-(L|_{E^c}) = n^-(L) - n^-(E^s \oplus E^u) = 0,$$

and thus $L|_{E^c / \ker L} > 0$. \square

Since E^c is invariant under JL , we can restrict the linearized Euler equation on E^c . The linear inviscid damping still holds true for initial data in E^c . Denote P_1 to be the projection of X to $\ker L$. By the same proof of Theorem 4.1 and Corollary 4.1, we have the following:

Theorem 4.2. *Suppose $U(y)$ is in class \mathcal{K}^+ and $\alpha < \alpha_{\max}$. Then :*

(i) *If $\ker L = \{0\}$, then*

$$\frac{1}{T} \int_0^T \|u(t)\|_{L^2}^2 dt \rightarrow 0, \quad \text{when } T \rightarrow \infty,$$

for any solution $\omega(t)$ of (4.1) with $\omega(0) \in E^c$. Here, E^c is the center space defined in (4.15).

(ii) *If $\ker L \neq \{0\}$, then*

$$\frac{1}{T} \int_0^T \|u_1(t)\|_{L^2}^2 dt \rightarrow 0, \quad \text{when } T \rightarrow \infty,$$

where $u_1(t)$ is the velocity corresponding to the vorticity $(I - P_1)\omega(t)$ with $\omega(0) \in E^c$.

Remark 4.5. Above theorem suggests that the dynamics of solutions of the linearized Euler equation on the center space E^c is similar to the stable case.

The invariant decomposition $X = E^s \oplus E^u \oplus E^c$ can be used to prove the exponential trichotomy of the semigroup e^{tJL} . We refer to Theorem 2.2 in [15] for the precise statement. The next natural step is to construct invariant manifolds (stable, unstable and center) for the nonlinear Euler equation, which will give a complete description of the local dynamics near $u_0 = (U(y), 0)$. The stable and unstable manifolds near any unstable shear flow were constructed in [14]. The construction of center manifold is under investigation. Once constructed, on such center manifold, the positivity of $L|_{E^c}$ (Lemma 4.2) could be used to prove nonlinear stability of solutions on the center manifold.

Remark 4.6. Recently, in [16], the stability of class \mathcal{K}^+ shear flows under Coriolis forces was studied. By using the Hamiltonian formulation and the instability index formula, the sharp stability condition can be obtained for some shear flows. Moreover, the linear damping as in Theorems 4.1 and 4.2 was proved for non-shear initial data.

Appendix

Lemma 4.3. *Let $U(y) \in C^2(y_1, y_2)$, where $-\infty < y_1 < y_2 < \infty$. Consider the Rayleigh equation*

$$\left(-\frac{d^2}{dy^2} + \alpha^2 + \frac{U''}{U - c} \right) \psi = 0, \tag{4.16}$$

with the periodic boundary condition

$$\psi(y_1) = \psi(y_2), \quad \psi'(y_1) = \psi'(y_2),$$

or the Dirichlet boundary condition

$$\psi (y_1) = \psi (y_2) = 0.$$

If (4.16) has a neutral solution with $\alpha > 0$, $c \in \mathbf{R}$ and $\psi \in H^2 (y_1, y_2)$, then c must be an inflection value of U .

Proof. First, we show that c must be in the range of $U (y)$. Suppose otherwise $c > \max U$ or $c < \min U$. Assume $c > \max U$. For the Dirichlet boundary condition, since $U - c < 0$ in $[y_1, y_2]$, by the proof of Lemma 3.5 of [10], $\psi \equiv 0$ in $[y_1, y_2]$, which is a contradiction. For the periodic boundary condition, (4.16) implies that the operator $L_0 = -\frac{d^2}{dy^2} - \frac{U''}{U-c}$ has a negative eigenvalue $-\alpha^2$. Let $\lambda_0 \leq -\alpha^2 < 0$ be the smallest eigenvalue of L_0 , then the corresponding eigenfunction ϕ can be taken to be positive. The equation $L_0\phi = \lambda_0\phi$ can be written as

$$((U - c) \phi' - U' \phi)' = -\lambda_0 (U - c) \phi.$$

Integrating above from y_1 to y_2 and using the periodic boundary condition, we have

$$\int_{y_1}^{y_2} (U - c) \phi dy = 0,$$

which is a contradiction again. Therefore c must be in the range of U .

Let $z_1 < z_2 < \dots < z_{k_c}$ ($k_c \geq 1$) be the zeros of $U (y) - c$ in $[y_1, y_2]$. We claim that there exists $1 \leq k \leq k_c$ such that $\psi (z_k) \neq 0$. For the Dirichlet boundary condition, this follows by Lemma 3.5 of [10]. For the periodic boundary condition, suppose otherwise $\psi (z_k) = 0$ for all $k = 1, 2, \dots, k_c$. Let $z_{k_c+1} = z_1 + y_2 - y_1$ which is the translation of z_1 by one period. Then $U - c$ takes the same sign on each interval $[z_i, z_{i+1}]$, $i = 1, 2, \dots, k_c$, and $\psi = 0$ at the end points. By the proof of Lemma 3.5 of [10], it follows that $\psi \equiv 0$ in all the intervals $[z_i, z_{i+1}]$. Thus $\psi \equiv 0$ in $[y_1, y_2]$, which is a contradiction. Let $1 \leq k_0 \leq k_c$ be such that $\psi (z_{k_0}) \neq 0$. Then we must have $U'' (z_{k_0}) = 0$, that is, $c = U (z_{k_0})$ is an inflection value. Suppose otherwise, then $U'' (z_{k_0}) \neq 0$ and thus $\frac{U''}{U-c} \psi$ is not in L^2_{loc} near z_{k_0} , which is in contradiction to the Rayleigh equation (4.16) and the assumption that $\psi \in H^2$. This finishes the proof of the Lemma. \square

Remark 4.7. The above lemma shows that for general shear flows $U (y)$, any H^2 neutral mode must have its phase speed c to be one of the inflection values. This fact is used in section 4 to exclude embedded eigenvalues and to obtain the instability index formula (4.12) and the positivity of $L|_{E^c}$. In [10,12], it was shown for a class of shear flows (called class \mathcal{F} in [10]) that any neutral limiting mode (i.e. the limit of a sequence of unstable modes) must be in H^2 and therefore the phase speed must be inflection values. These neutral limiting modes are important for finding linear stability/instability criteria since they give the transition points for stability/instability.

The flows $U (y)$ in class \mathcal{F} (see [10] for definition) include any monotone flow, class \mathcal{K}^+ flows, and more generally, any $U (y)$ satisfying an ODE $U'' = k (y) g(U)$ for some $k > 0$ and any g . However, for shear flows not in class \mathcal{F} , the limiting neutral modes might be singular (i.e. not in H^2). Such singular neutral modes might have their phase speeds as c other than the inflection values.

Acknowledgements. Lin is supported in part by NSF Grants DMS-1411803 and DMS-1715201. Part of this work started when Xu visited Georgia Tech during 2013-2014 by the support of JiNan University's oversea visiting fund. He thanks Georgia Tech for its hospitality during the visit. After this paper was finished, we learned that more quantitative results for the linear enhanced damping of Kolmogorov flows were obtained in [26]. LIN THANKS DONGYI WEI, ZHIFEI ZHANG and WEIREN ZHAO for their comments and discussions. The authors also thank the anonymous referee for useful comments.

Compliance with Ethical Standards

Conflict of interest There are no conflicts of interest for the research in this paper.

References

1. ARNOLD, V.I.: On conditions for non-linear stability of plane stationary curvilinear flows of an ideal fluid. (*Russian*) *Dokl. Akad. Nauk SSSR* **162**,975–978 1965
2. ARNOLD, V.I.: An a priori estimate in the theory of hydrodynamic stability. (*Russian*) *Izv. Vysš. Učebn. Zaved. Matematika* **54**, 3–5 1966
3. BECK, M, WAYNE, C.E.: Metastability and rapid convergence to quasi-stationary bar states for the two-dimensional Navier–Stokes equations. *Proc. Roy. Soc. Edinburgh Sect. A* **143**, 905–927 2013
4. BEDROSSIAN, J., MASMOUDI, N.: Inviscid damping and the asymptotic stability of planar shear flows in the 2D Euler equations. *Publ. Math. Inst. Hautes Études Sci.* **122**, 195–300 2015
5. BOUCHET, F., VENAILLE, A.: Statistical mechanics of two-dimensional and geophysical flows. *Phys. Rep.* **5**, 227-295 2012
6. BOUCHET, F., SIMONNET, E.: Random changes of flow topology in two-dimensional and geophysical turbulence. *Phys. Rev. Lett.* **102**, 094504 2009
7. CYCON, H.L., FROESE, R.G., KIRSCH, W., SIMON, B.: *Schrödinger Operators with Application to Quantum Mechanics and Global Geometry*. Texts and Monographs in Physics. Springer, Berlin, 1987.
8. HOWARD, L.N.: The number of unstable modes in hydrodynamic stability problems. *J. Mécanique* **3**, 433–443 1964
9. CONSTANTIN, P., KISELEV, A., RYZHIK, L., ZLATOŠ, A.: Diffusion and mixing in fluid flow. *Ann. Math.* **168**(2), 643–674 2008
10. LIN, Z.: Instability of some ideal plane flows. *SIAM J. Math. Anal.* **35**, 318–356 2003
11. LIN, Z.: Some stability and instability criteria for ideal plane flows. *Commun. Math. Phys.* **246**, 87–112 2004
12. LIN, Z.: *Some Recent Results on Instability of Ideal Plane Flows, Nonlinear Partial Differential Equations and Related Analysis*. Contemp. Math., vol. 371, pp. 217–229 Amer. Math. Soc., Providence, 2005.
13. LIN, Z., ZENG, C.: Inviscid dynamic structures near Couette flow. *Arch. Ration. Mech. Anal.* **200**, 1075–1097 2011
14. LIN, Z., ZENG, C.: Unstable manifolds of Euler equations. *Commun. Pure Appl. Math.* **66**, 1803-1936 2013
15. LIN, Z., ZENG, C.: *Instability, index theorem, and exponential trichotomy for Linear Hamiltonian PDEs*. Submitted, [arXiv:1703.04016](https://arxiv.org/abs/1703.04016).
16. LIN, Z., YANG, J., ZHU, H.: Barotropic instability of shear flows. Submitted [arXiv:1801.00950](https://arxiv.org/abs/1801.00950).
17. MARCHIORO, C., PULVIRENTI, M.: *Mathematical Theory of Incompressible Nonviscous Fluids*. Springer, Berlin 1994.
18. MATTHAEUS, W.H., STRIBLING, W.T., MARTINEZ, D., OUGHTON, S., MONTGOMERY, D.: Decaying, two-dimensional, Navier–Stokes turbulence at very long times. *Physica D* **51**, 531–538 1991

19. ORR, W.M.: Stability and instability of steady motions of a perfect liquid. *Proc. Ir. Acad. Sect. A Math. Astron. Phys. Sci.* **27**, 9-66 1907
20. ROBERT, R.: *Statistical Hydrodynamics (Onsager Revisited)*. Handbook of mathematical fluid dynamics, vol. II, pp. 1–54. North-Holland, Amsterdam, 2003.
21. TABELING, P.: Two-dimensional turbulence: a physicist approach. *Phys. Rep.* **362**, 1–62 2002
22. VILLANI, C., *Hypocoercivity*. Memoirs of the American Mathematical Society. American Mathematical Society, 2009.
23. YIN, Z., MONTGOMERY, D.C., CLERCX, H.J.H.: Alternative statistical-mechanical descriptions of decaying two-dimensional turbulence in terms of “patches” and “points”. *Phys. Fluids* **15**, 1937–1953 2003
24. WEI, D., ZHANG, Z., ZHAO, W.: Linear inviscid damping for a class of monotone shear flow in Sobolev spaces. *Commun. Pure. Appl. Math.* **71**, 617-687 2018
25. WEI, D., ZHANG, Z., ZHAO, W.: *Linear Inviscid damping and vorticity depletion for shear flows*. [arXiv:1704.00428](https://arxiv.org/abs/1704.00428).
26. WEI, D., ZHANG, Z., ZHAO, W.: *Linear inviscid damping and enhanced dissipation for the Kolmogorov flow*. [arXiv:1711.01822](https://arxiv.org/abs/1711.01822).
27. ZILLINGER, C.: Linear inviscid damping for monotone shear flows in a finite periodic channel, boundary effects, blow-up and critical Sobolev regularity, *Arch. Ration. Mech. Anal.* **221**, 1449–1509 2016

ZHIWU LIN
 School of Mathematics,
 Georgia Institute of Technology,
 Atlanta,
 GA,
 30332,
 USA.
 e-mail: zlin@math.gatech.edu

and

MING XU
 Department of Mathematics,
 JiNan University,
 Guangzhou,
 510632,
 China.

(Received July 2, 2017 / Accepted September 11, 2018)

Published online September 24, 2018

© Springer-Verlag GmbH Germany, part of Springer Nature (2018)