



Global Existence and Non-Uniqueness for 3D Navier–Stokes Equations with Space-Time White Noise

MARTINA HOFMANOVÁ, RONGCHAN ZHU  & XIANGCHAN ZHU

Communicated by V. Vicol

Abstract

We establish that global-in-time existence and non-uniqueness of probabilistically strong solutions to the three dimensional Navier–Stokes system driven by space-time white noise. In this setting, solutions are expected to have space regularity of at most $-1/2 - \kappa$ for any $\kappa > 0$. Consequently, the convective term is ill-defined analytically and probabilistic renormalization is required. Up until now, only local well-posedness has been known. With the help of paracontrolled calculus we decompose the system in a way which makes it amenable to convex integration. By a careful analysis of the regularity of each term, we develop an iterative procedure which yields global non-unique probabilistically strong paracontrolled solutions. Our result applies to any divergence free initial condition in $L^2 \cup B_{\infty, \infty}^{-1+\kappa}$, $\kappa > 0$, and also implies non-uniqueness in law.

1. Introduction

Thermal fluctuations are omnipresent on the molecular level of fluids. As such, fluids are not deterministic but rather stochastic and continually changing. In order

M.H. is grateful for funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (grant agreement No. 949981) and the financial support by the DFG through the CRC 1283 “Taming uncertainty and profiting from randomness and low regularity in analysis, stochastics and their applications”. R.Z. and X.Z. are grateful to the financial supports by National Key R&D Program of China (No. 2022YFA1006300). R.Z. is grateful to the financial supports of the NSFC (No. 12271030). X.Z. is grateful to the financial supports in part by National Key R&D Program of China (No. 2020YFA0712700) and the NSFC (No. 12090014, 12288201) and the support by key Lab of Random Complex Structures and Data Science, Youth Innovation Promotion Association (2020003), Chinese Academy of Science. Rongchan Zhu is the corresponding author

to incorporate these effects into the description of large scale dynamics, Landau and Lifshitz [47] proposed a Navier–Stokes system perturbed by stochastic flux terms. These are given by a divergence of delta correlated space-time Gaussian random fields included in the momentum equation. Mathematically, it is extremely challenging to make sense of such a system. Indeed, due to the irregularity of the noise combined with the nonlinearity of the system, the equations become critical in dimension 2 and supercritical in dimension higher in the sense of the theory of regularity structures by Hairer [42]: introducing an ultraviolet cut-off, i.e. mollifying the noise, and trying to remove the cut-off would require an infinite number of renormalizations. Therefore, neither the classical stochastic analysis tools nor the pathwise theories of regularity structures [42] and paracontrolled distributions by Gubinelli, Imkeller and Perkowski [35] are applicable. Nevertheless, despite these theoretical difficulties, the Landau–Lifshitz–Navier–Stokes system has been verified numerically for various equilibrium as well as non-equilibrium systems (see e.g. [28] and references therein).

In this paper, we contribute to the rigorous mathematical understanding of microscopic perturbations in fluid dynamics. To avoid the issue of criticality mentioned above, we consider the delta correlated space-time Gaussian noise as a forcing, not as a flux. This models forcing acting on the molecular level, which translated to the macroscopic level necessarily becomes delta correlated. Indeed, any two points in the large scale dynamics are extremely far apart on the molecular level, so their associated noises must be uncorrelated. Moreover, such a noise also appears in a scaling limit of point vortex approximation and the vorticity form of the 2D Euler equations perturbed by a certain transport type noise (cf. [30,31,49]). In fact, the scaling limit is given by the vorticity form of the 2D Navier–Stokes system driven by the curl of space-time white noise, which in the velocity-pressure variables reads as 2D Navier–Stokes equations driven by space-time white noise.

The corresponding gain of one derivative in comparison to the Landau–Lifshitz setting makes the system mathematically accessible as it remains subcritical up to space dimension 3. However, even this problem has resisted rigorous mathematical analysis for a long time due to its irregularity. More precisely, the space-time white noise in spatial dimension d can be shown to be a random distribution of space-time regularity $-(d+2)/2-\kappa$ under the parabolic scaling for any $\kappa > 0$. Accordingly, in view of Schauder’s estimates a solution is expected to be two degrees of regularity better, i.e. at most $-d/2+1-\kappa$. Hence, already in $d = 2$ solutions are not functions. Consequently, the product in the convective term is analytically ill-defined and probabilistic arguments are required in order to make sense of the equations.

We consider the three dimensional Navier–Stokes system with periodic boundary conditions driven by a space-time white noise

$$\begin{aligned} du + \operatorname{div}(u \otimes u) dt + \nabla p dt &= \Delta u dt + dB, \\ \operatorname{div} u &= 0, \\ u(0) &= u_0, \end{aligned} \tag{1.1}$$

where B is a cylindrical Wiener process on some stochastic basis $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbf{P})$. The time derivative of B is the space-time white noise. Our main results reads as follows:

Theorem 1.1. *For any given divergence free initial condition $u_0 \in L^2 \cup B_{\infty, \infty}^{-1+\kappa}$ \mathbf{P} -a.s., $\kappa > 0$, there exist infinitely many global-in-time probabilistically strong solutions solving (1.1) in a paracontrolled sense.*

Remark 1.2. In what follows we prove the result for the initial condition u_0 in $B_{\infty, \infty}^{-1+\kappa}$ or $u_0 - z(0) \in L^2$ for simplicity, where z is the stationary solution to the linear equation. If we choose $z(0) = 0$, a small modification of the proof implies that the result also holds for the initial condition $u_0 \in L^2$.

The main ideas behind the definition of paracontrolled solution are explained in Sect. 1.3 and the detailed presentation can be found in Sect. 4.1. Probabilistically strong solutions means that the solutions are adapted to the normal filtration $(\mathcal{F}_t)_{t \geq 0}$ generated by the cylindrical Wiener process.

Corollary 1.3. *Non-uniqueness in law holds for (1.1) for every given initial law supported on divergence free vector fields in $L^2 \cup B_{\infty, \infty}^{-1+\kappa}$, $\kappa > 0$.*

1.1. Singular SPDEs

In two space dimensions, the problem was solved locally in time in the seminal paper by Da Prato and Debussche [20]. Furthermore, using the properties of the Gaussian invariant measure, it was possible to obtain global-in-time existence for a.e. initial condition with respect to the invariant measure. By the strong Feller property in [62] global-in-time existence for every initial condition could be derived. Recently, for a related two dimensional critical problem, tightness of approximate stationary solutions and non-triviality of the limit has been established in a weak coupling regime by Cannizzaro and Kiedrowski [16].

The more irregular three dimensional setting remained open for much longer as substantially new ideas were required. These came in a parallel development with the theory of regularity structures by Hairer [42] and with the paracontrolled distributions introduced by Gubinelli, Imkeller and Perkowski [35]. These theories permit to treat a large number of singular subcritical SPDEs (cf. [2, 5, 15]) including the Kardar–Parisi–Zhang (KPZ) equation, the generalized parabolic Anderson model and the stochastic quantization equations for quantum fields (see [12, 13, 38, 41] and references therein). In particular they led to a local well-posedness theory for the Navier–Stokes system (1.1) in three dimensions by Zhu and Zhu [61].

The question of global existence is even more challenging. Roughly speaking, in the field of singular SPDEs the only available global existence results rely either on a strong drift present in the system or a particular transform for certain nonlinearities or on properties of an invariant measure.

- Suitable a priori estimates have been established for the dynamical Φ^4 stochastic quantization model by Mourrat and Weber [52, 53] and Gubinelli and Hofmanová [33, 34] (see also [57, 58] for the vector valued case). All these results make an essential use of the strong damping term $-\phi^3$.
- In [38, 54, 63], a priori estimates and paracontrolled solutions to the KPZ equation and singular Hamilton–Jacobi–Bellman equation were obtained by using Cole–Hopf’s or Zvonkin’s transform and maximum principle.

- Moreover, using the probabilistic notion of energy solutions [36,37,39] or studying the associated infinitesimal generator and Kolmogorov equation [40], it is possible to construct global solutions to KPZ equation, but this depends on the invariant Gaussian measure (i.e. the law of Brownian motion or spatial white noise).
- We mention that in [19,55] global martingale solutions were constructed for geometric stochastic heat equations by using a Dirichlet form approach. This relies on an integration by parts formula for the known invariant measure.

No such results are available for the 3D Navier–Stokes system with space-time white noise.

- There is no strong drift helping to stabilize the evolution.
- Due to the appearance of the divergence free condition and the corresponding pressure term, it is impossible to apply maximum principle or Cole–Hopf’s transform.
- The existence of an invariant measure is an open problem.
- No global energy (or other) estimates are available due to irregularity of solutions (see below for more explanation on this point).

Furthermore, even if the regularity of the noise is increased, global existence is not known. More precisely, consider the following regularized problem which interpolates between the case of a trace-class noise and the space-time white noise

$$\begin{aligned} du + \operatorname{div}(u \otimes u) dt + \nabla p dt &= \Delta u dt + (-\Delta)^{-\gamma/2} dB, \\ \operatorname{div} u &= 0. \end{aligned} \quad (1.2)$$

Here B is the cylindrical Wiener process. The case of $\gamma > 3/2$ corresponds to a trace-class noise whereas $\gamma = 0$ is the space-time white noise.

For $\gamma > 3/2$, global existence of probabilistically weak Leray solutions is classical (see Flandoli, Gatarek [29]). Their uniqueness remains an outstanding open problem. The authors in [29] also constructed stationary solutions via Krylov–Bogoliubov’s argument and energy estimates. Ergodicity and strong Feller property was proved by Da Prato and Debussche [21] and also by Flandoli and Romito [32]. A Markov transition semigroup was then constructed by Debussche, Odasso [23] and by Flandoli and Romito [32]. However, further structure and properties of the invariant measure are still unclear. Recently, in [43] we established non-uniqueness in law in a class of analytically weak (not Leray) solutions using the method of convex integration. In [45], we additionally presented non-uniqueness of Markov solutions and global-in-time existence and non-uniqueness of probabilistically strong and analytically weak solutions.

If $\gamma \leq 3/2$ the usual energy estimates pertinent to the notion of Leray solution are not available. Indeed, as the noise is no longer trace-class, Itô’s formula cannot be applied. However, for $\gamma \in (1/2, 3/2]$ it is still possible to prove global existence of probabilistically weak solutions. Namely, decomposing the velocity u into the sum of its stochastic part z and its nonlinear part v (as shown also in Sect. 3.1 below), one can derive a global energy estimate for v . It can be then combined with compactness and Skorokhod representation theorem to deduce global existence.

Nevertheless, since it is necessary to change the probability space in the course of the proof, these solutions are only probabilistically weak.

Up to now, even global existence of probabilistically weak solutions was open in the case of $\gamma \leq 1/2$. Only local well-posedness in the spirit of Zhu, Zhu [61] seemed to be possible. The difficulty can be seen as follows. If e.g. $\gamma = 1/2$, we decompose $u = v + z$ with z solving the linear stochastic equation

$$dz + \nabla p^z dt = \Delta z dt + (-\Delta)^{-1/4} dB, \quad \operatorname{div} z = 0,$$

and v solving the nonlinear equation

$$\partial_t v + \operatorname{div}((v + z) \otimes (v + z)) + \nabla p^v = \Delta v, \quad \operatorname{div} v = 0.$$

As $z \in C_T C^{-\kappa}$, the best regularity for v is given by $C_T C^{1-\kappa}$. Hence we cannot expect the energy estimate for v since this would require the $L_T^2 H^1$ -norm of v . We may include further decomposition like in the case of the Φ^4 model [33, 34, 52, 53], but this would lead to new nonlinear terms which cannot be absorbed as in the Φ^4 model. Due to this, we also cannot obtain energy inequality and uniform estimates as in the case of a trace-class case. Consequently, we cannot derive global solutions by a usual compactness argument. Note that it is also possible to consider global solutions for small initial data (cf. [8]). However, this would destroy adaptedness of the solutions since the initial data would depend on the whole path of the driving noise.

1.2. Convex integration

In the present paper, we focus on the case of space-time white noise, i.e. $\gamma = 0$. Simplified versions of our proofs as outlined in Sect. 3 also provide the results in the more regular cases of $\gamma > 0$. Our idea is to apply the method of convex integration in order to construct global-in-time solutions. This is an iterative procedure which permits to construct solutions explicitly scale by scale. It makes an essential use of the form of the nonlinearity which propagates oscillations and reduces an error term, the so-called Reynolds stress, in order to approach a solution as one proceeds through the iteration. As typical for the convex integration constructions, the same method gives rise to infinitely many solutions.

Convex integration was introduced into fluid dynamics by De Lellis and Székelyhidi Jr. [25–27]. This method has already led to a number of groundbreaking results concerning the incompressible Euler equations, culminating in the proof of Onsager's conjecture by Isett [46] and by Buckmaster, De Lellis, Székelyhidi Jr. and Vicol [4]. Also the question of well/ill-posedness of the three dimensional Navier–Stokes equations has experienced an immense breakthrough: Buckmaster and Vicol [10] established non-uniqueness of weak solutions with finite kinetic energy, Buckmaster, Colombo and Vicol [3] were able to connect two arbitrary strong solutions via a weak solution. Bureczak, Modena and Székelyhidi Jr. [6] then obtained a number of new ill-posedness results for power-law fluids and in particular also non-uniqueness of weak solutions to the Navier–Stokes equations for every given divergence free initial condition in L^2 . Sharp non-uniqueness results for the

Navier–Stokes equations in dimension $d \geq 2$ were obtained by Cheskidov and Luo [17, 18]. We refer to the reviews [9, 11] for further details and references.

All these convex integration results very much rely on the L^2 -setting. Namely, the constructed solutions belong (at least) to L^2 and the iteration converges strongly in L^2 , hence one can pass to the limit in the quadratic nonlinearity. As the energy inequality is available in this setting, it is also understood as a natural selection criterion for physical solutions. Convex integration yields such solutions for Euler equations (see [26], and [44] for the stochastic setting), or when the diffusion is weak, e.g. for power-law fluids with small parameter p (see [6]) and for hypodissipative Navier–Stokes equations for small α (see [14]). However, constructing Leray solutions by convex integration to the Navier–Stokes system seems to be out of reach at the moment. By a different method, a first non-uniqueness result for Leray solutions was established recently by Albritton, Brué and Colombo [1] for the Navier–Stokes system with a force.

Compared to the classical uniform estimates and the compactness argument, convex integration provides a new way of constructing solutions. This turns out to be particularly useful in the stochastic setting as uniqueness of Leray solutions is unknown and there has been no result of existence of global probabilistically strong solutions before. In [45], we proved such a result for a trace-class noise by convex integration. In less regular settings, as, for instance, for $\gamma \leq 1/2$ discussed above, there are no Leray solutions to compete with in the first place. Furthermore, there are no alternative globally defined solutions whatsoever (neither probabilistically strong nor probabilistically weak). In this paper, we use convex integration to construct global probabilistically strong solutions in this setting when the energy inequality is out of reach. The question of how to select physical solutions in this case remains open.

1.3. Decomposition

We introduce a decomposition of the Navier–Stokes system (1.1), which makes also this singular setting amenable to convex integration. Recall that solutions are only expected to have regularity $-1/2 - \kappa$, $\kappa > 0$, hence the quadratic term $u \otimes u$ is far from being well-defined analytically. The common idea in the field of singular SPDEs is to prescribe a particular form of a solution u so that the nonlinearity can be made sense of. In the first step, we write

$$u = z + z^{\vee} + h.$$

The first term z solves the stochastic heat equation

$$dz + \nabla p^z dt = \Delta z dt + dB, \quad \operatorname{div} z = 0,$$

and permits to isolate the most irregular part of u , the rest being more regular. Note that by Schauder estimates, z belongs to $B_{\infty, \infty}^{-1/2 - \kappa}$ and hence the product $z \otimes z$ is not well defined in the classical sense. As z is Gaussian, we can understand $z \otimes z$

as a Wick product using renormalization. In particular, for a suitable mollification z_ε of z , there are diverging constants $C_\varepsilon \in \mathbb{R}^{3 \times 3}$, $C_\varepsilon^{ij} \rightarrow \infty$, so that

$$z^{\mathcal{V}} = \lim_{\varepsilon \rightarrow 0} z_\varepsilon \otimes z_\varepsilon - C_\varepsilon$$

is well defined and belongs to $B_{\infty, \infty}^{-1-\kappa}$. More details of the probabilistic constructions are included in Sect. 4.1. In order to isolate the corresponding (still irregular) part of the solution u , we then define

$$\partial_t z^{\mathcal{V}} + \operatorname{div}(z^{\mathcal{V}}) + \nabla p^2 = \Delta z^{\mathcal{V}}, \quad \operatorname{div} z^{\mathcal{V}} = 0,$$

which by Schauder estimates belongs to $B_{\infty, \infty}^{-\kappa}$.

It is then seen that the remainder h is already function valued but its regularity is necessarily limited by $1/2 - \kappa$. Even further decomposition cannot improve this regularity since products of z with the unknown always appear in the equation. A way how to overcome this issue stems from the work by Gubinelli, Imkeller and Perkowski [35] on the parabolic Anderson model and was applied to (1.1) by Zhu, Zhu in [61]: one postulates a paracontrolled ansatz which describes a further structure of the solution h ; it reads as

$$h = -\mathbb{P}[h \prec \mathcal{I}\nabla z] + \vartheta - (z^{\mathcal{Y}} + z^{\mathcal{Z}}).$$

Here, $z^{\mathcal{Y}}, z^{\mathcal{Z}}$ are additional stochastic objects constructed by renormalization, \mathcal{I} is the heat operator, \mathbb{P} the Helmholtz projection and finally \prec denotes a paraproduct as introduced by Bony [7]. This permits to cancel the two most irregular terms in the equation for h so that the remainder ϑ becomes more regular, namely, $1 - \kappa$. Furthermore, by a commutator lemma it permits to make sense of the analytically ill-defined product $h \otimes z$. We refer to Sect. 2.2 for basic definitions of paracontrolled calculus and to Sect. 4.2 for more details on the notion of our paracontrolled solution. In view of the regularity of solutions, we also see that energy inequality is impossible in this case.

The above decomposition is sufficient to prove local well-posedness as done in [61]. However, a much more refined analysis is indispensable to apply convex integration. Therefore, we split further $h = v^1 + v^2$ where v^1 represents the irregular part and v^2 the regular one. In addition, the equation for v^1 is linear whereas the one for v^2 contains the quadratic nonlinearity. Similarly to the above discussion of the more regular cases (1.2) with $\gamma \leq 1/2$, even with this decomposition into $v^1 + v^2$, it is not possible to derive global estimates via the energy method. Our idea is instead to apply convex integration on the level of v^2 . However, the equation for v^2 is coupled with the equation for v^1 . Therefore, we put forward a joint iterative procedure approximating both equations at once. The Reynolds stress \hat{R}_q is only included in the equation for v_q^2 , where $q \in \mathbb{N}_0$ is the iteration parameter. Consequently, the construction of the new iteration v_{q+1}^2 relies only on the previous stress \hat{R}_q . Here, we employ the intermittent jets by Buckmaster, Colombo and Vicol [3] (see also [9] and our previous works [43, 45]). As the next step, we solve the equation for v_{q+1}^1 exactly by a fixed point argument. See Fig. 1 for a sketch of our procedure.

In order to make this strategy possible, it is necessary to find the decomposition of the equation for h into the system for v^1 and v^2 and to define the corresponding equations for the iterations v_q^1 and v_q^2 . This, together with the construction of each approximate velocity v_{q+1}^2 through the intermittent jets, has to be done in a way to decrease the corresponding Reynolds stress \mathring{R}_{q+1} as $q \rightarrow \infty$. Especially the control of \mathring{R}_{q+1} requires a careful analysis of each of the terms appearing in the equation for h . We have to balance various competing factors such as regularity, integrability, blow-up as $t \rightarrow 0$ and blow-up as $q \rightarrow \infty$ of various terms. The divergencies need to be compensated by small quantities. We rely on a decomposition of each product into the two paraproducts and the resonant term, because each of these parts behaves differently and requires a different treatment. Roughly speaking, irregular terms are included into v^1 while regular ones into v^2 , but the precise splitting is delicate. Further issues and required ideas are summarized as follows:

- In the first step, we aim at constructing convex integration solutions up to a stopping time. The stopping time can be chosen arbitrarily large with arbitrarily large probability and is used to have uniform in ω bounds for the stochastic objects. The reason for this is adaptedness: without a stopping time, the parameters would depend on the whole path and the constructed solutions could not be adapted to the given filtration $(\mathcal{F}_t)_{t \geq 0}$. Thus, they would not be probabilistically strong.
- In the second step, we overcome this limitation by extending the constructed solutions by other convex integration solutions. To this end, it is necessary to obtain convex integration solutions for any given divergence free initial condition in L^2 . Hence, we aim at starting v^1 as well as v_q^1 from the given initial condition, whereas v^2 and v_q^2 all start from zero. This simplifies the begin of the iterative procedure. To keep the same initial value during the iteration, the oscillations can only be added for positive times and we approach $t = 0$ as $q \rightarrow \infty$.
- A paracontrolled ansatz for v^1 and accordingly also for each v_q^1 needs to be included in order to make sense of the resonant part of the product $v^1 \otimes z$ and $v_q^1 \otimes z$. At each iteration step q , the equation for v_q^1 is coupled with the equation for the corresponding remainder v_q^\sharp (taking the role of ϑ above).
- Since the initial value of v^1 , v_q^1 and v_q^\sharp is only in L^2 , they have singularity at time zero when considered in more regular function spaces. But higher regularity is indispensable in order to control various terms, both in the equation for v_q^1 and v_q^\sharp but also in the formula for \mathring{R}_{q+1} . For this reason, we work with blow-up norms in time, but this brings an extra blow-up in the convex integration part and in particular in the estimate of \mathring{R}_{q+1} .
- We introduce additional (uniform in q) localizers $\Delta_{>R} + \Delta_{\leq R}$ in terms of Littlewood–Paley blocks. The part $\Delta_{>R}$ is always included into v^1 and helps to simplify the estimates of v_q^1 and v_q^\sharp as it provides an arbitrarily small constant and avoids the need for a Gronwall lemma.
- Some terms are regular and could be, in principle, included into v^2 , but they require regularity of v_q^1 which does not hold true uniformly as $t \rightarrow 0$. While the

fixed point equation for v_q^1 can be solved using suitable blow-up norms in time to overcome the singularity at $t = 0$, having such blow-ups in the equation for v_q^2 cannot be controlled in the convex integration. Hence, we further decompose these terms by using $\Delta_{>R} + \Delta_{\leq R}$ and include their irregular parts into v_q^1 .

- Moreover, in order to control the blow-up of certain terms, we include q -dependent localizers $\Delta_{\leq f(q)}$ for a suitable $f(q) \rightarrow \infty$ into the equation for v_q^1 . Thanks to this, we are able to add irregularity scale by scale in a controlled way. The opposite approach, namely including the irregular terms fully at the beginning of the iteration, does not seem to be possible.

The detailed decomposition is presented in Sect. 4.3. As a preparation, we explain the main ideas on the simpler settings of (1.2) with $\gamma > 0$ in Sect. 3.

1.4. Final Introductory Remarks

Previously, convex integration has always been used to deduce non-uniqueness of solutions in settings where energy inequality is available. It has also been used to obtain first existence results for weak solutions in situations where compactness does not guarantee the passage to the limit in the convective term, namely, for Euler equations (see [25–27]), and for power-law fluids with small parameter p (see [6]). As already mentioned above, these results even lead to infinitely many weak solutions satisfying the energy inequality.

Our result shows that convex integration can also be used to construct global solutions when the energy inequality is out of reach, in particular in the field of singular SPDEs. We hope that our technique could also be applied to other singular SPDEs, especially in cases where no strong damping is at hand. We also mention that the convex integration can be applied to other PDEs like transport and continuity equation [50, 51], 3D Hall–MHD system [22] and hyperviscous Navier–Stokes equations [48]. Our approach may also be applied to the corresponding singular versions of the latter two.

The nonlinearity in the Navier–Stokes system looks similarly to the one in the Langevin dynamic for the Yang–Mills measure, i.e. stochastic quantization of the Yang–Mills field. This was considered in [12, 56] where local-in-time solutions were constructed. However, existence of global solutions remains open. The idea is to use the dynamics and PDE techniques to study properties of the field. Formally, these equations have the law of the associated field as an invariant measure. In the case of the stochastic quantization of the Euclidean Φ^4 field theory, it was indeed possible to use the dynamics to construct and study properties of the corresponding measure (see [34] and [59]). As global existence for the Langevin dynamic for the Yang–Mills measure is out of reach by the classical PDE techniques, we hope that our technique can shed some light on this problem.

Finally, we point out that in the field of regularization by noise, it is believed that more noise, in the sense of more irregular noise, in the Navier–Stokes equations may imply uniqueness. Certain regularizing effect could indeed be proved on the level of the so-called strong Feller property established for (1.1) by Zhu, Zhu [62]. Nevertheless, our results show that even in this case we still have non-uniqueness.

Furthermore, also a weaker notion of uniqueness, namely, uniqueness in law is disproved.

Organization of the paper In Sect. 2 we introduce our notation and present preliminary results on Besov spaces, paraproducts and paracontrolled calculus. Section 3 is devoted to the more regular settings of (1.2) with $\gamma > 0$ and we discuss the main ideas of our decomposition. In particular, it is shown how decreasing the parameter γ , i.e. making the problem more irregular, necessarily requires further ideas and a more refined decomposition. In Sect. 4 we recall the construction of stochastic objects and introduce the notion of paracontrolled solution. We also present a formal decomposition of the system into the system for v^1 and v^2 as discussed above. The set-up of the iterative convex integration procedure and proofs of our main results are shown in Sect. 5. We give estimates of v_q^1 and v_q^\sharp in Sect. 6. Section 7 is devoted to the core of the convex integration construction, namely, the iteration Proposition 5.1. Finally, in Appendix A we recall the construction of intermittent jets and in “Appendix B” we prove auxiliary Schauder estimates.

2. Preliminaries

Throughout the paper, we use the notation $a \lesssim b$ if there exists a constant $c > 0$ such that $a \leq cb$, and we write $a \simeq b$ if $a \lesssim b$ and $b \lesssim a$.

2.1. Function Spaces

Given a Banach space E with a norm $\|\cdot\|_E$ and $T > 0$, we write $C_T E = C([0, T]; E)$ for the space of continuous functions from $[0, T]$ to E , equipped with the supremum norm $\|f\|_{C_T E} = \sup_{t \in [0, T]} \|f(t)\|_E$. For $p \in [1, \infty]$ we write $L_T^p E = L^p([0, T]; E)$ for the space of L^p -integrable functions from $[0, T]$ to E , equipped with the usual L^p -norm. We use $(\Delta_i)_{i \geq -1}$ to denote the Littlewood–Paley blocks corresponding to a dyadic partition of unity. Besov spaces on the torus with general indices $\alpha \in \mathbb{R}$, $p, q \in [1, \infty]$ are defined as the completion of $C^\infty(\mathbb{T}^d)$ with respect to the norm

$$\|u\|_{B_{p,q}^\alpha} := \left(\sum_{j \geq -1} 2^{j\alpha q} \|\Delta_j u\|_{L^p}^q \right)^{1/q}.$$

The Hölder–Besov space C^α is given by $C^\alpha = B_{\infty,\infty}^\alpha$ and we also set $H^\alpha = B_{2,2}^\alpha$, $\alpha \in \mathbb{R}$. To deal with the singularity at time zero we introduce the following blow-up norms: for $\alpha \in (0, 1)$, $p \in [1, \infty]$

$$\begin{aligned} \|f\|_{C_{T,\gamma}^\alpha L^p} &:= \sup_{0 \leq t \leq T} t^\gamma \|f(t)\|_{L^p} + \sup_{0 \leq s < t \leq T} s^\gamma \frac{\|f(t) - f(s)\|_{L^p}}{|t - s|^\alpha}, \\ \|f\|_{C_{T,\gamma} B_{p,\infty}^\alpha} &:= \sup_{0 \leq t \leq T} t^\gamma \|f(t)\|_{B_{p,\infty}^\alpha}. \end{aligned}$$

For $T > 0$ and a domain $D \subset \mathbb{R}^+$ we denote by $C_{T,x}^N$ and $C_{D,x}^N$, respectively, the space of C^N -functions on $[0, T] \times \mathbb{T}^3$ and on $D \times \mathbb{T}^3$, respectively, $N \in \mathbb{N}_0 := \mathbb{N} \cup \{0\}$. The spaces are equipped with the norms

$$\|f\|_{C_{T,x}^N} = \sum_{\substack{0 \leq n+|\alpha| \leq N \\ n \in \mathbb{N}_0, \alpha \in \mathbb{N}_0^3}} \|\partial_t^n D^\alpha f\|_{L_T^\infty L^\infty},$$

$$\|f\|_{C_{D,x}^N} = \sum_{\substack{0 \leq n+|\alpha| \leq N \\ n \in \mathbb{N}_0, \alpha \in \mathbb{N}_0^3}} \sup_{t \in D} \|\partial_t^n D^\alpha f\|_{L^\infty}.$$

Set $\Lambda = (1 - \Delta)^{1/2}$. For $s \geq 0, p \in [1, +\infty]$ we use $W^{s,p}$ to denote the subspace of L^p , consisting of all f which can be written in the form $f = \Lambda^{-s}g, g \in L^p$ and the $W^{s,p}$ norm of f is defined to be the L^p norm of g , i.e. $\|f\|_{W^{s,p}} := \|\Lambda^s f\|_{L^p}$. For $s < 0, p \in (1, \infty), W^{s,p}$ is the dual space of $W^{-s,q}$ with $\frac{1}{p} + \frac{1}{q} = 1$.

The following embedding results will be frequently used (we refer to e.g. [35, Lemma A.2] for the first one and to [60, Theorem 4.6.1] for the second one).

Lemma 2.1. (1) *Let $1 \leq p_1 \leq p_2 \leq \infty$ and $1 \leq q_1 \leq q_2 \leq \infty$, and let $\alpha \in \mathbb{R}$. Then $B_{p_1,q_1}^\alpha \subset B_{p_2,q_2}^{\alpha-d(1/p_1-1/p_2)}$.*
 (2) *Let $s \in \mathbb{R}, 1 < p < \infty, \epsilon > 0$. Then $W^{s,2} = B_{2,2}^s = H^s$, and $B_{p,1}^s \subset W^{s,p} \subset B_{p,\infty}^s \subset B_{p,1}^{s-\epsilon}$.*

2.2. Paraproducts, Commutators and Localizers

Paraproducts were introduced by Bony in [7] and they permit to decompose a product of two distributions into three parts which behave differently in terms of regularity. More precisely, using the Littlewood-Paley blocks, the product fg of two Schwartz distributions $f, g \in \mathcal{S}'(\mathbb{T}^d)$ can be formally decomposed as

$$fg = f \prec g + f \circ g + f \succ g,$$

with

$$f \prec g = g \succ f = \sum_{j \geq -1} \sum_{i < j-1} \Delta_i f \Delta_j g, \quad f \circ g = \sum_{|i-j| \leq 1} \Delta_i f \Delta_j g.$$

Here, the paraproducts \prec and \succ are always well-defined and critical is the resonant product denoted by \circ . In general, it is only well-defined provided the sum of the regularities of f and g in terms of Besov spaces is strictly positive. Moreover, we have the following paraproduct estimates from [7] (see also [35, Lemma 2.1], [53, Proposition A.7]):

Lemma 2.2. *Let $\beta \in \mathbb{R}, p, p_1, p_2, q \in [1, \infty]$ such that $\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2}$. Then it holds that*

$$\|f \prec g\|_{B_{p,q}^\beta} \lesssim \|f\|_{L^{p_1}} \|g\|_{B_{p_2,q}^\beta},$$

and if $\alpha < 0$ then

$$\|f \prec g\|_{B_{p,q}^{\alpha+\beta}} \lesssim \|f\|_{B_{p_1,q}^\alpha} \|g\|_{B_{p_2,q}^\beta}.$$

If $\alpha + \beta > 0$ then it holds that

$$\|f \circ g\|_{B_{p,q}^{\alpha+\beta}} \lesssim \|f\|_{B_{p_1,q}^\alpha} \|g\|_{B_{p_2,q}^\beta}.$$

We denote $\succ = \circ + \succ$, $\preccurlyeq = \circ + \prec$. The key tool of the paracontrolled calculus introduced in [35] is the following commutator lemma from [35, Lemma 2.4] (see also [53, Proposition A.9]).

Lemma 2.3. *Assume that $\alpha \in (0, 1)$ and $\beta, \gamma \in \mathbb{R}$ are such that $\alpha + \beta + \gamma > 0$ and $\beta + \gamma < 0$ and $p, p_1, p_2 \in [1, \infty]$ satisfy $\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2}$. Then there exist a bounded trilinear operator*

$$\text{com}(f, g, h) : B_{p_1,\infty}^\alpha \times C^\beta \times B_{p_2,\infty}^\gamma \rightarrow B_{p,\infty}^{\alpha+\beta+\gamma}$$

satisfying

$$\|\text{com}(f, g, h)\|_{B_{p,\infty}^{\alpha+\beta+\gamma}} \lesssim \|f\|_{B_{p_1,\infty}^\alpha} \|g\|_{C^\beta} \|h\|_{B_{p_2,\infty}^\gamma}$$

such that for smooth functions f, g, h it holds that

$$\text{com}(f, g, h) = (f \prec g) \circ h - f(g \circ h).$$

We also recall the following two lemmas for the Helmholtz projection \mathbb{P} from [61, Lemma 3.5, Lemma 3.6].

Lemma 2.4. *Assume that $\alpha \in (0, 1), \beta \in \mathbb{R}$ and $p \in [1, \infty]$. Then, for every $k, l = 1, 2, 3$,*

$$\|[\mathbb{P}^{kl}, f \prec]g\|_{B_{p,\infty}^{\alpha+\beta}} \lesssim \|f\|_{B_{p,\infty}^\alpha} \|g\|_{C^\beta}.$$

Lemma 2.5. *Assume that $\alpha \in \mathbb{R}$ and $p \in [1, \infty]$. Then for every $k, l = 1, 2, 3$*

$$\|\mathbb{P}^{kl} f\|_{B_{p,\infty}^\alpha} \lesssim \|f\|_{B_{p,\infty}^\alpha}.$$

Analogously to the the real-valued case, we may define paraproducts for vector-valued distributions. More precisely, for two vector-valued distributions $f, g \in \mathcal{S}'(\mathbb{T}^d; \mathbb{R}^m)$, we use the following tensor product notation

$$\begin{aligned} f \otimes g &= (f_i g_j)_{i,j=1}^m = f \otimes g + f \circledast g + f \otimes g \\ &= (f_i \prec g_j)_{i,j=1}^m + (f_i \circ g_j)_{i,j=1}^m + (f_i \succ g_j)_{i,j=1}^m \end{aligned}$$

and note that Lemma 2.2 carries over mutatis mutandis. We also denote

$$\otimes = \circledast + \otimes, \quad \circledast = \circledast + \otimes.$$

When there is no danger of confusion, we apply the simple paraproducts also within matrix–vector multiplication, i.e. for $f \in \mathcal{S}'(\mathbb{T}^d; \mathbb{R}^{m \times m})$ and $g \in \mathcal{S}'(\mathbb{T}^d; \mathbb{R}^m)$ we define using the Einstein summation convention

$$\begin{aligned} (f \succ g)_{i=1}^m &= (g \prec f)_{i=1}^m = \left(f^{ij} \succ g^j \right)_{i=1}^m, \\ (f \circ g)_{i=1}^m &= (g \circ f)_{i=1}^m = \left(f^{ij} \circ g^j \right)_{i=1}^m. \end{aligned}$$

Similarly to Lemma 2.3, we may also define a matrix-valued commutator as continuous extensions of

$$\begin{aligned} (f, g, h) &\mapsto (\text{com}(f, g, h))_{i,j=1}^m = (f \prec g) \otimes h - f \cdot (g \otimes h) \\ &= (f^k \prec g^{ik}) \circ h^j - f^k (g^{ik} \circ h^j), \\ (f, g, h) &\mapsto (\text{com}^*(f, g, h))_{i,j=1}^m = h \otimes (f \prec g) - (h \otimes g) \cdot f \\ &= h^i \circ (f^k \prec g^{jk}) - (h^i \circ g^{jk}) f^k, \end{aligned}$$

which is well-defined for smooth functions $f, h : \mathbb{T}^d \rightarrow \mathbb{R}^m, g : \mathbb{T}^d \rightarrow \mathbb{R}^{m \times m}$ and takes values in $\mathcal{S}'(\mathbb{T}^d; \mathbb{R}^{m \times m})$. A counterpart of the bound in Lemma 2.3 holds true in this setting as well.

Finally, we introduce localizers in terms of Littlewood–Paley expansions. Let $J \in \mathbb{N}_0$. For $f \in \mathcal{S}'(\mathbb{T}^d)$ we define

$$\Delta_{>J} f = \sum_{j>J} \Delta_j f, \quad \Delta_{\leq J} f = \sum_{j \leq J} \Delta_j f.$$

Then it holds, in particular for $\alpha \leq \beta \leq \gamma$, that

$$\|\Delta_{>J} f\|_{C^\alpha} \lesssim 2^{-J(\beta-\alpha)} \|f\|_{C^\beta}, \quad \|\Delta_{\leq J} f\|_{C^\gamma} \lesssim 2^{J(\gamma-\beta)} \|f\|_{C^\beta}. \tag{2.1}$$

3. More Regular Stochastic Perturbations

In our previous works [43,45] we considered the case of a trace class noise. We proved non-uniqueness in law as well as non-uniqueness of Markov solutions and existence and non-uniqueness of global-in-time probabilistically strong solutions. These results apply to the Navier–Stokes system

$$du + \text{div}(u \otimes u)dt + \nabla p dt = \Delta u dt + (-\Delta)^{-\gamma/2} dB, \quad \text{div } u = 0, \tag{3.1}$$

where $B = (B^1, B^2, B^3)$ is a vector-valued L^2 -cylindrical Wiener process on some stochastic basis $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbf{P})$ and $\gamma > 3/2$. Here the filtration $(\mathcal{F}_t)_{t \geq 0}$ is the normal filtration generated by the Wiener process B . As a main result of the present paper, we treat the case of space-time white noise, i.e. $\gamma = 0$. But already the more regular case of $\gamma \in (0, 3/2]$ presents interesting new challenges. In this section we want to outline some of the main ideas on examples of these more regular noises.

If the noise is not trace class, Itô’s formula cannot be applied in order to obtain an energy inequality. The case of $\gamma = 3/2$ is therefore the threshold where (and below

which) stochastic counterparts of Leray solutions on the level of u no longer make sense. Furthermore, the expected regularity of solutions depends on the regularity of the noise. This can be seen by looking at the linear counterpart of (3.1)

$$dz + \nabla p^z dt = \Delta z dt + (-\Delta)^{-\gamma/2} dB, \quad \operatorname{div} z = 0, \tag{3.2}$$

and realizing that $z \in C_T C^{\gamma-1/2-\kappa} \cap C_T^{1/2-\delta} C^{\gamma-3/2+2\delta-\kappa}$ \mathbf{P} -a.s. for $\kappa, \delta > 0$ small. This can be obtained by Schauder estimates from the fact that the space-time white noise dB/dt is a random distribution of space-time regularity $-5/2 - \kappa$ for $\kappa > 0$ with parabolic scaling. Thus, if $\gamma \leq 1/2$ the solution is not even function-valued and the quadratic nonlinearity in (3.1) is not well-defined analytically. Nevertheless, one can use probability theory and renormalization to define this nonlinearity using Wick products.

We distinguish the following cases with an increasing level of difficulty:

- (1) $\gamma \in (1/2, 3/2]$: solutions are function-valued hence the convective term is well-defined but the energy inequality for u cannot be obtained.
- (2) $\gamma \in (1/6, 1/2]$: solutions become distribution-valued, renormalization is needed to define the product.
- (3) $\gamma \in (0, 1/6]$: further decomposition is required in order to make sense of all the required products.
- (4) $\gamma = 0$: the case of space-time white noise, a so-called paracontrolled ansatz is required and we present a detailed proof in subsequent sections.

We stress that while global existence of martingale (i.e. probabilistically weak) solutions in the case $\gamma \in (1/2, 3/2]$ can be obtained by compactness and Skorokhod representation theorem, existence of probabilistically strong solutions was unknown. For $\gamma \leq 1/2$ even global existence of martingale solutions was an open problem.

In the remainder of this section, we focus on the first three regimes and explain how to make them amenable to convex integration and hence to global-in-time existence and non-uniqueness results.

3.1. The Case of $\gamma \in (1/2, 3/2]$

Even though the energy inequality cannot be computed here, the approach of [43,45] can be applied with minimal modifications. In particular, one uses the decomposition $u = z + v$ where z solves (3.2) and

$$\mathcal{L}v + \operatorname{div}((v + z) \otimes (v + z)) + \nabla p^v = 0, \quad \operatorname{div} v = 0. \tag{3.3}$$

Here and in the sequel, we use the notation $\mathcal{L} = \partial_t - \Delta$. The convex integration scheme is then given via the iteration system

$$\mathcal{L}v_q + \operatorname{div}((v_q + z_q) \otimes (v_q + z_q)) + \nabla p^q = \operatorname{div} \mathring{R}_q, \quad \operatorname{div} v_q = 0,$$

where $z_q = \Delta_{\leq f(q)} z$ with $\Delta_{\leq f(q)}$ being a cut-off of the Littlewood–Paley expansion. With a suitable definition of the stopping times, this permits to add noise scale by scale as one proceeds through the iteration. Setting the convex integration up

with an initial iteration v_0 as in [43], one can prove that the constructed solution v is distinct from the Leray solution to (3.3) starting from the same initial value. This implies existence of an initial value so that non-uniqueness holds for (3.1) as well. Using the probabilistic extension of solutions from [43], non-uniqueness in law on any time interval $[0, T]$, $T > 0$, follows. Following the ideas of [45] it is also possible to construct solutions with a prescribed L^2 initial condition and to obtain existence and non-uniqueness of global-in-time probabilistically strong solutions.

3.2. The Case of $\gamma \in (1/6, 1/2]$

This case is more delicate and further decomposition is required. More precisely, in addition to $u = z + v$ as above, we split v into its irregular and regular part, i.e. $v = v^1 + v^2$. The equation for v^1 contains all the irregular terms of the product $(v + z) \otimes (v + z)$, whereas the regular ones are put in v^2 . Additionally, the equation for v^1 shall be linear so that it can be solved by a fixed point argument. As a rule of thumb, we color the irregular terms magenta and the regular ones blue. The decomposition can be done as follows. The product $z \otimes z$ needs to be constructed by renormalization as a Wick product denoted by z^{\vee} and it is of spatial regularity $C^{2(\gamma-1/2)-\kappa}$. For the moment, we ignore this fact and proceed formally. We come back to the rigorous definition of the stochastic objects in Sect. 4.1.

So we have the first magenta term $z \otimes z$. Then we write with the help of para-products and Littlewood–Paley projectors

$$(v^1 + v^2) \otimes z = (v^1 + v^2) \otimes \Delta_{>R} z + (v^1 + v^2) \otimes \Delta_{>R} z + (v^1 + v^2) \otimes \Delta_{\leq R} z,$$

and treat the symmetric term $z \otimes (v^1 + v^2)$ the same way. Here, $\otimes = \circledast + \circledcirc$ and we included a suitable cut-off R to be chosen appropriately. This eventually simplifies the fixed point argument used to establish, for a given convex integration iteration v_q^2 , the existence and uniqueness of v_q^1 . Finally, we let $(v^1 + v^2) \otimes (v^1 + v^2)$. This leads to

$$\begin{aligned} \mathcal{L}v^1 + \operatorname{div} (z \otimes z + (v^1 + v^2) \otimes \Delta_{>R} z + \Delta_{>R} z \otimes (v^1 + v^2)) + \nabla p^1 &= 0, \quad \operatorname{div} v^1 = 0, \\ \mathcal{L}v^2 + \operatorname{div} ((v^1 + v^2) \otimes \Delta_{>R} z + \Delta_{>R} z \otimes (v^1 + v^2) + (v^1 + v^2) \otimes \Delta_{\leq R} z) \\ + \operatorname{div} (\Delta_{\leq R} z \otimes (v^1 + v^2) + (v^1 + v^2) \otimes \Delta_{\leq R} z) + \nabla p^2 &= 0, \quad \operatorname{div} v^2 = 0. \end{aligned}$$

We set up a convex integration scheme as an approximation of the above system of equations for v^1 and v^2 . In particular, we include further Littlewood–Paley projectors and let

$$\begin{aligned} \mathcal{L}v_q^1 + \operatorname{div} (z \otimes z + (v_q^1 + v_q^2) \otimes \Delta_{\leq f(q)} \Delta_{>R} z + \Delta_{\leq f(q)} \Delta_{>R} z \otimes (v_q^1 + v_q^2)) + \nabla p_q^1 &= 0, \quad \operatorname{div} v_q^1 = 0, \\ \mathcal{L}v_q^2 + \operatorname{div} ((v_q^1 + v_q^2) \otimes \Delta_{>R} z + \Delta_{>R} z \otimes (v_q^1 + v_q^2) + (v_q^1 + v_q^2) \otimes \Delta_{\leq R} z) \\ + \operatorname{div} (\Delta_{\leq R} z \otimes (v_q^1 + v_q^2) + (v_q^1 + v_q^2) \otimes \Delta_{\leq R} z) + \nabla p_q^2 &= \operatorname{div} \hat{R}_q, \quad \operatorname{div} v_q^2 = 0. \end{aligned}$$

Note that the Reynolds stress \hat{R}_q is only included in the equation for v_q^2 . Indeed, v_q^2 contains the quadratic nonlinearity which is used in the convex integration to reduce the stress.

At each iteration step $q + 1$, we first use the previous stress \mathring{R}_q in order to define the principle part of the corrector $w_{q+1}^{(p)}$, the incompressibility corrector $w_{q+1}^{(c)}$ and the time corrector $w_{q+1}^{(t)}$ in terms of the intermittent jets, see ‘‘Appendix A’’. This gives the next iteration v_{q+1}^2 and consequently we obtain v_{q+1}^1 by a fixed point argument, cf. Figure 1. The localizers $\Delta_{\leq f(q)}$ in the equation of v_q^1 are used to control the blow up of a certain norm of v_q^1 as $q \rightarrow \infty$.

3.3. The Case of $\gamma \in (0, 1/6]$

In this regime, also the resonant product $v^1 \otimes z$ becomes problematic. This can be overcome by introducing an additional stochastic object which permits to cancel the worst term, i.e. $z \otimes z$, from the equation for v^1 . To be precise, let

$$\mathcal{L}z_1 + \operatorname{div}(z \otimes z) + \nabla p^{z_1} = 0, \quad \operatorname{div} z_1 = 0, \tag{3.4}$$

and define $u = z + v = z + z_1 + v^1 + v^2$. Recall that in the rigorous analysis $z \otimes z$ needs to be replaced by the Wick product z^{\vee} of regularity $C^{2(\gamma-1/2)-\kappa}$. Consequently, z_1 then becomes our second stochastic object, later denoted as z^{\vee} with regularity $C^{2\gamma-\kappa}$. In addition, also the products $z_1 \otimes z$ and $z \otimes z_1$ need to be defined via renormalization as $z^{\vee} \otimes z$ and $z \otimes z^{\vee}$, respectively. We again ignore this fact for a moment and continue with the formal decomposition. The reader is referred to Sect. 4.1 below for more details on the stochastic construction.

Proceeding as above, this leads to

$$\begin{aligned} \mathcal{L}v^1 + \operatorname{div}(z_1 \otimes z + z \otimes z_1 + z_1 \otimes z_1 + (v^1 + v^2) \otimes \Delta_{>Rz} + \Delta_{>Rz} \otimes (v^1 + v^2)) + \nabla p^1 &= 0, \\ \operatorname{div} v^1 &= 0, \\ \mathcal{L}v^2 + \operatorname{div}((v^1 + v^2) \otimes \Delta_{>Rz} + \Delta_{>Rz} \otimes (v^1 + v^2) + (v^1 + v^2) \otimes (\Delta_{\leq Rz} + z_1)) \\ + \operatorname{div}((z_1 + \Delta_{\leq Rz}) \otimes (v^1 + v^2) + (v^1 + v^2) \otimes (v^1 + v^2)) + \nabla p^2 &= 0, \quad \operatorname{div} v^2 = 0, \end{aligned}$$

together with the convex integration scheme

$$\begin{aligned} \mathcal{L}v_q^1 + \operatorname{div}(z_1 \otimes z + z \otimes z_1 + z_1 \otimes z_1 + (v_q^1 + v_q^2) \otimes \Delta_{\leq f(q)} + \Delta_{>Rz} + \Delta_{\leq f(q)} \otimes \Delta_{>Rz} \otimes (v_q^1 + v_q^2)) \\ + \nabla p_q^1 &= 0, \quad \operatorname{div} v_q^1 = 0, \\ \mathcal{L}v_q^2 + \operatorname{div}((v_q^1 + v_q^2) \otimes \Delta_{>Rz} + \Delta_{>Rz} \otimes (v_q^1 + v_q^2) + (v_q^1 + v_q^2) \otimes (\Delta_{\leq Rz} + z_1)) \\ + \operatorname{div}((\Delta_{\leq Rz} + z_1) \otimes (v_q^1 + v_q^2) + (v_q^1 + v_q^2) \otimes (v_q^1 + v_q^2)) + \nabla p_q^2 &= \operatorname{div} \mathring{R}_q, \quad \operatorname{div} v_q^2 = 0. \end{aligned}$$

With this definition, one can obtain existence of an initial condition for which there are non-unique solutions before a stopping time. Since there are no global Leray solutions, for global existence it is necessary to extend these solutions by other convex integration solutions. To this end, an improved convex integration construction is needed, which gives the result for any prescribed initial condition in L^2 . In particular, the term

$$v^1 \otimes \Delta_{>Rz}$$

and its symmetric counterpart appearing in the equation for v^2 require regularity of v^1 . Accordingly, this introduces blow-up norms in time to overcome the singularity at $t = 0$ into the convex integration, which is not convenient. We therefore refine the decomposition above by writing

$$(v^1 + v^2) \otimes \Delta_{>Rz} = v^1 \otimes \Delta_{>Rz} + v^2 \otimes \Delta_{>Rz}.$$

We include these terms into the equations for v^1, v^2 accordingly respecting the colors. Then in the equations for v_q^1 and v_q^2 we rewrite

$$(v_q^1 + v_q^2) \otimes \Delta_{>Rz} = v_q^1 \otimes \Delta_{>Rz} + v_q^2 \otimes \Delta_{>Rz}.$$

The symmetric terms are treated the same way.

4. Paracontrolled Solutions

It turns out that even further expansion would not help in order to treat the case of space-time white noise, i.e. $\gamma = 0$. Indeed, there would always be ill-defined products. As understood in the field of singular SPDEs, a paracontrolled ansatz needs to be included. It postulates a particular structure of solutions and permits to make sense of the analytically ill-defined products using probabilistic tools. In the sequel, we first introduce the stochastic objects needed for the rigorous formulation of the Navier–Stokes system (1.1). Then we formulate the notion of paracontrolled solution incorporating the paracontrolled ansatz. Finally, we give a formal decomposition combined with paracontrolled ansatz in the spirit of Sect. 3.

4.1. Stochastic Objects

Let us recall that due to Theorem 1.1 [61], the equation (3.1) with $\gamma = 0$ is locally well-posed for initial conditions in C^η for $\eta \in (-1, -1/2)$. The solution u belongs to $C([0, \sigma]; C^\eta)$ where σ is a strictly positive stopping time so that

$$\|u_\varepsilon - u\|_{C_\sigma C^\eta} \rightarrow 0$$

in probability. Here, u_ε denotes the solution to the regularized Navier–Stokes system

$$\partial_t u_\varepsilon + \operatorname{div}(u_\varepsilon \otimes u_\varepsilon) + \nabla p_\varepsilon = \Delta u_\varepsilon + \zeta_\varepsilon, \quad \operatorname{div} u_\varepsilon = 0,$$

where ζ_ε is a mollification of the space-time white noise $\zeta = dB/dt$. In particular, the stochastic objects needed in our proof here were constructed in [61].

To summarize the main ideas, let z_ε be the stationary solution to

$$\partial_t z_\varepsilon + \nabla p^{z_\varepsilon} = \Delta z_\varepsilon + \zeta_\varepsilon, \quad \operatorname{div} z_\varepsilon = 0.$$

Then $z_\varepsilon \rightarrow z$ in $L^p(\Omega; C_T C^{-1/2-\kappa})$ for every $p \in [1, \infty)$. Moreover, the renormalized product $z \mathring{\vee}$ can be defined as a Wick product in the sense that there exist diverging constants $C_\varepsilon \in \mathbb{R}^{3 \times 3}, C_\varepsilon^{ij} \rightarrow \infty$, so that

$$z_\varepsilon \mathring{\vee} := z_\varepsilon \otimes z_\varepsilon - C_\varepsilon$$

has a well-defined limit in $L^p(\Omega; C_T C^{-1-\kappa})$ for every $p \in [1, \infty)$. In fact, $C_\varepsilon = \mathbf{E}[z_\varepsilon \otimes z_\varepsilon]$. We also introduce the following stochastic objects. Let

$$\partial_t z_\varepsilon^\mathbb{Y} - \Delta z_\varepsilon^\mathbb{Y} = -\mathbb{P}\text{div}(z_\varepsilon^\mathbb{V}), \quad \text{div } z_\varepsilon^\mathbb{Y} = 0, \quad z_\varepsilon^\mathbb{Y}(0) = 0,$$

with \mathbb{P} being the Leray projection operator, and define

$$\begin{aligned} z_\varepsilon^\mathbb{Z} &:= z_\varepsilon^\mathbb{Y} \otimes z_\varepsilon, & z_\varepsilon^\mathbb{N} &:= z_\varepsilon \otimes z_\varepsilon^\mathbb{Y}, & z_\varepsilon^\mathbb{L} &:= \mathbb{P}\mathcal{I}(\nabla z_\varepsilon) \odot z_\varepsilon = \left(\mathbb{P}\mathcal{I}(\nabla z_\varepsilon)^{ik} \circ z_\varepsilon^j \right)_{i,j,k=1}^3, \\ z_\varepsilon^\mathbb{M} &:= \mathbb{P}\mathcal{I}\text{div}(z_\varepsilon^\mathbb{Z}), & z_\varepsilon^\mathbb{O} &:= \mathbb{P}\mathcal{I}\text{div}(z_\varepsilon^\mathbb{N}), \\ z_\varepsilon^\mathbb{P} &:= z_\varepsilon^\mathbb{Y} \otimes z_\varepsilon^\mathbb{Y} - C_{1,\varepsilon}(t), & z_\varepsilon^\mathbb{Q} &:= z_\varepsilon^\mathbb{Z} \odot z_\varepsilon + z_\varepsilon^\mathbb{N} \odot z_\varepsilon - C_{2,\varepsilon}(t), \end{aligned}$$

where $\mathcal{I}f(t) = \int_0^t e^{(t-r)\Delta} f(r)dr$ and

$$\begin{aligned} C_{1,\varepsilon}(t) &= \mathbf{E}[(z_\varepsilon^\mathbb{Y} \otimes z_\varepsilon^\mathbb{Y})(t, 0)] \rightarrow \infty, \\ C_{2,\varepsilon}(t) &= \mathbf{E}[(z_\varepsilon^\mathbb{Z} \odot z_\varepsilon)(t, 0)] + \mathbf{E}[(z_\varepsilon^\mathbb{N} \odot z_\varepsilon)(t, 0)] \rightarrow \infty \end{aligned}$$

as $\varepsilon \rightarrow 0$.

For connecting the solutions in Sect. 5 we also introduce two stochastic objects $z_\varepsilon^\mathbb{R}(r; s)$ and $z_\varepsilon^\mathbb{S}(r; s)$. They are defined the same way as $z_\varepsilon^\mathbb{Z}$ and $z_\varepsilon^\mathbb{L}$ but replacing the last integration operator \mathcal{I} by $\mathcal{I}_{s,r} = \int_s^r e^{(r-l)\Delta} dl$ and the renormalized diverging constants $C_{i,\varepsilon}(r; s)$ are defined as the expectation of the corresponding terms for $i = 1, 2$.

We recall the following result from [61]:

Proposition 4.1. *For every $\kappa > 0$ and some $0 < \delta < 1/4$, there exist random distributions*

$$\mathbb{Z} := (z, z^\mathbb{V}, z^\mathbb{Y}, z^\mathbb{Z}, z^\mathbb{N}, z^\mathbb{M}, z^\mathbb{P}, z^\mathbb{Q}, z^\mathbb{R}, z^\mathbb{S}) \tag{4.1}$$

such that if τ_ε is a component in

$$\mathbb{Z}_\varepsilon := (z_\varepsilon, z_\varepsilon^\mathbb{V}, z_\varepsilon^\mathbb{Y}, z_\varepsilon^\mathbb{Z}, z_\varepsilon^\mathbb{N}, z_\varepsilon^\mathbb{M}, z_\varepsilon^\mathbb{P}, z_\varepsilon^\mathbb{Q}, z_\varepsilon^\mathbb{R}, z_\varepsilon^\mathbb{S})$$

and τ is the corresponding component in \mathbb{Z} then $\tau_\varepsilon \rightarrow \tau$ in $C_T C^{\alpha_\tau} \cap C_T^{\delta/2} C^{\alpha_\tau - \delta}$ a.s. as $\varepsilon \rightarrow 0$, where the regularities α_τ are summarized in Table 1. Furthermore, for every $p \in [1, \infty)$

$$\sup_{\varepsilon \geq 0} \mathbf{E} \|\tau_\varepsilon\|_{C_T C^{\alpha_\tau}}^p + \sup_{\varepsilon \geq 0} \mathbf{E} \|\tau_\varepsilon\|_{C_T^{\delta/2} C^{\alpha_\tau - \delta}}^p \lesssim 1.$$

For $\tau_\varepsilon(r; s) = z_\varepsilon^\mathbb{R}(r; s)$ and $\tau_\varepsilon(r; s) = z_\varepsilon^\mathbb{S}(r; s)$ there exist random distributions $\tau(r; s) = z^\mathbb{R}(r; s)$ and $\tau(r; s) = z^\mathbb{S}(r; s)$ such that

$$\sup_{\varepsilon \geq 0} \mathbf{E} \sup_{0 \leq s \leq r \leq T} \|\tau_\varepsilon\|_{C^{\alpha_\tau}}^p \lesssim 1,$$

Table 1. Regularity of stochastic objects

τ	z	$z^{\mathcal{V}}$	$z^{\mathcal{V}}$	$z^{\mathcal{V}}$	$z^{\mathcal{V}}$	$z^{\mathcal{V}}$	$z^{\mathcal{V}}$	$z^{\mathcal{V}}$
α_τ	$-\frac{1}{2} - \kappa$	$-\kappa$	$-1 - \kappa$	$-\frac{1}{2} - \kappa$	$-\frac{1}{2} - \kappa$	$-\kappa$	$-\kappa$	$-\kappa$

and

$$\sup_{\varepsilon \geq 0} \mathbf{E} \sup_{0 \leq s \leq r \leq T} \|\tau_\varepsilon - \tau\|_{C^{\alpha_\tau}}^p \rightarrow 0,$$

as $\varepsilon \rightarrow 0$.

The renormalization of $\tau(r; s)$ can be done by a similar argument as [61].

Remark 4.2. We emphasize that the renormalization constants $C_\varepsilon, C_{i,\varepsilon}, i = 1, 2$, only depend on t . Hence, due to the divergence in the nonlinear term, they do not appear in the approximate Navier–Stokes system driven by the mollified noise ζ_ε . If we modified the $C_\varepsilon, C_{i,\varepsilon}, i = 1, 2$, by adding a finite constant, some of the limit random distributions τ would change. For the rest of the paper, we fix the stochastic objects τ and prove existence of infinitely many solutions h to (4.2) with these fixed stochastic data.

Finally, we note that similarly to our decomposition, the local solution u obtained in Theorem 1.1 in [61] decomposes as $u = z + z_1 + h$ with a suitable paracontrolled ansatz for h . In particular, by Schauder estimates, the part h possesses a positive spatial regularity at positive times, i.e. it belongs in particular to L^2 . For this reason, it is sufficient to restrict our attention to initial conditions for the convex integration scheme in L^2 . Indeed, for an initial condition in $C^\eta, \eta \in (-1, -1/2)$, we can always start with the local solution from [61] and start with convex integration only at a positive (random) time.

4.2. Notion of Solution

To begin with, we write, formally,

$$u = z + v = z + z_1 + h,$$

where z and z_1 solve (3.2) and (3.4), respectively. Then the equation for h reads as

$$\begin{aligned} \mathcal{L}h + \operatorname{div}((h + z_1) \otimes (h + z_1) + z \otimes (h + z_1) + (h + z_1) \otimes z) + \nabla p^h &= 0, \\ \operatorname{div}h &= 0, \end{aligned} \tag{4.2}$$

which shall be further rewritten as a system for v^1 and v^2 in Sect. 4.3. Indeed, the multiplication of h and z is not well-defined as the expected sum of their regularities is not strictly positive for the resonant product $h \otimes z$ to be well-defined, cf. Lemma 2.2. Collecting the terms which make h too irregular leads to the following paracontrolled ansatz

$$h = -\mathbb{P}[h \prec \mathcal{I}\nabla z] + \vartheta - \mathbb{P}\mathcal{I} \operatorname{div}(z \otimes z_1 + z_1 \otimes z),$$

where $\mathcal{I}f(t) = \int_0^t P_{t-s} f(s) ds$.

Then ϑ becomes more regular than h since

$$\vartheta = h + \mathbb{P}\mathcal{I} \operatorname{div}(z \otimes z_1 + z_1 \otimes z) + \mathbb{P}\mathcal{I}[h \prec \nabla z] - \mathbb{P}[\mathcal{I}, h \prec] \nabla z, \tag{4.3}$$

where $[\mathcal{I}, h \prec] \nabla z$ denotes the commutator between \mathcal{I} and $h \prec$ given by

$$[\mathcal{I}, h \prec] \nabla z = \mathcal{I}[h \prec \nabla z] - h \prec \mathcal{I} \nabla z.$$

We use the same notation for other commutators as well. The second and the third terms on the right hand side of (4.3) cancel the irregular terms in h whereas the last term has better regularity by Lemma B.4. Using the commutator Lemmas B.4, 2.3 and 2.4, we write formally $h \circledast z$ as

$$\begin{aligned} h \circledast z &= -\mathbb{P}[h \prec \mathcal{I} \nabla z] \circledast z + \vartheta \circledast z - (\mathbb{P}\mathcal{I} \operatorname{div}(z \otimes z_1 + z_1 \otimes z)) \circledast z \\ &= -([\mathbb{P}, h \prec] \mathcal{I} \nabla z) \circledast z - \operatorname{com}(h, \mathbb{P}\mathcal{I} \nabla z, z) - h \cdot (\mathbb{P}\mathcal{I} \nabla z \circledast z) \\ &\quad + \vartheta \circledast z - (\mathbb{P}\mathcal{I} \operatorname{div}(z \otimes z_1 + z_1 \otimes z)) \circledast z. \end{aligned}$$

The above orange terms are still ill-defined and need to be replaced by the corresponding stochastic objects. Hence the rigorous paracontrolled ansatz and the definition of $h \circledast z$ read as

$$h = -\mathbb{P}[h \prec \mathcal{I} \nabla z] + \vartheta - (z \overset{\vee}{\mathcal{Y}} + z \overset{\vee}{\mathcal{Y}}), \tag{4.4}$$

$$h \circledast z := -([\mathbb{P}, h \prec] \mathcal{I} \nabla z) \circledast z - \operatorname{com}(h, \mathbb{P}\mathcal{I} \nabla z, z) - h \cdot z \overset{\vee}{\mathcal{Y}} + \vartheta \circledast z - z \overset{\vee}{\mathcal{Y}}, \tag{4.5}$$

Let us now formulate the definition of paracontrolled solution to (4.2). To this end, we recall that the given cylindrical Wiener process B is defined on a stochastic basis $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbf{P})$ and this is also where all the stochastic objects in Sect. 4.1 are constructed.

Definition 4.3. We say that a pair of $(\mathcal{F}_t)_{t \geq 0}$ -adapted processes

$$\begin{aligned} (h, \vartheta) &\in \left(L^2(0, T, L^2) \cap C_{T, 1/2-\kappa}^{1/10} L^{5/3} \cap L^1(0, T, B_{5/3, \infty}^{1/5}) \cap C_T H^{-1} \right) \\ &\quad \times \left(L^1(0, T, B_{1, \infty}^{3/5-\kappa}) \cap C_T H^{-1} \right) \end{aligned}$$

\mathbf{P} -a.s. with $\kappa > 0$ is a paracontrolled solution to (4.2) provided (4.4) holds and the equation (4.2) holds in the analytically weak sense with the resonant product $h \circledast z$ given by (4.5).

Remark 4.4. (i) We note that paracontrolled solutions are probabilistically strong, that is, they are adapted to the given filtration $(\mathcal{F}_t)_{t \geq 0}$.

(ii) The space $C_{T, 1/2-\kappa}^{1/10}$ with singularity at time zero is used in Definition 4.3 in order to cover the case of irregular initial conditions, namely, divergence free initial conditions in $C^{-1+2\kappa}$ for $\kappa > 0$.

(iii) The function spaces to which (h, ϑ) belong together with Proposition 4.1 guarantee that all the terms in (4.4)–(4.5) are well-defined. This in turn permits to make sense of all the terms in (4.2).

4.3. Formal Decomposition

We continue with the formal decomposition of the equation in order to find the desired paracontrolled ansatz. Moreover, in order to be able to apply convex integration, we need to introduce a further decomposition into v^1, v^2 . In this subsection, we ignore the fact that various products of z are not well-defined. We proceed as if these were well-defined and we replace their values in the end by the corresponding stochastic objects constructed in Sect. 4.1.

Let $h = v^1 + v^2$, (4.2) rewrites as

$$\begin{aligned} &\mathcal{L}(v^1 + v^2) + \operatorname{div}((v^1 + v^2 + z_1) \otimes (v^1 + v^2 + z_1)) \\ &\quad + z \otimes (v^1 + v^2 + z_1) + (v^1 + v^2 + z_1) \otimes z \\ &\quad + \nabla(p^1 + p^2) = 0, \quad \operatorname{div}(v^1 + v^2) = 0. \end{aligned} \tag{4.6}$$

The regular terms (encoded in blue) shall be put in the equation for v^2 , whereas the irregular ones (in magenta) into v^1 , so that in the end

$$\mathcal{L}v^1 + \operatorname{div}(z_1 \otimes z_1 + V^1 + V^{1,*}) + \nabla p^1 = 0, \quad \operatorname{div} v^1 = 0, \tag{4.7}$$

$$\mathcal{L}v^2 + \operatorname{div}((v^1 + v^2) \otimes (v^1 + v^2) + V^2 + V^{2,*}) + \nabla p^2 = 0, \quad \operatorname{div} v^2 = 0. \tag{4.8}$$

Here $V^{1,*}$ and $V^{2,*}$, respectively, denotes the transpose of V^1 and V^2 , respectively. First, we assign

$$(v^1 + v^2) \otimes (v^1 + v^2), \quad z_1 \otimes z_1 + z \otimes z_1 + z_1 \otimes z.$$

Then we decompose

$$\begin{aligned} (v^1 + v^2) \otimes z_1 &= (v^1 + v^2) \otimes \Delta_{>R}z_1 + (v^1 + v^2) \otimes \Delta_{\leq R}z_1 \\ &= (v^1 + v^2) \otimes \Delta_{>R}z_1 + v^1 \otimes \Delta_{>R}z_1 + v^2 \otimes \Delta_{>R}z_1 \\ &\quad + (v^1 + v^2) \otimes \Delta_{\leq R}z_1. \end{aligned}$$

These terms are included into V^1 and V^2 , whereas the symmetric counterparts

$$\begin{aligned} z_1 \otimes (v^1 + v^2) &= \Delta_{>R}z_1 \otimes (v^1 + v^2) + \Delta_{>R}z_1 \otimes v^1 \\ &\quad + \Delta_{>R}z_1 \otimes v^2 + \Delta_{\leq R}z_1 \otimes (v^1 + v^2) \end{aligned}$$

are in $V^{1,*}$ and $V^{2,*}$. Note that we colored $v^1 \otimes \Delta_{>R}z_1$ magenta. Indeed, it requires regularity of v^1 which is not true uniformly as $t \rightarrow 0$, since the initial condition is only in L^2 . Hence if colored blue, this term cannot be controlled in the convex integration scheme. This applies also to other terms below.

In the rest of the computation we only discuss the terms in V^1 and V^2 , the symmetric terms being included automatically. We split

$$\begin{aligned} (v^1 + v^2) \otimes z &= (v^1 + v^2) \otimes z + (v^1 + v^2) \otimes z + (v^1 + v^2) \otimes z \\ &= (v^1 + v^2) \otimes \Delta_{>R}z + v^1 \otimes \Delta_{>R}z \\ &\quad + v^2 \otimes \Delta_{>R}z + (v^1 + v^2) (\otimes + \otimes) \Delta_{\leq R}z + v^2 \otimes z + v^1 \otimes z. \end{aligned}$$

For the resonant product $v^1 \circledast z$ we shall use a paracontrolled ansatz which permits to cancel the worst terms in V^1 . In particular, the multiplication of v^1 and z is not well-defined as the expected sum of their regularities is not strictly positive making the resonant product $v^1 \circledast z$ ill-defined. First of all, there is the stochastic term $\text{div}(z \otimes z_1 + z_1 \otimes z)$ which makes v^1 too irregular. Second of all, it is the following paraproduct between v^1 and z

$$\text{div} \left(\Delta_{>R} z \otimes (v^1 + v^2) \right) = (v^1 + v^2) \prec \Delta_{>R} \nabla z,$$

which creates problems. This leads to the paracontrolled ansatz

$$\begin{aligned} v^1 &= -\mathbb{P}[(v^1 + v^2) \prec \Delta_{>R} \mathcal{I} \nabla z] + v^\sharp \\ &\quad - \mathbb{P} \mathcal{I} \text{div}(z \otimes z_1 + z_1 \otimes z), \quad \text{div } v^\sharp = 0, \end{aligned} \tag{4.9}$$

where $\mathcal{I}f(t) = \int_0^t e^{(t-s)\Delta} f(s) ds$. Thus,

$$\begin{aligned} v^\sharp &= v^1 + \mathbb{P} \mathcal{I} \text{div}(z \otimes z_1 + z_1 \otimes z) \\ &\quad + \mathcal{I} \mathbb{P}[(v^1 + v^2) \prec \Delta_{>R} \nabla z] - \mathbb{P}[\mathcal{I}, (v^1 + v^2) \prec] \Delta_{>R} \nabla z. \end{aligned} \tag{4.10}$$

This permits to cancel the two terms

$$-\mathbb{P} \mathcal{I} \text{div}(z \otimes z_1 + z_1 \otimes z) - \mathcal{I} \mathbb{P}[(v^1 + v^2) \prec \Delta_{>R} \nabla z]$$

from v^1 . Consequently, v^\sharp has a better regularity than v^1 and hence, in view of the commutator lemma, Lemma 2.3, the resonant product $v^1 \circledast z$ can be rigorously defined. To this end, we compute

$$\begin{aligned} \mathbb{P}[(v^1 + v^2) \prec \Delta_{>R} \mathcal{I} \nabla z] \circledast z &= \mathbb{P}[(v^1 + v^2) \prec \mathcal{I} \nabla z] \circledast z - \mathbb{P}[(v^1 + v^2) \prec \Delta_{\leq R} \mathcal{I} \nabla z] \circledast z \\ &= (\mathbb{P}, (v^1 + v^2) \prec] \mathcal{I} \nabla z \circledast z + [(v^1 + v^2) \prec \mathbb{P} \mathcal{I} \nabla z] \circledast z \\ &\quad - \mathbb{P}[(v^1 + v^2) \prec \Delta_{\leq R} \mathcal{I} \nabla z] \circledast z \\ &= (\mathbb{P}, (v^1 + v^2) \prec] \mathcal{I} \nabla z \circledast z + \text{com}(v^1 + v^2, \mathbb{P} \mathcal{I} \nabla z, z) \\ &\quad + (v^1 + v^2) \cdot (\mathbb{P} \mathcal{I} \nabla z \circledast z) \\ &\quad - \mathbb{P}[(v^1 + v^2) \prec \Delta_{\leq R} \mathcal{I} \nabla z] \circledast z. \end{aligned}$$

The above commutator between the Helmholtz projection and the paraproduct is understood componentwise as

$$(\mathbb{P}, (v^1 + v^2) \prec] \mathcal{I} \nabla z)^i = \mathbb{P}^{ij} [(v^1 + v^2)^k \prec \mathcal{I} \partial_k z^j] - (v^1 + v^2)^k \prec \mathbb{P}^{ij} \mathcal{I} \partial_k z^j,$$

and accordingly,

$$((v^1 + v^2) \prec \mathbb{P} \mathcal{I} \nabla z)^i = (v^1 + v^2)^k \prec \mathbb{P}^{ij} \mathcal{I} \partial_k z^j.$$

We deduce

$$\begin{aligned} v^1 \circledast z &= -\mathbb{P}[(v^1 + v^2) \prec \Delta_{>R} \mathcal{I} \nabla z] \circledast z \\ &\quad + v^\sharp \circledast z - (\mathbb{P} \mathcal{I} \text{div}(z \otimes z_1 + z_1 \otimes z)) \circledast z \\ &= -(\mathbb{P}, (v^1 + v^2) \prec] \mathcal{I} \nabla z \circledast z - \text{com}(v^1 + v^2, \mathbb{P} \mathcal{I} \nabla z, z) \\ &\quad - (v^1 + v^2) \cdot (\mathbb{P} \mathcal{I} \nabla z \circledast z) \end{aligned}$$

$$\begin{aligned}
 & -\mathbb{P}[(v^1 + v^2) \prec \Delta_{\leq R} \mathcal{I} \nabla z] \odot z \\
 & + v^\sharp \odot z - (\mathbb{P} \mathcal{I} \operatorname{div}(z \otimes z_1 + z_1 \otimes z)) \odot z.
 \end{aligned}$$

In order to avoid the singularity at time zero in the convex integration, we include an additional splitting into $\Delta_{>R}$ and $\Delta_{\leq R}$ of the remaining (uncolored) terms above as follows:

$$\begin{aligned}
 -(\mathbb{P}, (v^1 + v^2) \prec \mathcal{I} \nabla z) \odot z &= -(\mathbb{P}, v^1 \prec \mathcal{I} \nabla z) \odot \Delta_{>R} z \\
 &\quad - (\mathbb{P}, v^1 \prec \mathcal{I} \nabla z) \odot \Delta_{\leq R} z - (\mathbb{P}, v^2 \prec \mathcal{I} \nabla z) \odot z, \\
 -\operatorname{com}(v^1 + v^2, \mathbb{P} \mathcal{I} \nabla z, z) &= -\operatorname{com}(v^1, \mathbb{P} \mathcal{I} \nabla z, \Delta_{>R} z) \\
 &\quad - \operatorname{com}(v^1, \mathbb{P} \mathcal{I} \nabla z, \Delta_{\leq R} z) - \operatorname{com}(v^2, \mathbb{P} \mathcal{I} \nabla z, z), \\
 -(v^1 + v^2) \cdot (\mathbb{P} \mathcal{I} \nabla z \odot z) &= -(v^1 + v^2) \prec \Delta_{>R} (\mathbb{P} \mathcal{I} \nabla z \odot z) - v^1 \succ \Delta_{>R} (\mathbb{P} \mathcal{I} \nabla z \odot z) \\
 &\quad - v^2 \succ \Delta_{>R} (\mathbb{P} \mathcal{I} \nabla z \odot z) - (v^1 + v^2) \cdot \Delta_{\leq R} (\mathbb{P} \mathcal{I} \nabla z \odot z), \\
 v^\sharp \odot z &= v^\sharp \odot \Delta_{>R} z + v^\sharp \odot \Delta_{\leq R} z.
 \end{aligned}$$

Finally, collecting all the terms leads us to

$$\begin{aligned}
 V^1 &= z_1 \otimes z + (v^1 + v^2) \otimes \Delta_{>R} z_1 + v^1 \otimes \Delta_{>R} z_1 + (v^1 + v^2) \otimes \Delta_{>R} z + v^1 \otimes \Delta_{>R} z \\
 &\quad - (\mathbb{P} \mathcal{I} \operatorname{div}(z \otimes z_1 + z_1 \otimes z)) \odot z - (\mathbb{P}, v^1 \prec \mathcal{I} \nabla z) \odot \Delta_{>R} z - \operatorname{com}(v^1, \mathbb{P} \mathcal{I} \nabla z, \Delta_{>R} z) \\
 &\quad - (v^1 + v^2) \prec \Delta_{>R} (\mathbb{P} \mathcal{I} \nabla z \odot z) - v^1 \succ \Delta_{>R} (\mathbb{P} \mathcal{I} \nabla z \odot z) + v^\sharp \odot \Delta_{>R} z, \\
 V^2 &= v^2 \otimes \Delta_{>R} z_1 + (v^1 + v^2) \otimes \Delta_{\leq R} z_1 \\
 &\quad + v^2 \otimes \Delta_{>R} z + (v^1 + v^2) \otimes \Delta_{\leq R} z + v^2 \odot z \\
 &\quad - \mathbb{P}[(v^1 + v^2) \prec \Delta_{\leq R} \mathcal{I} \nabla z] \odot z - (\mathbb{P}, v^2 \prec \mathcal{I} \nabla z) \odot z - (\mathbb{P}, v^1 \prec \mathcal{I} \nabla z) \odot \Delta_{\leq R} z \\
 &\quad - \operatorname{com}(v^1, \mathbb{P} \mathcal{I} \nabla z, \Delta_{\leq R} z) - \operatorname{com}(v^2, \mathbb{P} \mathcal{I} \nabla z, z) - v^2 \succ \Delta_{>R} (\mathbb{P} \mathcal{I} \nabla z \odot z) \\
 &\quad - (v^1 + v^2) \cdot \Delta_{\leq R} (\mathbb{P} \mathcal{I} \nabla z \odot z) + v^\sharp \odot \Delta_{\leq R} z.
 \end{aligned}$$

It will be seen below, that letting

$$h = v^1 + v^2, \quad \vartheta = v^\sharp + v^2 + \mathbb{P}[(v^1 + v^2) \prec \Delta_{\leq R} \mathcal{I} \nabla z]$$

satisfies the requirements of Definition 4.3. The idea then is to apply convex integration on the level of the equation (4.8) for v^2 . In particular, we need to make sure that the convex integration gives v^2 of the required regularity for ϑ .

5. Convex integration Set-up and Results

The goal of this section is to construct infinitely many probabilistically strong paracontrolled solutions (h, ϑ) and deduce global existence and non-uniqueness for the system (4.2). The equation (4.2) for h splits formally into the coupled system (4.7), (4.8) for v^1 and v^2 and it remains to include the stochastic objects. This leads to

$$\begin{aligned}
 \mathcal{L}v^1 + \operatorname{div}(z \overset{\vee}{\nabla} + V^1 + V^{1,*}) + \nabla p^1 &= 0, \\
 \mathcal{L}v^2 + \operatorname{div}((v^1 + v^2) \otimes (v^1 + v^2) + V^2 + V^{2,*}) + \nabla p^2 &= 0, \\
 \operatorname{div} v^1 = \operatorname{div} v^2 = 0, \quad v^1(0) = v_0, \quad v^2(0) = 0, & \quad (5.1)
 \end{aligned}$$

with

$$\begin{aligned}
 V^1 &= z \overset{\vee}{\mathcal{Y}} + (v^1 + v^2) \otimes \Delta_{>Rz} \overset{\vee}{\mathcal{Y}} + v^1 \otimes \Delta_{>Rz} \overset{\vee}{\mathcal{Y}} + (v^1 + v^2) \otimes \Delta_{>Rz} + v^1 \otimes \Delta_{>Rz} \\
 &\quad - z \overset{\vee}{\mathcal{Y}} - ([\mathbb{P}, v^1 \prec] \mathcal{I} \nabla z) \otimes \Delta_{>Rz} - \text{com}(v^1, \mathbb{P} \mathcal{I} \nabla z, \Delta_{>Rz}) \\
 &\quad - (v^1 + v^2) \prec \Delta_{>Rz} \overset{\vee}{\mathcal{Y}} - v^1 \succ \Delta_{>Rz} \overset{\vee}{\mathcal{Y}} + v^\# \otimes \Delta_{>Rz}
 \end{aligned}$$

and

$$\begin{aligned}
 V^2 &= v^2 \otimes \Delta_{>Rz} \overset{\vee}{\mathcal{Y}} + (v^1 + v^2) \otimes \Delta_{\leq Rz} \overset{\vee}{\mathcal{Y}} \\
 &\quad + v^2 \otimes \Delta_{>Rz} + (v^1 + v^2) (\otimes + \otimes) \Delta_{\leq Rz} + v^2 \otimes z \\
 &\quad - \mathbb{P}[(v^1 + v^2) \prec \Delta_{\leq R} \mathcal{I} \nabla z] \otimes z - ([\mathbb{P}, v^2 \prec] \mathcal{I} \nabla z) \otimes z - ([\mathbb{P}, v^1 \prec] \mathcal{I} \nabla z) \otimes \Delta_{\leq Rz} \\
 &\quad - \text{com}(v^1, \mathbb{P} \mathcal{I} \nabla z, \Delta_{\leq Rz}) - \text{com}(v^2, \mathbb{P} \mathcal{I} \nabla z, z) - v^2 \succ \Delta_{>Rz} \overset{\vee}{\mathcal{Y}} \\
 &\quad - (v^1 + v^2) \cdot \Delta_{\leq Rz} \overset{\vee}{\mathcal{Y}} + v^\# \otimes \Delta_{\leq Rz}.
 \end{aligned}$$

The paracontrolled ansatz for v^1 reads as

$$v^1 = -\mathbb{P}[(v^1 + v^2) \prec \Delta_{>R} \mathcal{I} \nabla z] + v^\# - (z \overset{\vee}{\mathcal{Y}} + z \overset{\vee}{\mathcal{Y}}). \tag{5.2}$$

These equations need to be considered together within the convex integration scheme and we put forward a joint iterative procedure.

The convex integration iteration is indexed by a parameter $q \in \mathbb{N}_0$. It will be seen that the Reynolds stress \mathring{R}_q is only required for the approximations v_q^2 of v^2 , whereas the approximations v_q^1 of v^1 are obtained by a fixed point argument. We consider an increasing sequence $\{\lambda_q\}_{q \in \mathbb{N}_0} \subset \mathbb{N}$ which diverges to ∞ , and a sequence $\{\delta_q\}_{q \in \mathbb{N}_0} \subset (0, 1)$ which decreases to 0. We choose $a \in \mathbb{N}$, $b \in \mathbb{N}$, $\beta \in (0, 1)$ and let

$$\lambda_q = a^{(b^q)}, \quad \delta_q = \lambda_1^{2\beta} \lambda_q^{-2\beta},$$

where β will be chosen sufficiently small and a as well as b sufficiently large. At each step q , a triple $(v_q^1, v_q^2, \mathring{R}_q)$ is constructed solving the system

$$\begin{aligned}
 \mathcal{L}v_q^1 + \text{div}(z \overset{\vee}{\mathcal{Y}} + V_q^1 + V_q^{1,*}) + \nabla p_q^1 &= 0, \\
 \mathcal{L}v_q^2 + \text{div}((v_q^1 + v_q^2) \otimes (v_q^1 + v_q^2) + V_q^2 + V_q^{2,*}) + \nabla p_q^2 &= \text{div } \mathring{R}_q, \\
 \text{div } v_q^1 = \text{div } v_q^2 = 0, \quad v_q^1(0) = v_0, \quad v_q^2(0) = 0,
 \end{aligned} \tag{5.3}$$

where

$$\begin{aligned}
 V_q^1 &= z \overset{\vee}{\mathcal{Y}} + (v_q^1 + v_q^2) \otimes \Delta_{\leq f(q)} \Delta_{>R}(z + z \overset{\vee}{\mathcal{Y}}) + v_q^1 \otimes \Delta_{>Rz} \overset{\vee}{\mathcal{Y}} + v_q^1 \otimes \Delta_{\leq f(q)} \Delta_{>Rz} \\
 &\quad - ([\mathbb{P}, v_q^1 \prec] \mathcal{I} \nabla z) \otimes \Delta_{>Rz} - \text{com}(v_q^1, \mathbb{P} \mathcal{I} \nabla z, \Delta_{>Rz}) - z \overset{\vee}{\mathcal{Y}} \\
 &\quad - (v_q^1 + v_q^2) \prec \Delta_{\leq f(q)} \Delta_{>Rz} \overset{\vee}{\mathcal{Y}} - v_q^1 \succ \Delta_{>Rz} \overset{\vee}{\mathcal{Y}} + v_q^\# \otimes \Delta_{>Rz}
 \end{aligned}$$

and

$$V_q^2 = v_q^2 \otimes \Delta_{>Rz} \overset{\vee}{\mathcal{Y}} + (v_q^1 + v_q^2) \otimes \Delta_{\leq Rz} \overset{\vee}{\mathcal{Y}}$$




Fig. 1. Iteration scheme

$$\begin{aligned}
 &+v_q^2 \otimes \Delta_{>Rz} + (v_q^1 + v_q^2) (\otimes + \otimes) \Delta_{\leq Rz} + v_q^2 \otimes z \\
 &-\mathbb{P}[(v_q^1 + v_q^2) < \Delta_{\leq R\mathcal{I}\nabla z}] \otimes z \\
 &-(\mathbb{P}, v_q^2 < \mathcal{I}\nabla z) \otimes z - (\mathbb{P}, v_q^1 < \mathcal{I}\nabla z) \otimes \Delta_{\leq Rz} \\
 &-\text{com}(v_q^1, \mathbb{P}\mathcal{I}\nabla z, \Delta_{\leq Rz}) - \text{com}(v_q^2, \mathbb{P}\mathcal{I}\nabla z, z) - v_q^2 \succcurlyeq \Delta_{>Rz} \checkmark \\
 &-(v_q^1 + v_q^2) \cdot \Delta_{\leq Rz} \checkmark + v_q^\sharp \otimes \Delta_{\leq Rz}.
 \end{aligned}$$

Here, V_q^1 and V_q^2 are obtained from V^1 and V^2 , respectively, by replacing v^1, v^2, v^\sharp by v_q^1, v_q^2, v_q^\sharp and adding the projector $\Delta_{\leq f(q)}$, where $2^{f(q)} = \lambda_q^\theta, \theta = 10/21$, into the second, the fourth and the eighth term in V_q^1 . These projectors are used to control the blow-up of certain norms of v_q^1 and v_q^2 and also to prove the convergence of v_q^1 as $q \rightarrow \infty$. We shall therefore require the parameter a to be a power of 2^{21} and $b \in \mathbb{N}$. The parameter R is chosen in (6.7) below.

Analogously to the paracontrolled ansatz (5.2), v_q^1, v_q^2 and v_q^\sharp are linked via

$$v_q^1 = -\mathbb{P}[(v_q^1 + v_q^2) < \mathcal{I}(\nabla \Delta_{>R} \Delta_{\leq f(q)} z)] + v_q^\sharp - (z \checkmark + z \checkmark). \tag{5.4}$$

Our main goal is to prove convergence of v_q^1, v_q^2 and v_q^\sharp as $q \rightarrow \infty$ and to show that their limits satisfy (5.1), (5.2) in order to recover a paracontrolled solution to (4.2) in the sense of Definition 4.3.

See Fig. 1 for our iteration scheme. More precisely, we use v_q^2 to determine v_q^1 and v_q^\sharp by Schauder estimates. Then \mathring{R}_q is determined by v_q^1, v_q^\sharp and v_q^2 . The next velocity v_{q+1}^2 is only determined by \mathring{R}_q via a convex integration argument.

As the next step, we define a stopping time which controls suitable norms of all the required stochastic objects. Namely, for $L > 1$ we let

$$\begin{aligned}
 T_L &:= T_L^1 \wedge T_L^2 \wedge T_L^3 \wedge T_L^4 \wedge T_L^5 \wedge L^{1/2}, \\
 T_L^1 &:= \inf \left\{ t \geq 0, \|z(t)\|_{C^{-1/2-k}} \geq L^{1/2} \right\} \wedge \inf \left\{ t \geq 0, \|z\|_{C_t^{1/10} C^{-7/10-k}} \geq L^{1/2} \right\}, \\
 T_L^2 &:= \inf \left\{ t \geq 0, \|z \checkmark(t)\|_{C^{-k}} \geq L \right\} \wedge \inf \left\{ t \geq 0, \|z \checkmark\|_{C_t^{1/10} C^{-1/5-k}} \geq L \right\}, \\
 T_L^3 &:= \inf \left\{ t \geq 0, \|z \checkmark \checkmark(t)\|_{C^{-k}} \geq L \right\} \wedge \inf \left\{ t \geq 0, \|z \checkmark \checkmark(t)\|_{C^{-1/2-k}} + \|z \checkmark \checkmark(t)\|_{C^{-1/2-k}} \geq L \right\}, \\
 T_L^4 &:= \inf \left\{ t \geq 0, \sup_{0 \leq s < t} \|z \checkmark \checkmark(r; s)\|_{C^{-k}} \geq L \right\} \wedge \inf \left\{ t \geq 0, \|z \checkmark \checkmark\|_{C_t^{1/10} C^{-1/5-k}} \geq L \right\}, \\
 T_L^5 &:= \inf \left\{ t \geq 0, \sup_{0 \leq s < t} \|z \checkmark \checkmark \checkmark(r; s)\|_{C^{-k}} \geq L \right\}, \tag{5.5}
 \end{aligned}$$

where we denoted by $z^{\mathcal{Y}}(r; s)$ and $z^{\mathcal{V}}(r; s)$ the stochastic objects obtained the same way as $z^{\mathcal{Y}}(r)$ and $z^{\mathcal{V}}(r)$ but replacing the last integration operator $\mathcal{I} = \mathcal{I}_{0,r}$ by $\mathcal{I}_{s,r} = \int_s^r e^{(r-l)\Delta} dl$. It follows from Proposition 4.1 that the stopping time T_L is \mathbf{P} -a.s. strictly positive and it holds that $T_L \uparrow \infty$ as $L \rightarrow \infty$ \mathbf{P} -a.s.

We intend to solve (5.1) for any given divergence free initial condition $v_0 \in L^2 \cup C^{-1+\kappa}$ measurable with respect to \mathcal{F}_0 . However, in the first step, we take the following additional assumption: Let $N \geq 1$ be given and assume that \mathbf{P} -a.s.

$$\|v_0\|_{L^2} \leq N. \tag{5.6}$$

We keep this additional assumption on the initial condition throughout the convex integration step in Proposition 5.1. In Theorem 5.4 it is relaxed to $v_0 \in L^2$ \mathbf{P} -a.s and, finally, Corollary 5.6 proves the result if $v_0 \in C^{-1+\kappa}$ \mathbf{P} -a.s. We also suppose that there is a deterministic constant $M_L(N)^{1/2} \geq L^6 N + L^{29}$. In the following we write M_L instead of $M_L(N)$ for simplicity.

Let $\alpha \in (0, 1)$ be a small parameter to be chosen below. By induction on q we assume the following bounds for the iterations v_q^2 : if $t \in [0, T_L]$ then, for $p = \frac{32}{32-7\alpha}$,

$$\begin{aligned} \|v_q^2\|_{C_t W^{2/3,p}} &\leq a^{-\alpha/2} M_L^{1/2} (1 + \sum_{1 \leq r \leq q} \delta_r^{1/2}) \leq 3M_L^{1/2} a^{-\alpha/2}, \\ \|v_q^2\|_{C_t^{1/10} L^{5/3}} + \|v_q^2\|_{C_t W^{1/5,5/3}} &\leq a^{-\alpha/2} M_L^{1/2} (1 + \sum_{1 \leq r \leq q} \delta_r^{1/2}) \leq 3M_L^{1/2} a^{-\alpha/2}. \end{aligned} \tag{5.7}$$

Later on, we use the factor $a^{-\alpha/2}$ to absorb an implicit constant. Here we defined $\sum_{1 \leq r \leq 0} := 0$. In addition, we used $\sum_{r \geq 1} \delta_r^{1/2} \leq \sum_{r \geq 1} a^{b\beta - r b\beta} = \frac{1}{1 - a^{-b\beta}} \leq 2$ which boils down to the requirement that

$$a^{b\beta} \geq 2, \tag{5.8}$$

which we assume from now on. We also assume that L is large enough such that the implicit constant in (5.20) and (7.54) below can be absorbed by L . Moreover, for such L we can always choose a large enough such that $L \leq a^{\alpha/16}$. We denote

$$\sigma_q = 2^{-q}, q \in \mathbb{N}_0 \cup \{-1\}, \quad \gamma_q = 2^{-q}, q \in \mathbb{N}_0 \setminus \{3\}, \quad \gamma_3 = K,$$

for $K > 0$ arbitrary. This constant will be used in order to distinguish different solutions.

The key result is the following iterative proposition, which we prove below in Sect. 7:

Proposition 5.1. *Let $N \geq 1$ and let $L > 1$ sufficiently large. There exists a choice of parameters a, α, b, β such that the following holds true: Let $(v_q^1, v_q^2, \mathring{R}_q)$ for some $q \in \mathbb{N}_0$ be an $(\mathcal{F}_t)_{t \geq 0}$ -adapted solution to (5.3), (5.4) satisfying (5.7) and*

$$\|v_q^2(t)\|_{L^2} \leq \begin{cases} M_0(M_L^{1/2} \sum_{r=1}^q \delta_r^{1/2} + \sum_{r=1}^q \gamma_r^{1/2}) + 3M_0(M_L(1 + 3q))^{1/2}, & t \in (\frac{\sigma_q - 1}{2} \wedge T_L, T_L], \\ 0, & t \in [0, \frac{\sigma_q - 1}{2} \wedge T_L], \end{cases} \tag{5.9}$$

for a universal constant M_0 ,

$$\|v_q^2\|_{C_{t,x}^1} \leq \lambda_q^4 M_L^{1/2}, \quad t \in [0, T_L], \tag{5.10}$$

$$\|\mathring{R}_q(t)\|_{L^1} \leq \delta_{q+1} M_L, \quad t \in (\sigma_{q-1} \wedge T_L, T_L], \tag{5.11}$$

$$\|\mathring{R}_q(t)\|_{L^1} \leq M_L(1 + 3q), \quad t \in [0, T_L]. \tag{5.12}$$

Then there exists an $(\mathcal{F}_t)_{t \geq 0}$ -adapted process $(v_{q+1}^1, v_{q+1}^2, \mathring{R}_{q+1})$ which solves (5.3), (5.4) on the level $q + 1$ and satisfies

$$\begin{aligned} & \|v_{q+1}^2(t) - v_q^2(t)\|_{L^2} \\ & \leq \begin{cases} M_0(M_L^{1/2} \delta_{q+1}^{1/2} + \gamma_{q+1}^{1/2}), & t \in (4\sigma_q \wedge T_L, T_L], \\ M_0((M_L(1 + 3q))^{1/2} + \gamma_{q+1}^{1/2}), & t \in (\frac{\sigma_q}{2} \wedge T_L, 4\sigma_q \wedge T_L], \\ 0, & t \in [0, \frac{\sigma_q}{2} \wedge T_L], \end{cases} \end{aligned} \tag{5.13}$$

$$\begin{aligned} & \|\mathring{R}_{q+1}(t)\|_{L^1} \\ & \leq \begin{cases} M_L \delta_{q+2}, & t \in (\sigma_q \wedge T_L, T_L], \\ M_L \delta_{q+2} + \sup_{s \in [(t-\sigma_q/2) \vee 0, t]} \|\mathring{R}_q(s)\|_{L^1}, & t \in (\frac{\sigma_q}{2} \wedge T_L, \sigma_q \wedge T_L], \\ \sup_{s \in [(t-\sigma_q/2) \vee 0, t]} \|\mathring{R}_q(s)\|_{L^1} + 3M_L & t \in [0, \frac{\sigma_q}{2} \wedge T_L]. \end{cases} \end{aligned} \tag{5.14}$$

Consequently, $(v_{q+1}^2, \mathring{R}_{q+1})$ obeys (5.7), (5.9), (5.10), (5.11) and (5.12) at the level $q + 1$. Furthermore, for $1 < p = \frac{32}{32-7\alpha}$, $t \in [0, T_L]$ it holds that

$$\|v_{q+1}^2(t) - v_q^2(t)\|_{W^{2/3,p}} \leq M_L^{1/2} \delta_{q+1}^{1/2} a^{-\alpha/2}, \tag{5.15}$$

$$\|v_{q+1}^2 - v_q^2\|_{C_t^{1/10} L^{5/3}} + \|v_{q+1}^2 - v_q^2\|_{C_t W^{1/5,5/3}} \leq M_L^{1/2} \delta_{q+1}^{1/2} a^{-\alpha/2}, \tag{5.16}$$

and for $t \in (4\sigma_q \wedge T_L, T_L]$ we have

$$|\|v_{q+1}^2\|_{L^2}^2 - \|v_q^2\|_{L^2}^2 - 3\gamma_{q+1}| \leq 7M_L \delta_{q+1}. \tag{5.17}$$

Note that no bounds on v_q^1, v_{q+1}^1 were included in the statement of Proposition 5.1. Indeed, the definition of the new velocity v_{q+1}^2 does not require v_{q+1}^1 . Then, having v_{q+1}^2 at hand, all the necessary bounds for $v_{q+1}^1, v_{q+1}^\sharp$ follow from Sect. 6 below. In particular, in Sect. 6 we prove

Proposition 5.2. *Under the assumptions of Proposition 5.1, it holds for $\kappa > 0$ and $t \in [0, T_L]$*

$$\|v_q^1\|_{C_{t,1/6} B_{5/3}^{1/3-2\kappa}} + \|v_q^1\|_{C_{t,1/6}^{1/6-\kappa} L^{5/3}} + \|v_q^1\|_{C_t L^2} \lesssim a^{-\alpha/4} M_L^{1/2} + L^3 N + L^4, \tag{5.18}$$

$$\|v_q^\sharp\|_{C_{t,3/10} B_{5/3}^{3/5-\kappa}} + \|v_q^\sharp\|_{C_{t,3/10}^{1/20} B_{5/3}^{11/20-2\kappa}} + \|v_q^\sharp\|_{C_t L^2} \lesssim a^{-\alpha/8} M_L^{1/2} + L^5 N + L^6,$$

$$\|v_{q+1}^1 - v_q^1\|_{C_t L^2} + \|v_{q+1}^1 - v_q^1\|_{C_t B_{5/3}^{1/3-\kappa}} + \|v_{q+1}^1 - v_q^1\|_{C_t^{1/6-\kappa} L^{5/3}}$$

$$\begin{aligned} &\lesssim a^{-\alpha/2} M_L^{1/2} \delta_{q+1}^{1/2} + M_L^{1/2} \lambda_q^{-\theta/20}, \\ \|v_{q+1}^\sharp - v_q^\sharp\|_{C_{t,3/10} B_{\delta_{q+1}}^{3/5-\kappa}} + \|v_{q+1}^\sharp - v_q^\sharp\|_{C_t L^2} &\lesssim a^{-\alpha/2} M_L^{1/2} \delta_{q+1}^{1/2} + M_L^{1/2} \lambda_q^{-\theta/20}. \end{aligned} \tag{5.19}$$

Here $\theta = 10/21$ and the implicit constants are always universal and independent of q .

In what follows we always use $\kappa > 0$ to denote a small constant.

Remark 5.3. The best regularity we can expect for v^1 is $1/2 - \kappa$ whereas for v^\sharp it is $1 - \kappa$. It will be seen in Sect. 6 that their integrability is determined by v^2 and hence it comes from the convex integration argument. Here, we observe a competition between regularity and integrability, cf. (5.15) and (5.16) and their proofs in Sects. 7.4.3 and 7.4.5. For convenience, we have chosen integrability $5/3$ and space regularity $1/3 - 2\kappa$ for v^1 and $3/5 - \kappa$ for v^\sharp . The time weights are dictated by the desired space regularity as the initial value for v_q^1 and v_q^\sharp only belongs to L^2 . We also note that the bounds in Proposition 5.2 do not rely on the $W^{2/3,p}$ estimate of v_q^2 or the difference $v_{q+1}^2 - v_q^2$. Indeed, this is only needed to make sense of the resonant product $v^2 \circledast z$ in the limit and to control the corresponding part of the Reynolds stress.

We intend to start the iteration from $v_0^2 \equiv 0$ on $[0, T_L]$. Then (5.7), (5.9) and (5.10) hold. In that case, \mathring{R}_0 is the trace-free part of the matrix

$$v_0^1 \otimes v_0^1 + V_0^2 + V_0^{2,*},$$

where

$$\begin{aligned} V_0^2 &= v_0^1 \otimes \Delta_{\leq R z} \mathring{Y} + v_0^1 (\otimes + \oslash) \Delta_{\leq R z} \\ &\quad - \mathbb{P}[v_0^1 \prec \Delta_{\leq R} \mathcal{I} \nabla z] \circledast z - ([\mathbb{P}, v_0^1 \prec] \mathcal{I} \nabla z) \circledast \Delta_{\leq R z} \\ &\quad - \text{com}(v_0^1, \mathbb{P} \mathcal{I} \nabla z, \Delta_{\leq R z}) - v_0^1 \cdot \Delta_{\leq R z} \mathring{y} + v_0^\sharp \circledast \Delta_{\leq R z}. \end{aligned}$$

By (5.18) and (5.19), paraproduct estimates Lemma 2.2, commutator estimates Lemmas 2.3, 2.4, we have

$$\begin{aligned} \|R_0(t)\|_{L^1} &\lesssim \|v_0^1\|_{L^2} (\|\Delta_{\leq R z} \mathring{Y}\|_{C^\kappa} + \|\Delta_{\leq R z} \mathring{y}\|_{C^\kappa} + \|\Delta_{\leq R z}\|_{C^\kappa}) \\ &\quad + \|v_0^1\|_{L^2}^2 + \|v_0^\sharp\|_{L^2} \|\Delta_{\leq R z}\|_{C^\kappa} \\ &\quad + \|v_0^1\|_{L^2} \|z\|_{C^{-1/2-\kappa}} \|\Delta_{\leq R z}\|_{C^{-1/2+2\kappa}} \\ &\lesssim a^{-\alpha/8} M_L + L^6 N^2 + L^8 + L(a^{-\alpha/8} M_L^{1/2} + L^5 N + L^6) 2^{(1/2+2\kappa)R} \\ &\leq M_L. \end{aligned} \tag{5.20}$$

Here we used the value of R from (6.7) in Sect. 6 and the implicit constant can be absorbed by taking a and L large enough. Thus (5.11) as well as (5.12) are satisfied on the level $q = 0$, since $\delta_1 = 1$.

We deduce the following result:

Theorem 5.4. *There exists a \mathbf{P} -a.s. strictly positive stopping time T_L , arbitrarily large by choosing L large, such that for any \mathcal{F}_0 -measurable divergence free initial condition $v_0 \in L^2$ \mathbf{P} -a.s. the following holds true: There exists an $(\mathcal{F}_t)_{t \geq 0}$ -adapted process (v^1, v^2, v^\sharp) such that for $\kappa > 0$*

$$\begin{aligned} v^1 &\in C([0, T_L]; L^2) \cap L^1(0, T_L; B_{5/3, \infty}^{1/3-2\kappa}) \cap C_{T_L, 1/6}^{1/6-\kappa} L^{5/3}, \\ v^2 &\in L^p(0, T_L; L^2) \cap C([0, T_L], W^{2/3, 1} \cap W^{1/5, 5/3}) \cap C_{T_L}^{1/10} L^{5/3}, \\ v^\sharp &\in C([0, T_L]; L^2) \cap L^1(0, T_L; B_{5/3, \infty}^{3/5-\kappa}), \end{aligned}$$

\mathbf{P} -a.s. for all $p \in [1, \infty)$, and it is an analytically weak solution to (5.1) with $v^1(0) = v_0, v^2(0) = 0$ and satisfying (5.2). Furthermore, there are infinitely many such solutions and also infinitely many paracontrolled solutions $(h, \vartheta) = (v^1 + v^2, v^\sharp + v^2 + \mathbb{P}[(v^1 + v^2) \prec \Delta_{\leq R} \mathcal{I} \nabla z])$ on $[0, T_L]$ satisfying

$$\begin{aligned} h &\in C([0, T_L]; L^{5/3}) \cap L^1(0, T_L; B_{5/3, \infty}^{1/5}) \cap C_{T_L, 1/6}^{1/10} L^{5/3} \cap L^p(0, T_L; L^2), \\ \vartheta &\in C([0, T_L]; L^{5/3}) \cap L^1(0, T_L; B_{1, \infty}^{3/5-\kappa}), \end{aligned}$$

\mathbf{P} -a.s. for all $p \in [1, \infty)$, where R depends on L and is chosen in (6.7) below.

Proof. Letting $v_0^2 \equiv 0$, we repeatedly apply Proposition 5.1 and obtain $(\mathcal{F}_t)_{t \geq 0}$ -adapted processes $(v_q^1, v_q^2, \mathring{R}_q), q \in \mathbb{N}$, such that

$$v_q^2 \rightarrow v^2 \quad \text{in } C([0, T_L], W^{2/3, 1} \cap W^{1/5, 5/3}) \cap C_{T_L}^{1/10} L^{5/3}$$

as a consequence of (5.15), (5.16) and (5.7). In view of Proposition 5.2, it follows that

$$\begin{aligned} v_q^1 &\rightarrow v^1 \quad \text{in } C([0, T_L]; L^2) \cap L^1(0, T_L; B_{5/3, \infty}^{1/3-2\kappa}) \cap C_{T_L, 1/6}^{1/6-\kappa} L^{5/3}, \\ v_q^\sharp &\rightarrow v^\sharp \quad \text{in } C([0, T_L]; L^2) \cap L^1(0, T_L; B_{5/3, \infty}^{3/5-\kappa}). \end{aligned}$$

Then (v^1, v^2, v^\sharp) are $(\mathcal{F}_t)_{t \geq 0}$ -adapted. Moreover, using (5.13) we have, for every $p \in [1, \infty)$,

$$\begin{aligned} \int_0^{T_L} \|v_{q+1}^2 - v_q^2\|_{L^2}^p dt &\leq \int_{\sigma_q/2 \wedge T_L}^{4\sigma_q \wedge T_L} \|v_{q+1}^2 - v_q^2\|_{L^2}^p dt + \int_{4\sigma_q \wedge T_L}^{T_L} \|v_{q+1}^2 - v_q^2\|_{L^2}^p dt \\ &\lesssim \int_{\sigma_q/2 \wedge T_L}^{4\sigma_q \wedge T_L} M_0^p ((M_L(1+3q))^{1/2} + \gamma_{q+1}^{1/2})^p dt \\ &\quad + \int_{4\sigma_q \wedge T_L}^{T_L} M_0^p (M_L^{1/2} \delta_{q+1}^{1/2} + \gamma_{q+1}^{1/2})^p dt \\ &\lesssim M_0^p \left(2^{-q} ((M_L(1+3q))^{1/2} + \gamma_{q+1}^{1/2})^p + T_L (M_L^{1/2} \delta_{q+1}^{1/2} + \gamma_{q+1}^{1/2})^p \right). \end{aligned}$$

Thus, the sequence $v_q^2, q \in \mathbb{N}$, is Cauchy hence converging in $L^p(0, T_L; L^2)$ for all $p \in [1, \infty)$. Accordingly, $v_q^2 \rightarrow v^2$ also in $L^p(0, T_L; L^2)$. Furthermore, by (5.11), (5.12) we know, for all $p \in [1, \infty)$,

$$\int_0^{T_L} \|\mathring{R}_q(t)\|_{L^1}^p dt \lesssim M_L^p \delta_{q+1}^p T_L + (M_L(1+3q))^p 2^{-q} \rightarrow 0, \quad \text{as } q \rightarrow \infty.$$

Thus, the process (v^1, v^2, v^\sharp) satisfies (5.1) and (5.2) before T_L in the analytically weak sense. Since $v_q^1(0) = v_0$ and $v_q^2(0) = 0$ for all $q \in \mathbb{N}_0$ we deduce that $v^1(0) = v_0$ and $v^2(0) = 0$. Thus (h, ϑ) defined above solves (4.2) in the sense of Definition 4.3.

Next, we prove non-uniqueness of the constructed solutions. In view of (5.17), we have, on $t \in (4\sigma_0 \wedge T_L, T_L]$,

$$\begin{aligned} \left| \|v^2\|_{L^2}^2 - 3K \right| &\leq \left| \sum_{q=0}^{\infty} (\|v_{q+1}^2\|_{L^2}^2 - \|v_q^2\|_{L^2}^2 - 3\gamma_{q+1}) \right| + 3 \sum_{q \neq 2} \gamma_{q+1} \\ &\leq 7M_L \sum_{q=0}^{\infty} \delta_{q+1} + 3 \sum_{q \neq 2} \gamma_{q+1} \leq 7M_L \sum_{q=0}^{\infty} \delta_{q+1} + 3 \sum_{q \neq 2} \gamma_{q+1} \leq c, \end{aligned} \tag{5.21}$$

where the constant $c > 0$ is independent of K and the parameters a, α . This implies non-uniqueness by choosing different K . More precisely, for a given $L \geq 1$ sufficiently large it holds $\mathbf{P}(4\sigma_0 < T_L) > 0$. The parameters L, N determine $M_L(N)$ and consequently by choosing different $K = K(L, N)$ and $K' = K'(L, N)$ so that $3|K - K'| > 2c$ we deduce that the corresponding solutions v_K^2 and $v_{K'}^2$ have different L^2 -norms on the set $\{4\sigma_0 < T_L\}$. We claim that the sums $v_K^1 + v_K^2$ and $v_{K'}^1 + v_{K'}^2$ are different as well. Indeed, it is easy to see from (5.18) and (5.21) that

$$\sqrt{3K - c} - M_L^{1/2} \leq \|v^1 + v^2\|_{L^2} \leq M_L^{1/2} + \sqrt{3K + c}.$$

Choosing K' such that $\sqrt{3K' - c} - M_L^{1/2} > M_L^{1/2} + \sqrt{3K + c}$ gives different solutions.

For a general divergence free initial condition $v_0 \in L^2$ \mathbf{P} -a.s., we define

$$\Omega_N := \{N - 1 \leq \|v_0\|_{L^2} < N\} \in \mathcal{F}_0.$$

Then the first part of this proof gives the existence of infinitely many paracontrolled solutions (h^N, ϑ^N) on each Ω_N . Letting

$$h := \sum_{N \in \mathbb{N}} h^N 1_{\Omega_N}, \quad \vartheta := \sum_{N \in \mathbb{N}} \vartheta^N 1_{\Omega_N}$$

concludes the proof. Note that this also uses the fact that the stochastic objects are defined in advance and then the rest of the construction proceeds pathwise. \square

By an argument similar to [45, Theorem 1.1] we may extend the paracontrolled solutions obtained in Theorem 5.4 by other paracontrolled solutions in order to obtain global existence and non-uniqueness.

Theorem 5.5. *Let $v_0 \in L^2$ \mathbf{P} -a.s. be an \mathcal{F}_0 -measurable divergence free initial condition. There exist infinitely many paracontrolled solutions (h, ϑ) to (4.2) on $[0, \infty)$. Moreover, it holds that*

$$h \in L_{\text{loc}}^p([0, \infty); L^2) \cap C([0, \infty), L^{5/3}) \quad \mathbf{P}\text{-a.s. for all } p \in [1, \infty).$$

Proof. By Theorem 5.4 we constructed a paracontrolled solution $h = v^1 + v^2$ before the stopping time T_L starting from the given initial condition $h(0) = v_0 \in L^2$ \mathbf{P} -a.s. Since $T_L > 0$ \mathbf{P} -a.s., we know that for \mathbf{P} -a.e. ω there exists $q_0(\omega)$ such that $4\sigma_{q_0(\omega)} < T_L(\omega)$. By (5.13) we find that

$$\begin{aligned} \|v^2(T_L)\|_{L^2} &\leq \sum_{0 \leq q < q_0} \|v_{q+1}^2(T_L) - v_q^2(T_L)\|_{L^2} + \sum_{q \geq q_0} \|v_{q+1}^2(T_L) - v_q^2(T_L)\|_{L^2} \\ &\lesssim M_0 q_0 (M_L(1 + q_0))^{1/2} + M_0 (K^{1/2} + 1) + M_0 M_L^{1/2} < \infty. \end{aligned}$$

This implies that $\|v^2(T_L)\|_{L^2} < \infty$ \mathbf{P} -a.s. Since also $v^1(T_L) \in L^2$ \mathbf{P} -a.s., we can use the value $(v^1 + v^2)(T_L)$ as a new initial condition for h in Theorem 5.4.

More precisely, we consider $\hat{h}(0) = (v^1 + v^2)(T_L)$ and define $\hat{z}(t) = z(t + T_L)$, and similarly we define the stochastic objects

$$\begin{aligned} \hat{z}_1(t) &:= z \hat{\mathcal{Y}}(t + T_L), \quad \hat{z}_1 \otimes \hat{z}(t) := z \hat{\mathcal{Y}}(t + T_L), \\ \hat{z} \otimes \hat{z}_1(t) &:= z \hat{\mathcal{Y}}(t + T_L), \\ \hat{z} \hat{\mathcal{V}}(t) &:= z \hat{\mathcal{V}}(t + T_L), \quad \mathcal{I}(\nabla \hat{z})(t) := \mathcal{I}_{T_L, T_L+t}(\nabla z), \\ \mathbb{P}\mathcal{I}(\nabla \hat{z}) \otimes \hat{z}(t) &= \mathbb{P}\mathcal{I}_{T_L, T_L+t}(\nabla z) \otimes z(t + T_L) = z \hat{\mathcal{Y}}(t + T_L; T_L), \\ \mathbb{P}\mathcal{I}(\operatorname{div}(\hat{z} \otimes \hat{z}_1 + \hat{z}_1 \otimes \hat{z})) \otimes \hat{z}(t) &= \mathbb{P}\mathcal{I}_{T_L, T_L+t}(\operatorname{div}(z \hat{\mathcal{Y}} + z \hat{\mathcal{Y}})) \otimes z(t + T_L) \\ &= z \hat{\mathcal{Y}}(t + T_L; T_L). \end{aligned}$$

Then we define stopping time \hat{T}_L similar as in (5.5) with T_L^4 and T_L^5 replaced by

$$\begin{aligned} \hat{T}_L^4 &:= \inf \left\{ t \geq 0, \|z \hat{\mathcal{Y}}(T_L + t; T_L)\|_{C^{-\kappa}} \geq L \right\} \wedge \inf \left\{ t \geq 0, \|z \hat{\mathcal{Y}}\|_{C_{t+T_L}^{1/10} C^{-1/5-\kappa}} \geq L \right\}, \\ \hat{T}_L^5 &:= \inf \left\{ t \geq 0, \|z \hat{\mathcal{Y}}(T_L + t; T_L)\|_{C^{-\kappa}} \geq L \right\}. \end{aligned}$$

Then $\hat{T}_{L+1} \geq T_{L+1} - T_L$.

Consequently, we obtain solutions

$$\left(\hat{h} = \hat{v}^1 + \hat{v}^2, \hat{\vartheta} = \hat{v}^\sharp + \hat{v}^2 + \mathbb{P}[(\hat{v}^1 + \hat{v}^2) \prec \Delta_{\leq R} \mathcal{I}(\nabla \hat{z})] \right)$$

before \hat{T}_{L+1} adapted to \mathcal{F}_{t+T_L} . Here R is chosen as in (6.7) in Sect. 6 but in terms of $L + 1$ instead of L . Moreover, by Proposition 5.2 and (5.7) it holds that $\hat{h}(0) \in B_{5/3, \infty}^{1/5}$. Hence, there is no singularity near zero of \hat{h} and similarly as in Sect. 6 we obtain $\hat{h} \in C_T^{1/10} L^{5/3}$. Then we set $h(t) = (v^1 + v^2)1_{t \leq T_L} + \hat{h}(t - T_L)1_{T_{L+1} \geq t > T_L}$. Then, for $p \geq 1$,

$$h \in C([0, T_{L+1}]; L^{5/3}) \cap L^1(0, T_{L+1}; B_{5/3, \infty}^{1/5}) \cap C_{T_{L+1}, 1/6}^{1/10} L^{5/3} \cap L^p(0, T_{L+1}; L^2).$$

By the same argument as in the proof of [45, Theorem 1.1], h is adapted to $(\mathcal{F}_t)_{t \geq 0}$ and satisfies the equation (4.2) before $[0, T_{L+1}]$. Indeed, we have for $t > T_L$ that

$$h(t) = (v_1 + v_2)(T_L) - \mathbb{P} \int_{T_L}^t e^{(t-s)\Delta} \left[\operatorname{div} \left(h \otimes h + z \hat{\mathcal{Y}} \otimes h + h \otimes z \hat{\mathcal{Y}} + z \hat{\mathcal{V}} \right) \right]$$

$$+ h \otimes z + z \otimes h + z \overset{\vee}{\mathcal{Y}} + z \overset{\vee}{\mathcal{Y}} \Big] ds,$$

with the paracontrolled ansatz

$$h(t) = -\mathbb{P}[h(t) \prec \mathcal{I}_{T_L,t} \nabla z] + \hat{\vartheta}(t - T_L) - \mathcal{I}_{T_L,t}(z \overset{\vee}{\mathcal{Y}} + z \overset{\vee}{\mathcal{Y}}),$$

and for $s > T_L$

$$\begin{aligned} h \otimes z(s) &= -\mathbb{P}[h \prec \mathcal{I}_{T_L,s} \nabla z] \otimes z(s) + \hat{\vartheta}(s - T_L) \otimes z(s) - z \overset{\vee}{\mathcal{Y}}(s; T_L) \\ &= -([\mathbb{P}, h \prec] \mathcal{I}_{T_L,s} \nabla z) \otimes z(s) - \text{com}(h(s), \mathbb{P} \mathcal{I}_{T_L,s} \nabla z, z(s)) - h(s) \cdot z \overset{\vee}{\mathcal{Y}}(s; T_L) \\ &\quad + \hat{\vartheta}(s - T_L) \otimes z(s) - z \overset{\vee}{\mathcal{Y}}(s; T_L). \end{aligned}$$

Now, we define

$$\begin{aligned} \vartheta &= (v^\sharp + v^2 + \mathbb{P}[(v^1 + v^2) \prec \Delta_{\leq R} \mathcal{I}(\nabla z)]) 1_{t \leq T_L} \\ &\quad + \left(\hat{\vartheta}(t - T_L) + \mathbb{P}[h(t) \prec e^{(t-T_L)\Delta} \mathcal{I} \nabla z(T_L)] + e^{(t-T_L)\Delta} (z \overset{\vee}{\mathcal{Y}} + z \overset{\vee}{\mathcal{Y}})(T_L) \right) 1_{t > T_L}. \end{aligned}$$

It is easy to see that $\vartheta \in C([0, T_{L+1}], L^{5/3}) \cap L^1(0, T_{L+1}, B_{1,\infty}^{3/5-\kappa})$. Then, for $t, s > T_L$ it holds that

$$\begin{aligned} h(t) &= -\mathbb{P}[h(t) \prec (\mathcal{I} \nabla z)(t)] + \vartheta(t) - (z \overset{\vee}{\mathcal{Y}} + z \overset{\vee}{\mathcal{Y}})(t), \\ h \otimes z(s) &= -([\mathbb{P}, h(s) \prec] (\mathcal{I} \nabla z)(s)) \otimes z(s) \\ &\quad - \text{com}(h(s), \mathbb{P}(\mathcal{I} \nabla z)(s), z(s)) - h(s) \cdot z \overset{\vee}{\mathcal{Y}}(s) \\ &\quad + \vartheta(s) \otimes z(s) - z \overset{\vee}{\mathcal{Y}}(s). \end{aligned}$$

Here we used that from the renormalization it holds that

$$z \overset{\vee}{\mathcal{Y}}(s) = z \overset{\vee}{\mathcal{Y}}(s, T_L) + e^{(s-T_L)\Delta} (z \overset{\vee}{\mathcal{Y}} + z \overset{\vee}{\mathcal{Y}})(T_L) \otimes z(s)$$

and similarly for other terms.

Thus, (h, ϑ) satisfies the equation (4.2), as well as (4.4) and (4.5) before T_{L+1} . Now, we can iterate the above steps, i.e. starting from $h(T_{L+k})$ and constructing solutions $(h_{k+1}, \vartheta_{k+1})$ before the stopping time T_{L+k+1} . Define $h = h_1 1_{t \leq T_L} + \sum_{k=1}^\infty h_k 1_{\{T_{L+k-1} < t \leq T_{L+k}\}}$, and obtain that $h \in L^p_{\text{loc}}([0, \infty); L^2) \cap C([0, \infty); L^{5/3})$, for all $p \in [1, \infty)$. Similarly, we define ϑ and we obtain that (h, ϑ) is a paracontrolled solution. We emphasize that h does not blow up at any finite time T since for any time T we could find k_0 such $T \leq T_{L+k_0}$ and the infinite sum becomes a finite sum. The desired norm of h only depends on L, k_0 and the initial data. Furthermore, as in the proof of Theorem 5.4 we obtain infinitely many such solutions by choosing different K . □

Corollary 5.6. *Let $v_0 \in L^2 \cup C^{-1+\kappa}$ with $\kappa > 0$ \mathbf{P} -a.s. be a \mathcal{F}_0 -measurable divergence free initial condition. Then there exist infinitely many paracontrolled solutions (h, ϑ) to (4.2) on $[0, \infty)$.*

Proof. If $v_0 \in C^{-1+\kappa}$ with $\kappa > 0$, by [61] there exists a stopping time $0 < \sigma \leq T_L$ and a local paracontrolled solution (h, ϑ) to (4.2) in the sense of Definition 4.3. Now, $h(\sigma) \in C^{1/2-\kappa}$, $\kappa > 0$, and we can start from $h(\sigma)$ and obtain infinitely many global paracontrolled solutions by using Theorem 5.5. Moreover, there is no singularity at $h(\sigma)$ and $h \in C_{T,1/2-\kappa}^{1/10} L^{5/3}$. Note that due to singularity at zero, we only have $h \in L^2(0, T, L^2) \cap C_{T,1/2-\kappa}^{1/10} L^{5/3} \cap C_T H^{-1} \cap L^1(0, T, B_{5/3,\infty}^{1/5})$. \square

Accordingly, Theorem 1.1 is proved.

Finally, by exactly the same argument as in [45, Corollary 1.2], Corollary 1.3 follows.

6. Estimate of v_q^1 and v_q^\sharp

In this section, we work under the assumptions of Proposition 5.1. The main aim is to prove the bounds (5.18), (5.19) as well as for $\kappa > 0$ and $t \in [0, T_L]$

$$\begin{aligned} & \|v_{q+1}^1 - v_q^1\|_{C_t L^2} + \|v_{q+1}^1 - v_q^1\|_{C_t B_{5/3,\infty}^{1/3-2\kappa}} + \|v_{q+1}^1 - v_q^1\|_{C_t^{1/6-\kappa} L^{5/3}} \\ & \lesssim \|v_{q+1}^2 - v_q^2\|_{C_t B_{5/3,\infty}^{1/5}} + \|v_{q+1}^2 - v_q^2\|_{C_t^{1/10} L^{5/3}} + M_L^{1/2} \lambda_q^{-\theta/20}, \end{aligned} \tag{6.1}$$

$$\begin{aligned} & \|v_{q+1}^\sharp - v_q^\sharp\|_{C_{t,3/10} B_{5/3,\infty}^{3/5-\kappa}} + \|v_{q+1}^\sharp - v_q^\sharp\|_{C_t L^2} \\ & \lesssim \|v_{q+1}^2 - v_q^2\|_{C_t B_{5/3,\infty}^{1/5}} + \|v_{q+1}^2 - v_q^2\|_{C_t^{1/10} L^{5/3}} + M_L^{1/2} \lambda_q^{-\theta/20}, \end{aligned} \tag{6.2}$$

and

$$\|v_q^1\|_{C_{t,3/8} L^4} \lesssim M_L^{1/2} \lambda_q^{\theta(7/10+2\kappa)}. \tag{6.3}$$

As a consequence of (5.16), this proves Proposition 5.2. Moreover, we recall that the equation for v_q^1 is linear. Hence, for a given v_q^2 we obtain the existence and uniqueness of solution v_q^1 to (5.3) by a fixed point argument together with the uniform estimate derived in the sequel. Also, if v_q^2 is $(\mathcal{F}_t)_{t \geq 0}$ -adapted, so are (v_q^1, v_q^\sharp) . This in particular gives the existence of v_{q+1}^1 in Proposition 5.1, once the new velocity v_{q+1}^2 was constructed in Sect. 7.

In the following, we make use of the localizers $\Delta_{>R}$ present in the equation for v_q^1 in (5.3). Namely, by an appropriate choice of R we can always apply (2.1) to get a small constant in front of terms which contain v_q^1 . We are therefore able to absorb them into the left hand sides of the estimates without a Gronwall argument. Due to singularity at $t = 0$, we use the weighted in time norms $C_{t,\gamma}$ for several different $\gamma \geq 0$, see Sect. 2 for their definition. In what follows, all the estimates are pathwise and valid before the stopping time T_L .

6.1. Estimate of v_q^1 in $C_{t,1/6} B_{5/3,\infty}^{1/3-2\kappa}$ and $C_{t,1/6}^{1/6-\kappa} L^{5/3}$

We intend to apply Lemma B.2 and notice that by Remark B.3 each application yields a factor L , independently of the chosen time weights in the range $\gamma, \delta \in \{0, 1/6, 3/10\}$. However, we note that the difficult terms for Lemma B.2 are those where we need to decrease the weight, i.e. $\gamma < \delta$. For those we need to make sure that the condition

$$\gamma - \delta - \alpha/2 + \beta/2 + 1 > 0 \tag{6.4}$$

from Lemma B.2 is satisfied. It will be seen below that this is always achieved since these terms do not require such a gain in space regularity, i.e. the difference $\alpha - \beta$ compensates the negativity of the difference $\gamma - \delta$.

Hence, we shall bound each term appearing in V_q^1 as well as z^{\heartsuit} in appropriate (possibly time-weighted) function spaces with spatial regularity at least $B_{5/3,\infty}^{-2/3-\kappa}$. The terms in $V_q^{1,*}$ are estimated the same way.

Recall that by the definition of the stopping time (5.5) we have

$$\|z\|_{C_t C^{-1/2-\kappa}}^2 + \|z^{\heartsuit}\|_{C_t C^{-\kappa}} + \|z^{\spadesuit}\|_{C_t C^{-1/2-\kappa}} + \|z^{\clubsuit}\|_{C_t C^{-\kappa}} + \|z^{\diamondsuit}\|_{C_t C^{-\kappa}} \lesssim L.$$

By the paraproduct estimate Lemma 2.2 we have

$$\begin{aligned} & \| (v_q^1 + v_q^2) \otimes \Delta_{\leq f(q)} \Delta_{> RZ} \|_{C_{t,1/6} B_{5/3,\infty}^{-2/3-\kappa}} \\ & \lesssim \sup_{s \in [0,t]} s^{1/6} (\|v_q^1(s)\|_{L^{5/3}} + \|v_q^2(s)\|_{L^{5/3}}) \|\Delta_{> RZ}\|_{C_t C^{-2/3-\kappa}} \\ & \lesssim (\|v_q^1\|_{C_{t,1/6} L^{5/3}} + \|v_q^2\|_{C_{t,1/6} L^{5/3}}) L 2^{-R/6}, \\ & \| (v_q^1 + v_q^2) \otimes \Delta_{\leq f(q)} \Delta_{> RZ} \heartsuit \|_{C_{t,1/6} B_{5/3,\infty}^{-1/6-\kappa}} + \| (v_q^1 + v_q^2) \\ & < \Delta_{\leq f(q)} \Delta_{> RZ} \spadesuit \|_{C_{t,1/6} B_{5/3,\infty}^{-1/6-\kappa}} \\ & \lesssim \sup_{s \in [0,t]} s^{1/6} (\|v_q^1(s)\|_{L^{5/3}} + \|v_q^2(s)\|_{L^{5/3}}) (\|\Delta_{> RZ} \heartsuit\|_{C_t C^{-1/6-\kappa}} \\ & + \|\Delta_{> RZ} \spadesuit\|_{C_t C^{-1/6-\kappa}}) \\ & \lesssim (\|v_q^1\|_{C_{t,1/6} L^{5/3}} + \|v_q^2\|_{C_{t,1/6} L^{5/3}}) L 2^{-R/6}, \end{aligned}$$

and

$$\begin{aligned} & \|v_q^1 \otimes \Delta_{> RZ} \heartsuit\|_{C_{t,1/6} B_{5/3,\infty}^{\kappa}} + \|v_q^1 \heartsuit \otimes \Delta_{> RZ} \spadesuit\|_{C_{t,1/6} B_{5/3,\infty}^{\kappa}} \\ & + \|v_q^1 \otimes \Delta_{\leq f(q)} \Delta_{> RZ}\|_{C_{t,1/6} B_{5/3,\infty}^{-1/3-3\kappa}} \\ & \lesssim \sup_{s \in [0,t]} s^{1/6} \|v_q^1\|_{B_{5/3,\infty}^{1/3-2\kappa}} (\|\Delta_{> RZ} \heartsuit\|_{C_t C^{-1/6-\kappa}} \\ & + \|\Delta_{> RZ} \spadesuit\|_{C_t C^{-1/6-\kappa}} + \|\Delta_{> RZ}\|_{C_t C^{-2/3-\kappa}}) \\ & \lesssim \|v_q^1\|_{C_{t,1/6} B_{5/3,\infty}^{1/3-2\kappa}} L 2^{-R/6}. \end{aligned}$$

For the two commutators we use the commutator estimates, Lemmas 2.4 and 2.3, to obtain

$$\begin{aligned} & \|([\mathbb{P}, v_q^1 \prec] \mathcal{I}(\nabla z)) \odot \Delta_{>Rz}\|_{C_{t,1/6} B_{5/3,\infty}^\kappa} + \|\text{com}(v_q^1, \mathbb{P}\mathcal{I}(\nabla z), \Delta_{>Rz})\|_{C_{t,1/6} B_{5/3,\infty}^\kappa} \\ & \lesssim \|v_q^1\|_{C_{t,1/6} B_{5/3,\infty}^{1/3-2\kappa}} \|\Delta_{>Rz}\|_{C_t C^{-2/3-\kappa}} \|z\|_{C_t C^{-1/2-\kappa}} \lesssim \|v_q^1\|_{C_{t,1/6} B_{5/3,\infty}^{1/3-2\kappa}} L 2^{-R/6}. \end{aligned}$$

For the last term containing v_q^\sharp we use the paraproduct estimate and Lemma B.2. In particular, since v_q^\sharp requires a higher time weight $t^{3/10}$, we shall verify the condition (6.4). It turns out that this is satisfied as $\gamma = 1/6, \delta = 3/10, \alpha = 1/3 - 2\kappa, \beta = -1 + 1/20 - 2\kappa$ hence (6.4) holds. Accordingly, we obtain

$$\begin{aligned} & \|\mathcal{I} \text{div}(v_q^\sharp \odot \Delta_{>Rz})\|_{C_{t,1/6} B_{5/3,\infty}^{1/3-2\kappa}} + \|\mathcal{I} \text{div}(v_q^\sharp \odot \Delta_{>Rz})\|_{C_{t,1/6}^{1/6-\kappa} L^{5/3}} \\ & \lesssim L \|\text{div}(v_q^\sharp \odot \Delta_{>Rz})\|_{C_{t,3/10} B_{5/3,\infty}^{-1+1/20-2\kappa}} \\ & \lesssim L \|v_q^\sharp \odot \Delta_{>Rz}\|_{C_{t,3/10} B_{5/3,\infty}^{1/20-2\kappa}} \\ & \lesssim L \|v_q^\sharp\|_{C_{t,3/10} B_{5/3,\infty}^{3/5-\kappa}} \|\Delta_{>Rz}\|_{C_t C^{-11/20-\kappa}} \\ & \lesssim L^2 \|v_q^\sharp\|_{C_{t,3/10} B_{5/3,\infty}^{3/5-\kappa}} 2^{-R/20}. \end{aligned}$$

For the initial value part we have, by Lemma B.1 and (5.6), for $t \in (0, T_L]$, that

$$\begin{aligned} & \|e^{t\Delta} v_0\|_{B_{5/3,\infty}^{1/3-2\kappa}} \lesssim L^{1/2} t^{-1/6+\kappa} N, \\ & \|(e^{t\Delta} - e^{s\Delta})v_0\|_{L^{5/3}} \lesssim |t - s|^{1/6-\kappa} s^{-1/6+\kappa} N, \quad |t - s| \leq 1, \quad 0 < s < t. \end{aligned}$$

Summarizing all the above estimates and using Besov embedding Lemma 2.1, we obtain

$$\begin{aligned} & \|v_q^1\|_{C_{t,1/6} B_{5/3,\infty}^{1/3-2\kappa}} + \|v_q^1\|_{C_{t,1/6}^{1/6-\kappa} L^{5/3}} \\ & \lesssim L \left(L + N + L 2^{-R/20} (\|v_q^1\|_{C_{t,1/6} B_{5/3,\infty}^{1/3-2\kappa}} + \|v_q^\sharp\|_{C_{t,3/10} B_{5/3,\infty}^{3/5-\kappa}}) \right) \quad (6.5) \\ & + L 2^{-R/6} \|v_q^2\|_{C_{t,1/6} L^{5/3}}. \end{aligned}$$

Here, we used (5.6) and, as mentioned above, the extra factor L comes from Lemma B.2.

6.2. Estimate of v_q^\sharp in $C_{t,3/10} B_{5/3,\infty}^{3/5-\kappa}$ and $C_{t,3/10}^{1/20} B_{5/3,\infty}^{11/20-2\kappa}$

Let us proceed with the estimate of v_q^\sharp . Here, there are no difficulties coming from changing the time weight as all the terms require either a lower or the same weight. In view of (5.3), the paracontrolled ansatz (5.4), and since

$$\begin{aligned} & \text{div} \left((v_q^1 + v_q^2) \otimes \Delta_{\leq f(q)} \Delta_{>Rz} \right) = \Delta_{\leq f(q)} \Delta_{>Rz} \succ \nabla (v_q^1 + v_q^2), \\ & \text{div} \left(\Delta_{\leq f(q)} \Delta_{>Rz} \otimes (v_q^1 + v_q^2) \right) = (v_q^1 + v_q^2) \prec \nabla \Delta_{\leq f(q)} \Delta_{>Rz}, \end{aligned}$$

we obtain

$$\begin{aligned}
 v_q^\sharp &= v_q^1 + \mathbb{P}[(v_q^1 + v_q^2) \prec \mathcal{I}(\nabla \Delta_{>R} \Delta_{\leq f(q)} z)] + (z^{\check{Y}} + z^{\check{Y}^*}) \\
 &= v_q^1(0) - \mathcal{I} \mathbb{P} \operatorname{div} (z^{\check{Y}} + V_q^\sharp + V_q^{\sharp,*}) \\
 &\quad - \mathbb{P}[\mathcal{I}, (v_q^1 + v_q^2) \prec](\nabla \Delta_{>R} \Delta_{\leq f(q)} z) \\
 &\quad - \mathcal{I} \mathbb{P}[\nabla(v_q^1 + v_q^2) \prec \Delta_{\leq f(q)} \Delta_{>R} z], \tag{6.6}
 \end{aligned}$$

where

$$\begin{aligned}
 V_q^\sharp &= (v_q^1 + v_q^2) \otimes \Delta_{\leq f(q)} \Delta_{>R} z^{\check{Y}} + v_q^1 \otimes \Delta_{>R} z^{\check{Y}} + v_q^1 \otimes \Delta_{\leq f(q)} \Delta_{>R} z \\
 &\quad - ([\mathbb{P}, v_q^1 \prec] \mathcal{I} \nabla z) \otimes \Delta_{>R} z - \operatorname{com}(v_q^1, \mathbb{P} \mathcal{I} \nabla z, \Delta_{>R} z) - z^{\check{Z}} \\
 &\quad - (v_q^1 + v_q^2) \prec \Delta_{\leq f(q)} \Delta_{>R} z^{\check{Y}^*} - v_q^1 \succ \Delta_{>R} z^{\check{Y}^*} + v_q^\sharp \otimes \Delta_{>R} z.
 \end{aligned}$$

In what follows we estimate each term on the right hand side of (6.6).

From the above estimate we already know that $z^{\check{Y}}$ as well as all the terms in V_q^\sharp except for $v_q^\sharp \otimes \Delta_{>R} z$ are bounded in $B_{5/3, \infty}^{-1/3-3\kappa}$. We also show by paraproduct estimates from Lemma 2.2 that

$$\begin{aligned}
 \|v_q^\sharp \otimes \Delta_{>R} z\|_{C_{t, 3/10} B_{5/3, \infty}^{1/20-2\kappa}} &\lesssim \|v_q^\sharp\|_{C_{t, 3/10} B_{5/3, \infty}^{3/5-\kappa}} \|\Delta_{>R} z\|_{C_t C^{-11/20-\kappa}} \\
 &\lesssim L^{1/2} \|v_q^\sharp\|_{C_{t, 3/10} B_{5/3, \infty}^{3/5-\kappa}} 2^{-R/20}.
 \end{aligned}$$

Moreover, Lemma 2.2 also implies

$$\begin{aligned}
 &\|\mathbb{P}[\nabla(v_q^1 + v_q^2) \prec (\Delta_{>R} \Delta_{\leq f(q)} z)]\|_{C_{t, 1/6} B_{5/3, \infty}^{-27/20-\kappa}} \\
 &\lesssim \left(\sup_{s \in [0, t]} s^{1/6} \|v_q^1(s)\|_{B_{5/3, \infty}^{1/3-2\kappa}} + L^{1/6} \|v_q^2\|_{C_t B_{5/3, \infty}^{1/5}} \right) \|\Delta_{>R} z\|_{C_t C^{-11/20-\kappa}} \\
 &\lesssim (\|v_q^1\|_{C_{t, 1/6} B_{5/3, \infty}^{1/3-2\kappa}} L^{1/2} + L \|v_q^2\|_{C_t B_{5/3, \infty}^{1/5}}) 2^{-R/20},
 \end{aligned}$$

which can then be plugged in the Schauder estimate, Lemma B.2. Next, we note that Lemma B.4 can be applied to the remaining term in (6.6) which also gives a factor of L . Hence, we use interpolation to get

$$\begin{aligned}
 &\|\mathbb{P}[\mathcal{I}, (v_q^1 + v_q^2) \prec](\nabla \Delta_{>R} \Delta_{\leq f(q)} z)\|_{C_{t, 3/10} B_{5/3, \infty}^{3/5}} \\
 &\quad + \|\mathbb{P}[\mathcal{I}, (v_q^1 + v_q^2) \prec](\nabla \Delta_{>R} \Delta_{\leq f(q)} z)\|_{C_{t, 3/10}^{1/20} B_{5/3, \infty}^{11/20-2\kappa}} \\
 &\lesssim L (\|v_q^1\|_{C_{t, 1/6} B_{5/3, \infty}^{1/3-2\kappa}} + \|v_q^2\|_{C_t B_{5/3, \infty}^{1/5}} + \|v_q^1\|_{C_{t, 1/6}^{1/6-\kappa}} L^{5/3}) \\
 &\quad + \|v_q^2\|_{C_t^{1/10} L^{5/3}} \|\Delta_{>R} z\|_{C_t C^{-11/20-\kappa}} \\
 &\lesssim L^2 2^{-R/20} (\|v_q^1\|_{C_{t, 1/6} B_{5/3, \infty}^{1/3-2\kappa}} + \|v_q^2\|_{C_t B_{5/3, \infty}^{1/5}} \\
 &\quad + \|v_q^1\|_{C_{t, 1/6}^{1/6-\kappa}} L^{5/3} + \|v_q^2\|_{C_t^{1/10} L^{5/3}}).
 \end{aligned}$$

Combining the above estimates and using Schauder estimate Lemma B.2 and interpolation, we have

$$\begin{aligned} & \|v_q^\sharp\|_{C_{t,3/10}B_{5/3,\infty}^{3/5-\kappa}} + \|v_q^\sharp\|_{C_{t,3/10}^{1/20}B_{5/3,\infty}^{11/20-2\kappa}} \\ & \lesssim L^2 + LN + L^2 2^{-R/20} \left(\|v_q^1\|_{C_{t,1/6}B_{5/3,\infty}^{1/3-2\kappa}} + \|v_q^2\|_{C_t B_{5/3,\infty}^{1/5}} \right. \\ & \quad \left. + \|v_q^1\|_{C_{t,1/6}^{1/6-\kappa}L^{5/3}} + \|v_q^2\|_{C_t^{1/10}L^{5/3}} + \|v_q^\sharp\|_{C_{t,3/10}B_{5/3,\infty}^{3/5-\kappa}} \right) \\ & \quad + L^2 2^{-R/6} \|v_q^2\|_{C_{t,1/6}L^{5/3}}, \end{aligned}$$

which combined with (6.5) implies that

$$\begin{aligned} & \|v_q^1\|_{C_{t,1/6}B_{5/3,\infty}^{1/3-2\kappa}} + \|v_q^1\|_{C_{t,1/6}^{1/6-\kappa}L^{5/3}} + \|v_q^\sharp\|_{C_{t,3/10}B_{5/3,\infty}^{3/5-\kappa}} + \|v_q^\sharp\|_{C_{t,3/10}^{1/20}B_{5/3,\infty}^{11/20-2\kappa}} \\ & \lesssim L^2 + LN + L^2 2^{-R/20} \left(\|v_q^1\|_{C_{t,1/6}B_{5/3,\infty}^{1/3-2\kappa}} + \|v_q^2\|_{C_t B_{5/3,\infty}^{1/5}} \right. \\ & \quad \left. + \|v_q^1\|_{C_{t,1/6}^{1/6-\kappa}L^{5/3}} + \|v_q^2\|_{C_t^{1/10}L^{5/3}} + \|v_q^\sharp\|_{C_{t,3/10}B_{5/3,\infty}^{3/5-\kappa}} \right) \\ & \quad + L^2 2^{-R/6} \|v_q^2\|_{C_{t,1/6}L^{5/3}}. \end{aligned}$$

Then we choose R such that

$$2^{R/20} = 4CL^2, \tag{6.7}$$

with C being the implicit constant and use (5.7) to obtain

$$\begin{aligned} & \|v_q^1\|_{C_{t,1/6}B_{5/3,\infty}^{1/3-2\kappa}} + \|v_q^1\|_{C_{t,1/6}^{1/6-\kappa}L^{5/3}} + \|v_q^\sharp(s)\|_{C_{t,3/10}B_{5/3,\infty}^{3/5-\kappa}} + \|v_q^\sharp\|_{C_{t,3/10}^{1/20}B_{5/3,\infty}^{11/20-2\kappa}} \\ & \lesssim L^2 + LN + L^2 a^{-\alpha/2} M_L^{1/2}, \end{aligned} \tag{6.8}$$

which implies the first part of (5.18) and (5.19).

6.3. Estimate of v_q^1 in $C_t L^2$

Here, most of the terms are similar as in (6.5) but we need to be careful about the compatibility condition (6.4). As mentioned above, terms except for $v_q^\sharp \otimes \Delta_{>R} z$ and $(v_q^1 + v_q^2) \otimes \Delta_{\leq f(q)} \Delta_{>R} z$ have spatial regularity

$$B_{5/3,\infty}^{-1/3-3\kappa} \subset B_{2,\infty}^{-1/3-3/10-3\kappa}$$

by Lemma 2.1. Thus, the corresponding $C_t L^2$ norm can be bounded by Lemma B.2 with $\gamma = 0, \delta = 1/6, \alpha = \kappa, \beta = -1/3 - 3/10 - 3\kappa - 1$ as follows:

$$\begin{aligned} \|\mathcal{I} \operatorname{div}(\cdots)\|_{C_t L^2} & \lesssim \|\mathcal{I} \operatorname{div}(\cdots)\|_{C_t B_{2,\infty}^\kappa} \\ & \lesssim L \|\operatorname{div}(\cdots)\|_{C_{t,1/6} B_{2,\infty}^{-1/3-3/10-3\kappa-1}} \\ & \lesssim L^2 + L^2 \|v_q^1\|_{C_{t,1/6} B_{5/3,\infty}^{1/3-2\kappa}} + L^2 \|v_q^2\|_{C_t L^{5/3}}. \end{aligned}$$

Here, in \dots we collected all the terms from $V_q^1 + V_q^{1,*}$ except for the above mentioned two and $z^{\check{V}}$ and their symmetric counterparts.

Next, using paraproduct estimates and Lemma 2.1, the term corresponding to $v_q^\sharp \otimes \Delta_{>Rz}$ can be bounded in view of the embedding $B_{5/3,\infty}^{3/5-\kappa-1/2-\kappa} \subset B_{2,\infty}^{3/5-1/2-2\kappa-3/10} = B_{2,\infty}^{-1/5-2\kappa}$ by Lemma B.2 with $\gamma = 0, \delta = 3/10, \alpha = \kappa, \beta = -1/5 - 2\kappa - 1$ as

$$\begin{aligned} \|\mathcal{I} \operatorname{div}(v_q^\sharp \otimes \Delta_{>Rz})\|_{C_t L^2} &\lesssim \|\mathcal{I} \operatorname{div}(v_q^\sharp \otimes \Delta_{>Rz})\|_{C_t B_{2,\infty}^\kappa} \\ &\lesssim L \|\operatorname{div}(v_q^\sharp \otimes \Delta_{>Rz})\|_{C_{t,3/10} B_{2,\infty}^{-1/5-2\kappa-1}} \\ &\lesssim L \|v_q^\sharp \otimes \Delta_{>Rz}\|_{C_{t,3/10} B_{5/3,\infty}^{3/5-1/2-2\kappa}} \\ &\lesssim L \|v_q^\sharp\|_{C_{t,3/10} B_{5/3,\infty}^{3/5-\kappa}} \|z\|_{C_t C^{-1/2-\kappa}} \\ &\lesssim L^2 \|v_q^\sharp\|_{C_{t,3/10} B_{5/3,\infty}^{3/5-\kappa}}. \end{aligned}$$

We also use Lemma 2.2 and embedding $B_{5/3,\infty}^{-1+3/10+\kappa} \subset H^{-1}$ to get

$$\begin{aligned} \|(v_q^1 + v_q^2) \otimes \Delta_{\leq f(q)} \Delta_{>Rz}\|_{C_t H^{-1}} &\lesssim \|v_q^1\|_{C_t L^2} \|\Delta_{>Rz}\|_{C_t C^{-2/3-\kappa}} + \|v_q^2\|_{C_t L^{5/3}} \|z\|_{C_t C^{-1/2-\kappa}} \\ &\lesssim L^{1/2} (\|v_q^1\|_{C_t L^2} 2^{-R/6} + \|v_q^2\|_{C_t L^{5/3}}). \end{aligned}$$

Thus combining the above estimates and (5.7), (6.8) we obtain

$$\begin{aligned} &\|v_q^1\|_{C_t L^2} \\ &\lesssim L^2 + LN + L^2 (\|v_q^1\|_{C_{t,1/6} B_{5/3,\infty}^{1/3-2\kappa}} + \|v_q^2\|_{C_t L^{5/3}} + \|v_q^\sharp\|_{C_{t,3/10} B_{5/3,\infty}^{3/5-\kappa}}) \\ &\quad + L^2 \|v_q^1\|_{C_t L^2} 2^{-R/6} \\ &\lesssim L^4 + L^3 N + a^{-\alpha/2} L^4 M_L^{1/2} + L^2 a^{-\alpha/2} M_L^{1/2} + L^2 \|v_q^1\|_{C_t L^2} 2^{-R/6} \\ &\lesssim L^4 + L^3 N + a^{-\alpha/4} M_L^{1/2} + L^2 \|v_q^1\|_{C_t L^2} 2^{-R/6}, \end{aligned}$$

using the fact that $L^4 \leq a^{\alpha/4}$. Hence the last part of (5.18) follows.

6.4. Estimate of v_q^\sharp in $C_t L^2$

We apply the paracontrolled ansatz (5.4), the Besov embedding Lemma 2.1, and paraproduct estimates Lemma 2.2 as well as (5.7) and (5.5) together with the Schauder estimate to control $z^{\check{V}}$ and $z^{\check{V}}$. We deduce

$$\begin{aligned} \|v_q^\sharp\|_{C_t L^2} &\lesssim L \|v_q^1\|_{C_t L^2} + \|v_q^2\|_{\mathcal{I}(\nabla \Delta_{>R} \Delta_{\leq f(q)} z)}\|_{C_t B_{5/3,\infty}^{3/10+\kappa}} + L^2 \\ &\lesssim L (\|v_q^2\|_{C_t L^{5/3}} + \|v_q^1\|_{C_t L^2}) + L^2 \\ &\lesssim L a^{-\alpha/2} M_L^{1/2} + L (L^4 + L^3 N + a^{-\alpha/4} M_L^{1/2}) + L^2 \\ &\lesssim M_L^{1/2} a^{-\alpha/8} + L^5 N + L^6. \end{aligned}$$

The last part of (5.19) follows.

6.5. Estimate of the Difference $v_{q+1}^1 - v_q^1$ in $C_t B_{5/3,\infty}^{1/3-2\kappa}$ and $C_t^{1/6-\kappa} L^{5/3}$

Most of the terms can be estimated similarly as in (6.5). We do not have to consider the initial data as it vanishes and therefore we can even control directly the C_t -norms without any weight. The main change comes from the additional difference $\Delta_{\leq f(q+1)} - \Delta_{\leq f(q)}$. First, we use Lemma 2.2 to bound the terms with paraproducts $\otimes, <$ containing $\Delta_{\leq f(q+1)} - \Delta_{\leq f(q)}$ by Schauder estimates as

$$L \|v_q^1 + v_q^2\|_{C_t L^{5/3}} \|(\Delta_{\leq f(q+1)} - \Delta_{\leq f(q)}) \Delta_{>R}(z + z^\vee)\|_{C_t C^{-2/3-\kappa}} \lesssim M_L^{1/2} \lambda_q^{-\theta/6},$$

$$L \|v_q^1 + v_q^2\|_{C_t L^{5/3}} \|(\Delta_{\leq f(q+1)} - \Delta_{\leq f(q)}) \Delta_{>R} z^{\flat\delta'}\|_{C_t C^{-2/3-\kappa}} \lesssim M_L^{1/2} \lambda_q^{-\theta/6}.$$

The part of \otimes containing $\Delta_{\leq f(q+1)} - \Delta_{\leq f(q)}$ can be bounded by Schauder estimates Lemma B.2 with $\gamma = 0, \delta = 1/6, \alpha = 1/3 - 2\kappa, \beta = 1/3 - 2\kappa - 11/20 - \kappa - 1$ as

$$L \|v_q^1\|_{C_{t,1/6} B_{5/3,\infty}^{1/3-2\kappa}} \|(\Delta_{\leq f(q+1)} - \Delta_{\leq f(q)}) z\|_{C_t C^{-11/20-\kappa}} \lesssim M_L^{1/2} \lambda_q^{-\theta/20}.$$

Thus, in view of (6.5) we obtain

$$\begin{aligned} & \|v_{q+1}^1 - v_q^1\|_{C_t B_{5/3,\infty}^{1/3-2\kappa}} + \|v_{q+1}^1 - v_q^1\|_{C_t^{1/6-\kappa} L^{5/3}} \\ & \lesssim L^2 2^{-R/20} \left(\|v_{q+1}^1 - v_q^1\|_{C_t B_{5/3,\infty}^{1/3-2\kappa}} \right. \\ & \quad \left. + \|v_{q+1}^2 - v_q^2\|_{C_t L^{5/3}} + \|v_{q+1}^\sharp - v_q^\sharp\|_{C_{t,3/10} B_{5/3,\infty}^{3/5-\kappa}} \right) \\ & \quad + M_L^{1/2} \lambda_q^{-\theta/20}, \end{aligned} \tag{6.9}$$

where for v_q^\sharp part we used $\gamma = 0, \delta = 3/10, \alpha = 1/3 - 2\kappa, \beta = 3/5 - \kappa - 11/20 - 1$. Therefore, in order to deduce the first part of (6.1), we shall estimate the difference $v_{q+1}^\sharp - v_q^\sharp$.

6.6. Estimate of the Difference $v_{q+1}^\sharp - v_q^\sharp$ in $C_{t,3/10} B_{5/3,\infty}^{3/5-\kappa}$

Terms in V_q^\sharp can be bounded similarly as above. We only concentrate on

$$\mathbb{P}[\mathcal{I}, (v_q^1 + v_q^2) <](\nabla \Delta_{>R} \Delta_{\leq f(q)} z), \quad \mathcal{I}\mathbb{P}[\nabla(v_q^1 + v_q^2) < \Delta_{\leq f(q)} \Delta_{>R} z].$$

Similarly as before, most terms could be estimated as the estimates for v_q^\sharp with v_q replaced by $v_{q+1} - v_q$. We consider the terms containing $\Delta_{\leq f(q+1)} - \Delta_{\leq f(q)}$ and use

$$\|(\Delta_{\leq f(q+1)} - \Delta_{\leq f(q)}) z\|_{C_t C^{-11/20-\kappa}} \lesssim \lambda_q^{-\theta/20} L^{1/2}.$$

Thus, using (5.18) the $C_{t,3/10} B_{5/3,\infty}^{3/5-\kappa}$ -norm of these terms can be bounded by

$$\lambda_q^{-\theta/20} L^2 (\|v_q^1\|_{C_{t,1/6} B_{5/3,\infty}^{1/3-2\kappa}} + \|v_q^2\|_{C_t B_{5/3,\infty}^{1/5}} + \|v_q^1\|_{C_{t,1/6}^{1/6-\kappa} L^{5/3}} + \|v_q^2\|_{C_t^{1/10} L^{5/3}})$$

$$\lesssim \lambda_q^{-\theta/20} M_L^{1/2}.$$

Here we used Lemma B.4. Hence, we obtain

$$\begin{aligned} & \|v_{q+1}^\sharp - v_q^\sharp\|_{C_{t,3/10} B_{5/3,\infty}^{3/5-\kappa}} \\ & \lesssim \lambda_q^{-\theta/20} M_L^{1/2} + L^2 \left(\|v_{q+1}^1 - v_q^1\|_{C_t B_{5/3,\infty}^{1/3-2\kappa}} + \|v_{q+1}^2 - v_q^2\|_{C_t B_{5/3,\infty}^{1/5}} \right. \\ & \quad + \|v_{q+1}^1 - v_q^1\|_{C_t^{1/6-\kappa} L^{5/3}} + \|v_{q+1}^2 - v_q^2\|_{C_t^{1/10} L^{5/3}} \\ & \quad \left. + \|v_{q+1}^\sharp - v_q^\sharp\|_{C_{t,3/10} B_{5/3,\infty}^{3/5-\kappa}} \right) 2^{-R/20}. \end{aligned} \tag{6.10}$$

Combining this bound with (6.9) we deduce a first part of (6.1) and (6.2) and it remains to estimate the differences $v_{q+1}^1 - v_q^1$ and $v_{q+1}^\sharp - v_q^\sharp$ in $C_t L^2$.

6.7. Estimate of the Difference $v_{q+1}^1 - v_q^1$ and $v_{q+1}^\sharp - v_q^\sharp$ in $C_t L^2$

By the Besov embedding Lemma 2.1 it holds that

$$\|v_{q+1}^1 - v_q^1\|_{C_t L^2} \lesssim \|v_{q+1}^1 - v_q^1\|_{C_t B_{5/3,\infty}^{1/3-2\kappa}},$$

which in view of the first part of (6.1) implies the second part of (6.1). Moreover, by (5.4), the Besov embedding Lemma 2.1, and paraproduct estimates Lemma 2.2 we have

$$\begin{aligned} & \|(v_q^1 + v_q^2) \prec \mathcal{I}(\nabla \Delta_{>R}(\Delta_{\leq f(q)} - \Delta_{\leq f(q+1)})z)\|_{C_t B_{5/3,\infty}^{1/3-2\kappa}} \\ & \lesssim L(\|v_q^1\|_{C_t L^{5/3}} + \|v_q^2\|_{C_t L^{5/3}}) \|(\Delta_{\leq f(q)} - \Delta_{\leq f(q+1)})z\|_{C_t C^{-2/3-\kappa}} \\ & \lesssim L^{3/2}(\|v_q^1\|_{C_t L^2} + \|v_q^2\|_{C_t L^{5/3}}) \lambda_q^{-\theta/6}, \end{aligned}$$

which, due to $B_{5/3,\infty}^{1/3-2\kappa} \subset L^2$ and (6.1), implies

$$\begin{aligned} & \|v_{q+1}^\sharp - v_q^\sharp\|_{C_t L^2} \\ & \lesssim \|v_{q+1}^1 - v_q^1\|_{C_t L^2} + \|v_{q+1}^2 - v_q^2\|_{C_t L^{5/3}} + L^{3/2}(\|v_q^1\|_{C_t L^2} + \|v_q^2\|_{C_t L^{5/3}}) \lambda_q^{-\theta/6} \\ & \lesssim \|v_{q+1}^2 - v_q^2\|_{C_t B_{5/3,\infty}^{1/5}} + \|v_{q+1}^2 - v_q^2\|_{C_t^{1/10} L^{5/3}} + M_L^{1/2} \lambda_q^{-\theta/20}. \end{aligned}$$

This gives the remaining estimate of (6.2).

6.8. Estimate of v_q^1 in $C_{t,3/8} L^4$

This norm may a priori blow up during the iteration. We also estimate each term separately and apply the Schauder estimate Lemma B.2 which then gives an additional factor L . We have

$$\|\Delta_{>R} \Delta_{\leq f(q)}(z + z^{\mathbf{Y}})\|_{C_t C^{1/5+\kappa}} + \|\Delta_{>R} \Delta_{\leq f(q)} z^{\mathbf{Y}}\|_{C_t C^{1/5+\kappa}} \lesssim L \lambda_q^{(7/10+2\kappa)\theta}.$$

Hence, by Lemma 2.2 and (5.18), we have

$$\begin{aligned} & \| (v_q^1 + v_q^2) \otimes \Delta_{>R} \Delta_{\leq f(q)} (z + z^{\vee}) \|_{C_t B_{4,\infty}^{-1+\kappa}} \\ & + \| (v_q^1 + v_q^2) \prec \Delta_{>R} \Delta_{\leq f(q)} z^{\flat} \|_{C_t B_{4,\infty}^{-1+\kappa}} \\ & \lesssim \| v_q^1 + v_q^2 \|_{C_t B_{4,\infty}^{-6/5}} \left(\| \Delta_{>R} \Delta_{\leq f(q)} (z + z^{\vee}) \|_{C_t C^{1/5+\kappa}} \right. \\ & \left. + \| \Delta_{>R} \Delta_{\leq f(q)} z^{\flat} \|_{C_t C^{1/5+\kappa}} \right) \\ & \lesssim L \lambda_q^{(7/10+2\kappa)\theta} \| v_q^1 + v_q^2 \|_{C_t L^{5/3}}, \\ & \| v_q^1 \otimes \Delta_{>R} \Delta_{\leq f(q)} z \|_{C_{t,1/6} B_{4,\infty}^{-1+\kappa}} \\ & \lesssim \| v_q^1 \|_{C_{t,1/6} B_{4,\infty}^{-1+\kappa}} \| \Delta_{>R} \Delta_{\leq f(q)} z \|_{C_t L^\infty} \\ & \lesssim L^{1/2} \lambda_q^{(1/2+2\kappa)\theta} \| v_q^1 \|_{C_{t,1/6} B_{5/3,\infty}^{1/3-2\kappa}}, \end{aligned}$$

and

$$\begin{aligned} & \| v_q^1 \otimes \Delta_{>R} z^{\vee} \|_{C_{t,1/6} B_{4,\infty}^{-1+\kappa}} + \| v_q^1 \succ \Delta_{>R} z^{\flat} \|_{C_{t,1/6} B_{4,\infty}^{-1+\kappa}} \\ & \lesssim \| v_q^1 \otimes \Delta_{>R} z^{\vee} \|_{C_{t,1/6} B_{5/3,\infty}^{1/3-3\kappa}} + \| v_q^1 \succ \Delta_{>R} z^{\flat} \|_{C_{t,1/6} B_{5/3,\infty}^{1/3-3\kappa}} \\ & \lesssim \| v_q^1 \|_{C_{t,1/6} B_{5/3,\infty}^{1/3-2\kappa}} (\| z^{\vee} \|_{C_t C^{-\kappa}} + \| z^{\flat} \|_{C_t C^{-\kappa}}) \\ & \lesssim \| v_q^1 \|_{C_{t,1/6} B_{5/3,\infty}^{1/3-2\kappa}} L. \end{aligned}$$

Using Lemma B.2 and (5.19) we have

$$\begin{aligned} & \| \mathcal{I} \operatorname{div} (v_q^{\sharp} \otimes \Delta_{>R} z) \|_{C_{t,3/10} L^4} \lesssim \| \mathcal{I} \operatorname{div} (v_q^{\sharp} \otimes \Delta_{>R} z) \|_{C_{t,3/10} B_{4,\infty}^{\kappa}} \\ & \lesssim L \| v_q^{\sharp} \otimes \Delta_{>R} z \|_{C_{t,3/10} B_{4,\infty}^{-1+2\kappa}} \\ & \lesssim L \| v_q^{\sharp} \|_{C_{t,3/10} B_{5/3,\infty}^{3/5-\kappa}} \| \Delta_{>R} z \|_{C_t C^{-1/2-\kappa}}. \end{aligned}$$

By that commutator estimates of Lemma 2.3 we have

$$\begin{aligned} & \| ([\mathbb{P}, v_q^1 \prec] \mathcal{I}(\nabla z)) \otimes \Delta_{>R} z \|_{C_{t,1/6} B_{5/3,\infty}^{1/3-4\kappa}} + \| \operatorname{com} (v_q^1, \mathbb{P} \mathcal{I}(\nabla z), \Delta_{>R} z) \|_{C_{t,1/6} B_{5/3,\infty}^{1/3-4\kappa}} \\ & \lesssim \| v_q^1 \|_{C_{t,1/6} B_{5/3,\infty}^{1/3-2\kappa}} \| z \|_{C_t C^{-1/2-\kappa}}^2 \lesssim \| v_q^1 \|_{C_{t,1/6} B_{5/3,\infty}^{1/3-2\kappa}} L. \end{aligned}$$

Finally, for the initial value part we apply Lemma 9 in [24] to obtain

$$\| e^{t\Delta} v_0 \|_{L^4} \lesssim t^{-3/8} \| v_0 \|_{L^2}.$$

Combining the above estimates and applying the Schauder estimate Lemma B.2, the Besov embedding Lemma 2.1 as well as (5.18) and (5.19) and the definition of M_L we obtain (6.3).

Note that we only control the L^4 -norm instead of e.g. L^∞ because the para-product

$$v_q^1 \otimes \Delta_{\leq f(q)} \Delta_{>R} z$$

only belongs to $B_{5/3,\infty}^{1/3-2\kappa}$.

7. The Main Iteration: Proof of Proposition 5.1

7.1. Choice of Parameters

In the sequel, additional parameters will be indispensable and their value has to be carefully chosen in order to respect all the compatibility conditions appearing in the estimations below. First, for a sufficiently small $\alpha \in (0, 1)$ to be chosen below, we let $\ell \in (0, 1)$ be a small parameter satisfying

$$\ell^{4/5} \lambda_q^4 \leq \lambda_{q+1}^{-\alpha}, \quad \ell^{-1} \leq \lambda_{q+1}^{2\alpha}, \tag{7.1}$$

In particular, we define

$$\ell := \lambda_{q+1}^{-\frac{3\alpha}{2}} \lambda_q^{-2}. \tag{7.2}$$

In the sequel, we use the bounds

$$\alpha > 244\beta b, \quad 1 > 168\beta b^2, \quad \frac{1}{35} - 33\alpha > 2\beta b, \quad \alpha b > 128 \tag{7.3}$$

which can be obtained by choosing α small such that $\frac{1}{35} - 33\alpha > \alpha$, and choosing $b \in \mathbb{N}$ large enough such that $\alpha b > 128$ and finally choosing β small such that $\alpha > 244\beta b, 1 > 168\beta b^2$. Various estimates of this form are needed for the final control the new stress \mathring{R}_{q+1} . Hence, we shall choose α small first and b large, then β small enough. The last free parameter is a which is power of 2^{21} and satisfies the lower bounds given through

$$a > 4M_L + K, \quad L \leq a^{\alpha/16}.$$

Then by our condition we have

$$M_L(1 + 3q) + K \leq \lambda_q^{1/42} < \ell^{-2/183} < \lambda_{q+1}^{\alpha-2\beta}, \quad \sigma_q^{-1} < \ell^{-1/61}. \tag{7.4}$$

In the sequel, we increase a in order to absorb various implicit and universal constants.

We may freely increase the value of a provided we make β smaller at the same time.

7.2. Mollification

We intend to replace v_q^2 by a mollified velocity field v_ℓ . To this end, we extend

$$\begin{aligned} z(t) &= z(0), \quad z^{\dot{Y}}(t) = z^{\dot{Y}}(0) = 0, \quad z^{\dot{Y}'}(t) = 0, \\ \mathcal{I}(\nabla z)(t) &= 0, \quad v_q^i(t) = v_q^i(0), \quad \mathring{R}_q(t) = \mathring{R}_q(0) \quad \text{for } t < 0. \end{aligned}$$

As v_q^2 equals to zero near zero, $\partial_t v_q^2(0) = 0$, which implies by our extension that the equation holds also for $t < 0$. Let $\{\phi_\varepsilon\}_{\varepsilon>0}$ be a family of standard mollifiers

on \mathbb{R}^3 , and let $\{\varphi_\varepsilon\}_{\varepsilon>0}$ be a family of standard mollifiers with support on \mathbb{R}^+ . We define a mollification of v_q, \mathring{R}_q in space and time by convolution as

$$v_\ell = (v_q^2 *_x \phi_\ell) *_t \varphi_\ell, \quad \mathring{R}_\ell = (\mathring{R}_q *_x \phi_\ell) *_t \varphi_\ell,$$

where $\phi_\ell = \frac{1}{\ell^3} \phi(\frac{\cdot}{\ell})$ and $\varphi_\ell = \frac{1}{\ell} \varphi(\frac{\cdot}{\ell})$. Since the mollifier φ_ℓ is supported on \mathbb{R}^+ , it is easy to see that z_ℓ is $(\mathcal{F}_t)_{t \geq 0}$ -adapted and so are v_ℓ and \mathring{R}_ℓ . Then using the equation for v_q^2 we obtain that $(v_\ell, \mathring{R}_\ell)$ satisfies

$$\begin{aligned} \partial_t v_\ell - \Delta v_\ell + \operatorname{div} N + \nabla p_\ell &= \operatorname{div}(\mathring{R}_\ell + R_{\text{com}}) \\ \operatorname{div} v_\ell &= 0, \end{aligned} \tag{7.5}$$

where N is the trace-free part of the matrix

$$N = (v_q^1 + v_q^2) \otimes (v_q^1 + v_q^2) + V_q^2 + V_q^{2,*}$$

and

$$R_{\text{com}} = N - N *_x \phi_\ell *_t \varphi_\ell.$$

By using (5.7) (5.10) and (7.2) we know for $t \in [0, T_L]$

$$\begin{aligned} \|v_q^2 - v_\ell\|_{C_t^{1/10} L^{5/3}} + \|v_q^2 - v_\ell\|_{C_t W^{1/5,5/3}} &\lesssim \ell^{4/5} \|v_q^2\|_{C_{t,x}^1} \leq \ell^{4/5} \lambda_q^4 M_L^{1/2} \\ &\leq M_L^{1/2} \lambda_{q+1}^{-\alpha} \leq \frac{1}{4} M_L^{1/2} \delta_{q+1}^{1/2} a^{-\alpha/2}, \end{aligned} \tag{7.6}$$

$$\begin{aligned} \|v_q^2 - v_\ell\|_{C_t L^2} &\lesssim \ell \|v_q^2\|_{C_{t,x}^1} \leq \ell \lambda_q^4 M_L^{1/2} \\ &\leq M_L^{1/2} \lambda_{q+1}^{-\alpha} \leq \frac{1}{4} M_L^{1/2} \delta_{q+1}^{1/2}, \end{aligned} \tag{7.7}$$

and for $p \in [1, \infty]$

$$\begin{aligned} \|v_q^2 - v_\ell\|_{C_t W^{2/3,p}} &\lesssim \ell^{1/3} \|v_q^2\|_{C_{t,x}^1} \leq \ell^{1/3} \lambda_q^4 M_L^{1/2} \\ &\leq M_L^{1/2} \lambda_{q+1}^{-15\alpha/32} \leq \frac{1}{4} M_L^{1/2} \delta_{q+1}^{1/2} a^{-\alpha/2}, \end{aligned} \tag{7.8}$$

where we used the fact that $\alpha b > 128$ and $\alpha > 3\beta$ and we chose a large enough in order to absorb the implicit constant. In addition,

$$\|v_\ell\|_{C_{t,x}^N} \lesssim \ell^{-N+1} \|v_q\|_{C_{t,x}^1} \leq \ell^{-N+1} \lambda_q^4 M_L^{1/2} \lesssim M_L^{1/2} \ell^{-N} \lambda_{q+1}^{-\alpha} \tag{7.9}$$

holds for $t \in [0, T_L]$ and it holds for $t \in (\frac{\sigma_q}{2} \wedge T_L, T_L]$

$$\|v_\ell(t)\|_{L^2} \leq \|v_q^2\|_{C_t L^2} \lesssim M_0 (M_L^{1/2} + K^{1/2}) + 3M_0 M_L^{1/2} (1 + 3q)^{1/2} \tag{7.10}$$

with some universal implicit constant.

7.3. Construction of v_{q+1}^2 .

Let us proceed with the construction of the perturbation w_{q+1} , which then defines the next iteration by $v_{q+1}^2 := v_\ell + w_{q+1}$. To this end, we make use of the intermittent jets [9, Section 7.4], which we recall in Appendix A. In particular, the building blocks $W(\xi) = W_{\xi, r_\perp, r_\parallel, \lambda, \mu}$ for $\xi \in \Lambda$ are defined in (A.3) and the set Λ is introduced in Lemma A.1. The necessary estimates are collected in (A.7). For the intermittent jets we choose the following parameters:

$$\begin{aligned} \lambda &= \lambda_{q+1}, & r_\parallel &= \lambda_{q+1}^{-4/7}, & r_\perp &= r_\parallel^{-1/4} \lambda_{q+1}^{-1} = \lambda_{q+1}^{-6/7}, \\ \mu &= \lambda_{q+1} r_\parallel r_\perp^{-1} = \lambda_{q+1}^{9/7}. \end{aligned} \tag{7.11}$$

Since a is power of 2^{21} , $\lambda_{q+1} r_\perp = a^{(b^{q+1})/7} \in \mathbb{N}$.

Now we follow [45, Section 5.2] and introduce ρ as

$$\rho := 2\sqrt{\ell^2 + |\mathring{R}_\ell|^2} + \frac{\gamma_{q+1}}{(2\pi)^3},$$

which implies for $p \geq 1$

$$\|\rho(t)\|_{L^p} \leq 2\ell(2\pi)^{3/p} + 2\|\mathring{R}_\ell(t)\|_{L^p} + \gamma_{q+1}. \tag{7.12}$$

In view of (5.11), which holds on $(2\sigma_q \wedge T_L, T_L]$, and since $\text{supp}\varphi_\ell \subset [0, \ell]$, we obtain, for $t \in (4\sigma_q \wedge T_L, T_L]$, that

$$\|\rho\|_{C_{[4\sigma_q \wedge T_L, t], x}^0} \lesssim \ell^{-4} \delta_{q+1} M_L + \gamma_{q+1}, \tag{7.13}$$

where we also used the embedding $W^{4,1} \subset L^\infty$. Then, we deduce similarly as [45, (3.25)] for $N \geq 1$ and $t \in (4\sigma_q \wedge T_L, T_L]$

$$\begin{aligned} \|\rho\|_{C_{[4\sigma_q \wedge T_L, t], x}^N} &\lesssim \ell^{-4-N} M_L \delta_{q+1} + \ell^{-N+1} (\ell^{-5} M_L \delta_{q+1})^N \\ &+ \gamma_{q+1} \lesssim \ell^{2-7N} \delta_{q+1} M_L + \gamma_{q+1}. \end{aligned} \tag{7.14}$$

For a general $t \in [0, T_L]$, we have by (5.12),

$$\|\rho\|_{C_{t, x}^0} \lesssim \ell^{-4} M_L (1 + 3q) + \gamma_{q+1}, \tag{7.15}$$

and for $N \geq 1$,

$$\|\rho\|_{C_{t, x}^N} \lesssim \ell^{2-7N} M_L (1 + 3q) + \gamma_{q+1}, \tag{7.16}$$

where we used $M_L (1 + 3q) \leq \ell^{-1}$.

Next, we define the amplitude functions

$$\begin{aligned} a(\xi)(\omega, t, x) &:= a_{\xi, q+1}(\omega, t, x) \\ &:= \rho(\omega, t, x)^{1/2} \gamma_\xi \left(\text{Id} - \frac{\mathring{R}_\ell(\omega, t, x)}{\rho(\omega, t, x)} \right) (2\pi)^{-3/4}, \end{aligned} \tag{7.17}$$

where γ_ξ is introduced in Lemma A.1. Since ρ and \mathring{R}_ℓ are $(\mathcal{F}_t)_{t \geq 0}$ -adapted, we know that also $a_{(\xi)}$ is $(\mathcal{F}_t)_{t \geq 0}$ -adapted. By (A.5) we have

$$(2\pi)^{-\frac{3}{2}} \sum_{\xi \in \Lambda} a_{(\xi)}^2 \int_{\mathbb{T}^3} W_{(\xi)} \otimes W_{(\xi)} dx = \rho \text{Id} - \mathring{R}_\ell. \tag{7.18}$$

By using (7.12), for $t \in (4\sigma_q \wedge T_L, T_L]$,

$$\begin{aligned} \|a_{(\xi)}(t)\|_{L^2} &\leq \|\rho(t)\|_{L^1}^{1/2} \|\gamma_\xi\|_{C^0(B_{1/2}(\text{Id}))} \\ &\leq \frac{M}{8|\Lambda|(1+8\pi^3)^{1/2}} \left(2(2\pi)^3 \ell + 2\delta_{q+1} M_L + \gamma_{q+1}\right)^{1/2} \\ &\leq \frac{M}{4|\Lambda|} (M_L^{1/2} \delta_{q+1}^{1/2} + \gamma_{q+1}^{1/2}), \end{aligned} \tag{7.19}$$

and for $t \in [0, T_L]$,

$$\|a_{(\xi)}(t)\|_{C_t L^2} \leq \frac{M}{4|\Lambda|} (M_L^{1/2} (1+3q)^{1/2} + \gamma_{q+1}^{1/2}),$$

where M denotes the universal constant from Lemma A.1. From (7.13), (7.14), and similarly to [45, (3.30)], we deduce for $t \in (4\sigma_q \wedge T_L, T_L]$ that

$$\|\rho^{1/2}\|_{C_{[4\sigma_q \wedge T_L, t], x}^0} \lesssim \ell^{-2} \delta_{q+1}^{1/2} M_L^{1/2} + \gamma_{q+1}^{1/2},$$

and for $m = 1, \dots, N$, using $K \leq \ell^{-1}$,

$$\begin{aligned} \|\rho^{1/2}\|_{C_{[4\sigma_q \wedge T_L, t], x}^m} &\lesssim \ell^{1-7m} \delta_{q+1}^{1/2} M_L^{1/2} + \ell^{1/2-m} (\gamma_{q+1} + \ell^{-5} \delta_{q+1} M_L)^m \\ &\leq \ell^{1-7m} (\delta_{q+1}^{1/2} M_L^{1/2} + \gamma_{q+1}^{1/2}). \end{aligned}$$

This implies, for $N \in \mathbb{N}_0$, as in [45, (3.34)],

$$\|a_{(\xi)}\|_{C_{[4\sigma_q \wedge T_L, t], x}^N} \lesssim \ell^{-8-7N} (\delta_{q+1}^{1/2} M_L^{1/2} + \gamma_{q+1}^{1/2}). \tag{7.20}$$

For a general $t \in [0, T_L]$ we have, for $N \in \mathbb{N}_0$,

$$\|a_{(\xi)}\|_{C_{t,x}^N} \lesssim \ell^{-8-7N} (M_L^{1/2} (1+3q)^{1/2} + \gamma_{q+1}^{1/2}), \tag{7.21}$$

where we used $M_L(1+3q) + K \leq \ell^{-1}$.

Let us introduce a smooth cut-off function

$$\chi(t) = \begin{cases} 0, & t \leq \frac{\sigma_q}{2}, \\ \in (0, 1), & t \in (\frac{\sigma_q}{2}, \sigma_q), \\ 1, & t \geq \sigma_q. \end{cases}$$

Note that $\|\chi'\|_{C_t^0} \leq 2^{q+1}$ which has to be taken into account in the estimates of the $C_t^{1/10} L^{5/3}$ and $C_{t,x}^1$ -norms in (7.39)- (7.45) below.

With these preparations in hand, we define the principal part $w_{q+1}^{(p)}$ of the perturbation w_{q+1} as

$$w_{q+1}^{(p)} := \sum_{\xi \in \Lambda} a_{(\xi)} W_{(\xi)}. \tag{7.22}$$

Since the coefficients $a_{(\xi)}$ are $(\mathcal{F}_t)_{t \geq 0}$ -adapted and $W_{(\xi)}$ is a deterministic function we deduce that $w_{q+1}^{(p)}$ is also $(\mathcal{F}_t)_{t \geq 0}$ -adapted. Moreover, according to (7.18) and (A.4) it follows that

$$w_{q+1}^{(p)} \otimes w_{q+1}^{(p)} + \mathring{R}_\ell = \sum_{\xi \in \Lambda} a_{(\xi)}^2 \mathbb{P}_{\neq 0}(W_{(\xi)} \otimes W_{(\xi)}) + \rho \text{Id}, \tag{7.23}$$

where we use the notation $\mathbb{P}_{\neq 0} f := f - \mathcal{F}f(0) = f - (2\pi)^{-3/2} \int_{\mathbb{T}^3} f dx$.

We also define the incompressibility corrector by

$$w_{q+1}^{(c)} := \sum_{\xi \in \Lambda} \text{curl}(\nabla a_{(\xi)} \times V_{(\xi)}) + \nabla a_{(\xi)} \times \text{curl} V_{(\xi)} + a_{(\xi)} W_{(\xi)}^{(c)}, \tag{7.24}$$

with $W_{(\xi)}^{(c)}$ and $V_{(\xi)}$ being given in (A.6). Since $a_{(\xi)}$ is $(\mathcal{F}_t)_{t \geq 0}$ -adapted and $W_{(\xi)}$, $W_{(\xi)}^{(c)}$ and $V_{(\xi)}$ are deterministic functions we know that $w_{q+1}^{(c)}$ is also $(\mathcal{F}_t)_{t \geq 0}$ -adapted. By a direct computation we deduce that

$$w_{q+1}^{(p)} + w_{q+1}^{(c)} = \sum_{\xi \in \Lambda} \text{curl} \text{curl}(a_{(\xi)} V_{(\xi)}),$$

hence

$$\text{div}(w_{q+1}^{(p)} + w_{q+1}^{(c)}) = 0.$$

We also introduce a temporal corrector

$$w_{q+1}^{(t)} := -\frac{1}{\mu} \sum_{\xi \in \Lambda} \mathbb{P} \mathbb{P}_{\neq 0} \left(a_{(\xi)}^2 \phi_{(\xi)}^2 \psi_{(\xi)}^2 \xi \right), \tag{7.25}$$

where \mathbb{P} is the Helmholtz projection. Similarly to above $w_{q+1}^{(t)}$ is $(\mathcal{F}_t)_{t \geq 0}$ -adapted and by similar computation as [9, (7.38)] we obtain

$$\begin{aligned} & \partial_t w_{q+1}^{(t)} + \sum_{\xi \in \Lambda} \mathbb{P}_{\neq 0} \left(a_{(\xi)}^2 \text{div}(W_{(\xi)} \otimes W_{(\xi)}) \right) \\ &= -\frac{1}{\mu} \sum_{\xi \in \Lambda} \mathbb{P} \mathbb{P}_{\neq 0} \partial_t \left(a_{(\xi)}^2 \phi_{(\xi)}^2 \psi_{(\xi)}^2 \xi \right) + \frac{1}{\mu} \sum_{\xi \in \Lambda} \mathbb{P}_{\neq 0} \left(a_{(\xi)}^2 \partial_t (\phi_{(\xi)}^2 \psi_{(\xi)}^2 \xi) \right) \tag{7.26} \\ &= (\text{Id} - \mathbb{P}) \frac{1}{\mu} \sum_{\xi \in \Lambda} \mathbb{P}_{\neq 0} \partial_t \left(a_{(\xi)}^2 \phi_{(\xi)}^2 \psi_{(\xi)}^2 \xi \right) - \frac{1}{\mu} \sum_{\xi \in \Lambda} \mathbb{P}_{\neq 0} \left(\partial_t a_{(\xi)}^2 (\phi_{(\xi)}^2 \psi_{(\xi)}^2 \xi) \right). \end{aligned}$$

Note that the first term on the right hand side can be viewed as a pressure term.

We define the truncated perturbations $\tilde{w}_{q+1}^{(p)}, \tilde{w}_{q+1}^{(c)}, \tilde{w}_{q+1}^{(t)}$ as follows:

$$\tilde{w}_{q+1}^{(p)} := w_{q+1}^{(p)}\chi, \quad \tilde{w}_{q+1}^{(c)} := w_{q+1}^{(c)}\chi, \quad \tilde{w}_{q+1}^{(t)} := w_{q+1}^{(t)}\chi^2.$$

Define $w_{q+1} := \tilde{w}_{q+1}^{(p)} + \tilde{w}_{q+1}^{(c)} + \tilde{w}_{q+1}^{(t)}$ and

$$v_{q+1}^2 = v_\ell + w_{q+1} = v_\ell + \tilde{w}_{q+1}^{(p)} + \tilde{w}_{q+1}^{(c)} + \tilde{w}_{q+1}^{(t)}.$$

We note that by construction v_{q+1}^2 is $(\mathcal{F}_t)_{t \geq 0}$ adapted.

7.4. Verification of the Inductive Estimates for v_{q+1}^2

By (7.19) and (7.20) and similar argument as [45] we obtain for $t \in (4\sigma_q \wedge T_L, T_L]$ and some universal constant $M_0 \geq 1$

$$\begin{aligned} \|\tilde{w}_{q+1}^{(p)}(t)\|_{L^2} &\lesssim \sum_{\xi \in \Lambda} \frac{1}{4|\Lambda|} M(M_L^{1/2}\delta_{q+1}^{1/2} + \gamma_{q+1}^{1/2}) \|W_{(\xi)}\|_{C_t L^2} \\ &\leq \frac{M_0}{2} (M_L^{1/2}\delta_{q+1}^{1/2} + \gamma_{q+1}^{1/2}), \end{aligned} \tag{7.27}$$

where we used $150\alpha < \frac{1}{7}$ and for $t \in (\frac{\sigma_q}{2} \wedge T_L, 4\sigma_q \wedge T_L]$

$$\|\tilde{w}_{q+1}^{(p)}(t)\|_{L^2} \leq \frac{M_0}{2} ((M_L(1 + 3q))^{1/2} + \gamma_{q+1}^{1/2}). \tag{7.28}$$

Similarly as in [45, (3.43)-(3.46)], we apply (A.7) and (7.20) for general L^p -norms to deduce for $t \in (4\sigma_q \wedge T_L, T_L], p \in (1, \infty)$

$$\|\tilde{w}_{q+1}^{(p)}(t)\|_{L^p} \lesssim \ell^{-8} (M_L^{1/2}\delta_{q+1}^{1/2} + \gamma_{q+1}^{1/2}) r_\perp^{2/p-1} r_\parallel^{1/p-1/2}, \tag{7.29}$$

$$\|\tilde{w}_{q+1}^{(c)}(t)\|_{L^p} \lesssim \ell^{-22} (M_L^{1/2}\delta_{q+1}^{1/2} + \gamma_{q+1}^{1/2}) r_\perp^{2/p} r_\parallel^{1/p-3/2}, \tag{7.30}$$

and

$$\|\tilde{w}_{q+1}^{(t)}(t)\|_{L^p} \lesssim \ell^{-16} (M_L\delta_{q+1} + \gamma_{q+1}) r_\perp^{2/p-1} r_\parallel^{1/p-2} \lambda_{q+1}^{-1}, \tag{7.31}$$

$$\|\tilde{w}_{q+1}^{(c)}(t)\|_{L^p} + \|\tilde{w}_{q+1}^{(t)}(t)\|_{L^p} \lesssim \ell^{-8} (M_L^{1/2}\delta_{q+1}^{1/2} + \gamma_{q+1}^{1/2}) r_\perp^{2/p-1} r_\parallel^{1/p-1/2} \tag{7.32}$$

For a general $t \in (\frac{\sigma_q}{2} \wedge T_L, 4\sigma_q \wedge T_L]$ we have

$$\|\tilde{w}_{q+1}^{(p)}(t)\|_{L^p} \lesssim \ell^{-8} ((M_L(1 + 3q))^{1/2} + \gamma_{q+1}^{1/2}) r_\perp^{2/p-1} r_\parallel^{1/p-1/2}, \tag{7.33}$$

$$\|\tilde{w}_{q+1}^{(c)}(t)\|_{L^p} \lesssim \ell^{-22} ((M_L(1 + 3q))^{1/2} + \gamma_{q+1}^{1/2}) r_\perp^{2/p} r_\parallel^{1/p-3/2}, \tag{7.34}$$

and

$$\|\tilde{w}_{q+1}^{(t)}(t)\|_{L^p} \lesssim \ell^{-16} ((M_L(1 + 3q)) + \gamma_{q+1}) r_\perp^{2/p-1} r_\parallel^{1/p-2} \lambda_{q+1}^{-1}, \tag{7.35}$$

$$\|\tilde{w}_{q+1}^{(c)}(t)\|_{L^p} + \|\tilde{w}_{q+1}^{(t)}(t)\|_{L^p} \lesssim \ell^{-8}((M_L(1 + 3q))^{1/2} + \gamma_{q+1}^{1/2})r_{\perp}^{2/p-1}r_{\parallel}^{1/p-1/2}. \tag{7.36}$$

Combining (7.27), (7.30) and (7.31) we obtain, for $t \in (4\sigma_q \wedge T_L, T_L]$,

$$\begin{aligned} \|w_{q+1}(t)\|_{L^2} &\leq (M_L^{1/2}\delta_{q+1}^{1/2} + \gamma_{q+1}^{1/2}) \\ &\quad \left(\frac{M_0}{2} + C\lambda_{q+1}^{44\alpha-2/7} + C(M_L^{1/2}\delta_{q+1}^{1/2} + \gamma_{q+1}^{1/2})\lambda_{q+1}^{32\alpha-1/7} \right) \tag{7.37} \\ &\leq \frac{3}{4}M_0(M_L^{1/2}\delta_{q+1}^{1/2} + \gamma_{q+1}^{1/2}), \end{aligned}$$

and for $t \in (\frac{\sigma_q}{2} \wedge T_L, 4\sigma_q \wedge T_L]$

$$\|w_{q+1}(t)\|_{L^2} \leq \frac{3}{4}M_0((M_L(1 + 3q))^{1/2} + \gamma_{q+1}^{1/2}), \tag{7.38}$$

where we used $M_L(1 + 3q) + K < \ell^{-1}$ and the condition on α .

With these bounds, we have all in hand to complete the proof of Proposition 5.1. We split the details into several subsections.

7.4.1. Proof of (5.13) First, (7.37), together with (7.7), yields, for $t \in (4\sigma_q \wedge T_L, T_L]$,

$$\begin{aligned} \|v_{q+1}^2(t) - v_q^2(t)\|_{L^2} &\leq \|w_{q+1}(t)\|_{L^2} + \|v_{\ell}(t) - v_q^2(t)\|_{L^2} \\ &\leq M_0(M_L^{1/2}\delta_{q+1}^{1/2} + \gamma_{q+1}^{1/2}). \end{aligned}$$

For $t \in (\frac{1}{2}\sigma_q \wedge T_L, 4\sigma_q \wedge T_L]$ we use (7.38), (7.7) to obtain

$$\begin{aligned} \|v_{q+1}^2(t) - v_q^2(t)\|_{L^2} &\leq \|w_{q+1}(t)\|_{L^2} + \|v_{\ell}(t) - v_q^2(t)\|_{L^2} \\ &\leq M_0((M_L(1 + 3q))^{1/2} + \gamma_{q+1}^{1/2}). \end{aligned}$$

For $t \in [0, \frac{1}{2}\sigma_q \wedge T_L]$ it holds that $\chi(t) = 0$, as well as $v_q^2(t) = 0$, by (5.9), implying that

$$\|v_{q+1}^2 - v_q^2\|_{C_t L^2} = \|v_{\ell} - v_q^2\|_{C_t L^2} = 0.$$

Hence (5.13) follows.

7.4.2. Proof that (5.13) Implies (5.9) on the Level $q + 1$ From (5.13) we find for $t \in [0, \frac{1}{2}\sigma_q \wedge T_L]$

$$\|v_{q+1}^2\|_{C_t L^2} \leq \sum_{r=0}^q \|v_{r+1}^2 - v_r^2\|_{C_t L^2} = 0,$$

proving the second bound in (5.9) on the level $q + 1$. For the first bound in (5.9) on the level $q + 1$, we obtain, in view of (5.13), for $t \in (\frac{1}{2}\sigma_q \wedge T_L, T_L]$,

$$\begin{aligned} \|v_{q+1}^2(t)\|_{L^2} &\leq \sum_{0 \leq r \leq q} \|v_{r+1}^2(t) - v_r^2(t)\|_{L^2} \\ &\leq M_0 \left(M_L^{1/2} \sum_{0 \leq r \leq q} \delta_{r+1}^{1/2} \right. \\ &\quad \left. + \sum_{0 \leq r \leq q} (M_L(1 + 3r))^{1/2} \mathbf{1}_{t \in (\frac{\sigma_r}{2} \wedge T_L, 4\sigma_r \wedge T_L]} + \sum_{0 \leq r \leq q} \gamma_{r+1}^{1/2} \right) \\ &\leq M_0 \left(M_L^{1/2} \sum_{0 \leq r \leq q} \delta_{r+1}^{1/2} + 3(M_L(1 + 3q))^{1/2} + \sum_{0 \leq r \leq q} \gamma_{r+1}^{1/2} \right), \end{aligned}$$

where we used the fact that by the definition of $\sigma_r = 2^{-r}$ each $t \in [0, T_L]$ only belongs to three intervals $(\frac{\sigma_r}{2} \wedge T_L, 4\sigma_r \wedge T_L]$. Hence (5.9) follows.

7.4.3. Proof of (5.16) and the Second Inequality in (5.7) on the Level $q + 1$ In this section, we see in particular how the definition of intermittent jets determines the integrability $5/3$ which we use throughout the paper. It holds by (7.21), (A.7) and the choice of parameters in (7.3)

$$\begin{aligned} &\|\tilde{w}_{q+1}^{(p)}\|_{C_t^{1/10} L^{5/3}} + \|\tilde{w}_{q+1}^{(p)}\|_{C_t W^{1/5, 5/3}} \\ &\lesssim \sum_{\xi \in \Lambda} \|a(\xi)\|_{C_{t,x}^1} (\|W(\xi)\|_{C_t^{1/10} L^{5/3}} + \|W(\xi)\|_{C_t W^{1/5, 5/3}}) 2^{q+1} \\ &\lesssim (M_L^{1/2} (1 + 3q)^{1/2} + \gamma_{q+1}^{1/2}) \ell^{-15} r_{\perp}^{1/5} r_{\parallel}^{1/10} ((r_{\perp} \lambda_{q+1} \mu / r_{\parallel})^{1/10} + \lambda_{q+1}^{1/5}) 2^{q+1} \\ &\lesssim M_L^{1/2} \lambda_{q+1}^{32\alpha - 1/35}, \tag{7.39} \end{aligned}$$

$$\begin{aligned} &\|\tilde{w}_{q+1}^{(c)}\|_{C_t^{1/10} L^{5/3}} + \|\tilde{w}_{q+1}^{(c)}\|_{C_t W^{1/5, 5/3}} \\ &\lesssim \sum_{\xi \in \Lambda} 2^{q+1} \left(\|a(\xi)\|_{C_{t,x}^1} (\|W(\xi)^{(c)}\|_{C_t^{1/10} L^{5/3}} + \|W(\xi)^{(c)}\|_{C_t W^{1/5, 5/3}}) \right. \\ &\quad \left. + \|a(\xi)\|_{C_{t,x}^3} (\|V(\xi)\|_{C_t^{1/10} W^{1, 5/3}} + \|V(\xi)\|_{C_t W^{6/5, 5/3}}) \right) \\ &\lesssim (M_L^{1/2} (1 + 3q)^{1/2} + \gamma_{q+1}^{1/2}) \ell^{-29} r_{\perp}^{1/5} r_{\parallel}^{1/10} \\ &\quad \left(r_{\perp} r_{\parallel}^{-1} + \lambda_{q+1}^{-1} \right) \left((r_{\perp} \lambda_{q+1} \mu / r_{\parallel})^{1/10} + \lambda_{q+1}^{1/5} \right) 2^{q+1} \end{aligned}$$

$$\begin{aligned} &\lesssim M_L^{1/2} \ell^{-30} r_{\perp}^{6/5} r_{\parallel}^{-9/10} \left((r_{\perp} \lambda_{q+1} \mu / r_{\parallel})^{1/10} + \lambda_{q+1}^{1/5} \right) \\ &\lesssim M_L^{1/2} \lambda_{q+1}^{60\alpha-11/35}, \end{aligned} \tag{7.40}$$

$$\begin{aligned} &\|\tilde{w}_{q+1}^{(t)}\|_{C_t^{1/10} L^{5/3}} + \|\tilde{w}_{q+1}^{(t)}\|_{C_t W^{1/5,5/3}} \\ &\leq \frac{1}{\mu} 2^{q+1} \sum_{\xi \in \Lambda} \left(\|a(\xi)\|_{C_{t,x}^0} \|a(\xi)\|_{C_{t,x}^1} \|\phi(\xi)\|_{L^{10/3}}^2 \|\psi(\xi)\|_{C_t L^{10/3}}^2 \right. \\ &\quad + \|a(\xi)\|_{C_{t,x}^0}^2 \|\phi(\xi)\|_{L^{10/3}} \|\phi(\xi)\|_{W^{1/5,10/3}} \|\psi(\xi)\|_{C_t L^{10/3}}^2 \\ &\quad + \|a(\xi)\|_{C_{t,x}^0}^2 \|\phi(\xi)\|_{L^{10/3}}^2 \|\psi(\xi)\|_{C_t W^{1/5,10/3}} \|\psi(\xi)\|_{C_t L^{10/3}} \\ &\quad \left. + \|a(\xi)\|_{C_{t,x}^0}^2 \|\phi(\xi)\|_{L^{10/3}}^2 \|\psi(\xi)\|_{C_t^{1/10} L^{10/3}} \|\psi(\xi)\|_{C_t L^{10/3}} \right) \\ &\lesssim (M_L(1 + 3q) + \gamma_{q+1}) \ell^{-23} r_{\perp}^{1/5} r_{\parallel}^{-7/5} (\mu^{-1} r_{\perp}^{-1} r_{\parallel}) \\ &\quad \times \left((r_{\perp} \lambda_{q+1} \mu / r_{\parallel})^{1/10} + \lambda_{q+1}^{1/5} + (\lambda_{q+1} r_{\perp} r_{\parallel}^{-1})^{1/5} \right) 2^{q+1} \\ &\lesssim M_L \lambda_{q+1}^{48\alpha-6/35}, \end{aligned} \tag{7.41}$$

and

$$\|w_{q+1}\|_{C_t^{1/10} L^{5/3}} + \|w_{q+1}\|_{C_t W^{1/5,5/3}} \lesssim M_L^{1/2} \lambda_{q+1}^{32\alpha-1/35} \leq \frac{3}{4} M_L^{1/2} \delta_{q+1} a^{-\alpha/2} \tag{7.42}$$

In the last inequality above we used (7.3). Hence, (5.16) follows from (7.6). The second inequality in (5.7) on the level $q + 1$ follows as well.

7.4.4. Proof of (5.10) on the Level $q + 1$ Using (7.21) and similar as [45, Section 3.1.4] we find, for $t \in [0, T_L]$,

$$\|\tilde{w}_{q+1}^{(p)}\|_{C_{t,x}^1} \lesssim \ell^{-15} ((M_L(1 + 3q))^{1/2} + \gamma_{q+1}^{1/2}) r_{\perp}^{-1} r_{\parallel}^{-1/2} \lambda_{q+1}^2, \tag{7.43}$$

$$\|\tilde{w}_{q+1}^{(c)}\|_{C_{t,x}^1} \lesssim \ell^{-29} ((M_L(1 + 3q))^{1/2} + \gamma_{q+1}^{1/2}) r_{\parallel}^{-3/2} \lambda_{q+1}^2, \tag{7.44}$$

and

$$\|\tilde{w}_{q+1}^{(t)}\|_{C_{t,x}^1} \lesssim \ell^{-24} (M_L(1 + 3q) + \gamma_{q+1}) r_{\perp}^{-1} r_{\parallel}^{-2} \lambda_{q+1}^{1+\alpha}. \tag{7.45}$$

In particular, we see that the fact that the time derivative of χ behaves like $2\sigma_q^{-1} \lesssim \ell^{-1}$ does not pose any problems as the $C_{t,x}^0$ -norms of $\tilde{w}_{q+1}^{(p)}$, $\tilde{w}_{q+1}^{(c)}$ and $\tilde{w}_{q+1}^{(t)}$ always contain smaller powers of ℓ^{-1} .

Combining (7.9) and (7.43), (7.44), (7.45) with (7.1) we obtain, for $t \in [0, T_L]$,

$$\begin{aligned} \|v_{q+1}^2\|_{C_{t,x}^1} &\leq \|v_{\ell}\|_{C_{t,x}^1} + \|w_{q+1}\|_{C_{t,x}^1} \\ &\leq (M_L(1 + 3q) + \gamma_{q+1})^{1/2} \\ &\quad \left(\lambda_{q+1}^{\alpha} + C \lambda_{q+1}^{30\alpha+22/7} + C \lambda_{q+1}^{58\alpha+20/7} + C \lambda_{q+1}^{50\alpha+3} \right) \\ &\leq M_L^{1/2} \lambda_{q+1}^4, \end{aligned}$$

where we used $M_L(1 + 3q) + \gamma_{q+1} \leq \ell^{-1}$. This implies (5.10).

7.4.5. Proof of (5.15) and the First Inequality in (5.7) on the Level $q + 1$
 Similarly, we derive the following estimates: for $t \in [0, T_L]$ it follows from (7.1), (7.21) and (A.7) that

$$\begin{aligned} \|\tilde{w}_{q+1}^{(p)} + \tilde{w}_{q+1}^{(c)}\|_{C_t W^{1,p}} &\leq \sum_{\xi \in \Lambda} \|\text{curl curl}(a(\xi) V(\xi))\|_{C_t W^{1,p}} \\ &\lesssim \sum_{\xi \in \Lambda} \|a(\xi)\|_{C_{t,x}^3} \|V(\xi)\|_{C_t W^{3,p}} \\ &\lesssim \ell^{-29} ((M_L(1 + 3q))^{1/2} + \gamma_{q+1}^{1/2}) r_{\perp}^{2/p-1} r_{\parallel}^{1/p-1/2} \lambda_{q+1}, \end{aligned} \tag{7.46}$$

and

$$\begin{aligned} \|\tilde{w}_{q+1}^{(t)}\|_{C_t W^{1,p}} &\leq \frac{1}{\mu} \sum_{\xi \in \Lambda} \left(\|a(\xi)\|_{C_{t,x}^0} \|a(\xi)\|_{C_{t,x}^1} \|\phi(\xi)\|_{L^{2p}}^2 \|\psi(\xi)\|_{C_t L^{2p}}^2 \right. \\ &\quad + \|a(\xi)\|_{C_{t,x}^0}^2 \|\phi(\xi)\|_{L^{2p}} \|\nabla \phi(\xi)\|_{L^{2p}} \|\psi(\xi)\|_{C_t L^{2p}}^2 \\ &\quad \left. + \|a(\xi)\|_{C_{t,x}^0}^2 \|\phi(\xi)\|_{L^{2p}}^2 \|\nabla \psi(\xi)\|_{C_t L^{2p}} \|\psi(\xi)\|_{C_t L^{2p}} \right) \\ &\lesssim \ell^{-23} (M_L(1 + 3q) + \gamma_{q+1}) r_{\perp}^{2/p-2} r_{\parallel}^{1/p-1} \lambda_{q+1}^{-2/7}. \end{aligned} \tag{7.47}$$

We also have, for $p = \frac{32}{32-7\alpha}, r_{\perp}^{2/p-2} r_{\parallel}^{1/p-1} \leq \lambda_{q+1}^{\alpha}$ and

$$\begin{aligned} \|w_{q+1}\|_{C_t W^{1,p}} &\lesssim (M_L(1 + 3q) + \gamma_{q+1}) \ell^{-29} \lambda_{q+1}^{\alpha-1/7} \lesssim M_L^{1/2} \lambda_{q+1}^{60\alpha-1/7} \leq \\ &\frac{3}{4} M_L^{1/2} \delta_{q+1}^{1/2} a^{-\alpha/2}, \end{aligned} \tag{7.48}$$

where we used the condition for α, β and (7.3) in the second step, which combined with (7.8) implies (5.15) and hence the first inequality of (5.7).

7.5. Proof of (5.17)

We control the energy similarly as in [45, Section 3.1.5]. By definition, we find that

$$\begin{aligned} & \left| \|v_{q+1}^2\|_{L^2}^2 - \|v_q^2\|_{L^2}^2 - 3\gamma_{q+1} \right| \\ & \leq \left| \|\tilde{w}_{q+1}^{(p)}\|_{L^2}^2 - 3\gamma_{q+1} \right| + \|\tilde{w}_{q+1}^{(c)} + \tilde{w}_{q+1}^{(t)}\|_{L^2}^2 \\ & \quad + 2\|v_{\ell}(\tilde{w}_{q+1}^{(c)} + \tilde{w}_{q+1}^{(t)})\|_{L^1} + 2\|v_{\ell} \tilde{w}_{q+1}^{(p)}\|_{L^1} \\ & \quad + 2\|\tilde{w}_{q+1}^{(p)}(w_{q+1}^{(c)} + \tilde{w}_{q+1}^{(t)})\|_{L^1} + \left| \|v_{\ell}\|_{L^2}^2 - \|v_q^2\|_{L^2}^2 \right|. \end{aligned} \tag{7.49}$$

Let us begin with the bound of the first term on the right hand side of (7.49). We use (7.23) and the fact that \mathring{R}_{ℓ} is traceless to deduce for $t \in (4\sigma_q \wedge T_L, T_L]$

$$|\tilde{w}_{q+1}^{(p)}|^2 - \frac{3\gamma_{q+1}}{(2\pi)^3} = 6\sqrt{\ell^2 + |\mathring{R}_{\ell}|^2} + \sum_{\xi \in \Lambda} a_{(\xi)}^2 P_{\neq 0} |W_{(\xi)}|^2,$$

hence

$$\| \tilde{w}_{q+1}^{(p)} \|_{L^2}^2 - 3\gamma_{q+1} \leq 6 \cdot (2\pi)^3 \ell + 6 \| \dot{R}_\ell \|_{L^1} + \sum_{\xi \in \Lambda} \left| \int a_{(\xi)}^2 P_{\neq 0} |W_{(\xi)}|^2 \right|. \tag{7.50}$$

Here we estimate each term separately. Using (7.2) we find that

$$6 \cdot (2\pi)^3 \ell \leq 6 \cdot (2\pi)^3 \lambda_{q+1}^{-3\alpha/2} \leq \frac{1}{48} \lambda_{q+1}^{-2\beta} M_L \leq \frac{1}{48} \delta_{q+1} M_L,$$

which requires $2\beta < 3\alpha/2$ and choosing a large to absorb the constant. Using (5.11) on \dot{R}_q and $\text{supp} \varphi_\ell \subset [0, \ell]$ we know, for $t \in (4\sigma_q \wedge T_L, T_L]$, that

$$6 \| \dot{R}_\ell(t) \|_{L^1} \leq 6 \delta_{q+1} M_L.$$

For the last term in (7.50) we use a similar argument as to [45, Section 3.1.5] to get since $M_L \geq 1$, that

$$\begin{aligned} \sum_{\xi \in \Lambda} \left| \int a_{(\xi)}^2 \mathbb{P}_{\neq 0} |W_{(\xi)}|^2 \right| &\lesssim \lambda_{q+1}^{158\alpha-1/7} (M_L(1+3q) + K) \lesssim \lambda_{q+1}^{160\alpha-1/7} \\ &\leq \frac{1}{24} \lambda_1^{2\beta} \lambda_{q+1}^{-2\beta} M_L = \frac{1}{24} \delta_{q+1} M_L, \end{aligned}$$

where we used $M_L(1+3q) + K \leq \ell^{-1} \leq \lambda_{q+1}^{2\alpha}$ as well as $160\alpha + 2\beta < 1/7$. This completes the bound of (7.50).

Going back to (7.49), we control the remaining terms as follow: using the estimates (7.30), (7.31) and (7.1) we have, for $t \in (4\sigma_q \wedge T_L, T_L]$,

$$\begin{aligned} \| \tilde{w}_{q+1}^{(c)} + \tilde{w}_{q+1}^{(t)} \|_{L^2}^2 &\lesssim (M_L + \gamma_{q+1}) \lambda_{q+1}^{88\alpha-4/7} + (M_L^2 + \gamma_{q+1}^2) \lambda_{q+1}^{64\alpha-2/7} \\ &\leq \frac{1}{48} \lambda_{q+1}^{-2\beta} M_L \leq \frac{\delta_{q+1}}{48} M_L, \end{aligned}$$

where we use $M_L + \gamma_{q+1} \leq \ell^{-1}$ to control $M_L + \gamma_{q+1}$. Similarly we use (7.10) together with (7.27) to get, for $t \in (4\sigma_q \wedge T_L, T_L]$, that

$$\begin{aligned} &2 \| v_\ell (\tilde{w}_{q+1}^{(c)} + \tilde{w}_{q+1}^{(t)}) \|_{L^1} + 2 \| \tilde{w}_{q+1}^{(p)} (\tilde{w}_{q+1}^{(c)} + \tilde{w}_{q+1}^{(t)}) \|_{L^1} \\ &\lesssim M_0 ((M_L(1+3q))^{1/2} + K^{1/2}) \| \tilde{w}_{q+1}^{(c)} + \tilde{w}_{q+1}^{(t)} \|_{L^2} \\ &\lesssim M_0 ((M_L(1+3q))^{1/2} + K^{1/2}) \\ &\quad \left((M_L^{1/2} + \gamma_{q+1}^{1/2}) \lambda_{q+1}^{44\alpha-2/7} + (M_L + \gamma_{q+1}) \lambda_{q+1}^{32\alpha-1/7} \right) \\ &\leq \frac{1}{48} \lambda_{q+1}^{-2\beta} M_L \leq \frac{\delta_{q+1}}{48} M_L, \end{aligned}$$

where we used $M_L(1+3q) + K \leq \ell^{-1}$ and we possibly increased a to absorb M_0 . We use (7.1) and (7.29) and $\| v_\ell \|_{C_{t,x}^1} \leq \| v_q^2 \|_{C_{t,x}^1}$ to get, for every $\kappa > 0$,

$$2 \| v_\ell \tilde{w}_{q+1}^{(p)} \|_{L^1} \lesssim \| v_\ell \|_{L^\infty} \| \tilde{w}_{q+1}^{(p)} \|_{L^1} \lesssim M_L^{1/2} \lambda_q^4 \ell^{-8} (M_L^{1/2} \delta_{q+1}^{1/2} + \gamma_{q+1}^{1/2}) r_\perp^{1-\kappa} r_\parallel^{\frac{1}{2}(1-\kappa)}$$

$$\lesssim (M_L + K)\lambda_{q+1}^{17\alpha - \frac{8}{7}(1-\kappa)} \leq \frac{1}{96}\lambda_{q+1}^{-2\beta}M_L \leq \frac{\delta_{q+1}}{96}M_L.$$

For the last terms by (7.10) we get

$$\begin{aligned} |\|v_\ell\|_{L^2}^2 - \|v_q^2\|_{L^2}^2| &\leq \|v_\ell - v_q^2\|_{L^2}(\|v_\ell\|_{L^2} + \|v_q^2\|_{L^2}) \\ &\lesssim \ell\lambda_q^4 M_L^{1/2} M_0(M_L(1 + 3q) + K)^{1/2} \\ &\leq \frac{1}{96}\lambda_{q+1}^{-2\beta}M_L \leq \frac{\delta_{q+1}}{96}M_L, \end{aligned}$$

which requires $M_L(1 + 3q) + K < \lambda_{q+1}^{\alpha-2\beta}$ as in (7.4) and a large enough to absorb the extra constant.

Combining the above estimate (5.17) follows.

7.6. Definition of the Reynolds Stress \mathring{R}_{q+1}

Considering the equation for the difference $v_{q+1}^2 - v_\ell$, we obtain the formula for the new Reynolds stress

$$\begin{aligned} &\text{div}\mathring{R}_{q+1} - \nabla p_{q+1}^2 \\ &= \underbrace{-\Delta w_{q+1} + \partial_t(\tilde{w}_{q+1}^{(p)} + \tilde{w}_{q+1}^{(c)}) + \text{div}((v_\ell + v_{q+1}^1) \otimes w_{q+1} + w_{q+1} \otimes (v_\ell + v_{q+1}^1))}_{\text{div}(R_{\text{lin}}) + \nabla p_{\text{lin}}} \\ &\quad + \underbrace{\text{div}((\tilde{w}_{q+1}^{(c)} + \tilde{w}_{q+1}^{(t)}) \otimes w_{q+1} + \tilde{w}_{q+1}^{(p)} \otimes (\tilde{w}_{q+1}^{(c)} + \tilde{w}_{q+1}^{(t)}))}_{\text{div}(R_{\text{cor}}) + \nabla p_{\text{cor}}} \\ &\quad + \underbrace{\text{div}(\tilde{w}_{q+1}^{(p)} \otimes \tilde{w}_{q+1}^{(p)} + \mathring{R}_\ell) + \partial_t \tilde{w}_{q+1}^{(t)}}_{\text{div}(R_{\text{osc}}) + \nabla p_{\text{osc}}} \\ &\quad + \underbrace{\text{div}((V_{q+1}^2 - V_q^2) + (V_{q+1}^{2,*} - V_q^{2,*}))}_{\text{div}R_{\text{com1}} + \nabla p_{\text{com1}}} \\ &\quad + \underbrace{\text{div}((v_\ell + v_{q+1}^1) \otimes (v_\ell + v_{q+1}^1) - (v_q^1 + v_q^2) \otimes (v_q^1 + v_q^2))}_{\text{div}(R_{\text{com2}}) + \nabla p_{\text{com2}}} \\ &\quad + \text{div}(R_{\text{com}}) - \nabla p_\ell, \end{aligned} \tag{7.51}$$

where, using the notation $v_q = v_q^1 + v_q^2$,

$$\begin{aligned}
 &V_{q+1}^2 - V_q^2 \\
 &= (v_{q+1}^2 - v_q^2) \otimes \Delta_{>Rz} \mathcal{Y} + (v_{q+1} - v_q) \otimes \Delta_{\leq Rz} \mathcal{Y} \\
 &\quad + (v_{q+1}^2 - v_q^2) \otimes \Delta_{>Rz} + (v_{q+1} - v_q) (\otimes + \otimes) \Delta_{\leq Rz} + (v_{q+1}^2 - v_q^2) \otimes z \\
 &\quad - \mathbb{P}[(v_{q+1} - v_q) \prec \Delta_{\leq R} \mathcal{I} \nabla z] \otimes z - ([\mathbb{P}, (v_{q+1}^2 - v_q^2) \prec \mathcal{I} \nabla z] \otimes z \\
 &\quad - ([\mathbb{P}, (v_{q+1}^1 - v_q^1) \prec \mathcal{I} \nabla z] \otimes \Delta_{\leq Rz} \\
 &\quad - \text{com}(v_{q+1}^1 - v_q^1, \mathbb{P} \mathcal{I} \nabla z, \Delta_{\leq Rz}) \\
 &\quad - \text{com}(v_{q+1}^2 - v_q^2, \mathbb{P} \mathcal{I} \nabla z, z) - (v_{q+1}^2 - v_q^2) \succ \Delta_{>Rz} \mathcal{Y} \\
 &\quad - (v_{q+1} - v_q) \cdot \Delta_{\leq Rz} \mathcal{Y} + (v_{q+1}^\# - v_q^\#) \otimes \Delta_{\leq Rz}.
 \end{aligned}$$

Applying the inverse divergence operator \mathcal{R} we define

$$\begin{aligned}
 R_{\text{lin}} &:= -\mathcal{R} \Delta w_{q+1} + \mathcal{R} \partial_t (\tilde{w}_{q+1}^{(p)} + \tilde{w}_{q+1}^{(c)}) + (v_\ell + v_{q+1}^1) \\
 &\quad \otimes w_{q+1} + w_{q+1} \otimes (v_\ell + v_{q+1}^1), \\
 R_{\text{cor}} &:= (\tilde{w}_{q+1}^{(c)} + \tilde{w}_{q+1}^{(t)}) \otimes w_{q+1} + \tilde{w}_{q+1}^{(p)} \otimes (\tilde{w}_{q+1}^{(c)} + \tilde{w}_{q+1}^{(t)}),
 \end{aligned}$$

R_{com1} is the trace-free part of the matrix

$$(V_{q+1}^2 - V_q^2) + (V_{q+1}^{2,*} - V_q^{2,*}),$$

and

$$R_{\text{com2}} := (v_\ell + v_{q+1}^1) \otimes (v_\ell + v_{q+1}^1) - (v_q^2 + v_q^1) \otimes (v_q^2 + v_q^1).$$

Similarly as in [45], using (7.23), (7.26), the oscillation error is given by

$$\begin{aligned}
 R_{\text{osc}} &:= \chi^2 \sum_{\xi \in \Lambda} \mathcal{R} \left(\nabla a_{(\xi)}^2 \mathbb{P}_{\neq 0}(W_{(\xi)}) \otimes W_{(\xi)} \right) - \frac{\chi^2}{\mu} \sum_{\xi \in \Lambda} \mathcal{R} \left(\partial_t a_{(\xi)}^2 (\phi_{(\xi)}^2 \psi_{(\xi)}^2 \xi) \right) \\
 &\quad + \mathcal{R} w_{q+1}^{(t)} \partial_t \chi^2 + (1 - \chi^2) \mathring{R}_\ell \\
 &=: R_{\text{osc}}^{(x)} + R_{\text{osc}}^{(t)} + R_{\text{cut}}^{(1)}.
 \end{aligned}$$

Finally we define the Reynolds stress on the level $q + 1$ by

$$\mathring{R}_{q+1} := R_{\text{lin}} + R_{\text{cor}} + R_{\text{osc}} + R_{\text{com}} + R_{\text{com1}} + R_{\text{com2}}.$$

We observe that by construction, \mathring{R}_{q+1} is $(\mathcal{F}_t)_{t \geq 0}$ -adapted.

7.7. Verification of the Inductive Estimates for \mathring{R}_{q+1}

We shall establish the three bounds in (5.14). As the oscillations w_{q+1} were fully added for $t \in (\sigma_q \wedge T_L, T_L]$, this is the good interval where the desired smallness of \mathring{R}_{q+1} is achieved. In the middle interval $t \in (\frac{\sigma_q}{2} \wedge T_L, \sigma_q \wedge T_L]$, there is a part of \mathring{R}_{q+1} involving the cut-off $1 - \chi^2$, which can only be bounded by the previous stress \mathring{R}_q . In the time interval $t \in [0, \frac{\sigma_q}{2} \wedge T_L]$, there are no oscillations which could decrease the Reynolds stress and hence we can only prove a polynomial blow-up. Nevertheless, this eventually leads to convergence in L^p in time as this bad time interval is shrinking exponentially, cf. the proof of Theorem 5.4. Here, it is essential that we do not use regularity of v_q^1 and v_q^\sharp to avoid the blow-up in time.

Case I. Let $t \in (\sigma_q \wedge T_L, T_L]$. If $T_L \leq \sigma_q$ then there is nothing to estimate here, hence we assume that $\sigma_q < T_L$ and $t \in (\sigma_q, T_L]$. In this regime, it holds $\chi = 1$ and so the truncation does not play any role in the estimates. We estimate each term in the definition of \mathring{R}_{q+1} separately. We choose $p = \frac{32}{32-7\alpha} > 1$ so that in particular that $r_\perp^{2/p-2} r_\parallel^{1/p-1} \leq \lambda_{q+1}^\alpha$. For the linear error we apply (5.10) to obtain

$$\begin{aligned} \|\mathcal{R}_{\text{lin}}(t)\|_{L^1} &\lesssim \|\mathcal{R}\Delta w_{q+1}\|_{L^p} + \|\mathcal{R}\partial_t(w_{q+1}^{(p)} + w_{q+1}^{(c)})\|_{L^p} \\ &\quad + \|(v_\ell + v_{q+1}^1) \otimes w_{q+1} + w_{q+1} \otimes (v_\ell + v_{q+1}^1)\|_{L^1} \\ &\lesssim \|w_{q+1}\|_{W^{1,p}} + \sum_{\xi \in \Lambda} \|\partial_t \text{curl}(a(\xi) V(\xi))\|_{L^p} \\ &\quad + (M_L^{1/2} \lambda_q^4 + \|v_{q+1}^1(t)\|_{L^4}) \|w_{q+1}\|_{L^{4/3}}, \end{aligned}$$

where, by (A.7) and (7.21),

$$\begin{aligned} &\sum_{\xi \in \Lambda} \|\partial_t \text{curl}(a(\xi) V(\xi))\|_{C_t L^p} \\ &\leq \sum_{\xi \in \Lambda} \left(\|a(\xi)\|_{C_t C_x^1} \|\partial_t V(\xi)\|_{C_t W^{1,p}} + \|\partial_t a(\xi)\|_{C_t C_x^1} \|V(\xi)\|_{C_t W^{1,p}} \right) \\ &\lesssim (M_L(1 + 3q) + \gamma_{q+1})^{1/2} \ell^{-15} r_\perp^{2/p} r_\parallel^{1/p-3/2} \mu \\ &\quad + (M_L(1 + 3q) + \gamma_{q+1})^{1/2} \ell^{-22} r_\perp^{2/p-1} r_\parallel^{1/p-1/2} \lambda_{q+1}^{-1}. \end{aligned}$$

Remark 7.1. For the product $v_{q+1}^1 \otimes w_{q+1}$ we used the L^4 -norm of v_q^1 instead of L^2 in order to lower the required integrability of w_{q+1} . Indeed, w_{q+1} is not small in L^2 for $t \in (\sigma_q, T_L]$, cf. (7.37), (7.38). On the other hand, as a consequence of (6.3), increasing the integrability of v_{q+1}^1 leads to a blow-up in two respects: there is a blow up as $t \rightarrow 0$ but also the time-weighted norm in $C_{t,3/8} L^4$ has only a diverging bound as $q \rightarrow \infty$. We show below that both these divergencies are compensated by the smallness of w_{q+1} in $L^{4/3}$ and by using the fact that $t > \sigma_q$.

In view of (7.48), (7.33), (7.36) as well as (6.3) applied on the level $q + 1$, we deduce, for $t \in (\sigma_q, T_L]$,

$$\begin{aligned} \|R_{\text{lin}}(t)\|_{L^p} &\lesssim M_L^{1/2} \lambda_{q+1}^{60\alpha-1/7} + (M_L(1+3q) + \gamma_{q+1})^{1/2} \left(\ell^{-22} \lambda_{q+1}^{\alpha-1/7} \right. \\ &\quad \left. + M_L^{1/2} \ell^{-8} r_{\perp}^{1/2} r_{\parallel}^{1/4} (\lambda_q^4 + \sigma_q^{-3/8} \lambda_{q+1}^{1/3+\kappa}) \right) \\ &\lesssim M_L^{1/2} \lambda_{q+1}^{60\alpha-1/7} + (M_L(1+3q) + \gamma_{q+1})^{1/2} M_L^{1/2} \lambda_{q+1}^{16\alpha-5/21+\kappa} \sigma_q^{-3/8} \\ &\leq \frac{M_L \delta_{q+2}}{5}. \end{aligned}$$

Here, we have taken a sufficiently large and β sufficiently small.

The estimates of R_{cor} and R_{osc} are the same as the corresponding bounds in [45]. The strongest requirement comes from the bound of R_{cor} , namely, we have

$$\begin{aligned} \|R_{\text{cor}}(t)\|_{L^p} &\lesssim ((M_L(1+3q))^{1/2} + \gamma_{q+1}^{1/2})^3 \lambda_{q+1}^{49\alpha-1/7} \leq \ell^{-3/2} \lambda_{q+1}^{49\alpha-1/7} \\ &\leq \frac{M_L}{5} \lambda_{q+1}^{-2\beta b} \leq \frac{M_L}{5} \delta_{q+2}, \end{aligned}$$

which is satisfied provided $\lambda_{q+1}^{52\alpha-1/7} \leq \frac{M_L}{5} \lambda_{q+1}^{-2\beta b}$.

We use standard mollification estimates in order to bound R_{com} . More precisely, R_{com} has to vanish sufficiently fast in order to fulfill the first bound in (5.14). Hence, we need to use regularity of each term in N which then by mollification estimates leads to the desired decay. To this end, there is a number of terms which require spatial regularity of v_q^1 as well as v_q^\sharp . But these norms blow up as $t \rightarrow 0$, cf. Proposition 5.2. Thus, we make use of the fact that $t > \sigma_q$ and that the corresponding blow-up of order $\sigma_q^{-1/6}$ and $\sigma_q^{-3/10}$, respectively, can be absorbed by the smallness of ℓ due to our choice of the parameters in (7.4).

Let us now consider each term in R_{com} separately. For $I_1 := (v_q^1 + v_q^2) \otimes \Delta_{\leq RZ} \check{Y} - (v_q^1 + v_q^2) \cdot \Delta_{\leq RZ} \check{y}$ we use Proposition 5.2 and (5.7) to get that

$$\begin{aligned} &\|I_1 - I_1 *_x \phi_\ell *_t \varphi_\ell\|_{L^1} \\ &\lesssim \ell^{1/10} (\|v_q^2\|_{C_t^{1/10} L^{5/3}} \\ &\quad + \|v_q^1\|_{C_{[\sigma_q/2, t]}^{1/6-\kappa} L^{5/3}}) (\|\Delta_{\leq RZ} \check{Y}\|_{C_t^{1/10} L^\infty} + \|\Delta_{\leq RZ} \check{y}\|_{C_t^{1/10} L^\infty}) \\ &\quad + \ell^{1/10} (\|v_q^2\|_{C_t W^{1/5, 5/3}} \\ &\quad + \|v_q^1\|_{C_{[\sigma_q/2, t]} B_{5/3, \infty}^{1/6-\kappa}}) (\|\Delta_{\leq RZ} \check{Y}\|_{C_t C^{1/10}} \\ &\quad + \|\Delta_{\leq RZ} \check{y}\|_{C_t C^{1/10}}) \\ &\lesssim \ell^{1/10} \sigma_q^{-1/6} (M_L^{1/2} a^{-\alpha/4} + L^3 N + L^4) L^{10}, \end{aligned}$$

where we used

$$\begin{aligned} &\|\Delta_{\leq RZ} \check{Y}\|_{C_t C^{1/10}} + \|\Delta_{\leq RZ} \check{y}\|_{C_t^{1/10} L^\infty} \\ &\lesssim 2^{(1/5+2\kappa)R} (\|\check{z}^\check{Y}\|_{C_t^{1/10} C^{-1/5-\kappa}} + \|\check{z}^\check{y}\|_{C_t C^{-\kappa}}) \lesssim L^{10}, \end{aligned}$$

and similarly for $z^{\check{\vee}}$. The same estimate also holds for the symmetric counterpart. For $I_2 := (v_q^1 + v_q^2) \otimes (v_q^1 + v_q^2)$ we use (5.9) and (5.18) to get that

$$\begin{aligned} & \|I_2 - I_2 *_x \phi_\ell *_t \varphi_\ell\|_{L^1} \\ & \lesssim \left(\ell \|v_q^2\|_{C_{t,x}^1} + \ell^{1/61} \|v_q^1\|_{C_{[\sigma_q/2,t]}^{1/61} L^2} + \ell^{1/31} \|v_q^1\|_{C_{[\sigma_q/2,t]} H^{1/30-3\kappa}} \right) \\ & (\|v_q^1\|_{C_t L^2} + \|v_q^2\|_{C_t L^2}) \\ & \lesssim \sigma_q^{-1/6} \ell^{1/61} (M_L a^{-\alpha/8} + L^6 N^2 + L^8) \\ & \quad + (\ell \lambda_q^4 + \sigma_q^{-1/6} \ell^{1/61}) M_L^{1/2} (M_L^{1/2} (1 + 3q)^{1/2} + K^{1/2}) \\ & \lesssim M_L^{1/2} (M_L (1 + 3q) + K)^{1/2} (\ell \lambda_q^4 + \ell^{1/61} \sigma_q^{-1/6}), \end{aligned}$$

where we used interpolation and Proposition 5.2 and the embedding Lemma 2.1 to get that

$$\begin{aligned} \|v_q^1\|_{C_{t,1/6}^{1/61} L^2} & \lesssim \|v_q^1\|_{C_{t,1/6} H^{1/30-3\kappa}}^{\frac{9}{10(1-9\kappa)}} \|v_q^1\|_{C_{t,1/6}^{1/6-\kappa} H^{-3/10}}^{\frac{1-90\kappa}{10(1-9\kappa)}} \lesssim \\ & \|v_q^1\|_{C_{t,1/6} B_{5/3,\infty}^{1/3-2\kappa}}^{\frac{9}{10(1-9\kappa)}} \|v_q^1\|_{C_{t,1/6}^{1/6-\kappa} L^{5/3}}^{\frac{1-90\kappa}{10(1-9\kappa)}} \\ & \lesssim M_L^{1/2} a^{-\alpha/4} + L^3 N + L^4 \lesssim M_L^{1/2}, \end{aligned}$$

provided that κ is chosen sufficiently small. For $I_3 := v_q^2 \otimes \Delta_{>Rz} \check{\vee} - v_q^2 \otimes \Delta_{>Rz} \check{\vee}$ we use paraproduct estimates Lemma 2.2 and (5.10) to get that

$$\begin{aligned} & \|I_3 - I_3 *_x \phi_\ell *_t \varphi_\ell\|_{L^1} \\ & \lesssim \ell^{1/10} \|v_q^2\|_{C_{t,x}^1} (\|z^{\check{\vee}}\|_{C_t C^{-\kappa}} + \|z^{\check{\vee}}\|_{C_t C^{-\kappa}} + \|z^{\check{\vee}}\|_{C_t^{1/10} C^{-1/5-\kappa}} + \|z^{\check{\vee}}\|_{C_t^{1/10} C^{-1/5-\kappa}}) \\ & \lesssim \ell^{1/10} \lambda_q^4 L M_L^{1/2}. \end{aligned}$$

For $I_4 := (v_q^1 + v_q^2) (\otimes + \otimes) \Delta_{\leq Rz}$ we use paraproduct estimates Lemma 2.2 and (5.10) to get that

$$\begin{aligned} & \|I_4 - I_4 *_x \phi_\ell *_t \varphi_\ell\|_{L^1} \\ & \lesssim \ell^{1/10} (\|v_q^2\|_{C_{t,x}^1} + \|v_q^1\|_{C_{[\sigma_q/2,t]}^{1/10} B_{5/3,\infty}^{1/10}}) (\|\Delta_{\leq Rz}\|_{C_t^{1/10} L^\infty} + \|\Delta_{\leq Rz}\|_{C_t C^{1/10}}) \\ & \lesssim \ell^{1/10} L^{29} \left(M_L^{1/2} \lambda_q^4 + \sigma_q^{-1/6} (a^{-\alpha/4} M_L^{1/2} + L^3 N + L^4) \right), \end{aligned}$$

where we used

$$\|\Delta_{\leq Rz}\|_{C_t C^{1/10}} + \|\Delta_{\leq Rz}\|_{C_t^{1/10} L^\infty} \lesssim 2^{R(7/10+2\kappa)} L^{1/2} \leq L^{29},$$

and by interpolation and Proposition 5.2,

$$\|v_q^1\|_{C_{t,1/6}^{1/10} B_{5/3,\infty}^{1/10}} \lesssim \|v_q^1\|_{C_{t,1/6} B_{5/3,\infty}^{1/3-2\kappa}} + \|v_q^1\|_{C_{t,1/6}^{1/6-\kappa} L^{5/3}} \lesssim M_L^{1/2} a^{-\alpha/4} + L^3 N + L^4. \tag{7.52}$$

For $I_5 := v_q^2 \circlearrowleft \Delta_{>Rz} + v_q^2 \circledast z + v_q^\sharp \circledast \Delta_{\leq Rz}$ we use paraproduct estimates Lemma 2.2, Proposition 5.2 and (5.10) to get that

$$\begin{aligned} & \|I_5 - I_5 *_x \phi_\ell *_t \varphi_\ell\|_{L^1} \\ & \lesssim \ell^{1/24} (\|v_q^2\|_{C_{t,x}^1} + \|v_q^\sharp\|_{C_{[\sigma_q/2,t]} B_{5/3,\infty}^{3/5-\kappa}} \\ & \quad + \|v_q^\sharp\|_{C_{[\sigma_q/2,t]} B_{5/3,\infty}^{11/20-\kappa}}) (\|z\|_{C_t^{1/24} C^{-7/12-\kappa}} + \|z\|_{C_t C^{-1/2-\kappa}}) \\ & \lesssim \ell^{1/24} L M_L^{1/2} (\lambda_q^4 + \sigma_q^{-3/10} (a^{-\alpha/8} M_L^{1/2} + L^5 N + L^6)) \lesssim \ell^{1/24} M_L (\lambda_q^4 + \sigma_q^{-3/10}). \end{aligned}$$

Here we used $L \leq a^{\alpha/16}$ by the choice of the parameters and interpolation to bound $\|z\|_{C_t^{1/24} C^{-7/12-\kappa}}$ by

$$\|z\|_{C_t C^{-1/2-\kappa}} + \|z\|_{C_t^{1/10} C^{-7/10-\kappa}} \lesssim L^{1/2}.$$

For $I_6 := \mathbb{P}[v_q \prec \Delta_{\leq R\mathcal{I}(\nabla z)}] \circledast z$ we use (5.7), (5.18) and Lemma 2.2 to get that

$$\begin{aligned} & \|I_6 - I_6 *_x \phi_\ell *_t \varphi_\ell\|_{L^1} \\ & \lesssim \ell^{1/10} (\|v_q^2\|_{C_t^{1/10} L^{5/3}} + \|v_q^1\|_{C_{[\sigma_q/2,t]} L^{5/3}}) \\ & \quad \times \left(\|z\|_{C_t C^{-1/2-\kappa}} (\|\Delta_{\leq R\mathcal{I}\nabla z}\|_{C_t C^{3/5+2\kappa}} + \|\Delta_{\leq R\mathcal{I}\nabla z}\|_{C_t^{1/10} C^{1/2+2\kappa}}) \right. \\ & \quad \left. + \|z\|_{C_t^{1/10} C^{-7/10-\kappa}} \|\Delta_{\leq R\mathcal{I}\nabla z}\|_{C_t C^{7/10+2\kappa}} \right) \\ & \lesssim \ell^{1/10} L^{10} \sigma_q^{-1/6} (M_L^{1/2} a^{-\alpha/4} + L^3 N + L^4) \lesssim \ell^{1/10} M_L \sigma_q^{-1/6}. \end{aligned}$$

Here we used

$$\|\Delta_{\leq R\mathcal{I}\nabla z}\|_{C_t C^{7/10+2\kappa}} + \|\Delta_{\leq R\mathcal{I}\nabla z}\|_{C_t^{1/10} C^{1/2+2\kappa}} \lesssim 2^{R(1/5+3\kappa)} \|z\|_{C_t C^{-1/2-\kappa}} \lesssim L^9.$$

In view of the two commutator estimates, Lemma 2.3 and Lemma 2.4, the remaining terms containing v_q^2 could be controlled by

$$\ell^{1/10} \|v_q^2\|_{C_{t,x}^1} (\|z\|_{C_t C^{-1/2-\kappa}}^2 + \|z\|_{C_t C^{-1/2-\kappa}} \|z\|_{C_t^{1/10} C^{-7/10-\kappa}}) \lesssim \ell^{1/10} \lambda_q^4 M_L^{1/2} L,$$

and the remaining terms containing v_q^1 could be controlled by

$$\begin{aligned} & \ell^{1/10} \|v_q^1\|_{C_{[\sigma_q/2,t]} B_{5/3,\infty}^{1/10}} \\ & \left(\|z\|_{C_t C^{-1/2-\kappa}} \|\Delta_{\leq Rz}\|_{C_t C^{-1/2+1/5+2\kappa}} + \|z\|_{C_t C^{-1/2-\kappa}} \|\Delta_{\leq Rz}\|_{C_t^{1/10} C^{-1/2+2\kappa}} \right) \\ & \lesssim \ell^{1/10} L^{10} \sigma_q^{-1/6} (M_L^{1/2} a^{-\alpha/4} + L^3 N + L^4) \lesssim \ell^{1/10} M_L \sigma_q^{-1/6}. \end{aligned}$$

Here we used (5.18) and (7.52) and

$$\begin{aligned} & \|\Delta_{\leq Rz}\|_{C_t C^{-1/2+1/5+2\kappa}} + \|\Delta_{\leq Rz}\|_{C_t^{1/10} C^{-1/2+2\kappa}} \\ & \lesssim 2^{R(1/5+3\kappa)} (\|z\|_{C_t C^{-1/2-\kappa}} + \|z\|_{C_t^{1/10} C^{-7/10-\kappa}}) \lesssim L^9. \end{aligned}$$

Therefore, we have, for $t \in (\sigma_q, T_L]$, that

$$\begin{aligned} \|R_{\text{com}}(t)\|_{L^1} &\lesssim M_L^{1/2}(\ell^{1/61}\sigma_q^{-1/6} + \ell\lambda_q^4)(M_L(1 + 3q) + K)^{1/2} \\ &\quad + \ell^{1/24}M_L\lambda_q^4 + \ell^{1/24}M_L\sigma_q^{-3/10} \\ &\leq \frac{M_L\delta_{q+2}}{5}, \end{aligned}$$

where we used the choice of ℓ in (7.2) and the conditions

$$\alpha > 244\beta b, \quad \alpha b > 128, \quad (M_L(1 + 3q) + K)^{1/2} \leq \ell^{-1/183}, \quad \sigma_q^{-1/3} < \ell^{-1/183},$$

which can indeed be achieved by our conditions on the parameters.

Next, by the choice of α, β and b using paraproduct estimates from Lemma 2.2 we can bound $R_{\text{com}1}$ uniformly over the interval $[0, t]$ for $p = \frac{32}{32-7\alpha}$ to get that

$$\begin{aligned} \|R_{\text{com}1}\|_{C_t L^1} &\lesssim (\|v_{q+1}^2 - v_q^2\|_{C_t W^{2/3,p}} + \|v_{q+1}^1 - v_q^1\|_{C_t B_{5/3,\infty}^{1/3-2\kappa}} \\ &\quad + \|v_{q+1}^\sharp - v_q^\sharp\|_{C_t L^{5/3}})L^{21} \\ &\lesssim L^{21}M_L^{1/2}(\lambda_q^{-1/42} + \lambda_{q+1}^{-15\alpha/32} + \lambda_{q+1}^{60\alpha-1/7} + \lambda_{q+1}^{32\alpha-1/35}) \\ &\leq \frac{M_L\delta_{q+2}}{10}, \end{aligned}$$

where we used (7.8) and (7.48) to control $\|v_{q+1}^2 - v_q^2\|_{C_t W^{2/3,p}}$, (6.1), (7.6) and (7.42) to control $\|v_{q+1}^1 - v_q^1\|_{C_t B_{5/3,\infty}^{1/3-2\kappa}}$ and also (6.2), (7.6) and (7.42) to control $\|v_{q+1}^\sharp - v_q^\sharp\|_{C_t L^{5/3}}$. Moreover, we applied $1 > 168\beta b^2$ and $-32\alpha + \frac{1}{35} > 2\beta b$, the condition on α, β, b and

$$\|\Delta_{\leq R} z\|_{C_t C^\kappa} \lesssim 2^{R(1/2+2\kappa)} \|z\|_{C_t C^{-1/2-\kappa}} \lesssim L^{21}. \tag{7.53}$$

Furthermore, we use Proposition 5.2, (6.1), (5.9) and (7.42) to estimate also $R_{\text{com}2}$ uniformly over $[0, t]$ to obtain

$$\begin{aligned} \|R_{\text{com}2}\|_{C_t L^1} &\lesssim (M_L(1 + 3q) + K)^{1/2}(\|v_{q+1}^1 - v_q^1\|_{C_t L^2} + \|v_\ell - v_q^2\|_{C_t L^2}) \\ &\lesssim (M_L(1 + 3q) + K)^{1/2}M_L^{1/2}(\lambda_q^{-1/42} + \lambda_{q+1}^{-\alpha} + \lambda_{q+1}^{32\alpha-1/35}) \\ &\leq \frac{M_L\delta_{q+2}}{10}, \end{aligned}$$

Here we used $\alpha > 244\beta b, 1 > 168\beta b^2$ and $(M_L(1 + 3q) + K)^{1/2} \leq \lambda_q^{1/84} < \ell^{-1/183}$.

Remark 7.2. We note that it was essential in the estimate of $R_{\text{com}1}$ and $R_{\text{com}2}$ that no time weights were required for the difference $v_{q+1}^1 - v_q^1$ and $v_{q+1}^\sharp - v_q^\sharp$. Indeed, there would be no way to absorb the weight as time approaches zero. We also note that the bounds of $R_{\text{com}1}$ and $R_{\text{com}2}$ hold directly for all $t \in [0, T_L]$.

Summarizing all the above estimates we obtain

$$\|\mathring{R}_{q+1}(t)\|_{L^1} \leq M_L \delta_{q+2},$$

which is the desired bound.

Case II. Let $t \in (\frac{\sigma_q}{2} \wedge T_L, \sigma_q \wedge T_L]$. If $T_L \leq \frac{\sigma_q}{2}$ then there is nothing to estimate, hence we may assume $\frac{\sigma_q}{2} < T_L$ and $t \in (\frac{\sigma_q}{2}, \sigma_q \wedge T_L]$. Then we decompose $\mathring{R}_\ell = \chi^2 \mathring{R}_\ell + (1 - \chi^2) \mathring{R}_\ell$. The first part $\chi^2 \mathring{R}_\ell$ is canceled (up to the oscillation error $\chi^2 R_{\text{osc}}$) by $\tilde{w}_{q+1}^{(p)} \otimes \tilde{w}_{q+1}^{(p)} = \chi^2 w_{q+1}^{(p)} \otimes w_{q+1}^{(p)}$ and $\chi^2 \partial_t w_{q+1}^{(t)} = \partial_t \tilde{w}_{q+1}^{(t)} - (\chi^2)' w_{q+1}^{(t)}$. Thus, in this case, in the definition of \mathring{R}_{q+1} most terms are similar to **Case I.** and can be estimated similarly as above. Therefore, we only have to consider $(1 - \chi^2) \mathring{R}_\ell$, and

$$\text{div } R_{\text{cut}} := \chi'(t)(w_{q+1}^{(p)}(t) + w_{q+1}^{(c)}(t)) + (\chi^2)'(t)w_{q+1}^{(t)}(t).$$

We know that

$$\|(1 - \chi^2) \mathring{R}_\ell(t)\|_{L^1} \leq \sup_{s \in [t-\ell, t]} \|\mathring{R}_q(s)\|_{L^1}.$$

For R_{cut} we realize that the bounds (7.33) and (7.36) also hold for $w_{q+1}^{(p)}, w_{q+1}^{(c)}$ and $w_{q+1}^{(t)}$. Then we have, for $\kappa > 0$,

$$\begin{aligned} \|R_{\text{cut}}(t)\|_{L^1} &\leq \|\chi'(t)(w_{q+1}^{(p)}(t) + w_{q+1}^{(c)}(t))\|_{L^1} + \|(\chi^2)'(t)w_{q+1}^{(t)}(t)\|_{L^1} \\ &\lesssim \frac{1}{\sigma_q} \|w_{q+1}^{(p)}(t)\|_{L^1} + \frac{1}{\sigma_q} \|w_{q+1}^{(c)}(t)\|_{L^1} + \frac{1}{\sigma_q} \|w_{q+1}^{(t)}(t)\|_{L^1} \\ &\leq \frac{1}{\sigma_q} \ell^{-8} ((M_L + 3qM_L)^{1/2} + \gamma_{q+1}^{1/2}) r_\perp^{1-2\kappa} r_\parallel^{1/2-\kappa} \leq \frac{M_L}{10} \delta_{q+2}, \end{aligned}$$

where we use $\sigma_q^{-1} \leq \ell^{-1}$ and $M_L^{1/2}(1 + 3q)^{1/2} + \gamma_{q+1}^{1/2} \leq \ell^{-1}$. For R_{com} , where the required space regularity of v_q^1 and v_q^\sharp leads to a blow-up in time, we again use the fact that $t \geq \sigma_q/2$ and $4\ell \leq \sigma_q$ to have a bound similar to the first case.

Case III. For $t \in [0, \frac{\sigma_q}{2} \wedge T_L]$ we know $v_\ell(t) = v_{q+1}^2(t) = 0$ and so the Reynolds stress reduces to

$$\mathring{R}_{q+1} = \mathring{R}_\ell + R_{\text{com}1} + R_{\text{com}} + R_{\text{com}2}.$$

The bounds for $R_{\text{com}1}$ and $R_{\text{com}2}$ hold as in **Case I.** Unlike **Case I.** and **Case II.**, here we cannot use regularity of v_q^1 due to the blow-up at $t = 0$ as t is no longer bounded away from zero. On the other hand, we do not have to show smallness of R_{com} but only a polynomial blow-up, see (5.14). Therefore, we do not try to use the mollification estimates. Instead, we bound $\|R_{\text{com}}(t)\|_{L^1}$ directly using Lemma 2.2 and (7.53), (5.18), (5.19) as

$$\begin{aligned} \|R_{\text{com}}(t)\|_{L^1} &\lesssim \|v_q^1\|_{C_t L^2} (\|\Delta_{\leq R} z^\vee\|_{C_t L^\infty} + \|\Delta_{\leq R} z^\flat\|_{C_t L^\infty} + \|\Delta_{\leq R} z\|_{C_t C^\kappa}) \\ &\quad + \|v_q^1\|_{C_t L^2}^2 + \|v_q^\sharp\|_{C_t L^{5/3}} \|\Delta_{\leq R} z\|_{C_t C^\kappa} \end{aligned}$$

$$\begin{aligned}
 &+ \|v_q^1\|_{C_t L^2} \|z\|_{C_t C^{-1/2-\kappa}} \|\Delta_{\leq R} z\|_{C_t C^{-1/2+2\kappa}} \\
 &\leq 2M_L,
 \end{aligned}
 \tag{7.54}$$

where we used (7.53) and the implicit constant can be absorbed by taking a and L large enough.

Remark 7.3. We point out that for the two commutators in R_{com} we did not apply the commutator estimates Lemmas 2.3 and 2.4, as these would require regularity of v_q^1 . Instead, we estimated each term by the paraproduct estimates directly.

As a result, it follows that

$$\|\mathring{R}_{q+1}(t)\|_{L^1} \leq \sup_{s \in [t-\ell, t]} \|\mathring{R}_q(s)\|_{L^1} + 3M_L,$$

which completes the proof of (5.14) and also (5.11) and (5.12).

Data Availability Data sharing not applicable to this article as no datasets were generated or analyzed during the current study.

Declarations

Conflict of interest The authors have no competing interests to declare that are relevant to the content of this article.

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.

Appendix A: Intermittent Jets

In this part we recall the construction of intermittent jets from [9, Section 7.4]. We point out that the construction is entirely deterministic, that is, none of the functions below depends on ω . Let us begin with the following geometric lemma which can be found in [9, Lemma 6.6]:

Lemma A.1. *Denote by $\overline{B_{1/2}}(\text{Id})$ the closed ball of radius 1/2 around the identity matrix Id , in the space of 3×3 symmetric matrices. There exists $\Lambda \subset \mathbb{S}^2 \cap \mathbb{Q}^3$ such that for each $\xi \in \Lambda$ there exists a C^∞ -function $\gamma_\xi : \overline{B_{1/2}}(\text{Id}) \rightarrow \mathbb{R}$ such that*

$$R = \sum_{\xi \in \Lambda} \gamma_\xi^2(R) (\xi \otimes \xi)$$

for every symmetric matrix satisfying $|R - \text{Id}| \leq 1/2$. For $C_\Lambda = 8|\Lambda|(1 + 8\pi^3)^{1/2}$, where $|\Lambda|$ is the cardinality of the set Λ , we define the constant

$$M = C_\Lambda \sup_{\xi \in \Lambda} (\|\gamma_\xi\|_{C^0} + \sum_{|j| \leq N} \|D^j \gamma_\xi\|_{C^0}).$$

For each $\xi \in \Lambda$ let us define $A_\xi \in \mathbb{S}^2 \cap \mathbb{Q}^3$ to be an orthogonal vector to ξ . Then for each $\xi \in \Lambda$ we have that $\{\xi, A_\xi, \xi \times A_\xi\} \subset \mathbb{S}^2 \cap \mathbb{Q}^3$ form an orthonormal basis for \mathbb{R}^3 . We label by n_* the smallest natural such that

$$\{n_*\xi, n_*A_\xi, n_*\xi \times A_\xi\} \subset \mathbb{Z}^3$$

for every $\xi \in \Lambda$.

Let $\Phi : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a smooth function with support in a ball of radius 1. We normalize Φ such that $\phi = -\Delta\Phi$ obeys

$$\frac{1}{4\pi^2} \int_{\mathbb{R}^2} \phi^2(x_1, x_2) dx_1 dx_2 = 1. \tag{A.1}$$

By definition we know $\int_{\mathbb{R}^2} \phi dx = 0$. Define $\psi : \mathbb{R} \rightarrow \mathbb{R}$ to be a smooth, mean zero function with support in the ball of radius 1 satisfying

$$\frac{1}{2\pi} \int_{\mathbb{R}} \psi^2(x_3) dx_3 = 1. \tag{A.2}$$

For parameters $r_\perp, r_\parallel > 0$ such that

$$r_\perp \ll r_\parallel \ll 1,$$

we define the rescaled cut-off functions

$$\begin{aligned} \phi_{r_\perp}(x_1, x_2) &= \frac{1}{r_\perp} \phi\left(\frac{x_1}{r_\perp}, \frac{x_2}{r_\perp}\right), & \Phi_{r_\perp}(x_1, x_2) &= \frac{1}{r_\perp} \Phi\left(\frac{x_1}{r_\perp}, \frac{x_2}{r_\perp}\right), \\ \psi_{r_\parallel}(x_3) &= \frac{1}{r_\parallel^{1/2}} \psi\left(\frac{x_3}{r_\parallel}\right). \end{aligned}$$

We periodize $\phi_{r_\perp}, \Phi_{r_\perp}$ and ψ_{r_\parallel} so that they are viewed as periodic functions on $\mathbb{T}^2, \mathbb{T}^2$ and \mathbb{T} respectively.

Consider a large real number λ such that $\lambda r_\perp \in \mathbb{N}$, and a large time oscillation parameter $\mu > 0$. For every $\xi \in \Lambda$ we introduce

$$\begin{aligned} \psi_{(\xi)}(t, x) &:= \psi_{\xi, r_\perp, r_\parallel, \lambda, \mu}(t, x) := \psi_{r_\parallel}(n_* r_\perp \lambda(x \cdot \xi + \mu t)) \\ \Phi_{(\xi)}(x) &:= \Phi_{\xi, r_\perp, \lambda}(x) := \Phi_{r_\perp}(n_* r_\perp \lambda(x - \alpha_\xi) \cdot A_\xi, n_* r_\perp \lambda(x - \alpha_\xi) \cdot (\xi \times A_\xi)) \\ \phi_{(\xi)}(x) &:= \phi_{\xi, r_\perp, \lambda}(x) := \phi_{r_\perp}(n_* r_\perp \lambda(x - \alpha_\xi) \cdot A_\xi, n_* r_\perp \lambda(x - \alpha_\xi) \cdot (\xi \times A_\xi)), \end{aligned}$$

where $\alpha_\xi \in \mathbb{R}^3$ are shifts to ensure that $\{\Phi_{(\xi)}\}_{\xi \in \Lambda}$ have mutually disjoint support. The intermittent jets $W_{(\xi)} : \mathbb{T}^3 \times \mathbb{R} \rightarrow \mathbb{R}^3$ are defined as in [9, Section 7.4].

$$W_{(\xi)}(t, x) := W_{\xi, r_\perp, r_\parallel, \lambda, \mu}(t, x) := \xi \psi_{(\xi)}(t, x) \phi_{(\xi)}(x). \tag{A.3}$$

By the choice of α_ξ we have that

$$W_{(\xi)} \otimes W_{(\xi')} \equiv 0, \text{ for } \xi \neq \xi' \in \Lambda, \tag{A.4}$$

and by the normalizations (A.2) we obtain

$$\frac{1}{(2\pi)^3} \int_{\mathbb{T}^3} W_{(\xi)}(t, x) \otimes W_{(\xi)}(t, x) dx = \xi \otimes \xi.$$

These facts combined with Lemma A.1 imply that

$$\sum_{\xi \in \Lambda} \gamma_\xi^2(R) \frac{1}{(2\pi)^3} \int_{\mathbb{T}^3} W_{(\xi)}(t, x) \otimes W_{(\xi)}(t, x) dx = R, \tag{A.5}$$

for every symmetric matrix R satisfying $|R - \text{Id}| \leq 1/2$. Since $W_{(\xi)}$ are not divergence free, we introduce the corrector term

$$W_{(\xi)}^{(c)} := \frac{1}{n_*^2 \lambda^2} \nabla \psi_{(\xi)} \times \text{curl}(\Phi_{(\xi)} \xi) = \text{curl} \text{ curl } V_{(\xi)} - W_{(\xi)}. \tag{A.6}$$

with

$$V_{(\xi)}(t, x) := \frac{1}{n_*^2 \lambda^2} \xi \psi_{(\xi)}(t, x) \Phi_{(\xi)}(x).$$

Thus we have

$$\text{div} \left(W_{(\xi)} + W_{(\xi)}^{(c)} \right) \equiv 0.$$

Next, we recall the key bounds from [9, Section 7.4]. For $N, M \geq 0$ and $p \in [1, \infty]$ it holds, provided that $r_{\parallel}^{-1} \ll r_{\perp}^{-1} \ll \lambda$, that

$$\begin{aligned} & \|\nabla^N \partial_t^M \psi_{(\xi)}\|_{C_t L^p} \\ & \lesssim r_{\parallel}^{1/p-1/2} \left(\frac{r_{\perp} \lambda}{r_{\parallel}} \right)^N \left(\frac{r_{\perp} \lambda \mu}{r_{\parallel}} \right)^M, \\ & \|\nabla^N \phi_{(\xi)}\|_{L^p} + \|\nabla^N \Phi_{(\xi)}\|_{L^p} \lesssim r_{\perp}^{2/p-1} \lambda^N, \\ & \|\nabla^N \partial_t^M W_{(\xi)}\|_{C_t L^p} + \frac{r_{\parallel}}{r_{\perp}} \|\nabla^N \partial_t^M W_{(\xi)}^{(c)}\|_{C_t L^p} + \lambda^2 \|\nabla^N \partial_t^M V_{(\xi)}\|_{C_t L^p} \\ & \lesssim r_{\perp}^{2/p-1} r_{\parallel}^{1/p-1/2} \lambda^N \left(\frac{r_{\perp} \lambda \mu}{r_{\parallel}} \right)^M, \end{aligned} \tag{A.7}$$

where the implicit constants may depend on p, N and M , but are independent of $\lambda, r_{\perp}, r_{\parallel}, \mu$.

Appendix B: Estimates for the Heat Operator

To deal with the singularity at zero we introduce the following two norms: for $\alpha \in (0, 1)$, $p \in [1, \infty]$, $\gamma \geq 0$

$$\begin{aligned} \|f\|_{C_{T,\gamma}^\alpha L^p} &:= \sup_{0 \leq t \leq T} t^\gamma \|f(t)\|_{L^p} + \sup_{0 \leq s < t \leq T} s^\gamma \frac{\|f(t) - f(s)\|_{L^p}}{|t - s|^\alpha}, \\ \|f\|_{C_{T,\gamma} B_{p,\infty}^\alpha} &:= \sup_{0 \leq t \leq T} t^\gamma \|f(t)\|_{B_{p,\infty}^\alpha}. \end{aligned}$$

First, we recall the basic estimates for the heat semigroup $P_t := e^{t\Delta}$ from [63, Lemma 2.8]. Let $T \geq 1$.

Lemma B.1. *For any $\theta > 0$ and $\alpha \in \mathbb{R}$, there is a constant $C = C(\alpha, \theta) > 0$ such that, for $p, q \in [1, \infty]$ and all $t \in (0, T]$,*

$$\|P_t f\|_{B_{p,q}^{\theta+\alpha}} \lesssim_C T^{\theta/2} t^{-\theta/2} \|f\|_{B_{p,q}^\alpha}. \tag{B.1}$$

For any $0 < \theta < 2$, there is a constant $C = C(\theta) > 0$ such that, for all $t \in [0, 1]$,

$$\|P_t f - f\|_{L^p} \lesssim_C t^{\theta/2} \|f\|_{B_{p,\infty}^\theta}. \tag{B.2}$$

Then we prove the following for $\mathcal{I}f = \int_0^\cdot P_{t-s} f \, ds$:

Lemma B.2. *Let $\alpha \in (0, 2)$, $\beta \in (\alpha - 2, 0)$, $\gamma, \delta \in [0, 1)$, $p \in [1, \infty]$ so that*

$$\gamma - \delta - \alpha/2 + \beta/2 + 1 > 0.$$

Then

$$\|\mathcal{I}f\|_{C_{T,\gamma}^{\alpha/2} L^p} + \|\mathcal{I}f\|_{C_{T,\gamma} B_{p,\infty}^\alpha} \lesssim T^{\gamma-\delta+1} \|f\|_{C_{T,\delta} B_{p,\infty}^\beta}.$$

Proof. By Lemma B.1, we have for $0 \leq t \leq T$

$$t^\gamma \|\mathcal{I}f(t)\|_{B_{p,\infty}^\alpha} \lesssim T^{(\alpha-\beta)/2} t^\gamma \int_0^t (t-s)^{-(\alpha-\beta)/2} s^{-\delta} \|f\|_{C_{T,\delta} B_{p,\infty}^\beta},$$

where

$$\begin{aligned} & T^{(\alpha-\beta)/2} t^\gamma \int_0^t (t-s)^{-(\alpha-\beta)/2} s^{-\delta} \, ds \\ &= T^{(\alpha-\beta)/2} t^\gamma \int_0^{t/2} (t-s)^{-(\alpha-\beta)/2} s^{-\delta} \, ds + T^{(\alpha-\beta)/2} t^\gamma \\ & \int_{t/2}^t (t-s)^{-(\alpha-\beta)/2} s^{-\delta} \, ds \\ & \lesssim T^{(\alpha-\beta)/2} t^{\gamma-(\alpha-\beta)/2} \int_0^{t/2} s^{-\delta} \, ds + T^{(\alpha-\beta)/2} t^{\gamma-\delta} \int_{t/2}^t (t-s)^{-(\alpha-\beta)/2} \, ds \\ & \lesssim T^{(\alpha-\beta)/2} t^{\gamma-\delta-(\alpha-\beta)/2+1} \leq T^{\gamma-\delta+1}, \end{aligned}$$

provided that $\gamma - \delta - (\alpha - \beta)/2 + 1 \geq 0$.

To bound the other norm, we use Lemma B.1 again to get for $0 \leq s \leq t \leq T$, $|t - s| \leq 1$, $\kappa > 0$ small enough

$$\begin{aligned} s^\gamma \|\mathcal{I}f(t) - \mathcal{I}f(s)\|_{L^p} &\lesssim s^\gamma \int_s^t \|P_{t-r} f(r)\|_{B_{p,\infty}^\kappa} \, dr + s^\gamma \|(P_{t-s} - I)\mathcal{I}f(s)\|_{L^p} \\ &\lesssim s^\gamma \int_s^t (t-r)^{-(\kappa-\beta)/2} r^{-\delta} \, dr \|f\|_{C_{T,\delta} B_{p,\infty}^\beta} \\ &\quad + s^\gamma (t-s)^{\alpha/2} \|\mathcal{I}f(s)\|_{B_{p,\infty}^\alpha}. \end{aligned}$$

For the first term, we write $r = s + x(t - s)$ and obtain, if $\gamma \geq \delta$, that

$$\begin{aligned} s^\gamma \int_s^t (t-r)^{-(\kappa-\beta)/2} r^{-\delta} \, dr &\leq s^{\gamma-\delta} \int_s^t (t-r)^{-(\kappa-\beta)/2} \, dr \\ &= s^{\gamma-\delta} (t-s)^{1-\kappa/2+\beta/2} \int_0^1 (1-x)^{-(\kappa-\beta)/2} \, dx \lesssim T^{\gamma-\delta} (t-s)^{\alpha/2} \end{aligned}$$

since $\beta/2 + 1 > \alpha/2$; whereas if $\gamma < \delta$ we get

$$\begin{aligned} s^\gamma \int_s^t (t-r)^{-(\kappa-\beta)/2} r^{-\delta} \, dr &= s^\gamma (t-s)^{1-\kappa/2+\beta/2} \\ &\quad \int_0^1 (1-x)^{-(\kappa-\beta)/2} (s+x(t-s))^{-\delta+\gamma} (s+x(t-s))^{-\gamma} \, dx \\ &\leq (t-s)^{1-\kappa/2+\beta/2} \int_0^1 (1-x)^{-(\kappa-\beta)/2} (s+x(t-s))^{-\delta+\gamma} \, dx \\ &\leq (t-s)^{1-\kappa/2+\beta/2-\delta+\gamma} \int_0^1 (1-x)^{-(\kappa-\beta)/2} x^{-\delta+\gamma} \, dx \\ &\lesssim (t-s)^{\alpha/2}, \end{aligned}$$

provided that $-\alpha/2 + 1 - \delta + \gamma + \beta/2 > 0$,

For the second term we use the previous estimate. Therefore, for $|t - s| \leq 1$,

$$s^\gamma \|\mathcal{I}f(t) - \mathcal{I}f(s)\|_{L^p} \lesssim (t-s)^{\alpha/2} T^{1-\delta+\gamma} \|f\|_{C_{T,\delta} B_{p,\infty}^\alpha}.$$

Since $B_{p,\infty}^\alpha \subset L^p$, we estimate the remaining term in the Hölder norm in time using the first estimate above

$$t^\gamma \|\mathcal{I}f(t)\|_{L^p} \lesssim t^\gamma \|\mathcal{I}f(t)\|_{B_{p,\infty}^\alpha} \lesssim T^{\gamma-\delta+1} \|f\|_{C_{T,\delta} B_{p,\infty}^\beta}$$

and the proof is complete. □

Remark B.3. We observe that in Sect. 6, we need to use Lemma B.2 for various combinations of $\gamma, \delta \in \{0, 1/6, 3/10\}$. As a consequence, the power of T in the statement of Lemma B.2 is always bounded by 2. Since due to the definition of the stopping time (5.5) we have $T \leq L^{1/2}$ we obtain a factor L for any application of the Schauder estimate.

Lemma B.4. *Let $\alpha \in (0, 1)$, $p \in [1, \infty]$, $\gamma \in (0, 1)$, $\beta \in \mathbb{R}$. Then for any $\kappa > 0$, $T \geq 1$*

$$\begin{aligned} & \|[\mathcal{I}, f \prec]g\|_{C_{T,\gamma}^{\alpha/4} B_{p,\infty}^{\alpha/2+\beta+2-\kappa}} + \|[\mathcal{I}, f \prec]g\|_{C_{T,\gamma} B_{p,\infty}^{\alpha+\beta+2-\kappa}} \\ & \lesssim T^{1+\alpha/2} (\|f\|_{C_{T,\gamma}^{\alpha/2} L^p} + \|f\|_{C_{T,\gamma} B_{p,\infty}^{\alpha}}) \|g\|_{C_T C^\beta}. \end{aligned}$$

Proof. We have

$$\begin{aligned} [\mathcal{I}, f \prec]g(t) &= \int_0^t e^{(t-s)\Delta} [(f(s) - f(t)) \prec g(s)] ds \\ & \quad + \int_0^t [e^{(t-s)\Delta}, f(t) \prec]g(s) ds = I_1(t) + I_2(t). \end{aligned}$$

Then, by (B.1), we have

$$\begin{aligned} t^\gamma \|I_1(t)\|_{B_{p,\infty}^{\alpha+\beta+2-\kappa}} &\lesssim t^\gamma T^{1-\kappa/2+\alpha/2} \int_0^t (t-s)^{-1+\kappa/2} s^{-\gamma} ds \|f\|_{C_{T,\gamma}^{\alpha/2} L^p} \|g\|_{C_T C^\beta} \\ &\lesssim T^{1+\alpha/2} \|f\|_{C_{T,\gamma}^{\alpha/2} L^p} \|g\|_{C_T C^\beta}, \end{aligned}$$

and by [13, Lemma A.1] (see also [53]),

$$\begin{aligned} t^\gamma \|I_2(t)\|_{B_{p,\infty}^{\alpha+\beta+2-\kappa}} &\lesssim t^\gamma \int_0^t (t-s)^{-1+\kappa/2} ds \|f(t)\|_{B_{p,\infty}^{\alpha}} \|g\|_{C_T C^\beta} \\ &\lesssim T^\kappa \|f\|_{C_{T,\gamma} B_{p,\infty}^{\alpha}} \|g\|_{C_T C^\beta}. \end{aligned}$$

Moreover, for $0 \leq t_1 < t_2 \leq T$ satisfying $|t_2 - t_1| \leq 1$,

$$\begin{aligned} I_1(t_2) - I_1(t_1) &= \int_0^{t_1} \left(e^{(t_2-s)\Delta} [(f(s) - f(t_2)) \prec g(s)] \right. \\ & \quad \left. - e^{(t_1-s)\Delta} [(f(s) - f(t_1)) \prec g(s)] \right) ds \\ & \quad + \int_{t_1}^{t_2} e^{(t_2-s)\Delta} [(f(s) - f(t_2)) \prec g(s)] ds, \end{aligned}$$

which, by Lemma B.1, implies that

$$\begin{aligned} & t_1^\gamma \|I_1(t_2) - I_1(t_1)\|_{B_{p,\infty}^{\alpha/2+\beta+2-\kappa}} \\ & \lesssim T^{1-\kappa/2+\alpha/2} \|f\|_{C_{T,\gamma}^{\alpha/2} L^p} \|g\|_{C_T C^\beta} \\ & \quad \times t_1^\gamma \left(|t_1 - t_2|^{\alpha/4} \int_0^{t_1} (t_1 - s)^{-1+\kappa/2} s^{-\gamma} ds + \int_{t_1}^{t_2} (t_2 - s)^{-1+\alpha/4+\kappa/2} s^{-\gamma} ds \right) \\ & \lesssim T^{1+\alpha/2} \|f\|_{C_{T,\gamma}^{\alpha/2} L^p} \|g\|_{C_T C^\beta} |t_1 - t_2|^{\alpha/4}. \end{aligned}$$

Also, it holds that

$$I_2(t_2) - I_2(t_1)$$

$$\begin{aligned}
 &= \int_0^{t_1} [e^{(t_2-s)\Delta} - e^{(t_1-s)\Delta}, f(t_1) \prec]g(s)ds \\
 &+ \int_0^{t_1} [e^{(t_2-s)\Delta}, (f(t_2) - f(t_1)) \prec]g(s)ds \\
 &+ \int_{t_1}^{t_2} [e^{(t_2-s)\Delta}, f(t_2) \prec]g(s)ds = J_1 + J_2 + J_3.
 \end{aligned}$$

Then we obtain, by [13, Lemma A.1] (see also [53]),

$$\begin{aligned}
 &t_1^\gamma \|J_2 + J_3\|_{B_{p,\infty}^{\alpha/2+\beta+2-\kappa}} \\
 &\lesssim (\|f\|_{C_{T,\gamma}^{\alpha/4} B_{p,\infty}^{\alpha/2}} + \|f\|_{C_{T,\gamma} B_{p,\infty}^\alpha}) \|g\|_{C_T C^\beta} \\
 &\quad \times \left(|t_1 - t_2|^{\alpha/4} \int_0^{t_1} (t_1 - s)^{-1+\kappa/2} ds + \int_{t_1}^{t_2} (t_2 - s)^{-1+\alpha/4+\kappa} ds \right) \\
 &\lesssim (\|f\|_{C_{T,\gamma}^{\alpha/2} L^p} + \|f\|_{C_{T,\gamma} B_{p,\infty}^\alpha}) \|g\|_{C_T C^\beta} |t_1 - t_2|^{\alpha/4} T^\kappa.
 \end{aligned}$$

Since $D^m((1 - e^{-2^{2j}(t_2-t_1)|\xi|^2})e^{-2^{2j}(t_1-s)|\xi|^2}) \lesssim (|t_2 - t_1|2^{2j})^\delta e^{-c2^{2j}(t_1-s)}$ for $m \in \mathbb{N}_0$, $\delta \in (0, 1)$ and ξ in an annulus, by similar argument as [13, Lemma A.1] (see also [53]) we obtain

$$\begin{aligned}
 &\| [e^{(t_2-s)\Delta} - e^{(t_1-s)\Delta}, f(t_1) \prec]g(s) \|_{B_{p,\infty}^{\alpha/2+\beta+2-\kappa}} \\
 &\lesssim |t_2 - t_1|^{\alpha/4} \|f(t_1)\|_{B_{p,\infty}^\alpha} (t_1 - s)^{-1+\kappa/2} \|g(s)\|_{C^\beta}.
 \end{aligned}$$

This implies

$$\begin{aligned}
 &t_1^\gamma \|J_1\|_{B_{p,\infty}^{\alpha/2+\beta+2-\kappa}} \lesssim \|f\|_{C_{T,\gamma} B_{p,\infty}^\alpha} \|g\|_{C_T C^\beta} |t_1 - t_2|^{\alpha/4} \int_0^{t_1} (t_1 - r)^{-1+\kappa/2} dr \\
 &\lesssim \|f\|_{C_{T,\gamma} B_{p,\infty}^\alpha} \|g\|_{C_T C^\beta} |t_1 - t_2|^{\alpha/4} T^\kappa,
 \end{aligned}$$

and the proof is complete. □

References

1. ALBRITTON, D., BRUÉ, E., COLOMBO, M.: Non-uniqueness of Leray solutions of the forced Navier–Stokes equations. [arXiv:2112.03116](https://arxiv.org/abs/2112.03116) (2021)
2. BRUNED, Y., CHANDRA, A., CHEVYREV, I., HAIRER, M.: Renormalising SPDEs in regularity structures. *J. Eur. Math. Soc.* **23**, 869–947, 2021
3. BUCKMASTER, T., COLOMBO, M., VICOL, V.: Wild solutions of the navier-stokes equations whose singular sets in time have hausdorff dimension strictly less than 1. [arXiv:1809.00600](https://arxiv.org/abs/1809.00600) (2018)
4. BUCKMASTER, T., DE LELLIS, C., SZÉKELYHIDI, L., JR., VICOL, V.: Onsager’s conjecture for admissible weak solutions. *Commun. Pure Appl. Math.* **72**(2), 229–274, 2019
5. BRUNED, Y., HAIRER, M., ZAMBOTTI, L.: Algebraic renormalisation of regularity structures. *Invent. Math.* **215**(3), 1039–1156, 2019
6. BURCZAK, J., MODENA, S., SZÉKELYHIDI, L.: Non uniqueness of power-law flows. [arXiv:2007.08011](https://arxiv.org/abs/2007.08011) (2020)

7. BONY, J.-M.: Calcul symbolique et propagation des singularités pour les équations aux dérivées partielles non linéaires. *Ann. Sci. École Norm. Sup. (4)* **14**(2), 209–246, 1981
8. BARBU, V., RÖCKNER, M.: Global solutions to random 3D vorticity equations for small initial data. *J. Differ. Equ.* **263**, 5395–5411, 2017
9. BUCKMASTER, T., VICOL, V.: Convex integration and phenomenologies in turbulence. *EMS Surv. Math. Sci.* **6**(1–2), 173–263, 2019
10. BUCKMASTER, T., VICOL, V.: Nonuniqueness of weak solutions to the Navier–Stokes equation. *Ann. Math.* **189**(1), 101–144, 2019
11. BUCKMASTER, T., VICOL, V.: Convex integration constructions in hydrodynamics. *Bull. Amer. Math. Soc.* **58**(1), 1–44, 2021
12. CHANDRA, A., CHEVYREV, I., HAIRER, M., SHEN, H.: Langevin dynamic for the 2D Yang–Mills measure. arXiv preprint [arXiv:2006.04987](https://arxiv.org/abs/2006.04987) (2020)
13. CATELLIER, R., CHOUK, K.: Paracontrolled distributions and the 3-dimensional stochastic quantization equation. *Ann. Probab.* **46**(5), 2621–2679, 2018
14. COLOMBO, M., DE LELLIS, C., DE ROSA, L.: Ill-posedness of Leray solutions for the hypodissipative Navier–Stokes equations. *Commun. Math. Phys.* **362**, 659–688, 2018
15. CHANDRA, A., HAIRER, M.: An analytic BPHZ theorem for regularity structures. [arxiv:1612.08138](https://arxiv.org/abs/1612.08138).
16. CANNIZZARO, G., KIEDROWSKI, J.: Stationary stochastic Navier–Stokes on the plane at and above criticality [arXiv:2110.03959](https://arxiv.org/abs/2110.03959)
17. CHESKIDOV, A., LUO, X.: Sharp nonuniqueness for the Navier–Stokes equations. [arXiv:2009.06596](https://arxiv.org/abs/2009.06596) (2020)
18. CHESKIDOV, A., LUO, X.: L^2 -critical nonuniqueness for the 2D Navier–Stokes equations [arXiv:2105.12117](https://arxiv.org/abs/2105.12117) (2021)
19. CHEN, X., WU, B., ZHU, R., ZHU, X.: Stochastic heat equations for infinite strings with values in a manifold. *Trans. Amer. Math. Soc.* **374**(1), 407–452, 2021
20. DA PRATO, G., DEBUSSCHE, A.: Two-Dimensional Navier–Stokes Equations Driven by a Space–Time White Noise. *J. Funct. Anal.* **196**, 180–210, 2002
21. DA PRATO, G., DEBUSSCHE, A.: Ergodicity for the 3D stochastic Navier–Stokes equations. *J. Math. Pures Appl.* **82**, 877–947, 2003
22. DAI, M.: Non-uniqueness of Leray–Hopf weak solutions of the 3d Hall–MHD system. [arXiv:1812.11311](https://arxiv.org/abs/1812.11311) (2018)
23. DEBUSSCHE, A., ODASSO, C.: Markov solutions for the 3D stochastic Navier–Stokes equations with state dependent noise. *J. Evol. Equ.* **6**(2), 305–324, 2006
24. DEBUSSCHE, A., VOVELLE, J.: Invariant measure of scalar first-order conservation laws with stochastic forcing. *Probab. Theory Related Fields* **163**(3–4), 575–611, 2015
25. DE LELLIS, C., SZÉKELYHIDI, L., JR.: The Euler equations as a differential inclusion. *Ann. Math.* **170**(3), 1417–1436, 2009
26. DE LELLIS, C., SZÉKELYHIDI, L., JR.: On admissibility criteria for weak solutions of the Euler equations. *Arch. Ration. Mech. Anal.* **195**(1), 225–260, 2010
27. DE LELLIS, C., SZÉKELYHIDI, L., JR.: Dissipative continuous Euler flows. *Invent. Math.* **193**(2), 377–407, 2013
28. DONEV, A., VANDEN-EIJNDEN, E., GARCIA, A., BELL, J.: On the accuracy of finite-volume schemes for fluctuating hydrodynamics. *Commun. Appl. Math. Comput. Sci.* **5**(2), 149–97, 2010
29. FLANDOLI, F., GATAREK, D.: Martingale and stationary solutions for stochastic Navier–Stokes equations *Probab. Theory Related Fields* **102**, 367–391, 1995
30. FLANDOLI, F., LUO, D.: Convergence of transport noise to Ornstein–Uhlenbeck for 2D Euler equations under the enstrophy measure. *Ann. Probab.* **48**(1), 264–295, 2020
31. FLANDOLI, F., LUO, D.: Point vortex approximation for 2D Navier–Stokes equations driven by space-time white noise. *J. Math. Anal. Appl.* **493**(2), 124560, 2021
32. FLANDOLI, F., ROMITO, M.: Markov selections for the 3D stochastic Navier–Stokes equations. *Probab. Theory Related Fields* **140**(3–4), 407–458, 2008
33. GUBINELLI, M., HOFMANOVÁ, M.: Global solutions to elliptic and parabolic Φ^4 models in Euclidean space. *Comm. Math. Phys.* **368**(3), 1201–1266, 2019

34. GUBINELLI, M., HOFMANOVÁ, M.: A PDE construction of the Euclidean Φ^4 quantum field theory. *Comm. Math. Phys.* **384**, 1–75, 2021
35. GUBINELLI, M., IMKELLER, P., PERKOWSKI, N.: Paracontrolled distributions and singular PDEs. *Forum Math.* **3**(6), 2015.
36. GUBINELLI, M., JARA, M.: Regularization by noise and stochastic Burgers equations. *Stoch. Partial Differ. Equ. Anal. Comput.* **1**(2), 325–350, 2013
37. GONCALVES, P., JARA, M.: Nonlinear fluctuations of weakly asymmetric interacting particle systems. *Arch. Ration. Mech. Anal.* **212**(2), 597–644, 2014
38. GUBINELLI, M., PERKOWSKI, N.: KPZ reloaded. *Comm. Math. Phys.* **349**(1), 165–269, 2017
39. GUBINELLI, M., PERKOWSKI, N.: Energy solutions of KPZ are unique. *J. Amer. Math. Soc.* **31**(2), 427–471, 2018
40. GUBINELLI, M., PERKOWSKI, N.: The infinitesimal generator of the stochastic Burgers equation. *Probab. Theory Related Fields* **178**, 1067–1124, 2020
41. HAIRER, M.: Solving the KPZ equation. *Ann. Math.* **178**(2), 559–664, 2013
42. HAIRER, M.: A theory of regularity structures. *Invent. Math.* **198**(2), 269–504, 2014
43. HOFMANOVÁ, M., ZHU, R., ZHU, X.: Non-uniqueness in law of stochastic 3D Navier–Stokes equations. [arXiv:1912.11841v2](https://arxiv.org/abs/1912.11841v2), (2019)
44. HOFMANOVÁ, M., ZHU, R., ZHU, X.: On ill- and well-posedness of dissipative martingale solutions to stochastic 3D Euler equations. *Commun. Pure Appl. Math.* **LXXV**, 2446–2510, 2022
45. HOFMANOVÁ, M., ZHU, R., ZHU, X.: Global-in-time probabilistically strong and Markov solutions to stochastic 3D Navier–Stokes equations: existence and non-uniqueness. *Ann. Probab.* **51**(2), 524–579, 2023
46. ISETT, P.: A proof of Onsager’s conjecture. *Ann. Math.* **188**(3), 871–963, 2018
47. LANDAU, L.D., LIFSHITZ, E.M.: Fluid mechanics. Translated from the Russian by JB Sykes and WH Reid, Course of Theoretical Physics (1987)
48. LUO, T., TITI, E.: Non-uniqueness of weak solutions to hyperviscous Navier–Stokes equations—on sharpness of J.-L. Lions exponent. [arXiv:1808.07595](https://arxiv.org/abs/1808.07595) (2018)
49. LUO, D., ZHU, R.: Stochastic mSQG equations with multiplicative transport noises: white noise solutions and scaling limit Process. *Stochastic Appl.* **140**, 236–286, 2021
50. MODENA, S., SZÉKELYHIDI, JR, L.: Non-uniqueness for the transport equation with Sobolev vector fields. [arXiv:1712.03867](https://arxiv.org/abs/1712.03867) (2017)
51. MODENA, S., SZÉKELYHIDI, JR, L.: Non-renormalized solutions to the continuity equation. [arXiv:1806.09145](https://arxiv.org/abs/1806.09145) (2018)
52. MOURRAT, J.-C., WEBER, H.: Global well-posedness of the dynamic Φ^4 model in the plane. *Ann. Probab.* **45**(4), 2398–2476, 2017
53. MOURRAT, J.-C., WEBER, H.: The dynamic Φ_3^4 model comes down from infinity. *Comm. Math. Phys.* **356**(3), 673–753, 2017
54. PERKOWSKI, N., ROSATI, T.C.: The KPZ equation on the real line. *Electron. J. Probab.* **24**(117), 1–56, 2019
55. RÖCKNER, M., WU, B., ZHU, R., ZHU, X.: Stochastic heat equations with values in a manifold via Dirichlet forms. *SIAM J. Math. Anal.* **52**(3), 2237–2274, 2020
56. SHEN, H.: Stochastic quantization of an Abelian gauge theory. *Comm. Math. Phys.* **384**, 1445–1512, 2021
57. SHEN, H., SMITH, S., ZHU, R., ZHU, X.: Large N limit of the $O(N)$ linear sigma model via stochastic quantization. *Ann. Probab.* **50**(1), 131–202, 2022
58. SHEN, H., ZHU, R., ZHU, X.: Large N limit of the $O(N)$ linear sigma model in 3D. *Comm. Math. Phys.* **394**(3), 953–1009, 2022
59. SHEN, H., ZHU, R., ZHU, X.: An SPDE approach to perturbation theory of Φ_2^4 : asymptoticity and short distance behavior. *arXiv preprint* [arXiv:2108.11312](https://arxiv.org/abs/2108.11312) (2021)
60. TRIEBEL, H.: Interpolation theory, function spaces, differential operators, vol. 18. North-Holland Mathematical Library. North-Holland Publishing Co., Amsterdam-New York (1978)

61. ZHU, R., ZHU, X.: Three-dimensional Navier–Stokes equations driven by space-time white noise. *J. Differ. Equ.* **259**, 4443–4508, 2015
62. ZHU, R., ZHU, X.: Strong-Feller property for Navier–Stokes equations driven by space-time white noise, [arXiv:1709.09306](https://arxiv.org/abs/1709.09306) (2017)
63. ZHANG, X., ZHU, R., ZHU, X.: Singular HJB equations with applications to KPZ on the real line. *Probab. Theory Relat. Fields* **183**(3–4), 789–869, 2022

MARTINA HOFMANOVÁ & RONGCHAN ZHU

Fakultät für Mathematik,
Universität Bielefeld,
33501 Bielefeld
Germany.

e-mail: hofmanova@math.uni-bielefeld.de

and

RONGCHAN ZHU

Department of Mathematics,
Beijing Institute of Technology,
Beijing
100081 China.

e-mail: zhurongchan@126.com

and

XIANGCHAN ZHU

Academy of Mathematics and Systems Science,
Chinese Academy of Sciences,
Beijing
100190 China.

e-mail: zhuxiangchan@126.com

(Received August 6, 2022 / Accepted March 17, 2023)

Published online April 26, 2023

© The Author(s), under exclusive licence to Springer-Verlag GmbH, DE, part of Springer Nature (2023)