



Dynamic asymptotic dimension: relation to dynamics, topology, coarse geometry, and C^* -algebras

Erik Guentner¹ · Rufus Willett¹ · Guoliang Yu^{2,3}

Received: 27 October 2015 / Revised: 25 February 2016 / Published online: 28 March 2016
© Springer-Verlag Berlin Heidelberg 2016

Abstract We introduce dynamic asymptotic dimension, a notion of dimension for actions of discrete groups on locally compact spaces, and more generally for locally compact étale groupoids. We study our notion for minimal actions of the integer group, its relation with conditions used by Bartels, Lück, and Reich in the context of controlled topology, and its connections with Gromov’s theory of asymptotic dimension. We also show that dynamic asymptotic dimension gives bounds on the nuclear dimension of Winter and Zacharias for C^* -algebras associated to dynamical systems. Dynamic asymptotic dimension also has implications for K -theory and manifold topology: these will be drawn out in subsequent work.

1 Introduction

The main aim of this paper is to introduce *dynamic asymptotic dimension*, a property of topological dynamical systems. Precisely, we are interested in actions of discrete groups on locally compact spaces: throughout the paper, we say ‘ $\Gamma \curvearrowright X$ is an action’ as shorthand for saying that Γ is a discrete group acting by homeomorphisms on a locally compact Hausdorff topological space X .

✉ Rufus Willett
rufus@math.hawaii.edu

Erik Guentner
erik@math.hawaii.edu

Guoliang Yu
guoliangyu@math.tamu.edu

¹ University of Hawai‘i at Mānoa, Honolulu, HI, USA

² Texas A&M University, College Station, USA

³ Shanghai Center for Mathematical Sciences, Shanghai, China

To give an idea of our main definition, we will state it here in a specialized form. First, we need a preliminary on ‘broken’ orbit equivalence relations.

Definition 1.1 Let $\Gamma \curvearrowright X$ be an action. For a subset E of Γ and an open subset U of X , let $\sim_{U,E}$ be the equivalence relation on U generated by E : precisely, for $x, y \in U$, $x \sim_{U,E} y$ if there is a finite sequence

$$x = x_0, x_1, \dots, x_n = y$$

of points in U such that for each $i \in \{1, \dots, n\}$ there exists $g \in E \cup E^{-1} \cup \{e\}$ such that $gx_{i-1} = x_i$.

Note that if $U = X$ then $\sim_{U,E}$ is just the equivalence relation of being in the same orbit for the subgroup $\langle E \rangle$ of Γ generated by E . However, if U is a proper subset of X then $\sim_{U,E}$ equivalence classes will generally be smaller than the intersection of U with the $\langle E \rangle$ -orbits.

Here is our main definition.

Definition 1.2 Let $\Gamma \curvearrowright X$ be an action, where we assume for simplicity that X is compact, and the action is free.¹ The *dynamic asymptotic dimension* of $\Gamma \curvearrowright X$ is the smallest $d \in \mathbb{N}$ with the following property: for each finite subset E of Γ , there is an open cover $\{U_0, \dots, U_d\}$ of X such that for each $i \in \{0, \dots, d\}$, the equivalence relation $\sim_{U_i,E}$ on U_i has uniformly finite equivalence classes.

Heuristically, we think that $\Gamma \curvearrowright X$ has dynamic asymptotic dimension at most d , if for any (large) finite subset E of Γ , the action can be ‘broken’ into at most $d + 1$ parts, and on each part the ‘action generated by E ’ has only ‘finite complexity’. We generalize this definition to non-free actions on non-compact spaces in Definition 2.1 below, and then to locally compact, Hausdorff, étale groupoids in Definition 5.1.

Our main motivation for introducing this property is its implications for K -theory of associated algebras (and more general categories) and thus for manifold topology. These implications come via the use of controlled cutting-and-pasting, or Mayer–Vietoris, techniques pioneered by the third author [44] in the setting of asymptotic dimension, and developed in a more general context by the first and third authors in collaboration with Tessera [15]. Other important motivations come from work of Farrell and Jones [12], of Bartels, Lück and Reich [3], and of Bartels and Lück [2] in controlled topology. We will explore these aspects in other work [16].

We believe however, that dynamic asymptotic dimension will admit many interesting examples, and be useful in other contexts: it is the purpose of this paper to explore some of these other aspects. Specifically, we will develop some of the main motivating examples and discuss some consequences for the structure theory of C^* -algebras.

1.1 Examples

Our main theorems on examples are as follows.

¹ This means that if $gx = x$ for some $g \in \Gamma$ and $x \in X$, then $g = e$ is the identity element of Γ .

- Theorem 1.3** (i) *Let $\mathbb{Z} \curvearrowright X$ be a free, minimal \mathbb{Z} action on a compact space. Then the dynamic asymptotic dimension of $\mathbb{Z} \curvearrowright X$ is one.*
- (ii) *Let $\Gamma \curvearrowright X$ be an action satisfying a ‘Bartels–Lück–Reich type condition of dimension d ’ for the family of finite subgroups. Then the dynamic asymptotic dimension of $\Gamma \curvearrowright X$ is at most d .*
- (iii) *The canonical action $\Gamma \curvearrowright \beta\Gamma$ of any (countable) discrete group on its Stone–Čech compactification is equal to the asymptotic dimension of Γ in the sense of Gromov. More generally, if X is a bounded geometry coarse space and $G(X)$ the associated coarse groupoid, then the dynamic asymptotic dimension of $G(X)$ equals the asymptotic dimension of X .*
- (iv) *Let Γ be a countable group with asymptotic dimension d . Then Γ admits a free minimal action on the Cantor set with dynamic asymptotic dimension at most d .*

See Theorems 3.1, 4.7, 6.4 and Corollary 6.6 below for parts (i), (ii), (iii) and (iv) respectively (and more explanation of the terminology involved in each part).

Part (i) says that the actions that are perhaps most interesting from the point of view of classical topological dynamics fall under the purview of dynamic asymptotic dimension in a natural and simple way. On the other hand, part (iv) implies that many interesting classes of groups—for example, word hyperbolic groups [32], CAT(0) cubical groups [43], lattices in Lie groups, and many solvable groups [4]—admit at least some actions with finite dynamic asymptotic dimension. Parts (ii) and (iii) were our principal motivations, as they give the connections to controlled K -theory and controlled topology which underlie our work in those directions.

1.2 Implications

The main implications we explore in this paper are to the structure theory of C^* -algebras: in particular to nuclear dimension in the sense of Winter and Zacharias [42], a property that has been very important in Elliott’s classification program [9] and elsewhere. The main result we have here is Theorem 8.6 below: this says that under a minor technical hypothesis, the nuclear dimension of the reduced C^* -algebra of a free étale groupoid can be bounded in terms of the dynamic asymptotic dimension of the groupoid, and the covering dimension of the groupoid’s unit space. Rather than repeat Theorem 8.6 here, we just give some corollaries. See Sect. 8 for more details.

- Theorem 1.4** (i) *Let $\mathbb{Z} \curvearrowright X$ be a free, minimal action on a second countable compact space of covering dimension N . Then the nuclear dimension of the crossed product $C(X) \rtimes \mathbb{Z}$ is at most $2N + 1$.*
- (ii) *Let X be a bounded geometry coarse space. Then the nuclear dimension of the uniform Roe algebra $C_u^*(X)$ is at most the asymptotic dimension of X .*
- (iii) *Any countable group Γ admits a free, minimal action on the Cantor set X such that the associated reduced crossed product $C(X) \rtimes_r \Gamma$ has nuclear dimension at most the asymptotic dimension of Γ .*

Parts (i) and (ii) are originally due to Toms and Winter [39, Section 3] and Winter and Zacharias [42, Section 8] respectively; moreover, our proofs are in some sense

close to the original ones (if heavily disguised). Nonetheless, we think there is some interest in explicitly bringing these results under one dynamical framework. The final result seems to be new, and says that ‘many’ groups admit simple crossed products with finite nuclear dimension. Thanks to spectacular recent advances in C^* -algebra theory [10,38], it seems that one now knows that this implies these crossed products fall under the purview of the Elliott program.

1.3 Outline of the paper

In Sect. 2 we introduce our main definition in the case of group actions, and mention some basic consequences. In Sect. 3 we study minimal \mathbb{Z} actions on compact spaces using ideas of Putnam [27]. In Sect. 4 we discuss the connection to the work of Bartels, Lück, and Reich [1,3] in controlled topology. In Sect. 5, we extend the main definition to étale groupoids with applications to coarse geometry in the next section in mind. In Sect. 6 we explore the connections to coarse geometry, and use this to construct the examples in part (iv) of Theorem 1.3, as well as to clarify the relationship of the Bartels–Lück–Reich conditions to asymptotic dimension. Section 7 is devoted to a technical construction of almost invariant partitions of unity. Finally, Sect. 8 discusses the implications to nuclear dimension of groupoid C^* -algebras, and in particular all the parts of Theorem 1.4.

We have tried to write the paper in a ‘modular’ way, so that sections can be read independently of each other to a large extent. In particular, groupoids are not mentioned until Sect. 5, and (noncommutative) C^* -algebras are not mentioned before Sect. 8.

2 Dynamic asymptotic dimension for group actions

In this section, we introduce our main definition.

Definition 2.1 An action $\Gamma \curvearrowright X$ has *dynamic asymptotic dimension* d if d is the smallest natural number with the following property: for any compact subset K of X and finite subset E of Γ there are open subsets U_0, \dots, U_d of X that cover K such that for each $i \in \{0, \dots, d\}$, the set

$$\left\{ g \in \Gamma \left| \begin{array}{l} \text{there exist } x \in U_i \text{ and } g_n, \dots, g_1 \in E \text{ such that} \\ g = g_n \cdots g_2 g_1 \text{ and for all } k \in \{1, \dots, n\}, \\ g_k \cdots g_1 x \in U_i \end{array} \right. \right\} \quad (2.1)$$

is finite.

The action has *finite dynamic asymptotic dimension* if it has dynamic asymptotic dimension d for some d , and *infinite dynamic asymptotic dimension* otherwise.

Remark 2.2 (i) An action $\Gamma \curvearrowright X$ on a compact space X has dynamic asymptotic dimension 0 if and only if Γ is locally finite, i.e., any finite subset of Γ generates a finite subgroup.

(ii) Recall that an action $\Gamma \curvearrowright X$ is *proper* if for any compact subset K of X , the set $\{g \in \Gamma \mid gK \cap K \neq \emptyset\}$ is finite. It is *locally proper* if every finitely generated

subgroup of Γ acts properly on X . Point (i) says that an action on a compact space has dynamic asymptotic dimension zero if and only if it is locally proper. In general, it is still true that local properness implies dynamic asymptotic dimension zero: indeed, let a finite $E \subseteq \Gamma$ and compact $K \subseteq X$ be given. Let U_0 be any relatively compact open set that contains K (such a U_0 exists as X is locally compact). Then if $\langle E \rangle$ is the subgroup of Γ generated by E , the set in line (2.1) above is contained in

$$\{g \in \langle E \rangle \mid g^{-1}U_0 \cap U_0 \neq \emptyset\}$$

and is thus finite by local properness. The converse is false, however: for example, the action of \mathbb{Z} on \mathbb{R}^2 considered in [22, Chapter 5, Example 8.3] has dynamic asymptotic dimension zero but is not (locally) proper. It is perhaps most natural to characterize dynamic asymptotic dimension zero actions in terms of groupoids: see Example 5.3 below.

- (iii) If an action has finite dynamic asymptotic dimension, then all point stabilizers must be locally finite. In particular, an action of a torsion free group with finite dynamic asymptotic dimension is free.
- (iv) If $\Gamma \curvearrowright X$ is a free action with X compact, then Definition 2.1 above is equivalent to Definition 1.2 from the introduction.

To see this, assume first that $\Gamma \curvearrowright X$ satisfies the condition in Definition 2.1, and let a finite subset E of Γ be given. Take $K = X$, and a cover U_0, \dots, U_d as in Definition 2.1 for this K and E . Then if F_i is the finite set in line (2.1), all $\sim_{U_i, E}$ equivalence classes are contained in $F_i \cdot x$ for some x , and thus have uniformly bounded finite cardinalities.

Conversely say a finite subset E of Γ is given and $\{U_0, \dots, U_d\}$ is a cover of X with the properties in Definition 1.2, so in particular all $\sim_{U_i, E}$ equivalence classes have cardinality at most some integer N . Then each of the sets F_i in line (2.1) is contained in $(E \cup E^{-1} \cup \{e\})^N$, and thus has cardinality at most $(2|E| + 1)^N$.

- (v) See Corollary 7.2 below for a stronger-looking equivalent definition of dynamic asymptotic dimension in terms of almost-invariant partitions of unity.

3 Example: minimal \mathbb{Z} -actions

In this section we give our first non-trivial examples: minimal actions of \mathbb{Z} on compact spaces (recall that an action $\Gamma \curvearrowright X$ is *minimal* if all Γ -orbits are dense).

The proof is inspired by ideas of Ian Putnam that were originally used to build interesting AF-algebras associated to minimal actions of \mathbb{Z} on the Cantor set [27] (and subsequently much developed by Putnam and others). We would like to thank Prof. Putnam for suggesting that there might be a connection between his work and our notion of dynamic asymptotic dimension.

Theorem 3.1 *Let $\mathbb{Z} \curvearrowright X$ be a minimal \mathbb{Z} action on an infinite compact space X . Then the dynamic asymptotic dimension of the action is one.*

Proof As \mathbb{Z} is not locally finite, Remark 2.2 part (i) implies that the dynamic asymptotic dimension of $\mathbb{Z} \curvearrowright X$ is not zero, so it suffices to bound the dynamic asymptotic dimension above by one.

Let a finite subset E of \mathbb{Z} be given; we may as well assume that E is an ‘interval’ $[-N, N] \cap \mathbb{Z}$ for some $N \in \mathbb{N}$. As the action is minimal and X is infinite, the action is free. It follows that we can find a non-empty open subset U of X such that

$$n \cdot U \cap U = \emptyset$$

for all $n \in [-5N, 5N] \setminus \{0\}$. Let V be any non-empty open subset of X such that $\overline{V} \subseteq U$. Define

$$U_0 := \bigcup_{n=-N}^N n \cdot U, \quad U_1 := X \setminus \bigcup_{n=-N}^N n \cdot \overline{V}.$$

Clearly $\{U_0, U_1\}$ is an open cover of X . To finish the proof, it suffices to show that the sets

$$\left\{ n \in \mathbb{Z} \left| \begin{array}{l} \text{there exist } x \in U_i \text{ and } n_1, \dots, n_m \in E \text{ such that} \\ n = n_1 + \dots + n_m \text{ and for all } k \in \{1, \dots, m\}, \\ (n_k + \dots + n_1)x \in U_i \end{array} \right. \right\} \quad (3.1)$$

are finite for $i = 0, 1$.

First look at U_0 . We claim that in this case the set in line (3.1) is contained in $[-3N, 3N]$. Indeed, assume for contradiction that there exist $n_1, \dots, n_m \in [-N, N]$ with $|n_1 + \dots + n_m| > 3N$ and $x \in U_0$ with $(n_k + \dots + n_1)x \in U_0$ for all $k \in \{1, \dots, m\}$. Write $x = n_0x_0$ for some $x_0 \in U$ and $n_0 \in [-N, N]$ (such exist by definition of U_0). As each n_k is in $[-N, N]$ and $|n_m + \dots + n_1| > 3N$, there must exist k such that $3N \geq |n_k + \dots + n_1| > 2N$, whence

$$4N \geq |n_k + \dots + n_1 + n_0| > N. \quad (3.2)$$

As

$$(n_k + \dots + n_1)x = (n_k + \dots + n_1 + n_0)x_0$$

is in U_0 , there are $n'_0 \in [-N, N]$ and $x'_0 \in U$ with

$$(n_k + \dots + n_1 + n_0)x_0 = n'_0x'_0,$$

whence

$$(n_k + \dots + n_1 + n_0 - n'_0)x_0 = x'_0$$

Hence as x_0 and x'_0 are in U we have that

$$(n_k + \dots + n_1 + n_0 - n'_0) \cdot U \cap U \neq \emptyset.$$

However, line (3.2) implies that $5N \geq |n_k + \dots + n_1 + n_0 - n'_0| > 0$, so this contradicts that $U \cap n \cdot U = \emptyset$ for $n \in [-5N, 5N] \setminus \{0\}$.

We now look at the set in line (3.1) for U_1 . We first claim that there exists $M \in \mathbb{N}$ such that for all $x \in U_1$ there exists $m_- \in [-M, 0)$ and $m_+ \in (0, M]$ with $m_-x \in V$ and $m_+x \in V$. Indeed, for each $M \in \mathbb{N}$ define

$$W_M := \left\{ x \in X \mid \begin{array}{l} \text{there are } m_- \in [-M, 0), m_+ \in (0, M] \\ \text{such that } m_-x, m_+x \in V \end{array} \right\}.$$

It follows from the fact that V is open that each W_M is open. Moreover, minimality of the action implies that for each $x \in X$, the ‘half-orbits’ $\mathbb{N} \cdot x$ and $(-\mathbb{N}) \cdot x$ are dense: if not, the limit points of one of these sets would be a closed \mathbb{Z} -invariant subset. Hence the fact that V is open implies that each $x \in X$ is in W_M for some M , and so $\{W_M \mid M \in \mathbb{N}\}$ is an open cover of X . Compactness of X and the fact that $W_{M_1} \subseteq W_{M_2}$ for $M_1 \leq M_2$ implies that X is contained in W_M for some M , and this implies the claim.

To complete the proof, we will now show that for U_1 , the set in line (3.1) is contained in $[-M - N, M + N]$. Assume for contradiction this fails, so there exist $n_1, \dots, n_m \in E$ with $|n_1 + \dots + n_m| > M + N$ and $x \in U_1$ such that $(n_1 + \dots + n_k)x$ is in U_1 for all $k \in \{1, \dots, m\}$. Assume for simplicity that $n_1 + \dots + n_m > M + N$; the case $n_1 + \dots + n_m < -M - N$ can be handled similarly. Let $m_+ \in (0, M]$ be such that $m_+x \in V$. Then there exists $k \in \{1, \dots, m\}$ with $(n_1 + \dots + n_k) - m_+ \in [-N, N]$. In particular,

$$(n_1 + \dots + n_k)x = (m_+ + n)x$$

for some $n \in [-N, N]$. However as m_+x is in V , $(m_+ + n)x$ is in $X \setminus U_1$ by definition of U_1 , which is a contradiction. □

4 Example: Bartels–Lück–Reich conditions

Our main goal in this section is to study some properties of group actions that were important in the work of Bartels et al. [1,3] on the Farrell–Jones conjecture and show that that they imply that the action has finite dynamic asymptotic dimension.

We will need to establish some conventions on simplicial complexes. Let V be a set, thought of a discrete topological space. The *space of probability measures* on V is

$$P(V) := \{\mu \in l^1(V) \mid \mu(v) \geq 0 \text{ for all } v \in V \text{ and } \|\mu\|_1 = 1\},$$

equipped with the metric

$$d\left(\sum_{v \in V} t_v v, \sum_{v \in V} s_v v\right) := \sum_{v \in V} |t_v - s_v|.$$

coming from the l^1 -norm. Write $P_n(V)$ for the subset of $P(V)$ consisting of measures supported on at most $n + 1$ points and define

$$P_f(V) := \bigcup_{n=0}^{\infty} P_n(V)$$

to be the subspace of finitely supported probability measures. Elements of $P(V)$ will usually be written as formal sums

$$\mu = \sum_{v \in V} t_v v,$$

where $\mu(v) = t_v$ is in $[0, 1]$ and $\sum_v t_v = 1$. We identify V with $P_0(V)$ in the obvious way.

A *simplicial complex* is a subspace C of some $P_f(V)$ as above such that $V \subseteq C$, and with the property that whenever

$$\mu = \sum_{v \in V} t_v v$$

is an element of C and $S = \{v \in V \mid t_v \neq 0\}$ is the support of μ , then

$$\left\{ \sum_{v \in V} t_v v \in P_f(V) \mid t_v = 0 \text{ for } v \notin S \right\}$$

is contained in C . The *vertex set* of C is V . A simplicial complex C is equipped with the restriction of the l^1 -metric defined above. If C is a simplicial complex and $n \in \mathbb{N}$, an n -*simplex* in C is a subset that is equal to the convex hull of some set of $n + 1$ vertices. The n -*skeleton* of C is defined to be $C_n := C \cap P_n(V)$ (so in particular $C_0 = V$ and $C_{-1} = \emptyset$). The *dimension* of C is the smallest d such that $C = C_d$ (or infinity if no such d exists).

If Γ is a discrete group and C a simplicial complex with vertex set C , then a *simplicial action* of Γ on C is an action that is induced from some action of Γ on V via the formula

$$g \left(\sum_{v \in V} t_v v \right) = \sum_{v \in V} t_v (gv).$$

Note that a simplicial action is isometric. All actions on simplicial complexes will be assumed simplicial.

We will need the following technical lemma at a couple of points below: roughly, it says that simplicial complexes admit covers with rather rigid combinatorial properties.

Lemma 4.1 *Let C be a simplicial complex of dimension at most d and equipped with a simplicial action by Γ . For a subset A of C and $\delta > 0$, write*

$$N_\delta(A) := \{x \in C \mid d(x, A) < \delta\}$$

for the δ -neighbourhood of A . For each $i \in \{0, \dots, d\}$ define

$$V_i := N_{\frac{1}{3}10^{-i}}(C_i) \setminus \overline{N_{\frac{5}{2}10^{-i}}(C_{i-1})}.$$

Then the collection $\{V_0, \dots, V_d\}$ is an open cover of C by Γ -invariant subsets.

Moreover, for each $i \in \{0, \dots, d\}$ and i -simplex Δ , define

$$V_{i \Delta} := N_{\frac{1}{3}10^{-i}}(\Delta) \setminus \overline{N_{\frac{5}{2}10^{-i}}(C_{i-1})}.$$

Then

$$V_i = \bigcup_{\Delta \text{ an } i\text{-simplex}} V_{i \Delta},$$

the Γ action permutes the distinct $V_{i \Delta}$, and for distinct i -simplices Δ and Δ' ,

$$d(V_{i \Delta}, V_{i \Delta'}) \geq \frac{1}{3}10^{-i} \geq \frac{1}{3}10^{-d}. \tag{4.1}$$

Proof Clearly each V_i is open, and each is Γ -invariant as Γ acts simplicially (whence it preserves each i -skeleton and the metric). To show that $\{V_0, \dots, V_d\}$ covers C , it suffices to show that $C_i \subseteq V_0 \cup \dots \cup V_i$ for each i , which is clear by induction. The decomposition $V_i = \bigcup_{\Delta \text{ an } i\text{-simplex}} V_{i \Delta}$ is also clear as $C_i = \bigcup_{\Delta \text{ an } i\text{-simplex}} \Delta$. Moreover, as the action is simplicial Γ permutes the sets $V_{i \Delta}$.

Finally, say for contradiction that Δ, Δ' are distinct i -simplices such that $d(V_{i \Delta}, V_{i \Delta'}) < \frac{1}{3}10^{-i}$. Then there exists $\mu \in V_{i \Delta}$ such that $d(\mu, \Delta') < \frac{2}{3}10^{-i}$. Note that

$$\frac{1}{3}10^{-i} > d(\mu, \Delta) \geq \sum_{v \notin \Delta} t_v, \quad \text{and} \quad \frac{2}{3}10^{-i} > d(\mu, \Delta') \geq \sum_{v \notin \Delta'} t_v$$

whence

$$T := \sum_{v \notin \Delta \cap \Delta'} t_v \leq \sum_{v \notin \Delta} t_v + \sum_{v \notin \Delta'} t_v < 10^{-i}.$$

Hence in particular $T < 1$ and so we may define

$$v := \sum_{v \in \Delta \cap \Delta'} \frac{t_v}{1 - T} v,$$

which is in C_{i-1} as Δ and Δ' are distinct. Note that

$$d(\mu, C_{i-1}) \leq d(\mu, v) = \sum_{v \in \Delta \cap \Delta'} t_v \left(\frac{1}{1 - T} - 1 \right) + \sum_{v \notin \Delta \cap \Delta'} t_v = 2T < 2 \cdot 10^{-i}.$$

However, as μ is in $V_i \Delta$ we have $\mu \notin \overline{N_{\frac{5}{2}} 10^{-i}(C_{i-1})}$, and putting this together with the line above gives

$$\frac{5}{2} 10^{-i} < d(\mu, C_{i-1}) < 2 \cdot 10^{-i},$$

which is the desired contradiction. □

A *family* of subgroups of a discrete group Γ is a collection of subgroups satisfying the following conditions:

- \mathcal{F} is closed under conjugation;
- \mathcal{F} is closed under taking subgroups;
- \mathcal{F} is closed under taking finite index supergroups.

The examples that are important for applications are the family of finite subgroups, and that of virtually cyclic subgroups.

Definition 4.2 Let C be a simplicial complex equipped with a simplicial action of a discrete group Γ . Let \mathcal{F} be a family of subgroups of Γ satisfying the conditions above. Then C is called a (Γ, \mathcal{F}) -*complex* if the stabilizer of every vertex in C is an element of \mathcal{F} .

We will only use the following definition in the case that Y is a simplicial complex equipped with a simplicial action of Γ .

Definition 4.3 Let X be a topological space and Y a metric space, and assume that Γ is a discrete group acting on X by homeomorphisms, and on Y by isometries. Let E be a subset of Γ and $\epsilon > 0$. A map $f : X \rightarrow Y$ is (E, ϵ) -*equivariant* if

$$\sup_{x \in X} d_Y(f(gx), gf(x)) < \epsilon.$$

for all $g \in E$.

As the last of our preliminaries before getting to the conditions of Bartels–Lück–Reich, we have another technical lemma.

Lemma 4.4 Let $f : X \rightarrow C$ be a continuous (E, ϵ) -equivariant map as in Definition 4.3, where X is compact and E is finite. Then there exists a finite subset S of the vertex set of C and an (E, ϵ) -equivariant map $f' : X \rightarrow C$ such that $f'(X) \subseteq P(S) \cap C$.

Proof Note that for any fixed $\delta > 0$, the collection

$$\{N_\delta(P(S) \cap C) \mid S \text{ a finite set of vertices}\}$$

is an open cover of C . It follows that if $K \subseteq C$ is compact, then for any $\delta > 0$ there exists a finite subset S of V such that $K \subseteq N_\delta(P(S) \cap C)$. Define

$$\delta := \min \left\{ 1, \frac{1}{2} \min_{g \in E} \left\{ \epsilon - \sup_{x \in X} d_C(f(gx), gf(x)) \right\} \right\} > 0$$

and let S be a finite subset of V such that $f(X) \subseteq N_{\delta/2}(P(S) \cap C)$. We may write $f(x) = \sum_{v \in V} t_v(x)v$ where each $t_v : X \rightarrow [0, 1]$ is a continuous function. Note that for any $\mu = \sum_{v \in S} s_v v \in P(S)$,

$$d(\mu, f(x)) = \sum_{v \in S} |s_v - t_v(x)| + \sum_{v \notin S} t_v(x) \geq \sum_{v \notin S} t_v(x).$$

Taking the infimum over all such μ and using that $f(X) \subseteq N_{\delta/2}(P(S) \cap C)$ gives

$$\sum_{v \notin S} t_v(x) < \delta/2.$$

Hence in particular the formula $T(x) := \sum_{v \in S} t_v(x)$ defines a continuous function $T : X \rightarrow (1 - \delta/2, 1]$. Define

$$f'(x) := \sum_{v \in S} \frac{t_v(x)}{T(x)}v,$$

so $f' : X \rightarrow C \cap P(S)$ is a continuous function. Then for any $x \in X$,

$$d_C(f(x), f'(x)) = \sum_{v \in S} t_v(x) \left(\frac{1}{T(x)} - 1 \right) + \sum_{v \notin S} t_v(x) = 2(1 - T(x)) < \delta.$$

It follows that $f' : X \rightarrow C$ has the desired properties. □

We now come to the conditions that are our main object of study in this section. The first condition in the following proposition is essentially taken from Bartels' survey paper [1, Theorem A, page 9], and the second from the paper of Bartels–Lück–Reich on equivariant covers of hyperbolic groups [3, Theorem 1.2]. The result is very closely connected to [37, Lemma 4.4], and is to some extent already implicit in [3]; nonetheless, we do not think a complete proof exists anywhere in the literature, so give one here.

Proposition 4.5 *Say $\Gamma \curvearrowright X$ is an action with X compact, and \mathcal{F} is a family of subgroups of Γ . The following are equivalent.*

- (i) *There exists d such that for all finite $E \subseteq \Gamma$ and all $\epsilon > 0$ there exists a (Γ, \mathcal{F}) -complex C of dimension at most d and a continuous (E, ϵ) -equivariant map*

$$f : X \rightarrow C.$$

- (ii) *There exists d such that for each finite subset E of Γ there exists a Γ equivariant open cover \mathcal{U} of $X \times \Gamma$ such that the following hold.*
 - (A) *For every $U \in \mathcal{U}$ and $g \in \Gamma$, $gU = U$, or $gU \cap U = \emptyset$.*
 - (B) *For every $U \in \mathcal{U}$, $\{g \in \Gamma \mid gU = U\} \in \mathcal{F}$.*
 - (C) *The multiplicity of \mathcal{U} is at most $d + 1$.*

(D) \mathcal{U}/Γ is finite.

(E) For every $g \in \Gamma$ and $x \in X$ there exists $U \in \mathcal{U}$ such that $\{x\} \times gE \subseteq U$.

Proof Assume condition (i). Let $\epsilon = \frac{1}{6}10^{-d}$. Fix a finite subset E of Γ , and let C, f be as in condition (i) for this choice of E and ϵ . Using Lemma 4.4, we may assume that the image of f is contained in $P(S) \cap C$ for some finite subset S of the vertex set of C . Define

$$\phi : X \times \Gamma \rightarrow C \cap P(\Gamma \cdot S), \quad (x, g) \mapsto gf(g^{-1}x).$$

Then ϕ is equivariant, and for $g, h \in \Gamma$ with $g^{-1}h \in E$ satisfies

$$\begin{aligned} d(\phi(x, g), \phi(x, h)) &= d(gf(g^{-1}x), hf(h^{-1}x)) \\ &= d(f(g^{-1}x), g^{-1}hf(h^{-1}x)) \\ &< \epsilon. \end{aligned} \tag{4.2}$$

Now, let $\{V_0, \dots, V_d\}$ be the cover of C given by Lemma 4.1. With notation from Lemma 4.1, define

$$\mathcal{U} := \{\phi^{-1}(N_{\frac{1}{6}}(V_i \Delta)) \mid i \in \{0, \dots, d\} \text{ and } \Delta \text{ an } i\text{-simplex}\}.$$

The open cover \mathcal{U} of $X \times \Gamma$ is equivariant as it is the pullback of an equivariant cover by an equivariant map. It satisfies condition (A) as Γ permutes the disjoint sets $N_{\frac{1}{6}}(V_i \Delta)$ for each fixed i ; it satisfies condition (B) as C is a (Γ, \mathcal{F}) -complex; it satisfies condition (C) as each point in C can intersect at most one $N_{\frac{1}{6}}(V_i \Delta)$ for each i ; and it satisfies condition (D) as S is finite. Moreover, as $\{V_0, \dots, V_d\}$ covers C , the Lebesgue number of the cover

$$\mathcal{V} := \{N_{\frac{1}{6}}(V_i \Delta) \mid i \in \{0, \dots, d\} \text{ and } \Delta \text{ an } i\text{-simplex}\}.$$

of C is at least $\epsilon = \frac{1}{6}10^{-d}$. Hence the condition

$$g^{-1}h \in E \implies d(\phi(x, g), \phi(x, h)) < \epsilon$$

from line (4.2) above implies for any x and g there exists $V \in \mathcal{V}$ such that

$$\phi(\{x\} \times gE) \subseteq V;$$

this implies condition (E) for \mathcal{U} .

Now assume condition (ii). Let ϵ, E be given; replacing E by $E \cup E^{-1} \cup \{e\}$, we may assume that E is symmetric and contains the identity element. Let d be as in (ii), and take a cover \mathcal{U} of $X \times \Gamma$ as in (ii) for the finite set $E^n = \{g \in \Gamma \mid g = g_1 \dots g_n \text{ for some } g_1, \dots, g_n \in E\}$ where n is such that

$$\frac{(2d + 2)(4d + 6)}{n} < \epsilon.$$

Let C be the *nerve* of this cover: precisely, C is the subcomplex of $P(\mathcal{U})$ consisting of the union of the convex hulls of all subsets $\{U_0, \dots, U_m\}$ of the vertex set \mathcal{U} such that $\bigcap_{i=0}^m U_i \neq \emptyset$. Note that C is a (Γ, \mathcal{F}) -complex by conditions (A) and (B); it is moreover of dimension at most d by (C).

The next part of the argument is a ‘topological version’ of [7, Proposition 4.1]. For each $U \in \mathcal{U}$ and each $m \in \{0, \dots, n\}$ define the E^m -interior of U to be

$$U^{(m)} := \{(x, g) \in U \mid \{x\} \times gE^m \subseteq U\}.$$

This gives rise to a nested sequence of open sets

$$U^{(n)} \subseteq U^{(n-1)} \subseteq \dots \subseteq U^{(1)} \subseteq U^{(0)} = U.$$

Note that for each $m \in \{0, \dots, n\}$, $\{U^{(m)}\}_{U \in \mathcal{U}}$ is an open cover of $X \times \Gamma$ by condition (E). Let $\{V_U\}_{U \in \mathcal{U}}$ be an open cover of $X \times \Gamma$ such that for each U , $\overline{V_U} \subseteq U^{(n)}$; as $X \times \Gamma$ is normal, standard arguments in general topology imply that such a ‘shrunkened’ cover exists.

Fix $U \in \mathcal{U}$ for the moment. Define $V^{(n)} := V_U$. Define

$$W^{(n)} := \{(x, gh) \in X \times \Gamma \mid (x, g) \in V^{(n)}, h \in E\}$$

and note that if (x, g) is in $\overline{W^{(n)}}$, then (x, gh) is in $V^{(n)}$ for some $h \in E$, and thus

$$\{x\} \times gE^{n-1} \subseteq \{x\} \times ghE^n \subseteq U.$$

This says that $\overline{W^{(n)}} \subseteq U^{(n-1)}$, and thus by normality of $X \times \Gamma$ there is an open set $V^{(n-1)}$ with

$$\overline{W^{(n)}} \subseteq V^{(n-1)} \subseteq \overline{V^{(n-1)}} \subseteq U^{(n-1)}.$$

Now set

$$W^{(n-1)} := \{(x, gh) \in X \times \Gamma \mid (x, g) \in V^{(n-1)}, h \in E\},$$

and similarly we have $\overline{W^{(n-1)}} \subseteq U^{(n-2)}$ whence there is an open set $V^{(n-2)}$ with

$$\overline{W^{(n-1)}} \subseteq V^{(n-2)} \subseteq \overline{V^{(n-2)}} \subseteq U^{(n-2)}.$$

Continuing this process gives a nested sequence

$$V^{(n)} \subseteq V^{(n-1)} \subseteq \dots \subseteq V^{(1)} \subseteq V^{(0)}$$

of open sets such that: for each $m \in \{1, \dots, n\}$, $\overline{V^{(m)}} \subseteq V^{(m-1)}$; for each $m \in \{0, \dots, n\}$, $\overline{V^{(m)}} \subseteq U^{(m)}$; and for each $m \in \{1, \dots, n\}$,

$$\{(x, gh) \in X \times \Gamma \mid (x, g) \in V^{(m)}, h \in E\} \subseteq V^{(m-1)}. \tag{4.3}$$

Now, for each $m \in \{1, \dots, n\}$, Urysohn’s lemma implies there exists a continuous function $\psi_U^{(m)} : X \times \Gamma \rightarrow [0, 1]$ such that

$$\psi_U^{(m)}(V^{(m)}) = \{1\} \quad \text{and} \quad \psi_U^{(m)}((X \times \Gamma) \setminus V^{(m-1)}) = \{0\}.$$

Define

$$\psi_U := \sum_{m=1}^n \psi_U^{(m)}.$$

Note that if $(x, g) \in V^{(m-1)} \setminus V^{(m)}$ for some $m \in \{1, \dots, n\}$ then $\psi_U(x, g) \in [m - 1, m]$ and moreover $\psi_U(x, g) = 0$ whenever $(x, g) \notin V^{(0)}$. Hence by line (4.3) we have that for any $(x, g) \in X \times \Gamma$

$$h \in E \quad \Rightarrow \quad |\psi_U(x, g) - \psi_U(x, gh)| \leq 2. \tag{4.4}$$

At this point, we unfix U , and define $\phi_U : X \times \Gamma \rightarrow [0, 1]$ by the formula

$$\phi_U = \frac{\psi_U}{\sum_{V \in \mathcal{U}} \psi_V}.$$

As $\{V_U\}_{U \in \mathcal{U}}$ is an open cover of $X \times \Gamma$, $\{\phi_U\}_{U \in \mathcal{U}}$ is a well-defined partition of unity on $X \times \Gamma$. Moreover, for any $x \in X$, any $g, h \in G$ such that $g^{-1}h \in E$, and any $U \in \mathcal{U}$ we have that $|\phi_U(x, g) - \phi_U(x, h)|$ is bounded above by

$$\left| \frac{\psi_U(x, g) - \psi_U(x, h)}{\sum_{V \in \mathcal{U}} \psi_V(x, g)} \right| + \frac{\psi_U(x, h)}{\sum_{V \in \mathcal{U}} \psi_V(x, h)} \left| \frac{\sum_{V \in \mathcal{U}} \psi_V(x, g) - \sum_{V \in \mathcal{U}} \psi_V(x, h)}{\sum_{V \in \mathcal{U}} \psi_V(x, g)} \right|$$

which in turn is bounded above by

$$\left| \frac{\psi_U(x, g) - \psi_U(x, h)}{\sum_{V \in \mathcal{U}} \psi_V(x, g)} \right| + \left| \frac{\sum_{V \in \mathcal{U}} \psi_V(x, g) - \sum_{V \in \mathcal{U}} \psi_V(x, h)}{\sum_{V \in \mathcal{U}} \psi_V(x, g)} \right|.$$

By construction, at least one of the terms in the sum $\sum_{V \in \mathcal{U}} \psi_V(x, g)$ equals n . Combining this with line (4.4) and the fact that the cover has multiplicity at most $d + 1$ implies that the above is bounded by

$$\frac{2}{n} + \frac{4(d + 1)}{n} = \frac{4d + 6}{n}.$$

Hence for $x \in X$ and $g, h \in \Gamma$ with $g^{-1}h \in E$

$$\sum_{U \in \mathcal{U}} |\phi_U(x, g) - \phi_U(x, h)| < \frac{(2d + 2)(4d + 6)}{n}.$$

as at most $2d + 2$ of the terms can be non-zero by the multiplicity restriction.

Finally, define

$$f : X \rightarrow C, \quad x \mapsto \sum_{U \in \mathcal{U}} \phi_U(x, e)U,$$

which is continuous as each ϕ_U is. Note that for $g \in E, x \in X,$

$$d(f(gx), gf(x)) = \sum_{U \in \mathcal{U}} |\phi_U(gx, e) - \phi_U(gx, g)| < \frac{(2d + 2)(4d + 6)}{n} < \epsilon$$

by choice of n ; thus we may conclude (i). □

Definition 4.6 If an action $\Gamma \curvearrowright X$ and family \mathcal{F} satisfy the conditions in Proposition 4.5, we say that the action is *d-BLR for \mathcal{F}* .

The following theorem shows that the conditions on an action in Proposition 4.5 imply that the action has dynamic asymptotic dimension at most d . We do not know whether the converse is true: some evidence for a converse is provided by Corollary 6.5 below that relates the d -BLR condition and dynamic asymptotic dimension to asymptotic dimension in the sense of coarse geometry.

Theorem 4.7 *Let $\Gamma \curvearrowright X$ be an action with X compact, and let \mathcal{F} be the family of finite subgroups of Γ . If $\Gamma \curvearrowright X$ is a d -BLR action for \mathcal{F} , then it has dynamic asymptotic dimension at most d .*

Proof Assume condition (i) in Proposition 4.5 for the family \mathcal{F} of finite subgroups of Γ . Let a finite subset E of Γ be given; we may assume that E contains the identity and that $E = E^{-1}$. Let $\epsilon = \frac{1}{3}10^{-d}$. Then there exists a (Γ, \mathcal{F}) -complex of dimension at most d and an (E, ϵ) -equivariant continuous map

$$f : X \rightarrow C$$

as in Definition 4.3. Using Remark 4.4, we may assume that there exists a finite subset S of the vertex set of C such that $f(X)$ is contained in $C \cap P(S)$. Define also

$$F := \{g \in \Gamma \mid gS \cap S \neq \emptyset\}. \tag{4.5}$$

Note that as S is finite and as the stabilizer in Γ of each vertex in S is finite, F is a finite subset of Γ .

Now, let $\{V_0, \dots, V_d\}$ be the open cover of C as in Lemma 4.1; we will freely use the notation from Lemma 4.1 in the rest of the proof. For each $i \in \{0, \dots, d\}$ define $U_i := f^{-1}(V_i)$ and for each i -simplex Δ , define $U_{i \Delta} := f^{-1}(V_{i \Delta})$. Note that $\{U_0, \dots, U_d\}$ is an open cover of X , and that each U_i is the disjoint union of the sets $U_{i \Delta}$ as Δ ranges over all i -simplices. We claim that the following holds for any i and i -simplex Δ :

$$x \in U_{i \Delta}, \quad gx \in U_i, \quad g \in E \quad \Rightarrow \quad gx \in U_{i g\Delta}. \tag{4.6}$$

Indeed, with notation as in the line above, we know that gx is in $U_{i \Delta'}$ for some i -simplex Δ' . As f is (E, ϵ) -equivariant we have

$$d(f(gx), gf(x)) < \epsilon;$$

As $gf(x)$ is in $V_{i g\Delta}$ and $f(gx)$ is in $V_{i \Delta'}$, line (4.1) from Lemma 4.1 now forces $g\Delta = \Delta'$ and thus gx is in $U_{i g\Delta}$ as claimed.

To complete the proof, say $x \in U_i$ for some i and $g = g_n \cdots g_1$ with $g_k \in E$ and $g_k \cdots g_1 x \in U_i$ for each $k \in \{1, \dots, n\}$. If x is in $U_{i \Delta}$ for some i -simplex Δ , then repeated applications of line (4.6) force gx to be in $U_{i g\Delta}$, so in particular $f(gx) \in V_{i g\Delta} \cap f(X)$. As both $f(gx)$ and $f(x)$ are supported on vertices in S , this in turn forces $gS \cap S$ to be non-empty, and thus g is in the finite set F from line (4.5) above. This shows that $\Gamma \curvearrowright X$ satisfies the conditions in Definition 2.1 so we are done. □

Remark 4.8 In earlier versions of this work, we considered the following definition. An action $\Gamma \curvearrowright X$ with X compact is *d-amenable* if for all finite subsets E of Γ and all $\epsilon > 0$ there exists an (E, ϵ) -equivariant map

$$f : X \rightarrow P_d(\Gamma).$$

This definition is motivated by the notion of an amenable action, which is the same definition without the extra restriction on d . We also said the *amenability dimension* of an action is the smallest d such that the action is d -amenable. This notion was used in recent work of Szabó, Wu, and Zacharias, where they gave very interesting relations to Rokhlin-type conditions for actions of residually finite groups: see [37], particularly Section 4.

Clearly a d -amenable action is d -BLR; it is not difficult to see that the converse is true for torsion free groups, but false (for example) for finite groups in general (take X a point, and Γ to a finite group of cardinality larger than d). Thus in particular, Theorem 4.7 also implies that d -amenable actions have dynamic asymptotic dimension at most d (and thus by our other results, d -amenability has consequences for nuclear dimension).

5 Dynamic asymptotic dimension for groupoids

In this section we reformulate the definition of dynamic asymptotic dimension for groupoids. Our main goal is to explore connections to coarse geometry in Sect. 6, but also the extra generality seemed interesting for the results in Sect. 8.

For us, groupoids are always locally compact and Hausdorff; we generally leave these assumptions implicit from now on. We will also assume that groupoids are étale, but do not leave this (less standard) assumption implicit. Accessible, as well as reasonably self-contained and concise, introductions to this class of groupoids and their associated C^* -algebras can be found in [30, Section 2.3] and [6, Section 5.6].

We use the following notation for groupoids. A groupoid is denoted G , and its unit space by $G^{(0)}$. We identify $G^{(0)}$ with a closed and open subspace of G in the canonical

way. The range and source maps are denoted $r, s : G \rightarrow G^{(0)}$. An ordered pair (g, h) of elements from G is *composable* if $s(g) = r(h)$, and the composition is written gh in this case. The inverse of $g \in G$ is denoted g^{-1} .

Definition 5.1 Let G be an étale groupoid. Then G has *dynamic asymptotic dimension* $d \in \mathbb{N}$ if d is the smallest number with the following property: for every open relatively compact subset K of G there are open subsets U_0, \dots, U_d of $G^{(0)}$ that cover $s(K) \cup r(K)$ such that for each i , the set $\{g \in K \mid s(g), r(g) \in U_i\}$ is contained in a relatively compact subgroupoid of G .

Note that we may equivalently ask that for each i , the subgroupoid G_i generated by the set $\{g \in K \mid s(g), r(g) \in U_i\}$ is relatively compact. It is often convenient to note that this G_i is automatically open. This follows from the next quite general (and presumably well-known) lemma, which will be used several times in the remainder of the paper.

Lemma 5.2 *Let G be an étale groupoid.*

- (i) *If K, H are open (respectively compact, respectively relatively compact) subsets of G , then*

$$K \cdot H := \{kh \in G \mid k \in K, h \in H, s(k) = r(h)\}$$

is open (respectively compact, respectively relatively compact).

- (ii) *If $K \subseteq G$ is open, then the subgroupoid of G generated by K is also open, and is itself an étale groupoid.*

Proof For part (i), assume first that K and H are open, and let kh be a point of KH . As G is étale there are neighborhoods $V \subseteq K$ of k and $W \subseteq H$ of h such that r and s both restrict to homeomorphisms on V and W . Define $V' := (s|_V)^{-1}(s(V) \cap r(W))$, $W' := (r|_W)^{-1}(s(V) \cap r(W))$, and

$$U := \{k'h' \mid k \in V', h \in W', s(k') = r(h')\};$$

note that U is a subset of KH . As s and r both restrict to homeomorphisms on V' and W' , r restricts to a homeomorphism from U to $r(V')$. In particular, U contains an open neighborhood of kh , so we are done. Assume next that K and H are compact. It is clear from the corresponding property for K and H that any net in $K \cdot H$ has a convergent subnet, so $K \cdot H$ is compact. Finally, assume that K and H are relatively compact. Clearly $K \cdot H \subseteq \overline{K} \cdot \overline{H}$, and we have already shown that the latter set is compact.

For (ii), note that if H is any open subset of G then

$$H^n := \{h_n \cdots h_1 \mid h_k \in H \text{ and } s(h_k) = r(h_{k-1}) \text{ for all } k \in \{1, \dots, n\}\}$$

is open by (i). If K is open, then so too is $K \cup K^{-1} \cup r(K) \cup s(K)$ and the subgroup generated by K is equal as a set to

$$\bigcup_{n=1}^{\infty} (K \cup K^{-1} \cup r(K) \cup s(K))^n$$

and thus open. An open subgroupoid of an étale groupoid is easily seen to be itself étale, so this completes the proof. \square

We give a first, easy, example: it is a groupoid analogue of Remark 2.2, parts (i) and (ii).

Example 5.3 A groupoid is *locally finite* if it is the union of open, relatively compact subgroupoids. It is not difficult to check directly from Definition 5.1 that a groupoid has dynamic asymptotic dimension zero if and only if it is locally finite in this sense.

In particular, if $G = G^{(0)}$ is just a space, then G has dynamic asymptotic dimension zero.

We now show that Definition 5.1 generalizes our earlier definition for group actions. An action $\Gamma \curvearrowright X$ gives rise to an associated *transformation groupoid* $\Gamma \rtimes X$ as follows. To fix notation, recall that as a set $\Gamma \rtimes X$ is defined by

$$\Gamma \rtimes X := \{(gx, g, x) \mid x \in X, g \in \Gamma\}.$$

Note that the projection onto the last two variables is a bijection from $\Gamma \rtimes X$ to $\Gamma \times X$; $\Gamma \rtimes X$ is given the topology that makes this bijection a homeomorphism. The unit space of $G^{(0)}$ is $\{(x, e, x) \mid x \in X\}$ and the source and range maps are given by

$$r(gx, g, x) = gx, \quad s(gx, g, x) = x.$$

Composition and inverse are defined by

$$(ghx, g, hx)(hx, h, x) = (ghx, gh, x), \quad (gx, g, x)^{-1} = (x, g^{-1}, gx)$$

respectively.

Lemma 5.4 *An action $\Gamma \curvearrowright X$ has dynamic asymptotic dimension d in the sense of Definition 2.1 if and only if the corresponding transformation groupoid $G = \Gamma \rtimes X$ has dynamic asymptotic dimension d in the sense of Definition 5.1.*

Proof Say an action $\Gamma \curvearrowright X$ has dynamic asymptotic dimension d , and let K be an open relatively compact subset of the transformation groupoid $\Gamma \rtimes X$. Any open relatively compact subset of G is contained in a (relatively compact, open) set of the form $\{(gx, g, x) \in G \mid x \in K', g \in E\}$ for some finite subset E of Γ such that $e \in E$ and $E = E^{-1}$, and some open relatively compact subset K' of X . Hence we may assume that K has this form. Let U'_0, \dots, U'_d be open subsets of $G^{(0)}$ that cover $\overline{K'}$ and are such that the set

$$F_i := \left\{ g \in \Gamma \left| \begin{array}{l} \text{there exist } x \in U'_i \text{ and } g_n, \dots, g_1 \in E \text{ such that} \\ g = g_n \cdots g_2 g_1 \text{ and for each } k \in \{1, \dots, m\}, \\ g_k \cdots g_1 x \in U'_i \end{array} \right. \right\}$$

is finite for each i . Define $U_i := U'_i \cap K'$, so U_1, \dots, U_d are open subsets of $G^{(0)} = X$ that cover K' . Then the subgroupoid of G generated by

$$\{(gx, g, x) \in G \mid x, gx, \in U_i\}$$

is contained in $\{(gx, g, x) \in G \mid g \in F_i, x \in K'\}$, and is thus relatively compact.

The converse can be proved in a very similar way. □

6 Example: coarse spaces with finite asymptotic dimension

In this section we show that the coarse groupoid $G(X)$ associated to a bounded geometry coarse space X has dynamic asymptotic dimension d if and only if X has asymptotic dimension d . This example (in the special case X is the coarse space underlying a group Γ) was our original motivation, and also motivates the terminology ‘dynamic asymptotic dimension’.

We recall the following definition: compare for example [31, Chapter 2].

Definition 6.1 Let X be a set. A *coarse structure* on X is a collection \mathcal{E} of subsets of $X \times X$ called *controlled sets* such that:

- (i) the diagonal $\{(x, x) \mid x \in X\}$ is contained in \mathcal{E} ;
- (ii) if E is in \mathcal{E} and F is a subset of E , then F is in \mathcal{E} ;
- (iii) if E and F are in \mathcal{E} , then their union $E \cup F$ is in \mathcal{E} ;
- (iv) if E is in \mathcal{E} then

$$E^{-1} := \{(x, y) \in X \times X \mid (y, x) \in E\}$$

is in \mathcal{E} ;

- (v) if E and F are in \mathcal{E} , then

$$E \circ F := \{(x, y) \in X \times X \mid \text{there exists } z \in X \text{ such that } (x, z) \in E, (z, y) \in F\}$$

is in \mathcal{E} .

A set X equipped with a coarse structure is called a *coarse space*.

A coarse space X has *bounded geometry* if for each controlled set E the cardinals

$$\sup_{x \in X} |\{y \in X \mid (x, y) \in E\}|, \sup_{x \in X} |\{y \in X \mid (y, x) \in E\}|$$

are finite.

The reader should keep the following example of a coarse space in mind. If (X, d) is a discrete metric space, a *controlled tube* is defined to be a set of the form

$$E := \{(x, y) \in X \times X \mid d(x, y) < r\}$$

for some $r \geq 0$, and the controlled sets are defined to be those subsets E of $X \times X$ that are contained in a controlled tube. The bounded geometry condition corresponds to the following assumption: for any $r > 0$, there is a uniform bound on the cardinality of all balls of radius r in X .

We recall the following definition, due originally to Gromov [14, Section 1.E] in the special case that the coarse structure comes from a metric as above. See for example [31, Chapter 9], [24, Chapter 2], or [4, Section 12] for more information.

Definition 6.2 Let X be a coarse space and E a controlled set for X . A cover $\mathcal{U} = \{U_i\}_{i \in I}$ of X is:

- (i) E -separated if whenever $i \neq j$, $U_i \times U_j \cap E = \emptyset$;
- (ii) E -bounded if each set $U_i \times U_i$ is contained in E .

Definition 6.3 Let X be a coarse space. Then X has *asymptotic dimension* d if d is the smallest number with the following property: for any controlled set E there exists a controlled set F and a cover $\mathcal{U} = \{U_i\}_{i \in I}$ of X such that \mathcal{U} is F -bounded and such that \mathcal{U} admits a decomposition

$$\mathcal{U} = \mathcal{U}_0 \sqcup \dots \sqcup \mathcal{U}_d$$

such that each \mathcal{U}_i is E -separated.

Now, let X be a bounded geometry coarse space. Skandalis et al. [35] (see also [31, Chapter 10]) associate a groupoid to X as follows. For each controlled set E for X , let \overline{E} denote its closure in the product $\beta X \times \beta X$ of the Stone-Ćech compactification of X with itself. Define

$$G(X) := \bigcup_{E \in \mathcal{E}} \overline{E},$$

and equip $G(X)$ with the weak topology coming from this union, i.e., a subset U of $G(X)$ is open if and only if its intersection with each \overline{E} is open in the natural compact topology on \overline{E} . This is a locally compact Hausdorff topology for which each \overline{E} is a compact open subset of $G(X)$. It is not difficult to see that if (ω, η) are in \overline{E} , and (η, ζ) are in \overline{F} , then (ω, ζ) is in $\overline{E \circ F}$, whence it follows that $G(X)$ inherits a groupoid structure from the pair groupoid structure on $\beta X \times \beta X$ (its topology, however, is not the same as the one it inherits from $\beta X \times \beta X$). The groupoid operations satisfy all the necessary continuity axioms to show that $G(X)$ is a (locally compact, Hausdorff) étale groupoid.

Theorem 6.4 A bounded geometry coarse space X has asymptotic dimension d if and only if the associated coarse groupoid $G(X)$ has dynamic asymptotic dimension d .

Moreover, if X has asymptotic dimension at most d , then $G(X)$ has dynamic asymptotic dimension at most d in the following slightly stronger form: for any open, relatively compact subset K of $G(X)$, there exists an open cover $\{U_0, \dots, U_d\}$ of $G(X)^{(0)}$ such that the subset

$$\{g \in K \mid s(g), r(g) \in U_i\}$$

is contained in a compact open subgroupoid of $G(X)$.

Proof Assume first that $G(X)$ has dynamic asymptotic dimension d . Fix a controlled set E as in the definition of asymptotic dimension, which we may assume contains the diagonal, and let $K = \overline{E}$, a compact open subset of $G(X)$ that contains the unit space $G(X)^{(0)} = \beta X$. Take an open cover U_0, \dots, U_d of $G(X)^{(0)} = \beta X$ with the properties in Definition 5.1 for this K , so for each $i \in \{0, \dots, d\}$ the subgroupoid G_i generated by the set

$$\{g \in \overline{E} \mid s(g), r(g) \in U_i\}$$

is relatively compact. In particular, by definition of the topology on $G(X)$ there exists a controlled set $F \subseteq X \times X$ such that each G_i is contained in $\overline{F} \subseteq G(X)$. Fixing i for the moment, let \sim be the equivalence relation on U_i induced by G_i , so $x \sim y$ if there exists $g \in G_i$ with $s(g) = x$ and $r(g) = y$. Write the equivalence classes for this relation as $\{U_i^j \mid j \in J_i\}$ and define $\mathcal{U}_i := \{U_i^j \cap X \mid j \in J_i\}$ to be the collection of equivalence classes for this relation. Now let $\mathcal{U} = \mathcal{U}_0 \sqcup \dots \sqcup \mathcal{U}_d$ be the collection of all these subsets of X (which covers X as $\{U_0, \dots, U_d\}$ covers βX).

We claim that \mathcal{U} , decomposed as $\mathcal{U} = \mathcal{U}_0 \sqcup \dots \sqcup \mathcal{U}_d$, has the properties required by Definition 6.3. Indeed, for F -boundedness, note that the equivalence relation induced on each $U_i^j \cap X$ by G_i is entirely contained in F (as G_i is contained in \overline{F}). For E -separatedness, note that if for $j \neq k$ there was some (x, y) in $(U_i^j \times U_i^k) \cap E$, then we would have that $x \sim y$, contradicting the fact that U_i^j and U_i^k are distinct equivalence classes.

Conversely, say G has asymptotic dimension at most d . Let K be a compact subset of $G(X)$. The definition of the topology on $G(X)$ implies that $K \subseteq \overline{E}$ for some controlled set E . Let $\mathcal{U} = \mathcal{U}_0 \sqcup \dots \sqcup \mathcal{U}_d$ and F be as in the definition of finite asymptotic dimension for this E . For each i , set

$$U_i := \overline{\bigsqcup_{U \in \mathcal{U}_i} U},$$

which is a compact open subset of βX ; as \mathcal{U} is a cover of X , $\{U_0, \dots, U_d\}$ is a cover of βX . Note that for each i ,

$$G_i := \bigsqcup_{U \in \mathcal{U}_i} U \times U$$

is a subgroupoid of the pair groupoid that is contained (as a set) in F , and by continuity of the groupoid operations, the subgroupoid of $G(X)$ generated by

$$\{g \in E \mid r(g), s(g) \in U_i\}$$

is contained in $\overline{G_i}$, a subgroupoid of $G(X)$ contained (as a set) in the compact subset \overline{F} . Hence the dynamic asymptotic dimension of $G(X)$ is at most d .

The final statement is clear from the construction in the proof. □

Finally in this section, we give two consequences. The first discusses the relationship of the above to the Bartels–Lück–Reich conditions of Sect. 4; the second shows that groups of finite asymptotic dimension all admit free, amenable, minimal actions on the Cantor set which have finite dynamic asymptotic dimension.

Theorem 6.5 *Let Γ be a discrete group. The following are equivalent:*

- (i) Γ admits an action $\Gamma \curvearrowright X$ on a compact space that is d -BLR with respect to the family of finite subgroups.
- (ii) Γ admits an action $\Gamma \curvearrowright X$ on a compact space with dynamic asymptotic dimension at most d .
- (iii) The canonical action of Γ on $\beta\Gamma$ has dynamic asymptotic dimension at most d .
- (iv) Γ equipped with the canonical left-invariant coarse structure for which the controlled sets are

$$\mathcal{E} = \{E \subseteq \Gamma \times \Gamma \mid \{s^{-1}t \in \Gamma \mid (s, t) \in E\} \text{ is finite}\},$$

has asymptotic dimension at most d .

- (v) The canonical action of Γ on $\beta\Gamma$ is d -BLR with respect to the family of finite subgroups.

Proof The fact that (i) implies (ii) (for the same X) is Theorem 4.7.

For (ii) implies (iii), assume that $\Gamma \curvearrowright X$ is an action on a compact space with dynamic asymptotic dimension at most d . Let $\Gamma \rightarrow X$ be any orbit map, and let $\phi : \beta\Gamma \rightarrow X$ be the canonical extension to the Stone–Čech compactification of Γ . Pulling back covers along ϕ shows that $\Gamma \curvearrowright \beta\Gamma$ also has dynamic asymptotic dimension at most d .

For (iii) implies (iv), note that the transformation groupoid $\Gamma \ltimes \beta\Gamma$ has dynamic asymptotic dimension at most d by Lemma 5.4. On the other hand, this transformation groupoid canonically identifies with the coarse groupoid of Γ by [35, Proposition 3.4], so the result follows from Theorem 6.4.

The fact that (iv) implies (v) is essentially due to Higson and Roe, and very similar to the proof of Proposition 4.5. The result follows from combining the arguments of [18, Lemma 4.3] and [18, Section 3]: indeed, these give an (E, ϵ) -equivariant map $\beta\Gamma \rightarrow P_d(\Gamma)$ for any finite $E \subseteq \Gamma$ and any $\epsilon > 0$.

Finally, (v) implies (i) is trivial. □

The following theorem is a result of combining Theorem 6.4 with ideas of Rørdam and Sierakowski from [33, Section 6].

Theorem 6.6 *Let Γ be a countably infinite discrete group with asymptotic dimension d . Then Γ admits a free, minimal action on the Cantor set which has dynamic asymptotic dimension at most d .*

Proof Using Theorem 6.5, the action of Γ on $\beta\Gamma$ is d -BLR for the family \mathcal{F} of finite subgroups of Γ . Let

$$E_1 \subseteq E_2 \subseteq E_3 \subseteq \dots$$

be a nested sequence of finite subsets of Γ such that $\Gamma = \bigcup E_n$. For each n , let $f_n : \beta\Gamma \rightarrow C_n$ be a $(E_n, 1/n)$ -equivariant map to a (Γ, \mathcal{F}) simplicial complex of dimension at most d . As $\beta\Gamma$ is compact, we may assume by Remark 4.4 that the image of each f_n is contained in a set of the form $P(S_n) \cap C_n$, where S_n is a finite subset of the vertex set of C_n . Hence, replacing C_n by $C_n \cap P(\Gamma \cdot S_n)$, we may assume that there are only finitely many Γ -orbits of vertices of C_n , and thus in particular that the vertex set V_n of C_n is countable.

Now, let A denote a unital Γ -invariant C^* -subalgebra of $l^\infty(\Gamma)$. Let \widehat{A} denote the spectrum of A , which by Gelfand duality identifies with a quotient space of $\beta\Gamma$ such that the quotient map $\beta\Gamma \rightarrow \widehat{A}$ is equivariant. For each n , write

$$f_n(x) = \sum_{v \in V_n} t_v(x)v,$$

where each $t_v : \beta\Gamma \rightarrow [0, 1]$ is continuous. Define $\Omega_1 := \{t_v \mid v \in V_n, n \in \mathbb{N}\}$, which is a countable subset of $l^\infty(\Gamma)$. If A contains Ω_1 , then the action $\Gamma \curvearrowright \widehat{A}$ is d -BLR: indeed, the functions t_v then descend to the quotient space \widehat{A} of $\beta\Gamma$, whence the maps $f_n : \beta\Gamma \rightarrow C_n$ descend to $f_n : \widehat{A} \rightarrow C_n$ (this argument is inspired by [17, Lemma 3.5]). On the other hand, [33, Lemma 6.4] shows that there is a countable subset Ω_2 of $l^\infty(\Gamma)$ such that if A contains Ω_2 , then the action of Γ on \widehat{A} is free.

Define $\Omega := \Omega_1 \cup \Omega_2$, a countable subset of $l^\infty(\Gamma)$. Rørdam and Sierakowski [33, Lemma 6.7] show there is a countable Γ -invariant collection of projections \mathcal{P} such that the C^* -algebra generated by \mathcal{P} contains Ω . Let A be the C^* -algebra generated by \mathcal{P} and the unit of $l^\infty(G)$. It follows from the above discussion that \widehat{A} is a Γ -space, and that the action of Γ is free and d -BLR. Moreover, as A is unital and generated by countably many projections, the space \widehat{A} is compact, metrizable, and totally disconnected.

We now have that Γ admits a free, d -BLR action on a totally disconnected, metrizable compact space $Y := \widehat{A}$. Note that if Z is any non-empty closed Γ -invariant subset of Y , then Z and the induced Γ action on it will still have all these properties. Inclusion defines a partial order on the closed non-empty Γ -invariant subsets of Y ; compactness implies that any descending chain for this order has non-empty intersection, and thus Zorn's lemma implies there exists a minimal element X . It follows easily that the induced Γ action on X is minimal. We now have that Γ admits a free, minimal, d -BLR action on a totally disconnected, metrizable compact space X .

To complete the proof, it suffices to show that this X is a copy of the Cantor set, and for this it suffices to show that it has no isolated points. Note then that if $x \in X$ is isolated then the orbit of $\Gamma \cdot x$ is open, and thus $X \setminus \Gamma \cdot x$ is empty by minimality. As the action is free and Γ is infinite, the open cover $\{\{gx\} \mid g \in \Gamma\}$ of $X = \Gamma \cdot x$ has no finite subcover which contradicts compactness. Hence no isolated point can exist, and we are done. □

7 Partitions of unity

In this section we prove a technical result showing that dynamic asymptotic dimension gives rise to partitions of unity which are ‘almost’ invariant in an appropriate sense. This will be important in Sect. 8.

Proposition 7.1 *Let G be an étale groupoid with compact unit space, and with dynamic asymptotic dimension d . Then for any open relatively compact subset K of G and any $\epsilon > 0$ there exists an open cover $\{U_0, \dots, U_d\}$ of $r(K) \cup s(K)$ with the following properties.*

(i) *For each i , the set*

$$\{g \in K \mid s(g), r(g) \in U_i\}$$

is contained in an (open and) relatively compact subgroupoid of G .

(ii) *For each $x \in r(K) \cup s(K)$, the ‘partial orbit’ $s(r^{-1}(x) \cap K)$ is completely contained in some U_i .*

(iii) *There exists a collection of continuous functions $\{\phi_i : G^{(0)} \rightarrow [0, 1]\}_{i=0}^d$ such that the support of each ϕ_i is contained in U_i , such that for all $x \in r(K) \cup s(K)$ we have $\sum_{i=0}^d (\phi_i(x))^2 = 1$, and such that for any $g \in K$ and each i*

$$|\phi_i(s(g)) - \phi_i(r(g))| < \epsilon.$$

For the convenience of readers who are mainly interested in the case of group actions on compact spaces, we spell out here what this says in the special case where G is a transformation groupoid in the language of actions. The proposition is not really any easier to prove in this special case, however.

Corollary 7.2 *Let $\Gamma \curvearrowright X$ be an action with X compact, and with dynamic asymptotic dimension d . Then for any finite subset E of Γ and any $\epsilon > 0$ there exists an open cover $\{U_0, \dots, U_d\}$ of X with the following properties.*

(i) *For each i , the set*

$$\left\{ g \in \Gamma \left| \begin{array}{l} \exists x \in U_i \text{ and } g_n, \dots, g_1 \in E \text{ such that } g = g_n \cdots g_1 \\ \text{and } \forall k \in \{1, \dots, n\}, g_k \cdots g_1 x \in U_i \end{array} \right. \right\}$$

is finite.

(ii) *For each $x \in X$, the collection $E \cdot x := \{g^{-1}x \mid g \in E\}$ is completely contained in some U_i .*

(iii) *There exists a collection of continuous functions $\{\phi_i : X \rightarrow [0, 1]\}_{i=0}^d$ on X such that the support of each ϕ_i is contained in U_i , such that $\sum_{i=0}^d (\phi_i(x))^2 = 1$ for all $x \in X$, and such that for any $g \in E$ and each i*

$$\sup_{x \in X} |\phi_i(gx) - \phi_i(x)| < \epsilon.$$

□

The proof of Proposition 7.1 is related to that of Proposition 4.5. We wanted to keep the two proofs separate for the convenience of readers who are only interested in one or the other case, and as there are enough significant differences that it did not seem possible to make a ‘combined’ proof that was much shorter.

We start with a lemma.

Lemma 7.3 *Say G is an étale groupoid with dynamic asymptotic dimension d . Then for any open relatively compact subset K of G there is a cover $\{U_0, \dots, U_d\}$ of $r(K) \cup s(K)$ by relatively compact open subsets of $G^{(0)}$ such that for each i , the set $\{g \in K \mid s(g), r(g) \in U_i\}$ generates an (open and) relatively compact subgroupoid of G , and moreover so that for each $x \in G^{(0)}$ the set $s(r^{-1}(x) \cap K)$ is completely contained in some U_i .*

Proof Replacing K with $K \cup K^{-1} \cup r(K) \cup s(K)$ (which is open and relatively compact as r and s are open, continuous maps), we may assume that $K = K \cup K^{-1} \cup r(K) \cup s(K)$. Define

$$K^3 := \{g \in G \mid \text{there are } g_1, g_2, g_3 \in K \text{ such that } g = g_1g_2g_3\},$$

which is again an open relatively compact subset of G by Lemma 5.2. Let $\{V_0, \dots, V_d\}$ be an open cover of $r(K^3) \cup s(K^3)$ such that the subgroupoid G_i of G generated by

$$\{g \in K^3 \mid s(g), r(g) \in V_i\} \tag{7.1}$$

is relatively compact. For $i \in \{0, \dots, d\}$, define

$$U_i := s(K \cap r^{-1}(V_i)) \cap (r(K) \cup s(K)),$$

which is open and relatively compact. Note that as $r(K) \cup s(K) \subseteq K$, each U_i contains $V_i \cap (r(K) \cup s(K))$; thus we have

$$\begin{aligned} \bigcup_{i=0}^d U_i &\supseteq \bigcup_{i=0}^d V_i \cap (r(K) \cup s(K)) \supseteq (r(K^3) \cup s(K^3)) \cap (r(K) \cup s(K)) \\ &= r(K) \cup s(K). \end{aligned}$$

It remains to show that for each fixed i , the set $\{g \in K \mid s(g), r(g) \in U_i\}$ generates a relatively compact subgroupoid of G . Indeed, say g is in the subgroupoid generated by this set. Then (recalling that $K = K \cup K^{-1} \cup r(K) \cup s(K)$) there exists a finite sequence g_n, \dots, g_1 of composable elements from $\{g \in K \mid s(g), r(g) \in U_i\}$ such that $g = g_n \cdots g_1$. As for each $k \in \{1, \dots, n\}$ we have that $s(g_k) \in U_i$, there exists $h_k \in K$ such that $r(h_k) \in V_i$, and $s(h_k) = s(g_k)$. Moreover, as $s(g_n) \in U_i$, there exists $h_{n+1} \in K$ such that $s(h_{n+1}) = s(g_n)$ and such that $r(h_{n+1}) \in V_i$. For each $k \in \{1, \dots, n\}$, define

$$g'_k := h_{k+1}g_k h_k^{-1},$$

and note that $g'_k \in \{g \in K^3 \mid s(g), r(g) \in V_i\}$. On the other hand, $g = h_{n+1}^{-1}g'_n \cdots g'_1h_1$. If G_i is the subgroupoid of G generated by the set in line (7.1), it follows that g is in the subset

$$K \cdot G_i \cdot K := \{h \in G \mid \text{there are } h_0, h_2 \in K \text{ and } h_1 \in G_i \\ \text{such that } h = h_0h_1h_2\};$$

by relative-compactness of G_i and K , and Lemma 5.2, this set is relatively compact, so we are done. □

We need one more technical lemma before giving the proof of Proposition 7.1; in some sense it is an iterated version of Lemma 7.3 above.

Lemma 7.4 *Say G is an étale groupoid with dynamic asymptotic dimension d . Let K be an open relatively compact subset of G , and let N be a fixed natural number. Then for each $i \in \{0, \dots, d\}$ there is a nested collection*

$$U_i^{(0)} \subseteq U_i^{(1)} \subseteq \dots \subseteq U_i^{(N+1)}$$

of open, relatively compact subsets of $G^{(0)}$ with the following properties.

- (i) *The collection $\{U_0^{(0)}, \dots, U_d^{(0)}\}$ covers $\overline{r(K)} \cup \overline{s(K)}$.*
- (ii) *For all i, n , $\overline{U_i^{(n)}} \subseteq U_i^{(n+1)}$.*
- (iii) *For all i, n ,*

$$s(K \cap r^{-1}(U_i^{(n)})) \subseteq U_i^{(n+1)}.$$

- (iv) *For all i , the set*

$$\{g \in K \mid s(g), r(g) \in U_i^{(N+1)}\}$$

generates a relatively compact subgroupoid of G .

Proof Replacing K with $K \cup K^{-1} \cup r(K) \cup s(K)$, we may assume that $K = K \cup K^{-1} \cup r(K) \cup s(K)$. For each n , define

$$\overline{K}^n := \{g_n \cdots g_1 \mid g_k \in \overline{K} \text{ and } s(g_{k+1}) = r(g_k) \text{ for all } k\}.$$

Note that \overline{K}^n is compact for all n by Lemma 5.2, and that $\overline{K}^n \subseteq \overline{K}^{n+1}$ for all n by our assumption that $K = K \cup K^{-1} \cup r(K) \cup s(K)$. As G is locally compact, there exists an open, relatively compact subset $K' \subseteq G$ that contains \overline{K}^{N+1} . Let $\{V_0, \dots, V_d\}$ be an open cover of $r(K') \cup s(K')$ with the properties in Lemma 7.3 for the open relatively compact set K' . Replacing each V_i with its intersection with $r(K') \cup s(K')$, we may assume that each V_i is contained in $r(K') \cup s(K')$.

Now, for each $i \in \{0, \dots, d\}$ and each $n \in \{0, \dots, N + 1\}$, define

$$V_i^{(n)} := \{x \in V_i \mid s(\overline{K}^{N+1-n} \cap r^{-1}(x)) \subseteq V_i\}.$$

As the sets \overline{K}^n are nested, this gives a nested sequence of subsets

$$V_i^{(0)} \subseteq V_i^{(1)} \subseteq \dots \subseteq V_i^{(N+1)}$$

of $G^{(0)}$. As for each $x \in G^{(0)}$, $s(r^{-1}(x) \cap K') \subseteq V_i$ for some i , the collection $\{V_0^{(0)}, \dots, V_d^{(0)}\}$ covers $r(K') \cup s(K')$. We claim moreover that each $V_i^{(n)}$ is open. Indeed, if this does not happen, then we may find a net (x_λ) in $G^{(0)}$ converging to some $x \in V_i^{(n)}$ such that for each λ there exists $g_\lambda \in \overline{K}^{N+1-n} \cap r^{-1}(x_\lambda)$ with $s(g_\lambda) \notin V_i$. As \overline{K}^{N+1-n} is compact we may assume by passing to a subnet that (g_λ) converges to some $g \in \overline{K}^{N+1-n}$. As r is continuous, g is in $\overline{K}^{N+1-n} \cap r^{-1}(x)$, and as V_i is open, $s(g)$ is not in V_i . This contradicts that x is in $V_i^{(n)}$, so $V_i^{(n)}$ is open as claimed.

We may now define the sets $U_i^{(n)}$. As the set $\overline{r(K') \cup s(K')}$ is compact, it is normal. It follows that there are open subsets $U_0^{(0)}, \dots, U_d^{(0)}$ of $r(K') \cup s(K')$ such that $\overline{U_i^{(0)}} \subseteq V_i^{(0)}$ for each i and such that the collection $\{U_0^{(0)}, \dots, U_d^{(0)}\}$ covers $\overline{r(K') \cup s(K')}$. Note that for each i ,

$$\begin{aligned} \overline{s(K \cap r^{-1}(U_i^{(0)}))} &\subseteq \overline{s(K \cap r^{-1}(U_i^{(0)}))} \subseteq \overline{s(\overline{K} \cap r^{-1}(U_i^{(0)}))} \\ &\subseteq \overline{s(\overline{K} \cap r^{-1}(\overline{U_i^{(0)}}))} \subseteq \overline{s(\overline{K} \cap r^{-1}(V_i^{(0)}))}. \end{aligned}$$

It is not difficult to see that this last set is contained in $V_i^{(1)}$, however. Moreover, $\overline{U_i^{(0)}}$ is contained in $V_i^{(0)} \subseteq V_i^{(1)}$ by assumption. Hence by normality of the compact set $\overline{V_i}$, there exists an open set $U_i^{(1)}$ containing $\overline{s(K \cap r^{-1}(U_i^{(0)}))} \cup \overline{U_i^{(0)}}$ such that $\overline{U_i^{(1)}} \subseteq V_i^{(1)}$. Continuing in this way, we see that

$$\overline{s(K \cap r^{-1}(U_i^{(1)}))} \subseteq V_i^{(2)},$$

and thus there is an open set $U_i^{(2)}$ containing $\overline{s(K \cap r^{-1}(U_i^{(1)}))} \cup \overline{U_i^{(1)}}$, with closure contained in $V_i^{(2)}$, and so on.

The process above gives a nested sequence of open subsets

$$U_i^{(0)} \subseteq U_i^{(1)} \subseteq \dots \subseteq U_i^{(N+1)}$$

for each i such that

$$\overline{s(K \cap r^{-1}(U_i^{(n)}))} \subseteq U_i^{(n+1)} \quad \text{and} \quad \overline{U_i^{(n)}} \subseteq V_i^{(n)}$$

for all n . Note that each $U_i^{(n)}$ is relatively compact as $V_i^{(n)} \subseteq V_i \subseteq r(K') \cup s(K')$ for all n . It remains to check properties (i) through (iv) from the statement. Properties (i), (ii), and (iii) are obvious from the way we chose our sets. Property (iv) follows as

$$\{g \in K \mid s(g), r(g) \in U_i^{(N+1)}\} \subseteq \{g \in K' \mid s(g), r(g) \in V_i\},$$

and the set on the right generates a relatively compact subgroupoid by choice of the sets V_0, \dots, V_d . □

Proof of Proposition 7.1 Let K be an open, relatively compact subset of G and $\epsilon > 0$ be given. Replacing K with $K \cup K^{-1} \cup r(K) \cup s(K)$, we may assume that $K = K^{-1}$, and that K contains its image under r and s . Let N be any natural number larger than 2 such that $(\sqrt{2}(1+\sqrt{d+1}))/\sqrt{N} < \epsilon$, and let $\{U_i^{(n)} \mid i \in \{0, \dots, d\}, n \in \{0, \dots, N+1\}\}$ have the properties in Lemma 7.4. For each $i = 0, \dots, d$, set $U_i = U_i^{(N+1)}$. Condition (i) from Lemma 7.4 combined with the assumptions from the first paragraph of that lemma imply that $\{U_0, \dots, U_d\}$ is an open cover of $r(K) \cup s(K)$. Condition (ii) from the statement of Proposition 7.1 follows condition (iii) from Lemma 7.4 and the fact that $\{U_0^{(0)}, \dots, U_d^{(0)}\}$ covers $r(K) \cup s(K)$. Condition (i) from the statement of Proposition 7.1 follows directly from condition (iv) from Lemma 7.4. It remains to construct a partition of unity with the properties in condition (iii) from the statement of Proposition 7.1.

For each $i \in \{0, \dots, d\}$ and $n \in \{1, \dots, N\}$, let

$$\psi_i^{(n)} : G^{(0)} \rightarrow [0, 1]$$

be any continuous function which is constantly equal to one on $U_i^{(n-1)}$ and constantly equal to zero on $G^{(0)} \setminus U_i^{(n)}$; such functions exist by condition (ii) from Lemma 7.4 and Urysohn’s lemma. For $i \in \{0, \dots, d\}$ set

$$\psi_i = \frac{1}{N} \sum_{n=1}^N \psi_i^{(n)}$$

and define

$$\phi_i = \frac{\psi_i}{\max \left\{ \sqrt{\sum_{j=0}^d \psi_j^2}, 1 \right\}}.$$

Note that condition (i) from Lemma 7.4 implies that for all $x \in r(K) \cup s(K)$, there is at least one $i \in \{0, \dots, d\}$ such that $\psi_i(x) = 1$, whence

$$\sum_{i=0}^d (\phi_i(x))^2 = 1$$

for all $x \in r(K) \cup s(K)$.

Fix now $g \in K$ and $i \in \{0, \dots, d\}$; to complete the proof we must show that

$$|\phi_i(s(g)) - \phi_i(r(g))| < \epsilon. \tag{7.2}$$

For notational convenience, for each $j \in \{0, \dots, d\}$, set $U_j^{(n)} = G^{(0)}$ for $n \geq N + 2$. Define

$$M = M_j := \min\{n \mid r(g) \in U_j^{(n)}\}.$$

Note that

$$\psi_j^{(n)}(r(g)) = \begin{cases} 1 & n \geq M + 1 \\ 0 & n \leq M - 1 \end{cases}$$

and $0 \leq \psi(r(g)) \leq 1$ whence

$$\frac{N - (M + 1)}{N} \leq \psi_j(r(g)) \leq \frac{N - M}{N}. \tag{7.3}$$

Note also that condition (iii) from Lemma 7.4 combined with the fact that $K = K^{-1}$ implies that $s(g)$ is in $U_j^{(M+1)} \setminus U_j^{(M-2)}$ whence

$$\psi_j^{(n)}(s(g)) = \begin{cases} 1 & n \geq M + 2 \\ 0 & n \leq M - 2 \end{cases};$$

as $0 \leq \psi_j^{(n)}(s(g)) \leq 1$ for all values of n , this forces

$$\frac{N - (M + 2)}{N} \leq \psi_j(s(g)) \leq \frac{N - (M - 1)}{N}. \tag{7.4}$$

Combining lines (7.3) and (7.4), we may conclude that

$$|\psi_j(r(g)) - \psi_j(s(g))| \leq \frac{2}{N} \tag{7.5}$$

One the other hand, at least one of the ψ_j is equal to one on each of $s(g)$ and $r(g)$, and therefore

$$1 \leq \sum_{j=0}^d \psi_j(r(g)) \quad \text{and} \quad 1 \leq \sum_{j=0}^d \psi_j(s(g)). \tag{7.6}$$

Hence for our fixed choice of $i \in \{0, \dots, d\}$ and $g \in K$,

$$|\phi_i(r(g)) - \phi_i(s(g))| = \left| \frac{\psi_i(r(g))}{\sqrt{\sum_{j=0}^d \psi_j^2(r(g))}} - \frac{\psi_i(s(g))}{\sqrt{\sum_{j=0}^d \psi_j^2(s(g))}} \right|$$

and so

$$\begin{aligned}
 |\phi_i(r(g)) - \phi_i(s(g))| &\leq \frac{1}{\sqrt{\sum_{j=0}^d \psi_j^2(r(g))}} |\psi_i(r(g)) - \psi_i(s(g))| \\
 &\quad + |\psi_i(s(g))| \left| \frac{1}{\sqrt{\sum_{j=0}^d \psi_j^2(r(g))}} - \frac{1}{\sqrt{\sum_{j=0}^d \psi_j^2(s(g))}} \right|. \tag{7.7}
 \end{aligned}$$

Using lines (7.5) and (7.6) we have

$$\frac{1}{\sqrt{\sum_{j=0}^d \psi_j^2(r(g))}} |\psi_i(r(g)) - \psi_i(s(g))| \leq \frac{2}{N}. \tag{7.8}$$

On the other hand,

$$\begin{aligned}
 &|\psi_i(s(g))| \left| \frac{1}{\sqrt{\sum_{j=0}^d \psi_j^2(r(g))}} - \frac{1}{\sqrt{\sum_{j=0}^d \psi_j^2(s(g))}} \right| \\
 &= \frac{1}{\sqrt{\sum_{j=0}^d \psi_j^2(r(g))}} \frac{1}{\sqrt{\sum_{j=0}^d \psi_j^2(s(g))}} \left| \sqrt{\sum_{j=0}^d \psi_j^2(s(g))} - \sqrt{\sum_{j=0}^d \psi_j^2(r(g))} \right| \\
 &\leq \left| \sqrt{\sum_{j=0}^d \psi_j^2(s(g))} - \sqrt{\sum_{j=0}^d \psi_j^2(r(g))} \right| \\
 &\leq \sqrt{\left| \sum_{j=0}^d \psi_j^2(s(g)) - \sum_{j=0}^d \psi_j^2(r(g)) \right|} \\
 &\leq \sqrt{\sum_{j=0}^d |\psi_j(s(g)) - \psi_j(r(g))| |\psi_j(s(g)) + \psi_j(r(g))|}.
 \end{aligned}$$

Using line (7.5) again thus gives

$$|\psi_i(s(g))| \left| \frac{1}{\sqrt{\sum_{j=0}^d \psi_j^2(r(g))}} - \frac{1}{\sqrt{\sum_{j=0}^d \psi_j^2(s(g))}} \right| \leq \sqrt{\frac{4(d+1)}{N}}. \tag{7.9}$$

Combining lines (7.7), (7.8) and (7.9) therefore gives

$$\begin{aligned}
 |\phi_i(r(g)) - \phi_i(s(g))| &\leq \frac{2}{N} + \sqrt{\frac{4(d+1)}{N}} \leq \sqrt{\frac{2}{N}} + \sqrt{\frac{4(d+1)}{N}} \\
 &= \frac{\sqrt{2}(1 + \sqrt{d+1})}{\sqrt{N}}.
 \end{aligned}$$

which is smaller than ϵ by choice of N . □

8 Nuclear dimension

As usual, we adopt the conventions that all groupoids are locally compact and Hausdorff. Recall that a groupoid G is *free* (such groupoids are also called *principal*) if for each $x \in G^{(0)}$, the *isotropy group* defined by

$$G_x^x := \{g \in G \mid s(g) = r(g) = x\}$$


is trivial.

In this section, we study nuclear dimension for the reduced groupoid C^* -algebra $C_r^*(G)$ of a free groupoid G of finite dynamic asymptotic dimension. Modulo a minor technical assumption that is satisfied for example when G is second countable, our main result—Theorem 8.6 below—says that the nuclear dimension of $C_r^*(G)$ is bounded above by $(d + 1)(N + 1) - 1$, where d is the dynamic asymptotic dimension of G , and N is the covering dimension of $G^{(0)}$. The result is inspired by (and implies) the Winter-Zacharias result that the nuclear dimension of the uniform Roe algebra associated to a bounded geometry metric space is at most the asymptotic dimension of the space [42, Theorem 8.5]. See the end of the section for some other corollaries.

We first recall the definitions of nuclear dimension, covering dimension, and of the reduced C^* -algebra of a groupoid. More details on nuclear dimension can be found in the paper of Winter and Zacharias [42] that introduces the notion. We do not really need a definition of covering dimension, but record it as there is some ambiguity about which definition one should use for non-metrizable spaces. Accessible introductions to groupoid C^* -algebras, that largely focus on the case of interest here, can be found in [30, Section 2.3] and [6, Section 5.6].

Definition 8.1 A completely positive map $\phi : A \rightarrow B$ is *order zero* if it preserves orthogonality: in other words, if a_1, a_2 are positive elements of A such that $a_1 a_2 = 0$, then $\phi(a_1)\phi(a_2) = 0$.

Let A be a C^* -algebra. The *nuclear dimension* of A is the smallest integer $d \in \mathbb{N}$ with the following property. For any finite subset \mathcal{F} of A and any $\epsilon > 0$ there exist finite dimensional C^* -algebras F_0, \dots, F_d and contractive completely positive maps



such that each Ψ_i is order zero, and such that for all $a \in \mathcal{F}$,

$$\left\| \sum_{i=0}^d (\Psi_i \circ \Phi_i)(a) - a \right\| < \epsilon \|a\|.$$

Definition 8.2 Let X be a paracompact (Hausdorff) topological space. The *covering dimension* of X is the smallest integer $d \in \mathbb{N}$ such that every open cover of X has a refinement \mathcal{V} such that \mathcal{V} splits into a disjoint union

$$\mathcal{V} = \mathcal{V}_0 \sqcup \cdots \sqcup \mathcal{V}_d$$

such that whenever $V \neq W$ are distinct elements of some \mathcal{V}_i , then $V \cap W = \emptyset$.

The definition above is equivalent to several other commonly used definitions of covering dimension on the class of paracompact Hausdorff spaces: compare for example [21, Proposition 1.6] or [26, Remark 4.5].

Definition 8.3 Let G be an étale groupoid. Let $C_c(G)$ denote the vector space of compactly supported complex-valued functions on G , made into a $*$ -algebra via the convolution product and adjoint defined by

$$(f_1 f_2)(g) = \sum_{g_1 g_2 = g} f_1(g_1) f_2(g_2), \quad f^*(g) = \overline{f(g^{-1})}.$$

For each $x \in G^{(0)}$ let $l^2(s^{-1}(x))$ denote the Hilbert space of square-summable functions on the source fibre of x , and define a $*$ -representation π_x of $C_c(G)$ on $l^2(s^{-1}(x))$ by

$$(\pi_x(f)\xi)(g) = \sum_{g_1 g_2 = g} f(g_1)\xi(g_2).$$

The *reduced groupoid C^* -algebra of G* is the completion of $C_c(G)$ for the norm

$$\|f\| := \sup_{x \in G^{(0)}} \|\pi_x(f)\|.$$

Before we state the main theorem, we need one (ad-hoc) definition.

Definition 8.4 Let G be an étale groupoid, and H be an open subgroupoid of G . The subgroupoid H is *small* if it is either compact, or second countable and relatively compact.

A groupoid G has *strong dynamic asymptotic dimension at most d* if for any open relatively compact subset K of G there exists an open cover $\{U_0, \dots, U_d\}$ of $s(K) \cup r(K)$ such that for each i , the set

$$\{g \in K \mid s(g), r(g) \in U_i\}$$

is contained in a small subgroupoid of G .

Remark 8.5 (i) A second countable groupoid with dynamic asymptotic dimension at most d automatically has strong dynamic asymptotic dimension at most d . Thus readers who are only interested in the second countable case can just ignore the word ‘strong’ in all the statements below, and replace ‘small’ with ‘open and relatively compact’.

(ii) Theorem 6.4 implies that if X is a bounded geometry metric space of asymptotic dimension d , then the coarse groupoid $G(X)$ has strong dynamic asymptotic dimension d .

- (iii) Say G has strong dynamic asymptotic dimension d . Then Proposition 7.1, point (i) can be strengthened to say that

$$\{g \in K \mid s(g), r(g) \in U_i\}$$

is contained in a small subgroupoid of G for each i : indeed, exactly the same proof gives the stronger statement.

Theorem 8.6 *Let G be a free, étale, groupoid. Assume that G has strong dynamic asymptotic dimension at most d (Definition 8.4), and moreover that the unit space $G^{(0)}$ has topological covering dimension at most N .*

Then the nuclear dimension of the reduced groupoid C^ -algebra $C_r^*(G)$ is at most $(N + 1)(d + 1) - 1$.*

As well as the results subsumed by this theorem discussed in the corollaries at the end of this section, the reader might compare it to [37, Theorem 4.6], which deduces analogous estimates on nuclear dimension from Rokhlin type conditions, and a condition related to asymptotic dimension. We do not know the extent of the overlap between Theorem 8.6 above and [37, Theorem 4.6]; the relationship between the hypotheses of the two results seems worth investigating more carefully.

Most of the rest of this section is devoted to the proof of Theorem 8.6; we give some corollaries at the end. There are two main steps to the proof: an analysis of small subgroupoids, and a reduction to these subgroupoids.

8.1 Small subgroupoids

The goal of this subsection is to prove the following fact about small subgroupoids.

Proposition 8.7 *Let G be a free, étale groupoid. Let H be a small (open) subgroupoid of G in the sense of Definition 8.4, and assume that its unit space $H^{(0)}$ has covering dimension N . Then $C_r^*(H)$ has nuclear dimension² N .*

We will prove this separately in the case where H is compact, and in the case where H is relatively compact and second countable. Before this point though, we need some basic definitions and lemmas that will be used in both cases.

Definition 8.8 Let H be a (locally compact, Hausdorff) groupoid. Let \sim be the equivalence relation on $H^{(0)}$ induced by H , i.e., $x \sim y$ if there is $h \in H$ with $s(h) = x$ and $r(h) = y$. For $x \in H^{(0)}$, let $[x]$ denote its equivalence class. For each $m \in \mathbb{N}$, define

$$H_m^{(0)} := \{x \in H^{(0)} \mid |[x]| = m\}.$$

Finally, let $H^{(0)}/H$ and $H_m^{(0)}/H$ denote the spaces of equivalence classes equipped with the quotient topology in inherited from $H^{(0)}$.

² As will be clear from the proof, one can replace nuclear dimension with decomposition rank here, but we will not need this distinction.

Lemma 8.9 *Let H be a topological groupoid. Then the quotient map $\pi : H^{(0)} \rightarrow H^{(0)}/H$ is open.*

Proof Say U is an open subset of $H^{(0)}$. Then as s is continuous and r is open, the set $r(s^{-1}(U))$ is open in $H^{(0)}$. However, it is clear that $\pi^{-1}(\pi(U)) = r(s^{-1}(U))$, and thus by definition of the quotient topology, $\pi(U)$ is open. \square

Lemma 8.10 *Let G be an étale groupoid and H be a relatively compact subgroupoid of G . Then there exists $M \in \mathbb{N}$ such that $H_m^{(0)}$ is empty for all $m > M$.*

Proof Note that as G is étale, each $g \in G$ is contained in an open neighbourhood U on which both r and s are injective. As the closure \overline{H} is compact, there are finitely many of these neighbourhoods covering \overline{H} , say U_1, \dots, U_M . For each $x \in H^{(0)}$ we have $|[x]| \leq |(s|_H)^{-1}(x)|$; as $(s|_H)^{-1}(x)$ can intersect each U_i at most once, however, $|(s|_H)^{-1}(x)|$ is bounded above by M . \square

We will need the following classical theorem from dimension theory: see [26, Proposition 2.16].

Proposition 8.11 *Let X and Y be paracompact Hausdorff topological spaces. Let $\pi : X \rightarrow Y$ be a continuous open surjection such that for all $y \in Y$, $\pi^{-1}(y)$ is finite. Then the covering dimensions of X and Y are equal.* \square

At this point, we specialize to the case of Proposition 8.7 where H is compact; we will come back to the second countable and relatively compact case later.

Lemma 8.12 *Let H be a compact étale groupoid, and assume that the covering dimension of $H^{(0)}$ is N . Then $H^{(0)}/H$ is compact and Hausdorff, and the covering dimension of $H^{(0)}/H$ is exactly N .*

Proof As H is compact and $H^{(0)}$ is a closed subspace of H , $H^{(0)}$ is compact. Hence $H^{(0)}/H$ is a quotient space of a compact space, so compact. To see that $H^{(0)}/H$ is Hausdorff, note that the equivalence relation \sim on $H^{(0)}$ induced by H is equal to

$$(r \times s)(H) \subseteq H^{(0)} \times H^{(0)},$$

and is thus compact and in particular closed as $H^{(0)} \times H^{(0)}$ is Hausdorff. It is a standard fact that the quotient of a compact space by an equivalence relation that is closed in this sense is Hausdorff: see for example [40, Proposition 2.1]. The claim on covering dimension now follows immediately from Proposition 8.11 applied to the quotient map $\pi : H^{(0)} \rightarrow H^{(0)}/H$. \square

Proof of Proposition 8.7 when H is compact As H is compact and G is free, H is a free and proper groupoid. Hence $C_r^*(H)$ is Morita equivalent to $C(H^{(0)}/H)$ by [23, Example 2.5 and Theorem 2.8].³ Hence the nuclear dimension of $C_r^*(H)$ is equal to the covering dimension of $H^{(0)}/H$ by [42, Proposition 2.4 and Corollary 2.8], and this is N by Lemma 8.12. \square

³ The cited paper only covers the second countable case, but the second countability assumption is unnecessary when the groupoid is étale: see [13].

We now turn to the case of Proposition 8.7 when H is second countable with compact closure. We start by recalling a result of Winter, which needs the following notation. If A is a C^* -algebra then $\text{Prim}(A)$ denotes the collection of all kernels of irreducible representations of A , equipped with the hull-kernel topology: this is defined by saying that a subset S of $\text{Prim}(A)$ is closed if there exists an ideal I of A such that $S = \{J \in \text{Prim}(A) \mid J \supseteq I\}$. Recall moreover that for $m \in \mathbb{N}$, $\text{Prim}_m(A)$ denotes the subspace of $\text{Prim}(A)$ consisting of kernels of m -dimensional representations; by [8, Proposition 3.6.4], $\text{Prim}_m(A)$ is locally compact and Hausdorff when equipped with the subspace topology.

Here then is a (weak version of) Winter’s result: see [41, Theorem 1.6].

Theorem 8.13 *Let A be a separable C^* -algebra such that all irreducible representations of A have dimension at most M for some $M \in \mathbb{N}$, and assume that N is the maximal covering dimension of the spaces $\text{Prim}_m(A)$ for $m \in \{1, \dots, M\}$. Then the nuclear dimension of A is N .*

Proof Winter shows in [41, Theorem 1.6] that the decomposition rank of $C_r^*(H)$ is exactly N under the stated assumptions. Decomposition rank is (trivially—see [42, Remarks 2.2 (ii)]) an upper bound for the nuclear dimension, and it follows from [42, Proposition 2.9 and Corollary 2.10] that the nuclear dimension of $C_r^*(H)$ is bounded below by the maximum of the covering dimensions of the spaces $\text{Prim}_m(C_r^*(H))$.

Clearly the remaining case of Proposition 8.7 follows from this theorem of Winter and the following result.

Lemma 8.14 *Let H be a relatively compact, open subgroupoid of a free étale groupoid. Then $\text{Prim}(C_r^*(H))$ is naturally homeomorphic to $H^{(0)}/H$, via a homeomorphism that takes $\text{Prim}_m(A)$ to $H_m^{(0)}/H$. Moreover, the spaces $H_m^{(0)}/H$ are locally compact and Hausdorff, and the maximum of their covering dimensions equals the covering dimension of $H^{(0)}$.*

For the proof of the lemma, we need a little more notation, and some preliminary lemmas about ideals in groupoid C^* -algebras. Much of the following material seems likely to be standard for experts in groupoid C^* -algebras: for example, the results of [28, Pages 101–103] and [29, Corollary 4.9] are closely connected to what follows. However, we could not find complete proofs of exactly what we need in the literature, so give direct arguments below.

Definition 8.15 A subset U of $H^{(0)}$ is *invariant* if whenever $x \in U$ and $x \sim y$, we have $y \in U$.

We write $\mathcal{O}(H^{(0)})$ for the collection of all invariant open sets in $H^{(0)}$, and $\mathcal{I}(C_r^*(H))$ for the collection of all ideals in $C_r^*(H)$. Both of these sets are equipped with the partial orders defined by inclusion.

Lemma 8.16 *Let H be an étale groupoid. For a subset S of $C_r^*(H)$, write $\langle S \rangle$ for the ideal in $C_r^*(H)$ generated by S . Provisionally define maps by*

$$\Phi : \mathcal{I}(C_r^*(H)) \rightarrow \mathcal{O}(H^{(0)}), \quad I \mapsto \text{Prim}(I \cap C_0(H^{(0)}))$$

and

$$\Psi : \mathcal{O}(H^{(0)}) \rightarrow \mathcal{I}(C_r^*(H)), \quad U \mapsto \langle C_0(U) \rangle.$$

Then Φ and Ψ are well-defined and $\Phi(\Psi(U)) = U$ for all $U \in \mathcal{O}(H^{(0)})$.

Proof It is clear that Ψ is well-defined. To see that Φ is well-defined, we must show that the primitive ideal space $I \cap C_0(H^{(0)})$, which canonically identifies with the open subset

$$\{x \in H^{(0)} \mid \text{there exists } f \in I \cap C_0(H^{(0)}) \text{ with } f(x) \neq 0\}$$

of $H^{(0)}$, is invariant. Say then x is in this subset, and $x \sim y$, so there is $h \in H$ with $r(h) = x$ and $s(h) = y$. Let $f \in I \cap C_0(H^{(0)})$ be such that $f(x) \neq 0$. Let $U \ni h$ be an open subset of H such that the restrictions of r and s to U are injective. Let $\phi : U \rightarrow [0, 1]$ be any compactly supported function such that $\phi(h) = 1$ (so in particular, ϕ is an element of $C_r^*(H)$). Then $\phi f \phi^*$ is in $C_0(H^{(0)}) \cap I$, and is non-zero on y ; thus y is in the primitive ideal space of $C_0(H^{(0)}) \cap I$ as required.

We now check that for $U \in \mathcal{O}(H^{(0)})$ that $\Phi(\Psi(U)) = U$. This is equivalent to showing that

$$\langle C_0(U) \rangle \cap C_0(H^{(0)}) = C_0(U);$$

as the inclusion $\langle C_0(U) \rangle \cap C_0(H^{(0)}) \supseteq C_0(U)$ is obvious, it remains to show the converse inclusion. Say then f is an element of $\langle C_0(U) \rangle \cap C_0(H^{(0)})$ and $x \in H^{(0)}$ is such that $f(x) \neq 0$; we will show that x is in U . As f is in $\langle C_0(U) \rangle$ we may approximate it by elements of the algebraic ideal generated by $C_0(U)$, and thus in particular there must exist $f_i \in C_c(U)$ and $\phi_i, \psi_i \in C_c(H)$ for $i \in \{1, \dots, n\}$ such that

$$\sum_{i=1}^n (\phi_i f_i \psi_i)(x) \neq 0.$$

Hence for some fixed i , we have $(\phi_i f_i \psi_i)(x) \neq 0$. This says that

$$\sum_{g,h,k \in H, x=ghk} \phi_i(g) f_i(h) \psi_i(k) \neq 0.$$

As f_i is supported in U , this implies that there must be $h \in U$ and $k \in H$ with $h = r(k)$ and $x = s(k)$. Hence $x \sim h$, and thus x is in U by invariance. □

Lemma 8.17 *Let H be an open relatively compact subgroupoid of a free étale groupoid G . With notation as in Lemma 8.16, we have $\Psi(\Phi(I)) = I$ for all $I \in \mathcal{I}(C_r^*(H))$.*

Proof Given an ideal I in $C_r^*(H)$, we need to show that $\langle I \cap C_0(H^{(0)}) \rangle = I$. The inclusion $I \supseteq \langle I \cap C_0(H^{(0)}) \rangle$ is obvious, so it remains to show that $\langle I \cap C_0(H^{(0)}) \rangle \supseteq I$.

Say then f is an element of I . Let $\epsilon > 0$ and f_0 be an element of $C_c(H)$ such that $\|f - f_0\|_{C_r^*(H)} < \epsilon$ (note that f_0 need not be an element of I); write $K \subseteq H$ for the support of f_0 .

Fix for the moment $x \in H^{(0)}$ and write $s^{-1}(x) = \{h_1, \dots, h_m\}$ (this set is finite by Lemma 8.10). For each h_i choose an open neighborhood V_i of h_i on which r and s restrict to homeomorphisms and which is such that $V_i \cap V_j = \emptyset$ for all $i \neq j$. As $r(h_i) \neq r(h_j)$ for $i \neq j$, we may further assume that the sets $r(V_1), \dots, r(V_m)$ are mutually disjoint. Set $V_x = \bigcap_{i=1}^m s(V_i)$, an open neighborhood of x .

We now have an open cover $\{V_x\}_{x \in H^{(0)}}$ of $H^{(0)}$, and in particular of the compact subset $s(K)$. Hence by standard results about existence of partitions of unity there is a finite collection $\{\phi_i : H^{(0)} \rightarrow [0, 1] \mid i \in \{1, \dots, n\}\}$ of continuous functions such that each ϕ_i is supported in some compact subset of some V_x , and such that $\sum_{i=1}^n \phi_i(x) = 1$ for all $x \in s(K)$. It follows that $f_0(\sum_{i=1}^n \phi_i) = f_0$ (where the product is the convolution on $C_c(H)$). As the norm of $C_r^*(H)$ restricts to the supremum norm on on the C^* -subalgebra $C_0(H^{(0)})$ this implies that

$$\left\| f\left(\sum_{i=1}^n \phi_i\right) - f \right\|_{C_r^*(H)} \leq \|f - f_0\|_{C_r^*(H)} + \left\| \sum_{i=1}^n \phi_i \right\|_{C_r^*(H)} \|f - f_0\|_{C_r^*(H)} < 2\epsilon.$$

As ϵ was arbitrary, to complete the proof it suffices to show that each element $f\phi_i$ of I is actually in $(I \cap C_0(H^{(0)}))$.

Fix then $\phi = \phi_i$, which is supported in a compact subset K_x of some V_x . Say $s^{-1}(x) = \{h_1, \dots, h_m\}$, and write $s^{-1}(V_x) = \bigsqcup_{j=1}^m V_j$, where V_j is an open neighborhood of h_j such that s and r restrict to homeomorphisms on each V_j , and so that the sets $r(V_1), \dots, r(V_m)$ are mutually disjoint (the existence of such V_j follows from the construction of V_x). For each j , let $\psi_j : H \rightarrow [0, 1]$ be a continuous function supported in V_j , and such that $\psi_j(s^{-1}(K_x) \cap V_j) = \{1\}$ (such a ψ_j exists by Urysohn's lemma). Then $\sum_{j=1}^m \psi_j^* f \phi$ is an element of $C_0(H^{(0)}) \cap I$. Moreover, $f\phi = \sum_{j=1}^m \psi_j \psi_j^* f \phi$, so we are done. □

Lemma 8.18 *Say H is an open relatively compact subgroup of a free étale groupoid G . Then the correspondence from Lemmas 8.16 and 8.17 restricts to a bijection between the collection of primitive ideals in $C_r^*(H)$ and the collection of subsets of $H^{(0)}$ of the form $H^{(0)} \setminus [x]$ for some $x \in H^{(0)}$.*

Proof Note that by Lemma 8.10, any equivalence class $[x]$ of some $x \in H^{(0)}$ is finite, so closed. Hence all maximal open invariant sets are of the form $H^{(0)} \setminus [x]$ for some $x \in H^{(0)}$. Hence by Lemmas 8.16 and 8.17, the maximal ideals in $\mathcal{I}(C_r^*(H))$ are exactly those of the form $\Psi(H^{(0)} \setminus [x])$ for some $x \in H^{(0)}$. Maximal ideals are primitive by [8, Theorem 2.9.7(ii)], so to complete the proof, it suffices to prove that any non-maximal ideal in $C_r^*(H)$ is not primitive.

We first claim that if $[x], [y]$ are distinct equivalence classes in $H^{(0)}$, then there exist $U, V \in \mathcal{O}(H^{(0)})$ such that $[x] \subseteq U, [y] \subseteq V$, and such that for all $z \in U \cap V$, $\|z\| \geq \|x\| + \|y\|$. Indeed, let M be as in Lemma 8.10. Let $s : H \rightarrow H^{(0)}$ be the source map for H (not for the ambient groupoid G), and write $s^{-1}(x) = \{h_1, \dots, h_m\}$ and $s^{-1}(y) = \{g_1, \dots, g_n\}$, where $\|x\| = m, \|y\| = n$ (this is possible as the

groupoid H is free). As $[x] \cap [y] = \emptyset$ and H is Hausdorff there are open sets $U_i \ni g_i$ and $V_j \ni h_j$ such that s restricts to a homeomorphism on each U_i and V_j , such that $U_i \cap U_j = \emptyset = V_i \cap V_j$ for $i \neq j$, and such that $U_i \cap V_j = \emptyset$ for all i, j . Define

$$U_0 := \bigcap_{i=1}^m s(U_i) \quad \text{and} \quad V_0 := \bigcap_{j=1}^n s(V_j)$$

and set

$$U := r(s^{-1}(U_0)) \quad \text{and} \quad V := r(s^{-1}(V_0)).$$

Clearly U and V are open and invariant, and $[x] \subseteq U, [y] \subseteq V$. Consider now $z \in U \cap V$. Then $s^{-1}(z)$ must intersect all the sets U_i and V_j ; as these sets are disjoint, this forces $m + n \leq |s^{-1}(z)|$, and as H is a free groupoid, this forces

$$|[z]| = |s^{-1}(z)| \geq m + n = |[x]| + |[y]|,$$

completing the proof of the claim.

Now, say $V \in \mathcal{O}(H^{(0)})$ is not maximal, so there are $[x] \neq [y]$ with $[x], [y] \subseteq H^{(0)} \setminus V$. Say without loss of generality $|[x]| \geq |[y]| \geq 1$. The claim above implies there exist $U_x, U_y \in \mathcal{O}(H^{(0)})$ such that $[x] \subseteq U_x$ and $[y] \subseteq U_y$, and so that for any $z \in U_x \cap U_y, |[z]| > |[x]|$. If $U_x \cap U_y = \emptyset$, it is not difficult to see that the images of the ideals $\Psi(U_x)$ and $\Psi(U_y)$ are non-zero and orthogonal in $C_r^*(H)/\Psi(V)$ whence $\Psi(V)$ is not primitive⁴ as required. On the other hand, if there is some $z \in U_x \cap U_y$, then we may repeat the process with $[z]$ replacing $[x]$ and $[x]$ replacing $[y]$. Using Lemma 8.10, this process must finish eventually to give orthogonal non-zero ideals in $C_r^*(H)/\Psi(V)$, which is thus not primitive. \square

Proof of Lemma 8.14 It follows from Lemma 8.18 that the map

$$H^{(0)}/H \rightarrow \text{Prim}(A), \quad [x] \mapsto \Psi(H^{(0)} \setminus [x])$$

is a bijection. Lemmas 8.16 and 8.17, the definition of the quotient topology on $H^{(0)}/H$, and the definition of the hull-kernel topology on $\text{Prim}(A)$ imply that this map is a homeomorphism. Moreover, it is not difficult to see that $C_r^*(H)/\Psi(H^{(0)} \setminus [x])$ is isomorphic to the C^* -algebra $M_{|[x]|}(\mathbb{C})$ of $|[x]| \times |[x]|$ matrices over \mathbb{C} , whence it follows that this homeomorphism takes $H_m^{(0)}/H$ onto $\text{Prim}_m(A)$.

Note that the spaces $\text{Prim}_m(C_r^*(H))$ are locally compact and Hausdorff ([8, Proposition 3.6.4]), and that for each $m, \text{Prim}_m(C_r^*(H))$ is open in $\sqcup_{n \leq m} \text{Prim}_n(C_r^*(H))$ ([8, Proposition 3.6.3]). Hence in particular $H_m^{(0)}$ is open in $\sqcup_{n \leq m} H_n^{(0)}$ (one could also prove this directly, of course).

⁴ If A is a C^* -algebra faithfully represented on a Hilbert space H , and I, J are non-zero orthogonal ideals in A , then $I \cdot H$ and $J \cdot H$ are A -invariant non-zero subspaces of H ; in particular, the representation is reducible.

Let N denote the covering dimension of $H^{(0)}$, which we may assume is finite. As covering dimension does not increase on taking open subsets in our context (this follows for example from [42, Proposition 2.5]), it follows inductively that the covering dimension of $H_m^{(0)}$ is at most N for each m ; on the other hand, it follows from [26, Proposition 5.2] (plus second countability) that the covering dimension of $H^{(0)}$ is at most the maximum of the covering dimensions of $H_m^{(0)}$ for $m \in \{1, \dots, M\}$. Hence the maximum of the covering dimensions of $H_m^{(0)}$, $m \in \{1, \dots, M\}$, is exactly N . The claim on the covering dimension of $H_m^{(0)}/H$ now follows from Proposition 8.11 applied to the quotient map $\pi : H^{(0)} \rightarrow H^{(0)}/H$, which is open by Lemma 8.9. \square

8.2 Completion of the proof of Theorem 8.6

For this subsection, G is as in the assumptions of Theorem 8.6.

For the next two lemmas, if ϕ and f are in $C_c(G)$, let $f \cdot \phi$ denote their pointwise product in $C_c(G)$, i.e., $(\phi \cdot f)(g) = \phi(g)f(g)$. Also, for a subset K of G , we will write $C_K^*(G)$ for the subspace of $C_r^*(G)$ consisting of all elements supported in K .

Lemma 8.19 *Let K be a compact subset of G . Then there exists a constant $M > 0$ such that for all $\phi \in C_K^*(G)$ we have*

$$\|\phi \cdot f\|_{C_r^*(G)} \leq M \sup_{g \in G} |\phi(g)| \|f\|_{C_r^*(G)}.$$

Proof With notation as in Definition 8.3, it suffices to prove that there exists $M > 0$ such that for all $f \in C_c(G)$ and all $x \in G^{(0)}$,

$$\|\pi_x(\phi \cdot f)\| \leq M \sup_{g \in G} |\phi(g)| \|\pi_x(f)\|.$$

As G is étale, for each $g \in G$ there is an open neighbourhood U of G such that both r and s are injective when restricted to U ; as K is compact, there is a collection U_1, \dots, U_M of open subsets of G with this property such that $K \subseteq \bigcup_{i=1}^M U_i$. We may write ϕ as a sum $\phi = \phi_1 + \dots + \phi_M$, where each ϕ_i is supported in some U_i as above, and satisfies $\sup_{g \in G} |\phi_i(g)| \leq \sup_{g \in G} |\phi(g)|$. It thus suffices to prove that if $\psi \in C_c(G)$ is supported in an open set U such that r, s are injective when restricted to U , then

$$\|\pi_x(\psi \cdot f)\| \leq \sup_{g \in G} |\psi(g)| \|\pi_x(f)\|.$$

We now prove this. Indeed, computing for $\xi \in l^2(s^{-1}(x))$

$$\|\pi_x(\psi \cdot f)\xi\|^2 = \sum_{g \in s^{-1}(x)} \left| \sum_{g_1 g_2 = g} \psi(g_1) f(g_1) \xi(g_2) \right|^2.$$

For each $g \in G$, write g_U for the unique element in $r^{-1}(r(g)) \cap U$ (if this exists). Then the sum above becomes

$$\begin{aligned} & \sum_{\{g \in s^{-1}(x) \mid r^{-1}(r(g)) \cap U \neq \emptyset\}} |\psi(g_U) f(g_U) \xi(g_U^{-1}g)|^2 \\ & \leq \sup_{g \in G} |\psi(g)|^2 \sup \left\{ |f(g)|^2 \sum_{g \in s^{-1}(x)} |\xi(g)|^2 \mid g \in \bigcup_{s(h)=x} s^{-1}(r(h)) \right\} \\ & = \sup_{g \in G} |\psi(g)|^2 \sup \left\{ |f(g)|^2 \|\xi\|^2 \mid g \in \bigcup_{s(h)=x} s^{-1}(r(h)) \right\}. \end{aligned}$$

As the expression $\sup\{|f(g)|^2 \mid g \in \cup_{s(h)=x} s^{-1}(r(h))\}$ is easily seen to be a lower bound for $\|\pi_x(f)\|$, we are done. \square

Lemma 8.20 *For any $\epsilon > 0$ and compact subset K of G , there exists $\delta > 0$ such that if $\phi \in C_c(G^{(0)})$ satisfies*

$$\sup_{g \in K} |\phi(r(g)) - \phi(s(g))| < \delta,$$

then the commutator $[f, \phi]$ has norm at most $\epsilon \|\phi\| \|f\|$ for any $f \in C_K(G)$.

Proof With assumptions as above, the commutator is the element of $C_c(G)$ given by

$$[f, \phi](g) = f(g)\phi(s(g)) - \phi(r(g))f(g)$$

in other words, it is the function $(\phi \circ s - \phi \circ r) \cdot f$, where ‘ \cdot ’ denotes pointwise multiplication. Fix now a compact set K' containing an open neighbourhood of K , and let ψ be any function that agrees with $\phi \circ s - \phi \circ r$ on K , vanishes outside K' and is bounded above by $\sup_{g \in K} |(\phi \circ s - \phi \circ r)(g)|$. Lemma 8.19 then implies that there exists M such that for any $f \in C_K^*(G)$

$$\begin{aligned} \|[f, \phi]\|_{C_r^*(G)} &= \|\psi \cdot f\|_{C_r^*(G)} \leq M \sup_{g \in G} |\psi(g)| \|f\|_{C_r^*(G)} \\ &\leq M \sup_{g \in K} |(\phi \circ s - \phi \circ r)(g)| \|f\|_{C_r^*(G)}; \end{aligned}$$

taking $\delta = \epsilon/M$ thus works. \square

Proof of Theorem 8.6 Let a finite subset \mathcal{F} of $C_r^*(G)$ and $\epsilon > 0$ be given. As $C_c(G)$ is dense in $C_r^*(G)$ we may assume that \mathcal{F} is a subset of $C_c(G)$; let K be a compact subset of G such that $K = K^{-1}$ and such that each element $f \in \mathcal{F}$ is supported in K . Let $\widetilde{C_r^*(G)}$ denote the unitization of $C_r^*(G)$, and let $\widetilde{C_K^*(G)}$ denote the closed subspace of $\widetilde{C_r^*(G)}$ spanned by $\widetilde{C_K^*(G)}$ and the identity of $\widetilde{C_r^*(G)}$; as $K = K^{-1}$, this is an operator subsystem of $\widetilde{C_r^*(G)}$. Using Proposition 7.1 (see also Remark 8.5 part (iii)), there exists a collection $\{\phi_i : G^{(0)} \rightarrow [0, 1] \mid i \in \{0, \dots, d\}\}$ of continuous compactly supported functions on $G^{(0)}$ with the following properties:

- (i) for each $x \in r(K) \cup s(K)$ we have $\sum_{i=0}^d \phi_i^2(x) = 1$;
- (ii) for each i there is a small open subgroupoid H_i of G such that for any $f \in \widetilde{C_K^*(G)}$, the element $\phi_i f \phi_i$ is contained in the sub- C^* -algebra $C_r^*(H_i)$ of $C_r^*(G)$;
- (iii) for each i and $g \in K$,

$$|\phi_i(r(g)) - \phi_i(s(g))| < \frac{\epsilon}{2(d+1)}.$$

From the third point above and Lemma 8.20 we have

$$\|[f, \phi_i]\| < \frac{\epsilon}{2(d+1)} \|f\| \tag{8.1}$$

for any $f \in C_K^*(G)$, so in particular for all $f \in \mathcal{F}$.

Now, for each $i \in \{0, \dots, d\}$, the formula

$$\Phi_i : \widetilde{C_K^*(G)} \rightarrow C_r^*(H_i), \quad f \mapsto \phi_i f \phi_i$$

defines a contractive completely positive map. Let

$$\Psi_i : C_r^*(H_i) \rightarrow C_r^*(G)$$

be the canonical inclusion $*$ -homomorphism. For each $i \in \{0, \dots, d\}$ we now have a triangle of maps

$$\begin{array}{ccc}
 \widetilde{C_K^*(G)} & & C_r^*(G) \\
 \searrow \Phi_i & & \nearrow \Psi_i \\
 & C_r^*(H_i) &
 \end{array}
 \tag{8.2}$$

where each Φ_i is contractive and completely positive, and each Ψ_i is a $*$ -homomorphism, so in particular order zero. Moreover, for every $f \in \mathcal{F}$

$$\sum_{i=0}^d \Psi_i(\Phi_i(f)) = \sum_{i=0}^d \phi_i f \phi_i = \sum_{i=0}^d \phi_i^2 f + \sum_{i=0}^d \phi_i [f, \phi_i] = f + \sum_{i=0}^d \phi_i [f, \phi_i],$$

whence line (8.1) implies that

$$\left\| \sum_{i=0}^d \Psi_i(\Phi_i(f)) - f \right\| < \epsilon/2$$

for all $f \in \mathcal{F}$.

On the other hand, we have already shown that each $C_r^*(H_i)$ has nuclear dimension at most the covering dimension of $H_i^{(0)}$ in Proposition 8.7, and this is at most N as covering dimension does not increase under taking open subsets (this follows for example from [42, Proposition 2.5]). Combining the triangle in line (8.2) with approximations to each $C_r^*(H_i)$ arising from the definition of nuclear dimension gives triangles

$$\begin{array}{ccc} \widetilde{C_K^*(G)} & & C_r^*(G), \quad i \in \{0, \dots, d\}, j \in \{0, \dots, N\} \\ & \searrow \Phi_{ij} & \nearrow \Psi_{ij} \\ & F_{ij} & \end{array}$$

where each Φ_{ij} is contractive and completely positive, each Ψ_{ij} is contractive and order zero, each F_{ij} is a finite dimensional C^* -algebra, and

$$\left\| \sum_{i=0}^d \sum_{j=0}^N \Psi_{ij}(\Phi_{ij}(f)) - f \right\| < \epsilon$$

for all $f \in \mathcal{F}$. Finally, the finite dimensional version of Arveson’s extension theorem (see for example [6, Section 1.6]) implies that each Φ_{ij} extends to a contractive completely positive map

$$\Phi_{ij} : \widetilde{C_r^*(G)} \rightarrow F_{ij},$$

and restricting each of these maps to $C_r^*(G)$ gives approximating triangles

$$\begin{array}{ccc} C_r^*(G) & & C_r^*(G), \quad i \in \{0, \dots, d\}, j \in \{0, \dots, N\} \\ & \searrow \Phi_{ij} & \nearrow \Psi_{ij} \\ & F_{ij} & \end{array}$$

as required by the definition of nuclear dimension $(d + 1)(N + 1) - 1$. □

8.3 Consequences

Finally, we spell out a few consequences. Many of these are known results, but we think that combining the proofs under one common dynamical framework has some interest.

The first result is much easier to check directly! See Winter [42, Remark 2.2(iii)] and Kirchberg–Winter [21, Example 4.1]; nonetheless, it seemed interesting that it fits directly into our framework.

Corollary 8.21 *Separable AF C^* -algebras have nuclear dimension zero.*

Proof Renault [28, Proposition III.1.15] has shown that any (separable) AF C^* -algebra arises as the C^* -algebra of a locally finite groupoid (in the sense of Example 5.3) with unit space of covering dimension zero. The result thus follows directly from Example 5.3 and Theorem 8.6. \square

The next corollary is due to Winter–Zacharias: see [42, Section 8].

Corollary 8.22 *Let X be a bounded geometry coarse space of asymptotic dimension d . Then the uniform Roe algebra $C_u^*(X)$ has nuclear dimension at most d .*

Proof The uniform Roe algebra of X is naturally isomorphic to the reduced groupoid C^* -algebra of the coarse groupoid $G(X)$: see [31, Proposition 10.29] for a proof. The result now follows on combining Theorem 6.4, Theorem 8.6 (see also Remark 8.5, part (ii)), and the fact that the unit space of $G(X)$ is βX , which has covering dimension zero. \square

The next corollary is due to Toms–Winter [39, Section 3].

Corollary 8.23 *Let $\mathbb{Z} \curvearrowright X$ be a minimal action of \mathbb{Z} on an infinite second countable compact space X of covering dimension N . Then the nuclear dimension of $C(X) \rtimes_r \mathbb{Z}$ is at most $2N + 1$.*

Proof Combine Theorem 3.1, Lemma 5.4, Theorem 8.6, and the well-known (and easily checked) fact that the reduced groupoid C^* -algebra of a transformation groupoid $\Gamma \ltimes Y$ identifies naturally with $C(Y) \rtimes_r \Gamma$. \square

It is worth noting explicitly that the proofs of Corollary 8.22 and Corollary 8.23 in the original references given above are quite similar to our proof of Theorem 8.6: all involve constructing ‘almost invariant’ partitions of unity, and using these to ‘cut down’ to subhomogeneous C^* -algebras whose nuclear dimension can be directly estimated. This only became apparent to us ‘after the fact’, but we hope it helps to clarify how this style of argument that estimates nuclear dimension from geometric and / or dynamic assumptions is built up.

The following result seems to be new as stated. Note however that if $\Gamma \curvearrowright X$ is a minimal, free, amenable action of a *non-amenable* group on the Cantor set, then $C(X) \rtimes_r \Gamma$ is a Kirchberg algebra in the UCT class; the main result of [34] (see also [5]) thus implies that the nuclear dimension of $C(X) \rtimes_r \Gamma$ is one. As any exact (non-amenable) group admits such an action (see [33, Theorem 6.11]), and as finite asymptotic dimension implies exactness (see [18, Lemma 4.3] and [25]) but not conversely, the corollary below is not optimal, neither with respect to the class of groups covered, nor with respect to the estimate on nuclear dimension.

Corollary 8.24 *Let Γ be a countable discrete group with finite asymptotic dimension. Then Γ admits a minimal, free action on the Cantor set X such that $C(X) \rtimes_r \Gamma$ has nuclear dimension bounded above by the asymptotic dimension of Γ .*

Proof Combine Lemma 5.4, Theorems 6.6, 8.6, and again that the reduced groupoid C^* -algebra of a transformation groupoid $\Gamma \ltimes Y$ identifies naturally with $C(Y) \rtimes_r \Gamma$. \square

We should note that there are many other results in the literature that give estimates on nuclear dimension or other C^* -algebraic regularity properties based on conditions on actions: see for example [11, 19, 20, 36, 37]. Many of these results go further than ours in at least some ways: for example, [36] proves fairly general finite dimensionality results for \mathbb{Z}^n -actions, [20] treats non-free \mathbb{Z} -actions, [11] treats some \mathbb{Z} -actions on non-finite dimensional spaces, and several works deal with some actions on non-commutative C^* -algebras. It would be interesting to clarify the relationships holding between the various conditions involved in these results and ours.

The final result is not strictly a ‘corollary’ as such, but follows from exactly the same method of proof; it does not require any separability assumptions as these were only used in the above in the appeal to Theorem 8.13, and to avoid complications from general topology with respect to dimension theory. We guess it is possible to prove it more directly, and without the assumption of freeness, but did not pursue this.

Corollary 8.25 *Let G be a free étale groupoid with finite dynamic asymptotic dimension. Then G is amenable.*

Proof The method of proof of Theorem 8.6 implies in particular that $C_r^*(G)$ is nuclear (whether or not the unit space of G is finite dimensional). Hence G is amenable by [6, Theorem 5.6.18]. \square

Acknowledgements This paper has been some time in gestation, and has benefited from conversations with several people. In particular, we would like to thank Arthur Bartels, Siegfried Echterhoff, David Kerr, Ian Putnam, Daniel Ramras, Wilhelm Winter, and Jianchao Wu for useful comments, penetrating questions, and/or patient explanations. E. Guentner, R. Willett would like to thank Texas A&M University and the Shanghai Center for Mathematical Sciences for their hospitality during some of the work on this paper. E. Guentner was partially supported by a grant from the Simons Foundation (#245398). R. Willett was partially supported by NSF Grant DMS-1401126. G. Yu was partially supported by NSF Grant DMS-1362772 and NSFC Grant NSFC11420101001.

References

1. Bartels, A: On proofs of the Farrell-Jones conjecture. [arXiv:1210.1044v1](https://arxiv.org/abs/1210.1044v1) (2012)
2. Bartels, A., Lück, W.: The Borel conjecture for hyperbolic and CAT(0)-groups. *Ann. Math.* **175**(2), 631–689 (2012)
3. Bartels, A., Lück, W., Reich, H.: Equivariant covers for hyperbolic groups. *Geom. Topol.* **12**, 1799–1882 (2008)
4. Bell, G., Dranishnikov, A.: Asymptotic dimension. *Topology Appl.* **155**(12), 1265–1296 (2008)
5. Bosa, J., Brown, N., Sato, Y., Tikuisis, A., White, S., Winter, W.: Covering dimension of C^* -algebras and 2-coloured classification. [arXiv:1506.03974](https://arxiv.org/abs/1506.03974) (2015)
6. Brown, N., Ozawa, N.: C^* -Algebras and Finite-Dimensional Approximations. Graduate Studies in Mathematics, vol. 88. American Mathematical Society, Providence (2008)
7. Dadarlat, M., Guentner, E.: Uniform embeddability of relatively hyperbolic groups. *J. Reine Angew. Math.* **612**, 1–15 (2007)
8. Dixmier, J.: C^* -Algebras. North Holland Publishing Company, Amsterdam (1977)
9. Elliott, G.: The classification problem for amenable C^* -algebras. *Proc. Int. Congr. Math.* **1**(2), 922–932 (1995)
10. Elliott, G., Gong, G., Lin, H., Niu, Z.: On the classification of simple amenable C^* -algebras with finite decomposition rank, II. [arXiv:1507.03437](https://arxiv.org/abs/1507.03437) (2015)
11. Elliott, G., Niu, Z.: The C^* -algebra of a minimal homeomorphism of zero mean dimension. [arXiv:1406.2383v2](https://arxiv.org/abs/1406.2383v2) (2014)
12. Farrell, T., Jones, L.: K -theory and dynamics I. *Ann. Math.* **124**(2), 531–569 (1986)

13. Felix, R.: Renault's equivalence theorem for C^* -algebras of étale groupoids. Master's thesis, University of Hawai'i at Mānoa (2014)
14. Gromov, M.: Asymptotic invariants of infinite groups. In: Niblo, G., Roller, M. (eds.) *Geometric Group Theory*, vol. 2. London Mathematical Society, London (1993)
15. Guentner, E., Tessera, R., Yu, G.: A notion of geometric complexity and its application to topological rigidity. *Invent. Math.* **189**(2), 315–357 (2012)
16. Guentner, E., Willett, R., Yu, G.: Dynamic asymptotic dimension and controlled operator K-theory. Working draft (2015)
17. Higson, N.: Bivariant K -theory and the Novikov conjecture. *Geom. Funct. Anal.* **10**, 563–581 (2000)
18. Higson, N., Roe, J.: Amenable group actions and the Novikov conjecture. *J. Reine Angew. Math.* **519**, 143–153 (2000)
19. Hirshberg, I., Winter, W., Zacharias, J.: Rokhlin dimension and C^* -dynamics. *Comm. Math. Phys.* **335**, 637–670 (2015)
20. Hirshberg, I., Wu, J.: The nuclear dimension of C^* -algebras associated to homeomorphisms. [arXiv:1509.01508](https://arxiv.org/abs/1509.01508). With an appendix by Gabor Szabó (2015)
21. Kirchberg, E., Winter, W.: Covering dimension and quasidiagonality. *Intern. J. Math.* **15**, 63–85 (2004)
22. Massey, W.: *Algebraic Topology: An Introduction*. Springer, Berlin (1977)
23. Muhly, P., Renault, J., Williams, D.: Equivalence and isomorphism for groupoid C^* -algebras. *J. Oper. Theory* **17**, 3–22 (1987)
24. Nowak, P., Yu, G.: Large scale geometry. In: EMS Textbooks in Mathematics, European Mathematical Society (EMS), Zürich (2012)
25. Ozawa, N.: Amenable actions and exactness for discrete groups. *C. R. Acad. Sci. Paris Sér. I Math.* **330**, 691–695 (2000)
26. Pears, A.: *Dimension Theory of General Spaces*. Cambridge University Press, Cambridge (1975)
27. Putnam, I.: The C^* -algebras associated with minimal homeomorphisms of the Cantor set. *Pac. J. Math.* **136**(2), 329–353 (1989)
28. Renault, J.: *A Groupoid Approach to C^* -Algebras*. Lecture Notes in Mathematics, vol. 793. Springer, Berlin (1980)
29. Renault, J.: The ideal structure of groupoid crossed product C^* -algebras. *J. Oper. Theory* **25**, 3–36 (1991)
30. Renault, J.: C^* -algebras and dynamical systems. *Publicações Matemáticas do IMPA, 27º Colóquio Brasileiro de Matemática*. Instituto Nacional de Matemática Pura e Aplicada (2009)
31. Roe, J.: *Lectures on Coarse Geometry*. University Lecture Series, vol. 31. American Mathematical Society, Providence (2003)
32. Roe, J.: Hyperbolic groups have finite asymptotic dimension. *Proc. Am. Math. Soc.* **133**(9), 2489–2490 (2005)
33. Rørdam, M., Sierakowski, A.: Purely infinite C^* -algebras arising from crossed products. *Ergod. Theory Dynam. Syst.* **32**, 273–293 (2012)
34. Ruiz, E., Sims, A., Sørensen, A.: UCT-Kirchberg algebras have nuclear dimension one. *Adv. Math.* **279**, 1–28 (2015)
35. Skandalis, G., Tu, J.-L., Yu, G.: The coarse Baum–Connes conjecture and groupoids. *Topology* **41**, 807–834 (2002)
36. Szabó, G.: The Rokhlin dimension of topological \mathbb{Z}^m -actions. *Proc. Lond. Math. Soc.* **110**(3), 673–694 (2015)
37. G. Szabó, Zacharias, J., Wu, J.: Rokhlin dimension for actions of residually finite groups. [arXiv:1408.6096v2](https://arxiv.org/abs/1408.6096v2) (2014)
38. Tikuisis, A., White, S., Winter, W.: Quasidiagonality of nuclear C^* -algebras. [arXiv:1509.08318](https://arxiv.org/abs/1509.08318) (2015)
39. Toms, A., Winter, W.: Minimal dynamics and K -theoretic rigidity: Elliott's conjecture. *Geom. Funct. Anal.* **23**, 467–481 (2013)
40. Vick, J.: *Homology Theory: An Introduction to Algebraic Topology*. Academic Press, Cambridge (1973)
41. Winter, W.: Decomposition rank of subhomogeneous C^* -algebras. *Proc. Lond. Math. Soc.* **89**, 427–456 (2004)
42. Winter, W., Zacharias, J.: The nuclear dimension of C^* -algebras. *Adv. Math.* **224**(2), 461–498 (2010)
43. Wright, N.: Finite asymptotic dimension for CAT(0) cube complexes. *Geom. Topol.* **1**, 527–554 (2012)
44. Yu, G.: The Novikov conjecture for groups with finite asymptotic dimension. *Ann. Math.* **147**(2), 325–355 (1998)