



Dual braid monoids, Mikado braids and positivity in Hecke algebras

François Digne¹ · Thomas Gobet²

Received: 23 September 2015 / Accepted: 9 May 2016 / Published online: 4 June 2016
© Springer-Verlag Berlin Heidelberg 2016

Abstract We study the rational permutation braids, that is the elements of an Artin-Tits group of spherical type which can be written $x^{-1}y$ where x and y are prefixes of the Garside element of the braid monoid. We give a geometric characterization of these braids in type A_n and B_n and then show that in spherical types different from D_n the simple elements of the dual braid monoid (for arbitrary choice of Coxeter element) embedded in the braid group are rational permutation braids (we conjecture this to hold also in type D_n). This property implies positivity properties of the polynomials arising in the linear expansion of their images in the Iwahori-Hecke algebra when expressed in the Kazhdan-Lusztig basis. In type A_n , it implies positivity properties of their images in the Temperley-Lieb algebra when expressed in the diagram basis.

Contents

1	Introduction	216
2	Classical braid monoid	218
3	Dual braid monoids	218
3.1	Coxeter elements	218
3.2	Dual braid monoids	220
3.3	Hurwitz action on reduced decompositions	220
3.4	Groups of fractions of dual braid monoids are Artin-Tits groups	221
3.5	A formula for dual atoms	222
4	Rational permutation braids	223
4.1	Square-free braids	223
4.2	Rational permutation braids	223
5	Mikado braids of type A_n	224

✉ Thomas Gobet
gobet@mathematik.uni-kl.de

François Digne
digne@u-picardie.fr

¹ LAMFA, Université de Picardie Jules Verne, 33 Rue Saint-Leu, 80039 Amiens Cedex 1, France

² TU Kaiserslautern, Fachbereich Mathematik, Postfach 3049, 67653 Kaiserslautern, Germany

5.1 Square-free braids	224
5.2 Mikado braids	225
5.3 Simple dual braids are Mikado braids	227
5.3.1 Noncrossing partitions	227
5.3.2 Dual braids and graphical representation	228
5.3.3 Removing the strands of simple dual braids	230
5.3.4 Additional properties in case the Coxeter element is linear	230
6 Mikado braids of type B_n	231
6.1 Coxeter and Artin-Tits groups of type B_n	231
6.2 Mikado braids of type B_n	231
6.3 Simple dual braids are Mikado braids	232
6.3.1 Graphical representation of simple elements	232
6.3.2 Removing pairs of strands of simple dual braids	232
7 Dihedral and exceptional types	233
8 Positivity properties	233
8.1 Iwahori-Hecke algebra of a Coxeter system	234
8.2 Positivity properties of simple dual braids	234
8.3 Temperley-Lieb algebra and monomial basis	235
8.4 Projection of the canonical basis	236
8.5 Zinno basis	236
8.6 Positivity consequences	237
References	237

1 Introduction

This paper is motivated by positivity properties arising when expanding simple dual braids in the canonical basis of the Iwahori-Hecke algebra.

More precisely, a *dual braid monoid* associated to a finite Coxeter group and a choice of Coxeter element is a Garside monoid [16] whose group of fractions is isomorphic to the corresponding Artin-Tits group. The dual braid monoids were introduced by Bessis [4], extending definitions by Birman-Ko-Lee [6] and Bessis-Michel and the first author [5]. As Garside monoids, they possess a finite set of *simple elements* which generates the whole monoid and forms a lattice under the left-divisibility order and they embed into their group of fractions. Similar constructions have been made for some infinite Coxeter groups and for complex reflection groups, but in this paper, unless explicitly stated, “dual braid monoid” will mean a dual braid monoid associated with a finite Coxeter group.

In the framework of Artin-Tits groups, the standard example of a Garside monoid is the braid or Artin-Tits monoid, also known as positive or classical braid monoid. Its set of simple elements is the canonical positive lift of the Coxeter group in the Artin-Tits group. The embedding property in this specific case was shown by Brieskorn-Saito [8] and by Deligne [17]. In the dual approach, the set of simple generators of the Coxeter group is replaced in the spherical case by the whole set of reflections, that is, by the conjugates of the simple generators. A key tool in this approach is an action of the braid group on the set of reduced factorizations of a Coxeter element as a product of reflections, called the Hurwitz action. This action is known to be transitive [3, 4], providing an analogue in the dual approach of the Tits-Matsumoto property.

The dual braid monoid is then an analogue to the classical braid monoid, but having as set of generators a copy of the set of reflections of the Coxeter group. In the classical setting, the lattice of simple elements is isomorphic to the lattice obtained by endowing the Coxeter group with the left weak order defined by the classical length function. By mimicking this approach in the dual setting, that is, by replacing the classical length func-

tion by the length function with respect to the whole set of reflections, one obtains an order called *reflection* or *absolute order* on the Coxeter group. The absolute order does not in general endow the Coxeter group with a lattice structure, but by restricting such an order to the order ideal of a Coxeter element, one obtains a lattice [4, 11]. The dual braid monoid is built using this property. Its lattice of simple elements is isomorphic to the order ideal of a given Coxeter element ordered by the restriction of the absolute order [4].

There is a morphism from the Artin-Tits group to the group of invertible elements of the Iwahori-Hecke algebra attached to the corresponding Coxeter group. The images of the simple elements of the classical braid monoid through this morphism yield a basis of the Iwahori-Hecke algebra, the so-called *standard basis*. It is therefore natural to ask about properties of the images of the simple elements of the dual Garside structure of an Artin-Tits group in the associated Iwahori-Hecke algebra.

In type A_n , it is known that their images in the Temperley-Lieb algebra yield a basis [32, 39, 40] such that the base change to the diagram basis is triangular [25, 40].

One of the main points of this paper consists of understanding how one can express the simple dual braids in terms of the classical braid group generators. Indeed, the isomorphism between the group of fractions of a dual braid monoid and the Artin-Tits group is proven by a case-by-case analysis, and there is no known general formula for the simple dual braids in the braid group, even for the atoms. We recall the construction of the classical and dual braid monoids in Sects. 2 and 3 where we also prove some new results on dual Coxeter systems. In particular we give a formula for the reflections in terms of the simple reflections (Proposition 3.4) which lifts well to the Artin-Tits group, allowing us to get a uniform formula for the dual atoms in terms of the classical generators (Proposition 3.13) and this with a case-free proof.

We then introduce in Sect. 4 a finite set of braids, which we call *rational permutation braids*. These braids already appear in a paper by Dehornoy [14] in type A_n and in unpublished work of Dyer [19] for Artin-Tits groups attached to arbitrary (not necessarily finite) Coxeter groups. They may be defined in spherical types as the braids of the form $\mathbf{x}\mathbf{y}^{-1}$ where \mathbf{x} and \mathbf{y} are simple braids for the classical Garside structure. The images of such braids in the Iwahori-Hecke algebra turn out to have positive Kazhdan-Lusztig expansions; this is shown by Dyer-Lehrer [20] for finite Weyl groups using perverse sheaves and by Dyer [18] for arbitrary finite Coxeter groups using Soergel bimodules. These braids are closely related to the so-called *mixed braid relations* introduced by Dyer [19].

We give uniform Garside-theoretic characterizations of the rational permutation braids in the spherical case in Proposition 4.3, and then turn to a case-by-case analysis in types A_n (Sect. 5) and B_n (Sect. 6) to characterize them in terms of geometrical braids (Propositions 5.7 and 6.2). In these cases, they turn out to be what we call the *Mikado braids*, that is, the braids where one can inductively remove a strand which is above all the other strands. With this geometric characterization, we can show that all the simple dual braids, for any choice of Coxeter element, are rational permutation braids (Theorems 5.12 and 6.6)—the argument given here is therefore topological. We also prove the same result for dihedral groups and, by computer, for exceptional types (Theorem 7.1); we conjecture the result to also hold in type D_n (Conjecture 8.7).

It follows that the simple dual braids have a positive Kazhdan-Lusztig expansion in all types except possibly type D_n (Theorem 8.6). Positivity results in the Temperley-Lieb algebra (of type A_n) can also be derived (Theorem 8.16).

2 Classical braid monoid

Starting with a finite Coxeter System (W, S) , one can define the associated Artin-Tits monoid or *classical braid monoid* $B^+(W)$ and for each choice of a Coxeter element c an associated *dual braid monoid* $B_c^*(W)$ [4].

The classical braid monoid is defined by the following presentation:

$$B^+(W) = \langle S \mid \underbrace{s_1 s_2 \dots}_{m_{s_1, s_2}} = \underbrace{s_2 s_1 \dots}_{m_{s_1, s_2}} \text{ for } s_1, s_2 \in S \rangle^+$$

where S is a set in bijection $s \mapsto s$ with S and m_{s_1, s_2} is the order of $s_1 s_2$ in W .

The monoid $B^+(W)$ is a *Garside monoid* (see for instance [15, I, 2.1] for the definition), hence it satisfies the Ore conditions and embeds in its group of fractions, the braid group $B(W)$ of W . This group has the same presentation as $B^+(W)$ but as a group:

$$B(W) = \langle S \mid \underbrace{s_1 s_2 \dots}_{m_{s_1, s_2}} = \underbrace{s_2 s_1 \dots}_{m_{s_1, s_2}} \text{ for } s_1, s_2 \in S \rangle.$$

The group W is a quotient of $B(W)$ by the normal subgroup generated by the squares of the elements of S . We denote by $p : B(W) \rightarrow W$ this surjective morphism. It has a set-wise section obtained by lifting S -shortest expressions. We denote by \mathbf{W} the image of this section. This set \mathbf{W} is the set of simples of the Garside monoid $B^+(W)$. The Garside element of $B^+(W)$ is the lift $\Delta = w_0$ of the longest element w_0 of W .

By definition of Garside monoids the left- (*resp.* right-) divisibility gives a lattice structure to $B^+(W)$ which restricts to a lattice structure on \mathbf{W} . Moreover by construction of \mathbf{W} , the map p provides an isomorphism of lattices from \mathbf{W} to W endowed with the Coxeter theoretic left- (*resp.* right-) weak order.

3 Dual braid monoids

3.1 Coxeter elements

The dual braid monoids are defined using Coxeter elements. Before defining these monoids we prove some properties of the Coxeter elements that we will need.

Definition 3.1 Let (W, S) be a Coxeter system. An element $c \in W$ is a *standard Coxeter element of (W, S)* if it is a product of the elements of S in some order. An element $c \in W$ is a *Coxeter element* if there exists $S' \subset T = \bigcup_{w \in W} w S w^{-1}$ such that (W, S') is a Coxeter system and c is the product of the elements of S' in some order, that is, c is a standard Coxeter element of (W, S') . The set T is the set of *reflections* of (W, S) .

Remark 3.2 If S' is as above then by [9, Lemma 3.7] one has $T = \bigcup_{w \in W} w S' w^{-1}$. Moreover if W is finite and S' is as above, then by [9, Theorem 3.10], the Coxeter systems (W, S) and (W, S') are isomorphic. In particular, the type of (W, S) depends only on T . If (W, S) is infinite irreducible with no infinite entry in its Coxeter matrix, a stronger property holds: any S' such that (W, S') is a Coxeter system is conjugate to S [24, Theorem 1]. The example of the dihedral group $I_2(5)$ with $S = \{s, t\}$ and $S' = \{s, ststs\}$ shows that when W is finite S and S' need not be conjugate.

Definition 3.3 Let (W, S) be a Coxeter system.

- (1) We write ℓ_S , resp. ℓ_T , for the length function on W with respect to S , resp. T .
- (2) We say that $x \in W$ divides $y \in W$, written $x \preceq_T y$, if $\ell_T(x^{-1}y) + \ell_T(x) = \ell_T(y)$. We write $\text{Div}(y) := \{x \in W \mid x \preceq_T y\}$ and call the elements of $\text{Div}(y)$ the *divisors* of y .

Note that the definitions of ℓ_S and ℓ_T use the fact that each one of the sets S and T positively generates W . Note also that, since T is invariant under conjugation, the relation $\ell_T(x^{-1}y) + \ell_T(x) = \ell_T(y)$ is equivalent to $\ell_T(yx^{-1}) + \ell_T(x) = \ell_T(y)$: there is no need to distinguish between left- and right-divisibility.

The following proposition shows how to recover all reflections from a Coxeter element when W is finite. Recall that if c is a Coxeter element in a Coxeter system of rank n , one has $\ell_T(c) = n$ (see e.g., [3, Lemma 1.2]).

Proposition 3.4 *Let (W, S) be a finite Coxeter system with $S = \{s_1, \dots, s_n\}$. Let $c = s_1s_2 \cdots s_n$. Then the set of reflections of W is given by*

$$\{c^k s_1 s_2 \dots s_i s_{i-1} \dots s_1 c^{-k} \mid k \in \mathbb{N}, 1 \leq i \leq n\}.$$

Proof Let T_c be the set of the statement and T be the set of reflections of (W, S) . We have $T_c \subset T$. If c is such that $\ell_S(c^i) = i\ell_S(c)$ for any $i \leq |T|/h$, where $h = 2|T|/n$ is the Coxeter number (the order of c), then T_c is equal to T (this is a consequence of e.g., the proof of [12, 3.9]; see also [7, Chapter V, §6 exercise 2] in the case where c is bipartite), whence the result holds in that case. We now remark that if $c' = s_{i+1} \dots s_n s_1 \dots s_i$, with $1 \leq i \leq n - 1$ then $T_{c'}$ contains $(s_i \dots s_2 s_1) T_c (s_i \dots s_2 s_1)^{-1}$ so has cardinality at least $|T_c|$, whence the result holds: indeed, since W is finite, all standard Coxeter elements of (W, S) are cyclically conjugate (that is by a sequence of conjugations like the one deriving c' from c above): see e.g., [23, Theorem 3.1.4]. □

We take from [3, Section 1] the second item of the following definition.

- Definition 3.5** (1) A *parabolic subgroup* W' of a Coxeter system (W, S) is a subgroup generated by a conjugate S' of a subset of S . We shall say that (W', S') is a *parabolic subsystem* of (W, S) .
- (2) Let (W, S) be a Coxeter system with set of reflections T . Let $x \in W$. Then x is a *parabolic Coxeter element* in W if there exists a subset $S' = \{s'_1, \dots, s'_n\} \subset T$ such that $x = s'_1 \cdots s'_m$ for some $m \leq n$ and (W, S') is a Coxeter system.

Corollary 3.6 *Let (W, S) be a finite Coxeter system and let $x \in W$. The following are equivalent:*

- (1) *There exists a Coxeter element $c \in W$ such that $x \preceq_T c$.*
- (2) *The element x is a parabolic Coxeter element.*

Proof Assume that $x \preceq_T c$ for some Coxeter element c . Let $S' = \{s_1, \dots, s_n\}$ be a simple Coxeter system of (W, S) such that $c = s_1 \dots s_n$ and let T be the set of reflections of W . We argue by reverse induction on $\ell_T(x)$. First assume that $\ell_T(x) = n - 1$, so that there exists a decomposition of c into a product of n reflections $c = t_1 \dots t_n$ such that $x = t_2 \dots t_n$. By the above proposition we have $t_1 = c^k s_1 \dots s_i s_{i+1}$ for some $k \in \mathbb{N}$ and some $1 \leq i \leq n - 1$, where for two elements a and b in a group, we write ${}^a b$ for aba^{-1} . Let $y = s_i \dots s_1 c^{-k}$; we have ${}^y c = s_{i+1} \dots s_n s_1 \dots s_i$ and ${}^y t_1 = s_{i+1}$. Then ${}^y x = s_{i+2} \dots s_n s_1 \dots s_i$ is a standard Coxeter element in the standard parabolic subgroup W_1 generated by $S_1 = \{s_1, \dots, s_i, s_{i+2}, \dots, s_n\}$ so that x is a standard Coxeter element in the parabolic subsystem ${}^{y^{-1}}(W_1, S_1)$. Since a

conjugate of a simple system is again a simple system, we deduce that x is a parabolic Coxeter element. Now the reflections of a parabolic subgroup are precisely the reflections of W which lie in this subgroup, so that the induction can go on: a divisor of x in W is in the parabolic subsystem of which x is a standard Coxeter element. The converse is immediate. \square

Remark 3.7 In his case, that is for finite Coxeter groups, Bessis gives [4, Section 1.4] an alternative definition of parabolic Coxeter elements and shows [4, Lemma 1.4.3] that an element x of W is a parabolic Coxeter element in his sense if and only if there exists a bipartite Coxeter element c such that $x \preccurlyeq_T c$. Since in finite Coxeter groups any Coxeter element is conjugate to a bipartite one, one can drop the word bipartite from this definition. This last fact and the above lemma show that Bessis approach and the one of [3], that is of Definition 3.5 (2), actually lead to equivalent definitions for finite Coxeter groups; this equivalence is not obvious if we compare the definitions from Bessis and [3].

3.2 Dual braid monoids

The dual braid monoids are defined as follows: let (W, S) be a Coxeter system and let T denote the set of reflections of (W, S) . Let c be a standard Coxeter element. The dual braid monoid B_c^* associated to c is defined by generators and relations as follows. Let D_c be a set in one-to-one correspondence with the set $\text{Div}(c)$ of divisors of c . Then $B_c^*(W)$ is generated by D_c with only relations $\mathbf{x} \mathbf{y} = \mathbf{z}$ for $\mathbf{x}, \mathbf{y}, \mathbf{z}$ in D_c such that $x \mathbf{y} = \mathbf{z} \in \text{Div}(c)$ and $\ell_T(\mathbf{z}) = \ell_T(x) + \ell_T(y)$, where $x, y, z \in \text{Div}(c)$ are the elements corresponding to $\mathbf{x}, \mathbf{y}, \mathbf{z}$ respectively. The canonical bijection $D_c \rightarrow \text{Div}(c)$ extends to a (surjective) morphism of monoids from $B_c^*(W)$ to W . The above definition is valid for any Coxeter system but in the sequel we study only dual braid monoids associated to finite Coxeter groups.

In the spherical case the monoid $B_c^*(W)$ is a Garside monoid with D_c as its set of simples. In this case the monoid $B_c^*(W)$ has a presentation with a generating set smaller than D_c : first it can be shown that any reflection divides c (see [4, Lemma 1.3.3]); if we denote by T_c the lift of T in $D_c \subset B_c^*(W)$ we have

Proposition 3.8 ([4], Theorem 2.1.4) *In the spherical case the dual braid monoid has the following presentation:*

$$B_c^*(W) = \langle T_c \mid \mathbf{t}_1 \mathbf{t}_2 = \mathbf{t}_2 \mathbf{t}_3 \text{ for } \mathbf{t}_1, \mathbf{t}_2, \mathbf{t}_3 \text{ in } T_c \text{ with } t_1 t_2 = t_2 t_3 \preccurlyeq_T c \rangle^+.$$

The relations in the above presentation of $B_c^*(W)$ are called the *dual braid relations*.

Proof We recall the proof. To this end, we need to recall a few facts on the Hurwitz action on sets of T -reduced decompositions of elements.

3.3 Hurwitz action on reduced decompositions

We write \mathcal{B}_n for the Artin braid group on n strands, that is, the Artin-Tits group $B(W)$ where W is of type A_{n-1} .

Definition 3.9 (*Hurwitz action*) Let G be a group and g be in G ; the braid group \mathcal{B}_n acts on the set of n -tuples of elements of G whose product is equal to g as follows: if $\sigma_1, \dots, \sigma_{n-1}$ are the generators of the braid group \mathcal{B}_n , the *Hurwitz action* of σ_i maps a sequence (t_1, \dots, t_n) such that $g = t_1 t_2 \dots t_n$ to the sequence $(t_1, \dots, t_{i-1}, t_i t_{i+1} t_i^{-1}, t_i, t_{i+2}, \dots, t_n)$.

When considering the Hurwitz action, we will always denote the generators of \mathcal{B}_n by $\sigma_1, \dots, \sigma_{n-1}$. Otherwise we write the generators of $B(W)$ for W of type A_{n-1} as s_1, \dots, s_{n-1} . In the case of reduced factorizations of a Coxeter element into reflections we have:

Theorem 3.10 ([3], Theorem 1.3) *Let (W, S) be a (not necessarily finite) Coxeter system with T its set of reflections. If x is a parabolic Coxeter element in W , then the Hurwitz action is transitive on the set of decompositions of x into a product of $\ell_T(x)$ reflections.*

Note that for finite Coxeter groups, the assumption that x is a parabolic Coxeter element can be replaced by $x \preceq_T c$ for some Coxeter element c of (W, S) thanks to Corollary 3.6.

Let c be a standard Coxeter element in a Coxeter system (W, S) of rank n . By definition of B_c^* , any decomposition $c = t_1 \dots t_n$ into a product of n reflections is lifted in $B_c^*(W)$ to $\mathbf{c} = \mathbf{t}_1 \dots \mathbf{t}_n$ where \mathbf{c} is the lift of c in \mathbf{D}_c . Hence if x divides c , writing $x = t_1 \dots t_{\ell_T(x)}$ with $t_i \in T$ and $c = t_1 \dots t_n$, we see that the lift of x is a product of $\ell_T(x)$ elements of \mathbf{T}_c , so that \mathbf{T}_c generates $B_c^*(W)$. Moreover by Theorem 3.10 and Corollary 3.6 one can pass from any decomposition of $\mathbf{x} \in \mathbf{D}_c$ to any other one by dual braid relations. This gives Proposition 3.8. □

3.4 Groups of fractions of dual braid monoids are Artin-Tits groups

The following is proved in [4]:

Proposition 3.11 *The group of fractions of $B_c^*(W)$ is isomorphic to $B(W)$ and the restriction of the projection $p : B(W) \rightarrow W$ is the canonical surjection of the dual braid monoid onto W .*

We explain how this isomorphism is defined and sketch a proof that it is an isomorphism.

Sketch of proof Let us enumerate the elements s_1, \dots, s_n of S in such a way that $c = s_1 \dots s_n$. We have $S \subset T$ and an easy computation using the dual braid relations shows that the lift S' of S in $\mathbf{T}_c \subset B_c^*(W)$ satisfies the braid relations. This gives a morphism of monoids $B^+(W) \rightarrow B_c^*(W)$ mapping s_i to s'_i where s'_i is the lift of s_i in S' . This extends to a group morphism from $B(W)$ to the group of fractions of $B_c^*(W)$.

Since $c = s_1 \dots s_n$, whence $\mathbf{c} = s'_1 \dots s'_n$, Theorem 3.10 implies that all elements of \mathbf{T}_c can be obtained from S' using dual braid relations, so that S' generates the group of fractions of $B_c^*(W)$, hence the above morphism from $B(W)$ to this group is surjective.

To show that it is injective one first considers the particular case of a Coxeter element c which is bipartite, that is, such that either for $m = \lfloor n/2 \rfloor$ or for $m = \lfloor (n+1)/2 \rfloor$ the elements s_1, s_2, \dots, s_m pairwise commute and so do s_{m+1}, \dots, s_n . Bessis [4] proves that in this case the set $\cup_{k \in \mathbb{Z}} \mathbf{c}^k \mathbf{S} \mathbf{c}^{-k}$ is a lift in $B(W)$ of T and that its elements satisfy the dual braid relations (this last fact by a case by case analysis, see [4, Fact 2.2.4]). Hence we get a morphism in the other direction such that the composition from $B(W)$ to itself is the identity (since it is the identity on S).

To get the result for an arbitrary standard Coxeter element, one can use the fact that the lifts in $B^+(W)$ of all Coxeter elements of a given Coxeter system are conjugate in $B(W)$ (see [7, Chapter V, § 6 Lemme 1]) and that the Hurwitz action commutes with conjugation. □

We will now identify $B_c^*(W)$ with its image in $B(W)$ and S' with S . We then see \mathbf{T}_c as a subset of $B(W)$. Using this identification, we note that, by the above proof, the lift of the Coxeter element $c = s_1 \dots s_n$ in $B_c^*(W)$ is the same as its lift to the classical braid monoid, that is $\mathbf{c} = s_1 \dots s_n \in \mathbf{W}$. Theorem 3.10 implies:

Corollary 3.12 *The set of decompositions of c into a product of $\text{rank}(W)$ elements of T_c in $B(W)$ is a single orbit under the Hurwitz action and the restriction of $\pi : B(W) \rightarrow W$ to T_c induces a bijection from this Hurwitz orbit to the set of decompositions of c into $\text{rank}(W)$ reflections.*

We recall the proof for sake of completeness.

Proof Let $\text{Dec}_n(c)$ (resp. $\text{Dec}_n(\mathbf{c})$) be the set of decomposition of c (resp. \mathbf{c}) into n reflections (resp. into n elements of T_c) where $n = \text{rank}(W)$. The morphism π bijectively maps T_c to T , hence maps $\text{Dec}_n(\mathbf{c})$ to $\text{Dec}_n(c)$. By definition of $B_c^*(W)$ a decomposition $(t_1, \dots, t_n) \in \text{Dec}_n(c)$ is lifted in $B(W)$ to $(\mathbf{t}_1, \dots, \mathbf{t}_n) \in \text{Dec}_n(\mathbf{c})$. Hence π induces a bijection from $\text{Dec}_n(\mathbf{c})$ to $\text{Dec}_n(c)$.

By the dual braid relations, the Hurwitz action in $B(W)$ preserves $\text{Dec}_n(\mathbf{c})$. Since π is compatible with the Hurwitz action and by Theorem 3.10 $\text{Dec}_n(c)$ is a single orbit, we get the result. □

A question is then to understand which elements of $B(W)$ correspond to the subset D_c of $B_c^*(W)$ through the embedding of $B_c^*(W)$ into $B(W)$. This may be of interest for both computational reasons and for a possible case-free proof of Proposition 3.11.

We will address this question in Sects. 5 and 6 when W is of type A_n or B_n . In the case of reflections the following subsection gives an answer.

3.5 A formula for dual atoms

The following proposition is the analogue for the braid monoid of Proposition 3.4 and the proof is parallel.

Proposition 3.13 *Let $c = s_1 \dots s_n$ be a standard Coxeter element, then, through the embedding of $B_c^*(W)$ into $B(W)$, we have*

$$T_c = \{s_1 s_2 \dots s_i s_{i+1} s_i^{-1} s_{i-1}^{-1} \dots s_1^{-1}, \mid 0 \leq i < 2|T|\},$$

where the index i in s_i is taken modulo n .

Proof Let T'_c be the set of elements $s_1 s_2 \dots s_i s_{i+1} s_i^{-1} s_{i-1}^{-1} \dots s_1^{-1}$ with $i \geq 0$. The elements of T'_c are components of the elements of the Hurwitz orbit of (s_1, \dots, s_n) : this is clear if $i \leq n$ and the elements with $i > n$ are conjugates from the previous ones by a power of c . Hence by Corollary 3.12 we have $T'_c \subset T_c$. Since by definition $|T_c| = |T|$, for proving $T_c = T'_c$ it is sufficient to prove $|T'_c| \geq |T|$. If c is a Coxeter element such that $\ell_S(c^i) = i \ell_S(c)$ for any $i \leq |T|/h$, where $h = 2|T|/n$ is the Coxeter number (the order of c), e.g., if c is bipartite, then the image in W of T'_c is the full set T of reflections of W as seen in the proof of Proposition 3.4, so that the result holds in that case. We now remark that if $\mathbf{c} = s_1 \dots s_n$ and $\mathbf{c}' = s_i \dots s_n s_1 \dots s_{i-1}$ then $T'_{\mathbf{c}'}$ contains $(s_i \dots s_n) T'_c (s_i \dots s_n)^{-1}$ so has cardinality at least $|T'_c|$, whence the result holds, using the fact that all standard Coxeter elements of a given Coxeter system are cyclically conjugate. By [7, Chapter V, § 6 exercise 2] if c is bipartite, we have $\mathbf{c}^h = \mathbf{w}_0^2$, where \mathbf{w}_0 is the lift in $B^+(W)$ of the longest element of W . Since \mathbf{w}_0^2 is central in $B(W)$, one gets the same set T'_c of elements $s_1 s_2 \dots s_i s_{i+1} s_i^{-1} s_{i-1}^{-1} \dots s_1^{-1}$ for $0 \leq i < 2|T|$ as for $i \in \mathbb{N}$. □

4 Rational permutation braids

4.1 Square-free braids

Let (W, S) be a finite Coxeter system. Recall that we denote by $p : B(W) \rightarrow W$ the canonical surjection.

Definition 4.1 An element $\beta \in B(W)$ is a *square-free braid* if it can be represented by a braid word

$$s_{i_1}^{\varepsilon_1} \cdots s_{i_k}^{\varepsilon_k}$$

with $\varepsilon_j \in \{-1, 1\}$ such that $k = \ell_S(p(\beta))$. We denote by $B(W)^{\text{sqf}} \subset B(W)$ the set of square-free braids.

One has $W = B^+(W) \cap B(W)^{\text{sqf}}$ which is also the set of prefixes of the Garside element Δ (see [16, Example 1]). It follows from the definition that any square-free braid is obtained by replacing the various s_{i_j} for $j = 1, \dots, k$ in a reduced S -decomposition $s_{i_1} \cdots s_{i_k}$ of an element $w \in W$ by $s_{i_j}^{\pm 1}$.

4.2 Rational permutation braids

As any group of fractions of a cancellative monoid $B(W)$ is endowed with a left-invariant partial order that we denote by \preceq defined by $u \preceq v$ if $u^{-1}v \in B^+(W)$ which extends the left-divisibility relation on $B^+(W)$.

Definition 4.2 An element $\beta \in B(W)$ is a *rational permutation braid* if it lies in the interval $[\Delta^{-1}, \Delta]$ for the partial order \preceq .

The following proposition can be seen as a particular case of [14, Proposition 5.3]. It explains the terminology as the elements of W are usually called permutation braids.

Proposition 4.3 *Let $\beta \in B(W)$. The following are equivalent:*

- (1) *The braid β is a rational permutation braid,*
- (2) *There exist $x, y \in W$ such that $\beta = x^{-1}y$,*
- (3) *There exist $x, y \in W$ such that $\beta = xy^{-1}$.*

Proof We prove the equivalence of (1) and (2). The equivalence of (1) and (3) is similar.

Since $B^+(W)$ satisfies the Ore conditions, any element of $B(W)$ can be uniquely written as $x^{-1}y$ with x and y in $B^+(W)$ having no common left-divisor. We have $\Delta^{-1} \preceq x^{-1}y \preceq \Delta$ if and only if $x^{-1}y = \Delta^{-1}a = \Delta b^{-1}$ with a, b in $B^+(W)$. So (1) implies that a is a divisor of Δ^2 which by the general properties of Garside monoids (see [15, V, Proposition 3.24]) means that $a = a_1a_2$ with $a_1, a_2 \in W$, whence $x^{-1}y = (a_1^{-1}\Delta)^{-1}a_2$. The Ore conditions imply then that x and y are right-divisors of respectively $a_1^{-1}\Delta$ and a_2 in $B^+(W)$, so that x and y are in W . Conversely, if x and y are in W , that is divide Δ , then $x^{-1}y \preceq x^{-1}\Delta \preceq \Delta$ and $\Delta^{-1} \preceq \Delta^{-1}y \preceq x^{-1}y$. □

We thank Matthew Dyer for having pointed to us the property stated in the following lemma which is a particular case of [19, Section 9.4]. We reprove it here in our particular case.

Lemma 4.4 *If β is a rational permutation braid and if $s_1s_2 \cdots s_k$ is any reduced expression of its image in W , then for $i = 1, \dots, k$ there exists $\varepsilon_i = \pm 1$ such that $\beta = s_1^{\varepsilon_1}s_2^{\varepsilon_2} \cdots s_k^{\varepsilon_k}$. In particular, any rational permutation braid is a square-free braid.*

Proof We prove by induction on k that if $s_1 s_2 \dots s_k$ is a reduced decomposition of an element of W , then for $y \in W$ the element $s_1^{\varepsilon_1} \dots s_k^{\varepsilon_k} y$, where $\varepsilon_i = 1$ if $\ell_S(s_i s_{i+1} \dots s_k y) = \ell_S(s_{i+1} \dots s_k y) + 1$ and $\varepsilon_i = -1$ otherwise is in W . This concludes the proof, since if $\beta = x y^{-1}$ is a rational permutation braid with x and y in W and if $s_1 \dots s_k$ is a reduced decomposition of the image of β in W , then $s_1^{\varepsilon_1} \dots s_k^{\varepsilon_k} y$ is in W and has same image as x , hence is equal to x , so that $\beta = x y^{-1} = s_1^{\varepsilon_1} \dots s_k^{\varepsilon_k}$ is a square-free braid.

By induction hypothesis $x' = s_2^{\varepsilon_1} \dots s_k^{\varepsilon_k} y$ is in W . If $\varepsilon_1 = 1$ then $s_1 x'$ is in W . If $\varepsilon_1 = -1$ then, by the exchange lemma, x' has a reduced expression of the form $s_1 s'_2 \dots s'_m$ so that $s_1^{\varepsilon_1} x' = s'_2 \dots s'_m$ is again in W . □

Example 4.5 Let W be of type A_2 , with simple generating reflections s_1 and s_2 ; the element $w := s_1 s_2 s_1$ in W has exactly two reduced expressions $s_1 s_2 s_1 = s_2 s_1 s_2$. There are six square-free braids having w as image. For example, $\beta = s_1^{-1} s_2 s_1$. Here the reduced expression $s_1 s_2 s_1$ has been lifted to the braid word $s_1^{-1} s_2 s_1$; Lemma 4.4 says that we can lift any reduced expression, in particular we can also lift the reduced expression $s_2 s_1 s_2$ of w in which case one has $\beta = s_2 s_1 s_2^{-1}$.

Remark 4.6 Note that the set of rational permutation braids is the set of braids where “one can apply (mixed) braid relations in reduced words as in the Coxeter group”. Let us be more precise: if β is a rational permutation braid, then one can pass from any shortest expression of β with respect to $S \cup S^{-1}$ to any other one by mixed braid relations as defined in [19], that is, braid relations possibly involving inverses. Moreover, to any reduced expression of the image of β corresponds a shortest braid word for β . Precisely given any two reduced expressions of the image of a braid β in W which differ by a braid relation, then by Lemma 4.4 both reduced expressions can be lifted to reduced braid words for β which differ by a mixed braid relation. For example the relation $s_1^{-1} s_2 s_1 = s_2 s_1 s_2^{-1}$ in Example 4.5 is a mixed braid relation.

5 Mikado braids of type A_n

In this section (W, S) is a Coxeter system of type A_n . The group W is identified with the symmetric group \mathfrak{S}_{n+1} on $\{1, 2, \dots, n+1\}$ and S with the set of simple transpositions, writing $s_i := (i, i + 1)$. The Artin-Tits group $B(W)$ is identified with \mathcal{B}_{n+1} .

5.1 Square-free braids

We give some geometrical properties of square-free braids of type A_n . Firstly, notice that $\beta \in \mathcal{B}_{n+1}$ is a square-free braid if and only if there is a braid diagram for β where any two strands cross at most once. We call a braid diagram *reduced* if it has the minimal number of crossings.

Lemma 5.1 *Removing any strand in a diagram of a square-free braid $\beta \in \mathcal{B}_{n+1}$ gives a diagram for a square-free braid $\beta' \in \mathcal{B}_n$.*

Proof Let D be a diagram for β and D' be the diagram obtained by removing a given strand. Since β is a square-free braid, there is an isotopy which allows one to pass from D to a diagram \tilde{D} where any two strands cross at most once. If we forget the strand we want to remove, our isotopy deforms D' in a diagram \tilde{D}' obtained from \tilde{D} by removing the strand, and in \tilde{D}' any two strands among the remaining n strands cross each other at most once since they were crossing each other at most once in \tilde{D} . □

5.2 Mikado braids

Definition 5.2 A strand of a given reduced diagram D of a braid $\beta \in \mathcal{B}_{n+1}$ which is over the other strands it crosses is called *good*.

Remark 5.3 A strand is good in a reduced diagram for a braid $\beta \in \mathcal{B}_{n+1}^{\text{sqf}}$ if and only if it is good in any reduced diagram for β : indeed, we first claim that the number of crossing between any two strands is constant on the set of reduced diagrams of a square-free braid. To see this, notice that this property holds for the reduced diagrams of the permutation obtained as image of β in the symmetric group, and that the reduced diagrams for the various square-free braids having this permutation as image are obtained by replacing the crossings in the permutation diagrams by positive or negative crossings. Hence the claim holds. Now since β is a square-free braid, any two strands cross at most once, and obviously one cannot continuously deform a positive crossing in a negative one. Hence the number and type of crossing between any two strands is constant on the set of reduced diagrams for β . It follows that a strand which is above all the other in some reduced diagram must be above all the others in all the reduced diagrams.

Definition 5.4 We define *Mikado braids* recursively as


- (1) The braid e is a Mikado braid in \mathcal{B}_1 .
- (2) A braid $\beta \in \mathcal{B}_{n+1}$ is a Mikado braid if in any reduced diagram for β , there exists at least one good strand and if removing any such strand yields a Mikado braid in \mathcal{B}_n .

We denote by $\mathcal{B}_{n+1}^{\text{Mik}}$ the set of Mikado braids. An example of a Mikado braid in \mathcal{B}_7 is given in Fig. 1.

Remark 5.5 Notice that Definition 5.4 is equivalent to the definition of an f -realizable braid from [14, Section 2].

Remark 5.6 If $\beta \in \mathcal{B}_{n+1}^{\text{sqf}}$ with good i th strand such that removing it yields a braid $\beta' \in \mathcal{B}_n^{\text{Mik}}$, then $\beta \in \mathcal{B}_{n+1}^{\text{Mik}}$ (i.e., we can assume that the inductive condition in point (2) of Definition 5.4 is true for one strand instead of any strand). We prove it by induction on rank: indeed, assume that β has another good j th strand. We must show that $\tilde{\beta} \in \mathcal{B}_n$ obtained by removing the j th strand is in $\mathcal{B}_n^{\text{Mik}}$. The i th and j th strand do not cross. The j th strand of β may have become the $(j-1)$ th strand of β' but it is still good. In particular, we can remove it, yielding a braid $\beta'' \in \mathcal{B}_{n-1}^{\text{Mik}}$. But β'' is also obtained from $\tilde{\beta}$ by removing the strand corresponding to the i th strand in β (which may be the $(i-1)$ th strand of $\tilde{\beta}$), hence by induction $\tilde{\beta} \in \mathcal{B}_n^{\text{Mik}}$ since $\beta'' \in \mathcal{B}_{n-1}^{\text{Mik}}$ is.

Fig. 1 A Mikado braid in \mathcal{B}_7






Fig. 2 Illustration of the inductive process used in the proof of Proposition 5.7

Proposition 5.7 *Let $\beta \in \mathcal{B}_{n+1}$. Then β is a Mikado braid if and only if $\beta = \mathbf{x}^{-1} \mathbf{y}$ for some $\mathbf{x}, \mathbf{y} \in \mathbf{W}$, hence by Proposition 4.3 if and only if β a rational permutation braid.*

Proof First, assume that β is a rational permutation braid. We argue by induction on n . The trivial braid e has the claimed form. Now assume $\beta \in \mathcal{B}_{n+1}^{\text{Mik}}$ and consider any reduced braid diagram for it. Then there exists at least one good strand. Removing the rightmost such strand we get a braid $\beta' \in \mathcal{B}_n^{\text{Mik}}$ which by induction can be written as $\mathbf{x}'^{-1} \mathbf{y}'$ for $\mathbf{x}', \mathbf{y}' \in \mathbf{W}'$, where $\mathbf{W}' = \langle s_1, \dots, s_{n-1} \rangle$. Assuming that the strand removed joins the i th point of the sequence above to the j th point of the sequence below, one then has that $\beta = s_i^{-1} s_{i+1}^{-1} \dots s_n^{-1} \mathbf{x}'^{-1} \mathbf{y}' s_n s_{n-1} \dots s_j$ (see Fig. 2), where \mathbf{x}' and \mathbf{y}' are seen in \mathcal{B}_{n+1} under the embedding $\iota_n : \mathcal{B}_n \hookrightarrow \mathcal{B}_{n+1}$. But for any k , the parabolic Coxeter element $s_n s_{n-1} \dots s_k$ is left-reduced with respect to \mathbf{W}' . In particular for $k = i, j$, it implies that both $\mathbf{y}' s_n s_{n-1} \dots s_j$ and $\mathbf{x}' s_n s_{n-1} \dots s_i$ lie in \mathbf{W} .

Conversely, assume that $\beta = \mathbf{x}^{-1} \mathbf{y}$ with $\mathbf{x}, \mathbf{y} \in \mathbf{W}$. Note that in a reduced braid diagram for a (positive) permutation braid, *i.e.*, an element of \mathbf{W} , for all i the i th strand is above the strands numbered 1 to $i - 1$. Hence in \mathbf{x}^{-1} the strand which ends at position $n + 1$ is above all the others, the one ending at position n is just below, and so on. It implies that when concatenating reduced diagrams for \mathbf{x}^{-1} and \mathbf{y} , in the resulting diagram, the $(n + 1)$ th strand of \mathbf{y} is above all the others, the n th strand of \mathbf{y} is just below, and so on. Hence the Mikado property is satisfied, so that β is a Mikado braid. \square

This gives in particular a non-inductive algebraic characterization of rational permutation braids. To summarize, putting Propositions 4.3, 5.7 and Remark 5.5 together we get:

Theorem 5.8 *Let $\beta \in \mathcal{B}_{n+1}$. The following are equivalent:*

- (1) *The braid β is a Mikado braid.*
- (2) *The braid β is a rational permutation braid.*
- (3) *There exist $\mathbf{x}, \mathbf{y} \in \mathbf{W}$ such that $\beta = \mathbf{x}^{-1} \mathbf{y}$.*
- (4) *There exist $\mathbf{x}, \mathbf{y} \in \mathbf{W}$ such that $\beta = \mathbf{x} \mathbf{y}^{-1}$.*
- (5) *The braid β is f -realizable in the sense of [14].*

Remark 5.9 If one is interested in enumerative combinatorics, it is natural to ask for the number $\text{Mik}(n)$ of Mikado braids in \mathcal{B}_n . Using Theorem 5.8, we see that counting Mikado braids is equivalent to counting pairs of permutations $(x, y) \in \mathfrak{S}_n \times \mathfrak{S}_n$ with no common left descents. Indeed, there may be distinct pairs (\mathbf{x}, \mathbf{y}) and $(\mathbf{x}', \mathbf{y}') \in \mathbf{W} \times \mathbf{W}$ such that $\mathbf{x}^{-1} \mathbf{y} = \mathbf{x}'^{-1} \mathbf{y}'$, but taking \mathbf{x} and \mathbf{y} with no common left divisor in $\mathcal{B}(\mathbf{W})^+$ gives unicity. These pairs have been counted in [13] and correspond to the coefficients of a power series given by a Bessel function. The first values are $\text{Mik}(1) = 1, \text{Mik}(2) = 3, \text{Mik}(3) = 19, \text{Mik}(4) = 211, \dots$ There does not seem to exist a simple closed formula for $\text{Mik}(n)$.

5.3 Simple dual braids are Mikado braids

The aim of this Section is to show that the images in \mathcal{B}_{n+1} of the simple elements of a dual braid monoid B_c^* , where c is a standard Coxeter element, are Mikado braids. To this end we first describe a model for the group of fractions of B_c^* using braids in a cylinder and then explain how the embedding of B_c^* into the Artin braid group \mathcal{B}_{n+1} can be viewed using this model.

5.3.1 Noncrossing partitions

Given a standard Coxeter element c , we graphically describe the elements of $\text{Div}(c)$ as follows. The Coxeter elements in W are exactly the $(n+1)$ -cycles; a Coxeter element c is standard if and only if it is an $(n+1)$ -cycle $c = (i_1, i_2, \dots, i_k, \dots, i_{n+1})$ with $i_1 = 1, i_k = n+1$, with the property that the sequence $i_1 i_2 \dots i_k$ increasing and the sequence $i_k \dots i_n i_{n+1}$ decreasing (see [27, Lemma 8.2]). To represent the elements of $\text{Div}(c)$ we use $n+1$ points labeled by $i_1, i_2, \dots, i_k, \dots, i_{n+1}$ in clockwise order on a circle. Like in the case where $c = s_1 s_2 \dots s_n$, the elements of $\text{Div}(c)$ can be represented as a union of polygons having vertices the marked points as follows: to each cycle occurring in the decomposition of $x \in \text{Div}(c)$ into a product of disjoint cycles, one associates the polygon obtained as convex hull of the set of points on the circle labeled by the elements in the support of the cycle (in particular a polygon can be reduced to an edge or even a single point). Such a polygon with vertices labelled (i_1, i_2, \dots, i_k) in clockwise order corresponds to the cycle (i_1, i_2, \dots, i_k) which is the unique cycle dividing c with set of vertices $\{i_1, \dots, i_k\}$. The polygons obtained are pairwise disjoint; equivalently the partition defined by the cycle decomposition of x is noncrossing for this choice of labeling of the circle depending on c . To emphasize that the property to be a noncrossing partition depends on the labeling we will speak of a noncrossing partition of the sequence $(i_1, i_2, \dots, i_k, \dots, i_{n+1})$. So we have defined a bijection between $\text{Div}(c)$ and the set of noncrossing partitions of the sequence $(i_1, i_2, \dots, i_k, \dots, i_{n+1})$.

It will be convenient for the proofs to draw the point with label 1 at the top of the circle, the point with label $n+1$ at the bottom, the points with label in $i_k \dots i_n i_{n+1}$ on the left and the points with label in $i_1 i_2 \dots i_k$ on the right, each point having a specific height depending on its label: that is, if P, Q are two points with respective labels $i, j \in \{1, \dots, n+1\}, i < j$, then P is higher than Q . An example is given in Fig. 3. In case we represent a noncrossing partition we may use curvilinear polygons instead or regular polygons for a more comfortable reading (see Fig. 6, on the left).




Fig. 3 Example of a labeling of the vertices given by the standard Coxeter element $c = s_2 s_1 s_3 s_5 s_4 = (1, 3, 4, 6, 5, 2)$

5.3.2 Dual braids and graphical representation

We follow in this part [5], in particular *loc. cit.* Section 1 and Corollary 3.5, and refer to it for the results.

Definition 5.10 A dual braid is an element of the group of fractions of B_c^* . A simple dual braid is a dual braid which is a simple element of $B_c^* \subset \text{Frac}(B_c^*)$ as described in Sect. 3.2.

To describe the simple dual braids we consider braids where the starting points and the ending points of the strands are on two parallel circles. The points of the two circles are labeled as in Sect. 5.3.1. More precisely, we consider one unit circle in a plane with one complex coordinate at level $t = 0$ and a second unit circle in another such plane at level $t = 1$. So the starting points of the strands have coordinates $(z_k, 0)_{k=1, \dots, n+1}$ with $|z_k| = 1$, and the ending points have coordinates $(z_k, 1)$. The Coxeter element is $c = (i_1, \dots, i_{n+1})$ as in Sect. 5.3.1, when going along the circle in clockwise order starting from $(z_{i_1}, 0)$ the startpoints of the strands appear in the order $(z_{i_1}, 0), (z_{i_2}, 0), \dots, (z_{i_{n+1}}, 0)$, and the same for the endpoints, replacing the second coordinate by 1.

For each pair $\{i, j\}$ one lifts the reflection (i, j) to the braid $\delta_{i,j}$ where the strand starting with $(z_k, 0)$ for $k \neq i, j$ has fixed first coordinate and the strand starting with z_i (resp. z_j) is represented by the path $t \mapsto (t, (z_i + z_j)/2 + (z_i - z_j)/2(\cos(\pi t) + i \varepsilon \sin(\pi t)))$ (resp. same with i and j exchanged) where ε is small enough (see Fig. 4). We draw the figures with the level $t = 0$ above and the level $t = 1$ below.

These braids satisfy the relations in the dual presentation of Proposition 3.8 for the braid group B_{n+1} , that is $\delta_{i,j}\delta_{j,k} = \delta_{j,k}\delta_{k,i}$ when z_i, z_j, z_k are in clockwise order and $\delta_{i,j}\delta_{k,l} = \delta_{k,l}\delta_{i,j}$ when i, j, k, l are 4 distinct points with $[i, j]$ and $[k, l]$ noncrossing (these are exactly the cases where the product of two reflections divide the Coxeter element). The simple dual braids are in one-to-one correspondence with the noncrossing partitions of the sequence (i_1, \dots, i_{n+1}) as described in Sect. 5.3.1. The braid diagram associated to a simple dual braid represented by a polygon $(z_{j_1}, \dots, z_{j_k})$ where the points are in clockwise ordering is the product $\delta_{j_1, j_2}\delta_{j_2, j_3} \dots \delta_{j_{k-1}, j_k}$. In the resulting braid diagram the strand starting with $(z_{j_m}, 0)$ ends with $(z_{j_{m-1}}, 1)$ for $1 < m \leq k$ while the strand starting with $(z_{j_1}, 0)$ ends with $(z_{j_k}, 1)$ (see Fig. 5).

Fig. 4 Braid diagram for the dual braid $\delta_{i,j}$




Fig. 5 The braid associated to a quadrangle





Fig. 6 Passing from a dual to a classical braid

If the Coxeter element is standard the dual braids $\delta_{i,i+1}$ satisfy the usual braid relations and the identification of the group of fractions of the dual braid monoid with the braid group \mathcal{B}_{n+1} maps $\delta_{i,i+1}$ to σ_i . Graphically to recover the usual braid diagram of the image in \mathcal{B}_{n+1} of a simple dual braid β , one has to put the points $z_{i_1}, \dots, z_{i_{n+1}}$ on the circle in such a way that the imaginary parts are in decreasing order and “look from the right” to the braid. This gives a diagram for an Artin braid, which is a horizontal mirror diagram of a diagram for the image in \mathcal{B}_{n+1} of β . Equivalently starting from the noncrossing partition representation of β , one orders the polygons in counterclockwise order and then projects everything to the right; this gives the braid seen from the bottom (see Fig. 6).

The point in this description of the embedding which will be crucial later and follows from our graphical descriptions of the embedding above is that if a polygon in the noncrossing partition representation of β has no polygon at its right (we assume that an edge and a single point are polygons), then the strands from this polygon are above all the strands coming from other polygons in the classical Artin braid representation of the image of β , as one can see in Fig. 6 with the polygon reduced to the edge labeled by 4: on the very right, the strand labeled by 4 appears above all the other strands.

Remark 5.11 Note that the point where the Coxeter element needs to be standard is the identification of $\delta_{i,i+1}$ with σ_i . If the Coxeter element is not standard the braids $\delta_{i,i+1}$ do not satisfy the braid relations.

5.3.3 Removing the strands of simple dual braids

The aim here is to use the graphical representation of simple elements from the previous section to show that simple elements of dual braid monoids are rational permutation braids. The argumentation is therefore topological.

Any $u \in D_c$ always has at least one polygon. The polygons of the graphical representation correspond to the cycles (or blocks) of the corresponding permutation (or noncrossing partition). Consider any block represented by either a polygon or a point which we denote by P and which is right to any other polygon or point, that is, with no other block at its right in the picture (with our conventions described above). It always exists, since the polygons are disjoint and with vertices on the circle. One can then inductively remove all the strands corresponding to edges of P one after the other such that the removed strand at each step is good, starting with any strand having no other strand from P at its right (it always exists and one can keep going inductively since P is a polygon): such a strand is necessarily good. After having removed it one can find a strand which is good at this step, and so on.

After having removed all the strands from P , we go on with another polygon Q which is right to all the remaining polygons, and so on, until all the strands have been removed. Hence we showed:

Theorem 5.12 *Let c be a Coxeter element in a Coxeter group (W, S) of type A_n . Any element of $D_c \subset B_c^* \subset B(W)$ is a Mikado braid, and thus a rational permutation braid.*

5.3.4 Additional properties in case the Coxeter element is linear

We denote by $<$ the Bruhat order on W . In the next proposition, we show that in case $c = s_1 s_2 \cdots s_n$, one can always remove a strand in a braid representing $u \in D_c$ such that the resulting braid v lies in $D_{c'}$ for $c' = s_1 s_2 \cdots s_{n-1}$; this allows us to say a bit more in that case.

Proposition 5.13 *Let $c = s_1 s_2 \cdots s_n$, $u \in D_c$. Then $u \neq e$ can be written in the form $x^{-1}y$ with $x, y \in W$ and $p(x) < p(y)$.*

Proof We argue by induction on the rank. We prove the statement with the inequality $<$ replaced by \leq . It is then clear that the inequality is an equality if and only if $u = e$. The result is trivially true for $n = 1$. Assume $n > 1$. With our choice of Coxeter element, all indices lie on the right part of the circle. If there is a block of u reduced to a single index i , then the i th strand of u must be unbraided, and moreover since all the points lie on the right of the circle, the i th strand lies over all the other strands. We then argue as in the proof of Proposition 5.7; that is, one has $u = s_i^{-1} s_{i+1}^{-1} \cdots s_n^{-1} u' s_n s_{n-1} \cdots s_i$ where $u' \in B_n$ is the braid obtained from u by removing the i th strand. But u' is then in $D_{c'}$, where $c' = s_1 \cdots s_{n-1}$: indeed, it is graphically obtained from the noncrossing representation of u by removing the point with label i and subtracting 1 from each label larger than i , still yielding a diagram of a simple element u' for the smaller linear Coxeter element c' . By induction $u' = x'^{-1}y'$ with $p(x') < p(y')$. Since $s_n \cdots s_i$ is left-reduced with respect to the parabolic subgroup $\langle s_1, \dots, s_{n-1} \rangle$ we get that $p(x' s_n \cdots s_i) < p(y' s_n \cdots s_i)$, hence we have the claimed property with $x = x' s_n \cdots s_i$, $y = y' s_n \cdots s_i$.

Hence we can assume that there is no unbraided strand. But in that case, since the Coxeter element is linear, there must be a polygon in the graphical representation of u having an edge $(i - 1, i)$, that is an i th strand ending at $i - 1$ which is over all the other strands that it crosses. We argue as above, writing this time $u = s_i^{-1} s_{i+1}^{-1} \cdots s_n^{-1} u' s_n s_{n-1} \cdots s_{i-1}$. The element u'

is the simple element for c' obtained by contracting the edge $(i - 1, i)$, hence also identifying the points with labels i and $i - 1$ and subtracting 1 from any point with label bigger than i . Arguing as above we get the claim. \square

6 Mikado braids of type B_n

6.1 Coxeter and Artin-Tits groups of type B_n

The Coxeter group W of type B_n is the group of fixed points in the Coxeter group W' of type A_{2n-1} under the diagram automorphism τ which exchanges s_i and s_{2n-i} . This fact lifts to the corresponding Artin-Tits groups (see e.g., [33, Corollary 4.4]). So we shall see the Artin-Tits group $B(W)$ of type B_n as the group of fixed points in the Artin-Tits group of type A_{2n-1} under the diagram automorphism which we still denote by τ , that is, $B(W')^\tau = B(W)$. Topologically this means that the Artin-Tits group $B(W)$ can be seen as the group of symmetric braids in \mathcal{B}_{2n} (the symmetry exchanges the i th and the $(2n + 1 - i)$ th strands and exchanges under-/over-crossings).

We write \mathcal{B}_{2n}^B for the braids of type B_n identified with \mathcal{B}_{2n}^τ . For convenience, since \mathcal{B}_{2n}^B consists of symmetric braids, we label the strands of \mathcal{B}_{2n} by $-n, -n + 1, \dots, -1, 1, \dots, n$ rather than by $1, 2, \dots, 2n$. We write t_0, \dots, t_{n-1} for the Coxeter generators of W with relations

$$\begin{aligned} t_0 t_1 t_0 t_1 &= t_1 t_0 t_1 t_0, \\ t_i t_{i+1} t_i &= t_{i+1} t_i t_{i+1} \text{ if } 1 \leq i \leq n - 2, \\ t_i t_j &= t_j t_i \text{ if } |i - j| > 1. \end{aligned}$$

Inside W' one has $t_0 = s_n$ and $t_i = s_i s_{2n-i}$ for $i = 1, \dots, n - 1$, where $s_1 = (-n, -n + 1), s_2 = (-n + 1, -n + 2), \dots, s_n = (-1, 1), s_{n+1} = (1, 2), \dots, s_{2n-1} = (n - 1, n)$ in \mathfrak{S}_{2n} seen as the group of permutations of $\{-n, \dots, -1, 1, \dots, n\}$.

This lifts to the braid groups: the generators of $B(W)$ seen in \mathcal{B}_{2n} are $t_0 = s_n$ and $t_i = s_i s_{2n-i}$ for $i = 1, \dots, n - 1$.

6.2 Mikado braids of type B_n

Definition 6.1 A braid $\beta \in \mathcal{B}_{2n}^B$ is a *Mikado braid* (of type B_n) if when viewed in \mathcal{B}_{2n} , starting from any diagram D for β one can inductively remove pairs of symmetric strands, one being above all the other strands (so that its symmetric is under all the other strands).

We write $\mathcal{B}_{2n}^{B, \text{Mik}}$ for the set of Mikado braids of type B_n . It follows from the definition that $\mathcal{B}_{2n}^{B, \text{Mik}} = \mathcal{B}_{2n}^B \cap \mathcal{B}_{2n}^{\text{Mik}}$. Hence we can use the geometric properties of Mikado braids of type A_n given in Sect. 5.2.

Proposition 6.2 Let $\beta \in \mathcal{B}_{2n}^B$; then β is a Mikado braid of type B_n if and only if $\beta = x^{-1}y$ for some $x, y \in W$, hence by Proposition 4.3 if and only if β a rational permutation braid.

Note that here W refers to the lift of the Coxeter group W of type B_n in the braid group of type B_n .

Proof A braid $\beta \in \mathcal{B}_{2n}$ is in $\mathcal{B}_{2n}^{B, \text{Mik}}$ if and only if it is a Mikado braid in \mathcal{B}_{2n} and it is fixed by τ . By Proposition 5.7, this is equivalent to β being of the form $x^{-1}y$ with x and y in W' and β fixed by τ . Since there are gcds in Artin-Tits monoids of spherical type, the expression $x^{-1}y$

is unique under the condition that x and y have no common left-divisor (in the Artin-Tits monoid of type A_{2n-1}). Since τ induces an automorphism of the monoid, if x and y satisfy the unicity condition, then $x^{-1}y$ is τ -fixed if and only if each one of x and y is τ -fixed, that is, is in $W^\tau = W$. □

This gives in particular a non-inductive algebraic characterization of rational permutation braids of type B_n . To summarize, putting Propositions 4.3 and 6.2 together we get:

Theorem 6.3 *Let $\beta \in B(W) \cong \mathcal{B}_{2n}^B$. The following are equivalent:*

- (1) *The braid β is a Mikado braid.*
- (2) *The braid β is a rational permutation braid.*
- (3) *There exist $x, y \in W$ such that $\beta = x^{-1}y$.*
- (4) *There exist $x, y \in W$ such that $\beta = xy^{-1}$.*

Problem 6.4 What is the number of Mikado braids of type B_n ? Unlike in type A_n (see Remark 5.9), we have no hint towards an answer to this question.

6.3 Simple dual braids are Mikado braids

6.3.1 Graphical representation of simple elements

Through the embedding $W = W^\tau \subset W'$, the standard Coxeter elements of W identify with the τ -fixed standard Coxeter elements of W' . Moreover if T' denotes the set of reflections of W' and T the set of reflections of W , and if c is a τ -fixed standard Coxeter element of W' , then for x and y in W one has $x \preceq_T y \preceq_T c$ if and only if $x \preceq_{T'} y \preceq_{T'} c$ (see [10, Lemma 4.8]).

Remark 6.5 Note that a reflection of W is not a reflection of W' in general.

Since the dual braid monoid of type A_{2n-1} is defined by a presentation with generators the left-divisors of the chosen Coxeter element c and with relations given by $\ell_{T'}$ -shortest decompositions of c , the above observations imply that the dual braid monoid of type B_n identifies with the fixed points of τ in the dual braid monoid of type A_{2n-1} for the same Coxeter element.

The τ -fixed standard Coxeter elements of W' are the $2n$ -cycles of the form $c = (i_1, i_2, \dots, i_n, -i_1, -i_2, \dots, -i_n)$ with $\{i_1, i_2, \dots, i_n\} = \{1, 2, \dots, n\}$ and the sequence $i_1 i_2 \dots i_n$ first increasing, then decreasing.

We also get that the elements of $\text{Div}(c)$ for a Coxeter element $c \in W$ are in one-to-one correspondence with the τ -fixed noncrossing partitions of $2n$ points labelled $-n, \dots, -1, 1, 2, \dots, n$ on a circle with the clockwise order on the labels in the order given by c (see Fig. 7).


These symmetric partitions are called *noncrossing partitions of type B_n* associated to c (see [35]).

Such a partition corresponds to a τ -fixed braid in \mathcal{B}_{2n} by the same process as explained in Sect. 5.3.2.

6.3.2 Removing pairs of strands of simple dual braids

The argumentation to show inductively that any simple dual braid $x \in \mathcal{D}_c$ of type B_n is a Mikado braid can be led as in type A_n , that is, by looking at the graphical representation of

Fig. 7 Example of a labeling of the vertices given by the Coxeter element $c = t_1 t_2 t_0 t_4 t_3$ and of a noncrossing partition of type B_5 for this Coxeter element. The corresponding element of $\text{Div}(c)$ is $(-5, -4, -1)(5, 4, 1)(2, 3, -2, -3)$



x , equivalently the number of blocks of the noncrossing partition $x \in \text{Div}(c)$, and removing pairs of strands. We start from a polygon P which is to the right of any other polygon. Either P has a symmetric polygon $\bar{P} \neq P$ such that j indexes a vertex of P if and only if $-j$ indexes a vertex of \bar{P} ; or P is its own symmetric, in particular P has vertices on both sides of the circle and an index j indexes a vertex of P if and only if $-j$ indexes a vertex of P .

In the first case, we remove inductively pairs of strands of P , one corresponding to an edge of P and its symmetric which is an edge of \bar{P} as we did in type A_n , in such an order that at each step, the removed strand from P is above all the others (which implies that the corresponding symmetric strand which is also removed is below all the others), that is, the corresponding edge should be right to any other edge of P . In the second case, we do exactly the same, except that pairs of strands of P are removed simultaneously.

After having removed all the strands from P , we go on with another polygon Q which is right to all the remaining polygons, and so on, until all the strands have been removed. Hence we showed:

Theorem 6.6 *Let c be a Coxeter element in a Coxeter group (W, S) of type B_n . Any element of $\mathcal{D}_c \subset B_c^* \subset B(W)$ is a Mikado braid of type B_n , and thus a rational permutation braid.*

7 Dihedral and exceptional types

Theorem 7.1 *Let c be a Coxeter element in a Coxeter system (W, S) of type $H_3, H_4, I_2(m)$ ($m \geq 3$), E_n ($n = 6, 7, 8$) or F_4 ; any element of $\mathcal{D}_c \subset B_c^* \subset B(W)$ is a rational permutation braid.*

Proof For exceptional types we have checked the result using the program CHEVIE [34]. Let us prove it for dihedral types. By Proposition 4.3., it suffices to show that any element in \mathcal{D}_c has the form $\mathbf{x} \mathbf{y}^{-1}$ for $\mathbf{x}, \mathbf{y} \in \mathbf{W}$. Since $S = \{s, t\}$ there are only two possible choices of Coxeter element. Let $c = st$. Then the elements of \mathcal{D}_c are the identity element, \mathbf{c} and the elements of \mathcal{T}_c . By considering the Hurwitz action on (s, t) we see that the elements of \mathcal{T}_c are exactly the $\underbrace{(sts \cdots)}_k \underbrace{(sts \cdots)}_{k-1}^{-1}$ with $1 \leq k \leq m$, which concludes the proof. \square

8 Positivity properties

The whole paper is motivated by a positivity conjecture on the expansion of the images of simple dual braids in a canonical basis of the Iwahori-Hecke algebra of the Coxeter group. Using work from the previous sections we prove these properties here for finite irreducible Coxeter groups of type other than type D_n .

8.1 Iwahori-Hecke algebra of a Coxeter system

Definition 8.1 Let (W, S) be a Coxeter system. For $s, t \in S$, recall that $m_{s,t}$ denotes the order of st in W . The Iwahori-Hecke algebra $H(W) = H(W, S)$ of (W, S) is the associative, unital $\mathbb{Z}[v, v^{-1}]$ -algebra generated by a copy $\{T_s \mid s \in S\}$ of S with relations

$$\begin{aligned}
T_s^2 &= (v^{-2} - 1)T_s + v^{-2} & \forall s \in S, \\
\underbrace{T_s T_t \cdots}_{m_{s,t} \text{ copies}} &= \underbrace{T_t T_s \cdots}_{m_{s,t} \text{ copies}}, & \forall s, t \in S.
\end{aligned}$$

The algebra $H(W)$ has a standard basis $\{T_w\}_{w \in W}$ where T_w is the image of $w \in W$ under the unique group morphism $a : B(W) \rightarrow H(W)^\times$ such that $s \mapsto T_s$ for all $s \in S$. It also has two canonical bases $\{C_w\}_{w \in W}$ and $\{C'_w\}_{w \in W}$ defined in [31]. The algebra $H(W)$ has a unique semilinear involution j_H such that $j_H(T_s) = -v^2 T_s$ for all $s \in S$ and $j_H(v) = v^{-1}$. The two canonical bases are then related by the equalities

$$C_w = (-1)^{\ell_S(w)} j_H(C'_w), \quad \forall w \in W. \tag{8.2}$$

A reference for this material is [29, Section 7.9].

8.2 Positivity properties of simple dual braids

There are many positivity statements involving the canonical bases of $H(W)$. In this section we prove positivity results on the expansion of images of simple dual braids in $H(W)$ when they are expressed in the basis $\{C_w\}$. This comes as a corollary of the results of the previous sections and of the following:

Theorem 8.3 ([20, Corollary 2.9]) *Let (W, S) be a finite Weyl group, then for all $x, y \in W$, one has*

$$T_x^{-1} T_y \in \sum_{w \in W} \mathbb{N}[v, v^{-1}] C_w.$$

Remark 8.4 Dyer and Lehrer showed that in case (W, S) is a finite Coxeter group, the statement of the above theorem (for all x, y) is equivalent to the statement that $C'_x T_y$ has a positive expansion in the standard basis (for all x, y); they then showed that this last statement holds for Weyl groups ([20, Proposition 2.7 and Theorem 2.8]). Dyer later showed in [18, Conjecture 7(b)]¹ (using partially unpublished results) that the Kazhdan-Lusztig positivity conjecture (now proven in [21] as a corollary of Soergel conjecture [36]) implies the positivity of the expansion of $C'_x T_y$ in the standard basis for all finite Coxeter groups. Hence we can assume that Theorem 8.3 holds for all finite Coxeter groups.

Theorem 8.3 together with Proposition 4.3 yields:

Proposition 8.5 *Let $\beta \in B(W)$ be a rational permutation braid. Then*

$$a(\beta) \in \sum_{w \in W} \mathbb{N}[v, v^{-1}] C_w.$$

Proof This is an immediate consequence of Theorem 8.3, Proposition 4.3 and the fact that $a(w) = T_w$ for any $w \in W$. □

¹ The authors thank Matthew Dyer for pointing out this fact to the second author.

Theorem 8.6 *Let (W, S) be a finite irreducible Coxeter system of type other than D_n and let c be any Coxeter element in W ; then for any $\mathbf{u} \in \mathbf{D}_c$, one has that*

$$a(\mathbf{u}) \in \sum_{w \in W} \mathbb{N}[v, v^{-1}]C_w.$$

Proof This follows from Theorems 5.12, 6.6, 7.1 and Proposition 8.5. □

Notice that the proof of this theorem required to prove that simple dual braids can be written in the form $\mathbf{x}^{-1}\mathbf{y}$, which was done in types A_n and B_n by using the geometry of Artin braids.

We are not able to prove this property for type D_n , but both computations and the fact that it holds in any other spherical type leads us to conjecture that it again holds:

Conjecture 8.7 *Let (W, S) be a Coxeter system of type D_n . Let c be any Coxeter element in W ; then any element of $\mathbf{D}_c \subset B_c^* \subset B(W)$ is a rational permutation braid.*

Note that if this conjecture is true we will have the same positivity property for type D_n as for all other spherical types.

Problem 8.8 Does there exist a model for the braid group of type D_n by Artin-like braids which could be used to prove that simple elements of dual braid monoids are rational permutation braids? A model for type D_n can be found in [1], but the analogues of the Mikado braids in this model are not the expected ones.

Problem 8.9 Is there a uniform approach to show that simple elements of dual braid monoids are rational permutation braids?

8.3 Temperley-Lieb algebra and monomial basis

Since elements of \mathbf{W} , that is, images of the simple elements for the classical Garside structure provide a basis of the Iwahori-Hecke algebra, one might investigate the linear independence of images of elements of \mathbf{D}_c in $H(W)$. The set \mathbf{D}_c is too small to give a basis of $H(W)$: indeed, the algebra $H(W)$ has rank $|W|$ while $\text{Div}(c)$ is in general a strict subset of W counted by the generalized Catalan number of type W (see for instance [2]). Nevertheless, in type A_n , the set \mathbf{D}_c gives a basis of a remarkable quotient of $H(W)$, the *Temperley-Lieb algebra*. From now on, (W, S) is a Coxeter system of type A_n .

Definition 8.10 The *Temperley-Lieb algebra* TL_n is the associative, unital $\mathbb{Z}[v, v^{-1}]$ -algebra obtained as quotient of $H(W)$ by the two-sided ideal generated by the elements

$$\sum_{w \in \langle s_i, s_{i+1} \rangle} T_w,$$

for $i = 1, \dots, n - 1$, where $s_i = (i, i + 1)$. We write $\theta : H(W) \twoheadrightarrow \text{TL}_n$ for the quotient map. Alternatively, one gets an isomorphic algebra by taking the quotient by the two-sided ideal generated by the elements

$$\sum_{w \in \langle s_i, s_{i+1} \rangle} (-1)^{\ell_S(w)} v^{2\ell_S(w)} T_w,$$

for $i = 1, \dots, n - 1$ (see e.g., [28, Section 2.3 and Remark 2.4]). We write $\theta' : H(W) \twoheadrightarrow \text{TL}_n$ for this alternative quotient map.

The algebra TL_n has a presentation by generators b_{s_1}, \dots, b_{s_n} and relations

$$\begin{aligned} b_{s_i} b_{s_{i\pm 1}} b_{s_i} &= b_{s_i} && \forall i, i \pm 1 \in \{1, \dots, n\}, \\ b_{s_i} b_{s_j} &= b_{s_j} b_{s_i} && \forall i, j \in \{1, \dots, n\} \text{ with } |i - j| > 1, \\ b_{s_i}^2 &= (v + v^{-1})b_{s_i} && \forall i \in \{1, \dots, n\}. \end{aligned}$$

We have (see [28, Section 2.3 and Remark 2.4])

$$\theta(T_{s_i}) = v^{-1}b_{s_i} - 1, \theta'(T_{s_i}) = v^{-2} - v^{-1}b_{s_i}. \tag{8.11}$$

Definition 8.12 An element $w \in W$ is *fully commutative* if one can pass from any reduced S -decomposition of w to any other one by applying only relations of the form $s_i s_j = s_j s_i$ for $|i - j| > 1$. We denote by W_f the set of fully commutative elements. For details on fully commutative elements we refer to [37].

The algebra TL_n has a basis indexed by the elements of W_f . It is built as follows; given any $w \in W_f$ with reduced expression $st \cdots u$, the above presentation shows that the product $b_s b_t \cdots b_u$ does not depend on the choice of the reduced expression and therefore we can denote it by b_w . The set $\{b_w\}_{w \in W_f}$ turns out to yield a basis of TL_n (see [30, Corollary 5.32]).

8.4 Projection of the canonical basis

The basis $\{b_w\}_{w \in W_f}$ turns out to be related to the canonical basis $\{C'_w\}_{w \in W}$ of $H(W)$ as follows:

Theorem 8.13 ([22], Section 3) *The basis $\{b_w\}_{w \in W_f}$ is the projection of $\{C'_w\}_{w \in W}$ by θ . In symbols, for $w \in W_f$, one has $\theta(C'_w) = b_w$ while $\theta(C'_w) = 0$ for $w \notin W_f$.*

8.5 Zinno basis

The Temperley-Lieb algebra comes equipped with a semilinear involution j_{TL_n} such that $j_{TL_n}(v) = v^{-1}$ and $j_{TL_n}(b_s) = b_s$ for any $s \in S$. To realize the algebra TL_n as a quotient of $\mathbb{Z}[v, v^{-1}][\mathcal{B}_n]$, we use the composition of $a' : B(W) \rightarrow H(W), s \mapsto vT_s$ with θ' , that is, by (8.11) any generator $s_i \in \mathcal{B}_n$ is mapped to $v^{-1} - b_{s_i}$. We set $\omega := \theta' \circ a'$.

Theorem 8.14 ([40], Theorem 2, [32], Theorem 1) *Let $c = s_n s_{n-1} \cdots s_1$; then $\omega(D_c)$ yields a basis of TL_n .*

The above Theorem is stated differently in [40] since dual braid monoids had not been introduced when that paper appeared. Notice that the conventions used in [40] correspond to the quotient map $\theta \circ a$, which sends s_i to $v^{-1}b_{s_i} - 1$. Since one passes from one to the other by composing with a semilinear automorphism of $\mathbb{Z}[v, v^{-1}]$ -modules and we are here interested in bases, this does not affect the result.

This result is also generalized to arbitrary standard Coxeter elements in [39, Corollary 5.2.9] and [25, Theorem 3.8.28 and Remark 3.8.29]:

Theorem 8.15 *For any standard Coxeter element c , the set $\omega(D_c)$ yields a basis of TL_n .*

Hence the images of the simple elements of the Garside monoids B_c^* yield a basis of TL_n , in much the same way that the images of the simple elements of the Garside monoid $B^+(W)$ yield the standard basis of $H(W)$.

Note that one may have $x \in \text{Div}(c) \cap \text{Div}(c')$ for some $c' \neq c$, in which case the corresponding elements of \mathbf{D}_c and $\mathbf{D}_{c'}$, hence also the corresponding basis elements of TL_n are not equal in general. For example, in type A_2 with $c = s_1s_2$, $c' = s_2s_1$, the reflection $s_1s_2s_1$ lies in $\text{Div}(c) \cap \text{Div}(c')$ (as any reflection). The corresponding element of \mathbf{D}_c , *resp.* $\mathbf{D}_{c'}$ is $s_1s_2s_1^{-1}$, *resp.* $s_1^{-1}s_2s_1$. The two elements are not equal.

8.6 Positivity consequences

The two quotient maps from Sect. 8.3 are related by $\theta = j_{\text{TL}_n} \circ \theta' \circ j_H$. Using this fact together with Theorems 8.6, 8.13 and Equality (8.2) we get:

Theorem 8.16 *Let c be any standard Coxeter element, let $\{Z_x\}_{x \in \text{Div}(c)}$ be the corresponding basis of TL_n , i.e., $Z_x := \omega(\mathbf{x})$ for $x \in \text{Div}(c)$. Then for any $x \in \text{Div}(c)$, one has*

$$Z_x \in \sum_{w \in W_f} \mathbb{N}[v, v^{-1}](-1)^{\ell_S(w)} b_w.$$

One can show that there exist total orders on the sets $\text{Div}(c)$ and W_f such that the base change from $\{b_w\}_{w \in W_f}$ to $\{Z_x\}_{x \in \text{Div}(c)}$ is upper triangular. Zinno shows it for $c = s_1s_2 \cdots s_n$ in [40] and the result is extended in the second author's thesis [25] to arbitrary c . As noticed in [26], for $c = s_1s_2 \cdots s_n$, the order given by Zinno can be refined into the Bruhat order on $\text{Div}(c)$, which is studied extensively and combinatorially in [27], where the orders to consider for other choices of Coxeter elements are also described. Closed formulas for some of the coefficients (in the inverse matrix) are given in [26] in case the Coxeter element is $c = s_1s_2 \cdots s_n$ (which can be obviously adapted for $c = s_ns_{n-1} \cdots s_1$), but we do not have a closed formula for them in full generality.

Remark 8.17 In type B_n , Theorem 8.15 also holds (see [38]). In that case, the diagram basis of the Temperley-Lieb quotient is not the projection of the canonical basis of the corresponding Iwahori-Hecke algebra (see [22]), hence there is no analogue of Theorem 8.16 in that case. In type D_n , the cardinality of $|\mathbf{D}_c|$ is bigger than the dimension of the Temperley-Lieb algebra, but the images of the elements of \mathbf{D}_c in it still generate the whole algebra, as shown in [39].

Acknowledgements We thank Patrick Dehornoy, Matthew Dyer and Jean Michel for useful discussions. We also thank the anonymous referee for his careful reading of the manuscript and many interesting comments and remarks.

References

1. Allcock, D.: Braid pictures for Artin groups. *Trans. Am. Math. Soc.* **354**, 345–3474 (2002)
2. Armstrong, D.: Generalized noncrossing partitions and combinatorics of Coxeter groups. *Memoirs of the American Mathematical Society*, vol. 202, no. 949 (2009)
3. Baumeister, B., Dyer, M., Stump, C., Wegener, P.: A note on the transitive Hurwitz action on decompositions of parabolic Coxeter elements. *Proc. Am. Math. Soc., Ser. B* **1**, 149–154 (2014)
4. Bessis, D.: The dual braid monoid. *Ann. Sci. École Normale Supérieure* **36**, 647–683 (2003)
5. Bessis, D., Digne, F., Michel, J.: Springer theory in braid groups and the Birman-Ko-Lee monoid. *Pacific J. Math.* **205**, 287–309 (2002)
6. Birman, J., Ko, K.H., Lee, S.J.: A new approach to the word and conjugacy problems in the Braid groups. *Adv. Math.* **139**, 322–353 (1998)
7. Bourbaki, N.: *Groupes et algèbres de Lie*, chapitres 4,5 et 6, Masson (1981)
8. Brieskorn, E., Saito, K.: Artin-Gruppen und Coxeter-Gruppen. *Invent. Math.* **17**, 245–271 (1972)
9. Brady, N., McCammond, J., Mühlherr, B., Neumann, W.: Rigidity of Coxeter groups and Artin groups. *Geom. Dedic.* **94**, 91–109 (2002)

10. Brady, T., Watt, C.: $K(\pi, 1)$'s for Artin groups of finite type. *Geom. Dedic.* **94**, 225–230 (2002)
11. Brady, T., Watt, C.: Noncrossing partitions lattices in finite real reflection groups. *Trans. Am. Math. Soc.* **360**, 1983–2005 (2008)
12. Broué, M., Michel, J.: Sur certains éléments réguliers des groupes de Weyl et les variétés de Deligne-Lusztig associées. *Prog. Math.* **141**, 73–139 (1997)
13. Carlitz, L., Scoville, R., Vaughan, T.: Enumeration of pairs of permutations and sequences. *Bull. Am. Math. Soc.* **80**, 881–884 (1974)
14. Dehornoy, P.: Three-dimensional realizations of braids. *J. Lond. Math. Soc.* **60**, 108–132 (1999)
15. Dehornoy, P., Digne, F., Krammer, D., Godelle, E., Michel, J.: Foundations of Garside theory. *Tracts in Mathematics*, vol. 22, pp. 690. European Mathematical Society (2015)
16. Dehornoy, P., Paris, L.: Gaussian groups and Garside groups, two generalizations of Artin groups. *Proc. Lond. Math. Soc.* **79**, 569–604 (1999)
17. Deligne, P.: Les immeubles des groupes de tresses généralisés. *Invent. Math.* **17**, 273–302 (1972)
18. Dyer, M.J.: Representation theories from Coxeter groups, Representations of groups (Banff, AB, 1994), CMS Conf. Proc. 16, 105–139 Amer. Math. Soc, Providence, RI, (1995)
19. Dyer, M.J.: Modules for the dual nil Hecke ring. <http://www3.nd.edu/dyer/papers/nilhecke>
20. Dyer, M.J., Lehrer, G.I.: On positivity in Hecke algebras. *Geom. Dedic.* **35**, 115–125 (1990)
21. Elias, B., Williamson, G.: The Hodge theory of Soergel bimodules. *Ann. Math.* **180**, 1089–1136 (2014)
22. Fan, C.K., Green, R.M.: Monomials and Temperley-Lieb algebras. *J. Algebra* **190**, 498–517 (1997)
23. Geck, M., Pfeiffer, G.: Characters of finite Coxeter groups and Iwahori-Hecke algebras, London Math. Soc. Monographs, New Series 21, Oxford University Press (2000)
24. Franzsen, W.N., Howlett, R.B., Mühlherr, B.: Reflections in abstracts Coxeter groups. *Comment. Math. Helv.* **81**, 665–697 (2006)
25. Gobet, T.: Bases of Temperley-Lieb algebras, Ph.D. thesis, Université de Picardie, (2014)
26. Gobet, T.: Noncrossing partitions, fully commutative elements and bases of the Temperley-Lieb algebra. *J. Knot Theory Ramif.* **26**, 27 (2016)
27. Gobet, T., Williams, N.: Noncrossing partitions and Bruhat order. *Eur. J. Comb.* **53**, 8–34 (2016)
28. Halverson, T., Mazzocco, M., Ram, A.: Commuting families in Hecke and Temperley-Lieb algebras. *Nagoya Math. J.* **195**, 125–152 (2009)
29. Humphreys, J.: Reflection Groups and Coxeter Groups, Cambridge Studies in Advanced Mathematics, vol. 29. Cambridge University Press, Cambridge (1990)
30. Kassel, C., Turaev, V.: Braid Groups, Graduate Texts in Mathematics, vol. 247. Springer, New York (2008)
31. Kazhdan, D., Lusztig, G.: Representations of Coxeter groups and Hecke algebras. *Invent. Math.* **53**, 165–184 (1979)
32. Lee, E.K., Lee, S.J.: Dual presentation and linear basis of the Temperley-Lieb algebras. *J. Korean Math. Soc.* **47**, 445–454 (2010)
33. Michel, J.: A note on words in braid monoids. *J. Algebra* **215**, 366–377 (1999)
34. Michel, J.: The development version of the CHEVIE package of GAP3. *J. Algebra* **435**, 308–336 (2015)
35. Reiner, V.: Non-crossing partitions for classical reflection groups. *Discret. Math.* **177**, 195–222 (1997)
36. Soergel, W.: Kazhdan-Lusztig polynomials and indecomposable bimodules over polynomial rings. *J. Inst. Math. Jussieu* **6**, 501–525 (2007)
37. Stembridge, J.R.: On the fully commutative elements of Coxeter groups. *J. Algebr. Comb.* **5**, 353–385 (1996)
38. Vincenti, C.: Algèbre de Temperley-Lieb de type B. *C. R. Math. Acad. Sci. Paris* **342**, 233–236 (2006)
39. Vincenti, C.: Monoïde dual, antichaînes de racines et algèbres de Temperley-Lieb, Ph.D. thesis, Université de Picardie(2007)
40. Zinno, M.G.: A Temperley-Lieb basis coming from the braid group. *J. Knot Theory Ramif.* **11**, 575–599 (2002)