



A Bloch–Ogus theorem for henselian local rings in mixed characteristic

Johannes Schmidt¹ · Florian Strunk²

Received: 22 October 2019 / Accepted: 23 January 2023 / Published online: 27 February 2023
© The Author(s) 2023

Abstract

We show a conditional exactness statement for the Nisnevich Gersten complex associated to an \mathbb{A}^1 -invariant cohomology theory with Nisnevich descent for smooth schemes over a Dedekind ring with only infinite residue fields. As an application we derive a Nisnevich analogue of the Bloch–Ogus theorem for étale cohomology over a henselian discrete valuation ring with infinite residue field.

1 Introduction

Given an \mathbb{A}^1 -invariant cohomology theory E for smooth varieties X over a field k with Nisnevich descent, Colliot-Thélène, Hoobler and Kahn proved in [4] the exactness of the associated Gersten complex

$$\begin{aligned} 0 \rightarrow H^n(Y, E) \rightarrow \bigoplus_{z \in Y^{(0)}} H_{\{z\}}^n(Y, E) \rightarrow \bigoplus_{z \in Y^{(1)}} H_{\{z\}}^{n+1}(Y, E) \rightarrow \dots \\ \dots \rightarrow \bigoplus_{z \in Y^{(d-1)}} H_{\{z\}}^{n+d-1}(Y, E) \rightarrow \bigoplus_{z \in Y^{(d)}} H_{\{z\}}^{n+d}(Y, E) \rightarrow 0, \end{aligned} \quad (1a)$$

where $Y = \text{Spec}(\mathcal{O}_{X,x})$ is the local scheme at a point x and d is the dimension of X . The main ingredient of their proof is a geometric presentation theorem [4, Theorem 3.1.1] for a closed immersion $Z \hookrightarrow X$ which is due to Gabber. If E is algebraic K -theory, this result implies the Gersten conjecture for smooth schemes over a field, originally proved by Quillen [11, Theorem 5.11]. Taking E as étale cohomology with constant torsion coefficients defined over k , one obtains the Bloch–Ogus theorem [3].

The authors are supported by the SFB/CRC 1085 *Higher Invariants* (Regensburg) funded by the DFG and the DFG-Forschergruppe 1920 *Symmetrie, Geometrie und Arithmetik* (Heidelberg–Darmstadt).

✉ Florian Strunk
florian.strunk@ur.de

Johannes Schmidt
jschmidt@mathi.uni-heidelberg.de

¹ Mathematisches Institut, Universität Heidelberg, Heidelberg 69120, Germany

² Fakultät für Mathematik, Universität Regensburg, Regensburg 93040, Germany

In the mixed characteristic case of a discrete valuation ring with infinite residue field, an analogue of Gabber’s geometric presentation theorem for a closed immersion $Z \hookrightarrow X$ was shown in [12, Theorem 2.1]. However, there are two crucial differences to the equal characteristic case: Firstly, one has to require that the closed subscheme Z does not contain any irreducible component of the special fibre of X . Secondly, the presentation is not Zariski- but only Nisnevich-local in X .

In this paper, our goal is to adopt the techniques of Colliot-Thélène, Hoobler and Kahn to the mixed characteristic case using the more restricted version of the presentation theorem. Our main result is the following (see Theorem 5.13, below).

Theorem *Let S be a Dedekind scheme with only infinite residue fields and E an \mathbb{A}^1 -invariant cohomology theory for smooth schemes of finite type over S with Nisnevich descent. Let X/S be such a smooth scheme of dimension d , $x \in X$ a point and $Y = \text{Spec}(\mathcal{O}_{X,x}^h)$ the Henselian local scheme at x .*

- (1) *The Gersten complex (1a) is exact possible except at the first and third (non-trivial) spot.*
- (2) *If for each point x of X the forget support map for the special fibre Y_x*

$$\mathbb{R}\Gamma_{Y_x}(Y_{\text{Nis}}, E) \rightarrow \Gamma(Y, E)$$

is trivial, then the Gersten complex is exact everywhere.

In presence of the second condition, one obtains the usual resolution of the Nisnevich sheafification of the cohomology given by E by flabby Nisnevich sheaves. If E is algebraic K -theory, the theorem was known before, see [2] and [8].

As an application of the theorem above, we derive the following analogue of the Bloch–Ogus theorem in mixed characteristic (see Corollary 6.10).

Theorem *Let $Y = \text{Spec}(\mathcal{O}_{X,x}^h)$ be a Henselian local scheme of a d -dimensional smooth scheme X of finite type over a Henselian discrete valuation ring \mathfrak{o} with infinite residue field of characteristic p . Let K be a locally constant constructible sheaf of \mathbb{Z}/m -modules for m prime to p on the small étale site of $\text{Spec}(\mathfrak{o})$. Then the Gersten complex*

$$0 \rightarrow H^n(Y_{\text{ét}}, K) \rightarrow \bigoplus_{z \in Y^{(0)}} H^n(k(z), K) \rightarrow \cdots \rightarrow \bigoplus_{z \in Y^{(d)}} H^{n-d}(k(z), K(-d)) \rightarrow 0.$$

is exact. Here $H^i(k(z), -)$ denotes the Galois-cohomology of the field $k(z)$.

We remark that in [7] Geisser derived the exactness of the above Gersten complex from the Bloch–Kato–Conjecture even for a (Zariski) local scheme $Y = \text{Spec}(\mathcal{O}_{X,x})$ but only for coefficients $K = \mu_m^{\otimes r}$ where $n \leq r$. Our method of proof is more elementary, at least if the residue and quotient field of \mathfrak{o} are perfect (see Remark 6.9 and Remark 6.11).

The organization of the paper is as follows. In Sect. 2 we recall some known results on basechange of presheaves of spectra. We include a short reminder on elementary properties of the codimension for schemes not necessarily over a field. In Sect. 3 we define the coniveau filtration for a spectrum with Nisnevich descent and show that the filtration quotients are flabby Nisnevich sheaves. In Sect. 4 the Nisnevich Gersten complex is introduced. Up to this point, the \mathbb{A}^1 -invariance property has not been used yet. Sect. 5 contains an effaceability result which makes use of \mathbb{A}^1 -invariance and the geometric presentation theorem. This leads to our main theorem. The final Sect. 6 deals with the application to étale cohomology with torsion coefficients.

2 Preliminaries

Let S be a base scheme, which is always assumed to be noetherian and of finite dimension. Let moreover Sm_S be the category of smooth schemes of finite type over S and $\text{Spt}_{S^1}(\text{Sm}_S)$ the category of presheaves of spectra on Sm_S . For an object $X \in \text{Sm}_S$ we define the category $\text{Spt}_{S^1}(X_{\text{Nis}})$ analogously where X_{Nis} denotes the small Nisnevich site on X .

The details of the following model structures play no essential role for this text and we refer to the preliminary section of [12], which summarizes material from [1], [5, Sec.2] and [9, 10], for further explanation and references.

We consider the (stable) object-wise model structure on $\text{Spt}_{S^1}(\text{Sm}_S)$. Its homotopy category $\mathcal{SH}_{S^1}(\text{Sm}_S)$ is a triangulated category with exact triangles given by the homotopy (co)fibre sequences. The left Bousfield localization at the equivalences on Nisnevich stalks is called the (stable) Nisnevich local model structure with a fibrant replacement functor L_{Nis} . Likewise, in the case of the small site X_{Nis} , we define the (stable) Nisnevich local model structure on $\text{Spt}_{S^1}(X_{\text{Nis}})$ analogously. In the case of the big site Sm_S , a further left Bousfield localization yields the (stable) \mathbb{A}^1 -Nisnevich local model structure with a fibrant replacement functor L_{mot} . We are working in the non-localized model structure only and use the fibrant replacement functors to obtain statements about the localizations. Hence, whenever we speak of an exact triangle or a homotopy cofibre, we mean the respective terms for the object-wise model structure.

2.1 Basechange

A morphism $f : X \rightarrow Y$ of noetherian schemes of finite Krull dimension induces a covering preserving functor $\bar{f} : \text{Sm}_Y \rightarrow \text{Sm}_X$ by pullback. Precomposition \bar{f}_* with \bar{f} is the right adjoint of a Quillen adjunction

$$\bar{f}^* : \text{Spt}_{S^1}(\text{Sm}_Y) \rightleftarrows \text{Spt}_{S^1}(\text{Sm}_X) : \bar{f}_*$$

for each of the model structures from above (see again the preliminary section of [12] for more details and references). By abuse of notation, we write $f_* := \bar{f}_*$ and $f^* := \bar{f}^*$. If the morphism f is an object of Sm_Y itself, there is an adjunction

$$\underline{f} : \text{Sm}_X \rightleftarrows \text{Sm}_Y : \bar{f}$$

with the left adjoint given by post-composition. Again, precomposition with \underline{f} is the right adjoint of a Quillen adjunction

$$\underline{f}^* : \text{Spt}_{S^1}(\text{Sm}_X) \rightleftarrows \text{Spt}_{S^1}(\text{Sm}_Y) : \underline{f}_*$$

for each of the model structures. We clearly have $\underline{f}_* = \bar{f}^* = f^*$ and set $f_{\sharp} := \underline{f}^*$.

Likewise, for a morphism $f : X \rightarrow Y$, we obtain Quillen adjunctions

$$\text{Spt}_{S^1}(X_{\text{Nis}}) \begin{matrix} \xrightarrow{f_{\sharp}} \\ \rightleftarrows \\ \xleftarrow{f^*} \end{matrix} \text{Spt}_{S^1}(Y_{\text{Nis}}) \begin{matrix} \xrightarrow{f^*} \\ \rightleftarrows \\ \xleftarrow{f_*} \end{matrix} \text{Spt}_{S^1}(X_{\text{Nis}}) \tag{2a}$$

for the object-wise and the Nisnevich local model structure where again for the first one we have to assume that f is an object of Y_{Nis} whereas the second always exists.

For an object $X \rightarrow Y$ in Sm_Y , there is a canonical covering preserving inclusion functor $\delta_{X/Y} : X_{\text{Nis}} \rightarrow \text{Sm}_Y$. Precomposition with this functor yields the right adjoint of a Quillen adjunction

$$\delta_{X/Y}^* : \text{Spt}_{S^1}(X_{\text{Nis}}) \rightleftarrows \text{Spt}_{S^1}(\text{Sm}_Y) : \delta_{X/Y,*} \tag{2b}$$

for the object-wise and the Nisnevich local model structures. The inclusion functor $\delta_{X/Y}$ factorizes as

$$X_{\text{Nis}} \xrightarrow{\delta_X} \text{Sm}_X \xrightarrow{f} \text{Sm}_Y$$

(where we set $\delta_X := \delta_{X/X}$) and the adjunction (2b) factorizes as

$$\text{Spt}_{S^1}(X_{\text{Nis}}) \xrightleftharpoons[\delta_{X,*}]{\delta_X^*} \text{Spt}_{S^1}(\text{Sm}_X) \xrightleftharpoons[f^*]{f_!} \text{Spt}_{S^1}(\text{Sm}_Y).$$

If moreover X is an object of Y_{Nis} , there is a commutative diagram

$$\begin{CD} X_{\text{Nis}} @>\delta_X>> \text{Sm}_X \\ @VfVV @VVfV \\ Y_{\text{Nis}} @>\delta_Y>> \text{Sm}_Y \end{CD} \tag{2c}$$

inducing the diagramm

$$\begin{CD} \text{Spt}_{S^1}(X_{\text{Nis}}) @<\delta_X^*<< \text{Spt}_{S^1}(\text{Sm}_X) \\ @Vf_!VV @VVf_!V \\ \text{Spt}_{S^1}(Y_{\text{Nis}}) @<\delta_Y^*<< \text{Spt}_{S^1}(\text{Sm}_Y) \end{CD}$$

of Quillen adjunctions with diagonal (2b). In particular, for an étale morphism $f : X \rightarrow Y$ the restriction δ_* to the respective small sites commutes with f^* . In particular, these observations imply the following lemma.

Lemma 2.1 *Let $X \in \text{Sm}_S$ and $g : \tilde{X} \rightarrow X$ étale. Then $g^*\delta_{X/S,*} \cong \delta_{\tilde{X}/S,*}$. In particular, the unit of the adjunction $g^* \dashv g_*$ induces a canonical map*

$$\delta_{X/S,*}E \rightarrow g_*\delta_{\tilde{X}/S,*}E.$$

for $E \in \text{Spt}_{S^1}(\text{Sm}_S)$.

Remark 2.2 Let $f : X \rightarrow Y$ be any morphism between noetherian schemes of finite Krull dimension. A diagram analogous to (2c) with \tilde{f} in place of f shows that the restriction δ_* to the respective small sites commutes with f_* .

Definition 2.3 Let $E \in \text{Spt}_{S^1}(\text{Sm}_S)$ or $E \in \text{Spt}_{S^1}(X_{\text{Nis}})$ be a spectrum. For $n \in \mathbb{Z}$, we let $\pi_n(E)$ denote the presheaf $U \mapsto \pi_n(E(U))$ of abelian groups on Sm_S or X_{Nis} . Moreover, one sets $E^n(X) := \pi_{-n}(E(X))$. This common convention lets us synonymously use the term *cohomology theory* for such an object E .

Lemma 2.4 *Let $E \in \text{Spt}_{S^1}(X_{\text{Nis}})$ be a spectrum, $z \in X$ a point and consider the canonical morphism $\mathfrak{z} : \text{Spec}(\mathcal{O}_{X,z}) \rightarrow X$. Then the canonical morphism*

$$\mathfrak{z}_*\mathfrak{z}^*\pi_n(E) \rightarrow \pi_n(\mathfrak{z}_*\mathfrak{z}^*E)$$

is an isomorphism of presheaves on X_{Nis} for every $n \in \mathbb{Z}$.

Proof For any morphism f , we have an isomorphism $f_*\pi_n(E) \cong \pi_n(f_*E)$ as π_n and f_* are defined object-wise. Suppose for a moment that \mathfrak{z} were an object $f : U \rightarrow X$ of the site X_{Nis} .

In this case $f^*(F) \cong F \times_X U$. Since homotopy presheaves and pullbacks of presheaves are calculated object-wise, we obtain

$$f^*\pi_0(E) \cong \pi_0(E) \times_X U \cong \pi_0(E \times_X U) \cong \pi_0(f^*E).$$

We may write $\pi_n(F)$ alternatively as $\pi_0(\underline{\text{Hom}}(S^n, F))$, where $\underline{\text{Hom}}$ denotes the internal mapping space of presheaves. For this internal mapping space, there are isomorphisms

$$f_*(f^*\underline{\text{Hom}}(S^n, E)) \xrightarrow{\cong} f_*(\underline{\text{Hom}}(f^*(S^n), f^*(E))) \xrightarrow{\cong} \underline{\text{Hom}}(S^n, f_*f^*(E))$$

where for the first we used that f was assumed to be in X_{Nis} . Altogether, we have

$$\begin{aligned} f_*f^*\pi_n(E) &\cong f_*f^*\pi_0 \underline{\text{Hom}}(S^n, E) \\ &\cong \pi_0 f_*f^* \underline{\text{Hom}}(S^n, E) \\ &\cong \pi_0 \underline{\text{Hom}}(S^n, f_*f^*(E)) \\ &\cong \pi_n(f_*f^*E) \end{aligned}$$

in the case of f being an object of the site X_{Nis} .

For the case of the essentially open immersion $\mathfrak{z}: \text{Spec}(\mathcal{O}_{X,z}) \rightarrow X$ of the lemma, we write \mathfrak{z} as the cofiltered limit of the diagram $D_{(-)}: \mathcal{I} \rightarrow X_{\text{Nis}}$ given by the affine Zariski neighbourhoods of z in X . Then by the proof of [12, Lemma 1.5] one has a canonical natural isomorphism

$$\mathfrak{z}_*\mathfrak{z}^*(F) \cong \text{colim}_{i \in \mathcal{I}} d_{i,*}d_i^*(F)$$

where $d_i: D_i \rightarrow X$ is the structural morphism which is an open immersion. The result now follows from the case handled above and from the fact that homotopy groups commute with filtered colimits. □

2.2 Codimension

In this subsection, we recall basic notations on the codimension for the convenience of the reader.

Let X be a scheme and $Z \subseteq X$ an irreducible closed subset. Define

$$\text{codim}(Z, X) := \sup\{s \in \mathbb{N}_\infty \mid Z = Z_s \subsetneq \cdots \subsetneq Z_0 \subseteq X \text{ with } Z_i \subseteq X \text{ irred. cl.}\}.$$

For an arbitrary closed subset $Z \subseteq X$ we set

$$\text{codim}(Z, X) := \inf\{\text{codim}(Z', X) \mid Z' \subseteq Z \text{ is an irreducible component}\},$$

where by convention $\text{codim}(\emptyset, X) = \infty$, as the *codimension of Z in X* . One has

$$\text{codim}(Z, X) = \inf_{z \in Z} \dim(\mathcal{O}_{X,z}).$$

If Z is irreducible closed with generic point η_Z , then $\text{codim}(Z, X) = \dim(\mathcal{O}_{X,\eta_Z})$. Recall that a scheme X is called *catenary*, if for every two irreducible closed subsets $Z \subseteq Z' \subseteq X$ every maximal chain $Z = Z_s \subsetneq \cdots \subsetneq Z_0 = Z'$ of irreducible closed subsets has the same finite length. Examples of such are schemes (locally) of finite type over a field or over a one dimensional noetherian domain, e.g., a discrete valuation ring.

We have the following three easy lemmas.

Lemma 2.5 *Let X be a scheme and $Z \subseteq Z' \subseteq X$ two irreducible closed subsets with $\text{codim}(Z', X) = s$. Then*

$$Z \neq Z' \Leftrightarrow \text{codim}(Z, X) \geq s + 1.$$

Lemma 2.6 *Let X be a scheme and $Z_1, Z_2 \subseteq X$ two closed subsets with both $\text{codim}(Z_1, X) \geq s$ and $\text{codim}(Z_2, X) \geq s$. Then $\text{codim}(Z_1 \cup Z_2, X) \geq s$.*

Lemma 2.7 *Let X be an irreducible catenary scheme and $Z \subseteq Z' \subseteq X$ two irreducible closed subsets. Then*

$$\text{codim}(Z, X) = \text{codim}(Z, Z') + \text{codim}(Z', X).$$

For an integer $s \geq 0$, define

$$X^{(s)} = \{z \in X \mid \text{codim}(\overline{\{z\}}, X) = s\}$$

and say that z has codimension s in X if $z \in X^{(s)}$. Note, that $\overline{\{z\}}$ is always an irreducible closed subset of X and z is its generic point. One has $X^{(s)} = \emptyset$ for $s > \dim(X)$ as $\dim(X) = \sup_{x \in X} \dim(\mathcal{O}_{X,x})$. We have $\text{codim}(Z, X) = 0$ if and only if Z contains a whole irreducible component of X . Hence a point $z \in X$ is a generic point of an irreducible component of X if and only if $\dim(\mathcal{O}_{X,z}) = 0$. Thus, $X^{(0)}$ is precisely the set of generic points of irreducible components of X .

Remark 2.8 Please note that the inequality

$$\dim(Z) + \text{codim}(Z, X) \leq \dim(X)$$

is not always an equality, even for catenary schemes.

3 The coniveau filtration

3.1 In this section we define the coniveau filtration for a Nisnevich local fibrant spectrum on Sm_S . We fix a spectrum $E \in \text{Spt}_{S^1}(\text{Sm}_S)$.

Definition 3.2 Let $X/S \in \text{Sm}_S, Z \subseteq X$ a closed subset with complementary open immersion $j: (X \setminus Z) \hookrightarrow X$. Let E_X be the spectrum $\delta_{X/S,*}E$ in $\text{Spt}_{S^1}(X_{\text{Nis}})$. Lemma 2.1 induces a morphism

$$\eta_j: E_X \rightarrow j_*j^*E_X = j_*E_{X \setminus Z} \tag{3a}$$

in $\text{Spt}_{S^1}(X_{\text{Nis}})$ whose homotopy fibre, we denote by $E_{Z/X} \in \text{Spt}_{S^1}(X_{\text{Nis}})$.

Remark 3.3 Note that $E_{X/X} = E_X$ and $E_{\emptyset/X} = *$ in $\text{Spt}_{S^1}(X_{\text{Nis}})$.

Remark 3.4 Throughout Sects. 3 and 4, we could as well work with arbitrary spectra $E_X \in \text{Spt}_{S^1}(X_{\text{Nis}})$, not necessarily of the form $\delta_{X/S,*}E$ for a spectrum $E \in \text{Spt}_{S^1}(\text{Sm}_S)$. In this case, $E_{\tilde{X}}$ will be defined as f^*E_X for an étale morphism $f: \tilde{X} \rightarrow X$ of finite type.

3.5 We fix an étale morphism $f: \tilde{X} \rightarrow X$ of finite type and a closed subset $Z \subseteq X$. Let $\tilde{Z} = Z \times_X \tilde{X}$ denote the pullback of Z along f . We get an induced morphism $\tilde{f}: (\tilde{X} \setminus \tilde{Z}) \rightarrow (X \setminus Z)$ on the open complements. By Lemma 2.1, $f^*E_X \cong E_{\tilde{X}}$ and we have a canonical morphism $E_X \rightarrow f_*E_{\tilde{X}}$.

Lemma 3.6 *In the situation of 3.5, we have $f^*E_{Z/X} \simeq E_{\tilde{Z}/\tilde{X}}$ in $\text{Spt}_{S^1}(\tilde{X}_{\text{Nis}})$.*

Proof Let $j : (X \setminus Z) \hookrightarrow X$ and $\tilde{j} : (\tilde{X} \setminus \tilde{Z}) \hookrightarrow \tilde{X}$. First, we apply the homotopy exact functor f^* to the homotopy fibre sequence $E_{Z/X} \rightarrow E_X \rightarrow j_*j^*E_X$. Base change (since $f \in X_{\text{Nis}}$) for the pullback square

$$\begin{array}{ccc} \tilde{X} \setminus \tilde{Z} & \xrightarrow{\tilde{j}} & \tilde{X} \\ \tilde{f} \downarrow & & \downarrow f \\ X \setminus Z & \xrightarrow{j} & X \end{array}$$

yields an equivalence $f^*j_* \simeq \tilde{j}_*\tilde{f}^*$ and it remains to observe that $\tilde{f}^*j^* \simeq (j\tilde{f})^* \simeq (f\tilde{j})^* \simeq \tilde{j}^*f^*$, which is easy. □

Lemma 3.7 (Forget support map) *Let Z and Z' be closed subsets of X with $Z \subseteq Z' \subseteq X$ and let $E_X \in \text{Spt}_{S^1}(X_{\text{Nis}})$ be a spectrum. Then there is a canonical forget support map $E_{Z/X} \rightarrow E_{Z'/X}$. Further, this map sits in a canonical exact triangle*

$$E_{Z/X} \rightarrow E_{Z'/X} \rightarrow j_*E_{(Z' \setminus Z)/(X \setminus Z)}$$

of objects from $\text{Spt}_{S^1}(X_{\text{Nis}})$, where $j : (X \setminus Z) \hookrightarrow X$.

Proof Note first that $j' : (X \setminus Z') \hookrightarrow X$ factorizes as

$$X \setminus Z' = (X \setminus Z) \setminus (Z' \setminus Z) \xrightarrow{k} X \setminus Z \xrightarrow{j} X,$$

and therefore the unit $\text{id} \rightarrow k_*k^*$ of the adjunction $k^* \dashv k_*$ induces a morphism $j_*j'^* \rightarrow j_*k_*k^*j'^* \simeq j'_*j'^*$. This map is compatible with the units of the adjunctions $j'^* \dashv j'_*$ and $j^* \dashv j_*$, thus inducing the forget support map $E_{Z/X} \rightarrow E_{Z'/X}$ on homotopy fibres.

For the exact triangle consider the diagram

$$\begin{array}{ccccc} F[-1] & \longrightarrow & * & \longrightarrow & F \\ \downarrow & & \downarrow & & \downarrow \\ E_{Z/X} & \longrightarrow & E_X & \longrightarrow & j_*j^*E_X \\ \text{forget support} \downarrow & & \downarrow \text{id} & & \downarrow \\ E_{Z'/X} & \longrightarrow & E_X & \longrightarrow & j'_*j'^*E_X \end{array}$$

of distinguished triangles. It remains to identify F with $j_*E_{(Z' \setminus Z)/(X \setminus Z)}$. Applying the exact functor j_* to the exact triangle

$$E_{(Z' \setminus Z)/(X \setminus Z)} \rightarrow E_{(X \setminus Z)} \rightarrow k_*k^*E_{(X \setminus Z)},$$

yields the right vertical exact triangle

$$F = j_*E_{(Z' \setminus Z)/(X \setminus Z)} \rightarrow j_*j^*E_X \rightarrow j_*k_*k^*j^*E_X,$$

where we used the definition $j'^*E_X \simeq E_{(X \setminus Z)}$. □

Remark 3.8 In particular, the previous Lemma 3.7 induces a long exact sequence

$$\dots \rightarrow E_Z^n(X) \rightarrow E_{Z'}^n(X) \rightarrow E_{Z' \setminus Z}^n(X \setminus Z) \rightarrow E_Z^{n+1}(X) \rightarrow \dots$$

3.9 Recall that a Nisnevich distinguished square

$$\begin{array}{ccc} \tilde{U} & \longrightarrow & \tilde{X} \\ \downarrow & & \downarrow f \\ U & \xrightarrow{j} & X \end{array}$$

is a pullback square such that j is an open immersion, f is an étale morphism of finite type and $(X - j(U))_{\text{red}} \times_X \tilde{X} \rightarrow (X - j(U))_{\text{red}}$ is an isomorphism.

3.10 Recall moreover, that an object-wise fibrant spectrum $E \in \text{Spt}_{S^1}(\text{Sm}_S)$ (i.e., every evaluation $E(X)$ is an ordinary Ω -spectrum) is *Nisnevich local fibrant* if and only if $E(\emptyset) = *$ and for each Nisnevich distinguished square Q as in 3.9, the square $E(Q)$ is a homotopy pullback square (or equivalently a homotopy pushout square). Equivalently, the sequence

$$E(X) \rightarrow E(\tilde{X}) \oplus E(U) \rightarrow E(\tilde{U})$$

is a distinguished triangle and hence induces long exact sequences on homotopy groups. The same observation holds for the Nisnevich local fibrant objects of $\text{Spt}_{S^1}(X_{\text{Nis}})$ and the right adjoint $\delta_{X/S,*} : \text{Spt}_{S^1}(\text{Sm}_S) \rightarrow \text{Spt}_{S^1}(X_{\text{Nis}})$ of the adjunction (2b) preserves Nisnevich local fibrant objects.

Lemma 3.11 *An object-wise fibrant spectrum $E \in \text{Spt}_{S^1}(\text{Sm}_S)$ is Nisnevich local fibrant if and only if for all Nisnevich distinguished squares as in 3.9, the induced morphism*

$$E_{Z/X} \rightarrow f_* f^* E_{Z/X} \simeq f_* E_{\tilde{Z}/\tilde{X}}$$

(see Lemma 3.6) is an equivalence. Here, $Z := X \setminus U$ and $\tilde{Z} := (\tilde{X} \setminus \tilde{U}) \cong f^{-1}(Z)$.

Proof This follows immediately from the fact the a square of spectra is a homotopy pullback square if and only if the homotopy fibres of the horizontal morphisms are equivalent. \square

Lemma 3.12 *Let $E \in \text{Spt}_{S^1}(\text{Sm}_S)$ be a Nisnevich local fibrant spectrum. Let Z_1 and Z_2 be closed subsets of X , and set $Z_{12} = Z_1 \cap Z_2$ and $Z = Z_1 \cup Z_2$. Then the forget support maps*

$$\begin{array}{ccc} E_{Z_{12}/X} & \longrightarrow & E_{Z_2/X} \\ \downarrow & & \downarrow \\ E_{Z_1/X} & \longrightarrow & E_{Z/X} \end{array} \tag{3b}$$

form a homotopy (co)fibre square in $E \in \text{Spt}_{S^1}(X_{\text{Nis}})$.

Proof First, observe that the respective open immersions form a Nisnevich distinguished square

$$\begin{array}{ccc} X \setminus Z & \longrightarrow & X \setminus Z_2 \\ \downarrow & & \downarrow \\ X \setminus Z_1 & \longrightarrow & X \setminus Z_{12} \end{array}$$

Denote by $j : X \setminus Z \hookrightarrow X$ and $j_i : X \setminus Z_i \hookrightarrow X$ for $i = 1, i = 2$ or $i = 12$ the complementary open immersions. Since E is Nisnevich local fibrant, it follows that

$$\begin{array}{ccc} j_{12,*} j_{12}^* E_X & \longrightarrow & j_{2,*} j_2^* E_X \\ \downarrow & & \downarrow \\ j_{1,*} j_1^* E_X & \longrightarrow & j_* j^* E_X \end{array}$$

is a homotopy pullback square. Mapping into this square from the square with edges id_{E_X} , which is a homotopy pullback square for trivial reasons, and taking homotopy fibres, yields the square (3b). Thus, (3b) is a homotopy pullback square, too. \square

Corollary 3.13 *Let $E \in \text{Spt}_{S^1}(\text{Sm}_S)$ be a Nisnevich local fibrant spectrum and $Z_1, \dots, Z_r \subseteq X$ be disjoint closed subsets. Then*

$$\bigoplus_{i=1}^r E_{Z_i/X} \simeq E_{(\bigsqcup_{i=1}^r Z_i/X)}.$$

Definition 3.14 Let $E_X \in \text{Spt}_{S^1}(X_{\text{Nis}})$ be a spectrum. For an integer $s \geq 0$, we define the spectrum

$$E_{X^{(s)}} := \text{colim}_{\substack{Z \subseteq X \text{ closed} \\ \text{codim}(Z, X) \geq s}} E_{Z/X}$$

in $\text{Spt}_{S^1}(X_{\text{Nis}})$. The structure maps for the colimit are the forget support maps (see Lemma 3.7).

Remark 3.15 Informally, one should think of the colimit in the previous Definition 3.14 as “making the Z ’s bigger”. The index category is filtered as one can take the union of two closed sets.

3.16 Since a closed subset of X of codimension $\geq (s + 1)$ is in particular a closed subset of codimension $\geq s$, we get a filtration

$$* \rightarrow E_{X^{(d)}} \rightarrow \dots \rightarrow E_{X^{(s+1)}} \rightarrow E_{X^{(s)}} \rightarrow \dots \rightarrow E_{X^{(0)}} \cong E_X$$

of presheaves of spectra on X_{Nis} . For the last equivalence, observe that the colimit in Definition 3.14 has a terminal object $Z = X$ in the case $s = 0$.

Definition 3.17 We denote the homotopy cofibre of $E_{X^{(s+1)}} \rightarrow E_{X^{(s)}}$ by $E_{X^{(s/s+1)}}$.

3.18 As usual, one can associate a spectral sequence (more precisely, a presheaf on X_{Nis} of spectral sequences) to such a situation: Applying π_n for an integer n to the filtration of 3.16 yields a finite filtration

$$0 \subseteq \text{im } \pi_n(E_{X^{(d)}} \rightarrow E_X) \subseteq \dots \subseteq \text{im } \pi_n(E_{X^{(1)}} \rightarrow E_X) \subseteq \pi_n(E_X)$$

and the associated spectral sequence (in the category of presheaves of abelian groups on X_{Nis})

$$E_{p,q}^1 = \pi_{p+q}(E_{X^{(p/p+1)}}) \Rightarrow \pi_{p+q}(E_X)$$

is degenerate (and hence always converges in the strongest sense) as the filtration above is bounded. Reindexing and rephrasing along Definition 2.3, we get

$$E_1^{p,q} = E_{X^{(p/p+1)}}^{p+q} \Rightarrow E^{p+q}.$$

The constructed spectral sequence is not yet the coniveau spectral sequence. To obtain the latter, we will sheafify the whole situation (after taking homotopy groups as above) and identify the homotopy cofibres $E_{X^{(p/p+1)}}$ with certain coproducts.

3.19 For a point $z \in X$ we denote by \mathfrak{z} in the following the canonical morphism $\mathfrak{z}: \text{Spec}(\mathcal{O}_{X,z}) \rightarrow X$.

Proposition 3.20 *Let $E \in \text{Spt}_{S^1}(\text{Sm}_S)$ be a Nisnevich local fibrant spectrum. Then for every integer $s \geq 0$, we have an equivalence*

$$E_{X^{(s/s+1)}} \simeq \bigoplus_{z \in X^{(s)}} \mathfrak{J}^* \mathfrak{J}^* E_{Z/X},$$

where $\mathfrak{J}^* : \text{Spt}_{S^1}(X_{\text{Nis}}) \rightleftarrows \text{Spt}_{S^1}(\text{Spec}(\mathcal{O}_{X,z})_{\text{Nis}}) : \mathfrak{J}_*$ is the adjunction (2a) and where we set $Z := \{z\}$ in each summand by abuse of notation.

Proof For closed subsets Z and Z' of X with $Z \subseteq Z'$, Lemma 3.7 yields an exact triangle

$$E_{Z/X} \rightarrow E_{Z'/X} \rightarrow j_* E_{(Z' \setminus Z)/(X \setminus Z)}.$$

Taking filtered colimits yields an exact triangle

$$E_{X^{(s+1)}} \rightarrow E_{X^{(s)}} \rightarrow \underset{\substack{Z, Z' \subseteq X \text{ closed} \\ Z \subseteq Z' \\ \text{codim}(Z', X) \geq s \\ \text{codim}(Z, X) \geq s+1}}{\text{colim}} j_* E_{(Z' \setminus Z)/(X \setminus Z)} \tag{3c}$$

of objects from $\text{Spt}_{S^1}(X_{\text{Nis}})$. In particular, the right-hand side is equivalent to $E_{X^{(s/s+1)}}$. The colimit in (3c) runs over the filtered category of pairs (Z, Z') where $Z \subseteq Z'$ for $Z, Z' \subseteq X$ closed subsets of the indicated codimensions with an arrow $(Z, Z') \rightarrow (\hat{Z}, \hat{Z}')$ if and only if both $Z \subseteq \hat{Z}$ and $Z' \subseteq \hat{Z}'$.

We will now rewrite this colimit. Fix a pair (Z, Z') . Since X is noetherian, Z' is noetherian as a topological space and hence the union of its finite number of irreducible components Z'_1, \dots, Z'_r , each of codimension $\geq s$. It follows, that all the intersections $Z'_i \cap Z'_j$ for $i \neq j$ are of codimension $\geq s + 1$ by Lemma 2.5. Set

$$\hat{Z} := Z \cup \bigcup_{i \neq j} (Z'_i \cap Z'_j) \cup \bigcup_{\substack{i \text{ such that} \\ \text{codim}(Z'_i, X) \geq s+1}} Z'_i.$$

By Lemma 2.6, \hat{Z} has codimension $\geq s + 1$ and $(\hat{Z}, \hat{Z}' := Z')$ receives a map from our original pair (Z, Z') . Let $T \subseteq X^{(s)}$ be the set of generic points of those Z'_i of codimension s . Then $\overline{T} \cup \hat{Z} = \hat{Z}'$. Further, $\hat{U} = X \setminus \hat{Z} \subseteq X$ is an open *separating* neighbourhood of T . By this we mean that $\overline{T} \cap \hat{U}$ splits into a disjoint union of closures of points of T in \hat{U} . Combining these observations, we get a cofinal functor from the category of pairs (T, U) with $T \subseteq X^{(s)}$ a finite subset and $U \subseteq X$ an open separating neighbourhood of T into our original index category by mapping a pair (T, U) to the pair $(X \setminus U, \overline{T} \cup (X \setminus U))$. In particular,

$$\underset{\substack{Z, Z' \subseteq X \text{ closed} \\ Z \subseteq Z' \\ \text{codim}(Z', X) \geq s \\ \text{codim}(Z, X) \geq s+1}}{\text{colim}} j_* E_{(Z' \setminus Z)/(X \setminus Z)} \simeq \underset{\substack{T \subseteq X^{(s)} \text{ finite} \\ T \subseteq U \subseteq X \text{ open sep nbh}}}{\text{colim}} j_* E_{(\overline{T} \cap U)/U}.$$

As U is a separating neighbourhood of T , Corollary 3.13 gives a splitting

$$E_{(\overline{T} \cap U)/U} \simeq \bigoplus_{z \in T} E_{(\overline{\{z\}} \cap U)/U}.$$

Note that the open separating neighbourhoods of T are cofinal in all open neighbourhoods of T . In particular, we get

$$\begin{aligned} \operatorname{colim}_{\substack{T \subseteq X^{(s)} \text{ finite} \\ T \subseteq U \subseteq X \text{ open sep nbh}}} j_* E_{(\overline{T} \cap U)/U} &\simeq \operatorname{colim}_{\substack{T \subseteq X^{(s)} \text{ finite} \\ T \subseteq U \subseteq X \text{ open nbh}}} \bigoplus_{z \in T} j_* E_{(\overline{\{z\}} \cap U)/U} \\ &\simeq \bigoplus_{z \in X^{(s)}} \operatorname{colim}_{z \in U \subseteq X \text{ open nbh}} j_* E_{(\overline{\{z\}} \cap U)/U} \end{aligned}$$

Finally, by Lemma 3.6, $E_{(\overline{\{z\}} \cap U)/U} \simeq j^* E_{\overline{\{z\}}/X}$ and the claim follows. □

Corollary 3.21 *Let $E \in \operatorname{Spt}_{S^1}(\operatorname{Sm}_S)$ be a Nisnevich local fibrant spectrum. Then for every integer $s \geq 0$ and every integer q , we have an isomorphism*

$$E_{X^{(s/s+1)}}^q \cong \bigoplus_{z \in X^{(s)}} \mathfrak{z} * \mathfrak{z}^* E_{Z/X}^q.$$

Proof. By Proposition 3.20 $E_{X^{(s/s+1)}} \simeq \bigoplus_{z \in X^{(s)}} \mathfrak{z} * \mathfrak{z}^* E_{Z/X}$. Using Lemma 2.4 we compute

$$\begin{aligned} \pi_{-q} \left(\bigoplus_{z \in X^{(s)}} \mathfrak{z} * \mathfrak{z}^* E_{Z/X} \right) &\cong \bigoplus_{z \in X^{(s)}} \pi_{-q} (\mathfrak{z} * \mathfrak{z}^* E_{Z/X}) \\ &\cong \bigoplus_{z \in X^{(s)}} \mathfrak{z} * \mathfrak{z}^* \pi_{-q} (E_{Z/X}). \end{aligned} \quad \square$$

3.22 Recall that a sheaf F of abelian groups on the site X_{Nis} is called *flabby*, if the presheaf $H^q(-, F)$ on X_{Nis} is zero for $q \neq 0$. A flabby sheaf is in particular acyclic, i.e., $H^q(X, F) = 0$ for $q \neq 0$.

Proposition 3.23 *Let $E \in \operatorname{Spt}_{S^1}(\operatorname{Sm}_S)$ be a Nisnevich local fibrant spectrum, $z \in X^{(s)}$ with $Z := \overline{\{z\}}$ and q an integer. Then the presheaf*

$$\mathfrak{z} * \mathfrak{z}^* E_{Z/X}^q$$

of abelian groups is a flabby sheaf on X_{Nis} .

Proof Let q be an integer. Let $V \rightarrow X$ be étale of finite type with (set-theoretical) fibre $V(z)$ over the point $z \in X$. For a point $v \in V(z)$, we set $V_v^{\text{loc}} := \operatorname{Spec}(\mathcal{O}_{V,v})$ and $V_z^{\text{loc}} := V \times_X X_z^{\text{loc}}$. Using the identification $Z \times_X V_z^{\text{loc}} = \coprod_{v \in V(z)} v$, we have

$$\begin{aligned} \mathfrak{z} * \mathfrak{z}^* E_{Z/X}^q(V) &\cong \pi_{-q} (\mathfrak{z} * \mathfrak{z}^* E_{Z/X}(V)) && \text{(by Lemma 2.4)} \\ &\cong \pi_{-q} \left(E_{Z \times_X V_z^{\text{loc}}/V_z^{\text{loc}}}(V_z^{\text{loc}}) \right) && \text{(by Lemma 3.6)} \\ &\cong \pi_{-q} \left(E_{(\coprod_{v \in V(z)} v)/V_z^{\text{loc}}}(V_z^{\text{loc}}) \right) \\ &\cong \pi_{-q} \left(\bigoplus_{v \in V(z)} E_{v/V_z^{\text{loc}}}(V_z^{\text{loc}}) \right) && \text{(by Corollary 3.13)} \\ &\cong \bigoplus_{v \in V(z)} E_{v/V_v^{\text{loc}}}^q(V_v^{\text{loc}}). \end{aligned}$$

Let us now prove the sheaf property of $\mathfrak{z} * \mathfrak{z}^* E_{Z/X}^q$. Using Lemma 3.6, we may restrict us to Nisnevich covers $V \rightarrow X$ of X . Writing $W := V \times_X V$, we have to show that

$$\begin{array}{ccccc} 0 \rightarrow \mathfrak{z} * \mathfrak{z}^* E_{Z/X}^q(X) & \longrightarrow & \mathfrak{z} * \mathfrak{z}^* E_{Z/X}^q(V) & \xrightarrow{\operatorname{pr}_1^* - \operatorname{pr}_2^*} & \mathfrak{z} * \mathfrak{z}^* E_{Z/X}^q(W) \\ \parallel & & \parallel & & \parallel \\ E_{z/X_z^{\text{loc}}}^q(X_z^{\text{loc}}) & \rightarrow & \bigoplus_{v \in V(z)} E_{v/V_v^{\text{loc}}}^q(V_v^{\text{loc}}) & \rightarrow & \bigoplus_{w \in W(z)} E_{w/W_w^{\text{loc}}}^q(W_w^{\text{loc}}) \end{array}$$

is an exact sequence. Since $V \rightarrow X$ is a Nisnevich cover, we can find a point v_0 in the fibre $V(z)$ with residue field $k(v_0) = k(z)$. In particular, by Lemma 3.11, the composition

$$E_{z/X_z^{\text{loc}}}^q(X_z^{\text{loc}}) \longrightarrow \bigoplus_{v \in V(z)} E_{v/V_v^{\text{loc}}}^q(V_v^{\text{loc}}) \xrightarrow{\text{can}} E_{v_0/V_{v_0}^{\text{loc}}}^q(V_{v_0}^{\text{loc}})$$

is an isomorphism, which settles the exactness at $\mathfrak{z}^* \mathfrak{z}^* E_{Z/X}^q(X)$.

For the exactness at $\mathfrak{z}^* \mathfrak{z}^* E_{Z/X}^q(V)$, observe that $\bigoplus_{v \in V(z)} E_{v/V_v^{\text{loc}}}^q(V_v^{\text{loc}})$ splits into a direct sum of $E_{z/X_z^{\text{loc}}}^q(X_z^{\text{loc}})$ and $\bigoplus_{v_0 \neq v \in V(z)} E_{v/V_v^{\text{loc}}}^q(V_v^{\text{loc}})$. Hence, it is enough to show that the restricted map $\bigoplus_{v_0 \neq v \in V(z)} E_{v/V_v^{\text{loc}}}^q(V_v^{\text{loc}}) \rightarrow \bigoplus_{w \in W(z)} E_{w/W_w^{\text{loc}}}^q(W_w^{\text{loc}})$ is a monomorphism. To this end, it suffices to show that

$$\text{pr}_1^* - \text{pr}_2^*: E_{v/V_v^{\text{loc}}}^q(V_v^{\text{loc}}) \longrightarrow E_{v \otimes v_0/W_{v \otimes v_0}^{\text{loc}}}(W_{v \otimes v_0}^{\text{loc}}) \oplus E_{v_0 \otimes v/W_{v_0 \otimes v}^{\text{loc}}}(W_{v_0 \otimes v}^{\text{loc}})$$

is a monomorphism for each v different from v_0 . But even the projection

$$\text{pr}_1^*: E_{v/V_v^{\text{loc}}}^q(V_v^{\text{loc}}) \longrightarrow E_{v \otimes v_0/W_{v \otimes v_0}^{\text{loc}}}(W_{v \otimes v_0}^{\text{loc}})$$

is an isomorphism by Lemma 3.11: Indeed, the equality $k(v \otimes v_0) = k(v)$ follows from $k(v_0) = k(z)$, so $\text{pr}_1: (W_{v \otimes v_0}^{\text{loc}}, v \otimes v_0) \rightarrow (V_v^{\text{loc}}, v)$ is (essentially) a Nisnevich neighbourhood. This finishes the proof of the sheaf property of $\mathfrak{z}^* \mathfrak{z}^* E_{Z/X}^q$.

In order to show the flabbiness, let us first show that $\mathfrak{z}^* E_{Z/X}^q$ is flabby: Again, we have $\mathfrak{z}^* E_{Z/X}^q = E_{z/X_z^{\text{loc}}}^q$ by Lemma 3.6. Let $j: U \hookrightarrow X_z^{\text{loc}}$ be the open complement of the closed point $z \in X_z^{\text{loc}}$. Then $j^* E_{z/X_z^{\text{loc}}}^q$ is trivial by construction. Hence, $E_{z/X_z^{\text{loc}}}^q$ is supported on z , i.e., $E_{z/X_z^{\text{loc}}}^q = z_* z^* E_{z/X_z^{\text{loc}}}^q$ is flabby as a skyscraper-sheaf. For the flabbiness of $\mathfrak{z}^* \mathfrak{z}^* E_{Z/X}^q$, we have to show that $H^i(V_{\text{Nis}}, \mathfrak{z}^* \mathfrak{z}^* E_{Z/X}^q)$ is trivial for all $V \rightarrow X$ étale of finite type and $i > 0$. Since $\mathfrak{z}^* E_{Z/X}^q$ is flabby, it is $\mathbb{R}\mathfrak{z}^*$ -acyclic, i.e., $\mathfrak{z}^* \mathfrak{z}^* E_{Z/X}^q \simeq \mathbb{R}\mathfrak{z}^* \mathfrak{z}^* E_{Z/X}^q$. In particular, we have

$$H^i(V_{\text{Nis}}, \mathfrak{z}^* \mathfrak{z}^* E_{Z/X}^q) \cong H^i(V_{\text{Nis}}, \mathbb{R}\mathfrak{z}^* \mathfrak{z}^* E_{Z/X}^q) \cong H^i(V_{z, \text{Nis}}^{\text{loc}}, \mathfrak{z}^* E_{Z/X}^q),$$

but the latter group is trivial since $V_{z, \text{Nis}}^{\text{loc}} \rightarrow X_z^{\text{loc}}$ is étale and $\mathfrak{z}^* E_{Z/X}^q$ is flabby. □

Corollary 3.24 *Let $E \in \text{Spt}_{S^1}(\text{Sm}_S)$ be a Nisnevich local fibrant spectrum. Then for every integer $s \geq 0$ and every integer q , the presheaf $E_{X^{(s/s+1)}}^q$ is a flabby sheaf on X_{Nis} .*

Proof By Corollary 3.21, we have $E_{X^{(s/s+1)}}^q \simeq \bigoplus_{z \in X^{(s)}} \mathfrak{z}^* \mathfrak{z}^* E_{Z/X}^q$. Here the direct sum is the direct sum of presheaves. But X_{Nis} is noetherian (see e.g. [13, Proposition 5.2]), so the direct sum of sheaves is the direct sum of presheaves and the claim follows from Proposition 3.23. □

4 The Nisnevich Gersten complex

Let us remark that the Zariski-analogues of the following Lemma 4.1 and Proposition 4.6 were considered initially by Quillen for the special case of algebraic K -Theory in [11, §7. Prop 5.6].

Lemma 4.1 *Let $E \in \text{Spt}_{S^1}(\text{Sm}_S)$ be a spectrum and n an integer. The cofibre sequences of Definition 3.17 yield a complex of presheaves on X_{Nis} of abelian groups*

$$0 \rightarrow E_X^n \xrightarrow{e} E_{X^{(0/1)}}^n \xrightarrow{d^0} E_{X^{(1/2)}}^{n+1} \xrightarrow{d^1} \dots \xrightarrow{d^{d-2}} E_{X^{(d-1/d)}}^{n+d-1} \xrightarrow{d^{d-1}} E_{X^{(d)}}^{n+d} \rightarrow 0.$$

The (Nisnevich) sheafification $(-)^{\sim}$ of this complex is exact at the first spot $(E_X^n)^{\sim}$ if and only if the canonical map


$$(\alpha_0^n)^{\sim} : (E_{X^{(1)}}^n)^{\sim} \rightarrow (E_{X^{(0)}}^n)^{\sim} \tag{4a}$$

is zero and it is exact at the spot $(E_{X^{(s/s+1)}}^{n+s})^{\sim}$ for $s \geq 0$ if both the canonical maps

$$\begin{aligned} (\alpha_{s-1}^{n+s})^{\sim} &: (E_{X^{(s)}}^{n+s})^{\sim} \rightarrow (E_{X^{(s-1)}}^{n+s})^{\sim} \text{ and} \\ (\alpha_{s+1}^{n+s+1})^{\sim} &: (E_{X^{(s+2)}}^{n+s+1})^{\sim} \rightarrow (E_{X^{(s+1)}}^{n+s+1})^{\sim} \end{aligned} \tag{4b}$$

are zero (where the first condition is empty for $s = 0$).

Proof The long exact sequences on homotopy groups associated to the cofibre sequences $E_{X^{(s+1)}} \rightarrow E_{X^{(s)}} \rightarrow E_{X^{(s/s+1)}}$ from Definition 3.17 for $s \geq 0$ yield a diagram



and we define the middle horizontal sequence as indicated. This sequence is clearly a complex as the diagonal lines are complexes. The remaining statement follows immediately from sheafification $(-)^{\sim}$ applied to the whole diagram. \square

4.2 For a Nisnevich local fibrant spectrum $E \in \text{Spt}_{S^1}(\text{Sm}_S)$ we can rewrite the complex of presheaves on X_{Nis} of abelian groups from the previous Lemma 4.1 with the help of Corollary 3.21 as

$$\begin{aligned} 0 \rightarrow E_X^n \xrightarrow{e} \bigoplus_{z \in X^{(0)}} \mathfrak{z} * \mathfrak{z}^* E_{Z/X}^n \xrightarrow{d^0} \bigoplus_{z \in X^{(1)}} \mathfrak{z} * \mathfrak{z}^* E_{Z/X}^{n+1} \xrightarrow{d^1} \dots \\ \dots \xrightarrow{d^{d-2}} \bigoplus_{z \in X^{(d-1)}} \mathfrak{z} * \mathfrak{z}^* E_{Z/X}^{n+d-1} \xrightarrow{d^{d-1}} \bigoplus_{z \in X^{(d)}} \mathfrak{z} * \mathfrak{z}^* E_{Z/X}^{n+d} \rightarrow 0. \end{aligned} \tag{4c}$$

Definition 4.3 For every integer n , we define the Nisnevich Gersten complex $\mathcal{G}^\bullet(E, n)$ of E and homotopical degree n as the complex with entries

$$\mathcal{G}^s(E, n) := \bigoplus_{z \in X^{(s)}} \mathfrak{z} * \mathfrak{z}^* E_{Z/X}^{n+s}$$

for $s \geq 0$ and zero otherwise. The differentials d^s are defined as in 4.2.

4.4 For every integer n , we can reformulate (4c) as a map

$$E_X^n \xrightarrow{e} \mathcal{G}^\bullet(E, n)$$

into a complex of flabby sheaves (see Corollary 3.24) of abelian groups.

4.5 In abuse of notation, we will just write $E(X_x^h)$ for the stalk of E at a point x of X . Here of course X_x^h denotes the Henselian local scheme $\text{Spec}(\mathcal{O}_{X,x}^h)$.

Proposition 4.6 *Let $E \in \text{Spt}_{S^1}(\text{Sm}_S)$ be a Nisnevich local fibrant spectrum and n an integer. There is a complex of sheaves on X_{Nis} of abelian groups*

$$0 \rightarrow (E_X^n)^\sim \xrightarrow{\tilde{e}} \mathcal{G}^0(E, n) \xrightarrow{d^0} \mathcal{G}^1(E, n) \xrightarrow{d^1} \dots \xrightarrow{d^{d-1}} \mathcal{G}^d(E, n) \rightarrow 0$$

where all but the first entry are flabby Nisnevich sheaves. This complex is

- (1) exact at the first spot $(E_X^n)^\sim$ if and only if, for each point x of X and all $Z \subseteq X$ closed with $\text{codim}(Z, X) \geq 1$, the forget support map

$$E_{Z/X}^n(X_x^h) \rightarrow E_X^n(X_x^h)$$

is trivial and

- (2) exact at $\mathcal{G}^s(E, n)$ for $s \geq 0$ if, Nisnevich-locally on X ,

- (i) if $s = 1$, for each point x of X and all $Z \subseteq X$ closed with $\text{codim}(Z, X) \geq 1$, the forget support map

$$E_{Z/X}^{n+1}(X_x^h) \rightarrow E_X^{n+1}(X_x^h)$$

is trivial and

- (ii) if $s > 1$, for all $Z \subseteq X$ closed with $\text{codim}(Z, X) \geq s$, there exists $Z' \subseteq Z \subseteq X$ closed with $\text{codim}(Z', X) \geq s - 1$ such that the forget support map

$$E_{Z/X}^{n+s}(X) \rightarrow E_{Z'/X}^{n+s}(X)$$

is trivial and

- (iii) for all $s \geq 0$ and all $Z \subseteq X$ closed with $\text{codim}(Z, X) \geq s + 2$, there exists $Z \subseteq Z' \subseteq X$ closed with $\text{codim}(Z', X) \geq s + 1$ such that the forget support map

$$E_{Z/X}^{n+s+1}(X) \rightarrow E_{Z'/X}^{n+s+1}(X)$$

is trivial.

Proof The complex is obtained by applying the sheafification functor to the complex (4c). By Corollary 3.24, all but the first entry are flabby Nisnevich sheaves. The exactness conditions are just expanded versions of (4a) and (4b). □

5 Effaceability

5.1 In this chapter, let S be the spectrum of a Henselian discrete valuation ring \mathfrak{o} with infinite residue field \mathbb{F} of characteristic p and quotient field k . Note that we make use of this hypothesis only from Lemma 5.8 onwards.

Construction 5.2 Let $E \in \text{Spt}_{S^1}(\text{Sm}_S)$ be a spectrum and $f : \tilde{X} \rightarrow X$ a morphism in Sm_S . We consider the morphism

$$\eta_f : E_X \rightarrow f_* E_{\tilde{X}} \tag{5a}$$

in $\text{Spt}_{S^1}(X_{\text{Nis}})$ given on an étale morphism $V \rightarrow X$ of finite type as the map

$$E_X(V) = E(V) \rightarrow E(V \times_X \tilde{X}) = E_{\tilde{X}}(V \times_X \tilde{X})$$

induced by the projection. This clearly generalizes the construction (3a) where the morphism f was assumed to be étale. Indeed, in this case we have $f^*(E_X) \cong E_{\tilde{X}}$.

Construction 5.3 Next, for $E \in \text{Spt}_{S^1}(\text{Sm}_S)$, a closed subset $Z \subseteq X$ and a pullback diagram

$$\begin{array}{ccc} \tilde{X} \setminus \tilde{Z} & \xrightarrow{\tilde{j}} & \tilde{X} \\ \downarrow \tilde{f} & & \downarrow f \\ X \setminus Z & \xrightarrow{j} & X \end{array} \tag{5b}$$

we want to define a morphism

$$\eta_f : E_{Z/X} \rightarrow f_* E_{\tilde{Z}/\tilde{X}} \tag{5c}$$

that coincides with (5a) for $Z = X$. First note that the commutative diagram (5b) induces the base-change morphism

$$f^* j_* \xrightarrow{\text{(b.c.)}} \tilde{j}_* \tilde{f}^*$$

Further, by adjunction, Construction 5.2 induces a map

$$\tilde{f}^* j^* E_X \cong \tilde{f}^* E_{X \setminus Z} \xrightarrow{\eta_f^b} E_{\tilde{X} \setminus \tilde{Z}} \cong \tilde{j}^* E_{\tilde{X}}$$

Composition with the unit yields a morphism

$$j_* j^* E_X \xrightarrow{\eta} f_* f^* j_* j^* E_X \xrightarrow{f_*(\text{b.c.})j^*} f_* \tilde{j}_* \tilde{f}^* j^* E_X \xrightarrow{f_* \tilde{j}_* \eta_f^b} f_* \tilde{j}_* \tilde{j}^* E_{\tilde{X}} \tag{5d}$$

which is seen to fit into a commutative square

$$\begin{array}{ccc} E_X & \xrightarrow{\eta_j} & j_* j^* E_X \\ \downarrow \eta_f & & \downarrow \\ f_* E_{\tilde{X}} & \xrightarrow{f_* \eta_j} & f_* \tilde{j}_* \tilde{j}^* E_{\tilde{X}} \end{array}$$

inducing the desired map η_f by taking horizontal homotopy fibres.

Lemma 5.4 *Let*

$$X_2 \xrightarrow{f_2} X_1 \xrightarrow{f_1} X$$

be two morphisms of noetherian schemes of finite Krull dimension, $Z \subseteq X$ a closed subset and $Z_1 := Z \times_X X_1, Z_2 := Z \times_X X_2$ the respective base changes. Then we have a commutative triangle

$$\begin{array}{ccc} E_{Z/X} & \xrightarrow{\eta_{f_1}} & f_{1,*} E_{Z_1/X_1} \\ & \searrow \eta_{f_1 f_2} & \downarrow f_{1,*} \eta_{f_2} \\ & & f_{1,*} f_{2,*} E_{Z_2/X_2} \end{array}$$

in $\text{Spt}_{S^1}(X_{\text{Nis}})$ of the respective morphisms (5c).

Proof By adjointness it suffices to show the commutativity of the outer square of the diagram

$$\begin{array}{ccccc}
 f_2^* f_1^* j_*(EU) & \xrightarrow{f_2^*(b.c.)} & f_2^* j_{1,*} f_{1|U}^*(EU) & \xrightarrow{f_2^* j_{1,*} \eta_{f_{1|U}}^b} & f_2^* j_{1,*} EU_1 \xrightarrow{(b.c.)} j_{2,*} f_{2|U_1}^*(EU_1) \\
 \downarrow (b.c.) & \swarrow & \downarrow (b.c.) f_{1|U}^* & \xrightarrow{j_{2,*} \eta_{(f_1 f_2)|U}^b} & \downarrow j_{2,*} \eta_{f_{2|U_1}}^b \\
 j_{2,*} (f_1 f_2)_{|U}^*(EU) & \xrightarrow{\hspace{10em}} & & & j_{2,*} (EU_2)
 \end{array}$$

where $j : U \hookrightarrow X$, $j_1 : U_1 \hookrightarrow X_1$ and $j_2 : U_2 \hookrightarrow X_2$ are the respective open complements of Z, Z_1 and Z_2 and where $f_{1|U} : U_1 \rightarrow U$, $f_{2|U_1}$ and $(f_1 f_2)_{|U}$ are the respective restrictions. The triangle on the left-hand side commutes as base change morphisms are compatible with composition and the commutativity of the remaining part is easily seen. □

5.5 Recall, that a Nisnevich local fibrant spectrum $E \in \text{Spt}_{S^1}(S)$ is an \mathbb{A}^1 -Nisnevich local fibrant spectrum if $E(X) \rightarrow E(X \times \mathbb{A}^1)$ is an equivalence for all $X \in \text{Sm}_S$.

Lemma 5.6 *Let $E \in \text{Spt}_{S^1}(\text{Sm}_S)$ be an \mathbb{A}^1 -Nisnevich local fibrant spectrum. Let $X \in \text{Sm}_S$ be a scheme, $Z \subseteq X$ a closed subset and $\pi : \mathbb{A}_X^1 \rightarrow X$ the projection. Then the canonical map (c.f. (5c))*

$$E_{Z/X} \xrightarrow{\eta_\pi} \pi_* E_{\mathbb{A}_Z^1/\mathbb{A}_X^1}$$

is a weak equivalence.

Proof By construction of the map in question as a homotopy fibre, it suffices to show that the two maps

$$E_X \xrightarrow{\eta_\pi} \pi_* E_{\mathbb{A}_X^1}$$

and

$$j_* j^* E_X \rightarrow \pi_* \tilde{j}_* \tilde{j}^* E_{\mathbb{A}_X^1}$$

from (5d) are both object-wise weak equivalences, This can be checked directly by evaluation on an object $V \rightarrow X$ of the site X_{Nis} . □

Lemma 5.7 *Let $E \in \text{Spt}_{S^1}(\text{Sm}_S)$ be an \mathbb{A}^1 -Nisnevich local fibrant spectrum. Let $X \in \text{Sm}_S$ be a scheme and $s : X \hookrightarrow \mathbb{A}_X^1$ a section of the projection $\pi : \mathbb{A}_X^1 \rightarrow X$. Then there is a commutative diagram*

$$\begin{array}{ccc}
 E_{Z/X} & \xrightarrow{\eta_\pi} & \pi_* E_{\mathbb{A}_Z^1/\mathbb{A}_X^1} \\
 & \searrow & \downarrow \pi_* \eta_s \\
 & & \pi_* s_* E_{Z/X}
 \end{array}$$

of weak equivalences. In particular, for another section $s' : X \hookrightarrow \mathbb{A}_X^1$ of the projection, the morphisms $\pi_* \eta_s$ and $\pi_* \eta_{s'}$ are equal in the homotopy category.

Proof This follows from the previous Lemmas 5.4 and 5.6. □

Lemma 5.8 *Let $E \in \text{Spt}_{S^1}(\text{Sm}_S)$ be an \mathbb{A}^1 -Nisnevich local fibrant spectrum. Let $V \in \text{Sm}_S$ and $Z \hookrightarrow \mathbb{A}_V^1$ a closed subscheme such that $Z \hookrightarrow \mathbb{A}_V^1 \xrightarrow{\pi} V$ is finite. Let $\bar{Z} := \pi(Z)_{\text{red}}$ be*

the reduced image. Then $\text{codim}(\mathbb{A}^1_{\bar{Z}}, \mathbb{A}^1_V) = \text{codim}(Z, \mathbb{A}^1_V) - 1$ and the forget support map (see Lemma 3.7) induces the trivial morphism

$$\pi_* E_{Z/\mathbb{A}^1_V} \rightarrow \pi_* E_{\mathbb{A}^1_{\bar{Z}}/\mathbb{A}^1_V}$$

in the homotopy category.

Proof Consider the diagram

$$\begin{array}{ccc} \mathbb{A}^1_V & \xrightarrow{\pi} & V \\ j \downarrow & \nearrow \bar{\pi} & \\ \mathbb{P}^1_V & & \end{array}$$

where the non-vertical maps are the projections and j is the canonical open immersion. Let us first prove that the triangle

$$\begin{array}{ccc} \bar{\pi}_* E_{\mathbb{P}^1_{\bar{Z}}/\mathbb{P}^1_V} & \xrightarrow{\bar{\pi}_* \eta_{\bar{s}_\infty}} & \bar{\pi}_* \bar{s}_\infty,* E_{\bar{Z}/V} = E_{\bar{Z}/V} \\ & \searrow \bar{\pi}_* \eta_j & \downarrow \eta\pi \\ & & \pi_* E_{\mathbb{A}^1_{\bar{Z}}/\mathbb{A}^1_V} \end{array} \tag{5e}$$

commutes in the homotopy category, where $\bar{s}_\infty : V \hookrightarrow \mathbb{P}^1_V$ is the section at infinity. Let $\bar{s}_0 : V \xrightarrow{s_0} \mathbb{A}^1_V \xrightarrow{j} \mathbb{P}^1_V$ denote the zero-section. Since by Lemma 5.7 the morphism

$$\pi_* \eta_{s_0} : \pi_* E_{\mathbb{A}^1_{\bar{Z}}/\mathbb{A}^1_V} \rightarrow \pi_* s_{0,*} E_{\bar{Z}/V} = E_{\bar{Z}/V}$$

is a weak equivalence, it suffices to show that the outer triangle of the enlarged diagram

$$\begin{array}{ccc} \bar{\pi}_* E_{\mathbb{P}^1_{\bar{Z}}/\mathbb{P}^1_V} & \xrightarrow{\bar{\pi}_* \eta_{\bar{s}_\infty}} & E_{\bar{Z}/V} \\ & \searrow \bar{\pi}_* \eta_j & \downarrow \eta\pi \\ & & \pi_* E_{\mathbb{A}^1_{\bar{Z}}/\mathbb{A}^1_V} \\ & \searrow \bar{\pi}_* \eta_{\bar{s}_0} & \simeq \downarrow \pi_* \eta_{s_0} \\ & & E_{\bar{Z}/V} \end{array}$$

commutes. Indeed, the bottom triangle is obtained by applying $\bar{\pi}_*$ to a commutative triangle considered in Lemma 5.4 for $\bar{s}_0 = j s_0$. By the same Lemma 5.4 applied to $\text{id} = \pi s_0$, the right vertical composition is the identity. Hence, it suffices to show that $\bar{\pi}_* \eta_{\bar{s}_0} = \bar{\pi}_* \eta_{\bar{s}_\infty}$ holds in the homotopy category.

Since the sections \bar{s}_0 and $\bar{s}_\infty : V \rightarrow \mathbb{P}^1_V$ both factorize through the open immersion $j' : \mathbb{P}^1_V \setminus s_1(V) \hookrightarrow \mathbb{P}^1_V$ via s'_0 and $s'_\infty : V \rightarrow \mathbb{P}^1_V \setminus s_1(V)$, we have a commutative diagram

$$\begin{array}{ccc} \bar{\pi}_* E_{\mathbb{P}^1_{\bar{Z}}/\mathbb{P}^1_V} & \xrightarrow{\bar{\pi}_* j'} & \bar{\pi}_* j'_* E_{(\mathbb{P}^1_{\bar{Z}} \setminus s_1(V))/(\mathbb{P}^1_V \setminus s_1(V))} \\ & \searrow \bar{\pi}_* \eta_{\bar{s}_0} & \swarrow \pi'_* \eta_{s'_0} \\ & & E_{\bar{Z}/V} \end{array}$$

(and likewise for \bar{s}_∞ and s'_∞). Here, $\pi' : \mathbb{P}^1_V \setminus s_1(V) \rightarrow V$ is the projection. As $\mathbb{P}^1_V \setminus s_1(V) \cong \mathbb{A}^1_V$ and $\mathbb{P}^1_{\bar{Z}} \setminus s_1(V) \cong \mathbb{A}^1_{\bar{Z}}$, we obtain $\pi'_* \eta_{s'_0} = \pi'_* \eta_{s'_\infty}$ by Lemma 5.7, thus $\bar{\pi}_* \eta_{\bar{s}_0} = \bar{\pi}_* \eta_{\bar{s}_\infty}$. Summing up, this yields the commutativity of diagram (5e).

In order to show that the morphism in question

$$\pi_* E_{Z/\mathbb{A}_V^1} = \bar{\pi}_* j_* E_{Z/\mathbb{A}_V^1} \rightarrow \bar{\pi}_* j_* E_{\mathbb{A}_Z^1/\mathbb{A}_V^1} = \pi_* E_{\mathbb{A}_Z^1/\mathbb{A}_V^1}$$

is trivial in the homotopy category, we consider the diagram

$$\begin{array}{ccccc}
 \bar{\pi}_* j_* E_{Z/\mathbb{A}_V^1} & \longrightarrow & \bar{\pi}_* j_* E_{\mathbb{A}_Z^1/\mathbb{A}_V^1} & & \\
 \uparrow \simeq & & \uparrow & \swarrow & \\
 \bar{\pi}_* \eta_j & & \bar{\pi}_* \eta_j & & E_{\bar{Z}/V} \\
 \uparrow & & \uparrow \bar{\pi}_* \bar{s}_\infty & \nearrow & \swarrow \bar{\pi}_* j''_* \eta_{s_\infty} \\
 \pi_* E_{Z/\mathbb{P}_V^1} & \longrightarrow & \pi_* E_{\mathbb{P}_Z^1/\mathbb{P}_V^1} & \xrightarrow{\bar{\pi}_* \eta_{j''}} & \pi_* j''_* E_{(\mathbb{P}_Z^1 \setminus Z)/(\mathbb{P}_V^1 \setminus Z)}
 \end{array}$$

where the middle triangle is (5e). The left horizontal maps are induced by the respective forget support maps. For the right triangle, we note that $\bar{s}_\infty : V \hookrightarrow \mathbb{P}_V^1$ factorizes through $j'' : \mathbb{P}_V^1 \setminus Z \hookrightarrow \mathbb{P}_V^1$ via $s''_\infty : V \hookrightarrow \mathbb{P}_V^1 \setminus Z$. The commutativity of the square on the left-hand side is clear. The triangle on the right-hand side commutes again by Lemma 5.4. We observe that the lower horizontal line is given by $\bar{\pi}_*$ applied to the exact triangle of Lemma 3.7. In particular, it is an exact triangle itself and therefore the composition is trivial. Finally, the left vertical arrow is a weak equivalence by the excision Lemma 3.11. Hence the morphism $\pi_* E_{Z/\mathbb{A}_V^1} \rightarrow \pi_* E_{\mathbb{A}_Z^1/\mathbb{A}_V^1}$ in question is trivial in the homotopy category.

For the assertion $\text{codim}(\mathbb{A}_Z^1, \mathbb{A}_V^1) = \text{codim}(Z, \mathbb{A}_V^1) - 1$ we can argue component-wise on Z so we may assume that Z is irreducible. Further, we can replace \mathbb{A}_V^1 by a base change along a flat morphism $V' \rightarrow V$. In particular, we may assume that V is a local scheme with closed point \bar{Z} . As Z is finite over \bar{Z} , it is just a finite union of points in the curve $\mathbb{A}_{\bar{Z}}^1$. Thus, $\text{codim}(Z, \mathbb{A}_{\bar{Z}}^1) = 1$ and the assertion follows by Lemma 2.7. \square

5.9 In [3, Definition 4.4], the authors define the notion of an étale sheaf A to be effaceable at a point $x \in X$. (There is a corresponding dual definition of homological effaceability in [3, Definition 4.4].) The connection of this definition with the next Proposition 5.10 and its Corollary 5.12 is as follows. As we will explain in more detail in 6.2, there is a spectrum E which satisfies $E_X^n(U) = H^n(U, A)$ and $E_{Z/X}^n(U) = H_Z^n(U, A)$ for an étale sheaf A . With this terminology and under the required assumptions, the next Proposition 5.10 shows that A is “Nisnevich-locally” effaceable at the point $x \in X$.

Proposition 5.10 *Let $E \in \text{Spt}_{\mathcal{S}^1}(\text{Sm}_{\mathcal{S}})$ be an \mathbb{A}^1 -Nisnevich local fibrant spectrum. Let $X \in \text{Sm}_{\mathcal{S}}$, $Z \hookrightarrow X$ be a closed subscheme and $x \in X$ be a point. If x lies in the special fibre X_σ , assume that Z_σ does not contain any connected components of X_σ . Then, Nisnevich-locally on X around x , there exists a $V \in \text{Sm}_{\mathcal{S}}$, a smooth relative curve $p : X \rightarrow V$ with Z finite over V and a closed subscheme $Z' \hookrightarrow X$ containing Z such that $\text{codim}(Z', X) = \text{codim}(Z, X) - 1$ and the forget support map induces the trivial morphism*

$$p_* E_{Z/X} \rightarrow p_* E_{Z'/X}$$

in the homotopy category. In particular, $E_{Z/X}(X) \rightarrow E_{Z'/X}(X)$ is trivial in this case.

Proof Possibly after shrinking X Nisnevich-locally around x , we find a Nisnevich distinguished square

$$\begin{CD} X \setminus Z @>>> X \\ @VVV @VVfV \\ \mathbb{A}_V^1 \setminus f(Z) @>>> \mathbb{A}_V^1 \end{CD} \tag{5f}$$

such that $Z \hookrightarrow X \xrightarrow{f} \mathbb{A}_V^1 \xrightarrow{\pi} V$ is finite by [12, Theorem 2.1]. Let $p: X \xrightarrow{f} \mathbb{A}_V^1 \xrightarrow{\pi} V$ denote the composition and set $\bar{Z} := p(Z)_{\text{red}}$ and $Z' := p^{-1}(\bar{Z})$. Since f and π and hence the composition p is flat, the assertion about the codimensions holds true. By the excision Lemma 3.11, the upper horizontal morphism of the diagram

$$\begin{CD} E_{f(Z)/\mathbb{A}_V^1} @>\cong>> f_*E_{Z/X} \\ @VVV @VVV \\ E_{\mathbb{A}_Z^1/\mathbb{A}_V^1} @>>> f_*E_{Z'/X} \end{CD}$$

is an equivalence, where the vertical maps are the respective forget support maps and $f^{-1}f(Z) = Z$. Application of π_* yields the commutative diagram

$$\begin{CD} \pi_*E_{f(Z)/\mathbb{A}_V^1} @>\cong>> \pi_*E_{Z/X} \\ @VVV @VVV \\ \pi_*E_{\mathbb{A}_Z^1/\mathbb{A}_V^1} @>>> \pi_*E_{Z'/X}. \end{CD}$$

The left vertical morphism is trivial by the previous Lemma 5.8. Hence the right vertical morphism is trivial which proves the claim. \square

5.11 Denote by $X_{x,\eta}^h$ the generic fibre $\text{Spec}(\mathcal{O}_{X,x}^h \otimes_{\circ} k)$ of the Henselian local scheme at x . Similar to 4.5, by $E(X_{x,\eta}^h)$ we mean $\text{colim}_{(W,w)} E(W_\eta)$, where (W, w) runs through the Nisnevich neighbourhoods of x and W_η is the generic fibre.

Corollary 5.12 *Under the assumptions of Proposition 5.10, the forget support map*

$$E_{Z/X}(X_{x,\eta}^h) \rightarrow E_X(X_{x,\eta}^h)$$

is trivial.

Proof By [12, Theorem 2.1], there is a cofinal family of Nisnevich neighbourhoods (W, w) of x admitting a Nisnevich distinguished square of the form (5f) with the additional finiteness assumption. We even claim that for such neighbourhoods (W, w) , the forget support map $E_{Z/X}(W_\eta) \rightarrow E_X(W_\eta)$ is trivial. To show this, we may assume $W = X$, i.e., we assume X admits a Nisnevich distinguished square as in (5f) with Z/V finite. On the generic fibres, we still have a distinguished square

$$\begin{CD} X_\eta \setminus Z_\eta @>>> X_\eta \\ @VVV @VVf_\eta V \\ \mathbb{A}_{V_\eta}^1 \setminus f_\eta(Z_\eta) @>>> \mathbb{A}_{V_\eta}^1 \end{CD}$$

and as pullback, Z_η/V_η is still finite. Accordingly, the arguments in the proof of Proposition 5.10 go through for $Z_\eta \subseteq X_\eta$, as well. In particular, the forget support map $E_{Z/X}(X_\eta) \rightarrow E_X(X_\eta)$ is indeed trivial. \square

Theorem 5.13 *Let S be a Dedekind scheme with only infinite residue fields. Moreover, let $E \in \text{Spt}_{S^1}(\text{Sm}_S)$ be an \mathbb{A}^1 -Nisnevich local fibrant spectrum and $X \in \text{Sm}_S$ of dimension d . The complex of sheaves over the small Nisnevich site X_{Nis}*

$$\begin{aligned}
 0 \rightarrow (E_X^n)^\sim \xrightarrow{\tilde{e}} \bigoplus_{z \in X^{(0)}} \mathfrak{z} * \mathfrak{z}^* E_{Z/X}^n \xrightarrow{d^0} \bigoplus_{z \in X^{(1)}} \mathfrak{z} * \mathfrak{z}^* E_{Z/X}^{n+1} \xrightarrow{d^1} \dots \\
 \dots \xrightarrow{d^{d-2}} \bigoplus_{z \in X^{(d-1)}} \mathfrak{z} * \mathfrak{z}^* E_{Z/X}^{n+d-1} \xrightarrow{d^{d-1}} \bigoplus_{z \in X^{(d)}} \mathfrak{z} * \mathfrak{z}^* E_{Z/X}^{n+d} \rightarrow 0
 \end{aligned}$$

*is exact, possible except at the spots $(E_X^n)^\sim$ and $\bigoplus_{z \in X^{(1)}} \mathfrak{z} * \mathfrak{z}^* E_{Z/X}^{n+1}$. Moreover, if for each point x of X the forget support map for the special fibre*

$$E_{X_\sigma/X}(X_x^h) \rightarrow E_X(X_x^h)$$

is trivial, then it is exact everywhere and thus a resolution of $(E_X^n)^\sim$ by flabby Nisnevich sheaves. In this case, we have

$$H^p(Y, (E_X^n)^\sim) \cong H^p(\mathcal{G}^\bullet(E, n)(Y)).$$

for $Y \in X_{\text{Nis}}$ which vanishes for $p > d$.

Proof Since exactness is checked stalk-wise and we can compute the stalk at a point $x \in X$ after henselization of the local scheme obtained from S at the image of x , we may assume, that S is the spectrum of a Henselian discrete valuation ring with infinite residue field. Now the first result follows from Proposition 4.6 and Proposition 5.10.

Suppose the forget support maps $E_{X_\sigma/X}(X_x^h) \rightarrow E_X(X_x^h)$ are trivial for all points x . By our assumption and Propositions 4.6 and 5.10, it is enough so show that the forget support map $E_{Z/X}^n(X_x^h) \rightarrow E_X^n(X_x^h)$ is trivial for closed subsets $X_\sigma \subsetneq Z \subsetneq X$. We may replace X by the Henselian local scheme X_x^h . Write $Z = Z_1 \cup Z_2$ with $Z_1 = X_\sigma$ and $X_\sigma \not\subseteq Z_2$. Let $U = X \setminus Z$ and $U_i = X \setminus Z_i$ be the respective open complements. Observe that $U_1 = X_\eta$ and $U = U_{2,\eta}$ are just the generic fibres. Consider the exact triangles

$$E_{Z_2/X}(U_1) \rightarrow E_X(U_1) \rightarrow E_X(U).$$

and

$$E_{Z_1/X}(X) \rightarrow E_X(X) \rightarrow E_X(U_1).$$

By our assumption, the forget support map in the latter triangle is trivial, so the restriction map $E_X(X) \rightarrow E_X(U_1)$ admits a retraction r_1 . By Corollary 5.12, the forget support map in the former triangle is trivial, so the restriction map $E_X(U_1) \rightarrow E_X(U)$ admits a retraction r_2 . Set $r := r_1 \circ r_2: E_X(U) \rightarrow E_X(X)$. By construction, r is a retraction of the restriction map $E_X(X) \rightarrow E_X(U)$. Thus, using the exact triangle

$$E_{Z/X}(X) \rightarrow E_X(X) \rightarrow E_X(U),$$

we get that the forget support map $E_{Z/X}(X) \rightarrow E_X(X)$ is indeed trivial. □

6 A Bloch–Ogus theorem for étale cohomology

In this section we want to apply Theorem 5.13 to étale cohomology. Let us first fix the situation:

6.1 We are in the situation of 5.1. For the whole section, we fix an essentially smooth scheme X/S , connected and of finite dimension. Let us denote the structural morphism by $p_X : X \rightarrow S$. We fix a coefficient group $\Lambda := \mathbb{Z}/m$ for an integer $m > 0$ prime to p . We work in the derived category $\mathcal{D}_c^b(X_{\text{et}}, \Lambda)$ of bounded (above and below) complexes all of whose cohomology sheaves are constructible sheaves of Λ -modules. By an l.c.c. complex K^\bullet , we mean a complex $K^\bullet \in \mathcal{D}_c^b(X_{\text{et}}, \Lambda)$ with locally constant cohomology sheaves $H^q(K^\bullet)$ for all q .

6.2 Let $\varepsilon : X_{\text{et}} \rightarrow X_{\text{Nis}}$ be the canonical morphism of sites. Note that $\mathbb{R}\Gamma(X_{\text{et}}, -) \simeq \mathbb{R}\Gamma(X_{\text{Nis}}, \mathbb{R}\varepsilon_*(-))$. By abuse of notation, let us denote by ε also the corresponding morphism $\text{Sm}_{S,\text{et}} \rightarrow \text{Sm}_{S,\text{Nis}}$ of the smooth sites. For an l.c.c. complex K^\bullet in $\mathcal{D}_c^b(S_{\text{et}}, \Lambda)$, we denote by K^\bullet also the complex in $\mathcal{D}^b(\text{Sm}_{S,\text{et}}, \Lambda)$ that restricts to $p_X^* K^\bullet$ on each small site X_{et} . Further, we fix a Nisnevich local fibrant spectrum $E(K^\bullet) \in \text{Spt}_{S^1}(\text{Sm}_S)$ corresponding to $\mathbb{R}\varepsilon_* K^\bullet$ under the Dold–Kan correspondence.

Lemma 6.3 *The spectrum $E(K^\bullet)$ is \mathbb{A}^1 -local.*

Proof Indeed, the projection $\pi : \mathbb{A}_X^1 \rightarrow X$ induces a quasi-isomorphism $p_X^* K^\bullet \rightarrow \mathbb{R}\pi_* \pi^* p_X^* K^\bullet$ (e.g. [6, Corollary 7.7.4]) and hence a quasi-isomorphism on cohomology

$$\mathbb{R}\Gamma(X_{\text{Nis}}, \mathbb{R}\varepsilon_* p_X^* K^\bullet) \rightarrow \mathbb{R}\Gamma(\mathbb{A}_{X,\text{Nis}}^1, \mathbb{R}\varepsilon_* p_{\mathbb{A}_X^1}^* K^\bullet).$$

Under the Dold–Kan correspondence this translates to our claim. □

6.4 In order to apply Theorem 5.13 to the \mathbb{A}^1 -local spectrum $E(K^\bullet)$, we need to show that the forget support maps

$$E(K^\bullet)_{X_\sigma/X}(X_x^h) \rightarrow E(K^\bullet)_X(X_x^h)$$

vanish for all points x in X . Here, X_x^h denotes the spectrum of the henselization of the local ring of X at x . Unravelling the definitions, these maps are just the forget support maps

$$\mathbb{R}\Gamma_{X_x,\sigma}^h(X_{x,\text{et}}^h, p_X^* K^\bullet) \rightarrow \mathbb{R}\Gamma(X_{x,\text{et}}^h, p_X^* K^\bullet)$$

of étale cohomology.

In the following, we will make use of Gabber’s absolute purity theorem – but not in its full strength. The following easy special case will be sufficient for our cause:

Lemma 6.5 *In the situation of 6.1, let $i : Z \hookrightarrow X$ be a closed subscheme of codimension c , contained in the special fibre of X/S . Assume Z/\mathbb{F} is smooth and connected. Then the canonical morphism $\mathbb{R}i^! K^\bullet|_X \rightarrow K^\bullet|_Z(-c)[-2c]$ is a quasi-isomorphism for all l.c.c. complexes $K^\bullet \in \mathcal{D}_c^b(S_{\text{et}}, \Lambda)$.*

Proof Say, X/S and Z/\mathbb{F} have relative dimension m and n respectively. In particular, $c = m - n + 1$. Consider the commutative diagram

$$\begin{array}{ccc} Z & \xleftarrow{i} & X \\ \downarrow p_Z & & \downarrow p_X \\ \text{Spec}(\mathbb{F}) & \xleftarrow{\sigma} & S. \end{array}$$

By Poincaré-duality for X/S (respectively Z/\mathbb{F}), $\mathbb{R}p_X^! K^\bullet \simeq p_X^* K^\bullet(m)[2m]$ (respectively $\mathbb{R}p_Z^! \sigma^* K^\bullet \simeq p_Z^* \sigma^* K^\bullet(n)[2n]$). Further, by the special case of absolute purity for the closed

point in S (which is an easy exercise – e.g. the proof of [6, Lemma 8.3.6] goes through unchanged for l.c.c. sheaves and hence for l.c.c. complexes), $\mathbb{R}\sigma^!K^\bullet \simeq \sigma^*K^\bullet(-1)[-2]$. Summing up, we get

$$\begin{aligned} \mathbb{R}i^!p_X^*K^\bullet &\simeq \mathbb{R}i^!\mathbb{R}p_X^!K^\bullet(-m)[-2m] \\ &\simeq \mathbb{R}p_Z^!\mathbb{R}\sigma^!K^\bullet(-m)[-2m] \\ &\simeq \mathbb{R}p_Z^!\sigma^*K^\bullet(-m-1)[-2m-2] \\ &\simeq p_Z^*\sigma^*K^\bullet(-c)[-2c], \end{aligned}$$

finishing the proof. □

Lemma 6.6 *In the situation of 6.1, assume that X is Henselian local with closed point x in the special fibre of X/S . Then the canonical morphism $\sigma_{X,*}\mathbb{R}\sigma_X^!\Lambda \rightarrow \Lambda$ induces the trivial morphism in $\mathcal{D}^b(k(x)_{\text{et}}, \Lambda)$:*

$$x^*\sigma_{X,*}\mathbb{R}\sigma_X^!\Lambda \xrightarrow{\simeq 0} x^*\Lambda \simeq \Lambda.$$

In particular, the canonical map $\mathbb{R}\Gamma_{X_\sigma}(X_{\text{et}}, \Lambda) \rightarrow \mathbb{R}\Gamma(X_{\text{et}}, \Lambda)$ is trivial.

Proof The second claim follows from the first. Indeed, as X is local Henselian $\mathbb{R}\Gamma(X_{\text{et}}, -) \simeq \mathbb{R}\Gamma(k(x)_{\text{et}}, x^*(-))$. For the first claim, it is enough to show that the Tate-twist

$$x^*(\sigma_{X,*}\mathbb{R}\sigma_X^!\Lambda(1) \rightarrow \Lambda(1)) \tag{6a}$$

is trivial in $\mathcal{D}^b(k(x)_{\text{et}}, \Lambda)$. By Lemma 6.5, $\mathbb{R}\sigma_X^!\Lambda(1) \simeq \Lambda[-2]$. In particular, the sheaf-cohomology of (6a) in degree 2 is given by

$$\Lambda = H^0(X_{\sigma,\text{et}}, \Lambda) \simeq H_{X_\sigma}^2(X_{\text{et}}, \Lambda(1)) \rightarrow H^2(X_{\text{et}}, \Lambda(1)), \quad 1 \mapsto \hat{c}_1[\mathcal{O}(X_\sigma)], \tag{6b}$$

i.e., is trivial as X is a local scheme. Further,

$$x^*\sigma_{X,*}\mathbb{R}\sigma_X^!\Lambda(1) \simeq x^*\sigma_{X,*}\Lambda[-2] \simeq \Lambda[-2]$$

which implies

$$\text{Hom}_{\mathcal{D}^b(k(x)_{\text{et}}, \Lambda)}(x^*\sigma_{X,*}\mathbb{R}\sigma_X^!\Lambda(1), x^*\Lambda(1)) \cong H^2(k(x)_{\text{et}}, \Lambda(1))$$

and (6a) corresponds to a class contained in the image of (6b) (more precisely, (6a) corresponds to the class $\hat{c}_1[\mathcal{O}(X_\sigma)]$), hence it is trivial. □

Corollary 6.7 *In the situation of 6.1, assume that X is Henselian local with closed point x in the special fibre of X/S . Let $K^\bullet \in \mathcal{D}_c^b(S_{\text{et}}, \Lambda)$ be a l.c.c. complex. Then the canonical morphism $\sigma_{X,*}\mathbb{R}\sigma_X^!K^\bullet|_X \rightarrow K^\bullet|_X$ induces the trivial morphism in $\mathcal{D}^b(k(x)_{\text{et}}, \Lambda)$:*

$$x^*\sigma_{X,*}\mathbb{R}\sigma_X^!K^\bullet|_X \xrightarrow{\simeq 0} x^*K^\bullet|_X.$$

In particular, the canonical map $\mathbb{R}\Gamma_{X_\sigma}(X_{\text{et}}, K^\bullet|_X) \rightarrow \mathbb{R}\Gamma(X_{\text{et}}, K^\bullet|_X)$ is trivial.

Proof By Lemma 6.6, x^* applied to

$$(\sigma_{X,*}\mathbb{R}\sigma_X^!\Lambda \rightarrow \Lambda) \otimes^{\mathbb{L}} p_X^*K^\bullet \tag{6c}$$

is trivial. By the projection formula and Lemma 6.5, (6c) is isomorphic to the canonical morphism $\sigma_{X,*}\mathbb{R}\sigma_X^!p_X^*K^\bullet \rightarrow p_X^*K^\bullet$, so the claim follows. □

Combining Theorem 5.13 and Corollary 6.7, we get:

Theorem 6.8 *Let S be the spectrum of a Henselian discrete valuation ring with infinite residue field \mathbb{F} . Let X/S be smooth, $d = \dim(X)$ and K^\bullet an l.c.c. complex in $\mathcal{D}_c^b(S_{\text{et}}, \Lambda)$. Then the Nisnevich Gersten complex $\mathcal{G}^\bullet(E(K^\bullet), n)$ is a flasque resolution of the Nisnevich sheafification $\mathbb{R}^n \varepsilon_* K^\bullet|_X$ of étale cohomology with coefficients K^\bullet . In particular, we get the exact sequence*

$$0 \rightarrow \mathbb{R}^n \varepsilon_* K^\bullet|_X \rightarrow \bigoplus_{z \in X^{(0)}} z_* H^n(k(z), K^\bullet|_{k(z)}) \rightarrow \dots$$

$$\dots \rightarrow \bigoplus_{z \in X^{(d)}} z_* H^{n-d}(k(z), K^\bullet|_{k(z)}(-d)) \rightarrow 0.$$

Proof The spectrum $E(K^\bullet)$ is \mathbb{A}^1 -local by Lemma 6.3. Combining Theorem 5.13 and Corollary 6.7, we get that $\mathcal{G}^\bullet(E(K^\bullet), n)$ is a flasque resolution of $\mathbb{R}^n \varepsilon_* K^\bullet|_X$.

Let us compute $\mathcal{G}^s(E(K^\bullet), n) = \bigoplus_{z \in X^{(s)}} \mathfrak{z}^* \mathfrak{z}^* E(K^\bullet)_{Z/X}^{n+s}$: In the proof of Proposition 3.23 we saw that $\mathfrak{z}^* E(K^\bullet)_{Z/X}^{n+s} = z_* z^* E(K^\bullet)_{z/X_{z,\text{loc}}}^{n+s}$. Unravelling the definitions, $z^* E(K^\bullet)_{z/X_{z,\text{loc}}}^{n+s} \cong H_z^{n+s}(X_{z,\text{et}}^{\text{loc}}, K^\bullet)$. By absolute purity, $H_z^{n+s}(X_{z,\text{et}}^{\text{loc}}, K^\bullet) \cong H^{n-s}(k(z), K^\bullet(-s))$, which finishes the proof. □

Remark 6.9 We can avoid absolute purity in its full strength if we assume k and \mathbb{F} to be perfect: Computing $\mathfrak{z}^* E(K^\bullet)_{Z/X}^{n+s}$ under this assumption, we may assume Z to be smooth over k (if z is contained in the generic fibre of X/S) or smooth over \mathbb{F} (if z is contained in the special fibre of X/S) by generic smoothness. In both cases, $\mathfrak{z}^* E(K^\bullet)_{Z/X}^{n+s} \cong z_* H^{n-s}(k(z), K^\bullet(-s))$, either by relative purity or by Lemma 6.5.

Taking Nisnevich stalks, we get:

Corollary 6.10 *Let S be the spectrum of a Henselian discrete valuation ring with infinite residue field \mathbb{F} . Let X/S be smooth of finite type, $d = \dim(X)$ and K^\bullet an l.c.c. complex in $\mathcal{D}_c^b(S_{\text{et}}, \Lambda)$. Let x be a point of X and $Y = X_x^h$ the Nisnevich local scheme at x . Then there is an exact sequence*

$$0 \rightarrow H^n(Y_{\text{et}}, K^\bullet|_Y) \xrightarrow{e} \bigoplus_{z \in Y^{(0)}} H^n(k(z), z^* K^\bullet|_Y) \xrightarrow{d^0} \dots$$

$$\dots \xrightarrow{d^{d-1}} \bigoplus_{z \in Y^{(d)}} H^{n-d}(k(z), z^* K^\bullet|_Y(-d)) \rightarrow 0.$$

Remark 6.11 Using the Bloch–Kato–Conjecture, in [7] Geisser proved the exactness of the Gersten complex in degree n for X/S smooth even for $\varepsilon: X_{\text{et}} \rightarrow X_{\text{Zar}}$, but only for coefficients $K^\bullet = \Lambda(r) = \mu_m^{\otimes r}$ for $n \leq r$. If S is not strictly Henselian, this assumption excludes $H^{2r}(Y_{\text{et}}, \Lambda(r))$ for $r > 0$, i.e., the targets of the cycle class maps.

Funding Open Access funding enabled and organized by Projekt DEAL.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Ayoub, J.: Les six opérations de Grothendieck et le formalisme des cycles évanescents dans le monde motivique. I. *Astérisque* (2007), no. 314, x+466 pp. (2008)
2. Bloch, S.: A note on Gersten's conjecture in the mixed characteristic case, *Applications of algebraic K-theory to algebraic geometry and number theory, Part I, II* (Boulder, Colo., 1983), 75–78, *Contemp. Math.* 55, Amer. Math. Soc., Providence, RI (1986)
3. Bloch, S., Ogus, A.: Gersten's conjecture and the homology of schemes. *Ann. Sci. École Norm. Sup.* (4) 7, 181–201 (1974)
4. Colliot-Thélène, J.-L., Hoobler, R.T., Kahn, B.: The Bloch-Ogus-Gabber theorem, *Algebraic K-theory* (Toronto, ON, 1996), vol. 16. *Fields Inst. Commun. Amer. Math. Soc.*, Providence, RI (1997)
5. Dundas, B.I., Röndigs, O., Østvær, P.A.: Motivic functors, *Doc. Math.* 8 (2003)
6. Fu, L.: *Etale cohomology theory, nankai tracts in mathematics*, Vol. 13, World Scientific Publishing Co. Pte. Ltd., Hackensack (2011)
7. Geisser, T.: Motivic cohomology over Dedekind rings. *Math. Z.* (4) 248, 773–794 (2004)
8. Gillet, H., Levine, M.: The relative form of Gersten's conjecture over a discrete valuation ring: the smooth case. *J. Pure Appl. Algebra* (1) 46, 59–71 (1987)
9. Gillet, H., Levine, M.: Spectra and symmetric spectra in general model categories. *J. Pure Appl. Algebra* 165(1), 63–127 (2001)
10. Jardine, J.F.: Motivic symmetric spectra. *Doc. Math.* 5, 445–552 (2000)
11. Quillen, D.: Higher algebraic K-theory. I, *Algebraic K-theory, I: Higher K-theories* (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972), 85–147. *Lecture Notes in Math.*, Vol. 341, Springer, Berlin (1973)
12. Schmidt, J., Strunk, F.: Stable \mathbb{A}^1 -connectivity over Dedekind schemes. *Ann. K-Theory* (2) 3, 331–367 (2018)
13. Voevodsky, V., Röndigs, O., Østvær, P.A.: *Voevodsky's Nordfjordeid Lectures: Motivic Homotopy, Motivic Homotopy Theory, Lectures at a Summer School in Nordfjordeid, Norway, August 2002*. Springer, Berlin, Heidelberg, New York (2007)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.