



# An adaptive finite element DtN method for the elastic wave scattering problem

Peijun Li<sup>1</sup> · Xiaokai Yuan<sup>2</sup>

Received: 8 March 2019 / Revised: 19 December 2021 / Accepted: 17 February 2022 /  
Published online: 19 March 2022

© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2022

## Abstract

Consider the scattering of an incident wave by a rigid obstacle, which is immersed in a homogeneous and isotropic elastic medium in two dimensions. Based on a Dirichlet-to-Neumann (DtN) operator, an exact transparent boundary condition is introduced and the scattering problem is formulated as a boundary value problem of the elastic wave equation in a bounded domain. By developing a new duality argument, an a posteriori error estimate is derived for the discrete problem by using the finite element method with the truncated DtN operator. The a posteriori error estimate consists of the finite element approximation error and the truncation error of the DtN operator, where the latter decays exponentially with respect to the truncation parameter. An adaptive finite element algorithm is proposed to solve the elastic obstacle scattering problem, where the truncation parameter is determined through the truncation error and the mesh elements for local refinements are chosen through the finite element discretization error. Numerical experiments are presented to demonstrate the effectiveness of the proposed method.

**Mathematics Subject Classification** 65N12 · 65N15 · 65N30 · 78A45

---

The research of PL is supported in part by the NSF grant DMS-1912704.

---

✉ Peijun Li  
lpeijun@math.purdue.edu  
Xiaokai Yuan  
yuanxk@jlu.edu.cn

<sup>1</sup> Department of Mathematics, Purdue University, West Lafayette, IN 47907, USA

<sup>2</sup> School of Mathematics, Jilin University, Changchun 130012, Jilin, People's Republic of China

## 1 Introduction

A basic problem in classical scattering theory is the scattering of time-harmonic waves by a bounded and impenetrable medium, which is known as the obstacle scattering problem. It has played a crucial role in diverse scientific areas such as radar and sonar, geophysical exploration, medical imaging, and nondestructive testing. Motivated by these significant applications, the obstacle scattering problem has been widely studied for acoustic and electromagnetic waves. Consequently, a great deal of results are available concerning its solution [18, 41, 42]. Recently, the scattering problems for elastic waves have received ever-increasing attention due to the important applications in seismology and geophysics [3, 38, 39]. For instance, they are fundamental to detect the fractures in sedimentary rocks for the production of underground gas and liquids. Compared with acoustic and electromagnetic waves, elastic waves are less studied due to the coexistence of compressional waves and shear waves that have different wavenumbers [16, 36].

The obstacle scattering problem is usually formulated as an exterior boundary value problem imposed in an open domain. The unbounded physical domain needs to be truncated into a bounded computational domain for the convenience of mathematical analysis or numerical computation. Therefore, an appropriate boundary condition is required on the boundary of the truncated domain to avoid artificial wave reflection. Such a boundary condition is called the transparent boundary condition (TBC) or non-reflecting boundary condition. It is one of the important and active subjects in the research area of wave propagation [6, 19–24]. Since Berenger proposed a perfectly matched layer (PML) technique to solve the time-dependent Maxwell equations [7], the research on the PML has undergone a tremendous development due to its effectiveness and simplicity. Various constructions of PML have been proposed and studied for a wide range of scattering problems on acoustic and electromagnetic wave propagation [5, 10, 15, 27, 30, 44]. The basic idea of the PML technique is to surround the domain of interest by a layer of finite thickness fictitious medium that attenuates the waves coming from inside of the computational domain. When the waves reach the outer boundary of the PML region, their values are so small that the homogeneous Dirichlet boundary conditions can be imposed.

A posteriori error estimates are computable quantities which measure the solution errors of discrete problems. They are essential in designing algorithms for mesh modification which aim to equidistribute the computational effort and optimize the computation. The a posteriori error estimates based adaptive finite element methods have the ability of error control and asymptotically optimal approximation property [2]. They have become a class of important numerical tools for solving differential equations, especially for those where the solutions have singularity or multiscale phenomena. Combined with the PML technique, an efficient adaptive finite element method was developed in [11] for solving the two-dimensional diffraction grating problem, where the medium has a one-dimensional periodic structure and the model equation is the two-dimensional Helmholtz equation. It was shown that the a posteriori error estimate consists of the finite element discretization error and the PML truncation error which decays exponentially with respect to the PML parameters such as the thickness of the layer and the medium properties. Due to the superior numerical

performance, the adaptive PML method was quickly extended to solve the two- and three-dimensional obstacle scattering problems [10, 12] and the three-dimensional diffraction grating problem [4], where either the two-dimensional Helmholtz equation or the three-dimensional Maxwell equations were considered. Although the PML method has been developed to solve various elastic wave propagation problems in engineering and geophysics soon after it was introduced [14, 17, 25, 35], the rigorous mathematical studies were only recently done for elastic waves because of the complexity of the model equation [8, 13, 33, 34].

As a viable alternative, the finite element DtN method has been proposed to solve the obstacle scattering problems [29, 32, 40], the diffraction grating problems [31, 45], and the open cavity scattering problem [47], respectively, where the transparent boundary conditions are used to truncate the domains. In this new approach, the layer of artificial medium is not needed to enclose the domain of interest, which makes it different from the PML method. The transparent boundary conditions are based on nonlocal Dirichlet-to-Neumann (DtN) operators and are given as infinite Fourier series. Since the transparent boundary conditions are exact, the artificial boundary can be put as close as possible to the scattering structures, which can reduce the size of the computational domain. Numerically, the infinite series need to be truncated into a sum of finitely many terms by choosing an appropriate truncation parameter  $N$ . It is known that the convergence of the truncated DtN map could be arbitrarily slow to the original DtN map in the operator norm. The a posteriori error analysis of the PML method cannot be applied directly to the DtN method since the DtN map of the truncated PML problem converges exponentially fast to the DtN map of the untruncated PML problem. To overcome this issue, a duality argument had to be developed to obtain the a posteriori error estimate between the solution of the scattering problem and the finite element solution. Comparably, the a posteriori error estimates consist of the finite element discretization error and the DtN truncation error, which decays exponentially with respect to the truncation parameter  $N$ . The numerical experiments demonstrate that the adaptive DtN method has a competitive behavior to the adaptive PML method. Recently, an interesting adaptive finite element method was developed for the diffraction grating problem [48]. The method combines the PML and few-mode DtN truncations so that those Fourier modes which cannot be well absorbed by the PML can pass through the boundary without reflections.

In this paper, we present an adaptive finite element DtN method and carry out its mathematical analysis for the elastic wave scattering problem. The goal is threefold: (1) prove the exponential convergence of the truncated DtN operator; (2) give a complete a posteriori error estimate; (3) develop an effective adaptive finite element algorithm. This paper significantly extends the work on the acoustic scattering problem [29], where the Helmholtz equation was considered. Apparently, the techniques differ greatly from the existing work because of the complicated transparent boundary condition associated with the elastic wave equation.

Specifically, we consider a rigid obstacle which is immersed in a homogeneous and isotropic elastic medium in two dimensions. The Helmholtz decomposition is utilized to formulate the exterior boundary value problem of the elastic wave equation into a coupled exterior boundary value problem of the Helmholtz equation. By using a Dirichlet-to-Neumann (DtN) operator, an exact transparent boundary condition, which

is given as a Fourier series, is introduced to reduce the original scattering problem into a boundary value problem of the elastic wave equation in a bounded domain. The discrete problem is studied by using the finite element method with the truncated DtN operator. Based on the Helmholtz decomposition, a new duality argument is developed to obtain an a posteriori error estimate between the solution of the original scattering problem and the discrete problem. The a posteriori error estimate consists of the finite element approximation error and the truncation error of the DtN operator which is shown to decay exponentially with respect to the truncation parameter. The estimate is used to design the adaptive finite element algorithm to choose elements for refinements and to determine the truncation parameter  $N$ . Since the truncation error decays exponentially with respect to  $N$ , the choice of the truncation parameter  $N$  is not sensitive to the given tolerance. Numerical experiments are presented to demonstrate the effectiveness of the proposed method.

The paper is organized as follows. In Sect. 2, the elastic wave equation is introduced for the scattering by a rigid obstacle; a boundary value problem is formulated by using the transparent boundary condition; the corresponding weak formulation is discussed. In Sect. 3, the discrete problem is considered by using the finite element approximation with the truncated DtN operator. Section 4 is devoted to the a posteriori error analysis and serves as the basis of the adaptive algorithm. In Sect. 5, we discuss the numerical implementation of the adaptive algorithm and present two numerical examples to illustrate the performance of the proposed method. The paper is concluded with some general remarks and directions for future work in Sect. 6.

## 2 Problem formulation


Consider a two-dimensional elastically rigid obstacle  $D$  with Lipschitz continuous boundary  $\partial D$ , as seen in Fig. 1. Denote by  $\nu = (\nu_1, \nu_2)^\top$  and  $\tau = (\tau_1, \tau_2)^\top$  the unit normal and tangent vectors on  $\partial D$ , where  $\tau_1 = \nu_2$  and  $\tau_2 = -\nu_1$ . The exterior domain  $\mathbb{R}^2 \setminus \overline{D}$  is assumed to be filled with a homogeneous and isotropic elastic medium with a unit mass density. Let  $B_R = \{\mathbf{x} = (x, y)^\top \in \mathbb{R}^2 : |\mathbf{x}| < R\}$  and  $B_{\hat{R}} = \{\mathbf{x} \in \mathbb{R}^2 : |\mathbf{x}| < \hat{R}\}$  be the balls with radii  $R$  and  $\hat{R}$ , respectively, where  $R > \hat{R} > 0$ . Denote by  $\partial B_R$  and  $\partial B_{\hat{R}}$  the boundaries of  $B_R$  and  $B_{\hat{R}}$ , respectively. Let  $\hat{R}$  be large enough such that  $\overline{D} \subset B_{\hat{R}} \subset B_R$ . Denote by  $\Omega = B_R \setminus \overline{D}$  the bounded domain where the boundary value problem will be formulated.

Let the obstacle be illuminated by an incident wave  $\mathbf{u}^{\text{inc}}$ , which can be either a point source or a plane wave. The displacement of the total field  $\mathbf{u}$  satisfies the two-dimensional elastic wave equation

$$\mu \Delta \mathbf{u} + (\lambda + \mu) \nabla \nabla \cdot \mathbf{u} + \omega^2 \mathbf{u} = 0 \quad \text{in } \mathbb{R}^2 \setminus \overline{D}, \quad (2.1)$$

where  $\omega > 0$  is the angular frequency and  $\lambda, \mu$  are the Lamé constants satisfying  $\mu > 0, \lambda + \mu > 0$ . Since the obstacle is assumed to be rigid, the displacement of the

**Fig. 1** Schematic of the elastic wave scattering problem



total field vanishes on the boundary of the obstacle, i.e.,

$$\mathbf{u} = 0 \quad \text{on } \partial D. \tag{2.2}$$

The scattered field is defined as  $\mathbf{u}^s = \mathbf{u} - \mathbf{u}^{\text{inc}}$  and it is required to satisfy the Kupradze–Sommerfeld radiation condition

$$\lim_{r \rightarrow \infty} r^{1/2}(\partial_r \mathbf{u}_p^s - i\kappa_1 \mathbf{u}_p^s) = 0, \quad \lim_{r \rightarrow \infty} r^{1/2}(\partial_r \mathbf{u}_s^s - i\kappa_2 \mathbf{u}_s^s) = 0, \quad r = |\mathbf{x}|, \tag{2.3}$$

where

$$\mathbf{u}_p^s = -\frac{1}{\kappa_1^2} \nabla \nabla \cdot \mathbf{u}^s, \quad \mathbf{u}_s^s = \frac{1}{\kappa_2^2} \mathbf{curl} \mathbf{curl} \mathbf{u}^s,$$

are the compressional and shear wave components of  $\mathbf{u}^s$ , respectively. Here

$$\kappa_1 = \frac{\omega}{(\lambda + 2\mu)^{1/2}}, \quad \kappa_2 = \frac{\omega}{\mu^{1/2}}$$

are known as the compressional wavenumber and the shear wavenumber, respectively. Clearly we have  $\kappa_1 < \kappa_2$  since  $\mu > 0, \lambda + \mu > 0$ . Given a vector function  $\mathbf{u} = (u_1, u_2)^\top$  and a scalar function  $u$ , the scalar and vector curl operators are defined by

$$\mathbf{curl} \mathbf{u} = \partial_x u_2 - \partial_y u_1, \quad \mathbf{curl} u = (\partial_y u, -\partial_x u)^\top.$$

For any solution  $\mathbf{u}$  of (2.1), we introduce the Helmholtz decomposition

$$\mathbf{u} = \nabla \phi + \mathbf{curl} \psi, \tag{2.4}$$

where  $\phi, \psi$  are called the scalar potential functions. Substituting (2.4) into (2.1), the corresponding potential functions  $\phi^s, \psi^s$  for the scattered field  $\mathbf{u}^s$  satisfy the Helmholtz equations

$$\Delta \phi^s + \kappa_1^2 \phi^s = 0, \quad \Delta \psi^s + \kappa_2^2 \psi^s = 0 \quad \text{in } \mathbb{R}^2 \setminus \overline{D}. \tag{2.5}$$

Taking the dot product of (2.2) with  $\nu$  and  $\tau$ , respectively, we get

$$\partial_\nu \phi^s - \partial_\tau \psi^s = f_1, \quad \partial_\nu \psi^s + \partial_\tau \phi^s = f_2 \quad \text{on } \partial D, \tag{2.6}$$

where  $f_1 = -\mathbf{u}^{\text{inc}} \cdot \nu$  and  $f_2 = -\mathbf{u}^{\text{inc}} \cdot \tau$ .

In addition, the potential functions  $\phi^s, \psi^s$  satisfy the Sommerfeld radiation conditions

$$\lim_{r \rightarrow \infty} r^{1/2}(\partial_r \phi^s - \kappa_1 \phi^s) = 0, \quad \lim_{r \rightarrow \infty} r^{1/2}(\partial_r \psi^s - \kappa_2 \psi^s) = 0. \tag{2.7}$$

By the Helmholtz decomposition, it can be shown that the boundary value problems (2.1)–(2.3) and (2.5)–(2.7) are equivalent. The result is stated in the following lemma and the proof can be found in [38].

**Lemma 2.1** *Let  $\mathbf{u}^s$  be the scattered field corresponding to the solution of the boundary value problem (2.1)–(2.3). Then  $\phi^s = -\kappa_1^{-2} \nabla \cdot \mathbf{u}^s, \psi^s = \kappa_2^{-2} \text{curl} \mathbf{u}^s$  are the solutions of the coupled boundary value problem (2.5)–(2.7). Conversely, if  $\phi^s, \psi^s$  are the solutions of the boundary value problem (2.5)–(2.7), let  $\mathbf{u}^s = \nabla \phi^s + \text{curl} \psi^s$ , then  $\mathbf{u} = \mathbf{u}^s + \mathbf{u}^{\text{inc}}$  is the solution of the boundary value problem (2.1)–(2.3).*

Denote by  $L^2(\Omega)$  the usual Hilbert space of square integrable functions. Let  $H^1(\Omega)$  be the standard Sobolev space equipped with the norm

$$\|u\|_{H^1(\Omega)} = \left( \|u\|_{L^2(\Omega)}^2 + \|\nabla u\|_{L^2(\Omega)}^2 \right)^{1/2}.$$

Define  $H^1_{\partial D}(\Omega) = \{u \in H^1(\Omega) : u = 0 \text{ on } \partial D\}$ . For any function  $u \in L^2(\partial B_R)$ , it admits the Fourier series expansion

$$u(R, \theta) = \sum_{n \in \mathbb{Z}} \hat{u}_n(R) e^{in\theta}, \quad \hat{u}_n(R) = \frac{1}{2\pi} \int_0^{2\pi} u(R, \theta) e^{-in\theta} d\theta.$$

The trace space  $H^s(\partial B_R), s \in \mathbb{R}$  is defined by

$$H^s(\partial B_R) = \{u \in L^2(\partial B_R) : \|u\|_{H^s(\partial B_R)} < \infty\},$$

where  $H^s(\partial B_R)$  norm is given by

$$\|u\|_{H^s(\partial B_R)} = \left( 2\pi \sum_{n \in \mathbb{Z}} (1 + n^2)^s |\hat{u}_n(R)|^2 \right)^{1/2}.$$

Let  $\mathbf{H}^1(\Omega) = H^1(\Omega)^2$  and  $\mathbf{H}^1_{\partial D}(\Omega) = H^1_{\partial D}(\Omega)^2$  be the Cartesian product spaces equipped with the corresponding 2-norms of  $H^1(\Omega)$  and  $H^1_{\partial D}(\Omega)$ , respectively. Throughout the paper, we take the notation of  $a \lesssim b$  to stand for  $a \leq Cb$ , where  $C$  is a positive constant whose value is not required but should be clear from the context.

The elastic wave scattering problem (2.2)–(2.3) is formulated in the open domain  $\mathbb{R}^2 \setminus \overline{D}$ , which needs to be truncated into the bounded domain  $\Omega$ . An appropriate boundary condition is required on  $\partial B_R$ .

Define a boundary operator on  $\partial B_R$  as follows

$$\mathcal{B}v = \mu \partial_r v + (\lambda + \mu) \nabla \cdot v e_r,$$

where  $e_r$  is the unit outward normal vector on  $\partial B_R$ . It is shown in [38] that the scattered field  $\mathbf{u}^s$  satisfies the transparent boundary condition on  $\partial B_R$ :

$$\mathcal{B}\mathbf{u}^s = (\mathcal{T}\mathbf{u}^s)(R, \theta) := \sum_{n \in \mathbb{Z}} M_n \mathbf{u}_n^s(R) e^{in\theta}, \quad \mathbf{u}^s(R, \theta) = \sum_{n \in \mathbb{Z}} \mathbf{u}_n^s(R) e^{in\theta}, \quad (2.8)$$

where  $\mathcal{T}$  is called the Dirichlet-to-Neumann (DtN) operator and  $M_n$  is a  $2 \times 2$  matrix whose entries are given in (A.4) and (A.6). By a simple calculation, the total field  $\mathbf{u}$  satisfies the transparent boundary condition

$$\mathcal{B}\mathbf{u} = \mathcal{T}\mathbf{u} + \mathbf{g} \quad \text{on } \partial B_R, \quad (2.9)$$

where  $\mathbf{g} := \mathcal{B}\mathbf{u}^{\text{inc}} - \mathcal{T}\mathbf{u}^{\text{inc}}$ .

Based on the transparent boundary condition (2.9), the variational problem for (2.1)–(2.3) is to find  $\mathbf{u} \in \mathbf{H}^1_{\partial D}(\Omega)$  such that

$$b(\mathbf{u}, \mathbf{v}) = \int_{\partial B_R} \mathbf{g} \cdot \bar{\mathbf{v}} ds \quad \forall \mathbf{v} \in \mathbf{H}^1_{\partial D}(\Omega), \quad (2.10)$$

where the sesquilinear form  $b : \mathbf{H}^1_{\partial D}(\Omega) \times \mathbf{H}^1_{\partial D}(\Omega) \rightarrow \mathbb{C}$  is defined as

$$\begin{aligned} b(\mathbf{u}, \mathbf{v}) &= \mu \int_{\Omega} \nabla \mathbf{u} : \nabla \bar{\mathbf{v}} dx + (\lambda + \mu) \int_{\Omega} (\nabla \cdot \mathbf{u}) (\nabla \cdot \bar{\mathbf{v}}) dx \\ &\quad - \omega^2 \int_{\Omega} \mathbf{u} \cdot \bar{\mathbf{v}} dx - \int_{\partial B_R} \mathcal{T}\mathbf{u} \cdot \bar{\mathbf{v}} ds. \end{aligned} \quad (2.11)$$

Here  $A : B = \text{tr}(AB^T)$  is the Frobenius inner product of square matrices  $A$  and  $B$ .

Following [38], we may show that the variational problem (2.10) has a unique weak solution  $\mathbf{u} \in \mathbf{H}^1_{\partial D}(\Omega)$  for any frequency  $\omega$  and the solution satisfies the estimate

$$\|\mathbf{u}\|_{\mathbf{H}^1(\Omega)} \lesssim \|\mathbf{g}\|_{\mathbf{H}^{-1/2}(\partial B_R)} \lesssim \|\mathbf{u}^{\text{inc}}\|_{\mathbf{H}^1(\Omega)}. \quad (2.12)$$

It follows from the general theory in [1] that there exists a constant  $\gamma > 0$  such that the following inf-sup condition holds

$$\sup_{0 \neq \mathbf{v} \in \mathbf{H}^1_{\partial D}(\Omega)} \frac{|b(\mathbf{u}, \mathbf{v})|}{\|\mathbf{v}\|_{\mathbf{H}^1(\Omega)}} \geq \gamma \|\mathbf{u}\|_{\mathbf{H}^1(\Omega)} \quad \forall \mathbf{u} \in \mathbf{H}^1_{\partial D}(\Omega).$$

### 3 The discrete problem

In the variational problem (2.10), the DtN operator  $\mathcal{T}$  is given as an infinite series. In computation, the infinite series needs to be truncated into a finite sum. Given a sufficiently large  $N$ , we define the truncated DtN operator

$$\mathcal{T}_N \mathbf{u} = \sum_{|n| \leq N} M_n \mathbf{u}_n(R) e^{in\theta}. \tag{3.1}$$

Hence  $\mathbf{g}$  also needs to be approximated as  $\mathbf{g}_N = \mathcal{B}\mathbf{u}^{\text{inc}} - \mathcal{T}_N \mathbf{u}^{\text{inc}}$ . Using (3.1), we have the truncated problem: find  $\mathbf{u}_N \in \mathbf{H}^1_{\partial D}(\Omega)$  such that

$$b_N(\mathbf{u}_N, \mathbf{v}) = \int_{\partial B_R} \mathbf{g}_N \cdot \bar{\mathbf{v}} \, ds \quad \forall \mathbf{v} \in \mathbf{H}^1_{\partial D}(\Omega), \tag{3.2}$$

where the sesquilinear form  $b_N : \mathbf{H}^1_{\partial D}(\Omega) \times \mathbf{H}^1_{\partial D}(\Omega) \rightarrow \mathbb{C}$  is defined as

$$\begin{aligned} b_N(\mathbf{u}, \mathbf{v}) &= \mu \int_{\Omega} \nabla \mathbf{u} : \nabla \bar{\mathbf{v}} \, dx + (\lambda + \mu) \int_{\Omega} (\nabla \cdot \mathbf{u}) (\nabla \cdot \bar{\mathbf{v}}) \, dx \\ &\quad - \omega^2 \int_{\Omega} \mathbf{u} \cdot \bar{\mathbf{v}} \, dx - \int_{\partial B_R} \mathcal{T}_N \mathbf{u} \cdot \bar{\mathbf{v}} \, ds. \end{aligned} \tag{3.3}$$

Let us consider the discrete problem of (2.10) by using the finite element approximation. Let  $\mathcal{M}_h$  be a regular triangulation of  $\Omega$ , where  $h$  denotes the maximum diameter of all the elements in  $\mathcal{M}_h$ . For simplicity, we assume that the boundary  $\partial D$  is polygonal and ignore the approximation error of the boundary  $\partial B_R$ , which allows us to focus on deducing the a posteriori error estimate. Thus any edge  $e \in \mathcal{M}_h$  is a subset of  $\partial\Omega$  if it has two boundary vertices.

Let  $\Omega_h := \cup_{K \in \mathcal{M}_h} K$  and  $\tilde{\mathbf{V}}_h \subset \mathbf{H}^1(\Omega_h)$  be a conforming finite element space, i.e.,

$$\tilde{\mathbf{V}}_h := \left\{ \mathbf{v} \in C(\overline{\Omega_h})^2 : \mathbf{v}|_K \in P_m(K)^2 \text{ for any } K \in \mathcal{M}_h \right\},$$

where  $m$  is a positive integer and  $P_m(K)$  denotes the set of all polynomials of degree no more than  $m$ . Introduce an isoparametric-equivalent finite element space

$$\mathbf{V}_h := \left\{ \mathbf{v}(\mathbf{F}^{-1}(\mathbf{x})) : \mathbf{x} \in \mathbf{F}(\Omega_h), \mathbf{v} \in \tilde{\mathbf{V}}_h \right\},$$

where  $\mathbf{F} \in \tilde{\mathbf{V}}_h$  is a one-to-one continuous mapping constructed in [37]. We refer to [9, 37] for more details about the construction and properties of  $\mathbf{F}$ . The finite element approximation to the variational problem (2.10) is to find  $\mathbf{u}^h \in \mathbf{V}_{h,\partial D}$  such that

$$b(\mathbf{u}^h, \mathbf{v}^h) = \int_{\partial B_R} \mathbf{g} \cdot \bar{\mathbf{v}}^h \, ds \quad \forall \mathbf{v}^h \in \mathbf{V}_{h,\partial D}, \tag{3.4}$$



where  $V_{h,\partial D} = \{v \in V_h : v = 0 \text{ on } \partial D\}$ .

The truncated finite element approximation to the variational problem (3.2) is to find  $u_N^h \in V_{h,\partial D}$  such that

$$b_N(u_N^h, v^h) = \int_{\partial B_R} g_N \cdot \overline{v^h} ds \quad \forall v^h \in V_{h,\partial D}. \tag{3.5}$$

Following the argument in [28, 38], we may show that for sufficiently large  $N$  the variational problem (3.2) is well-posed. Moreover, for sufficiently small  $h$ , the discrete inf-sup condition of the sesquilinear form  $b_N$  may be established by following the approach in [43]. Based on the general theory in [1], the truncated variational problem (3.5) can be shown to have a unique solution  $u_N^h \in V_h$ . The details are omitted since our focus is the a posteriori error estimate.

### 4 The a posteriori error analysis

For any triangular element  $K \in \mathcal{M}_h$ , we denote by  $h_K$  its diameter. Let  $\mathcal{B}_h$  denote the set of all the edges of  $K$ . For any edge  $e \in \mathcal{B}_h$ , let  $h_e$  be its length. For any interior edge  $e$  which is the common side of triangular elements  $K_1, K_2 \in \mathcal{M}_h$ , we define the jump residual across  $e$  as

$$J_e = - \left( \mu \nabla u_N^h|_{K_1} \cdot \nu_1 + (\lambda + \mu) \nabla \cdot u_N^h|_{K_1} \nu_1 + \mu \nabla u_N^h|_{K_2} \cdot \nu_2 + (\lambda + \mu) \nabla \cdot u_N^h|_{K_2} \nu_2 \right),$$

where  $\nu_j$  is the unit outward normal vector on the boundary of  $K_j, j = 1, 2$ . For any boundary edge  $e \subset \partial B_R$ , we define the jump residual

$$J_e = 2 \left( \mathcal{R} u_N^h + g_N - \mu (\nabla u_N^h \cdot e_r) - (\lambda + \mu) (\nabla \cdot u_N^h) e_r \right).$$

For any triangular element  $K \in \mathcal{M}_h$ , denote by  $\eta_K$  the local error estimator which is given by

$$\eta_K = h_K \| \mathcal{R} u_N^h \|_{L^2(K)} + \left( \frac{1}{2} \sum_{e \in \partial K} h_e \| J_e \|_{L^2(e)}^2 \right)^{1/2},$$

where  $\mathcal{R}$  is the residual operator defined by

$$\mathcal{R} u = \mu \Delta u + (\lambda + \mu) \nabla (\nabla \cdot u) + \omega^2 u.$$

For convenience, we introduce a weighted norm  $\| \cdot \|_{H^1(\Omega)}$  which is given by

$$\| u \|_{H^1(\Omega)}^2 = \mu \int_{\Omega} |\nabla u|^2 dx + (\lambda + \mu) \int_{\Omega} |\nabla \cdot u|^2 dx + \omega^2 \int_{\Omega} |u|^2 dx. \tag{4.1}$$

It can be verified for any  $\mathbf{u} \in \mathbf{H}^1(\Omega)$  that

$$\min(\mu, \omega^2) \|\mathbf{u}\|_{\mathbf{H}^1(\Omega)}^2 \leq \|\mathbf{u}\|_{\mathbf{H}^1(\Omega)}^2 \leq \max(2\lambda + 3\mu, \omega^2) \|\mathbf{u}\|_{\mathbf{H}^1(\Omega)}^2, \tag{4.2}$$

which implies that the two norms  $\|\cdot\|_{\mathbf{H}^1(\Omega)}$  and  $\|\!\| \cdot \|\!\|_{\mathbf{H}^1(\Omega)}$  are equivalent.

Now we state the main result of this paper.

**Theorem 4.1** *Let  $\mathbf{u}$  and  $\mathbf{u}_N^h$  be the solutions of the variational problems (2.10) and (3.5), respectively. Then for sufficiently large  $N$ , the following a posteriori error estimate holds:*

$$\|\mathbf{u} - \mathbf{u}_N^h\|_{\mathbf{H}^1(\Omega)} \lesssim \left( \sum_{K \in \mathcal{M}_h} \eta_K^2 \right)^{1/2} + N \left( \frac{\hat{R}}{R} \right)^N \|\mathbf{u}^{\text{inc}}\|_{\mathbf{H}^1(\Omega)}.$$

We point out that the a posteriori error estimate consists of two parts: the first part arises from the finite element discretization error; and the second part comes from the truncation error of the DtN operator. Apparently, the DtN truncation error decreases exponentially with respect to  $N$  since  $\hat{R} < R$ .

In the rest of the paper, we shall prove the a posteriori error estimate in Theorem 4.1. First, we present the result on trace regularity for functions in  $H^1(\Omega)$ . The proof can be found in [29].

**Lemma 4.2** *For any  $u \in H^1(\Omega)$ , the following estimates hold:*

$$\|u\|_{H^{1/2}(\partial B_R)} \lesssim \|u\|_{H^1(\Omega)}, \quad \|u\|_{H^{1/2}(\partial B_{\hat{R}})} \lesssim \|u\|_{H^1(\Omega)}.$$

Let  $\boldsymbol{\xi} = \mathbf{u} - \mathbf{u}_N^h$ , where  $\mathbf{u}$  and  $\mathbf{u}_N^h$  are the solutions of the problems (2.10) and (3.5), respectively. Combining (4.1), (2.11), and (3.3), we have from straightforward calculations that

$$\begin{aligned} \|\!\| \boldsymbol{\xi} \|\!\|_{\mathbf{H}^1(\Omega)}^2 &= \mu \int_{\Omega} \nabla \boldsymbol{\xi} : \nabla \bar{\boldsymbol{\xi}} \, d\mathbf{x} + (\lambda + \mu) \int_{\Omega} (\nabla \cdot \boldsymbol{\xi}) (\nabla \cdot \bar{\boldsymbol{\xi}}) \, d\mathbf{x} + \omega^2 \int_{\Omega} \boldsymbol{\xi} \cdot \bar{\boldsymbol{\xi}} \, d\mathbf{x} \\ &= \Re b(\boldsymbol{\xi}, \boldsymbol{\xi}) + 2\omega^2 \int_{\Omega} |\boldsymbol{\xi}|^2 \, d\mathbf{x} + \Re \int_{\partial B_R} \mathcal{T} \boldsymbol{\xi} \cdot \bar{\boldsymbol{\xi}} \, ds \\ &= \Re b(\boldsymbol{\xi}, \boldsymbol{\xi}) + \Re \int_{\partial B_R} (\mathcal{T} - \mathcal{T}_N) \boldsymbol{\xi} \cdot \bar{\boldsymbol{\xi}} \, ds + 2\omega^2 \int_{\Omega} |\boldsymbol{\xi}|^2 \, d\mathbf{x} \\ &\quad + \Re \int_{\partial B_R} \mathcal{T}_N \boldsymbol{\xi} \cdot \bar{\boldsymbol{\xi}} \, ds, \end{aligned} \tag{4.3}$$

which is the error representation formula.

In the following, we discuss the four terms in (4.3) one by one. Lemma 4.5 presents the a posteriori error estimate for the truncation of the DtN operator on the scattered field  $\mathbf{u}^s$ . Lemma 4.6 gives the a posteriori error estimate for the total field  $\mathbf{u}$  on both of the finite element approximation and DtN operator truncation.

**Lemma 4.3** *Let  $0 < \kappa_1 < \kappa_2$  and  $0 < \hat{R} < R$ . For sufficiently large  $|n|$ , the following estimate holds for  $j = 1, 2$ :*

$$\left| \frac{H_n^{(j)}(\kappa_1 R)}{H_n^{(j)}(\kappa_1 \hat{R})} - \frac{H_n^{(j)}(\kappa_2 R)}{H_n^{(j)}(\kappa_2 \hat{R})} \right| \leq \frac{\kappa_2(\kappa_2 - \kappa_1)}{|n| - 1} (R^2 - \hat{R}^2) \left(\frac{\hat{R}}{R}\right)^{|n|},$$

where  $H_n^{(1)}$  and  $H_n^{(2)}$  are the Hankel functions of the first and second kind with order  $n$ , respectively.

**Proof** Since the Hankel functions of the first and second kind are complex conjugate to each other, we only need to show the proof for the Hankel function of the first kind.

Let  $J_n$  and  $Y_n$  be the Bessel functions of the first and second kind with order  $n$ , respectively. Since  $J_{-n}(z) = (-1)^n J_n(z)$ ,  $Y_{-n}(z) = (-1)^n Y_n(z)$ , it suffices to show the result for positive  $n$ . For a fixed  $z > 0$ , they admit the following asymptotic properties [46]:

$$J_n(z) \sim \frac{1}{\sqrt{2\pi n}} \left(\frac{ez}{2n}\right)^n, \quad Y_n(z) \sim -\sqrt{\frac{2}{\pi n}} \left(\frac{ez}{2n}\right)^{-n}, \quad n \rightarrow \infty. \tag{4.4}$$

Define  $S(z) = J_n(z)/Y_n(z)$ . A simple calculation yields

$$\begin{aligned} \frac{H_n^{(1)}(zR)}{H_n^{(1)}(z\hat{R})} &= \frac{J_n(zR) + iY_n(zR)}{J_n(z\hat{R}) + iY_n(z\hat{R})} = \frac{Y_n(zR)}{Y_n(z\hat{R})} \frac{1 - i\frac{J_n(zR)}{Y_n(zR)}}{1 - i\frac{J_n(z\hat{R})}{Y_n(z\hat{R})}} \\ &= \frac{Y_n(zR)}{Y_n(z\hat{R})} \frac{1 - iS_n(zR)}{1 - iS_n(z\hat{R})} = \frac{Y_n(zR)}{Y_n(z\hat{R})} + i\frac{Y_n(zR)}{Y_n(z\hat{R})} \frac{S_n(z\hat{R}) - S_n(zR)}{1 - iS_n(z\hat{R})}. \end{aligned} \tag{4.5}$$

By (4.4)–(4.5), we have

$$S_n(z) = \frac{J_n(z)}{Y_n(z)} \sim \frac{\frac{1}{\sqrt{2\pi n}} \left(\frac{ez}{2n}\right)^n}{-\sqrt{\frac{2}{\pi n}} \left(\frac{ez}{2n}\right)^{-n}} = -\frac{1}{2} \left(\frac{ez}{2n}\right)^{2n}, \quad n \rightarrow \infty.$$

and

$$\begin{aligned} \left| \frac{H_n^{(1)}(\kappa_1 R)}{H_n^{(1)}(\kappa_1 \hat{R})} - \frac{H_n^{(1)}(\kappa_2 R)}{H_n^{(1)}(\kappa_2 \hat{R})} \right| &\leq \left| \frac{Y_n(\kappa_1 R)}{Y_n(\kappa_1 \hat{R})} - \frac{Y_n(\kappa_2 R)}{Y_n(\kappa_2 \hat{R})} \right| + \left| \frac{Y_n(\kappa_1 R)}{Y_n(\kappa_1 \hat{R})} \frac{S_n(\kappa_1 \hat{R})}{1 - iS_n(\kappa_1 \hat{R})} \right| \\ &+ \left| \frac{Y_n(\kappa_1 R)}{Y_n(\kappa_1 \hat{R})} \frac{S_n(\kappa_1 R)}{1 - iS_n(\kappa_1 \hat{R})} \right| + \left| \frac{Y_n(\kappa_2 R)}{Y_n(\kappa_2 \hat{R})} \frac{S_n(\kappa_2 \hat{R})}{1 - iS_n(\kappa_2 \hat{R})} \right| + \left| \frac{Y_n(\kappa_2 R)}{Y_n(\kappa_2 \hat{R})} \frac{S_n(\kappa_2 R)}{1 - iS_n(\kappa_2 \hat{R})} \right|. \end{aligned}$$

It is easy to verify that

$$\left| \frac{S_n(zR)}{1 - iS_n(z\hat{R})} \right| \leq \left( \frac{e z R}{2n} \right)^{2n}, \quad \left| \frac{S_n(z\hat{R})}{1 - iS_n(z\hat{R})} \right| \leq \left( \frac{e z \hat{R}}{2n} \right)^{2n}$$

and

$$\frac{RzY'_n(zR)}{Y_n(zR)} \sim \frac{z^2 R^2}{2(n-1)} - n, \quad \frac{Y_n(zR)}{Y_n(z\hat{R})} \sim \left( \frac{\hat{R}}{R} \right)^{|n|}.$$

Combining the above estimates, we have for  $R > \hat{R}$  and  $\kappa_2 > \kappa_1$  that

$$\begin{aligned} & \left| \frac{Y_n(\kappa_1 R)}{Y_n(\kappa_1 \hat{R})} \frac{S_n(\kappa_1 \hat{R})}{1 - iS_n(\kappa_1 \hat{R})} \right| + \left| \frac{Y_n(\kappa_1 R)}{Y_n(\kappa_1 \hat{R})} \frac{S_n(\kappa_1 R)}{1 - iS_n(\kappa_1 \hat{R})} \right| + \left| \frac{Y_n(\kappa_2 R)}{Y_n(\kappa_2 \hat{R})} \frac{S_n(\kappa_2 \hat{R})}{1 - iS_n(\kappa_2 \hat{R})} \right| \\ & + \left| \frac{Y_n(\kappa_2 R)}{Y_n(\kappa_2 \hat{R})} \frac{S_n(\kappa_2 R)}{1 - iS_n(\kappa_2 \hat{R})} \right| \leq 2 \left( \frac{e \kappa_2 R}{2n} \right)^{2n} \left( \left| \frac{Y_n(\kappa_1 R)}{Y_n(\kappa_1 \hat{R})} \right| + \left| \frac{Y_n(\kappa_2 R)}{Y_n(\kappa_2 \hat{R})} \right| \right). \end{aligned}$$

Define  $F(z) = Y_n(zR)/Y_n(z\hat{R})$ . By the mean value theorem, there exists  $\xi \in (\kappa_1, \kappa_2)$  such that

$$\begin{aligned} F(\kappa_1) - F(\kappa_2) &= F'(\xi)(\kappa_1 - \kappa_2) \\ &= \frac{RY'_n(\xi R)Y_n(\xi \hat{R}) - \hat{R}Y_n(\xi R)Y'_n(\xi \hat{R})}{Y_n(\xi \hat{R})^2}(\kappa_1 - \kappa_2) \\ &= \left( \frac{R\xi Y'_n(\xi R)}{Y_n(\xi R)} - \frac{\hat{R}\xi Y'_n(\xi \hat{R})}{Y_n(\xi \hat{R})} \right) \frac{Y_n(\xi R)}{Y_n(\xi \hat{R})} \frac{\kappa_1 - \kappa_2}{\xi} \\ &\sim \frac{\xi(\kappa_1 - \kappa_2)}{2(n-1)} (R^2 - \hat{R}^2) \frac{Y_n(\xi R)}{Y_n(\xi \hat{R})}. \end{aligned}$$

Therefore,

$$\begin{aligned} \left| \frac{H_n^{(1)}(\kappa_1 R)}{H_n^{(1)}(\kappa_1 \hat{R})} - \frac{H_n^{(1)}(\kappa_2 R)}{H_n^{(1)}(\kappa_2 \hat{R})} \right| &\leq \left| \frac{\xi(\kappa_1 - \kappa_2)}{2(n-1)} \right| \left| (R^2 - \hat{R}^2) \frac{Y_n(\xi R)}{Y_n(\xi \hat{R})} \right| \\ &\quad + 2 \left( \frac{e \kappa_2 R}{2n} \right)^{2n} \left( \left| \frac{Y_n(\kappa_1 R)}{Y_n(\kappa_1 \hat{R})} \right| + \left| \frac{Y_n(\kappa_2 R)}{Y_n(\kappa_2 \hat{R})} \right| \right) \\ &\leq \frac{\kappa_2(\kappa_2 - \kappa_1)}{|n| - 1} (R^2 - \hat{R}^2) \left( \frac{\hat{R}}{R} \right)^{|n|}, \end{aligned}$$

which completes the proof. □

**Remark 4.4** In the proof of Lemma 4.3, we use the asymptotic properties of the Bessel functions (4.4) for the case when  $z$  is fixed and  $n \rightarrow \infty$ . It is worth mentioning that the truncation number should be increased with respect to  $\kappa_2 R$  in order to keep the term  $Y_n(z)$  dominating in  $H_n^{(1)}(z)$ , which is a technique issue. The result may be improved by making a more sophisticated analysis on the Bessel functions.

**Lemma 4.5** Let  $\mathbf{u}^s \in \mathbf{H}^1(\Omega)$  be the scattered field corresponding to the solution of the variational problem (2.10). For any  $\mathbf{v} \in \mathbf{H}^1(\Omega)$ , the following estimate holds:

$$\left| \int_{\partial B_R} (\mathcal{T} - \mathcal{T}_N) \mathbf{u}^s \cdot \bar{\mathbf{v}} \, ds \right| \lesssim N \left( \frac{\hat{R}}{R} \right)^N \|\mathbf{u}^{\text{inc}}\|_{\mathbf{H}^1(\Omega)} \|\mathbf{v}\|_{\mathbf{H}^1(\Omega)}.$$

**Proof** Recalling the Helmholtz decomposition  $\mathbf{u}^s = \nabla \phi^s + \text{curl} \psi^s$ , we have from the Fourier series expansions in (A.1) that

$$\phi_n^s(R) = \frac{H_n^{(1)}(\kappa_1 R)}{H_n^{(1)}(\kappa_1 \hat{R})} \phi_n^s(\hat{R}), \quad \psi_n^s(R) = \frac{H_n^{(1)}(\kappa_2 R)}{H_n^{(1)}(\kappa_2 \hat{R})} \psi_n^s(\hat{R}).$$

Comparing the Fourier coefficients of  $\mathbf{u}$  and  $\phi, \psi$  in the Helmholtz decomposition gives

$$\begin{aligned} \mathbf{u}_n^s(R) &= \begin{bmatrix} \alpha_{1n}(R) & \frac{in}{R} \\ \frac{in}{R} & -\alpha_{2n}(R) \end{bmatrix} \begin{bmatrix} \phi_n^s(R) \\ \psi_n^s(R) \end{bmatrix} \\ &= \begin{bmatrix} \alpha_{1n}(R) & \frac{in}{R} \\ \frac{in}{R} & -\alpha_{2n}(R) \end{bmatrix} \begin{bmatrix} \frac{H_n^{(1)}(\kappa_1 R)}{H_n^{(1)}(\kappa_1 \hat{R})} & 0 \\ 0 & \frac{H_n^{(1)}(\kappa_2 R)}{H_n^{(1)}(\kappa_2 \hat{R})} \end{bmatrix} \begin{bmatrix} \phi_n^s(\hat{R}) \\ \psi_n^s(\hat{R}) \end{bmatrix} \\ &= \begin{bmatrix} \alpha_{1n}(R) & \frac{in}{R} \\ \frac{in}{R} & -\alpha_{2n}(R) \end{bmatrix} \begin{bmatrix} \frac{H_n^{(1)}(\kappa_1 R)}{H_n^{(1)}(\kappa_1 \hat{R})} & 0 \\ 0 & \frac{H_n^{(1)}(\kappa_2 R)}{H_n^{(1)}(\kappa_2 \hat{R})} \end{bmatrix} \begin{bmatrix} -\alpha_{2n}(\hat{R}) & -\frac{in}{\hat{R}} \\ -\frac{in}{\hat{R}} & \alpha_{1n}(\hat{R}) \end{bmatrix} \frac{\mathbf{u}_n^s(\hat{R})}{\Lambda_n(\hat{R})} \\ &= -\frac{1}{\Lambda_n(\hat{R})} \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \mathbf{u}_n^s(\hat{R}), \end{aligned}$$

where  $\alpha_{jn}, \Lambda_n$  is given in (A.5) and

$$\begin{aligned} A_{11} &= \frac{H_n^{(1)}(\kappa_1 R)}{H_n^{(1)}(\kappa_1 \hat{R})} \alpha_{1n}(R) \alpha_{2n}(\hat{R}) - \frac{n^2}{R \hat{R}} \frac{H_n^{(1)}(\kappa_2 R)}{H_n^{(1)}(\kappa_2 \hat{R})}, \\ A_{12} &= \frac{H_n^{(1)}(\kappa_1 R)}{H_n^{(1)}(\kappa_1 \hat{R})} \alpha_{1n}(R) \frac{in}{\hat{R}} - \frac{in}{R} \alpha_{1n}(\hat{R}) \frac{H_n^{(1)}(\kappa_2 R)}{H_n^{(1)}(\kappa_2 \hat{R})}, \\ A_{21} &= \frac{H_n^{(1)}(\kappa_1 R)}{H_n^{(1)}(\kappa_1 \hat{R})} \alpha_{2n}(\hat{R}) \frac{in}{R} - \frac{in}{\hat{R}} \alpha_{2n}(R) \frac{H_n^{(1)}(\kappa_2 R)}{H_n^{(1)}(\kappa_2 \hat{R})}, \\ A_{22} &= \frac{H_n^{(1)}(\kappa_2 R)}{H_n^{(1)}(\kappa_2 \hat{R})} \alpha_{1n}(\hat{R}) \alpha_{2n}(R) - \frac{n^2}{R \hat{R}} \frac{H_n^{(1)}(\kappa_1 R)}{H_n^{(1)}(\kappa_1 \hat{R})}. \end{aligned}$$

By Lemma 4.3 and (A.7), we have

$$\begin{aligned}
 |A_{11}| &\leq \left| \frac{H_n^{(1)}(\kappa_1 R)}{H_n^{(1)}(\kappa_1 \hat{R})} \right| \left| \alpha_{1n}(R) \alpha_{2n}(\hat{R}) - \frac{n^2}{R \hat{R}} \right| + \frac{n^2}{R \hat{R}} \left| \frac{H_n^{(1)}(\kappa_1 R)}{H_n^{(1)}(\kappa_1 \hat{R})} - \frac{H_n^{(1)}(\kappa_2 R)}{H_n^{(1)}(\kappa_2 \hat{R})} \right| \\
 &\leq \left( \frac{\hat{R}}{R} \right)^{|n|} \left| \left( \frac{\kappa_1^2 R}{2(n-1)} - \frac{n}{R} \right) \left( \frac{\kappa_2^2 \hat{R}}{2(n-1)} - \frac{n}{\hat{R}} \right) - \frac{n^2}{R \hat{R}} \right| \\
 &\quad + \frac{n^2}{R \hat{R}} \frac{\kappa_2(\kappa_2 - \kappa_1)}{|n| - 1} (R^2 - \hat{R}^2) \left( \frac{\hat{R}}{R} \right)^{|n|} \\
 &\leq \kappa_2(\kappa_2 - \kappa_1) (R^2 - \hat{R}^2) \frac{2|n|}{R \hat{R}} \left( \frac{\hat{R}}{R} \right)^{|n|} \lesssim |n| \left( \frac{\hat{R}}{R} \right)^{|n|},
 \end{aligned}$$

Similarly, it can be shown that

$$|A_{ij}| \lesssim |n| \left( \frac{\hat{R}}{R} \right)^{|n|}, \quad i, j = 1, 2.$$

The details are omitted for brevity.

Combining the above estimates and Lemma A.1, we obtain

$$|\mathbf{u}_n^s(R)| \lesssim |n| \left( \frac{\hat{R}}{R} \right)^{|n|} |\mathbf{u}_n^s(\hat{R})|. \tag{4.6}$$

It follows from (A.6)–(A.7) that

$$\begin{aligned}
 &\left| \int_{\partial B_R} (\mathcal{T} - \mathcal{T}_N) \mathbf{u}^s \cdot \bar{\mathbf{v}} \, ds \right| \\
 &= \left| 2\pi R \sum_{|n|>N} \left\{ \left[ \left( -\frac{\mu}{R} + \frac{\omega^2}{\Lambda_n} \alpha_{2n}(R) \right) u_n^{s,r}(R) + in \left( -\frac{\mu}{R} + \frac{\omega^2}{\Lambda_n} \frac{1}{R} \right) u_n^{s,\theta}(R) \right] \overline{v_n^r(R)} \right. \right. \\
 &\quad \left. \left. + \left[ -in \left( -\frac{\mu}{R} + \frac{\omega^2}{\Lambda_n} \frac{1}{R} \right) u_n^{s,r}(R) + \left( -\frac{\mu}{R} + \frac{\omega^2}{\Lambda_n} \alpha_{1n}(R) \right) u_n^{s,\theta}(R) \right] \overline{v_n^\theta(R)} \right\} \right| \\
 &\lesssim \sum_{|n|>N} \left[ \sqrt{|n|} (|u_n^{s,r}(R)| + |u_n^{s,\theta}(R)|) \right] \left[ \sqrt{|n|} (|\overline{v_n^r(R)}| + |\overline{v_n^\theta(R)}|) \right] \\
 &\lesssim \left( \sum_{|n|>N} \left[ |n| (|u_n^{s,r}(R)|^2 + |u_n^{s,\theta}(R)|^2) \right] \right)^{1/2} \\
 &\quad \times \left( \sum_{|n|>N} \left[ |n| (|\overline{v_n^r(R)}|^2 + |\overline{v_n^\theta(R)}|^2) \right] \right)^{1/2}. \tag{4.7}
 \end{aligned}$$

Substituting (2.12) and (4.6) into (4.7) and using Lemma 4.2, we obtain

$$\begin{aligned} & \left| \int_{\partial B_R} (\mathcal{T} - \mathcal{T}_N) \mathbf{u}^s \cdot \bar{\mathbf{v}} \, ds \right| \\ & \lesssim \left( \sum_{|n|>N} \left[ |n|^3 \left( \frac{\hat{R}}{R} \right)^{2|n|} (|u_n^{s,r}(\hat{R})|^2 + |u_n^{s,\theta}(\hat{R})|^2) \right] \right)^{1/2} \|\mathbf{v}\|_{\mathbf{H}^{1/2}(\partial B_R)} \\ & \lesssim \max_{|n|>N} \left( |n| \left( \frac{\hat{R}}{R} \right)^{|n|} \right) \|\mathbf{u}^s\|_{\mathbf{H}^{1/2}(\partial B_{\hat{R}})} \|\mathbf{v}\|_{\mathbf{H}^{1/2}(\partial B_R)} \\ & \lesssim \max_{|n|>N} \left( |n| \left( \frac{\hat{R}}{R} \right)^{|n|} \right) \|\mathbf{u}^s\|_{\mathbf{H}^1(\Omega)} \|\mathbf{v}\|_{\mathbf{H}^1(\Omega)} \\ & \lesssim \max_{|n|>N} \left( |n| \left( \frac{\hat{R}}{R} \right)^{|n|} \right) \|\mathbf{u}^{\text{inc}}\|_{\mathbf{H}^1(\Omega)} \|\mathbf{v}\|_{\mathbf{H}^1(\Omega)}, \end{aligned}$$

which completes the proof by noting that  $|n| \left( \frac{\hat{R}}{R} \right)^{|n|}$  is decreasing for sufficiently large  $|n|$ . □

In Lemma 4.5, it is shown that the truncation error of the DtN operator on the scattered field decay exponentially with respect to the truncation parameter  $N$ . The result implies that  $N$  can be small in practice.

**Lemma 4.6** *Let  $\mathbf{v}$  be any function in  $\mathbf{H}^1_{\partial D}(\Omega)$ , we have*

$$\begin{aligned} & \left| b(\boldsymbol{\xi}, \mathbf{v}) + \int_{\partial B_R} (\mathcal{T} - \mathcal{T}_N) \boldsymbol{\xi} \cdot \bar{\mathbf{v}} \, ds \right| \\ & \lesssim \left( \left( \sum_{K \in \mathcal{M}_h} \eta_K^2 \right)^{1/2} + N \left( \frac{\hat{R}}{R} \right)^N \|\mathbf{u}^{\text{inc}}\|_{\mathbf{H}^1(\Omega)} \right) \|\mathbf{v}\|_{\mathbf{H}^1(\Omega)}. \end{aligned}$$

**Proof** For any function  $\mathbf{v}$  in  $\mathbf{H}^1_{\partial D}(\Omega)$  and  $\mathbf{v}^h$  in  $V_{h,\partial D}$ , we have

$$\begin{aligned} & b(\boldsymbol{\xi}, \mathbf{v}) + \int_{\partial B_R} (\mathcal{T} - \mathcal{T}_N) \boldsymbol{\xi} \cdot \bar{\mathbf{v}} \, ds \\ & = b(\mathbf{u}, \mathbf{v}) - b_N^h(\mathbf{u}_N^h, \mathbf{v}^h) + b_N^h(\mathbf{u}_N^h, \mathbf{v}^h) - b(\mathbf{u}_N^h, \mathbf{v}) + \int_{\partial B_R} (\mathcal{T} - \mathcal{T}_N) \boldsymbol{\xi} \cdot \bar{\mathbf{v}} \, ds \\ & = \int_{\partial B_R} \mathbf{g}_N \cdot (\bar{\mathbf{v}} - \bar{\mathbf{v}}^h) \, ds - b_N^h(\mathbf{u}_N^h, \mathbf{v} - \mathbf{v}^h) \\ & \quad + \int_{\partial B_R} (\mathbf{g} - \mathbf{g}_N) \cdot \bar{\mathbf{v}} \, ds + \int_{\partial B_R} (\mathcal{T} - \mathcal{T}_N) \mathbf{u} \cdot \bar{\mathbf{v}} \, ds. \end{aligned}$$

For any  $\mathbf{v}^h \in \mathbf{V}_{h,\partial D}$ , it follows from the integration by parts that

$$\begin{aligned}
 & -b_N(\mathbf{u}_N^h, \mathbf{v} - \mathbf{v}^h) + \int_{\partial B_R} \mathbf{g}_N \cdot (\bar{\mathbf{v}} - \bar{\mathbf{v}}^h) \, ds \\
 &= - \sum_{K \in \mathcal{M}_h} \left\{ \mu \int_K \nabla \mathbf{u}_N^h : \nabla (\bar{\mathbf{v}} - \bar{\mathbf{v}}^h) \, dx + (\lambda + \mu) \int_K (\nabla \cdot \mathbf{u}_N^h) \nabla \cdot (\bar{\mathbf{v}} - \bar{\mathbf{v}}^h) \, dx \right\} \\
 & \quad - \sum_{K \in \mathcal{M}_h} \left\{ -\omega^2 \int_K \mathbf{u}_N^h \cdot (\bar{\mathbf{v}} - \bar{\mathbf{v}}^h) \, dx - \int_{\partial B_R \cap \partial K} (\mathcal{T}_N \mathbf{u}_N^h + \mathbf{g}_N) \cdot (\bar{\mathbf{v}} - \bar{\mathbf{v}}^h) \, ds \right\} \\
 &= \sum_{K \in \mathcal{M}_h} \left\{ - \int_{\partial K} \left[ \mu \nabla \mathbf{u}_N^h \cdot \mathbf{v} + (\lambda + \mu) (\nabla \cdot \mathbf{u}_N^h) \mathbf{v} \right] \cdot (\bar{\mathbf{v}} - \bar{\mathbf{v}}^h) \, ds \right. \\
 & \quad \left. + \int_{\partial B_R \cap \partial K} (\mathcal{T}_N \mathbf{u}_N^h + \mathbf{g}_N) \cdot (\bar{\mathbf{v}} - \bar{\mathbf{v}}^h) \, ds \right\} \\
 & \quad + \sum_{K \in \mathcal{M}_h} \int_K \left[ \mu \Delta \mathbf{u}_N^h + (\lambda + \mu) \nabla \nabla \cdot \mathbf{u}_N^h + \omega^2 \mathbf{u}_N^h \right] \cdot (\bar{\mathbf{v}} - \bar{\mathbf{v}}^h) \, dx \\
 &= \sum_{K \in \mathcal{M}_h} \left[ \int_K \mathcal{B} \mathbf{u}_N^h \cdot (\bar{\mathbf{v}} - \bar{\mathbf{v}}^h) \, dx + \sum_{e \in \partial K} \frac{1}{2} \int_e J_e \cdot (\bar{\mathbf{v}} - \bar{\mathbf{v}}^h) \, ds \right].
 \end{aligned}$$

We take  $\mathbf{v}^h = \Pi_h \mathbf{v} \in \mathbf{V}_{h,\partial D}$ , where  $\Pi_h$  is the Clément-type interpolation operator and has the following interpolation estimates (cf. [12]):

$$\|\mathbf{v} - \Pi_h \mathbf{v}\|_{L^2(K)} \lesssim h_K \|\nabla \mathbf{v}\|_{L^2(\tilde{K})}, \quad \|\mathbf{v} - \Pi_h \mathbf{v}\|_{L^2(e)} \lesssim h_e^{1/2} \|\mathbf{v}\|_{\mathbf{H}^1(\tilde{K}_e)}.$$

Here  $\tilde{K}$  is the union of all the triangular elements in  $\mathcal{M}_h$ , which have nonempty intersection with the element  $K$ , and  $\tilde{K}_e$  is the union of  $\{\tilde{K} : e \subset K \in \mathcal{M}_h\}$ . It follows from the definitions of  $\mathbf{g}$  and  $\mathbf{g}_N$  that

$$\begin{aligned}
 \int_{\partial B_R} (\mathbf{g} - \mathbf{g}_N) \cdot \bar{\mathbf{v}} \, ds &= \int_{\partial B_R} \left[ (\mathcal{B} \mathbf{u}^{\text{inc}} - \mathcal{T} \mathbf{u}^{\text{inc}}) - (\mathcal{B} \mathbf{u}^{\text{inc}} - \mathcal{T}_N \mathbf{u}^{\text{inc}}) \right] \cdot \bar{\mathbf{v}} \, ds \\
 &= \int_{\partial B_R} (\mathcal{T}_N - \mathcal{T}) \mathbf{u}^{\text{inc}} \cdot \bar{\mathbf{v}} \, ds.
 \end{aligned}$$

By Lemma 4.5, we have

$$\begin{aligned}
 & \left| \int_{\partial B_R} (\mathbf{g} - \mathbf{g}_N) \cdot \bar{\mathbf{v}} \, ds + \int_{\partial B_R} (\mathcal{T} - \mathcal{T}_N) \mathbf{u} \cdot \bar{\mathbf{v}} \, ds \right| \\
 &= \left| \int_{\partial B_R} (\mathcal{T} - \mathcal{T}_N) \mathbf{u}^s \cdot \bar{\mathbf{v}} \, ds \right| \lesssim N \left( \frac{\hat{R}}{R} \right)^N \|\mathbf{u}^{\text{inc}}\|_{\mathbf{H}^1(\Omega)} \|\mathbf{v}\|_{\mathbf{H}^1(\Omega)}.
 \end{aligned}$$

The proof is completed by combining the above estimates. □

The following lemma is to estimate the last term in (4.3).



**Lemma 4.7** For any  $\delta > 0$ , there exists a positive constant  $C(\delta)$  independent of  $N$  such that

$$\Re \int_{\partial B_R} \mathcal{T}_N \xi \cdot \bar{\xi} ds \leq C(\delta) \|\xi\|_{L^2(B_R \setminus \overline{B_{\hat{R}}})}^2 + \left(\frac{R}{\hat{R}}\right) \delta \|\xi\|_{H^1(B_R \setminus \overline{B_{\hat{R}}})}^2.$$

**Proof** Using (3.1), we get from a simple calculation that

$$\Re \int_{\partial B_R} \mathcal{T}_N \xi \cdot \bar{\xi} ds = 2\pi R \Re \sum_{|n| \leq N} (M_n \xi_n) \cdot \bar{\xi}_n.$$

Denote  $\hat{M}_n = (M_n + M_n^*)/2$ . Then  $\Re (M_n \xi_n) \cdot \bar{\xi}_n = (\hat{M}_n \xi_n) \cdot \bar{\xi}_n$ . It is shown in [38] that  $\hat{M}_n$  is negative definite for sufficiently large  $|n|$ , i.e., there exists  $N_0 > 0$  such that  $(\hat{M}_n \xi_n) \cdot \bar{\xi}_n \leq 0$  for any  $|n| > N_0$ . Hence

$$\Re \int_{\partial B_R} \mathcal{T}_N \xi \cdot \bar{\xi} ds = 2\pi R \sum_{|n| \leq \min(N_0, N)} (\hat{M}_n \xi_n) \cdot \bar{\xi}_n + 2\pi R \sum_{N \geq |n| > \min(N_0, N)} (\hat{M}_n \xi_n) \cdot \bar{\xi}_n \tag{4.8}$$

Here we define

$$\sum_{N > |n| > \min(N_0, N)} (\hat{M}_n \xi_n) \cdot \bar{\xi}_n = 0, \quad N_0 > N.$$

Since the second part in (4.8) is non-positive, we only need to estimate the first part which consists of finite terms. Moreover we have

$$\begin{aligned} \Re \int_{\partial B_R} \mathcal{T}_N \xi \cdot \bar{\xi} ds &\leq 2\pi R \sum_{|n| \leq \min(N_0, N)} (\hat{M}_n \xi_n) \cdot \bar{\xi}_n \\ &\leq C \sum_{|n| \leq \min(N_0, N)} |\xi_n|^2 \leq C \|\xi\|_{L^2(\partial B_R)}^2. \end{aligned} \tag{4.9}$$

Consider the annulus

$$B_R \setminus B_{\hat{R}} = \{(r, \theta) : \hat{R} < r < R, 0 < \theta < 2\pi\}.$$

For any  $u \in H^1(B_R \setminus \overline{B_{\hat{R}}})$  and  $\delta > 0$ , it follows from the Cauchy–Schwarz inequality that

$$\begin{aligned} (R - \hat{R})|u(R)|^2 &= \int_{\hat{R}}^R |u(r)|^2 dr + \int_{\hat{R}}^R \int_r^R \frac{d}{dt} |u(t)|^2 dt dr \\ &\leq \int_{\hat{R}}^R |u(r)|^2 dr + (R - \hat{R}) \int_{\hat{R}}^R 2|u(r)||u'(r)| dr \end{aligned}$$

$$\begin{aligned}
 &= \int_{\hat{R}}^R |u(r)|^2 dr + (R - \hat{R}) \int_{\hat{R}}^R 2 \frac{|u(r)|}{\sqrt{\delta}} \sqrt{\delta} |u'(r)| dr \\
 &\leq \int_{\hat{R}}^R |u(r)|^2 dr + \delta^{-1} (R - \hat{R}) \int_{\hat{R}}^R |u(r)|^2 dr + \delta (R - \hat{R}) \int_{\hat{R}}^R |u'(r)|^2 dr,
 \end{aligned}$$

which gives

$$|u(R)|^2 \leq \left[ \delta^{-1} + (R - \hat{R})^{-1} \right] \int_{\hat{R}}^R |u(r)|^2 dr + \delta \int_{\hat{R}}^R |u'(r)|^2 dr. \tag{4.10}$$

On the other hand, we have

$$\|\nabla u\|_{L^2(B_R \setminus \overline{B_{\hat{R}}})}^2 = 2\pi \sum_{n \in \mathbb{Z}} \int_{\hat{R}}^R \left( r |u'_n(r)|^2 + \frac{n^2}{r} |u_n(r)|^2 \right) dr, \tag{4.11}$$

$$\|u\|_{L^2(B_R \setminus \overline{B_{\hat{R}}})}^2 = 2\pi \sum_{n \in \mathbb{Z}} \int_{\hat{R}}^R r |u_n(r)|^2 dr. \tag{4.12}$$

Using (4.10)–(4.12), we have for any  $u \in H^1(B_R \setminus \overline{B_{\hat{R}}})$  that

$$\begin{aligned}
 \|u\|_{L^2(\partial B_R)}^2 &= 2\pi R \sum_{n \in \mathbb{Z}} |u_n(R)|^2 \\
 &\leq 2\pi R \left[ \delta^{-1} + (R - \hat{R})^{-1} \right] \sum_{n \in \mathbb{Z}} \int_{\hat{R}}^R |u_n(r)|^2 dr \\
 &\quad + 2\pi R \delta \sum_{n \in \mathbb{Z}} \int_{\hat{R}}^R |u'_n(r)|^2 dr \\
 &\leq 2\pi \left[ \delta^{-1} + (R - \hat{R})^{-1} \right] \left( \frac{R}{\hat{R}} \right) \sum_{n \in \mathbb{Z}} \int_{\hat{R}}^R r |u_n(r)|^2 dr \\
 &\quad + 2\pi \delta \left( \frac{R}{\hat{R}} \right) \sum_{n \in \mathbb{Z}} \int_{\hat{R}}^R \left( r |u'_n(r)|^2 + \frac{n^2}{r} |u_n(r)|^2 \right) dr \\
 &= 2\pi \left[ \delta^{-1} + (R - \hat{R})^{-1} \right] \left( \frac{R}{\hat{R}} \right) \|u\|_{L^2(B_R \setminus \overline{B_{\hat{R}}})}^2 + \delta \left( \frac{R}{\hat{R}} \right) \|\nabla u\|_{L^2(B_R \setminus \overline{B_{\hat{R}}})}^2 \\
 &\leq C(\delta) \|u\|_{L^2(B_R \setminus \overline{B_{\hat{R}}})}^2 + \left( \frac{R}{\hat{R}} \right) \delta \|\nabla u\|_{L^2(B_R \setminus \overline{B_{\hat{R}}})}^2. \tag{4.13}
 \end{aligned}$$

Therefore, combining (4.9) and (4.13) yields

$$\begin{aligned}
 \Re \int_{\partial B_R} \mathcal{F}_N \xi \cdot \bar{\xi} ds &\leq C \|\xi\|_{L^2(\partial B_R)}^2 \leq C(\delta) \|\xi\|_{L^2(B_R \setminus \overline{B_{\hat{R}}})}^2 + \left( \frac{R}{\hat{R}} \right) \delta \int_{\Omega} |\nabla \xi|^2 dx \\
 &\leq C(\delta) \|\xi\|_{L^2(B_R \setminus \overline{B_{\hat{R}}})}^2 + \left( \frac{R}{\hat{R}} \right) \delta \|\xi\|_{H^1(B_R \setminus \overline{B_{\hat{R}}})}^2,
 \end{aligned}$$

which completes the proof. □

To estimate the third term on the right hand side of (4.3), we consider the dual problem

$$b(\mathbf{v}, \mathbf{p}) = \int_{\Omega} \mathbf{v} \cdot \bar{\xi} \, d\mathbf{x}, \quad \forall \mathbf{v} \in \mathbf{H}^1_{\partial D}(\Omega). \tag{4.14}$$

It is easy to check that  $\mathbf{p}$  is the solution of the following boundary value problem

$$\begin{cases} \mu \Delta \mathbf{p} + (\lambda + \mu) \nabla \nabla \cdot \mathbf{p} + \omega^2 \mathbf{p} = -\xi & \text{in } \Omega, \\ \mathbf{p} = 0 & \text{on } \partial D, \\ \mathcal{B} \mathbf{p} = \mathcal{T}^* \mathbf{p} & \text{on } \partial B_R, \end{cases} \tag{4.15}$$

where  $\mathcal{T}^*$  is the adjoint operator to the DtN operator  $\mathcal{T}$ . Letting  $\mathbf{v} = \xi$  in (4.14), we obtain

$$\|\xi\|^2_{L^2(\Omega)} = b(\xi, \mathbf{p}) + \int_{\partial B_R} (\mathcal{T} - \mathcal{T}_N) \xi \cdot \bar{\mathbf{p}} \, ds - \int_{\partial B_R} (\mathcal{T} - \mathcal{T}_N) \xi \cdot \bar{\mathbf{p}} \, ds. \tag{4.16}$$

To evaluate (4.16), we need to explicitly solve system (4.15), which is very complicate due to the coupling of the compressional and shear wave components. We consider the Helmholtz decomposition to the boundary value problem (4.15). Let

$$\xi = \nabla \xi_1 + \mathbf{curl} \xi_2, \tag{4.17}$$

where  $\xi_j, j = 1, 2$  has the Fourier series expansion

$$\xi_j(r, \theta) = \sum_{n \in \mathbb{Z}} \xi_{jn}(r) e^{in\theta}, \quad \hat{R} < r < R.$$

Meanwhile, we assume that

$$\xi(r, \theta) = \sum_{n \in \mathbb{Z}} (\xi_n^r(r) \mathbf{e}_r + \xi_n^\theta(r) \mathbf{e}_\theta) e^{in\theta}. \tag{4.18}$$

Using the Fourier series expansions and the Helmholtz decomposition, we get

$$\begin{aligned} \xi(r, \theta) &= \sum_{n \in \mathbb{Z}} [\xi_n^r(r) \mathbf{e}_r + \xi_n^\theta(r) \mathbf{e}_\theta] e^{in\theta} = \nabla \xi_1 + \mathbf{curl} \xi_2 \\ &= \sum_{n \in \mathbb{Z}} \left[ \xi'_{1n}(r) \mathbf{e}_r + \frac{in}{r} \xi_{1n}(r) \mathbf{e}_\theta + \frac{in}{r} \xi_{2n}(r) \mathbf{e}_r - \xi'_{2n}(r) \mathbf{e}_\theta \right] e^{in\theta} \\ &= \sum_{n \in \mathbb{Z}} \left[ \left( \xi'_{1n}(r) + \frac{in}{r} \xi_{2n}(r) \right) \mathbf{e}_r + \left( \frac{in}{r} \xi_{1n}(r) - \xi'_{2n}(r) \right) \mathbf{e}_\theta \right] e^{in\theta}, \end{aligned}$$

which shows that the Fourier coefficients  $\xi_{1n}, \xi_{2n}$  satisfy

$$\xi'_{1n}(r) + \frac{in}{r}\xi_{2n}(r) = \xi_n^r(r), \quad \frac{in}{r}\xi_{1n}(r) - \xi'_{2n}(r) = \xi_n^\theta(r), \quad r \in (\hat{R}, R).$$

**Lemma 4.8** *The system for the Fourier coefficients*

$$\begin{cases} \xi'_{1n}(r) + \frac{in}{r}\xi_{2n}(r) = \xi_n^r(r), & r \in (\hat{R}, R), \\ \frac{in}{r}\xi_{1n}(r) - \xi'_{2n}(r) = \xi_n^\theta(r), & r \in (\hat{R}, R), \\ \xi_{1n}(R) = 0, \quad \xi_{2n}(R) = 0, & r = R, \end{cases} \quad (4.19)$$

has a unique solution given by

$$\xi_{1n}(r) = -\frac{1}{2} \int_r^R \left[ \left(\frac{r}{t}\right)^n + \left(\frac{t}{r}\right)^n \right] \xi_n^r(t) dt + \frac{i}{2} \int_r^R \left[ \left(\frac{r}{t}\right)^n - \left(\frac{t}{r}\right)^n \right] \xi_n^\theta(t) dt, \quad (4.20)$$

$$\xi_{2n}(r) = \frac{i}{2} \int_r^R \left[ \left(\frac{t}{r}\right)^n - \left(\frac{r}{t}\right)^n \right] \xi_n^r(t) dt - \frac{1}{2} \int_r^R \left[ \left(\frac{r}{t}\right)^n + \left(\frac{t}{r}\right)^n \right] \xi_n^\theta(t) dt. \quad (4.21)$$

**Proof** Denote

$$A_n(r) = \begin{bmatrix} 0 & -\frac{in}{r} \\ \frac{in}{r} & 0 \end{bmatrix}.$$

By the standard theory of the first order differential system, the fundamental solution  $\Phi_n(r)$  is

$$\begin{aligned} \Phi_n(r) &= e^{\int_{\hat{R}}^r A_n(\tau) d\tau} = \exp \left( \begin{bmatrix} 0 & -in \ln \frac{r}{\hat{R}} \\ in \ln \frac{r}{\hat{R}} & 0 \end{bmatrix} \right) \\ &= \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{i}{\sqrt{2}} \\ \frac{i}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix} \begin{bmatrix} \left(\frac{r}{\hat{R}}\right)^n & 0 \\ 0 & \left(\frac{r}{\hat{R}}\right)^{-n} \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{2}} & -\frac{i}{\sqrt{2}} \\ -\frac{i}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix}. \end{aligned}$$

The inverse of  $\Phi_n$  is

$$\Phi_n^{-1}(r) = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{i}{\sqrt{2}} \\ \frac{i}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix} \begin{bmatrix} \left(\frac{r}{\hat{R}}\right)^{-n} & 0 \\ 0 & \left(\frac{r}{\hat{R}}\right)^n \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{2}} & -\frac{i}{\sqrt{2}} \\ -\frac{i}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix}.$$

Using the method of variation of parameters, we let

$$(\xi_{1n}(r), \xi_{2n}(r))^\top = \Phi_n(r) C_n(r),$$

where the unknown vector  $C_n(r)$  satisfies

$$\begin{aligned}
 C'_n(r) &= \Phi_n^{-1}(r)(\xi_n^r(r), \xi_n^\theta(r))^\top \\
 &= \frac{1}{2} \begin{bmatrix} \left[ \left(\frac{r}{\hat{R}}\right)^{-n} + \left(\frac{r}{\hat{R}}\right)^n \right] \xi_n^r(r) + i \left[ \left(\frac{r}{\hat{R}}\right)^n - \left(\frac{r}{\hat{R}}\right)^{-n} \right] \xi_n^\theta(r) \\ i \left[ \left(\frac{r}{\hat{R}}\right)^{-n} - \left(\frac{r}{\hat{R}}\right)^n \right] \xi_n^r(r) + \left[ \left(\frac{r}{\hat{R}}\right)^{-n} + \left(\frac{r}{\hat{R}}\right)^n \right] \xi_n^\theta(r) \end{bmatrix}. \tag{4.22}
 \end{aligned}$$

Using the boundary condition yields

$$(\xi_{1n}(R), \xi_{2n}(R))^\top = \Phi_n(R)C_n(R) = (0, 0)^\top,$$

which implies that  $C_n(R) = (0, 0)^\top$ . Then

$$C_n(r) = - \int_r^R C'_n(t)dt. \tag{4.23}$$

Combining (4.22) and (4.23), we have

$$C_n(r) = -\frac{1}{2} \begin{bmatrix} \int_r^R \left[ \left(\frac{t}{\hat{R}}\right)^{-n} + \left(\frac{t}{\hat{R}}\right)^n \right] \xi_n^r(t)dt + i \int_r^R \left[ \left(\frac{t}{\hat{R}}\right)^n - \left(\frac{t}{\hat{R}}\right)^{-n} \right] \xi_n^\theta(t)dt \\ i \int_r^R \left[ \left(\frac{t}{\hat{R}}\right)^{-n} - \left(\frac{t}{\hat{R}}\right)^n \right] \xi_n^r(t)dt + \int_r^R \left[ \left(\frac{t}{\hat{R}}\right)^n + \left(\frac{t}{\hat{R}}\right)^{-n} \right] \xi_n^\theta(t)dt. \end{bmatrix}.$$

Substituting  $C_n(r)$  into the general solution, we obtain

$$\begin{aligned}
 \xi_{1n}(r) &= -\frac{1}{2} \left(\frac{r}{\hat{R}}\right)^n \int_r^R \left(\frac{t}{\hat{R}}\right)^{-n} \xi_n^r(t)dt + \frac{i}{2} \left(\frac{r}{\hat{R}}\right)^n \int_r^R \left(\frac{t}{\hat{R}}\right)^{-n} \xi_n^\theta(t)dt \\
 &\quad -\frac{1}{2} \left(\frac{r}{\hat{R}}\right)^{-n} \int_r^R \left(\frac{t}{\hat{R}}\right)^n \xi_n^r(t)dt - \frac{i}{2} \left(\frac{r}{\hat{R}}\right)^{-n} \int_r^R \left(\frac{t}{\hat{R}}\right)^n \xi_n^\theta(t)dt \\
 \xi_{2n}(r) &= -\frac{i}{2} \left(\frac{r}{\hat{R}}\right)^n \int_r^R \left(\frac{t}{\hat{R}}\right)^{-n} \xi_n^r(t)dt + \frac{i}{2} \left(\frac{r}{\hat{R}}\right)^{-n} \int_r^R \left(\frac{t}{\hat{R}}\right)^n \xi_n^r(t)dt \\
 &\quad -\frac{1}{2} \left(\frac{r}{\hat{R}}\right)^n \int_r^R \left(\frac{t}{\hat{R}}\right)^{-n} \xi_n^\theta(t)dt - \frac{1}{2} \left(\frac{r}{\hat{R}}\right)^{-n} \int_r^R \left(\frac{t}{\hat{R}}\right)^n \xi_n^\theta(t)dt,
 \end{aligned}$$

which completes the proof. □

Let  $\mathbf{p}$  be the solution of the dual problem (4.15). Then  $\mathbf{p}$  satisfies the following boundary value problem in  $B_R \setminus \overline{B_{\hat{R}}}$ :

$$\begin{cases} \mu \Delta \mathbf{p} + (\lambda + \mu) \nabla \nabla \cdot \mathbf{p} + \omega^2 \mathbf{p} = -\boldsymbol{\xi} & \text{in } B_R \setminus \overline{B_{\hat{R}}}, \\ \mathbf{p}(\hat{R}, \theta) = \mathbf{p}(\hat{R}, \theta) & \text{on } \partial B_{\hat{R}}, \\ \mathcal{B} \mathbf{p} = \mathcal{T}^* \mathbf{p} & \text{on } \partial B_R. \end{cases} \tag{4.24}$$

Introduce the Helmholtz decomposition for  $\mathbf{p}$ :

$$\mathbf{p} = \nabla q_1 + \mathbf{curl} q_2, \tag{4.25}$$

where  $q_j, j = 1, 2$  admits the Fourier series expansion

$$q_j(r, \theta) = \sum_{n \in \mathbb{Z}} q_{jn}(r) e^{in\theta}.$$

Let  $\xi_{jn}, j = 1, 2$  be the solution of the system (4.19). Consider the second order system for  $q_{jn}, j = 1, 2$ :

$$\begin{cases} q''_{jn}(r) + \frac{1}{r}q'_{jn}(r) + (\kappa_j^2 - (\frac{n}{R})^2)q_{jn}(r) = c_j \xi_{jn}(r), & r \in (\hat{R}, R), \\ q_{jn}(\hat{R}) = q_{jn}(\hat{R}), & r = \hat{R}, \\ q'_{jn}(R) = \bar{\alpha}_{jn}q_{jn}(R), & r = R, \end{cases} \tag{4.26}$$

where  $c_1 = -1/(\lambda + 2\mu), c_2 = -1/\mu$ , and  $\alpha_{jn}$  is given in (A.5). The boundary condition  $q'_{jn}(R) = \bar{\alpha}_{jn}q_{jn}(R)$  comes from (A.2), i.e.,  $q_j$  satisfies the boundary condition

$$\partial_r q_j = \mathcal{T}_j^* q_j := \sum_{n \in \mathbb{Z}} \bar{\alpha}_{jn} q_{jn}(R) e^{in\theta} \quad \text{on } \partial B_R,$$

where  $\mathcal{T}_j^*$  is the adjoint operator to the DtN operator  $\mathcal{T}_j$ .

**Lemma 4.9** *The boundary value problem (4.24) and the second order system (4.26) are equivalent under the Helmholtz decomposition (4.25).*

**Proof** It suffices to show if the Fourier coefficients  $q_{jn}$  satisfy the second order system (4.26), then  $\mathbf{p} = \nabla q_1 + \mathbf{curl} q_2$  is the solution of (4.24).

In the polar coordinates, we let

$$\mathbf{p}(r, \theta) = \sum_{n \in \mathbb{Z}} (p_n^r(r) \mathbf{e}_r + p_n^\theta(r) \mathbf{e}_\theta) e^{in\theta}, \quad r \in (\hat{R}, R). \tag{4.27}$$

It follows from the Helmholtz decomposition that

$$p_n^r(r) = q'_{1n}(r) + \frac{in}{r} q_{2n}(r), \quad p_n^\theta(r) = \frac{in}{r} q_{1n}(r) - q'_{2n}(r). \tag{4.28}$$

Using (4.27)–(4.28), we have from a straightforward calculation that

$$\begin{aligned} \mathcal{B}\mathbf{p} &= (\mu \partial_r \mathbf{p} + (\lambda + \mu) \nabla \cdot \mathbf{p} \mathbf{e}_r)|_{r=R} \\ &= \sum_{n \in \mathbb{Z}} \left[ (\lambda + 2\mu) q''_{1n}(R) + (\lambda + \mu) \frac{1}{R} q'_{1n}(R) - (\lambda + \mu) \frac{n^2}{R^2} q_{1n}(R) \right] e^{in\theta} \mathbf{e}_r \end{aligned}$$

$$\begin{aligned}
 & + \sum_{n \in \mathbb{Z}} \left[ \mu \frac{in}{R} q'_{1n}(R) - \mu \frac{in}{R^2} q_{1n}(R) \right] e^{in\theta} \mathbf{e}_\theta \\
 & + \sum_{n \in \mathbb{Z}} \left[ \mu \frac{in}{R} q'_{2n}(R) - \mu \frac{in}{R^2} q_{2n}(R) \right] e^{in\theta} \mathbf{e}_r + \sum_{n \in \mathbb{Z}} -\mu q''_{2n}(R) e^{in\theta} \mathbf{e}_\theta.
 \end{aligned}$$

On the other hand, it is easy to verify that

$$\begin{aligned}
 \mathcal{T}^* \mathbf{p} & = \sum_{n \in \mathbb{Z}} \left\{ \left[ \overline{M_{11}^{(n)}} p_n^r(R) + \overline{M_{21}^{(n)}} p_n^\theta(R) \right] \mathbf{e}_r + \left[ \overline{M_{12}^{(n)}} p_n^r(R) + \overline{M_{22}^{(n)}} p_n^\theta(R) \right] \mathbf{e}_\theta \right\} e^{in\theta} \\
 & = \sum_{n \in \mathbb{Z}} \left\{ \overline{M_{11}^{(n)}} \left[ q'_{1n}(R) + \frac{in}{R} q_{2n}(R) \right] + \overline{M_{21}^{(n)}} \left[ \frac{in}{R} q_{1n}(R) - q'_{2n}(R) \right] \right\} \mathbf{e}_r e^{in\theta} \\
 & \quad + \sum_{n \in \mathbb{Z}} \left\{ \overline{M_{12}^{(n)}} \left[ q'_{1n}(R) + \frac{in}{R} q_{2n}(R) \right] + \overline{M_{22}^{(n)}} \left[ \frac{in}{R} q_{1n}(R) - q'_{2n}(R) \right] \right\} \mathbf{e}_\theta e^{in\theta},
 \end{aligned}$$

where  $M_{ij}^{(n)}$ ,  $i, j = 1, 2$  are given in (A.4).

Using the boundary condition  $q'_{jn}(R) = \overline{\alpha_{jn}} q_{jn}(R)$ , we get

$$\begin{aligned}
 & \left( \mu \frac{in}{R} - \overline{M_{12}^{(n)}} \right) q'_{1n}(R) - \left( \overline{M_{22}^{(n)}} \frac{in}{R} + \mu \frac{in}{R^2} \right) q_{1n}(R) \\
 & = \left( \mu \frac{in}{R} - in \frac{\mu}{R} + \omega^2 \frac{in}{R} \frac{1}{\Lambda_n(R)} \right) q'_{1n}(R) \\
 & \quad - \left( -\frac{\mu}{R} \frac{in}{R} + \omega^2 \frac{in}{R} \frac{\overline{\alpha_{1n}}}{\Lambda_n(R)} + \mu \frac{in}{R^2} \right) q_{1n}(R) \\
 & = \omega^2 \frac{in}{R} \frac{1}{\Lambda_n(R)} (q'_{1n}(R) - \overline{\alpha_{1n}} q_{1n}(R)) = 0
 \end{aligned}$$

and

$$\begin{aligned}
 & \left( \mu \frac{in}{R} + \overline{M_{21}^{(n)}} \right) q'_{2n}(R) - \left( \overline{M_{11}^{(n)}} \frac{in}{R} + \mu \frac{in}{R^2} \right) q_{2n}(R) \\
 & = \left( \mu \frac{in}{R} - in \frac{\mu}{R} + in \frac{\omega^2}{R} \frac{1}{\Lambda_n(R)} \right) q'_{2n}(R) \\
 & \quad - \left( \mu \frac{in}{R^2} - \frac{\mu}{R} \frac{in}{R} + \omega^2 \frac{in}{R} \frac{\overline{\alpha_{2n}}}{\Lambda_n(R)} \right) q_{2n}(R) \\
 & = in \frac{\omega^2}{R} \frac{1}{\Lambda_n(R)} (q'_{2n}(R) - \overline{\alpha_{2n}} q_{2n}(R)) = 0.
 \end{aligned}$$

Since  $q_{2n}$  satisfies the second order equation

$$q''_{2n}(r) + \frac{1}{r} q'_{2n}(r) + \left( \kappa_2^2 - \left( \frac{n}{r} \right)^2 \right) q_{2n}(r) = -\frac{1}{\mu} \xi_{2n}, \quad r \in (\hat{R}, R),$$

we obtain from the boundary condition  $\xi_{2n}(R) = 0$  that

$$\begin{aligned} & -\mu q''_{2n}(R) - \left( \overline{M_{12}^{(n)}} \frac{in}{R} q_{2n}(R) - \overline{M_{22}^{(n)}} q'_{2n}(R) \right) \\ &= -\mu q''_{2n}(R) - \frac{in}{R} \left( in \frac{\mu}{R} - \omega^2 \frac{in}{R} \frac{1}{\Lambda_n(R)} \right) q_{2n}(R) + \left( -\frac{\mu}{R} + \omega^2 \frac{\overline{\alpha_{1n}}}{\Lambda_n(R)} \right) q'_{2n}(R) \\ &= \xi_{2n}(R) + \mu \kappa_2^2 q_{2n}(R) + \omega^2 \left( \frac{in}{R} \right)^2 \frac{1}{\Lambda_n(R)} q_{2n}(R) + \omega^2 \frac{\overline{\alpha_{1n}}}{\Lambda_n(R)} q'_{2n}(R) \\ &= \xi_{2n}(R) + \frac{\omega^2}{\Lambda_n(R)} \left( \left( \frac{n}{R} \right)^2 q_{2n}(R) - \overline{\alpha_{1n} \alpha_{2n}} q_{2n}(R) - \left( \frac{n}{R} \right)^2 q_{2n}(R) + \overline{\alpha_{1n}} q'_{2n}(R) \right) \\ &= \xi_{2n}(R) + \omega^2 \frac{\overline{\alpha_{1n}}}{\Lambda_n(R)} (-\overline{\alpha_{2n}} q_{2n}(R) + q'_{2n}(R)) = 0. \end{aligned}$$

Similarly, combining the equation

$$q''_{1n}(r) + \frac{1}{r} q'_{1n}(r) + \left( \kappa_1^2 - \left( \frac{n}{r} \right)^2 \right) q_{1n}(r) = -\frac{1}{\lambda + 2\mu} \xi_{1n}(r), \quad r \in (\hat{R}, R)$$

and the boundary condition  $\xi_{1n}(R) = 0$ , we have

$$\begin{aligned} & (\lambda + 2\mu) q''_{1n}(R) + (\lambda + \mu) \frac{1}{R} q'_{1n}(R) - (\lambda + \mu) \frac{n^2}{R^2} q_{1n}(R) \\ & \quad - \overline{M_{11}^{(n)}} q'_{1n}(R) - \frac{in}{R} \overline{M_{21}^{(n)}} q_{1n}(R) \\ &= (\lambda + 2\mu) \left[ -\frac{1}{\lambda + 2\mu} \xi_{1n}(R) - \frac{1}{R} q'_{1n}(R) - \left( \kappa_1^2 - \left( \frac{n}{R} \right)^2 \right) q_{1n}(R) \right] \\ & \quad + \left( \frac{\lambda + 2\mu}{R} - \omega^2 \frac{\overline{\alpha_{2n}}}{\Lambda_n(R)} \right) q'_{1n}(R) \\ & \quad + \left( -(\lambda + 2\mu) \left( \frac{n}{R} \right)^2 + \omega^2 \frac{1}{\Lambda_n(R)} \left( \frac{n}{R} \right)^2 \right) q_{1n}(R) \\ &= -\xi_{1n}(R) - \omega^2 \frac{\overline{\alpha_{2n}}}{\Lambda_n(R)} q'_{1n}(R) + \left( -\omega^2 + \omega^2 \frac{1}{\Lambda_n(R)} \left( \frac{n}{R} \right)^2 \right) q_{1n}(R) \\ &= -\xi_{1n}(R) - \frac{\omega^2}{\Lambda_n(R)} \left( \overline{\alpha_{2n}} q'_{1n}(R) + \left( \frac{n}{R} \right)^2 q_{1n}(R) - \overline{\alpha_{1n} \alpha_{2n}} q_{1n}(R) \right. \\ & \quad \left. - \left( \frac{n}{R} \right)^2 q_{1n}(R) \right) \\ &= -\xi_{1n}(R) - \frac{\omega^2}{\Lambda_n(R)} \overline{\alpha_{2n}} [q'_{1n}(R) - \overline{\alpha_{1n}} q_{1n}(R)] = 0. \end{aligned}$$

Hence we prove that  $\mathcal{B}\mathbf{p} = \mathcal{T}^*\mathbf{p}$  on  $\partial B_R$ .

Moreover, we get from the Helmholtz decomposition that

$$\mu \Delta \mathbf{p} + (\lambda + \mu) \nabla \nabla \cdot \mathbf{p} + \omega^2 \mathbf{p}$$



$$\begin{aligned} &= \nabla \left( (\lambda + 2\mu)\Delta q_1 + \omega^2 q_1 \right) + \mathbf{curl} \left( \mu \Delta q_2 + \omega^2 q_2 \right) \\ &= -\nabla \xi_1 - \mathbf{curl} \xi_2 = -\xi, \end{aligned}$$

which completes the proof. □

Based on Lemma 4.8 and Lemma 4.9, we have the asymptotic properties of the solution to the dual problem (4.24) for large  $|n|$ .

**Theorem 4.10** *Let  $p$  be the solution of (4.24) and admit the Fourier series expansion*

$$p(r, \theta) = \sum_{n \in \mathbb{Z}} \left( p_n^r(r) \mathbf{e}_r + p_n^\theta(r) \mathbf{e}_\theta \right) e^{in\theta}.$$

For sufficient large  $|n|$ , the Fourier coefficients  $p_n^r, p_n^\theta$  satisfy the estimate

$$\begin{aligned} |p_n^r(R)|^2 + |p_n^\theta(R)|^2 &\lesssim n^2 \left( \frac{\hat{R}}{R} \right)^{2|n|+2} \left( |p_n^r(\hat{R})|^2 + |p_n^\theta(\hat{R})|^2 \right) \\ &\quad + \frac{1}{|n|^2} \left( \|\xi_n^r\|_{L^\infty(\hat{R}, R)}^2 + \|\xi_n^\theta\|_{L^\infty(\hat{R}, R)}^2 \right), \end{aligned}$$

where  $\xi_n^r, \xi_n^\theta$  are the Fourier coefficients of  $\xi$  in the polar coordinates and are given in (4.18).

**Proof** It follows from straightforward calculations that the second order systems (4.26) have a unique solution, which is given by

$$\begin{aligned} q_{1n}(r) &= \beta_{1n}(r)q_{1n}(\hat{R}) + \frac{i\pi}{4} \int_{\hat{R}}^r t W_{1n}(r, t) \xi_{1n}(t) dt \\ &\quad + \frac{i\pi}{4} \int_{\hat{R}}^R t \beta_{1n}(t) W_{1n}(\hat{R}, r) \xi_{1n}(t) dt, \end{aligned} \tag{4.29}$$

$$\begin{aligned} q_{2n}(r) &= \beta_{2n}(r)q_{2n}(\hat{R}) + \frac{i\pi}{4} \int_{\hat{R}}^r t W_{2n}(r, t) \xi_{2n}(t) dt \\ &\quad + \frac{i\pi}{4} \int_{\hat{R}}^R t \beta_{2n}(t) W_{2n}(\hat{R}, r) \xi_{2n}(t) dt, \end{aligned} \tag{4.30}$$

where

$$\beta_{jn}(r) = \frac{H_n^{(2)}(\kappa_j r)}{H_n^{(2)}(\kappa_j \hat{R})}, \quad W_{jn}(r, t) = H_n^{(1)}(\kappa_j r) H_n^{(2)}(\kappa_j t) - H_n^{(1)}(\kappa_j t) H_n^{(2)}(\kappa_j r).$$

Taking the derivative of (4.29)–(4.30) with respect to  $r$  gives

$$q'_{1n}(r) = \beta'_{1n}(r)q_{1n}(\hat{R}) + \frac{i\pi}{4} \int_{\hat{R}}^r t \partial_r W_{1n}(r, t) \xi_{1n}(t) dt$$

$$+ \frac{i\pi}{4} \int_{\hat{R}}^R t \beta_{1n}(t) \partial_t W_{1n}(\hat{R}, r) \xi_{1n}(t) dt, \tag{4.31}$$

$$q'_{2n}(r) = \beta'_{2n}(r) q_{2n}(\hat{R}) + \frac{i\pi}{4} \int_{\hat{R}}^r t \partial_r W_{2n}(r, t) \xi_{2n}(t) dt + \frac{i\pi}{4} \int_{\hat{R}}^R t \beta_{2n}(t) \partial_t W_{2n}(\hat{R}, r) \xi_{2n}(t) dt. \tag{4.32}$$

Evaluating (4.29)–(4.30) and (4.31)–(4.32) at  $r = R$  and  $r = \hat{R}$ , respectively, we may verify that

$$\begin{aligned} q_{1n}(R) &= \beta_{1n}(R) q_{1n}(\hat{R}) + \frac{i\pi}{4} \int_{\hat{R}}^R t \beta_{1n}(R) W_{1n}(\hat{R}, t) \xi_{1n}(t) dt, \\ q_{2n}(R) &= \beta_{2n}(R) q_{2n}(\hat{R}) + \frac{i\pi}{4} \int_{\hat{R}}^R t \beta_{2n}(R) W_{2n}(\hat{R}, t) \xi_{2n}(t) dt, \\ q'_{1n}(\hat{R}) &= \beta'_{1n}(\hat{R}) q_{1n}(\hat{R}) + \frac{1}{\hat{R}} \int_{\hat{R}}^R t \beta_{1n}(t) \xi_{1n}(t) dt, \\ q'_{2n}(\hat{R}) &= \beta'_{2n}(\hat{R}) q_{2n}(\hat{R}) + \frac{1}{\hat{R}} \int_{\hat{R}}^R t \beta_{2n}(t) \xi_{2n}(t) dt. \end{aligned}$$

It follows from the Helmholtz decomposition that

$$p_n^r(r) = q'_{1n}(r) + \frac{in}{r} q_{2n}(r), \quad p_n^\theta(r) = \frac{in}{r} q_{1n}(r) - q'_{2n}(r). \tag{4.33}$$

Evaluating (4.33) at  $r = R$ , noting  $\beta'_{jn}(R) = \overline{\alpha_{jn}(R)}$  and  $q'_{jn}(R) = \overline{\alpha_{jn}(R)} q_{jn}(R)$ , we obtain

$$\begin{bmatrix} p_n^r(R) \\ p_n^\theta(R) \end{bmatrix} = U_n(R) \begin{bmatrix} q_{1n}(\hat{R}) \\ q_{2n}(\hat{R}) \end{bmatrix} + \frac{i\pi}{4} U_n(R) \begin{bmatrix} \int_{\hat{R}}^R t W_{1n}(\hat{R}, t) \xi_{1n}(t) dt \\ \int_{\hat{R}}^R t W_{2n}(\hat{R}, t) \xi_{2n}(t) dt \end{bmatrix}, \tag{4.34}$$

where

$$U_n(R) = \begin{bmatrix} \overline{\alpha_{1n}(R)} & \frac{in}{R} \\ \frac{in}{R} & -\alpha_{2n}(R) \end{bmatrix} \begin{bmatrix} \beta_{1n}(R) & 0 \\ 0 & \beta_{2n}(R) \end{bmatrix}.$$

Similarly, evaluating (4.33) at  $r = \hat{R}$  and noting  $\beta'_{jn}(\hat{R}) = \overline{\alpha_{jn}(\hat{R})}$  yield that

$$\begin{bmatrix} p_n^r(\hat{R}) \\ p_n^\theta(\hat{R}) \end{bmatrix} = K_n(\hat{R}) \begin{bmatrix} q_{1n}(\hat{R}) \\ q_{2n}(\hat{R}) \end{bmatrix} + \begin{bmatrix} \eta_1 \\ \eta_2 \end{bmatrix}, \tag{4.35}$$

where

$$K_n(\hat{R}) = \begin{bmatrix} \overline{\alpha_{1n}(\hat{R})} & \frac{in}{\hat{R}} \\ \frac{in}{\hat{R}} & -\alpha_{2n}(\hat{R}) \end{bmatrix},$$

and

$$\eta_{1n} = \frac{1}{\hat{R}} \int_{\hat{R}}^R t \beta_{1n}(t) \xi_{1n}(t) dt, \quad \eta_{2n} = -\frac{1}{\hat{R}} \int_{\hat{R}}^R t \beta_{2n}(t) \xi_{2n}(t) dt.$$

Solving (4.35) for  $q_{1n}(\hat{R}), q_{2n}(\hat{R})$  in terms of  $p_n^r(\hat{R}), p_n^\theta(\hat{R})$  gives

$$\begin{bmatrix} q_{1n}(\hat{R}) \\ q_{2n}(\hat{R}) \end{bmatrix} = \frac{V_n(\hat{R})}{\Lambda_n(\hat{R})} \begin{bmatrix} p_n^r(\hat{R}) - \eta_{1n} \\ p_n^\theta(\hat{R}) - \eta_{2n} \end{bmatrix}, \tag{4.36}$$

where

$$\Lambda_n(\hat{R}) = \left(\frac{n}{\hat{R}}\right)^2 - \alpha_{1n}(\hat{R})\alpha_{2n}(\hat{R}), \quad V_n(\hat{R}) = \begin{bmatrix} -\overline{\alpha_{2n}(\hat{R})} & -\frac{in}{\hat{R}} \\ -\frac{in}{\hat{R}} & \alpha_{1n}(\hat{R}) \end{bmatrix}.$$

Substituting (4.36) into (4.34) yields

$$\begin{bmatrix} p_n^r(R) \\ p_n^\theta(R) \end{bmatrix} = \frac{U_n(R)V_n(\hat{R})}{\Lambda_n(\hat{R})} \begin{bmatrix} p_n^r(\hat{R}) \\ p_n^\theta(\hat{R}) \end{bmatrix} + \frac{i\pi}{4} U_n(R) \begin{bmatrix} \int_{\hat{R}}^R t W_{1n}(\hat{R}, t) \xi_{1n}(t) dt \\ \int_{\hat{R}}^R t W_{2n}(\hat{R}, t) \xi_{2n}(t) dt \end{bmatrix} - \frac{U_n(R)V_n(\hat{R})}{\Lambda_n(\hat{R})} \begin{bmatrix} \eta_{1n} \\ \eta_{2n} \end{bmatrix}. \tag{4.37}$$

Following the proofs in Lemmas 4.5 and A.1, we may similarly show that for sufficiently large  $|n|$

$$\left| \frac{U_n(R)V_n(\hat{R})}{\Lambda_n(\hat{R})} \right| \lesssim |n| \left(\frac{\hat{R}}{R}\right)^{|n|}.$$

For fixed  $t$  and sufficiently large  $|n|$ , using (4.20) and (4.21), we may easily show

$$|\xi_{1n}(t)| \lesssim \left( \|\xi_n^r\|_{L^\infty([\hat{R}, R])} + \|\xi_n^\theta\|_{L^\infty([\hat{R}, R])} \right) \int_t^R \left(\frac{r}{t}\right)^{|n|} dr, \tag{4.38}$$

$$|\xi_{2n}(t)| \lesssim \left( \|\xi_n^r\|_{L^\infty([\hat{R}, R])} + \|\xi_n^\theta\|_{L^\infty([\hat{R}, R])} \right) \int_t^R \left(\frac{r}{t}\right)^{|n|} dr. \tag{4.39}$$

By (4.38)–(4.39) and

$$W_{jn}(\hat{R}, t) \sim -\frac{2i}{\pi|n|} \left[ \left( \frac{t}{\hat{R}} \right)^{|n|} - \left( \frac{\hat{R}}{t} \right)^{|n|} \right], \quad \beta_{jn}(t) \sim \left( \frac{\hat{R}}{t} \right)^{|n|},$$

we get

$$\begin{aligned} \left| \int_{\hat{R}}^R t W_{jn}(\hat{R}, t) \xi_{jn}(t) dt \right| &\lesssim \left( \|\xi_n^r\|_{L^\infty([\hat{R}, R])} + \|\xi_n^\theta\|_{L^\infty([\hat{R}, R])} \right) \\ &\quad \frac{1}{|n|} \int_{\hat{R}}^R t \left( \frac{t}{\hat{R}} \right)^{|n|} \int_t^R \left( \frac{r}{t} \right)^{|n|} dr dt \\ &= \left( \|\xi_n^r\|_{L^\infty([\hat{R}, R])} + \|\xi_n^\theta\|_{L^\infty([\hat{R}, R])} \right) \frac{1}{|n|(|n|+1)} \\ &\quad \times \int_{\hat{R}}^R t \left( \frac{1}{\hat{R}} \right)^{|n|} \left( R^{|n|+1} - t^{|n|+1} \right) dt \\ &\lesssim \left( \|\xi_n^r\|_{L^\infty([\hat{R}, R])} + \|\xi_n^\theta\|_{L^\infty([\hat{R}, R])} \right) \frac{1}{|n|^2} \left( \frac{R}{\hat{R}} \right)^{|n|} \end{aligned}$$

and

$$\begin{aligned} \left| \frac{1}{\hat{R}} \int_{\hat{R}}^R t \beta_{jn}(t) \xi_{jn}(t) dt \right| &\lesssim \left( \|\xi_n^r\|_{L^\infty([\hat{R}, R])} + \|\xi_n^\theta\|_{L^\infty([\hat{R}, R])} \right) \\ &\quad \int_{\hat{R}}^R t \left( \frac{\hat{R}}{t} \right)^{|n|} \int_t^R \left( \frac{r}{t} \right)^{|n|} dr dt \\ &= \left( \|\xi_n^r\|_{L^\infty([\hat{R}, R])} + \|\xi_n^\theta\|_{L^\infty([\hat{R}, R])} \right) \frac{1}{1+|n|} \\ &\quad \times \int_{\hat{R}}^R \left( \frac{\hat{R}}{t} \right)^{|n|} \frac{R^{|n|+1} - t^{|n|+1}}{t^{|n|-1}} dt \\ &\lesssim \left( \|\xi_n^r\|_{L^\infty([\hat{R}, R])} + \|\xi_n^\theta\|_{L^\infty([\hat{R}, R])} \right) \frac{1}{|n|^2} \left( \frac{R}{\hat{R}} \right)^{|n|}. \end{aligned}$$

Substituting the above estimates into (4.37), we obtain

$$\begin{aligned} |p_n^r(R)|^2 + |p_n^\theta(R)|^2 &\lesssim n^2 \left( \frac{\hat{R}}{R} \right)^{2|n|+2} \left( |p_n^r(\hat{R})|^2 + |p_n^\theta(\hat{R})|^2 \right) \\ &\quad + \frac{1}{|n|^2} \left( \|\xi_n^r\|_{L^\infty([\hat{R}, R])}^2 + \|\xi_n^\theta\|_{L^\infty([\hat{R}, R])}^2 \right), \end{aligned}$$

which completes the proof.  $\square$

Using Theorem 4.10, we may estimate the last term in (4.16).

**Lemma 4.11** *Let  $\mathbf{p}$  be the solution of the dual problem (4.24). For sufficiently large  $N$ , the following estimate holds*

$$\left| \int_{\partial B_R} (\mathcal{T} - \mathcal{T}_N) \boldsymbol{\xi} \cdot \bar{\mathbf{p}} \, ds \right| \lesssim \frac{1}{N} \|\boldsymbol{\xi}\|_{\mathbf{H}^1(\Omega)}^2.$$

**Proof** Using the definitions of the DtN operators  $\mathcal{T}$ ,  $\mathcal{T}_N$  and following the discussion in (4.7), we have

$$\begin{aligned} \left| \int_{\partial B_R} (\mathcal{T} - \mathcal{T}_N) \boldsymbol{\xi} \cdot \bar{\mathbf{p}} \, ds \right| &\leq 2\pi R \sum_{|n|>N} |(M_n \boldsymbol{\xi}_n(R)) \cdot \bar{\mathbf{p}}_n(R)| \\ &\lesssim 2\pi R \sum_{|n|>N} |n| (|\xi_n^r(R)| + |\xi_n^\theta(R)|) (|p_n^r(R)| + |p_n^\theta(R)|) \\ &= 2\pi R \sum_{|n|>N} \frac{|n|}{|n|^{3/2} (1+n^2)^{1/4}} (1+n^2)^{1/4} \\ &\quad \times (|\xi_n^r(R)| + |\xi_n^\theta(R)|) |n|^{3/2} (|p_n^r(R)| + |p_n^\theta(R)|) \\ &\lesssim \frac{1}{N} \left[ \sum_{|n|>N} (1+n^2)^{1/2} (|\xi_n^r(R)| + |\xi_n^\theta(R)|)^2 \right]^{1/2} \\ &\quad \times \left[ \sum_{|n|>N} |n|^3 (|p_n^r(R)| + |p_n^\theta(R)|)^2 \right]^{1/2} \\ &\lesssim N^{-1} \|\boldsymbol{\xi}\|_{H^{1/2}(\partial B_R)} \left[ \sum_{|n|>N} |n|^3 (|p_n^r(R)|^2 + |p_n^\theta(R)|^2) \right]^{1/2} \\ &\lesssim N^{-1} \|\boldsymbol{\xi}\|_{H^1(\Omega)} \left[ \sum_{|n|>N} |n|^3 (|p_n^r(R)|^2 + |p_n^\theta(R)|^2) \right]^{1/2}. \end{aligned} \tag{4.40}$$

Following [29], we let  $t \in [\hat{R}, R]$  and assume, without loss of generality, that  $t$  is closer to the left endpoint  $\hat{R}$  than the right endpoint  $R$ . Denote  $\zeta = R - \hat{R}$ . Then we have  $R - t \geq \frac{\zeta}{2}$ . Thus

$$\begin{aligned} |\xi_n^{(r,\theta)}(t)|^2 &= \frac{1}{R-t} \int_R^t \left( (R-s) |\xi_n^{(r,\theta)}(s)|^2 \right)' \, ds \\ &= \frac{1}{R-t} \int_R^t \left( -|\xi_n^{(r,\theta)}(s)|^2 + 2(R-s) \Re(\xi_n^{(r,\theta)'}(s) \overline{\xi_n^{(r,\theta)}(s)}) \right) \, ds \\ &\leq \frac{1}{R-t} \int_t^R |\xi_n^{(r,\theta)}(s)|^2 \, ds + 2 \int_{\hat{R}}^R |\xi_n^{(r,\theta)}(s)| |\xi_n^{(r,\theta)'}(s)| \, ds, \end{aligned}$$

which implies that

$$\begin{aligned} \|\xi_n^{(r,\theta)}\|_{L^\infty([\hat{R}, R])}^2 &\leq \frac{2}{\xi} \|\xi_n^{(r,\theta)}\|_{L^2([\hat{R}, R])}^2 + 2\|\xi_n^{(r,\theta)}\|_{L^2([\hat{R}, R])} \|\xi_n^{(r,\theta)'}\|_{L^2([\hat{R}, R])} \\ &\leq \left(\frac{2}{\xi} + |n|\right) \|\xi_n^{(r,\theta)}\|_{L^2([\hat{R}, R])}^2 + |n|^{-1} \|\xi_n^{(r,\theta)'}\|_{L^2([\hat{R}, R])}^2. \end{aligned}$$

Using Lemma 4.10 and the Cauchy–Schwarz inequality, we get

$$\begin{aligned} &\sum_{|n|>N} |n|^3 \left( |p_n^r(R)|^2 + |p_n^\theta(R)|^2 \right) \\ &\lesssim \sum_{|n|>N} |n|^3 \left\{ n^2 \left(\frac{\hat{R}}{R}\right)^{2|n|+2} \left( |p_n^r(\hat{R})|^2 + |p_n^\theta(\hat{R})|^2 \right) \right. \\ &\quad \left. + \frac{1}{|n|^2} \left( \|\xi_n^r\|_{L^\infty([\hat{R}, R])}^2 + \|\xi_n^\theta\|_{L^\infty([\hat{R}, R])}^2 \right) \right\} \\ &\lesssim \sum_{|n|>N} |n|^5 \left(\frac{\hat{R}}{R}\right)^{|2n|} \left( |p_n^r(\hat{R})|^2 + |p_n^\theta(\hat{R})|^2 \right) \\ &\quad + \sum_{|n|>N} |n| \left( \|\xi_n^r\|_{L^\infty([\hat{R}, R])}^2 + \|\xi_n^\theta\|_{L^\infty([\hat{R}, R])}^2 \right) \\ &:= I_1 + I_2. \end{aligned}$$

Noting that the function  $t^4 e^{-2t}$  is bounded on  $(0, +\infty)$ , we have

$$\begin{aligned} I_1 &\lesssim \max_{|n|>N} \left( n^4 \left(\frac{\hat{R}}{R}\right)^{|2n|} \right) \sum_{|n|>N} |n| \left( |p_n^r(\hat{R})|^2 + |p_n^\theta(\hat{R})|^2 \right) \\ &\lesssim \|p\|_{H^{1/2}(\partial B_{\hat{R}})}^2 \lesssim \|\xi\|_{H^1(\Omega)}^2, \end{aligned}$$

where the last inequality uses the stability of the dual problem (4.24). For  $I_2$ , we can show that

$$\begin{aligned} I_2 &\lesssim \sum_{|n|>N} \left[ |n| \left(\frac{2}{\xi} + |n|\right) \left( \|\xi_n^r\|_{L^2([\hat{R}, R])}^2 + \|\xi_n^\theta\|_{L^2([\hat{R}, R])}^2 \right) \right. \\ &\quad \left. + \left( \|\xi_n^{r'}\|_{L^2([\hat{R}, R])}^2 + \|\xi_n^{\theta'}\|_{L^2([\hat{R}, R])}^2 \right) \right] \\ &\leq \sum_{|n|>N} \left[ \left(\frac{2}{\xi} |n| + n^2\right) \|\xi_n\|_{L^2([\hat{R}, R])}^2 + \|\xi_n'\|_{L^2([\hat{R}, R])}^2 \right]. \end{aligned}$$

On the other hand, a simple calculation yields

$$\begin{aligned} \|\xi_n^{(r,\theta)}\|_{H^1(B_R \setminus \overline{B_{\hat{R}}})}^2 &= 2\pi \sum_{n \in \mathbb{Z}} \int_{\hat{R}}^R \left[ \left( r + \frac{n^2}{r} \right) |\xi_n^{(r,\theta)}(r)|^2 + r |\xi_n^{(r,\theta)'}(r)|^2 \right] dr \\ &\geq 2\pi \sum_{n \in \mathbb{Z}} \int_{\hat{R}}^R \left[ \left( \hat{R} + \frac{n^2}{R} \right) |\xi_n^{(r,\theta)}(r)|^2 + \hat{R} |\xi_n^{(r,\theta)'}(r)|^2 \right] dr. \end{aligned}$$

It is easy to note that

$$\frac{2}{\zeta} |n| + n^2 \lesssim \hat{R} + \frac{n^2}{R}.$$

Combining the above estimates, we obtain

$$I_2 \lesssim \|\xi\|_{H^1(B_R \setminus B_{\hat{R}})}^2 \leq \|\xi\|_{H^1(\Omega)}^2,$$

which gives

$$\sum_{|n| > N} |n|^3 (|p_n^r(R)| + |p_n^\theta(R)|)^2 \lesssim \|\xi\|_{H^1(\Omega)}^2.$$

Substituting the above inequality into (4.40), we get

$$\left| \int_{\partial B_R} (\mathcal{T} - \mathcal{T}_N) \xi \cdot \bar{p} ds \right| \lesssim \frac{1}{N} \|\xi\|_{H^1(\Omega)}^2, \tag{4.41}$$

which completes the proof. □

Now, we are ready to present the proof of Theorem 4.1.

**Proof** By Lemma 4.6 and Lemma 4.7, we obtain

$$\begin{aligned} \|\xi\|_{H^1(\Omega)}^2 &= \Re b(\xi, \xi) + \Re \int_{\partial B_R} (\mathcal{T} - \mathcal{T}_N) \xi \cdot \bar{\xi} ds \\ &\quad + 2\omega^2 \int_{\Omega} \xi \cdot \bar{\xi} dx + \Re \int_{\partial B_R} \mathcal{T}_N \xi \cdot \bar{\xi} ds \\ &\leq C_1 \left[ \left( \sum_{K \in \mathcal{M}_h} \eta_K^2 \right)^{1/2} + N \left( \frac{\hat{R}}{R} \right)^N \|\mathbf{u}^{\text{inc}}\|_{H^1(\Omega)} \right] \|\xi\|_{H^1(\Omega)} \\ &\quad + (C_2 + C(\delta)) \|\xi\|_{L^2(\Omega)}^2 + \left( \frac{R}{\hat{R}} \right) \delta \|\xi\|_{H^1(\Omega)}^2. \end{aligned}$$

Using (4.2) and choosing  $\delta$  such that  $\frac{R}{\hat{R}} \frac{\delta}{\min(\mu, \omega^2)} < \frac{1}{2}$ , we get

$$\begin{aligned} \|\xi\|_{\mathbf{H}^1(\Omega)}^2 &\leq 2C_1 \left[ \left( \sum_{K \in \mathcal{M}_h} \eta_K^2 \right)^{1/2} + N \left( \frac{\hat{R}}{R} \right)^N \|\mathbf{u}^{\text{inc}}\|_{\mathbf{H}^1(\Omega)} \right] \|\xi\|_{\mathbf{H}^1(\Omega)} \\ &\quad + 2(C_2 + C(\delta)) \|\xi\|_{L^2(\Omega)}^2. \end{aligned} \tag{4.42}$$

It follows from (4.2), (4.16), Lemmas 4.6 and 4.11 that we have

$$\begin{aligned} \|\xi\|_{L^2(\Omega)}^2 &= b(\xi, \mathbf{p}) + \int_{\partial B_R} (\mathcal{T} - \mathcal{T}_N) \xi \cdot \bar{\mathbf{p}} ds - \int_{\partial B_R} (\mathcal{T} - \mathcal{T}_N) \xi \cdot \bar{\mathbf{p}} ds \\ &\lesssim \left[ \left( \sum_{K \in \mathcal{M}_h} \eta_K^2 \right)^{1/2} + N \left( \frac{\hat{R}}{R} \right)^N \|\mathbf{u}^{\text{inc}}\|_{\mathbf{H}^1(\Omega)} \right] \|\xi\|_{\mathbf{H}^1(\Omega)} + \frac{1}{N} \|\xi\|_{\mathbf{H}^1(\Omega)}^2. \end{aligned}$$

Substituting the above estimate into (4.42) and taking sufficiently large  $N$  such that

$$\frac{2(C_2 + C(\delta))}{N} \frac{1}{\min(\mu, \omega^2)} < 1,$$

we obtain

$$\|\mathbf{u} - \mathbf{u}_N^h\|_{\mathbf{H}^1(\Omega)} \lesssim \left( \sum_{K \in \mathcal{M}_h} \eta_K^2 \right)^{1/2} + N \left( \frac{\hat{R}}{R} \right)^N \|\mathbf{u}^{\text{inc}}\|_{\mathbf{H}^1(\Omega)}.$$

which completes the proof of theorem. □

## 5 Implementation and numerical experiments

In this section, we discuss the algorithmic implementation of the adaptive finite element DtN method and present two numerical examples to demonstrate the effectiveness of the proposed method.

### 5.1 Adaptive algorithm

Based on the a posteriori error estimate from Theorem 4.1, we use the FreeFem [26] to implement the adaptive algorithm of the linear finite element formulation. It is shown in Theorem 4.1 that the a posteriori error estimator consists of two parts: the finite element discretization error  $\epsilon_h$  and the DtN truncation error  $\epsilon_N$  which depends on the truncation number  $N$ . Explicitly

$$\epsilon_h = \left( \sum_{T \in \mathcal{M}_h} \eta_T^2 \right)^{1/2}, \quad \epsilon_N = N \left( \frac{\hat{R}}{R} \right)^N \|\mathbf{u}^{\text{inc}}\|_{\mathbf{H}^1(\Omega)}. \tag{5.1}$$



**Table 1** The adaptive finite element DtN method for the elastic wave scattering problem

---

(1)	Given the tolerance $\epsilon > 0, \theta \in (0, 1)$ ;
(2)	Fix the computational domain $\Omega = B_R \setminus \overline{D}$ by choosing the radius $R$ ;
(3)	Choose $\hat{R}$ and $N$ such that $\epsilon_N \leq 10^{-8}$ ;
(4)	Construct an initial triangulation $\mathcal{M}_h$ over $\Omega$ and compute error estimators;
(5)	While $\epsilon_h > \epsilon$ do
(6)	Refine the mesh $\mathcal{M}_h$ according to the strategy: <div style="text-align: center; margin-left: 40px;"> <math>\text{if } \eta_{\hat{T}} &gt; \theta \max_{T \in \mathcal{M}_h} \eta_T, \text{ then refine the element } \hat{T} \in \mathcal{M}_h;</math> </div>
(7)	Denote refined mesh still by $\mathcal{M}_h$ , solve the discrete problem (3.4) on the new mesh $\mathcal{M}_h$ ;
(8)	Compute the corresponding error estimators;
(9)	End while.

---

In the implementation, we choose  $\hat{R}$ ,  $R$ , and  $N$  based on (5.1) to make sure that the finite element discretization error is not polluted by the DtN truncation error, i.e.,  $\epsilon_N$  is required to be very small compared to  $\epsilon_h$ , for example,  $\epsilon_N \leq 10^{-8}$ . For simplicity, in the following numerical experiments,  $\hat{R}$  is chosen such that the obstacle is exactly contained in the disk  $B_{\hat{R}}$ , and  $N$  is taken to be the smallest positive integer such that  $\epsilon_N \leq 10^{-8}$ . The algorithm is shown in Table 1 for the adaptive finite element DtN method for solving the elastic wave scattering problem.

### 5.2 Numerical experiments

We report two examples to demonstrate the performance of the proposed method. The first example is a disk and has an analytical solution; the second example is a U-shaped obstacle which is commonly used to test numerical solutions for the wave scattering problems. In each example, we plot the magnitude of the numerical solution to give an intuition where the mesh should be refined, and also plot the actual mesh obtained by our algorithm to show the agreement. The a posteriori error is plotted against the number of nodal points to show the convergence rate. In the first example, we compare the numerical results by using the uniform and adaptive meshes to illustrate the effectiveness of the adaptive algorithm.

**Example 1** This example is constructed such that it has an exact solution. Let the obstacle  $D = B_{0.5}$  be a disk with radius 0.5 and take  $\Omega = B_1 \setminus \overline{B}_{0.5}$ , i.e.,  $\hat{R} = 0.5, R = 1$ . If we choose the incident wave as


$$\mathbf{u}^{\text{inc}}(\mathbf{x}) = -\frac{\kappa_1 H_0^{(1)'(\kappa_1 r)}}{r} \begin{pmatrix} x \\ y \end{pmatrix} - \frac{\kappa_2 H_0^{(1)'(\kappa_2 r)}}{r} \begin{pmatrix} y \\ -x \end{pmatrix}, \quad r = (x^2 + y^2)^{1/2},$$

then it is easy to check that the exact solution is

$$\mathbf{u}(\mathbf{x}) = \frac{\kappa_1 H_0^{(1)'(\kappa_1 r)}}{r} \begin{pmatrix} x \\ y \end{pmatrix} + \frac{\kappa_2 H_0^{(1)'(\kappa_2 r)}}{r} \begin{pmatrix} y \\ -x \end{pmatrix},$$

**Table 2** Comparison of numerical results using adaptive mesh and uniform mesh refinements for Example 1


Adaptive mesh			Uniform mesh		
DoF <sub>h</sub>	<i>e<sub>h</sub></i>	ϵ <sub>h</sub>	DoF <sub>h</sub>	<i>e<sub>h</sub></i>	ϵ <sub>h</sub>
1745	0.4632	3.9693	1745	0.4632	3.9693
2984	0.3256	2.6723	2667	0.3717	3.2365
5559	0.2253	1.9293	5857	0.2494	2.0625
9030	0.1778	1.5054	10630	0.1851	1.5856
15,407	0.1384	1.1686	20224	0.1330	1.1257




**Fig. 2** Quasi-optimality of the a priori and a posteriori error estimates for Example 1

where  $\kappa_1$  and  $\kappa_2$  are the compressional wave number and shear wave number, respectively.

In Table 2, numerical results are shown for the adaptive mesh refinement and the uniform mesh refinement, where DoF<sub>h</sub> stands for the degree of freedom or the number of nodal points of the mesh  $\mathcal{M}_h$ ,  $\epsilon_h$  is the a posteriori error estimate, and  $e_h = \|\mathbf{u} - \mathbf{u}_N^h\|_{H^1(\Omega)}$  is the a priori error. It can be seen that the adaptive mesh refinement requires fewer DoF<sub>h</sub> than the uniform mesh refinement to reach the same level of accuracy, which shows the advantage of using the adaptive mesh refinement. Figure 2 displays the curves of  $\log e_h$  and  $\log \epsilon_h$  versus  $\log$  DoF<sub>h</sub> for the uniform and adaptive mesh refinements with  $\omega = \pi, \lambda = 2, \mu = 1$ , i.e.,  $\kappa_1 = \pi/2, \kappa_2 = \pi$ . It indicates that the meshes and the associated numerical complexity are quasi-optimal, i.e.,  $\|\mathbf{u} - \mathbf{u}_N^h\|_{H^1(\Omega)} = O(\text{DoF}_h^{-1/2})$  holds asymptotically. Figure 3 plots the magnitude of the numerical solution and an adaptively refined mesh with 15,407 elements. We can see that the solution oscillates on the edge of the obstacle but it is smooth away from the obstacle. This feature is caught by the algorithm. The mesh is adaptively refined around the obstacle and is coarse away from the obstacle.




**Fig. 3** The numerical solution of Example 1. (Left) the magnitude of the numerical solution; (right) an adaptively refined mesh with 15,407 elements



**Fig. 4** Quasi-optimality of the a posteriori error estimates with different frequencies for Example 2

**Example 2** This example does not have an analytical solution. We consider a compressional plane incident wave  $\mathbf{u}^{\text{inc}}(\mathbf{x}) = \mathbf{d}e^{i\kappa_1 \mathbf{x} \cdot \mathbf{d}}$  with the incident direction  $\mathbf{d} = (1, 0)^\top$ . The obstacle is U-shaped and is contained in the rectangular domain  $\{\mathbf{x} \in \mathbb{R}^2 : -2 < x < 2.2, -0.7 < y < 0.7\}$ . Due to the problem geometry, the solution contains singularity around the corners of the obstacle. We take  $R = 3, \hat{R} = 2.31$ . Figure 4 shows the curve of  $\log \epsilon_h$  versus  $\log \text{DoF}_h$  at different frequencies  $\omega = 1, \pi, 2\pi$ . It demonstrates that the decay of the a posteriori error estimates is  $O(\text{DoF}_h^{-1/2})$ . Figure 5 plots the contour of the magnitude of the numerical solution and its corresponding mesh by using the parameters  $\omega = \pi, \lambda = 2, \mu = 1$ . Again, the algorithm does capture the solution feature and adaptively refines the mesh around the corners of the obstacle where the solution displays singularity.



**Fig. 5** The numerical solution of Example 2. (Left) The contour plot of the magnitude of the solution; (right) an adaptively refined mesh with 12,329 elements

## 6 Conclusion

In this paper, we present an adaptive finite element DtN method for the elastic obstacle scattering problem. Based on the Helmholtz decomposition, a new duality argument is developed to obtain the a posteriori error estimate. It does not only take into account of the finite element discretization error but also includes the truncation error of the DtN operator. We show that the truncation error decays exponentially with respect to the truncation parameter. The a posteriori error estimate for the solution of the discrete problem serves as a basis for the adaptive finite element approximation. Numerical results show that the proposed method is accurate and effective. This work provides a viable alternative to the adaptive finite element PML method to solve the elastic obstacle scattering problem. The method can be applied to solve many other wave propagation problems where the transparent boundary conditions are used for open domain truncation. Future work includes extending our analysis to the adaptive finite element DtN method for solving the three-dimensional elastic obstacle scattering problem, where a more complicated transparent boundary condition needs to be considered.

**Acknowledgements** We would like to thank the two anonymous referees for their insightful comments and suggestions that have helped us improve the results of the paper.

## Appendix A. Transparent boundary conditions

In this section, we show the transparent boundary conditions for the scalar potential functions  $\phi^s$ ,  $\psi^s$  and the displacement of the scattered field  $\mathbf{u}^s$  on  $\partial B_R$ .

In the exterior domain  $\mathbb{R}^2 \setminus \bar{B}_R$ , the solutions of the Helmholtz equations (2.5) have the Fourier series expansions in the polar coordinates:

$$\phi^s(r, \theta) = \sum_{n \in \mathbb{Z}} \frac{H_n^{(1)}(\kappa_1 r)}{H_n^{(1)}(\kappa_1 R)} \phi_n^s(R) e^{in\theta}, \quad \psi^s(r, \theta) = \sum_{n \in \mathbb{Z}} \frac{H_n^{(1)}(\kappa_2 r)}{H_n^{(1)}(\kappa_2 R)} \psi_n^s(R) e^{in\theta} \quad (\text{A.1})$$

where  $H_n^{(1)}$  is the Hankel function of the first kind with order  $n$ . Taking the normal derivative of (A.1), we obtain the transparent boundary condition for the scalar potentials  $\phi^s, \psi^s$  on  $\partial B_R$ :

$$\begin{aligned} \partial_r \phi^s &= \mathcal{T}_1 \phi^s := \sum_{n \in \mathbb{Z}} \frac{\kappa_1 H_n^{(1)' }(\kappa_1 R)}{H_n^{(1)}(\kappa_1 R)} \phi_n^s(R) e^{in\theta}, \\ \partial_r \psi^s &= \mathcal{T}_2 \psi^s := \sum_{n \in \mathbb{Z}} \frac{\kappa_2 H_n^{(1)' }(\kappa_2 R)}{H_n^{(1)}(\kappa_2 R)} \psi_n^s(R) e^{in\theta}. \end{aligned} \tag{A.2}$$

The polar coordinates  $(r, \theta)$  are related to the Cartesian coordinates  $\mathbf{x} = (x, y)$  by  $x = r \cos \theta, y = r \sin \theta$  with the local orthonormal basis  $\{\mathbf{e}_r, \mathbf{e}_\theta\}$ , where  $\mathbf{e}_r = (\cos \theta, \sin \theta)^\top, \mathbf{e}_\theta = (-\sin \theta, \cos \theta)^\top$ .

Define a boundary operator for the displacement of the scattered wave

$$\mathcal{B} \mathbf{u}^s = \mu \partial_r \mathbf{u}^s + (\lambda + \mu)(\nabla \cdot \mathbf{u}^s) \mathbf{e}_r \quad \text{on } \partial B_R.$$

Based on the Helmholtz decomposition (2.5) and the transparent boundary condition (A.2), it is shown in [38] that the scattered field  $\mathbf{u}$  satisfies the transparent boundary condition

$$\mathcal{B} \mathbf{u}^s = (\mathcal{T} \mathbf{u}^s)(R, \theta) := \sum_{n \in \mathbb{Z}} M_n \mathbf{u}_n^s(R) e^{in\theta} \quad \text{on } \partial B_R, \tag{A.3}$$

where

$$\mathbf{u}^s(R, \theta) = \sum_{n \in \mathbb{Z}} \mathbf{u}_n^s(R) e^{in\theta} = \sum_{n \in \mathbb{Z}} (u_n^{s,r}(R) \mathbf{e}_r + u_n^{s,\theta}(R) \mathbf{e}_\theta) e^{in\theta}$$

and  $M_n$  is a  $2 \times 2$  matrix defined by

$$M_n = \begin{bmatrix} M_{11}^{(n)} & M_{12}^{(n)} \\ M_{21}^{(n)} & M_{22}^{(n)} \end{bmatrix} = \frac{1}{\Lambda_n(R)} \begin{bmatrix} N_{11}^{(n)} & N_{12}^{(n)} \\ N_{21}^{(n)} & N_{22}^{(n)} \end{bmatrix}. \tag{A.4}$$

Here

$$\Lambda_n(R) = \left(\frac{n}{R}\right)^2 - \alpha_{1n}(R)\alpha_{2n}(R), \quad \alpha_{jn}(R) = \frac{\kappa_j H_n^{(1)' }(\kappa_j R)}{H_n^{(1)}(\kappa_j R)}, \tag{A.5}$$

and

$$\begin{aligned} N_{11}^{(n)} &= \mu \left(\frac{n}{R}\right)^2 \left(\alpha_{2n}(R) - \frac{1}{R}\right) - \alpha_{2n}(R) \left[ (\lambda + 2\mu) \frac{\kappa_1^2 H_n^{(1)' }(\kappa_1 R)}{H_n^{(1)}(\kappa_1 R)} \right. \\ &\quad \left. + (\lambda + \mu) \left(\frac{1}{R} \alpha_{1n}(R) - \left(\frac{n}{R}\right)^2\right) \right], \end{aligned}$$

$$\begin{aligned}
N_{12}^{(n)} &= \mu \frac{in}{R} \alpha_{1n}(R) \left( \alpha_{2n}(R) - \frac{1}{R} \right) - \frac{in}{R} \left[ (\lambda + 2\mu) \frac{\kappa_1^2 H_n^{(1)''}(\kappa_1 R)}{H_n^{(1)}(\kappa_1 R)} \right. \\
&\quad \left. + (\lambda + \mu) \left( \frac{1}{R} \alpha_{1n}(R) - \left( \frac{n}{R} \right)^2 \right) \right], \\
N_{21}^{(n)} &= -\mu \frac{in}{R} \alpha_{2n}(R) \left( \alpha_{1n}(R) - \frac{1}{R} \right) + \mu \frac{in}{R} \frac{\kappa_2^2 H_n^{(1)''}(\kappa_2 R)}{H_n^{(1)}(\kappa_2 R)}, \\
N_{22}^{(n)} &= \mu \left( \frac{n}{R} \right)^2 \left( \alpha_{1n}(R) - \frac{1}{R} \right) - \mu \alpha_{1n}(R) \frac{\kappa_2^2 H_n^{(1)''}(\kappa_2 R)}{H_n^{(1)}(\kappa_2 R)}.
\end{aligned}$$

The matrix entries  $N_{ij}^{(n)}$ ,  $i, j = 1, 2$  can be further simplified. Recall that the Hankel function  $H_n^{(1)}(z)$  satisfies the Bessel differential equation

$$z^2 H_n^{(1)''}(z) + z H_n^{(1)'}(z) + (z^2 - n^2) H_n^{(1)}(z) = 0.$$

We have from straightforward calculations that

$$\begin{aligned}
N_{11}^{(n)} &= -\alpha_{2n}(R) \left[ (\lambda + 2\mu) \left[ -\frac{1}{R^2} \left( \kappa_j R \frac{H_n^{(1)' }(\kappa_j R)}{H_n^{(1)}(\kappa_j R)} + ((\kappa_j R)^2 - n^2) \right) \right] \right. \\
&\quad \left. + (\lambda + \mu) \left( \frac{1}{R} \alpha_{1n}(R) - \left( \frac{n}{R} \right)^2 \right) \right] + \mu \left( \frac{n}{R} \right)^2 \left( \alpha_{2n}(R) - \frac{1}{R} \right) \\
&= -\alpha_{2n}(R) \left[ -\left( \frac{\lambda + 2\mu}{R} \right) \alpha_{1n}(R) - (\lambda + 2\mu) \kappa_1^2 \right. \\
&\quad \left. + (\lambda + 2\mu) \left( \frac{n}{R} \right)^2 + \left( \frac{\lambda + \mu}{R} \right) \alpha_{1n}(R) \right. \\
&\quad \left. - (\lambda + \mu) \left( \frac{n}{R} \right)^2 \right] + \mu \left( \frac{n}{R} \right)^2 \left( \alpha_{2n}(R) - \frac{1}{R} \right) \\
&= -\frac{\mu}{R} \left[ \left( \frac{n}{R} \right)^2 - \alpha_{1n}(R) \alpha_{2n}(R) \right] + \alpha_{2n}(R) \omega^2 \\
&= -\frac{\mu}{R} \Lambda_n(R) + \alpha_{2n}(R) \omega^2, \\
N_{12}^{(n)} &= -\frac{in}{R} \left[ (\lambda + 2\mu) \left[ -\frac{1}{R^2} \left( \kappa_j R \frac{H_n^{(1)' }(\kappa_j R)}{H_n^{(1)}(\kappa_j R)} + ((\kappa_j R)^2 - n^2) \right) \right] \right. \\
&\quad \left. + (\lambda + \mu) \left( \frac{1}{R} \alpha_{1n}(R) - \left( \frac{n}{R} \right)^2 \right) \right] \\
&\quad + \frac{in\mu}{R} \alpha_{1n}(R) \alpha_{2n}(R) - \mu \frac{in}{R^2} \alpha_{1n}(R) \\
&= -\frac{in}{R} \left[ -\frac{\mu}{R} \alpha_{1n}(R) + \mu \left( \frac{n}{R} \right)^2 - (\lambda + 2\mu) \kappa_1^2 \right]
\end{aligned}$$

$$\begin{aligned}
 & + \frac{in\mu}{R} \alpha_{1n}(R) \alpha_{2n}(R) - \frac{in}{R^2} \mu \alpha_{1n}(R) \\
 & = -\frac{in\mu}{R} \Lambda_n(R) + \frac{in}{R} \omega^2, \\
 N_{21}^{(n)} & = -\mu \frac{in}{R} \alpha_{2n}(R) \alpha_{1n}(R) + \frac{in\mu}{R^2} \alpha_{2n}(R) \\
 & \quad + \mu \frac{in}{R} \left( \frac{-1}{R^2} \right) \left( R \alpha_{2n}(R) + (\kappa_2 R)^2 - n^2 \right) \\
 & = -\mu \frac{in}{R} \alpha_{1n}(R) \alpha_{2n}(R) + \frac{in\mu}{R^2} \alpha_{2n}(R) - \mu \frac{in}{R^2} \alpha_{2n}(R) - \frac{in\mu}{R} \kappa_2^2 + i\mu \left( \frac{n}{R} \right)^3 \\
 & = \frac{i\mu n}{R} \Lambda_n(R) - \frac{in}{R} \omega^2, \\
 N_{22}^{(n)} & = \mu \left( \frac{n}{R} \right)^2 \alpha_{1n}(R) - \frac{\mu}{R} \left( \frac{n}{R} \right)^2 - \mu \alpha_{1n}(R) \frac{-1}{R^2} \left( R \alpha_{2n}(R) + (\kappa_2 R)^2 - n^2 \right) \\
 & = \mu \left( \frac{n}{R} \right)^2 \alpha_{1n}(R) - \frac{\mu}{R} \left( \frac{n}{R} \right)^2 + \frac{\mu}{R} \alpha_{1n}(R) \alpha_{2n}(R) \\
 & \quad + \alpha_{1n}(R) \mu \kappa_2^2 - \mu \left( \frac{n}{R} \right)^2 \alpha_{1n}(R) \\
 & = -\frac{\mu}{R} \left( \left( \frac{n}{R} \right)^2 - \alpha_{1n}(R) \alpha_{2n}(R) \right) + \alpha_{1n}(R) \omega^2 \\
 & = -\frac{\mu}{R} \Lambda_n(R) + \alpha_{1n}(R) \omega^2.
 \end{aligned}$$

Substituting the above into (A.3), we obtain

$$\begin{aligned}
 \mathcal{B}u^s & = \mathcal{T}u^s = \sum_{n \in \mathbb{Z}} \frac{1}{\Lambda_n} \left\{ \left[ \left( -\frac{\mu}{R} \Lambda_n(R) + \alpha_{2n}(R) \omega^2 \right) u_n^{s,r}(R) \right. \right. \\
 & \quad + \left( -\frac{in\mu}{R} \Lambda_n(R) + \frac{in}{R} \omega^2 \right) u_n^{s,\theta}(R) \Big] e_r \\
 & \quad + \left[ \left( \frac{i\mu n}{R} \Lambda_n(R) - \frac{in}{R} \omega^2 \right) u_n^{s,r}(R) \right. \\
 & \quad \left. \left. + \left( -\frac{\mu}{R} \Lambda_n(R) + \alpha_{1n}(R) \omega^2 \right) u_n^{s,\theta}(R) \right] e_\theta \right\} e^{in\theta}. \tag{A.6}
 \end{aligned}$$

**Lemma A.1** *Let  $z > 0$ . For sufficiently large  $|n|$ ,  $\Lambda_n(z)$  admits the following asymptotic property*

$$\Lambda_n(z) = \frac{1}{2} (\kappa_1^2 + \kappa_2^2) + \mathcal{O}\left(\frac{1}{|n|}\right).$$

**Proof** Using the asymptotic expansions of the Hankel functions [46]

$$\frac{H_n^{(1)'}(z)}{H_n^{(1)}(z)} = -\frac{|n|}{z} + \frac{z}{2|n|} + \mathcal{O}\left(\frac{1}{|n|^2}\right),$$

we have

$$\alpha_{jn}(z) = \frac{\kappa_j H_n^{(1)' }(\kappa_j z)}{H_n^{(1)}(\kappa_j z)} = -\frac{|n|}{z} + \frac{\kappa_j^2 z}{2|n|} + \mathcal{O}\left(\frac{1}{|n|^2}\right). \quad (\text{A.7})$$

A simple calculation yields that

$$\Lambda_n(z) = \left(\frac{n}{z}\right)^2 - \alpha_{1n}(z)\alpha_{2n}(z) = \frac{1}{2}(\kappa_1^2 + \kappa_2^2) + \mathcal{O}\left(\frac{1}{|n|}\right),$$

which completes the proof.  $\square$

## References

1. Babuška, I., Aziz, A.: Survey lectures on mathematical foundations of the finite element method. In: Aziz, A. (ed.) *The Mathematical Foundation of the Finite Element Method with Application to the Partial Differential Equations*, pp. 5–359. Academic Press, New York (1973)
2. Babuška, I., Werner, C.: Error estimates for adaptive finite element computations. *SIAM J. Numer. Anal.* **15**, 736–754 (1978)
3. Bao, G., Hu, G., Sun, J., Yin, T.: Direct and inverse elastic scattering from anisotropic media. *J. Math. Pures Appl.* **117**, 263–301 (2018)
4. Bao, G., Li, P., Wu, H.: An adaptive edge element method with perfectly matched absorbing layers for wave scattering by periodic structures. *Math. Comput.* **79**, 1–34 (2010)
5. Bao, G., Wu, H.: Convergence analysis of the perfectly matched layer problems for time harmonic Maxwell's equations. *SIAM J. Numer. Anal.* **43**, 2121–2143 (2005)
6. Bayliss, A., Turkel, E.: Radiation boundary conditions for numerical simulation of waves. *Commun. Pure Appl. Math.* **33**, 707–725 (1980)
7. Berenger, J.P.: A perfectly matched layer for the absorption of electromagnetic waves. *J. Comput. Phys.* **114**, 185–200 (1994)
8. Bramble, J.H., Pasciak, J.E., Treneer, D.: Analysis of a finite PML approximation to the three dimensional elastic wave scattering problem. *Math. Comput.* **79**, 2079–2101 (2010)
9. Brenner, S.C., Scott, L.R.: *The mathematical theory of finite element method*. In: *Computers & Mathematics with Applications* (2003)
10. Chen, J., Chen, Z.: An adaptive perfectly matched layer technique for 3-D time-harmonic electromagnetic scattering problems. *Math. Comput.* **77**, 673–698 (2008)
11. Chen, Z., Wu, H.: An adaptive finite element method with perfectly matched absorbing layers for the wave scattering by periodic structures. *SIAM J. Numer. Anal.* **41**, 799–826 (2003)
12. Chen, Z., Liu, X.: An adaptive perfectly matched layer technique for time-harmonic scattering problems. *SIAM J. Numer. Anal.* **43**, 645–671 (2005)
13. Chen, Z., Xiang, X., Zhang, X.: Convergence of the PML method for elastic wave scattering problems. *Math. Comput.* **85**, 2687–2714 (2016)
14. Chew, W., Liu, Q.: Perfectly matched layers for elastodynamics: a new absorbing boundary condition. *J. Comput. Acoust.* **4**, 341–359 (1996)
15. Chew, W., Weedon, W.: A 3D perfectly matched medium for modified Maxwell's equations with stretched coordinates. *Microwave Opt. Technol. Lett.* **13**, 599–604 (1994)
16. Ciarlet, P.G.: *Mathematical Elasticity, Vol. I: Three-Dimensional Elasticity, Studies in Mathematics and Its Applications*. North-Holland, Amsterdam (1988)
17. Collino, F., Tsogka, C.: Application of the PML absorbing layer model to the linear elastodynamics problem in anisotropic heterogeneous media. *Geophysics* **66**, 294–307 (2001)
18. Colton, D., Kress, R.: *Integral Equation Methods in Scattering Theory*. Wiley, New York (1983)
19. Engquist, B., Majda, A.: Absorbing boundary conditions for the numerical simulation of waves. *Math. Comput.* **31**, 629–651 (1977)
20. Gächter, G.K., Grote, M.J.: Dirichlet-to-Neumann map for three-dimensional elastic waves. *Wave Motion* **37**, 293–311 (2003)



21. Givoli, D., Keller, J.B.: Non-reflecting boundary conditions for elastic waves. *Wave Motion* **12**, 261–279 (1990)
22. Grote, M., Kirsch, C.: Dirichlet-to-Neumann boundary conditions for multiple scattering problems. *J. Comput. Phys.* **201**, 630–650 (2004)
23. Grote, M., Keller, J.: On nonreflecting boundary conditions. *J. Comput. Phys.* **122**, 231–243 (1995)
24. Hagstrom, T.: Radiation boundary conditions for the numerical simulation of waves. *Acta Numer.* **8**, 47–106 (1999)
25. Hastings, F.D., Schneider, J.B., Broschat, S.L.: Application of the perfectly matched layer (PML) absorbing boundary condition to elastic wave propagation. *J. Acoust. Soc. Am.* **100**, 3061–3069 (1996)
26. Hecht, F.: New development in FreeFem++. *J. Numer. Math.* **20**, 251–265 (2012)
27. Hohage, T., Schmidt, F., Zschiedrich, L.: Solving time-harmonic scattering problems based on the pole condition. II: convergence of the PML method. *SIAM J. Math. Anal.* **35**, 547–560 (2003)
28. Hsiao, G.C., Nigam, N., Pasiak, J.E., Xu, L.: Error analysis of the DtN-FEM for the scattering problem in acoustic via Fourier analysis. *J. Comput. Appl. Math.* **235**, 4949–4965 (2011)
29. Jiang, X., Li, P., Lv, J., Zheng, W.: An adaptive finite element method for the wave scattering with transparent boundary condition. *J. Sci. Comput.* **72**, 936–956 (2017)
30. Jiang, X., Li, P.: An adaptive finite element PML method for the acoustic-elastic interaction in three dimensions. *Commun. Comput. Phys.* **22**, 1486–1507 (2017)
31. Jiang, X., Li, P., Lv, J., Wang, Z., Wu, H., Zheng, W.: An adaptive finite element DtN method for Maxwell's equations in bi-periodic structures. *IMA J. Numer. Anal.* <https://doi.org/10.1093/imanum/drab052>
32. Jiang, X., Li, P., Zheng, W.: Numerical solution of acoustic scattering by an adaptive DtN finite element method. *Commun. Comput. Phys.* **13**, 1277–1244 (2013)
33. Jiang, X., Li, P., Lv, J., Zheng, W.: An adaptive finite element PML method for the elastic wave scattering problem in periodic structures. *ESAIM Math. Model. Numer. Anal.* **51**, 2017–2047 (2017)
34. Jiang, X., Li, P., Lv, J., Zheng, W.: Convergence of the PML solution for elastic wave scattering by bi-periodic structures. *Commun. Math. Sci.* **16**, 985–1014 (2018)
35. Komatitsch, D., Tromp, J.: A perfectly matched layer absorbing boundary condition for the second-order seismic wave equation. *Geophys. J. Int.* **154**, 146–153 (2003)
36. Landau, L.D., Lifshitz, E.M.: *Theory of Elasticity*, *Theory of Elasticity*. Pergamon Press, Oxford (1986)
37. Lenoir, M.: Optimal isoparametric finite elements and error estimates for domains involving curved boundaries. *SIAM J. Numer. Anal.* **23**, 562–580 (1986)
38. Li, P., Wang, Y., Wang, Z., Zhao, Y.: Inverse obstacle scattering for elastic waves. *Inverse Prob.* **32**, 115018 (2016)
39. Li, P., Yuan, X.: Inverse obstacle scattering for elastic waves in three dimensions. *Inverse Probl. Imaging* **13**, 545–573 (2019)
40. Li, Y., Zheng, W., Zhu, X.: A CIP-FEM for high frequency scattering problem with the truncated DtN boundary condition. *CSIAM Trans. Appl. Math.* **1**, 530–560 (2020)
41. Monk, P.: *Finite Element Methods for Maxwell's Equations*. Oxford University Press, Oxford (2003)
42. Nédélec, J.-C.: *Acoustic and Electromagnetic Equations Integral Representations for Harmonic Problems*. Springer, New York (2001)
43. Schatz, A.H.: An observation concerning Ritz-Galerkin methods with indefinite bilinear forms. *Math. Comput.* **28**, 959–962 (1974)
44. Turkel, E., Yefet, A.: Absorbing PML boundary layers for wave-like equations. *Appl. Numer. Math.* **27**, 533–557 (1998)
45. Wang, Z., Bao, G., Li, J., Li, P., Wu, H.: An adaptive finite element method for the diffraction grating problem with transparent boundary condition. *SIAM J. Numer. Anal.* **53**, 1585–1607 (2015)
46. Watson, G.N.: *A Treatise on the Theory of Bessel Functions*. Cambridge University Press, Cambridge (1995)
47. Yuan, X., Bao, G., Li, P.: An adaptive finite element DtN method for the open cavity scattering problems. *CSIAM Trans. Appl. Math.* **1**, 316–345 (2020)
48. Zhou, W., Wu, H.: An adaptive finite element method for the diffraction grating problem with PML and few-mode DtN truncations. *J. Sci. Comput.* **76**, 1813–1838 (2018)