

The Lorenz Attractor is Mixing

Stefano Luzzatto¹, Ian Melbourne², Frederic Paccaut³

¹ Department of Mathematics, Imperial College London, SW7 2AZ, U.K.

² Department of Mathematics and Statistics, University of Surrey, Guildford GU2 5XH, U.K.

³ LAMFA CNRS UMR 6140, University of Picardie, 33, rue Saint Leu, 80039 Amiens, France

Received: 8 October 2004 / Accepted: 25 March 2005
Published online: 9 August 2005 – © Springer-Verlag 2005

Abstract: We study a class of geometric Lorenz flows, introduced independently by Afraimovič, Bykov & Sil'nikov and by Guckenheimer & Williams, and give a verifiable condition for such flows to be mixing. As a consequence, we show that the classical Lorenz attractor is mixing.

1. Introduction and Statement of Results

1.1. The Lorenz equations. Many systems of nonlinear differential equations that were first studied almost 50 years ago and which were motivated mainly by problems in geophysical and astrophysical fluid dynamics and dynamical meteorology [2, 8, 15, 23], remain difficult to understand rigorously to the present day. In 1963, Lorenz [17] introduced the following system of differential equations:

$$\begin{aligned}\dot{x} &= 10(y - x), \\ \dot{y} &= 28x - y - xz, \\ \dot{z} &= xy - \frac{8}{3}z.\end{aligned}\tag{1.1}$$

Approximate numerical studies of these equations led Lorenz to emphasise the possibility and importance of *sensitive dependence on initial conditions* even in such simplified models of natural phenomena. A combination of results obtained over the last 25 years [1, 14, 20, 25, 30] and culminating in the work of Tucker [27, 28] gives the following statement (see [26, 29] for detailed surveys):

The Lorenz equations admit a robust attractor A which supports a “physical” ergodic invariant probability measure ν with a positive Lyapunov exponent.

Recall that the measure ν is called physical, or Sinai-Bowen-Ruelle (SRB), if for Lebesgue almost every solution $u(t) \in \mathbb{R}^3$ starting close to A and all continuous functions $h : \mathbb{R}^3 \rightarrow \mathbb{R}$,

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T h(u(t)) dt = \int_A h d\nu.$$

In this paper we take a further step in the understanding of the statistical properties of the Lorenz attractor. A measure ν is mixing for a flow Φ_t if

$$\nu(\Phi_t(A) \cap B) \rightarrow \nu(A)\nu(B)$$

for all measurable sets A, B , as $t \rightarrow \infty$. We say that the Lorenz attractor is mixing if the SRB measure ν mentioned above is mixing.

We prove the following

Theorem 1. *The Lorenz attractor is mixing.*

In fact, our result shows that the Lorenz attractor is *stably mixing*: sufficiently small C^1 perturbations of the flow are mixing.

There are relatively few explicit examples of flows that have been proved to be mixing. Anosov [4] showed that geodesic flows for compact manifolds of negative curvature are mixing, and this was generalised [16] to include contact flows. Moreover, mixing persists under C^1 perturbations. For codimension one Anosov flows [21] and for Anosov flows with a global infranil cross-section [5], the set of mixing flows is C^1 open, but the corresponding result for general Anosov flows is not known. Recently [11] it was shown that mixing holds for a C^2 -open and C^r -dense set of C^r uniformly hyperbolic (Axiom A) flows for all $r \geq 2$. (In these references, mixing is proved for any equilibrium measure for a Hölder potential.) However, the conditions for stable mixing in [11] are not easily verifiable.

The Lorenz attractor is an example of a *singular hyperbolic attractor* [18] (uniformly hyperbolic, except for a singularity due to the attractor containing an equilibrium). Somewhat surprisingly, we show that the singular nature of the Lorenz attractor assists in the search for a verifiable condition for mixing.

1.2. Geometric Lorenz attractors. Afraïmovič, Bykov and Sil'nikov [1] and Guckenheimer and Williams [14, 30] introduced a geometric model that is an abstraction of the numerically-observed features possessed by solutions to (1.1). Tucker [27, 28] proved that the geometric model is valid, so the Lorenz equations define a geometric Lorenz flow.

Accordingly, our approach to Theorem 1 is to establish mixing for geometric Lorenz flows satisfying certain hypotheses, and then to verify from [27, 28] that the hypotheses are satisfied for the Lorenz equations. Roughly speaking, a *geometric Lorenz flow* is the natural extension of a *geometric Lorenz semiflow* which is itself a suspension flow built over a certain type of one-dimensional expanding map f . Moreover, the roof function r has a logarithmic singularity (due to the equilibrium for the flow). Precise definitions are given in Sect. 2. In the literature, a standard assumption is that the map f is *locally eventually onto* (*l.e.o.*), see Definition 3. We prove that this is a sufficient condition for the corresponding flow to be mixing.

Theorem 2. *Let Φ_t be a geometric Lorenz flow. Suppose that the associated one-dimensional map f is l.e.o. Then Φ_t is mixing (and even Bernoulli).*

Remark 1.1. It was shown in [19] that geometric Lorenz flows are generically mixing (albeit in the C^1 topology). Stability of mixing is not proved in [19], nor is it shown there that the actual Lorenz attractor is mixing.

1.3. Outline of the proof of Theorem 2. The geometric Lorenz flow Φ_t possesses a strong stable foliation and quotienting by the foliation yields a geometric Lorenz semiflow f_t . It suffices to prove that the semiflow f_t is mixing. Moreover, weak mixing of f_t implies that the original flow is mixing (in fact Bernoulli) by a general result of Ratner [22].

As mentioned above, f_t is a suspension semiflow over a one-dimensional map f , where the roof function r possesses a logarithmic singularity. Label the singular point 0. Using spectral characterisations of weak mixing, it suffices to show that the cohomological equation $\psi \circ f = e^{iar} \psi$ has no measurable solutions for all $a > 0$. A recent Livšić regularity theorem by Bruin *et al.* [6] shows that if such an eigenfunction ψ exists, then it must be continuous. This, coupled with the regularity of f and r away from 0 and the singularity of r at 0, is used to obtain a contradiction.

The remainder of the paper is structured as follows. In Sect. 2, we discuss the geometric Lorenz flow and its relation to the Lorenz attractor. In particular, we indicate how Theorem 1 follows from Theorem 2. In Sect. 3, we prove Theorem 2. In Sect. 4, we discuss extensions of our main results and some related future directions.

2. The Lorenz Attractor is a Geometric Lorenz Flow

In this section we collect and organise several results from the existing literature on the relation between the Lorenz attractor and geometric Lorenz flows. We recall first of all some basic relevant facts about the Lorenz equations (1.1), see [26]. The origin is an equilibrium of saddle type with two negative (stable) and one positive (unstable) eigenvalues $\lambda_{ss} < \lambda_s < 0 < \lambda_u$. It is also the case that $\lambda_u > |\lambda_s|$.

Suppose that a finite number of nonresonance conditions are satisfied so that the vector field is smoothly linearisable in a neighbourhood of 0. In these coordinates, the flow near 0 has the form $(x_1, x_2, x_3) \mapsto (e^{\lambda_u t} x_1, e^{\lambda_{ss} t} x_2, e^{\lambda_s t} x_3)$. By a linear rescaling, we can suppose that the domain of linearisation of the flow includes the cube $[-1, 1]^3$. Define $\Sigma = \{(x_1, x_2, x_3) : |x_1|, |x_2| \leq 1, x_3 = 1\}$ and $\tilde{\Sigma} = \{(x_1, x_2, x_3) : x_1 = \pm 1, |x_2|, |x_3| \leq 1\}$. Then we define the first hit map $P_0 : \Sigma \rightarrow \tilde{\Sigma}$ by

$$P_0(x_1, x_2, 1) = (e^{\lambda_u t_0} x_1, e^{\lambda_{ss} t_0} x_2, e^{\lambda_s t_0}) = (\text{sgn } x_1, \tilde{x}_2, \tilde{x}_3),$$

where t_0 is the “time of flight” from Σ to $\tilde{\Sigma}$. Solving $e^{\lambda_u t_0} x_1 = \text{sgn } x_1$ for $t_0 = -(\ln |x_1|)/\lambda_u$, we obtain

$$P_0(x_1, x_2, 1) = (\text{sgn } x_1, |x_1|^\beta x_2, |x_1|^\alpha),$$

where $\alpha = |\lambda_s|/\lambda_u \in (0, 1)$ and $\beta = |\lambda_{ss}|/\lambda_u > 0$. Note that $P_0 : \Sigma \rightarrow \tilde{\Sigma}$ is well-defined on $\Sigma \setminus W^s(0)$, where $W^s(0) = \{(0, x_2, 1) : |x_2| \leq 1\}$, and that t_0 has a logarithmic singularity at $x_1 = 0$.

Tucker [27, 28] proves that there exists a compact trapping region $N \subset \Sigma$ such that the Poincaré first return map $P : N \setminus W^s(0) \rightarrow N$ is well defined. We can decompose

$P = P_1 \circ P_0$, where P_1 is the first-hit map between $\tilde{\Sigma}$ and Σ . Note that P_1 is a diffeomorphism where it is defined, and that the time of flight t_1 for P_1 is bounded. Hence the time of flight $t = t_0 + t_1$ for the full return map P is smooth except for a logarithmic singularity at $x_1 = 0$. Moreover, the following crucial *hyperbolicity estimate* holds.

Lemma 2.1 ([28]). *The return map P admits a forward invariant cone field. In other words, there exists a cone $\mathcal{C}(u)$ inside Σ at each point $u \in N \setminus W^s(0)$ such that $(dP)_u \mathcal{C}(u)$ is strictly contained inside $\mathcal{C}(Pu)$.*

Moreover, there exist constants $c > 0$, $\tau > 1$, such that for each $u \in N \setminus W^s(0)$,

$$\|(dP^n)_u v\| \geq c\tau^n \|v\|,$$

for every $v \in \mathcal{C}(u)$ and $n \geq 1$.

A consequence of Lemma 2.1 is that the return map P has an invariant $C^{1+\varepsilon}$ stable foliation for some $\varepsilon > 0$. Let $I = [-1, 1]$. We obtain a singular one-dimensional map $f : I \rightarrow I$ by quotienting along stable leaves. Let $J = I \setminus \{0\}$. Then it is immediate that on J :

- f(i)** f is $C^{1+\varepsilon}$,
- f(ii)** $|f^{(n)}| \geq c\tau^n$ for all $n \geq 1$,
- f(iii)** $C^{-1}|x|^{\alpha-1} \leq f'(x) \leq C|x|^{\alpha-1}$,

where $c > 0$, $\tau > 1$ are the constants in Lemma 2.1, $\alpha = |\lambda_s|/\lambda_u \in (0, 1)$, and $C \geq 1$ is a constant.

Remark 2.2. An explicit construction of f can be obtained as follows. Let $I \subset N$ denote a global cross-section to the stable foliation with $0 \in I$. Coordinates on I can be chosen so that $I = [-1, 1]$. For each $x \in I$, let $W^s(x) \subset N$ denote the corresponding stable leaf. Then $P(W^s(x)) \subset W^s(Px)$. Define f_x to be the unique intersection point of $W^s(Px)$ with I .

The underlying flow Φ_t possesses an invariant strong stable foliation corresponding to the stable foliation for P . Quotienting the flow along strong stable leaves yields a semiflow f_t with (noninvertible) Poincaré map $f : I \rightarrow I$ as above and return time function $r : I \rightarrow \mathbb{R}^+$ with a logarithmic singularity at 0. More precisely, r satisfies

- r(i)** $r(x) \rightarrow \infty$ as $x \rightarrow 0$,
- r(ii)** $|r(x) - r(y)| \leq C|\ln|x| - \ln|y||$ for all $x, y > 0$ and all $x, y < 0$.

Note that it follows from **r(ii)** that r is Lebesgue integrable and r is C^1 on J .

We have described how to pass from the original three-dimensional flow defined by the Lorenz equations (1.1) to a two-dimensional Poincaré map $P : \Sigma \rightarrow \Sigma$ and then to a one-dimensional expanding map $f : I \rightarrow I$. The process is reversible: taking the natural extension of $f : I \rightarrow I$ recovers the Poincaré map $P : N \rightarrow N$ and taking the suspension of the P by the roof function $t : N \rightarrow \mathbb{R}^+$ recovers the original Lorenz flow. Alternatively, these steps can be carried out in the opposite order to recover the Lorenz flow as the natural extension of the suspension semiflow of the map $f : I \rightarrow \mathbb{R}$ by the roof function $r : I \rightarrow \mathbb{R}^+$. The latter viewpoint is the *geometric Lorenz flow* construction [14, 30]. For completeness, the notions of suspension and natural extension are recalled at the end of this section.

Definition 1. *Let $f : I \rightarrow I$. Assume that $f(0)$ is undefined, with $f(0^+) = -1$ and $f(0^-) = +1$. It is assumed that $f(1) \in (0, 1)$ and $f(-1) \in (-1, 0)$. (See Fig. 1.) If conditions **f(i)–f(iii)** are satisfied, then f is a **Lorenz-like expanding map**.*




Fig. 1. The graph of a Lorenz-like expanding map $f : [-1, 1] \rightarrow [-1, 1]$

Definition 2. A semiflow f_t is called a geometric Lorenz semiflow if it is the suspension over a Lorenz-like expanding map $f : I \rightarrow I$ by a roof function $r : I \rightarrow \mathbb{R}^+$ satisfying conditions **r(i)**, **r(ii)**. A flow Φ_t is called a **geometric Lorenz flow** if it is the natural extension of a geometric Lorenz semiflow.

The results of Tucker described above imply that the Lorenz equations indeed define a geometric Lorenz flow. Moreover, as outlined below, Tucker showed that the associated expanding map $f : I \rightarrow I$ is transitive for the Lorenz equations, thus establishing the existence of the Lorenz attractor.

In the following definition, we stress that $f(0)$ is undefined. In particular, the endpoints ± 1 have no preimages.

Definition 3. The map $f : I \rightarrow I$ is **locally eventually onto (l.e.o.)** if for any open set $U \subset J$, there exists $k \geq 0$ such that $f^k U$ contains $(0, 1)$.

Remark 2.3. Since $f(0, 1)$ contains $(-1, 0)$ and vice versa, we could equally use $(-1, 0)$ instead of $(0, 1)$ in the definition of l.e.o. Clearly, the l.e.o. property implies that f is topologically transitive. The exact formulation of l.e.o. varies in the literature, but our definition agrees with the one in [12].

Remark 2.4. It was anticipated in [30] that **f(ii)** would hold with $c = 1, \tau > \sqrt{2}$. It is easy to check that this is a sufficient condition for f to be l.e.o.

Surprisingly [28], for the actual Lorenz equations it turns out that $f'(x) < 1$ for certain values of $x \in I$. Nevertheless, it is still the case that f is l.e.o., see Tucker [28].

It can be shown that Lorenz-like expanding maps satisfying the l.e.o. condition have a unique ergodic probability measure μ that is equivalent to Lebesgue (see for example Sect. 3). The suspended measure $\nu = \mu^r$ defines an SRB measure for the geometric Lorenz flow. In particular, Theorem 1 follows from Theorem 2. The latter result is proved in Sect. 3.

Remark 2.5. The approach in this paper establishes weak mixing for geometric Lorenz semiflows. It then follows from Ratner [22] that such semiflows, and the corresponding flows, are Bernoulli. In particular, they are mixing.

Suspensions and natural extensions. As promised, we end this section by recalling the notions of suspension and natural extension. Let (X, μ) be a probability space and $f : X \rightarrow X$ a (noninvertible) measure preserving transformation. Let $r : X \rightarrow \mathbb{R}^+$ be

an L^1 roof function. Define the suspension $X^r = \{(x, u) : x \in X, u \in [0, r(x)]\} / \sim$ where $(x, r(x)) \sim (fx, 0)$. Form the suspension semiflow $f_t : X^r \rightarrow X^r$ given by $f_t(x, u) = (x, u + t)$ computed modulo identifications. An invariant probability measure is given by $\mu^r = \mu \times \ell / \int r \, dm$, where ℓ is Lebesgue measure.

Next suppose that $f_t : \Omega \rightarrow \Omega$ is a semiflow preserving a probability measure ν . Form the natural extension (or inverse limit) $\widehat{\Omega}$ consisting of curves $\{x(s)\}_{s \geq 0}$ in Ω satisfying $f_t(x(s)) = x(s - t)$ for all $s \geq t \geq 0$. An invertible flow $\Phi_t : \widehat{\Omega} \rightarrow \widehat{\Omega}$ is given by $\Phi_t\{x(s)\}_{s \geq 0} = \{x(s + t)\}_{s \geq 0}$. The projection $\pi : \widehat{\Omega} \rightarrow \Omega, \{x(s)\}_{s \geq 0} \mapsto x(0)$ defines a semiconjugacy between Φ_t and f_t , and there is a unique Φ_t -invariant measure ν on $\widehat{\Omega}$ such that π is measure preserving.

3. Mixing for Geometric Lorenz Flows

In this section, we prove Theorem 2. Namely, we show that if $f : I \rightarrow I$ is an expanding map of Lorenz-type (satisfying **f(i)**–**f(iii)**) and $r : I \rightarrow \mathbb{R}^+$ is a logarithmic roof function (satisfying **r(i)**, **r(ii)**), then the corresponding geometric Lorenz flow is mixing. By Remark 2.5, it is enough to prove that the corresponding geometric Lorenz semiflow is weak mixing.

Let $f : I \rightarrow I$ be a Lorenz-like expanding map. As shown in [9] and [10], f admits an induced map F that is Gibbs-Markov. More precisely there is an open interval $Y \subset I$ containing 0 and (modulo a set of Lebesgue measure zero) there is a partition $\mathcal{P} = \{\omega\}$ of Y consisting of intervals, and a return function $R : Y \rightarrow \mathbb{N}$ constant on partition elements such that the induced map $F(x) = f^{R(x)}(x)$ restricts to a diffeomorphism $F|_\omega : \omega \rightarrow Y$ for each partition element ω , and such that the following conditions are satisfied:

- (a) There exists $\lambda > 1$ such that $|DF_\omega| \geq \lambda$ for each ω .
- (b) For all ω , $\log g_\omega$ is Hölder (uniformly in ω), where g is the Jacobian of the inverse of $F|_\omega : \omega \rightarrow Y$.
- (c) R is Lebesgue integrable.

A standard Folklore Theorem in dynamics says that conditions (a)–(c) imply the existence of a unique ergodic invariant measure μ for $f : I \rightarrow I$ that is absolutely continuous with respect to Lebesgue and whose support includes Y . In addition, the construction of [9, 10] satisfies

- (d) $0 \notin \overline{f^k \omega}$ for $0 \leq k < R(\omega)$ for each ω .

The following Livšic regularity result is due to Bruin *et al.* [6].

Lemma 3.1. *Suppose that $f : I \rightarrow I$ admits an induced map $F : Y \rightarrow Y$ satisfying conditions (a)–(d) above, with absolutely continuous ergodic measure μ . Let $r : I \rightarrow \mathbb{R}^+$ be a roof function satisfying condition **r(ii)**. Let $\psi : I \rightarrow S^1$ be a μ -measurable function satisfying*

$$e^{ir} = (\psi \circ f)\psi^{-1}, \quad a.e.$$

Then ψ has a version that is Hölder on Y .

Sketch proof. We provide the details required for the reader to pass between the formulation in [6] and the statement of the lemma. By conditions (a)–(c), the map $f : I \rightarrow I$ can be modelled by a Young tower [32] with base Y . The appropriate measure for the

tower has a Hölder density, so by condition (d), Bruin *et al.* [6, Theorem 2] applies to guarantee that ψ is Hölder on any fixed partition element of Y and hence on the whole of Y ([6, Remark 3]). The tower metric in [6] coincides with the Euclidean metric when restricted to Y , so ψ is Hölder in the original metric. \square

Remark 3.2. Related Livšic regularity results can be found in [13]. The arguments in [6, 13] are different from each other leading to distinct results, and the approach in [6] turns out to more convenient for our purposes.

Lemma 3.3. *Suppose that the geometric Lorenz semiflow f_t is not weak mixing. Then there exists a constant $a > 0$ and a measurable eigenfunction $\psi : X \rightarrow S^1$ continuous on $\bigcup_{k \geq 0} f^k Y$, such that*

$$e^{iar} = (\psi \circ f)\psi^{-1} \text{ a.e.} \tag{3.1}$$

Proof. It can be taken as the definition of weak mixing that $\phi \circ f_t = e^{iat}\phi$ has no measurable solutions $\phi : X^r \rightarrow S^1$ for $a > 0$. Suppose that ϕ is such a measurable solution. It follows from Fubini that there exists $\epsilon > 0$ with $r > \epsilon$ a.e. such that $\phi \circ f_t(x, \epsilon) = e^{iat}\phi(x, \epsilon)$ for almost every $x \in X$. Set $t = r(x)$ and $\psi(x) = \phi(x, \epsilon)$. Since $f_{r(x)}(x, \epsilon) = (fx, \epsilon)$ we obtain that $\psi \circ f = e^{iar}\psi$. Thus ψ is a measurable solution to Eq. (3.1). By Lemma 3.1, there is a solution ψ that is Hölder continuous on Y .

It is now straightforward to show that ψ is continuous on $\bigcup_{k \geq 0} f^k Y$. Suppose that $z = f^k y$, where $y \in Y$. Since $f(0)$ is undefined, it is certainly the case that $f^j y \neq 0$ for $j = 0, \dots, k - 1$. Hence we can choose an open set $U \subset Y$ containing y such that $0 \notin f^j U$ for $0 \leq j \leq k - 1$. In particular, there exists $\gamma \in (0, 1)$ such that ψ is C^γ on U and at the same time e^{iar_k} is C^1 on U .

Let $z_i = f^k y_i$ where $y_i \in U$ for $i = 1, 2$. Applying Eq. (3.1) in the form $\psi \circ f^k = e^{iar_k}\psi$, we obtain

$$\psi(z_1)\psi(z_2)^{-1} = \psi(y_1)\psi(y_2)^{-1} e^{iar_k(y_1)} e^{-iar_k(y_2)},$$

so by **f(ii)**,

$$|\psi(z_1)\psi(z_2)^{-1}| \leq D|y_1 - y_2|^\gamma \leq D(c\lambda^k)^{-\gamma}|z_1 - z_2|^\gamma.$$

Hence ψ is Hölder on $f^k U$ and so is certainly continuous at z as required. \square

Proof of Theorem 2. Suppose that f_t is not weak mixing. By Lemma 3.3, there exists $a > 0$ and a measurable eigenfunction $\psi : I \rightarrow S^1$ satisfying (3.1), such that ψ is continuous on $\bigcup_{k \geq 0} f^k Y$. Since f is l.e.o., ψ is continuous on $(-1, 1)$. In particular, ψ is continuous at $f(\pm 1)$.

Iterate equation (3.1) to obtain

$$e^{iar} e^{iar \circ f} = (\psi \circ f^2)\psi^{-1}. \tag{3.2}$$

We evaluate this equation along a sequence $x_n > 0$ with $x_n \rightarrow 0$. We have the limits $\psi(x_n) \rightarrow \psi(0)$, $\psi(f^2 x_n) \rightarrow \psi(f(-1))$, and $r(fx_n) \rightarrow r(-1)$. Hence, the right-hand-side of Eq. (3.2) and the second factor on the left-hand-side converge as $n \rightarrow \infty$. To obtain a contradiction, it suffices to choose $x_n \rightarrow 0$ so that $e^{iar(x_n)}$ does not converge. For n sufficiently large, choose $b_n > r(\epsilon/2^n)$ such that $e^{iab_n} = (-1)^n$. Since r is continuous on $I \setminus \{0\}$ and $r(x) \rightarrow \infty$ as $x \rightarrow 0$, there exists $x_n > 0$ such that $r(x_n) = b_n$. As required, $x_n \rightarrow 0$ and $e^{iar(x_n)}$ does not converge. \square

4. Extensions and Future Directions

In this paper, we have focused on geometric Lorenz attractors satisfying a locally eventually onto condition. In this section, we discuss how to relax the l.e.o. condition, and we describe how our results extend to larger classes of singular attractors. We end by mentioning some future directions.

The l.e.o. assumption occurs in much of the literature since it is a verifiable property of the flow that guarantees topological transitivity of f . Moreover, it holds for the Lorenz equations (1.1) themselves [28]. However, topological transitivity for $f : I \rightarrow I$ is not crucial in this paper. We could simply work with the unique ergodic component whose support includes the set Y introduced in Sect. 3. Passing to the natural extension of the suspension over this component yields a geometric Lorenz attractor with an SRB measure as before.

The proof of mixing for this attractor also relies to some extent on the l.e.o. property, but it is clear that the proof goes through under a much weaker condition: namely that a preimage of a forward image of 1 or -1 lies in Y . (That is, there exist integers $k, \ell \geq 0$ such that $f^\ell(1) \in f^k Y$ or $f^\ell(-1) \in f^k Y$.) This condition is open and dense within the class of Lorenz-like expanding maps, and in principle, it can be verified by a finite computation.

Next, we note that the specific structure of Lorenz-like expanding maps is not so crucial for the proof of mixing. Properties **f(i)**–**f(iii)** could be relaxed, or altered completely, provided f admits an induced map $F : Y \rightarrow Y$ satisfying conditions (a)–(d). The existence of a Gibbs-Markov induced map is a very general condition, see [7, 3]. It clearly holds for uniformly expanding maps and, we have made use of the recent work [9, 10] for Lorenz-like expanding maps.

Condition **f(ii)** in the definition of the Lorenz-like expanding map corresponds to the eigenvalue condition $\lambda_u > |\lambda_s|$ which is valid for the Lorenz attractor. The resulting class of geometric Lorenz attractors are sometimes called expanding. There are also *contracting geometric Lorenz attractors* [24] that arise when $\lambda_u < |\lambda_s|$. It seems likely that such attractors can be shown to be mixing using the ideas in this paper.

In situations where Lemma 2.1 fails, it may not be possible to reduce to a one-dimensional expanding map. However, it is plausible that the Poincaré map P could be modelled by an “unquotiented” Young tower as in [31] to which the ideas in this paper might still be applicable.

In this paper, we have established mixing. An interesting open question is to prove results on the speed of mixing. In a different direction, it would be interesting to derive statistical limit laws such as the central limit theorem and invariance principles for the Lorenz attractor.

Acknowledgements. This research was supported in part by EPSRC Grant GR/S11862/01. IM is greatly indebted to the University of Houston for the use of e-mail, given that pine is currently not supported on the University of Surrey network.

References

1. Afraïmovič, V.S., Bykov, V.V., Sil’nikov, L.P.: The origin and structure of the Lorenz attractor. Dokl. Akad. Nauk SSSR **234**, 336–339 (1977)
2. Allan, D.W.: On the behaviour of systems of coupled dynamos. Proc. Cambridge Philos. Soc. **58**, 671–693 (1962)
3. Alves, J.F., Luzzatto, S., Pinheiro, V.: Lyapunov exponents and rates of mixing for one-dimensional maps. Ergodic Th. & Dyn. Syst. **24**, 637–657 (2004)

4. Anosov, D.V.: Geodesic flows on closed Riemann manifolds with negative curvature. Proceedings of the Steklov Institute of Mathematics, No. 90 (1967). Translated from the Russian by S. Feder Providence, R.I.: American Mathematical Society, 1969
5. Bowen, R.: Mixing Anosov flows. *Topology* **15**, 77–79 (1976)
6. Bruin, H., Holland, M., Nicol, M.: Livsic regularity for Markov systems. *Ergodic Theory Dynam. Systems* (2005) to appear
7. Bruin, H., Luzzatto, S., van Strien S.: Decay of correlations in one-dimensional dynamics. *Ann. Sci. Éc. Norm. Sup.* **36**, 621–646 (2003)
8. Bullard, E.C.: The stability of a homopolar dynamo. *Cambridge Phil. Soc* **51**, 744–760 (1955)
9. Diaz-Ordaz, K.: Decay of correlations for non-Hölder observables for expanding Lorenz-like one-dimensional maps. (2005) is: To appear in *Disc. And Cont. Dyn. Syst.*
10. Diaz-Ordaz, K., Holland, M., Luzzatto, S.: Statistical properties of one-dimensional maps with critical points and singularities. (2005) Preprint
11. Field, M., Melbourne, I., Török, A.: Stability of mixing and rapid mixing for hyperbolic flows. <http://www.maths.surrey.ac.uk/personal/st/I.Melbourne/papers/rapid.pdf>
12. Glendinning, P., Sparrow, C.: Prime and renormalisable kneading invariants and the dynamics of expanding Lorenz maps. *Phys. D.* **62**, 22–50 (1993) Homoclinic chaos (Brussels, 1991).
13. Gouëzel, S.: Regularity of coboundaries for non uniformly expanding Markov maps. <http://org/list/math.DS/0502458>, 2005 is: To appear in *Proc AMS*.
14. Guckenheimer, J., Williams, R.F.: Structural stability of Lorenz attractors. *Inst. Hautes Études Sci. Publ. Math.* **50**, 59–72 (1979)
15. Hide, R.: Geomagnetism, ‘vacillation’, atmospheric predictability and deterministic chaos. In: ‘Paths of Discovery’, scripta varia, Vatican city: Pontifical Acad. of Sci. Press, 2005
16. Katok, A. (in collaboration with K. Burns): Infinitesimal Lyapunov functions, invariant cone families and stochastic properties of smooth dynamical systems. *Ergodic Theory Dynam. Systems* **14**, 757–785 (1994)
17. Lorenz, E.D.: Deterministic nonperiodic flow. *J. Atmosph. Sci.* **20**, 130–141 (1963)
18. Morales, C.A., Pacifico, M.J., Pujals, E.R.: Singular hyperbolic systems. *Proc. Amer. Math. Soc.* **127**, 3393–3401 (1999)
19. Morales, C.A., Pacifico, M.J.: Mixing attractors for 3-flows. *Nonlinearity* **14**, 359–378 (2001)
20. Pesin, Ya.: Dynamical systems with generalized hyperbolic attractors: hyperbolic, ergodic and topological properties. *Ergod. Th. & Dynam. Sys.* **12**, 123–151 (1992)
21. Plante, Joseph F.: Anosov flows. *Amer. J. Math.* **94**, 729–754 (1972)
22. Ratner, M.: Bernoulli flow over maps of the interval. *Israel J. Math.* **31**, 298–314 (1978)
23. Rikitake, T.: Oscillations of a system of disk dynamos. *Proc. Cambridge Philos. Soc.* **54**, 89–105 (1958)
24. Rovella, A.: The dynamics of perturbations of the contracting Lorenz attractor. *Bol. Soc. Brasil. Mat. (N.S.)* **24**, 233–259 (1993)
25. Sataev, E.A.: Invariant measures for hyperbolic mappings with singularities. *Usp. Mat. Nauk.* **47**, 147–202, 240 (1992) (Russian)
26. Sparrow, C.: The Lorenz equations: bifurcations, chaos and strange attractors. *Applied Mathematical Sciences*, Vol. **41**, Berlin: Springer Verlag, 1982
27. Tucker, W.: The Lorenz attractor exists. *C. R. Acad. Sci. Paris Sér. I Math.* **328**, 1197–1202 (1999)
28. Tucker, W.: A rigorous ODE solver and Smale’s 14th problem. *Found. Comput. Math.* **2**, 53–117 (2002)
29. Viana, M.: Dynamical systems: moving into the next century. In: *Mathematics unlimited—2001 and beyond*, Berlin: Springer, 2001, pp. 1167–1178
30. Williams, R.F.: The structure of Lorenz attractors. *Inst. Hautes Études Sci. Publ. Math.* **50**, 321–342 (1979)
31. Young, L.-S.: Statistical properties of dynamical systems with some hyperbolicity. *Ann. of Math. (2)* **147**, 585–650 (1998)
32. Young, L.-S., Recurrence times and rates of mixing. *Israel J. Math.* **110**, 153–188 (1999)