




Quaternionic Heisenberg Group and Heterotic String Solutions with Non-Constant Dilaton in Dimensions 7 and 5

Marisa Fernández¹ , Stefan Ivanov^{2,3,4}, Luis Ugarte⁵, Dimiter Vassilev⁶

¹ Facultad de Ciencia y Tecnología, Departamento de Matemáticas, Universidad del País Vasco, Apartado 644, 48080 Bilbao, Spain. E-mail: marisa.fernandez@ehu.es

² Faculty of Mathematics and Informatics, University of Sofia “St. Kl. Ohridski”, Blvd. James Bourchier 5, 1164 Sofia, Bulgaria

³ Institute of Mathematics and Informatics, Bulgarian Academy of Sciences, Sofia, Bulgaria

⁴ Department of Mathematics, University of Pennsylvania, DRL 209 South 33rd Street, Philadelphia, PA 19104-6395, USA. E-mail: ivanovsp@fmi.uni-sofia.bg

⁵ Departamento de Matemáticas, I.U.M.A., Universidad de Zaragoza, Campus Plaza San Francisco, 50009 Zaragoza, Spain. E-mail: ugarte@unizar.es

⁶ Department of Mathematics and Statistics, University of New Mexico, Albuquerque, NM 87131-0001, USA. E-mail: vassilev@math.unm.edu

Received: 17 November 2014 / Accepted: 28 February 2015

Published online: 10 June 2015 – © Springer-Verlag Berlin Heidelberg 2015

Abstract: New smooth solutions of the Strominger system with non vanishing flux, non-trivial instanton and non-constant dilaton based on the quaternionic Heisenberg group are constructed. We show that through appropriate contractions the solutions found in the G_2 -heterotic case converge to the heterotic solutions on 6-dimensional inner non-Kähler spaces previously found by the authors and, moreover, to new heterotic solutions with non-constant dilatons in dimension 5. All found solutions satisfy the heterotic equations of motion up to first order of α' .

Contents

1. Introduction	200
2. The Supersymmetry Equations and the Geometric Model	203
2.1 Dimension 7	203
2.2 Dimension 5	204
2.2.1 Almost contact structure point of view.	204
2.2.2 The $SU(2)$ -structure point of view.	204
3. The Quaternionic Heisenberg Group	205
3.1 The first Pontrjagin form of the $(-)$ -connection	207
4. A Conformally Compact Solution with Negative α'	210
5. A Complete Solution with Positive α'	212
6. Solutions Through Contractions	214
6.1 Six dimensional solutions	214
6.2 Five dimensional solutions	215
References	217

1. Introduction

We investigate smooth solutions with non-trivial fluxes to the heterotic equations of motion, preserving at least one supersymmetry up to the first order of the string tension α' in dimensions seven and five. Using the quaternionic Heisenberg group, we propose an explicit construction leading to new smooth solutions with non-constant dilatons of the Strominger system comprised of the Killing spinor equations and the Green-Schwarz anomaly cancellation condition. The found solutions satisfy the heterotic equations of motion up to the first order of α' .

Another goal of the paper is to point out that through contractions of the Lie algebra of the quaternionic Heisenberg group, the geometric structures, the partial differential equations, and their solutions found in the G_2 -heterotic case converge to the heterotic solutions on the 6-dimensional inner non-Kähler spaces found in [26] and to the new 5-dimensional heterotic solutions with non-constant dilaton.

The bosonic fields of the ten-dimensional supergravity, which arises as low energy effective theory of the heterotic string, are the spacetime metric g , the NS three-form field strength (flux) H , the dilaton ϕ and the gauge connection A with curvature 2-form F^A . The bosonic geometry is of the form $\mathbb{R}^{1,9-d} \times M^d$, where the bosonic fields are non-trivial only on M^d , $d \leq 8$. We consider the two connections $\nabla^\pm = \nabla^g \pm \frac{1}{2}H$, where ∇^g is the Levi-Civita connection of the Riemannian metric g . Both connections preserve the metric, $\nabla^\pm g = 0$, and have totally skew-symmetric torsion $\pm H$, respectively. We denote by R^g , R^\pm the corresponding curvature.

We consider the heterotic supergravity theory with an α' expansion, where $1/2\pi\alpha'$ is the heterotic string tension. The bosonic part of the ten-dimensional supergravity action in the string frame is ([13,47], $R = R^-$)

$$S = \frac{1}{2k^2} \int d^{10}x \sqrt{-g} e^{-2\phi} \left[Scal^g + 4(\nabla^g \phi)^2 - \frac{1}{2}|H|^2 - \frac{\alpha'}{4} \left(Tr|F^A|^2 - Tr|R|^2 \right) \right]. \quad (1.1)$$

The string frame field equations (the equations of motion induced from the action (1.1)) of the heterotic string up to the first order of α' in sigma model perturbation theory are [45,47]

$$\begin{aligned} Ric_{ij}^g - \frac{1}{4} H_{imn} H_j^{mn} + 2\nabla_i^g \nabla_j^g \phi - \frac{\alpha'}{4} \left[(F^A)_{imab} (F^A)_j^{mab} - R_{imnq} R_j^{mnq} \right] &= 0, \\ \nabla_i^g (e^{-2\phi} H_{jk}^i) &= 0, \quad \nabla_i^+ (e^{-2\phi} (F^A)_j^i) = 0, \end{aligned} \quad (1.2)$$

in the notations of [37]. The field equation of the dilaton ϕ is implied from the first two equations above.

The Green-Schwarz anomaly cancellation mechanism requires that the three-form Bianchi identity receives an α' correction of the form

$$dH = \frac{\alpha'}{4} 8\pi^2 (p_1(M^d) - p_1(E)) = \frac{\alpha'}{4} \left(Tr(R \wedge R) - Tr(F^A \wedge F^A) \right), \quad (1.3)$$

where $p_1(M^d)$ and $p_1(E)$ are the first Pontrjagin forms of M^d with respect to a connection ∇ with curvature R and a vector bundle E with connection A , respectively.

A class of heterotic-string backgrounds for which the Bianchi identity of the three-form H receives a correction of type (1.3) are those with (2,0) world-volume supersymmetry. Such models were considered in [48]. The target-space geometry of (2,0)-supersymmetric sigma models has been extensively investigated in [44,48,68]. Recently,

there is revived interest in these models [17, 20, 33–35, 37, 52, 53] as string backgrounds and in connection with heterotic-string compactification with fluxes mainly in dimension six [2–11, 14, 16, 22, 30, 31, 36, 39–42, 56, 59, 60, 62–64].

Equations (1.3), (1.1) and (1.2) involve a subtlety due to the choice of the connection ∇ on TM^d , since anomalies can be canceled independently of the choice [46]. Different connections correspond to different regularization schemes in the two-dimensional worldsheet non-linear sigma model. Hence the background fields given for the particular choice of ∇ must be related to those for a different choice by a field redefinition [65]. Connections on M^d proposed to investigate the anomaly cancellation (1.3) are ∇^g [35, 68], ∇^+ [17, 19, 24], ∇^- [13, 16, 37, 46, 49, 50, 54, 55, 58, 61], and the Chern connection ∇^c when $d = 6$ [9, 30, 31, 56, 68].

A heterotic geometry preserves supersymmetry iff in ten dimensions there exists at least one Majorana-Weyl spinor ϵ such that the following Killing-spinor equations hold [13, 68]

$$\begin{aligned}\delta_\lambda &= \nabla_m \epsilon = \left(\nabla_m^g + \frac{1}{4} H_{mnp} \Gamma^{np} \right) \epsilon = \nabla^+ \epsilon = 0, \\ \delta_\Psi &= \left(\Gamma^m \partial_m \phi - \frac{1}{12} H_{mnp} \Gamma^{mnp} \right) \epsilon = (d\phi - \frac{1}{2} H) \cdot \epsilon = 0, \\ \delta_\xi &= F_{mn}^A \Gamma^{mn} \epsilon = F^A \cdot \epsilon = 0,\end{aligned}\tag{1.4}$$

where λ , Ψ , ξ are the gravitino, the dilatino and the gaugino fields, Γ_i generate the Clifford algebra $\{\Gamma_i, \Gamma_j\} = 2g_{ij}$ and \cdot means Clifford action of forms on spinors.

The system of Killing spinor Eq. (1.4) together with the anomaly cancellation condition (1.3) is known as the *Strominger system* [68]. The last equation in (1.4) is the instanton condition, which means that the curvature F^A is contained in a Lie algebra of a Lie group that is a stabilizer of a non-trivial spinor. In dimension 7 this group is G_2 . Denoting the G_2 three-form by Θ , the G_2 -instanton condition has the form

$$\sum_{k,l=1}^7 (F^A)^i_j (E_k, E_l) \Theta(E_k, E_l, E_m) = 0.\tag{1.5}$$

In the presence of a curvature term $Tr(R \wedge R)$, the solutions of the Strominger system (1.4), (1.3) obey the second and the third equations of motion (the second and the third equations in (1.2)) but do not always satisfy the Einstein equations of motion (see [23–25], where a sufficient quadratic condition on R is found). It was proved in [51] that, in dimensions 5, 6, 7 and 8, the solutions of the Strominger system ((1.4) and (1.3)) also solve the heterotic supersymmetric equations of motion (1.2) if and only if R is an instanton (see [58, 62] for higher dimensions and different proofs). In particular, in dimension 7, R is required to be a G_2 -instanton.

The physically relevant connection on the tangent bundle to be considered in (1.3), (1.1), (1.2) is the $(-)$ -connection [13, 46]. One reason is that the curvature R^- of the $(-)$ -connection is an instanton up to the first order of α' , which is a consequence of the first equation in (1.4), (1.3) and the well known identity

$$R^+(X, Y, Z, U) - R^-(Z, U, X, Y) = \frac{1}{2} dH(X, Y, Z, U).\tag{1.6}$$

Indeed, (1.3) together with (1.6) imply $R^+(X, Y, Z, U) - R^-(Z, U, X, Y) = O(\alpha')$ and the first equation in (1.4) yields that the holonomy group of ∇^+ is contained in G_2 , i.e., the curvature 2-form $R^+(X, Y) \subset \mathfrak{g}_2$ and therefore R^- satisfies the instanton condition

(1.5) up to the first order of α' . Hence, a solution to the Strominger system with first Pontrjagin form of the $(-)$ -connection always satisfies the heterotic equations of motion (1.2) up to the first order of α' (see e.g. [58] and references therein).

In dimension $d = 7$ heterotic/type I solutions with non-zero fluxes to the equations of motion preserving at least one supersymmetry were constructed in [43]. These solutions are noncompact, conformal to a flat space, and satisfy (1.4) and (1.3) with vanishing curvature term, $R = 0$. Noncompact solutions to (1.4) and (1.3) in dimension 7 are also presented in [50]. The first compact heterotic/type I solutions with non-zero fluxes and constant dilaton to the equations of motion preserving at least one supersymmetry (satisfying (1.4) and (1.3)) in dimension seven were constructed in [25].

In dimension $d = 5$, if the field strength vanishes, $H = 0$, then the 5-dimensional case reduces to dimension four, since any five dimensional Riemannian spin manifold admitting ∇^g -parallel spinor is reducible. Non-compact solutions on circle bundle over 4-dimensional base endowed with a hyper Kähler metric (when the 4-dimensional metric is Eguchi-Hanson, Taub-NUT or Atiyah-Hitchin) have appeared in [12, 32, 57, 64, 66], the compact cases are discussed in [35], where a cohomological obstruction is presented. The first compact heterotic/type I solutions with non-zero fluxes and constant dilaton to the equations of motion preserving at least one supersymmetry (satisfying (1.4) and (1.3)) in dimension five are constructed in [23].

In this paper, we construct smooth solutions with non vanishing flux and non-constant dilaton to the Strominger system using the first Pontrjagin form of the $(-)$ -connection on a 7-dimensional complete non-compact manifold equipped with conformally cocalibrated G_2 -structures of pure type coupled with carefully chosen instanton bundle. The source of the construction is the already constructed smooth compact solutions to the Strominger system with constant dilaton on nilmanifolds presented in [25] and the ideas outlined there to consider special three-torus bundles over either conformally \mathbb{T}^4 or $K3$ manifold.

Our first family of solutions are complete G_2 manifolds, which are \mathbb{T}^3 -bundles over conformally compact asymptotically hyperbolic metric on \mathbb{T}^4 with conformal boundary at infinity a flat torus \mathbb{T}^3 . Using the first Pontrjagin form of the $(-)$ -connection, together with the first Pontrjagin form of a carefully chosen instanton, we show that the anomaly cancellation condition is satisfied with a *negative* α' and a non-constant dilaton which is a real slice of an elliptic function of order two. See Remark 4.4 for the primary obstacle to finding a compact solution using these techniques.

In Sect. 5 we present another smooth non-compact complete solution to the Strominger system with non-vanishing torsion, non-trivial instanton and non-constant dilaton using the first Pontrjagin form of the $(-)$ -connection with positive string tension on certain \mathbb{T}^3 -bundle over \mathbb{R}^4 . The non-constant dilaton function here is determined by the fundamental solution of the Laplacian on \mathbb{R}^4 .

Conventions. The connection 1-forms ω_{ji} of a metric connection ∇ , $\nabla g = 0$, with respect to a local orthonormal basis $\{E_1, \dots, E_d\}$ are given by $\omega_{ji}(E_k) = g(\nabla_{E_k} E_j, E_i)$, since we write $\nabla_X E_j = \omega_j^s(X) E_s$.

The curvature 2-forms Ω_j^i of ∇ are given in terms of the connection 1-forms ω_j^i by $\Omega_j^i = d\omega_j^i + \omega_k^j \wedge \omega_j^k$, $\Omega_{ji} = d\omega_{ji} + \omega_{ki} \wedge \omega_{jk}$, $R_{ijk}^l = \Omega_k^l(E_i, E_j)$, $R_{ijkl} = R_{ijk}^s g_{ls}$.

The first Pontrjagin class is represented by the 4-form $8\pi^2 p_1(\nabla) = \sum_{1 \leq i < j \leq d} \Omega_j^i \wedge \Omega_j^i$.

2. The Supersymmetry Equations and the Geometric Model

Geometrically, the vanishing of the gravitino variation is equivalent to the existence of a non-trivial real spinor parallel with respect to the metric connection ∇^+ with totally skew-symmetric torsion $T = H$. The presence of ∇^+ -parallel spinor leads to restriction of the holonomy group $Hol(\nabla^+)$ of the torsion connection ∇^+ .

2.1. Dimension 7. In dimension seven $Hol(\nabla^+)$ has to be contained in the exceptional group G_2 [27, 28, 33, 35]. The precise conditions to have a solution to the gravitino Killing spinor equation in dimension 7 were found in [27]. Namely, there exists a non-trivial parallel spinor with respect to a G_2 -connection with torsion 3-form T if and only if there exists an integrable G_2 -structure Θ , i.e. $d*\Theta = \theta^7 \wedge *\Theta$, where $\theta^7 = -\frac{1}{3}*(d\Theta \wedge \Theta) = \frac{1}{3}*(d*\Theta \wedge \Theta)$ is the Lee form. In this case, the connection ∇^+ is unique and the torsion 3-form T is given by the formula [27]

$$H = T = \frac{1}{6}(d\Theta, *\Theta) \Theta - *d\Theta + *(\theta^7 \wedge \Theta).$$

The necessary conditions to have a solution to the system of dilatino and gravitino Killing spinor equations (the first two equations in (1.4)) in dimension seven were derived in [27, 28, 33], and the sufficiency was proved in [27, 28]. The general existence result [27, 28] states that there exists a non-trivial solution to both dilatino and gravitino Killing spinor equations (the first two equations in (1.4)) in dimension 7 if and only if there exists a globally conformal co-calibrated G_2 -structure (Θ, g) of pure type and with exact Lee form θ^7 , i.e. a G_2 -structure Θ satisfying the equations

$$d*\Theta = \theta^7 \wedge *\Theta, \quad d\Theta \wedge \Theta = 0, \quad \theta^7 = -2d\phi. \tag{2.1}$$

Consequently, the torsion 3-form (the flux H) is given by $H = T = -*d\Theta - 2*(d\phi \wedge \Theta)$ and the Riemannian scalar curvature satisfies $s^g = 8\|d\phi\|^2 - \frac{1}{12}\|T\|^2 - 6\delta d\phi$. The Eq. (2.1) hold exactly when the G_2 -structure $(\bar{\Theta} = e^{-\frac{3}{2}\phi}\Theta, \bar{g} = e^{-\phi}g)$ obeys the equations $d\bar{*}\bar{\Theta} = d\bar{\Theta} \wedge \bar{\Theta} = 0$, i.e., it is co-calibrated of pure type.

A geometric model which fits the above structures was proposed in [25] as a certain \mathbb{T}^3 -bundle over a Calabi-Yau surface. For this, let $\Gamma_i, 1 \leq i \leq 3$, be three closed anti-self-dual 2-forms on a Calabi-Yau surface M^4 , which represent integral cohomology classes. Denote by ω_1 and by $\omega_2 + \sqrt{-1}\omega_3$ the (closed) Kähler form and the holomorphic volume form on M^4 , respectively. Then, there is a compact 7-dimensional manifold $M^{1,1,1}$ which is the total space of a \mathbb{T}^3 -bundle over M^4 and has a G_2 -structure

$$\Theta = \omega_1 \wedge \eta_1 + \omega_2 \wedge \eta_2 - \omega_3 \wedge \eta_3 + \eta_1 \wedge \eta_2 \wedge \eta_3,$$

solving the first two Killing spinor equations in (1.4) with constant dilaton in dimension 7, where $\eta_i, 1 \leq i \leq 3$, is a 1-form on $M^{1,1,1}$ such that $d\eta_i = \Gamma_i, 1 \leq i \leq 3$.

For any smooth function f on M^4 , the G_2 -structure on $M^{1,1,1}$ given by

$$\Theta_f = e^{2f} \left[\omega_1 \wedge \eta_1 + \omega_2 \wedge \eta_2 - \omega_3 \wedge \eta_3 \right] + \eta_1 \wedge \eta_2 \wedge \eta_3$$

solves the first two Killing spinor equations in (1.4) with non-constant dilaton $\phi = -2f$. The corresponding Riemannian metric has the form

$$g_f = e^{2f} g_{cy} + \eta_1 \otimes \eta_1 + \eta_2 \otimes \eta_2 + \eta_3 \otimes \eta_3.$$

To achieve a smooth solution to the Strominger system we still have to determine an auxiliary vector bundle with an instanton and a linear connection on $M^{1,1,1}$ in order to satisfy the anomaly cancellation condition (1.3).

2.2. Dimension 5. The existence of ∇^+ -parallel spinor in dimension 5 determines an almost contact metric structure and, equivalently, a reduction of the structure group $SO(5)$ to $SU(2)$. The properties of the almost contact metric structure as well as solutions to the gravitino and dilatino Killing-spinor equations are investigated in [27,29] and presented in terms of reduction to $SU(2)$ in [23].

2.2.1. Almost contact structure point of view. We recall that an almost contact metric structure consists of an odd dimensional manifold M^{2k+1} equipped with a Riemannian metric g , vector field ξ of length one, its dual 1-form η as well as an endomorphism ψ of the tangent bundle such that

$$\psi(\xi) = 0, \quad \psi^2 = -id + \eta \otimes \xi, \quad g(\psi., \psi.) = g(., .) - \eta \otimes \eta.$$

The Reeb vector field ξ is determined by the equations $\eta(\xi) = 1, \quad \xi \lrcorner d\eta = 0$, where \lrcorner denotes the interior multiplication. The Nijenhuis tensor N , the fundamental form F and the Lee form θ of an almost contact metric structure are defined by

$$N = [\psi., \psi.] + \psi^2[., .] - \psi[\psi., .] - \psi[., \psi.] + d\eta \otimes \xi, \quad F(., .) = g(., \psi.),$$

$$\theta = \frac{1}{2}F \lrcorner dF.$$

It was shown in [29] that the gravitino and the dilatino equations admit a solution in dimension five if and only if the Nijenhuis tensor is totally skew-symmetric, the Reeb vector field ξ is a Killing vector field and the equalities $2d\phi = \theta$ and $*_{\mathbb{H}}d\eta = -d\eta$ hold, where $*_{\mathbb{H}}$ denotes the Hodge operator acting on the 4-dimensional orthogonal complement \mathbb{H} of the vector ξ , $\mathbb{H} = \text{Ker } \eta$.

2.2.2. The $SU(2)$ -structure point of view. The reduction of the structure group $SO(5)$ to $SU(2)$ is described in terms of forms by Conti and Salamon in [18] (see also [32]) as follows: an $SU(2)$ -structure on a 5-dimensional manifold M is $(\eta, F = \omega_1, \omega_2, \omega_3)$, where η is a 1-form dual to ξ via the metric and $\omega_s, s = 1, 2, 3$, are 2-forms on M satisfying $\omega_s \wedge \omega_t = \delta_{st} v, \quad v \wedge \eta \neq 0$ for some 4-form v , and $X \lrcorner \omega_1 = Y \lrcorner \omega_2 \Rightarrow \omega_3(X, Y) \geq 0$. The 2-forms $\omega_s, s = 1, 2, 3$, can be chosen to form a basis of the \mathbb{H} -self-dual 2-forms [18].

It was shown in [23] that the first two equations in (1.4) admit a solution in dimension five exactly when there exists a five dimensional manifold M endowed with an $SU(2)$ -structure $(\eta, F = \omega_1, \omega_2, \omega_3)$ satisfying the structure equations:

$$d\omega_s = 2df \wedge \omega_s, \quad *_{\mathbb{H}}d\eta = -d\eta, \quad df(\xi) = 0. \tag{2.2}$$

The flux H is given by [27,29]

$$H = T = \eta \wedge d\eta + 2d^\psi f \wedge F, \quad \text{where } d^\psi f(X) = -df(\psi X). \tag{2.3}$$

The dilaton ϕ is equal to $\phi = 2f$.

In other words, the gravitino and dilatino equations in dimension five are satisfied if and only if the manifold is “special” or \mathbb{H} -conformal to a quasi-Sasakian manifold with \mathbb{H} -anti-self-dual exterior derivative of the almost contact form and the metric has the form

$$g_f = e^{2f} g_{\mathbb{H}} + \eta \otimes \eta.$$

It was proposed in [25] to investigate S^1 bundles over a conformally hyper-Kähler manifold. This ansatz guaranties solution to the first two equations in (1.4). To achieve a smooth solution to the Strominger system we still have to determine a linear connection on the tangent bundle and an auxiliary vector bundle with an $SU(2)$ -instanton, i.e., a connection A with curvature 2-form F^A satisfying

$$(F^A)^i_j(\psi E_k, \psi E_l) = (F^A)^i_j(E_k, E_l), \quad \sum_{k=1}^5 (F^A)^i_j(E_k, \psi E_k) = 0 \quad (2.4)$$

so that the anomaly cancellation condition (1.3) is satisfied.

3. The Quaternionic Heisenberg Group

The seven dimensional quaternionic Heisenberg group $G(\mathbb{H})$ is the connected simply connected Lie group with a group multiplication $[\cdot, \cdot]$ determined by the Lie algebra $\mathfrak{g}(\mathbb{H})$ with structure equations

$$\begin{aligned} d\gamma^1 = d\gamma^2 = d\gamma^3 = d\gamma^4 = 0, \quad d\gamma^5 = \gamma^{12} - \gamma^{34}, \quad d\gamma^6 = \gamma^{13} + \gamma^{24}, \\ d\gamma^7 = \gamma^{14} - \gamma^{23}. \end{aligned} \quad (3.1)$$

In order to obtain results in dimensions less than seven through contractions of $\mathfrak{g}(\mathbb{H})$ it will be convenient to consider the orbit of $G(\mathbb{H})$ under the natural action of $GL(3, \mathbb{R})$ on the $span\{\gamma^5, \gamma^6, \gamma^7\}$. Accordingly let K_A be a seven-dimensional real Lie group with Lie bracket $[x, x']_A = A[A^{-1}x, A^{-1}x']$ for $A \in GL(3, \mathbb{R})$ defined by a basis of left-invariant 1-forms $\{e^1, \dots, e^7\}$ such that $e^i = \gamma^i$ for $1 \leq i \leq 4$ and $(e^5 \ e^6 \ e^7) = A(\gamma^5 \ \gamma^6 \ \gamma^7)^T$. Hence, the structure equations of the Lie algebra \mathfrak{K}_A of the group K_A are

$$de^1 = de^2 = de^3 = de^4 = 0, \quad de^{4+i} = \sum_{j=1}^3 a_{ij} \sigma_j, \quad i = 1, 2, 3, \quad (3.2)$$

where $\sigma_1 = e^{12} - e^{34}$, $\sigma_2 = e^{13} + e^{24}$, $\sigma_3 = e^{14} - e^{23}$ are the three anti-self-dual forms on \mathbb{R}^4 and

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}. \quad (3.3)$$

We will denote the norm of A by $|A|$, $|A|^2 = \sum_{i,j=1}^3 a_{ij}^2$.

Since \mathfrak{K}_A is isomorphic to $\mathfrak{g}(\mathbb{H})$, if K_A is connected and simply connected it is isomorphic to $G(\mathbb{H})$. Furthermore, any lattice Γ_A gives rise to a (compact) nilmanifold $M_A = K_A/\Gamma_A$, which is a \mathbb{T}^3 -bundle over a \mathbb{T}^4 whose connection 1-form is the anti-self-dual curvature on the four torus.

Following [25] we consider the G_2 -structure on the Lie group K_A defined by the 3-form

$$\Theta = \omega_1 \wedge e^7 + \omega_2 \wedge e^5 - \omega_3 \wedge e^6 + e^{567}, \quad (3.4)$$

where

$$\omega_1 = e^{12} + e^{34}, \quad \omega_2 = e^{13} - e^{24}, \quad \omega_3 = e^{14} + e^{23}$$

are the three closed self-dual 2-forms on \mathbb{R}^4 . The corresponding Hodge dual 4-form $*\Theta$ is given by

$$*\Theta = \omega_1 \wedge e^{56} + \omega_2 \wedge e^{67} + \omega_3 \wedge e^{57} + \frac{1}{2}\omega_1 \wedge \omega_1. \quad (3.5)$$

It is easy to check using (3.2) and the property $\sigma_i \wedge \omega_j = 0$ for $1 \leq i, j \leq 3$ that

$$d*\Theta = 0, \quad d\Theta \wedge \Theta = 0, \quad (3.6)$$

i.e. Θ is co-calibrated of pure type. According to [27,28] this G_2 -structure solves the gravitino and dilatino equations with constant dilaton.

Let f be a smooth function on \mathbb{R}^4 . Following [25] we consider the G_2 form given by

$$\bar{\Theta} = e^{2f} \left[\omega_1 \wedge e^7 + \omega_2 \wedge e^5 - \omega_3 \wedge e^6 \right] + e^{567}. \quad (3.7)$$

The corresponding metric \bar{g} on K_A has an orthonormal basis of 1-forms defined by

$$\bar{e}^1 = e^f e^1, \quad \bar{e}^2 = e^f e^2, \quad \bar{e}^3 = e^f e^3, \quad \bar{e}^4 = e^f e^4, \quad \bar{e}^5 = e^5, \quad \bar{e}^6 = e^6, \quad \bar{e}^7 = e^7. \quad (3.8)$$

With respect to \bar{g} , the self-dual forms $\bar{\omega}_i$ and anti-self-dual forms $\bar{\sigma}_i$ are given by

$$\bar{\omega}_i = e^{2f} \omega_i, \quad \bar{\sigma}_i = e^{2f} \sigma_i, \quad i = 1, 2, 3. \quad (3.9)$$

The corresponding Hodge dual 4-form $\bar{*}\bar{\Theta}$ is

$$\bar{*}\bar{\Theta} = e^{2f} \left[\omega_1 \wedge e^{56} + \omega_2 \wedge e^{67} + \omega_3 \wedge e^{57} + \frac{e^{2f}}{2} \omega_1 \wedge \omega_1 \right]. \quad (3.10)$$

It was shown in [25, Theorem 6.1] using (3.6) that

$$d\bar{*}\bar{\Theta} = 2df \wedge \bar{*}\bar{\Theta}, \quad d\bar{\Theta} \wedge \bar{\Theta} = 0. \quad (3.11)$$

Then the Lie form $\bar{\theta}$ is given by

$$\bar{\theta} = 2df \quad (3.12)$$

and the G_2 -structure $\bar{\Theta}$ solves the gravitino and dilatino equations with non-constant dilaton $\phi = -2f$ [27,28].

According to [27,28], the torsion of the (+)-connection ∇^+ is the 3-form

$$T = -*d\Theta + *(\theta \wedge \Theta). \quad (3.13)$$

We calculate from (3.2) and (3.7) that

$$d\bar{\Theta} = 2df \wedge \bar{\Theta} - 2df \wedge e^{567} + de^{567}. \quad (3.14)$$

A substitution of (3.14) in (3.13) together with (3.12) give

$$\begin{aligned} \bar{T} &= \bar{*}(2df \wedge e^{567} - de^{567}) = e^{-f} \left[-2f_1 \bar{e}^{234} + 2f_2 \bar{e}^{134} - 2f_3 \bar{e}^{124} + 2f_4 \bar{e}^{123} \right] \\ &\quad + e^{-2f} \left[(a_{11} \bar{\sigma}_1 + a_{12} \bar{\sigma}_2 + a_{13} \bar{\sigma}_3) \wedge \bar{e}^5 + (a_{21} \bar{\sigma}_1 + a_{22} \bar{\sigma}_2 + a_{23} \bar{\sigma}_3) \wedge \bar{e}^6 \right. \\ &\quad \left. + (a_{31} \bar{\sigma}_1 + a_{32} \bar{\sigma}_2 + a_{33} \bar{\sigma}_3) \wedge \bar{e}^7 \right], \end{aligned} \quad (3.15)$$

where $f_i = \frac{\partial f}{\partial x_i}$, $1 \leq i \leq 4$, $\bar{\sigma}_1 = \bar{e}^{12} - \bar{e}^{34}$, $\bar{\sigma}_2 = \bar{e}^{13} + \bar{e}^{24}$ and $\bar{\sigma}_3 = \bar{e}^{14} - \bar{e}^{23}$. Letting $f_{ij} = \frac{\partial^2 f}{\partial x_j \partial x_i}$, $1 \leq i, j \leq 4$, a short calculation gives

$$d\bar{T} = -e^{-4f} \left[\Delta e^{2f} + 2|A|^2 \right] \bar{e}^{1234} = - \left[\Delta e^{2f} + 2|A|^2 \right] e^{1234}, \quad (3.16)$$

where $\Delta e^{2f} = (e^{2f})_{11} + (e^{2f})_{22} + (e^{2f})_{33} + (e^{2f})_{44}$ is the standard Laplacian on \mathbb{R}^4 .

3.1. The first Pontrjagin form of the (-)-connection. From Koszul's formula, we have that the Levi-Civita connection 1-forms $(\omega^{\bar{g}})^{\bar{i}}_{\bar{j}}$ of the metric \bar{g} are given by

$$\begin{aligned} (\omega^{\bar{g}})^{\bar{i}}_{\bar{j}}(\bar{e}_k) &= -\frac{1}{2} \left(\bar{g}(\bar{e}_i, [\bar{e}_j, \bar{e}_k]) - \bar{g}(\bar{e}_k, [\bar{e}_i, \bar{e}_j]) + \bar{g}(\bar{e}_j, [\bar{e}_k, \bar{e}_i]) \right) \\ &= \frac{1}{2} \left(d\bar{e}^i(\bar{e}_j, \bar{e}_k) - d\bar{e}^k(\bar{e}_i, \bar{e}_j) + d\bar{e}^j(\bar{e}_k, \bar{e}_i) \right) \end{aligned} \quad (3.17)$$

taking into account $\bar{g}(\bar{e}_i, [\bar{e}_j, \bar{e}_k]) = -d\bar{e}^i(\bar{e}_j, \bar{e}_k)$. With the help of (3.17) we compute the expressions for the connection 1-forms $(\omega^-)^{\bar{i}}_{\bar{j}}$ of the connection ∇^- ,

$$(\omega^-)^{\bar{i}}_{\bar{j}} = (\omega^{\bar{g}})^{\bar{i}}_{\bar{j}} - \frac{1}{2} (\bar{T})^{\bar{i}}_{\bar{j}}, \quad \text{where} \quad (\bar{T})^{\bar{i}}_{\bar{j}}(\bar{e}_k) = \bar{T}(\bar{e}_i, \bar{e}_j, \bar{e}_k). \quad (3.18)$$

Now, (3.18), (3.17) and (3.15) show that the possibly non-zero connection 1-forms $(\omega^-)^{\bar{i}}_{\bar{j}}$ are given in terms of the basis $\{\bar{e}^1, \dots, \bar{e}^7\}$ by the identities

$$\begin{aligned} (\omega^-)^{\bar{1}}_{\bar{2}} &= (\omega^-)^{\bar{3}}_{\bar{4}} = e^{-f} \left(f_2 \bar{e}^1 - f_1 \bar{e}^2 + f_4 \bar{e}^3 - f_3 \bar{e}^4 \right), \\ (\omega^-)^{\bar{1}}_{\bar{3}} &= -(\omega^-)^{\bar{2}}_{\bar{4}} = e^{-f} \left(f_3 \bar{e}^1 - f_4 \bar{e}^2 - f_1 \bar{e}^3 + f_2 \bar{e}^4 \right), \\ (\omega^-)^{\bar{1}}_{\bar{4}} &= (\omega^-)^{\bar{2}}_{\bar{3}} = e^{-f} \left(f_4 \bar{e}^1 + f_3 \bar{e}^2 - f_2 \bar{e}^3 - f_1 \bar{e}^4 \right), \\ (\omega^-)^{\bar{1}}_{\bar{5}} &= e^{-2f} \left(-a_{11} \bar{e}^2 - a_{12} \bar{e}^3 - a_{13} \bar{e}^4 \right), \quad (\omega^-)^{\bar{1}}_{\bar{6}} = e^{-2f} \left(-a_{21} \bar{e}^2 - a_{22} \bar{e}^3 - a_{23} \bar{e}^4 \right), \\ (\omega^-)^{\bar{1}}_{\bar{7}} &= e^{-2f} \left(-a_{31} \bar{e}^2 - a_{32} \bar{e}^3 - a_{33} \bar{e}^4 \right), \quad (\omega^-)^{\bar{2}}_{\bar{5}} = e^{-2f} \left(a_{11} \bar{e}^1 + a_{13} \bar{e}^3 - a_{12} \bar{e}^4 \right), \\ (\omega^-)^{\bar{2}}_{\bar{6}} &= e^{-2f} \left(a_{21} \bar{e}^1 + a_{23} \bar{e}^3 - a_{22} \bar{e}^4 \right), \quad (\omega^-)^{\bar{2}}_{\bar{7}} = e^{-2f} \left(a_{31} \bar{e}^1 + a_{33} \bar{e}^3 - a_{32} \bar{e}^4 \right), \\ (\omega^-)^{\bar{3}}_{\bar{5}} &= e^{-2f} \left(a_{12} \bar{e}^1 - a_{13} \bar{e}^2 + a_{11} \bar{e}^4 \right), \quad (\omega^-)^{\bar{3}}_{\bar{6}} = e^{-2f} \left(a_{22} \bar{e}^1 - a_{23} \bar{e}^2 + a_{21} \bar{e}^4 \right), \\ (\omega^-)^{\bar{3}}_{\bar{7}} &= e^{-2f} \left(a_{32} \bar{e}^1 - a_{33} \bar{e}^2 + a_{31} \bar{e}^4 \right), \quad (\omega^-)^{\bar{4}}_{\bar{5}} = e^{-2f} \left(a_{13} \bar{e}^1 + a_{12} \bar{e}^2 - a_{11} \bar{e}^3 \right), \\ (\omega^-)^{\bar{4}}_{\bar{6}} &= e^{-2f} \left(a_{23} \bar{e}^1 + a_{22} \bar{e}^2 - a_{21} \bar{e}^3 \right), \quad (\omega^-)^{\bar{4}}_{\bar{7}} = e^{-2f} \left(a_{33} \bar{e}^1 + a_{32} \bar{e}^2 - a_{31} \bar{e}^3 \right). \end{aligned} \quad (3.19)$$

A long straightforward calculation using (3.19) gives in terms of the basis $\{\bar{e}^1, \dots, \bar{e}^7\}$ the following formulas for the curvature 2-forms $(\Omega^-)^{\bar{i}}_{\bar{j}}$ of the connection ∇^- :

$$\begin{aligned}
(\Omega^-)_{\frac{1}{2}}^{\bar{1}} &= -e^{-2f} \left[f_{11} + f_{22} + 2f_3^2 + 2f_4^2 + (a_{11}^2 + a_{21}^2 + a_{31}^2)e^{-2f} \right] \bar{e}^{12} \\
&+ e^{-2f} \left[f_{14} - f_{23} - 2f_1 f_4 + 2f_2 f_3 - (a_{11}a_{12} + a_{21}a_{22} + a_{31}a_{32})e^{-2f} \right] \bar{\sigma}_2 \\
&- e^{-2f} \left[f_{13} + f_{24} - 2f_1 f_3 - 2f_2 f_4 + (a_{11}a_{13} + a_{21}a_{23} + a_{31}a_{33})e^{-2f} \right] \bar{\sigma}_3 \\
&- e^{-2f} \left[f_{33} + f_{44} + 2f_1^2 + 2f_2^2 + (a_{12}^2 + a_{22}^2 + a_{32}^2 + a_{13}^2 + a_{23}^2 + a_{33}^2)e^{-2f} \right] \bar{e}^{34},
\end{aligned}$$

$$\begin{aligned}
(\Omega^-)_{\frac{1}{3}}^{\bar{1}} &= -e^{-2f} \left[f_{14} + f_{23} - 2f_1 f_4 - 2f_2 f_3 + (a_{11}a_{12} + a_{21}a_{22} + a_{31}a_{32})e^{-2f} \right] \bar{\sigma}_1 \\
&- e^{-2f} \left[f_{11} + f_{33} + 2f_2^2 + 2f_4^2 + (a_{12}^2 + a_{22}^2 + a_{32}^2)e^{-2f} \right] \bar{e}^{13} \\
&+ e^{-2f} \left[f_{12} - f_{34} - 2f_1 f_2 + 2f_3 f_4 + (a_{12}a_{13} + a_{22}a_{23} + a_{32}a_{33})e^{-2f} \right] \bar{\sigma}_3 \\
&+ e^{-2f} \left[f_{22} + f_{44} + 2f_1^2 + 2f_3^2 + (a_{11}^2 + a_{21}^2 + a_{31}^2 + a_{13}^2 + a_{23}^2 + a_{33}^2)e^{-2f} \right] \bar{e}^{24},
\end{aligned}$$

$$\begin{aligned}
(\Omega^-)_{\frac{1}{4}}^{\bar{1}} &= e^{-2f} \left[f_{13} - f_{24} - 2f_1 f_3 + 2f_2 f_4 - (a_{11}a_{13} + a_{21}a_{23} + a_{31}a_{33})e^{-2f} \right] \bar{\sigma}_1 \\
&- e^{-2f} \left[f_{12} + f_{34} - 2f_1 f_2 - 2f_3 f_4 + (a_{12}a_{13} + a_{22}a_{23} + a_{32}a_{33})e^{-2f} \right] \bar{\sigma}_2 \\
&- e^{-2f} \left[f_{11} + f_{44} + 2f_2^2 + 2f_3^2 + (a_{13}^2 + a_{23}^2 + a_{33}^2)e^{-2f} \right] \bar{e}^{14} \\
&- e^{-2f} \left[f_{22} + f_{33} + 2f_1^2 + 2f_4^2 + (a_{11}^2 + a_{21}^2 + a_{31}^2 + a_{12}^2 + a_{22}^2 + a_{32}^2)e^{-2f} \right] \bar{e}^{23},
\end{aligned}$$

$$\begin{aligned}
(\Omega^-)_{\frac{1}{5}}^{\bar{1}} &= 2e^{-3f} [(a_{11}f_1 - a_{13}f_3 + a_{12}f_4)\bar{\sigma}_1 + (a_{12}f_1 + a_{13}f_2 - a_{11}f_4)\bar{\sigma}_2 \\
&+ (a_{13}f_1 - a_{12}f_2 + a_{11}f_3)\bar{\sigma}_3],
\end{aligned}$$

$$\begin{aligned}
(\Omega^-)_{\frac{1}{6}}^{\bar{1}} &= 2e^{-3f} [(a_{21}f_1 - a_{23}f_3 + a_{22}f_4)\bar{\sigma}_1 + (a_{22}f_1 + a_{23}f_2 - a_{21}f_4)\bar{\sigma}_2 \\
&+ (a_{23}f_1 - a_{22}f_2 + a_{21}f_3)\bar{\sigma}_3],
\end{aligned}$$

$$\begin{aligned}
(\Omega^-)_{\frac{1}{7}}^{\bar{1}} &= 2e^{-3f} [(a_{31}f_1 - a_{33}f_3 + a_{32}f_4)\bar{\sigma}_1 + (a_{32}f_1 + a_{33}f_2 - a_{31}f_4)\bar{\sigma}_2 \\
&+ (a_{33}f_1 - a_{32}f_2 + a_{31}f_3)\bar{\sigma}_3],
\end{aligned}$$

$$\begin{aligned}
(\Omega^-)_{\frac{2}{3}}^{\bar{2}} &= e^{-2f} \left[f_{13} - f_{24} - 2f_1 f_3 + 2f_2 f_4 + (a_{11}a_{13} + a_{21}a_{23} + a_{31}a_{33})e^{-2f} \right] \bar{\sigma}_1 \\
&- e^{-2f} \left[f_{12} + f_{34} - 2f_1 f_2 - 2f_3 f_4 - (a_{12}a_{13} + a_{22}a_{23} + a_{32}a_{33})e^{-2f} \right] \bar{\sigma}_2 \\
&- e^{-2f} \left[f_{11} + f_{44} + 2f_2^2 + 2f_3^2 + (a_{11}^2 + a_{21}^2 + a_{31}^2 + a_{12}^2 + a_{22}^2 + a_{32}^2)e^{-2f} \right] \bar{e}^{14} \\
&- e^{-2f} \left[f_{22} + f_{33} + 2f_1^2 + 2f_4^2 + (a_{13}^2 + a_{23}^2 + a_{33}^2)e^{-2f} \right] \bar{e}^{23},
\end{aligned}$$

$$\begin{aligned}
(\Omega^-)_{\frac{2}{4}}^{\bar{2}} &= e^{-2f} \left[f_{14} + f_{23} - 2f_1 f_4 - 2f_2 f_3 - (a_{11}a_{12} + a_{21}a_{22} + a_{31}a_{32})e^{-2f} \right] \bar{\sigma}_1 \\
&+ e^{-2f} \left[f_{11} + f_{33} + 2f_2^2 + 2f_4^2 + (a_{11}^2 + a_{21}^2 + a_{31}^2 + a_{13}^2 + a_{23}^2 + a_{33}^2)e^{-2f} \right] \bar{e}^{13} \\
&- e^{-2f} \left[f_{12} - f_{34} - 2f_1 f_2 + 2f_3 f_4 + (a_{12}a_{13} + a_{22}a_{23} + a_{32}a_{33})e^{-2f} \right] \bar{\sigma}_3 \\
&- e^{-2f} \left[f_{22} + f_{44} + 2f_1^2 + 2f_3^2 + (a_{12}^2 + a_{22}^2 + a_{32}^2)e^{-2f} \right] \bar{e}^{24},
\end{aligned}$$

$$\begin{aligned}
 (\Omega^-)_{\bar{5}}^{\bar{2}} &= 2e^{-3f} [(a_{11}f_2 - a_{12}f_3 - a_{13}f_4)\bar{\sigma}_1 - (a_{13}f_1 - a_{12}f_2 - a_{11}f_3)\bar{\sigma}_2 \\
 &\quad + (a_{12}f_1 + a_{13}f_2 + a_{11}f_4)\bar{\sigma}_3], \\
 (\Omega^-)_{\bar{6}}^{\bar{2}} &= 2e^{-3f} [(a_{21}f_2 - a_{22}f_3 - a_{23}f_4)\bar{\sigma}_1 - (a_{23}f_1 - a_{22}f_2 - a_{21}f_3)\bar{\sigma}_2 \\
 &\quad + (a_{22}f_1 + a_{23}f_2 + a_{21}f_4)\bar{\sigma}_3], \\
 (\Omega^-)_{\bar{7}}^{\bar{2}} &= 2e^{-3f} [(a_{31}f_2 - a_{32}f_3 - a_{33}f_4)\bar{\sigma}_1 - (a_{33}f_1 - a_{32}f_2 - a_{31}f_3)\bar{\sigma}_2 \\
 &\quad + (a_{32}f_1 + a_{33}f_2 + a_{31}f_4)\bar{\sigma}_3], \\
 (\Omega^-)_{\bar{4}}^{\bar{3}} &= -e^{-2f} [f_{11} + f_{22} + 2f_3^2 + 2f_4^2 + (a_{12}^2 + a_{22}^2 + a_{32}^2 + a_{13}^2 + a_{23}^2 + a_{33}^2)e^{-2f}]e^{12} \\
 &\quad + e^{-2f} [f_{14} - f_{23} - 2f_1f_4 + 2f_2f_3 + (a_{11}a_{12} + a_{21}a_{22} + a_{31}a_{32})e^{-2f}]\bar{\sigma}_2 \\
 &\quad - e^{-2f} [f_{13} + f_{24} - 2f_1f_3 - 2f_2f_4 - (a_{11}a_{13} + a_{21}a_{23} + a_{31}a_{33})e^{-2f}]\bar{\sigma}_3 \\
 &\quad - e^{-2f} [f_{33} + f_{44} + 2f_1^2 + 2f_2^2 + (a_{11}^2 + a_{21}^2 + a_{31}^2)e^{-2f}]e^{34}, \\
 (\Omega^-)_{\bar{5}}^{\bar{3}} &= 2e^{-3f} [(a_{13}f_1 + a_{12}f_2 + a_{11}f_3)\bar{\sigma}_1 - (a_{11}f_2 - a_{12}f_3 + a_{13}f_4)\bar{\sigma}_2 \\
 &\quad - (a_{11}f_1 - a_{13}f_3 - a_{12}f_4)\bar{\sigma}_3], \\
 (\Omega^-)_{\bar{6}}^{\bar{3}} &= 2e^{-3f} [(a_{23}f_1 + a_{22}f_2 + a_{21}f_3)\bar{\sigma}_1 - (a_{21}f_2 - a_{22}f_3 + a_{23}f_4)\bar{\sigma}_2 \\
 &\quad - (a_{21}f_1 - a_{23}f_3 - a_{22}f_4)\bar{\sigma}_3], \\
 (\Omega^-)_{\bar{7}}^{\bar{3}} &= 2e^{-3f} [(a_{33}f_1 + a_{32}f_2 + a_{31}f_3)\bar{\sigma}_1 - (a_{31}f_2 - a_{32}f_3 + a_{33}f_4)\bar{\sigma}_2 \\
 &\quad - (a_{31}f_1 - a_{33}f_3 - a_{32}f_4)\bar{\sigma}_3], \\
 (\Omega^-)_{\bar{5}}^{\bar{4}} &= 2e^{-3f} [-(a_{12}f_1 - a_{13}f_2 - a_{11}f_4)\bar{\sigma}_1 + (a_{11}f_1 + a_{13}f_3 + a_{12}f_4)\bar{\sigma}_2 \\
 &\quad - (a_{11}f_2 + a_{12}f_3 - a_{13}f_4)\bar{\sigma}_3], \\
 (\Omega^-)_{\bar{6}}^{\bar{4}} &= 2e^{-3f} [-(a_{22}f_1 - a_{23}f_2 - a_{21}f_4)\bar{\sigma}_1 + (a_{21}f_1 + a_{23}f_3 + a_{22}f_4)\bar{\sigma}_2 \\
 &\quad - (a_{21}f_2 + a_{22}f_3 - a_{23}f_4)\bar{\sigma}_3], \\
 (\Omega^-)_{\bar{7}}^{\bar{4}} &= 2e^{-3f} [-(a_{32}f_1 - a_{33}f_2 - a_{31}f_4)\bar{\sigma}_1 + (a_{31}f_1 + a_{33}f_3 + a_{32}f_4)\bar{\sigma}_2 \\
 &\quad - (a_{31}f_2 + a_{32}f_3 - a_{33}f_4)\bar{\sigma}_3], \\
 (\Omega^-)_{\bar{6}}^{\bar{5}} &= 2e^{-4f} [(a_{12}a_{23} - a_{13}a_{22})\bar{\sigma}_1 - (a_{11}a_{23} - a_{13}a_{21})\bar{\sigma}_2 + (a_{11}a_{22} - a_{12}a_{21})\bar{\sigma}_3], \\
 (\Omega^-)_{\bar{7}}^{\bar{5}} &= 2e^{-4f} [(a_{12}a_{33} - a_{13}a_{32})\bar{\sigma}_1 - (a_{11}a_{33} - a_{13}a_{31})\bar{\sigma}_2 + (a_{11}a_{32} - a_{12}a_{31})\bar{\sigma}_3], \\
 (\Omega^-)_{\bar{7}}^{\bar{6}} &= 2e^{-4f} [(a_{22}a_{33} - a_{23}a_{32})\bar{\sigma}_1 - (a_{21}a_{33} - a_{23}a_{31})\bar{\sigma}_2 + (a_{21}a_{32} - a_{22}a_{31})\bar{\sigma}_3].
 \end{aligned}$$

A long calculation based on the formulas for the curvature 2-form $(\Omega^-)_{\bar{j}}^{\bar{i}}$ of ∇^- gives.

Proposition 3.1. *The first Pontrjagin form of ∇^- is a scalar multiple of e^{1234} given by*

$$\pi^2 p_1(\nabla^-) = \left[\mathcal{F}_2[f] + \Delta_4 f - \frac{3}{8}|A|^2 \Delta e^{-2f} \right] e^{1234}, \tag{3.20}$$

where $\mathcal{F}_2[f]$ is the 2-Hessian of f , i.e., the sum of all principle 2×2 -minors of the Hessian, and $\Delta_4 f = \text{div}(|\nabla f|^2 \nabla f)$ is the 4-Laplacian of f .

The above Proposition shows, in particular, that even though the curvature 2-forms of ∇^- are quadratic in the gradient of the dilaton, the Pontrjagin form of ∇^- is also quadratic in these terms. Furthermore, if f depends on two of the variables then $\mathcal{F}_2[f] = \det(\text{Hess}f)$ while if f is a function of one variable $\mathcal{F}_2[f]$ vanishes.

4. A Conformally Compact Solution with Negative α'

In this section we give our first main result. Recall that K_A is the connected simply connected Lie group with Lie algebra \mathfrak{K}_A determined by (3.2). Due to the results recalled in Sect. 2.1 the remaining part is to solve the anomaly cancellation condition. This will be achieved for the G_2 structure (3.7) with the torsion term (3.16), the Pontrjagin form (3.20) of the ∇^- connection, and the G_2 -instanton defined below.

Proposition 4.1. *Let D_Λ , $\Lambda = (\lambda_{ij}) \in \mathfrak{gl}_3(\mathbb{R})$, be the linear connection on the Lie group K_A whose possibly non-zero 1-forms are given as follows*

$$\begin{aligned} (\omega^{D_\Lambda})_{\bar{2}}^{\bar{1}} &= -(\omega^{D_\Lambda})_{\bar{1}}^{\bar{2}} = -(\omega^{D_\Lambda})_{\bar{4}}^{\bar{3}} = (\omega^{D_\Lambda})_{\bar{3}}^{\bar{4}} = \lambda_{11} \bar{e}^5 + \lambda_{12} \bar{e}^6 + \lambda_{13} \bar{e}^7, \\ (\omega^{D_\Lambda})_{\bar{3}}^{\bar{1}} &= -(\omega^{D_\Lambda})_{\bar{1}}^{\bar{3}} = (\omega^{D_\Lambda})_{\bar{4}}^{\bar{2}} = -(\omega^{D_\Lambda})_{\bar{2}}^{\bar{4}} = \lambda_{21} \bar{e}^5 + \lambda_{22} \bar{e}^6 + \lambda_{23} \bar{e}^7, \\ (\omega^{D_\Lambda})_{\bar{4}}^{\bar{1}} &= -(\omega^{D_\Lambda})_{\bar{1}}^{\bar{4}} = -(\omega^{D_\Lambda})_{\bar{3}}^{\bar{2}} = (\omega^{D_\Lambda})_{\bar{2}}^{\bar{3}} = \lambda_{31} \bar{e}^5 + \lambda_{32} \bar{e}^6 + \lambda_{33} \bar{e}^7. \end{aligned}$$

Then, D_Λ is a G_2 -instanton with respect to the G_2 structure defined by (3.7) which preserves the metric if and only if $\text{rank}(\Lambda) \leq 1$.

Proof. Let us use the notation $\Lambda_{ijkl} = \lambda_{ik}\lambda_{jl} - \lambda_{jk}\lambda_{il} = \det \begin{pmatrix} \lambda_{ik} & \lambda_{il} \\ \lambda_{jk} & \lambda_{jl} \end{pmatrix}$ for the 2×2 minors of Λ . A direct calculation using (3.2) shows that the possibly non-zero curvature forms $(\Omega^{D_\Lambda})_{\bar{j}}^{\bar{i}}$ of the connection D_Λ are:

$$\begin{aligned} (\Omega^{D_\Lambda})_{\bar{2}}^{\bar{1}} &= -(\Omega^{D_\Lambda})_{\bar{1}}^{\bar{2}} = -(\Omega^{D_\Lambda})_{\bar{4}}^{\bar{3}} = (\Omega^{D_\Lambda})_{\bar{3}}^{\bar{4}} = e^{-2f} (a_{11}\lambda_{11} + a_{21}\lambda_{12} + a_{31}\lambda_{13}) \bar{\sigma}_1 \\ &\quad + e^{-2f} (a_{12}\lambda_{11} + a_{22}\lambda_{12} + a_{32}\lambda_{13}) \bar{\sigma}_2 + e^{-2f} (a_{13}\lambda_{11} + a_{23}\lambda_{12} + a_{33}\lambda_{13}) \bar{\sigma}_3 \\ &\quad + 2\Lambda_{2312} \bar{e}^{56} + 2\Lambda_{2313} \bar{e}^{57} + 2\Lambda_{2323} \bar{e}^{67}, \\ (\Omega^{D_\Lambda})_{\bar{3}}^{\bar{1}} &= -(\Omega^{D_\Lambda})_{\bar{1}}^{\bar{3}} = (\Omega^{D_\Lambda})_{\bar{4}}^{\bar{2}} = -(\Omega^{D_\Lambda})_{\bar{2}}^{\bar{4}} = e^{-2f} (a_{11}\lambda_{21} + a_{21}\lambda_{22} + a_{31}\lambda_{23}) \bar{\sigma}_1 \\ &\quad + e^{-2f} (a_{12}\lambda_{21} + a_{22}\lambda_{22} + a_{32}\lambda_{23}) \bar{\sigma}_2 + e^{-2f} (a_{13}\lambda_{21} + a_{23}\lambda_{22} + a_{33}\lambda_{23}) \bar{\sigma}_3 \\ &\quad - 2\Lambda_{1312} \bar{e}^{56} - 2\Lambda_{1313} \bar{e}^{57} - 2\Lambda_{1323} \bar{e}^{67}, \\ (\Omega^{D_\Lambda})_{\bar{4}}^{\bar{1}} &= -(\Omega^{D_\Lambda})_{\bar{1}}^{\bar{4}} = -(\Omega^{D_\Lambda})_{\bar{3}}^{\bar{2}} = (\Omega^{D_\Lambda})_{\bar{2}}^{\bar{3}} = e^{-2f} (a_{11}\lambda_{31} + a_{21}\lambda_{32} + a_{31}\lambda_{33}) \bar{\sigma}_1 \\ &\quad + e^{-2f} (a_{12}\lambda_{31} + a_{22}\lambda_{32} + a_{32}\lambda_{33}) \bar{\sigma}_2 + e^{-2f} (a_{13}\lambda_{31} + a_{23}\lambda_{32} + a_{33}\lambda_{33}) \bar{\sigma}_3 \\ &\quad + 2\Lambda_{1212} \bar{e}^{56} + 2\Lambda_{1213} \bar{e}^{57} + 2\Lambda_{1223} \bar{e}^{67}. \end{aligned}$$

Now, it is straightforward to see that D_Λ satisfies (1.5) if and only if all the 2×2 minors Λ_{ijkl} of the matrix Λ vanish. Therefore, D_Λ is a G_2 -instanton if and only if $\text{rank}(\Lambda) \leq 1$. \square

Corollary 4.2. *For a matrix Λ of rank one, $\Lambda = (\lambda_{ij}) \in \mathfrak{gl}_3(\mathbb{R})$, let D_Λ be the G_2 -instanton defined in Proposition 4.1. Then, the first Pontrjagin form $p_1(D_\Lambda)$ of the G_2 -instanton D_Λ is given by*

$$8\pi^2 p_1(D_\Lambda) = -4\lambda^2 e^{1234}, \quad (4.1)$$

where $\lambda = |\Lambda A|$ is the norm of the product matrix ΛA .

Proof. Since the 2×2 minors Λ_{ijkl} are all zero, the formulas for the curvature forms $(\Omega^{D_\Lambda})_{\bar{j}}^{\bar{i}}$ given in the proof of Proposition 4.1 imply the claimed identity. \square

We turn to the proof of our first main result whose formulation uses (3.2), (3.7), (3.19) and Proposition 4.1.

Theorem 4.3. *The conformally compact manifold $M^7 = (\Gamma \backslash K_A, \bar{\Theta}, \nabla^-, D_\Lambda, f)$ is a G_2 -manifold which solves the Strominger system with non-constant dilaton f , non-trivial flux $H = \bar{T}$, non-flat instanton D_Λ using the first Pontrjagin form of ∇^- and negative α' . The dilaton f depends on one variable and is determined as a real slice of the Weierstrass' elliptic function.*

The conformally compact manifold $M^7 = (\Gamma \backslash K_A, \bar{\Theta}, \nabla^-, D_\Lambda, f)$ satisfies the heterotic equations of motion (1.2) up to first order of α' .

Proof. By the construction in Sect. 2.1 we are left with solving the anomaly cancellation condition $d\bar{T} = \frac{\alpha'}{4} 8\pi^2 (p_1(\nabla^-) - p_1(D_\Lambda))$, which in our case taking into account (3.16), (3.20) and (4.1) becomes the single non-linear equation

$$\Delta e^{2f} + 2|A|^2 + \frac{\alpha'}{4} [8\mathcal{F}_2[f] + 8\Delta_4 f - 3|A|^2 \Delta e^{-2f} + 4\lambda^2] = 0. \tag{4.2}$$

Remark 4.4. We note explicitly that the existence of a periodic solution of Eq. (4.2) on \mathbb{R}^4 would give a compact solution of Strominger system with non-constant dilaton.

Up to relabeling the constants, this is the same equation as the one obtained through the anomaly cancellation that appeared in [26, Sect. 4.2]. Accordingly, we assume that the function f depends on one variable, $f = f(x^1)$, and for a negative α' we choose $2|A|^2 + \alpha'\lambda^2 = 0$, i.e., we let $\alpha' = -\alpha^2$ so that $2|A|^2 = \alpha^2\lambda^2$. This simplifies (4.2) to the ordinary differential equation

$$(e^{2f})' + \frac{3}{4}\alpha^2|A|^2 (e^{-2f})' - 2\alpha^2 f'^3 = C_0 = const. \tag{4.3}$$

A solution of the last equation for $C_0 = 0$ was found in [26, Sect. 4.2]. For ease of reading we repeat the key steps of the derivation in order to obtain a seven dimensional solution of the Strominger system. The substitution $u = \alpha^{-2}e^{2f}$ allows us to write (4.3) in the form

$$(e^{2f})' + \frac{3}{4}\alpha^2|A|^2 (e^{-2f})' - 2\alpha^2 f'^3 = \frac{\alpha^2 u'}{4u^3} (4u^3 - 3\frac{|A|^2}{\alpha^2} u - u^2).$$

For $C_0 = 0$ we shall solve the following ordinary differential equation for the function $u = u(x^1) > 0$

$$u^2 = 4u^3 - 3\frac{|A|^2}{\alpha^2} u = 4u(u-d)(u+d), \quad d = \sqrt{3|A|^2}/\alpha. \tag{4.4}$$

Replacing the real derivative with the complex derivative leads to the Weierstrass' equation

$$\left(\frac{d\mathcal{P}}{dz}\right)^2 = 4\mathcal{P}(\mathcal{P}-d)(\mathcal{P}+d) \tag{4.5}$$

for the doubly periodic Weierstrass \mathcal{P} function with a pole at the origin. As well known, [1, 21], near the origin \mathcal{P} has the expansion

$$\mathcal{P}(z) = \frac{1}{z^2} + \frac{d^2}{5}z^2 + d_1z^6 + \dots,$$

which has no z^4 term and only even powers of z . Furthermore, see [1, 21], letting τ_\pm be the basic half-periods such that τ_+ is real and τ_- is purely imaginary we have that \mathcal{P} is

real valued on the lines $\Re z = m\tau_+$ or $\Im z = im\tau_-$, $m \in \mathbb{Z}$. In the fundamental region centered at the origin, where \mathcal{P} has a pole of order two, we have that $\mathcal{P}(z)$ decreases from $+\infty$ to a to 0 to $-a$ to $-\infty$ as z varies along the sides of the half-period rectangle from 0 to τ_+ to $\tau_+ + \tau_-$ to τ_- to 0 .

Thus, $u(x^1) = \mathcal{P}(x^1)$ defines a non-negative $2\tau_+$ -periodic function with singularities at the points $2n\tau_+$, $n \in \mathbb{Z}$, which solves the real Eq. (4.4). From the Laurent expansion of the Weierstrass' function it follows

$$u(x_1) = \frac{1}{(x^1)^2} \left(1 + \frac{d^2}{5}(x^1)^4 + \dots \right).$$

By construction, $f = \frac{1}{2} \ln(\alpha^2 u)$ is a periodic function with singularities on the real line which is a solution to Eq. (4.2). Therefore the G_2 -structure defined by $\bar{\Theta}$ descends to the 7-dimensional nilmanifold $M^7 = \Gamma \backslash K_A$ with singularity, determined by the singularity of u , where K_A is the 2-step nilpotent Lie group with Lie algebra \mathfrak{K}_A , defined by (3.2), and Γ is a lattice with the same period as f , i.e., $2\tau_+$ in all variables. In fact, as seen from the asymptotic behavior of u , M^7 is the total space of a \mathbb{T}^3 -bundle over the asymptotically hyperbolic manifold M^4 with metric

$$\bar{g}_H = u(x^1) \left((dx^1)^2 + (dx^2)^2 + (dx^3)^2 + (dx^4)^2 \right),$$

which is a conformally compact 4-torus with conformal boundary at infinity a flat 3-torus. Thus, we conclude that there is a complete solution with non-constant dilaton, non-trivial instanton and flux and with a negative α' parameter.

The last statement follows from the fact that the $(-)$ -connection is an instanton up to the first order of α' . This completes the proof of Theorem 4.3. \square

From the apparent \mathbb{Z}_2 -symmetry of u determined by the symmetry with respect to the line $x^1 = \tau_+$ we also obtain a solution on the quotient M^7/\mathbb{Z}_2 .

5. A Complete Solution with Positive α'

In this section we exhibit a solution of the Strominger system using again the G_2 -structure (3.7) by solving the anomaly cancellation condition with torsion term (3.16), the Pontrjagin form (3.20) of the ∇^- connection, and the G_2 -instanton defined below with the help of Lemma 5.1.

Lemma 5.1. *The $(-)$ -connection of the G_2 -structure $\bar{\Theta}$ is a G_2 instanton with respect to $\bar{\Theta}$ if and only if the torsion 3-form is closed, $d\bar{T} = 0$, i.e. the dilaton function f satisfies the equality*

$$\Delta e^{2f} + 2|A|^2 = 0. \tag{5.1}$$

Proof. Let $\{\bar{e}_1, \dots, \bar{e}_7\}$ be the orthonormal basis dual to $\{\bar{e}^1, \dots, \bar{e}^7\}$. Using (1.6) we investigate the G_2 instanton condition (1.5) for R^- as follows

$$\begin{aligned} 0 &= \sum_{i,j=1}^7 R^-(\bar{e}_i, \bar{e}_j, \bar{e}_l, \bar{e}_m) \bar{\Theta}(\bar{e}_i, \bar{e}_j, \bar{e}_k) \\ &= \sum_{i,j=1}^7 [R^+ - d\bar{T}](\bar{e}_i, \bar{e}_j, \bar{e}_l, \bar{e}_m) \bar{\Theta}(\bar{e}_i, \bar{e}_j, \bar{e}_k) \\ &= - \sum_{i,j=1}^7 d\bar{T}(\bar{e}_i, \bar{e}_j, \bar{e}_l, \bar{e}_m) \bar{\Theta}(\bar{e}_i, \bar{e}_j, \bar{e}_k), \end{aligned} \tag{5.2}$$

where we used the fact that the holonomy of ∇^+ is contained in G_2 , i.e. $\sum_{i,j=1}^7 R^+(\bar{e}_i, \bar{e}_j, \bar{e}_l, \bar{e}_m)\bar{\Theta}(\bar{e}_i, \bar{e}_j, \bar{e}_k) = 0$. Now, applying (3.16) and (3.7) we conclude that (5.2) is satisfied if and only if (5.1) holds. \square

Let D_B be the ∇^- connection obtained by replacing A with the matrix B in Lemma 5.1, but allowing B to be singular, $B \in \mathfrak{gl}_3(\mathbb{R})$. Hence, the connection D_B is a G_2 -instanton with respect to the G_2 -structure defined by (3.7) iff the dilaton function satisfies

$$\Delta e^{2f} = -2|B|^2. \tag{5.3}$$

Equation (3.20) shows that the difference between the first Pontrjagin forms of ∇^- and D_B is given by the formula

$$8\pi^2 \left(p_1(\nabla^-) - p_1(D_B) \right) = -3 \left(|A|^2 - |B|^2 \right) \left(\Delta e^{-2f} \right) e^{1234}. \tag{5.4}$$

Therefore, recalling (3.16) and taking into account (5.4), the anomaly cancellation condition is

$$\begin{aligned} d\bar{T} - \frac{\alpha'}{4} 8\pi^2 \left(p_1(\nabla^-) - p_1(D_B) \right) \\ = - \left[\Delta e^{2f} + 2|A|^2 - \frac{3}{4}\alpha' \left(|A|^2 - |B|^2 \right) \left(\Delta e^{-2f} \right) \right] e^{1234} = 0 \end{aligned}$$

coupled with (5.3). Notice that at this point the analysis can proceed exactly as in [26, Sect. 5.2]. As a result we obtain the following results depending on the difference $|A|^2 - |B|^2$ being zero or non-zero.

For $B = O$, where O is the zero matrix in $\mathfrak{gl}_3(\mathbb{R})$, and a fixed $e \in \mathbb{R}^4$ we let

$$e^{2f} = \frac{3\alpha'}{4|x - e|^2}, \quad x \in \mathbb{R}^4. \tag{5.5}$$

Using logarithmic radial coordinates near the singularity (as e.g. in [15]) it follows that the 4-D metric induced on \mathbb{R}^4 is actually complete. In fact, taking the singularity at the origin, in the coordinate $t = \sqrt{3\alpha'}/2 \ln(4|x|^2/3\alpha') = -\sqrt{3\alpha'} f$, we have that the dilaton and the 4-D metric can be expressed as follows

$$f = -t\sqrt{3\alpha'}, \quad \bar{g}_H = \sum_{i=1}^4 e^{2f} (e^i)^2 = dt^2 + 3\alpha' ds_3^2,$$

where ds_3^2 is the metric on the unit three-dimensional sphere in the four dimensional Euclidean space. The completeness of the horizontal metric implies that the metric $\bar{g} = \bar{g}_H + (e^5)^2 + (e^6)^2 + (e^7)^2$ is also complete. Thus, we proved.

Theorem 5.2. *The non-compact complete simply connected manifold $(K_A, \bar{\Theta}, \nabla^-, D_O, f)$ described above is a complete G_2 manifold which solves the Strominger system with non-constant dilaton f determined by (5.5), non-zero flux $H = \bar{T}$ and non-flat instanton D_O using the first Pontrjagin form of ∇^- and positive α' . Furthermore, $(K_A, \bar{\Theta}, \nabla^-, D_O, f)$ also solves the heterotic equations of motion (1.2) up to the first order of α' .*

On the other hand, in the case $|A|^2 = |B|^2 \neq 0$ the anomaly condition is trivially satisfied for any α' , provided the torsion is closed, see Lemma 5.1. In this case the solution is given by the solutions of (5.1). Furthermore, both ∇^- and D_B are G_2 -instantons. For example, a particular solution is obtained by taking

$$e^{2f} = \frac{|A|^2}{4}(1 - |x|^2)$$

defined in the unit ball.

6. Solutions Through Contractions

In this section we consider appropriate contractions of the quaternionic Heisenberg algebra (3.2) leading to deformations of the geometric structures, the partial differential equations and their solutions found in Sects. 4 and 5 in the G_2 -heterotic case. We show that these G_2 solutions converge to the heterotic solutions on the 6-dimensional inner non-Kähler spaces constructed in [26]. Furthermore, this method allows us to find new heterotic solutions with non-constant dilaton in dimension 5.

6.1. Six dimensional solutions. Using the classification results of [69], it was shown in [26] that 2-step nilmanifolds which are \mathbb{T}^2 -bundles over \mathbb{T}^4 , with connection 1-form the anti-self-dual curvature of \mathbb{T}^4 , are precisely the 6-dimensional nilmanifolds admitting invariant balanced Hermitian metrics with Abelian complex structure J , i.e., $[JX, JY] = [X, Y]$. Moreover, in this case the Lie algebra underlying M is isomorphic to \mathfrak{h}_3 or \mathfrak{h}_5 . Here, \mathfrak{h}_3 is the Lie algebra underlying the nilmanifold given by the product of the 5-dimensional generalized Heisenberg nilmanifold by S^1 , while \mathfrak{h}_5 is the Lie algebra underlying the Iwasawa manifold. The structure equations of the Lie algebra \mathfrak{h}_5 are

$$de^1 = de^2 = de^3 = de^4 = 0, \quad de^5 = b\sigma_2, \quad de^6 = a\sigma_1 - b\sigma_3, \quad (6.1)$$

while \mathfrak{h}_3 is given by

$$de^1 = de^2 = de^3 = de^4 = 0, \quad de^5 = 0, \quad de^6 = a\sigma_1,$$

where $a, b \in \mathbb{R}^*$ and σ_i are the anti-self-dual forms on \mathbb{R}^4 , see after (3.2). Clearly \mathfrak{h}_3 is a contraction of \mathfrak{h}_5 and both are contractions of $\mathfrak{g}(\mathbb{H})$, see (3.2).

It is remarkable that the geometric structures, the partial differential equations and their solutions found in Sects. 4 and 5 converge to the heterotic solutions on the 6-dimensional inner non-Kähler spaces found in [26] as we explain next in details for \mathfrak{h}_5 . The $SU(3)$ -structure and corresponding solution based on \mathfrak{h}_3 is handled analogously.

Clearly \mathfrak{h}_5 is a contraction of \mathfrak{K}_A when $\varepsilon \rightarrow 0$ using, for example,

$$A_\varepsilon \stackrel{def}{=} \begin{pmatrix} 0 & b & 0 \\ a & 0 & -b \\ 0 & 0 & \varepsilon \end{pmatrix}.$$

Notice that by (3.8) we have $\bar{e}^7 = e_\varepsilon^7 = \varepsilon\gamma^7 \rightarrow 0$ as $\varepsilon \rightarrow 0$. With the above choice of A_ε we write the G_2 -form (3.7) in the usual way as

$$\bar{\Theta}_\varepsilon = \bar{F} \wedge e_\varepsilon^7 + \bar{\Psi}^+, \quad \bar{F} = e^{2f}\omega_1 + e^{5f}, \quad \bar{\Psi}^+ = e^{2f}(\omega_2 \wedge e^5 - \omega_3 \wedge e^6)$$

using (3.8) and indicating with subscript ε the dependence on ε through the matrix A_ε . In addition, we let $\tilde{\Psi}^- = e^{2f}(\omega_2 \wedge e^6 + \omega^3 \wedge e^5)$. In the limit $\varepsilon \rightarrow 0$, the forms \tilde{F} , $\tilde{\Psi}^\pm$ define an $SU(3)$ -structure $(\tilde{F}, \tilde{\Psi}^\pm)$ on a six dimensional space, obtained through the ansatz proposed in [38] from a \mathbb{T}^2 bundle over \mathbb{T}^4 (corresponding to $f = 0$), see [26, Sect. 3.2] for details in the case of \mathfrak{h}_5 . Therefore, this $SU(3)$ structure solves the first two Killing spinor equations. Furthermore, the Pontrjagin form of the ∇^- connection is given again by (3.20) as shown in [26, Sect. 3]. In fact, the connection forms (3.8) and the corresponding curvature 2-forms (notice that $(\Omega_\varepsilon^-)^{\tilde{i}} \rightarrow 0$ for all i) converge to those of the ∇^- connection of the $SU(3)$ case. Similarly, the seven dimensional anomaly cancellation conditions of Sects. 4 and 5 turn into the anomaly cancellation conditions for the corresponding six dimensional structures. As a consequence we obtain the six-dimensional solutions with non-constant dilaton found in [26].

6.2. *Five dimensional solutions.* We begin with recalling the five dimensional Lie algebra $\mathfrak{h}(2, 1)$ [23] with structure equations

$$de^j = 0, \quad j = 1, 2, 3, 4, \quad de^5 = \sum_{i=1}^3 a_i \sigma_i, \quad a_i \in \mathbb{R}, \quad (a_1, a_2, a_3) \neq (0, 0, 0). \tag{6.2}$$

Without loss of generality we will suppose next that $a_1 \neq 0$. Clearly $\mathfrak{h}(2, 1)$ is a contraction of \mathfrak{K}_A , see (3.2), using, for example,

$$A_\varepsilon \stackrel{def}{=} \begin{pmatrix} a_1 & a_2 & a_3 \\ 0 & \varepsilon & 0 \\ 0 & 0 & \varepsilon \end{pmatrix}$$

and letting $\varepsilon \rightarrow 0$. Notice that by (3.8) we have

$$\bar{e}^i = e^i_\varepsilon = \varepsilon \gamma^i \rightarrow 0, \quad i = 6, 7 \tag{6.3}$$

when $\varepsilon \rightarrow 0$.

It was shown in [23, Sect. 4] that the $SU(2)$ -structure $(e^5, \omega_1, \omega_2, \omega_3)$ is the unique family of left invariant solutions (with constant dilaton) to the first two Killing spinor equations on a five dimensional Lie group. Furthermore, [23] continued on showing that for $\nabla = \nabla^+$ or $\nabla = \nabla^g$ and suitably defined instantons one can obtain compact (nil-manifolds) heterotic solutions with constant dilaton. However, since the first Pontrjagin form of the connection ∇^- vanishes there is no compact solution with constant dilaton to the heterotic supersymmetry equations satisfying the anomaly cancellation condition with $\nabla = \nabla^-$.

From the current point of view, we consider the case $\nabla = \nabla^-$ as a contraction limit of the G_2 -solutions in Sects. 4 and 5. As a result we will obtain five dimensional solutions with non-constant dilaton. Indeed, applying (6.3) and (6.2) to (3.15) we obtain the expression for the torsion in dimension five described in (2.3). In other words, the torsion in dimension five is obtained as a dimensional reduction of the torsion in dimension seven. Furthermore, the Pontrjagin form of the ∇^- connection is given again by (3.20) taking into account (6.2). In fact, the connection forms (3.8) and the corresponding curvature 2-forms (notice that $(\Omega_\varepsilon^-)^{\tilde{i}} \rightarrow 0$ and $(\Omega_\varepsilon^-)^{\tilde{j}} \rightarrow 0$ for all i) converge to those

of the ∇^- connection of the $SU(2)$ structure in dimension five. Similarly, the seven dimensional anomaly cancellation conditions of Sects. 4 and 5 turn into the anomaly cancellation conditions for the corresponding five dimensional structures. At this point we turn to the construction of the five dimensional solutions with non-constant dilaton.

The five dimensional version of Theorem 4.3 is Theorem 6.2 below. In the statement of Theorem 6.2 we use (3.9), i.e., $\bar{\omega}_i = e^{2f} \omega_i$, $i = 1, 2, 3, 4$. Let $H(2, 1)$ be the five dimensional connected simply connected Lie group $H(2, 1)$ with Lie algebra $\mathfrak{h}(2, 1)$, We consider a lattice Γ in the Lie group $H(2, 1)$ with period $2\tau_+$ in all variables, where $2\tau_+$ is the period of the Weierstrass' \mathcal{P} function (4.5). The $SU(2)$ instanton D_Λ below corresponds to the instanton obtained from the one in Proposition 4.1 by setting the last two columns equal to zero as well as letting $\varepsilon \rightarrow 0$, see (6.3).

Lemma 6.1. *Let D_Λ , $\Lambda = (\lambda_1, \lambda_2, \lambda_3) \in \mathbb{R}^3$, be the linear connection on the Lie group $H(2, 1)$ whose possibly non-zero 1-forms are given as follows*

$$\begin{aligned} (\omega^{D_\Lambda})_{\bar{2}}^{\bar{1}} &= -(\omega^{D_\Lambda})_{\bar{1}}^{\bar{2}} = -(\omega^{D_\Lambda})_{\bar{4}}^{\bar{3}} = (\omega^{D_\Lambda})_{\bar{3}}^{\bar{4}} = \lambda_1 \bar{e}^5, \\ (\omega^{D_\Lambda})_{\bar{3}}^{\bar{1}} &= -(\omega^{D_\Lambda})_{\bar{1}}^{\bar{3}} = (\omega^{D_\Lambda})_{\bar{4}}^{\bar{2}} = -(\omega^{D_\Lambda})_{\bar{2}}^{\bar{4}} = \lambda_2 \bar{e}^5, \\ (\omega^{D_\Lambda})_{\bar{4}}^{\bar{1}} &= -(\omega^{D_\Lambda})_{\bar{1}}^{\bar{4}} = -(\omega^{D_\Lambda})_{\bar{3}}^{\bar{2}} = (\omega^{D_\Lambda})_{\bar{2}}^{\bar{3}} = \lambda_3 \bar{e}^5. \end{aligned}$$

Then, D_Λ is an $SU(2)$ -instanton with respect to the $SU(2)$ structure defined by $(e^5, \bar{\omega}_1, \bar{\omega}_2, \bar{\omega}_3)$.

We skip the proof which is similar to the proof of Proposition 4.1. The five dimensional version of Theorem 4.3 follows.

Theorem 6.2. *Let $(e^5, \bar{\omega}_1, \bar{\omega}_2, \bar{\omega}_3)$ be the $SU(2)$ -structure on the Lie group $H(2, 1)$. The conformally compact five manifold $M^5 = (\Gamma \backslash H(2, 1), \eta^5, \bar{\omega}_1, \bar{\omega}_2, \bar{\omega}_3, \nabla^-, D_\Lambda, f)$ is a conformally quasi-Sasakian five manifold which solves the Strominger system with non-constant dilaton f , non-trivial flux $H = \bar{T}$ and non-flat instanton D_Λ using the first Pontrjagin form of ∇^- and negative α' . The dilaton f depends on one variable and is determined as a real slice of the Weierstrass' elliptic function. In addition, M^5 satisfies the heterotic equations of motion (1.2) up to first order of α' .*

In order to obtain the five dimensional version of Theorem 5.2 we use the following property of the ∇^- connection whose 1-forms are

$$\begin{aligned} (\omega^-)_{\bar{2}}^{\bar{1}} &= (\omega^-)_{\bar{4}}^{\bar{3}} = e^{-f} \left(f_2 \bar{e}^1 - f_1 \bar{e}^2 + f_4 \bar{e}^3 - f_3 \bar{e}^4 \right), \\ (\omega^-)_{\bar{3}}^{\bar{1}} &= -(\omega^-)_{\bar{4}}^{\bar{2}} = e^{-f} \left(f_3 \bar{e}^1 - f_4 \bar{e}^2 - f_1 \bar{e}^3 + f_2 \bar{e}^4 \right), \\ (\omega^-)_{\bar{4}}^{\bar{1}} &= (\omega^-)_{\bar{3}}^{\bar{2}} = e^{-f} \left(f_4 \bar{e}^1 + f_3 \bar{e}^2 - f_2 \bar{e}^3 - f_1 \bar{e}^4 \right), \\ (\omega^-)_{\bar{5}}^{\bar{1}} &= e^{-2f} \left(-a_{11} \bar{e}^2 - a_{12} \bar{e}^3 - a_{13} \bar{e}^4 \right), \quad (\omega^-)_{\bar{5}}^{\bar{2}} = e^{-2f} \left(a_{11} \bar{e}^1 + a_{13} \bar{e}^3 - a_{12} \bar{e}^4 \right), \\ (\omega^-)_{\bar{5}}^{\bar{3}} &= e^{-2f} \left(a_{12} \bar{e}^1 - a_{13} \bar{e}^2 + a_{11} \bar{e}^4 \right), \quad (\omega^-)_{\bar{5}}^{\bar{4}} = e^{-2f} \left(a_{13} \bar{e}^1 + a_{12} \bar{e}^2 - a_{11} \bar{e}^3 \right), \end{aligned}$$

which are obtained from (3.19) taking into account (6.3).

Lemma 6.3. *The $(-)$ -connection of the $SU(2)$ structure $(e^5, \bar{\omega}_1, \bar{\omega}_2, \bar{\omega}_3)$ is an $SU(2)$ instanton iff the torsion 3-form is closed, $d\bar{T} = 0$, i.e., the dilaton function f satisfies Eq. (5.1).*

The proof of Lemma 6.3 is very similar to the proof of Lemma 5.1 and involves a direct calculation. Let D_O be the $SU(2)$ instanton constructed by Lemma 6.3 in the case $A = O$ -the zero:

Theorem 6.4. *The non-compact simply connected five manifold $(H(2, 1), e^5, \bar{\omega}_1, \bar{\omega}_2, \bar{\omega}_3, \nabla^-, D_O, f)$ is a complete conformally quasi-Sasakian five manifold which solves the Strominger system with non-constant dilaton f determined by (5.5), non-trivial flux $H = \bar{T}$, non-flat instanton D_O using the first Pontrjagin form of ∇^- and positive α' .*

The complete five manifold $(H(2, 1), e^5, \bar{\omega}_1, \bar{\omega}_2, \bar{\omega}_3, \nabla^-, D_O, f)$ satisfies the heterotic equations of motion (1.2) up to first order of α' .

Acknowledgements. The work was partially supported through Project MICINN (Spain) MTM2011-28326-C02-01/02, and Project of UPV/EHU ref. UFI11/52. S.I. is partially supported by Contract DFNI I02/4/12.12.2014 and Contract 148/2015 with the Sofia University “St.Kl.Ohridski”. D.V. was partially supported by the Simons Foundation grant #279381. S.I. and D.V. thank the University of Zaragoza and the University of the Basque Country for the hospitality and financial support provided while visiting the respective Departments of Mathematics.

References

- Ahlfors, L.V.: Complex analysis: an introduction of the theory of analytic functions of one complex variable. McGraw-Hill Book Co., New York, Toronto, London (1966)
- Anderson, L.B., Gray, J., Sharpe, E.: Algebroids, Heterotic Moduli Spaces and the Strominger system. *JHEP* **1407**, 037 (2014)
- Andreas, B., García-Fernández, M.: Heterotic non-Kähler geometries via polystable bundles on Calabi-Yau threefolds. *J. Geom. Phys.* **62**, 183–188 (2012)
- Andreas, B., García-Fernández, M.: Solutions of the Strominger system via stable bundles on Calabi-Yau threefolds. *Commun. Math. Phys.* **332**, 1381–1383 (2014)
- Andreas, B., García-Fernández, M.: Note on solutions of the Strominger system from unitary representations of cocompact lattices of $SL(2, \mathbb{C})$. *Commun. Math. Phys.* **315**, 153–168 (2012)
- Becker, K., Becker, M., Dasgupta, K., Green, P.S.: Compactifications of heterotic theory on non-Kähler complex manifolds: I. *JHEP* **0304**, 007 (2003)
- Becker, K., Becker, M., Dasgupta, K., Green, P.S., Sharpe, E.: Compactifications of heterotic strings on non-Kähler complex manifolds: II. *Nuclear Phys. B* **678**, 19–100 (2004)
- Becker, K., Becker, M., Dasgupta, K., Prokushkin, S.: Properties of heterotic vacua from superpotentials. *Nuclear Phys. B* **666**, 144–174 (2003)
- Becker, K., Becker, M., Fu, J.-X., Tseng, L.-S., Yau, S.-T.: Anomaly cancellation and smooth non-Kähler solutions in heterotic string theory. *Nuclear Phys. B* **751**, 108–128 (2006)
- Becker, K., Bertinato, C., Chung, Y.-C., Guo, G.: Supersymmetry breaking, heterotic strings and fluxes. *Nuclear Phys. B* **823**, 428–447 (2009)
- Becker, K., Sethi, S.: Torsional heterotic geometries. *Nuclear Phys. B* **820**, 1–31 (2009)
- Bena, I., Bobev, N., Warner, N.: Bubbles on manifolds with $U(1)$ isometry. *JHEP* **0708**, 004 (2007)
- Bergshoeff, E.A., Roo, M.de : The quartic effective action of the heterotic string and supersymmetry. *Nuclear Phys. B* **328**, 439–468 (1989)
- Biswas, I., Mukherjee, A.: Solutions of Strominger system from unitary representations of cocompact lattices of $SL(2, \mathbb{C})$, *Commun. Math. Phys.* **322**, 373–384 (2013)
- Callan, C.G., Harvey, J.A., Strominger, A.: Worldsheet approach to heterotic instantons and solitons. *Nuclear Phys. B* **359**, 611–634 (1991)
- Cardoso, G.L., Curio, G., Dall’Agata, G., Lüster, D.: BPS action and superpotential for heterotic string compactifications with fluxes. *JHEP* **0310**, 004 (2003)
- Cardoso, G.L., Curio, G., Dall’Agata, G., Lüster, D., Manousselis, P., Zoupanos, G.: Non-Kähler string backgrounds and their five torsion classes. *Nuclear Phys. B* **652**, 5–34 (2003)
- Conti, D., Salamon, S.: Killing spinors in dimension 5. *Trans. Amer. Math. Soc.* **359**, 5319–5343 (2007)
- Dasgupta, K., Firouzjahi, H., Gwyn, R.: On the warped heterotic axion. *JHEP* **0806**, 056 (2008)
- Wit, B.de, Smit, D.J., Hari Dass, N.D.: Residual supersymmetry of compactified $D=10$ supergravity. *Nuclear Phys. B* **283**, 165–191 (1987)
- Erdélyi, A., Magnus, W., Oberhettinger, F., Tricomi, F.G.: Higher transcendental functions. Vol. I, II and III. Robert E. Krieger Publishing Co., Inc., Melbourne (1981)

22. Fei, T., Yau, S.-T.: Invariant Solutions to the Strominger system on Complex Lie Groups and Their Quotients. [arXiv:1407.7641](https://arxiv.org/abs/1407.7641)
23. Fernández, M., Ivanov, S., Ugarte, L., Villacampa, R.: Compact supersymmetric solutions of the heterotic equations of motion in dimension 5. *Nuclear Phys.* **B 820**, 483–502 (2009)
24. Fernández, M., Ivanov, S., Ugarte, L., Villacampa, R.: Non-Kähler heterotic-string compactifications with non-zero fluxes and constant dilaton. *Commun. Math. Phys.* **288**, 677–697 (2009)
25. Fernández, M., Ivanov, S., Ugarte, L., Villacampa, R.: Compact supersymmetric solutions of the heterotic equations of motion in dimensions 7 and 8. *Adv. Theor. Math. Phys.* **15**, 245–284 (2011)
26. Fernández, M., Ivanov, S., Ugarte, L., Vassilev, D.: Non-Kähler heterotic string solutions with non-zero fluxes and non-constant dilaton. *JHEP* **1406**, 073 (2014)
27. Friedrich, Th., Ivanov, S.: Parallel spinors and connections with skew-symmetric torsion in string theory. *Asian J. Math.* **6**, 303–336 (2002)
28. Friedrich, Th., Ivanov, S.: Killing spinor equations in dimension 7 and geometry of integrable G_2 manifolds. *J. Geom. Phys.* **48**, 1–11 (2003)
29. Friedrich, Th., Ivanov, S.: Almost contact manifolds, connections with torsion and parallel spinors. *J. Reine Ang. Math.* **559**, 217–236 (2003)
30. Fu, J.-X., Tseng, L.-S., Yau, S.-T.: Local heterotic torsional models. *Commun. Math. Phys.* **289**, 1151–1169 (2009)
31. Fu, J.-X., Yau, S.-T.: The theory of superstring with flux on non-Kähler manifolds and the complex Monge-Ampère equation. *J. Diff. Geom.* **78**, 369–428 (2008)
32. Gauntlett, J.P., Gutowski, J., Hull, C., Pakis, S., Reall, H.: All supersymmetric solutions of minimal supergravity in five dimensions. *Class. Quantum Grav.* **20**, 4587–4634 (2003)
33. Gauntlett, J.P., Kim, N., Martelli, D., Waldram, D.: Fivebranes wrapped on SLAG three-cycles and related geometry. *JHEP* **0111**, 018 (2001)
34. Gauntlett, J.P., Martelli, D., Pakis, S., Waldram, D.: G -structures and wrapped NS5-branes. *Commun. Math. Phys.* **247**, 421–445 (2004)
35. Gauntlett, J.P., Martelli, D., Waldram, D.: Superstrings with intrinsic torsion. *Phys. Rev. D* **69**, 086002 (2004)
36. Giddings, S.B., Kachru, S., Polchinski, J.: Hierarchies from fluxes in string compactifications. *Phys. Rev. D* **66**, 106006 (2002)
37. Gillard, J., Papadopoulos, G., Tsimpis, D.: Anomaly, fluxes and (2, 0) heterotic-string compactifications. *JHEP* **0306**, 035 (2003)
38. Goldstein, E., Prokushkin, S.: Geometric model for complex non-Kähler manifolds with $SU(3)$ structure. *Commun. Math. Phys.* **251**, 65–78 (2004)
39. Gran, U., Lohrmann, P., Papadopoulos, G.: The spinorial geometry of supersymmetric heterotic string backgrounds. *JHEP* **0602**, 063 (2006)
40. Gran, U., Papadopoulos, G., Roest, D., Sloane, P.: Geometry of all supersymmetric type I backgrounds. *JHEP* **0708**, 074 (2007)
41. Gran, U., Papadopoulos, G., Roest, D.: Supersymmetric heterotic string backgrounds. *Phys. Lett. B* **656**, 119–126 (2007)
42. Gran, U., Papadopoulos, G.: Solution of heterotic Killing spinor equations and special geometry. Special metrics and supersymmetry. In: *AIP Conf. Proc., Amer. Inst. Phys.*, vol. 1093, pp. 144–161. Melville (2009)
43. Günaydin, M., Nikolai, H.: Seven-dimensional octonionic Yang-Mills instanton and its extension to an heterotic string soliton. *Phys. Lett. B* **353**, 169 (1991)
44. Howe, P.S., Papadopoulos, G.: Ultraviolet behavior of two-dimensional supersymmetric non-linear sigma models. *Nuclear Phys. B* **289**, 264–276 (1987)
45. Hull, C.M.: Compactifications of the heterotic superstring. *Phys. Lett. B* **178**, 357–364 (1986)
46. Hull, C.M.: Anomalies, ambiguities and superstrings. *Phys. Lett. B* **167**, 51–55 (1986)
47. Hull, C.M., Townsend, P.K.: The two loop beta function for sigma models with torsion. *Phys. Lett. B* **191**, 115–121 (1987)
48. Hull, C.M., Witten, E.: Supersymmetric sigma models and the heterotic string. *Phys. Lett. B* **160**, 398–402 (1985)
49. Imazato, H., Mizoguchi, S., Yata, M.: Taub-NUT crystal. *Int. J. Mod. Phys. A* **26**, 5143–5169 (2011)
50. Ivanov, P., Ivanov, S.: $SU(3)$ -instantons and G_2 , $Spin(7)$ -Heterotic string solitons. *Commun. Math. Phys.* **259**, 79–102 (2005)
51. Ivanov, S.: Heterotic supersymmetry, anomaly cancellation and equations of motion. *Phys. Lett. B* **685**, 190–196 (2010)
52. Ivanov, S., Papadopoulos, G.: Vanishing theorems and string backgrounds. *Class. Quantum Grav.* **18**, 1089–1110 (2001)
53. Ivanov, S., Papadopoulos, G.: A no-go theorem for string warped compactifications. *Phys. Lett. B* **497**, 309–316 (2001)

54. Kimura, T., Yi, P.: Comments on heterotic flux compactifications. *JHEP* **0607**, 030 (2006)
55. Kimura, T., Mizoguchi, S.: Chiral generations on intersecting 5-branes in heterotic string theory. *JHEP* **1004**, 028 (2010)
56. Li, J., Yau, S.-T.: The existence of supersymmetric string theory with torsion. *J. Diff. Geom.* **70**, 143–181 (2005)
57. Lu, H., Vazquez-Poritz, J.F.: Resolution of overlapping branes. *Phys. Lett.* **B 534**, 155 (2002)
58. Martelli, D., Sparks, J.: Non-Kähler heterotic rotations. *Adv. Theor. Math. Phys.* **15**, 131–174 (2011)
59. Maxfield, T., Sethi, S.: Domain Walls, Triples and Acceleration. *JHEP* **1408**, 066 (2014)
60. Melnikov, I.V., Minasian, R., Sethi, S.: Heterotic fluxes and supersymmetry. *JHEP* **1406**, 174 (2014)
61. Mizoguchi, S., Yata M.: Family unification via quasi-Nambu-Goldstone fermions in string theory. *Prog. Theor. Exp. Phys.* 053B01 (2013)
62. de la Ossa, X., Svanes, E.E.: Holomorphic bundles and the moduli space of $N=1$ supersymmetric heterotic compactifications. *JHEP* **1410**, 123 (2014)
63. de la Ossa, X., Svanes, E.E.: Connections, Field Redefinitions and Heterotic Supergravity. *JHEP* **1412**, 008 (2014)
64. Papadopoulos, G.: New half supersymmetric solutions of the heterotic string. *Class. Quantum Grav.* **26**, 135001 (2009)
65. Sen, A.: $(2, 0)$ supersymmetry and space-time supersymmetry in the heterotic string theory. *Nuclear Phys.* **B 278**, 289–308 (1986)
66. Stotyn, S., Mann, R.: Supergravity on an Atiyah-Hitchin base. *JHEP* **0806**, 087 (2008)
67. Strominger, A.: Heterotic solitons. *Nuclear Phys.* **B 343**, 167–184 (1990) [Erratum-ibid. **B 353** (1991) 565]
68. Strominger, A.: Superstrings with torsion. *Nuclear Phys.* **B 274**, 253–284 (1986)
69. Ugarte, L., Villacampa, R.: Balanced Hermitian geometry on 6-dimensional nilmanifolds. *Forum Math.* **27**, 1025–1070 (2015)

Communicated by N. A. Nekrasov