



Weak Solutions to the Navier–Stokes Inequality with Arbitrary Energy Profiles

Wojciech S. Ożański

Department of Mathematics, University of Southern California, Los Angeles, USA.
E-mail: ozanski@usc.edu

Received: 18 September 2018 / Accepted: 6 May 2019
Published online: 30 October 2019 – © Springer-Verlag GmbH Germany, part of Springer Nature 2019

Abstract: In a recent paper, Buckmaster and Vicol (Ann Math (2) 189(1):101–144, 2019) used the method of convex integration to construct weak solutions u to the 3D incompressible Navier–Stokes equations such that $\|u(t)\|_{L^2} = e(t)$ for a given non-negative and smooth energy profile $e: [0, T] \rightarrow \mathbb{R}$. However, it is not known whether it is possible to extend this method to construct nonunique *suitable weak solutions* (that is weak solutions satisfying the strong energy inequality (SEI) and the *local energy inequality* (LEI)), Leray–Hopf weak solutions (that is weak solutions satisfying the SEI), or at least to exclude energy profiles that are not nonincreasing. In this paper we are concerned with weak solutions to the Navier–Stokes inequality on \mathbb{R}^3 , that is vector fields that satisfy both the SEI and the LEI (but not necessarily solve the Navier–Stokes equations). Given $T > 0$ and a nonincreasing energy profile $e: [0, T] \rightarrow [0, \infty)$ we construct weak solution to the Navier–Stokes inequality that are localised in space and whose energy profile $\|u(t)\|_{L^2(\mathbb{R}^3)}$ stays arbitrarily close to $e(t)$ for all $t \in [0, T]$. Our method applies only to nonincreasing energy profiles. The relevance of such solutions is that, despite not satisfying the Navier–Stokes equations, they satisfy the partial regularity theory of Caffarelli et al. (Commun Pure Appl Math 35(6):771–831, 1982). In fact, Scheffer’s constructions of weak solutions to the Navier–Stokes inequality with blow-ups (Commun Math Phys 101(1):47–85, 1985; Commun Math Phys 110(4): 525–551, 1987) show that the Caffarelli, Kohn & Nirenberg’s theory is sharp for such solutions. Our approach gives an indication of a number of ideas used by Scheffer. Moreover, it can be used to obtain a stronger result than Scheffer’s. Namely, we obtain weak solutions to the Navier–Stokes inequality with both blow-up and a prescribed energy profile.

1. Introduction

The Navier–Stokes equations,

$$\begin{aligned}\partial_t u - \nu \Delta u + (u \cdot \nabla)u + \nabla p &= 0, \\ \operatorname{div} u &= 0,\end{aligned}$$

where u denotes the velocity of a fluid, p the scalar pressure and $\nu > 0$ the viscosity, comprise a fundamental model for viscous, incompressible flows. In the case of the whole space \mathbb{R}^3 the pressure function is given (at each time instant t) by the formula

$$p := \sum_{i,j=1}^3 \partial_{ij} \Psi * (u_i u_j), \quad (1.1)$$

where $\Psi(x) := (4\pi|x|)^{-1}$ denotes the fundamental solution of the Laplace equation in \mathbb{R}^3 and “ $*$ ” denotes the convolution. The formula above, which we shall refer to simply as the *pressure function corresponding to u* , can be derived by calculating the divergence of the Navier–Stokes equation.

The fundamental mathematical theory of the Navier–Stokes equations goes back to the pioneering work of Leray [11] (see [14] for a comprehensive review of this paper in more modern language), who used a Picard iteration scheme to prove existence and uniqueness of local-in-time strong solutions. Moreover, [11] proved the global-in-time existence (without uniqueness) of weak solutions satisfying the strong energy inequality,

$$\|u(t)\|^2 + 2\nu \int_s^t \|\nabla u(\tau)\|^2 d\tau \leq \|u(s)\|^2 \quad (1.2)$$

for almost every $s \geq 0$ and every $t > s$ (often called *Leray–Hopf weak solutions*). Here $\|\cdot\|$ denotes the $L^2(\mathbb{R}^3)$ norm. (Hopf [8] proved an analogous result in the case of a bounded smooth domain.) Although the fundamental question of global-in-time existence and uniqueness of strong solutions remains unresolved, many significant results contributed to the theory of the Navier–Stokes equations during the second half of the twentieth century. One such contribution is the partial regularity theory introduced by Scheffer [18–22] and subsequently developed by Caffarelli, Kohn and Nirenberg [4]; see also Lin [12], Ladyzhenskaya and Seregin [10], Vasseur [25] and Kukavica [9] for alternative approaches. This theory is concerned with so-called *suitable weak solutions*, that is Leray–Hopf weak solutions that are also weak solutions of the *Navier–Stokes inequality* (NSI).

Definition 1.1 (*Weak solution to the Navier–Stokes inequality*). A divergence-free vector field $u: \mathbb{R}^3 \times (0, \infty)$ satisfying $\sup_{t>0} \|u(t)\| < \infty$, $\nabla u \in L^2(\mathbb{R}^3 \times (0, \infty))$ is a *weak solution of the Navier–Stokes inequality* with viscosity $\nu > 0$ if it satisfies the inequality

$$u \cdot (\partial_t u - \nu \Delta u + (u \cdot \nabla)u + \nabla p) \leq 0 \quad (1.3)$$

in a weak sense, that is

$$2\nu \int_0^\infty \int_{\mathbb{R}^3} |\nabla u|^2 \varphi \leq \int_0^\infty \int_{\mathbb{R}^3} (|u|^2 (\partial_t \varphi + \nu \Delta \varphi) + (|u|^2 + 2p)(u \cdot \nabla) \varphi) \quad (1.4)$$

for all non-negative $\varphi \in C_0^\infty(\mathbb{R}^3 \times (0, \infty))$, where p is the pressure function corresponding to u (recall (1.1)).

The last inequality is usually called the *local energy inequality*. The existence of global-in-time suitable weak solutions given divergence-free initial data $u_0 \in L^2$ was proved by Scheffer [20] in the case of the whole space \mathbb{R}^3 and by Caffarelli et al. [4] in the case of a bounded domain.

In order to see that (1.4) is a weak form of the NSI (1.3), note that the NSI can be rewritten, for smooth u and p , in the form

$$\frac{1}{2}\partial_t|u|^2 - \frac{\nu}{2}\Delta|u|^2 + \nu|\nabla u|^2 + u \cdot \nabla \left(\frac{1}{2}|u|^2 + p \right) \leq 0, \quad (1.5)$$

where we used the calculus identity $u \cdot \Delta u = \Delta(|u|^2/2) - |\nabla u|^2$. Multiplication by 2φ and integration by parts gives (1.4).

Furthermore, setting

$$f := \partial_t u - \nu \Delta u + (u \cdot \nabla)u + \nabla p,$$

one can think of the Navier–Stokes inequality (1.4) as the inhomogeneous Navier–Stokes equations with forcing f ,

$$\partial_t u - \nu \Delta u + (u \cdot \nabla)u + \nabla p = f,$$

where f acts against the direction of the flow u , that is $f \cdot u \leq 0$.

The partial regularity theory gives sufficient conditions for local regularity of suitable weak solutions in space-time. Namely, letting $Q_r(z) := B_r(x) \times (t - r^2, t)$, a space-time cylinder based¹ at $z = (x, t)$, the central result of this theory, proved by Caffarelli et al. [4], is the following.

Theorem 1.2 (Partial regularity of the Navier–Stokes equations). *Let $u_0 \in L^2(\mathbb{R}^3)$ be weakly divergence-free and let u be a suitable weak solution of the Navier–Stokes equations on \mathbb{R}^3 with initial condition u_0 . There exists a universal constant $\varepsilon_0 > 0$ such that if*

$$\frac{1}{r^2} \int_{Q_r} |u|^3 + |p|^{3/2} < \varepsilon_0 \quad (1.6)$$

for some cylinder $Q_r = Q_r(z)$, $r > 0$, then u is bounded in $Q_{r/2}(z)$.

Moreover there exists a universal constant $\varepsilon_1 > 0$ such that if

$$\limsup_{r \rightarrow 0} \frac{1}{r} \int_{Q_r} |\nabla u|^2 < \varepsilon_1 \quad (1.7)$$

then u is bounded in a cylinder $Q_\rho(z)$ for some $\rho > 0$.

(The theorem is valid also for suitable weak solutions on a bounded smooth domain.) Here $\varepsilon_0, \varepsilon_1 > 0$ are certain universal constants (sufficiently small). We note that the proof of the above theorem does not actually use the fact that u is a suitable weak solution, but merely a weak solution to the NSI (which is not the case, however, in the subsequent alternative proofs due to Lin [12], Ladyzhenskaya and Seregin [10] and Vasseur [25] mentioned above).

The partial regularity theorem (Theorem 1.2) is a key ingredient in the $L_{3,\infty}$ regularity criterion for the three-dimensional Navier–Stokes equations (see [6]) and the uniqueness of Lagrangian trajectories for suitable weak solutions [17]; similar ideas have also been used for other models, such as the surface growth model $\partial_t u + u_{xxx} + \partial_{xx} u_x^2 = 0$ [15], which can serve as a “one-dimensional model” of the Navier–Stokes equations [1, 2].

¹ Note that here we use the convention of “nonanticipating” cylinders; namely that Q is based at a point (x, t) when (x, t) lies on the upper lid of the cylinder.

A key fact about the partial regularity theory is that the quantities involved in the local regularity criteria (that is $|u|^3$, $|p|^{3/2}$ and $|\nabla u|^2$), are known to be globally integrable for any vector field satisfying $\sup_{t>0} \|u(t)\| < \infty$, $\nabla u \in L^2(\mathbb{R}^3 \times (0, \infty))$ (which follows by interpolation, see for example, Lemma 3.5 and inequality (5.7) in Robinson et al. [16]); thus in particular for any Leray–Hopf weak solution (by (1.2)). Therefore Theorem 1.2 shows that, in a sense, if these quantities localise near a given point $z \in \mathbb{R}^3 \times (0, \infty)$ in a way that is “not too bad”, then z is not a singular point, and thus there cannot be “too many” singular points. In fact, by letting $S \subset \mathbb{R}^3 \times (0, \infty)$ denote the singular set, that is

$$S := \{(x, t) \in \mathbb{R}^3 \times (0, \infty) : u \text{ is unbounded in any neighbourhood of } (x, t)\}, \quad (1.8)$$

this can be made precise by estimating the “dimension” of S . Namely, a simple consequence of (1.6) and (1.7) is that

$$d_B(S) \leq 5/3, \quad \text{and} \quad d_H(S) \leq 1, \quad (1.9)$$

respectively,² see Theorem 15.8 and Theorem 16.2 in Robinson et al. [16]. Here d_B denotes the *box-counting dimension* (also called the *fractal dimension* or the *Minkowski dimension*) and d_H denotes the *Hausdorff dimension*. The relevant definitions can be found in Falconer [7], who also proves (in Proposition 3.4) the important property that $d_H(K) \leq d_B(K)$ for any compact set K .

Very recently, Buckmaster and Vicol [3] proved nonuniqueness of weak solutions to the Navier–Stokes equations on the torus \mathbb{T}^3 (rather than on \mathbb{R}^3). Their solutions belong to the class $C([0, T]; L^2(\mathbb{T}^3))$, but they do not belong to the class $L^2((0, T); H^1(\mathbb{T}^3))$. Thus in particular these do not satisfy the energy inequality (1.2), and so they are neither Leray–Hopf weak solutions nor weak solutions of the NSI. Moreover, the constructions of Buckmaster and Vicol [3] include weak solutions with increasing energy $\|u(t)\|$.

In this article we work towards the same goal as Buckmaster and Vicol [3], but from a different direction. Given an open set $W \subset \mathbb{R}^3$ and a nonincreasing energy profile $e: [0, T] \rightarrow [0, \infty)$ we construct a weak solution to the NSI such that its energy stays arbitrarily close to e and its support is contained in W for all times. Namely we prove the following theorem.

Theorem 1.3 (Weak solutions to the NSI with arbitrary energy profile). *Given an open set $W \subset \mathbb{R}^3$, $\varepsilon > 0$, $T > 0$ and a nonincreasing function $e: [0, T] \rightarrow [0, \infty)$ there exist $\nu_0 > 0$ and a weak solution u of the NSI for all $\nu \in [0, \nu_0]$ such that $\text{supp } u(t) \subset W$ for all $t \in [0, T]$ and*

$$\| \|u(t)\| - e(t) \| \leq \varepsilon \quad \text{for all } t \in [0, T]. \quad (1.10)$$

We point out that the vector field u given by the above theorem satisfies the NSI for all values of viscosity $\nu \in [0, \nu_0]$. However, we emphasize that it does not satisfy the Navier–Stokes equations (but merely the NSI).

Our approach is inspired by some ideas of Scheffer [23, 24], who showed that the bound $d_H(S) \leq 1$ is sharp for weak solutions of the NSI (of course, it is not known whether it is sharp for suitable weak solutions of the NSE). His 1985 result is the following.

² In fact, (1.7) implies a stronger estimate than $d_H(S) \leq 1$; namely that $\mathcal{P}^1(S) = 0$, where $\mathcal{P}^1(S)$ is the *parabolic Hausdorff measure* of S (see Theorem 16.2 in Robinson et al. [16] for details).

Theorem 1.4 (Weak solution of NSI with point singularity). *There exist $v_0 > 0$ and a vector field $u: \mathbb{R}^3 \times [0, \infty) \rightarrow \mathbb{R}^3$ that is a weak solution of the Navier–Stokes inequality with any $v \in [0, v_0]$ such that $u(t) \in C^\infty$, $\text{supp } u(t) \subset G$ for all t for some compact set $G \Subset \mathbb{R}$ (independent of t). Moreover u is unbounded in every neighbourhood of (x_0, T_0) , for some $x_0 \in \mathbb{R}^3$, $T_0 > 0$.*

It is clear, using an appropriate rescaling, that the statement of the above theorem is equivalent to the one where $v = 1$ and $(x_0, T_0) = (0, 1)$. Indeed, if u is the velocity field given by the theorem then $\sqrt{T_0/v_0}u(x_0 + \sqrt{T_0/v_0}x, T_0t)$ satisfies Theorem 1.4 with $v_0 = 1$, $(x_0, T_0) = (0, 1)$.

In a subsequent paper Scheffer [24] constructed weak solutions of the Navier–Stokes inequality that blow up on a Cantor set $S \times \{T_0\}$ with $d_H(S) \geq \xi$ for any preassigned $\xi \in (0, 1)$.

Theorem 1.5 (Nearly one-dimensional singular set). *Given $\xi \in (0, 1)$ there exists $v_0 > 0$, a compact set $G \Subset \mathbb{R}^3$ and a function $u: \mathbb{R}^3 \times [0, \infty) \rightarrow \mathbb{R}^3$ that is a weak solution to the Navier–Stokes inequality such that $u(t) \in C^\infty$, $\text{supp } u(t) \subset G$ for all t , and*

$$\xi \leq d_H(S) \leq 1,$$

where S is the singular set (recall (1.8)).

The author’s previous work, [13] provides a simpler presentation of Scheffer’s constructions of u from Theorems 1.4 and 1.5 and provides a new light on these constructions. In particular he introduces the concepts of a *structure* (which we exploit in this article, see below), the *pressure interaction function* and the *geometric arrangement*, which articulate the main tools used by Scheffer to obtain a blow-up, but also describe, in a sense, the geometry of the NSI and expose a number of degrees of freedom available in constructing weak solutions to the NSI. Furthermore, it is shown in [13] how one can obtain a blow-up on a Cantor set (Theorem 1.5) by a straightforward generalisation of the blow-up at a single point (Theorem 1.4).

It turns out that the construction from Theorem 1.3 can be combined with Scheffer’s constructions to yield a weak solution to the Navier–Stokes inequality with both the blow-up and the prescribed energy profile.

Theorem 1.6 (Weak solutions to the NSI with blow-up and arbitrary energy profile). *Given an open set $W \subset \mathbb{R}^3$, $\varepsilon > 0$, $T > 0$ and a nonincreasing function $e: [0, T] \rightarrow [0, \infty)$ such that $e(t) \rightarrow 0$ as $t \rightarrow T$ there exists $v_0 > 0$ and a weak solution u of the NSI for all $v \in [0, v_0]$ such that $\text{supp } u(t) \subset W$ and*

$$\| |u(t)| - e(t) \| \leq \varepsilon \quad \text{for all } t \in [0, T],$$

and the singular set S of u is of the form

$$S = S' \times \{T\},$$

where $S' \subset \mathbb{R}^3$ is a Cantor set with $d_H(S') \in [\xi, 1]$ for any preassigned $\xi \in (0, 1)$.

The structure of the article is as follows. In Sect. 2 we introduce some preliminary ideas including the notion of a *structure* (v, f, ϕ) on an open subset U of the upper half-plane

$$\mathbb{R}_+^2 := \{(x_1, x_2) : x_2 > 0\}.$$

In Sect. 3 we briefly sketch how the concept of a structure is used in the constructions of Scheffer (but we will refer the reader to Ożański [13] for the full proof). We then illustrate some useful properties of structures of the form $(0, f, \phi)$ and we show how they can be used to generate weak solutions to the NSI on arbitrarily long time intervals. In Sect. 4 we prove our main result, Theorem 1.3, and in Sect. 5 we prove Theorem 1.6. In the final section (Sect. 6) we prove Lemma 4.2, which is an important ingredient of the proof of Theorem 1.3.

2. Preliminaries

We will denote the $L^2(\mathbb{R}^3)$ norm by $\|\cdot\|$. We denote the space of indefinitely differentiable functions with compact support in a set U by $C_0^\infty(U)$. We denote the indicator function of a set U by χ_U . We frequently use the convention

$$h_t(\cdot) \equiv h(\cdot, t),$$

that is the subscript t denotes dependence on t (rather than the t -derivative, which we denote by ∂_t).

We say that a vector field $u: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is *axisymmetric* if $u(R_\theta x) = R_\theta(u(x))$ for any $\theta \in [0, 2\pi)$, $x \in \mathbb{R}^3$, where

$$R_\theta(x_1, x_2, x_3) := (x_1, x_2 \cos \phi - x_3 \sin \phi, x_2 \sin \phi + x_3 \cos \phi)$$

is the rotation operation around the x_1 axis. We say that a scalar function $q: \mathbb{R}^3 \rightarrow \mathbb{R}$ is *axisymmetric* if

$$q(R_\theta x) = q(x) \quad \text{for } \phi \in [0, 2\pi), x \in \mathbb{R}^3.$$

Observe that if a vector field $u \in C^2$ and a scalar function $q \in C^1$ are axisymmetric then the vector function $(u \cdot \nabla)u$ and the scalar functions

$$|u|^2, \quad \operatorname{div} u, \quad u \cdot \nabla |u|^2, \quad u \cdot \nabla q, \quad u \cdot \Delta u \quad \text{and} \quad \sum_{i,j=1}^3 \partial_i u_j \partial_j u_i \quad (2.1)$$

are axisymmetric, see Section 3.6.2 in Ożański [13] for details.

Let $U \in \mathbb{R}_+^2$. Set

$$R(U) := \{x \in \mathbb{R}^3 : x = R_\phi(y, 0) \text{ for some } \phi \in [0, 2\pi), y \in U\}, \quad (2.2)$$

the *rotation of U* .

Given $v = (v_1, v_2) \in C_0^\infty(U; \mathbb{R}^2)$ and $f: \mathbb{R}^2 \rightarrow [0, \infty)$ supported in \bar{U} and such that $f > |v|$ we define $u[v, f]: R(\bar{U}) \rightarrow \mathbb{R}^3$ to be the axisymmetric vector field such that

$$u[v, f](x_1, x_2, 0) := \left(v_1(x_1, x_2), v_2(x_1, x_2), \sqrt{f(x_1, x_2)^2 - |v(x_1, x_2)|^2} \right)$$

for $x_2 > 0$. In other words

$$u[v, f](x_1, \rho, \phi) = v_1(x_1, \rho) \widehat{x}_1 + v_2(x_1, \rho) \widehat{\rho} + \sqrt{f(x_1, \rho)^2 - |v(x_1, \rho)|^2} \widehat{\phi}, \quad (2.3)$$

where the cylindrical coordinates x_1, ρ, ϕ are defined using the representation

$$\begin{cases} x_1 = x_1, \\ x_2 = \rho \cos \phi, \\ x_3 = \rho \sin \phi \end{cases}$$

and the cylindrical basis vectors $\widehat{x}_1, \widehat{\rho}, \widehat{\phi}$ are

$$\begin{cases} \widehat{x}_1(x_1, \rho, \phi) := (1, 0, 0), \\ \widehat{\rho}(x_1, \rho, \phi) := (0, \cos \phi, \sin \phi), \\ \widehat{\phi}(x_1, \rho, \phi) := (0, -\sin \phi, \cos \phi). \end{cases} \quad (2.4)$$

Note that such a definition immediately gives

$$|u[v, f]| = f.$$

Moreover, it satisfies some other useful properties, which we state in a lemma.

Lemma 2.1 (Properties of $u[v, f]$).

(i) *The vector field $u[v, f]$ is divergence free if and only if v satisfies*

$$\operatorname{div}(x_2 v(x_1, x_2)) = 0 \quad \text{for all } (x_1, x_2) \in \mathbb{R}_+^2.$$

(ii) *If $v \equiv 0$ then*

$$\Delta u[0, f](x_1, \rho, \phi) = Lf(x_1, \rho)\widehat{\phi},$$

where

$$Lf(x_1, x_2) := \Delta f(x_1, x_2) + \frac{1}{x_2} \partial_{x_2} f(x_1, x_2) - \frac{1}{x_2^2} f(x_1, x_2). \quad (2.5)$$

In particular

$$\Delta u[0, f](x_1, x_2, 0) = (0, 0, Lf(x_1, x_2)). \quad (2.6)$$

(iii) *For all $x_1, x_2 \in \mathbb{R}$*

$$\partial_{x_3} |u[v, f]|(x_1, x_2, 0) = 0. \quad (2.7)$$

Proof. These are easy consequences of the definition (and the properties of cylindrical coordinates), see Lemma 3.2 in Ożański [13] for details. \square

Using part (ii) we can see that the term $u[0, f] \cdot \Delta u[0, f]$ (recall the Navier–Stokes inequality (1.3)), which is axisymmetric (recall (2.1)), can be made non-negative by ensuring that Lf is non-negative, since

$$u[0, f](x_1, x_2, 0) \cdot \Delta u[0, f](x_1, x_2, 0) = f(x_1, x_2) Lf(x_1, x_2) \quad (2.8)$$

and f is non-negative by definition. It is not clear how to construct f such that $Lf \geq 0$ everywhere, but there exists a generic way of constructing f which guarantees this property at points sufficiently close to the boundary of U if U is a rectangle. In order to state such a construction we denote (given $\eta > 0$) the “ η -subset” of U by U_η , that is

$$U_\eta := \{x \in U : \operatorname{dist}(x, \partial U) > \eta\}.$$

We have the following result.

Lemma 2.2 (The edge effects). *Let $U \in \mathbb{R}_+^2$ be an open rectangle, that is $U = (a_1, b_1) \times (a_2, b_2)$ for some $a_1, a_2, b_1, b_2 \in \mathbb{R}$ with $b_1 > a_1, b_2 > a_2 > 0$. Given $\eta > 0$ there exists $\delta \in (0, \eta)$ and $f \in C_0^\infty(\mathbb{R}_+^2; [0, 1])$ such that*

$$\begin{aligned} \text{supp } f &= \overline{U}, \quad f > 0 \text{ in } U \quad \text{with} \quad f = 1 \text{ on } U_\eta, \\ Lf &> 0 \quad \text{in } U \setminus U_\delta. \end{aligned}$$

Proof. See Lemma 3.15 in Ożański [13] for the proof (which is based on Section 5 in Scheffer [23]). \square

In other words, we can construct f that equals 1 on the given η -subset of U such that $Lf > 0$ outside of a sufficiently large δ -subset. We will later (in Lemma 4.2) refine this lemma to show that δ can be chosen proportional to η and that f is bounded away from 0 on U_δ .

We define $p^*[v, f]: \mathbb{R}^3 \rightarrow \mathbb{R}$ to be the pressure function corresponding to $u[v, f]$, that is

$$p^*[v, f](x) := \int_{\mathbb{R}^3} \sum_{i,j=1}^3 \frac{\partial_i u_j[v, f](y) \partial_j u_i[v, f](y)}{4\pi |x - y|} dy, \quad (2.9)$$

and we denote its restriction to \mathbb{R}^2 by $p[v, f]$,

$$p[v, f](x_1, x_2) := p^*[v, f](x_1, x_2, 0). \quad (2.10)$$

Since $u[v, f]$ is axisymmetric, the same is true of $p^*[v, f]$ (recall (2.1); see also (3.22) in Ożański [13] for a detailed verification of this fact). In particular

$$\partial_{x_3} p^*[v, f](x_1, x_2, 0) = 0 \quad \text{for all } x_1, x_2 \in \mathbb{R}, \quad (2.11)$$

as in Lemma 2.1 (iii) above.

2.1. A structure. We say that a triple (v, f, ϕ) is a *structure on* $U \in \mathbb{R}_+^2$ if $v \in C_0^\infty(U; \mathbb{R}^2)$, $f \in C_0^\infty(\mathbb{R}_+^2; [0, \infty))$, $\phi \in C_0^\infty(U; [0, 1])$ are such that $\text{supp } f = \overline{U}$,

$$\begin{aligned} \text{supp } v &\subset \{\phi = 1\}, \quad \text{div}(x_2 v(x_1, x_2)) = 0 \quad \text{in } U \\ \text{and } f &> |v| \quad \text{in } U \quad \text{with} \quad Lf > 0 \quad \text{in } U \setminus \{\phi = 1\}. \end{aligned}$$

Note that, given a structure (v, f, ϕ) , we obtain an axisymmetric divergence-free vector field $u[v, f]$ that is supported in $R(\overline{U})$ (which is, in particular, away from the x_1 axis), and such that

$$|u[v, f](x, 0)| = f(x) \quad \text{for } x \in \mathbb{R}_+^2.$$

Moreover we note that (av, f, ϕ) is a structure for any $a \in (-1, 1)$ whenever (v, f, ϕ) is, and that, given disjoint $U_1, U_2 \in \mathbb{R}_+^2$ and the corresponding structures (v_1, f_1, ϕ_1) , (v_2, f_2, ϕ_2) , the triple $(v_1 + v_2, f_1 + f_2, \phi_1 + \phi_2)$ is a structure on $U_1 \cup U_2$. Observe that the role of the cutoff function ϕ in the definition of a structure is to cut off the edge effects as well as “cut in” the support of v . Namely, in $R(\{\phi < 1\})$ (recall that R denotes the rotation, see (2.2)) we have $Lf \geq 0$ and $v = 0$, and so

$$u[v, f] \cdot \Delta u[v, f] \geq 0 \quad (2.12)$$

and

$$u[v, f] \cdot \nabla q = 0 \quad (2.13)$$

for any axisymmetric function $q: \mathbb{R}^3 \rightarrow \mathbb{R}$. This last property (which follows from (2.11)) is particularly useful when taking $q := |u[v, f]|^2 + 2p[v, f]$ as this gives one of the terms in the Navier–Stokes inequality (1.5).

2.2. A recipe for a structure. Using Lemma 2.2 one can construct structures on sets $U \Subset \mathbb{R}_+^2$ in the shape of a rectangle (which is the only shape we will consider in this article) in a generic way. This can be done using the following steps.

- First construct $w: U \rightarrow \mathbb{R}^2$ that is weakly divergence free (that is $\int_U w \cdot \nabla \psi = 0$ for every $\psi \in C_0^\infty(U)$) and compactly supported in U .
For example one can take $w := (x_2, x_1) \chi_{1 < |(x_1, x_2)| < 2}$, after an appropriate rescaling and translation (so that $\text{supp } w$ fits inside U); such a w is weakly divergence free due to the fact that $w \cdot n$ vanishes on the boundary of its support, where n denotes the respective normal vector to the boundary.
- Next, set $v := (J_\epsilon w)/x_2$, where J_ϵ denotes the standard mollification and $\epsilon > 0$ is small enough so that $\text{supp } v \Subset U$.
- Then construct f by using Lemma 2.2 (with any $\eta > 0$) and multiplying by a constant sufficiently large so that $f > |v|$ in U .
- Finally let $\phi \in C_0^\infty(U; [0, 1])$ be such that $\{\phi = 1\}$ contains U_δ (from Lemma 2.2) and $\text{supp } v$.

3. Applications of Structures

In this section we point out two important applications of the concept of a structure.

3.1. The construction of Scheffer. Here we show how the concept of a structure is used in the Scheffer construction, Theorem 1.4, which we will only use later in proving Theorem 1.6.

We show below how Theorem 1.4 can be proved in a straightforward way using the following theorem.

Theorem 3.1. *There exist a set $U \Subset \mathbb{R}_+^2$, a structure (v, f, ϕ) and $\mathcal{T} > 0$ with the following property: there exist smooth time-dependent extensions v_t, f_t ($t \in [0, \mathcal{T}]$) of v, f , respectively, such that $v_0 = v, f_0 = f, (v_t, f_t, \phi)$ is a structure on U for each $t \in [0, \mathcal{T}]$. Moreover, for some $v_0 > 0$ the vector field*

$$u(t) := u[v_t, f_t]$$

satisfies the NSI (1.3) in the classical sense for all $v \in [0, v_0]$ and $t \in [0, \mathcal{T}]$ as well as admits a large gain in magnitude of the form

$$|u(\tau x + z, \mathcal{T})| \geq \tau^{-1} |u(x, 0)|, \quad x \in \mathbb{R}^3, \quad (3.1)$$

for some $\tau \in (0, 1), z \in \mathbb{R}^3$.

Proof. See Section 3.3 in Ożański [13] (particularly Proposition 3.8 therein) for a detailed proof. \square

In fact, the set U (from the theorem above) is of the form $U = U_1 \cup U_2$ for some disjoint $U_1, U_2 \subseteq \mathbb{R}_+^2$ and $(v, f, \phi) = (v_1 + v_2, f_1 + f_2, \phi_1 + \phi_2)$, where $(v_1, f_1, \phi_1), (v_2, f_2, \phi_2)$ are some structures on U_1, U_2 , respectively. The elaborate part of the proof of Theorem 3.1 is devoted to the careful arrangement of U_1, U_2 and a construction of the corresponding structures and $\mathcal{T} > 0$ which magnifies certain interaction between U_1 and U_2 via the pressure function, and thus allows (3.1). We refer the reader to Sections 3.3 and 3.4 in Ożański [13] for the full proof of Theorem 3.1. We note, however, that the part of the theorem about the NSI is not that difficult. In fact we show in Lemma 3.3 below that any structure gives rise to infinitely many classical solutions of the NSI (on arbitrarily long time intervals) with $u[v, f]$ as the initial condition.

In order to prove Theorem 1.4 we will make use of an alternative form of the local energy inequality. Namely, the local energy inequality (1.4) is satisfied if the *local energy inequality on the time interval* $[S, S']$,

$$\begin{aligned} & \int_{\mathbb{R}^3} |u(x, S')|^2 \varphi \, dx - \int_{\mathbb{R}^3} |u(x, S)|^2 \varphi \, dx + 2\nu \int_S^{S'} \int_{\mathbb{R}^3} |\nabla u|^2 \varphi \\ & \leq \int_S^{S'} \int_{\mathbb{R}^3} (|u|^2 + 2p) u \cdot \nabla \varphi + \int_S^{S'} \int_{\mathbb{R}^3} |u|^2 (\partial_t \varphi + \nu \Delta \varphi), \end{aligned} \quad (3.2)$$

holds for all $S, S' > 0$ with $S < S'$, which is clear by taking S, S' such that $\text{supp } \varphi \subset \mathbb{R}^3 \times (S, S')$. An advantage of this alternative form of the local energy inequality is that it demonstrates how to combine weak solutions of the Navier–Stokes inequality one after another. Namely, (3.2) shows that a necessary and sufficient condition for two vector fields $u^{(1)}: \mathbb{R}^3 \times [t_0, t_1] \rightarrow \mathbb{R}^3$, $u^{(2)}: \mathbb{R}^3 \times [t_1, t_2] \rightarrow \mathbb{R}^3$ satisfying the local energy inequality on the time intervals $[t_0, t_1], [t_1, t_2]$, respectively, to combine (one after another) into a vector field satisfying the local energy inequality on the time interval $[t_0, t_2]$ is that

$$|u^{(2)}(x, t_1)| \leq |u^{(1)}(x, t_1)| \quad \text{for a.e. } x \in \mathbb{R}^3. \quad (3.3)$$

Using the above property and Theorem 3.1 we can employ a simple switching procedure to obtain Scheffer’s construction of the blow-up at a single point (i.e. the claim of Theorem 1.4). Namely, considering

$$u^{(1)}(x, t) := \tau^{-1} u(\Gamma^{-1}(x), \tau^{-2}(t - \mathcal{T})),$$

where $\Gamma(x) := \tau x + z$, we see that $u^{(1)}$ satisfies the Navier–Stokes inequality (1.3) in a classical sense for all $v \in [0, v_0]$ and $t \in [\mathcal{T}, (1 + \tau^2)\mathcal{T}]$, $\text{supp } u^{(1)}(t) = \Gamma(G)$ for all $t \in [\mathcal{T}, (1 + \tau^2)\mathcal{T}]$ and that (3.1) gives

$$\left| u^{(1)}(x, \mathcal{T}) \right| \leq |u(x, \mathcal{T})|, \quad x \in \mathbb{R}^3 \quad (3.4)$$

(and so $u, u^{(1)}$ can be combined “one after another”, recall (3.3)). Thus, since $u^{(1)}$ is larger in magnitude than u (by the factor of τ) and its time of existence is $[\mathcal{T}, (1 + \tau^2)\mathcal{T}]$, we see that by iterating such a switching we can obtain a vector field u that grows indefinitely in magnitude, while its support shrinks to a point (and thus will satisfy all the claims of Theorem 1.4), see Fig. 1. To be more precise we let $t_0 := 0$,

$$t_j := \mathcal{T} \sum_{k=0}^{j-1} \tau^{2k} \quad \text{for } j \geq 1, \quad (3.5)$$

$T_0 := \lim_{j \rightarrow \infty} t_j = T/(1 - \tau^2)$, $u^{(0)} := u$, and

$$u^{(j)}(x, t) := \tau^{-j} u \left(\Gamma^{-j}(x), \tau^{-2j}(t - t_j) \right), \quad j \geq 1, \quad (3.6)$$

see Fig. 1. As in (3.4), (3.1) gives that

$$\text{supp } u^{(j)}(t) = \Gamma^j(G) \quad \text{for } t \in [t_j, t_{j+1}] \quad (3.7)$$

and that the magnitude of the consecutive vector fields shrinks at every switching time, that is

$$\left| u^{(j)}(x, t_j) \right| \leq \left| u^{(j-1)}(x, t_j) \right|, \quad x \in \mathbb{R}^3, j \geq 1, \quad (3.8)$$

see Fig. 1.

Thus letting

$$u(t) := \begin{cases} u^{(j)}(t) & \text{if } t \in [t_j, t_{j+1}) \text{ for some } j \geq 0, \\ 0 & \text{if } t \geq T_0, \end{cases} \quad (3.9)$$

we obtain a vector field that satisfies all claims of Theorem 1.4 with $x_0 := z/(1 - \tau)$. Note that $u \in L^\infty((0, \infty); L^2(\mathbb{R}^3))$ and $\nabla u \in L^2(\mathbb{R}^3 \times (0, \infty))$ (which is required by the definition of weak solutions to the NSI, Definition 1.1) by construction (due to the rescaling (3.6) and the fact that $u^{(0)} = u$ is smooth on $R(\overline{U}) \times [0, T]$).

Observe that by construction

$$\|u(t)\|_p \rightarrow 0 \quad \text{as } t \rightarrow T_0^- \quad \text{for all } p \in [1, 3), \quad (3.10)$$

since $\tau \in (0, 1)$. Indeed we write for any $t \in [t_j, t_{j+1}]$, $j \geq 0$,

$$\begin{aligned} \|u(t)\|_p &= \|u^{(j)}(t)\|_p \leq \sup_{s \in [t_j, t_{j+1}]} \|u^{(j)}(s)\|_p \\ &= \tau^{-j(1-3/p)} \sup_{s \in [t_0, t_1]} \|u^{(0)}(s)\|_p \rightarrow 0 \quad \text{as } j \rightarrow \infty. \end{aligned}$$




Fig. 1. The switching procedure: the blow-up of $\|u(t)\|_\infty$ (left) and the shrinking support of $u(t)$ (right) as $t \rightarrow T_0^-$

3.2. *Structures of the Form* $(0, f, \phi)$. Let $U \Subset \mathbb{R}_+^2$. We now focus on the structures on U of the form $(0, f, \phi)$ and, for convenience, we set

$$u[f] := u[0, f].$$

Roughly speaking, $u[f]$ is a swirl-only axisymmetric vector field with (pointwise) magnitude f . Note that for all f_1, f_2 with $f_1 < f_2$

$$\left\| u \left[\sqrt{f_2^2 - f_1^2} \right] \right\|^2 = \|u[f_2]\|^2 - \|u[f_1]\|^2, \quad (3.11)$$

which is a useful property that we will use later (in (4.14) and (4.21)). As in (2.13) we see that

$$u[f] \cdot \nabla \left(|u[f]|^2 + 2p[f] \right) = 0 \quad \text{in } \mathbb{R}^3, \quad (3.12)$$

for any $f \in C_0^\infty(\mathbb{R}_+^2; [0, \infty))$. Using this property we can show that given any structure (v, f, ϕ) on a set $U \Subset \mathbb{R}_+^2$ there exists a time-dependent extension f_t of f such that $(0, f_t, \phi)$ is a structure on U and gives rise to a classical solution to the NSI (for all sufficiently small viscosities) that is almost constant in time. We make this precise in the following lemma, which we will use later.

Lemma 3.2. *Given $\varepsilon > 0, T > 0, U \Subset \mathbb{R}_+^2$ and a structure (v, f, ϕ) there exists $v_0 > 0$ and an axisymmetric classical solution u to the NSI for all $v \in [0, v_0], t \in [0, T]$ that is supported in $R(\bar{U})$ with $u(0) = u[f]$ and*

$$\|u(t) - u[f]\|_q \leq \varepsilon \quad \text{for all } t \in [0, T], q \in [1, \infty]. \quad (3.13)$$

Proof. Let

$$u(t) := u[f_t],$$

where

$$f_t^2 := f^2 - \delta t \phi$$

and $\delta > 0$ is sufficiently small such that $f_t > 0$ in U for all $t \in [0, T]$ (Note this is possible since f is continuous and $\text{supp } \phi \Subset \text{supp } f$). Clearly $u(0) = u[f]$ and (3.13) follows for $q \in \{1, \infty\}$ by taking δ sufficiently small. If $q \in (1, \infty)$ then (3.13) follows using Lebesgue interpolation.

It remains to verify that $u(t)$ satisfies the NSI. To this end let $v_0 > 0$ be sufficiently small such that

$$v_0 |u[f_t](x) \cdot \Delta u[f_t](x)| \leq \frac{\delta}{2} \quad \text{for } x \in \mathbb{R}^3, t \in [0, T]. \quad (3.14)$$

Due to the axisymmetry of u it is enough to verify the NSI only for points of the form $(x, 0, t)$, for $x \in \bar{U}, t \in [0, T]$. Setting p to be the pressure function corresponding to u (that is $p(t) := p^*[0, f_t]$) we use (3.12) to write

$$\begin{aligned} \partial_t |u(x, 0, t)|^2 &= -\delta \phi(x) \\ &= -\delta \phi(x) - u(x, 0, t) \cdot \nabla \left(|u(x, 0, t)|^2 + 2p(x, 0, t) \right) \\ &\leq 2v u(x, 0, t) \cdot \Delta u(x, 0, t) - u(x, 0, t) \cdot \nabla \left(|u(x, 0, t)|^2 + 2p(x, 0, t) \right), \end{aligned} \quad (3.15)$$

as required, where, in the last step, we used (2.12) for x such that $\phi(x) < 1$ and (3.14) for x such that $\phi(x) = 1$. \square

Observe that the lemma does not make any use of v . One similarly obtains the same result, but with the claim on the initial condition $u(0) = u[f]$ replaced by a condition at a final time, namely by the pointwise inequality $|u(T)| \geq |u[f]|$ everywhere in \mathbb{R}^3 . We thus obtain the following lemma, which we will use to prove Theorem 1.6.

Lemma 3.3. *Given $\varepsilon > 0$, $T > 0$, $U \Subset P$ and a structure (v, f, ϕ) there exists $\nu_0 > 0$ and an axisymmetric classical solution u to the NSI for all $\nu \in [0, \nu_0]$ that is supported in $R(\bar{U})$,*

$$|u(x, T)| \geq |u[f](x)| \quad \text{for all } x \in \mathbb{R}^3 \quad (3.16)$$

and

$$\|u(t) - u[f]\|_p \leq \varepsilon \quad \text{for all } t \in [0, T], p \in [1, \infty]. \quad (3.17)$$

Proof. The lemma follows in the same way as Lemma 3.2 after replacing “ f ” in the above proof by “ $(1 + \varepsilon)f$ ” for sufficiently small $\varepsilon > 0$ and then taking $\delta > 0$ (and so also ν_0) smaller. \square

Finally, observe that if $f_{1,t}, f_{2,t} \in C_0^\infty(\mathbb{R}_+^2; [0, \infty))$ are disjointly supported (for each t) then

$$p^*[0, f_{1,t} + f_{2,t}] = p^*[0, f_{1,t}] + p^*[0, f_{2,t}]$$

and so

$$u[f_{1,t} + f_{2,t}] \text{ satisfies the NSI in the classical sense} \quad (3.18)$$

whenever each of $u[f_{1,t}]$ and $u[f_{2,t}]$ does. Indeed, this is because the term

$$u(x_1, x_2, 0) \cdot \nabla p(x_1, x_2, 0) = u_3(x_1, x_2, 0) \partial_3 p(x_1, x_2, 0) \quad (3.19)$$

in the NSI vanishes (due to (2.11)). Note that (3.18) does not necessarily hold for structures (v, f, ϕ) with $v \neq 0$, as in this case the term $u \cdot \nabla p$ does not simplify as in (3.19). We will use (3.18) as a substitute for linearity of the NSI in the proof of Theorem 1.6 in Sect. 5.

4. Proof of Theorem 1.3

In this section we prove Theorem 1.3; namely given an open set $W \subset \mathbb{R}^3$, $\varepsilon > 0$, $T > 0$ and a continuous, nonincreasing function $e: [0, T] \rightarrow [0, \infty)$ there exist $\nu_0 > 0$ and a weak solution u of the NSI for all $\nu \in [0, \nu_0]$ such that $\text{supp } u(t) \subset W$ for all $t \in [0, T]$ and

$$\| \|u(t)\| - e(t) \| \leq \varepsilon \quad \text{for all } t \in [0, T]. \quad (4.1)$$

(Recall that $\| \cdot \|$ denotes the $L^2(\mathbb{R}^3)$ norm.)

We will assume that $e(t)$ is continuous. If $e(t)$ is discontinuous then one can easily incorporate the times at which $e(t)$ has jumps into the switching procedure. This will become clear from the proof, and we give a more detailed explanation in Sect. 4.1 below.

We can assume that $e(T) = 0$, as otherwise one could extend e continuously beyond T into a function decaying to 0 in finite time $T' > T$. Moreover, by translation in space we can assume that W intersects the x_1 axis. Let $U \Subset \mathbb{R}_+^2$ be such that $R(\bar{U}) \subset W$.

We will construct an axisymmetric weak solution to the NSI (for all sufficiently small viscosities) such that $u(t) \in C_0^\infty(\mathbb{R}^3)$, $\text{supp } u(t) \subset R(\bar{U})$ and

$$|\|u(t)\| - e(t)| \leq \varepsilon$$

for all $t \in [0, T]$.

Before the proof we comment on its strategy in an informal manner. Suppose for the moment that we would like to use a similar approach as in the proof of Lemma 3.2, that is define some rectangle $U \Subset \mathbb{R}_+^2$, a structure (v, f, ϕ) on it and $u(t) := u[f_t]$, where

$$f_t^2 := f^2 - (C - De(t)^2)\phi, \quad (4.2)$$

for some constants $C, D > 0$, such that

$$\|u(t)\| \approx e(t)$$

at least for small t . In fact we could use the recipe from Sect. 2.2 to construct (v, f, ϕ) . In order to proceed with the calculation (that is to guarantee the NSI) we would need to guarantee that $(e(t)^2)'$ is bounded above by some negative constant (so that the term with the Laplacian could be absorbed for x such that $\phi(x) = 1$; recall the last step of (3.15)), which is not a problem, as the following lemma demonstrates.

Lemma 4.1. *Given $\varepsilon > 0$ and a continuous and nonincreasing function $e: [0, T] \rightarrow [0, \infty)$ there exist $\zeta > 0$ and $\tilde{e}: [0, T] \rightarrow [0, \infty)$ such that $\tilde{e} \in C^\infty([0, T])$, and*

$$e(t) \leq \tilde{e}(t) \leq e(t) + \varepsilon, \quad \frac{d}{dt} \tilde{e}(t)^2 \leq -\zeta \quad \text{for } t \in [0, T].$$

Proof. Extend $e(t)$ by $e(T)$ for $t > T$ and by $e(0)$ for $t < 0$. Let $J_\delta e^2$ denote a mollification of e^2 . Since e^2 is uniformly continuous $J_\delta e^2$ converges to e^2 in the supremum norm as $\delta \rightarrow 0$, and so $\|J_\delta e^2 - e^2\|_{L^\infty(\mathbb{R})} < \varepsilon/4$ for sufficiently small δ . Then the function

$$\tilde{e}(t) := \sqrt{J_\delta e^2(t) + (\varepsilon/2 - \varepsilon t/4T)}$$

satisfies the claim of the lemma with $\zeta := \varepsilon/4T$. □

The problem with (4.2) is that its right-hand side can become negative for small times³ (so that $(0, f_t, \phi)$ would no longer be a structure, and so $u[f_t]$ would not be well-defined). We will overcome this problem by utilising the property (3.3). Namely, at time t_1 when the right-hand side of (4.2) becomes zero we will “trim” U to obtain a smaller set U^1 , on which the right-hand side of (4.2) does not vanish, and we will define a new structure $(0, f_1, \phi_1)$, with $f_1^2 \leq f^2 + (C - De(t_1)^2)\phi$. We will then continue the same way (as in (4.2)) to define $u(t) := u[f_{1,t}]$ for $t \geq t_1$ where

$$f_{1,t}^2 := f^2 - (C_1 - D_1 e(t)^2)\phi_1$$

for appropriately chosen C_1, D_1 . Note that such a continuation satisfies the local energy inequality, since (3.3) is satisfied. We will then continue in the same way to define

³ Note that the point $x \in U$ at which the right-hand side of (4.2) will become negative is located close to the ∂U since only for such x $\phi(x) = 1$ but $f(x) < \max f$.

U^2, U^3, \dots , structures $(0, f_2, \phi_2), (0, f_3, \phi_3), \dots$, and $u(t) := u[f_{k,t}]$ for $t \in [t_k, t_{k+1}]$, where

$$f_{k,t}^2 := f_k^2 - (C_k - D_k e(t)^2) \phi_k, \quad (4.3)$$

and $C_k, D_k > 0$ are chosen appropriately, until we reach time $t = T$.

Such a procedure might look innocent, but note that there is a potentially fatal flaw. Namely, we need to use an existence result such as Lemma 2.2 in order to construct f_k as well as $\delta_k > 0$; recall that δ_k controls the edge effect (that is $Lf_k \geq 0$ in $U^k \setminus U_{\delta_k}^k$) and that, according to the recipe from Sect. 2.2, ϕ_k is chosen so that $\phi_k = 1$ on $U_{\delta_k}^k$. However, Lemma 2.2 gives no control of δ_k , and so it seems possible that δ_k shrinks rapidly as k increases, and consequently

$$\inf_{U_{\delta_k}^k} f_k \rightarrow 0 \text{ rapidly as } k \text{ increases.}$$

Thus (since $\phi_k = 1$ on $U_{\delta_k}^k$) the length of the time interval $[t_k, t_{k+1}]$ would shrink rapidly to 0 as k increases (as the right-hand side of (4.3) would become negative for some x), and so it is not clear whether the union of all intervals,

$$\bigcup_{k \geq 0} [t_k, t_{k+1}],$$

would cover $[0, T]$.

In order to overcome this problem we prove a sharper version of Lemma 2.2 which states that we can choose $\delta = c'\eta$ and f such that $f > c$ in U_δ , where the constants $c, c' \in (0, 1)$ do not depend on the size of U .

Lemma 4.2 (The cut-off function with the edge effect on a rectangle). *Let $a > 0$ and $U \Subset \mathbb{R}_+^2$ be an open rectangle that is located at least a away from the x_1 axis, that is $U = (a_1, b_1) \times (a_2, b_2)$ for some $a_1, a_2, b_1, b_2 \in \mathbb{R}$ with $b_1 > a_1, b_2 > a_2 > a$. Given $\eta \in (0, \min\{1, (b_1 - a_1)/2, (b_2 - a_2)/2\})$ there exists $f \in C_0^\infty(\mathbb{R}_+^2; [0, 1])$ such that*

$$\begin{aligned} \text{supp } f &= \bar{U}, \quad f > 0 \text{ in } U \text{ with } f = 1 \text{ on } U_\eta, \\ Lf &> 0 \quad \text{in } U \setminus U_{c'\eta}, \text{ with } f > c \text{ in } U_{c'\eta/2}, \end{aligned}$$

where $c, c' \in (0, 1/2)$ depend only on a .

Proof. We prove the lemma in Appendix 6. □

The above lemma allows us to ensure that the time interval $[0, T]$ can be covered by only finitely many intervals $[t_k, t_{k+1}]$.

We now make the above strategy rigorous.

Proof of Theorem 1.3. (Recall that we also assume that $e(T) = 0$ and that $U \subset \mathbb{R}_+^2$ is such that $R(\bar{U}) \subset W$.) Fix $a > 0$ such that $\text{dist}(U, x_1\text{-axis}) \geq a$. By applying Lemma 4.1 we can assume that e^2 is differentiable on $[0, T]$ with $(e^2(t))' \leq -\zeta$ for all $t \in [0, T]$, where $\zeta > 0$. Let K be the smallest positive integer such that

$$(1 - c^2)^K e(0)^2 < \varepsilon^2,$$

where $c = c(a) \in (0, 1/2)$ is the constant from Lemma 4.2. For $k \in \{1, \dots, K\}$ let $t_k \in [0, T]$ be such that

$$e(t_k)^2 = (1 - c^2)^k e(0)^2. \quad (4.4)$$

(Note t_k is uniquely determined since $e(t)^2$ is strictly decreasing, $(e(t)^2)' \leq -\zeta$.) Let also $t_0 := 0$. Observe that the choice of K implies that

$$e(t)^2 \geq \varepsilon^2/2 \quad \text{for } t \in [t_0, t_K]. \quad (4.5)$$

Indeed, since $e(t)$ is decreasing and $c^2 < 1/2$,

$$e(t)^2 \geq e(t_K)^2 = (1 - c^2)(1 - c^2)^{K-1} e(0)^2 \geq (1 - c^2)\varepsilon^2 \geq \varepsilon^2/2,$$

as required, where we used the definition of K in the second inequality.

We set

$$d := \min_{k \in \{0, \dots, K-1\}} (t_{k+1} - t_k).$$

Given $k \in \{0, \dots, K-1\}$ we will construct a classical solution u_k to the NSI for all $v \in [0, v_0]$ (where v_0 is fixed in (4.19) below) on time interval $[t_k, t_{k+1}]$ (respectively) such that

$$\left| \|u_k(t)\|^2 - e(t)^2 \right| \leq \varepsilon^2/4 \quad \text{for } t \in [t_k, t_{k+1}], \quad (4.6)$$

and that

$$|u_{k+1}(t_{k+1})| \leq |u_k(t_{k+1})| \quad \text{a.e. in } \mathbb{R}^3 \quad \text{for } k = 0, \dots, K-2 \quad (4.7)$$

and

Then the claim of the theorem follows by defining

$$u(t) := \begin{cases} u_k(t) & t \in [t_k, t_{k+1}), k \in \{0, \dots, K-1\}, \\ 0 & t \geq t_K. \end{cases}$$

Indeed, (4.7) implies that we can switch from u_k to u_{k+1} at time t_{k+1} ($k = 0, \dots, K-2$), so that u is a weak solution of the NSI for all $v \in [0, v_0]$, $t \in [0, T]$. Moreover (4.6) implies (4.1), since

$$\left| \|u(t)\| - e(t) \right| = \left| \|u(t)\|^2 - e(t)^2 \right| / \left(\|u(t)\| + e(t) \right) \leq \varepsilon^2/4e(t) \leq \varepsilon \quad \text{for } t \in [t_0, t_K], \quad (4.8)$$

where we used (4.5) in the last inequality, and the claim for $t \in [t_K, T]$ follows trivially.

In order to construct u_k (for $k = 0, \dots, K-1$) we first fix $\mu > 0$ such that

$$\mu \|u[\chi_U]\| = e(0) \quad (4.9)$$

and we set $\eta > 0$ sufficiently small such that

$$\|u[\chi_{U \setminus U_{K\eta}}]\|^2 < \frac{\min\{\varepsilon^2, d\zeta\}}{8\mu^2}. \quad (4.10)$$

Note that (4.4) and (4.9) give

$$e(t_k) = (1 - c^2)^k \mu^2 \|u[\chi_U]\|^2. \quad (4.11)$$

We now let $U^k := U_{k\eta}$ and apply Lemma 4.2 to obtain $c, c' \in (0, 1/2)$ and $f_k \in C_0^\infty(P; [0, 1])$ ($k = 0, \dots, K - 1$) such that

$$\begin{aligned} \text{supp } f_k &= \overline{U^k}, \quad f_k > 0 \text{ in } U^k \text{ with } f_k = 1 \text{ on } \overline{U_\eta^k} = \overline{U^{k+1}}, \\ Lf_k &> 0 \quad \text{in } U^k \setminus U_{c'\eta}^k, \text{ with } f_k > c \text{ in } U_{c'\eta}^k. \end{aligned} \quad (4.12)$$

Recall that c, c' are independent of k . Let $\phi_k \in C_0^\infty(U^k; [0, 1])$ be such that

$$\text{supp } \phi_k \subset U_{c'\eta/2}^k \quad \text{and} \quad \phi_k = 1 \text{ on } U_{c'\eta}^k. \quad (4.13)$$

Note that (4.10) implies that

$$\left| \|u[f_k]\|^2 - \|u[\phi_k]\|^2 \right|, \left| \|u[\chi_U]\|^2 - \|u[\phi_k]\|^2 \right| \leq \frac{\min\{\varepsilon^2, d\zeta\}}{8\mu^2} \quad (4.14)$$

for all $k = 0, \dots, K - 1$. Indeed, as for the first of these quantities (the second one is analogous), note that since $\chi_{U_{k\eta}} \leq f_k, \phi_k \leq \chi_U$ we have $|f_k^2 - \phi_k^2| \leq \chi_{U \setminus U_{k\eta}}$. Thus

$$\left| |u[f_k]|^2 - |u[\phi_k]|^2 \right| = \left| u \left[\sqrt{|f_k^2 - \phi_k^2|} \right] \right|^2 \leq |u[\chi_{U \setminus U_{k\eta}}]|^2$$

and (4.14) follows by integrating over \mathbb{R}^3 and using (4.10).

We will consider an affine modification $E_k(t)^2$ of $e(t)^2$ on the time interval $[t_k, t_{k+1}]$ such that

$$E_k(t_k)^2 = (1 - c^2)^k \mu^2 \|u[\phi_k]\|^2 \quad \text{and} \quad E_k(t_{k+1})^2 = (1 - c^2) E_k(t_k)^2. \quad (4.15)$$

(Recall $e(t)$ satisfies the above conditions with $\|u[\phi_k]\|$ replaced by $\|u[\chi_U]\|$, see (4.11).) Namely we set

$$E_k(t)^2 := e(t)^2 - (1 - c^2)^k \mu^2 \left(\|u[\chi_U]\|^2 - \|u[\phi_k]\|^2 \right) \left(1 - c^2 \frac{t - t_k}{t_{k+1} - t_k} \right).$$

Roughly speaking, E_k is a convenient modification of e that allows us to satisfy (4.7) while not causing any trouble to either (4.6) or the NSI. For example, we see that

$$\begin{aligned} \left| E_k(t)^2 - e(t)^2 \right| &= (1 - c^2)^k \mu^2 \left(\|u[\chi_U]\|^2 - \|u[\phi_k]\|^2 \right) \left(1 - c^2 \frac{t - t_k}{t_{k+1} - t_k} \right) \\ &\leq (1 - c^2)^k \varepsilon^2 / 8 \\ &\leq \varepsilon^2 / 8 \end{aligned} \quad (4.16) \quad \text{for } t \in [t_k, t_{k+1}]$$

where we used (4.14). This implies in particular that $E_k(t)$ is well-defined (as $e(t)^2 \geq \varepsilon^2/2$, recall (4.5)). Moreover, using (4.14) again

$$\begin{aligned} (E_k(t)^2)' &= (e(t)^2)' + (1 - c^2)^k \mu^2 \left(\|u[\chi_U]\|^2 - \|u[\phi_k]\|^2 \right) \frac{c^2}{t_{k+1} - t_k} \\ &\leq -\zeta + (1 - c^2)^k c^2 d\zeta / 8(t_{k+1} - t_k) \\ &\leq -\zeta + \zeta / 8 \\ &< -\zeta / 2 \end{aligned} \quad (4.17) \quad \text{for } t \in [t_k, t_{k+1}].$$

We can now define u_k by writing

$$u_k(t) := u[f_{k,t}],$$

where

$$f_{k,t}^2 := (1 - c^2)^k \mu^2 f_k^2 - \left((1 - c^2)^k \mu^2 - \frac{E_k(t)^2}{\|u[\phi_k]\|^2} \right) \phi_k^2.$$

Observe that, due to the monotonicity of E_k (shown above) and (4.15), the last term above can be bounded above and below

$$0 \leq \left((1 - c^2)^k \mu^2 - \frac{E_k(t)^2}{\|u[\phi_k]\|^2} \right) \phi_k^2 \leq c^2 (1 - c^2)^k \mu^2 \phi_k^2 \quad (4.18)$$

for all $t \in [t_k, t_{k+1}]$. (This is the solution to the problem we discussed informally before the proof.)

This means, in particular, that $f_{k,t}^2$ is nonnegative in U^k (that is $f_{k,t}$ is well-defined by the above formula). Indeed, this is trivial for $x \in U^k \setminus U_{c'\eta/2}^k$ (as $\phi_k(x) = 0$ for such x), and for $x \in U_{c'\eta/2}^k$ we have $f_k^2(x) > c^2$ (recall Lemma 4.2) and so

$$f_{k,t}^2(x) > (1 - c^2)^k \mu^2 c^2 (1 - \phi_k) \geq 0,$$

as required.

Let $\nu_0 > 0$ be sufficiently small such that

$$\nu_0 \|u[f_{k,t}] \cdot \Delta u[f_{k,t}]\|_\infty \leq \frac{\zeta}{4\|u[\chi_U]\|^2} \quad \text{for } t \in [t_k, t_{k+1}], k = 0, \dots, K - 1. \quad (4.19)$$

Having fixed ν_0 we show that u_k is a classical solution of the NSI with any $\nu \in [0, \nu_0]$ on the time interval $[t_k, t_{k+1}]$. Namely for each such ν we can use the monotonicity of $E_k(t)^2$ (recall (4.17)) to obtain

$$\begin{aligned} \partial_t |u_k(x, 0, t)|^2 &= \partial_t E_k(t)^2 \frac{\phi_k(x)}{\|u[\phi_k]\|^2} \\ &\leq -\zeta \frac{\phi_k(x)}{2\|u[\chi_U]\|^2} \\ &= -\zeta \frac{\phi_k(x)}{2\|u[\chi_U]\|^2} - u_k(x, 0, t) \cdot \nabla \left(|u_k(x, 0, t)|^2 + 2p_k(x, 0, t) \right) \\ &\leq 2\nu u_k(x, 0, t) \cdot \Delta u_k(x, 0, t) - u_k(x, 0, t) \cdot \nabla \left(|u_k(x, 0, t)|^2 + 2p_k(x, 0, t) \right), \end{aligned} \quad (4.20)$$

as required, where we used (3.12) in the third step and, in the last step, we used (2.12) for x such that $\phi_k(x) < 1$ and (4.19) for x such that $\phi_k(x) = 1$.

It remains to verify (4.6) and (4.7). As for (4.6) we use observation (3.11) to write

$$\begin{aligned} \|u_k(t)\|^2 &= \|u[f_{k,t}]\|^2 = (1 - c^2)^k \mu^2 \|u[f_k]\|^2 \\ &\quad - \left((1 - c^2)^k \mu^2 \|u[\phi_k]\|^2 - E_k(t)^2 \right). \end{aligned} \quad (4.21)$$

Thus

$$\left| \|u_k(t)\|^2 - E_k(t)^2 \right| = (1 - c^2)^k \mu^2 \left| \|u[f_k]\|^2 - \|u[\phi_k]\|^2 \right| \leq \varepsilon^2/8,$$

where we used (4.14). This and (4.16) give (4.6), as required.

As for (4.7) it suffices to show the claim on $R(\overline{U(\mu_{k+1})\eta})$ (that is on the support of u_{k+1}). Moreover, since both u_k and u_{k+1} are axially symmetric (with the same axis of symmetry, the Ox_1 axis), it is enough to show the claim at the points of the form $(x, 0)$, where $x = (x_1, x_2) \in \overline{U^{k+1}}$. Recalling (from (4.12), (4.13)) that for such x $f_k(x) = \phi_k(x) = 1 \geq f_{k+1}(x)$ we obtain

$$\begin{aligned} |u_{k+1}(x, 0, t_{k+1})|^2 &= f_{k+1, t_{k+1}}^2(x) \\ &= (1 - c^2)^{k+1} \mu^2 f_{k+1}^2(x) - \left((1 - c^2)^{k+1} \mu^2 - \frac{E_{k+1}(t_{k+1})^2}{\|u[\phi_{k+1}]\|^2} \right) \phi_{k+1}(x)^2 \\ &\leq (1 - c^2)^{k+1} \mu^2 \\ &= (1 - c^2)^k \mu^2 - c^2 (1 - c^2)^k \mu^2 \\ &= (1 - c^2)^k \mu^2 f_k^2(x) - c^2 (1 - c^2)^k \mu^2 \phi_k^2(x) \\ &\leq (1 - c^2)^k \mu^2 f_k^2(x) - \left((1 - c^2)^k \mu^2 - \frac{E_k(t_{k+1})^2}{\|u[\phi_k]\|^2} \right) \phi_k(x)^2 \\ &= f_{k, t_{k+1}}^2(x) \\ &= |u_k(x, 0, t_{k+1})|^2 \end{aligned}$$

where we used (4.18) twice. □

4.1. The case of discontinuous $e(t)$. Here we comment on how to modify the proof of Theorem 1.3 to the case when $e(t)$ is discontinuous.

Since $e(t)$ is nonincreasing, it has $M \leq \lceil 3e(0)/\varepsilon \rceil$ jumps by at least $\varepsilon/3$, where $\lceil w \rceil$ stands for the smallest integer larger or equal $w \in \mathbb{R}$. One can modify Lemma 4.1 to be able to assume that e in Theorem 1.3 is piecewise smooth with $(e(t)^2)' \leq -\zeta$, and has at most M jumps. For such e Theorem 1.3 remains valid, by incorporating the jumps into the choice of t_k 's (so that, in particular, the cardinality of $\{t_k\}$ would be $M + K$, rather than K). The proof then follows in the same way as above.

5. Proof of Theorem 1.6

The construction of a weak solution to the NSI with blow-up on a Cantor set and with an arbitrary energy profile (Theorem 1.6) is similar to the proof of the following weaker result, where the blow-up on a Cantor set is replaced by a blow-up on a single point $x_0 \in \mathbb{R}^3$.

Proposition 5.1. *Given an open set $W \subset \mathbb{R}^3$, $\varepsilon > 0$, $T > 0$ and a nonincreasing function $e: [0, T] \rightarrow [0, \infty)$ such that $e(t) \rightarrow 0$ as $t \rightarrow T$ there exists $v_0 > 0$ and a weak solution u of the NSI for all $v \in [0, v_0]$ such that $\text{supp } u(t) \subset W$ and*

$$\|u(t)\| - e(t) \leq \varepsilon \quad \text{for all } t \in [0, T],$$

and that u is unbounded in any neighbourhood of (x_0, T) for some $x_0 \in W$.

Proof. By translation we can assume that W intersects the x_1 axis. Since W is open, there exists $\bar{x} = (x_1, 0, 0)$ and $R > 0$ such that $B(\bar{x}, R) \subset W$. Let $T' \in [0, T]$ be the first time such that $e(t) \leq \varepsilon/3$ for $t \in [T', T]$. Let u be given by (3.9) and let u_0 be its rescaling (i.e. $u_0(x, t) := \lambda u(\lambda x + x', \lambda^2 t + t')$ for sufficiently large $\lambda > 0$ and appropriately chosen $x' \in \mathbb{R}^3, t' \in \mathbb{R}$) such that u_0 is defined on time interval $[T'', T]$ for some $T'' \in (T', T)$ (rather than on $[0, T_0]$, which was the case for u), $u_0(T'')$ is axisymmetric (recall u was constructed by switching between axisymmetric vector fields $u^{(j)}$, which have different axes of symmetry, see (3.9)),

$$\text{supp } u_0(t) \subset B(\bar{x}, R) \quad \text{and} \quad \|u_0(t)\| \leq \varepsilon/3 \quad \text{for all } t \in [T'', T],$$

and that $u_0(t)$ blows up (at a point inside $B(\bar{x}, R)$) as $t \rightarrow T$. Note that u_0 is axisymmetric. We will denote by (v, f, ϕ) the structure corresponding to $u_0(T'')$, that is

$$u_0(T'') = u[v, f],$$

and we let $U := \{f > 0\}$ (i.e. the set on which the structure (v, f, ϕ) is based).

We now apply Lemma 3.3 with $\varepsilon/3, T''$ and $U_1, (v_1, f_1, \phi_1)$ to obtain an axisymmetric classical solution u_1 to the NSI on time interval $[0, T'']$ (with, possibly, lower values of viscosity than u_0) that is supported in $R(\bar{U})$, $\|u_1(t) - u_0(T'')\| \leq \varepsilon/3$ for all $t \in [0, T'']$ and

$$|u_1(x, T'')| \geq |u_0(x, T'')| \quad \text{for all } x \in \mathbb{R}^3.$$

The last property guarantees that u_1 and u_0 can be combined (u_1 for times less than T'' and u_0 for times greater or equal T'') to form a weak solution of the NSI on $[0, T]$.

Case 1. $T' = 0$ (i.e. $e(0) \leq \varepsilon$).

Then

$$u(t) := \begin{cases} u_1(t) & t \in [0, T''], \\ u_0(t - T'') & t \in [T'', T] \end{cases}$$

satisfies all the claims of Proposition 5.1.

Case 2. $T' > 0$ (i.e. when the energy profile is not small for all times).

In this case we construct another weak solution to the NSI on $[0, T']$ that is disjointly supported with u_1 and whose role is to, roughly speaking, waste all the nontrivial energy (i.e. cause the energy to decrease to ε). Namely, we fix a rectangle $U_2 \Subset \mathbb{R}_+^2$ that is disjoint with U_1 and we apply Theorem 1.3 with $\varepsilon/3, T', U_2$ and $e_2 := e - \varepsilon/3$ to obtain u_2 . We extend $u_2(t)$ by zero for $t \geq T'$. Then (using (3.18)) we see that

$$u(t) := \begin{cases} u_1(t) + u_2(t) & t \in [0, T'], \\ u_0(t - T'') & t \in [T'', T] \end{cases}$$

satisfies all the claims of Proposition 5.1. □

We now turn to the proof of Theorem 1.6. For this purpose we will need to use Scheffer's construction of a weak solution to the NSI with the singular set S satisfying $d_H(S) \in [\xi, 1]$ (that is Theorem 1.5), similarly as we used u (defined in (3.9)) above.

To this end we first introduce some handy notation related to constructions of Cantor sets.

5.1. Constructing a Cantor set. In this section, which is based on Section 4.1 from Ożański [13], we discuss the general concept of constructing Cantor sets.

The problem of constructing Cantor sets is usually demonstrated in a one-dimensional setting using intervals, as in the following proposition.

Proposition 5.2. *Let $I \subset \mathbb{R}$ be an interval and let $\tau \in (0, 1)$, $M \in \mathbb{N}$ be such that $\tau M < 1$. Let $C_0 := I$ and consider the iteration in which in the j -th step ($j \geq 1$) the set C_j is obtained by replacing each interval J contained in the set C_{j-1} by M equidistant copies of τJ , each of which is contained in J , see for example Fig. 2. Then the limiting object*

$$C := \bigcap_{j \geq 0} C_j$$

is a Cantor set whose Hausdorff dimension equals $-\log M / \log \tau$.

Proof. See Example 4.5 in Falconer [7] for a proof. □

Thus if $\tau \in (0, 1)$, $M \in \mathbb{N}$ satisfy

$$\tau^\xi M \geq 1 \quad \text{for some } \xi \in (0, 1),$$

we obtain a Cantor set C with

$$d_H(C) \geq \xi. \tag{5.1}$$

Note that both the above inequality and the constraint $\tau M < 1$ (which is necessary for the iteration described in the proposition above, see also Fig. 2) can be satisfied only for $\xi < 1$. In the remainder of this section we extend the result from the proposition above to the three-dimensional setting.

Let $G \subset \mathbb{R}^3$ be a compact set, $\tau \in (0, 1)$, $M \in \mathbb{N}$, $z = (z_1, z_2, 0) \in G$, $X > 0$ be such that

$$\tau^\xi M \geq 1, \quad \tau M < 1 \tag{5.2}$$

and

$$\begin{aligned} \{\Gamma_n(G)\}_{n=1, \dots, M} \text{ is a family of pairwise disjoint subsets of } G, \\ \text{with } \text{conv}\{\Gamma_n(G) : n = 1, \dots, M\} \subset G, \end{aligned} \tag{5.3}$$

where “conv” denotes the convex hull and

$$\Gamma_n(x) := \tau x + z + (n - 1)(X, 0, 0).$$

Equivalently,

$$\Gamma_n(x_1, x_2, x_3) = (\beta_n(x_1), \gamma(x_2), \tau x_3), \tag{5.4}$$

where

$$\begin{cases} \beta_n(x) := \tau x + z_1 + (n - 1)X, \\ \gamma(x) := \tau x + z_2, \end{cases} \quad x \in \mathbb{R}, n = 1, \dots, M.$$

Now for $j \geq 1$ let

$$M(j) := \{m = (m_1, \dots, m_j) : m_1, \dots, m_j \in \{1, \dots, M\}\}$$




Fig. 2. A construction of a Cantor set C on a line (here $M = 3, j = 0, 1, 2$)

denote the set of multi-indices m . Note that in particular $M(1) = \{1, \dots, M\}$. Informally speaking, each multiindex $m \in M(j)$ plays the role of a “coordinate” which let us identify any component of the set obtained in the j -th step of the construction of the Cantor set. Namely, letting

$$\pi_m := \beta_{m_1} \circ \dots \circ \beta_{m_j}, \quad m \in M(j),$$

that is

$$\pi_m(x) = \tau^j x + z_1 \frac{1 - \tau^j}{1 - \tau} + X \sum_{k=1}^j \tau^{k-1} (m_k - 1), \quad x \in \mathbb{R} \tag{5.5}$$

we see that the set C_j obtained in the j -th step of the construction of the Cantor set C (from the proposition above) can be expressed simply as

$$C_j := \bigcup_{m \in M(j)} \pi_m(I),$$

see Fig. 2. Moreover, each $\pi_m(I)$ can be identified by, roughly speaking, first choosing the m_1 -th subinterval, then m_2 -th subinterval, ... , up to m_j -th interval, where $m = (m_1, \dots, m_j)$. This is demonstrated in Fig. 2 in the case when $m = (1, 2) \in M(2)$.

In order to proceed with our construction of a Cantor set in three dimensions let

$$\Gamma_m(x_1, x_2, x_3) := \left(\pi_m(x_1), \gamma^j(x_2), \tau^j x_3 \right). \tag{5.6}$$

Note that such a definition reduces to (5.4) in the case $j = 1$. If $j = 0$ then let $M(0)$ consist of only one element m_0 and let $\pi_{m_0} := \text{id}$. Moreover, if $m \in M(j)$ and $\bar{m} \in M(j - 1)$ is its sub-multiindex, that is $\bar{m} = (m_1, \dots, m_{j-1})$ ($\bar{m} = m_0$ if $j = 1$), then (5.3) gives

$$\Gamma_m(G) = \Gamma_{\bar{m}}(\Gamma_{m_j}(G)) \subset \Gamma_{\bar{m}}(G), \tag{5.7}$$

which is a three-dimensional equivalent of the relation $\pi_m(I) \subset \pi_{\bar{m}}(I)$ (see Fig. 2). The above inclusion and (5.3) gives that

$$\Gamma_m(G) \cap \Gamma_{\tilde{m}}(G) = \emptyset \quad \text{for } m, \tilde{m} \in M(j), \quad j \geq 1, \quad \text{with } m \neq \tilde{m}. \tag{5.8}$$

Another consequence of (5.7) is that the family of sets

$$\left\{ \bigcup_{m \in M(j)} \Gamma_m(G) \right\}_j \quad \text{decreases as } j \text{ increases.} \tag{5.9}$$

Moreover, given j , each of the sets $\Gamma_m(G)$, $m \in M(j)$, is separated from the rest by at least $\tau^{j-1}\zeta$, where $\zeta > 0$ is the distance between $\Gamma_n(G)$ and $\Gamma_{n+1}(G)$, $n = 1, \dots, M-1$ (recall (5.3)).

Taking the intersection in j we obtain

$$S' := \bigcap_{j \geq 0} \bigcup_{m \in M(j)} \Gamma_m(G), \quad (5.10)$$

and we now show that

$$\xi \leq d_H(S') \leq 1. \quad (5.11)$$

Noting that S' is a subset of a line, the upper bound is trivial. As for the lower bound note that

$$S' \supset \bigcap_{j \geq 0} \bigcup_{m \in M(j)} \Gamma_m(\text{conv}\{\Gamma_n(G) : n = 1, \dots, M\}) =: S''.$$

Thus, letting $I \subset \mathbb{R}$ be the orthogonal projection of $\text{conv}\{\Gamma_n(G) : n = 1, \dots, M\}$ onto the x_1 axis, we see that I is an interval (as the projection of a convex set; this is the reason why we put the extra requirement for the convex hull in (5.3)). Thus the orthogonal projection of S'' onto the x_1 axis is

$$\bigcap_{j \geq 0} \bigcup_{m \in M(j)} \pi_m(I) = C,$$

where C is as in the proposition above. Thus, since the orthogonal projection onto the x_1 axis is a Lipschitz map, we obtain $d_H(S'') \geq d_H(C)$ (as a property of Hausdorff dimension, see, for example, Proposition 3.3 in Falconer [7]). Consequently

$$d_H(S') \geq d_H(S'') \geq d_H(C) \geq \xi,$$

as required (recall (5.1) for the last inequality).

5.2. Sketch of the Scheffer's construction with a blow-up on a Cantor set. Based on the discussion of Cantor sets above, we now briefly sketch the proof of Theorem 1.5. To this end we fix $\xi \in (0, 1)$ and we state the analogue of Theorem 3.1 in the case of the blow-up on a Cantor set.

Theorem 5.3. *There exist a set $U \Subset P$, a structure (v, f, ϕ) , $T > 0$, $M \in \mathbb{N}$, $\tau \in (0, 1)$, $z = (z_1, z_2, 0) \in G := R(\overline{U})$, $X > 0$, $\nu_0 > 0$ with the following properties: relations (5.2) and (5.3) are satisfied and, for each $m \in M(j)$, $j \geq 0$, there exist smooth time-dependent extensions $v_t^{(m)}, f_t^{(m)}$ ($t \in [0, T]$) of v, f , respectively, such that $v_0^{(m)} = v$, $f_0^{(m)} = f$, $(v_t^{(m)}, f_t^{(m)}, \phi)$ is a structure on U for each $t \in [0, T]$, $u[v_t^{(m)}, f_t^{(m)}]$ is bounded on $\mathbb{R}^3 \times (0, T)$ and $\nabla u[v_t^{(m)}, f_t^{(m)}]$ is bounded in $L^2(\mathbb{R}^3 \times (0, T))$, independently of $m \in M(j)$, $j \geq 0$. Moreover*

$$w^{(j)}(x, t) := \sum_{m \in M(j)} u[v_t^{(m)}, f_t^{(m)}] \left(\pi_m^{-1}(\tau^j x_1), x_2, x_3 \right) \quad (5.12)$$

satisfies the NSI (1.3) in the classical sense for all $v \in [0, \nu_0]$ and $t \in [0, T]$, and

$$\left| w^{(j)}(\tau^{-j} \pi_m(y_1), \gamma(y_2), \tau y_3, T) \right| \geq \tau^{-1} |u[v, f](y)| \quad \text{for } y \in \mathbb{R}^3, m \in M(j+1). \quad (5.13)$$

Proof. See Section 4 in Ożański [13]; there the so-called *geometric arrangement* in the beginning of Section 4.2 gives U , (v, f, ϕ) , T_0 , M , τ , z and $X > 0$, and Proposition 4.3 constructs $w^{(j)}$ (which is denoted by $v^{(j)}$). \square

Observe that the claim of Theorem 3.1 (that is the vector field $u(t)$ in the statement of Theorem 3.1) is recovered by letting $M := 1$ and $u(t) := w^{(0)}(t)$.

Given the theorem above we can easily obtain Scheffer's construction with a blow-up on a Cantor set (that is a solution u to Theorem 1.5).

Indeed, let

$$u^{(j)}(x_1, x_2, x_3, t) := \tau^{-j} w^{(j)}(\tau^{-j} x_1, \gamma^{-j}(x_2), \tau^{-j} x_3, \tau^{-2j}(t - t_j)), \quad (5.14)$$

where $t_0 := 0$ and $t_j := T \sum_{k=0}^{j-1} \tau^{2k}$, as in (3.5). Observe that

$$\text{supp } u^{(j)}(t) = \bigcup_{m \in M(j)} \Gamma_m(G), \quad t \in [t_j, t_{j+1}]$$

(instead of $\Gamma_1^j(G)$, which is the case in the Scheffer's construction with point blow-up; recall (3.7)), which shrinks (as $t \rightarrow T_0^-$) to the Cantor set S' (recall (5.10)), whose Hausdorff dimension is greater or equal ξ (recall (5.11)). In fact, generalising the arguments from Sect. 3.1 we can show that $u^{(j)}$ satisfies the NSI in the classical sense for all $v \in [0, v_0]$ and $t \in [t_j, t_{j+1}]$,

$$\left| u^{(j)}(x, t_j) \right| \leq \left| u^{(j-1)}(x, t_j) \right|, \quad x \in \mathbb{R}^3, j \geq 1, \quad (5.15)$$

and that consequently the vector field

$$u(t) := \begin{cases} u^{(j)}(t) & \text{if } t \in [t_j, t_{j+1}] \text{ for some } j \geq 0, \\ 0 & \text{if } t \geq T_0 \end{cases} \quad (5.16)$$

satisfies all the claims of Theorem 1.5. We refer the reader to Section 4.2 in Ożański [13] to a more detailed explanation. Here we prove merely (5.15), which motivates the appearance of the rescalings that were used in (5.13) (i.e. the appearance of π_m , τ , γ).

It is sufficient to consider $x \in \bigcup_{m \in M(j)} \Gamma_m(G)$, as otherwise the claim is trivial. Thus suppose that $x = \Gamma_m(y)$ for some $m \in M(j)$ and $y \in G$. We obtain

$$\begin{aligned} \left| u^{(j)}(x, t_j) \right| &= \tau^{-j} \left| w^{(j)}(\tau^{-j} x_1, \gamma^{-j}(x_2), \tau^{-j} x_3, 0) \right| \\ &= \tau^{-j} \sum_{\tilde{m} \in M(j)} \left| u[v, f] \left(\Gamma_{\tilde{m}}^{-1}(x) \right) \right| \\ &= \tau^{-j} |u[v, f](y)| \\ &\leq \tau^{-(j-1)} \left| w^{(j-1)} \left(\tau^{-(j-1)} \pi_m(y_1), \gamma(y_2), \tau y_3, T \right) \right| \\ &= \left| u^{(j-1)} \left(\pi_m(y_1), \gamma^j(y_2), \tau^j y_3, t_j \right) \right| \\ &= \left| u^{(j-1)}(x, t_j) \right|, \end{aligned}$$

as required, where we used (5.8) (so that $\Gamma_{\tilde{m}}^{-1}(\Gamma_m(y)) = y \chi_{\tilde{m}=m}$) in the third equality and (5.13) in the inequality (recall also the definitions (5.14), (5.12), (5.6) of $u^{(j)}$, $w^{(j)}$, Γ_m , respectively).

Furthermore, we note that $u \in L^\infty((0, \infty); L^2(\mathbb{R}^3))$ and $\nabla u \in L^2(\mathbb{R}^3 \times (0, \infty))$ (which is required by the definition of weak solutions to the NSI, Definition 1.1). Indeed, $u^{(j)}$ consists of M^j vector fields, each scaled by τ^{-j} , and so the claim follows from the fact that $M\tau < 1$ (so that $\sup_{t \in [t_j, t_{j+1}]} \|u^{(j)}(t)\|^2 \approx (M\tau)^j$ decreases to zero as $j \rightarrow \infty$, and $\sum_{j \geq 0} \int_{t_j}^{t_{j+1}} \|\nabla u^{(j)}(s)\|^2 ds \approx \sum_{j \geq 0} (M\tau)^j$ converges).

5.3. *Proof of Theorem 1.6.* Given u constructed in the previous section, Theorem 1.6 follows in the same way as Proposition 5.1.

6. A Sharpening of the Edge Effect Lemma (Lemma 2.2)

Here we prove Lemma 4.2 (the sharpening of the “edge effects” Lemma 2.2), which was used in the proof of Theorem 1.3.

In order to prove the lemma we will need a certain generalised Mean Value Theorem. For $g: \mathbb{R} \rightarrow \mathbb{R}$ let $g[a, b]$ denote the finite difference of g on $[a, b]$,

$$g[a, b] := \frac{g(a) - g(b)}{a - b}$$

and let $g[a, b, c]$ denote the finite difference of $g[\cdot, b]$ on $[a, c]$,

$$g[a, b, c] := \left(\frac{g(a) - g(b)}{a - b} - \frac{g(c) - g(b)}{c - b} \right) / (a - c).$$

Lemma 6.1 (Generalised mean value theorem). *If $a < b < c$, g is continuous in $[a, c]$ and twice differentiable in (a, c) then there exists $\xi \in (a, c)$ such that $g[a, b, c] = g''(\xi)/2$.*

Proof. We follow the argument of Theorem 4.2 in Conte and de Boor [5]. Let

$$p(x) := g[a, b, c](x - b)(x - c) + g[b, c](x - c) + g(c).$$

Then p is a quadratic polynomial approximating g at a, b, c , that is $p(a) = g(a)$, $p(b) = g(b)$, $p(c) = g(c)$. Thus the error function $e(x) := g(x) - p(x)$ has at least 3 zeros in $[a, c]$. A repeated application of Rolle’s theorem gives that e'' has at least one zero in (a, c) . In other words, there exists $\xi \in (a, c)$ such that $g''(\xi) = p''(\xi) = 2g[a, b, c]$. \square

Corollary 6.2. *If $g \in C^3(a - \delta, a + \delta)$ is such that $g = 0$ on $(a - \delta, a]$ and $g''' > 0$ on $(a, a + \delta)$ for some $a \in \mathbb{R}$, $\delta > 0$ then*

$$\begin{cases} g''(x) > 0, \\ 0 < g'(x) < (x - a)g''(x), \\ g(x) < (x - a)^2 g''(x) \end{cases} \quad \text{for } x \in (a, a + \delta).$$

Proof. Since $g''' > 0$ on $(a, a + \delta)$ we see that g'' is increasing on this interval and so also positive (as $g''(x) = 0$ for $x \leq a$). This also gives the first inequality in the second claim, while the second inequality follows from the Mean Value Theorem, $g'(x) =$

$(x-a)g''(\xi) < (x-a)g''(x)$, where $\xi \in (a, x)$. The last claim follows from the lemma above by noting that $2a-x \in (a-\delta, a]$ (so that $g(2a-x) = g(a) = 0$), and so

$$\begin{aligned} g(x) &= g(2a-x) - 2g(a) + g(x) = 2(x-a)^2 g[2a-x, a, x] \\ &= (x-a)^2 g''(\xi) < (x-a)^2 g''(x), \end{aligned}$$

where $\xi \in (2a-x, x)$. □

We can now prove Lemma 4.2; that is, given $a > 0$, an open rectangle $U \in \mathbb{R}_+^2$ that is at least a away from the x_1 axis (i.e. $U = (a_1, b_1) \times (a_2, b_2)$ with $a_2 > a$) and $\eta \in (0, \min\{1, (b_1 - a_1)/2, (b_2 - a_2)/2\})$ we construct $f \in C_0^\infty(\mathbb{R}_+^2; [0, 1])$ such that

$$\begin{aligned} \text{supp } f &= \bar{U}, \quad f > 0 \text{ in } U \text{ with } f = 1 \text{ on } U_\eta, \\ Lf &> 0 \quad \text{in } U \setminus U_{c'\eta}, \quad \text{with } f > c \text{ in } U_{c'\eta/2}, \end{aligned}$$

where $c, c' \in (0, 1/2)$ depend only on a .

Proof of Lemma 4.2. Without loss of generality we can assume that $a < 1$. Let $h \in C^\infty(\mathbb{R}; [0, 1])$ be a nondecreasing function such that

$$h(x) = \begin{cases} 0 & x \leq 0, \\ e^{-1/x^2} & x \in (0, 1/2), \\ 1 & x \geq 1. \end{cases}$$

Let

$$C_h := \|h\|_{C^2([0,1])} \in [1, \infty).$$

Observe that $h''' > 0$ on $(0, 1/2)$. Let $h_\eta(x) := h(x/\eta)$ and

$$f(x_1, x_2) := f_1(x_1)f_2(x_2),$$

where

$$f_i(x) := h_\eta(x - a_i)h_\eta(b_i - x), \quad i = 1, 2,$$

see Fig. 3.

Clearly

$$f_i''' > 0 \text{ on } (a_i, a_i + \eta/2) \quad \text{and} \quad f_i''' < 0 \text{ on } (b_i - \eta/2, b_i), \quad i = 1, 2.$$

Moreover $\text{supp } f = \bar{U}$, $f > 0$ in U , and $f = 1$ on U_η . We will show that

$$Lf > 0 \quad \text{on} \quad U \setminus U_{\eta'} \tag{6.1}$$




Fig. 3. The f_i 's, $i = 1, 2$

for

$$\eta' := c' \eta, \quad (6.2)$$

where

$$c' := \frac{a}{6\sqrt{C_h}} e^{-9/a^2} \in (0, 1/6). \quad (6.3)$$

Note that, since $c'/2 < 1/2$, we have that $f > (h_\eta(c'\eta/2))^2 = (e^{-4/(c')^2})^2 =: c$ in $U_{c'\eta/2}$. Thus the proof of the lemma is complete when we show (6.1).

To this end let

$$\eta'' := \frac{a\eta}{3}. \quad (6.4)$$

Obviously $\eta' \leq \eta'' \leq \eta \leq 1$. Letting

$$\begin{aligned} g_1(x_1) &:= f_1''(x_1), \\ g_2(x_2) &:= f_2''(x_2) + f_2'(x_2)/x_2 - f_2(x_2)/x_2^2, \end{aligned}$$

we see that

$$\begin{aligned} Lf(x_1, x_2) &= f_1''(x_1)f_2(x_2) + f_1(x_1)f_2''(x_2) + f_1(x_1)f_2'(x_2)/x_2 - f_1(x_1)f_2(x_2)/x_2^2 \\ &= g_1(x_1)f_2(x_2) + f_1(x_1)g_2(x_2). \end{aligned}$$

(Recall (2.5).) We need to show that the expression on the right-hand side above is positive in $U \setminus U_{\eta'}$. For this we first show the *claim*:

$$g_2 > f_2''/4 > 0 \quad \text{on } (a_2, a_2 + \eta'') \cup (b_2 - \eta'', b_2). \quad (6.5)$$

The claim follows from the corollary of the generalised Mean Value Theorem (see Corollary 6.2), which gives that $f_2'(x_2) > 0$ and $f_2(x_2) < (x_2 - a_2)^2 f_2''(x_2)$ for $x_2 \in (a_2, a_2 + \eta'')$ (since f_2 is given by the rescaled exponential function e^{-1/x_2^2} due to $\eta'' < \eta/2$). Thus

$$\begin{aligned} g_2(x_2) &> f_2''(x_2) - f_2(x_2)/x_2^2 \\ &> f_2''(x_2) \left(1 - \left(\frac{x_2 - a_2}{x_2} \right)^2 \right) \\ &> f_2''(x_2) \left(1 - \left(\frac{\eta''}{a_2} \right)^2 \right) \\ &> \frac{8}{9} f_2''(x_2) \\ &> \frac{1}{4} f_2''(x_2) > 0 \end{aligned}$$




Fig. 4. The “ η'' -corners” and “ η' -strips”

for such x_2 , where we also used the fact that $\eta'' < a_2/3$. On the other hand, applying Corollary 6.2 to $g_2(b_2 - \cdot)$ we obtain $f_2'(x_2) > (x_2 - b_2)f_2''(x_2)$ and $f_2(x_2) < (x_2 - b_2)^2 f_2''(x_2)$ for $x_2 \in (b_2 - \eta'', b_2)$, and so

$$\begin{aligned} g_2(x_2) &= f_2''(x_2) + f_2'(x_2)/x_2 - f_2(x_2)/x_2^2 \\ &> f_2''(x_2) \left(1 + \frac{x_2 - b_2}{x_2} - \left(\frac{x_2 - b_2}{x_2} \right)^2 \right) \\ &> f_2''(x_2) \left(1 - \frac{\eta''}{b_2 - \eta''} - \left(\frac{\eta''}{b_2 - \eta''} \right)^2 \right) \\ &> f_2''(x_2)/4 > 0 \end{aligned}$$

for such x_2 , where we also used the fact that $\eta''/(b_2 - \eta'') < 1/2$ (as $b_2 > a_2 > 3\eta''$), and so the *claim* follows.

Using the *claim* we see that g_i, f_i are positive on $(a_i, a_i + \eta'') \cup (b_i - \eta'', b_i)$, $i = 1, 2$. Thus

$$Lf > 0 \quad \text{in } ((a_1, a_1 + \eta'') \cup (b_1 - \eta'', b_1)) \times ((a_2, a_2 + \eta'') \cup (b_2 - \eta'', b_2)),$$

that is in the “ η'' -corners” of U , see Fig. 4.

Now let

$$\begin{aligned} m &:= e^{-9/a^2}, \\ M &:= \frac{3C_h}{\eta^2 a^2}. \end{aligned}$$

A direct calculation gives that

$$f_i \geq m, |g_i| \leq M \quad \text{in } [a_i + \eta'', b_i - \eta''], i = 1, 2.$$

Moreover,

$$\frac{m}{4} - (\eta')^2 M > 0.$$

Indeed, the left-hand side is simply $e^{-9/a^2}(1/4 - 3/36) > 0$.

We will show that

$$Lf > 0 \quad \text{in } [a_1 + \eta'', b_1 - \eta''] \times ((a_2, a_2 + \eta') \cup (b_2 - \eta', b_2))$$

$$\text{and in } ((a_1, a_1 + \eta') \cup (b_1 - \eta', b_1)) \times [a_2 + \eta'', b_2 - \eta''], \quad (6.6)$$

that is in the “ η' -strips” at ∂U between the η'' -corners, see Fig. 4. This will prove (6.1) (and so finish the proof) as the η' -strips together with the η'' -corners contain $U \setminus U_{\eta'}$.

In order to prove (6.6) let first $x_1 \in [a_1 + \eta'', b_1 - \eta'']$ and $x_2 \in (a_2, a_2 + \eta')$. Then $g_1(x_1) > -M$, $g_2(x_2) > f_2''(x_2)/4$ (from (6.5)), $f_2(x_2) < (x_2 - a_2)^2 f_2''(x_2)$ (from Corollary 6.2), $f_1(x_1) > m$, and so

$$Lf(x_1, x_2) = g_1(x_1)f_2(x_2) + f_1(x_1)g_2(x_2) > -Mf_2(x_2) + f_1(x_1)f_2''(x_2)/4$$

$$> f_2''(x_2) \left(-M(x_2 - a_2)^2 + m/4 \right) > f_2''(x_2) \left(m/4 - M(\eta')^2 \right) > 0.$$

As for $x_2 \in (b_2 - \eta', b_2)$, simply replace a_2 in the above calculation by b_2 . The opposite case, that is the case $x_1 \in (a_1, a_1 + \eta') \cup (b_1 - \eta', b_1)$, $x_2 \in [a_2 + \eta'', b_2 - \eta'']$, follows in the same way. \square

Acknowledgements. The author is very grateful to James Robinson for his careful reading of a draft of this article and his numerous comments, which significantly improved its quality. The author was supported partially by EPSRC as part of the MASDOC DTC at the University of Warwick, Grant No. EP/HO23364/1, and partially by postdoctoral funding from ERC 616797.

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

- Blömker, D., Romito, M.: Regularity and blow up in a surface growth model. *Dyn. Part. Differ. Equ.* **6**(3), 227–252 (2009)
- Blömker, D., Romito, M.: Local existence and uniqueness in the largest critical space for a surface growth model. *Nonlinear Diff. Equ. Appl.* **19**(3), 365–381 (2012)
- Buckmaster, T., Vicol, V.: Nonuniqueness of weak solutions to the Navier–Stokes equation. *Ann. Math.* (2) **189**(1), 101–144 (2019)
- Caffarelli, L., Kohn, R., Nirenberg, L.: Partial regularity of suitable weak solutions of the Navier–Stokes equations. *Commun. Pure Appl. Math.* **35**(6), 771–831 (1982)
- Conte, S.D., de Boor, C.: *Elementary Numerical Analysis: An Algorithmic Approach*. McGraw-Hill Book Co., New York (1972)
- Escauriaza, L., Seregin, G.A., Šverák, V.: $L_{3,\infty}$ -solutions of Navier–Stokes equations and backward uniqueness. *Russian Math. Surv.* **58**(2), 211–250 (2003)
- Falconer, K.: *Fractal Geometry—Mathematical Foundations and Applications*, 3rd edn. Wiley, Chichester (2014)
- Hopf, E.: Über die Anfangswertaufgabe für die hydrodynamischen Grundgleichungen. *Math. Nachr.* **4**, 213–231 (1951). English translation by Andreas Klöckner
- Kukavica, I.: Partial regularity results for solutions of the Navier–Stokes system. In: *Partial Differential Equations and Fluid Mechanics*, Vol. 364 of London Math. Soc. Lecture Note Ser., pp. 121–145. Cambridge University Press, Cambridge (2009)
- Ladyzhenskaya, O.A., Seregin, G.A.: On partial regularity of suitable weak solutions to the three-dimensional Navier–Stokes equations. *J. Math. Fluid Mech.* **1**(4), 356–387 (1999)
- Leray, J.: Sur le mouvement d’un liquide visqueux emplissant l’espace. *Acta Math.* **63**, 193–248. (An English translation due to Robert Terrell is available at <http://www.math.cornell.edu/~bterrell/leray.pdf> and [arXiv:1604.02484](https://arxiv.org/abs/1604.02484))
- Lin, F.: A new proof of the Caffarelli–Kohn–Nirenberg theorem. *Commun. Pure Appl. Math.* **51**(3), 241–257 (1998)

13. Ożański, W.S.: The Partial Regularity Theory of Caffarelli, Kohn, and Nirenberg and Its Sharpness. Lecture Notes in Mathematical Fluid Mechanics. Springer/Birkhäuser, Berlin (2019)
14. Ożański, W.S., Pooley, B.C.: Leray's fundamental work on the Navier–Stokes equations: a modern review of “Sur le mouvement d'un liquide visqueux emplissant l'espace”. In: Fefferman, C., Robinson, J.C., Rodrigo, J.L. (eds.) *Partial Differential Equations in Fluid Mechanics*. LMS Lecture Notes Series. Cambridge University Press, Cambridge (2018)
15. Ożański, W.S., Robinson, J.C.: Partial regularity for a surface growth model. *SIAM J. Math. Anal.* **51**(1), 228–255 (2019)
16. Robinson, J.C., Rodrigo, J.L., Sadowski, W.: *The Three-Dimensional Navier–Stokes Equations*, Vol. 157 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge (2016)
17. Robinson, J.C., Sadowski, W.: Almost-everywhere uniqueness of Lagrangian trajectories for suitable weak solutions of the three-dimensional Navier–Stokes equations. *Nonlinearity* **22**(9), 2093–2099 (2009)
18. Scheffer, V.: Partial regularity of solutions to the Navier–Stokes equations. *Pac. J. Math.* **66**(2), 535–552 (1976a)
19. Scheffer, V.: Turbulence and Hausdorff dimension. In: *Turbulence and Navier–Stokes equations* (Proceedings Conference, Univ. Paris-Sud, Orsay, 1975), Springer LNM, vol. 565, pp. 174–183. Springer, Berlin (1976b)
20. Scheffer, V.: Hausdorff measure and the Navier–Stokes equations. *Commun. Math. Phys.* **55**(2), 97–112 (1977)
21. Scheffer, V.: The Navier–Stokes equations in space dimension four. *Commun. Math. Phys.* **61**(1), 41–68 (1978)
22. Scheffer, V.: The Navier–Stokes equations on a bounded domain. *Commun. Math. Phys.* **73**(1), 1–42 (1980)
23. Scheffer, V.: A solution to the Navier–Stokes inequality with an internal singularity. *Commun. Math. Phys.* **101**(1), 47–85 (1985)
24. Scheffer, V.: Nearly one-dimensional singularities of solutions to the Navier–Stokes inequality. *Commun. Math. Phys.* **110**(4), 525–551 (1987)
25. Vasseur, A.F.: A new proof of partial regularity of solutions to Navier–Stokes equations. *Nonlinear Differ. Equ. Appl.* **14**(5–6), 753–785 (2007)

Communicated by C. De Lellis