

Monopoles and the Gibbons–Manton Metric

Roger Bielawski

Max-Planck-Institut für Mathematik, Gottfried-Claren-Strasse 26, 53225 Bonn, Germany.
E-mail: rogerb@mpim-bonn.mpg.de

Received: 6 June 1997 / Accepted: 7 October 1997

Abstract: We show that, in the region where monopoles are well separated, the L^2 -metric on the moduli space of n -monopoles is exponentially close to the T^n -invariant hyperkähler metric proposed by Gibbons and Manton. The proof is based on a description of the Gibbons–Manton metric as a metric on a certain moduli space of solutions to Nahm’s equations, and on twistor methods. In particular, we show how the twistor description of monopole metrics determines the asymptotic metric.

The construction of the Gibbons–Manton metric in terms of Nahm’s equations yields a class of interesting (pseudo)-hyperkähler metrics. For example we show, for each semisimple Lie group G and a maximal torus $T \leq G$, the existence of a $G \times T$ -invariant (pseudo)-hyperkähler manifold whose hyperkähler quotients by T are precisely Kronheimer’s hyperkähler metrics on $G^{\mathbb{C}}/T^{\mathbb{C}}$. A similar result holds for Kronheimer’s ALE-spaces.

The moduli space M_n of (framed) static $SU(2)$ -monopoles of charge n , i.e. solutions to Bogomolny equations $d_A \Phi = *F$, carries a natural hyperkähler metric [1]. The geodesic motion in this metric is a good approximation to the dynamics of low energy monopoles [26, 33]. For the charge $n = 2$ the metric has been determined explicitly by Atiyah and Hitchin [1], and it follows from their explicit formula that when the two monopoles are well separated, the metric becomes (exponentially fast) the Euclidean Taub-NUT metric with a negative mass parameter. It was also shown by N. Manton [27] that this asymptotic metric can be determined by treating well-separated monopoles as dyons. The equations of motion for a pair of dyons in \mathbb{R}^3 are found to be equivalent to the equations for geodesic motion on Taub-NUT space.

For an arbitrary charge n , it was shown in [3] that, when the individual monopoles are well-separated, the L^2 -metric is close (as the inverse of the separation distance) to the flat Euclidean metric. Gibbons and Manton [14] have then calculated the Lagrangian for the motion of n dyons in \mathbb{R}^3 and shown that it is equivalent to the Lagrangian for geodesic motion in a hyperkähler metric on a torus bundle over the configuration space $\tilde{C}_n(\mathbb{R}^3)$.

This metric is T^n -invariant and has a simple algebraic form. Gibbons and Manton have conjectured, by analogy with the $n = 2$ case, that the exact n -monopole metric differs from their metric by an exponentially small amount as the separation gets large. We shall prove this conjecture here.

Our strategy is as follows. We construct a certain moduli space \tilde{M}_n of solutions to Nahm's equations which carries a T^n -invariant hyperkähler metric. Using twistor methods we identify this metric as the Gibbons–Manton metric. Finally, we show that the metrics on \tilde{M}_n and M_n are exponentially close. This proof adapts equally well to the asymptotic behaviour of $SU(N)$ -monopole metrics with maximal symmetry breaking, as will be shown elsewhere.

The asymptotic picture can be explained in the twistor setting. We recall that a monopole is determined (up to framing) by a curve S – the spectral curve – in $T\mathbb{C}P^1$, which satisfies certain conditions [16]. One of these is triviality of the line bundle L^{-2} over S , and a nonzero section of this bundle is the other ingredient needed to determine the metric [19, 1]. Asymptotically we have now the following situation. When the individual monopoles become well separated the spectral curve of the n -monopole degenerates (exponentially fast) into the union of spectral curves S_i of individual monopoles, while the section of L^{-2} becomes (also exponentially fast) n meromorphic sections of L^{-2} over the individual S_i . The zeros and poles of these sections occur only at the intersection points of the curves S_i . This information (and the topology of the asymptotic region of M_n) is, as we show in the last section, sufficient to conclude that the asymptotic metric is the Gibbons–Manton metric.

The construction of the moduli space of solutions to Nahm's equations which gives the Gibbons–Manton metric admits various generalizations. Some of them are described in Sect. 4. Let us recall that Kronheimer [23] has shown existence of hyperkähler structures $M(\tau_1, \tau_2, \tau_3)$ on $G^{\mathbb{C}}/T^{\mathbb{C}}$, where G is a compact semisimple Lie group and $T \leq G$ is a maximal torus. These structures are parameterized by the cohomology classes $\tau_1, \tau_2, \tau_3 \in \text{Lie}(T)$ of the three Kähler forms. We show (in Sect. 4) that there is a (pseudo)-hyperkähler manifold M_G with a tri-Hamiltonian action of T such that, if $\mu : M_G \rightarrow \text{Lie}(T) \otimes \mathbb{R}^3$ is the hyperkähler moment map, then the hyperkähler quotient $\mu^{-1}(\tau_1, \tau_2, \tau_3)/T$ of M_G by T is precisely Kronheimer's $M(\tau_1, \tau_2, \tau_3)$. A similar construction can be done for Kronheimer's ALE-spaces.

The article is organized as follows. In Sects. 1 and 2 we recall the definitions of the Gibbons–Manton and monopole metrics. In Sect. 3 we introduce the moduli space \tilde{M}_n of solutions to Nahm's equations and give heuristic arguments why the metric on \tilde{M}_n should be exponentially close to the monopole metric. In Sect. 4, as a preliminary step to study \tilde{M}_n we introduce yet another moduli space of solutions to Nahm's equations, somewhat simpler than \tilde{M}_n . In that section we also discuss the relation with Kronheimer's metrics mentioned above. In Sect. 5 we identify \tilde{M}_n as a differential complex, and finally complex-symplectic manifold. In Sect. 6 we calculate the twistor space of \tilde{M}_n and identify its hyperkähler metric as the Gibbons–Manton metric. In Sect. 7 we finally show that the monopole metric and the metric on \tilde{M}_n are exponentially close. The short Sect. 8 shows how one can read off the Gibbons–Manton metric, as the asymptotic form of the monopole metric, from the twistor description of the latter.

1. The Gibbons–Manton metric

The Gibbons–Manton metric [14] is an example of $4n$ -dimensional (pseudo)-hyperkähler metric admitting a tri-Hamiltonian (hence isometric) action of the n -dimensional torus

T^n . Such metrics have particularly nice properties and were studied by several authors [25, 18, 32]. The Gibbons–Manton metric was described as a hyperkähler quotient of a flat quaternionic vector space by Gibbons and Rychenkova in [15]. We recall here this description, which we slightly modify to better suit our purposes. We start with flat hyperkähler metrics g_1 and g_2 on $M_1 = (S^1 \times \mathbb{R}^3)^n$ and $M_2 = \mathbb{H}^{n(n-1)/2}$. We consider a pseudo-hyperkähler metric on the product manifold $M = M_1 \times M_2$ given by $g = g_1 - g_2$. The complex structures on \mathbb{H} are given by the right multiplication by quaternions i, j, k . The metric g_1 is invariant under the obvious action (by translations) of $T^n = (S^1)^n$ and the metric g_2 is invariant under the left diagonal action of $T^{n(n-1)/2}$. We consider a homomorphism $\phi : T^{n(n-1)/2} \rightarrow T^n$ given by

$$(t_{ij})_{i < j} \mapsto \left(\prod_{j=i+1}^n t_{ij} \prod_{j=1}^{i-1} t_{ji}^{-1} \right)_{i=1, \dots, n} .$$

This defines an action of $T^{n(n-1)/2}$ on $M = M_1 \times M_2$ by $t \cdot (m_1, m_2) = (\phi(t) \cdot m_1, t \cdot m_2)$. Gibbons and Rychenkova have shown that the hyperkähler quotient of (M, g) by this action of $T^{n(n-1)/2}$ is the Gibbons–Manton metric.

We remark that, if we choose coordinates (t_i, \mathbf{x}_i) on M_1 , $t_i \in S^1$ and $\mathbf{x}_i \in \mathbb{R}^3$, and quaternionic coordinates q_{ij} , $i < j$, on $\mathbb{H}^{n(n-1)/2}$, then the moment map equations are:

$$\frac{1}{2} q_{ij} i \bar{q}_{ij} = \mathbf{x}_i - \mathbf{x}_j. \tag{1.1}$$

As long as $\mathbf{x}_i \neq \mathbf{x}_j$ for $i \neq j$, the torus $T^{n(n-1)/2}$ acts freely on the zero-set of the moment map. The quotient of this set by $T^{n(n-1)/2}$ is a smooth hyperkähler manifold which we denote by M_{GM} . The action of T^n on M_1 induces a free tri-Hamiltonian action on M_{GM} for which the moment map is just $(\mathbf{x}_1, \dots, \mathbf{x}_n)$. This makes M_{GM} into a T^n -bundle over the configuration space $\check{C}_n(\mathbb{R}^3)$ of n distinct points in \mathbb{R}^3 . We shall now determine this bundle. We recall that a basis of $H_2(\check{C}_n(\mathbb{R}^3), \mathbb{Z})$ is given by the $k(k-1)/2$ 2-spheres,

$$S_{ij}^2 = \{ \mathbf{x}_k \in \mathbb{R}^3; |\mathbf{x}_i - \mathbf{x}_j| = \text{const}, \mathbf{x}_k = \text{const} \text{ if } k \neq i, j \}, \tag{1.2}$$

where $i < j$. We have

Proposition 1.1. *The hyperkähler moment map for the action of T^n makes M_{GM} into a T^n -bundle over $\check{C}_n(\mathbb{R}^3)$ determined by the element (s_1, \dots, s_n) of $H^2(\check{C}_n(\mathbb{R}^3), \mathbb{Z}^n)$ given by*

$$s_k(S_{ij}^2) = \begin{cases} -1 & \text{if } k = i \\ 1 & \text{if } k = j \\ 0 & \text{otherwise.} \end{cases}$$

Proof. From the formula (1.1) it follows that restricting the bundle to a fixed S_{ij}^2 is equivalent to considering the case $n = 2$. In other words $s_k(S_{ij}^2) = 0$ if $k \neq i, j$ and we have to consider only one quaternionic coordinate q_{ij} . The zero-set of the moment map is $\frac{1}{2} q_{ij} i \bar{q}_{ij} = \mathbf{x}_i - \mathbf{x}_j$ and the circle S^1 by which we quotient acts by $t \cdot (q_{ij}, (t_i, \mathbf{x}_i), (t_j, \mathbf{x}_j)) = (t q_{ij}, (t t_i, \mathbf{x}_i), (t^{-1} t_j, \mathbf{x}_j))$. The quotient can be obtained by setting $t_i = 1$ and the induced action of the i^{th} generator s_i of T^n is then given by left multiplication by s_i^{-1} on q_{ij} . Since the map $q_{ij} \rightarrow \frac{1}{2} q_{ij} i \bar{q}_{ij}$ with the left action of S^1 on $\{q_{ij} \in \mathbb{H}; |q_{ij}| = 1\}$ is the Hopf bundle, it follows that $s_i(S_{ij}^2) = -1$. A similar argument shows that $s_j(S_{ij}^2) = 1$. \square

In particular, $(\bar{t}, \bar{\mathbf{x}}) = (t_i, \mathbf{x}_i)$ form local coordinates on M_{GM} . The metric tensor can be then written in the form [32]:

$$g = \Phi d\bar{\mathbf{x}} \cdot d\bar{\mathbf{x}} + \Phi^{-1}(d\bar{t} + A)^2,$$

where the matrix Φ and the 1-form A depend only on the \mathbf{x}_i and satisfy certain linear PDE's. In particular, Φ determines the metric. For the Gibbons–Manton metric

$$4\Phi_{ij} = \begin{cases} 1 - \sum_{k \neq i} \frac{1}{\|\mathbf{x}_i - \mathbf{x}_k\|} & \text{if } i = j \\ \frac{1}{\|\mathbf{x}_i - \mathbf{x}_j\|} & \text{if } i \neq j. \end{cases}$$

2. Nahm's Equations and Monopole Metrics

We shall recall in this section the description of the L^2 -metric on the moduli space of charge n $SU(2)$ -monopoles in terms of Nahm's equations. A proof that the Nahm transform [30, 16] between the two moduli spaces is an isometry was given by Nakajima in [31].

One starts with the space \mathcal{A} of quadruples (T_0, T_1, T_2, T_3) of smooth $\mathfrak{u}(n)$ -valued functions on $(-1, 1)$ such that T_1, T_2, T_3 have simple poles at ± 1 with residues $\frac{1}{2}\rho(\sigma_i), i = 1, 2, 3$, where $\rho : \mathfrak{su}(2) \rightarrow \mathfrak{u}(n)$ is the standard irreducible n -dimensional representation of $\mathfrak{su}(2)$ and σ_i are the Pauli matrices. Equipped with the L^2 -norm (given by a biinvariant inner product on $\mathfrak{u}(n)$), \mathcal{A} becomes a flat quaternionic affine space. There is an isometric and triholomorphic action of the gauge group \mathcal{G} of $U(n)$ -valued functions $g : [-1, 1] \rightarrow U(n)$ which are 1 at ± 1 :

$$\begin{aligned} T_0 &\mapsto \text{Ad}(g)T_0 - \dot{g}g^{-1}, \\ T_i &\mapsto \text{Ad}(g)T_i, \quad i = 1, 2, 3. \end{aligned} \tag{2.1}$$

The zero-set of the hyperkähler moment map for this action is then described by Nahm's equations [30]:

$$\dot{T}_i + [T_0, T_i] + \frac{1}{2} \sum_{j,k=1,2,3} \epsilon_{ijk} [T_j, T_k] = 0, \quad i = 1, 2, 3. \tag{2.2}$$

The quotient of the space of solutions by \mathcal{G} is a smooth hyperkähler manifold M_n of dimension $4n$. By the above mentioned result of Nakajima, M_n is the moduli space of (framed) charge n $SU(2)$ -monopoles. With respect to any complex structure M_n is biholomorphic to the space of based rational maps of degree n on $\mathbb{C}P^1$ [13].

If we replace $U(n)$ by $SU(n)$ (resp. by $PSU(n)$) in the above description, we obtain the moduli space of strongly centered (resp. centered) $SU(2)$ -monopoles of charge n .

Remark 2.1. A similar construction can be done for any compact Lie group G . We require $\rho : \mathfrak{su}(2) \rightarrow \mathfrak{g}$ to be a Lie algebra homomorphism whose image lies in the regular part of \mathfrak{g} . We obtain a smooth hyperkähler manifold of dimension $4 \text{rank } G$ which can be identified with a totally geodesic submanifold of a certain moduli space of $SU(N)$ -monopoles (with a minimal symmetry breaking). Alternatively, as a complex manifold, it is a desingularization of $(\mathfrak{h}^{\mathbb{C}} \times T^{\mathbb{C}}) / W$, where $T^{\mathbb{C}}$ is a maximal torus in $G^{\mathbb{C}}$, $\mathfrak{h}^{\mathbb{C}}$ its Lie algebra, and W the corresponding Weyl group [6].

The tangent space to M_n can be described as the space of solutions to the linearized Nahm’s equations and satisfying the condition of being orthogonal (in the L^2 -metric) to vectors arising from infinitesimal gauge transformations. In other words the tangent space to M_n at a solution (T_0, T_1, T_2, T_3) can be identified with the set of solutions (t_0, t_1, t_2, t_3) to the following system of linear equations:

$$\begin{aligned} \dot{t}_0 + [T_0, t_0] + [T_1, t_1] + [T_2, t_2] + [T_3, t_3] &= 0, \\ \dot{t}_1 + [T_0, t_1] - [T_1, t_0] + [T_2, t_3] - [T_3, t_2] &= 0, \\ \dot{t}_2 + [T_0, t_2] - [T_1, t_3] - [T_2, t_0] + [T_3, t_1] &= 0, \\ \dot{t}_3 + [T_0, t_3] + [T_1, t_2] - [T_2, t_1] - [T_3, t_0] &= 0. \end{aligned} \tag{2.3}$$

The metric is defined by

$$\|(t_0, t_1, t_2, t_3)\|^2 = \frac{1}{2} \int_{-1}^1 \sum_0^3 \|t_i\|^2. \tag{2.4}$$

The three anti-commuting complex structures can be seen by writing a tangent vector as $t_0 + it_1 + jt_2 + kt_3$.

3. The Asymptotic Moduli Space

We shall now construct a one-parameter family of moduli spaces $\tilde{M}_n(c)$, $c \in \mathbb{R}$, of solutions to Nahm’s equations carrying (pseudo-)hyperkähler metrics. We shall see later on that these metrics are the Gibbons–Manton metric with different mass parameters.

We consider the subspace Ω_1 of exponentially fast decaying functions in $C^1[0, \infty]$, i.e.:

$$\Omega_1 = \left\{ f : [0, \infty] \rightarrow \mathfrak{u}(n); \exists_{\eta>0} \sup_t (e^{\eta t} \|f(t)\| + e^{\eta t} \|df/dt\|) < +\infty \right\}. \tag{3.1}$$

As in the previous section, $\rho : \mathfrak{su}(2) \rightarrow \mathfrak{u}(n)$ is the standard irreducible n -dimensional representation of $\mathfrak{su}(2)$ (in particular, $\rho(\sigma_1)$ is a diagonal matrix). We denote by \mathfrak{h} the (Cartan) subalgebra of $\mathfrak{u}(n)$ consisting of diagonal matrices.

Let $\tilde{\mathcal{A}}_n$ be the space of C^1 -functions (T_0, T_1, T_2, T_3) defined on $(0, +\infty]$ and satisfying (cf. [23]):

- (i) T_1, T_2, T_3 have simple poles at 0 with $\text{res } T_i = \frac{1}{2}\rho(\sigma_i)$;
- (ii) $T_i(+\infty) \in \mathfrak{h}$ for $i = 0, \dots, 3$;
- (iii) $(T_1(+\infty), T_2(+\infty), T_3(+\infty))$ is a regular triple, i.e. its centralizer is \mathfrak{h} ;
- (iv) $(T_i(t) - T_i(+\infty)) \in \Omega_1$ for $i = 0, 1, 2, 3$.

Next we shall define the relevant gauge group. The Lie algebra of our gauge group $\mathcal{G}(c)$ is the space of C^2 -paths $\rho : [0, +\infty) \rightarrow \mathfrak{u}(n)$ such that

- (i) $\rho(0) = 0$ and $\dot{\rho}$ has a limit in \mathfrak{h} at $+\infty$;
- (ii) $(\dot{\rho} - \dot{\rho}(+\infty)) \in \Omega_1$, and $[\tau, \rho] \in \Omega_1$ for any regular element $\tau \in \mathfrak{h}$;
- (iii) $c\dot{\rho}(+\infty) + \lim_{t \rightarrow +\infty} (\rho(t) - t\dot{\rho}(+\infty)) = 0$.

It is the Lie algebra of the Lie group

$$\mathcal{G}(c) = \{g : [0, +\infty) \rightarrow U(n); g(0) = 1, s(g) := \lim \dot{g}g^{-1} \in \mathfrak{h}, (\tau - \text{Ad}(g)\tau) \in \Omega_1, (\dot{g}g^{-1} - s(g)) \in \Omega_1, \exp(cs(g)) \lim (g(t) \exp(-ts(g))) = 1\}.$$

Remark. The last condition in the definition of $\mathcal{G}(c)$ means that $g(t)$ is asymptotic to $\exp(ht - ch)$ for some diagonal h .

We introduce a family of metrics on $\tilde{\mathcal{A}}_n$. Let (t_0, t_1, t_2, t_3) be a tangent vector to the space $\tilde{\mathcal{A}}_n$ at a point (T_0, T_1, T_2, T_3) . The functions t_i are now regular at 0, $i = 0, \dots, 3$. We put

$$\|(t_0, t_1, t_2, t_3)\|_c^2 = c \sum_0^3 \|t_i(+\infty)\|^2 + \int_0^{+\infty} \sum_0^3 (\|t_i(s)\|^2 - \|t_i(+\infty)\|^2) ds. \tag{3.2}$$

We observe that the group $\mathcal{G}(c)$ acting by (2.1) preserves the metric $\|\cdot\|_c$ and the three complex structure of the flat hyperkähler manifold $\tilde{\mathcal{A}}_n$. We define $\tilde{M}_n(c)$ as the (formal) hyperkähler quotient of $\tilde{\mathcal{A}}_n$ by $\mathcal{G}(c)$ (with respect to the metric $\|\cdot\|_c$). The zero set of the moment map is given by (2.2) (here condition (iii) in the definition of $\text{Lie}(\mathcal{G}(c))$ is essential) and so $\tilde{M}_n(c)$ is defined as the moduli space of solutions to Nahm’s equations:

$$\tilde{M}_n(c) = \{\text{solutions to (2.2) in } \tilde{\mathcal{A}}_n\} / \mathcal{G}(c).$$

Remark. If $c > 0$, then the metric (3.2) on $\tilde{M}_n(c)$ will be seen to be positive definite if $(T_1(+\infty), T_2(+\infty), T_3(+\infty))$ is sufficiently far from the walls of Weyl chambers. On the other hand, if $c < 0$, then the metric will be shown to be everywhere negative definite. Therefore, for $c < 0$ we should really replace $\|\cdot\|_c$ with its negative; it is, however more convenient to consider the metrics $\|\cdot\|_c$.

We observe that sending a solution T_i to the solution $rT_i(rt)$ for any $r > 0$ induces a homothety of factor r between $\tilde{M}_n(c)$ and $\tilde{M}_n(rc)$.

Before we begin the detailed study of $\tilde{M}_n(c)$, let us explain why we expect this metric to be exponentially close to the monopole metric. It is known [4] that the solutions to Nahm’s equations on $(0, 2)$ corresponding to a well-separated monopole are exponentially close to being constant away from the boundary points (i.e. on any $[\epsilon, 2 - \epsilon]$). The same is true for solutions on the half line $(0, +\infty)$: as long as the triple $(T_1(+\infty), T_2(+\infty), T_3(+\infty))$ is regular, the solutions are exponentially close to being constant away from 0 [23] (it is helpful to notice that the space of regular triples is the same as the space $\tilde{C}_n(\mathbb{R}^3)$ of distinct points in \mathbb{R}^3). Our strategy is to take two solutions, on half-lines $(0, \infty)$ and $(-\infty, 2)$ with the same values at $\pm\infty$, cut them off at $t = 1$ and use this non-smooth solution on $(0, 2)$ (with correct boundary behaviour) to obtain an exact solution to the monopole Nahm data. The exact solution will differ from the approximate one by an exponentially small amount. Furthermore the part of the half-line solutions which we have cut off is exponentially close to being constant and, for $c = 1$, contributes an exponentially small amount to the metric $\|\cdot\|_c$ (all estimates are uniform and can be differentiated). This can be seen from the fact that we can rewrite (3.2) as

$$\|(t_0, t_1, t_2, t_3)\|_c^2 = \int_0^c \sum_0^3 \|t_i(s)\|^2 + \int_c^{+\infty} \sum_0^3 (\|t_i(s)\|^2 - \|t_i(+\infty)\|^2) ds. \tag{3.3}$$

The first term, together with the corresponding term for the solution on $(-\infty, 2)$, is exponentially close to the monopole metric (for $c = 1$).

4. Moduli Space of Regular Semisimple Adjoint Orbits

In order to obtain information about $\tilde{M}_n(c)$ we need to consider first another moduli space of solutions to Nahm’s equations, defined analogously, except that we require the solutions to be smooth at $t = 0$. This space, which can be defined for an arbitrary compact Lie group G , is of some interest as all hyperkähler structures on $G^{\mathbb{C}}/T^{\mathbb{C}}$ (here $T^{\mathbb{C}}$ is a maximal torus) due to Kronheimer [23] can be obtained from it as hyperkähler quotients (see Theorem 4.3 below). A reader who is primarily interested in monopoles should think of G as $U(n)$.

Let us first recall how Kronheimer constructs hyperkähler metrics on $G^{\mathbb{C}}/T^{\mathbb{C}}$. Let \mathfrak{h} be the Lie algebra of $T^{\mathbb{C}}$ and let $(\tau_1, \tau_2, \tau_3) \in \mathfrak{h}^3$ be a regular triple, i.e. one whose centralizer is \mathfrak{h} . For a fixed $\eta > 0$, consider the Banach space

$$\Omega_1^\eta = \left\{ f : [0, \infty) \rightarrow \mathfrak{g}; \sup_t (e^{\eta t} \|f(t)\| + e^{\eta t} \|df/dt\|) < +\infty \right\}$$

with the norm $\|f\| = \sup_t (e^{\eta t} \|f(t)\| + e^{\eta t} \|df/dt\|)$. Define $\mathcal{A}^\eta(\tau_1, \tau_2, \tau_3)$ as the space of C^1 -functions $(T_0, T_1, T_2, T_3) : (0, +\infty) \rightarrow \mathfrak{g}$ which satisfy:

$$\{T_0(t), (T_i(t) - \tau_i); i = 1, 2, 3\} \subset \Omega_1^\eta.$$

Define also \mathcal{G}^η by replacing Ω_1 with Ω_1^η in the definition of \mathcal{G} given in the previous section. Kronheimer shows then that for small enough η ,

$$M(\tau_1, \tau_2, \tau_3) = \{\text{solutions to (2.2) in } \mathcal{A}^\eta(\tau_1, \tau_2, \tau_3)\} / \mathcal{G}^\eta,$$

equipped with the L^2 metric is a smooth hyperkähler manifold, diffeomorphic to $G^{\mathbb{C}}/T^{\mathbb{C}}$. Furthermore, if (τ_2, τ_3) is regular, then $M(\tau_1, \tau_2, \tau_3)$ is biholomorphic, with respect to the complex structure I , to the complex adjoint orbit of $\tau_2 + i\tau_3$.

We observe that the union of all $M(\tau_1, \tau_2, \tau_3)$ has a natural topology and it is, in fact, a smooth manifold. We shall show now that there is a T -bundle over this union which carries a (pseudo)-hyperkähler metric. We define the space \mathcal{A}_G by omitting the condition (i) in the definition of $\tilde{\mathcal{A}}_n$ in the previous section. Instead we require that the T_i are smooth at $t = 0$ for $i = 0, 1, 2, 3$. We define $M_G(c)$, $c \in \mathbb{R}$, as the (formal) hyperkähler quotient of \mathcal{A}_G by $\mathcal{G}(c)$ with respect to the metric (3.2). We have:

Proposition 4.1. *$M_G(c)$ equipped with the metric (3.2) is a smooth hyperkähler manifold. The tangent space at a solution (T_0, T_1, T_2, T_3) is described by Eqs. (2.3).*

We remark that the metric 3.2 may be degenerate at some points. However the hypercomplex structure is defined everywhere.

Proof. Define $M_G^\eta(c)$ by replacing Ω with Ω^η in the definition of $M_G(c)$. By the exponential decay property of solutions to Nahm’s equations ([23], Lemma 3.4), a neighbourhood of a particular element in $M_G(c)$ is canonically identified with its neighbourhood in $M_G^\eta(c)$ for small enough η . Therefore we can use the transversality arguments of [23], Lemma 3.8 and Proposition 3.9 (with a slight modification due to condition (iii) in the definition of $\text{Lie}(\mathcal{G}(c))$) to deduce the smoothness. The fact that the metric is hyperkähler is, formally, the consequence of the fact that $M_G(c)$ is a hyperkähler quotient. One can, in fact, check directly that the three Kähler forms are closed. We shall also, later on, identify the complex structures and the complex symplectic forms proving their closedness. \square

We observe now that the action on \mathcal{A}_G of gauge transformations which are asymptotic to $\exp(-th + \lambda h)$, $h \in \mathfrak{h}$, $\lambda \in \mathbb{R}$, induce a free isometric action of $T = \exp(\mathfrak{h})$ on $M_G(c)$. In fact this action is tri-Hamiltonian and a simple calculation shows

Proposition 4.2. *The hyperkähler moment map $\mu = (\mu_1, \mu_2, \mu_3)$ for the action of T on $M_G(c)$ is given by $\mu_i(T_0, T_1, T_2, T_3) = T_i(+\infty)$ for $i = 1, 2, 3$.*

As an immediate corollary we have:

Theorem 4.3. *Let (τ_1, τ_2, τ_3) be a regular triple in \mathfrak{h}^3 . The hyperkähler quotient $\mu^{-1}(\tau_1, \tau_2, \tau_3)/T$ of $M_G(c)$ by the torus T is isometric to Kronheimer’s $M(\tau_1, \tau_2, \tau_3)$.*

We have also a tri-Hamiltonian action of G on $M_G(c)$ given by the gauge transformations with arbitrary values at $t = 0$. The hyperkähler moment map for this action is $(T_1(0), T_2(0), T_3(0))$.

We have two other group actions on $M_G(c)$. There is a free isometric and triholomorphic action of the Weyl group $W = N(T)/T$ given by the gauge transformations which become constant (and in W) exponentially fast.

Finally there is a free isometric $SU(2)$ -action which rotates the complex structures. As a consequence it has a globally defined Kähler potential for each Kähler form (cf. [18]). The potential for ω_2 (or ω_3) is given by the moment map for the action of a circle in $SU(2)$ which preserves I . This is easily seen to be

$$K_J = c \sum_{i=2}^3 \|T_i(+\infty)\|^2 + \int_0^{+\infty} \sum_{i=2}^3 \left(\|T_i(s)\|^2 - \|T_i(+\infty)\|^2 \right) ds.$$

Remark 4.4. There is a similar (pseudo)-hyperkähler manifold with a torus action such that the hyperkähler quotients by this torus are isometric to Kronheimer’s ALE-metrics on the minimal resolution of a given Kleinian singularity \mathbb{C}^2/Γ [24]. This manifold is defined as M_G except that the T_i have poles at $t = 0$ with the residues defined by a subregular homomorphism $\mathfrak{su}(2) \rightarrow \mathfrak{g}$ (cf. [6, 5]).

Remark 4.5. One can observe that $M_G(0)$ is a cone metric (with the $\mathbb{R}_{>0}$ -action given by $T_i(t) \mapsto rT_i(rt)$) and in fact, it is an \mathbb{H}^* -bundle over a pseudo-quaternion-Kähler manifold (cf. [34]).

Remark 4.6. It is instructive to consider the Kähler analogue of $M_G(c)$. The Kähler metrics on G/T (cf. [2]) can be described (cf. [7]) as the natural L^2 -metrics on the moduli space of solutions to $\hat{T}_1 = [T_1, T_0]$ with $T_0(t), (T_1(t) - \tau) \in \Omega_1$ for a fixed regular element τ of \mathfrak{h} (this gives the Kähler form whose cohomology class is τ). We can now do a construction similar to that of $M_G(c)$ to obtain moduli spaces $\hat{M}_G(c)$ with a $(G \times T)$ -invariant pseudo-Kähler metric whose Kähler quotients by T are precisely the Kähler metrics on G/T . In this case it is easy to compute both the topology and the complex structure of $\hat{M}_G(c)$: $\hat{M}_G(c)$ is diffeomorphic to $G \times \{\text{regular elements of } \mathfrak{h}\}$, and the complex structure at $(1, h)$ is given by $I(v + w, p) = (I_0 v - p, w)$, where $v \perp \mathfrak{h}$, $w, p \in \mathfrak{h}$ and I_0 is the complex structure on G/T .

5. $\tilde{M}_n(c)$ as a Manifold

We now return to the space $\tilde{M}_n(c)$ defined in Sect. 3. Our first task is to show that this space is smooth. We shall show that $\tilde{M}_n(c)$ is a smooth hyperkähler quotient of the

product of the space $M_{U(n)}(c - 1)$ considered in the previous section and of another moduli space of solutions to Nahm’s equations. This latter space, denoted by N_n , is given by $u(n)$ -valued solutions to Nahm’s equations defined on $(0, 1]$ smooth at $t = 1$ and with the same poles as $\tilde{M}_n(c)$ at $t = 0$. The gauge group consists of gauge transformations which are identity at $t = 0, 1$. Equipped with the metric (2.4) this is a smooth hyperkähler manifold [6, 11]. It admits a tri-Hamiltonian action of $U(n)$ given by gauge transformations with arbitrary values at $t = 1$. In addition, we consider the space $M_{U(n)}(c - 1)$ defined in the previous section. We identify it this time with the space of solutions on $[1, +\infty]$ via the map $T_i(t) \mapsto T_i(t + 1)$ (so that the gauge transformations behave now, near $+\infty$, as elements of $\mathcal{G}(c)$).

It is easy to observe that the space $\tilde{M}_n(c)$ is the hyperkähler quotient of $N_n \times M_{U(n)}(c - 1)$ by the diagonal action of $U(n)$ (cf. [6]; the moment map equations simply match the functions T_1, T_2, T_3 at $t = 1$; after that, quotienting by G means that the remaining gauge transformations are smooth at $t = 1$). Using this description of $\tilde{M}_n(c)$ we can finally show

Proposition 5.1. *$\tilde{M}_n(c)$ equipped with the metric (3.2) is a smooth hyperkähler manifold. The tangent space at a solution (T_0, T_1, T_2, T_3) is described by the Eqs. (2.3).*

Proof. Since the metric (3.2) may be degenerate, we still have to show that the moment map equations on $N_n \times M_{U(n)}(c - 1)$ are everywhere transversal. Consider a particular point in $M_{U(n)}(c - 1)$ which we represent by a solution $m = (T_0, T_1, T_2, T_3)$ with $T_0(+\infty) = 0$ and $T_i(+\infty) = \tau_i, i = 1, 2, 3$. Let μ be the hyperkähler moment map for the action of G on $N_n \times M_{U(n)}$. We observe that the image of $d\mu|_m$ contains the image of $d\mu'|_m, \mu'$ being the hyperkähler moment map for the action of G on $N_n \times M(\tau_1, \tau_2, \tau_3)$ (Kronheimer’s definition of $M(\tau_1, \tau_2, \tau_3)$ was recalled in the previous section). The metric on $N_n \times M(\tau_1, \tau_2, \tau_3)$ is non-degenerate and, as G acts freely, $d\mu'|_m$ is surjective. Thus $d\mu$ is surjective at each point in $N_n \times M_{U(n)}(c - 1)$ and $\tilde{M}_n(c)$ is smooth. \square

We observe that, as in the case of $M_{U(n)}(c)$, $\tilde{M}_n(c)$ has isometric actions of the torus T^n (defined as the diagonal subgroup of $U(n)$), of the symmetric group S_n , and of $SU(2)$. In particular, the hyperkähler moment map for the action of T^n is still given by the values of T_1, T_2, T_3 at infinity (cf. Proposition 4.2).

We can describe the topology of $\tilde{M}_n(c)$:

Proposition 5.2. *$\tilde{M}_n(c)$ is a principal T^n -bundle over the configuration space $\tilde{C}_n(\mathbb{R}^3)$ of n distinct points in \mathbb{R}^3 .*

We postpone identifying this bundle until the next section (Proposition 6.3).

Proof. The space $\tilde{C}_n(\mathbb{R}^3)$ is the space of regular triples in the subalgebra of diagonal matrices and the moment map μ for the action of T^n gives us a projection $\tilde{M}_n(c) \rightarrow \tilde{C}_n(\mathbb{R}^3)$. Let us consider a fixed regular triple (τ_1, τ_2, τ_3) and all elements of $\tilde{M}_n(c)$ with $T_i(+\infty) = \tau_i, i = 1, 2, 3$, i.e. $\mu^{-1}(\tau_1, \tau_2, \tau_3)$. For each such solution we can make T_0 identically 0 by some gauge transformation g with $g(0) = 1$. This is defined uniquely up to the action of $\mathcal{G} \times T^n$ and so the set of T^n -orbits projecting via μ to (τ_1, τ_2, τ_3) can be identified with the set of solutions to Nahm’s equations with $T_0 \equiv 0, T_1, T_2, T_3$ having the appropriate residues at $t = 0$ and being conjugate to τ_1, τ_2, τ_3 at infinity. By the considerations at the beginning of this section this space is the hyperkähler quotient of $N_n \times M(\tau_1, \tau_2, \tau_3)$ by $U(n)$. The arguments of [6] show that the corresponding complex-symplectic quotient can be identified with the intersection of a regular semisimple adjoint

orbit of $GL(n, \mathbb{C})$ with the slice to the regular nilpotent orbit. This intersection is a single point. Finally, in order to identify in this case the hyperkähler quotient with the complex-symplectic one we can adapt the argument in the proof of Proposition 2.20 in [20]. \square

Our next task is to describe the complex structure of $\tilde{M}_n(c)$ (because of the action of $SU(2)$ all complex structures are equivalent). As usual (cf. [13]), if we choose a complex structure, say I , we can introduce complex coordinates on the moduli space of solutions to Nahm's equations by writing $\alpha = T_0 + iT_1$ and $\beta = T_2 + iT_3$. The Nahm equations can be then written as one complex and one real equation:

$$\frac{d\beta}{dt} = [\beta, \alpha], \tag{5.1}$$

$$\frac{d}{dt}(\alpha + \alpha^*) = [\alpha^*, \alpha] + [\beta^*, \beta]. \tag{5.2}$$

By the remark made at the beginning of this section, $\tilde{M}_n(c)$ is the hyperkähler quotient of the product manifold $N_n \times M_{U(n)}(c-1)$. We shall show that as a complex symplectic manifold $\tilde{M}_n(c)$ is the complex-symplectic quotient of $N_n \times M_{U(n)}(c-1)$. Let us recall the complex structure of N_n [13, 19, 6, 12]. Let e_1, \dots, e_n denote the standard basis of \mathbb{C}^n . There is a unique solution w_1 of the equation

$$\frac{dw}{dt} = -\alpha w \tag{5.3}$$

with

$$\lim_{t \rightarrow 0} \left(t^{-(n-1)/2} w_1(t) - e_1 \right) = 0. \tag{5.4}$$

Setting $w_i(t) = \beta^{i-1}(t)w_1(t)$, we obtain a solution to (5.3) with

$$\lim_{t \rightarrow 0} \left(t^{i-(n+1)/2} w_i(t) - e_i \right) = 0.$$

The complex gauge transformation $g(t)$ with $g^{-1} = (w_1, \dots, w_n)$ makes α identically zero and sends $\beta(t)$ to the constant matrix

$$B(\beta_1, \dots, \beta_n) = \begin{pmatrix} 0 & \dots & 0 & (-1)^{n+1} S_n \\ 1 & \ddots & & (-1)^n S_{n-1} \\ & \ddots & \ddots & \vdots \\ 0 & \dots & 1 & S_1 \end{pmatrix}. \tag{5.5}$$

Here β_i denote the (constant) eigenvalues of $\beta(t)$ and S_i is the i^{th} elementary symmetric polynomial in $\{\beta_1, \dots, \beta_n\}$.

The mapping $(\alpha, \beta) \rightarrow (g(1), B)$ gives a biholomorphism between (N_n, I) and $GL(n, \mathbb{C}) \times \mathbb{C}^n$ [6].

We describe the complex structure of $\tilde{M}_n(c)$ as follows:

Proposition 5.3. *There exists a T^n -equivariant biholomorphism between $\tilde{M}_n(c)$ and an open subset of*

$$\left(\coprod_n \{[g, b] \in GL(n, \mathbb{C}) \times_N (\mathfrak{d} + \mathfrak{n}); gbg^{-1} \text{ is of the form (5.5)}\} \right) / \sim,$$

where \mathfrak{d} denotes diagonal matrices, the union is over unipotent algebras \mathfrak{n} (with respect to \mathfrak{d}) and $N = \exp \mathfrak{n}$. Furthermore, the relation \sim is given as follows: $[g, d+n] \sim [g', d'+n']$ if and only if $n \in \mathfrak{n}, n' \in \mathfrak{n}'$, and either $\mathfrak{n}' \subset \mathfrak{n}$ and there exists an $m \in N$ such that $gm^{-1} = g', \text{Ad}(m)(d+n) = d'+n'$ or vice versa (i.e. $\mathfrak{n} \subset \mathfrak{n}'$ etc.).

Remark. It will follow from the description of the twistor space that this biholomorphism is actually onto. Proving this right now would require showing that the T^n -action on $\tilde{M}_n(c)$ extends to the global action of $(\mathbb{C}^*)^n$. This, in turn, requires showing existence of solutions to a mixed Dirichlet-Robin problem on the half-line - something that seems quite tricky.

Proof. Fix a unipotent algebra \mathfrak{n} and consider the set of all solutions $(\alpha, \beta) = (T_0 + iT_1, T_2 + iT_3)$ on $[1, +\infty)$ such that the intersection of the sum of positive eigenvalues of $\text{ad}(iT_1(+\infty))$ with $C(\beta(+\infty))$ is contained in \mathfrak{n} . Let $M(\mathfrak{n}; c-1)$ be the corresponding subset of $M_{U(\mathfrak{n})}(c)$. We observe that, since $(T_1(+\infty), T_2(+\infty), T_3(+\infty))$ is a regular triple, the projection of $T_1(+\infty)$ onto $\mathfrak{d}^{\mathbb{C}} \cap C(\beta(+\infty))$ is a regular element, and so \mathfrak{n} contains the unipotent radical of a Borel subalgebra of $C(\beta(+\infty))$ for any element of $M(\mathfrak{n}; c-1)$. Using gauge freedom, we always make $T_0(+\infty) = 0$ and, by Proposition 4.1 of Biquard [8], such a representative is of the form $g(\alpha(+\infty), \beta(+\infty) + \text{Ad}(\exp\{-\alpha(+\infty)t\})n)$, where $n \in \mathfrak{n}$ and g is a bounded $Gl(n, \mathbb{C})$ -valued gauge transformation. The transformation g is defined modulo $\exp\{-\alpha(+\infty)t\}g_0 \exp\{\alpha(+\infty)t\}$ with $g_0 \in P = \exp(\mathfrak{d} + \mathfrak{n})$. Since $T_0(+\infty) = 0$ and T_0 is decaying exponentially fast, g has a limit (in $T^{\mathbb{C}}$) at $+\infty$. If we replace $g(t)$ by $g'(t) = g(t)g(+\infty)^{-1} \exp\{-\alpha(+\infty)t + c\alpha(+\infty)\}$, then $(\alpha, \beta) = (g'(0), \beta(+\infty) + n')$ for an $n' \in \mathfrak{n}$. The transformation g' , which satisfies (at infinity) the boundary condition of an element of $\mathcal{G}(c-1)^{\mathbb{C}}$, is now defined modulo constant gauge transformations in N . Moreover $g'(1)$ is independent of $\mathcal{G}(c-1)$ and we obtain a map $\phi : M(\mathfrak{n}) \rightarrow Gl(n, \mathbb{C}) \times_N (\mathfrak{d} + \mathfrak{n})$ by sending (α, β) to $(g'(1), \beta(+\infty) + n')$. Considering the infinitesimal version of this construction shows that ϕ is holomorphic.

Since ϕ is $U(\mathfrak{n})$ -equivariant, it is (locally) $Gl(n, \mathbb{C})$ -equivariant. We can adapt the argument of Proposition 2.20 in [19] to show that $\tilde{M}_n(c)$ is the complex-symplectic quotient of $N_n \times M_{U(\mathfrak{n})}(c-1)$ by (local action of) $Gl(n, \mathbb{C})$. Let us restrict attention to $N_n \times M(\mathfrak{n})$. The complex symplectic moment map at the point (g, B) of N_n is $-g^{-1}Bg$ (here $g \in Gl(n, \mathbb{C})$ and B is of the form (5.5)) and the complex symplectic moment map at the point corresponding to $[g', \beta_d + n]$ is $g'(\beta_d + n)g'^{-1}$ (here β_d is diagonal and $n \in \mathfrak{n}$). The moment map equation for the diagonal action of $Gl(n, \mathbb{C})$ is $g^{-1}Bg = g'(\beta_d + n)g'^{-1}$. If we now quotient by $Gl(n, \mathbb{C})$, i.e. send g to identity, we shall end up with the set of $[g', b] \in Gl(n, \mathbb{C}) \times_N (\mathfrak{d} + \mathfrak{n})$ such that $g'bg'^{-1} = B$ (B is determined by the diagonal part of b). This identifies the charts described in this proposition. By going through the procedure we can conclude that the charts for different \mathfrak{n} are matched as claimed.

So far we have shown that there is a holomorphic map ϕ from $\tilde{M}_n(c)$ to the manifold M described in the statement. We still have to show that ϕ is 1-1. By construction our map is T^n -equivariant, and so $(\mathbb{C}^*)^n$ -equivariant (where the action is defined). Since the $(\mathbb{C}^*)^n$ -action on M is free, it is free on $\tilde{M}_n(c)$. Furthermore the $(\mathbb{C}^*)^n$ -action on M leaves invariant sets of the form $M \cap (Gl(n, \mathbb{C}) \times_N (d + \mathfrak{n}))$, $d \in \mathfrak{d}$. Each such set is a single orbit of $(\mathbb{C}^*)^n$ and so ϕ is 1-1. \square

The above description of $\tilde{M}_n(c)$ is rather complicated. We remark that the open dense subset where $\beta(+\infty)$ is regular corresponds to $\mathfrak{n} = 0$, i.e. to

$$\{(\beta_d, g); \beta_d = \text{diag}(\beta_1, \dots, \beta_n), \beta_i \neq \beta_j \text{ if } i \neq j, g\beta_d g^{-1} = B(\beta_1, \dots, \beta_n)\}.$$

We shall denote the corresponding subset of $\tilde{M}_n(c)$ by $\tilde{M}_n^{\text{reg}}(c)$. We observe that an element g of $Gl(n, \mathbb{C})$ which sends $\text{diag}(\beta_1, \dots, \beta_n)$ to $B(\beta_1, \dots, \beta_n)$ is of the form

$$g = V(\beta_1, \dots, \beta_n)^{-1} \text{diag}(u_1, \dots, u_n), \tag{5.6}$$

where $u_i \neq 0$ and $V(\beta_1, \dots, \beta_n)$ is the Vandermonde matrix, i.e. $V_{ij} = (\beta_i)^{j-1}$. We can calculate the complex symplectic form $\omega = \omega_2 + i\omega_3$ on $\tilde{M}_n^{\text{reg}}(c)$:

Proposition 5.4. *The complex symplectic form ω on $\tilde{M}_n^{\text{reg}}(c)$ is given, in coordinates $\beta_i, u_i, i = 1, \dots, n$, by*

$$\sum_{i=1}^n \frac{du_i}{u_i} \wedge d\beta_i - \sum_{i < j} \frac{d\beta_i \wedge d\beta_j}{\beta_i - \beta_j}. \tag{5.7}$$

Proof. First, we calculate ω on the subset of $M_{U(n)}(c-1)$, where $\beta(+\infty)$ is regular. This subset is biholomorphic to $Gl(n, \mathbb{C}) \times \{\text{regular elements of } \mathfrak{h}^{\mathbb{C}}\}$ and according to the proof of Proposition 5.3, an element (α, β) of this set corresponding to $(g, \beta_d) \in Gl(n, \mathbb{C}) \times \mathfrak{h}$ can be written as $(\alpha, \beta) = (-\dot{g}(t)g^{-1}, g(t)\beta_d g(t)^{-1})$, where $g(t)$ is a complex gauge transformation with $g(0) = g$. Therefore a tangent vector $(a(t), b(t))$ at (α, β) can be written as

$$(a, b) = (-g\dot{\rho}g^{-1}, g(b_d + [\rho, \beta_d]g^{-1})), \tag{5.8}$$

where ρ is dual to $g^{-1}dg$ and b_d is dual to $d\beta_d$. The complex symplectic form on $M_{U(n)}(c-1)$ is given by

$$\omega = (c-1) \text{tr}(d\alpha(+\infty) \wedge d\beta(+\infty)) + \int_0^{+\infty} \text{tr}(d\alpha \wedge d\beta - d\alpha(+\infty) \wedge d\beta(+\infty)).$$

For two tangent vectors (a, b) and (\hat{a}, \hat{b}) , corresponding, via (5.8), to (ρ, b_d) and $(\hat{\rho}, \hat{b}_d)$ we obtain

$$\omega = -\text{tr}(b_d \hat{\rho} - \hat{\rho} \hat{b}_d - [\rho, \hat{\rho}]\beta_d),$$

where $\rho = \rho(0), \hat{\rho} = \hat{\rho}(0)$. To calculate the symplectic form on $\tilde{M}_n^{\text{reg}}(c)$ it remains to substitute (5.6) for g . Let us write u for $\text{diag}(u_1, \dots, u_n)$. Then ρ becomes dual to $u^{-1}du - u^{-1}dVV^{-1}u$. Let us write ν for the tangent vector dual to $u^{-1}du$ and Υ for the tangent vector dual to dVV^{-1} . Since ν is diagonal and the i^{th} row of Υ is of the form $b_i s$ (here we write $b_d = \text{diag}(b_1, \dots, b_n)$), for a covector s , we can write ω as

$$\omega = -\text{tr}(b_d \hat{\nu} - \hat{\nu} \hat{b}_d - [\Upsilon, \hat{\Upsilon}]\beta_d).$$

It remains to calculate $\text{tr}[\Upsilon, \hat{\Upsilon}]\beta_d$. Let us write W_{ij} for the $(i, j)^{\text{th}}$ entry of V^{-1} , i.e.

$$W_{ij} = (-1)^{n-i} S_{n-i}(\beta_1, \dots, \hat{\beta}_j, \dots, \beta_n) / \prod_{k \neq j} (\beta_j - \beta_k), \tag{5.9}$$

S_k being the k^{th} elementary symmetric polynomial ($S_0 = 1$). We calculate the $(i, i)^{\text{th}}$ entry of $[\Upsilon, \hat{\Upsilon}]$ as

$$\sum_j (b_i \hat{b}_j - \hat{b}_i b_j) \left(\sum_k (k-1) \beta_i^{k-2} W_{kj} \right) \left(\sum_k (k-1) \beta_j^{k-2} W_{ki} \right).$$

This means that

$$\text{tr}[\Upsilon, \hat{\Upsilon}]_{\beta_d} = \sum_{i < j} (b_i \hat{b}_j - \hat{b}_i b_j)(\beta_i - \beta_j) \left(\sum_k (k-1)\beta_i^{k-2} W_{kj} \right) \left(\sum_k (k-1)\beta_j^{k-2} W_{ki} \right).$$

Formula (5.7) will be proven if we can show (for $i \neq j$) the following identity:

$$\left(\sum_k (k-1)\beta_i^{k-2} W_{kj} \right) \left(\sum_k (k-1)\beta_j^{k-2} W_{ki} \right) = \frac{-1}{(\beta_i - \beta_j)^2}. \tag{5.10}$$

According to (5.9) we have

$$\sum_k (k-1)\beta_i^{k-2} W_{kj} = \frac{\sum_k (k-1)\beta_i^{k-2} (-1)^{n-k} S_{n-k}(\beta_1, \dots, \hat{\beta}_j, \dots, \beta_n)}{\prod_{s \neq j} (\beta_j - \beta_s)}. \tag{5.11}$$

We compute the numerator of this expression. We set $p = n - 1$ and $(a_1, \dots, a_p) = (\beta_1, \dots, \hat{\beta}_j, \dots, \beta_n)$. Then the numerator can be written as

$$\sum_{s=0}^p (p-s)(-1)^s a_i^{p-1-s} S_s(a_1, \dots, a_p) = \frac{d}{dt} \left(\sum_{s=0}^p (-1)^s S_s t^{p-s} \right)_{t=a_i}.$$

Since $\sum S_s t^s = \prod (1 + a_s t)$, we can rewrite the expression under the derivative as

$$\sum_{s=0}^p (-1)^s S_s t^{p-s} = \prod_{s=0}^p (t - a_s).$$

Taking the derivative and substituting a_i for t , finally gives

$$\sum_{s=0}^p (p-s)(-1)^s a_i^{p-1-s} = \prod_{s \neq i} (a_i - a_s).$$

Going back to (5.11), we have

$$\sum_k (k-1)\beta_i^{k-2} W_{kj} = \frac{\prod_{s \neq i, j} (\beta_i - \beta_s)}{\prod_{s \neq j} (\beta_j - \beta_s)},$$

from which (5.10) follows. \square

Remark 5.5. Setting

$$p_i = u_i / \prod_{j > i} (\beta_i - \beta_j),$$

the formula (5.7) can be rewritten as

$$\omega = \sum_{i=1}^n \frac{dp_i}{p_i} \wedge d\beta_i.$$

6. The Twistor Space and the Metric on $\tilde{M}_n(c)$

We shall now identify the twistor space $Z(c)$ of $\tilde{M}_n(c)$. As a first step, we observe, after Hitchin et al. [18], that the hyperkähler moment map μ for the T^n -action defines a moment map, also denoted by μ , for the complex-symplectic form along the fibers $Z(c) \rightarrow \mathbb{C}P^1$. This μ is a map from $Z(c)$ to $O(2) \otimes \mathbb{C}^n$. We shall first identify the open subset $Z^{\text{reg}}(c)$ of $Z(c)$ defined as the set

$$Z^{\text{reg}}(c) = \mu^{-1}(O(2) \otimes \mathbb{C}^n - O(2) \otimes \Delta), \tag{6.1}$$

where Δ is the generalized diagonal in \mathbb{C}^n . In terms of the coordinates $(\beta_1, \dots, \beta_n)$ and (u_1, \dots, u_n) given by (5.6), $Z^{\text{reg}}(c)$ has the following description:

Proposition 6.1. *$Z^{\text{reg}}(c)$ is obtained by taking two copies of $\mathbb{C} \times (\mathbb{C}^n - \Delta) \times (\mathbb{C}^*)^n$ with coordinates (ζ, β_i, u_i) and $(\tilde{\zeta}, \tilde{\beta}_i, \tilde{u}_i)$, $i = 1, \dots, n$, and identifying over $\zeta \neq 0$ by*

$$\begin{aligned} \tilde{\zeta} &= \zeta^{-1}, \\ \tilde{\beta}_i &= \zeta^{-2}\beta_i, \\ \tilde{u}_i &= \zeta^{-(n-1)} \exp\{-c\beta_i/\zeta\}u_i. \end{aligned}$$

The real structure is given by

$$\begin{aligned} \zeta &\mapsto -1/\bar{\zeta}, \\ \beta_i &\mapsto -\bar{\beta}_i/\bar{\zeta}^2, \\ u_i &\mapsto \bar{u}_i^{-1} (1/\bar{\zeta})^{n-1} \prod_{j \neq i} (\bar{\beta}_i - \bar{\beta}_j) e^{c\bar{\beta}_i/\bar{\zeta}}. \end{aligned}$$

Finally, the complex symplectic form along the fibers is given by (5.7).

Proof. For any hyperkähler moduli space of solutions to Nahm’s equations one can trivialize the twistor space by choosing an affine coordinate ζ on $\mathbb{C}P^1$ and then putting $\eta = \beta + (\alpha + \alpha^*)\zeta - \beta^*\zeta^2$, $u = \alpha - \beta^*\zeta$ for $\zeta \neq \infty$, and $\tilde{\eta} = \beta/\zeta^2 + (\alpha + \alpha^*)/\zeta - \beta^*$, $\tilde{u} = -\alpha^* - \beta/\zeta$ for $\zeta \neq 0$. Then, over $\zeta \neq 0, \infty$, we have $\tilde{\eta} = \eta/\zeta^2$, $\tilde{u} = u - \eta/\zeta$. Moreover, the real structure is $\zeta \mapsto -1/\bar{\zeta}$, $\eta \mapsto -\eta^*/\bar{\zeta}^2$, $u \mapsto -u^* + \eta^*/\bar{\zeta}$ (cf. [12, 9]).

We now have to go through the procedure in the proof of Proposition 5.3 to describe Z^{reg} in coordinates (ζ, β_i, u_i) and $(\tilde{\zeta}, \tilde{\beta}_i, \tilde{u}_i)$. First we describe the twistor space of N_n in coordinates (g, B) and (\tilde{g}, \tilde{B}) defined right after (5.5) (cf. [12]). Going through the procedure assigning (g, B) to (α, β) , we see that $\tilde{B} = B(\beta_1/\zeta^{-2}, \dots, \beta_n/\zeta^{-2})$. On the other hand g is given by $g = g(1)$, where $g(t)$ is a complex gauge transformation such that $\frac{d}{dt}g^{-1} = -ug^{-1}$. This means that $g(t)$ makes u identically zero. We observe that $\exp\{-Bt/\zeta\}g(t)$ makes \tilde{u} identically zero and $\tilde{\eta}$ into B/ζ^2 . The initial value for the solution g^{-1} depends on ζ and so we can write $\tilde{g}(t) = U \exp\{-Bt/\zeta\}g(t)$ for some constant matrix U . If we are to get the form (5.5), we must have $U = U'd(\zeta)$, where

$$d(\zeta) = \text{diag} (\zeta^{-(n-1)}, \zeta^{-(n-3)}, \dots, \zeta^{n-1}). \tag{6.2}$$

In addition U' commutes with $B(\beta_1/\zeta^{-2}, \dots, \beta_n/\zeta^{-2})$. Moreover, the initial value for the equation $\frac{d}{dt}g^{-1} = -\alpha g^{-1}$ depends only on the residues of $u, \eta, \tilde{u}, \tilde{\eta}$ and therefore U' does not depend on B . Since the initial values belong to $SU(n)$, we also have $U' \in SU(n)$. It follows that U' belongs to the center of $SU(n)$. This is only an ambiguity in the choice of trivialization and it does not affect the twistor space. Similar considerations show that the real structure sends $B(\beta_1, \dots, \beta_n)$ to $B(-\bar{\beta}_1/\bar{\zeta}^{-2}, \dots, -\bar{\beta}_n/\bar{\zeta}^{-2})$ and g to $r(\zeta) \exp\{B^*/\bar{\zeta}\} (g^*)^{-1}$, where

$$r_{ij}(\zeta) = \begin{cases} 0 & \text{if } i + j \neq n \\ (-1)^{j-1} \bar{\zeta}^{n+1-2j} & \text{if } i + j = n \end{cases}.$$

This time the remaining ambiguity is given by a real element in the center of $SU(n)$, i.e. -1 if n is even.

We now go through a similar procedure for the subset of $M_{U(n)}(c - 1)$, where $\beta(+\infty)$ is regular. We have assigned in the proof of Proposition 5.3 to each element of this set a pair $(g, \beta(+\infty))$. We already know how $\beta(+\infty)$ changes (as it is given by the complex moment map for a torus action). The proof of Proposition 5.3 shows that the other coordinates, g on $\{\zeta \neq \infty\}$ and \tilde{g} on $\{\zeta \neq 0\}$, are related by $\tilde{g} = g \exp\{-(c - 1)\beta(+\infty)/\zeta\}$. The real structure sends g to $(g^*)^{-1} \exp\{(c - 1)\beta(+\infty)^*/\bar{\zeta}\}$.

Finally we have to go to the complex-symplectic quotient as in the proof of Proposition 5.3. We end up with (g, β_d) and $(\tilde{g}, \tilde{\beta}_d)$, where $\beta_d = \text{diag}(\beta_1, \dots, \beta_n)$ and $g\beta_d g^{-1} = B(\beta_1, \dots, \beta_n)$ (and similarly for $(\tilde{g}, \tilde{\beta}_d)$). We see that β_i and $\tilde{\beta}_i$ are related as stated and $\tilde{g} = d(\zeta) \exp\{-B/\zeta\} g \exp\{-(c - 1)\beta_d/\zeta\}$. Since $\exp\{-B/\zeta\} g = g \exp\{-\beta_d/\zeta\}$, $\tilde{g} = d(\zeta) g \exp\{-c\beta_d/\zeta\}$. If we now go to the coordinates u_i, \tilde{u}_i defined by (5.6), we see that they change as required, since the (i, j) th entry of V^{-1} is given by (5.9) and the β_i change as prescribed (i.e. as sections of $O(2)$). A similar argument shows that the real structure is, up to a sign, the one described in the statement (it is enough to compare the last row in $r(\zeta)(V^{-1} \text{diag}\{u_i\})^{*-1} \text{diag}\{e^{c\beta_i/\zeta}\}$ and in $V^{-1}(-\tilde{\beta}_1/\bar{\zeta}^{-2}, \dots, -\tilde{\beta}_n/\bar{\zeta}^{-2}) \text{diag}\{u'_i\}$). We shall see shortly (Proposition 6.2) that the negative of the real structure described in the statement does not admit any sections (a section would be equivalent to a complex number with imaginary modulus). The formula for the complex symplectic structure is a direct consequence of Proposition 5.4. \square

We now wish to find the full twistor space and the metric on $\tilde{M}_n(c)$ and this means finding a family of real sections. We know their projections to $O(2) \otimes \mathbb{C}^n$: they are given by $(\beta + (\alpha + \alpha^*)\zeta - \beta^*)(+\infty)$ (cf. [18]) and are parameterized by n distinct points in \mathbb{R}^3 with coordinates $(x_i, \text{Re } z_i, \text{Im } z_i), i = 1, \dots, n$, where $x_i = \sqrt{-1}T_1(+\infty)$, $z_i = \beta(+\infty)$. In other words we have n curves $S_i = \{(\zeta, \eta); \eta = z_i + 2x_i\zeta - \bar{z}_i\zeta^2\}$ in $T\mathbb{C}P^1$ (here η is the fiber coordinate). According to Proposition 6.1 the u_i coordinate of a real section of $Z(c)$ changes as a non-zero section of the bundle $L^c(k - 1)$ (with the transition function $\zeta^{k-1}e^{c\eta/\zeta}$ from ∞ to 0) over S_i . This is true only away from the intersection points of the curves S_i and we have to understand what happens to the section at these points. Two curves $S_i = \{(\zeta, \eta); \eta = z_i + 2x_i\zeta - \bar{z}_i\zeta^2\}$ and $S_j = \{(\zeta, \eta); \eta = z_j + 2x_j\zeta - \bar{z}_j\zeta^2\}$ intersect in a pair of distinct points a_{ij} and a_{ji} , where

$$a_{ij} = \frac{(x_i - x_j) + r_{ij}}{\bar{z}_i - \bar{z}_j}, \quad r_{ij} = \sqrt{(x_i - x_j)^2 + |z_i - z_j|^2}. \tag{6.3}$$

We have:

Proposition 6.2. *The real sections of the twistor space $Z(c)$ of $\tilde{M}_n(c)$ are given, over $\zeta \neq \infty$, by $(\beta_1(\zeta), \dots, \beta_n(\zeta), u_1(\zeta), \dots, u_n(\zeta))$, where*

$$\begin{aligned} \beta_i(\zeta) &= z_i + 2x_i\zeta - \bar{z}_i\zeta^2, \\ u_i(\zeta) &= A_i \prod_{j \neq i} (\zeta - a_{ji}) e^{c(x_i - \bar{z}_i\zeta)}, \end{aligned}$$

where (x_i, z_i) , $i = 1, \dots, n$, are distinct points in $\mathbb{R} \times \mathbb{C}$ and A_i are complex numbers satisfying

$$A_i \bar{A}_i = \prod_{j \neq i} (x_i - x_j + r_{ij}).$$

Remark. Given Proposition 5.2, this finally shows that the biholomorphism of Proposition 5.3 is onto.

Proof. Consider a real section s of $Z(c)$ (corresponding to a solution (T_0, T_1, T_2, T_3)) which projects to a given real section $(\beta_1(\zeta), \dots, \beta_n(\zeta))$ of $O(2) \otimes \mathbb{C}^n$. For a generic section the intersection points of the β_s are all distinct. We consider the point a_{ji} at which β_i intersects β_j and let us assume that no other β_s intersect there. We recall that $\sqrt{-1}T_1(\zeta) = \frac{1}{2}(\alpha + \alpha^*) - \beta^* \zeta$ and, hence, $\sqrt{-1}T_1(\zeta)_{(+\infty)_{ss}} = x_s - \bar{z}_s \zeta$. This means that $\sqrt{-1}T_1(a_{ji})_{(+\infty)_{jj}} < \sqrt{-1}T_1(a_{ji})_{(+\infty)_{ii}}$, and so, with respect to the complex structure corresponding to $a_{ji} \in \mathbb{C}P^1$, the solution (T_0, T_1, T_2, T_3) belongs to the chart described in Proposition 5.3 with \mathfrak{n} generated by the matrix with the only non-zero entry having coordinates (i, j) . Let us write s as $(\beta_i(\zeta), u_i(\zeta))$, $i = 1, \dots, n$, in a neighbourhood of a_{ji} , $\zeta \neq a_{ji}$ (notice that the procedure of Proposition 6.1 does assign well-defined complex numbers $u_1(\zeta), \dots, u_n(\zeta)$ to each $\zeta \neq a_{ji}$). According to the proof of Proposition 5.3 there is an element $m(\zeta) \in N = \exp \mathfrak{n}$ such that the following expression

$$V(\beta_1(\zeta), \dots, \beta_n(\zeta))^{-1} \text{diag}(u_1(\zeta), \dots, u_n(\zeta))m(\zeta)$$

has an invertible limit at $\zeta = a_{ji}$.

Let $W_{kl}(\zeta)$ denote the (k, l) th entry of $V(\beta_1(\zeta), \dots, \beta_n(\zeta))^{-1}$ and let $p(\zeta)$ denote the only non-zero non-diagonal entry of $\text{diag}(u_1(\zeta), \dots, u_n(\zeta))m(\zeta)$ ($p(\zeta)$ is the (i, j) th entry). We then have that $W_{kj}u_j + W_{ki}p$ and $W_{ki}u_i$ have a finite limit at $\zeta = a_{ji}$, for all $k = 1, \dots, n$. From the formula (5.9) a finite limit for $W_{ni}u_i$ implies that $u_i(a_{ji}) = 0$, while the nonvanishing of the last row of $V^{-1} \text{diag}(u_s)m$ means that a_{ji} is a single zero of u_i . If more than two sections $\beta_s(\zeta)$ meet at a_{ji} the considerations are similar but involve larger n . We can conclude the a_{ji} contribute precisely $n - 1$ zeros of u_i (counting multiplicities) and, given Proposition 6.1, this proves the formula for $u_i(\zeta)$ as soon as we show that u_i has no other zeros, or, equivalently, no poles. To prove this latter statement it is enough to show that u_j does not have a pole at a_{ji} . We go back to the situation when \mathfrak{n} is one-dimensional, and where we concluded that $W_{kj}u_j + W_{ki}p$ has a finite limit at $\zeta = a_{ji}$ for all $k = 1, \dots, n$. We can write $W_{nj}u_j + W_{ni}p$ as $(fu_j + gp)/(\beta_i - \beta_j)$ where f and g have finite limits at $\zeta = a_{ji}$. We then have

$$W_{n-1,j}u_j + W_{n-1,i}p = - \left(fu_j \left(\sum_{s \neq j} \beta_s \right) + gp \left(\sum_{s \neq i} \beta_s \right) \right) \frac{1}{\beta_i - \beta_j},$$

which can be rewritten as

$$-fu_j - \left(\sum_{s \neq i} \beta_s \right) \frac{fu_j + gp}{\beta_i - \beta_j}.$$

Since the second term has a finite limit, so does fu_j and hence u_j . Again, if more than two sections $\beta_s(\zeta)$ meet at a_{ji} the considerations are similar but involve larger n . Thus

we have shown the second formula of the statement. The last formula follows from the reality condition and the fact that $a_{ji} = -1/\bar{a}_{ij}$ (this calculation also eliminates the ± 1 ambiguity in the choice of the real structure in the proof of 6.1). \square

We can finally identify $\tilde{M}_n(c)$ as a T^n -bundle over the configuration space $\tilde{C}_n(\mathbb{R}^3)$ of n distinct points \mathbf{x}_i in \mathbb{R}^3 .

Proposition 6.3. $\tilde{M}_n(c)$ is equivalent to the T^n -bundle described in Proposition 1.1.

Proof. From the last formula in Proposition 6.2 it follows that $A_i \neq 0$ if, for all $j \neq i$, $z_i \neq z_j$ or $x_i > x_j$. On the other hand, if we put

$$A_I = A_i \prod_{j \in I} a_{ji},$$

for any subset I of $\{j; j \neq i\}$, then we have

$$A_I \bar{A}_I = \prod_{\substack{j \neq i \\ j \notin I}} (x_i - x_j + r_{ij}) \prod_{j \in I} (x_j - x_i + r_{ij}).$$

Let us choose sets I_1, \dots, I_n such that $I_i \subset \{j; j \neq i\}$ and $j \in I_i \Leftrightarrow i \notin I_j$. Define $U(I_1, \dots, I_n)$ as the complement of the subset $\{(x_i, z_i)_{i=1, \dots, n}; I_i^c = \{j; z_i = z_j \text{ and } x_i < x_j\}\}$ (I_i^c denotes the complement of I_i in $\{j; j \neq i\}$). The sets $U(I_1, \dots, I_n)$ cover $\tilde{C}_n(\mathbb{R}^3)$ and over each of them the bundle $\tilde{M}_n(c)$ is trivialized by coordinates $(x_i, z_i, A_{I_i}/|A_{I_i}|)$. To determine the bundle, choose $i < j$. The bundle restricted to S^2_{ij} is given by the transition function from $U(I_1, \dots, I_n)$, where $j \notin I(i)$ to $U(I'_1, \dots, I'_n)$, where $I'_i = I_i \cup \{j\}$, $I'_j = I_j - \{i\}$, $I'_k = I_k$ for $k \neq i, j$. Let ϕ_k be the transition function for the k^{th} generator of T^n , i.e. the transition function from $A_{I_k}/|A_{I_k}|$ to $A_{I'_k}/|A_{I'_k}|$. We see that $\phi_k = 1$ if $k \neq i, j$, and $\phi_i = a_{ji}/|a_{ji}|$, $\phi_j = |a_{ji}|/a_{ji}$. Therefore $\phi_i = (z_j - z_i)/|z_j - z_i|$ and $\phi_j = \phi_i^{-1}$. It remains to identify the circle bundle over the sphere $x^2 + |z|^2 = \text{const}$ given by the transition function $z/|z|$ from the region $U_0 = \{z \neq 0 \text{ or } x > 0\}$ to the region $U_1 = \{z \neq 0 \text{ or } x < 0\}$. Let us write the unit 3-sphere as $\{(u, v) \in \mathbb{C}^2; |u|^2 + |v|^2 = 1\}$. The Hopf bundle is given the S^1 action $t \cdot (u, v) = (tu, t^{-1}v)$ and the projection $S^3 \rightarrow S^2$ by the map $x = |u|^2 - |v|^2$, $z = 2uv$. Over U_0 this bundle is trivialized by $(x, z, u/|u|)$ and over U_1 by $(x, z, |v|/v)$. The transition function is $|z|/z$. Thus $[\phi_i] = -1 \in H^1(S^2_{ij}, S^1)$. \square

We can now calculate the metric on $\tilde{M}_n(c)$. By the remark at the end of Sect. 3, it is enough to know the metric for $c = -1, 0, 1$, as the others are obtained by homothety. We shall calculate the metric for $c = 1$. The metric for $c = -1$ is the everywhere negative definite version of the Gibbons–Manton metric (this can be seen from the $c = 1$ calculation) and the one for $c = 0$ the and negative-definite cone over a 3-Sasakian manifold.

Theorem 6.4. $\tilde{M}_n(1)$ is isomorphic, as a hyperkähler manifold, to the Gibbons–Manton manifold M_{GM} defined in Sect. 1.

Proof. We know from the previous proposition that the two spaces are diffeomorphic. We shall show that the twistor description of $\tilde{M}_n(1)$ and of the Gibbons–Manton metric coincide. We recall from Sect. 1 that the latter is a hyperkähler quotient of $M = M_1 \times M_2$ by a torus, where $M_1 = (S^1 \times \mathbb{R}^3)^n$ and $M_2 = \mathbb{H}^{n(n-1)/2}$. With respect to any

complex structure $M_1 = (\mathbb{C}^*)^n \times \mathbb{C}^n$ and $M_2 = \mathbb{C}^{n(n-1)/2} \times \mathbb{C}^{n(n-1)/2}$. Let us write the corresponding complex coordinates as (p_i, β_i) , $i = 1, \dots, n$, on M_1 and as (v_{ij}, w_{ij}) , $i < j$, on M_2 . The complex-symplectic forms corresponding to metrics g_1 and g_2 are given by

$$\sum_{i=1}^n \frac{dp_i}{p_i} \wedge d\beta_i, \quad (6.4)$$

$$\sum_{i < j} dv_{ij} \wedge dw_{ij}. \quad (6.5)$$

The real sections of the twistor space Z_1 of M_1 are written, over $\zeta \neq \infty$, as

$$\beta_i(\zeta) = z_i + 2x_i\zeta - \bar{z}_i\zeta^2, \quad p_i(\zeta) = B_i e^{x_i - \bar{z}_i\zeta}, \quad (6.6)$$

where $B_i \bar{B}_i = 1$. The real sections of the twistor space Z_2 of M_2 are (cf. [2], chapter 13.F):

$$v_{ij}(\zeta) = C_{ij}(\zeta - a_{ij}), \quad w_{ij}(\zeta) = D_{ij}(\zeta - a_{ji}), \quad (6.7)$$

where a_{ij}, a_{ji} are roots of $v_{ij}w_{ij} = z_{ij} + 2x_{ij}\zeta - \bar{z}_{ij}\zeta^2$ for some $(x_{ij}, z_{ij}) \in \mathbb{R} \times \mathbb{C}$, i.e.

$$a_{ij} = \frac{x_{ij} + \sqrt{x_{ij}^2 + |z_{ij}|^2}}{\bar{z}_{ij}}, \quad a_{ji} = \frac{x_{ij} - \sqrt{x_{ij}^2 + |z_{ij}|^2}}{\bar{z}_{ij}}$$

and

$$C_{ij}\bar{C}_{ij} = -x_{ij} + \sqrt{x_{ij}^2 + |z_{ij}|^2}, \quad D_{ij}\bar{D}_{ij} = x_{ij} + \sqrt{x_{ij}^2 + |z_{ij}|^2}.$$

Here the particular choice of sections is forced either by the fact the metric is positive definite or by requiring that the S^1 -action $t \cdot (v_{ij}, w_{ij}) = (tv_{ij}, t^{-1}w_{ij})$ determines the Hopf bundle over the 2-sphere $x_{ij}^2 + |z_{ij}|^2 = 1$ (this calculation was done in the proof of Proposition 6.3). To obtain the twistor description of the Gibbons–Manton metric we have to perform the complex-symplectic quotient construction along the fibers of $Z_1 \oplus Z_2$ with respect to the difference of the forms (6.4) and (6.5). As in Sect. 1, the moment map equations are $v_{ij}w_{ij} = \beta_i - \beta_j$ and so the a_{ij}, a_{ji} are given by (6.3). Since we already know that the manifolds are diffeomorphic, it is sufficient to determine the metric on an open dense subset, e.g. on the set where all v_{ij} are non-zero. Quotienting this set by $(\mathbb{C}^*)^{n(n-1)/2}$ is equivalent to sending all v_{ij} to 1. This is achieved by acting by the element $(v_{ij})^{-1}$ of $(\mathbb{C}^*)^{n(n-1)/2}$. By the description of the torus action given in Sect. 1, this sends $p_i(\zeta)$ to

$$B_i \frac{\prod_{j < i} C_{ji}(\zeta - a_{ji})}{\prod_{j > i} C_{ij}(\zeta - a_{ij})} e^{x_i - \bar{z}_i\zeta} = E_i \frac{\prod_{j < i} (\zeta - a_{ji})}{\prod_{j > i} (\zeta - a_{ij})} e^{x_i - \bar{z}_i\zeta}, \quad (6.8)$$

where

$$E_i \bar{E}_i = \frac{\prod_{j < i} (x_i - x_j + r_{ij})}{\prod_{j > i} (x_j - x_i + r_{ij})}. \quad (6.9)$$

These and the β_i give the real sections for the Gibbons–Manton metric and the symplectic form is (6.4). We now compare this with the description of $Z(1)$ given in Proposition

6.2. According to Remark 5.5 we should set $p_i = u_i / \prod_{j>i}(\beta_i - \beta_j)$ in order to have the same symplectic form. We obtain

$$p_i(\zeta) = \frac{A_i}{\prod_{j>i}(\bar{z}_j - \bar{z}_i)} \frac{\prod_{j<i}(\zeta - a_{ji})}{\prod_{j>i}(\zeta - a_{ij})} e^{x_i - \bar{z}_i \zeta}.$$

All we have to do is to compare the norm of $A_i / \prod_{j>i}(\bar{z}_j - \bar{z}_i)$ with the norm of E_i . We have, from Proposition 6.2 and Eq. (6.9),

$$\begin{aligned} \frac{A_i \bar{A}_i}{\prod_{j>i} |z_j - z_i|^2} &= \frac{\prod_{j \neq i} (x_i - x_j + r_{ij})}{\prod_{j>i} |z_j - z_i|^2} = \prod_{j<i} (x_i - x_j + r_{ij}) \prod_{j>i} \frac{(x_i - x_j + r_{ij})}{|z_j - z_i|^2} \\ &= \frac{\prod_{j<i} (x_i - x_j + r_{ij})}{\prod_{j>i} (x_j - x_i + r_{ij})} = E_i \bar{E}_i, \end{aligned}$$

which proves the theorem. \square

We shall finish the section with a remark that Propositions 6.2 and 6.3 can be generalized to define hyperkähler metrics on a class of T^n -bundles over $\check{C}_n(\mathbb{R}^3)$. We have:

Theorem 6.5. *Let P be a T^n -bundle over $\check{C}_n(\mathbb{R}^3)$ determined by an element (s_1, \dots, s_n) of $H^2(\check{C}_n(\mathbb{R}^3), \mathbb{Z}^n)$ satisfying $s_k(S_{ij}^2) = 0$ if $k \neq i, j$ and $s_i(S_{ij}^2) = -s_j(S_{ij}^2)$. Then P carries a family of (pseudo)-hyperkähler metrics such that the real sections of the twistor space are given, over $\zeta \neq \infty$, by $(\beta_1(\zeta), \dots, \beta_n(\zeta), u_1(\zeta), \dots, u_n(\zeta))$, where*

$$\begin{aligned} \beta_i(\zeta) &= z_i + 2x_i \zeta - \bar{z}_i \zeta^2, \\ u_i(\zeta) &= A_i \prod_{j \neq i} (\zeta - a_{ji})^{s_{ij}} e^{c(x_i - \bar{z}_i \zeta)}, \end{aligned}$$

where c is a real constant, (x_i, z_i) , $i = 1, \dots, n$, are distinct points in $\mathbb{R} \times \mathbb{C}$, $s_{ij} = |s_i(S_{ij}^2)|$, and A_i are complex numbers satisfying

$$A_i \bar{A}_i = \prod_{j \neq i} (x_i - x_j + r_{ij})^{s_{ij}}.$$

This description determines a hypercomplex structure on P . A (pseudo)-hyperkähler metric can be then calculated using any complex-symplectic form along the fibers, given as a section of $\Lambda^2 T_F^* \otimes O(2)$, e.g. the form (5.7). These metrics will correspond to the motion of n dyons in \mathbb{R}^3 interacting in different ways (cf. [14]).

Remark. The calculation of the metric given above shows that the Taub-NUT metric (cf. [2]) has two very different descriptions in terms of Nahm’s equations: 1) it is the metric on the totally geodesic submanifold $\tilde{M}_2^0(-1)$ of $\tilde{M}_2(-1)$ defined by considering $\mathfrak{su}(2)$ -valued solutions to Nahm’s equations and $SU(2)$ -valued gauge transformations; 2) it is the metric on the moduli space of $SU(3)$ -monopoles of charge $(1, 1)$ [10, 29].

7. Asymptotic Comparison of the Metrics

We shall now show that the Gibbons–Manton metric and the monopole metric are asymptotically exponentially close. The asymptotic region, where the individual monopoles are separated, of the monopole space M_n is diffeomorphic to P/S_n , where P is a torus bundle over the configuration space $\check{C}_n(\mathbb{R}^3)$ and S_n the symmetric group. The bundle P is not, however, the bundle of Proposition 6.3. Rather, as we shall see shortly, it is the quotient of that bundle by a $(\mathbb{Z}_2)^n$ -subgroup of T^n . In other words it is the bundle determined by an $s \in H^2(\check{C}_n(\mathbb{R}^3), \mathbb{Z}^n)$ with all s_k being twice those in Proposition 6.3.

We shall compare the metric on M_n with the metric on the hyperkähler quotient of $\tilde{M}_n(1) \times \tilde{M}_n(1)$ by the diagonal T^n -action. We do this in order to have solutions to Nahm’s equations with poles at both ends of the interval $[-1, 1]$. For any c, c' , let us write $\tilde{M}_n(c, c')$ for the hyperkähler quotient of $\tilde{M}_n(c) \times \tilde{M}_n(c')$ by the diagonal action of T^n . The action of T^n given by $t \cdot (m, m') = (tm, m')$ induces a tri-Hamiltonian action of T^n on $\tilde{M}_n(c, c')$ which makes $\tilde{M}_n(c, c')$ into a T^n -bundle over $\check{C}_n(\mathbb{R}^3)$. We have

Lemma 7.1. *$\tilde{M}_n(c, c')$ is isomorphic, as a hyperkähler manifold, to $\tilde{M}_n(c+c')/(\mathbb{Z}_2)^n$, where $(\mathbb{Z}_2)^n = \{t \in T^n; t^2 = 1\}$.*

Proof. Let μ, μ' be the moment maps for the action of T^n on $\tilde{M}_n(c), \tilde{M}_n(c')$ respectively. The moment map for the diagonal T^n -action on the product is $\mu + \mu'$. If we go back to the proof of Proposition 6.3 and use the same notation, we can see that the zero-set of this moment map is a $(T^n \times T^n)$ -bundle over $\check{C}_n(\mathbb{R}^3)$ which restricted to each $S_{i_j}^2$ is given by transition functions $(\phi_1, \dots, \phi_n, \phi_1^{-1}, \dots, \phi_n^{-1})$ (the point being that $U(I'_1, \dots, I'_n) = -U(I_1, \dots, I_n)$). Hence, if we quotient by T^n , by sending the second T^n to 1 over each $U(I_1, \dots, I_n)$, we end up with a T^n -bundle for which the transition functions are $\phi_k^2, k = 1, \dots, n$. This proves the differential-geometric part of the statement. To obtain the isometry we repeat this argument for the twistor space of $\tilde{M}_n(c) \times \tilde{M}_n(c')$, performing the complex-symplectic quotient along the fibers as in the proof of Theorem 6.4. \square

From now on, we shall consider $\tilde{M}_n(1, 1)$ with *half* (compare formula (2.4)) of the metric given by the above lemma. In other words, locally the metric is still the Gibbons–Manton metric. We can identify $\tilde{M}_n(1, 1)$ with the moduli space of pairs $((T_0, T_1, T_2, T_3), (T'_0, T'_1, T'_2, T'_3))$ of solutions to Nahm’s equations, defined respectively on $[-1, \infty]$ and on $[-\infty, 1]$, such that $T_i(+\infty) = T'_i(-\infty)$ for $i = 0, 1, 2, 3$, and the residues of T_i at -1 and of T'_i at $+1, i = 1, 2, 3$, define the standard n -dimensional irreducible representation of $\mathfrak{su}(2)$. The group of gauge transformations $\mathcal{G}(1, 1)$ is now defined as pairs (g, g') such that $g(t+1), g'(-t+1) \in \mathcal{G}(c)$ for some c and $s = \lim_{t \rightarrow +\infty} \dot{g}g^{-1} = \lim_{t \rightarrow -\infty} \dot{g}'g'^{-1}$. The tangent space consists of pairs $((t_0, t_1, t_2, t_3), (t'_0, t'_1, t'_2, t'_3))$ defined on $[-1, \infty]$ and on $[-\infty, 1]$, respectively, with $t_i(+\infty) = t'_i(-\infty)$ and satisfying Eqs. (2.3). The metric on $\tilde{M}_n(1, 1)$ can be written as

$$\begin{aligned} \frac{1}{2} \sum_0^3 \|t_{i(+\infty)}\|^2 + \frac{1}{2} \int_{-1}^{+\infty} \sum_0^3 (\|t_i(s)\|^2 - \|t_{i(+\infty)}\|^2) ds + \\ \frac{1}{2} \sum_0^3 \|t'_{i(-\infty)}\|^2 + \frac{1}{2} \int_{-\infty}^1 \sum_0^3 (\|t'_i(s)\|^2 - \|t'_{i(-\infty)}\|^2) ds. \end{aligned}$$

We can rewrite this as

$$\begin{aligned} & \frac{1}{2} \int_0^{+\infty} \sum_0^3 (\|t_i(s)\|^2 - \|t_i(+\infty)\|^2) ds + \frac{1}{2} \int_{-\infty}^0 \sum_0^3 (\|t'_i(s)\|^2 - \|t_i(+\infty)\|^2) ds \\ & + \frac{1}{2} \int_{-1}^0 \sum_0^3 \|t_i(s)\|^2 ds + \frac{1}{2} \int_0^1 \sum_0^3 \|t'_i(s)\|^2 ds. \end{aligned} \tag{7.1}$$

Let us fix a complex structure, say I and write as in Sect. 5, α for $T_0 + iT_1$, β for $T_2 + iT_3$. We write an element of $\tilde{M}_n(1, 1)$ as a pair $((\alpha_-, \beta_-), (\alpha_+, \beta_+))$. We shall write β_i for the (i, i) th entry of $\beta_-(+\infty) = \beta_+(-\infty)$ and denote by $\tilde{M}_n^{\text{reg}}(1, 1)$ the subset of $\tilde{M}_n(1, 1)$ where all β_i are distinct. Similarly, we write M_n^{reg} for the subset of (α, β) in (M_n, I) where the eigenvalues of β are distinct. We shall prove:

Theorem 7.2. *There exists a biholomorphism ϕ from $\tilde{M}_n^{\text{reg}}(1, 1)/S_n$ to M_n^{reg} such that*

$$|\phi^*g - g'| = O(e^{-cR}), \tag{7.2}$$

where g, g' denote the monopole and Gibbons–Manton metric respectively, $c = c(n)$ is a constant, and R is the separation distance of particles in $C_n(\mathbb{R}^3)$, i.e.

$$R = \min\{|\mathbf{x}_i - \mathbf{x}_j|; i \neq j\}. \tag{7.3}$$

The same estimate holds for the Riemannian curvature tensor.

Since such a biholomorphism will be defined for any complex structure and the union of $\tilde{M}_n^{\text{reg}}(1, 1)$ for different complex structures is all of $\tilde{M}_n(1, 1)$, we conclude that the monopole and the Gibbons–Manton metrics are exponentially close in the asymptotic region of the monopole moduli space.

The remainder of the section is devoted to proving this theorem. We need the following lemma:

Lemma 7.3. *Let $C > 0$. The space $\tilde{M}_n^{\text{reg}}(1)$ is biholomorphic to the quotient of the space of solutions (α, β) to Eq. (5.1) which have the correct boundary behaviour at $t = 0$ and are constant (hence diagonal) for $t \geq C$ by the group of complex gauge transformations $g : [0, +\infty) \rightarrow Gl(n, \mathbb{C})$ with $g(0) = 1$ and $g(t) = \exp(ht - h)$ for some diagonal h for $t \geq C$.*

Proof. Let (α, β) be an element of $\tilde{M}_n^{\text{reg}}(1)$ and let $\alpha_d = \alpha(+\infty), \beta_d = \beta(+\infty)$. According to the proof of Proposition 5.3, there is a unique complex gauge transformation g defined on $[C/2, +\infty)$ with $g(+\infty) = 1$ such that $(\alpha, \beta) = g(\alpha_d, \beta_d)$. Let $\hat{g} : [C/2, \infty) \rightarrow Gl(n, \mathbb{C})$ be a smooth path with the values and the first derivatives of \hat{g} and g coinciding at $t = C/2$ and with $\hat{g}(t) = 1$ and for $t \geq C$. We obtain a solution $(\hat{\alpha}, \hat{\beta})$ to the complex Nahm equation (5.1) by setting

$$(\hat{\alpha}, \hat{\beta})(t) = \begin{cases} (\alpha, \beta)(t) & \text{if } t < C \\ \hat{g}(t)(\alpha_d, \beta_d) & \text{if } t \geq C. \end{cases} \tag{7.4}$$

This is a solution of the type described in the statement of this lemma. The proof of 5.3 shows further that it is only $g(C/2)\exp\{(1 - C/2)\alpha_d\}$ (and a solution to (5.1) on $[0, C/2)$) that determines the element of $\tilde{M}_n^{\text{reg}}(1)$. Therefore we obtain a well defined holomorphic map from $\tilde{M}_n^{\text{reg}}(1)$ to the moduli space described in the statement. Let us define the inverse map. Let $(\hat{\alpha}, \hat{\beta})$ be an element of the moduli space described in the

statement. As in [23] we can find a bounded complex gauge transformation g_0 such that $g_0(\hat{\alpha}, \hat{\beta})$ is an element of $\tilde{M}_n^{\text{reg}}(1)$.

We can assume that g_0 has a limit h at $+\infty$ (this follows from the convexity property of g_0 [13], since we can assume that $g_0(t)$ is hermitian for all t). According to Proposition 5.3 or 6.2 the action of T^n on $\tilde{M}_n(1)$ extends to a global action $(\mathbb{C}^*)^n$ with respect to the complex structure I (or any other). Let (α, β) be the element of $\tilde{M}_n^{\text{reg}}(1)$ obtained from $g_0(\hat{\alpha}, \hat{\beta})$ by the action of $h^{-1} \in (\mathbb{C}^*)^n$. Then $(\alpha, \beta) = g(\hat{\alpha}, \hat{\beta})$ and $g \in \mathcal{G}^{\mathbb{C}}(1)$. This gives the inverse mapping. \square

We can now construct a biholomorphism between $\tilde{M}_n^{\text{reg}}(1, 1)/S_n$ and M_n^{reg} . From the above lemma, $\tilde{M}_n^{\text{reg}}(1, 1)$ is biholomorphic to the quotient of the space of pairs $((\alpha_-, \beta_-), (\alpha_+, \beta_+))$ defined on $[-1, +\infty)$ and on $(-\infty, 1]$ respectively such that $(\alpha_-, \beta_-)(t + 1)$ and $(\alpha_+, \beta_+)(1 - t)$ are as in the above lemma, $(\alpha_-, \beta_-)(+\infty) = (\alpha_+, \beta_+)(-\infty)$ by the group of pairs (g_-, g_+) with $g_-(-1) = g_+(1) = 1$ and such that there are diagonal h, p with $g_-(t) = \exp(th - p)$ for $t > -r$, $g_+(t) = \exp(th - p)$ for $t < r$ ($r \in (0, 1)$ is fixed but arbitrary). We define a solution $(\hat{\alpha}, \hat{\beta})$ to the complex Nahm equation (5.1) on $(-1, 1)$ by

$$(\hat{\alpha}, \hat{\beta})(t) = \begin{cases} (\alpha_-, \beta_-)(t) & \text{if } t < 0 \\ (\alpha_+, \beta_+)(t) & \text{if } t \geq 0. \end{cases} \tag{7.5}$$

The $\mathcal{G}^{\mathbb{C}}$ -orbit of this solution (see Sect. 2 for the definition of \mathcal{G}) contains a unique element of M_n [13, 20]. Furthermore, the action of a (g_-, g_+) translates into the action of $g \in \mathcal{G}^{\mathbb{C}}$, where $g(t) = g_-(t)$ for $t < 0$ and $g(t) = g_+(t)$ for $t \geq 0$. Therefore we have a well defined holomorphic map ϕ_r from $\tilde{M}_n^{\text{reg}}(1, 1)$ to M_n . If we now have an element (α, β) of M_n^{reg} , we can diagonalize β on $[-r, r]$ and make α diagonal and constant on $[-r, r]$. Let $(\tilde{\alpha}, \tilde{\beta})$ be the resulting solution to the complex Nahm equation. We obtain an element of $\tilde{M}_n^{\text{reg}}(1, 1)$ by setting

$$(\alpha_-, \beta_-)(t) = \begin{cases} (\tilde{\alpha}, \tilde{\beta})(t) & \text{for } t < 0 \\ (\tilde{\alpha}, \tilde{\beta})(0) & \text{for } t \geq 0 \end{cases}$$

and similarly for (α_+, β_+) . This defines the inverse to ϕ_r up to the ordering of eigenvalues of β . In other words ϕ_r induces a biholomorphism between $\tilde{M}_n^{\text{reg}}(1, 1)/S_n$ and M_n^{reg} . Furthermore, for a fixed element $((\alpha_-, \beta_-), (\alpha_+, \beta_+))$ of $\tilde{M}_n^{\text{reg}}(1, 1)$ and two parameters r, r' , the resulting $(\hat{\alpha}, \hat{\beta})$ of (7.5) are $\mathcal{G}^{\mathbb{C}}$ -equivalent and therefore $\phi_r, \phi_{r'}$ induce the same biholomorphism ϕ .

Let us now prove the estimate (7.3). Fortunately, much of the analysis has been already done in [3]. First of all, we recall ([23], Lemma 3.4) that solutions to Nahm's equations which have a regular triple as a limit at infinity, approach this limit exponentially fast, of order $O(e^{-cR})$ (that is T_1, T_2, T_3 do and we can always make T_0 to have such decay by using the gauge freedom). The proofs of Propositions 3.11–3.14 in [3] show that the same holds for tangent vectors (t_0, t_1, t_2, t_3) . Let us now see what happens to a tangent vector v under the map ϕ . The gauge transformations (g, g') which make the element $((\alpha_-, \beta_-), (\alpha_+, \beta_+))$ of $\tilde{M}_n^{\text{reg}}(1, 1)$ constant and equal to the common value at infinity on $[-1 + C/2, +\infty)$ and $(-\infty, 1 - C/2]$ are exponentially close to the identity. In the next stage of the construction of ϕ – formula (7.4) – we have smoothed out the solutions which can be again done by gauge transformations exponentially close to 1.

Therefore the resulting tangent vector \hat{v} is exponentially close to the original one in the metric (7.1). We have then restricted the solutions (formula (7.5)) to obtain a solution $(\hat{\alpha}, \hat{\beta})$ to the complex Nahm equation on $[-1, 1]$. Let p denote this operation of restriction. The first line of the formula (7.1) is exponentially small and therefore the norm of \hat{v} in (7.1) and the norm of $dp(\hat{v})$ in (2.4) are exponentially close. The solution $(\hat{\alpha}, \hat{\beta})$ will not satisfy the real Nahm equation, however, we will have

$$F(\hat{\alpha}, \hat{\beta}) := \frac{d}{dt} (\hat{\alpha} + \hat{\alpha}^*) + [\hat{\alpha}, \hat{\alpha}^*] + [\hat{\beta}, \hat{\beta}^*] = O(e^{-cR}).$$

Lemma 2.10 in [13] implies now that we can solve the real equation by a complex gauge transformation bounded as $O(e^{-cR})$. We can now show that the vector $d\phi(v)$ tangent to M_n (which is obtained from $dp(\hat{v})$) is exponentially close to $dp(\hat{v})$ by following the analysis of Sect. 3 in [3] step by step, replacing the $O(1/R)$ estimates by $O(e^{-cR})$. This proves the estimate (7.3). For the curvature estimates we do the same using the analysis of Sect. 4 in [3]. This proves Theorem 7.2.

8. Twistor Description of Monopoles and the Gibbons–Manton Metric

We shall show in this section how the twistor description of monopole metrics determines the asymptotic metric. We recall [13] that the moduli space of n -monopoles is biholomorphic to the space of based rational maps $p(z)/q(z)$ on $\mathbb{C}P^1$ of degree n (based means that $\deg p < \deg q$). On the set, where the roots β_1, \dots, β_n of $q(z)$ are distinct, these roots and the values $p_i = p(\beta_i)$ of p form local coordinates and the complex-symplectic form can be written as [1]:

$$\sum_{i=1}^n d\beta_i \wedge \frac{dp_i}{p_i}. \tag{8.1}$$

The metric is determined by the real sections $p(z, \zeta)/q(z, \zeta)$. Their description is provided in [19]. The denominator $q(z, \zeta)$ is given by a curve S – the spectral curve of the monopole – in $T\mathbb{C}P^1$ [16]. This curve satisfies several conditions, one of which is the triviality of the line bundle L^{-2} restricted to S , and Hurtubise [19] shows that the numerator $p(z, \zeta)$ is given by a nonzero section of this bundle. (The values $p_i(\zeta)$ are given by the values of this section at the intersection points $\beta_i(\zeta)$ of S with $T_\zeta\mathbb{C}P^1$.)

What happens when the individual monopoles separate? First of all, the spectral curve approaches the union of spectral curves of individual monopoles exponentially fast [4]. These curves S_i are of the form $\eta_i = z_i + 2x_i\zeta - \bar{z}_i\zeta^2$, $i = 1, \dots, n$, where $(x_i, \text{Re } z_i, \text{Im } z_i)$ are locations of 1-monopoles (particles). What happens to the section of L^{-2} ? We make a heuristic assumption (which we know to be true from Sect. 6) that the section acquires zeros and poles at the intersection points of the S_i (more precisely the only singularities of $p_i(\zeta)$ occur at the intersection points of S_i with other S_j). As we shall see this is sufficient to determine the asymptotic metric.

First of all the real structure on the bundle L^{-2} is $u \mapsto \bar{u}^{-1}e^{-2\bar{\eta}/\zeta}$ and therefore if p_i has a zero at one of the points of $S_i \cap S_j$, then it has a pole of the same order at the other, and vice versa. Furthermore, since the metric and hence the real sections are invariant under the action of the symmetric group, we must have

$$p_i(\zeta) = A_i \prod_{j \neq i} \left(\frac{\zeta - a_{ij}}{\zeta - a_{ji}} \right)^k e^{-2(x_i - \bar{z}_i\zeta)}, \quad i = 1, \dots, n,$$

where a_{ij}, a_{ji} are the two points in $S_i \cap S_j$ given by (6.3) and k is an integer. The reality condition implies that

$$A_i \bar{A}_i = \prod_{j \neq i} a_{ji}^k \bar{a}_{ji}^k.$$

One can now calculate the asymptotic metric, using (8.1). The sign of k will determine the signature, while $|k|$ is simply a constant multiple. The actual value of k is determined by the topology of the asymptotic region of M_n , and comparing with Proposition 6.3 and the remarks at the beginning of Sect. 7 we conclude that $k = 1$ (in the coordinates of Proposition 6.2, $p_i = \prod_{j:j \neq i} (\beta_i - \beta_j)/u_i^2$).

We remark that the above analysis can be easily done for other compact Lie groups G . The twistor description of metrics on moduli spaces of G -monopoles with maximal symmetry breaking is known from the work of Murray [28] and Hurtubise and Murray [21, 22] and from this the asymptotic metric can be calculated. We shall do the exact analysis in the case of $G = SU(N)$ in a subsequent paper.

Acknowledgement. This work was carried out at the Max-Planck-Institut für Mathematik in Bonn. I wish to thank the Institute's directors and staff for their hospitality and for creating a stimulating research atmosphere. I also thank Michael Murray for comments.

Note added in proof

The metrics of section 4 (or, at least, their positive-definite counterparts) can be also obtained as finite-dimensional hyperkähler quotients. This follows from the construction of Kobak and Swann (Internat. J. Math. **7**, 193–210 (1996)) of a nilpotent $G^{\mathbb{C}}$ -orbit as a hyperkähler quotient of a vector space V by a product U of unitary groups. The hyperkähler quotient P of V by the semisimple part of U is a positive-definite analogue of the spaces $M_G(c)$ (i.e. Theorem 4.3 holds for P). Presumably the spaces $M_G(c)$ can also be obtained this way by changing the signature of the metric on V .

References

1. Atiyah, M.F. and Hitchin, N.J.: *The geometry and dynamics of magnetic monopoles*. Princeton: Princeton University Press, 1988
2. Besse, A.: *Einstein manifolds*. Berlin–Heidelberg–New York: Springer, 1987
3. Bielawski, R.: Asymptotic behaviour of $SU(2)$ monopole metrics. *J. reine angew. Math.* **468**, 139–165 (1995)
4. Bielawski, R.: Monopoles, particles and rational functions. *Ann. Glob. Anal. Geom.* **14**, 123–145 (1996)
5. Bielawski, R.: On the hyperkähler metrics associated to singularities of nilpotent varieties. *Ann. Glob. Anal. Geom.* **14**, 177–191 (1996)
6. Bielawski, R.: Hyperkähler structures and group actions. *J. London Math. Soc.* **55**, 400–414 (1997)
7. Bielawski, R.: Invariant hyperkähler metrics with a homogeneous complex structure. *Math. Proc. Cam. Phil. Soc.* **122**, 473–482 (1997)
8. Biquard, O.: Sur les équations de Nahm et les orbites coadjointes des groupes de Lie semi-simples complexes. *Math. Ann.* **304**, 253–276 (1996)
9. Biquard, O.: *Twistors des orbites coadjointes*. Ecole Polytechnique preprint (1997)
10. Connell, S.: The dynamics of the $SU(3)(1, 1)$ magnetic monopoles. Ph.D. thesis, The Flinders University of South Australia (1991)
11. Dancer, A.S.: Nahm's equations and hyperkähler geometry. *Commun. Math. Phys.* **158**, 545–568 (1993)
12. Dancer, A.S.: A family of hyperkähler manifolds. *Quart. J. Math. Oxford* **45**, 463–478 (1994)

13. Donaldson, S.K.: Nahm's equations and the classification of monopoles. *Commun. Math. Phys.* **96**, 387–407 (1984)
14. Gibbons, G.W. and Manton, N.S.: The moduli space metric for well-separated BPS monopoles. *Phys. Lett. B* **356**, 32–38 (1995)
15. Gibbons, G.W. and Rychenkova, P.: HyperKähler quotient construction of BPS monopole moduli spaces. *Commun. Math. Phys.* **186** (1997), 581–599
16. Hitchin, N.J.: On the construction of monopoles. *Commun. Math. Phys.* **89**, 145–190 (1983)
17. Hitchin, N.J.: Polygons and gravitons. *Math. Proc. Camb. Phil. Soc.* **83**, 465–476 (1979)
18. Hitchin, N.J., Karlhede, A., Lindström, U. and Roček, M.: Hyperkähler metrics and supersymmetry. *Commun. Math. Phys.* **108**, 535–586 (1985)
19. Hurtubise, J.C.: Monopoles and rational maps: a note on a theorem of Donaldson. *Commun. Math. Phys.* **100**, 191–196 (1985)
20. Hurtubise, J.C.: The classification of monopoles for the classical groups. *Commun. Math. Phys.* **120**, 613–641 (1989)
21. Hurtubise, J.C. and Murray, M.K.: On the construction of monopoles for the classical groups. *Commun. Math. Phys.* **122**, 35–89 (1989)
22. Hurtubise, J.C. and Murray, M.K.: Monopoles and their spectral data. *Commun. Math. Phys.* **133**, 487–508 (1990)
23. Kronheimer, P.B.: A hyper-kählerian structure on coadjoint orbits of a semisimple complex group. *J. London Math. Soc.* **42**, 193–208 (1990)
24. Kronheimer, P.B.: The construction of ALE spaces as hyper-Kähler quotients. *J. Diff. Geom.* **29**, 665–683 (1989)
25. Lindström, U. and Roček, M.: Scalar tensor duality and $N = 1, 2$ nonlinear σ -models. *Nucl. Phys.* **222B**, 285–308 (1983)
26. Manton, N.S.: A remark on the scattering of BPS monopoles. *Phys. Lett. B* **110**, 54–56 (1982)
27. Manton, N.S.: Monopole interactions at long range, *Phys. Lett. B* **154**, 397–400 (1985)
28. Murray, M.K.: Non-abelian magnetic monopoles. *Commun. Math. Phys.* **96**, 539–565 (1984)
29. Murray, M.K.: A note on the $(1, 1, \dots, 1)$ monopole metric. *J. Geom. Phys.* **23**, 31–41 (1997)
30. Nahm, W.: The construction of all self-dual monopoles by the ADHM method. In *Monopoles in quantum field theory*. Singapore: World Scientific, 1982
31. Nakajima, H.: Monopoles and Nahm's equations. In *Einstein metrics and Yang-Mills connections* New York: Marcel Dekker, 1993
32. Pedersen, H. and Poon, Y.S.: Hyper-Kähler metrics and a generalization of the Bogomolny equations. *Comm. Math. Phys.* **117**, 569–580 (1988)
33. Stuart, D.: The geodesic approximation for the Yang-Mills-Higgs equations. *Commun. Math. Phys.* **166**, 149–190 (1994)
34. Swann, A.: Hyperkähler and quaternionic Kähler geometry. *Math. Ann.* **289**, 421–450 (1991)

Communicated by H. Nicolai