

Monodromy and vanishing cycles in toric surfaces

Nick Salter¹

Received: 23 October 2017 / Accepted: 5 December 2018 /
Published online: 12 December 2018
© Springer-Verlag GmbH Germany, part of Springer Nature 2018

Abstract Given an ample line bundle on a toric surface, a question of Donaldson asks which simple closed curves can be vanishing cycles for nodal degenerations of smooth curves in the complete linear system. This paper provides a complete answer. This is accomplished by reformulating the problem in terms of the mapping class group-valued monodromy of the linear system, and giving a precise determination of this monodromy group.

1 Introduction

Let X be a smooth toric surface and \mathcal{L} an ample line bundle on X . In the complete linear system $|\mathcal{L}|$, there is a hypersurface \mathcal{D} known as the *discriminant locus* consisting of the singular curves $C \in |\mathcal{L}|$. The complement

$$\mathcal{M}(\mathcal{L}) := |\mathcal{L}| \setminus \mathcal{D}$$

therefore supports a tautological family of closed Riemann surfaces of some genus $g(\mathcal{L})$. Topologically, this is a fiber bundle $\pi: \mathcal{E}(\mathcal{L}) \rightarrow \mathcal{M}(\mathcal{L})$ with fiber $\Sigma_{g(\mathcal{L})}$. Consequently, there is a *monodromy representation*

This material is based upon work supported by the National Science Foundation under Award No. DMS-1703181.

✉ Nick Salter
nks@math.columbia.edu

¹ Department of Mathematics, Columbia University, 2990 Broadway, New York, NY 10027, USA

$$\mu_{\mathcal{L}}: \pi_1(\mathcal{M}(\mathcal{L}), C_0) \rightarrow \text{Mod}(C_0).$$

Here, $C_0 \in \mathcal{M}(\mathcal{L})$ is a fixed curve, and $\text{Mod}(C_0) := \pi_0(\text{Diff}^+(C_0))$ denotes the *mapping class group* of C_0 (see Sect. 2.1). Under $\mu_{\mathcal{L}}$, a based loop $\gamma \in \pi_1(\mathcal{M}(\mathcal{L}), C_0)$ is mapped to (the isotopy class of) the diffeomorphism $\mu_{\mathcal{L}}(\gamma) \in \text{Diff}(C_0)$ obtained by “parallel transport” of C_0 along γ . For details, see, e.g., [7, Section 5.6.1].

In this paper, we give a nearly complete answer to the following fundamental question. Define

$$\Gamma_{\mathcal{L}} := \text{Im}(\mu_{\mathcal{L}}) \leq \text{Mod}(\Sigma_{g(\mathcal{L})}).$$

Question 1.1 *What is $\Gamma_{\mathcal{L}}$? When is it a finite-index subgroup of $\text{Mod}(\Sigma_{g(\mathcal{L})})$? Can one give a precise characterization of $\Gamma_{\mathcal{L}}$?*

Question 1.1 is closely related to a question posed by Donaldson [6]. Fix a curve $C_0 \in \mathcal{M}(\mathcal{L})$ and an identification $C_0 \cong \Sigma_{g(\mathcal{L})}$. Define a *vanishing cycle* for \mathcal{L} as a simple closed curve γ on C_0 for which there is a degeneration of C_0 to a curve C' with a single node, such that γ becomes null-homotopic on C' . If c is a vanishing cycle, then necessarily the Dehn twist T_c lies in $\Gamma_{\mathcal{L}}$; it arises from a loop in $\mathcal{M}(\mathcal{L})$ encircling the nodal curve in $|\mathcal{L}|$.

Question 1.2 (Donaldson) *For \mathcal{L} an ample line bundle on a smooth toric surface X , which curves (on a fixed C_0) are vanishing cycles?*

A first insight into Questions 1.1 and 1.2 is to observe the presence of an invariant “higher spin structure”. Let K_X denote the canonical bundle of X . The *adjoint line bundle* of \mathcal{L} is the line bundle $\mathcal{L} \otimes K_X$. Define $r \in \mathbb{N}$ to be the highest root of $\mathcal{L} \otimes K_X$ in $\text{Pic}(X)$. As explained in Proposition 10.2, associated to $\mathcal{L} \otimes K_X$ is a $\mathbb{Z}/r\mathbb{Z}$ -valued *spin structure* $\phi_{\mathcal{L}}$, and the associated stabilizer subgroup $\text{Mod}(\Sigma_{g(\mathcal{L})})[\phi_{\mathcal{L}}]$ (see Definition 3.14). Proposition 10.2 asserts that necessarily $\Gamma_{\mathcal{L}} \leq \text{Mod}(\Sigma_{g(\mathcal{L})})[\phi_{\mathcal{L}}]$. The function $\phi_{\mathcal{L}}$ gives rise to a notion of *admissible curve* and the associated subgroup $\mathcal{T}_{\phi_{\mathcal{L}}} \leq \text{Mod}(\Sigma_{g(\mathcal{L})})[\phi_{\mathcal{L}}]$ of *admissible twists* (see Definition 3.16). If a curve c is a vanishing cycle, it is necessarily admissible; see Lemma 3.15. Our main theorem asserts that these necessary conditions are also sufficient (at least “virtually” so, in the case r is even).

Theorem A *Let \mathcal{L} be an ample line bundle on a smooth toric surface X for which the generic fiber is not hyperelliptic. Assume $r > 1$ or else $g(\mathcal{L}) \geq 5$.*

- *If r is odd, then $\Gamma_{\mathcal{L}} = \text{Mod}(\Sigma_{g(\mathcal{L})})[\phi_{\mathcal{L}}]$.*
- *If r is even, then $\Gamma_{\mathcal{L}} \leq \text{Mod}(\Sigma_{g(\mathcal{L})})$ is a finite-index subgroup that contains $\mathcal{T}_{\phi_{\mathcal{L}}}$.*

In either case, $[\text{Mod}(\Sigma_{g(\mathcal{L})}) : \Gamma_{\mathcal{L}}]$ is finite. Moreover, Question 1.2 admits the following complete answer: a curve γ is a vanishing cycle if and only if γ is an admissible curve.

We remark that many familiar algebraic surfaces such as $\mathbb{C}P^2$ and $\mathbb{C}P^1 \times \mathbb{C}P^1$ are smooth toric surfaces. For instance, as a special case of Theorem A we obtain the following theorem concerning smooth plane curves. The case $d = 5$ was addressed in [17], while the cases $d \leq 4$ are either classical or trivial.

Theorem 1.3 Set $g = \binom{d-1}{2}$, and define

$$\Gamma_d \leq \text{Mod}(\Sigma_g)$$

to be the monodromy group of the family of smooth curves in $\mathbb{C}P^2$ of degree d , i.e. the group $\Gamma_{\mathcal{L}}$ for the line bundle $\mathcal{L} = \mathcal{O}(d)$ on $\mathbb{C}P^2$. Then there exists a $\mathbb{Z}/(d - 3)\mathbb{Z}$ -valued spin structure ϕ_d such that the following hold.

- If d is even, then $\Gamma_d = \text{Mod}(\Sigma_g)[\phi_d]$.
- If d is odd, then Γ_d is of finite index in $\text{Mod}(\Sigma_g)[\phi_d]$, where Γ_d contains the subgroup \mathcal{T}_{ϕ_d} of admissible twists.

Theorem A also addresses a conjecture that was independently formulated by the author in [17] in the case of $X = \mathbb{C}P^2$, and in full generality by Crétois and Lang [3].

Conjecture 1.4 For any pair (X, \mathcal{L}) as above, there is an equality

$$\Gamma_{\mathcal{L}} = \text{Mod}(\Sigma_{g(\mathcal{L})})[\phi_{\mathcal{L}}].$$

Theorem A resolves Conjecture 1.4 in the affirmative whenever r is odd, and shows that in the case r even, $\Gamma_{\mathcal{L}}$ is at least of finite index in $\text{Mod}(\Sigma_{g(\mathcal{L})})[\phi_{\mathcal{L}}]$.

Theorem A is proved using a combination of methods from toric geometry and the theory of the mapping class group. On the toric end of the spectrum, we make essential use of the powerful results developed by Crétois and Lang [3]. The centerpiece of their theory is a combinatorial model for a curve $C_0 \in \mathcal{M}(\mathcal{L})$ based around a convex lattice polygon. Their results give a description of vanishing cycles in terms of lattice points and line segments, and allow one to produce many elements of $\Gamma_{\mathcal{L}}$. Crétois–Lang developed their methods in order to address Question 1.2 and Conjecture 1.4 in the case $r \leq 2$, and obtained complete answers in these cases. See [3] for the case $r = 1$, and [4] for the case $r = 2$, as well as the case where the general fiber is hyperelliptic.

On the mapping class group side, we carry out an extensive investigation of the groups $\text{Mod}(\Sigma_g)[\phi]$ and \mathcal{T}_{ϕ} mentioned above. We remark here that the theory of higher spin structures does not require the presence of a specific

ample line bundle \mathcal{L} , and so we adjust notation accordingly and refer to Riemann surfaces Σ_g , spin structures ϕ , etc. Our main result here is a general criterion for a collection of Dehn twists to generate (a finite-index subgroup of) $\text{Mod}(\Sigma_g)[\phi]$, given in Theorem 9.5.

Outline of the paper The bulk of the paper (Sects. 2–9) is devoted to developing the mapping class group technology necessary to show that the vanishing cycles investigated by Crétois–Lang generate a finite-index subgroup of the mapping class group. This culminates in Theorem 9.5. Portions of Theorem 9.5 are established earlier in Propositions 5.1 and 6.2.

Sections 2–4 contain preliminary results that are used throughout the paper. Section 2 collects the necessary background on mapping class groups; these results are standard and are included so as to fix notation and terminology, and to serve as a guide to the reader approaching the paper from a background in toric geometry. Section 3 presents the basic theory of higher spin structures, building off the foundational work of Humphries and Johnson [9]. Section 4 describes the action of the mapping class group on the set of higher spin structures. This yields several crucial corollaries (Corollaries 4.5, 4.10, 4.11) concerning the existence of configurations of curves with prescribed properties which are used extensively in subsequent sections.

Theorem 9.5 gives a criterion for a collection of Dehn twists to generate the so-called *admissible subgroup* \mathcal{T}_ϕ associated to a higher spin structure ϕ . A study of the admissible subgroup is sufficient to answer Question 1.2. The reader interested only in this portion of Theorem A can skip Sects. 5 and 6 and jump directly from Sects. 4–7.

The proof of Theorem 9.5 is carried out in Sects. 7–9. Section 7 establishes the connectivity of certain simplicial complexes acted on by the stabilizer subgroup of a higher spin structure. These results are used in the argument of Sect. 8, and also underlie the method by which the admissible subgroup is used to study the set of vanishing cycles. Section 8 is devoted to a study of certain subgroups of the admissible subgroup; the main result Proposition 8.2 furnishes a generating set for \mathcal{T}_ϕ in terms of these subgroups. Section 9 introduces the notion of a *network*; ultimately a network is a technical device used to factor the generators given in Proposition 8.2 into products of Dehn twists. Theorem 9.5 gives a sufficient condition, formulated in the language of networks, for a collection of Dehn twists to generate a subgroup containing the admissible subgroup.

The portion of Theorem A that goes beyond Question 1.2 concerns establishing that the admissible subgroup is finite-index in the mapping class group. This is the content of Sects. 5 and 6, which treat the case where the $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure under study has r odd or even, respectively. The arguments for these two cases are substantially different, owing to the fact that in the case of

r even, the higher spin structure has an Arf invariant which must be accounted for in various guises.

The net result of Sects. 2–9 is a criterion for a finite collection of Dehn twists to generate a finite-index subgroup of the mapping class group. In the final two sections, these results are applied in the setting of monodromy groups of linear systems on toric surfaces. Section 10 contains the necessary background material on toric surfaces, concentrating on the work of Crétois–Lang describing a particular finite collection of vanishing cycles. Section 11 exhibits a network amongst the set of vanishing cycles discussed in Sect. 10 and verifies that this network satisfies the hypotheses of Theorem 9.5 in order to obtain Theorem A.

2 Mapping class groups

This section collects background material on mapping class groups that will be used throughout the arguments in Sects. 3–9. Most of the material can be found in [7] and so will only be touched on briefly. The exception to this is the D_n relation of Sect. 2.3, which will consequently be dealt with in greater detail.

2.1 Basics

The material in this section is almost certainly well-known to a reader conversant in mapping class groups, but is included so as to fix notation and terminology.

Genus, boundary, punctures All surfaces under consideration are oriented and of finite type. A surface of genus g with n punctures and b boundary components is denoted by $\Sigma_{g,b}^n$. When one or more of $b, n = 0$, the corresponding decoration will be omitted.

Intersection numbers Let a, b be simple closed curves on a surface S . Often we will confuse the distinction between a simple closed curve and its isotopy class. The *geometric intersection number* between a, b will be notated $i(a, b)$ (see [7, Section 1.2.3]). For *oriented* simple closed curves a, b , the algebraic intersection number is denoted $\langle a, b \rangle$. Of course, algebraic intersection depends only on the homology classes $[a], [b] \in H_1(S; \mathbb{Z})$.

Mapping class groups Let $\Sigma_{g,b}^n$ be a surface. The *mapping class group* of $\Sigma_{g,b}^n$, written $\text{Mod}(\Sigma_{g,b}^n)$, is defined as




Fig. 1 The standard generators for $\Sigma_{g,3}^3$

$$\text{Mod}(\Sigma_{g,b}^n) := \pi_0 \left(\text{Diff}^+(\Sigma_{g,b}^n, \partial \Sigma_{g,b}^n) \right),$$

where $\text{Diff}^+(\Sigma_{g,b}^n, \partial \Sigma_{g,b}^n)$ denotes the group of orientation-preserving diffeomorphisms of $\Sigma_{g,b}^n$ that restrict to the identity on the boundary of $\Sigma_{g,b}^n$ and fix the punctures *pointwise* (not merely *setwise*, as some authors adopt).

The standard generators For a simple closed curve a on $\Sigma_{g,b}^n$, the *left-handed* Dehn twist about a is written T_a . For $g \geq 2$, the *standard generators* form a generating set for $\text{Mod}(\Sigma_{g,b}^n)$ consisting of the Dehn twists about the curves $a_1, a_2, b_1, \dots, b_g, c_1, \dots, c_{g-1}, d_1, \dots, d_{b+n-1}$ shown in Fig. 1.

The change-of-coordinates principle The classification of surfaces theorem asserts that if S, S' are two (connected and orientable) surfaces of finite type with the same genus, number of punctures, and number of boundary components, then there is a diffeomorphism $f: S \rightarrow S'$. This is often exploited in the study of mapping class groups in the guise of the “change-of-coordinates principle”. It is difficult to write down a single, all-encompassing statement of the change-of-coordinates principle, but informally, it states that *any configuration of curves, arcs, and/or subsurfaces of a surface S is determined up to diffeomorphism by combinatorial information alone*. In the present paper, the change-of-coordinates principle will often be invoked tacitly. The reader interested in a more thorough discussion of the change-of-coordinates principle is referred to [7, Section 1.3].

One consequence of the change-of-coordinates principle is that it becomes easy to understand the $\text{Mod}(S)$ orbits of many different kinds of configurations. As an example, we discuss here the action on *geometric symplectic bases* for S .

Definition 2.1 Let S be a surface of genus $g \geq 0$ with $n \geq 0$ boundary components and $b \geq 0$ punctures. A *geometric symplectic basis* for S is a collection of oriented simple closed curves $\mathcal{B} = \{\alpha_1, \beta_1, \dots, \alpha_g, \beta_g\}$ satisfying the following properties:

- (1) $i(a_i, b_i) = 1$ for each $i = 1, \dots, g$, and all other pairs of elements of \mathcal{B} are disjoint,
- (2) $\langle [a_i], [b_i] \rangle = 1$ for each $i = 1, \dots, g$.

Remark 2.2 The (homology classes of the) curves in a geometric symplectic basis form a basis for $H_1(S; \mathbb{Z})$ in the sense of linear algebra only when $n + b \leq 1$. In this paper, geometric symplectic bases are used to study $\mathbb{Z}/r\mathbb{Z}$ -valued spin structures. Proposition 3.8 and Theorem 3.9 together imply that a $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure is determined by its “signature” (Definition 4.1) in combination with its values on a geometric symplectic basis.

The following is a typical statement that is proved using the change-of-coordinates principle.

Lemma 2.3 *Let \mathcal{B} and \mathcal{B}' be two geometric symplectic bases for S . Then there is a diffeomorphism $f: S \rightarrow S$ such that $f(\mathcal{B}) = \mathcal{B}'$.*

2.2 The Birman exact sequence

A reference for this subsection is [7, Section 4.2]. Consider a surface $\Sigma_{g,b}^n$ with $n \geq 1$ and $2g + b + n \geq 4$. There is an inclusion $\Sigma_{g,b}^n \hookrightarrow \Sigma_{g,b}^{n-1}$ obtained by filling p in. This induces the *Birman exact sequence*

$$1 \rightarrow \pi_1(\Sigma_{g,b}^{n-1}, p) \rightarrow \text{Mod}(\Sigma_{g,b}^n) \rightarrow \text{Mod}(\Sigma_{g,b}^{n-1}) \rightarrow 1. \tag{1}$$

There is a slight variation on the Birman exact sequence where one fills in a boundary component with a closed disk, originally due to Johnson. In order to formulate this, we recall that the unit tangent bundle to a surface S is written UTS . Then the inclusion $\Sigma_{g,b}^n \rightarrow \Sigma_{g,b-1}^n$ induces the short exact sequence

$$1 \rightarrow \pi_1(UT\Sigma_{g,b-1}^n, \tilde{p}) \rightarrow \text{Mod}(\Sigma_{g,b}^n) \rightarrow \text{Mod}(\Sigma_{g,b-1}^n) \rightarrow 1, \tag{2}$$

where \tilde{p} is a unit tangent vector based at p . In both situations, the kernels admit descriptions in terms of Dehn twists. Consider first the case of (1). Let α be an *embedded, oriented* simple closed curve based at p , corresponding to an element $\alpha \in \pi_1(\Sigma_{g,b}^{n-1}, p)$. Let α_L (resp. α_R) denote the left (resp. right) side of a neighborhood of α . Both α_L, α_R are simple closed curves on $\Sigma_{g,b}^n$. Then $\alpha \in \pi_1(\Sigma_{g,b}^{n-1}, p)$ corresponds to $T_{\alpha_L} T_{\alpha_R}^{-1} \in \text{Mod}(\Sigma_{g,b}^n)$. The embedding $P: \pi_1(\Sigma_{g,b}^{n-1}, p) \rightarrow \text{Mod}(\Sigma_{g,b}^n)$ is known as the *point-pushing map*, and $\pi_1(\Sigma_{g,b}^{n-1}, p)$ is often referred to as the *point-pushing subgroup* of $\text{Mod}(\Sigma_{g,b}^n)$.

It is a basic topological fact that for any surface $\Sigma_{g,b}^{n-1}$, there exists a collection of simple closed curves $\alpha_1, \dots, \alpha_k$ based at p , such that $\{\alpha_1, \dots, \alpha_k\}$

generates $\pi_1(\Sigma_{g,b}^{n-1}, p)$. In practice, this means that to exhibit $\pi_1(\Sigma_{g,b}^n, p)$ as a subgroup of some group $H \leq \text{Mod}(\Sigma_{g,b}^n)$, it suffices to exhibit this finite collection of multitwists.

In the case of (2), everything is much the same. Let $\Sigma_{g,b}^n \hookrightarrow \Sigma_{g,b-1}^n$ be an inclusion corresponding to capping off a boundary component Δ of $\Sigma_{g,b}^n$. Let $p \in \Sigma_{g,b-1}^n$ be a point on the interior of this new disk, and \tilde{p} a tangent vector at p . Suppose that $\tilde{\alpha} \in \pi_1(UT\Sigma_{g,b-1}^n, \tilde{p})$ corresponds to a framed simple closed curve α based at \tilde{p} . We define α_L and α_R as before. Then

$$P(\tilde{\alpha}) = T_{\alpha_L} T_{\alpha_R}^{-1} T_{\Delta}^k$$

where $k \in \mathbb{Z}$ is the winding number of the tangent vector field specified by $\tilde{\alpha}$, relative to the tangential framing of the underlying curve α . The subgroup $\pi_1(UT\Sigma_{g,b-1}^n, \tilde{p})$ is known as the *disk-pushing subgroup* of $\text{Mod}(\Sigma_{g,b}^n)$.

There is an analogous set of “geometric” generators for $\pi_1(UT\Sigma_{g,b-1}^n, \tilde{p})$. Let $\alpha_1, \dots, \alpha_k$ be a collection of C^1 -embedded simple closed curves on $\Sigma_{g,b-1}^n$ based at p such that $\pi_1(\Sigma_{g,b-1}^n, p) = \langle \alpha_1, \dots, \alpha_k \rangle$ as above. Each α_i determines an element $\tilde{\alpha}_i \in \pi_1(\Sigma_{g,b-1}^n, \tilde{p})$ via the so-called *Johnson lift*, whereby α_i is framed using the forward-pointing tangent vector. Suppose that each $\tilde{\alpha}_i$ is based at some common tangent vector \tilde{p} . Then $\pi_1(UT\Sigma_{g,b-1}^n, \tilde{p})$ has a generating set of the following form:

$$\pi_1(UT\Sigma_{g,b-1}^n, \tilde{p}) = \langle \tilde{\alpha}_1, \dots, \tilde{\alpha}_k, \zeta \rangle,$$

where ζ is the loop around the S^1 fiber in the fibration $S^1 \rightarrow UT\Sigma_{g,b-1}^n \rightarrow \Sigma_{g,b-1}^n$. In terms of Dehn twists, the Johnson lifts $\tilde{\alpha}_i$ correspond to mapping classes $T_{\alpha_{i,L}} T_{\alpha_{i,R}}^{-1}$ as before, while ζ corresponds to T_{Δ} .

2.3 Relations

In this subsection we collect various relations in the mapping class group that will be used throughout the paper.

The braid relation Suppose a, b are simple closed curves satisfying $i(a, b) = 1$. Then the corresponding Dehn twists satisfy the *braid relation*:

$$T_a T_b T_a = T_b T_a T_b.$$

We will also employ the following alternative form, formulated in terms of the curves a, b themselves:

$$T_a T_b(a) = b.$$

The chain relation A chain of simple closed curves is a sequence (a_1, \dots, a_n) of simple closed curves such that $i(a_i, a_{i+1}) = 1$ and $i(a_i, a_j) = 0$ otherwise. Let ν denote a regular neighborhood of a chain of length n , where the representative curves a_1, \dots, a_n are in minimal position. When n is odd, $\partial\nu$ has two components Δ_1 and Δ_2 ; for n even, $\partial\nu = \Delta$ is a single (necessarily separating) curve. Abusing terminology, we will speak of the boundary of a chain itself, by which we mean the boundary of ν . Given a subsurface S with 1 or 2 boundary components, a chain a_1, \dots, a_n of curves on S is maximal if there is a deformation retraction of S onto $a_1 \cup \dots \cup a_n$. The following appears as [7, Proposition 4.12].

Proposition 2.4 (Chain relation) *For n odd,*

$$(T_{a_1} \dots T_{a_n})^{n+1} = T_{\Delta_1} T_{\Delta_2},$$

and for n even,

$$(T_{a_1} \dots T_{a_n})^{2n+2} = T_{\Delta}.$$

Remark 2.5 The intersection pattern of a chain of n simple closed curves is recorded by the Dynkin diagram of type A_n , where vertices in the graph are adjacent if the corresponding curves intersect, and are nonadjacent if the curves are disjoint. Such a chain of curves determines a homomorphism from the Artin group $A(A_n)$ of type A_n into the mapping class group $\text{Mod}(\nu)$, where generators of $A(A_n)$ are sent to Dehn twists about the corresponding curves.

Under this homomorphism, the chain relation is a consequence of the fact that $A(A_n)$ has nontrivial center. The twist(s) about the boundary component(s) appearing on the right-hand side of the expressions in Proposition 2.4 are elements of the center of $\text{Mod}(\nu)$, while the left-hand side merely gives the expression for a generator of $Z(A(A_n))$ as a word in the standard generators of $A(A_n)$. In [15, Section 2.4], Matsumoto explains how to determine the precise expression for this central element as a Dehn multitwist; this is the principle underlying the “ D_n relation” given in Proposition 2.6 below.

The D_n relation There is an analogous (though less ubiquitous) relation that arises from a configuration of curves whose intersection pattern is modeled on the Dynkin diagram of type D_n . Proposition 2.6 below is the specialization of [15, Proposition 2.4] to the case of an Artin group of type D_n . The case of n odd is treated explicitly in [15, Theorem 1.5], while the case of n even is given an alternate proof in [17, Proposition 4.5].




Fig. 2 The configuration of curves used in the D_n relation. Note the presence of the unlabeled curve c_{2g+1} on the far right side

Proposition 2.6 (D_n relation) *Let $n \geq 3$ be given, and express $n = 2g + 1$ or $n = 2g + 2$ according to whether n is odd or even. With reference to Fig. 2, let H_n be the group generated by elements of the form T_x , with $x \in \mathcal{D}_n$ one of the curves below:*

$$\mathcal{D}_n = \{a, a', c_1, \dots, c_{n-2}\}.$$

Then for $n = 2g + 1$ odd,

$$T_{\Delta_0}^{2g-1} T_{\Delta_2} \in H_n,$$

and for $n = 2g + 2$ even,

$$T_{\Delta_0}^g T_{\Delta_1} T_{\Delta_1'} \in H_n.$$

The D_n relation has some useful consequences which we record in Corollary 2.7 below. It is necessary to first describe the curves C_k that will appear in the statement. For $1 \leq k \leq g + 1$, let v_k be a regular neighborhood of the subconfiguration $\mathcal{D}_{2k+1} \subset \mathcal{D}_n$. Each such v_k is a surface of genus k with two boundary components. One of these is Δ_0 ; the other is defined to be the curve C_k . Note in particular that $C_g = \Delta_2$ and that C_{g+1} is the unlabeled boundary component of the ambient surface on the far right side of Fig. 2.

Corollary 2.7 *Fix notation as in Proposition 2.6, and for $1 \leq \ell \leq 2g + 3$, consider the configurations*

$$\mathcal{D}_\ell = \{a, a', c_1, \dots, c_{\ell-2}\}$$

as in Fig. 2. Let H_{2g+3}^+ be the group generated by H_{2g+3} and the Dehn twist T_{Δ_1} . Then the following assertions hold:

- (1) $T_{\Delta_1'} \in H_{2g+3}^+$,

(2) $T_{C_k}^m \in H_{2g+3}^+$ for any $1 \leq k \leq g + 1$ and any m such that $(2k - 1)m$ divides g .

Proof The proof of (1) follows from an important simple principle. Given a mapping class f and a simple closed curve d , there is a relation

$$fT_d f^{-1} = T_{f(d)}.$$

It follows that if $f, T_d \in H_{2g+3}^+$, then also $T_{f(d)} \in H_{2g+3}^+$. To establish (1), we will find $f \in H_{2g+3}^+$ such that $f(c_{2g+1}) = \Delta'_1$. This will be accomplished by means of the braid relation.

The curves $a, a', c_1, \dots, c_{2g}$ are arranged in the configuration of the D_{2g+2} relation; the boundary components correspond to $\Delta_0, \Delta_1, \Delta'_1$. By the D_{2g+2} -relation (Proposition 2.6),

$$T_{\Delta_0}^g T_{\Delta_1} T_{\Delta'_1} \in H_{2g+3}^+,$$

and since $T_{\Delta_1} \in H_{2g+3}^+$ by assumption, also $T_{\Delta_0}^g T_{\Delta'_1} \in H_{2g+3}^+$. Since Δ_0 is disjoint from both c_{2g+1} and Δ'_1 , the braid relation implies that

$$T_{c_{2g+1}} T_{\Delta_0}^g T_{\Delta'_1}(c_{2g+1}) = T_{c_{2g+1}} T_{\Delta'_1}(c_{2g+1}) = \Delta'_1.$$

Since $(T_{\Delta_0}^g T_{\Delta'_1}) \in H_{2g+3}^+$, this shows $T_{\Delta'_1} \in H_{2g+3}^+$ as required.

We observe that (2) follows from the D_{2k-1} relation (as applied to the subconfiguration \mathcal{D}_{2k-1}) and the claim that $T_{\Delta_0}^g \in H_{2g+3}^+$; this latter assertion follows from the D_{2g+2} relation (applied to \mathcal{D}_{2g+2}) and (1). \square

2.4 The Torelli group

Most of the material in this subsection can be found in [7, Chapter 6], but see also [12]. We begin by observing that the action of $\text{Mod}(\Sigma_g)$ on $H_1(\Sigma_g; \mathbb{Z})$ preserves the algebraic intersection pairing $\langle \cdot, \cdot \rangle$, leading to the *symplectic representation*

$$\Psi: \text{Mod}(\Sigma_g) \rightarrow \text{Sp}(2g, \mathbb{Z}). \tag{3}$$

This is classically known to be a surjection. The *Torelli group*, notated \mathcal{I}_g , is the kernel of this representation:

$$\mathcal{I}_g := \ker(\Psi).$$

Bounding pairs and separating twists There are two types of elements in \mathcal{I}_g that will be of particular importance. Suppose that c, d are simple closed curves such that $c \cup d$ bounds a subsurface $S \cong \Sigma_{h,2}$. Then $T_c T_d^{-1} \in \mathcal{I}_g$ is known as a *bounding pair map*. The *genus* of a bounding pair map is slightly ambiguous: if $c \cup d$ bounds a surface $\Sigma_{h,2}$, then also $c \cup d$ bounds a surface $\Sigma_{g-h-1,2}$ on the other side. One defines the genus of $T_c T_d^{-1}$ as $\min\{h, g - h - 1\}$. The second important class of elements is the class of *separating twists*—these are Dehn twists T_c for c a separating curve. The *genus* of a separating twist T_c that bounds a subsurface of genus h is defined as $g(c) = \min\{h, g - h\}$.

The Johnson homomorphism A fundamental tool in the study of \mathcal{I}_g is the *Johnson homomorphism*, due to Johnson [10]. This is a surjective homomorphism

$$\tau: \mathcal{I}_g \rightarrow \wedge^3 H_{\mathbb{Z}} / H_{\mathbb{Z}}, \tag{4}$$

where for convenience we define $H_A := H_1(\Sigma_g; A)$ for some abelian group A . The embedding $H_{\mathbb{Z}} \hookrightarrow \wedge^3 H_{\mathbb{Z}}$ is defined via

$$z \mapsto z \wedge (x_1 \wedge y_1 + \cdots + x_g \wedge y_g),$$

where $\{x_1, \dots, y_g\}$ is a symplectic basis for $H_{\mathbb{Z}}$. Recall that a symplectic basis must satisfy $\langle x_i, y_i \rangle = 1$ and $\langle x_i, x_j \rangle = \langle x_i, y_j \rangle = 0$ for $i \neq j$.

We will not need to know a precise definition of τ , but it will be useful to know some basic properties of τ , including how to compute τ on bounding pair maps and separating twists.

Lemma 2.8 (Johnson [10])

- (1) τ is $\text{Sp}(2g; \mathbb{Z})$ -equivariant, with respect to the conjugation action on \mathcal{I}_g and the evident action on $\wedge^3 H_{\mathbb{Z}} / H_{\mathbb{Z}}$.
- (2) $\tau(T_c) = 0$ for any separating twist T_c .
- (3) Let $c \cup d$ bound a subsurface $\Sigma_{h,2}$. Choose any further subsurface $\Sigma_{h,1} \subset \Sigma_{h,2}$, and let $\{x_1, y_1, \dots, x_h, y_h\}$ be a symplectic basis for $H_1(\Sigma_{h,1}; \mathbb{Z})$. Then

$$\tau(T_c T_d^{-1}) = (x_1 \wedge y_1 + \cdots + x_h \wedge y_h) \wedge [c],$$

where c is oriented with $\Sigma_{h,2}$ to the left. In the case $h = 1$, if α, β, γ is a maximal chain on $\Sigma_{1,2}$, then

$$\tau(T_c T_d^{-1}) = [\alpha] \wedge [\beta] \wedge [\gamma].$$

The Johnson kernel The *Johnson kernel*, written \mathcal{K}_g , is the kernel of the Johnson homomorphism:

$$\mathcal{K}_g := \ker(\tau).$$

A fundamental theorem of Johnson gives an alternate characterization of \mathcal{K}_g in terms of separating twists.

Theorem 2.9 (Johnson [13]) *Let \mathcal{T}_g be the subgroup of \mathcal{K}_g generated by all separating twists of genus at most two. Then for all $g \geq 3$,*

$$\mathcal{T}_g = \mathcal{K}_g.$$

3 Spin structures

In this section we introduce and study higher spin structures and their stabilizer subgroups. Section 3.1 defines higher spin structures and presents the work of Humphries–Johnson that gives a cohomological formulation of a higher spin structure. Section 3.2 discusses some cut-and-paste operations on simple closed curves and how these operations interact with higher spin structures. Section 3.3 defines spin structure stabilizer groups and some important elements of these groups. Finally Sect. 3.4 explains the connection between higher spin structures and the classical theory of spin structures as quadratic forms on vector spaces over $\mathbb{Z}/2\mathbb{Z}$.

3.1 Spin structures

Let S be a surface of genus $g \geq 0$. For simplicity, we assume in this section that S can have boundary components but not punctures; for surfaces with puncture, one can simply remove an open neighborhood of the puncture to produce a surface with boundary. Let \mathcal{S} denote the set of isotopy classes of oriented simple closed curves on S . In keeping with standard practice, the term “curve” will often be used to refer to an isotopy class of curves. Crucially, curves are not required to be essential [see property (2) of Definition 3.1]. The following definition is due to Humphries and Johnson [9]; see Remark 3.2 for a discussion of how to reconcile their definition with the one given here.

Definition 3.1 (*spin structure*) A $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure on S is a function $\phi: \mathcal{S} \rightarrow \mathbb{Z}/r\mathbb{Z}$ satisfying the following two properties.

(1) (Twist-linearity) Let $c, d \in \mathcal{S}$ be arbitrary. Then

$$\phi(T_c(d)) = \phi(d) + \langle d, c \rangle \phi(c) \pmod{r}.$$

- (2) (Normalization) For ζ the boundary of an embedded disk $D \subset S$, oriented with D to the left, $\phi(\zeta) = 1$.

Remark 3.2 The definition of a $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure presented in Definition 3.1 is superficially different from that given by Humphries and Johnson [9] in several respects. First, it should be noted that Humphries–Johnson study a more general notion of “twist-linear function”; only spin structures are needed in the present paper. Secondly, in Definition 3.1, simple closed curves are considered up to the equivalence relation of isotopy. This is an a priori different equivalence relation than the notion of “ L -direct homotopy” defined in [9, p. 366]. The precise definition of L -directness is cumbersome, but if two simple closed curves c and d are L -directly homotopic, then they are in particular homotopic in the ordinary sense. It is well-known that homotopy and isotopy determine the same equivalence relation on simple closed curves, see e.g. [7, Proposition 1.10]. Moreover, an isotopy is an instance of an L -direct homotopy, so that these notions coincide in our setting.

Remark 3.3 In the literature, higher spin structures go by various names and have various definitions; the term “ r -spin structure” is especially common. It is not a priori clear how to reconcile the definition given here with others. See Remark 3.7 for a brief discussion, or [17, Sections 2–3] for a fuller treatment.

Convention 3.4 Often we will speak of the value $\phi(c)$ where ϕ is some $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure and c is a curve *without a specified orientation*. Such a statement should be understood to mean that there is some unspecified orientation on c for which $\phi(c)$ has the stated value.

The Johnson lift Recall from the discussion in Sect. 2.2 the notion of the *Johnson lift*. In [9], Johnson–Humphries use the Johnson lift to give a homological formulation of a $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure. The following is an amalgamation of the Remark following Theorem 2.1 and Theorem 2.5 of [9].

Theorem 3.5 (Humphries–Johnson) *Let S be a surface. An element $\psi \in H^1(UTS; \mathbb{Z}/r\mathbb{Z})$ determines a $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure via*

$$\alpha \mapsto \psi(\tilde{\alpha}),$$

where α is a simple closed curve on S and $\tilde{\alpha}$ is the Johnson lift. This determines a $1 - 1$ correspondence

$$\{\phi \text{ a } \mathbb{Z}/r\mathbb{Z}\text{-valued spin structure on } S\} \leftrightarrow \{\phi \in H^1(UTS; \mathbb{Z}/r\mathbb{Z}) \mid \phi(\zeta) = 1\}.$$

Remark 3.6 From the standard presentation

$$\pi_1(UT\Sigma_g) = \left\langle a_1, b_1, \dots, a_g, b_g, \zeta \mid \prod_{i=1}^g [a_i, b_i] = \zeta^{2-2g} \right\rangle$$

and the Universal Coefficient Theorem, one sees that

$$H^1(UT\Sigma_g; A) \cong \text{Hom}(\pi_1(UT\Sigma_g), A) \cong \text{Hom}(\mathbb{Z}^{2g} \oplus \mathbb{Z}/(2g - 2)\mathbb{Z}, A).$$

The factor $\mathbb{Z}/(2g - 2)\mathbb{Z}$ in $H_1(UT\Sigma_g; \mathbb{Z}) = \mathbb{Z}^{2g} \oplus \mathbb{Z}/(2g - 2)\mathbb{Z}$ is generated by the class of $\tilde{\zeta}$, the Johnson lift of the non-essential curve ζ . In the case $A = \mathbb{Z}/r\mathbb{Z}$, it follows that there exists a spin structure if and only if $r \mid (2g - 2)$.

Remark 3.7 Via covering space theory, $\mathbb{Z}/r\mathbb{Z}$ -valued spin structures on Σ_g are in correspondence with cyclic r -fold coverings $\widetilde{UT\Sigma_g} \rightarrow UT\Sigma_g$ that restrict to connected coverings of the fiber S^1 . In the setting of linear systems on toric surfaces, such coverings arise from the presence of roots of the canonical line bundle of the generic fiber. See Proposition 10.2 and the references mentioned therein for more details.

An important consequence of Theorem 3.5 is the fact that $\mathbb{Z}/r\mathbb{Z}$ -valued spin structures satisfy a property known as the *homological coherence criterion*. This follows by combining Theorem 3.5 with [9, Lemma 2.4].

Proposition 3.8 (Homological coherence criterion) *Let ϕ be a $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure on a surface S , and let $S' \subset S$ be a subsurface with Euler characteristic $\chi(S') = m$. Suppose $\partial(S') = c_1 \cup \dots \cup c_k$, and all c_i are oriented so that S' is to the left. Then $\sum \phi(c_i) = m$.*

Theorem 3.5 shows that $\mathbb{Z}/r\mathbb{Z}$ -valued spin structures are determined by a finite amount of data. In the sequel it will be useful to have an explicit criterion for the equality of two $\mathbb{Z}/r\mathbb{Z}$ -valued spin structures. The following appears as [9, Corollary 2.6].

Theorem 3.9 (Humphries–Johnson) *Let S be a surface of genus $g \geq 0$. Let $\mathcal{B} = \{\gamma_1, \dots, \gamma_k\}$ be a set of oriented simple closed curves such that the set $\{[\gamma_1], \dots, [\gamma_k]\}$ forms a basis for $H_1(\Sigma_g; \mathbb{Z})$. Suppose ϕ and ψ are $\mathbb{Z}/r\mathbb{Z}$ -valued spin structures on S . Then $\phi = \psi$ if and only if $\phi(\gamma_i) = \psi(\gamma_i)$ for each $\gamma_i \in \mathcal{B}$.*

3.2 Operations on curves

In what follows, we will make use of two procedures for constructing new simple closed curves from old. Here, we define these operations and collect some facts about how they interact with spin structures.




Fig. 3 The smoothing operation




Fig. 4 The curve-arc sum operation

Definition 3.10 (*Smoothing, curve sum*) Let $\mathcal{C} = \{c_1, \dots, c_n\}$ be a collection of oriented embedded simple closed curves on a surface S . Suppose that all intersections between elements of \mathcal{C} are transverse. The *smoothing* of \mathcal{C} is the embedded multicurve obtained from \mathcal{C} by smoothly resolving all intersections in the unique orientation-preserving way. See Fig. 3.

Now suppose α and β are oriented simple closed curves. For natural numbers m, n , define the *curve sum* $m\alpha + n\beta$ as the smoothing of m parallel copies of α with n copies of β . In case $m < 0$ or $n < 0$, the curve sum $m\alpha + n\beta$ can be defined as before, with the orientation on α (resp. β) reversed if $m < 0$ (resp. $n < 0$). See Fig. 4.

By choosing arbitrary representatives in minimal position, both of these operations are well-defined on the level of isotopy classes.

Lemma 3.11 *Let α, β be oriented simple closed curves in minimal position, and let ϕ be a $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure. Then for any integers m, n ,*

$$\phi(m\alpha + n\beta) = m\phi(\alpha) + n\phi(\beta).$$

If in addition, $i(\alpha, \beta) = 1$ and $\gcd(m, n) = 1$, then $m\alpha + n\beta$ has a single component.

Proof The first assertion follows directly from the identification of ϕ with an element of $H^1(UTS; \mathbb{Z}/r\mathbb{Z})$ given in Theorem 3.5, while the second is straightforward to verify. □

Definition 3.12 (*Curve-arc sum*) Let α and β be disjoint oriented simple closed curves on S , and let ϵ be an arc connecting α to β whose interior is disjoint from $\alpha \cup \beta$. A regular neighborhood ν of $\alpha \cup \epsilon \cup \beta$ is homeomorphic to $\Sigma_{0,3}$. Two of the boundary components of ν are homotopic to α and

β , respectively. The *curve-arc-sum* $\alpha +_\epsilon \beta$ is the third boundary component of v . Again, the curve-arc sum descends to the level of isotopy classes.

Lemma 3.13 *Let $\alpha, \beta, \epsilon, v$ be as above. Orient α, β so that ϵ connects the left sides of α, β , and orient $\alpha +_\epsilon \beta$ so that the subsurface v is to the right. Then for ϕ a $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure,*

$$\phi(\alpha +_\epsilon \beta) = \phi(\alpha) + \phi(\beta) + 1.$$

In addition, on the level of homology, $[\alpha +_\epsilon \beta] = [\alpha] + [\beta]$.

Proof Observe that $\chi(v) = -1$. By the homological coherence criterion (Proposition 3.8),

$$-1 = \phi(\alpha) + \phi(\beta) + \phi(-(\alpha +_\epsilon \beta)),$$

where $-(\alpha +_\epsilon \beta)$ denotes the curve $\alpha +_\epsilon \beta$ with orientation opposite to that specified above. By the case $(m, n) = (-1, 0)$ of Lemma 3.11, it follows that $\phi(-(\alpha +_\epsilon \beta)) = -\phi(\alpha +_\epsilon \beta)$, from which the first claim follows. The second claim is an immediate consequence of the orientation conventions. □

3.3 The group $\text{Mod}(S)[\phi]$; first examples of elements

For any surface S , there is an obvious (left) action of $\text{Mod}(S)$ on the set of $\mathbb{Z}/r\mathbb{Z}$ -valued spin structures: for $f \in \text{Mod}(S)$ and $c \in \mathcal{S}$, define $(f \cdot \phi)(c) = \phi(f^{-1}(c))$. Similarly, if $f: S \rightarrow S'$ is a diffeomorphism and ϕ is a $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure on S' , there is a pullback $f^*(\phi)$ defined on S via $(f^*\phi)(c) = \phi(f(c))$.

Definition 3.14 (*Stabilizer subgroup*) Let ϕ be a spin structure on a surface S . The *stabilizer subgroup* of ϕ , written $\text{Mod}(S)[\phi]$, is defined as

$$\text{Mod}(S)[\phi] = \{f \in \text{Mod}(S) \mid (f \cdot \phi) = \phi\}.$$

Let ϕ be a $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure on a surface S . Below we discuss some fundamental examples of elements in $\text{Mod}(S)[\phi]$.

Dehn twist powers and admissible twists The twist-linearity formula of Definition 3.1 immediately implies the following characterization of Dehn twists in $\text{Mod}(S)[\phi]$.

Lemma 3.15 *Let c be a simple closed curve on S . If c is separating, then $T_c \in \text{Mod}(S)[\phi]$. If c is nonseparating, then $T_c^k \in \text{Mod}(S)[\phi]$ if and only if $k\phi(c) \equiv 0 \pmod{r}$. In particular, for c nonseparating, $T_c \in \text{Mod}(S)[\phi]$ if and only if $\phi(c) = 0$.*

Definition 3.16 (*Admissible*) A nonseparating curve c with $\phi(c) = 0$ is called an *admissible curve*. The associated element $T_c \in \text{Mod}(S)[\phi]$ is called an *admissible twist*. The group generated by all admissible twists is written \mathcal{T}_ϕ , and is called the *admissible subgroup*.

Fundamental multitwists Let $P \cong \Sigma_{0,3}$ be a pair of pants with boundary curves α, β, γ . Suppose that $\phi(\alpha) = a, \phi(\beta) = b$, and that $\phi(\gamma) = c$, with all curves oriented so that P lies to the left. By the homological coherence property, $a + b + c = -1$.

Definition 3.17 Let P and ϕ be as above. A ϕ -*bounding multitwist* associated to P , denoted $T_P(x, y, z)$, is given by

$$T_P(x, y, z) = T_\alpha^x T_\beta^y T_\gamma^z$$

for any choice of integers x, y, z such that $T_P \in \text{Mod}(S)[\phi]$.

By the above, $T_P(r, r, r)$ is a ϕ -bounding multitwist for any P and ϕ , but for special values of a, b, c , there are much simpler examples.

Lemma 3.18 Let P be as above, and suppose that $b = -a$, so that $c = -1$. Then $T_P(1, -1, b) = T_\alpha T_\beta^{-1} T_\gamma^b$ is a ϕ -bounding multitwist. The element $T_P(1, -1, b)$ is called a *fundamental multitwist* for P and is denoted T_P .

Proof Let d be any curve on S ; we must show that $\phi(d) = \phi(T_\alpha T_\beta^{-1} T_\gamma^b(d))$. As α, β, γ are all disjoint, the twist-linearity property, in combination with the fact that $[\alpha + \beta + \gamma] = 0$ in $H_1(S)$, gives

$$\begin{aligned} \phi(T_\alpha T_\beta^{-1} T_\gamma^b(d)) &= \phi(d) + \langle d, \alpha \rangle a - \langle d, \beta \rangle b - \langle d, \gamma \rangle b \\ &= \phi(d) - \langle d, \alpha + \beta + \gamma \rangle b \\ &= \phi(d). \end{aligned}$$

□

Remark 3.19 Of course, if $T_P(1, -1, b)$ is a fundamental multitwist, then so is $T_P(1, -1, b + kr)$ for any $k \in \mathbb{Z}$. An important special case is when $\phi(\alpha) = \phi(\beta) = 0$. Then $T_\alpha T_\beta^{-1}$ is a fundamental multitwist.

3.4 “Classical” spin structures

Spin structures in the sense of Definition 3.1 generalize the more familiar notion of a “classical” spin structure. In our setting, a classical spin structure is a spin structure valued in $\mathbb{Z}/2\mathbb{Z}$. We pause here to briefly review the theory of classical spin structures and the connection with our definition. These results,

especially the theory of the Arf invariant, will play a crucial role in the analysis of $\mathbb{Z}/r\mathbb{Z}$ -valued spin structures for r even to be begun in Proposition 4.9 and Corollary 4.10, and returned to in Sect. 6.

Let V be a vector space over the field $\mathbb{Z}/2\mathbb{Z}$ equipped with a nondegenerate symplectic pairing $\langle \cdot, \cdot \rangle$ (i.e. a nondegenerate bilinear pairing satisfying $\langle x, x \rangle = 0$ for all $x \in V$). The motivating example is $V = H_1(\Sigma_g; \mathbb{Z}/2\mathbb{Z})$ with the intersection pairing. A $\mathbb{Z}/2\mathbb{Z}$ quadratic form relative to $\langle \cdot, \cdot \rangle$ is a function $q: V \rightarrow \mathbb{Z}/2\mathbb{Z}$ such that for any $x, y \in V$, the equation

$$q(x + y) = q(x) + q(y) + \langle x, y \rangle \tag{5}$$

holds.

Let $\mathcal{B} = \{x_1, y_1, \dots, x_g, y_g\}$ be a symplectic basis for V . It is clear that q is determined by its values on \mathcal{B} . Define $Q(V, \langle \cdot, \cdot \rangle)$ as the set of $\mathbb{Z}/2\mathbb{Z}$ quadratic forms on V relative to $\langle \cdot, \cdot \rangle$; then a choice of \mathcal{B} provides a bijection

$$Q(V, \langle \cdot, \cdot \rangle) \cong (\mathbb{Z}/2\mathbb{Z})^{2g}.$$

There is an evident action of the group $\text{Sp}(V, \langle \cdot, \cdot \rangle)$ of $\langle \cdot, \cdot \rangle$ -preserving automorphisms on $Q(V, \langle \cdot, \cdot \rangle)$.

To understand the set of orbits, we introduce the *Arf invariant*. The Arf invariant of q is the element of $\mathbb{Z}/2\mathbb{Z}$ defined by the following formula:

$$\text{Arf}(q) := \sum_{i=1}^g q(x_i)q(y_i).$$

q is said to be *even* or *odd* according to whether $\text{Arf}(q) = 0, 1$ respectively; in this way we will speak of the *parity* of a spin structure. The following records some well-known properties of the Arf invariant.

Lemma 3.20 *Let $(V, \langle \cdot, \cdot \rangle)$ be a symplectic vector space over $\mathbb{Z}/2\mathbb{Z}$, and let $q, q' \in Q(V, \langle \cdot, \cdot \rangle)$ be quadratic forms.*

- (1) *Arf(q) is well-defined independently of the choice of symplectic basis,*
- (2) *q and q' are in the same orbit of $\text{Sp}(V, \langle \cdot, \cdot \rangle)$ if and only if $\text{Arf}(q) = \text{Arf}(q')$.*

Suppose now that ϕ is a $\mathbb{Z}/2d\mathbb{Z}$ -valued spin structure in the sense of Definition 3.1. The reduction $\mathbb{Z}/2d\mathbb{Z} \rightarrow \mathbb{Z}/2\mathbb{Z}$ associates to ϕ an underlying $\mathbb{Z}/2\mathbb{Z}$ -valued spin structure which we denote $\bar{\phi}$. A priori, $\bar{\phi}$ is defined on the set \mathcal{S} of isotopy classes of oriented curves on Σ_g . It follows from [11, Theorem 1A] that $\bar{\phi}$ factors through the map $[\cdot]: \mathcal{S} \rightarrow H_1(\Sigma_g; \mathbb{Z}/2\mathbb{Z})$. The induced map

$$\bar{\phi}: H_1(\Sigma_g; \mathbb{Z}/2\mathbb{Z}) \rightarrow \mathbb{Z}/2\mathbb{Z}$$

is not quite a classical spin structure, but it follows from [11, Theorem 1A] that the function

$$q_\phi := \bar{\phi} + 1 \quad (6)$$

does determine a classical spin structure.

In the remainder of this paper we will exclusively use the term “spin structure” in the sense of Definition 3.1. The reader versed in classical spin structures should be aware that certain formulas appear different in this setting. For instance, a Dehn twist about some nonseparating curve c preserves a $\mathbb{Z}/2\mathbb{Z}$ -valued spin structure ϕ if and only if $\phi(c) = 0$, whereas a transvection about some nonzero $v \in V$ preserves a quadratic form q if and only if $q(v) = 1$. Likewise, if ϕ is a $\mathbb{Z}/2d\mathbb{Z}$ -valued spin structure, the formula for the Arf invariant $\text{Arf}(\phi)$ of the underlying classical spin structure is given by

$$\text{Arf}(\phi) = \sum_{i=1}^g (\bar{\phi}(x_i) + 1)(\bar{\phi}(y_i) + 1) \pmod{2}. \quad (7)$$

4 The action of the mapping class group on spin structures

In what follows, we will need to understand the action of $\text{Mod}(\Sigma_g)$ on the set of $\mathbb{Z}/r\mathbb{Z}$ -valued spin structures. Following the discussion in Sect. 3.4, when r is even, the Arf invariant shows there are at least two orbits of $\text{Mod}(S)$ on the set of $\mathbb{Z}/r\mathbb{Z}$ -valued spin structures, but it is not clear what happens for odd r , nor whether there are further invariants leading to more orbits. The goal of this section is to give a complete description of this action. In the case of r odd, the mapping class group action on the set of $\mathbb{Z}/r\mathbb{Z}$ -valued spin structures is described in Proposition 4.2, and for r even it is described in Proposition 4.9. Both results can be understood as asserting that there are no “higher Arf invariants”.

4.1 Odd r

In the case of r odd, we will need to consider surfaces with multiple boundary components. Before formulating the results, we define the notion of the *signature* of a $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure.

Definition 4.1 (*Signature of a $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure*) Let S be a surface equipped with a $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure ϕ . Enumerate the boundary components as $\Delta_1, \dots, \Delta_n$, each one oriented so that S is to the left. The *signature of S rel ϕ* is defined as the n -tuple of values $\text{sig}(S, \phi) = (\phi(\Delta_1), \dots, \phi(\Delta_n))$. We will also speak of the signature of an individual Δ_k , defined as $\phi(\Delta_k)$.

Proposition 4.2 *Fix an odd integer r . Let S be a surface, and let ϕ and ψ be $\mathbb{Z}/r\mathbb{Z}$ -valued spin structures on S satisfying $\text{sig}(\phi) = \text{sig}(\psi)$. Suppose that either $g(S) \neq 1$ or else $g = 1$ and there is at least one boundary component with signature $\phi(c_1) = \psi(c_1) = k$ for some k such that $k + 1$ generates $\mathbb{Z}/r\mathbb{Z}$. Then there is a mapping class $f \in \text{Mod}(S)$ such that $f^*(\psi) = \phi$.*

Proof The proof is by induction on the genus $g(S)$. If $g(S) = 0$, then every curve c on S is separating, so that the homological coherence criterion (Proposition 3.8) implies that $\phi(c)$ and $\psi(c)$ can be computed just from the signature. In this case, it follows that in fact $\phi = \psi$.

For $g(S) \geq 1$, let α_0, β_0 be curves on S satisfying $i(\alpha_0, \beta_0) = 1$. Choose nonzero integers $a, b \in \mathbb{Z}$ such that $a \equiv \phi(\alpha_0)$ and $b \equiv \phi(\beta_0) \pmod{r}$. Let $d = \text{gcd}(a, b)$, and define $x = a/d, y = b/d$; by construction, x, y are coprime. Define the curve $\alpha_1 = y\alpha_0 - x\beta_0$ in the sense of Definition 3.10. By Lemma 3.11, $\phi(\alpha_1) = 0$.

Choose any curve γ_0 satisfying $i(\alpha_1, \gamma_0) = 1$. We claim there exists some separating oriented curve c on S that is disjoint from $\gamma_0 \cup \alpha_1$ and such that $\phi(c) = k$ for k such that $k + 1$ generates $\mathbb{Z}/r\mathbb{Z}$. In the case $g(S) = 1$ this is true by hypothesis, while for $g(S) \geq 2$, the curve c can be taken to be the neighborhood of some subsurface $T \subset S$ with $T \cong \Sigma_{1,1}$ and T disjoint from $\alpha_1 \cup \gamma_0$. In this case, orient c so that T lies to the right. By the homological coherence property, such a c satisfies $\phi(c) = 1$, and since r is odd, the claim follows.

Either c is isotopic to a boundary component of S and is oriented with S lying to the right, or else (by the change-of-coordinates principle), there exists an arc ϵ_0 from the left side of γ_0 to the left side of c that is disjoint from α_1 . In the former case, there exists an arc ϵ_0 from the right side of γ_0 to the right side of c that is disjoint from α_1 . Via Lemma 3.13, the curve-arc sum $\gamma_1 = \gamma_0 +_{\epsilon_0} c$ satisfies $\phi(\gamma_1) = \phi(\gamma_0) - (k + 1)$ in the former case, and $\phi(\gamma_1) = \phi(\gamma_0) + (k + 1)$ in the latter case. Since the curve c is null-homologous, there is an equality $[\gamma_1] = [\gamma_0]$. A further appeal to the change-of-coordinates principle shows that there is another arc ϵ_1 from the left side of γ_1 to the left of c , again disjoint from α_1 . This process can therefore be repeated indefinitely, giving rise to curves γ_m satisfying $\phi(\gamma_m) = \phi(\gamma_0) + m(k + 1)$. By hypothesis, $k + 1 \in \mathbb{Z}/r\mathbb{Z}$ is a generator, so that $\phi(\gamma_m) = 0$ for some m . Set $\beta_1 = \gamma_m$ for such an m . By construction, $i(\alpha_1, \beta_1) = 1$.

Likewise, construct curves α'_1, β'_1 satisfying $i(\alpha'_1, \beta'_1) = 1$ and $\psi(\alpha'_1) = \psi(\beta'_1) = 0$. Take (open) regular neighborhoods T_1 and T'_1 of $\alpha_1 \cup \beta_1$ and $\alpha'_1 \cup \beta'_1$, respectively. There is a diffeomorphism $f_1: T_1 \xrightarrow{\cong} T'_1$ for which $f_1(\alpha_1) = \alpha'_1$ and $f_1(\beta_1) = \beta'_1$. Define $c_1 = \partial \overline{T_1}$ and $c'_1 = \partial \overline{T'_1}$. Then $\phi(c_1) = 1$ when c_1 is oriented with T_1 on the right, and similarly for c'_1 . The curve c_1 is therefore a boundary component of $S \setminus T_1$ with signature $\phi(c_1) = 1$, and likewise for c'_1 . This shows that the inductive hypothesis is satisfied, and so there exists a

diffeomorphism $f_2: S \setminus T_1 \rightarrow S \setminus T'_1$ taking c_1 to c'_1 and fixing each remaining mutual boundary component, such that

$$f_2^*(\psi|_{S \setminus T'_1}) = \phi|_{S \setminus T_1}.$$

The diffeomorphisms f_1 and f_2 can be chosen in such a way as to extend to a diffeomorphism $f: S \rightarrow S$. Let $\mathcal{B} = \{\alpha_1, \beta_1, \dots, \alpha_g, \beta_g\}$ be a geometric symplectic basis for S , with α_1, β_1 the same curves as above. Necessarily α_k, β_k are curves on $S \setminus T_1$ for $k \geq 2$. By construction, the spin structures ϕ and $f^*(\psi)$ take the same values on each element of \mathcal{B} , and $\text{sig}(S, \phi) = \text{sig}(S, f^*(\psi))$. It then follows from Proposition 3.9 that $\phi = f^*(\psi)$ as claimed. \square

Proposition 4.2 has several corollaries that will be used extensively in the remainder of the paper. These play the role of a change-of-coordinates principle for surfaces in the presence of a $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure. The first of these was established in the second paragraph of the proof of Proposition 4.2. We remark that the assumption that r is odd played no role in the argument.

Corollary 4.3 *Let r be an integer and let ϕ be a $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure on a surface S . Let $S' \subset S$ be a subsurface of genus $h \geq 1$. Then there is some admissible curve $a \subset S'$ that is not parallel to a boundary component.*

This in turn leads to another useful result that will allow us to construct curves with prescribed intersection properties and arbitrary ϕ -values.

Corollary 4.4 *Let r be an integer and let ϕ be a $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure on a surface S . Let a, c_1, \dots, c_k be a collection of simple closed curves. Assume that there is some connected subsurface $T \subset S$ of positive genus disjoint from a, c_1, \dots, c_k , and that there is an arc ϵ connecting a to ∂T that is disjoint from all c_i . Then for $\ell \in \mathbb{Z}/r\mathbb{Z}$ arbitrary, there is a simple closed curve a_ℓ for which $i(a_\ell, c_i) = i(a, c_i)$ for $i = 1, \dots, k$, and for which $\phi(a_\ell) = \ell$.*

Proof By Corollary 4.3, there exists an admissible curve $b \subset T$ that is not boundary-parallel. The arc ϵ can be concatenated with an arc joining ∂T to b ; denote this extended arc by ϵ' . Set $\ell_0 = \phi(a)$ (where a is oriented with ϵ' lying to the left), and define $a_{\ell_0} := a$. Define $a_{\ell_0+1} := a_{\ell_0} +_{\epsilon'} b$. By Lemma 3.13, $\phi(a_{\ell_0+1}) = \phi(a_{\ell_0}) + 1 = \ell_0 + 1$.

To see that $i(a_{\ell_0+1}, c_i) = i(a, c_i)$, we appeal to the *bigon criterion* of [7, Proposition 1.7]. Choose representative curves for a, c_1, \dots, c_k , pairwise in minimal position. The bigon criterion asserts that a, c_i are in minimal position if and only if the configuration $a \cup c_i$ does not bound any *bigons*, i.e. an embedded disk whose boundary is the union of an arc of a and an arc of c_i meeting in exactly two points. The curve-arc sum a_{ℓ_0+1} meets each c_i in exactly the same set of points as a_{ℓ_0} . To conclude, it thus suffices to see that no bigons were introduced by the summing procedure. The only arc of a_{ℓ_0+1}

that is not also an arc of a_{ℓ_0} is the one along which the summing procedure is performed; denote the original arc of α_{ℓ_0} by α and the modified arc by α' . Suppose that there is an arc γ of c_i such that $\alpha' \cup \gamma$ bounds a bigon. As $\alpha' = \alpha +_{\epsilon'} b$, it must be the case that the curve $\alpha \cup \gamma$ is isotopic to b . But by assumption, $b \subset T$ is not boundary-parallel, so this cannot happen.

To construct a_ℓ for $\ell \in \mathbb{Z}/r\mathbb{Z}$ arbitrary, one simply repeats the above construction, producing, for any $t \geq 0$, a curve a_{ℓ_0+t} with the same intersection properties as a_{ℓ_0} and satisfying $\phi(a_{\ell_0+t}) = \ell_0 + t$. □

For the remaining corollaries of Proposition 4.2, we re-instate the requirement that r be odd.

Corollary 4.5 *Let r be an odd integer and let ϕ be a $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure on a surface S . Let $S' \subset S$ be a subsurface of genus $h \geq 1$, and suppose that if $h = 1$, then S' includes some boundary component of signature k such that $k + 1$ generates $\mathbb{Z}/r\mathbb{Z}$.*

- (1) *For all $x \in \mathbb{Z}/r\mathbb{Z}$, there exists some nonseparating curve c supported on S' satisfying $\phi(c) = x$,*
- (2) *For any $2h$ -tuple $(i_1, j_1, \dots, i_h, j_h)$ of elements of $\mathbb{Z}/r\mathbb{Z}$, there is some geometric symplectic basis $\mathcal{B} = \{a_1, b_1, \dots, a_h, b_h\}$ for S' with $\phi(a_\ell) = i_\ell$ and $\phi(b_\ell) = j_\ell$ for all $1 \leq \ell \leq h$,*
- (3) *For any $2h$ -tuple (k_1, \dots, k_{2h}) of elements of $\mathbb{Z}/r\mathbb{Z}$, there is some chain (a_1, \dots, a_{2h}) of curves on S' such that $\phi(a_\ell) = k_\ell$ for all $1 \leq \ell \leq 2h$.*

Proof Certainly (1) follows from (2). To establish (3), choose any geometric symplectic basis $\mathcal{B} = \{a'_\ell, b'_\ell\}$ on S' . There is *some* spin structure ψ on S' for which $\psi(a'_\ell) = i_\ell$ and $\psi(b'_\ell) = j_\ell$. By Proposition 4.2, there is a diffeomorphism f of S' such that $f^*(\psi) = \phi$. Then $\mathcal{B} = f^{-1}(\mathcal{B}')$ has the required properties.

We will deduce (3) from (2). Given the $2h$ -tuple (k_1, \dots, k_{2h}) , define a $2h$ -tuple $(i_1, j_1, \dots, i_h, j_h)$ as follows: set $i_\ell = 1 - \ell + \sum_{t=1}^\ell k_{2t-1}$, and set $j_\ell = k_{2\ell}$. By (2), there exists a geometric symplectic basis $\mathcal{B} = \{c_\ell, d_\ell\}$ on S' whose ϕ -values realize the tuple $(i_1, j_1, \dots, i_h, j_h)$. Any geometric symplectic basis can be “completed” into a chain as follows: for $\ell = 1, \dots, h - 1$, let f_ℓ be a simple closed curve satisfying $i(f_\ell, d_\ell) = i(f_\ell, d_{\ell+1}) = 1$ and $i(f_\ell, x) = 0$ for all other elements $x \in \mathcal{B}$. As \mathcal{B} is a geometric symplectic basis, this imposes the homological relation $[f_\ell] = [c_{\ell+1}] - [c_\ell]$, and the intersection conditions imposed on the set of curves $\{f_\ell\}$ imply that this homology is realized geometrically: $c_\ell \cup f_\ell \cup c_{\ell+1}$ must bound a pair of pants P_ℓ for each $\ell = 1, \dots, h - 1$. The orientations can be arranged so that P_ℓ lies to the right of c_ℓ and f_ℓ and to the left of $c_{\ell+1}$.

Applying the homological coherence property to each P_ℓ , it follows that $\phi(f_\ell) = k_{2\ell+1}$. By construction, the curves $c_1, d_1, f_1, d_2, f_2, d_3, \dots, f_{h-1}, d_h$

form a chain of length $2h$; denote this chain by C . By construction, $\phi(c_1) = i_1 = k_1$, and $\phi(d_\ell) = k_{2\ell}$. Altogether, this shows that C has the required properties. \square

4.2 Even r

Following the discussion in Sect. 3.4, we see that the Arf invariant distinguishes at least two orbits of $\text{Mod}(\Sigma_g)$ on the set of $\mathbb{Z}/r\mathbb{Z}$ -valued spin structures. To see that there are *exactly* two orbits, in Definition 4.6 we formulate two “model” $\mathbb{Z}/r\mathbb{Z}$ -valued spin structures $\phi_{\text{even}}^{\mathcal{B}}$ and $\phi_{\text{odd}}^{\mathcal{B}}$ of prescribed Arf invariant, and in Proposition 4.9 we show that every $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure is equivalent to one of $\phi_{\text{even}}^{\mathcal{B}}$ or $\phi_{\text{odd}}^{\mathcal{B}}$. We restrict attention here to the case where the surface S has at most one boundary component. The general setting of multiple boundary components introduces considerable subtlety owing to the failure for the intersection pairing to determine a symplectic form, and our results require only the case of at most one boundary component.

Definition 4.6 Let S be a surface of genus $g \geq 1$ with at most one boundary component. Fix a geometric symplectic basis $\mathcal{B} = \{\alpha_1, \beta_1, \dots, \alpha_g, \beta_g\}$. Define $\phi_{\text{even}}^{\mathcal{B}}$ and $\phi_{\text{odd}}^{\mathcal{B}}$ as the $\mathbb{Z}/r\mathbb{Z}$ -valued spin structures such that $\phi_{\text{even}}^{\mathcal{B}}(\gamma) = \phi_{\text{odd}}^{\mathcal{B}}(\gamma) = 0$ for all $\gamma \in \mathcal{B} \setminus \{\beta_g\}$, and where $\phi_{\text{even}}^{\mathcal{B}}(\beta_g)$ and $\phi_{\text{odd}}^{\mathcal{B}}(\beta_g)$ are chosen to be 0 or 1 as necessary so that $\text{Arf}(\phi_{\text{even}}^{\mathcal{B}}) = 0$ and $\text{Arf}(\phi_{\text{odd}}^{\mathcal{B}}) = 1$.

In spite of the evident dependence on geometric symplectic basis, as \mathcal{B} ranges over the set of all geometric symplectic bases, the elements $\phi_{\text{odd}}^{\mathcal{B}}$ lie in a single orbit of $\text{Mod}(S)$ (and the same is also true of $\phi_{\text{even}}^{\mathcal{B}}$). The following is immediate via the change-of-coordinates principle.

Lemma 4.7 Let \mathcal{B} and \mathcal{B}' be geometric symplectic bases. Then there is a diffeomorphism $f: S \rightarrow S$ such that $f(\mathcal{B}) = \mathcal{B}'$. Consequently, $f^*(\phi_{\text{even}}^{\mathcal{B}'}) = \phi_{\text{even}}^{\mathcal{B}}$ and $f^*(\phi_{\text{odd}}^{\mathcal{B}'}) = \phi_{\text{odd}}^{\mathcal{B}}$.

Definition 4.8 Let S be a surface of genus $g \geq 1$ with at most one boundary component endowed with a $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure ϕ . We say that ϕ is *even* if there is a geometric symplectic basis \mathcal{B} such that $\phi = \phi_{\text{even}}^{\mathcal{B}}$, and we say that ϕ is *odd* if $\phi = \phi_{\text{odd}}^{\mathcal{B}}$.

Proposition 4.9 Fix an even integer r . Let S be a surface of genus $g \geq 2$ with at most one boundary component. Let ϕ be a $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure on S . Then in the sense of Definition 4.8, either ϕ is even, or else ϕ is odd.

Proof The argument makes use of the techniques of the proof of Proposition 4.2. Let $\mathcal{B} = \{\alpha_1, \beta_1, \dots, \alpha_g, \beta_g\}$ be a geometric symplectic basis, and let S_i denote the genus-1 subsurface determined by α_i, β_i ; define D_i as the boundary

curve of S_i . Exactly as in Proposition 4.2, each pair α_i, β_i can be replaced by new curves α'_i, β'_i supported on S_i and satisfying $i(\alpha'_i, \beta'_i) = 1$, such that α'_i is admissible. Denote the corresponding geometric symplectic basis by \mathcal{B}' . For an arc ϵ connecting β'_1 to D_2 and disjoint from all other D_i , the curve-arc sum $\beta'_1 +_\epsilon D_2$ satisfies $\phi(\beta'_1 +_\epsilon D_2) = \phi(\beta'_1) + 2$. By repeatedly performing this curve-arc sum using an arc ϵ disjoint from $\mathcal{B}' \setminus \{\beta'_2\}$ (as in Proposition 4.2), β'_2 can be replaced with a curve β''_2 such that $\phi(\beta''_2) = 0$ or 1 . By performing an analogous operation on all β'_i , one obtains a geometric symplectic basis $\mathcal{B}'' = \{\alpha'_1, \beta''_1, \dots, \alpha'_g, \beta''_g\}$ such that $\phi(\alpha'_i) = 0$ and $\phi(\beta''_i) = 0$ or 1 .

It remains to further alter each $\beta''_1, \dots, \beta''_{g-1}$ so that $\phi(\beta''_i) = 0$ in this range. For $1 \leq i \leq g - 1$, let γ_i be a collection of disjoint curves such that $\beta_1, \gamma_1, \dots, \beta_{g-1}, \gamma_{g-1}, \beta_g$ forms a chain of length $2g - 1$, and such that each γ_i is disjoint from all α'_j . Then necessarily $\alpha_i, \gamma_i, \alpha_{i+1}$ forms a pair of pants, and so $\phi(\gamma_i) = -1$. If $\phi(\beta''_i) = 1$, then $\phi(T_{\gamma_i}(\beta''_i)) = 0$. Replace β''_1, β''_2 by $T_{\gamma_1}(\beta''_1), T_{\gamma_1}(\beta''_2)$, respectively. Repeat, applying $T_{\gamma_2}^k$ to $T_{\gamma_1}(\mathcal{B}'')$ for k such that $\phi(T_{\gamma_2}^k T_{\gamma_1}(\beta''_2)) = 0$. Proceed in this way, taking each β''_i for $i \leq g - 1$ to some β'''_i with $\phi(\beta'''_i) = 0$. At the end, the geometric symplectic basis $\mathcal{B}''' = \{\alpha'_1, \beta'''_1, \dots, \alpha'_g, \beta'''_g\}$ will satisfy $\phi(\gamma) = 0$ for all $\gamma \in \mathcal{B}'''$ except possibly $\gamma = \beta'''_g$. By repeating the curve-arc summing procedure, β'''_g can be altered to satisfy $\psi(\beta'''_g) = 0$ or 1 as required. Define $\tilde{\mathcal{B}}$ to be this geometric symplectic basis. Applying Theorem 3.9, we see that $\phi = \phi_{\tilde{\mathcal{B}}}^{\text{even}}$ or $\phi_{\tilde{\mathcal{B}}}^{\text{odd}}$ as required. \square

There is an analogue of Corollary 4.5 for r even, although the Arf invariant provides an obstruction that was not present in the case of odd r .

Corollary 4.10 *Let r be an even integer, and let $S' \subset S$ be a subsurface of genus $h \geq 2$ endowed with a $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure ϕ . Then the following assertions hold:*

- (1) *For all $x \in \mathbb{Z}/r\mathbb{Z}$, there exists some nonseparating curve c supported on S' satisfying $\phi(c) = x$.*
- (2) *For a given $2h$ -tuple $(i_1, j_1, \dots, i_h, j_h)$ of elements of $\mathbb{Z}/r\mathbb{Z}$, there is some geometric symplectic basis $\mathcal{B} = \{a_1, b_1, \dots, a_h, b_h\}$ for S' with $\phi(a_\ell) = i_\ell$ and $\phi(b_\ell) = j_\ell$ for $1 \leq \ell \leq h$ if and only if the parity of the spin structure defined by these conditions agrees with the parity of the restriction $\phi|_{S'}$ to S' .*
- (3) *For any $(2h - 2)$ -tuple $(i_1, j_1, \dots, i_{h-1}, j_{h-1})$ of elements of $\mathbb{Z}/r\mathbb{Z}$, there is some geometric symplectic basis $\mathcal{B} = \{a_1, b_1, \dots, a_h, b_h\}$ for S' with $\phi(a_\ell) = i_\ell$ and $\phi(b_\ell) = j_\ell$ for $1 \leq \ell \leq h - 1$.*
- (4) *For a given $2h$ -tuple (k_1, \dots, k_{2h}) of elements of $\mathbb{Z}/r\mathbb{Z}$, there is some chain (a_1, \dots, a_{2h}) of curves on S' such that $\phi(a_\ell) = k_\ell$ for all $1 \leq \ell \leq 2h$*

if and only if the parity of the spin structure defined by these conditions agrees with the parity of the restriction $\phi|_{S'}$ to S' .

- (5) For any $(2h - 2)$ -tuple (k_1, \dots, k_{2h-2}) of elements of $\mathbb{Z}/r\mathbb{Z}$, there is some chain (a_1, \dots, a_{2h-2}) of curves on S' such that $\phi(a_\ell) = k_\ell$ for all $1 \leq \ell \leq 2h - 2$.

Proof The proof is essentially identical to that of Corollary 4.5. The arguments for (2) and (3) are slightly novel; the remaining points follow their counterparts in Corollary 4.5 verbatim. To establish (2), let $B' = \{a'_\ell, b'_\ell\}$ be a geometric symplectic basis on S' . Let S'' be a subsurface of S' containing each curve in B' that has only one boundary component. Given $(i_1, j_1, \dots, i_h, j_h)$, there is some spin structure ψ on S'' for which $\psi(a'_\ell) = i_\ell$ and $\psi(b'_\ell) = j_\ell$ for $1 \leq \ell \leq h$. By Proposition 4.9, there is an element $f \in \text{Mod}(S'')$ for which $f^*(\psi) = \phi$ if and only if the Arf invariants of ϕ and ψ agree; if they do, then $B = f^{-1}(B')$ has the required properties.

(3) will be obtained from (2). Let $\epsilon \in \mathbb{Z}/2\mathbb{Z}$ denote the Arf invariant of ϕ , and define the quantity

$$\eta = \sum_{\ell=1}^{h-1} (i_\ell + 1)(j_\ell + 1) \pmod{2}.$$

As the formula (7) for the Arf invariant shows, given any $(2h - 2)$ -tuple $(i_1, j_1, \dots, i_{h-1}, j_{h-1})$ and any value $\epsilon \in \mathbb{Z}/2\mathbb{Z}$, there is a choice of $i_h, j_h \in \mathbb{Z}/r\mathbb{Z}$ for which $\eta + (i_h + 1)(j_h + 1) \equiv \epsilon \pmod{2}$. The result now follows by applying (2) to the tuple $(i_1, j_1, \dots, i_h, j_h)$. □

We will also require a result establishing the existence of configurations \mathcal{D}_n as in the D_n relation (Proposition 2.6).

Corollary 4.11 *Let $r = 2d$ be an even integer, and let Σ_g be a closed surface endowed with a $\mathbb{Z}/2d\mathbb{Z}$ -valued spin structure ϕ . Let Δ be a curve on Σ_g that separates Σ_g into subsurfaces S_1, S_2 for which the genus $g(S_1) \geq d + 1$. Set $n = 2g(S_1) - 2d + 1$. Then there exists a configuration $a, a', c_1, \dots, c_{n-2}$ of curves on S_1 arranged in the \mathcal{D}_n configuration, such that the elements a, a' , and c_i are admissible for all i , and such that $\Delta = \Delta_2$ as in Fig. 2.*

Proof By Corollary 4.10.5, there exists a chain a, c_1, \dots, c_{n-2} of admissible curves on S_1 . Let $a' \subset S_1$ be chosen so that $a \cup a' \cup \Delta$ bounds a subsurface of genus $g(S_1) - d - 1$ containing c_i for $i \geq 2$, and such that $i(a', c_1) = 1$. The other side of $a \cup a'$ bounds a subsurface of genus d , and so the homological coherence property implies that a' is admissible. By construction, the curves $a, a', c_1, \dots, c_{n-2}$ form the configuration \mathcal{D}_n of the D_n relation, and the boundary component Δ_2 of Fig. 2 is given here by Δ . □

5 r odd: generating $\text{Mod}(\Sigma_g)[\phi]$ by Dehn twists

Let ϕ be a $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure on a closed surface Σ_g . Throughout this section we assume that $r \mid (2g - 2)$ (so that, following Remark 3.6, Σ_g admits a $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure) and that r is odd. Recall from Definition 3.16 that the admissible subgroup is defined via

$$\mathcal{T}_\phi = \langle T_a \mid a \text{ nonseparating curve}, \phi(a) = 0 \rangle.$$

By construction, $\mathcal{T}_\phi \leq \text{Mod}(\Sigma_g)[\phi]$. The main result of this section is that for r odd, this containment is an equality.

Proposition 5.1 *For any $g \geq 3$ and for any odd integer r satisfying $r < g - 1$, there is an equality*

$$\mathcal{T}_\phi = \text{Mod}(\Sigma_g)[\phi].$$

Before beginning with the proof, we will first establish some properties of the group \mathcal{T}_ϕ which will be used throughout this section and the next.

Lemma 5.2 *Let ϕ be a $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure on a surface Σ_g with $r < g - 1$ and $g \geq 5$. Let c be any nonseparating simple closed curve on Σ_g . Suppose that r is odd, or else that r is even and $\phi(c) \equiv 1 \pmod{2}$. Then $T_c^r \in \mathcal{T}_\phi$.*

Proof Let c be as in the statement of Lemma 5.2. Our first objective is to construct a configuration of admissible curves \mathcal{D}_{2r+3} as in Corollary 2.7 for which $c = C_k$. By hypothesis, there is an expression of the form $\phi(c) = 2k - 1 \pmod{r}$ for some integer $1 \leq k \leq r$. Invoking Corollary 4.5.3 or 4.10.5 as appropriate, the hypothesis $r < g - 1$ implies that there is a chain a, c_1, \dots, c_{2k-1} of admissible curves disjoint from c , and there is a chain $c_{2k+1}, \dots, c_{2r+1}$ of admissible curves disjoint from c and from a, c_1, \dots, c_{2k-1} . Let a' be a curve such that $a \cup a' \cup c$ bounds a surface of genus $k - 1$ containing c_2, \dots, c_{2k-1} , and satisfying $i(a', c_1) = 1$ and $i(a', c_i) = 0$ for $2k + 1 \leq i \leq 2r + 1$. The homological coherence property implies that a' is admissible.

To complete the construction, it remains only to find the curve c_{2k} . Such a curve c_{2k} must be admissible, and c_{2k} must have the following intersection properties:

$$\begin{aligned} i(c_{2k}, c_{2k\pm 1}) &= 1, & i(c_{2k}, a) &= i(c_{2k}, a') = i(c_{2k}, c_i) = 0 \text{ for } |i - 2k| > 1, \\ i(c_{2k}, c) &= 2. \end{aligned} \tag{8}$$

Let c'_{2k} be any curve satisfying the intersection properties (8). If we can show that the complement of a regular neighborhood of the configuration

$\mathcal{D}'_{2r+3} := a, a', c_1, \dots, c_{2k-1}, c_{2k'}, c_{2k+1}, \dots, c_{2r+1}$ is a surface of positive genus, then the existence of c_{2k} will follow from Corollary 4.4.

The configuration \mathcal{D}'_{2r+3} is contained in a surface of genus $r + 1$ with two boundary components. Each boundary component is homologous to the nonseparating curve c , so the complement has genus $g - r - 2$. We must show that this quantity is positive. Establishing $g - r - 2 \geq 1$ is a matter of simple arithmetic. Writing $r = \frac{2g-2}{m}$ for some $m \geq 3$, we have

$$g - r - 2 = \frac{m - 2}{m}(g - 1) - 1 \geq \frac{g}{3} - 1 > 0,$$

since $g \geq 5$ by hypothesis.

Recalling that the group H_{2r+3}^+ from Corollary 2.7 is defined to be the group generated by the Dehn twists about the elements of $\mathcal{D}_{2r+3} \cup \{\Delta_1\}$, it follows that if each element of \mathcal{D}_{2r+3} is admissible, then $H_{2r+3}^+ \leq \mathcal{T}_\phi$. We have constructed the curves $a, a', c_1, \dots, c_{2r+1}$ so as to be admissible; homological coherence implies that also Δ_1 is admissible. Corollary 2.7.2 then implies that $T_{C_k}^r \in \mathcal{T}_\phi$ for any $1 \leq k \leq r + 1$. □

Lemma 5.3 *Let ϕ be a $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure on a surface Σ_g , and let $v \in H_1(\Sigma_g; \mathbb{Z})$ be any primitive homology class. If r is odd, then for any $k \in \mathbb{Z}/r\mathbb{Z}$, there exists a curve c for which $[c] = v$ and $\phi(c) = k$. If r is even, then for any $k \in \mathbb{Z}/r\mathbb{Z}$ such that $\phi \pmod{2}(v) \equiv k \pmod{2}$, there exists a curve c for which $[c] = v$ and $\phi(c) = k$.*

Proof Let c_0 be any (oriented) curve on Σ_g with $[c_0] = v$; set $\phi(c_0) = k_0$. Let c_1 be a curve disjoint from c_0 such that $c_0 \cup c_1$ bounds a subsurface of genus 1, oriented to the left of c_0 . Then $\phi(c_1) = k_0 + 2$ when oriented with the subsurface to the right, and $[c_0] = [c_1]$. This construction can be repeated, giving rise to curves c_m with $\phi(c_m) = k_0 + 2m$. If r is odd, then the set of values $k_0 + 2m$ for various values of m exhausts $\mathbb{Z}/r\mathbb{Z}$, and if r is even, then the set of values $k_0 + 2m$ exhausts the coset $k_0 + 2\mathbb{Z}/r\mathbb{Z}$. The claim follows by taking $c = c_m$ for the appropriate value of m . □

Proof of Proposition 5.1 The method is to compare the intersections of \mathcal{T}_ϕ and $\text{Mod}(\Sigma_g)[\phi]$ with \mathcal{I}_g and \mathcal{K}_g . We first present a high-level overview of the logical structure of the proof that explains how Proposition 5.1 follows from ancillary results; these results are then obtained in Steps 1–4.

Overview Recall from (3) the symplectic representation $\Psi: \text{Mod}(\Sigma_g) \rightarrow \text{Sp}(2g, \mathbb{Z})$ with kernel given by the Torelli group \mathcal{I}_g . To show that $\mathcal{T}_\phi = \text{Mod}(\Sigma_g)[\phi]$, it suffices to show that (I) $\Psi(\mathcal{T}_\phi) = \Psi(\text{Mod}(\Sigma_g)[\phi])$ and that (II) $\mathcal{T}_\phi \cap \mathcal{I}_g = \text{Mod}(\Sigma_g)[\phi] \cap \mathcal{I}_g$.

The equality of (I) is obtained in Step 1 as Lemma 5.4. The proof of (II) is carried out in Steps 2–4. The method is to study the restriction of the Johnson homomorphism to the groups $\mathcal{T}_\phi \cap \mathcal{I}_g$ and $\text{Mod}(\Sigma_g)[\phi] \cap \mathcal{I}_g$. Recall from (4) that the Johnson homomorphism is the surjective homomorphism

$$\tau: \mathcal{I}_g \rightarrow \wedge^3 H_{\mathbb{Z}}/H_{\mathbb{Z}},$$

and that the kernel is written \mathcal{K}_g . To establish (II), it suffices to show that (i) $\tau(\mathcal{T}_\phi \cap \mathcal{I}_g) = \tau(\text{Mod}(\Sigma_g)[\phi] \cap \mathcal{I}_g)$ and that (ii) $\mathcal{T}_\phi \cap \mathcal{K}_g = \text{Mod}(\Sigma_g)[\phi] \cap \mathcal{K}_g$. The equality of (i) is carried in Steps 2 and 3. The main result of Step 2, Lemma 5.7, establishes an upper bound on the image $\tau(\text{Mod}(\Sigma_g)[\phi] \cap \mathcal{I}_g)$, and the main result of Step 3, Lemma 5.8, shows that the subgroup $\tau(\mathcal{T}_\phi \cap \mathcal{I}_g)$ realizes this upper bound. Finally (ii) is established in Step 4: Lemma 5.9 shows that there is a containment $\mathcal{K}_g \leq \mathcal{T}_\phi$.

Step 1: The symplectic quotient The first step is to understand the image of \mathcal{T}_ϕ and $\text{Mod}(\Sigma_g)[\phi]$ in the symplectic group $\text{Sp}(2g, \mathbb{Z})$.

Lemma 5.4 *For r odd, the symplectic representation $\Psi: \text{Mod}(\Sigma_g) \rightarrow \text{Sp}(2g, \mathbb{Z})$ restricts to a surjection*

$$\Psi: \mathcal{T}_\phi \twoheadrightarrow \text{Sp}(2g, \mathbb{Z}).$$

It follows that also $\Psi: \text{Mod}(\Sigma_g)[\phi] \twoheadrightarrow \text{Sp}(2g, \mathbb{Z})$ is a surjection.

Proof Let $v \in H_1(\Sigma_g; \mathbb{Z})$ be a primitive element. By Lemma 5.3, there is some curve c with $[c] = v$ and $\phi(c) = 0$. The result follows from this, since $\text{Sp}(2g, \mathbb{Z})$ is generated by the set of transvections T_v given by $x \mapsto x + \langle x, v \rangle v$ for $v \in H_1(\Sigma_g; \mathbb{Z})$ primitive, and $\Psi(T_c) = T_{[c]}$. □

Step 2: $\text{Mod}(\Sigma_g)[\phi]$ and the Johnson homomorphism Our next objective is Lemma 5.7 below. This concerns the image of $\text{Mod}(\Sigma_g)[\phi] \cap \mathcal{I}_g$ under the Johnson homomorphism. In order to formulate the result, it is necessary to first study a different quotient of \mathcal{I}_g first constructed by Chillingworth [1, 2]. Chillingworth’s work is formulated using the notion of a “winding number function”; as explained in [9, Introduction], a winding number function is a particular instance of a spin structure. The properties of a winding number function that Chillingworth exploits in his work are common to all spin structures, and so we formulate his results in this larger context. See also [10, Section 5] for a brief summary of Chillingworth’s work. Recall in the statement below that \mathcal{S} is defined to be the set of isotopy classes of oriented simple closed curves on a surface Σ_g .

Theorem 5.5 (Chillingworth) *Let ϕ be a $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure on a closed surface Σ_g . Let c be the function $c: \mathcal{I}_g \times \mathcal{S} \rightarrow \mathbb{Z}/r\mathbb{Z}$ defined by the formula*

$$c(f, \gamma) = \phi(f(\gamma)) - \phi(\gamma).$$

Then the value $c(f, \gamma)$ depends only on the homology class $[\gamma] \in H_{\mathbb{Z}}$, and c descends to a homomorphism

$$c: \mathcal{I}_g \rightarrow \text{Hom}(H_{\mathbb{Z}}, \mathbb{Z}/r\mathbb{Z}) \cong H^1(\Sigma_g; \mathbb{Z}/r\mathbb{Z}).$$

In particular, c does not depend on the choice of $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure.

In [10], Johnson related Chillingworth’s homomorphism to the Johnson homomorphism. To formulate the precise connection, we require the following well-known lemma; see e.g [10, Sections 5,6].

Lemma 5.6 *There is a $\text{Sp}(2g, \mathbb{Z})$ -equivariant surjection*

$$C: \wedge^3 H_{\mathbb{Z}}/H_{\mathbb{Z}} \rightarrow H_{\mathbb{Z}/(g-1)\mathbb{Z}}$$

given by the “contraction”

$$C(x \wedge y \wedge z) = \langle x, y \rangle z + \langle y, z \rangle x + \langle z, x \rangle y \pmod{g-1}. \tag{9}$$

It follows that for any $r \mid (g-1)$, there is a $\text{Sp}(2g, \mathbb{Z})$ -equivariant surjection

$$C_r: \wedge^3 H_{\mathbb{Z}}/H_{\mathbb{Z}} \rightarrow H_{\mathbb{Z}/r\mathbb{Z}}$$

given by post-composing C with the reduction mod r . We can now formulate the main result of Step 3.

Lemma 5.7 *Let ϕ be a $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure on a surface of genus g , with $g \geq 3$ and r odd. Then $C_r \circ \tau = 0$ on $\text{Mod}(\Sigma_g)[\phi] \cap \mathcal{I}_g$.*

Proof According to [10, Theorem 3], the composition $C_r \circ \tau$ coincides (up to an application of Poincaré duality) with the mod- r Chillingworth homomorphism $c: \mathcal{I}_g \rightarrow H^1(\Sigma_g; \mathbb{Z}/r\mathbb{Z})$. The formula for c given in Theorem 5.5 shows that c measures how $f \in \mathcal{I}_g$ alters the set of values $\{\phi(\gamma) \mid \gamma \in \mathcal{S}\}$; it therefore follows immediately that the restriction of c to $\text{Mod}(\Sigma_g)[\phi] \cap \mathcal{I}_g$ is trivial. \square

Step 3 \mathcal{T}_{ϕ} and the Johnson homomorphism In the previous step, we showed that there is a containment

$$\tau(\text{Mod}(\Sigma_g)[\phi] \cap \mathcal{I}_g) \leq \ker(C_r \circ \tau).$$

Our next result establishes that this containment is an equality, even when restricted to the subgroup $\tau(\mathcal{T}_\phi \cap \mathcal{I}_g)$.

Lemma 5.8 *For $r < g - 1$ odd and for $g \geq 3$, the Johnson homomorphism τ gives a surjection*

$$\tau: \mathcal{T}_\phi \cap \mathcal{I}_g \twoheadrightarrow \ker(C_r \circ \tau).$$

Proof Define $K := \ker(C_r)$. We must show that $\mathcal{T}_\phi \cap \mathcal{I}_g$ surjects onto K under τ . The first step will be to determine a generating set for K , and then we will exhibit each generator within $\tau(\mathcal{T}_\phi \cap \mathcal{I}_g)$.

To determine a generating set for K , we consider the short exact sequence

$$1 \rightarrow K \rightarrow \wedge^3 H_{\mathbb{Z}}/H_{\mathbb{Z}} \rightarrow H_{\mathbb{Z}/r\mathbb{Z}} \rightarrow 1.$$

determined by C_r . By lifting a set of relations $\{r_i\}$ for $H_{\mathbb{Z}/r\mathbb{Z}}$ to $\wedge^3 H_{\mathbb{Z}}/H_{\mathbb{Z}}$, we will obtain a set of generators $\{\tilde{r}_i\}$ for K . Let $\mathcal{B} = \{x_1, y_1, \dots, x_g, y_g\}$ be a symplectic basis for $H_{\mathbb{Z}}$. There is an associated basis $\wedge^3 \mathcal{B} \subset \wedge^3 H_{\mathbb{Z}}$ given by

$$\wedge^3 \mathcal{B} := \{z_1 \wedge z_2 \wedge z_3 \mid z_i \in \mathcal{B} \text{ distinct}\}$$

Thus also $\wedge^3 H_{\mathbb{Z}}/H_{\mathbb{Z}}$ is generated by the image of $\wedge^3 \mathcal{B}$.

To determine the relations r_i , we must understand $C_r(z_1 \wedge z_2 \wedge z_3)$ for the various possibilities for $\{z_1, z_2, z_3\}$. There are two orbits of generators under the action of $\text{Sp}(2g, \mathbb{Z})$. The first orbit consists of elements of the form $z \wedge x_i \wedge y_i$ (necessarily with $z \neq x_i, y_i$), and the second orbit consists of elements of the form $z_i \wedge z_j \wedge z_k$ with each $z_\ell \in \{x_\ell, y_\ell\}$ and with i, j, k mutually distinct.

The image of $z \wedge x_i \wedge y_i$ in $H_{\mathbb{Z}/r\mathbb{Z}}$ is

$$C_r(z \wedge x_i \wedge y_i) = z,$$


while $C_r(z_i \wedge z_j \wedge z_k) = 0$ for elements of the second type. Define A to be the abelian group generated by the symbols $C_r(z_1 \wedge z_2 \wedge z_3)$ for $z_1 \wedge z_2 \wedge z_3 \in \wedge^3 \mathcal{B}$, subject to the relations (R1)–(R3) below:

- (R1) $rC_r(z \wedge x_i \wedge y_i) = 0$
- (R2) $C_r(z \wedge x_i \wedge y_i) - C_r(z \wedge x_j \wedge y_j) = 0$
- (R3) $C_r(z_i \wedge z_j \wedge z_k) = 0$ for $\{i, j, k\} \subset \{1, \dots, g\}$ distinct.

It can be easily verified that there is an isomorphism $A \cong H_{\mathbb{Z}/r\mathbb{Z}}$, so that the relations (R1)–(R3) can be lifted to $\wedge^3 H_{\mathbb{Z}}/H_{\mathbb{Z}}$ to give a generating set for K as desired. The corresponding generators are given below.

- (G1) $rz \wedge x_i \wedge y_i$

Fig. 5 The configuration of curves used to exhibit (G2)



- (G2) $z \wedge (x_i \wedge y_i - x_j \wedge y_j)$
- (G3) $z_i \wedge z_j \wedge z_k$ for $\{i, j, k\} \subset \{1, \dots, g\}$ distinct.

Having determined a generating set for K , it remains to exhibit each such generator in the form $\tau(f)$ for $f \in \mathcal{T}_\phi \cap \mathcal{I}_g$. These will be handled on a case-by-case basis. We start with (G1). By Lemma 2.8, there exist curves c, d that determine a genus-1 bounding pair map with $\tau(T_c T_d^{-1}) = z \wedge x_i \wedge y_i$. By Lemma 5.2, $T_c^r, T_d^r \in \mathcal{T}_\phi$, so that $T_c^r T_d^{-r} \in \mathcal{T}_\phi$ is an element with the required properties.

Next we consider (G2). Let c be a curve with $[c] = z$ and $\phi(c) = 0$. By the change-of-coordinates principle, there exist curves a, b with the following properties: (1) $a \cup b$ bounds a subsurface S of genus 2, (2) $c \subset S$, (3) $[a] = [b] = [c]$, and c separates S into two subsurfaces S_1, S_2 each of genus 1, (4) x_i, y_i determine a symplectic basis for S_1 and x_j, y_j determine a symplectic basis for S_2 . Such a configuration is shown in Fig. 5. By homological coherence, $\phi(a) = \phi(b) = -2$ when a, b are oriented with S to the left. By Lemma 2.8,

$$\tau(T_a T_c^{-1}) = z \wedge x_i \wedge y_i$$

and

$$\tau(T_b T_c^{-1}) = -z \wedge x_j \wedge y_j.$$

Therefore, it is necessary to show $T_a T_b T_c^{-2} \in \mathcal{T}_\phi$. By hypothesis, $T_c \in \mathcal{T}_\phi$. By Corollary 4.5.3, there exists a chain a_1, \dots, a_5 of curves on S for which $\phi(a_i) = 0$. By the chain relation (Proposition 2.4), $T_a T_b \in \mathcal{T}_\phi$, and the result follows.

It remains to exhibit generators of type (G3). Any such generator is equivalent under the action of $\text{Sp}(2g, \mathbb{Z})$ to $y_1 \wedge y_2 \wedge y_3$. By combining the $\text{Sp}(2g, \mathbb{Z})$ -equivariance of τ (Lemma 2.8.1) with the result of Lemma 5.4, it suffices to exhibit only $y_1 \wedge y_2 \wedge y_3$. Figure 6 shows the two 3-chains $C_1 = (c_6 1, c_2, c_3)$ and $C_2 = (c'_1, c_2, c'_1 + \epsilon d)$. Observe that d is a boundary




Fig. 6 Top: The relevant portion of the geometric symplectic basis \mathcal{B} . Bottom: The configuration of curves used to exhibit (G3). Orientations have been suppressed wherever possible

component for regular neighborhoods of both C_1 and C_2 ; let e_1, e_2 denote the other boundary component of C_1, C_2 , respectively.

By Corollary 4.5.2, there exists a geometric symplectic basis \mathcal{B} that contains the elements c_1, c_2, b, f as depicted in the top portion of Fig. 6, with homology classes and ϕ -values given in the table below. The remaining entries in the table have been filled in using the homological coherence property. (A value of $*$ indicates that the value is irrelevant and/or underdetermined, and if an orientation is left unspecified, this is in accordance with Convention 3.4).

Curve:	c_1	c_2	c_3	c'_1	b	f	d	$c'_1 + \epsilon d$
Homology class:	x_1	y_1	$y_2 - x_1$	$x_1 - y_3$	y_3	$-y_2$	y_2	$y_2 + y_3 - x_1$
ϕ -value:	0	*	0	*	*	-1	-2	*

By Lemma 2.8,

$$\begin{aligned} \tau(T_d T_{e_1}^{-1}) &= x_1 \wedge y_1 \wedge y_2, \\ \tau(T_d T_{e_2}^{-1}) &= (x_1 - y_3) \wedge y_1 \wedge y_2. \end{aligned}$$

It follows that $\tau(T_{e_1}^{-1} T_{e_2}) = y_3 \wedge y_1 \wedge y_2$. As $d \cup e_1$ and $d \cup e_2$ each bound subsurfaces of genus 1 and $\phi(d) = -2$ when d is oriented with these subsurfaces to the left, the homological coherence property implies that e_1 and e_2 are admissible. The result follows. □

Step 4: The Johnson kernel The final piece of the analysis concerns the relationship between $\text{Mod}(\Sigma_g)[\phi]$ and the Johnson kernel \mathcal{K}_g .

Lemma 5.9 *Let ϕ be a $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure with r odd. If $g \geq 3$, then \mathcal{T}_ϕ contains the Johnson kernel \mathcal{K}_g . It follows that also*

$$\mathcal{K}_g \leq \text{Mod}(\Sigma_g)[\phi].$$

Proof According to Johnson’s Theorem 2.9, \mathcal{K}_g has a generating set consisting of the set of all T_c for c a separating curve. Each such c divides Σ_g into two subsurfaces S, S' , and since $g \geq 3$, without loss of generality we can assume that $g(S) > 1$. By Corollary 4.5.3, there exists a chain $a_1, \dots, a_{2g(S)}$ of curves on S such that $\phi(a_i) = 0$ for all i . By hypothesis, $T_{a_i} \in \mathcal{T}_\phi$ for all i . By the chain relation (Proposition 2.4), it follows that $T_c \in \mathcal{T}_\phi$ as required. □

This concludes the proof of Proposition 5.1. □

6 r even: \mathcal{T}_ϕ has finite index in $\text{Mod}(\Sigma_g)$

We continue to assume that $r \mid (2g - 2)$, but now we take $r = 2d$ to be even. For r even, we cannot give a complete characterization of \mathcal{T}_ϕ as in Proposition 5.1, but we will show that \mathcal{T}_ϕ has finite index in $\text{Mod}(\Sigma_g)$. The minimal genus for which the ensuing arguments apply has a rather intricate dependence on r , encapsulated in the definition below.

Definition 6.1 For an integer $d \geq 1$, define $k(d)$ as follows:

$$k(d) = \begin{cases} 2 & d \text{ odd or } d \geq 6 \text{ even} \\ 6 & d = 2 \\ 5 & d = 4. \end{cases}$$

Suppose $r = 2d$ is an even integer. Define

$$g(r) = k(d)d + 1.$$

Proposition 6.2 *Let $r = 2d$ be an even integer. Suppose $g \geq g(r)$ and that $r < g - 1$. Then \mathcal{T}_ϕ is a finite-index subgroup of $\text{Mod}(\Sigma_g)$.*

The presence of an underlying $\mathbb{Z}/2\mathbb{Z}$ spin structure makes proving the analogues of Lemmas 5.8 and 5.9 substantially more difficult. At present, we do not know how to establish the analogue of Lemma 5.8, owing to the fact that the Arf invariant provides an obstruction to finding the configurations of curves on subsurfaces needed for the arguments therein. Thus we content ourselves with showing that $\mathcal{T}_\phi \leq \text{Mod}(\Sigma_g)$ is finite-index.

Proof of Proposition 6.2 The proof of Proposition 6.2 follows a similar outline to that of Proposition 5.1. We begin with an overview of the proof.

Overview To establish finiteness of the index $[\text{Mod}(\Sigma_g) : \mathcal{T}_\phi]$, it suffices to show that the indices $[\text{Sp}(2g, \mathbb{Z}) : \Psi(\mathcal{T}_\phi)]$ and $[\mathcal{I}_g : \mathcal{T}_\phi \cap \mathcal{I}_g]$ are both finite. Finiteness of $[\text{Sp}(2g, \mathbb{Z}) : \Psi(\mathcal{T}_\phi)]$ is established in Lemma 6.4 of Step 1, which moreover gives a complete description of the subgroup $\Psi(\mathcal{T}_\phi)$.

Finiteness of $[\mathcal{I}_g : \mathcal{T}_\phi \cap \mathcal{I}_g]$ is obtained in Steps 2 and 3, again by using the Johnson homomorphism to analyze the intersection $\mathcal{T}_\phi \cap \mathcal{I}_g$ as in Steps 2-4 of the proof of Proposition 5.1. The main result of Step 2 is Lemma 6.6, which shows that $\tau(\mathcal{T}_\phi \cap \mathcal{I}_g)$ has finite index in $\wedge^3 H_{\mathbb{Z}}/H_{\mathbb{Z}}$. Step 3 completes the argument by showing the containment $\mathcal{K}_g \leq \mathcal{T}_\phi$; this is obtained as Lemma 6.7. We advise the reader that Step 3 is substantially more complicated than its counterpart Step 4 of the proof of Proposition 5.1, and will require an explanatory outline of its own.

Step 1: The symplectic quotient The case of r even is no more difficult than for r odd. Let q be a $\mathbb{Z}/2\mathbb{Z}$ -valued spin structure. An *anisotropic transvection* is a transvection

$$T_v(w) = w + \langle w, v \rangle v$$

for a primitive $v \in H_1(\Sigma_g; \mathbb{Z})$ such that $q(v) = 0$.

The following theorem is surely well-known to experts but we were unable to find a reference. A special case is treated in [5, Proposition 14].

Theorem 6.3 (Folklore) *Let q be a $\mathbb{Z}/2\mathbb{Z}$ -valued spin structure on Σ_g for $g \geq 3$, and let $\text{Sp}(2g, \mathbb{Z})[q]$ denote the subgroup of $\text{Sp}(2g, \mathbb{Z})$ that fixes q . Then $\text{Sp}(2g, \mathbb{Z})[q]$ is generated by the collection of anisotropic transvections*

$$\{T_v \mid v \in H_1(\Sigma_g; \mathbb{Z}) \text{ primitive, } q(v) = 0\}.$$

Proof The action of $\text{Sp}(2g, \mathbb{Z})$ on the set of spin structures factors through the quotient $f: \text{Sp}(2g, \mathbb{Z}) \rightarrow \text{Sp}(2g, \mathbb{Z}/2\mathbb{Z})$. Define $\text{Sp}(2g, \mathbb{Z})[2] := \ker(f)$. Thus, there is a short exact sequence

$$1 \rightarrow \mathrm{Sp}(2g, \mathbb{Z})[2] \rightarrow \mathrm{Sp}(2g, \mathbb{Z})[q] \rightarrow \mathrm{Sp}(2g, \mathbb{Z}/2\mathbb{Z})[q] \rightarrow 1,$$

with $\mathrm{Sp}(2g, \mathbb{Z}/2\mathbb{Z})[q]$ denoting the stabilizer of q in $\mathrm{Sp}(2g, \mathbb{Z}/2\mathbb{Z})$. According to [8, Theorem 14.16], the group $\mathrm{Sp}(2g, \mathbb{Z}/2\mathbb{Z})[q]$ is generated by the images of all anisotropic transvections. So it remains to see only that the subgroup of $\mathrm{Sp}(2g, \mathbb{Z})[q]$ generated by anisotropic transvections contains $\mathrm{Sp}(2g, \mathbb{Z})[2]$. According to [14, Lemma 5], the group $\mathrm{Sp}(2g, \mathbb{Z})[2]$ is generated by the collection of “square transvections” T_w^2 , where w ranges over all primitive $w \in H_1(\Sigma_g; \mathbb{Z})$.

If $q(w) = 0$ then $T_w \in \mathrm{Sp}(2g, \mathbb{Z})[q]$ and so there is nothing to show. Assume now that $q(w) = 1$. It is easy to produce (e.g. by the change-of-coordinates principle on Σ_g) vectors $v_1, v_2, v_3 \in H_1(\Sigma_g; \mathbb{Z})$ with the following properties:

- (1) $q(v_i) = 0$ for all i ,
- (2) $\langle v_1, v_2 \rangle = \langle v_2, v_3 \rangle = 1$ and $\langle v_1, v_3 \rangle = 0$,
- (3) $\langle v_i, w \rangle = 0$ for all i ,
- (4) $v_1 + v_3 = w$.

The chain relation in $\mathrm{Mod}(\Sigma_g)$ (Proposition 2.4) descends to show the relation

$$(T_{v_1} T_{v_2} T_{v_3})^4 = T_w^2.$$

Since the left-hand side is a product of anisotropic transvections, it follows that for w arbitrary, $T_w^2 \in \mathrm{Sp}(2g, \mathbb{Z})[q]$ as required. □

The following is the main result of Step 1.

Lemma 6.4 *Let ϕ be a $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure for r an even integer, and let*

$$q := \phi \pmod{2}$$

denote the associated $\mathbb{Z}/2\mathbb{Z}$ -valued spin structure. The symplectic representation $\Psi: \mathrm{Mod}(\Sigma_g) \rightarrow \mathrm{Sp}(2g, \mathbb{Z})$ restricts to a surjection

$$\Psi: \mathcal{T}_\phi \twoheadrightarrow \mathrm{Sp}(2g, \mathbb{Z})[q],$$

where $\mathrm{Sp}(2g, \mathbb{Z})[q]$ denotes the stabilizer of q in $\mathrm{Sp}(2g, \mathbb{Z})$.

Proof As \mathcal{T}_ϕ preserves the $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure ϕ , it also preserves the mod-2 reduction q . Thus $\mathcal{T}_\phi \leq \mathrm{Mod}(\Sigma_g)[q]$ and so $\Psi(\mathcal{T}_\phi) \leq \mathrm{Sp}(2g, \mathbb{Z})[q]$. Let $v \in H_1(\Sigma_g; \mathbb{Z})$ be a primitive element satisfying $q(v) = 0$. By Lemma 5.3, there is some curve c with $[c] = v$ and $\phi(c) = 0$. As $T_c \in \mathrm{Mod}(\Sigma_g)[\phi]$ and $\Psi(T_c) = T_v$, the result now follows from Theorem 6.3. □

Step 2: The Johnson homomorphism We remind the reader that while the value $\phi(c)$ on a simple closed curve depends on more than the homology class $[c] \in H_1(\Sigma_g; \mathbb{Z})$, the discussion of Sect. 3.4 establishes that the mod-2 reduction $q(c)$ *does* depend only on the homology class $[c]$ (indeed, the coefficients here can be taken to be $\mathbb{Z}/2\mathbb{Z}$). Thus the arguments in Step 3 can be carried out entirely in the homological setting.

For the duration of Step 2, we adopt the following notation. As usual, define

$$q := \phi \pmod{2}.$$

There exists a symplectic basis $\{x_1, y_1, \dots, x_g, y_g\}$ for $H_1(\Sigma_g; \mathbb{Z})$ such that $q(x_i) = 0$ for $1 \leq i \leq g$ and $q(y_j) = 0$ for $1 \leq j \leq g - 1$; with such a basis, $\text{Arf}(q)$ depends only on g and on $q(y_g)$.

Before proceeding to the main result of Step 2 (Lemma 6.6), we begin with an algebraic lemma.

Lemma 6.5 *Set $v := x_1 \wedge y_1 \wedge x_4$. Let $V \leq \wedge^3 H_{\mathbb{Z}}$ denote the submodule generated by the set*

$$\{gv \mid g \in \text{Sp}(2g, \mathbb{Z})[q]\}.$$

Then $V = \wedge^3 H_{\mathbb{Z}}$ for $g \geq 5$.

Proof As remarked in Lemma 5.8, $\wedge^3 H_{\mathbb{Z}}$ is generated by elements of the form $z_i \wedge z_j \wedge z_k$ with each $z_i \in \{x_1, y_1, \dots, x_g, y_g\}$. To begin with, we will exhibit generators for the submodule of $\wedge^3 H_{\mathbb{Z}}$ spanned by generators $z_i \wedge z_j \wedge z_k$ for which $z_i, z_j, z_k \in \{x_1, y_1, \dots, x_{g-1}, y_{g-1}\}$. The restriction of $\text{Sp}(2g; \mathbb{Z})[q]$ to this submodule is independent of the parity of q . For $i \neq j \leq g - 1$, define $S_{i,j} \in \text{Sp}(2g, \mathbb{Z})$ via

$$\begin{aligned} S_{i,j}(x_i) &= x_j, & S_{i,j}(y_i) &= y_j, \\ S_{i,j}(x_j) &= x_i, & S_{i,j}(y_j) &= y_i, \end{aligned}$$

with all other generators fixed. As $q(x_k) = q(y_k) = 0$ for $k \leq g - 1$, in fact $S_{i,j}$ is an element of $\text{Sp}(2g, \mathbb{Z})[q]$. Applying $S_{i,j}$ for $i, j \neq 4$ to v shows that V contains all generators of the form $x_i \wedge y_i \wedge x_4$. Applying $S_{i,4}$ to $x_j \wedge y_j \wedge x_4$ for $i \neq j$ shows that V contains all generators of the form $x_j \wedge y_j \wedge x_i$ for $j \neq 4$; then applying $S_{j,4}$ to $x_j \wedge y_j \wedge x_i$ shows that V contains all elements of the form $x_j \wedge y_j \wedge x_i$.

For $1 \leq i \leq g - 1$, define $R_i \in \text{Sp}(2g, \mathbb{Z})$ via

$$R_i(x_i) = y_i, \quad R_i(y_i) = -x_i$$

with all other generators fixed. Again, the condition $q(x_k) = q(y_k) = 0$ for $k \leq g - 1$ implies that R_i is an element of $\text{Sp}(2g, \mathbb{Z})[q]$. Applying R_i to $x_j \wedge y_j \wedge x_i$ shows that also V contains all elements of the form $x_j \wedge y_j \wedge y_i$.

It remains to exhibit generators of the form $z_i \wedge z_j \wedge z_k$ with $z_\ell \in \{x_\ell, y_\ell\}$ and $i, j, k < g$ all distinct. Consider the transvection $T_{x_4-x_1} \in \text{Sp}(2g, \mathbb{Z})[q]$. Applied to $x_1 \wedge y_1 \wedge x_2$, this shows that

$$x_1 \wedge (y_1 + x_4) \wedge x_2 \in V,$$

hence also $x_1 \wedge x_2 \wedge x_4 \in V$. Now by repeated applications of the elements $S_{i,j}$ and R_i , one can produce all remaining generators.

In the case $q(y_g) = 0$, the elements $S_{i,g}$ and R_g are contained in $\text{Sp}(2g, \mathbb{Z})[q]$, and so the above argument extends to complete this case. It remains to consider the case where $q(y_g) = 1$. In this case, the formula (5) defining a $\mathbb{Z}/2\mathbb{Z}$ -valued quadratic form shows that $q(y_{g-1} + y_g) = 0$. Applying $T_{y_{g-1}+y_g}$ to the elements $x_1 \wedge x_2 \wedge x_{g-1}$ and $x_1 \wedge y_1 \wedge x_{g-1}$ shows that $x_1 \wedge x_2 \wedge y_g$ and $x_1 \wedge y_1 \wedge x_g$ are elements of V . Applying $S_{i,j}$ and R_i for $i, j \leq g - 1$ produces all elements of the form $z_i \wedge z_j \wedge y_g$ with $z_\ell \in \{x_\ell, y_\ell\}$ ($i, j \leq g - 1$). Then applying T_{x_g} to these elements shows that also each $z_i \wedge z_j \wedge x_g \in V$.

By (5), we have $q(x_1 + x_g - y_g) = 0$. Applying $T_{x_1+x_g-y_g}^{-1}$ to $y_1 \wedge y_2 \wedge y_g$ gives

$$w = (y_1 + x_1 + x_g - y_g) \wedge y_2 \wedge (x_1 + x_g);$$

expanding this product yields the expression $w = -y_2 \wedge x_g \wedge y_g + w'$, with w' expressed entirely in terms of generators already known to be elements of V . Applying $S_{i,j}$ and R_i as in the above paragraph shows that all the remaining generators $z_i \wedge x_g \wedge y_g$ are elements of V . □

The following is the main result of Step 2.

Lemma 6.6 *For $g \geq 5$, the image $\tau(\mathcal{T}_\phi \cap \mathcal{I}_g)$ under the Johnson homomorphism is a finite-index subgroup of $\wedge^3 H_{\mathbb{Z}}/H_{\mathbb{Z}}$.*

Proof As stated in Lemma 2.8.1, the homomorphism $\tau: \mathcal{I}_g \rightarrow \wedge^3 H_{\mathbb{Z}}/H_{\mathbb{Z}}$ is $\text{Sp}(2g, \mathbb{Z})$ -equivariant. The strategy for the proof of Lemma 6.6 is to first exhibit a single nonzero element of $\tau(\mathcal{T}_\phi \cap \mathcal{I}_g)$, and then to exploit this equivariance.

By Corollary 4.10.5, there exists a 3-chain of admissible curves a_1, a_2, a_3 such that

$$[a_1] = x_1, \quad [a_2] = y_1, \quad [a_3] = x_4 - x_1.$$

Let ν be a regular neighborhood of this chain, and denote the boundary curves as b, b' . As a_1 and a_2 are admissible, homological coherence implies that

$\phi(b) = \phi(b') = -1$ when oriented so that v lies to the left of both b, b' . By Lemma 5.2, T_b^r is an element of \mathcal{T}_ϕ . It follows by the chain relation (Proposition 2.4) that the bounding pair map $T_b^r T_{b'}^{-r} \in \mathcal{T}_\phi$. One sees that $[b] = [a_1] + [a_3] = x_4$. By Lemma 2.8.3,

$$\tau(T_b^r T_{b'}^{-r}) = r(x_1 \wedge y_1 \wedge x_4).$$

By Lemma 6.4 and the equivariance of τ with respect to $\text{Sp}(2g, \mathbb{Z})$ (and *a fortiori* with respect to $\text{Sp}(2g, \mathbb{Z})[q]$), it follows that $\tau(\mathcal{T}_\phi \cap \mathcal{I}_g)$ contains the \mathbb{Z} -span of the entire $\text{Sp}(2g, \mathbb{Z})[q]$ -orbit of $v := r(x_1 \wedge y_1 \wedge x_4)$. Lemma 6.6 now follows from Lemma 6.5. □

Step 3: The Johnson kernel In this section, we establish the following result.

Lemma 6.7 *Let ϕ be a $\mathbb{Z}/2d\mathbb{Z}$ -valued spin structure on Σ_g . Assume that g satisfies the hypotheses of Proposition 6.2. Then \mathcal{T}_ϕ contains the Johnson kernel \mathcal{K}_g .*

Before beginning the proof, we explain the difficulties imposed by the assumption that $r = 2d$ is even.

The Arf invariant as obstruction The mechanism of proof for Lemma 5.9 was the chain relation (Proposition 2.4): if $S \subset \Sigma_g$ has one boundary component, we exploited Corollary 4.5 to produce a maximal chain $\{a_i\}$ of curves on S with $\phi(a_i) = 0$, and then used the chain relation to express $T_{\partial S}$ in terms of the admissible twists $\{T_{a_i}\}$. Now suppose ϕ is a $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure for r even, and let $q = \phi \pmod{2}$ denote the mod-2 reduction. For any subsurface $S \subset \Sigma_g$ with one boundary component, q restricts to give a $\mathbb{Z}/2\mathbb{Z}$ -valued spin structure $q|_S$ on S . The Arf invariant of $q|_S$, written here as $\epsilon(S)$, provides an obstruction to the existence of a maximal chain $\{a_i\}$ of admissible curves on S , since such a chain determines the value $\epsilon(S)$ solely as a function of $g(S)$.

Suppose $c \subset \Sigma_g$ is a separating curve that divides Σ_g into disjoint surfaces S, S' . Such a c is called *easy* if at least one of S, S' supports a maximal chain of admissible curves, and is *hard* otherwise. By Corollary 4.10.4 and the chain relation (Proposition 2.4), if c is easy, then $T_c \in \mathcal{T}_\phi$.

Outline of proof of Lemma 6.7 We begin with Lemma 6.8, which characterizes those subsurfaces supporting a maximal chain of admissible curves in terms of the Arf invariant. This in particular shows the relevance of the genus of the subsurface mod 4, which in turn forces us to treat the cases $r \equiv 0, r \equiv 2 \pmod{4}$ separately. We therefore establish Lemma 6.7 by combining Lemmas 6.10 and 6.13, which treat the cases of $r \equiv 2 \pmod{4}$ and $r \equiv 0 \pmod{4}$, respectively.

These are handled in Substeps 1 and 2, respectively. In each case, we first show that all separating twists of particular genera are elements of \mathcal{T}_ϕ . In Substep 1, Lemma 6.9 shows that all separating twists of genus d lie in \mathcal{T}_ϕ . In Substep 2, Lemma 6.11 shows that all separating twists of genus $h \equiv d + 2 \pmod{2d}$ lie in \mathcal{T}_ϕ , and Lemma 6.12 establishes the same result for separating twists of genus $h \equiv d + 4 \pmod{2d}$. Lemmas 6.10 and 6.13 then follow from these preliminary results and an application of the D_n relation (Proposition 2.6).

Lemma 6.8 *Let $S \subset \Sigma$ be a subsurface with single boundary component. Assume the genus $g(S) \geq 2$. Then there is a maximal chain of admissible curves on S if and only if one of the following conditions hold:*

- $g(S) \equiv 1$ or $2 \pmod{4}$ and $\epsilon(S) = 1$,
- $g(S) \equiv 3$ or $0 \pmod{4}$ and $\epsilon(S) = 0$.

Proof Suppose S supports a maximal chain $a_1, \dots, a_{2g(S)}$ of admissible curves. Since the chain determines a basis for $H_1(S; \mathbb{Z})$, the conditions $\phi(a_i) = 0$ completely determine ϕ . One can easily compute $\epsilon(S)$ from this and see that the above conditions are necessary. Sufficiency follows from Corollary 4.10.4. □

Substep 1: d odd The objective of Substep 1 is Lemma 6.10 below. The first step is to see that all separating twists T_c of genus d are elements of \mathcal{T}_ϕ , regardless of whether c is easy.

Lemma 6.9 *Let $S \subset \Sigma_g$ be a subsurface of genus d with a single boundary component c . If ϕ is a $\mathbb{Z}/2d\mathbb{Z}$ -valued spin structure with d odd, then $T_c \in \mathcal{T}_\phi$.*

Proof If c is easy then there is nothing to show. Assume therefore that c is hard. If c is oriented so that S lies to the right, then $\phi(c) = -(1 - 2d) \equiv -1 \pmod{2d}$. The assumption that $r = 2d < g - 1$ implies that $\Sigma_g \setminus S$ has genus at least 2. We claim that there exists a 3-chain of admissible curves x, y, z on $\Sigma_g \setminus S$ such that $c \cup x \cup z$ forms a pair of pants. To see this, we invoke Corollary 4.3 to let $x \subset \Sigma_g \setminus S$ be an admissible curve. Let $z \subset \Sigma_g \setminus S$ be any curve such that $c \cup x \cup z$ bounds a pair of pants; admissibility of z follows by the homological coherence property, as c is oriented with $\Sigma_g \setminus S$ to the left. To construct y , let $y' \subset \Sigma_g \setminus S$ be any curve such that x, y', z forms a chain. By Corollary 4.4, y' can be replaced with an admissible curve y with the same intersection properties.

Let S' denote the connected surface of genus $d + 1$ containing S and $x \cup y \cup z$. If \mathcal{B} is a basis for $H_1(S; \mathbb{Z})$, then $\mathcal{B} \cup \{x, y\}$ forms a basis for $H_1(S'; \mathbb{Z})$. Applying the formula (7) for the Arf invariant, it follows that $\epsilon(S') = \epsilon(S) + 1$.

Since c is hard and $d = g(S)$ is odd, Lemma 6.8 implies that $\epsilon(S) = 0$ if $g(S) \equiv 1 \pmod{4}$ and that $\epsilon(S) = 1$ otherwise. Recalling that $\epsilon(S') =$

$\epsilon(S) + 1$, in the first case, $g(S') \equiv 2 \pmod{4}$ and $\epsilon(S') = 1$, and in the second case, $g(S') \equiv 0 \pmod{4}$ and $\epsilon(S') = 0$. Lemma 6.8 then implies that $c' := \partial S'$ must be easy, and so $T_{c'} \in \mathcal{T}_\phi$. Applying the chain relation (Proposition 2.4) to x, y, z shows that $T_c T_{c'} \in \mathcal{T}_\phi$; this implies that also $T_c \in \mathcal{T}_\phi$. \square

Lemma 6.10 *Let ϕ be a $\mathbb{Z}/2d\mathbb{Z}$ -valued spin structure on Σ_g with d odd. Assume that g satisfies the hypotheses of Proposition 6.2. Then \mathcal{T}_ϕ contains the Johnson kernel \mathcal{K}_g .*

Proof By Theorem 2.9, it suffices to show that $T_c \in \mathcal{T}_\phi$ for all separating curves c of arbitrary genus. To do this, we combine Lemma 6.9 with the D_n relation (Proposition 2.6). Suppose c is a separating curve on Σ_g . Since $g = kd + 1$ with $k \geq 2$, at least one side of c must be a subsurface S of genus $g(S) \geq d + 1$. Set $n := 2g(S) - 2d + 1$. By Corollary 4.11, there is a configuration \mathcal{D}_n of admissible curves as in the D_n relation for which $\Delta_2 = c$. The other boundary component Δ_0 bounds a subsurface of genus d . Applying the D_n relation, we have $T_{\Delta_0}^{n-1} T_c \in \mathcal{T}_\phi$. But since Δ_0 bounds a surface of genus d , also $T_{\Delta_0} \in \mathcal{T}_\phi$ by Lemma 6.9. Thus $\mathcal{K}_g \leq \mathcal{T}_\phi$ in this case. \square

Substep 2: d even The objective is to establish Lemma 6.13. The argument here follows a similar outline to that of Substep 1 but now requires the two preliminary Lemmas 6.11 and 6.12.

Lemma 6.11 *Let $S \subset \Sigma_g$ be a subsurface of genus $g(S) \geq 5$ with a single boundary component c , such that $g(S) \equiv d + 2 \pmod{2d}$. If ϕ is a $\mathbb{Z}/2d\mathbb{Z}$ -valued spin structure with d even, then $T_c \in \mathcal{T}_\phi$.*

Proof Orient c so that S lies to the left. Then

$$\phi(c) = 1 - 2g(S) \equiv 1 - 2(d + 2) \equiv -3 \pmod{2d}.$$

By Corollary 4.10.5, there exists a chain a_1, \dots, a_6 of admissible curves on S . Let a_7 be any curve on S such that $i(a_7, a_k) = 1$ for $k = 6$ and is zero for $k \leq 5$, and such that $c \cup a_1 \cup a_3 \cup a_5 \cup a_7$ bounds a subsurface of S homeomorphic to $\Sigma_{0,5}$. By homological coherence, a_7 is admissible.

Let S' denote the subsurface of S homeomorphic to $\Sigma_{g(S)-3,1}$ determined by the complement of the chain a_1, \dots, a_7 . Applying the formula (7) for the Arf invariant, one finds that $\epsilon(S') = \epsilon(S)$. On the other hand, $g(S') \equiv g(S) + 1 \pmod{4}$. By hypothesis, $g(S)$ is even, and so referring to Lemma 6.8, if c is hard, then $c' := \partial S'$ must be easy. The arguments given at the conclusion of Lemma 6.9 now apply to give the result. \square

Lemma 6.12 *Let $S \subset \Sigma$ be a subsurface of genus $g(S) \geq 9$ with a single boundary component c , such that $g(S) \equiv d + 4 \pmod{2d}$. If ϕ is a $\mathbb{Z}/2d\mathbb{Z}$ -valued spin structure with d even, then $T_c \in \mathcal{T}_\phi$.*

Proof This is proved along similar lines to Lemma 6.11. Arguing as in the first paragraph of the proof of Lemma 6.11, there exists a chain a_1, \dots, a_{15} of admissible curves on S such that $c \cup a_1 \cup a_3 \cup \dots \cup a_{15}$ bounds a subsurface of S homeomorphic to $\Sigma_{0,9}$. Let S' denote the subsurface of S homeomorphic to $\Sigma_{g(S)-7,1}$ determined by the complement of the chain a_1, \dots, a_{15} . The rest of the argument proceeds as in Lemma 6.11: one shows that if c is hard, necessarily $c' := \partial S'$ must be easy, and the result follows as before by the chain relation (Proposition 2.4). \square

Lemma 6.13 *Let ϕ be a $\mathbb{Z}/2d\mathbb{Z}$ -valued spin structure on Σ_g with d even. Assume that g satisfies the hypotheses of Proposition 6.2. Then \mathcal{T}_ϕ contains the Johnson kernel \mathcal{K}_g .*

Proof According to Johnson’s Theorem 2.9, in order to show that $\mathcal{K}_g \leq \mathcal{T}_\phi$, it suffices to exhibit all separating twists of genus 1 and 2 as elements of \mathcal{T}_ϕ . To do this, we again appeal to the D_n relation (Proposition 2.6). Suppose c is a separating curve on Σ_g with $g(c) \leq 2$. By hypothesis, $g \geq kd + 1$ with d even and $k \geq 2$. Since the genus of one side of c is at most 2, the genus h of the other side of c is at least $kd - 1 \geq 2d - 1$. If $d \geq 6$, then $2d - 1 \geq 11$. If $d = 4$, then by assumption $k \geq 5$, and so $h \geq 19$. If $d = 2$ then we assume $k \geq 6$, so that $h \geq 11$.

In all three of these cases, Corollary 4.11 implies that there exists an $n \geq 4$ and a configuration $a, a', c_1, \dots, c_{2n-1}$ of admissible curves in the configuration of the D_{2n+1} relation, with $\Delta_2 = c$ and C_4 bounding a subsurface of genus $g(S) \equiv d + 4 \pmod{2d}$ disjoint from S , such that the hypotheses of Lemmas 6.11 and 6.12 hold.

By the D_k relation (for $k = 2n + 1, 5, 9$ respectively), $T_{\Delta_0}^{2n-1} T_c$ and $T_{\Delta_0}^3 T_{C_2}$ and $T_{\Delta_0}^7 T_{C_4}$ are all elements of \mathcal{T}_ϕ . By Lemma 6.11, $T_{C_2} \in \mathcal{T}_\phi$ as well, hence $T_{\Delta_0}^3 \in \mathcal{T}_\phi$. Likewise, Lemma 6.12 shows that $T_{C_4} \in \mathcal{T}_\phi$, hence $T_{\Delta_0}^7 \in \mathcal{T}_\phi$. Combining these last two results shows that $T_{\Delta_0} \in \mathcal{T}_\phi$, and ultimately that $T_c \in \mathcal{T}_\phi$ as required. \square

This concludes the proof of Proposition 6.2. \square

7 Connectivity of some complexes

This section is devoted to establishing the connectivity of the simplicial complexes $\mathcal{C}_{sep,2}(\Sigma_g)$ and $\mathcal{C}_\phi^1(\Sigma_g)$ to be defined below. The first of these will be an important ingredient in the proof of Proposition 8.2, and the second will feature in the proof of Theorem A. The mechanism by which these will be seen to be connected is the so-called *Putman trick*. The version given below is slightly less general than the full theorem as stated in [16], but will suffice for our purposes.




Fig. 7 The configuration of curves needed for Lemma 7.3

Theorem 7.1 (The Putman trick) *Let X be a simplicial graph, and let G act on X by simplicial automorphisms. Suppose that the action of G on the set of vertices $X^{(0)}$ is transitive. Fix some base vertex $v \in X^{(0)}$. Let $\Sigma = \Sigma^{-1}$ be a symmetric set of generators for G , and suppose that for each $s \in \Sigma$, there is a path in X connecting v to $s \cdot v$. Then X is connected.*

Definition 7.2 $C_{sep,2}(\Sigma_g)$ is the simplicial graph where vertices correspond to (isotopy classes of) separating curves c bounding a subsurface homeomorphic to $\Sigma_{2,1}$, and where c and d are adjacent in $C_{sep,2}(\Sigma_g)$ whenever c and d are disjoint in Σ_g .

Lemma 7.3 $C_{sep,2}(\Sigma_g)$ is connected for $g \geq 5$.

Proof This is a straightforward consequence of Theorem 7.1. With reference to Fig. 7 and the standard generating set of Fig. 1, observe that only the generator $T_{c_2}^\pm$ does not fix the base vertex c . In this case, the genus 2 subsurface determined by d is disjoint from both c and $T_{c_2}^\pm(c)$, and so there is a path $c, d, T_{c_2}^\pm(c)$ in $C_{sep,2}(\Sigma_g)$. □

Definition 7.4 Let ϕ be a $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure on a surface Σ_g . The graph $C_\phi(\Sigma_g)$ has vertices consisting of the admissible curves for ϕ , where a and b are adjacent whenever $i(a, b) = 0$. The graph $C_\phi^1(\Sigma_g)$ has the same vertex set as $C_\phi(\Sigma_g)$, but vertices a, b are adjacent whenever $i(a, b) = 1$.

Lemma 7.5 $C_\phi^1(\Sigma_g)$ is connected for $g \geq 5$.

Proof The first step is to establish the connectivity of $C_\phi(\Sigma_g)$. Let a, b be vertices. Choose subsurfaces S_a, S_b containing a, b respectively, each homeomorphic to $\Sigma_{2,1}$. By Lemma 7.3, there is a path S_{a_0}, \dots, S_{a_n} in $C_{sep,2}(\Sigma_g)$ with $a \subset S_{a_0}$ and $b \subset S_{a_n}$, with each S_{a_i} disjoint from $S_{a_{i+1}}$. By Corollary 4.3, on each S_{a_i} there exists some admissible curve a_i . By construction, $a = a_0, a_1, \dots, a_n = b$ is a path in $C_\phi(\Sigma_g)$ connecting a to b .

The connectivity of $C_\phi^1(\Sigma_g)$ now follows readily. Given a path $a = a_0, \dots, a_n = b$ in $C_\phi(\Sigma_g)$, Corollary 4.4 implies that for each i , there exists some admissible curve c_i such that $i(a_i, c_i) = i(a_{i+1}, c_i) = 1$. The path $a_0, c_0, a_1, c_1, \dots, c_{n-1}, a_n$ connects a to b in $C_\phi^1(\Sigma_g)$. □

8 Subsurface push subgroups and \mathcal{T}_ϕ

As discussed in the introduction, the main technical result on the groups $\text{Mod}(\Sigma_g)[\phi]$ and \mathcal{T}_ϕ that we require is a criterion for a collection of Dehn twists to generate \mathcal{T}_ϕ , given below as Theorem 9.5. This is the first of two sections dedicated to proving Theorem 9.5. Here, we formulate and prove the intermediate result Proposition 8.2, which gives a generating set for \mathcal{T}_ϕ not consisting entirely of Dehn twists. The results here concern a class of subgroups known as *spin subsurface push subgroups*; these are introduced in Sects. 8.1 and 8.2.

8.1 Subsurface push subgroups

Recall the classical *inclusion map*, as discussed in [7, Theorem 3.18]. Let $S' \subset S$ be a subsurface either of genus $g(S') \geq 2$ with $n \geq 1$ boundary components, or else of genus $g(S') = 1$ with $n \geq 2$ boundary components. Assume that no component of $\partial S'$ bounds a closed disk in S . Let a_1, \dots, a_k denote the boundary components of S' that bound punctured disks in S , let $b_1, b'_1, \dots, b_\ell, b'_\ell$ denote the pairs of boundary components of S' that cobound an annulus in S , and c_1, \dots, c_m denote the remaining boundary components. Let $i_*: \text{Mod}(S') \rightarrow \text{Mod}(S)$ denote the map on mapping class groups arising from the inclusion $i: S' \hookrightarrow S$. Then

$$\ker(i_*) = \left\langle T_{a_1}, \dots, T_{a_k}, T_{b_1} T_{b'_1}^{-1}, \dots, T_{b_\ell} T_{b'_\ell}^{-1} \right\rangle.$$

Let Δ be a boundary component of S' , and suppose that Δ does not bound a punctured disk in S . Let $\overline{S'}$ denote the surface obtained from S' by capping off Δ with a closed disk. According to (2), there is a subgroup of $\text{Mod}(S')$ isomorphic to $\pi_1(UT\overline{S'})$. The *subsurface push subgroup* for (S', Δ) is defined to be the image of $\pi_1(UT\overline{S'})$ under the inclusion $i_*: \text{Mod}(S') \rightarrow \text{Mod}(S)$. This will be written $\Pi(S', \Delta)$, or simply $\Pi(S')$ if the boundary component does not need to be emphasized.

We remark here that i_* restricts to an *injection* $\pi_1(UT\overline{S'}) \hookrightarrow \text{Mod}(S)$, even when there exists some other boundary component Δ' of S' such that $\Delta \cup \Delta'$ cobounds an annulus on S . To see this, observe that $\pi_1(UT\overline{S'}) \leq \text{Mod}(S')$ is characterized by the property that $f \in \pi_1(UT\overline{S'})$ if and only if f becomes isotopic to the identity when extended to $\overline{S'}$. It is easy to see that no element of $\ker(i_*)$ has this property.

8.2 Spin subsurface push subgroups

Let $S' \subset S$ be a subsurface with some boundary component Δ satisfying $\phi(\Delta) = -1$. The following lemma shows that $\text{Mod}(S)[\phi]$ contains a finite-index subgroup of $\Pi(S', \Delta)$. This subgroup, written $\tilde{\Pi}(S', \Delta)$, is called a *spin subsurface push subgroup*. Before proceeding with the rest of the section, the reader may wish to review the notion of a *fundamental multitwist* defined in Sect. 3.3.

Lemma 8.1 *Let $S' \subset S$ be a subsurface with some boundary component Δ satisfying $\phi(\Delta) = -1$. Then there is a finite-index subgroup $\tilde{\Pi}(S', \Delta) \leq \text{Mod}(S)[\phi] \cap \Pi(S', \Delta)$ characterized by the diagram given below, whose rows are short exact sequences:*

$$\begin{array}{ccccccc}
 1 & \longrightarrow & \langle T_{\Delta}^r \rangle & \longrightarrow & \tilde{\Pi}(S', \Delta) & \longrightarrow & \pi_1(\overline{S'}) \longrightarrow 1 \\
 & & \downarrow & & \downarrow & & \parallel \\
 1 & \longrightarrow & \langle T_{\Delta} \rangle & \longrightarrow & \Pi(S', \Delta) & \longrightarrow & \pi_1(\overline{S'}) \longrightarrow 1.
 \end{array} \tag{10}$$

The subgroup $\tilde{\Pi}(S', \Delta)$ contains all fundamental multitwists for pairs of pants $P \subset S'$ of the form $P = a \cup b \cup \Delta$.

Proof Following the discussion of Sect. 2.2, there exists a “geometric” generating set for $\pi_1(UT\overline{S'}, \Delta)$ of the following form:

$$\pi_1(UT\overline{S'}) = \langle \tilde{\alpha}_1, \dots, \tilde{\alpha}_k, \zeta \rangle. \tag{11}$$

Here α_i is some simple closed curve on $\overline{S'}$ based at Δ , and $\tilde{\alpha}_i$ denotes the Johnson lift to $\pi_1(UT\overline{S'})$. As before, ζ denotes the loop around the fiber. As an element of $\text{Mod}(S')$, each $\tilde{\alpha}_i$ is of the form $T_{\alpha_{i,L}} T_{\alpha_{i,R}}$, where $\alpha_{i,L}$ denotes the curve on S' lying to the left of α_i and $\alpha_{i,R}$ lies to the right. It follows that $P_i = \alpha_{i,L} \cup \alpha_{i,R} \cup \Delta$ forms a pair of pants on S' . Following Lemma 3.18, the fundamental multitwist

$$T_{P_i} = T_{\alpha_{i,L}} T_{\alpha_{i,R}}^{-1} T_{\Delta}^{\phi(\alpha_{i,R})}$$

lies in $\text{Mod}(S)[\phi] \cap \Pi(S', \Delta)$. Embedding $\pi_1(UT\overline{S'})$ into $\text{Mod}(S')$, the generating set of (11) can be replaced by the following generating set for $\Pi(S', \Delta)$:

$$\Pi(S', \Delta) = \langle T_{P_1}, \dots, T_{P_k}, T_{\Delta} \rangle.$$

Define

$$\tilde{\Pi}(S', \Delta) = \langle T_{P_1}, \dots, T_{P_k}, T_{\Delta}^r \rangle.$$

By construction, $\tilde{\Pi}(S', \Delta) \leq \text{Mod}(S)[\phi]$. Under the projection $\Pi(S', \Delta) \rightarrow \pi_1(\overline{S'})$, the set $\{T_{P_i}\}$ maps onto a generating set for $\pi_1(\overline{S'})$. It follows that $\tilde{\Pi}(S', \Delta)$ surjects onto $\pi_1(\overline{S'})$. As $T_\Delta^m \in \text{Mod}(S)[\phi]$ if and only if $r \mid m$, it follows that $\tilde{\Pi}(S', \Delta)$ is indeed characterized by the diagram (10) as claimed.

For the second claim, let $P = a \cup b \cup \Delta$ be a pair of pants on S' . The curves a, b are isotopic on $\overline{S'}$ and cobound an annulus containing the basepoint. It follows that $T_a T_b^{-1} \in \pi_1(\overline{S'})$. Via (10), there is some lift $T_a T_b^{-1} T_\Delta^k \in \tilde{\Pi}(S', \Delta)$, and as $T_\Delta^r \in \tilde{\Pi}(S', \Delta)$ as well, it follows that all fundamental multitwists for P are elements of $\tilde{\Pi}(S', \Delta)$ as claimed. \square

For the purposes of this paper, we will most often be concerned with subsurface push subgroups for a special class of subsurfaces. Let $b \subset \Sigma_g$ be a nonseparating closed curve satisfying $\phi(b) = -1$. The boundary component Δ of $\Sigma_g \setminus \{b\}$ corresponding to the left side of b satisfies $\phi(\Delta) = -1$, and to ease notation, we write $\tilde{\Pi}(\Sigma_g \setminus \{b\})$ to refer to this spin subsurface push subgroup.

8.3 Generating admissible twists

We have arrived at the key result of the section.

Proposition 8.2 *Let ϕ be a $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure on a closed surface Σ_g for $g \geq 5$ and any integer r . Let (a_0, a_1, b) be an ordered 3-chain of curves with $\phi(a_0) = \phi(a_1) = 0$ and $\phi(b) = -1$. Let $H \leq \text{Mod}(\Sigma_g)$ be a subgroup containing T_{a_0}, T_{a_1} and the spin subsurface push group $\tilde{\Pi}(\Sigma_g \setminus \{b\})$. Then H contains T_ϕ .*

The proof will require the preliminary Lemma 8.3, for which we introduce some terminology. For a subgroup $H \leq \text{Mod}(\Sigma_g)$, we say that a simple closed curve a is an H -curve if $T_a \in H$. We also say that curves a, b are H -equivalent if there exists some $f \in H$ with $f(a) = b$. If a and $b = f(a)$ are H -equivalent and $\tilde{\Pi}(\Sigma_g \setminus \{a\}) \leq H$, then also $\tilde{\Pi}(\Sigma_g \setminus \{b\}) = f \tilde{\Pi}(\Sigma_g \setminus \{a\}) f^{-1}$ is a subgroup of H .

The following lemma establishes some sufficient conditions for H -equivalence of curves.

Lemma 8.3 *Let Σ_g be a surface of genus $g \geq 5$. Let a_0, a_1, b be an ordered 3-chain of curves with $\phi(a_0) = \phi(a_1) = 0$ and $\phi(b) = -1$. Let $H \leq \text{Mod}(\Sigma_g)$ be a subgroup containing T_{a_0}, T_{a_1} and $\tilde{\Pi}(\Sigma_g \setminus \{b\})$.*

- (1) *Let b' be an oriented curve satisfying $\phi(b') = -1$ such that $i(b, b') = 0$ and $i(a_1, b') = 1$. Then b and b' are H -equivalent. It follows that $\tilde{\Pi}(\Sigma_g \setminus \{b'\}) \leq H$.*
- (2) *Let a' be any nonseparating curve satisfying $\phi(a') = 0$ such that a' is disjoint from the configuration $a_0 \cup a_1 \cup b$. Then a' is an H -curve.*

- (3) Let b' be any nonseparating curve satisfying $\phi(b') = -1$ such that b' is disjoint from the configuration $a_0 \cup a_1 \cup b$. Then b and b' are H -equivalent, and hence $\tilde{\Pi}(\Sigma_g \setminus \{b'\}) \leq H$.

Proof(1) If $b = b'$ there is nothing to prove. Otherwise, given a_1, b, b' , we define a curve b'' as follows. Let ϵ be the portion of a_1 connecting the left side of b to one of the sides of b' ; then b'' is defined as the curve-arc sum $b'' := b +_\epsilon b'$. By construction $b \cup b' \cup b''$ bounds a pair of pants P lying to the left of b , and $i(a_1, b'') = 0$. By Lemma 3.13, there exists an orientation of b'' such that $\phi(b'') = -1$. This can be determined as follows: b'' is oriented with P lying to the right if and only if P lies to the left of b' . If P lies to the left of b' , then the element $T_b T_{b'} T_{b''}^{-1}$ is a fundamental multitwist and hence an element of $\tilde{\Pi}(\Sigma_g \setminus \{b\}) \leq H$. Otherwise, $T_b T_{b'}^{-1} T_{b''}$ is a fundamental multitwist. In the first case, the braid relation implies that

$$T_{a_1}(T_b T_{b'} T_{b''}^{-1})T_{a_1}(b) = b',$$

while in the second case,

$$T_{a_1}^{-1}(T_b T_{b'}^{-1} T_{b''})T_{a_1}(b) = b'.$$

In either case, the indicated element lies in H , showing the H -equivalence between b, b' .

- (2) Let ϵ be an arc connecting a_0 to a' that is disjoint from $a_1 \cup b$, and define $b' := a_0 +_\epsilon a'$. It is possible that $b' = b$, but this will not pose any difficulty. Then $a_0 \cup a' \cup b'$ forms a pair of pants and b' satisfies the intersection conditions $i(b, b') = 0$ and $i(a_1, b') = 1$. By the homological coherence property, $\phi(b') = -1$. By the second assertion of (1), $\tilde{\Pi}(\Sigma_g \setminus \{b'\}) \leq H$. As $a_0 \cup a' \cup b'$ forms a pair of pants, it follows that $T_{a_0} T_{a'}^{-1}$ is a fundamental multitwist, and so $T_{a_0} T_{a'}^{-1} \in \tilde{\Pi}(\Sigma_g \setminus \{b'\}) \leq H$. As $T_{a_0} \in H$ by hypothesis, this shows that $T_{a'} \in H$ as desired.
- (3) Given b' , Corollary 4.4 implies that there exists an admissible curve a' that is disjoint from $a_0 \cup a_1 \cup b$ and for which $i(a', b') = 1$. Corollary 4.4 also establishes the existence of a curve b'' , satisfying $\phi(b'') = -1$, with the following intersection properties:

$$i(b, b'') = i(b', b'') = i(a_0, b'') = 0; \quad i(a', b'') = i(a_1, b'') = 1.$$

By (1), b and b'' are H -equivalent. By (2), a' is an H -curve, so that by (1) again, b'' and b' are H -equivalent, showing the result. □

Proof of Proposition 8.2 Let a be any admissible curve. There is some genus 2 subsurface $S' \cong \Sigma_{2,1}$ containing a , and there is also some genus 2 subsurface

$S \cong \Sigma_{2,1}$ that contains the curves a_0, a_1, b . By Lemma 7.3, there is a path $S_0 = S - S_1 - \dots - S_n = S'$ of subsurfaces homeomorphic to $\Sigma_{2,1}$ with boundary components ∂S_i and ∂S_{i+1} disjoint for $i = 1, \dots, n - 1$, hence $S_i \cap S_{i+1} = \emptyset$ for $i = 1, \dots, n - 1$.

For $i = 1, \dots, n$, let a_{2i} be an admissible curve contained in S_i ; we take $a_{2n} = a$. We claim that there exist curves a_{2i+1} and b_i on S_i such that a_{2i}, a_{2i+1}, b_i forms a chain, and $\phi(a_{2i+1}) = 0, \phi(b_i) = -1$. To see this, let $T \subset S_i$ be a subsurface of genus 1 that does not contain a_{2i} . By Corollary 4.3, there is an admissible curve a' contained in T . Let ϵ be an arc connecting a_{2i} and a' ; then $b_i := a_{2i} +_\epsilon a'$ satisfies $\phi(b_i) = -1$ for a suitable choice of orientation. Let c be any curve on S_i such that $i(c, a_{2i}) = i(c, b_i) = 1$. Then $a_{2i+1} := T_{b_i}^{\phi(c)}(c)$ is admissible, and a_{2i}, a_{2i+1}, b_i forms a chain as required.

We assume for the sake of induction that a_{2i}, a_{2i+1} are H -curves and that $\tilde{\Pi}(\Sigma_g \setminus \{b_i\}) \leq H$. Then by Lemma 8.3.2, also a_{2i+2}, a_{2i+3} are H -curves, and $\tilde{\Pi}(\Sigma_g \setminus \{b_{i+1}\}) \leq H$. The base case $i = 0$ holds by hypothesis, taking $b_0 = b$. The claim now follows by induction. □

9 Networks

In this section we deduce Theorem 9.5 from Proposition 8.2. The key notion is that of a *network* of curves. In Sect. 9.1, we establish the basic theory of networks, and in Sect. 9.2, we state and prove Theorem 9.5. Departing from our conventions elsewhere in the paper, in this section we work with individual curves and *not* merely their isotopy classes.

9.1 Networks and their basic theory

Definition 9.1 Let $S = \Sigma_{g,b}^n$ be a surface, viewed as a compact surface with marked points. A *network* on S is any collection $\mathcal{N} = \{a_1, \dots, a_n\}$ of simple closed curves on S , disjoint from any marked points, such that $\#(a_i \cap a_j) \leq 1$ for all pairs of curves $a_i, a_j \in \mathcal{N}$, and such that there are no triple intersections. A network \mathcal{N} has an associated *intersection graph* $\Gamma_{\mathcal{N}}$, whose vertices correspond to curves $x \in \mathcal{N}$, with vertices x, y adjacent if and only if $\#(x \cap y) = 1$. A network is said to be *connected* if $\Gamma_{\mathcal{N}}$ is connected, and *arboreal* if $\Gamma_{\mathcal{N}}$ is a tree. A network is *filling* if

$$S \setminus \bigcup_{a \in \mathcal{N}} a$$

is a disjoint union of disks and boundary-parallel annuli; each component is allowed to contain at most one marked point of S .

The data of a network encodes both an abstract finite set of curves as well as a topological subspace of the surface S . To avoid confusing these, let the symbol \mathcal{N} denote this finite set, and let $\widehat{\mathcal{N}}$ denote the space. When \mathcal{N} is arboreal, there is a simple generating set for $\pi_1(\widehat{\mathcal{N}})$. To describe it, endow $\widehat{\mathcal{N}}$ with the structure of a CW complex, and let \mathcal{T} be a spanning tree for this CW complex.

Lemma 9.2 *Let \mathcal{N} be an arboreal network. Then there is a 1–1 correspondence between the set of edges $\widehat{\mathcal{N}} \setminus \mathcal{T}$, and the set \mathcal{N} .*

Proof Each edge of $\widehat{\mathcal{N}}$ is contained in a unique element of \mathcal{N} . For a given $a \in \mathcal{N}$, let $a_1, \dots, a_{n(a)}$ denote these edges, ordered so that adjacent edges are numbered consecutively. For each $a \in \mathcal{N}$, the sequence $a_1, \dots, a_{n(a)}$ forms a cycle in $\widehat{\mathcal{N}}$. Thus for each $a \in \mathcal{N}$, there is at least one edge a_1 (without loss of generality) that is not contained in \mathcal{T} .

It remains to show that for each $a \in \mathcal{N}$, there is *exactly* one edge not contained in \mathcal{T} . Equivalently, we must show that the intersection $a \cap \mathcal{T}$ is connected as a topological space. The assumption that \mathcal{N} is arboreal implies that $\Gamma_{\mathcal{N}}$ has the following property: let $\Gamma_{\mathcal{N}}(a)$ be the graph obtained from $\Gamma_{\mathcal{N}}$ by removing all edges incident to a . Then each vertex b adjacent to a in $\Gamma_{\mathcal{N}}$ determines a *distinct* component of $\Gamma_{\mathcal{N}}(a)$.

Let $v, w \in a$ be vertices of $\widehat{\mathcal{N}}$, and let e_1, \dots, e_n be the unique geodesic path in \mathcal{T} connecting v to w . It suffices to show that this path is contained in a . If this is not the case, let k_1 (resp. k_2) be the minimal (resp. maximal) integer such that e_{k_1} (resp. e_{k_2}) is not contained in a . Then exactly one vertex v_1 of e_{k_1} (resp. v_2 of e_{k_2}) lies on a , and the other lies on some adjacent curve b_1 (resp. b_2). As $i(a, b_1) = i(a, b_2) = 1$ and the path e_1, \dots, e_n visits each vertex in $\widehat{\mathcal{N}}$ at most once, it follows that b_1 and b_2 are distinct elements of \mathcal{N} . As explained in the above paragraph, the arboreality assumption implies that every path in $\widehat{\mathcal{N}}$ connecting a point in b_1 to a point in b_2 must pass through a . Any such path must pass through v_1 and v_2 : this shows that if the path e_1, \dots, e_n enters b_1 , it must pass through v_1 at least twice, contrary to assumption. \square

Via Lemma 9.2, each $a \in \mathcal{N}$ determines a *unique* based loop $P(a)$ by following the unique path in \mathcal{T} from the basepoint to a . Lemma 9.3 below now follows by a standard application of the Seifert–Van Kampen theorem.

Lemma 9.3 *Let \mathcal{N} be an arboreal network. Then $\pi_1(\widehat{\mathcal{N}})$ is generated by the set of loops $\{P(a) \mid a \in \mathcal{N}\}$. If \mathcal{N} is moreover filling, then the map $\pi_1(\widehat{\mathcal{N}}) \rightarrow \pi_1(S)$ is a surjection, and so $\pi_1(S)$ is also generated by this collection of loops.*

$\pi_1(S)$ is a normal subgroup of $\text{Mod}(S)$: if $\alpha \in \pi_1(S)$ is a mapping class corresponding to a based loop and $f \in \text{Mod}(S)$ is arbitrary, then conjugation by f takes α to the mapping class corresponding to the based loop $f(\alpha)$. In

the context of the “network presentation” of $\pi_1(S)$ arising from the surjection $\pi_1(\widehat{\mathcal{N}}) \rightarrow \pi_1(S)$, this means that $\pi_1(S)$ has a very simple *normal* generating set as a subgroup of $\text{Mod}(S)$, as the following makes precise.

Lemma 9.4 *Let $\widehat{\mathcal{N}} \subset S$ be an arboreal filling network. Let $H \leq \text{Mod}(S)$ be a subgroup containing T_a for each $a \in \mathcal{N}$. If H also contains $P(a_1) \in \pi_1(S)$ for some $a_1 \in \mathcal{N}$, then H contains the entire point-pushing subgroup $\pi_1(S)$.*

Proof As recorded in Lemma 9.3, $\pi_1(\widehat{\mathcal{N}})$, and hence also $\pi_1(S)$, is generated by the collection of elements $P(a)$ for $a \in \mathcal{N}$. We will proceed by induction. Define connected subnetworks

$$\mathcal{N}_0 \subset \mathcal{N}_1 \subset \dots \subset \mathcal{N}_n = \mathcal{N}$$

as follows: \mathcal{N}_k consists of all those curves a at a distance of at most k from the base vertex $a_1 \in \Gamma_{\mathcal{N}}$ (viewing $\Gamma_{\mathcal{N}}$ as a metric space for which each edge has length 1). We suppose that $\pi_1(\widehat{\mathcal{N}}_k) \leq H$; the base case $k = 0$ holds by hypothesis.

Let $a \in \mathcal{N}_{k+1} \setminus \mathcal{N}_k$ be arbitrary. Let $a' \in \mathcal{N}_k$ be adjacent to a . By the braid relation,

$$T_{a'}T_a(a') = a,$$

and hence $P(a) = (T_{a'}T_a)P(a')(T_{a'}T_a)^{-1} \in H$. This completes the inductive step. □

9.2 Network generating sets for \mathcal{T}_ϕ

Having established some of the basic theory of networks, we can now formulate and prove the key technical result of the paper. For hypotheses (2) and (3), the reader may wish to consult Fig. 2 and the surrounding discussion of the D_n relation (Proposition 2.6) and the configuration \mathcal{D}_n . For an example of a network satisfying the hypotheses of Theorem 9.5, see Fig. 8.

Theorem 9.5 *Let ϕ be a $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure on a closed surface Σ_g , with $1 \leq r < g - 1$. Let $\mathcal{N} = \{a_n\}$ be a connected filling network of curves on Σ_g with the following properties:*

- (1) *Every element a_n is admissible,*
- (2) *There is a collection a_1, \dots, a_{2r+4} of elements of \mathcal{N} such that a_1, \dots, a_{2r+3} are arranged in the configuration of the curves \mathcal{D}_{2r+3} of the D_{2r+3} relation, and a_{2r+4} corresponds to the boundary component Δ_1 associated to the subconfiguration \mathcal{D}_{2r+2} .*
- (3) *Let $b \subset \Sigma_g$ correspond to the curve Δ_0 of the D_{2r+3} relation, as appearing in Fig. 2. Then \mathcal{N} must contain some curve d with $i(d, b) = 1$.*

(4) Let $\mathcal{N}' \subset \mathcal{N}$ be the subnetwork consisting of all curves in \mathcal{N} disjoint from b . Then \mathcal{N}' must be an arboreal filling network for $\Sigma_g \setminus \{b\}$.

If $g \geq 5$, then $\langle T_{a_i}, a_i \in \mathcal{N} \rangle$ contains the admissible subgroup \mathcal{T}_ϕ .

Moreover, if r is odd, then

$$\langle T_{a_i}, a_i \in \mathcal{N} \rangle = \text{Mod}(\Sigma_g)[\phi].$$

If $r = 2d$ is even and $g \geq g(r)$ for the function $g(r)$ of Definition 6.1, then $\langle T_{a_i}, a_i \in \mathcal{N} \rangle$ is a subgroup of finite index in $\text{Mod}(\Sigma_g)[\phi]$.

Proof Define

$$H = \langle T_{a_i}, a_i \in \mathcal{N} \rangle.$$

By hypothesis (1), $H \leq \mathcal{T}_\phi$. We establish the opposite containment $\mathcal{T}_\phi \leq H$. The remaining assertions in the statement of Theorem 9.5 follow by an appeal to Propositions 5.1 or 6.2 as appropriate. The containment $\mathcal{T}_\phi \leq H$ will follow from Proposition 8.2. To see that the hypotheses of Proposition 8.2 are satisfied by H , it is necessary to establish a containment $\tilde{\Pi}(\Sigma_g \setminus \{b\}) \leq H$, and to find suitable curves corresponding to a_0, a_1 in the statement of Proposition 8.2.

Consider the curves $\{a_1, \dots, a_{2r+4}\} \subset \mathcal{N}$ corresponding to $\mathcal{D}_{2r+3} \cup \{\Delta_1\}$ as in Corollary 2.7, as posited by hypothesis (2). Without loss of generality, assume that $a_1, a_2, a_3 \in \mathcal{N}$ correspond to the curves a, c_1, a' of \mathcal{D}_{2r+3} , so that $b \subset \Sigma_g$ is one of the boundary components of the chain a_1, a_2, a_3 . Let d be the curve with $i(d, b) = 1$ posited by hypothesis (3), and let P be the pair of pants bounded by a_1, a_3, b . The intersection $d \cap P$ must be a single arc, since $d \in \mathcal{N}$ and so $\#(d \cap a_1) \leq 1$ and $\#(d \cap a_3) \leq 1$. Without loss of generality, assume $\#(d \cap a_1) = 1$ and $\#(d \cap a_3) = 0$. Then the 3-chain a_1, d, b on Σ_g corresponds to the 3-chain a_0, a_1, b of Proposition 8.2, since $\phi(b) = -1$ by the homological coherence property. By assumption, $T_d, T_{a_1} \in H$, so it remains only to establish $\tilde{\Pi}(\Sigma_g \setminus \{b\}) \leq H$.

By hypothesis (4), the restriction \mathcal{N}' determines an arboreal filling network on $\Sigma_g \setminus \{b\}$. The same is therefore true on the surface $\overline{\Sigma_g \setminus \{b\}}$ obtained by filling in the boundary component corresponding to the left side of b (where b is oriented so that a_1, a_3 lie to the left). The surface $\overline{\Sigma_g \setminus \{b\}}$ is connected, since the hypothesis $i(d, b) = 1$ implies that d is nonseparating. We treat $\overline{\Sigma_g \setminus \{b\}}$ as a surface $\Sigma_{g-1,1}^1$, with the marked point corresponding to the filled-in left side of b .

We will show that $\tilde{\Pi}(\Sigma_g \setminus \{b\}) \leq H$ by appealing to Lemma 8.1. We must therefore show that T'_b is an element of H , and show that the image of the map

$$\langle T_a, a \in \mathcal{N}' \rangle \rightarrow \text{Mod}(\overline{\Sigma_g \setminus \{b\}})$$

contains the point-pushing subgroup $\pi_1(\overline{\Sigma_g \setminus \{b\}})$. Applying Corollary 2.7.2, we obtain $T_b^r \in H$. To exhibit $\pi_1(\overline{\Sigma_g \setminus \{b\}})$, we will appeal to Lemma 9.4. The element $T_{a_1} T_{a_3}^{-1}$ corresponds to an element $P(a_1) \in \pi_1(\overline{\Sigma_g \setminus \{b\}})$. By Lemma 9.4, it follows that the entire point-pushing subgroup $\pi_1(\overline{\Sigma_g \setminus \{b\}})$ is contained in the subgroup $\langle T_a, a \in \mathcal{N}' \rangle \leq H$. □

10 Linear systems in toric surfaces

The purpose of this section is to give a minimal account of the work of Crétois–Lang in [3]. We do not attempt to give a detailed summary of the theory of toric surfaces; the interested reader is referred to [3] and the references therein.

Consider the integer lattice $\mathbb{Z}^2 \subset \mathbb{R}^2$. A *lattice polygon* Δ is the convex hull of a finite collection $\{v_1, \dots, v_n\}$ of $n \geq 3$ elements $v_i \in \mathbb{Z}^2$, not all collinear. Given a polygon Δ which contains at least one lattice point in the interior $\text{int}(\Delta)$, the *adjoint polygon* Δ_a is defined to be the convex hull of $\text{int}(\Delta) \cap \mathbb{Z}^2$.

The following proposition is a concise summary of the correspondence between line bundles on toric surfaces and polygons. For details, see [3, Section 3]. In item (1), a *unimodular transformation* of \mathbb{R}^2 is an affine map $A: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ (necessarily invertible) such that $A\mathbb{Z}^2 = \mathbb{Z}^2$.

Proposition 10.1 *Let X be a smooth toric surface.*

- (1) *Associated to any nef line bundle \mathcal{L} on X is a convex lattice polygon $\Delta_{\mathcal{L}}$, well-defined up to unimodular transformations.*
- (2) *If \mathcal{L} is nef, then the roots of \mathcal{L} (i.e. the line bundles \mathcal{S} for which $n\mathcal{S} = \mathcal{L}$ for some integer n) are in correspondence with the dilates $\frac{1}{n}\Delta_{\mathcal{L}}$ for which $\frac{1}{n}\Delta_{\mathcal{L}}$ is a lattice polygon.*
- (3) *Suppose that \mathcal{L} is ample and that $\text{int}(\Delta_{\mathcal{L}}) \cap \mathbb{Z}^2$ is nonempty. Then the adjoint line bundle $\mathcal{L} \otimes K_X$ is nef, and $\Delta_{\mathcal{L} \otimes K_X} = (\Delta_{\mathcal{L}})_a$.*
- (4) *Let \mathcal{L} be ample. The genus $g(\mathcal{L})$ of a smooth $C \in |\mathcal{L}|$ is given by the formula*

$$g(\mathcal{L}) = \#(\text{int}(\Delta_{\mathcal{L}}) \cap \mathbb{Z}^2) = \#((\Delta_{\mathcal{L}})_a \cap \mathbb{Z}^2).$$

- (5) *Let \mathcal{L} be ample. A generic fiber $C \in |\mathcal{L}|$ is hyperelliptic if and only if $(\Delta_{\mathcal{L}})_a$ is a line segment.*

The following proposition indicates the connection between the divisibility properties of $\mathcal{L} \otimes K_X$ as an element of $\text{Pic}(X)$ (or after Proposition 10.1.2, the divisibility of $(\Delta_{\mathcal{L}})_a$), and the presence of invariant higher spin structures. It is a folklore theorem; see [17, Theorem 1.1] and [3, Proposition 2.7] for written accounts.

Proposition 10.2 *Let \mathcal{L} be an ample line bundle on a smooth toric surface X . For any r such that the adjoint line bundle $\mathcal{L} \otimes K_X$ admits a r th root in $\text{Pic}(X)$, there exists a (unique) $\mathbb{Z}/r\mathbb{Z}$ -valued spin structure ϕ preserved by the monodromy $\mu_{\mathcal{L}}$:*

$$\Gamma_{\mathcal{L}} \leq \text{Mod}(\Sigma_{g(\mathcal{L})})[\phi].$$

Proposition 10.1 suggests that it might be profitable to “model” a smooth $C \in |\mathcal{L}|$ on the lattice polygon $\Delta_{\mathcal{L}}$.

Construction 10.3 (*Inflation procedure*) Let Δ be a lattice polygon. Let $B(r, x)$ denote the open ball of radius r centered at $x \in \mathbb{R}^2$. Define the surface with boundary

$$\Delta^\circ := \Delta \setminus \bigcup_{v \in \text{int}(\Delta) \cap \mathbb{Z}^2} B(v, 1/4).$$

The *inflation* of Δ is the surface C_Δ obtained as the double of Δ° along its boundary. It is a closed oriented surface of genus $g = \#(\text{int}(\Delta) \cap \mathbb{Z}^2)$. In particular, for $\Delta = \Delta_{\mathcal{L}}$ for some ample \mathcal{L} , the inflation C_Δ has genus $g(\mathcal{L})$. See Fig. 8 for the example of $\mathcal{O}(6)$ on $\mathbb{C}\mathbb{P}^2$.

The first indication of the utility of the inflation procedure is provided by the following theorem of Crétois–Lang. For an inflation C_Δ , define an *A-curve* to be any simple closed curve on C_Δ that corresponds to the circle of radius $1/4$ centered at an interior lattice point of Δ .

Theorem 10.4 [3, Theorem 3] *Let \mathcal{L} be an ample line bundle on a smooth toric surface X . There is a homeomorphism $f: C_0 \rightarrow C_{\Delta_{\mathcal{L}}}$ identifying a smooth $C_0 \in |\mathcal{L}|$ with $C_{\Delta_{\mathcal{L}}}$, such that every A-curve $a \subset C_{\Delta_{\mathcal{L}}}$ is a vanishing cycle, and*

$$T_a \in \Gamma_{\mathcal{L}}.$$

Crétois–Lang also determine a second family of elements of $\Gamma_{\mathcal{L}}$ arising from the combinatorics of Δ . A *primitive integer segment* is a line segment $\sigma \subset \mathbb{R}^2$ whose endpoints lie on \mathbb{Z}^2 and whose interior is disjoint from \mathbb{Z}^2 . A primitive integer segment determines a line in \mathbb{R}^2 in the obvious way. When a lattice polygon Δ is fixed, it will be understood that a primitive integer segment connects lattice points $v, w \in \Delta \cap \mathbb{Z}^2$, and such that v and w do not lie along the same edge of Δ . Under the inflation procedure, a primitive integer segment corresponds to a simple closed curve. For a primitive integer segment σ , we write T_σ for the corresponding Dehn twist.

Suppose that Δ is a lattice polygon, let $d \geq 1$ be an integer. We say that Δ is *divisible by d* if the dilate $\frac{1}{d}\Delta$ is again a lattice polygon. If Δ is divisible by d , then after translating Δ so that one vertex lies in the sublattice $d\mathbb{Z}^2$, the remaining vertices do also. We write

$$\Delta(d) := \Delta \cap d\mathbb{Z}^2,$$

relative to any such embedding. The following is a combination of Propositions 7.13 and 7.16 of [3].

Theorem 10.5 (Crétois–Lang) *Let \mathcal{L} be an ample line bundle on a smooth toric surface X . Suppose that the adjoint polygon $(\Delta_{\mathcal{L}})_a$ is divisible by d . Suppose that σ is a primitive integer segment such that the line it generates intersects $(\Delta_{\mathcal{L}})_a(d)$. Then, with respect to the identification $f: C_0 \rightarrow C_{\Delta_{\mathcal{L}}}$ of Theorem 10.4, we have that σ is a vanishing cycle and $T_{\sigma} \in \Gamma_{\mathcal{L}}$.*

Taken together, Theorems 10.4 and 10.5 produce a large family of Dehn twists in $\Gamma_{\mathcal{L}}$. In the next section, we will see that they provide sufficiently many elements to satisfy the hypotheses of Theorem 9.5, which will lead to a proof of Theorem A.

11 Proof of Theorem A

Fix a toric surface X and an ample line bundle \mathcal{L} . This determines the polygons $\Delta_{\mathcal{L}}$ and $(\Delta_{\mathcal{L}})_a$, as well as the monodromy group $\Gamma_{\mathcal{L}}$. For convenience, we will drop reference to \mathcal{L} from the notation, and speak of Δ , Δ_a , Γ , etc. We also shorten notation for the inflation curve $C_{\Delta_{\mathcal{L}}}$, and refer simply to C instead.

By hypothesis, r is the highest root of the line bundle $\mathcal{L} \otimes K_X$. Proposition 10.1.2 implies that Δ_a is r -divisible. Our first objective is to find a network \mathcal{N} satisfying the hypotheses of Theorem 9.5. This will show all but the last assertion of Theorem A. Once this has been accomplished, we will see that the answer to Question 1.2 readily follows.

Genus hypotheses We first address the genus assumptions of Theorem 9.5. Recalling that Δ_a is assumed to be r -divisible, a calculation using Pick’s formula implies that for $r > 1$,

$$g \geq \frac{(r + 1)(r + 2)}{2}.$$

This shows that $g \geq 5$ for all $r > 1$ and that $g \geq r + 1$ for $r = 2d$ even. For $r = 2d = 4$, this gives $g \geq 15$, and for $r = 2d = 8$ this gives $g \geq 45$. In all cases, the hypothesis $g \geq g(r)$ of Theorem 9.5 holds.

The remaining assumption to be addressed is the requirement that $r < g - 1$. As noted in Remark 3.6, r must divide $2g - 2$, so we must only show that the cases $r = 2g - 2$ and $r = g - 1$ do not occur in the study of linear systems on toric surfaces. Suppose first that $r = 2g - 2$. This implies that the adjoint polygon Δ_a contains precisely g lattice points, but is also $2g - 2$ -divisible. This is an absurdity: let e be an edge of the lattice polygon $\frac{1}{2g-2}\Delta_a$; then the dilate $(2g - 2)e$ contains at least $2g - 1 > g$ lattice points. In the case $r = g - 1$, a similar analysis shows that in fact Δ_a must equal the $g - 1$ -fold dilation of a primitive integer segment. By Proposition 10.1.5, this implies that the general fiber of the linear system is hyperelliptic, which we have excluded from consideration.

Constructing the network \mathcal{N} Recall that according to Theorem 10.4, each integer point $v \in \Delta_a \cap \mathbb{Z}^2$ determines a vanishing cycle in Γ ; we introduce the notation $A(v)$ to refer to the curve associated to v . When we have a specific identification of Δ with a lattice polygon, we will use the notation $A(x, y)$ to refer to the A -curve at the integer point (x, y) . Similarly, given a primitive integer segment σ , we let $B(\sigma)$ denote the associated simple closed curve on C . When Δ is identified with a lattice polygon, we write $B((x, y), (z, w))$ for the B -curve associated to the primitive integer segment connecting (x, y) and (z, w) . We refer to these as A -curves and B -curves, respectively.

To define the network \mathcal{N} , it will be useful to introduce some terminology. Let σ be a primitive integer segment, and let $L(\sigma)$ be the line determined by σ . For an integer point v , we say that σ *points towards* v if $v \in L(\sigma)$. We also introduce the notion of a κ -*standard embedding*. Let κ be a vertex of Δ_a . A κ -standard embedding is an embedding of Δ into \mathbb{R}^2 such that κ corresponds to $(0, 0)$ and such that the edges of Δ_a incident to κ lie along the x and y axes. Any embedding $\Delta \subset \mathbb{R}^2$ can be made κ -standard by applying a suitable unimodular transformation. Following Proposition 10.1.1, we are free to apply unimodular transformations as needed.

Let κ be a vertex of Δ_a . Let \mathcal{N} be the network consisting of the following curves:

- (1) All A -curves.
- (2) The curve $B(\sigma)$, where σ is defined as follows. Let κ' be a vertex of Δ_a adjacent to κ . Let e' be the edge of Δ_a containing κ' and not containing κ , and let $w \in \partial\Delta_a$ be the integer point lying on e' that is connected to κ' by a primitive integer segment σ .
- (3) The curve $B(\tau)$ defined as follows. Under a κ -standard embedding of Δ , necessarily $(0, -1) \in \partial\Delta$. Since Δ_a is assumed to be d -divisible, the edge of Δ_a lying along the x -axis extends at least as far as $(d, 0)$. We take τ to be the primitive integer segment identified with $B((d, 0), (0, -1))$ in this embedding of Δ .




Fig. 8 Example: $(X, \mathcal{L}) = (\mathbb{C}P^2, \mathcal{O}(6))$; here $r = 3$. Left: the lattice polygons Δ and Δ_a (shaded). Right: the inflation construction, and the network \mathcal{N} , depicted in both shades of blue. The curves a_1, \dots, a_{10} of Theorems 9.5.2 and 9.5.3 are shown in the light shade of blue. Note the curve b (in red) is not part of the network, but does correspond to the curve b of Theorem 9.5.3 (color figure online)

- (4) All B -curves associated to primitive integer segments pointing towards κ , but such that the associated line does not pass through the interior of the segments σ or τ or the primitive integer segment connecting $(-1, 1)$ to $(0, 1)$.

See Fig. 8 for a picture of \mathcal{N} in the case of the line bundle $\mathcal{O}(6)$ on $\mathbb{C}P^2$.

Remark 11.1 As can be seen in Fig. 8, certain elements of \mathcal{N} are mutually isotopic. This harmless excess is introduced only to make the definition of \mathcal{N} more tidy.

In anticipation of an appeal to Theorem 9.5, we also define the subnetwork

$$\mathcal{N}' := \mathcal{N} \setminus \{A(0, 1)\}. \tag{12}$$

First properties of \mathcal{N} We first claim that \mathcal{N} is a network. Indeed, all A -curves are mutually disjoint. The set of primitive integer segments under consideration meet only at integer points in Δ_a , and hence the associated B -curves are also mutually disjoint. Suppose σ has endpoints v, w . Then $i(A(v), B(\sigma)) = i(A(w), B(\sigma)) = 1$, and $i(A(u), B(\sigma)) = 0$ for any other integer point u . Thus \mathcal{N} is a network.

Indeed, \mathcal{N} is a connected network, as follows from the description of $\Gamma_{\mathcal{N}}$ and $\Gamma_{\mathcal{N}'}$ given below.

Lemma 11.2 *The graph $\Gamma_{\mathcal{N}}$ has the homotopy type of S^1 , and $\Gamma_{\mathcal{N}'}$ is a tree, i.e. \mathcal{N}' is arboreal.*

Proof We first establish that $\Gamma_{\mathcal{N}}$ is connected. It suffices to show that every $c \in \mathcal{N}$ is connected to $A(\kappa)$. We first consider the case of an A -curve $A(v)$. If $v \in \Delta_a$ is some other integer point, there is a line segment L connecting v to

κ . This decomposes as a union of primitive integer segments σ_i based at the integer points v_j lying on L . Each such segment determines a B -curve in \mathcal{N} , and it is clear that there is a path from $A(v)$ to $A(\kappa)$ alternating between $B(\sigma_i)$ and $A(v_j)$. The argument for a B -curve (including the exceptional elements $B(\sigma)$ and $B(\tau)$) is similarly straightforward.

We next claim that the subnetwork

$$\mathcal{N}'' := \mathcal{N} \setminus B(\sigma)$$

is arboreal. The curves $B(\sigma)$ and $B(\tau)$ are the only B -curves in \mathcal{N} that do not lie on a line passing through κ . Thus the network consisting only of curves of type (1) and (4) is arboreal by construction. As $B(\tau)$ intersects only $A(d, 0)$, this shows that the network consisting of curves of type (1), (3), and (4) is also arboreal, but this network is \mathcal{N}'' by definition.

The curve $B(\sigma)$ intersects only the A -curves $A(\kappa')$ and $A(w)$. Thus $\Gamma_{\mathcal{N}}$ is obtained from the tree $\Gamma_{\mathcal{N}''}$ by adding one new vertex that is connected to two edges, so that $\Gamma_{\mathcal{N}} \simeq S^1$ as claimed.

It will follow from this that \mathcal{N}' is also arboreal. The path in $\Gamma_{\mathcal{N}''}$ connecting κ' to w follows the y -axis down to κ , then proceeds out along the line connecting κ to w ; in particular, it passes through the vertex $(0, 1)$. Thus, removing $A(0, 1)$ to create the network \mathcal{N}' removes the single circuit in $\Gamma_{\mathcal{N}}$, so that $\Gamma_{\mathcal{N}'}$ is a tree as claimed. □

We claim that \mathcal{N} is filling. This will be established in the next two lemmas. Recall the definition of Δ° from the definition of the inflation procedure in Construction 10.3.

Lemma 11.3 *Let $S \subset \Delta$ denote the union of all primitive integer segments associated to B -curves in \mathcal{N} . Then*

- (1) *Each component of $\Delta^\circ \setminus S$ is simply-connected.*
- (2) *For each component D of $\Delta^\circ \setminus S$, the intersection $D \cap \partial\Delta$ has at most one component.*

Proof We begin by observing that there are homotopy equivalences $\Delta^\circ \simeq \Delta \setminus (\Delta_a \cap \mathbb{Z}^2)$ and $\Delta^\circ \setminus S \simeq \Delta \setminus (S \cup (\Delta_a \cap \mathbb{Z}^2))$. It will be tidier to work with this latter space, and so we formulate our arguments in this setting.

Embed Δ into \mathbb{R}^2 and consider S as a planar graph contained in Δ . Basic properties of convexity imply that for any integer point $v \in \mathbb{Z}^2 \cap \Delta_a$, the line connecting v and κ does not intersect either of $B(\sigma)$ or $B(\tau)$. Hence this line determines a union of primitive integer segments in \mathcal{N} , and upon the removal of these segments over all v , there is an equality

$$\Delta \setminus (S \cup (\mathbb{Z}^2 \cap \Delta_a)) = \Delta \setminus S.$$

To prove (1), it therefore suffices to show that $H_1(\Delta \setminus S; \mathbb{Z}) = 0$. There is a map of pairs $f: (\Delta, \partial\Delta) \rightarrow (S^2, *)$, where $*$ $\in S^2$ is an arbitrary basepoint. f induces a homeomorphism

$$f: \Delta \setminus \partial\Delta \rightarrow S^2 \setminus \{*\}.$$

Since the segment $B(\tau)$ (among many others) intersects $\partial\Delta$, it follows that f induces a homotopy equivalence

$$f: \Delta \setminus S \rightarrow S^2 \setminus f(S),$$

and hence there is an isomorphism

$$f_*: \tilde{H}_1(\Delta \setminus S; \mathbb{Z}) \rightarrow \tilde{H}_1(S^2 \setminus f(S); \mathbb{Z}).$$

By Alexander duality, $\tilde{H}_1(S^2 \setminus f(S); \mathbb{Z}) \cong \tilde{H}^0(f(S); \mathbb{Z}) = 0$, the latter holding because S is connected by construction. This proves (1).

For (2), consider the subconfiguration $S' \subset S$ consisting of all primitive integer segments lying along a line connecting κ to any integer point $v \in \partial\Delta$. This provides a subdivision of Δ into convex sets, each of which has a vertex at κ . Convexity then implies that each component D'_i of $\Delta \setminus S'$ intersects $\partial\Delta$ in at most one component. The subdivision of Δ induced by S is a refinement of that induced by S' . Since all segments in S that intersect $\partial\Delta$ are elements of S' , there is an equality

$$\partial\Delta \setminus (S \cap \partial\Delta) = \partial\Delta \setminus (S' \cap \partial\Delta).$$

Thus each component of $\partial\Delta \setminus (S \cap \partial\Delta)$ corresponds to a *distinct* component of $\Delta \setminus S$, and (2) follows. □

Lemma 11.4 *Each component of $C \setminus \mathcal{N}$ is simply-connected, i.e. \mathcal{N} is filling.*

Proof By construction, the “deflation” map $p: C \rightarrow \Delta$ takes components of $C \setminus \mathcal{N}$ to components of $\Delta \setminus S$, where S continues to denote the union of all primitive integer segments associated to B -curves in \mathcal{N} . This map on components is at most 2-to-1, and is *exactly* 2-to-1 in the case where the component D of $\Delta \setminus S$ does not contain a lattice point in its interior and does not intersect $\partial\Delta$. In this 2-to-1 case, each component \tilde{D}_1, \tilde{D}_2 of $C \setminus \mathcal{N}$ is mapped homeomorphically by p onto the component of $\Delta \setminus S$ corresponding to D . Since D is assumed to contain no lattice points in the interior, it follows that the corresponding component of $\Delta \setminus S$ is simply-connected, and hence \tilde{D}_1, \tilde{D}_2 are as well.

Lemma 11.3.1 implies that no component of $\Delta \setminus S$ contains an interior lattice point, and so it remains only to be seen that every component of $C \setminus \mathcal{N}$ that

corresponds to a component of $\Delta \setminus S$ intersecting $\partial\Delta$ is simply-connected. Let $\tilde{D} \subset C \setminus \mathcal{N}$ be such a component, and let D be the corresponding component of $\Delta^\circ \setminus S$. Observe that \tilde{D} is constructed by attaching two copies of D along $D \cap \partial\Delta$. It follows that \tilde{D} is simply-connected if and only if $D \cap \partial\Delta$ is connected. The result now follows by Lemma 11.3.2. \square

Applicability of Theorem 9.5 It remains to verify the properties (1)–(4) of Theorem 9.5. By Theorem 10.4, for an A -curve $A(v)$, the associated Dehn twist $T_{A(v)} \in \Gamma$. By Theorem 10.5, any curve $B(\xi) \in \mathcal{N}$ arising from a primitive integer segment ξ also satisfies $T_{B(\xi)} \in \Gamma$. It follows from the definitions that any curve c in any connected network is necessarily non-separating. For a nonseparating curve $c \subset C$, the Dehn twist $T_c \in \Gamma$ only if the associated spin structure satisfies $\phi(c) = 0$. Hence (1) holds.

For (2), we take a κ -standard embedding of Δ . It is now easy to find a collection of curves \mathcal{S}_{2r+4} determining the configuration $\mathcal{D}_{2r+3} \cup \{\Delta_1\}$ of Corollary 2.7. We take $a = B((0, 0), (0, -1))$ and $a' = B((0, 0), (0, 1))$. Since Δ_a is assumed to be r -divisible, the edge of Δ_a lying along the x -axis extends at least as far as $(r, 0)$. For $1 \leq k \leq r + 1$, we can therefore take c_{2k-1} to be $A(0, k - 1)$, and for $1 \leq k \leq r$, we take c_{2k} to be $B((k - 1, 0), (k, 0))$. We take $a_{r+1} = B(\tau) = B((r, 0), (0, -1))$. The segments connecting $(0, -1), (0, 0), (1, 0), \dots, (d, 0), (0, -1)$ separate Δ into two components, hence under the inflation procedure, the associated B -curves separate C . From the construction it is clear that the curves bound a subsurface of genus 0 with $r + 2$ boundary components, as required for the configuration $\mathcal{D}_{2r+3} \cup \{\Delta_1\}$ of Corollary 2.7.

For (3), we observe that from the construction, the Δ_0 curve of the configuration \mathcal{D}_{2r+3} corresponds to $b := B((-1, 1), (0, 1))$ on Δ . One sees that $A(0, 1)$ intersects this curve, and is an element of \mathcal{N} as needed.

For (4), we begin by observing that *only* the element $A(0, 1) \in \mathcal{N}$ intersects $B((-1, 1), (0, 1))$. Enumerate the components of $C \setminus \mathcal{N}$ as $\{D_i\}$. We claim that there are exactly three disks D_1, D_2, D_3 in $C \setminus \mathcal{N}$ with boundary lying on $A(0, 1)$, and that $b \subset \overline{D_1} \subset C$. Indeed, using the notation of item (2) in the definition of \mathcal{N} , the disks D_2 and D_3 arise via inflation from the component of $\Delta \setminus S$ bounded by the triangle formed by e', σ , and the primitive integer segment(s) connecting κ to w . Neither D_2 nor D_3 intersects b , and the only curve in \mathcal{N} intersecting b is $A(0, 1)$; this implies that $b \subset \overline{D_1}$ as claimed.

Thus, in $(C \setminus \mathcal{N}) \cup A(0, 1)$, the disks D_2 and D_3 are joined into a single disk D^+ , while D_1 has two portions of its boundary joined to create an annulus with core curve b . Upon passing to $(C \setminus \{b\}) \setminus \mathcal{N}'$, this annulus is cut open to create two annular regions bounded by b , while the disks D^+ and D_i for $i \geq 4$ are unaffected. Thus \mathcal{N}' does determine a filling network on $C \setminus \{b\}$ as required. Arboreality of \mathcal{N}' was established in Lemma 11.2.

From admissible twists to vanishing cycles In order to address Question 1.2, it is necessary to better understand the relationship between admissible twists and vanishing cycles. A first remark is that any vanishing cycle is necessarily an admissible curve, so it remains only to show the converse. We observe that if α is a loop in $\mathcal{M}(\mathcal{L})$ based at C_0 that determines a vanishing cycle, then any conjugate $\beta\alpha\beta^{-1}$ also determines a vanishing cycle. To complete the argument, it therefore suffices to establish the following claim.

Lemma 11.5 *Let a be any admissible curve on C_0 . Then T_a is conjugate in Γ to some twist T_c for c a vanishing cycle.*

Proof An admissible curve a determines a vertex in the graph $\mathcal{C}_\phi^1(C_0)$ of Sect. 7. Theorems 10.4 and 10.5 together imply that Γ has a generating set consisting entirely of vanishing cycles. Thus the set of vertices in $\mathcal{C}_\phi^1(C_0)$ corresponding to vanishing cycles is nonempty.

We claim that if $a \in \mathcal{C}_\phi^1(C_0)$ is adjacent to some vanishing cycle c , then a is also a vanishing cycle. Indeed, the condition that a and c are adjacent in $\mathcal{C}_\phi^1(C_0)$ is equivalent to $i(a, c) = 1$, and hence by the braid relation,

$$T_a = (T_c T_a) T_c (T_c T_a)^{-1}.$$

As $T_c, T_a \in \Gamma$ by the first part of Theorem A, the above observation implies that T_a is a vanishing cycle. The claim now follows from the connectivity of $\mathcal{C}_\phi^1(C_0)$ established in Lemma 7.5.

This concludes the proof of Theorem A. □

Acknowledgements The author would like to extend his warmest thanks to R. Crétois and L. Lang for helpful discussions of their work. He would also like to acknowledge C. McMullen for some insightful comments on a preliminary draft, and M. Nichols for a productive conversation. A special thanks is due to an anonymous referee for a very careful reading of the preprint and for many useful suggestions, both mathematical and expository.

References

1. Chillingworth, D.: Winding numbers on surfaces. I. *Math. Ann.* **196**, 218–249 (1972)
2. Chillingworth, D.: Winding numbers on surfaces. II. *Math. Ann.* **199**, 131–153 (1972)
3. Crétois, R., Lang, L.: The Vanishing Cycles of Curves in Toric Surfaces I. Preprint. [arXiv:1701.00608v2.pdf](https://arxiv.org/abs/1701.00608v2) (2017)
4. Crétois, R., Lang, L.: The Vanishing Cycles of Curves in Toric Surfaces II. Preprint. [arXiv:1706.07252.pdf](https://arxiv.org/abs/1706.07252) (2017)
5. Dieudonné, J.: *Sur les groupes classiques*. Hermann, Paris (1973). Troisième édition revue et corrigée, Publications de l'Institut de Mathématique de l'Université de Strasbourg, VI, Actualités Scientifiques et Industrielles, No. 1040
6. Donaldson, S.: Polynomials, vanishing cycles and Floer homology. In: *Mathematics: Frontiers and Perspectives*, pp. 55–64. American Mathematical Society, Providence (2000)

7. Farb, B., Margalit, D.: A Primer on Mapping Class Groups, Volume 49 of Princeton Mathematical Series. Princeton University Press, Princeton, NJ (2012)
8. Grove, L.C.: Classical Groups and Geometric Algebra, Volume 39 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI (2002)
9. Humphries, S., Johnson, D.: A generalization of winding number functions on surfaces. *Proc. Lond. Math. Soc.* (3) **58**(2), 366–386 (1989)
10. Johnson, D.: An abelian quotient of the mapping class group \mathcal{I}_g . *Math. Ann.* **249**(3), 225–242 (1980)
11. Johnson, D.: Spin structures and quadratic forms on surfaces. *J. Lond. Math. Soc.* (2) **22**(2), 365–373 (1980)
12. Johnson, D.: A survey of the Torelli group. In: *Low-Dimensional Topology* (San Francisco, Calif., 1981), Volume 20 of Contemporary Mathematics, pp. 165–179. American Mathematical Society, Providence (1983)
13. Johnson, D.: The structure of the Torelli group. II. A characterization of the group generated by twists on bounding curves. *Topology* **24**(2), 113–126 (1985)
14. Johnson, D.: The structure of the Torelli group. III. The abelianization of \mathcal{T} . *Topology* **24**(2), 127–144 (1985)
15. Matsumoto, M.: A presentation of mapping class groups in terms of Artin groups and geometric monodromy of singularities. *Math. Ann.* **316**(3), 401–418 (2000)
16. Putman, A.: A note on the connectivity of certain complexes associated to surfaces. *Enseign. Math.* (2) **54**(3–4), 287–301 (2008)
17. Salter, N.: On the Monodromy Group of the Family of Smooth Plane Curves. Preprint. [arXiv:1610.04920.pdf](https://arxiv.org/abs/1610.04920) (2016)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.